

a
Device-based recomputation vs GBD 2023 reported (pooled RR source)
DALYs
Deaths
Global
Brazil
China
Germany
India
Japan
Nigeria
USA
x5.9
x4.0
x6.5
x6.1
x5.7
x4.2
x8.6
x5.5
x5.7
x3.8
x6.4
x5.8
x4.8
x5.3
x8.5
x5.8
100
10,000
1,000,000
100,000,000
LPA-attributable burden, 2022 (log scale)
b
Gap by SDI quintile (2022; pooled RR source)
PAF (attributable fraction of deaths)
0.20
0.15
0.10
0.05
0.00
x9.3
x5.1
x5.3
x4.9
x5.8
Low SDI (n=50)
Low-middle SDI (n=25)
Middle SDI (n=18)
High-middle SDI (n=23)
High SDI (n=88)
GBD 2023 reported
This study
c
Fold (this study vs GBD 2023) by country and cause (2022; 204 countries, 8 causes; pooled RR source)
Lower extremity peripheral arterial disease
Breast cancer
Tuberculosis
Colon and rectum cancer
Chronic kidney disease
Diabetes mellitus
Stroke
Ischemic heart disease
This study / GBD 2023
40
20
10
5
2
1
Low SDI (n=50)
Low-middle SDI (n=25)
Middle SDI (n=18)
High-middle SDI (n=23)
High SDI (n=88)
Colour scale clipped at fold = 43 (33 of 1632 tiles, 2.0%); all folds exceed 1.
d
Gap fully explained by two definitional differences (Global, 2022; residual +0.72%)
PAF (attributable fraction of deaths)
0.200
0.150
0.100
0.050
0.000
0.174
fold 5.72
0.020
x0.11
0.030
x1.54
x1.007
0.030
fold 1.00
This study (primary)
RR magnitude
TMREL definition
Residual
GBD 2023 (reported)
Study estimate
RR magnitude
TMREL definition
Residual
GBD 2023 (reported)