Table 2. Hazard ratios (95% CI) for device-measured activity (MIMS/day) and mortality, NHANES 2011-2014 (n = 7,007)

outcome	model	term	hr	p
All-cause mortality	Model 1 (minimally adjusted)	Q2	0.67 (0.54, 0.83)	<0.001
All-cause mortality	Model 1 (minimally adjusted)	Q3	0.52 (0.39, 0.69)	<0.001
All-cause mortality	Model 1 (minimally adjusted)	Q4	0.49 (0.34, 0.69)	0.002
All-cause mortality	Model 1 (minimally adjusted)	P for trend	-	<0.001
All-cause mortality	Model 1 (minimally adjusted)	Per IQR (4879 MIMS/day)	0.62 (0.53, 0.72)	<0.001
All-cause mortality	Model 2 (fully adjusted)	Q2	0.66 (0.53, 0.81)	<0.001
All-cause mortality	Model 2 (fully adjusted)	Q3	0.52 (0.39, 0.69)	<0.001
All-cause mortality	Model 2 (fully adjusted)	Q4	0.48 (0.34, 0.68)	0.003
All-cause mortality	Model 2 (fully adjusted)	P for trend	-	<0.001
All-cause mortality	Model 2 (fully adjusted)	Per IQR (4879 MIMS/day)	0.61 (0.52, 0.71)	<0.001
Cardiovascular mortality	Model 1 (minimally adjusted)	Q2	0.62 (0.42, 0.93)	0.054
Cardiovascular mortality	Model 1 (minimally adjusted)	Q3	0.49 (0.28, 0.86)	0.053
Cardiovascular mortality	Model 1 (minimally adjusted)	Q4	0.32 (0.14, 0.71)	0.018
Cardiovascular mortality	Model 1 (minimally adjusted)	P for trend	-	<0.001
Cardiovascular mortality	Model 1 (minimally adjusted)	Per IQR (4879 MIMS/day)	0.56 (0.42, 0.75)	<0.001
Cardiovascular mortality	Model 2 (fully adjusted)	Q2	0.63 (0.42, 0.94)	0.055
Cardiovascular mortality	Model 2 (fully adjusted)	Q3	0.51 (0.29, 0.90)	0.064
Cardiovascular mortality	Model 2 (fully adjusted)	Q4	0.34 (0.15, 0.76)	0.028
Cardiovascular mortality	Model 2 (fully adjusted)	P for trend	-	0.001
Cardiovascular mortality	Model 2 (fully adjusted)	Per IQR (4879 MIMS/day)	0.57 (0.42, 0.78)	<0.001

Weighted Cox proportional hazards models (svycoxph); reference = lowest quartile. Model 1 adjusts for age, sex, race, education, poverty-income ratio, smoking, alcohol, diabetes, cardiovascular history; Model 2 additionally adjusts for BMI and systolic blood