

期末考试第二部分

一、find_name_value函数

1. 功能描述：

find_name_value函数用于解析并分离数据文件名称中的变量名和对应的数值。用户调用find_name_value函数，输入代表文件名的字符串，函数将返回一个元组 (name,value)

2. 设置测试用例：

1.1 标准正数

```
assert find_name_value('temorary0.1') == ('temporary', 0.1)
```

1.2 标准负数

```
assert find_name_value('melody0.5n') == ('melody', -0.5)
```

1.3 无符号整数

```
assert find_name_value('temp2010') == ('temp', 2010)
```

1.4 带符号正数

```
assert find_name_value('tp+0.2') == ('tp', 0.2)
```

1.5 带符号负数

```
assert find_name_value('tp-0.1') == ('tp', -0.1)
```

2.1 纯变量名，无数字

```
assert find_name_value('tanpeng') == ('tanpeng', None)
```

2.2 小数点开头的数字

```
assert find_name_value('rongrui.2010') == ('rongrui', 0.2010)
```

2.3 数值为零

```
assert find_name_value('life0') == ('life', 0.0)
```

3.1 空字符串

```
assert find_name_value('') == ('', None)
```

3.2 文件名以数字开头

```
assert find_name_value('123value') == ('123value', None)
```

3. 利用函数找到目标文件名的值

1. 'phi0.1 xN14.2 kappa0.5n'

```
print(find_name_value('phi0.1 xN14.2 kappa0.5n'))  
('phi', 0.1)
```

2. 'a1_b14n_n0_c02'

```
('a', 1.0)
```

二、用Matlab作图

1. Matlab代码

```
% 环面参数  
R = 3; % 大圆半径  
r = 1; % 小圆半径  
  
% 定义参数范围  
theta = linspace(0, 2*pi, 100); % 小圆的角度范围  
phi = linspace(0, 2*pi, 100); % 大圆的角度范围  
  
% 创建网格  
[Theta, Phi] = meshgrid(theta, phi);  
  
% 计算 x, y, z 坐标  
X = (R + r * cos(Theta)) .* cos(Phi);
```

```

Y = (R + r * cos(Theta)) .* sin(Phi);
Z = r * sin(Theta);

% 绘制环面
figure;
surf(X, Y, Z);

% 图像美化
shading interp;      % 插值填充
colormap jet;        % 颜色映射
axis equal;          % 等比例坐标
title('环面 (Torus) 的三维图像');
xlabel('X');
ylabel('Y');
zlabel('Z');

```

! 结果图(<https://users/desktop/结果图.fig>)

三、用Mathematica求值

1. 求无穷级数的和 $\left[\sum_{n=1}^{\infty} \frac{1}{n^3 + n^2} \right]$

```
Sum[1/(n^3 + n^2), n, 1, Infinity{ }]
```

结果为： $\left[\frac{\pi^2}{6} - 1 \approx 0.644934 \right]$

2. 求定积分的值 $\left[\int_0^{\infty} \frac{\sqrt{x} \ln x}{(x+1)^2} dx \right]$

```
Integrate[Sqrt[x] * Log[x] / (x+1)^2, {x, 0, Infinity}]
```

结果为： $\left[-\frac{\pi}{4} \approx -0.785398 \right]$

四、用markdown写出文本内容

Q: Find the solution of the following equation with respect to (θ) :

A: $[A \cos \theta + B \sin \theta + C = 0]$

let $(x_1 = \cos \theta)$ and $(x_2 = \sin \theta)$, then the solution is given by the intersection of the circle and the line:

$$[x_1^2 + x_2^2 = 1]$$

$$[A x_1 + B x_2 + C = 0]$$

We reformulate the equations in a parametric form:

$$[|x|^2 = 1]$$

$$[x(t) = a + t b]$$

where $(x = (x_1, x_2))$, $(a = (0, -C / B))$, $(b = (-C / A, C / B))$, and (t) is a parameter. The intersection points satisfy the following equation:

$[|a + t b|^2 = 1]$ which can be solved for (t) to find the intersection points:

$$[t_{1,2} = \frac{-a \cdot b \pm \sqrt{(a \cdot b)^2 - |b|^2 (|a|^2 - 1)}}{|b|^2}]$$