

## PART 2

### Q1.

详见 `part2_q1.ipynb`

### Q2.

Input:

```
Integrate[(Sin[x] - Sin[3*x] + Sin[5*x])/(Cos[x] + Cos[3*x] + Cos[5*x]),x]
```

Output:

```
-Log[Cos[x]]
```

---

In[1]=

Integrate

(

Sin

[

x

]

-

Sin

[

3 \* x

]

+

Sin

[

5 \* x

]

)

/

(

Cos

[

x

]

+

Cos

[

3 \* x

]

+

Cos

[

5 \* x

]

)

,

x

]

积分

正弦

正弦

正弦

余弦

余弦

余弦

Out[1]=

-Log[Cos[x]]

绘图

x 的导数

x 的积分

转换为指数形式

更多...

🔄

⚙️

💬


x

y


z

### Q3.

```
[x,y,z]=meshgrid(-1:0.1:1); isosurface(x,y,z,x.^2+y.^2+z.^2-1,0); axis equal;  
colormap summer
```



```
[x,y,z] = meshgrid(-4:0.1:4,-2:0.1:2,-2:0.1:2); a = 2; b = 1; v = (x.^2/a^2 +  
y.^2/b^2 + z.^2/b^2); isosurface(x,y,z,v,1); axis equal; colormap([1 0 0])
```



Q4.

## Lorenz Attractor

The Lorenz attractor is an [attractor](#) that arises in a simplified system of equations describing the two-dimensional flow of fluid. In the early 1960s, Lorenz accidentally discovered the chaotic behavior of this system when he found that, for a simplified system, periodic solutions of the form

$$\psi = \psi_0 \sin\left(\frac{\pi ax}{H}\right) \sin\left(\frac{\pi z}{H}\right)$$

$$\theta = \theta_0 \cos\left(\frac{\pi ax}{H}\right) \sin\left(\frac{\pi z}{H}\right)$$

grew for Rayleigh numbers larger than the critical value,  $Ra > Ra_c$ . Furthermore, vastly different results were obtained for very small changes in the initial values, representing one of the earliest discoveries of the so-called [butterfly effect](#).

Lorenz obtained the simplified equations

$$\dot{X} = \sigma(Y - X)$$

$$\dot{Y} = X(\rho - Z) - Y$$

$$\dot{Z} = XY - \beta Z$$

now known as the Lorenz equations.