

Accounting for Covariates (i.e. nuisances) with Multivariate Permutation tests

Livio Finos

University of Padova

May 16, 2022



(minimal) Bibliography

- Anderson M. Winkler, Gerard R. Ridgway, Matthew A. Webster, Stephen M. Smith, Thomas E. Nichols (2014) Permutation inference for the general linear model. *NeuroImage*, 92, doi.org/10.1016/j.neuroimage.2014.01.060
- Sara Kherad-Pajouh, Olivier Renaud (2010) An exact permutation method for testing any effect in balanced and unbalanced fixed effect ANOVA. *Computational Statistics and Data Analysis*. doi:10.1016/j.csda.2010.02.015
- Aldo Solari, Livio Finos and Jelle Goeman (2014) Rotation-based multiple testing in the multivariate linear model. *Biometrics*, 70, doi.org/10.1111/biom.12238



Motivating example

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients
- $p = 12625$ gene expression profiles
- covariate of interest: group (B or T-cell type patients)
- differentially expressed genes between the two groups?

$$\text{gene} = \gamma_0 + \beta \cdot \text{group} + \text{error}$$



Motivating example

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients
- $p = 12625$ gene expression profiles
- covariate of interest: group (B or T-cell type patients)
- differentially expressed genes between the two groups?

$$\text{gene} = \gamma_0 + \beta \cdot \text{group} + \text{error}$$

$$\begin{bmatrix} \text{gene}_1 \\ \vdots \\ \text{gene}_p \end{bmatrix} = \begin{bmatrix} \gamma_{01} \\ \vdots \\ \gamma_{0p} \end{bmatrix} + \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} \cdot \text{group} + \begin{bmatrix} \text{error}_1 \\ \vdots \\ \text{error}_p \end{bmatrix}$$



Multivariate linear model

$$\mathbf{Y} = \mathbf{1}\mathbf{G}_{1 \times p} + \mathbf{X}\mathbf{B}_{1 \times p} + \mathbf{E}$$

- \mathbf{Y} : $(n \times p)$ matrix of responses
- \mathbf{X} : $(n \times 1)$ matrix of covariates (group)
- $\mathbf{E} \sim (\mathbf{0}_{n \times p}, \mathbf{I}_n \otimes \mathbf{\Sigma})$ matrix of errors
 $\mathbf{\Sigma}$: $(p \times p)$ gene-gene covariance matrix

Marginal model (j -th gene)

$$\mathbf{y}_j = \mathbf{1}\gamma_{0j} + \mathbf{X}\beta_j + \varepsilon_j, \quad \varepsilon_j \sim (\mathbf{0}_{n \times 1}, \sigma_j^2 \mathbf{I}_n)$$

Multiple hypotheses

$$H_j : \beta_j = \mathbf{0}, \quad \forall \gamma_j \quad j = 1, \dots, p$$



The Multiple Testing Problem

$$\begin{bmatrix} \text{gene}_1 \\ \vdots \\ \text{gene}_p \end{bmatrix} = \begin{bmatrix} \gamma_{01} \\ \vdots \\ \gamma_{0p} \end{bmatrix} + \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} \cdot \text{group} + \begin{bmatrix} \text{error}_1 \\ \vdots \\ \text{error}_p \end{bmatrix}$$



The Multiple Testing Problem

$$\begin{bmatrix} \text{gene}_1 \\ \vdots \\ \text{gene}_p \end{bmatrix} = \begin{bmatrix} \gamma_{01} \\ \vdots \\ \gamma_{0p} \end{bmatrix} + \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} \cdot \text{group} + \begin{bmatrix} \text{error}_1 \\ \vdots \\ \text{error}_p \end{bmatrix}$$

$H_{0j} : \beta_j = 0$ (= two groups are equal) $j = 1, \dots, p$



The Multiple Testing Problem


$$\begin{bmatrix} \text{gene}_1 \\ \vdots \\ \text{gene}_p \end{bmatrix} = \begin{bmatrix} \gamma_{01} \\ \vdots \\ \gamma_{0p} \end{bmatrix} + \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} \cdot \text{group} + \begin{bmatrix} \text{error}_1 \\ \vdots \\ \text{error}_p \end{bmatrix}$$

$H_{0j} : \beta_j = 0$ (= two groups are equal) $j = 1, \dots, p$

one p-value for each gene (p)



Type I error if not correcting for multiple testing



... a very common problem

The problem of multiplicity control arises when more than one (statistical) hypothesis is tested (12625 in this case, one for each gene).

This problem is very common in many (other) fields of medicine:

- clinical trials with multiple endpoints
- neuroimaging experiments

and in many other fields like:

- psycho-sociological
- ecological
- quality control
- many others...



Familywise Error Rate (FWER)

FWER:

probability of one or more false positives (i.e. wrong rejections)

The most well-known method controlling the FWER:

Holm procedure (Holm, 1979).

Based on step-wise application of Bonferroni inequality:

hence: reject genes with $p_i \leq \alpha / (\# \text{ of genes})$, $i = 1, \dots, 12625$



Familywise Error Rate (FWER)

FWER:

probability of one or more false positives (i.e. wrong rejections)

The most well-known method controlling the FWER:

[Holm](#) procedure (Holm, 1979).

Based on step-wise application of Bonferroni inequality:

hence: reject genes with $p_i \leq \alpha / (\# \text{ of genes})$, $i = 1, \dots, 12625$

Is valid for every form of [dependence](#) among p-values.



Familywise Error Rate (FWER)

FWER:

probability of one or more false positives (i.e. wrong rejections)

The most well-known method controlling the FWER:

Holm procedure (Holm, 1979).

Based on step-wise application of Bonferroni inequality:

hence: reject genes with $p_i \leq \alpha / (\# \text{ of genes})$, $i = 1, \dots, 12625$

Is valid for every form of **dependence** among p-values.

BUT becomes **very conservative** when dependencies are strong.



Familywise Error Rate (FWER)

FWER:

probability of one or more false positives (i.e. wrong rejections)

The most well-known method controlling the FWER:

[Holm](#) procedure (Holm, 1979).

Based on step-wise application of Bonferroni inequality:

hence: reject genes with $p_i \leq \alpha / (\# \text{ of genes})$, $i = 1, \dots, 12625$

Is valid for every form of [dependence](#) among p-values.

BUT becomes [very conservative](#) when dependencies are strong.

Permutation tests are often a solution.



Outline

1 Dealing with dependence among tests

2 Confounding



(Multivariate) Permutation tests

Under $H_0 : \bigcap_{j=1}^p H_j$,

$$\mathbf{Y} \stackrel{d}{=} \Pi \mathbf{Y}$$

for every permutation matrix Π (*null-invariant transformation*) i.e.

Exchangeability: $f(\mathbf{Y}) = f(\Pi \mathbf{Y})$



(Multivariate) Permutation tests

Under $H_0 : \bigcap_{j=1}^p H_j$,

$$\mathbf{Y} \stackrel{d}{=} \Pi \mathbf{Y}$$

for every permutation matrix Π (*null-invariant transformation*) i.e.

Exchangeability: $f(\mathbf{Y}) = f(\Pi \mathbf{Y})$

When it holds

re-sampling null datasets is possible

$$n\text{-vector } \mathbf{X} = \begin{cases} -1/n_B & \text{if B patient,} \\ +1/n_T & \text{if T patient.} \end{cases}$$

(vector of) Test Statistic: $t_{obs} = \mathbf{X}^T \mathbf{Y}$

(vector of) Test Statistic of permuted data: $t_{\Pi} = \mathbf{X}^T \Pi \mathbf{Y}$



Joint distributions of P-values

- Observed statistic is any among possible permutations ($\Pi = \mathbb{I}$)
- compute test statistics on every possible permutation or sample them (e.g. 10 000 random permutations).



Joint distributions of P-values

- Observed statistic is any among possible permutations ($\Pi = \mathbb{I}$)
- compute test statistics on every possible permutation or sample them (e.g. 10 000 random permutations).
- This provides the joint distribution of the test statistics




Joint distributions of P-values

- Observed statistic is any among possible permutations ($\Pi = \mathbb{I}$)
- compute test statistics on every possible permutation or sample them (e.g. 10 000 random permutations).
- This provides the joint distribution of the test statistics
- Compute the joint distribution of p-values from joint dist. of test statistic
(compute p-values for observed test stat. and for all permuted test stats.)



Joint distributions of P-values



- Hypothesis:

$$H : H_1 \cap H_2$$

- min-p test:

$$\Pr_H \left\{ \min(p_1, p_2) < c \right\} \leq \alpha$$

- Bonferroni inequality:

$$c = \alpha/2$$



Holm vs Westfall & Young min-p



Holm vs Westfall & Young min-p

Holm: Repeat Bonferroni inequality,



Holm vs Westfall & Young min-p

Holm: Repeat Bonferroni inequality,
Westfall & Young: Repeat min-p step,



Holm vs Westfall & Young min-p

Holm: Repeat Bonferroni inequality,
Westfall & Young: Repeat min-p step,

both provides a list of adjusted p-values

W&Y is usually more powerful

(i.e. lower adjusted p-values, more rejections) than Holm



Holm vs Westfall & Young min-p

Holm: Repeat Bonferroni inequality,
Westfall & Young: Repeat min-p step,

both provides a list of adjusted p-values

W&Y is usually more powerful

(i.e. lower adjusted p-values, more rejections) than Holm

this is because permutation tests account for dependence.



Holm vs Westfall & Young min-p

Holm: Repeat Bonferroni inequality,
Westfall & Young: Repeat min-p step,

both provides a list of adjusted p-values

W&Y is usually more powerful

(i.e. lower adjusted p-values, more rejections) than Holm

this is because permutation tests account for dependence.

You can perform it on R using

`library(multtest)` or `library(flip)`



Outline

1 Dealing with dependence among tests

2 Confounding



Motivating example

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients
- $p = 12625$ genes
- covariate of interest: group (B or T-cell type patients)
- **confounder**: sex
- differentially expressed genes between the two groups?

$$\text{gene} = \gamma_0 + \gamma_1 \cdot \text{sex} + \beta \cdot \text{group} + \text{error}$$

$$\begin{bmatrix} \text{gene}_1 \\ \vdots \\ \text{gene}_p \end{bmatrix} = \begin{bmatrix} \gamma_{01} \\ \vdots \\ \gamma_{0p} \end{bmatrix} + \begin{bmatrix} \gamma_{11} \\ \vdots \\ \gamma_{1p} \end{bmatrix} \cdot \text{sex} + \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} \cdot \text{group} + \begin{bmatrix} \text{error}_1 \\ \vdots \\ \text{error}_p \end{bmatrix}$$



Multivariate linear model

$$\mathbf{Y} = \mathbf{Z}\mathbf{G} + \mathbf{X}\mathbf{B} + \mathbf{E}$$

- \mathbf{Y} : $(n \times p)$ matrix of responses
- \mathbf{X} : $(n \times q)$ matrix of covariates (group)
- \mathbf{Z} : $(n \times 2)$ intercept and sex
- $\mathbf{E} \sim (\mathbf{0}_{n \times p}, \mathbf{I}_n \otimes \mathbf{\Sigma})$ matrix of errors
 $\mathbf{\Sigma}$: $(p \times p)$ gene-gene covariance matrix

Marginal model (j -th gene)

$$\mathbf{y}_j = \mathbf{Z}\boldsymbol{\gamma}_j + \mathbf{X}\boldsymbol{\beta}_j + \boldsymbol{\varepsilon}_j, \quad \boldsymbol{\varepsilon}_j \sim (\mathbf{0}_{n \times 1}, \sigma_j^2 \mathbf{I}_n)$$

Multiple hypotheses

$$H_j : \boldsymbol{\beta}_j = \mathbf{0}, \quad \forall \boldsymbol{\gamma}_j \quad j = 1, \dots, p$$




Ignoring the confounders


Drop **Z** from the model:

$$\mathbf{Y} = \mathbf{X}\mathbf{B} + \mathbf{E}$$

false negative ($\beta \neq 0, \hat{\beta} \approx 0$)



false positive ($\beta = 0, \hat{\beta} \neq 0$)



Goal

Permutation methods are very useful in multiple testing since they easily **deal with dependencies** even when $p \gg n$. e.g.

- Westfall & Young min-p (controls the FamilyWise Error Rate)
- Meinshausen method (controls the proportion of False Rejections)
- Statistical NonParametric Mapping (SnPM, controls the FWER at cluster-level)

There are no standard solutions **accounting for covariates**
Permutation of the observed response \mathbf{Y} is not a valid solution.
We need a valid method to account for confounders.



Adjusting for confounders

Residuals of \mathbf{Z} i.e.

Project \mathbf{Y} into the subspace \perp to $\text{span}(\mathbf{Z})$:

$$\mathbf{Y} = \mathbf{ZG} + \mathbf{XB} + \mathbf{E}$$

$$\begin{aligned}(\mathbf{I}_n - \mathbf{H})\mathbf{Y} &= (\mathbf{I}_n - \mathbf{H})\mathbf{ZG} + (\mathbf{I}_n - \mathbf{H})\mathbf{XB} + (\mathbf{I}_n - \mathbf{H})\mathbf{E} \\ &\quad ((\mathbf{I}_n - \mathbf{H})\mathbf{ZG} = 0)\end{aligned}$$

$$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{XB} + (\mathbf{I}_n - \mathbf{H})\mathbf{E}$$

where

- $\mathbf{H} = \mathbf{Z}(\mathbf{Z}^T\mathbf{Z})^{-1}\mathbf{Z}^T$ is the $n \times n$ projection matrix
- $(\mathbf{I}_n - \mathbf{H})$ is the $n \times n$ 'residualizing' matrix



Exchangeability

- Null model: $(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{E} \sim (\mathbf{0}_{n \times p}, (\mathbf{I}_n - \mathbf{H}) \otimes \boldsymbol{\Sigma})$
- $(\mathbf{I}_n - \mathbf{H})\mathbf{Y} \stackrel{d}{\neq} \Pi(\mathbf{I}_n - \mathbf{H})\mathbf{Y}$
where Π : permutation matrix (*null-invariant transformation*)



Exchangeability


- Null model: $(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{E} \sim (\mathbf{0}_{n \times p}, (\mathbf{I}_n - \mathbf{H}) \otimes \boldsymbol{\Sigma})$

- $(\mathbf{I}_n - \mathbf{H})\mathbf{Y} \stackrel{d}{\neq} \Pi(\mathbf{I}_n - \mathbf{H})\mathbf{Y}$

where Π : permutation matrix (*null-invariant transformation*)

i.e. observations are not exchangeable anymore :(

a look to $(\mathbf{I}_n - \mathbf{H})$:



solutions to be discussed:

- permutation for factor (i.e. discrete)
- permutation/rotation for covariates (i.e. continuous + discrete)




Nuisance Factors (i.e. Discrete)

In our example, sex is a 2-levels factor



Nuisance Factors (i.e. Discrete)

In our example, sex is a 2-levels factor
a look to H_0



Nuisance Factors (i.e. Discrete)

Solution:

permutations within each strata of confounder:
(i.e. within Female and Male)



Nuisance Factors (i.e. Discrete)

Solution:

permutations within each strata of confounder:
(i.e. within Female and Male)

Even more:

Exchangeability within strata implies **EXACT** control of the type I error,
allowed different models (e.g. heteroscedastic errors) between strata.



Nuisance Factors (i.e. Discrete)

Solution:

permutations within each strata of confounder:
(i.e. within Female and Male)

Even more:

Exchangeability within strata implies **EXACT control of the type I error**,

allowed different models (e.g. heteroscedastic errors) between strata.

Special case:

exact solution for paired samples (and one-sample) test with heteroscedastic errors (few more slides upon request)



Paired samples

$$\mathbf{Y} = \mathbf{ZG} + \mathbf{XB} + \mathbf{E}$$

- \mathbf{X} : treatments $\mathbf{X} = (-1, +1, -1, +1, \dots, -1, +1)$
- \mathbf{Z} : $n/2$ dummy variables, one for each subject.




Paired samples

$$\mathbf{Y} = \mathbf{Z}\mathbf{G} + \mathbf{X}\mathbf{B} + \mathbf{E}$$

- \mathbf{X} : treatments $\mathbf{X} = (-1, +1, -1, +1, \dots, -1, +1)$
- \mathbf{Z} : $n/2$ dummy variables, one for each subject.

a look to H_0



Paired samples

Permutation solution:

- Π : flip responses only **within** the same subject
- test statistic: $\mathbf{X}^T(\mathbf{I}_n - \mathbf{H})\Pi(\mathbf{I}_n - \mathbf{H})\mathbf{Y}$
- (equivalent to standard permutation test for paired sample)
- subject-specific model (e.g. variance) is allowed.
- (still) Exact control of Type I error.



Paired samples

Permutation solution:

- Π : flip responses only **within** the same subject
- test statistic: $\mathbf{X}^T(\mathbf{I}_n - \mathbf{H})\Pi(\mathbf{I}_n - \mathbf{H})\mathbf{Y}$
- (equivalent to standard permutation test for paired sample)
- subject-specific model (e.g. variance) is allowed.
- (still) Exact control of Type I error.


parametric vs nonparametric: a small simulation

- $n = 10$,
- normal, Homoscedastic errors and
- normal, Heteroscedastic errors
(non linear growth of variance from 1 to n)




Paired samples - Homoscedastic



Homoscedastic Errors



Homoscedastic Errors



Paired samples - Heteroscedastic



$t = \frac{\bar{x}_{diff} - \mu_{diff}}{\sqrt{\hat{\sigma}_{diff}^2/n}}$: parametric t estimates $\hat{\sigma}_{diff}^2$,
while permutation test does not!



Nuisance Covariates (i.e. Quantitative)

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients
- $p = 12625$ genes
- covariate of interest: group (B or T-cell type patients)
- **quantitative confounder(s)**: age (+ sex)
- differentially expressed genes between the two groups?

$$\text{gene} = \gamma_0 + \gamma_1 \cdot \text{age} + \beta \cdot \text{group} + \text{error}$$



Multivariate linear model

The Multivariate linear model of course is the same:

$$\mathbf{Y} = \mathbf{ZG} + \mathbf{XB} + \mathbf{E}$$

- \mathbf{Y} : $(n \times p)$ matrix of responses
- \mathbf{X} : $(n \times q)$ matrix of covariates (group)
- \mathbf{Z} : $(n \times c)$ matrix of confounders, $c < n$
- $\mathbf{E} \sim (\mathbf{0}_{n \times p}, \mathbf{I}_n \otimes \mathbf{\Sigma})$ matrix of errors
 $\mathbf{\Sigma}$: $(p \times p)$ gene-gene covariance matrix

Marginal model (j -th gene)

$$\mathbf{y}_j = \mathbf{Z}\gamma_j + \mathbf{X}\beta_j + \varepsilon_j, \quad \varepsilon_j \sim (\mathbf{0}_{n \times 1}, \sigma_j^2 \mathbf{I}_n)$$

Multiple hypotheses

$$H_j : \beta_j = \mathbf{0}, \quad \forall \gamma_j \quad j = 1, \dots, p$$



Loss of exchangeability

For continuous variables:

$(\mathbf{I}_n - \mathbf{H})$ is not a block matrix, there are no strata
(NO permutations within strata)



Loss of exchangeability

For continuous variables:

$(\mathbf{I}_n - \mathbf{H})$ is not a block matrix, there are no strata
(NO permutations within strata)

under the null model:

$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{E} \sim (\mathbf{0}_{n \times p}, (\mathbf{I}_n - \mathbf{H}) \otimes \mathbf{\Sigma})$ matrix of errors



Loss of exchangeability

For continuous variables:


$(\mathbf{I}_n - \mathbf{H})$ is not a block matrix, there are no strata
(NO permutations within strata)

under the null model:

$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{E} \sim (\mathbf{0}_{n \times p}, (\mathbf{I}_n - \mathbf{H}) \otimes \mathbf{\Sigma})$ matrix of errors
rows are not exchangeable.



a look to H_0



Recovering the (weak) randomization hypothesis

- $(\mathbf{I}_n - \mathbf{H}) = \mathbf{Q}\mathbf{Q}^\top$ with $\mathbf{Q}^\top\mathbf{Q} = \mathbf{I}_{n-c}$
 $(\mathbf{I}_n - \mathbf{H})$ idempotent (i.e. eigenvalues 1 or 0),
 \mathbf{Q} matrix of $n - c$ (orthogonal) eigenvectors of $(\mathbf{I}_n - \mathbf{H})$.



Recovering the (weak) randomization hypothesis

- $(\mathbf{I}_n - \mathbf{H}) = \mathbf{Q}\mathbf{Q}^\top$ with $\mathbf{Q}^\top\mathbf{Q} = \mathbf{I}_{n-c}$
 $(\mathbf{I}_n - \mathbf{H})$ idempotent (i.e. eigenvalues 1 or 0),
 \mathbf{Q} matrix of $n - c$ (orthogonal) eigenvectors of $(\mathbf{I}_n - \mathbf{H})$.
- The model:

$$\mathbf{Y} = \mathbf{Z}\mathbf{G} + \mathbf{X}\mathbf{B} + \mathbf{E}$$

$$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{Z}\mathbf{G} + (\mathbf{I}_n - \mathbf{H})\mathbf{X}\mathbf{B} + (\mathbf{I}_n - \mathbf{H})\mathbf{E}$$

$$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{X}\mathbf{B} + (\mathbf{I}_n - \mathbf{H})\mathbf{E}$$



Recovering the (weak) randomization hypothesis

- $(\mathbf{I}_n - \mathbf{H}) = \mathbf{Q}\mathbf{Q}^\top$ with $\mathbf{Q}^\top\mathbf{Q} = \mathbf{I}_{n-c}$
 $(\mathbf{I}_n - \mathbf{H})$ idempotent (i.e. eigenvalues 1 or 0),
 \mathbf{Q} matrix of $n - c$ (orthogonal) eigenvectors of $(\mathbf{I}_n - \mathbf{H})$.
- The model:

$$\mathbf{Y} = \mathbf{Z}\mathbf{G} + \mathbf{X}\mathbf{B} + \mathbf{E}$$

$$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{Z}\mathbf{G} + (\mathbf{I}_n - \mathbf{H})\mathbf{X}\mathbf{B} + (\mathbf{I}_n - \mathbf{H})\mathbf{E}$$

$$(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = (\mathbf{I}_n - \mathbf{H})\mathbf{X}\mathbf{B} + (\mathbf{I}_n - \mathbf{H})\mathbf{E}$$

$$\mathbf{Q}^\top(\mathbf{I}_n - \mathbf{H})\mathbf{Y} = \mathbf{Q}^\top(\mathbf{I}_n - \mathbf{H})\mathbf{X}\mathbf{B} + \mathbf{Q}^\top(\mathbf{I}_n - \mathbf{H})\mathbf{E}$$

$$\mathbf{Q}^\top\mathbf{Y} = \mathbf{Q}^\top\mathbf{X}\mathbf{B} + \mathbf{Q}^\top\mathbf{E}$$

$$\tilde{\mathbf{Y}} = \tilde{\mathbf{X}}\mathbf{B} + \tilde{\mathbf{E}}$$



Recovering the (weak) randomization hypothesis

- Under the **Null model**:

$$\begin{aligned} \mathbf{Q}^\top (\mathbf{I}_n - \mathbf{H}) \mathbf{Y} &= \mathbf{Q}^\top (\mathbf{I}_n - \mathbf{H}) \mathbf{E} \\ \mathbf{Q}^\top \mathbf{Y} &= \mathbf{Q}^\top \mathbf{E} \\ \tilde{\mathbf{Y}} &= \tilde{\mathbf{E}} \end{aligned}$$

- $\tilde{\mathbf{Y}} \sim (\mathbf{0}_{(n-c) \times p}, \mathbf{I}_{n-c} \otimes \Sigma)$
 $\tilde{\Pi} \tilde{\mathbf{Y}} \sim (\mathbf{0}_{(n-c) \times p}, \tilde{\Pi} \mathbf{I}_{n-c} \tilde{\Pi}^\top \otimes \Sigma)$ (**n-c orthogonal** rows) i.e.
permute the orthogonalized residuals $\tilde{\mathbf{Y}}$ instead of $(\mathbf{I}_n - \mathbf{H}) \mathbf{Y}$
- $\tilde{\mathbf{Y}}$ and $\tilde{\Pi} \tilde{\mathbf{Y}}$ have the **same first two moments**,
therefore we can perform approximated permutation tests.
- as a consequence, **asymptotic control of the type I error**
(hint: test stat is asymptotically normal by CLT + normal distribution has null higher moments)



Rotation Tests


Since we renounced to exactness, we get a broader class of transformations that **preserve first two moments**:

rotation test:


- Same as permutation but use rotations $\tilde{\mathbf{O}}$ instead of permutations $\tilde{\Pi}$.
- i.e. random generate orthogonal basis ($\tilde{\mathbf{O}}^T \tilde{\mathbf{O}} = \mathbf{I}_{n-c}$)
- test statistic: $\tilde{\mathbf{X}}^T \tilde{\mathbf{O}} \tilde{\mathbf{Y}}$
- Langsrud (2005), Perry and Owen (2010)




Rotations vs Permutations




Rotations vs Permutations



Rotations vs Permutations



Rotations vs Permutations



Some properties

- Rotations of the orthogonalized residuals extends Permutations (i.e. orthogonal matrices with not all zeros and ones).
- Both tests are approximated (and asymptotically exact and consistent),
- When $\tilde{\mathbf{Y}}$ *left-spherically distributed* (e.g. normal), $\tilde{\mathbf{Y}} \stackrel{d}{=} \tilde{\mathbf{O}}\tilde{\mathbf{Y}}$ i.e. exact test.



Some properties

- Rotations of the orthogonalized residuals extends Permutations (i.e. orthogonal matrices with not all zeros and ones).
- Both tests are approximated (and asymptotically exact and consistent),
- When $\tilde{\mathbf{Y}}$ *left-spherically distributed* (e.g. normal), $\tilde{\mathbf{Y}} \stackrel{d}{=} \tilde{\mathbf{O}}\tilde{\mathbf{Y}}$ i.e. exact test.
- Easy to generate from the joint distribution i.e. dependence among tests is dealt.



... back to motivating example: ALL results

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients, $p = 12625$ genes
- covariate of interest: group (B or T-cell type patients)
- confounder(s): age + sex



... back to motivating example: ALL results

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients, $p = 12625$ genes
- covariate of interest: group (B or T-cell type patients)
- `confounder(s)`: age + sex
- `library(flip)`
`pvalues = flip(Y, X=~group, Z=~age+sex,`
`data=ALLdata, testType='rotation', perms=10000)`



... back to motivating example: ALL results

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients, $p = 12625$ genes
- covariate of interest: group (B or T-cell type patients)
- `confounder(s)`: age + sex
- `library(flip)`
`pvalues = flip(Y, X=~group, Z=~age+sex,`
`data=ALLdata, testType='rotation', perms=10000)`

Application to Multiplicity control:

- Familywise Error Rate:
parametric & Holm vs permutation & min-p



... back to motivating example: ALL results

- Acute Lymphoblastic Leukemia data (Chiaretti et al., 2004)
- $n = 128$ patients, $p = 12625$ genes
- covariate of interest: group (B or T-cell type patients)
- `confounder(s)`: age + sex
- `library(flip)`
`pvalues = flip(Y, X=~group, Z=~age+sex,`
`data=ALLdata, testType='rotation', perms=10000)`


Application to Multiplicity control:

- Familywise Error Rate:
parametric & Holm vs permutation & min-p
- simultaneous CI for number of rejected hypos:
parametric & Simes vs permutation & Meinshausen




Familywise Error Rate (Holm vs min-p)

Raw p-values




Adjusted p-values




Familywise Error Rate (Holm vs min-p)

Adjusted p-values



Familywise Error Rate (Holm vs min-p)


```
res=flip.adjust(pvalues,'maxT')
```




Simes vs Meinshausen

```
library(cherry); curveMeinshausen(pvalues)
```

of false rejections
(lower bound .95)



Prop. of false rejections
(lower bound .95)



Take-home message

- Permutations and Rotations
 - easily deal with dependence, even when $p \gg n$,
 - more powerful multiplicity control



Take-home message

- Permutations and Rotations
 - easily deal with dependence, even when $p \gg n$,
 - more powerful multiplicity control
- Adjusting for confounders



Take-home message

- Permutations and Rotations
 - easily deal with dependence, even when $p \gg n$,
 - more powerful multiplicity control
- Adjusting for confounders
 - discrete:** permutations within strata provide exact tests



Take-home message

- Permutations and Rotations
 - easily deal with dependence, even when $p \gg n$,
 - more powerful multiplicity control
- Adjusting for confounders
 - discrete:** permutations within strata provide exact tests and require less assumptions on data



Take-home message

- Permutations and Rotations
 - easily deal with dependence, even when $p \gg n$,
 - more powerful multiplicity control
- Adjusting for confounders
 - discrete:** permutations within strata provide exact tests and require less assumptions on data
 - continuous:** use orthogonal residuals (with permutations or rotations):
 - Approximated test in general
 - Exact for multivariate left-spherical (e.g. normal)
- simple algorithm even for high dimensional data
- **R packages** 'flip' and 'cherry'

