

Five-Term Arithmetic Progressions and Cubes in a Pure Cubic Field

Codex (GPT-5.6-sol)

September 2026

Abstract

For a nonconstant rational five-term arithmetic progression, we ask how many nonzero terms can, after multiplication by one common rational scalar, be cubes in a single nontrivial pure cubic field. We prove that the exact maximum is four. The proof begins by determining the kernel of the map from rational cube classes to the Kummer group of $\mathbf{Q}(\sqrt[3]{D})$. After normalizing radicands and quotienting by affine color change and reversal, we obtain exactly 25 five-position color orbits. Ten orbits are excluded by known theorems on rational cubes in arithmetic progression. A prime-support lemma reduces the remaining 15 orbits to 60 explicit systems of three diagonal cubic equations. For every system, exhaustive finite-field enumeration gives an obstruction at a displayed prime of good reduction; a versioned certificate records all $p^2 - 1$ nonzero parameter pairs. The progression $(-3, -1, 1, 3, 5)$ supplies four counted terms over $\mathbf{Q}(\sqrt[3]{3})$. Thus $R_{(3,1)}^\times(5) = 4$. This finite classification excludes every five-hit color pattern but does not classify all rational progressions attaining four hits. We additionally prove that two positive-rank genus-one curves supply infinite families in six distinct four-hit orbits. Four arise from $2X^3 - 3Y^3 + Z^3 = 0$; the two further orbits $((0, 1, 2, 4), 0010)$ and $((0, 1, 3, 4), 0010)$ arise from $3X^3 - 4Y^3 + Z^3 = 0$. These six branches are infinite. Separately, the genus-four model $((0, 1, 2, 4), 0102)$ contains the explicit nondegenerate maximizer $(64, 36, 8, -20, -48)$ over $\mathbf{Q}(\sqrt[3]{6})$; its elliptic quotient has a certified non-torsion point, but no infinite family on the source curve is claimed. Thus seven of the initial 31 four-hit models are closed for existence and 24 remain open.

Keywords: arithmetic progression; pure cubic field; Kummer class; perfect cube; local obstruction.

Mathematics Subject Classification (2020): Primary 11B25; Secondary 11D41, 11R16, 11Y50.

1 Definition and normalizations

Let $A_i = a + id$ for $0 \leq i < 5$, where $a, d \in \mathbf{Q}$ and $d \neq 0$. For a nontrivial class $[D] \in \mathbf{Q}^*/\mathbf{Q}^{*3}$ put $K_D = \mathbf{Q}(\sqrt[3]{D})$. Define $R_{(3,1)}^\times(5)$ as the maximum number of nonzero A_i for which there are one D and one $\lambda \in \mathbf{Q}^*$ such that $\lambda A_i \in K_D^3$ at every counted position.

The radicand may be chosen as a positive cube-free integer: -1 is already a rational cube, and multiplying D by a rational cube changes no field. Moreover D and D^2 generate the same field. We quotient progressions by common rational scaling and by reversal $(a, d) \mapsto (a + 4d, -d)$. Zero has no class in $\mathbf{Q}^*/\mathbf{Q}^{*3}$ and is never counted. If D is a rational cube, then $K_D = \mathbf{Q}$ and the problem is the ordinary rational-cube problem, whose five-term maximum is three [2]; this degree-one degeneration cannot attain our maximum.

2 The pure-cubic Kummer kernel

Lemma 2.1. *If $[D] \neq 1$ in $\mathbf{Q}^*/\mathbf{Q}^{*3}$, then*

$$\ker(\mathbf{Q}^*/\mathbf{Q}^{*3} \longrightarrow K_D^*/K_D^{*3}) = \langle [D] \rangle = \{1, [D], [D]^2\}.$$

Equivalently, a nonzero rational r is a cube in K_D if and only if $[r] \in \langle [D] \rangle$.

Proof. Write $\alpha^3 = D$ and $y = a + b\alpha + c\alpha^2$. Reduction of y^3 in the basis $1, \alpha, \alpha^2$ gives nonconstant coefficients

$$3(a^2b + Dac^2 + Db^2c), \quad 3(a^2c + ab^2 + Dbc^2).$$

If $a = 0$, their vanishing implies $bc = 0$, so y lies on either the $\mathbf{Q}\alpha$ or $\mathbf{Q}\alpha^2$ axis. If $a \neq 0$, put $B = b/a$, $C = c/a$. The two equations are

$$B + DC^2 + DB^2C = 0, \quad C + B^2 + DBC^2 = 0.$$

Their resultants, eliminating C and B respectively, are

$$BD(B^3D - 1)^2, \quad C(C^3D^2 - 1)^2.$$

Since D is not a rational cube, the first forces $B = 0$, and then the first equation forces $C = 0$. Hence an element of K_D with rational cube lies in exactly one of $\mathbf{Q}$, $\mathbf{Q}\alpha$, $\mathbf{Q}\alpha^2$. Cubing proves the asserted three classes, and the converse is immediate. $\square$

Thus, after the common scaling, every counted term has a unique color $c_i \in \mathbf{F}_3$ and can be written

$$A_i = D^{c_i} x_i^3, \quad x_i \in \mathbf{Q}^*. \quad (1)$$

Changing the common scale by D^k , replacing D by D^2 , and reversing the progression act on color words by

$$c_i \mapsto c_i + k, \quad c_i \mapsto -c_i, \quad (c_0, \dots, c_4) \mapsto (c_4, \dots, c_0).$$

3 The 25 color orbits

Exact enumeration gives 25 orbits of $\mathbf{F}_3^5$ under these operations. For completeness, Burnside's lemma gives the same count without trusting the choice of representatives: the fixed-point counts of the 12 group elements are

$$243, 27, 9, 9, 9, 1, 1, 1, 0, 0, 0, 0.$$

Their sum is 300, and hence there are $300/12 = 25$ orbits. Nine contain a monochromatic three-term arithmetic progression of indices among

$$(0, 1, 2), (1, 2, 3), (2, 3, 4), (0, 2, 4).$$

Such an orbit would give three nonconstant rational cubes in arithmetic progression after scaling, contrary to Darmon–Merel [1]. Exactly one further orbit, represented by 00100, survives this test but has four terms of one color, at positions 0, 1, 3, 4. It contradicts the exact ordinary bound $P_5(3) = 3$ of Hajdu–Tengely [2]. The remaining representatives are listed below. There is no hidden integrality assumption here: choosing an integer L that clears the finitely many rational cube roots and the progression parameters, multiplication of the entire progression by the common cube L^3 makes both the progression and those roots integral. Cube positions and nonconstancy are therefore preserved.

$$\begin{aligned} &00101, 00102, 00110, 00112, 00120, 00121, 00122, 01001, 01002, 01012, \\ &01021, 01102, 01120, 01201, 01210. \end{aligned} \quad (1)$$

4 Prime support and the 60 curves

Clear denominators and divide by the common gcd, obtaining a primitive integer progression $A_i = a + id$. Its classes have the affine form $g + c_i\delta$ in $\mathbf{Q}^*/\mathbf{Q}^{*3}$. For $i \neq j$,

$$\gcd(A_i, A_j) \mid |i - j|d,$$

because $\gcd(a, d) = 1$. Therefore a prime $p > 3$ divides at most one of the five terms.

Lemma 4.1 (prime-support lemma). *For each of the 15 words in (1), every $p > 3$ occurs in every A_i with exponent divisible by three. Consequently both g and δ are supported only at 2 and 3.*

Proof. If a $p > 3$ exponent were nonzero modulo three, the five valuation residues would be supported at one position. But they must be the values of the affine function $c \mapsto g_p + c\delta_p$ on the displayed color word. For a two-color word in (1), both color blocks have size at least two, so a singleton support is impossible. For a three-color word, an affine function over $\mathbf{F}_3$ vanishing at the other two colors vanishes identically. This is again impossible. Hence all five residues vanish. Taking differences of two used colors proves the assertion for δ , and then for g . $\square$

After another common rational scaling removes g , the four possible one-dimensional subspaces of the $\{2, 3\}$ -supported class plane have representatives

$$D = 2, 3, 6, 18.$$

For a word $c = (c_0, \dots, c_4)$ and one such D , all five hits give the concrete projective curve in $\mathbf{P}^4$

$$D^{c_i} x_i^3 - 2D^{c_{i+1}} x_{i+1}^3 + D^{c_{i+2}} x_{i+2}^3 = 0 \quad (0 \leq i \leq 2). \quad (2)$$

At every prime used below, $p \geq 7$ and $p \nmid D$. These conditions really give good reduction. Put $u_i = D^{c_i} x_i^2$. After deleting the invertible factor 3, the Jacobian matrix of the three equations is

$$\begin{pmatrix} u_0 & -2u_1 & u_2 & 0 & 0 \\ 0 & u_1 & -2u_2 & u_3 & 0 \\ 0 & 0 & u_2 & -2u_3 & u_4 \end{pmatrix}.$$

A projective solution has at most one $u_i = 0$: two zero terms in the induced five-term progression would force both its initial term and common difference to vanish because $p > 4$. If no u_i is zero, columns 0, 1, 2 give a nonzero minor. If the unique zero is respectively u_0, u_1, u_2, u_3, u_4 , columns 124, 024, 014, 014, 013 give

$$3u_1u_2u_4, -2u_0u_2u_4, u_0u_1u_4, u_0u_1u_4, -2u_0u_1u_3.$$

Thus the Jacobian has rank 3 at every geometric point, proving smoothness.

Table 1 gives, in word order (1), a prime with no nonzero projective point on (2); columns correspond to $D = 2, 3, 6, 18$. Every listed prime is coprime to $3D$. The certificate enumerates all $(a, d) \in \mathbf{F}_p^2 \setminus \{(0, 0)\}$ and

word	2	3	6	18	word	2	3	6	18
00101	7	7	31	7	00102	7	7	13	7
00110	7	7	13	7	00112	7	7	13	7
00120	7	13	13	31	00121	7	19	61	31
00122	13	13	13	43	01001	7	7	13	7
01002	7	7	61	7	01012	13	7	13	7
01021	7	13	13	31	01102	13	7	13	7
01120	7	7	61	7	01201	7	7	13	7
01210	7	13	31	31					

Table 1: Good primes obstructing the 60 remaining word–radicand models

tests whether $a + id \in D^{c_i} \mathbf{F}_p^3$ for all five positions. This is equivalent to enumerating the projective solutions of (2), including solutions with some zero coordinates. Absence modulo p excludes a rational point.

The complete finite computation is identified by

```
PAPER_CUBE_KUMMER5_CERTIFICATE.json,
schema paper-cube-pure-cubic-kummer-n5-v2,
SHA-256
7c5ff0bd36ebfc3bace0ca5898625b2fd7f03d318c754c6650d3a2483dd83977.
```

Its generator and tests are located by the root-level file

`PAPER_CUBE_KUMMER5_SUPPLEMENT_MANIFEST.json`.

Each of the 60 cells scans all $p^2 - 1$ nonzero parameter pairs, for 23520 pairs in total. The current working tree is public at <https://github.com/lixiang90/vibemath>. At submission, its exact payload must additionally be frozen at the immutable DOI or release URL stated in the data-availability record.

5 Exact maximum and scope of classification

Theorem 5.1. *With nonzero terms and nontrivial pure cubic fields as in Section 1,*

$$R_{(3,1)}^\times(5) = 4.$$

Proof. A five-hit progression has one of the 25 color orbits. The first ten are excluded in Section 3. Lemma 4.1 and the 60 good-prime obstructions exclude the remaining 15. Hence the maximum is at most four. For the reverse inequality, in $K = \mathbf{Q}(\alpha)$ with $\alpha^3 = 3$, the nonzero progression

$$-3, -1, 1, 3, 5$$

has its first four terms equal to the cubes of $-\alpha, -1, 1, \alpha$. □

The theorem includes a complete classification and exclusion of every *five-hit color/position class*. The next two sections close seven of the 31 four-hit models for existence: six have proved infinite families, and the seventh has one rigorously verified nondegenerate point. They do not classify every rational progression attaining four hits. The other 24 models remain distinct Diophantine problems, and we do not replace their rational-point classification by a bounded search.

If zero were counted, $-2, -1, 0, 1, 2$ would give five hits in $\mathbf{Q}(\sqrt[3]{2})$; this is exactly why the superscript $\times$ is necessary.

6 Six four-hit branches on two positive-rank cubics

6.1 The diagonal-cubic cluster at $u^3 - 243$

We strengthen Theorem 5.1 in four of the 31 four-hit color/position models left open above. Begin with positions 0, 1, 3 rational cubes. Writing them as X^3, Y^3, Z^3 , respectively, the AP identity $A_3 = 3A_1 - 2A_0$ gives the smooth plane cubic

$$C : \quad 2X^3 - 3Y^3 + Z^3 = 0. \tag{3}$$

The associated five-term progression is

$$(X^3, Y^3, 2Y^3 - X^3, Z^3, 4Y^3 - 3X^3). \tag{4}$$

The three partial derivatives of (3) have no common projective zero, so C is a genus-one curve. Put $O = (1 : 1 : 1)$ and use it as origin.

Theorem 6.1. *The curve (3) has positive Mordell–Weil rank. Consequently, in each of the four distinct color/position orbits*

$$\begin{aligned} ((0, 1, 3, 4), 0001), & \quad ((0, 1, 2, 3), 0010), \\ ((0, 1, 2, 3), 0100), & \quad ((0, 1, 2, 4), 0111), \end{aligned}$$

there are infinitely many inequivalent nonconstant rational five-term arithmetic progressions, with no zero term, having exactly four counted positions in a nontrivial pure cubic field.

Proof. For $(X : Y : Z) \in C$, set

$$\begin{aligned} N &= 6X^3Y^3 + 3Y^3Z^3 - 2Z^3X^3, \\ T &= (2X^3 + 3Y^3)(-3Y^3 - Z^3)(Z^3 - 2X^3). \end{aligned}$$

On the open set $XYZ \neq 0$, define

$$\phi(X : Y : Z) = \left(\frac{N}{X^2Y^2Z^2}, -\frac{T}{2X^3Y^3Z^3} \right) = (u, v). \quad (5)$$

A direct expansion, followed by use of (3), gives

$$v^2 = u^3 - 243.$$

Equivalently, after clearing denominators the numerator of $v^2 - u^3 + 243$ is divisible by $2X^3 - 3Y^3 + Z^3$; the exact quotient is checked by the accompanying symbolic test. Thus (5) is a rational map to

$$E : v^2 = u^3 - 243.$$

A rational map from a smooth projective curve to the proper curve E extends over its apparent exceptional points. Notice also that $C(\mathbf{Q})$ has no point with $XYZ = 0$: such a point would make one of $3, -2, 3/2$ a rational cube.

Let

$$P = (4 : 1 : -5) \in C(\mathbf{Q}), \quad Q = (7, 10) \in E(\mathbf{Q}).$$

Exact substitution and the tangent formula on E give

$$\phi(O) = (7, -10) = -Q, \quad \phi(P) = \left(\frac{16009}{400}, -\frac{2021723}{8000} \right) = 2Q.$$

The integral equation for E has discriminant $-16 \cdot 27 \cdot 243^2 = -2^4 3^{13}$. Since Q has nonzero second coordinate and 10^2 does not divide this discriminant, the Nagell–Lutz theorem shows that Q has infinite order.

Translate ϕ by $-\phi(O)$. The resulting nonconstant morphism sends origins to origins and hence is an isogeny of genus-one curves. Its value at P is $2Q - (-Q) = 3Q$, which is non-torsion. Therefore P has infinite order in $C(\mathbf{Q})$, and the points

$$P_n = O + n(P - O) \quad (n \in \mathbf{Z})$$

are pairwise distinct.

For $n \neq 0$, substitute $P_n = (X_n : Y_n : Z_n)$ in (4). None of X_n, Y_n, Z_n can vanish, as observed above; $2Y_n^3 - X_n^3 = 0$ would make 2 a rational cube, while $4Y_n^3 - 3X_n^3 = 0$ would make $4/3$ one. Moreover the common difference vanishes only when $X_n = Y_n = Z_n$, that is, at $P_n = O$.

The last term $A_4 = 4Y_n^3 - 3X_n^3$ is not a rational cube. Otherwise positions 0, 1, 3, 4 would be four rational cubes in a nonconstant five-term AP, contradicting Hajdu–Tengely’s bound $P_5(3) = 3$. Choose a positive cube-free representative $D_{4,n}$ of $[A_4]$. Then $A_4 = D_{4,n}W_{4,n}^3$ for some $W_{4,n} \in \mathbf{Q}^*$, and $K_{D_{4,n}}$ is genuinely cubic. Thus positions 0, 1, 3, 4 have colors 0001. Theorem 5.1 prevents position 2 from being a fifth hit.

The middle term $A_2 = 2Y_n^3 - X_n^3$ is likewise not a rational cube, since otherwise positions 0, 1, 2, 3 would be four rational cubes, again contradicting $P_5(3) = 3$. If $D_{2,n}$ is the positive cube-free representative of $[A_2]$, then $A_2 = D_{2,n}W_{2,n}^3$ with $W_{2,n} \in \mathbf{Q}^*$ and $K_{D_{2,n}}$ genuinely cubic. Positions 0, 1, 2, 3 therefore have colors 0010, and Theorem 5.1 prevents position 4 from being a fifth hit. These two models are not identified by affine color change and reversal: the first omits the central position 2, whereas the second omits an endpoint (position 4, or position 0 after reversal).

Two further AP constructions use the same point $(X_n : Y_n : Z_n)$ of C :

$$(Z_n^3, (Z_n^3 + Y_n^3)/2, Y_n^3, X_n^3, 2Y_n^3 - Z_n^3), \quad (6)$$

$$(2X_n^3 - Y_n^3, X_n^3, Y_n^3, 2Y_n^3 - X_n^3, Z_n^3). \quad (7)$$

The first is an AP because $X^3 - 3Y^3 + 2Z^3 = 0$ after swapping X and Z in (3); its rational-cube positions are 0, 2, 3. Its singleton A_1 cannot vanish: otherwise $Z = -Y$ and the cubic would give $X^3 = 2Y^3$. Nor can its missing term A_4 vanish, since that would give $Z^3 = 2Y^3$. If A_1 were a rational cube, positions 0, 1, 2, 3 would contradict $P_5(3) = 3$. Its nontrivial cube class therefore produces the orbit $((0, 1, 2, 3), 0100)$, and Theorem 5.1 excludes a fifth hit.

In the second AP the rational-cube positions are 1, 2, 4. Its singleton A_0 and missing term A_3 are nonzero, since their vanishing would make 2 a rational cube. The singleton is not a rational cube by the same $P_5(3) = 3$ argument. Before color normalization its word is 1000; the affine color map $c \mapsto 2c + 1$ gives the representative 0111. Again Theorem 5.1 excludes a fifth hit. The exact orbit enumeration verifies that all four displayed models are pairwise distinct.

Finally, distinct projective points of $C(\mathbf{Q})$ give distinct progressions up to common rational scaling within each construction. Indeed, comparison of any fixed nonzero rational-cube position makes the scale a rational cube, after which injectivity of cubing on $\mathbf{Q}$ recovers the same projective coordinates. Reversal has fibers of size at most two, so infinitely many equivalence classes remain after reversal. $\square$

For the first nonconstant point P , formula (4) gives

$$(64, 1, -62, -125, -188),$$

with common difference -63 . If $\alpha^3 = 188$ and $\beta^3 = 62$, then the 0001 roots at positions 0, 1, 3, 4 are 4, 1, $-5, -\alpha$, while the 0010 roots at positions 0, 1, 2, 3 are 4, 1, $-\beta, -5$. These are two distinct color/position models carried by the same rational AP. They differ from the lower-bound progression in Theorem 5.1: that progression has four consecutive counted positions in the orbit 0110 over $\mathbf{Q}(\sqrt[3]{3})$, while the present one realizes both displayed orbits.

The same point P in (6) and (7) gives the reversed pair

$$(-125, -62, 1, 64, 127), \quad (127, 64, 1, -62, -125).$$

The first realizes 0100 over $\mathbf{Q}(\sqrt[3]{62})$, while the second realizes 0111 over $\mathbf{Q}(\sqrt[3]{127})$.

The construction is an integer-parameter Mordell–Weil family, made explicit by the exact chord law in the supplement. It is not a rational $\mathbf{P}^1$ parametrization, which a smooth genus-one curve cannot admit. We prove positive rank but do not claim the exact rank, torsion subgroup, or a complete set of generators for $C(\mathbf{Q})$.

6.2 The second diagonal-cubic cluster

The remaining two $3 + 1$ models have repeated-color positions 0, 1, 4. Writing these entries as X^3, Y^3, Z^3 , respectively, gives

$$C' : \quad 3X^3 - 4Y^3 + Z^3 = 0 \tag{8}$$

and the progression

$$(X^3, Y^3, 2Y^3 - X^3, 3Y^3 - 2X^3, Z^3). \tag{9}$$

The partial derivatives $9X^2, -12Y^2, 3Z^2$ show that C' is smooth. Use $O' = (1 : 1 : 1)$ as origin.

Theorem 6.2. *The curve (8) has positive Mordell–Weil rank. Consequently each of*

$$((0, 1, 2, 4), 0010), \quad ((0, 1, 3, 4), 0010)$$

contains infinitely many inequivalent nonconstant rational five-term arithmetic progressions, with no zero term, having exactly four counted positions in a nontrivial pure cubic field.

Proof. For $(X : Y : Z) \in C'$, put

$$\begin{aligned} N' &= 12X^3Y^3 + 4Y^3Z^3 - 3Z^3X^3, \\ T' &= (3X^3 + 4Y^3)(-4Y^3 - Z^3)(Z^3 - 3X^3). \end{aligned}$$

On $XYZ \neq 0$ define

$$\phi'(X : Y : Z) = \left(\frac{N'}{X^2 Y^2 Z^2}, -\frac{T'}{2X^3 Y^3 Z^3} \right) = (u, v).$$

The exact symbolic certificate verifies

$$(T')^2 - 4(N')^3 + 3888X^6 Y^6 Z^6 = (3X^3 - 4Y^3 + Z^3)H'(X, Y, Z).$$

Thus ϕ' maps to

$$E' : v^2 = u^3 - 972.$$

There is no rational point of C' with $XYZ = 0$, since such a point would make one of $4, -3, 4/3$ a rational cube. The displayed rational map therefore has no rational exceptional point; more generally it extends from the smooth projective curve C' to the proper curve E' .

Let

$$P' = (5 : 2 : -7) \in C'(\mathbf{Q}), \quad Q' = (13, 35) \in E'(\mathbf{Q}).$$

Exact substitution and the tangent law on E' give

$$\phi'(O') = (13, -35) = -Q', \quad \phi'(P') = \left(\frac{129649}{4900}, -\frac{45441143}{343000} \right) = 2Q'.$$

The integral equation for E' has discriminant

$$-16 \cdot 27 \cdot 972^2 = -2^8 3^{13}.$$

Because $35^2 \nmid -2^8 3^{13}$, Nagell–Lutz shows that Q' has infinite order. Translating ϕ' by $-\phi'(O')$ gives a nonconstant morphism sending origins to origins, hence an isogeny. It sends $P' - O'$ to $2Q' - (-Q') = 3Q'$, so $P' - O'$ is non-torsion.

Put $P'_n = O' + n(P' - O') = (X_n : Y_n : Z_n)$. These points are pairwise distinct. For $n \neq 0$, (9) is nonconstant. Its three rational-cube entries are nonzero because $C'(\mathbf{Q})$ has no point with $XYZ = 0$; its two middle entries are nonzero because their vanishing would make 2 or $3/2$ a rational cube. Neither middle entry is a rational cube, since that would produce four rational cubes in a five-term rational AP, contrary to $P_5(3) = 3$.

Choose positive cube-free representatives $D_{2,n}$ and $D_{3,n}$ of the classes of the entries in positions 2 and 3. The Kummer-kernel lemma shows that over $\mathbf{Q}(\sqrt[3]{D_{2,n}})$ the positions 0, 1, 2, 4 have colors 0010, while over $\mathbf{Q}(\sqrt[3]{D_{3,n}})$ the positions 0, 1, 3, 4 have colors 0010. Both fields are nontrivial. Theorem 5.1 excludes a fifth hit in either field.

Finally, proportional progressions from (9) make the common scale a rational cube by comparison at position 0; comparison at positions 1 and 4 then recovers the same projective point of C' . Reversal has fibers of size at most two, so infinitely many equivalence classes remain. $\square$

The point P' gives

$$(125, 8, -109, -226, -343).$$

Over $\mathbf{Q}(\sqrt[3]{109})$ this realizes $((0, 1, 2, 4), 0010)$, and over $\mathbf{Q}(\sqrt[3]{226})$ it realizes $((0, 1, 3, 4), 0010)$. The exact supplement checks the two cube-class vectors

$$(0, 0, 1, *, 0), \quad (0, 0, *, 1, 0),$$

where $*$ denotes a non-hit.

7 A seventh four-hit model: strict existence

Theorem 7.1 (Existence in the 0102 model). *The genus-four model $((0, 1, 2, 4), 0102)$ contains a nonconstant rational five-term arithmetic progression with no zero term and exactly four cube positions in the genuine cubic field $\mathbf{Q}(\sqrt[3]{6})$. Its natural elliptic quotient has positive Mordell–Weil rank.*

Proof. Put $A_0 = X^3$ and $A_2 = Y^3$. Then

$$A_1 = (X^3 + Y^3)/2, \quad A_4 = 2Y^3 - X^3.$$

The singleton positions 1 and 4 have inverse Kummer classes precisely on the weighted-projective curve

$$(X^3 + Y^3)(2Y^3 - X^3) = 2W^3. \quad (10)$$

This is the smooth genus-four cyclic cover reconstructed in the supplement. The point $(X : Y : W) = (2 : 1 : -3)$ satisfies (10) and gives

$$(8, 9/2, 1, -5/2, -6).$$

Multiplication by the rational cube 8 preserves every cube condition and gives the integer progression

$$(64, 36, 8, -20, -48), \quad d = -28.$$

Let $\alpha^3 = 6$. The polynomial $T^3 - 6$ is Eisenstein at 2, so $[\mathbf{Q}(\alpha) : \mathbf{Q}] = 3$, and

$$64 = 4^3, \quad 36 = (\alpha^2)^3, \quad 8 = 2^3, \quad -48 = (-2\alpha)^3.$$

Thus positions 0, 1, 2, 4 are cubes. Their classes modulo $\langle [6] \rangle$ are 0201; the permitted color automorphism $c \mapsto 2c$ gives the representative 0102.

The omitted term is not a cube. By Lemma 2.1, a nonzero rational cube in $\mathbf{Q}(\alpha)$ has class 1, 6, or 36 modulo rational cubes. Every such class has 5-adic valuation zero modulo 3, whereas $v_5(-20) = 1$. All five displayed terms are nonzero and pairwise distinct, so this is a legal nondegenerate four-hit progression.

For completeness, on $Y \neq 0$ define

$$u = -2W/Y^2, \quad v = 2(X/Y)^3 - 1.$$

Equation (10) gives the exact identity

$$v^2 = u^3 + 9.$$

The source point maps to $(u, v) = (6, 15)$. This integral Weierstrass equation has discriminant $-16 \cdot 27 \cdot 9^2 = -34992$, and $15^2 = 225 \nmid 34992$. Nagell–Lutz therefore proves that $(6, 15)$ is non-torsion. This proves positive rank of the quotient only; it does not imply infinitely many rational lifts to the genus-four source. $\square$

8 Relation to earlier work and claim boundary

Darmon–Merel and Hajdu–Tengely provide the two rational-power obstructions used in the five-hit proof. The closest methodological precedent we found is Gonzalez-Jimenez–Xarles’s treatment of five squares over quadratic fields [4]. Their reduction already assigns Kummer types to the positions, identifies common scaling and reversal, and studies the resulting curves over $\mathbf{Q}$ by rational-point and local-solubility methods. Thus we do not claim originality for the general type/color, scaling, reversal, curve, or local- obstruction strategy. Their problem requires all five terms to be squares over a quadratic field, whereas ours concerns exponent three, ternary Kummer colors, partial hits, and a variable pure cubic field.

Other work on powers in number fields usually asks for all terms to be powers of elements of the field: Gonzalez-Jimenez treats three cubes over quadratic fields [3], Xarles proves qualitative degree-dependent bounds for powers in progressions [7], and Bremner–Siksek exclude five squares over cubic fields [8]. Hajdu–Tengely and Bruin–Gyory–Hajdu–Tengely study literal powers in primitive integer progressions [5, 6]. Those problems do not impose our combination of rational AP entries, one common rational scale, partial hits, and one variable pure cubic field.

Accordingly, the precise strengthened claim here is only Theorems 5.1, 6.1, 6.2, and 7.1: the exact five-term maximum, exhaustive exclusion of all five-hit color classes, positive-rank infinite families in six stated four-hit orbits, and strict existence in one further orbit. We make no priority claim and no classification claim for the other 24 four-hit models. In particular, the unreviewed rank-zero quotient notes from Round 11 are not part of any manuscript theorem. The supplement reconstructs the 29-model Round-09 input as four plane cubics, nine genus-4 bidegree curves, and sixteen genus-4 cyclic covers; its coordinate-permutation

clusters are not asserted to be complete $\mathbf{Q}$ -isomorphism classes. Our search of accessible publisher pages, arXiv records, zbMATH metadata, author publication pages, the normalized progression, and the displayed genus-four equation located no equivalent construction. This is not a priority assertion. MathSciNet was blocked by institutional authentication, and the complete forward-citation chain and full-text equation search remain to be checked.

Statements and Declarations

Funding. No funding statement is supplied because this research artifact is not being submitted.

Competing interests. No competing-interests declaration is made because no submission is being made.

Author contributions. Codex (GPT-5.6-sol): conceptualization, formal analysis, methodology, software, validation, writing—original draft, and writing—review and editing.

Data and code availability. The exact source, tests, and finite certificate are available in the public working repository at <https://github.com/lixiang90/vibemath>. The accepted payload must also be deposited at PERSISTENT ARCHIVE DOI/URL -- NOT ASSIGNED; no immutable record has been supplied because no submission is being made.

Use of generative AI. The sole listed author of this research artifact is Codex (GPT-5.6-sol), an AI model. It performed exploratory mathematical research, drafting, code generation, exact validation, and preparation of the reproducibility package. No human author, affiliation, corresponding-author address, or submission is asserted.

Ethics approval, consent to participate, and consent for publication. Not applicable: this is a mathematical study involving no human participants, animals, or personal data.

Supplementary information. Online Resource 1, PAPER_CUBE_KUMMER5_ARCHIVE_CANDIDATE.zip (ZIP), contains the exact source, PDF, generator, tests, 60-model certificate, manifest, data dictionary, and release metadata used for the finite classification. It is a local candidate and has not been deposited. The Round-11 existence certificate is distributed separately in the working tree.

References

- [1] H. Darmon and L. Merel, *Winding quotient and some variants of Fermat’s Last Theorem*, J. Reine Angew. Math. **490** (1997), 81–100, <https://doi.org/10.1515/crll.1997.490.81>.
- [2] L. Hajdu and S. Tengely, *Powers in arithmetic progressions*, Ramanujan J. **55** (2021), 965–986, <https://doi.org/10.1007/s11139-020-00331-5>.
- [3] E. Gonzalez-Jimenez, *Three cubes in arithmetic progression over quadratic fields*, Arch. Math. (Basel) **95** (2010), 233–241, <https://doi.org/10.1007/s00013-010-0166-5>.
- [4] E. Gonzalez-Jimenez and X. Xarles, *Five squares in arithmetic progression over quadratic fields*, Rev. Mat. Iberoam. **29** (2013), 1211–1238, <https://doi.org/10.4171/RMI/754>.
- [5] L. Hajdu and S. Tengely, *Arithmetic progressions of squares, cubes and n -th powers*, Funct. Approx. Comment. Math. **41** (2009), 129–138, <https://doi.org/10.7169/facm/1261157805>.
- [6] N. Bruin, K. Gyory, L. Hajdu, and S. Tengely, *Arithmetic progressions consisting of unlike powers*, Indag. Math. (N.S.) **17** (2006), 539–555, [https://doi.org/10.1016/S0019-3577\(07\)00002-X](https://doi.org/10.1016/S0019-3577(07)00002-X).

- [7] X. Xarles, *Squares in arithmetic progression over number fields*, J. Number Theory **132** (2012), 379–389, <https://doi.org/10.1016/j.jnt.2011.07.010>.
- [8] A. Bremner and S. Siksek, *Squares in arithmetic progression over cubic fields*, Int. J. Number Theory **12** (2016), 1409–1414, <https://doi.org/10.1142/S179304211650086X>.