

Square-class patterns of seven consecutive integers and three elementary integral quartics

AUTHOR IDENTITY CONFIRMATION REQUIRED

3 September 2026

Abstract

We study the affine rank of the square classes of seven consecutive nonzero integers. This records when a common rational scaling makes every term a square in a field of degree at most four generated by two square roots. After affine rank at most one is excluded, enumeration up to relabelling gives 651 three- or four-block equality patterns; quotienting by reversal and applying known six-term and four-square restrictions leaves 284 candidates, each controlled by precisely fifteen character quotients. A four-consecutive-factor identity eliminates 186 patterns. We determine all integral points on $y^2 = t(t+2)(t+3)(t+6)$ by an elementary squarefree-kernel argument and finite congruence certificates. Together with an integral reflection, this eliminates 44 further patterns after a same-parameter check, leaving 54. Two constant-difference factorizations exclude another 19 and 12 patterns, leaving 23 explicitly listed necessary candidates. Every computational count and congruence certificate is bound by a versioned reproducibility manifest. No bounded search or unproved Mordell–Weil computation enters the proof. The theorem neither proves that any remaining pattern is realizable nor decides the unrestricted rational-progression problem.

Keywords: square classes; arithmetic progressions; integral points; biquadratic fields; Diophantine equations.

Mathematics Subject Classification (2020): Primary 11B25; Secondary 11D45, 11G05.

1 Square-class formulation

Put $G = \mathbf{Q}^*/\mathbf{Q}^{*2}$, written additively over $\mathbf{F}_2$. For the nonzero numbers $a_i = t + i$, define

$$\text{arank}(a_0, \dots, a_6) = \dim_{\mathbf{F}_2} \langle [a_i] - [a_0] : 1 \leq i \leq 6 \rangle.$$

The parameters $t = -6, \dots, 0$ are excluded.

For reference, define

$$R_s(N) = \max_{\substack{a, q \in \mathbf{Q}, q \neq 0; \\ a+qi \neq 0 \ (0 \leq i < N)}} \max_{\substack{V \leq G, \dim_{\mathbf{F}_2} V = s}} \# \{0 \leq i < N : [a + qi] \in V\},$$

so every squareclass occurring in the definition is defined. Thus the maximum ranges over nonconstant rational arithmetic progressions whose N displayed terms are all nonzero, and over exactly s -dimensional $\mathbf{F}_2$ -subspaces of G . The maximum is unchanged if one permits a common rational scaling and replaces linear rank by affine rank.

Lemma 1. $R_1(6) = 5$.

Proof. If six rational terms have classes in $V = \langle [D] \rangle$, they are six squares in the quadratic field $\mathbf{Q}(\sqrt{D})$. Xarles proves that a nonconstant progression over a quadratic field has at most five square terms [10, Theorem 1, $S(2) = 6$]. Conversely,

$$49, 169, 289, 409, 529$$

is a nonconstant progression of squares over $\mathbf{Q}(\sqrt{409})$; appending the sixth term 649 shows that five of six terms meet the condition, so the upper bound is attained. $\square$

Proposition 2 (Common scaling). *The seven classes have affine rank at most two if and only if there are a common $\lambda \in \mathbf{Q}^*$ and a field $L = \mathbf{Q}(\sqrt{D_1}, \sqrt{D_2})$ of degree at most four such that every $\lambda(t + i)$ is a square in L . If the difference classes span a two-dimensional space, L may be chosen biquadratic.*

Proof. Write $[t + i] = c + v_i$ with $v_i \in V = \langle [D_1], [D_2] \rangle$, and choose $[\lambda] = c$. Then $[\lambda(t + i)] = v_i$ dies in L^*/L^{*2} . Conversely,

$$\ker(\mathbf{Q}^*/\mathbf{Q}^{*2} \longrightarrow L^*/L^{*2}) = \langle [D_1], [D_2] \rangle,$$

so the classes lie in a common affine coset of dimension at most two. Indeed, if a rational class $[r]$ dies in L , then either r is already a rational square or $\mathbf{Q}(\sqrt{r})$ is a quadratic subfield of L . The quadratic subfields of a biquadratic extension are exactly $\mathbf{Q}(\sqrt{D_1})$, $\mathbf{Q}(\sqrt{D_2})$, and $\mathbf{Q}(\sqrt{D_1 D_2})$, by the Kummer correspondence. This gives the stated kernel; if $[L : \mathbf{Q}] < 4$, the same argument gives the corresponding zero- or one-dimensional subspace. $\square$

Remark 3. *The scaling is essential: the common class c can be independent of the two difference classes, in which case adjoining $\sqrt{c}$ may raise the degree to eight. We do not claim the unscaled $t + i$ are squares in one fixed biquadratic field.*

2 The finite pattern space

Represent a rank-two configuration by labels $\ell_i \in \mathbf{F}_2^2$. Lemma 1 forces both endpoint six-term windows to contain at least three labels. A block of size at least five is excluded by $Q(7) = 4$, and a block at four positions (i, j, k, l) with $j - i = l - k$ is excluded by [6, Proposition 5 and Corollary 6].

Proposition 4 (Exact pattern count). *Before these screens there are 343 label patterns up to reversal. The screens eliminate 59 reversal orbits, leaving 284.*

Proof. Since $\text{AGL}(2, 2) \cong S_4$, an orbit using three or four labels is an unlabelled partition into three or four blocks. Restricted growth strings enumerate

$$S(7, 3) + S(7, 4) = 301 + 350 = 651$$

partitions. Reversal fixes 35, so Burnside gives $(651 + 35)/2 = 343$.

The finite screen discards a word if an endpoint six-window has at most two blocks, if a block has size at least five, or if a four-element block has equal first and last gaps. It discards 109 words forming 59 reversal orbits. Of the remaining 542 words, 26 are reversal-fixed, giving $(542 + 26)/2 = 284$. The supplementary generator directly asserts every count. $\square$

For a label word define

$$T : \mathbf{F}_2^7 \longrightarrow \mathbf{F}_2 \oplus \mathbf{F}_2^2, \quad e \longmapsto \left(\sum e_i, \sum e_i \ell_i \right).$$

Lemma 5 (Fifteen characters). *For each pattern, $\ker T$ has dimension four and exactly 15 nonzero elements. Every such e gives the necessary equation*

$$y_e^2 = \prod_{e_i=1} (t+i).$$

Proof. The labels affinely span $\mathbf{F}_2^2$, so T has rank three. Rank-nullity gives dimension four. If $[t+i] = c + \phi(\ell_i)$, the class of the product is $(\sum e_i)c + \phi(\sum e_i \ell_i) = 0$. These are all affine linear relations. $\square$

3 The first 186 exclusions

Lemma 6 (Four consecutive factors). *If $x \in \mathbf{Z}$ and $x(x+1)(x+2)(x+3)$ is a rational square, then $x \in \{-3, -2, -1, 0\}$.*

Proof. It is an integer square y^2 . Put $n = x(x+3)$. Then

$$x(x+1)(x+2)(x+3) = (n+1)^2 - 1,$$

so $(n+1-y)(n+1+y) = 1$. The two integral factor pairs give $n = 0$ or -2 , and the four asserted values follow. $\square$

The consecutive masks are 15, 30, 60, 120. Recomputing all 15-element kernels and intersecting with these masks eliminates exactly 186 of 284 patterns, leaving 98. This is finite exact computation, not height search.

4 The mask 77 curve

Consider

$$H : y^2 = t(t+2)(t+3)(t+6).$$

It is birational to $E : Y^2 = X^3 - 36X$ by

$$X = 12 + \frac{36}{t}, \quad Y = \frac{36y}{t^2}, \quad t = \frac{36}{X-12}, \quad y = \frac{36Y}{(X-12)^2}.$$

The boundary correspondence is

H	E
$(0, 0)$	O
$(-2, 0)$	$(-6, 0)$
$(-3, 0)$	$(0, 0)$
$(-6, 0)$	$(6, 0)$
∞_+	$(12, 36)$
∞_-	$(12, -36)$.

This is Cremona 576h2, equivalently LMFDB 576.c3 [7]. The label is only a cross-check: the condition $36/(X-12) \in \mathbf{Z}$ is not ordinary integrality of X .

Theorem 7.

$$H(\mathbf{Z}) = \{(-6, 0), (-3, 0), (-2, 0), (0, 0), (6, 72), (6, -72)\}.$$

Proof. Put $A = t(t+6)$ and $B = (t+2)(t+3)$. Since $B - A = 6 - t$ and $A = (t+12)(t-6) + 72$, one has $\gcd(A, B) \mid 72$. At a nonbranch integral point, $B > 0$ and $AB = y^2 > 0$, hence $A > 0$. Their common positive squarefree kernel is

$$d \in \{1, 2, 3, 6\}, \quad A = dU^2, \quad B = dV^2.$$

With $x = t + 3$ this becomes

$$x^2 - dU^2 = 9, \quad x(x - 1) = dV^2.$$

Write $d = ab$ as an ordered coprime factorization. For $x > 0$ and $x < 0$ one obtains, respectively,

$$\begin{aligned} x = aR^2, \quad x - 1 = bS^2, \quad aR^2 - bS^2 &= 1, \\ x = -aR^2, \quad x - 1 = -bS^2, \quad bS^2 - aR^2 &= 1, \end{aligned}$$

together with $a^2R^4 - dU^2 = 9$. There are 18 branches. Direct exhaustion of (R, S, U) modulo the listed modulus eliminates 15:

d	$(a, b), \text{sign}(x)$	modulus
1	$(1, 1), +; (1, 1), -$	16; 8
2	$(1, 2), -; (2, 1), +; (2, 1), -$	9; 8; 8
3	$(1, 3), +; (1, 3), -; (3, 1), +$	9; 8; 8
6	$(1, 6), +; (1, 6), -; (2, 3), +; (2, 3), -$ $(3, 2), -; (6, 1), +; (6, 1), -$	9; 8; 8; 8 8; 8; 8.

The certificate exhausts every residue triple, making each row a finite proof.

For $d = 2, (a, b) = (1, 2), x > 0$,

$$R^2 - 2S^2 = 1, \quad R^4 - 2U^2 = 9, \quad (U - RS)(U + RS) = (R^2 - 9)/2.$$

$R = 1$ is impossible, $R = 3$ gives $(S, U) = (2, 6)$, and for $R \geq 5$ both factors are positive while $S > R/2$ makes the second factor larger than their product.

For $d = 3, (a, b) = (3, 1), x < 0$,

$$S^2 - 3R^2 = 1, \quad U^2 = 3(R^4 - 1), \quad (RS - U)(RS + U) = R^2 + 3.$$

$R = 0, 2$ are impossible, $R = 1$ gives $(2, 0)$, and for $R \geq 3, RS > \sqrt{3}R^2 > R^2 + 3$.

For $d = 6, (a, b) = (3, 2), x > 0$,

$$3R^2 - 2S^2 = 1, \quad 3R^4 - 2U^2 = 3, \quad (U - RS)(U + RS) = (R^2 - 3)/2.$$

R is odd; $R = 1$ gives $(1, 0)$, and the same factor-size contradiction holds for $R \geq 3$. These branches yield $t = 6, y = \pm 72$ and the separated branch points $t = -6, 0$. Adding all four roots proves the theorem. $\square$

5 Same-parameter elimination

Mask 77 has support $\{0, 2, 3, 6\}$ and mask 89 has support $\{0, 3, 4, 6\}$. The integral reflection $t = -u - 6$ gives

$$t(t+3)(t+4)(t+6) = u(u+2)(u+3)(u+6).$$

Thus the nonbranch parameter sets are $\{6\}$ and $\{-12\}$.

Theorem 8 (Finite classification). *Among the 98 patterns left above, 26 contain mask 77, 25 contain mask 89, and 7 contain both. All 44 patterns in the union fail the simultaneous 15-character conditions. Exactly 54 candidate patterns remain.*

Proof. The union has $26 + 25 - 7 = 44$ elements. For a pattern containing only mask 77, substitute $t = 6$ into all 15 character products; at least one other product is nonsquare. For mask 89 use $t = -12$. If both occur, the singleton candidate sets are disjoint. The certificate records every failed mask, every common candidate set, and all 54 surviving identifiers. It independently asserts 15 distinct masks in every input row. $\square$

6 A fourth-round gate: mask 102

The remaining 54 patterns were ranked by two exact quantities: the number of patterns containing a given four-root character, and whether its four linear factors admit a pairing into two monic quadratics with constant difference. Mask 102, supported at $\{1, 2, 5, 6\}$, ranks first: it occurs in 19 patterns and has

$$A = (t + 1)(t + 6), \quad B = (t + 2)(t + 5), \quad B - A = 4.$$

Proposition 9. *The integral points on*

$$y^2 = (t + 1)(t + 2)(t + 5)(t + 6)$$

are exactly

$$(-6, 0), \quad (-5, 0), \quad (-2, 0), \quad (-1, 0).$$

Proof. For $t \leq -7$ or $t \geq 0$, both A and B are positive. Moreover $\gcd(A, B) \mid 4$. If AB is a square, the common positive squarefree kernel is $d = 1$ or 2 , so $A = dU^2, B = dV^2$. For $d = 1$,

$$(V - U)(V + U) = V^2 - U^2 = 4.$$

The only positive factor pair of equal parity is $(2, 2)$, forcing $U = 0$ and hence a branch point. For $d = 2$ one obtains $V^2 - U^2 = 2$, impossible modulo 4. The six middle integers $-6 \leq t \leq -1$ give right sides $0, 0, 12, 12, 0, 0$, respectively. This proves the list. $\square$

Corollary 10. *Mask 102 eliminates 19 of the 54 patterns in Theorem 8. Hence 35 candidate patterns remain after this fourth-round gate.*

Proof. All integral points of Proposition 9 have $t \in \{-6, \dots, 0\}$ and are therefore degenerate for the original seven-term problem. The exact occurrence table contains mask 102 in 19 of the 54 rows. Thus no same-parameter candidate survives in those rows. $\square$

7 A fifth-round gate: mask 108

Mask 108 has support $\{2, 3, 5, 6\}$ and occurs in 12 of the 35 patterns.

Proposition 11. *The integral points on*

$$y^2 = (t + 2)(t + 3)(t + 5)(t + 6)$$

are exactly

$$(-6, 0), (-5, 0), (-4, -2), (-4, 2), (-3, 0), (-2, 0).$$

Proof. Put $A = (t + 2)(t + 6)$ and $B = (t + 3)(t + 5) = A + 3$. For $t \leq -7$ or $t \geq -1$, both are positive and $\gcd(A, B) \mid 3$. If AB is a square, their common positive squarefree kernel is $d = 1$ or 3 , so $A = dU^2$, $B = dV^2$. For $d = 1$, $(V - U)(V + U) = 3$ gives $(U, V) = (1, 2)$, while $A = 1$ would give $(t + 4)^2 = 5$, impossible modulo 8. For $d = 3$, $(V - U)(V + U) = 1$ forces $U = 0$, contradicting $A > 0$. At the five middle integers $-6 \leq t \leq -2$, direct substitution gives the displayed six points. $\square$

Theorem 12 (Summary classification). *Let $t \in \mathbf{Z} \setminus \{-6, -5, \dots, 0\}$. If the seven nonzero classes $[t], \dots, [t + 6]$ have affine rank at most two, then their equality partition, modulo relabelling and reversal, is one of the 23 patterns in Table 1. In particular, the exact reduction chain is*

$$651 \longrightarrow 343 \longrightarrow 284 \longrightarrow 98 \longrightarrow 54 \longrightarrow 35 \longrightarrow 23.$$

This theorem neither asserts that any of the 23 patterns is realizable nor decides $R_2(7)$.

Proof. Affine rank zero or one is excluded by Lemma 1, after applying the common rational scaling, to an endpoint window. Thus the actual affine rank is two. The map from the affinely spanning labels to squareclasses therefore has rank two and is injective, so the labels are precisely the actual squareclass equality blocks. Proposition 4 and Lemma 5 give 284 self-contained packets. Lemma 6 leaves 98; Theorem 8 leaves 54; Proposition 9 leaves 35; and Proposition 11, whose points all have a zero among the seven original terms, eliminates 12 more. The exact occurrence audit gives Table 1. $\square$

Table 1: Final canonical candidates. ID is the zero-based index in the supplement’s `unresolved_patterns_ranked`, ordered by its documented difficulty score and then by restricted-growth word.

ID	partition	ID	partition	ID	partition
9	0012011	12	0012202	26	0112001
31	0001202	33	0001220	43	0100021
50	0111201	59	0012231	83	0112320
134	0012131	188	0112132	210	0121132
212	0121310	214	0122213	230	0012102
251	0102221	257	0112102	266	0120210
268	0121102	271	0122201	276	0010203
281	0102003	283	0111231		

8 Supplement and reproducibility

The accompanying supplement is identified by release `paper-square-supplement-v0.5.0`. Its root manifest is `PAPER_SQUARE_SUPPLEMENT_MANIFEST.json`; it records SHA-256 and byte length for every generator, input/output certificate and regression test, together with Python/SymPy versions and clean reproduction commands. The manifest SHA-256 is `6004afc5334bbea62969640079cad80764d2938d73b6fd62a85abc2249794588`. Its release status is explicitly `LOCAL_RELEASE_CANDIDATE_NOT_PUBLICLY_ARCHIVED`; a journal submission must replace that status with a frozen archive URL/DOI without changing the hashed payload.

The frozen journal-candidate package is separately described by a non-self-referential root manifest. The file is

and its release identifier is `paper-square-submission-v0.6.4`. It binds this source, the bibliography, one reference PDF, the supplement manifest, and all submission prose. The reference PDF is byte-bound as an archived artifact, but PDF metadata timestamps mean that an independent rebuild is required to agree in mathematical content and diagnostics, not necessarily byte for byte.

The full regression command is stored verbatim in the manifest. The finite congruence certificate stores, for each of the 15 excluded mask-77 branches, the modulus and exhaustive residue predicate. The remaining three branches carry symbolic factor identities and thresholds.

Online Resource 1 (supplementary code and certificates). The supplementary ZIP contains the exact Python generators, regression tests, JSON certificates and nested manifest used to audit the finite pattern counts and the mask-77, mask-89, mask-102 and mask-108 exclusions. It contains no additional mathematical claim beyond the theorem and claim boundary stated in this article. Before submission, the file metadata and upload form must add the approved article title, journal name, author names, affiliations and corresponding-author e-mail address.

9 Prior-art boundary and limitations

Work over number fields [10, 5, 1, 3, 4] asks for terms that are themselves squares in a prescribed extension. Our affine formulation allows Proposition 2’s common rational scaling. Nearby algorithmic work concerns S -integral points [9], simultaneous Pell equations [8], and integral points in progression on congruent-number curves [2]. Exact-quartic, S -integral-transform and Pell-system searches found no published statement identical to Theorems 7 and 8. This is a *not found* report, not proof of novelty.

The manifest and nested certificates bind SAFE and Round-04 inputs by SHA-256 and store the 18 branches, structured factor identities, 44 same-parameter audits, and 54 remaining identifiers. The fourth-round certificate separately stores the mask-102 ranking, proof data, the 19 excluded identifiers and the 35 intermediate survivors; the mask-108 certificate stores its complete point proof data, 12 exclusions and 23 final candidates. An end-to-end test regenerates 651–343–284 and all $284 \cdot 15 = 4260$ characters. We do not prove that the final 23 patterns are realized, or that no seven-term rank-two progression exists. The 2026 papers have additional rank and class-number hypotheses; none is inferred from a label here.

Statements and Declarations

Funding. AUTHOR CONFIRMATION REQUIRED. The human authors must insert and approve a complete and accurate funding statement before submission; no funding status is inferred here.

Competing interests. AUTHOR CONFIRMATION REQUIRED. The human authors must insert and approve a complete statement of relevant financial and non-financial interests before submission; no absence of interests is inferred here.

Author identity and contributions. AUTHOR CONFIRMATION REQUIRED.

No author identity, affiliation, corresponding-author information or contribution allocation is inferred. Every listed author must satisfy the journal’s authorship criteria, approve the final manuscript and accept accountability for the work.

Use of large language models and AI-assisted tools.

Large language models were used in this project for mathematical exploration, generation and revision of computer code, literature-search assistance, and generation and revision of manuscript and submission text. No AI system is an author. Before submission, the human authors must independently verify every proof, computation, code path, citation and statement, confirm that this disclosure accurately describes the use made, approve the final text and accept full responsibility for its integrity.

Data and code availability.

The code, certificates and manifests are currently only a local release candidate and have not been deposited in a public repository. **AUTHOR CONFIRMATION REQUIRED.** Before submission, the human authors must approve an immutable public deposit, insert its genuine persistent identifier here, and verify the downloaded archive against the published SHA-256 manifest. No DOI or public URL is claimed in this draft.

Ethics approval.

Not applicable: this mathematical study involved no human participants, animals or personal data.

References

- [1] Andrew Bremner and Samir Siksek. Squares in arithmetic progression over cubic fields. *International Journal of Number Theory*, 12(5):1409–1414, 2016.
- [2] Andrew Bremner, Joseph H. Silverman, and Nikos Tzanakis. Integral points in arithmetic progression on $y^2 = x(x^2 - n^2)$. *Journal of Number Theory*, 80(2):187–208, 2000.
- [3] Enrique González-Jiménez. Squares in arithmetic progression over quadratic extensions of number fields. *International Journal of Number Theory*, 22(6):1141–1158, 2026.
- [4] Enrique González-Jiménez and Nguyen Xuan Tho. Squares in arithmetic progression over certain non-primitive quartic number fields. *Journal of Number Theory*, 282:13–30, 2026.
- [5] Enrique González-Jiménez and Xavier Xarles. Five squares in arithmetic progression over quadratic fields. *Revista Matemática Iberoamericana*, 29(4):1211–1238, 2013.
- [6] Enrique González-Jiménez and Xavier Xarles. On a conjecture of rudin on squares in arithmetic progressions. *LMS Journal of Computation and Mathematics*, 17(1):58–76, 2014.
- [7] LMFDB Collaboration. Elliptic curve 576.c3, 2026. Accessed 3 September 2026.
- [8] David W. Masser and John H. Rickert. Simultaneous pell equations. *Journal of Number Theory*, 61(1):52–66, 1996.
- [9] Attila Pethő, Horst G. Zimmer, Josef Gebel, and Emanuel Herrmann. Computing all S -integral points on elliptic curves. *Mathematical Proceedings of the Cambridge Philosophical Society*, 127(3):383–402, 1999.
- [10] Xavier Xarles. Squares in arithmetic progression over number fields. *Journal of Number Theory*, 132(3):379–389, 2012.