

ARTHUR LI

ABSTRACT ALGEBRA

Introduction

THIS COLLECTION of notes serve as a guide to mastering abstract algebra with content from undergraduate to graduate level course. The notes combine knowledge from different sources, including course notes and textbooks used in the courses.

The proofs for Theorems, Propositions and Lemmas will be added when I have completed the entire skeleton.

Prerequisites

These notes will assume no familiarity with any aspects of abstract algebra, and builds upon the foundation from Group Theory to more abstract topics such as Categories and Commutative Algebra. A good starting point will be the series on [Visual Group Theory](#) by [Professor Matthew Macauley](#).

Familiarity with basic styles of proof is assumed (contradiction, contrapositive, etc.).

Organization and Sources

This section will be edited as the notes progress towards completion.

Contents

1	<i>Preliminaries</i>	5
1.1	<i>Introductory Ideas and Definitions</i>	5
2	<i>Group Theory</i>	7
2.1	<i>Basic Axioms</i>	7
2.2	<i>Homomorphisms and Subgroups</i>	7
2.3	<i>Cyclic Groups</i>	7
2.4	<i>Cosets</i>	7
2.5	<i>Normality, Quotient Groups</i>	7
2.6	<i>Isomorphism Theorems</i>	7
2.7	<i>Symmetric, Alternating and Dihedral Groups</i>	7
2.8	<i>Categories, Products, Coproducts, Free Objects</i>	7
2.9	<i>Direct Products, Direct Sums</i>	7
2.10	<i>Free Groups, Free Products</i>	7
2.11	<i>Matrix Groups</i>	7
3	<i>Group Structures</i>	8
3.1	<i>Free Abelian Groups</i>	8
3.2	<i>Finitely Generated Abelian Groups</i>	8
3.3	<i>Krull-Schmidt Theorem</i>	8
3.4	<i>Group Action</i>	8
3.5	<i>The Sylow Theorems</i>	8
3.6	<i>Semidirect Products</i>	8
3.7	<i>Normal and Subnormal Series</i>	8

4	<i>Ring Theory</i>	9
4.1	<i>Basic Axioms</i>	9
4.2	<i>Examples of Rings</i>	12
4.3	<i>Ring Homomorphisms</i>	14
4.4	<i>Ring Isomorphisms</i>	18
4.5	<i>Ideals, Rings of Fractions, Local Rings</i>	19
4.6	<i>Euclidean Domains, PID, UFD</i>	19
5	<i>Modules</i>	20
5.1	<i>Basic Axioms</i>	20
6	<i>Category Theory</i>	21
6.1	<i>Basic Axioms</i>	21

1 Preliminaries

1.1 Introductory Ideas and Definitions

Definition 1.1.1. *Class* is a collection A of objects (elements) such that given any object x it is possible to determine if x is a member of A .

Definition 1.1.2. *Axiom of extensionality* asserts that two classes with the same elements are equal.
(Formally, $[x \in A \iff x \in B] \Rightarrow A = B$).

Definition 1.1.3. A class is defined to be a *set* if and only if there exists a class B such that $A \in B$.
A class that is not a set is called a *proper set*.

Definition 1.1.4. *Axiom of class formation* asserts that for any statement $P(y)$ in the first predicate calculus involve a variable y , there exists a class A such that $x \in A$ if and only if x is a set and the statement $P(x)$ is true. The class is denoted $\{x|P(x)\}$.

Definition 1.1.5. A class A is a *subclass* of class B ($B \supset A$) provided $\forall x \in A, x \in A \iff x \in B$.
A subclass A of a class B that is itself a set is called a *subset* of B .
The *empty or null set* (denoted \emptyset) is the set with no elements.

Definition 1.1.6. *Power axiom* asserts that for every set A the class $P(A)$ of all subsets of A is itself a set. $P(A)$ is the *power set* of A , denoted 2^A .

Definition 1.1.7. A *family of sets* indexed by (nonempty) class I is a collection of sets A_i , one for each $i \in I$ (denoted $\{A_i|i \in I\}$).

The *union* is defined as $\bigcup_{i \in I} A_i = \{x|x \in A_i \text{ for some } i \in I\}$.

The *intersection* is defined as $\bigcap_{i \in I} A_i = \{x|x \in A_i \text{ for every } i \in I\}$.

If $A \cap B = \emptyset$, then A and B are disjoint.

Definition 1.1.8. The *relative complement* of A in B is the following subclass of B : $B - A = \{x|x \in B \text{ and } x \notin A\}$. If all classes under discussion are subsets of some fixed set U (the universe of discussion), then $U - A = A'$ is the *complement* of A .

Definition 1.1.9. Given classes A and B , a **function / map / mapping** f from A to B (written $f : A \rightarrow B$ assigns to each $a \in A$ exactly one element $b \in B$).

Then b is the value of function at a , or the **image** of a , written $f(a)$.

A is the **domain** of the function, written $\text{dom } f$, and B is the **range** or **codomain**.

Two functions are **equal** if they have the same domain and range, and have the same value for each element of their common domain.

Definition 1.1.10. If $f : A \rightarrow B$ is a function and $S \subset A$, the function from S to B given by $a \mapsto f(a)$, for $a \in S$, is **restriction** of f to S , denoted $f|_S : S \rightarrow B$.

If $S \in A$, the function $1_A|_S : S \rightarrow A$ is the **inclusion map** of S into A .

Definition 1.1.11. Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be functions. The **composite** of f and g is the function $A \rightarrow C$ given by $a \mapsto g(f(a))$, $a \in A$. This is denoted $g \circ f$ or simply gf .

Definition 1.1.12. The **diagram of functions** is said to be commutative if $gf = h$, or if $kh = gf$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow h & \swarrow g \\ & C & \end{array} \quad \begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow h & & \downarrow g \\ C & \xrightarrow{k} & D \end{array} \quad (1.1)$$

Definition 1.1.13. Let $f : A \rightarrow B$ be a function. If $S \in A$, the **image of S under f** (denoted $f(S)$) is the class $\{b \in B | b = f(a) \text{ for some } a \in S\}$.

The class $f(A)$ is the **image of f** , denoted $\text{Im } f$.

If $T \subset B$, the **inverse image of T** under f (denoted $f^{-1}(T)$), is the class $\{a \in A | f(a) \in T\}$.

Definition 1.1.14. A function $f : A \rightarrow B$ is said to be **injective** (or one-to-one) provided $\forall a, a' \in A, a \neq a' \Rightarrow f(a) \neq f(a')$, or $f(a) = f(a') \Rightarrow a = a'$.

A function f is **surjective** (or on-to) provided $f(A) \approx B$; in other words, for each $b \in B$, $b = f(a)$ for some $a \in A$.

A function f is **bijective** (or one-to-one correspondence) if it is both injective and surjective.

Definition 1.1.15. The map $g : B \rightarrow A$ is a **left inverse** of f if $gf = 1_A$.

The map $h : B \rightarrow A$ is a **right inverse** of f if $fb = 1_B$.

If a map $f : A \rightarrow B$ has both a left inverse g and a right inverse h , then $g = g1_B = g(fh) = (gf)h = 1_Ah = h$, and $g = h$ is the **two-sided inverse**.

2 *Group Theory*

2.1 *Basic Axioms*

2.2 *Homomorphisms and Subgroups*

2.3 *Cyclic Groups*

2.4 *Cosets*

2.5 *Normality, Quotient Groups*

2.6 *Isomorphism Theorems*

2.7 *Symmetric, Alternating and Dihedral Groups*

2.8 *Categories, Products, Coproducts, Free Objects*

2.9 *Direct Products, Direct Sums*

2.10 *Free Groups, Free Products*

2.11 *Matrix Groups*

3 *Group Structures*

3.1 *Free Abelian Groups*

3.2 *Finitely Generated Abelian Groups*

3.3 *Krull-Schmidt Theorem*

3.4 *Group Action*

3.5 *The Sylow Theorems*

3.6 *Semidirect Products*

3.7 *Normal and Subnormal Series*

4 Ring Theory

4.1 Basic Axioms

Definition 4.1.1. A **ring** is a nonempty set R with two binary operations $+$ (addition) and \times (multiplication), $(R, +, \times)$, such that:

- (i) $(R, +)$ is an additive abelian group with 0 as the additive identity
- (ii) the binary operation \times is associative: $(a \times b) \times c = a \times (b \times c)$, $\forall a, b, c \in R$
- (iii) left and right distributive laws: $(a + b) \times c = (a \times c) + (b \times c)$, $a \times (b + c) = (a \times b) + (a \times c)$, $\forall a, b, c \in R$.

Definition 4.1.2. If in addition to definition of ring, $a \times b = b \times a \forall a, b \in R$, then R is a **commutative ring**.

Definition 4.1.3. The ring R has a **multiplicative identity** if there is an element $1_R \in R$ such that $1_R \times a = a \times 1_R = a$, $\forall a \in R$.

The ring R has a **additive identity** if there is an element $0_R \in R$ such that $a - b = a + (-b) = 0_R$, where $-b$ is the **additive inverse**.

Definition 4.1.4. A **division ring** R is a ring such that:

- (i) R has a multiplicative identity 1_R ;
- (ii) $1_R \neq 0_R$; and
- (iii) \forall nonzero element $a \in R \setminus \{0\}$ has a unique multiplicative inverse a^{-1} such that $aa^{-1} = 1 = a^{-1}a$

Definition 4.1.5. A **field** is a division ring which is commutative.

If R is a division ring (field), then (R, \times) is a (commutative) **multiplicative group**, $R^\times = R \setminus \{0\}$.

Definition 4.1.6. Let $F = (F, +, \times)$ be a field. A nonempty subset $E \subseteq F$ is a **subfield** if:

- (i) $(E, +)$ is an additive subgroup of $(F, +)$;
- (ii) E is closed under multiplication \times : $a, b \in E \Rightarrow a \times b \in E$;
- (iii) $1_F \in E$; and
- (iv) $a \in E \setminus \{0\} \Rightarrow a^{-1} \in E$

Remark 4.1.7. The *trivial ring* is $\{0\}$.

The *integer ring* is $(\mathbb{Z}, +, \times)$ with 1, but is neither a division ring or field.

$n\mathbb{Z} = \{ns | s \in \mathbb{Z}\}$ is a subring of \mathbb{Z} .

$(\mathbb{Z}/n\mathbb{Z}, +, \times)$ is a commutative ring with 1 for $n \geq 2$.

Remark 4.1.8. The 2-dimensional vector space $\mathbb{Q}[\sqrt{D}] = \mathbb{Q} + \mathbb{Q}\sqrt{D} = \{a + b\sqrt{D} | a, b \in \mathbb{Q}\}$ with \mathbb{Q} -basis $\{1, \sqrt{D}\}$ is a *Quadratic Field*.

Define $\mathbb{Q}(\sqrt{D}) = \left\{ \frac{a+b\sqrt{D}}{c+d\sqrt{D}} | a, b, c, d \in \mathbb{Q}, c+d\sqrt{D} \neq 0 \right\}$. Then $\mathbb{Q}(\sqrt{D}) = \mathbb{Q}[\sqrt{D}]$.

More generally, for a field F , $\mathbb{Q}(F) = \left\{ \frac{\alpha}{\beta} = \alpha\beta^{-1} | \alpha, \beta \in F, \beta \neq 0 \right\} = F$.

Remark 4.1.9. Let $H = \mathbb{R} + \mathbb{R}i + \mathbb{R}j + \mathbb{R}k = \{a + bi + cj + dk | a, b, c, d \in \mathbb{R}\}$ be the 4-dimensional vector space over \mathbb{R} with \mathbb{R} -basis $(1, i, j, k)$.

The multiplication is extended linearly by distributive law: $i^2 = j^2 = k^2 = -1$, $ij = k = -ji$, $jk = i = -kj$, $ki = j = -ik$. Then H is a *Real Quaternion Ring*.

$H_{\mathbb{Q}} = \mathbb{Q} + \mathbb{Q}i + \mathbb{Q}j + \mathbb{Q}k = \{a + bi + cj + dk | a, b, c, d \in \mathbb{Q}\}$ is the *Rational Hamilton Quaternion Ring*.

Remark 4.1.10. Let $\mathbb{R}V[x] = \{f : \mathbb{R} \rightarrow \mathbb{R}\}$ be the set of all real-valued functions. Let $x \mapsto c(x) = c$ be a constant function.

For $f, g \in \mathbb{R}V[x]$, the natural addition is $x \mapsto (f + g)(x) = f(x) + g(x)$.

The multiplication (not composition) is $x \mapsto (fg)(x) = f(x)g(x)$.

The $(\mathbb{R}V[x], +, \times)$ is a commutative (*real valued-function*) *ring* with multiplicative identity 1 being the constant function 1.

Definition 4.1.11. Let R be a ring with $1 \neq 0$. An element $u \in R$ is a *unit* if it has a multiplicative identity inverse u' such that $uu' = 1 = u'u$.

The *set of all units* of R are $U(R) = \{u \in R | u \text{ is a unit}\}$.

The *multiplicative group of units of the ring* R is $(U(R), \times)$.

Remark 4.1.12. More generally, let X be a set and R be a ring. Let $X_{t_0}R := \{f : X \rightarrow R\}$ be the set of all maps between X and R . Then for $f, g \in X_{t_0}R$, there are natural addition $f + g$ and multiplication fg ($x \mapsto f(x)g(x)$) as in previous remark.

Then $(X_{t_0}R, +, \times)$ is a ring, called the *R-Valued Function Ring*.

If R has 1 then so does $X_{t_0}R$. If R is commutative then so does $X_{t_0}R$.

Every $c \in R$ defines a constant function (an element in $X_{t_0}R$, $c : X \rightarrow R$; $x \mapsto c(x) = c$).

Identify R with the subset of $X_{t_0}R$ of constant function. Then R is a subring of $X_{t_0}R$.

Remark 4.1.13. Let $n \geq 2$. Then $U(\mathbb{Z}/n\mathbb{Z})$ is a commutative multiplicative group of order $|U(\mathbb{Z}/n\mathbb{Z})| = \varphi(n)$. Hence $\varphi(n)$ is the *Euler's φ -function*, $\varphi(n) = |\{1 \leq s \leq n | \gcd(s, n) = 1\}|$.

Definition 4.1.14. An *Integral Domain* is a commutative ring with $1 \neq 0$ such that $\forall a, b \in R$, $ab = 0 \Rightarrow a = 0$ or $b = 0$, or equivalently, $\forall a, b \in R$, $a \neq 0$, $b \neq 0 \Rightarrow ab \neq 0$.

\mathbb{Z} is an integral domain.

Every field is an integral domain.

Definition 4.1.15. Let R be a ring. A nonzero element $a \in R$ is a **zero divisor** if there is a nonzero $b \in R$ such that either $ab = 0$ or $ba = 0$.

A commutative ring R with 1 is an integral domain if and only if R has no zero divisors.

Proposition 4.1.16. Let R be a ring with $1 \neq 0$. Then R is an integral domain if and only if the cancellation law holds: $\forall a, b, c \in R, c \neq 0, ca = cb \Rightarrow a = b$.

Corollary 4.1.17. Let R be a finite integral domain, i.e., R is an integral domain with the cardinality $|R| < \infty$. Then R is a field.

Proposition 4.1.18. Let $n \geq 2$. Then the following are equivalent:

- (i) $\mathbb{Z}/n\mathbb{Z}$ is a field
- (ii) $\mathbb{Z}/n\mathbb{Z}$ is an integral domain
- (iii) n is a prime

Definition 4.1.19. Let R be a ring. A nonempty subset $S \subseteq R$ is a **subring** of R if:

- (i) $(S, +)$ is an additive subgroup of $(R, +)$ and
- (ii) S is closed under multiplication

$$\mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$$

Proposition 4.1.20. (Subring Criterion) Let R be a ring and $S \subseteq R$ a nonempty subset. Then the following are equivalent:

- (i) S is a subring of R
- (ii) S is closed under subtracting and multiplication: $a, b \in S \Rightarrow ab \in S; a - b = a + (-b) \in S$

Remark 4.1.21. Being a subring is a transitive condition. If R is a subring of S and S is a subring of T , then R is a subring of T .

If both S_i are subrings of R and $S_1 \subseteq S_2$, then S_1 is a subring of S_2 .

Remark 4.1.22. (Subring without 1) If R is a ring with $1 = 1_R$ then a subring $S \subseteq R$ may not contain 1 , i.e., $m\mathbb{Z} = \{ms \mid s \in \mathbb{Z}, |m| \geq 2\}$ is a subring of \mathbb{Z} which does not contain 1 .

Remark 4.1.23. (Intersection of subrings) Let R_α ($\alpha \in \Sigma$) be a (not necessarily finite or countable) collection of subrings of a ring R . Then the intersection $\bigcap_{\alpha \in \Sigma} R_\alpha$ is a subring of R .

Generally, the union of subrings may not be a subring.

Remark 4.1.24. (*Union of ascending subrings*) Let $R_1 \subseteq R_2 \subseteq \cdots$ be an ascending chain of subrings R_i of a ring R . Then the union $\bigcup_{i=1}^{\infty} R_i$ is a subring of R .

Remark 4.1.25. (*Addition of subrings*) Let R be a ring and let R_i be subrings of R .

Then the addition $R_1 + \cdots + R_n$ is closed under subtraction, but may not be closed under multiplication, hence may not be a subring of R .

Remark 4.1.26. (*Integral domain is a subring of a field*)

Let F be a field. Let $R \subseteq F$ be a subring such that $1 \in R$. Then R is an integral domain.

Every integral domain R is a subring of some field $\mathbb{Q}(R)$ (the fractional field of R).

Remark 4.1.27. (*Product of Rings*) let $n \geq 1$ and let $R_i = (R_i, +, \times)$ ($i = 1, \dots, n$) be rings.

Then the direct product is a ring, $R = R_1 \times \cdots \times R_n$. (The direct product is $(a_1, \dots, a_n) \times (a'_1, \dots, a'_n) = (a_1 a'_1, \dots, a_n a'_n)$).

The unit subgroups has the relation $U(R) = U(R_1) \times \cdots \times U(R_n)$

4.2 Examples of Rings

Definition 4.2.1. The (*polynomial ring $R[x]$ over a ring R*) is $(R[x], +, \times)$,

where $R[x] = \{\sum_{j=0}^d b_j x^j \mid d \geq 0, b_j \in R\}$.

There are natural addition and multiplication operations for polynomials.

Remark 4.2.2. Let R be a commutative ring with 1. Let $S := R[x]$ be the polynomial ring over R .

(i) R is a subring of S which consists of constant polynomial functions.

(ii) $0_S = 0_R$

(iii) S contains $1 = 1_S$, and $1_S = 1_R$.

Proposition 4.2.3. (*Polynomial ring over integral domain*) Let R be an integral domain. Let $f(x), g(x) \in R[x]$. Then

(i) $\deg(f(x)g(x)) = \deg(f(x)) + \deg(g(x))$

(ii) $U(R[x]) = U(R)$. Namely, $g(x)$ is a unit of $R[x]$ if and only if $g = a_0 \in R$ (constant polynomial) with a_0 a unit in R .

(iii) $R[x]$ is an integral domain

Remark 4.2.4. The (*matrix ring of $n \times n$ square matrices with entries in the ring R*) is defined as $(M_n(R), +, \times)$, where

$$M_n(R) = \left\{ A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix} \mid a_{ij} \in R \right\}$$

If $A = (a_{ij}), B = (b_{ij}) \in M_n(F)$, then $A + B = (a_{ij} + b_{ij}), AB = (c_{ij})$ where $c_{ij} = \sum_{k=1}^n a_{ik}b_{kj}$.
 $A = (a_{ij}) = \text{Diag}[a_{11}, \dots, a_{nn}]$ is a diagonal matrix if $a_{ij} = 0$ ($i \neq j$).
 $A = (a_{ij}) = \text{Diag}(a_1, \dots, a_n)$ is a scalar matrix if $a_{ii} = a \in R \forall i$, and $a_{ij} = 0$ ($i \neq j$).
 $A = (a_{ij})$ is an upper triangular matrix if $a_{ij} = 0$ ($i < j$). The lower triangular matrix is defined similarly.

Remark 4.2.5. Let R be a ring and $S = M_n(R)$ the matrix ring with entries in R . Then

- (i) $0_S = (a_{ij})$ where $a_{ij} = 0$ (the zero matrix)
- (ii) If R has $1 = 1_R$, then S also has $1 = 1_S$ with $1_S = \text{Diag}[1_R, \dots, 1_R]$
- (iii) The set $S_{c_n}(R) = \{\text{Diag}[a, \dots, a] | a_i \in R\}$ of all scalar matrices in $M_n(R)$ is a subring of $M_n(R)$. There is a natural ring isomorphism $R \cong S_{c_n}(R)$.
- (iv) The set $D_n(R) = \{\text{Diag}[a_1, \dots, a_n] | a_i \in R\}$ of all diagonal matrices in $M_n(R)$ is a subring of $M_n(R)$. There is a natural ring isomorphism $D_n(R) \cong R^n := R \times \dots \times R$ (n times).
- (v) The set $UT_n(R) := \{(a_{ij}) | (a_{ij} \in R, (a_{ij} = 0 (\forall i > j)))\}$ of all upper triangular matrices in $M_n(R)$ is a subring of $M_n(R)$. Similarly, the set $LT_n(R)$ of all lower triangular matrices in $M_n(R)$ is a subring of $M_n(R)$.
- (vi) If R is a subring of R , then $M_n(T)$ is a subring of $M_n(R)$
- (vii) Even if R is commutative, $M_n(R)$ may not be commutative when $n \geq 2$.
- (viii) If $n \geq 2$, then $M_n(R)$ is not an integral domain (even when R is a field).

Definition 4.2.6. Let R be a ring with 1. Set $GL_n(R) := U(M_n(R))$ the set of all units in $M_n(R)$. Then $GL_n(R)$ is a multiplicative group called the **general linear group of degree n over R** .

Definition 4.2.7. Let R be a commutative ring with 1. Define determinant $\det(A) = |A|$, Let $SL_n(R) := \{A \in M_n(R) | \det(A) = 1\}$ be the set of all matrices in $M_n(R)$ with determinants equal to 1. Then $SL_n(R)$ is a multiplicative subgroup of $GL_n(R)$ called the **special linear group of degree n over R** .

Definition 4.2.8. (Group Rings $R[G]$)

Let R be a commutative ring with $1 \neq 0$. Let $G = \{g_1, \dots, g_n\}$ be a finite multiplicative group of order n . Then $R[G]$ is a **group ring**, where $R[G] = Rg_1 + \dots + Rg_n = \{a_1g_1 + \dots + a_ng_n | a_i \in R\}$.
Natural addition is defined as $(\sum_{i=1}^n a_i g_i) + (\sum_{i=1}^n b_i g_i) = (\sum_{i=1}^n (a_i + b_i) g_i)$.
Multiplication is defined as $(\sum_{i=1}^n a_i g_i) \times (\sum_{j=1}^n b_j g_j) := (\sum_{k=1}^n c_k g_k)$ where $c_k = \sum_{g_i g_j = g_k} a_i b_j$ with the sum running $\forall (i, j)$ with $g_i g_j = g_k$.

Remark 4.2.9. Let R be a commutative ring with $1 \neq 0$, G a multiplicative group, and $R[G]$ the group ring. Then

- (i) $R[G]$ is a commutative ring if and only if G is commutative (=abelian) group
- (ii) $R[G]$ has the multiplicative identity $1 = 1_R e_G$

Remark 4.2.10. Let $R[G]$ be a group ring.

(i) There is a natural injective ring homomorphism $R \rightarrow R[G]; r \mapsto r e_G$. Identify R with the image Re_G of this injective homomorphism.

- (ii) For every $g \in G$, the element $1_R g$ is a unit in $R[G]$
- (iii) There is a natural injective group homomorphism $G \rightarrow U(G[R]); g \mapsto 1_R g$. Identify G with the image $1_R G$ of this injective homomorphism.
- (iv) If S is a subring of R , then $S[G]$ is a subring of $R[G]$. If H is a subgroup of G , then $R[H]$ is a subring of $R[G]$.
- (v) $T = \{\sum_{i=1}^n a_i g_i \in R[G] \mid \sum_{i=1}^n a_i = 0\}$ is a subring of $R[G]$ (an ideal of $R[G]$)

Remark 4.2.11. When R is a division ring or field, then $R[G]$ (as an additive group) is a vector space over R of dimension equal to $|G|$ with basis $\{g_1, \dots, g_n\} = G$. Hence $R[G] = Rg_1 + \dots + Rg_n = Rg_1 \oplus \dots \oplus Rg_n$, the direct sum of 1-dimensional vector subspaces Rg_i over R .

4.3 Ring Homomorphisms

Definition 4.3.1. Let R, S be rings. A map $\varphi : R \rightarrow S$ is a **ring homomorphism** if it respects the additive and multiplicative structures.

$$\begin{aligned}\varphi(a + b) &= \varphi(a) + \varphi(b) \quad \forall a, b \in R \\ \varphi(ab) &= \varphi(a)\varphi(b) \quad \forall a, b \in R\end{aligned}$$

Definition 4.3.2. Let R, S be rings. A map $\varphi : R \rightarrow S$ is a **ring isomorphism** if it is a ring homomorphism and bijective. This is denoted $\varphi : R \xrightarrow{\sim} S$. Rings R and S is **isomorphic**, denoted $R \cong S$ or $R \simeq S$.

Definition 4.3.3. The **kernel** of a ring homomorphism φ is defined as $\ker \varphi = \varphi^{-1}(0_S) = \{a \in R \mid \varphi(a) = 0_S\}$.

Remark 4.3.4. (Examples of homomorphism)

- (i) Let R, S be rings. The map $R \rightarrow S, a \mapsto 0$ is a **zero or trivial map / homomorphism**.
- (ii) Suppose R_1 is a subring of a ring R . The map $\iota : R_1 \rightarrow R, a \mapsto a$ is a **inclusion homomorphism**.
- (iii) Let $n \in \mathbb{Z}$. The quotient map $\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}, s \mapsto \bar{s} = [s]_n$ is a **quotient homomorphism** between additive groups $(\mathbb{Z}, +)$ and $((\mathbb{Z})/n(\mathbb{Z}), +)$.
- (iv) Let X be a set, R a ring, and $X_{t_0}R = \{f : X \rightarrow R\}$ the ring of all maps from X to R . Fix an element $c \in R$. Then $E_c : X_{t_0}R \rightarrow R; f \mapsto E_c(f) := f(c)$ is a **function evaluation map**, called the **Evaluation at c** .

Proposition 4.3.5. Let R, S be rings and $\varphi : R \rightarrow S$ be a ring homomorphism. Let $R_1 \subseteq R$ be a subring. Then

- (i) The φ -image $\varphi(R_1) = \{b \in S \mid b = \varphi(a) \text{ for some } a \in R_1\}$ is a subring of S .
- (ii) $\ker \varphi$ is a subring of R such that $\forall a \in R, \forall k \in \ker \varphi \Rightarrow ak \in \ker \varphi$. In other words, $\ker \varphi$ is a subring of R , $R(\ker \varphi) \subseteq \ker \varphi$ and $(\ker \varphi)R \subseteq \ker \varphi$.

Definition 4.3.6. Let R be a ring, $I \subseteq R$ a subset and $r \in R$. The subset $I \subseteq R$ is a **left-ideal** of R if:

- (i) I is a subring of R ; and

(ii) I is closed under left multiplication by elements from R : $rI \subseteq I$ ($\forall r \in R$), i.e., $RI \subseteq I$.

Definition 4.3.7. Let R be a ring, $I \subseteq R$ a subset and $r \in R$. The subset $I \subseteq R$ is a **right-ideal** of R if:

(i) I is a subring of R ; and

(ii) I is closed under right multiplication by elements from R : $Ir \subseteq I$ ($\forall r \in R$), i.e., $IR \subseteq I$.

Definition 4.3.8. Let R be a ring, $I \subseteq R$ a subset and $r \in R$. The subset $I \subseteq R$ is a **(two-sided) ideal** of R if is both a left-ideal and right-ideal. In other words, $RI \subseteq I$ and $IR \subseteq I$.

Proposition 4.3.9. (Ideal Criterion) Let R be a ring and I a nonempty subset of R . The following are equivalent:

(i) I is a two-sided ideal of R ;

(ii) $\forall r \in R, \forall a, b \in R \Rightarrow ra, ar, a - b \in I$

(iii) (If R is commutative) $\forall r \in R, \forall a, b \in R \Rightarrow ra, a - b \in I$

(iv) (If R is commutative with 1) $\forall r \in R, \forall a, b \in R \Rightarrow a + rb \in I$

Proposition 4.3.10. Let R_α ($\alpha \in \Sigma$) be a family of subrings of a ring R . Let J_α be a left (resp. 2-sided) ideal of R_α . Then the intersection $\bigcap_{\alpha \in \Sigma} J_\alpha$ is a left (resp. 2-sided) ideal of the subring $\bigcap_{\alpha \in \Sigma} R_\alpha$.

Corollary 4.3.11. Let J_α ($\alpha \in \Sigma$) be a family of left (resp. 2-sided) ideals of a ring R . Then the intersection $\bigcap_{\alpha \in \Sigma} J_\alpha$ is also a left (resp. 2-sided) ideal of R .

Proposition 4.3.12. Let J_α ($\alpha \in \Sigma$) be a finite family of left (resp. 2-sided) ideals of a ring R . Then the addition $\sum_{\alpha \in \Sigma} J_\alpha$ is also a left (resp. 2-sided) ideal of R .

More generally, if J_α ($\alpha \in \Sigma$) is an infinite (countable or uncountable) family of left (resp. 2-sided) ideals of a ring R , then the subset $\{\sum x_\alpha | x_\alpha \in J_\alpha, x_\alpha \neq 0 \text{ for only finitely many } \alpha\}$ is also a left (resp. 2-sided) ideal of R .

Definition 4.3.13. Let X be a subset of a ring R . Let J_α ($\alpha \in \Sigma$) be all the ideals of R with $J_\alpha \supseteq X$.

Then the intersection $\bigcap_{\alpha \in \Sigma} J_\alpha$ is the **ideal generated by X** , denoted (X) .

This (X) is the smallest among all ideals of R containing X .

If $X = \{r_1, \dots, r_n\}$, then write $(X) = (r_1, \dots, r_n)$.

Definition 4.3.14. For $r \in R$, the ideal (r) generated by a single element r is the **principal ideal of ring R** .

Definition 4.3.15. Let R be a ring. An ideal I is **finitely generated** if $I = (r_1, \dots, r_n)$ for some $r_i \in R$.

Proposition 4.3.16. Let R be a ring; X, Y, X_i the subsets of R ; and $r_j \in R$.

- (i) Let J be an ideal of R . Then $(X) \subseteq J$ if and only if $X \subseteq J$.
- (ii) The equality of ideals holds: $(X_1 \cup \cdots \cup X_n) = (X_1) + \cdots + (X_n)$.
- (iii) In particular, $(r_1, \dots, r_n) = (r_1) + \cdots + (r_n)$.

Proposition 4.3.17. Let R be a ring. Let $B \subseteq R$ and $a, a_1, \dots, a_n \in R$.

- (i) $RB = \{\sum_{i=1}^s r_i b_i \mid r_i \in R, b_i \in B, s \geq 1\}$ is a left-ideal of R , but may not be a 2-sided ideal.
- (ii) More generally, $R\{a_1, \dots, a_n\} = Ra_1 + \cdots + Ra_n = \{\sum_{i=1}^n r_i a_i \mid r_i \in R\}$ are left-ideals of R , but they may not be 2-sided ideals.
- (iii) The ideal (a) generated by a is given by $(a) = \mathbb{Z}a + aR + Ra + RaR$. An arbitrary element of (a) is of the form $ma + ar + r'a + \sum_{i=1}^n r_i a r'_i$ where $m \in \mathbb{Z}; r, r', r_i, r'_i \in R; n \geq 1$.
- (iv) If R contains 1, then $(a) = RaR$, and an arbitrary element of (a) is of the form $\sum_{i=1}^n r_i a r'_i$ where $r_i, r'_i \in R; n \geq 1$.
- (v) If R is commutative and contains 1, then $(a) = aR = Ra = ra \mid r \in R$. An arbitrary element of (a) is of the form ra where $r \in R$.

Proposition 4.3.18. Let R be a ring with $1 \neq 0$ and I an ideal of R . Then the following are equivalent:

- (i) $I = R$
- (ii) $1 \in I$
- (iii) I contains a unit.

Proposition 4.3.19. Suppose R is a ring with 1. Let $X \subseteq R$ be a subset, and $b_1, \dots, b_n \in R$. Then

- (i) the ideal generated by X is $(X) = RXR = \{\sum_{i=1}^s r_i a_i r'_i \mid a_i \in X; r_i, r'_i \in R; s \geq 1\}$, the smallest among all ideals of R containing X .
- (ii) the ideal generated by $\{b_1, \dots, b_n\}$ is given by $(b_1, \dots, b_n) = (b_1) + \cdots + (b_n) = Rb_1R + \cdots + Rb_nR$, the smallest among all ideals of R containing $\{b_1, \dots, b_n\}$.

Proposition 4.3.20. Let J_α ($\alpha \in \Sigma$) be a family of left (resp. 2-sided) ideals of a ring R .

Then the inclusion is $R(\bigcup_{\alpha \in \Sigma} J_\alpha) \subseteq \{\sum_{\alpha \in \Sigma} a_\alpha \mid a_\alpha \in J_\alpha; a_\alpha \neq 0 \text{ for only finitely many } \alpha\}$, where the RHS is a left (resp. 2-sided) ideal of R , and the smallest among those of R containing all J_α , where LHS = RHS when R contains 1.

If R contains 1 and Σ is finite, then $R(\bigcup_{\alpha \in \Sigma} J_\alpha) = \sum_{\alpha \in \Sigma} J_\alpha$.

Proposition 4.3.21. Let J, J_1, \dots, J_n be ideals of a ring.

Then $J_1 \cdots J_n = \{\sum_{l=1}^k a_1(l) \cdots a_n(l) \mid a_i(l) \in J_i, k \geq 1\}$ and it is an ideal of R .

in particular, $J^n = J \cdots J = \{\sum_{l=1}^k a_1(l) \cdots a_n(l) \mid a_i(l) \in J, k \geq 1\}$ and it is an ideal of R .

Proposition 4.3.22. Let $R = R_1 \times \cdots \times R_n$ be a direct product of rings.

Then $S_i = \{0_{R_1}\} \times \cdots \times \{0_{R_{i-1}}\} \times R_i \times \{0_{R_{i+1}}\} \times \cdots \times \{0_{R_n}\}$ is an ideal (2-sided) of R .

Furthermore, $R = \sum_{i=1}^n S_i$.

Proposition 4.3.23. Let $\varphi : R \rightarrow S$ be a ring homomorphism.
Then $\ker \varphi$ is an ideal of R .

Definition 4.3.24. Let R be a ring and $I \subseteq R$ an ideal. Then $(I, +)$ is a normal subgroup of additive group $(R, +)$.
The *quotient additive group* is $R/I = \{\bar{r} = r + I \mid r \in R\}$, with well-defined addition $\bar{r} + \bar{s} := r + s$.

Theorem 4.3.25. Let R be a ring and $I \subseteq R$ an ideal. Then

(i) for cosets $\bar{r}, \bar{s} \in R/I$, the multiplication $\bar{r} \times \bar{s} := \overline{rs}$ is a well-defined binary operation on R/I , i.e., this multiplication does not depend on the choice of representatives r, s of the cosets.

(ii) $(R/I, +, \times)$ is a ring with $0_{R/I} = \bar{0}_R$.

(iii) $\bar{r} = 0_{R/I} (= \bar{0}_R)$ if and only if $r \in I$.

Definition 4.3.26. Let R be a ring and $I \subseteq R$ an ideal.
Then the ring $(R/I, +, \times)$ is the *quotient ring* of R by I .

Remark 4.3.27. Let R be a ring and $(I, +)$ a subgroup of the additive group $(R, +)$.

Then I is an ideal of R if and only if the multiplication \times on the additive quotient group $(R/I, +)$ is well-defined so that $(R/I, +, \times)$ is a ring.

Definition 4.3.28. Let R be a ring, $I \subseteq R$ an ideal, and R/I the quotient ring.

The *surjective quotient map* $\gamma : R \rightarrow R/I; r \mapsto \bar{r} = r + I$ from the additive group $(R, +)$ to the additive group $(R/I, +)$ is a ring homomorphism such that $\ker \gamma = I$.

The *quotient ring homomorphism* refers to γ .

Remark 4.3.29. (*Equivalence concepts of kernel and ideal*)

The kernel of every ring homomorphism is an ideal.

Every ideal is equal to the kernel of some (surjective) homomorphism.

Definition 4.3.30. Let R be a commutative ring and I an ideal.

An element $a \in R$ is *nilpotent* if $a^n = 0$ for some $n \geq 1$ (depending on a).

The set of all nilpotent elements of R is the *nilradical of R* , $\text{nil}(R) := \{a \in R \mid a^n = 0, \text{ for some } n \geq 1\}$.

In fact, $\text{nil}(R)$ is an ideal of R , and $\text{nil}(R/\text{nil}(R)) = 0$.

Definition 4.3.31. Let R be a commutative ring and I an ideal.

The set of *radical of I* is $\text{rad}(I) = \{r \in R \mid r^n \in I, \text{ for some } n \geq 1\}$.

In fact, $\text{rad}(I)$ is an ideal of R containing I such that $\text{rad}(I)/I = \text{nil}(R/I)$.

Definition 4.3.32. Let R be a commutative ring and J an ideal.
 J is a radical if $\text{rad}(J) = J$. Every prime idea of R is ideal.

Definition 4.3.33. Let R be a commutative ring and I an ideal. When R contains 1 and $I \subset R$,
define $\text{Jac}(I) = \bigcap_{M: \max, M \supseteq I} M$, where M runs in the set of all maximal ideals of R containing I .
In fact, $\text{Jac}(I)$ is an ideal of R containing the radical $\text{rad}(I)$ of I .
 $\text{Jac}(0)$ is the *Jacobson radical of R* .
Thus $\text{Jac}(I)$ is the pre-image of $\text{Jac}(0_{R/I})$ via $R \rightarrow R/I$.

Remark 4.3.34. Let R be a commutative ring and I an ideal. Then $\text{nil}(R/I^n) \supseteq I/I^n$, and $\text{rad}(I^n) \supseteq I$ (the inclusions might be strict).

Remark 4.3.35. For the polynomial ring $F[x]$ over field F , if $I = (x)$ is the principal ideal generated by x , then $I^n = (x^n)$. Hence $\text{nil}(F[x]/I^n) = I/I^n$ and $\text{rad}(I^n) = I$.

Remark 4.3.36. The Jacobson radical of $\mathbb{Z}/12\mathbb{Z}$ is $6\mathbb{Z}/12\mathbb{Z}$, included in the intersection (of two maximal ideals) $(2\mathbb{Z}/12\mathbb{Z}) \cap (3\mathbb{Z}/12\mathbb{Z})$.

The Jacobson radical of the polynomial ring $F[x]$ over field F is 0, which is contained in the intersection (of two maximal ideals) $(x) \cap (x-1)$.

4.4 Ring Isomorphisms

Definition 4.4.1. (*First Isomorphism Theorem*)

Let $\varphi : R \rightarrow S$ be a ring homomorphism, $\gamma : R \rightarrow R/\ker \varphi$ the (surjective) quotient ring homomorphism, and $\bar{\varphi} : R/\ker \varphi \xrightarrow{\sim} \varphi(R)$; $\bar{r} \mapsto \bar{\varphi}(\bar{r}) := \varphi(r)$ a (well-defined) ring homomorphism..

Then $\varphi = \bar{\varphi} \circ \gamma$.

$$\begin{array}{ccc}
 R & \xrightarrow{\varphi} & \varphi(R) \\
 \searrow \gamma & & \nearrow \bar{\varphi} \\
 & R/\ker \varphi &
 \end{array} \tag{4.1}$$

Remark 4.4.2. Let R, S be a commutative ring with 1 and $\varphi : R \rightarrow S$ a ring homomorphism. Then φ induces a ring homomorphism $\tilde{\varphi} : R[x] \rightarrow S[x]$; $f(x) = \sum a_i x^i \mapsto \tilde{\varphi}(f(x)) = \sum \varphi(a_i) x^i$.

Furthermore, if $J = \ker \varphi$, then $\ker \tilde{\varphi} = J[x] = \{ \sum_{i=1}^n a_i x^i \mid a_i \in J, n \geq 1 \}$ is the polynomial ring with coefficients in J . Finally, $J[x] = JR[x]$ and $J[x]$ is the ideal of $R[x]$ generated by J , i.e., $J[x] = (J)$.

Remark 4.4.3. Let R be a commutative ring with 1 and I an ideal of R .

Then there is an isomorphism $R[x]/I[x] \cong (R/I)[x]$.

Remark 4.4.4. If $\varphi : R \rightarrow S$ is a ring homomorphism, then it induces a homomorphism (between matrix rings):
 $\varphi_n : M_n(R) \rightarrow M_n(S); A = (r_{ij}) \mapsto \varphi_n(A) := (\varphi(r_{ij}))$.

Remark 4.4.5. Let $G = \langle g_1, \dots, g_n \rangle$ be a multiplicative group of order $|G| = n$, R a ring, and $R[G] = Rg_1 + \dots + Rg_n$ the group ring.
 Then the map $\text{Tr} : R[G] \rightarrow R; \sum_{i=1}^n r_i g_i \mapsto \sum_{i=1}^n r_i$ is a ring isomorphism.

Remark 4.4.6. (One-sided Ideals)

Let $n \geq 2$ and $M_n(R)$ a matrix ring over a ring R . Let $L_k = \{A = (a_{ij}) \in M_n(R) | a_{ij} = 0, \forall j \neq k\}$.

Then L_k is a left ideal of $M_n(R)$, but not a right ideal of $M_n(R)$ when R contains 1_R .

Similarly, let $R_k = \{A = (a_{ij}) \in M_n(R) | a_{ij} = 0, \forall i \neq k\}$.

Then R_k is a right ideal of $M_n(R)$, but not a left ideal of $M_n(R)$ when R contains 1_R .

More generally, let $1 \leq k_1 < \dots < k_r \leq n$ with $r < n$.

Let $L_{k_1, \dots, k_r} = \{A = (a_{ij}) \in M_n(R) | a_{ij} = 0, \forall j \notin \{k_1, \dots, k_r\}\}$.

Then L_{k_1, \dots, k_r} is a left ideal of $M_n(R)$, but not a right ideal of $M_n(R)$ when R contains 1_R .

Let $R_{k_1, \dots, k_r} = \{A = (a_{ij}) \in M_n(R) | a_{ij} = 0, \forall i \notin \{k_1, \dots, k_r\}\}$.

Then R_{k_1, \dots, k_r} is a right ideal of $M_n(R)$, but not a left ideal of $M_n(R)$ when R contains 1_R .

Definition 4.4.7. (Second Isomorphism Theorem)

Let R be a ring, $R_1 \subseteq R$ a subring, and $J \subseteq R$ an ideal. Then:

(i) $R_1 + J$ is a subring of R

(ii) $R_1 \cap J$ is an ideal of R

(iii) There is an isomorphism $\varphi : R_1 / (R_1 \cap J) \xrightarrow{\sim} (R_1 + J) / J; \bar{r} = r + (R_1 \cap J) \mapsto \varphi(\bar{r}) := \bar{r} = r + J$.

Definition 4.4.8. (Third Isomorphism Theorem)

Let R be a ring, and $I \subseteq J$ ideals of R . Then:

(i) J/I is an ideal of the quotient ring R/I

(ii) There is an isomorphism $\varphi : R/J \xrightarrow{\sim} (R/I)/(J/I); \bar{r} = r + J \mapsto \bar{r} + J/I = (r + I) + J/I$

Definition 4.4.9. (Fourth Isomorphism Theorem)/Correspondence Theorem for Rings

Let R be a ring, $I \subseteq R$ an ideal, and $\gamma : R \rightarrow R/I$ the (surjective) quotient ring homomorphism.

Let Σ_1 be the set of subrings of R containing $I = \ker \gamma$, and Σ_2 be the set of subrings of R/I . Then:

(i) if $R_1 \in \Sigma_1$, then $\gamma(R_1) = R_1/I \in \Sigma_2$. Conversely, if $R'_1 \in \Sigma_2$, then $R'_1 = R_1/I$ with $R_1 := \gamma^{-1}(R'_1) = \{r \in R | \gamma(r) \in R'_1\} \in \Sigma_1$.

(ii) The map $f : \Sigma_1 \rightarrow \Sigma_2; R_1 \mapsto R_1/I$ is a well-defined bijection.

(iii) $J_1 \in \Sigma_1$ is an ideal of R if and only if J_1/I is an ideal of R/I . If this is the case, then $R/J_1 \cong (R/I)/(J_1/I)$

(iv) For $R_i \in \Sigma_1$, $R_1 \subseteq R_2$ holds if and only if $R_1/I \subseteq R_2/I$ holds.

4.5 Ideals, Rings of Fractions, Local Rings

4.6 Euclidean Domains, PID, UFD

5 *Modules*

5.1 *Basic Axioms*

6 *Category Theory*

6.1 *Basic Axioms*