# Mathematics

Arthur Li

July 19, 2026

# Contents

**Notation List** 1

## I Set Theory, Logic, and Foundations 10

**1 Sets** 10
1.1 Introduction to Sets . . . 10
1.2 Cartesian Products . . . 10
1.3 Subsets and Power Sets . . . 10
1.4 Union, Intersection, Difference, and Complement . . . 11
1.5 Indexed Families and Products . . . 11
1.6 Russell's Paradox and Axioms . . . 12

**2 Logic and Proofs** 12
2.1 Statements and Well-Formed Language . . . 12
2.2 Connectives and Negation . . . 12
2.3 Quantifiers and Bound Variables . . . 12
2.4 Proof Grammar . . . 12
2.5 Proof by Contradiction . . . 13
2.6 Mathematical Induction and Well-Ordering . . . 13

**3 Counting and Finite Sums** 13
3.1 Lists and Counting Principles . . . 13
3.2 Permutations, Subsets, and Binomial Coefficients . . . 14
3.3 Inclusion-Exclusion and Combinatorial Proof . . . 15
3.4 Finite Sums . . . 16

**4 Relations, Orders, and Lattices** 16
4.1 Relations and Equivalence . . . 16
4.2 Partitions and Quotient Sets . . . 16
4.3 Ordered Sets, Well-Orders, and Lattices . . . 17
4.4 Relations Between Sets and Closures . . . 18

**5 Functions** 18
5.1 Functions, Images, and Preimages . . . 18
5.2 Injective, Surjective, and Bijective Maps . . . 19
5.3 Composition, Restriction, and Inverse Functions . . . 20

**6 Cardinality, Choice, and Infinite Sets** 20
6.1 Same Cardinality, Finite, Countable, and Uncountable . . . 20
6.2 Diagonalization and Power Sets . . . 24
6.3 Schroeder–Bernstein and Cardinal Comparison . . . 24
6.4 Cardinal Arithmetic . . . 25
6.5 Choice and Zorn's Lemma . . . 25

**7 Axiomatic and Advanced Set Theory** 26
7.1 Extensionality, Specification, and Replacement . . . 26
7.2 Ordinals, Transfinite Recursion, and Well-Foundedness . . . 26
7.3 Models, Constructibility, and Independence . . . 26
7.4 Filters, Ultrafilters, and Sets of Reals . . . 26

## II Structural Linear Algebra 27

**1 Vector Spaces and Linear Structure** 27
1.1 Vector Spaces . . . 27
1.2 Subspaces and Spans . . . 28
1.3 Linear Independence and Bases . . . 29
1.4 Dimension and Coordinates . . . 32
1.5 Linear Maps and Operators . . . 33

1.6 Kernels, Ranges, and Map Properties . . . 35
1.7 Rank-Nullity and Dimension Criteria . . . 36

**2 Direct Sums, Quotients, and Decomposition 38**
2.1 Sums and Direct Sums . . . 38
2.2 Quotient Spaces . . . 41
2.3 Decomposition and Complementation . . . 43
2.4 Algebraic Projections and Resolutions . . . 46

**3 Matrix Representations, Change of Basis, Block Matrices, and Operations 47**
3.1 Matrix Algebra and Linear Systems . . . 47
3.2 Coordinates and Matrix Representations . . . 49
3.3 Matrix-Induced Maps and Rank . . . 50
3.4 Invertibility and the Matrix-Map Dictionary . . . 51
3.5 Change of Basis and Coordinate Transformations . . . 52
3.6 Rank and Invertibility Consequences . . . 52
3.7 Block Matrix Algebra . . . 53
3.8 Matrix Operations and Identities . . . 54

**4 Duality and Multilinear Structure 55**
4.1 Dual Spaces and Dual Bases . . . 55
4.2 Double Duals and Dual Maps . . . 56
4.3 Annihilators and Transposes . . . 58
4.4 Multilinear Structure . . . 60

## III Linear Maps and Matrix Representations 63

**1 Linear Maps and Operations 63**
1.1 Further Linear Map Operations and Examples . . . 63

**2 Kernels, Images, Rank, and Nullity 63**
2.1 Shift Operators and Kernel-Range Examples . . . 63
2.2 Further Rank-Nullity and Invertibility Results . . . 63
2.3 Further Quotients and Isomorphism Theorems . . . 63
2.4 Further Finite-Dimensional Splitting and Isomorphism Tests . . . 63

**3 Matrix Representations 63**
3.1 Further Matrix Algebra and Linear Systems . . . 63
3.2 Further Coordinates and Matrix Representations . . . 63
3.3 Further Matrix-Induced Maps and Rank . . . 63
3.4 Further Invertibility and the Matrix-Map Dictionary . . . 63

**4 Row Reduction and Elementary Matrices 63**
4.1 Row Reduction and Elementary Matrices . . . 63

**5 Change of Basis 65**
5.1 Further Change of Basis and Coordinate Transformations . . . 65

**6 Matrix Equivalence and Similarity 65**
6.1 Matrix Equivalence and Rank Normal Forms . . . 65
6.2 Similarity and Operator Coordinates . . . 66

## IV Geometric and Spectral Linear Algebra 68

**1 Inner Product Geometry 68**
1.1 Inner Products and Norms . . . 68
1.2 Unit Vectors and Orthogonality . . . 72
1.3 Orthonormal Bases and Gram–Schmidt . . . 73
1.4 Orthonormal Coordinates and Inner-Product Identities . . . 75
1.5 Gram Matrices . . . 77
1.6 Finite-Dimensional Riesz Representation . . . 77

**2 Orthogonality, Projections, and Least Squares** **78**
2.1 Orthogonal Complements . . . 78
2.2 Orthogonal Projections . . . 80
2.3 Least Squares and Metric Optimization . . . 82
2.4 Adjoints and Fundamental Subspaces . . . 85
2.5 Self-Adjoint Operators and Orthogonal Projections . . . 88
2.6 Isometries and Unitary Matrices . . . 89
2.7 Householder Reflections . . . 91

**3 Bilinear and Quadratic Forms** **92**
3.1 Bilinear Form Types and Congruence Invariants . . . 92
3.2 Quadratic Forms and Polar Forms . . . 93
3.3 Orthogonality, Radicals, and Nonsingular Metric Spaces . . . 94
3.4 Radicals, Hyperbolic Spaces, and Witt Theory . . . 95
3.5 Symplectic Classification and Transvections . . . 99
3.6 Orthogonal Classification over Common Fields . . . 100
3.7 Quadratic Refinements over $\mathbb{F}_2$ . . . 102
3.8 Orthogonal Isometries and Symmetry Groups . . . 102

**4 Affine, Projective, and Convex Geometry** **104**
4.1 Affine Flats, Hulls, and Incidence . . . 104
4.2 Affine Independence and Barycentric Coordinates . . . 106
4.3 Affine Maps and Affine Groups . . . 107
4.4 Linear Projective Geometry and Affine Completion . . . 108
4.5 Convex Combinations, Cones, and Hulls . . . 109
4.6 Hyperplanes, Nearest Points, and Separation . . . 110
4.7 Positive Orthants, Alternatives, and Farkas's Lemma . . . 111

**5 Eigenstructure and Invariant Subspaces** **113**
5.1 Eigenvalues, Eigenspaces, and Spectral Criteria . . . 113
5.2 Invariant Subspaces, Reducing Pairs, and Quotient Operators . . . 114
5.3 Triangularization over Complex and Real Fields . . . 117
5.4 Diagonalization, Commutants, and Simultaneous Forms . . . 121

**6 Spectral Theorems and Operator Functions** **124**
6.1 Normal, Self-Adjoint, and Spectral Theorems . . . 124
6.2 Normal Operator Consequences . . . 128
6.3 Operator Functions . . . 132

## V Polynomial Structure and Canonical Forms 136

**1 Operator Polynomials** **136**
1.1 Operator Polynomials and Evaluation . . . 136
1.2 Polynomial Functional Calculus . . . 137

**2 Characteristic and Minimal Polynomials** **137**
2.1 Characteristic Polynomial Foundations . . . 137
2.2 Minimal Polynomial Foundations . . . 139

**3 Cayley-Hamilton and Annihilators** **139**
3.1 Cayley-Hamilton and Minimal Polynomials . . . 139
3.2 Diagonalizability and Minimal Polynomial Tests . . . 141
3.3 Annihilators and Spectral Control . . . 142
3.4 Polynomial Spectral Mapping . . . 143
3.5 Polynomial Invariance and Block Separation . . . 143

**4 Generalized Eigenspaces** **144**
4.1 Generalized Eigenspaces and Nilpotent Structure . . . 144

**5 Primary Decomposition** **146**
5.1 Primary Decomposition and Spectral Splitting . . . 146
5.2 Real Spectral Data via Complexification . . . 148
5.3 Normal Primary Structure . . . 149

**6 Jordan and Canonical Forms** **150**
6.1 Jordan Chains and Nilpotent Blocks . . . 150
6.2 Complex Jordan Form and Block Sizes . . . 151
6.3 Products AB and BA and Jordan Structure . . . 153
6.4 Nonderogatory Matrices and Commutants . . . 154
6.5 Jordan Operator Functions and Square Roots . . . 155

**7 Transpose and Conjugate Similarity** **158**
7.1 Transpose Similarity and Conjugate Similarity . . . 158

**VI Matrix Analysis and Numerical Linear Algebra** **161**

**1 Trace, Determinant, Rank, and Matrix Invariants** **161**
1.1 Trace and Determinant . . . 161
1.2 Minors, Cofactors, and Rank . . . 165
1.3 Commutators and Special Matrix Classes . . . 168

**2 Positive Definite and Structured Matrices** **170**
2.1 Positive Semidefinite and Positive Definite Matrices . . . 170
2.2 Principal Minors and Inertia . . . 173
2.3 Hermitian and Normal Matrix Structure . . . 175

**3 Matrix Products, Vectorization, and Spectral Radius** **176**
3.1 Kronecker Products and Vectorization . . . 176
3.2 Hadamard Products . . . 177
3.3 Spectral Radius and Matrix Powers . . . 178

**4 Matrix Decompositions** **180**
4.1 LU, QR, and Cholesky-Type Decompositions . . . 180
4.2 Polar and Singular Value Decompositions . . . 182

**5 Norms, Conditioning, Perturbation, and Stability** **187**
5.1 Matrix Norms . . . 187
5.2 Low-Rank Approximation and Pseudoinverses . . . 189
5.3 Conditioning . . . 192
5.4 Perturbation Bounds . . . 193
5.5 Diagonal Dominance and Definiteness . . . 194
5.6 Rayleigh Quotients and Minimax . . . 195
5.7 Eigenvalue and Singular-Value Interlacing . . . 196
5.8 Weyl Eigenvalue Bounds . . . 198
5.9 Brauer Rank-One Spectral Shifts . . . 198

**6 Numerical Linear Algebra** **199**
6.1 Direct Linear-System Solves . . . 199
6.2 QR Least Squares . . . 199
6.3 Banded and Sparse Direct Methods . . . 199
6.4 Krylov Subspace Methods . . . 200
6.5 Stationary Iterations and Splittings . . . 200
6.6 Hessenberg Reduction and QR Eigenvalue Methods . . . 201
6.7 Power and Shift-Invert Iterations . . . 202
6.8 Numerical SVD Methods . . . 203
6.9 Discrete Fourier Transforms and Circulant Matrices . . . 203
6.10 Fast Fourier Transform . . . 204

**VII Groups, Rings, and Fields** **205**

**1 Algebraic Language and Categories** **205**
1.1 Universal Properties and Equivalence-Based Structure . . . 205
1.2 Categories and Functors . . . 205
1.3 Algebraic Structures as Objects . . . 205
1.4 Exactness and Diagrams . . . 205

**2 Group Theory** **205**
2.1 Groups, Subgroups, and Cyclic Groups . . . 205
2.2 Group Homomorphisms, Kernels, and Images . . . 206
2.3 Cosets, Normality, and Structure Theorems . . . 206
2.4 Group Actions, Orbits, and Stabilizers . . . 206
2.5 Conjugacy Classes and the Class Equation . . . 206
2.6 Sylow Theory and Finite Groups . . . 207
2.7 Representation-Theoretic Preview . . . 207

**3 Rings, Ideals, and Polynomial Rings** **207**
3.1 Rings, Identities, Fields, and Examples . . . 207
3.2 Ring Homomorphisms and Subrings . . . 207
3.3 Ideals and Generated Ideals . . . 208
3.4 Quotient Rings and Maximal Ideals . . . 208
3.5 Integral Domains and Fields of Quotients . . . 209
3.6 Ring Characteristic . . . 209
3.7 Algebras over Fields . . . 210
3.8 Units, Divisibility, and Factorization Domains . . . 210
3.9 Polynomial Rings, Division, and GCDs . . . 210
3.10 Irreducible Polynomials and Unique Factorization . . . 211
3.11 Principal Ideal Domains and Unique Factorization . . . 212
3.12 Roots of Unity and Cyclotomic Polynomials . . . 213

**4 Fields and Galois Theory** **213**
4.1 Field Extensions . . . 213
4.2 Splitting Fields and Normality . . . 213
4.3 Galois Groups . . . 213
4.4 Finite Fields and Characteristic . . . 213
4.5 Classical Applications . . . 213

**VIII Modules, Multilinear Algebra, and Homological Methods** **214**

**1 Module Theory** **214**
1.1 Modules and Submodules . . . 214
1.2 Generated, Cyclic, and Finitely Generated Modules . . . 215
1.3 Linear Independence in Modules . . . 215
1.4 Module Homomorphisms, Kernels, and Images . . . 216
1.5 Quotient Modules, Correspondence, and Isomorphism . . . 216
1.6 Torsion and Annihilators . . . 217
1.7 Direct Sums and Complements . . . 218
1.8 Retracts, One-Sided Inverses, and Split Maps . . . 219

**2 Noetherian Modules and PID Structure Theorems** **220**
2.1 Free Modules, Bases, and Universal Property . . . 220
2.2 Rank and Coordinate Models of Free Modules . . . 221
2.3 Free Quotients and Splittings . . . 222
2.4 Noetherian Modules . . . 223
2.5 Orders and Cyclic Modules over a PID . . . 223
2.6 Submodules of Finite Free PID Modules . . . 224
2.7 Smith Normal Form and Module Presentations . . . 225
2.8 Finitely Generated PID Modules . . . 226
2.9 Primary Decomposition and Cyclic Primary Modules . . . 226
2.10 Elementary Divisors and Invariant Factors . . . 227

**3 Linear Operators as Modules** **229**
3.1 Polynomial Modules from Operators . . . 229
3.2 Invariant Factors . . . 231
3.3 Elementary Divisors . . . 232
3.4 Operator Classification . . . 233

**4 Associative Algebras** **236**
4.1 Associative Algebra Structures . . . 236
4.2 Modules over Algebras . . . 238
4.3 Zero Divisors and Spectra of Algebraic Elements . . . 239
4.4 Division Algebras . . . 240

**5 Tensor Products** **241**
5.1 Tensor Products and Universal Properties . . . 241
5.2 Tensor Product Bases and Maps . . . 242
5.3 Iterated Tensor Products and Tensor Spaces . . . 244

**6 Symmetric and Exterior Algebras** **246**
6.1 Symmetric Algebras . . . 246
6.2 Exterior Algebras . . . 247
6.3 Multilinear Algebra Revisited . . . 248

**7 Chain Complexes and Homology** **248**
7.1 Chain Complexes . . . 248
7.2 Homology . . . 248
7.3 Derived Functors Preview . . . 248

**IX Commutative Algebra** **249**

**1 Prime Ideals and Spectra** **249**
1.1 Prime Ideals . . . 249
1.2 Spectra . . . 249

**2 Localization and Local Rings** **249**
2.1 Localization . . . 249
2.2 Local Rings . . . 249

**3 Noetherian Rings and Modules** **249**
3.1 Noetherian Rings . . . 249
3.2 Noetherian Modules . . . 249

**4 Integral Dependence and Dimension** **249**
4.1 Integral Dependence . . . 249
4.2 Dimension Theory . . . 249

**5 Regularity and Singularities** **249**
5.1 Regularity . . . 249
5.2 Singularities Preview . . . 249

**X Algebraic Geometry** **250**

**1 Affine Algebraic Sets** **250**
1.1 Affine Algebraic Sets . . . 250
1.2 Ideals and Coordinate Rings . . . 250

**2 Morphisms and Coordinate Rings** **250**
2.1 Regular Maps . . . 250
2.2 Coordinate Ring Functoriality . . . 250

**3 Projective Geometry** **250**
3.1 Projective Space . . . 250
3.2 Projective Varieties . . . 250

**4 Schemes** **250**
4.1 Affine Schemes . . . 250
4.2 Gluing and Scheme Morphisms . . . 250

**5 Sheaves and Cohomology** **250**
5.1 Sheaves . . . 250
5.2 Cohomology Preview . . . 250

**6 Smoothness, Tangent Spaces, and Dimension** **250**
6.1 Smoothness . . . 250
6.2 Tangent Spaces . . . 250
6.3 Dimension . . . 250

**7 Divisors, Line Bundles, and Curves** **250**
7.1 Divisors . . . 250
7.2 Line Bundles . . . 250
7.3 Curves . . . 250

## XI Real and Topological Analysis 251

**1 Number Systems and Completeness** **251**
1.1 Ordered Fields, Suprema, and Real Structure . . . 251
1.2 Complex Numbers and Absolute Value . . . 256

**2 Real Sequences and Completeness Methods** **257**
2.1 Sequences, Cauchy Criteria, and Completeness . . . 257
2.2 Divergence, Upper and Lower Limits, and Sequence Estimates . . . 260

**3 Metric Spaces and Euclidean Topology** **263**
3.1 Metrics, Open Sets, and Euclidean Topology . . . 263
3.2 Relative Topology . . . 268
3.3 Density, Separability, and Closure Tests . . . 268
3.4 Bounded Sets and Metric Sequences . . . 269

**4 Compactness and Completeness** **272**
4.1 Compactness, Completeness, and Total Boundedness . . . 272
4.2 Heine–Borel and Euclidean Compactness . . . 278
4.3 Baire Category . . . 280
4.4 Perfect Sets and the Cantor Set . . . 283
4.5 Complete Metric Spaces and Completions . . . 284

**5 Connectedness and Continuity** **286**
5.1 Connectedness and Continuity . . . 286
5.2 Uniform Continuity and Compactness . . . 293
5.3 One-Sided Limits, Monotone Functions, and Extended Limits . . . 295

**6 General Topology for Analysis** **297**
6.1 Topological Spaces, Bases, and Separation . . . 297
6.2 Continuity and Topological Constructions . . . 299
6.3 Topological Compactness and Normality Tools . . . 301
6.4 Sequences, Countability, and Metrization . . . 304
6.5 Topological Connectedness and Paths . . . 308

**7 Finite-Dimensional Topological Vector Spaces** **310**
7.1 Natural Topology and Continuity . . . 310

## XII Calculus, Series, and Approximation 312

**1 One-Variable Differentiation and Riemann Integration** **312**
1.1 Differentiation and Basic Theorems . . . 312
1.2 Riemann Integration . . . 317

**2 Riemann-Stieltjes and Improper Integration** **318**
2.1 Riemann-Stieltjes Integrability . . . 318
2.2 Fundamental Theorems of Calculus . . . 321
2.3 Step Integrators and Change of Variable . . . 321
2.4 Improper Integrals . . . 322
2.5 Vector-Valued Integrals and Curves . . . 323
2.6 Further Topics in Integration . . . 324

**3 Several-Variable Differentiation** **324**
3.1 Partial and Total Derivatives . . . 324
3.2 Differentiability and Linear Approximation . . . 326

**4 Inverse, Implicit, and Rank Theorems** **328**
4.1 Inverse Function Theorem . . . 328
4.2 Implicit Function Theorem . . . 330
4.3 Rank Theorem . . . 331
4.4 Further Topics in Local Differential Theory . . . 331

**5 Differential Forms and Stokes Theory** **332**
5.1 Multiple Integration and Change of Variables . . . 332
5.2 Differential Forms and Manifolds . . . 335
5.3 Stokes Theory . . . 337
5.4 Further Topics in Differential Forms and Stokes Theory . . . 338

**6 Infinite Series and Summability** **339**
6.1 Series Tests . . . 339

**7 Uniform Convergence** **345**
7.1 Uniform Limits . . . 345
7.2 Interchange of Limits . . . 348
7.3 Further Topics in Uniform Convergence . . . 348

**8 Function-Space Compactness** **349**
8.1 Equicontinuity . . . 349
8.2 Arzela-Ascoli-Type Compactness . . . 350
8.3 Further Topics in Function-Space Compactness . . . 351

**9 Power Series and Approximation** **352**
9.1 Power Series . . . 352
9.2 Approximation Methods . . . 356
9.3 The Umbral Calculus . . . 359
9.4 Umbral Recurrences and Variants . . . 366

**10 Fourier Structure** **367**
10.1 Fourier Orthogonality . . . 367
10.2 Fourier Approximation . . . 368
10.3 Further Topics in Fourier Analysis . . . 370

**XIII Measure Foundations and Extension** **371**

**1 Sigma-Algebras and Measurable Spaces** **371**
1.1 Sigma-Algebras and Measurable Structures . . . 371

**2 Measures, Charges, and Basic Examples** **372**
2.1 Measures, Charges, and Basic Constructions . . . 372

**3 Outer Measures and Caratheodory Extension** **374**
3.1 Outer Measures and Extension Theorems . . . 374
3.2 Euclidean Cells and Cell Outer Measure . . . 379
3.3 Caratheodory Measurability and Lebesgue Measure . . . 381
3.4 Caratheodory-Hahn Extension . . . 383
3.5 Metric Outer Measures and Hausdorff Measure . . . 386

**4 Borel, Lebesgue, and Stieltjes Measures** **388**
4.1 Borel, Lebesgue, and Stieltjes Measure Constructions . . . 388

**5 Continuity and Regularity of Measures** **391**
5.1 Measure Continuity and Regularity . . . 391
5.2 Radon Measures on Locally Compact Spaces . . . 394
5.3 Invariant Measures on Compact Groups . . . 397

**6 Regular Approximation and Caratheodory Tests** **399**
6.1 Borel Approximation and Regular Measurable Sets . . . 399
6.2 Compact and Cell Approximation . . . 402
6.3 Caratheodory Tests and Inner Measure . . . 403

**7 Vitali, Steinhaus, and Nonmeasurable Sets** **404**
7.1 Vitali, Steinhaus, and Nonmeasurability . . . 404

**8 Cantor–Lebesgue Singular Function** **406**
8.1 Cantor-Lebesgue Singular Function Construction . . . 406

**XIV Measure, Integration, and Functional Analysis** **408**

**1 Measurable Functions and Lebesgue Integration** **408**
1.1 Measurable Functions . . . 408
1.2 Lebesgue Integration . . . 412
1.3 Convergence Theorems . . . 418
1.4 Parameter Integrals and Differentiation . . . 422
1.5 Riemann-Lebesgue Comparison . . . 424
1.6 Littlewood and Lusin Approximation . . . 425
1.7 Convex Functions and Jensen's Inequality . . . 427

**2 Product Measures and Fubini Theory** **428**
2.1 Product Measures . . . 428
2.2 Fubini and Tonelli Theorems . . . 430
2.3 Linear Changes of Variables and Euclidean Invariance . . . 434

**3 Radon-Nikodym and Differentiation** **436**
3.1 Signed and Absolutely Continuous Measures . . . 436
3.2 Radon-Nikodym Theorem . . . 439
3.3 Differentiation of Measures . . . 442
3.4 Monotone and Bounded-Variation Differentiation . . . 444
3.5 Absolute Continuity and BV Decomposition . . . 447
3.6 Nikodym Metric and Setwise Measure Limits . . . 451

**4 $L^p$ Spaces and Modes of Convergence** **453**
4.1 $L^p$ Spaces . . . 453
4.2 Modes of Convergence . . . 462
4.3 Composition and Convergence Modes . . . 468
4.4 Uniform Integrability and Vitali Convergence . . . 470
4.5 Approximation and Separability in $L^p$ . . . 472
4.6 Weak Convergence in $L^p$ . . . 473

**5 Banach and Hilbert Spaces** **478**
5.1 Normed and Banach Spaces . . . 478
5.2 Hilbert Spaces . . . 482
5.3 Core Functional Analytic Theorems . . . 491
5.4 Open Mapping, Closed Graph, and Complemented Subspaces . . . 493

**6 Weak Topologies, Compact Operators, and Spectral Theory** **495**
6.1 Weak Topologies and Duality . . . 495
6.2 Compact Operators . . . 500
6.3 Spectral Theory of Operators . . . 504
6.4 Measure Duality and Riesz Representation . . . 507

**XV Probability Foundations and Distributions** **509**

**1 Probability Spaces and Random Variables** **509**
1.1 Probability Spaces and Event Algebra . . . 509
1.2 Finite Probability, Counting, and Independence . . . 511
1.3 Random Variables, Laws, and Distribution Functions . . . 513
1.4 Standard Distribution Families and Approximations . . . 515

**2 Joint Distributions, Multivariate Laws, and Transformations** **517**
2.1 Joint Distributions and Independence . . . 517
2.2 Multivariate Models and Copulas . . . 518
2.3 Transformations of Random Variables . . . 520

**3 Expectation and Integration in Probability** **521**
3.1 Expectation, Variance, and LOTUS . . . 521
3.2 Moments, Sample Statistics, and Random Vectors . . . 522
3.3 Probability Inequalities . . . 524
3.4 Moment Generating Functions . . . 525

**4 Order Statistics, Extrema, and Moment Methods** **526**
4.1 Order Statistics, Extrema, and Moment Methods . . . 526

**5 Normal-Derived Distributions** **528**
5.1 Normal-Derived Distribution Families . . . 528

**XVI Probability and Stochastic Systems** **531**

**1 Conditional Probability and Conditional Expectation** **531**
1.1 Conditional Probability . . . 531
1.2 Conditional Expectation . . . 534
1.3 Conditional Moment Methods and Prediction . . . 536

**2 Probabilistic Convergence** **537**
2.1 Modes of Probabilistic Convergence . . . 537
2.2 Probability on Metric Spaces . . . 539

**3 Limit Theorems** **539**
3.1 Laws of Large Numbers . . . 539
3.2 Central Limit Theory . . . 540

**4 Stochastic Processes** **543**
4.1 Process-Level Probability . . . 543
4.2 Filtrations and Adapted Processes . . . 543
4.3 Martingales in Discrete and Continuous Time . . . 543
4.4 Markov Chains and Markov Processes . . . 543
4.5 Stationary Processes and Ergodicity . . . 547
4.6 Gaussian Processes . . . 547
4.7 Point Processes and Renewal Structure . . . 547

**5 Stochastic Calculus and SDEs** **548**
5.1 Brownian Motion . . . 548
5.2 Stochastic Integration . . . 548
5.3 Itô Calculus . . . 548
5.4 Stochastic Differential Equations . . . 548
5.5 Change of Measure and Girsanov Theory . . . 548
5.6 Diffusions and PDE Connections . . . 548
5.7 Stochastic Flows and Advanced Dynamics . . . 548

**XVII Dynamical and PDE Systems** **549**

**1 Dynamical Systems and ODEs** **549**
1.1 ODE Foundations and Flows . . . 549
1.2 Linear Systems and Stability . . . 559
1.3 Nonlinear Stability Theory . . . 565
1.4 Bifurcation Theory . . . 567
1.5 Dynamical Systems Structure . . . 570
1.6 Semigroups and Evolution Viewpoint . . . 572
1.7 Measure-Preserving Dynamics and Ergodicity . . . 572

**2 Partial Differential Equations** **575**
2.1 First-Order PDE and Characteristics . . . 575
2.2 Elliptic Equations . . . 575
2.3 Parabolic Equations . . . 575
2.4 Hyperbolic Equations . . . 575

**3 Weak Solutions and Evolution Equations** **575**
3.1 Sobolev Spaces for PDE . . . 575
3.2 Weak Solutions and Variational Methods . . . 575
3.3 Evolution Equations . . . 575

**XVIII Numerical and Stochastic Computation** **576**

**1 Numerical Analysis** **576**
1.1 Floating Point and Error Analysis . . . 576
1.2 Nonlinear Equations and Optimization Basics . . . 576
1.3 Interpolation and Approximation . . . 576
1.4 Numerical Differentiation and Quadrature . . . 576
1.5 Numerical Stability and Convergence . . . 576

**2 Numerical Differential Systems** **576**
2.1 Numerical ODEs . . . 576
2.2 Geometric Numerical Integration . . . 578
2.3 Finite Difference PDE Methods . . . 578
2.4 Finite Element Methods . . . 578
2.5 Numerical SDEs . . . 578
2.6 Computational Stability Across ODE/PDE/SDE . . . 578

**3 Monte Carlo and Stochastic Computation** **578**
3.1 Monte Carlo Foundations . . . 578
3.2 Variance Reduction . . . 579
3.3 Markov Chain Monte Carlo . . . 579
3.4 Sequential Monte Carlo . . . 580
3.5 Rare-Event Simulation . . . 580
3.6 Stochastic Optimization . . . 580

**XIX Statistical Inference and Time Series** **582**

**1 Statistical Models and Experiments** **582**
1.1 Statistical Experiments and Likelihoods . . . 582
1.2 Sufficiency, Exponential Families, and Local Asymptotics . . . 583
1.3 Design, Randomization, and Observational Comparisons . . . 584

**2 Estimation Theory** **585**
2.1 Estimators, Bias, Variance, and Risk . . . 585
2.2 Information Bounds and Asymptotic Efficiency . . . 585
2.3 Method of Moments and Parametric Likelihood Examples . . . 586

**3 Hypothesis Testing and Confidence** **587**
3.1 Tests, Likelihood Ratios, and Error Control . . . 587
3.2 Confidence Sets, Pivots, and Asymptotic Intervals . . . 590
3.3 Goodness-of-Fit and Diagnostic Plots . . . 591
3.4 Normal Sampling Pivots . . . 593
3.5 Two-Sample Normal Inference . . . 593
3.6 Paired-Sample Inference . . . 595

**4 Decision Theory** **596**
4.1 Loss, Risk, Bayes, and Minimax Rules . . . 596
4.2 Shrinkage, Empirical Bayes, and Regularization . . . 597

**5 Bayesian Inference** **597**
5.1 Priors, Posteriors, and Conjugacy . . . 597
5.2 Computation, Prediction, and Asymptotic Bayes . . . 598
5.3 Bayesian Normal Comparisons . . . 599

**6 Nonparametric and High-Dimensional Statistics** **600**
6.1 Empirical Processes, Density, and Regression Smoothing . . . 600
6.2 Sparse High-Dimensional Inference . . . 604
6.3 Rank-Based Comparative Tests . . . 605

**7 Time Series Models** **606**
7.1 Stationarity, Autocovariance, and Linear Processes . . . 606
7.2 ARIMA, State-Space, and Volatility Models . . . 607

**8 Sequential Inference and Resampling** **608**
8.1 Resampling, Bootstrap, and Permutation Methods . . . 608
8.2 Sequential Experiments, Filtering, and Online Inference . . . 609
8.3 Two-Sample Resampling Methods . . . 610

## XX Statistical Data Analysis and Applied Models 611

**1 Survey Sampling** **611**
1.1 Finite Population Parameters and Simple Random Sampling . . . 611
1.2 Normal Approximation and Confidence Intervals . . . 613
1.3 Ratio Estimation . . . 614
1.4 Stratified Random Sampling . . . 616

**2 Exploratory Data Analysis and Data Summaries** **618**
2.1 Empirical Distribution and Survival Summaries . . . 618
2.2 Quantile and Shape Displays . . . 618
2.3 Location, Scale, and Robust Summaries . . . 620
2.4 Boxplots, Scatterplots, and Relationship Summaries . . . 620

**3 Categorical Data Analysis** **622**
3.1 Fisher's Exact Test and Fixed Margins . . . 622
3.2 Chi-Square Tests for Contingency Tables . . . 623
3.3 Matched Pairs and McNemar's Test . . . 624
3.4 Odds Ratios and Case-Control Sampling . . . 625
3.5 Higher-Order Tables and Sampling Design . . . 626

**4 Analysis of Variance and Designed Experiments** **627**
4.1 One-Way Layouts and the $F$ Test . . . 627
4.2 Multiple Comparisons After ANOVA . . . 629
4.3 Rank-Based One-Way ANOVA . . . 630
4.4 Two-Way Layouts and Interaction . . . 631
4.5 Randomised Blocks and Friedman's Test . . . 633

**5 Linear Models and Regression** **634**
5.1 Least Squares Curve Fitting and Linearization . . . 634
5.2 Simple Linear Regression Inference and Diagnostics . . . 635
5.3 Matrix Linear Models and Normal Equations . . . 637
5.4 Inference, Prediction, and Multiple Regression . . . 638
5.5 Random Design, Bootstrap, and Extensions . . . 640

## XXI Statistical Learning and Deep Learning 641

**1 Learning Problems and Risk** **641**
1.1 Prediction, Loss Functions, and Risk . . . 641
1.2 Empirical Risk Minimization . . . 642
1.3 Surrogate Losses and Calibration . . . 642
1.4 Bias-Variance and Approximation-Estimation Tradeoffs . . . 643

**2 Generalization Theory** **643**
2.1 Uniform Convergence and VC Theory . . . 643
2.2 Rademacher Complexity and Covering Numbers . . . 644
2.3 Concentration and Stability . . . 645
2.4 PAC-Bayes and Information-Theoretic Bounds . . . 646

**3 Unsupervised and Latent-Variable Learning** **646**
3.1 Clustering and Mixture Models . . . 646
3.2 Dimension Reduction and Manifold Learning . . . 650
3.3 Matrix Factorization and Topic Models . . . 652
3.4 Representation Learning Objectives . . . 653
3.5 Association Rules and Density-Ratio Views . . . 654
3.6 Graph Ranking and PageRank . . . 655
3.7 Latent Factor and Component Analysis . . . 656

**4 Probabilistic Graphical and Causal Learning** **657**
4.1 Directed and Undirected Graphical Models . . . 657
4.2 Inference and Learning in Graphical Models . . . 659
4.3 Causal Graphs, Interventions, and Identifiability . . . 660
4.4 Counterfactuals and Policy Evaluation . . . 660

**5 Neural Network Foundations** **661**
5.1 Network Architectures and Expressivity . . . 661
5.2 Backpropagation and Automatic Differentiation . . . 662
5.3 Optimization for Deep Learning . . . 663
5.4 Regularization, Normalization, and Generalization . . . 663

**6 Modern Deep Models** **664**
6.1 Convolutional and Geometric Architectures . . . 664
6.2 Sequence Models and Attention . . . 665
6.3 Transformers and Large-Scale Training . . . 665
6.4 Generative Models . . . 666
6.5 Reinforcement Learning and Decision Systems . . . 666

**7 Learning Systems and Evaluation** **667**
7.1 Data Splitting, Validation, and Benchmarking . . . 667
7.2 Robustness, Shift, and Distributional Uncertainty . . . 668
7.3 Interpretability, Fairness, and Safety . . . 669
7.4 Deployment, Monitoring, and Feedback Loops . . . 669

**XXII Classical Supervised Learning** **671**

**1 Linear Prediction and Generalized Linear Models** **671**
1.1 Linear Predictors and Least Squares . . . 671
1.2 Linear Classification and Generalized Linear Models . . . 671
1.3 Discriminant Analysis and Naive Bayes . . . 672

**2 Regularisation and Model Selection** **673**
2.1 Penalized Linear Estimation . . . 673
2.2 Validation and Selection Bias . . . 674
2.3 Derived Input Directions and Regularization Paths . . . 674
2.4 Effective Degrees and Information Criteria . . . 676

**3 Basis Expansions, Splines, and Wavelets** **676**
3.1 Basis Expansions, Splines, and Wavelet Features . . . 676

**4 Kernel Methods and RKHS Theory** **677**
4.1 Kernel Methods and RKHS Representations . . . 677

**5 Support Vector Machines and Margin Methods** **679**
5.1 Margins and SVM Primal Problems . . . 679
5.2 Dual Problems and Kernelization . . . 680
5.3 Support Vector Regression and Perceptrons . . . 681

**6 Trees and Ensemble Averaging** **682**
6.1 Tree Models and Bagging . . . 682
6.2 CART Splits and Pruning . . . 682
6.3 Ensembles, Random Forests, and Stacking . . . 683

**7 Additive Models and Adaptive Basis Methods** **684**
7.1 Generalized Additive Models . . . 685
7.2 Adaptive Basis and Rule Methods . . . 685

**8 Boosting and Additive Trees** **686**
8.1 Stagewise Boosting . . . 686
8.2 AdaBoost and Exponential Loss . . . 687
8.3 Gradient Boosted Trees and Rule Ensembles . . . 688

**9 High-Dimensional Supervised Learning** **690**
9.1 High-Dimensional Linear Rules . . . 690
9.2 Sparse and Structured Penalties . . . 691
9.3 Latent Factors and Distance-Based Classifiers . . . 692

**10 Prototype and Nearest-Neighbor Methods** **692**
10.1 Prototype Classifiers . . . 693
10.2 Nearest Neighbors . . . 693
10.3 Invariant Distances and Adaptive Neighborhoods . . . 694

**11 Flexible and Mixture Discriminant Analysis** **695**
11.1 Flexible and Mixture Discriminant Rules . . . 695

# Notation List

## Collision Index and Scope Rules

| | |
|---|---|
| bare $A, B, \ldots$; $\mathcal{A}, \mathcal{B}, \ldots$ | unadorned Latin and calligraphic letters denote local objects unless a decorated form is listed below; local declarations in a theorem, proof, or subsection take precedence |
| $B,\ B_r(x),\ \overline{B}_r(x),\ \mathrm{B}(x,y),\ \mathcal{B}(\mathbb{R}^n)$ | matrix or set variable / open ball / closed ball / beta function / Borel $\sigma$-algebra; font, arguments, and ambient subject determine the referent |
| $I_{\mathcal{V}},\ I_n,\ I^k,\ I_R$ | identity operator, identity matrix, $k$-cell, and power-series interval |
| $P,\ \mathcal{P}(X),\ \mathscr{P},\ P_{\mathcal{U}},\ P_j$ | locally declared projection or map / power set / umbral polynomial algebra / orthogonal projection / spectral projection |
| $\mathcal{F},\ \mathscr{F},\ \mathcal{F}(K),\ \mathcal{F}(\mathcal{V})$ | local family of functions or operators / umbral formal-series algebra / finite subsets of $K$ / finite-rank operators |
| $\mathcal{L}(\mathcal{V},\mathcal{W}),\ \mathcal{L}^n,\ \mathscr{L}^1,\ \mathrm{LS}(A,\mathbf{y}),\ \mathcal{L}(\theta,\alpha)$ | linear maps / Lebesgue measurable sets / integrable functions / least-squares solution set / local Lagrangian or loss notation |
| $R(f,\Pi,\{c_i\}),\ R,\ R_k(A),\ A=QR$ | Riemann sum, locally declared radius or scalar, row-sum quantity, and QR factorization; arity and argument type disambiguate the notation |
| $\Delta x_i,\ \Delta A,\ \Delta(A),\ \Delta(\mathcal{V}),\ \Delta_{k-1}$ | increment, matrix perturbation, defect from normality, metric discriminant, and coefficient simplex |
| $\Omega,\ \Omega_T,\ \Omega_n,\ \mathrm{vol}(\Omega)$ | local domain or set / finite spectrum of $T$ / primitive roots of unity or DFT diagonal, as specified by context / volume domain |
| $Q^k,\ Q_n,\ \Phi_n$ | standard $k$-simplex / normalized Fourier matrix / cyclotomic polynomial; cyclotomic polynomials use $\Phi_n$ |
| $\overline{z},\ \overline{E},\ \overline{\mathcal{A}},\ A^*,\ T^*\omega,\ f',\ \mathcal{V}'$ | conjugation, closure, uniform closure, adjoint or conjugate transpose, pullback, derivative, and dual space; the decorated object determines the operation |

## Global Conventions and Basic Notation

*Convention.* Notation is scoped by subject, argument type, font, decoration, and any local definition in force. Unadorned letters denote local objects by default; persistent meanings are carried by subscripts, superscripts, operators, fonts, or declared argument patterns. When a glyph is reused, the distinguishing feature is recorded below.

*Set, Function, and Logical Notation*

| | |
|---|---|
| $x \in E,\ x \notin E$ | membership / non-membership |
| $E \subseteq F,\ \emptyset$ | subset inclusion, not necessarily proper / empty set |
| $X \times Y,\ f : X \to Y$ | Cartesian product / function with domain $X$ and codomain $Y$ |
| $\Longrightarrow,\ \Longleftrightarrow$ | logical implication / logical equivalence |
| $:=$ | definition by equality |

*Number, Size, and Scalar Notation*

| | |
|---|---|
| $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ | natural numbers, integers, rationals, reals, and complexes |
| $\mathbb{F},\ \mathbb{F}_0$ | ambient scalar field unless fixed as $\mathbb{R}$ or $\mathbb{C}$ / prime subfield of $\mathbb{F}$; probabilities use explicit Pr or locally declared probability spaces |
| $\lvert x\rvert,\ \lvert E\rvert,\ \lVert x\rVert$ | absolute value or complex modulus for scalars; interval length or cardinality according to argument type; norm of a vector, function, or operator |
| $a, b, c$ | locally declared scalars or interval endpoints |

## Linear and Operator Structures Notation

*Vector Spaces and Lists*

| | |
|---|---|
| $\mathbb{F},\ \mathbb{F}_0$ | ambient scalar field, typically $\mathbb{R}$ or $\mathbb{C}$ / prime subfield of $\mathbb{F}$ |
| $V(n,q),\ \mathbb{F}_q^n$ | $n$-dimensional coordinate space over the finite field with $q$ elements |
| $\mathbb{F}^{(\mathbb{N})}$ | vector space of finitely supported sequences over $\mathbb{F}$ |
| $\mathbf{u},\mathbf{v},\mathbf{w},\mathbf{0}$ | vectors / zero vector |
| $\mathcal{U},\mathcal{V},\mathcal{W}$ | vector spaces or subspaces; other calligraphic letters such as $\mathcal{A},\mathcal{B},\mathcal{F},\mathcal{M}$ are local unless listed below; umbral algebras use script $\mathscr{P},\mathscr{F}$ |
| $I_{\mathcal{V}},\ I_n,\ 0$ | identity operator on $\mathcal{V}$ / identity matrix / zero map or zero matrix; distinct from the cell $I^k$ and interval $I_R$ |
| $\beta=(\mathbf{v}_1,\dots,\mathbf{v}_m)$ | ordered vector list; a basis iff it is linearly independent and spans |
| $(),\ \beta_{\widehat{j}}$ | empty list / list $\beta$ with the $j$th entry omitted |
| $(\beta,\mathbf{v}),\ \beta_{j\leftarrow\mathbf{v}}$ | list with $\mathbf{v}$ appended / list with the $j$th entry replaced by $\mathbf{v}$ |
| $\mathcal{B}$ | locally declared basis set, such as a Hamel, Hilbert, module, or affine basis; $\mathcal{B}(\mathbb{R}^n)$ denotes the Borel $\sigma$-algebra in analysis |
| $\operatorname{span}(E),\ \dim(\mathcal{V})$ | linear span of $E$ / dimension of $\mathcal{V}$ |
| $\operatorname{codim}_{\mathcal{V}}(\mathcal{U})$ | codimension of $\mathcal{U}$ in $\mathcal{V}$; equal to $\dim(\mathcal{V}/\mathcal{U})$ in finite dimension |
| $\bigwedge_{i\in I}\mathcal{U}_i,\ \bigvee_{i\in I}\mathcal{U}_i$ | meet / join of a family of subspaces |
| $\sum_{i\in I}\mathcal{U}_i$ | sum of an indexed family of subspaces; equivalently the join |
| $\mathcal{U}+\mathcal{W},\ \mathcal{U}\oplus\mathcal{W},\ \mathcal{V}/\mathcal{U}$ | sum, direct sum, quotient space |
| $\mathcal{V}_1\times\cdots\times\mathcal{V}_m$ | product vector space with componentwise operations |
| $\prod_{i\in I}\mathcal{V}_i,\ \bigoplus_{i\in I}^{\text{ext}}\mathcal{V}_i$ | external direct product / external direct sum of finite-support families |
| $\mathcal{V}\cong\mathcal{W},\ \mathbf{e}_{j,k}$ | isomorphic vector spaces / basis vector supported in one product component |
| $\mathbf{v}+\mathcal{U},\ \pi(\mathbf{v})$ | coset modulo $\mathcal{U}$ / locally declared quotient map $\mathcal{V}\to\mathcal{V}/\mathcal{U}$ |

*Linear Maps and Duality*

| | |
|---|---|
| $T:\mathcal{V}\to\mathcal{W},\ \mathcal{L}(\mathcal{V},\mathcal{W})$ | linear map / vector space of linear maps; distinct from measure-theoretic $\mathcal{L}^n$, least-squares $\operatorname{LS}(A,\mathbf{y})$, and local loss notation $\mathcal{L}(\theta)$ |
| $ST,\ T\vert_{\mathcal{U}},\ T/\mathcal{U}$ | composition $S\circ T$ / restriction to $\mathcal{U}$ / induced quotient operator |
| $T^{-1},\ T^m,\ p(T)$ | inverse of an invertible map or operator / operator power / polynomial functional calculus |
| $\ker(T),\ \operatorname{range}(T)$ | kernel and range of $T$ |
| $\mathcal{V}_{\mathbb{C}},\ \operatorname{cpx},\ T_{\mathbb{C}}$ | complexification of a real vector space / canonical embedding $\mathbf{v}\mapsto\mathbf{v}+i\mathbf{0}$ / complexification of a real operator |
| $\operatorname{graph}(T),\ [S,T]$ | graph of a map / commutator $ST-TS$ |
| $\operatorname{Comm}(\mathcal{F}),\ \operatorname{Comm}(A)$ | commutant of a family / commutant of a single matrix or operator |
| $A\sim_{\text{row}}B,\ S\sim T$ | row-equivalent matrices / operator similarity when declared; distinct from cardinal, a.e., and coefficient equivalence |
| $\mathcal{V}',\ \beta',\ T'$ | dual space, dual basis, and dual map; prime denotes duality in this context |
| $\varphi,\psi,\ \mathcal{U}^0$ | linear functionals / annihilator |
| $\mathcal{V}'',\ \Lambda_{\mathcal{V}}$ | double dual / natural evaluation map $\mathcal{V}\to\mathcal{V}''$ |
| $[\mathbf{v}]_\beta,\ [T]_{\gamma\leftarrow\beta}$ | coordinate vector in basis $\beta$ / matrix of $T$ from $\beta$-coordinates to $\gamma$-coordinates |
| $C_{\gamma\leftarrow\beta}$ | change-of-basis matrix sending $\beta$-coordinate columns to $\gamma$-coordinate columns |

*Matrices and Entrywise Structure*

| | |
|---|---|
| $A, B, C,\ \mathrm{Mat}(m, n, \mathbb{F})$ | matrix variables / space of $m \times n$ matrices over $\mathbb{F}$; decorated forms such as $C_f$ and $C_{ij}$ have separate meanings |
| $A_{ij},\ A_{j,\cdot},\ A_{\cdot,k}$ | entry in row $i$ and column $j$ / $j$th row / $k$th column |
| $[a_{ij}],\ E_{j,k},\ A\mathbf{x}$ | matrix with entries $a_{ij}$ / matrix unit / matrix-vector product |
| $A^{-1},\ A^t,\ A^T,\ A^*$ | inverse matrix when it exists / transpose / alternate transpose notation / conjugate transpose or adjoint |
| $\mathrm{diag}(\lambda_1, \ldots, \lambda_n)$ | diagonal matrix with diagonal entries $\lambda_1, \ldots, \lambda_n$ |
| $\overline{z},\ \mathrm{Re}\, z,\ \mathrm{Im}\, z$ | complex conjugate / real part / imaginary part; for sets, overline denotes closure |
| $\mathrm{tr}(A), \det(A)$ | trace and determinant |
| $\mathfrak{S}_n,\ \mathrm{sign}(\sigma),\ \mathrm{adj}(A)$ | permutation group / permutation sign / adjugate matrix |
| $M_{ij},\ C_{ij},\ A_j(\mathbf{b})$ | minor matrix / cofactor / matrix obtained from $A$ by replacing column $j$ with $\mathbf{b}$ |
| $\mathrm{col}(A),\ \mathrm{row}(A)$ | column space and row space of $A$ |
| $\mathrm{rank}(A),\ \mathrm{nullity}(A)$ | rank and nullity of $A$ |
| $A \oplus B,\ A \otimes B$ | matrix direct sum / Kronecker product; vector-space direct sums use subspace notation |
| $A \otimes I_q + I_n \otimes B$ | Kronecker sum of square matrices $A$ and $B$ in compatible dimensions |
| $A \circ B,\ \mathrm{vec}(A)$ | Hadamard product / column-stacking vectorisation |
| $M/A,\ M/D$ | Schur complement of the block $A$ or $D$ in a compatible block matrix $M$; distinct from quotient-space notation |
| $B,\ L = B^t B$ | directed incidence matrix / associated incidence Laplacian in a graph context |

*Inner Product Geometry*

| | |
|---|---|
| $\|\mathbf{v}\|,\ \|T\|_{\mathrm{op}}$ | vector norm / operator norm |
| $\|\cdot\|_1,\ \|\cdot\|_2,\ \|\cdot\|_\infty$ | coordinate one-norm / Euclidean or spectral two-norm according to argument type / sup norm |
| $\|\cdot\|_F$ | Frobenius norm |
| $\langle \mathbf{u}, \mathbf{v} \rangle,\ \langle A, B \rangle_F$ | inner product / Frobenius inner product |
| $\mathbf{u} \perp \mathbf{v},\ \mathcal{U}^\perp,\ \delta_{ij}$ | orthogonal vectors, orthogonal complement, Kronecker delta |
| $\mathrm{proj}_{\mathbf{v}}(\mathbf{u}),\ P_{\mathcal{U}}$ | orthogonal projection of $\mathbf{u}$ onto $\mathrm{span}(\mathbf{v})$ / orthogonal projection onto $\mathcal{U}$ |
| $\mathrm{LS}(A, \mathbf{y})$ | least-squares solution set $\arg\min_{\mathbf{x} \in \mathbb{F}^n} \|A\mathbf{x} - \mathbf{y}\|_2$ for $A \in \mathrm{Mat}(m, n, \mathbb{F})$; roman notation avoids collision with $\mathcal{L}$ and $\mathcal{L}^n$ |
| $d(\mathbf{v}, \mathcal{U})$ | distance $\inf_{\mathbf{u} \in \mathcal{U}} \|\mathbf{v} - \mathbf{u}\|$ from $\mathbf{v}$ to $\mathcal{U}$ |
| $\mathrm{Gram}(\mathbf{u}_1, \ldots, \mathbf{u}_n)$ | Gram matrix of $\mathbf{u}_1, \ldots, \mathbf{u}_n$ |
| $\mathrm{GramDet}(\mathbf{u}_1, \ldots, \mathbf{u}_n)$ | Gram determinant |
| $P_{\mathbf{u}},\ U_{\mathbf{w}}$ | rank-one projection matrix / Householder matrix, with defining vector indicated by the subscript |
| $\gamma = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$ | ordered basis; orthonormal when declared in an inner-product context |
| $\mathbf{e} = (1, \ldots, 1)^t$ | all-ones vector |
| $\mathrm{Ray}_A(\mathbf{x}),\ \mathbf{x}^* A \mathbf{x}$ | Rayleigh quotient notation; for unit $\mathbf{x}$, $\mathrm{Ray}_A(\mathbf{x}) = \mathbf{x}^* A \mathbf{x}$ / Hermitian quadratic form |
| $\mathrm{Sym}(B),\ \mathrm{Skew}(B),\ \mathrm{Alt}(B)$ | predicates that the bilinear form $B$ is symmetric, skew-symmetric, or alternating |
| $M_\beta,\ \Delta(\mathcal{V}),\ \mathrm{rad}(\mathcal{V})$ | matrix of a metric bilinear form in basis $\beta$ / discriminant set or square class / radical of a metric vector space |
| $\mathcal{V} = \mathcal{S} \odot \mathcal{T},\ \mathcal{H},\ w(\mathcal{V})$ | orthogonal direct sum / hyperbolic space or plane in metric-vector-space contexts / Witt index; affine hyperplanes use $\mathcal{H}(\mathbf{N}, b)$ |
| $\mathcal{O}(\mathcal{V}),\ \mathrm{Sp}(\mathcal{V}),\ \tau_{\mathbf{v},a},\ \sigma_{\mathbf{u}}$ | orthogonal group / symplectic group / symplectic transvection / orthogonal symmetry |

*Affine, Projective, and Convex Geometry*

| | |
|---|---|
| $\mathcal{A}(\mathcal{V}),\ \mathcal{A}_0(\mathcal{V}),\ \mathrm{Sub}(\mathcal{V})$ | flats in $\mathcal{V}$ / $\mathcal{A}(\mathcal{V}) \cup \{\emptyset\}$ / subspaces of $\mathcal{V}$; $\mathcal{A}$ without arguments remains local |
| $\mathrm{affspan}(E),\ \mathrm{affhull}(E)$ | affine span / least flat containing $E$ |
| $X \parallel Y,\ X \vee Y$ | parallel flats / join of flats or projective subspaces; meet is intersection in the relevant lattice |
| $\mathbb{R}^n_+,\ \mathbb{R}^n_{++},\ \mathbf{x} \geq \mathbf{0},\ \mathbf{x} > \mathbf{0},\ \mathbf{x} \gg \mathbf{0}$ | nonzero nonnegative orthant / coordinatewise positive orthant / coordinatewise nonnegative, nonzero nonnegative, and coordinatewise positive inequalities |
| $\mathrm{conv}(S),\ \Delta_{k-1},\ C_A$ | convex hull / coefficient simplex $\{(t_i) : t_i \geq 0,\ \sum_i t_i = 1\}$ / nonnegative cone generated by the columns of $A$; distinct from cofactor and companion notation |
| $\mathcal{H}(\mathbf{N}, b),\ \mathcal{H}_\pm(\mathbf{N}, b),\ \langle \mathbf{N}, X \rangle < b$ | affine hyperplane / closed half-spaces / setwise inequality, meaning $\langle \mathbf{N}, \mathbf{x} \rangle < b$ for all $\mathbf{x} \in X$ |

*Spectral and Polynomial Structure*

| | |
|---|---|
| $E_\lambda(T),\ \mathrm{spec}(T)$ | eigenspace $\ker(T - \lambda I_\mathcal{V})$ / spectrum |
| $\mathrm{Inv}(T)$ | lattice or set of $T$-invariant subspaces |
| $\lambda_i(A),\ m_A(\lambda)$ | eigenvalues in the locally declared order / algebraic multiplicity of $\lambda$ as a root of $p_A$; distinct from measure notation $m(E)$ |
| $\mathcal{G}_\lambda(T),\ J_k(\lambda)$ | generalized eigenspace for $\lambda$ / Jordan block of size $k$ and eigenvalue $\lambda$ |
| $p_A,\ q_T,\ \chi_T$ | matrix characteristic polynomial / operator characteristic polynomial / characteristic polynomial determined by elementary divisors |
| $\mu_A,\ \mu_T,\ w_p(\lambda, A)$ | minimal polynomial of a matrix or operator / Weyr count of $\lambda$-Jordan blocks of size at least $p$; distinct from measure notation $\mu(E)$ |
| $P_j,\ T = \sum_j \mu_j P_j$ | spectral projections / spectral decomposition with distinct eigenvalues $\mu_j$ |
| $\sqrt{T},\ A^{1/2},\ \rho(A)$ | positive square root of a positive operator or matrix / spectral radius of $A$; distinct from the interval parameter in $K_\rho$ |
| $\mathbb{F}[z]$ | polynomials in $z$ over $\mathbb{F}$ |
| $\mathscr{P} = \mathbb{C}[x],\ \mathscr{P}',\ \mathscr{F} = \mathbb{C}[[t]]$ | umbral polynomial algebra / algebraic dual / formal power-series algebra; script notation distinguishes these from $\mathcal{P}(X)$ and local families $\mathcal{F}$ |
| $\mathcal{V}_T,\ \mathcal{C}_T(\mathbf{v}),\ \mathrm{ann}_T(\mathbf{v})$ | $\mathbb{F}[z]$-module induced by $T$ / $T$-cyclic subspace generated by $\mathbf{v}$ / annihilator ideal of $\mathbf{v}$ |
| $m_\mathbf{v},\ C_f,\ \mathrm{InvFact}(T),\ \mathrm{ElemDiv}(T)$ | order polynomial of $\mathbf{v}$ under $T$ / companion matrix of $f$ / invariant factors / elementary divisors |
| $e^{tA}$ | matrix exponential $\sum_{k \geq 0} (tA)^k / k!$ |

*Matrix Analysis and Numerical Structure*

| | |
|---|---|
| $A \succeq 0,\ A \succ 0$ | Hermitian positive semidefinite / Hermitian positive definite |
| $\mathrm{in}(A),\ n_+(A), n_-(A), n_0(A)$ | inertia of a Hermitian matrix / positive, negative, and zero eigenvalue counts |
| $A = QR,\ A = LL^*,\ A = LDL^t$ | QR factorization / Cholesky factorization / symmetric or Hermitian $LDL^t$ factorization |
| $A = V\Sigma W^*,\ A = X\Sigma_r Y^*$ | singular value decomposition / compact SVD with rank $r$ |
| $\sigma_j(A),\ \sigma_{\max}(A),\ \sigma_{\min}(A)$ | singular values and extreme singular values |
| $\lvert T\rvert,\ \lvert A\rvert,\ \mathrm{vol}(\Omega)$ | operator or matrix modulus / volume of a set; scalar absolute value remains $\lvert x\rvert$ |
| $A^\dagger,\ A_k$ | Moore–Penrose pseudoinverse / rank-$k$ truncated SVD approximation in an SVD context |
| $\kappa_2(A),\ \Delta A,\ \Delta\mathbf{x}$ | spectral condition number of an invertible matrix / matrix and solution perturbations |
| $\Delta(A)$ | defect from normality; distinct from perturbation notation $\Delta A$ |
| $R_k(A),\ R_k^{\mathrm{off}}(A),\ R_{\max}(A)$ | absolute row sum $\sum_j \lvert a_{kj}\rvert$ / deleted row sum $\sum_{j\neq k} \lvert a_{kj}\rvert$ / maximum absolute row sum |
| $\mathrm{Ger}_k(A),\ \mathrm{Ger}(A)$ | $k$th Gersgorin disk / union of all Gersgorin disks |
| $\mathrm{tridiag}(a, b, c)$ | tridiagonal matrix with subdiagonal $a$, diagonal $b$, and superdiagonal $c$ |
| $w,\ F_n,\ Q_n,\ \omega_n$ | half-bandwidth / unnormalised Fourier matrix / normalised Fourier matrix / primitive $n$th root used for the DFT |
| $\mathcal{K}_m(A, \mathbf{b})$ | Krylov subspace $\mathrm{span}(\mathbf{b}, A\mathbf{b}, \ldots, A^{m-1}\mathbf{b})$ |
| $Q_m,\ H_m$ | Krylov basis matrix / projected Hessenberg or tridiagonal matrix |
| $A = S - T,\ B = S^{-1}T,\ \mathbf{e}_k$ | matrix splitting / stationary iteration matrix / $k$th error vector |
| $A = D + L + U,\ B_J,\ B_{\mathrm{GS}}$ | diagonal, lower, and upper splitting / Jacobi and Gauss–Seidel iteration matrices |
| $\omega,\ S_\omega,\ T_\omega,\ B_\omega$ | SOR parameter / SOR splitting matrices / SOR iteration matrix; distinct from the differential form $\omega$ |
| $M,\ M^{-1}A,\ AM^{-1}$ | preconditioner / left and right preconditioned matrices |

## Algebraic Structures Notation

*Algebraic Language and Categories*

| | |
|---|---|
| $\mathcal{C},\ A \xrightarrow{f} B,\ \mathrm{id}_A$ | ambient category / morphism $f : A \to B$ / identity morphism of $A$; $\mathcal{C}$ remains local unless declared |
| $\mathrm{Hom}_{\mathcal{C}}(A, B),\ g \circ f$ | set of morphisms $A \to B$ in $\mathcal{C}$ / composition for compatible morphisms |
| $A \simeq B,\ F : \mathcal{C} \to \mathcal{D}$ | isomorphic objects / functor with domain $\mathcal{C}$ and codomain $\mathcal{D}$ |

*Groups, Actions, Rings, and Ideals*

| | |
|---|---|
| $G, H, K,\ e_G,\ \langle S\rangle$ | groups or subgroups / identity element of $G$ / substructure generated by $S$ in the ambient algebraic category |
| $H \leq G,\ H \trianglelefteq G,\ G/H,\ [G : H]$ | subgroup / normal subgroup / quotient group / subgroup index |
| $\mathrm{ord}(g),\ \mathrm{Aut}(G)$ | order of an element / automorphism group |
| $R, S,\ 0_R,\ 1_R,\ R^\times$ | rings / zero and identity elements of $R$ / unit group of $R$ |
| $Z(R),\ \mathrm{char}(R)$ | center of $R$ / characteristic of $R$ |
| $I, J,\ \mathfrak{a}, \mathfrak{p},\ R/I$ | ideals; fraktur letters denote distinguished ideals, such as prime ideals, when declared / quotient ring |
| $R[x],\ \mathbb{F}[z],\ \deg f$ | polynomial rings in the displayed indeterminate / degree of a polynomial |
| $U_n,\ \Omega_n,\ \Phi_n(x),\ \varphi(n)$ | complex $n$th roots of unity / primitive $n$th roots / $n$th cyclotomic polynomial / Euler totient; $\Omega_n$ is algebraic root notation, distinct from domain notation |

*Modules and Associative Algebras*

| | |
|---|---|
| $M, N,\ S \le M,\ M/N$ | modules / submodule $S$ of $M$ / quotient module |
| $\mathrm{Hom}_R(M,N),\ \mathrm{End}_R(M)$ | set of $R$-module homomorphisms $M \to N$ / endomorphism ring of $M$ |
| $\ker\tau,\ \mathrm{im}\,\tau,\ \overline{\tau}$ | kernel / image / quotient-induced map $M/\ker\tau \to \mathrm{im}\,\tau$ for a homomorphism $\tau : M \to N$ |
| $\bigoplus_i M_i,\ M = N \oplus H$ | external direct sum of modules / internal direct-sum decomposition |
| $\mathcal{C}_S(N),\ \mathcal{L}_\sigma,\ \mathcal{R}_\sigma$ | set $\{H \le N : N = S \oplus H\}$ of complements / left inverses of $\sigma$ / right inverses of $\sigma$; subscript distinguishes these from $\mathcal{L}$ and $\mathcal{L}^n$ |
| $\mathcal{F}(\mathcal{V})$ | finite-rank operators $\{T \in \mathcal{L}(\mathcal{V}) : \dim \mathrm{im}\, T < \infty\}$ in associative-algebra contexts; distinct from local function-family notation |

## Algebraic Geometry and Commutative Algebra Notation

*Status.* This part of the notes is still mostly scaffolded. The entries below reserve the notation promised by the section headings and keep it distinct from already-active algebra and analysis notation.

*Commutative Algebra*

| | |
|---|---|
| $\mathrm{Spec}\, R,\ \mathfrak{p} \in \mathrm{Spec}\, R$ | set of prime ideals of a commutative ring $R$ / prime ideal $\mathfrak{p} \subseteq R$ |
| $D(f),\ V(I)$ | standard open set $\{\mathfrak{p} \in \mathrm{Spec}\, R : f \notin \mathfrak{p}\}$ / Zariski closed set $\{\mathfrak{p} : I \subseteq \mathfrak{p}\}$ |
| $S^{-1}R,\ R_{\mathfrak{p}},\ k(\mathfrak{p})$ | localization at a multiplicative set $S$ / localization $(R \setminus \mathfrak{p})^{-1}R$ / residue field $R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}$ |
| $\mathrm{rad}(I),\ \sqrt{I}$ | radical ideal $\{r \in R : r^n \in I \text{ for some } n \ge 1\}$; use $\mathrm{rad}(\mathcal{V})$ for metric-vector-space radicals |

*Affine and Projective Geometry*

| | |
|---|---|
| $\mathbb{A}^n_k,\ \mathbb{P}^n_k$ | affine $n$-space over a field $k$ / projective $n$-space over $k$; geometric notation distinct from probability notation |
| $I(X),\ Z(I),\ k[X]$ | vanishing ideal $\{f \in k[x_1,\dots,x_n] : f|_X = 0\}$ / zero locus of an ideal / coordinate ring, typically $k[x_1,\dots,x_n]/I(X)$ |
| $f^{\#},\ \varphi^*$ | coordinate-ring pullback, e.g. $f^{\#} : k[Y] \to k[X]$ for $f : X \to Y$; distinct from adjoints and differential-form pullbacks by context |

## Analysis and Continuous Structure Notation

*Sets and Functions in Analysis*

| | |
|---|---|
| $\mathcal{P}(X)$ | power set $\{A : A \subseteq X\}$; projection symbols use $P_{\mathcal{U}}$ or $P_j$ |
| $f(A),\ f^{-1}(B)$ | image $\{f(a) : a \in A\}$ / preimage $\{x : f(x) \in B\}$ under $f$ |
| $\mathrm{range}(f),\ f\|_A$ | range $f(\mathrm{dom}\, f)$ / restriction of $f$ to $A$ |
| $\{A_i\}_{i\in I},\ \bigcup_{i\in I} A_i,\ \bigcap_{i\in I} A_i$ | indexed family of sets / set of points lying in some $A_i$ / set of points lying in every $A_i$ |
| $g \circ f,\ \mathrm{id}_X$ | composition of functions with compatible domain and codomain / identity function on $X$ |
| $J_n,\ \#E,\ A \sim_{\#} B,\ [x]$ | finite initial segment $J_0 = \emptyset$, $J_n = \{1,\dots,n\}$ / cardinality of a finite set / existence of a bijection $A \to B$ / equivalence class of $x$ for the specified relation |

*Order and Completeness*

| | |
|---|---|
| $\sup E,\ \inf E$ | least upper bound / greatest lower bound in the ambient ordered set, with values in $\overline{\mathbb{R}}$ when extended values are allowed |
| $\|x\|,\ \|a-b\|$ | real absolute value or complex modulus / usual real-line distance $d(a,b) = \|a-b\|$ |
| $\overline{\mathbb{R}}$ | extended real line $\mathbb{R} \cup \{-\infty, +\infty\}$ ordered by $-\infty < x < +\infty$ for $x \in \mathbb{R}$; the bar is conventional, not topological closure |
| $x < y,\ x \le y$ | strict order / non-strict order in the ambient ordered set |

*Sequences and Series*

| | |
|---|---|
| $(a_n),\ a_n \to a$ | sequence indexed by $n \in \mathbb{N}$ / convergence of the sequence to $a$ |
| $(a_{n_k}),\ n_1 < n_2 < \cdots$ | subsequence indexed by a strictly increasing sequence $(n_k)$ of positive integers |
| $s_n = \sum_{k=1}^{n} a_k$ | partial-sum sequence associated to $\sum a_k$ |
| $\limsup a_n = s^*,\ \liminf a_n = s_*$ | upper and lower limits in $\overline{\mathbb{R}}$, equivalently $\inf_N \sup_{n\geq N} a_n$ and $\sup_N \inf_{n\geq N} a_n$ for real sequences |
| $\sum_{n=1}^{\infty} a_n,\ \prod_{n=1}^{\infty} a_n$ | limit of partial sums / limit of finite partial products when defined |

*Metric and Topological Notation*

| | |
|---|---|
| $d(x,y),\ B_r(x)$ | metric distance / open ball $\{y : d(x,y) < r\}$ |
| $B_\varepsilon(a) = V_\varepsilon(a)$ | real-line open ball $(a-\varepsilon, a+\varepsilon)$; $V_\varepsilon(a)$ is Abbott's notation |
| $\overline{B}_r(x),\ \{G_\alpha\}_{\alpha\in A}$ | closed ball $\{y : d(x,y) \leq r\}$ / indexed open cover |
| $d_E,\ d_{\max},\ d_{\text{sum}}$ | Euclidean, maximum, and sum product metrics on a product space |
| $E^\circ,\ \overline{E},\ E^{\text{acc}}$ | interior / topological closure / set of limit points of $E$ |
| $\text{bd}(E),\ E^c$ | boundary $\overline{E} \setminus E^\circ$ / complement in the ambient space |
| $\operatorname{diam} E$ | diameter $\sup\{d(p,q) : p,q \in E\}$ of a nonempty metric set |
| $X \simeq Y$ | homeomorphic metric spaces, i.e. related by a bijection with continuous inverse |
| $A \cap \overline{B} = \emptyset,\ \overline{A} \cap B = \emptyset$ | definition of separated subsets $A$ and $B$ |
| $F_\sigma,\ G_\delta,$ meager/thin/first category | countable union of closed sets / countable intersection of open sets / subset of a countable union of nowhere-dense sets |
| residual/thick, generic | set containing a countable intersection of dense open sets / property whose truth set is residual |

*Functions, Differentiation, and Integration*

| | |
|---|---|
| $\lim_{x\to p} f(x),\ f(x+),\ f(x-)$ | limit of $f$ at $p$ / right and left limits at $x$ |
| $C(X),\ C([a,b]),\ C_b(X),\ \|f\|_\infty$ | continuous functions on $X$ and $[a,b]$ / bounded continuous functions / supremum norm |
| $f_n \to f,\ f_n \rightrightarrows f$ | pointwise convergence / uniform convergence on the ambient domain |
| $f'(x),\ Df(a),\ D_j f_i,\ \partial_i f,\ J_f(x)$ | one-variable derivative, total derivative at $a$, component partial $D_j f_i$, scalar partial, and Jacobian determinant; $J_f$ is distinct from $J_n$ and $J_k(\lambda)$ |
| $\nabla f,\ D_{\mathbf{u}} f,\ C^r,\ C^\infty,\ C^\omega$ | gradient, directional derivative, and classes $C^r$, $C^\infty$, $C^\omega$ |
| $\Pi = \{x_0, \ldots, x_n\},\ c_i,\ \Delta x_i,\ \Delta\alpha_i$ | partition of $[a,b]$, tags $c_i \in [x_{i-1}, x_i]$, and increments $\Delta x_i = x_i - x_{i-1}$, $\Delta\alpha_i = \alpha(x_i) - \alpha(x_{i-1})$ |
| $R(f,\Pi,\{c_i\}),\ \delta,\ \delta(x)$ | $R(f,\Pi,\{c_i\}) = \sum_{i=1}^{n} f(c_i)\Delta x_i$ / constant mesh bound / gauge; uppercase $R(\cdots)$ is local to Riemann sums |
| $U(\Pi,f,\alpha),\ L(\Pi,f,\alpha)$ | Riemann–Stieltjes upper and lower sums with integrator $\alpha$ |
| $f \in \mathcal{R},\ f \in \mathcal{R}(\alpha)$ | ordinary Riemann integrability / Riemann–Stieltjes integrability with respect to $\alpha$ |
| $\int_a^b f,\ \int_a^b f\,d\alpha,\ \int_E f\,d\mu$ | ordinary Riemann integral / Riemann–Stieltjes integral / measure integral |
| $\int_a^\infty f,\ \int_{-\infty}^\infty f$ | one-sided and two-sided improper integrals as limits over finite truncations |
| $\ell(\Pi,\gamma),\ \ell(\gamma)$ | polygonal length of $\gamma$ along $\Pi$ / curve length |
| $I^k,\ Q^k,\ \operatorname{supp}(f)$ | $k$-cell, standard $k$-simplex, and closed support $\overline{\{x : f(x) \neq 0\}}$; $I^k$ is not an identity operator |
| $dx_I,\ dx_I \wedge dx_J$ | basic form indexed by $I$ / wedge product of basic forms |
| $\omega,\ d\omega,\ T^*\omega,\ \partial_{\text{ch}}\Psi$ | differential form, exterior derivative, pullback by $T$, and chain boundary |
| $\int_\Psi \omega,\ \partial_{\text{ch}}[\mathbf{p}_0,\ldots,\mathbf{p}_k]$ | integral of $\omega$ over the chain $\Psi$ / oriented boundary $\sum_{j=0}^{k}(-1)^j[\mathbf{p}_0,\ldots,\widehat{\mathbf{p}_j},\ldots,\mathbf{p}_k]$ |
| $dV,\ dA,\ ds$ | volume, area, and arc-length elements |

*Power Series, Fourier, and Special Functions*

| | |
|---|---|
| $\sum_{n\geq 0} c_n(x-a)^n,\ R$ | power series centered at $a$ / radius of convergence in this context, distinct from $R(\cdots)$ and the $R$ in $QR$ |
| $I_R,\ K_\rho$ | centered interval $I_R = (-R, R)$ / compact subinterval $K_\rho = [-\rho, \rho] \subset I_R$ for $0 < \rho < R$ |
| $\widehat{f}(m),\ s_N(f;x),\ s_N(f)$ | $\widehat{f}(m) = (2\pi)^{-1}\int_{-\pi}^{\pi} f(x)e^{-imx}\,dx$ / $s_N(f;x) = \sum_{\lvert m\rvert\leq N}\widehat{f}(m)e^{imx}$, viewed pointwise or as a function |
| $f \sim_{\text{coef}} \sum \widehat{f}(n)e^{inx},\ f \sim_{\text{coef}} \sum c_n\phi_n$ | formal Fourier or orthonormal coefficient expansion; no convergence assertion is implied |
| $D_N(t)$ | Dirichlet kernel $D_N(t) = \sum_{m=-N}^{N} e^{imt}$; uppercase $D_N$ is not derivative notation |
| $\mathsf{E},\ \mathsf{L},\ \mathsf{C},\ \mathsf{S}$ | power-series exponential, inverse logarithm, cosine, and sine |
| $\Gamma(x),\ \mathrm{B}(x,y)$ | gamma function / beta function in roman type, distinct from balls $B_r(x)$ and variables $B$ |

*Measure and Function Spaces*

| | |
|---|---|
| $E \in \mathcal{M},\ E \in \mathcal{M}(\mu)$ | $E$ measurable in the ambient $\sigma$-algebra / Rudin measurable set generated by $\mu$ |
| $\mathcal{B}(\mathbb{R}^n),\ \mathcal{L}^n$ | Borel $\sigma$-algebra on $\mathbb{R}^n$ / Lebesgue measurable subsets of $\mathbb{R}^n$; not a basis $\mathcal{B}$, a linear-map space $\mathcal{L}$, or a least-squares set LS |
| $\mathcal{E},\ \mathcal{M}_F(\mu),\ A\triangle B,\ d_\mu(A,B)$ | elementary sets, finitely $\mu$-measurable sets, symmetric difference, and $d_\mu(A,B) = \mu^*(A\triangle B)$; $\mathcal{M}$ is measure-theoretic here |
| $\mu^*(E),\ m^*(E),\ \mathbf{1}_E$ | $\mu$-outer measure / Lebesgue outer measure / indicator $\mathbf{1}_E(x) = 1$ for $x \in E$ and 0 otherwise |
| $\mathrm{vol}_n(B),\ m_n^*(\Omega)$ | $n$-dimensional box volume / box-cover outer measure $\inf\{\sum_{j\in J}\mathrm{vol}_n(B_j) : (B_j)_{j\in J} \text{ covers } \Omega,\ J \text{ at most countable}\}$ |
| $m(E),\ \mu(E)$ | Lebesgue or elementary volume of $E$ / measure of $E$; not algebraic multiplicity $m_A(\lambda)$ or minimal polynomial $\mu_A$ |
| $\lambda^+,\ \lambda^-,\ \lvert\lambda\rvert,\ \lambda \ll \mu,\ \lambda \perp \mu,\ \frac{d\lambda}{d\mu}$ | positive variation, negative variation, total variation, absolute continuity, mutual singularity, and Radon–Nikodym derivative |
| $\bowtie,\ \{f \bowtie a\}$ | relation in $\{>, \geq, <, \leq\}$ / level set $\{x : f(x) \bowtie a\}$ |
| $f^+,\ f^-,\ \mathcal{S}^+(f),\ \mathcal{I}_E(s)$ | $f^+ = \max(f, 0),\ f^- = -\min(f, 0)$ / simple minorants of $f$ / simple-function integral on $E$ |
| $f \sim_{\text{a.e.}} g \text{ on } E,\ \text{a.e.}$ | $\mu(\{x \in E : f(x) \neq g(x)\}) = 0$ / outside a null set |
| $E_x,\ E^y,\ f_x,\ f^y$ | product slices $E_x = \{y : (x,y) \in E\}$, $E^y = \{x : (x,y) \in E\}$, and $f_x(y) = f^y(x) = f(x,y)$ |
| $\mathcal{U}f,\ F_a$ | undergraph $\{(x,t) : 0 < t < f(x)\}$ for $f \geq 0$ / superlevel set $F_a = \{x : f(x) > a\}$; $\mathcal{U}f$ is not a vector subspace |
| $[f : A],\ [B : Q],\ \mathrm{dp}(E)$ | $[f : A] = m(A)^{-1}\int_A f$ / $[B : Q] = m(B \cap Q)/m(Q)$ / density-point set of $E$ |
| $\mathscr{L}^1(\mu),\ \mathscr{L}^1_{\text{loc}}(\mathbb{R}^n)$ | measurable $f$ with $\int \lvert f\rvert\,d\mu < +\infty$ / locally integrable functions; script $\mathscr{L}$ is distinct from $\mathcal{L}^n$ and $\mathcal{L}$ |
| $L^p(\mu),\ L^p(E),\ \lVert f\rVert_p$ | Lebesgue space modulo a.e. equality with respect to $\mu$ or the ambient measure on $E$ / $\lVert f\rVert_p = (\int \lvert f\rvert^p\,d\mu)^{1/p}$ |

## Probability and Stochastic Structure Notation

*Probability Spaces and Distributions*

| | |
|---|---|
| $(\Omega, \mathcal{F}, \mathbb{P})$, $\Pr(A)$ | probability space / event probability $\Pr(A) = \mathbb{P}(A)$ for $A \in \mathcal{F}$; explicit probability notation keeps $\mathbb{P}_k^n$ and $\mathbb{F}_0$ unambiguous |
| $X, Y$, $X_*\mathbb{P}$, $\mathcal{L}(X)$ | random variables / pushforward law $X_*\mathbb{P}(B) = \mathbb{P}(X^{-1}(B))$ / law of $X$ when locally declared; not Lebesgue $\mathcal{L}^n$ |
| $X \sim F$, $X \overset{\text{dist}}{=} Y$, $X_1, \dots, X_n$ iid | $X$ has distribution $F$ / $X$ and $Y$ have the same distribution / independent and identically distributed sample |
| $F(t)$, $g(t)$, $\Phi(t)$ | distribution function $F(t) = \Pr(X \le t)$ / density $g$ / standard normal distribution function in probabilistic contexts |
| $\mathbb{E}[X]$, $\operatorname{Var}(X)$, $\operatorname{Cov}(X, Y)$ | expectation / $\operatorname{Var}(X) = \mathbb{E}[(X - \mathbb{E}X)^2]$ / $\operatorname{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}X)(Y - \mathbb{E}Y)]$, when finite |
| $\operatorname{Corr}(X, Y)$, $\rho$, $\sigma_{XY}$ | correlation coefficient / local correlation parameter / covariance shorthand $\sigma_{XY} = \operatorname{Cov}(X, Y)$ when declared |
| $\psi_X(t)$ | moment-generating function $\psi_X(t) = \mathbb{E}e^{tX}$ on its domain of finiteness |
| $X \perp\!\!\!\perp Y$ | independence of random variables, equivalently independence of $\sigma(X)$ and $\sigma(Y)$ when defined |
| $X \perp\!\!\!\perp Y \mid Z$ | conditional independence of $X$ and $Y$ given $Z$ |
| $X_n \xrightarrow{\mathbb{P}} X$, $X_n \rightsquigarrow X$, $X_n \xrightarrow{L^p} X$, $X_n \xrightarrow{\text{a.s.}} X$, $X_n \xrightarrow{qm} X$ | convergence in probability / convergence in distribution / convergence in $L^p$ / almost-sure convergence / convergence in quadratic mean |

*Stochastic Processes and Markov Structure*

| | |
|---|---|
| $(X_t)_{t \in T}$, $\mathcal{F}_t$ | stochastic process indexed by $T$ / filtration $(\mathcal{F}_t)$ when locally declared; this $\mathcal{F}_t$ is not the umbral algebra $\mathscr{F}$ |
| $\mathbf{x}$, $\mathbf{e}$, $A\mathbf{e} = \mathbf{e}$ | probability vector $\mathbf{x} \ge 0$, $\sum_i x_i = 1$ / all-ones vector $\mathbf{e} = (1, \dots, 1)^t$ / row-stochastic Markov convention $A = [a_{ij}] \ge 0$, $A\mathbf{e} = \mathbf{e}$ |
| $\pi$, $\pi A = \pi$, $A^p$ | stationary distribution written as a row probability vector / stationarity equation / $p$-step Markov matrix |
| $P_t$, $p_t(x, y)$ | transition semigroup / transition probability, kernel, or density from $x$ to $y$, when introduced |

## Numerical and Computational Mathematics Notation

*Status.* Apart from numerical linear algebra, already covered under Linear and Operator Structures, the numerical and computational chapters remain placeholders. The entries below reserve notation only; they do not introduce PDE or machine-learning conventions.

*Floating Point, Approximation, and Error*

| | |
|---|---|
| $\operatorname{fl}(x)$, $u$, $\varepsilon_{\text{mach}}$ | floating-point rounded value, unit roundoff, and machine epsilon |
| $x_h$, $e_h$, $\mathcal{O}(h^p)$ | approximation at discretization scale $h$, corresponding error, and Landau order as $h \to 0$ |
| $\widehat{x}$, $\Delta x$, $r = b - A\widehat{x}$ | computed approximation, perturbation or error in $x$, and residual for $Ax = b$ |

*Monte Carlo and Stochastic Computation*

| | |
|---|---|
| $X_1, \dots, X_N$, $\widehat{\mu}_N$ | samples / sample-mean estimator $\widehat{\mu}_N = N^{-1} \sum_{i=1}^N X_i$ when estimating $\mu = \mathbb{E}X$ |
| $\operatorname{Var}(\widehat{\mu}_N)$, $\operatorname{SE}(\widehat{\mu}_N)$ | estimator variance / standard error $\operatorname{SE}(\widehat{\mu}_N) = \sqrt{\operatorname{Var}(\widehat{\mu}_N)}$ |
| $w_i$, $\widehat{I}_N$ | importance weights / Monte Carlo estimate of an integral or expectation, when locally introduced |

# Part I
# Set Theory, Logic, and Foundations

Foundational prerequisites collect the logical, set-theoretic, functional, order-theoretic, and cardinal language used before the dedicated algebraic and analytic foundations sections.

## 1 Sets

Sets provide the membership language used throughout the notes: elements, subsets, products, power sets, operations, and indexed families.

### 1.1 Introduction to Sets

**Definition 1.1.1.** *Sets and Membership*
A *set* is a collection. Write $x \in E$, $x \notin E$, and $\emptyset$.

**Remark 1.1.2.** *Set-Builder Notation*
$\{x \in X : P(x)\}$ means all $x \in X$ satisfying $P(x)$.
*Key Insight*: Expand membership conditions.

### 1.2 Cartesian Products

**Definition 1.2.1.** *Cartesian Product*
For sets $X$ and $Y$, the *Cartesian product* is $X \times Y := \{(x, y) : x \in X,\ y \in Y\}$.
The order matters: the first coordinate comes from $X$, and the second comes from $Y$.




Figure 1: $A \times B$: first coordinate in $A$, second in $B$.

### 1.3 Subsets and Power Sets

**Definition 1.3.1.** *Basic Subset Terminology*
Let $A, B$ be sets. Write $A \subseteq B$; $A = B$ iff $A \subseteq B$ and $B \subseteq A$.

(i) $A$ is a *proper subset* of $B$, written $A \subsetneq B$, iff $A \subseteq B$ and $A \neq B$.
(ii) $A$ is a *singleton* iff $\exists x$ such that $A = \{x\}$.
(iii) $A$ is *nonempty* iff $A \neq \emptyset$.

**Definition 1.3.2.** *Power Set*
For a set $X$, the *power set* is $\mathcal{P}(X) := \{S : S \subseteq X\}$.

**Method 1.3.3.** *Proof by Two Containments*
To prove $A = B$, prove both inclusions.

(i) To prove $A \subseteq B$, let $x \in A$ be arbitrary and use the definitions and hypotheses to show $x \in B$.
(ii) To prove $B \subseteq A$, let $x \in B$ be arbitrary and use the definitions and hypotheses to show $x \in A$.
(iii) Once both containments are proved, conclude $A = B$ by extensionality of sets.

If both sides are described by membership conditions, this is a chain, $x \in A \iff \cdots \iff x \in B$.

**Example 1.3.4.** *Small Power Sets*

(i) If $X = \emptyset$, then $\mathcal{P}(X) = \{\emptyset\}$.

(ii) If $X = \{a, b\}$ with $a \neq b$, then $\mathcal{P}(X) = \{\emptyset, \{a\}, \{b\}, X\}$.

Hence $S \in \mathcal{P}(X)$ iff $S \subseteq X$; membership in $\mathcal{P}(X)$ is containment in $X$.

## 1.4 Union, Intersection, Difference, and Complement

**Definition 1.4.1.** *Binary Set Operations*
For $A, B \subseteq X$,

$$\begin{aligned} A \cup B &:= \{x \in X : x \in A \text{ or } x \in B\}, \\ A \cap B &:= \{x \in X : x \in A \text{ and } x \in B\}, \\ A \setminus B &:= \{x \in X : x \in A \text{ and } x \notin B\}, \\ A^c &:= X \setminus A. \end{aligned}$$

**Definition 1.4.2.** *Symmetric Difference*
For sets $A$ and $B$, the *symmetric difference* is $A \triangle B := (A \setminus B) \cup (B \setminus A) = (A \cup B) \setminus (A \cap B)$.
It records the elements belonging to exactly one of $A$ and $B$.

**Proposition 1.4.3.** *Basic Set Algebra*
*Let $A, B, C \subseteq X$. Then*

*(i)* $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$;
*(ii)* $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$;
*(iii)* $(A \cup B)^c = A^c \cap B^c$;
*(iv)* $(A \cap B)^c = A^c \cup B^c$.

*Proof.* It suffices to prove (i) and (iii); the others are similar. Let $x \in X$. Then

(i) $x \in A \cap (B \cup C) \iff x \in A$ and $(x \in B$ or $x \in C) \iff (x \in A \cap B)$ or $(x \in A \cap C) \iff x \in (A \cap B) \cup (A \cap C)$
(iii) $x \in (A \cup B)^c \iff x \notin A \cup B \iff x \notin A$ and $x \notin B \iff x \in A^c \cap B^c$

*Structural Commentary*: Set algebra is membership logic.
*Key Insight*: Translate sets into conditions on $x$. □




Figure 2: De Morgan's law says that being outside a union is the same as being outside each set.

## 1.5 Indexed Families and Products

**Definition 1.5.1.** *Families, Indexed Families, Unions, and Intersections*
Let $X$ be a set and let $I$ be an index set.

(i) A *family* or *collection* of sets is a set whose members are sets.
(ii) An *indexed family* in $X$ over $I$ is a map $A : I \to \mathcal{P}(X)$; write $A_i := A(i)$ and $\{A_i\}_{i \in I}$.
(iii) For an indexed family $\{A_i\}_{i \in I}$ in $X$, $\bigcup_{i \in I} A_i := \{x \in X : \exists i \in I,\ x \in A_i\}$.
(iv) For an indexed family $\{A_i\}_{i \in I}$ in $X$, $\bigcap_{i \in I} A_i := \{x \in X : \forall i \in I,\ x \in A_i\}$.

Union corresponds to $\exists i \in I$; intersection corresponds to $\forall i \in I$.

**Definition 1.5.2.** *Disjoint Families and Parametrizations*
Let $\mathcal{F}$ be a family of sets and let $\{A_i\}_{i \in I}$ be an indexed family.

(i) A family $\mathcal{F}$ is *pairwise disjoint* iff $E, F \in \mathcal{F}$ and $E \neq F$ imply $E \cap F = \emptyset$.
(ii) An indexed family $\{A_i\}_{i \in I}$ *parametrizes* a family $\mathcal{F}$ iff $\mathcal{F} = \{A_i : i \in I\}$.

**Theorem 1.5.3.** *Arbitrary De Morgan Laws*
*Let $X$ be a set. For any collection $\{E_\alpha\}$ of subsets of $X$,* $\left(\bigcup_\alpha E_\alpha\right)^c = \bigcap_\alpha E_\alpha^c$ *and* $\left(\bigcap_\alpha E_\alpha\right)^c = \bigcup_\alpha E_\alpha^c$.

*Proof.* The first identity follows from membership: $x \in \left(\bigcup_\alpha E_\alpha\right)^c \iff x \notin E_\alpha \ \forall \alpha \iff x \in \bigcap_\alpha E_\alpha^c$.
The second follows by applying the first identity to the complements $E_\alpha^c$, or by the same membership argument.
*Structural Commentary*: Complements exchange unions and intersections.
*Key Insight*: Negating "for some" gives "for every", and conversely. □

**Definition 1.5.4.** *Cartesian Product of an Indexed Family*
Let $\{E_i\}_{i\in I}$ be an indexed family of sets and set $U := \bigcup_{i\in I} E_i$.
The *Cartesian product* is $\prod_{i\in I} E_i := \{x : I \to U : \forall i \in I,\ x(i) \in E_i\}$.

(i) If $I = \{1, 2\}$, then $x \mapsto (x(1), x(2))$ gives the canonical bijection $\prod_{i\in\{1,2\}} E_i \cong E_1 \times E_2$.

(ii) If $E_i = E$ for every $i \in I$, then $\prod_{i\in I} E_i = E^I = \{x : I \to E\}$.

Different parametrizations of the same family may give different indexed products.

**Remark 1.5.5.** *Index Variables*
For sets $(A_n)$, $\bigcup_n A_n$ means membership in some $A_n$; $\bigcap_n A_n$, membership in every $A_n$.
In $\bigcup_{i\in I} A_i$ and $\bigcap_{i\in I} A_i$, the index $i$ is bound: for any fresh bound index $j$, $\bigcup_{i\in I} A_i = \bigcup_{j\in I} A_j$ and $\bigcap_{i\in I} A_i = \bigcap_{j\in I} A_j$.

## 1.6 Russell's Paradox and Axioms

*Placeholder*: Add Russell's paradox as a motivation for restricted comprehension, then state the informal role of extensionality, separation, pairing, union, power set, infinity, replacement, foundation, and choice.

# 2 Logic and Proofs

Logic fixes the grammar of mathematical statements before the notes use that grammar for definitions, theorems, and proofs.

## 2.1 Statements and Well-Formed Language

**Principle 2.1.1.** *Well-Formed Statements*
A mathematical statement is well formed only after every symbol has a declared role and every expression has a defined meaning.

## 2.2 Connectives and Negation

**Principle 2.2.1.** *Connectives and Negation*
The word "or" is inclusive in mathematics, and the basic negation rules are

$$\neg(a \wedge b) \iff \neg a \vee \neg b, \qquad \neg(a \vee b) \iff \neg a \wedge \neg b, \qquad \neg(a \Rightarrow b) \iff a \wedge \neg b$$

## 2.3 Quantifiers and Bound Variables

**Principle 2.3.1.** *Language Carries Structure*
Objects, hypotheses, conclusions, and quantifier order matter. A variable introduced by a quantifier, summation sign, integral, set-builder condition, or similar construction is *bound* inside that construction; changing its name does not change the statement as long as its scope is unchanged and no free variable is captured. For instance,

$$\forall x \in X\ \exists y \in Y\ P(x, y), \qquad \exists y \in Y\ \forall x \in X\ P(x, y)$$

are different: in the first $y$ may depend on $x$; in the second one $y$ works for all $x$.

## 2.4 Proof Grammar

**Method 2.4.1.** *Proof Grammar*
In theorem statements, put the hypotheses before the conclusion and keep quantifiers in the order meant.
In particular, switching $\forall$ and $\exists$ usually changes the statement; see Principle 2.3.1.
Before applying a rule, also check that the displayed expressions are well formed; an undefined expression makes the sentence ill formed, not merely false.
*Key Insight*: Before proving or negating a statement, parse its quantifiers and logical connectives.

## 2.5 Proof by Contradiction

**Method 2.5.1.** *Proof by Contradiction*
By Principle 2.2.1, a proof by contradiction of $a \Rightarrow b$ assumes both $a$ and $\neg b$ and derives an impossibility.

## 2.6 Mathematical Induction and Well-Ordering

**Principle 2.6.1.** *Mathematical Induction*
Let $P_1, P_2, \ldots$ be statements indexed by positive integers. To prove $P_n$ for every $n \in \mathbb{N}$, it is enough to prove:

(i) *basis*: $P_1$ is true;
(ii) *induction step*: for every $n \in \mathbb{N}$, $P_n$ implies $P_{n+1}$.

The induction step is an implication; it does not prove $P_{n+1}$ in isolation.
The basis starts the chain, and the implication carries truth from one statement to the next.

**Principle 2.6.2.** *Well-Ordering of $\mathbb{N}$*
Every nonempty subset of $\mathbb{N}$ has a least element, justifying first unused indices, first repeated values, and increasing listings of infinite subsets of $\mathbb{N}$.

# 3 Counting and Finite Sums

Counting records finite constructions before cardinality and infinite sets are treated abstractly.

## 3.1 Lists and Counting Principles

Counting arguments turn structured finite choices into cardinalities. In probability, this is the bridge from a uniform finite sample space to numerical probabilities.

**Definition 3.1.1.** *Lists, Words, and Product Sets* Let $S_1, \ldots, S_p$ be finite sets.

(i) A *list* of length $p$ with $i$th entry in $S_i$ is an element $(s_1, \ldots, s_p) \in S_1 \times \cdots \times S_p$.
(ii) If $S_1 = \cdots = S_p = S$, a list in $S^p$ may be called a *word* of length $p$ over the alphabet $S$.
(iii) A counting model is *ordered* when changing the order of the selected objects changes the outcome.

**Theorem 3.1.2.** *Multiplication Principle*
*Let $p \geq 1$ and let $S_1, \ldots, S_p$ be finite sets with $|S_i| = n_i$. Then $|S_1 \times \cdots \times S_p| = \prod_{i=1}^{p} n_i$.*
*Equivalently, a $p$-stage experiment whose $i$th stage has $n_i$ possible choices after every history has $\prod_{i=1}^{p} n_i$ ordered outcomes; in particular, two stages of sizes $m, n$ give $mn$ ordered outcomes.*

*Proof.* The case $p = 1$ is immediate.
For $p = 2$, write $|S_1| = m$ and $|S_2| = n$, and choose enumerations $S_1 = \{a_1, \ldots, a_m\}$ and $S_2 = \{b_1, \ldots, b_n\}$.
The map $\{1, \ldots, m\} \times \{1, \ldots, n\} \to S_1 \times S_2$, $(i, j) \mapsto (a_i, b_j)$, is a bijection, so $|S_1 \times S_2| = mn$.
For $p > 2$, use induction: $|S_1 \times \cdots \times S_p| = |(S_1 \times \cdots \times S_{p-1}) \times S_p| = (\prod_{i=1}^{p-1} n_i) n_p = \prod_{i=1}^{p} n_i$.
*Structural Commentary*: Outcomes form a finite product, or a constant-branching tree.
*Key Insight*: Counts multiply because every history has equally many extensions. □

**Example 3.1.3.** *Finite Words by the Multiplication Principle*

(i) A playing card can be described by a suit and a face value, so there are $4 \cdot 13 = 52$ suit-value combinations.
(ii) There are $2^8 = 256$ binary words of length 8.
(iii) A DNA word of length $N$ over the alphabet $\{A, C, G, T\}$ has $4^N$ possible sequences.
A codon of length 3 therefore has $4^3 = 64$ possible words.

*Key Insight*: Independent slots multiply; word counts grow exponentially with length.

**Proposition 3.1.4.** *Addition, Subtraction, and Division Principles*

*(i) Addition: if finite sets $A_1, \ldots, A_r$ are pairwise disjoint, then $|\bigcup_{i=1}^{r} A_i| = \sum_{i=1}^{r} |A_i|$.*
*(ii) Subtraction: if $B \subseteq A$ and $A$ is finite, then $|A \setminus B| = |A| - |B|$.*
*(iii) Division: if $f : Y \to X$ is surjective, $X, Y$ are finite, $d \geq 1$, and $\left|f^{-1}(\{x\})\right| = d$ for every $x \in X$. Then $|Y| = d\,|X|$ and $|X| = |Y|/d$.*

*Proof.*

(i) Let $n_i = |A_i|$ and enumerate $A_i = \{a_{i,1}, \ldots, a_{i,n_i}\}$. Since the $A_i$ are pairwise disjoint, $(i, k) \mapsto a_{i,k}$ is a bijection $\{(i, k) : 1 \leq i \leq r,\ 1 \leq k \leq n_i\} \to \bigcup_{i=1}^{r} A_i$, so $|\bigcup_{i=1}^{r} A_i| = \sum_{i=1}^{r} n_i = \sum_{i=1}^{r} |A_i|$.
(ii) Since $A$ is the disjoint union of $B$ and $A \setminus B$, part (i) gives $|A| = |B| + |A \setminus B|$, so $|A \setminus B| = |A| - |B|$.

(iii) For $x \neq x'$, $f^{-1}(\{x\}) \cap f^{-1}(\{x'\}) = \varnothing$, and $Y = \bigcup_{x \in X} f^{-1}(\{x\})$. Thus the fibers partition $Y$, so by (i), $|Y| = \sum_{x \in X} \left|f^{-1}(\{x\})\right| = \sum_{x \in X} d = d\,|X|$; since $d \geq 1$, $|X| = |Y|/d$.

*Structural Commentary*: Count parts; quotient equal fibers.
*Key Insight*: Make representations disjoint, then add their fiber sizes. □

## 3.2 Permutations, Subsets, and Binomial Coefficients

**Definition 3.2.1.** *Permutations and Sign*
Let $n \geq 1$. Let $\mathfrak{S}_n$ denote the set of permutations of $\{1, \ldots, n\}$.
*Sign* of $\sigma \in \mathfrak{S}_n$ is $\operatorname{sign}(\sigma) = 1$ if number of pairs $j < k$ with $\sigma(j) > \sigma(k)$ is even, and $\operatorname{sign}(\sigma) = -1$ otherwise.

**Definition 3.2.2.** *Factorials and Binomial Coefficients*

(i) The *factorial* is defined by $0! := 1$ and, for $n \in \mathbb{N}$ with $n \geq 1$, $n! := n(n-1)\cdots 3 \cdot 2 \cdot 1$.
(ii) For integers $0 \leq r \leq n$, the *falling factorial* is $(n)_r := n(n-1)\cdots(n-r+1)$, with $(n)_0 := 1$.
(iii) For integers $0 \leq k \leq n$, the *binomial coefficient* is $\binom{n}{k} := \frac{n!}{k!(n-k)!}$.
(iv) The symbol $\binom{n}{k}$ is read *$n$ choose $k$*.

**Proposition 3.2.3.** *Ordered Sampling With and Without Replacement*
*Let $S$ be finite with $|S| = n$, let $0 \leq r \leq n$, and set $D_r(S) := \{(s_1, \ldots, s_r) \in S^r : s_i \neq s_j$ for all $1 \leq i < j \leq r\}$. Use $S^0 = \{()\}$ and empty products equal 1.*

*(i) With replacement: the ordered samples are $S^r$, and $|S^r| = n^r$.*
*(ii) Without replacement: the ordered samples are $D_r(S)$, and $|D_r(S)| = (n)_r$, where $(n)_0 = 1$ and $(n)_r = n(n-1)\cdots(n-r+1)$ for $r \geq 1$.*
*(iii) Full orderings: $|D_n(S)| = n!$.*

*Proof.*

(i) If $r = 0$, then $\left|S^0\right| = 1 = n^0$; if $r \geq 1$, Theorem 3.1.2 gives $|S^r| = \prod_{k=1}^{r} n = n^r$.
(ii) For $k = 1, \ldots, r$, after choosing distinct $s_1, \ldots, s_{k-1}$, the entry $s_k$ has $n - k + 1$ choices, so $|D_r(S)| = \prod_{k=1}^{r}(n-k+1) = (n)_r$.
(iii) Taking $r = n$ in (ii) gives $|D_n(S)| = (n)_n = n!$.

*Structural Commentary*: Replacement counts all tuples; no replacement counts injective tuples.
*Key Insight*: Order gives slots; replacement determines whether choices stay fixed or decrease. □

**Proposition 3.2.4.** *Unordered Samples Without Replacement*
*Let $S$ be finite with $|S| = n$, let $0 \leq r \leq n$, and set $\mathcal{P}_r(S) := \{A \subseteq S : |A| = r\}$. Then $|\mathcal{P}_r(S)| = \binom{n}{r} = \frac{(n)_r}{r!} = \frac{n!}{(n-r)!r!}$.*

*Proof.* Let $\pi : D_r(S) \to \mathcal{P}_r(S)$ be $\pi(s_1, \ldots, s_r) = \{s_1, \ldots, s_r\}$.
This map is surjective, and for each $A \in \mathcal{P}_r(S)$, $\pi^{-1}(\{A\})$ consists of the $r!$ orderings of $A$.
By Proposition 3.2.3, $|D_r(S)| = (n)_r$; by the division principle, $|\mathcal{P}_r(S)| = (n)_r/r!$.
Since $(n)_r = n!/(n-r)!$, the claimed identity follows.
*Structural Commentary*: Unordered samples are ordered injective samples modulo reordering.
*Key Insight*: Count ordered first, then divide by the $r!$ orders of each subset. □

**Example 3.2.5.** *Committees*
The number of ways to choose a committee of 3 people from a class of 20 people is $\binom{20}{3} = \frac{20!}{3!17!} = \frac{20 \cdot 19 \cdot 18}{3 \cdot 2 \cdot 1} = 1140$.

**Proposition 3.2.6.** *Edge Values and Symmetry*
*For $n \in \mathbb{N}$ and $k \in \mathbb{Z}$ with $0 \leq k \leq n$:*

*(i) Edge values: $\binom{n}{0} = \binom{n}{n} = 1$.*
*(ii) Symmetry: $\binom{n}{k} = \binom{n}{n-k}$.*

*Proof.* Let $n \in \mathbb{N}$ and $k \in \mathbb{Z}$ with $0 \leq k \leq n$.

(i) Since $0! := 1$, $\binom{n}{0} = \frac{n!}{0!n!} = 1$ and $\binom{n}{n} = \frac{n!}{n!0!} = 1$.
(ii) Since $0 \leq n - k \leq n$, $\binom{n}{n-k} = \frac{n!}{(n-k)!(n-(n-k))!} = \frac{n!}{(n-k)!k!} = \frac{n!}{k!(n-k)!} = \binom{n}{k}$.

*Structural Commentary*: Choosing the selected $k$ objects is equivalent to choosing the omitted $n - k$ objects.
*Key Insight*: Complements turn $k$-subsets into $(n-k)$-subsets. □

**Proposition 3.2.7.** *Binomial Theorem and Subset Count*
*For $n \in \mathbb{N}$ and commuting $a, b$, $(a+b)^n = \sum_{k=0}^{n} \binom{n}{k} a^k b^{n-k}$.*
*If $S$ is finite with $|S| = n$, then $|\mathcal{P}(S)| = 2^n = \sum_{k=0}^{n} \binom{n}{k}$.*

*Proof.* Write $(a+b)^n = \prod_{j=1}^{n}(a+b)$.
For each $I \subseteq \{1,\dots,n\}$, choose $a$ from factors in $I$ and $b$ from the rest, giving $a^{|I|}b^{n-|I|}$.
Grouping by $k = |I|$ gives $(a+b)^n = \sum_{k=0}^{n}\binom{n}{k}a^k b^{n-k}$.
For $|S| = n$, the sets $\mathcal{P}_k(S) := \{A \subseteq S : |A| = k\}$ partition $\mathcal{P}(S)$, so $|\mathcal{P}(S)| = \sum_{k=0}^{n}|\mathcal{P}_k(S)| = \sum_{k=0}^{n}\binom{n}{k}$.
Also $|\mathcal{P}(S)| = 2^n$ by choosing in/out for each element.
*Structural Commentary*: Expansion and subset counting both group binary choices by size.
*Key Insight*: A term is determined by which positions choose $a$. □

**Definition 3.2.8.** *Multisets*
Let $S$ be a set. A *multiset on $S$* is a pair $(S_0, m)$ such that

(i) $S_0 \subseteq S$ is the *underlying set*;
(ii) $m : S_0 \to \mathbb{Z}_{>0}$ is the *multiplicity function*;
(iii) $m(s)$ is the *multiplicity* of $s \in S_0$;
(iv) if $S_0$ is finite, the *size* is $|m| := \sum_{s\in S_0} m(s)$.

Equivalently, record $(S_0, m)$ as $\{(s, m(s)) : s \in S_0\}$.

**Example 3.2.9.** *Multiset Bookkeeping*
The list $\{a,a,b,b,b,c\}$ is the multiset on $\{a,b,c\}$ with multiplicities $2,3,1$. This language is useful whenever order is irrelevant but repetition matters, such as eigenvalues counted with algebraic multiplicity.

**Definition 3.2.10.** *Multinomial Coefficients*
For nonnegative integers $n_1,\dots,n_r$ with $n_1+\cdots+n_r = n$, define $\binom{n}{n_1\, n_2\cdots n_r} := \frac{n!}{n_1!n_2!\cdots n_r!}$.

**Proposition 3.2.11.** *Multinomial Count and Expansion*
*Let $r \geq 1$, let $\mathbf{n} = (n_1,\dots,n_r) \in \mathbb{N}_0^r$ with $n = \sum_{i=1}^{r} n_i$, and let $x_1,\dots,x_r$ commute.*
*If $\mathcal{M}_{\mathbf{n}}$ is the set of labeled partitions $(A_1,\dots,A_r)$ of $\{1,\dots,n\}$ with $|A_i| = n_i$,*
*then $|\mathcal{M}_{\mathbf{n}}| = \binom{n}{n_1\, n_2\cdots n_r} = \frac{n!}{n_1!n_2!\cdots n_r!}$, and $(x_1+\cdots+x_r)^n = \sum_{m_1+\cdots+m_r=n}\binom{n}{m_1\, m_2\cdots m_r}x_1^{m_1}\cdots x_r^{m_r}$.*

*Proof.* Choose the parts successively. After $A_1,\dots,A_{j-1}$ are chosen, there are $n - \sum_{i=1}^{j-1} n_i$ positions left, so $A_j$ has $\binom{n-\sum_{i=1}^{j-1}n_i}{n_j}$ choices. Thus $|\mathcal{M}_{\mathbf{n}}| = \prod_{j=1}^{r}\binom{n-\sum_{i=1}^{j-1}n_i}{n_j} = \frac{n!}{n_1!\cdots n_r!}$. For the expansion, write each term in $\prod_{k=1}^{n}(x_1+\cdots+x_r)$ as a function $\phi : \{1,\dots,n\} \to \{1,\dots,r\}$, giving $\prod_{k=1}^{n} x_{\phi(k)}$. If $m_i = \left|\phi^{-1}(\{i\})\right|$, then the number of such $\phi$ is $\binom{n}{m_1\, m_2\cdots m_r}$; grouping by $(m_1,\dots,m_r)$ gives the identity.
*Structural Commentary*: Multinomial coefficients count labeled partitions of positions.
*Key Insight*: Group product terms by how many times each label is chosen. □

**Example 3.2.12.** *Repeated Letters, Committees, and Pairings*

(i) The number of arrangements of the multiset $\{A,A,G,G,G,G,C,C,C\}$ is $\binom{9}{2\,4\,3} = 9!/(2!4!3!)$.
(ii) A committee of seven split into labeled subcommittees of sizes $3,2,2$ can be done in $\binom{7}{3\,2\,2} = 210$ ways.
(iii) If $2m$ people are paired without assigning the pairs to labeled positions, the count is $(2m)!/(m!2^m)$.

## 3.3 Inclusion-Exclusion and Combinatorial Proof

**Proposition 3.3.1.** *Finite Inclusion-Exclusion for Two and Three Sets*
*Let $A, B, C$ be finite sets, and write $\mathbf{1}_E(x)$ for membership in $E$.*

*(i)* $|A \cup B| = |A| + |B| - |A \cap B|$.
*(ii)* $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$.

*Proof.*

(i) Let $U = A \cup B$. For $x \in U$, $\mathbf{1}_U(x) = \mathbf{1}_A(x) + \mathbf{1}_B(x) - \mathbf{1}_{A\cap B}(x)$. Sum over $U$; $|U| = |A| + |B| - |A \cap B|$.
(ii) Let $U = A \cup B \cup C$ and $m(x) := \mathbf{1}_A(x) + \mathbf{1}_B(x) + \mathbf{1}_C(x)$.
For $x \in U$, $m(x) \in \{1,2,3\}$ and $m(x) - \binom{m(x)}{2} + \binom{m(x)}{3} = 1$.
Thus each $x \in U$ contributes once to $|A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$, QED.

*Structural Commentary*: Inclusion-exclusion weights each membership pattern to total weight 1.
*Key Insight*: Add single memberships, subtract overcounts, then restore triple overlaps. □

**Method 3.3.2.** *Combinatorial Proof by Classification*
To prove $N_L = N_R$ for finite counts:

(i) choose a finite object set $\Omega$ and a pattern map $p : \Omega \to P$;
(ii) write both sides as sums over fibers, $N_L = \sum_{\alpha\in P} w_L(\alpha)\left|p^{-1}(\{\alpha\})\right|$ and $N_R = \sum_{\alpha\in P} w_R(\alpha)\left|p^{-1}(\{\alpha\})\right|$;
(iii) verify $w_L(\alpha) = w_R(\alpha)$ for every $\alpha \in P$.

## 3.4 Finite Sums

**Definition 3.4.1.** *Finite Sums Indexed by Integers*

Let $a_i \in \mathbb{C}$ be assigned for integers $m \le i \le n$. Define $\sum_{i=m}^{n} a_i := a_m + a_{m+1} + \cdots + a_n$.

If $m > n$, the sum over the empty index set is 0.

**Proposition 3.4.2.** *Basic Laws for Finite Sums*

*Let $a_i, b_i \in \mathbb{C}$ and $c \in \mathbb{C}$ be assigned for $m \le i \le n$.*

*(i) If $m \le r < n$, then $\sum_{i=m}^{n} a_i = \sum_{i=m}^{r} a_i + \sum_{i=r+1}^{n} a_i$.*

*(ii) $\sum_{i=m}^{n} (a_i + b_i) = \sum_{i=m}^{n} a_i + \sum_{i=m}^{n} b_i$ and $\sum_{i=m}^{n} c a_i = c \sum_{i=m}^{n} a_i$.*

*(iii) $\left| \sum_{i=m}^{n} a_i \right| \le \sum_{i=m}^{n} |a_i|$.*

*(iv) If $a_i, b_i \in \mathbb{R}$ and $a_i \le b_i$ for every $i$, then $\sum_{i=m}^{n} a_i \le \sum_{i=m}^{n} b_i$.*

*Proof.*

(i) By Definition 3.4.1 and associativity of addition in $\mathbb{C}$, $\sum_{i=m}^{n} a_i = \sum_{i=m}^{r} a_i + \sum_{i=r+1}^{n} a_i$.

(ii) This follows from associativity, commutativity, and distributivity in the field $\mathbb{C}$.

(iii) Induct on $n - m$. The case $n = m$ is immediate.
If the claim holds up to $n$, then the triangle inequality gives

$$\left| \sum_{i=m}^{n+1} a_i \right| \le \left| \sum_{i=m}^{n} a_i \right| + |a_{n+1}| \le \sum_{i=m}^{n} |a_i| + |a_{n+1}| = \sum_{i=m}^{n+1} |a_i| .$$

(iv) Induct on $n - m$. The case $n = m$ is the hypothesis $a_m \le b_m$.
If the claim holds up to $n$, then compatibility of the order on $\mathbb{R}$ with addition gives

$$\sum_{i=m}^{n+1} a_i = \sum_{i=m}^{n} a_i + a_{n+1} \le \sum_{i=m}^{n} b_i + b_{n+1} = \sum_{i=m}^{n+1} b_i .$$

*Structural Commentary*: Finite sums are iterated addition, so their laws are inherited from the scalar field.
*Key Insight*: No convergence hypothesis is needed for a finite sum. □

# 4 Relations, Orders, and Lattices

Relations package comparisons, identifications, order structures, and quotient constructions.

## 4.1 Relations and Equivalence

**Definition 4.1.1.** *Relations*

Let $X, Y$ be sets. A *relation* from $X$ to $Y$ is a subset $R \subseteq X \times Y$; write $xRy$ iff $(x, y) \in R$.

**Definition 4.1.2.** *Equivalence Relations*

Let $X$ be a set. A relation $\sim$ on $X$ is an *equivalence relation* iff, for all $x, y, z \in X$,
$x \sim x$, $x \sim y \Rightarrow y \sim x$, and $(x \sim y \wedge y \sim z) \Rightarrow x \sim z$.

**Definition 4.1.3.** *Equivalence Classes, Invariants, and Canonical Forms*

Let $X, Y$ be sets.

(i) If $\sim$ is an equivalence relation on $X$, the equivalence class of $x \in X$ is $[x]_\sim := \{y \in X : y \sim x\}$.
(ii) A map $f : X \to Y$ is an *invariant* of $\sim$ iff $x \sim y \Rightarrow f(x) = f(y)$ for all $x, y \in X$.
(iii) A map $f : X \to Y$ is a *complete invariant* of $\sim$ iff $x \sim y \iff f(x) = f(y)$ for all $x, y \in X$.
(iv) A subset $\mathcal{C} \subseteq X$ is a *set of canonical forms* for $\sim$ iff $|\mathcal{C} \cap [x]_\sim| = 1$ for every $x \in X$.

## 4.2 Partitions and Quotient Sets

**Proposition 4.2.1.** *Equivalence Classes Partition the Set*

*Let $\sim$ be an equivalence relation on $X$. Then the equivalence classes form a partition of $X$.*

*Proof.* Every $x \in X$ lies in $[x]$ since $x \sim x$. If $[x] \cap [y] \neq \emptyset$, choose $z$ in the intersection.
Then $x \sim z \sim y$, so $u \in [x]$ implies $u \sim y$ and $u \in [y]$. Hence $[x] \subseteq [y]$, and symmetry gives equality.
*Structural Commentary*: Equivalence means partition.
*Key Insight*: Touching classes coincide. □




Figure 3: Equivalence classes form disjoint blocks that cover the underlying set.

**Proposition 4.2.2.** *Partitions Determine Equivalence Relations*
*Let $\mathcal{F}$ be a partition of $X$. For $x \in X$, let $B_x \in \mathcal{F}$ be the unique block with $x \in B_x$.*
*Define a relation $\sim_{\mathcal{F}}$ on $X$ by $x \sim_{\mathcal{F}} y$ iff $B_x = B_y$.*

*(i) $\sim_{\mathcal{F}}$ is an equivalence relation on $X$.*
*(ii) For every $x \in X$, $[x]_{\sim_{\mathcal{F}}} = B_x$; hence the equivalence classes are exactly the members of $\mathcal{F}$.*

*Proof.*

(i) For $x \in X$, $B_x = B_x$, so $x \sim_{\mathcal{F}} x$. If $x \sim_{\mathcal{F}} y$, then $B_x = B_y$, hence $B_y = B_x$ and $y \sim_{\mathcal{F}} x$. If $x \sim_{\mathcal{F}} y$ and $y \sim_{\mathcal{F}} z$, then $B_x = B_y = B_z$, hence $x \sim_{\mathcal{F}} z$.
(ii) Fix $x \in X$. If $y \in [x]_{\sim_{\mathcal{F}}}$, then $B_y = B_x$, so $y \in B_x$. If $y \in B_x$, then the unique block containing $y$ is $B_x$, so $B_y = B_x$ and $y \in [x]_{\sim_{\mathcal{F}}}$. Thus $[x]_{\sim_{\mathcal{F}}} = B_x$. Conversely, if $B \in \mathcal{F}$, choose $x \in B$; then $B = B_x = [x]_{\sim_{\mathcal{F}}}$.

*Structural Commentary*: A partition is the data of its block-membership relation.
*Key Insight*: Transitivity is equality of blocks. □

**Example 4.2.3.** *Rational Equivalence on $\mathbb{R}$*
On $\mathbb{R}$, define $x \sim_{\mathbb{Q}} y$ iff $x - y \in \mathbb{Q}$.

(i) Reflexive: $x - x = 0 \in \mathbb{Q}$.
(ii) Symmetric: $x - y \in \mathbb{Q}$ implies $y - x = -(x - y) \in \mathbb{Q}$.
(iii) Transitive: $x - y, y - z \in \mathbb{Q}$ imply $x - z = (x - y) + (y - z) \in \mathbb{Q}$.

Thus $\sim_{\mathbb{Q}}$ is an equivalence relation, and $[x]_{\sim_{\mathbb{Q}}} = x + \mathbb{Q} = \{x + q : q \in \mathbb{Q}\}$.
A set $R \subseteq \mathbb{R}$ is a representative set iff $|R \cap (x + \mathbb{Q})| = 1$ for every $x \in \mathbb{R}$.

**Definition 4.2.4.** *Quotient Set*
Let $\sim$ be an equivalence relation on $X$. The *quotient set* is $X/\!\sim := \{[x]_\sim : x \in X\}$.
A map $g : X/\!\sim \to Y$ defined by representatives, $g([x]_\sim) := f(x)$, is well-defined iff $x \sim y$ implies $f(x) = f(y)$.

## 4.3 Ordered Sets, Well-Orders, and Lattices

**Definition 4.3.1.** *Ordered Set Vocabulary*
Let $S$ be a set, let $E, C \subseteq S$, and let $\leq_S$ be a relation on $S$. Write $x <_S y$ iff $x \leq_S y$ and $x \neq y$.

(i) $\leq_S$ is a *partial order* if it is reflexive, antisymmetric, and transitive; then $(S, \leq_S)$ is a *poset*.
(ii) $\leq_S$ is an *order* if it is a partial order and any $x, y \in S$ are comparable; then $(S, \leq_S)$ is an *ordered set*.
(iii) $C$ is a *chain* if $\leq_S |_C$ is an order.
(iv) $S$ is *well-ordered* if every nonempty $Y \subseteq S$ has $\ell \in Y$ with $\ell \leq_S y$ for all $y \in Y$.
(v) $\beta \in S$ is an *upper bound* of $E$ if $x \leq_S \beta$ for all $x \in E$; lower bounds reverse the inequality.
(vi) $E$ is *bounded above* if it has an upper bound; bounded below is dual.
(vii) $\sup E$ is the least upper bound of $E$, and $\inf E$ is the greatest lower bound of $E$, when these exist.
(viii) $m \in E$ is *maximal* if no $x \in E$ satisfies $m <_S x$; minimal is dual.
(ix) $S$ has the *least-upper-bound property* if every nonempty bounded-above $E \subseteq S$ has $\sup E \in S$.




Figure 4: The supremum is the least upper bound, while the infimum is the greatest lower bound.

**Example 4.3.2.** *Suprema Need Not Be Elements*
In $\mathbb{Q}$, the set $E_1 = \{r \in \mathbb{Q} : r < 0\}$ and the set $E_2 = \{r \in \mathbb{Q} : r \leq 0\}$ both have supremum 0, but $0 \notin E_1$ and $0 \in E_2$. If $E = \{1/n : n = 1, 2, 3, \ldots\}$, then $\sup E = 1$ while $\inf E = 0$, and the infimum is not a member of $E$.

**Theorem 4.3.3.** *Least Upper Bounds Give Greatest Lower Bounds*
*If $S$ has the least-upper-bound property, then every nonempty bounded-below set $B \subseteq S$ has an infimum.*

*Proof.* Let $L$ be the set of lower bounds of $B$. Then $L$ is nonempty and every $x \in B$ is an upper bound of $L$, so $\alpha = \sup L$ exists. Since every $x \in B$ bounds $L$ above, $\alpha \leq x$ for all $x \in B$; hence $\alpha$ is a lower bound of $B$. If $\beta > \alpha$, then $\beta \notin L$, so $\beta$ is not a lower bound. Thus $\alpha = \inf B$.
*Structural Commentary*: Infima are suprema of lower bounds.
*Key Insight*: Completeness above gives completeness below. □

**Definition 4.3.4.** *Lattice Vocabulary*
Let $(L, \leq)$ be a partially ordered set.

(i) A *meet* of $x, y \in L$ is $\inf\{x, y\}$ when it exists, often written $x \wedge y$.
(ii) A *join* of $x, y \in L$ is $\sup\{x, y\}$ when it exists, often written $x \vee y$.
(iii) A *lattice* is a partially ordered set in which every two elements have a meet and a join.
(iv) A lattice is *complete* if every subset $E \subseteq L$ has an infimum and a supremum.
(v) A *bottom* element is an element $\bot \in L$ with $\bot \leq x$ for every $x \in L$.
A *top* element is an element $\top \in L$ with $x \leq \top$ for every $x \in L$.
(vi) A lattice is *modular* if $x \leq z$ implies $z \wedge (y \vee x) = (z \wedge y) \vee x$ for all $x, y, z \in L$.

## 4.4 Relations Between Sets and Closures

*Placeholder*: Add relations between different sets, inverse relations, relation composition, reflexive/transitive closure, and order-isomorphism examples.

# 5 Functions

Functions are relations with unique outputs; their images, preimages, restrictions, compositions, and inverses organize nearly every later construction.

## 5.1 Functions, Images, and Preimages

**Definition 5.1.1.** *Functions, Images, and Preimages*
Let $f : X \to Y$ be a *function*, assigning each $x \in X$ a unique $f(x) \in Y$.

(i) $X = \mathrm{dom}(f)$ is the *domain* and $Y$ is the *codomain*;
(ii) $\mathrm{range}(f) := f(X) := \{f(x) : x \in X\}$ is the *range* of $f$;
(iii) for $A \subseteq X$, $f(A) := \{f(x) : x \in A\}$ is the *image* of $A$;
(iv) for $B \subseteq Y$, $f^{-1}(B) := \{x \in X : f(x) \in B\}$ is the *preimage* of $B$; no inverse function is required.

**Definition 5.1.2.** *Support of a Function*
Let $f : X \to Y$. If $Y$ has distinguished 0, $\mathrm{supp}(f) := \{x \in X : f(x) \neq 0\}$.




Figure 5: Image–preimage containment: $f(A) \subseteq B \iff A \subseteq f^{-1}(B)$.

**Proposition 5.1.3.** *Image and Preimage Calculus*
*Let $f : X \to Y$, let $I \neq \emptyset$, and let $A_i \subseteq X$ and $B_i \subseteq Y$ for each $i \in I$.*

*(i)* $f\left(\bigcup_{i\in I} A_i\right) = \bigcup_{i\in I} f(A_i)$.
*(ii)* $f\left(\bigcap_{i\in I} A_i\right) \subseteq \bigcap_{i\in I} f(A_i)$.
*(iii)* $f^{-1}\left(\bigcup_{i\in I} B_i\right) = \bigcup_{i\in I} f^{-1}(B_i)$.
*(iv)* $f^{-1}\left(\bigcap_{i\in I} B_i\right) = \bigcap_{i\in I} f^{-1}(B_i)$.
*(v) For every $B \subseteq Y$, $f^{-1}(Y \setminus B) = X \setminus f^{-1}(B)$.*
*(vi) For all $B, C \subseteq Y$, $f^{-1}(B \setminus C) = f^{-1}(B) \setminus f^{-1}(C)$.*
*(vii) For every $B \subseteq Y$, $f(f^{-1}(B)) = B \cap \mathrm{range}(f)$.*

*Proof.*

(i) For $y \in Y$, $y \in f\left(\bigcup_{i\in I} A_i\right) \iff \exists i \in I,\ \exists x \in A_i,\ y = f(x) \iff y \in \bigcup_{i\in I} f(A_i)$.

(ii) If $y \in f\left(\bigcap_{i\in I} A_i\right)$, then $\exists x \in \bigcap_{i\in I} A_i$ such that $y = f(x)$. Hence $x \in A_i$ for every $i \in I$, so $y \in \bigcap_{i\in I} f(A_i)$.

(iii) For $x \in X$, $x \in f^{-1}\left(\bigcup_{i\in I} B_i\right) \iff \exists i \in I,\ f(x) \in B_i \iff x \in \bigcup_{i\in I} f^{-1}(B_i)$.

(iv) For $x \in X$, $x \in f^{-1}\left(\bigcap_{i\in I} B_i\right) \iff \forall i \in I,\ f(x) \in B_i \iff x \in \bigcap_{i\in I} f^{-1}(B_i)$.

(v) For $x \in X$, $x \in f^{-1}(Y \setminus B) \iff f(x) \notin B \iff x \notin f^{-1}(B) \iff x \in X \setminus f^{-1}(B)$.

(vi) For $x \in X$, $x \in f^{-1}(B \setminus C) \iff f(x) \in B,\ f(x) \notin C \iff x \in f^{-1}(B) \setminus f^{-1}(C)$.

(vii) For $y \in Y$, $y \in f(f^{-1}(B)) \iff \exists x \in X,\ f(x) \in B,\ y = f(x) \iff y \in B \cap \mathrm{range}(f)$.

*Structural Commentary*: Preimages preserve Boolean operations.
*Key Insight*: Membership pulls back through $f$. □

## 5.2 Injective, Surjective, and Bijective Maps

**Definition 5.2.1.** *Injective, Surjective, and Bijective Maps*
Let $f : X \to Y$.

i. $f$ is *injective* if $f(x_1) = f(x_2)$ implies $x_1 = x_2$.
ii. $f$ is *surjective* if $\mathrm{range}(f) = Y$.
iii. $f$ is *bijective* if it is both injective and surjective.




Figure 6: Surjective: covers codomain; injective: no collisions; bijective: both.

**Example 5.2.2.** *Image and Preimage for the Square Map*
Let $f : \mathbb{R} \to \mathbb{R}$ be given by $f(x) = x^2$. Then $f([-2,1]) = [0,4]$, $f^{-1}([1,4]) = [-2,-1] \cup [1,2]$. The map is not injective because $f(1) = f(-1)$, and it is not surjective as a map $\mathbb{R} \to \mathbb{R}$ because no real number maps to $-1$.

## 5.3 Composition, Restriction, and Inverse Functions

**Definition 5.3.1.** *Composition, Restriction, Extension, and Identity*
Let $f : X \to Y$.

(i) for $A \subseteq X$, $f|_A : A \to Y$ is the *restriction* of $f$ to $A$;
(ii) if $X \subseteq U$, an *extension* of $f$ to $U$ is $\widetilde{f} : U \to Y$ with $\widetilde{f}|_X = f$;
(iii) if $g : C \to Z$ and $f(X) \subseteq C$, then $g \circ f : X \to Z$ is given by $(g \circ f)(x) = g(f(x))$;
(iv) $\mathrm{id}_X : X \to X$ denotes the *identity function*.

**Theorem 5.3.2.** *Bijective Means Invertible*
*A function $f : X \to Y$ is bijective iff there exists a function $g : Y \to X$ such that $g \circ f = \mathrm{id}_X$, $f \circ g = \mathrm{id}_Y$. When such a function exists, it is unique and is denoted $f^{-1} : Y \to X$.*

*Proof.*

($\Rightarrow$) If $f$ is bijective, define $g(y)$ as the unique $x$ with $f(x) = y$. Then $g \circ f = \mathrm{id}_X$ and $f \circ g = \mathrm{id}_Y$.
($\Leftarrow$) If such a $g$ exists, applying $g$ to $f(x_1) = f(x_2)$ gives $x_1 = x_2$, and $y = f(g(y))$ gives surjectivity.

If $g$ and $h$ are two-sided inverses, then $g(y) = g(f(h(y))) = h(y)$, so the inverse is unique.
*Structural Commentary*: A true inverse undoes $f$ on both sides.
*Key Insight*: Bijective means exactly one preimage per output. □

**Proposition 5.3.3.** *Composition and Restriction Calculus*
*Let $f : X \to Y$, $g : Y \to Z$, and $h : Z \to W$.*

*(i)* $h \circ (g \circ f) = (h \circ g) \circ f$.
*(ii)* $\mathrm{id}_Y \circ f = f$ *and* $f \circ \mathrm{id}_X = f$.
*(iii) For every* $A \subseteq X$, $(g \circ f)(A) = g(f(A))$.
*(iv) For every* $C \subseteq Z$, $(g \circ f)^{-1}(C) = f^{-1}(g^{-1}(C))$.
*(v) For every* $A \subseteq X$, $(g \circ f)|_A = g \circ (f|_A)$.
*(vi) If $f$ and $g$ are bijective, then $g \circ f$ is bijective and* $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$.

*Proof.*

(i) For $x \in X$, $(h \circ (g \circ f))(x) = h(g(f(x))) = ((h \circ g) \circ f)(x)$.
(ii) For $x \in X$, $(\mathrm{id}_Y \circ f)(x) = f(x)$ and $(f \circ \mathrm{id}_X)(x) = f(x)$.
(iii) For $z \in Z$, $z \in (g \circ f)(A) \iff \exists x \in A,\ z = g(f(x)) \iff z \in g(f(A))$.
(iv) For $x \in X$, $x \in (g \circ f)^{-1}(C) \iff g(f(x)) \in C \iff f(x) \in g^{-1}(C) \iff x \in f^{-1}(g^{-1}(C))$.
(v) For $x \in A$, $((g \circ f)|_A)(x) = g(f(x)) = (g \circ (f|_A))(x)$.
(vi) If $f$ and $g$ are bijective, then $f^{-1} : Y \to X$ and $g^{-1} : Z \to Y$ exist. For $x \in X$, $(f^{-1} \circ g^{-1})((g \circ f)(x)) = x$; for $z \in Z$, $(g \circ f)((f^{-1} \circ g^{-1})(z)) = z$. Thus $f^{-1} \circ g^{-1}$ is the inverse of $g \circ f$.

*Structural Commentary*: Composition is pointwise; inverses reverse order.
*Key Insight*: Evaluate at an element. □

# 6 Cardinality, Choice, and Infinite Sets

Cardinality compares sets by functions rather than by informal size.

## 6.1 Same Cardinality, Finite, Countable, and Uncountable

**Definition 6.1.1.** *Same Cardinality*
For sets $A, B$, define $A \sim_{\#} B$ iff there exists a bijection $f : A \to B$.
Then $A$ and $B$ have *same cardinality*, or are *equivalent*. The relation $\sim_{\#}$ is an equivalence relation.
$\mathrm{id}_A$ proves reflexivity, inverses of bijections prove symmetry, and composition of bijections proves transitivity.

**Definition 6.1.2.** *Finite, Countable, and Uncountable*
Set $J_0 := \emptyset$, and for $n \geq 1$ set $J_n := \{1, 2, \ldots, n\}$. A set $A$ is

(i) *finite* if $A \sim_{\#} J_n$ for some $n \in \mathbb{N}$;
(ii) *infinite* if it is not finite;
(iii) *countable* if $A \sim_{\#} \mathbb{N}$;

(iv) *uncountable* if it is neither finite nor countable;
(v) *at most countable* if it is finite or countable.

Countable sets are also called enumerable or denumerable.

**Proposition 6.1.3.** *Finite Cardinality is Well Defined*
*Let $m, n \in \mathbb{N}$. If $A \sim_{\#} J_m$ and $A \sim_{\#} J_n$, then $m = n$.*
*Hence for finite $A$, the value $\#A := n$ with $A \sim_{\#} J_n$ is well-defined.*

*Proof.* By transitivity of $\sim_{\#}$, it suffices to prove $J_m \sim_{\#} J_n \Rightarrow m = n$.
Induct on $m$. If $m = 0$, then $J_m = \emptyset$, so any bijection $J_m \to J_n$ forces $J_n = \emptyset$ and $n = 0$.
Assume the claim for $m - 1$, with $m \geq 1$, and let $f : J_m \to J_n$ be bijective. Then $n \geq 1$.
Removing $m$ and $f(m)$ gives a bijection $J_m \setminus \{m\} \to J_n \setminus \{f(m)\}$. Since these sets have the standard relabelings $J_{m-1}$ and $J_{n-1}$, the induction hypothesis gives $m - 1 = n - 1$, hence $m = n$.
*Structural Commentary*: Finite size is bijection-invariant.
*Key Insight*: Remove one matched element. □

**Definition 6.1.4.** *Summation over a Finite Set*
Let $X$ be a finite set with $\#X = n$, and let $f : X \to \mathbb{C}$.
Choose any bijection $g : J_n \to X$, where $J_n = \{1, \ldots, n\}$, and set $\sum_{x \in X} f(x) := \sum_{i=1}^{n} f(g(i))$.

When $X = \emptyset$, this is the empty sum 0.
*Key Insight*: Proposition 6.1.5 shows that the value is independent of the chosen bijection.

**Proposition 6.1.5.** *Finite Set Summation Laws*
*Let $X, Y$ be finite sets.*

*(i) Well-definedness: different bijections $J_{\#X} \to X$ give the same value in Definition 6.1.4.*
*(ii) Substitution: if $h : Y \to X$ is a bijection and $f : X \to \mathbb{C}$, then $\sum_{x \in X} f(x) = \sum_{y \in Y} f(h(y))$.*
*(iii) Linearity: for $f, g : X \to \mathbb{C}$ and $c \in \mathbb{C}$, $\sum_{x \in X} (f(x)+g(x)) = \sum_{x \in X} f(x) + \sum_{x \in X} g(x)$ and $\sum_{x \in X} cf(x) = c \sum_{x \in X} f(x)$.*
*(iv) Disjoint unions: if $X \cap Y = \emptyset$ and $f : X \cup Y \to \mathbb{C}$, then $\sum_{z \in X \cup Y} f(z) = \sum_{x \in X} f(x) + \sum_{y \in Y} f(y)$.*
*(v) Finite Fubini: if $f : X \times Y \to \mathbb{C}$, then $\sum_{x \in X} \sum_{y \in Y} f(x, y) = \sum_{(x,y) \in X \times Y} f(x, y) = \sum_{y \in Y} \sum_{x \in X} f(x, y)$.*

*Proof.*

(i) For $f : X \to \mathbb{C}$, let $n = \#X$ and let $g, k : J_n \to X$ be bijections. Induct on $n$. The case $n = 0$ is empty. If $n \geq 1$, let $j \in J_n$ satisfy $k(j) = g(n)$. Removing $n$ from $g$ and $j$ from $k$ gives bijections onto $X \setminus \{g(n)\}$; after relabeling by $J_{n-1}$, the induction hypothesis makes the remaining sums equal. Adding the common last term $f(g(n)) = f(k(j))$ gives equality of the two sums.

(ii) Let $g : J_{\#Y} \to Y$ be a bijection. Then $h \circ g : J_{\#Y} \to X$ is a bijection, so $\#Y = \#X$ by Proposition 6.1.3. Hence, by (i) and Definition 6.1.4, $\sum_{x \in X} f(x) = \sum_{i=1}^{\#Y} f(h(g(i))) = \sum_{y \in Y} f(h(y))$.

(iii) Choose a bijection $\gamma : J_{\#X} \to X$.
Proposition 3.4.2 gives $\sum_{x \in X} (f(x) + g(x)) = \sum_{x \in X} f(x) + \sum_{x \in X} g(x)$ and $\sum_{x \in X} cf(x) = c \sum_{x \in X} f(x)$

(iv) Let $\alpha : J_{\#X} \to X$ and $\beta : J_{\#Y} \to Y$ be bijections, and enumerate $X \cup Y$ by listing $\alpha$ first and then $\beta$. Since $X \cap Y = \emptyset$, this is a bijection, and Proposition 3.4.2 gives $\sum_{z \in X \cup Y} f(z) = \sum_{x \in X} f(x) + \sum_{y \in Y} f(y)$.

(v) Let $\alpha : J_{\#X} \to X$ and $\beta : J_{\#Y} \to Y$ be bijections.
The maps $\ell((i-1)\#Y + j) = (\alpha(i), \beta(j))$ and $r((j-1)\#X + i) = (\alpha(i), \beta(j))$ for $i \in J_{\#X}$ and $j \in J_{\#Y}$ give the two lexicographic bijections $J_{(\#X)(\#Y)} \to X \times Y$.
Applying Definition 6.1.4, Proposition 3.4.2, and (i) yields $\sum_{x \in X} \sum_{y \in Y} f(x, y) = \sum_{(x,y) \in X \times Y} f(x, y) = \sum_{y \in Y} \sum_{x \in X} f(x, y)$.

*Structural Commentary*: A finite set sum is an order-free finite sum.
*Key Insight*: Bijections are bookkeeping devices; the value does not depend on the bookkeeping order. □

**Definition 6.1.6.** *Sequences as Enumerations*
Let $A$ be a set.

(i) A *sequence* in $A$ is a map $x : \mathbb{N} \to A$, written $(x_n)_{n \in \mathbb{N}}$ with $x_n := x(n)$.
(ii) A sequence $x : \mathbb{N} \to A$ *enumerates* $A$ iff $x(\mathbb{N}) = A$.
(iii) A sequence $x : \mathbb{N} \to A$ is a *distinct enumeration* of $A$ iff $x : \mathbb{N} \to A$ is bijective.

Thus $A$ is countable iff it has a distinct enumeration.

**Example 6.1.7.** *The Integers Are Countable*
The set $\mathbb{Z}$ is countable. One explicit enumeration is given by

$$f(n) = \begin{cases} -\frac{n}{2}, & n \text{ even}, \\ \frac{n+1}{2}, & n \text{ odd}, \end{cases} \qquad n \in \mathbb{N}.$$

This lists $\mathbb{Z}$ as $0, 1, -1, 2, -2, 3, -3, \ldots$.

**Theorem 6.1.8.** *Infinite Subsets of Countable Sets*
*Let $A$ be countable and let $E \subseteq A$ be infinite. Then $E$ is countable.*

*Proof.* Let $x : \mathbb{N} \to A$ be a distinct enumeration of $A$. Since $E$ is infinite, $I_E := \{n \in \mathbb{N} : x_n \in E\}$ is infinite. Write $I_E = \{n_1 < n_2 < \cdots\}$ and define $g : \mathbb{N} \to E$ by $g(k) := x_{n_k}$. The terms $(x_n)$ are distinct, so $g$ is injective. If $y \in E$, then $y = x_n$ for a unique $n \in \mathbb{N}$; since $n \in I_E$, $n = n_k$ for some $k$, and $y = g(k)$.
Thus $g$ is bijective, so $E$ is countable.
*Structural Commentary*: Subsets select indices.
*Key Insight*: Keep the hitting indices. □

**Proposition 6.1.9.** *Subsets and Images of At Most Countable Sets*
*Let $A$ be at most countable.*

*(i) If $B \subseteq A$, then $B$ is at most countable.*
*(ii) If $f : A \to Y$ is a function, then $f(A)$ is at most countable.*

*Proof.* If $A$ is finite, both claims are immediate. Otherwise write $A = \{a_n : n \in \mathbb{N}\}$ with the $a_n$ distinct.

(i) If $B$ is infinite, let $I_B := \{n : a_n \in B\} = \{n_1 < n_2 < \cdots\}$.
Then $k \mapsto a_{n_k}$ bijects $\mathbb{N}$ onto $B$; the finite case is trivial.
(ii) Let $I := \{n : f(a_n) \notin \{f(a_m) : m < n\}\}$. By (i), $I$ is at most countable, $n \mapsto f(a_n)$ bijects $I$ onto $f(A)$.

*Structural Commentary*: Subsets select indices; images select first appearances.
*Key Insight*: Subsets and images preserve at-most-countability. □

**Proposition 6.1.10.** *At Most Countable Sets Are Exactly Enumerable Images*
*For a set $B$, the following are equivalent:*

*(i) $B$ is at most countable.*
*(ii) $B = \emptyset$ or there exists a surjection $f : \mathbb{N} \to B$.*

*Thus a nonempty set is at most countable iff it is the image of a sequence.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) If $B = \emptyset$, (ii) holds. If $B$ is countable, choose a bijection $f : \mathbb{N} \to B$. If $B = \{b_1, \ldots, b_m\}$ is finite and nonempty, define $f(0) = b_1$, $f(n) = b_n$ for $1 \le n \le m$, and $f(n) = b_m$ for $n > m$.
In both nonempty cases, $f : \mathbb{N} \to B$ is surjective.
(*ii*) $\Rightarrow$ (*i*) If $B = \emptyset$, then $B$ is finite.
If $f : \mathbb{N} \to B$ is surjective, then $B = f(\mathbb{N})$ is at most countable by Proposition 6.1.9.

*Structural Commentary*: Surjections list with repetition.
*Key Insight*: At most countable means sequence image. □




Figure 7: Diagonal scan of countable rows into one sequence, skipping repeats.

**Theorem 6.1.11.** *Countable Union of Countable Sets*
*Let $(E_n)_{n \ge 1}$ be countable sets and set $S := \bigcup_{n=1}^{\infty} E_n$. Then $S$ is countable.*

*Proof.* For each $n \geq 1$, choose a bijection $x_n : \mathbb{N} \to E_n$ and write $x_{n,k} := x_n(k)$.
Let $D := \{(n,k) : n \geq 1,\ k \in \mathbb{N}\}$. Diagonal order gives a surjection $d : \mathbb{N} \to D$.
Set $a(t) := x_{n,k}$ for $d(t) = (n,k)$; then $a(\mathbb{N}) = S$, so $S$ is at most countable by Proposition 6.1.10.
Since $E_1 \subseteq S$ and $E_1$ is countable, $S$ is infinite. Hence $S$ is countable.
*Structural Commentary*: Diagonal scanning enumerates pairs.
*Key Insight*: Scan the two-index list. □

**Corollary 6.1.12.** *At Most Countable Unions*
*If $A$ is at most countable and $B_\alpha$ is at most countable for each $\alpha \in A$, then $\bigcup_{\alpha \in A} B_\alpha$ is at most countable.*

*Proof.* Let $T := \bigcup_{\alpha \in A} B_\alpha$. If $T = \emptyset$, we are done.
Otherwise write the nonempty $B_\alpha$ as $(B_{\alpha_j})_{j \geq 1}$, with repetitions if needed, and enumerate $B_{\alpha_j}$ by $(b_{jk})_{k \in \mathbb{N}}$.
A diagonal scan of $(b_{jk})$ has image $T$, so Proposition 6.1.9 makes $T$ at most countable.
*Structural Commentary*: Finite pieces cause no difficulty; they can be embedded into countable rows.
*Key Insight*: At most countable indexing + at most countable indexed sets gives at most countable total set. □

**Theorem 6.1.13.** *Finite Tuples over a Countable Set*
*Let $A$ be countable. For each fixed $n \in \mathbb{N}$ with $n \geq 1$, the set of all $n$-tuples $(a_1, \ldots, a_n)$ with $a_k \in A$ is countable.*

*Proof.* Let $B_n := \{(a_1, \ldots, a_n) : a_k \in A\}$, $B_1 = A$. If $B_{n-1}$ is countable, $B_n = \bigcup_{b \in B_{n-1}} \{(b,a) : a \in A\}$, each set $\{(b,a) : a \in A\}$ is equivalent to $A$. Hence $B_n$ is countable by Theorem 6.1.11.
*Structural Commentary*: Build an $n$-tuple by appending one coordinate to an $(n-1)$-tuple.
*Key Insight*: Induction plus countable unions controls finite products. □




Figure 8: Diagonal enumeration of $\mathbb{N} \times \mathbb{N}$ by finite $i + j$ diagonals.

**Corollary 6.1.14.** *Countable Cartesian Products*
*Let $A, B$ be sets.*

*(i) If $A$ and $B$ are at most countable, then $A \times B$ is at most countable.*
*(ii) If $A$ and $B$ are countable and nonempty, then $A \times B$ is countable.*

*In particular, $\mathbb{N} \times \mathbb{N}$ is countable.*

*Proof.*

(i) For $a \in A$, set $S_a := \{a\} \times B$. The map $B \to S_a$, $b \mapsto (a,b)$, is bijective, so each $S_a$ is at most countable. Since $A \times B = \bigcup_{a \in A} S_a$ and $A$ is at most countable, Corollary 6.1.12 gives that $A \times B$ is at most countable.

(ii) By (i), $A \times B$ is at most countable. Choose $b_0 \in B$. The map $A \to A \times B$, $a \mapsto (a, b_0)$, is injective; since $A$ is infinite, $A \times B$ is infinite. Hence $A \times B$ is countable. Taking $A = B = \mathbb{N}$ gives $\mathbb{N} \times \mathbb{N}$ countable.

*Structural Commentary*: Products decompose into slices.
*Key Insight*: Count slices, then fix a coordinate. □

**Corollary 6.1.15.** *The Rationals Are Countable*
*The set $\mathbb{Q}$ is countable.*

*Proof.* Let $D := \{(p,q) \in \mathbb{Z} \times \mathbb{Z} : q \neq 0\}$ and define $\pi : D \to \mathbb{Q}$ by $\pi(p,q) := p/q$.
Since $D \subseteq \mathbb{Z} \times \mathbb{Z}$, Corollary 6.1.14 and Proposition 6.1.9 make $D$ at most countable.
Since $\pi(D) = \mathbb{Q}$, $\mathbb{Q}$ is at most countable. The injection $\mathbb{Z} \to \mathbb{Q}$, $n \mapsto n$, shows $\mathbb{Q}$ is infinite; hence $\mathbb{Q}$ is countable.
*Structural Commentary*: Fractions are images of integer pairs.
*Key Insight*: Count pairs, then quotient. □

**Method 6.1.16.** *Cardinality Proof Patterns*

(i) *Same cardinality*: To prove $|A| = |B|$, construct a bijection $f : A \to B$.
(ii) *Countable*: To prove $A$ is countable, construct a bijection $f : \mathbb{N} \to A$.

(iii) *Uncountable*: To prove $A$ is uncountable, for arbitrary $s : \mathbb{N} \to A$ construct $a_s \in A \setminus s[\mathbb{N}]$.

*Key Idea*: Maps certify cardinality; witnesses $a_s \in A \setminus s[\mathbb{N}]$ refute enumerations.

## 6.2 Diagonalization and Power Sets

**Theorem 6.2.1.** *Cantor's Theorem*
*Let $X$ be a set. No map $f : X \to \mathcal{P}(X)$ is surjective. Hence $X \not\sim_{\#} \mathcal{P}(X)$.*

*Proof.* Fix $f : X \to \mathcal{P}(X)$ and set $D_f := \{x \in X : x \notin f(x)\}$.
If $f$ were surjective, then $D_f = f(a)$ for some $a \in X$. Hence $a \in D_f \iff a \notin f(a) \iff a \notin D_f$, a contradiction.
Thus no $f : X \to \mathcal{P}(X)$ is surjective, so no bijection $X \to \mathcal{P}(X)$ exists.
*Structural Commentary*: The diagonal subset escapes $f$.
*Key Insight*: Flip membership at $x$. □

**Theorem 6.2.2.** *Binary Sequences Are Uncountable*
*Let $A$ be the set of all sequences whose terms are the digits 0 and 1. Then $A$ is uncountable.*

*Proof.* Suppose $A$ were countable, $A = \{s_1, s_2, \ldots\}$ with $s_n = (s_n(k))_{k\in\mathbb{N}}$. Define $s \in A$ by $s(k) := 1 - s_k(k)$.
Then $s(n) \neq s_n(n)$ for every $n$, so $s \neq s_n$ for all $n$. Hence $s \notin A$, a contradiction. Thus $A$ is uncountable.
*Structural Commentary*: The construction attacks the diagonal of a proposed enumeration.
*Key Insight*: A sequence can be built to escape every row of a countable list. □

$$
\begin{array}{lcl}
s_1 & = & \boxed{a_{11}}\ a_{12}\ a_{13}\ a_{14}\ a_{15} \\
s_2 & = & a_{21}\ \boxed{a_{22}}\ a_{23}\ a_{24}\ a_{25} \\
s_3 & = & a_{31}\ a_{32}\ \boxed{a_{33}}\ a_{34}\ a_{35}\ \cdots \\
s_4 & = & a_{41}\ a_{42}\ a_{43}\ \boxed{a_{44}}\ a_{45} \\
s_5 & = & a_{51}\ a_{52}\ a_{53}\ a_{54}\ \boxed{a_{55}}
\end{array}
$$

Figure 9: Cantor diagonalization: $s(n) = 1 - a_{nn}$ differs from row $s_n$ at position $n$.

## 6.3 Schroeder–Bernstein and Cardinal Comparison

**Theorem 6.3.1.** *Schroeder–Bernstein Theorem*
*Let $A, B$ be sets, and let $f : A \to B$ and $g : B \to A$ be injections. Then $A \sim_{\#} B$.*

*Proof.* Set $A_0 := A \setminus g(B)$. For $n \geq 0$, define $B_n := f(A_n)$ and $A_{n+1} := g(B_n)$.
Put $U_A := \bigcup_{n\geq 0} A_n$, $U_B := \bigcup_{n\geq 0} B_n$, $A_* := A \setminus U_A$, and $B_* := B \setminus U_B$.
The sets $A_n$ are pairwise disjoint: if $A_i \cap A_j \neq \emptyset$ with $i < j$, then $i = 0$ contradicts $A_j \subseteq g(B)$.
If $i > 0$, injectivity of $g$ and $f$ gives $A_{i-1} \cap A_{j-1} \neq \emptyset$, and iteration reduces to the $i = 0$ contradiction.
Hence the sets $B_n = f(A_n)$ are pairwise disjoint.
The restriction $g_* := g|_{B_*} : B_* \to A_*$ is bijective. Indeed, $g(B_*) \subseteq A_*$; conversely, if $a \in A_*$, then $a \notin A_0$, so $a = g(b)$ for a unique $b \in B$, and $b \notin U_B$, since $b \in B_n$ would imply $a \in A_{n+1}$.
Define $h : A \to B$ by $h(x) := f(x)$ for $x \in U_A$ and $h(x) := g_*^{-1}(x)$ for $x \in A_*$. Since $A = U_A \sqcup A_*$, $B = U_B \sqcup B_*$, $f$ maps each $A_n$ bijectively onto $B_n$, and $g_*^{-1}$ maps $A_*$ bijectively onto $B_*$, the map $h$ is bijective.
*Structural Commentary*: Matched layers plus matched remainder.
*Key Insight*: Use $f$ on layers, $g_*^{-1}$ on the remainder. □

**Definition 6.3.2.** *Cardinal Comparison and Arithmetic*
Let $A, B$ be sets, and let $\kappa = |A|$, $\lambda = |B|$.

(i) *Cardinal comparison*: $|A| \leq |B|$ iff $\exists$ an injection $A \to B$.
(ii) *Strict cardinal comparison*: $|A| < |B|$ iff $|A| \leq |B|$ and $A \not\sim_{\#} B$.
(iii) If $A \cap B = \emptyset$, the *cardinal sum* is $\kappa + \lambda := |A \cup B|$.
(iv) The *cardinal product* is $\kappa\lambda := |A \times B|$.
(v) The *function set* is $A^B := \{f : B \to A\}$, and the *cardinal power* is $\kappa^\lambda := \left|A^B\right|$.

**Definition 6.3.3.** *Continuum Cardinal*
The *continuum cardinal* is $\mathfrak{c} := 2^{\aleph_0} := \left|\{0,1\}^{\mathbb{N}}\right|$. Equivalently, $\mathfrak{c} = |\mathcal{P}(\mathbb{N})|$ via $s \mapsto \{n \in \mathbb{N} : s(n) = 1\}$.

**Perspective 6.3.4.** *Continuum Cardinality*
Theorem 6.2.2 and Theorem 6.2.1 give $\aleph_0 < \mathfrak{c}$.
Thus the continuum is the first cardinal jump produced by binary choice over $\mathbb{N}$.

## 6.4 Cardinal Arithmetic

**Proposition 6.4.1.** *Cardinal Arithmetic Reference Laws*
*Let $\kappa, \lambda, \mu$ be cardinal numbers.*

*(i)* $\kappa + \lambda = \lambda + \kappa$ *and* $\kappa\lambda = \lambda\kappa$;
*(ii)* $\kappa + (\lambda + \mu) = (\kappa + \lambda) + \mu$ *and* $\kappa(\lambda\mu) = (\kappa\lambda)\mu$;
*(iii)* $\kappa(\lambda + \mu) = \kappa\lambda + \kappa\mu$;
*(iv)* $\kappa^{\lambda+\mu} = \kappa^\lambda \kappa^\mu$, $(\kappa^\lambda)^\mu = \kappa^{\lambda\mu}$, *and* $(\kappa\lambda)^\mu = \kappa^\mu \lambda^\mu$.

*Proof.* Choose pairwise disjoint sets $A, B, C$ with $|A| = \kappa$, $|B| = \lambda$, and $|C| = \mu$.

(i) Use $A \cup B \cong B \cup A$ and $A \times B \cong B \times A$.
(ii) Use $A \cup (B \cup C) \cong (A \cup B) \cup C$ and $A \times (B \times C) \cong (A \times B) \times C$.
(iii) Use $A \times (B \cup C) \cong (A \times B) \cup (A \times C)$.
(iv) Use $A^{B \cup C} \cong A^B \times A^C$, $(A^B)^C \cong A^{B \times C}$, and $(A \times B)^C \cong A^C \times B^C$.

Taking cardinalities gives (i)–(iv).
*Structural Commentary*: The laws are canonical bijections.
*Key Insight*: Cardinal arithmetic is set construction up to bijection. □

**Theorem 6.4.2.** *Infinite Cardinal Absorption*
*Let $\kappa, \lambda$ be cardinals and set $\mu := \max\{\kappa, \lambda\}$. Assume $\mu$ is infinite.*

*(i)* $\kappa + \lambda = \mu$.
*(ii) If* $\kappa, \lambda > 0$, *then* $\kappa\lambda = \mu$.

*In particular,* $\aleph_0 + \aleph_0 = \aleph_0 \cdot \aleph_0 = \aleph_0$.

*Proof.* Use the standard infinite square law $\mu + \mu = \mu^2 = \mu$.

(i) Since $\kappa, \lambda \le \mu$, $\mu \le \kappa + \lambda \le \mu + \mu = \mu$, hence $\kappa + \lambda = \mu$.
(ii) Assume $\kappa, \lambda > 0$. Choose nonempty sets $A, B$ with $|A| = \kappa$ and $|B| = \lambda$.
If $\mu = \kappa$, fix $b_0 \in B$ and embed $A$ by $a \mapsto (a, b_0)$; if $\mu = \lambda$, fix $a_0 \in A$ and embed $B$ by $b \mapsto (a_0, b)$.
Thus $\mu \le \kappa\lambda$. Also $\kappa\lambda \le \mu^2 = \mu$, hence $\kappa\lambda = \mu$.

*Structural Commentary*: Infinite cardinals absorb bounded pieces.
*Key Insight*: Squeeze by $\mu^2 = \mu$. □

## 6.5 Choice and Zorn's Lemma

**Definition 6.5.1.** *Choice Function*
Let $\mathcal{F}$ be a set of sets with $F \neq \varnothing$ for all $F \in \mathcal{F}$, and let $(X_i)_{i \in I}$ satisfy $X_i \neq \varnothing$ for all $i \in I$.

(i) A *choice function* on $\mathcal{F}$ is a map $c : \mathcal{F} \to \bigcup_{F \in \mathcal{F}} F$ such that $c(F) \in F$ for all $F \in \mathcal{F}$.
(ii) An *indexed choice function* for $(X_i)_{i \in I}$ is a map $c : I \to \bigcup_{i \in I} X_i$ such that $c(i) \in X_i$ for all $i \in I$.

**Principle 6.5.2.** *Choice Principle*
Let $(X_i)_{i \in I}$ be an indexed family with $X_i \neq \varnothing$ for all $i \in I$.
The principle asserts $\exists c : I \to \bigcup_{i \in I} X_i$ with $c(i) \in X_i$ for all $i \in I$; equivalently, $\prod_{i \in I} X_i \neq \varnothing$.
Construct $c$ when explicit; invoke choice only for arbitrary simultaneous choices.

**Principle 6.5.3.** *Zorn's Lemma*
Let $(X, \le)$ be a partially ordered set with $X \neq \varnothing$. If every chain $C \subseteq X$ has an upper bound $u \in X$, meaning $c \le u$ for all $c \in C$, then there exists $m \in X$ such that $m \le x$ implies $x = m$ for all $x \in X$.

**Definition 6.5.4.** *Inclusion and Containment Orders*
Let $\mathcal{F}$ be a family of sets and let $\mathcal{A} \subseteq \mathcal{F}$.

(i) *Inclusion*: $A \le B$ iff $A \subseteq B$.
An upper bound of $\mathcal{A}$ is $U \in \mathcal{F}$ with $A \subseteq U$ for all $A \in \mathcal{A}$
$M \in \mathcal{F}$ is maximal iff $M \subseteq B \in \mathcal{F}$ implies $B = M$.
(ii) *Reverse inclusion*: $A \le B$ iff $B \subseteq A$.
An upper bound of $\mathcal{A}$ is $U \in \mathcal{F}$ with $U \subseteq A$ for all $A \in \mathcal{A}$;
$M \in \mathcal{F}$ is maximal iff $B \subseteq M$ with $B \in \mathcal{F}$ implies $B = M$.

**Method 6.5.5.** *Zorn Maximality Template*
To prove existence by maximality, encode candidate objects as a poset $(P, \le)$.

(i) Define $P$ so that a maximal $m \in P$ is a desired object. Then Prove $P \neq \varnothing$.
(ii) For an arbitrary chain $C \subseteq P$, construct $u \in P$ with $p \le u$ for all $p \in C$.

(iii) Take a maximal $m \in P$ and verify the target property.

For inclusion orders, the upper-bound candidate is often $\bigcup C$; for reverse-inclusion orders, it is often $\bigcap C$ or another object contained in every member of $C$. The key verification is always membership in $P$.

# 7 Axiomatic and Advanced Set Theory

This section reserves space for the deeper set-theoretic material suggested by the set-theory and metatheory contents: axioms, ordinals, models, forcing, large cardinals, and sets of reals.

## 7.1 Extensionality, Specification, and Replacement

*Placeholder*: Add ZF-style axiom statements, informal motivations, and the separation/replacement distinction.

## 7.2 Ordinals, Transfinite Recursion, and Well-Foundedness

*Placeholder*: Add well-founded relations, transitive sets, ordinal numbers, ordinal arithmetic, transfinite induction, and transfinite recursion.

## 7.3 Models, Constructibility, and Independence

*Placeholder*: Add models of set theory, the constructible universe, forcing previews, consistency strength, and independence statements.

## 7.4 Filters, Ultrafilters, and Sets of Reals

*Placeholder*: Add filters, ideals, ultrafilters, stationary/closed-unbounded sets, Borel and projective sets, descriptive set theory, and large-cardinal gateways.

# Part II
# Structural Linear Algebra

## 1 Vector Spaces and Linear Structure

### 1.1 Vector Spaces

Vector spaces encode addition and scalar multiplication. Subspaces, spans, bases, dimension, and coordinates describe how linear structure is generated and measured.

**Definition 1.1.1.** *Vector Space*
Let $\mathbb{F}$ be a field. A vector space over $\mathbb{F}$ is a set $\mathcal{V}$ with vector addition and scalar multiplication such that, $\forall\, \mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and $\forall\, a, b \in \mathbb{F}$, the following axioms hold:

i. vector addition is commutative: $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$;
ii. vector addition is associative: $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$;
iii. there exists an additive identity $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{0} + \mathbf{v} = \mathbf{v}$;
iv. for each $\mathbf{v} \in \mathcal{V}$ there exists an additive inverse $-\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}$;
v. scalar multiplication has identity: $1\mathbf{v} = \mathbf{v}$;
vi. scalar multiplication is compatible with multiplication in $\mathbb{F}$: $(ab)\mathbf{v} = a(b\mathbf{v})$;
vii. scalar multiplication distributes over vector addition: $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$;
viii. scalar multiplication distributes over addition in $\mathbb{F}$: $(a + b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}$.

If $\mathcal{V} = \{\mathbf{0}\}$, then $\mathcal{V}$ is a *zero vector space*; otherwise $\mathcal{V}$ is a *nonzero vector space*.

**Theorem 1.1.2.** *Basic Consequences of Vector Space Axioms*
*Let $\mathcal{V}$ be a vector space over $\mathbb{F}$. Then:*

*(i) the additive identity is unique;*
*(ii) for each $\mathbf{v} \in \mathcal{V}$, the additive inverse of $\mathbf{v}$ is unique;*
*(iii) $\forall \mathbf{v} \in \mathcal{V}$, $0\mathbf{v} = \mathbf{0}$;*
*(iv) $\forall a \in \mathbb{F}$, $a\mathbf{0} = \mathbf{0}$;*
*(v) $\forall a \in \mathbb{F}$ and $\mathbf{v} \in \mathcal{V}$, $a\mathbf{v} = \mathbf{0} \iff a = 0$ or $\mathbf{v} = \mathbf{0}$.*

*Proof.*

(i) If $\mathbf{0}$ and $\mathbf{0}'$ are additive identities, then $\mathbf{0} = \mathbf{0} + \mathbf{0}' = \mathbf{0}'$.
(ii) If $\mathbf{w}, \mathbf{w}'$ are additive inverses of $\mathbf{v}$, then $\mathbf{w} = \mathbf{w} + \mathbf{0} = \mathbf{w} + (\mathbf{v} + \mathbf{w}') = (\mathbf{w} + \mathbf{v}) + \mathbf{w}' = \mathbf{0} + \mathbf{w}' = \mathbf{w}'$.
(iii) $0\mathbf{v} = (0 + 0)\mathbf{v} = 0\mathbf{v} + 0\mathbf{v}$; adding $-(0\mathbf{v})$ gives $0\mathbf{v} = \mathbf{0}$.
(iv) $a\mathbf{0} = a(\mathbf{0} + \mathbf{0}) = a\mathbf{0} + a\mathbf{0}$; adding $-(a\mathbf{0})$ gives $a\mathbf{0} = \mathbf{0}$.
(v) ($\Rightarrow$) Suppose $a\mathbf{v} = \mathbf{0}$. If $a = 0$, the first alternative holds.
If $a \neq 0$, then $\mathbf{v} = 1\mathbf{v} = (a^{-1}a)\mathbf{v} = a^{-1}(a\mathbf{v}) = a^{-1}\mathbf{0} = \mathbf{0}$.
(v) ($\Leftarrow$) If $a = 0$ or $\mathbf{v} = \mathbf{0}$, then (iii) or (iv) gives $a\mathbf{v} = \mathbf{0}$.

*Structural Commentary*: These identities make cancellation, zeros, and inverses available.
*Key Insight*: The vector space axioms force the zero and inverse behaviour to be rigid, not optional structure. □

**Corollary 1.1.3.** *Scalar Multiplication by $-1$ Is Negation*
*Let $\mathcal{V}$ be a vector space over $\mathbb{F}$. Then $\forall\, \mathbf{v} \in \mathcal{V}$, $(-1)\mathbf{v} = -\mathbf{v}$.*

*Proof.* For $\mathbf{v} \in \mathcal{V}$, $(-1)\mathbf{v} + \mathbf{v} = (-1)\mathbf{v} + 1\mathbf{v} = (-1 + 1)\mathbf{v} = 0\mathbf{v} = \mathbf{0}$.
Hence $(-1)\mathbf{v}$ is an additive inverse of $\mathbf{v}$; by uniqueness, $(-1)\mathbf{v} = -\mathbf{v}$.
*Structural Commentary*: Scalar multiplication by $-1$ is exactly the same operation as taking an additive inverse.
*Key Insight*: Negation is not an extra operation; it is already encoded by the field action. □

**Definition 1.1.4.** *Finitely Nonzero Sequence Space*
Let $\mathbb{F}^{(\mathbb{N})}$ denote the set of sequences $(x_1, x_2, \ldots)$ in $\mathbb{F}$ with finite support, i.e. $x_j = 0$ for all but finitely many $j \in \mathbb{N}$. Addition and scalar multiplication are componentwise.

**Proposition 1.1.5.** *Finitely Nonzero Sequences Form a Vector Space*
*The set $\mathbb{F}^{(\mathbb{N})}$ is a vector space over $\mathbb{F}$ under componentwise addition and scalar multiplication.*

*Proof.* The zero sequence has finite support.
If $\mathbf{x}, \mathbf{y} \in \mathbb{F}^{(\mathbb{N})}$ and $a \in \mathbb{F}$, then $a\mathbf{x} + \mathbf{y}$ can be nonzero only where $\mathbf{x}$ or $\mathbf{y}$ is nonzero, a finite set of indices.
Hence $a\mathbf{x} + \mathbf{y} \in \mathbb{F}^{(\mathbb{N})}$, and the vector space axioms are inherited componentwise from $\mathbb{F}$.
*Structural Commentary*: Finite support is preserved by finite linear combinations.
*Key Insight*: Componentwise operations cannot create infinitely many new nonzero entries. □

**Example 1.1.6.** *Standard Vector Space Models*
Let $\mathbb{F}$ be a field and let $X$ be a set. The following are $\mathbb{F}$-vector spaces.

(i) $\mathbb{F}^X = \{f : X \to \mathbb{F}\}$, with pointwise operations $(f+g)(x) = f(x) + g(x)$ and $(af)(x) = af(x)$.
(ii) $\mathrm{Mat}(m, n, \mathbb{F})$ and $\mathbb{F}^n$, with entrywise operations. For $\mathbb{F} = \mathbb{F}_q$, set $V(n, q) := \mathbb{F}_q^n$.
(iii) $\mathrm{Seq}(\mathbb{F}) = \mathbb{F}^{\mathbb{N}}$, with componentwise operations; $\mathbb{F}^{(\mathbb{N})}$ is its finite-support subspace.
(iv) Over $\mathbb{C}$, the sequence spaces $c_0 = \{(z_n) : z_n \to 0\}$, $\ell^\infty = \{(z_n) : \sup_n |z_n| < \infty\}$, and, for $1 \le p < \infty$, $\ell^p = \left\{(z_n) : \sum_{n=1}^{\infty} |z_n|^p < \infty\right\}$ are vector spaces under componentwise operations; for $\ell^p$, closure under addition follows from Minkowski's inequality.

*Structural Commentary*: The vector-space axioms are independent of finite-dimensional coordinates; function and sequence spaces give canonical infinite-dimensional models.
*Key Insight*: Pointwise or componentwise operations inherit linear structure from the scalar field.

**Definition 1.1.7.** *Product of Vector Spaces*
Let $\mathcal{V}_1, \dots, \mathcal{V}_m$ be vector spaces over $\mathbb{F}$. Their *product* is $\mathcal{V}_1 \times \cdots \times \mathcal{V}_m := \{(\mathbf{v}_1, \dots, \mathbf{v}_m) : \mathbf{v}_j \in \mathcal{V}_j\}$.
Addition and scalar multiplication are defined componentwise:
$(\mathbf{u}_1, \dots, \mathbf{u}_m) + (\mathbf{v}_1, \dots, \mathbf{v}_m) := (\mathbf{u}_1 + \mathbf{v}_1, \dots, \mathbf{u}_m + \mathbf{v}_m)$ and $a(\mathbf{v}_1, \dots, \mathbf{v}_m) := (a\mathbf{v}_1, \dots, a\mathbf{v}_m)$.

**Proposition 1.1.8.** *Products Are Vector Spaces*
*If $\mathcal{V}_1, \dots, \mathcal{V}_m$ are vector spaces over $\mathbb{F}$, then $\mathcal{V}_1 \times \cdots \times \mathcal{V}_m$ is a vector space under componentwise operations.*

*Proof.* The additive identity is $(\mathbf{0}_1, \dots, \mathbf{0}_m)$, where $\mathbf{0}_j$ is the zero vector of $\mathcal{V}_j$, and additive inverse of $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is $(-\mathbf{v}_1, \dots, -\mathbf{v}_m)$. All remaining vector space axioms hold componentwise because they hold in each $\mathcal{V}_j$.
*Structural Commentary*: Product spaces collect several vector spaces into one coordinatewise structure.
*Key Insight*: Every vector space axiom reduces to the same axiom in each component. □

## 1.2 Subspaces and Spans

**Definition 1.2.1.** *Subspace*
Let $\mathcal{V}$ be a vector space over $\mathbb{F}$. A subset $\mathcal{U} \subseteq \mathcal{V}$ is a *subspace*, written $\mathcal{U} \le \mathcal{V}$, if $\mathcal{U}$ is a vector space over $\mathbb{F}$ under the operations inherited from $\mathcal{V}$.

**Theorem 1.2.2.** *Subspace Test*
*Let $\mathcal{V}$ be a vector space over $\mathbb{F}$ and let $\mathcal{U}$ be a nonempty subset of $\mathcal{V}$.*
*Then $\mathcal{U}$ is a subspace of $\mathcal{V}$ iff $\forall\, \mathbf{u}, \mathbf{v} \in \mathcal{U}$ and $\forall\, c \in \mathbb{F}$, one has $c\mathbf{u} + \mathbf{v} \in \mathcal{U}$.*

*Proof.*
(⇒) If $\mathcal{U} \le \mathcal{V}$, then closure under scalar multiplication and addition gives $c\mathbf{u} + \mathbf{v} \in \mathcal{U}$ for all $\mathbf{u}, \mathbf{v} \in \mathcal{U}$ and $c \in \mathbb{F}$.
(⇐) Assume the stated closure condition. Choose $\mathbf{u}_0 \in \mathcal{U}$. Then $\mathbf{0} = (-1)\mathbf{u}_0 + \mathbf{u}_0 \in \mathcal{U}$. For $\mathbf{u}, \mathbf{v} \in \mathcal{U}$ and $c \in \mathbb{F}$, the identities $-\mathbf{u} = (-1)\mathbf{u} + \mathbf{0}$, $c\mathbf{u} = c\mathbf{u} + \mathbf{0}$, and $\mathbf{u} + \mathbf{v} = 1\mathbf{u} + \mathbf{v}$ give closure under additive inverses, scalar multiplication, and addition. The remaining axioms are inherited from $\mathcal{V}$, so $\mathcal{U} \le \mathcal{V}$.

*Structural Commentary*: One closure condition captures the subspace axioms.
*Key Insight*: To prove a subset is a subspace, prove it survives one affine-looking linear combination. □

**Proposition 1.2.3.** *Intersection of Subspaces*
*Let $\mathcal{U}$ and $\mathcal{W}$ be subspaces of $\mathcal{V}$. Then $\mathcal{U} \cap \mathcal{W} = \{\mathbf{v} : \mathbf{v} \in \mathcal{U}$ and $\mathbf{v} \in \mathcal{W}\}$ is a subspace of $\mathcal{V}$.*

*Proof.* Since $\mathbf{0} \in \mathcal{U}$ and $\mathbf{0} \in \mathcal{W}$, we have $\mathbf{0} \in \mathcal{U} \cap \mathcal{W}$, so the intersection is nonempty.
If $\mathbf{u}, \mathbf{v} \in \mathcal{U} \cap \mathcal{W}$ and $c \in \mathbb{F}$, then $c\mathbf{u} + \mathbf{v}$ lies in both $\mathcal{U}$ and $\mathcal{W}$ because each is a subspace.
Thus $c\mathbf{u} + \mathbf{v} \in \mathcal{U} \cap \mathcal{W}$, and Theorem 1.2.2 gives $\mathcal{U} \cap \mathcal{W} \le \mathcal{V}$.
*Structural Commentary*: Intersections inherit shared closure.
*Key Insight*: Common vectors remain closed under linear combinations. □

**Definition 1.2.4.** *Lists, Linear Combinations, and Span*

(i) A *list* in $\mathcal{V}$ is $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_m)$; the empty list is $()$.
(ii) For $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_m)$ and $a_1, \dots, a_m \in \mathbb{F}$, a *linear combination* of $\beta$ is $\sum_{j=1}^{m} a_j \mathbf{v}_j$.
(iii) For $E \subseteq \mathcal{V}$, the *span* of $E$ is $\mathrm{span}(E) := \left\{\sum_{j=1}^{m} a_j \mathbf{v}_j : m \in \mathbb{N},\ a_j \in \mathbb{F},\ \mathbf{v}_j \in E\right\}$, with $\mathrm{span}(\emptyset) = \{\mathbf{0}\}$.
(iv) For $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_m)$, the *span* of $\beta$ is $\mathrm{span}(\beta) := \mathrm{span}(\{\mathbf{v}_1, \dots, \mathbf{v}_m\})$.
(v) $E \subseteq \mathcal{V}$ *spans* $\mathcal{V}$ iff $\mathrm{span}(E) = \mathcal{V}$; a list $\beta$ *spans* $\mathcal{V}$ iff $\mathrm{span}(\beta) = \mathcal{V}$.

**Theorem 1.2.5.** *Span is the Smallest Containing Subspace*
*Let $E \subseteq \mathcal{V}$. Then* $\operatorname{span}(E)$ *is the smallest subspace of* $\mathcal{V}$ *containing* $E$.

*Proof.* If $E = \emptyset$, then $\operatorname{span}(E) = \{\mathbf{0}\}$, which is a subspace. Suppose $E \neq \emptyset$. For $\mathbf{x}, \mathbf{y} \in \operatorname{span}(E)$ and $c \in \mathbb{F}$, both $\mathbf{x}$ and $\mathbf{y}$ are finite linear combinations of vectors in $E$, so $c\mathbf{x} + \mathbf{y}$ is also such a combination. Hence $c\mathbf{x} + \mathbf{y} \in \operatorname{span}(E)$, and Theorem 1.2.2 makes $\operatorname{span}(E)$ a subspace.
Each $\mathbf{v} \in E$ satisfies $\mathbf{v} = 1\mathbf{v} \in \operatorname{span}(E)$, so $E \subseteq \operatorname{span}(E)$. If $\mathcal{U} \leq \mathcal{V}$ and $E \subseteq \mathcal{U}$, then closure under finite linear combinations gives $\operatorname{span}(E) \subseteq \mathcal{U}$. Thus $\operatorname{span}(E)$ is the smallest subspace containing $E$.
*Structural Commentary*: Span closes a set of vectors under finite linear combinations, giving exactly the subspace forced by those vectors.
*Key Insight*: Nothing beyond linear closure is added. Every containing subspace must already contain span. □

**Proposition 1.2.6.** *Monotonicity of Span*
*Let $\mathcal{U}$ and $\mathcal{W}$ be subsets of $\mathcal{V}$. If $\mathcal{U} \subseteq \mathcal{W}$, then* $\operatorname{span}(\mathcal{U}) \subseteq \operatorname{span}(\mathcal{W})$.

*Proof.* If $\mathcal{U} = \emptyset$, then $\operatorname{span}(\mathcal{U}) = \{\mathbf{0}\} \subseteq \operatorname{span}(\mathcal{W})$.
Suppose $\mathcal{U} \neq \emptyset$. Every vector in $\operatorname{span}(\mathcal{U})$ is a finite linear combination of vectors in $\mathcal{U}$; since $\mathcal{U} \subseteq \mathcal{W}$, the same expression is a finite linear combination of vectors in $\mathcal{W}$. Hence $\operatorname{span}(\mathcal{U}) \subseteq \operatorname{span}(\mathcal{W})$.
*Structural Commentary*: Enlarging the generating set can only enlarge the collection of linear combinations.
*Key Insight*: Span is monotone with respect to inclusion of generating sets. □

**Method 1.2.7.** *Subspace Diagnostic*
Let $S \subseteq \mathcal{V}$.

(i) To prove $S \leq \mathcal{V}$, show $S \neq \varnothing$ and $\forall\, \mathbf{u}, \mathbf{v} \in S,\ \forall\, c \in \mathbb{F},\ c\mathbf{u} + \mathbf{v} \in S$.
(ii) To disprove $S \leq \mathcal{V}$, exhibit $\mathbf{u}, \mathbf{v} \in S$ and $c \in \mathbb{F}$ with $c\mathbf{u} + \mathbf{v} \notin S$; $\mathbf{0} \in S$ alone is insufficient.
(iii) The generated subspace is $\langle S \rangle = \operatorname{span}(S) = \left\{ \sum_{i=1}^{n} a_i \mathbf{s}_i : n \in \mathbb{N},\ a_i \in \mathbb{F},\ \mathbf{s}_i \in S \right\}$.

**Definition 1.2.8.** *Linear Code and Hamming Weight*

(i) The *ambient code space* is $V(n, q) := \mathbb{F}_q^n$.
(ii) A *linear code* of length $n$ over $\mathbb{F}_q$ is a *subspace* $C \leq V(n, q)$.
(iii) For $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{F}_q^n$, its *Hamming weight* is $\operatorname{wt}(\mathbf{x}) := \#\{j : 1 \leq j \leq n,\ x_j \neq 0\}$.

**Example 1.2.9.** *Even-Weight Code*
In $V(n, 2) = \mathbb{F}_2^n$, let $E_n = \left\{ \mathbf{x} \in \mathbb{F}_2^n : \sum_{j=1}^{n} x_j = 0 \right\}$. The condition $\sum_{j=1}^{n} x_j = 0$ is equivalent to $\operatorname{wt}(\mathbf{x})$ being even.
If $\mathbf{x}, \mathbf{y} \in E_n$ and $c \in \mathbb{F}_2$, then $\sum_{j=1}^{n} (cx_j + y_j) = c \sum_{j=1}^{n} x_j + \sum_{j=1}^{n} y_j = 0$, so $c\mathbf{x} + \mathbf{y} \in E_n$. Hence $E_n \leq V(n, 2)$ by Theorem 1.2.2, and $E_n$ is a linear code.
*Structural Commentary*: The even-weight code is the parity-zero subspace in $V(n, 2)$.
*Key Insight*: Parity is linear over $\mathbb{F}_2$.

## 1.3 Linear Independence and Bases

**Definition 1.3.1.** *Linear Independence, Bases, and Dimension*

(i) A list $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is *linearly independent* if $\sum_{j=1}^{m} a_j \mathbf{v}_j = \mathbf{0} \Rightarrow a_1 = \cdots = a_m = 0$.
Otherwise it is *linearly dependent*. The empty list is linearly independent.
(ii) A list $\beta$ is a *basis* of $\mathcal{V}$ if $\beta$ is linearly independent and $\operatorname{span}(\beta) = \mathcal{V}$.
(iii) $\mathcal{V}$ is *finite-dimensional* if some finite list $\beta$ satisfies $\operatorname{span}(\beta) = \mathcal{V}$; otherwise $\mathcal{V}$ is *infinite-dimensional*.

**Proposition 1.3.2.** *Unique Coefficients for Independent Lists*
*Let $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ be a linearly independent list in $\mathcal{V}$.*
*Then $a_1\mathbf{v}_1 + \cdots + a_m\mathbf{v}_m = b_1\mathbf{v}_1 + \cdots + b_m\mathbf{v}_m$ iff $a_j = b_j$ for each $j = 1, \ldots, m$.*

*Proof.*

(⇒) Subtracting the two sides gives $(a_1 - b_1)\mathbf{v}_1 + \cdots + (a_m - b_m)\mathbf{v}_m = \mathbf{0}$.
Linear independence forces $a_j - b_j = 0$ for each $j$, hence $a_j = b_j$.
(⇐) If $a_j = b_j$ for each $j$, then the two linear combinations have identical coefficients and are equal.

*Structural Commentary*: Linear independence makes coefficients in a linear combination unique.
*Key Insight*: Independent vectors allow no hidden coefficient cancellation. □

**Proposition 1.3.3.** *Unique Coordinate Representation*
*$\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$ iff each $\mathbf{v} \in \mathcal{V}$ has a unique coefficient vector $(a_1, \ldots, a_n) \in \mathbb{F}^n$ with $\mathbf{v} = \sum_{j=1}^{n} a_j \mathbf{v}_j$.*

*Proof.*

($\Rightarrow$) Spanning gives a representation of each $\mathbf{v} \in \mathcal{V}$. If $\mathbf{v} = \sum_{j=1}^{n} a_j \mathbf{v}_j = \sum_{j=1}^{n} b_j \mathbf{v}_j$, then Proposition 1.3.2 gives $a_j = b_j$ for each $j$.

($\Leftarrow$) Existence gives $\operatorname{span}(\beta) = \mathcal{V}$. If $\sum_{j=1}^{n} a_j \mathbf{v}_j = \mathbf{0}$, compare with $\mathbf{0} = \sum_{j=1}^{n} 0\mathbf{v}_j$; uniqueness gives $a_j = 0$ for each $j$, so $\beta$ is independent.

*Structural Commentary*: A basis is exactly existence and uniqueness of coordinates.
*Key Insight*: Coordinates are basis coefficients. □

**Proposition 1.3.4.** *Appending to a Dependent List*
*Let $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ be a linearly dependent list in $\mathcal{V}$.*
*Then for each $\mathbf{v} \in \mathcal{V}$, the list $(\mathbf{v}_1, \ldots, \mathbf{v}_m, \mathbf{v})$ is linearly dependent.*

*Proof.* Since $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly dependent, $\exists$ scalars $a_1, \ldots, a_m$, not all zero, with $a_1\mathbf{v}_1 + \cdots + a_m\mathbf{v}_m = \mathbf{0}$. Appending coefficient 0 gives $a_1\mathbf{v}_1 + \cdots + a_m\mathbf{v}_m + 0\mathbf{v} = \mathbf{0}$, and the scalars $a_1, \ldots, a_m, 0$ are not all zero.
*Structural Commentary*: Longer lists inherit dependence; set new coefficients to 0.
*Key Insight*: Once a list has a redundancy, adding more vectors cannot remove it. □

**Definition 1.3.5.** *Omitting a Vector from a List*
Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ be a list in $\mathcal{V}$ with $m \geq 2$.
For $j \in \{1, \ldots, m\}$, the list obtained from $\beta$ by omitting $\mathbf{v}_j$ is denoted $(\mathbf{v}_1, \ldots, \widehat{\mathbf{v}_j}, \ldots, \mathbf{v}_m)$.

**Proposition 1.3.6.** *Omitting from an Independent List*
*If $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly independent and $m \geq 2$, then $\beta_{\widehat{j}} := (\mathbf{v}_1, \ldots, \widehat{\mathbf{v}_j}, \ldots, \mathbf{v}_m)$ is linearly independent for each $j \in \{1, \ldots, m\}$.*

*Proof.* Fix $j$. Any relation $\sum_{i \neq j} a_i \mathbf{v}_i = \mathbf{0}$ on $\beta_{\widehat{j}}$ becomes a relation on $\beta$ after setting $a_j = 0$. Independence of $\beta$ gives $a_i = 0$ for all $i$, so the sublist is independent.
*Structural Commentary*: Deletion cannot create cancellation.
*Key Insight*: Every sublist of an independent list remains independent. □

**Proposition 1.3.7.** *Adjoining and Removing Vectors from Lists*
*Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ be a list in $\mathcal{V}$. Write $(\beta, \mathbf{v}) := (\mathbf{v}_1, \ldots, \mathbf{v}_m, \mathbf{v})$ and $\beta_{\widehat{j}} := (\mathbf{v}_1, \ldots, \widehat{\mathbf{v}_j}, \ldots, \mathbf{v}_m)$.*

*(i) If $\beta$ is linearly independent and $\mathbf{v} \notin \operatorname{span}(\beta)$, then $(\beta, \mathbf{v})$ is linearly independent.*

*(ii) If $\sum_{i=1}^{m} a_i \mathbf{v}_i = \mathbf{0}$ is nontrivial and $a_j \neq 0$, then $\operatorname{span}(\beta_{\widehat{j}}) = \operatorname{span}(\beta)$.*
*Hence $\operatorname{span}(\beta) = \mathcal{V}$ implies $\operatorname{span}(\beta_{\widehat{j}}) = \mathcal{V}$.*

*Proof.*

(i) Suppose $\sum_{i=1}^{m} a_i \mathbf{v}_i + a\mathbf{v} = \mathbf{0}$. If $a \neq 0$, then $\mathbf{v} = -a^{-1} \sum_{i=1}^{m} a_i \mathbf{v}_i \in \operatorname{span}(\beta)$, a contradiction.
Hence $a = 0$, and independence of $\beta$ gives $a_1 = \cdots = a_m = 0$.

(ii) Solving the relation for $\mathbf{v}_j$ gives $\mathbf{v}_j = -a_j^{-1} \sum_{i \neq j} a_i \mathbf{v}_i \in \operatorname{span}(\beta_{\widehat{j}})$. Thus every vector of $\beta$ lies in $\operatorname{span}(\beta_{\widehat{j}})$, so $\operatorname{span}(\beta) \subseteq \operatorname{span}(\beta_{\widehat{j}})$. The reverse inclusion is immediate because $\beta_{\widehat{j}}$ is a sublist of $\beta$.

*Structural Commentary*: New directions preserve independence; redundant vectors preserve span.
*Key Insight*: Add new directions; remove redundant ones. □

**Lemma 1.3.8.** *Replacement Lemma*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$, let $\beta = (\mathbf{u}_1, \ldots, \mathbf{u}_r)$ span $\mathcal{V}$, and let $\mathbf{0} \neq \mathbf{v} = \sum_{i=1}^{r} c_i \mathbf{u}_i$.*
*For $j \in \{1, \ldots, r\}$, set $\beta_{j \leftarrow \mathbf{v}} := (\mathbf{u}_1, \ldots, \mathbf{u}_{j-1}, \mathbf{v}, \mathbf{u}_{j+1}, \ldots, \mathbf{u}_r)$.*

*(i) $c_j \neq 0$ for some $j$.*
*(ii) If $c_j \neq 0$, then $\operatorname{span}(\beta_{j \leftarrow \mathbf{v}}) = \mathcal{V}$; if $\beta$ is a basis, then $\beta_{j \leftarrow \mathbf{v}}$ is a basis.*
*(iii) If $r \geq 2$, $\beta$ is a basis, and $\mathbf{v} \notin \operatorname{span}(\mathbf{u}_1, \ldots, \mathbf{u}_k)$ for some $1 \leq k < r$, then $\beta_{j \leftarrow \mathbf{v}}$ is a basis for some $j > k$.*

*Proof.*

(i) If all $c_j = 0$, then $\mathbf{v} = \mathbf{0}$.

(ii) Since $\mathbf{u}_j = c_j^{-1}\left(\mathbf{v} - \sum_{i\neq j} c_i\mathbf{u}_i\right)$, every vector of $\beta$ lies in $\operatorname{span}(\beta_{j\leftarrow\mathbf{v}})$.
Conversely, $\mathbf{v} \in \operatorname{span}(\beta)$, so every vector of $\beta_{j\leftarrow\mathbf{v}}$ lies in $\operatorname{span}(\beta)$. Hence $\operatorname{span}(\beta_{j\leftarrow\mathbf{v}}) = \mathcal{V}$.
If $\beta$ is a basis and $\sum_{i<j} b_i\mathbf{u}_i + a\mathbf{v} + \sum_{i>j} b_i\mathbf{u}_i = \mathbf{0}$, substitution gives $\sum_{i\neq j}(b_i + ac_i)\mathbf{u}_i + ac_j\mathbf{u}_j = \mathbf{0}$. Independence of $\beta$ gives $ac_j = 0$ and $b_i + ac_i = 0$ for $i \neq j$.
Since $c_j \neq 0$, $a = 0$, and then every $b_i = 0$. Thus $\beta_{j\leftarrow\mathbf{v}}$ is linearly independent.

(iii) Since $\mathbf{v} \notin \operatorname{span}(\mathbf{u}_1, \ldots, \mathbf{u}_k)$, some coefficient among $c_{k+1}, \ldots, c_r$ is nonzero.
Choose $j > k$ with $c_j \neq 0$; by (ii), $\beta_{j\leftarrow\mathbf{v}}$ is a basis.

*Structural Commentary*: Replacement preserves span.
*Key Insight*: Replace where the coefficient is nonzero. □

**Lemma 1.3.9.** *Linear Dependence Lemma*
*Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ be linearly dependent with $\mathbf{v}_1 \neq \mathbf{0}$, and set $\beta_{\widehat{j}} := (\mathbf{v}_1, \ldots, \widehat{\mathbf{v}_j}, \ldots, \mathbf{v}_m)$.*
*Then some $j \in \{2, \ldots, m\}$ satisfies $\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$ and $\operatorname{span}(\beta_{\widehat{j}}) = \operatorname{span}(\beta)$.*

*Proof.* Choose a nontrivial relation $\sum_{i=1}^{m} a_i\mathbf{v}_i = \mathbf{0}$ and let $j = \max\{i : a_i \neq 0\}$.
If $j = 1$, then $a_1\mathbf{v}_1 = \mathbf{0}$ with $a_1 \neq 0$, so $\mathbf{v}_1 = \mathbf{0}$, contrary to hypothesis. Hence $j \geq 2$.
Since $a_i = 0$ for $i > j$, we have $\mathbf{v}_j = -a_j^{-1}\sum_{i=1}^{j-1} a_i\mathbf{v}_i$, so $\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$.
Substituting this expression for $\mathbf{v}_j$ rewrites every linear combination of $\beta$ without $\mathbf{v}_j$, so $\operatorname{span}(\beta) \subseteq \operatorname{span}(\beta_{\widehat{j}})$.
The reverse inclusion follows because $\beta_{\widehat{j}}$ is a sublist of $\beta$. Hence the spans are equal.
*Structural Commentary*: The last nonzero coefficient gives the removable vector.
*Key Insight*: Dependence gives a later vector removable without losing span. □

**Theorem 1.3.10.** *Independent Lists Are No Longer Than Spanning Lists*
*If $\mathcal{V}$ is finite-dimensional, $\alpha = (\mathbf{u}_1, \ldots, \mathbf{u}_m)$ is linearly independent, and $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$ spans $\mathcal{V}$, then $m \leq n$.*

*Proof.* Suppose $m > n$.
If $n = 0$, then $\mathcal{V} = \operatorname{span}(\gamma) = \{\mathbf{0}\}$, contradicting independence of the nonempty list $\alpha$; hence $n \geq 1$.
Construct spanning lists $\gamma_k$ of length $n$ that contain $\mathbf{u}_1, \ldots, \mathbf{u}_k$ for $0 \leq k \leq n$. Set $\gamma_0 = \gamma$. Given $\gamma_k$ with $k < n$, write $\mathbf{u}_{k+1}$ in $\operatorname{span}(\gamma_k)$. Since $\mathbf{u}_{k+1} \notin \operatorname{span}(\mathbf{u}_1, \ldots, \mathbf{u}_k)$, some coefficient on an entry of $\gamma_k$ outside $\mathbf{u}_1, \ldots, \mathbf{u}_k$ is nonzero. Replacing that entry by $\mathbf{u}_{k+1}$ preserves span by Lemma 1.3.8(ii); call the result $\gamma_{k+1}$.
Then $\gamma_n$ spans $\mathcal{V}$ and contains $\mathbf{u}_1, \ldots, \mathbf{u}_n$, so $\operatorname{span}(\mathbf{u}_1, \ldots, \mathbf{u}_n) = \mathcal{V}$.
Since $m > n$, this puts $\mathbf{u}_{n+1}$ in the span of earlier vectors, contradicting independence of $\alpha$.
*Structural Commentary*: Independent length is bounded by spanning length.
*Key Insight*: Independent lists cannot be longer than spanning lists in finite dimensions. □

**Theorem 1.3.11.** *Reducing Spanning Lists*
*Every spanning list in a finite-dimensional vector space contains a sublist that is a basis.*

*Proof.* Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be a spanning list of $\mathcal{V}$. If $\beta$ is independent, it is a basis. Otherwise remove one vector without changing span: omit $\mathbf{v}_1$ if $\mathbf{v}_1 = \mathbf{0}$, and otherwise use Lemma 1.3.9 to remove a suitable $\mathbf{v}_j$. The resulting sublist still spans $\mathcal{V}$ and has length one less. Repeat this step. The length strictly decreases, so the process stops with a spanning independent sublist, hence a basis.
*Structural Commentary*: Delete redundant vectors while preserving span.
*Key Insight*: Spanning gives enough vectors; reduction deletes exactly the unnecessary ones. □

**Theorem 1.3.12.** *Extending Independent Lists*
*Every linearly independent list in a finite-dimensional vector space can be extended to a basis.*

*Proof.* Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ be linearly independent in $\mathcal{V}$, and choose a spanning list $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$ of $\mathcal{V}$.
Initialize $\mathcal{B} = \beta$. Inspect $\mathbf{w}_1, \ldots, \mathbf{w}_n$ in order, appending $\mathbf{w}_j$ exactly when $\mathbf{w}_j \notin \operatorname{span}(\mathcal{B})$. Each append preserves independence by Proposition 1.3.7.
After the inspection, each $\mathbf{w}_j$ lies in $\operatorname{span}(\mathcal{B})$: it was either already there or was appended. Since the $\mathbf{w}_j$ span $\mathcal{V}$ and $\operatorname{span}(\mathcal{B})$ is a subspace, every vector of $\mathcal{V}$ lies in $\operatorname{span}(\mathcal{B})$.
Thus $\mathcal{B}$ is a spanning independent list, hence a basis.
*Structural Commentary*: An independent list has no redundancy, but may miss directions. Appending only vectors outside its span fills those directions while preserving independence.
*Key Insight*: A finite-dimensional independent list becomes a basis once enough new directions are added. □

**Corollary 1.3.13.** *Bases and Subspaces in Finite Dimension*
*If $\mathcal{V}$ is finite-dimensional, then $\mathcal{V}$ has a basis. Moreover, $\mathcal{U} \leq \mathcal{V}$ implies $\mathcal{U}$ is finite-dimensional.*

*Proof.* Choose a finite spanning list $\gamma$ of $\mathcal{V}$. By Theorem 1.3.11, some sublist of $\gamma$ is a basis of $\mathcal{V}$.
Let $\mathcal{U} \leq \mathcal{V}$. If $\mathcal{U} = \{\mathbf{0}\}$, then $()$ is a basis. Otherwise start with $\beta = ()$ and append vectors from $\mathcal{U} \setminus \operatorname{span}(\beta)$ while possible. Proposition 1.3.7 preserves independence, while Theorem 1.3.10 bounds the length by the length of $\gamma$; hence the process terminates. At termination, $\mathcal{U} \subseteq \operatorname{span}(\beta)$, so $\beta$ is a finite basis of $\mathcal{U}$.
*Structural Commentary*: Finite-dimensional spaces have finite coordinate systems, and subspaces cannot contain longer independent lists than the ambient space permits.
*Key Insight*: Finite-dimensionality passes downward to subspaces. □

**Theorem 1.3.14.** *Basis Length Is Well-Defined*
*Any two bases of a finite-dimensional vector space have the same length.*

*Proof.* Let $\beta_1 = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ and $\beta_2 = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$ be bases of $\mathcal{V}$. Theorem 1.3.10 applied to $\beta_1$ against $\beta_2$ gives $m \leq n$, and reversing the roles gives $n \leq m$. Hence $m = n$.
*Structural Commentary*: Bases are simultaneously independent and spanning, so the comparison theorem traps their lengths in both directions. This makes dimension well-defined.
*Key Insight*: Dimension is an intrinsic number, not an artefact of a particular basis. □

## 1.4 Dimension and Coordinates

**Definition 1.4.1.** *Dimension*

(i) If $\mathcal{V} = \{\mathbf{0}\}$, then $\dim(\mathcal{V}) := 0$.
(ii) If $\mathcal{V}$ has a basis of length $n \geq 1$, then $\mathcal{V}$ is *$n$-dimensional* and $\dim(\mathcal{V}) := n$.

Theorem 1.3.14 makes $\dim(\mathcal{V})$ well-defined.

**Proposition 1.4.2.** *Subspace Dimension Criterion*
*Let $\mathcal{V}$ be finite-dimensional and let $\mathcal{U} \leq \mathcal{V}$. Then $\dim(\mathcal{U}) \leq \dim(\mathcal{V})$, and equality holds iff $\mathcal{U} = \mathcal{V}$.*

*Proof.* Let $\delta = (\mathbf{u}_1, \ldots, \mathbf{u}_m)$ be a basis of $\mathcal{U}$ and let $\beta$ be a basis of $\mathcal{V}$, where $n = \dim(\mathcal{V})$.
Since $\delta$ is independent in $\mathcal{V}$ and $\beta$ spans $\mathcal{V}$, Theorem 1.3.10 gives $m \leq n$.
($\Rightarrow$) If $\mathcal{U} = \mathcal{V}$, then any basis of $\mathcal{U}$ is a basis of $\mathcal{V}$, so $\dim(\mathcal{U}) = \dim(\mathcal{V})$.
($\Leftarrow$) If $\dim(\mathcal{U}) = \dim(\mathcal{V})$, then $\delta$ is independent in $\mathcal{V}$ of length $n$. By Theorem 1.3.12, extending $\delta$ to a basis of $\mathcal{V}$ adds no vector, since every basis has length $n$. Hence $\mathcal{U} = \operatorname{span}(\delta) = \mathcal{V}$.
*Structural Commentary*: Dimension bounds subspaces; equality forces fullness.
*Key Insight*: Equal dimension subspaces fill the ambient space. □

**Proposition 1.4.3.** *Full-Dimension Basis Criteria*
*Let $\mathcal{V}$ be finite-dimensional and set $n := \dim(\mathcal{V})$. Then:*

*(i) if $\gamma$ spans $\mathcal{V}$ and has length $n$, then $\gamma$ is a basis;*
*(ii) if $\alpha$ is linearly independent in $\mathcal{V}$ and has length $n$, then $\alpha$ is a basis.*

*Proof.*

(i) By Theorem 1.3.11, $\gamma$ contains a basis sublist. Since every basis has length $n$, no proper sublist can have length $n$; hence $\gamma$ itself is a basis.
(ii) Extend $\alpha$ to a basis of $\mathcal{V}$ by Theorem 1.3.12. Since every basis has length $n$ and $\alpha$ already has length $n$, no vector is added; hence $\alpha$ is a basis.

*Structural Commentary*: Full-length spanning or independent lists cannot be shortened or extended.
*Key Insight*: In dimension $n$, $n$ independent or $n$ spanning vectors already form a basis. □

**Theorem 1.4.4.** *Basis Characterizations*
*Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be a finite list in $\mathcal{V}$, and write $\beta^\frown \mathbf{x} = (\mathbf{v}_1, \ldots, \mathbf{v}_n, \mathbf{x})$. Then the following are equivalent:*

*(i) $\beta$ is a basis of $\mathcal{V}$;*
*(ii) each $\mathbf{x} \in \mathcal{V}$ has a unique coefficient vector $(a_1, \ldots, a_n) \in \mathbb{F}^n$ with $\mathbf{x} = \sum_{j=1}^{n} a_j \mathbf{v}_j$;*
*(iii) $\operatorname{span}(\beta) = \mathcal{V}$, and no proper sublist of $\beta$ spans $\mathcal{V}$;*
*(iv) $\beta$ is linearly independent, and $\beta^\frown \mathbf{x}$ is linearly dependent for each $\mathbf{x} \in \mathcal{V}$.*

*Proof.*

(i) $\Rightarrow$ (ii) This is Proposition 1.3.3.

(*ii*) $\Rightarrow$ (*iii*) Existence gives $\operatorname{span}(\beta) = \mathcal{V}$.
If a proper sublist $\gamma \subsetneq \beta$ spanned $\mathcal{V}$ and omitted the occurrence $\mathbf{v}_j$, then $\mathbf{v}_j$ would have both its standard $\beta$-coordinate representation and one using only $\gamma$, contradicting uniqueness.
(*iii*) $\Rightarrow$ (*iv*) If $\beta$ were dependent, Proposition 1.3.7 would remove some $\mathbf{v}_j$ without changing $\operatorname{span}(\beta)$, contradicting minimality. Thus $\beta$ is independent.
Since $\operatorname{span}(\beta) = \mathcal{V}$, writing $\mathbf{x}$ as a linear combination of $\beta$ gives a nontrivial relation on $\beta^\frown\mathbf{x}$.
(*iv*) $\Rightarrow$ (*i*) If $\operatorname{span}(\beta) \neq \mathcal{V}$, choose $\mathbf{x} \notin \operatorname{span}(\beta)$. Then Proposition 1.3.7 makes $\beta^\frown\mathbf{x}$ independent, contradicting (iv). Hence $\beta$ spans $\mathcal{V}$, and with independence from (iv), it is a basis.

*Structural Commentary*: A basis is simultaneously just enough to span and too large to extend independently.
*Key Insight*: Spanning and independence are dual minimality and maximality conditions. □

**Definition 1.4.5.** *Hamel Basis*
A subset $\mathcal{B} \subseteq \mathcal{V}$ is a *Hamel basis* of $\mathcal{V}$ if it is linearly independent and $\operatorname{span}(\mathcal{B}) = \mathcal{V}$. Equivalently, each $\mathbf{v} \in \mathcal{V}$ has a unique finitely supported coefficient function $a : \mathcal{B} \to \mathbb{F}$ with $\mathbf{v} = \sum_{\mathbf{b}\in\mathcal{B}} a(\mathbf{b})\mathbf{b}$.

Finite ordered bases are orderings of finite Hamel bases.

**Theorem 1.4.6.** *Hamel Basis Extension*
*Let $I \subseteq S \subseteq \mathcal{V}$, with $I$ linearly independent and $\operatorname{span}(S) = \mathcal{V}$.*
*Assuming Zorn's lemma, there exists a Hamel basis $\mathcal{B}$ of $\mathcal{V}$ such that $I \subseteq \mathcal{B} \subseteq S$. Consequently, every nonzero vector space has a basis; every independent set extends to a basis; every spanning set contains a basis.*

*Proof.* Let $\mathscr{P} = \{E : I \subseteq E \subseteq S,\ E \text{ linearly independent}\}$, ordered by inclusion. Then $I \in \mathscr{P}$.
If $\mathcal{C} \subseteq \mathscr{P}$ is a nonempty chain, put $U = \bigcup_{E\in\mathcal{C}} E$. In any finite relation $\sum_{k=1}^{m} a_k\mathbf{u}_k = \mathbf{0}$ in $U$, the finitely many vectors $\mathbf{u}_k$ lie in one member of the chain; independence there gives $a_k = 0$ for all $k$.
Thus $U \in \mathscr{P}$ and is an upper bound for $\mathcal{C}$. The empty chain is bounded by $I$.
By Zorn's lemma, Principle 6.5.3, choose maximal $\mathcal{B} \in \mathscr{P}$.
If $s \in S \setminus \operatorname{span}(\mathcal{B})$, then $\mathcal{B} \cup \{s\}$ is linearly independent, contradicting maximality. Hence $S \subseteq \operatorname{span}(\mathcal{B})$, so $\operatorname{span}(\mathcal{B}) = \mathcal{V}$ and $\mathcal{B}$ is a Hamel basis. The consequences follow by taking $S = \mathcal{V}$ or $I = \emptyset$.
*Structural Commentary*: Finite extension uses exchange; arbitrary extension uses maximality.
*Key Insight*: Maximal independence inside a spanning set forces spanning. □

**Theorem 1.4.7.** *Dimension as Basis Cardinality*
*If $\mathcal{B}$ and $\mathcal{C}$ are Hamel bases of $\mathcal{V}$, then $|\mathcal{B}| = |\mathcal{C}|$.*

*Proof.* If one basis is finite, say $|\mathcal{B}| = n$, then $\mathcal{C}$ cannot contain $n+1$ distinct vectors, since such a finite sublist would be independent while $\mathcal{B}$ spans $\mathcal{V}$, contradicting Theorem 1.3.10. Hence $\mathcal{C}$ is finite, and Theorem 1.3.14 gives $|\mathcal{C}| = n$. Thus assume both bases are infinite. For $\mathbf{c} \in \mathcal{C}$, let $\operatorname{supp}_{\mathcal{B}}(\mathbf{c}) \subseteq \mathcal{B}$ be its finite $\mathcal{B}$-support.
Every $\mathbf{b} \in \mathcal{B}$ appears in some $\operatorname{supp}_{\mathcal{B}}(\mathbf{c})$: otherwise all vectors in $\operatorname{span}(\mathcal{C})$ would have zero $\mathbf{b}$-coordinate, contradicting $\operatorname{span}(\mathcal{C}) = \mathcal{V}$. Hence $\mathcal{B} \subseteq \bigcup_{\mathbf{c}\in\mathcal{C}} \operatorname{supp}_{\mathcal{B}}(\mathbf{c})$. Thus $|\mathcal{B}| \leq \sum_{\mathbf{c}\in\mathcal{C}} |\operatorname{supp}_{\mathcal{B}}(\mathbf{c})| \leq |\mathcal{C}|\,\aleph_0 = |\mathcal{C}|$ by Theorem 6.4.2.
Interchanging $\mathcal{B}$ and $\mathcal{C}$ gives $|\mathcal{C}| \leq |\mathcal{B}|$; Theorem 6.3.1 gives equality.
*Structural Commentary*: Even in infinite dimensions, a basis size is intrinsic.
*Key Insight*: Each basis vector is detected inside finite supports relative to the other basis. □

**Definition 1.4.8.** *Hamel Dimension*
For any Hamel basis $\mathcal{B}$ of $\mathcal{V}$, set $\dim(\mathcal{V}) := |\mathcal{B}|$. This is well-defined by Theorem 1.4.7.

## 1.5 Linear Maps and Operators

Linear maps preserve vector-space structure. Kernels, ranges, rank-nullity, matrices, change of basis, and similarity describe them without making coordinates primary.

**Definition 1.5.1.** *Linear Map*
Let $\mathcal{V}$ and $\mathcal{W}$ be vector spaces over the same field $\mathbb{F}$.
A function $T : \mathcal{V} \to \mathcal{W}$ is *linear* if $T(a\mathbf{u} + b\mathbf{v}) = aT\mathbf{u} + bT\mathbf{v}$ for all $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ and $a, b \in \mathbb{F}$.
The set of all linear maps from $\mathcal{V}$ to $\mathcal{W}$ is denoted $\mathcal{L}(\mathcal{V}, \mathcal{W})$. If $\mathcal{V} = \mathcal{W}$, elements of $\mathcal{L}(\mathcal{V})$ are *operators* on $\mathcal{V}$.

**Definition 1.5.2.** *Zero and Identity Maps*
For vector spaces $\mathcal{V}$ and $\mathcal{W}$, the *zero map* from $\mathcal{V}$ to $\mathcal{W}$ is $0_{\mathcal{V},\mathcal{W}} : \mathcal{V} \to \mathcal{W}$, where $0_{\mathcal{V},\mathcal{W}}\mathbf{v} = \mathbf{0}_{\mathcal{W}}$ for all $\mathbf{v} \in \mathcal{V}$.
The *identity map* on $\mathcal{V}$ is $I_{\mathcal{V}} : \mathcal{V} \to \mathcal{V}$, where $I_{\mathcal{V}}\mathbf{v} = \mathbf{v}$ for all $\mathbf{v} \in \mathcal{V}$.

**Proposition 1.5.3.** *Linear Maps Send Zero to Zero*
*If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $T\mathbf{0}_{\mathcal{V}} = \mathbf{0}_{\mathcal{W}}$.*

*Proof.* Linearity gives $T\mathbf{0}_{\mathcal{V}} = T(\mathbf{0}_{\mathcal{V}} + \mathbf{0}_{\mathcal{V}}) = T\mathbf{0}_{\mathcal{V}} + T\mathbf{0}_{\mathcal{V}}$. Adding $-T\mathbf{0}_{\mathcal{V}}$ to both sides gives $T\mathbf{0}_{\mathcal{V}} = \mathbf{0}_{\mathcal{W}}$.
*Structural Commentary*: Linearity preserves zero.
*Key Insight*: A linear map cannot move zero away from zero. □

**Definition 1.5.4.** *Operations on Linear Maps*
For $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ and $a \in \mathbb{F}$, define pointwise $(S+T)\mathbf{v} := S\mathbf{v} + T\mathbf{v}$ and $(aT)\mathbf{v} := a(T\mathbf{v})$ for every $\mathbf{v} \in \mathcal{V}$.

**Proposition 1.5.5.** *Linear Maps Form a Vector Space*
*With the operations above, $\mathcal{L}(\mathcal{V}, \mathcal{W})$ is a vector space over $\mathbb{F}$.*

*Proof.* If $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ and $a, c \in \mathbb{F}$, then for $\mathbf{u}, \mathbf{v} \in \mathcal{V}$,

$$(aS+T)(c\mathbf{u}+\mathbf{v}) = a(cS\mathbf{u}+S\mathbf{v}) + (cT\mathbf{u}+T\mathbf{v}) = c(aS+T)\mathbf{u} + (aS+T)\mathbf{v}.$$

Thus $\mathcal{L}(\mathcal{V}, \mathcal{W})$ is closed under linear combinations. The zero map is the additive identity, $-T$ is the additive inverse of $T$, and the remaining vector-space axioms hold pointwise because they hold in $\mathcal{W}$.
*Structural Commentary*: Linear map operations are pointwise.
*Key Insight*: Pointwise sums and scalar multiples preserve linearity. □

**Definition 1.5.6.** *Product of Linear Maps*
For $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ and $S \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, the *product* $ST \in \mathcal{L}(\mathcal{U}, \mathcal{W})$ is defined by $(ST)\mathbf{u} := S(T\mathbf{u})$ for all $\mathbf{u} \in \mathcal{U}$.

**Proposition 1.5.7.** *Algebra of Linear Map Products*
*If $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ and $S \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $ST \in \mathcal{L}(\mathcal{U}, \mathcal{W})$. Whenever defined, $(RS)T = R(ST)$, $TI_{\mathcal{U}} = T = I_{\mathcal{V}}T$, $(S_1+S_2)T = S_1T + S_2T$, and $S(T_1+T_2) = ST_1 + ST_2$.*

*Proof.* For $\mathbf{u}, \mathbf{x} \in \mathcal{U}$ and $a \in \mathbb{F}$, $(ST)(a\mathbf{u}+\mathbf{x}) = S(aT\mathbf{u}+T\mathbf{x}) = aS(T\mathbf{u}) + S(T\mathbf{x}) = a(ST)\mathbf{u} + (ST)\mathbf{x}$, hence $ST \in \mathcal{L}(\mathcal{U}, \mathcal{W})$. For each remaining identity $A = B$ and admissible $\mathbf{z}$, $A\mathbf{z} = B\mathbf{z}$ by associativity of evaluation, identity maps, or distributivity in the target space; hence $A = B$.
*Structural Commentary*: Composition chains linear actions, while addition distributes through evaluation.
*Key Insight*: Linear map algebra is function composition plus pointwise vector-space algebra. □

**Definition 1.5.8.** *Graph of a Function*
For a function $T : \mathcal{V} \to \mathcal{W}$, the *graph* of $T$ is $\operatorname{graph}(T) := \{(\mathbf{v}, T\mathbf{v}) : \mathbf{v} \in \mathcal{V}\} \subseteq \mathcal{V} \times \mathcal{W}$.

**Proposition 1.5.9.** *Linearity and Graphs*
*A function $T : \mathcal{V} \to \mathcal{W}$ is linear if and only if* $\operatorname{graph}(T)$ *is a subspace of $\mathcal{V} \times \mathcal{W}$.*

*Proof.*

($\Rightarrow$) If $T$ is linear, then $(\mathbf{0}, \mathbf{0}) \in \operatorname{graph}(T)$.
If $(\mathbf{u}, T\mathbf{u}), (\mathbf{v}, T\mathbf{v}) \in \operatorname{graph}(T)$ and $a \in \mathbb{F}$, then $a(\mathbf{u}, T\mathbf{u}) + (\mathbf{v}, T\mathbf{v}) = (a\mathbf{u}+\mathbf{v}, T(a\mathbf{u}+\mathbf{v})) \in \operatorname{graph}(T)$.
Thus $\operatorname{graph}(T)$ is a subspace.

($\Leftarrow$) If $\operatorname{graph}(T)$ is a subspace, then for $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ and $a \in \mathbb{F}$, the vector $a(\mathbf{u}, T\mathbf{u}) + (\mathbf{v}, T\mathbf{v}) = (a\mathbf{u}+\mathbf{v}, aT\mathbf{u}+T\mathbf{v})$ lies in $\operatorname{graph}(T)$. Hence $T(a\mathbf{u}+\mathbf{v}) = aT\mathbf{u} + T\mathbf{v}$, so $T$ is linear.

*Structural Commentary*: A graph records a function as a subset of a product space.
*Key Insight*: Closure of the graph is exactly linearity of the function. □

**Principle 1.5.10.** *Coordinate-Free Principle*
A linear map is completely determined by what it does to a basis. Coordinates do not create the map; they only encode it once bases have been chosen.

**Proposition 1.5.11.** *Linear Maps Are Determined by a Basis*
*If $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$ and $\mathbf{w}_1, \ldots, \mathbf{w}_n \in \mathcal{W}$ are arbitrary, then $\exists$ a unique linear map $T : \mathcal{V} \to \mathcal{W}$ such that $T(\mathbf{v}_j) = \mathbf{w}_j$ for $j = 1, \ldots, n$.*

*Proof.* Every $\mathbf{v} \in \mathcal{V}$ has a unique expression $\mathbf{v} = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$. Define $T(\mathbf{v}) := a_1\mathbf{w}_1 + \cdots + a_n\mathbf{w}_n$.
This is well-defined because basis coordinates are unique, and linearity follows from coordinate expansions. Any linear map with the prescribed basis values must agree with this formula, so it is unique.
*Structural Commentary*: A basis reduces a linear map to finitely many choices.
*Key Insight*: Once basis images are fixed, linearity determines every other value. □

## 1.6 Kernels, Ranges, and Map Properties

**Definition 1.6.1.** *Kernel and Range*
For $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, define $\ker(T) := \{\mathbf{v} \in \mathcal{V} : T\mathbf{v} = \mathbf{0}_{\mathcal{W}}\}$ and $\text{range}(T) := \{T\mathbf{v} : \mathbf{v} \in \mathcal{V}\}$.
The kernel is also called the *nullspace* of $T$.

**Definition 1.6.2.** *Injective and Surjective Linear Maps*
A function $T : \mathcal{V} \to \mathcal{W}$ is *injective* if $T\mathbf{u} = T\mathbf{v}$ implies $\mathbf{u} = \mathbf{v}$. It is *surjective* if $\text{range}(T) = \mathcal{W}$.

**Proposition 1.6.3.** *Injectivity and the Kernel*
*A linear map $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ is injective if and only if $\ker(T) = \{\mathbf{0}\}$.*

*Proof.*
($\Rightarrow$) If $T$ is injective and $\mathbf{v} \in \ker(T)$, then $T\mathbf{v} = \mathbf{0} = T\mathbf{0}$, so $\mathbf{v} = \mathbf{0}$.
($\Leftarrow$) If $\ker(T) = \{\mathbf{0}\}$ and $T\mathbf{v} = T\mathbf{w}$, then $T(\mathbf{v} - \mathbf{w}) = \mathbf{0}$. Hence $\mathbf{v} - \mathbf{w} \in \ker(T)$, so $\mathbf{v} = \mathbf{w}$.
*Structural Commentary*: The kernel measures exactly where a linear map collapses vectors to zero.
*Key Insight*: Injectivity is equivalent to collapsing only the zero vector. □

**Proposition 1.6.4.** *Kernel and Range Are Subspaces*
*If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $\ker(T)$ is a subspace of $\mathcal{V}$ and $\text{range}(T)$ is a subspace of $\mathcal{W}$.*

*Proof.* Since $T\mathbf{0}_{\mathcal{V}} = \mathbf{0}_{\mathcal{W}}$, the kernel is nonempty.
If $\mathbf{u}, \mathbf{v} \in \ker(T)$ and $a \in \mathbb{F}$, then $T(a\mathbf{u} + \mathbf{v}) = aT\mathbf{u} + T\mathbf{v} = \mathbf{0}_{\mathcal{W}}$, so $a\mathbf{u} + \mathbf{v} \in \ker(T)$.
For the range, if $T\mathbf{u}, T\mathbf{v} \in \text{range}(T)$ and $a \in \mathbb{F}$, then $aT\mathbf{u} + T\mathbf{v} = T(a\mathbf{u} + \mathbf{v}) \in \text{range}(T)$.
*Structural Commentary*: The kernel and range are the two canonical subspaces attached to a linear map.
*Key Insight*: Linearity makes both preimages of zero and images of vectors closed under linear combinations. □




Figure 10: For a linear map $T : \mathcal{V} \to \mathcal{W}$, the range of $T$ may be a proper subspace of $\mathcal{W}$, and the kernel may contain nonzero vectors. The conditions $\text{range}(T) = \mathcal{W}$ and $\ker(T) = \{\mathbf{0}\}$ characterize surjectivity and injectivity, respectively.

**Definition 1.6.5.** *Invertible Linear Map and Isomorphism*
A linear map $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ is *invertible* if there exists $S \in \mathcal{L}(\mathcal{W}, \mathcal{V})$ such that $ST = I_{\mathcal{V}}$ and $TS = I_{\mathcal{W}}$.
An invertible linear map is called an *isomorphism*, and if one exists we write $\mathcal{V} \cong \mathcal{W}$.

**Proposition 1.6.6.** *Inverse Is Unique*
*An invertible linear map has a unique inverse. Thus if $T$ is invertible, its inverse is denoted $T^{-1}$.*

*Proof.* Suppose $S_1, S_2 \in \mathcal{L}(\mathcal{W}, \mathcal{V})$ are inverses of $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.
Then $S_1 = S_1 I_{\mathcal{W}} = S_1(TS_2) = (S_1T)S_2 = I_{\mathcal{V}}S_2 = S_2$. Hence the inverse is unique.
*Structural Commentary*: A two-sided inverse is forced by the composition identities.
*Key Insight*: Associativity lets the two candidate inverses meet around $T$ and collapse to the same map. □

**Proposition 1.6.7.** *Inverse of a Product Reverses Order*
*Let $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ and $S \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ be invertible. Then $ST$ is invertible and $(ST)^{-1} = T^{-1}S^{-1}$.*

*Proof.* Since $T^{-1} \in \mathcal{L}(\mathcal{V}, \mathcal{U})$ and $S^{-1} \in \mathcal{L}(\mathcal{W}, \mathcal{V})$, the product $T^{-1}S^{-1} \in \mathcal{L}(\mathcal{W}, \mathcal{U})$ is defined.
Moreover, $(T^{-1}S^{-1})(ST) = T^{-1}(S^{-1}S)T = T^{-1}T = I_{\mathcal{U}}$ and $(ST)(T^{-1}S^{-1}) = S(TT^{-1})S^{-1} = SS^{-1} = I_{\mathcal{W}}$.
Thus $T^{-1}S^{-1}$ is a two-sided inverse of $ST$, so $(ST)^{-1} = T^{-1}S^{-1}$.
*Structural Commentary*: To undo a composition, undo the last map first.
*Key Insight*: Composition order reverses under inversion. □

**Proposition 1.6.8.** *Invertibility and Bijectivity*
*A linear map is invertible if and only if it is both injective and surjective.*

*Proof.*

($\Rightarrow$) Let $S$ be an inverse of $T$. If $T\mathbf{v} = T\mathbf{w}$, then $\mathbf{v} = ST\mathbf{v} = ST\mathbf{w} = \mathbf{w}$, so $T$ is injective. Also every $\mathbf{w}$ in the codomain satisfies $\mathbf{w} = T(S\mathbf{w})$, so $T$ is surjective.

($\Leftarrow$) Suppose $T$ is injective and surjective. Define $S\mathbf{w}$ to be the unique vector $\mathbf{v}$ with $T\mathbf{v} = \mathbf{w}$. Since both sides below map under $T$ to $a\mathbf{w}_1 + \mathbf{w}_2$, $S(a\mathbf{w}_1 + \mathbf{w}_2) = aS\mathbf{w}_1 + S\mathbf{w}_2$. Thus $S$ is linear and satisfies $ST = I$ and $TS = I$, so $T$ is invertible.

*Structural Commentary*: Bijectivity gives a unique reverse map, and linearity makes it a linear inverse.
*Key Insight*: A linear map is invertible exactly when it is one-to-one and onto. □

**Definition 1.6.9.** *Morphism Terminology for Linear Maps*
For $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$:

(i) $T$ is a *homomorphism*.
(ii) If $\mathcal{V} = \mathcal{W}$, then $T$ is an *endomorphism*.
(iii) If $\ker(T) = \{\mathbf{0}\}$, then $T$ is a *monomorphism*, or *embedding*.
(iv) If $\text{range}(T) = \mathcal{W}$, then $T$ is an *epimorphism*.
(v) If $\ker(T) = \{\mathbf{0}\}$ and $\text{range}(T) = \mathcal{W}$, then $T$ is an *isomorphism*.
(vi) If $\mathcal{V} = \mathcal{W}$ and $T$ is invertible, then $T$ is an *automorphism*.

## 1.7 Rank-Nullity and Dimension Criteria

**Theorem 1.7.1.** *Range from Basis Images*
*Let $\mathcal{V}$ be finite-dimensional with basis $(\mathbf{v}_1, \dots, \mathbf{v}_n)$, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then* $\text{range}(T) = \text{span}(T\mathbf{v}_1, \dots, T\mathbf{v}_n)$. *In particular,* $\text{range}(T)$ *is finite-dimensional and* $\dim(\text{range}(T)) \leq n$.

*Proof.* If $\mathbf{v} = c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n$, then $T\mathbf{v} = c_1T\mathbf{v}_1 + \cdots + c_nT\mathbf{v}_n$.
Thus every vector in $\text{range}(T)$ lies in $\text{span}(T\mathbf{v}_1, \dots, T\mathbf{v}_n)$, and the reverse inclusion is immediate from the definition of range. The dimension bound follows because this spanning list has length $n$.
*Structural Commentary*: The range is generated by the images of any basis of the domain.
*Key Insight*: Linear maps carry a basis-spanning formula directly into a range-spanning formula. □

**Definition 1.7.2.** *Rank and Nullity*
For a finite-dimensional domain, the rank of $T$ is $\dim(\text{range}(T))$ and the nullity of $T$ is $\dim(\ker(T))$.

**Theorem 1.7.3.** *Rank-Nullity Theorem*
*If $\mathcal{V}$ is finite-dim and $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then* $\text{range}(T)$ *is finite-dim and* $\dim(\mathcal{V}) = \dim(\ker(T)) + \dim(\text{range}(T))$.

*Proof.* By Proposition 1.6.4 and Corollary 1.3.13, $\ker(T)$ has a basis $(\mathbf{u}_1, \dots, \mathbf{u}_m)$.
Extend it to a basis $(\mathbf{u}_1, \dots, \mathbf{u}_m, \mathbf{w}_1, \dots, \mathbf{w}_r)$ of $\mathcal{V}$ by Theorem 1.3.12.
If $\mathbf{v} = \sum_{i=1}^{m} a_i\mathbf{u}_i + \sum_{j=1}^{r} b_j\mathbf{w}_j$, then $T\mathbf{v} = \sum_{j=1}^{r} b_jT\mathbf{w}_j$, because $T\mathbf{u}_i = \mathbf{0}$. Hence $(T\mathbf{w}_1, \dots, T\mathbf{w}_r)$ spans $\text{range}(T)$.
If $\sum_{j=1}^{r} c_jT\mathbf{w}_j = \mathbf{0}$, then $\sum_{j=1}^{r} c_j\mathbf{w}_j \in \ker(T)$, so for some scalars $d_i$, $\sum_{j=1}^{r} c_j\mathbf{w}_j = \sum_{i=1}^{m} d_i\mathbf{u}_i$.
Thus $\sum_{j=1}^{r} c_j\mathbf{w}_j - \sum_{i=1}^{m} d_i\mathbf{u}_i = \mathbf{0}$, and linear independence of extended basis forces every $c_j$ and $d_i$ to be zero.
Hence $(T\mathbf{w}_1, \dots, T\mathbf{w}_r)$ is a basis of $\text{range}(T)$. Thus $\dim(\ker(T)) = m$, $\dim(\text{range}(T)) = r$, and $\dim(\mathcal{V}) = m + r$.
*Structural Commentary*: An adapted basis splits $\mathcal{V}$ into kernel directions and directions detected by $T$.
*Key Insight*: Rank-nullity says every domain dimension is either killed by $T$ or appears in its range. □

**Corollary 1.7.4.** *Maps to Smaller Spaces Are Not Injective*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional vector spaces with* $\dim(\mathcal{V}) > \dim(\mathcal{W})$.
*Then no linear map from $\mathcal{V}$ to $\mathcal{W}$ is injective.*

*Proof.* Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. By Theorem 1.7.3, $\dim(\ker(T)) = \dim(\mathcal{V}) - \dim(\text{range}(T)) \geq \dim(\mathcal{V}) - \dim(\mathcal{W}) > 0$.
Thus $\ker(T)$ contains a nonzero vector, so $T$ is not injective by Proposition 1.6.3.
*Structural Commentary*: Excess domain dimension forces collapse.
*Key Insight*: The dimension deficit must appear inside the kernel. □

**Corollary 1.7.5.** *Maps to Larger Spaces Are Not Surjective*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional vector spaces with* $\dim(\mathcal{V}) < \dim(\mathcal{W})$.
*Then no linear map from $\mathcal{V}$ to $\mathcal{W}$ is surjective.*

*Proof.* Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. By Theorem 1.7.3, $\dim(\text{range}(T)) = \dim(\mathcal{V}) - \dim(\ker(T)) \leq \dim(\mathcal{V}) < \dim(\mathcal{W})$. Hence $\text{range}(T)$ is a proper subspace of $\mathcal{W}$, so $T$ is not surjective.
*Structural Commentary*: A smaller domain cannot cover a larger codomain.
*Key Insight*: The range can never have dimension larger than the domain. □

**Corollary 1.7.6.** *Equal-Dimension Kernel-Range Criterion*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional vector spaces with $\dim(\mathcal{V}) = \dim(\mathcal{W})$, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then $\ker(T) = \{\mathbf{0}\}$ if and only if $\text{range}(T) = \mathcal{W}$. Equivalently, $T$ is injective if and only if $T$ is surjective.*

*Proof.*
($\Rightarrow$) If $\ker(T) = \{\mathbf{0}\}$, then rank-nullity gives $\dim(\text{range}(T)) = \dim(\mathcal{V}) = \dim(\mathcal{W})$.
Since $\text{range}(T)$ is a subspace of $\mathcal{W}$ with the same dimension, $\text{range}(T) = \mathcal{W}$.
($\Leftarrow$) If $\text{range}(T) = \mathcal{W}$, then rank-nullity gives $\dim(\ker(T)) = \dim(\mathcal{V}) - \dim(\mathcal{W}) = 0$, hence $\ker(T) = \{\mathbf{0}\}$.
*Structural Commentary*: Equal finite dimensions make injectivity and surjectivity equivalent.
*Key Insight*: In equal dimensions, having no kernel loss is exactly having full range. □

**Proposition 1.7.7.** *Isomorphisms Preserve Bases*
*Let $T : \mathcal{V} \to \mathcal{W}$ be an isomorphism and let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be a finite list in $\mathcal{V}$. Set $T\beta = (T\mathbf{v}_1, \ldots, T\mathbf{v}_n)$. The following are equivalent:*

*(i) $\beta$ is a basis of $\mathcal{V}$.*
*(ii) $T\beta$ is a basis of $\mathcal{W}$.*

*Proof.*
($i$) $\Rightarrow$ ($ii$) If $\mathbf{w} \in \mathcal{W}$, choose $\mathbf{v} \in \mathcal{V}$ with $\mathbf{w} = T\mathbf{v}$ and write $\mathbf{v} = \sum_j a_j \mathbf{v}_j$. Then $\mathbf{w} = \sum_j a_j T\mathbf{v}_j$, so $T\beta$ spans $\mathcal{W}$. If $\sum_j a_j T\mathbf{v}_j = \mathbf{0}$, applying $T^{-1}$ gives $\sum_j a_j \mathbf{v}_j = \mathbf{0}$; hence all $a_j = 0$. Thus $T\beta$ is a basis.
($ii$) $\Rightarrow$ ($i$) Assume $T\beta$ is a basis of $\mathcal{W}$. If $\mathbf{v} \in \mathcal{V}$, write $T\mathbf{v} = \sum_j a_j T\mathbf{v}_j$; applying $T^{-1}$ gives $\mathbf{v} = \sum_j a_j \mathbf{v}_j$, so $\beta$ spans $\mathcal{V}$. If $\sum_j a_j \mathbf{v}_j = \mathbf{0}$, applying $T$ gives $\sum_j a_j T\mathbf{v}_j = \mathbf{0}$; hence all $a_j = 0$. Thus $\beta$ is linearly independent, and therefore a basis of $\mathcal{V}$.
*Structural Commentary*: Isomorphisms preserve exactly the spanning and independence data that define a basis.
*Key Insight*: An isomorphism changes the ambient space, not the linear relations among vectors. □

**Theorem 1.7.8.** *Finite-Dimensional Invertibility Criteria*
*If $\mathcal{V}$ is finite-dimensional and $T \in \mathcal{L}(\mathcal{V})$, then the following are equivalent:*

*i. $T$ is invertible;*
*ii. $T$ is injective;*
*iii. $T$ is surjective.*

*Proof.*
($i$) $\Rightarrow$ ($ii$) follows from Proposition 1.6.8.
($ii$) $\Rightarrow$ ($iii$): If $T$ is injective, then $\ker(T) = \{\mathbf{0}\}$.
Rank-nullity gives $\dim(\text{range}(T)) = \dim(\mathcal{V})$, hence $\text{range}(T) = \mathcal{V}$, so $T$ is surjective.
($iii$) $\Rightarrow$ ($i$): If $T$ is surjective, rank-nullity gives $\dim(\ker(T)) = 0$, so $T$ is injective.
Thus $T$ is bijective, hence invertible by Proposition 1.6.8.
*Structural Commentary*: For operators on finite-dimensional spaces, kernel and range dimensions trade off exactly. Injectivity and surjectivity therefore force each other.
*Key Insight*: In equal finite dimensions, losing no information is the same as reaching the whole space. □

**Perspective 1.7.9.** *Kernel Obstruction to Invertibility*
By Theorem 1.7.8 and Proposition 1.6.3, a finite-dimensional operator is invertible exactly when its kernel is zero. Any nonzero $\mathbf{v} \in \ker T$ therefore blocks invertibility.

**Theorem 1.7.10.** *Isomorphism and Dimension*
*Two finite-dimensional vector spaces are isomorphic if and only if they have the same dimension.*

*Proof.*
($\Rightarrow$) If $T : \mathcal{V} \to \mathcal{W}$ is an isomorphism and $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$, then Proposition 1.7.7 shows that $(T\mathbf{v}_1, \ldots, T\mathbf{v}_n)$ is a basis of $\mathcal{W}$.
($\Leftarrow$) If both spaces have dimension $n$, choose bases $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ and $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$.
By Proposition 1.5.11, there is a linear map $T$ with $T\mathbf{v}_j = \mathbf{w}_j$ for every $j$. Then $T(\sum a_j \mathbf{v}_j) = \sum a_j \mathbf{w}_j$, so $\ker(T) = \{\mathbf{0}\}$ and $\text{range}(T) = \mathcal{W}$. Hence $T$ is an isomorphism.

*Structural Commentary*: Dimension classifies finite-dimensional spaces up to isomorphism.
*Key Insight*: Matching bases give an isomorphism; any isomorphism carries bases to bases. □

# 2 Direct Sums, Quotients, and Decomposition

## 2.1 Sums and Direct Sums

Direct sums split spaces into complementary pieces; quotients collapse selected directions. Dimension formulas keep both constructions controlled.

**Definition 2.1.1.** *Sums of Subspaces*
Let $\mathcal{U}_1, \dots, \mathcal{U}_m \leq \mathcal{V}$.

(i) The *sum* of $\mathcal{U}_1, \dots, \mathcal{U}_m$ is $\mathcal{U}_1 + \cdots + \mathcal{U}_m := \{\mathbf{u}_1 + \cdots + \mathbf{u}_m : \mathbf{u}_j \in \mathcal{U}_j\}$.
(ii) For subspaces $\mathcal{U}, \mathcal{W} \leq \mathcal{V}$, $\mathcal{U} + \mathcal{W} = \operatorname{span}(\mathcal{U} \cup \mathcal{W}) = \{\mathbf{u} + \mathbf{w} : \mathbf{u} \in \mathcal{U},\ \mathbf{w} \in \mathcal{W}\}$.
(iii) The sum is *direct* if every vector in it has a unique decomposition as $\mathbf{u}_1 + \cdots + \mathbf{u}_m$ with $\mathbf{u}_j \in \mathcal{U}_j$; in this case it is denoted $\mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$.

**Proposition 2.1.2.** *Sum of Subspaces Is the Smallest Containing Subspace*
*Suppose $\mathcal{U}_1, \dots, \mathcal{U}_m$ are subspaces of $\mathcal{V}$. Then $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ is the smallest subspace of $\mathcal{V}$ containing $\mathcal{U}_1, \dots, \mathcal{U}_m$.*

*Proof.* Since $\mathbf{0} \in \mathcal{U}_j$ for each $j$, we have $\mathbf{0} \in \mathcal{U}_1 + \cdots + \mathcal{U}_m$. If $\mathbf{x} = \mathbf{u}_1 + \cdots + \mathbf{u}_m$ and $\mathbf{y} = \mathbf{v}_1 + \cdots + \mathbf{v}_m$ lie in $\mathcal{U}_1 + \cdots + \mathcal{U}_m$, then for any $c \in \mathbb{F}$, $c\mathbf{x} + \mathbf{y} = (c\mathbf{u}_1 + \mathbf{v}_1) + \cdots + (c\mathbf{u}_m + \mathbf{v}_m) \in \mathcal{U}_1 + \cdots + \mathcal{U}_m$, because $c\mathbf{u}_j + \mathbf{v}_j \in \mathcal{U}_j$ for each $j$. By Theorem 1.2.2, $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ is a subspace of $\mathcal{V}$.
Each $\mathcal{U}_j$ is contained in $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ by taking all other summands to be $\mathbf{0}$. If $\mathcal{W}$ is any subspace of $\mathcal{V}$ containing $\mathcal{U}_1, \dots, \mathcal{U}_m$, then closure under addition gives $\mathbf{u}_1 + \cdots + \mathbf{u}_m \in \mathcal{W}$ whenever $\mathbf{u}_j \in \mathcal{U}_j$.
Hence $\mathcal{U}_1 + \cdots + \mathcal{U}_m \subseteq \mathcal{W}$, so the sum is the smallest such subspace.
*Structural Commentary*: Sums of subspaces are obtained by closing several subspaces under finite addition. Minimality follows because every containing subspace is already closed under those sums.
*Key Insight*: The sum adds no extra structure beyond the linear combinations forced by the given subspaces. □

**Perspective 2.1.3.** *Sum Is Not Union*
The union $\mathcal{U} \cup \mathcal{W}$ keeps vectors from either subspace; the sum $\mathcal{U} + \mathcal{W}$ adds vectors from both. Unless one subspace contains the other, $\mathcal{U} \cup \mathcal{W}$ is not closed under addition.

**Theorem 2.1.4.** *Finite Unions of Proper Subspaces*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be over an infinite field $\mathbb{F}$. If $\mathcal{U}_1, \dots, \mathcal{U}_n \subsetneq \mathcal{V}$ are subspaces, then $\mathcal{V} \neq \bigcup_{j=1}^{n} \mathcal{U}_j$.*

*Proof.* Assume for contradiction that $\mathcal{V} = \bigcup_{j=1}^{n} \mathcal{U}_j$. After deleting redundant subspaces and reindexing, assume the cover is minimal, so $\mathcal{U}_i \not\subseteq \bigcup_{j \neq i} \mathcal{U}_j$. If $n = 1$, then $\mathcal{V} = \mathcal{U}_1$, contradicting $\mathcal{U}_1 \subsetneq \mathcal{V}$; hence $n \geq 2$.

Choose $\mathbf{w} \in \mathcal{U}_1 \setminus \bigcup_{j=2}^{n} \mathcal{U}_j$ and $\mathbf{v} \in \mathcal{V} \setminus \mathcal{U}_1$. Since $\mathbf{0} \in \mathcal{U}_j$ for every $j$, we have $\mathbf{w} \neq \mathbf{0}$.

Let $A = \{\mathbf{v} + r\mathbf{w} : r \in \mathbb{F}\}$. Then $|A| = |\mathbb{F}| = \infty$, $A \cap \mathcal{U}_1 = \emptyset$, and for $j \geq 2$, $|A \cap \mathcal{U}_j| \leq 1$, since two points $\mathbf{v} + r_1\mathbf{w}, \mathbf{v} + r_2\mathbf{w} \in \mathcal{U}_j$ with $r_1 \neq r_2$ imply $\mathbf{w} \in \mathcal{U}_j$.

Hence $\left| A \cap \bigcup_{j=1}^{n} \mathcal{U}_j \right| \leq n - 1 < \infty$, contradicting $A \subseteq \mathcal{V} = \bigcup_{j=1}^{n} \mathcal{U}_j$.
*Structural Commentary*: Over an infinite field, finite proper subspaces cannot cover a well-chosen affine line.
*Key Insight*: A single one-parameter translate has too many points for finitely many proper intersections. □

**Perspective 2.1.5.** *Sums Always Exist; Directness Is Extra*
For subspaces $\mathcal{U}, \mathcal{W} \subseteq \mathcal{V}$, the sum $\mathcal{U} + \mathcal{W}$ is always a subspace by Proposition 2.1.2. Writing $\mathcal{U} \oplus \mathcal{W}$ asserts the extra condition $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$, equivalently uniqueness of the decomposition $\mathbf{u} + \mathbf{w}$.

**Proposition 2.1.6.** *Two-Subspace Directness Criterion*
*Let $\mathcal{U}$ and $\mathcal{W}$ be subspaces of $\mathcal{V}$. Then $\mathcal{U} + \mathcal{W}$ is direct if and only if $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$.*

*Proof.*

(⇒) Suppose $\mathcal{U} + \mathcal{W}$ is direct. If $\mathbf{x} \in \mathcal{U} \cap \mathcal{W}$, then $\mathbf{x} = \mathbf{x} + \mathbf{0} = \mathbf{0} + \mathbf{x}$ gives two decompositions of $\mathbf{x}$ with one term in each subspace. Uniqueness gives $\mathbf{x} = \mathbf{0}$, so $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$.

($\Leftarrow$) Suppose $\mathcal{U}\cap\mathcal{W}=\{\mathbf{0}\}$. Every vector in $\mathcal{U}+\mathcal{W}$ has at least one representation $\mathbf{u}+\mathbf{w}$. If $\mathbf{u}_1+\mathbf{w}_1=\mathbf{u}_2+\mathbf{w}_2$, with $\mathbf{u}_1,\mathbf{u}_2\in\mathcal{U}$ and $\mathbf{w}_1,\mathbf{w}_2\in\mathcal{W}$, then $\mathbf{u}_1-\mathbf{u}_2=\mathbf{w}_2-\mathbf{w}_1\in\mathcal{U}\cap\mathcal{W}$, hence equals $\mathbf{0}$.
Thus $\mathbf{u}_1=\mathbf{u}_2$ and $\mathbf{w}_1=\mathbf{w}_2$, so the sum is direct.

*Structural Commentary*: Trivial intersection prevents cancellation between summands.
*Key Insight*: Directness means unique decomposition. □

**Definition 2.1.7.** *Subspace Poset, Meet, and Join*

(i) The *subspace poset* of $\mathcal{V}$ is $\mathrm{Sub}(\mathcal{V}):=\{\mathcal{U}\subseteq\mathcal{V}:\mathcal{U}\leq\mathcal{V}\}$, ordered by inclusion.
(ii) For $\{\mathcal{U}_i:i\in I\}\subseteq\mathrm{Sub}(\mathcal{V})$, the *meet* is $\bigwedge_{i\in I}\mathcal{U}_i:=\bigcap_{i\in I}\mathcal{U}_i$.
(iii) For $\{\mathcal{U}_i:i\in I\}\subseteq\mathrm{Sub}(\mathcal{V})$, the *join* is $\bigvee_{i\in I}\mathcal{U}_i:=\sum_{i\in I}\mathcal{U}_i:=\mathrm{span}\left(\bigcup_{i\in I}\mathcal{U}_i\right)$.
(iv) The empty meet is $\mathcal{V}$, and the empty join is $\{\mathbf{0}\}$.

**Proposition 2.1.8.** *Subspace Meet and Join*
*For subspaces $\{\mathcal{U}_i:i\in I\}$ of $\mathcal{V}$, ordered by inclusion, $\bigwedge_{i\in I}\mathcal{U}_i=\inf\{\mathcal{U}_i:i\in I\}$; and $\bigvee_{i\in I}\mathcal{U}_i=\sup\{\mathcal{U}_i:i\in I\}$.*

*Proof.* If $I=\emptyset$, conventions give vacuous bounds $\mathcal{V}$ and $\{\mathbf{0}\}$. Assume $I\neq\emptyset$.
The meet $M=\bigcap_{i\in I}\mathcal{U}_i$ is a subspace by Proposition 1.2.3; moreover $M\subseteq\mathcal{U}_i$ for all $i$, and any subspace $\mathcal{W}$ with $\mathcal{W}\subseteq\mathcal{U}_i$ for all $i$ satisfies $\mathcal{W}\subseteq M$. The join $J=\mathrm{span}\left(\bigcup_{i\in I}\mathcal{U}_i\right)$ is a subspace by Theorem 1.2.5; moreover $\mathcal{U}_i\subseteq J$ for all $i$, and any subspace $\mathcal{W}$ with $\mathcal{U}_i\subseteq\mathcal{W}$ for all $i$ contains $\bigcup_{i\in I}\mathcal{U}_i$, hence contains $J$.
*Structural Commentary*: Meet enforces common constraints; join closes the combined directions under span.
*Key Insight*: Meet is intersection, while join is span of the union. □

**Theorem 2.1.9.** *Subspace Lattice*
*The subspace poset $\mathrm{Sub}(\mathcal{V})$ is a complete modular lattice.*
*Explicitly, for every $\{\mathcal{U}_i:i\in I\}\subseteq\mathrm{Sub}(\mathcal{V})$, $\bot=\{\mathbf{0}\}$, $\top=\mathcal{V}$, $\bigwedge_{i\in I}\mathcal{U}_i=\bigcap_{i\in I}\mathcal{U}_i$, $\bigvee_{i\in I}\mathcal{U}_i=\sum_{i\in I}\mathcal{U}_i=\mathrm{span}\left(\bigcup_{i\in I}\mathcal{U}_i\right)$.*
*Moreover, if $\mathcal{U}\subseteq\mathcal{S}$, then $\mathcal{S}\cap(\mathcal{T}+\mathcal{U})=(\mathcal{S}\cap\mathcal{T})+\mathcal{U}$.*

*Proof.* Proposition 2.1.8 gives arbitrary meets and joins with the displayed formulas; the empty meet and join give $\top=\mathcal{V}$ and $\bot=\{\mathbf{0}\}$. It remains to prove modularity. Assume $\mathcal{U}\subseteq\mathcal{S}$. If $\mathbf{x}\in(\mathcal{S}\cap\mathcal{T})+\mathcal{U}$, then $\mathbf{x}=\mathbf{y}+\mathbf{u}$ with $\mathbf{y}\in\mathcal{S}\cap\mathcal{T}$ and $\mathbf{u}\in\mathcal{U}$, hence $\mathbf{x}\in\mathcal{S}\cap(\mathcal{T}+\mathcal{U})$. Conversely, if $\mathbf{x}\in\mathcal{S}\cap(\mathcal{T}+\mathcal{U})$, write $\mathbf{x}=\mathbf{t}+\mathbf{u}$ with $\mathbf{t}\in\mathcal{T}$ and $\mathbf{u}\in\mathcal{U}$. Since $\mathbf{x}\in\mathcal{S}$ and $\mathbf{u}\in\mathcal{U}\subseteq\mathcal{S}$, $\mathbf{t}=\mathbf{x}-\mathbf{u}\in\mathcal{S}\cap\mathcal{T}$, so $\mathbf{x}\in(\mathcal{S}\cap\mathcal{T})+\mathcal{U}$.
*Structural Commentary*: The subspace lattice is complete and modular, not generally distributive.
*Key Insight*: If one summand already lies inside $\mathcal{S}$, intersecting with $\mathcal{S}$ can be pushed through the sum. □




Figure 11: For two subspaces, intersection is the meet and sum is the join under inclusion.

**Proposition 2.1.10.** *Equivalent Conditions for a Direct Sum*
*For subspaces $\mathcal{U}_1,\ldots,\mathcal{U}_m\subseteq\mathcal{V}$, the following are equivalent:*

*(i) $\mathcal{V}=\mathcal{U}_1\oplus\cdots\oplus\mathcal{U}_m$;*
*(ii) every $\mathbf{v}\in\mathcal{V}$ has a unique representation $\mathbf{v}=\mathbf{u}_1+\cdots+\mathbf{u}_m$ with $\mathbf{u}_j\in\mathcal{U}_j$;*
*(iii) $\mathcal{V}=\mathcal{U}_1+\cdots+\mathcal{U}_m$, and if $\mathbf{u}_1+\cdots+\mathbf{u}_m=\mathbf{0}$ with $\mathbf{u}_j\in\mathcal{U}_j$, then $\mathbf{u}_1=\cdots=\mathbf{u}_m=\mathbf{0}$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) This follows from the definition of direct sum.
(*ii*) $\Rightarrow$ (*iii*) Every vector lies in $\mathcal{U}_1+\cdots+\mathcal{U}_m$, so $\mathcal{V}=\mathcal{U}_1+\cdots+\mathcal{U}_m$. If $\mathbf{u}_1+\cdots+\mathbf{u}_m=\mathbf{0}$, then this representation of $\mathbf{0}$ must equal $\mathbf{0}+\cdots+\mathbf{0}$ by uniqueness, so each $\mathbf{u}_j=\mathbf{0}$.
(*iii*) $\Rightarrow$ (*i*) Since $\mathcal{V}=\mathcal{U}_1+\cdots+\mathcal{U}_m$, every $\mathbf{v}\in\mathcal{V}$ has at least one representation.
If $\mathbf{v}=\mathbf{u}_1+\cdots+\mathbf{u}_m=\mathbf{w}_1+\cdots+\mathbf{w}_m$, then $(\mathbf{u}_1-\mathbf{w}_1)+\cdots+(\mathbf{u}_m-\mathbf{w}_m)=\mathbf{0}$. By (iii), $\mathbf{u}_j-\mathbf{w}_j=\mathbf{0}$ for every $j$, so the representation is unique. Hence $\mathcal{V}=\mathcal{U}_1\oplus\cdots\oplus\mathcal{U}_m$.

*Structural Commentary*: Direct sums require both spanning by the summands and no cancellation among them. The zero-representation condition isolates the no-cancellation part.
*Key Insight*: Uniqueness is equivalent to saying the only way the summands add to zero is trivially. □

**Theorem 2.1.11.** *Dimension of a Product*
*If $\mathcal{V}_1, \dots, \mathcal{V}_m$ are finite-dimensional vector spaces over $\mathbb{F}$, then $\mathcal{V}_1 \times \cdots \times \mathcal{V}_m$ is finite-dimensional, and $\dim(\mathcal{V}_1 \times \cdots \times \mathcal{V}_m) = \dim(\mathcal{V}_1) + \cdots + \dim(\mathcal{V}_m)$.*

*Proof.* For each $j$, let $(\mathbf{v}_{j,1}, \dots, \mathbf{v}_{j,n_j})$ be a basis of $\mathcal{V}_j$. For every $j$ and $k$, let $\mathbf{e}_{j,k}$ denote the element of $\mathcal{V}_1 \times \cdots \times \mathcal{V}_m$ whose $j$th component is $\mathbf{v}_{j,k}$ and whose other components are zero.
The list of all $\mathbf{e}_{j,k}$ spans the product because each component can be expanded in its own basis. If a linear combination of the $\mathbf{e}_{j,k}$ is zero, then looking at the $j$th component gives a linear relation among $(\mathbf{v}_{j,1}, \dots, \mathbf{v}_{j,n_j})$, so all coefficients in that component are zero. Hence the list is linearly independent and therefore is a basis of the product. Its length is $n_1 + \cdots + n_m$.
*Structural Commentary*: A basis for a product is formed by placing each component basis into its own slot.
*Key Insight*: Product dimension counts the independent directions contributed by each factor. □

**Definition 2.1.12.** *External Product*
Let $\{\mathcal{V}_i : i \in I\}$ be $\mathbb{F}$-vector spaces.

(i) The *external direct product* is $\prod_{i \in I} \mathcal{V}_i = \{\mathbf{v} : I \to \bigcup_{i \in I} \mathcal{V}_i : \mathbf{v}(i) \in \mathcal{V}_i\}$.

(ii) Operations are componentwise: $(\mathbf{u} + \mathbf{v})(i) = \mathbf{u}(i) + \mathbf{v}(i)$ and $(a\mathbf{v})(i) = a\mathbf{v}(i)$.

**Definition 2.1.13.** *Support and Finite Sums of a Family*
Let $I$ be an index set. For each $i \in I$, let $\mathcal{X}_i$ be a vector space with zero vector $\mathbf{0}_i$.

(i) *Support.* For $\mathbf{x} = (\mathbf{x}_i)_{i \in I} \in \prod_{i \in I} \mathcal{X}_i$, define $\operatorname{supp}(\mathbf{x}) := \{i \in I : \mathbf{x}_i \neq \mathbf{0}_i\}$.

(ii) The indexed family $\mathbf{x}$ is *finitely supported* iff $|\operatorname{supp}(\mathbf{x})| < \infty$.

(iii) *Finite sum.* If all $\mathcal{X}_i$ lie in a common vector space and $\mathbf{x}$ is finitely supported, define $\sum_{i \in I} \mathbf{x}_i := \sum_{i \in \operatorname{supp}(\mathbf{x})} \mathbf{x}_i$.

**Definition 2.1.14.** *External Direct Sum*
Let $\{\mathcal{V}_i : i \in I\}$ be $\mathbb{F}$-vector spaces. The *external direct sum* is $\bigoplus_{i \in I}^{\text{ext}} \mathcal{V}_i = \{\mathbf{v} \in \prod_{i \in I} \mathcal{V}_i : |\operatorname{supp}(\mathbf{v})| < \infty\}$. If $|I| < \infty$, then $\bigoplus_{i \in I}^{\text{ext}} \mathcal{V}_i = \prod_{i \in I} \mathcal{V}_i$.

**Proposition 2.1.15.** *External Sums Are Finite-Support Products*
*With componentwise operations, $\prod_{i \in I} \mathcal{V}_i$ is an $\mathbb{F}$-vector space, $\bigoplus_{i \in I}^{\text{ext}} \mathcal{V}_i = \{\mathbf{v} \in \prod_{i \in I} \mathcal{V}_i : |\operatorname{supp}(\mathbf{v})| < \infty\} \leq \prod_{i \in I} \mathcal{V}_i$.*

*Proof.* Let $P = \prod_{i \in I} \mathcal{V}_i$. For $\mathbf{u}, \mathbf{v} \in P$ and $a, b \in \mathbb{F}$, $(a\mathbf{u} + b\mathbf{v})(i) = a\mathbf{u}(i) + b\mathbf{v}(i) \in \mathcal{V}_i$.
The zero $0_P(i) = 0_i$, inverses $(-\mathbf{v})(i) = -\mathbf{v}(i)$, and all axioms are coordinatewise.
Let $S = \bigoplus_{i \in I}^{\text{ext}} \mathcal{V}_i$. If $\mathbf{u}, \mathbf{v} \in S$ and $a, b \in \mathbb{F}$, then $\operatorname{supp}(a\mathbf{u} + b\mathbf{v}) \subseteq \operatorname{supp}(\mathbf{u}) \cup \operatorname{supp}(\mathbf{v})$, so $a\mathbf{u} + b\mathbf{v} \in S$.
Also $0_P \in S$. Hence $S \leq P$.
*Structural Commentary*: $\prod_i \mathcal{V}_i$: arbitrary support; $\bigoplus_i^{\text{ext}} \mathcal{V}_i$: finite support.
*Key Insight*: Finite support is exactly what finite linear combinations preserve. □

**Definition 2.1.16.** *Internal Direct Sum of a Family*
For $\{\mathcal{U}_i : i \in I\} \subseteq \operatorname{Sub}(\mathcal{V})$, $\mathcal{V} = \bigoplus_{i \in I} \mathcal{U}_i$ means every $\mathbf{v} \in \mathcal{V}$ has a unique finitely supported $\mathbf{u} = (\mathbf{u}_i)_{i \in I}$ with $\mathbf{u}_i \in \mathcal{U}_i$ and $\mathbf{v} = \sum_{i \in I} \mathbf{u}_i$.
Equivalently, $\mathcal{V} = \sum_{i \in I} \mathcal{U}_i$ and every finite relation $\sum_{i \in F} \mathbf{u}_i = \mathbf{0}$, with $F \subseteq I$ finite and $\mathbf{u}_i \in \mathcal{U}_i$, has $\mathbf{u}_i = \mathbf{0}$ $\forall i \in F$.

**Proposition 2.1.17.** *Family Direct-Sum Criterion*
*For $\{\mathcal{U}_i\}_{i \in I} \subseteq \operatorname{Sub}(\mathcal{V})$, $\sum_{i \in I} \mathcal{U}_i$ is direct $\iff \forall i \in I$, $\mathcal{U}_i \cap \sum_{j \in I \setminus \{i\}} \mathcal{U}_j = \{\mathbf{0}\}$.*

*Proof.*

($\Rightarrow$) If $\mathbf{x} \in \mathcal{U}_i \cap \sum_{j \in I \setminus \{i\}} \mathcal{U}_j$, then $\mathbf{x} = \sum_{j \in F} \mathbf{u}_j$ for some finite $F \subseteq I \setminus \{i\}$ and $\mathbf{u}_j \in \mathcal{U}_j$.
Directness applied to $\mathbf{x} - \sum_{j \in F} \mathbf{u}_j = \mathbf{0}$ gives $\mathbf{x} = \mathbf{0}$.

($\Leftarrow$) If $\sum_{k=1}^{m} \mathbf{u}_{i_k} = \mathbf{0}$ with distinct $i_k$ and $\mathbf{u}_{i_k} \in \mathcal{U}_{i_k}$, then for each $k$, $\mathbf{u}_{i_k} = -\sum_{\ell \neq k} \mathbf{u}_{i_\ell} \in \mathcal{U}_{i_k} \cap \sum_{j \in I \setminus \{i_k\}} \mathcal{U}_j = \{\mathbf{0}\}$. Thus all finite zero-relations are trivial, so the sum is direct.

*Structural Commentary*: Family directness requires each summand to miss all others together.
*Key Insight*: Directness is subspace independence. □

**Proposition 2.1.18.** *Direct Sum Map*
*Suppose $\mathcal{U}_1, \ldots, \mathcal{U}_m$ are subspaces of $\mathcal{V}$. Define $\Gamma : \mathcal{U}_1 \times \cdots \times \mathcal{U}_m \to \mathcal{U}_1 + \cdots + \mathcal{U}_m$ by $\Gamma(\mathbf{u}_1, \ldots, \mathbf{u}_m) = \mathbf{u}_1 + \cdots + \mathbf{u}_m$. Then $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ is a direct sum if and only if $\Gamma$ is injective.*

*Proof.*

($\Rightarrow$) Suppose $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ is direct. If $\Gamma(\mathbf{u}_1, \ldots, \mathbf{u}_m) = \mathbf{0}$, then $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$, so directness gives $\mathbf{u}_j = \mathbf{0}$ for all $j$. Hence $\ker \Gamma = \{(\mathbf{0}, \ldots, \mathbf{0})\}$, and $\Gamma$ is injective.

($\Leftarrow$) Suppose $\Gamma$ is injective. If $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$ with $\mathbf{u}_j \in \mathcal{U}_j$, then $\Gamma(\mathbf{u}_1, \ldots, \mathbf{u}_m) = \Gamma(\mathbf{0}, \ldots, \mathbf{0})$. Injectivity gives $\mathbf{u}_j = \mathbf{0}$ for all $j$, so every zero decomposition is trivial. Hence the sum is direct.

*Structural Commentary*: The addition map identifies decompositions precisely through its kernel.
*Key Insight*: Directness is injectivity of $\Gamma$. □




Figure 12: Intersection, sum, and product of subspaces.

## 2.2 Quotient Spaces

**Definition 2.2.1.** *Affine Subset and Parallel Subspaces*
Let $\mathcal{U}$ be a subspace of $\mathcal{V}$. For $\mathbf{v} \in \mathcal{V}$, the subset $\mathbf{v} + \mathcal{U} := \{\mathbf{v} + \mathbf{u} : \mathbf{u} \in \mathcal{U}\}$ is an *affine subset* of $\mathcal{V}$ parallel to $\mathcal{U}$.

**Proposition 2.2.2.** *Parallel Affine Subsets Are Equal or Disjoint*
*Let $\mathcal{U}$ be a subspace of $\mathcal{V}$ and let $\mathbf{v}, \mathbf{w} \in \mathcal{V}$. The following are equivalent:*

*(i)* $\mathbf{v} - \mathbf{w} \in \mathcal{U}$;
*(ii)* $\mathbf{v} + \mathcal{U} = \mathbf{w} + \mathcal{U}$;
*(iii)* $(\mathbf{v} + \mathcal{U}) \cap (\mathbf{w} + \mathcal{U}) \neq \emptyset$.

*Proof.*

*(i)* $\Rightarrow$ *(ii)* If $\mathbf{v} - \mathbf{w} \in \mathcal{U}$ and $\mathbf{u} \in \mathcal{U}$, then $\mathbf{v} + \mathbf{u} = \mathbf{w} + ((\mathbf{v} - \mathbf{w}) + \mathbf{u}) \in \mathbf{w} + \mathcal{U}$. Thus $\mathbf{v} + \mathcal{U} \subseteq \mathbf{w} + \mathcal{U}$, and the reverse inclusion is similar.

*(ii)* $\Rightarrow$ *(iii)* Equal affine subsets have the same elements.

*(iii)* $\Rightarrow$ *(i)* If $\mathbf{x} \in (\mathbf{v} + \mathcal{U}) \cap (\mathbf{w} + \mathcal{U})$, then $\mathbf{x} = \mathbf{v} + \mathbf{u}_1 = \mathbf{w} + \mathbf{u}_2$ for some $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}$. Hence $\mathbf{v} - \mathbf{w} = \mathbf{u}_2 - \mathbf{u}_1 \in \mathcal{U}$.

*Structural Commentary*: Parallel affine subsets either coincide or do not meet.
*Key Insight*: Intersection is controlled by the difference of base points. □

**Definition 2.2.3.** *Quotient Space*
Let $\mathcal{U}$ be a subspace of $\mathcal{V}$.
The *quotient space* $\mathcal{V}/\mathcal{U}$ is the set of all affine subsets of $\mathcal{V}$ parallel to $\mathcal{U}$: $\mathcal{V}/\mathcal{U} := \{\mathbf{v} + \mathcal{U} : \mathbf{v} \in \mathcal{V}\}$.

**Proposition 2.2.4.** *Quotient Relation Is a Linear Congruence*
*Let $\mathcal{U} \leq \mathcal{V}$ and define $\mathbf{v} \equiv_{\mathcal{U}} \mathbf{w} \iff \mathbf{v} - \mathbf{w} \in \mathcal{U}$.*
*Then $\equiv_{\mathcal{U}}$ is an equivalence relation and, for $a, b \in \mathbb{F}$, $\mathbf{v}_1 \equiv_{\mathcal{U}} \mathbf{v}_2$, $\mathbf{w}_1 \equiv_{\mathcal{U}} \mathbf{w}_2 \implies a\mathbf{v}_1 + b\mathbf{w}_1 \equiv_{\mathcal{U}} a\mathbf{v}_2 + b\mathbf{w}_2$.*

*Proof.* Since $\mathcal{U} \leq \mathcal{V}$, $\mathbf{0} \in \mathcal{U}$, $-\mathcal{U} = \mathcal{U}$, and $\mathcal{U} + \mathcal{U} \subseteq \mathcal{U}$; hence $\equiv_{\mathcal{U}}$ is reflexive, symmetric, and transitive.
If $\mathbf{v}_1 \equiv_{\mathcal{U}} \mathbf{v}_2$ and $\mathbf{w}_1 \equiv_{\mathcal{U}} \mathbf{w}_2$, then $(a\mathbf{v}_1 + b\mathbf{w}_1) - (a\mathbf{v}_2 + b\mathbf{w}_2) = a(\mathbf{v}_1 - \mathbf{v}_2) + b(\mathbf{w}_1 - \mathbf{w}_2) \in \mathcal{U}$.
*Structural Commentary*: Subspace congruence makes quotient operations well-defined.
*Key Insight*: Subspace closure is exactly the well-definedness mechanism for quotient operations. □

**Perspective 2.2.5.** *Cosets as Equivalence Classes*
The quotient relation is $\mathbf{v} \equiv_{\mathcal{U}} \mathbf{w}$ when $\mathbf{v} - \mathbf{w} \in \mathcal{U}$. Proposition 2.2.2 shows that the equivalence class of $\mathbf{v}$ is exactly the coset $\mathbf{v} + \mathcal{U}$. Thus the quotient space remembers vectors only up to directions in $\mathcal{U}$.

**Proposition 2.2.6.** *Structure of the Quotient*
*Let $\mathcal{U}$ be a subspace of $\mathcal{V}$. Addition and scalar multiplication on $\mathcal{V}/\mathcal{U}$ defined by $(\mathbf{v}+\mathcal{U})+(\mathbf{w}+\mathcal{U}) := (\mathbf{v}+\mathbf{w})+\mathcal{U}$ and $a(\mathbf{v}+\mathcal{U}) := a\mathbf{v}+\mathcal{U}$ are well-defined, and with these operations $\mathcal{V}/\mathcal{U}$ is a vector space.*

*Proof.* If $\mathbf{v}+\mathcal{U} = \mathbf{v}'+\mathcal{U}$ and $\mathbf{w}+\mathcal{U} = \mathbf{w}'+\mathcal{U}$, then $\mathbf{v}-\mathbf{v}' \in \mathcal{U}$ and $\mathbf{w}-\mathbf{w}' \in \mathcal{U}$.
Since $\mathcal{U}$ is a subspace, $(\mathbf{v}+\mathbf{w})-(\mathbf{v}'+\mathbf{w}') \in \mathcal{U}$, and $a\mathbf{v}-a\mathbf{v}' \in \mathcal{U}$ so the operations do not depend on the chosen representatives. The vector space axioms are inherited from $\mathcal{V}$ under these operations.
*Structural Commentary*: Quotients preserve operations modulo $\mathcal{U}$.
*Key Insight*: Cosets identify vectors differing by elements of $\mathcal{U}$. □

**Definition 2.2.7.** *Quotient Map*
Let $\mathcal{U}$ be a subspace of $\mathcal{V}$. The *quotient map* is the map $\pi : \mathcal{V} \to \mathcal{V}/\mathcal{U}$ defined by $\pi(\mathbf{v}) = \mathbf{v}+\mathcal{U}$.

**Proposition 2.2.8.** *Properties of the Quotient Map*
*The quotient map $\pi$ is linear, surjective, and $\ker(\pi) = \mathcal{U}$.*

*Proof.* For $\mathbf{v}, \mathbf{w} \in \mathcal{V}$ and $a \in \mathbb{F}$, $\pi(a\mathbf{v}+\mathbf{w}) = (a\mathbf{v}+\mathbf{w})+\mathcal{U} = a(\mathbf{v}+\mathcal{U})+(\mathbf{w}+\mathcal{U}) = a\pi(\mathbf{v})+\pi(\mathbf{w})$ so $\pi$ is linear. It is surjective because every coset in $\mathcal{V}/\mathcal{U}$ has the form $\mathbf{v}+\mathcal{U} = \pi(\mathbf{v})$. Finally, $\ker(\pi) = \{\mathbf{v} \in \mathcal{V} : \mathbf{v}+\mathcal{U} = \mathcal{U}\} = \mathcal{U}$.
*Structural Commentary*: The quotient map collapses $\mathcal{U}$ to zero and records cosets.
*Key Insight*: Quotient maps realise subspaces as kernels. □

**Theorem 2.2.9.** *Dimension of a Quotient*
*If $\mathcal{V}$ is finite-dimensional and $\mathcal{U}$ is a subspace, then $\dim(\mathcal{V}/\mathcal{U}) = \dim(\mathcal{V}) - \dim(\mathcal{U})$.*

*Proof.* Choose a basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ of $\mathcal{U}$ and extend it to a basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_k)$ of $\mathcal{V}$.
The cosets $\mathbf{w}_1+\mathcal{U}, \ldots, \mathbf{w}_k+\mathcal{U}$ span $\mathcal{V}/\mathcal{U}$ because the $\mathbf{u}_j$-terms vanish in the quotient.
If $a_1(\mathbf{w}_1+\mathcal{U}) + \cdots + a_k(\mathbf{w}_k+\mathcal{U}) = \mathcal{U}$, then $a_1\mathbf{w}_1 + \cdots + a_k\mathbf{w}_k \in \mathcal{U}$, so a linear relation in the extended basis forces $a_1 = \cdots = a_k = 0$. Hence these cosets form a basis of $\mathcal{V}/\mathcal{U}$, and its length is $k = \dim(\mathcal{V}) - \dim(\mathcal{U})$.
*Structural Commentary*: Quotiening removes the directions in $\mathcal{U}$ and keeps the added basis directions.
*Key Insight*: Passing to $\mathcal{V}/\mathcal{U}$ subtracts the collapsed dimension. □

**Theorem 2.2.10.** *Universal Property of the Quotient Map*
*Let $\mathcal{U} \leq \mathcal{V}$, $\pi : \mathcal{V} \to \mathcal{V}/\mathcal{U}$, and $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ with $\mathcal{U} \leq \ker(T)$.*
*Then $\exists!\, \overline{T} \in \mathcal{L}(\mathcal{V}/\mathcal{U}, \mathcal{W})$ such that $\overline{T}\pi = T$, $\overline{T}(\mathbf{v}+\mathcal{U}) = T\mathbf{v}$, and $\ker(\overline{T}) = \ker(T)/\mathcal{U}$, $\operatorname{range}(\overline{T}) = \operatorname{range}(T)$.*

*Proof.* Define $\overline{T}(\mathbf{v}+\mathcal{U}) = T\mathbf{v}$. If $\mathbf{v}+\mathcal{U} = \mathbf{w}+\mathcal{U}$, then $\mathbf{v}-\mathbf{w} \in \mathcal{U} \leq \ker(T)$, so $T\mathbf{v} = T\mathbf{w}$; hence $\overline{T}$ is well-defined.
Also $\overline{T}\big(a(\mathbf{v}+\mathcal{U}) + b(\mathbf{w}+\mathcal{U})\big) = T(a\mathbf{v}+b\mathbf{w}) = aT\mathbf{v} + bT\mathbf{w}$.
By definition, $\overline{T}\pi = T$. If $S\pi = T$, then $S(\mathbf{v}+\mathcal{U}) = S\pi(\mathbf{v}) = T\mathbf{v} = \overline{T}(\mathbf{v}+\mathcal{U})$, so $S = \overline{T}$.
Also $\mathbf{v}+\mathcal{U} \in \ker(\overline{T}) \iff T\mathbf{v} = \mathbf{0} \iff \mathbf{v} \in \ker(T)$.
So $\ker(\overline{T}) = \ker(T)/\mathcal{U}$, and $\operatorname{range}(\overline{T}) = \{\overline{T}(\mathbf{v}+\mathcal{U}) : \mathbf{v} \in \mathcal{V}\} = \operatorname{range}(T)$.
*Structural Commentary*: Maps killing $\mathcal{U}$ factor through $\mathcal{V}/\mathcal{U}$.
*Key Insight*: Quotients are universal domains for maps with prescribed kernel ambiguity. □




Figure 13: If $\mathcal{U} \subseteq \ker(L)$, then $L = \overline{L}\pi$ uniquely.

**Theorem 2.2.11.** *Subspace Correspondence for Quotients*
*Let $\pi : \mathcal{V} \to \mathcal{V}/\mathcal{U}$ be the quotient map. The assignments $\mathcal{T} \longmapsto \mathcal{T}/\mathcal{U}$ and $\mathcal{Y} \longmapsto \pi^{-1}(\mathcal{Y})$ are inverse inclusion-preserving bijections between $\{\mathcal{T}$ subspace of $\mathcal{V} : \mathcal{U} \subseteq \mathcal{T}\}$ and $\{\mathcal{Y}$ subspace of $\mathcal{V}/\mathcal{U}\}$.*

*Proof.* If $\mathcal{U} \subseteq \mathcal{T} \subseteq \mathcal{V}$, then $\mathcal{T}/\mathcal{U}$ is a subspace of $\mathcal{V}/\mathcal{U}$.
If $\mathcal{Y} \subseteq \mathcal{V}/\mathcal{U}$ is a subspace, then $\pi^{-1}(\mathcal{Y})$ is a subspace of $\mathcal{V}$ and contains $\ker(\pi) = \mathcal{U}$.
For $\mathcal{U} \subseteq \mathcal{T} \subseteq \mathcal{V}$, $\pi^{-1}(\mathcal{T}/\mathcal{U}) = \mathcal{T}$, since $\mathbf{v}+\mathcal{U} = \mathbf{t}+\mathcal{U}$ with $\mathbf{t} \in \mathcal{T}$ implies $\mathbf{v}-\mathbf{t} \in \mathcal{U} \subseteq \mathcal{T}$.
For $\mathcal{Y} \subseteq \mathcal{V}/\mathcal{U}$, surjectivity gives $\pi\left(\pi^{-1}(\mathcal{Y})\right) = \mathcal{Y}$, that is, $\pi^{-1}(\mathcal{Y})/\mathcal{U} = \mathcal{Y}$
Both assignments preserve inclusion by monotonicity of images and preimages.
*Structural Commentary*: Subspaces above $\mathcal{U}$ are exactly the subspaces visible after collapsing $\mathcal{U}$.
*Key Insight*: The quotient map translates containment above $\mathcal{U}$ into containment inside $\mathcal{V}/\mathcal{U}$. □




Figure 14: Intermediate subspaces above $\mathcal{U}$ correspond to subspaces of the quotient.

**Theorem 2.2.12.** *Quotient by the Kernel*
*Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then $\widetilde{T}(\mathbf{v} + \ker(T)) := T\mathbf{v}$ determines a well-defined linear map $\widetilde{T} : \mathcal{V}/\ker(T) \to \mathcal{W}$. Also,*

*(i) $\widetilde{T}$ is injective;*
*(ii) $\operatorname{range}(\widetilde{T}) = \operatorname{range}(T)$;*
*(iii) $\mathcal{V}/\ker(T) \cong \operatorname{range}(T)$.*

*Proof.* If $\mathbf{v} + \ker(T) = \mathbf{v}' + \ker(T)$, then $\mathbf{v} - \mathbf{v}' \in \ker(T)$, so $T\mathbf{v} = T\mathbf{v}'$. Thus $\widetilde{T}$ is well-defined.
For cosets and $a \in \mathbb{F}$, $\widetilde{T}\big(a(\mathbf{v} + \ker(T)) + (\mathbf{v}' + \ker(T))\big) = T(a\mathbf{v} + \mathbf{v}') = aT\mathbf{v} + T\mathbf{v}'$, so $\widetilde{T}$ is linear.

(i) If $\widetilde{T}(\mathbf{v} + \ker(T)) = \mathbf{0}$, then $T\mathbf{v} = \mathbf{0}$, so $\mathbf{v} \in \ker(T)$ and $\mathbf{v} + \ker(T) = \ker(T)$, the zero coset. Hence $\ker(\widetilde{T}) = \{\ker(T)\}$ and $\widetilde{T}$ is injective.
(ii) For every coset $\mathbf{v} + \ker(T)$, the value $\widetilde{T}(\mathbf{v} + \ker(T)) = T\mathbf{v}$ lies in $\operatorname{range}(T)$. Conversely, every vector in $\operatorname{range}(T)$ has the form $T\mathbf{v} = \widetilde{T}(\mathbf{v} + \ker(T))$. Hence the ranges are equal.
(iii) Let $\widehat{T} : \mathcal{V}/\ker(T) \to \operatorname{range}(T)$ be the same formula as $\widetilde{T}$ with codomain restricted to $\operatorname{range}(T)$. The linearity proved above and parts (i)–(ii) show that $\widehat{T}$ is linear, injective, and surjective, so Proposition 1.6.8 gives $\mathcal{V}/\ker(T) \cong \operatorname{range}(T)$.

*Structural Commentary*: Quotienting by the kernel removes exactly the ambiguity of $T$.
*Key Insight*: Every linear map factors as a quotient followed by an isomorphism onto its range. □

**Theorem 2.2.13.** *Second Isomorphism Theorem*
*Let $\mathcal{S}$ and $\mathcal{T}$ be subspaces of $\mathcal{V}$. Then $(\mathcal{S} + \mathcal{T})/\mathcal{T} \cong \mathcal{S}/(\mathcal{S} \cap \mathcal{T})$.*

*Proof.* Define $L : \mathcal{S} \to (\mathcal{S} + \mathcal{T})/\mathcal{T}$ by $L\mathbf{s} = \mathbf{s} + \mathcal{T}$. Then $L$ is linear by the quotient operations.
Moreover, $\ker(L) = \{\mathbf{s} \in \mathcal{S} : \mathbf{s} + \mathcal{T} = \mathcal{T}\} = \mathcal{S} \cap \mathcal{T}$. If $(\mathbf{s} + \mathbf{t}) + \mathcal{T} \in (\mathcal{S} + \mathcal{T})/\mathcal{T}$, with $\mathbf{s} \in \mathcal{S}$ and $\mathbf{t} \in \mathcal{T}$, then $(\mathbf{s} + \mathbf{t}) + \mathcal{T} = \mathbf{s} + \mathcal{T} = L\mathbf{s}$, so $L$ is surjective. Theorem 2.2.12 gives the stated isomorphism.
*Structural Commentary*: Modding out by $\mathcal{T}$ leaves only the part of $\mathcal{S}$ not already inside $\mathcal{T}$.
*Key Insight*: In a quotient by $\mathcal{T}$, adding $\mathcal{T}$ to $\mathcal{S}$ costs nothing. □

**Theorem 2.2.14.** *Third Isomorphism Theorem*
*If $\mathcal{U} \subseteq \mathcal{T} \subseteq \mathcal{V}$ are subspaces, then $(\mathcal{V}/\mathcal{U})/(\mathcal{T}/\mathcal{U}) \cong \mathcal{V}/\mathcal{T}$.*

*Proof.* Define $L : \mathcal{V}/\mathcal{U} \to \mathcal{V}/\mathcal{T}$ by $L(\mathbf{v} + \mathcal{U}) = \mathbf{v} + \mathcal{T}$.
If $\mathbf{v} + \mathcal{U} = \mathbf{w} + \mathcal{U}$, then $\mathbf{v} - \mathbf{w} \in \mathcal{U} \subseteq \mathcal{T}$, so $L$ is well-defined.
It is linear by the quotient operations, and it is surjective since $L(\mathbf{v} + \mathcal{U}) = \mathbf{v} + \mathcal{T}$ for every coset $\mathbf{v} + \mathcal{T}$.
Moreover, $\ker(L) = \{\mathbf{v}+\mathcal{U} : \mathbf{v}+\mathcal{T} = \mathcal{T}\} = \{\mathbf{v}+\mathcal{U} : \mathbf{v} \in \mathcal{T}\} = \mathcal{T}/\mathcal{U}$. Theorem 2.2.12 gives $(\mathcal{V}/\mathcal{U})/(\mathcal{T}/\mathcal{U}) \cong \mathcal{V}/\mathcal{T}$.
*Structural Commentary*: Collapsing $\mathcal{U}$ and then collapsing $\mathcal{T}/\mathcal{U}$ is the same as collapsing $\mathcal{T}$ at once.
*Key Insight*: Quotients of quotients cancel only when the nesting is explicit. □

This block develops two complementary ways to organise a vector space: build from smaller pieces by spans and direct sums, or collapse a chosen subspace by passing to a quotient. Bases and dimension measure exactly what is added, split off, or made invisible.

## 2.3 Decomposition and Complementation

**Theorem 2.3.1.** *Dimension Formula for a Sum*
*If $\mathcal{U}, \mathcal{W}$ are finite-dimensional subspaces of $\mathcal{V}$, then $\dim(\mathcal{U} + \mathcal{W}) = \dim(\mathcal{U}) + \dim(\mathcal{W}) - \dim(\mathcal{U} \cap \mathcal{W})$.*

*Proof.* Choose a basis $(\mathbf{e}_1, \dots, \mathbf{e}_r)$ of $\mathcal{U} \cap \mathcal{W}$, extend it to a basis $(\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{u}_1, \dots, \mathbf{u}_p)$ of $\mathcal{U}$, and to a basis $(\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{w}_1, \dots, \mathbf{w}_q)$ of $\mathcal{W}$. Then $(\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{u}_1, \dots, \mathbf{u}_p, \mathbf{w}_1, \dots, \mathbf{w}_q)$ spans $\mathcal{U} + \mathcal{W}$ because every vector in the sum is the sum of one vector from $\mathcal{U}$ and one vector from $\mathcal{W}$.
To prove linear independence, suppose $a_1\mathbf{e}_1 + \cdots + a_r\mathbf{e}_r + b_1\mathbf{u}_1 + \cdots + b_p\mathbf{u}_p + c_1\mathbf{w}_1 + \cdots + c_q\mathbf{w}_q = \mathbf{0}$. Then $a_1\mathbf{e}_1 + \cdots + a_r\mathbf{e}_r + b_1\mathbf{u}_1 + \cdots + b_p\mathbf{u}_p$ lies in $\mathcal{U} \cap \mathcal{W}$. Since $(\mathbf{e}_1, \dots, \mathbf{e}_r)$ is a basis of $\mathcal{U} \cap \mathcal{W}$ and $(\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{u}_1, \dots, \mathbf{u}_p)$ is a basis of $\mathcal{U}$, we get $b_1 = \cdots = b_p = 0$. The remaining relation lies in the basis $(\mathbf{e}_1, \dots, \mathbf{e}_r, \mathbf{w}_1, \dots, \mathbf{w}_q)$ of $\mathcal{W}$,

so $a_1 = \cdots = a_r = c_1 = \cdots = c_q = 0$. Thus the displayed list is a basis of $\mathcal{U} + \mathcal{W}$.
Therefore $\dim(\mathcal{U} + \mathcal{W}) = r + p + q = (r + p) + (r + q) - r$.
*Structural Commentary*: The formula corrects double-counting: vectors in $\mathcal{U} \cap \mathcal{W}$ are counted once in each subspace. Extending an intersection basis separates shared directions from new directions.
*Key Insight*: The dimension of a sum is the sum of dimensions minus the overlap. □

**Corollary 2.3.2.** *Dimension of a Direct Sum*
*If $\mathcal{U}$ and $\mathcal{W}$ are finite-dimensional subspaces of $\mathcal{V}$ with $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$, then $\dim(\mathcal{U} \oplus \mathcal{W}) = \dim(\mathcal{U}) + \dim(\mathcal{W})$.*

*Proof.* Since $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$, the sum is direct and $\dim(\mathcal{U} \cap \mathcal{W}) = 0$. The result follows from Theorem 2.3.1.
*Structural Commentary*: When the overlap is trivial, there is no double-counting in the dimension formula. Direct sums are exactly the case where the overlap term vanishes.
*Key Insight*: Dimensions add exactly for direct sums. □

**Corollary 2.3.3.** *Large Subspaces Must Intersect*
*Let $\mathcal{U}$ and $\mathcal{W}$ be subspaces of a finite-dimensional vector space $\mathcal{V}$, and let $k$ be a positive integer.*

*(i) If $\dim(\mathcal{U}) + \dim(\mathcal{W}) > \dim(\mathcal{V})$, then $\mathcal{U} \cap \mathcal{W}$ contains a nonzero vector.*
*(ii) If $\dim(\mathcal{U}) + \dim(\mathcal{W}) \geq \dim(\mathcal{V}) + k$, then $\mathcal{U} \cap \mathcal{W}$ contains $k$ linearly independent vectors.*

*Proof.* Since $\mathcal{U} + \mathcal{W} \subseteq \mathcal{V}$, Theorem 2.3.1 gives
$\dim(\mathcal{U} \cap \mathcal{W}) = \dim(\mathcal{U}) + \dim(\mathcal{W}) - \dim(\mathcal{U} + \mathcal{W}) \geq \dim(\mathcal{U}) + \dim(\mathcal{W}) - \dim(\mathcal{V})$.
If $\dim(\mathcal{U}) + \dim(\mathcal{W}) > \dim(\mathcal{V})$, the right-hand side is at least 1, so the intersection contains a nonzero vector.
If $\dim(\mathcal{U}) + \dim(\mathcal{W}) \geq \dim(\mathcal{V}) + k$, then $\dim(\mathcal{U} \cap \mathcal{W}) \geq k$, so intersection contains $k$ linearly independent vectors.
*Structural Commentary*: Excess dimension appears as overlap.
*Key Insight*: Two large subspaces cannot fit inside $\mathcal{V}$ without sharing directions. □

**Theorem 2.3.4.** *Direct Sums and Dimension*
*Suppose $\mathcal{V}$ is finite-dimensional and $\mathcal{U}_1, \ldots, \mathcal{U}_m$ are subspaces of $\mathcal{V}$. Then $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ is a direct sum if and only if $\dim(\mathcal{U}_1 + \cdots + \mathcal{U}_m) = \dim(\mathcal{U}_1) + \cdots + \dim(\mathcal{U}_m)$.*

*Proof.* For each $j$, choose a basis $\beta_j$ of $\mathcal{U}_j$, and concatenate these bases into a list $\beta$.
This list spans $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ and has length $\dim(\mathcal{U}_1) + \cdots + \dim(\mathcal{U}_m)$.

($\Rightarrow$) If the sum is direct, any linear relation among the vectors in $\beta$ gives a zero decomposition $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$ with $\mathbf{u}_j \in \mathcal{U}_j$. Directness gives $\mathbf{u}_j = \mathbf{0}$ for every $j$, and independence of each $\beta_j$ gives all coefficients zero. Thus $\beta$ is a basis of the sum, and the dimensions add.

($\Leftarrow$) If the dimensions add, then $\beta$ is a spanning list whose length equals the dimension of the sum, so Proposition 1.4.3 makes $\beta$ a basis. A zero decomposition $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$ expands to a zero linear relation in this basis, so every $\mathbf{u}_j = \mathbf{0}$. Hence the sum is direct.

*Structural Commentary*: Direct sums add dimensions without hidden overlap.
*Key Insight*: Equality of dimensions means the concatenated bases have no linear redundancy. □

**Definition 2.3.5.** *Basis Adapted to a Direct Sum*
Suppose $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$ and each $\beta_j$ is a basis of $\mathcal{U}_j$. The concatenated basis $\beta = (\beta_1, \ldots, \beta_m)$ of $\mathcal{V}$ is called a *basis adapted to the direct-sum decomposition*.
Equivalently, the vectors in an adapted basis are grouped by the summands whose direct sum is $\mathcal{V}$.

**Proposition 2.3.6.** *Complements Exist in Finite Dimensions*
*If $\mathcal{V}$ is finite-dimensional and $\mathcal{U} \subseteq \mathcal{V}$ is a subspace, then there exists a subspace $\mathcal{W}$ such that $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.*

*Proof.* Choose a basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ of $\mathcal{U}$ and extend it to a basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_n)$ of $\mathcal{V}$.
Let $\mathcal{W} = \operatorname{span}(\mathbf{w}_1, \ldots, \mathbf{w}_n)$. Every vector in $\mathcal{V}$ is a linear combination of this extended basis, so $\mathcal{V} = \mathcal{U} + \mathcal{W}$. If $\mathbf{v} \in \mathcal{U} \cap \mathcal{W}$, then $\mathbf{v}$ has one representation using only $\mathbf{u}_1, \ldots, \mathbf{u}_m$ and another using only $\mathbf{w}_1, \ldots, \mathbf{w}_n$.
Uniqueness of coefficients in the basis of $\mathcal{V}$ forces $\mathbf{v} = \mathbf{0}$. Hence $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.
*Structural Commentary*: Complements split a space into a subspace and the remaining basis directions.
*Key Insight*: Extend a basis of $\mathcal{U}$ to split $\mathcal{V}$. □

**Theorem 2.3.7.** *Complements Exist*
*Assuming Zorn's lemma, for every $\mathcal{U} \leq \mathcal{V}$, $\exists\, \mathcal{W} \leq \mathcal{V}$ such that $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.*

*Proof.* By Theorem 1.4.6, choose a Hamel basis $\mathcal{B}_\mathcal{U}$ of $\mathcal{U}$ and a Hamel basis $\mathcal{B} \supseteq \mathcal{B}_\mathcal{U}$ of $\mathcal{V}$.
Put $\mathcal{C} = \mathcal{B} \setminus \mathcal{B}_\mathcal{U}$ and $\mathcal{W} = \operatorname{span}(\mathcal{C})$. For $\mathbf{v} \in \mathcal{V}$, its finite $\mathcal{B}$-expansion splits as $\mathbf{v} = \sum_{b \in F_\mathcal{U}} a_b b + \sum_{c \in F_\mathcal{C}} a_c c \in \mathcal{U} + \mathcal{W}$, with finite $F_\mathcal{U} \subseteq \mathcal{B}_\mathcal{U}$ and $F_\mathcal{C} \subseteq \mathcal{C}$. If $\mathbf{x} \in \mathcal{U} \cap \mathcal{W}$, then its $\mathcal{B}$-coordinates are supported in both $\mathcal{B}_\mathcal{U}$ and $\mathcal{C}$; uniqueness gives $\mathbf{x} = \mathbf{0}$. Hence $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.
*Structural Commentary*: Extend a basis of $\mathcal{U}$; span the new directions.
*Key Insight*: The uniqueness of basis expansion becomes the uniqueness of direct-sum decomposition. □

**Proposition 2.3.8.** *Rank-Nullity as a Complement Isomorphism*
*Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, and suppose $\mathcal{V} = \ker(T) \oplus \mathcal{M}$. Then $T|_\mathcal{M} : \mathcal{M} \to \operatorname{range}(T)$ is an isomorphism.*

*Proof.* Indeed, $\ker(T|_\mathcal{M}) = \mathcal{M} \cap \ker(T) = \{\mathbf{0}\}$, and every $T\mathbf{v}$ equals $T\mathbf{m}$ for the decomposition $\mathbf{v} = \mathbf{k} + \mathbf{m}$.
Thus rank-nullity counts $\mathcal{V} = \ker(T) \oplus \mathcal{M}$ with $\mathcal{M} \cong \operatorname{range}(T)$. □

**Proposition 2.3.9.** *One-Sided Inverses Split in Finite Dimensions*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional vector spaces and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.*

*(i) If $T$ is injective, then there exists $L \in \mathcal{L}(\mathcal{W}, \mathcal{V})$ such that $LT = I_\mathcal{V}$.*
*(ii) If $T$ is surjective, then there exists $R \in \mathcal{L}(\mathcal{W}, \mathcal{V})$ such that $TR = I_\mathcal{W}$.*

*Proof.*

(i) Choose $\mathcal{H} \subseteq \mathcal{W}$ with $\mathcal{W} = \operatorname{range}(T) \oplus \mathcal{H}$. For $\mathbf{w} = T\mathbf{v} + \mathbf{h}$, define $L\mathbf{w} = \mathbf{v}$. This is well-defined since the direct-sum decomposition is unique and $T$ is injective. It is linear, and $LT\mathbf{v} = \mathbf{v}$ for every $\mathbf{v} \in \mathcal{V}$.
(ii) Choose $\mathcal{H} \subseteq \mathcal{V}$ with $\mathcal{V} = \ker(T) \oplus \mathcal{H}$. Then $T|_\mathcal{H} : \mathcal{H} \to \mathcal{W}$ is injective since $\mathcal{H} \cap \ker(T) = \{\mathbf{0}\}$, and surjective since each $\mathbf{v} = \mathbf{k} + \mathbf{h}$ satisfies $T\mathbf{v} = T\mathbf{h}$. Let $R = (T|_\mathcal{H})^{-1}$, viewed as a map $\mathcal{W} \to \mathcal{V}$. Then $R$ is linear and $TR = I_\mathcal{W}$.

*Structural Commentary*: Complements supply the missing directions on which a one-sided inverse is chosen.
*Key Insight*: Finite-dimensional vector spaces split kernels and ranges. □

**Proposition 2.3.10.** *Quotient by a Complemented Subspace*
*Let $\mathcal{U}$ and $\mathcal{W}$ be subspaces of $\mathcal{V}$ with $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$. Then $J : \mathcal{W} \to \mathcal{V}/\mathcal{U}$, $J(\mathbf{w}) = \mathbf{w} + \mathcal{U}$ is an isomorphism. Hence every complement of $\mathcal{U}$ is isomorphic to $\mathcal{V}/\mathcal{U}$.*

*Proof.* The map is linear by the quotient operations from Proposition 2.2.6.
If $J(\mathbf{w}) = \mathcal{U}$, then $\mathbf{w} \in \mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$, so $J$ is injective.
If $\mathbf{v} + \mathcal{U} \in \mathcal{V}/\mathcal{U}$, write $\mathbf{v} = \mathbf{u} + \mathbf{w}$ with $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{W}$. Then $\mathbf{v} + \mathcal{U} = \mathbf{w} + \mathcal{U} = J(\mathbf{w})$, so $J$ is surjective.
*Structural Commentary*: A complement gives canonical representatives for quotient cosets.
*Key Insight*: Collapsing $\mathcal{U}$ leaves exactly the complementary directions visible. □

**Definition 2.3.11.** *Codimension*
If $\mathcal{U}$ is a subspace of $\mathcal{V}$ and $\dim(\mathcal{V}/\mathcal{U}) < \infty$, define $\operatorname{codim}_\mathcal{V}(\mathcal{U}) := \dim(\mathcal{V}/\mathcal{U})$.
Thus, if $\mathcal{V}$ is finite-dimensional, Theorem 2.2.9 gives $\operatorname{codim}_\mathcal{V}(\mathcal{U}) = \dim(\mathcal{V}) - \dim(\mathcal{U})$.
If $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, Proposition 2.3.10 gives $\operatorname{codim}_\mathcal{V}(\mathcal{U}) = \dim(\mathcal{W})$.

**Perspective 2.3.12.** *Quotients Split Over Fields*
In finite-dimensional vector spaces, choosing a complement $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$ identifies $\mathcal{V}/\mathcal{U}$ with $\mathcal{W}$. The quotient map collapses $\mathcal{U}$ and keeps the complementary directions.

**Proposition 2.3.13.** *Quotient of a Direct Sum*
*If $\mathcal{V} = \mathcal{V}_1 \oplus \mathcal{V}_2$ and $\mathcal{U}_j \leq \mathcal{V}_j$ for $j = 1, 2$, then $\frac{\mathcal{V}}{\mathcal{U}_1 \oplus \mathcal{U}_2} \cong \frac{\mathcal{V}_1}{\mathcal{U}_1} \oplus \frac{\mathcal{V}_2}{\mathcal{U}_2}$.*

*Proof.* Define, using the unique decomposition
$\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2$, $\Phi : \mathcal{V} \to (\mathcal{V}_1/\mathcal{U}_1) \oplus (\mathcal{V}_2/\mathcal{U}_2)$ and $\Phi(\mathbf{v}_1 + \mathbf{v}_2) = (\mathbf{v}_1 + \mathcal{U}_1, \mathbf{v}_2 + \mathcal{U}_2)$.
For $\mathbf{x} = \mathbf{x}_1 + \mathbf{x}_2$, $\mathbf{y} = \mathbf{y}_1 + \mathbf{y}_2$, and $a, b \in \mathbb{F}$, $\Phi(a\mathbf{x} + b\mathbf{y}) = a\Phi(\mathbf{x}) + b\Phi(\mathbf{y})$, so $\Phi$ is linear.
It is surjective since $(\mathbf{v}_1 + \mathcal{U}_1, \mathbf{v}_2 + \mathcal{U}_2) = \Phi(\mathbf{v}_1 + \mathbf{v}_2)$. Finally, $\ker(\Phi) = \{\mathbf{v}_1 + \mathbf{v}_2 : \mathbf{v}_1 \in \mathcal{U}_1,\ \mathbf{v}_2 \in \mathcal{U}_2\} = \mathcal{U}_1 \oplus \mathcal{U}_2$.
The result follows from Theorem 2.2.12.
*Structural Commentary*: Quotienting respects direct-sum coordinates.
*Key Insight*: Direct-sum coordinates make quotienting componentwise. □

**Perspective 2.3.14.** *Quotient and Complement Warnings*
If $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, then $\mathcal{W} \cong \mathcal{V}/\mathcal{U}$, but complements are not functorial in $\mathcal{U}$.
For $\mathcal{V}$ countably infinite-dimensional with basis $(\mathbf{e}_n)_{n \geq 1}$, put $\mathcal{S} = \operatorname{span}(\mathbf{e}_n : n \geq 2)$.
Then $\mathcal{S} \cong \mathcal{V}$ and $\mathcal{V} = \mathcal{V} \oplus \{\mathbf{0}\} = \mathcal{S} \oplus \mathbb{F}\mathbf{e}_1$; $\mathcal{V}/\mathcal{V} = \{\mathbf{0}\}$, $\mathcal{V}/\mathcal{S} \cong \mathbb{F}\mathbf{e}_1$.

## 2.4 Algebraic Projections and Resolutions

**Definition 2.4.1.** *Projection Along a Complement*
An operator $P \in \mathcal{L}(\mathcal{V})$ is an *algebraic projection*, or *idempotent operator*, if $P^2 = P$.
If $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, the *projection onto $\mathcal{U}$ along $\mathcal{W}$* is $\rho_{\mathcal{U},\mathcal{W}} \in \mathcal{L}(\mathcal{V})$ where $\rho_{\mathcal{U},\mathcal{W}}(\mathbf{u}+\mathbf{w}) := \mathbf{u}$ for $\mathbf{u} \in \mathcal{U}$, $\mathbf{w} \in \mathcal{W}$.

**Proposition 2.4.2.** *Algebraic Projections and Direct-Sum Splittings*
*For $P \in \mathcal{L}(\mathcal{V})$, $P$ is an algebraic projection if and only if $\mathcal{V} = \operatorname{range}(P) \oplus \ker(P)$ and $P = \rho_{\operatorname{range}(P),\ker(P)}$.*
*In particular, if $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$ and $\rho = \rho_{\mathcal{U},\mathcal{W}}$, then $\rho$ is an algebraic projection, $\operatorname{range}(\rho) = \mathcal{U}$, and $\ker(\rho) = \mathcal{W}$.*

*Proof.*
($\Rightarrow$) Suppose $P$ is an algebraic projection, so $P^2 = P$. For every $\mathbf{v} \in \mathcal{V}$, $\mathbf{v} = P\mathbf{v} + (\mathbf{v} - P\mathbf{v})$, with $P\mathbf{v} \in \operatorname{range}(P)$ and $\mathbf{v} - P\mathbf{v} \in \ker(P)$. If $\mathbf{x} \in \operatorname{range}(P) \cap \ker(P)$, then $\mathbf{x} = P\mathbf{y}$ and $\mathbf{0} = P\mathbf{x} = P^2\mathbf{y} = \mathbf{x}$. Thus the sum is direct. Also $P$ fixes $\operatorname{range}(P)$ and kills $\ker(P)$, so $P = \rho_{\operatorname{range}(P),\ker(P)}$.
($\Leftarrow$) Assume the right-hand side. For $\mathbf{v} = \mathbf{x} + \mathbf{y}$ with $\mathbf{x} \in \operatorname{range}(P)$ and $\mathbf{y} \in \ker(P)$, $P^2\mathbf{v} = P\mathbf{x} = \mathbf{x} = P\mathbf{v}$. Thus $P$ is an algebraic projection. For $\rho_{\mathcal{U},\mathcal{W}}$, the formula $\rho_{\mathcal{U},\mathcal{W}}(\mathbf{u}+\mathbf{w}) = \mathbf{u}$ gives range $\mathcal{U}$ and kernel $\mathcal{W}$, and the equivalence above shows that $\rho$ is an algebraic projection.

*Structural Commentary*: Idempotents keep one summand and kill its complement.
*Key Insight*: Direct-sum coordinates turn a projection into $I \oplus 0$. □

**Proposition 2.4.3.** *Complementary Algebraic Projections*
*If $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, then $\rho_{\mathcal{U},\mathcal{W}} + \rho_{\mathcal{W},\mathcal{U}} = I_{\mathcal{V}}$, and $\rho_{\mathcal{U},\mathcal{W}}\rho_{\mathcal{W},\mathcal{U}} = \rho_{\mathcal{W},\mathcal{U}}\rho_{\mathcal{U},\mathcal{W}} = 0$.*

*Proof.* For $\mathbf{v} = \mathbf{u} + \mathbf{w}$, $\mathbf{u} \in \mathcal{U}$, $\mathbf{w} \in \mathcal{W}$, $(\rho_{\mathcal{U},\mathcal{W}} + \rho_{\mathcal{W},\mathcal{U}})\mathbf{v} = \mathbf{u} + \mathbf{w} = \mathbf{v}$,
and $\rho_{\mathcal{U},\mathcal{W}}\rho_{\mathcal{W},\mathcal{U}}\mathbf{v} = \rho_{\mathcal{U},\mathcal{W}}\mathbf{w} = \mathbf{0}$; $\rho_{\mathcal{W},\mathcal{U}}\rho_{\mathcal{U},\mathcal{W}}\mathbf{v} = \rho_{\mathcal{W},\mathcal{U}}\mathbf{u} = \mathbf{0}$. Thus the stated operator identities hold.
*Structural Commentary*: A direct-sum decomposition gives two complementary idempotents.
*Key Insight*: Projection onto one summand kills the other summand. □

**Definition 2.4.4.** *Orthogonal Projections in the Algebraic Sense*
Two algebraic projections $\rho, \sigma \in \mathcal{L}(\mathcal{V})$ are *algebraically orthogonal*, written $\rho \perp \sigma$, if $\rho\sigma = \sigma\rho = 0$.

*Resolution of identity* on $\mathcal{V}$ is a finite family of algebraic projections $(\rho_i)_{i=1}^k$ with $\rho_i \perp \rho_j$ for $i \neq j$ and $\sum_{i=1}^k \rho_i = I_{\mathcal{V}}$.

**Theorem 2.4.5.** *Resolution Ranges Form a Direct Sum*
*If $(\rho_i)_{i=1}^k$ is a resolution of $I_{\mathcal{V}}$ and $R_i = \operatorname{range}(\rho_i)$, then $\mathcal{V} = \bigoplus_{i=1}^k R_i$, and $\ker(\rho_i) = \bigoplus_{j \neq i} R_j$.*

*Proof.* Since $\sum_{i=1}^k \rho_i = I_{\mathcal{V}}$, $\mathbf{v} = \sum_{i=1}^k \rho_i\mathbf{v} \in \sum_{i=1}^k R_i$.
If $\sum_{i=1}^k \mathbf{x}_i = \mathbf{0}$ with $\mathbf{x}_i \in R_i$, then for each $1 \leq m \leq k$, $\mathbf{0} = \rho_m \sum_{i=1}^k \mathbf{x}_i = \mathbf{x}_m$, as $\rho_m$ fixes $R_m$ and kills $R_i$ for $i \neq m$.
Thus $\mathcal{V} = \bigoplus_{i=1}^k R_i$. Also, for $\mathbf{v} = \sum_{j=1}^k \mathbf{x}_j$ with $\mathbf{x}_j \in R_j$, $\rho_i\mathbf{v} = \mathbf{x}_i$, so $\ker(\rho_i) = \bigoplus_{j \neq i} R_j$.
*Structural Commentary*: A resolution decomposes the identity into independent idempotent pieces.
*Key Insight*: Pairwise zero products are the operator form of directness. □

**Theorem 2.4.6.** *Direct Sums Give Resolutions*
*If $\mathcal{V} = \bigoplus_{i=1}^k \mathcal{U}_i$ and $\rho_i = \rho_{\mathcal{U}_i,\bigoplus_{j\neq i}\mathcal{U}_j}$, then $(\rho_i)_{i=1}^k$ is a resolution of $I_{\mathcal{V}}$.*

*Proof.* For $\mathbf{v} = \sum_{i=1}^k \mathbf{u}_i$, $\mathbf{u}_i \in \mathcal{U}_i$, $\sum_{i=1}^k \rho_i\mathbf{v} = \sum_{i=1}^k \mathbf{u}_i = \mathbf{v}$. Thus $\sum_{i=1}^k \rho_i = I_{\mathcal{V}}$.
If $i \neq j$, then $\rho_i\rho_j\mathbf{v} = \rho_i\mathbf{u}_j = \mathbf{0}$ and $\rho_j\rho_i\mathbf{v} = \rho_j\mathbf{u}_i = \mathbf{0}$, so $\rho_i \perp \rho_j$.
*Structural Commentary*: Direct-sum coordinates produce mutually annihilating coordinate projections.
*Key Insight*: Coordinate projections add to the identity and kill all other summands. □

**Theorem 2.4.7.** *Algebra of Projections*
*Let $\rho, \sigma \in \mathcal{L}(\mathcal{V})$ be algebraic projections and $\operatorname{char}(\mathbb{F}) \neq 2$.*

*(i) $\rho + \sigma$ is an algebraic projection $\iff \rho \perp \sigma$.*
*Then $\operatorname{range}(\rho+\sigma) = \operatorname{range}(\rho) \oplus \operatorname{range}(\sigma)$; $\ker(\rho+\sigma) = \ker(\rho) \cap \ker(\sigma)$*

*(ii) $\rho - \sigma$ is an algebraic projection $\iff \rho\sigma = \sigma\rho = \sigma$.*
*Then $\operatorname{range}(\rho-\sigma) = \operatorname{range}(\rho) \cap \ker(\sigma)$; $\ker(\rho-\sigma) = \ker(\rho) \oplus \operatorname{range}(\sigma)$.*

*(iii) If $\rho\sigma = \sigma\rho$, then $\rho\sigma$ is algebraic projection and $\operatorname{range}(\rho\sigma) = \operatorname{range}(\rho) \cap \operatorname{range}(\sigma)$; $\ker(\rho\sigma) = \ker(\rho) + \ker(\sigma)$.*

*Proof.*

(i) Put $A = \rho\sigma$, $B = \sigma\rho$, and $C = \rho\sigma\rho$. Since $(\rho+\sigma)^2 = \rho+\sigma \iff A+B = 0$,
$A+B=0$ gives $A+C = \rho(A+B) = 0$ and $C+B = (A+B)\rho = 0$, hence $B = A$ and $2A = 0$.
Thus $A = B = 0$. The converse is immediate. Assume $\rho \perp \sigma$. Then $\operatorname{range}(\rho+\sigma) \subseteq \operatorname{range}(\rho) + \operatorname{range}(\sigma)$, while $(\rho+\sigma)(\mathbf{x}+\mathbf{y}) = \mathbf{x}+\mathbf{y}$ for $\mathbf{x} \in \operatorname{range}(\rho)$, $\mathbf{y} \in \operatorname{range}(\sigma)$. Also $\operatorname{range}(\rho) \cap \operatorname{range}(\sigma) = \{\mathbf{0}\}$, since $\mathbf{z} \in \operatorname{range}(\rho) \cap \operatorname{range}(\sigma)$ and $\mathbf{z} = \sigma\mathbf{w}$ imply $\mathbf{z} = \rho\mathbf{z} = \rho\sigma\mathbf{w} = \mathbf{0}$. Finally, $(\rho+\sigma)\mathbf{v} = 0 \iff \rho\mathbf{v} = 0 = \sigma\mathbf{v}$.

(ii) Since $P^2 = P \iff (I_{\mathcal{V}} - P)^2 = I_{\mathcal{V}} - P$, $\rho - \sigma$ is an algebraic projection $\iff (I_{\mathcal{V}} - \rho) + \sigma$ is an algebraic projection. By (i), this is equivalent to $(I_{\mathcal{V}} - \rho)\sigma = \sigma(I_{\mathcal{V}} - \rho) = 0$, i.e. $\rho\sigma = \sigma\rho = \sigma$.
Assume $\rho\sigma = \sigma\rho = \sigma$ and set $Q = \rho - \sigma$. Then $Q\mathbf{v} \in \operatorname{range}(\rho) \cap \ker(\sigma)$; $Q\mathbf{x} = \mathbf{x}$, $\mathbf{x} \in \operatorname{range}(\rho) \cap \ker(\sigma)$, so $\operatorname{range}(Q) = \operatorname{range}(\rho) \cap \ker(\sigma)$. Also $Q$ kills both $\ker(\rho)$ and $\operatorname{range}(\sigma)$, and $\ker(\rho) \cap \operatorname{range}(\sigma) = \{\mathbf{0}\}$.
If $\mathbf{v} \in \ker(Q)$, then $\rho\mathbf{v} = \sigma\mathbf{v}$ and $\mathbf{v} = (\mathbf{v} - \rho\mathbf{v}) + \sigma\mathbf{v} \in \ker(\rho) \oplus \operatorname{range}(\sigma)$. Hence $\ker(Q) = \ker(\rho) \oplus \operatorname{range}(\sigma)$.

(iii) If $\rho\sigma = \sigma\rho$, then $(\rho\sigma)^2 = \rho^2\sigma^2 = \rho\sigma$. Moreover $\mathbf{x} \in \operatorname{range}(\rho\sigma) \iff \rho\mathbf{x} = \mathbf{x} = \sigma\mathbf{x}$,
so $\operatorname{range}(\rho\sigma) = \operatorname{range}(\rho) \cap \operatorname{range}(\sigma)$. Finally, $\ker(\rho) + \ker(\sigma) \subseteq \ker(\rho\sigma)$, and if $\rho\sigma\mathbf{v} = 0$,
then $\mathbf{v} = (\mathbf{v} - \sigma\mathbf{v}) + \sigma\mathbf{v}$; $\mathbf{v} - \sigma\mathbf{v} \in \ker(\sigma)$, $\sigma\mathbf{v} \in \ker(\rho)$. Thus $\ker(\rho\sigma) = \ker(\rho) + \ker(\sigma)$.

*Structural Commentary*: Operator identities encode range-kernel geometry.
*Key Insight*: Idempotence is a rigid equation; expanding it forces the cross terms to encode the geometry. □

# 3 Matrix Representations, Change of Basis, Block Matrices, and Operations

## 3.1 Matrix Algebra and Linear Systems

**Definition 3.1.1.** *Matrices and Matrix Operations*

(i) *Matrix space*: for $m, n \in \mathbb{Z}_{>0}$, $\operatorname{Mat}(m,n,\mathbb{F}) = \{[A_{j,k}]_{1\le j\le m, 1\le k\le n} : A_{j,k} \in \mathbb{F}\}$.
(ii) *Entries, rows, columns*: for $A = [A_{j,k}] \in \operatorname{Mat}(m,n,\mathbb{F})$,
$A_{j,k}$ is the $(j,k)$-entry, $A_{j,\cdot}$ is row $j$, and $A_{\cdot,k}$ is column $k$.
(iii) *Transpose*: for $A \in \operatorname{Mat}(m,n,\mathbb{F})$, $A^t \in \operatorname{Mat}(n,m,\mathbb{F})$ is defined by $(A^t)_{k,j} = A_{j,k}$.
(iv) *Main diagonal*: for $A = [A_{j,k}] \in \operatorname{Mat}(m,n,\mathbb{F})$ and $r = \min(m,n)$, $\operatorname{diag}(A) = (A_{1,1}, \dots, A_{r,r})$.
(v) *Symmetry*: for $A \in \operatorname{Mat}(n,n,\mathbb{F})$, $A$ is symmetric if $A^t = A$ and skew-symmetric if $A^t = -A$.
(vi) *Triangularity*: for $A \in \operatorname{Mat}(n,n,\mathbb{F})$, $A$ is upper triangular if $A_{j,k} = 0$ for $j > k$,
lower triangular if $A_{j,k} = 0$ for $j < k$, and diagonal if $A_{j,k} = 0$ for $j \neq k$.
(vii) *Linear operations*: for $A, C \in \operatorname{Mat}(m,n,\mathbb{F})$ and $a \in \mathbb{F}$, $(A+C)_{j,k} = A_{j,k} + C_{j,k}$ and $(aA)_{j,k} = aA_{j,k}$.
(viii) *Matrix product*: for $A \in \operatorname{Mat}(m,n,\mathbb{F})$, $B \in \operatorname{Mat}(n,p,\mathbb{F})$, $AB \in \operatorname{Mat}(m,p,\mathbb{F})$ is $(AB)_{j,k} := \sum_{r=1}^{n} A_{j,r}B_{r,k}$.
(ix) *Identity and inverse*: $I_n = [\delta_{j,k}] \in \operatorname{Mat}(n,n,\mathbb{F})$; for $A, B \in \operatorname{Mat}(n,n,\mathbb{F})$, $B$ is a two-sided inverse of $A$ if $AB = I_n = BA$, and $A$ is invertible if such $B$ exists.

**Lemma 3.1.2.** *Uniqueness of Two-Sided Inverses*
*Let $A, B, C \in \operatorname{Mat}(n,n,\mathbb{F})$. If $AB = I_n$ and $BC = I_n$, then $A = C$.*

*Proof.* Using $BC = I_n$ and $AB = I_n$, $A = AI_n = A(BC) = (AB)C = I_nC = C$.
*Structural Commentary*: A common middle inverse forces the left and right inverses to coincide.
*Key Insight*: Associativity turns $AB = I_n = BC$ into $A = C$. □

**Definition 3.1.3.** *Inverse Matrix Notation*
If $A \in \operatorname{Mat}(n,n,\mathbb{F})$ is invertible, Lemma 3.1.2 shows that its two-sided inverse is unique.
This inverse is denoted $A^{-1}$, and the set of invertible matrices in $\operatorname{Mat}(n,n,\mathbb{F})$ is denoted $\operatorname{Mat}(n,n,\mathbb{F})^{\times}$.

**Proposition 3.1.4.** *Matrix Spaces*
*For positive integers $m, n$, $\operatorname{Mat}(m,n,\mathbb{F})$ is a vector space over $\mathbb{F}$ with dimension $mn$.*

*Proof.* Entrywise addition and scalar multiplication satisfy the vector space axioms because $\mathbb{F}$ does. For $1 \le j \le m$ and $1 \le k \le n$, let $E_{j,k}$ be the matrix with 1 in entry $(j,k)$ and 0 elsewhere.

Every $A = [A_{j,k}]$ has the unique expansion $A = \sum_{j=1}^{m} \sum_{k=1}^{n} A_{j,k}E_{j,k}$.

Thus the $mn$ matrices $E_{j,k}$ form a basis, so $\dim(\operatorname{Mat}(m,n,\mathbb{F})) = mn$.
*Structural Commentary*: A matrix is a finite array of independent scalar slots.
*Key Insight*: The standard entry matrices isolate those slots one at a time. □

**Proposition 3.1.5.** *Matrix Product Entries Are Row Times Column*
*If $A \in \operatorname{Mat}(m,n,\mathbb{F})$ and $B \in \operatorname{Mat}(n,p,\mathbb{F})$, then $(AB)_{j,k} = A_{j,\cdot}B_{\cdot,k}$ for $1 \le j \le m$ and $1 \le k \le p$.*

*Proof.* The row-column product $A_{j,\cdot}B_{\cdot,k}$ is exactly $\sum_{r=1}^{n} A_{j,r}B_{r,k}$, which is the definition of $(AB)_{j,k}$.
*Structural Commentary*: Matrix multiplication is assembled entry-by-entry from row-column pairings.
*Key Insight*: Each entry of $AB$ sees one row of $A$ and one column of $B$. □

**Proposition 3.1.6.** *Matrix-Vector Products as Column Combinations*
*If* $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathrm{Mat}(m,n,\mathbb{F})$ *and* $\mathbf{x} = (x_1,\dots,x_n)^t \in \mathbb{F}^n$, *then* $A\mathbf{x} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n$.

*Proof.* The $j$th entry of $A\mathbf{x}$ is $\sum_{k=1}^{n} A_{j,k}x_k$, which is also the $j$th entry of $x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n$.
*Structural Commentary*: Multiplying by a column selects a linear combination of the columns of $A$.
*Key Insight*: The coordinates of $\mathbf{x}$ are precisely the column coefficients. □

**Definition 3.1.7.** *Column Space, Matrix Rank, and Nullity*
Let $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathrm{Mat}(m,n,\mathbb{F})$.

(i) $\mathrm{col}(A) := \mathrm{span}(\mathbf{a}_1,\dots,\mathbf{a}_n) \subseteq \mathbb{F}^m$, and $\mathrm{row}(A) := \mathrm{col}(A^t) \subseteq \mathbb{F}^n$.
(ii) $\mathrm{rank}(A) := \dim(\mathrm{col}(A))$; column rank is $\dim(\mathrm{col}(A))$, and row rank is $\dim(\mathrm{row}(A))$.
(iii) $A$ has *full rank* if $\mathrm{rank}(A) = \min(m,n)$, *full column rank* if $\mathrm{rank}(A) = n$, and *full row rank* if $\mathrm{rank}(A) = m$.
(iv) $\ker(A) := \{\mathbf{x} \in \mathbb{F}^n : A\mathbf{x} = \mathbf{0}\}$ is the *nullspace*, and $\mathrm{nullity}(A) := \dim(\ker(A))$.

**Proposition 3.1.8.** *Solvability and Complete Solution Sets*
*Let* $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathrm{Mat}(m,n,\mathbb{F})$ *and let* $\mathbf{b} \in \mathbb{F}^m$. *Then* $A\mathbf{x} = \mathbf{b}$ *is solvable if and only if* $\mathbf{b} \in \mathrm{col}(A)$.
*If* $\mathbf{x}_p$ *is one solution, then every solution is of the form* $\mathbf{x} = \mathbf{x}_p + \mathbf{x}_0$, *where* $\mathbf{x}_0 \in \ker(A)$.

*Proof.*

($\Rightarrow$) If $A\mathbf{x} = \mathbf{b}$ for some $\mathbf{x} = (x_1,\dots,x_n)^T$, then $\mathbf{b} = A\mathbf{x} = \sum_{j=1}^{n} x_j\mathbf{a}_j \in \mathrm{col}(A)$.

($\Leftarrow$) If $\mathbf{b} \in \mathrm{col}(A)$, then $\mathbf{b} = \sum_{j=1}^{n} x_j\mathbf{a}_j$ for some scalars $x_j$. Hence, for $\mathbf{x} = (x_1,\dots,x_n)^T$, we have $A\mathbf{x} = \mathbf{b}$.

Now suppose $A\mathbf{x}_p = \mathbf{b}$. For any $\mathbf{x}$, $A\mathbf{x} = \mathbf{b} \iff A(\mathbf{x} - \mathbf{x}_p) = \mathbf{0} \iff \mathbf{x} - \mathbf{x}_p \in \ker(A)$.
Thus every solution is exactly of the form $\mathbf{x} = \mathbf{x}_p + \mathbf{x}_0$, where $\mathbf{x}_0 \in \ker(A)$.
*Structural Commentary*: A consistent system is an affine shift of $\ker(A)$ by any particular solution.
*Key Insight*: Column space decides existence; kernel decides nonuniqueness. □

**Proposition 3.1.9.** *Matrix Products by Columns*
*If* $A \in \mathrm{Mat}(m,r,\mathbb{F})$ *and* $B = [\mathbf{b}_1 \ \cdots \ \mathbf{b}_n] \in \mathrm{Mat}(r,n,\mathbb{F})$, *then* $AB = [A\mathbf{b}_1 \ \cdots \ A\mathbf{b}_n]$.

*Proof.* For each $k$, the $k$th column of $AB$ has entries $\sum_{\ell=1}^{r} A_{j,\ell}B_{\ell,k}$, which are exactly the entries of $A\mathbf{b}_k$.
*Structural Commentary*: Matrix products apply $A$ to columns of $B$.
*Key Insight*: The columns of $AB$ are obtained by applying $A$ to the columns of $B$. □

**Proposition 3.1.10.** *Matrix Products as Sums of Outer Products*
*Let* $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_r] \in \mathrm{Mat}(m,r,\mathbb{F})$, *and write* $B \in \mathrm{Mat}(r,n,\mathbb{F})$ *by rows as*

$$B = \begin{bmatrix} \mathbf{b}_1^t \\ \vdots \\ \mathbf{b}_r^t \end{bmatrix}.$$

*Then* $AB = \mathbf{a}_1\mathbf{b}_1^t + \cdots + \mathbf{a}_r\mathbf{b}_r^t$.

*Proof.* The $(j,k)$ entry of $\mathbf{a}_\ell\mathbf{b}_\ell^t$ is $A_{j,\ell}B_{\ell,k}$. Summing over $\ell = 1,\dots,r$ gives $\sum_{\ell=1}^{r} A_{j,\ell}B_{\ell,k} = (AB)_{j,k}$.
*Structural Commentary*: Column-row products decompose a matrix product into single column-row pieces.
*Key Insight*: Each shared index $\ell$ contributes one outer product. □

**Proposition 3.1.11.** *Elementary Transpose Rules*
*Let* $A, C \in \mathrm{Mat}(m,n,\mathbb{F})$ *and let* $a \in \mathbb{F}$. *Then* $(A^t)^t = A$; $(A+C)^t = A^t + C^t$; *and* $(aA)^t = aA^t$.

*Proof.* For all admissible indices, $((A^t)^t)_{j,k} = (A^t)_{k,j} = A_{j,k}$, so $(A^t)^t = A$.
Also $((A+C)^t)_{j,k} = (A+C)_{k,j} = A_{k,j} + C_{k,j} = (A^t + C^t)_{j,k}$ and $((aA)^t)_{j,k} = (aA)_{k,j} = aA_{k,j} = (aA^t)_{j,k}$.
*Structural Commentary*: Transpose swaps indices; entrywise algebra is unchanged.
*Key Insight*: Transpose is linear and involutive. □

**Example 3.1.12.** *Symmetric and Skew-Symmetric Decomposition*
Assume $\mathrm{char}(\mathbb{F}) \neq 2$, $M = \mathrm{Mat}(n,n,\mathbb{F})$, let $\mathrm{Sym}_n(\mathbb{F}) = \{A \in M : A^t = A\}$, $\mathrm{Skew}_n(\mathbb{F}) = \{A \in M : A^t = -A\}$.
For the map $\tau : M \to M$ defined by $\tau(A) = A^t$, linearity follows from Proposition 3.1.11. Hence $\mathrm{Sym}_n(\mathbb{F}) = \ker(\tau - I)$ and $\mathrm{Skew}_n(\mathbb{F}) = \ker(\tau + I)$, so both are subspaces of $M$.
For every $A \in M$, $A = \frac{1}{2}(A + A^t) + \frac{1}{2}(A - A^t) \in \mathrm{Sym}_n(\mathbb{F}) + \mathrm{Skew}_n(\mathbb{F})$.

Also $A \in \mathrm{Sym}_n(\mathbb{F}) \cap \mathrm{Skew}_n(\mathbb{F})$ implies $A = A^t = -A$, hence $A = 0$. Thus $M = \mathrm{Sym}_n(\mathbb{F}) \oplus \mathrm{Skew}_n(\mathbb{F})$.
*Structural Commentary*: Transpose splits a matrix into its even and odd parts.
*Key Insight*: The directness of the split uses the ability to divide by 2.

**Proposition 3.1.13.** *Transpose Reverses Products*
*If $A \in \mathrm{Mat}(m,n,\mathbb{F})$ and $B \in \mathrm{Mat}(n,p,\mathbb{F})$, then $(AB)^t = B^t A^t$.*

*Proof.* For $1 \le j \le p$ and $1 \le k \le m$, $((AB)^t)_{j,k} = (AB)_{k,j} = \sum_{\ell=1}^{n} A_{k,\ell} B_{\ell,j} = \sum_{\ell=1}^{n} (B^t)_{j,\ell} (A^t)_{\ell,k} = (B^t A^t)_{j,k}$.
*Structural Commentary*: Transposition swaps the order of composition encoded by matrix multiplication.
*Key Insight*: Rows become columns, so the product order reverses. □

**Corollary 3.1.14.** *Matrix Inverses and Transposes Reverse Products*
*Let $A, B \in \mathrm{Mat}(n,n,\mathbb{F})$ be invertible. Then $AB$, $A^t$ are invertible, and $(AB)^{-1} = B^{-1}A^{-1}$; $(A^t)^{-1} = (A^{-1})^t$.*

*Proof.* First, $(B^{-1}A^{-1})(AB) = B^{-1}(A^{-1}A)B = B^{-1}B = I_n$; $(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AA^{-1} = I_n$. Hence $(AB)^{-1} = B^{-1}A^{-1}$. By Proposition 3.1.13, $(A^{-1})^t A^t = (AA^{-1})^t = I_n$; $A^t(A^{-1})^t = (A^{-1}A)^t = I_n$.
*Structural Commentary*: Inversion and transposition both reverse multiplication order.
*Key Insight*: Apply the defining inverse equations and reverse products. □

## 3.2 Coordinates and Matrix Representations

**Definition 3.2.1.** *Coordinate Vector*
Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of a finite-dimensional vector space $\mathcal{V}$.
If $\mathbf{v} = c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n$, then the *$\beta$-coordinate vector* of $\mathbf{v}$ is

$$[\mathbf{v}]_\beta := \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{F}^n.$$

The function $[\cdot]_\beta : \mathcal{V} \to \mathbb{F}^n$ is the *$\beta$-basis representation function*.

**Proposition 3.2.2.** *Coordinate Maps Are Linear Bijections*
*Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of a finite-dimensional vector space $\mathcal{V}$.*
*Then the coordinate map $[\cdot]_\beta : \mathcal{V} \to \mathbb{F}^n$ is linear and bijective.*

*Proof.* If $\mathbf{u} = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$ and $\mathbf{v} = b_1\mathbf{v}_1 + \cdots + b_n\mathbf{v}_n$, then for $c \in \mathbb{F}$,

$$[c\mathbf{u} + \mathbf{v}]_\beta = \begin{bmatrix} ca_1 + b_1 \\ \vdots \\ ca_n + b_n \end{bmatrix} = c[\mathbf{u}]_\beta + [\mathbf{v}]_\beta$$

Thus $[\cdot]_\beta$ is linear. It is injective because $[\mathbf{v}]_\beta = \mathbf{0}$ forces all basis coefficients of $\mathbf{v}$ to be zero, and it is surjective because every column $(c_1, \dots, c_n)^t \in \mathbb{F}^n$ equals $[c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n]_\beta$.
*Structural Commentary*: A basis converts abstract vectors into coordinate columns without losing structure.
*Key Insight*: Coordinates give a structure-preserving copy of the vector space, not extra information. □

**Definition 3.2.3.** *Matrix Representation*
Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of $\mathcal{V}$ and $\gamma = (\mathbf{w}_1, \dots, \mathbf{w}_m)$ a basis of $\mathcal{W}$. For $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, the matrix of $T$ relative to $(\beta, \gamma)$ is $[T]_{\gamma \leftarrow \beta} := \begin{bmatrix} [T(\mathbf{v}_1)]_\gamma & \cdots & [T(\mathbf{v}_n)]_\gamma \end{bmatrix} \in \mathrm{Mat}(m,n,\mathbb{F})$.

**Proposition 3.2.4.** *Matrices Respect Linear Combinations of Maps*
*Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of $\mathcal{V}$ and $\gamma$ be a basis of $\mathcal{W}$.*
*If $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ and $a \in \mathbb{F}$, then $[aS + T]_{\gamma \leftarrow \beta} = a[S]_{\gamma \leftarrow \beta} + [T]_{\gamma \leftarrow \beta}$.*

*Proof.* The $j$th column of $[aS + T]_{\gamma \leftarrow \beta}$ is $[(aS + T)\mathbf{v}_j]_\gamma = [aS\mathbf{v}_j + T\mathbf{v}_j]_\gamma = a[S\mathbf{v}_j]_\gamma + [T\mathbf{v}_j]_\gamma$, which is the $j$th column of $a[S]_{\gamma \leftarrow \beta} + [T]_{\gamma \leftarrow \beta}$.
*Structural Commentary*: Matrix representation records basis images column-by-column.
*Key Insight*: Linear combinations of maps become entrywise linear combinations of their matrices. □

**Proposition 3.2.5.** *Matrix Representation Acts on Coordinates*
*Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of $\mathcal{V}$, let $\gamma$ be a basis of $\mathcal{W}$, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.*
*Then for every $\mathbf{v} \in \mathcal{V}$, $[T(\mathbf{v})]_\gamma = [T]_{\gamma \leftarrow \beta}[\mathbf{v}]_\beta$.*

*Proof.* Write $\mathbf{v} = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$. Then linearity gives $T(\mathbf{v}) = a_1T(\mathbf{v}_1) + \cdots + a_nT(\mathbf{v}_n)$ and the right-hand side is exactly the linear combination of the columns of $[T]_{\gamma \leftarrow \beta}$ with coefficients $a_1, \dots, a_n$.
*Structural Commentary*: Matrix representation turns applying $T$ into multiplying coordinate columns.
*Key Insight*: The columns of $[T]_{\gamma \leftarrow \beta}$ are the coordinate images of the basis vectors. □

## 3.3 Matrix-Induced Maps and Rank

**Definition 3.3.1.** *Linear Map Induced by a Matrix*
For $A \in \mathrm{Mat}(m,n,\mathbb{F})$, the *linear map induced by $A$* is the map $T_A : \mathbb{F}^n \to \mathbb{F}^m$ defined by $T_A(\mathbf{x}) = A\mathbf{x}$.

**Proposition 3.3.2.** *Matrix-Induced Maps Are Linear*
*If $A \in \mathrm{Mat}(m,n,\mathbb{F})$, then $T_A : \mathbb{F}^n \to \mathbb{F}^m$ is linear.*

*Proof.* For $\mathbf{x},\mathbf{y} \in \mathbb{F}^n$ and $a \in \mathbb{F}$, $T_A(a\mathbf{x}+\mathbf{y}) = A(a\mathbf{x}+\mathbf{y}) = aA\mathbf{x} + A\mathbf{y} = aT_A(\mathbf{x}) + T_A(\mathbf{y})$.
*Structural Commentary*: Matrix multiplication preserves addition and scalar multiplication.
*Key Insight*: Every matrix defines a canonical linear map between coordinate spaces. □

**Corollary 3.3.3.** *Matrix Rank-Nullity*
*Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$. Then $\dim(\ker(A)) + \dim(\mathrm{col}(A)) = n$ and $\mathrm{nullity}(A) + \mathrm{rank}(A) = n$.*
*If $m = n$, then $\ker(A) = \{\mathbf{0}\}$ if and only if $\mathrm{col}(A) = \mathbb{F}^n$.*

*Proof.* Apply Theorem 1.7.3 to the induced linear map $T_A : \mathbb{F}^n \to \mathbb{F}^m$. By definition, $\ker(T_A) = \ker(A)$ and $\mathrm{range}(T_A) = \mathrm{col}(A)$, giving the displayed identities. When $m = n$, the remaining equivalence follows as:
($\Rightarrow$) If $\ker(A) = \{\mathbf{0}\}$, then $\dim(\mathrm{col}(A)) = n$. Since $\mathrm{col}(A) \subseteq \mathbb{F}^n$, Proposition 1.4.2 gives $\mathrm{col}(A) = \mathbb{F}^n$.
($\Leftarrow$) If $\mathrm{col}(A) = \mathbb{F}^n$, then $\dim(\mathrm{col}(A)) = n$, so the identity gives $\dim(\ker(A)) = 0$. Hence $\ker(A) = \{\mathbf{0}\}$.
*Structural Commentary*: Matrix rank-nullity is ordinary rank-nullity written in coordinates.
*Key Insight*: Kernel loss plus column-space dimension equals the number of columns. □

**Theorem 3.3.4.** *Row Rank Equals Column Rank*
*For $A \in \mathrm{Mat}(m,n,\mathbb{F})$, $\dim(\mathrm{row}(A)) = \dim(\mathrm{col}(A))$. Equivalently, $\mathrm{rank}(A^t) = \mathrm{rank}(A)$.*

*Proof.* Let $r = \mathrm{rank}(A)$. If $r = 0$, then $\mathrm{col}(A) = \{\mathbf{0}\}$, so every column of $A$ is zero and $A = 0$; the result follows. Suppose $r > 0$. Choose columns $\mathbf{a}_{j_1}, \ldots, \mathbf{a}_{j_r}$ forming a basis of $\mathrm{col}(A)$, and set $B = [\mathbf{a}_{j_1} \ \cdots \ \mathbf{a}_{j_r}] \in \mathrm{Mat}(m,r,\mathbb{F})$. For each column $\mathbf{a}_k$ of $A$, there is a unique $\mathbf{x}_k \in \mathbb{F}^r$ such that $\mathbf{a}_k = B\mathbf{x}_k$. With $X = [\mathbf{x}_1 \ \cdots \ \mathbf{x}_n] \in \mathrm{Mat}(r,n,\mathbb{F})$, we have $A = BX$. Therefore $A^t = X^tB^t$, so every column of $A^t$ is a linear combination of the $r$ columns of $X^t$. Hence $\mathrm{rank}(A^t) \le r = \mathrm{rank}(A)$. Applying the same argument to $A^t$ gives $\mathrm{rank}(A) \le \mathrm{rank}(A^t)$.
*Structural Commentary*: Column rank controls row rank by factoring $A$ through a basis of its column space.
*Key Insight*: Rows and columns measure the same intrinsic rank, viewed from transposed coordinates. □

**Perspective 3.3.5.** *Row and Column Spaces Have Different Hosts*
For $A \in \mathrm{Mat}(m,n,\mathbb{F})$, $\mathrm{row}(A) \subseteq \mathbb{F}^n$ and $\mathrm{col}(A) \subseteq \mathbb{F}^m$.
Hence, when $m \neq n$, Theorem 3.3.4 asserts only $\dim \mathrm{row}(A) = \dim \mathrm{col}(A)$, not equality of subspaces.

**Theorem 3.3.6.** *Four Fundamental Matrix Subspaces*
*Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$ and let $r = \mathrm{rank}(A)$. Then $\mathrm{row}(A), \ker(A) \subseteq \mathbb{F}^n$; $\mathrm{col}(A), \ker(A^t) \subseteq \mathbb{F}^m$, and $\dim(\mathrm{col}(A)) = \dim(\mathrm{row}(A)) = r$; $\dim(\ker(A)) = n - r$; $\dim(\ker(A^t)) = m - r$.*

*Proof.* The inclusions follow from the ambient dimensions of the rows and columns, and from the domains of $A$ and $A^t$. By definition, $\dim(\mathrm{col}(A)) = r$; by Theorem 3.3.4, $\dim(\mathrm{row}(A)) = r$.
Rank-nullity gives $\dim(\ker(A)) = n - \mathrm{rank}(A) = n - r$. Since $\mathrm{rank}(A^t) = \dim(\mathrm{col}(A^t)) = \dim(\mathrm{row}(A)) = r$, rank-nullity for $A^t : \mathbb{F}^m \to \mathbb{F}^n$ gives $\dim(\ker(A^t)) = m - \mathrm{rank}(A^t) = m - r$.
*Structural Commentary*: One rank determines all four fundamental dimensions.
*Key Insight*: Rank counts what survives; nullity counts what is killed. □




Figure 15: Four fundamental subspaces by host coordinate space.

**Corollary 3.3.7.** *Homogeneous Systems with More Variables Than Equations*
*If $A \in \mathrm{Mat}(m,n,\mathbb{F})$ with $n > m$, then the homogeneous system $A\mathbf{x} = \mathbf{0}$ has a nonzero solution.*

*Proof.* The induced map $T_A : \mathbb{F}^n \to \mathbb{F}^m$ has domain dimension larger than codomain dimension, so Corollary 1.7.4 shows that $T_A$ is not injective.
Hence $\ker(A) = \ker(T_A)$ contains a nonzero vector $\mathbf{x}$, and this vector satisfies $A\mathbf{x} = \mathbf{0}$.
*Structural Commentary*: Extra variables force a nontrivial kernel.
*Key Insight*: A nontrivial kernel is exactly a nonzero homogeneous solution. □

**Corollary 3.3.8.** *Inhomogeneous Systems with More Equations Than Variables*
*If $A \in \mathrm{Mat}(m,n,\mathbb{F})$ with $m > n$, then there exists $\mathbf{b} \in \mathbb{F}^m$ such that $A\mathbf{x} = \mathbf{b}$ has no solution.*

*Proof.* The induced map $T_A : \mathbb{F}^n \to \mathbb{F}^m$ has domain dimension smaller than codomain dimension, so Corollary 1.7.5 shows that $T_A$ is not surjective. Choose $\mathbf{b} \in \mathbb{F}^m \setminus \mathrm{range}(T_A)$. Then no $\mathbf{x} \in \mathbb{F}^n$ satisfies $A\mathbf{x} = \mathbf{b}$.
*Structural Commentary*: Too few variables cannot reach every possible right-hand side.
*Key Insight*: A missing range vector becomes an inconsistent inhomogeneous system. □

## 3.4 Invertibility and the Matrix-Map Dictionary

**Theorem 3.4.1.** *Columns of an Invertible Matrix Form a Basis*
*Let $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathrm{Mat}(n,n,\mathbb{F})$ be invertible. Then $(\mathbf{a}_1, \ldots, \mathbf{a}_n)$ is a basis of $\mathbb{F}^n$.*

*Proof.* The columns span $\mathbb{F}^n$ because for any $\mathbf{y} \in \mathbb{F}^n$, writing $\mathbf{x} = A^{-1}\mathbf{y}$ gives $\mathbf{y} = A\mathbf{x} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n$. They are linearly independent because if $c_1\mathbf{a}_1 + \cdots + c_n\mathbf{a}_n = \mathbf{0}$, then $A\mathbf{c} = \mathbf{0}$, so $\mathbf{c} = A^{-1}\mathbf{0} = \mathbf{0}$.
*Structural Commentary*: An invertible matrix is a coordinate isomorphism of $\mathbb{F}^n$. Its columns are the images of the standard coordinate directions.
*Key Insight*: Invertibility forces the columns to span and prevents nontrivial column relations. □

**Theorem 3.4.2.** *One-Sided Inverses of Square Matrices*
*Let $A, B \in \mathrm{Mat}(n,n,\mathbb{F})$. If $AB = I_n$, then $BA = I_n$.*

*Proof.* If $B\mathbf{x} = \mathbf{0}$, then $\mathbf{x} = AB\mathbf{x} = \mathbf{0}$, so the induced map $T_B : \mathbb{F}^n \to \mathbb{F}^n$ is injective. By Theorem 1.7.8, $T_B$ is surjective. For any $\mathbf{y} \in \mathbb{F}^n$, choose $\mathbf{x}$ with $\mathbf{y} = B\mathbf{x}$. Then $BA\mathbf{y} = BAB\mathbf{x} = B\mathbf{x} = \mathbf{y}$.
Thus $BA$ acts as the identity on every vector of $\mathbb{F}^n$, so $BA = I_n$.
*Structural Commentary*: In finite dimensions, injectivity and surjectivity coincide for square linear maps. A left inverse makes $B$ injective, hence onto, which forces the missing inverse relation.
*Key Insight*: A one-sided inverse of a square matrix is automatically two-sided. □

**Proposition 3.4.3.** *Composition and Matrix Multiplication*
*If $T \in \mathcal{L}(\mathcal{U},\mathcal{V})$ and $S \in \mathcal{L}(\mathcal{V},\mathcal{W})$, then $[ST]_{\delta\leftarrow\alpha} = [S]_{\delta\leftarrow\beta}[T]_{\beta\leftarrow\alpha}$ for any bases $\alpha$ of $\mathcal{U}$, $\beta$ of $\mathcal{V}$, and $\delta$ of $\mathcal{W}$.*

*Proof.* For every $\mathbf{u} \in \mathcal{U}$, $[(ST)\mathbf{u}]_\delta = [S]_{\delta\leftarrow\beta}[T\mathbf{u}]_\beta = [S]_{\delta\leftarrow\beta}[T]_{\beta\leftarrow\alpha}[\mathbf{u}]_\alpha$.
Since coordinate columns $[\mathbf{u}]_\alpha$ range over all of $\mathbb{F}^{\dim(\mathcal{U})}$, the representing matrices are equal.
*Structural Commentary*: Matrix multiplication is the coordinate shadow of composition of linear maps.
*Key Insight*: Applying $T$ then $S$ becomes multiplying first by the matrix of $T$, then by the matrix of $S$. □

**Theorem 3.4.4.** *Linear Maps and Matrices Are the Same Data*
*If $\dim(\mathcal{V}) = n$ and $\dim(\mathcal{W}) = m$, then choosing bases of $\mathcal{V}$ and $\mathcal{W}$ gives isomorphism $\mathcal{L}(\mathcal{V},\mathcal{W}) \cong \mathrm{Mat}(m,n,\mathbb{F})$. Consequently, $\dim(\mathcal{L}(\mathcal{V},\mathcal{W})) = mn$.*

*Proof.* Fix bases $\beta$ of $\mathcal{V}$ and $\gamma$ of $\mathcal{W}$. The map $T \mapsto [T]_{\gamma\leftarrow\beta}$ is linear. It is injective because a linear map whose matrix is zero sends every basis vector of $\mathcal{V}$ to $\mathbf{0}$, hence is the zero map. It is surjective because, for any $A = [a_{ij}] \in \mathrm{Mat}(m,n,\mathbb{F})$, there is a unique linear map satisfying $T(\mathbf{v}_j) = a_{1j}\mathbf{w}_1 + \cdots + a_{mj}\mathbf{w}_m$, $j = 1, \ldots, n$.
Thus $\mathcal{L}(\mathcal{V},\mathcal{W}) \cong \mathrm{Mat}(m,n,\mathbb{F})$, whose standard entry matrices form a basis of length $mn$.
*Structural Commentary*: Choosing bases converts every linear map into one and only one array of scalars.
*Key Insight*: A matrix is exactly the list of coordinate images of the domain basis vectors. □

**Perspective 3.4.5.** *Finite-Support Coordinate Model*
Proposition 3.2.2 identifies an $n$-dimensional vector space with $\mathbb{F}^n$, and Theorem 3.4.4 identifies linear maps with matrices after bases are chosen. The space $\mathbb{F}^{(\mathbb{N})}$ is the countable standard-coordinate model with finite support. This is the field analogue of the free-module notation $(R^B)_0$.

**Motivation 3.4.6.** *Coordinate Encoding*
This is the key bridge between abstraction and computation. The linear map is basis-free; the matrix is what the map looks like after coordinates have been chosen.

## 3.5 Change of Basis and Coordinate Transformations

**Definition 3.5.1.** *Change of Basis Matrix*
Let $\beta$ and $\gamma$ be two bases of the same vector space $\mathcal{V}$. The change-of-basis matrix from $\beta$-coordinates to $\gamma$-coordinates is $C_{\gamma\leftarrow\beta} := [I_{\mathcal{V}}]_{\gamma\leftarrow\beta}$. Thus $[\mathbf{v}]_\gamma = C_{\gamma\leftarrow\beta}[\mathbf{v}]_\beta$ for every $\mathbf{v} \in \mathcal{V}$.




Figure 16: Change of basis: $[\mathbf{v}]_\gamma = C_{\gamma\leftarrow\beta}[\mathbf{v}]_\beta$.

**Theorem 3.5.2.** *Change-of-Basis Matrices*
*Let $\mathcal{V}$ be $n$-dimensional and let $\beta$ be a basis of $\mathcal{V}$.*

*(i) If $\gamma$ is a basis of $\mathcal{V}$, then $C_{\gamma\leftarrow\beta}$ is invertible and $(C_{\gamma\leftarrow\beta})^{-1} = C_{\beta\leftarrow\gamma}$.*
*(ii) If $S \in \mathrm{Mat}(n, n, \mathbb{F})$ is invertible, then there is a basis $\gamma$ of $\mathcal{V}$ such that $S = C_{\gamma\leftarrow\beta}$.*

*Proof.*

(i) For any $\mathbf{v} \in \mathcal{V}$, $C_{\beta\leftarrow\gamma}C_{\gamma\leftarrow\beta}[\mathbf{v}]_\beta = [\mathbf{v}]_\beta$, $C_{\gamma\leftarrow\beta}C_{\beta\leftarrow\gamma}[\mathbf{v}]_\gamma = [\mathbf{v}]_\gamma$.
Since $[\cdot]_\beta$ and $[\cdot]_\gamma$ are onto $\mathbb{F}^n$, both products equal $I_n$. Hence $C_{\gamma\leftarrow\beta}$ is invertible with inverse $C_{\beta\leftarrow\gamma}$.
(ii) Write $S^{-1} = [\mathbf{s}_1 \ \cdots \ \mathbf{s}_n]$. For each $j$, choose the unique $\mathbf{w}_j \in \mathcal{V}$ with $[\mathbf{w}_j]_\beta = \mathbf{s}_j$, and set $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$.
Since columns of $S^{-1}$ form a basis of $\mathbb{F}^n$ and $[\cdot]_\beta$ is an isomorphism, $\gamma$ is a basis of $\mathcal{V}$.
Moreover $C_{\beta\leftarrow\gamma} = S^{-1}$, so $C_{\gamma\leftarrow\beta} = S$.

*Structural Commentary*: Change of basis is an invertible coordinate rewrite.
*Key Insight*: Every basis change is reversible, and every invertible matrix is one. □

**Theorem 3.5.3.** *Change of Bases for a Linear Map*
*Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Let $\beta, \beta'$ be bases of $\mathcal{V}$ and let $\gamma, \gamma'$ be bases of $\mathcal{W}$. Then $[T]_{\gamma'\leftarrow\beta'} = C_{\gamma'\leftarrow\gamma}[T]_{\gamma\leftarrow\beta}C_{\beta\leftarrow\beta'}$.*

*Proof.* For every $\mathbf{v} \in \mathcal{V}$, $[T\mathbf{v}]_{\gamma'} = C_{\gamma'\leftarrow\gamma}[T\mathbf{v}]_\gamma = C_{\gamma'\leftarrow\gamma}[T]_{\gamma\leftarrow\beta}[\mathbf{v}]_\beta = C_{\gamma'\leftarrow\gamma}[T]_{\gamma\leftarrow\beta}C_{\beta\leftarrow\beta'}[\mathbf{v}]_{\beta'}$.
Since $[\cdot]_{\beta'}$ is onto, the representing matrices are equal.
*Structural Commentary*: Basis changes act by $C_{\gamma'\leftarrow\gamma}$ on the left and $C_{\beta\leftarrow\beta'}$ on the right.
*Key Insight*: Domain changes act on input coordinates; codomain changes act on output coordinates. □

## 3.6 Rank and Invertibility Consequences

**Corollary 3.6.1.** *Basis Columns Give an Invertible Matrix*
*If $(\mathbf{a}_1, \ldots, \mathbf{a}_n)$ is a basis of $\mathbb{F}^n$, then $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathrm{Mat}(n, n, \mathbb{F})$ is invertible.*

*Proof.* Let $\epsilon = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be the standard basis and $\alpha = (\mathbf{a}_1, \ldots, \mathbf{a}_n)$. Then $C_{\epsilon\leftarrow\alpha} = \begin{bmatrix}[\mathbf{a}_1]_\epsilon & \cdots & [\mathbf{a}_n]_\epsilon\end{bmatrix} = A$.
Since $\alpha$ is a basis, Theorem 3.5.2 gives that $A$ is invertible.
*Structural Commentary*: Basis columns form a change-of-basis matrix.
*Key Insight*: Changing coordinates is reversible. □

**Theorem 3.6.2.** *Completing a Full-Column-Rank Matrix*
*If $X \in \mathrm{Mat}(m, n, \mathbb{F})$ has rank $n < m$, then $\exists\ X' \in \mathrm{Mat}(m, m-n, \mathbb{F})$ s.t. $\begin{bmatrix}X & X'\end{bmatrix} \in \mathrm{Mat}(m, m, \mathbb{F})$ is invertible.*

*Proof.* Write $X = [\mathbf{x}_1 \ \cdots \ \mathbf{x}_n]$. Since $\mathrm{rank}(X) = n$, the columns $\mathbf{x}_1, \ldots, \mathbf{x}_n$ are linearly independent in $\mathbb{F}^m$. By Theorem 1.3.12, extend this list to a basis $(\mathbf{x}_1, \ldots, \mathbf{x}_n, \mathbf{x}_{n+1}, \ldots, \mathbf{x}_m)$ of $\mathbb{F}^m$.
Set $X' = [\mathbf{x}_{n+1} \ \cdots \ \mathbf{x}_m]$. Corollary 3.6.1 shows that $\begin{bmatrix}X & X'\end{bmatrix}$ is invertible.
*Structural Commentary*: Full column rank means the existing columns are independent directions. Extending them to a basis turns the rectangular matrix into an invertible square matrix.
*Key Insight*: Independent columns can always be completed to a basis in finite dimensions. □

**Corollary 3.6.3.** *Invertibility and Full Rank*
*For $A \in \mathrm{Mat}(n, n, \mathbb{F})$, $A$ is invertible $\iff \mathrm{rank}(A) = n$.*

*Proof.*

($\Rightarrow$) If $A$ is invertible, its columns form a basis of $\mathbb{F}^n$ by Theorem 3.4.1; hence $\mathrm{rank}(A) = n$.
($\Leftarrow$) If $\mathrm{rank}(A) = n$, then $\mathrm{col}(A) = \mathbb{F}^n$.
Thus the columns of $A$ form a basis by Proposition 1.4.3, so Corollary 3.6.1 gives that $A$ is invertible.

*Structural Commentary*: Full rank is square-matrix invertibility.
*Key Insight*: Invertibility means the columns fill $\mathbb{F}^n$. □

**Corollary 3.6.4.** *Transpose Preserves Invertibility*
*For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, the matrix $A$ is invertible if and only if $A^t$ is invertible.*

*Proof.* By Theorem 3.3.4, $\mathrm{rank}(A) = \mathrm{rank}(A^t)$. Thus $\mathrm{rank}(A) = n$ if and only if $\mathrm{rank}(A^t) = n$.
Corollary 3.6.3 gives the result.
*Structural Commentary*: Transposition exchanges rows and columns without changing rank.
*Key Insight*: A square matrix is invertible exactly when either its rows or columns have full rank. □

**Theorem 3.6.5.** *Full-Rank Products Preserve Rank*
*Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$. If $X \in \mathrm{Mat}(p,m,\mathbb{F})$ has full column rank and $Y \in \mathrm{Mat}(n,q,\mathbb{F})$ has full row rank, then $\mathrm{rank}(A) = \mathrm{rank}(XAY)$. This holds when $X$ and $Y$ are square invertible matrices of appropriate sizes.*

*Proof.* Because $X$ has full column rank, the induced map $L_X : \mathbb{F}^m \to \mathbb{F}^p$, $L_X(\mathbf{u}) = X\mathbf{u}$, has kernel $\{\mathbf{0}\}$.
Thus $\mathbf{z} \in \ker(XA) \iff X(A\mathbf{z}) = \mathbf{0} \iff A\mathbf{z} = \mathbf{0} \iff \mathbf{z} \in \ker(A)$.
Since $XA$ and $A$ have the same number of columns, Corollary 3.3.3 gives $\mathrm{rank}(XA) = \mathrm{rank}(A)$.
Because $Y$ has full row rank, $Y^t$ has full column rank. Applying the left-multiplication result to $A^t$ and $Y^t$, then using Theorem 3.3.4, gives $\mathrm{rank}(AY) = \mathrm{rank}((AY)^t) = \mathrm{rank}(Y^tA^t) = \mathrm{rank}(A^t) = \mathrm{rank}(A)$.
Applying the left-multiplication result once more to $AY$ gives $\mathrm{rank}(XAY) = \mathrm{rank}(AY) = \mathrm{rank}(A)$.
*Structural Commentary*: Multiplication by a full-column-rank matrix on the left loses no output information, while full row rank on the right loses no input information.
*Key Insight*: Rank changes only when multiplication introduces a new kernel or collapses the row side. □

## 3.7 Block Matrix Algebra

**Definition 3.7.1.** *Block Matrices*
Suppose $\mathcal{V} = \mathcal{U}_1 \oplus \mathcal{U}_2$ and $\mathcal{W} = \mathcal{Z}_1 \oplus \mathcal{Z}_2$. If $T \in \mathcal{L}(\mathcal{V},\mathcal{W})$, relative to bases adapted to these decompositions in the sense of Definition 2.3.5, the matrix of $T$ has block form $[T] = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$, where each block records the part of $T$ mapping one summand of the domain into one summand of the codomain.

**Definition 3.7.2.** *Conformal Block Partitions*
For a block matrix $A = [A_{ij}]$, the submatrices $A_{ij}$ are called *blocks*. Two block partitions are *conformal for addition* when corresponding blocks have the same sizes. They are *conformal for multiplication* when every block product appearing in the usual row-column block multiplication is defined.

**Proposition 3.7.3.** *Stacks Intersect Kernels, Concatenations Sum Columns*
*Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$, $B \in \mathrm{Mat}(p,n,\mathbb{F})$, $C \in \mathrm{Mat}(m,r,\mathbb{F})$, and $D \in \mathrm{Mat}(m,s,\mathbb{F})$.*
*Then* $\ker \begin{bmatrix} A \\ B \end{bmatrix} = \ker(A) \cap \ker(B)$; *and* $\mathrm{col} \begin{bmatrix} C & D \end{bmatrix} = \mathrm{col}(C) + \mathrm{col}(D)$.

*Proof.* For $\mathbf{x} \in \mathbb{F}^n$, $\mathbf{x} \in \ker \begin{bmatrix} A \\ B \end{bmatrix} \iff \begin{bmatrix} A \\ B \end{bmatrix} \mathbf{x} = \mathbf{0} \iff A\mathbf{x} = \mathbf{0}$ and $B\mathbf{x} = \mathbf{0} \iff \mathbf{x} \in \ker(A) \cap \ker(B)$.

For $\mathbf{y} \in \mathbb{F}^m$, $\mathbf{y} \in \mathrm{col} \begin{bmatrix} C & D \end{bmatrix} \iff \exists\, \mathbf{u} \in \mathbb{F}^r,\ \mathbf{v} \in \mathbb{F}^s$ such that $\mathbf{y} = \begin{bmatrix} C & D \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix}$
$\iff \exists\, \mathbf{u}, \mathbf{v}$ such that $\mathbf{y} = C\mathbf{u} + D\mathbf{v} \iff \mathbf{y} \in \mathrm{col}(C) + \mathrm{col}(D)$.
*Structural Commentary*: Stacking intersects constraints; concatenation adds column directions.
*Key Insight*: Stacks encode intersections of nullspaces, while concatenations encode sums of column spaces. □

**Proposition 3.7.4.** *Block Matrix Arithmetic*
*For conformal $2 \times 2$ block matrices,*

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} + \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}+B_{11} & A_{12}+B_{12} \\ A_{21}+B_{21} & A_{22}+B_{22} \end{bmatrix}$$

*and*

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11}+A_{12}B_{21} & A_{11}B_{12}+A_{12}B_{22} \\ A_{21}B_{11}+A_{22}B_{21} & A_{21}B_{12}+A_{22}B_{22} \end{bmatrix}.$$

*Proof.* Each displayed block consists of the same entries obtained by ordinary entrywise addition or ordinary row-column multiplication, with the common indices grouped according to the partition.
*Structural Commentary*: This is ordinary matrix arithmetic with indices grouped into rectangular packages.
*Key Insight*: Conformality is exactly the condition that the grouped row-column products make sense. □

**Proposition 3.7.5.** *Block Centrosymmetry and Commutation*
*Let* $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathrm{Mat}(2n, 2n, \mathbb{F})$, $L_{2n} = \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix}$.
*Then* $M$ *has block centrosymmetric form* $A = D$ *and* $B = C$ *if and only if* $L_{2n}M = ML_{2n}$.

*Proof.* Block multiplication gives $L_{2n}M = \begin{bmatrix} C & D \\ A & B \end{bmatrix}$ and $ML_{2n} = \begin{bmatrix} B & A \\ D & C \end{bmatrix}$.
The two matrices are equal exactly when $A = D$ and $B = C$.
*Structural Commentary*: The matrix $L_{2n}$ swaps the two block coordinates. Commuting with that swap means $M$ treats the two block positions symmetrically.
*Key Insight*: Compute both block products; equality forces matching opposite blocks. □

**Definition 3.7.6.** *Direct Sum of Matrices*
For $A \in \mathrm{Mat}(m, n, \mathbb{F})$ and $B \in \mathrm{Mat}(p, q, \mathbb{F})$, the *direct sum* of $A$ and $B$ is $A \oplus B := \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in \mathrm{Mat}(m+p, n+q, \mathbb{F})$.

## 3.8 Matrix Operations and Identities

**Definition 3.8.1.** *Polynomial Evaluation on a Square Matrix*
If $A \in \mathrm{Mat}(n, n, \mathbb{F})$ and $p(z) = \sum_{j=0}^{d} c_j z^j \in \mathbb{F}[z]$, then $p(A) := \sum_{j=0}^{d} c_j A^j$, $A^0 := I_n$.

**Proposition 3.8.2.** *Algebra of Matrix Direct Sums*
*Whenever the expressions are defined,*
*(i)* $a(A \oplus B) + b(C \oplus D) = (aA + bC) \oplus (aB + bD)$;
*(ii)* $(A \oplus B)(C \oplus D) = AC \oplus BD$;
*(iii) if* $A$ *and* $B$ *are square, then* $(A \oplus B)^k = A^k \oplus B^k$ *for every positive integer* $k$;
*(iv) if* $A$ *and* $B$ *are square and* $p$ *is a polynomial, then* $p(A \oplus B) = p(A) \oplus p(B)$.

*Proof.* Block addition gives $a(A \oplus B) + b(C \oplus D) = \begin{bmatrix} aA + bC & 0 \\ 0 & aB + bD \end{bmatrix} = (aA + bC) \oplus (aB + bD)$.
Block multiplication gives $(A \oplus B)(C \oplus D) = \begin{bmatrix} AC & 0 \\ 0 & BD \end{bmatrix} = AC \oplus BD$.
For powers, the case $k = 1$ is the definition. If $(A \oplus B)^k = A^k \oplus B^k$, then part (ii) gives $(A \oplus B)^{k+1} = A^{k+1} \oplus B^{k+1}$. Hence (iii) follows by induction. For the polynomial statement, suppose $A \in \mathrm{Mat}(m, m, \mathbb{F})$ and $B \in \mathrm{Mat}(r, r, \mathbb{F})$. For $j = 0$, Definition 3.8.1 gives $(A \oplus B)^0 = I_{m+r} = I_m \oplus I_r = A^0 \oplus B^0$. If $p(z) = \sum_{j=0}^{d} c_j z^j$,
then $p(A \oplus B) = \sum_{j=0}^{d} c_j (A^j \oplus B^j) = \left( \sum_{j=0}^{d} c_j A^j \right) \oplus \left( \sum_{j=0}^{d} c_j B^j \right) = p(A) \oplus p(B)$.
*Structural Commentary*: A direct sum matrix keeps the two blocks independent under linear combinations, products, powers, and polynomials.
*Key Insight*: Block diagonal matrices behave componentwise. □

**Proposition 3.8.3.** *Diagonal Multiplication Scales Rows and Columns*
*Let* $A = [a_{ij}] \in \mathrm{Mat}(m, n, \mathbb{F})$, *let* $\Lambda = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$, *and let* $M = \mathrm{diag}(\mu_1, \ldots, \mu_m)$. *Then* $A\Lambda = [\lambda_j a_{ij}]$ *and* $MA = [\mu_i a_{ij}]$. *Thus right multiplication by* $\Lambda$ *scales columns of* $A$, *left multiplication by* $M$ *scales rows of* $A$.

*Proof.* The $(i, j)$ entry of $A\Lambda$ is $\sum_{r=1}^{n} a_{ir}\Lambda_{rj} = a_{ij}\lambda_j$, and the $(i, j)$ entry of $MA$ is $\sum_{r=1}^{m} M_{ir}a_{rj} = \mu_i a_{ij}$.
*Structural Commentary*: Diagonal matrices act by independent scalar weights on coordinate directions.
*Key Insight*: Multiplication on the right scales columns, while multiplication on the left scales rows. □

**Proposition 3.8.4.** *Inverse of an Upper Block-Triangular Matrix*
*Let* $Y \in \mathrm{Mat}(m, m, \mathbb{F})$ *and* $Z \in \mathrm{Mat}(n, n, \mathbb{F})$ *be invertible, and let* $X \in \mathrm{Mat}(m, n, \mathbb{F})$.
*Then* $\begin{bmatrix} Y & X \\ 0 & Z \end{bmatrix}^{-1} = \begin{bmatrix} Y^{-1} & -Y^{-1}XZ^{-1} \\ 0 & Z^{-1} \end{bmatrix}$.

*Proof.* Let $M = \begin{bmatrix} Y & X \\ 0 & Z \end{bmatrix}$ and $N = \begin{bmatrix} Y^{-1} & -Y^{-1}XZ^{-1} \\ 0 & Z^{-1} \end{bmatrix}$. Proposition 3.7.4 gives

$$MN = \begin{bmatrix} I_m & -XZ^{-1} + XZ^{-1} \\ 0 & I_n \end{bmatrix} = I_{m+n}, \qquad NM = \begin{bmatrix} I_m & Y^{-1}X - Y^{-1}X \\ 0 & I_n \end{bmatrix} = I_{m+n}$$

. Thus $N = M^{-1}$.
*Structural Commentary*: Upper block-triangular inversion preserves the zero lower-left block. The off-diagonal

term is forced by cancelling the original upper-right block.
*Key Insight*: Inverting a block triangular matrix inverts diagonal blocks and corrects the coupling block. □

**Theorem 3.8.5.** *Inverse of an Upper Triangular Matrix*
*Let $A = [a_{ij}] \in \mathrm{Mat}(n, n, \mathbb{F})$ be upper triangular with nonzero diagonal entries. Then $A$ is invertible, $A^{-1}$ is upper triangular, and the diagonal entries of $A^{-1}$ are $a_{11}^{-1}, \dots, a_{nn}^{-1}$.*

*Proof.* We argue by induction on $n$. The case $n = 1$ is immediate. For $n > 1$, write $A = \begin{bmatrix} B & \mathbf{x} \\ 0 & a_{nn} \end{bmatrix}$, where $B$ is $(n-1) \times (n-1)$ upper triangular with nonzero diagonal entries.
By induction, $B^{-1}$ is upper triangular with diagonal entries $a_{11}^{-1}, \dots, a_{n-1,n-1}^{-1}$.
Proposition 3.8.4 gives $A^{-1} = \begin{bmatrix} B^{-1} & -B^{-1}\mathbf{x}a_{nn}^{-1} \\ 0 & a_{nn}^{-1} \end{bmatrix}$, which is upper triangular and has the asserted diagonal.
*Structural Commentary*: An upper triangular matrix splits into a smaller upper triangular block and one final scalar block. The block inverse formula preserves this triangular structure.
*Key Insight*: Nonzero diagonals make blockwise triangular inversion possible. □

# 4 Duality and Multilinear Structure

## 4.1 Dual Spaces and Dual Bases

Duality studies vectors through scalar-valued tests. Functionals, dual maps, annihilators, transposes, bilinear forms, and tensor products extend linear structure into multilinear language.

**Definition 4.1.1.** *Linear Functional and Dual Space*
A *linear functional* on $\mathcal{V}$ is a linear map from $\mathcal{V}$ to $\mathbb{F}$. The *dual space* of $\mathcal{V}$ is $\mathcal{V}' := \mathcal{L}(\mathcal{V}, \mathbb{F})$.

**Example 4.1.2.** *Evaluation and Integral Functionals*
The evaluation map $\mathrm{ev}_0 : \mathbb{F}[x] \to \mathbb{F}$, $p \mapsto p(0)$ is a linear functional, since $\mathrm{ev}_0(\alpha p + \beta q) = \alpha p(0) + \beta q(0)$ for all $\alpha, \beta \in \mathbb{F}$ and $p, q \in \mathbb{F}[x]$. If $C[a, b]$ is the real vector space of continuous functions on $[a, b]$, then $I : C[a, b] \to \mathbb{R}$, $I(f) := \int_a^b f(x)\,dx$ is a linear functional, since $I(\alpha f + \beta g) = \alpha I(f) + \beta I(g)$ $\forall \alpha, \beta \in \mathbb{R}$ and $f, g \in C[a, b]$.

**Definition 4.1.3.** *Dual Basis*
Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of $\mathcal{V}$. The *dual basis* of $\beta$ is the list $\beta' = (\varphi_1, \dots, \varphi_n)$ in $\mathcal{V}'$ defined by

$$\varphi_j(\mathbf{v}_k) = \begin{cases} 1, & k = j, \\ 0, & k \neq j. \end{cases}$$

Existence and uniqueness of each $\varphi_j$ follow from Proposition 1.5.11.

**Theorem 4.1.4.** *Dual Basis and Dimension of the Dual Space*
*Let $\mathcal{V}$ be finite-dimensional with basis $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$, and let $\beta' = (\varphi_1, \dots, \varphi_n)$ be the dual basis. Then $\beta'$ is a basis of $\mathcal{V}'$. In particular, $\dim(\mathcal{V}') = \dim(\mathcal{V})$.*

*Proof.* First suppose $a_1\varphi_1 + \cdots + a_n\varphi_n = 0$. Evaluating at $\mathbf{v}_j$ gives $a_j = 0$ for each $j$, so $\beta'$ is linearly independent. Now let $\varphi \in \mathcal{V}'$. For $\mathbf{v} = c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n$, $\left(\sum_{j=1}^{n} \varphi(\mathbf{v}_j)\varphi_j\right)(\mathbf{v}) = \sum_{j=1}^{n} \varphi(\mathbf{v}_j)c_j = \varphi(\mathbf{v})$.
Thus $\varphi = \sum_{j=1}^{n} \varphi(\mathbf{v}_j)\varphi_j$, so $\beta'$ spans $\mathcal{V}'$.
*Structural Commentary*: A basis of $\mathcal{V}$ induces the dual coordinate system.
*Key Insight*: A linear functional is determined exactly by its values on a basis. □

**Proposition 4.1.5.** *Coordinate Functionals for a Hamel Basis*
*Let $\mathcal{B} = \{\mathbf{b}_i\}_{i \in I}$ be a Hamel basis of $\mathcal{V}$, define $\varepsilon_i \in \mathcal{V}'$ by $\varepsilon_i\left(\sum_{j \in F} a_j\mathbf{b}_j\right) := a_i$, $F \subseteq I$ finite, $a_i = 0$ if $i \notin F$.*
*Then $\{\varepsilon_i\}_{i \in I}$ is linearly independent in $\mathcal{V}'$. Moreover, the following are equivalent:*

*(i)* $\mathrm{span}\{\varepsilon_i : i \in I\} = \mathcal{V}'$;
*(ii)* $|I| < \infty$;
*(iii)* $\dim(\mathcal{V}) < \infty$.

*Proof.* If $\sum_{r=1}^{n} c_r\varepsilon_{i_r} = 0$ with distinct $i_r$, then $0 = \left(\sum_{r=1}^{n} c_r\varepsilon_{i_r}\right)(\mathbf{b}_{i_s}) = c_s$ $\forall s$. Hence $\{\varepsilon_i\}_{i \in I}$ linearly independent.

*(i)* $\Rightarrow$ *(ii)*. Suppose $\mathrm{span}\{\varepsilon_i : i \in I\} = \mathcal{V}'$. Put $\mathrm{supp}_{\mathcal{B}}(\psi) := \{i \in I : \psi(\mathbf{b}_i) \neq 0\}$.
If $I$ were infinite, define $\varphi \in \mathcal{V}'$ by $\varphi(\mathbf{b}_i) = 1$ for all $i \in I$.
Then $\mathrm{supp}_{\mathcal{B}}(\varphi) = I$, while $\mathrm{supp}_{\mathcal{B}}(\psi)$ is finite for every $\psi \in \mathrm{span}\{\varepsilon_i : i \in I\}$, a contradiction.

(*ii*) $\Rightarrow$ (*iii*). Since $\mathcal{B}$ is a basis, $\dim(\mathcal{V}) = |I| < \infty$.
(*iii*) $\Rightarrow$ (*i*). If $\dim(\mathcal{V}) < \infty$, then Theorem 4.1.4 gives $\operatorname{span}\{\varepsilon_i : i \in I\} = \mathcal{V}'$.

*Structural Commentary*: Coordinate spans are finite-support.
*Key Insight*: Infinite-dimensional coordinate functionals are independent, not spanning. □

**Theorem 4.1.6.** *Dimension of the Algebraic Dual*
*For every $\mathbb{F}$-vector space $\mathcal{V}$, $\dim(\mathcal{V}) \le \dim(\mathcal{V}')$. Moreover, the following are equivalent:*

*(i)* $\dim(\mathcal{V}) = \dim(\mathcal{V}')$;
*(ii)* $\dim(\mathcal{V}) < \infty$.

*Proof.* Let $\mathcal{B} = \{\mathbf{b}_i\}_{i\in I}$ be a Hamel basis. By Proposition 4.1.5, $\{\varepsilon_i\}_{i\in I} \subseteq \mathcal{V}'$ is linearly independent; hence $\dim(\mathcal{V}) = |I| \le \dim(\mathcal{V}')$. For the equality criterion:

(*i*) $\Rightarrow$ (*ii*). Suppose $\dim(\mathcal{V}) = \dim(\mathcal{V}')$. If $I$ were infinite, then $\mathcal{V} \cong (\mathbb{F}^I)_0$ and $\mathcal{V}' \cong \mathbb{F}^I$.
Let $\mathbb{F}_0 \subseteq \mathbb{F}$ be the prime subfield. Since $\mathbb{F}_0$ is countable, $\dim_{\mathbb{F}_0}\left((\mathbb{F}_0^I)_0\right) = |I|$ and $\left|\mathbb{F}_0^I\right| > |I|$ by Cantor, so $\dim_{\mathbb{F}_0}(\mathbb{F}_0^I) > |I|$; hence $\dim_{\mathbb{F}_0}\left((\mathbb{F}_0^I)_0\right) = |I| < \dim_{\mathbb{F}_0}(\mathbb{F}_0^I)$.
Every $\mathbb{F}_0$-linearly independent family in $\mathbb{F}_0^I$ is $\mathbb{F}$-linearly independent in $\mathbb{F}^I$, by expanding coefficients in an $\mathbb{F}_0$-basis of $\mathbb{F}$. Thus $\dim_{\mathbb{F}}(\mathcal{V}) = \dim_{\mathbb{F}}\left((\mathbb{F}^I)_0\right) < \dim_{\mathbb{F}}(\mathbb{F}^I) = \dim_{\mathbb{F}}(\mathcal{V}')$, a contradiction. Hence $\dim(\mathcal{V}) < \infty$.
(*ii*) $\Rightarrow$ (*i*). If $\dim(\mathcal{V}) < \infty$, then Theorem 4.1.4 gives $\dim(\mathcal{V}) = \dim(\mathcal{V}')$.

*Structural Commentary*: Algebraic duals have arbitrary support.
*Key Insight*: Infinite algebraic duals are bigger than their original spaces. □

**Proposition 4.1.7.** *Functionals Separate Points*
*Let $\mathcal{V}$ be a vector space.*

*(i) If $\mathbf{v} \neq \mathbf{0}$, then there exists $\varphi \in \mathcal{V}'$ such that $\varphi(\mathbf{v}) \neq 0$.*
*(ii) A vector $\mathbf{v} \in \mathcal{V}$ is zero if and only if $\varphi(\mathbf{v}) = 0$ for every $\varphi \in \mathcal{V}'$.*
*(iii) If $\mathcal{U} \le \mathcal{V}$ and $\mathbf{x} \notin \mathcal{U}$, then there exists $\varphi \in \mathcal{V}'$ such that $\varphi|_{\mathcal{U}} = 0$ and $\varphi(\mathbf{x}) = 1$.*
*(iv) If $\varphi \in \mathcal{V}'$ and $\varphi(\mathbf{x}) \neq 0$, then $\mathcal{V} = \operatorname{span}(\mathbf{x}) \oplus \ker(\varphi)$.*
*(v) If nonzero $\varphi, \psi \in \mathcal{V}'$ have the same kernel, then $\varphi = a\psi$ for some nonzero $a \in \mathbb{F}$.*

*Proof.*

(i) Extend $\{\mathbf{v}\}$ to a Hamel basis $\mathcal{B}$ of $\mathcal{V}$ by Theorem 1.4.6. Define $\varphi$ on $\mathcal{B}$ by $\varphi(\mathbf{v}) = 1$ and $\varphi(\mathbf{b}) = 0$ for $\mathbf{b} \in \mathcal{B} \setminus \{\mathbf{v}\}$, then extend linearly to finite basis expansions.
(ii) If $\mathbf{v} = \mathbf{0}$, then $\varphi(\mathbf{v}) = 0$ for every $\varphi \in \mathcal{V}'$. Conversely, if $\varphi(\mathbf{v}) = 0$ for every $\varphi \in \mathcal{V}'$ and $\mathbf{v} \neq \mathbf{0}$, then (i) gives some $\psi \in \mathcal{V}'$ with $\psi(\mathbf{v}) \neq 0$, a contradiction.
(iii) Choose a Hamel basis $\mathcal{B}_{\mathcal{U}}$ of $\mathcal{U}$. Since $\mathbf{x} \notin \mathcal{U} = \operatorname{span}(\mathcal{B}_{\mathcal{U}})$, the set $\mathcal{B}_{\mathcal{U}} \cup \{\mathbf{x}\}$ is linearly independent. Extend it to a Hamel basis $\mathcal{B}$ of $\mathcal{V}$ by Theorem 1.4.6. Define $\varphi$ on $\mathcal{B}$ by $\varphi(\mathbf{x}) = 1$ and $\varphi(\mathbf{b}) = 0$ for every $\mathbf{b} \in \mathcal{B} \setminus \{\mathbf{x}\}$, then extend linearly to finite basis expansions.
(iv) Since $\varphi(\mathbf{x}) \neq 0$, $\operatorname{span}(\mathbf{x}) \cap \ker(\varphi) = \{\mathbf{0}\}$.
For any $\mathbf{v} \in \mathcal{V}$, $\mathbf{v} = \frac{\varphi(\mathbf{v})}{\varphi(\mathbf{x})}\mathbf{x} + \left(\mathbf{v} - \frac{\varphi(\mathbf{v})}{\varphi(\mathbf{x})}\mathbf{x}\right)$, and the second summand lies in $\ker(\varphi)$.
(v) Let $\mathcal{K} = \ker(\varphi) = \ker(\psi)$ and choose $\mathbf{x} \notin \mathcal{K}$. By (iv), every $\mathbf{v} \in \mathcal{V}$ can be written $\mathbf{v} = c\mathbf{x} + \mathbf{k}$ with $\mathbf{k} \in \mathcal{K}$. Setting $a = \varphi(\mathbf{x})/\psi(\mathbf{x})$ gives $\varphi(\mathbf{v}) = c\varphi(\mathbf{x}) = a\,c\psi(\mathbf{x}) = a\psi(\mathbf{v})$.

*Structural Commentary*: Dual vectors separate points and define hyperplanes.
*Key Insight*: A nonzero functional splits the space into one visible direction plus its kernel. □

**Corollary 4.1.8.** *Kernels of Nonzero Functionals*
*If $0 \neq \varphi \in \mathcal{V}'$, then $\varphi$ is surjective onto $\mathbb{F}$, $\mathcal{V}/\ker(\varphi) \cong \mathbb{F}$, and $\operatorname{codim}_{\mathcal{V}}(\ker\varphi) = 1$.*

*Proof.* Choose $\mathbf{x}$ with $\varphi(\mathbf{x}) \neq 0$. For each $a \in \mathbb{F}$, $\varphi\big(a\varphi(\mathbf{x})^{-1}\mathbf{x}\big) = a$, so $\operatorname{im}\varphi = \mathbb{F}$.
Hence, by Theorem 2.2.12, $\mathcal{V}/\ker(\varphi) \cong \operatorname{im}\varphi = \mathbb{F}$ and $\operatorname{codim}_{\mathcal{V}}(\ker\varphi) = \dim(\mathcal{V}/\ker\varphi) = 1$.
*Structural Commentary*: A nonzero functional sees exactly one scalar direction and kills a hyperplane.
*Key Insight*: Kernels of nonzero functionals have codimension one. □

## 4.2 Double Duals and Dual Maps

**Definition 4.2.1.** *Algebraic Double Dual and Natural Map*

(i) The *algebraic double dual* of $\mathcal{V}$ is $\mathcal{V}'' := (\mathcal{V}')'$.
(ii) The *natural evaluation map* is $\Lambda_{\mathcal{V}} : \mathcal{V} \to \mathcal{V}''$; $\Lambda_{\mathcal{V}}(\mathbf{v})(\varphi) = \varphi(\mathbf{v})$.
(iii) $\mathcal{V}$ is *algebraically reflexive* if $\Lambda_{\mathcal{V}}$ is surjective.

**Theorem 4.2.2.** *Algebraic Reflexivity*
*Natural map $\Lambda_{\mathcal{V}} \in \mathcal{L}(\mathcal{V}, \mathcal{V}'')$ is linear and injective. It is an isomorphism if and only if $\mathcal{V}$ is finite-dimensional.*

*Proof.* For $\mathbf{v} \in \mathcal{V}$, $\Lambda_{\mathcal{V}}(\mathbf{v}) : \varphi \mapsto \varphi(\mathbf{v})$ is linear on $\mathcal{V}'$.
For $a, b \in \mathbb{F}$ and $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, $\Lambda_{\mathcal{V}}(a\mathbf{u} + b\mathbf{v})(\varphi) = a\Lambda_{\mathcal{V}}(\mathbf{u})(\varphi) + b\Lambda_{\mathcal{V}}(\mathbf{v})(\varphi)$, $\varphi \in \mathcal{V}'$, so $\Lambda_{\mathcal{V}} \in \mathcal{L}(\mathcal{V}, \mathcal{V}'')$.
If $\mathbf{v} \in \ker(\Lambda_{\mathcal{V}})$, then $\varphi(\mathbf{v}) = 0$ for all $\varphi \in \mathcal{V}'$, hence $\mathbf{v} = \mathbf{0}$ by Proposition 4.1.7. Thus $\ker(\Lambda_{\mathcal{V}}) = \{\mathbf{0}\}$.
For the isomorphism criterion:

($\Rightarrow$) If $\Lambda_{\mathcal{V}}$ is an isomorphism and $\dim(\mathcal{V}) = \infty$, then Theorem 4.1.6 gives $\dim(\mathcal{V}'') > \dim(\mathcal{V}') > \dim(\mathcal{V})$, contradicting $\mathcal{V} \cong \mathcal{V}''$. Hence $\dim(\mathcal{V}) < \infty$.
($\Leftarrow$) If $\dim(\mathcal{V}) < \infty$, then $\dim(\mathcal{V}'') = \dim(\mathcal{V}') = \dim(\mathcal{V})$ by Theorem 4.1.4. Since $\Lambda_{\mathcal{V}}$ is injective, it is an isomorphism.

*Structural Commentary*: $\Lambda_{\mathcal{V}}$ embeds vectors as evaluation functionals.
*Key Insight*: Finite-dimensional spaces are exactly the algebraically reflexive vector spaces. □

**Definition 4.2.3.** *Dual Map*
If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, the *dual map* of $T$ is the map $T' : \mathcal{W}' \to \mathcal{V}'$ defined by $T'\varphi := \varphi T$ for every $\varphi \in \mathcal{W}'$.
Equivalently, $(T'\varphi)(\mathbf{v}) = \varphi(T\mathbf{v})$.

**Proposition 4.2.4.** *Dual Maps Are Linear*
*If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $T' \in \mathcal{L}(\mathcal{W}', \mathcal{V}')$.*

*Proof.* For $\varphi, \psi \in \mathcal{W}'$, $a \in \mathbb{F}$, and $\mathbf{v} \in \mathcal{V}$, $(T'(a\varphi + \psi))(\mathbf{v}) = (a\varphi + \psi)(T\mathbf{v}) = a\varphi(T\mathbf{v}) + \psi(T\mathbf{v}) = (aT'\varphi + T'\psi)(\mathbf{v})$.
*Structural Commentary*: The dual map pulls functionals back along $T$ and preserves linear structure.
*Key Insight*: Dual maps reverse direction because a functional can test $T\mathbf{v}$ only after $T$ is applied. □

**Theorem 4.2.5.** *Algebra of Dual Maps*
*The dual operation has the following properties:*

*(i) If $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ and $a \in \mathbb{F}$, then $(aS + T)' = aS' + T'$.*
*(ii) If $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ and $S \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $(ST)' = T'S'$.*
*(iii) $(I_{\mathcal{V}})' = I_{\mathcal{V}'}$.*

*Proof.*

(i) For $\varphi \in \mathcal{W}'$ and $\mathbf{v} \in \mathcal{V}$, $((aS + T)'\varphi)(\mathbf{v}) = \varphi(aS\mathbf{v} + T\mathbf{v}) = a(S'\varphi)(\mathbf{v}) + (T'\varphi)(\mathbf{v})$, so $(aS + T)' = aS' + T'$.
(ii) For $\varphi \in \mathcal{W}'$ and $\mathbf{u} \in \mathcal{U}$, $((ST)'\varphi)(\mathbf{u}) = \varphi(S(T\mathbf{u})) = (T'(S'\varphi))(\mathbf{u})$, so $(ST)' = T'S'$.
(iii) For $\varphi \in \mathcal{V}'$ and $\mathbf{v} \in \mathcal{V}$, $(I'_{\mathcal{V}}\varphi)(\mathbf{v}) = \varphi(I_{\mathcal{V}}\mathbf{v}) = \varphi(\mathbf{v})$, so $(I_{\mathcal{V}})' = I_{\mathcal{V}'}$.

*Structural Commentary*: Duality is linear on maps and contravariant under composition.
*Key Insight*: Pullbacks compose in the opposite order. □

**Corollary 4.2.6.** *Double Dual Maps Identify with Original Maps*
*Let $\dim \mathcal{V}, \dim \mathcal{W} < \infty$ and $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.*
*Under $\Lambda_{\mathcal{V}} : \mathcal{V} \cong \mathcal{V}''$ and $\Lambda_{\mathcal{W}} : \mathcal{W} \cong \mathcal{W}''$, $T''\Lambda_{\mathcal{V}} = \Lambda_{\mathcal{W}}T$, so $T''$ identifies with $T$.*

*Proof.* For $\mathbf{v} \in \mathcal{V}$ and $\psi \in \mathcal{W}'$, $\big(T''\Lambda_{\mathcal{V}}(\mathbf{v})\big)(\psi) = \Lambda_{\mathcal{V}}(\mathbf{v})(T'\psi) = (T'\psi)(\mathbf{v}) = \psi(T\mathbf{v}) = \Lambda_{\mathcal{W}}(T\mathbf{v})(\psi)$.
Hence $T''\Lambda_{\mathcal{V}} = \Lambda_{\mathcal{W}}T$.
*Structural Commentary*: Double dualisation commutes with natural evaluation.
*Key Insight*: The double dual square commutes because both sides evaluate every functional the same way. □

**Perspective 4.2.7.** *Why the Double Dual Is Natural*
A finite-dimensional space $\mathcal{V}$ is isomorphic to $\mathcal{V}'$, but such an isomorphism usually requires choosing a basis. The map $\Lambda_{\mathcal{V}} : \mathcal{V} \to \mathcal{V}''$ requires no choice of basis.

**Corollary 4.2.8.** *Dual Map of an Inverse*
*If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ is an isomorphism, then $T'$ is an isomorphism and $(T^{-1})' = (T')^{-1}$*

*Proof.* Since $T^{-1} \in \mathcal{L}(\mathcal{W}, \mathcal{V})$, Theorem 4.2.5 gives $T'(T^{-1})' = (T^{-1}T)' = (I_{\mathcal{V}})' = I_{\mathcal{V}'}$
and $(T^{-1})'T' = (TT^{-1})' = (I_{\mathcal{W}})' = I_{\mathcal{W}'}$. Thus $(T^{-1})'$ is inverse of $T'$, $T'$ is isomorphism and $(T^{-1})' = (T')^{-1}$.
*Structural Commentary*: Duality preserves inverses contravariantly.
*Key Insight*: Invertibility survives duality. □

## 4.3 Annihilators and Transposes

**Definition 4.3.1.** *Annihilator*
If $\mathcal{U} \subseteq \mathcal{V}$, the *annihilator* of $\mathcal{U}$ is $\mathcal{U}^0 := \{\varphi \in \mathcal{V}' : \varphi(\mathbf{u}) = 0 \text{ for every } \mathbf{u} \in \mathcal{U}\}$.

**Proposition 4.3.2.** *Annihilators Are Subspaces*
*If $\mathcal{U} \subseteq \mathcal{V}$, then $\mathcal{U}^0$ is a subspace of $\mathcal{V}'$.*

*Proof.* The zero functional lies in $\mathcal{U}^0$. If $\varphi, \psi \in \mathcal{U}^0$, $a \in \mathbb{F}$, and $\mathbf{u} \in \mathcal{U}$, then $(a\varphi + \psi)(\mathbf{u}) = a\varphi(\mathbf{u}) + \psi(\mathbf{u}) = 0$. Thus $a\varphi + \psi \in \mathcal{U}^0$, and Theorem 1.2.2 shows that $\mathcal{U}^0$ is a subspace of $\mathcal{V}'$.
*Structural Commentary*: The annihilator collects all linear tests that vanish on a chosen set. Closure follows because vanishing is preserved by linear combinations of tests.
*Key Insight*: Annihilators are kernels of evaluation constraints. □

**Theorem 4.3.3.** *Dimension of the Annihilator*
*Let $\mathcal{V}$ be finite-dimensional and let $\mathcal{U}$ be a subspace of $\mathcal{V}$. Then* $\dim(\mathcal{U}) + \dim(\mathcal{U}^0) = \dim(\mathcal{V})$.

*Proof.* Choose a basis $(\mathbf{u}_1, \dots, \mathbf{u}_m)$ of $\mathcal{U}$ and extend it to a basis $(\mathbf{u}_1, \dots, \mathbf{u}_m, \mathbf{w}_1, \dots, \mathbf{w}_k)$ of $\mathcal{V}$.
Let $(\varphi_1, \dots, \varphi_m, \psi_1, \dots, \psi_k)$ be the corresponding dual basis.
By Theorem 4.1.4, every $\varphi \in \mathcal{V}'$ has the expansion $\varphi = \sum_{j=1}^{m} \varphi(\mathbf{u}_j)\varphi_j + \sum_{\ell=1}^{k} \varphi(\mathbf{w}_\ell)\psi_\ell$.
If $\varphi \in \mathcal{U}^0$, then $\varphi(\mathbf{u}_j) = 0$ for every $j$, so $\varphi \in \operatorname{span}(\psi_1, \dots, \psi_k)$.
Conversely, each $\psi_\ell$ vanishes on every $\mathbf{u}_j$, hence on $\mathcal{U} = \operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_m)$. Therefore $\mathcal{U}^0 = \operatorname{span}(\psi_1, \dots, \psi_k)$.
The list $(\psi_1, \dots, \psi_k)$ is linearly independent as a sublist of the dual basis, so $\dim(\mathcal{U}^0) = k = \dim(\mathcal{V}) - \dim(\mathcal{U})$.
*Structural Commentary*: Extending a basis of $\mathcal{U}$ separates directions inside $\mathcal{U}$ from complementary directions. The annihilator is dual to the complementary part.
*Key Insight*: To vanish on $\mathcal{U}$, a functional can only use dual coordinates attached to directions outside $\mathcal{U}$. □

**Theorem 4.3.4.** *Annihilator Lattice Identities*
*Let $\mathcal{V}$ be finite-dimensional, and let $\mathcal{S}, \mathcal{T}$ be subspaces of $\mathcal{V}$.*

*(i)* *If $\mathcal{S} \subseteq \mathcal{T}$, then $\mathcal{T}^0 \subseteq \mathcal{S}^0$.*
*(ii)* $(\mathcal{S} + \mathcal{T})^0 = \mathcal{S}^0 \cap \mathcal{T}^0$.
*(iii)* $(\mathcal{S} \cap \mathcal{T})^0 = \mathcal{S}^0 + \mathcal{T}^0$.

*Proof.*

(i) If $\varphi \in \mathcal{T}^0$ and $\mathbf{s} \in \mathcal{S}$, then $\mathbf{s} \in \mathcal{T}$, so $\varphi(\mathbf{s}) = 0$. Hence $\varphi \in \mathcal{S}^0$.
(ii) For $\varphi \in \mathcal{V}'$, $\varphi \in (\mathcal{S} + \mathcal{T})^0$ if and only if $\varphi(\mathbf{s} + \mathbf{t}) = 0$ for all $\mathbf{s} \in \mathcal{S}$ and $\mathbf{t} \in \mathcal{T}$, if and only if $\varphi \in \mathcal{S}^0 \cap \mathcal{T}^0$.
(iii) Since $\mathcal{S} \cap \mathcal{T} \subseteq \mathcal{S}, \mathcal{T}$, part (i) gives $\mathcal{S}^0 + \mathcal{T}^0 \subseteq (\mathcal{S} \cap \mathcal{T})^0$. By part (ii), Theorem 2.3.1, and Theorem 4.3.3, $\dim(\mathcal{S}^0 + \mathcal{T}^0) = \dim(\mathcal{S}^0) + \dim(\mathcal{T}^0) - \dim(\mathcal{S}^0 \cap \mathcal{T}^0) = \dim(\mathcal{V}) - \dim(\mathcal{S} \cap \mathcal{T}) = \dim\big((\mathcal{S} \cap \mathcal{T})^0\big)$. The inclusion is therefore equality.

*Structural Commentary*: Annihilation is an order-reversing lattice duality.
*Key Insight*: Vanishing on more vectors gives fewer functionals. □

**Theorem 4.3.5.** *Double Annihilator*
*Let $\dim \mathcal{V} < \infty$ and $\emptyset \neq M \subseteq \mathcal{V}$. Under $\Lambda_{\mathcal{V}} : \mathcal{V} \cong \mathcal{V}''$, $M^{00} = \Lambda_{\mathcal{V}}(\operatorname{span} M)$.*
*In particular, if $\mathcal{S} \leq \mathcal{V}$, then $\mathcal{S}^{00} = \Lambda_{\mathcal{V}}(\mathcal{S}) \cong \mathcal{S}$.*

*Proof.* If $\mathbf{v} \in \operatorname{span} M$ and $\varphi \in M^0$, then $\varphi(\mathbf{v}) = 0$, so $\Lambda_{\mathcal{V}}(\mathbf{v})(\varphi) = 0$. Hence $\Lambda_{\mathcal{V}}(\operatorname{span} M) \subseteq M^{00}$.
Conversely, let $\eta \in M^{00}$. By Theorem 4.2.2, $\eta = \Lambda_{\mathcal{V}}(\mathbf{v})$ for a unique $\mathbf{v} \in \mathcal{V}$.
If $\mathbf{v} \notin \operatorname{span} M$, Proposition 4.1.7(iii) gives $\varphi \in \mathcal{V}'$ with $\varphi|_{\operatorname{span} M} = 0$ and $\varphi(\mathbf{v}) = 1$.
Then $\varphi \in M^0$ but $\eta(\varphi) = \Lambda_{\mathcal{V}}(\mathbf{v})(\varphi) = 1$, contradicting $\eta \in M^{00}$. Thus $\mathbf{v} \in \operatorname{span} M$, so $M^{00} \subseteq \Lambda_{\mathcal{V}}(\operatorname{span} M)$.
*Structural Commentary*: Double annihilation returns the span under $\Lambda_{\mathcal{V}}$.
*Key Insight*: Linear functionals separate any vector outside a finite-dimensional span. □

**Theorem 4.3.6.** *Kernel and Range of the Dual Map*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then*

*(i)* $\ker(T') = (\operatorname{range}(T))^0$;
*(ii)* $\dim(\ker(T')) = \dim(\ker(T)) + \dim(\mathcal{W}) - \dim(\mathcal{V})$;
*(iii)* $\operatorname{range}(T') = (\ker(T))^0$;
*(iv)* $\dim(\operatorname{range}(T')) = \dim(\operatorname{range}(T))$.

*Proof.*

(i) For $\varphi \in \mathcal{W}'$, $T'\varphi = 0$ if and only if $\varphi(T\mathbf{v}) = 0$ for every $\mathbf{v} \in \mathcal{V}$, if and only if $\varphi$ vanishes on $\operatorname{range}(T)$.

(ii) By (i), Theorem 4.3.3, and rank-nullity,
$\dim(\ker(T')) = \dim(\mathcal{W}) - \dim(\text{range}(T)) = \dim(\ker(T)) + \dim(\mathcal{W}) - \dim(\mathcal{V})$.
(iii) If $\varphi = T'\psi$ and $\mathbf{v} \in \ker(T)$, then $\varphi(\mathbf{v}) = \psi(T\mathbf{v}) = 0$, so $\text{range}(T') \subseteq (\ker(T))^0$. Also, by rank-nullity for $T'$, (ii), Theorem 4.1.4, and Theorem 4.3.3, $\dim(\text{range}(T')) = \dim(\mathcal{V}) - \dim(\ker(T)) = \dim\left((\ker(T))^0\right)$. Thus $\text{range}(T') = (\ker(T))^0$.
(iv) By (iii), Theorem 4.3.3, and rank-nullity, $\dim(\text{range}(T')) = \dim(\mathcal{V}) - \dim(\ker(T)) = \dim(\text{range}(T))$

*Structural Commentary*: Duality exchanges kernels and ranges through annihilators.
*Key Insight*: A functional dies after pulling back by $T$ exactly when it vanishes on everything $T$ can reach. □

**Corollary 4.3.7.** *Surjectivity and Dual Injectivity*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then $T$ is surjective if and only if $T'$ is injective.*

*Proof.*
($\Rightarrow$) Suppose $T$ is surjective. Then $\text{range}(T) = \mathcal{W}$, so Theorem 4.3.6 gives $\ker(T') = \mathcal{W}^0 = \{0\}$.
($\Leftarrow$) Suppose $T'$ is injective. Then $\ker(T') = \{0\}$, so Theorem 4.3.6 gives $(\text{range}(T))^0 = \{0\}$. By Theorem 4.3.3, $\dim(\text{range}(T)) = \dim(\mathcal{W})$, hence $\text{range}(T) = \mathcal{W}$.

*Structural Commentary*: A map is onto exactly when no nonzero functional can vanish on its range. Dual injectivity detects this absence of invisible codomain directions.
*Key Insight*: Surjectivity of $T$ becomes injectivity of the pullback map $T'$. □

**Corollary 4.3.8.** *Injectivity and Dual Surjectivity*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then $T$ is injective if and only if $T'$ is surjective.*

*Proof.*
($\Rightarrow$) Suppose $T$ is injective. Then $\ker(T) = \{\mathbf{0}\}$, so Theorem 4.3.6 gives $\text{range}(T') = (\ker(T))^0 = \mathcal{V}'$.
($\Leftarrow$) Suppose $T'$ is surjective. Then $\text{range}(T') = \mathcal{V}'$, so Theorem 4.3.6 gives $(\ker(T))^0 = \mathcal{V}'$. By Theorem 4.3.3, this forces $\dim(\ker(T)) = 0$, hence $\ker(T) = \{\mathbf{0}\}$.

*Structural Commentary*: Injectivity says there are no nonzero domain directions killed by $T$. The dual statement says every test on the domain comes by pulling back a test from the codomain.
*Key Insight*: No kernel directions means no obstruction to extending functionals backward through $T$. □

**Corollary 4.3.9.** *Dual of a Quotient as an Annihilator*
*Let $\mathcal{V}$ be finite-dimensional, let $\mathcal{U} \leq \mathcal{V}$, and let $\pi : \mathcal{V} \to \mathcal{V}/\mathcal{U}$ be the quotient map. Then $\ker(\pi') = \{0\}$ and $\text{range}(\pi') = \mathcal{U}^0$. Hence $\pi'$ identifies $(\mathcal{V}/\mathcal{U})'$ with $\mathcal{U}^0$, so $(\mathcal{V}/\mathcal{U})' \cong \mathcal{U}^0$.*

*Proof.* By Theorem 2.2.9, $\mathcal{V}/\mathcal{U}$ is finite-dimensional. Since $\pi$ is surjective and $\ker(\pi) = \mathcal{U}$, Corollary 4.3.7 gives $\ker(\pi') = \{0\}$, and Theorem 4.3.6 gives $\text{range}(\pi') = (\ker(\pi))^0 = \mathcal{U}^0$.
*Structural Commentary*: A functional on the quotient is exactly a functional on $\mathcal{V}$ that cannot see $\mathcal{U}$.
*Key Insight*: Pulling back along the quotient map identifies $(\mathcal{V}/\mathcal{U})'$ with $\mathcal{U}^0$. □

**Theorem 4.3.10.** *Direct Sums and Annihilators*
*Let $\mathcal{V}$ be finite-dimensional, and suppose $\mathcal{V} = \mathcal{S} \oplus \mathcal{T}$.*

*(i) Extension by zero on $\mathcal{S}$ gives an isomorphism $\mathcal{T}' \cong \mathcal{S}^0$.*
*(ii) $\mathcal{V}' = \mathcal{S}^0 \oplus \mathcal{T}^0$.*

*Proof.*
(i) Define $E : \mathcal{T}' \to \mathcal{S}^0$ by $(E\varphi)(\mathbf{s} + \mathbf{t}) = \varphi(\mathbf{t})$. Since $\mathcal{V} = \mathcal{S} \oplus \mathcal{T}$, this is well-defined and linear. Restriction $R : \mathcal{S}^0 \to \mathcal{T}'$, $R\psi = \psi|_{\mathcal{T}}$, satisfies $RE = I_{\mathcal{T}'}$ and $ER = I_{\mathcal{S}^0}$, so $E$ is an isomorphism.
(ii) By Theorem 4.3.4, $\mathcal{S}^0 + \mathcal{T}^0 = (\mathcal{S} \cap \mathcal{T})^0 = \{\mathbf{0}\}^0 = \mathcal{V}'$, and $\mathcal{S}^0 \cap \mathcal{T}^0 = (\mathcal{S} + \mathcal{T})^0 = \mathcal{V}^0 = \{0\}$. Thus $\mathcal{V}' = \mathcal{S}^0 \oplus \mathcal{T}^0$.

*Structural Commentary*: A direct sum decomposes the dual by complementary annihilators.
*Key Insight*: Direct sums split the dual space by annihilators. □

**Theorem 4.3.11.** *Range Membership Dual Alternative*
*Let $\mathcal{V}, \mathcal{W}$ be finite-dimensional, $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, and $\mathbf{w} \in \mathcal{W}$. Exactly one holds:*

*(i) $\mathbf{w} \in \text{range}(T)$;*
*(ii) there exists $\psi \in \mathcal{W}'$ such that $\psi T = 0$ and $\psi(\mathbf{w}) \neq 0$.*

*Proof.*
(i) The alternatives are incompatible: if $\mathbf{w} = T\mathbf{v}$ and $\psi T = 0$, then $\psi(\mathbf{w}) = \psi(T\mathbf{v}) = 0$.

(ii) Suppose $\mathbf{w} \notin \operatorname{range}(T)$. By Proposition 4.1.7(iii), there exists $\psi \in \mathcal{W}'$ such that $\psi|_{\operatorname{range}(T)} = 0$ and $\psi(\mathbf{w}) = 1$.
Then $\psi$ vanishes on $\operatorname{range}(T)$, so $\psi T = 0$, while $\psi(\mathbf{w}) = 1$.

*Structural Commentary*: A missing range vector is detected by a dual test that vanishes on the whole range.
*Key Insight*: Either a vector is reached by the map, or a functional separates it from the range. □




Figure 17: A dual certificate detects a vector outside the range.

**Proposition 4.3.12.** *Standard Coordinate Dual Identification*
*For each positive integer $m$, the map*

$$\Theta : \mathbb{F}^m \to (\mathbb{F}^m)', \qquad \Theta(\mathbf{y})(\mathbf{z}) = \mathbf{y}^t\mathbf{z},$$

*is an isomorphism. Hence every $\psi \in (\mathbb{F}^m)'$ has a unique form $\psi(\mathbf{z}) = \mathbf{y}^t\mathbf{z}$ with $\mathbf{y} \in \mathbb{F}^m$.*

*Proof.* Linearity of $\Theta$ follows from bilinearity of the coordinate dot product. If $\Theta(\mathbf{y}) = 0$, then evaluating at the standard basis vector $\mathbf{e}_j$ gives $y_j = 0$ for every $j$, so $\mathbf{y} = \mathbf{0}$. Thus $\Theta$ is injective. Since $\dim(\mathbb{F}^m) = m = \dim((\mathbb{F}^m)')$ by Theorem 4.1.4, Corollary 1.7.6 gives that $\Theta$ is an isomorphism.
*Structural Commentary*: Coordinate column vectors represent all coordinate linear functionals by row multiplication.
*Key Insight*: The standard dual of $\mathbb{F}^m$ is represented by transposed coordinate vectors. □

**Corollary 4.3.13.** *Matrix Equation Alternative*
*Let $A \in \operatorname{Mat}(m, n, \mathbb{F})$ and $\mathbf{b} \in \mathbb{F}^m$. Exactly one holds:*

*(i) $A\mathbf{x} = \mathbf{b}$ for some $\mathbf{x} \in \mathbb{F}^n$;*
*(ii) $\mathbf{y}^tA = \mathbf{0}^t$ and $\mathbf{y}^t\mathbf{b} \neq 0$ for some $\mathbf{y} \in \mathbb{F}^m$.*

*Proof.*

(i) By Theorem 4.3.11, for $T_A : \mathbb{F}^n \to \mathbb{F}^m$ and $\mathbf{w} = \mathbf{b}$, $\mathbf{b} \in \operatorname{range}(T_A)$ is exactly $A\mathbf{x} = \mathbf{b}$ for some $\mathbf{x} \in \mathbb{F}^n$.
(ii) By Proposition 4.3.12, $\psi \in (\mathbb{F}^m)'$ has the form $\psi(\mathbf{z}) = \mathbf{y}^t\mathbf{z}$ for a unique $\mathbf{y} \in \mathbb{F}^m$.
Thus $\psi T_A = 0$ iff $\mathbf{y}^tA = \mathbf{0}^t$, and $\psi(\mathbf{b}) \neq 0$ iff $\mathbf{y}^t\mathbf{b} \neq 0$.

*Structural Commentary*: Inconsistency is certified by a left-null vector.
*Key Insight*: Left null vectors are the coordinate form of dual separating functionals. □

**Theorem 4.3.14.** *Matrix of the Dual Map Is the Transpose*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, and let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ and $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_m)$ be bases of $\mathcal{V}$ and $\mathcal{W}$. If $\beta'$ and $\gamma'$ are the corresponding dual bases, then $[T']_{\beta' \leftarrow \gamma'} = ([T]_{\gamma \leftarrow \beta})^t$.*

*Proof.* Let $A = [T]_{\gamma \leftarrow \beta}$ and $C = [T']_{\beta' \leftarrow \gamma'}$. Write $\gamma' = (\psi_1, \ldots, \psi_m)$ and $\beta' = (\varphi_1, \ldots, \varphi_n)$.
For $1 \leq j \leq m$, $T'\psi_j = \sum_{r=1}^{n} C_{r,j}\varphi_r$. Evaluating at $\mathbf{v}_k$ gives $C_{k,j} = (T'\psi_j)(\mathbf{v}_k) = \psi_j(T\mathbf{v}_k)$.
Since $T\mathbf{v}_k = \sum_{r=1}^{m} A_{r,k}\mathbf{w}_r$, we get $\psi_j(T\mathbf{v}_k) = A_{j,k}$. Hence $C_{k,j} = A_{j,k}$ for all $j, k$, so $C = A^t$.
*Structural Commentary*: Duality turns the matrix of a map into the matrix of its pullback on coordinate functionals. The row-column indexing is exactly transposition.
*Key Insight*: Coefficients of $T\mathbf{v}_k$ become values of the dual basis functionals on $T\mathbf{v}_k$. □

## 4.4 Multilinear Structure

**Definition 4.4.1.** *Bilinear Forms and Matrices*
A map $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ is *bilinear* if it is linear in each variable separately.
For bases $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ of $\mathcal{V}$ and $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$ of $\mathcal{W}$, the *matrix of $B$ in the bases $\beta, \gamma$* is $[B]_{\beta,\gamma} = [B(\mathbf{v}_i, \mathbf{w}_j)]$. If $\mathbf{x} = [\mathbf{v}]_\beta$ and $\mathbf{y} = [\mathbf{w}]_\gamma$, then $B(\mathbf{v}, \mathbf{w}) = \mathbf{x}^t[B]_{\beta,\gamma}\mathbf{y}$.

**Proposition 4.4.2.** *Bilinear Forms Linearize through Duals*
*Let $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ be bilinear, and define $\Phi_B : \mathcal{V} \to \mathcal{W}'$ where $(\Phi_B\mathbf{v})(\mathbf{w}) = B(\mathbf{v}, \mathbf{w})$, and $\Psi_B : \mathcal{W} \to \mathcal{V}'$ where $(\Psi_B\mathbf{w})(\mathbf{v}) = B(\mathbf{v}, \mathbf{w})$. Then $\Phi_B, \Psi_B$ are linear. For bases $\beta, \gamma$ and $M = [B]_{\beta,\gamma}$, $[\Phi_B]_{\gamma' \leftarrow \beta} = M^t$; and $[\Psi_B]_{\beta' \leftarrow \gamma} = M$.*

*Proof.* For fixed $\mathbf{v}$ and $\mathbf{w}$, bilinearity gives $\Phi_B\mathbf{v} \in \mathcal{W}'$ and $\Psi_B\mathbf{w} \in \mathcal{V}'$. For $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}$, $a \in \mathbb{F}$, and every $\mathbf{w} \in \mathcal{W}$, $(\Phi_B(a\mathbf{v}_1 + \mathbf{v}_2))(\mathbf{w}) = B(a\mathbf{v}_1 + \mathbf{v}_2, \mathbf{w}) = a(\Phi_B\mathbf{v}_1)(\mathbf{w}) + (\Phi_B\mathbf{v}_2)(\mathbf{w})$, so $\Phi_B$ is linear.
For $\mathbf{w}_1, \mathbf{w}_2 \in \mathcal{W}$, $a \in \mathbb{F}$, and every $\mathbf{v} \in \mathcal{V}$, $(\Psi_B(a\mathbf{w}_1 + \mathbf{w}_2))(\mathbf{v}) = B(\mathbf{v}, a\mathbf{w}_1 + \mathbf{w}_2) = a(\Psi_B\mathbf{w}_1)(\mathbf{v}) + (\Psi_B\mathbf{w}_2)(\mathbf{v})$, so $\Psi_B$ is linear. Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ and $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$.
The $j$th $\gamma'$-coordinate of $\Phi_B(\mathbf{v}_i)$ is $(\Phi_B\mathbf{v}_i)(\mathbf{w}_j) = B(\mathbf{v}_i, \mathbf{w}_j) = M_{i,j}$, so column $i$ of $[\Phi_B]_{\gamma' \leftarrow \beta}$ is row $i$ of $M$ transposed, hence $[\Phi_B]_{\gamma' \leftarrow \beta} = M^t$. Likewise, the $i$th $\beta'$-coordinate of $\Psi_B(\mathbf{w}_j)$ is $M_{i,j}$, so $[\Psi_B]_{\beta' \leftarrow \gamma} = M$.
*Structural Commentary*: A bilinear form becomes linear once one input is frozen. Tracking both freezes gives two dual-valued linear maps whose matrices are transposes of one another.
*Key Insight*: Bilinearity is linearity into a dual space. □

**Definition 4.4.3.** *Radicals and Nonsingularity of a Bilinear Pairing*
Let $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ be bilinear.

(i) The *left* and *right radicals* are

$$\operatorname{rad}_L(B) := \{\mathbf{v} \in \mathcal{V} : B(\mathbf{v}, \mathbf{w}) = 0 \text{ for every } \mathbf{w} \in \mathcal{W}\},$$
$$\operatorname{rad}_R(B) := \{\mathbf{w} \in \mathcal{W} : B(\mathbf{v}, \mathbf{w}) = 0 \text{ for every } \mathbf{v} \in \mathcal{V}\}.$$

(ii) For $S \subseteq \mathcal{V}$ and $T \subseteq \mathcal{W}$, the *right* and *left orthogonals* are

$$S^{\perp_R} := \{\mathbf{w} \in \mathcal{W} : B(\mathbf{s}, \mathbf{w}) = 0 \text{ for every } \mathbf{s} \in S\},$$
$${}^{\perp_L}T := \{\mathbf{v} \in \mathcal{V} : B(\mathbf{v}, \mathbf{t}) = 0 \text{ for every } \mathbf{t} \in T\}.$$

(iii) $B$ is *left nondegenerate* if $\operatorname{rad}_L(B) = \{\mathbf{0}\}$ and *right nondegenerate* if $\operatorname{rad}_R(B) = \{\mathbf{0}\}$.
(iv) $B$ is *nonsingular* or *perfect* if $\Phi_B : \mathcal{V} \to \mathcal{W}'$ and $\Psi_B : \mathcal{W} \to \mathcal{V}'$ from Proposition 4.4.2 are isomorphisms.

**Proposition 4.4.4.** *Radicals, Matrices, and Dual Maps*
*Let $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ be bilinear, $\dim \mathcal{V} = m$, $\dim \mathcal{W} = n$, and $M = [B]_{\beta,\gamma}$ in ordered bases $\beta, \gamma$. Then $\{[\mathbf{v}]_\beta : \mathbf{v} \in \operatorname{rad}_L(B)\} = \ker(M^t)$ and $\{[\mathbf{w}]_\gamma : \mathbf{w} \in \operatorname{rad}_R(B)\} = \ker(M)$, therefore $\dim \operatorname{rad}_L(B) = m - \operatorname{rank}(M)$ and $\dim \operatorname{rad}_R(B) = n - \operatorname{rank}(M)$. Moreover, $\ker(\Phi_B) = \operatorname{rad}_L(B)$ and $\ker(\Psi_B) = \operatorname{rad}_R(B)$.*

*Proof.* For $\mathbf{x} = [\mathbf{v}]_\beta$ and $\mathbf{y} = [\mathbf{w}]_\gamma$, $B(\mathbf{v}, \mathbf{w}) = \mathbf{x}^t M \mathbf{y}$.
Thus $\mathbf{v} \in \operatorname{rad}_L(B)$ if and only if $\mathbf{x}^t M \mathbf{y} = 0$ for all $\mathbf{y}$, if and only if $M^t\mathbf{x} = 0$. Similarly, $\mathbf{w} \in \operatorname{rad}_R(B)$ if and only if $M\mathbf{y} = 0$. This proves the coordinate-set identities. Rank-nullity gives the dimension formulas. Finally, $\mathbf{v} \in \ker(\Phi_B) \iff B(\mathbf{v}, \mathbf{w}) = 0$ for every $\mathbf{w} \in \mathcal{W} \iff \mathbf{v} \in \operatorname{rad}_L(B)$; similarly $\ker(\Psi_B) = \operatorname{rad}_R(B)$.
*Structural Commentary*: Radicals are the kernel obstructions in each slot.
*Key Insight*: In coordinates, radicals are matrix nullspaces. □

**Proposition 4.4.5.** *Two-Sided Basis Change for Pairings*
*Let $B : \mathcal{U} \times \mathcal{W} \to \mathbb{F}$ be bilinear, let $\alpha, \alpha'$ be bases of $\mathcal{U}$, and let $\beta, \beta'$ be bases of $\mathcal{W}$. Set $M_{\alpha,\beta} := [B]_{\alpha,\beta}$, $P = C_{\alpha \leftarrow \alpha'}$, and $Q = C_{\beta \leftarrow \beta'}$. Then $M_{\alpha',\beta'} = P^t M_{\alpha,\beta} Q$.*
*For square forms this is congruence; for rectangular pairings it is two-sided equivalence.*

*Proof.* By Definition 4.4.1, $[\mathbf{u}]_{\alpha'}^t M_{\alpha',\beta'} [\mathbf{w}]_{\beta'} = B(\mathbf{u}, \mathbf{w}) = (P[\mathbf{u}]_{\alpha'})^t M_{\alpha,\beta}(Q[\mathbf{w}]_{\beta'}) = [\mathbf{u}]_{\alpha'}^t P^t M_{\alpha,\beta} Q [\mathbf{w}]_{\beta'}$.
Thus $M_{\alpha',\beta'} = P^t M_{\alpha,\beta} Q$.
*Structural Commentary*: Pairing matrices change independently in each input slot.
*Key Insight*: The number of input slots determines how basis changes act on the matrix. □

**Proposition 4.4.6.** *Square Pairing Nonsingularity Criteria*
*Let $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ be bilinear with $\dim \mathcal{V} = \dim \mathcal{W} = n$, and let $M = [B]_{\beta,\gamma}$ in ordered bases $\beta, \gamma$. Then the following are equivalent:*

*(i) $B$ is left nondegenerate;*
*(ii) $B$ is right nondegenerate;*
*(iii) $M$ is invertible;*
*(iv) $B$ is perfect.*

*Proof.*

(i) $\Rightarrow$ (ii). If $\operatorname{rad}_L(B) = 0$, then by Proposition 4.4.4, $\ker(M^t) = 0 \Rightarrow \operatorname{rank}(M) = n \Rightarrow \ker(M) = 0$.
(ii) $\Rightarrow$ (iii). If $\operatorname{rad}_R(B) = 0$, then $\ker(M) = 0$ by Proposition 4.4.4; since $M \in \operatorname{Mat}(n, n, \mathbb{F})$, $M$ is invertible.

(*iii*) $\Rightarrow$ (*iv*). If $M$ is invertible, then $\ker(M) = \ker(M^t) = 0$, so $\mathrm{rad}_L(B) = \mathrm{rad}_R(B) = 0$.
Hence $\ker(\Phi_B) = \ker(\Psi_B) = 0$ by Proposition 4.4.4.
Since $\dim \mathcal{V} = \dim \mathcal{W}' = \dim \mathcal{W} = \dim \mathcal{V}' = n$, $\Phi_B$ and $\Psi_B$ are isomorphisms; hence $B$ is perfect.
(*iv*) $\Rightarrow$ (*i*). If $B$ is perfect, then $\Phi_B$ is an isomorphism. Thus $\mathrm{rad}_L(B) = \ker(\Phi_B) = 0$.

*Structural Commentary*: In square pairings, one-sided nondegeneracy is invertibility.
*Key Insight*: Perfect pairings are exactly invertible matrices viewed without coordinates. □

**Corollary 4.4.7.** *Perfect Pairings Represent Functionals*
*Let $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ be perfect.*
*Then $\forall \psi \in \mathcal{W}'\ \exists! \mathbf{v} \in \mathcal{V} : \psi(\mathbf{w}) = B(\mathbf{v}, \mathbf{w})$, $\mathbf{w} \in \mathcal{W}$; and $\forall \varphi \in \mathcal{V}'\ \exists! \mathbf{w} \in \mathcal{W} : \varphi(\mathbf{v}) = B(\mathbf{v}, \mathbf{w})$, $\mathbf{v} \in \mathcal{V}$.*

*Proof.* Since $B$ is perfect, $\Phi_B : \mathcal{V} \to \mathcal{W}'$ and $\Psi_B : \mathcal{W} \to \mathcal{V}'$ are isomorphisms.
Thus $\psi = \Phi_B(\mathbf{v})$ and $\varphi = \Psi_B(\mathbf{w})$ for unique $\mathbf{v}, \mathbf{w}$, which gives the two formulas.
*Structural Commentary*: Perfect pairings identify each space with the other's dual.
*Key Insight*: Functionals are represented by vectors exactly when the appropriate pairing map is onto. □

**Definition 4.4.8.** *Matrix Congruence*
Matrices $M, N \in \mathrm{Mat}(n, n, \mathbb{F})$ are *congruent* if $\exists\, P \in \mathrm{Mat}(n, n, \mathbb{F})$ invertible such that $N = P^t M P$.

**Proposition 4.4.9.** *Basis Change for a Square Bilinear Form*
*Let $\mathcal{V}$ be finite-dimensional, let $B : \mathcal{V} \times \mathcal{V} \to \mathbb{F}$ be bilinear, and let $\beta, \gamma$ be ordered bases of $\mathcal{V}$.*
*Set $M_\beta := [B]_{\beta,\beta}$; $M_\gamma := [B]_{\gamma,\gamma}$, and $C := C_{\beta \leftarrow \gamma}$.*
*Then $M_\gamma = C^t M_\beta C$. Hence $M_\beta$ and $M_\gamma$ are congruent.*

*Proof.* For $\mathbf{x}, \mathbf{y} \in \mathcal{V}$, $[\mathbf{x}]_\beta = C[\mathbf{x}]_\gamma$ and $[\mathbf{y}]_\beta = C[\mathbf{y}]_\gamma$. Therefore $B(\mathbf{x}, \mathbf{y}) = [\mathbf{x}]_\beta^t M_\beta [\mathbf{y}]_\beta = [\mathbf{x}]_\gamma^t C^t M_\beta C [\mathbf{y}]_\gamma$.
By uniqueness of the representing matrix in the basis $\gamma$, $M_\gamma = C^t M_\beta C$.
Since $C$ is invertible by Theorem 3.5.2, Definition 4.4.8 gives congruence.
*Structural Commentary*: Operators change by similarity; square bilinear forms change by congruence because both input slots are rewritten.
*Key Insight*: A basis change acts on a square bilinear form from both sides. □




Figure 18: Changing both slots gives $M_\gamma = C_{\beta \leftarrow \gamma}^t M_\beta C_{\beta \leftarrow \gamma}$.

**Perspective 4.4.10.** *Bilinear Forms, Inner Products, and Tensor Scope*
(i) Bilinear forms: $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$; inner products add $\mathbf{v} \neq \mathbf{0} \Rightarrow B(\mathbf{v}, \mathbf{v}) > 0$ and, over $\mathbb{C}$, $B(\mathbf{v}, \mathbf{w}) = \overline{B(\mathbf{w}, \mathbf{v})}$.
(ii) $\Phi_B : \mathcal{V} \to \mathcal{W}'$ and $\Psi_B : \mathcal{W} \to \mathcal{V}'$ encode $B$ in dual spaces, with $[\Phi_B]_{\gamma' \leftarrow \beta} = [B]_{\beta,\gamma}^t$ and $[\Psi_B]_{\beta' \leftarrow \gamma} = [B]_{\beta,\gamma}$.
(iii) $\mathcal{V} \oplus \mathcal{W}$ and $\mathcal{V} \times \mathcal{W}$ use single-coordinate slots; the finite tensor scaffold uses paired slots $(i, j)$.

# Part III
# Linear Maps and Matrix Representations

## 1 Linear Maps and Operations

### 1.1 Further Linear Map Operations and Examples

## 2 Kernels, Images, Rank, and Nullity

### 2.1 Shift Operators and Kernel-Range Examples

**Proposition 2.1.1.** *Shift Operators on Finitely Nonzero Sequences*
*Define* $R, S : \mathbb{F}^{(\mathbb{N})} \to \mathbb{F}^{(\mathbb{N})}$ *by* $R(x_1, x_2, \ldots) = (0, x_1, x_2, \ldots)$, $S(x_1, x_2, \ldots) = (x_2, x_3, \ldots)$. *Then* $R, S \in \mathcal{L}(\mathbb{F}^{(\mathbb{N})})$.
*Also,* $\ker(R) = \{\mathbf{0}\}$, $\text{range}(R) = \{(x_1, x_2, \ldots) \in \mathbb{F}^{(\mathbb{N})} : x_1 = 0\}$, $\ker(S) = \text{span}((1, 0, 0, \ldots))$, $\text{range}(S) = \mathbb{F}^{(\mathbb{N})}$.

*Proof.* Both $R$ and $S$ preserve finite support, so they map $\mathbb{F}^{(\mathbb{N})}$ into itself.
Since both maps act componentwise by shifting entries, $R(a\mathbf{x}+\mathbf{y}) = aR\mathbf{x}+R\mathbf{y}$ and $S(a\mathbf{x}+\mathbf{y}) = aS\mathbf{x}+S\mathbf{y}$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{F}^{(\mathbb{N})}$ and $a \in \mathbb{F}$. Thus $R$ and $S$ are linear.
The map $R$ has zero kernel as $R\mathbf{x} = \mathbf{0}$ forces every entry of $\mathbf{x}$ to be zero, and its range is exactly the set of finitely nonzero sequences with first entry zero. For $S$, $S\mathbf{x} = \mathbf{0}$ forces $x_2 = x_3 = \cdots = 0$, so $\ker(S) = \text{span}((1, 0, 0, \ldots))$.
Finally, for any $(y_1, y_2, \ldots) \in \mathbb{F}^{(\mathbb{N})}$, we have $S(0, y_1, y_2, \ldots) = (y_1, y_2, \ldots)$, so $S$ is surjective.
*Structural Commentary*: Shift operators expose behavior that finite-dimensional maps cannot have.
*Key Insight*: The right shift is injective but not surjective, while the left shift is surjective but not injective. □

**Example 2.1.2.** *Shift Operators Show Noncommutativity*
For the shifts $R, S \in \mathcal{L}(\mathbb{F}^{(\mathbb{N})})$ from Proposition 2.1.1, $SR = I_{\mathbb{F}^{(\mathbb{N})}}$ and $RS(x_1, x_2, \ldots) = (0, x_2, x_3, \ldots)$.
Hence $RS \neq SR$, for instance on $(1, 0, 0, \ldots)$. Therefore $\mathcal{L}(\mathbb{F}^{(\mathbb{N})})$ is not commutative under composition.

### 2.2 Further Rank-Nullity and Invertibility Results

### 2.3 Further Quotients and Isomorphism Theorems

### 2.4 Further Finite-Dimensional Splitting and Isomorphism Tests

## 3 Matrix Representations

### 3.1 Further Matrix Algebra and Linear Systems

### 3.2 Further Coordinates and Matrix Representations

### 3.3 Further Matrix-Induced Maps and Rank

### 3.4 Further Invertibility and the Matrix-Map Dictionary

## 4 Row Reduction and Elementary Matrices

### 4.1 Row Reduction and Elementary Matrices

**Definition 4.1.1.** *Elementary Row Operations and Elementary Matrices*
For row symbols $R_i$, the *elementary row operations* are:

(i) *Row scaling*: $R_i \leftarrow cR_i$, $c \in \mathbb{F}^\times$;
(ii) *Row swap*: $R_i \leftrightarrow R_j$, $i \neq j$;
(iii) *Row replacement*: $R_i \leftarrow R_i + cR_j$, $i \neq j$, $c \in \mathbb{F}$.

An *elementary matrix* is $E = \epsilon(I_m)$, where $\epsilon$ is one elementary row operation.

**Proposition 4.1.2.** *Elementary Matrices Perform Row Operations*
*Let* $A \in \text{Mat}(m, n, \mathbb{F})$.

*(i) If* $\epsilon$ *is an elementary row operation and* $E_\epsilon = \epsilon(I_m)$, *then* $E_\epsilon A = \epsilon(A)$.
*(ii)* $E_\epsilon^{-1} = E_{\epsilon^{-1}}$; *hence every elementary matrix is invertible with elementary inverse.*
*(iii) If* $F = E_\epsilon^t \in \text{Mat}(n, n, \mathbb{F})$, *then* $AF = (\epsilon(A^t))^t$.
*Hence right multiplication performs the corresponding column operation.*

*Proof.*

(i) For each row index $r$, $\text{row}_r(E_\epsilon A) = \text{row}_r(E_\epsilon)A$.
So the rows of $E_\epsilon A$ are exactly the $\epsilon$-linear combinations of the rows of $A$.
(ii) Since $\epsilon^{-1}$ is elementary, $E_{\epsilon^{-1}}E_\epsilon = E_{\epsilon^{-1}}\epsilon(I_m) = \epsilon^{-1}(\epsilon(I_m)) = I_m$, and similarly $E_\epsilon E_{\epsilon^{-1}} = I_m$.
(iii) By (i) and Proposition 3.1.13, $(AF)^t = E_\epsilon A^t = \epsilon(A^t)$, so $AF = (\epsilon(A^t))^t$.

*Structural Commentary*: Elementary matrices encode row and column operations by multiplication.
*Key Insight*: Row operations are left multiplication; column operations are right multiplication. □

**Definition 4.1.3.** *Row Equivalence and Reduced Row Echelon Form*
For $A, B \in \text{Mat}(m, n, \mathbb{F})$, write $A \sim_{\text{row}} B$ if $B = E_k \cdots E_1 A$ for elementary $E_i \in \text{Mat}(m, m, \mathbb{F})$ and some $k \geq 0$.
A matrix $R = [r_{ij}]$ is in *reduced row echelon form* if its nonzero rows have pivot columns $p_1, \ldots, p_s$ such that:

(i) rows $1, \ldots, s$ are nonzero and rows $s+1, \ldots, m$ are zero;
(ii) $r_{i,p_i} = 1$ and $r_{i,j} = 0$ for $j < p_i$;
(iii) $p_1 < \cdots < p_s$;
(iv) $r_{\ell,p_i} = 0$ for all $\ell \neq i$.

**Theorem 4.1.4.** *Reduced Row Echelon Form*
*For every $A \in \text{Mat}(m, n, \mathbb{F})$, there is a unique RREF matrix $\text{rref}(A)$ with $A \sim_{\text{row}} \text{rref}(A)$.*
*If $R = \text{rref}(A)$, then $R = E_k \cdots E_1 A$ for elementary $E_i \in \text{Mat}(m, m, \mathbb{F})$.*

*Proof.* Gaussian elimination successively chooses a pivot column, swaps/scales to a pivot 1, clears the pivot column, and continues on the remaining submatrix. Proposition 4.1.2 gives $R = E_k \cdots E_1 A$ with $R$ in RREF.
For uniqueness, row operations preserve $\ker(A) = \{\mathbf{x} : A\mathbf{x} = \mathbf{0}\}$. For an RREF matrix, the pivot set and pivot-variable formulas are determined by this kernel.
Hence two row-equivalent RREF matrices have the same pivot columns and rows.
*Structural Commentary*: RREF is the canonical row-operation representative of a matrix.
*Key Insight*: Elimination gives existence; the solution space pins down the reduced representative. □

**Lemma 4.1.5.** *Invertible Matrices Row-Reduce to the Identity*
*If $P \in \text{Mat}(m, m, \mathbb{F})$ is invertible, then $\text{rref}(P) = I_m$. Hence there are elementary matrices $E_1, \ldots, E_k \in \text{Mat}(m, m, \mathbb{F})$ such that $E_k \cdots E_1 P = I_m$.*

*Proof.* Let $R = \text{rref}(P)$. By Theorem 4.1.4, $R = E_k \cdots E_1 P$ for elementary matrices $E_i$. Since each $E_i$ is invertible by Proposition 4.1.2, left multiplication by $E_k \cdots E_1$ preserves the kernel, so $\ker(R) = \ker(P)$.
Because $P$ is invertible, $P\mathbf{x} = \mathbf{0}$ implies $\mathbf{x} = P^{-1}P\mathbf{x} = \mathbf{0}$; hence $\ker(R) = \{\mathbf{0}\}$.
Suppose the nonzero rows of $R$ have pivot columns $p_1 < \cdots < p_s$. If $s < m$, choose a non-pivot column $j$. Set the non-pivot variable $x_j = 1$, set all other non-pivot variables equal to 0, and define the pivot variables recursively from $i = s$ down to 1 so that the $i$th row equation is satisfied. This gives a nonzero $\mathbf{x}$ with $R\mathbf{x} = \mathbf{0}$, contradicting $\ker(R) = \{\mathbf{0}\}$. Thus $s = m$.
Since $p_1 < \cdots < p_m$ are $m$ columns in an $m \times m$ matrix, $p_i = i$ for every $i$. The RREF conditions then force the $i$th pivot column to be the $i$th standard coordinate column, so $R = I_m$. □

**Corollary 4.1.6.** *Row Equivalence Characterization*
*For $A, B \in \text{Mat}(m, n, \mathbb{F})$, the following are equivalent:*

*(i)* $A \sim_{\text{row}} B$.
*(ii)* $\text{rref}(A) = \text{rref}(B)$.
*(iii)* $\exists\, P \in \text{Mat}(m, m, \mathbb{F})^\times$ *such that* $B = PA$.

*Proof.*

*(i)* $\Rightarrow$ *(ii)* Row-equivalent matrices have the same unique RREF by Theorem 4.1.4.
*(ii)* $\Rightarrow$ *(iii)* If $R = \text{rref}(A) = \text{rref}(B)$, then $R = E_k \cdots E_1 A = F_\ell \cdots F_1 B$.
Since each $F_j$ is invertible, $B = (F_\ell \cdots F_1)^{-1} E_k \cdots E_1 A$.
*(iii)* $\Rightarrow$ *(i)* If $B = PA$ and $P \in \text{Mat}(m, m, \mathbb{F})^\times$, Lemma 4.1.5 gives elementary matrices with $E_k \cdots E_1 P = I_m$.
Hence $P = E_1^{-1} \cdots E_k^{-1}$, so $B$ is obtained from $A$ by elementary row operations.

*Structural Commentary*: Row equivalence is invertible left multiplication.
*Key Insight*: Products of elementary matrices are the coordinate record of row reduction. □

**Method 4.1.7.** *Augmented Inverse Computation Preview*
For $A \in \text{Mat}(n, n, \mathbb{F})$:

(i) Form $[\, A \; I_n \,]$.
(ii) Apply Gauss–Jordan reduction to the left block.

(iii) If $[\,A\ I_n\,] \sim [\,I_n\ B\,]$, then $B = A^{-1}$.
(iv) Otherwise $A$ is not invertible, by Corollary 3.6.3.

# 5 Change of Basis

## 5.1 Further Change of Basis and Coordinate Transformations

# 6 Matrix Equivalence and Similarity

## 6.1 Matrix Equivalence and Rank Normal Forms

**Definition 6.1.1.** *Matrix Equivalence*
For $A, B \in \mathrm{Mat}(m,n,\mathbb{F})$, write $A \sim_{\mathrm{eq}} B$ if $\exists\ P \in \mathrm{Mat}(m,m,\mathbb{F})^{\times}$ and $Q \in \mathrm{Mat}(n,n,\mathbb{F})^{\times}$ s.t. $B = PAQ$.

**Proposition 6.1.2.** *Matrix Equivalence and Elementary Operations*
*The relation $\sim_{\mathrm{eq}}$ is an equivalence relation on $\mathrm{Mat}(m,n,\mathbb{F})$. Moreover, $A \sim_{\mathrm{eq}} B$ if and only if $B$ can be obtained from $A$ by finitely many elementary row and column operations.*

*Proof.* Reflexivity follows from $A = I_m A I_n$. If $B = PAQ$ with $P, Q$ invertible, then $A = P^{-1}BQ^{-1}$, so symmetry holds. If $B = P_1AQ_1$ and $C = P_2BQ_2$, then $C = (P_2P_1)A(Q_1Q_2)$, and products of invertible matrices are invertible by Corollary 3.1.14; hence transitivity holds.
Each elementary row operation is left multiplication by an invertible elementary matrix by Proposition 4.1.2. Each elementary column operation is right multiplication by the transpose of an invertible elementary matrix, again by Proposition 4.1.2 and Corollary 3.1.14. Thus any finite sequence of elementary row and column operations has the form $B = PAQ$ with $P, Q$ invertible, so $A \sim_{\mathrm{eq}} B$.
Conversely, suppose $A \sim_{\mathrm{eq}} B$, so $B = PAQ$ with $P, Q$ invertible. Lemma 4.1.5 gives products of elementary matrices expressing $P$ and $Q$ as products of elementary factors. Left multiplication by the factors of $P$ is a sequence of elementary row operations, and right multiplication by the factors of $Q$ is a sequence of elementary column operations. Hence $B$ can be obtained from $A$ by finitely many elementary row and column operations. □

**Theorem 6.1.3.** *Equivalent Matrices Represent the Same Linear Map*
*Let $\dim \mathcal{V} = n$, $\dim \mathcal{W} = m$, and $A, B \in \mathrm{Mat}(m,n,\mathbb{F})$.*
*Then $A \sim_{\mathrm{eq}} B \iff \exists\, T \in \mathcal{L}(\mathcal{V},\mathcal{W})$ with $A = [T]_{\gamma\leftarrow\beta}$, $B = [T]_{\gamma'\leftarrow\beta'}$ for some bases $\beta, \beta'$ of $\mathcal{V}$ and $\gamma, \gamma'$ of $\mathcal{W}$.*

*Proof.*
(⇒) Let $B = PAQ$ with $P, Q$ invertible. Choose bases $\beta, \gamma$ and define $T \in \mathcal{L}(\mathcal{V},\mathcal{W})$ by $[T]_{\gamma\leftarrow\beta} = A$.
By Theorem 3.5.2, choose basis $\gamma'$ with $P = C_{\gamma'\leftarrow\gamma}$, and basis $\beta'$ with $Q^{-1} = C_{\beta'\leftarrow\beta}$. Then $Q = C_{\beta\leftarrow\beta'}$.
Then Theorem 3.5.3 gives $[T]_{\gamma'\leftarrow\beta'} = PAQ = B$.
(⇐) If $A = [T]_{\gamma\leftarrow\beta}$ and $B = [T]_{\gamma'\leftarrow\beta'}$, then $B = C_{\gamma'\leftarrow\gamma} A C_{\beta\leftarrow\beta'}$, so $A \sim_{\mathrm{eq}} B$.

*Structural Commentary*: Equivalence is two-sided basis change.
*Key Insight*: Same map; different coordinates. □

**Theorem 6.1.4.** *Rank Classifies Matrix Equivalence*
*For $0 \le r \le \min(m,n)$, set $J_r^{m,n} := \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} \in \mathrm{Mat}(m,n,\mathbb{F})$.*
*For $A, B \in \mathrm{Mat}(m,n,\mathbb{F})$, $A \sim_{\mathrm{eq}} J_{\mathrm{rank}(A)}^{m,n}$ and $A \sim_{\mathrm{eq}} B \iff \mathrm{rank}(A) = \mathrm{rank}(B)$.*

*Proof.*
(⇒) If $B = PAQ$ with $P, Q$ invertible, then $\mathrm{rank}(B) = \mathrm{rank}(A)$ by Theorem 3.6.5.
(⇐) For any $X$ with $\mathrm{rank}(X) = r$, let $T_X : \mathbb{F}^n \to \mathbb{F}^m$ be $T_X(\mathbf{x}) = X\mathbf{x}$.
Choose $\mathcal{M}_X$ with $\mathbb{F}^n = \ker(T_X) \oplus \mathcal{M}_X$; then $T_X|_{\mathcal{M}_X} : \mathcal{M}_X \to \mathrm{range}(T_X)$ is an isomorphism.
Choose a basis $(\mathbf{u}_1, \dots, \mathbf{u}_r)$ of $\mathcal{M}_X$, extend it by a basis of $\ker(T_X)$ to $\beta_X$, and choose $\gamma_X$ beginning with $T_X\mathbf{u}_1, \dots, T_X\mathbf{u}_r$. Then $[T_X]_{\gamma_X\leftarrow\beta_X} = J_r^{m,n}$, so $X \sim_{\mathrm{eq}} J_r^{m,n}$.
If $\mathrm{rank}(A) = \mathrm{rank}(B) = r$, then $A \sim_{\mathrm{eq}} J_r^{m,n}$ and $B \sim_{\mathrm{eq}} J_r^{m,n}$. By Proposition 6.1.2, symmetry gives $J_r^{m,n} \sim_{\mathrm{eq}} B$, and transitivity gives $A \sim_{\mathrm{eq}} B$.

*Structural Commentary*: Rank is the invariant of two-sided equivalence.
*Key Insight*: Adapt bases to kernel and range. □

**Corollary 6.1.5.** *Rank Normal Form for Linear Maps*
*Let $T \in \mathcal{L}(\mathcal{V},\mathcal{W})$, $\dim \mathcal{V} = n$, $\dim \mathcal{W} = m$, and $\mathrm{rank}(T) = r$. Then $\exists$ bases $\beta, \gamma$ with $[T]_{\gamma\leftarrow\beta} = J_r^{m,n}$.*

*Proof.* For initial bases $\beta_0, \gamma_0$, let $A = [T]_{\gamma_0 \leftarrow \beta_0}$.
By Theorem 6.1.4, $A \sim_{\text{eq}} J_r^{m,n}$; Theorem 6.1.3 gives bases $\beta, \gamma$ with $[T]_{\gamma \leftarrow \beta} = J_r^{m,n}$.
*Structural Commentary*: Rank gives the two-basis normal form.
*Key Insight*: Rank normalizes the map. □

**Corollary 6.1.6.** *Linear Maps Classified by Rank*
*For $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, write $S \sim_{\text{bc}} T$ if $\exists$ bases $\beta_S, \beta_T$ of $\mathcal{V}$ and $\gamma_S, \gamma_T$ of $\mathcal{W}$ such that $[S]_{\gamma_S \leftarrow \beta_S} = [T]_{\gamma_T \leftarrow \beta_T}$. Then $S \sim_{\text{bc}} T \iff \operatorname{rank}(S) = \operatorname{rank}(T)$*

*Proof.*

($\Rightarrow$) If $[S]_{\gamma_S \leftarrow \beta_S} = [T]_{\gamma_T \leftarrow \beta_T}$, then the two maps have the same matrix rank, hence the same rank.
($\Leftarrow$) If $\operatorname{rank}(S) = \operatorname{rank}(T) = r$, Corollary 6.1.5 gives bases with $[S]_{\gamma_S \leftarrow \beta_S} = J_r^{m,n} = [T]_{\gamma_T \leftarrow \beta_T}$.

*Structural Commentary*: Rank is the invariant of basis-change equivalence.
*Key Insight*: Same rank means same normal form. □

## 6.2 Similarity and Operator Coordinates

**Theorem 6.2.1.** *Change of Basis Formula*
*Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $\beta$ be a basis of $\mathcal{V}$.*

*(i) If $\gamma$ is a basis of $\mathcal{V}$ and $S = C_{\gamma \leftarrow \beta}$, then $[T]_{\gamma \leftarrow \gamma} = S[T]_{\beta \leftarrow \beta} S^{-1}$.*
*(ii) If $S \in \operatorname{Mat}(n, n, \mathbb{F})$ is invertible, then there is a basis $\gamma$ of $\mathcal{V}$ such that $[T]_{\gamma \leftarrow \gamma} = S[T]_{\beta \leftarrow \beta} S^{-1}$.*

*Proof.*

(i) Suppose $\gamma$ is a basis and $S = C_{\gamma \leftarrow \beta}$. By Theorem 3.5.2, $S^{-1} = C_{\beta \leftarrow \gamma}$.
For every $\mathbf{v} \in \mathcal{V}$, $[T\mathbf{v}]_\gamma = S[T\mathbf{v}]_\beta = S[T]_{\beta \leftarrow \beta}[\mathbf{v}]_\beta = S[T]_{\beta \leftarrow \beta} S^{-1}[\mathbf{v}]_\gamma$.
Since $[\cdot]_\gamma$ is onto $\mathbb{F}^n$, the representing matrices are equal.
(ii) If $S$ is invertible, Theorem 3.5.2 supplies a basis $\gamma$ with $S = C_{\gamma \leftarrow \beta}$, and part (i) applies.

*Structural Commentary*: Changing basis conjugates operator matrices.
*Key Insight*: Similarity is a coordinate rewrite. □

**Definition 6.2.2.** *Similarity*
Matrices $A, B \in \operatorname{Mat}(n, n, \mathbb{F})$ are similar if there exists an invertible matrix $P$ such that $B = PAP^{-1}$.

**Proposition 6.2.3.** *Similarity Is an Equivalence Relation*
*Similarity is an equivalence relation on $\operatorname{Mat}(n, n, \mathbb{F})$.*

*Proof.* For any $A$, $A = I_n A I_n^{-1}$, so similarity is reflexive. If $B = PAP^{-1}$, then $A = P^{-1}BP$, so it is symmetric.
If $B = PAP^{-1}$ and $C = QBQ^{-1}$, then $C = Q(PAP^{-1})Q^{-1} = (QP)A(QP)^{-1}$ so it is transitive.
*Structural Commentary*: Similarity groups matrices by change of basis.
*Key Insight*: Identity, inverse, and composition give the three equivalence properties. □

**Proposition 6.2.4.** *Upper Block-Triangular Similarity Shift*
*Let $B \in \operatorname{Mat}(m, m, \mathbb{F})$, let $D \in \operatorname{Mat}(n, n, \mathbb{F})$, and let $C, X \in \operatorname{Mat}(m, n, \mathbb{F})$.*
*Then $\begin{bmatrix} B & C \\ 0 & D \end{bmatrix}$ is similar to $\begin{bmatrix} B & C + XD - BX \\ 0 & D \end{bmatrix}$.*

*Proof.* The matrices $\begin{bmatrix} I_m & X \\ 0 & I_n \end{bmatrix}$ and $\begin{bmatrix} I_m & -X \\ 0 & I_n \end{bmatrix}$ are inverses in $\operatorname{Mat}(m+n, m+n, \mathbb{F})$, by block multiplication,

$$\begin{bmatrix} I_m & X \\ 0 & I_n \end{bmatrix} \begin{bmatrix} B & C \\ 0 & D \end{bmatrix} \begin{bmatrix} I_m & -X \\ 0 & I_n \end{bmatrix} = \begin{bmatrix} B & C + XD - BX \\ 0 & D \end{bmatrix}.$$

*Structural Commentary*: Unit upper block-triangular conjugation changes only the coupling block.
*Key Insight*: Similarity can adjust the connecting block while preserving the diagonal blocks. □

**Remark 6.2.5.** *Similarity Convention*
The forms $PAP^{-1}$ and $M^{-1}AM$ are the same convention with $P = M^{-1}$. In Theorem 6.2.1, this means choosing opposite coordinate-change directions.




Figure 19: One operator in two bases: $PA = BP$, so $B = PAP^{-1}$.

**Corollary 6.2.6.** *Similarity as One Operator in Two Bases*
*For $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$, the following are equivalent:*

*(i) $A$ and $B$ are similar.*
*(ii) There are an $n$-dim vector space $\mathcal{V}$, bases $\beta, \gamma$ of $\mathcal{V}$, and $T \in \mathcal{L}(\mathcal{V})$ such that $A = [T]_{\beta\leftarrow\beta}$ and $B = [T]_{\gamma\leftarrow\gamma}$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*): Suppose $B = PAP^{-1}$ with $P$ invertible.
Let $\mathcal{V} = \mathbb{F}^n$, let $\beta$ be the standard basis, and let $T_A(\mathbf{x}) = A\mathbf{x}$. By Theorem 3.5.2, choose $\gamma$ with $P = C_{\gamma\leftarrow\beta}$.
Then $[T_A]_{\beta\leftarrow\beta} = A$, and Theorem 6.2.1 gives $[T_A]_{\gamma\leftarrow\gamma} = B$.
(*ii*) $\Rightarrow$ (*i*): If $A = [T]_{\beta\leftarrow\beta}$ and $B = [T]_{\gamma\leftarrow\gamma}$, then $B = C_{\gamma\leftarrow\beta}AC_{\beta\leftarrow\gamma}$ with $C_{\gamma\leftarrow\beta}$ invertible.

*Structural Commentary*: Similar matrices are one operator in different bases.
*Key Insight*: Conjugation records a change of basis. □

**Perspective 6.2.7.** *Similarity as Same Operator, New Coordinates*
Similarity does not mean two operators are merely related; it means they are the same operator written in two different bases. This is the structural content behind change of basis.

**Definition 6.2.8.** *Similarity of Operators*
For $T \in \mathcal{L}(\mathcal{V})$ and $S \in \mathcal{L}(\mathcal{W})$, $T \sim S \iff \exists\, \Phi : \mathcal{V} \xrightarrow{\sim} \mathcal{W}$ such that $S = \Phi T \Phi^{-1}$.
If $\mathcal{V} = \mathcal{W}$, then $\Phi \in \mathrm{Aut}(\mathcal{V})$ and similarity is conjugacy in $\mathcal{L}(\mathcal{V})$.

**Proposition 6.2.9.** *Operator Similarity and Common Matrix Representatives*
*Let $\dim \mathcal{V} = \dim \mathcal{W} = n$, $T \in \mathcal{L}(\mathcal{V})$, and $S \in \mathcal{L}(\mathcal{W})$.*
*Then $T \sim S \iff \exists\, A \in \mathrm{Mat}(n, n, \mathbb{F})$ with $A = [T]_{\beta\leftarrow\beta} = [S]_{\gamma\leftarrow\gamma}$ for some bases $\beta$ of $\mathcal{V}$ and $\gamma$ of $\mathcal{W}$.*

*Proof.*

($\Rightarrow$) Let $S = \Phi T \Phi^{-1}$ and $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$. Put $\gamma = (\Phi\mathbf{v}_1, \ldots, \Phi\mathbf{v}_n)$.
Then $\gamma$ is a basis and, for all $j$, $S(\Phi\mathbf{v}_j) = \Phi T\mathbf{v}_j$, so $[S]_{\gamma\leftarrow\gamma} = [T]_{\beta\leftarrow\beta}$.
($\Leftarrow$) Suppose $[T]_{\beta\leftarrow\beta} = [S]_{\gamma\leftarrow\gamma}$. Define $\Phi : \mathcal{V} \to \mathcal{W}$ by $\Phi(\beta_j) = \gamma_j$.
Then $\Phi$ is an isomorphism and $\Phi T\beta_j = S\Phi\beta_j$ for every $j$, hence $\Phi T = S\Phi$. Thus $S = \Phi T \Phi^{-1}$.

*Structural Commentary*: Similarity is equality of matrices after identifying bases.
*Key Insight*: A common matrix is an operator-level conjugacy written in bases. □

**Theorem 6.2.10.** *Similarity and Scalar Shifts*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$.*

*(i) If $A$ and $B$ are similar, then $A - \lambda I_n$ and $B - \lambda I_n$ are similar for every $\lambda \in \mathbb{F}$.*
*(ii) If $A - \lambda I_n$ and $B - \lambda I_n$ are similar for some $\lambda \in \mathbb{F}$, then $A$ and $B$ are similar.*

*Proof.*

(i) If $B = PAP^{-1}$ with $P$ invertible, then $B - \lambda I_n = PAP^{-1} - \lambda PP^{-1} = P(A - \lambda I_n)P^{-1}$.
(ii) If $B - \lambda I_n = P(A - \lambda I_n)P^{-1}$ with $P$ invertible, then adding $\lambda I_n = \lambda PP^{-1}$ gives $B = PAP^{-1}$.

*Structural Commentary*: Scalar shifts commute with change of basis.
*Key Insight*: The identity matrix is basis-independent. □

# Part IV
# Geometric and Spectral Linear Algebra

## 1 Inner Product Geometry

### 1.1 Inner Products and Norms

Throughout this subsection, $\mathbb{F}$ is $\mathbb{R}$ or $\mathbb{C}$. Inner products turn linear spaces into geometric spaces with norms, orthogonality, orthonormal bases, and Gram matrices.

**Definition 1.1.1.** *Preliminary Terminology*

(i) For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, the *dot product* is $\mathbf{x} \cdot \mathbf{y} := x_1 y_1 + \cdots + x_n y_n$.
(ii) For $z = a + bi \in \mathbb{C}$, the *complex conjugate* is $\overline{z} := a - bi$, and the *modulus* is $|z| := \sqrt{a^2 + b^2}$.
(iii) A map $f : \mathcal{V} \to \mathbb{F}$ is *conjugate linear* if $f(\mathbf{u} + \mathbf{v}) = f(\mathbf{u}) + f(\mathbf{v})$ and $f(\lambda \mathbf{v}) = \overline{\lambda} f(\mathbf{v})$.
(iv) A function $B : \mathcal{V} \times \mathcal{V} \to \mathbb{F}$ is *sesquilinear* if it is linear in one slot and conjugate linear in the other.

**Motivation 1.1.2.** *From Dot Products to Inner Products*
For $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{R}^n$, the Euclidean length is $\|\mathbf{x}\| = \sqrt{x_1^2 + \cdots + x_n^2}$.
The expression under the square root is the dot product of $\mathbf{x}$ with itself. Over $\mathbb{C}$, conjugates are needed to keep squared length real and nonnegative: $\|\mathbf{z}\|^2 = |z_1|^2 + \cdots + |z_n|^2 = z_1\overline{z_1} + \cdots + z_n\overline{z_n}$.
Thus the standard inner product on $\mathbb{F}^n$ is $\langle \mathbf{x}, \mathbf{y} \rangle := x_1\overline{y_1} + \cdots + x_n\overline{y_n}$.

**Definition 1.1.3.** *Inner Product*
Let $\mathcal{V}$ be a vector space over $\mathbb{F}$. An *inner product* on $\mathcal{V}$ is a function that assigns to each pair $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ a scalar $\langle \mathbf{u}, \mathbf{v} \rangle \in \mathbb{F}$ satisfying, for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and all $\lambda \in \mathbb{F}$,

i. *positivity*: $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0$;
ii. *definiteness*: $\langle \mathbf{v}, \mathbf{v} \rangle = 0$ if and only if $\mathbf{v} = \mathbf{0}$;
iii. *additivity in the first slot*: $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$;
iv. *homogeneity in the first slot*: $\langle \lambda \mathbf{u}, \mathbf{v} \rangle = \lambda \langle \mathbf{u}, \mathbf{v} \rangle$;
v. *conjugate symmetry*: $\langle \mathbf{u}, \mathbf{v} \rangle = \overline{\langle \mathbf{v}, \mathbf{u} \rangle}$.

In the positivity axiom, $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0$ means that $\langle \mathbf{v}, \mathbf{v} \rangle$ is a real nonnegative scalar.

**Definition 1.1.4.** *Inner-Product Space*
An *inner-product space* is a vector space $\mathcal{V}$ together with an inner product on $\mathcal{V}$.

**Example 1.1.5.** *Standard Inner Product on $\mathbb{F}^n$*
For $\mathbf{x}, \mathbf{y} \in \mathbb{F}^n$, the standard inner product is $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{j=1}^{n} x_j \overline{y_j}$. When $\mathbb{F} = \mathbb{R}$, this is the usual dot product.

**Example 1.1.6.** *Function and Matrix Inner Products*
If $\mathcal{V} = C_{\mathbb{F}}[a, b]$, then $\langle f, g \rangle = \int_a^b f(t)\overline{g(t)}\, dt$ is an inner product.
If $\mathcal{V} = \mathrm{Mat}(m, n, \mathbb{F})$, then the *Frobenius inner product* is $\langle A, B \rangle_F := \sum_{i=1}^{m} \sum_{j=1}^{n} a_{ij}\overline{b_{ij}}$.

**Example 1.1.7.** *Square-Summable Sequences*
Set $\ell^2(\mathbb{F}) := \left\{ \mathbf{x} = (x_j)_{j \geq 1} : \sum_{j=1}^{\infty} |x_j|^2 < \infty \right\}$, where $\langle \mathbf{x}, \mathbf{y} \rangle := \sum_{j=1}^{\infty} x_j \overline{y_j}$. Since $|x_j\overline{y_j}| \leq (|x_j|^2 + |y_j|^2)/2$, the series converges absolutely; termwise operations give an inner product on $\ell^2(\mathbb{F})$, with $\mathbb{F}^{(\mathbb{N})} \subseteq \ell^2(\mathbb{F})$.

**Proposition 1.1.8.** *Basic Inner-Product Identities*
*Let $\mathcal{V}$ be an inner-product space. Then for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and $\lambda \in \mathbb{F}$,*

*i. for fixed $\mathbf{u}$, the map $\mathbf{v} \mapsto \langle \mathbf{v}, \mathbf{u} \rangle$ is linear;*
*ii. $\langle \mathbf{0}, \mathbf{u} \rangle = 0$ and $\langle \mathbf{u}, \mathbf{0} \rangle = 0$;*
*iii. $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$;*
*iv. $\langle \mathbf{u}, \lambda \mathbf{v} \rangle = \overline{\lambda} \langle \mathbf{u}, \mathbf{v} \rangle$.*

*Proof.*

(i) This is exactly additivity and homogeneity in the first slot.
(ii) A linear map sends $\mathbf{0}$ to 0, so $\langle \mathbf{0}, \mathbf{u} \rangle = 0$. Conjugate symmetry gives $\langle \mathbf{u}, \mathbf{0} \rangle = \overline{\langle \mathbf{0}, \mathbf{u} \rangle} = 0$.
(iii) By conjugate symmetry and additivity, $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \overline{\langle \mathbf{v} + \mathbf{w}, \mathbf{u} \rangle} = \overline{\langle \mathbf{v}, \mathbf{u} \rangle + \langle \mathbf{w}, \mathbf{u} \rangle} = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$.

(iv) By conjugate symmetry and homogeneity, $\langle \mathbf{u}, \lambda\mathbf{v}\rangle = \overline{\langle \lambda\mathbf{v}, \mathbf{u}\rangle} = \overline{\lambda \langle \mathbf{v}, \mathbf{u}\rangle} = \overline{\lambda}\langle \mathbf{u}, \mathbf{v}\rangle$.

*Structural Commentary*: First-slot linearity and conjugate symmetry force conjugate linearity in second slot.
*Key Insight*: Once one slot is linear, the other slot is forced to be conjugate linear. □

**Definition 1.1.9.** *Norm Induced by an Inner Product*
For $\mathbf{v} \in \mathcal{V}$, the *norm* of $\mathbf{v}$ is $\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v}\rangle}$.




Figure 20: The norm is vector length, with $\|\mathbf{x}\|^2 = \langle \mathbf{x}, \mathbf{x}\rangle$.

**Lemma 1.1.10.** *Inner Products Separate Vectors*
*For* $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$, $(\forall \mathbf{x} \in \mathcal{V},\ \langle \mathbf{x}, \mathbf{u}\rangle = \langle \mathbf{x}, \mathbf{v}\rangle) \Longrightarrow \mathbf{u} = \mathbf{v}$, *and* $(\forall \mathbf{x} \in \mathcal{V},\ \langle \mathbf{x}, \mathbf{w}\rangle = 0) \Longleftrightarrow \mathbf{w} = \mathbf{0}$.

*Proof.* Set $\mathbf{x} = \mathbf{u} - \mathbf{v}$. Then $\|\mathbf{u} - \mathbf{v}\|^2 = \langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v}\rangle = 0$, hence $\mathbf{u} = \mathbf{v}$.
Taking $\mathbf{u} = \mathbf{w}$ and $\mathbf{v} = \mathbf{0}$ gives the second assertion; the reverse implication is immediate.
*Structural Commentary*: Testing against all vectors detects the vector being tested.
*Key Insight*: A vector orthogonal to every vector, including itself, must be zero. □

**Example 1.1.11.** *Norms from Standard Inner Products*
In $\mathbb{F}^n$, the norm from the standard inner product is the *Euclidean norm* $\|(x_1, \ldots, x_n)\|_2 = \sqrt{|x_1|^2 + \cdots + |x_n|^2}$.
In $C_{\mathbb{F}}[a, b]$, with the integral inner product, the derived norm is the *$L^2$ norm* $\|f\|_2 = \sqrt{\int_a^b |f(t)|^2\, dt}$.
In $\mathrm{Mat}(m, n, \mathbb{F})$, with the Frobenius inner product, the *Frobenius norm* is $\|A\|_F = \sqrt{\langle A, A\rangle_F} = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2}$, $A = [a_{ij}]$. The Frobenius norm is also called the *Schur norm* or *Hilbert-Schmidt norm*.

**Proposition 1.1.12.** *Basic Properties of the Derived Norm*
*Let $\mathcal{V}$ be an inner-product space with derived norm $\|\cdot\|$. For all $\mathbf{v} \in \mathcal{V}$ and $\lambda \in \mathbb{F}$,*

*i.* $\|\mathbf{v}\|$ *is real and nonnegative;*
*ii.* $\|\mathbf{v}\| = 0$ *if and only if* $\mathbf{v} = \mathbf{0}$*;*
*iii.* $\|\lambda\mathbf{v}\| = |\lambda|\,\|\mathbf{v}\|$.

*Proof.*

(i) By positivity, $\langle \mathbf{v}, \mathbf{v}\rangle$ is real and nonnegative, so its square root is real and nonnegative.
(ii) By definition, $\|\mathbf{v}\| = 0$ if and only if $\langle \mathbf{v}, \mathbf{v}\rangle = 0$, which holds if and only if $\mathbf{v} = \mathbf{0}$ by definiteness.
(iii) Compute $\|\lambda\mathbf{v}\|^2 = \langle \lambda\mathbf{v}, \lambda\mathbf{v}\rangle = \lambda\overline{\lambda}\langle \mathbf{v}, \mathbf{v}\rangle = |\lambda|^2\|\mathbf{v}\|^2$. Taking nonnegative square roots, $\|\lambda\mathbf{v}\| = |\lambda|\,\|\mathbf{v}\|$.

*Structural Commentary*: Positivity gives length; sesquilinearity gives scaling.
*Key Insight*: Squared length is easier to manipulate than length itself. □

**Theorem 1.1.13.** *Cauchy-Schwarz Inequality*
*For all $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, $|\langle \mathbf{u}, \mathbf{v}\rangle| \le \|\mathbf{u}\|\,\|\mathbf{v}\|$. Equality holds if and only if one of $\mathbf{u}, \mathbf{v}$ is a scalar multiple of the other.*

*Proof.* If $\mathbf{v} = \mathbf{0}$, both sides are 0. Assume $\mathbf{v} \neq \mathbf{0}$.
For any $\lambda \in \mathbb{F}$, positivity gives $0 \le \|\mathbf{u} - \lambda\mathbf{v}\|^2 = \|\mathbf{u}\|^2 - \lambda\langle \mathbf{v}, \mathbf{u}\rangle - \overline{\lambda}\langle \mathbf{u}, \mathbf{v}\rangle + |\lambda|^2\|\mathbf{v}\|^2$.
Choose $\lambda = \frac{\langle \mathbf{u}, \mathbf{v}\rangle}{\|\mathbf{v}\|^2}$. Since $\langle \mathbf{v}, \mathbf{u}\rangle = \overline{\langle \mathbf{u}, \mathbf{v}\rangle}$, the inequality becomes $0 \le \|\mathbf{u}\|^2 - \frac{|\langle \mathbf{u}, \mathbf{v}\rangle|^2}{\|\mathbf{v}\|^2}$.
Multiplying by $\|\mathbf{v}\|^2$ and taking square roots proves the inequality. Equality in this argument is equivalent to $\|\mathbf{u} - \lambda\mathbf{v}\| = 0$, hence to $\mathbf{u} = \lambda\mathbf{v}$ for the chosen $\lambda$.
Conversely, if $\mathbf{u} = \lambda\mathbf{v}$, then $|\langle \mathbf{u}, \mathbf{v}\rangle| = |\lambda|\ \langle \mathbf{v}, \mathbf{v}\rangle = \|\lambda\mathbf{v}\|\,\|\mathbf{v}\| = \|\mathbf{u}\|\,\|\mathbf{v}\|$, so equality holds.
The case $\mathbf{v} = \mathbf{0}$ is also covered because $\mathbf{v} = 0\mathbf{u}$.
*Structural Commentary*: Positivity of squared length controls the mixed inner-product term.
*Key Insight*: The best scalar comparison between two vectors bounds their inner product. □

**Theorem 1.1.14.** *Triangle Inequality*
*For all* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, $\|\mathbf{u}+\mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$.
*Equality holds if and only if one of* $\mathbf{u}, \mathbf{v}$ *is a nonnegative scalar multiple of the other.*

*Proof.* We compute

$$\|\mathbf{u}+\mathbf{v}\|^2 = \langle \mathbf{u}+\mathbf{v}, \mathbf{u}+\mathbf{v}\rangle = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + \langle \mathbf{u}, \mathbf{v}\rangle + \langle \mathbf{v}, \mathbf{u}\rangle = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2\operatorname{Re}\langle \mathbf{u}, \mathbf{v}\rangle$$
$$\leq \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2|\langle \mathbf{u}, \mathbf{v}\rangle| \leq \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2\|\mathbf{u}\|\|\mathbf{v}\| = \big(\|\mathbf{u}\| + \|\mathbf{v}\|\big)^2.$$

Taking square roots gives the inequality. If either vector is $\mathbf{0}$, the equality condition is immediate.
Otherwise, equality in the two displayed inequalities holds exactly when $\operatorname{Re}\langle \mathbf{u}, \mathbf{v}\rangle = |\langle \mathbf{u}, \mathbf{v}\rangle|$ and equality holds in Cauchy-Schwarz. The first condition says $\langle \mathbf{u}, \mathbf{v}\rangle$ is real nonnegative, and the second says $\mathbf{u} = a\mathbf{v}$ for some $a \in \mathbb{F}$. Then $a\|\mathbf{v}\|^2 = \langle \mathbf{u}, \mathbf{v}\rangle$ is real nonnegative, so $a \geq 0$.
Conversely, if $\mathbf{u} = a\mathbf{v}$ with $a \geq 0$, then $\|\mathbf{u}+\mathbf{v}\| = (a+1)\|\mathbf{v}\| = \|\mathbf{u}\| + \|\mathbf{v}\|$.
*Structural Commentary*: Square the norm and bound the cross term by Cauchy-Schwarz.
*Key Insight*: Equality means the two vectors point in the same direction. □




Figure 21: Triangle inequality: $\|\mathbf{u}+\mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$.

**Proposition 1.1.15.** *Reverse Triangle Inequality*
*For all* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, $|\|\mathbf{u}\| - \|\mathbf{v}\|| \leq \|\mathbf{u}-\mathbf{v}\|$.

*Proof.* The triangle inequality gives $\|\mathbf{u}\| \leq \|\mathbf{u}-\mathbf{v}\| + \|\mathbf{v}\|$, hence $\|\mathbf{u}\| - \|\mathbf{v}\| \leq \|\mathbf{u}-\mathbf{v}\|$.
Interchanging $\mathbf{u}$ and $\mathbf{v}$ gives $\|\mathbf{v}\| - \|\mathbf{u}\| \leq \|\mathbf{u}-\mathbf{v}\|$. Combining the two inequalities gives the result.
*Structural Commentary*: Distances control changes in length.
*Key Insight*: The ordinary triangle inequality applied twice gives the reverse one. □

**Definition 1.1.16.** *Norm and Normed Vector Space*
Let $\mathcal{V}$ be a vector space over $\mathbb{F}$. A *norm* on $\mathcal{V}$ is a function $\|\cdot\| : \mathcal{V} \to [0, \infty)$ such that, $\forall \mathbf{u}, \mathbf{v} \in \mathcal{V}$ and $\lambda \in \mathbb{F}$,

i. $\|\mathbf{u}\| \geq 0$;
ii. $\|\mathbf{u}\| = 0$ if and only if $\mathbf{u} = \mathbf{0}$;
iii. $\|\lambda\mathbf{u}\| = |\lambda|\,\|\mathbf{u}\|$;
iv. $\|\mathbf{u}+\mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$.

A *normed vector space* is a vector space equipped with a norm. Its *unit ball* is $\{\mathbf{v} \in \mathcal{V} : \|\mathbf{v}\| \leq 1\}$.

**Definition 1.1.17.** *Distance Induced by an Inner Product*
Let $\mathcal{V}$ be an inner-product space. The *distance* between two vectors in $\mathcal{V}$ is $d(\mathbf{u}, \mathbf{v}) := \|\mathbf{u}-\mathbf{v}\|$.

**Proposition 1.1.18.** *Inner-Product Distance Is a Metric*
*The function* $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u}-\mathbf{v}\|$ *is a metric on* $\mathcal{V}$*: for all* $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$,

*i.* $d(\mathbf{u}, \mathbf{v}) \geq 0$, *and* $d(\mathbf{u}, \mathbf{v}) = 0$ *if and only if* $\mathbf{u} = \mathbf{v}$*;*
*ii.* $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$*;*
*iii.* $d(\mathbf{u}, \mathbf{v}) \leq d(\mathbf{u}, \mathbf{w}) + d(\mathbf{w}, \mathbf{v})$.

*Proof.*

(i) By Proposition 1.1.12, $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u}-\mathbf{v}\| \geq 0$, with equality iff $\mathbf{u}-\mathbf{v} = \mathbf{0}$.
(ii) $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u}-\mathbf{v}\| = \|-(\mathbf{v}-\mathbf{u})\| = \|\mathbf{v}-\mathbf{u}\| = d(\mathbf{v}, \mathbf{u})$.
(iii) Since $\mathbf{u}-\mathbf{v} = (\mathbf{u}-\mathbf{w}) + (\mathbf{w}-\mathbf{v})$, Theorem 1.1.14 gives $d(\mathbf{u}, \mathbf{v}) \leq d(\mathbf{u}, \mathbf{w}) + d(\mathbf{w}, \mathbf{v})$.

*Structural Commentary*: The norm turns displacement vectors into distances.
*Key Insight*: Metric triangle inequalities are norm triangle inequalities applied to differences. □

**Perspective 1.1.19.** *Metric Viewpoint*
Define $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u}-\mathbf{v}\|$. Then $\mathbf{v}_n \to \mathbf{v}$ iff $\|\mathbf{v}_n - \mathbf{v}\| \to 0$.
In finite-dimensional inner-product spaces, subspaces are closed, linear maps are continuous, and the ambient space is complete. Infinite-dimensional analogues require additional hypotheses; subsequent projection and basis results therefore state finite-dimensional assumptions explicitly.

**Theorem 1.1.20.** *Parallelogram Identity*
*For all* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, $\|\mathbf{u}+\mathbf{v}\|^2 + \|\mathbf{u}-\mathbf{v}\|^2 = 2\big(\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2\big)$.

*Proof.* Expanding both squared norms gives

$$\|\mathbf{u}+\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \langle \mathbf{u}, \mathbf{v}\rangle + \langle \mathbf{v}, \mathbf{u}\rangle + \|\mathbf{v}\|^2, \quad \|\mathbf{u}-\mathbf{v}\|^2 = \|\mathbf{u}\|^2 - \langle \mathbf{u}, \mathbf{v}\rangle - \langle \mathbf{v}, \mathbf{u}\rangle + \|\mathbf{v}\|^2.$$

Adding the two identities cancels the cross terms and yields the result.
*Structural Commentary*: Opposite cross terms cancel when the two diagonal norms are added.
*Key Insight*: Inner-product norms remember parallelogram geometry through cancellation. □




Figure 22: Parallelogram identity: $\|\mathbf{u}+\mathbf{v}\|^2 + \|\mathbf{u}-\mathbf{v}\|^2 = 2(\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2)$.

**Proposition 1.1.21.** *Coordinate Norms on* $\mathbb{F}^n$
*For* $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{F}^n$, $\|\mathbf{x}\|_1 = \sum_{j=1}^{n} |x_j|$, $\|\mathbf{x}\|_2 = \left(\sum_{j=1}^{n} |x_j|^2\right)^{1/2}$, $\|\mathbf{x}\|_\infty = \max_{1\le j\le n} |x_j|$ *are norms on* $\mathbb{F}^n$.

*Proof.*

(i) For $\|\cdot\|_1$, nonnegativity, definiteness, and homogeneity follow from the corresponding facts for modulus. The triangle inequality follows from $\|\mathbf{x}+\mathbf{y}\|_1 = \sum_{j=1}^{n} |x_j + y_j| \le \sum_{j=1}^{n} (|x_j| + |y_j|) = \|\mathbf{x}\|_1 + \|\mathbf{y}\|_1$.

(ii) For $\|\cdot\|_2$, the axioms follow from its standard inner product via Proposition 1.1.12 and Theorem 1.1.14.

(iii) For $\|\cdot\|_\infty$, nonnegativity, definiteness, and homogeneity are immediate from the maximum of finitely many moduli. If $k$ is chosen so that $\|\mathbf{x}+\mathbf{y}\|_\infty = |x_k + y_k|$, then $\|\mathbf{x}+\mathbf{y}\|_\infty \le |x_k| + |y_k| \le \|\mathbf{x}\|_\infty + \|\mathbf{y}\|_\infty$.

*Structural Commentary*: The $\ell_1$ norm sums coordinate magnitudes, $\ell_\infty$ keeps only the largest coordinate magnitude, and $\ell_2$ comes from the standard inner product.
*Key Insight*: All three obey the same abstract axioms, even though their unit balls have different shapes. □




$\|\cdot\|_1 \le 1$ $\|\cdot\|_2 \le 1$ $\|\cdot\|_\infty \le 1$

Figure 23: The unit balls of $\|\cdot\|_1$, $\|\cdot\|_2$, and $\|\cdot\|_\infty$ in $\mathbb{R}^2$.

**Corollary 1.1.22.** *$\ell_1$ and $\ell_\infty$ Are Not Inner-Product Norms*
*If $n \ge 2$, then the norms $\|\cdot\|_1$ and $\|\cdot\|_\infty$ on $\mathbb{F}^n$ are not induced by any inner product.*

*Proof.* Let $\mathbf{e}_1, \mathbf{e}_2$ be the first two standard coordinate vectors.
For $\|\cdot\|_1$, $\|\mathbf{e}_1+\mathbf{e}_2\|_1^2 + \|\mathbf{e}_1-\mathbf{e}_2\|_1^2 = 4+4 = 8$, but $2\|\mathbf{e}_1\|_1^2 + 2\|\mathbf{e}_2\|_1^2 = 4$.
For $\|\cdot\|_\infty$, $\|\mathbf{e}_1+\mathbf{e}_2\|_\infty^2 + \|\mathbf{e}_1-\mathbf{e}_2\|_\infty^2 = 1+1 = 2$ but $2\|\mathbf{e}_1\|_\infty^2 + 2\|\mathbf{e}_2\|_\infty^2 = 4$.
Both norms violate Theorem 1.1.20, which every inner-product norm must satisfy.
*Structural Commentary*: The parallelogram identity detects inner-product norms.
*Key Insight*: Different unit-ball geometry can rule out hidden inner-product structure. □

**Theorem 1.1.23.** *Polarization Identities*
*Let $\mathcal{V}$ be an inner-product space and let $\mathbf{u}, \mathbf{v} \in \mathcal{V}$.*

*i. If $\mathbb{F} = \mathbb{R}$, then* $\langle \mathbf{u}, \mathbf{v}\rangle = \frac{1}{4}\left(\|\mathbf{u}+\mathbf{v}\|^2 - \|\mathbf{u}-\mathbf{v}\|^2\right)$.

*ii. If* $\mathbb{F} = \mathbb{C}$, *then* $\langle \mathbf{u}, \mathbf{v} \rangle = \frac{1}{4}\left( \|\mathbf{u}+\mathbf{v}\|^2 - \|\mathbf{u}-\mathbf{v}\|^2 + i\,\|\mathbf{u}+i\mathbf{v}\|^2 - i\,\|\mathbf{u}-i\mathbf{v}\|^2 \right)$.

*Proof.* Let $z := \langle \mathbf{u}, \mathbf{v} \rangle$.

(i) If $\mathbb{F} = \mathbb{R}$, then expanding squared norms gives $\|\mathbf{u}+\mathbf{v}\|^2 - \|\mathbf{u}-\mathbf{v}\|^2 = 4\operatorname{Re} z = 4z$, proving (i).

(ii) If $\mathbb{F} = \mathbb{C}$, the first expansion gives $\|\mathbf{u}+\mathbf{v}\|^2 - \|\mathbf{u}-\mathbf{v}\|^2 = 4\operatorname{Re} z$. The same expansion with $i\mathbf{v}$ gives $\|\mathbf{u}+i\mathbf{v}\|^2 - \|\mathbf{u}-i\mathbf{v}\|^2 = 4\operatorname{Im} z$. Combining the real and imaginary parts gives (ii).

*Structural Commentary*: Two norm differences extract the real and imaginary parts.
*Key Insight*: The norm determines the inner product because it determines both parts of every cross term. □




Figure 24: Polarization identity: $\|\mathbf{u}+\mathbf{v}\|^2 - \|\mathbf{u}-\mathbf{v}\|^2 = 4\operatorname{Re}\langle \mathbf{u}, \mathbf{v} \rangle$.

**Proposition 1.1.24.** *Operator Zero Tests from Inner Products*
*Let* $T \in \mathcal{L}(\mathcal{V})$.

*i. If* $\langle T\mathbf{v}, \mathbf{w} \rangle = 0$ *for every* $\mathbf{v}, \mathbf{w} \in \mathcal{V}$, *then* $T = 0$.
*ii. If* $\mathbb{F} = \mathbb{C}$ *and* $\langle T\mathbf{v}, \mathbf{v} \rangle = 0$ *for every* $\mathbf{v} \in \mathcal{V}$, *then* $T = 0$.
*iii. The preceding diagonal test can fail over* $\mathbb{R}$.

*Proof.*

(i) Taking $\mathbf{w} = T\mathbf{v}$ gives $\|T\mathbf{v}\|^2 = 0$ for every $\mathbf{v}$.

(ii) Put $Q(\mathbf{x}) = \langle T\mathbf{x}, \mathbf{x} \rangle$. For $\mathbf{x}, \mathbf{y} \in \mathcal{V}$ and $r \in \mathbb{C}$, the hypothesis applied to $r\mathbf{x}+\mathbf{y}$ gives

$$0 = Q(r\mathbf{x}+\mathbf{y}) = |r|^2 Q(\mathbf{x}) + Q(\mathbf{y}) + r\langle T\mathbf{x}, \mathbf{y} \rangle + \overline{r}\langle T\mathbf{y}, \mathbf{x} \rangle = r\langle T\mathbf{x}, \mathbf{y} \rangle + \overline{r}\langle T\mathbf{y}, \mathbf{x} \rangle .$$

Taking $r = 1, i$ gives $\langle T\mathbf{x}, \mathbf{y} \rangle = 0$ for all $\mathbf{x}, \mathbf{y}$; by (i), $T = 0$.

(iii) On standard $\mathbb{R}^2$, a $90^\circ$ rotation has $\langle T\mathbf{v}, \mathbf{v} \rangle = 0$ for all $\mathbf{v}$, yet $T \neq 0$.

*Structural Commentary*: Complex scalar directions recover mixed inner products from diagonal values.
*Key Insight*: Over $\mathbb{R}$, a skew-symmetric operator can hide on the diagonal; over $\mathbb{C}$, polarization detects it. □

**Principle 1.1.25.** *Geometry from Algebra*
Most geometric statements in an inner-product space come from expanding $\|\mathbf{u} \pm \mathbf{v}\|^2$ and then interpreting the cross terms $\langle \mathbf{u}, \mathbf{v} \rangle$ and $\langle \mathbf{v}, \mathbf{u} \rangle$.

## 1.2 Unit Vectors and Orthogonality

**Definition 1.2.1.** *Unit Vector and Normalization*
A vector $\mathbf{u} \in \mathcal{V}$ is a *unit vector* if $\|\mathbf{u}\| = 1$. If $\mathbf{v} \neq \mathbf{0}$, then $\frac{\mathbf{v}}{\|\mathbf{v}\|}$ is called the *normalization* of $\mathbf{v}$.

**Proposition 1.2.2.** *Normalization Gives a Unit Vector*
*If* $\mathbf{v} \in \mathcal{V}$ *and* $\mathbf{v} \neq \mathbf{0}$, *then* $\frac{\mathbf{v}}{\|\mathbf{v}\|}$ *is a unit vector in the same direction as* $\mathbf{v}$.

*Proof.* By Proposition 1.1.12, $\|\mathbf{v}\| > 0$ and $\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = \frac{1}{\|\mathbf{v}\|}\|\mathbf{v}\| = 1$.
The vector $\frac{\mathbf{v}}{\|\mathbf{v}\|}$ is a positive scalar multiple of $\mathbf{v}$, so it lies in the same one-dimensional direction. *Structural Commentary*: Normalization changes scale but preserves direction.
*Key Insight*: Divide by the length to make the length equal to one. □

**Definition 1.2.3.** *Orthogonal Vectors*
Vectors $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ are *orthogonal*, written $\mathbf{u} \perp \mathbf{v}$, if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

**Proposition 1.2.4.** *Orthogonality and the Zero Vector*
*Let* $\mathcal{V}$ *be an inner-product space.*

*i.* $\mathbf{u} \perp \mathbf{v}$ *if and only if* $\mathbf{v} \perp \mathbf{u}$.
*ii.* $\mathbf{0}$ *is orthogonal to every vector in* $\mathcal{V}$.

*iii. If* $\mathbf{v}$ *is orthogonal to itself, then* $\mathbf{v} = \mathbf{0}$.

*Proof.*

(i) If $\langle \mathbf{u}, \mathbf{v} \rangle = 0$, then conjugate symmetry gives $\langle \mathbf{v}, \mathbf{u} \rangle = \overline{0} = 0$, and the converse is identical.
(ii) This follows from Proposition 1.1.8, which gives $\langle \mathbf{0}, \mathbf{u} \rangle = 0$ and $\langle \mathbf{u}, \mathbf{0} \rangle = 0$.
(iii) If $\mathbf{v} \perp \mathbf{v}$, then $\langle \mathbf{v}, \mathbf{v} \rangle = 0$, hence $\mathbf{v} = \mathbf{0}$ by definiteness.

*Structural Commentary*: Conjugate symmetry makes orthogonality symmetric, and definiteness isolates $\mathbf{0}$.
*Key Insight*: Conjugate symmetry preserves the equation "equals zero." □

**Proposition 1.2.5.** *Mutually Orthogonal Nonzero Vectors Are Independent*
*Mutually orthogonal nonzero vectors* $\mathbf{v}_1, \dots, \mathbf{v}_m$ *in an inner-product space are linearly independent.*

*Proof.* If $\sum_{j=1}^{m} c_j \mathbf{v}_j = \mathbf{0}$, then for each $k$, $0 = \left\langle \sum_{j=1}^{m} c_j \mathbf{v}_j, \mathbf{v}_k \right\rangle = c_k \langle \mathbf{v}_k, \mathbf{v}_k \rangle$.
Since $\langle \mathbf{v}_k, \mathbf{v}_k \rangle > 0$, $c_k = 0$. Thus all coefficients vanish.
*Structural Commentary*: Orthogonality isolates one coefficient at a time.
*Key Insight*: Pair a dependence relation with one vector from the list; every cross term disappears. □

## 1.3 Orthonormal Bases and Gram–Schmidt

**Definition 1.3.1.** *Orthonormal List*
A list $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ of vectors in an inner-product space $\mathcal{V}$ is *orthonormal* if

$$\langle \mathbf{e}_j, \mathbf{e}_k \rangle = \begin{cases} 1, & j = k, \\ 0, & j \neq k. \end{cases}$$

Equivalently, every $\mathbf{e}_j$ has norm 1 and distinct vectors in the list are orthogonal.

**Definition 1.3.2.** *Orthogonal and Orthonormal Sets*
An indexed family $(\mathbf{u}_i)_{i \in I}$ is *orthogonal* if $\langle \mathbf{u}_i, \mathbf{u}_j \rangle = 0$ for $i \neq j$, and *orthonormal* if $\langle \mathbf{u}_i, \mathbf{u}_j \rangle = \delta_{ij}$, $i, j \in I$.
A nonzero orthogonal family becomes orthonormal after replacing each $\mathbf{u}_i$ by $\mathbf{u}_i / \|\mathbf{u}_i\|$.

**Proposition 1.3.3.** *Norm of an Orthonormal Linear Combination*
*If* $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ *is an orthonormal list in* $\mathcal{V}$*, then* $\|a_1 \mathbf{e}_1 + \cdots + a_m \mathbf{e}_m\|^2 = |a_1|^2 + \cdots + |a_m|^2$ *for all* $a_1, \dots, a_m \in \mathbb{F}$.

*Proof.* Using orthonormality, $\left\| \sum_{j=1}^{m} a_j \mathbf{e}_j \right\|^2 = \left\langle \sum_{j=1}^{m} a_j \mathbf{e}_j, \sum_{k=1}^{m} a_k \mathbf{e}_k \right\rangle = \sum_{j=1}^{m} \sum_{k=1}^{m} a_j \overline{a_k} \langle \mathbf{e}_j, \mathbf{e}_k \rangle = \sum_{j=1}^{m} |a_j|^2$.
*Structural Commentary*: Orthonormality deletes every cross term in the squared-norm expansion.
*Key Insight*: In orthonormal coordinates, length is computed coefficient-by-coefficient. □

**Corollary 1.3.4.** *Orthonormal Lists Are Linearly Independent*
*Every orthonormal list of vectors in an inner-product space is linearly independent.*

*Proof.* If $a_1 \mathbf{e}_1 + \cdots + a_m \mathbf{e}_m = \mathbf{0}$, then Proposition 1.3.3 gives $0 = \|\mathbf{0}\|^2 = \|a_1 \mathbf{e}_1 + \cdots + a_m \mathbf{e}_m\|^2 = \sum_{j=1}^{m} |a_j|^2$.
Hence each $a_j = 0$, so the list is linearly independent.
*Structural Commentary*: The norm identity converts a linear relation into a sum of nonnegative scalars.
*Key Insight*: A zero orthonormal linear combination can only have zero coefficients. □

**Definition 1.3.5.** *Orthonormal Basis*
An *orthonormal basis* of $\mathcal{V}$ is a basis of $\mathcal{V}$ that is an orthonormal list.

**Proposition 1.3.6.** *Right-Length Orthonormal Lists Are Bases*
*If* $\mathcal{V}$ *is finite-dimensional and* $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ *is an orthonormal list in* $\mathcal{V}$ *with* $n = \dim(\mathcal{V})$*, then* $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ *is an orthonormal basis of* $\mathcal{V}$.

*Proof.* By Corollary 1.3.4, the list is linearly independent.
Since its length is $\dim(\mathcal{V})$, Proposition 1.4.3 implies that it is a basis.
*Structural Commentary*: Orthonormality gives independence; the correct length forces spanning.
*Key Insight*: In finite dimensions, $n$ independent vectors in an $n$-dimensional space already form a basis. □

**Theorem 1.3.7.** *Gram–Schmidt Process*
*Let* $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ *be a linearly independent list in an inner-product space* $\mathcal{V}$.
*Define* $\mathbf{u}_1 = \mathbf{v}_1$, $\mathbf{e}_1 = \frac{\mathbf{u}_1}{\|\mathbf{u}_1\|}$ *and, for* $2 \le j \le m$, $\mathbf{u}_j = \mathbf{v}_j - \sum_{k=1}^{j-1} \langle \mathbf{v}_j, \mathbf{e}_k \rangle \, \mathbf{e}_k$, $\mathbf{e}_j = \frac{\mathbf{u}_j}{\|\mathbf{u}_j\|}$.
*Then* $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ *is orthonormal and* $\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_j) = \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_j)$.

*Proof.* Proceed by induction on $j$. The case $j = 1$ is normalization of the nonzero vector $\mathbf{v}_1$.
Assume the result holds through $j-1$. Since $\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_{j-1}) = \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_{j-1})$ and the $\mathbf{v}$-list is linearly independent, $\mathbf{v}_j \notin \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_{j-1})$. Hence $\mathbf{u}_j \neq \mathbf{0}$, because $\mathbf{u}_j = \mathbf{0}$ would put $\mathbf{v}_j$ in that span.
For $k < j$, $\langle \mathbf{u}_j, \mathbf{e}_k \rangle = \langle \mathbf{v}_j, \mathbf{e}_k \rangle - \sum_{i=1}^{j-1} \langle \mathbf{v}_j, \mathbf{e}_i \rangle \langle \mathbf{e}_i, \mathbf{e}_k \rangle = \langle \mathbf{v}_j, \mathbf{e}_k \rangle - \langle \mathbf{v}_j, \mathbf{e}_k \rangle = 0$.
Thus $\mathbf{e}_j$ is a unit vector orthogonal to $\mathbf{e}_1, \dots, \mathbf{e}_{j-1}$.
Also $\mathbf{e}_j$ lies in $\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_j)$ because it is a nonzero scalar multiple of $\mathbf{u}_j$, and $\mathbf{u}_j$ is built from $\mathbf{v}_j$ and the previous $\mathbf{e}_k$, whose span agrees with $\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_{j-1})$.
Conversely, $\mathbf{v}_j = \mathbf{u}_j + \sum_{k=1}^{j-1} \langle \mathbf{v}_j, \mathbf{e}_k \rangle \, \mathbf{e}_k$ lies in $\operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_j)$.
Combining this with the induction hypothesis gives equality of the two spans at step $j$.
*Structural Commentary*: Subtract the previous-span component, then normalize.
*Key Insight*: Gram–Schmidt changes the basis vectors but preserves every initial span. □

**Theorem 1.3.8.** *Gram–Schmidt for Arbitrary Finite Lists*
*Let* $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ *be a finite list in an inner-product space.*
*Define* $\mathbf{u}_1 = \mathbf{v}_1$ *and, for* $2 \le k \le m$, $\mathbf{u}_k = \mathbf{v}_k - \sum_{\substack{1 \le j < k \\ \mathbf{u}_j \neq \mathbf{0}}} \frac{\langle \mathbf{v}_k, \mathbf{u}_j \rangle}{\langle \mathbf{u}_j, \mathbf{u}_j \rangle} \mathbf{u}_j$.
*Set* $U_k = \operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_k)$ *and* $V_k = \operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_k)$, *with* $U_0 = V_0 = \{\mathbf{0}\}$.
*Then* $(\mathbf{u}_1, \dots, \mathbf{u}_m)$ *is orthogonal,* $U_k = V_k$ *for* $1 \le k \le m$.

*Proof.* Induct on $k$. The case $k = 0$ is trivial. Assume $(\mathbf{u}_1, \dots, \mathbf{u}_{k-1})$ is orthogonal and $U_{k-1} = V_{k-1}$.
For $i < k$ with $\mathbf{u}_i \neq \mathbf{0}$, $\langle \mathbf{u}_k, \mathbf{u}_i \rangle = \langle \mathbf{v}_k, \mathbf{u}_i \rangle - \frac{\langle \mathbf{v}_k, \mathbf{u}_i \rangle}{\langle \mathbf{u}_i, \mathbf{u}_i \rangle} \langle \mathbf{u}_i, \mathbf{u}_i \rangle = 0$, while $\mathbf{u}_i = \mathbf{0}$ is trivial.
Hence the list remains orthogonal. Also $\mathbf{u}_k \in V_k$ and $\mathbf{v}_k \in U_k$, so $U_k = V_k$.
*Structural Commentary*: Gram–Schmidt preserves every prefix span.
*Key Insight*: Each step subtracts previous orthogonal directions. □

**Corollary 1.3.9.** *Zero Remainders Detect Dependence*
*Let* $(\mathbf{u}_1, \dots, \mathbf{u}_m)$ *be constructed from* $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ *by Theorem 1.3.8,*
*and set* $U_k = \operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_k)$, $V_k = \operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_k)$, *with* $U_0 = V_0 = \{\mathbf{0}\}$.
*Then* $\mathbf{u}_k = \mathbf{0} \iff \mathbf{v}_k \in V_{k-1}$, $1 \le k \le m$.

*Proof.* The case $k = 1$ follows from $\mathbf{u}_1 = \mathbf{v}_1$ and $V_0 = \{\mathbf{0}\}$. Let $2 \le k \le m$.
($\Rightarrow$) If $\mathbf{u}_k = \mathbf{0}$, the recursion gives $\mathbf{v}_k \in U_{k-1} = V_{k-1}$.
($\Leftarrow$) If $\mathbf{v}_k \in V_{k-1} = U_{k-1}$, write $\mathbf{v}_k = \sum_{\substack{1 \le j < k \\ \mathbf{u}_j \neq \mathbf{0}}} a_j \mathbf{u}_j$. Orthogonality gives $a_j = \frac{\langle \mathbf{v}_k, \mathbf{u}_j \rangle}{\langle \mathbf{u}_j, \mathbf{u}_j \rangle}$, recursion gives $\mathbf{u}_k = \mathbf{0}$.
*Structural Commentary*: Zero remainders certify dependence.
*Key Insight*: A nonzero remainder is a new direction. □




Figure 25: One Gram–Schmidt step: subtract the previous-span component, then normalize.

**Example 1.3.10.** *Legendre Polynomials from Gram–Schmidt*
Let $\mathbb{R}[x]$ have inner product $\langle p, q \rangle = \int_{-1}^{1} p(x) q(x) \, dx$, and apply Gram–Schmidt to $(1, x, x^2, x^3, \dots)$.
The first unnormalized vectors are $u_1(x) = 1$, $u_2(x) = x$, $u_3(x) = x^2 - \frac{1}{3}$, $u_4(x) = x^3 - \frac{3}{5}x$.
Thus $(u_n)$ is, up to nonzero scalar multiples, the Legendre sequence.
*Structural Commentary*: Orthogonal polynomials are orthogonalized monomials.
*Key Insight*: Changing the inner product changes the polynomial family produced by Gram–Schmidt.

**Lemma 1.3.11.** *Gram–Schmidt on Orthogonal Lists*
*If $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is a nonzero mutually orthogonal list, then Gram–Schmidt produces $\mathbf{e}_j = \frac{\mathbf{v}_j}{\|\mathbf{v}_j\|}$, $j = 1, \dots, m$. In particular, if the original list is orthonormal, Gram–Schmidt leaves it unchanged.*

*Proof.* Assume $\mathbf{e}_i = \mathbf{v}_i / \|\mathbf{v}_i\|$ for $i < j$. Since $\mathbf{v}_j \perp \mathbf{v}_i$, we have $\langle \mathbf{v}_j, \mathbf{e}_i \rangle = 0$ for every $i < j$.
Hence the Gram–Schmidt remainder at step $j$ is $\mathbf{u}_j = \mathbf{v}_j$, so $\mathbf{e}_j = \mathbf{v}_j / \|\mathbf{v}_j\|$.
If the original list is orthonormal, then $\|\mathbf{v}_j\| = 1$ for every $j$.
*Structural Commentary*: Orthogonal input has no previous-span component to subtract.
*Key Insight*: Gram–Schmidt only removes nonorthogonal overlap. □

**Corollary 1.3.12.** *Existence of Orthonormal Bases*
*Every finite-dimensional inner-product space has an orthonormal basis.*

*Proof.* Choose any basis of $\mathcal{V}$ and apply Theorem 1.3.7.
The resulting orthonormal list has the same span as the original basis, hence spans $\mathcal{V}$.
*Structural Commentary*: Gram–Schmidt orthonormalizes a basis without changing its span.
*Key Insight*: Finite-dimensional inner-product spaces always admit geometry-friendly coordinates. □

**Corollary 1.3.13.** *Extending an Orthonormal List*
*Every orthonormal list in a finite-dimensional inner-product space can be extended to an orthonormal basis.*

*Proof.* Let $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ be orthonormal. It is linearly independent by Corollary 1.3.4, so extend it to a basis $(\mathbf{e}_1, \dots, \mathbf{e}_m, \mathbf{v}_{m+1}, \dots, \mathbf{v}_n)$ by Theorem 1.3.12.
During Gram–Schmidt, the first $m$ remainders are the original $\mathbf{e}_j$ because the list is already orthonormal; after normalization they remain unchanged. The remaining vectors are orthonormalized by Theorem 1.3.7, and the final list spans $\mathcal{V}$ because Gram–Schmidt preserves the total span.
*Structural Commentary*: Extend to a basis, then orthonormalize the new part.
*Key Insight*: Existing orthonormal directions are already cleaned up, so Gram–Schmidt does not move them. □

## 1.4 Orthonormal Coordinates and Inner-Product Identities

**Theorem 1.4.1.** *Coordinates in an Orthonormal Basis*
*Let $\mathcal{V}$ be finite-dimensional and let $\beta = (\mathbf{e}_1, \dots, \mathbf{e}_n)$ be an orthonormal basis of $\mathcal{V}$.*

*Then for every $\mathbf{v} \in \mathcal{V}$, $\mathbf{v} = \sum_{j=1}^{n} \langle \mathbf{v}, \mathbf{e}_j \rangle \, \mathbf{e}_j$, $\|\mathbf{v}\|^2 = \sum_{j=1}^{n} \left| \langle \mathbf{v}, \mathbf{e}_j \rangle \right|^2$ and $[\mathbf{v}]_\beta = \begin{bmatrix} \langle \mathbf{v}, \mathbf{e}_1 \rangle \\ \vdots \\ \langle \mathbf{v}, \mathbf{e}_n \rangle \end{bmatrix}$.*

*Proof.* Since $\beta$ is a basis, write $\mathbf{v} = \sum_{j=1}^{n} a_j \mathbf{e}_j$. Taking inner product with $\mathbf{e}_k$ gives $\langle \mathbf{v}, \mathbf{e}_k \rangle = \sum_{j=1}^{n} a_j \langle \mathbf{e}_j, \mathbf{e}_k \rangle = a_k$.
Thus the displayed expansion and coordinate column follow.
Applying Proposition 1.3.3 to the same expansion gives $\|\mathbf{v}\|^2 = \sum_{j=1}^{n} |a_j|^2 = \sum_{j=1}^{n} \left| \langle \mathbf{v}, \mathbf{e}_j \rangle \right|^2$.
*Structural Commentary*: An orthonormal basis makes coordinate extraction an inner-product computation.
*Key Insight*: Pairing with $\mathbf{e}_k$ isolates the $k$th coordinate. □




Figure 26: An orthonormal basis resolves a vector into perpendicular coordinate components.

**Remark 1.4.2.** *Cross Terms*
In an expansion such as $\|\mathbf{u} + \mathbf{v}\|^2$, the terms involving both vectors, $\langle \mathbf{u}, \mathbf{v} \rangle$ and $\langle \mathbf{v}, \mathbf{u} \rangle$, are called *cross terms*. The terms $\|\mathbf{u}\|^2$ and $\|\mathbf{v}\|^2$ are the pure squared-length terms.

**Theorem 1.4.3.** *Pythagorean Theorem*
*If* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ *and* $\mathbf{u} \perp \mathbf{v}$, *then* $\|\mathbf{u}+\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$.

*Proof.* Since $\mathbf{u} \perp \mathbf{v}$, Proposition 1.2.4 gives $\langle \mathbf{v}, \mathbf{u} \rangle = 0$. Thus

$$\|\mathbf{u}+\mathbf{v}\|^2 = \langle \mathbf{u}+\mathbf{v}, \mathbf{u}+\mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{u} \rangle + \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{v}, \mathbf{u} \rangle + \langle \mathbf{v}, \mathbf{v} \rangle = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 .$$

*Structural Commentary*: Orthogonality cancels the cross terms in the squared-norm expansion.
*Key Insight*: Perpendicular components add in squared length, not in length. □

**Proposition 1.4.4.** *Pythagorean Converse over* $\mathbb{R}$
*In a real inner-product space,* $\mathbf{u} \perp \mathbf{v} \iff \|\mathbf{u}+\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$.

*Proof.* Use $\|\mathbf{u}+\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + 2\langle \mathbf{u}, \mathbf{v} \rangle + \|\mathbf{v}\|^2$.
($\Rightarrow$) If $\mathbf{u} \perp \mathbf{v}$, then $\langle \mathbf{u}, \mathbf{v} \rangle = 0$, so the identity follows.
($\Leftarrow$) If the identity holds, then $2\langle \mathbf{u}, \mathbf{v} \rangle = 0$, hence $\mathbf{u} \perp \mathbf{v}$.

*Structural Commentary*: In real inner-product spaces, the cross term has no hidden imaginary part.
*Key Insight*: The Pythagorean identity detects orthogonality exactly over $\mathbb{R}$. □

**Theorem 1.4.5.** *Bessel's Inequality*
*Let* $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ *be an orthonormal list in an inner-product space* $\mathcal{V}$.
*Then, for every* $\mathbf{v} \in \mathcal{V}$, $\sum_{j=1}^{m} \left|\langle \mathbf{v}, \mathbf{e}_j \rangle\right|^2 \leq \|\mathbf{v}\|^2$.

*Proof.* Set $\mathbf{u} = \sum_{j=1}^{m} \langle \mathbf{v}, \mathbf{e}_j \rangle \mathbf{e}_j$. For each $k$, orthonormality gives $\langle \mathbf{v}-\mathbf{u}, \mathbf{e}_k \rangle = \langle \mathbf{v}, \mathbf{e}_k \rangle - \sum_{j=1}^{m} \langle \mathbf{v}, \mathbf{e}_j \rangle \langle \mathbf{e}_j, \mathbf{e}_k \rangle = 0$.
Thus $\mathbf{v}-\mathbf{u}$ is orthogonal to $\mathbf{u}$. By Pythagorean Theorem 1.4.3 and Proposition 1.3.3,

$$\|\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}-\mathbf{u}\|^2 \geq \|\mathbf{u}\|^2 = \sum_{j=1}^{m} \left|\langle \mathbf{v}, \mathbf{e}_j \rangle\right|^2 .$$

*Structural Commentary*: The chosen directions give an orthogonal part of $\mathbf{v}$; the residual is perpendicular.
*Key Insight*: Partial orthonormal coordinates cannot contain more energy than the whole vector. □

**Corollary 1.4.6.** *Coefficients Against an Orthonormal Sequence Vanish*
*Let* $(\mathbf{e}_1, \mathbf{e}_2, \ldots)$ *be an orthonormal sequence in an inner-product space* $\mathcal{V}$, *meaning every finite sublist is orthonormal. Then, for every* $\mathbf{v} \in \mathcal{V}$, $\lim_{n\to\infty} \langle \mathbf{v}, \mathbf{e}_n \rangle = 0$.

*Proof.* Fix $\epsilon > 0$. By Theorem 1.4.5, every finite set of indices satisfying $\left|\langle \mathbf{v}, \mathbf{e}_j \rangle\right| \geq \epsilon$ has cardinality at most $\|\mathbf{v}\|^2 / \epsilon^2$. Hence only finitely many $j$ have $\left|\langle \mathbf{v}, \mathbf{e}_j \rangle\right| \geq \epsilon$. This holds for every $\epsilon > 0$, so $\langle \mathbf{v}, \mathbf{e}_n \rangle \to 0$.
*Structural Commentary*: Bessel bounds the number of large orthonormal coefficients.
*Key Insight*: Infinitely many large coefficients would force infinite energy. □

**Theorem 1.4.7.** *Parseval's Identity*
*Let* $\beta = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$ *be an orthonormal basis of an inner-product space* $\mathcal{V}$.
*Then, for all* $\mathbf{v}, \mathbf{w} \in \mathcal{V}$, $\langle \mathbf{v}, \mathbf{w} \rangle = \sum_{j=1}^{n} \langle \mathbf{v}, \mathbf{e}_j \rangle \overline{\langle \mathbf{w}, \mathbf{e}_j \rangle} = \langle [\mathbf{v}]_\beta, [\mathbf{w}]_\beta \rangle_{\mathbb{F}^n}$.

*Proof.* By Theorem 1.4.1, $\mathbf{v} = \sum_{j=1}^{n} \langle \mathbf{v}, \mathbf{e}_j \rangle \mathbf{e}_j$ and $\mathbf{w} = \sum_{k=1}^{n} \langle \mathbf{w}, \mathbf{e}_k \rangle \mathbf{e}_k$. By linearity in first slot and conjugate linearity in second, $\langle \mathbf{v}, \mathbf{w} \rangle = \sum_{j=1}^{n} \sum_{k=1}^{n} \langle \mathbf{v}, \mathbf{e}_j \rangle \overline{\langle \mathbf{w}, \mathbf{e}_k \rangle} \langle \mathbf{e}_j, \mathbf{e}_k \rangle = \sum_{j=1}^{n} \langle \mathbf{v}, \mathbf{e}_j \rangle \overline{\langle \mathbf{w}, \mathbf{e}_j \rangle}$.
The last expression is exactly the standard inner product of the coordinate columns in $\mathbb{F}^n$.
*Structural Commentary*: Orthonormal coordinates make inner products coordinatewise.
*Key Insight*: Orthonormal coordinates preserve angles and lengths, not just linear combinations. □

**Theorem 1.4.8.** *Matrix Entries from an Orthonormal Codomain Basis*
*Let* $\mathcal{V}$ *and* $\mathcal{W}$ *be finite-dimensional vector spaces over* $\mathbb{F}$, *with* $\mathcal{W}$ *an inner-product space. Let* $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ *be a basis of* $\mathcal{V}$ *and let* $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_m)$ *be an orthonormal basis of* $\mathcal{W}$. *If* $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, *then*

$$[T]_{\gamma \leftarrow \beta} = \begin{bmatrix} \langle T\mathbf{v}_1, \mathbf{w}_1 \rangle & \cdots & \langle T\mathbf{v}_n, \mathbf{w}_1 \rangle \\ \vdots & \ddots & \vdots \\ \langle T\mathbf{v}_1, \mathbf{w}_m \rangle & \cdots & \langle T\mathbf{v}_n, \mathbf{w}_m \rangle \end{bmatrix} .$$

*Conversely, for each* $A \in \mathrm{Mat}(m, n, \mathbb{F})$, *there is a unique* $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ *such that* $[T]_{\gamma \leftarrow \beta} = A$.

*Proof.* The $j$th column of $[T]_{\gamma\leftarrow\beta}$ is $[T\mathbf{v}_j]_\gamma$. Since $\gamma$ is orthonormal, Theorem 1.4.1 gives

$$[T\mathbf{v}_j]_\gamma = \begin{bmatrix} \langle T\mathbf{v}_j, \mathbf{w}_1\rangle \\ \vdots \\ \langle T\mathbf{v}_j, \mathbf{w}_m\rangle \end{bmatrix}.$$

This proves the displayed matrix formula. The converse is exactly the surjectivity and injectivity of the coordinate encoding in Theorem 3.4.4.
*Structural Commentary*: Columns are basis-image coordinates, read by inner products.
*Key Insight*: Each entry measures $T\mathbf{v}_j$ against one output direction. □




Figure 27: The $j$th column of $[T]_{\gamma\leftarrow\beta}$ is obtained by resolving $T\mathbf{v}_j$ against the orthonormal basis of $\mathcal{W}$.

**Corollary 1.4.9.** *Inner Products Through an Operator Matrix*
*Let $\beta = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be an orthonormal basis of a finite-dimensional inner-product space $\mathcal{V}$, and let $T \in \mathcal{L}(\mathcal{V})$. Then, for all $\mathbf{v}, \mathbf{w} \in \mathcal{V}$, $\langle T\mathbf{v}, \mathbf{w}\rangle = \langle [T]_{\beta\leftarrow\beta}[\mathbf{v}]_\beta, [\mathbf{w}]_\beta\rangle_{\mathbb{F}^n}$.*

*Proof.* By Proposition 3.2.5, $[T\mathbf{v}]_\beta = [T]_{\beta\leftarrow\beta}[\mathbf{v}]_\beta$. Applying Theorem 1.4.7 to $T\mathbf{v}$ and $\mathbf{w}$ gives the result.
*Structural Commentary*: Orthonormal coordinates turn $\langle T\mathbf{v}, \mathbf{w}\rangle$ into matrix computation.
*Key Insight*: Orthonormal coordinates make operator geometry computable by matrices. □

## 1.5 Gram Matrices

**Definition 1.5.1.** *Gram Matrix*
For vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$ in an inner-product space, the *Gram matrix* is

$$\operatorname{Gram}(\mathbf{u}_1, \ldots, \mathbf{u}_n) := \begin{bmatrix} \langle \mathbf{u}_1, \mathbf{u}_1\rangle & \langle \mathbf{u}_2, \mathbf{u}_1\rangle & \cdots & \langle \mathbf{u}_n, \mathbf{u}_1\rangle \\ \langle \mathbf{u}_1, \mathbf{u}_2\rangle & \langle \mathbf{u}_2, \mathbf{u}_2\rangle & \cdots & \langle \mathbf{u}_n, \mathbf{u}_2\rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle \mathbf{u}_1, \mathbf{u}_n\rangle & \langle \mathbf{u}_2, \mathbf{u}_n\rangle & \cdots & \langle \mathbf{u}_n, \mathbf{u}_n\rangle \end{bmatrix}.$$

**Proposition 1.5.2.** *Gram Matrix Quadratic Form*
*For any vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$, Gram matrix $G = \operatorname{Gram}(\mathbf{u}_1, \ldots, \mathbf{u}_n)$ is Hermitian and satisfies $\mathbf{c}^*G\mathbf{c} = \left\| \sum_{j=1}^n c_j\mathbf{u}_j \right\|^2$ for every $\mathbf{c} = [c_1 \ \cdots \ c_n]^t \in \mathbb{F}^n$. Hence the Gram quadratic form is nonnegative. If $\mathbf{u}_1, \ldots, \mathbf{u}_n$ are linearly independent, then it is positive for every nonzero $\mathbf{c}$.*

*Proof.* Hermitian symmetry follows from conjugate symmetry of the inner product. Expanding the squared norm gives $\left\| \sum_{j=1}^n c_j\mathbf{u}_j \right\|^2 = \sum_{i,j} \overline{c_i} \langle \mathbf{u}_j, \mathbf{u}_i\rangle c_j = \mathbf{c}^*G\mathbf{c}$. The right-hand side is a squared norm, so it is nonnegative. If $\mathbf{u}_1, \ldots, \mathbf{u}_n$ are linearly independent, the squared norm is 0 only when all $c_j = 0$.
*Structural Commentary*: A Gram matrix records every pairwise inner product in a list of vectors.
*Key Insight*: The quadratic form of a Gram matrix is exactly the squared norm of a linear combination. □

## 1.6 Finite-Dimensional Riesz Representation

**Theorem 1.6.1.** *Riesz Representation Theorem*
*Let $\mathcal{V}$ be a finite-dimensional inner-product space and let $\varphi : \mathcal{V} \to \mathbb{F}$ be a linear functional. Then there is a unique vector $\mathbf{u} \in \mathcal{V}$ such that $\varphi(\mathbf{v}) = \langle \mathbf{v}, \mathbf{u}\rangle$ for every $\mathbf{v} \in \mathcal{V}$.*

*Proof.* Let $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be an orthonormal basis of $\mathcal{V}$.
By Theorem 1.4.1, $\varphi(\mathbf{v}) = \varphi\left( \sum_{j=1}^n \langle \mathbf{v}, \mathbf{e}_j\rangle \mathbf{e}_j \right) = \sum_{j=1}^n \langle \mathbf{v}, \mathbf{e}_j\rangle \varphi(\mathbf{e}_j)$.

Set $\mathbf{u} = \sum_{j=1}^{n} \overline{\varphi(\mathbf{e}_j)}\,\mathbf{e}_j$. Since inner product is conjugate linear in second slot, $\langle \mathbf{v}, \mathbf{u} \rangle = \sum_{j=1}^{n} \langle \mathbf{v}, \mathbf{e}_j \rangle\, \varphi(\mathbf{e}_j) = \varphi(\mathbf{v})$.
For uniqueness, suppose $\langle \mathbf{v}, \mathbf{u}_1 \rangle = \langle \mathbf{v}, \mathbf{u}_2 \rangle \ \ \forall \mathbf{v} \in \mathcal{V}$. Taking $\mathbf{v} = \mathbf{u}_1 - \mathbf{u}_2$ gives $\|\mathbf{u}_1 - \mathbf{u}_2\|^2 = 0$, hence $\mathbf{u}_1 = \mathbf{u}_2$.
*Structural Commentary*: An orthonormal basis packages a functional as one representing vector.
*Key Insight*: Finite-dimensional functionals are inner products with fixed vectors. □

**Definition 1.6.2.** *Riesz Vector*
Let $\mathcal{V}$ be finite-dimensional and let $\varphi : \mathcal{V} \to \mathbb{F}$ be a linear functional. The unique vector $\mathbf{u} \in \mathcal{V}$ satisfying $\varphi(\mathbf{v}) = \langle \mathbf{v}, \mathbf{u} \rangle$ for every $\mathbf{v} \in \mathcal{V}$ is the *Riesz vector* representing $\varphi$.

**Theorem 1.6.3.** *Riesz Map*
*Let $\mathcal{V}$ be finite-dimensional inner-product space, $\tau : \mathcal{V} \to \mathcal{V}'$, $\tau(\mathbf{u}) = \langle \cdot, \mathbf{u} \rangle$. Then $\tau$ is bijective, conjugate-linear. Hence $R = \tau^{-1}$ is conjugate-linear, $R(a\varphi + \psi) = \overline{a}\, R(\varphi) + R(\psi)$, $a \in \mathbb{F}$, $\varphi, \psi \in \mathcal{V}'$.*
*Over $\mathbb{R}$ this is ordinary linearity.*

*Proof.* For $a \in \mathbb{F}$ and $\mathbf{u}, \mathbf{w} \in \mathcal{V}$, $\tau(a\mathbf{u} + \mathbf{w}) = \overline{a}\tau(\mathbf{u}) + \tau(\mathbf{w})$, so $\tau$ is conjugate-linear.
If $\tau(\mathbf{u}) = 0$, then $\|\mathbf{u}\|^2 = \tau(\mathbf{u})(\mathbf{u}) = 0$, so $\tau$ is injective. Surjectivity is Theorem 1.6.1.
For $a \in \mathbb{F}$, $\varphi, \psi \in \mathcal{V}'$, and $\mathbf{v} \in \mathcal{V}$, $\langle \mathbf{v}, \overline{a}R(\varphi) + R(\psi) \rangle = a \langle \mathbf{v}, R(\varphi) \rangle + \langle \mathbf{v}, R(\psi) \rangle = (a\varphi + \psi)(\mathbf{v})$.
Uniqueness gives $R(a\varphi + \psi) = \overline{a}R(\varphi) + R(\psi)$.
*Structural Commentary*: Riesz identifies $\mathcal{V}$ with $\mathcal{V}'$ conjugate-linearly.
*Key Insight*: The representing slot conjugates scalars. □




Figure 28: Riesz identifies a functional with its representing vector.

**Perspective 1.6.4.** *Riesz as Coordinate Duality*
The vector $\sum_{j=1}^{n} \overline{\varphi(\mathbf{e}_j)}\mathbf{e}_j$ seems to depend on the orthonormal basis, but Theorem 1.6.1 shows that the resulting representing vector is intrinsic.

# 2 Orthogonality, Projections, and Least Squares

## 2.1 Orthogonal Complements

Orthogonal complements turn orthogonality into a subspace operation; projections turn it into canonical decompositions and best approximations.

**Definition 2.1.1.** *Orthogonality and Orthogonal Direct Sums*
Let $\mathcal{X}, \mathcal{Y} \subseteq \mathcal{V}$ and let $\mathcal{S}, \mathcal{T} \leq \mathcal{V}$.

i. $\mathcal{X}$ and $\mathcal{Y}$ are *orthogonal*, written $\mathcal{X} \perp \mathcal{Y}$, if $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ for all $\mathbf{x} \in \mathcal{X}$ and $\mathbf{y} \in \mathcal{Y}$.
ii. $\mathcal{V}$ is the *orthogonal direct sum* of $\mathcal{S}$ and $\mathcal{T}$, written $\mathcal{V} = \mathcal{S} \odot \mathcal{T}$, if $\mathcal{V} = \mathcal{S} \oplus \mathcal{T}$ and $\mathcal{S} \perp \mathcal{T}$.

**Definition 2.1.2.** *Orthogonal Complement*
Let $\mathcal{U} \subseteq \mathcal{V}$. The *orthogonal complement* of $\mathcal{U}$ is $\mathcal{U}^{\perp} := \{\mathbf{v} \in \mathcal{V} : \langle \mathbf{v}, \mathbf{u} \rangle = 0 \text{ for all } \mathbf{u} \in \mathcal{U}\}$.

**Proposition 2.1.3.** *Basic Properties of Orthogonal Complements*
*For subsets $\mathcal{U}, \mathcal{W} \subseteq \mathcal{V}$,*

*(i) $\mathcal{U}^{\perp}$ is a subspace of $\mathcal{V}$;*
*(ii) $\{\mathbf{0}\}^{\perp} = \mathcal{V}$;*
*(iii) $\mathcal{V}^{\perp} = \{\mathbf{0}\}$;*
*(iv) $\mathcal{U} \cap \mathcal{U}^{\perp} \subseteq \{\mathbf{0}\}$;*
*(v) if $\mathcal{U} \subseteq \mathcal{W}$, then $\mathcal{W}^{\perp} \subseteq \mathcal{U}^{\perp}$.*

*Proof.*

(i) We have $\mathbf{0} \in \mathcal{U}^{\perp}$. If $\mathbf{v}, \mathbf{w} \in \mathcal{U}^{\perp}$ and $a \in \mathbb{F}$, then for every $\mathbf{u} \in \mathcal{U}$, $\langle a\mathbf{v} + \mathbf{w}, \mathbf{u} \rangle = a \langle \mathbf{v}, \mathbf{u} \rangle + \langle \mathbf{w}, \mathbf{u} \rangle = 0$.

(ii) For every $\mathbf{v} \in \mathcal{V}$, $\langle \mathbf{v}, \mathbf{0} \rangle = 0$, so $\mathbf{v} \in \{\mathbf{0}\}^\perp$.
(iii) If $\mathbf{v} \in \mathcal{V}^\perp$, then $\langle \mathbf{v}, \mathbf{v} \rangle = 0$, so $\mathbf{v} = \mathbf{0}$.
(iv) If $\mathbf{v} \in \mathcal{U} \cap \mathcal{U}^\perp$, then $\langle \mathbf{v}, \mathbf{v} \rangle = 0$, so $\mathbf{v} = \mathbf{0}$.
(v) If $\mathbf{v} \in \mathcal{W}^\perp$, then $\mathbf{v}$ is orthogonal to every vector in $\mathcal{W}$, hence to every vector in $\mathcal{U}$.

*Structural Commentary*: Orthogonal complements are solution spaces of homogeneous inner-product equations. Larger input sets impose more equations, so their complements become smaller.
*Key Insight*: Orthogonality constraints are linear constraints. □

**Remark 2.1.4.** *Definiteness Matters*
For a bilinear form $B$, $\mathcal{U}^\perp = \{\mathbf{v} : B(\mathbf{v}, \mathbf{u}) = 0 \ \forall \mathbf{u} \in \mathcal{U}\}$ is still a subspace.
Without positive definiteness, however, one may have $\mathcal{V}^\perp \neq \{\mathbf{0}\}$ or $\mathcal{U} \cap \mathcal{U}^\perp \neq \{\mathbf{0}\}$.
Thus $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$ is an inner-product result, not a consequence of bilinearity alone.

**Theorem 2.1.5.** *Direct Sum with the Orthogonal Complement*
*If $\mathcal{U}$ is a finite-dimensional subspace of $\mathcal{V}$, then $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$.*

*Proof.* Let $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ be an orthonormal basis of $\mathcal{U}$. For $\mathbf{v} \in \mathcal{V}$, define $\mathbf{u} = \sum_{j=1}^{m} \langle \mathbf{v}, \mathbf{e}_j \rangle \, \mathbf{e}_j$ and set $\mathbf{w} = \mathbf{v} - \mathbf{u}$.

Clearly $\mathbf{u} \in \mathcal{U}$. For each $k$, $\langle \mathbf{w}, \mathbf{e}_k \rangle = \langle \mathbf{v}, \mathbf{e}_k \rangle - \sum_{j=1}^{m} \langle \mathbf{v}, \mathbf{e}_j \rangle \langle \mathbf{e}_j, \mathbf{e}_k \rangle = 0$.
If $\mathbf{x} \in \mathcal{U}$, write $\mathbf{x} = \sum_{k=1}^{m} a_k \mathbf{e}_k$. Then $\langle \mathbf{w}, \mathbf{x} \rangle = \sum_{k=1}^{m} \overline{a_k} \langle \mathbf{w}, \mathbf{e}_k \rangle = 0$, so $\mathbf{w} \in \mathcal{U}^\perp$ and $\mathbf{v} = \mathbf{u} + \mathbf{w}$.
Hence $\mathcal{V} = \mathcal{U} + \mathcal{U}^\perp$. The intersection is $\{\mathbf{0}\}$ by Proposition 2.1.3, so the sum is direct.
*Structural Commentary*: Each vector splits into a $\mathcal{U}$-component and an orthogonal residual.
*Key Insight*: Projection plus residual gives the canonical direct-sum decomposition. □

**Proposition 2.1.6.** *Orthogonal Direct Sum Characterization*
*For subspaces $\mathcal{S}, \mathcal{T} \subseteq \mathcal{V}$, the following are equivalent:*

*i.* $\mathcal{V} = \mathcal{S} \odot \mathcal{T}$.
*ii.* $\mathcal{V} = \mathcal{S} \oplus \mathcal{T}$ *and* $\mathcal{T} = \mathcal{S}^\perp$.

*Proof.*

(*i*) $\Rightarrow$ (*ii*) If $\mathcal{V} = \mathcal{S} \odot \mathcal{T}$, then $\mathcal{V} = \mathcal{S} \oplus \mathcal{T}$ and $\mathcal{T} \subseteq \mathcal{S}^\perp$. For $\mathbf{v} \in \mathcal{S}^\perp$, write $\mathbf{v} = \mathbf{s} + \mathbf{t}$ with $\mathbf{s} \in \mathcal{S}$, $\mathbf{t} \in \mathcal{T}$. Then $\mathbf{s} = \mathbf{v} - \mathbf{t} \in \mathcal{S}^\perp$, so $\mathbf{s} \in \mathcal{S} \cap \mathcal{S}^\perp = \{\mathbf{0}\}$ and $\mathbf{v} \in \mathcal{T}$. Hence $\mathcal{T} = \mathcal{S}^\perp$.
(*ii*) $\Rightarrow$ (*i*) If $\mathcal{V} = \mathcal{S} \oplus \mathcal{T}$ and $\mathcal{T} = \mathcal{S}^\perp$, then $\mathcal{S} \perp \mathcal{T}$, so $\mathcal{V} = \mathcal{S} \odot \mathcal{T}$.

*Structural Commentary*: Orthogonal complements are maximal orthogonal partners.
*Key Insight*: A full orthogonal summand equals the complement. □

**Corollary 2.1.7.** *Dimension of the Orthogonal Complement*
*If $\mathcal{V}$ is finite-dimensional and $\mathcal{U}$ is a subspace of $\mathcal{V}$, then* $\dim(\mathcal{U}^\perp) = \dim(\mathcal{V}) - \dim(\mathcal{U})$.

*Proof.* Since $\mathcal{V}$ is finite-dimensional, $\mathcal{U}$ is finite-dimensional by Corollary 1.3.13. Theorem 2.1.5 gives $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$. Thus Corollary 2.3.2 gives $\dim(\mathcal{V}) = \dim(\mathcal{U}) + \dim(\mathcal{U}^\perp)$.
*Structural Commentary*: Dimensions add across the orthogonal split.
*Key Insight*: Orthogonal complements contain exactly the dimensions not used by $\mathcal{U}$. □

**Corollary 2.1.8.** *Double Orthogonal Complement*
*If $\mathcal{U}$ is a finite-dimensional subspace of $\mathcal{V}$, then $\mathcal{U} = (\mathcal{U}^\perp)^\perp$.*

*Proof.* If $\mathbf{u} \in \mathcal{U}$, then $\mathbf{u}$ is orthogonal to every vector in $\mathcal{U}^\perp$, so $\mathcal{U} \subseteq (\mathcal{U}^\perp)^\perp$. Conversely, let $\mathbf{v} \in (\mathcal{U}^\perp)^\perp$. By Theorem 2.1.5, write $\mathbf{v} = \mathbf{u} + \mathbf{w}$ with $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{U}^\perp$. Since $\mathcal{U} \subseteq (\mathcal{U}^\perp)^\perp$, we have $\mathbf{w} = \mathbf{v} - \mathbf{u} \in (\mathcal{U}^\perp)^\perp$.
Hence $\mathbf{w} \in \mathcal{U}^\perp \cap (\mathcal{U}^\perp)^\perp$, so $\mathbf{w} = \mathbf{0}$ and $\mathbf{v} = \mathbf{u} \in \mathcal{U}$.
*Structural Commentary*: The orthogonal split leaves no extra direction beyond $\mathcal{U}$.
*Key Insight*: Taking the orthogonal complement twice recovers the original finite-dimensional subspace. □

**Corollary 2.1.9.** *Double Complement of a Finite Span*
*If $\mathcal{X} \subseteq \mathcal{V}$ and* $\operatorname{span}(\mathcal{X})$ *is finite-dimensional, then* $(\mathcal{X}^\perp)^\perp = \operatorname{span}(\mathcal{X})$.

*Proof.* Let $W = \operatorname{span}(\mathcal{X})$. As $\mathcal{X}^\perp = W^\perp$, $W$ is finite-dimensional, by Corollary 2.1.8, $(\mathcal{X}^\perp)^\perp = (W^\perp)^\perp = W$.
*Structural Commentary*: Orthogonal complements depend only on the span of the tested set.
*Key Insight*: Double orthogonality recovers the finite-dimensional span, not merely the original subset. □

**Perspective 2.1.10.** *Orthogonal Versus Complementary*
Orthogonality gives $\mathcal{W} \subseteq \mathcal{U}^\perp$ and $\mathcal{U} \subseteq \mathcal{W}^\perp$, not necessarily equality. In finite dimensions, equality follows once $\dim(\mathcal{U}) + \dim(\mathcal{W}) = \dim(\mathcal{V})$ by Corollary 2.1.7.




Figure 29: Orthogonal subspaces are complements only when their sum fills $\mathcal{V}$.

## 2.2 Orthogonal Projections

**Proposition 2.2.1.** *Orthogonal Projection onto a Line*
*Let* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ *with* $\mathbf{v} \neq \mathbf{0}$. *Define* $c := \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{v}\|^2}$, $\mathbf{w} := \mathbf{u} - c\mathbf{v}$. *Then* $\mathbf{w} \perp \mathbf{v}$ *and* $\mathbf{u} = c\mathbf{v} + \mathbf{w}$.
*The vector* $\operatorname{proj}_{\mathbf{v}}(\mathbf{u}) := c\mathbf{v} = \left\langle \mathbf{u}, \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\rangle \frac{\mathbf{v}}{\|\mathbf{v}\|}$ *is the orthogonal projection of* $\mathbf{u}$ *onto* $\operatorname{span}(\mathbf{v})$.

*Proof.* The decomposition $\mathbf{u} = c\mathbf{v} + \mathbf{w}$ follows from the definition of $\mathbf{w}$.
Moreover, $\langle \mathbf{w}, \mathbf{v} \rangle = \langle \mathbf{u} - c\mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle - c \langle \mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle - \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{v}\|^2} \|\mathbf{v}\|^2 = 0$. Thus $\mathbf{w} \perp \mathbf{v}$.
*Structural Commentary*: Choose $c$ so the remainder $\mathbf{u} - c\mathbf{v}$ has zero inner product with $\mathbf{v}$.
*Key Insight*: Projection is the part detected by pairing with the direction vector. □




Figure 30: Writing $\mathbf{u}$ as a component parallel to $\mathbf{v}$ plus a component orthogonal to $\mathbf{v}$.

**Definition 2.2.2.** *Orthogonal Projection onto a Subspace*
Let $\mathcal{U}$ be a finite-dimensional subspace of $\mathcal{V}$.
For $\mathbf{v} \in \mathcal{V}$, write the unique decomposition $\mathbf{v} = \mathbf{u} + \mathbf{w}$, $\mathbf{u} \in \mathcal{U}$, $\mathbf{w} \in \mathcal{U}^\perp$, from Theorem 2.1.5.
The *orthogonal projection* of $\mathcal{V}$ onto $\mathcal{U}$ is the map $P_{\mathcal{U}} : \mathcal{V} \to \mathcal{V}$ defined by $P_{\mathcal{U}}\mathbf{v} = \mathbf{u}$.




Figure 31: Orthogonal projection decomposes $\mathbf{v}$ into $P_{\mathcal{U}}\mathbf{v}$ plus an orthogonal residual.

**Proposition 2.2.3.** *Properties of Orthogonal Projection*
*Let* $\mathcal{U}$ *be a finite-dimensional subspace of* $\mathcal{V}$. *Then:*

*i.* $P_{\mathcal{U}} \in \mathcal{L}(\mathcal{V})$;
*ii.* $P_{\mathcal{U}}\mathbf{u} = \mathbf{u}$ *for every* $\mathbf{u} \in \mathcal{U}$;
*iii.* $P_{\mathcal{U}}\mathbf{w} = \mathbf{0}$ *for every* $\mathbf{w} \in \mathcal{U}^\perp$;
*iv.* $\operatorname{range}(P_{\mathcal{U}}) = \mathcal{U}$;
*v.* $\ker(P_{\mathcal{U}}) = \mathcal{U}^\perp$, *i.e. the kernel/nullspace of* $P_{\mathcal{U}}$ *is* $\mathcal{U}^\perp$;
*vi.* $\mathbf{v} - P_{\mathcal{U}}\mathbf{v} \in \mathcal{U}^\perp$ *for every* $\mathbf{v} \in \mathcal{V}$;
*vii.* $P_{\mathcal{U}}^2 = P_{\mathcal{U}}$;
*viii.* $\|P_{\mathcal{U}}\mathbf{v}\| \leq \|\mathbf{v}\|$ *for every* $\mathbf{v} \in \mathcal{V}$;
*ix. if* $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ *is an orthonormal basis of* $\mathcal{U}$, *then* $P_{\mathcal{U}}\mathbf{v} = \sum_{j=1}^{m} \langle \mathbf{v}, \mathbf{e}_j \rangle \mathbf{e}_j$, $\mathbf{v} \in \mathcal{V}$.

*Proof.*

(i) For linearity, let $a \in \mathbb{F}$ and write $\mathbf{v}_r = \mathbf{u}_r + \mathbf{w}_r$ with $\mathbf{u}_r \in \mathcal{U}$ and $\mathbf{w}_r \in \mathcal{U}^\perp$ for $r = 1, 2$.
Then $a\mathbf{v}_1 + \mathbf{v}_2 = (a\mathbf{u}_1 + \mathbf{u}_2) + (a\mathbf{w}_1 + \mathbf{w}_2)$, where first summand lies in $\mathcal{U}$ and second in $\mathcal{U}^\perp$.
By uniqueness of the orthogonal decomposition, $P_{\mathcal{U}}(a\mathbf{v}_1 + \mathbf{v}_2) = aP_{\mathcal{U}}\mathbf{v}_1 + P_{\mathcal{U}}\mathbf{v}_2$.
(ii) This follows from $\mathbf{u} = \mathbf{u} + \mathbf{0}$ with $\mathbf{u} \in \mathcal{U}$ and $\mathbf{0} \in \mathcal{U}^\perp$.
(iii) This follows from $\mathbf{w} = \mathbf{0} + \mathbf{w}$ with $\mathbf{0} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{U}^\perp$.
(iv) The definition gives $\text{range}(P_{\mathcal{U}}) \subseteq \mathcal{U}$, and (ii) gives $\mathcal{U} \subseteq \text{range}(P_{\mathcal{U}})$.
(v) Part (iii) gives $\mathcal{U}^\perp \subseteq \ker(P_{\mathcal{U}})$.
Conversely, if $P_{\mathcal{U}}\mathbf{v} = \mathbf{0}$, then $\mathbf{v} = \mathbf{0} + (\mathbf{v} - P_{\mathcal{U}}\mathbf{v})$ with $\mathbf{v} - P_{\mathcal{U}}\mathbf{v} \in \mathcal{U}^\perp$, so $\mathbf{v} \in \mathcal{U}^\perp$.
(vi) This is exactly the residual term in the decomposition $\mathbf{v} = P_{\mathcal{U}}\mathbf{v} + (\mathbf{v} - P_{\mathcal{U}}\mathbf{v})$.
(vii) By (ii), $P_{\mathcal{U}}(P_{\mathcal{U}}\mathbf{v}) = P_{\mathcal{U}}\mathbf{v}$ for every $\mathbf{v} \in \mathcal{V}$.
(viii) Write $\mathbf{v} = \mathbf{u} + \mathbf{w}$ with $\mathbf{u} \in \mathcal{U}$, $\mathbf{w} \in \mathcal{U}^\perp$. Then $\mathbf{u} \perp \mathbf{w}$ and $\|\mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{w}\|^2 \geq \|\mathbf{u}\|^2 = \|P_{\mathcal{U}}\mathbf{v}\|^2$.
Since norms are nonnegative, $\|P_{\mathcal{U}}\mathbf{v}\| \leq \|\mathbf{v}\|$.
(ix) This is the explicit $\mathcal{U}$-component constructed in the proof of Theorem 2.1.5.

*Structural Commentary*: Orthogonal projection keeps the $\mathcal{U}$-component and discards the $\mathcal{U}^\perp$-component.
*Key Insight*: $P_{\mathcal{U}}$ is the identity on $\mathcal{U}$ and zero on $\mathcal{U}^\perp$. □

**Definition 2.2.4.** *Rank-One Projection Matrix*
Let $\mathbf{u} \in \mathbb{F}^n$ be a unit vector. The *rank-one projection matrix* onto $\text{span}(\mathbf{u})$ is $P_{\mathbf{u}} := \mathbf{u}\mathbf{u}^* \in \text{Mat}(n, n, \mathbb{F})$, where $^*$ denotes conjugate transpose. For every $\mathbf{x} \in \mathbb{F}^n$, $P_{\mathbf{u}}\mathbf{x} = \langle \mathbf{x}, \mathbf{u} \rangle_{\mathbb{F}^n} \mathbf{u}$.

**Proposition 2.2.5.** *Properties of Rank-One Projection Matrices*
*Let $\mathbf{u} \in \mathbb{F}^n$ be a unit vector. Then:*

*i.* $\text{range}(P_{\mathbf{u}}) = \text{span}(\mathbf{u})$;
*ii.* $\ker(P_{\mathbf{u}}) = \text{span}(\mathbf{u})^\perp$;
*iii.* $P_{\mathbf{u}}^* = P_{\mathbf{u}}$;
*iv.* $P_{\mathbf{u}}^2 = P_{\mathbf{u}}$.

*Proof.*

(i) The formula $P_{\mathbf{u}}\mathbf{x} = \langle \mathbf{x}, \mathbf{u} \rangle \mathbf{u}$ gives $\text{range}(P_{\mathbf{u}}) \subseteq \text{span}(\mathbf{u})$, and $P_{\mathbf{u}}\mathbf{u} = \mathbf{u}$ gives the reverse inclusion.
(ii) The same formula gives $P_{\mathbf{u}}\mathbf{x} = 0$ exactly when $\langle \mathbf{x}, \mathbf{u} \rangle \mathbf{u} = \mathbf{0}$.
Since $\mathbf{u} \neq \mathbf{0}$, this is equivalent to $\langle \mathbf{x}, \mathbf{u} \rangle = 0$, i.e. $\mathbf{x} \in \text{span}(\mathbf{u})^\perp$.
(iii) Since $P_{\mathbf{u}} = \mathbf{u}\mathbf{u}^*$, $P_{\mathbf{u}}^* = (\mathbf{u}\mathbf{u}^*)^* = \mathbf{u}\mathbf{u}^* = P_{\mathbf{u}}$.
(iv) Because $\mathbf{u}^*\mathbf{u} = \|\mathbf{u}\|_2^2 = 1$, $P_{\mathbf{u}}^2 = \mathbf{u}(\mathbf{u}^*\mathbf{u})\mathbf{u}^* = \mathbf{u}\mathbf{u}^* = P_{\mathbf{u}}$.

*Structural Commentary*: A rank-one projection keeps the $\mathbf{u}$ direction and kills its orthogonal complement.
*Key Insight*: The outer product $\mathbf{u}\mathbf{u}^*$ encodes projection onto one unit direction. □

**Theorem 2.2.6.** *Orthogonal Projection Minimizes Distance*
*Let $\mathcal{U}$ be a finite-dimensional subspace of $\mathcal{V}$. If $\mathbf{v} \in \mathcal{V}$ and $\mathbf{u} \in \mathcal{U}$, then $\|\mathbf{v} - P_{\mathcal{U}}\mathbf{v}\| \leq \|\mathbf{v} - \mathbf{u}\|$.*
*Equality holds if and only if $\mathbf{u} = P_{\mathcal{U}}\mathbf{v}$.*

*Proof.* Set $\mathbf{p} = P_{\mathcal{U}}\mathbf{v}$, $\mathbf{r} = \mathbf{v} - \mathbf{p} \in \mathcal{U}^\perp$, and $\mathbf{h} = \mathbf{p} - \mathbf{u} \in \mathcal{U}$.
Then $\mathbf{r} \perp \mathbf{h}$, so $\|\mathbf{v} - \mathbf{u}\|^2 = \|\mathbf{r} + \mathbf{h}\|^2 = \|\mathbf{r}\|^2 + \|\mathbf{h}\|^2 \geq \|\mathbf{r}\|^2 = \|\mathbf{v} - P_{\mathcal{U}}\mathbf{v}\|^2$.
($\Rightarrow$) If equality holds, then $\|\mathbf{h}\| = 0$, hence $\mathbf{u} = \mathbf{p} = P_{\mathcal{U}}\mathbf{v}$.
($\Leftarrow$) If $\mathbf{u} = P_{\mathcal{U}}\mathbf{v}$, then equality is immediate.

*Structural Commentary*: Projection error is orthogonal to every in-subspace correction.
*Key Insight*: The projection is the unique nearest point. □

**Definition 2.2.7.** *Fourier Coefficients and Expansion*
Let $\mathcal{O} = (\mathbf{e}_1, \ldots, \mathbf{e}_m)$ be an orthonormal list and $\mathbf{v} \in \mathcal{V}$.
The *Fourier coefficients* of $\mathbf{v}$ relative to $\mathcal{O}$ are $c_j(\mathbf{v}) := \langle \mathbf{v}, \mathbf{e}_j \rangle$, $1 \leq j \leq m$.
The *Fourier expansion* of $\mathbf{v}$ onto $\mathcal{O}$ is $\widehat{\mathbf{v}}_{\mathcal{O}} := \sum_{j=1}^{m} c_j(\mathbf{v})\mathbf{e}_j$.

**Theorem 2.2.8.** *Fourier Expansion onto a Finite Orthonormal Set*
*Let $\mathcal{O} = (\mathbf{e}_1, \ldots, \mathbf{e}_m)$ be orthonormal, let $\mathcal{S} = \text{span}(\mathcal{O})$, and let $\mathbf{v} \in \mathcal{V}$. Then:*

*i.* $\widehat{\mathbf{v}}_{\mathcal{O}}$ *is the unique* $\mathbf{s} \in \mathcal{S}$ *with* $\mathbf{v} - \mathbf{s} \in \mathcal{S}^\perp$;
*ii.* $\widehat{\mathbf{v}}_{\mathcal{O}} = \arg\min_{\mathbf{s} \in \mathcal{S}} \|\mathbf{v} - \mathbf{s}\|$;
*iii.* $\|\widehat{\mathbf{v}}_{\mathcal{O}}\| \leq \|\mathbf{v}\|$.

*Proof.*

(i) For $1 \le k \le m$, $\langle \mathbf{v} - \widehat{\mathbf{v}}_{\mathcal{O}}, \mathbf{e}_k \rangle = c_k(\mathbf{v}) - \sum_{j=1}^{m} c_j(\mathbf{v})\delta_{jk} = 0$. Hence $\mathbf{v} - \widehat{\mathbf{v}}_{\mathcal{O}} \in \mathcal{S}^{\perp}$.
If $\mathbf{s} \in \mathcal{S}$ also satisfies $\mathbf{v} - \mathbf{s} \in \mathcal{S}^{\perp}$, then $\mathbf{s} - \widehat{\mathbf{v}}_{\mathcal{O}} \in \mathcal{S} \cap \mathcal{S}^{\perp} = \{\mathbf{0}\}$.
(ii) By (i), $\widehat{\mathbf{v}}_{\mathcal{O}} = P_{\mathcal{S}}\mathbf{v}$; Theorem 2.2.6 gives the minimizer.
(iii) By Theorem 1.4.5, $\|\widehat{\mathbf{v}}_{\mathcal{O}}\|^2 = \sum_{j=1}^{m} |c_j(\mathbf{v})|^2 \le \|\mathbf{v}\|^2$, hence $\|\widehat{\mathbf{v}}_{\mathcal{O}}\| \le \|\mathbf{v}\|$.

*Structural Commentary*: Fourier expansion is orthogonal projection in orthonormal coordinates.
*Key Insight*: The coefficients are obtained by pairing with the basis vectors because all cross terms vanish. □

**Theorem 2.2.9.** *Characterizing Finite Orthonormal Bases*
*Let $\mathcal{O} = (\mathbf{e}_1, \dots, \mathbf{e}_m)$ be orthonormal, let $\mathcal{S} = \operatorname{span}(\mathcal{O})$, and write $c_j(\mathbf{v})$ and $\widehat{\mathbf{v}} := \widehat{\mathbf{v}}_{\mathcal{O}}$ as in Definition 2.2.7. The following are equivalent:*

*i.* $\mathcal{S} = \mathcal{V}$.
*ii.* $\mathcal{S}^{\perp} = \{\mathbf{0}\}$.
*iii.* $\widehat{\mathbf{v}} = \mathbf{v}$ *for all* $\mathbf{v} \in \mathcal{V}$.
*iv.* $\|\widehat{\mathbf{v}}\| = \|\mathbf{v}\|$ *for all* $\mathbf{v} \in \mathcal{V}$.
*v. For all* $\mathbf{v}, \mathbf{w} \in \mathcal{V}$, $\langle \mathbf{v}, \mathbf{w} \rangle = \sum_{j=1}^{m} c_j(\mathbf{v})\overline{c_j(\mathbf{w})}$.

*Proof.*

(*i*) $\Rightarrow$ (*ii*) If $\mathcal{S} = \mathcal{V}$, then $\mathcal{S}^{\perp} = \mathcal{V}^{\perp} = \{\mathbf{0}\}$.
(*ii*) $\Rightarrow$ (*iii*) By Theorem 2.2.8, $\mathbf{v} - \widehat{\mathbf{v}} \in \mathcal{S}^{\perp} = \{\mathbf{0}\}$.
(*iii*) $\Rightarrow$ (*i*) Since $\widehat{\mathbf{v}} \in \mathcal{S}$ for all $\mathbf{v}$, condition (*iii*) gives $\mathcal{V} \subseteq \mathcal{S}$, hence $\mathcal{S} = \mathcal{V}$.
(*iii*) $\Rightarrow$ (*iv*) If $\widehat{\mathbf{v}} = \mathbf{v}$, then $\|\widehat{\mathbf{v}}\| = \|\mathbf{v}\|$.
(*iv*) $\Rightarrow$ (*iii*) The decomposition $\mathbf{v} = \widehat{\mathbf{v}} + (\mathbf{v} - \widehat{\mathbf{v}})$ is orthogonal, so $\|\mathbf{v}\|^2 = \|\widehat{\mathbf{v}}\|^2 + \|\mathbf{v} - \widehat{\mathbf{v}}\|^2$.
Together with (*iv*), this gives $\mathbf{v} = \widehat{\mathbf{v}}$.
(*iii*) $\Rightarrow$ (*v*) If $\widehat{\mathbf{v}} = \mathbf{v}$ and $\widehat{\mathbf{w}} = \mathbf{w}$, then orthonormality gives $\langle \mathbf{v}, \mathbf{w} \rangle = \sum_{j=1}^{m} c_j(\mathbf{v})\overline{c_j(\mathbf{w})}$.
(*v*) $\Rightarrow$ (*iv*) Put $\mathbf{w} = \mathbf{v}$ and use $\|\widehat{\mathbf{v}}\|^2 = \sum_{j=1}^{m} |c_j(\mathbf{v})|^2$ to get $\|\mathbf{v}\| = \|\widehat{\mathbf{v}}\|$.

*Structural Commentary*: Each condition says projection onto $\mathcal{S}$ loses no component.
*Key Insight*: Basis means lossless projection. □

**Definition 2.2.10.** *Distance to a Subspace*
Let $\mathcal{U}$ be a nonempty subset of an inner-product space $\mathcal{V}$.
The *distance* from $\mathbf{v} \in \mathcal{V}$ to $\mathcal{U}$ is $d(\mathbf{v}, \mathcal{U}) := \inf_{\mathbf{u} \in \mathcal{U}} \|\mathbf{v} - \mathbf{u}\|$.
If $\mathcal{U}$ is finite-dimensional, Theorem 2.2.6 gives $d(\mathbf{v}, \mathcal{U}) = \|\mathbf{v} - P_{\mathcal{U}}\mathbf{v}\|$.

**Perspective 2.2.11.** *Why the Finite-Dimensional Hypothesis Matters*
In $\ell^2(\mathbb{F})$, let $\mathcal{S} = \mathbb{F}^{(\mathbb{N})} = \operatorname{span}\{\mathbf{e}_j : j \ge 1\}$.
If $\mathbf{x} \in \mathcal{S}^{\perp}$, then $x_j = \langle \mathbf{x}, \mathbf{e}_j \rangle = 0 \quad (\forall j)$, so $\mathcal{S}^{\perp} = \{\mathbf{0}\}$. Since $\mathcal{S} \subsetneq \ell^2(\mathbb{F})$, $\mathcal{S} \oplus \mathcal{S}^{\perp} \neq \ell^2(\mathbb{F})$.

## 2.3 Least Squares and Metric Optimization

Least squares is projection theory as optimization: normal equations, full-rank formulas, and minimum-norm solutions.

**Principle 2.3.1.** *Least Squares as Orthogonal Projection*
In an inner-product space, the best approximation to $\mathbf{v}$ from a finite-dimensional subspace $\mathcal{U}$ is $P_{\mathcal{U}}\mathbf{v}$. The residual $\mathbf{v} - P_{\mathcal{U}}\mathbf{v}$ is orthogonal to every allowable correction inside $\mathcal{U}$.

**Definition 2.3.2.** *Matrix Least-Squares Solution*
Let $A \in \operatorname{Mat}(m, n, \mathbb{F})$ and $\mathbf{y} \in \mathbb{F}^m$.
A vector $\mathbf{x}_0 \in \mathbb{F}^n$ is a *least-squares solution* of $A\mathbf{x} = \mathbf{y}$ if $\|A\mathbf{x}_0 - \mathbf{y}\|_2 = \min_{\mathbf{x} \in \mathbb{F}^n} \|A\mathbf{x} - \mathbf{y}\|_2$.
The vector $A\mathbf{x}_0$ is the fitted vector, and $\mathbf{y} - A\mathbf{x}_0$ is the residual.

**Theorem 2.3.3.** *Normal Equations for Projection*
*Let $\mathcal{U}$ be a finite-dimensional subspace of an inner-product space $\mathcal{V}$, and suppose $\mathcal{U} = \operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_n)$.*

*For $\mathbf{v} \in \mathcal{V}$, the projection has the form $P_{\mathcal{U}}\mathbf{v} = \sum_{j=1}^{n} c_j \mathbf{u}_j$, where $(c_1, \ldots, c_n)$ solves the normal equations*

$$\begin{bmatrix} \langle \mathbf{u}_1, \mathbf{u}_1 \rangle & \langle \mathbf{u}_2, \mathbf{u}_1 \rangle & \cdots & \langle \mathbf{u}_n, \mathbf{u}_1 \rangle \\ \langle \mathbf{u}_1, \mathbf{u}_2 \rangle & \langle \mathbf{u}_2, \mathbf{u}_2 \rangle & \cdots & \langle \mathbf{u}_n, \mathbf{u}_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle \mathbf{u}_1, \mathbf{u}_n \rangle & \langle \mathbf{u}_2, \mathbf{u}_n \rangle & \cdots & \langle \mathbf{u}_n, \mathbf{u}_n \rangle \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} = \begin{bmatrix} \langle \mathbf{v}, \mathbf{u}_1 \rangle \\ \langle \mathbf{v}, \mathbf{u}_2 \rangle \\ \vdots \\ \langle \mathbf{v}, \mathbf{u}_n \rangle \end{bmatrix}.$$

*This system is consistent. If $\mathbf{u}_1, \ldots, \mathbf{u}_n$ are linearly independent, then the solution is unique.*

*Proof.* Since $P_{\mathcal{U}}\mathbf{v} \in \mathcal{U}$, write $P_{\mathcal{U}}\mathbf{v} = \sum_{j=1}^{n} c_j \mathbf{u}_j$.

The residual $\mathbf{v} - P_{\mathcal{U}}\mathbf{v}$ lies in $\mathcal{U}^{\perp}$, so for each $i$, $0 = \left\langle \mathbf{v} - \sum_{j=1}^{n} c_j \mathbf{u}_j, \mathbf{u}_i \right\rangle = \langle \mathbf{v}, \mathbf{u}_i \rangle - \sum_{j=1}^{n} c_j \langle \mathbf{u}_j, \mathbf{u}_i \rangle$.
These equations are exactly the displayed system, so consistency follows from the existence of $P_{\mathcal{U}}\mathbf{v}$.
Conversely, if $(d_1, \ldots, d_n)$ solves displayed system and $\mathbf{q} := \sum_{j=1}^{n} d_j \mathbf{u}_j$, then $\langle \mathbf{v} - \mathbf{q}, \mathbf{u}_i \rangle = 0$ for every $i$.
Since $\mathbf{u}_i$ span $\mathcal{U}$, it follows that $\mathbf{v} - \mathbf{q} \in \mathcal{U}^{\perp}$, hence $\mathbf{q} = P_{\mathcal{U}}\mathbf{v}$ by uniqueness of the decomposition. If $\mathbf{u}_1, \ldots, \mathbf{u}_n$ are linearly independent, they form a basis of $\mathcal{U}$, so the coordinates of this projection in that basis are unique.
*Structural Commentary*: Normal equations express residual orthogonality.
*Key Insight*: The projection error is perpendicular to $\mathcal{U}$. □




Figure 32: Normal equations impose $\langle \mathbf{v} - P_{\mathcal{U}}\mathbf{v}, \mathbf{u}_i \rangle = 0$ for each spanning vector.

**Theorem 2.3.4.** *Matrix Least Squares and Normal Equations*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ and $\mathbf{y} \in \mathbb{F}^m$, where $A^*$ denotes the conjugate transpose of $A$.*
*A vector $\mathbf{x}_0 \in \mathbb{F}^n$ satisfies $\|\mathbf{y} - A\mathbf{x}_0\|_2 = \min_{\mathbf{x} \in \mathbb{F}^n} \|\mathbf{y} - A\mathbf{x}\|_2$ if and only if $A^*A\mathbf{x}_0 = A^*\mathbf{y}$.*
*The normal equations are always consistent, and their solution is unique if and only if $\mathrm{rank}(A) = n$.*

*Proof.* Let $\mathcal{U} = \mathrm{col}(A)$, and write $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n]$. Since $\{A\mathbf{x} : \mathbf{x} \in \mathbb{F}^n\} = \mathcal{U}$, Theorem 2.2.6 shows that $\mathbf{x}_0$ is least-squares optimal exactly when $A\mathbf{x}_0 = P_{\mathcal{U}}\mathbf{y}$, or equivalently when $\mathbf{y} - A\mathbf{x}_0 \in \mathcal{U}^{\perp}$.

($\Rightarrow$) If $\mathbf{x}_0$ is least-squares optimal, then $\mathbf{y} - A\mathbf{x}_0 \in \mathcal{U}^{\perp}$. Hence $\langle \mathbf{y} - A\mathbf{x}_0, \mathbf{a}_j \rangle = 0$, $j = 1, \ldots, n$.
These are the entries of $A^*(\mathbf{y} - A\mathbf{x}_0)$, so $A^*(\mathbf{y} - A\mathbf{x}_0) = \mathbf{0}$ and $A^*A\mathbf{x}_0 = A^*\mathbf{y}$.

($\Leftarrow$) If $A^*A\mathbf{x}_0 = A^*\mathbf{y}$, then $A^*(\mathbf{y} - A\mathbf{x}_0) = \mathbf{0}$. Thus $\langle \mathbf{y} - A\mathbf{x}_0, \mathbf{a}_j \rangle = 0$ for every $j$, so $\mathbf{y} - A\mathbf{x}_0$ is orthogonal to every linear combination of the columns of $A$. Hence $\mathbf{y} - A\mathbf{x}_0 \in \mathcal{U}^{\perp}$.
Hence $A\mathbf{x}_0 = P_{\mathcal{U}}\mathbf{y}$, so $\mathbf{x}_0$ is least-squares optimal.

Consistency follows by choosing $\mathbf{x}_0$ such that $A\mathbf{x}_0 = P_{\mathcal{U}}\mathbf{y}$.
If $\mathbf{x}_0, \mathbf{x}_1$ solve the normal equations, then $\|A(\mathbf{x}_0 - \mathbf{x}_1)\|_2^2 = \langle \mathbf{x}_0 - \mathbf{x}_1, A^*A(\mathbf{x}_0 - \mathbf{x}_1) \rangle = 0$, so $\mathbf{x}_0 - \mathbf{x}_1 \in \ker(A)$.
Conversely, if $\mathbf{z} \in \ker(A)$ and $\mathbf{x}_0$ solves the normal equations, then $\mathbf{x}_0 + \mathbf{z}$ also solves them. Thus uniqueness is equivalent to $\ker(A) = \{\mathbf{0}\}$, equivalently $\mathrm{rank}(A) = n$.
*Structural Commentary*: Least squares is orthogonal projection onto the column space of $A$.
*Key Insight*: The normal equations say exactly that the residual is orthogonal to every column of $A$. □




Figure 33: Fitted vector and orthogonal residual: $A\mathbf{x}_0 = P_{\mathrm{col}(A)}\mathbf{y}$ and $\mathbf{r} = \mathbf{y} - A\mathbf{x}_0 \in \mathrm{col}(A)^\perp = \ker(A^*)$.

**Proposition 2.3.5.** *Least-Squares Solution Set and Minimum Norm*
*Let* $A \in \mathrm{Mat}(m,n,\mathbb{F})$, $\mathbf{y} \in \mathbb{F}^m$, *and* $\mathrm{LS}(A,\mathbf{y}) = \arg\min_{\mathbf{x}\in\mathbb{F}^n} \|A\mathbf{x}-\mathbf{y}\|_2$. *If* $\mathbf{x}_0 \in \mathrm{LS}(A,\mathbf{y})$, *then* $\mathrm{LS}(A,\mathbf{y}) = \mathbf{x}_0 + \ker(A)$.
*Also* $\mathrm{LS}(A,\mathbf{y}) \cap (\ker A)^\perp = \{\mathbf{s}\}$, *where* $\mathbf{s} = P_{(\ker A)^\perp}\mathbf{x}_0$, *and* $\mathbf{s}$ *is unique minimizer of* $\|\cdot\|_2$ *on* $\mathrm{LS}(A,\mathbf{y})$.

*Proof.* Set $\mathbf{b} = P_{\mathrm{col}(A)}\mathbf{y}$. By Theorem 2.3.4, $\mathrm{LS}(A,\mathbf{y}) = \{\mathbf{x}\in\mathbb{F}^n : A\mathbf{x} = \mathbf{b}\}$.
Thus, for $\mathbf{x}_0 \in \mathrm{LS}(A,\mathbf{y})$, $\mathrm{LS}(A,\mathbf{y}) = \{\mathbf{x} : A(\mathbf{x}-\mathbf{x}_0) = \mathbf{0}\} = \mathbf{x}_0 + \ker(A)$.
Write $\mathbf{x}_0 = \mathbf{s}+\mathbf{k}$ with $\mathbf{s} \in (\ker A)^\perp$ and $\mathbf{k} \in \ker(A)$. Since $A\mathbf{s} = A\mathbf{x}_0$, $\mathbf{s} \in \mathrm{LS}(A,\mathbf{y}) \cap (\ker A)^\perp$.
If $\mathbf{u}$ is another element of this intersection, then $\mathbf{u}-\mathbf{s} \in \ker(A) \cap (\ker A)^\perp = \{\mathbf{0}\}$.
Finally, any $\mathbf{x} \in \mathrm{LS}(A,\mathbf{y})$ has $\mathbf{x} = \mathbf{s}+\mathbf{z}$ with $\mathbf{z} \in \ker(A)$, so $\|\mathbf{x}\|_2^2 = \|\mathbf{s}\|_2^2 + \|\mathbf{z}\|_2^2 \geq \|\mathbf{s}\|_2^2$, with equality iff $\mathbf{z} = \mathbf{0}$.
*Structural Commentary*: Least-squares solutions form an affine translate of the nullspace.
*Key Insight*: The minimum-norm representative is the one with no nullspace component. □

**Corollary 2.3.6.** *Projection Formula for Full Column Rank*
*Let* $A \in \mathrm{Mat}(m,n,\mathbb{F})$ *have rank* $n$.
*Then* $A^*A$ *is invertible and the orthogonal projection onto* $\mathrm{col}(A)$ *is* $P_{\mathrm{col}(A)} = A(A^*A)^{-1}A^*$.

*Proof.* If $A^*A\mathbf{x} = \mathbf{0}$, then $\|A\mathbf{x}\|_2^2 = \langle \mathbf{x}, A^*A\mathbf{x}\rangle = 0$, so $A\mathbf{x} = \mathbf{0}$.
Since $\mathrm{rank}(A) = n$, we get $\mathbf{x} = \mathbf{0}$. Hence $A^*A$ is invertible. By Theorem 2.3.4, the least-squares solution is $\mathbf{x}_0 = (A^*A)^{-1}A^*\mathbf{y}$. The fitted vector is $P_{\mathrm{col}(A)}\mathbf{y} = A\mathbf{x}_0 = A(A^*A)^{-1}A^*\mathbf{y}$.
Since this holds for every $\mathbf{y}$, the displayed operator formula follows.
*Structural Commentary*: Full column rank makes the normal-equation coordinates unique.
*Key Insight*: Solve for coefficients, then map back by $A$. □

**Definition 2.3.7.** *Minimum Norm Solution*
Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$ and $\mathbf{y} \in \mathrm{col}(A)$.
A solution $\mathbf{s}$ of $A\mathbf{x} = \mathbf{y}$ is a *minimum norm solution* if $\|\mathbf{s}\|_2 \leq \|\mathbf{x}\|_2$ for every solution $\mathbf{x}$ of $A\mathbf{x} = \mathbf{y}$.

**Theorem 2.3.8.** *Minimum Norm Solution of a Consistent Linear System*
*Let* $A \in \mathrm{Mat}(m,n,\mathbb{F})$, $\mathbf{y} \in \mathbb{F}^m$, *and suppose* $A\mathbf{x} = \mathbf{y}$ *is consistent.*

*i. There is a unique minimum norm solution* $\mathbf{s} \in \mathbb{F}^n$, *and* $\mathbf{s} \in \mathrm{col}(A^*)$.
*ii.* $\mathbf{s}$ *is the only solution of* $A\mathbf{x} = \mathbf{y}$ *that lies in* $\mathrm{col}(A^*)$.
*iii. The system* $AA^*\mathbf{u} = \mathbf{y}$ *is consistent. If* $AA^*\mathbf{u}_0 = \mathbf{y}$, *then* $\mathbf{s} = A^*\mathbf{u}_0$.

*Proof.* First note that $\mathrm{col}(A^*) = (\ker A)^\perp$. Indeed, if $\mathbf{z} = A^*\mathbf{u}$ and $\mathbf{x} \in \ker(A)$, then $\langle \mathbf{x}, \mathbf{z}\rangle = \langle A\mathbf{x}, \mathbf{u}\rangle = 0$, so $\mathbf{z}$ is orthogonal to every vector in $\ker(A)$. Thus $\mathrm{col}(A^*) \subseteq (\ker A)^\perp$. Also,

$$\dim(\mathrm{col}(A^*)) = \mathrm{rank}(A^*) = \mathrm{rank}(A) = n - \dim(\ker(A)) = \dim((\ker A)^\perp),$$

so the two subspaces are equal. Hence $\mathbb{F}^n = \mathrm{col}(A^*) \oplus \ker(A)$ orthogonally.

(i) Choose a solution $\mathbf{x}_0$ of $A\mathbf{x} = \mathbf{y}$ and write $\mathbf{x}_0 = \mathbf{s}+\mathbf{v}$, with $\mathbf{s} \in \mathrm{col}(A^*)$ and $\mathbf{v} \in \ker(A)$.
Then $A\mathbf{s} = A\mathbf{x}_0 = \mathbf{y}$. If $\mathbf{x}$ is any solution, then $\mathbf{x}-\mathbf{s} \in \ker(A)$, so every solution has the form $\mathbf{s}+\mathbf{w}$ with $\mathbf{w} \in \ker(A)$. Since $\mathbf{s} \perp \mathbf{w}$, $\|\mathbf{s}+\mathbf{w}\|_2^2 = \|\mathbf{s}\|_2^2 + \|\mathbf{w}\|_2^2 \geq \|\mathbf{s}\|_2^2$, with equality only when $\mathbf{w} = \mathbf{0}$.
Thus $\mathbf{s}$ is the unique solution of minimum norm.
(ii) If $\mathbf{s}' \in \mathrm{col}(A^*)$ is another solution, then $\mathbf{s}'-\mathbf{s} \in \ker(A) \cap \mathrm{col}(A^*) = \{\mathbf{0}\}$, so $\mathbf{s}' = \mathbf{s}$.
(iii) Since $\mathbf{s} \in \mathrm{col}(A^*)$, write $\mathbf{s} = A^*\mathbf{z}$. Then $AA^*\mathbf{z} = A\mathbf{s} = \mathbf{y}$, so $AA^*\mathbf{u} = \mathbf{y}$ is consistent.
Conversely, if $AA^*\mathbf{u}_0 = \mathbf{y}$, then $A^*\mathbf{u}_0$ is a solution in $\mathrm{col}(A^*)$, so $A^*\mathbf{u}_0 = \mathbf{s}$ by (ii).

*Structural Commentary*: Solutions are $\mathbf{s} + \ker(A)$; $\mathrm{col}(A^*)$ meets this set orthogonally.
*Key Insight*: The minimum norm solution has no nullspace component. □




Figure 34: Among all solutions $\mathbf{s} + \ker(A)$, the unique one in $\mathrm{col}(A^*) = (\ker A)^\perp$ has minimum norm.

**Perspective 2.3.9.** *Roadmap to the Pseudoinverse*
Theorem 2.3.4 projects $\mathbf{y}$ onto $\mathrm{col}(A)$; Proposition 2.3.5 selects the minimum-norm preimage in $(\ker A)^\perp$, identified with $\mathrm{col}(A^*)$ in Theorem 2.3.8.
Definition 5.2.4 and Theorem 5.2.9 package both choices as $A^\dagger \mathbf{y}$.

## 2.4 Adjoints and Fundamental Subspaces

Adjoints express dual action in inner-product language. They become conjugate transposes in orthonormal coordinates and exchange kernels, ranges, and orthogonal complements.

**Definition 2.4.1.** *Adjoint of a Linear Map*
Let $\mathcal{V}$ and $\mathcal{W}$ be inner-product spaces and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.
A function $S : \mathcal{W} \to \mathcal{V}$ is an *adjoint* of $T$ if $\langle T\mathbf{v}, \mathbf{w}\rangle_{\mathcal{W}} = \langle \mathbf{v}, S\mathbf{w}\rangle_{\mathcal{V}}$, $\mathbf{v} \in \mathcal{V}$, $\mathbf{w} \in \mathcal{W}$.

**Perspective 2.4.2.** *Dual Map and Inner-Product Adjoint*
For $T : \mathcal{V} \to \mathcal{W}$, the pullback $\varphi \mapsto \varphi \circ T$ is the *dual map* $T' : \mathcal{W}' \to \mathcal{V}'$. Also called *operator adjoint* and written $T^\times$. It is distinct from the inner-product adjoint $T^*$ of Definition 2.4.1, which requires inner products.

**Lemma 2.4.3.** *Uniqueness of Adjoints*
*A linear map has at most one adjoint.*

*Proof.* Suppose $S_1, S_2 : \mathcal{W} \to \mathcal{V}$ are adjoints of $T$. For each $\mathbf{w} \in \mathcal{W}$ and every $\mathbf{v} \in \mathcal{V}$, $\langle \mathbf{v}, S_1\mathbf{w} - S_2\mathbf{w}\rangle_{\mathcal{V}} = 0$.
Taking $\mathbf{v} = S_1\mathbf{w} - S_2\mathbf{w}$ gives $S_1\mathbf{w} = S_2\mathbf{w}$. Hence $S_1 = S_2$.
*Structural Commentary*: Same tests force candidate adjoints to agree.
*Key Insight*: Inner products separate vectors. □

**Theorem 2.4.4.** *Existence of the Adjoint*
*Let $\mathcal{V}$ be a finite-dimensional inner-product space, let $\mathcal{W}$ be an inner-product space, and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then $T$ has a unique adjoint $T^* : \mathcal{W} \to \mathcal{V}$, and $T^*$ is linear.*

*Proof.* For each $\mathbf{w} \in \mathcal{W}$, the map $\varphi_{\mathbf{w}} : \mathcal{V} \to \mathbb{F}$ defined by $\varphi_{\mathbf{w}}(\mathbf{v}) = \langle T\mathbf{v}, \mathbf{w}\rangle_{\mathcal{W}}$ is a linear functional. By Theorem 1.6.1, there is a unique vector $T^*\mathbf{w} \in \mathcal{V}$ such that $\langle T\mathbf{v}, \mathbf{w}\rangle_{\mathcal{W}} = \langle \mathbf{v}, T^*\mathbf{w}\rangle_{\mathcal{V}}$ for every $\mathbf{v} \in \mathcal{V}$.
Thus $T^*$ is an adjoint, and it is unique by Lemma 2.4.3. For $\mathbf{w}_1, \mathbf{w}_2 \in \mathcal{W}$ and $a \in \mathbb{F}$,

$$\langle \mathbf{v}, T^*(a\mathbf{w}_1 + \mathbf{w}_2)\rangle_{\mathcal{V}} = \langle T\mathbf{v}, a\mathbf{w}_1 + \mathbf{w}_2\rangle_{\mathcal{W}} = \overline{a}\,\langle T\mathbf{v}, \mathbf{w}_1\rangle_{\mathcal{W}} + \langle T\mathbf{v}, \mathbf{w}_2\rangle_{\mathcal{W}} = \langle \mathbf{v}, aT^*\mathbf{w}_1 + T^*\mathbf{w}_2\rangle_{\mathcal{V}}\,.$$

for every $\mathbf{v} \in \mathcal{V}$. Taking the difference of the two second-slot vectors and then setting $\mathbf{v}$ equal to that difference gives $T^*(a\mathbf{w}_1 + \mathbf{w}_2) = aT^*\mathbf{w}_1 + T^*\mathbf{w}_2$.
*Structural Commentary*: Each $\mathbf{w}$ gives a Riesz-represented functional on $\mathcal{V}$.
*Key Insight*: The adjoint is pointwise Riesz representation for $T$. □




Figure 35: Adjoint relation: $\langle T\mathbf{v}, \mathbf{w}\rangle_{\mathcal{W}} = \langle \mathbf{v}, T^*\mathbf{w}\rangle_{\mathcal{V}}$.

**Perspective 2.4.5.** *Hilbert Adjoints as Riesz-Transported Dual Maps*
Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional inner-product spaces, and let $R_{\mathcal{V}} : \mathcal{V}' \to \mathcal{V}$ and $R_{\mathcal{W}} : \mathcal{W}' \to \mathcal{W}$ send each functional to its Riesz vector from Definition 1.6.2. If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then the dual map $T' : \mathcal{W}' \to \mathcal{V}'$ from Definition 4.2.3 and the Hilbert-space adjoint $T^*$ satisfy $T' = \left(R_{\mathcal{V}}\right)^{-1} \circ T^* \circ R_{\mathcal{W}}$.
Indeed, for $\varphi \in \mathcal{W}'$ and $\mathbf{v} \in \mathcal{V}$, $(T'\varphi)(\mathbf{v}) = \varphi(T\mathbf{v}) = \langle T\mathbf{v}, R_{\mathcal{W}}\varphi\rangle_{\mathcal{W}} = \langle \mathbf{v}, T^* R_{\mathcal{W}}\varphi\rangle_{\mathcal{V}}$. Thus the algebraic pullback of functionals is the same operation as moving Riesz vectors by $T^*$ and then returning to functionals.
*Structural Commentary*: The dual map belongs to the algebraic theory of Definition 4.2.3; the adjoint appears only after the Riesz identification from Theorem 1.6.1.
*Key Insight*: $T'$ and $T^*$ encode the same reversed action after functionals have been identified with vectors.




Figure 36: Riesz transports the dual map to the adjoint: $R_{\mathcal{V}} T' = T^* R_{\mathcal{W}}$.

**Definition 2.4.6.** *Conjugate Transpose*
For $A = [a_{ij}] \in \mathrm{Mat}(m, n, \mathbb{F})$, the *conjugate transpose* of $A$ is $A^* = [\overline{a_{ji}}] \in \mathrm{Mat}(n, m, \mathbb{F})$.
Over $\mathbb{R}$, this is the ordinary transpose.

**Theorem 2.4.7.** *Matrix of the Adjoint*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional inner-product spaces. Let $\beta = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be an orthonormal basis of $\mathcal{V}$ and let $\gamma = (\mathbf{f}_1, \ldots, \mathbf{f}_m)$ be an orthonormal basis of $\mathcal{W}$. If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $[T^*]_{\beta \leftarrow \gamma} = \left([T]_{\gamma \leftarrow \beta}\right)^*$.*

*Proof.* Let $A = [T]_{\gamma \leftarrow \beta}$ and $B = [T^*]_{\beta \leftarrow \gamma}$.
By Theorem 1.4.8, $B_{i,j} = \langle T^* \mathbf{f}_j, \mathbf{e}_i\rangle_{\mathcal{V}} = \overline{\langle \mathbf{e}_i, T^* \mathbf{f}_j\rangle_{\mathcal{V}}} = \overline{\langle T\mathbf{e}_i, \mathbf{f}_j\rangle_{\mathcal{W}}} = \overline{A_{j,i}}$. Thus $B = A^*$.
*Structural Commentary*: Entries swap by moving $T$ across the inner product.
*Key Insight*: In orthonormal coordinates, taking adjoints is exactly taking conjugate transposes. □

**Proposition 2.4.8.** *Basic Adjoint Identities*
*For finite-dimensional inner-product spaces and composable linear maps:*

*i.* $(aS + T)^* = \overline{a}S^* + T^*$ *for* $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ *and* $a \in \mathbb{F}$;
*ii.* $(SR)^* = R^* S^*$ *for* $R \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ *and* $S \in \mathcal{L}(\mathcal{V}, \mathcal{W})$;
*iii.* $(T^*)^* = T$;
*iv.* $I_{\mathcal{V}}^* = I_{\mathcal{V}}$.

*Proof.* Verify the defining identity; uniqueness gives equality.

(i) For $\mathbf{v} \in \mathcal{V}$, $\mathbf{w} \in \mathcal{W}$, $\langle (aS + T)\mathbf{v}, \mathbf{w}\rangle = a\langle S\mathbf{v}, \mathbf{w}\rangle + \langle T\mathbf{v}, \mathbf{w}\rangle = \langle \mathbf{v}, \overline{a}S^*\mathbf{w} + T^*\mathbf{w}\rangle$, so $(aS+T)^* = \overline{a}S^* + T^*$.
(ii) For $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{W}$, $\langle SR\mathbf{u}, \mathbf{w}\rangle = \langle R\mathbf{u}, S^*\mathbf{w}\rangle = \langle \mathbf{u}, R^* S^*\mathbf{w}\rangle$, so $(SR)^* = R^* S^*$.
(iii) Conjugating $\langle T\mathbf{v}, \mathbf{w}\rangle = \langle \mathbf{v}, T^*\mathbf{w}\rangle$ gives $\langle T^*\mathbf{w}, \mathbf{v}\rangle = \langle \mathbf{w}, T\mathbf{v}\rangle$, so $T$ is the adjoint of $T^*$.
(iv) Since $\langle I\mathbf{v}, \mathbf{w}\rangle = \langle \mathbf{v}, I\mathbf{w}\rangle$, the identity map is its own adjoint.

*Structural Commentary*: Put both sides in one inner-product slot, then use uniqueness.
*Key Insight*: Like matrix transposes, adjoints reverse products and conjugate scalars. □

**Proposition 2.4.9.** *Adjoints of Inverses and Polynomials*
*Let $T \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space. For $p(z) = \sum_{j=0}^{m} a_j z^j \in \mathbb{F}[z]$, set $p^{\#}(z) = \sum_{j=0}^{m} \overline{a_j} z^j$.*

*(i) If $T$ is invertible, then $T^*$ is invertible and $(T^{-1})^* = (T^*)^{-1}$.*
*(ii) $p(T)^* = p^{\#}(T^*)$. In particular, $p \in \mathbb{R}[z] \Rightarrow p(T)^* = p(T^*)$.*

*Proof.*

(i) Proposition 2.4.8 gives $(T^{-1})^* T^* = (TT^{-1})^* = I_{\mathcal{V}}$ and $T^*(T^{-1})^* = (T^{-1}T)^* = I_{\mathcal{V}}$.
(ii) By conjugate-linearity in scalars and product reversal, $p(T)^* = \left(\sum_{j=0}^{m} a_j T^j\right)^* = \sum_{j=0}^{m} \overline{a_j}(T^*)^j = p^{\#}(T^*)$.

*Structural Commentary*: Adjoint reverses multiplication and conjugates scalars.
*Key Insight*: Adjoints send $p(T)$ to $p^{\#}(T^*)$. □

**Theorem 2.4.10.** *Adjoints and Invariant Orthogonal Complements*
*Let $T \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space and let $\mathcal{U} \leq \mathcal{V}$.*

*(i)* $\mathcal{U} \in \mathrm{Inv}(T) \iff \mathcal{U}^\perp \in \mathrm{Inv}(T^*)$.
*(ii)* $(\mathcal{U},\mathcal{U}^\perp)$ *reduces* $T \iff \mathcal{U} \in \mathrm{Inv}(T) \cap \mathrm{Inv}(T^*)$.
*(iii) If* $(\mathcal{U},\mathcal{U}^\perp)$ *reduces* $T$, *then* $(T|_\mathcal{U})^* = (T^*)|_\mathcal{U}$.

*Proof.*

(i) Since $\mathcal{U} = (\mathcal{U}^\perp)^\perp$, $\mathcal{U} \in \mathrm{Inv}(T) \iff \langle T\mathbf{u}, \mathbf{z}\rangle = 0 \quad (\mathbf{u} \in \mathcal{U},\ \mathbf{z} \in \mathcal{U}^\perp)$
$\iff \langle \mathbf{u}, T^*\mathbf{z}\rangle = 0$, where $\mathbf{u} \in \mathcal{U},\ \mathbf{z} \in \mathcal{U}^\perp \iff \mathcal{U}^\perp \in \mathrm{Inv}(T^*)$.
(ii) By definition and then (i) applied to $T^*$, $(\mathcal{U},\mathcal{U}^\perp)$ reduces $T \iff \mathcal{U},\mathcal{U}^\perp \in \mathrm{Inv}(T) \iff \mathcal{U} \in \mathrm{Inv}(T) \cap \mathrm{Inv}(T^*)$.
(iii) By (ii), $(T^*)|_\mathcal{U}$ is well-defined. For $\mathbf{u}, \mathbf{v} \in \mathcal{U}$,
$\langle T|_\mathcal{U}\mathbf{u}, \mathbf{v}\rangle_\mathcal{U} = \langle T\mathbf{u}, \mathbf{v}\rangle_\mathcal{V} = \langle \mathbf{u}, T^*\mathbf{v}\rangle_\mathcal{V} = \langle \mathbf{u}, (T^*)|_\mathcal{U}\mathbf{v}\rangle_\mathcal{U}$, so uniqueness of adjoints on $\mathcal{U}$ gives the claim.

*Structural Commentary*: Adjoints trade invariance with orthogonal-complement invariance.
*Key Insight*: A subspace reduces $T$ precisely when it is stable under both $T$ and $T^*$. □

**Proposition 2.4.11.** *Adjoint Kernel-Range Orthogonality*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional inner-product spaces and let $T \in \mathcal{L}(\mathcal{V},\mathcal{W})$.*
*Then* $\ker(T^*) = (\mathrm{range}(T))^\perp$, $\mathrm{range}(T^*) = (\ker(T))^\perp$, *and* $\ker(T) = (\mathrm{range}(T^*))^\perp$, $\mathrm{range}(T) = (\ker(T^*))^\perp$.
*Equivalently, the nullspace of an adjoint is the orthogonal complement of the range of the original map.*

*Proof.* For $\mathbf{w} \in \mathcal{W}$,

$$\begin{aligned}\mathbf{w} \in \ker(T^*) &\iff \langle \mathbf{v}, T^*\mathbf{w}\rangle_\mathcal{V} = 0 \quad \text{for every } \mathbf{v} \in \mathcal{V}\\ &\iff \langle T\mathbf{v}, \mathbf{w}\rangle_\mathcal{W} = 0 \quad \text{for every } \mathbf{v} \in \mathcal{V}\\ &\iff \mathbf{w} \in (\mathrm{range}(T))^\perp.\end{aligned}$$

Applying this identity to $T^*$ gives $\ker(T) = (\mathrm{range}(T^*))^\perp$.
Since all spaces are finite-dimensional, the ranges and kernels are finite-dimensional subspaces.
Taking orthogonal complements and using Corollary 2.1.8 gives

$$(\ker(T^*))^\perp = ((\mathrm{range}(T))^\perp)^\perp = \mathrm{range}(T), \qquad (\ker(T))^\perp = ((\mathrm{range}(T^*))^\perp)^\perp = \mathrm{range}(T^*)$$

*Structural Commentary*: Adjoints trade ranges for kernels, and orthogonal complements reverse that trade.
*Key Insight*: A vector is killed by $T^*$ exactly when it is orthogonal to everything $T$ can produce. □

**Proposition 2.4.12.** *Adjoint Products and Algebraic Projections*
*Let $\mathcal{V}$ and $\mathcal{W}$ be finite-dimensional inner-product spaces and let $T \in \mathcal{L}(\mathcal{V},\mathcal{W})$.*

*(i)* $\ker(T^*T) = \ker(T)$ *and* $\ker(TT^*) = \ker(T^*)$.
*(ii)* $\mathrm{range}(T^*T) = \mathrm{range}(T^*)$ *and* $\mathrm{range}(TT^*) = \mathrm{range}(T)$.
*(iii) If* $\mathcal{V} = \mathcal{U} \oplus \mathcal{Z}$, *then* $\left(\rho_{\mathcal{U},\mathcal{Z}}\right)^* = \rho_{\mathcal{Z}^\perp,\mathcal{U}^\perp}$.

*Proof.*

(i) For $\mathbf{v} \in \mathcal{V}$, $\mathbf{v} \in \ker(T^*T) \iff 0 = \langle T^*T\mathbf{v}, \mathbf{v}\rangle_\mathcal{V} = \|T\mathbf{v}\|^2 \iff \mathbf{v} \in \ker(T)$.
Apply this to $T^*$ for $\ker(TT^*) = \ker(T^*)$.
(ii) The inclusions $\mathrm{range}(T^*T) \subseteq \mathrm{range}(T^*)$ and $\mathrm{range}(TT^*) \subseteq \mathrm{range}(T)$ are immediate.
By (i), rank-nullity, and Proposition 2.4.11, $\dim \mathrm{range}(T^*T) = \dim \mathcal{V} - \dim \ker(T) = \dim \mathrm{range}(T^*)$, so equality holds; the $TT^*$ identity follows similarly.
(iii) Put $P = \rho_{\mathcal{U},\mathcal{Z}}$ and $Q = \rho_{\mathcal{Z}^\perp,\mathcal{U}^\perp}$. Since $\mathcal{Z}^\perp \cap \mathcal{U}^\perp = (\mathcal{U}+\mathcal{Z})^\perp = \{\mathbf{0}\}$ and the dimensions add, $\mathcal{V} = \mathcal{Z}^\perp \oplus \mathcal{U}^\perp$.
Write $\mathbf{x} = \mathbf{u}+\mathbf{z}$ and $\mathbf{y} = \mathbf{a}+\mathbf{b}$ with $\mathbf{u} \in \mathcal{U}$, $\mathbf{z} \in \mathcal{Z}$, $\mathbf{a} \in \mathcal{Z}^\perp$, $\mathbf{b} \in \mathcal{U}^\perp$.
Then $\langle P\mathbf{x}, \mathbf{y}\rangle = \langle \mathbf{u}, \mathbf{a}+\mathbf{b}\rangle = \langle \mathbf{u}, \mathbf{a}\rangle = \langle \mathbf{u}+\mathbf{z}, \mathbf{a}\rangle = \langle \mathbf{x}, Q\mathbf{y}\rangle$, so $P^* = Q$.

*Structural Commentary*: Oblique projection adjoints swap orthogonal-complement summands.
*Key Insight*: Orthogonal projections are exactly the self-adjoint special case $\mathcal{Z} = \mathcal{U}^\perp$. □

**Corollary 2.4.13.** *Matrix Fundamental Subspaces as Orthogonal Complements*
*Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$ act as $A : \mathbb{F}^n \to \mathbb{F}^m$ with standard inner products.*
*Then* $\ker(A) = \mathrm{col}(A^*)^\perp$ *and* $\mathrm{col}(A^*) = \ker(A)^\perp$; $\ker(A^*) = \mathrm{col}(A)^\perp$ *and* $\mathrm{col}(A) = \ker(A^*)^\perp$.
*Over $\mathbb{R}$, this says* $\ker(A) \perp \mathrm{row}(A)$ *in* $\mathbb{R}^n$ *and* $\ker(A^t) \perp \mathrm{col}(A)$ *in* $\mathbb{R}^m$, *with both pairs complementary.*

*Proof.* By Theorem 2.4.7, the adjoint of $A$ is represented by $A^*$.
Proposition 2.4.11 gives $\ker(A^*) = \mathrm{col}(A)^\perp$ and $\ker(A) = \mathrm{col}(A^*)^\perp$. Taking orthogonal complements and using Corollary 2.1.8 gives the two column-space identities. Over $\mathbb{R}$, $A^* = A^t$ and $\mathrm{col}(A^t) = \mathrm{row}(A)$.
*Structural Commentary*: The four fundamental subspaces form two orthogonal-complement pairs.
*Key Insight*: Adjoints turn missing directions into kernels. □

## 2.5 Self-Adjoint Operators and Orthogonal Projections

**Definition 2.5.1.** *Self-Adjoint and Hermitian*
An operator $T \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space is *self-adjoint* if $T^* = T$. A square matrix $A$ is *Hermitian* if $A^* = A$. In orthonormal coordinates, these are the same condition by Theorem 2.4.7.

**Lemma 2.5.2.** *Zero Complex Quadratic Form*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$.*
*If $\langle T\mathbf{v}, \mathbf{v}\rangle = 0$ for every $\mathbf{v} \in \mathcal{V}$, then $T = 0$.*

*Proof.* Set $Q(\mathbf{x}) = \langle T\mathbf{x}, \mathbf{x}\rangle$.
For $\mathbf{u}, \mathbf{w} \in \mathcal{V}$, expansion gives the polarization identity for the sesquilinear form $(\mathbf{u}, \mathbf{w}) \mapsto \langle T\mathbf{u}, \mathbf{w}\rangle$:

$$\langle T\mathbf{u}, \mathbf{w}\rangle = \frac{Q(\mathbf{u}+\mathbf{w}) - Q(\mathbf{u}-\mathbf{w})}{4} + i\,\frac{Q(\mathbf{u}+i\mathbf{w}) - Q(\mathbf{u}-i\mathbf{w})}{4}.$$

The hypothesis makes every term on the right zero, so $\langle T\mathbf{u}, \mathbf{w}\rangle = 0$ for all $\mathbf{u}, \mathbf{w}$.
Taking $\mathbf{w} = T\mathbf{u}$ gives $\|T\mathbf{u}\|^2 = 0$, hence $T\mathbf{u} = \mathbf{0}$ for every $\mathbf{u}$.
*Structural Commentary*: Over $\mathbb{C}$, diagonal values of a sesquilinear form determine all mixed values.
*Key Insight*: Polarization turns the one-vector condition into a two-vector condition. □

**Proposition 2.5.3.** *Self-Adjointness from Real Quadratic Values*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$.*
*Then $T$ is self-adjoint if and only if $\langle T\mathbf{v}, \mathbf{v}\rangle \in \mathbb{R}$ for every $\mathbf{v} \in \mathcal{V}$.*

*Proof.* For each $\mathbf{v}$, $\langle (T - T^*)\mathbf{v}, \mathbf{v}\rangle = \langle T\mathbf{v}, \mathbf{v}\rangle - \langle T^*\mathbf{v}, \mathbf{v}\rangle = \langle T\mathbf{v}, \mathbf{v}\rangle - \overline{\langle T\mathbf{v}, \mathbf{v}\rangle}$.
If all values $\langle T\mathbf{v}, \mathbf{v}\rangle$ are real, this expression is zero for every $\mathbf{v}$, and Lemma 2.5.2 gives $T - T^* = 0$.
Conversely, if $T = T^*$, then $\langle T\mathbf{v}, \mathbf{v}\rangle = \langle \mathbf{v}, T\mathbf{v}\rangle = \overline{\langle T\mathbf{v}, \mathbf{v}\rangle}$, so $\langle T\mathbf{v}, \mathbf{v}\rangle$ is real.
*Structural Commentary*: The imaginary part of $\langle T\mathbf{v}, \mathbf{v}\rangle$ is exactly the diagonal trace of $T - T^*$.
*Key Insight*: Over $\mathbb{C}$, real quadratic values force the operator to equal its adjoint. □

**Proposition 2.5.4.** *Zero Quadratic Form for Self-Adjoint Operators*
*Let $T$ be a self-adjoint operator on a finite-dimensional inner-product space $\mathcal{V}$.*
*If $\langle T\mathbf{v}, \mathbf{v}\rangle = 0$ for every $\mathbf{v} \in \mathcal{V}$, then $T = 0$.*

*Proof.* If $\mathbb{F} = \mathbb{C}$, this is Lemma 2.5.2. Assume $\mathbb{F} = \mathbb{R}$.
For $\mathbf{u}, \mathbf{w} \in \mathcal{V}$, self-adjointness gives $\langle T\mathbf{w}, \mathbf{u}\rangle = \langle \mathbf{w}, T\mathbf{u}\rangle = \langle T\mathbf{u}, \mathbf{w}\rangle$, so the real polarization identity gives

$$\langle T\mathbf{u}, \mathbf{w}\rangle = \frac{\langle T(\mathbf{u}+\mathbf{w}), \mathbf{u}+\mathbf{w}\rangle - \langle T(\mathbf{u}-\mathbf{w}), \mathbf{u}-\mathbf{w}\rangle}{4} = 0.$$

Taking $\mathbf{w} = T\mathbf{u}$ gives $T\mathbf{u} = \mathbf{0}$ for every $\mathbf{u}$.
*Structural Commentary*: Self-adjointness supplies the missing symmetry needed for real polarization.
*Key Insight*: A self-adjoint operator is determined by its quadratic form. □

**Definition 2.5.5.** *Orthogonal Projection Matrix*
A square matrix $P \in \mathrm{Mat}(n, n, \mathbb{F})$ is an *orthogonal projection matrix* if $P^2 = P$ and $P^* = P$.

**Theorem 2.5.6.** *Orthogonal Projections as Self-Adjoint Idempotents*
*Let $\mathcal{V}$ be finite-dimensional and let $P \in \mathcal{L}(\mathcal{V})$.*
*Then $P$ is the orthogonal projection onto* range$(P)$ *if and only if $P^2 = P$ and $P^* = P$.*

*Proof.*
($\Rightarrow$) If $P = P_{\mathcal{U}}$ for $\mathcal{U} = \mathrm{range}(P)$, then Proposition 2.2.3 gives $P^2 = P$. For $\mathbf{v} = \mathbf{u} + \mathbf{w}$ and $\mathbf{x} = \mathbf{a} + \mathbf{b}$ with $\mathbf{u}, \mathbf{a} \in \mathcal{U}$ and $\mathbf{w}, \mathbf{b} \in \mathcal{U}^\perp$, $\langle P\mathbf{v}, \mathbf{x}\rangle = \langle \mathbf{u}, \mathbf{a}+\mathbf{b}\rangle = \langle \mathbf{u}, \mathbf{a}\rangle = \langle \mathbf{u}+\mathbf{w}, \mathbf{a}\rangle = \langle \mathbf{v}, P\mathbf{x}\rangle$, so $P^* = P$.
($\Leftarrow$) Suppose $P^2 = P$ and $P^* = P$. For each $\mathbf{v}$, $\mathbf{v} = P\mathbf{v} + (\mathbf{v} - P\mathbf{v})$ with $P\mathbf{v} \in \mathrm{range}(P)$ and $P(\mathbf{v} - P\mathbf{v}) = \mathbf{0}$. Thus the residual lies in $\ker(P)$. Since $P = P^*$, Proposition 2.4.11 gives $\ker(P) = (\mathrm{range}(P))^\perp$. Hence every $\mathbf{v}$ has the orthogonal decomposition $\mathbf{v} = P\mathbf{v} + (\mathbf{v} - P\mathbf{v})$ with first part in range$(P)$, so $P$ is exactly the orthogonal projection onto range$(P)$.

*Structural Commentary*: Idempotence splits by range and kernel; self-adjointness makes the split orthogonal.
*Key Insight*: An idempotent is orthogonal exactly when $\ker(P) = (\mathrm{range}(P))^\perp$. □

## 2.6 Isometries and Unitary Matrices

**Definition 2.6.1.** *Isometry and Isometric Isomorphism*
Let $\mathcal{V}$ and $\mathcal{W}$ be inner-product spaces and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.

(i) $T$ is an *isometry* if $\|T\mathbf{v}\| = \|\mathbf{v}\|$ for all $\mathbf{v} \in \mathcal{V}$.
(ii) $T$ is an *isometric isomorphism* if $T$ is bijective and isometric.
The spaces $\mathcal{V}$ and $\mathcal{W}$ are *isometrically isomorphic* if such a $T$ exists.

**Proposition 2.6.2.** *Basic Properties of Isometries*
*Let $\mathcal{V}$ be an inner-product space and let $S, T \in \mathcal{L}(\mathcal{V})$ be isometries.*

*i. $ST$ is an isometry;*
*ii. $T$ is injective;*
*iii. if $\mathcal{V}$ is finite-dimensional, then $T$ is invertible and $T^{-1}$ is an isometry.*

*Proof.*

(i) For every $\mathbf{v}$, $\|ST\mathbf{v}\| = \|T\mathbf{v}\| = \|\mathbf{v}\|$.
(ii) If $T\mathbf{v} = \mathbf{0}$, then $\|\mathbf{v}\| = \|T\mathbf{v}\| = 0$, so $\mathbf{v} = \mathbf{0}$. Thus $\ker(T) = \{\mathbf{0}\}$.
(iii) In finite dimensions, injectivity implies invertibility by Theorem 1.7.8.
If $\mathbf{u} = T\mathbf{v}$, then $\left\|T^{-1}\mathbf{u}\right\| = \|\mathbf{v}\| = \|T\mathbf{v}\| = \|\mathbf{u}\|$.

*Structural Commentary*: Isometries preserve length, so they cannot collapse nonzero vectors.
*Key Insight*: In finite dimensions, length preservation already forces an invertible geometry-preserving map. □

**Example 2.6.3.** *An Isometry Need Not Be Onto*
Define $T : \ell^2(\mathbb{F}) \to \ell^2(\mathbb{F})$ by $T(x_1, x_2, \ldots) = (0, x_1, x_2, \ldots)$.
For $\mathbf{x} = (x_j)_{j \geq 1}$, $\|T\mathbf{x}\|_2^2 = \sum_{j \geq 1} |x_j|^2 = \|\mathbf{x}\|_2^2$, so $T$ is an isometry.
Since every vector in $\operatorname{range}(T)$ has first coordinate 0, $(1, 0, 0, \ldots) \notin \operatorname{range}(T)$; hence $T$ is not surjective.
*Structural Commentary*: Infinite-dimensional isometries can preserve all lengths while missing whole directions.
*Key Insight*: Injective does not imply surjective outside finite dimensions.

**Theorem 2.6.4.** *Isometries Preserve Inner Products*
*Let $\mathcal{V}$ and $\mathcal{W}$ be inner-product spaces and let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.*
*Then $T$ is an isometry if and only if $\langle T\mathbf{u}, T\mathbf{v}\rangle_{\mathcal{W}} = \langle \mathbf{u}, \mathbf{v}\rangle_{\mathcal{V}}$ for every $\mathbf{u}, \mathbf{v} \in \mathcal{V}$.*

*Proof.*

(⇒) Suppose $T$ is an isometry. Then every norm appearing in the polarization identities of Theorem 1.1.23 is unchanged after applying $T$. Therefore $\langle T\mathbf{u}, T\mathbf{v}\rangle_{\mathcal{W}} = \langle \mathbf{u}, \mathbf{v}\rangle_{\mathcal{V}}$.
(⇐) Suppose $T$ preserves inner products. Then $\|T\mathbf{u}\|^2 = \langle T\mathbf{u}, T\mathbf{u}\rangle_{\mathcal{W}} = \langle \mathbf{u}, \mathbf{u}\rangle_{\mathcal{V}} = \|\mathbf{u}\|^2$, so $T$ is an isometry.

*Structural Commentary*: Polarization converts preservation of norms into preservation of inner products.
*Key Insight*: Length preservation for all vectors also preserves angle information. □

**Remark 2.6.5.** *Isometry Conventions*

(i) For linear maps between inner-product spaces, norm preservation and inner-product preservation are equivalent by Theorem 2.6.4; hence either convention defines the same isometries.
(ii) For general bilinear forms, an isometry is often required to be bijective and form-preserving; if the form is singular, preservation alone need not imply injectivity.

**Theorem 2.6.6.** *Isometries and the Adjoint Product*
*Let $\mathcal{V}$ be finite-dimensional and let $T \in \mathcal{L}(\mathcal{V})$. Then $T$ is an isometry if and only if $T^*T = I_{\mathcal{V}}$.*

*Proof.*

(⇒) Suppose $T$ is an isometry. By Theorem 2.6.4, $\langle T^*T\mathbf{u}, \mathbf{v}\rangle = \langle T\mathbf{u}, T\mathbf{v}\rangle = \langle \mathbf{u}, \mathbf{v}\rangle$ for all $\mathbf{u}, \mathbf{v}$.
Thus $\langle (T^*T - I_{\mathcal{V}})\mathbf{u}, \mathbf{v}\rangle = 0$ for every $\mathbf{v}$, so $T^*T\mathbf{u} = \mathbf{u}$ for every $\mathbf{u}$.
(⇐) Suppose $T^*T = I_{\mathcal{V}}$. Then $\|T\mathbf{u}\|^2 = \langle T\mathbf{u}, T\mathbf{u}\rangle = \langle T^*T\mathbf{u}, \mathbf{u}\rangle = \|\mathbf{u}\|^2$ for every $\mathbf{u}$, so $T$ is an isometry.

*Structural Commentary*: The operator $T^*T$ records inner products after applying $T$.
*Key Insight*: $T$ is an isometry exactly when this Gram operator is the identity. □

**Definition 2.6.7.** *Unitary and Orthogonal*
An operator $U \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space is *unitary* if $U^*U = UU^* = I_{\mathcal{V}}$.
A matrix $U \in \operatorname{Mat}(n, n, \mathbb{F})$ is *unitary* if $U^*U = I_n$. If $\mathbb{F} = \mathbb{R}$, a unitary matrix is called *orthogonal*.




Figure 37: Unitary invariance: $\|U\mathbf{x}\| = \|\mathbf{x}\|$ and $\langle U\mathbf{x}, U\mathbf{y}\rangle = \langle \mathbf{x}, \mathbf{y}\rangle$.

**Theorem 2.6.8.** *Equivalent Conditions for a Unitary Matrix*
*Let $U = [\mathbf{u}_1 \ \cdots \ \mathbf{u}_n] \in \mathrm{Mat}(n, n, \mathbb{F})$. The following are equivalent:*

*i. $U$ is unitary, i.e. $U^*U = I_n$;*
*ii. the columns $\mathbf{u}_1, \ldots, \mathbf{u}_n$ are orthonormal in $\mathbb{F}^n$;*
*iii. $UU^* = I_n$;*
*iv. $U^*$ is unitary;*
*v. the rows of $U$ are orthonormal;*
*vi. $U$ is invertible and $U^{-1} = U^*$;*
*vii. $\|U\mathbf{x}\|_2 = \|\mathbf{x}\|_2$ for every $\mathbf{x} \in \mathbb{F}^n$;*
*viii. $\langle U\mathbf{x}, U\mathbf{y}\rangle_{\mathbb{F}^n} = \langle \mathbf{x}, \mathbf{y}\rangle_{\mathbb{F}^n}$ for every $\mathbf{x}, \mathbf{y} \in \mathbb{F}^n$.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)*. Since $(U^*U)_{ij} = \mathbf{u}_i^*\mathbf{u}_j = \langle \mathbf{u}_j, \mathbf{u}_i\rangle_{\mathbb{F}^n}$, equation $U^*U = I_n$ states that columns of $U$ are orthonormal.
*(ii)* $\Rightarrow$ *(iii)*. Orthonormal columns give $U^*U = I_n$. A left inverse of a square matrix is also a right inverse by Theorem 3.4.2, so $UU^* = I_n$.
*(iii)* $\Rightarrow$ *(iv)*. The equation $UU^* = I_n$ is $(U^*)^*U^* = I_n$, so $U^*$ is unitary.
*(iv)* $\Rightarrow$ *(v)*. Apply *(i)* $\Rightarrow$ *(ii)* to $U^*$; the columns of $U^*$ are the conjugates of the rows of $U$.
*(v)* $\Rightarrow$ *(vi)*. Orthonormal rows give $UU^* = I_n$. Thus $U^*$ is the inverse of $U$, so $U^{-1} = U^*$.
*(vi)* $\Rightarrow$ *(vii)*. If $U^{-1} = U^*$, then $U^*U = I_n$. Therefore $\|U\mathbf{x}\|_2^2 = \langle U\mathbf{x}, U\mathbf{x}\rangle_{\mathbb{F}^n} = \langle U^*U\mathbf{x}, \mathbf{x}\rangle_{\mathbb{F}^n} = \|\mathbf{x}\|_2^2$ for every $\mathbf{x} \in \mathbb{F}^n$. Since both norms are nonnegative, $\|U\mathbf{x}\|_2 = \|\mathbf{x}\|_2$.
*(vii)* $\Rightarrow$ *(viii)*. Apply Theorem 2.6.4 to $\mathbf{x} \mapsto U\mathbf{x}$.
*(viii)* $\Rightarrow$ *(i)*. For every $\mathbf{x}, \mathbf{y} \in \mathbb{F}^n$, $\langle (U^*U - I_n)\mathbf{x}, \mathbf{y}\rangle_{\mathbb{F}^n} = \langle U\mathbf{x}, U\mathbf{y}\rangle_{\mathbb{F}^n} - \langle \mathbf{x}, \mathbf{y}\rangle_{\mathbb{F}^n} = 0$. Taking $\mathbf{y} = (U^*U - I_n)\mathbf{x}$ shows $(U^*U - I_n)\mathbf{x} = \mathbf{0}$ for every $\mathbf{x}$, so $U^*U = I_n$.

*Structural Commentary*: A unitary matrix is an isometry written in orthonormal coordinates.
*Key Insight*: Columns, rows, inverses, norms, and inner products are different faces of the same condition. □

**Definition 2.6.9.** *Unitary and Permutation Similarity*
Square matrices $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$ are *unitarily similar* if $A = UBU^*$ for some unitary $U$. They are *orthogonally similar* if $\mathbb{F} = \mathbb{R}$ and $U$ can be chosen real orthogonal.
A *permutation matrix* is a square matrix with exactly one entry equal to 1 in each row and each column, and all other entries equal to 0. Matrices are *permutation similar* if $A = PBP^*$ for some permutation matrix $P$.

**Theorem 2.6.10.** *Unitary Coordinate Form of an Orthogonal Projection*
*Let $P \in \mathrm{Mat}(n, n, \mathbb{F})$ be an orthogonal projection matrix and let $r = \mathrm{rank}(P)$. Then $P$ is unitarily similar to $I_r \oplus 0_{n-r}$. Equivalently, there is a unitary matrix $U$ such that*

$$U^*PU = \begin{bmatrix} I_r & 0 \\ 0 & 0_{n-r} \end{bmatrix}.$$

*Proof.* Let $\mathcal{U} = \mathrm{col}(P)$. Identifying $P$ with its induced operator on $\mathbb{F}^n$, Theorem 2.5.6 says $P$ is the orthogonal projection onto $\mathcal{U}$, so $P$ is the identity on $\mathcal{U}$ and zero on $\mathcal{U}^\perp$. Choose an orthonormal basis $\mathbf{u}_1, \ldots, \mathbf{u}_r$ of $\mathcal{U}$ and extend it to an orthonormal basis $\mathbf{u}_1, \ldots, \mathbf{u}_n$ of $\mathbb{F}^n$. Let $U = [\mathbf{u}_1 \ \cdots \ \mathbf{u}_n]$.
Then $U$ is unitary. For $j \le r$, $P\mathbf{u}_j = \mathbf{u}_j$, and for $j > r$, $\mathbf{u}_j \in \mathcal{U}^\perp$ so $P\mathbf{u}_j = \mathbf{0}$. Therefore the $j$th column of $U^*PU$ is the coordinate column of $P\mathbf{u}_j$ in this orthonormal basis, giving the displayed block diagonal form.
*Structural Commentary*: Orthogonal projection splits the space into range and orthogonal kernel.
*Key Insight*: In a basis adapted to $\mathrm{col}(P) \oplus \mathrm{col}(P)^\perp$, projection keeps the first coordinates and kills the rest. □




Figure 38: Adapted basis: $U^*PU = I_r \oplus 0_{n-r}$, identity on $\mathrm{col}(P)$ and zero on $\mathrm{col}(P)^\perp$.

**Proposition 2.6.11.** *Algebra of Unitary Matrices*
*Let $U, V \in \mathrm{Mat}(n, n, \mathbb{F})$ and $W \in \mathrm{Mat}(m, m, \mathbb{F})$, where $V \oplus W$ denotes the block diagonal matrix with blocks $V$ and $W$.*

*i. If $U$ and $V$ are unitary, then $UV$ is unitary.*
*ii. $V \oplus W$ is unitary if and only if $V$ and $W$ are unitary.*

*Proof.*

(i) Compute $(UV)^*(UV) = V^*U^*UV = V^*V = I_n$.
(ii) Since $(V \oplus W)^*(V \oplus W) = V^*V \oplus W^*W$, this product equals $I_{n+m}$ exactly when $V^*V = I_n$ and $W^*W = I_m$.

*Structural Commentary*: Unitary structure is stable under composition and blockwise orthogonal sums.
*Key Insight*: The equation $U^*U = I$ makes every verification a short adjoint calculation. □

**Theorem 2.6.12.** *Extending Isometric Columns to a Unitary Matrix*
*If $X \in \mathrm{Mat}(m, n, \mathbb{F})$ and $X^*X = I_n$, then there exists $X' \in \mathrm{Mat}(m, m-n, \mathbb{F})$ such that $[X\ X']$ is unitary.*

*Proof.* Write $X = [\mathbf{x}_1\ \cdots\ \mathbf{x}_n]$. Since $(X^*X)_{ij} = \langle \mathbf{x}_j, \mathbf{x}_i \rangle_{\mathbb{F}^m}$, the condition $X^*X = I_n$ says $\mathbf{x}_1, \ldots, \mathbf{x}_n$ are orthonormal in $\mathbb{F}^m$. By Corollary 1.3.13, extend this list to an orthonormal basis $\mathbf{x}_1, \ldots, \mathbf{x}_m$ of $\mathbb{F}^m$.
Let $X' = [\mathbf{x}_{n+1}\ \cdots\ \mathbf{x}_m]$. Then the columns of $[X\ X']$ are orthonormal, so $[X\ X']$ is unitary.
*Structural Commentary*: Isometric columns are a partial orthonormal basis.
*Key Insight*: Complete the columns by extending the orthonormal list. □

**Proposition 2.6.13.** *Permutation Matrices Are Orthogonal*
*If $P = [\mathbf{e}_{\sigma(1)}\ \cdots\ \mathbf{e}_{\sigma(n)}]$ is a permutation matrix, then $P^*P = PP^* = I_n$. Hence $P$ is real orthogonal.*

*Proof.* The columns of $P$ are a rearrangement of the standard orthonormal basis of $\mathbb{R}^n$, hence they are orthonormal. Theorem 2.6.8 gives $P^*P = PP^* = I_n$.
*Structural Commentary*: A permutation matrix only reorders the standard basis.
*Key Insight*: Reordering orthonormal vectors preserves orthonormality. □

## 2.7 Householder Reflections

**Definition 2.7.1.** *Householder Matrix*
Let $\mathbf{w} \in \mathbb{F}^n$ be nonzero and set $\mathbf{u} = \mathbf{w}/\|\mathbf{w}\|_2$.
The *Householder matrix* associated to $\mathbf{w}$ is $U_{\mathbf{w}} := I_n - 2P_{\mathbf{u}} = I_n - 2\frac{\mathbf{w}\mathbf{w}^*}{\|\mathbf{w}\|_2^2}$.
It acts by $\mathbf{x} \mapsto U_{\mathbf{w}}\mathbf{x} = \mathbf{x} - 2\langle \mathbf{x}, \mathbf{u} \rangle_{\mathbb{F}^n} \mathbf{u}$.




Figure 39: A Householder matrix fixes $\mathbf{w}^\perp$ and flips the $\mathrm{span}(\mathbf{w})$ component.

**Theorem 2.7.2.** *Householder Matrices Are Unitary Reflections*
*Every Householder matrix $U_{\mathbf{w}}$ is unitary, Hermitian, and involutive: $U_{\mathbf{w}}^* U_{\mathbf{w}} = I_n$, $U_{\mathbf{w}}^* = U_{\mathbf{w}}$, $U_{\mathbf{w}}^2 = I_n$.*
*If $\mathbf{w} \in \mathbb{R}^n$, then $U_{\mathbf{w}}$ is real orthogonal and symmetric.*

*Proof.* Let $\mathbf{u} = \mathbf{w}/\|\mathbf{w}\|_2$. By Proposition 2.2.5, $P_{\mathbf{u}}^* = P_{\mathbf{u}}$ and $P_{\mathbf{u}}^2 = P_{\mathbf{u}}$.
Therefore $U_{\mathbf{w}}^* = I_n - 2P_{\mathbf{u}} = U_{\mathbf{w}}$, $U_{\mathbf{w}}^2 = (I_n - 2P_{\mathbf{u}})^2 = I_n - 4P_{\mathbf{u}} + 4P_{\mathbf{u}}^2 = I_n$.
Thus $U_{\mathbf{w}}^* U_{\mathbf{w}} = U_{\mathbf{w}}^2 = I_n$, so $U_{\mathbf{w}}$ is unitary. The real case is the same statement with conjugation removed.
*Structural Commentary*: A Householder matrix reflects across an orthogonal hyperplane.
*Key Insight*: The identity $P_{\mathbf{u}}^2 = P_{\mathbf{u}}$ makes $(I - 2P_{\mathbf{u}})^2 = I$. □

**Theorem 2.7.3.** *Householder Map Between Equal-Norm Vectors*
*Let $\mathbf{x}, \mathbf{y} \in \mathbb{F}^n$ satisfy $\|\mathbf{x}\|_2 = \|\mathbf{y}\|_2 \neq 0$. Define $\sigma = \begin{cases} 1, & \langle \mathbf{x}, \mathbf{y} \rangle_{\mathbb{F}^n} = 0, \\ -\dfrac{\overline{\langle \mathbf{x}, \mathbf{y} \rangle_{\mathbb{F}^n}}}{|\langle \mathbf{x}, \mathbf{y} \rangle_{\mathbb{F}^n}|}, & \langle \mathbf{x}, \mathbf{y} \rangle_{\mathbb{F}^n} \neq 0, \end{cases}$, where $\mathbf{w} = \mathbf{y} - \sigma\mathbf{x}$.*
*Then $\mathbf{w} \neq \mathbf{0}$, $|\sigma| = 1$, and $\sigma U_{\mathbf{w}} \mathbf{x} = \mathbf{y}$.*

*Proof.* Clearly $|\sigma| = 1$. Let $z = \langle \mathbf{x}, \mathbf{y} \rangle$ and $r = \|\mathbf{x}\|_2 = \|\mathbf{y}\|_2$. By construction, $\sigma z = -|z|$, with the convention that this equals 0 when $z = 0$. Hence $\|\mathbf{w}\|_2^2 = \|\mathbf{y} - \sigma\mathbf{x}\|_2^2 = 2r^2 - \sigma z - \overline{\sigma z} = 2\left(r^2 + |z|\right) > 0$, so $\mathbf{w} \neq \mathbf{0}$.
Since $\mathbf{y} = \mathbf{w} + \sigma\mathbf{x}$, $\sigma U_{\mathbf{w}} \mathbf{x} = \sigma\mathbf{x} - 2\sigma \frac{\langle \mathbf{x}, \mathbf{w} \rangle}{\|\mathbf{w}\|_2^2} \mathbf{w} = \mathbf{y} - \left(1 + 2\sigma \frac{\langle \mathbf{x}, \mathbf{w} \rangle}{\|\mathbf{w}\|_2^2}\right) \mathbf{w}$.
But $\langle \mathbf{x}, \mathbf{w} \rangle = z - \overline{\sigma} r^2$, so $2\sigma \langle \mathbf{x}, \mathbf{w} \rangle = 2\sigma z - 2r^2 = -2(r^2 + |z|) = -\|\mathbf{w}\|_2^2$.
Coefficient of $\mathbf{w}$ is hence zero, proving $\sigma U_{\mathbf{w}} \mathbf{x} = \mathbf{y}$.
*Structural Commentary*: The scalar $\sigma$ chooses the phase so that $\mathbf{y} - \sigma\mathbf{x}$ gives a stable reflection direction.
*Key Insight*: Householder reflections connect equal-norm vectors up to a unit scalar. □

**Corollary 2.7.4.** *Householder Unitary Annihilator*
*Let $\mathbf{x} = [x_1 \ \cdots \ x_n]^t \in \mathbb{F}^n$ be nonzero. Define $\sigma = \begin{cases} 1, & x_1 = 0, \\ -\dfrac{\overline{x_1}}{|x_1|}, & x_1 \neq 0, \end{cases}$, where $\mathbf{w} = \|\mathbf{x}\|_2 \mathbf{e}_1 - \sigma\mathbf{x}$.*
*Then $\sigma U_{\mathbf{w}} \mathbf{x} = \|\mathbf{x}\|_2 \mathbf{e}_1$. If $\mathbf{x}$ is a unit vector, then $\overline{\sigma} U_{\mathbf{w}}$ is unitary and has $\mathbf{x}$ as its first column.*

*Proof.* Apply Theorem 2.7.3 with $\mathbf{y} = \|\mathbf{x}\|_2 \mathbf{e}_1$, noting that $\langle \mathbf{x}, \mathbf{y} \rangle = x_1 \|\mathbf{x}\|_2$. If $\|\mathbf{x}\|_2 = 1$, then $(\sigma U_{\mathbf{w}})\mathbf{x} = \mathbf{e}_1$, so $(\sigma U_{\mathbf{w}})^* \mathbf{e}_1 = \mathbf{x}$. Since $(\sigma U_{\mathbf{w}})^* = \overline{\sigma} U_{\mathbf{w}}$, this says the first column of $\overline{\sigma} U_{\mathbf{w}}$ is $\mathbf{x}$.
*Structural Commentary*: The reflector concentrates all mass of $\mathbf{x}$ into the first coordinate.
*Key Insight*: Householder matrices produce explicit unitary matrices with a prescribed first column. □

# 3 Bilinear and Quadratic Forms

## 3.1 Bilinear Form Types and Congruence Invariants

Metric vector spaces keep the algebraic part of inner-product geometry and drop positivity. Throughout this subsection $\mathbb{F}$ is an arbitrary field and vector spaces are finite-dimensional unless explicitly stated otherwise. General bilinear pairings, their left and right radicals, representing matrices, and basis changes are established in Definition 4.4.1, Definition 4.4.3, Proposition 4.4.4, and Proposition 4.4.9. Here the two input spaces coincide, and the fixed bilinear form is written $\langle \mathbf{x}, \mathbf{y} \rangle$.

**Perspective 3.1.1.** *Square Pairings*
Henceforth $B : \mathcal{V} \times \mathcal{V} \to \mathbb{F}$ is a square pairing, usually symmetric, skew-symmetric, or alternating.
In these cases $\mathrm{rad}_L(B) = \mathrm{rad}_R(B)$, so one radical and one orthogonality notation suffice.

**Definition 3.1.2.** *Symmetric, Skew-Symmetric, and Alternating Forms*
Let $\mathcal{V}$ be a vector space over $\mathbb{F}$ and let $B : \mathcal{V} \times \mathcal{V} \to \mathbb{F}$ be bilinear.

(i) $B$ is a *bilinear form* on $\mathcal{V}$; when fixed, write $B(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, \mathbf{y} \rangle$.
(ii) $\mathrm{Sym}(B)$ means $B(\mathbf{x}, \mathbf{y}) = B(\mathbf{y}, \mathbf{x})$ for all $\mathbf{x}, \mathbf{y} \in \mathcal{V}$; then $B$ is *symmetric*.
(iii) $\mathrm{Skew}(B)$ means $B(\mathbf{x}, \mathbf{y}) = -B(\mathbf{y}, \mathbf{x})$ for all $\mathbf{x}, \mathbf{y} \in \mathcal{V}$; then $B$ is *skew-symmetric*.
(iv) $\mathrm{Alt}(B)$ means $B(\mathbf{x}, \mathbf{x}) = 0$ for all $\mathbf{x} \in \mathcal{V}$; then $B$ is *alternating*.
(v) A *metric vector space* is a pair $(\mathcal{V}, B)$ with $\mathrm{Sym}(B)$, $\mathrm{Skew}(B)$, or $\mathrm{Alt}(B)$.
The symmetric case is an *orthogonal geometry*; the alternating case is a *symplectic geometry*.

**Proposition 3.1.3.** *Characteristic Controls the Form Types*
*Let $B(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, \mathbf{y} \rangle$ be a bilinear form on $\mathcal{V}$.*

*(i)* $\mathrm{Alt}(B) \Rightarrow \mathrm{Skew}(B)$.

*(ii) If* $\operatorname{char}(\mathbb{F}) \neq 2$, *then* $\operatorname{Alt}(B) \iff \operatorname{Skew}(B)$, *and* $\operatorname{Sym}(B) \wedge \operatorname{Alt}(B) \Rightarrow B = 0$.
*(iii) If* $\operatorname{char}(\mathbb{F}) = 2$, *then* $\operatorname{Sym}(B) \iff \operatorname{Skew}(B)$ *and* $\operatorname{Alt}(B) \Rightarrow \operatorname{Sym}(B)$.

*Proof.*

(i) If $\operatorname{Alt}(B)$, then $0 = \langle \mathbf{x}+\mathbf{y}, \mathbf{x}+\mathbf{y}\rangle = \langle \mathbf{x},\mathbf{y}\rangle + \langle \mathbf{y},\mathbf{x}\rangle$, so $\langle \mathbf{x},\mathbf{y}\rangle = -\langle \mathbf{y},\mathbf{x}\rangle$.
(ii) Assume $\operatorname{char}(\mathbb{F}) \neq 2$. By (i), $\operatorname{Alt}(B) \Rightarrow \operatorname{Skew}(B)$. Conversely, $\operatorname{Skew}(B)$ gives $\langle \mathbf{x},\mathbf{x}\rangle = -\langle \mathbf{x},\mathbf{x}\rangle$, hence $2\langle \mathbf{x},\mathbf{x}\rangle = 0$ and $\operatorname{Alt}(B)$. If also $\operatorname{Sym}(B)$ and $\operatorname{Alt}(B)$, then $\langle \mathbf{x},\mathbf{y}\rangle = -\langle \mathbf{x},\mathbf{y}\rangle$, so $\langle \mathbf{x},\mathbf{y}\rangle = 0$ for all $\mathbf{x},\mathbf{y}$.
(iii) If $\operatorname{char}(\mathbb{F}) = 2$, then $-1 = 1$, so $\operatorname{Sym}(B) \iff \operatorname{Skew}(B)$.
Moreover $\operatorname{Alt}(B) \Rightarrow \operatorname{Skew}(B)$ by (i), hence $\operatorname{Alt}(B) \Rightarrow \operatorname{Sym}(B)$.

*Structural Commentary*: Skew-symmetry collapses by characteristic.
*Key Insight*: The equation $\langle \mathbf{x}+\mathbf{y}, \mathbf{x}+\mathbf{y}\rangle = 0$ polarizes alternation. □

**Definition 3.1.4.** *Matrix, Rank, and Discriminant of a Form*
Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be an ordered basis of a metric vector space $\mathcal{V}$.

(i) The *matrix of the form* in $\beta$ is $M_\beta := [\langle \mathbf{v}_i, \mathbf{v}_j\rangle]_{i,j}$.
(ii) The *rank* of the form is $\operatorname{rank}(M_\beta)$, independent of $\beta$.
(iii) The *discriminant set* is $\Delta(\mathcal{V}) := \{r^2 \det(M_\beta) : r \in \mathbb{F}^\times\}$.
(iv) If the form is nonsingular, then $\Delta(\mathcal{V})$ is the square class of $\det(M_\beta)$ in $\mathbb{F}^\times/(\mathbb{F}^\times)^2$.

**Proposition 3.1.5.** *Matrices Represent Forms up to Congruence*
*Let* $\beta, \gamma$ *be ordered bases of* $\mathcal{V}$ *and put* $C = C_{\beta\leftarrow\gamma}$.

*(i)* $\langle \mathbf{x},\mathbf{y}\rangle = [\mathbf{x}]_\beta^t M_\beta [\mathbf{y}]_\beta$.
*(ii)* $M_\gamma = C^t M_\beta C$.
*(iii) Matrices* $A, N$ *represent the same form on* $\mathcal{V}$ *if and only if* $N = P^t A P$ *for some invertible* $P$.

*Proof.*

(i) If $\mathbf{x} = \sum_i a_i \mathbf{v}_i$ and $\mathbf{y} = \sum_j b_j \mathbf{v}_j$, then $\langle \mathbf{x},\mathbf{y}\rangle = \sum_{i,j} a_i b_j \langle \mathbf{v}_i, \mathbf{v}_j\rangle = [\mathbf{x}]_\beta^t M_\beta [\mathbf{y}]_\beta$.
(ii) Since $[\mathbf{x}]_\beta = C[\mathbf{x}]_\gamma$ and $[\mathbf{y}]_\beta = C[\mathbf{y}]_\gamma$, $[\mathbf{x}]_\gamma^t M_\gamma [\mathbf{y}]_\gamma = \langle \mathbf{x},\mathbf{y}\rangle = [\mathbf{x}]_\gamma^t C^t M_\beta C [\mathbf{y}]_\gamma$, so $M_\gamma = C^t M_\beta C$.
(iii) If $A = M_\beta$ and $N = M_\gamma$ represent one form, then (ii) gives $N = C_{\beta\leftarrow\gamma}^t A C_{\beta\leftarrow\gamma}$.
Conversely, suppose $N = P^t A P$ with $P = [p_{ij}]$ invertible and $A = M_\beta$.
Define $\gamma = (\mathbf{w}_1, \dots, \mathbf{w}_n)$ by $\mathbf{w}_j = \sum_i p_{ij} \mathbf{v}_i$. Then $C_{\beta\leftarrow\gamma} = P$, so (ii) gives $M_\gamma = P^t A P = N$.

*Structural Commentary*: Congruence is basis change for forms.
*Key Insight*: Basis change acts in both slots: $M \mapsto P^t M P$. □

## 3.2 Quadratic Forms and Polar Forms

**Definition 3.2.1.** *Quadratic Forms and Polar Forms*
A *quadratic form* on $\mathcal{V}$ is a function $Q : \mathcal{V} \to \mathbb{F}$ such that:

(i) $Q(a\mathbf{v}) = a^2 Q(\mathbf{v})$ for all $a \in \mathbb{F}$ and $\mathbf{v} \in \mathcal{V}$;
(ii) the *polar form* $B_Q(\mathbf{u},\mathbf{v}) := Q(\mathbf{u}+\mathbf{v}) - Q(\mathbf{u}) - Q(\mathbf{v})$ is bilinear.

**Proposition 3.2.2.** *Quadratic Forms and Symmetric Bilinear Forms*
*For every quadratic form* $Q$, *its polar form* $B_Q$ *is symmetric.*

*If* $\operatorname{char}(\mathbb{F}) \neq 2$, *maps* $\{B : \operatorname{Sym}(B)\} \underset{Q\mapsto B_Q}{\overset{B\mapsto Q_B}{\rightleftarrows}} \{Q : Q \text{ quadratic}\}$ *and* $Q_B(\mathbf{x}) := \frac{1}{2}B(\mathbf{x},\mathbf{x})$ *are inverse correspondences between symmetric bilinear forms and quadratic forms.*

*Proof.* First, $B_Q(\mathbf{u},\mathbf{v}) = Q(\mathbf{u}+\mathbf{v}) - Q(\mathbf{u}) - Q(\mathbf{v}) = B_Q(\mathbf{v},\mathbf{u})$, so $B_Q$ is symmetric.
($\Rightarrow$) If $\operatorname{Sym}(B)$, then $Q_B(a\mathbf{x}) = a^2 Q_B(\mathbf{x})$ and $B_{Q_B}(\mathbf{u},\mathbf{v}) = \frac{1}{2}\big(B(\mathbf{u}+\mathbf{v},\mathbf{u}+\mathbf{v}) - B(\mathbf{u},\mathbf{u}) - B(\mathbf{v},\mathbf{v})\big) = B(\mathbf{u},\mathbf{v})$.
($\Leftarrow$) If $Q$ is quadratic, then $Q_{B_Q}(\mathbf{x}) = \frac{1}{2}B_Q(\mathbf{x},\mathbf{x}) = \frac{1}{2}\big(Q(2\mathbf{x}) - 2Q(\mathbf{x})\big) = Q(\mathbf{x})$.

*Structural Commentary*: Away from characteristic 2, quadratic and symmetric bilinear data coincide.
*Key Insight*: Dividing by 2 recovers diagonal data from polar data. □

**Proposition 3.2.3.** *Polar Forms in Characteristic Two*
*If* $\operatorname{char}(\mathbb{F}) = 2$, *then* $\operatorname{Alt}(B_Q)$ *for every quadratic form* $Q$ *on* $\mathcal{V}$.

*Proof.* For $\mathbf{x} \in \mathcal{V}$, $B_Q(\mathbf{x},\mathbf{x}) = Q(2\mathbf{x}) - Q(\mathbf{x}) - Q(\mathbf{x}) = Q(\mathbf{0}) = 0$. Thus $\operatorname{Alt}(B_Q)$.
*Structural Commentary*: In characteristic 2, every polar form is alternating.
*Key Insight*: The diagonal of the polar form vanishes automatically because both $2\mathbf{x}$ and $2Q(\mathbf{x})$ vanish. □

**Example 3.2.4.** *Characteristic Two Loses Diagonal Data*
Assume $\text{char}(\mathbb{F}) = 2$. On $\mathbb{F}^2$, let $Q_0(x,y) = xy$ and $Q_1(x,y) = xy + x^2$.
Then $B_{Q_0}((x,y),(u,v)) = xv + uy = B_{Q_1}((x,y),(u,v))$, and $Q_0(1,0) = 0 \neq 1 = Q_1(1,0)$.
Indeed, the added square has zero polar contribution, where $(x+u)^2 - x^2 - u^2 = 2xu = 0$.
More generally, for any linear $L : \mathcal{V} \to \mathbb{F}$, $Q$ and $Q + L^2$ have the same polar form.
*Structural Commentary*: In characteristic 2, squares are invisible to polarization.
*Key Insight*: Characteristic 2 quadratic forms need data beyond their polar forms.

**Example 3.2.5.** *Alternating versus Nonalternating Symmetric Forms*
Assume $\text{char}(\mathbb{F}) = 2$.
The symmetric matrix $H = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ defines a nonsingular alternating form on $\mathbb{F}^2$, since $\begin{bmatrix} x & y \end{bmatrix} H \begin{bmatrix} x \\ y \end{bmatrix} = 2xy = 0$.
Thus $H$ is not congruent to any nonzero diagonal form: alternating means all self-pairings vanish, while a nonzero diagonal form has some basis vector with nonzero self-pairing. By contrast, [1] is symmetric and nonalternating.
*Structural Commentary*: In characteristic 2, symmetric forms split into alternating and nonalternating parts.
*Key Insight*: Nonsingular alternating blocks are invisible on the diagonal.

## 3.3 Orthogonality, Radicals, and Nonsingular Metric Spaces

**Definition 3.3.1.** *Orthogonality, Isotropy, and Radical*
Let $\mathcal{V}$ be a metric vector space.

(i) $\mathbf{x} \perp \mathbf{y}$ means $\langle \mathbf{x}, \mathbf{y} \rangle = 0$; then $\mathbf{x}, \mathbf{y}$ are *orthogonal*. For $S \subseteq \mathcal{V}$, $S^\perp := \{\mathbf{v} \in \mathcal{V} : \langle \mathbf{v}, \mathbf{s} \rangle = 0 \ \forall \mathbf{s} \in S\}$
(ii) $\mathcal{S} \leq \mathcal{V}$ is *totally isotropic* or *totally degenerate* if $\mathcal{S} \subseteq \mathcal{S}^\perp$, and *nonsingular as a subspace* if $\mathcal{S} \cap \mathcal{S}^\perp = \{\mathbf{0}\}$.
(iii) $\mathbf{x} \neq \mathbf{0}$ is *isotropic* or *null* if $\langle \mathbf{x}, \mathbf{x} \rangle = 0$, and *nonisotropic* otherwise.
The space $\mathcal{V}$ is *isotropic* if such an $\mathbf{x}$ exists, and *anisotropic* otherwise.
(iv) $\mathbf{x}$ is *degenerate* if $\mathbf{x} \perp \mathcal{V}$. The *radical* is $\text{rad}(\mathcal{V}) := \mathcal{V}^\perp$
(v) $\mathcal{V}$ is *nonsingular* or *nondegenerate* if $\text{rad}(\mathcal{V}) = \{\mathbf{0}\}$, *singular* otherwise.
(vi) $\mathcal{V}$ is *totally singular* or *totally degenerate* if $\text{rad}(\mathcal{V}) = \mathcal{V}$.

**Proposition 3.3.2.** *Metric Radical as a Bilinear Radical*
*Let $B(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, \mathbf{y} \rangle$ be the form of a metric vector space $\mathcal{V}$. Then $\text{rad}_L(B) = \text{rad}_R(B) = \text{rad}(\mathcal{V})$.*
*Thus $\mathcal{V}$ is nonsingular if and only if $B$ is perfect.*

*Proof.*

(i) By definition, $\text{rad}(\mathcal{V}) = \{\mathbf{x} \in \mathcal{V} : \langle \mathbf{x}, \mathbf{y} \rangle = 0 \ \forall \mathbf{y} \in \mathcal{V}\} = \text{rad}_L(B)$
(ii) If $\mathbf{x} \in \text{rad}_L(B)$, then $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ for all $\mathbf{y}$. Since $B$ is symmetric or skew-symmetric, $\langle \mathbf{y}, \mathbf{x} \rangle = \pm \langle \mathbf{x}, \mathbf{y} \rangle = 0$, so $\mathbf{x} \in \text{rad}_R(B)$. The reverse inclusion is identical.
(iii) By Proposition 4.4.6, $B$ is perfect if and only if $\text{rad}_L(B) = \text{rad}_R(B) = \{\mathbf{0}\}$, if and only if $\text{rad}(\mathcal{V}) = \{\mathbf{0}\}$.

*Structural Commentary*: Metric forms are the square bilinear forms whose two null directions agree.
*Key Insight*: Metric geometry uses the common left-right radical. □

Proposition 4.4.4 identifies the coordinate radical with $\ker(M_\beta^t)$ and gives $\dim \text{rad}(\mathcal{V}) = \dim \mathcal{V} - \text{rank}(M_\beta)$.
Proposition 4.4.6 gives $\text{rad}(\mathcal{V}) = \{\mathbf{0}\} \iff M_\beta$ is invertible.

**Remark 3.3.3.** *Subspace Radicals*
For $\mathcal{S} \leq \mathcal{V}$, compute $\text{rad}(\mathcal{S})$ using $\langle \cdot, \cdot \rangle|_{\mathcal{S} \times \mathcal{S}}$, where $\text{rad}(\mathcal{S}) = \{\mathbf{s} \in \mathcal{S} : \langle \mathbf{s}, \mathbf{t} \rangle = 0 \ \forall \mathbf{t} \in \mathcal{S}\} = \mathcal{S} \cap \mathcal{S}^\perp$.
In general, $\text{rad}(\mathcal{S}) \neq \mathcal{S}^\perp$; the latter is ambient.

**Proposition 3.3.4.** *Riesz Map Isomorphism Criterion*
*Let $\mathcal{V}$ be a finite-dimensional metric vector space and define $\tau : \mathcal{V} \to \mathcal{V}'$ where $\tau(\mathbf{x})(\mathbf{v}) = \langle \mathbf{v}, \mathbf{x} \rangle$.*
*Then $\ker(\tau) = \text{rad}(\mathcal{V})$, and $\tau$ is an isomorphism $\iff \text{rad}(\mathcal{V}) = \{\mathbf{0}\}$.*

*Proof.* Bilinearity gives $\tau \in \mathcal{L}(\mathcal{V}, \mathcal{V}')$, and
$\mathbf{x} \in \ker(\tau) \iff \langle \mathbf{v}, \mathbf{x} \rangle = 0 \ \forall \mathbf{v} \in \mathcal{V} \iff \langle \mathbf{x}, \mathbf{v} \rangle = 0 \ \forall \mathbf{v} \in \mathcal{V} \iff \mathbf{x} \in \mathcal{V}^\perp = \text{rad}(\mathcal{V})$

($\Rightarrow$) If $\tau$ is an isomorphism, then $\ker(\tau) = \{\mathbf{0}\}$, hence $\text{rad}(\mathcal{V}) = \{\mathbf{0}\}$.
($\Leftarrow$) If $\text{rad}(\mathcal{V}) = \{\mathbf{0}\}$, then $\ker(\tau) = \{\mathbf{0}\}$. Since $\dim \mathcal{V} = \dim \mathcal{V}'$, $\tau$ is an isomorphism.

*Structural Commentary*: The Riesz map is invertible exactly off the radical.
*Key Insight*: Nonsingularity is injectivity of the Riesz map. □

**Theorem 3.3.5.** *Riesz Map for Nonsingular Metric Spaces*
*Let $\mathcal{V}$ be a finite-dimensional nonsingular metric vector space. Then $\forall f \in \mathcal{V}' \ \exists! \, \mathbf{x} \in \mathcal{V}$, $f(\mathbf{v}) = \langle \mathbf{v}, \mathbf{x} \rangle \ \ \forall \mathbf{v} \in \mathcal{V}$.*

*Proof.* Let $\tau : \mathcal{V} \to \mathcal{V}'$ be given by $\tau(\mathbf{x})(\mathbf{v}) = \langle \mathbf{v}, \mathbf{x} \rangle$. By Proposition 3.3.4, $\tau$ is an isomorphism.
Hence each $f \in \mathcal{V}'$ has a unique $\mathbf{x} \in \mathcal{V}$ with $\tau(\mathbf{x}) = f$.
*Structural Commentary*: Nonsingularity identifies vectors with covectors; positivity is irrelevant.
*Key Insight*: Riesz representation is fundamentally nonsingularity, not order or length. □

**Definition 3.3.6.** *Isometries and Classical Groups*
Let $\mathcal{V}, \mathcal{W}$ be metric vector spaces.

(i) A bijective linear map $\sigma : \mathcal{V} \to \mathcal{W}$ is an *isometry* if $\langle \sigma\mathbf{x}, \sigma\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle \ \forall \mathbf{x}, \mathbf{y} \in \mathcal{V}$.
(ii) The group of isometries $\mathcal{V} \to \mathcal{V}$ is denoted $\mathrm{Iso}(\mathcal{V})$. If $\mathcal{V}$ is nonsingular orthogonal, this is the *orthogonal group* $\mathcal{O}(\mathcal{V})$; if $\mathcal{V}$ is nonsingular symplectic, this is the *symplectic group* $\mathrm{Sp}(\mathcal{V})$.

**Proposition 3.3.7.** *Matrix Criterion for Isometries*
*Let $\beta$ be a basis of a metric vector space $\mathcal{V}$, let $M = M_\beta$, and let $A = [\sigma]_{\beta \leftarrow \beta}$.*
*Then $\sigma \in \mathrm{Iso}(\mathcal{V}) \iff \sigma$ is bijective and $A^t M A = M$. If $M$ is nonsingular, then $\det A = \pm 1$.*

*Proof.*

(⇒) If $\sigma$ is an isometry, then it is bijective and, for all coordinate vectors $X, Y$, $X^t A^t M A Y = (AX)^t M (AY) = X^t M Y$. Hence $A^t M A = M$.
(⇐) If $\sigma$ is bijective and $A^t M A = M$, then for all $\mathbf{x}, \mathbf{y} \in \mathcal{V}$ with coordinate vectors $X, Y$, $\langle \sigma\mathbf{x}, \sigma\mathbf{y} \rangle = (AX)^t M (AY) = X^t M Y = \langle \mathbf{x}, \mathbf{y} \rangle$, so $\sigma$ is an isometry.

If $M$ is invertible, taking determinants in $A^t M A = M$ gives $(\det A)^2 = 1$, hence $\det A = \pm 1$.
*Structural Commentary*: Isometries are exactly the coordinate changes that stabilize the representing matrix.
*Key Insight*: Orthogonal and symplectic groups are stabilizer groups under congruence. □

**Definition 3.3.8.** *Adjoint for a Nonsingular Bilinear Form*
Let $\mathcal{V}$ be nonsingular.
For $T \in \mathcal{L}(\mathcal{V})$, the *metric adjoint* is the unique $T^* \in \mathcal{L}(\mathcal{V})$ such that $\langle T\mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, T^*\mathbf{y} \rangle \ \forall \mathbf{x}, \mathbf{y} \in \mathcal{V}$.

**Proposition 3.3.9.** *Metric Adjoints and Isometries*
*Let $\mathcal{V}$ be nonsingular and let $T \in \mathcal{L}(\mathcal{V})$.*

*(i)* $\exists! \, T^* \in \mathcal{L}(\mathcal{V})$.
*(ii) If $\beta$ is ordered, $M = M_\beta$, and $A = [T]_\beta$, then $[T^*]_\beta = M^{-1} A^t M$.*
*(iii) $T$ is an isometry $\iff T$ is bijective and $T^*T = I$. For bijective $T$, this is equivalent to $T^{-1} = T^*$.*

*Proof.*

(i) For $\mathbf{y} \in \mathcal{V}$, set $f_\mathbf{y}(\mathbf{x}) = \langle T\mathbf{x}, \mathbf{y} \rangle$. Theorem 3.3.5 gives a unique $T^*\mathbf{y}$ such that $f_\mathbf{y}(\mathbf{x}) = \langle \mathbf{x}, T^*\mathbf{y} \rangle$ for all $\mathbf{x}$. Uniqueness gives linearity of $\mathbf{y} \mapsto T^*\mathbf{y}$.
(ii) With $A = [T]_\beta$ and $B = [T^*]_\beta$, the adjoint identity gives $A^t M = MB$, so $B = M^{-1} A^t M$.
(iii) $\forall \mathbf{x}, \mathbf{y} \in \mathcal{V}$, $\langle T\mathbf{x}, T\mathbf{y} \rangle = \langle \mathbf{x}, T^*T\mathbf{y} \rangle$. so form preservation $\iff \langle \mathbf{x}, (T^*T - I)\mathbf{y} \rangle = 0 \ \forall \mathbf{x}, \mathbf{y} \iff T^*T = I$. This gives the stated bijective criterion; if $T$ is bijective, then $T^*T = I \iff T^* = T^{-1}$.

*Structural Commentary*: Adjoint theory needs nonsingularity, not positivity.
*Key Insight*: Isometry means the adjoint is the inverse. □

**Perspective 3.3.10.** *The Classification Problem*
Metric vector spaces are isometric exactly when their representing matrices are congruent.
Thus classification up to isometry is equivalent to classification of the associated matrices up to congruence, i.e. to specifying canonical representatives or complete congruence invariants.

**Perspective 3.3.11.** *Congruence versus Similarity*
Similarity $B = P^{-1} A P$ is change of basis for an operator. Congruence $N = P^t M P$ is change of basis for a bilinear form, with the same basis change in both slots. Over $\mathbb{R}$ this is the real case of $*$-congruence. Hence inertia classifies forms, not similarity classes of operators.

## 3.4 Radicals, Hyperbolic Spaces, and Witt Theory

**Theorem 3.4.1.** *Dimensions of Orthogonal Complements*
*Let $\mathcal{S} \leq \mathcal{V}$. If $\mathrm{rad}(\mathcal{V}) = \{\mathbf{0}\}$ or $\mathrm{rad}(\mathcal{S}) = \{\mathbf{0}\}$, then $\dim(\mathcal{S}) + \dim(\mathcal{S}^\perp) = \dim(\mathcal{V})$.*
*If $\mathrm{rad}(\mathcal{V}) = \{\mathbf{0}\}$, then $\mathcal{S}^{\perp\perp} = \mathcal{S}$, $\mathrm{rad}(\mathcal{S}) = \{\mathbf{0}\} \iff \mathrm{rad}(\mathcal{S}^\perp) = \{\mathbf{0}\}$.*

*Proof.*

i. Define $\rho : \mathcal{V} \to \mathcal{S}'$ by $\rho(\mathbf{x})(\mathbf{s}) = \langle \mathbf{s}, \mathbf{x} \rangle$. Then $\ker(\rho) = \mathcal{S}^\perp$.

ii. If $\operatorname{rad}(\mathcal{V}) = \{\mathbf{0}\}$, extend $f \in \mathcal{S}'$ to $\widetilde{f} \in \mathcal{V}'$ and use Theorem 3.3.5. If $\operatorname{rad}(\mathcal{S}) = \{\mathbf{0}\}$, represent $f$ inside $\mathcal{S}$. Thus $\rho$ is onto, so rank-nullity gives $\dim \mathcal{V} = \dim \mathcal{S}^\perp + \dim \mathcal{S}$.
iii. If $\operatorname{rad}(\mathcal{V}) = \{\mathbf{0}\}$, then $\mathcal{S} \subseteq \mathcal{S}^{\perp\perp}$ and $\dim \mathcal{S}^{\perp\perp} = \dim \mathcal{V} - \dim \mathcal{S}^\perp = \dim \mathcal{S}$, hence $\mathcal{S}^{\perp\perp} = \mathcal{S}$.
($\Rightarrow$) If $\operatorname{rad}(\mathcal{S}) = \{\mathbf{0}\}$, then $\operatorname{rad}(\mathcal{S}^\perp) = \mathcal{S}^\perp \cap \mathcal{S}^{\perp\perp} = \mathcal{S}^\perp \cap \mathcal{S}$, so $\operatorname{rad}(\mathcal{S}^\perp) = \{\mathbf{0}\}$.
($\Leftarrow$) If $\operatorname{rad}(\mathcal{S}^\perp) = \{\mathbf{0}\}$, then $\mathcal{S}^\perp \cap \mathcal{S} = \{\mathbf{0}\}$, so $\operatorname{rad}(\mathcal{S}) = \mathcal{S} \cap \mathcal{S}^\perp = \{\mathbf{0}\}$.

*Structural Commentary*: Complement dimensions come from surjectivity of the restriction-pairing map.
*Key Insight*: Orthogonal complement dimensions are rank-nullity for the restriction-of-pairing map. □

**Definition 3.4.2.** *Orthogonal Direct Sum*
For subspaces $\mathcal{S}, \mathcal{T} \leq \mathcal{V}$, write $\mathcal{V} = \mathcal{S} \odot \mathcal{T} \iff \mathcal{V} = \mathcal{S} \oplus \mathcal{T}$ and $\mathcal{S} \perp \mathcal{T}$.
Then $\mathcal{V}$ is the *orthogonal direct sum* of $\mathcal{S}$ and $\mathcal{T}$.

**Theorem 3.4.3.** *Splitting off the Radical*
*Let $R = \operatorname{rad}(\mathcal{V})$. Then $\exists\, \mathcal{S} \leq \mathcal{V}$ such that $\mathcal{V} = R \odot \mathcal{S}$, $\operatorname{rad}(\mathcal{S}) = \{\mathbf{0}\}$, and $R$ is totally singular.*

*Proof.* Choose $\mathcal{S}$ with $\mathcal{V} = R \oplus \mathcal{S}$. Since $R \perp \mathcal{V}$, $\mathcal{V} = R \odot \mathcal{S}$ and $R \subseteq R^\perp$, so $R$ is totally singular. If $\mathbf{s} \in \operatorname{rad}(\mathcal{S})$, then $\mathbf{s} \perp \mathcal{S}$ and $\mathbf{s} \perp R$, hence $\mathbf{s} \in R \cap \mathcal{S} = \{\mathbf{0}\}$.
*Structural Commentary*: Singularity splits as an invisible summand.
*Key Insight*: Once the radical is removed, the remaining form is nonsingular. □

**Proposition 3.4.4.** *Metric Quotients*
*Let $\mathcal{U} \leq \mathcal{V}$. The rule $\langle \mathbf{x} + \mathcal{U}, \mathbf{y} + \mathcal{U} \rangle_{\mathcal{V}/\mathcal{U}} := \langle \mathbf{x}, \mathbf{y} \rangle$ is well-defined if and only if $\mathcal{U} \leq \operatorname{rad}(\mathcal{V})$.*
*Then $\operatorname{rad}(\mathcal{V}/\mathcal{U}) = \operatorname{rad}(\mathcal{V})/\mathcal{U}$. In particular, $\mathcal{V}/\operatorname{rad}(\mathcal{V})$ is nonsingular.*

*Proof.*
($\Rightarrow$) If the rule is well-defined, then $0 = \langle \mathbf{u} + \mathcal{U}, \mathbf{y} + \mathcal{U} \rangle_{\mathcal{V}/\mathcal{U}} = \langle \mathbf{u}, \mathbf{y} \rangle$ $\forall\, \mathbf{u} \in \mathcal{U},\ \mathbf{y} \in \mathcal{V}$, so $\mathcal{U} \leq \operatorname{rad}(\mathcal{V})$.
($\Leftarrow$) If $\mathcal{U} \leq \operatorname{rad}(\mathcal{V})$, replacing $\mathbf{x}, \mathbf{y}$ by $\mathbf{x}+\mathbf{u}, \mathbf{y}+\mathbf{v}$ changes $\langle \mathbf{x}, \mathbf{y} \rangle$ by $\langle \mathbf{u}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{v} \rangle = 0$, rule well-defined.
If $\mathcal{U} \leq \operatorname{rad}(\mathcal{V})$, then $\mathbf{x} + \mathcal{U} \in \operatorname{rad}(\mathcal{V}/\mathcal{U}) \iff \langle \mathbf{x}, \mathbf{y} \rangle = 0\ \forall\, \mathbf{y} \in \mathcal{V} \iff \mathbf{x} \in \operatorname{rad}(\mathcal{V})$, hence $\operatorname{rad}(\mathcal{V}/\mathcal{U}) = \operatorname{rad}(\mathcal{V})/\mathcal{U}$.
*Structural Commentary*: Quotients preserve the form exactly along invisible directions.
*Key Insight*: The radical is the largest subspace that can be collapsed without making the form ambiguous. □

**Proposition 3.4.5.** *Radicals and Orthogonal Direct Sums*
*If $\mathcal{V} = \mathcal{S} \odot \mathcal{T}$, then $\operatorname{rad}(\mathcal{V}) = \operatorname{rad}(\mathcal{S}) \odot \operatorname{rad}(\mathcal{T})$; $\mathcal{V}/\operatorname{rad}(\mathcal{V}) \cong \mathcal{S}/\operatorname{rad}(\mathcal{S}) \odot \mathcal{T}/\operatorname{rad}(\mathcal{T})$ as metric vector spaces.*
*Hence $\mathcal{V}$ is nonsingular if and only if $\mathcal{S}$ and $\mathcal{T}$ are nonsingular.*

*Proof.* For $\mathbf{v} = \mathbf{s} + \mathbf{t}$, $\mathbf{v} \in \operatorname{rad}(\mathcal{V}) \iff \mathbf{v} \perp \mathcal{V} \iff \mathbf{s} \perp \mathcal{S}$ and $\mathbf{t} \perp \mathcal{T} \iff \mathbf{s} \in \operatorname{rad}(\mathcal{S}),\ \mathbf{t} \in \operatorname{rad}(\mathcal{T})$.
Hence $\operatorname{rad}(\mathcal{V}) = \operatorname{rad}(\mathcal{S}) \odot \operatorname{rad}(\mathcal{T})$. The isomorphism is induced by $\Phi((\mathbf{s}+\mathbf{t}) + \operatorname{rad}(\mathcal{V})) = (\mathbf{s} + \operatorname{rad}(\mathcal{S}), \mathbf{t} + \operatorname{rad}(\mathcal{T}))$.
It is well-defined and bijective by the radical formula, and it preserves the form since cross terms vanish.
($\Rightarrow$) If $\mathcal{V}$ is nonsingular, then $\operatorname{rad}(\mathcal{S}) \odot \operatorname{rad}(\mathcal{T}) = \{\mathbf{0}\}$, so $\mathcal{S}$ and $\mathcal{T}$ are nonsingular.
($\Leftarrow$) If $\mathcal{S}$ and $\mathcal{T}$ are nonsingular, then $\operatorname{rad}(\mathcal{V}) = \operatorname{rad}(\mathcal{S}) \odot \operatorname{rad}(\mathcal{T}) = \{\mathbf{0}\}$.
*Structural Commentary*: Orthogonal direct sums split both visible and invisible parts.
*Key Insight*: The regular quotient of an orthogonal sum is the orthogonal sum of the regular quotients. □

**Corollary 3.4.6.** *Recovering a Singular Space from Its Regular Quotient*
*If $\dim \mathcal{V} = \dim \mathcal{W}$ and $\mathcal{V}/\operatorname{rad}(\mathcal{V}) \cong \mathcal{W}/\operatorname{rad}(\mathcal{W})$, then $\mathcal{V} \cong \mathcal{W}$ as metric vector spaces.*

*Proof.* By Theorem 3.4.3, choose $\mathcal{V} = \operatorname{rad}(\mathcal{V}) \odot \mathcal{S}$, $\mathcal{W} = \operatorname{rad}(\mathcal{W}) \odot \mathcal{T}$, with $\mathcal{S}, \mathcal{T}$ nonsingular.
The quotient isometry gives $\mathcal{S} \cong \mathcal{T}$. Since $\dim \mathcal{V} = \dim \mathcal{W}$, also $\dim \operatorname{rad}(\mathcal{V}) = \dim \operatorname{rad}(\mathcal{W})$; choose any vector-space isomorphism between the radicals, which is an isometry because both radicals are totally singular.
The orthogonal direct sum of these two isometries gives $\mathcal{V} \cong \mathcal{W}$.
*Structural Commentary*: A singular metric space is its regular quotient plus invisible dimension.
*Key Insight*: Once the regular quotient and total dimension are fixed, the radical has no further metric data. □

**Definition 3.4.7.** *Hyperbolic Pairs and Hyperbolic Spaces*
(i) A *hyperbolic pair* is $(\mathbf{u}, \mathbf{v})$ with $\langle \mathbf{u}, \mathbf{u} \rangle = \langle \mathbf{v}, \mathbf{v} \rangle = 0$ and $\langle \mathbf{u}, \mathbf{v} \rangle = 1$.
(ii) The plane $\operatorname{span}(\mathbf{u}, \mathbf{v})$ of a hyperbolic pair is a *hyperbolic plane*.
(iii) A *hyperbolic space* is an orthogonal direct sum of hyperbolic planes.
(iv) A basis $(\mathbf{u}_1, \mathbf{v}_1, \ldots, \mathbf{u}_k, \mathbf{v}_k)$ is a *hyperbolic basis* if each $(\mathbf{u}_i, \mathbf{v}_i)$ is hyperbolic and the planes $\operatorname{span}(\mathbf{u}_i, \mathbf{v}_i)$ are mutually orthogonal. In the symplectic case it is a *symplectic basis*.

**Theorem 3.4.8.** *Capturing Isotropic Vectors*
*Let $\mathcal{V}$ be nonsingular. If $\mathcal{V}$ is orthogonal, assume $\operatorname{char}(\mathbb{F}) \neq 2$.*
*For every isotropic $\mathbf{u} \in \mathcal{V}$, $\exists\, \mathbf{v} \in \mathcal{V}$ such that $(\mathbf{u}, \mathbf{v})$ is hyperbolic.*

*Proof.* Choose $\mathbf{x}$ with $c := \langle \mathbf{u}, \mathbf{x} \rangle \neq 0$. If the form is alternating, take $\mathbf{v} = c^{-1}\mathbf{x}$. Then $\langle \mathbf{u}, \mathbf{v} \rangle = 1$ and $\langle \mathbf{v}, \mathbf{v} \rangle = 0$. If the form is symmetric, set $\mathbf{v} = c^{-1}\mathbf{x} - \frac{\langle \mathbf{x}, \mathbf{x} \rangle}{2c^2}\mathbf{u}$. Using $\langle \mathbf{u}, \mathbf{u} \rangle = 0$ gives $\langle \mathbf{u}, \mathbf{v} \rangle = 1$ and $\langle \mathbf{v}, \mathbf{v} \rangle = 0$.
*Structural Commentary*: Nonsingularity supplies a partner; one adjustment makes it isotropic.
*Key Insight*: Isotropic directions in nonsingular spaces come in hyperbolic pairs. □

**Proposition 3.4.9.** *Recognizing Orthogonal Hyperbolic Planes*
*Let $\mathcal{H}$ be a 2-dimensional nonsingular orthogonal geometry with* $\mathrm{char}(\mathbb{F}) \neq 2$. *The following are equivalent:*

*(i) $\mathcal{H}$ is a hyperbolic plane;*
*(ii) $\exists\, 0 \neq \mathbf{u} \in \mathcal{H}$ with $\langle \mathbf{u}, \mathbf{u} \rangle = 0$;*
*(iii) $\Delta(\mathcal{H}) = (\mathbb{F}^\times)^2(-1)$ as a square class.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) A hyperbolic pair consists of nonzero isotropic vectors.
(*ii*) $\Rightarrow$ (*i*) By Theorem 3.4.8, choose $\mathbf{v}$ with $(\mathbf{u}, \mathbf{v})$ hyperbolic. Since $\dim \mathcal{H} = 2$, $\mathcal{H} = \mathrm{span}(\mathbf{u}, \mathbf{v})$.
(*i*) $\Rightarrow$ (*iii*) In a hyperbolic basis, $M = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, so $\det M = -1$ and $\Delta(\mathcal{H}) = (\mathbb{F}^\times)^2(-1)$.
(*iii*) $\Rightarrow$ (*ii*) Diagonalize $M = \mathrm{diag}(a, b)$ with $a, b \neq 0$. If $\Delta(\mathcal{H}) = (\mathbb{F}^\times)^2(-1)$, then $ab = -r^2$ for some $r \neq 0$. Hence $a(r/a)^2 + b = 0$. Thus $(r/a, 1)$ is isotropic in this basis.

*Structural Commentary*: In dimension 2, isotropy and discriminant detect hyperbolicity.
*Key Insight*: Hyperbolic means the form represents 0 nontrivially. □

**Theorem 3.4.10.** *Nonsingular Completions*
*Let $\mathcal{U}$ be a subspace of a nonsingular metric vector space $\mathcal{V}$, and assume* $\mathrm{char}(\mathbb{F}) \neq 2$ *in the orthogonal case. Write $\mathcal{U} = \mathrm{span}(\mathbf{u}_1, \ldots, \mathbf{u}_r) \odot \mathcal{W}$, where $(\mathbf{u}_1, \ldots, \mathbf{u}_r)$ is a basis of $\mathrm{rad}(\mathcal{U})$ and $\mathcal{W}$ is nonsingular. Then $\exists\, \mathbf{v}_1, \ldots, \mathbf{v}_r$ such that $\widehat{\mathcal{U}} = \mathcal{H}_r \odot \mathcal{W}$; $\mathcal{H}_r = \mathrm{span}(\mathbf{u}_1, \mathbf{v}_1, \ldots, \mathbf{u}_r, \mathbf{v}_r)$ is nonsingular, contains $\mathcal{U}$, and $\dim \widehat{\mathcal{U}} = \dim \mathcal{U} + \dim \mathrm{rad}(\mathcal{U})$.*
*The minimal nonsingular extensions of $\mathcal{U}$ are exactly such hyperbolic extensions, up to isometry.*

*Proof.*

(i) Since $\mathcal{W}$ is nonsingular, Theorem 3.4.1 gives $\mathcal{V} = \mathcal{W} \odot \mathcal{W}^\perp$. Thus $\mathbf{u}_i \in \mathcal{W}^\perp$ for all $i$.
(ii) Construct $\mathbf{v}_j$ inductively. Suppose $\mathcal{H}_{j-1}$ is built and set $\mathcal{S}_j = \mathrm{span}(\mathbf{u}_{j+1}, \ldots, \mathbf{u}_r) \odot \mathcal{W} \odot \mathcal{H}_{j-1}$. Then $\mathbf{u}_j \perp \mathcal{S}_j$ and $\mathbf{u}_j \notin \mathcal{S}_j = \mathcal{S}_j^{\perp\perp}$, so $\exists \mathbf{x} \in \mathcal{S}_j^\perp$ with $c := \langle \mathbf{u}_j, \mathbf{x} \rangle \neq 0$. Put

$$\mathbf{v}_j = \begin{cases} c^{-1}\mathbf{x}, & \text{alternating case,} \\ c^{-1}\mathbf{x} - \dfrac{\langle \mathbf{x}, \mathbf{x} \rangle}{2c^2}\mathbf{u}_j, & \text{orthogonal case.} \end{cases}$$

Then $\mathbf{v}_j \in \mathcal{S}_j^\perp$ and $(\mathbf{u}_j, \mathbf{v}_j)$ is hyperbolic; set $\mathcal{H}_j = \mathcal{H}_{j-1} \odot \mathrm{span}(\mathbf{u}_j, \mathbf{v}_j)$.
(iii) After $r$ steps, $\widehat{\mathcal{U}} = \mathcal{H}_r \odot \mathcal{W}$ is nonsingular and $\dim \widehat{\mathcal{U}} = \dim \mathcal{U} + r$.
(iv) Let $\mathcal{X} \supseteq \mathcal{U}$ be nonsingular and choose $\mathcal{X} = \mathcal{U} \oplus \mathcal{Y}$. If $0 \neq \mathbf{u} \in \mathrm{rad}(\mathcal{U})$ paired trivially with $\mathcal{Y}$, then $\mathbf{u} \in \mathrm{rad}(\mathcal{X})$, impossible. Hence the pairing $\mathrm{rad}(\mathcal{U}) \times \mathcal{Y} \to \mathbb{F}$ has zero left radical, so $\dim \mathcal{Y} \geq r$ and $\dim \mathcal{X} \geq \dim \mathcal{U} + r$. If equality holds, the construction inside $\mathcal{X}$ gives $\widehat{\mathcal{U}} \subseteq \mathcal{X}$ with equal dimension, hence $\widehat{\mathcal{U}} = \mathcal{X}$. Any two minimal completions are isometric since each is $\mathcal{W}\odot$ a hyperbolic $r$-space.

*Structural Commentary*: Add one hyperbolic partner per radical direction.
*Key Insight*: Minimal nonsingular completion is radical repair by hyperbolic planes. □

**Principle 3.4.11.** *Witt Extension and Cancellation*
Let $\mathcal{V} \cong \mathcal{V}'$ be nonsingular metric vector spaces of same kind, either symplectic or orthogonal with $\mathrm{char}(\mathbb{F}) \neq 2$. The relevant principles are:

(i) *Witt extension*: every isometry $\tau : \mathcal{S} \to \mathcal{T}$ between subspaces $\mathcal{S} \leq \mathcal{V}$ and $\mathcal{T} \leq \mathcal{V}'$ extends to an isometry $\overline{\tau} : \mathcal{V} \to \mathcal{V}'$ with $\overline{\tau}|_{\mathcal{S}} = \tau$;
(ii) *Witt cancellation*: for nonsingular subspaces $\mathcal{S} \leq \mathcal{V}$ and $\mathcal{T} \leq \mathcal{V}'$, $\mathcal{S} \cong \mathcal{T} \Rightarrow \mathcal{S}^\perp \cong \mathcal{T}^\perp$.

**Proposition 3.4.12.** *Witt Extension–Cancellation Bridge*
*Let $\mathcal{V} \cong \mathcal{V}'$ be nonsingular metric vector spaces of same kind, either symplectic or orthogonal with* $\mathrm{char}(\mathbb{F}) \neq 2$. *Using Theorem 3.4.10, Witt extension $\iff$ Witt cancellation.*

*Proof.*

($\Rightarrow$) Let $\mathcal{V} = \mathcal{S} \odot \mathcal{S}^\perp$, $\mathcal{V}' = \mathcal{T} \odot \mathcal{T}^\perp$, and let $\sigma : \mathcal{S} \to \mathcal{T}$ be an isometry. Extend $\sigma$ to $\widehat{\sigma} : \mathcal{V} \to \mathcal{V}'$. Then $\widehat{\sigma}(\mathcal{S}^\perp) = \mathcal{T}^\perp$, hence $\mathcal{S}^\perp \cong \mathcal{T}^\perp$.

($\Leftarrow$) Let $\sigma : \mathcal{S} \to \mathcal{T}$ be an isometry. By Theorem 3.4.10, choose minimal nonsingular completions $\widehat{\mathcal{S}} \supseteq \mathcal{S}$ and $\widehat{\mathcal{T}} \supseteq \mathcal{T}$, and extend $\sigma$ to $\widehat{\sigma} : \widehat{\mathcal{S}} \to \widehat{\mathcal{T}}$ by matching hyperbolic partners of $\operatorname{rad}(\mathcal{S})$ and $\operatorname{rad}(\mathcal{T})$.
Then $\mathcal{V} = \widehat{\mathcal{S}} \odot \widehat{\mathcal{S}}^{\perp}$, $\mathcal{V}' = \widehat{\mathcal{T}} \odot \widehat{\mathcal{T}}^{\perp}$. Cancellation gives $\eta : \widehat{\mathcal{S}}^{\perp} \to \widehat{\mathcal{T}}^{\perp}$, so $\widehat{\sigma} \odot \eta : \mathcal{V} \to \mathcal{V}'$ extends $\sigma$.

*Structural Commentary*: Complete first, then cancel complements.
*Key Insight*: Complete singular subspaces first; after that, the complement is the only missing piece. □

**Theorem 3.4.13.** *Witt Extension Theorem*
*Let $\mathcal{V} \cong \mathcal{W}$ be nonsingular metric vector spaces of same kind, either symplectic or orthogonal with $\operatorname{char}(\mathbb{F}) \neq 2$. For every subspace isometry $\sigma : \mathcal{S} \to \mathcal{T}$, $\mathcal{S} \leq \mathcal{V}$, $\mathcal{T} \leq \mathcal{W}$, there exists an isometry $\widehat{\sigma} : \mathcal{V} \to \mathcal{W}$ with $\widehat{\sigma}|_{\mathcal{S}} = \sigma$. Thus, if $\mathcal{V} = \mathcal{W}$, every subspace isometry is the restriction of an ambient isometry.*

*Proof.* Induct simultaneously on $(\dim \mathcal{V}, \dim \mathcal{S})$. If $\mathcal{S} = 0$, take any fixed isometry $\mathcal{V} \cong \mathcal{W}$.
Let $\mathcal{R} = \operatorname{rad}(\mathcal{S})$, $\mathcal{R}' = \sigma\mathcal{R} = \operatorname{rad}(\mathcal{T})$. By Theorem 3.4.10, choose minimal nonsingular completions $\widehat{\mathcal{S}} \supseteq \mathcal{S}$ and $\widehat{\mathcal{T}} \supseteq \mathcal{T}$; matching hyperbolic partners over $\mathcal{R}$ and $\mathcal{R}'$ extends $\sigma$ to $\widehat{\sigma} : \widehat{\mathcal{S}} \to \widehat{\mathcal{T}}$. Hence assume $\mathcal{S}, \mathcal{T}$ nonsingular.
Choose a nonsingular summand $\mathcal{A} \leq \mathcal{S}$, where

$$\mathcal{A} = \begin{cases} \operatorname{span}(\mathbf{s}, \mathbf{u}) \text{ hyperbolic}, & \text{symplectic case}, \\ \operatorname{span}(\mathbf{s}),\ \langle \mathbf{s}, \mathbf{s} \rangle \neq 0, & \text{orthogonal case}. \end{cases}$$

Set $\mathcal{B} = \sigma\mathcal{A}$, $\mathcal{S}_0 = \mathcal{S} \cap \mathcal{A}^{\perp}$, and $\mathcal{T}_0 = \mathcal{T} \cap \mathcal{B}^{\perp}$. One-summand case extends $\sigma|_{\mathcal{A}}$ to $\alpha : \mathcal{V} \to \mathcal{W}$; hence $\alpha(\mathcal{A}^{\perp}) = \mathcal{B}^{\perp}$ and $\mathcal{V} = \mathcal{A} \odot \mathcal{A}^{\perp}$, $\mathcal{W} = \mathcal{B} \odot \mathcal{B}^{\perp}$, $\mathcal{S} = \mathcal{A} \odot \mathcal{S}_0$, and $\mathcal{T} = \mathcal{B} \odot \mathcal{T}_0$. Since $\dim \mathcal{A}^{\perp} < \dim \mathcal{V}$, induction gives an isometry $\eta : \mathcal{A}^{\perp} \to \mathcal{B}^{\perp}$ with $\eta|_{\mathcal{S}_0} = \sigma|_{\mathcal{S}_0}$. Then $(\sigma|_{\mathcal{A}}) \odot \eta : \mathcal{V} \to \mathcal{W}$ extends $\sigma$.
*Structural Commentary*: Subspace isometries extend ambiently.
*Key Insight*: Complete radicals, then peel matched summands. □

**Corollary 3.4.14.** *Witt Cancellation*
*Let $\mathcal{V} = \mathcal{S} \odot \mathcal{A}$, $\mathcal{W} = \mathcal{T} \odot \mathcal{B}$ be nonsingular metric vector spaces of the same kind, either symplectic or orthogonal with $\operatorname{char}(\mathbb{F}) \neq 2$. If $\mathcal{V} \cong \mathcal{W}$ and $\mathcal{S} \cong \mathcal{T}$, then $\mathcal{A} \cong \mathcal{B}$.*

*Proof.* Let $\sigma : \mathcal{S} \to \mathcal{T}$ be an isometry. By Theorem 3.4.13, $\exists\, \widehat{\sigma} : \mathcal{V} \to \mathcal{W}$ with $\widehat{\sigma}|_{\mathcal{S}} = \sigma$.
Since $\mathcal{A} = \mathcal{S}^{\perp}$ and $\mathcal{B} = \mathcal{T}^{\perp}$, $\widehat{\sigma}(\mathcal{A}) = \widehat{\sigma}(\mathcal{S}^{\perp}) = (\widehat{\sigma}\mathcal{S})^{\perp} = \mathcal{T}^{\perp} = \mathcal{B}$. Thus $\widehat{\sigma}|_{\mathcal{A}} : \mathcal{A} \to \mathcal{B}$ is an isometry.
*Structural Commentary*: Matched summands cancel.
*Key Insight*: Extend, then restrict to complements. □

**Definition 3.4.15.** *Witt Index*
Let $\mathcal{V}$ be nonsingular orthogonal with $\operatorname{char}(\mathbb{F}) \neq 2$. The *Witt index* is $w(\mathcal{V}) := \max\{\dim \mathcal{L} : \mathcal{L} \leq \mathcal{V},\ \mathcal{L} \subseteq \mathcal{L}^{\perp}\}$.

**Theorem 3.4.16.** *Witt Index and Anisotropic Decomposition*
*Let $\mathcal{V}$ be orthogonal over $\mathbb{F}$ with $\operatorname{char}(\mathbb{F}) \neq 2$. Then:*

*(i) if $\mathcal{V}$ is nonsingular, every maximal totally isotropic subspace has dimension $w(\mathcal{V})$;*
*(ii) if $\mathcal{V}$ is nonsingular, every maximal hyperbolic subspace has dimension $2w(\mathcal{V})$;*
*(iii) for $R = \operatorname{rad}(\mathcal{V})$ and a nonsingular complement $\mathcal{U}$ of $R$, there exist $\mathcal{H} \leq \mathcal{U}$ hyperbolic and $\mathcal{S} \leq \mathcal{U}$ anisotropic such that $\mathcal{V} = \mathcal{S} \odot \mathcal{H} \odot R$, $\dim \mathcal{H} = 2w(\mathcal{U})$.*

*Proof.*

(i) Let $\mathcal{L}$ be maximal totally isotropic and choose $\mathcal{M}$ totally isotropic with $\dim \mathcal{M} = w(\mathcal{V})$.
Since $\dim \mathcal{L} \leq w(\mathcal{V})$, choose $\mathcal{M}_0 \leq \mathcal{M}$ with $\dim \mathcal{M}_0 = \dim \mathcal{L}$.
Any isomorphism $\mathcal{L} \to \mathcal{M}_0$ is an isometry, so Theorem 3.4.13 extends it to $\Phi \in \mathcal{O}(\mathcal{V})$.
Then $\mathcal{L} \subseteq \Phi^{-1}\mathcal{M}$; maximality gives $\mathcal{L} = \Phi^{-1}\mathcal{M}$, hence $\dim \mathcal{L} = w(\mathcal{V})$.

(ii) If $\mathcal{H}$ has $k$ hyperbolic planes, then $\mathcal{H}$ contains a $k$-dimensional totally isotropic subspace, so $k \leq w(\mathcal{V})$.
Conversely, Theorem 3.4.10 completes a $w(\mathcal{V})$-dimensional totally isotropic subspace to a hyperbolic subspace $\mathcal{H}_0$ of dimension $2w(\mathcal{V})$. If a maximal hyperbolic $\mathcal{H}$ had dimension $2k < 2w(\mathcal{V})$, choose hyperbolic $\mathcal{H}_1 \leq \mathcal{H}_0$ with $\dim \mathcal{H}_1 = 2k$ and extend an isometry $\mathcal{H} \to \mathcal{H}_1$ to $\Phi \in \mathcal{O}(\mathcal{V})$; then $\Phi^{-1}\mathcal{H}_0$ is a larger hyperbolic subspace containing $\mathcal{H}$, contradiction.

(iii) By Theorem 3.4.3, $\mathcal{V} = R \odot \mathcal{U}$ with $\mathcal{U}$ nonsingular. Choose maximal hyperbolic $\mathcal{H} \leq \mathcal{U}$ and set $\mathcal{S} = \mathcal{H}^{\perp} \cap \mathcal{U}$.
Then $\dim \mathcal{H} = 2w(\mathcal{U})$ by (ii), and $\mathcal{U} = \mathcal{H} \odot \mathcal{S}$. If $\mathcal{S}$ had a nonzero isotropic vector, Theorem 3.4.8 would give a hyperbolic plane in $\mathcal{S}$, enlarging $\mathcal{H}$; hence $\mathcal{S}$ is anisotropic.

*Structural Commentary*: Orthogonal spaces split into radical, hyperbolic part, and anisotropic residue.
*Key Insight*: The Witt index counts how many hyperbolic planes can be split off. □

## 3.5 Symplectic Classification and Transvections

**Theorem 3.5.1.** *Classification of Symplectic Geometries*
*Let $\mathcal{V}$ be a symplectic geometry, $R = \operatorname{rad}(\mathcal{V})$, and $n = \dim(\mathcal{V})$.*

*(i) If $R = 0$, then $\mathcal{V}$ is hyperbolic; hence $n$ is even.*
*(ii) $\mathcal{V} = R \odot \mathcal{H}$, where $R$ is totally singular and $\mathcal{H}$ is hyperbolic.*
*(iii) If $r = \operatorname{rank}(\mathcal{V}) = 2k$, then some basis has representing matrix $M_{\mathcal{V}} = X_{2k,n-2k} := \operatorname{diag}(J_{2k}, 0_{n-2k})$, $J_{2k} := \bigoplus_{i=1}^{k} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. For symplectic geometries, $\mathcal{V}, \mathcal{W}$, $\mathcal{V} \cong \mathcal{W} \iff (\dim \mathcal{V}, \operatorname{rank} \mathcal{V}) = (\dim \mathcal{W}, \operatorname{rank} \mathcal{W})$.*

*Proof.*

(i) The case $\mathcal{V} = 0$ is clear. If $R = 0$ and $\mathcal{V} \neq 0$, then every $0 \neq x \in \mathcal{V}$ is isotropic.
By Theorem 3.4.8, $x$ lies in a hyperbolic plane $\mathcal{H}_1 \subset \mathcal{V}$. Since $\mathcal{H}_1$ is nonsingular, Theorem 3.4.1 gives $\mathcal{V} = \mathcal{H}_1 \odot \mathcal{H}_1^{\perp}$. Induction on $n$ gives the claim.
(ii) By Theorem 3.4.3, $\mathcal{V} = R \odot \mathcal{H}$ with $\operatorname{rad}(\mathcal{H}) = 0$.
Hence $\mathcal{H}$ is hyperbolic by (i), and $R$ is totally singular since $R \perp \mathcal{V}$.
(iii) In (ii), $\dim \mathcal{H} = r = 2k$ and $\dim R = n - 2k$.
A symplectic basis of $\mathcal{H}$ followed by any basis of $R$ gives $M_{\mathcal{V}} = X_{2k,n-2k}$. Dimension and rank are isometry invariants; conversely, equal $(n, r)$ gives the same normal matrix, hence an isometry.

*Structural Commentary*: Alternating forms have only one nonsingular building block: the hyperbolic plane.
*Key Insight*: Symplectic classification is rank plus dimension. □

Theorem 3.4.13 and Corollary 3.4.14 specialize to nonsingular symplectic geometries.

**Definition 3.5.2.** *Symplectic Transvection*
Let $\mathcal{V}$ be a nonsingular symplectic geometry.
For $\mathbf{v} \in \mathcal{V} \setminus \{\mathbf{0}\}$, $a \in \mathbb{F}$, *symplectic transvection* with direction $\mathbf{v}$, parameter $a$ is $\tau_{\mathbf{v},a} : \mathcal{V} \to \mathcal{V}$, $\mathbf{x} \mapsto \mathbf{x} + a \langle \mathbf{x}, \mathbf{v} \rangle \mathbf{v}$.

**Proposition 3.5.3.** *Basic Properties of Symplectic Transvections*
*Let $\mathcal{V}$ be a nonsingular symplectic geometry, $0 \neq \mathbf{v} \in \mathcal{V}$, $a, b \in \mathbb{F}$, $c \in \mathbb{F}^{\times}$, and $\sigma \in \operatorname{Sp}(\mathcal{V})$. Then:*

*(i) $\tau_{\mathbf{v},a} \in \operatorname{Sp}(\mathcal{V})$;*
*(ii) $\tau_{\mathbf{v},a} = I$ if and only if $a = 0$;*
*(iii) $\tau_{\mathbf{v},a}\tau_{\mathbf{v},b} = \tau_{\mathbf{v},a+b}$ and $\tau_{\mathbf{v},a}^{-1} = \tau_{\mathbf{v},-a}$;*
*(iv) $\sigma\tau_{\mathbf{v},a}\sigma^{-1} = \tau_{\sigma\mathbf{v},a}$;*
*(v) $\tau_{c\mathbf{v},a} = \tau_{\mathbf{v},ac^2}$.*

*Proof.*

(i) For $\mathbf{x}, \mathbf{y} \in \mathcal{V}$, alternation and skew-symmetry give
$\langle \tau_{\mathbf{v},a}\mathbf{x}, \tau_{\mathbf{v},a}\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + a \langle \mathbf{x}, \mathbf{v} \rangle \langle \mathbf{v}, \mathbf{y} \rangle + a \langle \mathbf{y}, \mathbf{v} \rangle \langle \mathbf{x}, \mathbf{v} \rangle + a^2 \langle \mathbf{x}, \mathbf{v} \rangle \langle \mathbf{y}, \mathbf{v} \rangle \langle \mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle$.
Also $\tau_{\mathbf{v},-a}\tau_{\mathbf{v},a}(\mathbf{x}) = \mathbf{x} = \tau_{\mathbf{v},a}\tau_{\mathbf{v},-a}(\mathbf{x})$, so $\tau_{\mathbf{v},a}$ is invertible and hence symplectic.
(ii) If $a = 0$, then $\tau_{\mathbf{v},a} = I$. If $a \neq 0$, nonsingularity gives $\mathbf{x} \in \mathcal{V}$ with $\langle \mathbf{x}, \mathbf{v} \rangle \neq 0$, so $\tau_{\mathbf{v},a}\mathbf{x} - \mathbf{x} = a \langle \mathbf{x}, \mathbf{v} \rangle \mathbf{v} \neq \mathbf{0}$.
(iii) For $\mathbf{x} \in \mathcal{V}$, since $\langle \mathbf{v}, \mathbf{v} \rangle = 0$, $\tau_{\mathbf{v},a}\tau_{\mathbf{v},b}(\mathbf{x}) = \mathbf{x} + (a+b) \langle \mathbf{x}, \mathbf{v} \rangle \mathbf{v} = \tau_{\mathbf{v},a+b}(\mathbf{x})$. Taking $b = -a$ gives $\tau_{\mathbf{v},a}^{-1} = \tau_{\mathbf{v},-a}$.
(iv) For $\mathbf{x} \in \mathcal{V}$, $\sigma\tau_{\mathbf{v},a}\sigma^{-1}(\mathbf{x}) = \mathbf{x} + a \langle \sigma^{-1}\mathbf{x}, \mathbf{v} \rangle \sigma\mathbf{v} = \mathbf{x} + a \langle \mathbf{x}, \sigma\mathbf{v} \rangle \sigma\mathbf{v} = \tau_{\sigma\mathbf{v},a}(\mathbf{x})$.
(v) For $\mathbf{x} \in \mathcal{V}$, $\tau_{c\mathbf{v},a}(\mathbf{x}) = \mathbf{x} + a \langle \mathbf{x}, c\mathbf{v} \rangle c\mathbf{v} = \mathbf{x} + ac^2 \langle \mathbf{x}, \mathbf{v} \rangle \mathbf{v} = \tau_{\mathbf{v},ac^2}(\mathbf{x})$

*Structural Commentary*: Fix $\mathbf{v}^{\perp}$; shear along $\mathbb{F}\mathbf{v}$.
*Key Insight*: Alternation cancels the possible quadratic error term. □

**Theorem 3.5.4.** *Symplectic Transvections Generate*
*For a nonsingular symplectic geometry $\mathcal{V}$, $\operatorname{Sp}(\mathcal{V}) = \langle \tau_{\mathbf{v},a} : 0 \neq \mathbf{v} \in \mathcal{V},\ a \in \mathbb{F} \rangle$.*

*Proof.* Let $G$ denote the right-hand side. By Proposition 3.5.3, $G \leq \operatorname{Sp}(\mathcal{V})$.
If $\mathbf{x}, \mathbf{y} \neq \mathbf{0}$, $\langle \mathbf{x}, \mathbf{y} \rangle \neq 0$, then $\tau_{\mathbf{y}-\mathbf{x},\, \langle \mathbf{x}, \mathbf{y} \rangle^{-1}}(\mathbf{x}) = \mathbf{y}$. If $\langle \mathbf{x}, \mathbf{y} \rangle = 0$, choose $\mathbf{z}$ with $\langle \mathbf{x}, \mathbf{z} \rangle \langle \mathbf{z}, \mathbf{y} \rangle \neq 0$, $\mathbf{x} \mapsto \mathbf{z} \mapsto \mathbf{y}$.
Hence $G$ is transitive on nonzero vectors; the same argument in $\mathbf{u}^{\perp}/\mathbb{F}\mathbf{u}$ gives transitivity on hyperbolic pairs.
Let $\sigma \in \operatorname{Sp}(\mathcal{V})$ and induct on $n = \dim \mathcal{V}$. For $n = 2$, hyperbolic-pair transitivity gives $\sigma \in G$.
For $n > 2$, choose a hyperbolic plane $\mathcal{H} = \operatorname{span}(\mathbf{u}, \mathbf{v})$. Pair transitivity gives $\gamma \in G$ with $\gamma\sigma|_{\mathcal{H}} = I_{\mathcal{H}}$.
Then $\gamma\sigma$ restricts to an element of $\operatorname{Sp}(\mathcal{H}^{\perp})$, which is a product of transvections by induction; extended by the identity on $\mathcal{H}$, these are transvections of $\mathcal{V}$. Thus $\sigma \in G$.
*Structural Commentary*: Symplectic transformations are built from elementary shears along null directions.
*Key Insight*: Hyperbolic pairs are the movable frames of symplectic geometry. □

**Proposition 3.5.5.** *Symplectic Transformations Have Determinant One*
*If $\mathcal{V}$ is nonsingular symplectic, then $\operatorname{Sp}(\mathcal{V}) \subseteq \operatorname{SL}(\mathcal{V})$.*

*Proof.* By Theorem 3.5.4, it suffices to show $\det(\tau_{\mathbf{v},a}) = 1$. Let $\tau_{\mathbf{v},a} = I + a\,\mathbf{v} \otimes \lambda_{\mathbf{v}}$, $\lambda_{\mathbf{v}}(\mathbf{x}) = \langle \mathbf{x}, \mathbf{v} \rangle$.
Since $\lambda_{\mathbf{v}}(\mathbf{v}) = \langle \mathbf{v}, \mathbf{v} \rangle = 0$, $\tau_{\mathbf{v},a}$ fixes $\mathbb{F}\mathbf{v}$ and induces $I$ on $\mathcal{V}/\mathbb{F}\mathbf{v}$. Hence its matrix in any basis starting with $\mathbf{v}$ is unipotent triangular, so $\det(\tau_{\mathbf{v},a}) = 1$.
*Structural Commentary*: Generators rule out $\det = -1$.
*Key Insight*: Symplectic transvections have determinant 1, and they generate the symplectic group. □

## 3.6 Orthogonal Classification over Common Fields

**Theorem 3.6.1.** *Orthogonal Bases and Diagonal Congruence*
*Let $(\mathcal{V}, B)$ be an orthogonal geometry, $q(\mathbf{x}) := B(\mathbf{x}, \mathbf{x})$, $n = \dim \mathcal{V}$, and $r = \operatorname{rank} B$.*
*Then $\exists$ basis $\beta$ such that $M_\beta = \operatorname{diag}(a_1, \dots, a_r, 0_{n-r})$, $\quad a_i \in \mathbb{F}^\times \iff \neg\big(\operatorname{char}(\mathbb{F}) = 2 \ \wedge\ q \equiv 0 \ \wedge\ B \neq 0\big)$.*
*Equivalently, a symmetric matrix is congruent to a diagonal matrix iff it is not a nonzero alternating matrix in characteristic 2.*

*Proof.*

(⇒) If $M_\beta$ is diagonal and $\operatorname{char}(\mathbb{F}) = 2$ with $q \equiv 0$, then all diagonal entries of $M_\beta$ vanish. Hence $M_\beta = 0$, so $B = 0$.

(⇐) If $B = 0$, any basis works. Otherwise, the hypothesis and Proposition 3.1.3 give $q \not\equiv 0$.
Peel off nonisotropic lines, where $\mathcal{V} = \mathbb{F}\mathbf{u}_1 \odot \cdots \odot \mathbb{F}\mathbf{u}_s \odot \mathcal{W}$, $q(\mathbf{u}_i) = a_i \in \mathbb{F}^\times$, $q|_{\mathcal{W}} \equiv 0$.
If $\operatorname{char}(\mathbb{F}) \neq 2$, Proposition 3.1.3 gives $B|_{\mathcal{W}} = 0$, so append any basis of $\mathcal{W}$.
If $\operatorname{char}(\mathbb{F}) = 2$, then $\mathcal{W} = R \odot \mathcal{H}_1 \odot \cdots \odot \mathcal{H}_t$ by Theorem 3.5.1.
For each hyperbolic plane $\mathcal{H}_j = \operatorname{span}(\mathbf{e}, \mathbf{f})$, rescale so $B(\mathbf{e}, \mathbf{f}) = a = q(\mathbf{u})$ for some current nonisotropic $\mathbf{u}$.
Then $\mathbb{F}\mathbf{u} \odot \mathcal{H}_j = \mathbb{F}(\mathbf{u} + \mathbf{e}) \odot \mathbb{F}(\mathbf{u} + \mathbf{f}) \odot \mathbb{F}(\mathbf{u} + \mathbf{e} + \mathbf{f})$ is orthogonal with all three norms $a$.
Absorb all $\mathcal{H}_j$ and append any basis of $R$.

*Structural Commentary*: Only nonzero alternating characteristic-2 blocks obstruct diagonals.
*Key Insight*: Peel off nonisotropic lines until only the radical or characteristic-2 alternating residue remains. □

**Example 3.6.2.** *Characteristic Two Orthogonal-Symplectic Plane*
Assume $\operatorname{char}(\mathbb{F}) = 2$. Let $\mathcal{H} = \mathbb{F}\mathbf{e} \oplus \mathbb{F}\mathbf{f}$ with Gram matrix $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.
Then $B$ is symmetric, alternating, and nonsingular. For every $a, b \in \mathbb{F}$, $B(a\mathbf{e} + b\mathbf{f}, a\mathbf{e} + b\mathbf{f}) = 2ab = 0$, so $\mathcal{H}$ is both orthogonal and symplectic. If $0 \neq \mathbf{x} \in \mathcal{H}$, then $\{\mathbf{x}\}^\perp = \mathbb{F}\mathbf{x}$; hence no two independent nonzero vectors are orthogonal. Thus $\mathcal{H}$ has no orthogonal basis, the obstruction excluded in Theorem 3.6.1.

**Method 3.6.3.** *Lagrange Orthogonalization*
Let $\mathcal{V}$ be an orthogonal geometry with $\operatorname{char}(\mathbb{F}) \neq 2$, and set $q(\mathbf{x}) = \langle \mathbf{x}, \mathbf{x} \rangle$.
From a generating list $L = (\mathbf{v}_1, \dots, \mathbf{v}_n)$, construct an orthogonal basis as follows.

(i) In the current summand with generators $(\mathbf{x}_i)$, choose a pivot $\mathbf{u}$ with $q(\mathbf{u}) \neq 0$; if none exists, stop with a totally degenerate remainder. If all $\mathbf{x}_i$ are isotropic but the summand is not totally degenerate, take $\mathbf{u} = \mathbf{x}_i + \mathbf{x}_j$ with $\langle \mathbf{x}_i, \mathbf{x}_j \rangle \neq 0$.

(ii) Given mutually orthogonal pivots $\mathbf{u}_1, \dots, \mathbf{u}_k$, let $S_k = \operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_k)$.
Replace each generator $\mathbf{x}_i$ by $\mathbf{w}_i = \mathbf{x}_i - \sum_{j=1}^{k} \frac{\langle \mathbf{x}_i, \mathbf{u}_j \rangle}{q(\mathbf{u}_j)} \mathbf{u}_j$. Then the nonzero $\mathbf{w}_i$ generate $S_k^\perp$.

(iii) Iterate inside $S_k^\perp$ until the remaining summand $R$ is totally degenerate.
Output $\beta = (\mathbf{u}_1, \dots, \mathbf{u}_r, \mathbf{z}_1, \dots, \mathbf{z}_m)$, where $(\mathbf{z}_1, \dots, \mathbf{z}_m)$ is any basis of $R$.

Only nonisotropic pivots are allowed, since the projection step divides by $q(\mathbf{u}_j) \neq 0$.

**Theorem 3.6.4.** *Canonical Orthogonal Forms over Common Fields*
*Let $M = M^t \in M_n(\mathbb{F})$, let $r = \operatorname{rank} M$, and write $M \sim N$ for congruence. If $\operatorname{char}(\mathbb{F}) = 2$, assume $M$ is not nonzero alternating.*

*(i) If $\mathbb{F}$ is algebraically closed, then $M \sim I_r \oplus 0_{n-r}$; $r$ is complete.*
*(ii) If $\mathbb{F} = \mathbb{R}$, then $M \sim I_p \oplus (-I_m) \oplus 0_k$ with $p + m + k = n$; $(p, m)$ is complete.*
*(iii) If $\mathbb{F} = \mathbb{F}_q$ and $\operatorname{char}(\mathbb{F}) = 2$, then $M \sim I_r \oplus 0_{n-r}$.*
*(iv) If $\mathbb{F} = \mathbb{F}_q$ has odd characteristic and $d \in \mathbb{F}_q^\times$ is a fixed nonsquare, then for $r > 0$, $M \sim I_r \oplus 0_{n-r}$ or $M \sim I_{r-1} \oplus [d] \oplus 0_{n-r}$ with exactly one alternative; for $r = 0$, $M \sim 0_n$.*

*Proof.*

(i) By Theorem 3.6.1, write $M \sim \operatorname{diag}(a_1, \dots, a_r, 0_{n-r})$ with $a_i \neq 0$.
If $\mathbb{F}$ is algebraically closed, $a_i = s_i^2$, so scaling by $s_i^{-1}$ gives $M \sim I_r \oplus 0_{n-r}$.

(ii) Write $M \sim \operatorname{diag}(a_1, \dots, a_r, 0_k)$ over $\mathbb{R}$. Rescaling gives $a_i = \pm 1$, hence $M \sim I_p \oplus (-I_m) \oplus 0_k$.
Uniqueness of $(p, m)$ is Sylvester's inertia theorem, Theorem 2.2.9.

(iii) Write $M \sim \operatorname{diag}(a_1, \dots, a_r, 0_{n-r})$ over $\mathbb{F}_q$ with $\operatorname{char}(\mathbb{F}_q) = 2$.
Since $x \mapsto x^2$ is an automorphism, each $a_i$ is a square; hence $M \sim I_r \oplus 0_{n-r}$.

(iv) Write $M \sim \operatorname{diag}(a_1, \dots, a_r, 0_{n-r})$ over odd $\mathbb{F}_q$. Since $\mathbb{F}_q^\times/(\mathbb{F}_q^\times)^2 = \{1, d\}$, the nonsingular diagonal part is determined by its discriminant square class $\delta(M) := a_1 \cdots a_r \pmod{(\mathbb{F}_q^\times)^2}$, giving $I_r$ when $\delta(M) = 1$ and $I_{r-1} \oplus [d]$ when $\delta(M) = d$. Appending the zero block gives the stated forms, and $\delta$ separates them.

*Structural Commentary*: Once diagonalized, congruence classification becomes square-class bookkeeping.
*Key Insight*: The base field determines how many diagonal coefficients can be normalized to 1. □

**Definition 3.6.5.** *Universal Orthogonal Geometry*
An orthogonal geometry $\mathcal{V}$ over $\mathbb{F}$ is *universal* if its norm map $q(\mathbf{x}) := \langle \mathbf{x}, \mathbf{x} \rangle$ represents every scalar $q(\mathcal{V}) = \mathbb{F}$.

**Proposition 3.6.6.** *Finite-Field Square Classes and Universality*
*Let $\mathbb{F}_q$ be a finite field and let $S = (\mathbb{F}_q^\times)^2$.*

*(i) If $\operatorname{char}(\mathbb{F}_q) = 2$, then every element of $\mathbb{F}_q$ is a square.*
*(ii) If $\operatorname{char}(\mathbb{F}_q) \neq 2$, then $|S| = (q-1)/2$ and $\mathbb{F}_q^\times \setminus S = dS$ for any nonsquare $d$.*
*(iii) If $\mathcal{V}$ is orthogonal over odd $\mathbb{F}_q$ and contains a nonsingular subspace of dimension $\geq 2$, then $\mathcal{V}$ is universal.*

*Proof.*

(i) The map $\varphi : \mathbb{F}_q \to \mathbb{F}_q$, $\varphi(x) = x^2$, is the Frobenius automorphism in characteristic 2.

(ii) For $\operatorname{char}(\mathbb{F}_q) \neq 2$, the homomorphism $\varphi : \mathbb{F}_q^\times \to \mathbb{F}_q^\times$, $\varphi(x) = x^2$, has kernel $\{\pm 1\}$.
Hence $|S| = (q-1)/2$, and the nonsquares are the other coset $dS$.

(iii) Let $q(\mathbf{x}) = \langle \mathbf{x}, \mathbf{x} \rangle$. By Theorem 3.6.1, the nonsingular subspace contains orthogonal nonisotropic $\mathbf{u}, \mathbf{v}$; set $a = q(\mathbf{u})$, $b = q(\mathbf{v})$. For fixed $c \in \mathbb{F}_q$, the sets $A = a\{\alpha^2 : \alpha \in \mathbb{F}_q\}$ and $B = c - b\{\beta^2 : \beta \in \mathbb{F}_q\}$ both have $(q+1)/2$ elements, so $A \cap B \neq \varnothing$. Thus $c = a\alpha^2 + b\beta^2 = q(\alpha\mathbf{u} + \beta\mathbf{v})$ for some $\alpha, \beta \in \mathbb{F}_q$.

*Structural Commentary*: Over finite fields, cardinality replaces order arguments.
*Key Insight*: Two half-sized subsets of a finite field must overlap. □

**Example 3.6.7.** *Two Rank-Two Orthogonal Planes over $\mathbb{F}_5$*
In $\mathbb{F}_5$, $(\mathbb{F}_5^\times)^2 = \{1, 4\}$ and the nonsquare class is $\{2, 3\}$. Let $q_1(x, y) = x^2 + y^2$, $q_2(x, y) = x^2 + 2y^2$.
Then $q_1$ has Gram matrix $I_2$ and is hyperbolic, since $q_1(1, 2) = 1 + 4 = 0$.

The form $q_2$ has Gram matrix $\operatorname{diag}(1, 2)$ and is anisotropic: if $q_2(x, y) = 0$ with $y \neq 0$, then $\left(\frac{x}{y}\right)^2 = -2 = 3$, impossible since 3 is nonsquare; if $y = 0$, then $x = 0$.
Thus $I_2$ and $\operatorname{diag}(1, 2)$ realize the two nonsingular rank-2 classes in Theorem 3.6.4.
*Structural Commentary*: Odd finite-field classes are square-class classes.
*Key Insight*: In rank 2, the square class of the determinant decides whether the plane is hyperbolic or anisotropic.

**Lemma 3.6.8.** *Odd Finite-Field Isotropy Bound*
*Let $\mathbb{F}_q$ have odd characteristic, $\mathcal{V}$ be nonsingular orthogonal with $q(\mathbf{x}) := \langle \mathbf{x}, \mathbf{x} \rangle$. Let $\mathcal{H}$ be hyperbolic plane.*

*(i) If $\dim \mathcal{V} \geq 3$, then $\mathcal{V}$ is isotropic.*
*(ii) $\mathcal{V} \cong \mathcal{H}^{\odot s} \odot \mathcal{R}$, where $\mathcal{R}$ is nonsingular anisotropic and $\dim \mathcal{R} \leq 2$.*

*Proof.*

(i) By Theorem 3.6.1, choose a nonsingular plane $\mathcal{P} \subset \mathcal{V}$ and $0 \neq \mathbf{z} \in \mathcal{P}^\perp$. If $q(\mathbf{z}) = 0$, then $\mathbf{z}$ is isotropic. Otherwise Proposition 3.6.6 gives $\mathbf{p} \in \mathcal{P}$ with $q(\mathbf{p}) = -q(\mathbf{z})$, so $\mathbf{p} + \mathbf{z}$ is isotropic.

(ii) Induct on $\dim \mathcal{V}$. If $\mathcal{V}$ is anisotropic, take $s = 0$ and $\mathcal{R} = \mathcal{V}$; then (i) gives $\dim \mathcal{R} \leq 2$. Otherwise Theorem 3.4.8 gives $\mathcal{V} \cong \mathcal{H} \odot \mathcal{W}$ with $\mathcal{W}$ nonsingular and $\dim \mathcal{W} = \dim \mathcal{V} - 2$, so induction gives the decomposition.

*Structural Commentary*: Odd finite fields force isotropy in dimension $\geq 3$.
*Key Insight*: Hyperbolic splitting leaves only a residue of dimension at most 2. □

**Theorem 3.6.9.** *Odd Finite-Field Orthogonal Decomposition*
*Let $\mathbb{F}_q$ have odd characteristic, fix a nonsquare $d \in \mathbb{F}_q^\times$, and let $\mathcal{V}$ be nonsingular orthogonal.*
*Write $\mathcal{H}$ for a hyperbolic plane, $\langle a \rangle$ for a line of value $a$, and $\mathcal{A}_d := \langle 1 \rangle \odot \langle -d \rangle$.*

*(i) If $\dim \mathcal{V} = 2m + 1$, then $\exists\, a \in \mathbb{F}_q^\times$ such that $\mathcal{V} \cong \mathcal{H}^{\odot m} \odot \langle a \rangle$, and $a(\mathbb{F}_q^\times)^2 = (-1)^m \Delta(\mathcal{V})$.*
*(ii) If $\dim \mathcal{V} = 2m$, then exactly one holds: $\mathcal{V} \cong \mathcal{H}^{\odot m} \iff \Delta(\mathcal{V}) = (-1)^m (\mathbb{F}_q^\times)^2$*
*or $\mathcal{V} \cong \mathcal{H}^{\odot(m-1)} \odot \mathcal{A}_d \iff \Delta(\mathcal{V}) = (-1)^m d(\mathbb{F}_q^\times)^2$.*

*Proof.*

(i) By Lemma 3.6.8, $\mathcal{V} \cong \mathcal{H}^{\odot m} \odot \mathcal{R}$ with $\dim \mathcal{R} = 1$, so $\mathcal{R} \cong \langle a \rangle$.
Since $\Delta(\mathcal{H}) = -(\mathbb{F}_q^\times)^2$, $\Delta(\mathcal{V}) = (-1)^m a(\mathbb{F}_q^\times)^2$, hence $a(\mathbb{F}_q^\times)^2 = (-1)^m \Delta(\mathcal{V})$.

(ii) By Lemma 3.6.8, $\mathcal{V} \cong \mathcal{H}^{\odot s} \odot \mathcal{R}$ with $\dim \mathcal{R} = 0$ or 2. If $\dim \mathcal{R} = 0$, then $s = m$ and $\Delta(\mathcal{V}) = (-1)^m (\mathbb{F}_q^\times)^2$. If $\dim \mathcal{R} = 2$, then $s = m - 1$ and Proposition 3.4.9 shows that the anisotropic residue is the nonsquare class $\mathcal{A}_d$, with $\Delta(\mathcal{A}_d) = -d(\mathbb{F}_q^\times)^2$; hence $\Delta(\mathcal{V}) = (-1)^m d(\mathbb{F}_q^\times)^2$.

*Structural Commentary*: The anisotropic residue has dimension $\leq 2$.
*Key Insight*: Parity and discriminant determine the residue. □

**Definition 3.6.10.** *Minkowski Space*
*Minkowski space* is the orthogonal geometry $\mathbb{R}^{3,1} := \mathbb{R}^4$ with standard basis $(\mathbf{e}_1, \dots, \mathbf{e}_4)$ and bilinear form $\langle \mathbf{x}, \mathbf{y} \rangle = x_1 y_1 + x_2 y_2 + x_3 y_3 - x_4 y_4$.
Equivalently, its Gram matrix is $\operatorname{diag}(1, 1, 1, -1)$. It is nonsingular but not positive definite.

**Example 3.6.11.** *Null Vectors in Minkowski Space*
In $\mathbb{R}^{3,1}$, set $q(\mathbf{x}) := \langle \mathbf{x}, \mathbf{x} \rangle = x_1^2 + x_2^2 + x_3^2 - x_4^2$.
Null cone is $\mathcal{N} := \{\mathbf{x} \neq \mathbf{0} : q(\mathbf{x}) = 0\} = \{\mathbf{x} \neq \mathbf{0} : x_1^2 + x_2^2 + x_3^2 = x_4^2\}$. Every $\mathbf{x} \in \mathcal{N}$ is isotropic; i.e., $\mathbf{e}_1 + \mathbf{e}_4 \in \mathcal{N}$.
If $\mathbb{R}^{3,1}$ were isometric to Euclidean $\mathbb{R}^4$, the isometry matrix $A$ in the standard bases would satisfy $A^t I_4 A = \operatorname{diag}(1, 1, 1, -1)$. Taking determinants gives $(\det A)^2 = -1$, impossible over $\mathbb{R}$.

## 3.7 Quadratic Refinements over $\mathbb{F}_2$

**Example 3.7.1.** *Quadratic Refinements over $\mathbb{F}_2$*
On $V = \mathbb{F}_2^2$, set $B((x, y), (u, v)) = xv + yu$, $Q_0(x, y) = xy$, $Q_1(x, y) = x^2 + xy + y^2$. Then $B_{Q_0} = B_{Q_1} = B$, the unique nonsingular alternating plane over $\mathbb{F}_2$. However, $Q_0^{-1}(0) \setminus \{\mathbf{0}\} = \{(1, 0), (0, 1)\}$, $Q_1^{-1}(0) = \{\mathbf{0}\}$.
Hence $Q_0 \ncong Q_1$ although they have the same polar form.
*Structural Commentary*: One polar form, two refinements.
*Key Insight*: Track $Q$, not only $B_Q$.

**Definition 3.7.2.** *Arf Invariant over $\mathbb{F}_2$*
Let $Q : \mathcal{V} \to \mathbb{F}_2$ be quadratic.

(i) Its *polar form* is $B_Q(\mathbf{x}, \mathbf{y}) = Q(\mathbf{x} + \mathbf{y}) + Q(\mathbf{x}) + Q(\mathbf{y})$ and is assumed nonsingular.
(ii) For a *symplectic basis* $\beta = (\mathbf{u}_1, \mathbf{v}_1, \dots, \mathbf{u}_m, \mathbf{v}_m)$ of $B_Q$,
the *Arf invariant* is $\operatorname{Arf}(Q) := \sum_{i=1}^{m} Q(\mathbf{u}_i) Q(\mathbf{v}_i) \in \mathbb{F}_2$, independent of $\beta$.

**Theorem 3.7.3.** *Quadratic Refinements over $\mathbb{F}_2$*
*Let $Q : \mathcal{V} \to \mathbb{F}_2$ be quadratic with nonsingular polar form $B_Q$. Then $\dim \mathcal{V} = 2m$, and $Q \cong Q_{\operatorname{Arf}(Q)}$.*
*Here $Q_0(x_1, y_1, \dots, x_m, y_m) := \sum_{i=1}^{m} x_i y_i$ and, for $m \geq 1$, $Q_1(x_1, y_1, \dots, x_m, y_m) := x_1^2 + x_1 y_1 + y_1^2 + \sum_{i=2}^{m} x_i y_i$.*
*$Q_0$ has Arf invariant 0, and $Q_1$ has Arf invariant 1.*

*Proof.* As $B_Q$ nonsingular alternating, Theorem 3.5.1 gives symplectic basis $(\mathbf{u}_1, \mathbf{v}_1, \dots, \mathbf{u}_m, \mathbf{v}_m)$, $\dim \mathcal{V} = 2m$.
On each hyperbolic pair $(\mathbf{u}, \mathbf{v})$, if $Q(\mathbf{u})Q(\mathbf{v}) = 0$, a change of pair gives the refined plane $xy$.
If $Q(\mathbf{u}) = Q(\mathbf{v}) = 1$, the plane is $x^2 + xy + y^2$.
Thus $Q$ is an orthogonal sum of refined planes of types $H_Q := xy$ and $A_Q := x^2 + xy + y^2$.
Since $A_Q \odot A_Q \cong H_Q \odot H_Q$, all pairs of anisotropic refinements cancel.
Remaining parity is $\operatorname{Arf}(Q) = \sum_{i=1}^{m} Q(\mathbf{u}_i) Q(\mathbf{v}_i) \in \mathbb{F}_2$, so $Q \cong Q_0$ when $\operatorname{Arf}(Q) = 0$ and $Q \cong Q_1$ when $\operatorname{Arf}(Q) = 1$.
*Structural Commentary*: In characteristic 2, the same symplectic polar form supports two quadratic geometries.
*Key Insight*: The Arf invariant is the missing one-bit datum that the polar bilinear form cannot see. □

## 3.8 Orthogonal Isometries and Symmetry Groups

**Proposition 3.8.1.** *Isometries from Quadratic Values*
*Let $\mathcal{V}$ be an orthogonal geometry with $\operatorname{char}(\mathbb{F}) \neq 2$, let $q(\mathbf{x}) := \langle \mathbf{x}, \mathbf{x} \rangle$, and let $T : \mathcal{V} \to \mathcal{V}$ be bijective. Then $T$ is an isometry if and only if $q(T\mathbf{x}) = q(\mathbf{x})$ for all $\mathbf{x} \in \mathcal{V}$.*

*Proof.*
($\Rightarrow$) If $T$ is an isometry, then $q(T\mathbf{x}) = \langle T\mathbf{x}, T\mathbf{x} \rangle = \langle \mathbf{x}, \mathbf{x} \rangle = q(\mathbf{x})$.
($\Leftarrow$) If $q \circ T = q$, then $\langle T\mathbf{x}, T\mathbf{y} \rangle = \frac{q(T\mathbf{x}+T\mathbf{y}) - q(T\mathbf{x}) - q(T\mathbf{y})}{2} = \frac{q(\mathbf{x}+\mathbf{y}) - q(\mathbf{x}) - q(\mathbf{y})}{2} = \langle \mathbf{x}, \mathbf{y} \rangle$.
Thus $T$ preserves the form, so it is an isometry.

*Structural Commentary*: In odd characteristic, $q$ determines the orthogonal form.
*Key Insight*: Polarization turns preservation of lengths into preservation of inner products. □

**Definition 3.8.2.** *Rotations, Reflections, and Symmetries*
Let $\mathcal{V}$ be a nonsingular orthogonal geometry.

(i) For $\sigma \in \mathcal{O}(\mathcal{V})$, $\det \sigma = 1$ means $\sigma$ is a *rotation*, and $\det \sigma = -1$ means $\sigma$ is a *reflection*.

(ii) If $\operatorname{char}(\mathbb{F}) \neq 2$ and $q(\mathbf{u}) = \langle \mathbf{u}, \mathbf{u} \rangle \neq 0$, the *symmetry* in the line $\mathbb{F}\mathbf{u}$ is $\sigma_{\mathbf{u}}(\mathbf{x}) = \mathbf{x} - 2\frac{\langle \mathbf{x}, \mathbf{u} \rangle}{\langle \mathbf{u}, \mathbf{u} \rangle}\mathbf{u}$.
It satisfies $\sigma_{\mathbf{u}}\mathbf{u} = -\mathbf{u}$ and $\sigma_{\mathbf{u}}|_{\operatorname{span}(\mathbf{u})^\perp} = I$.

**Proposition 3.8.3.** *Matching Equal-Length Vectors by a Symmetry*
*Let $\mathcal{V}$ be nonsingular orthogonal with $\operatorname{char}(\mathbb{F}) \neq 2$, and set $q(\mathbf{x}) = \langle \mathbf{x}, \mathbf{x} \rangle$.*
*If $q(\mathbf{u}) = q(\mathbf{v}) \neq 0$, then $\exists$ a symmetry $\sigma$ with $\sigma\mathbf{u} = \mathbf{v}$ or $\sigma\mathbf{u} = -\mathbf{v}$.*

*Proof.* Let $\mathbf{a} = \mathbf{u} - \mathbf{v}$ and $\mathbf{b} = \mathbf{u} + \mathbf{v}$. Then $\langle \mathbf{a}, \mathbf{b} \rangle = q(\mathbf{u}) - q(\mathbf{v}) = 0$.
They cannot both be isotropic: if $q(\mathbf{a}) = q(\mathbf{b}) = 0$, then $q(2\mathbf{u}) = 0$, contradicting $q(\mathbf{u}) \neq 0$.
If $q(\mathbf{a}) \neq 0$, then $\sigma_{\mathbf{a}}\mathbf{u} = \mathbf{v}$; if $q(\mathbf{b}) \neq 0$, then $\sigma_{\mathbf{b}}\mathbf{u} = -\mathbf{v}$.
*Structural Commentary*: One symmetry matches equal-length lines.
*Key Insight*: Use $\mathbf{u} \pm \mathbf{v}$ as the reflecting direction. □

**Theorem 3.8.4.** *Orthogonal Transformations Are Products of Symmetries*
*Let $\mathcal{V}$ be nonsingular orthogonal with $\operatorname{char}(\mathbb{F}) \neq 2$, and set $q(\mathbf{x}) := \langle \mathbf{x}, \mathbf{x} \rangle$. Then $\mathcal{O}(\mathcal{V}) = \langle \sigma_{\mathbf{u}} : q(\mathbf{u}) \neq 0 \rangle$.*

*Proof.* Induct on $n = \dim \mathcal{V}$.
If $n = 0$, this is trivial; if $n = 1$, any $\tau \in \mathcal{O}(\mathcal{V})$ acts on a nonisotropic line by $\pm 1$, hence is $I$ or the symmetry of that line. For $n > 1$, choose $\mathbf{u}$ with $q(\mathbf{u}) \neq 0$. Since $q(\tau\mathbf{u}) = q(\mathbf{u})$, Proposition 3.8.3 gives a symmetry $\sigma$ with $\sigma\tau\mathbf{u} = \mathbf{u}$ or $-\mathbf{u}$. Composing further with $\sigma_{\mathbf{u}}$ if necessary, obtain a product $\rho$ of symmetries such that $\rho\tau$ fixes $\mathbf{u}$. Hence $\rho\tau$ preserves $\operatorname{span}(\mathbf{u})^\perp$, and its restriction lies in $\mathcal{O}(\operatorname{span}(\mathbf{u})^\perp)$.
By induction, this restriction is a product of symmetries in $\operatorname{span}(\mathbf{u})^\perp$, extended by $I$ on $\mathbb{F}\mathbf{u}$; these are symmetries of $\mathcal{V}$. Thus $\tau$ is a product of symmetries.
*Structural Commentary*: Symmetries are the orthogonal analogue of elementary generators.
*Key Insight*: Match one nonisotropic line, then reduce to its orthogonal complement. □

**Proposition 3.8.5.** *Fixed Spaces of Isometries*
*Let $\mathcal{V}$ be nonsingular and let $\tau \in \operatorname{Iso}(\mathcal{V})$. Then $\ker(I - \tau) = \operatorname{im}(I - \tau)^\perp$, and $\dim \ker(I - \tau) = \dim \operatorname{im}(I - \tau)^\perp$.*

*Proof.* If $\mathbf{x}$ is fixed by $\tau$, then for every $\mathbf{y}$, $\langle \mathbf{x}, (I - \tau)\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle - \langle \tau\mathbf{x}, \tau\mathbf{y} \rangle = 0$, so $\ker(I - \tau) \subseteq \operatorname{im}(I - \tau)^\perp$.
The dimensions agree by rank-nullity and Theorem 3.4.1, hence equality holds.
*Structural Commentary*: The fixed space of an isometry is the orthogonal complement of its displacement space.
*Key Insight*: Fixed vectors are exactly vectors orthogonal to all displacements. □

**Proposition 3.8.6.** *Symmetries and Fixed Hyperplanes*
*Let $\mathcal{V}$ be nonsingular orthogonal with $\operatorname{char}(\mathbb{F}) \neq 2$, and let $\tau \in \mathcal{O}(\mathcal{V})$ with $\tau \neq I$. Then $\tau = \sigma_{\mathbf{u}}$ for some nonisotropic $\mathbf{u}$ $\iff$ $\ker(I - \tau)$ is a nonsingular hyperplane. In this case $\ker(I - \sigma_{\mathbf{u}}) = \operatorname{span}(\mathbf{u})^\perp$.*

*Proof.*

($\Rightarrow$) The formula for $\sigma_{\mathbf{u}}$ gives $\sigma_{\mathbf{u}}\mathbf{u} = -\mathbf{u}$ and fixes $\operatorname{span}(\mathbf{u})^\perp$ pointwise.
Hence $\ker(I - \sigma_{\mathbf{u}}) = \operatorname{span}(\mathbf{u})^\perp$, a nonsingular hyperplane.

($\Leftarrow$) Let $\mathcal{H} = \ker(I - \tau)$ be a nonsingular hyperplane and write $\mathcal{V} = \mathcal{H} \odot \mathcal{L}$ with $\dim \mathcal{L} = 1$.
Since $\tau$ fixes $\mathcal{H}$ and is an isometry, $\tau\mathcal{L} = \mathcal{L}$. Thus $\tau|_{\mathcal{L}} = \lambda I$ with $\lambda^2 = 1$; as $\tau \neq I$ and $\operatorname{char}(\mathbb{F}) \neq 2$, $\lambda = -1$. For $0 \neq \mathbf{u} \in \mathcal{L}$, $\tau = \sigma_{\mathbf{u}}$.

*Structural Commentary*: Symmetries are exactly fixed-hyperplane isometries.
*Key Insight*: A symmetry fixes a nonsingular hyperplane and flips its orthogonal line. □

**Proposition 3.8.7.** *No Unipotent Isometries in Anisotropic Spaces*
*Let $\mathcal{V}$ be nonsingular anisotropic. If $\tau \in \operatorname{Iso}(\mathcal{V})$ is unipotent, then $\tau = I$.*

*Proof.* Put $N = \tau - I$. Then $N$ is nilpotent and, by Proposition 3.3.9, $N^* = \tau^* - I = \tau^{-1} - I \in N\mathbb{F}[N]$.
If $N \neq 0$, choose maximal $k \geq 1$ with $N^k \neq 0$ and choose $\mathbf{x}$ with $\mathbf{u} = N^k\mathbf{x} \neq 0$.
Since $N^{k+1} = 0$ and $(N^*)^k \in N^k\mathbb{F}[N]$, $\langle \mathbf{u}, \mathbf{u} \rangle = \langle N^k\mathbf{x}, N^k\mathbf{x} \rangle = \langle \mathbf{x}, (N^*)^k N^k\mathbf{x} \rangle = 0$, contradicting anisotropy.
Hence $N = 0$ and $\tau = I$.
*Structural Commentary*: Top nilpotent displacements are isotropic.
*Key Insight*: An anisotropic form leaves no room for unipotent shear. □

Theorem 3.4.13 and Corollary 3.4.14 specialize to nonsingular orthogonal geometries in characteristic not equal to 2.

# 4 Affine, Projective, and Convex Geometry

## 4.1 Affine Flats, Hulls, and Incidence

Affine geometry studies cosets of subspaces as geometric objects in their own right. Throughout this subsection, vector spaces are finite-dimensional over $\mathbb{F}$ unless explicitly stated otherwise.

**Definition 4.1.1.** *Flats* Let $\mathcal{S} \leq \mathcal{V}$ and $\mathbf{x} \in \mathcal{V}$, and set $\mathbf{x}+\mathcal{S} = \{\mathbf{x}+\mathbf{s} : \mathbf{s} \in \mathcal{S}\}$.

(i) The affine subset $\mathbf{x}+\mathcal{S}$ from Definition 2.2.1 is a *flat* in $\mathcal{V}$; $\mathcal{S}$ is its *base*, and $\mathbf{x}$ is a *flat representative*.
(ii) Let $\mathcal{A}(\mathcal{V})$ denote the set of all flats in $\mathcal{V}$, with $\dim(\mathcal{A}(\mathcal{V})) := \dim(\mathcal{V})$.
(iii) The *dimension* of $\mathbf{x}+\mathcal{S}$ is $\dim(\mathcal{S})$.
A 0-flat is a *point*, a 1-flat is a *line*, a 2-flat is a *plane*, and a codimension-one flat is a *hyperplane*.
(iv) Two flats $X = \mathbf{x}+\mathcal{S}$ and $Y = \mathbf{y}+\mathcal{T}$ are *parallel*, written $X \parallel Y$, if $\mathcal{S} \subseteq \mathcal{T}$ or $\mathcal{T} \subseteq \mathcal{S}$.

*Structural Commentary*: A flat has many representatives but one base: $\mathbf{x}+\mathcal{S} = \mathbf{y}+\mathcal{T} \Rightarrow \mathcal{S} = \mathcal{T}$.

**Proposition 4.1.2.** *Basic Incidence of Flats*
*Let $X = \mathbf{x}+\mathcal{S}$ and $Y = \mathbf{y}+\mathcal{T}$ be flats in $\mathcal{V}$, and write $\mathbf{w}+X := \{\mathbf{w}+\mathbf{u} : \mathbf{u} \in X\}$.*

*(i)* $(\exists\, \mathbf{w} \in \mathcal{V})(\mathbf{w}+X \subseteq Y) \iff \mathcal{S} \subseteq \mathcal{T}$.
*(ii)* $(\exists\, \mathbf{w} \in \mathcal{V})(\mathbf{w}+X = Y) \iff \mathcal{S} = \mathcal{T}$.
*(iii) If $X \cap Y \neq \emptyset$, then $X \subseteq Y \iff \mathcal{S} \subseteq \mathcal{T}$.*
*(iv) If $X \cap Y \neq \emptyset$, then $X = Y \iff \mathcal{S} = \mathcal{T}$.*
*(v) If $X \parallel Y$, then $X \subseteq Y$, or $Y \subseteq X$, or $X \cap Y = \emptyset$.*
*(vi)* $X \parallel Y \iff [(\exists\, \mathbf{w} \in \mathcal{V})(\mathbf{w}+X \subseteq Y)] \vee [(\exists\, \mathbf{w} \in \mathcal{V})(\mathbf{w}+Y \subseteq X)]$.

*Proof.*

(i) Since $\mathbf{w}+X = (\mathbf{w}+\mathbf{x})+\mathcal{S}$, every translate of $X$ has base $\mathcal{S}$. If $\mathbf{w}+X \subseteq Y$, choose $\mathbf{z} \in \mathbf{w}+X$; then $\mathbf{w}+X = \mathbf{z}+\mathcal{S}$ and $Y = \mathbf{z}+\mathcal{T}$, hence $\mathcal{S} \subseteq \mathcal{T}$. Conversely, if $\mathcal{S} \subseteq \mathcal{T}$, then $(\mathbf{y}-\mathbf{x})+X = \mathbf{y}+\mathcal{S} \subseteq Y$.
(ii) If $\mathbf{w}+X = Y$, then the bases are equal. Conversely, if $\mathcal{S} = \mathcal{T}$, then $(\mathbf{y}-\mathbf{x})+X = Y$.
(iii) Choose $\mathbf{z} \in X \cap Y$. Then $X = \mathbf{z}+\mathcal{S}$ and $Y = \mathbf{z}+\mathcal{T}$, so translating by $-\mathbf{z}$ gives $X \subseteq Y \iff \mathcal{S} \subseteq \mathcal{T}$.
(iv) With the same $\mathbf{z}$, translate by $-\mathbf{z}$ to get $X = Y \iff \mathcal{S} = \mathcal{T}$.
(v) If $X \parallel Y$, then $\mathcal{S} \subseteq \mathcal{T}$ or $\mathcal{T} \subseteq \mathcal{S}$. If $X \cap Y \neq \emptyset$, part (iii) gives $X \subseteq Y$ or $Y \subseteq X$; otherwise $X \cap Y = \emptyset$.
(vi) If $X \parallel Y$, then either $\mathcal{S} \subseteq \mathcal{T}$ or $\mathcal{T} \subseteq \mathcal{S}$, so part (i) gives a translate $X' \subseteq Y$ or a translate $Y' \subseteq X$. Conversely, either containment gives $\mathcal{S} \subseteq \mathcal{T}$ or $\mathcal{T} \subseteq \mathcal{S}$ by part (i), hence $X \parallel Y$.

*Structural Commentary*: Intersecting flats reduce affine incidence to subspace incidence.
*Key Insight*: The representative may move, but the base controls containment, equality, and parallelism. □

**Definition 4.1.3.** *Affine Combinations*

(i) For $\mathbf{x}_1, \ldots, \mathbf{x}_n \in \mathcal{V}$ and $r_1, \ldots, r_n \in \mathbb{F}$, the sum $\sum_{j=1}^n r_j\mathbf{x}_j$ is an *affine combination* if $\sum_{j=1}^n r_j = 1$.
(ii) A nonempty $E \subseteq \mathcal{V}$ is *affine closed* if $\sum_{j=1}^n r_j\mathbf{x}_j \in E$ whenever $\mathbf{x}_j \in E$, $r_j \in \mathbb{F}$, and $\sum_{j=1}^n r_j = 1$.
(iii) A nonempty $E \subseteq \mathcal{V}$ is *two-affine closed* if $r\mathbf{x}+(1-r)\mathbf{y} \in E$ for all $\mathbf{x}, \mathbf{y} \in E$ and $r \in \mathbb{F}$.
(iv) For $\mathbf{x} \neq \mathbf{y}$, the affine line through them is $\overline{\mathbf{xy}} := \{r\mathbf{x}+(1-r)\mathbf{y} : r \in \mathbb{F}\} = \mathbf{y}+\operatorname{span}(\mathbf{x}-\mathbf{y})$.
Equivalently, a nonempty $E \subseteq \mathcal{V}$ is two-affine closed iff $\overline{\mathbf{xy}} \subseteq E$ for all distinct $\mathbf{x}, \mathbf{y} \in E$.

**Theorem 4.1.4.** *Two-Point Affine Closure*
*Assume $\operatorname{char}(\mathbb{F}) \neq 2$. For every $\emptyset \neq E \subseteq \mathcal{V}$, $E$ is affine closed $\iff$ $E$ is two-affine closed.*

*Proof.*

(⇒) A two-point affine combination is an affine combination, so affine closure implies two-affine closure.
(⇐) Assume $E$ is two-affine closed.

Prove by induction on $n$ that every affine combination $\sum_{j=1}^n r_j\mathbf{x}_j$, with $\mathbf{x}_j \in E$ and $\sum_{j=1}^n r_j = 1$, lies in $E$.

The cases $n = 1, 2$ are immediate. Let $n \geq 3$ and set $\mathbf{z} := \sum_{j=1}^n r_j\mathbf{x}_j$, $\sum_{j=1}^n r_j = 1$.

If some $r_j \neq 1$, relabel so $r_1 \neq 1$ and put $\mathbf{u} := \sum_{j=2}^n \frac{r_j}{1-r_1}\mathbf{x}_j$. Then $\sum_{j=2}^n \frac{r_j}{1-r_1} = 1$, so $\mathbf{u} \in E$ by induction, and $\mathbf{z} = r_1\mathbf{x}_1 + (1-r_1)\mathbf{u} \in E$ by two-affine closure.
Otherwise all $r_j = 1$; since 2 is invertible, $\mathbf{z} = 2\left(\frac{1}{2}\mathbf{x}_1 + \frac{1}{2}\mathbf{x}_2\right) + r_3\mathbf{x}_3 + \cdots + r_n\mathbf{x}_n$, where $\frac{1}{2}\mathbf{x}_1 + \frac{1}{2}\mathbf{x}_2 \in E$ by two-affine closure; this is an $(n-1)$-term affine combination, hence lies in $E$ by induction.

*Structural Commentary*: For $\operatorname{char}(\mathbb{F}) \neq 2$, two-point closure generates finite affine closure.
*Key Insight*: The only exceptional induction case is handled by averaging two coefficient-1 terms. □

**Remark 4.1.5.** *Characteristic Two Warning*
Condition $\operatorname{char}(\mathbb{F}) \neq 2$ is necessary: in $\mathbb{F}_2^2$, $E := \{(0,0),(1,0),(0,1)\}$ is two-affine closed, but $\operatorname{aff}(E) = \mathbb{F}_2^2 \neq E$.

**Theorem 4.1.6.** *Flats Are the Affine-Closed Sets*
*For every $\emptyset \neq E \subseteq \mathcal{V}$, $E \in \mathcal{A}(\mathcal{V}) \iff$ $E$ is affine closed.*

*Proof.*

(i) Suppose $E = \mathbf{x} + \mathcal{S}$ with $\mathcal{S} \leq \mathcal{V}$. If $\mathbf{x}_j = \mathbf{x} + \mathbf{s}_j \in E$ and $\sum_j r_j = 1$,
then $\sum_j r_j \mathbf{x}_j = \sum_j r_j(\mathbf{x} + \mathbf{s}_j) = \mathbf{x} + \sum_j r_j \mathbf{s}_j \in \mathbf{x} + \mathcal{S}$. Thus $E$ is affine closed.

(ii) Suppose $E$ is affine closed. Choose $\mathbf{x} \in E$ and set $\mathcal{S} := E - \mathbf{x} = \{\mathbf{e} - \mathbf{x} : \mathbf{e} \in E\}$.
For $\mathbf{s}_1, \mathbf{s}_2 \in \mathcal{S}$, write $\mathbf{s}_j = \mathbf{x}_j - \mathbf{x}$ with $\mathbf{x}_j \in E$.
For $r_1, r_2 \in \mathbb{F}$, $r_1\mathbf{s}_1 + r_2\mathbf{s}_2 + \mathbf{x} = r_1\mathbf{x}_1 + r_2\mathbf{x}_2 + (1 - r_1 - r_2)\mathbf{x} \in E$, so $r_1\mathbf{s}_1 + r_2\mathbf{s}_2 \in \mathcal{S}$.
Hence $\mathcal{S} \leq \mathcal{V}$ by Theorem 1.2.2, and $E = \mathbf{x} + \mathcal{S}$ is a flat.

*Structural Commentary*: An affine-closed set becomes a subspace after choosing one point as an origin.
*Key Insight*: Flats are precisely translated subspaces. □

**Theorem 4.1.7.** *Flats Are the Two-Affine-Closed Sets*
*Assume* $\operatorname{char}(\mathbb{F}) \neq 2$. *For every $\emptyset \neq E \subseteq \mathcal{V}$, $E \in \mathcal{A}(\mathcal{V}) \iff$ $E$ is two-affine closed.*

*Proof.*

(i) If $E \in \mathcal{A}(\mathcal{V})$, then $E$ is affine closed by Theorem 4.1.6, hence two-affine closed by Theorem 4.1.4.

(ii) If $E$ is two-affine closed, then $E$ is affine closed by Theorem 4.1.4, hence $E \in \mathcal{A}(\mathcal{V})$ by Theorem 4.1.6.

*Structural Commentary*: In characteristic not 2, two-point closure detects flats.
*Key Insight*: Two-affine closure equals finite affine closure, hence flatness. □

**Definition 4.1.8.** *Affine Hull and Affine Span*
Let $\emptyset \neq E \subseteq \mathcal{V}$.

(i) The *affine hull* $\operatorname{affhull}(E)$ is the least flat $X \in \mathcal{A}(\mathcal{V})$ with $E \subseteq X$.

(ii) The *affine span* $\operatorname{affspan}(E)$ is the set of all finite affine combinations of points of $E$.

**Theorem 4.1.9.** *Affine Hull Formula*
*Let $\emptyset \neq E \subseteq \mathcal{V}$, $\mathbf{x} \in E$, and $\mathcal{S}_\mathbf{x} := \operatorname{span}(E - \mathbf{x})$.*
*Then* $\operatorname{affhull}(E) = \operatorname{affspan}(E) = \mathbf{x} + \mathcal{S}_\mathbf{x}$, *and hence* $\dim(\operatorname{affhull}(E)) = \dim(\mathcal{S}_\mathbf{x})$.

*Proof.* If $\mathbf{z} = \sum_{j=1}^{n} r_j \mathbf{e}_j$ with $\mathbf{e}_j \in E$ and $\sum_j r_j = 1$, then $\mathbf{z} = \mathbf{x} + \sum_{j=1}^{n} r_j(\mathbf{e}_j - \mathbf{x}) \in \mathbf{x} + \mathcal{S}_\mathbf{x}$, so $\operatorname{affspan}(E) \subseteq \mathbf{x} + \mathcal{S}_\mathbf{x}$.
Conversely, for $\sum_{j=1}^{n} a_j(\mathbf{e}_j - \mathbf{x}) \in \mathcal{S}_\mathbf{x}$, $\mathbf{x} + \sum_{j=1}^{n} a_j(\mathbf{e}_j - \mathbf{x}) = \left(1 - \sum_{j=1}^{n} a_j\right)\mathbf{x} + \sum_{j=1}^{n} a_j \mathbf{e}_j \in \operatorname{affspan}(E)$, hence $\operatorname{affspan}(E) = \mathbf{x} + \mathcal{S}_\mathbf{x}$. This set is a flat containing $E$; any flat containing $E$ is affine closed by Theorem 4.1.6, hence contains $\operatorname{affspan}(E)$. Therefore it is $\operatorname{affhull}(E)$. The dimension formula follows from Definition 4.1.1.
*Structural Commentary*: Affine span is ordinary span after temporarily choosing one point as the origin.
*Key Insight*: Differences from a base point record all directions in the affine hull. □

**Lemma 4.1.10.** *Intersections of Flats*
*Let $X_i = \mathbf{x}_i + \mathcal{S}_i$ be flats indexed by $i \in I$. If $\bigcap_{i \in I} X_i \neq \emptyset$ and $\mathbf{z} \in \bigcap_{i \in I} X_i$, then $\bigcap_{i \in I} X_i = \mathbf{z} + \bigcap_{i \in I} \mathcal{S}_i$.*

*Proof.* Since $\mathbf{z} \in X_i$, each flat has the form $X_i = \mathbf{z} + \mathcal{S}_i$. Hence $\mathbf{u} \in \bigcap_{i \in I} X_i \iff \mathbf{u} - \mathbf{z} \in \bigcap_{i \in I} \mathcal{S}_i$, QED.
*Structural Commentary*: A common point turns affine intersection into subspace intersection.
*Key Insight*: Translate to the common point; only the bases remain. □

**Theorem 4.1.11.** *Lattice of Flats*
*Let $\mathcal{A}_0(\mathcal{V}) := \mathcal{A}(\mathcal{V}) \cup \{\emptyset\}$, ordered by inclusion. Then $\mathcal{A}_0(\mathcal{V})$ is a complete lattice.*
*For every family $\mathcal{C} \subseteq \mathcal{A}_0(\mathcal{V})$, $\bigwedge \mathcal{C} = \bigcap_{X \in \mathcal{C}} X$, $\bigvee \mathcal{C} = \begin{cases} \operatorname{affhull}\left(\bigcup_{X \in \mathcal{C}} X\right), & \bigcup_{X \in \mathcal{C}} X \neq \emptyset, \\ \emptyset, & \bigcup_{X \in \mathcal{C}} X = \emptyset. \end{cases}$*

*Proof.*

(i) Arbitrary intersections of elements of $\mathcal{A}_0(\mathcal{V})$ lie in $\mathcal{A}_0(\mathcal{V})$: they are either empty, equal to $\mathcal{V}$ for the empty family, or flats by Lemma 4.1.10.

(ii) If $\bigcup_{X\in\mathcal{C}} X = \emptyset$, then $\emptyset$ is the least upper bound of $\mathcal{C}$.
Otherwise, affhull$(\bigcup_{X\in\mathcal{C}} X)$ is the least flat containing every $X \in \mathcal{C}$, so it is the join.

(iii) Since arbitrary meets and joins exist, $\mathcal{A}_0(\mathcal{V})$ is complete.

*Structural Commentary*: Adding the empty set repairs the failure of affine flats to be closed under intersection.
*Key Insight*: Meet is common incidence; join is the affine hull forced by union. □

**Remark 4.1.12.** *Empty Affine Hull Convention*
Within $\mathcal{A}_0(\mathcal{V})$, one may set affhull$(\emptyset) := \emptyset$ to write the join formula uniformly.

**Corollary 4.1.13.** *Join of Two Flats*
*Let $X = \mathbf{x} + \mathcal{S}$ and $Y = \mathbf{y} + \mathcal{T}$ be flats. Then $X \vee Y = \mathbf{x} + \operatorname{span}\big(\{\mathbf{x}-\mathbf{y}\} \cup \mathcal{S} \cup \mathcal{T}\big)$.*
*If $X \cap Y \neq \emptyset$, then $X \vee Y = \mathbf{x} + (\mathcal{S} + \mathcal{T})$.*

*Proof.* By Theorem 4.1.11, $X \vee Y = \text{affhull}(X \cup Y)$. Apply Theorem 4.1.9 to $X \cup Y$ with base point $\mathbf{x}$, then $(X \cup Y) - \mathbf{x} = \mathcal{S} \cup \big((\mathbf{y}-\mathbf{x}) + \mathcal{T}\big)$, whose span is $\operatorname{span}(\{\mathbf{x}-\mathbf{y}\} \cup \mathcal{S} \cup \mathcal{T})$.
If $X \cap Y \neq \emptyset$, then $\mathbf{x} - \mathbf{y} \in \mathcal{S} + \mathcal{T}$, so this span is $\mathcal{S} + \mathcal{T}$.
*Structural Commentary*: The join adds both bases and the displacement between representatives.
*Key Insight*: The displacement direction disappears exactly when the flats meet. □

**Corollary 4.1.14.** *Dimension of the Join of Two Flats*
*Let $X = \mathbf{x} + \mathcal{S}$ and $Y = \mathbf{y} + \mathcal{T}$ be flats.*

*(i) If $X \cap Y \neq \emptyset$, then $\dim(X \vee Y) = \dim(X) + \dim(Y) - \dim(X \cap Y)$.*
*(ii) If $X \cap Y = \emptyset$, then $\dim(X \vee Y) = \dim(\mathcal{S} + \mathcal{T}) + 1$.*

*Proof.*

(i) If $X \cap Y \neq \emptyset$, choose $\mathbf{z} \in X \cap Y$.
By Corollary 4.1.13 and Lemma 4.1.10, $X \vee Y = \mathbf{x} + (\mathcal{S} + \mathcal{T})$; $X \cap Y = \mathbf{z} + (\mathcal{S} \cap \mathcal{T})$. Thus the dimension formula for subspace sums gives $\dim(X \vee Y) = \dim(\mathcal{S} + \mathcal{T}) = \dim(\mathcal{S}) + \dim(\mathcal{T}) - \dim(\mathcal{S} \cap \mathcal{T})$.
Since $\dim X = \dim \mathcal{S}$, $\dim Y = \dim \mathcal{T}$, and $\dim(X \cap Y) = \dim(\mathcal{S} \cap \mathcal{T})$, this is (i).

(ii) If $X \cap Y = \emptyset$, Corollary 4.1.13 gives base $\mathcal{U} := \operatorname{span}(\{\mathbf{x}-\mathbf{y}\} \cup \mathcal{S} \cup \mathcal{T})$.
Since $\mathbf{x} - \mathbf{y} \notin \mathcal{S} + \mathcal{T}$, we have $\dim \mathcal{U} = \dim(\mathcal{S} + \mathcal{T}) + 1$, hence $\dim(X \vee Y) = \dim(\mathcal{S} + \mathcal{T}) + 1$.

*Structural Commentary*: Intersections subtract overlap; disjoint joins add one connecting direction.
*Key Insight*: The vector between representatives is new exactly when the flats are disjoint. □

## 4.2 Affine Independence and Barycentric Coordinates

**Definition 4.2.1.** *Affinely Independent Sets*
A nonempty set $E \subseteq \mathcal{V}$ is *affinely independent* if, for every $\mathbf{x} \in E$, the set $(E - \mathbf{x}) \setminus \{\mathbf{0}\}$ is linearly independent.

**Theorem 4.2.2.** *Affine Independence Criteria*
*For $\emptyset \neq E \subseteq \mathcal{V}$, conditions (i)–(iv) are equivalent. If $E$ is finite, they are also equivalent to (v).*

*(i) $E$ is affinely independent.*
*(ii) For every $\mathbf{x} \in E$, $\mathbf{x} \notin \text{affhull}(E \setminus \{\mathbf{x}\})$, with affhull$(\emptyset) = \emptyset$ in the singleton case.*
*(iii) $\forall$ distinct $\mathbf{x}_1, \ldots, \mathbf{x}_n \in E$ and $r_1, \ldots, r_n \in \mathbb{F}$, $\sum_{j=1}^{n} r_j \mathbf{x}_j = \mathbf{0}$ and $\sum_{j=1}^{n} r_j = 0$ imply $r_1 = \cdots = r_n = 0$.*
*(iv) $\forall$ distinct $\mathbf{x}_1, \ldots, \mathbf{x}_n \in E$ and affine coefficient vectors $\mathbf{r}, \mathbf{s} \in \mathbb{F}^n$,*
*$\sum_{j=1}^{n} r_j \mathbf{x}_j = \sum_{j=1}^{n} s_j \mathbf{x}_j$ and $\sum_{j=1}^{n} r_j = \sum_{j=1}^{n} s_j = 1$ imply $\mathbf{r} = \mathbf{s}$.*
*(v) If $E = \{\mathbf{x}_1, \ldots, \mathbf{x}_n\}$ is finite, then $\dim(\text{affhull}(E)) = n - 1$.*

*Proof.*

(i) $\Rightarrow$ (iii) Let $\sum_{j=1}^{n} r_j \mathbf{x}_j = \mathbf{0}$ with distinct $\mathbf{x}_j \in E$ and $\sum_j r_j = 0$. Then $\sum_{j=1}^{n-1} r_j(\mathbf{x}_j - \mathbf{x}_n) = \mathbf{0}$.
By (i), $r_1 = \cdots = r_{n-1} = 0$, and then $r_n = 0$.

(iii) $\Rightarrow$ (i) Fix $\mathbf{x} \in E$. Any relation $\sum_{j=1}^{m} a_j(\mathbf{y}_j - \mathbf{x}) = \mathbf{0}$, $\mathbf{y}_j \in E \setminus \{\mathbf{x}\}$ is the zero-sum relation
$\sum_{j=1}^{m} a_j \mathbf{y}_j - (\sum_j a_j)\mathbf{x} = \mathbf{0}$, so (iii) gives $a_1 = \cdots = a_m = 0$.

(iii) $\Rightarrow$ (ii) If $\mathbf{x}_k \in \text{affhull}(E \setminus \{\mathbf{x}_k\})$, then $\mathbf{x}_k = \sum_j r_j \mathbf{x}_j$ for $\mathbf{x}_j \neq \mathbf{x}_k$ and $\sum_j r_j = 1$,
giving a nontrivial zero-sum relation, contradiction.

(*ii*) $\Rightarrow$ (*iii*) If a zero-sum relation is nontrivial, choose $r_k \neq 0$.
Then $\mathbf{x}_k = \sum_{j\neq k} \left(-\frac{r_j}{r_k}\right)\mathbf{x}_j$, $\sum_{j\neq k} -\frac{r_j}{r_k} = 1$, so $\mathbf{x}_k \in \operatorname{affhull}(E \setminus \{\mathbf{x}_k\})$, contradicting (ii).
(*iii*) $\Rightarrow$ (*iv*) Subtract the two affine representations; the coefficient vector $\mathbf{r}-\mathbf{s}$ has sum 0, so (iii) gives $\mathbf{r}=\mathbf{s}$.
(*iv*) $\Rightarrow$ (*iii*) Let $\sum_j r_j\mathbf{x}_j = \mathbf{0}$ and $\sum_j r_j = 0$. Choose $\mathbf{a} \in \mathbb{F}^n$ with $\sum_j a_j = 1$.
Then $\mathbf{a}$ and $\mathbf{a}+\mathbf{r}$ are affine coefficient vectors for the same point. By (iv), $\mathbf{r}=\mathbf{0}$.
(*i*) $\Leftrightarrow$ (*v*) If $E = \{\mathbf{x}_1,\dots,\mathbf{x}_n\}$ and $\mathbf{x} \in E$, then Theorem 4.1.9 gives $\dim(\operatorname{affhull}(E)) = \dim(\operatorname{span}(E-\mathbf{x}))$.
Since $(E-\mathbf{x})\setminus\{\mathbf{0}\}$ has $n-1$ vectors, this dimension is $n-1$ iff those vectors are linearly independent.

*Structural Commentary*: Affine independence is linear independence after forgetting one chosen origin.
*Key Insight*: Uniqueness is absence of nontrivial zero-sum affine relations. □

**Definition 4.2.3.** *Affine Bases and Barycentric Coordinates* Let $A = \mathbf{x}+\mathcal{S}$ be an $n$-flat.

(i) A subset $\mathcal{B} \subseteq A$ is an *affine basis* of $A$ if $\mathcal{B}$ is affinely independent and $\operatorname{affspan}(\mathcal{B}) = A$; an *ordered affine basis* is an ordering of such a basis.
(ii) If $(\mathbf{s}_1,\dots,\mathbf{s}_n)$ is an ordered basis of $\mathcal{S}$, then $(\mathbf{x}+\mathbf{s}_1,\dots,\mathbf{x}+\mathbf{s}_n,\mathbf{x})$ is an ordered affine basis of $A$.
(iii) For an ordered affine basis $(\mathbf{p}_1,\dots,\mathbf{p}_{n+1})$, every $\mathbf{v} \in A$ has a unique expression $\mathbf{v} = \sum_{j=1}^{n+1} r_j\mathbf{p}_j$, $\sum_{j=1}^{n+1} r_j = 1$.
The scalars $(r_1,\dots,r_{n+1})$ are the *barycentric coordinates* of $\mathbf{v}$ in this basis.

**Example 4.2.4.** *Barycentric Coordinates on a Plane*
Let $A = \mathbf{x}+\operatorname{span}(\mathbf{v}_1,\mathbf{v}_2) \subseteq \mathbb{R}^3$, with $(\mathbf{v}_1,\mathbf{v}_2)$ linearly independent.
Set $\mathbf{p}_1 = \mathbf{x}+\mathbf{v}_1$, $\mathbf{p}_2 = \mathbf{x}+\mathbf{v}_2$, and $\mathbf{p}_3 = \mathbf{x}$. Then $(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3)$ is an ordered affine basis of $A$, and every $\mathbf{a} \in A$ has a unique barycentric expansion $\mathbf{a} = r_1\mathbf{p}_1 + r_2\mathbf{p}_2 + r_3\mathbf{p}_3$, $r_1+r_2+r_3 = 1$, with $r_1, r_2, r_3 \in \mathbb{R}$.

## 4.3 Affine Maps and Affine Groups

**Definition 4.3.1.** *Affine Transformation*
A map $f : \mathcal{V} \to \mathcal{V}$ is an *affine transformation* if it preserves finite affine combinations:
for all $\mathbf{x}_j \in \mathcal{V}$ and $r_j \in \mathbb{F}$ with $\sum_j r_j = 1$, $f\left(\sum_j r_j\mathbf{x}_j\right) = \sum_j r_j f(\mathbf{x}_j)$.

**Theorem 4.3.2.** *Two-Point Test for Affine Maps*
*Assume* $\operatorname{char}(\mathbb{F}) \neq 2$.
*A map $f : \mathcal{V} \to \mathcal{V}$ is affine if and only if $f(r\mathbf{x}+(1-r)\mathbf{y}) = rf(\mathbf{x})+(1-r)f(\mathbf{y})$ for all $\mathbf{x},\mathbf{y} \in \mathcal{V}$ and $r \in \mathbb{F}$.*

*Proof.*
($\Rightarrow$) Since $r+(1-r)=1$, affinity gives $f(r\mathbf{x}+(1-r)\mathbf{y}) = rf(\mathbf{x})+(1-r)f(\mathbf{y})$.
($\Leftarrow$) Assume displayed identity. Prove by induction on $n$ that $f\left(\sum_{j=1}^n r_j\mathbf{x}_j\right) = \sum_{j=1}^n r_j f(\mathbf{x}_j)$; $\sum_{j=1}^n r_j = 1$.
The cases $n = 1, 2$ are immediate. For $n \geq 3$, set $\mathbf{z} := \sum_{j=1}^n r_j\mathbf{x}_j$. If some $r_j \neq 1$, relabel so $r_1 \neq 1$ and put $\mathbf{u} := \sum_{j=2}^n \frac{r_j}{1-r_1}\mathbf{x}_j$. Then $f(\mathbf{z}) = f(r_1\mathbf{x}_1+(1-r_1)\mathbf{u}) = r_1f(\mathbf{x}_1)+(1-r_1)f(\mathbf{u}) = \sum_{j=1}^n r_j f(\mathbf{x}_j)$, by the two-point identity and induction. Otherwise all $r_j = 1$; since 2 is invertible, $\mathbf{z} = 2\left(\frac{1}{2}\mathbf{x}_1 + \frac{1}{2}\mathbf{x}_2\right) + r_3\mathbf{x}_3 + \cdots + r_n\mathbf{x}_n$, an $(n-1)$-term affine combination, so induction plus two-point identity gives same formula for $f(\mathbf{z})$.

*Structural Commentary*: In characteristic not 2, line preservation gives affine preservation.
*Key Insight*: The same two-point assembly that builds affine-closed sets also tests affine maps. □

**Definition 4.3.3.** *Translations*
For $\mathbf{a} \in \mathcal{V}$, the *translation by* $\mathbf{a}$ is the map $T_\mathbf{a} : \mathcal{V} \to \mathcal{V}$ defined by $T_\mathbf{a}(\mathbf{x}) = \mathbf{x}+\mathbf{a}$.

**Theorem 4.3.4.** *Affine Maps Are Linear Maps Followed by Translations*
*A map $f : \mathcal{V} \to \mathcal{V}$ is affine iff $f = T_\mathbf{a} \circ \tau$ for unique $\mathbf{a} \in \mathcal{V}$ and $\tau \in \mathcal{L}(\mathcal{V})$, namely $\mathbf{a} = f(\mathbf{0})$ and $\tau = T_{-\mathbf{a}} \circ f$.*

*Proof.*
($\Rightarrow$) Set $\mathbf{a} := f(\mathbf{0})$ and $\tau(\mathbf{x}) := f(\mathbf{x})-\mathbf{a}$. For $r, s \in \mathbb{F}$ and $\mathbf{x},\mathbf{y} \in \mathcal{V}$,
$\tau(r\mathbf{x}+s\mathbf{y}) = f(r\mathbf{x}+s\mathbf{y}+(1-r-s)\mathbf{0}) - \mathbf{a} = r\tau(\mathbf{x})+s\tau(\mathbf{y})$, so $\tau \in \mathcal{L}(\mathcal{V})$ and $f = T_\mathbf{a} \circ \tau$.
Uniqueness follows from $\mathbf{a} = f(\mathbf{0})$ and $\tau(\mathbf{x}) = f(\mathbf{x})-\mathbf{a}$.
($\Leftarrow$) If $f = T_\mathbf{a} \circ \tau$ with $\tau$ linear,
then for $\sum_j r_j = 1$, $f\left(\sum_j r_j\mathbf{x}_j\right) = \mathbf{a}+\tau\left(\sum_j r_j\mathbf{x}_j\right) = \sum_j r_j(\mathbf{a}+\tau\mathbf{x}_j) = \sum_j r_j f(\mathbf{x}_j)$, hence $f$ is affine.

*Structural Commentary*: An affine map is linear after moving its image of the origin back to the origin.
*Key Insight*: The failure of an affine map to be linear is exactly a translation. □

**Definition 4.3.5.** *Affine Group Data*

(i) The *general linear group* is $\mathrm{GL}(\mathcal{V}) := \{\tau \in \mathcal{L}(\mathcal{V}) : \tau \text{ is invertible}\}$.
(ii) The *affine group* is $\mathrm{Aff}(\mathcal{V}) := \{T_{\mathbf{a}} \circ \tau : \mathbf{a} \in \mathcal{V}, \tau \in \mathrm{GL}(\mathcal{V})\}$.
(iii) The *translation group* is $\mathrm{Trans}(\mathcal{V}) := \{T_{\mathbf{a}} : \mathbf{a} \in \mathcal{V}\}$.
(iv) The *linear-part map* is $\Lambda : \mathrm{Aff}(\mathcal{V}) \to \mathrm{GL}(\mathcal{V})$; $\Lambda(T_{\mathbf{a}} \circ \tau) = \tau$, by Theorem 4.3.4.

**Corollary 4.3.6.** *Bijective Affine Maps*
*For $\mathbf{a} \in \mathcal{V}$ and $\tau \in \mathcal{L}(\mathcal{V})$, $T_{\mathbf{a}} \circ \tau$ is bijective $\iff$ $\tau \in \mathrm{GL}(\mathcal{V})$.*

*Proof.*
($\Rightarrow$) If $T_{\mathbf{a}} \circ \tau$ is bijective, then $\tau = T_{-\mathbf{a}} \circ (T_{\mathbf{a}} \circ \tau)$ is bijective, hence $\tau \in \mathrm{GL}(\mathcal{V})$.
($\Leftarrow$) If $\tau \in \mathrm{GL}(\mathcal{V})$, then $T_{\mathbf{a}} \circ \tau$ is bijective with inverse $\tau^{-1} \circ T_{-\mathbf{a}}$.
*Structural Commentary*: Translation never affects invertibility; only the linear part does.
*Key Insight*: Bijective affine maps are exactly translations of linear isomorphisms. □

**Corollary 4.3.7.** *Affine Group and Translations*

*(i) For $\mathbf{a}, \mathbf{b} \in \mathcal{V}$ and $\tau, \sigma \in \mathcal{L}(\mathcal{V})$, $(T_{\mathbf{a}} \circ \tau) \circ (T_{\mathbf{b}} \circ \sigma) = T_{\mathbf{a}+\tau\mathbf{b}} \circ (\tau\sigma)$.*
*(ii) $\mathrm{Aff}(\mathcal{V})$ is a group, $\mathrm{Trans}(\mathcal{V}) \trianglelefteq \mathrm{Aff}(\mathcal{V})$, and $\Lambda$ is a surjective group homomorphism with $\ker\Lambda = \mathrm{Trans}(\mathcal{V})$.*
*(iii) Consequently, $\mathrm{Aff}(\mathcal{V})/\mathrm{Trans}(\mathcal{V}) \cong \mathrm{GL}(\mathcal{V})$.*

*Proof.*
i. For $\mathbf{x} \in \mathcal{V}$, $((T_{\mathbf{a}} \circ \tau) \circ (T_{\mathbf{b}} \circ \sigma))(\mathbf{x}) = \mathbf{a} + \tau\mathbf{b} + (\tau\sigma)\mathbf{x}$.
ii. For $\tau, \sigma \in \mathrm{GL}(\mathcal{V})$, (i) gives closure in $\mathrm{Aff}(\mathcal{V})$; the identity is $T_{\mathbf{0}} \circ I_{\mathcal{V}}$, and $(T_{\mathbf{a}} \circ \tau)^{-1} = T_{-\tau^{-1}\mathbf{a}} \circ \tau^{-1}$. Thus $\mathrm{Aff}(\mathcal{V})$ is a group. Also (i) gives $\Lambda(gh) = \Lambda(g)\Lambda(h)$, $\ker\Lambda = \{T_{\mathbf{a}} \circ I_{\mathcal{V}} : \mathbf{a} \in \mathcal{V}\} = \mathrm{Trans}(\mathcal{V})$, and $\Lambda$ is surjective since $\tau = T_{\mathbf{0}} \circ \tau$ for all $\tau \in \mathrm{GL}(\mathcal{V})$. Hence $\mathrm{Trans}(\mathcal{V}) \trianglelefteq \mathrm{Aff}(\mathcal{V})$.
iii. Since $\Lambda$ is surjective and $\ker\Lambda = \mathrm{Trans}(\mathcal{V})$, the first isomorphism theorem gives the induced isomorphism $\overline{\Lambda} : \mathrm{Aff}(\mathcal{V})/\mathrm{Trans}(\mathcal{V}) \to \mathrm{GL}(\mathcal{V})$; $\overline{\Lambda}(g\ \mathrm{Trans}(\mathcal{V})) = \Lambda(g)$.

*Structural Commentary*: The affine group is the translation part together with the invertible linear part.
*Key Insight*: Forgetting the translation records a homomorphism onto the linear automorphism group. □

## 4.4 Linear Projective Geometry and Affine Completion

**Definition 4.4.1.** *Projective Geometry of a Vector Space*

(i) Let $\mathrm{Sub}(\mathcal{V})$ be the set of subspaces of $\mathcal{V}$.
The *projective geometry* of $\mathcal{V}$ is $\mathrm{Sub}(\mathcal{V})$ with $\mathrm{pdim}(\mathcal{U}) := \dim(\mathcal{U}) - 1$.
(ii) A *projective point* is a 1-dimensional subspace; a *projective line* is a 2-dimensional subspace.
(iii) If $H = \mathbf{x} + \mathcal{K}$ is an affine hyperplane with $\mathbf{0} \notin H$ and $X \subseteq H$ is an affine flat, then $X$ is represented projectively by $\mathrm{span}(X)$, with $\mathrm{pdim}(\mathrm{span}(X)) = \dim(X)$.

**Perspective 4.4.2.** *Projective Completion of an Affine Hyperplane*
Let $H = \mathbf{x} + \mathcal{K}$ be a hyperplane with $\mathbf{0} \notin H$. The map $X \subseteq H \mapsto \mathrm{span}(X) \leq \mathcal{V}$ adjoins the missing directions in $\mathcal{K}$. Thus, when $\dim H = 2$, parallel affine lines in $H$ acquire a common projective point at infinity.




Figure 40: In the projective completion of $H = \mathbf{x} + \mathcal{K}$, an affine flat $X \subseteq H$ is represented by $\mathrm{span}(X)$; parallel flats meet in their common direction $D \leq \mathcal{K}$.

**Theorem 4.4.3.** *Linear Span Map from Affine to Projective Geometry*
*Let $H = \mathbf{x} + \mathcal{K}$ be an affine hyperplane with $\mathbf{0} \notin H$, and set $\mathcal{A}(H) := \{X \in \mathcal{A}(\mathcal{V}) : X \subseteq H\}$.*
*The map $P : \mathcal{A}(H) \to \mathrm{Sub}(\mathcal{V})$, $P(X) = \mathrm{span}(X)$ satisfies:*

*(i)* *$P$ is injective and $P^{-1}(U) = U \cap H$ for $U \in \mathrm{im}\, P$.*
*(ii)* *$\mathrm{im}\, P = \{\mathcal{U} \leq \mathcal{V} : \mathcal{U} \not\subseteq \mathcal{K}\}$.*
*(iii)* *$X \subseteq Y$ if and only if $P(X) \subseteq P(Y)$.*
*(iv)* *If $\bigcap_{i\in I} X_i \neq \emptyset$, then $P\left(\bigcap_{i\in I} X_i\right) = \bigcap_{i\in I} P(X_i)$.*
*(v)* *For any family $\{X_i : i \in I\} \subseteq \mathcal{A}(H)$, $P\left(\bigvee_{i\in I} X_i\right) = \bigvee_{i\in I} P(X_i)$, where $\bigvee_{i\in I} P(X_i)$ is computed in $\mathrm{Sub}(\mathcal{V})$, i.e. $\bigvee_{i\in I} P(X_i) = \sum_{i\in I} P(X_i)$.*
*(vi)* *$\mathrm{pdim}(P(X)) = \dim(X)$.*
*(vii)* *$X \parallel Y$ if and only if one of $P(X) \cap \mathcal{K}$ and $P(Y) \cap \mathcal{K}$ is contained in the other.*

*Proof.*

(i) Let $X = \mathbf{u} + \mathcal{S} \subseteq H$. Then $\mathcal{S} \leq \mathcal{K}$, $\mathbf{u} \notin \mathcal{K}$, and $P(X) = \mathbb{F}\mathbf{u} \oplus \mathcal{S}$.
If $\mathbf{z} \in P(X) \cap H$, then $\mathbf{z} = r\mathbf{u} + \mathbf{s} = \mathbf{u} + \mathbf{k}$ with $\mathbf{s} \in \mathcal{S}$ and $\mathbf{k} \in \mathcal{K}$. Hence $(1-r)\mathbf{u} \in \mathcal{K}$, so $r = 1$, and $P(X) \cap H = X$. Thus $P$ is injective and $P^{-1}(U) = U \cap H$ on $\mathrm{im}\, P$.
(ii) If $\mathcal{U} \not\subseteq \mathcal{K}$, choose $\mathbf{u} \in \mathcal{U} \setminus \mathcal{K}$. Since $\mathcal{V}/\mathcal{K}$ is 1-dimensional and $H = \mathbf{x} + \mathcal{K}$, there is $r \neq 0$ with $\mathbf{h} := r\mathbf{u} \in H$. Then $\mathcal{U} \cap H = \mathbf{h} + (\mathcal{U} \cap \mathcal{K})$; $P(\mathcal{U} \cap H) = \mathbb{F}\mathbf{h} + (\mathcal{U} \cap \mathcal{K}) = \mathcal{U}$. Thus $\mathrm{im}\, P = \{\mathcal{U} \leq \mathcal{V} : \mathcal{U} \not\subseteq \mathcal{K}\}$.
(iii) If $X \subseteq Y$, then $P(X) \subseteq P(Y)$. Conversely, if $P(X) \subseteq P(Y)$, then $X = P(X) \cap H \subseteq P(Y) \cap H = Y$.
(iv) If $\mathbf{z} \in \bigcap_i X_i$ and $X_i = \mathbf{z} + \mathcal{S}_i$, then Lemma 4.1.10 gives $\bigcap_i X_i = \mathbf{z} + \bigcap_i \mathcal{S}_i$.
Since $P(X_i) = \mathbb{F}\mathbf{z} \oplus \mathcal{S}_i$, $\bigcap_i P(X_i) = \mathbb{F}\mathbf{z} \oplus \bigcap_i \mathcal{S}_i = P\left(\bigcap_i X_i\right)$.
(v) By Theorem 4.1.11, $P\left(\bigvee_i X_i\right) = \mathrm{span}\left(\mathrm{affhull}\left(\bigcup_i X_i\right)\right) = \mathrm{span}\left(\bigcup_i X_i\right) = \bigvee_i P(X_i)$.
(vi) For $X = \mathbf{u} + \mathcal{S} \subseteq H$, $\mathrm{pdim}(P(X)) = \dim(\mathbb{F}\mathbf{u} \oplus \mathcal{S}) - 1 = \dim \mathcal{S} = \dim X$.
(vii) If $X = \mathbf{u} + \mathcal{S}$ and $Y = \mathbf{v} + \mathcal{T}$, then $P(X) \cap \mathcal{K} = \mathcal{S}$ and $P(Y) \cap \mathcal{K} = \mathcal{T}$.
Hence $X \parallel Y \iff \mathcal{S} \subseteq \mathcal{T}$ or $\mathcal{T} \subseteq \mathcal{S} \iff P(X) \cap \mathcal{K} \subseteq P(Y) \cap \mathcal{K}$ or $P(Y) \cap \mathcal{K} \subseteq P(X) \cap \mathcal{K}$.

*Structural Commentary*: $H$ selects affine representatives for projective subspaces outside $\mathcal{K}$.
*Key Insight*: Intersecting with $H$ undoes the linear span exactly because $H$ does not pass through the origin. □

## 4.5 Convex Combinations, Cones, and Hulls

**Definition 4.5.1.** *Convex Combinations, Cones, and Hulls*
Let $\mathbf{x}_1, \ldots, \mathbf{x}_k \in \mathbb{R}^n$.

i. A *convex combination* of these vectors is a vector $t_1\mathbf{x}_1 + \cdots + t_k\mathbf{x}_k$, $t_i \geq 0$, $\sum_{i=1}^{k} t_i = 1$.
ii. A set $C \subseteq \mathbb{R}^n$ is *convex* if the line segment between any two of its points lies in $C$.
iii. A set $K \subseteq \mathbb{R}^n$ is a *cone* if $\mathbf{x} \in K$ and $a \geq 0$ imply $a\mathbf{x} \in K$.
iv. *Convex hull* of finite set $S = \{\mathbf{x}_1, \ldots, \mathbf{x}_k\}$ is smallest convex subset of $\mathbb{R}^n$ containing $S$, denoted $\mathrm{conv}(S)$.

Closedness and compactness use the Euclidean metric conventions from Definitions 3.1.2 and 4.1.1.

**Proposition 4.5.2.** *Finite Convex Hulls*
*Let $S = \{\mathbf{x}_1, \ldots, \mathbf{x}_k\} \subseteq \mathbb{R}^n$ and $\Delta_{k-1} := \left\{(t_1, \ldots, t_k) \in \mathbb{R}^k : t_i \geq 0,\ \sum_{i=1}^{k} t_i = 1\right\}$.*
*Then $\mathrm{conv}(S) = \left\{\sum_{i=1}^{k} t_i\mathbf{x}_i : (t_1, \ldots, t_k) \in \Delta_{k-1}\right\}$. Moreover, $\mathrm{conv}(S)$ is compact.*

*Proof.* Let $T : \Delta_{k-1} \to \mathbb{R}^n$ be $T(t_1, \ldots, t_k) = \sum_{i=1}^{k} t_i\mathbf{x}_i$, and set $C := T(\Delta_{k-1})$. Taking $t_i = 1$ and $t_j = 0$ for $j \neq i$ gives $S \subseteq C$. If $K \subseteq \mathbb{R}^n$ is convex and $S \subseteq K$, then $C \subseteq K$, hence $C \subseteq \mathrm{conv}(S)$.
Conversely, $\Delta_{k-1}$ is convex and $T$ is affine, so $C$ is convex; since $S \subseteq C$, $\mathrm{conv}(S) \subseteq C$. Thus $\mathrm{conv}(S) = C$.
The simplex $\Delta_{k-1}$ is closed and bounded, hence compact by Heine–Borel, Theorem 4.2.5.
Since $T$ is continuous, $\mathrm{conv}(S) = T(\Delta_{k-1})$ is compact by Theorem 5.1.35.
*Structural Commentary*: A finite convex hull is the continuous image of a compact coefficient simplex.
*Key Insight*: Convexity is carried by the coefficients $t_i$, not by the ambient coordinates. □

**Proposition 4.5.3.** *Sums of Convex and Closed Sets*
*For $X, Y \subseteq \mathbb{R}^n$, set $X + Y := \{\mathbf{x} + \mathbf{y} : \mathbf{x} \in X,\ \mathbf{y} \in Y\}$. Then:*

*(i)* *$X, Y$ convex $\implies$ $X + Y$ convex.*

*(ii) $X$ compact and $Y$ closed $\implies$ $X+Y$ closed.*

*Proof.*

(i) Assume $X, Y$ are convex. Let $\mathbf{z}_r = \mathbf{x}_r + \mathbf{y}_r \in X+Y$ for $r = 0, 1$, and let $0 \le t \le 1$. Then $t\mathbf{z}_0 + (1-t)\mathbf{z}_1 = \big(t\mathbf{x}_0 + (1-t)\mathbf{x}_1\big) + \big(t\mathbf{y}_0 + (1-t)\mathbf{y}_1\big) \in X+Y$. Thus $X+Y$ is convex.

(ii) Assume $X$ is compact and $Y$ is closed. Let $\mathbf{z}_j \in X+Y$ and $\mathbf{z}_j \to \mathbf{z}$. Let $\mathbf{z}_j = \mathbf{x}_j + \mathbf{y}_j$ with $\mathbf{x}_j \in X$ and $\mathbf{y}_j \in Y$. Since $X$ is compact, a subsequence satisfies $\mathbf{x}_{j_\ell} \to \mathbf{x} \in X$. Then $\mathbf{y}_{j_\ell} = \mathbf{z}_{j_\ell} - \mathbf{x}_{j_\ell} \to \mathbf{z} - \mathbf{x} \in Y$, as $Y$ is closed. Hence $\mathbf{z} = \mathbf{x} + (\mathbf{z}-\mathbf{x}) \in X+Y$, so $X+Y$ is closed.

*Structural Commentary*: Compactness lets one hold the $X$-part still enough for closedness of $Y$ to apply.
*Key Insight*: A limit in a sum can be split by passing to a convergent compact subsequence. □

## 4.6 Hyperplanes, Nearest Points, and Separation

**Definition 4.6.1.** *Hyperplanes, Half-Spaces, and Separation*
Let $\mathbf{N} \in \mathbb{R}^n \setminus \{\mathbf{0}\}$ and $b \in \mathbb{R}$.

(i) The *affine hyperplane* with normal $\mathbf{N}$ and level $b$ is $\mathcal{H}(\mathbf{N}, b) := \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{N}, \mathbf{x}\rangle = b\}$.
(ii) The case $b = 0$ gives the *linear hyperplane* $\mathcal{H}(\mathbf{N}, 0) = \operatorname{span}(\mathbf{N})^\perp$.
(iii) The *closed half-spaces* determined by $\mathcal{H}(\mathbf{N}, b)$ are $\mathcal{H}_+(\mathbf{N}, b) := \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{N}, \mathbf{x}\rangle \ge b\}$ and $\mathcal{H}_-(\mathbf{N}, b) := \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{N}, \mathbf{x}\rangle \le b\}$. The *open half-spaces* are obtained by replacing $\ge, \le$ with $>, <$.
(iv) For $X \subseteq \mathbb{R}^n$ and $\star \in \{<, \le, >, \ge\}$, use the *set-inequality convention* $\langle \mathbf{N}, X\rangle \star b \iff \langle \mathbf{N}, \mathbf{x}\rangle \star b \ \forall \mathbf{x} \in X$.
(v) An *orientation* of $\mathcal{H}(\mathbf{N}, b)$ is $\sigma(\mathbf{N}, b) := (\sigma\mathbf{N}, \sigma b)$ for $\sigma \in \{\pm 1\}$.
(vi) Sets $X, Y \subseteq \mathbb{R}^n$ are *strictly separated* by $\mathcal{H}(\mathbf{N}, b)$ if $\exists \sigma \in \{\pm 1\}$ such that $\langle \sigma\mathbf{N}, X\rangle < \sigma b < \langle \sigma\mathbf{N}, Y\rangle$
(vii) Sets $X, Y \subseteq \mathbb{R}^n$ are *strongly separated* by $\mathcal{H}(\mathbf{N}, b)$ if $\exists \sigma \in \{\pm 1\}$ and $\varepsilon > 0$ such that $\langle \sigma\mathbf{N}, X\rangle < \sigma b - \varepsilon < \sigma b + \varepsilon < \langle \sigma\mathbf{N}, Y\rangle$.

**Remark 4.6.2.** *Hyperplane Geometry*
The half-spaces satisfy $\mathcal{H}_+(\mathbf{N}, b) \cap \mathcal{H}_-(\mathbf{N}, b) = \mathcal{H}(\mathbf{N}, b)$, and two open half-spaces with $\mathcal{H}(\mathbf{N}, b)$ partition $\mathbb{R}^n$. Moreover, $\mathbf{N} \in \mathcal{H}(\mathbf{N}, \|\mathbf{N}\|^2)$, by Cauchy–Schwarz (Theorem 1.1.13), $\mathbf{N}$ is closest point of this hyperplane to $\mathbf{0}$.

**Theorem 4.6.3.** *Nearest Points and Separating a Point*
*Let $C \subseteq \mathbb{R}^n$ be nonempty, closed, and convex.*

*(i) There exists a unique $\mathbf{N}_C \in C$ such that $\|\mathbf{N}_C\| = \min_{\mathbf{x} \in C} \|\mathbf{x}\|$.*
*(ii) If $\mathbf{b} \notin C$, set $D := C - \mathbf{b}$ and let $\mathbf{N} \in D$ satisfy $\|\mathbf{N}\| = \min_{\mathbf{z} \in D} \|\mathbf{z}\|$. Then $\mathbf{N} \neq \mathbf{0}$ and $\langle \mathbf{N}, C\rangle \ge \|\mathbf{N}\|^2 + \langle \mathbf{N}, \mathbf{b}\rangle > \langle \mathbf{N}, \mathbf{b}\rangle$. With $\alpha := \langle \mathbf{N}, \mathbf{b}\rangle + \frac{1}{2}\|\mathbf{N}\|^2$, the hyperplane $\mathcal{H}(\mathbf{N}, \alpha)$ strongly separates $\{\mathbf{b}\}$ and $C$.*

*Proof.*

(i) Choose $\mathbf{x}_0 \in C$ and $s \ge \|\mathbf{x}_0\|$. Then $K := C \cap \overline{B}_s(\mathbf{0})$ is nonempty and compact by Theorem 4.2.5. Since $\|\cdot\|$ is continuous, Theorem 5.1.37 gives $\mathbf{N}_C \in K$ minimizing $\|\cdot\|$ on $K$. If $\mathbf{x} \in C$ and $\|\mathbf{x}\| < \|\mathbf{N}_C\|$, then $\mathbf{x} \in K$, a contradiction; hence $\mathbf{N}_C$ minimizes on $C$. If $\mathbf{u} \neq \mathbf{v}$ are minimizers of norm $m$, then $\mathbf{w} := (\mathbf{u}+\mathbf{v})/2 \in C$ and, by the parallelogram identity, Theorem 1.1.20, $\|\mathbf{w}\|^2 = m^2 - \frac{1}{4}\|\mathbf{u}-\mathbf{v}\|^2 < m^2$, a contradiction.

(ii) The set $D = C - \mathbf{b}$ is nonempty, closed, and convex, so (i) gives its unique minimum-norm vector $\mathbf{N}$. Since $\mathbf{b} \notin C$, $\mathbf{0} \notin D$, hence $\mathbf{N} \neq \mathbf{0}$. For $\mathbf{z} \in D$, define $\phi(t) := \|\mathbf{N} + t(\mathbf{z} - \mathbf{N})\|^2$, $0 \le t \le 1$. Minimality gives $\phi'_+(0) \ge 0$, hence $\langle \mathbf{N}, \mathbf{z}\rangle \ge \|\mathbf{N}\|^2$. Taking $\mathbf{z} = \mathbf{x} - \mathbf{b}$ yields $\langle \mathbf{N}, \mathbf{x}\rangle \ge \|\mathbf{N}\|^2 + \langle \mathbf{N}, \mathbf{b}\rangle$ for all $\mathbf{x} \in C$. For $\alpha := \langle \mathbf{N}, \mathbf{b}\rangle + \|\mathbf{N}\|^2/2$ and $\varepsilon := \|\mathbf{N}\|^2/4$, then $\langle \mathbf{N}, \mathbf{b}\rangle < \alpha - \varepsilon < \alpha + \varepsilon < \langle \mathbf{N}, C\rangle$, so $\mathcal{H}(\mathbf{N}, \alpha)$ strongly separates $\{\mathbf{b}\}$ and $C$.

*Structural Commentary*: The nearest point in $C - \mathbf{b}$ supplies the normal vector to a separating hyperplane.
*Key Insight*: Minimal norm turns a metric problem into the linear inequality $\langle \mathbf{N}, \mathbf{z}\rangle \ge \|\mathbf{N}\|^2$. □




Figure 41: A nearest vector in $C - \mathbf{b}$ gives a normal direction and a strongly separating hyperplane.

**Theorem 4.6.4.** *Separating a Compact Convex Set from a Subspace*
*Let $C \subseteq \mathbb{R}^n$ be nonempty, compact, and convex, and let $S \leq \mathbb{R}^n$ satisfy $C \cap S = \emptyset$.*
*Then there exists $\mathbf{N} \in S^\perp \setminus \{\mathbf{0}\}$ such that $\langle \mathbf{N}, C\rangle \geq \|\mathbf{N}\|^2 > 0$.*
*Consequently, $\mathcal{H}(\mathbf{N}, \|\mathbf{N}\|^2/2)$ strongly separates $S$ and $C$.*

*Proof.*

(i) Set $D := S + C$. Since $S$ is closed in finite dimension, Proposition 4.5.3 gives $D$ closed and convex. Also $\mathbf{0} \notin D$, for $\mathbf{0} = \mathbf{s} + \mathbf{c}$ would imply $\mathbf{c} = -\mathbf{s} \in C \cap S$.

(ii) Apply Theorem 4.6.3 to $D$ and $\mathbf{0}$. Then $\exists \mathbf{N} \neq \mathbf{0}$ such that $\langle \mathbf{N}, \mathbf{s} + \mathbf{c}\rangle \geq \|\mathbf{N}\|^2$, $\mathbf{s} \in S$, $\mathbf{c} \in C$. Fix $\mathbf{c}_0 \in C$. If $a := \langle \mathbf{N}, \mathbf{s}\rangle \neq 0$ for some $\mathbf{s} \in S$, then $\langle \mathbf{N}, \lambda\mathbf{s} + \mathbf{c}_0\rangle = \lambda a + \langle \mathbf{N}, \mathbf{c}_0\rangle \to -\infty$ for $\lambda$ of opposite sign and $|\lambda| \to \infty$, a contradiction. Hence $\mathbf{N} \in S^\perp$. Taking $\mathbf{s} = \mathbf{0}$ gives $\langle \mathbf{N}, C\rangle \geq \|\mathbf{N}\|^2 > 0$. Since $\langle \mathbf{N}, \mathbf{s}\rangle = 0$ for all $\mathbf{s} \in S$, the choice $\varepsilon := \|\mathbf{N}\|^2/4$ gives strong separation by $\mathcal{H}(\mathbf{N}, \|\mathbf{N}\|^2/2)$.

*Structural Commentary*: The sum $S + C$ is a closed convex set disjoint from $\mathbf{0}$, so point separation applies.
*Key Insight*: Since the separating inequality holds for all translates by $S$, scaling inside $S$ forces $\mathbf{N}$ to annihilate $S$. □

## 4.7 Positive Orthants, Alternatives, and Farkas's Lemma

This subsection uses finite-dimensional real inner-product geometry to decide when linear systems have positive solutions. The guiding theme is that a missing positive solution is often detected by a separating hyperplane, hence by a dual vector certificate.

**Definition 4.7.1.** *Positive Orthants*
For $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{R}^n$, use the following order and orthant notation:

(i) $\mathbf{x} \geq \mathbf{0}$ iff $\mathbf{x}$ is *coordinatewise nonnegative*: $x_i \geq 0$ for every $i$.
(ii) $\mathbf{x} > \mathbf{0}$ iff $\mathbf{x}$ is *nonzero nonnegative*: $\mathbf{x} \geq \mathbf{0}$ and $\mathbf{x} \neq \mathbf{0}$.
(iii) $\mathbf{x} \gg \mathbf{0}$ iff $\mathbf{x}$ is *coordinatewise positive*: $x_i > 0$ for every $i$.
(iv) $\mathbb{R}^n_+ := \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x} > \mathbf{0}\}$ is the *nonzero nonnegative orthant*.
(v) $\mathbb{R}^n_{++} := \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x} \gg \mathbf{0}\}$ is the *coordinatewise positive interior*.

**Motivation 4.7.2.** *Positive Null Vectors as Probabilities*
Let $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathrm{Mat}(m, n, \mathbb{R})$ and let $\mathbf{x} = (x_1, \ldots, x_n) > \mathbf{0}$ satisfy $A\mathbf{x} = \mathbf{0}$.

(i) Set $s := x_1 + \cdots + x_n > 0$ and $\boldsymbol{\pi} := s^{-1}\mathbf{x}$.
(ii) Then $\pi_i \geq 0$ and $\sum_{i=1}^n \pi_i = 1$, so $\boldsymbol{\pi}$ is a probability vector.
(iii) Since $A\boldsymbol{\pi} = s^{-1}A\mathbf{x} = \mathbf{0}$, equivalently $\sum_{i=1}^n \pi_i \mathbf{a}_i = \mathbf{0}$.
(iv) If $\mathbf{x} \gg \mathbf{0}$, then $\pi_i > 0$ for every $i$.

Thus a positive null vector normalizes to probabilities under which the columns of $A$ have zero expected value.

**Principle 4.7.3.** *Positive Solutions Are Orthant Intersections*
For $A \in \mathrm{Mat}(m, n, \mathbb{R})$, $\ker(A) = \mathrm{row}(A)^\perp$ by Corollary 2.4.13.
Hence $\{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \mathbf{0},\ \mathbf{x} > \mathbf{0}\} = \mathrm{row}(A)^\perp \cap \mathbb{R}^n_+$; and $\{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \mathbf{0},\ \mathbf{x} \gg \mathbf{0}\} = \mathrm{row}(A)^\perp \cap \mathbb{R}^n_{++}$.
Thus *positive null vectors* are precisely orthant intersections with $\mathrm{row}(A)^\perp$.




Figure 42: In $\mathbb{R}^2$, a negative-slope line $S$ has $S^\perp$ meeting the strongly positive orthant.

**Theorem 4.7.4.** *Positive Orthant Alternatives for Subspaces*
*Let $S \leq \mathbb{R}^n$. Then:*

*(i)* $S \cap \mathbb{R}^n_+ = \emptyset \iff S^\perp \cap \mathbb{R}^n_{++} \neq \emptyset$.
*(ii)* $S \cap \mathbb{R}^n_{++} = \emptyset \iff S^\perp \cap \mathbb{R}^n_+ \neq \emptyset$.

*Proof.*

*(i)* ($\Rightarrow$) Let $\Delta := \operatorname{conv}\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$. By Proposition 4.5.2, $\Delta$ is compact and convex, and $\Delta \subseteq \mathbb{R}^n_+$.
Hence $\Delta \cap S = \emptyset$. Theorem 4.6.4 gives $\mathbf{N} \in S^\perp \setminus \{\mathbf{0}\}$ with $\langle \mathbf{N}, \mathbf{e}_i \rangle \geq \|\mathbf{N}\|^2 > 0$, $1 \leq i \leq n$. QED.
*(i)* ($\Leftarrow$) If $\mathbf{x} \in S \cap \mathbb{R}^n_+$ and $\mathbf{N} \in S^\perp \cap \mathbb{R}^n_{++}$, then $\langle \mathbf{N}, \mathbf{x} \rangle > 0$, contradicting $\mathbf{N} \perp S$.
*(ii)* ($\Rightarrow$) If $S = \{\mathbf{0}\}$, then $\mathbf{e}_1 \in S^\perp \cap \mathbb{R}^n_+$. Assume $S \neq \{\mathbf{0}\}$.
Let $\mathbf{B}_1, \dots, \mathbf{B}_k$ be a basis of $S$ and set $M = [\mathbf{B}_1 \ \cdots \ \mathbf{B}_k] \in \operatorname{Mat}(n, k, \mathbb{R})$, $\operatorname{rows}(M) = \{\mathbf{R}_1, \dots, \mathbf{R}_n\} \subseteq \mathbb{R}^k$.
For $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{a} \in S^\perp \iff \sum_{i=1}^n a_i \mathbf{R}_i = \mathbf{0}$.
Set $K := \operatorname{conv}\{\mathbf{R}_1, \dots, \mathbf{R}_n\}$. Hence $S^\perp \cap \mathbb{R}^n_+ \neq \emptyset \iff \mathbf{0} \in K$. If $\mathbf{0} \notin K$, Theorem 4.6.3 gives $\mathbf{u} \in \mathbb{R}^k$ with $\langle \mathbf{u}, \mathbf{R}_i \rangle > 0$ for all $i$. Then $\mathbf{v} := M\mathbf{u} \in S$ has coordinates $v_i = \langle \mathbf{u}, \mathbf{R}_i \rangle > 0$, so $\mathbf{v} \in S \cap \mathbb{R}^n_{++}$, a contradiction. Thus $\mathbf{0} \in K$, and $S^\perp \cap \mathbb{R}^n_+ \neq \emptyset$.

*Structural Commentary*: Use the simplex and row hull as compact witnesses.
*Key Insight*: Missing positivity yields an orthogonal positive certificate. □

**Theorem 4.7.5.** *Homogeneous Positive Alternatives*
*Let $A \in \operatorname{Mat}(m, n, \mathbb{R})$. Exactly one alternative holds in each pair:*

*(i)* $\ker(A) \cap \mathbb{R}^n_{++} \neq \emptyset$ *or* $\operatorname{col}(A^t) \cap \mathbb{R}^n_+ \neq \emptyset$.
*(ii)* $\ker(A) \cap \mathbb{R}^n_+ \neq \emptyset$ *or* $\operatorname{col}(A^t) \cap \mathbb{R}^n_{++} \neq \emptyset$.

*Proof.*

(i) Let $S := \operatorname{row}(A) = \operatorname{col}(A^t) \subseteq \mathbb{R}^n$. By Principle 4.7.3, $\ker(A) = S^\perp$.
Moreover, $A^t\mathbf{v} \in \operatorname{col}(A^t) = S$ for every $\mathbf{v} \in \mathbb{R}^m$, and every vector of $S$ is of this form.
Hence the first pair is equivalent to $S^\perp \cap \mathbb{R}^n_{++} \neq \emptyset$ or $S \cap \mathbb{R}^n_+ \neq \emptyset$.
Theorem 4.7.4(i) says exactly that these two alternatives are mutually exclusive and exhaustive.
(ii) With the same $S$, one has $\ker(A) \cap \mathbb{R}^n_+ = S^\perp \cap \mathbb{R}^n_+$, and $\{A^t\mathbf{v} : \mathbf{v} \in \mathbb{R}^m\} \cap \mathbb{R}^n_{++} = S \cap \mathbb{R}^n_{++}$.
Thus the second pair is equivalent to $S^\perp \cap \mathbb{R}^n_+ \neq \emptyset$ or $S \cap \mathbb{R}^n_{++} \neq \emptyset$.
Theorem 4.7.4(ii) says exactly that these two alternatives are mutually exclusive and exhaustive.

*Structural Commentary*: The row space gives the certificate side; the nullspace gives the solution side.
*Key Insight*: Positive solvability is an orthogonal-complement alternative. □

**Method 4.7.6.** *Certifying Positive Homogeneous Systems*
Let $A \in \operatorname{Mat}(m, n, \mathbb{R})$ and $S := \operatorname{row}(A) = \operatorname{col}(A^t)$.

(i) $\ker(A) \cap \mathbb{R}^n_{++} = \emptyset$ is certified by $\exists\, \mathbf{v} \in \mathbb{R}^m$ with $A^t\mathbf{v} > \mathbf{0}$.
(ii) $\ker(A) \cap \mathbb{R}^n_+ = \emptyset$ is certified by $\exists\, \mathbf{v} \in \mathbb{R}^m$ with $A^t\mathbf{v} \gg \mathbf{0}$.
(iii) Existence follows from row-space avoidance,
$S \cap \mathbb{R}^n_+ = \emptyset \Rightarrow \ker(A) \cap \mathbb{R}^n_{++} \neq \emptyset$; or $S \cap \mathbb{R}^n_{++} = \emptyset \Rightarrow \ker(A) \cap \mathbb{R}^n_+ \neq \emptyset$.

**Lemma 4.7.7.** *Nonnegative Matrix Cone*
*Let* $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \operatorname{Mat}(m, n, \mathbb{R})$ *and* $C_A := A(\mathbb{R}^n_{\geq 0}) = \left\{ \sum_{i=1}^n t_i \mathbf{a}_i : t_i \geq 0 \right\}$.
*Then $C_A$ is a closed convex cone in $\mathbb{R}^m$.*

*Proof.* For $\mathbf{x}_r = A\mathbf{y}_r \in C_A$ with $\mathbf{y}_r \geq \mathbf{0}$.
For $0 \leq t \leq 1$, $a \geq 0$, $t\mathbf{x}_1 + (1-t)\mathbf{x}_2 = A(t\mathbf{y}_1 + (1-t)\mathbf{y}_2) \in C_A$, where $a\mathbf{x}_1 = A(a\mathbf{y}_1) \in C_A$, $C_A$ is convex cone.
For closedness, choose for $\mathbf{x} \in C_A$ a representation $\mathbf{x} = \sum_{i\in I} t_i\mathbf{a}_i, \qquad t_i > 0$ with $I$ minimal.
If $\{\mathbf{a}_i : i \in I\}$ were dependent, choose $(d_i)_{i\in I} \neq 0$ with $\sum_{i\in I} d_i\mathbf{a}_i = \mathbf{0}$.
Replacing $d$ by $-d$ if necessary, some $d_i < 0$; then $\lambda := \min_{d_i<0} \frac{t_i}{-d_i}$ makes $t_i + \lambda d_i \geq 0$ for all $i$ and zero for at least one $i$, contradicting minimality. Hence the active columns are independent.
Thus $C_A = \bigcup_{\substack{I\subseteq\{1,\dots,n\} \\ \{\mathbf{a}_i : i\in I\} \text{ independent}}} K_I$, where $K_I := \left\{\sum_{i\in I} t_i\mathbf{a}_i : t_i \geq 0\right\}$.
For such $I$, the map $T_I : \mathbb{R}^I \to \operatorname{span}\{\mathbf{a}_i : i \in I\}$, where $T_I((t_i)) = \sum_{i\in I} t_i\mathbf{a}_i$, is a linear isomorphism, so $K_I = T_I(\mathbb{R}^I_{\geq 0})$ is closed in its finite-dimensional range, and the range is closed in $\mathbb{R}^m$.
Hence each $K_I$ is closed in $\mathbb{R}^m$, and the finite union above is closed.
*Structural Commentary*: Reduce $C_A$ to finitely many independent-column cones.
*Key Insight*: Minimal support turns generators independent, making each cone closed. □

**Theorem 4.7.8.** *Farkas's Lemma*
*Let $A \in \operatorname{Mat}(m, n, \mathbb{R})$, $\mathbf{b} \in \mathbb{R}^m \setminus \{\mathbf{0}\}$, and $C_A := \{A\mathbf{y} : \mathbf{y} \in \mathbb{R}^n,\ \mathbf{y} \geq \mathbf{0}\}$.*
*Exactly one of the following holds:*

*(i) $\mathbf{b} \in C_A$, equivalently $\exists\, \mathbf{u} > \mathbf{0}$ with $A\mathbf{u} = \mathbf{b}$.*
*(ii) $\exists\, \mathbf{v} \in \mathbb{R}^m$ such that $A^t\mathbf{v} \leq \mathbf{0}$ and $\langle \mathbf{v}, \mathbf{b}\rangle > 0$.*

*Proof.*

(i) If $\mathbf{b} = A\mathbf{u}$ with $\mathbf{u} \geq \mathbf{0}$ and $A^t\mathbf{v} \leq \mathbf{0}$, then $\langle \mathbf{v}, \mathbf{b}\rangle = \langle \mathbf{v}, A\mathbf{u}\rangle = \langle A^t\mathbf{v}, \mathbf{u}\rangle \leq 0$.
(ii) Suppose $\mathbf{b} \notin C_A$. By Lemma 4.7.7, $C_A$ is closed and convex. Since $\mathbf{0} \in C_A$, Theorem 4.6.3, after changing sign if necessary, gives $\mathbf{v} \in \mathbb{R}^m$ and $\alpha \in \mathbb{R}$ such that $\langle \mathbf{v}, C_A\rangle < \alpha < \langle \mathbf{v}, \mathbf{b}\rangle$.
Thus $\alpha > 0$ and $\langle \mathbf{v}, \mathbf{b}\rangle > 0$. For every $\mathbf{y} \geq \mathbf{0}$, $\langle A^t\mathbf{v}, \mathbf{y}\rangle = \langle \mathbf{v}, A\mathbf{y}\rangle < \alpha$.
If $(A^t\mathbf{v})_i > 0$, taking $\mathbf{y} = \lambda\mathbf{e}_i$ and $\lambda \to \infty$ contradicts the bounded inequality. Hence $A^t\mathbf{v} \leq \mathbf{0}$.

*Structural Commentary*: Nonrepresentability means $\mathbf{b} \notin C_A$.
*Key Insight*: Separation turns nonmembership in a cone into the dual inequality $A^t\mathbf{v} \leq 0$. □

**Example 4.7.9.** *Why the Weak Inequality Matters*
The conclusion $A^t\mathbf{v} \leq \mathbf{0}$ cannot generally be strengthened to $A^t\mathbf{v} \ll \mathbf{0}$. Let $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$.
If $\mathbf{u} \geq \mathbf{0}$, then $(A\mathbf{u})_1 = u_1 \geq 0$, so $A\mathbf{u} \neq \mathbf{b}$.
The certificate $\mathbf{v} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$ satisfies $A^t\mathbf{v} = (-1, 0, 0)^t \leq \mathbf{0}$ and $\langle \mathbf{v}, \mathbf{b}\rangle = 1 > 0$.
For arbitrary $\mathbf{w} = (r, s)^t$, $A^t\mathbf{w} = (r, s, -s)^t$, so $A^t\mathbf{w} \ll \mathbf{0}$ would force $s < 0 < -s$, impossible.
*Structural Commentary*: Opposite columns force a zero in every certificate.
*Key Insight*: Boundary separation is sometimes the best possible separation.

# 5 Eigenstructure and Invariant Subspaces

## 5.1 Eigenvalues, Eigenspaces, and Spectral Criteria

Eigenstructure begins by identifying the scalars for which a shifted operator has a nontrivial kernel.

**Definition 5.1.1.** *Eigenvalue and Eigenvector*
Let $T \in \mathcal{L}(\mathcal{V})$. A scalar $\lambda \in \mathbb{F}$ is an *eigenvalue* of $T$ if there exists $\mathbf{v} \in \mathcal{V}$ with $\mathbf{v} \neq \mathbf{0}$ and $T\mathbf{v} = \lambda\mathbf{v}$. Such a vector $\mathbf{v}$ is an *eigenvector* of $T$ corresponding to $\lambda$.

**Definition 5.1.2.** *Eigenspace*
Let $T \in \mathcal{L}(\mathcal{V})$ and $\lambda \in \mathbb{F}$. The *eigenspace* of $T$ corresponding to $\lambda$ is $E_\lambda(T) := \ker(T - \lambda I_\mathcal{V})$. Thus $\lambda$ is an eigenvalue of $T$ if and only if $E_\lambda(T) \neq \{\mathbf{0}\}$. Equivalently, $E_\lambda(T)$ is the nullspace of the shifted operator $T - \lambda I_\mathcal{V}$.

**Definition 5.1.3.** *Spectrum and Geometric Multiplicity*
Let $T \in \mathcal{L}(\mathcal{V})$. The *spectrum* of $T$ is $\operatorname{spec}(T) := \{\lambda \in \mathbb{F} : \lambda \text{ is an eigenvalue of } T\}$.
The *geometric multiplicity* of $\lambda \in \mathbb{F}$ is $\dim E_\lambda(T)$. For a square matrix $A \in \operatorname{Mat}(n, n, \mathbb{F})$, $E_\lambda(A) := \ker(A - \lambda I_n)$ and $\operatorname{spec}(A)$ denotes the spectrum of the induced operator on $\mathbb{F}^n$.

**Theorem 5.1.4.** *Equivalent Conditions for an Eigenvalue*
*Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $\lambda \in \mathbb{F}$. The following are equivalent:*

*(i)* $\lambda$ *is an eigenvalue of* $T$*;*
*(ii)* $T - \lambda I_{\mathcal{V}}$ *is not injective;*
*(iii)* $T - \lambda I_{\mathcal{V}}$ *is not surjective;*
*(iv)* $T - \lambda I_{\mathcal{V}}$ *is not invertible.*

*Proof.*

$(i) \Rightarrow (ii)$ If $\lambda$ is an eigenvalue, Definition 5.1.2 gives $\ker(T - \lambda I_{\mathcal{V}}) \neq \{\mathbf{0}\}$, so $T - \lambda I_{\mathcal{V}}$ is not injective.
$(ii) \Rightarrow (i)$ If $T - \lambda I_{\mathcal{V}}$ is not injective, then $\ker(T - \lambda I_{\mathcal{V}}) \neq \{\mathbf{0}\}$, so $\lambda$ is an eigenvalue by Definition 5.1.2.
$(ii) \Rightarrow (iii)$ If $T - \lambda I_{\mathcal{V}}$ were surjective, Theorem 1.7.8 would make it injective, a contradiction.
$(iii) \Rightarrow (iv)$ If $T - \lambda I_{\mathcal{V}}$ were invertible, then it would be surjective, a contradiction.
$(iv) \Rightarrow (ii)$ If $T - \lambda I_{\mathcal{V}}$ were injective, Theorem 1.7.8 would make it invertible, a contradiction.

*Structural Commentary*: In finite dimensions, noninvertibility has equivalent forms.
*Key Insight*: A scalar $\lambda$ is spectral precisely when $T - \lambda I_{\mathcal{V}}$ has a nontrivial kernel. □

**Theorem 5.1.5.** *Eigenvectors for Distinct Eigenvalues Are Independent*
*Let $T \in \mathcal{L}(\mathcal{V})$. Suppose $\lambda_1, \ldots, \lambda_m$ are distinct eigenvalues of $T$, and let $\mathbf{v}_1, \ldots, \mathbf{v}_m$ be corresponding eigenvectors. Then $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly independent.*

*Proof.* Suppose the list is linearly dependent. Let $k$ be smallest positive integer s.t. $\mathbf{v}_k \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$. Then $k \geq 2$, $(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$ is linearly independent, and $\exists$ scalars $a_1, \ldots, a_{k-1}$ s.t. $\mathbf{v}_k = a_1\mathbf{v}_1 + \cdots + a_{k-1}\mathbf{v}_{k-1}$. Applying $T$ gives $\lambda_k\mathbf{v}_k = a_1\lambda_1\mathbf{v}_1 + \cdots + a_{k-1}\lambda_{k-1}\mathbf{v}_{k-1}$. Multiplying first displayed equation by $\lambda_k$ and subtracting gives $0 = a_1(\lambda_k - \lambda_1)\mathbf{v}_1 + \cdots + a_{k-1}(\lambda_k - \lambda_{k-1})\mathbf{v}_{k-1}$. Since $\mathbf{v}_1, \ldots, \mathbf{v}_{k-1}$ are linearly independent and $\lambda_k \neq \lambda_i$ for $i < k$, all $a_i = 0$. Hence $\mathbf{v}_k = \mathbf{0}$, contradicting that $\mathbf{v}_k$ is an eigenvector.
*Structural Commentary*: A smallest dependence isolates the first new eigenvector that depends on earlier ones. Applying $T$ and subtracting separates the distinct eigenvalue factors.
*Key Insight*: Distinct eigenvalues make the same dependence relation impossible after applying $T$. □

**Corollary 5.1.6.** *Number of Distinct Eigenvalues*
*If $\mathcal{V}$ is finite-dimensional and $T \in \mathcal{L}(\mathcal{V})$, then $T$ has at most $\dim(\mathcal{V})$ distinct eigenvalues.*

*Proof.* If $T$ had more than $\dim(\mathcal{V})$ distinct eigenvalues, choose $m > \dim(\mathcal{V})$ of them, one eigenvector for each. By Theorem 5.1.5, these eigenvectors form linearly independent list of length $m$, contradicting Theorem 1.3.10.
*Structural Commentary*: Distinct eigenvalues give independent directions.
*Key Insight*: Count one eigenvector per eigenspace. □

**Corollary 5.1.7.** *Invertibility and the Zero Eigenvalue*
*Let $\mathcal{V}$ be finite-dimensional and let $T \in \mathcal{L}(\mathcal{V})$. Then $T$ is invertible if and only if $0 \notin \operatorname{spec}(T)$.*

*Proof.* Apply Theorem 5.1.4 with $\lambda = 0$. Then 0 is an eigenvalue exactly when $T - 0I_{\mathcal{V}} = T$ is not invertible.
*Structural Commentary*: The zero eigenvalue is the spectral form of singularity.
*Key Insight*: Invertibility is equivalent to having trivial kernel, and $E_0(T) = \ker T$. □

**Corollary 5.1.8.** *Matrix Invertibility and the Spectrum*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{F})$. Then $A$ is invertible if and only if $0 \notin \operatorname{spec}(A)$.*

*Proof.*

$(\Rightarrow)$ If $A$ is invertible, then $\ker(A) = \{\mathbf{0}\}$, so 0 is not an eigenvalue of $A$.
$(\Leftarrow)$ If $0 \notin \operatorname{spec}(A)$, then $\ker(A) = \{\mathbf{0}\}$. Corollary 3.3.3 gives full rank, hence $A$ is invertible.

*Structural Commentary*: Matrix invertibility is the coordinate form of trivial kernel.
*Key Insight*: A zero eigenvalue is exactly a nonzero solution of $A\mathbf{x} = \mathbf{0}$. □

## 5.2 Invariant Subspaces, Reducing Pairs, and Quotient Operators

**Definition 5.2.1.** *Invariant Subspace*
Let $T \in \mathcal{L}(\mathcal{V})$. A subspace $\mathcal{U}$ of $\mathcal{V}$ is *invariant under $T$* if $T\mathbf{u} \in \mathcal{U}$ for every $\mathbf{u} \in \mathcal{U}$.
The set of all $T$-invariant subspaces is $\operatorname{Inv}(T) := \{\mathcal{U} \leq \mathcal{V} : T\mathcal{U} \subseteq \mathcal{U}\}$.
*Key Insight*: $\mathcal{U}$ is invariant under $T$ iff $T|_{\mathcal{U}} \in \mathcal{L}(\mathcal{U})$.

**Proposition 5.2.2.** *One-Dimensional Invariant Subspaces and Eigenvectors*
*Let $T \in \mathcal{L}(\mathcal{V})$, $\mathbf{v} \neq \mathbf{0}$. Then $\operatorname{span}(\mathbf{v})$ is invariant under $T$ if and only if there exists $\lambda \in \mathbb{F}$ such that $T\mathbf{v} = \lambda\mathbf{v}$.*

*Proof.*

($\Rightarrow$) If $\operatorname{span}(\mathbf{v})$ is invariant under $T$, then $T\mathbf{v} \in \operatorname{span}(\mathbf{v})$, so $T\mathbf{v} = \lambda\mathbf{v}$ for some $\lambda \in \mathbb{F}$.
($\Leftarrow$) If $T\mathbf{v} = \lambda\mathbf{v}$ and $\mathbf{u} = a\mathbf{v} \in \operatorname{span}(\mathbf{v})$, then $T\mathbf{u} = aT\mathbf{v} = a\lambda\mathbf{v} \in \operatorname{span}(\mathbf{v})$.

*Structural Commentary*: One-dimensional invariance means rescaling one direction.
*Key Insight*: In dimension one, staying inside the subspace means being a scalar multiple of the same vector. □




Figure 43: An eigendirection is a one-dimensional invariant subspace.

**Proposition 5.2.3.** *Invariant Subspaces and Block Form*
*Let $T \in \mathcal{L}(\mathcal{V})$ and suppose $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.*

*(i) If $\mathcal{U}$ is invariant under $T$, then in a basis adapted to $\mathcal{U} \oplus \mathcal{W}$, matrix of $T$ has upper block-triangular form*

$$\begin{bmatrix} A & B \\ 0 & D \end{bmatrix}.$$

*(ii) If both $\mathcal{U}$ and $\mathcal{W}$ are invariant under $T$, then the matrix of $T$ has the block-diagonal form*

$$\begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix}.$$

*Proof.* Choose basis $(\mathbf{u}_1, \ldots, \mathbf{u}_p)$ of $\mathcal{U}$ and basis $(\mathbf{w}_1, \ldots, \mathbf{w}_q)$ of $\mathcal{W}$, and use concatenation as basis of $\mathcal{V}$.

(i) If $\mathcal{U}$ is invariant, then each $T\mathbf{u}_j$ lies in $\mathcal{U}$, so its coordinates along $\mathbf{w}_1, \ldots, \mathbf{w}_q$ are zero. These entries form the lower-left block.
(ii) If $\mathcal{W}$ is also invariant, then each $T\mathbf{w}_j$ lies in $\mathcal{W}$, so its coordinates along $\mathbf{u}_1, \ldots, \mathbf{u}_p$ are zero. These entries form the upper-right block.

*Structural Commentary*: Block form records how an operator moves vectors between chosen summands. Invariant summands force the corresponding off-diagonal movement to vanish.
*Key Insight*: Zero blocks are coordinate evidence of invariant subspaces. □

**Definition 5.2.4.** *Reducing Pair*
Let $T \in \mathcal{L}(\mathcal{V})$. A pair of subspaces $(\mathcal{U}, \mathcal{W})$ *reduces* $T$ if $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$; $T(\mathcal{U}) \subseteq \mathcal{U}$; and $T(\mathcal{W}) \subseteq \mathcal{W}$.
Then $T = T|_{\mathcal{U}} \oplus T|_{\mathcal{W}}$ with respect to $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.

**Proposition 5.2.5.** *Reducing Pairs and Projections*
*Let $T \in \mathcal{L}(\mathcal{V})$, let $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, and let $\rho = \rho_{\mathcal{U},\mathcal{W}}$. Then $(\mathcal{U}, \mathcal{W})$ reduces $T \iff T\rho = \rho T$.*

*Proof.*

($\Rightarrow$) Let $\mathbf{v} = \mathbf{u} + \mathbf{w}$ with $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{W}$.
Since both summands are invariant, $T\rho\mathbf{v} = T\mathbf{u}$ and $\rho T\mathbf{v} = \rho(T\mathbf{u} + T\mathbf{w}) = T\mathbf{u}$. Thus $T\rho = \rho T$.
($\Leftarrow$) Suppose $T\rho = \rho T$. If $\mathbf{u} \in \mathcal{U}$, then $\rho(T\mathbf{u}) = T(\rho\mathbf{u}) = T\mathbf{u}$, so $T\mathbf{u} \in \operatorname{range}(\rho) = \mathcal{U}$.
If $\mathbf{w} \in \mathcal{W}$, then $\rho(T\mathbf{w}) = T(\rho\mathbf{w}) = \mathbf{0}$, so $T\mathbf{w} \in \ker(\rho) = \mathcal{W}$. Hence $(\mathcal{U}, \mathcal{W})$ reduces $T$.

*Structural Commentary*: Commutation with $\rho$ means $T$ preserves the two projection components.
*Key Insight*: Reducing pairs are exactly decompositions whose projection commutes with $T$. □

**Proposition 5.2.6.** *Invariant Subspaces and Algebraic Projections*
*Let $T \in \mathcal{L}(\mathcal{V})$, let $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, and let $\rho = \rho_{\mathcal{U},\mathcal{W}}$. Then $\mathcal{U} \in \operatorname{Inv}(T) \iff \rho T\rho = T\rho$.*
*Consequently, if $\mathcal{U} \in \operatorname{Inv}(T)$, then $\rho_{\mathcal{U},\mathcal{Z}} T \rho_{\mathcal{U},\mathcal{Z}} = T\rho_{\mathcal{U},\mathcal{Z}}$ for every complement $\mathcal{Z}$ of $\mathcal{U}$.*

*Proof.* For $\mathbf{v} = \mathbf{u} + \mathbf{w}$ with $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{W}$, $\mathcal{U} \in \operatorname{Inv}(T)$ gives $(\rho T\rho - T\rho)\mathbf{v} = \rho T\mathbf{u} - T\mathbf{u} = \mathbf{0}$.
Thus $\rho T\rho = T\rho$. Conversely, if $\rho T\rho = T\rho$ and $\mathbf{u} \in \mathcal{U}$, then $\rho T\mathbf{u} = \rho T\rho\mathbf{u} = T\rho\mathbf{u} = T\mathbf{u}$.
Hence $T\mathbf{u} \in \operatorname{range}(\rho) = \mathcal{U}$, so $\mathcal{U} \in \operatorname{Inv}(T)$.

*Structural Commentary*: A projection onto $\mathcal{U}$ tests whether the $\mathcal{W}$-component of $T\mathbf{u}$ vanishes.
*Key Insight*: Invariance says applying $T$ after projecting into $\mathcal{U}$ needs no further correction by $\rho$. □

**Perspective 5.2.7.** *Matrix Viewpoint on Structure*
Basic block matrices already capture important geometry. Upper block-triangular form records the presence of an invariant subspace; block diagonal form records a genuine decomposition into independent invariant pieces.

**Theorem 5.2.8.** *Invariant Subspaces, Blocks, and Projections*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{F})$, let $\mathcal{U}$ be an $r$-dimensional subspace of $\mathbb{F}^n$ with $1 \le r \le n-1$, and let $Q = [Q_1\ Q_2]$ be unitary, where the columns of $Q_1$ and $Q_2$ are orthonormal bases of $\mathcal{U}$ and $\mathcal{U}^\perp$, respectively. Let $P = Q_1Q_1^*$ be the orthogonal projection onto $\mathcal{U}$. The following are equivalent:*

*(i)* $\mathcal{U}$ *is invariant under* $A$;
*(ii)* $Q^*AQ = \begin{bmatrix} B & X \\ 0 & C \end{bmatrix}$ *for some compatible matrices* $B, X, C$;
*(iii)* $PAP = AP$.

*Proof.* Let $M = Q^*AQ = \begin{bmatrix} B & X \\ Y & C \end{bmatrix}$ and $E = Q^*PQ = I_r \oplus 0_{n-r}$. Since $Q$ is unitary and $P = Q_1Q_1^*$, the matrix of $P$ in the $Q$-coordinates is $E$. Also $Y = Q_2^*AQ_1$, and $\ker(Q_2^*) = \mathrm{col}(Q_1) = \mathcal{U}$.

*(i)* $\Rightarrow$ *(ii)* If $A\mathcal{U} \subseteq \mathcal{U}$, then each column of $AQ_1$ lies in $\mathcal{U} = \ker(Q_2^*)$, so $Y = Q_2^*AQ_1 = 0$.

*(ii)* $\Rightarrow$ *(iii)* If $Y = 0$, then $EME = \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix}$ and $ME = \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix}$.
Thus $Q^*(PAP)Q = EME = ME = Q^*(AP)Q$, so $PAP = AP$.

*(iii)* $\Rightarrow$ *(i)* If $PAP = AP$, then $EME = ME$. Since $EME = \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix}$ and $ME = \begin{bmatrix} B & 0 \\ Y & 0 \end{bmatrix}$, we get $Y = 0$.
Hence the columns of $AQ_1$ lie in $\ker(Q_2^*) = \mathcal{U}$. Since the columns of $Q_1$ span $\mathcal{U}$, linearity gives $A\mathcal{U} \subseteq \mathcal{U}$.

*Structural Commentary*: In adapted coordinates, the lower-left block is the component of $A\mathcal{U}$ in $\mathcal{U}^\perp$.
*Key Insight*: $PAP = AP$ exactly kills this component. □

**Corollary 5.2.9.** *Reducing Subspaces and Commuting Projections*
*Maintain the notation of Theorem 5.2.8. The following are equivalent:*

*(i)* $\mathcal{U}$ *is invariant under both* $A$ *and* $A^*$;
*(ii)* $Q^*AQ = B \oplus C$ *for some* $B \in \mathrm{Mat}(r,r,\mathbb{F})$ *and* $C \in \mathrm{Mat}(n-r,n-r,\mathbb{F})$;
*(iii)* $PA = AP$.

*Proof.* Let $M = Q^*AQ = \begin{bmatrix} B & X \\ Y & C \end{bmatrix}$ and $E = Q^*PQ = I_r \oplus 0_{n-r}$.
Then $Q^*A^*Q = M^*$, $Q^*(PA)Q = EM$, and $Q^*(AP)Q = ME$.

*(i)* $\Rightarrow$ *(ii)* By Theorem 5.2.8, $A$-invariance gives $Y = 0$. Applying the same theorem to $A^*$ gives that the lower-left block of $M^*$ is zero, namely $X^* = 0$, so $X = 0$. Thus $M = B \oplus C$.

*(ii)* $\Rightarrow$ *(iii)* If $M = B \oplus C$, then $EM = ME$, hence $PA = AP$.

*(iii)* $\Rightarrow$ *(i)* If $PA = AP$, then $EM = ME$. Since $EM = \begin{bmatrix} B & X \\ 0 & 0 \end{bmatrix}$ and $ME = \begin{bmatrix} B & 0 \\ Y & 0 \end{bmatrix}$, we get $X = Y = 0$.
The equation $Y = 0$ gives $A$-invariance by Theorem 5.2.8. Since the lower-left block of $M^*$ is $X^* = 0$, applying the same theorem to $A^*$ gives $A^*$-invariance.

*Structural Commentary*: Reducing subspaces are exactly block diagonal decompositions in adapted coordinates.
*Key Insight*: Commutation with $P$ kills both off-diagonal blocks. □




Figure 44: Adapted coordinates: invariance gives upper block form; $PA = AP$ gives block diagonal form.

**Definition 5.2.10.** *Restriction and Quotient Operators*
Let $T \in \mathcal{L}(\mathcal{V})$ and let $\mathcal{U}$ be invariant under $T$.

(i) The *restriction operator* $T|_{\mathcal{U}} \in \mathcal{L}(\mathcal{U})$ is defined by $T|_{\mathcal{U}}(\mathbf{u}) = T\mathbf{u}$ for $\mathbf{u} \in \mathcal{U}$.
(ii) The *quotient operator* $T/\mathcal{U} \in \mathcal{L}(\mathcal{V}/\mathcal{U})$ is defined by $(T/\mathcal{U})(\mathbf{v} + \mathcal{U}) = T\mathbf{v} + \mathcal{U}$, $\mathbf{v} \in \mathcal{V}$.

**Proposition 5.2.11.** *Restriction and Quotient Operators Are Well-Defined*
*Let $T \in \mathcal{L}(\mathcal{V})$ and let $\mathcal{U}$ be invariant under $T$. Then $T|_{\mathcal{U}} \in \mathcal{L}(\mathcal{U})$ and $T/\mathcal{U} \in \mathcal{L}(\mathcal{V}/\mathcal{U})$.*

*Proof.* Since $\mathcal{U}$ is invariant under $T$, $T\mathbf{u} \in \mathcal{U}$ whenever $\mathbf{u} \in \mathcal{U}$, so $T|_{\mathcal{U}}$ maps $\mathcal{U}$ into itself. Its linearity is inherited from $T$. For the quotient operator, suppose $\mathbf{v} + \mathcal{U} = \mathbf{w} + \mathcal{U}$. Then $\mathbf{v} - \mathbf{w} \in \mathcal{U}$.
Invariance gives $T(\mathbf{v} - \mathbf{w}) \in \mathcal{U}$, so $T\mathbf{v} - T\mathbf{w} \in \mathcal{U}$ and hence $T\mathbf{v} + \mathcal{U} = T\mathbf{w} + \mathcal{U}$. Thus $T/\mathcal{U}$ is well-defined.
For $a \in \mathbb{F}$, $(T/\mathcal{U})(a(\mathbf{v} + \mathcal{U}) + (\mathbf{w} + \mathcal{U})) = T(a\mathbf{v} + \mathbf{w}) + \mathcal{U} = a(T\mathbf{v} + \mathcal{U}) + (T\mathbf{w} + \mathcal{U})$, so $T/\mathcal{U}$ is linear.
*Structural Commentary*: Restriction stays inside $\mathcal{U}$; quotient collapses $\mathcal{U}$.
*Key Insight*: Invariance makes both operators well-defined. □

**Perspective 5.2.12.** *Two Smaller Views of an Operator*
When $\mathcal{U}$ is invariant under $T$, the operators $T|_{\mathcal{U}}$ and $T/\mathcal{U}$ split attention between what happens inside $\mathcal{U}$ and what remains after $\mathcal{U}$ is collapsed. These two views often simplify an operator, even though they need not determine the whole operator by themselves.

## 5.3 Triangularization over Complex and Real Fields

**Definition 5.3.1.** *Diagonal and Upper-Triangular Matrices*
The *diagonal* of a square matrix is its upper-left to lower-right entries. A square matrix is *diagonal* if all off-diagonal entries are zero, and *upper triangular* if all entries below the diagonal are zero.

**Theorem 5.3.2.** *Complex Operators Have Eigenvalues*
*Every operator on a finite-dimensional, nonzero complex vector space has an eigenvalue.*

*Proof.* Let $T \in \mathcal{L}(\mathcal{V})$. By Proposition 1.1.6, $T$ has a nonconstant annihilating polynomial. Since $\mathcal{V}$ is complex and nonzero, Theorem 3.3.2 implies that one root of this polynomial is an eigenvalue of $T$.
*Structural Commentary*: Finite dimension gives an annihilating polynomial; over $\mathbb{C}$, it factors linearly.
*Key Insight*: One noninjective factor produces an eigenvector. □

**Theorem 5.3.3.** *Criteria for an Upper-Triangular Operator Matrix*
*Let $T \in \mathcal{L}(\mathcal{V})$ and let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be a basis of $\mathcal{V}$. The following are equivalent:*

*(i)* $[T]_{\beta \leftarrow \beta}$ *is upper triangular;*
*(ii)* $T\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$ *for each* $j = 1, \ldots, n$*;*
*(iii)* $\operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$ *is invariant under* $T$ *for each* $j = 1, \ldots, n$.

*Proof.*

(i) $\Rightarrow$ (ii) The $j$th column of $[T]_{\beta \leftarrow \beta}$ is the coordinate column of $T\mathbf{v}_j$ in the basis $\beta$. Since the matrix is upper triangular, this column has zero entries below position $j$, so $T\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$.
(ii) $\Rightarrow$ (i) If $T\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$ for every $j$, then the $j$th coordinate column of $T\mathbf{v}_j$ has zero entries below position $j$. Hence $[T]_{\beta \leftarrow \beta}$ is upper triangular.
(ii) $\Rightarrow$ (iii) If $\mathbf{u} \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$, then linearity gives $T\mathbf{u}$ as a linear combination of $T\mathbf{v}_1, \ldots, T\mathbf{v}_j$, all of which lie in $\operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$. Hence this span is invariant.
(iii) $\Rightarrow$ (ii) Apply invariance of $\operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$ to $\mathbf{v}_j$.

*Structural Commentary*: Upper triangularity records an invariant flag.
*Key Insight*: Columns below the diagonal vanish exactly when $T\mathbf{v}_j$ uses only earlier basis vectors and itself. □

**Corollary 5.3.4.** *Upper-Triangular Form in an Orthonormal Basis*
*Let $\mathcal{V}$ be a finite-dimensional inner-product space and let $T \in \mathcal{L}(\mathcal{V})$. If $T$ has an upper-triangular matrix with respect to some basis of $\mathcal{V}$, then $T$ has an upper-triangular matrix with respect to some orthonormal basis of $\mathcal{V}$.*

*Proof.* Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be a basis for which $[T]_{\beta \leftarrow \beta}$ is upper triangular. By Theorem 5.3.3, each subspace

$$\mathcal{U}_j = \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j), \qquad j = 1, \ldots, n,$$

is invariant under $T$. Apply Gram–Schmidt to $\beta$ and obtain an orthonormal basis $\gamma = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$ with

$$\operatorname{span}(\mathbf{e}_1, \ldots, \mathbf{e}_j) = \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j) = \mathcal{U}_j.$$

Thus each $\operatorname{span}(\mathbf{e}_1, \ldots, \mathbf{e}_j)$ is invariant under $T$, and Theorem 5.3.3 shows that $[T]_{\gamma \leftarrow \gamma}$ is upper triangular.
*Structural Commentary*: Upper triangularity is preservation of a nested flag of subspaces. Gram–Schmidt

preserves that flag while making the basis orthonormal.
*Key Insight*: Triangular form depends on the flag, not on the particular basis used inside the flag. □

**Theorem 5.3.5.** *Complex Upper Triangularisation*
*Let $\mathcal{V}$ be a finite-dimensional complex vector space and let $T \in \mathcal{L}(\mathcal{V})$. Then $T$ has an upper-triangular matrix with respect to some basis of $\mathcal{V}$.*

*Proof.* Induct on $n = \dim(\mathcal{V})$. If $n \le 1$, the result is immediate.
Suppose $n > 1$ and the result holds in smaller dimensions. Choose an eigenvector $\mathbf{v}_1$ of $T$ by Theorem 5.3.2, and set $\mathcal{U} = \operatorname{span}(\mathbf{v}_1)$. Then $\mathcal{U}$ is $T$-invariant, so the quotient operator $T/\mathcal{U}$ is defined on $\mathcal{V}/\mathcal{U}$, which has dimension $n-1$. By induction, choose a basis $(\mathbf{v}_2+\mathcal{U}, \dots, \mathbf{v}_n+\mathcal{U})$ of $\mathcal{V}/\mathcal{U}$ with respect to which $T/\mathcal{U}$ is upper triangular, and choose representatives $\mathbf{v}_2, \dots, \mathbf{v}_n \in \mathcal{V}$.
The lifted list $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$: any relation descends to the quotient basis, forcing all coefficients of $\mathbf{v}_2, \dots, \mathbf{v}_n$ to be zero, and then the coefficient of $\mathbf{v}_1$ is zero. Now $T\mathbf{v}_1 \in \operatorname{span}(\mathbf{v}_1)$.
For $j \ge 2$, upper triangularity of $T/\mathcal{U}$ gives $T\mathbf{v}_j + \mathcal{U} = (T/\mathcal{U})(\mathbf{v}_j + \mathcal{U}) \in \operatorname{span}(\mathbf{v}_2 + \mathcal{U}, \dots, \mathbf{v}_j + \mathcal{U})$, so $T\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_j)$. Hence Theorem 5.3.3 gives an upper-triangular matrix for $T$ in the basis $(\mathbf{v}_1, \dots, \mathbf{v}_n)$.
*Structural Commentary*: Lift an upper-triangular quotient basis above an eigenline.
*Key Insight*: The quotient builds the remaining invariant flag. □

**Definition 5.3.6.** *Complexification of a Real Vector Space*
Let $\mathcal{V}$ be a real vector space.

(i) The *complexification* of $\mathcal{V}$ is $\mathcal{V}_{\mathbb{C}} := \mathcal{V} \times \mathcal{V}$, with $(\mathbf{u}, \mathbf{v})$ denoted $\mathbf{u} + i\mathbf{v}$.
(ii) Addition and $\mathbb{C}$-scalar multiplication are

$$(\mathbf{u}_1 + i\mathbf{v}_1) + (\mathbf{u}_2 + i\mathbf{v}_2) = (\mathbf{u}_1 + \mathbf{u}_2) + i(\mathbf{v}_1 + \mathbf{v}_2),$$
$$(a + bi)(\mathbf{u} + i\mathbf{v}) = (a\mathbf{u} - b\mathbf{v}) + i(b\mathbf{u} + a\mathbf{v}), \qquad a, b \in \mathbb{R}.$$

(iii) *Complexification map* is real-linear embedding $\operatorname{cpx} : \mathcal{V} \hookrightarrow \mathcal{V}_{\mathbb{C}}$; $\operatorname{cpx}(\mathbf{u}) = \mathbf{u} + i\mathbf{0}$, where $\{\mathbf{u} + i\mathbf{0} : \mathbf{u} \in \mathcal{V}\}$.
(iv) For $\mathbf{z} = \mathbf{u} + i\mathbf{v} \in \mathcal{V}_{\mathbb{C}}$, define $\operatorname{Re}\mathbf{z} := \mathbf{u}$ and $\operatorname{Im}\mathbf{z} := \mathbf{v}$. Hence $\mathbf{z} = \mathbf{0}$ iff $\operatorname{Re}\mathbf{z} = \operatorname{Im}\mathbf{z} = \mathbf{0}$.

**Proposition 5.3.7.** *Complexification Preserves Bases*
*Let $\mathcal{V}$ be real. Then $\mathcal{V}_{\mathbb{C}}$ is complex, and every real basis $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ of $\mathcal{V}$ is a complex basis of $\mathcal{V}_{\mathbb{C}}$.*
*Thus $\dim_{\mathbb{C}}(\mathcal{V}_{\mathbb{C}}) = \dim_{\mathbb{R}}(\mathcal{V})$.*

*Proof.* The vector-space axioms follow componentwise from those of $\mathcal{V}$.
For $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, write $\mathbf{u} = \sum_{j=1}^{n} a_j\mathbf{v}_j$ and $\mathbf{v} = \sum_{j=1}^{n} b_j\mathbf{v}_j$, $a_j, b_j \in \mathbb{R}$. Then $\mathbf{u} + i\mathbf{v} = \sum_{j=1}^{n} (a_j + ib_j)\mathbf{v}_j$, so the list spans.
If $\sum_{j=1}^{n} \lambda_j\mathbf{v}_j = \mathbf{0}$ and $\lambda_j = a_j + ib_j$, then $\sum_{j=1}^{n} a_j\mathbf{v}_j = \mathbf{0}$ and $\sum_{j=1}^{n} b_j\mathbf{v}_j = \mathbf{0}$.
Real linear independence gives $a_j = b_j = 0$ for all $j$, hence $\lambda_j = 0$ for all $j$.
*Structural Commentary*: Complexification keeps the basis length unchanged.
*Key Insight*: Separate real and imaginary parts. □

**Corollary 5.3.8.** *Coordinates Survive Complexification*
*Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a real basis of $\mathcal{V}$ and set $\beta_{\mathbb{C}} := \operatorname{cpx}(\beta)$. Then, for all $\mathbf{v} \in \mathcal{V}$, $[\operatorname{cpx}(\mathbf{v})]_{\beta_{\mathbb{C}}} = [\mathbf{v}]_{\beta} \in \mathbb{R}^n \subseteq \mathbb{C}^n$.*

*Proof.* If $[\mathbf{v}]_{\beta} = (a_1, \dots, a_n)^t$, then $a_j \in \mathbb{R}$ and $\operatorname{cpx}(\mathbf{v}) = \mathbf{v} + i\mathbf{0} = \sum_{j=1}^{n} a_j(\mathbf{v}_j + i\mathbf{0}) = \sum_{j=1}^{n} a_j \operatorname{cpx}(\mathbf{v}_j)$.
Thus $[\operatorname{cpx}(\mathbf{v})]_{\beta_{\mathbb{C}}} = (a_1, \dots, a_n)^t = [\mathbf{v}]_{\beta}$.
*Structural Commentary*: The embedded real copy keeps exactly the same coordinate columns.
*Key Insight*: Complexification enlarges the scalar field, not the coordinates of real vectors. □

**Proposition 5.3.9.** *Complexification as a Real Space*
*If $\mathcal{V}$ has real basis $(\mathbf{v}_1, \dots, \mathbf{v}_n)$, then $\mathcal{V}_{\mathbb{C}}$, as a real vector space, has basis $(\mathbf{v}_1, \dots, \mathbf{v}_n, i\mathbf{v}_1, \dots, i\mathbf{v}_n)$.*
*Hence $\dim_{\mathbb{R}}(\mathcal{V}_{\mathbb{C}}) = 2\dim_{\mathbb{R}}(\mathcal{V})$, and $\operatorname{cpx} : \mathcal{V} \to \mathcal{V}_{\mathbb{C}}$, $\operatorname{cpx}(\mathbf{v}) = \mathbf{v} + i\mathbf{0}$, is real-linear and injective.*

*Proof.* For $\mathbf{u} = \sum a_j\mathbf{v}_j$ and $\mathbf{w} = \sum b_j\mathbf{v}_j$, $\mathbf{u} + i\mathbf{w} = \sum a_j\mathbf{v}_j + \sum b_j(i\mathbf{v}_j)$, so the displayed list spans $\mathcal{V}_{\mathbb{C}}$ over $\mathbb{R}$.
If $\sum a_j\mathbf{v}_j + \sum b_j(i\mathbf{v}_j) = \mathbf{0}$, then Re and Im give $\sum a_j\mathbf{v}_j = \sum b_j\mathbf{v}_j = \mathbf{0}$; all $a_j, b_j$ vanish. Thus this is real basis.
For $r \in \mathbb{R}$, $\operatorname{cpx}(r\mathbf{v} + \mathbf{w}) = r\operatorname{cpx}(\mathbf{v}) + \operatorname{cpx}(\mathbf{w})$, and $\ker(\operatorname{cpx}) = \{\mathbf{0}\}$, so cpx is real-linear and injective.
*Structural Commentary*: Complexification is two real copies of $\mathcal{V}$.
*Key Insight*: The real and imaginary parts are independent real copies of the same basis. □

**Definition 5.3.10.** *Complexification of a Real Operator*
Let $\mathcal{V}$ be a real vector space and let $T \in \mathcal{L}(\mathcal{V})$.
The *complexification* of $T$ is the operator $T_{\mathbb{C}} \in \mathcal{L}(\mathcal{V}_{\mathbb{C}})$ defined by $T_{\mathbb{C}}(\mathbf{u} + i\mathbf{v}) := T\mathbf{u} + iT\mathbf{v}$.

**Proposition 5.3.11.** *Complexification Preserves Operator Data*
*Let $\mathcal{V}$ be a real vector space and let $T \in \mathcal{L}(\mathcal{V})$.*

*(i)* $T_{\mathbb{C}} \in \mathcal{L}(\mathcal{V}_{\mathbb{C}})$.
*(ii)* $(T_{\mathbb{C}})^k(\mathbf{u} + i\mathbf{v}) = T^k\mathbf{u} + iT^k\mathbf{v}$ *for every integer* $k \geq 0$.
*(iii) If $\beta$ is a real basis of $\mathcal{V}$, then $[T_{\mathbb{C}}]_{\beta \leftarrow \beta} = [T]_{\beta \leftarrow \beta}$, now regarded as a complex matrix.*

*Proof.*

(i) Additivity is immediate. For $\alpha = a + bi \in \mathbb{C}$,

$$T_{\mathbb{C}}\big(\alpha(\mathbf{u} + i\mathbf{v})\big) = T(a\mathbf{u} - b\mathbf{v}) + iT(b\mathbf{u} + a\mathbf{v}) = (aT\mathbf{u} - bT\mathbf{v}) + i(bT\mathbf{u} + aT\mathbf{v}) = \alpha T_{\mathbb{C}}(\mathbf{u} + i\mathbf{v}),$$

so $T_{\mathbb{C}}$ is complex-linear.

(ii) For $k = 0$, the claim is the identity. If it holds for $k$, then

$$(T_{\mathbb{C}})^{k+1}(\mathbf{u} + i\mathbf{v}) = T_{\mathbb{C}}(T^k\mathbf{u} + iT^k\mathbf{v}) = T^{k+1}\mathbf{u} + iT^{k+1}\mathbf{v}.$$

Thus the formula holds for all $k \geq 0$.

(iii) If $T\mathbf{v}_j = \sum_i a_{ij}\mathbf{v}_i$ with $a_{ij} \in \mathbb{R}$, then $T_{\mathbb{C}}\mathbf{v}_j = \sum_i a_{ij}\mathbf{v}_i$. Hence the coordinate matrix is unchanged.

*Structural Commentary*: Complexification keeps the coordinate matrix unchanged.
*Key Insight*: Same real entries; larger scalar field. □

Figure 45: Complexification packages two real vectors: $T_{\mathbb{C}}(\mathbf{u} + i\mathbf{v}) = T\mathbf{u} + iT\mathbf{v}$.

**Method 5.3.12.** *Complexifying a Real Matrix*
To use complex eigenvalues for a real matrix $A \in \mathrm{Mat}(n, n, \mathbb{R})$:

(i) Regard the same real matrix $A$ as a complex matrix on $\mathbb{C}^n$.
(ii) Find complex eigenpairs or invariant data for this complexified operator.
(iii) If $A(\mathbf{u} + i\mathbf{v}) = (a + bi)(\mathbf{u} + i\mathbf{v})$, separate parts: $A\mathbf{u} = a\mathbf{u} - b\mathbf{v}$, $A\mathbf{v} = b\mathbf{u} + a\mathbf{v}$
(iv) Conclude that $\mathrm{span}(\mathbf{u}, \mathbf{v}) \subseteq \mathbb{R}^n$ is an invariant line or plane.

**Theorem 5.3.13.** *Invariant Line or Plane over $\mathbb{R}$*
*Every operator on a nonzero finite-dimensional real vector space has an invariant subspace of dimension* 1 *or* 2.

*Proof.* Let $\mathcal{V}$ be a nonzero finite-dimensional real vector space and let $T \in \mathcal{L}(\mathcal{V})$.
By Theorem 5.3.2, $T_{\mathbb{C}}$ has an eigenvalue $\lambda = a + bi$ with eigenvector $\mathbf{u} + i\mathbf{v} \neq \mathbf{0}$.
The equation $T_{\mathbb{C}}(\mathbf{u} + i\mathbf{v}) = (a + bi)(\mathbf{u} + i\mathbf{v})$ gives $T\mathbf{u} = a\mathbf{u} - b\mathbf{v}$ and $T\mathbf{v} = b\mathbf{u} + a\mathbf{v}$.
Thus $\mathcal{U} = \mathrm{span}(\mathbf{u}, \mathbf{v})$ is nonzero, has dimension 1 or 2, and is invariant under $T$.
*Structural Commentary*: A complex eigenvector decomposes into two real directions that are closed under $T$.
*Key Insight*: Nonreal eigenvalues become real invariant planes; real eigenvalues become invariant lines. □




Figure 46: Separate a complex eigenpair into a real invariant plane.

**Proposition 5.3.14.** *Real Eigenvalues of a Complexification*
*Let $\mathcal{V}$ be real, let $T \in \mathcal{L}(\mathcal{V})$, and let $\lambda \in \mathbb{R}$. Then $\lambda$ is an eigenvalue of $T$ if and only if it is an eigenvalue of $T_{\mathbb{C}}$.*

*Proof.*

($\Rightarrow$) If $T\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$, then $T_{\mathbb{C}}(\mathbf{v} + i\mathbf{0}) = \lambda(\mathbf{v} + i\mathbf{0})$.
($\Leftarrow$) If $T_{\mathbb{C}}(\mathbf{u} + i\mathbf{v}) = \lambda(\mathbf{u} + i\mathbf{v})$ and $\mathbf{u} + i\mathbf{v} \neq \mathbf{0}$, then $T\mathbf{u} = \lambda\mathbf{u}$ and $T\mathbf{v} = \lambda\mathbf{v}$.
Since $\mathbf{u}$ or $\mathbf{v}$ is nonzero, $T$ has eigenvalue $\lambda$.

*Structural Commentary*: Real eigenvalues survive complexification exactly.
*Key Insight*: For real $\lambda$, real and imaginary parts decouple. □

**Definition 5.3.15.** *Almost Upper-Triangular Real Matrices*
Let $A \in \mathrm{Mat}(n, n, \mathbb{R})$. The matrix $A$ is *almost upper triangular* if there exist $d_1, \dots, d_m \in \{1, 2\}$ with $\sum_j d_j = n$ such that, in the corresponding block form $A = [A_{ij}]_{i,j=1}^{m}$,

(i) $A_{ij} = 0$ for $i > j$;

(ii) each diagonal block is either $A_{jj} = [\lambda_j]$, $\lambda_j \in \mathbb{R}$, or $A_{jj} = \begin{bmatrix} a_j & -b_j \\ b_j & a_j \end{bmatrix}$, $a_j, b_j \in \mathbb{R}$, $b_j \neq 0$.

The matrix $A$ is in *real Schur block form* if only (i) holds for some such block sizes $d_j \in \{1, 2\}$.

**Theorem 5.3.16.** *Real Almost Upper Triangularisation*
*Let $\dim_{\mathbb{R}} \mathcal{V} < \infty$ and let $T \in \mathcal{L}(\mathcal{V})$. Then $\exists$ a basis $\beta$ of $\mathcal{V}$ such that $[T]_{\beta \leftarrow \beta}$ is almost upper triangular.*

*Proof.* Induct on $n = \dim_{\mathbb{R}} \mathcal{V}$. If $n \leq 1$, clear.
For $n \geq 2$, Theorem 5.3.2 gives an eigenpair $T_{\mathbb{C}}(\mathbf{u} + i\mathbf{v}) = (a + bi)(\mathbf{u} + i\mathbf{v})$, where $\mathbf{u} + i\mathbf{v} \neq \mathbf{0}$.
Thus $T\mathbf{u} = a\mathbf{u} - b\mathbf{v}$ and $T\mathbf{v} = b\mathbf{u} + a\mathbf{v}$.
If $b = 0$, choose $\mathbf{0} \neq \mathbf{x} \in \{\mathbf{u}, \mathbf{v}\}$, set $\mathcal{U} = \mathrm{span}(\mathbf{x})$, $\alpha = (\mathbf{x})$, so $[T|_{\mathcal{U}}]_{\alpha \leftarrow \alpha} = [a]$.
If $b \neq 0$, set $\mathcal{U} = \mathrm{span}(\mathbf{u}, \mathbf{v})$, $\alpha = (\mathbf{u}, -\mathbf{v})$; then $\mathbf{u}, \mathbf{v}$ are independent and $[T|_{\mathcal{U}}]_{\alpha \leftarrow \alpha} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$.
In either case, $T(\mathcal{U}) \subseteq \mathcal{U}$, $\dim \mathcal{U} \in \{1, 2\}$, and the first diagonal block is admissible.
By induction, $T/\mathcal{U}$ has an almost upper-triangular matrix in some quotient basis $\overline{\gamma}$.
Lift $\overline{\gamma}$ to $\gamma \subseteq \mathcal{V}$ and set $\beta = (\alpha, \gamma)$. Then $[T]_{\beta \leftarrow \beta}$ is block upper triangular with first block $[T|_{\mathcal{U}}]_{\alpha \leftarrow \alpha}$ and quotient block $[T/\mathcal{U}]_{\overline{\gamma} \leftarrow \overline{\gamma}}$, hence is almost upper triangular.
*Structural Commentary*: Real triangularisation starts with an invariant line or plane.
*Key Insight*: Complex eigenvectors split into real and imaginary parts; those parts form the first real block. □

**Corollary 5.3.17.** *Orthonormal Real Schur Block Form*
*Let $\dim_{\mathbb{R}} \mathcal{V} < \infty$ and let $T \in \mathcal{L}(\mathcal{V})$. Then $\exists$ an orthonormal basis $\gamma$ of $\mathcal{V}$ s.t. $[T]_{\gamma \leftarrow \gamma}$ is in real Schur block form.*

*Proof.* By Theorem 5.3.16, choose $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ with block sizes $d_j \in \{1, 2\}$.
Put $s_k = d_1 + \cdots + d_k$ and $\mathcal{U}_k = \mathrm{span}(\mathbf{v}_1, \dots, \mathbf{v}_{s_k})$. Then $T(\mathcal{U}_k) \subseteq \mathcal{U}_k$ for all $k$.
Apply Gram–Schmidt to $\beta$ and obtain an orthonormal basis $\gamma = (\mathbf{e}_1, \dots, \mathbf{e}_n)$ with $\mathrm{span}(\mathbf{e}_1, \dots, \mathbf{e}_{s_k}) = \mathcal{U}_k$, $1 \leq k \leq m$. Thus $[T]_{\gamma \leftarrow \gamma}$ is block upper triangular for the same $d_j \in \{1, 2\}$, hence is in real Schur block form.
*Structural Commentary*: Gram–Schmidt preserves the invariant block flag.
*Key Insight*: Orthonormalization changes blocks, not the flag. □

**Proposition 5.3.18.** *Invertibility from an Upper-Triangular Matrix*
*Suppose $T \in \mathcal{L}(\mathcal{V})$ has an upper-triangular matrix with diagonal entries $\lambda_1, \dots, \lambda_n$ with respect to some basis of $\mathcal{V}$. Then $T$ is invertible if and only if $\lambda_j \neq 0$ for every $j$.*

*Proof.* Let $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ be such a basis.

($\Leftarrow$) If every $\lambda_j \neq 0$, then upper-triangular matrix $[T]_{\beta \leftarrow \beta}$ is invertible by Theorem 3.8.5, so $T$ is invertible.

($\Rightarrow$) Suppose $T$ is invertible. If $\lambda_j = 0$ for some $j$, set $\mathcal{U}_j = \mathrm{span}(\mathbf{v}_1, \dots, \mathbf{v}_j)$ and $\mathcal{U}_0 = \{\mathbf{0}\}$. Upper triangularity gives $T(\mathcal{U}_j) \subseteq \mathcal{U}_{j-1}$ because the coefficient of $\mathbf{v}_j$ in $T\mathbf{v}_j$ is $\lambda_j = 0$ and $T\mathbf{v}_i \in \mathcal{U}_{j-1}$ for $i < j$.
Since $\dim(\mathcal{U}_j) > \dim(\mathcal{U}_{j-1})$, the map $\mathcal{U}_j \to \mathcal{U}_{j-1}$ induced by $T$ is not injective. Thus there is a nonzero $\mathbf{u} \in \mathcal{U}_j$ with $T\mathbf{u} = \mathbf{0}$, contradicting invertibility. Hence every $\lambda_j \neq 0$.

*Structural Commentary*: Diagonal entries control each flag layer.
*Key Insight*: A zero diagonal entry forces a kernel vector. □

**Proposition 5.3.19.** *Eigenvalues from an Upper-Triangular Matrix*
*Suppose $T \in \mathcal{L}(\mathcal{V})$ has an upper-triangular matrix with diagonal entries $\lambda_1, \dots, \lambda_n$ with respect to some basis of $\mathcal{V}$. Then the eigenvalues of $T$ are precisely $\lambda_1, \dots, \lambda_n$.*

*Proof.* For $\lambda \in \mathbb{F}$, the matrix of $T - \lambda I_{\mathcal{V}}$ in the same basis is upper triangular with diagonal entries $\lambda_1 - \lambda, \dots, \lambda_n - \lambda$. By Proposition 5.3.18, $T - \lambda I_{\mathcal{V}}$ is not invertible if and only if $\lambda = \lambda_j$ for some $j$. By Theorem 5.1.4, this is equivalent to $\lambda$ being an eigenvalue of $T$.
*Structural Commentary*: Scalar shifts preserve triangular form.
*Key Insight*: Eigenvalues occur exactly when one shifted diagonal entry becomes zero. □

**Perspective 5.3.20.** *Why Triangular Form Matters*
Upper-triangular form is the first useful simplification beyond finding one eigenvector. It preserves enough of the operator to read eigenvalues from the diagonal while still allowing nontrivial coupling above the diagonal.

## 5.4 Diagonalization, Commutants, and Simultaneous Forms

**Definition 5.4.1.** *Diagonalizable Operator*
An operator $T \in \mathcal{L}(\mathcal{V})$ is *diagonalizable* if $T$ has a diagonal matrix with respect to some basis of $\mathcal{V}$.

**Proposition 5.4.2.** *Eigenspaces for Distinct Eigenvalues Form a Direct Sum*
*Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and suppose $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of $T$. Then $E_{\lambda_1}(T) + \cdots + E_{\lambda_m}(T)$ is a direct sum. Consequently, $\dim E_{\lambda_1}(T) + \cdots + \dim E_{\lambda_m}(T) \leq \dim(\mathcal{V})$.*

*Proof.* Suppose $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$ with $\mathbf{u}_j \in E_{\lambda_j}(T)$. If some $\mathbf{u}_j$ were nonzero, then the nonzero vectors among $\mathbf{u}_1, \dots, \mathbf{u}_m$ would be eigenvectors corresponding to distinct eigenvalues, and Theorem 5.1.5 would make them linearly independent. This contradicts their sum being $\mathbf{0}$. Hence every $\mathbf{u}_j = \mathbf{0}$, so the sum is direct.
By Theorem 2.3.4, $\dim(E_{\lambda_1}(T) + \cdots + E_{\lambda_m}(T)) = \dim E_{\lambda_1}(T) + \cdots + \dim E_{\lambda_m}(T)$.
The left side is the dimension of a subspace of $\mathcal{V}$, so Proposition 1.4.2 gives the stated inequality.
*Structural Commentary*: Distinct eigenspaces form independent pieces.
*Key Insight*: A zero sum across them would contradict eigenvector independence. □

**Theorem 5.4.3.** *Conditions Equivalent to Diagonalizability*
*Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $\lambda_1, \dots, \lambda_m$ denote the distinct eigenvalues of $T$. Set $n = \dim(\mathcal{V})$. The following are equivalent:*

*(i)* *$T$ is diagonalizable;*
*(ii)* *$\mathcal{V}$ has a basis consisting of eigenvectors of $T$;*
*(iii)* *$\exists$ one-dimensional subspaces $\mathcal{U}_1, \dots, \mathcal{U}_n$ of $\mathcal{V}$, each invariant under $T$, such that $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_n$;*
*(iv)* *$\mathcal{V} = E_{\lambda_1}(T) \oplus \cdots \oplus E_{\lambda_m}(T)$;*
*(v)* *$\dim(\mathcal{V}) = \dim E_{\lambda_1}(T) + \cdots + \dim E_{\lambda_m}(T)$.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* If $T$ is diagonalizable, then $T$ has a diagonal matrix with respect to some basis $(\mathbf{v}_1, \dots, \mathbf{v}_n)$. Hence $T\mathbf{v}_j = a_j\mathbf{v}_j$ for each $j$, so this basis consists of eigenvectors of $T$.
*(ii)* $\Rightarrow$ *(i)* If $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ is a basis of eigenvectors of $T$, then $T\mathbf{v}_j = a_j\mathbf{v}_j$ for each $j$. Thus the matrix of $T$ with respect to this basis is diagonal.
*(ii)* $\Rightarrow$ *(iii)* Let $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ be an eigenbasis, and put $\mathcal{U}_j = \operatorname{span}(\mathbf{v}_j)$. Each $\mathcal{U}_j$ is one-dimensional and invariant under $T$ by Proposition 5.2.2; the basis decomposition gives $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_n$.
*(iii)* $\Rightarrow$ *(ii)* Choose nonzero $\mathbf{v}_j \in \mathcal{U}_j$. Proposition 5.2.2 makes each $\mathbf{v}_j$ an eigenvector of $T$, and the direct-sum decomposition makes $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ a basis of $\mathcal{V}$.
*(ii)* $\Rightarrow$ *(iv)* Every vector in an eigenbasis lies in one of $E_{\lambda_1}(T), \dots, E_{\lambda_m}(T)$, so these eigenspaces span $\mathcal{V}$. Proposition 5.4.2 shows their sum is direct.
*(iv)* $\Rightarrow$ *(v)* If $\mathcal{V} = E_{\lambda_1}(T) \oplus \cdots \oplus E_{\lambda_m}(T)$, then dimensions add by Theorem 2.3.4.
*(v)* $\Rightarrow$ *(ii)* For each $j$, choose a basis $\beta_j$ of $E_{\lambda_j}(T)$, and concatenate these bases into a list $\beta$. The length of $\beta$ is $\dim(\mathcal{V})$. If a linear combination of vectors in $\beta$ is zero, grouping terms by eigenspace gives $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$ with $\mathbf{u}_j \in E_{\lambda_j}(T)$. Proposition 5.4.2 gives $\mathbf{u}_j = \mathbf{0}$ for every $j$; since each $\beta_j$ is independent, all coefficients are zero. Thus $\beta$ is independent, Proposition 1.4.3 makes $\beta$ a basis of $\mathcal{V}$ consisting of eigenvectors of $T$.

*Structural Commentary*: Diagonalizability means the eigenspaces exhaust $\mathcal{V}$. Directness is automatic for distinct eigenspaces, so the real test is spanning, or equivalently the dimension count.
*Key Insight*: A diagonal basis is exactly an eigenbasis. □

**Corollary 5.4.4.** *Enough Eigenvalues Imply Diagonalizability*
*If $\mathcal{V}$ is finite-dimensional and $T \in \mathcal{L}(\mathcal{V})$ has $\dim(\mathcal{V})$ distinct eigenvalues, then $T$ is diagonalizable.*

*Proof.* Let $n = \dim(\mathcal{V})$, and let $\lambda_1, \dots, \lambda_n$ be distinct eigenvalues of $T$. For each $j$, choose an eigenvector $\mathbf{v}_j$ corresponding to $\lambda_j$. By Theorem 5.1.5, $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ is linearly independent. Since this list has length $n$, Proposition 1.4.3 makes it a basis of $\mathcal{V}$. Thus $\mathcal{V}$ has an eigenbasis, so $T$ is diagonalizable by Theorem 5.4.3.
*Structural Commentary*: One eigenvector from each distinct eigenvalue already gives a full-size independent list.
*Key Insight*: Once the independent eigenvectors reach $\dim(\mathcal{V})$, they must form a basis. □

**Perspective 5.4.5.** *Diagonalization as Exhaustion by Eigenspaces*
Distinct eigenspaces are already separated by Proposition 5.4.2. Thus diagonalizability is the stronger statement that these separated pieces account for every direction in $\mathcal{V}$.

**Theorem 5.4.6.** *Algebraic Spectral Resolution*
*Let $0 \neq \mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $\Lambda = \{\lambda_1, \dots, \lambda_m\} \subseteq \mathbb{F}$ with $\lambda_i \neq \lambda_j$ for $i \neq j$. The following are equivalent:*

*(i).* *$T$ is diagonalizable and $\operatorname{spec}(T) = \Lambda$.*

*(ii).* $\exists$ *nonzero projections* $\rho_1, \dots, \rho_m \in \mathcal{L}(\mathcal{V})$ *s.t.* $\rho_i\rho_j = 0$ $(i \neq j)$, $\sum_{i=1}^{m} \rho_i = I_{\mathcal{V}}$, $T = \sum_{i=1}^{m} \lambda_i \rho_i$.

*In this case, for each* $i$, $\operatorname{range}(\rho_i) = E_{\lambda_i}(T)$; $\ker(\rho_i) = \bigoplus_{j\neq i} E_{\lambda_j}(T)$; *and* $\rho_i = \prod_{j\neq i} \frac{T-\lambda_j I_{\mathcal{V}}}{\lambda_i-\lambda_j}$

*Proof.*

$(i) \Rightarrow (ii)$ By Theorem 5.4.3, $\mathcal{V} = \bigoplus_{i=1}^{m} E_{\lambda_i}(T)$. Let $\rho_i = \rho_{E_{\lambda_i}(T), \bigoplus_{j\neq i} E_{\lambda_j}(T)}$.

Then $\rho_i\rho_j = 0$ for $i \neq j$, $\sum_{i=1}^{m} \rho_i = I_{\mathcal{V}}$, and $T = \sum_{i=1}^{m} \lambda_i\rho_i$. Also $\operatorname{range}(\rho_i)$ and $\ker(\rho_i)$ are as displayed.

For $\ell_i(z) = \prod_{j\neq i} \frac{z-\lambda_j}{\lambda_i-\lambda_j}$, we have $\ell_i(\lambda_k) = \delta_{ik}$, hence $\ell_i(T) = \rho_i$.

$(ii) \Rightarrow (i)$ Put $\mathcal{V}_i = \operatorname{range}(\rho_i)$. By Theorem 2.4.5, $\mathcal{V} = \bigoplus_{i=1}^{m} \mathcal{V}_i$.

If $\mathbf{x} \in \mathcal{V}_i$, then $\rho_i\mathbf{x} = \mathbf{x}$ and $\rho_j\mathbf{x} = \mathbf{0}$ for $j \neq i$, so $T\mathbf{x} = \lambda_i\mathbf{x}$. Thus $\mathcal{V}_i \subseteq E_{\lambda_i}(T)$.

Conversely, if $T\mathbf{x} = \lambda_i\mathbf{x}$ and $\mathbf{x} = \sum_{j=1}^{m} \mathbf{x}_j$ with $\mathbf{x}_j \in \mathcal{V}_j$, then $\sum_{j=1}^{m} (\lambda_j - \lambda_i)\mathbf{x}_j = \mathbf{0}$, so $\mathbf{x}_j = \mathbf{0}$ for $j \neq i$.

Hence $\mathcal{V}_i = E_{\lambda_i}(T)$. Since each $\mathcal{V}_i \neq 0$, each $\lambda_i \in \operatorname{spec}(T)$; concatenating bases of the $\mathcal{V}_i$ gives an eigenbasis, so $T$ is diagonalizable and $\operatorname{spec}(T) = \Lambda$.

The kernel and polynomial formulas follow from $\mathcal{V} = \bigoplus_{i=1}^{m} \mathcal{V}_i$ and $\ell_i(\lambda_k) = \delta_{ik}$.

*Structural Commentary*: A diagonalizable operator is a weighted sum of the projections onto its eigenspaces.
*Key Insight*: Spectral projections are Lagrange polynomials in the operator. □

**Lemma 5.4.7.** *Diagonalizable Restrictions to Invariant Subspaces*
*Let* $T \in \mathcal{L}(\mathcal{V})$ *be diagonalizable and let* $\mathcal{U} \in \operatorname{Inv}(T)$. *For* $\lambda \in \operatorname{spec}(T)$, *set* $\mathcal{U}_\lambda = \mathcal{U} \cap E_\lambda(T)$.
*Then* $\mathcal{U} = \bigoplus_{\lambda\in\operatorname{spec}(T)} \mathcal{U}_\lambda$, *where* $T|_{\mathcal{U}}$ *is diagonalizable.*

*Proof.* Let $\rho_\lambda = p_\lambda(T)$ be the spectral projection from Theorem 5.4.6. Since $\mathcal{U} \in \operatorname{Inv}(T)$, $\rho_\lambda(\mathcal{U}) \subseteq \mathcal{U}$.
Hence, for $\mathbf{u} \in \mathcal{U}$, $\mathbf{u} = \sum_{\lambda\in\operatorname{spec}(T)} \rho_\lambda\mathbf{u}$, where $\rho_\lambda\mathbf{u} \in \mathcal{U}_\lambda$. Thus $\mathcal{U} = \sum_{\lambda\in\operatorname{spec}(T)} \mathcal{U}_\lambda$, and the sum is direct since $\mathcal{U}_\lambda \subseteq E_\lambda(T)$. Bases of the nonzero $\mathcal{U}_\lambda$ concatenate to an eigenbasis for $T|_{\mathcal{U}}$.
*Structural Commentary*: Invariant subspaces inherit spectral projections.
*Key Insight*: Polynomial spectral projections cannot move vectors out of a $T$-invariant subspace. □

**Theorem 5.4.8.** *Simultaneous Diagonalisation of Commuting Diagonalizable Operators*
*Let* $\dim \mathcal{V} < \infty$, *and let* $\emptyset \neq \mathcal{F} \subseteq \mathcal{L}(\mathcal{V})$ *be finite with every* $T \in \mathcal{F}$ *diagonalizable.*
*Then* $\forall S, T \in \mathcal{F}$, $ST = TS \iff \exists$ *a basis* $\beta$ *of* $\mathcal{V}$ *such that* $[T]_{\beta\leftarrow\beta}$ *is diagonal* $\forall T \in \mathcal{F}$.

*Proof.*

$(\Rightarrow)$ Induct on $n = \dim \mathcal{V}$. If every $T \in \mathcal{F}$ is scalar, any basis works. Otherwise choose nonscalar $S \in \mathcal{F}$.
By Theorem 5.4.6, $\mathcal{V} = \bigoplus_{\lambda\in\operatorname{spec}(S)} E_\lambda(S)$, with at least two nonzero summands of dimension $< n$.
If $T \in \mathcal{F}$ and $\mathbf{x} \in E_\lambda(S)$, then $S(T\mathbf{x}) = T(S\mathbf{x}) = \lambda T\mathbf{x}$, so $E_\lambda(S) \in \operatorname{Inv}(T)$.
By Lemma 5.4.7, each $T|_{E_\lambda(S)}$ is diagonalizable; the restricted family still commutes.
Induction gives a common eigenbasis of each $E_\lambda(S)$; concatenate over $\lambda$.

$(\Leftarrow)$ For $S, T \in \mathcal{F}$, set $D_S = [S]_{\beta\leftarrow\beta}$ and $D_T = [T]_{\beta\leftarrow\beta}$.
Since $D_S, D_T$ are diagonal, $[ST]_{\beta\leftarrow\beta} = D_SD_T = D_TD_S = [TS]_{\beta\leftarrow\beta}$. Thus $ST = TS$.

*Structural Commentary*: Commutation preserves eigenspaces.
*Key Insight*: Diagonalize one operator, then restrict. □

**Definition 5.4.9.** *Commutant*
For $\emptyset \neq \mathcal{F} \subseteq \operatorname{Mat}(n, n, \mathbb{F})$, the *commutant* of $\mathcal{F}$ is $\operatorname{Comm}(\mathcal{F}) := \{X \in \operatorname{Mat}(n, n, \mathbb{F}) : XA = AX \ \forall A \in \mathcal{F}\}$.
For a single matrix $A$, write $\operatorname{Comm}(A)$.

**Lemma 5.4.10.** *Commuting with a Block Scalar Diagonal Matrix*
*Let* $\lambda_1, \dots, \lambda_k \in \mathbb{F}$ *and let* $\Lambda = \lambda_1 I_{n_1} \oplus \cdots \oplus \lambda_k I_{n_k}$.
*Partition* $A = [A_{ij}] \in \operatorname{Mat}(n, n, \mathbb{F})$ *conformally with* $\Lambda$, *so* $A_{ij} \in \operatorname{Mat}(n_i, n_j, \mathbb{F})$.

*(i) If* $A$ *is block diagonal, then* $A\Lambda = \Lambda A$.
*(ii) If* $A\Lambda = \Lambda A$ *and* $\lambda_1, \dots, \lambda_k$ *are distinct, then* $A$ *is block diagonal.*

*Proof.* Block multiplication gives $(A\Lambda)_{ij} = \lambda_j A_{ij}$ and $(\Lambda A)_{ij} = \lambda_i A_{ij}$. If $A$ is block diagonal, then $A_{ij} = 0$ for $i \neq j$, and these two expressions agree for every block. Conversely, if $A\Lambda = \Lambda A$, then $(\lambda_j - \lambda_i)A_{ij} = 0$ for every $i, j$. Distinctness of the $\lambda_i$ forces $A_{ij} = 0$ whenever $i \neq j$, so $A$ is block diagonal.
*Structural Commentary*: Distinct scalar blocks force commuting matrices to respect the block decomposition.
*Key Insight*: Distinct block scalars make every off-diagonal block visible and force it to vanish. □

**Example 5.4.11.** *Distinctness Is Necessary*
The distinctness hypothesis in Lemma 5.4.10 cannot be dropped. If $\lambda_1 = \cdots = \lambda_k = 1$, then $\Lambda = I_n$ commutes with every matrix in $\mathrm{Mat}(n, n, \mathbb{F})$, including matrices that are not block diagonal for the chosen partition.

**Theorem 5.4.12.** *Commutant of a Diagonalizable Matrix*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ be diagonalizable. Suppose its distinct eigenvalues are $\mu_1, \ldots, \mu_d$, and set $n_i = \dim E_{\mu_i}(A)$. Write $A = S(\mu_1 I_{n_1} \oplus \cdots \oplus \mu_d I_{n_d})S^{-1}$. Then $\mathrm{Comm}(A) = \{S(X_1 \oplus \cdots \oplus X_d)S^{-1} : X_i \in \mathrm{Mat}(n_i, n_i, \mathbb{C})\}$. In particular, $\dim \mathrm{Comm}(A) = n_1^2 + \cdots + n_d^2$.*

*Proof.* Let $\Lambda = \mu_1 I_{n_1} \oplus \cdots \oplus \mu_d I_{n_d}$. Matrix $B$ commutes with $A$ if and only if $S^{-1}BS$ commutes with $\Lambda$.
By Lemma 5.4.10, this happens exactly when $S^{-1}BS = X_1 \oplus \cdots \oplus X_d$ with $X_i \in \mathrm{Mat}(n_i, n_i, \mathbb{C})$.
Dimension count follows because conjugation by $S$ is a vector-space isomorphism and $\dim \mathrm{Mat}(n_i, n_i, \mathbb{C}) = n_i^2$.
*Structural Commentary*: Diagonalizing $A$ reduces commutation to respecting equal-eigenvalue blocks.
*Key Insight*: Distinct eigenvalue blocks cannot mix under a commuting matrix. □

**Theorem 5.4.13.** *Common Eigenvector for a Commuting Family*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$, and let $\mathcal{F} \subseteq \mathcal{L}(\mathcal{V})$ be a nonempty commuting family: $ST = TS$ for all $S, T \in \mathcal{F}$. Then there exists $\mathbf{v} \neq \mathbf{0}$ that is an eigenvector of every operator in $\mathcal{F}$.*

*Proof.* First suppose $\mathcal{F} = \{T_1, \ldots, T_m\}$ is finite, and induct on $m$. The case $m = 1$ is Theorem 5.3.2.
For $m > 1$, choose $\lambda \in \mathrm{spec}(T_1)$ and set $\mathcal{U} = E_\lambda(T_1)$. Then $\mathcal{U} \neq \{\mathbf{0}\}$.
If $\mathbf{u} \in \mathcal{U}$ and $j \geq 2$, commutation gives $(T_1 - \lambda I_\mathcal{V})T_j\mathbf{u} = T_j(T_1 - \lambda I_\mathcal{V})\mathbf{u} = \mathbf{0}$, so $\mathcal{U}$ is invariant under $T_2, \ldots, T_m$.
Their restrictions to $\mathcal{U}$ still commute, and by induction have a common eigenvector $\mathbf{v} \in \mathcal{U}$, $\mathbf{v} \neq \mathbf{0}$.
Since $\mathbf{v} \in E_\lambda(T_1)$, it is also an eigenvector of $T_1$. For general $\mathcal{F}$, the subspace $\mathrm{span}(\mathcal{F})$ of $\mathcal{L}(\mathcal{V})$ is finite-dimensional. Choose a basis $T_1, \ldots, T_m$ of this span from the spanning set $\mathcal{F}$.
The finite case gives a nonzero $\mathbf{v}$ and scalars $\mu_j$ with $T_j\mathbf{v} = \mu_j\mathbf{v}$. If $S \in \mathcal{F}$, write $S = \sum_j a_j T_j$.
Then $S\mathbf{v} = (\sum_j a_j\mu_j)\mathbf{v}$, so $\mathbf{v}$ is an eigenvector of every member of $\mathcal{F}$.
*Structural Commentary*: Commutation makes one eigenspace invariant, reducing the family by one.
*Key Insight*: Find the common eigenvector inside an eigenspace. □

**Theorem 5.4.14.** *Simultaneous Schur Triangularisation*
*Let $\emptyset \neq \mathcal{F} \subseteq \mathrm{Mat}(n, n, \mathbb{C})$ be a commuting family.*
*Then there is a unitary $U \in \mathrm{Mat}(n, n, \mathbb{C})$ such that $U^*AU$ is upper triangular for every $A \in \mathcal{F}$.*
*If every matrix in $\mathcal{F}$ is real and has only real eigenvalues, then $U$ can be chosen real orthogonal.*

*Proof.* Induct on $n$. The case $n = 1$ is immediate.
For $n > 1$, Theorem 5.4.13 gives a common eigenvector; normalize it to a unit vector $\mathbf{x}$ and extend to an orthonormal basis. Let $V = [\mathbf{x}\ V_2]$ be the corresponding unitary matrix.
For every $A \in \mathcal{F}$, since $A\mathbf{x} = \lambda_A\mathbf{x}$, we have $V^*AV = \begin{bmatrix} \lambda_A & * \\ 0 & A' \end{bmatrix}$.
If $A, B \in \mathcal{F}$, then $AB = BA$. The lower-right block of $V^*ABV = (V^*AV)(V^*BV)$ is $A'B'$, while the lower-right block of $V^*BAV$ is $B'A'$. Hence $A'B' = B'A'$, so the compressed family is still commuting.
By induction, choose a unitary $W$ that simultaneously upper triangularises all compressed blocks.
Then $U = V(1 \oplus W)$ is unitary and $U^*AU$ is upper triangular for every $A \in \mathcal{F}$.
For the real case, run the same induction over $\mathbb{R}$. A common real eigenvector exists as follows. If all matrices are scalar, any real unit vector works. Otherwise choose a nonscalar $A_0 \in \mathcal{F}$.
Since $A_0$ has only real eigenvalues, some real eigenspace $\mathcal{U} = E_\lambda(A_0)$ is nonzero; since $A_0$ is not scalar, $\mathcal{U} \neq \mathbb{R}^n$.
Commutation gives $(A_0 - \lambda I)B\mathbf{u} = B(A_0 - \lambda I)\mathbf{u} = \mathbf{0}$ for $B \in \mathcal{F}$ and $\mathbf{u} \in \mathcal{U}$, so $\mathcal{U}$ is invariant under every $B$.
The restricted family on $\mathcal{U}$ is commuting and still has only real eigenvalues: in a basis adapted to $\mathcal{U}$ each $B$ has block form $\begin{bmatrix} B|_\mathcal{U} & * \\ 0 & * \end{bmatrix}$, so the characteristic polynomial of $B|_\mathcal{U}$ divides that of $B$. Induction on $\dim(\mathcal{U})$ gives a common real eigenvector. Extending by real orthonormal bases gives a real orthogonal $U$.
*Structural Commentary*: A common eigenline creates the first block; commutation descends to the compression.
*Key Insight*: Simultaneous triangularisation is Schur induction with a common eigenvector at every step. □




Figure 47: A shared eigenline gives a simultaneous first block; lower-right compressions still commute.

**Corollary 5.4.15.** *Eigenvalues of Sums and Products of Commuting Matrices*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{C})$ and $AB = BA$. Their eigenvalues can be ordered as $\lambda_1, \dots, \lambda_n$ and $\mu_1, \dots, \mu_n$ so that $A + B$ has eigenvalues $\lambda_i + \mu_i$ and $AB$ has eigenvalues $\lambda_i \mu_i$, counted with multiplicity.*

*Proof.* Choose simultaneous Schur forms $T = U^*AU$ and $R = U^*BU$. Set $\lambda_i = t_{ii}$ and $\mu_i = r_{ii}$. Then $T + R$ and $TR$ are upper triangular with diagonals $\lambda_i + \mu_i$ and $\lambda_i \mu_i$, and are similar to $A + B$ and $AB$.
*Structural Commentary*: Simultaneous Schur form aligns both spectra on one diagonal.
*Key Insight*: In common triangular coordinates, diagonals add and multiply. □

# 6 Spectral Theorems and Operator Functions

## 6.1 Normal, Self-Adjoint, and Spectral Theorems

Structured spectral theory studies operators whose geometry and spectrum align. Normal, self-adjoint, positive, and unitary maps admit orthogonal decompositions and spectral resolutions.

**Definition 6.1.1.** *Normal Operator*
An operator $T \in \mathcal{L}(\mathcal{V})$ on a finite-dim inner-product space is *normal* if it commutes with its adjoint: $TT^* = T^*T$.
A square matrix $A$ is *normal* if $A^*A = AA^*$.
Every self-adjoint operator, every Hermitian matrix, and every unitary matrix is normal.

**Theorem 6.1.2.** *Normal Operators and Equal Adjoint Norms*
*An operator $T \in \mathcal{L}(\mathcal{V})$ is normal if and only if $\|T\mathbf{v}\| = \|T^*\mathbf{v}\|$ for every $\mathbf{v} \in \mathcal{V}$.*

*Proof.* The operator $A = T^*T - TT^*$ is self-adjoint.
For every $\mathbf{v} \in \mathcal{V}$, $\langle A\mathbf{v}, \mathbf{v}\rangle = \langle T^*T\mathbf{v}, \mathbf{v}\rangle - \langle TT^*\mathbf{v}, \mathbf{v}\rangle = \|T\mathbf{v}\|^2 - \|T^*\mathbf{v}\|^2$.
Thus $T$ is normal exactly when $A = 0$, which by Proposition 2.5.4 is equivalent to the displayed quadratic form being zero for every $\mathbf{v}$. This is equivalent to $\|T\mathbf{v}\| = \|T^*\mathbf{v}\|$ for every $\mathbf{v}$.
*Structural Commentary*: Normality is measured by the self-adjoint commutator $T^*T - TT^*$.
*Key Insight*: A normal operator and its adjoint move every vector the same distance. □

**Theorem 6.1.3.** *Normal Operators Share Eigenvectors with Their Adjoints*
*Let $T \in \mathcal{L}(\mathcal{V})$ be normal. If $T\mathbf{v} = \lambda\mathbf{v}$ for some $\mathbf{v} \neq \mathbf{0}$, then $T^*\mathbf{v} = \overline{\lambda}\mathbf{v}$.*

*Proof.* The operator $T - \lambda I_{\mathcal{V}}$ is normal by Proposition 2.4.8 and scalar multiples of $I_{\mathcal{V}}$ commute with $T$ and $T^*$. Applying Theorem 6.1.2 to $T - \lambda I_{\mathcal{V}}$ gives $0 = \|(T - \lambda I_{\mathcal{V}})\mathbf{v}\| = \|(T - \lambda I_{\mathcal{V}})^*\mathbf{v}\| = \|(T^* - \overline{\lambda} I_{\mathcal{V}})\mathbf{v}\|$.
*Structural Commentary*: For normal operators, kernels of $T - \lambda I$ and $T^* - \overline{\lambda} I$ coincide.
*Key Insight*: Normality makes eigenvector information visible from both $T$ and $T^*$. □




Figure 48: For normal $T$, $T\mathbf{v} = \lambda\mathbf{v}$ implies $T^*\mathbf{v} = \overline{\lambda}\mathbf{v}$.

**Theorem 6.1.4.** *Orthogonal Eigenvectors for Normal Operators*
*Let $T \in \mathcal{L}(\mathcal{V})$ be normal. Eigenvectors of $T$ corresponding to distinct eigenvalues are orthogonal.*

*Proof.* Suppose $T\mathbf{u} = \alpha\mathbf{u}$ and $T\mathbf{v} = \beta\mathbf{v}$ with $\alpha \neq \beta$. By Theorem 6.1.3, $T^*\mathbf{v} = \overline{\beta}\mathbf{v}$.
Hence $\alpha \langle \mathbf{u}, \mathbf{v}\rangle = \langle T\mathbf{u}, \mathbf{v}\rangle = \langle \mathbf{u}, T^*\mathbf{v}\rangle = \langle \mathbf{u}, \overline{\beta}\mathbf{v}\rangle = \beta \langle \mathbf{u}, \mathbf{v}\rangle$. Since $\alpha - \beta \neq 0$, $\langle \mathbf{u}, \mathbf{v}\rangle = 0$.
*Structural Commentary*: Normality lets the operator move from one eigenvector to the other through the adjoint.
*Key Insight*: Distinct eigenvalue scalings are incompatible unless the inner product is zero. □

**Theorem 6.1.5.** *Eigenvalues of Self-Adjoint Operators Are Real*
*Every eigenvalue of a self-adjoint operator on a finite-dimensional inner-product space is real.*

*Proof.* Let $T = T^*$ and $T\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$. Then $\lambda \|\mathbf{v}\|^2 = \langle \lambda\mathbf{v}, \mathbf{v}\rangle = \langle T\mathbf{v}, \mathbf{v}\rangle = \langle \mathbf{v}, T\mathbf{v}\rangle = \langle \mathbf{v}, \lambda\mathbf{v}\rangle = \overline{\lambda}\|\mathbf{v}\|^2$. Since $\|\mathbf{v}\|^2 > 0$, $\lambda = \overline{\lambda}$, so $\lambda \in \mathbb{R}$.
*Structural Commentary*: Self-adjointness moves $T$ between inner-product slots without changing the value.
*Key Insight*: An eigenvalue must equal its conjugate because the eigenvector has positive squared norm. □

**Proposition 6.1.6.** *Invertible Quadratic Expressions*
*Let $T \in \mathcal{L}(\mathcal{V})$ be self-adjoint on a finite-dimensional inner-product space.*
*If $b, c \in \mathbb{R}$ and $b^2 < 4c$, then $T^2 + bT + cI_{\mathcal{V}}$ is invertible.*

*Proof.* Let $S = T^2 + bT + cI_{\mathcal{V}}$. For $\mathbf{v} \neq \mathbf{0}$,

$$\begin{aligned}\langle S\mathbf{v}, \mathbf{v}\rangle &= \|T\mathbf{v}\|^2 + b\langle T\mathbf{v}, \mathbf{v}\rangle + c\|\mathbf{v}\|^2 \geq \|T\mathbf{v}\|^2 - |b|\,\|T\mathbf{v}\|\,\|\mathbf{v}\| + c\|\mathbf{v}\|^2 \\ &= \left(\|T\mathbf{v}\| - \frac{|b|}{2}\|\mathbf{v}\|\right)^2 + \left(c - \frac{b^2}{4}\right)\|\mathbf{v}\|^2 > 0\end{aligned}$$

Thus $S\mathbf{v} \neq \mathbf{0}$ for every nonzero $\mathbf{v}$, so $\ker(S) = \{\mathbf{0}\}$. In finite dimensions, injectivity implies invertibility.
*Structural Commentary*: Negative discriminant keeps this self-adjoint quadratic positive.
*Key Insight*: Self-adjointness converts $\langle T^2\mathbf{v}, \mathbf{v}\rangle$ into $\|T\mathbf{v}\|^2$, so completing the square controls the kernel. □

**Lemma 6.1.7.** *Self-Adjoint Operators Have Eigenvalues*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be a finite-dimensional real inner-product space. Every self-adjoint $T \in \mathcal{L}(\mathcal{V})$ has an eigenvalue.*

*Proof.* Choose $\mathbf{v} \neq \mathbf{0}$ and set $n = \dim(\mathcal{V})$.
The list $\mathbf{v}, T\mathbf{v}, \ldots, T^n\mathbf{v}$ is linearly dependent, so some nonzero real polynomial $p$ satisfies $p(T)\mathbf{v} = \mathbf{0}$.
Factor $p$ over $\mathbb{R}$ as $p(x) = a\prod_{j=1}^{M}(x^2 + b_j x + c_j)\prod_{k=1}^{m}(x - \lambda_k)$, where $a \neq 0$, $b_j^2 < 4c_j$, $\lambda_k \in \mathbb{R}$.
By Proposition 6.1.6, each quadratic factor $Q_j(T) := T^2 + b_j T + c_j I_{\mathcal{V}}$ is invertible. Let $R(T) = \prod_j Q_j(T)$ and $L(T) = \prod_k (T - \lambda_k I_{\mathcal{V}})$. The factors commute because they are polynomials in $T$, and $R(T)$ is invertible.
If $m = 0$, then $p(T) = aR(T)$ would be invertible, contradicting $p(T)\mathbf{v} = \mathbf{0}$ with $\mathbf{v} \neq \mathbf{0}$. Hence $m \geq 1$.
Since $\mathbf{0} = p(T)\mathbf{v} = aR(T)L(T)\mathbf{v}$ and $aR(T)$ is invertible, $L(T)\mathbf{v} = \mathbf{0}$. Thus $L(T)$ has nontrivial kernel.
If every $T - \lambda_k I_{\mathcal{V}}$ were injective, their product $L(T)$ would be injective, a contradiction. Therefore some $T - \lambda_k I_{\mathcal{V}}$ is not injective, and $\lambda_k$ is an eigenvalue of $T$.
*Structural Commentary*: An annihilating polynomial remains; quadratic factors are invertible.
*Key Insight*: The nontrivial kernel must come from a real linear factor, hence from an eigenvalue. □

**Proposition 6.1.8.** *Self-Adjoint Operators and Invariant Subspaces*
*Let $T \in \mathcal{L}(\mathcal{V})$ be self-adjoint, and let $\mathcal{U}$ be a subspace of $\mathcal{V}$ invariant under $T$. Then:*

*i. $\mathcal{U}^\perp$ is invariant under $T$;*
*ii. $T|_{\mathcal{U}} \in \mathcal{L}(\mathcal{U})$ is self-adjoint;*
*iii. $T|_{\mathcal{U}^\perp} \in \mathcal{L}(\mathcal{U}^\perp)$ is self-adjoint.*

*Proof.*

(i) If $\mathbf{v} \in \mathcal{U}^\perp$ and $\mathbf{u} \in \mathcal{U}$, then $\langle T\mathbf{v}, \mathbf{u}\rangle = \langle \mathbf{v}, T\mathbf{u}\rangle = 0$ because $T\mathbf{u} \in \mathcal{U}$. Hence $T\mathbf{v} \in \mathcal{U}^\perp$.
(ii) For $\mathbf{u}, \mathbf{v} \in \mathcal{U}$, $\langle (T|_{\mathcal{U}})\mathbf{u}, \mathbf{v}\rangle = \langle T\mathbf{u}, \mathbf{v}\rangle = \langle \mathbf{u}, T\mathbf{v}\rangle = \langle \mathbf{u}, (T|_{\mathcal{U}})\mathbf{v}\rangle$. Thus $T|_{\mathcal{U}}$ is self-adjoint.
(iii) By (i), $\mathcal{U}^\perp$ is invariant, so (ii)'s argument applies to $T|_{\mathcal{U}^\perp}$.

*Structural Commentary*: Self-adjointness preserves orthogonal complements of invariant subspaces.
*Key Insight*: Move $T$ across the inner product; invariance of $\mathcal{U}$ then forces orthogonality to persist. □

**Theorem 6.1.9.** *Schur Triangularisation*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$. Then $T$ has an upper-triangular matrix with respect to some orthonormal basis of $\mathcal{V}$.*

*Proof.* Proceed by induction on $n = \dim(\mathcal{V})$. The cases $n = 0$ and $n = 1$ are immediate.
Suppose $n \geq 2$ and the result holds for smaller dimensions. By Theorem 5.3.2, choose a unit eigenvector $\mathbf{e}_1$ of $T$, say $T\mathbf{e}_1 = \lambda\mathbf{e}_1$. Let $\mathcal{U} = \operatorname{span}(\mathbf{e}_1)$ and define $S \in \mathcal{L}(\mathcal{U}^\perp)$ by $S\mathbf{w} = P_{\mathcal{U}^\perp}(T\mathbf{w})$.
By induction, choose an orthonormal basis $(\mathbf{e}_2, \ldots, \mathbf{e}_n)$ of $\mathcal{U}^\perp$ with respect to which $S$ has an upper-triangular matrix. Then $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ is an orthonormal basis of $\mathcal{V}$.
For $j \geq 2$, the coordinates of $T\mathbf{e}_j$ along $\mathbf{e}_2, \ldots, \mathbf{e}_n$ are exactly the coordinates of $S\mathbf{e}_j$, hence have no entries below row $j$. Also $T\mathbf{e}_1 = \lambda\mathbf{e}_1$. Therefore $[T]_{\gamma\leftarrow\gamma}$ is upper triangular for $\gamma = (\mathbf{e}_1, \ldots, \mathbf{e}_n)$.
*Structural Commentary*: Schur triangularisation builds an eigenline first, then triangularises the projected action

on its orthogonal complement.
*Key Insight*: The lower-right block is controlled by the compression $P_{\mathcal{U}^\perp} T|_{\mathcal{U}^\perp}$. □

**Theorem 6.1.10.** *Matrix Schur Triangularisation with Ordered Diagonal*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ have eigenvalues $\lambda_1, \dots, \lambda_n$ in a prescribed order. If $A\mathbf{x} = \lambda_1 \mathbf{x}$ and $\|\mathbf{x}\|_2 = 1$, then there is a unitary $U = [\mathbf{x}\ U_2]$ such that $U^* A U$ is upper triangular with diagonal $\lambda_1, \dots, \lambda_n$.*
*If $A$ is real, all $\lambda_i \in \mathbb{R}$, and $\mathbf{x} \in \mathbb{R}^n$, then $U$ may be chosen real orthogonal.*

*Proof.* Induct on $n$. The case $n = 1$ is immediate. Extend $\mathbf{x}$ to an orthonormal basis and set $V = [\mathbf{x}\ V_2]$.
Since $A\mathbf{x} = \lambda_1 \mathbf{x}$, the first column of $V^* A V$ is $\lambda_1 \mathbf{e}_1$, so $V^* A V = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$.
Block triangularity gives $p_{V^*AV}(z) = (z - \lambda_1) p_{A'}(z)$, so $A'$ has eigenvalues $\lambda_2, \dots, \lambda_n$, counted with algebraic multiplicity. By induction, choose unitary $W$ with $W^* A' W = T'$ upper triangular and diagonal $\lambda_2, \dots, \lambda_n$.
Then $U := V(I_1 \oplus W)$ has first column $\mathbf{x}$ and $U^* A U = \begin{bmatrix} \lambda_1 & * \\ 0 & T' \end{bmatrix}$.
The real case uses real orthonormal extensions; the block $A'$ is real, and the same characteristic-polynomial factorization leaves the real eigenvalue list $\lambda_2, \dots, \lambda_n$ for the induction step.
*Structural Commentary*: Isolate the chosen eigenline, then repeat on the lower-right block.
*Key Insight*: The diagonal order is controlled by which eigenline is peeled off first. □




Figure 49: Schur induction: isolate an eigenline, then triangularise the compressed block.

**Corollary 6.1.11.** *Eigenvalues of the Inverse*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ be invertible with eigenvalues $\lambda_1, \dots, \lambda_n$, counted with algebraic multiplicity. Then $A^{-1}$ has eigenvalues $\lambda_1^{-1}, \dots, \lambda_n^{-1}$, counted with algebraic multiplicity.*

*Proof.* Take a Schur form $A = UTU^*$ with diagonal $\lambda_1, \dots, \lambda_n$.
Then $A^{-1} = UT^{-1}U^*$, and $T^{-1}$ is upper triangular with diagonal $\lambda_1^{-1}, \dots, \lambda_n^{-1}$.
*Structural Commentary*: Invert the Schur form and read the diagonal.
*Key Insight*: The inverse spectrum is the reciprocal spectrum. □

**Lemma 6.1.12.** *Upper-Triangular Normal Matrices Are Diagonal*
*If $A \in \mathrm{Mat}(n, n, \mathbb{C})$ is upper triangular and normal, then $A$ is diagonal.*

*Proof.* Induct on $n$. The case $n = 1$ is immediate.
Since $A$ is upper triangular, $A\mathbf{e}_1 = a_{11}\mathbf{e}_1$, while $A^*\mathbf{e}_1 = \overline{a_{11}}\mathbf{e}_1 + \overline{a_{12}}\mathbf{e}_2 + \cdots + \overline{a_{1n}}\mathbf{e}_n$.
Normality gives $\|A\mathbf{e}_1\|_2 = \|A^*\mathbf{e}_1\|_2$, hence $a_{12} = \cdots = a_{1n} = 0$. Thus $A = a_{11} \oplus B$ with $B$ upper triangular.
The block diagonal normality equation gives $B^*B = BB^*$, so $B$ is normal. By induction, $B$ is diagonal.
*Structural Commentary*: Normality forbids upper-triangular row tails.
*Key Insight*: Compare the lengths of $A\mathbf{e}_1$ and $A^*\mathbf{e}_1$ to kill the first row, then induct. □

**Theorem 6.1.13.** *Complex Spectral Theorem*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$. The following are equivalent:*

*i. $T$ is normal;*
*ii. $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of $T$;*
*iii. $T$ has a diagonal matrix with respect to some orthonormal basis of $\mathcal{V}$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) By Theorem 6.1.9, choose an orthonormal basis $\gamma$ such that $A = [T]_{\gamma \leftarrow \gamma}$ is upper triangular. Since $T$ is normal, $A$ is normal. Lemma 6.1.12 gives that $A$ is diagonal, so $\gamma$ is an orthonormal basis of eigenvectors.
(*ii*) $\Rightarrow$ (*iii*) In an orthonormal eigenbasis, the matrix of $T$ is diagonal.
(*iii*) $\Rightarrow$ (*i*) If $[T]$ is diagonal in an orthonormal basis, then $[T^*] = [T]^*$ is diagonal as well. Diagonal matrices commute, so $TT^* = T^*T$; hence $T$ is normal.

*Structural Commentary*: Schur gives triangular form; normality kills the upper entries.
*Key Insight*: Equal norms of $T\mathbf{e}_j$ and $T^*\mathbf{e}_j$ force each row tail to vanish. □




Figure 50: Normality turns Schur triangular form diagonal by forcing row tails to vanish.

**Theorem 6.1.14.** *Real Spectral Theorem*
*Let $\mathcal{V}$ be a finite-dimensional real inner-product space and let $T \in \mathcal{L}(\mathcal{V})$. The following are equivalent:*

*i. $T$ is self-adjoint;*
*ii. $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of $T$;*
*iii. $T$ has a diagonal matrix with respect to some orthonormal basis of $\mathcal{V}$.*

*Proof.*

$(i) \Rightarrow (ii)$ Assume $T = T^*$ and induct on $n = \dim(\mathcal{V})$. For $n = 0$ there is nothing to prove, and for $n = 1$ every nonzero vector is an eigenvector after normalization.
Let $n \geq 2$. Lemma 6.1.7 gives a unit eigenvector $\mathbf{u}$, and set $\mathcal{U} = \operatorname{span}(\mathbf{u})$. The subspace $\mathcal{U}$ is $T$-invariant. By Proposition 6.1.8, $\mathcal{U}^\perp$ is $T$-invariant and the restriction $T|_{\mathcal{U}^\perp}$ is self-adjoint.
Since $\dim(\mathcal{U}^\perp) = n - 1$, the induction hypothesis gives an orthonormal eigenbasis of $\mathcal{U}^\perp$.
Adjoining $\mathbf{u}$ gives an orthonormal eigenbasis of $\mathcal{V}$ by Theorem 2.1.5.

$(ii) \Rightarrow (iii)$ In an orthonormal eigenbasis, the matrix of $T$ is diagonal.

$(iii) \Rightarrow (i)$ In an orthonormal basis where $[T]$ is real diagonal, $[T^*] = [T]^t = [T]$. Hence $T = T^*$.

*Structural Commentary*: Split off one eigendirection, then restrict to its orthogonal complement.
*Key Insight*: One eigenvector reduces the dimension. □

**Theorem 6.1.15.** *Spectral Resolution of a Normal Operator*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $\mu_1, \ldots, \mu_d$ be distinct scalars. For $T \in \mathcal{L}(\mathcal{V})$, consider:*

*(i) $T$ is normal and $\operatorname{spec}(T) \subseteq \{\mu_1, \ldots, \mu_d\}$.*
*(ii) $\exists$ pairwise orthogonal projections $P_1, \ldots, P_d \in \mathcal{L}(\mathcal{V})$ such that $P_i P_j = 0$ for $i \neq j$, $P_1 + \cdots + P_d = I_\mathcal{V}$, and $T = \mu_1 P_1 + \cdots + \mu_d P_d$*
*(iii) Any projections satisfying (ii) obey $\operatorname{range}(P_j) = E_{\mu_j}(T)$ whenever $\mu_j \in \operatorname{spec}(T)$.*

*Then (i) and (ii) are equivalent, and any projections satisfying (ii) satisfy (iii).*

*Proof.*

$(i) \Rightarrow (ii)$. By Theorem 6.1.13, $\mathcal{V} = \bigoplus_{\lambda \in \operatorname{spec}(T)} E_\lambda(T)$ as an orthogonal direct sum.
Let $P_j$ be the orthogonal projection onto $E_{\mu_j}(T)$, and set $P_j = 0$ when $\mu_j \notin \operatorname{spec}(T)$. Since $\operatorname{spec}(T) \subseteq \{\mu_1, \ldots, \mu_d\}$, these projections are pairwise orthogonal, sum to $I_\mathcal{V}$, and satisfy $T = \sum_j \mu_j P_j$.

$(ii) \Rightarrow (i)$. Since $P_j^* = P_j$ and $P_i P_j = 0$ for $i \neq j$, $T^* = \sum_{j=1}^{d} \overline{\mu_j} P_j$ and $T^* T = \sum_{j=1}^{d} |\mu_j|^2 P_j = T T^*$.
Thus $T$ is normal. If $\lambda \in \operatorname{spec}(T)$ and $\mathbf{0} \neq \mathbf{x} \in E_\lambda(T)$, then $0 = (T - \lambda I_\mathcal{V})\mathbf{x} = \sum_{j=1}^{d} (\mu_j - \lambda) P_j \mathbf{x}$.
The summands lie in the pairwise orthogonal ranges $\operatorname{range}(P_j)$, hence $0 = \sum_{j=1}^{d} |\mu_j - \lambda|^2 \|P_j \mathbf{x}\|^2$.
Because $\sum_j P_j = I_\mathcal{V}$, at least one $P_k \mathbf{x}$ is nonzero, so $\mu_k = \lambda$. Thus $\operatorname{spec}(T) \subseteq \{\mu_1, \ldots, \mu_d\}$.

$(ii) \Rightarrow (iii)$. For $\mathbf{x} \in \mathcal{V}$, $\mathbf{x} = \sum_{i=1}^{d} P_i \mathbf{x}$ and $(T - \mu_j I_\mathcal{V})\mathbf{x} = \sum_{i \neq j} (\mu_i - \mu_j) P_i \mathbf{x}$.
The summands are orthogonal, so $\|(T - \mu_j I_\mathcal{V})\mathbf{x}\|^2 = \sum_{i \neq j} |\mu_i - \mu_j|^2 \|P_i \mathbf{x}\|^2$, which vanishes exactly when $P_i \mathbf{x} = 0$ for all $i \neq j$, equivalently $\mathbf{x} = P_j \mathbf{x}$. Thus $E_{\mu_j}(T) = \operatorname{range}(P_j)$.

*Structural Commentary*: A normal operator decomposes into scalar actions on orthogonal eigenspaces.
*Key Insight*: Spectral projections isolate eigenspaces; the operator is the weighted sum of those projections. □

Figure 51: Spectral resolution: $\mathcal{V} = \bigoplus_j \operatorname{range}(P_j)$, $P_i P_j = 0$ for $i \neq j$, $\sum_j P_j = I_\mathcal{V}$, and $T = \sum_j \mu_j P_j$.

**Perspective 6.1.16.** *Spectral Resolutions Are Orthogonal Resolutions*
For the projections $P_j$ in Theorem 6.1.15, $P_j = P_j^* = P_j^2$; $P_iP_j = 0$ $(i \neq j)$; $\sum_j P_j = I_{\mathcal{V}}$.
Hence $\mathcal{V} = \bigoplus_j^{\perp} \operatorname{range}(P_j)$ is the orthogonal decomposition into $T$-invariant spectral subspaces.
*Structural Commentary*: Each $P_j$ is forced by its eigenspace: it is the orthogonal projection onto it.
*Key Insight*: Normality turns diagonalization into a projection-valued decomposition.

**Corollary 6.1.17.** *Functional Calculus from Spectral Resolution*

*Let $T = \sum_{j=1}^{d} \mu_j P_j$ be the spectral resolution of a normal operator and let $f \in \mathbb{C}[z]$. Then $f(T) = \sum_{j=1}^{d} f(\mu_j)P_j$.*

*In particular, for each $j$ there is a polynomial $p_j$ such that $p_j(T) = P_j$.*

*Proof.* Since $P_iP_j = 0$ for $i \neq j$ and $P_j^2 = P_j$, induction gives $T^m = \sum_j \mu_j^m P_j$ for every $m \geq 0$.
Linearity then gives the displayed formula for every polynomial $f$.
For the final assertion, choose the Lagrange polynomial $p_j$ satisfying $p_j(\mu_i) = \delta_{ij}$. Then $p_j(T) = P_j$.
*Structural Commentary*: Orthogonal spectral projections multiply like coordinate idempotents.
*Key Insight*: Polynomial functions of a normal operator act by scalar functions on each eigenspace. □

## 6.2 Normal Operator Consequences

**Perspective 6.2.1.** *Normal Means Diagonal in Orthogonal Coordinates*
Complex spectral theorem states that normal operators are precisely the operators whose full structure is visible in an orthonormal eigenbasis. This is why normality is the natural coordinate-free version of being diagonal.

**Proposition 6.2.2.** *Basic Spectral Properties of Normal Operators*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space, let $T \in \mathcal{L}(\mathcal{V})$ be normal, and let $p \in \mathbb{C}[z]$.*

*(i) $p(T)$ is normal and commutes with $T$.*
*(ii) $T^*$ is normal and $E_\lambda(T) = E_{\overline{\lambda}}(T^*)$ for every $\lambda \in \mathbb{C}$.*
*(iii) $\ker T = \ker T^*$ and $\operatorname{range}(T) = \operatorname{range}(T^*)$.*

*Proof.* Choose an orthonormal eigenbasis $\mathbf{e}_1, \ldots, \mathbf{e}_n$ with $T\mathbf{e}_j = \mu_j \mathbf{e}_j$.

(i) $p(T)\mathbf{e}_j = p(\mu_j)\mathbf{e}_j$, $p(T)$ is diagonal in this basis, hence normal. It commutes with $T$ by Proposition 1.1.5.
(ii) Since $T^*\mathbf{e}_j = \overline{\mu_j}\mathbf{e}_j$, $T^*$ is diagonal in the same basis, hence normal. Also $T\mathbf{v} = \lambda\mathbf{v}$ iff $T^*\mathbf{v} = \overline{\lambda}\mathbf{v}$.
(iii) Part (ii) with $\lambda = 0$ gives $\ker T = \ker T^*$. Both ranges are $\operatorname{span}\{\mathbf{e}_j : \mu_j \neq 0\}$.

*Structural Commentary*: Normality gives one orthonormal basis that diagonalises both $T$ and $T^*$.
*Key Insight*: Kernel and range are read from the nonzero diagonal entries in that basis. □

**Lemma 6.2.3.** *Adjoints of Normal Matrices as One Polynomial*
*Let $A_1, \ldots, A_k$ be normal complex matrices. Then $\exists$ a polynomial $p \in \mathbb{C}[z]$ s.t. $p(A_j) = A_j^*$ $\forall j = 1, \ldots, k$.*

*Proof.* Let $\Omega = \bigcup_{j=1}^{k} \operatorname{spec}(A_j)$. Since $\Omega$ is finite, choose a Lagrange interpolation polynomial $p$ satisfying $p(\lambda) = \overline{\lambda}$ for every $\lambda \in \Omega$. For each $A_j$, Corollary 6.1.17 gives $p(A_j) = A_j^*$.
*Structural Commentary*: One polynomial interpolates conjugation on all spectra.
*Key Insight*: For normal matrices, $A^*$ is $p(A)$ with $p(\lambda) = \overline{\lambda}$. □

**Theorem 6.2.4.** *Fuglede–Putnam Theorem*
*Let $A \in \operatorname{Mat}(m, m, \mathbb{C})$ and $B \in \operatorname{Mat}(n, n, \mathbb{C})$ be normal, and let $X \in \operatorname{Mat}(m, n, \mathbb{C})$.*
*Then $AX = XB \iff A^*X = XB^*$.*

*Proof.*

($\Rightarrow$) Assume $AX = XB$. Then $A^rX = XB^r$ for every $r \geq 0$, so $p(A)X = Xp(B)$ for every polynomial $p$. By Lemma 6.2.3, choose $p$ with $p(A) = A^*$ and $p(B) = B^*$. Thus $A^*X = XB^*$.
($\Leftarrow$) Assume $A^*X = XB^*$. Apply the same argument to the normal matrices $A^*$ and $B^*$, choosing $q$ with $q(A^*) = A$ and $q(B^*) = B$. Hence $AX = XB$.

*Structural Commentary*: An intertwining relation passes through every polynomial in the two normal matrices.
*Key Insight*: Normality makes adjoints polynomial, so intertwining transfers to adjoints. □

**Method 6.2.5.** *Finding a Spectral Decomposition of a Normal Matrix*
For a normal matrix $A \in \operatorname{Mat}(n, n, \mathbb{C})$:

i. Find the distinct eigenvalues $\mu_1, \ldots, \mu_d$.
ii. For each $j$, solve $(A - \mu_j I_n)\mathbf{x} = 0$ to get $E_{\mu_j}(A)$.

iii. Orthonormalise a basis of each eigenspace.
iv. Concatenate these orthonormal bases as the columns of $U$; then $U$ is unitary and $A = U\Lambda U^*$, where $\Lambda$ repeats each $\mu_j$ according to $\dim E_{\mu_j}(A)$.

**Theorem 6.2.6.** *Spectral Classification of Normal Operators*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$ be normal.*

*(i) $T$ is unitary if and only if $|\lambda| = 1$ for every $\lambda \in \operatorname{spec}(T)$.*
*(ii) $T$ is self-adjoint if and only if $\operatorname{spec}(T) \subseteq \mathbb{R}$.*
*(iii) $T$ is an orthogonal projection if and only if $\operatorname{spec}(T) \subseteq \{0, 1\}$.*
*(iv) $T$ is skew-adjoint, meaning $T^* = -T$, if and only if every $\lambda \in \operatorname{spec}(T)$ is purely imaginary.*

*Proof.* Write the spectral resolution $T = \sum_j \mu_j P_j$, where the $\mu_j$ are the distinct eigenvalues and $P_j \neq 0$.
Then $I_{\mathcal{V}} = \sum_j P_j$, $T^* = \sum_j \overline{\mu_j} P_j$, $T^*T = \sum_j |\mu_j|^2 P_j$, and $T^2 = \sum_j \mu_j^2 P_j$.
Since ranges of $P_j$ are nonzero and independent, an identity $\sum_j \alpha_j P_j = 0$ holds only when every $\alpha_j = 0$.
(i) $T$ is unitary iff $T^*T = I_{\mathcal{V}} = \sum_j P_j$, iff $|\mu_j| = 1$ for every $j$.
(ii) $T$ is self-adjoint iff $T = T^*$, iff $\mu_j = \overline{\mu_j}$ for every $j$.
(iii) $T$ is an orthogonal projection iff $T = T^*$ and $T^2 = T$, iff $\mu_j = \overline{\mu_j}$ and $\mu_j^2 = \mu_j$ $\forall j$, iff $\mu_j \in \{0, 1\}$ $\forall j$.
(iv) $T^* = -T$ iff $\overline{\mu_j} = -\mu_j$ for every $j$, iff every $\mu_j$ is purely imaginary.
*Structural Commentary*: For normal operators, structural classes are scalar conditions on spectral blocks.
*Key Insight*: For normal $T$, these properties reduce to scalar tests. □

**Proposition 6.2.7.** *Permutation Similarity Reorders Diagonal Entries*
*Let $\Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_n)$ and let $P = [\mathbf{e}_{\sigma(1)} \ \cdots \ \mathbf{e}_{\sigma(n)}]$ be a permutation matrix. Then $P^*\Lambda P = \operatorname{diag}(\lambda_{\sigma(1)}, \ldots, \lambda_{\sigma(n)})$.*

*Proof.* The $(i, j)$ entry of $P^*\Lambda P$ is $\mathbf{e}_{\sigma(i)}^* \Lambda \mathbf{e}_{\sigma(j)} = \lambda_{\sigma(j)} \mathbf{e}_{\sigma(i)}^* \mathbf{e}_{\sigma(j)} = \lambda_{\sigma(j)} \delta_{ij}$.
Thus all off-diagonal entries are zero, and the $j$th diagonal entry is $\lambda_{\sigma(j)}$.
*Structural Commentary*: Permutation similarity applies the same basis reordering to rows and columns.
*Key Insight*: Diagonal entries move exactly as the standard basis vectors are rearranged. □

**Corollary 6.2.8.** *Normal Matrices with the Same Spectrum Are Unitarily Similar*
*If $A, B \in \operatorname{Mat}(n, n, \mathbb{C})$ are normal and have the same eigenvalues counted with algebraic multiplicity, then $A$ and $B$ are unitarily similar.*

*Proof.* By Theorem 6.1.13, write $A = U\Lambda U^*$ and $B = V\Gamma V^*$ with $U, V$ unitary and $\Lambda, \Gamma$ diagonal.
Since eigenvalues agree with multiplicity, Proposition 6.2.7 gives a permutation matrix $P$ such that $\Gamma = P^*\Lambda P$.
Hence $B = (VP^*U^*)A(UPV^*)$, and $UPV^*$ is unitary.
*Structural Commentary*: Normal matrices are classified up to unitary similarity by their eigenvalue multiset.
*Key Insight*: After diagonalising both matrices, only a permutation of diagonal entries remains. □

**Proposition 6.2.9.** *Cartesian Decomposition and Normality*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space.*
*For every $T \in \mathcal{L}(\mathcal{V})$, there are unique self-adjoint operators $H, K \in \mathcal{L}(\mathcal{V})$ such that $T = H + iK$.*
*They are $H = \frac{1}{2}(T + T^*)$ and $K = \frac{1}{2i}(T - T^*)$. Moreover, $T$ is normal if and only if $HK = KH$.*

*Proof.* The displayed formulas give self-adjoint $H, K$ and $T = H + iK$. If $T = X + iY$ with $X, Y$ self-adjoint, then $T^* = X - iY$, so $X = \frac{1}{2}(T + T^*)$ and $Y = \frac{1}{2i}(T - T^*)$; uniqueness follows.
For $T = H + iK$, $T^*T = (H - iK)(H + iK) = H^2 + K^2 + i(HK - KH)$,
whereas $TT^* = (H + iK)(H - iK) = H^2 + K^2 - i(HK - KH)$. Thus $T^*T = TT^*$ if and only if $HK = KH$.
*Structural Commentary*: Every complex operator splits into Hermitian and skew-Hermitian parts.
*Key Insight*: Normality says the two Hermitian coordinates of $T$ commute. □




Figure 52: Cartesian decomposition: $T$ is normal exactly when $HK = KH$.

**Proposition 6.2.10.** *Normal Block Upper-Triangular Matrices Collapse*
*Let $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. Let $A = \begin{bmatrix} B & X \\ 0 & C \end{bmatrix} \in \mathrm{Mat}(n, n, \mathbb{F})$, where $B$ and $C$ are square.*
*Then $A$ is normal if and only if $X = 0$ and both $B$ and $C$ are normal.*

*Proof.*
($\Rightarrow$) Suppose $A$ is normal. Comparing the $(1,1)$ blocks of $A^*A$ and $AA^*$ gives $B^*B = BB^* + XX^*$. Taking traces and using cyclicity, $\mathrm{tr}(XX^*) = \mathrm{tr}(B^*B) - \mathrm{tr}(BB^*) = 0$. Since $\mathrm{tr}(XX^*) = \|X\|_F^2$, we get $X = 0$. Thus $A = B \oplus C$, and $A^*A = AA^*$ reduces blockwise to $B^*B = BB^*$ and $C^*C = CC^*$.
($\Leftarrow$) If $X = 0$ and $B, C$ are normal, then $A = B \oplus C$, so blockwise multiplication gives $A^*A = AA^*$.
*Structural Commentary*: A normal operator cannot have one-way coupling between orthogonal blocks.
*Key Insight*: The off-diagonal block contributes the nonnegative trace $\|X\|_F^2$, so normality forces it to vanish. □




Figure 53: Normality removes the upper-right block in a block upper-triangular matrix.

**Theorem 6.2.11.** *Invariant Subspaces of Normal Matrices Reduce*
*Let $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ be normal, and let $\mathcal{U} \subseteq \mathbb{F}^n$ be invariant under $A$. Then $\mathcal{U}$ is invariant under $A^*$. If $P$ is the orthogonal projection onto $\mathcal{U}$, then $AP = PA$.*

*Proof.* The cases $\mathcal{U} = \{0\}$ and $\mathcal{U} = \mathbb{F}^n$ are immediate. Otherwise choose a unitary $Q = [Q_1\ Q_2]$ whose columns form orthonormal bases of $\mathcal{U}$ and $\mathcal{U}^\perp$. Invariance of $\mathcal{U}$ says $AQ_1$ has columns in $\mathcal{U} = \mathrm{col}(Q_1)$, hence $Q_2^*AQ_1 = 0$. Therefore $Q^*AQ = \begin{bmatrix} B & X \\ 0 & C \end{bmatrix}$. This matrix is normal because it is unitarily similar to $A$.
Proposition 6.2.10 gives $X = 0$, so $Q^*AQ = B \oplus C$. Hence $Q^*A^*Q = B^* \oplus C^*$, which shows $\mathcal{U}$ is $A^*$-invariant. Also $Q^*PQ = I_{\dim(\mathcal{U})} \oplus 0$, so $Q^*AQ$ commutes with $Q^*PQ$; conjugating back gives $AP = PA$.
*Structural Commentary*: In adapted coordinates, normal invariance becomes block diagonality.
*Key Insight*: Normal invariant subspaces are automatically reducing subspaces. □

**Corollary 6.2.12.** *Invariant Subspaces of Normal Operators Reduce*
*Let $\mathcal{V}$ be a finite-dimensional inner-product space, let $T \in \mathcal{L}(\mathcal{V})$ be normal, let $\mathcal{U}$ be invariant under $T$. Then:*
*(i) $\mathcal{U}^\perp$ is invariant under $T$;*
*(ii) $\mathcal{U}$ is invariant under $T^*$;*
*(iii) $(T|_{\mathcal{U}})^* = (T^*)|_{\mathcal{U}}$;*
*(iv) $T|_{\mathcal{U}}$ and $T|_{\mathcal{U}^\perp}$ are normal.*

*Proof.* Choose an orthonormal basis adapted to $\mathcal{U} \oplus \mathcal{U}^\perp$.
In these coordinates, Theorem 6.2.11 makes the matrix of $T$ block diagonal, say $A = B \oplus C$.
The block diagonal form shows $\mathcal{U}^\perp$ is invariant under $T$ and $\mathcal{U}$ is invariant under $T^*$. It also gives $(T|_{\mathcal{U}})^* = (T^*)|_{\mathcal{U}}$ because the adjoint block on $\mathcal{U}$ is $B^*$. Finally, normality of $A$ reduces blockwise to normality of $B$ and $C$.
*Structural Commentary*: Normal operator turns every invariant subspace into an orthogonal reducing summand.
*Key Insight*: Normality kills the one-way coupling between $\mathcal{U}$ and $\mathcal{U}^\perp$. □

**Perspective 6.2.13.** *Real Normal Operators Use Planes*
Over $\mathbb{R}$, normal operators need not have enough eigenvectors. The replacement for a missing conjugate eigenpair is a two-dimensional invariant plane carrying a rotation-dilation block.

**Proposition 6.2.14.** *Two-Dimensional Real Normal Blocks*
*Let $\dim(\mathcal{V}) = 2$ over $\mathbb{R}$ and let $T \in \mathcal{L}(\mathcal{V})$.*
*Then $T$ is normal but not self-adjoint if and only if, in every orthonormal basis, $[T] = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, $b \neq 0$.*
*Moreover, some orthonormal basis makes $b > 0$.*

*Proof.*
($\Rightarrow$) In an orthonormal basis, write $A = [T] = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$.
Normality gives $A^tA = AA^t$. Comparing the $(1,1)$ and $(2,2)$ entries gives $b^2 = c^2$. If $c = b$, then $A = A^t$,

impossible because $T$ is not self-adjoint. Thus $c = -b$ and $b \neq 0$. Comparing off-diagonal entries gives $-ab + bd = ab - bd$, so $d = a$. Since the original orthonormal basis was arbitrary, every orthonormal basis has the displayed form.

($\Leftarrow$) A matrix $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ with $b \neq 0$ satisfies $A^t A = AA^t$ and is not symmetric. Hence $T$ is normal but not self-adjoint. If $b < 0$, replace the second basis vector by its negative to make $b > 0$.

*Structural Commentary*: In real dimension 2, non-self-adjoint normality is exactly a scalar plus a rotation block.
*Key Insight*: Normality forces $c = -b$ and $d = a$. □

**Theorem 6.2.15.** *Real Normal Operator Block Form*
*Let $\mathcal{V}$ be finite-dimensional over $\mathbb{R}$ and let $T \in \mathcal{L}(\mathcal{V})$. Then $T$ is normal if and only if some orthonormal basis makes $[T]$ block diagonal, with each diagonal block either 1-by-1 or of the form $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, $b > 0$.*

*Proof.*

($\Rightarrow$) Induct on $n = \dim(\mathcal{V})$. The cases $n \leq 1$ are immediate.
For $n \geq 2$, Theorem 5.3.13 gives an invariant subspace $\mathcal{U}$ with $\dim(\mathcal{U}) \in \{1, 2\}$.
By Corollary 6.2.12, $\mathcal{U}^\perp$ is invariant under $T$ and both restrictions are normal.
If $\dim(\mathcal{U}) = 1$, take a unit vector in $\mathcal{U}$. If $\dim(\mathcal{U}) = 2$, use the Real Spectral Theorem 6.1.14 when $T|_{\mathcal{U}}$ is self-adjoint, giving scalar 1-by-1 blocks on $\mathcal{U}$, and use Proposition 6.2.14 otherwise.
By induction, choose the required orthonormal block basis on $\mathcal{U}^\perp$. Concatenating it with the basis chosen on $\mathcal{U}$ gives an orthonormal basis of $\mathcal{V}$ because $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$.

($\Leftarrow$) Each allowed block is normal. If $[T]$ is the block diagonal sum of these blocks, then $[T]^*[T]$ and $[T][T]^*$ are the corresponding block diagonal sums of $B^*B$ and $BB^*$, hence are equal block by block.

*Structural Commentary*: Normality lets invariant lines or planes split orthogonally.
*Key Insight*: Real normal operators split into scalar lines and rotation-dilation planes. □




Figure 54: Orthogonal line/plane split into scalar and rotation blocks.

**Theorem 6.2.16.** *Simultaneous Unitary Diagonalisation of Commuting Normal Operators*
*Let $\emptyset \neq \mathcal{F} \subseteq \mathcal{L}(\mathcal{V})$ be a normal family on a finite-dimensional complex inner-product space. Then $\mathcal{F}$ is pairwise commuting if and only if $\mathcal{V}$ has an orthonormal basis of common eigenvectors for $\mathcal{F}$.*

*Proof.*

($\Leftarrow$) In a common orthonormal eigenbasis, every $A \in \mathcal{F}$ is diagonal. Hence all operators in $\mathcal{F}$ commute.

($\Rightarrow$) Induct on $\dim(\mathcal{V})$. The zero-dimensional case is vacuous. If every $A \in \mathcal{F}$ is scalar, any orthonormal basis works. Otherwise choose a nonscalar $A_0 \in \mathcal{F}$ and write its spectral resolution as $A_0 = \sum_j \mu_j P_j$.
Since $A_0$ is nonscalar, at least two nonzero ranges $\operatorname{range}(P_j)$ occur, and each has dimension strictly smaller than $\dim(\mathcal{V})$. By Corollary 6.1.17, each $P_j$ is a polynomial in $A_0$.
If $B \in \mathcal{F}$, then $BA_0 = A_0B$, so $B$ commutes with every polynomial in $A_0$, in particular with $P_j$. Hence $B(\operatorname{range}(P_j)) \subseteq \operatorname{range}(P_j)$ because $P_jB\mathbf{x} = BP_j\mathbf{x} = B\mathbf{x}$ whenever $P_j\mathbf{x} = \mathbf{x}$.
The same commutation preserves $\ker(P_j) = \operatorname{range}(P_j)^\perp$.
In the decomposition $\mathcal{V} = \operatorname{range}(P_j) \oplus \ker(P_j)$, the matrix of $B$ is therefore block diagonal; since $B$ is normal, its diagonal restrictions are normal block by block.
Thus, on each nonzero $\operatorname{range}(P_j)$, the restricted family is again pairwise commuting and normal.
By induction it has a common orthonormal eigenbasis. Concatenating these bases over all $j$ gives a common orthonormal eigenbasis of $\mathcal{V}$.

*Structural Commentary*: Spectral projections reduce the whole commuting family.
*Key Insight*: Diagonalise one nonscalar member, then induct on its eigenspaces. □




Figure 55: For $A_0 = \sum_j \mu_j P_j$, commutativity gives $BP_j = P_jB$; the ranges reduce $\mathcal{F}$, and induction gives a common orthonormal eigenbasis.

**Corollary 6.2.17.** *Sums and Products of Commuting Normal Operators*
*Let $S, T \in \mathcal{L}(\mathcal{V})$ be commuting normal operators on a finite-dimensional complex inner-product space. Then $S+T$ and $ST$ are normal.*

*Proof.* By Theorem 6.2.16, $S$ and $T$ are diagonal in a common orthonormal basis.
Then $S+T$ and $ST$ are also diagonal in that basis, hence normal.
*Structural Commentary*: Commuting normal operators share orthonormal coordinates.
*Key Insight*: In common diagonal form, sums and products remain diagonal. □

**Theorem 6.2.18.** *Description of Complex Isometries*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space and let $S \in \mathcal{L}(\mathcal{V})$. Then $S$ is an isometry if and only if $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of $S$ whose eigenvalues all have absolute value 1.*

*Proof.* If $S\mathbf{e}_j = \lambda_j \mathbf{e}_j$ in an orthonormal basis with $|\lambda_j| = 1$, then for $\mathbf{v} = \sum_j c_j \mathbf{e}_j$,
$\|S\mathbf{v}\|^2 = \sum_j |\lambda_j c_j|^2 = \sum_j |c_j|^2 = \|\mathbf{v}\|^2$, so $S$ is an isometry. Conversely, if $S$ is an isometry, then $S^*S = I_{\mathcal{V}}$ by Theorem 2.6.6. Finite-dimensional invertibility gives $SS^* = I_{\mathcal{V}}$, so $S$ is normal. By Complex Spectral Theorem, choose orthonormal eigenbasis $S\mathbf{e}_j = \lambda_j \mathbf{e}_j$. Since $\mathbf{e}_j$ is unit and $S$ preserves norms, $|\lambda_j| = \|S\mathbf{e}_j\| = 1$.
*Structural Commentary*: Complex isometries are normal with unit-circle spectrum.
*Key Insight*: In an eigenbasis, isometry means $|\lambda_j| = 1$. □

**Example 6.2.19.** *Plane Rotations*
For $\theta \in \mathbb{R}$, the real matrix $R_\theta = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$ is an isometry of $\mathbb{R}^2$. If $\theta$ is not an integer multiple of $\pi$, then $R_\theta$ has no real eigenvalue, which is why real isometries require two-dimensional rotation blocks.

**Theorem 6.2.20.** *Real Isometry Block Form*
*Let $\mathcal{V}$ be finite-dimensional over $\mathbb{R}$ and let $S \in \mathcal{L}(\mathcal{V})$. Then $S$ is an isometry if and only if some orthonormal basis makes $[S]$ block diagonal, with each diagonal block one of $[1]$, $[-1]$, $\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$ with $0 < \theta < \pi$.*

*Proof.*
($\Rightarrow$) Since $S$ is an isometry, $S^*S = I_{\mathcal{V}}$ by Theorem 2.6.6. Finite-dimensional invertibility gives $SS^* = I_{\mathcal{V}}$, so $S$ is normal. Apply Theorem 6.2.15. A 1-by-1 block $[\lambda]$ has $\lambda^2 = 1$, so $\lambda = \pm 1$.
A 2-by-2 block $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ has $a^2 + b^2 = 1$ and $b > 0$, so $a = \cos\theta$ and $b = \sin\theta$ for a unique $\theta \in (0, \pi)$.
($\Leftarrow$) Each displayed block $B$ satisfies $B^tB = I$ on its coordinate subspace. Therefore the block diagonal matrix satisfies $[S]^t[S] = I$, so $S$ preserves norms and is an isometry.
*Structural Commentary*: Isometries are the unit-length special case of real normal blocks.
*Key Insight*: Scalar blocks reduce to $\pm 1$, and planar normal blocks reduce to rotations. □




Figure 56: Normal form: fixed/flipped lines and rotation planes.

## 6.3 Operator Functions

This subsection develops finite spectral functions of normal operators, positive square roots, commutativity through spectral projections, and matrix exponentials for diagonalizable matrices.

**Definition 6.3.1.** *Spectral Resolution Notation*
Let $T \in \mathcal{L}(\mathcal{V})$ be normal on a finite-dimensional complex inner-product space.
Define $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$, where $\Omega_T = \operatorname{spec}(T)$.

(i) *Spectral Set*: $\Omega_T = \operatorname{spec}(T)$.
(ii) *Spectral Projection*: $P_\lambda$ is the orthogonal projection onto $E_\lambda(T)$.
(iii) *Spectral Resolution*: $\{P_\lambda\}_{\lambda \in \Omega_T}$, equivalently $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$.

**Theorem 6.3.2.** *Finite Spectral Functional Calculus for Normal Operators*
*Let $T \in \mathcal{L}(\mathcal{V})$ be normal, with spectral resolution $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$.*
*The finite spectral functional calculus is $\mathbb{C}^{\Omega_T} \to \mathcal{L}(\mathcal{V})$, where $f \mapsto f(T) := \sum_{\lambda \in \Omega_T} f(\lambda) P_\lambda$.*
*If $p \in \mathbb{C}[z]$ and $p|_{\Omega_T} = f$, then $p(T) = f(T)$. Hence every $f \in \mathbb{C}^{\Omega_T}$ has a polynomial representative in $T$.*

*Proof.* For $p|_{\Omega_T} = f$, Corollary 6.1.17 gives $p(T) = \sum_{\lambda \in \Omega_T} p(\lambda) P_\lambda = \sum_{\lambda \in \Omega_T} f(\lambda) P_\lambda = f(T)$.
Since $\Omega_T$ is finite, Lagrange interpolation gives $p \in \mathbb{C}[z]$ with $p|_{\Omega_T} = f$.
*Structural Commentary*: Spectral sums reduce finite functions to polynomial evaluation.
*Key Insight*: Finite spectral data is interpolated by one polynomial. □

**Theorem 6.3.3.** *Rules for Normal Spectral Functions*
*Let $T \in \mathcal{L}(\mathcal{V})$ be normal, with spectral resolution $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$. For $f, g : \Omega_T \to \mathbb{C}$:*

*(i)* $(f+g)(T) = f(T) + g(T)$ *and* $(fg)(T) = f(T)g(T)$;
*(ii)* $f(T)^* = \overline{f}(T)$, *where* $\overline{f}(\lambda) = \overline{f(\lambda)}$;
*(iii)* $f(T)$ *is normal and* $\operatorname{spec}(f(T)) = f(\Omega_T)$.

*Proof.*

(i) Since $P_\lambda P_\mu = \delta_{\lambda\mu} P_\lambda$, $f(T)g(T) = \sum_{\lambda \in \Omega_T} f(\lambda) g(\lambda) P_\lambda = (fg)(T)$; and $(f+g)(T) = f(T) + g(T)$.

(ii) Since $P_\lambda^* = P_\lambda$, $f(T)^* = \sum_{\lambda \in \Omega_T} \overline{f(\lambda)} P_\lambda = \overline{f}(T)$.

(iii) For each $\lambda \in \Omega_T$, $f(T)|_{\operatorname{range}(P_\lambda)} = f(\lambda) I$.
Thus $f(T)$ is diagonal in an orthonormal eigenbasis, hence normal, and $\operatorname{spec}(f(T)) = f(\Omega_T)$.

*Structural Commentary*: Spectral projections make operator rules pointwise.
*Key Insight*: Spectral functions act as scalars on spectral subspaces. □

**Definition 6.3.4.** *Positive Operator*
$T \in \mathcal{L}(\mathcal{V})$ is *positive* if $T = T^*$ and $\langle T\mathbf{v}, \mathbf{v} \rangle \geq 0$ for all $\mathbf{v} \in \mathcal{V}$.

**Definition 6.3.5.** *Square Root of an Operator*
An operator $R \in \mathcal{L}(\mathcal{V})$ is a *square root* of $T \in \mathcal{L}(\mathcal{V})$ if $R^2 = T$.

**Theorem 6.3.6.** *Characterization of Positive Operators*
*For $T \in \mathcal{L}(\mathcal{V})$, the following are equivalent:*

*i.* $T$ *is positive;*
*ii.* $T$ *is self-adjoint and every eigenvalue of $T$ is nonnegative;*
*iii.* $T$ *has a positive square root;*
*iv.* $T$ *has a self-adjoint square root;*
*v. there exists $R \in \mathcal{L}(\mathcal{V})$ such that $T = R^*R$.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* If $T\mathbf{v} = \lambda \mathbf{v}$ and $\mathbf{v} \neq \mathbf{0}$, then $\lambda \in \mathbb{R}$ by Theorem 6.1.5, and $0 \leq \langle T\mathbf{v}, \mathbf{v} \rangle = \lambda \|\mathbf{v}\|^2$, so $\lambda \geq 0$.
*(ii)* $\Rightarrow$ *(iii)* By the real or complex spectral theorem, choose an orthonormal eigenbasis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ with $T\mathbf{e}_j = \lambda_j \mathbf{e}_j$ and $\lambda_j \geq 0$. Define $S\mathbf{e}_j = \sqrt{\lambda_j}\mathbf{e}_j$ and extend linearly.
Then $S$ is diagonal with nonnegative real diagonal entries in an orthonormal basis, hence self-adjoint and positive, and $S^2\mathbf{e}_j = \lambda_j \mathbf{e}_j = T\mathbf{e}_j$ for every $j$. Thus $S^2 = T$.
*(iii)* $\Rightarrow$ *(iv)* A positive square root is self-adjoint.
*(iv)* $\Rightarrow$ *(v)* If $T = R^2$ with $R = R^*$, then $T = R^*R$.
*(v)* $\Rightarrow$ *(i)* If $T = R^*R$, then $T^* = T$ and $\langle T\mathbf{v}, \mathbf{v} \rangle = \langle R\mathbf{v}, R\mathbf{v} \rangle = \|R\mathbf{v}\|^2 \geq 0$ for every $\mathbf{v}$, so $T$ is positive.

*Structural Commentary*: Positivity equals nonnegative spectrum and square-root factorization.
*Key Insight*: In an eigenbasis, the proof is scalar: $\lambda \geq 0$ iff $\sqrt{\lambda}$ exists. □

**Proposition 6.3.7.** *Unique Positive Square Root*
*Every positive operator on $\mathcal{V}$ has a unique positive square root.*

*Proof.* Existence follows from Theorem 6.3.6. For uniqueness, decompose $\mathcal{V} = E_{\lambda_1}(T) \oplus \cdots \oplus E_{\lambda_m}(T)$.
Let $R$ be a positive square root of $T$. Since $R^2 = T$, we have $RT = R^3 = TR$, so each $E_{\lambda_j}(T)$ is invariant under $R$. On $E_{\lambda_j}(T)$, the restriction $R_j := R|_{E_{\lambda_j}(T)}$ is positive and satisfies $R_j^2 = \lambda_j I$.
By the spectral theorem for the positive operator $R_j$, every eigenvalue $\rho$ of $R_j$ satisfies $\rho \geq 0$ and $\rho^2 = \lambda_j$,

hence $\rho = \sqrt{\lambda_j}$. Therefore $R_j = \sqrt{\lambda_j} I$ on $E_{\lambda_j}(T)$.
Thus every positive square root agrees with the same scalar operator on each eigenspace $E_{\lambda_j}(T)$, so it is unique.
*Structural Commentary*: Each eigenspace determines its own scalar root.
*Key Insight*: Positivity selects the nonnegative root. □

**Definition 6.3.8.** *Positive Square Root Notation*
If $T$ is positive, then $\sqrt{T}$ denotes the unique positive square root of $T$.

**Example 6.3.9.** *Basic Functions from the Spectral Calculus*
Let $T \in \mathcal{L}(\mathcal{V})$ be normal, with spectral resolution $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$. Then $T^* = \sum_{\lambda \in \Omega_T} \overline{\lambda} P_\lambda$.
If $0 \notin \Omega_T$, then $T^{-1} = \sum_{\lambda \in \Omega_T} \lambda^{-1} P_\lambda$. If $T \geq 0$, then $T^{1/2} = \sum_{\lambda \in \Omega_T} \lambda^{1/2} P_\lambda$, where $\lambda^{1/2} \geq 0$.

**Theorem 6.3.10.** *Commutativity via Spectral Projections*
*Let $T, S \in \mathcal{L}(\mathcal{V})$ be normal, with spectral resolutions $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$ and $S = \sum_{\mu \in \Omega_S} \mu Q_\mu$.*

*(i) For $R \in \mathcal{L}(\mathcal{V})$, $RT = TR$ if and only if $RP_\lambda = P_\lambda R$ for all $\lambda \in \Omega_T$.*
*(ii) $TS = ST$ if and only if $P_\lambda Q_\mu = Q_\mu P_\lambda$ for all $\lambda \in \Omega_T$ and $\mu \in \Omega_S$.*
*(iii) If $f : \Omega_T \to \mathbb{C}$ and $g : \Omega_S \to \mathbb{C}$ are injective, then $f(T)g(S) = g(S)f(T)$ if and only if $TS = ST$.*

*Proof.*

(i). If $RT = TR$, then $Rp(T) = p(T)R$ for all $p \in \mathbb{C}[z]$. Since $P_\lambda \in \mathbb{C}[T]$ by Corollary 6.1.17, $RP_\lambda = P_\lambda R$. Conversely, if $RP_\lambda = P_\lambda R$ for all $\lambda \in \Omega_T$, then $RT = \sum_{\lambda \in \Omega_T} \lambda R P_\lambda = \sum_{\lambda \in \Omega_T} \lambda P_\lambda R = TR$.

(ii). If $TS = ST$, then (i) with $R = S$ gives $SP_\lambda = P_\lambda S$. Since $Q_\mu \in \mathbb{C}[S]$, $P_\lambda Q_\mu = Q_\mu P_\lambda$ for all $\lambda, \mu$. Conversely, if $P_\lambda Q_\mu = Q_\mu P_\lambda$ for all $\lambda, \mu$, then $TS = \sum_{\lambda,\mu} \lambda\mu P_\lambda Q_\mu = \sum_{\lambda,\mu} \lambda\mu Q_\mu P_\lambda = ST$

(iii). Suppose $f$ and $g$ are injective. If $f(T)g(S) = g(S)f(T)$, choose $h : f(\Omega_T) \to \Omega_T$ and $k : g(\Omega_S) \to \Omega_S$ with $h \circ f = \mathrm{id}_{\Omega_T}$ and $k \circ g = \mathrm{id}_{\Omega_S}$. By Theorem 6.3.2, $T = h(f(T))$ and $S = k(g(S))$, so $TS = ST$. Conversely, if $TS = ST$, then (ii) gives $P_\lambda Q_\mu = Q_\mu P_\lambda$,
and $f(T)g(S) = \sum_{\lambda,\mu} f(\lambda)g(\mu) P_\lambda Q_\mu = \sum_{\lambda,\mu} f(\lambda)g(\mu) Q_\mu P_\lambda = g(S)f(T)$.

*Structural Commentary*: Spectral projections are the coordinate idempotents of a normal operator.
*Key Insight*: To commute with a normal operator is to preserve each of its eigenspaces. □

**Corollary 6.3.11.** *A Commuting Normal Pair Comes from One Normal Operator*
*Let $T, S \in \mathcal{L}(\mathcal{V})$ be normal on a finite-dimensional complex inner-product space.*
*Then $TS = ST \iff \exists\, H \in \mathcal{L}(\mathcal{V})$ normal, $p, q \in \mathbb{C}[z]$ such that $T = p(H)$, $S = q(H)$.*
*If $TS = ST$, then $H$ may be chosen in $\mathbb{C}[T, S]$.*

*Proof.*

($\Rightarrow$) Suppose $TS = ST$. Write $T = \sum_{\lambda \in \Omega_T} \lambda P_\lambda$ and $S = \sum_{\mu \in \Omega_S} \mu Q_\mu$.
By Theorem 6.3.10, $P_\lambda Q_\mu = Q_\mu P_\lambda$. Let $\mathcal{I} = \{(\lambda, \mu) \in \Omega_T \times \Omega_S : P_\lambda Q_\mu \neq 0\}$.
The projections $P_\lambda Q_\mu$, $(\lambda, \mu) \in \mathcal{I}$, are pairwise orthogonal and sum to $I_\mathcal{V}$.
Choose distinct scalars $\alpha_{\lambda\mu}$ for $(\lambda, \mu) \in \mathcal{I}$ and set $H = \sum_{(\lambda,\mu) \in \mathcal{I}} \alpha_{\lambda\mu} P_\lambda Q_\mu$.
Then $H$ is normal, and $H \in \mathbb{C}[T, S]$ since $P_\lambda \in \mathbb{C}[T]$ and $Q_\mu \in \mathbb{C}[S]$.
Choose $p, q \in \mathbb{C}[z]$ with $p(\alpha_{\lambda\mu}) = \lambda$ and $q(\alpha_{\lambda\mu}) = \mu$ on $\mathcal{I}$.
Hence $p(H) = \sum_{(\lambda,\mu) \in \mathcal{I}} \lambda P_\lambda Q_\mu = T$, and $q(H) = \sum_{(\lambda,\mu) \in \mathcal{I}} \mu P_\lambda Q_\mu = S$.

($\Leftarrow$) If $T = p(H)$ and $S = q(H)$, then $TS = p(H)q(H) = q(H)p(H) = ST$.

*Structural Commentary*: One scalar labels each common spectral block.
*Key Insight*: Joint spectral projections reduce a pair of commuting normal operators to finite scalar data. □

**Definition 6.3.12.** *Matrix Exponential*
For $A \in \mathrm{Mat}(n, n, \mathbb{C})$ and $t \in \mathbb{C}$, define $e^{tA} := \sum_{k=0}^{\infty} \frac{t^k A^k}{k!}$.

**Proposition 6.3.13.** *Pure Modes for Powers and Exponentials*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ be diagonalizable with $A = S\Lambda S^{-1}$, where $\Lambda = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$ and the columns of $S$ are $\mathbf{v}_1, \ldots, \mathbf{v}_n$. If $\mathbf{x}_0 = \sum_{j=1}^{n} c_j \mathbf{v}_j$, then $A^k \mathbf{x}_0 = \sum_{j=1}^{n} c_j \lambda_j^k \mathbf{v}_j$ for $k \geq 0$; and $e^{tA}\mathbf{x}_0 = \sum_{j=1}^{n} c_j e^{\lambda_j t} \mathbf{v}_j$ for $t \in \mathbb{C}$.*

*Proof.* Let $c = (c_1, \dots, c_n)^t$, so $\mathbf{x}_0 = Sc$.

Then $A^k = S\Lambda^k S^{-1}$, and Definition 6.3.12 gives $e^{tA} = S\left(\sum_{r=0}^{\infty} \frac{t^r \Lambda^r}{r!}\right) S^{-1} = S\,\mathrm{diag}(e^{\lambda_1 t}, \dots, e^{\lambda_n t})S^{-1}$.

Applying these two formulas to $Sc$ gives the stated identities.

*Structural Commentary*: Eigenbasis coordinates decouple powers and exponentials.

*Key Insight*: Diagonalization turns coupled evolution into independent scalar modes. □

# Part V
# Polynomial Structure and Canonical Forms

## 1 Operator Polynomials

### 1.1 Operator Polynomials and Evaluation

Operator polynomials encode powers and polynomial equations satisfied by a linear operator.

**Definition 1.1.1.** *Operator Powers*
Let $T \in \mathcal{L}(\mathcal{V})$. The *powers* of $T$ are $T^m := \underbrace{T \cdots T}_{m \text{ times}}$ for $m \in \mathbb{N}$, with $T^0 := I_{\mathcal{V}}$. If $T$ is invertible, set $T^{-m} := (T^{-1})^m$.

**Definition 1.1.2.** *Polynomial Applied to an Operator*
Let $T \in \mathcal{L}(\mathcal{V})$.
For $p(z) = \sum_{j=0}^{m} a_j z^j \in \mathbb{F}[z]$, the *polynomial $p$ applied to $T$* is $p(T) := \sum_{j=0}^{m} a_j T^j$.

**Definition 1.1.3.** *Annihilating Polynomial*
Let $T \in \mathcal{L}(\mathcal{V})$.
A polynomial $p \in \mathbb{F}[z]$ is an *annihilating polynomial* for $T$ if $p(T) = 0_{\mathcal{L}(\mathcal{V})}$; equivalently, $p$ *annihilates* $T$.

**Definition 1.1.4.** *Product of Polynomials*
If $p, q \in \mathbb{F}[z]$, their product $pq \in \mathbb{F}[z]$ is defined by $(pq)(z) = p(z)q(z)$ for $z \in \mathbb{F}$.

**Proposition 1.1.5.** *Multiplicative Properties of Operator Polynomials*
*Let $p, q \in \mathbb{F}[z]$ and $T \in \mathcal{L}(\mathcal{V})$. Then*
*(i)* $(pq)(T) = p(T)q(T)$;
*(ii)* $p(T)q(T) = q(T)p(T)$.

*Proof.*
(i) Write $p(z) = \sum_{j=0}^{m} a_j z^j$ and $q(z) = \sum_{k=0}^{n} b_k z^k$. Then $(pq)(z) = \sum_{j=0}^{m} \sum_{k=0}^{n} a_j b_k z^{j+k}$.
Substituting $T$ gives $(pq)(T) = \sum_{j=0}^{m} \sum_{k=0}^{n} a_j b_k T^{j+k} = \left( \sum_{j=0}^{m} a_j T^j \right) \left( \sum_{k=0}^{n} b_k T^k \right) = p(T)q(T)$.
(ii) Since polynomial multiplication is commutative, $pq = qp$.
By part (i), $p(T)q(T) = (pq)(T) = (qp)(T) = q(T)p(T)$.
*Structural Commentary*: Polynomial evaluation preserves products.
*Key Insight*: Expanding a polynomial product works the same with $T$ as with $z$. □

**Proposition 1.1.6.** *Existence of Annihilating Polynomials*
*Let $\mathcal{V}$ be finite-dimensional with $n = \dim(\mathcal{V}) > 0$, and let $T \in \mathcal{L}(\mathcal{V})$.*
*Then $T$ has a nonconstant annihilating polynomial of degree at most $n^2$.*

*Proof.* By Theorem 3.4.4, the vector space $\mathcal{L}(\mathcal{V})$ has dimension $n^2$, so the $n^2 + 1$ operators $I_{\mathcal{V}}, T, T^2, \ldots, T^{n^2}$ are linearly dependent. Hence there are scalars $a_0, \ldots, a_{n^2}$, not all zero, such that $\sum_{j=0}^{n^2} a_j T^j = 0$.
If $a_1 = \cdots = a_{n^2} = 0$, then $a_0 I_{\mathcal{V}} = 0$, forcing $a_0 = 0$ because $\mathcal{V} \neq \{\mathbf{0}\}$, a contradiction. Thus $p(z) = \sum_{j=0}^{n^2} a_j z^j$ is a nonconstant annihilating polynomial.
*Structural Commentary*: Finite-dimensionality makes the powers of one operator linearly dependent inside $\mathcal{L}(\mathcal{V})$.
*Key Insight*: A dependence among $I_{\mathcal{V}}, T, \ldots, T^{n^2}$ is exactly a polynomial equation satisfied by $T$. □

**Proposition 1.1.7.** *Polynomial Evaluation on Eigenvectors*
*Let $T \in \mathcal{L}(\mathcal{V})$, let $p \in \mathbb{F}[z]$, and suppose $T\mathbf{v} = \lambda \mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$. Then $p(T)\mathbf{v} = p(\lambda)\mathbf{v}$.*

*Proof.* Induct on $j$ to get $T^j \mathbf{v} = \lambda^j \mathbf{v}$ for every $j \geq 0$.
If $p(z) = \sum_{j=0}^{m} a_j z^j$, then $p(T)\mathbf{v} = \sum_{j=0}^{m} a_j T^j \mathbf{v} = \sum_{j=0}^{m} a_j \lambda^j \mathbf{v} = p(\lambda)\mathbf{v}$.
*Structural Commentary*: On a one-dimensional eigenspace, $T$ acts like the scalar $\lambda$.
*Key Insight*: Polynomial expressions in $T$ reduce to the same polynomial evaluated at $\lambda$. □

## 1.2 Polynomial Functional Calculus

**Theorem 1.2.1.** *Polynomial Functional Calculus for Diagonalizable Matrices*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{F})$ be diagonalizable and write $A = S\Lambda S^{-1}$, where $\Lambda = \mathrm{diag}(\lambda_1,\dots,\lambda_n)$.*
*If $p \in \mathbb{F}[z]$, then $p(A) = S\,\mathrm{diag}(p(\lambda_1),\dots,p(\lambda_n))S^{-1}$.*

*Proof.* For every $j \geq 0$, $A^j = S\Lambda^j S^{-1}$.

If $p(z) = \sum_j c_j z^j$, then $p(A) = \sum_j c_j A^j = S\left(\sum_j c_j \Lambda^j\right) S^{-1} = Sp(\Lambda)S^{-1}$.
Since $\Lambda$ is diagonal, $p(\Lambda) = \mathrm{diag}(p(\lambda_1),\dots,p(\lambda_n))$.
*Structural Commentary*: Diagonalization reduces $p(A)$ to scalar evaluation on eigenvalues.
*Key Insight*: Compute functions of $A$ by applying the function to the eigenvalues. □

# 2 Characteristic and Minimal Polynomials

## 2.1 Characteristic Polynomial Foundations

**Motivation 2.1.1.** *Two Spectral Polynomials*
The characteristic polynomial records eigenvalues with algebraic multiplicity.
The minimal polynomial records the shortest polynomial relation satisfied by the operator.

**Definition 2.1.2.** *Characteristic Polynomial*
For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, the *characteristic polynomial* of $A$ is $p_A(z) := \det(zI_n - A)$.

**Theorem 2.1.3.** *Basic Form of the Characteristic Polynomial*
*Let $A = [a_{ij}] \in \mathrm{Mat}(n,n,\mathbb{F})$. Then $p_A$ is monic of degree $n$ and $p_A(z) = z^n - (\mathrm{tr}\,A)z^{n-1} + \cdots + (-1)^n \det A$.*
*Moreover, for $\lambda \in \mathbb{F}$, $\lambda \in \mathrm{spec}(A)$ if and only if $p_A(\lambda) = 0$.*

*Proof.* The only determinant term of degree greater than $n-2$ comes from the identity permutation:

$$(z-a_{11})(z-a_{22})\cdots(z-a_{nn}) = z^n - (a_{11}+\cdots+a_{nn})z^{n-1} + \cdots .$$

Thus $p_A$ is monic and the coefficient of $z^{n-1}$ is $-\mathrm{tr}\,A$; also $p_A(0) = \det(-A) = (-1)^n \det A$. Finally, $p_A(\lambda) = 0$ if and only if $\lambda I_n - A$ is singular, equivalently $\ker(A - \lambda I_n) \neq \{\mathbf{0}\}$.
*Structural Commentary*: The diagonal product determines the top two coefficients.
*Key Insight*: Eigenvalues are precisely the roots of $\det(zI_n - A)$. □

**Definition 2.1.4.** *Algebraic Multiplicity*
For $\lambda \in \mathbb{F}$, let $m_A(\lambda)$ be the multiplicity of $\lambda$ as a root of $p_A$, with $m_A(\lambda) = 0$ if $\lambda \notin \mathrm{spec}(A)$.
For $\lambda \in \mathrm{spec}(A)$, this is the *algebraic multiplicity* of $\lambda$. Unqualified multiplicity means algebraic multiplicity.




Figure 57: Form $p_A(z) = \det(zI_n - A)$ and factor it; each root $\lambda$ is counted by $m_A(\lambda)$.

**Method 2.1.5.** *Characteristic Equation and Eigenspace Computation*
To compute eigenvalues and eigenspaces of $A \in \mathrm{Mat}(n,n,\mathbb{F})$:

(i) Form the characteristic polynomial $p_A(z) = \det(zI_n - A)$ from Definition 2.1.2.
(ii) Solve the characteristic equation $p_A(\lambda) = 0$. By Theorem 2.1.3, the solutions are exactly the eigenvalues in $\mathbb{F}$.
(iii) For each eigenvalue $\lambda$, row-reduce $A - \lambda I_n$ and compute

$$E_\lambda(A) = \ker(A - \lambda I_n).$$

(iv) Choose a basis of each eigenspace. The dimension of this basis is the geometric multiplicity of $\lambda$.

Algebraic multiplicity comes from the factorisation of $p_A$, while geometric multiplicity comes from the nullspace computation.

**Proposition 2.1.6.** *Upper-Triangular Eigenvalues*
*If $A = [a_{ij}] \in \mathrm{Mat}(n, n, \mathbb{F})$ is upper triangular, then $p_A(z) = \prod_{j=1}^{n} (z - a_{jj})$.*
*Thus $\mathrm{spec}(A) = \{a_{jj} : 1 \le j \le n\}$, and for $\lambda \in \mathrm{spec}(A)$, $m_A(\lambda) = \#\{j : a_{jj} = \lambda\}$.*

*Proof.* Since $zI_n - A$ is upper triangular with diagonal entries $z - a_{jj}$, $p_A(z) = \det(zI_n - A) = \prod_{j=1}^{n} (z - a_{jj})$.
Thus the roots of $p_A$ are exactly the diagonal entries, and the exponent of $(z - \lambda)$ is $\#\{j : a_{jj} = \lambda\}$.
*Structural Commentary*: Triangular form removes all off-diagonal determinant contributions.
*Key Insight*: Once a matrix is triangular, the spectrum is visible on the diagonal. □

**Theorem 2.1.7.** *Similarity Preserves Characteristic Data*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$ and suppose $B = SAS^{-1}$ for some invertible $S$. Then $p_B = p_A$. In particular, for each $\lambda \in \mathrm{spec}(A)$, $m_B(\lambda) = m_A(\lambda)$ and $\dim E_\lambda(B) = \dim E_\lambda(A)$.*

*Proof.* For every $z$, $p_B(z) = \det(zI_n - SAS^{-1}) = \det(S(zI_n - A)S^{-1}) = \det(S)\det(zI_n - A)\det(S)^{-1} = p_A(z)$.
Thus roots and root multiplicities agree, so $m_B(\lambda) = m_A(\lambda)$. Also $B - \lambda I_n = S(A - \lambda I_n)S^{-1}$.
Hence $\mathbf{x} \in \ker(A - \lambda I_n)$ if and only if $S\mathbf{x} \in \ker(B - \lambda I_n)$, $S$ restricts to isomorphism between two eigenspaces.
*Structural Commentary*: Similarity changes coordinates, not the shifted determinant or shifted kernel dimension.
*Key Insight*: Conjugation inserts $S$ and $S^{-1}$, which cancel in determinants and identify kernels. □

**Proposition 2.1.8.** *Block Upper-Triangular Characteristic Polynomial*
*Let $A = \begin{bmatrix} B & C \\ 0 & D \end{bmatrix}$, $B$ and $D$ are square. Then $p_A(z) = p_B(z)p_D(z)$, so $m_A(\lambda) = m_B(\lambda) + m_D(\lambda)$ $\forall \lambda \in \mathbb{F}$.*

*Proof.* The shifted matrix is block upper triangular, $zI - A = \begin{bmatrix} zI - B & -C \\ 0 & zI - D \end{bmatrix}$.
Theorem 1.1.11 gives $p_A(z) = \det(zI - B)\det(zI - D) = p_B(z)p_D(z)$.
Roots of a product are the roots of the factors, with multiplicity added.
*Structural Commentary*: The off-diagonal block couples vectors but not determinant factors.
*Key Insight*: In block upper-triangular form, spectral data lives on the diagonal blocks. □

**Theorem 2.1.9.** *Geometric Multiplicity Is at Most Algebraic Multiplicity*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ and let $\lambda \in \mathrm{spec}(A)$. Then $\dim E_\lambda(A) \le m_A(\lambda)$.*

*Proof.* Let $k = \dim E_\lambda(A)$ and choose a basis $\mathbf{x}_1, \ldots, \mathbf{x}_k$ of $E_\lambda(A)$. Extend it to a basis of $\mathbb{F}^n$, and let $S$ be the matrix whose columns are this basis. In the new basis,

$$S^{-1}AS = \begin{bmatrix} \lambda I_k & * \\ 0 & C \end{bmatrix}.$$

By Theorem 2.1.7 and Proposition 2.1.8, $p_A(z) = p_{S^{-1}AS}(z) = (z - \lambda)^k p_C(z)$. Thus $m_A(\lambda) \ge k = \dim E_\lambda(A)$.
*Structural Commentary*: An eigenspace-first basis exposes a $\lambda I_k$ block.
*Key Insight*: Each independent eigenvector forces one visible factor $z - \lambda$ in the characteristic polynomial. □

**Theorem 2.1.10.** *Diagonalizability and Multiplicity*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$. Then $A$ is diagonalizable if and only if $\dim E_\lambda(A) = m_A(\lambda)$ for every $\lambda \in \mathrm{spec}(A)$.*

*Proof.*
($\Rightarrow$) If $A$ is diagonalizable, then $A$ is similar to a diagonal matrix $D$. Theorem 2.1.7 gives $m_A(\lambda) = m_D(\lambda)$ and $\dim E_\lambda(A) = \dim E_\lambda(D)$. For diagonal $D$, both numbers count diagonal entries equal to $\lambda$.
($\Leftarrow$) Assume $\dim E_\lambda(A) = m_A(\lambda)$ for every eigenvalue.
If $\mu_1, \ldots, \mu_d$ are distinct eigenvalues, then $\sum_{i=1}^{d} m_A(\mu_i) = n$. Hence $\dim E_{\mu_1}(A) + \cdots + \dim E_{\mu_d}(A) = n$.
Theorem 5.4.3 gives an eigenbasis, so $A$ is diagonalizable.
*Structural Commentary*: Diagonalizability means root counts match eigenspace dimensions.
*Key Insight*: No missing eigendirections means an eigenbasis. □

**Definition 2.1.11.** *Characteristic Polynomial of an Operator*
Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{F}$, let $n = \dim(\mathcal{V})$, and let $T \in \mathcal{L}(\mathcal{V})$. For a basis $\beta$ of $\mathcal{V}$, the *characteristic polynomial* of $T$ is $q_T(z) := p_{[T]_{\beta \leftarrow \beta}}(z) = \det(zI_n - [T]_{\beta \leftarrow \beta})$.
This is independent of $\beta$ by Theorems 6.2.1 and 2.1.7.

**Proposition 2.1.12.** *Degree and Roots of $q_T$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{F}$, let $T \in \mathcal{L}(\mathcal{V})$, and let $q_T$ be the characteristic polynomial of $T$.*

*(i)* $\deg q_T = \dim(\mathcal{V})$.
*(ii) The roots of $q_T$ are precisely the eigenvalues of $T$.*

*Proof.* Choose a basis $\beta$ of $\mathcal{V}$ and set $A = [T]_{\beta \leftarrow \beta}$.

(i) Definition 2.1.11 gives $q_T = p_A$, so Theorem 2.1.3 gives $\deg q_T = \dim(\mathcal{V})$.
(ii) For $\mathbf{v} \in \mathcal{V}$, $T\mathbf{v} = \lambda \mathbf{v}$ if and only if $A[\mathbf{v}]_\beta = \lambda[\mathbf{v}]_\beta$. Since the coordinate map is bijective, $\operatorname{spec}(T) = \operatorname{spec}(A)$. Theorem 2.1.3 identifies this set with the roots of $p_A = q_T$.

*Structural Commentary*: Coordinates transfer degree and eigenvalue data from $A$ to $T$.
*Key Insight*: The characteristic polynomial is independent of the chosen basis. □

## 2.2 Minimal Polynomial Foundations

**Definition 2.2.1.** *Monic Polynomial*
A nonzero polynomial is *monic* if its highest-degree coefficient is 1.

**Theorem 2.2.2.** *Existence of the Minimal Polynomial*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dim over $\mathbb{F}$ and let $T \in \mathcal{L}(\mathcal{V})$. Then $T$ has a unique least-degree monic annihilator.*

*Proof.* Proposition 1.1.6 gives a nonzero annihilator; scaling makes it monic, so a least-degree monic annihilator exists. If $p$ and $q$ are two of them, then $(p-q)(T) = 0$ and $\deg(p-q) < \deg p$.
If $p \neq q$, scaling $p - q$ gives a smaller monic annihilator, impossible. Hence $p = q$.
*Structural Commentary*: Existence comes from finite-dimensional dependence; uniqueness from minimality.
*Key Insight*: Two least monic annihilators cannot differ. □

**Definition 2.2.3.** *Minimal Polynomial of an Operator*
Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional and let $T \in \mathcal{L}(\mathcal{V})$. The *minimal polynomial* of $T$, denoted $\mu_T$, is the unique monic polynomial of smallest degree such that $\mu_T(T) = 0$.

**Definition 2.2.4.** *Minimal Polynomial of a Matrix*
Let $A \in \operatorname{Mat}(n, n, \mathbb{F})$. The *minimal polynomial* of $A$, denoted $\mu_A$, is the unique monic polynomial of smallest positive degree such that $\mu_A(A) = 0$.

**Theorem 2.2.5.** *Annihilators Are Multiples of $\mu_T$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dim, $T \in \mathcal{L}(\mathcal{V})$, and $f \in \mathbb{F}[z]$ be nonzero. Then $f(T) = 0$ if and only if $\mu_T$ divides $f$.*

*Proof.*

($\Rightarrow$) Suppose $f(T) = 0$. Divide $f = s\mu_T + r$, where $r = 0$ or $\deg r < \deg \mu_T$.
Then $r(T) = 0$, so minimality forces $r = 0$. Hence $\mu_T$ divides $f$.
($\Leftarrow$) If $f = \mu_T s$, then $f(T) = \mu_T(T)s(T) = 0$.

*Structural Commentary*: Polynomial division reduces every annihilator to its remainder modulo $\mu_T$.
*Key Insight*: Minimality forces that remainder to be zero. □

# 3 Cayley-Hamilton and Annihilators

## 3.1 Cayley-Hamilton and Minimal Polynomials

**Theorem 3.1.1.** *Cayley–Hamilton Theorem*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{C})$ have characteristic polynomial $p_A(z) = \det(zI_n - A)$. Then $p_A(A) = 0$.*

*Proof.* Choose a Schur form $A = UTU^*$ with diagonal $\lambda_1, \ldots, \lambda_n$.
Since $p_A(A) = Up_A(T)U^*$ and $p_A(z) = \prod_{i=1}^n (z - \lambda_i)$, it suffices to prove $(T-\lambda_1 I_n)(T-\lambda_2 I_n)\cdots(T-\lambda_n I_n) = 0$.
Let $Q_j := \prod_{i=1}^{j} (T - \lambda_i I_n)$. We show that the first $j$ columns of $Q_j$ are zero. For $j = 1$, the first column of $T - \lambda_1 I_n$ is zero as $T$ is upper triangular with first diagonal entry $\lambda_1$. Assume the first $j$ columns of $Q_j$ are zero.
Since $T - \lambda_{j+1} I_n$ is upper triangular and its $(j+1, j+1)$ entry is zero, its first $j+1$ columns lie in $\operatorname{span}(\mathbf{e}_1, \ldots, \mathbf{e}_j)$.
By the induction hypothesis, $Q_j$ kills this span, so $Q_{j+1} = Q_j(T - \lambda_{j+1} I_n)$ has first $j+1$ zero columns.
At $j = n$, all columns of $Q_n$ are zero, so $Q_n = 0$.
*Structural Commentary*: Schur form reduces the theorem to a triangular column-killing induction.
*Key Insight*: Each factor $T - \lambda_i I_n$ kills the next leading column. □




Figure 58: In Schur form, the factors $T - \lambda_i I$ kill the leading columns one layer at a time.

**Theorem 3.1.2.** *Cayley–Hamilton Theorem for Operators*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$, let $T \in \mathcal{L}(\mathcal{V})$, and let $q_T$ be the characteristic polynomial of $T$. Then $q_T(T) = 0$.*

*Proof.* Choose a basis $\beta$ of $\mathcal{V}$ and set $A = [T]_{\beta \leftarrow \beta}$. Definition 2.1.11 gives $q_T = p_A$.
Propositions 3.2.4 and 3.4.3 give $[q_T(T)]_{\beta \leftarrow \beta} = q_T(A) = p_A(A) = 0$ by Theorem 3.1.1.
Theorem 3.4.4 then gives $q_T(T) = 0$.
*Structural Commentary*: Coordinates reduce the operator theorem to the matrix theorem.
*Key Insight*: Polynomial evaluation commutes with taking a matrix representation. □

**Corollary 3.1.3.** *Reduction Modulo the Characteristic Polynomial*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ and let $f \in \mathbb{C}[z]$. If $f = p_A q + r$ with $\deg r < n$, then $f(A) = r(A)$.*

*Proof.* Evaluate division identity at $A$: $f(A) = p_A(A)q(A) + r(A) = 0q(A) + r(A) = r(A)$.
Theorem 3.1.1 gives $p_A(A) = 0$.
*Structural Commentary*: Cayley–Hamilton makes polynomial expressions in $A$ reducible modulo $p_A$.
*Key Insight*: Every high power of $A$ can be rewritten using powers below $n$. □

**Corollary 3.1.4.** *Powers Span at Most n Dimensions*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$. Then $\mathrm{span}(I_n, A, A^2, \ldots) = \mathrm{span}(I_n, A, \ldots, A^{n-1})$, so this has dimension at most $n$.*

*Proof.* For each $k \geq 0$, Corollary 3.1.3 applied to $f(z) = z^{n+k}$ gives a polynomial $r_k$ of degree at most $n - 1$ such that $A^{n+k} = r_k(A)$. Thus every power of $A$ lies in $\mathrm{span}(I_n, A, \ldots, A^{n-1})$.
*Structural Commentary*: Polynomial reduction collapses the infinite power list to its first $n$ terms.
*Key Insight*: Cayley–Hamilton gives the recurrence relation for powers of $A$. □

**Corollary 3.1.5.** *Inverse from the Characteristic Polynomial*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ have $p_A(z) = z^n + c_{n-1}z^{n-1} + \cdots + c_1 z + c_0$.*
*If $A$ is invertible, then $A^{-1} = -c_0^{-1}(A^{n-1} + c_{n-1}A^{n-2} + \cdots + c_2 A + c_1 I_n)$.*

*Proof.* Since $A$ is invertible, $0 \notin \mathrm{spec}(A)$, so $c_0 = p_A(0) = (-1)^n \det A \neq 0$.
Cayley–Hamilton gives $A^n + c_{n-1}A^{n-1} + \cdots + c_1 A + c_0 I_n = 0$.
Move $c_0 I_n$ to the other side and multiply by $-c_0^{-1}A^{-1}$.
*Structural Commentary*: The characteristic equation contains an identity term exactly when $A$ is invertible.
*Key Insight*: Solve Cayley–Hamilton for $A^{-1}$ after factoring out one $A$. □

**Theorem 3.1.6.** *Basic Properties of the Minimal Polynomial*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$, and let $\lambda_1, \ldots, \lambda_d$ be the distinct eigenvalues of $A$.*

*(i)* $\deg \mu_A \leq n$.

*(ii) There are integers $q_i$ satisfying $1 \leq q_i \leq m_A(\lambda_i)$ such that $\mu_A(z) = \prod_{i=1}^{d} (z - \lambda_i)^{q_i}$; in particular, $\mu_A \mid p_A$.*

*Proof.*

(i) Apply Theorem 2.2.2 to the operator induced by $A$. By Theorem 3.1.1, $p_A(A) = 0$; hence minimality gives $\deg \mu_A \leq n$.

(ii) By Theorem 3.1.1, $p_A(A) = 0$. By Theorem 2.2.5, applied to the operator induced by $A$, $\mu_A \mid p_A = \prod_{i=1}^{d} (z - \lambda_i)^{m_A(\lambda_i)}$. Thus the roots of $\mu_A$ are eigenvalues with exponents at most the algebraic multiplicities. If $A\mathbf{v} = \lambda_i \mathbf{v}$ and $\mathbf{v} \neq \mathbf{0}$, then $\mu_A(\lambda_i)\mathbf{v} = \mu_A(A)\mathbf{v} = 0$, so every $\lambda_i$ occurs.

*Structural Commentary*: Cayley–Hamilton bounds degree and divisibility; eigenvectors identify the roots.
*Key Insight*: The least monic annihilator retains every eigenvalue with minimal exponents. □




Figure 59: $\mu_A$ is the least monic annihilator; arrows indicate that $\mu_A$ divides $p_A$ and every annihilator $f$.

**Theorem 3.1.7.** *Similarity Preserves the Minimal Polynomial*
*If $A, B \in \mathrm{Mat}(n, n, \mathbb{C})$ are similar, then $p_A = p_B$ and $\mu_A = \mu_B$.*

*Proof.* The identity $p_A = p_B$ is Theorem 2.1.7. Let $A = SBS^{-1}$, then $f(A) = Sf(B)S^{-1}$ for every polynomial $f$, so $A$ and $B$ have the same annihilating polynomials. Hence $\mu_A = \mu_B$.
*Structural Commentary*: Similarity preserves polynomial equations.
*Key Insight*: Conjugation commutes with polynomial evaluation. □

## 3.2 Diagonalizability and Minimal Polynomial Tests

**Theorem 3.2.1.** *Diagonalizability and the Minimal Polynomial*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ have distinct eigenvalues $\lambda_1, \ldots, \lambda_d$.*
*Then $A$ is diagonalizable if and only if $\mu_A(z) = (z - \lambda_1) \cdots (z - \lambda_d)$.*
*Equivalently, $A$ is diagonalizable if and only if $\mu_A$ is square-free.*

*Proof.* Set $p(z) := \prod_{i=1}^{d}(z - \lambda_i)$. By Theorem 3.1.6, $\mu_A(z) = \prod_{i=1}^{d}(z - \lambda_i)^{q_i}$ with $q_i \geq 1$, so $\mu_A$ is square-free if and only if $\mu_A = p$.

($\Rightarrow$) If $A = S\Lambda S^{-1}$ is diagonalizable, then $p(\Lambda) = 0$, so $p(A) = 0$.
Theorem 3.1.6 gives $\mu_A \mid p$; since every factor $z - \lambda_i$ occurs in $\mu_A$, we have $\mu_A = p$.

($\Leftarrow$) Suppose $\mu_A = p$. For $i \in \{1, \ldots, d\}$, set $\ell_i(z) := \prod_{j \neq i} \frac{z - \lambda_j}{\lambda_i - \lambda_j}$ and $\rho_i := \ell_i(A)$.
The polynomials $\ell_i \ell_j$ for $i \neq j$, $\sum_i \ell_i - 1$, and $z - \sum_i \lambda_i \ell_i$ vanish at $\lambda_1, \ldots, \lambda_d$, so each is divisible by $p$.
Since $p(A) = 0$, $\rho_i \rho_j = 0$ for $i \neq j$, $\sum_i \rho_i = I_n$, and $A = \sum_i \lambda_i \rho_i$. Thus $\rho_i^2 = \rho_i$.
If $\rho_i = 0$, then $\prod_{j \neq i}(A - \lambda_j I_n) = 0$, contradicting the minimality of the degree-$d$ polynomial $\mu_A = p$.
Hence every $\rho_i \neq 0$.
Theorem 5.4.6, applied to the operator induced by $A$, gives that $A$ is diagonalizable.

*Structural Commentary*: Lagrange polynomials turn square-free annihilation into a spectral resolution.
*Key Insight*: Distinct roots produce complementary eigenspace projections. □

**Corollary 3.2.2.** *Square-Free Annihilator Criterion*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$. If a polynomial $f$ with no repeated roots satisfies $f(A) = 0$, then $A$ is diagonalizable.*

*Proof.* Theorem 2.2.5 gives $\mu_A \mid f$. Thus $\mu_A$ is square-free, so Theorem 3.2.1 applies.
*Structural Commentary*: A square-free annihilator forces square-free $\mu_A$.
*Key Insight*: Square-free $\mu_A$ means diagonalizable. □

**Method 3.2.3.** *Computing $\mu_T$ from a Matrix*
Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ be the matrix of $T$ in some basis. Propositions 3.2.4 and 3.4.3 give $[p(T)] = p(A)$ for every $p \in \mathbb{F}[z]$; Theorem 3.4.4 then gives $\mu_T = \mu_A$.

(i) For $m = 1, \ldots, n^2$, test the monic candidate $p_m(z) = a_0 + a_1 z + \cdots + a_{m-1} z^{m-1} + z^m$.
(ii) Solve linear system by comparing matrix entries in $a_0 I + a_1 A + \cdots + a_{m-1} A^{m-1} = -A^m$ for $a_0, \ldots, a_{m-1}$.
(iii) If no solution exists, increase $m$. The first solvable candidate is $\mu_T$.

Proposition 1.1.6 guarantees termination by $m = n^2$.




Figure 60: Solve $a_0 I + \cdots + a_{m-1} A^{m-1} = -A^m$; the first solvable monic candidate is $\mu_T$.

**Method 3.2.4.** *Testing a Minimal Polynomial from $p_A$*
Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$, and suppose $p_A(z) = \prod_i (z - \lambda_i)^{a_i}$ is known.

i. Write each candidate as $m_{\mathbf{q}}(z) = \prod_i (z - \lambda_i)^{q_i}$, where $1 \leq q_i \leq a_i$.

ii. Test exponent vectors $\mathbf{q} = (q_i)$ in increasing total degree $\sum_i q_i$.
iii. For each $\mathbf{q}$, compute $m_{\mathbf{q}}(A) = \prod_i (A - \lambda_i I)^{q_i}$.
iv. The first candidate with $m_{\mathbf{q}}(A) = 0$ is $\mu_A$, by Theorem 3.1.6.

## 3.3 Annihilators and Spectral Control

**Corollary 3.3.1.** *$q_T$ Is a Multiple of $\mu_T$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$ and let $T \in \mathcal{L}(\mathcal{V})$.*
*Then the characteristic polynomial $q_T$ is a polynomial multiple of the minimal polynomial $\mu_T$.*

*Proof.* Theorem 3.1.2 gives $q_T(T) = 0$. Theorem 2.2.5 then implies that $\mu_T$ divides $q_T$.
*Structural Commentary*: Cayley–Hamilton turns the characteristic polynomial into an annihilator.
*Key Insight*: The minimal polynomial divides every annihilator, hence divides $q_T$. □

**Theorem 3.3.2.** *Annihilating Polynomials Control the Spectrum*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $p$ be a nonconstant annihilating polynomial for $T$.*

*(i) Every eigenvalue of $T$ is a root of $p$.*
*(ii) If $\mathbb{F} = \mathbb{C}$, then at least one root of $p$ is an eigenvalue of $T$.*

*Proof.*

(i) If $T\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$, then Proposition 1.1.7 gives $0 = p(T)\mathbf{v} = p(\lambda)\mathbf{v}$. Since $\mathbf{v} \neq \mathbf{0}$, $p(\lambda) = 0$.
(ii) Factor $p(z) = c(z - \lambda_1)\cdots(z - \lambda_m)$ over $\mathbb{C}$, with $c \neq 0$. Then $0 = p(T) = c(T - \lambda_1 I_{\mathcal{V}})\cdots(T - \lambda_m I_{\mathcal{V}})$. At least one factor $T - \lambda_j I_{\mathcal{V}}$ is not injective; otherwise product would be injective. Hence $\lambda_j$ is an eigenvalue.

*Structural Commentary*: Annihilating polynomial restricts spectrum to its roots; over $\mathbb{C}$, one root must appear.
*Key Insight*: Eigenvectors turn $p(T)$ into $p(\lambda)$. □

**Theorem 3.3.3.** *Roots of $\mu_T$ Are Eigenvalues*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional and let $T \in \mathcal{L}(\mathcal{V})$. Then the roots of $\mu_T$ are precisely the eigenvalues of $T$.*

*Proof.*

($\Rightarrow$) If $\mu_T(\lambda) = 0$, write $\mu_T(z) = (z - \lambda)g(z)$. Since $\deg g < \deg \mu_T$, minimality gives $g(T) \neq 0$; choose $\mathbf{v}$ with $\mathbf{u} = g(T)\mathbf{v} \neq \mathbf{0}$. Then $(T - \lambda I_{\mathcal{V}})\mathbf{u} = \mu_T(T)\mathbf{v} = \mathbf{0}$.
($\Leftarrow$) If $T\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$, then $0 = \mu_T(T)\mathbf{v} = \mu_T(\lambda)\mathbf{v}$, so $\mu_T(\lambda) = 0$.

*Structural Commentary*: A root splits off one shifted operator.
*Key Insight*: The quotient cannot already annihilate $T$. □

**Theorem 3.3.4.** *Minimal Polynomial of a Spectrally Disjoint Direct Sum*
*Let $A_i \in \mathrm{Mat}(n_i, n_i, \mathbb{C})$ for $i \in \{1, \ldots, d\}$, let $A = A_1 \oplus \cdots \oplus A_d$, and suppose the spectra of $A_1, \ldots, A_d$ are pairwise disjoint. Then $\mu_A = \mu_{A_1}\mu_{A_2}\cdots\mu_{A_d}$.*

*Proof.* Let $f = \prod_i \mu_{A_i}$. Since $f(A_i) = 0$ for all $i$, blockwise evaluation gives $f(A) = 0$, and therefore $\mu_A \mid f$.
Conversely, $\mu_A(A_i) = 0$ for each $i$, so $\mu_{A_i} \mid \mu_A$ for each $i$. The roots of $\mu_{A_i}$ are exactly $\mathrm{spec}(A_i)$ by Theorem 3.3.3, so the disjointness of the spectra makes the polynomials $\mu_{A_i}$ pairwise relatively prime.
Hence their product $f$ divides $\mu_A$. Since $\mu_A \mid f$ and $f \mid \mu_A$, and both are monic, $\mu_A = f$.
*Structural Commentary*: Direct sums are annihilated blockwise; disjoint spectra make the block factors coprime.
*Key Insight*: Each spectral component contributes its own minimal-polynomial factor. □

**Corollary 3.3.5.** *Minimal and Characteristic Polynomials Have the Same Roots*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$ and let $T \in \mathcal{L}(\mathcal{V})$.*
*Then $\{\lambda \in \mathbb{C} : \mu_T(\lambda) = 0\} = \{\lambda \in \mathbb{C} : q_T(\lambda) = 0\} = \mathrm{spec}(T)$.*

*Proof.* By Theorem 3.3.3, $\{\lambda \in \mathbb{C} : \mu_T(\lambda) = 0\} = \mathrm{spec}(T)$.
By Proposition 2.1.12, $\{\lambda \in \mathbb{C} : q_T(\lambda) = 0\} = \mathrm{spec}(T)$.
*Structural Commentary*: Different multiplicities, same spectral scalars.
*Key Insight*: Minimal polynomial roots and characteristic polynomial roots are both just eigenvalues. □

**Perspective 3.3.6.** *Characteristic vs. Minimal Polynomial*
The characteristic polynomial records eigenvalues with algebraic multiplicity. The minimal polynomial has the same roots, with the least exponents needed to annihilate the operator.

## 3.4 Polynomial Spectral Mapping

**Proposition 3.4.1.** *Eigenvalues of a Polynomial in a Matrix*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$, let $p \in \mathbb{C}[z]$, and let $\lambda_1,\dots,\lambda_n$ be the eigenvalues of $A$, counted with algebraic multiplicity. Then the eigenvalues of $p(A)$ are $p(\lambda_1),\dots,p(\lambda_n)$, counted with algebraic multiplicity.*

*Proof.* Choose a Schur form $U^*AU = T$ with diagonal $\lambda_1,\dots,\lambda_n$. Then $p(A) = Up(T)U^*$, and $p(T)$ is upper triangular with diagonal $p(\lambda_1),\dots,p(\lambda_n)$. Hence $p(A)$ has precisely these eigenvalues.
*Structural Commentary*: Schur form reduces the claim to the diagonal.
*Key Insight*: A polynomial preserves triangular form and applies $p$ to each diagonal entry. □

**Corollary 3.4.2.** *Polynomial Spectral Mapping*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$ and $p \in \mathbb{C}[z]$. Then $\mathrm{spec}(p(A)) = \{p(\lambda) : \lambda \in \mathrm{spec}(A)\}$.*

*Proof.* Let $\lambda_1,\dots,\lambda_n$ be the eigenvalues of $A$, counted with algebraic multiplicity.
By Proposition 3.4.1, the eigenvalues of $p(A)$, counted with algebraic multiplicity, are $p(\lambda_1),\dots,p(\lambda_n)$. Taking underlying sets gives the claim.
*Structural Commentary*: Polynomial evaluation pushes the spectrum forward through the scalar polynomial.
*Key Insight*: Once Schur form has exposed the diagonal, $p$ acts entry by entry there. □

**Corollary 3.4.3.** *Polynomial Spectral Mapping for Operators*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$, let $T \in \mathcal{L}(\mathcal{V})$, and let $p \in \mathbb{C}[z]$. Then $\mathrm{spec}(p(T)) = \{p(\lambda) : \lambda \in \mathrm{spec}(T)\}$.*

*Proof.* Choose a basis $\beta$ and set $A = [T]_{\beta\leftarrow\beta}$.
Then $[p(T)]_{\beta\leftarrow\beta} = p(A)$, and coordinate isomorphism gives $\mathrm{spec}(T) = \mathrm{spec}(A)$ and $\mathrm{spec}(p(T)) = \mathrm{spec}(p(A))$.
By Corollary 3.4.2, $\mathrm{spec}(p(T)) = \mathrm{spec}(p(A)) = \{p(\lambda) : \lambda \in \mathrm{spec}(A)\} = \{p(\lambda) : \lambda \in \mathrm{spec}(T)\}$.
*Structural Commentary*: Coordinates reduce the operator statement to the matrix statement.
*Key Insight*: Polynomial spectral mapping is an operator fact seen through any convenient basis. □

**Corollary 3.4.4.** *Polynomial Invertibility Test*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$ and $p \in \mathbb{C}[z]$. If $p(\lambda) \neq 0$ for every $\lambda \in \mathrm{spec}(A)$, then $p(A)$ is invertible.*

*Proof.* By Corollary 3.4.2, $0 \notin \mathrm{spec}(p(A))$. By Corollary 5.1.8, $p(A)$ is invertible.
*Structural Commentary*: Invertibility of $p(A)$ is checked on spectral values of $A$.
*Key Insight*: Avoiding zero on the spectrum prevents $p(A)$ from having a kernel. □

## 3.5 Polynomial Invariance and Block Separation

**Proposition 3.5.1.** *Kernel and Range of a Polynomial in an Operator Are Invariant*
*Let $T \in \mathcal{L}(\mathcal{V})$ and $p \in \mathbb{F}[z]$. Then $\ker(p(T))$ and $\mathrm{range}(p(T))$ are invariant under $T$.*

*Proof.* If $\mathbf{v} \in \ker(p(T))$, then $p(T)\mathbf{v} = \mathbf{0}$.
Since $p(T)$ commutes with $T$ by Proposition 1.1.5, $p(T)(T\mathbf{v}) = T(p(T)\mathbf{v}) = T\mathbf{0} = \mathbf{0}$.
If $\mathbf{v} \in \mathrm{range}(p(T))$, write $\mathbf{v} = p(T)\mathbf{u}$. Then $T\mathbf{v} = Tp(T)\mathbf{u} = p(T)(T\mathbf{u})$, so $T\mathbf{v} \in \mathrm{range}(p(T))$.
*Structural Commentary*: Polynomial kernels and ranges are stable under the original operator.
*Key Insight*: Since $T$ commutes with $p(T)$, it preserves both $\ker p(T)$ and $\mathrm{range}\, p(T)$. □

**Theorem 3.5.2.** *Sylvester Linear Matrix Equation*
*Let $A \in \mathrm{Mat}(m,m,\mathbb{C})$ and $B \in \mathrm{Mat}(n,n,\mathbb{C})$. If $\mathrm{spec}(A) \cap \mathrm{spec}(B) = \emptyset$, then for every $C \in \mathrm{Mat}(m,n,\mathbb{C})$ the equation $AX - XB = C$ has a unique solution $X \in \mathrm{Mat}(m,n,\mathbb{C})$.*

*Proof.* Let $L(X) = AX - XB$. Since $L$ has equal finite-dimensional domain and codomain, it suffices to show $\ker L = \{0\}$. If $AX = XB$, then $p(A)X = Xp(B)$ for every polynomial $p$.
With $p = p_B$, $p_B(B) = 0$ by Theorem 3.1.1, so $p_B(A)X = Xp_B(B) = 0$. For every $\lambda \in \mathrm{spec}(A)$, disjointness gives $p_B(\lambda) \neq 0$; hence $p_B(A)$ is invertible by Corollary 3.4.4. Thus $X = 0$, so $L$ is bijective.
*Structural Commentary*: Polynomial calculus passes through homogeneous solutions; disjoint spectra invert $p_B(A)$.
*Key Insight*: Spectral separation turns an intertwining relation into the zero relation. □




Figure 61: Disjoint spectra make $p_B(A)$ invertible, so the Sylvester map $L(X) = AX - XB$ is invertible.

**Corollary 3.5.3.** *Removing a Spectrally Separated Off-Diagonal Block*
*Let* $A \in \mathrm{Mat}(m,m,\mathbb{C})$, $B \in \mathrm{Mat}(n,n,\mathbb{C})$, $C \in \mathrm{Mat}(m,n,\mathbb{C})$, *and* $M = \begin{bmatrix} A & C \\ 0 & B \end{bmatrix}$. *If* $\mathrm{spec}(A) \cap \mathrm{spec}(B) = \emptyset$, *then* $M$ *is similar to* $A \oplus B$.

*Proof.* By Theorem 3.5.2, choose $X$ such that $AX - XB = -C$. Then

$$\begin{bmatrix} I & -X \\ 0 & I \end{bmatrix} \begin{bmatrix} A & C \\ 0 & B \end{bmatrix} \begin{bmatrix} I & X \\ 0 & I \end{bmatrix} = \begin{bmatrix} A & AX + C - XB \\ 0 & B \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}.$$

*Structural Commentary*: A block shear cancels the off-diagonal block via Sylvester's equation.
*Key Insight*: Spectrally separated diagonal blocks cannot carry essential upper-right coupling. □

**Theorem 3.5.4.** *Schur Form Grouped by Eigenvalues*
*Let* $A \in \mathrm{Mat}(n,n,\mathbb{C})$ *have distinct eigenvalues* $\lambda_1, \ldots, \lambda_d$ *with algebraic multiplicities* $n_1, \ldots, n_d$.
*Then* $A$ *is unitarily similar to a block upper-triangular matrix*

$$T = \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1d} \\ 0 & T_{22} & \cdots & T_{2d} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & T_{dd} \end{bmatrix}, \qquad T_{ii} \in \mathrm{Mat}(n_i, n_i, \mathbb{C}),$$

*where each* $T_{ii}$ *is upper triangular with all diagonal entries equal to* $\lambda_i$. *Moreover,* $A$ *is similar to* $T_{11} \oplus \cdots \oplus T_{dd}$.

*Proof.* Order the Schur diagonal as $\lambda_1, \ldots, \lambda_1, \lambda_2, \ldots, \lambda_2, \ldots, \lambda_d, \ldots, \lambda_d$.
By Theorem 6.1.10, $A$ has the displayed block Schur form. Since $\mathrm{spec}(T_{11})$ is disjoint from the trailing block spectrum, Corollary 3.5.3 splits off $T_{11}$. Iterating splits off every diagonal block.
*Structural Commentary*: Grouped Schur form isolates spectral blocks; Sylvester shears split disjoint groups.
*Key Insight*: Cross-group upper blocks are coordinate artefacts. □

**Corollary 3.5.5.** *Commuting with Disjoint Blocks*
*Let* $A_{ii} \in \mathrm{Mat}(n_i, n_i, \mathbb{C})$ *for* $i \in \{1, \ldots, d\}$, *set* $n = \sum_{i=1}^{d} n_i$ *and* $A = A_{11} \oplus \cdots \oplus A_{dd}$, *and suppose the spectra of* $A_{11}, \ldots, A_{dd}$ *are pairwise disjoint.*
*Let* $B = [B_{ij}] \in \mathrm{Mat}(n,n,\mathbb{C})$ *be partitioned conformally with* $A$. *If* $AB = BA$, *then* $B_{ij} = 0$ *for all* $i \neq j$.

*Proof.* Equating $(i,j)$ blocks in $AB = BA$ gives $A_{ii}B_{ij} - B_{ij}A_{jj} = 0$.
For $i \neq j$, the spectra of $A_{ii}$ and $A_{jj}$ are disjoint. By Theorem 3.5.2, the homogeneous equation has the unique solution 0, so $B_{ij} = 0$.
*Structural Commentary*: A commuting matrix cannot mix direct-sum blocks with disjoint spectra.
*Key Insight*: The off-diagonal block solves a homogeneous Sylvester equation, whose only solution is zero. □

# 4 Generalized Eigenspaces

## 4.1 Generalized Eigenspaces and Nilpotent Structure

Generalized eigenvectors repair the main defect of ordinary eigenvectors: an operator may not have enough eigendirections to form a basis. The replacement is to study vectors killed by powers of shifted operators.

**Definition 4.1.1.** *Generalized Eigenvectors and Generalized Eigenspaces*
Let $T \in \mathcal{L}(\mathcal{V})$, $\lambda \in \mathbb{F}$. Nonzero vector $\mathbf{v} \in \mathcal{V}$ is a *generalized eigenvector* of $T$ to $\lambda$ if $(T - \lambda I_{\mathcal{V}})^j \mathbf{v} = \mathbf{0}$ for some positive integer $j$. The *generalized eigenspace* corresponding to $\lambda$ is

$$\mathcal{G}_\lambda(T) := \{\mathbf{0}\} \cup \{\mathbf{v} \neq \mathbf{0} : \mathbf{v} \text{ is a generalized eigenvector of } T \text{ corresponding to } \lambda\}.$$

Every eigenvector is a generalized eigenvector by taking $j = 1$. The point is that higher powers of $T - \lambda I_{\mathcal{V}}$ may reveal directions that are not eigenvectors but are eventually pushed into the eigenspace.

**Proposition 4.1.2.** *Kernels of Powers Increase*
*Let* $T \in \mathcal{L}(\mathcal{V})$. *Then* $\{\mathbf{0}\} = \ker(T^0) \subseteq \ker(T) \subseteq \ker(T^2) \subseteq \cdots \subseteq \ker(T^k) \subseteq \ker(T^{k+1}) \subseteq \cdots$.

*Proof.* If $\mathbf{v} \in \ker(T^k)$, then $T^k\mathbf{v} = \mathbf{0}$, hence $T^{k+1}\mathbf{v} = T(T^k\mathbf{v}) = \mathbf{0}$. Thus $\ker(T^k) \subseteq \ker(T^{k+1})$ for every $k \geq 0$.
*Structural Commentary*: Powers of $T$ form an increasing chain of kernels.
*Key Insight*: Once a vector is killed by $T^k$, applying one more $T$ cannot revive it. □

**Proposition 4.1.3.** *Equality Forces Stabilization*
*Let* $T \in \mathcal{L}(\mathcal{V})$ *and* $m \geq 0$. *If* $\ker(T^m) = \ker(T^{m+1})$, *then* $\ker(T^m) = \ker(T^{m+1}) = \ker(T^{m+2}) = \cdots$.

*Proof.* By Proposition 4.1.2, it suffices to prove $\ker(T^{m+k+1}) \subseteq \ker(T^{m+k})$ for every $k \geq 0$.
Take $\mathbf{v} \in \ker(T^{m+k+1})$. Then $T^{m+1}(T^k\mathbf{v}) = T^{m+k+1}\mathbf{v} = \mathbf{0}$, so $T^k\mathbf{v} \in \ker(T^{m+1}) = \ker(T^m)$.
Hence $T^{m+k}\mathbf{v} = T^m(T^k\mathbf{v}) = \mathbf{0}$, proving the reverse inclusion.
*Structural Commentary*: Stabilization at one step propagates through the whole tail.
*Key Insight*: A later kernel vector can be shifted down into the already-stable kernel. □

**Theorem 4.1.4.** *Kernels Stop Growing by $\dim(\mathcal{V})$*
*If $\mathcal{V}$ is finite-dimensional, $T \in \mathcal{L}(\mathcal{V})$, and $n = \dim(\mathcal{V})$, then* $\ker(T^n) = \ker(T^{n+1}) = \ker(T^{n+2}) = \cdots$.

*Proof.* If $\ker(T^n) \neq \ker(T^{n+1})$, then all inclusions $\{\mathbf{0}\} = \ker(T^0) \subsetneq \ker(T) \subsetneq \cdots \subsetneq \ker(T^{n+1})$ are strict by Proposition 4.1.3. Each strict inclusion raises dimension by at least 1, so $\dim\ker(T^{n+1}) \geq n+1$, impossible because $\ker(T^{n+1}) \subseteq \mathcal{V}$. Thus $\ker(T^n) = \ker(T^{n+1})$, and the tail equality follows from Proposition 4.1.3.
*Structural Commentary*: Strict kernel growth is dimension-limited.
*Key Insight*: Dimension bounds force the kernel chain to stabilize no later than step $\dim(\mathcal{V})$. □




Figure 62: The kernel filtration grows until it stabilizes; after equality appears once, the tail is constant.

**Theorem 4.1.5.** *Stable Kernel-Range Decomposition*
*Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and set $n = \dim(\mathcal{V})$. Then* $\mathcal{V} = \ker(T^n) \oplus \operatorname{range}(T^n)$.

*Proof.* First show the intersection is trivial. If $\mathbf{v} \in \ker(T^n) \cap \operatorname{range}(T^n)$, then $\mathbf{v} = T^n\mathbf{u}$ and $T^n\mathbf{v} = \mathbf{0}$. Hence $T^{2n}\mathbf{u} = \mathbf{0}$, so $\mathbf{u} \in \ker(T^{2n}) = \ker(T^n)$ by Theorem 4.1.4. Thus $\mathbf{v} = T^n\mathbf{u} = \mathbf{0}$.
By Rank-Nullity Theorem 1.7.3, $\dim\ker(T^n) + \dim\operatorname{range}(T^n) = \dim(\mathcal{V})$.
The zero intersection and the dimension count imply the sum is direct and equals $\mathcal{V}$.
*Structural Commentary*: At the stable power, killed directions and surviving image directions separate cleanly.
*Key Insight*: Stabilization turns the usual rank-nullity count into an actual direct-sum decomposition. □

**Theorem 4.1.6.** *Description of Generalized Eigenspaces*
*Let $\mathcal{V}$ be finite-dimensional, $T \in \mathcal{L}(\mathcal{V})$, $\lambda \in \mathbb{F}$, and $n = \dim(\mathcal{V})$. Then* $\mathcal{G}_\lambda(T) = \ker\big((T - \lambda I_\mathcal{V})^n\big)$.
*In particular, $\mathcal{G}_\lambda(T)$ is a subspace of $\mathcal{V}$.*

*Proof.* Set $S = T - \lambda I_\mathcal{V}$.
($\Rightarrow$) If $\mathbf{v} \in \mathcal{G}_\lambda(T)$, then $S^j\mathbf{v} = \mathbf{0}$ for some $j \geq 1$. If $j \leq n$, then $\mathbf{v} \in \ker(S^n)$ by Proposition 4.1.2; if $j > n$, then $\ker(S^j) = \ker(S^n)$ by Theorem 4.1.4.
($\Leftarrow$) If $\mathbf{v} \in \ker(S^n)$, then either $\mathbf{v} = \mathbf{0}$ or $\mathbf{v}$ is a generalized eigenvector of $T$ corresponding to $\lambda$.
Thus $\mathcal{G}_\lambda(T) = \ker(S^n)$, which is a subspace.
*Structural Commentary*: Generalized eigenspaces are stable shifted kernels.
*Key Insight*: Finite dimension makes exponent $\dim(\mathcal{V})$ enough. □

**Proposition 4.1.7.** *Generalized Eigenspaces Are Invariant*
*Let $\mathcal{V}$ be finite-dimensional and $T \in \mathcal{L}(\mathcal{V})$. Then $\mathcal{G}_\lambda(T)$ is invariant under $T$ for every $\lambda \in \mathbb{F}$.*

*Proof.* By Theorem 4.1.6, $\mathcal{G}_\lambda(T) = \ker((T - \lambda I_\mathcal{V})^n)$, where $n = \dim(\mathcal{V})$.
This is the kernel of a polynomial in $T$, so it is invariant under $T$ by Proposition 3.5.1.
*Structural Commentary*: A generalized eigenspace is cut out by a polynomial equation in $T$.
*Key Insight*: Polynomial kernels commute with the operator that defines them. □

**Definition 4.1.8.** *Nilpotent Operator*
An operator $N \in \mathcal{L}(\mathcal{V})$ is *nilpotent* if $N^m = 0$ for some positive integer $m$.

**Corollary 4.1.9.** *Nilpotent Power Bound*
*If $\mathcal{V}$ is finite-dimensional, $N \in \mathcal{L}(\mathcal{V})$ is nilpotent, and $n = \dim(\mathcal{V})$, then* $N^n = 0$.

*Proof.* Since $N$ is nilpotent, $\ker(N^m) = \mathcal{V}$ for some $m$.
The kernel chain has stabilized by the $n$th step by Theorem 4.1.4, so $\ker(N^n) = \mathcal{V}$. Hence $N^n = 0$.
*Structural Commentary*: Nilpotence says some kernel in the chain is already the whole space.
*Key Insight*: The stabilization bound pulls that full-kernel conclusion back to exponent $\dim(\mathcal{V})$. □

**Lemma 4.1.10.** *Nonzero Scalar Plus Nilpotent Is Invertible*
*Let $N \in \mathcal{L}(\mathcal{V})$ be nilpotent and let $a \in \mathbb{F}$ with $a \neq 0$. Then $aI_{\mathcal{V}} + N$ is invertible.*

*Proof.* Choose $m$ such that $N^m = 0$. Then $(aI_{\mathcal{V}} + N)\left(a^{-1}\sum_{r=0}^{m-1}(-a^{-1}N)^r\right) = I_{\mathcal{V}} - (-a^{-1}N)^m = I_{\mathcal{V}}$, and the same computation in the reverse order gives a two-sided inverse.
*Structural Commentary*: A nilpotent perturbation has a finite geometric-series inverse.
*Key Insight*: Nilpotence truncates the Neumann series exactly. □

**Theorem 4.1.11.** *Generalized Eigenvectors for Distinct Eigenvalues Are Independent*
*Let $\mathcal{V}$ be finite-dimensional and $T \in \mathcal{L}(\mathcal{V})$. Suppose $\lambda_1, \ldots, \lambda_m$ are distinct eigenvalues of $T$, and let $\mathbf{v}_1, \ldots, \mathbf{v}_m$ be corresponding generalized eigenvectors. Then $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly independent.*

*Proof.* Let $n = \dim(\mathcal{V})$ and suppose $a_1\mathbf{v}_1 + \cdots + a_m\mathbf{v}_m = \mathbf{0}$.
Fix $r \in \{1, \ldots, m\}$ and apply $Q_r(T) := \prod_{i \neq r}(T - \lambda_i I_{\mathcal{V}})^n$ to the relation.
For $i \neq r$, $Q_r(T)\mathbf{v}_i = \mathbf{0}$ by Theorem 4.1.6, so $a_r Q_r(T)\mathbf{v}_r = \mathbf{0}$.
On $\mathcal{G}_{\lambda_r}(T)$, the operator $T - \lambda_i I_{\mathcal{V}}$ equals $(T - \lambda_r I_{\mathcal{V}}) + (\lambda_r - \lambda_i)I_{\mathcal{V}}$. The first term is nilpotent on $\mathcal{G}_{\lambda_r}(T)$ and $\lambda_r - \lambda_i \neq 0$, so each factor is invertible on $\mathcal{G}_{\lambda_r}(T)$ by Lemma 4.1.10. Thus $Q_r(T)\mathbf{v}_r \neq \mathbf{0}$, forcing $a_r = 0$. Since $r$ was arbitrary, all coefficients vanish.
*Structural Commentary*: Polynomial filters isolate one generalized eigenspace at a time.
*Key Insight*: Distinct eigenvalues let polynomial filters kill the other components and preserve the target one. □

**Theorem 4.1.12.** *Nilpotent Operators and Strictly Upper-Triangular Matrices*
*Let $\mathcal{V}$ be finite-dimensional and $N \in \mathcal{L}(\mathcal{V})$. Then $N$ is nilpotent if and only if $N$ has a strictly upper-triangular matrix in some basis.*

*Proof.* Let $n = \dim(\mathcal{V})$. By Corollary 4.1.9, $\{\mathbf{0}\} \subseteq \ker(N) \subseteq \ker(N^2) \subseteq \cdots \subseteq \ker(N^n) = \mathcal{V}$.

(⇒) Choose a basis of $\ker(N)$, extend it to a basis of $\ker(N^2)$, and continue until obtaining a basis of $\mathcal{V}$. If $\mathbf{v}$ first appears in $\ker(N^k)$, then $N\mathbf{v} \in \ker(N^{k-1})$, which is spanned by earlier basis vectors. Hence the matrix of $N$ is strictly upper triangular.

(⇐) If the matrix of $N$ is $n \times n$ strictly upper triangular, then $N$ sends each basis vector into the span of earlier basis vectors. After $n$ applications every basis vector is sent to $\mathbf{0}$, so $N^n = 0$.

*Structural Commentary*: Kernel filtration gives adapted basis; triangular form records motion to earlier layers.
*Key Insight*: Nilpotence is strict upper-triangularity in a suitable basis. □




Figure 63: Extending bases through the kernel filtration makes $N$ move vectors to earlier layers.

**Perspective 4.1.13.** *Eigenvalue Plus Nilpotent Part*
On $\mathcal{G}_\lambda(T)$, the operator $T$ behaves as $\lambda I_{\mathcal{V}}$ plus the nilpotent operator $T - \lambda I_{\mathcal{V}}$. Generalized eigenvectors are therefore ordinary eigenvectors enriched by nilpotent chain data.

# 5 Primary Decomposition

## 5.1 Primary Decomposition and Spectral Splitting

**Theorem 5.1.1.** *Primary Decomposition over $\mathbb{C}$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dim over $\mathbb{C}$, let $T \in \mathcal{L}(\mathcal{V})$, and let $\lambda_1, \ldots, \lambda_m$ be the distinct eigenvalues of $T$. Then*
*(i) $\mathcal{V} = \mathcal{G}_{\lambda_1}(T) \oplus \cdots \oplus \mathcal{G}_{\lambda_m}(T)$;*
*(ii) each $(T - \lambda_j I_{\mathcal{V}})|_{\mathcal{G}_{\lambda_j}(T)}$ is nilpotent.*

*Proof.* Let $n = \dim(\mathcal{V})$.

(i) Proceed by induction on $n$. If $n = 1$, then $T$ has single eigenvalue $\lambda_1$ and $\mathcal{V} = \mathcal{G}_{\lambda_1}(T)$.
Assume $n > 1$. Let $S = T - \lambda_1 I_{\mathcal{V}}$, $\mathcal{U} := \operatorname{range}(S^n)$.
By Theorems 4.1.5 and 4.1.6, $\mathcal{V} = \ker(S^n) \oplus \operatorname{range}(S^n) = \mathcal{G}_{\lambda_1}(T) \oplus \mathcal{U}$. The subspace $\mathcal{U}$ is $T$-invariant by Proposition 3.5.1, and $\dim(\mathcal{U}) < n$ because $\mathcal{G}_{\lambda_1}(T) \neq \{\mathbf{0}\}$. If $\mathcal{U} = \{\mathbf{0}\}$, then (i) follows.
Otherwise distinct eigenvalues of $T|_{\mathcal{U}}$ are $\lambda_2, \ldots, \lambda_m$: every eigenvalue of $T|_{\mathcal{U}}$ is an eigenvalue of $T$ but not $\lambda_1$, while an eigenvector $\mathbf{v}$ of $T$ for $\lambda_j$, $j \geq 2$, satisfies $S^n\mathbf{v} = (\lambda_j - \lambda_1)^n\mathbf{v}$ and hence lies in $\mathcal{U}$.

Induction gives $\mathcal{U} = \mathcal{G}_{\lambda_2}(T|_{\mathcal{U}}) \oplus \cdots \oplus \mathcal{G}_{\lambda_m}(T|_{\mathcal{U}})$. For $j \geq 2$, $\mathcal{G}_{\lambda_j}(T|_{\mathcal{U}}) = \mathcal{G}_{\lambda_j}(T)$.
The inclusion $\subseteq$ follows from the definition. Conversely, if $\mathbf{v} \in \mathcal{G}_{\lambda_j}(T)$, write $\mathbf{v} = \mathbf{g} + \mathbf{u}$ with $\mathbf{g} \in \mathcal{G}_{\lambda_1}(T)$ and $\mathbf{u} \in \mathcal{U}$. For some $r$, $(T - \lambda_j I_{\mathcal{V}})^r \mathbf{v} = \mathbf{0}$. Since both summands are $T$-invariant, the two terms $(T - \lambda_j I_{\mathcal{V}})^r \mathbf{g}$ and $(T - \lambda_j I_{\mathcal{V}})^r \mathbf{u}$ remain in the two direct summands and have sum zero, so both are zero. Thus $\mathbf{g} \in \mathcal{G}_{\lambda_1}(T) \cap \mathcal{G}_{\lambda_j}(T) = \{\mathbf{0}\}$ by Theorem 4.1.11, and $\mathbf{v} = \mathbf{u} \in \mathcal{G}_{\lambda_j}(T|_{\mathcal{U}})$. Hence (i) follows.

(ii) By Proposition 4.1.7, the restriction is an operator on $\mathcal{G}_{\lambda_j}(T)$. By Theorem 4.1.6, $(T - \lambda_j I_{\mathcal{V}})^n \mathbf{v} = \mathbf{0}\ \forall \mathbf{v} \in \mathcal{G}_{\lambda_j}(T)$. Hence $n$th power of $(T - \lambda_j I_{\mathcal{V}})|_{\mathcal{G}_{\lambda_j}(T)}$ is zero.

*Structural Commentary*: Peel off one generalized eigenspace, then induct on the invariant range.
*Key Insight*: Stable kernel-range splitting isolates one eigenvalue at a time. □




Figure 64: Primary decomposition: on $\mathcal{G}_{\lambda_j}(T)$, $T = \lambda_j I$ plus nilpotent.

**Corollary 5.1.2.** *Basis of Generalized Eigenvectors*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$ and $T \in \mathcal{L}(\mathcal{V})$. Then $\mathcal{V}$ has a generalized-eigenvector basis for $T$.*

*Proof.* Let $\lambda_1, \ldots, \lambda_m$ be the distinct eigenvalues of $T$. By Theorem 5.1.1, $\mathcal{V} = \mathcal{G}_{\lambda_1}(T) \oplus \cdots \oplus \mathcal{G}_{\lambda_m}(T)$. Choose a basis of each generalized eigenspace and concatenate the bases. Directness makes the concatenated list a basis of $\mathcal{V}$, and every nonzero vector in each list is a generalized eigenvector.
*Structural Commentary*: A direct sum basis is obtained by choosing bases in each summand.
*Key Insight*: Bases of the direct-sum pieces concatenate. □

**Theorem 5.1.3.** *Block Diagonal Form with Upper-Triangular Primary Blocks*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$, let $\lambda_1, \ldots, \lambda_m$ be the distinct eigenvalues of $T \in \mathcal{L}(\mathcal{V})$, and set $d_j = \dim \mathcal{G}_{\lambda_j}(T)$. Then $T$ has a matrix of the form*

$$\begin{bmatrix} A_1 & & 0 \\ & \ddots & \\ 0 & & A_m \end{bmatrix},$$

*where each $A_j \in \mathrm{Mat}(d_j, d_j, \mathbb{C})$ is upper triangular with diagonal $\lambda_j, \ldots, \lambda_j$.*

*Proof.* Set $\mathcal{U}_j = \mathcal{G}_{\lambda_j}(T)$ and $N_j = (T - \lambda_j I_{\mathcal{V}})|_{\mathcal{U}_j}$.
By Theorem 5.1.1, the $\mathcal{U}_j$ form a $T$-invariant direct-sum decomposition and each $N_j$ is nilpotent.
By Theorem 4.1.12, choose a basis of each $\mathcal{U}_j$ in which $N_j$ is strictly upper triangular. Then $T|_{\mathcal{U}_j} = \lambda_j I_{\mathcal{U}_j} + N_j$ has an upper triangular matrix with diagonal $\lambda_j, \ldots, \lambda_j$. Concatenating these bases gives the desired block diagonal matrix.
*Structural Commentary*: Primary decomposition gives the blocks; nilpotent triangularisation shapes them.
*Key Insight*: On each generalized eigenspace, $T$ is eigenvalue plus nilpotent. □




Figure 65: Primary blocks: $A_j = \lambda_j I + N_j$, with $N_j$ strictly upper triangular.

**Definition 5.1.4.** *Multiplicity of an Eigenvalue*
Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{C}$ and let $T \in \mathcal{L}(\mathcal{V})$.

The *multiplicity* of an eigenvalue $\lambda$ of $T$ is $\dim \mathcal{G}_\lambda(T) = \dim \ker\big((T-\lambda I_{\mathcal{V}})^{\dim(\mathcal{V})}\big)$.
By Theorem 5.1.3 and Proposition 2.1.6, this equals the multiplicity of $\lambda$ as a root of $q_T$.




Figure 66: Each primary summand contributes $(z-\lambda_j)^{d_j}$, where $d_j = \dim \mathcal{G}_{\lambda_j}(T)$.

**Proposition 5.1.5.** *Sum of Multiplicities*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be a finite-dimensional complex vector space and let $T \in \mathcal{L}(\mathcal{V})$. Then the sum of the multiplicities of all eigenvalues of $T$ equals* $\dim(\mathcal{V})$.

*Proof.* Let $\lambda_1, \ldots, \lambda_m$ be the distinct eigenvalues of $T$. By Theorem 5.1.1, $\mathcal{V} = \mathcal{G}_{\lambda_1}(T) \oplus \cdots \oplus \mathcal{G}_{\lambda_m}(T)$.
Taking dimensions, by Corollary 2.3.2, $\dim(\mathcal{V}) = \sum_{j=1}^{m} \dim \mathcal{G}_{\lambda_j}(T)$, exactly the sum of the multiplicities.
*Structural Commentary*: Multiplicity is dimension of the generalized-eigenspace summand.
*Key Insight*: Multiplicities are dimensions of the direct-sum pieces. □

## 5.2 Real Spectral Data via Complexification

**Lemma 5.2.1.** *Conjugation Transfers Generalized Eigenspaces*
*Let $\mathcal{V}$ be a real vector space, let $T \in \mathcal{L}(\mathcal{V})$, let $\lambda \in \mathbb{C}$, let $k \geq 0$, and let $\mathbf{u}, \mathbf{v} \in \mathcal{V}$. Then $(T_{\mathbb{C}} - \lambda I_{\mathcal{V}_{\mathbb{C}}})^k(\mathbf{u}+i\mathbf{v}) = \mathbf{0}$ $\iff$ $(T_{\mathbb{C}} - \overline{\lambda} I_{\mathcal{V}_{\mathbb{C}}})^k(\mathbf{u}-i\mathbf{v}) = \mathbf{0}$.*

*Proof.* Let $\Gamma(\mathbf{u}+i\mathbf{v}) = \mathbf{u}-i\mathbf{v}$.
Then $\Gamma$ is a conjugate-linear bijection, $\Gamma T_{\mathbb{C}} = T_{\mathbb{C}}\Gamma$, and $\Gamma\big((T_{\mathbb{C}} - \lambda I)^k \mathbf{x}\big) = (T_{\mathbb{C}} - \overline{\lambda} I)^k \Gamma \mathbf{x}$.
($\Rightarrow$) Apply the displayed identity to $\mathbf{x} = \mathbf{u} + i\mathbf{v}$.
($\Leftarrow$) Apply the first direction with $\overline{\lambda}$ in place of $\lambda$ and $\mathbf{u} - i\mathbf{v}$ in place of $\mathbf{u} + i\mathbf{v}$.
*Structural Commentary*: Conjugation commutes with $T_{\mathbb{C}}$ and conjugates scalars.
*Key Insight*: Conjugation pairs generalized eigenspaces. □

**Corollary 5.2.2.** *Conjugate Eigenvalues Have Equal Multiplicity*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{R}$ and let $T \in \mathcal{L}(\mathcal{V})$.*
*If $\lambda$ is an eigenvalue of $T_{\mathbb{C}}$, then $\overline{\lambda}$ is an eigenvalue of $T_{\mathbb{C}}$, and* $\dim_{\mathbb{C}} \mathcal{G}_\lambda(T_{\mathbb{C}}) = \dim_{\mathbb{C}} \mathcal{G}_{\overline{\lambda}}(T_{\mathbb{C}})$.

*Proof.* By Lemma 5.2.1 with $k = 1$, conjugation sends a nonzero eigenvector for $\lambda$ to a nonzero eigenvector for $\overline{\lambda}$. Thus $\overline{\lambda}$ is an eigenvalue. With $k = \dim_{\mathbb{C}} \mathcal{V}_{\mathbb{C}}$, the generalized eigenspaces are the corresponding stabilized kernels by Theorem 4.1.6. By Lemma 5.2.1, $\Gamma(\mathbf{u}+i\mathbf{v}) = \mathbf{u}-i\mathbf{v}$ restricts to conjugate-linear bijection $\mathcal{G}_\lambda(T_{\mathbb{C}}) \longrightarrow \mathcal{G}_{\overline{\lambda}}(T_{\mathbb{C}})$. Conjugate-linear bijections preserve complex dimension, so the multiplicities agree.
*Structural Commentary*: Conjugation mirrors generalized eigenspaces.
*Key Insight*: Nonreal spectral data of a real operator arrives in equal-sized conjugate pairs. □

**Corollary 5.2.3.** *Odd-Dimensional Real Operators Have Eigenvalues*
*Every operator on a nonzero odd-dimensional real vector space has a real eigenvalue.*

*Proof.* Let $T \in \mathcal{L}(\mathcal{V})$, where $\dim_{\mathbb{R}} \mathcal{V}$ is odd. By Proposition 5.3.7, $\dim_{\mathbb{C}} \mathcal{V}_{\mathbb{C}}$ is odd.
The sum of the multiplicities of all eigenvalues of $T_{\mathbb{C}}$ equals this odd dimension by Proposition 5.1.5. Nonreal eigenvalues contribute in conjugate pairs of equal multiplicity by Corollary 5.2.2, hence contribute an even total.
Therefore $T_{\mathbb{C}}$ has at least one real eigenvalue, which is an eigenvalue of $T$ by Proposition 5.3.14.
*Structural Commentary*: Odd dimension forces a real eigenvalue.
*Key Insight*: Parity forces at least one real spectral direction. □

**Theorem 5.2.4.** *Characteristic Polynomial over $\mathbb{R}$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be a finite-dimensional real vector space, let $T \in \mathcal{L}(\mathcal{V})$, and let $q_T$ be its characteristic polynomial.*
*(i)* $q_T \in \mathbb{R}[z]$.
*(ii)* $\deg q_T = \dim_{\mathbb{R}} \mathcal{V}$.

*(iii) The real zeros of $q_T$ are precisely the eigenvalues of $T$.*
*(iv) $q_T(T) = 0$.*

*Proof.* Choose a basis $\beta$ of $\mathcal{V}$ and its complexification $\beta_{\mathbb{C}}$. Since $[T_{\mathbb{C}}]_{\beta_{\mathbb{C}} \leftarrow \beta_{\mathbb{C}}} = [T]_{\beta \leftarrow \beta}$, Definition 2.1.11 gives $q_{T_{\mathbb{C}}} = q_T$.

(i) Factor $q_{T_{\mathbb{C}}}$ over $\mathbb{C}$. Real eigenvalues give real linear factors. If $\lambda \notin \mathbb{R}$ has multiplicity $m$, then $\overline{\lambda}$ has multiplicity $m$ by Corollary 5.2.2, and $(z-\lambda)^m(z-\overline{\lambda})^m = \left(z^2 - 2\operatorname{Re}(\lambda)z + |\lambda|^2\right)^m$, which is real.
(ii) By Propositions 2.1.12 and 5.3.7, $\deg q_T = \dim_{\mathbb{R}} \mathcal{V}$.
(iii) By Proposition 2.1.12, the real zeros of $q_T = q_{T_{\mathbb{C}}}$ are the real eigenvalues of $T_{\mathbb{C}}$; these are the eigenvalues of $T$ by Proposition 5.3.14.
(iv) By Theorem 3.1.2, $q_T(T_{\mathbb{C}}) = 0$. By (i) and Proposition 5.3.11, $q_T(T_{\mathbb{C}}) = (q_T(T))_{\mathbb{C}}$. If $q_T(T)$ were nonzero on some $\mathbf{u} \in \mathcal{V}$, then $(q_T(T))_{\mathbb{C}}(\mathbf{u} + i\mathbf{0}) = q_T(T)\mathbf{u} + i\mathbf{0}$ would be nonzero, a contradiction. Hence $q_T(T) = 0$.

*Structural Commentary*: Nonreal factors pair into real quadratics.
*Key Insight*: Complexify, then pair conjugates. □




Figure 67: Conjugate factors pair into real quadratics.

**Corollary 5.2.5.** *$q_T$ Is a Multiple of $\mu_T$ over $\mathbb{R}$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be a finite-dimensional real vector space and let $T \in \mathcal{L}(\mathcal{V})$. Then $\deg \mu_T \leq \dim_{\mathbb{R}} \mathcal{V}$, and $q_T$ is a polynomial multiple of $\mu_T$ in $\mathbb{R}[z]$.*

*Proof.* By Theorem 5.2.4, $q_T(T) = 0$.
By Theorem 2.2.5, $\mu_T$ divides $q_T$ in $\mathbb{R}[z]$. The degree bound follows from $\deg q_T = \dim_{\mathbb{R}} \mathcal{V}$.
*Structural Commentary*: Cayley–Hamilton makes $q_T$ a real annihilator of $T$.
*Key Insight*: The minimal polynomial divides every real annihilator. □

**Proposition 5.2.6.** *Minimal Polynomial Survives Complexification*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dimensional over $\mathbb{R}$ and let $T \in \mathcal{L}(\mathcal{V})$. Then $\mu_{T_{\mathbb{C}}} = \mu_T$ in $\mathbb{C}[z]$.*

*Proof.*
($\Rightarrow$) Since $\mu_T(T) = 0$, $\mu_T(T_{\mathbb{C}}) = (\mu_T(T))_{\mathbb{C}} = 0$ by Proposition 5.3.11. Hence $\mu_{T_{\mathbb{C}}}$ divides $\mu_T$ in $\mathbb{C}[z]$.
($\Leftarrow$) Let $q = \mu_{T_{\mathbb{C}}} = r + is$, with $r, s \in \mathbb{R}[z]$. Since $q$ is monic, $r$ is monic and $\deg r = \deg q$. For every $\mathbf{u} \in \mathcal{V}$, $0 = q(T_{\mathbb{C}})(\mathbf{u} + i\mathbf{0}) = r(T)\mathbf{u} + is(T)\mathbf{u}$. Thus $r(T) = 0$, so $\deg \mu_T \leq \deg r = \deg q$. The first part gives opposite degree inequality. As $q$ divides $\mu_T$ and both are monic with equal degree, $q = \mu_T$.

*Structural Commentary*: Annihilation passes to complexification and back by real-part extraction.
*Key Insight*: Evaluate $q(T_{\mathbb{C}})$ on $\mathbf{u} + i\mathbf{0}$. □

## 5.3 Normal Primary Structure

**Corollary 5.3.1.** *Normal Operators Have No Nilpotent Primary Part*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be a finite-dimensional complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$ be normal.*
*(i) If $S = p(T)$ for some $p \in \mathbb{C}[z]$, then $\ker(S^k) = \ker(S)$ for every $k \geq 1$.*
*(ii) The minimal polynomial of $T$ is square-free, where $\mu_T(z) = \prod_{\lambda \in \operatorname{spec}(T)} (z - \lambda)$.*

*Proof.* By Theorem 6.1.13, choose an orthonormal eigenbasis $\mathbf{e}_1, \ldots, \mathbf{e}_n$ with $T\mathbf{e}_j = \lambda_j \mathbf{e}_j$.

(i) For $S = p(T)$, $k \geq 1$, $S^k\mathbf{e}_j = p(\lambda_j)^k \mathbf{e}_j$; $\ker(S^k) = \operatorname{span}\{\mathbf{e}_j : p(\lambda_j)^k = 0\} = \operatorname{span}\{\mathbf{e}_j : p(\lambda_j) = 0\} = \ker(S)$.
(ii) Let $m(z) = \prod_{\lambda \in \operatorname{spec}(T)}(z - \lambda)$. Since $m(T)\mathbf{e}_j = 0$ for all $j$, $m(T) = 0$, so $\mu_T \mid m$. Conversely, every $\lambda \in \operatorname{spec}(T)$ is a root of $\mu_T$ by Theorem 3.3.3, hence $m \mid \mu_T$. Since both polynomials are monic, $\mu_T = m$.

*Structural Commentary*: Normality makes every primary component an eigenspace.
*Key Insight*: A diagonal operator has no hidden nilpotent part. □

**Proposition 5.3.2.** *Coprime Normal Primary Pieces Are Orthogonal*
*Let $\mathcal{V}$ be a finite-dimensional complex inner-product space, let $T \in \mathcal{L}(\mathcal{V})$ be normal, and let $\mathcal{S}, \mathcal{R} \leq \mathcal{V}$ be $T$-invariant.*
*If $p, q \in \mathbb{C}[z]$, $(p, q) = 1$, $p(T)|_{\mathcal{S}} = 0$, and $q(T)|_{\mathcal{R}} = 0$, then $\mathcal{S} \perp \mathcal{R}$.*

*Proof.* Choose $a, b \in \mathbb{C}[z]$ with $ap + bq = 1$. For $\mathbf{s} \in \mathcal{S}$, $\mathbf{s} = a(T)p(T)\mathbf{s} + b(T)q(T)\mathbf{s} = b(T)q(T)\mathbf{s}$.
By Corollary 6.2.12, $\mathcal{R}$ reduces $T$, so it reduces $b(T)$ and $q(T)$. Hence $b(T)^*\mathcal{R} \subseteq \mathcal{R}$ and $q(T)^*|_{\mathcal{R}} = (q(T)|_{\mathcal{R}})^* = 0$.
Thus, for $\mathbf{r} \in \mathcal{R}$, $\langle \mathbf{s}, \mathbf{r} \rangle = \langle b(T)q(T)\mathbf{s}, \mathbf{r} \rangle = \langle \mathbf{s}, q(T)^* b(T)^* \mathbf{r} \rangle = 0$.
*Structural Commentary*: Coprime annihilators orthogonally separate normal invariant pieces.
*Key Insight*: Normality makes invariant subspaces reduce, so annihilation also holds after taking adjoints. □

# 6 Jordan and Canonical Forms

## 6.1 Jordan Chains and Nilpotent Blocks

Jordan and canonical forms turn primary decomposition into explicit blocks. Jordan chains, complexification, and real block forms organize the nilpotent and real spectral structure. Each Jordan block records one eigenvalue together with the length of a nilpotent chain.

**Definition 6.1.1.** *Jordan Chains, Blocks, and Matrices*
Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$, let $\lambda \in \mathbb{C}$, and let $k \geq 1$. A *Jordan chain* for $A$ at $\lambda$ is a list $(\mathbf{x}_1, \ldots, \mathbf{x}_k)$ such that $(A - \lambda I_n)\mathbf{x}_1 = \mathbf{0}$ and $(A - \lambda I_n)\mathbf{x}_j = \mathbf{x}_{j-1}$ for $2 \leq j \leq k$.
The *nilpotent Jordan block* $J_k(0)$ is the $k \times k$ matrix with 1 on the first superdiagonal and 0 elsewhere:

$$J_k(0) = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & \cdots & 0 & 0 \end{bmatrix}.$$

Set $J_k(\lambda) := \lambda I_k + J_k(0)$. A *Jordan matrix* is a direct sum of Jordan blocks. A basis is a *Jordan basis* for an operator $T$ if the matrix of $T$ in that basis is a Jordan matrix.

**Perspective 6.1.2.** *Jordan Chains as Columns*
For a Jordan chain $(\mathbf{x}_1, \ldots, \mathbf{x}_k)$ and $X = [\mathbf{x}_1 \ \cdots \ \mathbf{x}_k]$, $AX = XJ_k(\lambda)$.
If all chains form the columns of a basis matrix $M$, then $AM = MJ$ and $M^{-1}AM = J$.




Figure 68: Jordan chain for $N = A - \lambda I$; columns give $AX = XJ_k(\lambda)$.

**Theorem 6.1.3.** *Nilpotent Jordan Chain Basis*
*Let $\mathcal{V}$ be finite-dim, $N \in \mathcal{L}(\mathcal{V})$ be nilpotent. Then $\exists$ vectors $\mathbf{v}_1, \ldots, \mathbf{v}_r \in \mathcal{V}$ and nonnegative integers $m_1, \ldots, m_r$ such that $\mathcal{B}_N = \left(N^{m_1}\mathbf{v}_1, \ldots, N\mathbf{v}_1, \mathbf{v}_1, \ldots, N^{m_r}\mathbf{v}_r, \ldots, N\mathbf{v}_r, \mathbf{v}_r\right)$ is a basis of $\mathcal{V}$ and $N^{m_j+1}\mathbf{v}_j = \mathbf{0}$ for every $j$.*

*Proof.* Induct on $n = \dim(\mathcal{V})$. If $N = 0$, any basis works with all $m_j = 0$.
Now assume $N \neq 0$. Then $N$ is not injective, so $\mathrm{range}(N)$ is a proper subspace, and $N|_{\mathrm{range}(N)}$ is nilpotent.
By induction, $\mathrm{range}(N)$ has a chain basis $\left(N^{m_1}\mathbf{u}_1, \ldots, N\mathbf{u}_1, \mathbf{u}_1, \ldots, N^{m_r}\mathbf{u}_r, \ldots, N\mathbf{u}_r, \mathbf{u}_r\right)$ with $N^{m_j+1}\mathbf{u}_j = \mathbf{0}$.
Choose $\mathbf{v}_j \in \mathcal{V}$ such that $N\mathbf{v}_j = \mathbf{u}_j$, and lift the chains to $\mathcal{L} = \left(N^{m_1+1}\mathbf{v}_1, \ldots, N\mathbf{v}_1, \mathbf{v}_1, \ldots, N^{m_r+1}\mathbf{v}_r, \ldots, N\mathbf{v}_r, \mathbf{v}_r\right)$.
The list $\mathcal{L}$ is linearly independent. Indeed, applying $N$ to a dependence among vectors in $\mathcal{L}$ gives a dependence among the range-chain basis, so all coefficients except possibly those on the leading vectors $N^{m_j+1}\mathbf{v}_j$ vanish.
The remaining relation is a relation among $N^{m_j+1}\mathbf{v}_j = N^{m_j}\mathbf{u}_j$, which are members of the range-chain basis, so those coefficients vanish too. Also $N(\mathrm{span}(\mathcal{L})) = \mathrm{range}(N)$: each range-chain basis vector $N^q\mathbf{u}_j$ is $N(N^q\mathbf{v}_j)$ for the corresponding lifted vector. Extend $\mathcal{L}$ to a basis by adding $\mathbf{w}_1, \ldots, \mathbf{w}_s$. For each $i$, choose $\mathbf{x}_i \in \mathrm{span}(\mathcal{L})$ such that $N\mathbf{x}_i = N\mathbf{w}_i$, and set $\mathbf{z}_i = \mathbf{w}_i - \mathbf{x}_i$. Then $N\mathbf{z}_i = \mathbf{0}$. Since each $\mathbf{x}_i$ lies in $\mathrm{span}(\mathcal{L})$, replacing $\mathbf{w}_i$ by $\mathbf{z}_i$ preserves the span and linear independence of the full basis. Thus $\mathcal{L}, \mathbf{z}_1, \ldots, \mathbf{z}_s$ is the desired chain basis.
*Structural Commentary*: Build chains in $\mathrm{range}(N)$, lift them once, then add kernel chains.
*Key Insight*: Applying $N$ turns lifted-chain dependences into known range-chain dependences. □




Figure 69: A nilpotent chain shifts left under $N$; in this basis order it becomes one Jordan block.

**Method 6.1.4.** *Filippov Column-Space Jordan Chains*
Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$, let $\lambda \in \mathrm{spec}(A)$, and set $N = A - \lambda I_n$ on the generalized eigenspace $\mathcal{G}_\lambda(A) = \ker(N^n)$.

(i) Work first inside $\mathrm{range}(N) \cap \mathcal{G}_\lambda(A)$, where $N$ is again nilpotent.
(ii) Build Jordan chains for this smaller nilpotent operator.
(iii) For each chain ending in $\mathbf{u}$, solve $N\mathbf{v} = \mathbf{u}$ in $\mathcal{G}_\lambda(A)$ and lift the chain by appending $\mathbf{v}$.
(iv) Add enough vectors from $\ker(N)$ to complete a basis of $\mathcal{G}_\lambda(A)$; these added vectors are length-one chains.
(v) Repeat the process recursively until all of $\mathcal{G}_\lambda(A)$ is covered, then concatenate the chains for all eigenvalues.

The lifting steps are valid by Theorem 6.1.3; the chain relations give $AM = MJ$ for the resulting basis matrix.

**Proposition 6.1.5.** *Powers of a Jordan Block*
*Let $k \geq 1$ and $\lambda \in \mathbb{C}$. For $p \geq 0$, $J_k(\lambda)^p = \sum_{r=0}^{\min(p,k-1)} \binom{p}{r} \lambda^{p-r} J_k(0)^r$. Thus $J_k(\lambda)^p \to 0$ if and only if $|\lambda| < 1$.*

*Proof.*

($\Rightarrow$) If $J_k(\lambda)^p \to 0$, then its diagonal entries $\lambda^p$ tend to 0, so $|\lambda| < 1$.
($\Leftarrow$) If $|\lambda| < 1$, then for each fixed $0 \leq r < k$ the scalar $\binom{p}{r}\lambda^{p-r}$ tends to 0, because a polynomial in $p$ is dominated by the exponential decay of $|\lambda|^p$.

*Structural Commentary*: Jordan powers are exponential-polynomial terms.
*Key Insight*: Polynomial growth cannot defeat exponential decay when $|\lambda| < 1$. □

## 6.2 Complex Jordan Form and Block Sizes

**Theorem 6.2.1.** *Jordan Form over $\mathbb{C}$*
*Let $\dim_{\mathbb{C}} \mathcal{V} = n \geq 1$ and let $T \in \mathcal{L}(\mathcal{V})$. Then $\exists$ a basis $\mathcal{B}$ of $\mathcal{V}$ and a Jordan matrix $J$ with $[T]_{\mathcal{B}} = J$.*

*Proof.* Let $\mathrm{spec}(T) = \{\lambda_1, \ldots, \lambda_m\}$, set $G_j = \mathcal{G}_{\lambda_j}(T)$, $N_j = (T - \lambda_j I_{\mathcal{V}})|_{G_j}$, and $T_j = T|_{G_j}$.
By Theorem 5.1.1, $\mathcal{V} = \bigoplus_{j=1}^{m} G_j$ and each $N_j$ is nilpotent. By Theorem 6.1.3, choose an $N_j$-chain basis $\mathcal{B}_j$ of $G_j$.
For an $N_j$-chain $\mathcal{C} \subset \mathcal{B}_j$ of length $r$, $[N_j]_{\mathcal{C}} = J_r(0)$ and $[T_j]_{\mathcal{C}} = \lambda_j I_r + J_r(0) = J_r(\lambda_j)$. Let $\mathcal{B} = \mathcal{B}_1 \sqcup \cdots \sqcup \mathcal{B}_m$.
By directness, $\mathcal{B}$ is a basis of $\mathcal{V}$, and $[T]_{\mathcal{B}}$ is a direct sum of Jordan blocks.
*Structural Commentary*: Primary decomposition reduces $T$ to nilpotent chain bases.
*Key Insight*: On $G_j$, $T = \lambda_j I + N_j$. □

**Theorem 6.2.2.** *Real Jordan Block Form*
*Let $n \in \mathbb{N}$, $A \in \mathrm{Mat}(n, n, \mathbb{R})$. For $a \in \mathbb{R}$, $b > 0$, set $R(a, b) := \left(\begin{smallmatrix} a & b \\ -b & a \end{smallmatrix}\right)$.*
*Let $\mathcal{J}_k(a, b) = [C_{rs}]_{r,s=1}^{k}$ with $C_{rr} = R(a, b)$, $C_{r,r+1} = I_2$ for $r < k$, and $C_{rs} = 0_2$ otherwise.*
*Then $\exists S \in \mathrm{GL}_n(\mathbb{R})$ such that $S^{-1}AS$ is a direct sum of blocks of the following types:*

*(i) $J_k(\lambda)$, where $k \in \mathbb{N}$ and $\lambda \in \mathbb{R}$;*
*(ii) $\mathcal{J}_k(a, b)$, where $k \in \mathbb{N}$, $a \in \mathbb{R}$, and $b > 0$.*

*Proof.* Complexify $A$ to $A_{\mathbb{C}}$. By Theorem 6.2.1, choose Jordan chains for $A_{\mathbb{C}}$.
By $\chi_A \in \mathbb{R}[t]$, $\lambda \notin \mathbb{R}$ implies $\overline{\lambda} \in \mathrm{spec}(A_{\mathbb{C}})$.

(i) For $\lambda \in \mathbb{R}$, let $G_\lambda^{\mathbb{R}} = \ker(A - \lambda I_n)^n$ and $N_\lambda = (A - \lambda I_n)|_{G_\lambda^{\mathbb{R}}}$.
Then $(G_\lambda^{\mathbb{R}})_{\mathbb{C}} = \ker(A_{\mathbb{C}} - \lambda I_n)^n$ and $N_\lambda$ is nilpotent.
By Theorem 6.1.3, take an $N_\lambda$-chain basis $\mathcal{B}_\lambda$; $[A]_{\mathcal{B}_\lambda}$ is a direct sum of $J_k(\lambda)$ blocks.
(ii) Let $\lambda = a + ib$, $b > 0$, and let $v = (v_1, \ldots, v_k)$ be a $\lambda$-chain. By Lemma 5.2.1, $\overline{v}$ is a $\overline{\lambda}$-chain.
Let $v_j = u_j + iw_j$; $U_j = [u_j\ w_j]$, $U_0 = 0$, $\mathcal{U}_v = (u_1, w_1, \ldots, u_k, w_k)$, $W_v = \mathrm{span}_{\mathbb{C}}(v, \overline{v}) \cap \mathbb{R}^n$.
By $(v, \overline{v}) \leftrightarrow \mathcal{U}_v$, $\mathcal{U}_v$ is real basis of $W_v$. By chain relations, $AU_j = U_j R(a, b) + U_{j-1}$; $[A]_{\mathcal{U}_v} = \mathcal{J}_k(a, b)$.

Let $\mathcal{B}$ concatenate all $\mathcal{B}_\lambda$ and one $\mathcal{U}_v$ for each pair $(\lambda, \overline{\lambda})$, $\mathrm{Im}\,\lambda > 0$.
By primary directness, $\mathcal{B}$ is a real basis of $\mathbb{R}^n$. With $S = [\mathcal{B}]$, $S^{-1}AS$ has the asserted blocks.
*Structural Commentary*: Real chains and conjugate-pair chains exhaust the primary decomposition.
*Key Insight*: A nonreal chain becomes one real block chain. □




Figure 70: Primary pieces $\mathcal{G}_{\lambda_j}(T)$ give nilpotent chains, hence Jordan blocks.

**Corollary 6.2.3.** *Diagonalizability in Jordan Form*
*Let $n \in \mathbb{N}$, $A \in \mathrm{Mat}(n, n, \mathbb{C})$, $J = \bigoplus_{\alpha=1}^{s} J_{k_\alpha}(\mu_\alpha)$ be Jordan form of $A$.*
*Set $\delta_{\lambda\alpha} := \delta_{\lambda,\mu_\alpha}$ and $b_J(\lambda) = \sum_{\alpha=1}^{s} \delta_{\lambda\alpha}$ for $\lambda \in \mathrm{spec}(A)$. Then:*

*(i)* $\dim E_\lambda(A) = b_J(\lambda)$, $\lambda \in \mathrm{spec}(A)$;
*(ii)* *$A$ is diagonalizable if and only if every Jordan block in $J$ has size 1.*

*Proof.* Choose $S \in \mathrm{GL}_n(\mathbb{C})$ such that $A = SJS^{-1}$.

(i) For $\lambda \in \mathrm{spec}(A)$, $S : E_\lambda(J) \cong E_\lambda(A)$ and $\dim E_\lambda(J_{k_\alpha}(\mu_\alpha)) = \delta_{\lambda\alpha}$. Thus $\dim E_\lambda(A) = \sum_{\alpha=1}^{s} \delta_{\lambda\alpha} = b_J(\lambda)$.

(ii) Put $a_J(\lambda) = \sum_{\alpha=1}^{s} k_\alpha \delta_{\lambda\alpha}$. By Theorem 2.1.10 and (i), $A$ is diagonalizable $\iff a_J = b_J$ on $\mathrm{spec}(A)$.
Since $a_J(\lambda) - b_J(\lambda) = \sum_{\alpha=1}^{s} (k_\alpha - 1)\delta_{\lambda\alpha}$ and $k_\alpha \geq 1$, this is equivalent to $k_\alpha = 1$ for every $\alpha$.

*Structural Commentary*: Similarity transfers eigenspaces; block kernels count directions.
*Key Insight*: Algebraic multiplicity counts block lengths. □

**Lemma 6.2.4.** *Powers of a Nilpotent Jordan Block*
*Let $k \in \mathbb{N}$, let $p \in \mathbb{N}_0$, and set $N = J_k(0) \in \mathrm{Mat}(k, k, \mathbb{C})$. Then:*

*(i)* $\mathrm{rank}(N^p) = \max(k - p, 0)$;
*(ii)* $\dim \ker(N^p) = \min(p, k)$.

*Proof.* Let $(\mathbf{e}_j)_{j=1}^{k}$ be the standard basis. Set $(q)_+ := \max(q, 0)$ and $\mathcal{E}_m = \mathrm{span}(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ for $0 \leq m \leq k$, with $\mathcal{E}_0 = \{\mathbf{0}\}$. By $N\mathbf{e}_j = \mathbf{e}_{j-1}$ for $j > 1$ and $N\mathbf{e}_1 = \mathbf{0}$, $N^p\mathbf{e}_j = \mathbf{e}_{j-p}$ for $j > p$ and $\mathbf{0}$ otherwise.

(i) $\mathrm{im}(N^p) = \mathcal{E}_{(k-p)_+}$, hence $\mathrm{rank}(N^p) = \dim \mathcal{E}_{(k-p)_+} = (k-p)_+ = \max(k-p, 0)$.
(ii) By rank-nullity and (i), $\dim \ker(N^p) = k - (k-p)_+ = \min(p, k)$.

*Structural Commentary*: Powers shift the chain and truncate its span.
*Key Insight*: Rank counts surviving positions. □

**Definition 6.2.5.** *Weyr Characteristic*
Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ and let $\lambda \in \mathrm{spec}(A)$.

(i) For $p \in \mathbb{N}_0$, define the shifted-power rank $r_p(\lambda, A) := \mathrm{rank}\big((A - \lambda I_n)^p\big)$.
(ii) For $p \in \mathbb{N}$, define the Weyr number $w_p(\lambda, A) := r_{p-1}(\lambda, A) - r_p(\lambda, A)$. If $s := \max\{p \in \mathbb{N} : w_p(\lambda, A) > 0\}$, then the finite sequence $\big(w_1(\lambda, A), \ldots, w_s(\lambda, A)\big)$ is the *Weyr characteristic* of $A$ associated with $\lambda$.

**Theorem 6.2.6.** *Jordan Block Sizes from Rank Drops*
*Let $n \in \mathbb{N}$, $A \in \mathrm{Mat}(n, n, \mathbb{C})$, $\lambda \in \mathrm{spec}(A)$, and $J = \bigoplus_{\alpha=1}^{s} J_{k_\alpha}(\mu_\alpha)$ be a Jordan form of $A$.*
*For $q \in \mathbb{N}_0$, set $r_q = \mathrm{rank}\big((A - \lambda I_n)^q\big)$; for $q \in \mathbb{N}$, set $w_q = r_{q-1} - r_q$.*
*For $p \in \mathbb{N}$, set $I_p = \{\alpha : \mu_\alpha = \lambda,\ k_\alpha \geq p\}$. Then:*

*(i)* $w_p = \#I_p$;
*(ii)* $w_p - w_{p+1} = \#(I_p \setminus I_{p+1}) = \#\{\alpha : \mu_\alpha = \lambda,\ k_\alpha = p\}$.

*Proof.* By similarity-invariance of rank, assume $A = J$. Let $\mathbf{1}_E$ be the indicator of $E$.
Set $B_\alpha = J_{k_\alpha}(\mu_\alpha)$ and $\rho_q(B) = \mathrm{rank}\big((B - \lambda I)^q\big)$.
Put $\Delta_{p,\alpha} = \rho_{p-1}(B_\alpha) - \rho_p(B_\alpha)$. By rank additivity, $w_p = \sum_{\alpha=1}^{s} \Delta_{p,\alpha}$.

(i) By invertibility for $\mu_\alpha \neq \lambda$ and by Lemma 6.2.4 for $\mu_\alpha = \lambda$, $\Delta_{p,\alpha} = \mathbf{1}_{I_p}(\alpha)$. Hence $w_p = \sum_{\alpha=1}^{s} \mathbf{1}_{I_p}(\alpha) = \#I_p$.
(ii) By (i), $w_p - w_{p+1} = \#I_p - \#I_{p+1} = \#(I_p \setminus I_{p+1}) = \#\{\alpha : \mu_\alpha = \lambda,\ k_\alpha = p\}$.

*Structural Commentary*: Rank drops split into blockwise indicators.
*Key Insight*: A block contributes until its length. □




Figure 71: For $r_p = \operatorname{rank}((A-\lambda I)^p)$, drops count surviving chains; $w_p - w_{p+1}$ counts size-$p$ blocks.

**Corollary 6.2.7.** *Uniqueness of Jordan Form*
*Let $n \in \mathbb{N}$, $A \in \operatorname{Mat}(n,n,\mathbb{C})$. Let $J, J'$ be Jordan matrices similar to $A$.*
*For $M \in \{J, J'\}$, let $\mathfrak{B}(M)$ be its multiset of Jordan blocks, set $m_M(\lambda,k) = \#\{B \in \mathfrak{B}(M) : B = J_k(\lambda)\}$. Then:*
*(i) $m_J(\lambda,k) = m_{J'}(\lambda,k)$ for all $(\lambda,k) \in \operatorname{spec}(A) \times \mathbb{N}$;*
*(ii) $\exists$ a block-permutation matrix $P$ such that $J' = P^{-1}JP$.*

*Proof.* Set $d_k^\lambda(M) = w_k(\lambda, M) - w_{k+1}(\lambda, M)$. By similarity invariance, $d_k^\lambda(J) = d_k^\lambda(J')\ \forall (\lambda,k) \in \operatorname{spec}(A) \times \mathbb{N}$.
(i) For $(\lambda,k) \in \operatorname{spec}(A) \times \mathbb{N}$, by Theorem 6.2.6, $m_J(\lambda,k) = d_k^\lambda(J) = d_k^\lambda(J') = m_{J'}(\lambda,k)$.
(ii) By (i), $\mathfrak{B}(J) = \mathfrak{B}(J')$, so $\exists$ a block-permutation matrix $P$ with $J' = P^{-1}JP$.
*Structural Commentary*: Shifted ranks determine block multiplicities.
*Key Insight*: Weyr data fixes Jordan blocks. □

**Method 6.2.8.** *Computing Jordan Blocks by Rank Drops*
Let $A \in \operatorname{Mat}(n,n,\mathbb{C})$ and let $\lambda \in \operatorname{spec}(A)$. To determine the $\lambda$-Jordan blocks:
(i) Compute $r_p(\lambda, A) = \operatorname{rank}\big((A-\lambda I_n)^p\big)$ for $p = 0, 1, 2, \ldots$ until $r_{p-1}(\lambda, A) = r_p(\lambda, A)$.
(ii) Set $w_p(\lambda, A) = r_{p-1}(\lambda, A) - r_p(\lambda, A)$. This is the number of $\lambda$-blocks of size at least $p$.
(iii) For $p \in \mathbb{N}$, take $w_p(\lambda, A) - w_{p+1}(\lambda, A)$ copies of $J_p(\lambda)$.
(iv) Repeat (i)–(iii) as $\lambda$ ranges over $\operatorname{spec}(A)$, and take the direct sum of all resulting blocks.
The validity of steps (ii)–(iii) follows from Theorem 6.2.6.

## 6.3 Products AB and BA and Jordan Structure

**Proposition 6.3.1.** *AB and BA Have the Same Characteristic Polynomial*
*For $A, B \in \operatorname{Mat}(n,n,\mathbb{F})$, $p_{AB} = p_{BA}$.*
*If $A \in \operatorname{GL}(n,\mathbb{F})$ or $B \in \operatorname{GL}(n,\mathbb{F})$, then $AB \sim_{\mathbb{F}} BA$; otherwise not necessarily.*

*Proof.* In $\mathbb{F}(z)$, by Sylvester's identity $\det(zI - AB) = z^n \det(I - z^{-1}AB) = z^n \det(I - z^{-1}BA) = \det(zI - BA)$, hence $p_{AB} = p_{BA}$. If $A$ is invertible, $BA = A^{-1}(AB)A$; if $B$ is invertible, $AB = B^{-1}(BA)B$.
Without invertibility, take $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.
Then $AB = B$, $BA = 0$, $p_{AB} = p_{BA} = z^2$, but $\operatorname{rank}(AB) = 1 \neq 0 = \operatorname{rank}(BA)$, so $AB \not\sim_{\mathbb{F}} BA$.
*Structural Commentary*: Characteristic polynomials see less than similarity.
*Key Insight*: $AB$ and $BA$ share determinant data; invertibility yields similarity. □

**Theorem 6.3.2.** *Nonzero Jordan Blocks of AB and BA*
*Let $A, B \in \operatorname{Mat}(n,n,\mathbb{C})$. For every $\lambda \neq 0$, the Jordan blocks of $AB$ and $BA$ at $\lambda$ coincide.*

*Proof.* Set $X = \begin{bmatrix} AB & A \\ 0 & 0 \end{bmatrix}$ and $Y = \begin{bmatrix} 0 & A \\ 0 & BA \end{bmatrix}$. Then $\begin{bmatrix} I_n & 0 \\ B & I_n \end{bmatrix} X \begin{bmatrix} I_n & 0 \\ -B & I_n \end{bmatrix} = Y$, so $X$ and $Y$ are similar.
For $\lambda \neq 0$ and $p \geq 1$, the shifted powers have block triangular forms

$$(X - \lambda I_{2n})^p = \begin{bmatrix} (AB - \lambda I_n)^p & * \\ 0 & (-\lambda)^p I_n \end{bmatrix}, \qquad (Y - \lambda I_{2n})^p = \begin{bmatrix} (-\lambda)^p I_n & * \\ 0 & (BA - \lambda I_n)^p \end{bmatrix}$$

The diagonal block $(-\lambda)^p I_n$ is invertible.
Block Gaussian elimination by invertible row and column operations removes the starred block, so

$$\operatorname{rank}\big((X - \lambda I_{2n})^p\big) = n + \operatorname{rank}\big((AB - \lambda I_n)^p\big),$$
$$\operatorname{rank}\big((Y - \lambda I_{2n})^p\big) = n + \operatorname{rank}\big((BA - \lambda I_n)^p\big).$$

Since $X$ and $Y$ are similar, these left sides agree for all $p$.
Hence the shifted rank drops of $AB$ and $BA$ at $\lambda$ agree. By Theorem 6.2.6, their Jordan blocks at $\lambda$ coincide.
*Structural Commentary*: Similar block embeddings make the nonzero shifted ranks match.
*Key Insight*: At $\lambda \neq 0$, the extra block is invertible and splits off. □




Figure 72: For $\lambda \neq 0$, the auxiliary similarity reduces the comparison to shifted ranks of $AB$ and $BA$.

**Corollary 6.3.3.** *Similarity Criterion for AB and BA*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{C})$. $AB$ is similar to $BA$ if and only if* $\operatorname{rank}\big((AB)^p\big) = \operatorname{rank}\big((BA)^p\big)$, $p = 1, \ldots, n$.

*Proof.*
(⇒) If $AB \sim BA$, then $(AB)^p \sim (BA)^p$ for all $p$, so their ranks are equal.
(⇐) Assume the rank equalities hold for $p = 1, \ldots, n$. By Theorem 6.3.2, the nonzero Jordan blocks coincide. Together with the equality at $p = 0$, the rank equalities give the same Weyr characteristic at $\lambda = 0$. By Theorem 6.2.6, the nilpotent blocks also agree, so the Jordan forms coincide up to block order.

*Structural Commentary*: Nonzero blocks agree automatically; only the nilpotent blocks must be tested.
*Key Insight*: Powers of $AB$ and $BA$ measure exactly the zero-Jordan-block data. □

**Method 6.3.4.** *Checking Whether AB and BA Are Similar*
For $A, B \in \mathrm{Mat}(n, n, \mathbb{C})$:

(i) The nonzero Jordan blocks of $AB$ and $BA$ already match.
(ii) Compute $\operatorname{rank}\big((AB)^p\big)$ and $\operatorname{rank}\big((BA)^p\big)$ for $p = 1, \ldots, n$.
(iii) If all ranks agree, then $AB \sim BA$; if some rank differs, then only the nilpotent Jordan structure differs.

Step (i) follows from Theorem 6.3.2, and step (iii) is Corollary 6.3.3.

## 6.4 Nonderogatory Matrices and Commutants

**Definition 6.4.1.** *Nonderogatory Matrix*
A matrix $A \in \mathrm{Mat}(n, n, \mathbb{C})$ is *nonderogatory* if its Jordan form has exactly one Jordan block for each distinct eigenvalue. Equivalently, $\dim E_\lambda(A) = 1$ for every $\lambda \in \operatorname{spec}(A)$. The equivalence follows from Corollary 6.2.3.

**Lemma 6.4.2.** *Commutant of One Jordan Block*
*If $J = J_m(\lambda)$ and $XJ = JX$, then $X = p(J)$ for some polynomial $p$ with $\deg p < m$.*

*Proof.* It suffices to treat $N = J_m(0)$, since $J_m(\lambda) = \lambda I_m + N$.
The vector $\mathbf{e}_m$ is cyclic for $N$: $(\mathbf{e}_m, N\mathbf{e}_m, \ldots, N^{m-1}\mathbf{e}_m)$ is the standard basis in reverse order.
Choose the unique polynomial $p$ with $\deg p < m$ such that $p(N)\mathbf{e}_m = X\mathbf{e}_m$.
For every $r = 0, \ldots, m-1$, $p(N)N^r\mathbf{e}_m = N^r p(N)\mathbf{e}_m = N^r X\mathbf{e}_m = XN^r\mathbf{e}_m$.
Thus $p(N)$ and $X$ agree on a basis, so $X = p(N) = q(J)$ for $q(z) = p(z - \lambda)$.
*Structural Commentary*: A single Jordan block is generated by one cyclic vector.
*Key Insight*: The cyclic vector determines the commutant. □

**Theorem 6.4.3.** *Nonderogatory Commutants Are Polynomials*
*If $A \in \mathrm{Mat}(n, n, \mathbb{C})$ is nonderogatory, then* $\operatorname{Comm}(A) = \{p(A) : p \in \mathbb{C}[z]\}$.

*Proof.* The inclusion $\{p(A) : p \in \mathbb{C}[z]\} \subseteq \operatorname{Comm}(A)$ follows from Proposition 1.1.5.
For reverse inclusion, by Theorem 6.2.1, $A = SJS^{-1}$ for a Jordan matrix $J = J_{m_1}(\lambda_1) \oplus \cdots \oplus J_{m_d}(\lambda_d)$ with distinct $\lambda_i$. Conjugation identifies $\operatorname{Comm}(A)$ with $\operatorname{Comm}(J)$ and satisfies $p(A) = Sp(J)S^{-1}$, so it suffices to prove the claim for $J$.
If $XJ = JX$ and $X = [X_{ij}]$ is partitioned conformally with $J$, the block equation is $X_{ij}J_{m_j}(\lambda_j) = J_{m_i}(\lambda_i)X_{ij}$.
By Corollary 3.5.5, $X_{ij} = 0$ for $i \neq j$. Each diagonal block $X_{ii}$ commutes with $J_{m_i}(\lambda_i)$, so by Lemma 6.4.2, $X_{ii} = p_i(J_{m_i}(\lambda_i))$ for some polynomial $p_i$ with $\deg p_i < m_i$. The ideals generated by $(z - \lambda_i)^{m_i}$ are pairwise coprime, so Hermite interpolation gives a polynomial $p$ with $p \equiv p_i \pmod{(z - \lambda_i)^{m_i}}$ for every $i$.
On the $i$th block, $p - p_i = (z - \lambda_i)^{m_i} q_i$ for some $q_i$, and $(J_{m_i}(\lambda_i) - \lambda_i I)^{m_i} = 0$. Hence $p(J_{m_i}(\lambda_i)) = p_i(J_{m_i}(\lambda_i)) = X_{ii}$ for every $i$, so $X = p(J)$. Conjugating back to $A$ proves the claim.

*Structural Commentary*: Disjoint eigenvalue blocks force block diagonality; one Jordan block per eigenvalue makes each diagonal commutant polynomial.
*Key Insight*: Hermite interpolation glues the separate block polynomials into one polynomial in $A$. □




Figure 73: Blockwise commutant polynomials glue by interpolation.

**Corollary 6.4.4.** *Simple Spectrum Commutants Are Polynomials*
*If $A \in \mathrm{Mat}(n,n,\mathbb{C})$ has $n$ distinct eigenvalues, then* $\mathrm{Comm}(A) = \{p(A) : p \in \mathbb{C}[z]\}$, $\dim \mathrm{Comm}(A) = n$.

*Proof.* Write $A = S\,\mathrm{diag}(\lambda_1,\dots,\lambda_n)S^{-1}$ with distinct $\lambda_i$. By Theorem 5.4.12, every $B \in \mathrm{Comm}(A)$ has the form $S\,\mathrm{diag}(\xi_1,\dots,\xi_n)S^{-1}$. Choose the Lagrange interpolation polynomial $p(z) = \sum_{i=1}^{n} \xi_i \prod_{j\neq i} \frac{z-\lambda_j}{\lambda_i-\lambda_j}$.

Then $p(\lambda_i) = \xi_i$ for every $i$. By Theorem 1.2.1, $B = p(A)$.
The dimension statement is the case $n_1 = \cdots = n_n = 1$ of Theorem 5.4.12.
*Structural Commentary*: Simple spectrum leaves one scalar choice per eigendirection.
*Key Insight*: Interpolation converts arbitrary diagonal data into a polynomial in $A$. □

## 6.5 Jordan Operator Functions and Square Roots

**Proposition 6.5.1.** *Exponential of a Jordan Block*
*For $k \in \mathbb{N}$ with $k \geq 1$, $\lambda \in \mathbb{C}$, and $t \in \mathbb{C}$, one has* $e^{tJ_k(\lambda)} = e^{\lambda t} \sum_{r=0}^{k-1} \frac{t^r}{r!} J_k(0)^r$.

*Proof.* Set $N = J_k(0)$, so $J_k(\lambda) = \lambda I_k + N$ and $N^k = 0$.
By binomial theorem, $(\lambda I_k + N)^m = \sum_{r=0}^{\min(m,k-1)} \binom{m}{r} \lambda^{m-r} N^r$.

Absolute convergence permits regrouping by powers of $N$. Thus $e^{tJ_k(\lambda)} = \sum_{r=0}^{k-1} \frac{t^r}{r!} N^r \sum_{s=0}^{\infty} \frac{(\lambda t)^s}{s!} = e^{\lambda t} \sum_{r=0}^{k-1} \frac{t^r}{r!} N^r$.
*Structural Commentary*: Expand the scalar and nilpotent parts together, then regroup by nilpotent degree.
*Key Insight*: Nilpotence truncates the matrix directions while the scalar series remains exponential. □

**Proposition 6.5.2.** *Commuting and Similarity Laws for the Exponential*
*Let $A, B \in \mathrm{Mat}(n,n,\mathbb{C})$ and $S \in GL_n(\mathbb{C})$. Then:*
*(i)* $AB = BA \implies e^{A+B} = e^A e^B$.
*(ii)* $e^{SAS^{-1}} = Se^A S^{-1}$.

*Proof.*

(i) Since $A$ and $B$ commute, $(A+B)^m = \sum_{r=0}^{m} \binom{m}{r} A^r B^{m-r}$.

Absolute convergence permits the Cauchy product, $e^A e^B = \sum_{m=0}^{\infty} \frac{1}{m!} \sum_{r=0}^{m} \binom{m}{r} A^r B^{m-r} = e^{A+B}$.

(ii) Put $X = SAS^{-1}$ and $H_m \colon X^m = SA^m S^{-1}$. Then $H_0$ is $I_n = SI_nS^{-1}$.
If $H_m$ holds, then $X^{m+1} = SA^m S^{-1} SAS^{-1} = SA^{m+1}S^{-1}$, so $H_{m+1}$ holds.

By induction, $H_m$ holds for all $m \in \mathbb{N}_0$, and $e^X = S\left(\sum_{m=0}^{\infty} \frac{A^m}{m!}\right) S^{-1} = Se^A S^{-1}$.

*Structural Commentary*: Apply power-series algebra only after verifying the required matrix identities.
*Key Insight*: Commutativity enables products; similarity passes through powers. □

**Theorem 6.5.3.** *Analytic Power Series on Jordan Blocks*
*Let $n \in \mathbb{N}$, $A \in \mathrm{Mat}(n,n,\mathbb{C})$, and $f(z) = \sum_{m=0}^{\infty} c_m z^m$ have radius $R > 0$ with $\rho(A) < R$.*
*Set $D_R := \{z \in \mathbb{C} : |z| < R\}$ and $N_k := J_k(0)$. For $(k,\lambda) \in \mathbb{N} \times D_R$, set $F_k(\lambda) := \sum_{r=0}^{k-1} \frac{f^{(r)}(\lambda)}{r!} N_k^r$.*
*For $J = \bigoplus_{\alpha=1}^{s} J_{k_\alpha}(\lambda_\alpha)$ with $\lambda_\alpha \in D_R$, set $f(J) := \bigoplus_{\alpha=1}^{s} F_{k_\alpha}(\lambda_\alpha)$. Then:*

*(i)* $\sum_{m=0}^{\infty} c_m J_k(\lambda)^m$ *converges to* $F_k(\lambda)$ *for every* $(k,\lambda) \in \mathbb{N} \times D_R$.

*(ii)* *The series* $f(A) := \sum_{m=0}^{\infty} c_m A^m$ *converges.*

*(iii)* *If* $A = SJS^{-1}$ *is a Jordan decomposition, then* $f(A) = Sf(J)S^{-1}$.

*Proof.* For $M \in \mathbb{N}_0$, set $P_M(X) := \sum_{m=0}^{M} c_m X^m$, and use $\binom{m}{r} = 0$ for $r > m$.

(i) Fix $(k,\lambda) \in \mathbb{N} \times D_R$ and set $N = N_k$. Since $N^k = 0$, $J_k(\lambda)^m = \sum_{r=0}^{k-1} \binom{m}{r} \lambda^{m-r} N^r$ for $m \in \mathbb{N}_0$.
For $0 \le r < k$, $\sum_{m=r}^{\infty} c_m \binom{m}{r} \lambda^{m-r} = \frac{f^{(r)}(\lambda)}{r!}$ by termwise differentiation.
Absolute convergence permits regrouping over $0 \le r < k$, so $P_M(J_k(\lambda)) \to F_k(\lambda)$.

(ii) By Theorem 6.2.1, $A = SJS^{-1}$ with $J = \bigoplus_{\alpha=1}^{s} J_{k_\alpha}(\lambda_\alpha)$. Since $\lambda_\alpha \in \mathrm{spec}(A)$ and $\rho(A) < R$, $\lambda_\alpha \in D_R$.
By (i), $P_M(J) \to f(J)$, and $P_M(A) = SP_M(J)S^{-1} \to Sf(J)S^{-1}$.

(iii) By (ii), $f(A) = \lim_{M\to\infty} P_M(A) = S\left(\lim_{M\to\infty} P_M(J)\right) S^{-1} = Sf(J)S^{-1}$.

*Structural Commentary*: Blockwise binomial expansion reduces $f(J_k(\lambda))$ to nilpotent powers.
*Key Insight*: Derivatives supply the chain coefficients. □

**Example 6.5.4.** *Closed Form for a $2 \times 2$ Exponential*
Let $A = \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)$, and set $\delta := \frac{a+d}{2}$ and $B := A - \delta I_2$. Choose $\Delta \in \mathbb{C}$ with $\Delta^2 = \left(\frac{a-d}{2}\right)^2 + bc$.
By Theorem 3.1.1, $B^2 = \Delta^2 I_2$. Since $A = \delta I_2 + B$, by Proposition 6.5.2(i) and even–odd decomposition of exponential series, $e^A = e^{\delta}\left(\cosh(\Delta) I_2 + \frac{\sinh(\Delta)}{\Delta} B\right)$. At $\Delta = 0$, $\frac{\sinh(\Delta)}{\Delta}$ is interpreted by continuity as 1.

**Theorem 6.5.5.** *Spectral Mapping and Jordan Structure of the Exponential*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$, let $A = SJS^{-1}$ be a Jordan decomposition, and set $E := \exp(A)$.*
*Write $J = \bigoplus_{\alpha=1}^{s} B_\alpha$, $B_\alpha = J_{k_\alpha}(\lambda_\alpha)$. For $\rho \in \mathbb{C}$, set $I_\rho := \{\alpha : e^{\lambda_\alpha} = \rho\}$, $\Lambda_\rho(A) := \{\lambda \in \mathrm{spec}(A) : e^{\lambda} = \rho\}$.*
*Let $\mathcal{S}_\rho(M)$ denote the multiset of $\rho$-Jordan block sizes of $M$. Then:*

*(i)* $\mathrm{spec}(E) = \{e^{\lambda} : \lambda \in \mathrm{spec}(A)\}$.

*(ii)* $\forall \rho \in \mathrm{spec}(E)$, $\mathcal{G}_\rho(E) = \bigoplus_{\lambda \in \Lambda_\rho(A)} \mathcal{G}_\lambda(A)$.

*(iii)* $\mathcal{S}_\rho(E) = \{k_\alpha : \alpha \in I_\rho\}$ *for every* $\rho \in \mathrm{spec}(E)$.

*(iv)* $\det(E) = e^{\mathrm{tr}(A)}$.

*Proof.* Set $\sigma(M) := \mathrm{spec}(M)$, $E_J := e^J$, $E_\alpha := e^{B_\alpha}$, and $\tau := \sum_{\alpha=1}^{s} k_\alpha \lambda_\alpha = \mathrm{tr}(A)$.
By Proposition 6.5.2(ii), $E = SE_JS^{-1}$; work with $J$. Let $V_\alpha$ be coordinate space of $B_\alpha$; $K_\rho := \bigoplus_{\alpha \in I_\rho} V_\alpha$.
Write $B_\alpha = \lambda_\alpha I_{k_\alpha} + N_\alpha$, $q_\alpha(z) := \sum_{r=0}^{k_\alpha - 1} \frac{z^r}{(r+1)!}$, $Q_\alpha := q_\alpha(N_\alpha)$. By Proposition 6.5.1, $E_\alpha - e^{\lambda_\alpha} I_{k_\alpha} = e^{\lambda_\alpha} N_\alpha Q_\alpha$.
With $M_\alpha := Q_\alpha - I_{k_\alpha} \in N_\alpha \mathbb{C}[N_\alpha]$, $M_\alpha^{k_\alpha} = 0$ and $Q_\alpha^{-1} = \sum_{j=0}^{k_\alpha - 1} (-M_\alpha)^j$. Also $Q_\alpha N_\alpha = N_\alpha Q_\alpha$.

(i) $\mathrm{diag}(E_\alpha) = e^{\lambda_\alpha} I_{k_\alpha}$, so $\sigma(E_\alpha) = \{e^{\lambda_\alpha}\}$.
By $E = SE_JS^{-1}$ and $E_J = \bigoplus_{\alpha=1}^{s} E_\alpha$, $\sigma(E) = \{e^{\lambda_\alpha} : 1 \le \alpha \le s\} = \{e^{\lambda} : \lambda \in \sigma(A)\}$.

(ii) If $\alpha \in I_\rho$, then $(E_\alpha - \rho I_{k_\alpha})^r = \rho^r N_\alpha^r Q_\alpha^r$ for $r \in \mathbb{N}$, so $\mathcal{G}_\rho(E_J|_{V_\alpha}) = V_\alpha$.
If $\alpha \notin I_\rho$, then $E_\alpha - \rho I_{k_\alpha} \in GL_{k_\alpha}(\mathbb{C})$. Thus $\mathcal{G}_\rho(E) = S\mathcal{G}_\rho(E_J) = SK_\rho = \bigoplus_{\lambda \in \Lambda_\rho(A)} \mathcal{G}_\lambda(A)$.

(iii) For $\alpha \in I_\rho$ and $r \in \mathbb{N}$, $\mathrm{rank}((E_\alpha - \rho I_{k_\alpha})^r) = \mathrm{rank}(N_\alpha^r)$. By Theorem 6.2.6, $\mathcal{S}_\rho(E_\alpha) = \{k_\alpha\}$.
For $\alpha \notin I_\rho$, $\mathcal{S}_\rho(E_\alpha) = \varnothing$. By similarity and direct sums, (iii) follows.

(iv) $\det(E) = \det(E_J) = \prod_{\alpha=1}^{s} e^{k_\alpha \lambda_\alpha} = e^{\tau} = e^{\mathrm{tr}(A)}$.

*Structural Commentary*: Exponentials merge block labels with equal $e^\lambda$.
*Key Insight*: $Q_\alpha$ is invertible, so nilpotent ranks are unchanged. □

**Definition 6.5.6.** *Matrix Logarithm*
Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$. A matrix $B \in \mathrm{Mat}(n,n,\mathbb{C})$ is a *matrix logarithm* of $A$ if $e^B = A$.

**Theorem 6.5.7.** *Existence of Complex Matrix Logarithms*
*Let $n \in \mathbb{N}$ and $A \in \mathrm{Mat}(n,n,\mathbb{C})$. Then $\exists B \in \mathrm{Mat}(n,n,\mathbb{C})$ with $e^B = A$ iff $A \in GL_n(\mathbb{C})$.*

*Proof.*
($\Rightarrow$) Suppose $A = e^B$. Since $B$ and $-B$ commute, by Proposition 6.5.2(i), $Ae^{-B} = I_n = e^{-B}A$; $A \in GL_n(\mathbb{C})$.
($\Leftarrow$) Suppose $A \in GL_n(\mathbb{C})$. Choose $A = SJS^{-1}$ with $J = \bigoplus_{\alpha=1}^{s} J_{k_\alpha}(\lambda_\alpha)$. Then $\lambda_\alpha \neq 0$ for all $\alpha$.

Choose $\ell_\alpha \in \mathbb{C}$, $e^{\ell_\alpha} = \lambda_\alpha$. Set $N_\alpha := J_{k_\alpha}(0)$, $X_\alpha := \lambda_\alpha^{-1} N_\alpha$. Set $Y_\alpha := \sum_{r=1}^{k_\alpha - 1} \frac{(-1)^{r+1}}{r} X_\alpha^r$, $L_\alpha := \ell_\alpha I_{k_\alpha} + Y_\alpha$.
Since $X_\alpha^{k_\alpha} = 0$, by $e^{\log(1+x)} = 1 + x$ in $\mathbb{C}[x]/(x^{k_\alpha})$, $e^{Y_\alpha} = I_{k_\alpha} + X_\alpha$.
Since $\ell_\alpha I_{k_\alpha}$ commutes with $Y_\alpha$, by Proposition 6.5.2(i), $e^{L_\alpha} = e^{\ell_\alpha}(I_{k_\alpha} + X_\alpha) = J_{k_\alpha}(\lambda_\alpha)$.
Set $L_J := \bigoplus_{\alpha=1}^{s} L_\alpha$ and $B := SL_JS^{-1}$. By Proposition 6.5.2(ii), $e^B = Se^{L_J}S^{-1} = SJS^{-1} = A$.

*Structural Commentary*: Combine scalar logs with nilpotent log polynomials.
*Key Insight*: Nonzero block eigenvalues permit logarithms. □

**Perspective 6.5.8.** *Real Matrix Logarithms*
For $A \in \mathrm{Mat}(n,n,\mathbb{C})$, by Theorem 6.5.7, $A = e^B$ for some complex $B$ iff $A \in GL_n(\mathbb{C})$.
Over $\mathbb{R}$, $A \in GL_n(\mathbb{R})$ is not sufficient.
If $A \in GL_n(\mathbb{R})$ and $\mathrm{spec}(A) \cap \mathbb{R} \subset (0,\infty)$, choose $\ell_\lambda = \log\lambda \in \mathbb{R}$ for real $\lambda$ and $\ell_{\overline{\lambda}} = \overline{\ell_\lambda}$ for nonreal pairs.
By conjugate pairing of real Jordan blocks, the block logarithms assemble to $B \in \mathrm{Mat}(n,n,\mathbb{R})$ with $e^B = A$.
For $A \in GL_n(\mathbb{R})$, $A^2$ has a real logarithm: $\mu < 0$ and $\mu \in \mathrm{spec}(A^2)$ imply $\{\lambda, \overline{\lambda}\} = \{\pm i\sqrt{-\mu}\}$ for some $\lambda \in \mathrm{spec}(A)$. By conjugate pairing, the negative real blocks of $A^2$ occur in pairs.
These are sufficient facts, not the full real criterion.

**Proposition 6.5.9.** *Jordan Resolvent and Primary Spectral Projectors*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$, set $\Sigma := \mathrm{spec}(A)$, and set $G_\nu := \mathcal{G}_\nu(A)$.*
*For $m \in \mathbb{N}$, $M \in \mathrm{Mat}(m,m,\mathbb{C})$, and $z \notin \mathrm{spec}(M)$, set $R_M(z) := (zI_m - M)^{-1}$.*
*By primary decomposition, $\mathbb{C}^n = \bigoplus_{\nu\in\Sigma} G_\nu$. For $\mu \in \Sigma$, define $P_\mu$ by $P_\mu|_{G_\nu} = \delta_{\mu\nu} I_{G_\nu}$.*
*For meromorphic $F$ near $\mu$, let $[F]_{\mu,-1}$ be its $(z-\mu)^{-1}$ coefficient. Then:*

*(i) For $k \in \mathbb{N}$, $\lambda \in \mathbb{C}$, $B = J_k(\lambda) = \lambda I_k + N$, and $z \neq \lambda$, $R_B(z) = \sum_{r=0}^{k-1} \frac{N^r}{(z-\lambda)^{r+1}}$.*
*(ii) $[R_A]_{\mu,-1} = P_\mu$ for every $\mu \in \Sigma$.*
*(iii) $P_\lambda P_\mu = \delta_{\lambda\mu} P_\mu$ for all $\lambda, \mu \in \Sigma$, and $AP_\mu = P_\mu A$ for all $\mu \in \Sigma$.*
*(iv) $\sum_{\mu\in\Sigma} P_\mu = I_n$.*

*Proof.*
(i) Let $B = J_k(\lambda) = \lambda I_k + N$ and set $a := (z-\lambda)^{-1}$.

By $zI_k - B = (z-\lambda)(I_k - aN)$, $N^k = 0$, and the finite geometric identity, $(I_k - aN)^{-1} = \sum_{r=0}^{k-1} a^r N^r$.

By $R_B(z) = a(I_k - aN)^{-1}$, $R_B(z) = a\sum_{r=0}^{k-1} a^r N^r = \sum_{r=0}^{k-1} \frac{N^r}{(z-\lambda)^{r+1}}$.

(ii) Choose $A = SJS^{-1}$ by Theorem 6.2.1, with $J = \bigoplus_{\alpha=1}^{s} J_{k_\alpha}(\lambda_\alpha)$. By similarity, $R_A(z) = SR_J(z)S^{-1}$.

Set $H_\nu := \mathcal{G}_\nu(J)$ and $C_\mu := [R_J]_{\mu,-1}$. By (i), $C_\mu|_{H_\nu} = \delta_{\mu\nu} I_{H_\nu}$. By conjugation, $[R_A]_{\mu,-1} = SC_\mu S^{-1} = P_\mu$.
(iii) Let $v = \sum_{\nu\in\Sigma} v_\nu$, $v_\nu \in G_\nu$. Then $P_\mu v = v_\mu$, $P_\lambda P_\mu v = \delta_{\lambda\mu} v_\mu = \delta_{\lambda\mu} P_\mu v$; $AG_\nu \subseteq G_\nu$, $AP_\mu v = Av_\mu = P_\mu A v$.

(iv) For $v = \sum_{\nu\in\Sigma} v_\nu$ as in (iii), $\left(\sum_{\mu\in\Sigma} P_\mu\right) v = \sum_{\mu\in\Sigma} v_\mu = v$. Since $v$ is arbitrary, (iv) follows.

*Structural Commentary*: Laurent coefficients isolate primary summands.
*Key Insight*: The simple-pole coefficient is the primary projection. □

**Definition 6.5.10.** *Contracting, Neutral, and Expanding Primary Subspaces*
Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$, and let $P_\lambda$ be the primary projection onto $\mathcal{G}_\lambda(A)$.
Set $\Sigma_- := \{\lambda \in \mathrm{spec}(A) : |\lambda| < 1\}$, $\Sigma_0 := \{\lambda \in \mathrm{spec}(A) : |\lambda| = 1\}$, and $\Sigma_+ := \{\lambda \in \mathrm{spec}(A) : |\lambda| > 1\}$. Define:

(i) $\mathcal{G}^-(A) := \bigoplus_{\lambda \in \Sigma_-} \mathcal{G}_\lambda(A)$ and $P_A^- := \sum_{\lambda \in \Sigma_-} P_\lambda$, the *contracting* part.

(ii) $\mathcal{G}^0(A) := \bigoplus_{\lambda \in \Sigma_0} \mathcal{G}_\lambda(A)$ and $P_A^0 := \sum_{\lambda \in \Sigma_0} P_\lambda$, the *neutral* part.

(iii) $\mathcal{G}^+(A) := \bigoplus_{\lambda \in \Sigma_+} \mathcal{G}_\lambda(A)$ and $P_A^+ := \sum_{\lambda \in \Sigma_+} P_\lambda$, the *expanding* part.

By primary decomposition, $\mathbb{C}^n = \mathcal{G}^-(A) \oplus \mathcal{G}^0(A) \oplus \mathcal{G}^+(A)$, and Proposition 6.5.9 identifies $P_A^-$, $P_A^0$, and $P_A^+$ as the corresponding projections.

**Lemma 6.5.11.** *Identity Plus Nilpotent Has a Square Root*
*Let $\mathcal{V}$ be a complex vector space and let $N \in \mathcal{L}(\mathcal{V})$ be nilpotent. Then $\exists\, R \in \mathcal{L}(\mathcal{V})$ such that $R^2 = I_\mathcal{V} + N$.*

*Proof.* Choose $m \in \mathbb{N}$ with $N^m = 0$. Set $a_0 = 1$. For $1 \le k < m$, set $s_k := \sum_{r=1}^{k-1} a_r a_{k-r}$ and $a_k := \frac{1}{2}(\delta_{k1} - s_k)$.
Set $R := \sum_{k=0}^{m-1} a_k N^k$. For $0 \le k < m$, set $c_k := \sum_{r=0}^{k} a_r a_{k-r}$. By the recursion, $c_0 = 1$, $c_1 = 1$ if $m \ge 2$, and $c_k = 0$ for $2 \le k < m$.
By $N^m = 0$, $R^2 = \sum_{k=0}^{m-1} c_k N^k = I_\mathcal{V} + N$.
*Structural Commentary*: Coefficient matching truncates at nilpotency.
*Key Insight*: Choose $a_k$ so $R^2$ has only degrees 0 and 1. □

Figure 74: Nilpotence truncates the square-root series; coefficients are chosen so $R^2 = I + N$.

**Theorem 6.5.12.** *Invertible Complex Operators Have Square Roots*
*Let $\mathcal{V}$ be finite-dimensional over $\mathbb{C}$ and let $T \in \mathcal{L}(\mathcal{V})$ be invertible. Then $\exists\, R \in \mathcal{L}(\mathcal{V})$ such that $R^2 = T$.*

*Proof.* If $\mathcal{V} = \{\mathbf{0}\}$, take $R = I_\mathcal{V}$. Assume $\mathcal{V} \neq \{\mathbf{0}\}$.
Let $\{\lambda_1, \ldots, \lambda_m\} = \mathrm{spec}(T)$ and set $\mathcal{U}_j := \mathcal{G}_{\lambda_j}(T)$.
By Theorem 5.1.1, $\mathcal{V} = \bigoplus_{j=1}^{m} \mathcal{U}_j$. By $T \in GL(\mathcal{V})$, $\lambda_j \neq 0$.
Set $N_j := (T - \lambda_j I_\mathcal{V})|_{\mathcal{U}_j}$ and $M_j := \lambda_j^{-1} N_j$. Then $M_j$ is nilpotent and $T|_{\mathcal{U}_j} = \lambda_j(I_{\mathcal{U}_j} + M_j)$.
By Lemma 6.5.11, choose $B_j \in \mathcal{L}(\mathcal{U}_j)$ with $B_j^2 = I_{\mathcal{U}_j} + M_j$.
Choose $\alpha_j \in \mathbb{C}$ with $\alpha_j^2 = \lambda_j$ and set $R_j := \alpha_j B_j$.
By these choices, $R_j^2 = \lambda_j B_j^2 = \lambda_j(I_{\mathcal{U}_j} + M_j) = T|_{\mathcal{U}_j}$.
Define $R := \bigoplus_{j=1}^{m} R_j$.
By $(R^2)|_{\mathcal{U}_j} = R_j^2 = T|_{\mathcal{U}_j}$ for every $j$, $R^2 = T$.
*Structural Commentary*: Square-root each primary restriction, then sum.
*Key Insight*: Nonzero eigenvalues separate scalar roots from nilpotent roots. □

# 7 Transpose and Conjugate Similarity

## 7.1 Transpose Similarity and Conjugate Similarity

**Theorem 7.1.1.** *Conjugate Similarity Criterion*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$. Then $A$ is similar to $\overline{A}$ if and only if each $J_k(\lambda)$ occurs in the Jordan form of $A$ with the same multiplicity as $J_k(\overline{\lambda})$.*

*Proof.* By Theorem 6.2.1, choose a Jordan form $J$ of $A$. Then $\overline{J}$ is a Jordan form of $\overline{A}$, obtained by replacing each $J_k(\lambda)$ by $J_k(\overline{\lambda})$.

($\Rightarrow$) If $A$ is similar to $\overline{A}$, then $J$ and $\overline{J}$ are similar.
By Corollary 6.2.7, their Jordan blocks have the same multiplicities.

($\Leftarrow$) If the multiplicities agree, then $J$ and $\overline{J}$ differ only by a permutation of blocks. Hence $A$ is similar to $\overline{A}$.

*Structural Commentary*: Conjugation changes eigenvalue labels but not block sizes.
*Key Insight*: Non-real Jordan blocks must occur in conjugate pairs. □

**Definition 7.1.2.** *Reversal Matrix*
For $k \geq 1$, define the *reversal matrix* $R_k \in \mathrm{Mat}(k, k, \mathbb{C})$ by $(R_k)_{ij} = \begin{cases} 1, & i+j = k+1, \\ 0, & i+j \neq k+1 \end{cases}$. Then $R_k^t = R_k$ and $R_k^2 = I_k$.

**Lemma 7.1.3.** *Reversal Conjugates a Jordan Block to Its Transpose*
*For every $k \geq 1$ and $\lambda \in \mathbb{C}$, $J_k(\lambda)^t = R_k J_k(\lambda) R_k$. Moreover, $R_k J_k(\lambda)$ and $J_k(\lambda) R_k$ are symmetric.*

*Proof.* Let $N = J_k(0)$. Since $R_k$ reverses standard basis order, $R_k N R_k = N^t$.
Adding scalar gives $R_k J_k(\lambda) R_k = J_k(\lambda)^t$. As $R_k = R_k^t = R_k^{-1}$, this identity gives $J_k(\lambda)^t R_k = R_k J_k(\lambda)$ and $R_k J_k(\lambda)^t = J_k(\lambda) R_k$, so $R_k J_k(\lambda)$ and $J_k(\lambda) R_k$ are symmetric.
*Structural Commentary*: Reversal changes the superdiagonal shift into the subdiagonal shift.
*Key Insight*: Transposition reads the same Jordan chain backwards. □

**Theorem 7.1.4.** *Transpose Similarity and Symmetric Factorisations*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$. Then:*

*(i) $A$ is similar to $A^t$ via a symmetric similarity matrix.*
*(ii) $A = BC = DE$, where $B, C, D, E$ are symmetric and $B, E$ are invertible.*

*Proof.* By Theorem 6.2.1, write $A = SJS^{-1}$ for a Jordan matrix $J$, and let $R$ be the direct sum of the reversal matrices matching the Jordan blocks of $J$. By Lemma 7.1.3, $J^t = RJR$, $R^t = R$, $R^2 = I_n$, and both $RJ$ and $JR$ are symmetric.

(i) Set $P = (S^{-1})^t R S^{-1}$. Then $P^t = P$, $P^{-1} = SRS^t$, and $PAP^{-1} = (S^{-1})^t RJRS^t = (S^{-1})^t J^t S^t = A^t$.
(ii) Using $R^2 = I_n$, $A = (SRS^t)\big((S^{-1})^t RJS^{-1}\big)$ and $A = (SJRS^t)\big((S^{-1})^t RS^{-1}\big)$.
The products equal $SJS^{-1}$ because $S^t(S^{-1})^t = I_n$ and $R^2 = I_n$. All four factors are symmetric because $R, RJ, JR$ are symmetric; $SRS^t$ and $(S^{-1})^t RS^{-1}$ are invertible.

*Structural Commentary*: Reduce to Jordan blocks, reverse chains, then conjugate back.
*Key Insight*: Symmetric involutions produce both transpose similarity and symmetric factors. □




Figure 75: Reversal gives $R_k J_k(0) R_k = J_k(0)^t$: the same chain read backwards.

**Corollary 7.1.5.** *Hermitian Products and Conjugate Products*
*Let $A, H, K \in \mathrm{Mat}(n, n, \mathbb{C})$, and suppose $H$ and $K$ are Hermitian.*
*Then $HK$ is similar to $KH$, and $A\overline{A}$ is similar to $\overline{A}A$.*

*Proof.* For each $p = 1, \ldots, n$, adjoints preserve rank, and $((HK)^p)^* = (KH)^p$. Hence $\mathrm{rank}\big((HK)^p\big) = \mathrm{rank}\big((KH)^p\big)$.
Similarly, entrywise conjugation preserves rank and gives $\overline{(A\overline{A})^p} = (\overline{A}A)^p$, so $\mathrm{rank}\big((A\overline{A})^p\big) = \mathrm{rank}\big((\overline{A}A)^p\big)$.
Both similarities follow from Corollary 6.3.3.
*Structural Commentary*: Adjoint and conjugation preserve rank and reverse product order.
*Key Insight*: The rank-power test turns order reversal into similarity. □

**Theorem 7.1.6.** *Conjugate Similarity and Hermitian Products*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$. The following are equivalent:*

*(i) $A$ is similar to $\overline{A}$.*
*(ii) $A$ is similar to $A^*$.*
*(iii) $A$ is similar to $A^*$ via a Hermitian similarity matrix.*
*(iv) $A = HK = LM$, where $H, K, L, M$ are Hermitian, and $H, M$ are invertible.*
*(v) $A = HK$ for some Hermitian matrices $H, K$.*

*Proof.*

(i) $\Rightarrow$ (ii) By Theorem 7.1.4 applied to $\overline{A}$, $\overline{A} \sim (\overline{A})^t = A^*$. Thus $A \sim A^*$.

(*ii*) $\Rightarrow$ (*iii*) Choose invertible $S$ such that $A = SA^*S^{-1}$, so $AS = SA^*$. For $\theta \in \mathbb{R}$, put $S_\theta = e^{i\theta}S$. Then $AS_\theta = S_\theta A^*$ and, by taking adjoints, $AS_\theta^* = S_\theta^* A^*$. Hence $A(S_\theta + S_\theta^*) = (S_\theta + S_\theta^*)A^*$. The matrix $H_\theta = S_\theta + S_\theta^*$ is Hermitian. Also $S_\theta^{-1}H_\theta = I_n + e^{-2i\theta}S^{-1}S^*$. Thus $H_\theta$ is singular exactly when $-e^{2i\theta}$ is an eigenvalue of $S^{-1}S^*$. There are only finitely many such values, so choose $\theta$ for which $H_\theta$ is invertible. Then $A = H_\theta A^* H_\theta^{-1}$.

(*iii*) $\Rightarrow$ (*iv*) Let $R$ be invertible Hermitian and satisfy $A = RA^*R^{-1}$. Then $AR = RA^*$ and $R^{-1}A = A^*R^{-1}$. Thus $A^*R^{-1}$ and $RA^*$ are Hermitian, and $A = R(A^*R^{-1}) = (RA^*)R^{-1}$. This gives the two asserted Hermitian factorizations, with the first factor in the first product and the second factor in the second product invertible.

(*iv*) $\Rightarrow$ (*v*) The first factorization in (iv) already has the form $A = HK$ with $H$ and $K$ Hermitian.

(*v*) $\Rightarrow$ (*i*) If $A = HK$ with $H, K$ Hermitian, then by Corollary 7.1.5, $HK \sim KH$. By Theorem 7.1.4, $KH \sim (KH)^t = H^tK^t = \overline{H}\,\overline{K} = \overline{A}$. Hence $A \sim \overline{A}$.

*Structural Commentary*: The equivalences move between conjugation, adjoints, and Hermitian products by combining transpose similarity with a Hermitianized similarity matrix.
*Key Insight*: A phase-rotated average $e^{i\theta}S + e^{-i\theta}S^*$ turns an arbitrary similarity to $A^*$ into a Hermitian one. □




Figure 76: Conjugate similarity, adjoint similarity, and Hermitian product factorisations are equivalent.

# Part VI
# Matrix Analysis and Numerical Linear Algebra

## 1 Trace, Determinant, Rank, and Matrix Invariants

### 1.1 Trace and Determinant

Matrix classes package operator structure in coordinates. Trace, determinant, rank, positivity, idempotents, nilpotents, spectral radius, and structured products give computable invariants.

**Definition 1.1.1.** *Trace and Determinant of a Matrix*
For $A = [A_{j,k}] \in \mathrm{Mat}(n, n, \mathbb{F})$, the *trace* of $A$ is $\mathrm{tr}(A) := A_{1,1} + \cdots + A_{n,n}$.
The *determinant* of $A$ is $\det A := \sum_{\sigma \in \mathfrak{S}_n} \mathrm{sign}(\sigma) A_{\sigma(1),1} \cdots A_{\sigma(n),n}$.

**Definition 1.1.2.** *Gram Determinant*
For vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$ in an inner-product space, define their *Gram determinant* by $\mathrm{GramDet}(\mathbf{u}_1, \ldots, \mathbf{u}_n) := \det \mathrm{Gram}(\mathbf{u}_1, \ldots, \mathbf{u}_n)$.

**Example 1.1.3.** *Trace Functional and the Frobenius Inner Product*
On $\mathrm{Mat}(n, n, \mathbb{F})$ with the Frobenius inner product from Example 1.1.6, the Riesz vector of the trace functional $A \mapsto \mathrm{tr}(A)$ is $I_n$, since $\langle A, I_n \rangle_F = \sum_{j=1}^{n} a_{jj} = \mathrm{tr}(A)$.

**Proposition 1.1.4.** *Trace Arithmetic*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$ and $r \in \mathbb{F}$. Then*
*(i)* $\mathrm{tr}(rA) = r\,\mathrm{tr}(A)$;
*(ii)* $\mathrm{tr}(A + B) = \mathrm{tr}(A) + \mathrm{tr}(B)$.

*Proof.* We prove the two identities separately.
(i) Summing the diagonal entries of $rA$ gives $\mathrm{tr}(rA) = \sum_{j=1}^{n} r a_{jj} = r \sum_{j=1}^{n} a_{jj} = r\,\mathrm{tr}(A)$.
(ii) Summing the diagonal entries of $A+B$ gives $\mathrm{tr}(A+B) = \sum_{j=1}^{n} (a_{jj}+b_{jj}) = \sum_{j=1}^{n} a_{jj} + \sum_{j=1}^{n} b_{jj} = \mathrm{tr}(A)+\mathrm{tr}(B)$.
*Structural Commentary*: Trace is linear because it is the sum of diagonal entries.
*Key Insight*: Scalar multiplication and addition pass through the diagonal sum. □

**Proposition 1.1.5.** *Trace Is Cyclic for Two Factors*
*If $A \in \mathrm{Mat}(m, n, \mathbb{F})$ and $B \in \mathrm{Mat}(n, m, \mathbb{F})$, then $\mathrm{tr}(AB) = \mathrm{tr}(BA)$.*

*Proof.* Expand both traces by entries: $\mathrm{tr}(AB) = \sum_{j=1}^{m} \sum_{k=1}^{n} A_{j,k} B_{k,j} = \sum_{k=1}^{n} \sum_{j=1}^{m} B_{k,j} A_{j,k} = \mathrm{tr}(BA)$.
*Structural Commentary*: Cyclicity is just reindexing the same double sum.
*Key Insight*: The two double sums contain exactly the same scalar terms. □

**Proposition 1.1.6.** *Cyclic Products*
*Let $m \geq 2$, $A_j \in \mathrm{Mat}(n_j, n_{j+1}, \mathbb{F})$ for $j = 1, \ldots, m$, $n_{m+1} = n_1$, $P_{p,q} := A_p A_{p+1} \cdots A_q$.*
*Then $\mathrm{tr}(P_{1,m}) = \mathrm{tr}(A_m P_{1,m-1})$. Consequently, for $A, B, C \in \mathrm{Mat}(n, n, \mathbb{F})$, $\mathrm{tr}(ABC) = \mathrm{tr}(BCA) = \mathrm{tr}(CAB)$.*

*Proof.* Since $P_{1,m-1} \in \mathrm{Mat}(n_1, n_m, \mathbb{F})$ and $A_m \in \mathrm{Mat}(n_m, n_1, \mathbb{F})$, by Proposition 1.1.5, $\mathrm{tr}(P_{1,m}) = \mathrm{tr}(P_{1,m-1} A_m) = \mathrm{tr}(A_m P_{1,m-1})$.
Applying this cyclic move twice to $ABC$ gives the three-factor identity.
*Structural Commentary*: Trace preserves cyclic order, not arbitrary order.
*Key Insight*: One cyclic move is $\mathrm{tr}(XY) = \mathrm{tr}(YX)$ with $X = P_{1,m-1}$ and $Y = A_m$. □

**Example 1.1.7.** *Cyclicity Is Not Arbitrary Commutativity*
Let $A = E_{1,2}$, $B = E_{2,1}$, and $C = E_{1,1}$ in $\mathrm{Mat}(2, 2, \mathbb{F})$. Then $\mathrm{tr}(ABC) = 1$, while $\mathrm{tr}(ACB) = 0$.
Thus trace permits cyclic rotations of factors, but not arbitrary rearrangements.

**Theorem 1.1.8.** *Basic Determinant Laws*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$.*
*(i)* $\det I_n = 1$.
*(ii)* $\det(A)$ *is linear as a function of each column of $A$ when the other columns are fixed.*

*(iii)* *Swapping two columns negates the determinant; equal columns force* $\det A = 0$.
*(iv)* *If $A$ is triangular, then* $\det A = A_{1,1} \cdots A_{n,n}$.
*(v)* $\det(AB) = \det(A)\det(B)$.
*(vi)* *$A$ is invertible if and only if* $\det A \neq 0$. *In that case* $\det(A^{-1}) = (\det A)^{-1}$.
*(vii)* $\det(A^t) = \det(A)$.

*Proof.*

(i) $\det I_n = \sum_{\sigma \in \mathfrak{S}_n} \operatorname{sign}(\sigma)(I_n)_{\sigma(1),1} \cdots (I_n)_{\sigma(n),n} = 1$ as product is nonzero only when $\sigma(k) = k\ \forall k$.

(ii) For fixed $k$, $\det(A_{\cdot,1}, \ldots, \mathbf{x}, \ldots, A_{\cdot,n}) = \sum_{\sigma \in \mathfrak{S}_n} \operatorname{sign}(\sigma) x_{\sigma(k)} \prod_{\ell \neq k} A_{\sigma(\ell),\ell}$, which is linear in $\mathbf{x}$.

(iii) If $B_{\cdot,k} = A_{\cdot,\tau(k)}$ for transposition $\tau$, then $\det B = \sum_{\rho \in \mathfrak{S}_n} \operatorname{sign}(\rho\tau) \prod_{k=1}^{n} A_{\rho(k),k} = -\det A$.
If the swapped columns are equal, then $B = A$, so $\det A = -\det A$ and $\det A = 0$.

(iv) If $A$ is upper triangular and $\prod_{k=1}^{n} A_{\sigma(k),k} \neq 0$, then $\sigma(k) \leq k$ for all $k$, hence $\sigma(k) = k$ for all $k$.
Thus $\det A = A_{1,1} \cdots A_{n,n}$. If $A$ is lower triangular, nonzero products require $\sigma(k) \geq k$ for all $k$, again forcing $\sigma(k) = k$ for all $k$, so the same diagonal product formula holds.

(v) If $F$ is alternating and multilinear in columns, then expanding each column in the standard basis gives $F(A_{\cdot,1}, \ldots, A_{\cdot,n}) = \sum_{i_1,\ldots,i_n} A_{i_1,1} \cdots A_{i_n,n} F(\mathbf{e}_{i_1}, \ldots, \mathbf{e}_{i_n}) = \sum_{\sigma \in \mathfrak{S}_n} \operatorname{sign}(\sigma) F(I_n) A_{\sigma(1),1} \cdots A_{\sigma(n),n}$.
The terms with repeated $i_j$ vanish by alternation, and the surviving ordered bases differ from $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ by sign of permutation. For $H(B) = \det(AB)$, map $H$ is alternating multilinear and $H(I_n) = \det A$.
Hence $\det(AB) = H(B) = H(I_n)\det B = \det(A)\det(B)$.

(vi) If $A$ is invertible, then $1 = \det I_n = \det(AA^{-1}) = \det(A)\det(A^{-1})$.
If $A_{\cdot,k} = \sum_{\ell \neq k} c_\ell A_{\cdot,\ell}$, then $\det A = \sum_{\ell \neq k} c_\ell \det(A_{\cdot,1}, \ldots, A_{\cdot,\ell}, \ldots, A_{\cdot,\ell}, \ldots, A_{\cdot,n}) = 0$.
Thus $\det A \neq 0$ iff the columns of $A$ are independent, iff $A$ is invertible.

(vii) Using the permutation definition and replacing $\sigma$ by $\sigma^{-1}$,
$\det(A^t) = \sum_{\sigma \in \mathfrak{S}_n} \operatorname{sign}(\sigma) A_{1,\sigma(1)} \cdots A_{n,\sigma(n)} = \sum_{\tau \in \mathfrak{S}_n} \operatorname{sign}(\tau) A_{\tau(1),1} \cdots A_{\tau(n),n} = \det(A)$; $\operatorname{sign}(\tau^{-1}) = \operatorname{sign}(\tau)$.

*Structural Commentary*: The determinant is the normalized alternating multilinear measure of columns.
*Key Insight*: Multilinearity expands columns; alternation kills repeated directions. □




Figure 77: Determinant is signed column-volume; swaps reverse orientation and dependence gives zero volume.

**Method 1.1.9.** *Determinant by Structure*
To compute or simplify a determinant:

(i) first look for triangular or block triangular form;
(ii) use column linearity to split complicated columns only when it reduces structure;
(iii) use column swaps to move useful pivots into position, tracking a factor of $-1$ for each swap;
(iv) detect zero determinants from dependent columns or repeated columns;
(v) for products or changes of basis, use multiplicativity instead of expanding permutations.

These steps use Theorem 1.1.8.




Figure 78: A linear map scales oriented volume by its determinant; singular maps collapse volume to zero.

**Lemma 1.1.10.** *Identity Blocks Do Not Change Determinant*
*Let* $A \in \mathrm{Mat}(m, m, \mathbb{F})$ *and let* $n \geq 0$. *Then* $\det(I_n \oplus A) = \det(A) = \det(A \oplus I_n)$.

*Proof.* In the permutation formula for $I_n \oplus A$, a nonzero term must satisfy $\sigma(k) = k$ for $1 \leq k \leq n$. Remaining choices are exactly permutations of $\{n+1, \ldots, n+m\}$, after relabeling, gives $\det(A)$. Hence $\det(I_n \oplus A) = \det(A)$.
The same argument applied to the last $n$ coordinates gives $\det(A \oplus I_n) = \det(A)$.
The case $n = 0$ is the empty identity block convention.
*Structural Commentary*: Identity blocks add directions without changing volume.
*Key Insight*: In the determinant expansion, the identity block forces the matching coordinates to be fixed. □

**Theorem 1.1.11.** *Determinant of an Upper Block-Triangular Matrix*
*Let* $A \in \mathrm{Mat}(r, r, \mathbb{F})$, $B \in \mathrm{Mat}(r, s, \mathbb{F})$, *and* $D \in \mathrm{Mat}(s, s, \mathbb{F})$. *Then* $\det \begin{bmatrix} A & B \\ 0 & D \end{bmatrix} = \det(A)\det(D)$.

*Proof.* Factor $\begin{bmatrix} A & B \\ 0 & D \end{bmatrix} = \begin{bmatrix} I_r & 0 \\ 0 & D \end{bmatrix} \begin{bmatrix} I_r & B \\ 0 & I_s \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & I_s \end{bmatrix}$.
The first and last factors have determinants $\det(D)$ and $\det(A)$ by Lemma 1.1.10.
The middle factor is upper triangular with every diagonal entry equal to 1, so its determinant is 1.
Multiplicativity of determinant gives the formula.
*Structural Commentary*: Block upper triangular matrices separate determinant contribution into diagonal blocks. The off-diagonal block does not affect the determinant.
*Key Insight*: Factor the matrix into two block diagonal pieces and one unit upper triangular piece. □

**Proposition 1.1.12.** *Schur Complement Determinant Formula*
*Let* $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$, $A \in \mathrm{Mat}(r, r, \mathbb{F})$, $D \in \mathrm{Mat}(s, s, \mathbb{F})$.
*If* $A$ *is invertible, then the Schur complement of* $A$ *in* $M$ *is* $M/A := D - CA^{-1}B$, *and* $\det(M) = \det(A)\det(M/A)$.

*Proof.* Block Gaussian elimination gives $\begin{bmatrix} I_r & 0 \\ -CA^{-1} & I_s \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} A & B \\ 0 & D - CA^{-1}B \end{bmatrix}$.
The elimination matrix is lower block-triangular with identity diagonal blocks, so its determinant is 1. Take determinants, apply Theorem 1.1.11 to right-hand block triangular matrix, $\det(M) = \det(A)\det(D - CA^{-1}B)$.
*Structural Commentary*: Schur complementation is block Gaussian elimination.
*Key Insight*: Eliminating $C$ splits the determinant into $\det(A)$ times the remaining block determinant. □

**Proposition 1.1.13.** *Schur Complement of the Lower-Right Block*
*Let* $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$, $A \in \mathrm{Mat}(r, r, \mathbb{F})$, $D \in \mathrm{Mat}(s, s, \mathbb{F})$.
*If* $D$ *is invertible, then the Schur complement of* $D$ *in* $M$ *is* $M/D := A - BD^{-1}C$, *and* $\det(M) = \det(D)\det(M/D)$.

*Proof.* Block Gaussian elimination gives $\begin{bmatrix} I_r & -BD^{-1} \\ 0 & I_s \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} A - BD^{-1}C & 0 \\ C & D \end{bmatrix}$.
The elimination matrix has determinant 1. The transpose of the matrix on the right is upper block-triangular, and $\det(X^t) = \det(X)$ by Theorem 1.1.8. By Theorem 1.1.11, $\det(M) = \det(A - BD^{-1}C)\det(D)$.
*Structural Commentary*: This is the lower-right version of Schur complementation. Eliminating the upper-right block isolates the determinant contribution of $D$.
*Key Insight*: Eliminate by blocks, then read the determinant from the diagonal blocks. □

**Proposition 1.1.14.** *Bordered Determinant Formulas*
*Let* $A \in \mathrm{Mat}(n, n, \mathbb{F})$, *let* $\mathbf{x}, \mathbf{y} \in \mathbb{F}^n$, *and let* $c \in \mathbb{F}$.

*(i) If* $c \neq 0$, *then* $\det \begin{bmatrix} c & \mathbf{x}^t \\ \mathbf{y} & A \end{bmatrix} = c \det\left(A - \frac{1}{c}\mathbf{y}\mathbf{x}^t\right)$.

*(ii) If* $A$ *is invertible, then* $\det \begin{bmatrix} A & \mathbf{x} \\ \mathbf{y}^t & c \end{bmatrix} = \left(c - \mathbf{y}^t A^{-1}\mathbf{x}\right)\det(A)$

*Proof.*

(i) This is Proposition 1.1.12 with the scalar block $c$ in the upper-left corner.
(ii) This is Proposition 1.1.12 with $A$ as the upper-left block and $D = c$.

*Structural Commentary*: Bordered determinants are Schur complements with scalar borders.
*Key Insight*: Treat the border as a $1 \times 1$ block and apply the Schur complement formula. □

**Theorem 1.1.15.** *Sylvester Determinant Identity*
*Let* $A \in \mathrm{Mat}(n, m, \mathbb{F})$ *and* $B \in \mathrm{Mat}(m, n, \mathbb{F})$. *Then* $\det(I_n + AB) = \det(I_m + BA)$.

*Proof.* Let $M = \begin{bmatrix} I_n & -A \\ B & I_m \end{bmatrix}$.
By Proposition 1.1.13 with lower-right block $I_m$, $\det(M) = \det(I_n + AB)$. By Proposition 1.1.12 with upper-left block $I_n$, $\det(M) = \det(I_m + BA)$. Hence the two determinants are equal.
*Structural Commentary*: Two Schur complements of one block matrix give Sylvester's identity.
*Key Insight*: Compare the same determinant after eliminating different identity blocks. □

**Theorem 1.1.16.** *Nonzero Eigenvalues of AB and BA*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{C})$, $B \in \mathrm{Mat}(n, m, \mathbb{C})$, and suppose $n \geq m$. Then $p_{BA}(z) = z^{n-m} p_{AB}(z)$. Consequently, for $\lambda \neq 0$, $m_{AB}(\lambda) = m_{BA}(\lambda)$. If $m_{AB}(0) = k$, then $m_{BA}(0) = k + n - m$.*

*Proof.* For $z \neq 0$, by Theorem 1.1.15 applied to $-z^{-1}B$ and $A$,

$$\det(zI_n - BA) = z^n \det(I_n - z^{-1}BA) = z^n \det(I_m - z^{-1}AB) = z^{n-m} \det(zI_m - AB).$$

Both sides are polynomials in $z$, so the identity holds for all $z$. The claimed multiplicities follow by factoring $p_{AB}$ and comparing with $p_{BA} = z^{n-m} p_{AB}$.
*Structural Commentary*: Sylvester's determinant identity moves the rectangular product across the determinant.
*Key Insight*: Passing from $AB$ to $BA$ only adds the necessary extra zero roots when the sizes differ. □

**Theorem 1.1.17.** *Trace and Determinant Are Similarity Invariants*
*If $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$ are similar, then $\mathrm{tr}(A) = \mathrm{tr}(B)$, $\det(A) = \det(B)$.*

*Proof.* Write $B = PAP^{-1}$ with $P$ invertible.
Proposition 1.1.5 gives $\mathrm{tr}(B) = \mathrm{tr}\big(P(AP^{-1})\big) = \mathrm{tr}\big((AP^{-1})P\big) = \mathrm{tr}(A)$.
Theorem 1.1.8 gives $\det(B) = \det(P)\det(A)\det(P^{-1}) = \det(A)$.
*Structural Commentary*: Trace and determinant ignore the coordinate system used to write an operator.
*Key Insight*: Conjugating by $P$ inserts $P$ and $P^{-1}$, which cancel inside trace and determinant. □

**Perspective 1.1.18.** *Similarity Invariants and Canonical Forms*
Similarity is equality up to change of basis. Trace, determinant, characteristic polynomials, and minimal polynomials are similarity invariants by Theorems 1.1.17, 2.1.7, and 3.1.7, but they are not generally complete. Canonical forms, such as the Jordan form of Theorem 6.2.1, choose one structured representative of each similarity class when possible.

**Principle 1.1.19.** *Trace and Determinant Are Coordinate-Free*
Trace and determinant are first computed from a matrix, but their real structural meaning is operator-level: changing basis conjugates the matrix, and conjugation preserves both quantities.

**Theorem 1.1.20.** *Trace and Determinant of an Operator Are Basis-Independent*
*Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $\beta, \gamma$ be bases of $\mathcal{V}$. Then $\mathrm{tr}\big([T]_{\beta \leftarrow \beta}\big) = \mathrm{tr}\big([T]_{\gamma \leftarrow \gamma}\big)$, $\det\big([T]_{\beta \leftarrow \beta}\big) = \det\big([T]_{\gamma \leftarrow \gamma}\big)$.*

*Proof.* By Theorem 6.2.1, the two matrices are similar: $[T]_{\gamma \leftarrow \gamma} = C_{\gamma \leftarrow \beta} [T]_{\beta \leftarrow \beta} C_{\beta \leftarrow \gamma}$.
Theorem 1.1.17 now gives equality of traces and determinants.
*Structural Commentary*: Basis changes conjugate matrices; trace and determinant remain operator invariants.
*Key Insight*: Similar matrices have the same trace and determinant. □




Figure 79: Basis changes conjugate matrices; trace and determinant are unchanged.

**Definition 1.1.21.** *Trace and Determinant of an Operator*
Let $\mathcal{V}$ be finite-dim, let $T \in \mathcal{L}(\mathcal{V})$. Choose any basis $\beta$ of $\mathcal{V}$. Define $\mathrm{tr}\, T := \mathrm{tr}\big([T]_{\beta \leftarrow \beta}\big)$, $\det T := \det\big([T]_{\beta \leftarrow \beta}\big)$.
Theorem 1.1.20 shows these definitions do not depend on chosen basis.

**Proposition 1.1.22.** *Unitary Matrices and Isometries Have Determinant of Modulus One*
*Let $U \in \mathrm{Mat}(n, n, \mathbb{F})$ be unitary. Then $|\det U| = 1$. More generally, if $S \in \mathcal{L}(\mathcal{V})$ is an isometry on a finite-dimensional inner-product space over $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, then $|\det S| = 1$.*

*Proof.* The permutation definition gives $\det(U^*) = \overline{\det U}$. By Theorem 1.1.8, taking determinants in $U^*U = I_n$ gives $1 = \det(U^*)\det(U) = \overline{\det U}\det U = |\det U|^2$. Hence $|\det U| = 1$.
For an isometry $S$, choose an orthonormal basis and let $U$ be the matrix of $S$ in that basis. Then $U$ is unitary by Theorem 2.6.6, and $\det S = \det U$.
*Structural Commentary*: In orthonormal coordinates, isometries are unitary or orthogonal, so the determinant calculation is the matrix one.
*Key Insight*: Determinants turn $U^*U = I$ into $|\det U|^2 = 1$. □

**Proposition 1.1.23.** *Trace and Determinant from $q_T$*
*Let $\mathcal{V} \neq \{\mathbf{0}\}$ be finite-dim over $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, let $T \in \mathcal{L}(\mathcal{V})$, $q_T(z) = z^n + c_{n-1}z^{n-1} + \cdots + c_1 z + c_0$, $n = \dim \mathcal{V}$. Then $\operatorname{tr} T = -c_{n-1}$ and $\det T = (-1)^n c_0$. If $\mathbb{F} = \mathbb{C}$ and $\lambda_1, \ldots, \lambda_n$ list the eigenvalues of $T$ with algebraic multiplicity, then $\operatorname{tr} T = \lambda_1 + \cdots + \lambda_n$, $\det T = \lambda_1 \cdots \lambda_n$.*

*Proof.* Choose a basis $\beta$ and set $A = [T]_{\beta \leftarrow \beta}$.
Then $q_T = p_A$, and by Definition 1.1.21, $\operatorname{tr} T = \operatorname{tr} A$ and $\det T = \det A$.
Theorem 2.1.3 gives $\operatorname{tr} T = -c_{n-1}$ and $\det T = (-1)^n c_0$.
If $\mathbb{F} = \mathbb{C}$, then comparing $q_T(z) = (z - \lambda_1)\cdots(z - \lambda_n)$ with the displayed form of $q_T$ gives the formulas.
*Structural Commentary*: Characteristic polynomial stores trace and determinant in first and last coefficients.
*Key Insight*: Compare coefficients of $\det(zI - T)$. □

**Proposition 1.1.24.** *Trace and Determinant from Eigenvalues*
*Let $\lambda_1, \ldots, \lambda_n$ list the eigenvalues of $A \in \operatorname{Mat}(n, n, \mathbb{C})$ according to $m_A$. Then $\operatorname{tr} A = \sum_{j=1}^{n} \lambda_j$ and $\det A = \prod_{j=1}^{n} \lambda_j$.*

*Proof.* By Fundamental Theorem of Algebra, $p_A(z) = \prod_{j=1}^{n}(z - \lambda_j) = z^n - \left(\sum_{j=1}^{n} \lambda_j\right) z^{n-1} + \cdots + (-1)^n \prod_{j=1}^{n} \lambda_j$.
Theorem 2.1.3 gives $p_A(z) = z^n - (\operatorname{tr} A)z^{n-1} + \cdots + (-1)^n \det A$. Uniqueness of coefficients gives the identities.
*Structural Commentary*: Coefficients of $p_A$ encode symmetric functions of the eigenvalues.
*Key Insight*: Trace is the first root sum and determinant is the full root product. □

**Proposition 1.1.25.** *Transpose, Conjugate, and Adjoint Spectra*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{C})$. Then $p_{A^t}(z) = p_A(z)$, $p_{\overline{A}}(z) = \overline{p_A(\overline{z})}$, and $p_{A^*}(z) = \overline{p_A(\overline{z})}$.*
*Consequently, if $\lambda \in \operatorname{spec}(A)$, then $m_{A^t}(\lambda) = m_A(\lambda)$ and $m_{\overline{A}}(\overline{\lambda}) = m_{A^*}(\overline{\lambda}) = m_A(\lambda)$.*

*Proof.* Use determinant identities:

$$p_{A^t}(z) = \det(zI_n - A^t) = \det((zI_n - A)^t) = p_A(z),$$
$$p_{\overline{A}}(z) = \det(zI_n - \overline{A}) = \det(\overline{\overline{z}I_n - A}) = \overline{\det(\overline{z}I_n - A)} = \overline{p_A(\overline{z})}.$$

Since $A^* = \overline{A}^t$, the first identity applied to $\overline{A}$ gives the formula for $A^*$.
Multiplicity is preserved because conjugating a polynomial sends a factor $(z - \lambda)^k$ to $(z - \overline{\lambda})^k$.
*Structural Commentary*: Transpose preserves $p_A$; conjugation conjugates its roots.
*Key Insight*: The adjoint combines conjugation with transposition, so its spectrum is conjugated. □

## 1.2 Minors, Cofactors, and Rank

**Definition 1.2.1.** *Minors, Cofactors, and Adjugate*
For $A \in \operatorname{Mat}(n, n, \mathbb{F})$, let $M_{ij}$ delete row $i$ and column $j$ from $A$; set $\det(\emptyset_{0\times 0}) = 1$. Define $C_{ij} := (-1)^{i+j}\det(M_{ij})$. Then $C = [C_{ij}]$ is the *cofactor matrix*, and $\operatorname{adj}(A) := C^t$ is the *adjugate*.

**Lemma 1.2.2.** *Cofactor Expansion*
*For $A = [a_{ij}] \in \operatorname{Mat}(n, n, \mathbb{F})$, $\det A = \sum_{k=1}^{n} a_{ik}C_{ik}$ and $\det A = \sum_{k=1}^{n} a_{kj}C_{kj}$.*

*Proof.* Using $\det A = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{r=1}^{n} a_{r,\sigma(r)}$, group terms by $\sigma(i) = k$.
For fixed $k$, the remaining permutation sum equals $(-1)^{i+k}\det(M_{ik}) = C_{ik}$, giving the row formula. Applying this to $A^t$ and using Theorem 1.1.8 gives the column formula.
*Structural Commentary*: Cofactor expansion is the big determinant formula collected around one row or column.
*Key Insight*: Fix one entry; the rest is its minor. □

**Theorem 1.2.3.** *Adjugate Identity and Cramer's Rule*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{F})$ and let $\mathbf{b} \in \mathbb{F}^n$. Then $A\operatorname{adj}(A) = \operatorname{adj}(A)A = (\det A)I_n$. If $\det A \neq 0$, then, for $A\mathbf{x} = \mathbf{b}$, $A^{-1} = \frac{1}{\det A}\operatorname{adj}(A)$ and $x_j = \frac{\det A_j(\mathbf{b})}{\det A}$ for $1 \leq j \leq n$, where $A_j(\mathbf{b})$ replaces column $j$ of $A$ by $\mathbf{b}$.*

*Proof.* Since $(\operatorname{adj} A)_{kj} = C_{jk}$, $(A \operatorname{adj} A)_{ij} = \sum_k a_{ik} C_{jk}$.
For $i = j$ this is $\det A$ by Lemma 1.2.2; for $i \neq j$ it is the cofactor expansion of a matrix with rows $i$ and $j$ equal, hence 0. Thus $A \operatorname{adj}(A) = (\det A) I_n$. The same argument applied to $A^t$ gives $\operatorname{adj}(A) A = (\det A) I_n$. If $\det A \neq 0$, division gives the inverse formula. Finally, $(\operatorname{adj} A\,\mathbf{b})_j = \sum_i C_{ij} b_i = \det A_j(\mathbf{b})$, by expansion down column $j$, so $x = A^{-1}\mathbf{b}$ gives Cramer's rule.
*Structural Commentary*: Cofactors assemble all one-column replacements of $A$ into a single inverse formula.
*Key Insight*: Off-diagonal products vanish because they are determinants with two equal rows or columns. □

**Remark 1.2.4.** *Adjugate Form of the Bordered Formula*
Proposition 1.1.14 gives the adjugate form $\det \begin{bmatrix} A & \mathbf{x} \\ \mathbf{y}^t & c \end{bmatrix} = c \det(A) - \mathbf{y}^t \operatorname{adj}(A) \mathbf{x}$. No invertibility of $A$ is required.

**Theorem 1.2.5.** *Cauchy–Binet Formula*
*Let $A \in \operatorname{Mat}(k, n, \mathbb{F})$, $B \in \operatorname{Mat}(n, k, \mathbb{F})$, and $k \leq n$. For $J = \{j_1 < \cdots < j_k\} \subseteq \{1, \ldots, n\}$, let $A^J$ use the columns of $A$ in $J$ and $B_J$ use the rows of $B$ in $J$.*
*Then $\det(AB) = \sum_J \det(A^J) \det(B_J)$, where $J$ ranges over all $k$-subsets.*

*Proof.* Put $S(A, B) := \sum_J \det(A^J) \det(B_J)$. For an elementary $n \times n$ matrix $E$, determinant multilinearity and row/column operation rules give $S(A, B) = S(AE, E^{-1}B)$, and $AB = (AE)(E^{-1}B)$.
If $\operatorname{rank} A < k$, both sides vanish. Otherwise reduce $A$ by elementary column operations to $A' = [C\ 0]$ with $C$ invertible, and transform $B$ oppositely to $B'$.
Then $AB = A'B'$ and $\det(A'B') = \det(C) \det(B'_{J_0}) = S(A', B')$, $J_0 = \{1, \ldots, k\}$, as every other $k$-minor of $A'$ is zero. Hence $\det(AB) = S(A, B)$.
*Structural Commentary*: Cauchy–Binet is determinant multiplicativity for rectangular middle dimensions.
*Key Insight*: The middle space contributes one $k$-coordinate shadow at a time. □

**Proposition 1.2.6.** *Transpose Preserves Spectrum and Geometric Multiplicity*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{F})$ and $\lambda \in \mathbb{F}$. Then $\lambda \in \operatorname{spec}(A) \iff \lambda \in \operatorname{spec}(A^t)$, and $\dim E_\lambda(A) = \dim E_\lambda(A^t)$.*

*Proof.* Since $(A - \lambda I_n)^t = A^t - \lambda I_n$, Theorem 3.3.4 gives $\operatorname{rank}(A - \lambda I_n) = \operatorname{rank}(A^t - \lambda I_n)$.
Matrix rank-nullity then gives $\dim E_\lambda(A) = n - \operatorname{rank}(A - \lambda I_n) = n - \operatorname{rank}(A^t - \lambda I_n) = \dim E_\lambda(A^t)$.
($\Rightarrow$) If $\lambda \in \operatorname{spec}(A)$, then $E_\lambda(A) \neq \{\mathbf{0}\}$, so $E_\lambda(A^t) \neq \{\mathbf{0}\}$ and $\lambda \in \operatorname{spec}(A^t)$.
($\Leftarrow$) The same argument with $A$ and $A^t$ interchanged gives the converse.
*Structural Commentary*: Transposition preserves rank, so shifted kernels have the same dimension.
*Key Insight*: Spectrum and geometric multiplicity are kernel facts for $A - \lambda I_n$. □

**Corollary 1.2.7.** *Square Matrix Algebraic Invertibility Dictionary*
*Let $A = [\mathbf{a}_1\ \cdots\ \mathbf{a}_n] \in \operatorname{Mat}(n, n, \mathbb{F})$. The following are equivalent:*

*(i) $A$ is invertible;*
*(ii) $\mathbf{a}_1, \ldots, \mathbf{a}_n$ form a basis of $\mathbb{F}^n$;*
*(iii) $\mathbf{a}_1, \ldots, \mathbf{a}_n$ are linearly independent;*
*(iv) $\mathbf{a}_1, \ldots, \mathbf{a}_n$ span $\mathbb{F}^n$;*
*(v) for every $\mathbf{b} \in \mathbb{F}^n$, the system $A\mathbf{x} = \mathbf{b}$ has a unique solution;*
*(vi) the homogeneous system $A\mathbf{x} = \mathbf{0}$ has only the zero solution.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) By Theorem 3.4.1, the columns of an invertible matrix form a basis.
(*ii*) $\Rightarrow$ (*iii*) A basis is linearly independent.
(*iii*) $\Rightarrow$ (*iv*) In $\mathbb{F}^n$, $n$ linearly independent vectors form a basis, hence span.
(*iv*) $\Rightarrow$ (*v*) In $\mathbb{F}^n$, $n$ spanning vectors form a basis, so every $\mathbf{b}$ has unique coordinates $\mathbf{x}$ in $A\mathbf{x} = \mathbf{b}$.
(*v*) $\Rightarrow$ (*vi*) Apply (v) to $\mathbf{b} = \mathbf{0}$; since $\mathbf{x} = \mathbf{0}$ is a solution, it is the only one.
(*vi*) $\Rightarrow$ (*i*) Condition (vi) says the columns are independent.
Hence they form a basis of $\mathbb{F}^n$, and Corollary 3.6.1 gives invertibility.

*Structural Commentary*: Square invertibility is column basishood.
*Key Insight*: Invertible columns give unique coordinates. □

**Corollary 1.2.8.** *Nonsingularity by Rank, Determinant, and Spectrum*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{F})$. The following are equivalent:*

*(i) $A$ is invertible;*
*(ii) $\operatorname{rank}(A) = n$;*

*(iii)* $\det A \neq 0$;
*(iv)* $0 \notin \operatorname{spec}(A)$.

*Over $\mathbb{C}$, this is also equivalent to all eigenvalues of $A$ being nonzero.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) This is Corollary 3.6.3.
(*ii*) $\Rightarrow$ (*iii*) Full rank gives invertibility by Corollary 3.6.3, hence $\det A \neq 0$ by Theorem 1.1.8.
(*iii*) $\Rightarrow$ (*iv*) If $\det A \neq 0$, then $A$ is invertible by Theorem 1.1.8; hence $0 \notin \operatorname{spec}(A)$ by Corollary 5.1.8.
(*iv*) $\Rightarrow$ (*i*) This is Corollary 5.1.8.

Over $\mathbb{C}$, Theorem 2.1.3 identifies $\lambda_1, \ldots, \lambda_n$ as the roots of $p_A$, so (iv) is equivalent to $\lambda_j \neq 0$ for all $j$.
*Structural Commentary*: Rank, determinant, and spectrum detect singularity.
*Key Insight*: Singularity can be seen as rank loss, zero determinant, or the zero eigenvalue. □

**Theorem 1.2.9.** *Rank Bounds for Products and Concatenations*
*Let $A \in \operatorname{Mat}(m,k,\mathbb{F})$, $B \in \operatorname{Mat}(k,n,\mathbb{F})$, and $C \in \operatorname{Mat}(m,p,\mathbb{F})$. Then $\operatorname{rank}(AB) \leq \min(\operatorname{rank}(A), \operatorname{rank}(B))$ and $\operatorname{rank}(AB) = \operatorname{rank}(A)$ if and only if $\operatorname{col}(AB) = \operatorname{col}(A)$.*
*Also, $\max(\operatorname{rank}(A), \operatorname{rank}(C)) \leq \operatorname{rank}\begin{bmatrix} A & C \end{bmatrix}$ and $\operatorname{rank}\begin{bmatrix} A & C \end{bmatrix} = \operatorname{rank}(A)$ if and only if $\operatorname{col}(A) = \operatorname{col}(A) + \operatorname{col}(C)$.*

*Proof.* Each column of $AB$ is a linear combination of the columns of $A$, so $\operatorname{col}(AB) \subseteq \operatorname{col}(A)$. Therefore $\operatorname{rank}(AB) \leq \operatorname{rank}(A)$; since both spaces are finite-dimensional, equality holds exactly when $\operatorname{col}(AB) = \operatorname{col}(A)$.
Applying this same argument to $(AB)^t = B^t A^t$ and using Theorem 3.3.4 gives $\operatorname{rank}(AB) \leq \operatorname{rank}(B)$.
For the block matrix, its columns are precisely the columns of $A$ and $C$, hence $\operatorname{col}\begin{bmatrix} A & C \end{bmatrix} = \operatorname{col}(A) + \operatorname{col}(C)$.
This space contains both $\operatorname{col}(A)$ and $\operatorname{col}(C)$, giving the lower bound. Equality with $\operatorname{rank}(A)$ occurs exactly when the containing subspace has the same dimension as $\operatorname{col}(A)$, equivalently when $\operatorname{col}(A) + \operatorname{col}(C) = \operatorname{col}(A)$.
*Structural Commentary*: Products can only reduce accessible column directions, while horizontal concatenation can only add column directions.
*Key Insight*: Rank tracks whether a matrix operation preserves, loses, or adds column-space dimension. □

**Theorem 1.2.10.** *Sylvester Rank Inequality*
*Let $A \in \operatorname{Mat}(m,k,\mathbb{F})$ and $B \in \operatorname{Mat}(k,n,\mathbb{F})$. Then $\operatorname{rank}(A) + \operatorname{rank}(B) - k \leq \operatorname{rank}(AB)$.*

*Proof.* Let $R = \operatorname{col}(B) \subseteq \mathbb{F}^k$. The columns of $AB$ are $A$ applied to the columns of $B$, so $\operatorname{col}(AB) = A(R)$.
Apply rank-nullity to the restriction $T_A|_R : R \to \mathbb{F}^m$. Its range is $A(R) = \operatorname{col}(AB)$ and its kernel is $R \cap \ker(A)$.
Hence $\operatorname{rank}(AB) = \dim(A(R)) = \dim(R) - \dim(R \cap \ker(A)) \geq \operatorname{rank}(B) - \dim(\ker(A))$.
Since $A$ has $k$ columns, Corollary 3.3.3 gives $\dim(\ker(A)) = k - \operatorname{rank}(A)$. Substitution gives desired inequality.
*Structural Commentary*: Multiplication by $A$ can lose only the part of $\operatorname{col}(B)$ lying in $\ker(A)$. Sylvester's inequality is this loss estimate in dimensions.
*Key Insight*: The shared middle dimension $k$ bounds how much rank can disappear in the product. □

**Theorem 1.2.11.** *Frobenius Rank Inequality*
*Let $A \in \operatorname{Mat}(m,k,\mathbb{F})$, $B \in \operatorname{Mat}(k,p,\mathbb{F})$, and $C \in \operatorname{Mat}(p,n,\mathbb{F})$.*
*Then $\operatorname{rank}(AB) + \operatorname{rank}(BC) \leq \operatorname{rank}(B) + \operatorname{rank}(ABC)$.*

*Proof.* Let $R = \operatorname{col}(B)$ and $S = \operatorname{col}(BC)$. Since every column of $BC$ is a linear combination of columns of $B$, we have $S \subseteq R$. The induced map $T_A$ maps $R$ onto $\operatorname{col}(AB)$ and maps $S$ onto $\operatorname{col}(ABC)$.
Rank-nullity on the restrictions $T_A|_R$ and $T_A|_S$ gives

$$\operatorname{rank}(AB) = \dim R - \dim(R \cap \ker(A)), \quad \operatorname{rank}(ABC) = \dim S - \dim(S \cap \ker(A)).$$

Since $S \cap \ker(A) \subseteq R \cap \ker(A)$, we have $\dim(S \cap \ker(A)) \leq \dim(R \cap \ker(A))$.
Thus $\dim R - \dim(R \cap \ker(A)) + \dim S \leq \dim R + \dim S - \dim(S \cap \ker(A))$.
Substituting $\dim R = \operatorname{rank}(B)$ and $\dim S = \operatorname{rank}(BC)$ gives the stated inequality.
*Structural Commentary*: Frobenius tracks $A$ on nested column spaces.
*Key Insight*: The smaller column space cannot lose more dimensions to $\ker(A)$ than the larger one. □

**Corollary 1.2.12.** *Rank Inequalities from Frobenius*
*The following are special cases of Theorem 1.2.11:*

*(i)* $\operatorname{rank}(AB) \leq \min(\operatorname{rank}(A), \operatorname{rank}(B))$;
*(ii)* $\operatorname{rank}(A) + \operatorname{rank}(B) - k \leq \operatorname{rank}(AB)$.

*Proof.*

(i) Apply Frobenius to conformable factors $X, Y, Z$. Taking $Z = 0$ gives $\operatorname{rank}(XY) \leq \operatorname{rank}(Y)$, because $\operatorname{rank}(Y0) = \operatorname{rank}(XY0) = 0$. Taking $X = 0$ gives $\operatorname{rank}(YZ) \leq \operatorname{rank}(Y)$.
Renaming the factors gives $\operatorname{rank}(AB) \leq \operatorname{rank}(B)$ and $\operatorname{rank}(AB) \leq \operatorname{rank}(A)$.

(ii) Taking the middle matrix to be $I_k$ gives $\operatorname{rank}(A)+\operatorname{rank}(B) \le k+\operatorname{rank}(AB)$, which is Sylvester's inequality.

*Structural Commentary*: Frobenius is the three-factor parent inequality. The usual upper and lower product-rank bounds appear by making one factor zero or identity.
*Key Insight*: Zero factors isolate upper bounds; an identity middle factor yields the Sylvester lower bound. □

**Proposition 1.2.13.** *Sylvester's Law of Nullity*
*For a matrix $X$, let $\operatorname{nullity}(X) = \dim(\ker(X))$; this is the dimension of its nullspace.*

*(i) If $A \in \operatorname{Mat}(m,k,\mathbb{F})$ and $B \in \operatorname{Mat}(k,n,\mathbb{F})$, then $\operatorname{nullity}(AB) \le \operatorname{nullity}(A) + \operatorname{nullity}(B)$.*
*(ii) If $A, B \in \operatorname{Mat}(n,n,\mathbb{F})$, then $\max(\operatorname{nullity}(A), \operatorname{nullity}(B)) \le \operatorname{nullity}(AB)$.*

*Proof.*

(i) By rank-nullity and Theorem 1.2.10,

$$\begin{aligned}\operatorname{nullity}(AB) &= n - \operatorname{rank}(AB) \le n - \operatorname{rank}(A) - \operatorname{rank}(B) + k \\ &= (k - \operatorname{rank}(A)) + (n - \operatorname{rank}(B)) = \operatorname{nullity}(A) + \operatorname{nullity}(B).\end{aligned}$$

(ii) For square $A, B$, $\operatorname{rank}(AB) \le \operatorname{rank}(A)$, $\operatorname{rank}(AB) \le \operatorname{rank}(B)$. Subtract from $n$ gives desired lower bounds.

*Structural Commentary*: Nullity is rank loss from the domain.
*Key Insight*: Use rank-nullity to translate rank bounds into kernel bounds. □

**Theorem 1.2.14.** *Rank Triangle Inequality*
*If $A, B \in \operatorname{Mat}(m,n,\mathbb{F})$, then $|\operatorname{rank}(A) - \operatorname{rank}(B)| \le \operatorname{rank}(A+B) \le \operatorname{rank}(A) + \operatorname{rank}(B)$.*

*Proof.* Since $\operatorname{col}(A+B) \subseteq \operatorname{col}(A) + \operatorname{col}(B)$, Theorem 2.3.1 gives $\operatorname{rank}(A+B) \le \dim(\operatorname{col}(A) + \operatorname{col}(B)) \le \operatorname{rank}(A) + \operatorname{rank}(B)$. Applying this upper bound to $A = (A+B) - B$ gives $\operatorname{rank}(A) \le \operatorname{rank}(A+B) + \operatorname{rank}(B)$, so $\operatorname{rank}(A) - \operatorname{rank}(B) \le \operatorname{rank}(A+B)$. Applying it to $B = (A+B) - A$ gives $\operatorname{rank}(B) - \operatorname{rank}(A) \le \operatorname{rank}(A+B)$. Together these are exactly $|\operatorname{rank}(A) - \operatorname{rank}(B)| \le \operatorname{rank}(A+B)$.
*Structural Commentary*: The rank of a sum is controlled by the sum of the two column spaces. The reverse bounds come by solving for either summand.
*Key Insight*: Rank behaves like a dimension-valued size under addition. □

## 1.3 Commutators and Special Matrix Classes

**Definition 1.3.1.** *Commutator*
For $S, T \in \mathcal{L}(\mathcal{V})$, the *commutator* of $S$ and $T$ is $[S,T] := ST - TS$.
For $A, B \in \operatorname{Mat}(n,n,\mathbb{F})$, the *matrix commutator* is $[A,B] := AB - BA$.

**Proposition 1.3.2.** *Trace Obstruction for Commutators*
*If $A, B \in \operatorname{Mat}(n,n,\mathbb{F})$, then $\operatorname{tr}(AB - BA) = 0$. Hence any matrix similar to a commutator has trace zero.*

*Proof.* By linearity of trace and Proposition 1.1.5, $\operatorname{tr}(AB - BA) = \operatorname{tr}(AB) - \operatorname{tr}(BA) = 0$.
If $C$ is similar to $AB - BA$, then Theorem 1.1.17 gives $\operatorname{tr}(C) = \operatorname{tr}(AB - BA) = 0$.
*Structural Commentary*: Trace kills commutators because it is cyclic.
*Key Insight*: $AB$ and $BA$ have the same trace, so their difference has trace zero. □

**Corollary 1.3.3.** *The Identity Is Not a Commutator*
*Assume $\dim \mathcal{V} \cdot 1_{\mathbb{F}} \neq 0$. There do not exist $S, T \in \mathcal{L}(\mathcal{V})$ such that $ST - TS = I_{\mathcal{V}}$.*

*Proof.* If $ST - TS = I_{\mathcal{V}}$, choose a basis and let $A = [S]$ and $B = [T]$ in that basis. Then $AB - BA = I_n$.
Taking traces and using Proposition 1.1.5, $0 = \operatorname{tr}(AB) - \operatorname{tr}(BA) = \operatorname{tr}(I_n) = n \cdot 1_{\mathbb{F}}$, contradicting the hypothesis.
*Structural Commentary*: Commutators have zero trace, while the identity has dimension as its trace.
*Key Insight*: Trace detects a scalar obstruction invisible from multiplication alone. □

**Method 1.3.4.** *Trace–Determinant Invariant Method*
To get eigenvalue data without solving for eigenvalues:

(i) choose a basis in which $[T]_{\beta \leftarrow \beta}$ is easiest to compute;
(ii) compute $\operatorname{tr} T$ and $\det T$ from this matrix;
(iii) if the eigenvalues are $\lambda_1, \dots, \lambda_n$, conclude $\lambda_1 + \cdots + \lambda_n = \operatorname{tr} T$, and $\lambda_1 \cdots \lambda_n = \det T$;
(iv) for equations $ST - TS = A$, first test $\operatorname{tr} A = 0$; otherwise no solution exists.

**Lemma 1.3.5.** *Scalar Matrices and One-Step Orbits*
*Let $A \in \operatorname{Mat}(n,n,\mathbb{F})$ with $n \ge 2$.*
*Then $(\mathbf{x}, A\mathbf{x})$ is linearly dependent for every $\mathbf{x} \in \mathbb{F}^n$ if and only if $A$ is a scalar matrix.*

*Proof.*

($\Rightarrow$) Suppose $A = cI_n$. Then $A\mathbf{x} = c\mathbf{x}$ for every $\mathbf{x}$, so $(\mathbf{x}, A\mathbf{x})$ is linearly dependent.

($\Leftarrow$) Suppose $(\mathbf{x}, A\mathbf{x})$ is linearly dependent for every $\mathbf{x}$.
For each standard basis vector $\mathbf{e}_i$, $\exists a_i \in \mathbb{F}$ such that $A\mathbf{e}_i = a_i\mathbf{e}_i$. For $i \geq 2$, dependence of $\mathbf{e}_1 + \mathbf{e}_i$ and $A(\mathbf{e}_1 + \mathbf{e}_i) = a_1\mathbf{e}_1 + a_i\mathbf{e}_i$ means $a_1\mathbf{e}_1 + a_i\mathbf{e}_i = b(\mathbf{e}_1 + \mathbf{e}_i)$ for some $b \in \mathbb{F}$. Comparing the independent vectors $\mathbf{e}_1, \mathbf{e}_i$ gives $a_1 = b = a_i$. Hence every $a_i$ equals $a_1$, so $A = a_1 I_n$.

*Structural Commentary*: A scalar matrix preserves every one-dimensional coordinate direction and every diagonal line between two coordinates. These diagonal lines force all coordinate scalars to agree.
*Key Insight*: If every vector is an eigenvector, then there can be only one eigenvalue. □

**Lemma 1.3.6.** *Nonscalar Matrices Have a Zero Diagonal Entry After Similarity*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ with $n \geq 2$.*
*If $A$ is not a scalar matrix, then $A$ is similar over $\mathbb{F}$ to a matrix whose $(1,1)$ entry is $0$.*

*Proof.* By Lemma 1.3.5, there is $\mathbf{x} \in \mathbb{F}^n$ such that $(\mathbf{x}, A\mathbf{x})$ is linearly independent. Extend $(\mathbf{x}, A\mathbf{x})$ to a basis $\beta$ of $\mathbb{F}^n$. In the basis $\beta$, the first column of the matrix of $A$ is the coordinate vector of $A\mathbf{x}$, namely $(0, 1, 0, \ldots, 0)^t$. Thus the $(1,1)$ entry is 0, and this matrix is similar to $A$.
*Structural Commentary*: Nonscalarity produces a vector whose first iterate points in a genuinely new direction. Using these two vectors as the start of a basis forces a zero in the first diagonal position.
*Key Insight*: Put $\mathbf{x}$ first and $A\mathbf{x}$ second in the basis. □

**Theorem 1.3.7.** *Similarity to a Constant-Diagonal Matrix*
*Let $n \geq 1$, let $A \in \mathrm{Mat}(n, n, \mathbb{F})$, and assume $k \cdot 1_{\mathbb{F}} \neq 0$ for every $1 \leq k \leq n$. Then $A$ is similar over $\mathbb{F}$ to a matrix whose diagonal entries are all equal to $\frac{1}{n}\operatorname{tr}(A)$.*

*Proof.* Set $\tau = \frac{1}{n}\operatorname{tr}(A)$ and $B = A - \tau I_n$. Then $\operatorname{tr}(B) = 0$. It is enough to show that every trace-zero matrix is similar to a matrix with zero diagonal, because Theorem 6.2.10 then adds $\tau I_n$ back to the diagonal.
To prove the trace-zero claim by induction on $n$. If $n = 1$, a trace-zero matrix is $[0]$. Let $n \geq 2$ and assume the claim holds for smaller sizes. Let $\operatorname{tr}(B) = 0$. If $B$ is scalar, then $B = cI_n$ and $0 = \operatorname{tr}(B) = nc$.
Since $n \cdot 1_{\mathbb{F}} \neq 0$, we get $c = 0$, so $B$ already has zero diagonal.
If $B$ is not scalar, Lemma 1.3.6 makes $B$ similar to $\begin{bmatrix} 0 & * \\ * & B_1 \end{bmatrix}$, and $\operatorname{tr}(B_1) = 0$.
By induction, choose an invertible $S_1$ such that $S_1^{-1}B_1S_1$ has zero diagonal. Conjugating the displayed block matrix by $1 \oplus S_1$ leaves the first diagonal entry equal to 0 and replaces the lower-right block by $S_1^{-1}B_1S_1$.
Hence the resulting similar matrix has every diagonal entry equal to 0.
*Structural Commentary*: The proof removes one zero diagonal entry at a time.
*Key Insight*: Similarity creates the first zero; trace zero handles the smaller block. □

**Theorem 1.3.8.** *Shoda's Theorem in the Distinct-Diagonal Case*
*Let $n \geq 1$ and assume $k \cdot 1_{\mathbb{F}} \neq 0$ for every $1 \leq k \leq n$. For $A \in \mathrm{Mat}(n, n, \mathbb{F})$, the matrix $A$ is a commutator in $\mathrm{Mat}(n, n, \mathbb{F})$ if and only if $\operatorname{tr}(A) = 0$. In particular, no nonzero scalar matrix is a commutator.*

*Proof.*

($\Rightarrow$) Proposition 1.3.2 gives $\operatorname{tr}(A) = 0$.
Thus $cI_n$ can be a commutator only when $nc = \operatorname{tr}(cI_n) = 0$, hence only when $c = 0$.

($\Leftarrow$) Suppose $\operatorname{tr}(A) = 0$. By Theorem 1.3.7, there is an invertible $S$ such that $B = S^{-1}AS = [b_{ij}]$ has zero diagonal. Let $X = \operatorname{diag}(1, 2, \ldots, n)$, viewing the integers as scalars in $\mathbb{F}$, and define $Y = [y_{ij}]$ by

$$y_{ij} = \begin{cases} (i-j)^{-1}b_{ij}, & i \neq j, \\ 0, & i = j. \end{cases}$$

The hypothesis ensures that $i - j \neq 0$ in $\mathbb{F}$ whenever $i \neq j$. Thus $XY - YX = [(i-j)y_{ij}] = B$.
Hence $A = (SXS^{-1})(SYS^{-1}) - (SYS^{-1})(SXS^{-1})$, so $A$ is a commutator.

*Structural Commentary*: Trace is the obstruction; similarity exposes zero diagonal form.
*Key Insight*: Cyclicity forces trace zero; distinct diagonals realize it. □

**Remark 1.3.9.** *Field Hypothesis in the Diagonal Proof*
The proof above uses that the integers $1, \ldots, n$ are distinct and that each smaller block size remains nonzero in $\mathbb{F}$. In small characteristic, this diagonal proof must be replaced by a more delicate argument.

**Definition 1.3.10.** *Idempotent and Nilpotent Matrices*
Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$.

(i) $A$ is *idempotent* if $A^2 = A$.

(ii) $A$ is *nilpotent* if $A^k = 0$ for some positive integer $k$.

An idempotent matrix is an algebraic projection: it fixes its range and kills its kernel.

**Proposition 1.3.11.** *Basic Invariants of Idempotents*
*If $P \in \mathrm{Mat}(n,n,\mathbb{F})$ satisfies $P^2 = P$, then $\mathbb{F}^n = \mathrm{range}(P) \oplus \ker(P)$, $P$ is diagonalizable, and its only possible eigenvalues are $0$ and $1$. Moreover $\mathrm{tr}(P) = \mathrm{rank}(P)$.*
*If $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, then $P$ is the orthogonal projection onto $\mathrm{range}(P)$ if and only if $P = P^*$.*

*Proof.* For every $\mathbf{x}$, write $\mathbf{x} = P\mathbf{x} + (\mathbf{x} - P\mathbf{x})$. The first term lies in $\mathrm{range}(P)$, while $P(\mathbf{x} - P\mathbf{x}) = P\mathbf{x} - P^2\mathbf{x} = 0$, so the second term lies in $\ker(P)$. The intersection is zero because $\mathbf{x} = P\mathbf{y}$ and $P\mathbf{x} = 0$ imply $\mathbf{x} = P^2\mathbf{y} = P\mathbf{x} = 0$.
In a basis adapted to $\mathrm{range}(P) \oplus \ker(P)$, the matrix of $P$ is $I_r \oplus 0$, where $r = \mathrm{rank}(P)$.
Hence $P$ is diagonalizable, its eigenvalues lie in $\{0, 1\}$, and $\mathrm{tr}(P) = r$.
For the final claim, self-adjoint idempotents are exactly orthogonal projections by Theorem 2.5.6.
*Structural Commentary*: Idempotents split the space into fixed directions and killed directions.
*Key Insight*: In adapted coordinates, every idempotent is just $I_r \oplus 0$. □

# 2 Positive Definite and Structured Matrices

## 2.1 Positive Semidefinite and Positive Definite Matrices

**Definition 2.1.1.** *Positive Semidefinite and Positive Definite Matrices*
Let $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$ and let $A \in \mathrm{Mat}(n,n,\mathbb{F})$.
The matrix $A$ is *positive semidefinite*, written $A \succeq 0$, if $A = A^*$ and $\mathbf{x}^* A\mathbf{x} \geq 0$ for every $\mathbf{x} \in \mathbb{F}^n$.
It is *positive definite*, written $A \succ 0$, if $A = A^*$ and $\mathbf{x}^* A\mathbf{x} > 0$ for every nonzero $\mathbf{x} \in \mathbb{F}^n$.
Negative semidefinite and negative definite mean $-A \succeq 0$ and $-A \succ 0$, respectively.

**Proposition 2.1.2.** *Rectangular Gram Products Are Positive Semidefinite*
*Let $R \in \mathrm{Mat}(m,n,\mathbb{F})$, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. Then $R^*R \succeq 0$, $RR^* \succeq 0$, and $(R^*R)_{j,j} = \|R_{\cdot,j}\|_2^2$; $(RR^*)_{i,i} = \|R_{i,\cdot}\|_2^2$.*
*Full column rank gives $R^*R \succ 0$; full row rank gives $RR^* \succ 0$.*

*Proof.* The matrices are self-adjoint by Proposition 2.4.8.
For $\mathbf{x} \in \mathbb{F}^n$ and $\mathbf{y} \in \mathbb{F}^m$, $\mathbf{x}^* R^* R\mathbf{x} = \|R\mathbf{x}\|_2^2 \geq 0$ and $\mathbf{y}^* RR^*\mathbf{y} = \|R^*\mathbf{y}\|_2^2 \geq 0$. Hence both products are positive semidefinite. Taking $\mathbf{x} = \mathbf{e}_j$ and $\mathbf{y} = \mathbf{e}_i$ gives the diagonal formulas. If $R$ has full column rank, then $R\mathbf{x} \neq 0$ for $\mathbf{x} \neq 0$, so $R^*R \succ 0$. If $R$ has full row rank, apply the same argument to $R^*$.
*Structural Commentary*: Multiplying a matrix by its adjoint produces a Gram matrix of columns or rows.
*Key Insight*: The quadratic form is a squared norm, so no cancellation can make it negative. □

**Theorem 2.1.3.** *Equivalent Forms of Positive Semidefiniteness*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{F})$, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. The following are equivalent:*

*(i)* $A \succeq 0$.
*(ii)* *$A$ is Hermitian and all eigenvalues of $A$ are nonnegative.*
*(iii)* $A = B^*B$ *for some* $B \in \mathrm{Mat}(m,n,\mathbb{F})$.
*(iv)* *$A$ has a positive semidefinite square root.*

*In this case the positive semidefinite square root is unique.*
*If $A = U\Lambda U^*$ with $\Lambda = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$ and $\lambda_i \geq 0$, then $A^{1/2} = U\,\mathrm{diag}(\sqrt{\lambda_1}, \ldots, \sqrt{\lambda_n})U^*$.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* This is the matrix form of Theorem 6.3.6.
*(ii)* $\Rightarrow$ *(iv)* By the spectral theorem, write $A = U\Lambda U^*$, where $U$ is unitary and $\Lambda = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$ with $\lambda_i \geq 0$. Then $A^{1/2} = U\,\mathrm{diag}(\sqrt{\lambda_1}, \ldots, \sqrt{\lambda_n})U^*$ is positive semidefinite and satisfies $(A^{1/2})^2 = A$.
*(iv)* $\Rightarrow$ *(iii)* If $C \succeq 0$ and $C^2 = A$, then $C = C^*$ and $A = C^*C$, so (iii) holds with $m = n$.
*(iii)* $\Rightarrow$ *(i)* If $A = B^*B$, then $A^* = A$ and $\mathbf{x}^* A\mathbf{x} = \mathbf{x}^* B^* B\mathbf{x} = \|B\mathbf{x}\|_2^2 \geq 0$, so $A \succeq 0$.

The positive semidefinite square root is unique by Proposition 6.3.7.
*Structural Commentary*: Positive semidefinite matrices are Gram-type matrices with nonnegative spectrum.
*Key Insight*: Positivity is best checked either on eigenvalues or by finding a factor $B$ with $A = B^*B$. □

**Theorem 2.1.4.** *Kernel, Rank, and Columns of $B^*B$*
*Let $B \in \mathrm{Mat}(m,n,\mathbb{F})$, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, and set $A = B^*B$. Then:*

*(i)* $A \succeq 0$;
*(ii)* $\ker(A) = \ker(B)$;
*(iii)* $\mathrm{rank}(A) = \mathrm{rank}(B)$;

*(iv)* $\operatorname{col}(A) = \operatorname{col}(B^*)$;
*(v)* $A = 0$ *if and only if* $B = 0$.

*Proof.*

(i) Positive semidefiniteness follows from Theorem 2.1.3.
(ii) If $B\mathbf{x} = 0$, then $A\mathbf{x} = 0$. Conversely, if $A\mathbf{x} = 0$, then $0 = \mathbf{x}^* A\mathbf{x} = \|B\mathbf{x}\|_2^2$ so $B\mathbf{x} = 0$.
(iii) Since $A$ and $B$ both have $n$ columns, rank-nullity and (ii) give $\operatorname{rank}(A) = \operatorname{rank}(B)$.
(iv) Since $A = A^*$, Proposition 2.4.11 and (ii) give $\operatorname{col}(A) = \ker(A)^\perp = \ker(B)^\perp = \operatorname{col}(B^*)$.
(v) By (iii), $\operatorname{rank}(A) = 0$ if and only if $\operatorname{rank}(B) = 0$.
A matrix has rank zero exactly when it is the zero matrix, so $A = 0$ if and only if $B = 0$.

*Structural Commentary*: The Gram product $B^*B$ preserves the kernel of $B$.
*Key Insight*: The identity $\mathbf{x}^* B^* B\mathbf{x} = \|B\mathbf{x}\|_2^2$ is the whole mechanism. □




Figure 80: For $A = B^*B$, the squared-norm identity gives $\ker A = \ker B$.

**Corollary 2.1.5.** *Zero Quadratic Form Detects the Kernel*
*If* $A \in \operatorname{Mat}(n, n, \mathbb{F})$ *is positive semidefinite, where* $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, *then* $\mathbf{x}^* A\mathbf{x} = 0$ *if and only if* $A\mathbf{x} = 0$.

*Proof.* Write $A = B^*B$.

($\Rightarrow$) If $\mathbf{x}^* A\mathbf{x} = 0$, then $\|B\mathbf{x}\|_2^2 = 0$, so $B\mathbf{x} = 0$ and hence $A\mathbf{x} = B^*B\mathbf{x} = 0$.
($\Leftarrow$) If $A\mathbf{x} = 0$, then $\mathbf{x}^* A\mathbf{x} = 0$.

*Structural Commentary*: A zero PSD quadratic value has no hidden cancellation.
*Key Insight*: Zero length-square means the factored vector itself is zero. □

**Method 2.1.6.** *PSD Spectral Factorisation Method*
To analyse a Hermitian matrix $A$:

(i) diagonalise $A = U\Lambda U^*$;
(ii) check the signs of the diagonal entries of $\Lambda$;
(iii) if $\Lambda \geq 0$, set $A^{1/2} = U\Lambda^{1/2}U^*$ and use $A = (A^{1/2})^* A^{1/2}$;
(iv) translate kernel, rank, and zero-quadratic-form questions into the factor $A^{1/2}$.

**Theorem 2.1.7.** *Equivalent Forms of Positive Definiteness*
*Let* $A \in \operatorname{Mat}(n, n, \mathbb{F})$, *where* $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. *The following are equivalent:*

*(i)* $A \succ 0$.
*(ii)* $A$ *is Hermitian and all eigenvalues of* $A$ *are positive.*
*(iii)* $A \succeq 0$ *and* $A$ *is invertible.*
*(iv)* $A = B^*B$ *for some invertible* $B \in \operatorname{Mat}(n, n, \mathbb{F})$.
*(v)* $A$ *is invertible and* $A^{-1} \succ 0$.

*Proof.*

*(i)* $\Rightarrow$ *(ii)* By definition $A$ is Hermitian. If $(\lambda, \mathbf{v})$ is an eigenpair with $\mathbf{v} \neq \mathbf{0}$, then $\lambda \|\mathbf{v}\|_2^2 = \mathbf{v}^* A\mathbf{v} > 0$, so every eigenvalue is positive.
*(ii)* $\Rightarrow$ *(iii)* Positive eigenvalues imply $A \succeq 0$ by Theorem 2.1.3; they also exclude 0, so $A$ is invertible.
*(iii)* $\Rightarrow$ *(iv)* By Theorem 2.1.3, $A$ has a positive semidefinite square root $A^{1/2}$.
Since $A$ is invertible, $A^{1/2}$ is invertible, and $A = (A^{1/2})^* A^{1/2}$.
*(iv)* $\Rightarrow$ *(i)* If $A = B^*B$ and $B$ is invertible, then for every nonzero $\mathbf{x}$, $\mathbf{x}^* A\mathbf{x} = \|B\mathbf{x}\|_2^2 > 0$.
*(ii)* $\Leftrightarrow$ *(v)* Hermitian-ness is preserved by inversion, and eigenvalues of $A^{-1}$ are reciprocal eigenvalues of $A$.
Thus positive eigenvalues of $A$ are equivalent to positive eigenvalues of $A^{-1}$.

*Structural Commentary*: Positive definite matrices are the invertible part of the positive semidefinite cone.
*Key Insight*: Positive definite means every eigenvalue is strictly on the positive side of zero. □

**Proposition 2.1.8.** *Trace and Determinant of Positive Semidefinite Matrices*
*Let* $A \in \operatorname{Mat}(n, n, \mathbb{F})$ *be positive semidefinite, where* $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. *Then* $\operatorname{tr} A \geq 0$, *with equality if and only if* $A = 0$, *and* $\det A \geq 0$, *with strict inequality if and only if* $A$ *is positive definite.*

*Proof.* List the eigenvalues of $A$ as $\lambda_1, \dots, \lambda_n$. By Theorem 2.1.3, each $\lambda_j \geq 0$.
Hence $\operatorname{tr} A = \sum_{j=1}^{n} \lambda_j \geq 0$ and $\det A = \prod_{j=1}^{n} \lambda_j \geq 0$.
The trace is zero exactly when every $\lambda_j = 0$, which for a Hermitian matrix means $A = 0$. The determinant is positive exactly when every $\lambda_j > 0$, which is positive definiteness by Theorem 2.1.7.
*Structural Commentary*: Trace and determinant aggregate the nonnegative eigenvalues by sum and product.
*Key Insight*: Equality is controlled by which eigenvalues vanish. □

**Theorem 2.1.9.** *Stability Rules for Positive Semidefinite Matrices*
*Let $A, B \succeq 0$ over $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. Whenever the expressions are defined:*

*(i) $aA + bB \succeq 0$ for $a, b \geq 0$; if $a, b > 0$ and one of $A, B$ is positive definite, then $aA + bB \succ 0$.*
*(ii) $S^*AS \succeq 0$; moreover, $S^*AS \succ 0$ if and only if $S\mathbf{x} \notin \ker(A)$ for all $\mathbf{x} \neq \mathbf{0}$.*
*(iii) $A \oplus B \succeq 0$, with $A \oplus B \succ 0$ if and only if $A \succ 0$ and $B \succ 0$.*

*Proof.*

(i) $\mathbf{x}^*(aA + bB)\mathbf{x} = a\mathbf{x}^*A\mathbf{x} + b\mathbf{x}^*B\mathbf{x} \geq 0$. If, for example, $A \succ 0$ and $a > 0$, then for every nonzero $\mathbf{x}$ the first summand is positive and the second is nonnegative, so $aA + bB \succ 0$; the case $B \succ 0$ is identical.
(ii) $\mathbf{x}^*S^*AS\mathbf{x} = (S\mathbf{x})^*A(S\mathbf{x}) \geq 0$. By Corollary 2.1.5, equality occurs exactly when $S\mathbf{x} \in \ker(A)$.
(iii) $(\mathbf{x} \oplus \mathbf{y})^*(A \oplus B)(\mathbf{x} \oplus \mathbf{y}) = \mathbf{x}^*A\mathbf{x} + \mathbf{y}^*B\mathbf{y} \geq 0$. This expression is positive for every nonzero $\mathbf{x} \oplus \mathbf{y}$ exactly when each summand is positive on its own nonzero vectors, namely when $A \succ 0$ and $B \succ 0$.

*Structural Commentary*: The PSD cone is stable under sums, pullbacks, and direct sums.
*Key Insight*: Test the quadratic form after each construction. □

**Proposition 2.1.10.** *Explicit Square Root for a $2 \times 2$ Positive Semidefinite Matrix*
*Let $A \in \operatorname{Mat}(2, 2, \mathbb{F})$, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, be positive semidefinite and nonzero. Set $\tau = \left(\operatorname{tr} A + 2\sqrt{\det A}\right)^{1/2}$. Then $\tau > 0$ and $A^{1/2} = \frac{1}{\tau}\left(A + \sqrt{\det A}\, I_2\right)$.*

*Proof.* Let $C = \frac{1}{\tau}\left(A + \sqrt{\det A}\, I_2\right)$. By Proposition 2.1.8, $\det A \geq 0$ and $\operatorname{tr} A > 0$, hence $\tau > 0$; by Theorem 2.1.9, $C \succeq 0$. Cayley–Hamilton gives $A^2 - (\operatorname{tr} A)A + (\det A)I_2 = 0$, so

$$C^2 = \frac{1}{\tau^2}\left(A^2 + 2\sqrt{\det A}\, A + (\det A)I_2\right) = \frac{1}{\tau^2}\left((\operatorname{tr} A)A + 2\sqrt{\det A}\, A\right) = A.$$

Thus $C = A^{1/2}$ by uniqueness of the positive semidefinite square root.
*Structural Commentary*: In dimension 2, Cayley–Hamilton makes $A^{1/2}$ linear in $A$.
*Key Insight*: Square the linear candidate and use Cayley–Hamilton. □

**Definition 2.1.11.** *Directed Incidence Matrix*
For edges $e = u \to v$ on vertices $1, \dots, n$, $B \in \operatorname{Mat}(m, n, \mathbb{R})$ has row $e$ equal to $-\mathbf{e}_u^t + \mathbf{e}_v^t$.




$$B = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 \\ -1 & 0 & 0 & 1 \end{bmatrix}$$

Figure 81: A directed incidence matrix records oriented edge differences from vertices to edges.

**Example 2.1.12.** *A Directed Incidence Matrix*
For Figure 81, $B = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 \\ -1 & 0 & 0 & 1 \end{bmatrix}$. For node potentials $\mathbf{x} = (x_1, x_2, x_3, x_4)^t$, $B\mathbf{x} = \begin{bmatrix} x_2 - x_1 \\ x_3 - x_2 \\ x_3 - x_4 \\ x_4 - x_1 \end{bmatrix}$.
Hence $\ker(B) = \operatorname{span}\{(1, 1, 1, 1)^t\}$, and $\operatorname{rank}(B) = 3$. The loop vector $\mathbf{y} = (1, 1, -1, -1)^t$ spans $\ker(B^t)$.
Thus $\mathbf{b} \in \operatorname{col}(B)$ if and only if $\mathbf{y}^t\mathbf{b} = b_1 + b_2 - b_3 - b_4 = 0$. This is Kirchhoff loop compatibility.
*Structural Commentary*: Incidence subspaces encode graph constraints.
*Key Insight*: A sparse matrix can encode both the algebra of a linear map and the combinatorics of a graph.

**Perspective 2.1.13.** *Incidence Data and Least Squares*
For an incidence matrix $B$ as in Example 2.1.12, $B\mathbf{x} = \mathbf{b}$ seeks node potentials matching edge differences. If loop consistency fails, least squares projects $\mathbf{b}$ onto $\operatorname{col}(B)$, where $B^*(B\mathbf{x}_0 - \mathbf{b}) = \mathbf{0}$ (Theorem 2.3.4 on $B$). The constant-vector kernel means potentials are unique only after fixing one node or choosing minimum norm.

**Proposition 2.1.14.** *Incidence Laplacians Are Positive Semidefinite*
*Let $B$ be a directed incidence matrix and set $L = B^tB$. Then $L \succeq 0$ and $\mathbf{x}^tL\mathbf{x} = \sum_{e=(u\to v)} (x_v - x_u)^2$, with $\ker(L)$ equal to the vectors constant on each connected component.*

*Proof.* Since $L = B^tB$, Proposition 2.1.2 gives $L \succeq 0$.
For $e = (u \to v)$, $(B\mathbf{x})_e = x_v - x_u$, so $\mathbf{x}^tL\mathbf{x} = \|B\mathbf{x}\|_2^2 = \sum_{e=(u\to v)} (x_v - x_u)^2$. Thus $\mathbf{x} \in \ker(L)$ if and only if $B\mathbf{x} = \mathbf{0}$, equivalently $x_u = x_v$ on every edge, which means $\mathbf{x}$ is constant on each connected component.
*Structural Commentary*: Laplacians sum squared edge differences.
*Key Insight*: A graph Laplacian is a Gram product, so positivity is automatic. □

## 2.2 Principal Minors and Inertia

**Definition 2.2.1.** *Principal and Leading Principal Minors*
For $A \in \mathrm{Mat}(n, n, \mathbb{C})$, a *principal minor* is the determinant of a principal submatrix of $A$.
A *leading principal minor* is the determinant of the leading $k \times k$ principal submatrix for some $k = 1, \dots, n$.

**Corollary 2.2.2.** *Principal Blocks and Column Inclusion*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ be PSD, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, and partition $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$, $A_{11} \in \mathrm{Mat}(k, k, \mathbb{F})$.*
*Then $A_{11} \succeq 0$, $\mathrm{col}(A_{12}) \subseteq \mathrm{col}(A_{11})$, and $\mathrm{rank}(A_{12}) \le \mathrm{rank}(A_{11})$. If $A \succ 0$, then $A_{11} \succ 0$.*

*Proof.* Write $A = B^*B$ and partition $B = [B_1 \ B_2]$ conformally. Then $A_{11} = B_1^*B_1$ and $A_{12} = B_1^*B_2$.
Hence $A_{11} \succeq 0$ and $\mathrm{col}(A_{12}) = \mathrm{col}(B_1^*B_2) \subseteq \mathrm{col}(B_1^*) = \mathrm{col}(B_1^*B_1) = \mathrm{col}(A_{11})$, equality $\mathrm{col}(B_1^*) = \mathrm{col}(B_1^*B_1)$ follows from Theorem 2.1.4. The rank inequality follows by taking dimensions
If $A \succ 0$ and $\mathbf{x} \neq 0$ in $\mathbb{F}^k$, then $\mathbf{x}^*A_{11}\mathbf{x} = \begin{bmatrix} \mathbf{x} \\ 0 \end{bmatrix}^* A \begin{bmatrix} \mathbf{x} \\ 0 \end{bmatrix} > 0$, so $A_{11} \succ 0$.
*Structural Commentary*: In a positive block matrix, cross-block columns lie in principal block's column space.
*Key Insight*: Factor first; then column inclusion is immediate from $A_{12} = B_1^*B_2$. □




Figure 82: For $A \succeq 0$, factorisation forces $\mathrm{col}(A_{12}) \subseteq \mathrm{col}(A_{11})$.

**Theorem 2.2.3.** *Sylvester's Criterion*
*A Hermitian matrix $A \in \mathrm{Mat}(n, n, \mathbb{C})$ is positive definite if and only if all leading principal minors are positive.*

*Proof.*
($\Rightarrow$) If $A \succ 0$, then each leading principal submatrix is positive definite by Corollary 2.2.2. Its eigenvalues are positive, so its determinant is positive.
($\Leftarrow$) Induct on $n$. If $n = 1$, the single leading principal minor is the only diagonal entry, so positivity of that minor is exactly positive definiteness. Assume $n \ge 2$ and write $A = \begin{bmatrix} B & \mathbf{y} \\ \mathbf{y}^* & a \end{bmatrix}$, $B \in \mathrm{Mat}(n-1, n-1, \mathbb{C})$
The leading principal minors of $B$ are positive, so $B \succ 0$.
Since $\det(B) > 0$ and $\det(A) > 0$, Proposition 1.1.12 gives $a - \mathbf{y}^*B^{-1}\mathbf{y} = \frac{\det(A)}{\det(B)} > 0$.
For $\mathbf{u} \in \mathbb{C}^{n-1}$ and $t \in \mathbb{C}$, $\begin{bmatrix} \mathbf{u} \\ t \end{bmatrix}^* A \begin{bmatrix} \mathbf{u} \\ t \end{bmatrix} = (\mathbf{u} + B^{-1}\mathbf{y}t)^*B(\mathbf{u} + B^{-1}\mathbf{y}t) + \left(a - \mathbf{y}^*B^{-1}\mathbf{y}\right)|t|^2$.
First term is nonnegative as $B \succ 0$, and second coefficient is positive. If displayed vector is nonzero, then either $t \neq 0$ and second term is positive, or $t = 0$ and then $\mathbf{u} \neq 0$, making first term positive. Thus $A \succ 0$.
*Structural Commentary*: Leading minors make each leading block positive and final Schur complement positive.
*Key Insight*: Complete the square with the leading block. □

**Method 2.2.4.** $2 \times 2$ *Quadratic Form Test*
Let $A = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$ be real symmetric, $\Delta = ac - b^2$, and $q(x, y) = \begin{bmatrix} x & y \end{bmatrix} A \begin{bmatrix} x \\ y \end{bmatrix} = ax^2 + 2bxy + cy^2$.

For $a \neq 0$, $q(x,y) = a\left(x + \frac{b}{a}y\right)^2 + \frac{\Delta}{a}y^2$.
If $a = 0$, swap variables when $c \neq 0$; if $a = c = 0$, use $q = 2bxy$. Thus:

(i) $q \succ 0$ if and only if $a > 0$ and $\Delta > 0$;
(ii) $q \prec 0$ if and only if $a < 0$ and $\Delta > 0$;
(iii) $q$ is indefinite if and only if $\Delta < 0$;
(iv) if $\Delta = 0$, then $q \succeq 0$ iff $a, c \geq 0$, and $q \preceq 0$ iff $a, c \leq 0$.

This is the $2 \times 2$ form of Sylvester's criterion and inertia.
*Structural Commentary*: Completing the square is the $2 \times 2$ shadow of the $LDL^*$ and Schur-complement tests.
*Key Insight*: $\Delta$ and the leading sign classify the form.

**Theorem 2.2.5.** *Principal Axes for Real Quadratic Forms*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{R})$ be symmetric. Then:*

*(i) there are a real orthogonal $Q$ and real $\lambda_1, \ldots, \lambda_n$ such that, with $\mathbf{y} = Q^t\mathbf{x}$, $\mathbf{x}^t A\mathbf{x} = \sum_{j=1}^{n} \lambda_j y_j^2$.*
*(ii) $A \succeq 0$ if and only if every $\lambda_j \geq 0$;*
*(iii) $A \succ 0$ if and only if every $\lambda_j > 0$;*
*(iv) $A$ is indefinite if and only if some $\lambda_j > 0$ and some $\lambda_k < 0$.*

*Proof.*

(i) By the real spectral theorem, write $A = Q\,\mathrm{diag}(\lambda_1, \ldots, \lambda_n)Q^t$.
Then $\mathbf{x}^t A\mathbf{x} = \mathbf{x}^t Q\,\mathrm{diag}(\lambda_j)Q^t\mathbf{x} = \mathbf{y}^t\,\mathrm{diag}(\lambda_j)\mathbf{y} = \sum_{j=1}^{n} \lambda_j y_j^2$.
(ii) The displayed sum is nonnegative for all $\mathbf{y}$ exactly when each coefficient $\lambda_j$ is nonnegative.
(iii) The displayed sum is positive for all nonzero $\mathbf{y}$ exactly when each coefficient $\lambda_j$ is positive.
(iv) The diagonal form takes both signs exactly when its coefficients include both signs.

*Structural Commentary*: Principal axes turn the form into signed squares.
*Key Insight*: Eigenvalue signs classify the quadratic form. □

**Method 2.2.6.** *Quadratic Form Principal-Axes Method*
To analyse a real quadratic form:

(i) write it as $\mathbf{x}^t A\mathbf{x}$ with $A = A^t$; split each cross coefficient equally between $a_{ij}$ and $a_{ji}$;
(ii) diagonalise $A = Q\Lambda Q^t$;
(iii) change variables by $\mathbf{y} = Q^t\mathbf{x}$;
(iv) classify the form from the signs of the eigenvalues in $\Lambda$.




Figure 83: Orthogonal coordinates reveal the principal axes of a quadratic form.

**Definition 2.2.7.** *$*$-Congruence and Hermitian Inertia*
Two matrices $A, B \in \mathrm{Mat}(n, n, \mathbb{C})$ are *$*$-congruent* if $A = SBS^*$ for some invertible $S \in \mathrm{Mat}(n, n, \mathbb{C})$.
For Hermitian $A$, its *inertia* is $\mathrm{in}(A) = (n_+(A), n_-(A), n_0(A))$, where these are the numbers of positive, negative, and zero eigenvalues of $A$, counted with multiplicity.

**Lemma 2.2.8.** *Positive Subspaces and Inertia*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ be Hermitian. The maximal dimensions of subspaces on which $\mathbf{x}^* A\mathbf{x}$ is positive or negative for every nonzero $\mathbf{x}$ are $n_+(A)$ and $n_-(A)$, respectively.*

*Proof.* Let $P_+$, $P_{\leq 0}$ be positive, nonpositive spectral subspaces. Then $\dim P_+ = n_+(A)$, $P_+$ is positive for $A$. If $\dim \mathcal{U} > n_+(A)$, then $\dim \mathcal{U} + \dim P_{\leq 0} > n_+(A) + n_-(A) + n_0(A) = n$, so $\mathcal{U} \cap P_{\leq 0}$ contains a nonzero vector. On this vector quadratic form is nonpositive, so $\mathcal{U}$ cannot be positive. Apply this to $-A$ for negative case.

*Structural Commentary*: The positive spectral subspace is maximal; the nonpositive part blocks enlargement.
*Key Insight*: Inertia counts maximal sign-definite dimensions. □

**Theorem 2.2.9.** *Sylvester's Inertia Theorem*
*Hermitian matrices $A, B$ of the same size are $*$-congruent if and only if $n_+(A) = n_+(B)$ and $n_-(A) = n_-(B)$.*

*Proof.*
($\Rightarrow$) Suppose $A = SBS^*$ with $S$ invertible. Then $\mathbf{x}^*A\mathbf{x} = (S^*\mathbf{x})^*B(S^*\mathbf{x})$.
Thus $S^*$ transports positive and negative subspaces, so Lemma 2.2.8 gives $n_\pm(A) = n_\pm(B)$.
($\Leftarrow$) Let the common inertia be $(p, q, r)$ and set $D = I_p \oplus (-I_q) \oplus 0_r$.
By the spectral theorem, each of $A$ and $B$ is $*$-congruent to $D$, hence they are $*$-congruent to each other.

*Structural Commentary*: $*$-congruence is a coordinate change for Hermitian quadratic forms.
*Key Insight*: Spectral decomposition reduces the form to signed squares and zero directions. □




Figure 84: $*$-congruence reduces a Hermitian form to signed coordinate axes.

## 2.3 Hermitian and Normal Matrix Structure

**Theorem 2.3.1.** *Simultaneous Congruence with One Positive Definite Form*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$ be Hermitian, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, and let $A \succ 0$.*
*Then, for some invertible $S$ and real diagonal $\Lambda$, $A = SS^*$ and $B = S\Lambda S^*$.*
*The diagonal entries of $\Lambda$ are the eigenvalues of $A^{-1}B$.*

*Proof.* Let $C = A^{-1/2}BA^{-1/2} = U\Lambda U^*$ be spectral decomposition, $S = A^{1/2}U$. Then $SS^* = A$ and $S\Lambda S^* = B$.
Also $A^{-1}B = A^{-1/2}CA^{1/2}$, so $A^{-1}B$ is similar to $C$ and has eigenvalues given by the diagonal entries of $\Lambda$.
*Structural Commentary*: The positive definite form $A$ normalizes $B$ to a Hermitian matrix.
*Key Insight*: Normalize, diagonalise, then transform back. □

**Corollary 2.3.2.** *Products with One Positive Definite Factor*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, and let $A \succ 0$.*

*(i) If $B$ is Hermitian, then $AB$ is diagonalizable with real spectrum.*
*(ii) If $B \succeq 0$, then $AB$ is diagonalizable with nonnegative real spectrum; if $B \neq 0$, one eigenvalue is positive.*
*(iii) If $B \succ 0$, then $AB$ is diagonalizable with positive real spectrum.*

*Proof.* Let $C = A^{1/2}BA^{1/2}$. Since $A^{-1/2}(AB)A^{1/2} = C$, the matrices $AB$ and $C$ are similar.

(i) If $B$ is Hermitian, then $C$ is Hermitian, hence diagonalizable with real spectrum.
(ii) If $B \succeq 0$, then $C \succeq 0$ by Theorem 2.1.9, so its spectrum is nonnegative.
If $B \neq 0$, then $C \neq 0$, hence some eigenvalue is positive.
(iii) If $B \succ 0$, then $C \succ 0$ by Theorem 2.1.9, so all eigenvalues are positive.

*Structural Commentary*: A positive definite factor makes $AB$ similar to a Hermitian congruence.
*Key Insight*: Use $AB \sim A^{1/2}BA^{1/2}$. □

**Definition 2.3.3.** *Phase Inertia Matrix*
A *phase inertia matrix* is a diagonal complex matrix whose nonzero diagonal entries have modulus 1.

**Proposition 2.3.4.** *Phase Inertia Form for Normal Matrices*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ be normal.*
*Then $A$ is $*$-congruent to a phase inertia matrix $D$, whose entries record the spectral rays of $A$.*

*Proof.* Write $A = U\,\mathrm{diag}(\lambda_j)U^*$. Write $\lambda_j = r_j d_j$, where $r_j > 0$ and $|d_j| = 1$ if $\lambda_j \neq 0$, while $r_j = 1$ and $d_j = 0$ if $\lambda_j = 0$. Set $D = \mathrm{diag}(d_j)$, $S = U\,\mathrm{diag}(\sqrt{r_j})$. Then $S$ is invertible and $A = SDS^*$.
The entries $d_j = e^{i\theta}$ count exactly the eigenvalues on the ray $\{\rho e^{i\theta} : \rho > 0\}$.
*Structural Commentary*: Normal diagonalization separates modulus from phase; $*$-congruence absorbs moduli.
*Key Insight*: Only spectral phases survive. □

**Perspective 2.3.5.** *Hermitian Inertia as Real Phase Inertia*
For Hermitian matrices, the only nonzero spectral rays are the positive and negative real axes.
Sylvester's inertia theorem is therefore the real-ray version of phase inertia.

# 3 Matrix Products, Vectorization, and Spectral Radius

## 3.1 Kronecker Products and Vectorization

**Definition 3.1.1.** *Kronecker Product*
Let $A = [a_{ij}] \in \mathrm{Mat}(m, n, \mathbb{F})$ and $B \in \mathrm{Mat}(p, q, \mathbb{F})$. The *Kronecker product* of $A$ and $B$ is the block matrix

$$A \otimes B := \begin{bmatrix} a_{11}B & a_{12}B & \cdots & a_{1n}B \\ a_{21}B & a_{22}B & \cdots & a_{2n}B \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}B & a_{m2}B & \cdots & a_{mn}B \end{bmatrix} \in \mathrm{Mat}(mp, nq, \mathbb{F}).$$

The Kronecker product is also called the tensor product of matrices.

**Proposition 3.1.2.** *Basic Kronecker Identities*
*Whenever the expressions are defined,*

*(i)* $c(A \otimes B) = (cA) \otimes B = A \otimes (cB)$;
*(ii)* $(A + B) \otimes C = A \otimes C + B \otimes C$ *and* $A \otimes (B + C) = A \otimes B + A \otimes C$;
*(iii)* $(A \otimes B) \otimes C = A \otimes (B \otimes C)$;
*(iv)* $(A \otimes B)^t = A^t \otimes B^t$;
*(v)* $I_m \otimes I_n = I_{mn}$.

*If* $\mathbb{F} = \mathbb{C}$, *then also* $\overline{A \otimes B} = \overline{A} \otimes \overline{B}$ *and* $(A \otimes B)^* = A^* \otimes B^*$.

*Proof.*

(i) The $(i, j)$ block of $c(A \otimes B)$ is $c(a_{ij}B) = (ca_{ij})B = a_{ij}(cB)$, which is $(i, j)$ block of $(cA) \otimes B$ and $A \otimes (cB)$.
(ii) The $(i, j)$ block of $(A + B) \otimes C$ is $(a_{ij} + b_{ij})C = a_{ij}C + b_{ij}C$, so $(A + B) \otimes C = A \otimes C + B \otimes C$. The proof of $A \otimes (B + C) = A \otimes B + A \otimes C$ is identical inside each block.
(iii) If $B = [b_{rs}]$, then both $(A \otimes B) \otimes C$ and $A \otimes (B \otimes C)$ have blocks $a_{ij}b_{rs}C$ in the same relative positions.
(iv) The $(j, i)$ block of $(A \otimes B)^t$ is $(a_{ij}B)^t = a_{ij}B^t$, which is the $(j, i)$ block of $A^t \otimes B^t$.
(v) The block matrix $I_m \otimes I_n$ has $I_n$ on its diagonal blocks and zero elsewhere, hence equals $I_{mn}$.

If $\mathbb{F} = \mathbb{C}$, conjugation is checked blockwise, adjoint identity follows from conjugation together with part (iv).
*Structural Commentary*: Kronecker products replace scalar entries by whole scaled blocks. Standard algebraic rules survive because each block is still controlled by scalar distributivity and transposition.
*Key Insight*: Check identities block-by-block; each block reduces to an ordinary scalar-matrix identity. □

**Theorem 3.1.3.** *Mixed Product Property*
*Let* $A \in \mathrm{Mat}(m, n, \mathbb{F})$, $B \in \mathrm{Mat}(p, q, \mathbb{F})$, $C \in \mathrm{Mat}(n, r, \mathbb{F})$, $D \in \mathrm{Mat}(q, s, \mathbb{F})$. $(A \otimes B)(C \otimes D) = (AC) \otimes (BD)$.

*Proof.* Write $A = [a_{ik}]$ and $C = [c_{kj}]$. The $(i, j)$ block of $(A \otimes B)(C \otimes D)$ is
$\sum_{k=1}^{n} (a_{ik}B)(c_{kj}D) = \left(\sum_{k=1}^{n} a_{ik}c_{kj}\right) BD = (AC)_{ij}BD$, which is exactly the $(i, j)$ block of $(AC) \otimes (BD)$.
*Structural Commentary*: The outer indices multiply as $AC$, while the inner block action multiplies as $BD$.
*Key Insight*: Kronecker multiplication separates into ordinary multiplication in each factor. □

**Corollary 3.1.4.** *Inverse of a Kronecker Product*
*If* $A \in \mathrm{Mat}(m, m, \mathbb{F})$ *and* $B \in \mathrm{Mat}(n, n, \mathbb{F})$ *are invertible, then* $A \otimes B$ *is invertible and* $(A \otimes B)^{-1} = A^{-1} \otimes B^{-1}$.

*Proof.* By Theorem 3.1.3, $(A \otimes B)(A^{-1} \otimes B^{-1}) = I_m \otimes I_n = I_{mn}$ and the reverse product is identical.
*Structural Commentary*: Inverting a Kronecker product inverts each independent factor.
*Key Insight*: The mixed product property turns the inverse check into two ordinary inverse checks. □

**Proposition 3.1.5.** *Kronecker Product and Sum Spectra in the Diagonalizable Case*
*Let $A = P\Lambda P^{-1}$, $B = QMQ^{-1}$, where $\Lambda = \operatorname{diag}(\lambda_1, \dots, \lambda_n)$ and $M = \operatorname{diag}(\mu_1, \dots, \mu_q)$.*
*Then $A \otimes B \sim \Lambda \otimes M$; and $A \otimes I_q + I_n \otimes B \sim \Lambda \otimes I_q + I_n \otimes M$.*
*Their diagonal entries are respectively $\lambda_i \mu_j$ and $\lambda_i + \mu_j$ for $1 \le i \le n$, $1 \le j \le q$.*

*Proof.* By Theorem 3.1.3 and $(P \otimes Q)^{-1} = P^{-1} \otimes Q^{-1}$, $A \otimes B = (P \otimes Q)(\Lambda \otimes M)(P^{-1} \otimes Q^{-1})$.
Similarly, $A \otimes I_q + I_n \otimes B = (P \otimes Q)(\Lambda \otimes I_q + I_n \otimes M)(P^{-1} \otimes Q^{-1})$.
The middle matrices are diagonal with the stated $(i, j)$-indexed entries.
*Structural Commentary*: Eigenvectors tensor together, and Kronecker sums add the two scalar eigenvalues.
*Key Insight*: Diagonalize each factor first; the Kronecker operation becomes entrywise on eigenvalue pairs. □




Figure 85: Vertical plus horizontal second differences give the five-point stencil.

**Definition 3.1.6.** *Vectorisation*
Let $X = [\mathbf{x}_1 \ \mathbf{x}_2 \ \cdots \ \mathbf{x}_n] \in \operatorname{Mat}(m, n, \mathbb{F})$. The *vectorisation* of $X$ is

$$\operatorname{vec}(X) := \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_n \end{bmatrix} \in \mathbb{F}^{mn}.$$

Thus vec stacks the columns of $X$ vertically.

**Theorem 3.1.7.** *Vectorisation of a Triple Product*
*Let $A \in \operatorname{Mat}(m, n, \mathbb{F})$, $X \in \operatorname{Mat}(n, p, \mathbb{F})$, and $B \in \operatorname{Mat}(p, q, \mathbb{F})$. Then $\operatorname{vec}(AXB) = (B^t \otimes A)\operatorname{vec}(X)$.*

*Proof.* Write $B = [\mathbf{b}_1 \ \cdots \ \mathbf{b}_q]$ and $X = [\mathbf{x}_1 \ \cdots \ \mathbf{x}_p]$.
The $k$th column of $AXB$ is $AX\mathbf{b}_k = A \sum_{i=1}^{p} b_{ik}\mathbf{x}_i = \sum_{i=1}^{p} b_{ik} A\mathbf{x}_i = (\mathbf{b}_k^t \otimes A)\operatorname{vec}(X)$.
Stacking these $q$ columns gives $(B^t \otimes A)\operatorname{vec}(X)$.
*Structural Commentary*: Vectorisation rewrites two-sided multiplication as one matrix action.
*Key Insight*: Left multiplication contributes $A$; right multiplication contributes $B^t$. □

**Perspective 3.1.8.** *Vectorising a Minimal-Polynomial Test*
For $A \in \operatorname{Mat}(n, n, \mathbb{F})$ and a monic candidate $p_m(z) = a_0 + a_1 z + \cdots + a_{m-1} z^{m-1} + z^m$,
the matrix equation $a_0 I + a_1 A + \cdots + a_{m-1} A^{m-1} = -A^m$ can be written as the ordinary linear system

$$\begin{bmatrix} \operatorname{vec}(I) & \operatorname{vec}(A) & \cdots & \operatorname{vec}(A^{m-1}) \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_{m-1} \end{bmatrix} = -\operatorname{vec}(A^m)$$

*Structural Commentary*: Vectorisation turns matrix identities into linear systems.
*Key Insight*: Polynomial membership becomes a coefficient system in vectorized powers.

## 3.2 Hadamard Products

**Definition 3.2.1.** *Hadamard Product*
Let $A = [a_{ij}], B = [b_{ij}] \in \operatorname{Mat}(m, n, \mathbb{F})$.
The *Hadamard product*, or *entrywise product*, of $A$ and $B$ is $A \circ B := [a_{ij} b_{ij}] \in \operatorname{Mat}(m, n, \mathbb{F})$.

**Proposition 3.2.2.** *Basic Hadamard Identities*
*For matrices of compatible sizes:*

*(i)* $A \circ B = B \circ A$;
*(ii)* $A \circ (\gamma B + C) = \gamma(A \circ B) + A \circ C$;
*(iii)* $(A \circ B)^* = A^* \circ B^*$;
*(iv)* *if $E$ is the all-ones matrix, then $A \circ E = A$;*

*(v) if* $D = \operatorname{diag}(d_1, \ldots, d_n)$, *then* $A \circ D = \operatorname{diag}(a_{11}d_1, \ldots, a_{nn}d_n)$.

*Proof.*

(i) $(A \circ B)_{ij} = a_{ij}b_{ij} = b_{ij}a_{ij} = (B \circ A)_{ij}$.
(ii) $\big(A \circ (\gamma B + C)\big)_{ij} = a_{ij}(\gamma b_{ij} + c_{ij}) = \gamma a_{ij}b_{ij} + a_{ij}c_{ij}$.
(iii) $\big((A \circ B)^*\big)_{ij} = \overline{a_{ji}b_{ji}} = \overline{a_{ji}}\,\overline{b_{ji}} = (A^* \circ B^*)_{ij}$.
(iv) $(A \circ E)_{ij} = a_{ij}$.
(v) Since $D$ has zero off-diagonal entries, $A \circ D$ is diagonal with entries $a_{ii}d_i$.

*Structural Commentary*: Hadamard multiplication is entrywise, with no row-column summation.
*Key Insight*: Prove Hadamard identities entry by entry. □

**Theorem 3.2.3.** *Schur Product Theorem*
*If* $A, B \in \operatorname{Mat}(n, n, \mathbb{F})$ *are positive semidefinite, where* $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, *then* $A \circ B$ *is positive semidefinite.*

*Proof.* By Theorem 2.1.3, regard $A$ and $B$ as Gram matrices: $a_{ij} = \langle \mathbf{u}_j, \mathbf{u}_i \rangle$, $b_{ij} = \langle \mathbf{v}_j, \mathbf{v}_i \rangle$ for suitable vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$ and $\mathbf{v}_1, \ldots, \mathbf{v}_n$. Then $(A \circ B)_{ij} = \langle \mathbf{u}_j, \mathbf{u}_i \rangle \langle \mathbf{v}_j, \mathbf{v}_i \rangle = \langle \mathbf{u}_j \otimes \mathbf{v}_j, \mathbf{u}_i \otimes \mathbf{v}_i \rangle$.
Thus $A \circ B$ is the Gram matrix of the vectors $\mathbf{u}_j \otimes \mathbf{v}_j$, hence is positive semidefinite by Proposition 1.5.2.
*Structural Commentary*: The Schur product is the Gram matrix of paired tensor vectors.
*Key Insight*: Pair the $j$th vector from each Gram representation and take tensor products. □




Figure 86: For Gram matrices, entrywise products become tensor inner products, so $A \circ B$ is Gram.

**Corollary 3.2.4.** *Positive Definite Hadamard Products*
*Let* $A, B \in \operatorname{Mat}(n, n, \mathbb{F})$ *be positive semidefinite, where* $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$.

*(i) If* $A$ *is positive definite and every diagonal entry of* $B$ *is positive, then* $A \circ B$ *is positive definite.*
*(ii) If* $A$ *and* $B$ *are positive definite, then* $A \circ B$ *is positive definite.*

*Proof.*

(i) Let $\lambda_{\min} > 0$ be the smallest eigenvalue of $A$. Then $A - \lambda_{\min} I_n \succeq 0$, so $(A - \lambda_{\min} I_n) \circ B \succeq 0$ by Theorem 3.2.3. Thus, for $\mathbf{x} \neq \mathbf{0}$, $\mathbf{x}^*(A \circ B)\mathbf{x} = \mathbf{x}^*((A - \lambda_{\min} I_n) \circ B)\mathbf{x} \lambda_{\min} \sum_{j=1}^{n} b_{jj} |x_j|^2 > 0$.
(ii) If $B \succ 0$, then $b_{jj} = \mathbf{e}_j^* B \mathbf{e}_j > 0$ for every $j$, so (i) applies.

*Structural Commentary*: Subtract the spectral floor of $A$; the diagonal of $B$ gives strictness.
*Key Insight*: Use $I_n \circ B = \operatorname{diag}(b_{11}, \ldots, b_{nn})$. □

**Method 3.2.5.** *Schur Product Positivity Method*
To prove an entrywise product is positive semidefinite:

(i) factor or interpret each matrix as a Gram matrix;
(ii) pair the corresponding vectors;
(iii) take tensor products of the paired vectors;
(iv) identify the Hadamard product as the Gram matrix of those tensor products.

## 3.3 Spectral Radius and Matrix Powers

**Theorem 3.3.1.** *Ranks of Matrix Powers Stabilize*
*Let* $A \in \operatorname{Mat}(n, n, \mathbb{F})$. *Then:*

*(i)* $\operatorname{rank}(A^k) \geq \operatorname{rank}(A^{k+1})$ *for* $k = 1, 2, \ldots$;
*(ii) if* $\operatorname{rank}(A^k) = \operatorname{rank}(A^{k+1})$, *then* $\operatorname{col}(A^k) = \operatorname{col}(A^{k+p})$ *and* $\operatorname{rank}(A^k) = \operatorname{rank}(A^{k+p})$ *for every* $p = 1, 2, \ldots$;
*(iii) if* $A$ *is not invertible, then there is a least positive integer* $q \leq n$ *such that* $\operatorname{rank}(A^q) = \operatorname{rank}(A^{q+1})$.

*Proof.*

(i) Since $A^{k+1} = A^k A$, each column of $A^{k+1}$ is a linear combination of the columns of $A^k$. Hence $\operatorname{col}(A^{k+1}) \subseteq \operatorname{col}(A^k)$, so $\operatorname{rank}(A^{k+1}) \leq \operatorname{rank}(A^k)$.

(ii) The inclusion $\operatorname{col}(A^{k+1}) \subseteq \operatorname{col}(A^k)$ and equality of dimensions give $\operatorname{col}(A^{k+1}) = \operatorname{col}(A^k)$. Let $U = \operatorname{col}(A^k)$. Then multiplication by $A$ maps $U$ onto $U$, so induction gives $\operatorname{col}(A^{k+p}) = U$ for every $p \geq 1$.

(iii) If $A$ is not invertible, then Corollary 3.6.3 gives $\operatorname{rank}(A) \leq n-1$. If all inequalities $\operatorname{rank}(A^k) > \operatorname{rank}(A^{k+1})$ held for $k = 1, \ldots, n$, then $\operatorname{rank}(A^{n+1}) \leq \operatorname{rank}(A) - n < 0$, impossible. Hence equality occurs for some $q \leq n$; choose the least such $q$.

*Structural Commentary*: The column spaces of powers form a decreasing chain. In finite dimensions, a decreasing chain cannot keep losing dimension forever.
*Key Insight*: Once two consecutive ranks agree, the image has stabilized under multiplication by $A$. □

**Definition 3.3.2.** *Index of a Matrix*
If $A \in \operatorname{Mat}(n, n, \mathbb{F})$ is invertible, define its *index* to be 0. If $A$ is not invertible, its index is the least positive integer $q$ such that $\operatorname{rank}(A^q) = \operatorname{rank}(A^{q+1})$.

**Definition 3.3.3.** *Spectral Radius and Convergent Matrix*
For $A \in \operatorname{Mat}(n, n, \mathbb{C})$, the *spectral radius* is $\rho(A) := \max\{|\lambda| : \lambda \in \operatorname{spec}(A)\}$.
We say $A^p \to 0$ if every entry of $A^p$ converges to 0 as $p \to \infty$.

**Proposition 3.3.4.** *Basic Invariants of Nilpotents*
*For $N \in \operatorname{Mat}(n, n, \mathbb{F})$, the following are equivalent:*

*(i) $N$ is nilpotent;*
*(ii) the characteristic polynomial of $N$ is $t^n$;*
*(iii) $N^n = 0$.*

*In this case, $\operatorname{tr}(N) = 0$, $\det(N) = 0$ for $n \geq 1$, and the spectral radius of $N$ over $\mathbb{C}$ is 0.*

*Proof.* If $N^k = 0$, then every eigenvalue $\lambda$ of $N$ over an algebraic closure satisfies $\lambda^k = 0$, hence $\lambda = 0$. Thus the characteristic polynomial is $t^n$. If the characteristic polynomial is $t^n$, Cayley–Hamilton gives $N^n = 0$. The implication $N^n = 0 \Rightarrow$ nilpotent is immediate. The trace, determinant, and spectral-radius statements follow from the eigenvalue data and from Definitions 1.1.1 and 3.3.3.
*Structural Commentary*: Nilpotent matrices have no spectral mass away from zero.
*Key Insight*: Cayley–Hamilton turns the characteristic polynomial $t^n$ into the equation $N^n = 0$. □

**Theorem 3.3.5.** *Matrix Powers Converge to Zero*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{C})$. Then $A^p \to 0$ if and only if $\rho(A) < 1$.*

*Proof.* Choose a Jordan form $A = SJS^{-1}$.
Since $S$ and $S^{-1}$ are fixed, $A^p \to 0$ if and only if every Jordan block power in $J^p$ converges to 0.
($\Rightarrow$) Each block power tends to 0, so Proposition 6.1.5 gives $|\lambda| < 1$ for every eigenvalue. Thus $\rho(A) < 1$.
($\Leftarrow$) If $\rho(A) < 1$, then every eigenvalue satisfies $|\lambda| < 1$. Proposition 6.1.5 makes each block power tend to 0.
*Structural Commentary*: Similarity reduces matrix-power convergence to Jordan block-power convergence.
*Key Insight*: The largest eigenvalue modulus controls the asymptotic size of all blocks. □




Figure 87: The gray circle is $|\lambda| = 1$: inside it, powers decay; on or outside it, they do not.

**Definition 3.3.6.** *Power Bounded Matrix*
A matrix $A \in \operatorname{Mat}(n, n, \mathbb{C})$ is *power bounded* if for each entry position $(i, j)$, the sequence of entries of $A^p$ in that position is bounded for $p = 1, 2, \ldots$.

**Theorem 3.3.7.** *Power Bounded Jordan Criterion*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{C})$. The following are equivalent:*

*(i) $A$ is power bounded.*
*(ii) $\rho(A) \leq 1$, and every eigenvalue $\lambda$ with $|\lambda| = 1$ has only $1 \times 1$ Jordan blocks.*
*(iii) $\rho(A) \leq 1$, and $\dim E_\lambda(A) = m_A(\lambda)$ for every $\lambda \in \operatorname{spec}(A)$ with $|\lambda| = 1$.*

*Proof.* Work in Jordan form.

(*i*) $\Rightarrow$ (*ii*): If $|\lambda| > 1$, the diagonal entries $\lambda^p$ are unbounded.
If $|\lambda| = 1$ and $J_k(\lambda)$ has $k \geq 2$, then the first superdiagonal contains $p\lambda^{p-1}$, also unbounded.

(*ii*) $\Rightarrow$ (*i*): Blocks with $|\lambda| < 1$ converge to 0 by Proposition 6.1.5; $1 \times 1$ blocks with $|\lambda| = 1$ are bounded. Finite direct sums and fixed similarities preserve boundedness.

(*ii*) $\Leftrightarrow$ (*iii*): For fixed $\lambda$, $\dim E_\lambda(A)$ counts Jordan blocks, while $m_A(\lambda)$ counts their total size. Equality holds exactly when every block has size 1.

*Structural Commentary*: Unit-circle eigenvalues are harmless only when their nilpotent parts vanish.
*Key Insight*: Nontrivial Jordan blocks on the unit circle create polynomial growth. □

# 4 Matrix Decompositions

## 4.1 LU, QR, and Cholesky-Type Decompositions

Decompositions expose usable structure in a matrix. LU, QR, Cholesky, SVD, compact SVD, and polar decomposition separate elimination, geometry, positivity, rank, and stretching.

**Perspective 4.1.1.** *Matrix Factorisation Roadmap*
Factorisations expose structure by task: elimination gives $LU$ and $LDL^t$ (Theorem 4.1.5, Corollary 4.1.6); positivity gives Cholesky (Theorem 4.1.12); geometry gives $QR$ (Theorem 4.1.8); singular directions give SVD, compact SVD, and the pseudoinverse (Theorems 4.2.10, 4.2.14, Definition 5.2.4); polar decomposition separates motion from stretch (Theorem 4.2.19). Spectral factorisations appeared earlier: diagonalisation, spectral theorems, Jordan form, and Schur form (Theorems 5.4.3, 6.1.13, 6.1.14, 6.2.1, 6.1.10).
*Key Insight*: Choose the factorisation that exposes the needed structure.

**Definition 4.1.2.** *Full-Rank Factorization*
A factorization $A = XY$ with $X$ full column rank and $Y$ full row rank is called a *full-rank factorization* of $A$.

**Theorem 4.1.3.** *Full-Rank Factorization*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ have rank $r \geq 1$, and let the columns of $X \in \mathrm{Mat}(m, r, \mathbb{F})$ be a basis of* $\mathrm{col}(A)$. *Then there is a unique $Y \in \mathrm{Mat}(r, n, \mathbb{F})$ such that $A = XY$, and* $\mathrm{rank}(X) = \mathrm{rank}(Y) = r$.

*Proof.* Write $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n]$. Since the columns of $X$ form a basis of $\mathrm{col}(A)$, for each $j$ there is a unique $\mathbf{y}_j \in \mathbb{F}^r$ such that $\mathbf{a}_j = X\mathbf{y}_j$. Set $Y = [\mathbf{y}_1 \ \cdots \ \mathbf{y}_n]$. Then $A = [X\mathbf{y}_1 \ \cdots \ X\mathbf{y}_n] = XY$ and uniqueness of the coordinates $\mathbf{y}_j$ gives uniqueness of $Y$. The columns of $X$ are independent, so $\mathrm{rank}(X) = r$.
By Theorem 1.2.9, $r = \mathrm{rank}(A) = \mathrm{rank}(XY) \leq \mathrm{rank}(Y) \leq r$ because $Y$ has only $r$ rows. Hence $\mathrm{rank}(Y) = r$.
*Structural Commentary*: A matrix factors through a basis of its column space.
*Key Insight*: Full-rank factorization separates column directions from their coordinates. □

**Definition 4.1.4.** *LU Factorisation*
For $A \in \mathrm{Mat}(n, n, \mathbb{F})$, an *LU factorisation* is a product $A = LU$, where $L$ is lower triangular and $U$ is upper triangular. It is *unit lower triangular* if every diagonal entry of $L$ is 1.

**Theorem 4.1.5.** *LU Factorisation from Nonzero Leading Principal Minors*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$. If every leading principal minor of $A$ is nonzero, then $A = LU$ for a unique unit lower triangular $L$ and a unique upper triangular $U$.*

*Proof.* Induct on $n$. The case $n = 1$ is immediate. Write $A = \begin{bmatrix} a & \mathbf{r} \\ \mathbf{c} & B \end{bmatrix}$, $a \neq 0$.
Set $\boldsymbol{\ell} = a^{-1}\mathbf{c}$ and $S = B - a^{-1}\mathbf{c}\mathbf{r}$. For each leading $k \times k$ principal block of $A$, the block determinant formula gives determinant $a$ times the corresponding leading $(k-1) \times (k-1)$ principal minor of $S$. Hence all leading principal minors of $S$ are nonzero. By induction, $S = L_1U_1$ with $L_1$ unit lower triangular and $U_1$ upper triangular.
Then $A = \begin{bmatrix} 1 & 0 \\ \boldsymbol{\ell} & L_1 \end{bmatrix} \begin{bmatrix} a & \mathbf{r} \\ 0 & U_1 \end{bmatrix}$. For uniqueness, comparing the first row and first column of $LU$ determines $a$, $\mathbf{r}$, and $\boldsymbol{\ell}$, then reduces uniqueness to the lower-right Schur complement.
*Structural Commentary*: Gaussian elimination peels off one pivot and passes to a Schur complement.
*Key Insight*: Nonzero leading principal minors are exactly the no-row-swap pivot condition. □

**Corollary 4.1.6.** *Symmetric LU Gives an $LDL^t$ Factorisation*
*Let $A = A^t \in \mathrm{Mat}(n, n, \mathbb{R})$ have nonzero leading principal minors.*
*Then $A = LDL^t$, with $L$ unit lower triangular and $D$ diagonal invertible.*

*Proof.* By Theorem 4.1.5, write $A = LU$ uniquely with $L$ unit lower triangular and $U$ upper triangular. Since the leading principal minors are nonzero, each $u_{ii} \neq 0$. Put $D = \operatorname{diag}(u_{11}, \dots, u_{nn})$ and $U_0 = D^{-1}U$; then $U_0$ is unit upper triangular and $A = LDU_0$. Since $A = A^t$, also $A = U_0^t D L^t$. This is another LU factorisation with unit lower factor $U_0^t$, so uniqueness gives $L = U_0^t$. Hence $A = LDL^t$.
*Structural Commentary*: Symmetry makes the upper triangular unit factor the transpose of the lower one.
*Key Insight*: Once the pivots are separated into $D$, the two triangular halves mirror each other. □

**Definition 4.1.7.** *Wide and Narrow QR Factorisation*
For $A \in \operatorname{Mat}(m, n, \mathbb{F})$ with $m \geq n$, factorisation $A = V \begin{bmatrix} R \\ 0 \end{bmatrix}$, $V \in \operatorname{Mat}(m, m, \mathbb{F})$ unitary, $R \in \operatorname{Mat}(n, n, \mathbb{F})$ upper triangular is called a *wide QR factorisation*. Writing $V = [Q\ Q']$ gives the *narrow QR factorisation* $A = QR$.

**Theorem 4.1.8.** *QR Factorisation*
*Let $A \in \operatorname{Mat}(m, n, \mathbb{F})$ and suppose $m \geq n$.*

*(i) $\exists$ a unitary $V$ and an upper triangular $R$ with real nonnegative diagonal entries such that $A = V \begin{bmatrix} R \\ 0 \end{bmatrix}$. Writing $V = [Q\ Q']$ with $Q$ the first $n$ columns gives $A = QR$, with $Q$ orthonormal columns.*
*(ii) If $\operatorname{rank}(A) = n$, then narrow $A = QR$ is unique when $R$ has positive diagonal entries.*

*Proof.*

(i) At step $k$, after the first $k-1$ columns have been triangularized, apply Corollary 2.7.4 to the tail of the $k$th column. This gives a unitary $I_{k-1} \oplus U_k$ that sends that tail to $r_{kk}\mathbf{e}_1$ with $r_{kk} \geq 0$.
Since the first $k-1$ entries of every earlier column already contain all possible nonzero entries, acting only on rows $k, \dots, m$ preserves the earlier triangular columns.
After $n$ steps, $U_n \cdots U_1 A = \begin{bmatrix} R \\ 0 \end{bmatrix}$, where $R$ is upper triangular with nonnegative diagonal entries.
Let $V = (U_n \cdots U_1)^*$. Then $A = V \begin{bmatrix} R \\ 0 \end{bmatrix}$, and $V = [Q\ Q']$ gives $A = QR$ with $Q$ orthonormal columns.
If $\operatorname{rank}(A) = n$, then $n = \operatorname{rank}(A) = \operatorname{rank}(R)$, so the triangular matrix $R$ is invertible and its nonnegative diagonal entries are positive.

(ii) Suppose $A = Q_1R_1 = Q_2R_2$, where $Q_i^*Q_i = I_n$ and the $R_i$ are upper triangular with positive diagonal entries. Then $A^*A = R_1^*R_1 = R_2^*R_2$. Let $D = R_1R_2^{-1}$. Substituting $R_1 = DR_2$ gives $D^*D = I_n$.
Thus the columns of $D$ are orthonormal. Since $D$ is upper triangular, the first column has only one possible nonzero entry; orthogonality with it forces the rest of the first row to vanish.
Inducting on the lower-right principal submatrix shows $D$ is diagonal, and $D^*D = I_n$ gives diagonal entries of modulus 1. These diagonal entries are ratios of positive diagonal entries of $R_1$ and $R_2$, so they are positive. Hence $D = I_n$. Therefore $R_1 = R_2$, and then $Q_1 = Q_2$.

*Structural Commentary*: Householder steps triangularise columns; full rank makes the diagonal positive.
*Key Insight*: Positive diagonals remove the sign/phase ambiguity. □




Figure 88: QR writes $A = QR$ while preserving prefix column spaces: $\operatorname{col}(A_k) = \operatorname{col}(Q_k)$.

**Perspective 4.1.9.** *Gram–Schmidt and Householder QR*
Let $A = [\mathbf{a}_1\ \cdots\ \mathbf{a}_n]$ have full column rank.
Gram–Schmidt gives orthonormal columns $\mathbf{q}_1, \dots, \mathbf{q}_n$ s.t. $\operatorname{span}(\mathbf{a}_1, \dots, \mathbf{a}_k) = \operatorname{span}(\mathbf{q}_1, \dots, \mathbf{q}_k)$, $k = 1, \dots, n$.
Hence $A = QR$ with $Q^*Q = I$ and $R$ upper triangular. The Householder construction gives the same factorization via unitary transformations, which is the numerically stable form.

**Perspective 4.1.10.** *Rank-Deficient QR from Gram–Schmidt*
Let $A = [\mathbf{a}_1\ \cdots\ \mathbf{a}_n] \in \operatorname{Mat}(m, n, \mathbb{F})$, $m \geq n$, and let $\mathbf{u}_j$ be the arbitrary-list Gram–Schmidt remainders from Theorem 1.3.8. Set $\mathcal{J} = \{j : \mathbf{u}_j \neq \mathbf{0}\}$. For $j \in \mathcal{J}$ put $\mathbf{q}_j = \mathbf{u}_j / \|\mathbf{u}_j\|$; for $j \notin \mathcal{J}$ choose $\mathbf{q}_j \in \operatorname{col}(A)^\perp$ so that $Q = [\mathbf{q}_1\ \cdots\ \mathbf{q}_n]$ has orthonormal columns. With $R = Q^*A$, one has $A = QR$, $R$ upper triangular, and $r_{jj} = 0$ exactly when $\mathbf{u}_j = \mathbf{0}$. Thus uniqueness fails unless $\mathcal{J} = \{1, \dots, n\}$ and $r_{jj} > 0$ for all $j$.
*Structural Commentary*: Rank defect becomes zero QR pivots.
*Key Insight*: The full-rank hypothesis in uniqueness is a statement about nonzero orthogonal remainders.

**Corollary 4.1.11.** *Prefix Column Spaces in QR Factorisation*
*Let $A = QR$ be the narrow QR factorisation of $A \in \mathrm{Mat}(m, n, \mathbb{F})$ with $\mathrm{rank}(A) = n$ and positive diagonal entries in $R$. For $k = 1, \ldots, n$, let $A_k$ and $Q_k$ be the first $k$ columns of $A$ and $Q$, respectively. Then $\mathrm{col}(A_k) = \mathrm{col}(Q_k)$. Thus the columns of $Q_k$ form an orthonormal basis of $\mathrm{col}(A_k)$.*

*Proof.* Let $R_k$ be the leading $k \times k$ principal submatrix of $R$. Since $R$ is upper triangular with positive diagonal entries, $R_k$ is invertible. The first $k$ columns of $A = QR$ satisfy $A_k = Q_k R_k$, because upper triangularity prevents later columns of $Q$ from contributing. Hence $\mathrm{col}(A_k) \subseteq \mathrm{col}(Q_k)$ and $Q_k = A_k R_k^{-1}$ gives the reverse inclusion. The columns of $Q_k$ are orthonormal because $Q$ has orthonormal columns.
*Structural Commentary*: QR preserves prefix column spaces while orthonormalising them.
*Key Insight*: Invertible $R_k$ only changes coordinates within that space. □

**Theorem 4.1.12.** *Cholesky Factorisation*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$, where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. If $A \succeq 0$, then $A = LL^*$ for some lower triangular $L$ with nonnegative diagonal entries. If $A \succ 0$, then $L$ is unique and has positive diagonal entries.*

*Proof.* Let $B = A^{1/2}$ and take a QR factorisation $B = QR$ with $R$ upper triangular and diagonal entries nonnegative. Then $A = B^*B = R^*R$, so $L = R^*$ works.
If $A \succ 0$, then $L$ is invertible, so its diagonal entries are positive. For uniqueness, suppose $A = L_1 L_1^* = L_2 L_2^*$ with $L_i$ lower triangular and positive diagonal. Set $R_i = L_i^*$. Then $R_i$ is upper triangular with positive diagonal and $B^2 = A = R_i^* R_i$. Define $Q_i = B R_i^{-1}$. Then $Q_i^* Q_i = (R_i^{-1})^* B^2 R_i^{-1} = (R_i^{-1})^* R_i^* R_i R_i^{-1} = I_n$, so $B = Q_i R_i$ are QR factorisations with positive diagonal. Uniqueness of QR gives $R_1 = R_2$, hence $L_1 = L_2$.
*Structural Commentary*: Cholesky is QR applied to $A^{1/2}$.
*Key Insight*: Triangularise the square root, then set $L = R^*$. □




Figure 89: Apply QR to $B = A^{1/2}$; then $A = R^*R = LL^*$, with uniqueness in the positive definite case.

## 4.2 Polar and Singular Value Decompositions

**Theorem 4.2.1.** *Polar Decomposition*
*For every $T \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space, $\exists$ an isometry $S \in \mathcal{L}(\mathcal{V})$ s.t. $T = S\sqrt{T^*T}$.*

*Proof.* Let $P = \sqrt{T^*T}$. Since $T^*T$ is positive, $P$ is well defined by Proposition 6.3.7.
For every $\mathbf{v} \in \mathcal{V}$, $\|T\mathbf{v}\|^2 = \langle T^*T\mathbf{v}, \mathbf{v}\rangle = \langle P^2\mathbf{v}, \mathbf{v}\rangle = \|P\mathbf{v}\|^2$.
Define $S_1 : \mathrm{range}(P) \to \mathrm{range}(T)$ by $S_1(P\mathbf{v}) = T\mathbf{v}$. The norm identity shows this is well defined: if $P\mathbf{v}_1 = P\mathbf{v}_2$, then $\|T(\mathbf{v}_1 - \mathbf{v}_2)\| = \|P(\mathbf{v}_1 - \mathbf{v}_2)\| = 0$. It also shows $S_1$ is an isometry onto $\mathrm{range}(T)$.
The same norm identity gives $\ker(P) = \ker(T)$, so $\mathrm{rank}(P) = \mathrm{rank}(T)$ by rank-nullity. Hence $\mathrm{range}(P)^\perp$ and $\mathrm{range}(T)^\perp$ have the same dimension. Choose any isometry between these complements and combine it orthogonally with $S_1$ to obtain an isometry $S$ on all of $\mathcal{V}$.
Then $SP\mathbf{v} = S_1(P\mathbf{v}) = T\mathbf{v}$ for every $\mathbf{v}$, so $T = S\sqrt{T^*T}$.
*Structural Commentary*: The positive factor $\sqrt{T^*T}$ reproduces exactly the lengths produced by $T$.
*Key Insight*: Once the metric stretching has been extracted, the remaining map is forced to be an isometry. □

**Corollary 4.2.2.** *Uniqueness in Operator Polar Decomposition*
*Let $T \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space, and put $|T| := \sqrt{T^*T}$ and $|T^*| := \sqrt{TT^*}$.*

*(i) If $T = SP$, $S^*S = I_{\mathcal{V}}$, and $P \succeq 0$, then $P = |T|$. Hence the positive right polar factor is unique.*
*(ii) $\exists$ left polar decomposition $T = |T^*|\, R$ with $R$ an isometry, and $|T^*|$ is unique positive left polar factor.*
*(iii) If $T$ is invertible, then both isometric factors are unique: $S = T\,|T|^{-1}$ and $R = |T^*|^{-1}\,T$.*

*Proof.*

(i) If $T = SP$, then $T^*T = PS^*SP = P^2$. Since $P \succeq 0$, Proposition 6.3.7 gives $P = \sqrt{T^*T} = |T|$.
(ii) Apply Theorem 4.2.1 to $T^*$. There is an isometry $S_0$ such that $T^* = S_0\,|T^*|$.
Taking adjoints gives $T = |T^*|\, S_0^*$. Set $R = S_0^*$. If also $T = QR$ with $Q \succeq 0$ and $R$ an isometry, then $RR^* = I_{\mathcal{V}}$ in finite dimensions, so $TT^* = QRR^*Q = Q^2$. Hence $Q = |T^*|$.
(iii) If $T$ is invertible, then $|T|$ and $|T^*|$ are positive definite, hence invertible.
Therefore any right and left polar decompositions force $S = T\,|T|^{-1}$, $R = |T^*|^{-1}\,T$.

*Structural Commentary*: The positive polar factor is canonical.
*Key Insight*: Invertibility removes the kernel complement, so the whole factorisation becomes unique. □




Figure 90: Polar decomposition builds $S_1$ using $\|T\mathbf{v}\| = \left\|\sqrt{T^*T}\mathbf{v}\right\|$.

**Perspective 4.2.3.** *Polar Decomposition Separates Stretch and Direction*
In $T = S\sqrt{T^*T}$, the positive factor $\sqrt{T^*T}$ determines the metric stretch and the isometry $S$ determines the direction. Thus polar decomposition is the operator analogue of modulus times phase.
*Structural Commentary*: For normal $T$, the polar factors commute and share spectral data.
*Key Insight*: Polar decomposition factors an operator into positive geometry and isometric motion.

**Definition 4.2.4.** *Singular Values*
For finite-dimensional inner-product spaces $\mathcal{U}, \mathcal{V}$ and $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$, the *singular values* of $T$ are the eigenvalues of $\sqrt{T^*T} \in \mathcal{L}(\mathcal{U})$, where each eigenvalue $\lambda$ is repeated $\dim E_\lambda(\sqrt{T^*T})$.

**Theorem 4.2.5.** *Singular Value Decomposition*
*Let $T \in \mathcal{L}(\mathcal{V})$ have singular values $s_1, \dots, s_n$. Then there exist orthonormal bases $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ and $(\mathbf{f}_1, \dots, \mathbf{f}_n)$ of $\mathcal{V}$ such that $T\mathbf{v} = s_1 \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{f}_1 + \cdots + s_n \langle \mathbf{v}, \mathbf{e}_n \rangle \mathbf{f}_n$ for every $\mathbf{v} \in \mathcal{V}$.*

*Proof.* Let $|T| = \sqrt{T^*T}$. Since $|T|$ is positive, the real or complex spectral theorem gives an orthonormal basis $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ with $|T|\,\mathbf{e}_j = s_j \mathbf{e}_j$. Thus $|T|\,\mathbf{v} = \sum_{j=1}^{n} s_j \langle \mathbf{v}, \mathbf{e}_j \rangle \mathbf{e}_j$.
By Theorem 4.2.1, $T = S\,|T|$ for some isometry $S$. Put $\mathbf{f}_j = S\mathbf{e}_j$. The vectors $\mathbf{f}_1, \dots, \mathbf{f}_n$ form an orthonormal basis, and applying $S$ to the displayed expansion gives the desired formula.
*Structural Commentary*: SVD diagonalises $|T|$, then transports its eigenvectors by $S$.
*Key Insight*: The $s_j$ are stretches; the $\mathbf{f}_j$ are their landing directions. □




Figure 91: Since $T = S\sqrt{T^*T}$, each $\mathbf{e}_j$ maps to $s_j\mathbf{f}_j$.

**Proposition 4.2.6.** *Singular Values from $T^*T$*
*The singular values of $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ are the nonnegative square roots of the eigenvalues of $T^*T$, with the same eigenspace multiplicities.*

*Proof.* By the spectral theorem, choose an orthonormal basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ and nonnegative scalars $\lambda_1, \dots, \lambda_n$ such that $T^*T\mathbf{e}_j = \lambda_j \mathbf{e}_j$. The unique positive square root of $T^*T$ sends $\mathbf{e}_j$ to $\sqrt{\lambda_j}\mathbf{e}_j$.
Thus the singular values are $\sqrt{\lambda_j}$, with the same eigenspace multiplicities.
*Structural Commentary*: The positive square root keeps eigenspaces and sends $\lambda$ to $\sqrt{\lambda}$.
*Key Insight*: Singular values can be read from $T^*T$ directly. □

**Theorem 4.2.7.** *Singular Vectors for a Linear Map*
*Let $\mathcal{U}, \mathcal{V}$ be finite-dimensional inner-product spaces over $\mathbb{R}$ or $\mathbb{C}$, and let $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ have rank $r$.*
*Put $n = \dim \mathcal{U}$, $m = \dim \mathcal{V}$, and $\mathcal{U}_+ = \operatorname{range}(T^*)$, $\mathcal{U}_0 = \ker(T)$; $\mathcal{V}_+ = \operatorname{range}(T)$, $\mathcal{V}_0 = \ker(T^*)$.*
*Then $\exists$ ordered orthonormal bases $\beta = (\mathbf{u}_1, \dots, \mathbf{u}_r, \mathbf{u}_{r+1}, \dots, \mathbf{u}_n)$; $\gamma = (\mathbf{v}_1, \dots, \mathbf{v}_r, \mathbf{v}_{r+1}, \dots, \mathbf{v}_m)$*
*with $\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_r) = \mathcal{U}_+$; $\operatorname{span}(\mathbf{u}_{r+1}, \dots, \mathbf{u}_n) = \mathcal{U}_0$; $\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_r) = \mathcal{V}_+$; $\operatorname{span}(\mathbf{v}_{r+1}, \dots, \mathbf{v}_m) = \mathcal{V}_0$,*
*and numbers $s_1, \dots, s_r > 0$ such that $T\mathbf{u}_k = s_k\mathbf{v}_k$; $T^*\mathbf{v}_k = s_k\mathbf{u}_k$, for $1 \le k \le r$.*
*The $s_k$ are the positive singular values; $\mathbf{u}_k$ and $\mathbf{v}_k$ are right and left singular vectors.*

*Proof.* Let $A = T^*T$. Then $A \succeq 0$, $A = A^*$, and $\ker(A) = \ker(T)$. By spectral theorem, choose rthonormal eigenbasis $\mathbf{u}_1, \dots, \mathbf{u}_n$ s.t. $A\mathbf{u}_k = s_k^2\mathbf{u}_k$ for $1 \le k \le r$; $A\mathbf{u}_k = 0$ for $k > r$; $s_k > 0$.
Thus $\operatorname{span}(\mathbf{u}_{r+1}, \dots, \mathbf{u}_n) = \mathcal{U}_0$, and Proposition 2.4.11 gives $\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_r) = \mathcal{U}_+$. For $1 \le k \le r$,
set $\mathbf{v}_k = s_k^{-1}T\mathbf{u}_k$. Then $\langle \mathbf{v}_i, \mathbf{v}_j \rangle = \frac{1}{s_i s_j} \langle T\mathbf{u}_i, T\mathbf{u}_j \rangle = \frac{1}{s_i s_j} \langle T^*T\mathbf{u}_i, \mathbf{u}_j \rangle = \delta_{ij}$; $T^*\mathbf{v}_k = s_k^{-1}A\mathbf{u}_k = s_k\mathbf{u}_k$.
Hence $\mathbf{v}_1, \dots, \mathbf{v}_r$ are orthonormal and span $\mathcal{V}_+ = T(\mathcal{U}_+)$. Complete them by an orthonormal basis of $\mathcal{V}_0 = \mathcal{V}_+^\perp$.
*Structural Commentary*: $T^*T$ gives right directions; $T$ gives left directions.
*Key Insight*: Away from the two kernels, $T$ and $T^*$ act symmetrically by the same positive scalars. □




Figure 92: The singular-vector bases split the domain and codomain into active singular subspaces and kernels.

**Perspective 4.2.8.** *Operator-to-Matrix SVD Bridge*
In the bases from Theorem 4.2.7, the matrix of $T$ is rectangular diagonal with diagonal entries $s_1, \dots, s_r$ and zeros elsewhere. Returning to standard orthonormal bases gives the matrix form $A = V\Sigma W^*$ in Theorem 4.2.10.

**Definition 4.2.9.** *Matrix Singular Values*
Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$, let $q = \min(m, n)$, and let $r = \mathrm{rank}(A)$. The *positive singular values* of $A$ are the positive eigenvalues of $|A| = (A^*A)^{1/2}$, written $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > 0$.
The full list of singular values is $\sigma_1, \dots, \sigma_r, \sigma_{r+1}, \dots, \sigma_q$, where $\sigma_{r+1} = \cdots = \sigma_q = 0$.

**Theorem 4.2.10.** *Matrix Singular Value Decomposition*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ have singular values $\sigma_1 \geq \cdots \geq \sigma_q \geq 0$, where $q = \min(m, n)$. Then $\exists$ unitary matrices $V \in \mathrm{Mat}(m, m, \mathbb{F})$, $W \in \mathrm{Mat}(n, n, \mathbb{F})$ such that $A = V\Sigma W^*$, $\Sigma = \begin{bmatrix} \mathrm{diag}(\sigma_1, \dots, \sigma_q) & 0 \\ 0 & 0 \end{bmatrix} \in \mathrm{Mat}(m, n, \mathbb{R})$ with the evident deletion of empty blocks.*

*Proof.* If $A = 0$, take $\Sigma = 0$ and choose any unitary $V, W$. Assume $r = \mathrm{rank}(A) \geq 1$.
Diagonalise $A^*A$ as $W^*A^*AW = \mathrm{diag}(\sigma_1^2, \dots, \sigma_r^2, 0, \dots, 0)$, where $\mathbf{w}_1, \dots, \mathbf{w}_n$ are the columns of $W$.
For $1 \leq j \leq r$, set $\mathbf{v}_j = \sigma_j^{-1} A\mathbf{w}_j$. Then $\mathbf{v}_i^*\mathbf{v}_j = \frac{1}{\sigma_i\sigma_j}\mathbf{w}_i^*A^*A\mathbf{w}_j = \delta_{ij}$ so $\mathbf{v}_1, \dots, \mathbf{v}_r$ are orthonormal.
Complete them to an orthonormal basis of $\mathbb{F}^m$ and let $V$ be the matrix with these columns.
For $j > r$, $\|A\mathbf{w}_j\|_2^2 = \mathbf{w}_j^*A^*A\mathbf{w}_j = 0$, hence $A\mathbf{w}_j = 0$. Thus $AW = V\Sigma$, equivalently $A = V\Sigma W^*$.
*Structural Commentary*: $A^*A$ determines the right singular directions and singular values.
*Key Insight*: Apply $A$ and normalize to get the left singular directions. □




Figure 93: SVD: right-unitary motion, diagonal stretch, left-unitary motion.

**Proposition 4.2.11.** *Basic Singular Value Identities*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ have singular values $\sigma_1 \geq \cdots \geq \sigma_q \geq 0$, where $q = \min(m, n)$ and $r = \mathrm{rank}(A)$. Let $c \in \mathbb{F}$.*

*(i) The positive eigenvalues of $A^*A$ and $AA^*$ are $\sigma_1^2, \dots, \sigma_r^2$.*
*(ii) $\sum_{j=1}^q \sigma_j^2 = \mathrm{tr}(A^*A) = \mathrm{tr}(AA^*) = \|A\|_F^2$, where $\|A\|_F^2 := \mathrm{tr}(A^*A)$.*
*(iii) $A$ and $A^*$ have the same singular values. Over $\mathbb{C}$, so do $A^t$ and $\overline{A}$.*
*(iv) The singular values of $cA$ are $|c|\,\sigma_1, \dots, |c|\,\sigma_q$.*

*Proof.* Let $A = V\Sigma W^*$ be an SVD. Then $A^*A = W(\Sigma^*\Sigma)W^*$ and $AA^* = V(\Sigma\Sigma^*)V^*$.

(i) The nonzero diagonal entries of $\Sigma^*\Sigma$ and $\Sigma\Sigma^*$ are $\sigma_1^2, \dots, \sigma_r^2$.
(ii) Since $A^*A$ and $\Sigma^*\Sigma$ are unitarily similar, and $AA^*$ and $\Sigma\Sigma^*$ are unitarily similar,
$\mathrm{tr}(A^*A) = \mathrm{tr}(\Sigma^*\Sigma) = \sum_{j=1}^{q} \sigma_j^2 = \mathrm{tr}(\Sigma\Sigma^*) = \mathrm{tr}(AA^*)$. The final equality is the definition of $\|A\|_F$.
(iii) For $A^*$, the two Gram matrices are interchanged. Over $\mathbb{C}$, $(A^t)^*A^t = (AA^*)^t$ and $(\overline{A})^*\overline{A} = (A^*A)^t$.
Since a matrix and its transpose have the same characteristic polynomial, the spectra agree.

(iv) Since $(cA)^*(cA) = |c|^2 A^*A$, singular values scale by $|c|$.

*Structural Commentary*: Unitary factors do not change Gram spectra; $\Sigma$ contains all singular data.
*Key Insight*: Reduce each claim to $\Sigma^*\Sigma$ and $\Sigma\Sigma^*$. □

**Corollary 4.2.12.** *Square Invertibility and Singular Values*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$, $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, have singular values $\sigma_1 \geq \cdots \geq \sigma_n \geq 0$. The following are equivalent:*

*(i) $A$ is invertible;*
*(ii) $\mathrm{rank}(A) = n$;*
*(iii) $\sigma_n > 0$;*
*(iv) $A^*A \succ 0$.*

*If $r = \mathrm{rank}(A) < n$, then $\sigma_1, \ldots, \sigma_r > 0$ and $\sigma_{r+1} = \cdots = \sigma_n = 0$.*

*Proof.*

(i) $\Rightarrow$ (ii) By Corollary 3.6.3, invertibility gives full rank.
(ii) $\Rightarrow$ (iii) By Definition 4.2.9, the number of positive singular values is $\mathrm{rank}(A)$.
(iii) $\Rightarrow$ (iv) If $\sigma_n > 0$, then $A$ has full column rank, so Proposition 2.1.2 gives $A^*A \succ 0$.
(iv) $\Rightarrow$ (i) If $A\mathbf{x} = \mathbf{0}$, then $\mathbf{x}^*A^*A\mathbf{x} = 0$; PSD gives $\mathbf{x} = \mathbf{0}$. Thus $\ker(A) = \{\mathbf{0}\}$, so $A$ is invertible.

The rank-defect statement follows again from Definition 4.2.9.
*Structural Commentary*: SVD sees nonsingularity as the absence of zero stretch directions.
*Key Insight*: A zero singular value is exactly a nonzero vector sent to zero. □

**Definition 4.2.13.** *Compact Singular Value Decomposition*
Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ have rank $r \geq 1$.
A factorisation $A = X\Sigma_r Y^*$ is a *compact singular value decomposition* if $X \in \mathrm{Mat}(m, r, \mathbb{F})$ and $Y \in \mathrm{Mat}(n, r, \mathbb{F})$ have orthonormal columns and $\Sigma_r = \mathrm{diag}(\sigma_1, \ldots, \sigma_r)$ with $\sigma_1 \geq \cdots \geq \sigma_r > 0$.

**Theorem 4.2.14.** *Compact Singular Value Decomposition*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ have rank $r \geq 1$.*

*(i) If $A = V\Sigma W^*$ is an SVD and $X, Y$ are the first $r$ columns of $V, W$, then $A = X\Sigma_r Y^*$.*
*(ii) Conversely, if $X, Y$ have orthonormal columns and $\Sigma_r = \mathrm{diag}(\sigma_1, \ldots, \sigma_r)$ with $\sigma_1 \geq \cdots \geq \sigma_r > 0$, then $X\Sigma_r Y^*$ has positive singular values $\sigma_1, \ldots, \sigma_r$.*

*Proof.*

(i) Partition $V = [X\ X_0]$, $W = [Y\ Y_0]$, and $\Sigma = \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix}$. Multiplication gives $A = V\Sigma W^* = X\Sigma_r Y^*$.

(ii) Let $B = X\Sigma_r Y^*$. Since $X^*X = Y^*Y = I_r$, $B^*B = Y\Sigma_r^2 Y^*$.
Completing $Y$ to a unitary matrix $\widetilde{Y} = [Y\ Y_0]$ gives $\widetilde{Y}^*B^*B\widetilde{Y} = \Sigma_r^2 \oplus 0$.
Thus the positive eigenvalues of $B^*B$ are $\sigma_1^2, \ldots, \sigma_r^2$, so the positive singular values of $B$ are $\sigma_1, \ldots, \sigma_r$.

*Structural Commentary*: Compact SVD keeps only the nonzero singular directions.
*Key Insight*: The rank-$r$ data $X, \Sigma_r, Y$ reconstructs $A$. □

**Theorem 4.2.15.** *Column and Kernel Spaces from Compact SVD*
*Let $A = X\Sigma_r Y^*$ be a compact SVD. Then $\mathrm{col}(A) = \mathrm{col}(X) = (\ker A^*)^\perp$, and $\ker(A) = \mathrm{col}(Y)^\perp = (\mathrm{col}(A^*))^\perp$.*

*Proof.* Since $A = X\Sigma_r Y^*$ and $AY\Sigma_r^{-1} = X$, $\mathrm{col}(A) = \mathrm{col}(X)$.
For $\mathbf{z} \in \mathbb{F}^n$, $A\mathbf{z} = 0 \iff X\Sigma_r Y^*\mathbf{z} = 0 \iff Y^*\mathbf{z} = 0 \iff \mathbf{z} \perp \mathrm{col}(Y)$, because $X$ has independent columns and $\Sigma_r$ is invertible. The orthogonal-complement identities follow from Proposition 2.4.11.
*Structural Commentary*: Compact SVD exposes image directions $X$ and null directions $\mathrm{col}(Y)^\perp$.
*Key Insight*: $A$ acts only on the right singular subspace $\mathrm{col}(Y)$. □




Figure 94: Compact SVD: $A$ maps $\mathrm{col}(Y)$ onto $\mathrm{col}(X)$ and kills $\ker(A) = \mathrm{col}(Y)^\perp$.

**Theorem 4.2.16.** *Uniqueness of Compact SVD up to Singular Subspaces*
*Let $A = X_1\Sigma_r Y_1^* = X_2\Sigma_r Y_2^*$ be compact SVDs with the same $\Sigma_r$. Then $X_1 = X_2U$, $Y_1 = Y_2U$ for some unitary $U$ satisfying $U\Sigma_r = \Sigma_r U$. Hence $U = U_1 \oplus \cdots \oplus U_d$ along the equal singular-value blocks of $\Sigma_r$.*

*Proof.* Theorem 4.2.15 gives $\operatorname{col}(X_1) = \operatorname{col}(X_2)$ and $\operatorname{col}(Y_1) = \operatorname{col}(Y_2)$. Since each $X_i$ and $Y_i$ has orthonormal columns, there are unitary $U, V \in \operatorname{Mat}(r, r, \mathbb{F})$ such that $X_1 = X_2U$ and $Y_1 = Y_2V$. Substitution and multiplication on the left by $X_2^*$ and on the right by $Y_2$ give $\Sigma_r = U\Sigma_r V^*$, hence $U\Sigma_r = \Sigma_r V$.
Therefore $U\Sigma_r^2 U^* = \Sigma_r^2$, so $U$ commutes with $\Sigma_r^2$. Lemma 5.4.10 makes $U$ block diagonal on equal singular-value blocks. Hence $U\Sigma_r = \Sigma_r U$, and $U\Sigma_r = \Sigma_r V$ forces $V = U$ because $\Sigma_r$ is invertible.
*Structural Commentary*: Compact SVD is unique up to unitary changes inside equal singular-value blocks.
*Key Insight*: Distinct singular values fix subspaces; repeated ones only fix the subspace, not its basis. □

**Definition 4.2.17.** *Matrix Modulus*
For $A \in \operatorname{Mat}(m, n, \mathbb{F})$, the *modulus* of $A$ is $|A| := (A^*A)^{1/2} \in \operatorname{Mat}(n, n, \mathbb{F})$.
Thus $|A^*| = (AA^*)^{1/2} \in \operatorname{Mat}(m, m, \mathbb{F})$.

**Proposition 4.2.18.** *Modulus from SVD*
*Let $A = V\Sigma W^*$ be an SVD. Then $|A| = W(\Sigma^*\Sigma)^{1/2}W^*$ and $|A^*| = V(\Sigma\Sigma^*)^{1/2}V^*$.*

*Proof.* Since $A^*A = W(\Sigma^*\Sigma)W^*$ and $AA^* = V(\Sigma\Sigma^*)V^*$, the displayed matrices are positive semidefinite square roots of $A^*A$ and $AA^*$, respectively. Uniqueness of the positive semidefinite square root gives the claim.
*Structural Commentary*: The modulus keeps only the singular-value stretch.
*Key Insight*: In an SVD, $|A|$ is the right singular-vector version of the diagonal stretch. □

**Theorem 4.2.19.** *Rectangular Polar Decomposition*
*Let $A = V\Sigma W^* \in \operatorname{Mat}(m, n, \mathbb{F})$ be an SVD.*

*(i) If $m \geq n$, let $V_n$ be the first $n$ columns of $V$ and set $U = V_nW^*$. Then $U^*U = I_n$ and $A = U|A|$.*
*(ii) If $m \leq n$, let $W_m$ be the first $m$ columns of $W$ and set $U = VW_m^*$. Then $UU^* = I_m$ and $A = |A^*|U$.*
*(iii) If $m = n$, then $U = VW^*$ is unitary and $A = U|A| = |A^*|U$.*

*Proof.*

(i) Proposition 4.2.18 gives $|A| = W(\Sigma^*\Sigma)^{1/2}W^*$.
Since $V_n(\Sigma^*\Sigma)^{1/2} = V\Sigma$, $U|A| = V_nW^*W(\Sigma^*\Sigma)^{1/2}W^* = V\Sigma W^* = A$. Also $U^*U = WV_n^*V_nW^* = I_n$.
(ii) Apply (i) to $A^*$ with the SVD $A^* = W\Sigma^*V^*$, obtaining $A^* = U_0|A^*|$ with $U_0 = W_mV^*$ and $U_0^*U_0 = I_m$.
Taking adjoints gives $A = |A^*|U_0^* = |A^*|VW_m^*$, and $U_0^*(U_0^*)^* = U_0^*U_0 = I_m$. Thus $U = VW_m^*$ works.
(iii) This is the common case of (i) and (ii), with $U = VW^*$ unitary.

*Structural Commentary*: Polar decomposition compresses the two SVD unitary factors into one partial isometry.
*Key Insight*: SVD records every singular direction; polar form records stretch plus motion. □

**Method 4.2.20.** *SVD and Polar Computation Method*
To compute an SVD and the associated polar factors of $A \in \operatorname{Mat}(m, n, \mathbb{F})$:

(i) Diagonalise $A^*A$; the square roots of its positive eigenvalues are the positive singular values.
(ii) Choose orthonormal right singular vectors $\mathbf{w}_j$ for the positive eigenvalues and set $\mathbf{v}_j = \sigma_j^{-1}A\mathbf{w}_j$.
(iii) Complete the $\mathbf{v}_j$ and $\mathbf{w}_j$ to unitary matrices $V, W$ to obtain $A = V\Sigma W^*$.
Omit zero-singular directions for compact SVD.
(iv) Use $|A| = W(\Sigma^*\Sigma)^{1/2}W^*$ and merge $V, W$ as in Theorem 4.2.19 to obtain the polar factor.

**Theorem 4.2.21.** *Uniqueness of the Positive Polar Factor*
*Let $A \in \operatorname{Mat}(m, n, \mathbb{F})$.*

*(i) If $m \geq n$ and $A = UB$ with $U^*U = I_n$ and $B \succeq 0$, then $B = |A|$.*
*If $\operatorname{rank}(A) = n$, then $U = A|A|^{-1}$ is unique.*
*(ii) If $m \leq n$ and $A = CU$ with $UU^* = I_m$ and $C \succeq 0$, then $C = |A^*|$.*
*If $\operatorname{rank}(A) = m$, then $U = |A^*|^{-1}A$ is unique.*

*Proof.*

(i) $A^*A = BU^*UB = B^2$, $B = |A|$ by uniqueness of positive square root. If $\operatorname{rank}(A) = n$, then $\ker(A) = \{\mathbf{0}\}$.
Since $\mathbf{x}^*A^*A\mathbf{x} = \|A\mathbf{x}\|_2^2$, this gives $\ker(A^*A) = \{\mathbf{0}\}$ and $|A|$ is invertible. Thus $U = A|A|^{-1}$.
(ii) $AA^* = CUU^*C = C^2$, so $C = |A^*|$. If $\operatorname{rank}(A) = m$, then $A^*$ has full column rank, so $\ker(A^*) = \{\mathbf{0}\}$.
Since $\mathbf{y}^*AA^*\mathbf{y} = \|A^*\mathbf{y}\|_2^2$, this gives $\ker(AA^*) = \{\mathbf{0}\}$ and $|A^*|$ is invertible. Thus $U = |A^*|^{-1}A$.

*Structural Commentary*: The positive factor is fixed by the Gram matrix.
*Key Insight*: Full rank makes the partial isometry unique. □

**Theorem 4.2.22.** *Normality and Polar Commutation*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ and suppose $A = U|A|$ with $U$ unitary. Then $A$ is normal if and only if $U|A| = |A|U$.*

*Proof.*
($\Rightarrow$) If $A$ is normal, then $|A|^2 = A^*A = AA^* = U|A|^2U^*$. Thus $U|A|^2U^* = |A|^2$. The matrix $U|A|U^*$ is PSD and squares to $|A|^2$, so uniqueness of positive square root gives $U|A|U^* = |A|$. Hence $U|A| = |A|U$.
($\Leftarrow$) If $U|A| = |A|U$, then $AA^* = U|A|^2U^* = |A|^2 = A^*A$.

*Structural Commentary*: Normality says that the polar motion preserves the stretching eigenspaces.
*Key Insight*: Normality is commutation of polar motion and stretch. □

**Proposition 4.2.23.** *Absolute Determinant Comes from the Positive Polar Factor*
*For $T \in \mathcal{L}(\mathcal{V})$ on a finite-dimensional inner-product space, $|\det T| = \det\sqrt{T^*T}$.*

*Proof.* By Theorem 4.2.1, $T = S\sqrt{T^*T}$ for some isometry $S$. Hence $|\det T| = |\det S|\left|\det\sqrt{T^*T}\right| = \det\sqrt{T^*T}$, where Proposition 1.1.22 gives $|\det S| = 1$, and positivity of $\sqrt{T^*T}$ gives nonnegative determinant.
*Structural Commentary*: Polar decomposition isolates stretch; only the positive factor changes volume.
*Key Insight*: Determinant sign or phase belongs to the isometric part; volume belongs to the positive part. □

# 5 Norms, Conditioning, Perturbation, and Stability

## 5.1 Matrix Norms

Matrix norms measure stretch, energy, approximation quality, and sensitivity. Singular values organize norms, low-rank approximation, pseudoinverses, conditioning, and error bounds.

**Definition 5.1.1.** *Spectral Norm and Extreme Singular Values*
For $A \in \mathrm{Mat}(m, n, \mathbb{F})$, define the *spectral norm* $\|A\|_2 := \max_{\|\mathbf{x}\|_2=1} \|A\mathbf{x}\|_2$.
If the singular values of $A$ are $\sigma_1 \geq \cdots \geq \sigma_q \geq 0$, $q = \min(m, n)$, set $\sigma_{\max}(A) := \sigma_1$.
When $A$ is square, $\sigma_{\min}(A) := \sigma_n$.

**Theorem 5.1.2.** *Spectral Norm is the Largest Singular Value*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$ have singular values $\sigma_1 \geq \cdots \geq \sigma_q \geq 0$. Then $\|A\|_2 = \sigma_{\max}(A)$.*
*Moreover, if $A = V\Sigma W^*$ is an SVD and $\mathbf{w}_1$ is the first column of $W$, then $\|A\mathbf{w}_1\|_2 = \sigma_1$.*

*Proof.* Let $\mathbf{w}_1, \ldots, \mathbf{w}_n$ be the columns of $W$. For $\mathbf{x} = \sum_{j=1}^n c_j\mathbf{w}_j$ with $\|\mathbf{x}\|_2 = 1$, unitary invariance of the Euclidean norm gives $\|A\mathbf{x}\|_2^2 = \|\Sigma W^*\mathbf{x}\|_2^2 = \sum_{j=1}^q \sigma_j^2|c_j|^2 \leq \sigma_1^2\sum_{j=1}^n |c_j|^2 = \sigma_1^2$.
Thus $\|A\|_2 \leq \sigma_1$. Taking $\mathbf{x} = \mathbf{w}_1$ gives equality.
*Structural Commentary*: The right singular basis diagonalises the stretching computation.
*Key Insight*: The largest stretch occurs in a top right singular direction. □

**Proposition 5.1.3.** *Basic Properties of the Spectral Norm*
*For compatible matrices $A, B$ and scalar $c$:*
*(i)* $\|A\|_2 \geq 0$, *and* $\|A\|_2 = 0$ *if and only if* $A = 0$;
*(ii)* $\|cA\|_2 = |c|\,\|A\|_2$;
*(iii)* $\|A + B\|_2 \leq \|A\|_2 + \|B\|_2$;
*(iv)* $\|AB\|_2 \leq \|A\|_2\,\|B\|_2$.

*Proof.*
(i) Since $\|A\mathbf{x}\|_2 \geq 0$, $\|A\|_2 \geq 0$. Also $\|A\|_2 = 0$ iff $A\mathbf{x} = 0$ for every unit $\mathbf{x}$, iff $A = 0$.
(ii) $\|cA\|_2 = \max_{\|\mathbf{x}\|_2=1}\|cA\mathbf{x}\|_2 = |c|\,\|A\|_2$.
(iii) For $\|\mathbf{x}\|_2 = 1$, $\|(A + B)\mathbf{x}\|_2 \leq \|A\mathbf{x}\|_2 + \|B\mathbf{x}\|_2 \leq \|A\|_2 + \|B\|_2$. Maximize over unit $\mathbf{x}$.
(iv) For $\|\mathbf{x}\|_2 = 1$, $\|AB\mathbf{x}\|_2 \leq \|A\|_2\,\|B\mathbf{x}\|_2 \leq \|A\|_2\,\|B\|_2$. Maximize over unit $\mathbf{x}$.

*Structural Commentary*: The spectral norm is the operator norm induced by Euclidean geometry.
*Key Insight*: Bound each output vector first, then take the worst case. □

**Definition 5.1.4.** *Operator Norm on $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$*
For $A \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$, define $\|A\|_{\mathrm{op}} := \sup_{\|\mathbf{h}\|_2\leq 1}\|A\mathbf{h}\|_2$.
Equivalently, after standard bases identify $A$ with its matrix, $\|A\|_{\mathrm{op}}$ is the spectral norm from Definition 5.1.1.
Thus $\|A\mathbf{h}\|_2 \leq \|A\|_{\mathrm{op}}\,\|\mathbf{h}\|_2$, and sums and products obey the norm estimates in Proposition 5.1.3.

The induced metric on $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ is $d(A, B) := \|A - B\|_{\text{op}}$; this is the metric used for continuity of derivative maps in Definition 3.2.2.

**Proposition 5.1.5.** *Frobenius and Spectral Norm Comparison*
*Let $A \in \text{Mat}(m, n, \mathbb{F})$ have singular values $\sigma_1 \geq \cdots \geq \sigma_q \geq 0$. Then $\|A\|_F^2 = \sum_{j=1}^q \sigma_j^2$ and $\|A\|_2 \leq \|A\|_F$. If $A \neq 0$, equality $\|A\|_2 = \|A\|_F$ holds if and only if $\text{rank}(A) = 1$.*

*Proof.* The Frobenius identity is Proposition 4.2.11. Hence $\|A\|_2^2 = \sigma_1^2 \leq \sum_{j=1}^q \sigma_j^2 = \|A\|_F^2$. Assume $A \neq 0$.
($\Rightarrow$) If $\|A\|_2 = \|A\|_F$, then $\sum_{j=2}^q \sigma_j^2 = 0$, so only $\sigma_1$ is nonzero and $\text{rank}(A) = 1$.
($\Leftarrow$) If $\text{rank}(A) = 1$, then $\sigma_2 = \cdots = \sigma_q = 0$, so $\|A\|_2 = \|A\|_F$.
*Structural Commentary*: Frobenius counts all singular energy; spectral keeps the largest part.
*Key Insight*: Equality means exactly one nonzero singular value. □

**Theorem 5.1.6.** *Submultiplicativity of Frobenius and Spectral Norms*
*Let $A \in \text{Mat}(m, n, \mathbb{F})$, $B \in \text{Mat}(n, k, \mathbb{F})$. Then $\|AB\|_F \leq \|A\|_2 \|B\|_F \leq \|A\|_F \|B\|_F$ and $\|AB\|_2 \leq \|A\|_2 \|B\|_2$.*

*Proof.* Write $B = [\mathbf{b}_1 \ \cdots \ \mathbf{b}_k]$. Then $\|AB\|_F^2 = \sum_{j=1}^k \|A\mathbf{b}_j\|_2^2 \leq \|A\|_2^2 \sum_{j=1}^k \|\mathbf{b}_j\|_2^2 = \|A\|_2^2 \|B\|_F^2$.
Proposition 5.1.5 gives $\|A\|_2 \leq \|A\|_F$. The spectral-norm inequality is Proposition 5.1.3.
*Structural Commentary*: Frobenius norm is column energy; each column of $AB$ is $A$ applied to a column of $B$.
*Key Insight*: Control every column by the spectral norm of $A$. □

**Proposition 5.1.7.** *Unitary Invariance of Singular Values*
*Let $A \in \text{Mat}(m, n, \mathbb{F})$, and let $U \in \text{Mat}(m, m, \mathbb{F})$ and $V \in \text{Mat}(n, n, \mathbb{F})$ be unitary. Then $A$ and $UAV$ have the same singular values. In particular, $\|UAV\|_2 = \|A\|_2$ and $\|UAV\|_F = \|A\|_F$.*

*Proof.* Since $(UAV)^*(UAV) = V^*A^*AV$, the Gram matrices of $A$ and $UAV$ are unitarily similar, hence have the same eigenvalues. Singular values are the nonnegative square roots of these eigenvalues. The two norm identities follow from Theorem 5.1.2 and Proposition 5.1.5.
*Structural Commentary*: Left and right unitary factors change coordinates, not singular-value data.
*Key Insight*: Singular values are spectral data of $A^*A$, and unitary conjugation preserves that spectrum. □

**Example 5.1.8.** *Similarity Need Not Preserve Singular Values*
Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$. Then $B = DAD^{-1}$ with $D = \text{diag}(2, 1)$, so $A$ and $B$ are similar.
However $A^*A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$, $B^*B = \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix}$ have different eigenvalues, so $A$ and $B$ have different singular values.
Singular values are preserved by unitary left-right equivalence, not by arbitrary similarity.

**Theorem 5.1.9.** *Schur's Inequality and Normality*
*Let $A \in \text{Mat}(n, n, \mathbb{C})$ have eigenvalues $\lambda_1, \ldots, \lambda_n$, counted algebraically.*
*Then $\sum_{j=1}^n |\lambda_j|^2 \leq \|A\|_F^2$ with equality if and only if $A$ is normal.*

*Proof.* Let $A = UTU^*$ be a Schur form, where $T = [t_{ij}]$ is upper triangular and $t_{jj} = \lambda_j$.
Then $\|A\|_F^2 = \text{tr}(T^*T) = \sum_{j=1}^n |\lambda_j|^2 + \sum_{i<j} |t_{ij}|^2$. Inequality holds. Equality holds exactly when $T$ is diagonal.
($\Rightarrow$) If equality holds, then $T$ is diagonal, so $A$ is unitarily diagonalizable and hence normal.
($\Leftarrow$) If $A$ is normal, then $T$ is normal and upper triangular, hence diagonal by Lemma 6.1.12.
*Structural Commentary*: Schur form splits Frobenius norm into eigenvalue mass plus off-diagonal mass.
*Key Insight*: Normality means the Schur off-diagonal part vanishes. □

**Definition 5.1.10.** *Defect from Normality*
Let $A \in \text{Mat}(n, n, \mathbb{C})$ have eigenvalues $\lambda_1, \ldots, \lambda_n$, counted with algebraic multiplicity.
The *defect from normality* of $A$ is $\Delta(A) := \|A\|_F^2 - \sum_{j=1}^n |\lambda_j|^2$.
By Theorem 5.1.9, $\Delta(A) \geq 0$, with equality if and only if $A$ is normal.

**Theorem 5.1.11.** *Normality Transfers from AB to BA*
*Let $A, B \in \text{Mat}(n, n, \mathbb{C})$. If $A$, $B$, and $AB$ are normal, then $BA$ is normal.*

*Proof.* By Theorem 1.1.16, $AB$ and $BA$ have the same eigenvalues counted with algebraic multiplicity. Also,

$$\begin{aligned}\operatorname{tr}\big((BA)^*(BA)\big) &= \operatorname{tr}(A^*B^*BA) = \operatorname{tr}(AA^*B^*B)\\ &= \operatorname{tr}(A^*AB^*B) = \operatorname{tr}(A^*ABB^*) = \operatorname{tr}(B^*A^*AB) = \operatorname{tr}\big((AB)^*(AB)\big),\end{aligned}$$

using cyclicity of trace and normality of $A$ and $B$. Thus $\|BA\|_F^2 = \|AB\|_F^2$, so $\Delta(BA) = \Delta(AB)$. Since $AB$ is normal, $\Delta(AB) = 0$; hence $\Delta(BA) = 0$, and Theorem 5.1.9 gives that $BA$ is normal.
*Structural Commentary*: Same spectrum and Frobenius norm give the same Schur defect.
*Key Insight*: Schur defect turns normality of $AB$ into normality of $BA$. □

**Theorem 5.1.12.** *Eigenvalues Versus Singular Values*
*Let $A \in \operatorname{Mat}(n,n,\mathbb{C})$ have eigenvalues $\lambda_1,\dots,\lambda_n$ ordered so that $|\lambda_1| \geq \cdots \geq |\lambda_n|$, and singular values $\sigma_1 \geq \cdots \geq \sigma_n \geq 0$. Then:*

*(i)* $|\det A| = \prod_{j=1}^n |\lambda_j| = \prod_{j=1}^n \sigma_j$;
*(ii)* $\rho(A) \leq \|A\|_2$;
*(iii)* $\sigma_n \leq |\lambda_n|$;
*(iv)* *$A$ is normal if and only if $|\lambda_j| = \sigma_j$ for every $j$;*
*(v)* *$A$ is positive semidefinite if and only if $\lambda_j = \sigma_j$ for every $j$.*

*Proof.*

(i) The eigenvalue product gives $|\det A| = \prod_j |\lambda_j|$.
If $A = V\Sigma W^*$ is an SVD, then $|\det V| = |\det W| = 1$, so $|\det A| = |\det \Sigma| = \prod_j \sigma_j$.
(ii) If $A\mathbf{x} = \lambda\mathbf{x}$ with $\|\mathbf{x}\|_2 = 1$, then $|\lambda| = \|A\mathbf{x}\|_2 \leq \|A\|_2$. Maximize over eigenvalues.
(iii) If $A$ is singular, both sides are 0 or the inequality is immediate.
If $A$ is invertible, apply (ii) to $A^{-1}$: $\frac{1}{|\lambda_n|} = \rho(A^{-1}) \leq \|A^{-1}\|_2 = \frac{1}{\sigma_n}$.
(iv) If $A$ is normal, the spectral theorem gives $\sigma_j = |\lambda_j|$. Conversely, equality gives $\sum_j |\lambda_j|^2 = \sum_j \sigma_j^2 = \|A\|_F^2$, so Theorem 5.1.9 gives normality.
(v) If $A \succeq 0$, then $A$ is normal and $\lambda_j \geq 0$, so (iv) gives $\lambda_j = \sigma_j$. Conversely, equality gives normality with nonnegative real eigenvalues, hence $A \succeq 0$ by Theorem 2.1.3.

*Structural Commentary*: Normality aligns eigenvalue moduli with singular values; positivity removes the moduli.
*Key Insight*: Normality is the bridge that turns spectral data into singular-value data. □

**Theorem 5.1.13.** *Trace Bound by Singular Values*
*If $A \in \operatorname{Mat}(n,n,\mathbb{C})$, then $|\operatorname{tr} A| \leq \sum_{j=1}^n \sigma_j(A)$. Consequently, $\sum_{j=1}^n |\lambda_j(A)| \leq \sum_{j=1}^n \sigma_j(A)$ and $\sum_{j=1}^n |a_{jj}| \leq \sum_{j=1}^n \sigma_j(A)$.*

*Proof.* Let $A = V\Sigma W^*$ be an SVD.
Writing $\mathbf{v}_k, \mathbf{w}_k$ for the columns of $V, W$, $\operatorname{tr} A = \sum_{k=1}^n \sigma_k \mathbf{w}_k^*\mathbf{v}_k$, $|\mathbf{w}_k^*\mathbf{v}_k| \leq 1$, so $|\operatorname{tr} A| \leq \sum_k \sigma_k$.
For eigenvalues, take a Schur form $T$ of $A$ and a diagonal unitary $D$ with $\operatorname{tr}(DT) = \sum_j |\lambda_j|$.
For diagonal entries, take diagonal unitary $D$ with $\operatorname{tr}(DA) = \sum_j |a_{jj}|$.
In both cases, unitary invariance of singular values and the trace bound give the result.
*Structural Commentary*: Phase rotations turn trace estimates into eigenvalue and diagonal estimates.
*Key Insight*: Prove $|\operatorname{tr} A| \leq \sum_j \sigma_j$ once, then rotate phases. □

## 5.2 Low-Rank Approximation and Pseudoinverses

**Proposition 5.2.1.** *Rank-One SVD Terms and Truncation Error*
*Let $A = X\Sigma_r Y^* = \sum_{j=1}^r \sigma_j \mathbf{x}_j \mathbf{y}_j^*$ be a compact SVD, with $\sigma_1 \geq \cdots \geq \sigma_r > 0$.*
*For $0 \leq k \leq r$, set $A_k := \sum_{j=1}^k \sigma_j \mathbf{x}_j \mathbf{y}_j^*$, $A_0 := 0$.*
*Then the rank-one summands are mutually orthogonal in the Frobenius inner product, and*

$$\|A\|_F^2 = \sum_{j=1}^r \sigma_j^2, \qquad \|A - A_k\|_F^2 = \sum_{j=k+1}^r \sigma_j^2, \qquad \|A - A_k\|_2 = \begin{cases}\sigma_{k+1}, & k < r,\\ 0, & k = r.\end{cases}$$

*Proof.* For $i,j \in \{1,\dots,r\}$, $\langle \sigma_i \mathbf{x}_i\mathbf{y}_i^*, \sigma_j\mathbf{x}_j\mathbf{y}_j^*\rangle_F = \sigma_i\sigma_j \operatorname{tr}(\mathbf{y}_i\mathbf{x}_i^*\mathbf{x}_j\mathbf{y}_j^*) = \sigma_i\sigma_j\delta_{ij}$.
Pythagoras in the Frobenius inner product gives the two Frobenius formulas. The residual $A - A_k$ has compact SVD $\sum_{j=k+1}^r \sigma_j \mathbf{x}_j\mathbf{y}_j^*$, so Theorem 5.1.2 gives the spectral-norm formula.

*Structural Commentary*: Compact SVD decomposes a matrix into orthogonal rank-one energy packets.
*Key Insight*: Truncation error is exactly the energy left in the discarded singular directions. □




Figure 95: Blue bars form $A_k$; red bars are discarded, with $\|A - A_k\|_F^2 = \sum_{j>k} \sigma_j^2$.

**Theorem 5.2.2.** *Eckart–Young Low-Rank Approximation*
*Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$ have singular values $\sigma_1 \geq \cdots \geq \sigma_q \geq 0$, where $q = \min(m,n)$ and $\sigma_j = 0$ for $j > q$. If $A_k$ is the rank-$k$ truncated SVD, $0 \leq k < q$, and $B \in \mathrm{Mat}(m,n,\mathbb{F})$ has $\mathrm{rank}(B) \leq k$, then*

$$\|A - A_k\|_2 \leq \|A - B\|_2\,, \qquad \|A - A_k\|_F \leq \|A - B\|_F\,.$$

*Proof.* Let $\mathbf{w}_1, \ldots, \mathbf{w}_n$ be right singular vectors for $A$.

i. Since $\mathrm{rank}(B) \leq k$, rank-nullity gives $\dim \ker(B) \geq n - k$. The subspaces $\ker(B)$ and $\mathrm{span}(\mathbf{w}_1, \ldots, \mathbf{w}_{k+1})$ in $\mathbb{F}^n$ have dimensions summing at least $n+1$, so their intersection contains a nonzero vector.
Choose unit $\mathbf{x}$ in this intersection. Let $\mathbf{x} = \sum_{i=1}^{k+1} c_i \mathbf{w}_i$ gives $\sum_i |c_i|^2 = 1$ and $\|A\mathbf{x}\|_2^2 = \sum_{i=1}^{k+1} \sigma_i^2 |c_i|^2 \geq \sigma_{k+1}^2$.
Thus $\|A - B\|_2 \geq \|(A-B)\mathbf{x}\|_2 = \|A\mathbf{x}\|_2 \geq \sigma_{k+1} = \|A - A_k\|_2$.

ii. Choose an $(n-k)$-dimensional subspace $\mathcal{U} \subseteq \ker(B)$, with projection $P$ and orthonormal basis $\mathbf{z}_1, \ldots, \mathbf{z}_{n-k}$.
Since $B\mathbf{z}_j = 0$, $\|A - B\|_F^2 \geq \sum_{j=1}^{n-k} \|(A-B)\mathbf{z}_j\|_2^2 = \sum_{j=1}^{n-k} \|A\mathbf{z}_j\|_2^2 = \sum_{i=1}^{n} \sigma_i^2 \|P\mathbf{w}_i\|_2^2$.
Put $\alpha_i = \|P\mathbf{w}_i\|_2^2$. Then $0 \leq \alpha_i \leq 1$ and $\sum_i \alpha_i = \mathrm{tr}(P) = n - k$.
Among all such weights, the sum $\sum_i \sigma_i^2 \alpha_i$ is minimized by placing weight 1 on the $n-k$ smallest values of the decreasing sequence $\sigma_i^2$, hence $\sum_i \sigma_i^2 \alpha_i \geq \sum_{i=k+1}^{n} \sigma_i^2 = \|A - A_k\|_F^2$.

*Structural Commentary*: Rank-$k$ competitors vanish on at least $n-k$ directions.
*Key Insight*: The optimal loss is precisely the singular-value tail. □

**Method 5.2.3.** *Truncated SVD Approximation Method*
To compress or approximate $A$ by rank $k$:

(i) compute or estimate the leading singular triples $(\sigma_j, \mathbf{x}_j, \mathbf{y}_j)$;
(ii) keep $A_k = \sum_{j=1}^{k} \sigma_j \mathbf{x}_j \mathbf{y}_j^*$;
(iii) use $\|A - A_k\|_F^2 = \sum_{j>k} \sigma_j^2$ and $\|A - A_k\|_2 = \sigma_{k+1}$ to quantify the lost information.

**Definition 5.2.4.** *Moore–Penrose Pseudoinverse*
Let $A \in \mathrm{Mat}(m,n,\mathbb{F})$. If $A = 0$, define $A^\dagger = 0_{n,m}$.
If $\mathrm{rank}(A) = r \geq 1$ and $A = X\Sigma_r Y^*$ is a compact SVD, define $A^\dagger := Y\Sigma_r^{-1}X^* \in \mathrm{Mat}(n,m,\mathbb{F})$.
This is the *pseudoinverse* or *Moore–Penrose pseudoinverse* of $A$.

**Proposition 5.2.5.** *Pseudoinverse is Well Defined*
*The definition of $A^\dagger$ does not depend on the chosen compact SVD of $A$.*

*Proof.* Let $A = X_1\Sigma_r Y_1^* = X_2\Sigma_r Y_2^*$. By Theorem 4.2.16, $X_1 = X_2 U$, $Y_1 = Y_2 U$, and $U\Sigma_r^{-1}U^* = \Sigma_r^{-1}$.
Hence $Y_1\Sigma_r^{-1}X_1^* = Y_2 U\Sigma_r^{-1}U^*X_2^* = Y_2\Sigma_r^{-1}X_2^*$.
*Structural Commentary*: Compact SVD ambiguity cancels in $Y\Sigma_r^{-1}X^*$.
*Key Insight*: Inverting the singular values preserves the same block symmetries. □

**Corollary 5.2.6.** *Pseudoinverse from a Full SVD*
*Let $A = V\Sigma W^*$ be a full SVD of $A \in \mathrm{Mat}(m,n,\mathbb{F})$, with rank $r$ and positive singular values $\sigma_1, \ldots, \sigma_r$.*
*Write $V = [X\ X_0]$, $W = [Y\ Y_0]$, and $\Sigma_r = \mathrm{diag}(\sigma_1, \ldots, \sigma_r)$.*
*In conformal blocks, $\Sigma = \begin{bmatrix} \Sigma_r & 0 \\ 0 & 0 \end{bmatrix}$, $\Sigma^\dagger := \begin{bmatrix} \Sigma_r^{-1} & 0 \\ 0 & 0 \end{bmatrix} \in \mathrm{Mat}(n,m,\mathbb{F})$. Then $A^\dagger = W\Sigma^\dagger V^*$.*

*Proof.* If $r = 0$, then $A = 0$ and $W\Sigma^\dagger V^* = 0$.
If $r > 0$, the block form gives $A = X\Sigma_r Y^*$; $A^\dagger = Y\Sigma_r^{-1}X^*$; and $W\Sigma^\dagger V^* = Y\Sigma_r^{-1}X^*$. Thus $A^\dagger = W\Sigma^\dagger V^*$.
*Structural Commentary*: Full SVD pads compact SVD by zero blocks.
*Key Insight*: The pseudoinverse flips $V$ and $W$ and inverts only the positive singular values. □

**Theorem 5.2.7.** *Basic Properties of the Pseudoinverse*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$. Then:*

*(i)* $\mathrm{col}(A^\dagger) = \mathrm{col}(A^*)$ *and* $\ker(A^\dagger) = \ker(A^*)$;
*(ii)* $AA^\dagger$ *is the orthogonal projection of* $\mathbb{F}^m$ *onto* $\mathrm{col}(A)$;
*(iii)* $A^\dagger A$ *is the orthogonal projection of* $\mathbb{F}^n$ *onto* $\mathrm{col}(A^*)$;
*(iv)* $AA^\dagger A = A$ *and* $A^\dagger AA^\dagger = A^\dagger$;
*(v)* $(A^\dagger)^\dagger = A$;
*(vi) if* $m = n$ *and* $A$ *is invertible, then* $A^\dagger = A^{-1}$.

*Proof.* The zero case is immediate. Let $A = X\Sigma_r Y^*$ be a compact SVD; then $A^\dagger = Y\Sigma_r^{-1}X^*$.

(i) $\mathrm{col}(A^\dagger) = \mathrm{col}(Y) = \mathrm{col}(A^*)$ and $\ker(A^\dagger) = \ker(X^*) = \ker(A^*)$.
(ii) $AA^\dagger = XX^*$, the orthogonal projection onto $\mathrm{col}(X) = \mathrm{col}(A)$.
(iii) $A^\dagger A = YY^*$, the orthogonal projection onto $\mathrm{col}(Y) = \mathrm{col}(A^*)$.
(iv) These follow from $XX^*X = X$ and $YY^*Y = Y$.
(v) The compact SVD of $A^\dagger$ is $Y\Sigma_r^{-1}X^*$, so $(A^\dagger)^\dagger = A$.
(vi) If $A$ is square and invertible, then $AA^\dagger = A^\dagger A = I_n$, hence $A^\dagger = A^{-1}$.

*Structural Commentary*: The pseudoinverse inverts nonzero singular directions and projects away null directions.
*Key Insight*: Replace each $\sigma_j > 0$ by $\sigma_j^{-1}$ on the same singular subspaces. □




Figure 96: $A$ and $A^\dagger$ invert the nonzero singular subspaces; they vanish on $\ker(A)$ and $\ker(A^*)$, respectively.

**Theorem 5.2.8.** *Penrose Characterization of the Pseudoinverse*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$, $\mathcal{C} = \mathrm{col}(A)$, and $\mathcal{D} = \mathrm{col}(A^*) = (\ker A)^\perp$. Then $A^\dagger$ is unique $B \in \mathrm{Mat}(n, m, \mathbb{F})$ satisfying*

*(i)* $ABA = A$;
*(ii)* $BAB = B$;
*(iii)* $AB$ *is Hermitian;*
*(iv)* $BA$ *is Hermitian.*

*Proof.*

(i) Theorem 5.2.7 gives $AA^\dagger A = A$; $A^\dagger AA^\dagger = A^\dagger$, and $AA^\dagger$, $A^\dagger A$ are orthogonal projections. Hence $A^\dagger$ satisfies the four equations.
(ii) Let $B$ satisfy the four equations. Then $(AB)^2 = ABAB = AB$; $(AB)^* = AB$. Also $\mathrm{range}(AB) = \mathcal{C}$, since $\mathrm{range}(AB) \subseteq \mathcal{C}$ and $AB(A\mathbf{x}) = ABA\mathbf{x} = A\mathbf{x}$. By Theorem 2.5.6, $AB = P_{\mathcal{C}}$.
(iii) $(BA)^2 = BABA = BA$; $(BA)^* = BA$. Also $\ker(BA) = \ker(A)$, as $BA\mathbf{x} = 0 \Rightarrow A\mathbf{x} = ABA\mathbf{x} = 0$ and the reverse inclusion is immediate. By Theorem 2.5.6, $BA = P_{\mathcal{D}}$.
(iv) For $\mathbf{y} \in \mathbb{F}^m$, write $\mathbf{y} = \mathbf{p} + \mathbf{z}$ with $\mathbf{p} \in \mathcal{C}$ and $\mathbf{z} \in \mathcal{C}^\perp = \ker(A^*)$. Since $AB\mathbf{z} = 0$, $B\mathbf{z} = BAB\mathbf{z} = 0$. If $\mathbf{p} = A\mathbf{x}$, then $B\mathbf{p} = BA\mathbf{x} = P_{\mathcal{D}}\mathbf{x}$, independent of the choice of $\mathbf{x}$ because $\ker(P_{\mathcal{D}}) = \mathcal{D}^\perp = \ker(A)$. Thus the four equations determine $B$ uniquely. Since $A^\dagger$ satisfies them, $B = A^\dagger$.

*Structural Commentary*: Penrose equations force the two projection factors.
*Key Insight*: Once $AB$ and $BA$ are projections, $B$ is forced. □

**Theorem 5.2.9.** *Pseudoinverse and Minimum-Norm Least Squares*
*Let $A \in \mathrm{Mat}(m, n, \mathbb{F})$, let $\mathbf{y} \in \mathbb{F}^m$, and set $\mathbf{x}_0 = A^\dagger\mathbf{y}$.*

*(i) If $A\mathbf{x} = \mathbf{y}$ is consistent, then $A\mathbf{x}_0 = \mathbf{y}$ and $\mathbf{x}_0$ is the unique solution of smallest Euclidean norm.*
*(ii) For arbitrary $\mathbf{y}$, $\mathbf{x}_0$ is a least-squares solution of $A\mathbf{x} \approx \mathbf{y}$, and it is the unique least-squares solution of smallest Euclidean norm.*

*Proof.* Let $P = AA^\dagger = P_{\mathrm{col}(A)}$ and note that $\mathbf{x}_0 \in \mathrm{col}(A^\dagger) = \mathrm{col}(A^*) = (\ker A)^\perp$.

(i) If $\mathbf{y} \in \mathrm{col}(A)$, then $A\mathbf{x}_0 = P\mathbf{y} = \mathbf{y}$. For any solution $\mathbf{x}$, $\mathbf{x} - \mathbf{x}_0 \in \ker(A)$, so $\|\mathbf{x}\|_2^2 = \|\mathbf{x}_0 + (\mathbf{x} - \mathbf{x}_0)\|_2^2 = \|\mathbf{x}_0\|_2^2 + \|\mathbf{x} - \mathbf{x}_0\|_2^2$ with equality only when $\mathbf{x} = \mathbf{x}_0$.

(ii) For arbitrary $\mathbf{y}$, $A\mathbf{x}_0 = P\mathbf{y}$, so $\mathbf{y} - A\mathbf{x}_0 \perp \operatorname{col}(A)$ and $\mathbf{x}_0$ is least-squares optimal by Theorem 2.3.4. Any least-squares solution $\mathbf{x}$ satisfies $A\mathbf{x} = P\mathbf{y} = A\mathbf{x}_0$, hence the same orthogonal decomposition gives $\|\mathbf{x}\|_2 \geq \|\mathbf{x}_0\|_2$, with equality only for $\mathbf{x} = \mathbf{x}_0$.

*Structural Commentary*: Least squares projects onto $\operatorname{col}(A)$, then removes the $\ker(A)$ component.
*Key Insight*: $A^\dagger\mathbf{y}$ is the fitted solution orthogonal to $\ker(A)$. □

**Method 5.2.10.** *Pseudoinverse Solution Method*
For a system $A\mathbf{x} \approx \mathbf{y}$:

(i) compute a compact SVD $A = X\Sigma_r Y^*$;
(ii) form $A^\dagger = Y\Sigma_r^{-1}X^*$;
(iii) set $\mathbf{x}_0 = A^\dagger\mathbf{y}$;
(iv) interpret $A\mathbf{x}_0 = P_{\operatorname{col}(A)}\mathbf{y}$ and $\mathbf{x}_0 \in (\ker A)^\perp$.

## 5.3 Conditioning

**Definition 5.3.1.** *Spectral Condition Number*
If $A \in \operatorname{Mat}(n, n, \mathbb{F})$ is invertible, its *spectral condition number* is $\kappa_2(A) := \|A\|_2 \left\|A^{-1}\right\|_2$.
Equivalently, $\kappa_2(A) = \frac{\sigma_{\max}(A)}{\sigma_{\min}(A)}$. If $A$ is singular, $\kappa_2(A)$ is undefined.

**Proposition 5.3.2.** *Properties of the Spectral Condition Number*
*Let $A, B \in \operatorname{Mat}(n, n, \mathbb{F})$ be invertible.*

*(i)* $\kappa_2(AB) \leq \kappa_2(A)\kappa_2(B)$;
*(ii)* $\kappa_2(A) \geq 1$, *with equality if and only if $A$ is a nonzero scalar multiple of a unitary matrix;*
*(iii)* $\kappa_2(cA) = \kappa_2(A)$ *for every nonzero scalar $c$;*
*(iv)* $\kappa_2(A) = \kappa_2(A^*) = \kappa_2(\overline{A}) = \kappa_2(A^t) = \kappa_2(A^{-1})$ *over $\mathbb{C}$, with the evident real special cases;*
*(v)* $\kappa_2(A^*A) = \kappa_2(A)^2$;
*(vi) if $U, V$ are unitary, then* $\kappa_2(UAV) = \kappa_2(A)$.

*Proof.*

(i) Use submultiplicativity: $\kappa_2(AB) = \|AB\|_2 \left\|B^{-1}A^{-1}\right\|_2 \leq \|A\|_2 \|B\|_2 \left\|B^{-1}\right\|_2 \left\|A^{-1}\right\|_2$.
(ii) Since $I_n = AA^{-1}$, $1 = \|I_n\|_2 \leq \kappa_2(A)$. Equality means $\sigma_{\max}(A) = \sigma_{\min}(A)$, equivalently every singular value is the same positive number $s$. If $A = V\Sigma W^*$, then $\Sigma = sI_n$, so $A = sVW^*$ is a nonzero scalar multiple of a unitary matrix. The converse is immediate.
(iii) This follows from $\|cA\|_2 = |c| \|A\|_2$ and $(cA)^{-1} = c^{-1}A^{-1}$.
(iv) These matrices have the same singular values as $A$, except $A^{-1}$ has singular values $\sigma_n^{-1} \geq \cdots \geq \sigma_1^{-1}$.
(v) The singular values of $A^*A$ are $\sigma_1(A)^2, \ldots, \sigma_n(A)^2$.
(vi) This follows from Proposition 5.1.7.

*Structural Commentary*: Condition number compares largest and smallest singular stretches.
*Key Insight*: A well-conditioned matrix stretches every direction by nearly the same factor. □

**Corollary 5.3.3.** *Positive Definite Condition Number by Eigenvalues*
*Let $A \succ 0$ have eigenvalues $0 < \lambda_1 \leq \cdots \leq \lambda_n$. Then $\|A\|_2 = \lambda_n$; $\left\|A^{-1}\right\|_2 = \frac{1}{\lambda_1}$; and $\kappa_2(A) = \frac{\lambda_n}{\lambda_1}$.*

*Proof.* By Theorems 2.1.7 and 5.1.12, the singular values of $A$ are $\lambda_1, \ldots, \lambda_n$.
Hence Theorem 5.1.2 gives $\|A\|_2 = \lambda_n$. Since $\operatorname{spec}(A^{-1}) = \{\lambda_1^{-1}, \ldots, \lambda_n^{-1}\}$, same argument gives $\left\|A^{-1}\right\|_2 = \lambda_1^{-1}$, and multiplying yields $\kappa_2(A) = \lambda_n/\lambda_1$.
*Structural Commentary*: For $A \succ 0$, singular and eigenvalue stretches coincide.
*Key Insight*: Conditioning compares the largest and smallest positive eigendirections. □

**Perspective 5.3.4.** *Determinant is Not Conditioning*
Determinants measure volume, not sensitivity.
$\kappa_2(cA) = \kappa_2(A)$ for $c \neq 0$ (Proposition 5.3.2), but $\det(cA) = c^n \det(A)$.

**Theorem 5.3.5.** *Condition Number and Linear-System Error*
*Let $A \in \operatorname{Mat}(n, n, \mathbb{F})$ be invertible, let $\mathbf{y} \neq \mathbf{0}$, and let $A\mathbf{x} = \mathbf{y}$.*
*If $A(\Delta\mathbf{x}) = \Delta\mathbf{y}$, then* $\frac{\|\Delta\mathbf{x}\|_2}{\|\mathbf{x}\|_2} \leq \kappa_2(A)\frac{\|\Delta\mathbf{y}\|_2}{\|\mathbf{y}\|_2}$

*Proof.* From $\mathbf{y} = A\mathbf{x}$, we have $\|\mathbf{y}\|_2 \leq \|A\|_2 \|\mathbf{x}\|_2$, hence $1/\|\mathbf{x}\|_2 \leq \|A\|_2 / \|\mathbf{y}\|_2$. Also $\Delta\mathbf{x} = A^{-1}\Delta\mathbf{y}$, so $\|\Delta\mathbf{x}\|_2 \leq \left\|A^{-1}\right\|_2 \|\Delta\mathbf{y}\|_2$. Multiplying the two estimates gives the displayed bound.
*Structural Commentary*: Solution error is controlled by the ratio of extreme singular stretches.
*Key Insight*: $\kappa_2(A)$ measures worst-case error magnification. □

**Theorem 5.3.6.** *Condition Number and Matrix Perturbation*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{F})$ be invertible. Suppose $A\mathbf{x} = \mathbf{y}$; $(A+\Delta A)(\mathbf{x}+\Delta\mathbf{x}) = \mathbf{y}$, and $\mathbf{x}+\Delta\mathbf{x} \neq \mathbf{0}$. Then $\frac{\|\Delta\mathbf{x}\|_2}{\|\mathbf{x}+\Delta\mathbf{x}\|_2} \leq \kappa_2(A)\frac{\|\Delta A\|_2}{\|A\|_2}$.*

*Proof.* Subtracting gives $A\Delta\mathbf{x} + \Delta A(\mathbf{x}+\Delta\mathbf{x}) = \mathbf{0}$, so $\Delta\mathbf{x} = -A^{-1}\Delta A(\mathbf{x}+\Delta\mathbf{x})$.
Thus $\|\Delta\mathbf{x}\|_2 \leq \left\|A^{-1}\right\|_2 \|\Delta A\|_2 \|\mathbf{x}+\Delta\mathbf{x}\|_2 = \kappa_2(A)\frac{\|\Delta A\|_2}{\|A\|_2}\|\mathbf{x}+\Delta\mathbf{x}\|_2$. Divide by $\|\mathbf{x}+\Delta\mathbf{x}\|_2 \neq 0$.
*Structural Commentary*: Perturbing the matrix changes the operator applied to the perturbed solution.
*Key Insight*: The same condition number controls matrix errors and right-hand-side errors. □




Figure 97: $A$ maps a unit error ball to an ellipse; $\kappa_2(A) = \sigma_{\max}/\sigma_{\min}$ is the long/short semiaxis ratio.

## 5.4 Perturbation Bounds

Perturbation theory studies how matrix information changes under small errors. Bauer–Fike, Gersgorin, interlacing, Weyl inequalities, and pseudospectra measure robustness.

**Lemma 5.4.1.** *Small Spectral-Norm Perturbations of the Identity*
*If $E \in \mathrm{Mat}(n,n,\mathbb{C})$ and $\|E\|_2 < 1$, then $I_n + E$ is invertible.*

*Proof.* Theorem 5.1.12 gives $\rho(E) \leq \|E\|_2 < 1$, so $-1 \notin \mathrm{spec}(E)$.
Hence $0 \notin \mathrm{spec}(I_n+E)$, and $I_n+E$ is invertible.
*Structural Commentary*: Spectral-norm smallness traps every eigenvalue inside the unit disk.
*Key Insight*: If $-1$ is not an eigenvalue of $E$, then $I+E$ has no kernel. □

**Theorem 5.4.2.** *Bauer–Fike Eigenvalue Perturbation Bound*
*Let $A \in \mathrm{Mat}(n,n,\mathbb{C})$ be diagonalizable, with $A = S\Lambda S^{-1}$ and $\Lambda = \mathrm{diag}(\lambda_1,\ldots,\lambda_n)$. If $\Delta A \in \mathrm{Mat}(n,n,\mathbb{C})$ and $\mu \in \mathrm{spec}(A+\Delta A)$, then $\min_{1\leq j\leq n}|\mu-\lambda_j| \leq \kappa_2(S)\|\Delta A\|_2$.*

*Proof.* If $\mu = \lambda_j$ for some $j$, the claim is immediate. Otherwise $\Lambda - \mu I_n$ is invertible.
Since $\mu$ is an eigenvalue of $A+\Delta A$, the matrix $A+\Delta A-\mu I_n$ is singular.
Conjugating gives $S^{-1}(A+\Delta A-\mu I_n)S = (\Lambda-\mu I_n)\left(I_n + (\Lambda-\mu I_n)^{-1}S^{-1}\Delta AS\right)$ so the second factor is singular.
Lemma 5.4.1 implies $1 \leq \left\|(\Lambda-\mu I_n)^{-1}S^{-1}\Delta AS\right\|_2 \leq \frac{\left\|S^{-1}\right\|_2\|\Delta A\|_2\|S\|_2}{\min_j|\mu-\lambda_j|}$. Rearranging gives the bound.
*Structural Commentary*: Diagonalize, then test how perturbation breaks shifted invertibility.
*Key Insight*: Eigenvalue sensitivity is controlled by how ill-conditioned the eigenvector matrix $S$ is. □

**Method 5.4.3.** *Bauer–Fike Perturbation Method*
To bound eigenvalue movement for a diagonalizable matrix:

(i) diagonalize $A = S\Lambda S^{-1}$;
(ii) estimate $\|\Delta A\|_2$;
(iii) compute or bound $\kappa_2(S)$;
(iv) place every eigenvalue of $A+\Delta A$ inside the union of disks $|z-\lambda_j| \leq \kappa_2(S)\|\Delta A\|_2$.

**Definition 5.4.4.** *Gersgorin Disks*
Let $A = [a_{ij}] \in \mathrm{Mat}(n,n,\mathbb{C})$. The *deleted absolute row sum* in row $k$ is $R_k^{\mathrm{off}}(A) := \sum_{j\neq k}|a_{kj}|$.

The *$k$th Gersgorin disk* of $A$ is $\mathrm{Ger}_k(A) := \{z \in \mathbb{C} : |z-a_{kk}| \leq R_k^{\mathrm{off}}(A)\}$, where $\mathrm{Ger}(A) := \bigcup_{k=1}^{n}\mathrm{Ger}_k(A)$.

**Theorem 5.4.5.** *Gersgorin Disk Theorem*
*Let $A = [a_{ij}] \in \mathrm{Mat}(n,n,\mathbb{C})$. Then $\mathrm{spec}(A) \subseteq \mathrm{Ger}(A) = \bigcup_{k=1}^{n}\mathrm{Ger}_k(A)$.*

*Proof.* Let $\lambda \in \operatorname{spec}(A)$ and choose an eigenvector $\mathbf{x} = [x_1 \ \cdots \ x_n]^t$.
Pick $k$ such that $|x_k| = \max_j |x_j|$; then $x_k \neq 0$. The $k$th row of $A\mathbf{x} = \lambda\mathbf{x}$ gives $(\lambda - a_{kk})x_k = \sum_{j\neq k} a_{kj}x_j$.

Divide by $x_k$ and use $|x_j/x_k| \leq 1$: $|\lambda - a_{kk}| = \left|\sum_{j\neq k} a_{kj}\frac{x_j}{x_k}\right| \leq \sum_{j\neq k} |a_{kj}|\left|\frac{x_j}{x_k}\right| \leq R_k^{\text{off}}(A)$.

Thus $\lambda \in \operatorname{Ger}_k(A) \subseteq \operatorname{Ger}(A)$.
*Structural Commentary*: Normalize by a largest component; all row ratios have modulus at most one.
*Key Insight*: Row $k$ traps $\lambda$ in the disk centered at $a_{kk}$ with radius $R_k^{\text{off}}(A)$. □

**Corollary 5.4.6.** *Row-Column Gersgorin Refinement*
*Let* $A \in \operatorname{Mat}(n, n, \mathbb{C})$. *Then* $\operatorname{spec}(A) \subseteq \operatorname{Ger}(A) \cap \operatorname{Ger}(A^t)$.
*If* $R_k(A) = \sum_{j=1}^{n} |a_{kj}|$ *and* $R_{\max}(A) = \max_{1\leq k\leq n} R_k(A)$, *then* $\operatorname{spec}(A) \subseteq \{z \in \mathbb{C} : |z| \leq \min(R_{\max}(A), R_{\max}(A^t))\}$.

*Proof.* Theorem 5.4.5 gives $\operatorname{spec}(A) \subseteq \operatorname{Ger}(A)$.
Proposition 1.2.6 gives $\operatorname{spec}(A) = \operatorname{spec}(A^t)$, so applying Gersgorin to $A^t$ gives $\operatorname{spec}(A) \subseteq \operatorname{Ger}(A^t)$.
Also $\operatorname{Ger}_k(A) \subseteq \{z : |z| \leq |a_{kk}| + R_k^{\text{off}}(A) = R_k(A)\}$, and similarly for $A^t$.
*Structural Commentary*: Row disks and column disks are both valid because $A$ and $A^t$ have the same spectrum.
*Key Insight*: Intersecting independent enclosures usually sharpens the eigenvalue search region. □

**Method 5.4.7.** *Gersgorin Localisation Method*
For $A = [a_{ij}]$, Gersgorin localisation gives a disk enclosure for the spectrum.

(i) For each row $k$, place a disk at the diagonal entry $a_{kk}$ with radius $R_k^{\text{off}}(A) = \sum_{j\neq k} |a_{kj}|$.
The row enclosure is $\operatorname{Ger}(A) = \bigcup_{k=1}^{n} \operatorname{Ger}_k(A)$.
(ii) Theorem 5.4.5 guarantees that every eigenvalue of $A$ lies in $\operatorname{Ger}(A)$.
(iii) For a sharper enclosure, repeat the same construction for $A^t$. The centers are still $a_{kk}$, while the radii becomes deleted column sums of $A$. Then use $\operatorname{spec}(A) \subseteq \operatorname{Ger}(A) \cap \operatorname{Ger}(A^t)$. This is the intersection of two unions of disks; an eigenvalue need not lie in the overlap of each individual row-column disk pair.




Figure 98: Blue row disks contain the spectrum; green dashed column disks give the sharper row-column enclosure $\operatorname{Ger}(A) \cap \operatorname{Ger}(A^t)$. Red points illustrate possible eigenvalue locations.

**Perspective 5.4.8.** *Pseudospectral Intuition*
For a matrix $A$, the $\varepsilon$-pseudospectrum records complex numbers $z$ for which $A - zI$ is singular or nearly singular. Equivalently, it asks where a perturbation of size at most $\varepsilon$ can move the spectrum. Bauer–Fike gives a clean enclosure for diagonalizable matrices, while Gersgorin and norm bounds give cruder but often cheaper enclosures. The main lesson is that eigenvalues of nonnormal matrices can be much more sensitive than their locations alone suggest.

## 5.5 Diagonal Dominance and Definiteness

**Definition 5.5.1.** *Diagonal Dominance*
Let $A = [a_{ij}] \in \operatorname{Mat}(n, n, \mathbb{C})$. Matrix $A$ is *diagonally dominant* if $|a_{kk}| \geq R_k^{\text{off}}(A)$ for every $k$, and *strictly diagonally dominant* if $|a_{kk}| > R_k^{\text{off}}(A)$ for every $k$.

**Corollary 5.5.2.** *Strict Diagonal Dominance Implies Invertibility*
*Let* $A \in \mathrm{Mat}(n, n, \mathbb{C})$.

*(i) If $A$ is strictly diagonally dominant, then $A$ is invertible.*
*(ii) If $A$ has real spectrum and nonnegative diagonal entries, then diagonal dominance gives* $\mathrm{spec}(A) \subseteq [0, \infty)$*; strict diagonal dominance gives* $\mathrm{spec}(A) \subseteq (0, \infty)$.

*Proof.*

(i) Strict dominance gives $0 \notin \mathrm{Ger}_k(A)$ for every $k$, since $|a_{kk}| > R_k^{\mathrm{off}}(A)$. Thus $0 \notin \mathrm{spec}(A)$ by Gersgorin, so $A$ is invertible by Corollary 5.1.8.
(ii) If $a_{kk} \geq 0$ and $a_{kk} \geq R_k^{\mathrm{off}}(A)$, then each $\mathrm{Ger}_k(A)$ lies in the closed right half-plane. In the strict case, each disk lies in the open right half-plane. Gersgorin and the real-spectrum hypothesis give the claim.

*Structural Commentary*: Gersgorin converts diagonal dominance into disk placement.
*Key Insight*: The disks control whether eigenvalues can reach 0 or the left half-plane. □

**Corollary 5.5.3.** *Diagonal Dominance and Positive Semidefiniteness*
*Let* $A = [a_{ij}] \in \mathrm{Mat}(n, n, \mathbb{C})$ *be Hermitian with nonnegative diagonal entries.*

*(i) If $A$ is diagonally dominant, then $A$ is positive semidefinite.*
*(ii) If $A$ is diagonally dominant and invertible, then $A$ is positive definite.*
*(iii) If $A$ is strictly diagonally dominant, then $A$ is positive definite.*

*Proof.* Hermitian matrices have real spectrum.

(i) Corollary 5.5.2 gives $\mathrm{spec}(A) \subseteq [0, \infty)$; hence $A \succeq 0$ by Theorem 2.1.3.
(ii) If $A$ is invertible, then $0 \notin \mathrm{spec}(A)$, so every eigenvalue is positive; hence $A \succ 0$.
(iii) Strict diagonal dominance implies invertibility by Corollary 5.5.2; apply (ii).

*Structural Commentary*: Gersgorin traps the real spectrum in $[0, \infty)$; invertibility removes 0.
*Key Insight*: For Hermitian matrices, positivity is spectral location. □

**Theorem 5.5.4.** *Weak Diagonal Dominance Criterion*
*Let* $A = [a_{ij}] \in \mathrm{Mat}(n, n, \mathbb{C})$ *be diagonally dominant with no zero entries.*
*If* $|a_{kk}| > R_k^{\mathrm{off}}(A)$ *for some $k$, then $A$ is invertible.*

*Proof.* Assume $A\mathbf{x} = 0$ with $\mathbf{x} \neq \mathbf{0}$, and choose $k$ with $|x_k| = \|\mathbf{x}\|_\infty$.

Row $k$ gives $|a_{kk}| \, \|\mathbf{x}\|_\infty = \left| -\sum_{j \neq k} a_{kj} x_j \right| \leq \sum_{j \neq k} |a_{kj}| \, |x_j| \leq R_k^{\mathrm{off}}(A) \, \|\mathbf{x}\|_\infty \leq |a_{kk}| \, \|\mathbf{x}\|_\infty$.

Hence all inequalities are equalities. Since no $a_{kj}$ vanishes, $|x_j| = \|\mathbf{x}\|_\infty$ for every $j$. Applying the same chain to any row $i$ gives $|a_{ii}| = R_i'(A)$ for every $i$, contradicting the strict row. Thus $\ker(A) = \{\mathbf{0}\}$, so $A$ is invertible.
*Structural Commentary*: A zero vector in the kernel would force equality in every Gersgorin row.
*Key Insight*: One strict row breaks that equality pattern. □

## 5.6 Rayleigh Quotients and Minimax

**Definition 5.6.1.** *Increasing Hermitian Eigenvalue Order*
For a Hermitian matrix $A \in \mathrm{Mat}(n, n, \mathbb{C})$, write $\lambda_1(A) \leq \lambda_2(A) \leq \cdots \leq \lambda_n(A)$ for its eigenvalues counted with algebraic multiplicity. If only one Hermitian matrix is under discussion, we write $\lambda_i$ instead of $\lambda_i(A)$.

**Theorem 5.6.2.** *Rayleigh Quotient on a Spectral Interval*
*Let* $A \in \mathrm{Mat}(n, n, \mathbb{C})$ *be Hermitian with increasing eigenvalues* $\lambda_1 \leq \cdots \leq \lambda_n$ *and corresponding orthonormal eigenvectors* $\mathbf{u}_1, \ldots, \mathbf{u}_n$*. For a unit vector, write* $\mathrm{Ray}_A(\mathbf{x}) := \mathbf{x}^* A \mathbf{x}$*. If* $1 \leq p \leq q \leq n$ *and* $\mathbf{x} \in \mathrm{span}(\mathbf{u}_p, \ldots, \mathbf{u}_q)$ *is a unit vector, then* $\lambda_p \leq \mathrm{Ray}_A(\mathbf{x}) \leq \lambda_q$.

*Proof.* Write $\mathbf{x} = \sum_{i=p}^{q} c_i \mathbf{u}_i$. Orthonormality gives $\sum_{i=p}^{q} |c_i|^2 = 1$, and $\mathbf{x}^* A \mathbf{x} = \sum_{i=p}^{q} \lambda_i \, |c_i|^2$.

This is a convex combination of $\lambda_p, \ldots, \lambda_q$, so it lies between the smallest and largest of them.
*Structural Commentary*: In an eigenbasis, the Rayleigh quotient is an eigenvalue average.
*Key Insight*: Spectral coordinates turn a quadratic form into weighted scalar data. □

**Corollary 5.6.3.** *Rayleigh Extrema and Diagonal Entry Bounds*
*Let* $A \in \mathrm{Mat}(n, n, \mathbb{C})$ *be Hermitian.*

*(i) If* $\mathcal{U}_{p,q} = \mathrm{span}(\mathbf{u}_p, \ldots, \mathbf{u}_q)$*, then* $\min_{\substack{\mathbf{x} \in \mathcal{U}_{p,q} \\ \|\mathbf{x}\|_2 = 1}} \mathbf{x}^* A \mathbf{x} = \lambda_p(A)$*, and* $\max_{\substack{\mathbf{x} \in \mathcal{U}_{p,q} \\ \|\mathbf{x}\|_2 = 1}} \mathbf{x}^* A \mathbf{x} = \lambda_q(A)$.

*(ii) For every unit vector* $\mathbf{x}$, $\lambda_1(A) \le \mathbf{x}^* A\mathbf{x} \le \lambda_n(A)$.
*(iii) If* $A = [a_{ij}]$, *then* $\lambda_1(A) \le a_{ii} \le \lambda_n(A)$ *for* $i = 1, \dots, n$.

*Proof.*

(i) Theorem 5.6.2 gives the inequalities on $\mathcal{U}_{p,q}$, with equality at $\mathbf{u}_p$ and $\mathbf{u}_q$.
(ii) This is (i) with $p = 1$ and $q = n$.
(iii) Apply (ii) to $\mathbf{x} = \mathbf{e}_i$, since $\mathbf{e}_i^* A\mathbf{e}_i = a_{ii}$.

*Structural Commentary*: Rayleigh extrema on extreme eigendirections; diagonal entries are coordinate quotients.
*Key Insight*: The whole numerical range of a Hermitian matrix sits inside its spectral interval. □




Figure 99: For $\mathbf{x} \in \mathcal{U}_{p,q}$, $\mathrm{Ray}_A(\mathbf{x})$ is a convex combination of $\lambda_p, \dots, \lambda_q$.

**Theorem 5.6.4.** *Rayleigh Minimax Principle*
*Let* $A = A^* \in \mathrm{Mat}(n, n, \mathbb{C})$ *have eigenvalues* $\lambda_1 \le \cdots \le \lambda_n$, *and set* $R_A(\mathbf{x}) = \mathbf{x}^* A\mathbf{x}/(\mathbf{x}^*\mathbf{x})$ *for* $\mathbf{x} \ne \mathbf{0}$.
*For* $1 \le k \le n$, $\lambda_k = \min\limits_{\substack{\mathcal{S} \subseteq \mathbb{C}^n \\ \dim \mathcal{S} = k}} \max\limits_{\mathbf{0} \ne \mathbf{x} \in \mathcal{S}} R_A(\mathbf{x}) = \max\limits_{\substack{\mathcal{S} \subseteq \mathbb{C}^n \\ \dim \mathcal{S} = n-k+1}} \min\limits_{\mathbf{0} \ne \mathbf{x} \in \mathcal{S}} R_A(\mathbf{x})$.

*Proof.* Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be an orthonormal eigenbasis in increasing order.
Set $E_k = \mathrm{span}(\mathbf{u}_1, \dots, \mathbf{u}_k)$; $F_k = \mathrm{span}(\mathbf{u}_k, \dots, \mathbf{u}_n)$.
By Theorem 5.6.2, $\max_{\mathbf{0} \ne \mathbf{x} \in E_k} R_A(\mathbf{x}) = \lambda_k$ and $\min_{\mathbf{0} \ne \mathbf{x} \in F_k} R_A(\mathbf{x}) = \lambda_k$.
If $\dim \mathcal{S} = k$, then $\mathcal{S} \cap F_k \ne \{\mathbf{0}\}$, so $\max_{\mathbf{0} \ne \mathbf{x} \in \mathcal{S}} R_A(\mathbf{x}) \ge \lambda_k$; equality is attained by $E_k$.
If $\dim \mathcal{T} = n - k + 1$, then $\mathcal{T} \cap E_k \ne \{\mathbf{0}\}$, so $\min_{\mathbf{0} \ne \mathbf{x} \in \mathcal{T}} R_A(\mathbf{x}) \le \lambda_k$; equality is attained by $F_k$.
*Structural Commentary*: Dimension forces intersections with spectral flags.
*Key Insight*: Minimax is Rayleigh's quotient plus dimension counting. □

**Method 5.6.5.** *Rayleigh Eigenbasis Method*
To bound a Hermitian quadratic form:

(i) diagonalise $A$ with an orthonormal eigenbasis;
(ii) expand the unit vector in that basis;
(iii) rewrite $\mathbf{x}^* A\mathbf{x}$ as $\sum_i \lambda_i |c_i|^2$;
(iv) use the eigenvalue range allowed by the support of $\mathbf{x}$.

## 5.7 Eigenvalue and Singular-Value Interlacing

**Theorem 5.7.1.** *Rank-One Positive Perturbation Interlacing*
*Let* $A, E \in \mathrm{Mat}(n, n, \mathbb{C})$ *be Hermitian, with* $E \succeq 0$ *and* $\mathrm{rank}(E) \le 1$.
*Then* $\lambda_i(A) \le \lambda_i(A + E) \le \lambda_{i+1}(A) \quad (1 \le i < n)$, $\lambda_n(A) \le \lambda_n(A + E)$
*Equivalently,* $\lambda_i(A - E) \le \lambda_i(A) \le \lambda_{i+1}(A - E) \quad (1 \le i < n)$, $\lambda_n(A - E) \le \lambda_n(A)$.

*Proof.* Let $\mathbf{u}_j$ and $\mathbf{v}_j$ be orthonormal eigenvectors of $A$ and $A + E$, respectively, in increasing eigenvalue order.
For the lower bound, set $\mathcal{U} = \mathrm{span}(\mathbf{u}_i, \dots, \mathbf{u}_n)$, $\mathcal{V} = \mathrm{span}(\mathbf{v}_1, \dots, \mathbf{v}_i)$. Since $\dim \mathcal{U} + \dim \mathcal{V} = n + 1$, choose a unit $\mathbf{x} \in \mathcal{U} \cap \mathcal{V}$. Then $\lambda_i(A) \le \mathbf{x}^* A\mathbf{x} \le \mathbf{x}^*(A + E)\mathbf{x} \le \lambda_i(A + E)$ by Theorem 5.6.2 and $E \succeq 0$.
For the upper bound, set $\mathcal{U} = \mathrm{span}(\mathbf{u}_1, \dots, \mathbf{u}_{i+1})$, $\mathcal{V} = \mathrm{span}(\mathbf{v}_i, \dots, \mathbf{v}_n)$, $\mathcal{W} = \ker(E)$. Here $\dim(\mathcal{U} \cap \mathcal{V}) \ge 2$ and $\dim \mathcal{W} \ge n - 1$, so choose a unit $\mathbf{x} \in \mathcal{U} \cap \mathcal{V} \cap \mathcal{W}$. Thus $\lambda_i(A + E) \le \mathbf{x}^*(A + E)\mathbf{x} = \mathbf{x}^* A\mathbf{x} \le \lambda_{i+1}(A)$.
The statements for $A - E$ follow by applying the proved result to $A - E$ and $E$.
*Structural Commentary*: Interlacing comes from test vectors in overlapping spectral subspaces.
*Key Insight*: Rank one leaves a large kernel, so the upper-bound vector can avoid the perturbation. □

**Corollary 5.7.2.** *Rank-Two Indefinite Interlacing*
*Let* $A, H \in \mathrm{Mat}(n, n, \mathbb{C})$ *be Hermitian,* $n \ge 2$. *Suppose* $\mathrm{rank}(H) \le 2$ *and* $H$ *has at most one positive and one negative eigenvalue. Then*

$$\lambda_1(A + H) \le \lambda_2(A), \qquad \lambda_{i-1}(A) \le \lambda_i(A + H) \le \lambda_{i+1}(A) \quad (2 \le i < n), \qquad \lambda_{n-1}(A) \le \lambda_n(A + H).$$

*Proof.* Write $H = E_+ - E_-$ with $E_\pm \succeq 0$ and $\operatorname{rank}(E_\pm) \le 1$. Set $B = A - E_-$; then $A = B + E_-$ and $A+H = B+E_+$. For $2 \le i < n$, rank-one interlacing gives $\lambda_{i-1}(A) \le \lambda_i(B) \le \lambda_i(A+H) \le \lambda_{i+1}(B) \le \lambda_{i+1}(A)$. The endpoints follow similarly: $\lambda_1(A+H) \le \lambda_2(B) \le \lambda_2(A)$, and $\lambda_{n-1}(A) \le \lambda_n(B) \le \lambda_n(A+H)$.
*Structural Commentary*: Such an $H$ is a difference of two rank-one PSD perturbations.
*Key Insight*: Interlace once with $E_-$, then once with $E_+$. □

**Theorem 5.7.3.** *Cauchy Interlacing: One Principal Deletion*
*Let* $A = \begin{bmatrix} B & \mathbf{y} \\ \mathbf{y}^* & c \end{bmatrix} \in \operatorname{Mat}(n+1, n+1, \mathbb{C})$ *be Hermitian, with* $B \in \operatorname{Mat}(n, n, \mathbb{C})$.
*Then, for* $1 \le i \le n$, $\lambda_i(A) \le \lambda_i(B) \le \lambda_{i+1}(A)$.

*Proof.* Let $\mathbf{u}_j$ and $\mathbf{v}_j$ be orthonormal eigenvectors of $A$ and $B$, respectively, in increasing eigenvalue order. Identify $\mathbb{C}^n$ with $\{(\mathbf{z}, 0)\} \subseteq \mathbb{C}^{n+1}$, so $(\mathbf{z},0)^* A (\mathbf{z},0) = \mathbf{z}^* B \mathbf{z}$.
For the lower bound, set $\mathcal{S} = \operatorname{span}((\mathbf{v}_1,0), \dots, (\mathbf{v}_i,0))$ and $\mathcal{T} = \operatorname{span}(\mathbf{u}_i, \dots, \mathbf{u}_{n+1})$. Their dimensions are $i$ and $n+2-i$, so $\dim \mathcal{S} + \dim \mathcal{T} > n+1$ and $\mathcal{S} \cap \mathcal{T}$ contains a nonzero vector.
Choose a unit $\mathbf{x} = (\mathbf{z}, 0)$ in the intersection. The spectral expansion in the $\mathbf{u}_j$ gives $\mathbf{x}^* A \mathbf{x} \ge \lambda_i(A)$, while the expansion in the $\mathbf{v}_j$ gives $\mathbf{z}^* B \mathbf{z} \le \lambda_i(B)$. Hence $\lambda_i(A) \le \lambda_i(B)$.
For the upper bound, take $\mathcal{S}' = \operatorname{span}((\mathbf{v}_i,0), \dots, (\mathbf{v}_n,0))$ and $\mathcal{T}' = \operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_{i+1})$. Again the dimensions sum to $n+2$, so choose a unit $\mathbf{x} = (\mathbf{z}, 0) \in \mathcal{S}' \cap \mathcal{T}'$.
Now $\mathbf{z}^* B \mathbf{z} \ge \lambda_i(B)$ and $\mathbf{x}^* A \mathbf{x} \le \lambda_{i+1}(A)$, giving $\lambda_i(B) \le \lambda_{i+1}(A)$.
*Structural Commentary*: A principal submatrix is a coordinate slice of the larger quadratic form.
*Key Insight*: Deleting one coordinate costs one dimension, so eigenvalues shift by at most one index. □

**Corollary 5.7.4.** *Singular Values Interlace After Deleting a Row or Column*
*Let* $B$ *be obtained from* $A \in \operatorname{Mat}(n, n, \mathbb{C})$ *by deleting one row or column.*
*With singular values ordered decreasingly,* $\sigma_i(A) \ge \sigma_i(B) \ge \sigma_{i+1}(A) \quad (1 \le i < n)$.

*Proof.* For a row deletion, permute rows so that $PA = \begin{bmatrix} B \\ \mathbf{y}^* \end{bmatrix}$.
Then $PA$ has the same singular values as $A$, and $BB^*$ is a principal submatrix of $(PA)(PA)^*$.
Cauchy interlacing applied to these Gram matrices gives the result after taking square roots and reversing the increasing eigenvalue order. A column deletion follows by applying the row case to $A^*$.
*Structural Commentary*: Row deletion makes $BB^*$ a principal submatrix of a permuted $AA^*$.
*Key Insight*: Singular-value interlacing is Cauchy interlacing applied to Gram matrices. □

**Corollary 5.7.5.** *Cauchy Interlacing for Principal Submatrices*
*Let* $B \in \operatorname{Mat}(n, n, \mathbb{C})$ *be a principal submatrix of Hermitian* $A \in \operatorname{Mat}(n+m, n+m, \mathbb{C})$.
*Then, for* $1 \le i \le n$, $\lambda_i(A) \le \lambda_i(B) \le \lambda_{i+m}(A)$.

*Proof.* After a permutation similarity, $B$ is leading.
Delete the other $m$ coordinates one at a time and apply Theorem 5.7.3 at each step.
*Structural Commentary*: Principal submatrices come from repeated coordinate deletion.
*Key Insight*: Each deletion shifts eigenvalue indices by at most one. □




Figure 100: Rank-one and principal-submatrix interlacing place new eigenvalues between old ones.

**Theorem 5.7.6.** *Schur Majorization of the Diagonal*
*Let* $A \in \operatorname{Mat}(n, n, \mathbb{C})$ *be Hermitian, with increasing eigenvalues* $\lambda_i(A)$ *and increasing diagonal entries* $d_i$.
*Then* $\sum_{i=1}^{k} \lambda_i(A) \le \sum_{i=1}^{k} d_i$, $(1 \le k \le n)$ *with equality for* $k = n$.

*Proof.* Permute coordinates so that $a_{ii} = d_i$, and let $A_k$ be the leading $k \times k$ principal submatrix.
Cauchy interlacing gives $\lambda_i(A) \le \lambda_i(A_k)$ for $1 \le i \le k$, hence $\sum_{i=1}^{k} \lambda_i(A) \le \sum_{i=1}^{k} \lambda_i(A_k) = \operatorname{tr}(A_k) = \sum_{i=1}^{k} d_i$.
For $k = n$, this is $\operatorname{tr}(A) = \operatorname{tr}(A)$.
*Structural Commentary*: Ordered diagonal sums are traces of leading principal blocks.
*Key Insight*: Diagonal majorization is Cauchy interlacing plus trace. □

**Method 5.7.7.** *Eigenvalue Interlacing Method*
To prove or use an interlacing estimate:

(i) order Hermitian eigenvalues increasingly;
(ii) identify dimension lost: rank-one perturbation, one deleted coordinate, $m$-coordinate principal submatrix;
(iii) choose test vectors by intersecting spectral subspaces with the unaffected subspace;
(iv) convert the Rayleigh quotient bounds into eigenvalue inequalities.

## 5.8 Weyl Eigenvalue Bounds

**Theorem 5.8.1.** *Weyl Eigenvalue Inequalities*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{C})$ be Hermitian. For $1 \le i \le n$,*

$$\lambda_i(A+B) \le \lambda_{i+j}(A) + \lambda_{n-j}(B), \qquad 0 \le j \le n-i,$$
$$\lambda_{i-j+1}(A) + \lambda_j(B) \le \lambda_i(A+B), \qquad 1 \le j \le i.$$

*Proof.* Let $\mathbf{u}_k, \mathbf{v}_k, \mathbf{w}_k$ be increasing eigenbases for $A, A+B, B$.
Fix $i$ and $0 \le j \le n-i$, and set $\mathcal{U} = \mathrm{span}(\mathbf{u}_1, \dots, \mathbf{u}_{i+j})$, $\mathcal{V} = \mathrm{span}(\mathbf{v}_i, \dots, \mathbf{v}_n)$, $\mathcal{W} = \mathrm{span}(\mathbf{w}_1, \dots, \mathbf{w}_{n-j})$.
Their dimensions sum to $2n+1$, so choose a unit $\mathbf{x} \in \mathcal{U} \cap \mathcal{V} \cap \mathcal{W}$.
Then $\lambda_i(A+B) \le \mathbf{x}^*(A+B)\mathbf{x} = \mathbf{x}^*A\mathbf{x} + \mathbf{x}^*B\mathbf{x} \le \lambda_{i+j}(A) + \lambda_{n-j}(B)$.
The lower bound follows by applying this upper bound to $-A$ and $-B$ and using $\lambda_k(-C) = -\lambda_{n-k+1}(C)$.
*Structural Commentary*: Weyl bounds come from intersecting the spectral flags of $A$, $B$, and $A+B$.
*Key Insight*: One shared test vector controls all three Rayleigh quotients. □

**Method 5.8.2.** *Weyl Bound Method*
To bound eigenvalues of $A+B$:

(i) choose the target index $i$;
(ii) use the upper Weyl family with $j = 0, \dots, n-i$ to get upper bounds;
(iii) use the lower Weyl family with $j = 1, \dots, i$ to get lower bounds;
(iv) for positive semidefinite or low-rank $B$, insert the known zero or nonnegative eigenvalues to recover monotonicity or interlacing.

## 5.9 Brauer Rank-One Spectral Shifts

**Theorem 5.9.1.** *Brauer Rank-One Eigenvalue Shift*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$, let $A\mathbf{x} = \lambda\mathbf{x}$ with $\mathbf{x} \neq \mathbf{0}$, and let $\mathbf{y} \in \mathbb{C}^n$. If $\lambda, \lambda_2, \dots, \lambda_n$ are the eigenvalues of $A$ with multiplicity, then the eigenvalues of $A + \mathbf{x}\mathbf{y}^*$ are $\lambda + \mathbf{y}^*\mathbf{x}, \lambda_2, \dots, \lambda_n$.*
*Moreover, $\mathbf{x}$ is an eigenvector of $A + \mathbf{x}\mathbf{y}^*$ corresponding to $\lambda + \mathbf{y}^*\mathbf{x}$.*

*Proof.* Put $\mathbf{u} = \mathbf{x}/\|\mathbf{x}\|_2$. By Schur triangularisation, choose a unitary $U = [\mathbf{u}\ U_2]$ such that $U^*AU = \begin{bmatrix} \lambda & * \\ 0 & T' \end{bmatrix}$, where the diagonal entries of $T'$ are $\lambda_2, \dots, \lambda_n$. Also $U^*(\mathbf{x}\mathbf{y}^*)U = \begin{bmatrix} \mathbf{y}^*\mathbf{x} & * \\ 0 & 0 \end{bmatrix}$. Therefore $U^*(A + \mathbf{x}\mathbf{y}^*)U$ is upper triangular with diagonal entries $\lambda + \mathbf{y}^*\mathbf{x}, \lambda_2, \dots, \lambda_n$. Finally, $(A + \mathbf{x}\mathbf{y}^*)\mathbf{x} = A\mathbf{x} + \mathbf{x}(\mathbf{y}^*\mathbf{x}) = (\lambda + \mathbf{y}^*\mathbf{x})\mathbf{x}$.
*Structural Commentary*: A rank-one eigenvector update changes only the first Schur diagonal entry.
*Key Insight*: Align the update with the first coordinate; only one diagonal term shifts. □




Figure 101: Brauer's update shifts one Schur diagonal entry.

**Corollary 5.9.2.** *Damped Rank-One Eigenvalue Reset*
*Let $A \in \mathrm{Mat}(n, n, \mathbb{C})$ have eigenpair $(\lambda, \mathbf{x})$, let $\mathbf{y}^*\mathbf{x} = 1$, and let $\tau \in \mathbb{C}$. If $\lambda, \lambda_2, \dots, \lambda_n$ are the eigenvalues of $A$, then $A_\tau := \tau A + (1-\tau)\lambda\mathbf{x}\mathbf{y}^*$ has eigenvalues $\lambda, \tau\lambda_2, \dots, \tau\lambda_n$.*

*Proof.* The eigenvalues of $\tau A$ are $\tau\lambda, \tau\lambda_2, \dots, \tau\lambda_n$. Since $A_\tau = \tau A + \mathbf{x}((1-\overline{\tau})\overline{\lambda}\mathbf{y})^*$, Theorem 5.9.1 shifts only $\tau\lambda$ by $(1-\tau)\lambda\mathbf{y}^*\mathbf{x} = (1-\tau)\lambda$. Thus $\tau\lambda$ becomes $\lambda$, while the other eigenvalues stay $\tau\lambda_i$.
*Structural Commentary*: Damping scales the spectrum; Brauer's update restores one eigenvalue.
*Key Insight*: Brauer's theorem changes one spectral coordinate without moving the rest. □

# 6 Numerical Linear Algebra

## 6.1 Direct Linear-System Solves

Numerical linear algebra turns matrix structure into algorithms for direct solves, least squares, eigenvalues, SVD, structured transforms, Krylov methods, iteration, and preconditioning.

**Perspective 6.1.1.** *Solve with Factors Before Forming Inverses*
For $A\mathbf{x} = \mathbf{b}$, use available factors such as LU, QR, or Cholesky to solve triangular or orthogonal subproblems. This avoids forming $A^{-1}$ and preserves sparse or triangular structure.

**Perspective 6.1.2.** *Iterative Refinement by Residual Correction*
Given an approximate solution $\mathbf{x}_k$ of $A\mathbf{x} = \mathbf{b}$, set $\mathbf{r}_k = \mathbf{b} - A\mathbf{x}_k$.
Solve $A\mathbf{d}_k = \mathbf{r}_k$ with the same factors or preconditioner, then update $\mathbf{x}_{k+1} = \mathbf{x}_k + \mathbf{d}_k$.
Exact arithmetic corrects in one step; floating point helps when $\mathbf{r}_k$ is accurate and conditioning is moderate.

## 6.2 QR Least Squares

**Proposition 6.2.1.** *QR Route for Least Squares*
*Let $A = QR \in \mathrm{Mat}(m, n, \mathbb{F})$ be a full-column-rank narrow QR factorisation. For $\mathbf{y} \in \mathbb{F}^m$, the least-squares solution solves $R\mathbf{x} = Q^*\mathbf{y}$. Moreover $P_{\mathrm{col}(A)} = QQ^*$.*

*Proof.* The normal equations reduce by $A = QR$ and $Q^*Q = I_n$ to $R^*R\mathbf{x} = R^*Q^*\mathbf{y}$. Since $R$ is invertible, this is equivalent to $R\mathbf{x} = Q^*\mathbf{y}$. Also $\mathrm{col}(A) = \mathrm{col}(Q)$, and the columns of $Q$ are orthonormal, so $P_{\mathrm{col}(A)} = QQ^*$.
*Structural Commentary*: QR replaces $\mathrm{col}(A)$ by an orthonormal basis $Q$.
*Key Insight*: Project onto $Q$, then solve triangularly. □

Successive orthogonalization fits in an orthogonal basis for the same column space.

**Algorithm 1:** Least Squares by Successive Orthogonalization

**Require:** Full-column-rank design columns $x_0, \ldots, x_p$ and response vector $y$
**Ensure:** Least-squares coefficients $\widehat{\beta}$
  **for** $j = 0, \ldots, p$ **do**
    $z_j \leftarrow x_j$
    **for** $\ell = 0, \ldots, j-1$ **do**
      $r_{\ell j} \leftarrow q_\ell^T x_j$
      $z_j \leftarrow z_j - r_{\ell j} q_\ell$
    **end for**
    $r_{jj} \leftarrow \|z_j\|_2$
    $q_j \leftarrow z_j / r_{jj}$
  **end for**
  Form $Q = [q_0, \ldots, q_p]$ and the upper-triangular matrix $R = (r_{\ell j})$
  Solve $R\widehat{\beta} = Q^T y$ by back-substitution

**Perspective 6.2.2.** *Why Normal Equations Can Be Numerically Fragile*
For full-column-rank least squares, solving $A^*A\mathbf{x} = A^*\mathbf{y}$ squares the spectral condition number because $\kappa_2(A^*A) = \kappa_2(A)^2$. QR-based methods avoid this squaring and therefore preserve conditioning better.

**Perspective 6.2.3.** *Solve Least Squares with Factors*
For $X \in \mathbb{R}^{n \times p}$ with $\mathrm{rank}(X) = p$, normal equations use $X^tX\widehat{\beta} = X^tY$ but $\kappa_2(X^tX) = \kappa_2(X)^2$.
Prefer $X = QR$ with $Q^tQ = I_p$ and $R$ nonsingular, giving $\widehat{\beta} = R^{-1}Q^tY$.
Use Cholesky on $X^tX$ only when $X^tX$ is well conditioned.

## 6.3 Banded and Sparse Direct Methods

**Definition 6.3.1.** *Band and Tridiagonal Matrices*
For $A = [a_{ij}] \in \mathrm{Mat}(n, n, \mathbb{F})$ and $1 \le w \le n$, say that $A$ has *half-bandwidth* $w$ if $a_{ij} = 0$ whenever $|i - j| \ge w$. Thus $w = 1$ gives a diagonal matrix, while $w = 2$ gives a *tridiagonal* matrix.

**Method 6.3.2.** *Banded Elimination Preview*
Let $A \in \mathrm{Mat}(n, n, \mathbb{F})$ have half-bandwidth $w$.

(i) Eliminate without row exchanges that leave the band.
(ii) At pivot $k$, update only entries with $|i - k| < w$ and $|j - k| < w$.

(iii) Obtain $A = LU$ with $L$ and $U$ of comparable bandwidth.

Cost: $O(nw^2)$ operations; for tridiagonal matrices ($w = 2$), $O(n)$.




Figure 102: A tridiagonal matrix can factor as bidiagonal $L$ times bidiagonal $U$, retaining local band structure.

**Perspective 6.3.3.** *Five-Point Stencils Are Banded but Less Local*

The 1D second-difference pattern $-1, 2, -1$ is tridiagonal; the row-ordered $N \times N$ five-point stencil is still sparse, but vertical neighbors create diagonals offset by about $N$. Thus bandwidth grows with grid width, and incomplete LU keeps a prescribed sparse pattern for a cheap approximate solve or preconditioner.




Figure 103: Row ordering gives matrix bands at offsets $0, \pm 1, \pm N$.

## 6.4 Krylov Subspace Methods

**Definition 6.4.1.** *Krylov Subspace*

For $A \in \mathrm{Mat}(n, n, \mathbb{F})$ and a vector $\mathbf{b} \in \mathbb{F}^n$, the $m$th *Krylov subspace* generated by $(A, \mathbf{b})$ is $\mathcal{K}_m(A, \mathbf{b}) := \mathrm{span}\{\mathbf{b}, A\mathbf{b}, \ldots, A^{m-1}\mathbf{b}\}$.

**Perspective 6.4.2.** *Krylov Methods Preview*

Krylov methods work in $\mathcal{K}_m(A, \mathbf{b})$: Arnoldi and Lanczos build bases, while CG and GMRES impose projection or residual-minimization conditions using only matrix-vector products.

**Method 6.4.3.** *Arnoldi, Lanczos, and CG Preview*

Let $\mathcal{K}_m = \mathcal{K}_m(A, \mathbf{b})$ and let $Q_m = [\mathbf{q}_1 \ \cdots \ \mathbf{q}_m]$ be an orthonormal basis.

(i) *Arnoldi*: $AQ_m = Q_m H_m + h_{m+1,m}\mathbf{q}_{m+1}\mathbf{e}_m^t$, with $H_m$ upper Hessenberg.
(ii) *Lanczos*: if $A = A^*$, then $H_m$ is tridiagonal and $A\mathbf{q}_j = \beta_{j-1}\mathbf{q}_{j-1} + \alpha_j\mathbf{q}_j + \beta_j\mathbf{q}_{j+1}$.
(iii) *CG*: if $A \succ 0$, choose $\mathbf{x}_k \in \mathbf{x}_0 + \mathcal{K}_k(A, \mathbf{r}_0)$ with $\mathbf{r}_k \perp \mathcal{K}_k(A, \mathbf{r}_0)$; search directions satisfy $\mathbf{p}_i^* A\mathbf{p}_j = 0$, $i \neq j$.

Each step uses one matrix-vector product.

## 6.5 Stationary Iterations and Splittings

**Definition 6.5.1.** *Matrix Splitting Iteration*

Let $A = S - T$ with $S$ invertible, and set $B := S^{-1}T$.
The *splitting iteration* for $A\mathbf{x} = \mathbf{b}$ is $S\mathbf{x}_{k+1} = T\mathbf{x}_k + \mathbf{b}$, and $\mathbf{x}_{k+1} = B\mathbf{x}_k + S^{-1}\mathbf{b}$.
For a solution $\mathbf{x}_*$, the error $\mathbf{e}_k := \mathbf{x}_k - \mathbf{x}_*$ satisfies $\mathbf{e}_{k+1} = B\mathbf{e}_k$.

**Theorem 6.5.2.** *Splitting Convergence Criterion*

*Let $A = S - T$, with $S$ invertible, and set $B = S^{-1}T$. For $\mathbf{x}_{k+1} = B\mathbf{x}_k + S^{-1}\mathbf{b}$:*

*(i) if $A$ is invertible, then $\mathbf{x}_k \to A^{-1}\mathbf{b}$ for all $\mathbf{x}_0, \mathbf{b}$ if and only if $\rho(B) < 1$;*
*(ii) $\rho(B) < 1$ implies $A$ is invertible.*

*Proof.*

(i) Let $\mathbf{x}_* = A^{-1}\mathbf{b}$ and $\mathbf{e}_k = \mathbf{x}_k - \mathbf{x}_*$. Then $\mathbf{e}_k = B^k\mathbf{e}_0$, so convergence for all $\mathbf{x}_0, \mathbf{b}$ is equivalent to $B^k \to 0$, equivalently $\rho(B) < 1$ by Theorem 3.3.5.

(ii) If $\rho(B) < 1$, then $1 \notin \operatorname{spec}(B)$, so $I_n - B$ is invertible. Since $A = S - T = S(I_n - B)$, $A$ is invertible.

*Structural Commentary*: A stationary method is a matrix-power problem in disguise.
*Key Insight*: The iteration matrix $B = S^{-1}T$ controls the errors. □

**Corollary 6.5.3.** *Singular Matrices Obstruct Splitting Convergence*
*Let $A = S - T$, with $S$ invertible. If $A$ is singular, then $1 \in \operatorname{spec}(S^{-1}T)$.*
*Hence $\rho(S^{-1}T) \geq 1$, so $\rho(S^{-1}T) < 1$ cannot hold.*

*Proof.* Choose $\mathbf{0} \neq \mathbf{x} \in \ker A$. Then $(S-T)\mathbf{x} = \mathbf{0}$, so $S\mathbf{x} = T\mathbf{x}$ and $S^{-1}T\mathbf{x} = \mathbf{x}$. Thus 1 is eigenvalue of $S^{-1}T$.
*Structural Commentary*: Singular $A$ forces an undamped error mode.
*Key Insight*: A null vector of $A$ becomes a fixed error mode. □

**Method 6.5.4.** *Stationary Iterative Method*
For $A\mathbf{x} = \mathbf{b}$, a stationary iteration is:

(i) Choose fixed $M$ and $\mathbf{c}$.
(ii) Update $\mathbf{x}_{k+1} = M\mathbf{x}_k + \mathbf{c}$.
(iii) If $\mathbf{x}_*$ exists, use $\mathbf{x}_{k+1} - \mathbf{x}_* = M(\mathbf{x}_k - \mathbf{x}_*)$ and Theorem 3.3.5 for convergence.

**Method 6.5.5.** *Jacobi, Gauss–Seidel, and SOR Splittings*
Let $A = D + L + U$, with $D$ diagonal, $L$ strictly lower triangular, and $U$ strictly upper triangular. For a splitting $A = S - T$, use $S\mathbf{x}_{k+1} = T\mathbf{x}_k + \mathbf{b}$ and $B = S^{-1}T$.

(i) *Jacobi*: $S = D$, $T = -(L+U)$, $B_J = -D^{-1}(L+U)$.
(ii) *Gauss–Seidel*: $S = D + L$, $T = -U$, $B_{\mathrm{GS}} = -(D+L)^{-1}U$.
(iii) *SOR*: for $\omega \neq 0$, $S_\omega = \frac{1}{\omega}D + L$; $T_\omega = \left(\frac{1}{\omega} - 1\right)D - U$; and $B_\omega = S_\omega^{-1}T_\omega$.

Convergence is controlled by $\rho(B)$; $\omega = 1$ gives Gauss–Seidel.

**Corollary 6.5.6.** *Strict Diagonal Dominance Gives Jacobi Convergence*
*If $A = [a_{ij}] \in \operatorname{Mat}(n, n, \mathbb{C})$ is strictly row diagonally dominant, then Jacobi converges for every right-hand side.*

*Proof.* Here $a_{ii} \neq 0$ and $B_J = -D^{-1}(L+U) = I_n - D^{-1}A$. Thus $(B_J)_{ii} = 0$ and $\sum_{j \neq i} |(B_J)_{ij}| = \sum_{j \neq i} |a_{ij}| / |a_{ii}| < 1$.
Gersgorin gives $\operatorname{spec}(B_J) \subset \{z : |z| < 1\}$, so $\rho(B_J) < 1$; Theorem 6.5.2 applies.
*Structural Commentary*: Diagonal dominance puts the Jacobi spectrum in the unit disk.
*Key Insight*: Small off-diagonal row sums force the error matrix to have spectral radius below 1. □

**Example 6.5.7.** *Two-by-Two Jacobi and Gauss–Seidel*
For $A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$, $\mathbf{b} \in \mathbb{R}^2$, write $A = D + L + U$.
The Jacobi iteration matrix is $B_J = -D^{-1}(L+U) = \begin{bmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & 0 \end{bmatrix}$, so $\rho(B_J) = \frac{1}{2}$.
The Gauss–Seidel iteration matrix is $B_{\mathrm{GS}} = -(D+L)^{-1}U = \begin{bmatrix} 0 & -\frac{1}{2} \\ 0 & \frac{1}{4} \end{bmatrix}$, so $\rho(B_{\mathrm{GS}}) = \frac{1}{4}$.
Both converge by Theorem 6.5.2, and smaller spectral radius predicts faster asymptotic decay for Gauss–Seidel.

**Example 6.5.8.** *Poisson Tridiagonal Stationary Rates*
For $A_n = \operatorname{tridiag}(-1, 2, -1)$, Jacobi gives $B_J = I_n - \frac{1}{2}A_n$; $\lambda_k(B_J) = \cos\frac{k\pi}{n+1}$; and $\rho(B_J) = \cos\frac{\pi}{n+1}$.
For the same ordering, Gauss–Seidel satisfies $\rho(B_{\mathrm{GS}}) = \rho(B_J)^2 = \cos^2\frac{\pi}{n+1}$. As $n \to \infty$, both radii tend to 1; fine grids need acceleration or preconditioning.

**Perspective 6.5.9.** *Preconditioning Preview*
A preconditioner replaces $A\mathbf{x} = \mathbf{b}$ by an equivalent system such as $M^{-1}A\mathbf{x} = M^{-1}\mathbf{b}$ or $AM^{-1}\mathbf{y} = \mathbf{b}$, with $\mathbf{x} = M^{-1}\mathbf{y}$. It should improve conditioning or convergence while being much cheaper to apply than $A^{-1}$.

## 6.6 Hessenberg Reduction and QR Eigenvalue Methods

**Definition 6.6.1.** *Upper Hessenberg Matrix*
Let $A = [a_{ij}] \in \operatorname{Mat}(n, n, \mathbb{F})$. Then $A$ is *upper Hessenberg* if $a_{ij} = 0$ whenever $i > j + 1$.
Equivalently, all entries below the first subdiagonal are zero.

Numerical linear algebra also replaces a matrix by a unitarily similar one with many forced zero entries. Upper Hessenberg form is the standard preliminary reduction before QR-type eigenvalue algorithms.



Figure 104: Upper Hessenberg pattern: red is the first subdiagonal; blue entries below it are forced zero.

**Theorem 6.6.2.** *Unitary Upper Hessenberg Reduction*
*Every $A \in \mathrm{Mat}(n, n, \mathbb{F})$ is unitarily similar to upper Hessenberg matrix with real nonnegative subdiagonal entries.*

*Proof.* Set $A_0 = A$. For $k = 1, \ldots, n-2$, suppose columns $1, \ldots, k-1$ of $A_{k-1}$ already have zeros below the first subdiagonal. Let $\mathbf{w}$ be the tail of column $k$ in rows $k+1, \ldots, n$.
By Corollary 2.7.4, choose a unitary $V_k$ with $V_k^* \mathbf{w} = \alpha \mathbf{e}_1$, $\alpha \geq 0$.
Put $U_k = I_k \oplus V_k$ and $A_k = U_k^* A_{k-1} U_k$. Right multiplication by $U_k$ changes only columns $k+1, \ldots, n$, so columns $1, \ldots, k-1$ keep their existing Hessenberg zeros. Left multiplication by $U_k^*$ acts only on rows $k+1, \ldots, n$; those rows are already zero in columns $1, \ldots, k-1$, so the zeros remain. In column $k$, the tail becomes $V_k^* \mathbf{w} = \alpha \mathbf{e}_1$, so entries below row $k+1$ are zero and the new subdiagonal entry is real nonnegative.
After $n-2$ steps, $A_{n-2} = U^* A U$ is upper Hessenberg, where $U = U_1 \cdots U_{n-2}$ is unitary.
*Structural Commentary*: Each trailing unitary creates one Hessenberg column and preserves earlier ones.
*Key Insight*: Kill one column tail at a time. □

**Perspective 6.6.3.** *Why Hessenberg Form Matters*
Hessenberg form preserves spectral data by unitary similarity, making QR-type eigenvalue algorithms cheaper.

**Method 6.6.4.** *QR Eigenvalue Algorithm Preview*
For a square matrix $A_0$, the basic QR eigenvalue iteration is:

(i) In practice, first reduce $A_0$ to Hessenberg form by Theorem 6.6.2.
(ii) Factor $A_k = Q_k R_k$, with $Q_k$ unitary and $R_k$ upper triangular.
(iii) Set $A_{k+1} = R_k Q_k = Q_k^* A_k Q_k$, using shifts to speed convergence.

Each step is a unitary similarity, so the spectrum is preserved.

**Method 6.6.5.** *Hessenberg QR Step by Plane Rotations*
Let $H \in \mathrm{Mat}(n, n, \mathbb{C})$ be upper Hessenberg and let $\mu \in \mathbb{C}$.

(i) Set $B_0 = H - \mu I_n$.
(ii) For $j = 1, \ldots, n-1$, choose a unitary plane rotation $G_j$ acting on coordinates $j, j+1$, so that $(G_j^* B_{j-1})_{j+1,j} = 0$ and $B_j = G_j^* B_{j-1}$.
(iii) Put $Q = G_1 \cdots G_{n-1}$ and $R = B_{n-1}$; then $H - \mu I_n = QR$.
(iv) Define the shifted QR update $H_+ = RQ + \mu I_n = Q^* H Q$

Adjacent rotations implement the Hessenberg step by local bulge chasing.

## 6.7 Power and Shift-Invert Iterations

**Theorem 6.7.1.** *Power Method Dominant-Eigenvector Preview*
*Let $A\mathbf{v}_j = \lambda_j \mathbf{v}_j$, where $\mathbf{v}_1, \ldots, \mathbf{v}_n$ is a basis of $\mathbb{C}^n$, and assume $|\lambda_1| > \max_{j \geq 2} |\lambda_j|$.*
*If $\mathbf{u}_0 = \sum_{j=1}^{n} c_j \mathbf{v}_j$ with $c_1 \neq 0$, then $\lambda_1^{-k} A^k \mathbf{u}_0 \longrightarrow c_1 \mathbf{v}_1$. Thus the directions of $A^k \mathbf{u}_0$ converge to $\mathrm{span}\{\mathbf{v}_1\}$.*

*Proof.* Set $\rho = \max_{j \geq 2} |\lambda_j / \lambda_1| < 1$. Since $A^k \mathbf{v}_j = \lambda_j^k \mathbf{v}_j$, $\lambda_1^{-k} A^k \mathbf{u}_0 = c_1 \mathbf{v}_1 + \sum_{j=2}^{n} c_j \left(\frac{\lambda_j}{\lambda_1}\right)^k \mathbf{v}_j$.

The remainder has norm at most $\rho^k \sum_{j=2}^{n} |c_j| \, \|\mathbf{v}_j\|$, hence tends to 0.
*Structural Commentary*: The power method is spectral filtering by repeated multiplication.
*Key Insight*: A spectral gap makes every non-dominant component decay relative to the dominant one. □

**Method 6.7.2.** *Inverse and Shifted Power Iteration Preview*
Let $\mathcal{N}(\mathbf{z}) := \mathbf{z} / \|\mathbf{z}\|_2$ for $\mathbf{z} \neq \mathbf{0}$.

(i) *Inverse*: if $0 \notin \mathrm{spec}(A)$, solve $A\mathbf{y}_{k+1} = \mathbf{x}_k$ and set $\mathbf{x}_{k+1} = \mathcal{N}(\mathbf{y}_{k+1})$.
This is power iteration on $A^{-1}$ and targets minimal $|\lambda|$.

(ii) *Shift-invert*: for $\sigma \notin \operatorname{spec}(A)$, $(A-\sigma I)\mathbf{y}_{k+1} = \mathbf{x}_k$, $\mathbf{x}_{k+1} = \mathcal{N}(\mathbf{y}_{k+1})$. Eigenvalues become $(\lambda-\sigma)^{-1}$, targeting $\lambda$ nearest $\sigma$.
(iii) *Shifted power*: $\mathbf{x}_{k+1} = \mathcal{N}\big((A-\sigma I)\mathbf{x}_k\big)$; eigenvalues become $\lambda-\sigma$, emphasizing large $|\lambda-\sigma|$.

Use factors of $A$ or $A-\sigma I$ for solves; do not form explicit inverses.

## 6.8 Numerical SVD Methods

**Method 6.8.1.** *SVD Numerics Preview*
Stable numerical SVD computation proceeds as:

(i) Avoid forming $A^*A$ when possible, since it can square the condition number.
(ii) Use orthogonal or unitary transformations to reduce $A$ to bidiagonal form.
(iii) Compute singular values from the bidiagonal problem, then use them for low-rank approximation, pseudoinverse truncation, and least-squares diagnostics through Theorem 5.2.2 and Definition 5.3.1.

## 6.9 Discrete Fourier Transforms and Circulant Matrices

**Definition 6.9.1.** *Discrete Fourier Matrix*
For $n \geq 1$, set $\omega_n = e^{2\pi i/n}$ and $F_n := [\omega_n^{jk}]_{j,k=0}^{n-1}$.
Then $F_n\mathbf{c}$ is the unnormalised DFT of $\mathbf{c} \in \mathbb{C}^n$; $Q_n := n^{-1/2}F_n$ is the normalized Fourier matrix.

**Definition 6.9.2.** *Circulant Matrix*
On the 0-indexed basis of $\mathbb{C}^n$, define the *cyclic shift* $P_n\mathbf{e}_j = \mathbf{e}_{(j-1) \bmod n}$.
A matrix $C \in \operatorname{Mat}(n,n,\mathbb{C})$ is *circulant* if, for some $a_0,\dots,a_{n-1} \in \mathbb{C}$,

$$C = a_0 I_n + a_1 P_n + \cdots + a_{n-1}P_n^{n-1} = p(P_n),$$
$$p(z) = a_0 + a_1 z + \cdots + a_{n-1}z^{n-1}$$

**Theorem 6.9.3.** *Fourier Basis for Cyclic Shifts and Circulants*
*Let $n \geq 1$, let $\omega_n := e^{2\pi i/n}$, and let $P_n$ be the cyclic shift.*
*For $0 \leq k < n$, set $\mathbf{f}_k := (\omega_n^{jk})_{j=0}^{n-1}$; $F_n := [\mathbf{f}_0 \ \cdots \ \mathbf{f}_{n-1}]$; and $\Omega_n := \operatorname{diag}(1,\omega_n,\dots,\omega_n^{n-1})$.*
*Then $F_n^*F_n = nI_n$, so $Q_n := n^{-1/2}F_n$ is unitary, and $P_nF_n = F_n\Omega_n$.*
*If $C = p(P_n)$ is circulant, then $Q_n^*CQ_n = p(\Omega_n) = \operatorname{diag}\big(p(1),p(\omega_n),\dots,p(\omega_n^{n-1})\big)$.*

*Proof.* For $0 \leq k,\ell \leq n-1$, $\langle \mathbf{f}_k, \mathbf{f}_\ell\rangle = \sum_{r=0}^{n-1}(\omega_n^{k-\ell})^r = n\delta_{k\ell}$. For $k=\ell$ the sum is $n$; for $k \neq \ell$, $W = \omega_n^{k-\ell}$ has $W \neq 1$ and $W^n = 1$, so $(1-W)\sum_{r=0}^{n-1} W^r = 0$. Hence $F_n^*F_n = nI_n$, and Theorem 2.6.8 makes $Q_n$ unitary.
Also $P_n\mathbf{f}_k = \omega_n^k\mathbf{f}_k$, so $Q_n^*P_nQ_n = \Omega_n$; applying $p$ gives the formula.
*Structural Commentary*: Fourier coordinates diagonalize every circulant.
*Key Insight*: Orthogonality of roots of unity turns a cyclic matrix problem into scalar evaluation. □




Figure 105: Fourier diagonalizes $P_n$; polynomial evaluation diagonalizes $C = p(P_n)$.

**Proposition 6.9.4.** *Inverse of the Fourier Matrix*
*The Fourier matrix satisfies $F_n^*F_n = nI_n$, and therefore $F_n^{-1} = \frac{1}{n}F_n^*$.*

*Proof.* Theorem 6.9.3 gives $Q_n^*Q_n = I_n$ for $Q_n = n^{-1/2}F_n$. Multiplying by $n$ gives $F_n^*F_n = nI_n$, so $n^{-1}F_n^*$ is a left inverse of $F_n$. Since $F_n$ is square, Theorem 3.4.2 makes it the inverse.
*Structural Commentary*: Fourier inversion is conjugate transpose plus one scalar normalization.
*Key Insight*: The inverse is simple because the Fourier columns are orthogonal. □

## 6.10 Fast Fourier Transform

**Method 6.10.1.** *Radix-Two FFT Split*
Let $n = 2m$ and $\mathbf{y} = F_n\mathbf{c}$.

(i) Split $\mathbf{c}' := (c_{2r})_{r=0}^{m-1}$ and $\mathbf{c}'' := (c_{2r+1})_{r=0}^{m-1}$.
(ii) Compute $\mathbf{y}' := F_m\mathbf{c}'$ and $\mathbf{y}'' := F_m\mathbf{c}''$.
(iii) With $d_j := \omega_n^j$, and using $\omega_n^2 = \omega_m$ and $\omega_n^m = -1$, set $y_j = y'_j + d_j y''_j$; and $y_{j+m} = y'_j - d_j y''_j$ for $0 \le j < m$.
(iv) For $n = 2^\ell$, repeat (i)–(iii) for $\ell$ stages.
$n^2$ dense products become about $\frac{1}{2} n \log_2 n$ twiddle multiplications plus additions.




Figure 106: For $n = 4$: split, apply two $F_2$ transforms, then $B_4\binom{\mathbf{y}'}{\mathbf{y}''} = \binom{\mathbf{y}'+D\mathbf{y}''}{\mathbf{y}'-D\mathbf{y}''}$ with $D = \mathrm{diag}(1, i)$.

# Part VII
# Groups, Rings, and Fields

## 1 Algebraic Language and Categories

### 1.1 Universal Properties and Equivalence-Based Structure

The elementary set, order, choice, function, and bijection vocabulary used here is collected in the foundational prerequisites. This subsection continues with algebraic applications of universal properties and equivalence-based structure.

### 1.2 Categories and Functors

### 1.3 Algebraic Structures as Objects

### 1.4 Exactness and Diagrams

## 2 Group Theory

### 2.1 Groups, Subgroups, and Cyclic Groups

Groups are the algebraic language of reversible operations. They appear here as the additive structure underlying rings and modules, and later as symmetry groups acting on mathematical objects.

**Definition 2.1.1.** *Groups*
A *group* is a nonempty set $G$ with a binary operation $(a, b) \mapsto ab$ such that:

(i) $(ab)c = a(bc)$ for all $a, b, c \in G$;
(ii) there is an element $e \in G$ with $eg = ge = g$ for all $g \in G$;
(iii) for each $g \in G$, there is $g^{-1} \in G$ with $gg^{-1} = g^{-1}g = e$.

The group is *abelian* if $ab = ba$ for all $a, b \in G$.
In an abelian group the operation is often written additively, with identity $0$ and inverse $-a$.

**Example 2.1.2.** *Basic Groups*

(i) The bijections $S \to S$ form a group under composition; it is usually nonabelian.
(ii) $\mathrm{Mat}(m, n, \mathbb{F})$ is an abelian group under matrix addition.
(iii) $\mathrm{GL}_n(\mathbb{F})$, the invertible $n \times n$ matrices, is a group under multiplication.
(iv) $\mathbb{Z}/n\mathbb{Z}$ is a finite abelian group under addition modulo $n$.

**Definition 2.1.3.** *Subgroup*
A *subgroup* of a group $G$ is a nonempty subset $H \subseteq G$ that is itself a group under operation inherited from $G$.

**Proposition 2.1.4.** *Subgroup Test*
*Let $G$ be a group and let $H \subseteq G$ be nonempty. Then $H \leq G \iff ab^{-1} \in H \ \forall a, b \in H$.*

*Proof.*
($\Rightarrow$) If $H \leq G$, then $b^{-1} \in H$ and $ab^{-1} \in H$ for all $a, b \in H$.
($\Leftarrow$) Assume $ab^{-1} \in H$ for all $a, b \in H$. Choose $h \in H$. Then $e_G = hh^{-1} \in H$. For $b \in H$, $b^{-1} = e_G b^{-1} \in H$. For $a, c \in H$, since $c^{-1} \in H$, we have $ac = a(c^{-1})^{-1} \in H$. Hence $H \leq G$.

*Structural Commentary*: Closure under one expression packages identity, inverses, and products.
*Key Insight*: In a group, subtraction-style closure is enough. □

**Definition 2.1.5.** *Generated and Cyclic Groups*
Let $G$ be a group.

(i) For $a \in G$, the *subgroup generated by $a$* is $\langle a \rangle := \{a^k : k \in \mathbb{Z}\} \leq G$.
(ii) $G$ is *cyclic* if $G = \langle a \rangle$ for some $a \in G$.
If it exists, $\mathrm{ord}(a) := \min\{n \in \mathbb{N}_{>0} : a^n = e_G\}$; then $\langle a \rangle = \{e_G, a, \ldots, a^{\mathrm{ord}(a)-1}\}$.

## 2.2 Group Homomorphisms, Kernels, and Images

**Definition 2.2.1.** *Group Homomorphisms, Kernels, and Images*
Let $G, H$ be groups.

(i) A *homomorphism* $\varphi : G \to H$ satisfies $\varphi(ab) = \varphi(a)\varphi(b)$ for all $a, b \in G$.
(ii) For a homomorphism $\varphi : G \to H$, $\ker \varphi := \{g \in G : \varphi(g) = e_H\}$; and $\operatorname{im} \varphi := \varphi(G)$.
An *isomorphism* is a bijective homomorphism.

**Proposition 2.2.2.** *Basic Homomorphism Properties*
*Let $\varphi : G \to H$ be a group homomorphism. Then:*

*(i)* $\varphi(e_G) = e_H$;
*(ii)* $\varphi(g^{-1}) = \varphi(g)^{-1}$ *for all* $g \in G$;
*(iii)* $\ker \varphi \leq G$;
*(iv)* $\operatorname{im} \varphi \leq H$.

*Proof.*

(i) Since $\varphi(e_G) = \varphi(e_G e_G) = \varphi(e_G)^2$, cancellation gives $\varphi(e_G) = e_H$.
(ii) For $g \in G$, $\varphi(g)\varphi(g^{-1}) = \varphi(gg^{-1}) = e_H$, hence $\varphi(g^{-1}) = \varphi(g)^{-1}$.
(iii) If $a, b \in \ker \varphi$, then $\varphi(ab^{-1}) = \varphi(a)\varphi(b)^{-1} = e_H$, so $\ker \varphi \leq G$ by Proposition 2.1.4.
(iv) If $x = \varphi(a)$ and $y = \varphi(b)$ lie in $\operatorname{im} \varphi$, then $xy^{-1} = \varphi(ab^{-1}) \in \operatorname{im} \varphi$, so $\operatorname{im} \varphi \leq H$ by Proposition 2.1.4.

*Structural Commentary*: Homomorphisms preserve all group structure once they preserve multiplication.
*Key Insight*: $\varphi(ab) = \varphi(a)\varphi(b)$ preserves $e$, inverses, kernels, and images. □

**Proposition 2.2.3.** *Kernel Criterion for Injectivity*
*Let $\varphi : G \to H$ be a group homomorphism. Then $\varphi$ injective $\iff$ $\ker \varphi = \{e_G\}$.*

*Proof.*

($\Rightarrow$) If $\varphi$ injective, $g \in \ker \varphi$, then $\varphi(g) = e_H = \varphi(e_G)$ by Proposition 2.2.2, so $g = e_G$. Hence $\ker \varphi = \{e_G\}$.
($\Leftarrow$) If $\ker \varphi = \{e_G\}$ and $\varphi(a) = \varphi(b)$, then $\varphi(ab^{-1}) = \varphi(a)\varphi(b)^{-1} = e_H$, so $ab^{-1} = e_G$ and $a = b$.

*Structural Commentary*: Injectivity is detected by the identity fiber $\ker \varphi$.
*Key Insight*: The kernel measures exactly the failure of injectivity. □

## 2.3 Cosets, Normality, and Structure Theorems

## 2.4 Group Actions, Orbits, and Stabilizers

A group action packages the idea that elements of a group act as reversible motions of a set. The conjugation action is the finite-group ingredient used later in Wedderburn's theorem.

**Definition 2.4.1.** *Actions, Orbits, and Stabilizers*
Let $G$ be a group and let $\mathcal{X}$ be a set.

(i) *Left action* of $G$ on $\mathcal{X}$ is map $G \times \mathcal{X} \to \mathcal{X}$, $(g, x) \mapsto gx$, s.t. $e_G x = x$ and $(gh)x = g(hx)$ $\forall g, h \in G$, $x \in \mathcal{X}$.
(ii) The *orbit* of $x \in \mathcal{X}$ is $Gx := \{gx : g \in G\}$.
(iii) The *stabilizer* of $x \in \mathcal{X}$ is $G_x := \{g \in G : gx = x\}$.

**Theorem 2.4.2.** *Orbit-Stabilizer*
*Let a finite group $G$ act on $\mathcal{X}$. For each $x \in \mathcal{X}$, $|Gx| = [G : G_x] = \frac{|G|}{|G_x|}$.*

*Proof.* Define $\Phi : G/G_x \to Gx$, $\Phi(gG_x) = gx$.
For $g, h \in G$, $gG_x = hG_x \iff h^{-1}g \in G_x \iff h^{-1}gx = x \iff gx = hx$. Thus $\Phi$ is well defined and injective. It is surjective by definition of $Gx$, hence bijective, so $|Gx| = |G/G_x| = [G : G_x]$.
*Structural Commentary*: Orbits are parametrized by cosets of the stabilizer.
*Key Insight*: Same orbit point $\iff$ same stabilizer coset. □

## 2.5 Conjugacy Classes and the Class Equation

**Definition 2.5.1.** *Conjugacy Classes and Centralizers*
Let $G$ be a group.

(i) The *conjugation action* of $G$ on itself is $g \cdot x := gxg^{-1}$.
(ii) The *conjugacy class* of $x \in G$ is $\{x\}^G := \{gxg^{-1} : g \in G\}$.
(iii) The *centralizer* of $x \in G$ is $C_G(x) := \{g \in G : gx = xg\}$.

(iv) The *center* of $G$ is $Z(G) := \{g \in G : gx = xg \text{ for all } x \in G\}$.

**Theorem 2.5.2.** *Class Equation*
*Let $G$ be finite, $S \subseteq G$ contain one representative of each noncentral conjugacy class. Then $|G| = |Z(G)| + \sum_{x \in S} [G : C_G(x)]$.*

*Proof.* The conjugacy classes form a partition $G = Z(G) \sqcup \bigsqcup_{x \in S} \{x\}^G$.
Indeed, $\left|\{x\}^G\right| = 1 \iff gxg^{-1} = x$ for all $g \in G \iff x \in Z(G)$. For $x \in S$, Theorem 2.4.2 applied to conjugation gives $\left|\{x\}^G\right| = [G : C_G(x)]$. Taking cardinalities yields the formula.
*Structural Commentary*: $G$ decomposes into central singleton classes and noncentral conjugacy classes.
*Key Insight*: Count conjugacy classes by orbit-stabilizer. □

## 2.6 Sylow Theory and Finite Groups

## 2.7 Representation-Theoretic Preview

# 3 Rings, Ideals, and Polynomial Rings

## 3.1 Rings, Identities, Fields, and Examples

Rings add multiplication to an abelian group.

**Definition 3.1.1.** *Rings, Identities, and Fields*
Let $R$ be a set with operations $+$ and $\cdot$.

(i) $R$ is a *ring* if $(R, +)$ is an abelian group, multiplication is associative, and $a(b+c) = ab + ac$, $(a+b)c = ac + bc$ for all $a, b, c \in R$.
(ii) A ring $R$ is *commutative* if $ab = ba$ for all $a, b \in R$.
(iii) A ring $R$ has *identity* if there is $1_R \in R$ such that $1_R a = a 1_R = a$ for all $a \in R$.
(iv) A *field* is a commutative ring with identity $1_R \neq 0$ such that every $a \in R \setminus \{0\}$ has $a^{-1} \in R$.

**Example 3.1.2.** *Basic Rings*

(i) $(\mathbb{Z}, +, \cdot)$ and $(\mathbb{F}[x], +, \cdot)$ are commutative rings with identity 1.
(ii) $\mathrm{Mat}(n, n, \mathbb{F})$ is a unital ring under matrix addition and multiplication, with identity $I_n$; it is noncommutative for $n \geq 2$.
(iii) $2\mathbb{Z}$ is a commutative ring under inherited addition and multiplication, but has no identity.
(iv) $\mathbb{Z}/n\mathbb{Z}$ is a commutative ring under addition and multiplication modulo $n$, with identity $1 + n\mathbb{Z}$.

## 3.2 Ring Homomorphisms and Subrings

**Definition 3.2.1.** *Ring Homomorphisms and Subrings*
Let $R, S$ be rings with identity.

(i) A *ring homomorphism* $\varphi : R \to S$ preserves $+$, $\cdot$, and 1, where $\varphi(a+b) = \varphi(a) + \varphi(b)$, $\varphi(ab) = \varphi(a)\varphi(b)$, $\varphi(1_R) = 1_S$ for all $a, b \in R$.
(ii) A *subring* of $R$ is a subset $T \subseteq R$ that is a ring under the inherited operations and has identity $1_R$.

**Proposition 3.2.2.** *Subring Test*
*Let $R$ be a ring with identity and let $S \subseteq R$. Then $S \leq R \iff 1_R \in S$, $a - b \in S$, $ab \in S$ $\forall a, b \in S$.*

*Proof.*
($\Rightarrow$) If $S \leq R$, then $1_R \in S$ and $S$ is closed under subtraction and multiplication.
($\Leftarrow$) Assume $1_R \in S$, $a - b \in S$, and $ab \in S$ for all $a, b \in S$. Then $S$ is nonempty, $(S, +) \leq (R, +)$ by Proposition 2.1.4, and multiplication restricts to $S$. All ring laws and the identity $1_R$ are inherited from $R$, so $S \leq R$.

*Structural Commentary*: A subring is a subset stable under the operations and containing the ambient unit.
*Key Insight*: Subtraction handles the additive group part in one stroke. □

## 3.3 Ideals and Generated Ideals

Ideals are the subsets by which quotient rings can be formed, just as subspaces are the subsets by which quotient vector spaces can be formed.

**Definition 3.3.1.** *Ideals and Generated Ideals*
Let $R$ be a ring.

(i) A nonempty subset $\mathcal{I} \subseteq R$ is a *two-sided ideal* if $a - b \in \mathcal{I}\ \forall a, b \in \mathcal{I}$, and $ra, ar \in \mathcal{I}\ \forall a \in \mathcal{I},\ r \in R$.

(ii) If $R$ is commutative with identity, $S \subseteq R$, *ideal generated by $S$* is $\langle S \rangle := \left\{ \sum_{j=1}^{n} r_j s_j : n \geq 1,\ r_j \in R,\ s_j \in S \right\}$.

(iii) *Finite generated ideal notation*: if $S = \{s_1, \ldots, s_n\}$, write $\langle s_1, \ldots, s_n \rangle := \langle S \rangle$.

**Example 3.3.2.** *Principal Ideals in Polynomial Rings*
In $\mathbb{F}[x]$, $\langle p(x) \rangle = \{q(x)p(x) : q(x) \in \mathbb{F}[x]\}$ is the *principal* ideal generated by $p(x)$.

**Proposition 3.3.3.** *Generated Ideals Are Smallest*
*Let $R$ be a commutative ring with identity and let $S \subseteq R$.*
*Then $\langle S \rangle$ is an ideal, $S \subseteq \langle S \rangle$, and for every ideal $\mathcal{J} \subseteq R$, $S \subseteq \mathcal{J} \implies \langle S \rangle \subseteq \mathcal{J}$.*

*Proof.* Let $x = \sum_{j=1}^{m} r_j s_j$ and $y = \sum_{k=1}^{n} t_k u_k$ lie in $\langle S \rangle$.
Then $x - y = \sum_{j=1}^{m} r_j s_j + \sum_{k=1}^{n} (-t_k) u_k \in \langle S \rangle$; $ax = \sum_{j=1}^{m} (ar_j) s_j \in \langle S \rangle$ for all $a \in R$, so $\langle S \rangle$ is an ideal.
Since $s = 1_R s \in \langle S \rangle$, $S \subseteq \langle S \rangle$. If $\mathcal{J}$ is an ideal with $S \subseteq \mathcal{J}$, then $r_j s_j \in \mathcal{J}$ and finite sums remain in $\mathcal{J}$, QED.
*Structural Commentary*: Generated ideals close a set under exactly the operations ideals must allow.
*Key Insight*: To force $S$ into an ideal, one must include all ring-linear combinations of elements of $S$. □

**Proposition 3.3.4.** *Intersections and Chain Unions of Ideals*
*Let $R$ be a ring.*

*(i) If $\{\mathcal{I}_\alpha\}_{\alpha \in A}$ is a nonempty family of ideals of $R$, then $\bigcap_{\alpha \in A} \mathcal{I}_\alpha$ is an ideal of $R$.*
*(ii) If $\{\mathcal{I}_\alpha\}_{\alpha \in A}$ is a chain of ideals of $R$ under inclusion, then $\bigcup_{\alpha \in A} \mathcal{I}_\alpha$ is an ideal of $R$.*

*Proof.*

(i) Let $\mathcal{I} = \bigcap_{\alpha \in A} \mathcal{I}_\alpha$. If $a, b \in \mathcal{I}$ and $r \in R$, then $a - b, ra, ar \in \mathcal{I}_\alpha$ for every $\alpha \in A$, hence $a - b, ra, ar \in \mathcal{I}$.
(ii) Let $\mathcal{U} = \bigcup_{\alpha \in A} \mathcal{I}_\alpha$. If $a, b \in \mathcal{U}$, then $a \in \mathcal{I}_\alpha$ and $b \in \mathcal{I}_\beta$ for some $\alpha, \beta$. Since the family is a chain, one of $\mathcal{I}_\alpha, \mathcal{I}_\beta$ contains both $a, b$; hence $a - b \in \mathcal{U}$. If $a \in \mathcal{U}$ and $r \in R$, then $a \in \mathcal{I}_\alpha$ for some $\alpha$, so $ra, ar \in \mathcal{I}_\alpha \subseteq \mathcal{U}$.

*Structural Commentary*: Chains avoid the usual failure of unions to preserve closure.
*Key Insight*: Comparability places any finite test inside one ideal. □

## 3.4 Quotient Rings and Maximal Ideals

**Definition 3.4.1.** *Quotient Rings*
Let $R$ be a ring and let $\mathcal{I} \trianglelefteq R$ be a two-sided ideal.

(i) The underlying set is the additive quotient group $R/\mathcal{I} = \{a + \mathcal{I} : a \in R\}$.
(ii) Addition and multiplication are $(a + \mathcal{I}) + (b + \mathcal{I}) = (a + b) + \mathcal{I}$ and $(a + \mathcal{I})(b + \mathcal{I}) = ab + \mathcal{I}$.
(iii) With these operations, $R/\mathcal{I}$ is the *quotient ring* of $R$ by $\mathcal{I}$.

**Proposition 3.4.2.** *Ideals Make Quotient Multiplication Well Defined*
*Let $\mathcal{I} \leq (R, +)$. The rule $(a + \mathcal{I})(b + \mathcal{I}) := ab + \mathcal{I}$ is well defined on $R/\mathcal{I}$ iff $\mathcal{I} \trianglelefteq R$.*

*Proof.*

($\Rightarrow$) If the rule is well defined and $i \in \mathcal{I}$, then $i + \mathcal{I} = 0 + \mathcal{I}$.
For $r \in R$, $ri + \mathcal{I} = r0 + \mathcal{I}$, and $ir + \mathcal{I} = 0r + \mathcal{I}$, so $ri, ir \in \mathcal{I}$. Thus $\mathcal{I} \trianglelefteq R$.
($\Leftarrow$) If $\mathcal{I} \trianglelefteq R$ and $a' = a + i$, $b' = b + j$ with $i, j \in \mathcal{I}$, then $a'b' - ab = aj + ib + ij \in \mathcal{I}$.
Hence $a'b' + \mathcal{I} = ab + \mathcal{I}$, so the product is independent of representatives.

*Structural Commentary*: Ideals are exactly the additive subgroups compatible with multiplication of cosets.
*Key Insight*: Coset multiplication is well defined exactly for ideals. □

**Definition 3.4.3.** *Maximal Ideals*
Let $R$ be a ring and let $\mathcal{M} \trianglelefteq R$.

(i) $\mathcal{M}$ is *proper* if $\mathcal{M} \neq R$.
(ii) $\mathcal{M}$ is *maximal* if $\mathcal{M} \neq R$ and, for every ideal $\mathcal{J} \trianglelefteq R$, $\mathcal{M} \subseteq \mathcal{J} \subseteq R \implies \mathcal{J} = \mathcal{M}$ or $\mathcal{J} = R$.

**Theorem 3.4.4.** *Maximal Ideals and Fields*
*Let $R$ be a commutative ring with identity and let $\mathcal{I} \trianglelefteq R$. Then $R/\mathcal{I}$ is a field $\iff$ $\mathcal{I}$ is maximal*

*Proof.*
($\Rightarrow$) If $R/\mathcal{I}$ is a field and $\mathcal{I} \subsetneq \mathcal{J} \trianglelefteq R$, then $\mathcal{J}/\mathcal{I}$ is a nonzero ideal of $R/\mathcal{I}$.
Hence $\mathcal{J}/\mathcal{I} = R/\mathcal{I}$, so $\mathcal{J} = R$. Thus $\mathcal{I}$ is maximal.
($\Leftarrow$) If $\mathcal{I}$ is maximal and $0 \neq a + \mathcal{I} \in R/\mathcal{I}$, then $a \notin \mathcal{I}$.
The ideal $\mathcal{I} + \langle a \rangle$ strictly contains $\mathcal{I}$, so maximality gives $\mathcal{I} + \langle a \rangle = R$. Thus $1 = i + ra$ for some $i \in \mathcal{I}$, $r \in R$, and $(r + \mathcal{I})(a + \mathcal{I}) = 1 + \mathcal{I}$. Hence every nonzero coset is invertible.
*Structural Commentary*: Maximal ideals are exactly kernels whose quotients have no residual ideal structure.
*Key Insight*: Maximality makes every nonzero coset invertible. □

**Theorem 3.4.5.** *Existence of Maximal Ideals*
*If $R$ is a commutative ring with identity and $R \neq 0$, then $R$ has a maximal ideal.*

*Proof.* Let $\mathcal{P} := \{\mathcal{I} \trianglelefteq R : \mathcal{I} \neq R\}$, ordered by inclusion. Since $R \neq 0$, $\{0\} \in \mathcal{P}$.
If $\mathcal{C} \subseteq \mathcal{P}$ is a chain, then $\mathcal{U} := \bigcup_{\mathcal{I} \in \mathcal{C}} \mathcal{I}$ is an ideal by Proposition 3.3.4.
Also $\mathcal{U} \neq R$: if $1_R \in \mathcal{U}$, then $1_R \in \mathcal{I}$ for some $\mathcal{I} \in \mathcal{C}$, forcing $\mathcal{I} = R$, contradiction. Thus $\mathcal{U} \in \mathcal{P}$ is an upper bound for $\mathcal{C}$. By Zorn's lemma, Principle 6.5.3, $\mathcal{P}$ has a maximal element, i.e. $R$ has a maximal ideal.
*Structural Commentary*: Maximal ideals are produced by maximality, not by an explicit formula in general.
*Key Insight*: Chain unions keep properness because containing 1 is a finite obstruction. □

## 3.5 Integral Domains and Fields of Quotients

**Definition 3.5.1.** *Zero Divisors and Integral Domains*
Let $R$ be a ring.

(i) A *zero divisor* is a nonzero $a \in R$ such that $\exists\, 0 \neq b \in R$ with $ab = 0$ or $ba = 0$.
(ii) An *integral domain* is a commutative ring with identity $1_R \neq 0$ and no zero divisors.

**Proposition 3.5.2.** *Cancellation in Integral Domains*
*Let $R$ be a commutative ring with identity $1_R \neq 0$.*
*Then $R$ is an integral domain $\iff$ $(\forall r, x, y \in R)(r \neq 0 \wedge rx = ry \Rightarrow x = y)$.*

*Proof.*
($\Rightarrow$) If $rx = ry$ and $r \neq 0$, then $r(x - y) = 0$; no zero divisors gives $x - y = 0$.
($\Leftarrow$) If $ab = 0$ and $a \neq 0$, then $ab = a0$; cancellation gives $b = 0$, so no zero divisors exist.
*Structural Commentary*: Domains are exactly cancellative commutative rings.
*Key Insight*: Zero divisors obstruct cancellation. □

**Definition 3.5.3.** *Field of Quotients*
Let $R$ be an integral domain. On $R \times (R \setminus \{0\})$, set $(a, b) \sim (c, d) \iff ad = bc$.
The *field of quotients* of $R$ is $(R \times (R \setminus \{0\}))/\sim$; write $[(a, b)] = a/b$. For $b, d \neq 0$, $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd}$; $\frac{a}{b}\frac{c}{d} = \frac{ac}{bd}$.

**Proposition 3.5.4.** *Domains Embed in Fields*
*Let $R$ be an integral domain with field of quotients $K$.*
*Then $\iota : R \hookrightarrow K$; $\iota(a) = a/1$ is an injective ring homomorphism.*

*Proof.* By Proposition 3.5.2, the relation and operations in Definition 3.5.3 are well defined.
For $a, b \in R$, $\iota(a + b) = \frac{a+b}{1} = \frac{a}{1} + \frac{b}{1}$; $\iota(ab) = \frac{ab}{1} = \frac{a}{1}\frac{b}{1}$; and $\iota(1_R) = \frac{1_R}{1}$..
Thus $\iota$ is a ring homomorphism. If $\iota(a) = \iota(b)$, then $a/1 = b/1$, so $a = b$; hence $\iota$ is injective.
*Structural Commentary*: The field of quotients formally adds denominators to a domain.
*Key Insight*: Integral domains are exactly the setting where fraction arithmetic is consistent. □

## 3.6 Ring Characteristic

**Definition 3.6.1.** *Characteristic*
Let $R$ be a ring with identity. Its *characteristic* is $\operatorname{char}(R) := \begin{cases} \min\{n \in \mathbb{N}_{>0} : n1_R = 0_R\}, & \text{if this set is nonempty,} \\ 0, & \text{otherwise.} \end{cases}$.

**Proposition 3.6.2.** *Finite Domains Have Prime Characteristic*
*Let $R$ be a finite ring with identity. Then $\operatorname{char}(R) \neq 0$. If $R$ is an integral domain, then $\operatorname{char}(R)$ is prime.*

*Proof.* Since $R$ is finite, the map $n \mapsto n1_R$ from $\mathbb{N}$ to $R$ is not injective.
Thus $i1_R = j1_R$ for some $i < j$, so $(j-i)1_R = 0_R$ and $\operatorname{char}(R) > 0$.
Assume $R$ is a domain and set $c = \operatorname{char}(R)$. Since $1_R \neq 0_R$, $c > 1$.
If $c = ab$ with $1 < a, b < c$, then $(a1_R)(b1_R) = ab1_R = c1_R = 0_R$.
By minimality of $c$, $a1_R \neq 0_R$ and $b1_R \neq 0_R$, contradicting that $R$ has no zero divisors. Hence $c$ is prime.
*Structural Commentary*: Characteristic records the additive order of the unit.
*Key Insight*: In a domain, a composite characteristic would factor zero nontrivially. □

## 3.7 Algebras over Fields

**Definition 3.7.1.** *Algebras over a Field*
Let $\mathbb{F}$ be a field.
An *$\mathbb{F}$-algebra* is an $\mathbb{F}$-vector space $\mathcal{A}$ with a ring product satisfying $\lambda(ab) = (\lambda a)b = a(\lambda b)$ $\forall \lambda \in \mathbb{F}$ and $a, b \in \mathcal{A}$.

**Example 3.7.2.** *Matrix and Polynomial Algebras*
$\operatorname{Mat}(n, n, \mathbb{F})$ and $\mathbb{F}[x]$ are algebras over $\mathbb{F}$. The first is generally noncommutative; the second is commutative.

## 3.8 Units, Divisibility, and Factorization Domains

**Definition 3.8.1.** *Units, Associates, Prime and Irreducible Elements*
Let $R$ be an integral domain.

(i) A *unit* is an element $u \in R$ with $uv = 1$ for some $v \in R$.
(ii) Nonzero elements $a, b \in R$ are *associates*, written $a \sim b$, if $a = ub$ for some unit $u$.
(iii) A nonzero nonunit $p \in R$ is *prime* if $p \mid ab$ implies $p \mid a$ or $p \mid b$.
(iv) A nonzero nonunit $r \in R$ is *irreducible* if $r = ab$ implies $a$ or $b$ is a unit.

**Proposition 3.8.2.** *Divisibility and Principal Ideals*
*Let $R$ be an integral domain and let $r, s \in R$ be nonzero.*

*(i)* $r \in R^\times \iff \langle r \rangle = R$.
*(ii)* $r \sim s \iff \langle r \rangle = \langle s \rangle$.
*(iii)* $r \mid s \iff \langle s \rangle \subseteq \langle r \rangle$.

*Proof.*

*(i)* ($\Rightarrow$) If $rv = 1$, then $1 \in \langle r \rangle$, so $\langle r \rangle = R$.
*(i)* ($\Leftarrow$) If $\langle r \rangle = R$, then $1 = ar$ for some $a \in R$, hence $r \in R^\times$.
*(ii)* ($\Rightarrow$) If $r = us$ with $u \in R^\times$, then $r \in \langle s \rangle$ and $s = u^{-1}r \in \langle r \rangle$, so $\langle r \rangle = \langle s \rangle$.
*(ii)* ($\Leftarrow$) If $\langle r \rangle = \langle s \rangle$, then $r = as$ and $s = br$ for some $a, b \in R$.
Thus $r = abr$, so $ab = 1$ by cancellation; hence $a \in R^\times$ and $r \sim s$.
*(iii)* ($\Rightarrow$) If $s = ar$, then every element of $\langle s \rangle$ is of the form $xs = xar \in \langle r \rangle$.
*(iii)* ($\Leftarrow$) If $\langle s \rangle \subseteq \langle r \rangle$, then $s \in \langle r \rangle$, so $s = ar$ for some $a \in R$; hence $r \mid s$.

*Structural Commentary*: Divisibility reverses inclusion of principal ideals.
*Key Insight*: Principal ideals translate multiplicative divisibility into order by containment. □

**Definition 3.8.3.** *PIDs and UFDs*
Let $R$ be an integral domain.

(i) $R$ is a *principal ideal domain* if every ideal $\mathcal{I} \trianglelefteq R$ has $\mathcal{I} = \langle a \rangle$ for some $a \in R$.
(ii) $R$ is a *unique factorization domain* if every $0 \neq a \in R \setminus R^\times$ factors into irreducibles, uniquely up to reordering and associates.

## 3.9 Polynomial Rings, Division, and GCDs

Polynomial rings over fields behave like the integers in the ways linear algebra needs: division with remainder, gcds, Bezout identities, irreducible factorization, and principal ideals.

**Definition 3.9.1.** *Polynomials over a Field*
Let $\mathbb{F}$ be a field.

(i) $\mathbb{F}[x]$ denotes the ring of polynomials in one variable over $\mathbb{F}$.
(ii) If $0 \neq p \in \mathbb{F}[x]$ and $p(x) = a_0 + a_1 x + \cdots + a_n x^n$, $a_n \neq 0$, then $\operatorname{lc}(p) := a_n$, $\deg p := n$.
Also, $p$ is *monic* iff $\operatorname{lc}(p) = 1$. Set $\deg 0 = -\infty$.
(iii) A nonzero $p \in \mathbb{F}[x]$ *splits over* $\mathbb{F}$ iff $p = \alpha \prod_{i=1}^{m} (x - \lambda_i)$ for some $m \geq 0$, $\alpha \in \mathbb{F}^\times$, and $\lambda_i \in \mathbb{F}$.

**Theorem 3.9.2.** *Polynomial Division Algorithm*
*Let $f, g \in \mathbb{F}[x]$ with $g \neq 0$. Then $\exists!\,(q, r) \in \mathbb{F}[x]^2$ such that $f = qg + r$, $r = 0$ or $\deg r < \deg g$.*

*Proof.* Let $d = \deg g$ and $\ell = \mathrm{lc}(g)$. Existence follows by induction on $\deg f$. If $f = 0$ or $\deg f < d$, take $(q, r) = (0, f)$. Otherwise, let $n = \deg f$, $a = \mathrm{lc}(f)$, and $f_1 = f - a\ell^{-1}x^{n-d}g$. Then $\deg f_1 < n$.
By induction, $f_1 = q_1 g + r$ with $r = 0$ or $\deg r < d$, hence $f = (q_1 + a\ell^{-1}x^{n-d})g + r$.
For uniqueness, if $qg + r = q'g + r'$ with admissible remainders, then $(q - q')g = r' - r$.
If $q \neq q'$, then $\deg((q - q')g) \geq d$, while $\deg(r' - r) < d$, contradiction. Hence $q = q'$ and then $r = r'$.
*Structural Commentary*: Division works because the leading coefficient of $g$ is invertible in the field.
*Key Insight*: Cancel leading terms until the remainder is too small to divide again. □

**Definition 3.9.3.** *Divisibility, GCDs, and Coprimality*
For $p, q \in \mathbb{F}[x]$, write $p \mid q$ if $q = hp$ for some $h \in \mathbb{F}[x]$.
A *greatest common divisor* of $f, g \in \mathbb{F}[x]$, not both zero, is the unique monic polynomial $\gcd(f, g)$ such that:

(i) $\gcd(f, g) \mid f$ and $\gcd(f, g) \mid g$;
(ii) every common divisor of $f$ and $g$ divides $\gcd(f, g)$.

The polynomials $f, g$ are *relatively prime* if $\gcd(f, g) = 1$.

**Theorem 3.9.4.** *Polynomial GCD and Bezout Identity*
*For $f, g \in \mathbb{F}[x]$, not both zero, $\exists$ a unique monic $d \in \mathbb{F}[x]$ s.t. $d \mid f$, $d \mid g$, $h \mid f$, $h \mid g \implies h \mid d$*
*Moreover, $d = \gcd(f, g)$ satisfies $d = af + bg$ for some $a, b \in \mathbb{F}[x]$.*
*Hence $\gcd(f, g) = 1$ iff $\exists\, a, b \in \mathbb{F}[x]$ with $af + bg = 1$.*

*Proof.* If one of $f, g$ is 0, the claim is immediate from the monic associate of the other.
Assume $f, g \neq 0$ and run the Euclidean algorithm: $r_{i-1} = q_i r_i + r_{i+1}$, where $r_{i+1} < \deg r_i$, starting with $(r_0, r_1) = (f, g)$ and ending with $r_N \neq 0$, $r_{N+1} = 0$. Then $r_N \mid r_{N-1}$, and upward substitution gives $r_N \mid f, g$.
Conversely, any common divisor of $f, g$ divides every $r_i$, hence divides $r_N$. Thus $d := \mathrm{lc}(r_N)^{-1} r_N$ is a monic gcd; uniqueness follows since two monic gcds divide each other.
Back-substitution gives $d = af + bg$ for some $a, b \in \mathbb{F}[x]$. Finally, $d = 1$ gives $af + bg = 1$; conversely, if $af + bg = 1$, every common divisor divides 1, so $d = 1$.
*Structural Commentary*: The Euclidean algorithm produces both the gcd and its Bezout certificate.
*Key Insight*: GCD data is ideal data in $\mathbb{F}[x]$. □

## 3.10 Irreducible Polynomials and Unique Factorization

**Definition 3.10.1.** *Irreducible Polynomials*
A nonconstant polynomial $p \in \mathbb{F}[x]$ is *irreducible over $\mathbb{F}$* if every factorization $p = ab$ in $\mathbb{F}[x]$ has $a$ or $b$ constant.

**Proposition 3.10.2.** *Irreducible Polynomials Are Prime in $\mathbb{F}[x]$*
*Let $p \in \mathbb{F}[x]$ be nonconstant. Then $p$ is irreducible over $\mathbb{F}$ iff $\forall f, g \in \mathbb{F}[x]$, $p \mid fg \implies p \mid f$ or $p \mid g$.*

*Proof.*

(⇒) Assume $p$ irreducible and $p \mid fg$. If $p \mid f$, done. Otherwise $\gcd(p, f) = 1$, so Theorem 3.9.4 gives $a, b \in \mathbb{F}[x]$ with $ap + bf = 1$. Multiplying by $g$ gives $apg + bfg = g$; since $p \mid apg$ and $p \mid bfg$, we get $p \mid g$.
(⇐) Assume the divisibility property and write $p = ab$. Since $p \mid ab$, either $p \mid a$ or $p \mid b$. If $p \mid a$, then $a = ph$, so $p = phb$ and cancellation gives $1 = hb$, hence $b \in \mathbb{F}^\times$. Similarly, $p \mid b$ forces $a \in \mathbb{F}^\times$. Thus $p$ is irreducible.

*Structural Commentary*: Over $\mathbb{F}[x]$, irreducible and prime are the same notion.
*Key Insight*: Bezout turns coprimality into a divisibility test. □

**Theorem 3.10.3.** *Unique Factorization in $\mathbb{F}[x]$*

*For every nonconstant $f \in \mathbb{F}[x]$, $\exists\, c \in \mathbb{F}^\times$ and monic irreducibles $p_1, \ldots, p_m \in \mathbb{F}[x]$ such that $f = c \prod_{i=1}^{m} p_i$.*

*If $f = d \prod_{j=1}^{n} q_j$ is another such factorization, then $m = n$, $c = d$, and $p_i = q_{\sigma(i)}$ for some $\sigma \in S_m$.*

*Proof.*

(i) *Existence*: Induct on $\deg f$. If $f$ is irreducible, write $f = \mathrm{lc}(f)\big(\mathrm{lc}(f)^{-1} f\big)$.
If $f = uv$ with $0 < \deg u, \deg v < \deg f$, apply induction to $u$ and $v$ and concatenate the factors.
(ii) *Uniqueness*: Suppose $c \prod_{i=1}^{m} p_i = d \prod_{j=1}^{n} q_j$ with all $p_i, q_j$ monic irreducible. Since $p_1 \mid \prod_{j=1}^{n} q_j$, iterating Proposition 3.10.2 gives $p_1 \mid q_j$ for some $j$. As $p_1, q_j$ are monic irreducible, $p_1 = q_j$.
Cancel and continue by induction on $m$; the final scalar equality follows from leading coefficients.

*Structural Commentary*: Polynomial factorization over a field mirrors prime factorization in $\mathbb{Z}$.
*Key Insight*: The prime property of irreducibles drives uniqueness. □

## 3.11 Principal Ideal Domains and Unique Factorization

**Theorem 3.11.1.** *$\mathbb{Z}$ and $\mathbb{F}[x]$ Are Principal Ideal Domains*

*(i) If $0 \neq \mathcal{I} \trianglelefteq \mathbb{Z}$, then $\mathcal{I} = \langle d \rangle$ for $d = \min(\mathcal{I} \cap \mathbb{N}_{>0})$.*
*(ii) If $0 \neq \mathcal{I} \trianglelefteq \mathbb{F}[x]$, then $\mathcal{I} = \langle m \rangle$ for the unique monic $m \in \mathcal{I}$ of minimal degree.*

*Proof.*

(i) If $\mathcal{I} = 0$, then $\mathcal{I} = \langle 0 \rangle$. Otherwise choose $d = \min(\mathcal{I} \cap \mathbb{N}_{>0})$. For $a \in \mathcal{I}$, write $a = qd + r$ with $0 \leq r < d$. Since $r = a - qd \in \mathcal{I}$, minimality gives $r = 0$, so $\mathcal{I} = \langle d \rangle$.
(ii) If $\mathcal{I} = 0$, then $\mathcal{I} = \langle 0 \rangle$. Otherwise choose monic $m \in \mathcal{I}$ of minimal degree. For $p \in \mathcal{I}$, divide $p = qm + r$ with $r = 0$ or $\deg r < \deg m$. Since $r = p - qm \in \mathcal{I}$, minimality gives $r = 0$, so $\mathcal{I} = \langle m \rangle$.
If $m'$ is another monic minimal generator, then $m \mid m'$ and $m' \mid m$, hence $m = m'$.

*Structural Commentary*: Euclidean division turns arbitrary ideals into one-generator ideals.
*Key Insight*: Choose the smallest positive integer or smallest-degree monic polynomial and divide by it. □

**Corollary 3.11.2.** *GCDs Generate Polynomial Ideals*
*For $0 \neq p_1, \ldots, p_n \in \mathbb{F}[x]$ and monic $m = \gcd(p_1, \ldots, p_n)$, $\langle p_1, \ldots, p_n \rangle = \langle m \rangle$.*

*Proof.* By Theorem 3.11.1, $\langle p_1, \ldots, p_n \rangle = \langle g \rangle$ for a unique monic $g$. Since $p_i \in \langle g \rangle$, $g \mid p_i$ for all $i$.
If $q \mid p_i$ for all $i$, then $\langle p_1, \ldots, p_n \rangle \subseteq \langle q \rangle$, so $g \in \langle q \rangle$ and $q \mid g$. Hence $g = m$, and the claim follows.
*Structural Commentary*: In $\mathbb{F}[x]$, gcds and finitely generated ideals are the same data.
*Key Insight*: The gcd is the monic generator of the ideal generated by the polynomials. □

**Example 3.11.3.** *A Polynomial Ring That Is Not a PID*
In $\mathbb{F}[x, y]$, the ideal $\langle x, y \rangle$ is not principal. If $\langle x, y \rangle = \langle p \rangle$, then $p \mid x$ and $p \mid y$, so $p \in \mathbb{F}^\times$. Hence $\langle x, y \rangle = \mathbb{F}[x, y]$, contradicting $1 \notin \langle x, y \rangle$ since every element of $\langle x, y \rangle$ has zero constant term.

**Proposition 3.11.4.** *PIDs Satisfy the Ascending Chain Condition*
*If $R$ is a PID and $\mathcal{I}_1 \subseteq \mathcal{I}_2 \subseteq \cdots$ is a chain of ideals, then $\exists N$ such that*

$$\mathcal{I}_j = \mathcal{I}_N \qquad \forall j \geq N.$$

*Proof.* Let $\mathcal{I} = \bigcup_{j \geq 1} \mathcal{I}_j$. By Proposition 3.3.4, $\mathcal{I}$ is an ideal. Since $R$ is a PID, $\mathcal{I} = \langle a \rangle$. Choose $N$ with $a \in \mathcal{I}_N$.
Then $\mathcal{I} = \langle a \rangle \subseteq \mathcal{I}_N \subseteq \mathcal{I}$, so $\mathcal{I}_j = \mathcal{I}_N$ for all $j \geq N$.
*Structural Commentary*: The generator appears at a finite stage.
*Key Insight*: Principal generation converts infinite ideal growth into a finite containment. □

**Theorem 3.11.5.** *Prime, Irreducible, and Bezout in a PID*
*Let $R$ be a principal ideal domain.*

*(i) For $0 \neq r \in R \setminus R^\times$, $r$ is irreducible iff $\langle r \rangle$ is maximal among proper ideals.*
*(ii) Every irreducible $r \in R$ is prime.*
*(iii) For $a, b \in R$, $\gcd(a, b) = 1$ iff $\exists x, y \in R$ such that $xa + yb = 1$.*

*Proof.*

(i) If $r$ is irreducible and $\langle r \rangle \subseteq \mathcal{I} \subseteq R$, write $\mathcal{I} = \langle a \rangle$. Then $r = xa$, so irreducibility gives $\mathcal{I} = \langle r \rangle$ or $\mathcal{I} = R$. Conversely, if $r = ab$ with $a, b \notin R^\times$, then $\langle r \rangle \subsetneq \langle a \rangle \subsetneq R$.
(ii) Let $r$ be irreducible and $r \mid ab$. By (i), $\langle r, a \rangle = \langle r \rangle$ or $R$. In the first case $r \mid a$.
In the second, $1 = xr + ya$, so $b = xrb + yab$ and $r \mid b$.
(iii) If $xa + yb = 1$, every common divisor of $a, b$ is a unit. Conversely, let $\langle a, b \rangle = \langle d \rangle$.
If $\gcd(a, b) = 1$, then $d \in R^\times$, so $\langle a, b \rangle = R$ and $1 = xa + yb$ for some $x, y \in R$.

*Structural Commentary*: In a PID, factorization statements become ideal-containment statements.
*Key Insight*: Maximal principal ideals are the bridge from irreducible to prime. □

**Theorem 3.11.6.** *Every PID Is a UFD*
*If $R$ is a PID, then $R$ is a UFD.*

*Proof.* Let $0 \neq r \in R \setminus R^\times$. If factorization into nonunits never terminates, obtain $s_1, s_2, \ldots$ with $s_{j+1} \mid s_j$ properly. By Proposition 3.8.2,

$$\langle s_1 \rangle \subsetneq \langle s_2 \rangle \subsetneq \cdots,$$

contradicting Proposition 3.11.4; hence irreducible factorizations exist.
For uniqueness, let $p_1 \cdots p_m = q_1 \cdots q_n$ with all $p_i, q_j$ irreducible. By Theorem 3.11.5, $p_1$ is prime, so $p_1 \mid q_j$ for some $j$; irreducibility gives $p_1 \sim q_j$. Cancel associates and iterate. Thus $m = n$ and the factors agree up to order and associates.

*Structural Commentary*: ACC gives existence; irreducible-is-prime gives uniqueness.
*Key Insight*: PIDs have enough ideal control to force unique factorization. □

## 3.12 Roots of Unity and Cyclotomic Polynomials

**Definition 3.12.1.** *Roots of Unity and Cyclotomic Polynomials*
For $n \geq 1$:

(i) The complex $n$th roots of unity are $U_n := \{z \in \mathbb{C} : z^n = 1\}$.
(ii) A root $\omega \in U_n$ is *primitive* iff $\operatorname{ord}(\omega) = n$; set $\Omega_n := \{\omega \in U_n : \operatorname{ord}(\omega) = n\}$.
(iii) The $n$th *cyclotomic polynomial* is $\Phi_n(x) := \prod_{\omega \in \Omega_n} (x - \omega)$.
(iv) The number of primitive $n$th roots is $\varphi(n) := |\Omega_n|$.

**Proposition 3.12.2.** *Roots of Unity Factorization*
*Let $\zeta_n = e^{2\pi i/n}$. Then $U_n = \langle \zeta_n \rangle$; $|U_n| = n$; $\Omega_n = \{\zeta_n^k : 1 \leq k \leq n,\ \gcd(k,n) = 1\}$; $|\Omega_n| = \varphi(n)$. Moreover, $U_n = \coprod_{d|n} \Omega_d$ and $x^n - 1 = \prod_{d|n} \Phi_d(x)$.*

*Proof.* The roots of $x^n - 1$ are precisely $\zeta_n^k$ for $0 \leq k < n$, hence $U_n = \langle \zeta_n \rangle$ and $|U_n| = n$.
In a cyclic group, $\operatorname{ord}(\zeta_n^k) = n/\gcd(k,n)$, so $\zeta_n^k$ is primitive iff $\gcd(k,n) = 1$; thus $|\Omega_n| = \varphi(n)$.
Each $z \in U_n$ has unique order $d \mid n$, giving $U_n = \coprod_{d|n} \Omega_d$.
Therefore $x^n - 1 = \prod_{z \in U_n} (x - z) = \prod_{d|n} \prod_{\omega \in \Omega_d} (x - \omega) = \prod_{d|n} \Phi_d(x)$ *Structural Commentary*: Orders partition roots of unity by divisors.
*Key Insight*: Cyclicity turns root counting into order counting. □

**Proposition 3.12.3.** *Cyclotomic Polynomials Have Integer Coefficients*
*For every $n \geq 1$, $\Phi_n(x) \in \mathbb{Z}[x]$ is monic.*

*Proof.* Induct on $n$. For $n = 1$, $\Phi_1(x) = x - 1$.
Assume $\Phi_d \in \mathbb{Z}[x]$ for every proper divisor $d \mid n$, and set $P_n(x) := \prod_{\substack{d|n \\ d<n}} \Phi_d(x) \in \mathbb{Z}[x]$.
Then $P_n$ is monic, and Proposition 3.12.2 gives $x^n - 1 = \Phi_n(x) P_n(x)$.
Divide $x^n - 1$ by $P_n$ in $\mathbb{Z}[x]$: $x^n - 1 = hP_n + r$, where $h, r \in \mathbb{Z}[x]$ and $r = 0$ or $\deg r < \deg P_n$.
Uniqueness of division over $\mathbb{C}[x]$ gives $h = \Phi_n$ and $r = 0$. Hence $\Phi_n \in \mathbb{Z}[x]$, and $\Phi_n$ is monic by definition.
*Structural Commentary*: Proper divisors isolate the next cyclotomic factor.
*Key Insight*: Monic division keeps the quotient in $\mathbb{Z}[x]$. □

**Lemma 3.12.4.** *A Cyclotomic Size Estimate*
*For $n > 1$ and $z \in \mathbb{Z}_{\geq 2}$, $|\Phi_n(z)| > z - 1$.*

*Proof.* For $\omega \in \Omega_n$, $\omega \neq 1$ and $|\omega| = 1$, hence $|z - \omega| > z - 1$. Thus $|\Phi_n(z)| = \prod_{\omega \in \Omega_n} |z - \omega| > (z-1)^{\varphi(n)} \geq z - 1$.
*Structural Commentary*: Primitive factors all lie outside the closest point 1.
*Key Insight*: Each primitive root contributes a factor larger than $z - 1$. □

# 4 Fields and Galois Theory

## 4.1 Field Extensions

## 4.2 Splitting Fields and Normality

## 4.3 Galois Groups

## 4.4 Finite Fields and Characteristic

**Perspective 4.4.1.** *Finite Fields and Characteristic*
If $\operatorname{char}(F) = 2$, then $(\forall a \in F)\ 2a = 0$, so $a = -a$. A finite field has $|F| = p^n$ with $p$ prime and $n \geq 1$; conversely, each prime power $q = p^n$ determines a unique field $\mathbb{F}_q$ up to isomorphism.
Thus the classification is field-theoretic, while the characteristic constraint is ring-theoretic.

## 4.5 Classical Applications

# Part VIII
# Modules, Multilinear Algebra, and Homological Methods

## 1 Module Theory

### 1.1 Modules and Submodules

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity 1.
Modules are vector spaces with the scalar field replaced by a ring; the price is that nonzero scalars may fail to be invertible and may even kill nonzero vectors.

**Perspective 1.1.1.** *Polynomial Scalars from One Operator*
Let $\mathcal{V}$ be a vector space over $\mathbb{F}$ and let $T \in \mathcal{L}(\mathcal{V})$. Every polynomial $p(z) \in \mathbb{F}[z]$ acts on $\mathcal{V}$ by $p \cdot \mathbf{v} := p(T)\mathbf{v}$. Thus one operator turns $\mathcal{V}$ into an object with scalar ring $\mathbb{F}[z]$.

**Definition 1.1.2.** *R-Module*
An *R-module* is an abelian group $(M, +)$ equipped with a scalar action $R \times M \to M$, $(r, u) \mapsto ru$, such that for all $r, s \in R$ and $u, v \in M$,

(i) $r(u + v) = ru + rv$.
(ii) $(r + s)u = ru + su$.
(iii) $(rs)u = r(su)$.
(iv) $1u = u$.

The ring $R$ is the *base ring* of $M$.

**Example 1.1.3.** *Basic Module Models*

(i) $R^n$ is an $R$-module under componentwise addition and scalar multiplication.
(ii) $\mathrm{Mat}(m, n, R)$ is an $R$-module under entrywise operations.
(iii) $R$ is an $R$-module over itself by ring multiplication.
(iv) Every vector space over a field $\mathbb{F}$ is an $\mathbb{F}$-module.

*Structural Commentary*: Modules are abelian groups with compatible scalar actions.
*Key Insight*: The new behavior comes from the scalar ring, not from changing addition.

**Perspective 1.1.4.** *The Base Ring Controls the Geometry*
The structure of an $R$-module is governed by the arithmetic of $R$. When $R$ is a field, every nonzero scalar is invertible, giving the usual theory of bases, dimension, and complements. For a general ring, nonunits and zero divisors obstruct division and cancellation, so these notions need not retain their vector-space properties.

**Definition 1.1.5.** *Submodule*
A *submodule* of an $R$-module $M$ is a nonempty subset $N \subseteq M$ that is an $R$-module under the operations inherited from $M$, written $N \leq M$.

**Theorem 1.1.6.** *Submodule Test*
*Let $M$ be an $R$-module and let $\emptyset \neq N \subseteq M$. Then $N \leq M \iff \forall r, s \in R,\ \forall u, v \in N,\ ru + sv \in N$.*

*Proof.*

($\Rightarrow$) If $N \leq M$, then $ru, sv \in N$, hence $ru + sv \in N$.
($\Leftarrow$) Assume the displayed condition. Choose $n \in N$. Then $0 = 0n + 0n \in N$.
For $x, y \in N$ and $r \in R$, $x + y = 1x + 1y \in N$, $-x = (-1)x + 0x \in N$, $rx = rx + 0x \in N$.
Thus $N$ is an additive subgroup of $M$ closed under $R$-scalar multiplication, so the remaining module axioms are inherited from $M$ and $N \leq M$.

*Structural Commentary*: One closure test packages addition, additive inverses, and scalar multiplication.
*Key Insight*: Submodules are exactly the subsets closed under all two-term $R$-linear combinations. □

**Proposition 1.1.7.** *Intersections and Sums of Submodules*
*Let $M$ be an $R$-module.*
*If $A, B \leq M$, then $A \cap B \leq M$; $A + B := \{a + b : a \in A,\ b \in B\} \leq M$, and $A + B = \bigcap_{\substack{N \leq M \\ A \cup B \subseteq N}} N$.*

*Proof.* Since $0 \in A \cap B$ and $rx+sy \in A \cap B$ for all $x, y \in A \cap B$ and $r, s \in R$, the submodule test gives $A \cap B \leq M$. If $x = a + b$ and $y = a' + b'$ with $a, a' \in A$ and $b, b' \in B$, then $rx + sy = (ra + sa') + (rb + sb') \in A + B$. Thus $A + B \leq M$. Moreover, $A \cup B \subseteq A + B$, and if $N \leq M$ contains $A \cup B$, then $a + b \in N$ for all $a \in A$, $b \in B$; hence $A + B \subseteq N$. This proves the displayed intersection formula.
*Structural Commentary*: $A \cap B$ is the meet; $A + B$ is the join.
*Key Insight*: Sums are forced by closure under addition. □

**Proposition 1.1.8.** *Ideals Are Submodules of the Regular Module*
*View $R$ as the regular $R$-module ${}_RR$. For every $N \subseteq R$,*

$$N \leq {}_RR \iff N \trianglelefteq R.$$

*Proof.* For $N \subseteq R$,

$$N \leq {}_RR \iff N \leq (R, +) \text{ and } RN \subseteq N \iff N \trianglelefteq R,$$

by Definition 3.3.1.
*Structural Commentary*: Ideals are the module-theoretic subobjects of the ring acting on itself.
*Key Insight*: Ring multiplication becomes scalar multiplication. □

## 1.2 Generated, Cyclic, and Finitely Generated Modules

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity and $M$ denotes an $R$-module.

**Definition 1.2.1.** *Generated Submodule*

(i) For $S \subseteq M$, *submodule generated by $S$* is $\langle\langle S \rangle\rangle_R := \left\{ \sum_{j=1}^{n} r_j s_j : n \geq 1,\ r_j \in R,\ s_j \in S \right\}$; $\langle\langle \emptyset \rangle\rangle_R = \{0\}$.
(ii) A subset $S \subseteq M$ *spans* or *generates* $M$ if $\langle\langle S \rangle\rangle_R = M$.

**Proposition 1.2.2.** *Generated Submodule Is Smallest*
*For every $S \subseteq M$, $\langle\langle S \rangle\rangle_R = \bigcap_{\substack{N \leq M \\ S \subseteq N}} N$.*

*Proof.* Let $G := \langle\langle S \rangle\rangle_R$. By construction, $S \subseteq G$ and the submodule test gives $G \leq M$.
If $N \leq M$ and $S \subseteq N$, then $\sum_{j=1}^{n} r_j s_j \in N$, where $r_j \in R,\ s_j \in S$, so $G \subseteq N$.
*Structural Commentary*: Generation means closing a set under the operations the module axioms require.
*Key Insight*: The generated submodule adds no elements except those forced by finite $R$-linear combination. □

**Definition 1.2.3.** *Cyclic and Finitely Generated Modules*

(i) For $v \in M$, the *cyclic submodule generated by $v$* is $\langle\langle v \rangle\rangle_R = Rv = \{rv : r \in R\}$.
(ii) $M$ is *finitely generated* if $M = \langle\langle v_1, \ldots, v_n \rangle\rangle_R$ for some $v_1, \ldots, v_n \in M$.
(iii) $M$ is *cyclic* if $M = Rv$ for some $v \in M$.

**Example 1.2.4.** *A Finitely Generated Module with a Non-Finitely Generated Submodule*
Let $R = \mathbb{F}[x_1, x_2, \ldots]$. Then $R = \langle\langle 1 \rangle\rangle_R$ as an $R$-module.
Set $I := (x_1, x_2, \ldots) = \{f \in R : f(0, 0, \ldots) = 0\}$. Then $I \trianglelefteq R$, hence $I \leq R$, but $I$ is not finitely generated.
If $I = \langle\langle p_1, \ldots, p_n \rangle\rangle_R$, choose $k$ such that $x_k$ occurs in no $p_i$. Let $\varphi_k : R \to \mathbb{F}[x_k]$ send $x_k \mapsto x_k$ and $x_j \mapsto 0$ for $j \neq k$. Since $\varphi_k(p_i) = 0$ for all $i$, every element of $\langle\langle p_1, \ldots, p_n \rangle\rangle_R$ maps to 0, but $\varphi_k(x_k) = x_k \neq 0$, contradiction.

## 1.3 Linear Independence in Modules

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity and $M$ denotes an $R$-module.

**Definition 1.3.1.** *Linear Independence in a Module*

(i) Subset $S \subseteq M$ is *linearly independent over $R$* if,
$\forall$ distinct $v_1, \ldots, v_n \in S$, $r_1 v_1 + \cdots + r_n v_n = 0 \implies r_1 = \cdots = r_n = 0$.
(ii) A subset $S \subseteq M$ is *linearly dependent* if it is not linearly independent.

**Example 1.3.2.** *Dependence Without Solving for One Vector*
In the $\mathbb{Z}$-module $\mathbb{Z}$, $3 \cdot 2 - 2 \cdot 3 = 0$. Thus $\{2, 3\}$ is linearly dependent.
However, $2 \notin 3\mathbb{Z}$; $3 \notin 2\mathbb{Z}$, so dependence need not express one element as a scalar multiple of another.

**Proposition 1.3.3.** *Scaling Independent Sets over a Domain*
*Assume $R$ is an integral domain. If $\{v_1, \ldots, v_n\}$ is $R$-linearly independent and $0 \neq r \in R$, then $\{rv_1, \ldots, rv_n\}$ is $R$-linearly independent.*

*Proof.* Let $a_1, \dots, a_n \in R$ and suppose $a_1(rv_1) + \cdots + a_n(rv_n) = 0$. Then $(ra_1)v_1 + \cdots + (ra_n)v_n = 0$. Independence gives $ra_i = 0$ for all $i$. Since $R$ is a domain and $r \neq 0$, $a_i = 0$ for all $i$.
*Structural Commentary*: Domains preserve independence under nonzero scaling.
*Key Insight*: The failure of this statement is exactly zero-divisor behavior. □

## 1.4 Module Homomorphisms, Kernels, and Images

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity.

**Definition 1.4.1.** *Module Homomorphism*
Let $M, N$ be $R$-modules and let $\tau : M \to N$.

(i) $\tau$ is *$R$-linear* if, $\forall r, s \in R$ and $u, v \in M$, $\tau(ru+sv) = r\tau(u)+s\tau(v)$, and is then called an *$R$-homomorphism*.
(ii) The set of all $R$-homomorphisms $M \to N$ is $\mathrm{Hom}_R(M, N)$.
(iii) An $R$-homomorphism $M \to M$ is an *endomorphism*.
(iv) An injective, surjective, or bijective $R$-homomorphism is a *monomorphism*, *epimorphism*, or *isomorphism*.

**Proposition 1.4.2.** *Hom, Kernel, and Image*
*Let $M, N$ be $R$-modules and $\tau \in \mathrm{Hom}_R(M, N)$.*

*(i) With $(a\sigma + b\rho)(m) := a\sigma(m) + b\rho(m)$, $\mathrm{Hom}_R(M, N)$ is an $R$-module.*
*(ii) $\ker\tau := \tau^{-1}(\{0\}) \leq M$.*
*(iii) $\mathrm{im}\,\tau := \tau(M) \leq N$.*
*(iv) $\tau$ is a monomorphism $\iff \ker\tau = \{0\}$.*

*Proof.* By Theorem 1.1.6, for $A \subseteq P$ in an $R$-module $P$, $A \leq P \iff A \neq \emptyset$ and $\forall a, b \in R$, $\forall x, y \in A$, $ax+by \in A$.

(i) For $\sigma, \rho \in \mathrm{Hom}_R(M, N)$ and $a, b, r, s \in R$, $(a\sigma + b\rho)(ru + sv) = r(a\sigma + b\rho)(u) + s(a\sigma + b\rho)(v)$, where $u, v \in M$, using commutativity of $R$. Since $0 \in \mathrm{Hom}_R(M, N)$, the displayed criterion gives the claim inside the pointwise module of maps $M \to N$.
(ii) Since $0 \in \ker\tau$, and $x, y \in \ker\tau$, $a, b \in R$ imply $\tau(ax + by) = a\tau(x) + b\tau(y) = 0$, so the displayed criterion gives $\ker\tau \leq M$.
(iii) Since $0 = \tau(0) \in \mathrm{im}\,\tau$, and $y_i = \tau(x_i) \in \mathrm{im}\,\tau$ $(i = 1, 2)$, $a, b \in R$ imply $ay_1 + by_2 = \tau(ax_1 + bx_2) \in \mathrm{im}\,\tau$, so the displayed criterion gives $\mathrm{im}\,\tau \leq N$.
(iv) If $\tau$ is injective, then $\tau(m) = 0 = \tau(0)$ implies $m = 0$, so $\ker\tau = \{0\}$. Conversely, if $\ker\tau = \{0\}$ and $\tau(m) = \tau(m')$, then $\tau(m - m') = 0$, hence $m = m'$.

*Structural Commentary*: Kernels and images are the canonical submodules attached to a module map.
*Key Insight*: Homomorphisms preserve precisely the finite $R$-linear combinations that define submodules. □

## 1.5 Quotient Modules, Correspondence, and Isomorphism

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity and $M$ denotes an $R$-module.

**Definition 1.5.1.** *Quotient Module*
Let $S \leq M$. Define $u \equiv v \pmod{S} \iff u - v \in S$.
The *quotient module* is $M/S := \{u + S : u \in M\}$; $u + S := \{u + s : s \in S\}$,
with operations $(u + S) + (v + S) := (u + v) + S$; $r(u + S) := ru + S$.

**Proposition 1.5.2.** *Quotient Operations Are Well-Defined*
*Let $S \leq M$. The operations of Definition 1.5.1 are well-defined and make $M/S$ an $R$-module.*
*Moreover, $0_{M/S} = S$, $\pi : M \twoheadrightarrow M/S$, $\pi(u) := u + S$, $\ker\pi = S$.*

*Proof.* If $u \equiv u' \pmod{S}$ and $v \equiv v' \pmod{S}$, then $(u + v) - (u' + v') = (u - u') + (v - v') \in S$ and $ru - ru' = r(u - u') \in S$. Thus $+$ and $R$-scaling on cosets are representative-independent. The module axioms descend from $M$. Finally, $0_{M/S} = 0 + S = S$, $\pi$ is surjective, and $\ker\pi = \{u \in M : u + S = S\} = \{u \in M : u \in S\} = S$. *Structural Commentary*: Quotients collapse exactly the submodule $S$ to zero.
*Key Insight*: Two representatives define the same coset precisely when their difference is invisible modulo $S$. □

**Theorem 1.5.3.** *Correspondence Theorem for Modules*
*Let $S \leq M$ and let $\pi : M \twoheadrightarrow M/S$ be the quotient map. Then $\Phi : \{T \leq M : S \leq T\} \longrightarrow \{U \leq M/S\}$, where $\Phi(T) := T/S$ is an order isomorphism, with inverse $\Psi(U) := \pi^{-1}(U)$.*

*Proof.* If $S \leq T \leq M$, then $\Phi(T) = \pi(T) \leq M/S$.
If $U \leq M/S$, then $\Psi(U) = \pi^{-1}(U) \leq M$ and $S = \ker\pi \leq \Psi(U)$.
Moreover, $\Psi\Phi(T) = \pi^{-1}(T/S) = T$ and $\Phi\Psi(U) = \pi(\pi^{-1}(U)) = U$, where the second equality uses surjectivity of

$\pi$. Finally, $T_1 \le T_2 \implies T_1/S \le T_2/S$, and $U_1 \le U_2 \implies \pi^{-1}(U_1) \le \pi^{-1}(U_2)$. Thus $\Phi$ is an order isomorphism.
*Structural Commentary*: Submodules above $S$ are exactly the submodules visible after $S$ is collapsed.
*Key Insight*: Pull back along the quotient map to recover the original submodule. □

**Theorem 1.5.4.** *Module Isomorphism Theorems*
*The following canonical maps are $R$-module isomorphisms.*

*(i)* *If $\tau \in \mathrm{Hom}_R(M,N)$, then $\overline{\tau} : M/\ker\tau \xrightarrow{\sim} \mathrm{im}\,\tau$ and $\overline{\tau}(m+\ker\tau) := \tau(m)$.*
*(ii)* *If $S,T \le M$, then $\theta : S/(S\cap T) \xrightarrow{\sim} (S+T)/T$ and $\theta\big(s+(S\cap T)\big) := s+T$.*
*(iii)* *If $S \le T \le M$, then $\eta : (M/S)/(T/S) \xrightarrow{\sim} M/T$ and $\eta\big((m+S)+T/S\big) := m+T$.*

*Proof.*

(i) If $m-m' \in \ker\tau$, then $\tau(m)=\tau(m')$, so $\overline{\tau}$ is well-defined and $R$-linear.
Also $\mathrm{im}\,\overline{\tau} = \mathrm{im}\,\tau$ and $\ker\overline{\tau} = \{m+\ker\tau : \tau(m)=0\} = \{0_{M/\ker\tau}\}$.
(ii) Apply (i) to $\varphi \in \mathrm{Hom}_R(S,(S+T)/T)$, $\varphi(s) := s+T$.
Then $\mathrm{im}\,\varphi = (S+T)/T$ and $\ker\varphi = \{s\in S : s\in T\} = S\cap T$.
(iii) Apply (i) to $\psi \in \mathrm{Hom}_R(M/S, M/T)$, $\psi(m+S) := m+T$.
Then $\mathrm{im}\,\psi = M/T$ and $\ker\psi = \{m+S : m\in T\} = T/S$.

*Structural Commentary*: Each theorem identifies a quotient by the exact ambiguity in a natural map.
*Key Insight*: Image is domain modulo kernel, repeatedly. □

## 1.6 Torsion and Annihilators

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity and $M$ denotes an $R$-module.

**Definition 1.6.1.** *Torsion*

(i) A nonzero element $v \in M$ is a *torsion element* if $\exists\, 0 \ne r \in R$ such that $rv = 0$.
(ii) $M$ is *torsion-free* if it has no torsion elements.
(iii) $M$ is a *torsion module* if every nonzero element of $M$ is torsion.
(iv) The *torsion set* of $M$ is $M_{\mathrm{tor}} := \{0\} \cup \{v \in M : v \text{ is torsion}\}$

**Example 1.6.2.** *$\mathbb{Z}/n\mathbb{Z}$ as a Torsion Module*
For $n>1$, $\mathbb{Z}/n\mathbb{Z}$ is a $\mathbb{Z}$-module via $k\cdot\overline{a} = \overline{ka}$. For every $\overline{a} \in \mathbb{Z}/n\mathbb{Z}$, $n\overline{a} = \overline{0}$.
Hence $\mathbb{Z}/n\mathbb{Z}$ is a torsion module, and $\{\overline{a}\}$ is linearly dependent whenever $\overline{a} \ne \overline{0}$.

**Proposition 1.6.3.** *Torsion Submodule over an Integral Domain*
*If $R$ is an integral domain, then $M_{\mathrm{tor}} \le M$, $(M/M_{\mathrm{tor}})_{\mathrm{tor}} = \{0\}$.*

*Proof.* Let $x,y \in M_{\mathrm{tor}}$ and $r,s \in R$.
Choose $a,b \in R\setminus\{0\}$ with $ax=0$ and $by=0$, taking $a=1$ if $x=0$ and $b=1$ if $y=0$.
Since $R$ is a domain, $ab \ne 0$, and $ab(rx+sy) = rb(ax)+sa(by) = 0$. Thus $rx+sy \in M_{\mathrm{tor}}$, so $M_{\mathrm{tor}} \le M$.
Now let $0 \ne r \in R$ and suppose $r(m+M_{\mathrm{tor}}) = M_{\mathrm{tor}}$. Then $rm \in M_{\mathrm{tor}}$, so $srm = 0$ for some $0 \ne s \in R$; if $rm=0$, take $s=1$. Since $sr \ne 0$, $m \in M_{\mathrm{tor}}$, hence $m+M_{\mathrm{tor}} = M_{\mathrm{tor}}$.
*Structural Commentary*: Domains let annihilators be multiplied without becoming zero.
*Key Insight*: Domains make torsion closed and quotients torsion-free. □

**Definition 1.6.4.** *Annihilators*

(i) For $v \in M$, the *annihilator* of $v$ is $\mathrm{Ann}_R(v) := \{r \in R : rv = 0\}$.
(ii) For $N \le M$, the *annihilator* of $N$ is $\mathrm{Ann}_R(N) := \{r \in R : rN = \{0\}\}$.

**Proposition 1.6.5.** *Annihilator Properties*
*For $v \in M$ and $N \le M$:*

*(i)* *$\mathrm{Ann}_R(v) \trianglelefteq R$ and $\mathrm{Ann}_R(N) \trianglelefteq R$.*
*(ii)* *$v$ is torsion $\iff v \ne 0$ and $\mathrm{Ann}_R(v) \ne \{0\}$.*
*(iii)* *$A \le B \le M \implies \mathrm{Ann}_R(B) \subseteq \mathrm{Ann}_R(A)$.*

*Proof.*

*(i)* Since $0 \in \mathrm{Ann}_R(v)$ and, for $a,b \in \mathrm{Ann}_R(v)$ and $r,s \in R$, $(ra+sb)v = r(av)+s(bv) = 0$, $\mathrm{Ann}_R(v) \trianglelefteq R$.
The same computation for every $n \in N$ gives $\mathrm{Ann}_R(N) \trianglelefteq R$.
*(ii)* ($\Rightarrow$) If $v$ is torsion, then $v \ne 0$ and $\exists\, 0 \ne r \in R$ with $rv = 0$, so $\mathrm{Ann}_R(v) \ne \{0\}$.
*(ii)* ($\Leftarrow$) If $v \ne 0$ and $\mathrm{Ann}_R(v) \ne \{0\}$, choose $0 \ne r \in \mathrm{Ann}_R(v)$. Then $rv = 0$, so $v$ is torsion.
*(iii)* If $A \le B \le M$ and $r \in \mathrm{Ann}_R(B)$, then $rA = \{0\}$; hence $r \in \mathrm{Ann}_R(A)$.

*Structural Commentary*: Annihilators reverse inclusion.
*Key Insight*: Killing more vectors gives fewer scalars. □

**Proposition 1.6.6.** *Torsion Generators Give a Nonzero Annihilator over a Domain*
*Assume $R$ is an integral domain. If $M = \langle\langle u_1, \dots, u_n \rangle\rangle_R$, $u_i \in M_{\text{tor}}$ $\forall i$, then $\text{Ann}_R(M) \neq \{0\}$.*

*Proof.* Choose $0 \neq a_i \in R$ with $a_i u_i = 0$, and set $a := \prod_{i=1}^{n} a_i$. Since $R$ is a domain, $a \neq 0$.
For each $i$, $au_i = 0$, hence $aM = \{0\}$ because the $u_i$ generate $M$. Thus $0 \neq a \in \text{Ann}_R(M)$.
*Structural Commentary*: Finite generation lets many annihilators be compressed into one product.
*Key Insight*: A finite torsion generating set gives a global order for the module. □

## 1.7 Direct Sums and Complements

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity and $M$ denotes an $R$-module.

**Definition 1.7.1.** *Direct Sums and Direct Summands*

(i) For $R$-modules $M_1, \dots, M_n$, their *external direct sum* is $\bigoplus_{i=1}^{n} M_i := \{(m_i)_{i=1}^n : m_i \in M_i\}$ with componentwise $R$-module operations.
(ii) For submodules $\{S_i : i \in I\}$ of $M$, their *internal direct sum* is denoted $M = \bigoplus_{i \in I} S_i$, which means $M = \sum_{i \in I} S_i$ and $\forall i \in I$, $S_i \cap \sum_{j \neq i} S_j = \{0\}$.
(iii) A submodule $S \leq M$ is a *direct summand* if $\exists\, T \leq M$ such that $M = S \oplus T$. Such a $T$ is a *complement* of $S$ in $M$.

**Proposition 1.7.2.** *Direct Sum Characterizations*
*For a family $\{S_i : i \in I\}$ of distinct submodules of $M$, the following conditions are equivalent:*

*(i)* $\sum_{i \in I} S_i$ *is direct, i.e.* $\forall k \in I$, $S_k \cap \sum_{i \neq k} S_i = \{0\}$.
*(ii) For every* $F \subseteq_{\text{fin}} I$ *and* $x_i \in S_i$ $(i \in F)$, $\sum_{i \in F} x_i = 0 \implies x_i = 0\ \forall i \in F$.
*(iii) Every* $0 \neq x \in \sum_{i \in I} S_i$ *has a unique finite nonzero decomposition,* $\exists!\, (F, (x_i)_{i \in F})$ *where* $F \subseteq_{\text{fin}} I$; $0 \neq x_i \in S_i$; $x = \sum_{i \in F} x_i$.

*Proof.*

*(i)* $\Rightarrow$ *(ii)*. If (i) holds and $\sum_{i \in F} x_i = 0$, then any $x_k \neq 0$ gives $x_k = -\sum_{i \in F \setminus \{k\}} x_i \in S_k \cap \sum_{i \neq k} S_i$, a contradiction.
*(ii)* $\Rightarrow$ *(i)*. If $x \in S_k \cap \sum_{i \neq k} S_i$, write $x = \sum_{i \in F} x_i$ with $F \subseteq_{\text{fin}} I \setminus \{k\}$ and $x_i \in S_i$.
Then $x - \sum_{i \in F} x_i = 0$, so (ii) gives $x = 0$.
*(ii)* $\Rightarrow$ *(iii)*. Existence follows from the definition of $\sum_i S_i$ after deleting zero terms.
If $\sum_{i \in F} x_i = \sum_{j \in G} y_j$, $0 \neq x_i \in S_i$, $0 \neq y_j \in S_j$, then, over $H = F \cup G$, missing terms set to 0, $\sum_{h \in H} (x_h - y_h) = 0$.
By (ii), $x_h = y_h$ for all $h \in H$, hence $F = G$ and the terms agree.
*(iii)* $\Rightarrow$ *(ii)*. If $\sum_{i \in F} x_i = 0$, set $F_0 := \{i \in F : x_i \neq 0\}$.
If $F_0 \neq \emptyset$, choose $k \in F_0$. If $F_0 = \{k\}$, then $x_k = 0$, contradiction; otherwise $0 \neq x_k = -\sum_{i \in F_0 \setminus \{k\}} x_i$,
giving two finite nonzero decompositions of $x_k$, contradiction. Hence $F_0 = \emptyset$.

*Structural Commentary*: Directness is the absence of hidden cancellation among different summands.
*Key Insight*: Unique decomposition is controlled by the zero vector. □

**Example 1.7.3.** *Most Submodules of $\mathbb{Z}$ Are Not Complemented*
As a $\mathbb{Z}$-module, every submodule of $\mathbb{Z}$ is $n\mathbb{Z}$ for a unique $n \geq 0$. For $m, n > 0$, $m\mathbb{Z} \cap n\mathbb{Z} = \text{lcm}(m,n)\mathbb{Z} \neq \{0\}$. Thus $m\mathbb{Z} \oplus n\mathbb{Z} \neq \mathbb{Z}$ for all $m, n > 0$, so the only complemented submodules of $\mathbb{Z}$ are $\{0\}$ and $\mathbb{Z}$.

**Theorem 1.7.4.** *Complements and Quotients*
*If $M = S \oplus T$, then $\pi|_T : T \xrightarrow{\sim} M/S$, $t \mapsto t + S$, where $\pi : M \twoheadrightarrow M/S$ is the quotient map.*
*Hence any two complements of $S$ are isomorphic.*

*Proof.* $\ker(\pi|_T) = T \cap S = \{0\}$. Also, for $m = s + t$ with $s \in S$, $t \in T$, one has $m + S = t + S$; hence $\text{im}(\pi|_T) = M/S$. Thus $\pi|_T$ is an isomorphism, and all complements are isomorphic to $M/S$.
*Structural Commentary*: A complement gives a concrete copy of the quotient inside the original module.
*Key Insight*: Quotienting by $S$ remembers exactly the complementary coordinate. □

## 1.8 Retracts, One-Sided Inverses, and Split Maps

Throughout this subsection, $R$ denotes a nonzero commutative ring with identity.

**Theorem 1.8.1.** *Direct Summands Extend Epimorphisms*
*Let $N \leq M$ and let $M_1$ be an $R$-module.*

*(i) If $M = N \oplus H$ and $\sigma : N \twoheadrightarrow M_1$, then $\exists$ a unique $\overline{\sigma} \in \mathrm{Hom}_R(M, M_1)$ such that $\overline{\sigma}(n+h) := \sigma(n)$, $n \in N$, $h \in H$, $\overline{\sigma}|_N = \sigma$, $\overline{\sigma} : M \twoheadrightarrow M_1$, and $\ker \overline{\sigma} = \ker \sigma \oplus H$.*

*(ii) If $\sigma : N \xrightarrow{\sim} M_1$, then $\mathcal{E}_\sigma := \{\overline{\sigma} : M \twoheadrightarrow M_1 : \overline{\sigma}|_N = \sigma\} \longleftrightarrow \mathcal{C}_N(M) := \{H \leq M : M = N \oplus H\}$ via $\overline{\sigma} \mapsto \ker \overline{\sigma}$ and $H \mapsto \overline{\sigma}_H$, where $\overline{\sigma}_H(n+h) := \sigma(n)$.*

*Proof.*

(i) Direct-sum uniqueness makes $\overline{\sigma}$ well-defined, and componentwise computation gives $R$-linearity. Moreover, $\mathrm{im}\,\overline{\sigma} = \mathrm{im}\,\sigma = M_1$; $\ker \overline{\sigma} = \{n+h : \sigma(n) = 0\} = \ker \sigma \oplus H$. This shows uniqueness.

(ii) If $H \in \mathcal{C}_N(M)$, then (i) gives $\overline{\sigma}_H \in \mathcal{E}_\sigma$ and, since $\ker \sigma = \{0\}$, $\ker \overline{\sigma}_H = H$. Conversely, if $\overline{\sigma} \in \mathcal{E}_\sigma$, then $N \cap \ker \overline{\sigma} = \{0\}$, and for each $m \in M$, $n := \sigma^{-1}(\overline{\sigma}(m)) \in N$, $m - n \in \ker \overline{\sigma}$. Thus $M = N \oplus \ker \overline{\sigma}$, so the two displayed assignments are inverse.

*Structural Commentary*: A complement supplies the directions on which an extension may vanish.
*Key Insight*: Extending an isomorphism is equivalent to choosing its kernel complement. □

**Definition 1.8.2.** *Retract*
Let $N \leq M$.

(i) A *retraction* onto $N$ is a map $\rho \in \mathrm{Hom}_R(M, N)$ such that $\rho|_N = \mathrm{id}_N$.
(ii) The submodule $N$ is a *retract* of $M$ iff $\exists\, \rho \in \mathrm{Hom}_R(M, N)$ with $\rho|_N = \mathrm{id}_N$.

**Corollary 1.8.3.** *Retracts Are Direct Summands*
*For $N \leq M$, $N$ is a retract of $M \iff \exists\, H \leq M$ such that $M = N \oplus H$.*

*Proof.*

($\Rightarrow$) Let $\rho : M \to N$ satisfy $\rho|_N = \mathrm{id}_N$. For every $m \in M$, $m = \rho(m) + (m - \rho(m))$, $\rho(m) \in N$, $m - \rho(m) \in \ker \rho$. Moreover $N \cap \ker \rho = \{0\}$, so $M = N \oplus \ker \rho$.

($\Leftarrow$) If $M = N \oplus H$, define $\rho : M \to N$ by $\rho(n+h) := n$, $n \in N$, $h \in H$. Then $\rho \in \mathrm{Hom}_R(M, N)$ and $\rho|_N = \mathrm{id}_N$, so $N$ is a retract.

*Structural Commentary*: Retractions are projections onto submodules.
*Key Insight*: A submodule is split exactly when it can be recovered by an idempotent map. □

**Definition 1.8.4.** *One-Sided Inverses*
Let $\tau \in \mathrm{Hom}_R(A, B)$.

(i) A *left inverse* of $\tau$ is $\lambda \in \mathrm{Hom}_R(B, A)$ such that $\lambda\tau = \mathrm{id}_A$.
(ii) A *right inverse* of $\tau$ is $\rho \in \mathrm{Hom}_R(B, A)$ such that $\tau\rho = \mathrm{id}_B$.
(iii) A *two-sided inverse* of $\tau$ is $\upsilon \in \mathrm{Hom}_R(B, A)$ such that $\upsilon\tau = \mathrm{id}_A$ and $\tau\upsilon = \mathrm{id}_B$.

**Theorem 1.8.5.** *One-Sided Inverses and Complements*
*Let $\sigma : M \to M_1$ be an $R$-homomorphism.*
*For $S \leq N$, set $\mathcal{C}_S(N) := \{H \leq N : N = S \oplus H\}$, $\mathcal{L}_\sigma := \{\lambda \in \mathrm{Hom}_R(M_1, M) : \lambda\sigma = \mathrm{id}_M\}$, and $\mathcal{R}_\sigma := \{\rho \in \mathrm{Hom}_R(M_1, M) : \sigma\rho = \mathrm{id}_{M_1}\}$.*

*(i) If $\sigma$ is injective, then $\mathcal{L}_\sigma \xrightarrow{\sim} \mathcal{C}_{\mathrm{im}\,\sigma}(M_1)$ by $\lambda \mapsto \ker \lambda$.*

*(ii) If $\sigma$ is surjective, then $\mathcal{R}_\sigma \xrightarrow{\sim} \mathcal{C}_{\ker \sigma}(M)$ by $\rho \mapsto \mathrm{im}\,\rho$.*

*Proof.*

(i) For $H \in \mathcal{C}_{\mathrm{im}\,\sigma}(M_1)$, define $\lambda_H(\sigma m + h) := m$. Directness and injectivity give well-definedness; $\lambda_H\sigma = \mathrm{id}_M$ and $\ker \lambda_H = H$. Conversely, if $\lambda \in \mathcal{L}_\sigma$, then $\mathrm{im}\,\sigma \cap \ker \lambda = \{0\}$, since $\sigma m \in \ker \lambda \Rightarrow m = 0$, and $y = \sigma\lambda y + (y - \sigma\lambda y)$ with $y - \sigma\lambda y \in \ker \lambda$. Thus $M_1 = \mathrm{im}\,\sigma \oplus \ker \lambda$, and the constructions are inverse.

(ii) For $H \in \mathcal{C}_{\ker \sigma}(M)$, $\sigma|_H : H \to M_1$ is an isomorphism; let $\rho_H = (\sigma|_H)^{-1}$ followed by $H \hookrightarrow M$. Then $\rho_H \in \mathcal{R}_\sigma$ and $\mathrm{im}\,\rho_H = H$. Conversely, if $\rho \in \mathcal{R}_\sigma$, then $\rho$ is injective, $\mathrm{im}\,\rho \cap \ker \sigma = \{0\}$, and $m = (m - \rho\sigma m) + \rho\sigma m$ with $m - \rho\sigma m \in \ker \sigma$, $\rho\sigma m \in \mathrm{im}\,\rho$. Thus $M = \ker \sigma \oplus \mathrm{im}\,\rho$, and the constructions are inverse.

*Structural Commentary*: One-sided inverses exist exactly when the relevant image or kernel splits.
*Key Insight*: Injective maps need a complemented image; surjective maps need a complemented kernel. □

**Proposition 1.8.6.** *Split Epimorphisms Split the Domain*
*Let $\sigma : M \twoheadrightarrow N$ be an $R$-homomorphism and set $\mathcal{R}_\sigma := \{\rho \in \mathrm{Hom}_R(N, M) : \sigma\rho = \mathrm{id}_N\}$. These are equivalent:*

*(i)* $\mathcal{R}_\sigma \neq \emptyset$.
*(ii)* $\exists\, H \leq M$ *such that* $M = \ker\sigma \oplus H$.

*Moreover,* $\rho \in \mathcal{R}_\sigma \implies M = \ker\sigma \oplus \mathrm{im}\,\rho$.

*Proof.*

($\Rightarrow$) Let $\rho \in \mathcal{R}_\sigma$. If $\rho n = 0$, then $n = \sigma\rho n = 0$, so $\rho$ is injective.
If $\rho n \in \ker\sigma$, then $n = \sigma\rho n = 0$, hence $\mathrm{im}\,\rho \cap \ker\sigma = \{0\}$.
For $m \in M$, $m = (m - \rho\sigma m) + \rho\sigma m$, with $m - \rho\sigma m \in \ker\sigma$ and $\rho\sigma m \in \mathrm{im}\,\rho$. Thus $M = \ker\sigma \oplus \mathrm{im}\,\rho$.
($\Leftarrow$) Let $M = \ker\sigma \oplus H$. Then $\ker(\sigma|_H) = H \cap \ker\sigma = \{0\}$, and $\mathrm{im}(\sigma|_H) = N$ since $n = \sigma m$, $m = k + h$, gives $n = \sigma h$. Thus $\sigma|_H : H \xrightarrow{\sim} N$, and $\rho := \iota_H(\sigma|_H)^{-1} \in \mathcal{R}_\sigma$, where $\iota_H : H \hookrightarrow M$.

*Structural Commentary*: A split epimorphism is a quotient map with complemented kernel.
*Key Insight*: The right inverse chooses canonical representatives for the cosets of $\ker\sigma$. □

# 2 Noetherian Modules and PID Structure Theorems

## 2.1 Free Modules, Bases, and Universal Property

This section begins with the structure facts that make free modules and finite generation usable: rank, splitting maps onto free modules, Noetherian modules. It then develops the classification of finitely generated modules over a principal ideal domain. The guiding model is the abelian-group theorem over $\mathbb{Z}$ and, later, the polynomial-module classification behind rational canonical form.

**Definition 2.1.1.** *Free Module and Basis*

(i) A subset $\mathcal{B} \subseteq M$ is a *basis* of $M$ if $\mathcal{B}$ is $R$-linearly independent and $\langle\langle \mathcal{B} \rangle\rangle_R = M$.
(ii) An $R$-module $M$ is *free* if $M = \{0\}$ or $M$ has a basis.
(iii) If $\mathcal{B}$ is a basis of $M$, then $M$ is *free on* $\mathcal{B}$.

**Theorem 2.1.2.** *Basis Means Unique Finite Coordinates*
*A subset $\mathcal{B} \subseteq M$ is a basis if and only if $\forall v \in M,\ \exists!\,(F, (r_b)_{b\in F})$, with $F \subseteq_{\mathrm{fin}} \mathcal{B}$, $r_b \in R \setminus \{0\}$, $v = \sum_{b\in F} r_b b$.*

*The case $F = \emptyset$ gives the empty sum 0.*

*Proof.*

($\Rightarrow$) Spanning gives a finite expression; collecting equal basis elements and deleting zero coefficients gives the displayed form. If $\sum_{b\in F} r_b b = \sum_{b\in G} s_b b$, then $\sum_{b\in F\cup G} (r_b - s_b)b = 0$, where missing coefficients are 0.
Independence gives $r_b = s_b$ for all $b$, hence $F = G$ and the coefficients agree.
($\Leftarrow$) Existence gives $\langle\langle \mathcal{B} \rangle\rangle_R = M$. If $\sum_{b\in F} r_b b = 0$ and $F_0 := \{b \in F : r_b \neq 0\}$ is nonempty, then $0 = \sum_{b\in F_0} r_b b$ contradicts uniqueness of the empty expression. Thus all $r_b = 0$, so $\mathcal{B}$ is independent.

*Structural Commentary*: Bases give unique finite coordinates.
*Key Insight*: Existence is spanning; uniqueness is independence. □

**Proposition 2.1.3.** *Bases Are Minimal Spanning and Maximal Independent*
*If $\mathcal{B}$ is a basis of $M$, then $\forall b \in \mathcal{B}$, $\langle\langle \mathcal{B} \setminus \{b\} \rangle\rangle_R \neq M$, and $\forall v \in M \setminus \mathcal{B}$, $\mathcal{B} \cup \{v\}$ is $R$-linearly dependent.*

*Proof.* If $b \in \mathcal{B}$ and $b \in \langle\langle \mathcal{B} \setminus \{b\} \rangle\rangle_R$, then $b$ has two finite coordinate expressions, contradicting Theorem 2.1.2. Hence deleting $b$ destroys spanning. If $v \notin \mathcal{B}$, write $v = \sum_{b\in F} r_b b$ with $F \subseteq_{\mathrm{fin}} \mathcal{B}$.
Then $v - \sum_{b\in F} r_b b = 0$ is a nontrivial relation in $\mathcal{B} \cup \{v\}$, so $\mathcal{B} \cup \{v\}$ is dependent.
*Structural Commentary*: Basis still implies the familiar minimal/maximal properties.
*Key Insight*: The converses are vector-space privileges, not general module facts. □

**Theorem 2.1.4.** *Maps Out of a Free Module*
*Let $M$ be free on $\mathcal{B}$ and let $N$ be an $R$-module.*
*For every function $f : \mathcal{B} \to N$, there exists a unique $\widetilde{f} \in \mathrm{Hom}_R(M, N)$ such that $\widetilde{f}|_{\mathcal{B}} = f$.*

*Proof.* For $v = \sum_{b\in F} r_b b$ with $F \subseteq_{\mathrm{fin}} \mathcal{B}$, define $\widetilde{f}(v) := \sum_{b\in F} r_b f(b)$.
Theorem 2.1.2 makes the expression of $v$ unique, so $\widetilde{f}$ is well-defined.
For $x = \sum a_b b$ and $y = \sum c_b b$, combining supports gives $\widetilde{f}(rx + sy) = r\widetilde{f}(x) + s\widetilde{f}(y)$, so $\widetilde{f}$ is $R$-linear and clearly

extends $f$. If $\phi \in \mathrm{Hom}_R(M,N)$ also extends $f$, then $\phi\left(\sum_{b\in F} r_b b\right) = \sum_{b\in F} r_b f(b) = \widetilde{f}\left(\sum_{b\in F} r_b b\right)$, hence $\phi = \widetilde{f}$.

*Structural Commentary*: A free module is controlled entirely by its basis values.
*Key Insight*: Freeness is the universal permission to choose images of basis elements arbitrarily. □

**Example 2.1.5.** *A Free Module with a Nonfree Submodule*
Let $R = \mathbb{Z}\times\mathbb{Z}$ with componentwise multiplication. As an $R$-module, $R = R(1,1)$, so $R$ is free on $\{(1,1)\}$.
Set $N := \mathbb{Z}\times\{0\} \le R$. For every $0 \neq (a,0) \in N$, $(0,1)(a,0) = 0$.
Thus every nonzero element of $N$ is torsion, so $N$ has no nonempty independent subset and is not free.

**Example 2.1.6.** *A Quotient of a Free Module Need Not Be Free*
As a $\mathbb{Z}$-module, $\mathbb{Z}$ is free on $\{1\}$. For $n > 1$, the submodule $n\mathbb{Z}$ is cyclic, but the quotient $\mathbb{Z}/n\mathbb{Z}$ is torsion by Example 1.6.2. Hence $\mathbb{Z}/n\mathbb{Z}$ is not free as a $\mathbb{Z}$-module.

**Principle 2.1.7.** *Modules Are Not Just Vector Spaces with New Letters*
For a general ring $R$, the following vector-space implications fail:

(i) $N \le M \not\Rightarrow \exists\, H \le M$ with $M = N \oplus H$.
(ii) $M$ finitely generated and $N \le M \not\Rightarrow N$ finitely generated.
(iii) $M \neq 0 \not\Rightarrow \exists\, \emptyset \neq S \subseteq M$ linearly independent.
(iv) $S$ minimal spanning or maximal independent $\not\Rightarrow S$ is a basis.
(v) $M$ free and $N \le M \not\Rightarrow N$ free.
(vi) $F$ free and $S \le F \not\Rightarrow F/S$ free.

Over a field, these obstructions vanish because $R\setminus\{0\} \subseteq R^\times$.

## 2.2 Rank and Coordinate Models of Free Modules

**Perspective 2.2.1.** *Dimension as the Field Case of Rank*
An $\mathbb{F}$-vector space is a free $\mathbb{F}$-module. By Theorem 1.3.14, finite bases have invariant cardinality; over general rings, rank invariance requires additional hypotheses.

**Theorem 2.2.2.** *Rank Is Well Defined fo r Free Modules*
*Let $A$ benonzero commutative ring with identity, $M$ be free $A$-module, $\mathrm{Bas}_A(M)$ denote set of $A$-bases of $M$.*

*(i)* $B, B' \in \mathrm{Bas}_A(M) \Longrightarrow |B| = |B'|$.
*(ii)* $B \in \mathrm{Bas}_A(M)$, $S \subseteq M$, *and* $\langle\!\langle S\rangle\!\rangle_A = M \Longrightarrow |B| \le |S|$.

*Proof.*

(i) Choose a maximal ideal $\mathcal{I} \lhd A$ by Theorem 3.4.5.
Put $K := A/\mathcal{I}$, a field by Theorem 3.4.4, $\mathcal{I}M := \langle\!\langle im : i\in\mathcal{I},\ m\in M\rangle\!\rangle_A$, and $V := M/\mathcal{I}M$.
Since $\mathcal{I}V = 0$, $V$ is a $K$-vector space via $(a+\mathcal{I})(m+\mathcal{I}M) := am + \mathcal{I}M$.
(ii) Let $B \in \mathrm{Bas}_A(M)$ and set $\overline{B} := \{\overline{b} := b + \mathcal{I}M : b \in B\}$.
In $B$-coordinates, $\mathcal{I}M = \{\sum_{b\in F} i_b b : F \subseteq_{\mathrm{fin}} B,\ i_b \in \mathcal{I}\}$; hence $\overline{B}$ spans $V$.
If $\sum_{j=1}^{n} (a_j + \mathcal{I})\overline{b_j} = 0$ with distinct $b_j \in B$, then $\sum_{j=1}^{n} a_j b_j \in \mathcal{I}M$, so $a_j \in \mathcal{I}\ \forall j$; hence $\overline{B}$ is $K$-independent.
If $\overline{b} = \overline{b'}$ with $b \neq b'$, then $b - b' \in \mathcal{I}M$, forcing the $B$-coefficient $1 \in \mathcal{I}$, impossible. Thus $B \xrightarrow{\sim} \overline{B}$.
(iii) Therefore $|B| = \left|\overline{B}\right| = \dim_K V$ for every $B \in \mathrm{Bas}_A(M)$, so (i) follows from Theorem 1.4.7.
If $\langle\!\langle S\rangle\!\rangle_A = M$, then $\overline{S} := \{s + \mathcal{I}M : s \in S\}$ spans $V$, so $|B| = \dim_K V \le \left|\overline{S}\right| \le |S|$.

*Structural Commentary*: Modulo a maximal ideal, free module becomes a vector space with same basis size.
*Key Insight*: The field quotient $A/\mathcal{I}$ lets vector-space dimension certify module rank. □

**Definition 2.2.3.** *Rank of a Free Module*
Let $A$ be a nonzero commutative ring with identity. The *rank* of a free $A$-module $M$, denoted $\mathrm{rk}_A(M)$, is the cardinality of any basis of $M$. The zero module has rank 0.

**Example 2.2.4.** *Why Commutativity Matters for Rank*
Let $V = \bigoplus_{k\ge 0} \mathbb{F}e_k$ and $E = \mathcal{L}(V)$. As a left $E$-module, $E = E\,\mathrm{id}_V$, so $\{\mathrm{id}_V\}$ is a basis.
For each $n \ge 1$, define $\beta_s \in E$, $0 \le s < n$, by $\beta_s(e_{kn+t}) = \begin{cases} e_k, & t = s, \\ 0, & t \neq s, \end{cases}$ where $k \ge 0$, $0 \le t < n$.
If $\sum_{s=0}^{n-1} \alpha_s\beta_s = 0$, then evaluating at $e_{kn+t}$ gives $\alpha_t(e_k) = 0$ for all $k, t$, hence $\alpha_t = 0$ for all $t$.

Conversely, for $h \in E$, define $\alpha_s(e_k) := h(e_{kn+s})$; then $h = \sum_{s=0}^{n-1} \alpha_s \beta_s$.
Thus $\{\beta_0, \dots, \beta_{n-1}\}$ is an $E$-basis of $E$, so $E$ has bases of all positive finite cardinalities.
*Structural Commentary*: Noncommutative scalar rings can make a free module have incompatible basis sizes.
*Key Insight*: The rank theorem is a commutative-ring theorem, not a formal consequence of freeness alone.

**Theorem 2.2.5.** *Finite-Support Coordinate Modules*
*Let $A$ be a commutative ring with identity and let $B$ be a set. Put $A^{(B)} := \{f \in A^B : |\mathrm{supp}(f)| < \infty\}$.*
*For $b \in B$, let $\delta_b \in A^{(B)}$ be given by $\delta_b(b) = 1$ and $\delta_b(x) = 0$ for $x \neq b$.*
*Then $A^{(B)} = \bigoplus_{b \in B} A\delta_b$, so $\{\delta_b : b \in B\}$ is an $A$-basis. If $A \neq 0$, then $\mathrm{rk}_A(A^{(B)}) = |B|$.*

*Proof.* For $f, g \in A^{(B)}$ and $a \in A$, $\mathrm{supp}(f+g) \subseteq \mathrm{supp}(f) \cup \mathrm{supp}(g)$ and $\mathrm{supp}(af) \subseteq \mathrm{supp}(f)$, hence $A^{(B)} \leq A^B$.
For $f \in A^{(B)}$, $f = \sum_{b \in \mathrm{supp}(f)} f(b)\delta_b$, so the $\delta_b$ span. If $\sum_{j=1}^{m} a_j \delta_{b_j} = 0$ with $b_1, \dots, b_m$ distinct, then evaluation at $b_i$ gives $a_i = 0$ for every $i$. Thus the $\delta_b$ are independent, and the rank statement follows from the displayed basis.
*Structural Commentary*: Free modules are finite-support coordinate modules on their bases.
*Key Insight*: Finite support is the module-theoretic version of finite linear combination. □

**Proposition 2.2.6.** *Free Modules Are Coordinate Modules*
*Let $A$ be a commutative ring with identity and let $M$ be free on a basis $B$.*
*Then the coordinate map $\Phi_B : M \to A^{(B)}$ is an isomorphism.*
*If $A \neq 0$ and $M, N$ are free $A$-modules, then $M \cong N \iff \mathrm{rk}_A(M) = \mathrm{rk}_A(N)$.*

*Proof.* For $m = \sum_{b \in F} a_b b$, set $\Phi_B(m) = f_m$, with $f_m(b) = a_b$ for $b \in F$ and $f_m(b) = 0$ for $b \notin F$.
Unique coordinates give $\Phi_B : M \xrightarrow{\sim} A^{(B)}$, with inverse $f \mapsto \sum_{b \in \mathrm{supp}(f)} f(b)b$.

($\Rightarrow$) If $\psi : M \xrightarrow{\sim} N$ and $C$ is a basis of $M$, then $\psi(C)$ is a basis of $N$, so $\mathrm{rk}_A(M) = |C| = |\psi(C)| = \mathrm{rk}_A(N)$.
($\Leftarrow$) If $\mathrm{rk}_A(M) = \mathrm{rk}_A(N)$, choose bases $C, D$ with $|C| = |D|$.
Any bijection $C \to D$ extends, by Theorem 2.1.4, to $M \xrightarrow{\sim} N$.

*Structural Commentary*: A basis identifies a free module with a standard coordinate model.
*Key Insight*: Over a nonzero commutative ring, rank classifies free modules up to isomorphism. □

**Theorem 2.2.7.** *Independent Sets Are Bounded by Rank over a Domain*
*Let $A$ be integral domain with field of quotients $Q$, $M$ free over $A$, $S \subseteq M$ $A$-independent. Then $|S| \leq \mathrm{rk}_A(M)$.*

*Proof.* Choose an $A$-basis $B$ and identify $M = A^{(B)} \subseteq Q^{(B)}$.
If $\sum_{i=1}^{n} q_i s_i = 0$ with $q_i \in Q$ and distinct $s_i \in S$, choose $d \in A \setminus \{0\}$ with $dq_i \in A$ for all $i$.
Then $\sum_{i=1}^{n} (dq_i)s_i = 0$ in $M$, so $dq_i = 0$ by $A$-independence, hence $q_i = 0$ for all $i$.
Thus $S$ is $Q$-independent in $Q^{(B)}$, and $|S| \leq \dim_Q Q^{(B)} = |B| = \mathrm{rk}_A(M)$.
*Structural Commentary*: The field of quotients lets ordinary dimension bound module independence.
*Key Insight*: Domains allow denominator clearing without creating false zero relations. □

## 2.3 Free Quotients and Splittings

**Theorem 2.3.1.** *Free Quotients Split*
*Let $A$ be a commutative ring with identity.*

*(i) If $\sigma : M \twoheadrightarrow F$ and $F$ is free, then $\exists N \leq M$ such that $M = \ker(\sigma) \oplus N$ and $N \cong F$.*
*(ii) If $S \leq M$ and $M/S$ is free, then $\exists T \leq M$ such that $M = S \oplus T$ and $T \cong M/S$; hence $M \cong S \oplus M/S$.*
*(iii) If $A \neq 0$ and $M, S, M/S$ are free, then $\mathrm{rk}_A(M) = \mathrm{rk}_A(S) + \mathrm{rk}_A(M/S)$, with cardinal addition; in finite rank, $\mathrm{rk}_A(M/S) = \mathrm{rk}_A(M) - \mathrm{rk}_A(S)$.*

*Proof.*

(i) Let $B$ be a basis of $F$. For $b \in B$, choose $m_b \in M$ with $\sigma(m_b) = b$. By Theorem 2.1.4, $b \mapsto m_b$ extends to $\tau : F \to M$. Since $\sigma\tau = \mathrm{id}_F$, Theorem 1.8.5 gives $M = \ker(\sigma) \oplus \mathrm{im}(\tau)$ and $\mathrm{im}(\tau) \cong F$.
(ii) Apply (i) to $\pi : M \twoheadrightarrow M/S$. Then $M = S \oplus T$ for some $T \cong M/S$, so $M \cong S \oplus M/S$.
(iii) From (ii), $M = S \oplus T$ with $T \cong M/S$. If $B_S, B_T$ are bases of $S, T$, then $B_S \sqcup B_T$ is a basis of $M$, hence $\mathrm{rk}_A(M) = |B_S| + |B_T| = \mathrm{rk}_A(S) + \mathrm{rk}_A(M/S)$. The finite formula follows.

*Structural Commentary*: Free modules are projective: maps onto them have linear sections.
*Key Insight*: Choosing representatives for basis vectors of the quotient is enough to split the quotient map. □

## 2.4 Noetherian Modules

**Definition 2.4.1.** *Noetherian Modules and Rings*
Let $A$ be a commutative ring with identity.

(i) An $A$-module $M$ satisfies *ACC* if every chain $M_1 \leq M_2 \leq \cdots$ of submodules satisfies $\exists n_0, \forall n \geq n_0, M_n = M_{n_0}$. Such $M$ is *Noetherian*.
(ii) The ring $A$ is *Noetherian* if the regular $A$-module $A$ is Noetherian, equivalently if every ideal-chain $I_1 \subseteq I_2 \subseteq \cdots$ stabilizes.

**Theorem 2.4.2.** *Noetherian Means Finitely Generated Submodules*
*Let $M$ be an $A$-module.*

*(i) $M$ is Noetherian $\iff$ every $N \leq M$ is f.g.*
*(ii) If $A$ is commutative with identity, then $A$ is Noetherian $\iff$ every $\mathcal{I} \lhd A$ is f.g.*

*Proof.*

(i) ($\Rightarrow$) Let $N \leq M$. If $N$ is not f.g., choose $u_{n+1} \in N \setminus \langle\!\langle u_1, \ldots, u_n \rangle\!\rangle_A$ inductively. Then $\langle\!\langle u_1 \rangle\!\rangle_A < \langle\!\langle u_1, u_2 \rangle\!\rangle_A < \cdots$, contradicting ACC.
(i) ($\Leftarrow$) Let $M_1 \leq M_2 \leq \cdots$ and put $U = \bigcup_{n \geq 1} M_n$. Then $U \leq M$, so $U = \langle\!\langle u_1, \ldots, u_r \rangle\!\rangle_A$. Choose $k$ with $u_1, \ldots, u_r \in M_k$. Hence $U = M_k$ and $M_n = M_k$ for all $n \geq k$.
(ii) Apply (i) to the regular $A$-module $A$: its submodules are exactly the ideals of $A$.

*Structural Commentary*: ACC and finite generation are two faces of the same finiteness condition.
*Key Insight*: Failure of finite generation builds an infinite increasing chain one new generator at a time. □

## 2.5 Orders and Cyclic Modules over a PID

**Definition 2.5.1.** *Standing PID Notation*
In the PID-structure results, $R$ denotes a principal ideal domain unless otherwise stated.
For $r \in R$, write $\langle r \rangle = Rr$; for an $R$-module $M$ and $v \in M$, write $\langle\!\langle v \rangle\!\rangle_R = Rv$.

**Perspective 2.5.2.** *PID Arithmetic Controls Module Structure*
Fields have $F \setminus \{0\} \subseteq F^\times$. PIDs replace invertibility by factorization, gcds, and lcms.
This divisibility data classifies finitely generated modules.

**Definition 2.5.3.** *Order of a Submodule or Element*
Let $M$ be an $R$-module and $N \leq M$. Since $\mathrm{Ann}_R(N) = \{r \in R : rN = 0\} = \langle \alpha \rangle$, any such $\alpha$ is an *order* of $N$; write $o(N) = \alpha$, up to associates. For $v \in M$, set $o(v) := o(\langle\!\langle v \rangle\!\rangle_R)$.

**Proposition 2.5.4.** *Orders and Direct Sums*
*Let $M = A \oplus B$. If $o(A) = \alpha$ and $o(B) = \beta$, then $o(M) \sim \mathrm{lcm}(\alpha, \beta)$.*
*More generally, $o(\bigoplus_{i=1}^{n} M_i) \sim \mathrm{lcm}(o(M_1), \ldots, o(M_n))$.*

*Proof.* $r(A \oplus B) = 0 \iff rA = 0$ and $rB = 0$.
Hence $\mathrm{Ann}_R(A \oplus B) = \mathrm{Ann}_R(A) \cap \mathrm{Ann}_R(B) = \langle \alpha \rangle \cap \langle \beta \rangle = \langle \mathrm{lcm}(\alpha, \beta) \rangle$.
Thus $o(A \oplus B) \sim \mathrm{lcm}(\alpha, \beta)$, and induction gives the finite case.
*Structural Commentary*: Direct sums intersect annihilator conditions.
*Key Insight*: Lcms appear because one scalar must kill every summand at once. □

**Theorem 2.5.5.** *Cyclic Modules over a PID*
*Let $M$ be an $R$-module and let $v \in M$.*

*(i) If $\mathrm{Ann}_R(v) = \langle \alpha \rangle$, then $R/\langle \alpha \rangle \xrightarrow{\sim} \langle\!\langle v \rangle\!\rangle_R$, $r + \langle \alpha \rangle \mapsto rv$.*
*(ii) If $C = \langle\!\langle v \rangle\!\rangle_R$ and $S \leq C$, then $S = \langle\!\langle sv \rangle\!\rangle_R$ for some $s \in R$.*
*(iii) If $o(v) = \alpha$ and $\beta \in R$, then $o(\beta v) \sim \alpha / \gcd(\alpha, \beta)$, and $\langle\!\langle \beta v \rangle\!\rangle_R = \langle\!\langle v \rangle\!\rangle_R \iff \gcd(\alpha, \beta) \sim 1$.*

*Proof.*

(i) The map $\phi : R \to \langle\!\langle v \rangle\!\rangle_R$, $\phi(r) = rv$, is onto and $\ker \phi = \mathrm{Ann}_R(v) = \langle \alpha \rangle$. Hence $R/\langle \alpha \rangle \xrightarrow{\sim} \langle\!\langle v \rangle\!\rangle_R$.
(ii) For $S \leq C$, set $I = \{r \in R : rv \in S\} \lhd R$.
Since $R$ is a PID, $I = \langle s \rangle$ for some $s \in R$, and $S = \{rv : r \in I\} = \langle\!\langle sv \rangle\!\rangle_R$.
(iii) Put $d = \gcd(\alpha, \beta)$, $\alpha = d\alpha'$, and $\beta = d\beta'$ with $\gcd(\alpha', \beta') \sim 1$.
For $r \in R$, $r\beta v = 0 \iff \alpha \mid r\beta \iff \alpha' \mid r\beta' \iff \alpha' \mid r$, the last step by Bezout.
Thus $\mathrm{Ann}_R(\beta v) = \langle \alpha' \rangle = \langle \alpha / \gcd(\alpha, \beta) \rangle$. Finally, $\langle\!\langle \beta v \rangle\!\rangle_R = \langle\!\langle v \rangle\!\rangle_R \iff (\exists c \in R)(v = c\beta v)$
$\iff (\exists c \in R)(\alpha \mid 1 - c\beta) \iff \gcd(\alpha, \beta) \sim 1$.

*Structural Commentary*: A cyclic module is the base ring modulo the first relation satisfied by its generator.
*Key Insight*: Multiplying a cyclic generator divides its order by the gcd of the multiplier and the old order. □

**Theorem 2.5.6.** *Composing and Decomposing Cyclic Modules*
*Let $M$ be an $R$-module.*

*(i) If $o(u_i) = \alpha_i \neq 0$ and $\gcd(\alpha_i, \alpha_j) \sim 1$ for $i \neq j$, then $o(\sum_{i=1}^{n} u_i) \sim \prod_{i=1}^{n} \alpha_i$ and $\bigoplus_{i=1}^{n} \langle\langle u_i \rangle\rangle_R = \langle\langle \sum_{i=1}^{n} u_i \rangle\rangle_R$.*

*(ii) If $o(v) = \alpha = \prod_{i=1}^{n} \alpha_i \neq 0$ and $\gcd(\alpha_i, \alpha_j) \sim 1$ for $i \neq j$,*
*then $\exists v_1, \ldots, v_n \in M$ such that $v = \sum_{i=1}^{n} v_i$, $o(v_i) \sim \alpha_i$, and $\langle\langle v \rangle\rangle_R = \bigoplus_{i=1}^{n} \langle\langle v_i \rangle\rangle_R$.*

*Proof.*

(i) It suffices to prove $n = 2$ and iterate. Let $o(u) = \alpha$, $o(w) = \beta$, and $\gcd(\alpha, \beta) \sim 1$.
If $x \in \langle\langle u \rangle\rangle_R \cap \langle\langle w \rangle\rangle_R$, then $\alpha x = \beta x = 0$, so Bezout gives $x = 0$. If $r(u + w) = 0$, then $ru = -rw$ lies in this intersection; hence $ru = rw = 0$, so $\alpha \mid r$ and $\beta \mid r$, hence $\alpha\beta \mid r$. Since $\alpha\beta(u+w) = 0$, $o(u+w) \sim \alpha\beta$.
Choose $a, b \in R$ with $a\alpha + b\beta = 1$. Then $u = b\beta(u+w)$ and $w = a\alpha(u+w)$, so $\langle\langle u \rangle\rangle_R \oplus \langle\langle w \rangle\rangle_R = \langle\langle u+w \rangle\rangle_R$.

(ii) Put $\beta_i = \alpha/\alpha_i$. Since $\gcd(\beta_1, \ldots, \beta_n) \sim 1$, choose $c_i \in R$ with $\sum_{i=1}^{n} c_i\beta_i = 1$, and set $v_i = c_i\beta_i v$.
Then $v = \sum_{i=1}^{n} v_i$. Also $\gcd(c_i, \alpha_i) \sim 1$ and $\gcd(\alpha, \beta_i) \sim \beta_i$, so Theorem 2.5.5 gives $o(v_i) \sim \alpha_i$.
Part (i) gives $\langle\langle v \rangle\rangle_R = \bigoplus_{i=1}^{n} \langle\langle v_i \rangle\rangle_R$.

*Structural Commentary*: Relatively prime cyclic pieces merge into one cyclic piece.
*Key Insight*: Bezout identities recover each component from the single sum. □

**Proposition 2.5.7.** *Prime Divisors Give Prime-Order Cyclic Submodules*
*Let $M$ be f.g. torsion with order $\mu$. If $p \mid \mu$ is prime, then $\exists w \in M$ with $o(w) \sim p$; equivalently $Rw \cong R/\langle p \rangle$.*

*Proof.* Write $\mu = pq$. Since $q \notin \mathrm{Ann}_R(M) = \langle \mu \rangle$, choose $v \in M$ with $w := qv \neq 0$.
Then $pw = \mu v = 0$, so $o(w) \mid p$. Since $w \neq 0$, $o(w)$ is nonunit; as $p$ is prime, $o(w) \sim p$.
*Structural Commentary*: Every prime in the global order appears in some one-dimensional primary layer.
*Key Insight*: Divide the order by one prime and apply the remaining scalar to an element it does not kill. □

## 2.6 Submodules of Finite Free PID Modules

**Perspective 2.6.1.** *Noetherian Modules over a PID*
Over a PID, $N \leq R^n$ implies $N$ free with $\mathrm{rk}_R(N) \leq n$; Noetherianity is the f.g. shadow.

**Theorem 2.6.2.** *Submodules of Finite Free PID Modules*
*Let $F$ be free over $R$, $\mathrm{rk}_R(F) = n < \infty$, and $N \leq F$. Then $\exists m \leq n$ such that $N \cong R^m$.*
*Consequently, for $\mu_R(M) := \min\{|X| : \langle\langle X \rangle\rangle_R = M\}$, $\mu_R(M) \leq n$ implies $S \leq M \Longrightarrow \mu_R(S) \leq n$.*

*Proof.* Identify $F \simeq R^n$. For $1 \leq k \leq n$, set $I_k := \{x_k : x = (x_1, \ldots, x_k, 0, \ldots, 0) \in N\} \lhd R$ and write $I_k = \langle r_k \rangle$..
If $r_k \neq 0$, choose $u_k \in N \cap (R^k \times \{0\}^{n-k})$ with $(u_k)_k = r_k$. Let $K = \{k : r_k \neq 0\}$.
If $\sum_{k \in K} c_k u_k = 0$ and $j = \max\{k : c_k \neq 0\}$, then the $j$th coordinate gives $c_j r_j = 0$, hence $c_j = 0$, contradiction.
Thus $\{u_k : k \in K\}$ is $R$-independent. Set $\ell(0) = 0$ and $\ell(x) = \max \mathrm{supp}(x)$ for $0 \neq x \in N$.
If $j = \ell(x) > 0$, then $x_j \in I_j = \langle r_j \rangle$, say $x_j = cr_j$, and $\ell(x - cu_j) < j$.
Induction on $\ell(x)$ gives $N = \bigoplus_{k \in K} Ru_k \cong R^{|K|}$, so $\mathrm{rk}_R(N) = |K| \leq n$.
If $\mu_R(M) \leq n$, choose $\pi : R^n \twoheadrightarrow M$. For $S \leq M$, $P := \pi^{-1}(S) \leq R^n$ has $P = \langle\langle v_1, \ldots, v_m \rangle\rangle_R$ with $m \leq n$.
Hence $S = \langle\langle \pi(v_1), \ldots, \pi(v_m) \rangle\rangle_R$ and $\mu_R(S) \leq m \leq n$.
*Structural Commentary*: PIDs let one choose pivot generators in a submodule of $R^n$.
*Key Insight*: Pivot ideals $I_k$ yield triangular free generators. □

**Corollary 2.6.3.** *Finitely Generated PID Modules Are Noetherian*
*If $\mu_R(M) < \infty$, then $M$ is Noetherian.*

*Proof.* Let $n = \mu_R(M)$. By Theorem 2.6.2, $N \leq M \Longrightarrow \mu_R(N) \leq n < \infty$.
Thus every submodule of $M$ is f.g.; Theorem 2.4.2 gives $M$ Noetherian.
*Structural Commentary*: Finite generation of every submodule prevents infinite strict growth.
*Key Insight*: Generators of the union appear at a finite stage. □

**Proposition 2.6.4.** *A Full-Size Spanning Set Is a Basis*
*Let* $\mathrm{rk}_R(F) = n < \infty$. *If* $F = \langle\langle s_1, \dots, s_n \rangle\rangle_R$, *then* $(s_1, \dots, s_n)$ *is an* $R$*-basis of* $F$.

*Proof.* Choose a basis $(b_i)_{i=1}^n$ and define $\tau \in \mathrm{End}_R(F)$ by $\tau(b_i) = s_i$. Then $\tau$ is onto.
Since $F$ is free, $\exists \rho : F \to F$ with $\tau\rho = \mathrm{id}_F$; hence $F = \ker\tau \oplus \mathrm{im}\,\rho$ and $\mathrm{im}\,\rho \cong F$.
By Theorem 2.6.2, $\ker\tau \cong R^r$ for some $r \le n$. Thus $R^n \cong F \cong \ker\tau \oplus F \cong R^{r+n}$, so $r = 0$ by Theorem 2.2.2.
Hence $\ker\tau = 0$, $\tau$ is an isomorphism, and $(s_i) = \tau(b_i)$ is a basis.
*Structural Commentary*: Surjective finite-free endomorphisms over a PID are injective.
*Key Insight*: Rank leaves no room for a split kernel. □

## 2.7 Smith Normal Form and Module Presentations

**Perspective 2.7.1.** *Smith Form as Diagonalized Relations*
For a presentation $R^n \xrightarrow{A} R^m \to M \to 0$, Smith form replaces $A$ by a diagonal relation matrix, decoupling finite relations into cyclic constraints.

**Theorem 2.7.2.** *Smith Normal Form over a PID*
*Let* $A \in \mathrm{Mat}(m, n, R)$, *viewed as* $R^n \to R^m$.
$\exists$ *invertible* $P \in \mathrm{Mat}(m, m, R)$ *and* $Q \in \mathrm{Mat}(n, n, R)$ *such that* $PAQ = \mathrm{diag}(d_1, \dots, d_r, 0, \dots, 0)$,
*where* $d_i \in R \setminus \{0\}$, $d_1 \mid \cdots \mid d_r$, *and* $r = \mathrm{rk}_R(\mathrm{im}\,A)$. *The ideals* $\langle d_i \rangle$ *are unique.*

*Proof.*

(i) *Existence*: If $A = 0$, done. Set $I(A) = \langle a_{ij} \rangle = \langle d \rangle$. Elementary operations move a Bezout combination equal to $d$ to position $(1,1)$. If a pivot $p \nmid a$, Bezout on $(p, a)$ replaces $p$ by $\gcd(p, a)$, so $\langle p \rangle$ strictly increases; ACC in a PID forces termination. Then $p$ divides every entry, so clear its row and column. Induct on the complementary submatrix. Finally replace each bad adjacent pair $(a, b)$ by $(\gcd(a, b), \mathrm{lcm}(a, b))$ until $d_1 \mid \cdots \mid d_r$.

(ii) *Uniqueness*: Let $\Delta_k(A)$ be the ideal of $k \times k$ minors and $\Delta_0(A) = R$. Matrix equivalence preserves each $\Delta_k$. In Smith form, $\Delta_k(A) = \langle d_1 \cdots d_k \rangle$ for $1 \le k \le r$, and $\Delta_k(A) = 0$ for $k > r$. Since $R$ is a domain, $(\Delta_k(A) : \Delta_{k-1}(A)) = \langle d_k \rangle$, so $d_k$ is determined up to associates.

*Structural Commentary*: Smith form preserves rank plus divisibility data.
*Key Insight*: Determinantal ideals encode invariant factors. □

**Theorem 2.7.3.** *Adapted Bases for Submodules of Free Modules*
*Let* $N \le F$, $\mathrm{rk}_R(F) = n$, *and* $\mathrm{rk}_R(N) = k$.
*Then* $\exists (v_i)_{i=1}^n \in \mathrm{Bas}_R(F)$ *and* $r_1, \dots, r_k \in R \setminus \{0\}$ *such that* $r_1 \mid \cdots \mid r_k$ *and* $(r_i v_i)_{i=1}^k \in \mathrm{Bas}_R(N)$.

*Proof.* Choose bases of $N$ and $F$, and let $A$ represent $\iota : N \hookrightarrow F$.
Smith form gives $PAQ = \mathrm{diag}(r_1, \dots, r_k, 0, \dots, 0)$ with $r_i \neq 0$ and $r_1 \mid \cdots \mid r_k$, since $\iota$ is injective.
The row operations give a basis $(v_i)_{i=1}^n$ of $F$ and the column operations give a basis $(w_i)_{i=1}^k$ of $N$ with $\iota(w_i) = r_i v_i$. Hence $(r_i v_i)_{i=1}^k = (w_i)_{i=1}^k$ is a basis of $N$.
*Structural Commentary*: Inclusions diagonalize after choosing bases.
*Key Insight*: Diagonal entries are the required rescalings. □

**Corollary 2.7.4.** *Smith Form Presents Finitely Generated PID Modules*
*Let* $R^n \xrightarrow{A} R^m \to M \to 0$ *be a finite presentation.*
*If* $A \sim \mathrm{diag}(d_1, \dots, d_r, 0, \dots, 0)$ *in Smith form.* $I = \{i : 1 \le i \le r,\ d_i \notin R^\times\}$, *then* $M \cong R^{m-r} \oplus \bigoplus_{i \in I} R/\langle d_i \rangle$.

*Proof.* Equivalence $A \sim PAQ$ changes bases in $R^m, R^n$, so $\mathrm{coker}\,A \cong \mathrm{coker}(PAQ)$.
For the diagonal map, $d_i \in R^\times$ gives 0, $d_i \notin R^\times$ gives $R/\langle d_i \rangle$, and each zero row gives $R$.
*Structural Commentary*: Smith form turns presentations into direct sums.
*Key Insight*: Diagonal relations create cyclic torsion summands, and zero rows create free summands. □

**Example 2.7.5.** *Smith Form for a Finite Abelian Group*
Let $M = \mathbb{Z}^2/\langle\langle (6,0), (0,10), (2,4) \rangle\rangle_{\mathbb{Z}}$, with relation matrix $A = \begin{bmatrix} 6 & 0 & 2 \\ 0 & 10 & 4 \end{bmatrix}$. Then $\Delta_1(A) = \langle 6, 10, 2, 4 \rangle = \langle 2 \rangle$ and $\Delta_2(A) = \langle 60, 24, -20 \rangle = \langle 4 \rangle$. Hence $d_1 \sim 2$, $d_1 d_2 \sim 4$, so $d_2 \sim 2$ and $M \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$.
*Key Insight*: Determinantal ideals read off cyclic summands.

## 2.8 Finitely Generated PID Modules

**Theorem 2.8.1.** *Finitely Generated Torsion-Free Modules Are Free*
*For a f.g. $R$-module $M$, $M$ is free $\iff$ $M$ is torsion-free.*

*Proof.*

($\Rightarrow$) If $M \cong R^n$, then $am = 0$ with $a \neq 0$ forces every coordinate of $m$ to be 0, since $R$ is a domain.

($\Leftarrow$) Let $M$ be torsion-free and $G = \{u_1, \dots, u_n\}$ generate $M$.
Choose maximal $R$-independent $S = \{v_1, \dots, v_k\} \subseteq G$. For $w \in G \setminus S$, dependence of $S \cup \{w\}$ gives $a_w \neq 0$ and $b_{w,i} \in R$ with $a_w w + \sum_i b_{w,i} v_i = 0$. Put $a = \prod_{w \in G \setminus S} a_w$. Then $aM \subseteq \langle\langle S \rangle\rangle_R$, so $aM$ is free by Theorem 2.6.2. Since $M$ is torsion-free, $M \xrightarrow{\sim} aM$, $m \mapsto am$; hence $M$ is free.

*Structural Commentary*: Clear denominators into a free submodule.
*Key Insight*: One scalar embeds $M$ into a free sp an. □

**Theorem 2.8.2.** *Free Plus Torsion Decomposition*
*For f.g. $M$, $M = M_{\text{tor}} \oplus F$ with $F$ f.g. free. Moreover $M_{\text{tor}}$ is intrinsic and $F \cong M/M_{\text{tor}}$.*

*Proof.* By Proposition 1.6.3, $T := M_{\text{tor}} \leq M$ and $Q := M/T$ is f.g. torsion-free.
Thus $Q$ is free by Theorem 2.8.1. Since $Q$ is free, the quotient map $\pi : M \twoheadrightarrow Q$ splits; hence $M = T \oplus \operatorname{im}\sigma$ for any section $\sigma : Q \to M$, and $\operatorname{im}\sigma \cong Q$. If $M = T' \oplus G$ with $T'$ torsion and $G$ free, then $m = t' + g \in M_{\text{tor}}$ forces $g = 0$, so $T' = M_{\text{tor}}$. Every complement is isomorphic to $M/M_{\text{tor}}$.
*Structural Commentary*: F.g. PID modules split into torsion plus free.
*Key Insight*: The quotient by torsion is free, and freeness makes the quotient map split. □

## 2.9 Primary Decomposition and Cyclic Primary Modules

**Definition 2.9.1.** *Primary Module*
Let $p \in R$ be prime. A torsion $R$-module $M$ is *$p$-primary* if $o(M) \sim p^e$ for some $e \geq 1$.

**Theorem 2.9.2.** *Primary Decomposition*
*Let $M$ be f.g. torsion with $o(M) = \mu \sim \prod_{i=1}^{n} p_i^{e_i}$, where the $p_i$ are distinct nonassociate primes.*
*Put $q_i := p_i^{e_i}$, $\mu_i := \mu/q_i$, and $M_i := \mu_i M$. Then $M = \bigoplus_{i=1}^{n} M_i$, $M_i = \{v \in M : q_i v = 0\}$, and $o(M_i) \sim q_i$.*
*The decomposition is unique up to ordering.*

*Proof.* Since $\gcd(\mu_1, \dots, \mu_n) \sim 1$, choose $a_i \in R$ with $\sum_i a_i \mu_i = 1$. Hence $x = \sum_i a_i \mu_i x$, so $M = \sum_i M_i$.
Also $q_i M_i = \mu M = 0$. Conversely, if $q_i v = 0$, choose $a, b \in R$ with $a\mu_i + bq_i = 1$; then $v = a\mu_i v \in M_i$.
Thus $M_i = \{v : q_i v = 0\}$. If $\sum_i x_i = 0$ with $x_i \in M_i$, applying $\mu_i$ gives $\mu_i x_i = 0$ as $q_j \mid \mu_i$ for $j \neq i$.
Since $q_i x_i = 0$ and $\gcd(q_i, \mu_i) \sim 1$, Bezout gives $x_i = 0$; hence the sum is direct.
Now $o(M_i) \mid q_i$ and $\operatorname{lcm}_i o(M_i) \sim o(M) \sim \prod_i q_i$, so $o(M_i) \sim q_i$.
Finally each summand is the intrinsic set $\{v : q_i v = 0\}$, where $q_i$ is the $p_i$-part of $o(M)$.
*Structural Commentary*: Primary decomposition separates a torsion module prime by prime.
*Key Insight*: Bezout idempotents built from the coprime factors of the order project onto the primary pieces. □

**Lemma 2.9.3.** *The First $p$-Torsion Layer*
*Let $p \in R$ be prime and $M[p] := \ker(p : M \to M) = \{v \in M : pv = 0\}$.*
*Then $M[p]$ is an $R/\langle p \rangle$-vector space. If $M = S \oplus T$, then $M[p] = S[p] \oplus T[p]$.*

*Proof.* $\langle p \rangle$ is maximal by Theorem 3.11.5 and Theorem 3.4.4; hence $\kappa := R/\langle p \rangle$ is a field. Define $(r + \langle p \rangle)v := rv$.
If $r - r' \in \langle p \rangle$ and $v \in M[p]$, then $(r - r')v = 0$, so the action is well-defined; the axioms are inherited from $M$.
If $v = s + t \in M[p]$ under $M = S \oplus T$, then $0 = pv = ps + pt$ gives $ps = pt = 0$ by directness.
Thus $v \in S[p] \oplus T[p]$, and the reverse inclusion is immediate.
*Structural Commentary*: Killing by $p$ turns scalar multiplication modulo $p$ into field scalar multiplication.
*Key Insight*: The layer $M[p]$ counts the number of cyclic primary summands. □

**Theorem 2.9.4.** *Cyclic Decomposition of a Primary Module*
*Let $M$ be f.g. $p$-primary torsion with $o(M) \sim p^e$.*
*Then $M = \bigoplus_{i=1}^{m} \langle\langle v_i \rangle\rangle_R$ with $\operatorname{Ann}_R(v_i) = \langle p^{e_i} \rangle$ and $e = e_1 \geq e_2 \geq \cdots \geq e_m \geq 1$.*
*The annihilator chain $\langle p^{e_1} \rangle \subseteq \cdots \subseteq \langle p^{e_m} \rangle$ is intrinsic.*

*Proof.*

(i) *Existence*: Choose a finite presentation of $M$.
Corollary 2.7.4 gives $M \cong R^s \oplus \bigoplus_{\ell=1}^{t} R/\langle d_\ell \rangle$, with $d_\ell$ nonunit. Since $M$ is torsion, $s = 0$.
Since $M$ is $p$-primary, $d_\ell \sim p^{e_\ell}$ with $e_\ell \geq 1$. Reorder so $e_1 \geq \cdots \geq e_m$.
Proposition 2.5.4 gives $o(M) \sim p^{e_1}$, hence $e_1 = e$.

(ii) *Uniqueness*: Put $\kappa = R/\langle p \rangle$. By Lemma 2.9.3, each summand $R/\langle p^{e_i} \rangle$ contributes one $\kappa$-dimension to $M[p]$, so $\dim_\kappa M[p] = m$. Passing from $M$ to $pM$ replaces each $e_i > 1$ by $e_i - 1$ and removes each $e_i = 1$. Induction on $e$ recovers the exponent multiset, hence the annihilator chain.

*Structural Commentary*: A primary module splits into cyclic strings whose lengths are the exponents $e_i$.
*Key Insight*: The dimensions of the layers $M[p], pM[p], p^2M[p], \ldots$ determine the string lengths. □

**Proposition 2.9.5.** *Layer Counts Recover Primary Exponents*
*Let $\kappa = R/\langle p \rangle$ and $\lambda_k(M) := \dim_\kappa(p^{k-1}M/p^kM)$ for $k \geq 1$.*
*If $M \cong \bigoplus_{i=1}^{m} R/\langle p^{e_i} \rangle$ with $e_1 \geq \cdots \geq e_m \geq 1$, then $\lambda_k(M) = \#\{i : e_i \geq k\}$.*
*Hence the number of summands of exact exponent $k$ is $\lambda_k(M) - \lambda_{k+1}(M)$.*

*Proof.* For $C = R/\langle p^e \rangle$, $p^{k-1}C/p^kC$ is 0 if $k > e$ and is $\kappa$ if $k \leq e$, generated by the class of $p^{k-1}$.
Since direct sums commute with $p^{k-1}M$ and $p^kM$, dimensions add, giving $\lambda_k(M) = \#\{i : e_i \geq k\}$.
Thus $\lambda_k - \lambda_{k+1}$ counts exactly the summands ending at exponent $k$.
*Structural Commentary*: Successive $p$-layers determine the exponent list.
*Key Insight*: A cyclic summand of order $p^e$ contributes one dimension to exactly the first $e$ layers. □

## 2.10 Elementary Divisors and Invariant Factors

**Definition 2.10.1.** *Elementary Divisors*
For finitely generated torsion $M$, if its primary cyclic summands have annihilators $\langle p_i^{e_{i,j}} \rangle$,
set $\mathrm{ElemDiv}(M) := \{p_i^{e_{i,j}}\}_{i,j}$ as a multiset, up to associates.

**Theorem 2.10.2.** *Primary Cyclic Decomposition*
*Every f.g. torsion $R$-module $M$ admits $M = \bigoplus_i \bigoplus_{j=1}^{k_i} \langle\langle v_{i,j} \rangle\rangle_R$, where $\mathrm{Ann}_R(v_{i,j}) = \langle p_i^{e_{i,j}} \rangle$, $p_i$ are distinct nonassociate primes, and $e_{i,1} \geq \cdots \geq e_{i,k_i} \geq 1$. For f.g. torsion $M, N$, $M \cong N \iff \mathrm{ElemDiv}(M) = \mathrm{ElemDiv}(N)$.*

*Proof.*

($\Rightarrow$) Theorem 2.9.2 gives $M = \bigoplus_i M_{p_i}$ with each $M_{p_i}$ f.g. $p_i$-primary. Theorem 2.9.4 gives $M_{p_i} = \bigoplus_{j=1}^{k_i} \langle\langle v_{i,j} \rangle\rangle_R$ with $\mathrm{Ann}_R(v_{i,j}) = \langle p_i^{e_{i,j}} \rangle$. If $M \cong N$, the same uniqueness statements identify the primary pieces and their exponent chains; hence $\mathrm{ElemDiv}(M) = \mathrm{ElemDiv}(N)$.

($\Leftarrow$) If $\mathrm{ElemDiv}(M) = \mathrm{ElemDiv}(N)$, then both modules are isomorphic to the same direct sum $\bigoplus_{q \in \mathrm{ElemDiv}(M)} R/\langle q \rangle$, since a cyclic summand with annihilator $\langle q \rangle$ is $R/\langle q \rangle$ by Theorem 2.5.5. Thus $M \cong N$.

*Structural Commentary*: A torsion module is a direct sum of prime-power cyclic blocks.
*Key Insight*: Elementary divisors are the atoms of the PID module classification. □

**Proposition 2.10.3.** *Elementary Divisors of a Direct Sum*
*For f.g. torsion $A, B$, $\mathrm{ElemDiv}(A \oplus B) = \mathrm{ElemDiv}(A) \uplus \mathrm{ElemDiv}(B)$, where $\uplus$ is multiset union.*

*Proof.* If $A = \bigoplus_i A_i$ and $B = \bigoplus_j B_j$ are primary cyclic decompositions, then $A \oplus B = (\bigoplus_i A_i) \oplus (\bigoplus_j B_j)$ is one for $A \oplus B$. Thus the prime-power annihilator multiset is the multiset union.
*Structural Commentary*: Direct sums concatenate cyclic primary blocks.
*Key Insight*: Elementary divisors count blocks with multiplicity. □

**Definition 2.10.4.** *Invariant Factors*
For f.g. torsion $M$, write $\mathrm{ElemDiv}(M) = \{p_i^{e_{i,j}} : 1 \leq j \leq k_i\}_i$ with $e_{i,1} \geq \cdots \geq e_{i,k_i} \geq 1$.
Let $m = \max_i k_i$, set $e_{i,j} = 0$ for $j > k_i$, and define $d_j := \prod_i p_i^{e_{i,j}}$.
The nonunit $d_j$, up to associates, are the *invariant factors*; they satisfy $d_m \mid \cdots \mid d_1$.

**Method 2.10.5.** *From Elementary Divisors to Invariant Factors*

(i) Group $\mathrm{ElemDiv}(M)$ by prime: $\{p_i^{e_{i,j}}\}_j$.
(ii) Reorder each list so $e_{i,1} \geq e_{i,2} \geq \cdots$ and pad by $e_{i,j} = 0$ to common length.

(iii) Form $d_j = \prod_i p_i^{e_{i,j}}$.
(iv) Discard unit $d_j$; the rest are the invariant factors from Definition 2.10.4.

**Example 2.10.6.** *Packing Elementary Divisors into Invariant Factors*
Over $R = \mathbb{F}[z]$, let $\mathrm{ElemDiv}(M) = \{(z-1)^3, z-1, (z^2+1)^2, z^2+1\}$. The exponent rows are $(3,1)$ for $z-1$ and $(2,1)$ for $z^2+1$, so $d_1 = (z-1)^3(z^2+1)^2$ and $d_2 = (z-1)(z^2+1)$, with $d_2 \mid d_1$. Hence $M \cong R/\langle d_1\rangle \oplus R/\langle d_2\rangle$.
*Structural Commentary*: Elementary divisors separate primes; invariant factors pack rows.
*Key Insight*: Line up equal-height primary strings and multiply across the row.

**Example 2.10.7.** *Recombining Elementary Divisors*
Let $G = \mathbb{Z}/12\mathbb{Z} \oplus \mathbb{Z}/18\mathbb{Z}$. Then $\mathrm{ElemDiv}(G) = \{2, 2^2, 3, 3^2\}$. Recombining equal-height primary blocks gives $2 \cdot 3 = 6$ and $2^2 \cdot 3^2 = 36$; hence $G \cong \mathbb{Z}/6\mathbb{Z} \oplus \mathbb{Z}/36\mathbb{Z}$, with $6 \mid 36$.
*Structural Commentary*: Elementary divisors split by primes; invariant factors recombine them.
*Key Insight*: Prime-wise data equals one divisibility chain.

**Theorem 2.10.8.** *Invariant Factor Decomposition*
*For f.g. torsion $M$, $M \cong \bigoplus_{j=1}^m R/\langle d_j\rangle$ with $d_j \notin R^\times$ and $d_m \mid \cdots \mid d_1$.*
*Writing $\mathrm{Inv}(M) := (d_m, \ldots, d_1)$ up to associates, for f.g. torsion $M, N$, $M \cong N \iff \mathrm{Inv}(M) = \mathrm{Inv}(N)$.*

*Proof.* By Theorem 2.10.2, $M \cong \bigoplus_{q \in \mathrm{ElemDiv}(M)} R/\langle q\rangle$.
Let $\mathrm{ElemDiv}(M) = \{p_i^{e_{i,j}} : 1 \le j \le k_i\}_i$ with $e_{i,1} \ge \cdots \ge e_{i,k_i}$, pad by $e_{i,j} = 0$, and set $d_j = \prod_i p_i^{e_{i,j}}$.
For fixed $j$, the nonunit $p_i^{e_{i,j}}$ are pairwise coprime, so Theorem 2.5.6 gives $\bigoplus_{e_{i,j}>0} R/\langle p_i^{e_{i,j}}\rangle \cong R/\langle d_j\rangle$.
Hence $M \cong \bigoplus_j R/\langle d_j\rangle$ with $d_m \mid \cdots \mid d_1$.
($\Rightarrow$) If $M \cong N$, then $\mathrm{ElemDiv}(M) = \mathrm{ElemDiv}(N)$ by Theorem 2.10.2.
Grouping by height gives $\mathrm{Inv}(M) = \mathrm{Inv}(N)$.
($\Leftarrow$) If $\mathrm{Inv}(M) = \mathrm{Inv}(N)$, by factorization of each $d_j$, $\mathrm{ElemDiv}(M) = \mathrm{ElemDiv}(N)$; QED (Theorem 2.10.2).
*Structural Commentary*: Invariant factors pack prime-power blocks.
*Key Insight*: Same data, different packaging. □

**Corollary 2.10.9.** *Classification of Finitely Generated PID Modules*
*For f.g. $M$, $M \cong R^r \oplus \bigoplus_{j=1}^m R/\langle d_j\rangle$, where $r \ge 0$, $d_j \notin R^\times$, and $d_m \mid \cdots \mid d_1$.*
*The pair $(r, \mathrm{Inv}(M_{\mathrm{tor}}))$, equivalently $(r, \mathrm{ElemDiv}(M_{\mathrm{tor}}))$, is a complete invariant.*

*Proof.* Let $T := M_{\mathrm{tor}}$. Theorem 2.8.2 gives $M = T \oplus F$ with $F \cong R^r$.
Theorem 2.10.8 gives $T \cong \bigoplus_{j=1}^m R/\langle d_j\rangle$ with $d_m \mid \cdots \mid d_1$. Since $T$ is intrinsic and $F \cong M/T$, uniqueness and completeness follow from rank uniqueness and the invariant-factor theorem.
*Structural Commentary*: F.g. PID modules are free plus cyclic torsion.
*Key Insight*: The free rank records the zero-annihilator part; invariant factors record the torsion part. □

**Method 2.10.10.** *Reading a Module from Smith Form*
For finite presentation $R^n \xrightarrow{A} R^m \to M \to 0$ over PID, put $A \sim D = \mathrm{diag}(d_1, \ldots, d_r, 0, \ldots, 0)$, $d_1 \mid \cdots \mid d_r$:
(i) $d_i \in R^\times \implies$ no summand;
(ii) $d_i \notin R^\times \implies R/\langle d_i\rangle$;
(iii) each zero target coordinate contributes one $R$;
(iv) factor nonunit $d_i$ into prime powers for elementary divisors.
*Key Insight*: Smith form changes arbitrary relations into independent one-coordinate relations.

**Perspective 2.10.11.** *Largest Invariant Factor and Product*
If $M \cong \bigoplus_{j=1}^m R/\langle d_j\rangle$ with $d_m \mid \cdots \mid d_1$, then $\mathrm{Ann}_R(M) = \langle d_1\rangle$ and $\prod_{j=1}^m d_j \sim \prod_{q \in \mathrm{ElemDiv}(M)} q$.

**Theorem 2.10.12.** *Characterizing Cyclic Torsion Modules*
*Let $M$ be f.g. torsion with order $\mu \sim \prod_{i=1}^n p_i^{e_i}$, $p_i$ distinct nonassociate primes. These are equivalent:*
*(i) $M$ is cyclic.*
*(ii) $\forall i$, the primary component $M_{p_i}$ is cyclic.*
*(iii)* $\mathrm{ElemDiv}(M) = \{p_i^{e_i} : 1 \le i \le n\}$.
*(iv)* $M \cong R/\langle \mu\rangle$.

*Proof.*
(i) $\Rightarrow$ (ii) If $M = \langle\!\langle x\rangle\!\rangle_R$, Theorem 2.5.6 gives $M = \bigoplus_i \langle\!\langle x_i\rangle\!\rangle_R$ with $o(x_i) \sim p_i^{e_i}$; hence $M_{p_i} = \langle\!\langle x_i\rangle\!\rangle_R$.
(ii) $\Rightarrow$ (iii) By Theorem 2.10.2, cyclic $M_{p_i}$ of order $p_i^{e_i}$ has sole elementary divisor $p_i^{e_i}$; use Proposition 2.10.3.
(iii) $\Rightarrow$ (iv) By Theorem 2.10.8, there is one invariant factor $d_1 \sim \prod_i p_i^{e_i} \sim \mu$.

(*iv*) $\Rightarrow$ (*i*) Since $R/\langle\mu\rangle = \langle\!\langle 1+\langle\mu\rangle\rangle\!\rangle_R$, any isomorphism $R/\langle\mu\rangle \xrightarrow{\sim} M$ sends $1+\langle\mu\rangle$ to a generator of $M$.

*Structural Commentary*: Cyclic torsion means one string per prime.
*Key Insight*: Multiple elementary divisors for the same prime are the obstruction to one generator. □

**Definition 2.10.13.** *Indecomposable Module*
A nonzero $R$-module $M$ is *indecomposable* if $M = A \oplus B \Longrightarrow A = 0$ or $B = 0$.

**Theorem 2.10.14.** *Indecomposable Finitely Generated Torsion Modules*
*Let $0 \neq M$ be f.g. torsion. The following are equivalent:*

*(i) $M$ is indecomposable.*
*(ii) $M$ is primary cyclic.*
*(iii)* $\mathrm{ElemDiv}(M) = \{p^e\}$ *for some prime $p$ and $e \geq 1$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) By Theorem 2.10.2, write $M \cong \bigoplus_{\alpha\in A} C_\alpha$, where $0 \neq C_\alpha \cong R/\langle p_\alpha^{e_\alpha}\rangle$.
If $|A| \geq 2$, then $M \cong C_{\alpha_0} \oplus \bigoplus_{\alpha\neq\alpha_0} C_\alpha$ is a nontrivial decomposition, contradiction.
Since $M \neq 0$, $|A| = 1$; hence $M$ is primary cyclic.
(*ii*) $\Rightarrow$ (*iii*) If $M \cong R/\langle p^e\rangle$, then $\mathrm{ElemDiv}(M) = \{p^e\}$.
(*iii*) $\Rightarrow$ (*i*) If $M = A \oplus B$, Proposition 2.10.3 gives $\{p^e\} = \mathrm{ElemDiv}(A) \sqcup \mathrm{ElemDiv}(B)$.
Hence one multiset is empty, so one summand is 0.

*Structural Commentary*: Indecomposable torsion modules are precisely the single primary cyclic blocks.
*Key Insight*: Elementary divisors are the indecomposable pieces. □

# 3 Linear Operators as Modules

## 3.1 Polynomial Modules from Operators

Throughout this subsection, $\mathcal{V}$ is finite-dimensional over $\mathbb{F}$ unless explicitly stated otherwise.

**Perspective 3.1.1.** *An Operator as a Polynomial Action*
Fix $T \in \mathcal{L}(\mathcal{V})$. Evaluation at $T$ defines an $\mathbb{F}[z]$-action on $\mathcal{V}$ by $p \cdot \mathbf{v} := p(T)\mathbf{v}$.
Thus the orbit $\mathbf{v}, T\mathbf{v}, T^2\mathbf{v}, \ldots$ is encoded by the cyclic subspace $\{p(T)\mathbf{v} : p \in \mathbb{F}[z]\} = \mathrm{span}\{T^j\mathbf{v} : j \geq 0\}$.

**Definition 3.1.2.** *The Module Attached to an Operator*
Let $T \in \mathcal{L}(\mathcal{V})$. The *operator module* attached to $T$ is the $\mathbb{F}[z]$-module $\mathcal{V}_T$ with underlying additive group $\mathcal{V}$ and action $p(z) \cdot \mathbf{v} := p(T)\mathbf{v}$. For $A \in \mathrm{Mat}(n, n, \mathbb{F})$, define $(\mathbb{F}^n)_A$ analogously by $p(z) \cdot \mathbf{x} := p(A)\mathbf{x}$.

**Proposition 3.1.3.** *Operator Modules Are Finitely Generated Torsion Modules*
*Let $0 < \dim \mathcal{V} < \infty$ and $T \in \mathcal{L}(\mathcal{V})$. Then $\mathcal{V}_T$ is a finitely generated torsion $\mathbb{F}[z]$-module.*

*Proof.* Let $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be an $\mathbb{F}$-basis of $\mathcal{V}$. Since $\mathbb{F} \subseteq \mathbb{F}[z]$, this basis generates $\mathcal{V}_T$.
By Proposition 1.1.6, there exists $0 \neq p \in \mathbb{F}[z]$ with $p(T) = 0$. Thus $p \cdot \mathbf{v} = p(T)\mathbf{v} = \mathbf{0}$ for all $\mathbf{v} \in \mathcal{V}$, so $\mathcal{V}_T$ is torsion. Finally, $\mathbb{F}[z]$ is a PID by Theorem 3.11.1.
*Structural Commentary*: Finite-dimensionality gives a global polynomial relation for the operator.
*Key Insight*: Rational canonical form is the coordinate form of the structure theorem for $\mathcal{V}_T$. □

**Theorem 3.1.4.** *Submodules Are Invariant Subspaces*
*Let $T \in \mathcal{L}(\mathcal{V})$ and $\mathcal{S} \subseteq \mathcal{V}$. Then $\mathcal{S} \leq_{\mathbb{F}[z]} \mathcal{V}_T$ if and only if $\mathcal{S} \leq_{\mathbb{F}} \mathcal{V}$ and $T\mathcal{S} \subseteq \mathcal{S}$.*

*Proof.*

($\Rightarrow$) If $\mathcal{S} \leq_{\mathbb{F}[z]} \mathcal{V}_T$, then $\mathbb{F} \subseteq \mathbb{F}[z]$ gives $\mathcal{S} \leq_{\mathbb{F}} \mathcal{V}$. Since $z \cdot \mathbf{s} = T\mathbf{s}$ for $\mathbf{s} \in \mathcal{S}$, we have $T\mathcal{S} \subseteq \mathcal{S}$.
($\Leftarrow$) If $\mathcal{S} \leq_{\mathbb{F}} \mathcal{V}$ and $T\mathcal{S} \subseteq \mathcal{S}$, then $T^j\mathcal{S} \subseteq \mathcal{S}$ for all $j \geq 0$.
Hence for $p(z) = \sum_j a_j z^j$ and $\mathbf{s} \in \mathcal{S}$, $p \cdot \mathbf{s} = p(T)\mathbf{s} = \sum_j a_j T^j\mathbf{s} \in \mathcal{S}$, so $\mathcal{S} \leq_{\mathbb{F}[z]} \mathcal{V}_T$.

*Structural Commentary*: The single scalar $z$ remembers the operator.
*Key Insight*: Invariance under $T$ is exactly closure under all polynomial scalars. □

**Theorem 3.1.5.** *Similarity Is Module Isomorphism*
*Let $T, S \in \mathcal{L}(\mathcal{V})$. Then $\mathcal{V}_T \cong_{\mathbb{F}[z]} \mathcal{V}_S \Longleftrightarrow T \sim S$. More precisely, for $\Phi \in \mathrm{GL}(\mathcal{V})$, $\Phi : \mathcal{V}_T \to \mathcal{V}_S$ is $\mathbb{F}[z]$-linear if and only if $\Phi T = S\Phi$, equivalently $S = \Phi T \Phi^{-1}$.*

*Proof.*

($\Rightarrow$) If $\Phi : \mathcal{V}_T \to \mathcal{V}_S$ is an $\mathbb{F}[z]$-isomorphism, then $\Phi \in \mathrm{GL}(\mathcal{V})$, $\Phi(T\mathbf{v}) = \Phi(z \cdot \mathbf{v}) = z \cdot \Phi(\mathbf{v}) = S\Phi(\mathbf{v})$ $\forall \mathbf{v} \in \mathcal{V}$.
Hence $\Phi T = S\Phi$, so $S = \Phi T \Phi^{-1}$.

($\Leftarrow$) If $\Phi \in \mathrm{GL}(\mathcal{V})$ and $S = \Phi T \Phi^{-1}$, then $\Phi T = S\Phi$, hence $\Phi T^j = S^j \Phi$ for all $j \geq 0$.
Thus for $p(z) = \sum_j a_j z^j$, $\Phi(p(T)\mathbf{v}) = p(S)\Phi(\mathbf{v})$, so $\Phi : \mathcal{V}_T \to \mathcal{V}_S$ is an $\mathbb{F}[z]$-isomorphism.

*Structural Commentary*: A module isomorphism is precisely an intertwiner for the action of $z$.
*Key Insight*: Similarity is isomorphism of the associated $\mathbb{F}[z]$-modules. □

**Corollary 3.1.6.** *Matrix Similarity Is Matrix-Module Isomorphism*
*Let $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$. Then $A \sim B \iff (\mathbb{F}^n)_A \cong_{\mathbb{F}[z]} (\mathbb{F}^n)_B$. More precisely, for $X \in \mathrm{GL}_n(\mathbb{F})$, $X : (\mathbb{F}^n)_A \to (\mathbb{F}^n)_B$ is $\mathbb{F}[z]$-linear if and only if $XA = BX$, equivalently $B = XAX^{-1}$.*

*Proof.*

($\Rightarrow$) If $X : (\mathbb{F}^n)_A \to (\mathbb{F}^n)_B$ is an $\mathbb{F}[z]$-isomorphism, then $X \in \mathrm{GL}_n(\mathbb{F})$ and $X(A\mathbf{x}) = X(z \cdot \mathbf{x}) = z \cdot X\mathbf{x} = BX\mathbf{x}$ for all $\mathbf{x} \in \mathbb{F}^n$. Hence $XA = BX$, so $B = XAX^{-1}$.
($\Leftarrow$) If $X \in \mathrm{GL}_n(\mathbb{F})$ and $B = XAX^{-1}$, then $XA = BX$, hence $XA^j = B^j X$ for all $j \geq 0$. Thus $Xp(A) = p(B)X$ for all $p \in \mathbb{F}[z]$, so $X : (\mathbb{F}^n)_A \to (\mathbb{F}^n)_B$ is an $\mathbb{F}[z]$-isomorphism.

*Structural Commentary*: Similarity is invertible module intertwining.
*Key Insight*: The matrix is remembered by the scalar action of $z$. □

**Definition 3.1.7.** *Module Annihilator and Minimal Polynomial*

(i) The *annihilator* of $\mathcal{V}_T$ is $\mathrm{Ann}_{\mathbb{F}[z]}(\mathcal{V}_T) := \{p \in \mathbb{F}[z] : p(T)\mathcal{V} = \{\mathbf{0}\}\}$.
(ii) *Minimal polynomial* $\mu_T$ is unique monic generator of this nonzero principal ideal, so $\mathrm{Ann}_{\mathbb{F}[z]}(\mathcal{V}_T) = \langle \mu_T \rangle$.
(iii) For $A \in \mathrm{Mat}(n, n, \mathbb{F})$, define $\mu_A := \mu_{T_A}$, where $T_A(\mathbf{x}) = A\mathbf{x}$.

**Proposition 3.1.8.** *Minimal Polynomial as Module Order*
*Let $T \in \mathcal{L}(\mathcal{V})$. Then $\mu_T$ is the unique monic polynomial of minimal degree in $\{p \in \mathbb{F}[z] : p(T) = 0\}$. If $S \sim T$, then $\mu_S = \mu_T$; hence similar matrices have the same minimal polynomial.*

*Proof.* By Definition 3.1.7, $\mathrm{Ann}_{\mathbb{F}[z]}(\mathcal{V}_T) = \langle \mu_T \rangle$ and $p \in \mathrm{Ann}_{\mathbb{F}[z]}(\mathcal{V}_T)$ iff $p(T) = 0$. Thus $\mu_T$ is the unique monic annihilator of minimal degree. If $S = \Phi T \Phi^{-1}$, then $p(S) = \Phi p(T) \Phi^{-1}$ for all $p \in \mathbb{F}[z]$, so $\mathrm{Ann}_{\mathbb{F}[z]}(\mathcal{V}_S) = \mathrm{Ann}_{\mathbb{F}[z]}(\mathcal{V}_T)$ and $\mu_S = \mu_T$. The matrix case follows by applying this to the associated operators.
*Structural Commentary*: The minimal polynomial is the order of the whole operator module.
*Key Insight*: Similarity preserves the module and therefore preserves its annihilator ideal. □

**Definition 3.1.9.** *Cyclic Subspace and Vector Annihilator*
Let $T \in \mathcal{L}(\mathcal{V})$ and $\mathbf{v} \in \mathcal{V}$.

(i) The *$T$-cyclic subspace generated by $\mathbf{v}$* is $\mathcal{C}_T(\mathbf{v}) := \mathrm{span}\{T^j\mathbf{v} : j \geq 0\}$
(ii) The vector $\mathbf{v}$ is a *cyclic vector* for $T$ if $\mathcal{C}_T(\mathbf{v}) = \mathcal{V}$.
(iii) The *annihilator of $\mathbf{v}$ for $T$* is $\mathrm{ann}_T(\mathbf{v}) := \{p \in \mathbb{F}[z] : p(T)\mathbf{v} = \mathbf{0}\}$

**Proposition 3.1.10.** *Cyclic Subspaces Are Smallest Invariant Hosts*
*Let $T \in \mathcal{L}(\mathcal{V})$ and $\mathbf{v} \in \mathcal{V}$. Then $\mathcal{C}_T(\mathbf{v}) = \{p(T)\mathbf{v} : p \in \mathbb{F}[z]\}$.*
*Moreover, $\mathcal{C}_T(\mathbf{v})$ is the smallest $T$-invariant subspace containing $\mathbf{v}$, and $\mathrm{ann}_T(\mathbf{v})$ is an ideal of $\mathbb{F}[z]$.*

*Proof.* Equality follows as $p(T)\mathbf{v}$ is a finite linear combination of the orbit vectors $T^j\mathbf{v}$, and every such finite linear combination equals $p(T)\mathbf{v}$ for some $p \in \mathbb{F}[z]$. Also $T(p(T)\mathbf{v}) = (zp)(T)\mathbf{v}$, so $\mathcal{C}_T(\mathbf{v})$ is $T$-invariant.
If $\mathcal{U}$ is $T$-invariant and contains $\mathbf{v}$, then $T^j\mathbf{v} \in \mathcal{U}$ for every $j \geq 0$, hence $\mathcal{C}_T(\mathbf{v}) \subseteq \mathcal{U}$. Finally, if $p, q \in \mathrm{ann}_T(\mathbf{v})$ and $r \in \mathbb{F}[z]$, then $(p+q)(T)\mathbf{v} = \mathbf{0}$ and $(rp)(T)\mathbf{v} = r(T)p(T)\mathbf{v} = \mathbf{0}$ by Proposition 1.1.5. $\mathrm{ann}_T(\mathbf{v})$ is an ideal.
*Structural Commentary*: A cyclic subspace is the span of the orbit generated by repeatedly applying $T$.
*Key Insight*: Invariance is exactly what lets the whole orbit stay inside the same host subspace. □




Figure 107: The $T$-orbit of $\mathbf{v}$ spans $\mathcal{C}_T(\mathbf{v})$.

**Definition 3.1.11.** *Order of a Vector under an Operator*
Let $\mathcal{V}$ be finite-dimensional, let $T \in \mathcal{L}(\mathcal{V})$, and let $\mathbf{v} \neq \mathbf{0}$. The *order of $\mathbf{v}$ under $T$* is the unique monic polynomial $m_{\mathbf{v}} \in \mathrm{ann}_T(\mathbf{v})$ of least degree; equivalently, $\mathrm{ann}_T(\mathbf{v}) = (m_{\mathbf{v}})$

**Definition 3.1.12.** *Companion Matrix*
Let $f(z) = z^n + c_{n-1}z^{n-1} + \cdots + c_1 z + c_0$ be monic. For $n \geq 2$, the *companion matrix* of $f$ is

$$C_f := \begin{bmatrix} 0 & 0 & \cdots & 0 & -c_0 \\ 1 & 0 & \cdots & 0 & -c_1 \\ 0 & 1 & \cdots & 0 & -c_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -c_{n-1} \end{bmatrix}.$$

For $n = 1$, set $C_f = [-c_0]$.

**Theorem 3.1.13.** *Companion Matrix Realises a Polynomial*
*Let $f(z) = z^n + \sum_{i=0}^{n-1} c_i z^i$ be monic and let $C_f$ be its companion matrix. Then $p_{C_f} = \mu_{C_f} = f$.*

*Proof.* Let $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be the standard basis.
For $0 \leq j < n$, $C_f^j \mathbf{e}_1 = \mathbf{e}_{j+1}$, and last column gives $C_f^n \mathbf{e}_1 = -\sum_{i=0}^{n-1} c_i C_f^i \mathbf{e}_1$. Hence $f(C_f)\mathbf{e}_1 = 0$.
Since $\mathbf{e}_j = C_f^{j-1}\mathbf{e}_1$ and $f(C_f)C_f = C_f f(C_f)$, $f(C_f)\mathbf{e}_j = 0\ \forall j$, so $f(C_f) = 0$.
If $0 \neq g(z) = \sum_{i=0}^{m} b_i z^i$ with $m < n$, then $g(C_f)\mathbf{e}_1 = \sum_{i=0}^{m} b_i \mathbf{e}_{i+1} \neq \mathbf{0}$.
Thus no nonzero polynomial of degree $< n$ annihilates $C_f$, so $\deg \mu_{C_f} = n$.
Since $\mu_{C_f} \mid f$ and both are monic of degree $n$, $\mu_{C_f} = f$.
By Cayley–Hamilton, $\mu_{C_f} \mid p_{C_f}$; as $p_{C_f}$ and $f$ are monic of degree $n$, $p_{C_f} = f$.
*Structural Commentary*: The cyclic vector $\mathbf{e}_1$ generates the basis, and last column imposes relation $f(C_f) = 0$.
*Key Insight*: The companion matrix makes $f$ the first possible annihilating relation. □

**Theorem 3.1.14.** *Cyclic Subspaces as Polynomial Quotients*
*Let $\dim \mathcal{V} < \infty$, $T \in \mathcal{L}(\mathcal{V})$, $0 \neq \mathbf{v} \in \mathcal{V}$, and let $m_{\mathbf{v}}(z) = z^d + \sum_{i=0}^{d-1} c_i z^i$ be the order of $\mathbf{v}$.*
*Define $\Phi_{\mathbf{v}} : \mathbb{F}[z] \to \mathcal{C}_T(\mathbf{v})$ by $\Phi_{\mathbf{v}}(p) = p(T)\mathbf{v}$ and $\mathcal{B}_{\mathbf{v}} := (\mathbf{v}, T\mathbf{v}, \ldots, T^{d-1}\mathbf{v})$. Then*
*(i) $\operatorname{im} \Phi_{\mathbf{v}} = \mathcal{C}_T(\mathbf{v})$, $\ker \Phi_{\mathbf{v}} = \langle m_{\mathbf{v}} \rangle$, and $\mathcal{C}_T(\mathbf{v}) \cong \mathbb{F}[z]/\langle m_{\mathbf{v}} \rangle$;*
*(ii) $\mathcal{B}_{\mathbf{v}}$ is a basis of $\mathcal{C}_T(\mathbf{v})$;*
*(iii) $[T|_{\mathcal{C}_T(\mathbf{v})}]_{\mathcal{B}_{\mathbf{v}}} = C_{m_{\mathbf{v}}}$.*

*Proof.*
(i) By Proposition 3.1.10, $\operatorname{im} \Phi_{\mathbf{v}} = \mathcal{C}_T(\mathbf{v})$. $\ker \Phi_{\mathbf{v}} = \operatorname{ann}_T(\mathbf{v}) = \langle m_{\mathbf{v}} \rangle$, by Theorem 2.2.12 $\mathcal{C}_T(\mathbf{v}) \cong \mathbb{F}[z]/\langle m_{\mathbf{v}} \rangle$.
(ii) If $\sum_{j=0}^{d-1} a_j T^j \mathbf{v} = \mathbf{0}$, then $q(z) = \sum_{j=0}^{d-1} a_j z^j \in \operatorname{ann}_T(\mathbf{v})$ and $\deg q < d$, hence $q = 0$; thus $\mathcal{B}_{\mathbf{v}}$ is independent.
For $p = hm_{\mathbf{v}} + r$ with $\deg r < d$ or $r = 0$, $p(T)\mathbf{v} = r(T)\mathbf{v} \in \operatorname{span} \mathcal{B}_{\mathbf{v}}$, so $\mathcal{B}_{\mathbf{v}}$ spans.
(iii) For $0 \leq j < d-1$, $T(T^j\mathbf{v}) = T^{j+1}\mathbf{v}$. Since $m_{\mathbf{v}}(T)\mathbf{v} = \mathbf{0}$, $T^d\mathbf{v} = -\sum_{i=0}^{d-1} c_i T^i \mathbf{v}$, last-column relation of $C_{m_{\mathbf{v}}}$.

*Structural Commentary*: The first relation cuts out the quotient and companion block.
*Key Insight*: One orbit carries the whole cyclic subspace. □

**Perspective 3.1.15.** *The Translation Dictionary*
The module vocabulary for $\mathcal{V}_T$ and the vector-space vocabulary for $T$ match as follows:

| $\mathbb{F}[z]$-module $\mathcal{V}_T$ | $\mathbb{F}$-vector space $\mathcal{V}$ |
|---|---|
| $p(z) \cdot \mathbf{v}$ | $p(T)\mathbf{v}$ |
| submodule | $T$-invariant subspace |
| $\operatorname{Ann}_{\mathbb{F}[z]}(\mathcal{V}_T) = \langle \mu_T \rangle$ | minimal polynomial of $T$ |
| $\langle\langle \mathbf{v} \rangle\rangle_{\mathbb{F}[z]}$ | $\mathcal{C}_T(\mathbf{v})$ |
| $\mathbb{F}[z]/\langle m_{\mathbf{v}} \rangle$ | cyclic block with companion matrix $C_{m_{\mathbf{v}}}$ |

*Key Insight*: Once $z$ is interpreted as $T$, module structure becomes operator structure.

## 3.2 Invariant Factors

**Definition 3.2.1.** *Invariant Factors and Elementary Divisors*
Let $T \in \mathcal{L}(\mathcal{V})$.
(i) A *cyclic decomposition* is $\mathcal{V} = \bigoplus_{j=1}^{r} \mathcal{C}_j$, where each $\mathcal{C}_j$ is nonzero, $T$-invariant, and cyclic.
(ii) $\operatorname{InvFact}(T) := (f_1, \ldots, f_r)$, where $f_j$ are monic nonconstant invariant factors of $\mathcal{V}_T$; $f_r \mid f_{r-1} \mid \cdots \mid f_1$.

(iii) $\mathrm{ElemDiv}(T)$ is the multiset of monic prime powers in the $f_j$, counted with multiplicity.
(iv) *Elementary divisor basis* is $\mathcal{B}_{\mathrm{ED}} := \bigsqcup_\alpha(\mathbf{v}_\alpha, T\mathbf{v}_\alpha, \dots, T^{d_\alpha-1}\mathbf{v}_\alpha)$, where $d_\alpha = \deg m_{\mathbf{v}_\alpha}$; $\{m_{\mathbf{v}_\alpha}\}_\alpha = \mathrm{ElemDiv}(T)$.
(v) For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, set $\mathrm{InvFact}(A) := \mathrm{InvFact}(T_A)$ and $\mathrm{ElemDiv}(A) := \mathrm{ElemDiv}(T_A)$.

**Theorem 3.2.2.** *Invariant-Factor Skeleton for Operators*
*Let* $0 < \dim \mathcal{V} < \infty$, $T \in \mathcal{L}(\mathcal{V})$, *and* $\mathrm{InvFact}(T) = (f_1, \dots, f_r)$.
*Then* $f_r \mid \cdots \mid f_1$ *and there are* $T$*-cyclic subspaces* $\mathcal{C}_j$ *such that* $\mathcal{V} = \bigoplus_{j=1}^r \mathcal{C}_j$, $\mathcal{C}_j \cong \mathbb{F}[z]/\langle f_j\rangle$, $[T] = \bigoplus_{j=1}^r C_{f_j}$ *in the concatenated cyclic bases,* $\mu_T = f_1$, *and* $p_T = \prod_{j=1}^r f_j$.

*Proof.* By Proposition 3.1.3, $\mathcal{V}_T$ is finitely generated torsion over $\mathbb{F}[z]$.
Theorem 2.10.8 gives $\mathcal{V}_T \cong \bigoplus_{j=1}^r \mathbb{F}[z]/\langle f_j\rangle$ with $f_r \mid \cdots \mid f_1$, uniquely.
Transport cyclic generators to $T$-cyclic subspaces $\mathcal{C}_j$; Theorem 3.1.14 gives $\mathcal{C}_j \cong \mathbb{F}[z]/\langle f_j\rangle$ and block $C_{f_j}$.
Since $\mathrm{Ann}(\mathcal{V}_T) = \langle \mathrm{lcm}(f_1, \dots, f_r)\rangle = \langle f_1\rangle = \langle \mu_T\rangle$, $f_1 = \mu_T$.
Finally, Theorem 3.1.13 gives $p_{C_{f_j}} = f_j$, so $p_T = \prod_{j=1}^r f_j$.
*Structural Commentary*: Torsion module decomposition produces companion blocks.
*Key Insight*: Invariant factors are the polynomial-module version of choosing canonical blocks for similarity. □

## 3.3 Elementary Divisors

**Theorem 3.3.1.** *Primary Cyclic Decomposition for an Operator*
*Let* $\mu_T = \prod_{i=1}^s p_i^{e_i}$ *with* $p_i \in \mathbb{F}[z]$ *distinct monic irreducible, and set* $\mathcal{V}_{p_i} := \ker p_i(T)^{e_i}$. *Then*
*(i)* $\mathcal{V}_T = \bigoplus_{i=1}^s \mathcal{V}_{p_i}$;
*(ii)* $\mathcal{V}_{p_i} = \bigoplus_{j=1}^{k_i} \mathcal{C}_T(\mathbf{v}_{i,j})$, *with* $m_{\mathbf{v}_{i,j}} = p_i^{e_{i,j}}$ *and* $e_i = e_{i,1} \geq e_{i,2} \geq \cdots \geq e_{i,k_i}$;
*(iii)* $\mathrm{ElemDiv}(T) = \{p_i^{e_{i,j}} : 1 \leq i \leq s,\ 1 \leq j \leq k_i\}$, *corresponding cyclic bases form elementary divisor basis.*

*Proof.*
(i) By Theorem 2.10.2, $\mathcal{V}_T = \bigoplus_i \mathcal{V}_T[p_i^\infty]$; $\mathcal{V}_T[p_i^\infty] := \{\mathbf{x} \in \mathcal{V} : \exists N \geq 1,\ p_i(T)^N\mathbf{x} = 0\} = \ker p_i(T)^{e_i} = \mathcal{V}_{p_i}$.
(ii) By Theorem 2.10.2, $\mathcal{V}_{p_i} = \bigoplus_{j=1}^{k_i} \mathcal{C}_T(\mathbf{v}_{i,j})$, $m_{\mathbf{v}_{i,j}} = p_i^{e_{i,j}}$ and $e_i = e_{i,1} \geq \cdots \geq e_{i,k_i}$.
(iii) Definition 3.2.1 gives $\mathrm{ElemDiv}(T) = \{p_i^{e_{i,j}} : 1 \leq i \leq s,\ 1 \leq j \leq k_i\}$, and concatenating the bases of $\mathcal{C}_T(\mathbf{v}_{i,j})$ gives an elementary divisor basis.
*Structural Commentary*: Primary module pieces are polynomial-kernel pieces for the operator.
*Key Insight*: Elementary divisors are exactly the orders of the cyclic primary summands. □

**Theorem 3.3.2.** *Reducing Decompositions Combine the Data*
*Let* $\mathcal{V} = \mathcal{A} \oplus \mathcal{B}$ *with* $\mathcal{A}, \mathcal{B}$ $T$*-invariant, and set* $T_\mathcal{A} := T|_\mathcal{A}$, $T_\mathcal{B} := T|_\mathcal{B}$.
*Then* $\mu_T = \mathrm{lcm}(\mu_{T_\mathcal{A}}, \mu_{T_\mathcal{B}})$ *and* $\mathrm{ElemDiv}(T) = \mathrm{ElemDiv}(T_\mathcal{A}) \sqcup \mathrm{ElemDiv}(T_\mathcal{B})$, *where* $\sqcup$ *is multiset union.*
*The invariant factors are obtained by recombining the elementary divisors into a divisibility chain.*

*Proof.* By Theorem 3.1.4, $\mathcal{V}_T = \mathcal{A}_{T_\mathcal{A}} \oplus \mathcal{B}_{T_\mathcal{B}}$ as $\mathbb{F}[z]$-modules.
Hence $\mathrm{Ann}(\mathcal{V}_T) = \mathrm{Ann}(\mathcal{A}_{T_\mathcal{A}}) \cap \mathrm{Ann}(\mathcal{B}_{T_\mathcal{B}}) = \langle \mathrm{lcm}(\mu_{T_\mathcal{A}}, \mu_{T_\mathcal{B}})\rangle$.
Proposition 2.10.3 gives elementary-divisor identity.
*Structural Commentary*: Reducing decompositions split the operator module.
*Key Insight*: Direct sums take lcm on global orders and multiset union on elementary divisors. □

**Definition 3.3.3.** *Characteristic Polynomial from Elementary Divisors*
For $T \in \mathcal{L}(\mathcal{V})$, define $\chi_T(z) := \prod_{q \in \mathrm{ElemDiv}(T)} q(z)$, with product over the multiset $\mathrm{ElemDiv}(T)$.
For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, set $\chi_A := \chi_{T_A}$.

**Theorem 3.3.4.** *Minimal and Characteristic Polynomials from Elementary Divisors*
*Let* $\mathrm{ElemDiv}(T) = \{p_i^{e_{i,j}} : 1 \leq i \leq s,\ 1 \leq j \leq k_i\}$.
*Then* $\mu_T = \prod_i p_i^{\max_j e_{i,j}}$, $\chi_T = \prod_{i,j} p_i^{e_{i,j}}$, *and* $\deg \chi_T = \dim \mathcal{V}$.
*Hence* $\mu_T \mid \chi_T$, *and* $\mu_T, \chi_T$ *have the same irreducible factors.*

*Proof.* By Theorem 3.3.1, $\mathcal{V}_T = \bigoplus_{i,j} \mathcal{C}_T(\mathbf{v}_{i,j})$ with $m_{\mathbf{v}_{i,j}} = p_i^{e_{i,j}}$.
Hence $\mathrm{Ann}(\mathcal{V}_T) = \langle \mathrm{lcm}_{i,j}\, p_i^{e_{i,j}}\rangle = \langle \prod_i p_i^{\max_j e_{i,j}}\rangle$, so $\mu_T = \prod_i p_i^{\max_j e_{i,j}}$. Definition 3.3.3 gives $\chi_T = \prod_{i,j} p_i^{e_{i,j}}$.
By Theorem 3.1.14, $\dim \mathcal{C}_T(\mathbf{v}_{i,j}) = \deg p_i^{e_{i,j}}$; summing gives $\deg \chi_T = \dim \mathcal{V}$.
Divisibility and common irreducible factors are immediate.
*Structural Commentary*: $\mu_T$ records maximal exponents; $\chi_T$ records all exponents.
*Key Insight*: The pair $(\mu_T, \chi_T)$ forgets the block distribution. □

**Definition 3.3.5.** *Nonderogatory Operators*
$T \in \mathcal{L}(\mathcal{V})$ is *nonderogatory* if $\mu_T = \chi_T$, equivalently $\deg \mu_T = \dim \mathcal{V}$.
For $A \in \mathrm{Mat}(n, n, \mathbb{F})$, $A$ is nonderogatory if $T_A$ is.

**Theorem 3.3.6.** *Cyclic and Nonderogatory Are the Same*
*For $T \in \mathcal{L}(\mathcal{V})$, the following are equivalent:*

*(i)* $\mathcal{V}_T$ *is cyclic;*
*(ii)* $T$ *has a cyclic vector;*
*(iii)* $\mathrm{ElemDiv}(T) = \{p_1^{e_1}, \dots, p_s^{e_s}\}$*, one elementary divisor for each irreducible factor of* $\mu_T$*;*
*(iv)* $T$ *is nonderogatory.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* If $\mathcal{V}_T = \mathbb{F}[z]\mathbf{v}$, then $\mathcal{V} = \{p(T)\mathbf{v} : p \in \mathbb{F}[z]\} = \mathcal{C}_T(\mathbf{v})$, so $\mathbf{v}$ is cyclic.
*(ii)* $\Rightarrow$ *(i)* If $\mathcal{C}_T(\mathbf{v}) = \mathcal{V}$, then $\mathcal{V}_T = \mathbb{F}[z]\mathbf{v}$.
*(i)* $\Leftrightarrow$ *(iii)* Theorem 2.10.12 on $\mathcal{V}_T$. Cyclic torsion module has one cyclic summand for each primary factor.
*(iii)* $\Rightarrow$ *(iv)* If $\mathrm{ElemDiv}(T) = \{p_i^{e_i}\}_{i=1}^s$, then $\chi_T = \prod_i p_i^{e_i} = \mu_T$, so $T$ is nonderogatory.
*(iv)* $\Rightarrow$ *(iii)* Let $\mathrm{ElemDiv}(T) = \{p_i^{e_{i,j}}\}$. If $\mu_T = \chi_T$, then $\prod_i p_i^{\max_j e_{i,j}} = \prod_{i,j} p_i^{e_{i,j}}$, so $k_i = 1$ for all $i$.

*Structural Commentary*: Nonderogatory means all primary information can be generated by one vector.
*Key Insight*: Equality $\mu_T = \chi_T$ says that product and lcm of the cyclic orders coincide. □

**Perspective 3.3.7.** *Elementary Divisors and Jordan Blocks*
Over $\mathbb{C}$, elementary divisors are exactly $(z-\lambda)^k$, and $\mathbb{C}[z]/\langle (z-\lambda)^k \rangle$ corresponds to the Jordan block $J_k(\lambda)$. Thus $\mathrm{ElemDiv}(T)$ is the multiset of Jordan blocks.

## 3.4 Operator Classification

**Perspective 3.4.1.** *Operator Modules and Similarity*
For $T \in \mathcal{L}(\mathcal{V})$, $\mathcal{V}_T$ has action $p \cdot \mathbf{v} = p(T)\mathbf{v}$. Then $S \sim T \iff \mathcal{V}_S \cong_{\mathbb{F}[z]} \mathcal{V}_T$; $\mathbb{F}[z]/\langle f \rangle \iff C_f$ (Theorem 3.1.14).

**Theorem 3.4.2.** *Complete Similarity Invariants*
*Let $T, S \in \mathcal{L}(\mathcal{V})$ and $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$.*

*(i)* $T \sim S \iff \mathcal{V}_T \cong_{\mathbb{F}[z]} \mathcal{V}_S \iff \mathrm{ElemDiv}(T) = \mathrm{ElemDiv}(S) \iff \mathrm{InvFact}(T) = \mathrm{InvFact}(S)$.
*(ii)* $A \sim B \iff (\mathbb{F}^n)_A \cong_{\mathbb{F}[z]} (\mathbb{F}^n)_B \iff \mathrm{ElemDiv}(A) = \mathrm{ElemDiv}(B) \iff \mathrm{InvFact}(A) = \mathrm{InvFact}(B)$.
*(iii) If $A = [T]_{\mathcal{B}}$ for some basis $\mathcal{B}$ of $\mathcal{V}$, then* $\mathrm{ElemDiv}(A) = \mathrm{ElemDiv}(T)$ *and* $\mathrm{InvFact}(A) = \mathrm{InvFact}(T)$.

*Proof.*

*(i)* $(\Rightarrow)$ If $T \sim S$, then $\mathcal{V}_T \cong_{\mathbb{F}[z]} \mathcal{V}_S$ by Theorem 3.1.5.
Module isomorphism preserves the PID decomposition, hence preserves ElemDiv and InvFact.
*(i)* $(\Leftarrow)$ Equality of either ElemDiv or InvFact gives $\mathcal{V}_T \cong_{\mathbb{F}[z]} \mathcal{V}_S$ by uniqueness of the PID decomposition.
Thus $T \sim S$ by Theorem 3.1.5.
*(ii)* $(\Rightarrow)$ If $A \sim B$, then $(\mathbb{F}^n)_A \cong_{\mathbb{F}[z]} (\mathbb{F}^n)_B$ by Corollary 3.1.6; hence the divisor data agree.
*(ii)* $(\Leftarrow)$ Equality of either matrix divisor datum gives $(\mathbb{F}^n)_A \cong_{\mathbb{F}[z]} (\mathbb{F}^n)_B$, so $A \sim B$ by Corollary 3.1.6.
*(iii)* If $A = [T]_{\mathcal{B}}$, then $\Phi(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}}$ satisfies $\Phi(p(T)\mathbf{v}) = p(A)\Phi(\mathbf{v})$.
Hence $\Phi : \mathcal{V}_T \to (\mathbb{F}^n)_A$ is an $\mathbb{F}[z]$-isomorphism, so the divisor data agree.

*Structural Commentary*: Similarity is module isomorphism plus PID data.
*Key Insight*: Elementary divisors classify operator modules. □

**Lemma 3.4.3.** *Determinant of a Companion Block*
*Let $f(z) = z^n + \sum_{i=0}^{n-1} c_i z^i$ be monic. Then* $\det(zI_n - C_f) = f(z)$.

*Proof.* Set $D_n(c_0, \dots, c_{n-1}) := \det(zI_n - C_f)$. For $n = 1$, $D_1(c_0) = z + c_0$.
For $n \geq 2$, first-row expansion gives $D_n(c_0, \dots, c_{n-1}) = zD_{n-1}(c_1, \dots, c_{n-1}) + c_0$, since the $(1,1)$-minor is again companion-shaped and the last first-row cofactor contributes $c_0$. Induction gives $D_n = z^n + \sum_{i=0}^{n-1} c_i z^i = f(z)$.
*Structural Commentary*: The last column of $C_f$ stores the coefficients of the polynomial relation.
*Key Insight*: The shifted determinant reads that stored relation back as $f$. □

**Definition 3.4.4.** *Rational Canonical Block Forms*

(i) *Invariant-factor RCF*: $A = \mathrm{diag}(C_{s_1}, \dots, C_{s_r})$, where $s_j \in \mathbb{F}[z]$ are monic nonconstant and $s_r \mid \cdots \mid s_1$.
(ii) *Elementary-divisor RCF*: $A = \mathrm{diag}(C_{q_1}, \dots, C_{q_m})$, where each $q_j$ is power of monic irreducible polynomial.

**Theorem 3.4.5.** *Rational Canonical Form Skeleton*
*For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, there is a unique tuple $\mathrm{InvFact}(A) = (s_1,\dots,s_r)$ with $s_j \in \mathbb{F}[z]$ monic nonconstant, $s_r \mid \cdots \mid s_1$, and $A \sim_{\mathbb{F}} R_A := \mathrm{diag}(C_{s_1},\dots,C_{s_r})$. Moreover, $\mu_A = s_1$ and $p_A = \prod_{j=1}^{r} s_j$.*
*Equivalently, for $A, B \in \mathrm{Mat}(n,n,\mathbb{F})$, $A \sim_{\mathbb{F}} B \iff \mathrm{ElemDiv}(A) = \mathrm{ElemDiv}(B)$ as multisets.*

*Proof.* Let $T_A : \mathbb{F}^n \to \mathbb{F}^n$, $\mathbf{x} \mapsto A\mathbf{x}$.
Theorem 3.2.2 gives $\mathbb{F}^n = \bigoplus_{j=1}^{r} \mathcal{C}_j$, $\mathcal{C}_j \cong_{\mathbb{F}[z]} \mathbb{F}[z]/\langle s_j\rangle$, $s_r \mid \cdots \mid s_1$, $[T_A] = R_A$, $\mu_{T_A} = s_1$, and $p_{T_A} = \prod_{j=1}^{r} s_j$.
Hence $A \sim_{\mathbb{F}} R_A$, $\mu_A = s_1$, and $p_A = \prod_{j=1}^{r} s_j$; uniqueness of $\mathrm{InvFact}(A)$ is uniqueness in the same theorem.
($\Rightarrow$) If $A \sim_{\mathbb{F}} B$, then Theorem 3.4.2 gives $\mathrm{ElemDiv}(A) = \mathrm{ElemDiv}(B)$.
($\Leftarrow$) If $\mathrm{ElemDiv}(A) = \mathrm{ElemDiv}(B)$, then Theorem 3.4.2 gives $A \sim_{\mathbb{F}} B$.
*Structural Commentary*: Companion blocks classify similarity without requiring $p_A$ to split.
*Key Insight*: Similarity classes can be represented by polynomial data rather than by choosing roots. □

**Theorem 3.4.6.** *Rational Canonical Form: Two Versions*
*Let $T \in \mathcal{L}(\mathcal{V})$.*
*(i) If $\mathrm{ElemDiv}(T) = \{q_1,\dots,q_m\}$, then $\exists$ basis $\mathcal{R}$ with $[T]_{\mathcal{R}} = \mathrm{diag}(C_{q_1},\dots,C_{q_m})$. The form is unique up to block permutation.*
*(ii) If $\mathrm{InvFact}(T) = (s_1,\dots,s_r)$ and $s_r \mid \cdots \mid s_1$, then $\exists$ basis $\mathcal{B}$ with $[T]_{\mathcal{B}} = \mathrm{diag}(C_{s_1},\dots,C_{s_r})$. The form is unique in this order.*
*(iii) If $A \in \mathrm{Mat}(n,n,\mathbb{F})$, then the similarity class $[A]_{\sim_{\mathbb{F}}}$ contains both forms. Elementary-divisor form is unique up to block order, invariant-factor form by its divisibility chain.*

*Proof.*
(i) Theorem 3.3.1 gives $\mathcal{V} = \bigoplus_{\alpha=1}^{m} \mathcal{C}_\alpha$ with orders $q_\alpha$; Theorem 3.1.14 gives cyclic bases and blocks $C_{q_\alpha}$. Permuting summands conjugates by a permutation matrix.
(ii) Theorem 3.2.2 gives $\mathcal{V} = \bigoplus_{j=1}^{r} \mathcal{C}_j$, $\mathcal{C}_j \cong_{\mathbb{F}[z]} \mathbb{F}[z]/\langle s_j\rangle$, and cyclic bases with block matrix $\mathrm{diag}(C_{s_1},\dots,C_{s_r})$.
(iii) For $A$, apply (i),(ii) to $T_A : \mathbf{x} \mapsto A\mathbf{x}$; any chosen basis gives a similarity matrix. Uniqueness follows from Theorem 3.4.2.
*Structural Commentary*: Elementary divisors split primary blocks; invariant factors recombine them.
*Key Insight*: Both forms classify the same similarity classes, but they expose different polynomial data. □

**Method 3.4.7.** *Constructing an Operator from Elementary Divisors*
Let $Q = \{q_1,\dots,q_m\}$ be a finite multiset of monic prime powers in $\mathbb{F}[z]$, and set $n := \sum_{j=1}^{m} \deg q_j$.
(i) Set $R_Q := \mathrm{diag}(C_{q_1},\dots,C_{q_m}) \in \mathrm{Mat}(n,n,\mathbb{F})$.
(ii) Then $\mathrm{ElemDiv}(T_{R_Q}) = Q$; hence every such $Q$ is realised on $\mathbb{F}^n$.
(iii) Apply Method 2.10.5 to recombine $Q$ into $\mathrm{InvFact}(T_{R_Q}) = (s_1,\dots,s_r)$, then $\mathrm{diag}(C_{s_1},\dots,C_{s_r}) \sim_{\mathbb{F}} R_Q$.
*Structural Commentary*: RCF constructs operators from divisor data.
*Key Insight*: Companion blocks realise cyclic summands; block sums realise direct sums.

**Example 3.4.8.** *Rational Canonical Possibilities from Degree Counts*
Let $T \in \mathcal{L}(\mathbb{R}^7)$ and $\mu_T = (z-1)(z^2+1)^2$. For $Q = \mathrm{ElemDiv}(T)$, $\mathrm{lcm}\, Q = \mu_T$ and $\sum_{q\in Q} \deg q = 7$.
Hence $Q_1 = \{z-1, (z^2+1)^2, z^2+1\}$ or $Q_2 = \{z-1, z-1, z-1, (z^2+1)^2\}$.
Thus the elementary-divisor RCF is $R_i = \bigoplus_{q\in Q_i} C_q$ for $i = 1, 2$, up to block order.

**Theorem 3.4.9.** *Determinant Form of the Characteristic Polynomial*
*For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, $\chi_A(z) = \det(zI_n - A)$. If $A = [T]_{\mathcal{B}}$, then $\chi_T = \chi_A$.*

*Proof.* Let $Q := \mathrm{ElemDiv}(A)$ and choose elementary-divisor RCF $R = \bigoplus_{q\in Q} C_q \sim_{\mathbb{F}} A$.
By Lemma 3.4.3, $\det(zI - R) = \prod_{q\in Q} \det(zI - C_q) = \prod_{q\in Q} q = \chi_R$.
Since similarity preserves both $Q$ and $\det(zI - \cdot)$, $\chi_A = \chi_R = \det(zI - R) = \det(zI - A)$.
If $A = [T]_{\mathcal{B}}$, then $\mathrm{ElemDiv}(A) = \mathrm{ElemDiv}(T)$, hence $\chi_A = \chi_T$.
*Structural Commentary*: Rational canonical form reduces the determinant computation to companion blocks.
*Key Insight*: The abstract product of elementary divisors is the concrete shifted determinant. □

**Theorem 3.4.10.** *Changing the Base Field*
*Let $\mathbb{F} \subseteq \mathbb{K}$, $A \in \mathrm{Mat}(n,n,\mathbb{F})$, and $\mathrm{ElemDiv}_{\mathbb{F}}(A) = \{p_i^{e_{i,j}}\}_{i,j}$ with $p_i \in \mathbb{F}[z]$ monic irreducible.*
*If $p_i = \prod_{\ell=1}^{m_i} a_{i,\ell}^{d_{i,\ell}}$ in $\mathbb{K}[z]$, with $a_{i,\ell}$ pairwise distinct monic irreducible over $\mathbb{K}$, then $\mathrm{ElemDiv}_{\mathbb{K}}(A) = \{a_{i,\ell}^{d_{i,\ell}e_{i,j}}\}_{i,j,\ell}$.*
*Moreover, $\mathrm{InvFact}_{\mathbb{K}}(A) = \mathrm{InvFact}_{\mathbb{F}}(A)$.*

*Proof.* Take elementary-divisor RCF over $\mathbb{F}$: $R = \bigoplus_{i,j} C_{p_i^{e_{i,j}}} \sim_{\mathbb{F}} A$.
Since $R \in \mathrm{Mat}(n,n,\mathbb{K})$ and $p_i^{e_{i,j}} = \prod_{\ell=1}^{m_i} a_{i,\ell}^{d_{i,\ell} e_{i,j}}$ with coprime factors in $\mathbb{K}[z]$,
Theorem 2.5.6 gives $C_{p_i^{e_{i,j}}} \sim_{\mathbb{K}} \bigoplus_{\ell=1}^{m_i} C_{a_{i,\ell}^{d_{i,\ell} e_{i,j}}}$. Summing over $(i,j)$ yields $\mathrm{ElemDiv}_{\mathbb{K}}(A)$.
Recombination into the divisibility chain gives the original invariant factors.
*Structural Commentary*: Field extension refines elementary divisors.
*Key Insight*: Elementary divisors split; invariant factors persist. □

**Corollary 3.4.11.** *Nonderogatory Means One Companion Block*
*For $A \in \mathrm{Mat}(n,n,\mathbb{F})$, the following are equivalent:*

*(i) $A$ is nonderogatory, i.e. $\mu_A = \chi_A$;*
*(ii) $(\mathbb{F}^n)_A$ is cyclic;*
*(iii) $A \sim_{\mathbb{F}} C_{\chi_A}$.*

*Proof.*

(i) $\Rightarrow$ (ii) This is Theorem 3.3.6.
(ii) $\Rightarrow$ (iii) Cyclicity gives $\mathrm{InvFact}(A) = (s)$. By Theorem 3.2.2, $s = \mu_A = \chi_A$; hence $A \sim_{\mathbb{F}} C_{\chi_A}$.
(iii) $\Rightarrow$ (i) Since $\mu_{C_f} = f$ by Theorem 3.1.13, similarity gives $\mu_A = \mu_{C_{\chi_A}} = \chi_A$.

*Structural Commentary*: Nonderogatory means one cyclic block.
*Key Insight*: One cyclic generator gives one companion block. □

**Corollary 3.4.12.** *When the Elementary-Divisor Form Is Diagonal*
*Let $T \in \mathcal{L}(\mathcal{V})$, $Q = \mathrm{ElemDiv}(T)$, and $R_{\mathrm{ed}} = \bigoplus_{q \in Q} C_q$.*
*Then $R_{\mathrm{ed}}$ is diagonal iff $\deg q = 1$ for all $q \in Q$, iff $\mu_T = \prod_{\lambda \in \Lambda} (z - \lambda)$ for some finite $\Lambda \subseteq \mathbb{F}$.*

*Proof.*

($\Rightarrow$) If $R_{\mathrm{ed}}$ is diagonal, each block $C_q$ has size 1, hence $\deg q = 1$ for all $q \in Q$.
Thus $\mu_T = \mathrm{lcm}\, Q$ is a product of distinct linear factors.
($\Leftarrow$) If $\mu_T = \prod_{\lambda \in \Lambda}(z - \lambda)$, then each prime power $q \in Q$ divides $\mu_T$, so $q = z - \lambda$ for some $\lambda \in \Lambda$.
Hence each $C_q$ is $1 \times 1$, and $R_{\mathrm{ed}}$ is diagonal.

*Structural Commentary*: Diagonal RCF means all companion blocks are $1 \times 1$.
*Key Insight*: Nonlinear or repeated factors obstruct diagonal RCF. □

**Example 3.4.13.** *The Same Polynomial over Different Fields*
Let $\dim V = 4$ and $\mu_T = (z^2+1)(z^2-2)$. Since $\deg \mu_T = 4$, $T$ is nonderogatory and $\mathrm{InvFact}_K(T) = (\mu_T)$ for $K = \mathbb{Q}, \mathbb{R}, \mathbb{C}$; hence $R_{\mathrm{inv},K} = C_{\mu_T}$. Meanwhile

| $K$ | $\mathrm{ElemDiv}_K(T)$ |
|---|---|
| $\mathbb{Q}$ | $\{z^2+1,\ z^2-2\}$ |
| $\mathbb{R}$ | $\{z^2+1,\ z-\sqrt{2},\ z+\sqrt{2}\}$ |
| $\mathbb{C}$ | $\{z-i,\ z+i,\ z-\sqrt{2},\ z+\sqrt{2}\}$ |

Thus $R_{\mathrm{ed},K} = \bigoplus_{q \in \mathrm{ElemDiv}_K(T)} C_q$: scalar extension refines elementary divisors while preserving the invariant factor. Over $\mathbb{C}$, $R_{\mathrm{ed},\mathbb{C}}$ is diagonal.

**Perspective 3.4.14.** *Crosswalk with the Spectral Presentation*
The module and spectral presentations encode the same data at different splitting levels:

(i) diagonalizable $\iff$ $\mathrm{ElemDiv}(T)$ consists of linear factors, by Theorem 3.2.1;
(ii) $(z-\lambda)^k \iff J_k(\lambda)$ over an algebraically closed field, by Perspective 3.3.7 and Theorem 6.2.1;
(iii) rank drops recover the exponents $k$ via Proposition 2.9.5, with $p = z - \lambda$ and $M = \mathcal{V}_T$.

*Key Insight*: Rational form is Jordan form before irreducibles split.

**Method 3.4.15.** *Moving Between Module and Spectral Data*
For $T \in \mathcal{L}(\mathcal{V})$, translate the $\mathbb{F}[z]$-module data of $\mathcal{V}_T$ into spectral data as follows.

(i) Factor $\mu_T = \prod_{i=1}^{s} p_i^{e_i}$ over $\mathbb{F}$; the primary components are $\ker p_i(T)^{e_i}$.
(ii) Decompose each $\ker p_i(T)^{e_i}$ into cyclic summands; their orders $p_i^{e_{i,j}}$ are the elementary divisors.
(iii) Over $\mathbb{F}$, each elementary divisor $p_i^{e_{i,j}}$ gives a companion block $C_{p_i^{e_{i,j}}}$ in rational canonical form.
(iv) Over a splitting field for the $p_i$, each elementary divisor refines into powers $(z-\lambda)^k$ by Theorem 3.4.10; each resulting cyclic block is similar to $J_k(\lambda)$ by Theorem 3.1.14 and Theorem 6.1.3.
(v) $T$ is diagonalizable exactly when every elementary divisor is linear, i.e., Corollary 3.4.12 applies.

*Key Insight*: Rational and Jordan forms encode the same elementary-divisor data over different fields.




Figure 108: Rational-canonical dictionary: $[T]_\sim$, $[\mathcal{V}_T]_{\mathbb{F}[z]}$, $\{s_i\}$ or $\{q_j\}$, and $\mathrm{diag}(C_{s_i})$ are equivalent data.

# 4 Associative Algebras

## 4.1 Associative Algebra Structures

The module viewpoint in Definition 3.1.2 uses one operator at a time. An algebra keeps track of a whole vector space of operators or products at once. The guiding example is $\mathcal{L}(\mathcal{V})$, where addition and scalar multiplication are pointwise and multiplication is composition.

**Definition 4.1.1.** *Unital Associative Algebra*
Let $\mathbb{F}$ be a field.

(i) An *associative $\mathbb{F}$-algebra* is an $\mathbb{F}$-vector space $\mathcal{A}$ and unital ring $(\mathcal{A}, +, \cdot, 1_\mathcal{A})$ such that $r(ab) = (ra)b = a(rb)$ for all $r \in \mathbb{F}$ and $a, b \in \mathcal{A}$.
(ii) $\mathcal{A}$ is *finite-dimensional* iff $\dim_\mathbb{F} \mathcal{A} < \infty$, and *commutative* iff $ab = ba$ for all $a, b \in \mathcal{A}$.
(iii) $a \in \mathcal{A}$ is *invertible* iff $\exists b \in \mathcal{A}$ such that $ab = ba = 1_\mathcal{A}$.
(iv) In this subsection, *algebra* means unital associative $\mathbb{F}$-algebra unless stated otherwise.

**Remark 4.1.2.** *Convention*
Relative to Definition 3.7.1, assume multiplication is associative and unital.
Thus, *algebra* means unital associative $\mathbb{F}$-algebra.

**Definition 4.1.3.** *Center and Central Algebras*

(i) The *center* of an $\mathbb{F}$-algebra $\mathcal{A}$ is $Z(\mathcal{A}) := \{a \in \mathcal{A} : ax = xa \ \forall x \in \mathcal{A}\}$.
(ii) $\mathcal{A}$ is *central over $\mathbb{F}$* iff $Z(\mathcal{A}) = \mathbb{F}1_\mathcal{A} := \{r1_\mathcal{A} : r \in \mathbb{F}\}$.

**Proposition 4.1.4.** *The Scalar Field Sits in the Center*
*If $\mathcal{A}$ is an $\mathbb{F}$-algebra with $1_\mathcal{A} \neq 0$, then $\iota_\mathcal{A} : \mathbb{F} \hookrightarrow Z(\mathcal{A})$, $\quad r \mapsto r1_\mathcal{A}$, is an injective unital ring homomorphism. Conversely, if $R$ is a unital ring and $\mathbb{F} \leq Z(R)$ is a subfield, then $R$ is an $\mathbb{F}$-algebra via $r \cdot x := rx$.*

*Proof.*

($\Rightarrow$) Write $\iota = \iota_\mathcal{A}$. For $r, s \in \mathbb{F}$, $\iota(r+s) = (r+s)1_\mathcal{A} = \iota(r) + \iota(s)$, $\iota(rs) = (rs)1_\mathcal{A} = (r1_\mathcal{A})(s1_\mathcal{A})$, $\iota(1) = 1_\mathcal{A}$.
For $r \in \mathbb{F}$ and $a \in \mathcal{A}$, $\iota(r)a = (r1_\mathcal{A})a = r(1_\mathcal{A}a) = ra = r(a1_\mathcal{A}) = a(r1_\mathcal{A}) = a\iota(r)$, so $\operatorname{im}\iota \subseteq Z(\mathcal{A})$.
If $\iota(r) = 0$ and $r \neq 0$, then $1_\mathcal{A} = \iota(r^{-1})\iota(r) = 0$, contradiction; hence $\ker\iota = \{0\}$.
($\Leftarrow$) Define $r \cdot x = rx$. The field laws in $\mathbb{F} \subseteq R$ give the vector-space axioms, and $1_\mathbb{F} \cdot x = x$.
For $r \in \mathbb{F}$ and $x, y \in R$, centrality gives $r \cdot (xy) = rxy = (rx)y = x(ry) = x(r \cdot y)$, so the scalar action is compatible with the ring product.

*Structural Commentary*: An algebra structure is a choice of central scalar copy $\mathbb{F}1_\mathcal{A} \subseteq Z(\mathcal{A})$.
*Key Insight*: Scalars are precisely central multipliers of the unit. □

**Definition 4.1.5.** *Subalgebras, Ideals, Quotients, Homomorphisms, and Regular Representation*
Let $\mathcal{A}$ be an $\mathbb{F}$-algebra.

(i) A *subalgebra* is a subset $\mathcal{B} \subseteq \mathcal{A}$ such that $1_\mathcal{A} \in \mathcal{B}$, $\mathcal{B} \leq_\mathbb{F} \mathcal{A}$, and $b, b' \in \mathcal{B} \Rightarrow bb' \in \mathcal{B}$.
(ii) For $X \subseteq \mathcal{A}$, the *subalgebra generated by $X$* is $\langle X \rangle_{\text{subalg}} := \bigcap\{\mathcal{B} : \mathcal{B} \text{ is a subalgebra of } \mathcal{A},\ X \subseteq \mathcal{B}\}$.
Equivalently, $\langle X \rangle_{\text{subalg}} = \operatorname{span}_\mathbb{F}\{1_\mathcal{A},\ x_1 \cdots x_m : m \geq 1,\ x_i \in X\}$.
(iii) A *two-sided ideal* is a subset $\mathcal{I} \subseteq \mathcal{A}$ such that $\mathcal{I} \neq \emptyset$, $u, v \in \mathcal{I} \Rightarrow u - v \in \mathcal{I}$, and $a \in \mathcal{A},\ u \in \mathcal{I} \Rightarrow au, ua \in \mathcal{I}$.
(iv) For nonempty $X \subseteq \mathcal{A}$, the *ideal generated by $X$* is $\langle X \rangle_{\text{ideal}} := \{\sum_{j=1}^n a_j x_j b_j : n \geq 1,\ x_j \in X,\ a_j, b_j \in \mathcal{A}\}$.
(v) $\mathcal{A}$ is *simple* iff $\mathcal{A}^2 \neq 0$ and its only two-sided ideals are $0$ and $\mathcal{A}$.
(vi) If $\mathcal{I} \triangleleft \mathcal{A}$, then $\mathcal{A}/\mathcal{I}$ is an $\mathbb{F}$-algebra by $(a + \mathcal{I})(b + \mathcal{I}) := ab + \mathcal{I}$ and $r(a + \mathcal{I}) := ra + \mathcal{I}$.
(vii) For $\mathbb{F}$-algebras $\mathcal{A}, \mathcal{B}$, an *algebra homomorphism* is a map $\sigma : \mathcal{A} \to \mathcal{B}$ such that, for all $a, a' \in \mathcal{A}$ and $r \in \mathbb{F}$, $\sigma(a + a') = \sigma(a) + \sigma(a')$, $\sigma(aa') = \sigma(a)\sigma(a')$, $\sigma(1_\mathcal{A}) = 1_\mathcal{B}$, and $\sigma(ra) = r\sigma(a)$.
(viii) For $a \in \mathcal{A}$, the *left multiplication operator* is $L_a : \mathcal{A} \to \mathcal{A}$, $x \mapsto ax$.
The map $L_\mathcal{A} : \mathcal{A} \to \mathcal{L}(\mathcal{A})$, $a \mapsto L_a$, is the *left regular representation* of $\mathcal{A}$.

**Proposition 4.1.6.** *Kernel and First Isomorphism Theorem for Algebras*
*If $\sigma : \mathcal{A} \to \mathcal{B}$ is an $\mathbb{F}$-algebra homomorphism,*
*then $\ker\sigma \triangleleft \mathcal{A}$ and $\overline{\sigma} : \mathcal{A}/\ker\sigma \to \operatorname{im}\sigma$, $a + \ker\sigma \mapsto \sigma(a)$, is an $\mathbb{F}$-algebra isomorphism.*

*Proof.* Set $K := \ker \sigma$. Linearity gives $K \leq_{\mathbb{F}} \mathcal{A}$.
If $u \in K$ and $a \in \mathcal{A}$, then $\sigma(au) = \sigma(a)\sigma(u) = 0 = \sigma(u)\sigma(a) = \sigma(ua)$, hence $au, ua \in K$ and $K \triangleleft \mathcal{A}$.
Define $\overline{\sigma}(a + K) := \sigma(a)$. Since $a + K = a' + K \iff a - a' \in K \iff \sigma(a) = \sigma(a')$, $\overline{\sigma}$ is well defined and injective; it is surjective onto $\operatorname{im} \sigma$.
For $a, a' \in \mathcal{A}$ and $r \in \mathbb{F}$, $\overline{\sigma}((a + K)(a' + K)) = \sigma(aa') = \sigma(a)\sigma(a')$, $\overline{\sigma}(r(a + K)) = r\sigma(a)$, and $\overline{\sigma}(1_{\mathcal{A}} + K) = 1_{\mathcal{B}}$.
Thus $\overline{\sigma}$ is an $\mathbb{F}$-algebra isomorphism.
*Structural Commentary*: Quotienting by the kernel removes exactly the relations invisible to $\sigma$.
*Key Insight*: The image is the faithful algebra obtained after collapsing $\ker \sigma$. □

**Theorem 4.1.7.** *Left Regular Representation*
*For every $\mathbb{F}$-algebra $\mathcal{A}$, $L_{\mathcal{A}} : \mathcal{A} \hookrightarrow \mathcal{L}(\mathcal{A})$, $a \mapsto L_a$, is an $\mathbb{F}$-algebra monomorphism.*

*Proof.* For $a, b \in \mathcal{A}$, $r \in \mathbb{F}$, and $x \in \mathcal{A}$, $L_{a+b}(x) = (a + b)x = L_a(x) + L_b(x)$ and $L_{ra}(x) = (ra)x = rL_a(x)$, so $L_{\mathcal{A}}$ is $\mathbb{F}$-linear. Also $L_{ab}(x) = (ab)x = a(bx) = (L_aL_b)(x)$ and $L_{1_{\mathcal{A}}} = I_{\mathcal{A}}$, so products and units are preserved.
If $L_a = 0$, then $a = L_a(1_{\mathcal{A}}) = 0$, so $L_{\mathcal{A}}$ is injective.
*Structural Commentary*: Left multiplication realises $\mathcal{A}$ inside its own operator algebra.
*Key Insight*: The unit detects elements, since $L_a(1_{\mathcal{A}}) = a$. □




Figure 109: The left regular representation sends $a \in \mathcal{A}$ to $L_a : x \mapsto ax$; $L_{ab} = L_aL_b$ and $L_a(1_{\mathcal{A}}) = a$ give multiplicativity and injectivity.

**Method 4.1.8.** *Building Multiplication from Structure Constants*
Let $V = \bigoplus_{i \in I} \mathbb{F}b_i$. Choose $c_{ij}^k \in \mathbb{F}$ with $\{k : c_{ij}^k \neq 0\}$ finite for each $(i, j)$.

(i) Define $b_ib_j := \sum_{k \in I} c_{ij}^k b_k$ and extend bilinearly: $(\sum_i r_ib_i)(\sum_j s_jb_j) := \sum_{i,j,k} r_is_jc_{ij}^k b_k$.
(ii) The product is associative iff $\sum_{\ell \in I} c_{ij}^\ell c_{\ell k}^m = \sum_{\ell \in I} c_{jk}^\ell c_{i\ell}^m$ for all $i, j, k, m \in I$.
(iii) The product is commutative iff $c_{ij}^k = c_{ji}^k$ for all $i, j, k \in I$.
(iv) A basis vector $b_e$ is an identity iff $c_{ej}^m = \delta_j^m = c_{je}^m$ for all $j, m \in I$.

**Example 4.1.9.** *Basic Algebras*

(i) If $\mathbb{F} \subseteq \mathbb{E}$ are fields, then $\mathbb{E}$ is an $\mathbb{F}$-algebra by $r \cdot x = rx$ and multiplication in $\mathbb{E}$.
(ii) $\mathbb{F}[x]$ is a commutative $\mathbb{F}$-algebra with unit 1 and scalar embedding $r \mapsto r$.
(iii) $\operatorname{Mat}(n, n, \mathbb{F})$ is an $\mathbb{F}$-algebra with unit $I_n$ and product $AB$.
(iv) For an $\mathbb{F}$-vector space $\mathcal{V}$, $\mathcal{L}(\mathcal{V}) = \operatorname{End}_{\mathbb{F}}(\mathcal{V})$ is an $\mathbb{F}$-algebra with unit $I_{\mathcal{V}}$ and product $S \circ T$.

**Definition 4.1.10.** *Group Algebra*
For a group $G$ and field $\mathbb{F}$, the *group algebra* is $\mathbb{F}[G] := \{\sum_{g \in G} r_g g : r_g \in \mathbb{F},\ |\{g : r_g \neq 0\}| < \infty\}$.
It has coefficientwise vector-space operations and product $(\sum_g r_g g)(\sum_h s_h h) := \sum_{g,h} r_g s_h (gh)$.

**Example 4.1.11.** *Group Algebras*
In $\mathbb{F}[G]$, $G$ is an $\mathbb{F}$-basis, $e_G$ is the algebra identity, and associativity follows from $(gh)k = g(hk)$ by bilinearity.
If $G$ is finite, then $\dim_{\mathbb{F}} \mathbb{F}[G] = |G|$. Moreover, $\mathbb{F}[G]$ is commutative iff $G$ is abelian.

**Example 4.1.12.** *Cyclic Group Algebras and Circulant Shifts*
Let $G = \langle \alpha \rangle$, $|G| = n$, and let $\mathcal{B} = (1, \alpha, \ldots, \alpha^{n-1})$ be the ordered $\mathbb{F}$-basis of $\mathbb{F}[G]$.
For $0 \leq j, k < n$, $L_{\alpha^k}(\alpha^j) = \alpha^{j+k}$ with exponents read modulo $n$, so $L_{\alpha^k}$ is the $k$-fold cyclic shift of $\mathcal{B}$.
Let $P := [L_\alpha]_{\mathcal{B}}$, one has $[L_{\alpha^k}]_{\mathcal{B}} = P^k$, where

$$P = \begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix}.$$

Hence $[L_{\sum_{k=0}^{n-1} c_k \alpha^k}]_{\mathcal{B}} = \sum_{k=0}^{n-1} c_k P^k$, precisely the circulant matrices.

**Proposition 4.1.13.** *Endomorphisms and Matrices Are the Same Algebra after Choosing a Basis*
*Let $\dim \mathcal{V} = n$ and fix an ordered basis $\mathcal{B}$ of $\mathcal{V}$.*
*Then $\mu_{\mathcal{B}} : \mathcal{L}(\mathcal{V}) \to \operatorname{Mat}(n, n, \mathbb{F})$, $T \mapsto [T]_{\mathcal{B}}$, is an $\mathbb{F}$-algebra isomorphism.*

*Proof.* By Theorem 3.4.4, $\mu_{\mathcal{B}}$ is an $\mathbb{F}$-linear bijection.
For $S, T \in \mathcal{L}(\mathcal{V})$ and $x \in \mathcal{V}$, $[STx]_{\mathcal{B}} = [S]_{\mathcal{B}}[Tx]_{\mathcal{B}} = [S]_{\mathcal{B}}[T]_{\mathcal{B}}[x]_{\mathcal{B}}$, hence $\mu_{\mathcal{B}}(ST) = \mu_{\mathcal{B}}(S)\mu_{\mathcal{B}}(T)$.
Also $\mu_{\mathcal{B}}(I_{\mathcal{V}}) = I_n$; therefore $\mu_{\mathcal{B}}$ is an $\mathbb{F}$-algebra isomorphism.
*Structural Commentary*: A basis is a coordinate bridge from abstract endomorphisms to concrete matrices.
*Key Insight*: Composition becomes matrix multiplication after taking coordinate columns. □

**Theorem 4.1.14.** *Center and Ideals of an Endomorphism Algebra*
*Let $0 \neq \mathcal{V}$ be an $\mathbb{F}$-vector space, and set $\mathcal{F}(\mathcal{V}) := \{T \in \mathcal{L}(\mathcal{V}) : \dim \operatorname{im} T < \infty\}$.*

*(i) $Z(\mathcal{L}(\mathcal{V})) = \mathbb{F}I_{\mathcal{V}}$, so $\mathcal{L}(\mathcal{V})$ is central.*
*(ii) $\mathcal{F}(\mathcal{V}) \triangleleft \mathcal{L}(\mathcal{V})$, and $0 \neq \mathcal{J} \triangleleft \mathcal{L}(\mathcal{V}) \Rightarrow \mathcal{F}(\mathcal{V}) \subseteq \mathcal{J}$.*
*(iii) $\mathcal{L}(\mathcal{V})$ is simple iff $\dim \mathcal{V} < \infty$.*

*Proof.*

(i) Let $T \in Z(\mathcal{L}(\mathcal{V}))$. For $0 \neq v \in \mathcal{V}$ and $\varphi \in \mathcal{V}^*$ with $\varphi(v) = 1$, commutation with $v \otimes \varphi : x \mapsto \varphi(x)v$ gives $Tv = T(v \otimes \varphi)(v) = (v \otimes \varphi)T(v) = \varphi(Tv)v$, so every line is $T$-stable. Thus $Tv = \lambda_v v$ for $v \neq 0$, and applying this to independent $u, v$ and to $u + v$ gives $\lambda_u = \lambda_v$; hence $T = rI_{\mathcal{V}}$.
Scalars commute with all endomorphisms.

(ii) If $R, S \in \mathcal{L}(\mathcal{V})$ and $T \in \mathcal{F}(\mathcal{V})$, then $\operatorname{im}(RTS) \subseteq R(\operatorname{im} T)$, so $RTS \in \mathcal{F}(\mathcal{V})$ and $\mathcal{F}(\mathcal{V}) \triangleleft \mathcal{L}(\mathcal{V})$. If $0 \neq \mathcal{J} \triangleleft \mathcal{L}(\mathcal{V})$, choose $0 \neq S \in \mathcal{J}$, $e$ with $Se \neq 0$, and $\eta \in \mathcal{V}^*$ with $\eta(Se) = 1$. For any $v \in \mathcal{V}$ and $\varphi \in \mathcal{V}^*$, $v \otimes \varphi = (v \otimes \eta)S(e \otimes \varphi) \in \mathcal{J}$, so $\mathcal{J}$ contains all rank-one maps, hence all finite-rank maps.

(iii) If $\dim \mathcal{V} < \infty$, then $\mathcal{F}(\mathcal{V}) = \mathcal{L}(\mathcal{V})$, so (ii) makes every nonzero two-sided ideal equal to $\mathcal{L}(\mathcal{V})$.
If $\dim \mathcal{V} = \infty$, then $0 \neq \mathcal{F}(\mathcal{V}) \neq \mathcal{L}(\mathcal{V})$, so $\mathcal{L}(\mathcal{V})$ is not simple.

*Structural Commentary*: Rank-one operators are the atoms controlling both centrality and ideals in $\mathcal{L}(\mathcal{V})$.
*Key Insight*: Sandwich one nonzero map to obtain every rank-one map. □

**Perspective 4.1.15.** *Operator Connection*
$T \in \mathcal{L}(\mathcal{V})$ gives an $\mathbb{F}[x]$-module by $x \cdot v = T(v)$. For any algebra $\mathcal{A}$, the left regular map $L_{\mathcal{A}} : \mathcal{A} \hookrightarrow \mathcal{L}(\mathcal{A})$ realizes $\mathcal{A}$ as a faithful endomorphism algebra.

## 4.2 Modules over Algebras

**Definition 4.2.1.** *Algebraic Elements and Minimal Polynomials*
Let $\mathcal{A}$ be an $\mathbb{F}$-algebra and $a \in \mathcal{A}$.

(i) The *evaluation map at $a$* is $\operatorname{ev}_a : \mathbb{F}[x] \to \mathcal{A}$, $p \mapsto p(a)$.
(ii) $a$ is *algebraic over $\mathbb{F}$* iff $\ker \operatorname{ev}_a \neq 0$.
(iii) If $a$ is algebraic, its *minimal polynomial* $m_a$ is the unique monic generator of $\ker \operatorname{ev}_a$, equivalently the monic polynomial of least degree with $m_a(a) = 0$.

**Theorem 4.2.2.** *Finite-Dimensional Algebras Are Algebraic*
*If $\dim_{\mathbb{F}} \mathcal{A} < \infty$, then every $a \in \mathcal{A}$ is algebraic over $\mathbb{F}$. Moreover, the algebraic closure $\overline{\mathbb{Q}}_{\mathbb{C}}$ of $\mathbb{Q}$ inside $\mathbb{C}$ is an infinite-dimensional $\mathbb{Q}$-algebra whose elements are all algebraic.*

*Proof.* Let $\dim_{\mathbb{F}} \mathcal{A} = n$ and $a \in \mathcal{A}$. The $n+1$ vectors $1, a, \dots, a^n$ are linearly dependent, so $\sum_{j=0}^{n} c_j a^j = 0$ for some $(c_0, \dots, c_n) \neq 0$; hence $p(x) = \sum_{j=0}^{n} c_j x^j \neq 0$ satisfies $p(a) = 0$.
If $\alpha, \beta \in \overline{\mathbb{Q}}_{\mathbb{C}}$, then $\mathbb{Q}(\alpha, \beta)/\mathbb{Q}$ is finite, so $\alpha + \beta$, $\alpha\beta$, and $r\alpha$ for $r \in \mathbb{Q}$ are algebraic; hence $\overline{\mathbb{Q}}_{\mathbb{C}}$ is a $\mathbb{Q}$-algebra.
For each $n \geq 1$, $x^n - 2$ is Eisenstein at 2, so it has a root of degree $n$ in $\overline{\mathbb{Q}}_{\mathbb{C}}$; therefore $\dim_{\mathbb{Q}} \overline{\mathbb{Q}}_{\mathbb{C}} = \infty$.
*Structural Commentary*: Finite-dimensionality forces powers of each element into a linear relation.
*Key Insight*: Algebraic need not mean finite-dimensional as an algebra. □

**Proposition 4.2.3.** *The Algebra Generated by One Algebraic Element*
*Let $a \in \mathcal{A}$ be algebraic with minimal polynomial $m_a$. Then $\mathbb{F}[a] := \operatorname{im} \operatorname{ev}_a = \{p(a) : \deg p < \deg m_a\}$ and $\mathbb{F}[a] \cong \mathbb{F}[x]/\langle m_a \rangle$.*

*Proof.* $\operatorname{ev}_a : \mathbb{F}[x] \to \mathcal{A}$ is an $\mathbb{F}$-algebra homomorphism with image $\mathbb{F}[a]$.
Since $\ker \operatorname{ev}_a = \{p : p(a) = 0\}$ and $m_a$ is the monic generator of this kernel, $\ker \operatorname{ev}_a = \langle m_a \rangle$.
Proposition 4.1.6 gives $\mathbb{F}[a] \cong \mathbb{F}[x]/\langle m_a \rangle$. $\forall p \in \mathbb{F}[x]$, by division $p = qm_a + r$, $\deg r < \deg m_a$, hence $p(a) = r(a)$.
*Structural Commentary*: One algebraic element gives one polynomial quotient.
*Key Insight*: Evaluation turns polynomial division modulo $m_a$ into normal forms in $\mathbb{F}[a]$. □

**Theorem 4.2.4.** *Minimal Polynomial, Annihilator, and Invertibility*
*Let $a \in \mathcal{A}$ be algebraic and write $m_a(x) = x^n + \sum_{j=0}^{n-1} \alpha_j x^j$.*
*Then $\operatorname{ann}_{\mathbb{F}[x]}(a) := \{f \in \mathbb{F}[x] : f(a) = 0\} = \langle m_a \rangle$, and $a \in \mathcal{A}^\times \iff \alpha_0 \neq 0$.*

*Proof.* By Definition 4.2.1, $\operatorname{ann}_{\mathbb{F}[x]}(a) = \ker \operatorname{ev}_a = \langle m_a \rangle$.

($\Rightarrow$) If $a \in \mathcal{A}^\times$ and $\alpha_0 = 0$, then $m_a = xq$ with $\deg q < \deg m_a$, and $0 = m_a(a) = aq(a)$ gives $q(a) = 0$, contradicting minimality of $m_a$; hence $\alpha_0 \neq 0$.

($\Leftarrow$) If $\alpha_0 \neq 0$, then $0 = m_a(a) = a(a^{n-1} + \sum_{j=1}^{n-1} \alpha_j a^{j-1}) + \alpha_0$, so $b := -\alpha_0^{-1}(a^{n-1} + \sum_{j=1}^{n-1} \alpha_j a^{j-1})$ satisfies $ab = 1_\mathcal{A}$. Since $b \in \mathbb{F}[a]$, $ab = ba$, so $ba = 1_\mathcal{A}$ and $a \in \mathcal{A}^\times$.

*Structural Commentary*: Invertibility is read from the constant term of the minimal polynomial.
*Key Insight*: A nonzero constant term solves for $a^{-1}$ as a polynomial in $a$. □

## 4.3 Zero Divisors and Spectra of Algebraic Elements

**Definition 4.3.1.** *Zero Divisors and Spectrum*
Let $\mathcal{A}$ be an $\mathbb{F}$-algebra and $0 \neq a \in \mathcal{A}$.

(i) $a$ is a *left zero divisor* iff $\exists b \in \mathcal{A} \setminus \{0\}$ with $ab = 0$.
(ii) $a$ is a *right zero divisor* iff $\exists c \in \mathcal{A} \setminus \{0\}$ with $ca = 0$.
(iii) If $a$ is algebraic, *spectrum* is $\operatorname{spec}_\mathcal{A}(a) := \{r \in \mathbb{F} : m_a(r) = 0\}$; elements are algebraic *eigenvalues* of $a$.

**Theorem 4.3.2.** *Invertible if and only if Not a Zero Divisor*
*Let $a \in \mathcal{A}$ be algebraic. Then $a \in \mathcal{A}^\times$ iff $a$ is neither a left nor a right zero divisor.*

*Proof.*

($\Rightarrow$) If $a \in \mathcal{A}^\times$ and $ab = 0$, then $b = a^{-1}ab = 0$; if $ca = 0$, then $c = caa^{-1} = 0$. Hence $a$ is neither a left nor a right zero divisor.

($\Leftarrow$) Prove the contrapositive. If $a \notin \mathcal{A}^\times$, Theorem 4.2.4 gives $m_a(x) = xp(x)$ with $p \neq 0$ and $\deg p < \deg m_a$. Then $0 = m_a(a) = ap(a) = p(a)a$, while minimality gives $p(a) \neq 0$. Thus $a$ is both left, right zero divisor.

*Structural Commentary*: For algebraic elements, noninvertibility forces a polynomial kernel witness.
*Key Insight*: Factoring $m_a$ by $x$ manufactures the zero-divisor witness $p(a)$. □

**Corollary 4.3.3.** *One-Sided Inverses Become Two-Sided*
*Let $a \in \mathcal{A}$ be algebraic. If $\exists b \in \mathcal{A}$ with $ba = 1_\mathcal{A}$ or $ab = 1_\mathcal{A}$, then $a \in \mathcal{A}^\times$.*

*Proof.* Assume $ba = 1_\mathcal{A}$. If $ac = 0$, then $c = bac = 0$; if $ax = 0$, then $x = bax = 0$. Hence $a$ is neither a right nor a left zero divisor, so Theorem 4.3.2 gives $a \in \mathcal{A}^\times$. The case $ab = 1_\mathcal{A}$ is symmetric.
*Structural Commentary*: Algebraicity upgrades one-sided cancellation to two-sided invertibility.
*Key Insight*: A one-sided inverse rules out both zero-divisor obstructions. □

**Theorem 4.3.4.** *Roots Detected by Invertibility*
*Let $a \in \mathcal{A}$ be algebraic and $r \in \mathbb{F}$. Then $m_a(r) = 0 \iff a - r1_\mathcal{A} \notin \mathcal{A}^\times$. In particular, $a \in \mathcal{A}^\times \iff 0 \notin \operatorname{spec}_\mathcal{A}(a)$.*

*Proof.*

($\Rightarrow$) If $m_a(r) = 0$, write $m_a = (x - r)q$ with $\deg q < \deg m_a$. Then $0 = m_a(a) = (a - r1_\mathcal{A})q(a) = q(a)(a - r1_\mathcal{A})$ and $q(a) \neq 0$ by minimality, so $a - r1_\mathcal{A} \notin \mathcal{A}^\times$.

($\Leftarrow$) If $m_a(r) \neq 0$, then $\gcd(x - r, m_a) = 1$, so $\exists u, v \in \mathbb{F}[x]$ with $u(x)(x - r) + v(x)m_a(x) = 1$. Evaluating at $a$ gives $u(a)(a - r1_\mathcal{A}) = 1_\mathcal{A} = (a - r1_\mathcal{A})u(a)$, hence $a - r1_\mathcal{A} \in \mathcal{A}^\times$, contradiction.

Taking $r = 0$ gives $a \in \mathcal{A}^\times \iff m_a(0) \neq 0 \iff 0 \notin \operatorname{spec}_\mathcal{A}(a)$.
*Structural Commentary*: Roots of $m_a$ are precisely the nonunit scalar shifts of $a$.
*Key Insight*: Bézout turns $\gcd(x - r, m_a) = 1$ into an explicit inverse. □

**Theorem 4.3.5.** *Polynomial Spectral Mapping for Algebraic Elements*
*Let $\mathbb{F}$ be algebraically closed, $a \in \mathcal{A}$ algebraic, and $p \in \mathbb{F}[x]$.*
*Then $p(a)$ is algebraic and* $\operatorname{spec}_\mathcal{A}(p(a)) = p(\operatorname{spec}_\mathcal{A}(a)) := \{p(r) : r \in \operatorname{spec}_\mathcal{A}(a)\}$.

*Proof.* Since $\mathbb{F}[p(a)] \subseteq \mathbb{F}[a]$ and $\dim_\mathbb{F} \mathbb{F}[a] < \infty$, $p(a)$ is algebraic. If $p$ is constant, the claim is immediate.
If $\lambda \in \operatorname{spec}_\mathcal{A}(p(a))$, then $p(a) - \lambda 1_\mathcal{A} \notin \mathcal{A}^\times$. Factor $p(x) - \lambda = c\prod_{j=1}^{m}(x - r_j)^{e_j}$ over $\mathbb{F}$.
Since a product of units is a unit, $a - r_j 1_\mathcal{A} \notin \mathcal{A}^\times$ for some $j$. Theorem 4.3.4 gives $r_j \in \operatorname{spec}_\mathcal{A}(a)$ and $\lambda = p(r_j)$.
Conversely, if $r \in \operatorname{spec}_\mathcal{A}(a)$, then $p(x) - p(r) = (x - r)q(x)$, so $p(a) - p(r)1_\mathcal{A} = (a - r1_\mathcal{A})q(a)$ is noninvertible; otherwise $a - r1_\mathcal{A}$ has a one-sided inverse and Corollary 4.3.3 makes it invertible. Thus $p(r) \in \operatorname{spec}_\mathcal{A}(p(a))$.
*Structural Commentary*: Polynomial evaluation carries spectra by applying the same polynomial to roots.
*Key Insight*: Factor $p(x) - \lambda$ and detect a nonunit scalar shift of $a$. □

**Proposition 4.3.6.** *Spectra of ab and ba*
*If $\dim_\mathbb{F} \mathcal{A} < \infty$, $\mathbb{F}$ is algebraically closed, and $a, b \in \mathcal{A}$, then* $\operatorname{spec}_\mathcal{A}(ab) = \operatorname{spec}_\mathcal{A}(ba)$.

*Proof.* All elements are algebraic by Theorem 4.2.2. Let $r \in \mathbb{F}^\times$.
If $ba - r1_\mathcal{A} \in \mathcal{A}^\times$, set $u := a(ba - r1_\mathcal{A})^{-1}b - 1_\mathcal{A}$; then $(ab - r1_\mathcal{A})u = r1_\mathcal{A}$, so $ab - r1_\mathcal{A}$ has a right inverse, hence is invertible by Corollary 4.3.3. Thus $r \notin \operatorname{spec}_\mathcal{A}(ba) \Rightarrow r \notin \operatorname{spec}_\mathcal{A}(ab)$, and symmetry gives equality off 0.
If $ba \in \mathcal{A}^\times$ with inverse $y$, then $b(ay) = 1_\mathcal{A}$ and $(yb)a = 1_\mathcal{A}$, so $b, a \in \mathcal{A}^\times$ by Corollary 4.3.3; hence $ab \in \mathcal{A}^\times$. The converse is symmetric, so 0 belongs to both spectra or neither.
*Structural Commentary*: $\operatorname{spec}(ab) \setminus \{0\} = \operatorname{spec}(ba) \setminus \{0\}$, and algebraicity resolves the 0-case.
*Key Insight*: The resolvent identity transfers invertibility between $ba - r1_\mathcal{A}$ and $ab - r1_\mathcal{A}$ for $r \neq 0$. □

## 4.4 Division Algebras

**Definition 4.4.1.** *Division Algebras*
An associative $\mathbb{F}$-algebra $\mathcal{D}$ is a *division algebra* iff $\mathcal{D} \setminus \{0\} \subseteq \mathcal{D}^\times$.

**Example 4.4.2.** *Hamilton's Quaternions*
Hamilton's quaternion algebra is $\mathbb{H} := \mathbb{R}1 \oplus \mathbb{R}i \oplus \mathbb{R}j \oplus \mathbb{R}k$ with unit 1,
relations $i^2 = j^2 = k^2 = -1$, $ij = k = -ji$, $jk = i = -kj$, and $ki = j = -ik$, extended $\mathbb{R}$-bilinearly.
It is noncommutative since $ij = -ji$. For $q = r_0 + r_1 i + r_2 j + r_3 k$, define $\overline{q} := r_0 - r_1 i - r_2 j - r_3 k$ and $N(q) := q\overline{q} = r_0^2 + r_1^2 + r_2^2 + r_3^2$. If $q \neq 0$, then $N(q) > 0$ and $q^{-1} = N(q)^{-1}\overline{q}$, so $\mathbb{H}$ is a real division algebra.




Figure 110: Quaternion basis products: the cycle $i \to j \to k \to i$ gives $ij = k$, $jk = i$, and $ki = j$; reversed order negates the product. Also $i^2 = j^2 = k^2 = -1$, and 1 is the unit.

**Theorem 4.4.3.** *Division Algebras over Algebraically Closed Fields*
*If $\dim_\mathbb{F} \mathcal{D} < \infty$, $\mathcal{D}$ is a division $\mathbb{F}$-algebra, and $\mathbb{F}$ is algebraically closed, then $\mathcal{D} = \mathbb{F}1_\mathcal{D} \cong \mathbb{F}$.*

*Proof.* Let $a \in \mathcal{D}$. By Theorem 4.2.2, $a$ is algebraic with minimal polynomial $m_a$.
If $m_a = pq$ with $0 < \deg p, \deg q < \deg m_a$, then $0 = m_a(a) = p(a)q(a)$, while minimality gives $p(a), q(a) \neq 0$; this contradicts $\mathcal{D} \setminus \{0\} \subseteq \mathcal{D}^\times$. Thus $m_a$ is irreducible, hence linear over algebraically closed $\mathbb{F}$: $m_a = x - r$.
Therefore $a = r1_\mathcal{D}$, so $\mathcal{D} \subseteq \mathbb{F}1_\mathcal{D}$, and the reverse inclusion is the scalar copy.
*Structural Commentary*: Algebraic closure leaves only scalars.
*Key Insight*: Irreducible minimal polynomials collapse to linear polynomials. □

**Theorem 4.4.4.** *Wedderburn's Theorem*
*Every finite division algebra $\mathcal{D}$ is commutative; equivalently $\mathcal{D} = Z(\mathcal{D})$, hence $\mathcal{D}$ is a finite field.*

*Proof.* Let $F := Z(\mathcal{D})$, $|F| = z$, $[\mathcal{D} : F] = n$, and $G := \mathcal{D}^\times$.
Suppose $\mathcal{D}$ is noncommutative, and let $B$ be representatives of the noncentral conjugacy classes of $G$.
For $\beta \in B$, $C_\mathcal{D}(\beta)$ is a proper division subalgebra containing $F$, so $|C_\mathcal{D}(\beta)| = z^{k_\beta}$ with $1 < k_\beta < n$ and $k_\beta \mid n$.
By the class equation, Theorem 2.5.2, $z^n - 1 = z - 1 + \sum_{\beta \in B}(z^n - 1)/(z^{k_\beta} - 1)$.
Since $k_\beta \mid n$ and $k_\beta < n$, the primitive $n$th roots occur in $(x^n - 1)/(x^{k_\beta} - 1)$, so Propositions 3.12.2 and 3.12.3 give $Q_n(z) \mid (z^n - 1)/(z^{k_\beta} - 1)$ for each $\beta$, and also $Q_n(z) \mid z^n - 1$.
The class equation therefore gives $Q_n(z) \mid z - 1$, contradicting Lemma 3.12.4, which gives $|Q_n(z)| > z - 1$.
Thus $G$ has no noncentral conjugacy classes, so $G$ is abelian and $\mathcal{D}$ is commutative.
*Structural Commentary*: The class equation converts noncommutativity into cyclotomic divisibility.
*Key Insight*: The cyclotomic lower bound is too large to divide $z - 1$. □

**Theorem 4.4.5.** *Frobenius Classification over $\mathbb{R}$*
*If $\dim_\mathbb{R} \mathcal{D} < \infty$ and $\mathcal{D}$ is a real division algebra, then $\mathcal{D} \cong \mathbb{R}$, $\mathcal{D} \cong \mathbb{C}$, or $\mathcal{D} \cong \mathbb{H}$.*

*Proof.* For each $d \in \mathcal{D}$, $m_d$ is linear or irreducible quadratic over $\mathbb{R}$; completing the square gives $d = r + u$ with $r \in \mathbb{R}$ and $u^2 \in \mathbb{R}_{\leq 0}$. Set $\mathcal{D}_0 := \{u \in \mathcal{D} : u^2 \in \mathbb{R}_{\leq 0}\}$; the preceding decomposition and uniqueness give $\mathcal{D} = \mathbb{R} \oplus \mathcal{D}_0$. If $\mathcal{D}_0 = 0$, then $\mathcal{D} \cong \mathbb{R}$. If $\dim \mathcal{D}_0 = 1$, choose $i \in \mathcal{D}_0$ with $i^2 = -1$, so $\mathcal{D} = \mathbb{R} \oplus \mathbb{R}i \cong \mathbb{C}$.
If $\dim \mathcal{D}_0 > 1$, use $\langle u, v \rangle := -\frac{1}{2}(uv + vu)$ on $\mathcal{D}_0$ to choose orthonormal $i, j \in \mathcal{D}_0$ with $i^2 = j^2 = -1$; $ij + ji = 0$.

Then $k := ij$ satisfies $k^2 = -1$, $ij = k$, $jk = i$, $ki = j$, and reversed products change sign.
If $\ell \in \mathcal{D}_0$ is orthogonal to $i, j, k$, the same identities force $\ell = 0$, so $\mathcal{D} = \mathbb{R} \oplus \mathbb{R}i \oplus \mathbb{R}j \oplus \mathbb{R}k \cong \mathbb{H}$.
*Structural Commentary*: A real division algebra splits into scalars plus a Euclidean pure part.
*Key Insight*: Orthogonal imaginary units generate exactly $\mathbb{C}$ or $\mathbb{H}$. □

# 5 Tensor Products

## 5.1 Tensor Products and Universal Properties

Tensor products are the device that turns bilinear data into linear data. The guiding rule is the same one already used for quotient spaces in Theorem 2.2.10: a universal object is recognised by the unique way every compatible map factors through it.

**Definition 5.1.1.** *Bilinear Maps*
Let $\mathcal{U}, \mathcal{V}, \mathcal{W}$ be $\mathbb{F}$-vector spaces and $f : \mathcal{U} \times \mathcal{V} \to \mathcal{W}$.

(i) $f$ is *bilinear* if $\forall r, s \in \mathbb{F}$, $\mathbf{u}, \mathbf{u}' \in \mathcal{U}$, $\mathbf{v}, \mathbf{v}' \in \mathcal{V}$, $f(r\mathbf{u} + s\mathbf{u}', \mathbf{v}) = rf(\mathbf{u}, \mathbf{v}) + sf(\mathbf{u}', \mathbf{v})$ and $f(\mathbf{u}, r\mathbf{v} + s\mathbf{v}') = rf(\mathbf{u}, \mathbf{v}) + sf(\mathbf{u}, \mathbf{v}')$.
(ii) $\mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{W}) := \{f : \mathcal{U} \times \mathcal{V} \to \mathcal{W} : f \text{ bilinear}\}$.
(iii) If $\mathcal{W} = \mathbb{F}$, then $\mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathbb{F})$ is the space of *bilinear forms* from Definition 4.4.1.

**Example 5.1.2.** *Basic Bilinear Maps*

(i) If $A$ is an $\mathbb{F}$-algebra, then $\mu : A \times A \to A$, $\mu(a, b) := ab$, is bilinear.
(ii) Evaluation $\mathrm{ev} : \mathcal{L}(\mathcal{V}, \mathcal{W}) \times \mathcal{V} \to \mathcal{W}$, $\mathrm{ev}(T, \mathbf{v}) := T\mathbf{v}$, is bilinear.
(iii) If $\varphi \in \mathcal{V}'$ and $\psi \in \mathcal{W}'$, then $B_{\varphi,\psi} : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$, $B_{\varphi,\psi}(\mathbf{v}, \mathbf{w}) := \varphi(\mathbf{v})\psi(\mathbf{w})$, is bilinear.

**Definition 5.1.3.** *Tensor Product by Universal Property*
Let $\mathcal{U}, \mathcal{V}$ be $\mathbb{F}$-vector spaces.

(i) A pair $(\mathcal{T}, t)$ is a *tensor product* of $\mathcal{U}$ and $\mathcal{V}$ if $\mathcal{T}$ is an $\mathbb{F}$-vector space, $t \in \mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{T})$, and for every $\mathbb{F}$-vector space $\mathcal{W}$ and $f \in \mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{W})$, $\exists! \widetilde{f} \in \mathcal{L}(\mathcal{T}, \mathcal{W})$ such that $f = \widetilde{f} \circ t$.
(ii) The map $\widetilde{f}$ is the *mediating map* for $f$.
(iii) Write $\mathcal{T} = \mathcal{U} \otimes \mathcal{V}$ and $\mathbf{u} \otimes \mathbf{v} := t(\mathbf{u}, \mathbf{v})$; such tensors are *decomposable*.




Figure 111: A tensor product is universal for bilinear maps out of $\mathcal{U} \times \mathcal{V}$.

**Theorem 5.1.4.** *Coordinate-Free Quotient Construction*
*Let $\mathcal{U}, \mathcal{V}$ be $\mathbb{F}$-vector spaces and let $\mathbb{F}^{(\mathcal{U}\times\mathcal{V})}$ have basis $\{[\mathbf{u}, \mathbf{v}] : (\mathbf{u}, \mathbf{v}) \in \mathcal{U} \times \mathcal{V}\}$.*
*Let $\mathcal{R}$ be the span of all $r[\mathbf{u}_1, \mathbf{v}] + s[\mathbf{u}_2, \mathbf{v}] - [r\mathbf{u}_1 + s\mathbf{u}_2, \mathbf{v}]$; and $r[\mathbf{u}, \mathbf{v}_1] + s[\mathbf{u}, \mathbf{v}_2] - [\mathbf{u}, r\mathbf{v}_1 + s\mathbf{v}_2]$, where $r, s \in \mathbb{F}$, $\mathbf{u}, \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}$, and $\mathbf{v}, \mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}$. Then $\mathcal{U} \otimes \mathcal{V} := \mathbb{F}^{(\mathcal{U}\times\mathcal{V})}/\mathcal{R}$, $t(\mathbf{u}, \mathbf{v}) = [\mathbf{u}, \mathbf{v}] + \mathcal{R}$ is a tensor product of $\mathcal{U}$ and $\mathcal{V}$.*

*Proof.* Let $\pi : \mathbb{F}^{(\mathcal{U}\times\mathcal{V})} \to \mathcal{U} \otimes \mathcal{V}$ be the quotient map.
Since the generators of $\mathcal{R}$ vanish in the quotient, $t(\mathbf{u}, \mathbf{v}) = \pi[\mathbf{u}, \mathbf{v}]$ is bilinear.
Let $f \in \mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{W})$ and define $\alpha : \mathbb{F}^{(\mathcal{U}\times\mathcal{V})} \to \mathcal{W}$ by $\alpha[\mathbf{u}, \mathbf{v}] := f(\mathbf{u}, \mathbf{v})$.
Bilinearity gives $\mathcal{R} \subseteq \ker \alpha$, so Theorem 2.2.10 gives a unique $\widetilde{f} \in \mathcal{L}(\mathcal{U} \otimes \mathcal{V}, \mathcal{W})$ with $\widetilde{f} \circ \pi = \alpha$. Hence $\widetilde{f} \circ t = f$.
If $L \circ t = f$, then $L$ and $\widetilde{f}$ agree on the spanning set $\{[\mathbf{u}, \mathbf{v}] + \mathcal{R}\}$, so $L = \widetilde{f}$.
*Structural Commentary*: The quotient kills precisely the failure of the raw ordered-pair basis to be bilinear.
*Key Insight*: A tensor product is a free bilinear map, followed by no extra identifications. □




Figure 112: The free linear map $\alpha$ kills $\mathcal{R}$, so it descends uniquely to $\widetilde{f} : \mathcal{U} \otimes \mathcal{V} \to \mathcal{W}$.

**Corollary 5.1.5.** *Tensor Products Are Unique up to Isomorphism*
*If $(\mathcal{T}, t)$ and $(\mathcal{S}, s)$ are tensor products of $\mathcal{U}$ and $\mathcal{V}$,*
*then there is a unique isomorphism $\Phi : \mathcal{T} \to \mathcal{S}$ satisfying $\Phi(t(\mathbf{u}, \mathbf{v})) = s(\mathbf{u}, \mathbf{v})$, where $\mathbf{u} \in \mathcal{U},\ \mathbf{v} \in \mathcal{V}$.*

*Proof.* Apply universal property of $(\mathcal{T}, t)$ to bilinear map $s$ and universal property of $(\mathcal{S}, s)$ to bilinear map $t$. This gives linear maps $\Phi : \mathcal{T} \to \mathcal{S}$ and $\Psi : \mathcal{S} \to \mathcal{T}$. The composites $\Psi\Phi$ and $I_{\mathcal{T}}$ are both mediating maps for $t$, while $\Phi\Psi$ and $I_{\mathcal{S}}$ are both mediating maps for $s$. Uniqueness forces $\Psi\Phi = I_{\mathcal{T}}$ and $\Phi\Psi = I_{\mathcal{S}}$.
*Structural Commentary*: Universal objects are unique up to canonical isomorphism.
*Key Insight*: Two universal factorisations force the comparison maps to be inverse. □

**Theorem 5.1.6.** *Bilinearity Equals Linearity on a Tensor Product*
*For $\mathbb{F}$-vector spaces $\mathcal{U}, \mathcal{V}, \mathcal{W}$, the map $\Phi : \mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{W}) \to \mathcal{L}(\mathcal{U} \otimes \mathcal{V}, \mathcal{W})$, $\Phi(f) = \widetilde{f}$,*
*where $\widetilde{f}(\mathbf{u} \otimes \mathbf{v}) = f(\mathbf{u}, \mathbf{v})$, is an isomorphism.*

*Proof.* The universal property gives existence and uniqueness of $\widetilde{f}$.
Define $\Psi : \mathcal{L}(\mathcal{U} \otimes \mathcal{V}, \mathcal{W}) \to \mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{W})$ by $\Psi(L) := L \circ t$.
Then $\Psi(\Phi(f)) = \widetilde{f} \circ t = f$, and $\Phi(\Psi(L)) = L$ by uniqueness of the mediating map for $L \circ t$. Hence $\Phi$ is bijective.
For $a \in \mathbb{F}$ and $f, g$ bilinear, $\Phi(af + g)(\mathbf{u} \otimes \mathbf{v}) = af(\mathbf{u}, \mathbf{v}) + g(\mathbf{u}, \mathbf{v}) = (a\Phi(f) + \Phi(g))(\mathbf{u} \otimes \mathbf{v})$, so $\Phi$ is linear.
*Structural Commentary*: Tensor products package bilinear maps as ordinary linear maps.
*Key Insight*: To define a linear map out of $\mathcal{U} \otimes \mathcal{V}$, define a bilinear rule on $\mathcal{U} \times \mathcal{V}$. □

**Method 5.1.7.** *Defining Maps on Tensor Products*
To define $L \in \mathcal{L}(\mathcal{U} \otimes \mathcal{V}, \mathcal{W})$:

(i) define a candidate rule $f : \mathcal{U} \times \mathcal{V} \to \mathcal{W}$ by $f(\mathbf{u}, \mathbf{v})$;
(ii) prove $f \in \mathrm{Bil}_{\mathbb{F}}(\mathcal{U}, \mathcal{V}; \mathcal{W})$;
(iii) set $L := \Phi(f)$ from Theorem 5.1.6, so $L(\mathbf{u} \otimes \mathbf{v}) = f(\mathbf{u}, \mathbf{v})$.

Do not assign arbitrary values on decomposable tensors: they span $\mathcal{U} \otimes \mathcal{V}$ but are usually linearly dependent.

## 5.2 Tensor Product Bases and Maps

**Definition 5.2.1.** *Finite-Coordinate Tensor Product Model*
Let $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ and $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$ be ordered bases of $\mathcal{V}$ and $\mathcal{W}$.
Define the *coordinate tensor model* $\mathcal{V} \otimes_{\beta,\gamma} \mathcal{W} := \mathrm{Mat}(m, n, \mathbb{F})$ by $\mathbf{v}_i \otimes_{\beta,\gamma} \mathbf{w}_j := E_{i,j}$.
If $\mathbf{v} = \sum_{i=1}^{m} x_i \mathbf{v}_i$ and $\mathbf{w} = \sum_{j=1}^{n} y_j \mathbf{w}_j$, then $\mathbf{v} \otimes_{\beta,\gamma} \mathbf{w} := \sum_{i=1}^{m} \sum_{j=1}^{n} x_i y_j E_{i,j} \in \mathrm{Mat}(m, n, \mathbb{F})$.

**Proposition 5.2.2.** *Tensor Model Basis and Bilinear Evaluation*
*In the coordinate tensor model associated to bases $\beta = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$ and $\gamma = (\mathbf{w}_1, \ldots, \mathbf{w}_n)$:*

*(i) the matrices $E_{i,j} = \mathbf{v}_i \otimes_{\beta,\gamma} \mathbf{w}_j$ form a basis;*
*(ii) the map $(\mathbf{v}, \mathbf{w}) \mapsto \mathbf{v} \otimes_{\beta,\gamma} \mathbf{w}$ is bilinear;*
*(iii) every bilinear form $B : \mathcal{V} \times \mathcal{W} \to \mathbb{F}$ determines a unique linear functional $\Theta_B$ on $\mathrm{Mat}(m, n, \mathbb{F})$ such that $\Theta_B(\mathbf{v} \otimes_{\beta,\gamma} \mathbf{w}) = B(\mathbf{v}, \mathbf{w})$ for all $\mathbf{v} \in \mathcal{V}$ and $\mathbf{w} \in \mathcal{W}$.*

*Proof.*

(i) This is the standard entry-matrix basis of $\mathrm{Mat}(m, n, \mathbb{F})$ from Proposition 3.1.4.
(ii) The coordinates of $a\mathbf{v}_1 + \mathbf{v}_2$ are the same linear combination of the coordinates of $\mathbf{v}_1$ and $\mathbf{v}_2$, and the displayed formula is linear in those coordinates. The same argument applies in the second variable.
(iii) Define $\Theta_B(E_{i,j}) = B(\mathbf{v}_i, \mathbf{w}_j)$ on the basis from (i), and extend linearly by Proposition 1.5.11. If $\mathbf{v} = \sum_i x_i \mathbf{v}_i$ and $\mathbf{w} = \sum_j y_j \mathbf{w}_j$, then $\Theta_B(\mathbf{v} \otimes_{\beta,\gamma} \mathbf{w}) = \sum_{i=1}^{m} \sum_{j=1}^{n} x_i y_j B(\mathbf{v}_i, \mathbf{w}_j) = B(\mathbf{v}, \mathbf{w})$, by bilinearity of $B$.
Uniqueness follows since the values on basis matrices $E_{i,j}$ are forced by $\mathbf{v}_i, \mathbf{w}_j$.

*Structural Commentary*: The coordinate tensor model stores each pair of basis directions in its own matrix slot, so a bilinear form can be evaluated by a single linear functional on those slots.
*Key Insight*: Tensor coordinates turn products of coordinates into linear coordinates. □




Figure 113: A basis pair occupies the $(i, j)$ matrix slot.

**Theorem 5.2.3.** *Tensor Bases, Dimension, and Coordinate Rank*
*Let $\mathcal{U}$ and $\mathcal{V}$ be finite-dimensional with bases $\beta = (\mathbf{u}_1, \dots, \mathbf{u}_m)$ and $\gamma = (\mathbf{v}_1, \dots, \mathbf{v}_n)$.*

*(i) The tensors $\mathbf{u}_i \otimes \mathbf{v}_j$ form a basis of $\mathcal{U} \otimes \mathcal{V}$.*
*(ii)* $\dim(\mathcal{U} \otimes \mathcal{V}) = \dim(\mathcal{U}) \dim(\mathcal{V})$.
*(iii) Every $\mathbf{z} \in \mathcal{U} \otimes \mathcal{V}$ has a unique matrix $R_{\beta,\gamma}(\mathbf{z}) = [r_{ij}] \in \mathrm{Mat}(m, n, \mathbb{F})$ such that $\mathbf{z} = \sum_{i=1}^{m} \sum_{j=1}^{n} r_{ij}(\mathbf{u}_i \otimes \mathbf{v}_j)$.*
*If $\beta', \gamma'$ are other bases, then $\operatorname{rank} R_{\beta',\gamma'}(\mathbf{z}) = \operatorname{rank} R_{\beta,\gamma}(\mathbf{z})$.*

*Proof.*

(i) The coordinate tensor model of Definition 5.2.1 has the universal property in Definition 5.1.3. Hence Corollary 5.1.5 identifies $\mathcal{U} \otimes \mathcal{V} \cong \mathrm{Mat}(m, n, \mathbb{F})$ with $\mathbf{u}_i \otimes \mathbf{v}_j \mapsto E_{i,j}$, and $\{E_{i,j}\}$ is a basis.
(ii) Counting the $mn$ basis tensors gives $\dim(\mathcal{U} \otimes \mathcal{V}) = mn = \dim(\mathcal{U}) \dim(\mathcal{V})$.
(iii) The basis in (i) gives unique coefficients $r_{ij}$, hence $R_{\beta,\gamma}(\mathbf{z})$. If $\beta', \gamma'$ are other bases, then $R_{\beta',\gamma'}(\mathbf{z}) = P R_{\beta,\gamma}(\mathbf{z}) Q$ for invertible change-of-basis matrices $P, Q$, so $\operatorname{rank} R_{\beta',\gamma'}(\mathbf{z}) = \operatorname{rank} R_{\beta,\gamma}(\mathbf{z})$.

*Structural Commentary*: Tensor rank is coordinate-free matrix rank.
*Key Insight*: Changing bases performs row and column changes, which preserve matrix rank. □

**Definition 5.2.4.** *Tensor Coordinate Matrix and Rank*
Let $\mathcal{U}, \mathcal{V}$ be finite-dimensional, with bases $\beta = (\mathbf{u}_1, \dots, \mathbf{u}_m)$ and $\gamma = (\mathbf{v}_1, \dots, \mathbf{v}_n)$.

(i) For $\mathbf{z} = \sum_{i=1}^{m} \sum_{j=1}^{n} r_{ij}(\mathbf{u}_i \otimes \mathbf{v}_j)$, define the *coordinate matrix* $R_{\beta,\gamma}(\mathbf{z}) := [r_{ij}]$.
(ii) The *rank of the tensor* $\mathbf{z}$ is $\operatorname{rank}(\mathbf{z}) := \operatorname{rank} R_{\beta,\gamma}(\mathbf{z})$, independent of $\beta, \gamma$ by Theorem 5.2.3.

**Theorem 5.2.5.** *Zero Tests and Decomposable Rank*
*Let $\mathcal{U}$ and $\mathcal{V}$ be vector spaces.*

*(i)* $\forall \mathbf{u} \in \mathcal{U}, \mathbf{v} \in \mathcal{V}$, $\mathbf{0} \otimes \mathbf{v} = \mathbf{u} \otimes \mathbf{0} = \mathbf{0}$.
*(ii) If $\mathbf{u}_1, \dots, \mathbf{u}_r$ are linearly independent and $\sum_{i=1}^{r} \mathbf{u}_i \otimes \mathbf{v}_i = \mathbf{0}$, then $\mathbf{v}_1 = \cdots = \mathbf{v}_r = \mathbf{0}$.*
*(iii) If $\dim \mathcal{U}, \dim \mathcal{V} < \infty$, then $\mathbf{z} \neq \mathbf{0}$ is decomposable $\iff \operatorname{rank}(\mathbf{z}) = 1$.*
*More generally,* $\operatorname{rank}(\mathbf{z}) = \min \left\{ r \geq 0 : \mathbf{z} = \sum_{i=1}^{r} \mathbf{x}_i \otimes \mathbf{y}_i,\ \mathbf{x}_i \in \mathcal{U}, \mathbf{y}_i \in \mathcal{V} \right\}$.

*Proof.*

(i) The tensor map $t : \mathcal{U} \times \mathcal{V} \to \mathcal{U} \otimes \mathcal{V}$ is bilinear, hence linear in each coordinate; therefore $t(\mathbf{0}, \mathbf{v}) = t(\mathbf{u}, \mathbf{0}) = \mathbf{0}$.
(ii) For each $k$, extend the dual basis functional on $\operatorname{span}(\mathbf{u}_1, \dots, \mathbf{u}_r)$ to $\varphi_k \in \mathcal{U}'$ with $\varphi_k(\mathbf{u}_i) = \delta_{ki}$. The bilinear map $(\mathbf{u}, \mathbf{v}) \mapsto \varphi_k(\mathbf{u})\mathbf{v}$ induces $T_k \in \mathcal{L}(\mathcal{U} \otimes \mathcal{V}, \mathcal{V})$.
Applying $T_k$ gives $\mathbf{0} = T_k\left(\sum_{i=1}^{r} \mathbf{u}_i \otimes \mathbf{v}_i\right) = \sum_{i=1}^{r} \delta_{ki} \mathbf{v}_i = \mathbf{v}_k$.
(iii) Choose bases $\beta, \gamma$. If $\mathbf{z} = \mathbf{x} \otimes \mathbf{y} \neq \mathbf{0}$, then $R_{\beta,\gamma}(\mathbf{z}) = [\mathbf{x}]_\beta [\mathbf{y}]_\gamma^{\mathsf{T}}$, so $\operatorname{rank}(\mathbf{z}) = 1$.
Conversely, if $\operatorname{rank} R_{\beta,\gamma}(\mathbf{z}) = r$, write $R_{\beta,\gamma}(\mathbf{z}) = \sum_{i=1}^{r} a_i b_i^{\mathsf{T}}$ with columns $a_i, b_i$.
Translating $a_i, b_i$ back to vectors $\mathbf{x}_i, \mathbf{y}_i$ gives $\mathbf{z} = \sum_{i=1}^{r} \mathbf{x}_i \otimes \mathbf{y}_i$ as a sum of $r$ decomposable tensors.
Conversely, any sum of $s$ decomposable tensors gives a sum of $s$ rank-one coordinate matrices, so $\operatorname{rank}(\mathbf{z}) \leq s$; hence the displayed minimum equals $\operatorname{rank}(\mathbf{z})$.

*Structural Commentary*: Linear functionals can test one tensor coordinate at a time.
*Key Insight*: Tensor rank is matrix rank after choosing bases, but the number itself survives changing bases. □

**Theorem 5.2.6.** *Tensor Product of Linear Maps*
*Let $A \in \mathcal{L}(\mathcal{U},\mathcal{U}')$ and $B \in \mathcal{L}(\mathcal{V},\mathcal{V}')$.*
*There exists a unique $A \otimes B : \mathcal{U} \otimes \mathcal{V} \to \mathcal{U}' \otimes \mathcal{V}'$ such that $(A \otimes B)(\mathbf{u} \otimes \mathbf{v}) = A\mathbf{u} \otimes B\mathbf{v}$.*
*Moreover, $(A_2A_1) \otimes (B_2B_1) = (A_2 \otimes B_2)(A_1 \otimes B_1)$, $I_{\mathcal{U}} \otimes I_{\mathcal{V}} = I_{\mathcal{U}\otimes\mathcal{V}}$.*
*In finite dimensions, $[A \otimes B]^{\beta'\otimes\gamma'}_{\beta\otimes\gamma} = [A]^{\beta'}_{\beta} \otimes [B]^{\gamma'}_{\gamma}$ for lexicographically ordered tensor-product bases, where the right-hand $\otimes$ is the matrix Kronecker product.*

*Proof.* The map $(\mathbf{u},\mathbf{v}) \mapsto A\mathbf{u} \otimes B\mathbf{v}$ is bilinear, so Theorem 5.1.6 gives the unique $A \otimes B$.
The two functorial identities hold on decomposable tensors $((A_2 \otimes B_2)(A_1 \otimes B_1))(\mathbf{u} \otimes \mathbf{v}) = A_2A_1\mathbf{u} \otimes B_2B_1\mathbf{v}$ and $(I_{\mathcal{U}} \otimes I_{\mathcal{V}})(\mathbf{u} \otimes \mathbf{v}) = \mathbf{u} \otimes \mathbf{v}$, and decomposable tensors span. If $A\mathbf{u}_j = \sum_i a_{ij}\mathbf{u}'_i$ and $B\mathbf{v}_\ell = \sum_k b_{k\ell}\mathbf{v}'_k$, then $(A \otimes B)(\mathbf{u}_j \otimes \mathbf{v}_\ell) = \sum_{i,k} a_{ij}b_{k\ell}(\mathbf{u}'_i \otimes \mathbf{v}'_k)$, which is the Kronecker block-entry rule.
*Structural Commentary*: Tensoring maps acts independently in the two tensor coordinates.
*Key Insight*: The mixed product property for Kronecker products is the coordinate shadow of functoriality. □

**Theorem 5.2.7.** *Tensor Product of Spaces of Linear Maps*
*Let $\mathcal{U},\mathcal{V},\mathcal{X},\mathcal{Y}$ be $\mathbb{F}$-vector spaces. The assignment $(A,B) \mapsto A \otimes B$ induces a natural injective map $\Theta : \mathcal{L}(\mathcal{U},\mathcal{X}) \otimes \mathcal{L}(\mathcal{V},\mathcal{Y}) \to \mathcal{L}(\mathcal{U} \otimes \mathcal{V}, \mathcal{X} \otimes \mathcal{Y})$ with $\Theta(A \otimes B)(\mathbf{u} \otimes \mathbf{v}) = A\mathbf{u} \otimes B\mathbf{v}$.*
*If $\mathcal{U},\mathcal{V},\mathcal{X},\mathcal{Y}$ are finite-dimensional, then $\Theta$ is an isomorphism.*
*In particular, $\mathcal{U}' \otimes \mathcal{V}' \hookrightarrow (\mathcal{U} \otimes \mathcal{V})'$; $(\varphi \otimes \psi)(\mathbf{u} \otimes \mathbf{v}) = \varphi(\mathbf{u})\psi(\mathbf{v})$ is an isomorphism when $\dim\mathcal{U}, \dim\mathcal{V} < \infty$.*

*Proof.*

(i) By Theorem 5.2.6, $(A,B) \mapsto A \otimes B$ is bilinear, hence induces $\Theta$.

(ii) Let $0 \neq Z = \sum_{i=1}^{r} A_i \otimes B_i$ with $r$ minimal. Then $A_i \neq 0$ and $(B_i)$ is linearly independent.
Choose $\mathbf{u}$ with some $A_i\mathbf{u} \neq 0$, after reindexing let $A_1\mathbf{u},\dots,A_m\mathbf{u}$ be basis of $\operatorname{span}\{A_i\mathbf{u} : 1 \le i \le r\}$.
Absorb remaining coefficients into $B_i$ gives $C_1,\dots,C_m$, not all zero, s.t. $\Theta(Z)(\mathbf{u} \otimes \mathbf{v}) = \sum_{i=1}^{m} (A_i\mathbf{u}) \otimes C_i\mathbf{v}$.
If $\Theta(Z) = 0$, Theorem 5.2.5 gives $C_i\mathbf{v} = 0$ for all $i,\mathbf{v}$, contradiction. Thus $\ker\Theta = 0$.

(iii) If $\mathcal{U},\mathcal{V},\mathcal{X},\mathcal{Y}$ are finite-dimensional, then the domain and codomain of $\Theta$ have equal dimension by Theorem 5.2.3, so injectivity implies bijectivity. Taking $\mathcal{X} = \mathcal{Y} = \mathbb{F}$, $\mathcal{L}(\mathcal{U},\mathbb{F}) = \mathcal{U}'$, $\mathcal{L}(\mathcal{U} \otimes \mathcal{V},\mathbb{F}) = (\mathcal{U} \otimes \mathcal{V})'$ gives the dual embedding and finite-dimensional isomorphism.

*Structural Commentary*: Tensoring maps yields maps on tensor products.
*Key Insight*: The rule on decomposable tensors is enough, and finite dimensions make the embedding onto. □

**Proposition 5.2.8.** *Scalar Extension by Tensor Product*
*Let $K/\mathbb{F}$ be a field extension and let $\mathcal{V}_{\mathbb{F}}$ be an $\mathbb{F}$-vector space.*
*Then $K \otimes_{\mathbb{F}} \mathcal{V}_{\mathbb{F}}$ is a $K$-vector space via $\alpha \cdot (\beta \otimes \mathbf{v}) := (\alpha\beta) \otimes \mathbf{v}$. With $\mu : \mathcal{V}_{\mathbb{F}} \to K \otimes_{\mathbb{F}} \mathcal{V}_{\mathbb{F}}$, $\mu(\mathbf{v}) := 1 \otimes \mathbf{v}$, every $\mathbb{F}$-linear $f : \mathcal{V}_{\mathbb{F}} \to \mathcal{Y}$ into a $K$-vector space $\mathcal{Y}$ admits a unique $K$-linear $\widetilde{f} : K \otimes_{\mathbb{F}} \mathcal{V}_{\mathbb{F}} \to \mathcal{Y}$ such that $\widetilde{f} \circ \mu = f$.*

*Proof.* For $\alpha \in K$, $(\beta,\mathbf{v}) \mapsto (\alpha\beta) \otimes \mathbf{v}$ is $\mathbb{F}$-bilinear, hence induces $M_\alpha \in \mathcal{L}_{\mathbb{F}}(K \otimes_{\mathbb{F}} \mathcal{V}_{\mathbb{F}})$.
On decomposable tensors, $M_\alpha M_\lambda = M_{\alpha\lambda}$, $M_1 = I$, and distributivity holds; since decomposable tensors span, these define a $K$-vector-space structure.
For $f$, set $g(\beta,\mathbf{v}) := \beta f(\mathbf{v})$. Then $g$ is $\mathbb{F}$-bilinear, so it induces $\widetilde{f}$ with $\widetilde{f}(\beta \otimes \mathbf{v}) = \beta f(\mathbf{v})$. This is $K$-linear and gives $\widetilde{f} \circ \mu = f$. If $h$ is $K$-linear and $h \circ \mu = f$, then $h(\beta \otimes \mathbf{v}) = \beta h(1 \otimes \mathbf{v}) = \beta f(\mathbf{v})$, so $h = \widetilde{f}$.
*Structural Commentary*: Scalar extension tensors in the larger field.
*Key Insight*: Tensoring with $K$ upgrades which scalars are allowed to act. □

## 5.3 Iterated Tensor Products and Tensor Spaces

The tensor-product universal property extends from bilinear maps to maps with more variables, and then to tensor spaces and contractions.

**Definition 5.3.1.** *Multilinear Maps and Iterated Tensor Products*
Let $\mathcal{V}_1,\dots,\mathcal{V}_n,\mathcal{W}$ be $\mathbb{F}$-vector spaces.

(i) A map $f : \mathcal{V}_1 \times \cdots \times \mathcal{V}_n \to \mathcal{W}$ is *multilinear*, or *$n$-linear*, if for each $k$, the map $\mathbf{v}_k \mapsto f(\mathbf{v}_1,\dots,\mathbf{v}_k,\dots,\mathbf{v}_n)$ is linear with the other entries fixed.

(ii) $\operatorname{Mult}_{\mathbb{F}}(\mathcal{V}_1,\dots,\mathcal{V}_n;\mathcal{W}) := \{f : \mathcal{V}_1 \times \cdots \times \mathcal{V}_n \to \mathcal{W} : f \text{ multilinear}\}$.

(iii) If $\mathcal{W} = \mathbb{F}$, elements of $\operatorname{Mult}_{\mathbb{F}}(\mathcal{V}_1,\dots,\mathcal{V}_n;\mathbb{F})$ are *multilinear forms*.

(iv) A pair $(\mathcal{T},t)$ is an *iterated tensor product* of $\mathcal{V}_1,\dots,\mathcal{V}_n$ if $t \in \operatorname{Mult}_{\mathbb{F}}(\mathcal{V}_1,\dots,\mathcal{V}_n;\mathcal{T})$ and, for every $\mathbb{F}$-vector space $\mathcal{Y}$ and $f \in \operatorname{Mult}_{\mathbb{F}}(\mathcal{V}_1,\dots,\mathcal{V}_n;\mathcal{Y})$, $\exists!\widetilde{f} \in \mathcal{L}(\mathcal{T},\mathcal{Y})$ such that $f = \widetilde{f} \circ t$.

(v) Write $\mathcal{T} = \mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n$ and $\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_n := t(\mathbf{v}_1,\dots,\mathbf{v}_n)$.

**Example 5.3.2.** *Basic Multilinear Maps*

(i) If $A$ is an associative $\mathbb{F}$-algebra, then $m_n : A^n \to A$, $m_n(a_1, \dots, a_n) := a_1 \cdots a_n$, is $n$-linear.
(ii) If $\dim \mathcal{V} = n$, then $\det : \mathcal{V}^n \to \mathbb{F}$, applied to coordinate columns in any basis, is alternating $n$-linear form.




Figure 114: The iterated tensor product is universal for multilinear maps.

**Theorem 5.3.3.** *Basic Laws for Iterated Tensor Products*
*Let $\mathcal{V}_1, \dots, \mathcal{V}_n, \mathcal{V}, \mathcal{W}$ be $\mathbb{F}$-vector spaces.*

*(i)* $\mathrm{Mult}_{\mathbb{F}}(\mathcal{V}_1, \dots, \mathcal{V}_n; \mathcal{W}) \cong \mathcal{L}(\mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n, \mathcal{W})$ *naturally.*
*(ii) For $1 \le r < n$, the canonical $\alpha_r$ is $\alpha_r : (\mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_r) \otimes (\mathcal{V}_{r+1} \otimes \cdots \otimes \mathcal{V}_n) \cong \mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n$, where $\alpha_r((\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_r) \otimes (\mathbf{v}_{r+1} \otimes \cdots \otimes \mathbf{v}_n)) = \mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_n$.*
*(iii) For $\pi \in S_n$, the canonical $\tau_\pi$ is $\tau_\pi : \mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n \cong \mathcal{V}_{\pi(1)} \otimes \cdots \otimes \mathcal{V}_{\pi(n)}$, where $\tau_\pi(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_n) = \mathbf{v}_{\pi(1)} \otimes \cdots \otimes \mathbf{v}_{\pi(n)}$.*
*(iv) $\lambda : \mathbb{F} \otimes \mathcal{V} \cong \mathcal{V}$, $\lambda(r \otimes \mathbf{v}) = r\mathbf{v}$, and $\rho : \mathcal{V} \otimes \mathbb{F} \cong \mathcal{V}$, $\rho(\mathbf{v} \otimes r) = r\mathbf{v}$.*
*(v) If $\dim \mathcal{V}_i < \infty$ for all $i$, then $\dim(\mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n) = \prod_{i=1}^n \dim(\mathcal{V}_i)$.*

*Proof.*

(i) Let $t : \mathcal{V}_1 \times \cdots \times \mathcal{V}_n \to \mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n$ be canonical. Define $\Phi(f) = \widetilde{f}$, where $\widetilde{f}$ is the unique linear map satisfying $\widetilde{f}(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_n) = f(\mathbf{v}_1, \dots, \mathbf{v}_n)$; conversely define $\Psi(T) = T \circ t$. Then $\Psi \circ \Phi$ and $\Phi \circ \Psi$ are identity maps, so $\Phi$ is a linear bijection.
(ii) Set $A_r = (\mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_r) \otimes (\mathcal{V}_{r+1} \otimes \cdots \otimes \mathcal{V}_n)$ and $B = \mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n$. The rules defining $\alpha_r : A_r \to B$ and $\beta_r : B \to A_r$, $\beta_r(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_n) = (\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_r) \otimes (\mathbf{v}_{r+1} \otimes \cdots \otimes \mathbf{v}_n)$, are induced by multilinearity; $\alpha_r \circ \beta_r$ and $\beta_r \circ \alpha_r$ fix the spanning decomposable tensors, hence are identities.
(iii) Set $C_\pi = \mathcal{V}_{\pi(1)} \otimes \cdots \otimes \mathcal{V}_{\pi(n)}$. The map $p_\pi : \mathcal{V}_1 \times \cdots \times \mathcal{V}_n \to C_\pi$, $p_\pi(\mathbf{v}_1, \dots, \mathbf{v}_n) = \mathbf{v}_{\pi(1)} \otimes \cdots \otimes \mathbf{v}_{\pi(n)}$, is multilinear, so the universal property induces the unique $\tau_\pi$ with $\tau_\pi \circ t = p_\pi$.
The map $\sigma_\pi : C_\pi \to \mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n$, $\sigma_\pi(\mathbf{v}_{\pi(1)} \otimes \cdots \otimes \mathbf{v}_{\pi(n)}) = \mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_n$, is induced similarly; $\sigma_\pi \tau_\pi$ and $\tau_\pi \sigma_\pi$ fix decomposable spanning tensors, hence are identities.
(iv) The maps $m_L : \mathbb{F} \times \mathcal{V} \to \mathcal{V}$, $m_L(r, \mathbf{v}) = r\mathbf{v}$, and $m_R : \mathcal{V} \times \mathbb{F} \to \mathcal{V}$, $m_R(\mathbf{v}, r) = r\mathbf{v}$, are bilinear, hence induce $\lambda$ and $\rho$. Since $\lambda(1 \otimes \mathbf{v}) = \mathbf{v}$ and $1 \otimes r\mathbf{v} = r \otimes \mathbf{v}$, $\lambda^{-1}(\mathbf{v}) = 1 \otimes \mathbf{v}$.
Since $\rho(\mathbf{v} \otimes 1) = \mathbf{v}$ and $r\mathbf{v} \otimes 1 = \mathbf{v} \otimes r$, $\rho^{-1}(\mathbf{v}) = \mathbf{v} \otimes 1$.
(v) Choose finite bases $B_i$ of $\mathcal{V}_i$ and set $\mathcal{B} = \{\mathbf{b}_1 \otimes \cdots \otimes \mathbf{b}_n : \mathbf{b}_i \in B_i\}$. By repeated application of Theorem 5.2.3, $\mathcal{B}$ is a basis; hence $\dim(\mathcal{V}_1 \otimes \cdots \otimes \mathcal{V}_n) = \#\mathcal{B} = \prod_{i=1}^n \#B_i = \prod_{i=1}^n \dim \mathcal{V}_i$.

*Structural Commentary*: Tensor parenthesising and ordering are canonical bookkeeping.
*Key Insight*: Universal properties extend pure-tensor rules uniquely. □

**Definition 5.3.4.** *Tensor Spaces of Type $(p, q)$*
Let $\dim \mathcal{V} < \infty$.

(i) For $(p, q) \in \mathbb{Z}^2_{\ge 0} \setminus \{(0, 0)\}$, with empty tensor powers equal to $\mathbb{F}$, $\mathcal{T}^p_q(\mathcal{V}) := \mathcal{V}^{\otimes p} \otimes (\mathcal{V}')^{\otimes q}$.
(ii) $\mathcal{T}^p_q(\mathcal{V})$ is the *tensor space of type $(p, q)$*; $p$ is the *contravariant type*, $q$ the *covariant type*.
(iii) $\mathcal{T}^0_0(\mathcal{V}) := \mathbb{F}$.

**Proposition 5.3.5.** *Tensor Spaces as Multilinear Functionals*
*Let $\dim \mathcal{V} = n < \infty$ and $p, q \in \mathbb{Z}_{\ge 0}$.*

*(i)* $\dim \mathcal{T}^p_q(\mathcal{V}) = n^{p+q}$.
*(ii) Let $\iota : \mathcal{V} \to \mathcal{V}''$, $\iota(\mathbf{v})(\eta) = \eta(\mathbf{v})$. Via $\iota$, natural isomorphism is $\Theta_{p,q} : \mathcal{T}^p_q(\mathcal{V}) \xrightarrow{\sim} \mathrm{Mult}_{\mathbb{F}}((\mathcal{V}')^{\times p} \times \mathcal{V}^{\times q}; \mathbb{F})$; where $\Theta_{p,q}(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_p \otimes \varphi_1 \otimes \cdots \otimes \varphi_q)(\eta_1, \dots, \eta_p, \mathbf{x}_1, \dots, \mathbf{x}_q) = \prod_{i=1}^p \eta_i(\mathbf{v}_i) \prod_{j=1}^q \varphi_j(\mathbf{x}_j)$, with empty products equal to 1.*

*Proof.*

(i) By Theorem 5.3.3, $\dim \mathcal{V}' = n$, $\dim \mathcal{T}^p_q(\mathcal{V}) = \dim(\mathcal{V}^{\otimes p}) \dim((\mathcal{V}')^{\otimes q}) = n^p n^q = n^{p+q}$, with $n^0 = 1$.

(ii) Set $X = (\mathcal{V}')^{\otimes p} \otimes \mathcal{V}^{\otimes q}$. Since $\iota : \mathcal{V} \cong \mathcal{V}''$ and all factors are finite-dimensional, $\mathcal{T}^p_q(\mathcal{V}) = \mathcal{V}^{\otimes p} \otimes (\mathcal{V}')^{\otimes q} \cong ((\mathcal{V}')')^{\otimes p} \otimes \mathcal{V}'^{\otimes q} \cong X' \cong \mathrm{Mult}_{\mathbb{F}}((\mathcal{V}')^{\times p} \times \mathcal{V}^{\times q}; \mathbb{F})$. On decomposables, the composite is $\Theta_{p,q}$; spanning gives uniqueness.

*Structural Commentary*: Tensor elements are multilinear functionals after finite-dimensional duality.
*Key Insight*: Choosing which factors are active turns tensor factors into input slots. □

**Definition 5.3.6.** *Contraction*
Let $p, q \geq 1$. The *contraction* is the unique linear map $C : \mathcal{T}^p_q(\mathcal{V}) \to \mathcal{T}^{p-1}_{q-1}(\mathcal{V})$, where $C(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_p \otimes \varphi_1 \otimes \cdots \otimes \varphi_q) = \varphi_1(\mathbf{v}_1)\mathbf{v}_2 \otimes \cdots \otimes \mathbf{v}_p \otimes \varphi_2 \otimes \cdots \otimes \varphi_q$.
For $(i, j)$, $C_{ij}$ is $C$ conjugated by the factor permutation placing $\mathbf{v}_i, \varphi_j$ first.

**Example 5.3.7.** *Rank-One Operators and Trace*
Under $\mathcal{V} \otimes \mathcal{V}' \cong \mathcal{L}(\mathcal{V})$, $\mathbf{v} \otimes \varphi \mapsto T_{\mathbf{v},\varphi}$, where $T_{\mathbf{v},\varphi}(\mathbf{x}) = \varphi(\mathbf{x})\mathbf{v}$. Hence $C(\mathbf{v} \otimes \varphi) = \varphi(\mathbf{v}) = \mathrm{tr}(T_{\mathbf{v},\varphi})$.

# 6 Symmetric and Exterior Algebras

## 6.1 Symmetric Algebras

**Definition 6.1.1.** *Symmetric, Skew-Symmetric, and Alternating Maps*
Let $f : \mathcal{V}^{\times n} \to \mathcal{W}$ be multilinear.

(i) $f$ is *symmetric* if interchanging any two coordinate positions does not change its value.
(ii) $f$ is *skew-symmetric* if interchanging any two coordinate positions multiplies its value by $-1$.
(iii) $f$ is *alternating* if $f(\mathbf{v}_1, \ldots, \mathbf{v}_n) = 0$ whenever two coordinate entries are equal.

If $\mathrm{char}(\mathbb{F}) \neq 2$, alternating and skew-symmetric are equivalent.
If $\mathrm{char}(\mathbb{F}) = 2$, skew-symmetric is the same as symmetric, and alternating is the stronger condition.

**Definition 6.1.2.** *Permutation Action and Symmetric/Exterior Tensor Spaces*
Let $\dim \mathcal{V} < \infty$, $p \in \mathbb{Z}_{\geq 0}$, $\mathcal{V}^{\otimes 0} = \mathbb{F}$, $T_p := \mathcal{V}^{\otimes p}$, and $G_p := \mathfrak{S}_p$.

(i) For $\sigma \in G_p$, the *permutation action* is $\lambda_\sigma : T_p \to T_p$, determined by $\lambda_\sigma(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_p) = \mathbf{v}_{\sigma(1)} \otimes \cdots \otimes \mathbf{v}_{\sigma(p)}$.
(ii) The *symmetric tensors* are $\mathrm{ST}^p(\mathcal{V}) := \{\mathbf{z} \in T_p : \forall \sigma \in G_p, \lambda_\sigma \mathbf{z} = \mathbf{z}\}$.
(iii) The *antisymmetric tensors* are $\mathrm{AT}^p(\mathcal{V}) := \{\mathbf{z} \in T_p : \forall \sigma \in G_p, \lambda_\sigma \mathbf{z} = \mathrm{sign}(\sigma)\mathbf{z}\}$.
(iv) Set $R_\vee := \mathrm{span}\{\lambda_\sigma \mathbf{z} - \mathbf{z} : \sigma \in G_p, \mathbf{z} \in T_p\}$.
(v) The *symmetric power* is $\mathrm{Sym}^p(\mathcal{V}) := T_p / R_\vee$.
(vi) Set $A_p := \mathrm{span}\{\lambda_\sigma \mathbf{z} - \mathrm{sign}(\sigma)\mathbf{z} : \sigma \in G_p, \mathbf{z} \in T_p\}$ and $D_p := \mathrm{span}\{\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_p : \exists i < j, \mathbf{v}_i = \mathbf{v}_j\}$. The *exterior power* is $\bigwedge^p \mathcal{V} := T_p/(A_p + D_p)$.
(vii) For quotient map $q_\vee$, the *symmetric product* is $q_\vee(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_p) = \mathbf{v}_1 \vee \cdots \vee \mathbf{v}_p$
(viii) For quotient map $q_\wedge$, the *exterior product* is $q_\wedge(\mathbf{v}_1 \otimes \cdots \otimes \mathbf{v}_p) = \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_p$.

**Theorem 6.1.3.** *Universal Properties of Symmetric and Exterior Powers*
*Let $\mathcal{U}$ be a vector space, $M = \mathrm{Mult}_{\mathbb{F}}(\mathcal{V}, \ldots, \mathcal{V}; \mathcal{U})$, and $f^\sigma(\mathbf{v}_1, \ldots, \mathbf{v}_p) := f(\mathbf{v}_{\sigma(1)}, \ldots, \mathbf{v}_{\sigma(p)})$.*

*(i)* $\mathcal{L}(\mathrm{Sym}^p(\mathcal{V}), \mathcal{U}) \cong M_\vee$, *where* $M_\vee := \{f \in M : \forall \sigma \in \mathfrak{S}_p,\ f^\sigma = f\}$.
*(ii)* $\mathcal{L}(\bigwedge^p \mathcal{V}, \mathcal{U}) \cong M_\wedge$, *where* $M_\wedge := \{f \in M : \forall \sigma \in \mathfrak{S}_p,\ f^\sigma = \mathrm{sign}(\sigma) f,\ \forall (\mathbf{v}_1, \ldots, \mathbf{v}_p), (\exists i < j, \mathbf{v}_i = \mathbf{v}_j) \Rightarrow f(\mathbf{v}_1, \ldots, \mathbf{v}_p) = 0\}$.

*Proof.*

(i) Let $t : \mathcal{V}^{\times p} \to \mathcal{V}^{\otimes p}$ and $q_\vee : \mathcal{V}^{\otimes p} \to \mathrm{Sym}^p(\mathcal{V})$ be canonical. The map $\Phi_\vee(T) = T q_\vee t$ sends $\mathcal{L}(\mathrm{Sym}^p(\mathcal{V}), \mathcal{U})$ into $M_\vee$. Conversely, for $f \in M_\vee$, there is a unique $\widetilde{f} : \mathcal{V}^{\otimes p} \to \mathcal{U}$ with $f = \widetilde{f} t$; since $\widetilde{f}(R_\vee) = 0$, it factors uniquely through $q_\vee$ as $\widetilde{f} = \overline{f} q_\vee$, where $\overline{f} : \mathrm{Sym}^p(\mathcal{V}) \to \mathcal{U}$. Thus $f \mapsto \overline{f}$ is $\Phi_\vee^{-1}$.
(ii) Let $t : \mathcal{V}^{\times p} \to \mathcal{V}^{\otimes p}$ and $q_\wedge : \mathcal{V}^{\otimes p} \to \bigwedge^p \mathcal{V}$ be canonical. The map $\Phi_\wedge(T) = T q_\wedge t$ sends $\mathcal{L}(\bigwedge^p \mathcal{V}, \mathcal{U})$ into $M_\wedge$. Conversely, for $f \in M_\wedge$, the induced $\widetilde{f} : \mathcal{V}^{\otimes p} \to \mathcal{U}$ satisfies $\widetilde{f}(R_\wedge) = 0$, hence factors uniquely as $\widetilde{f} = \overline{f} q_\wedge$; thus $f \mapsto \overline{f}$ is $\Phi_\wedge^{-1}$.

*Structural Commentary*: Symmetric and exterior powers are tensor quotients by permutation relations.
*Key Insight*: Symmetric and alternating multilinear maps are ordinary linear maps out of the right quotient. □

**Theorem 6.1.4.** *Dimensions of Symmetric and Exterior Powers*
*Let $\dim \mathcal{V} = n < \infty$ and $p \in \mathbb{Z}_{\geq 0}$. Set $I_\vee = \{1 \leq i_1 \leq \cdots \leq i_p \leq n\}$ and $I_\wedge = \{1 \leq i_1 < \cdots < i_p \leq n\}$.*

*(i)* $\dim \mathrm{Sym}^p(\mathcal{V}) = \#I_\vee = \binom{n+p-1}{p}$.
*(ii)* $\dim \bigwedge^p \mathcal{V} = \#I_\wedge = \binom{n}{p}$, *with* $\binom{n}{p} = 0$ *for* $p > n$.

*Proof.* Let $E = (\mathbf{e}_1, \dots, \mathbf{e}_n)$ be a basis. For $I = (i_1, \dots, i_p)$, set $\mathbf{e}_I^{\vee} = \mathbf{e}_{i_1} \vee \cdots \vee \mathbf{e}_{i_p}$ and $\mathbf{e}_I^{\wedge} = \mathbf{e}_{i_1} \wedge \cdots \wedge \mathbf{e}_{i_p}$.

(i) $R_{\vee}$ identifies each word with its nondecreasing representative. Thus $B_{\vee} = \{\mathbf{e}_I^{\vee} : I \in I_{\vee}\}$ is the quotient basis, and $\dim \mathrm{Sym}^p(\mathcal{V}) = \#B_{\vee} = \#I_{\vee} = \binom{n+p-1}{p}$.

(ii) $R_{\wedge}$ kills repeated words and identifies each remaining word with its increasing representative up to sign. Thus $B_{\wedge} = \{\mathbf{e}_I^{\wedge} : I \in I_{\wedge}\}$ is the quotient basis, and $\dim \bigwedge^p \mathcal{V} = \#B_{\wedge} = \#I_{\wedge} = \binom{n}{p}$.

*Structural Commentary*: Symmetric powers count repetitions; exterior powers forbid repetitions.
*Key Insight*: Multisets give symmetric dimensions, subsets give exterior dimensions. □

**Proposition 6.1.5.** *Symmetrization in Characteristic Zero*
*Assume* $\mathrm{char}(\mathbb{F}) = 0$. *Define* $S, A \in \mathcal{L}(\mathcal{V}^{\otimes p})$ *by* $S := \frac{1}{p!} \sum_{\sigma \in \mathfrak{S}_p} \lambda_\sigma$ *and* $A := \frac{1}{p!} \sum_{\sigma \in \mathfrak{S}_p} \mathrm{sign}(\sigma) \lambda_\sigma$.

*(i)* $S^2 = S$ *and* $\mathrm{im}\, S = \mathrm{ST}^p(\mathcal{V})$.
*(ii)* $A^2 = A$ *and* $\mathrm{im}\, A = \mathrm{AT}^p(\mathcal{V})$.

*Proof.*

(i) For $\tau \in \mathfrak{S}_p$, $\lambda_\tau S = p!^{-1} \sum_\sigma \lambda_{\tau\sigma} = S$, hence $\mathrm{im}\, S \subseteq \mathrm{ST}^p(\mathcal{V})$.
If $\mathbf{z} \in \mathrm{ST}^p(\mathcal{V})$, then $S\mathbf{z} = \mathbf{z}$, so $\mathrm{im}\, S = \mathrm{ST}^p(\mathcal{V})$ and $S^2 = S$.

(ii) For $\tau \in \mathfrak{S}_p$, $\lambda_\tau A = p!^{-1} \sum_\sigma \mathrm{sign}(\sigma) \lambda_{\tau\sigma} = \mathrm{sign}(\tau) A$, hence $\mathrm{im}\, A \subseteq \mathrm{AT}^p(\mathcal{V})$.
If $\mathbf{z} \in \mathrm{AT}^p(\mathcal{V})$, then $A\mathbf{z} = p!^{-1} \sum_\sigma \mathrm{sign}(\sigma)^2 \mathbf{z} = \mathbf{z}$, so $\mathrm{im}\, A = \mathrm{AT}^p(\mathcal{V})$ and $A^2 = A$.

*Structural Commentary*: Averaging gives symmetry projections.
*Key Insight*: The factor $1/p!$ makes the averaging map fix the tensors it projects onto. □

**Definition 6.1.6.** *Tensor, Symmetric, and Exterior Algebras*
Let $\dim \mathcal{V} < \infty$.

(i) The *tensor algebra* is $\mathcal{T}(\mathcal{V}) := \bigoplus_{p \geq 0} \mathcal{V}^{\otimes p}$, with $\mathcal{V}^{\otimes 0} := \mathbb{F}$ and product induced by tensor concatenation.
(ii) The *symmetric algebra* is $\mathrm{Sym}(\mathcal{V}) := \bigoplus_{p \geq 0} \mathrm{Sym}^p(\mathcal{V})$, with product induced by $\vee$.
(iii) The *exterior algebra* is $\bigwedge \mathcal{V} := \bigoplus_{0 \leq p \leq \dim \mathcal{V}} \bigwedge^p \mathcal{V}$, with product induced by $\wedge$.
(iv) Each is *graded*: degree $p$ times degree $q$ lies in degree $p + q$.

## 6.2 Exterior Algebras

**Corollary 6.2.1.** *Determinant from the Exterior Universal Property*
*Let* $\dim \mathcal{V} = n$, $E = (\mathbf{e}_1, \dots, \mathbf{e}_n)$, $\Omega_E := \mathbf{e}_1 \wedge \cdots \wedge \mathbf{e}_n$, *and* $A_E(\mathbf{v}_1, \dots, \mathbf{v}_n) = ([\mathbf{v}_j]_i)_{i,j}$.

*(i) There is a unique* $\ell_E \in (\bigwedge^n \mathcal{V})'$ *with* $\ell_E(\Omega_E) = 1$*; hence* $d_E(\mathbf{v}_1, \dots, \mathbf{v}_n) := \ell_E(\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_n)$ *is the unique alternating form satisfying* $d_E(\mathbf{e}_1, \dots, \mathbf{e}_n) = 1$.

*(ii) For all* $\mathbf{v}_1, \dots, \mathbf{v}_n \in \mathcal{V}$, $d_E(\mathbf{v}_1, \dots, \mathbf{v}_n) = \sum_{\sigma \in \mathfrak{S}_n} \mathrm{sign}(\sigma) \prod_{j=1}^n [\mathbf{v}_j]_{\sigma(j)} = \det A_E(\mathbf{v}_1, \dots, \mathbf{v}_n)$.

*Proof.*

(i) By Theorem 6.1.4, $\bigwedge^n \mathcal{V} = \mathbb{F}\Omega_E$, so $\ell_E(\Omega_E) = 1$ fixes a unique $\ell_E \in (\bigwedge^n \mathcal{V})'$; via Theorem 6.1.3, $\ell_E$ corresponds to $d_E(\mathbf{v}_1, \dots, \mathbf{v}_n) = \ell_E(\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_n)$, hence $d_E$ is unique.

(ii) Let $\mathbf{v}_j = \sum_{i=1}^n a_{ij} \mathbf{e}_i$, $a_{ij} = [\mathbf{v}_j]_i$. Repeated index tuples have zero wedge; distinct tuples are $(\sigma(1), \dots, \sigma(n))$, with wedge $\mathrm{sign}(\sigma)\Omega_E$. Hence $\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_n = \sum_{\sigma \in \mathfrak{S}_n} \mathrm{sign}(\sigma) \prod_{j=1}^n a_{\sigma(j),j} \Omega_E$, and

$$d_E(\mathbf{v}_1, \dots, \mathbf{v}_n) = \ell_E(\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_n) = \sum_{\sigma \in \mathfrak{S}_n} \mathrm{sign}(\sigma) \prod_{j=1}^n a_{\sigma(j),j}$$
$$= \det(a_{ij}) = \det A_E(\mathbf{v}_1, \dots, \mathbf{v}_n).$$

*Structural Commentary*: Determinant is the basis-normalised alternating form.
*Key Insight*: Top exterior dimension one turns normalisation into determinant. □

**Corollary 6.2.2.** *Determinant Properties from Exterior Powers*
*Let* $A, B \in \mathrm{Mat}(n, n, \mathbb{F})$, *and set* $d(A) := d_E(A_{\bullet 1}, \dots, A_{\bullet n})$ *for* $E$ *the standard basis of* $\mathbb{F}^n$.

*(i)* $d(A^t) = d(A)$.
*(ii)* $d(AB) = d(A)d(B)$.
*(iii)* $A \in \mathrm{GL}(n, \mathbb{F}) \iff d(A) \neq 0$.

*Proof.* Let $\omega = \mathbf{e}_1 \wedge \cdots \wedge \mathbf{e}_n$ and let $T_A$ denote the linear map with matrix $A$.

(i) By Corollary 6.2.1, $d(A^t) = \sum_\sigma \mathrm{sign}(\sigma) \prod_{j=1}^n A_{j,\sigma(j)}$. With $\tau = \sigma^{-1}$, $\sum_\tau \mathrm{sign}(\tau) \prod_{i=1}^n A_{\tau(i),i} = d(A)$.

(ii) Since $(\bigwedge^n T_A)\omega = d(A)\omega$ and $T_{AB} = T_A \circ T_B$, applying $\bigwedge^n(T_A \circ T_B) = (\bigwedge^n T_A)(\bigwedge^n T_B)$ to $\omega$ gives $d(AB)\omega = d(A)d(B)\omega$, hence $d(AB) = d(A)d(B)$.

*(iii)* ($\Rightarrow$) If $A \in \mathrm{GL}(n, \mathbb{F})$, then $I = AA^{-1}$, so $1 = d(I) = d(A)d(A^{-1})$ by (ii); hence $d(A) \neq 0$.

*(iii)* ($\Leftarrow$) Let $C_j = A_{\bullet j}$. Suppose $\sum_{j=1}^n c_j C_j = 0$ and $c_k \neq 0$. Then $C_k = -c_k^{-1} \sum_{j \neq k} c_j C_j$, so multilinearity in the $k$th slot gives $d(A) = d(C_1, \ldots, C_k, \ldots, C_n) = -c_k^{-1} \sum_{j \neq k} c_j d(C_1, \ldots, C_{k-1}, C_j, C_{k+1}, \ldots, C_n) = 0$, since each summand has equal $j$th and $k$th entries.
Thus $d(A) \neq 0$ implies $C_1, \ldots, C_n$ are independent, hence $A \in \mathrm{GL}(n, \mathbb{F})$.

*Structural Commentary*: Determinant laws are top-exterior scalar laws.
*Key Insight*: Top exterior dimension one makes determinants multiplicative. □

## 6.3 Multilinear Algebra Revisited

# 7 Chain Complexes and Homology

## 7.1 Chain Complexes

## 7.2 Homology

## 7.3 Derived Functors Preview

# Part IX
# Commutative Algebra


## 1 Prime Ideals and Spectra

### 1.1 Prime Ideals

### 1.2 Spectra

## 2 Localization and Local Rings

### 2.1 Localization

### 2.2 Local Rings

## 3 Noetherian Rings and Modules

### 3.1 Noetherian Rings


**Theorem 3.1.1.** *Hilbert Basis Theorem*
*Let $A$ be commutative with identity. If $A$ is Noetherian, then $A[x]$ is Noetherian.*

*Proof.* It suffices to show every ideal $\mathcal{J} \lhd A[x]$ is f.g. If $\mathcal{J} = 0$, done.
For $g \neq 0$, write $\mathrm{lc}(g)$ and $\mathrm{lt}(g)$. Set $L = \langle \mathrm{lc}(g) : g \in \mathcal{J}, g \neq 0 \rangle \lhd A$. Since $A$ is Noetherian, choose $a_1, \dots, a_m$ generating $L$ and $f_i \in \mathcal{J}$ with $\mathrm{lc}(f_i) = a_i$. Replacing $f_i$ by powers of $x$ times $f_i$, assume $\deg f_i = d$ for all $i$.
For $0 \leq k < d$, set $L_k = \langle \mathrm{lc}(g) : g \in \mathcal{J}, \deg g = k \rangle$ and choose finite $P_k \subseteq \mathcal{J}$ with $\deg p = k$ for $p \in P_k$ and $\{\mathrm{lc}(p) : p \in P_k\}$ generating $L_k$. Put $G = \{f_1, \dots, f_m\} \cup \bigcup_{k=0}^{d-1} P_k$ and $\mathcal{A} = \langle G \rangle \lhd A[x]$. Then $\mathcal{A} \subseteq \mathcal{J}$.
For $g \in \mathcal{J} \setminus \mathcal{A}$ of minimal degree, set $r = \deg g$. If $r < d$, write $\mathrm{lc}(g) = \sum_{p \in P_r} c_p \, \mathrm{lc}(p)$ and set $h = \sum_{p \in P_r} c_p p$.
If $r \geq d$, write $\mathrm{lc}(g) = \sum_{i=1}^{m} c_i a_i$ and set $h = \sum_{i=1}^{m} c_i x^{r-d} f_i$. In either case $h \in \mathcal{A}$ and $\mathrm{lt}(h) = \mathrm{lt}(g)$.
Then $g - h \in \mathcal{J} \setminus \mathcal{A}$ and $\deg(g - h) < \deg g$, contradiction. Thus $\mathcal{J} = \mathcal{A}$ is f.g.
*Structural Commentary*: Ideals in $A[x]$ are controlled by finitely many leading coefficients in $A$.
*Key Insight*: Match the leading term, subtract, and induct on degree. □

### 3.2 Noetherian Modules

**Theorem 3.2.1.** *Finitely Generated Modules over Noetherian Rings*
*Let $A$ be a commutative ring with identity. Then $A$ is Noetherian $\iff$ every f.g. $A$-module is Noetherian.*

*Proof.*

($\Rightarrow$) Assume $A$ Noetherian. First prove by induction on $n$ that every $S \leq A^n$ is f.g.
The case $n = 1$ is Theorem 2.4.2. For $n > 1$, let $\pi : S \to A$ be last-coordinate projection.
Then $\pi(S) \lhd A$ is f.g. and $\ker(\pi) \leq A^{n-1}$ is f.g. by induction. Lifts of generators of $\pi(S)$, together with generators of $\ker(\pi)$, generate $S$. Thus $A^n$ is Noetherian. If $M$ is f.g., choose $A^n \twoheadrightarrow M$; for $N \leq M$, its preimage in $A^n$ is f.g., hence $N$ is f.g. Therefore $M$ is Noetherian.

($\Leftarrow$) Since $A = \langle\langle 1 \rangle\rangle_A$, the regular $A$-module $A$ is f.g.; by hypothesis it is Noetherian. Hence $A$ is Noetherian.

*Structural Commentary*: Noetherianity passes from rings to finite free modules to quotients.
*Key Insight*: Finitely generated modules are quotients of finite free modules. □


## 4 Integral Dependence and Dimension

### 4.1 Integral Dependence

### 4.2 Dimension Theory

## 5 Regularity and Singularities

### 5.1 Regularity

### 5.2 Singularities Preview

# Part X
# Algebraic Geometry


## 1 Affine Algebraic Sets

### 1.1 Affine Algebraic Sets

### 1.2 Ideals and Coordinate Rings

## 2 Morphisms and Coordinate Rings

### 2.1 Regular Maps

### 2.2 Coordinate Ring Functoriality

## 3 Projective Geometry

### 3.1 Projective Space

### 3.2 Projective Varieties

## 4 Schemes

### 4.1 Affine Schemes

### 4.2 Gluing and Scheme Morphisms

## 5 Sheaves and Cohomology

### 5.1 Sheaves

### 5.2 Cohomology Preview

## 6 Smoothness, Tangent Spaces, and Dimension

### 6.1 Smoothness

### 6.2 Tangent Spaces

### 6.3 Dimension

## 7 Divisors, Line Bundles, and Curves

### 7.1 Divisors

### 7.2 Line Bundles

### 7.3 Curves

# Part XI
# Real and Topological Analysis

## 1 Number Systems and Completeness

### 1.1 Ordered Fields, Suprema, and Real Structure

Number systems supply the ordered and algebraic background for Analysis: $\mathbb{Q}$ has gaps, $\mathbb{R}$ repairs them by completeness, and $\mathbb{C}$ keeps field algebra without order.

**Definition 1.1.1.** *Natural, Rational, Irrational, and Dense Subsets*
A set $E \subseteq \mathbb{R}$ is *inductive* iff $1 \in E$ and, for all $x \in \mathbb{R}$, $x \in E$ implies $x + 1 \in E$.
Let $\mathcal{I}$ be the family of inductive subsets of $\mathbb{R}$.

(i) $\mathbb{N} := \bigcap_{E \in \mathcal{I}} E$, the least inductive subset of $\mathbb{R}$.
(ii) $\mathbb{Z} := \mathbb{N} \cup \{0\} \cup \{-n : n \in \mathbb{N}\}$.
(iii) $\mathbb{Q} := \{m/n : m, n \in \mathbb{Z},\ n \neq 0\}$; $x \in \mathbb{R}$ is *irrational* iff $x \notin \mathbb{Q}$.
(iv) $D \subseteq \mathbb{R}$ is *dense in* $\mathbb{R}$ iff for all $a, b \in \mathbb{R}$ with $a < b$, there exists $d \in D$ with $a < d < b$.

**Definition 1.1.2.** *Fields and Ordered Fields*
Let $F$ be a set with operations $+, \cdot : F \times F \to F$ and elements $0, 1 \in F$ with $0 \neq 1$.

(i) A *field* has $(F, +, 0)$ and $(F \setminus \{0\}, \cdot, 1)$ abelian groups and $x(y + z) = xy + xz$ for all $x, y, z \in F$.
(ii) An *ordered field* is a field with total order $<$ satisfying $y < z \Rightarrow x + y < x + z$ and $0 < x, 0 < y \Rightarrow 0 < xy$.

**Principle 1.1.3.** *Completeness Axiom*
Every nonempty set $E \subseteq \mathbb{R}$ that is bounded above has a least upper bound in $\mathbb{R}$, denoted $\sup E$.
Equivalently, the real line has no missing cut between a nonempty bounded set and its upper bounds.

**Remark 1.1.4.** *Fields as Objects and Scalars*
In analysis, $\mathbb{Q}$ and $\mathbb{R}$ are ordered fields: $\mathbb{Q}$ lacks the least-upper-bound property, while $\mathbb{R}$ has it.

**Proposition 1.1.5.** *Basic Consequences of the Field Axioms*
*Let $F$ be a field. For all $a, b, c, x, y \in F$:*

*(i)* $a + c = b + c$ *implies* $a = b$; *if* $c \neq 0$, *then* $ac = bc$ *implies* $a = b$.
*(ii)* *The identities* $0, 1$ *and inverses* $-a$, $a^{-1}$ $(a \neq 0)$ *are unique.*
*(iii)* $0x = 0$.
*(iv)* $xy = 0$ *implies* $x = 0$ *or* $y = 0$.
*(v)* $(-x)y = -(xy) = x(-y)$ *and* $(-x)(-y) = xy$.

*Proof.*

(i) Add $-c$ to $a + c = b + c$. If $c \neq 0$, multiply $ac = bc$ by $c^{-1}$.
(ii) In any group $(G, *, e)$, if $e'$ is an identity, $e = e * e' = e'$.
If $u * a = e = a * v$, then $u = u * e = u * (a * v) = (u * a) * v = v$. Apply to $(F, +, 0)$ and $(F \setminus \{0\}, \cdot, 1)$.
(iii) Since $0x + 0x = (0 + 0)x = 0x$, cancellation gives $0x = 0$.
(iv) If $xy = 0$ and $x \neq 0$, multiply by $x^{-1}$ to get $y = 0$.
(v) $xy + (-x)y = (x + (-x))y = 0$, so $(-x)y = -(xy)$; $x(-y) = -(xy)$. Also $(-x)(-y) = -(x(-y)) = xy$.

*Structural Commentary*: Inverses cancel terms.
*Key Insight*: Negatives are unique additive inverses. □

**Proposition 1.1.6.** *Basic Inequality Rules in Ordered Fields*
*Let $F$ be an ordered field. For all $c, x, y \in F$:*

*(i)* $x > 0$ *iff* $-x < 0$.
*(ii)* *If* $c > 0$ *and* $x < y$, *then* $cx < cy$.
*(iii)* *If* $c < 0$ *and* $x < y$, *then* $cy < cx$.
*(iv)* *If* $x \neq 0$, *then* $x^2 > 0$; *in particular,* $1 > 0$.
*(v)* *If* $0 < x < y$, *then* $0 < 1/y < 1/x$.

*Proof.*

(i) If $x > 0$, add $-x$ to $0 < x$ to get $-x < 0$. If $-x < 0$, add $x$ to get $0 < x$.
(ii) Since $x < y$, $y - x > 0$. With $c > 0$, $c(y - x) > 0$, so $cy - cx > 0$ and $cx < cy$.
(iii) Since $c < 0$, (i) gives $-c > 0$. By (ii), $(-c)x < (-c)y$; hence $-cx < -cy$, so $cy < cx$.

(iv) If $x > 0$, then $x^2 > 0$. If $x < 0$, then $-x > 0$ by (i), so $(-x)^2 > 0$ and $x^2 = (-x)^2$. Since $x \neq 0$, totality gives one of these cases. Taking $x = 1$ gives $1 = 1^2 > 0$.

(v) By (iv), $(1/x)^2 > 0$ and $(1/y)^2 > 0$. Since $x, y > 0$, $1/x = x(1/x)^2 > 0$ and $1/y = y(1/y)^2 > 0$, so $(1/x)(1/y) > 0$. Apply (ii) to $x < y$ with multiplier $(1/x)(1/y)$ to get $1/y < 1/x$.

*Structural Commentary*: Positivity preserves order under multiplication.
*Key Insight*: Negative multiplication is positive multiplication by $-c$. □

**Proposition 1.1.7.** *The Rational Gap at $\sqrt{2}$*
*There is no rational $p$ such that $p^2 = 2$.*

*Proof.* Let $p = m/n$. Then $m^2 = 2n^2$, so $m$ is even; $m = 2k$. Hence $n^2 = 2k^2$, so $n$ is even, a contradiction.
*Structural Commentary*: This exhibits a gap in $\mathbb{Q}$.
*Key Insight*: $\sqrt{2}$ is not rational. □




Figure 115: The diagonal of a unit square has length $\sqrt{2}$, so the rational gap is geometric as well as algebraic.

**Example 1.1.8.** *No Largest Lower Side and No Smallest Upper Side*
Let $A := \{p \in \mathbb{Q} : p > 0,\ p^2 < 2\}$, and $B := \{p \in \mathbb{Q} : p > 0,\ p^2 > 2\}$.
For every $p > 0$ define $q := p - \frac{p^2-2}{p+2} = \frac{2p+2}{p+2}$. Then $q^2 - 2 = \frac{2(p^2-2)}{(p+2)^2}$. Hence if $p \in A$, then $q \in A$ and $q > p$; if $p \in B$, then $q \in B$ and $0 < q < p$. Thus $A$ has no largest element in $\mathbb{Q}$, and $B$ has no smallest element in $\mathbb{Q}$.

**Remark 1.1.9.** *Density Does Not Mean Completeness*
Between two rationals $r < s$ there is another rational, namely $(r + s)/2$. Nevertheless, $\mathbb{Q}$ is not complete.




Figure 116: Reflecting $B$ through 0 turns lower bounds of $B$ into upper bounds of $-B$, so $\inf B = -\sup(-B)$.

**Corollary 1.1.10.** *Infimum by Reflection*
*Let $B \subseteq \mathbb{R}$ be nonempty and bounded below, and define $-B := \{-b : b \in B\}$.*
*Then $-B$ is nonempty and bounded above, and $\inf B = -\sup(-B)$.*

*Proof.* If $m$ is a lower bound of $B$, then $-m$ is an upper bound of $-B$; hence $s := \sup(-B)$ exists.
For $b \in B$, $-b \leq s$, so $-s \leq b$; thus $-s$ is a lower bound of $B$.
If $t$ is any lower bound of $B$, then $-t$ is an upper bound of $-B$, so $s \leq -t$ and $t \leq -s$. Therefore $-s = \inf B$.
*Structural Commentary*: Negation reverses bounds.
*Key Insight*: $\inf B = -\sup(-B)$. □

**Definition 1.1.11.** *Real Absolute Value and Distance*
For $x \in \mathbb{R}$, define $|x| := \begin{cases} x, & x \geq 0, \\ -x, & x < 0. \end{cases}$
For $a, b \in \mathbb{R}$, the distance between $a$ and $b$ on the real line is $|a - b|$.

**Proposition 1.1.12.** *Real Absolute-Value Laws*
*For $x, y, a, b, c \in \mathbb{R}$,*

*(i) $|x| \geq 0$, $|x| = 0$ iff $x = 0$, and $-|x| \leq x \leq |x|$;*

*(ii)* $|xy| = |x|\,|y|$;
*(iii)* $|x+y| \le |x| + |y|$;
*(iv)* $|a-b| \le |a-c| + |c-b|$.

*Proof.*

(i) If $x \ge 0$, then $|x| = x$; if $x < 0$, then $|x| = -x > 0$. Hence $|x| \ge 0$, and $|x| = 0$ iff $x = 0$. The bounds $-|x| \le x \le |x|$ follow from the same two cases.
(ii) If $x, y$ have the same sign, then $xy \ge 0$ and $|xy| = xy = |x|\,|y|$. If they have opposite signs, then $xy < 0$ and $|xy| = -xy = |x|\,|y|$.
(iii) By (i), $-|x| \le x \le |x|$ and $-|y| \le y \le |y|$, so $-(|x| + |y|) \le x + y \le |x| + |y|$. Since $|x| + |y| \ge 0$, (i) applied to $x + y$ gives $|x + y| \le |x| + |y|$.
(iv) Apply (iii) to $a - b = (a - c) + (c - b)$.

*Structural Commentary*: Absolute value is two-sided order control.
*Key Insight*: Apply (iii) to $(a - c) + (c - b)$. □

**Perspective 1.1.13.** *Completeness Becomes a Theorem*
Dedekind cuts build completeness from $\mathbb{Q}$: real numbers are rational cuts, rational cuts embed $\mathbb{Q}$, and least upper bounds arise by taking unions of cuts.

**Definition 1.1.14.** *Dedekind Cuts and Their Order*
A *cut* is a set $\alpha \subseteq \mathbb{Q}$ such that

(i) $\alpha$ is nonempty and $\alpha \neq \mathbb{Q}$;
(ii) if $p \in \alpha$, $q \in \mathbb{Q}$, and $q < p$, then $q \in \alpha$;
(iii) if $p \in \alpha$, then some $r \in \alpha$ satisfies $p < r$.

Let $\mathscr{D}$ be the set of all cuts. For $\alpha, \beta \in \mathscr{D}$, define $\alpha < \beta$ to mean $\alpha \subsetneq \beta$.
Rational letters $p, q, r, \ldots$ denote elements of $\mathbb{Q}$; Greek letters $\alpha, \beta, \gamma, \ldots$ denote cuts.
A cut is the rational left side of a real number: a proper initial segment with no last point.




Figure 117: A Dedekind cut splits $\mathbb{Q}$ into $\alpha$ and $\mathbb{Q} \setminus \alpha$, with no last rational in $\alpha$.

**Remark 1.1.15.** *Left Cuts Versus Two-Sided Cuts*
Two-sided notation writes a cut as $A|B$, with $A = \alpha$ and $B = \mathbb{Q} \setminus \alpha$; Definition 1.1.14 records only $A$.
Thus "no largest element" is the same left-side condition.

**Definition 1.1.16.** *Arithmetic of Cuts*
For cuts $\alpha, \beta$, define $\alpha + \beta := \{r + s : r \in \alpha,\ s \in \beta\}$ and $0^* := \{q \in \mathbb{Q} : q < 0\}$.
Let candidate additive inverse of $\alpha$ be $-\alpha := \{p \in \mathbb{Q} : \exists r \in \mathbb{Q},\ r > 0,\ -p - r \notin \alpha\}$. Let $\mathscr{D}^+ := \{\alpha \in \mathscr{D} : \alpha > 0^*\}$.
For $\alpha, \beta \in \mathscr{D}^+$, define $\alpha\beta := \{p \in \mathbb{Q} : \exists r \in \alpha,\ \exists s \in \beta,\ r > 0,\ s > 0,\ p \le rs\}$ and $1^* := \{q \in \mathbb{Q} : q < 1\}$.
Extend multiplication to all cuts by setting $\alpha 0^* = 0^* \alpha = 0^*$ and, for nonzero signs,

$$\alpha\beta := \begin{cases} (-\alpha)(-\beta), & \alpha < 0^*,\ \beta < 0^*, \\ -\big((-\alpha)\beta\big), & \alpha < 0^*,\ \beta > 0^*, \\ -\big(\alpha(-\beta)\big), & \alpha > 0^*,\ \beta < 0^*. \end{cases}$$

*Motivation*: arithmetic uses rational approximations inside cuts; signs come after positive products.

**Definition 1.1.17.** *Rational Cuts*
For $r \in \mathbb{Q}$, define the *rational cut*, $r^* := \{p \in \mathbb{Q} : p < r\}$.
The map $r \mapsto r^*$ is intended to identify $\mathbb{Q}$ with a subfield of the cut field.

**Theorem 1.1.18.** *Existence of the Real Field*
*There exists an ordered field $\mathbb{R}$ with the least-upper-bound property. Moreover, $\mathbb{R}$ contains $\mathbb{Q}$ as a subfield.*

*Proof.* Construct $\mathbb{R}$ from $\mathbb{Q}$ using Definitions 1.1.14–1.1.17.

(i) *Order.* The relation $\alpha < \beta$ is transitive, and at most one of $\alpha < \beta$, $\alpha = \beta$, and $\beta < \alpha$ can hold. For trichotomy, suppose $\alpha \not\subseteq \beta$ and $\alpha \neq \beta$. Choose $p \in \alpha$ with $p \notin \beta$. If $q \in \beta$, then $q < p$, since $p \notin \beta$ and $\beta$ is downward closed. Hence $q \in \alpha$, so $\beta \subseteq \alpha$. Since $\alpha \neq \beta$, $\beta < \alpha$. Thus $\mathscr{D}$ is ordered.
*Commentary*: One missing rational decides containment.

(ii) *Least Upper Bounds.* Let $A \subseteq \mathscr{D}$ be nonempty and bounded above by a cut $\beta$. Define $\gamma := \bigcup_{\alpha \in A} \alpha$.
Since some $\alpha_0 \in A$ is nonempty, $\gamma$ is nonempty. Since every $\alpha \in A$ lies in $\beta$, $\gamma \subseteq \beta$, so $\gamma \neq \mathbb{Q}$. If $p \in \gamma$ and $q < p$, then $p \in \alpha_1$ for some $\alpha_1 \in A$, hence $q \in \alpha_1 \subseteq \gamma$. If $p \in \gamma$, choose $\alpha_1 \in A$ with $p \in \alpha_1$ and then choose $r \in \alpha_1$ with $p < r$; then $r \in \gamma$. Thus $\gamma$ is a cut.
Every $\alpha \in A$ satisfies $\alpha \leq \gamma$. If $\delta < \gamma$, choose $s \in \gamma \setminus \delta$. Then $s \in \alpha$ for some $\alpha \in A$, so $\delta < \alpha$ and $\delta$ is not an upper bound of $A$. Hence $\gamma = \sup A$.
*Motivation*: Supremum is union.
*Commentary*: Bounded unions stay proper and give the smallest upper bound.

(iii) *Addition.* The set $\alpha + \beta$ is a cut. It is nonempty by choosing one element from each summand.
If $r' \notin \alpha$ and $s' \notin \beta$, then every $r \in \alpha$ and $s \in \beta$ satisfies $r < r'$ and $s < s'$, so $r + s < r' + s'$; hence $r' + s' \notin \alpha + \beta$, and $\alpha + \beta$ is proper. If $p = r + s \in \alpha + \beta$ and $q < p$, then $q = r + (s + q - p)$ with $s+q-p < s$, so $s+q-p \in \beta$ and $q \in \alpha+\beta$. If $p = r+s$, choose $r_1 \in \alpha$ with $r < r_1$; then $p < r_1+s \in \alpha+\beta$. Thus $\alpha + \beta$ is a cut. Commutativity and associativity follow by rewriting representatives in $\mathbb{Q}$.
The cut $0^*$ is the additive identity: $\alpha + 0^* \subseteq \alpha$ by downward closure, while if $p \in \alpha$, choose $r \in \alpha$ with $p < r$ and write $p = r + (p - r)$ with $p - r < 0$.
The inverse $-\alpha$ is a cut, and $\alpha + (-\alpha) = 0^*$. The inclusion $\alpha + (-\alpha) \subseteq 0^*$ follows because $s \in -\alpha$ forces $-s \notin \alpha$, so any $r \in \alpha$ satisfies $r < -s$. Conversely, if $v < 0$, put $w = -v/2 > 0$ and choose an integer $n$ with $nw \in \alpha$ but $(n+1)w \notin \alpha$; then $p = -(n+2)w$ lies in $-\alpha$ and $v = nw + p \in \alpha + (-\alpha)$.
*Commentary*: Addition inherits its laws from $\mathbb{Q}$ once closure and inverses are checked.

(iv) *Order Compatibility with Addition.* The additive laws make $\mathscr{D}$ an additive group.
If $\beta < \gamma$, then $\alpha+\beta \subseteq \alpha+\gamma$. Equality would imply $\beta = \gamma$ after adding $-\alpha$ to both sides, so $\alpha+\beta < \alpha+\gamma$. Consequently, $\alpha > 0^*$ if and only if $-\alpha < 0^*$.
*Commentary*: Cancellation turns inclusion into strict inclusion.

(v) *Positive Multiplication.* If $\alpha, \beta \in \mathscr{D}^+$, then $\alpha\beta$ is a cut. Since each positive cut contains some positive rational, $\alpha\beta$ is nonempty. Choose positive $r' \notin \alpha$ and $s' \notin \beta$; then every positive $r \in \alpha$ and $s \in \beta$ satisfies $rs < r's'$, so $r's' \notin \alpha\beta$, and the product is proper. Downward closure is built into the condition $p \leq rs$. If $p \leq rs$ with $r, s > 0$, then either choose $q$ with $p < q < rs$, or, when needed, choose $r_1 \in \alpha$ with $r < r_1$ and take $q$ with $p < q \leq r_1 s$; hence the product has no largest element.
Commutativity follows from $rs = sr$. Associativity and distributivity are proved by unfolding membership and using the rational identities $(rs)t = r(st)$ and $r(s+t) = rs+rt$, with a positive rational slack inserted whenever a representative must be made strictly larger inside a cut. The identity is $1^*$: if $p \leq rs$ with $r \in \alpha$, $s < 1$, and $r, s > 0$, then $p < r$, so $p \in \alpha$; conversely, if $p \in \alpha$, choose $r \in \alpha$ with $p < r$ and then choose $s \in \mathbb{Q}$ with $0 < s < 1$ and $p \leq rs$. In particular, if $\alpha > 0^*$ and $\beta > 0^*$, then $\alpha\beta > 0^*$.
*Commentary*: Positivity avoids sign reversals.

(vi) *Signed Multiplication.* The definitions extend multiplication to all cuts by reducing to positive factors and additive inverses. Since $\delta = -(-\delta)$ and the sign rules for addition have already been proved, commutativity, associativity, and inverses reduce to the positive case.
For distributivity, split into signs of $\alpha, \beta, \gamma, \beta + \gamma$. For example, if $\alpha > 0^*$, $\beta < 0^*$, and $\beta + \gamma > 0^*$, then $\gamma = (\beta + \gamma) + (-\beta)$, so positive distributivity gives $\alpha\gamma = \alpha(\beta + \gamma) + \alpha(-\beta)$. Because $\alpha(-\beta) = -(\alpha\beta)$, adding $\alpha\beta$ to both sides gives $\alpha\beta + \alpha\gamma = \alpha(\beta + \gamma)$. The other sign cases are identical reductions after replacing negative terms by their additive inverses.
*Commentary*: Distributivity reduces to positive cases plus signs.

(vii) *Embedding $\mathbb{Q}$.* Each $r^*$ is a cut. For $r, s \in \mathbb{Q}$, $r^* + s^* = (r + s)^*$, $r^*s^* = (rs)^*$, and $r^* < s^* \iff r < s$.
For addition, if $p = u+v$ with $u < r$ and $v < s$, then $p < r+s$. Conversely, if $p < r+s$, choose $t > 0$ with $2t = r + s - p$ and put $r' = r - t$, $s' = s - t$; then $r' < r$, $s' < s$, and $p = r' + s'$. For positive products, if $p \leq uv$ with $0 < u < r$ and $0 < v < s$, then $p < rs$. Conversely, if $0 < p < rs$, choose positive $u < r$ and $v < s$ close enough to $r$ and $s$ that $p \leq uv$; if $p \leq 0$, use any positive $u < r$, $v < s$. The signed product cases follow from the definitions by additive inverses. The order equivalence follows because $r < s$ puts $r$ in $s^*$ but not in $r^*$, while $r^* < s^*$ gives some $p < s$ with $p \notin r^*$, hence $r \leq p < s$.
*Commentary*: The map $r \mapsto r^*$ preserves arithmetic and order.

Identify $\mathbb{Q}$ with the subfield $\mathbb{Q}^* := \{r^* : r \in \mathbb{Q}\} \subseteq \mathscr{D}$, and write $\mathbb{R}$ for $\mathscr{D}$. Items (i), (iii)–(vi) give an ordered field; item (ii) gives the least-upper-bound property; item (vii) embeds $\mathbb{Q}$ as a subfield.
Any two ordered fields with the least-upper-bound property are isomorphic, so this construction characterizes the real field up to ordered-field structure. □

**Perspective 1.1.19.** *Number Systems from Representatives*
Choose representatives $X$, an equivalence relation $\sim$, and classes $[x] \in X/\sim$: $\mathbb{Z} : (a, b) \mapsto [(a, b)]$, $\mathbb{Q} : (a, b)$ with $b \neq 0 \mapsto [(a, b)]$, and $\mathbb{R}$ via cuts or Cauchy data. Prove each operation or order is well defined, e.g. $x \sim x'$ and $y \sim y'$ imply $x \star y \sim x' \star y'$ and $x < y \Leftrightarrow x' < y'$; then use $X/\sim$ as the number system.
*Key Insight*: Laws must descend to classes.

**Theorem 1.1.20.** *Archimedean Property and Density of $\mathbb{Q}$*

*(i) If $x, y \in \mathbb{R}$ and $x > 0$, then there is a positive integer $n$ such that $nx > y$.*
*(ii) If $x, y \in \mathbb{R}$ and $x < y$, then there exists $p \in \mathbb{Q}$ such that $x < p < y$.*

*Proof.*

(i) Let $A = \{nx : n \in \mathbb{N}\}$. If no $nx$ exceeded $y$, then $A$ would be bounded above.
By Theorem 1.1.18, let $\alpha = \sup A$. Since $x > 0$, $\alpha - x < \alpha$, so $\alpha - x$ is not an upper bound.
Hence $\alpha - x < mx$ for some $m \in \mathbb{N}$, and $\alpha < (m+1)x \in A$, a contradiction.

(ii) Since $y - x > 0$, part (i) gives $n \in \mathbb{N}$ with $n(y-x) > 1$. Apply part (i) again to get $m_1, m_2 \in \mathbb{N}$ with $m_1 > nx$ and $m_2 > -nx$. Then some $m \in \mathbb{Z}$ satisfies $m - 1 \le nx < m$.
Thus $nx < m \le nx + 1 < ny$, so $x < m/n < y$ and $m/n \in \mathbb{Q}$.

*Structural Commentary*: Completeness makes $\{nx : n \in \mathbb{N}\}$ unbounded.
*Key Insight*: Archimedean unboundedness gives rational density. □




Figure 118: Choose $n$ so the spacing $1/n$ is smaller than $b - a$; then one grid point $m/n$ falls inside $(a, b)$.

**Theorem 1.1.21.** *Existence and Uniqueness of Positive nth Roots*
*For every $x \in \mathbb{R}$ with $x > 0$ and every positive integer $n$, $\exists$ a unique positive real number $y$ such that $y^n = x$.*

*Proof.*

(i) *Uniqueness.* If $0 < y_1 < y_2$, then $y_1^n < y_2^n$.

(ii) *Existence.* Let $E := \{t \in \mathbb{R} : t > 0,\ t^n < x\}$. $E \neq \varnothing$: for $t = x/(1+x)$, $0 < t < 1$ and $t^n \le t < x$.
Also $E$ is bounded above by $1 + x$, since $t > 1 + x$ implies $t^n \ge t > x$. By Theorem 1.1.18, let $y = \sup E$.
For $0 < a < b$, $b^n - a^n = (b-a)(b^{n-1} + b^{n-2}a + \cdots + a^{n-1}) < (b-a)nb^{n-1}$.
If $y^n < x$, choose $h$ with $0 < h < 1$ and $h < \frac{x - y^n}{n(y+1)^{n-1}}$.
Then $(y+h)^n - y^n < hn(y+h)^{n-1} < hn(y+1)^{n-1} < x - y^n$.
Thus $y + h \in E$, contradicting that $y$ is an upper bound. If $y^n > x$, set $k := \frac{y^n - x}{ny^{n-1}}$. Then $0 < k < y$.
For $t \ge y - k$, $y^n - t^n \le y^n - (y-k)^n < kny^{n-1} = y^n - x$, so $t^n > x$ and $t \notin E$.
Hence $y - k$ is an upper bound for $E$, contradicting $y = \sup E$. Therefore $y^n = x$.

*Structural Commentary*: The root is $\sup\{t > 0 : t^n < x\}$.
*Key Insight*: Completeness supplies the missing root. □

**Corollary 1.1.22.** *Density of Rational and Irrational Points*
*If $x < y$ in $\mathbb{R}$, then there exist $r \in \mathbb{Q}$ and $s \in \mathbb{R} \setminus \mathbb{Q}$ such that $x < r < y$ and $x < s < y$.*

*Proof.* The rational point $r$ is supplied by Theorem 1.1.20. Let $\eta = \sqrt{2}$.
Choose $n \in \mathbb{N}$ with $0 < \eta/n < y - x$, and choose $q \in \mathbb{Q}$ with $x - \eta/n < q < y - \eta/n$.
Then $s := q + \eta/n$ lies in $(x, y)$ and is irrational, since $s \in \mathbb{Q}$ would imply $\eta = n(s - q) \in \mathbb{Q}$.
*Structural Commentary*: Irrational density is rational density shifted by a small irrational amount.
*Key Insight*: Once one irrational is known, every interval contains irrational translates of rational points. □

**Corollary 1.1.23.** *Product Rule for Positive Roots*
*If $a, b > 0$ and $n$ is a positive integer, then $(ab)^{1/n} = a^{1/n} b^{1/n}$.*

*Proof.* Let $\alpha = a^{1/n}$ and $\beta = b^{1/n}$. Then $ab = \alpha^n \beta^n = (\alpha\beta)^n$ by commutativity of multiplication.
The uniqueness part of Theorem 1.1.21 gives the identity.
*Structural Commentary*: Use uniqueness after checking $n$th powers.
*Key Insight*: Root identities are uniqueness arguments. □

**Definition 1.1.24.** *Extended Real Number System*
The *extended real number system* is $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$.
The order on $\mathbb{R}$ is preserved, and $-\infty < x < +\infty$, $(x \in \mathbb{R})$.
Every subset of $\mathbb{R}$ has a supremum and an infimum in $\overline{\mathbb{R}}$ after the conventions $\sup \emptyset = -\infty$, $\inf \emptyset = +\infty$, $\sup E = +\infty$ when $E$ is unbounded above, and $\inf E = -\infty$ when $E$ is unbounded below.

**Remark 1.1.25.** *Extended Arithmetic Conventions*
$\overline{\mathbb{R}}$ is not a field. For $x \in \mathbb{R}$:

(i) $\infty + \infty = \infty$ and $-\infty - \infty = -\infty$.
(ii) $x + \infty = \infty$ and $x - \infty = -\infty$.
(iii) If $x > 0$, then $x \cdot \infty = \infty$ and $x \cdot (-\infty) = -\infty$; if $x < 0$, the signs reverse.

Expressions $\infty - \infty$ and $0 \cdot \infty$ are undefined.

**Remark 1.1.26.** *Unbounded Intervals*
For $a, b \in \mathbb{R}$:

(i) $(a, \infty) := \{x \in \mathbb{R} : a < x\}$ and $[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$.
(ii) $(-\infty, b) := \{x \in \mathbb{R} : x < b\}$ and $(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}$.

Also $(-\infty, \infty) := \mathbb{R}$.

## 1.2 Complex Numbers and Absolute Value

**Definition 1.2.1.** *Complex Numbers*
Let $\mathbb{C} := \mathbb{R}^2$. For $x = (a, b)$ and $y = (c, d)$ in $\mathbb{C}$:

(i) $x = y$ iff $a = c$ and $b = d$.
(ii) $x + y := (a + c, b + d)$.
(iii) $xy := (ac - bd, ad + bc)$.

Elements of $\mathbb{C}$ are *complex numbers*.

**Theorem 1.2.2.** *The Complex Numbers Form a Field*
*Let $\mathbb{C} = \mathbb{R}^2$ with $(a, b) + (c, d) = (a + c, b + d)$ and $(a, b)(c, d) = (ac - bd, ad + bc)$.*

*(i) $\mathbb{C}$ is a field.*
*(ii) The additive and multiplicative identities are $(0, 0)$ and $(1, 0)$.*
*(iii) If $x = (a, b) \neq (0, 0)$, then $x^{-1} = \left(\frac{a}{a^2+b^2}, \frac{-b}{a^2+b^2}\right)$.*

*Proof.*

(i) For $x, y, z \in \mathbb{C}$, associativity, commutativity, and distributivity follow by expanding coordinates in $\mathbb{R}$; $-(a, b) = (-a, -b)$, and (iii) gives $x^{-1}$ for $x \neq 0$.
(ii) Directly, $(a, b) + (0, 0) = (a, b)$ and $(a, b)(1, 0) = (a, b)$.
(iii) Since $(a, b) \neq (0, 0)$, $a^2 + b^2 > 0$. Multiplication gives $(a, b)\left(\frac{a}{a^2+b^2}, \frac{-b}{a^2+b^2}\right) = (1, 0)$.

*Structural Commentary*: Inverses use conjugate-over-norm.
*Key Insight*: $a^2 + b^2 > 0$ for $(a, b) \neq (0, 0)$. □

**Proposition 1.2.3.** *Real Numbers Inside Complex Numbers*
*Let $\iota : \mathbb{R} \to \mathbb{C}$ be $\iota(a) := (a, 0)$, and set $i := (0, 1)$.*

*(i) $\iota(a + b) = \iota(a) + \iota(b)$ and $\iota(ab) = \iota(a)\iota(b)$ for all $a, b \in \mathbb{R}$.*
*(ii) $\iota$ is injective, $\iota(0) = (0, 0)$, and $\iota(1) = (1, 0)$.*

*Where $a \in \mathbb{R}$ with $\iota(a) \in \mathbb{C}$; $i$ is the complex unit.*

*Proof.*

(i) $\iota(a) + \iota(b) = (a, 0) + (b, 0) = (a + b, 0) = \iota(a + b)$, and $\iota(a)\iota(b) = (a, 0)(b, 0) = (ab, 0) = \iota(ab)$.
(ii) If $\iota(a) = \iota(b)$, then $(a, 0) = (b, 0)$, so $a = b$. Also $\iota(0) = (0, 0)$ and $\iota(1) = (1, 0)$.

*Structural Commentary*: Real arithmetic embeds as the real axis.
*Key Insight*: $\iota$ preserves $+, \cdot, 0, 1$. □

**Proposition 1.2.4.** *Coordinate Form and $i^2$*
*For $i = (0, 1) \in \mathbb{C}$:*

*(i) $i^2 = -1$.*
*(ii) For all $a, b \in \mathbb{R}$, $(a, b) = a + bi$.*

*Proof.*

(i) $i^2 = (0, 1)(0, 1) = (-1, 0) = -1$.
(ii) $a + bi = (a, 0) + (b, 0)(0, 1) = (a, 0) + (0, b) = (a, b)$.

*Structural Commentary*: Pair coordinates become $a + bi$.
*Key Insight*: $i = (0, 1)$ encodes the second coordinate. □

**Definition 1.2.5.** *Conjugate, Real Part, and Imaginary Part*
Let $z = a + bi$ with $a, b \in \mathbb{R}$.

(i) The *complex conjugate* of $z$ is $\overline{z} := a - bi$.
(ii) The *real part* and *imaginary part* of $z$ are $\mathrm{Re}(z) := a$ and $\mathrm{Im}(z) := b$.

**Proposition 1.2.6.** *Basic Conjugation Identities*
*If $z, w \in \mathbb{C}$, then*

*(i)* $\overline{z+w} = \overline{z} + \overline{w}$;
*(ii)* $\overline{zw} = \overline{z}\,\overline{w}$;
*(iii)* $z + \overline{z} = 2\,\mathrm{Re}(z)$ *and* $z - \overline{z} = 2i\,\mathrm{Im}(z)$;
*(iv)* $z\overline{z}$ *is real and positive unless* $z = 0$.

*Proof.* Write $z = a + bi$ and $w = c + di$.

(i) $\overline{z+w} = (a+c) - (b+d)i = \overline{z} + \overline{w}$.
(ii) $\overline{zw} = (ac - bd) - (ad + bc)i = (a - bi)(c - di) = \overline{z}\,\overline{w}$.
(iii) $z + \overline{z} = 2a = 2\,\mathrm{Re}(z)$ and $z - \overline{z} = 2bi = 2i\,\mathrm{Im}(z)$.
(iv) $z\overline{z} = a^2 + b^2$, positive unless $a = b = 0$, i.e. unless $z = 0$.

*Structural Commentary*: Conjugation fixes Re and negates Im.
*Key Insight*: $z\overline{z} = a^2 + b^2 \geq 0$. □

**Definition 1.2.7.** *Absolute Value of a Complex Number*
For $z = a + bi \in \mathbb{C}$, define $|z| := \sqrt{z\overline{z}} = \sqrt{a^2 + b^2}$, the unique $r \in [0, \infty)$ with $r^2 = z\overline{z} = a^2 + b^2$.

**Theorem 1.2.8.** *Absolute Value Identities*
*For $z, w \in \mathbb{C}$,*

*(i)* $|z| > 0$ *unless* $z = 0$, *and* $|0| = 0$;
*(ii)* $|\overline{z}| = |z|$;
*(iii)* $|zw| = |z|\,|w|$;
*(iv)* $|\mathrm{Re}(z)| \leq |z|$;
*(v)* $|z + w| \leq |z| + |w|$.

*Proof.*

(i) If $z = a + bi$, then $|z| = (a^2 + b^2)^{1/2}$; this is positive unless $a = b = 0$, and $|0| = 0$.
(ii) $|\overline{z}|^2 = \overline{z}z = z\overline{z} = |z|^2$, so nonnegative square roots give $|\overline{z}| = |z|$.
(iii) $|zw|^2 = zw\overline{zw} = z\overline{z}w\overline{w} = |z|^2\,|w|^2$; take nonnegative square roots.
(iv) If $z = a + bi$, then $|\mathrm{Re}(z)| = |a| \leq (a^2 + b^2)^{1/2} = |z|$.
(v) $z\overline{w}$ and $\overline{z}w$ are conjugates, $|z+w|^2 = |z|^2 + 2\,\mathrm{Re}(z\overline{w}) + |w|^2 \leq |z|^2 + 2\,|z\overline{w}| + |w|^2 = (|z| + |w|)^2$, using (iii) and (iv). Take nonnegative square roots.

*Structural Commentary*: The complex absolute value behaves like length because $z\overline{z}$ is a squared length.
*Key Insight*: Triangle inequality is a real-part estimate plus multiplicativity. □

# 2 Real Sequences and Completeness Methods

## 2.1 Sequences, Cauchy Criteria, and Completeness

Sequences are the first testing ground for limits, order, and completeness. The main tools are subsequences, Cauchy criteria, monotone convergence, limsup/liminf, and Bolzano–Weierstrass.

**Definition 2.1.1.** *Convergent Real and Complex Sequences*
Let $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$, let $(s_n)$ be a sequence in $\mathbb{K}$, and let $s \in \mathbb{K}$.

(i) $(s_n)$ *converges* to $s$ iff for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $n \geq N$ implies $|s_n - s| < \varepsilon$.
(ii) Write $s_n \to s$ or $\lim_{n\to\infty} s_n = s$. If no $s \in \mathbb{K}$ satisfies (i), then $(s_n)$ *diverges*.

Figure 119: For all $n \geq N$, the terms $s_n$ stay inside $(s - \varepsilon, s + \varepsilon)$.

**Definition 2.1.2.** *Range and Boundedness of a Real Sequence*
Let $(s_n)$ be a real sequence.

(i) The *range* of $(s_n)$ is $\text{range}(s_n) := \{s_n : n \in \mathbb{N}\}$.
(ii) $(s_n)$ is *bounded* iff there exist $m, M \in \mathbb{R}$ such that $m \leq s_n \leq M$ for all $n \in \mathbb{N}$.

**Proposition 2.1.3.** *Basic Consequences of Sequence Convergence*
*Let $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$, and let $(s_n)$ and $(t_n)$ be sequences in $\mathbb{K}$.*

*(i) If $s_n \to s$ and $s_n \to t$, then $s = t$.*
*(ii) If $s_n \to s$, then $\{|s_n| : n \in \mathbb{N}\}$ is bounded.*
*(iii) If $\mathbb{K} = \mathbb{R}$, $s_n \leq t_n$ eventually, $s_n \to s$, and $t_n \to t$, then $s \leq t$.*
*In particular, $s_n \leq c$ eventually implies $s \leq c$.*

*Proof.*

(i) For $\varepsilon > 0$ and large $n$, $|s_n - s|, |s_n - t| < \varepsilon/2$, so $|s - t| < \varepsilon$; hence $s = t$.
(ii) For large $n$, $|s_n - s| < 1$, hence $|s_n| \leq |s| + 1$; the remaining finitely many $|s_n|$ are bounded.
(iii) If $s > t$, let $\varepsilon = (s - t)/3$. For large $n$, $s_n > s - \varepsilon > t + \varepsilon > t_n$, contradicting $s_n \leq t_n$.
Take $t_n = c$ for the final claim.

*Structural Commentary*: Limits are rigid: tails determine the value, stay bounded, and preserve eventual order.
*Key Insight*: The triangle inequality and an $\varepsilon/3$ gap control the basic consequences. □

**Theorem 2.1.4.** *Limit Algebra for Complex Sequences*
*Suppose $(s_n)$ and $(t_n)$ are complex sequences with $s_n \to s$ and $t_n \to t$. Then, for every $c \in \mathbb{C}$,*

*(i) $s_n + t_n \to s + t$;*
*(ii) $cs_n \to cs$ and $c + s_n \to c + s$;*
*(iii) $s_n t_n \to st$;*
*(iv) if $s \neq 0$ and $s_n \neq 0$ for all $n$, then $1/s_n \to 1/s$.*

*Proof.*

(i) Given $\varepsilon > 0$, choose $N$ such that $|s_n - s| < \varepsilon/2$, $|t_n - t| < \varepsilon/2$ for $n \geq N$.
Then $|(s_n + t_n) - (s + t)| \leq |s_n - s| + |t_n - t| < \varepsilon$.
(ii) Scalar multiplication uses $|cs_n - cs| = |c|\,|s_n - s|$; translation is (i) with constant sequence $c$.
(iii) Since $s_n \to s$, choose $N \geq 2$ with $|s_n - s| < 1$ for $n \geq N$ and set $M := \max\{|s_1|, \ldots, |s_{N-1}|, |s| + 1\}$.
Then $|s_n t_n - st| \leq |s_n|\,|t_n - t| + |t|\,|s_n - s| \leq M\,|t_n - t| + |t|\,|s_n - s| \to 0$.
(iv) If $s \neq 0$, then for all sufficiently large $n$, $|s_n| > |s|/2$. Hence $\left|\frac{1}{s_n} - \frac{1}{s}\right| = \frac{|s_n - s|}{|s_n||s|} \leq \frac{2}{|s|^2}\,|s_n - s| \to 0$.

*Structural Commentary*: Boundedness controls products; nonzero limits control reciprocals.
*Key Insight*: Product and reciprocal rules are estimates after the easy sum rule. □

**Definition 2.1.5.** *Subsequences and Subsequential Limits*
Let $(p_n)$ be a sequence and let $\sigma : \mathbb{N} \to \mathbb{N}$ be strictly increasing, with $n_k := \sigma(k)$.

(i) The sequence $(p_{n_k})_{k \in \mathbb{N}}$ is a *subsequence* of $(p_n)$.
(ii) A point $p$ is a *subsequential limit* of $(p_n)$ iff there exists such $\sigma$ with $p_{\sigma(k)} \to p$.

**Remark 2.1.6.** *Subsequences as Selection Maps*
View $(p_n)$ as a function $p : \mathbb{N} \to X$, $p(n) = p_n$.
For a strictly increasing $\sigma : \mathbb{N} \to \mathbb{N}$, the subsequence is $p \circ \sigma : \mathbb{N} \to X$, namely $k \mapsto p_{\sigma(k)}$.
*Key Insight*: A subsequence keeps the original order but is allowed to skip terms.

**Remark 2.1.7.** *Subsequences of Convergent Sequences*
If $p_n \to p$, then every subsequence of $(p_n)$ also converges to $p$. Conversely, if every subsequence of $(p_n)$ converges to $p$, then $(p_n)$ itself converges to $p$, since $(p_n)$ is one of its subsequences.

**Definition 2.1.8.** *Monotone Real Sequence*
Let $(s_n)$ be a real sequence.

(i) $(s_n)$ is *monotonically increasing* iff $s_n \leq s_{n+1}$ for all $n \in \mathbb{N}$.
(ii) $(s_n)$ is *monotonically decreasing* iff $s_n \geq s_{n+1}$ for all $n \in \mathbb{N}$.

The sequence is *monotonic* iff it is increasing or decreasing.

**Definition 2.1.9.** *Dominant Index*
For a real sequence $(s_n)$, an index $n \in \mathbb{N}$ is *dominant* iff $s_n \geq s_m$ for all $m > n$.

**Theorem 2.1.10.** *Monotone Subsequence Theorem*
*Every real sequence has a monotone subsequence.*

*Proof.* If the dominant indices $n_1 < n_2 < \cdots$ are infinite, then $s_{n_1} \geq s_{n_2} \geq \cdots$, giving a decreasing subsequence.
Otherwise choose $N$ beyond all dominant indices and set $n_1 > N$.
Given $n_k > N$, nondominance gives $n_{k+1} > n_k$ with $s_{n_{k+1}} > s_{n_k}$; hence $(s_{n_k})$ is increasing.
*Structural Commentary*: Dominant indices give decreasing case; nondominance recursively gives increasing case.
*Key Insight*: The proof organizes arbitrary real behavior into one ordered thread. □




Figure 120: For an increasing bounded sequence, $s = \sup\{s_n\}$ is the limit.

**Theorem 2.1.11.** *Monotone Convergence Theorem for Sequences*
*A monotonic real sequence converges if and only if it is bounded.*

*Proof.*

(⇒) If $s_n \to s$, choose $N$ with $|s_n - s| < 1$ for $n \geq N$.
The tail lies in $(s-1, s+1)$ and the initial set $\{s_1, \ldots, s_{N-1}\}$ is finite, so $(s_n)$ is bounded.

(⇐) Let $E := \{s_n : n \in \mathbb{N}\}$. If $(s_n)$ is increasing, set $s := \sup E$. Given $\varepsilon > 0$, choose $N$ with $s - \varepsilon < s_N \leq s$; then $s - \varepsilon < s_n \leq s$ for $n \geq N$, so $s_n \to s$. If $(s_n)$ is decreasing, set $s := \inf E$. Given $\varepsilon > 0$, choose $N$ with $s \leq s_N < s + \varepsilon$; then $s \leq s_n \leq s_N < s + \varepsilon$ for $n \geq N$, so $s_n \to s$.

*Structural Commentary*: Monotonicity traps bounded tails at $\sup E$ or $\inf E$.
*Key Insight*: The least upper bound is forced to be the limit. □

**Theorem 2.1.12.** *Bolzano–Weierstrass for Real Sequences*
*Every bounded real sequence has a convergent subsequence.*

*Proof.* Let $(s_{n_k})$ be a monotone subsequence.
If $m \leq s_n \leq M$ for all $n$, then $m \leq s_{n_k} \leq M$ for all $k$; hence $(s_{n_k})$ is bounded and monotone, so it converges.
*Structural Commentary*: Boundedness prevents escape; monotone subsequences force one ordered thread.
*Key Insight*: Monotone subsequence plus monotone convergence is the real-line compactness mechanism. □

**Definition 2.1.13.** *Cauchy Real Sequence*
A real sequence $(s_n)$ is *Cauchy* if, for every $\varepsilon > 0$, there is $N \in \mathbb{N}$ such that $m, n \geq N$ implies $|s_n - s_m| < \varepsilon$.

**Lemma 2.1.14.** *Cauchy Plus One Convergent Subsequence*
*If a real Cauchy sequence has a subsequence converging to $s \in \mathbb{R}$, then the whole sequence converges to $s$.*

*Proof.* Given $\varepsilon > 0$, choose $N_0$ so that $m, n \geq N_0$ implies $|s_m - s_n| < \varepsilon/2$.
Choose $j$ with $n_j \geq N_0$ and $\left|s_{n_j} - s\right| < \varepsilon/2$. Then for every $n \geq N_0$, $|s_n - s| \leq \left|s_n - s_{n_j}\right| + \left|s_{n_j} - s\right| < \varepsilon$.
*Structural Commentary*: Cauchy tails follow any convergent thread.
*Key Insight*: Mutual tail closeness turns one subsequential limit into the full limit. □

**Theorem 2.1.15.** *Real Cauchy Criterion*
*A real sequence converges in $\mathbb{R}$ if and only if it is Cauchy.*

*Proof.*

(⇒) If $s_n \to s$, choose $N$ with $|s_n - s| < \varepsilon/2$ for $n \geq N$. Then $m, n \geq N$ implies $|s_n - s_m| < \varepsilon$.

(⇐) If $(s_n)$ is Cauchy, choose $N$ with $|s_m - s_n| < 1$ for $m, n \geq N$. Then the tail lies in $(s_N - 1, s_N + 1)$, so $(s_n)$ is bounded. Bolzano–Weierstrass gives a convergent subsequence; Lemma 2.1.14 gives $s_n \to s$.

*Structural Commentary*: Completeness of $\mathbb{R}$ means Cauchy behavior has a real endpoint.
*Key Insight*: Cauchy boundedness plus a convergent subsequence forces the full limit. □

## 2.2 Divergence, Upper and Lower Limits, and Sequence Estimates

**Definition 2.2.1.** *Divergence to $\pm\infty$*
Let $s : \mathbb{N} \to \mathbb{R}$ and write $s_n := s(n)$.

(i) $s_n \to +\infty$ iff $\forall M \in \mathbb{R},\ \exists N \in \mathbb{N},\ \forall n \in \mathbb{N},\ n \geq N \Rightarrow s_n \geq M$.
(ii) $s_n \to -\infty$ iff $\forall M \in \mathbb{R},\ \exists N \in \mathbb{N},\ \forall n \in \mathbb{N},\ n \geq N \Rightarrow s_n \leq M$.

**Proposition 2.2.2.** *Monotone Sequences Have Extended Limits*
*Let $(s_n)$ be a real sequence.*

*(i) If $(s_n)$ is increasing, then $s_n \to \sup\{s_n : n \in \mathbb{N}\}$ in $\overline{\mathbb{R}}$.*
*(ii) If $(s_n)$ is decreasing, then $s_n \to \inf\{s_n : n \in \mathbb{N}\}$ in $\overline{\mathbb{R}}$.*

*Thus a monotone real sequence has a finite limit exactly when it is bounded.*

*Proof.*

(i) Assume $(s_n)$ is increasing and set $a := \sup\{s_n : n \in \mathbb{N}\} \in \overline{\mathbb{R}}$. If $a \in \mathbb{R}$, then for every $\varepsilon > 0$ there is $N \in \mathbb{N}$ with $a - \varepsilon < s_N \leq a$, and $n \geq N$ implies $a - \varepsilon < s_n \leq a$, so $s_n \to a$. If $a = +\infty$, then for every $M \in \mathbb{R}$ there is $N \in \mathbb{N}$ with $s_N \geq M$, and $n \geq N$ implies $s_n \geq M$, so $s_n \to +\infty$.
(ii) Assume $(s_n)$ is decreasing and set $b := \inf\{s_n : n \in \mathbb{N}\} \in \overline{\mathbb{R}}$. If $b \in \mathbb{R}$, then for every $\varepsilon > 0$ there is $N \in \mathbb{N}$ with $b \leq s_N < b + \varepsilon$, and $n \geq N$ implies $b \leq s_n \leq s_N < b + \varepsilon$, so $s_n \to b$. If $b = -\infty$, then for every $M \in \mathbb{R}$ there is $N \in \mathbb{N}$ with $s_N \leq M$, and $n \geq N$ implies $s_n \leq M$, so $s_n \to -\infty$.

In (i), $a \in \mathbb{R}$ iff $(s_n)$ is bounded; in (ii), $b \in \mathbb{R}$ iff $(s_n)$ is bounded.
Thus the extended limit is finite exactly when $(s_n)$ is bounded.
*Structural Commentary*: Monotone tails have one endpoint in $\overline{\mathbb{R}}$.
*Key Insight*: Finite endpoints give $\varepsilon$-traps; infinite endpoints give threshold tails. □

**Proposition 2.2.3.** *Unbounded Real Sequences Have Infinite Subsequences*
*A real sequence is unbounded iff some subsequence tends to $+\infty$ or $-\infty$.*

*Proof.*

($\Rightarrow$) If $(s_n)$ is unbounded above, every tail is unbounded above; with $n_0 := 0$, choose $n_k > n_{k-1}$ with $s_{n_k} \geq k$, hence $s_{n_k} \to +\infty$. If $(s_n)$ is unbounded below, every tail is unbounded below.
With $n_0 := 0$, choose $n_k > n_{k-1}$ with $s_{n_k} \leq -k$, hence $s_{n_k} \to -\infty$.
($\Leftarrow$) If $s_{n_k} \to +\infty$, then for every $M \in \mathbb{R}$ some $k$ satisfies $s_{n_k} \geq M$, so $(s_n)$ is unbounded above.
If $s_{n_k} \to -\infty$, then for every $M \in \mathbb{R}$ some $k$ satisfies $s_{n_k} \leq M$, so $(s_n)$ is unbounded below.

*Structural Commentary*: Unbounded tails supply the thresholds $\pm k$.
*Key Insight*: Large late terms form a divergent subsequence. □




Figure 121: For $s_n = [(-1)^n + 1]n$, odd terms (orange) stay at 0, while even terms (blue) diverge to $+\infty$.

**Definition 2.2.4.** *Upper and Lower Limits*
Let $(s_n)$ be a real sequence and set $E := E(s) := \{x \in \overline{\mathbb{R}} : \exists\, n_1 < n_2 < \cdots,\ s_{n_k} \to x\}$.

(i) The *upper limit* of $(s_n)$ is $\limsup_{n\to\infty} s_n := s^* := \sup_{\overline{\mathbb{R}}} E$.
(ii) The *lower limit* of $(s_n)$ is $\liminf_{n\to\infty} s_n := s_* := \inf_{\overline{\mathbb{R}}} E$.

**Theorem 2.2.5.** *Characterization of the Upper Limit*
*Let $(s_n)$ be real sequence. Set $E := \{x \in \overline{\mathbb{R}} : \exists\, n_1 < n_2 < \cdots,\ s_{n_k} \to x\}$, $s^* := \sup_{\overline{\mathbb{R}}} E$, $s_* := \inf_{\overline{\mathbb{R}}} E$. Then:*

*(i) $s^* \in E$.*
*(ii) If $x \in \mathbb{R}$ and $x > s^*$, then $s_n < x$ eventually.*

*(iii)* $s^*$ *is the unique* $y \in E$ *such that* $s_n < x$ *eventually for all* $x \in \mathbb{R}$ *with* $x > y$.

*Dually,* $s_*$ *is the unique* $y \in E$ *such that* $s_n > x$ *eventually for all* $x \in \mathbb{R}$ *with* $x < y$.

*Proof.*

(i) Set $n_0 := 0$. If $s^* = +\infty$, choose $y_k \in E$ with $y_k > k$; diagonal selection gives $n_k > n_{k-1}$ with $s_{n_k} > k$, hence $+\infty \in E$. If $s^* \in \mathbb{R}$, choose $y_k \in E \cap (s^* - 1/k, s^*]$; diagonal selection gives $n_k > n_{k-1}$ with $|s_{n_k} - y_k| < 1/k$, hence $s_{n_k} \to s^*$. If $s^* = -\infty$ and $-\infty \notin E$, some tail is bounded below; that tail is unbounded above or bounded, giving respectively $+\infty \in E$ or a finite element of $E$, a contradiction. QED.

(ii) If $x > s^*$ and $s_n \geq x$ infinitely often, some subsequence has extended subsequential limit $y \in E \cap [x, +\infty]$, contradicting $s^* = \sup_{\overline{\mathbb{R}}} E < x \leq y$. Thus $s_n < x$ eventually.

(iii) By (i)–(ii), $s^*$ satisfies the stated conditions. If $p < q$ both satisfy them, choose $x \in \mathbb{R}$ with $p < x < q$; eventual $s_n < x$ from $p$ contradicts the subsequence tending to $q$.

Apply the upper-limit result to $(-s_n)$; its subsequential-limit set is $-E$ and its upper limit is $-s_*$.
*Structural Commentary*: Limsup is the maximal subsequential limit and eventual ceiling.
*Key Insight*: Limsup combines subsequences with eventual upper bounds. □

**Proposition 2.2.6.** *Tail Suprema Formula*
*For a real sequence* $(s_n)$ *and* $N \in \mathbb{N}$, *set* $T_N := \{s_n : n \geq N\}$, $u_N := \sup_{\overline{\mathbb{R}}} T_N$, *and* $\ell_N := \inf_{\overline{\mathbb{R}}} T_N$.

*(i)* $u_{N+1} \leq u_N$ *and* $\ell_{N+1} \geq \ell_N$ *for all* $N \in \mathbb{N}$.
*(ii)* $\limsup_{n\to\infty} s_n = \inf_{N\in\mathbb{N}} u_N$ *and* $\liminf_{n\to\infty} s_n = \sup_{N\in\mathbb{N}} \ell_N$.

*Proof.*

(i) Since $T_{N+1} \subseteq T_N$, $\sup_{\overline{\mathbb{R}}} T_{N+1} \leq \sup_{\overline{\mathbb{R}}} T_N$ and $\inf_{\overline{\mathbb{R}}} T_{N+1} \geq \inf_{\overline{\mathbb{R}}} T_N$.

(ii) Set $s^* := \limsup s_n$, $s_* := \liminf s_n$, $u := \inf_{N\in\mathbb{N}} u_N$, and $\ell := \sup_{N\in\mathbb{N}} \ell_N$.
By Theorem 2.2.5, $x > s^* \Rightarrow u \leq x$ and $y < s^* \Rightarrow u \geq y$ for $x, y \in \mathbb{R}$, hence $u = s^*$.
Dually, $x < s_* \Rightarrow \ell \geq x$ and $y > s_* \Rightarrow \ell \leq y$ for $x, y \in \mathbb{R}$, hence $\ell = s_*$.

*Structural Commentary*: Tail envelopes converge to lim sup and lim inf.
*Key Insight*: Shrinking tails compute the extremal subsequential limits. □




Figure 122: Blue tail suprema $u_N$ decrease to $s^* = \limsup s_n$; orange tail infima $\ell_N$ increase to $s_* = \liminf s_n$.

**Theorem 2.2.7.** *Convergence and Equal Upper/Lower Limits*
*A real sequence* $(s_n)$ *converges to a finite number* $s \in \mathbb{R}$ *if and only if* $\limsup_{n\to\infty} s_n = \liminf_{n\to\infty} s_n = s$.

*Proof.*

($\Rightarrow$) If $s_n \to s$, every subsequence converges to $s$, and the tail is bounded, so no subsequence tends to $\pm\infty$. Thus $E = \{s\}$, hence $\limsup s_n = \liminf s_n = s$.

($\Leftarrow$) Suppose $\limsup s_n = \liminf s_n = s$. Given $\varepsilon > 0$, Theorem 2.2.5 gives $s_n < s+\varepsilon$ and $s_n > s-\varepsilon$ eventually. Hence $|s_n - s| < \varepsilon$ eventually, so $s_n \to s$.

*Structural Commentary*: Equality of the two extremal subsequential limits traps the whole tail.
*Key Insight*: Limsup supplies an eventual ceiling; liminf supplies an eventual floor. □

**Theorem 2.2.8.** *Order Comparison for Upper and Lower Limits*
*If* $s_n \leq t_n$ *for all* $n \geq N$, *where* $N$ *is fixed, then:*

*(i)* $\limsup\limits_{n\to\infty} s_n \leq \limsup\limits_{n\to\infty} t_n$.
*(ii)* $\liminf\limits_{n\to\infty} s_n \leq \liminf\limits_{n\to\infty} t_n$.

*Proof.* Discard finitely many terms, so $s_n \leq t_n$ for all $n$.

(i) Put $a = \limsup s_n$ and $b = \limsup t_n$. If $x < a$, then $s_n > x$ infinitely often, hence $t_n > x$ infinitely often; Theorem 2.2.5 gives $x \leq b$. Thus $a \leq b$.

(ii) Apply (i) to $-t_n \le -s_n$ to get $\limsup(-t_n) \le \limsup(-s_n)$, equivalently $\liminf s_n \le \liminf t_n$.

*Structural Commentary*: Eventual order is inherited by the extremal subsequential limits.
*Key Insight*: Limsup and liminf ignore finitely many initial terms. □

**Theorem 2.2.9.** *Pinching Theorem for Real Sequences*
*Let $(a_n)$, $(s_n)$, and $(b_n)$ be real sequences with $a_n \le s_n \le b_n$ for all sufficiently large $n$.*
*If $a_n \to s$ and $b_n \to s$, then $s_n \to s$.*

*Proof.* Let $\varepsilon > 0$. Choose $N_0$ s.t. $a_n \le s_n \le b_n$ for $n \ge N_0$; $N_1$ s.t. $s-\varepsilon < a_n$ for $n \ge N_1$; and $N_2$ s.t. $b_n < s+\varepsilon$ for $n \ge N_2$. If $N = \max\{N_0, N_1, N_2\}$, then $s - \varepsilon < s_n < s + \varepsilon$ for all $n \ge N$. Hence $s_n \to s$.
*Structural Commentary*: Two convergent bounds squeeze the middle sequence to the same limit.
*Key Insight*: Once both walls are inside the $\varepsilon$-neighborhood, the middle term has no room to escape. □




Figure 123: With $a_n \le s_n \le b_n$ and $a_n, b_n \to s$, the middle sequence also tends to $s$.

**Theorem 2.2.10.** *Special Sequence Limits*
*(i) If $p > 0$, then $\lim_{n\to\infty} n^{-p} = 0$.*
*(ii) If $p > 0$, then $\lim_{n\to\infty} \sqrt[n]{p} = 1$.*
*(iii) $\lim_{n\to\infty} \sqrt[n]{n} = 1$.*
*(iv) If $p > 0$ and $\alpha \in \mathbb{R}$, then $\lim_{n\to\infty} \frac{n^\alpha}{(1+p)^n} = 0$.*
*(v) If $|x| < 1$, then $\lim_{n\to\infty} x^n = 0$.*

*Proof.*
(i) Given $\varepsilon > 0$, choose $N > (1/\varepsilon)^{1/p}$ by Theorem 1.1.20. Then $n \ge N$ gives $n^{-p} < \varepsilon$.
(ii) If $p > 1$, set $x_n := \sqrt[n]{p} - 1$. Then $x_n > 0$ and $1 + nx_n \le (1 + x_n)^n = p$, so $0 < x_n \le (p-1)/n \to 0$. By Theorem 2.2.9, $x_n \to 0$. If $p = 1$, then $\sqrt[n]{p} = 1$ for every $n$. If $0 < p < 1$, then $1/p > 1$, so $\sqrt[n]{1/p} \to 1$ by the previous case. Since $\sqrt[n]{p} = 1/\sqrt[n]{1/p}$ and limits preserve reciprocals away from 0, $\sqrt[n]{p} \to 1$.
(iii) Put $x_n := \sqrt[n]{n} - 1$. Then $x_n \ge 0$, and for $n \ge 2$ the binomial theorem gives $n = (1 + x_n)^n \ge \frac{n(n-1)}{2} x_n^2$. Thus $0 \le x_n \le \sqrt{2/(n-1)}$, so Theorem 2.2.9 gives $x_n \to 0$.
(iv) Choose an integer $k > \alpha$ with $k > 0$. For $n > 2k$, $(1+p)^n \ge \binom{n}{k} p^k = \frac{n(n-1)\cdots(n-k+1)}{k!} p^k > \frac{n^k p^k}{2^k k!}$. Hence $0 < \frac{n^\alpha}{(1+p)^n} < \frac{2^k k!}{p^k} n^{\alpha-k}$, and $n^{\alpha-k} \to 0$ by (i). Apply Theorem 2.2.9.
(v) If $x \ne 0$, write $|x| = (1+p)^{-1}$ with $p > 0$ and apply (iv) with $\alpha = 0$ to get $|x|^n \to 0$. Since $-|x|^n \le x^n \le |x|^n$, Theorem 2.2.9 gives $x^n \to 0$; $x = 0$ is immediate.

*Structural Commentary*: Each limit is reduced to an elementary comparison sequence.
*Key Insight*: The binomial theorem supplies the right upper bounds for roots and exponential domination. □

**Proposition 2.2.11.** *Decimal Truncations Squeeze a Real*
*Let $x \ge 0$. For $n \ge 0$, set $s_n := \max\{k \in \mathbb{Z}_{\ge 0} : k10^{-n} \le x\}$.*
*Then $s_n 10^{-n} \le x < (s_n + 1)10^{-n}$; and $0 \le x - s_n 10^{-n} < 10^{-n}$; so $s_n 10^{-n} \to x$.*
*Moreover $s_{n+1} = 10s_n + a_{n+1}$ for some $a_{n+1} \in \{0, \dots, 9\}$.*

*Proof.* The maximum exists by the Archimedean property, Theorem 1.1.20, and well-ordering, Principle 2.6.2. Maximality gives the displayed inequalities. The squeeze and Theorem 2.2.10 give $s_n 10^{-n} \to x$.
At levels $n$ and $n+1$, $10s_n \le s_{n+1} < 10s_n + 10$, so $a_{n+1} := s_{n+1} - 10s_n \in \{0, \dots, 9\}$.
*Structural Commentary*: Decimal existence is an Archimedean grid argument followed by a squeeze.
*Key Insight*: Refining the mesh by a factor of 10 adds exactly one digit. □




Figure 124: At precision $10^{-n}$, the truncation $s_n 10^{-n}$ traps $x$ in a cell of width $10^{-n}$.

**Method 2.2.12.** *Sequence Proof Patterns*

(i) *Convergence*: fix $\varepsilon > 0$, then find $N$ so that all later terms lie in the relevant $\varepsilon$-ball.
(ii) *Uniqueness*: compare two proposed limits with $\varepsilon/2$ and use the triangle inequality.
(iii) *Subsequences*: choose $n_k \uparrow$ recursively with distances $< 1/k$.
(iv) *Cauchy arguments*: estimate the diameter of tails rather than guessing the limit first.
(v) *Monotone sequences*: use sup or inf and then prove eventual trapping.

*Key Insight*: Most sequence proofs are bookkeeping for quantifiers, indices, and one controlling inequality.

# 3 Metric Spaces and Euclidean Topology

## 3.1 Metrics, Open Sets, and Euclidean Topology

Metric topology studies sets through balls: openness, closure, and limit points come from distance. Euclidean estimates rely on norms, Cauchy–Schwarz, and the triangle inequality.

**Definition 3.1.1.** *Metric Space*
A *metric space* is a set $X$ equipped with a function $d : X \times X \to \mathbb{R}$ such that, for all $p, q, r \in X$,

i. $d(p,q) > 0$ if $p \neq q$, and $d(p,p) = 0$;
ii. $d(p,q) = d(q,p)$;
iii. $d(p,q) \le d(p,r) + d(r,q)$.

The function $d$ is called a *distance function* or *metric*.

**Definition 3.1.2.** *Metric Topology Terminology*
Let $X$ be a metric space, and let all points and sets below be understood as points of $X$ and subsets of $X$.

(i) For $p \in X$ and $r > 0$, the *open ball* centered at $p$ with radius $r$ is $B_r(p) := \{q \in X : d(p,q) < r\}$.
The corresponding *closed ball* is $\overline{B}_r(p) := \{q \in X : d(p,q) \le r\}$.
The *sphere* of radius $r$ about $p$ is $\{q \in X : d(p,q) = r\}$.
(ii) A *neighborhood* of $p$ is an open set $U \subseteq X$ with $p \in U$.
(iii) A point $p$ is a *limit point* of $E$ if every open ball $B_r(p)$ contains a point $q \neq p$ with $q \in E$.
(iv) If $p \in E$ and $p$ is not a limit point of $E$, then $p$ is an *isolated point* of $E$.
(v) A set $E$ is *closed* if every limit point of $E$ is a point of $E$.
(vi) A point $p$ is an *interior point* of $E$ if some neighborhood of $p$ is contained in $E$.
(vii) A set $E$ is *open* if every point of $E$ is an interior point of $E$.
(viii) The *complement* of $E$ is $E^c := X \setminus E$.
(ix) A set $E$ is *dense* in $X$ if every point of $X$ is either a point of $E$, a limit point of $E$, or both.

For $X = \mathbb{R}$, this gives $B_\varepsilon(a) = (a - \varepsilon, a + \varepsilon)$.
*Structural Commentary*: Balls generate the local tests; neighborhoods are open sets containing the point.

**Perspective 3.1.3.** *$\varepsilon$-Closeness Becomes Metric Language*
On $\mathbb{R}$, $\varepsilon$-close means $|x - y| < \varepsilon$, i.e. $d(x,y) < \varepsilon$ with $d(x,y) = |x - y|$; may be abstracted to metric balls.
*Key Insight*: Absolute value is distance on $\mathbb{R}$.

**Example 3.1.4.** *Euclidean Metric and Subspaces*
The Euclidean space $\mathbb{R}^k$ is a metric space with $d(\mathbf{x}, \mathbf{y}) := \|\mathbf{x} - \mathbf{y}\|_2, \qquad \mathbf{x}, \mathbf{y} \in \mathbb{R}^k$.
Every subset $Y$ of a metric space $X$ is a metric space by restricting the same distance function to points of $Y$.

**Example 3.1.5.** *Discrete Metric*
On any set $X$, define $d(x,y) = \begin{cases} 0, & x = y, \\ 1, & x \neq y. \end{cases}$
This is the *discrete metric*: distinct points are all distance 1 apart.

**Example 3.1.6.** *Hamming Distance*
For $x, y \in \{0,1\}^n$, define $h(x,y) := |\{i : x_i \neq y_i\}|$. Then $h$ is a metric.

*Proof.* $h \geq 0$, $h(x,y) = h(y,x)$, and $h(x,y) = 0 \iff x_i = y_i\ \forall i \iff x = y$.
For $x, y, z \in \{0,1\}^n$, $\{i : x_i \neq y_i\} \subseteq \{i : x_i \neq z_i\} \cup \{i : z_i \neq y_i\}$, so $h(x,y) \leq h(x,z) + h(z,y)$.
*Structural Commentary*: Hamming distance counts coordinate disagreements.
*Key Insight*: Each disagreement between $x$ and $y$ must be paid for through $z$ in at least one leg. □

**Definition 3.1.7.** *Euclidean Intervals, Cells, Balls, and Convexity*

(i) In $\mathbb{R}$, $(a,b) := \{x : a < x < b\}$, $[a,b] := \{x : a \leq x \leq b\}$, with $[a,b)$ and $(a,b]$ defined by mixed inequalities.
(ii) For $a_i < b_i$, a *k-cell* is $\prod_{i=1}^{k}[a_i,b_i] = \{\mathbf{x} \in \mathbb{R}^k : a_i \leq x_i \leq b_i,\ 1 \leq i \leq k\}$.
(iii) For $\mathbf{x} \in \mathbb{R}^k$ and $r > 0$, $B_r(\mathbf{x}) := \{\mathbf{y} : \|\mathbf{y} - \mathbf{x}\|_2 < r\}$ and $\overline{B}_r(\mathbf{x}) := \{\mathbf{y} : \|\mathbf{y} - \mathbf{x}\|_2 \leq r\}$.
(iv) $E \subseteq \mathbb{R}^k$ is *convex* iff $\lambda\mathbf{x} + (1-\lambda)\mathbf{y} \in E$ for all $\mathbf{x}, \mathbf{y} \in E$ and $0 \leq \lambda \leq 1$.
(v) The *integer lattice* is $\mathbb{Z}^k$, and the *first orthant* is $\{\mathbf{x} \in \mathbb{R}^k : x_i \geq 0\ \forall i\}$.
(vi) The *unit ball* is $B^k := \{\mathbf{x} \in \mathbb{R}^k : \|\mathbf{x}\|_2 \leq 1\}$, and the *unit sphere* is $S^{k-1} := \{\mathbf{x} \in \mathbb{R}^k : \|\mathbf{x}\|_2 = 1\}$.
(vii) A *convex combination* of $\mathbf{x}, \mathbf{y}$ is $\lambda\mathbf{x} + (1-\lambda)\mathbf{y}$ with $0 \leq \lambda \leq 1$; these points form the line segment $[\mathbf{x}, \mathbf{y}]$.

**Proposition 3.1.8.** *Balls and Cells Are Convex*
*Open balls, closed balls, and k-cells in $\mathbb{R}^k$ are convex.*

*Proof.* Let $\mathbf{y}, \mathbf{z} \in B_r(\mathbf{x})$ and $0 < \lambda < 1$.
Then $\|\lambda\mathbf{y} + (1-\lambda)\mathbf{z} - \mathbf{x}\|_2 = \|\lambda(\mathbf{y}-\mathbf{x}) + (1-\lambda)(\mathbf{z}-\mathbf{x})\|_2 \leq \lambda\|\mathbf{y}-\mathbf{x}\|_2 + (1-\lambda)\|\mathbf{z}-\mathbf{x}\|_2 < r$.
The same proof with $\leq r$ gives closed balls. For a $k$-cell, each coordinate of $\lambda\mathbf{y} + (1-\lambda)\mathbf{z}$ lies between the corresponding coordinates of $\mathbf{y}$ and $\mathbf{z}$, hence remains in the same interval $[a_i, b_i]$.
*Structural Commentary*: Convexity is checked by following the line segment between two points.
*Key Insight*: Balls use the triangle inequality; cells use coordinate inequalities. □

**Proposition 3.1.9.** *Bounded Compression of a Metric*
*If $(X,d)$ is a metric space, then $\rho(x,y) := d(x,y)/(1+d(x,y))$ is a bounded metric on $X$, and $\tau_\rho = \tau_d$.*

*Proof.* $0 \leq \rho < 1$, and definiteness and symmetry follow from $d$.
For $a, b \geq 0$, $(a+b)/(1+a+b) \leq a/(1+a) + b/(1+b)$.
Since $t/(1+t)$ is increasing, $d(x,z) \leq d(x,y) + d(y,z)$ gives $\rho(x,z) \leq \rho(x,y) + \rho(y,z)$.
For balls, $B_d(x,\eta) \subseteq B_\rho(x,\eta)$ for $0 < \eta < 1$, and $B_\rho(x,\eta) \subseteq B_d(x, \eta/(1-\eta))$; hence $\tau_\rho = \tau_d$.
*Structural Commentary*: Compressing distances into $[0,1)$ changes boundedness but not local neighborhoods.
*Key Insight*: Topology depends on small-scale control, not on the large-scale size of distances. □

**Definition 3.1.10.** *Isometries and Pseudometrics*
Let $(X,\rho)$ and $(Y,\sigma)$ be metric spaces, and let $F : X \to Y$.

(i) $F$ is an *isometry* from $(X,\rho)$ onto $(Y,\sigma)$ iff $F(X) = Y$ and $\forall x, y \in X$, $\sigma(Fx, Fy) = \rho(x,y)$.
(ii) $(X,\rho)$ and $(Y,\sigma)$ are *isometric*, written $(X,\rho) \cong (Y,\sigma)$, iff $\exists$ an isometry $F : (X,\rho) \to (Y,\sigma)$.

For a set $Z$, a *pseudometric* on $Z$ is a map $d : Z \times Z \to [0,\infty)$ such that $\forall x, y, z \in Z$, $d(x,x) = 0$, $d(x,y) = d(y,x)$, and $d(x,z) \leq d(x,y) + d(y,z)$; points $x \neq y$ may satisfy $d(x,y) = 0$.
*Structural Commentary*: Isometry preserves distance; pseudometrics permit zero-distance identification.

**Proposition 3.1.11.** *Metric Quotient of a Pseudometric*
*Let $X$ be a set and let $\rho$ be a pseudometric on $X$.*
*Define $x \sim_\rho y$ iff $\rho(x,y) = 0$, and define $\widetilde{\rho} : X/\!\sim_\rho \times X/\!\sim_\rho \to [0,\infty)$ by $\widetilde{\rho}([x],[y]) := \rho(x,y)$.*

*(i) $\sim_\rho$ is an equivalence relation on $X$.*
*(ii) $\widetilde{\rho}$ is a well-defined metric on $X/\!\sim_\rho$.*

*Proof.*

(i) For $x, y, z \in X$, $\rho(x,x) = 0$, $\rho(x,y) = 0 \Rightarrow \rho(y,x) = 0$;
$\rho(x,y) = \rho(y,z) = 0 \Rightarrow 0 \leq \rho(x,z) \leq \rho(x,y) + \rho(y,z) = 0$.
Hence $\sim_\rho$ is reflexive, symmetric, and transitive.
(ii) If $x \sim_\rho x'$ and $y \sim_\rho y'$, then $\rho(x,y) \leq \rho(x,x') + \rho(x',y') + \rho(y',y) = \rho(x',y')$, and the reverse inequality is identical; hence $\widetilde{\rho}$ is well-defined. For $[x],[y],[z] \in X/\!\sim_\rho$, nonnegativity, symmetry, and $\widetilde{\rho}([x],[z]) = \rho(x,z) \leq \rho(x,y) + \rho(y,z) = \widetilde{\rho}([x],[y]) + \widetilde{\rho}([y],[z])$ follow from $\rho$.
Finally, $\widetilde{\rho}([x],[y]) = 0$ iff $\rho(x,y) = 0$ iff $[x] = [y]$.

*Structural Commentary*: Collapse exactly zero-distance classes.
*Key Insight*: The quotient restores definiteness. □

**Example 3.1.12.** *Open, Closed, Both, or Neither*
In $\mathbb{R}$, $\emptyset, \mathbb{R}$ are open and closed, while $[a,b)$, $a < b$, is neither. If $d$ is discrete on $X$, then $B_{1/2}(x) = \{x\}$ for all $x \in X$, so every $E \subseteq X$ and $E^c$ are open; hence every $E \subseteq X$ is closed.

**Theorem 3.1.13.** *Every Open Ball Is Open*
*Every open ball in a metric space is an open set.*

*Proof.* Let $E = B_r(p)$, and choose $q \in E$. Put $h := r - d(p,q) > 0$.
If $s \in B_h(q)$, then $d(p,s) \le d(p,q) + d(q,s) < d(p,q) + h = r$.
Thus $B_h(q) \subseteq E$, so $q$ is an interior point of $E$. Since $q$ was arbitrary, $E$ is open.
*Structural Commentary*: Around any point inside a ball, the leftover distance to boundary gives a smaller ball.
*Key Insight*: The triangle inequality turns radius slack into openness. □




Figure 125: Every point inside a metric ball has a smaller ball still contained in the original ball.

**Theorem 3.1.14.** *Limit Points Force Infinitely Many Nearby Points*
*Let $(X,d)$ be a metric space, let $E \subseteq X$, and let $p \in X$. If $p$ is a limit point of $E$, then $N \cap E$ is infinite for every neighborhood $N$ of $p$.*

*Proof.* Let $N$ be a neighborhood of $p$ and suppose $N \cap E$ is finite. Since $N$ is open and $p \in N$, choose $\eta > 0$ with $B_\eta(p) \subseteq N$. Put $F := (N \cap E) \setminus \{p\}$. If $F = \emptyset$, then $B_\eta(p) \cap (E \setminus \{p\}) = \emptyset$, contradicting that $p$ is a limit point. If $F \neq \emptyset$, set $\delta := \min(\{\eta\} \cup \{d(p,q) : q \in F\}) > 0$. If $q \in B_\delta(p) \cap (E \setminus \{p\})$, then $q \in F$ and $d(p,q) < \delta \le d(p,q)$, impossible. Thus $B_\delta(p) \cap (E \setminus \{p\}) = \emptyset$, again contradicting that $p$ is a limit point.
*Structural Commentary*: A finite list of nearby exceptions has a smallest positive distance from $p$.
*Key Insight*: Limit points cannot be witnessed by only finitely many neighboring points. □

**Corollary 3.1.15.** *Finite Point Sets Have No Limit Points*
*A finite point set in a metric space has no limit points.*

*Proof.* Let $E \subseteq X$ be finite. If $p$ were a limit point of $E$, then by Theorem 3.1.14, every neighborhood of $p$ would contain infinitely many distinct points of $E$, impossible as $E$ is finite. Hence $E$ has no limit points.
*Structural Commentary*: This corollary is the finite obstruction to accumulation.
*Key Insight*: A limit point requires infinitely many nearby witnesses. □

**Theorem 3.1.16.** *Open Sets and Closed Complements*
*A set $E$ is open if and only if its complement $E^c$ is closed.*

*Proof.*
($\Rightarrow$) Suppose $E$ is open. If $x$ is a limit point of $E^c$ and $x \in E$, then some neighborhood $N$ of $x$ satisfies $N \subseteq E$, contradicting $N \cap (E^c \setminus \{x\}) \neq \emptyset$. Hence $x \in E^c$, so $E^c$ is closed.
($\Leftarrow$) Suppose $E^c$ is closed. If $x \in E$, then $x \notin E^c$, hence $x$ is not a limit point of $E^c$. Thus some neighborhood $N$ of $x$ has $N \cap E^c = \emptyset$, i.e. $N \subseteq E$. Hence $E$ is open.

*Structural Commentary*: Openness of a set is equivalent to closure of the outside.
*Key Insight*: Interior points of $E$ cannot be limit points of $E^c$. □

**Corollary 3.1.17.** *Closed Sets and Open Complements*
*A set $F$ is closed if and only if $F^c$ is open.*

*Proof.*
($\Rightarrow$) If $F$ is closed, then $(F^c)^c = F$ is closed; by Theorem 3.1.16 with $E = F^c$, $F^c$ is open.
($\Leftarrow$) If $F^c$ is open, then Theorem 3.1.16 gives $(F^c)^c$ closed; since $(F^c)^c = F$, $F$ is closed.

*Structural Commentary*: This is the same open–closed duality after taking complements.
*Key Insight*: Closedness of $F$ is openness of the outside $F^c$. □

**Corollary 3.1.18.** *Closed Balls Are Closed*
*For every $p \in X$ and $r > 0$, the closed ball $\overline{B}_r(p)$ is closed.*

*Proof.* Let $q \notin \overline{B}_r(p)$, so $d(p,q) > r$. Put $h := (d(p,q) - r)/2$.
If $s \in B_h(q)$, then $d(p,s) \geq d(p,q) - d(q,s) > d(p,q) - h > r$, so $s \notin \overline{B}_r(p)$.
Hence the complement of $\overline{B}_r(p)$ is open, and $\overline{B}_r(p)$ is closed.
*Structural Commentary*: Outside a closed ball, positive distance remains to the boundary.
*Key Insight*: Reverse triangle estimates certify closedness. □

**Theorem 3.1.19.** *Basic Topological Calculus*
*In a metric space:*

*(i) any union of open sets is open;*
*(ii) any intersection of closed sets is closed;*
*(iii) any finite intersection of open sets is open;*
*(iv) any finite union of closed sets is closed.*

*Proof.*

(i) Let $G := \bigcup_\alpha G_\alpha$. If $x \in G$, then $x \in G_\alpha$ for some $\alpha$.
Since $G_\alpha$ is open, some neighborhood of $x$ lies in $G_\alpha \subseteq G$. Hence $G$ is open.

(ii) Let $F := \bigcap_\alpha F_\alpha$. By Theorem 1.5.3, $F^c = \bigcup_\alpha F_\alpha^c$.
Each $F_\alpha^c$ is open by Corollary 3.1.17, so $F^c$ is open by (i). Hence $F$ is closed.

(iii) Let $H := \bigcap_{i=1}^n G_i$. If $x \in H$, let $r_i > 0$, $B_{r_i}(x) \subseteq G_i$, $r := \min_{1 \leq i \leq n} r_i$. Then $B_r(x) \subseteq H$, so $H$ is open.

(iv) Let $F := \bigcup_{i=1}^n F_i$. By Theorem 1.5.3, $F^c = \bigcap_{i=1}^n F_i^c$.
Each $F_i^c$ is open by Corollary 3.1.17, so $F^c$ is open by (iii). Hence $F$ is closed.

*Structural Commentary*: Arbitrary unions are easy for open sets; finite intersections require a minimum radius.
*Key Insight*: The finite minimum is the reason infinite intersections of open sets need not be open. □

**Theorem 3.1.20.** *Open Sets on the Real Line Split into Intervals*
*Let $\emptyset \neq G \subseteq \mathbb{R}$ be open.*

*(i) $G = \bigsqcup_{\alpha \in A} I_\alpha$ for a family of pairwise disjoint open intervals $I_\alpha \subseteq \mathbb{R}$, possibly unbounded.*
*(ii) The index set $A$ is at most countable.*

*Proof.*

(i) For $a, b \in \mathbb{R}$, set $I(a,b) := [\min\{a,b\}, \max\{a,b\}]$.
For $x, y \in G$, set $x \sim y$ iff $I(x,y) \subseteq G$, and let $C_x := [x]_\sim$. Then $\sim$ is an equivalence relation.
If $y \in C_x$ and $z \in I(x,y)$, then $I(x,z) \subseteq I(x,y) \subseteq G$, so $z \in C_x$; hence $C_x$ is an interval.
If $y \in C_x$, choose $r > 0$ with $(y-r, y+r) \subseteq G$; $z \in (y-r, y+r)$ implies $I(x,z) \subseteq I(x,y) \cup (y-r, y+r) \subseteq G$, so $z \in C_x$. Thus each $C_x$ is open, and the classes give a disjoint union of $G$.

(ii) Let $\mathcal{C} := G/\sim$ and enumerate $\mathbb{Q} = \{q_j : j \in \mathbb{N}\}$. Since each $C \in \mathcal{C}$ contains a bounded nonempty open interval, $C \cap \mathbb{Q} \neq \emptyset$. Define $\phi(C) := q_{\min\{j : q_j \in C\}}$. Disjointness gives $\phi$ injective, so $\mathcal{C}$ is at most countable.

*Structural Commentary*: Maximal open intervals are counted by rationals.
*Key Insight*: Density of $\mathbb{Q}$ turns disjoint interval geometry into countability. □

**Definition 3.1.21.** *Closure*
If $E \subseteq X$ and $E^{\text{acc}}$ denotes the set of all limit points of $E$ in $X$, then the *closure* of $E$ is $\overline{E} := E \cup E^{\text{acc}}$.

**Theorem 3.1.22.** *Closure Is the Smallest Closed Superset*
*If $X$ is a metric space and $E \subseteq X$, then:*

*(i) $\overline{E}$ is closed;*
*(ii) $\overline{E} = E$ if and only if $E$ is closed;*
*(iii) $\overline{E} \subseteq F$ for every closed set $F \subseteq X$ such that $E \subseteq F$.*

*Thus $\overline{E}$ is the smallest closed subset of $X$ that contains $E$.*

*Proof.*

(i) Let $p \notin \overline{E}$. Then $p \notin E$ and $p$ is not a limit point of $E$, so $B_r(p) \cap E = \emptyset$ for some $r > 0$.
If $q \in B_r(p)$, then $q \notin E$ and $B_{r-d(p,q)}(q) \subseteq B_r(p)$, so $q \notin E^{\text{acc}}$.
Hence $B_r(p) \subseteq (\overline{E})^c$, and $\overline{E}$ is closed by Corollary 3.1.17.

(ii) If $\overline{E} = E$, then $E$ is closed by (i). Conversely, if $E$ is closed, then $E^{\text{acc}} \subseteq E$, so $\overline{E} = E \cup E^{\text{acc}} = E$.

(iii) If $F$ is closed and $E \subseteq F$, then $E^{\text{acc}} \subseteq F^{\text{acc}} \subseteq F$. Thus $\overline{E} = E \cup E^{\text{acc}} \subseteq F$.

*Structural Commentary*: Closure adds exactly the limit points forced by $E$.
*Key Insight*: Closed supersets must contain all limit points, so they must contain $\overline{E}$. □




Figure 126: If $d = d(x, x_n) < r$, then any $0 < \rho < r - d$ gives $B_\rho(x_n) \subseteq B_r(x)$.

**Definition 3.1.23.** *Interior, Boundary, Adherence, and Distance*
Let $(X, d)$ be a metric space and $E, F \subseteq X$.

(i) $E^\circ := \{p \in E : \exists r > 0,\ B_r(p) \subseteq E\}$ is the *interior*.
(ii) $\text{ext}(E) := (E^c)^\circ = \{p \in X : \exists r > 0,\ B_r(p) \cap E = \emptyset\}$ is the *exterior*.
(iii) $\text{bd}(E) := \overline{E} \setminus E^\circ = \{p \in X : \forall r > 0,\ B_r(p) \cap E \neq \emptyset,\ B_r(p) \cap E^c \neq \emptyset\}$ is the *boundary*.
(iv) $p \in X$ is *adherent* to $E$ iff $\forall r > 0,\ B_r(p) \cap E \neq \emptyset$, equivalently $p \in \overline{E}$.
(v) For $p \in X$ and $\emptyset \neq E \subseteq X$, the *point-set distance* is $\text{dist}(p, E) := \inf\{d(p, q) : q \in E\}$.
(vi) For nonempty $E, F \subseteq X$, the *set-set distance* is $\text{dist}(E, F) := \inf\{d(x, y) : x \in E,\ y \in F\}$.




Figure 127: Interior, exterior, and boundary points are distinguished by how small balls meet $E$ and $E^c$.

**Proposition 3.1.24.** *Distance-to-Set Closure Criterion*
*Let $\emptyset \neq E \subseteq X$. For $p \in X$, $p \in \overline{E} \iff \text{dist}(p, E) = 0$.*

*Proof.*

($\Rightarrow$) If $p \in \overline{E}$, then $\forall \varepsilon > 0,\ B_\varepsilon(p) \cap E \neq \emptyset$; choose $q_\varepsilon \in E$ with $d(p, q_\varepsilon) < \varepsilon$.
Thus $0 \leq \text{dist}(p, E) \leq \varepsilon$ for all $\varepsilon > 0$, so $\text{dist}(p, E) = 0$.
($\Leftarrow$) If $\text{dist}(p, E) = 0$, then $\forall r > 0,\ \exists q \in E$ with $d(p, q) < r$.
Hence $B_r(p) \cap E \neq \emptyset$ for all $r > 0$, so $p$ is adherent to $E$, equivalently $p \in \overline{E}$.

*Structural Commentary*: Closure can be detected by vanishing distance to the set.
*Key Insight*: The infimum packages all small-ball tests into one number. □

**Remark 3.1.25.** *Set Distance Is Not a Metric on Subsets*
The function $\text{dist}(E, F)$ is useful, but it is not a metric on nonempty subsets.
In $\mathbb{R}$, for example, $\text{dist}((0, 1), (1, 2)) = 0$ although the sets are distinct and disjoint.

Figure 128: Adherent endpoints need not lie in $E$; isolated points are adherent but not limit points.

## 3.2 Relative Topology

**Definition 3.2.1.** *Relative Openness*
Suppose $E \subseteq Y \subseteq X$, where $X$ is a metric space. The set $E$ is *open relative to $Y$* if for each $p \in E$ there is an $r > 0$ such that $q \in E$ whenever $q \in Y$ and $d(p,q) < r$.

**Theorem 3.2.2.** *Relative Opens Are Intersections*
*Suppose $Y \subseteq X$. A subset $E$ of $Y$ is open relative to $Y$ if and only if $E = Y \cap G$ for some open subset $G$ of $X$.*

*Proof.*
($\Rightarrow$) For each $p \in E$, choose $r_p > 0$ with $B_{r_p}(p) \cap Y \subseteq E$, and set $G := \bigcup_{p \in E} B_{r_p}(p)$. Then $G$ is open in $X$, and $E \subseteq G \cap Y$. If $q \in G \cap Y$, then $q \in B_{r_p}(p) \cap Y$ for some $p \in E$, so $q \in E$. Thus $E = G \cap Y$.
($\Leftarrow$) If $E = G \cap Y$ with $G$ open in $X$, then for each $p \in E$ choose $r > 0$ with $B_r(p) \subseteq G$. Hence $B_r(p) \cap Y \subseteq G \cap Y = E$, so $E$ is open relative to $Y$.

*Structural Commentary*: Relative openness is ordinary openness viewed through the ambient subset $Y$.
*Key Insight*: Intersect open sets in $X$ with $Y$ to get exactly the open sets of $Y$. □

**Corollary 3.2.3.** *Relative Closed Sets Are Intersections*
*Let $(X,d)$ be a metric space, let $Y \subseteq X$, and let $F \subseteq Y$.*
*Then $F$ is closed relative to $Y$ iff $\exists C \subseteq X$ such that $C$ is closed in $X$ and $F = Y \cap C$.*

*Proof.*
($\Rightarrow$) If $F$ is closed relative to $Y$, then $Y \setminus F$ is open relative to $Y$. By Theorem 3.2.2, $\exists$ open $G \subseteq X$ such that $Y \setminus F = Y \cap G$. Put $C := X \setminus G$. Then $C$ is closed in $X$, and $F = Y \setminus (Y \setminus F) = Y \setminus (Y \cap G) = Y \cap C$.
($\Leftarrow$) If $F = Y \cap C$ with $C$ closed in $X$, then $X \setminus C$ is open in $X$ and $Y \setminus F = Y \setminus (Y \cap C) = Y \cap (X \setminus C)$. By Theorem 3.2.2, $Y \setminus F$ is open relative to $Y$, so $F$ is closed relative to $Y$.

*Structural Commentary*: Relative closedness is ordinary closedness seen through the ambient subset.
*Key Insight*: The closed version is the complement of the relative-open intersection theorem. □

**Example 3.2.4.** *Relative Topology Depends on the Ambient Space*
Let $X = \{(x,0) : x \in \mathbb{R}\} \subseteq \mathbb{R}^2$ with the metric inherited from $\mathbb{R}^2$. The set $E = \{(x,0) : -1 < x < 1\}$ is not open as a subset of $\mathbb{R}^2$, but it is open relative to $X$, since $E = X \cap \big((-1,1) \times (-1,1)\big)$.
Likewise, $[0,1)$ is not closed in $\mathbb{R}$, but it is closed relative to $(-1,1)$ because $[0,1) = (-1,1) \cap [0,\infty)$.

## 3.3 Density, Separability, and Closure Tests

**Proposition 3.3.1.** *Density and Closure*
*Let $X$ be a metric space and let $E \subseteq X$. Then $E$ is dense in $X$ if and only if $\overline{E} = X$.*

*Proof.*
($\Rightarrow$) Suppose $E$ is dense in $X$. Let $p \in X$. By Definition 3.1.2, $p$ is either a point of $E$ or a limit point of $E$. Hence $p \in E \cup E^{\text{acc}} = \overline{E}$ by Definition 3.1.21. Thus $X \subseteq \overline{E}$. Since $\overline{E} \subseteq X$, $\overline{E} = X$.
($\Leftarrow$) Suppose $\overline{E} = X$. Let $p \in X$. Then $p \in \overline{E} = E \cup E^{\text{acc}}$, so $p$ is either a point of $E$ or a limit point of $E$. Therefore $E$ is dense in $X$ by Definition 3.1.2.

*Structural Commentary*: Density says every point is already in $E$ or is forced by nearby points of $E$.
*Key Insight*: The closure packages exactly those two possibilities. □

**Proposition 3.3.2.** *Real-Line Density Interval Test*
*Let $E \subseteq \mathbb{R}$. Then $E$ is dense in $\mathbb{R}$ if and only if every nonempty open interval $(a,b)$ satisfies $(a,b) \cap E \neq \emptyset$.*

*Proof.*
($\Rightarrow$) Let $(a,b) \neq \emptyset$ and choose $p \in (a,b)$, $r := \frac{1}{2}\min\{p-a, b-p\}$. If $p \in E$, done. If $p \notin E$, density gives $p \in E^{\text{acc}}$, so choose $q \in (E \setminus \{p\}) \cap B_r(p)$. Since $B_r(p) \subseteq (a,b)$, $q \in (a,b) \cap E$.
($\Leftarrow$) Suppose $(a,b) \cap E \neq \emptyset$ for every nonempty open interval $(a,b)$.
For $p \in \mathbb{R}$, either $p \in E$, or $p \notin E$ and $\forall \varepsilon > 0$, choose $q \in (p-\varepsilon, p+\varepsilon) \cap E$.
Then $q \neq p$ and $q \in B_\varepsilon(p)$, so $p \in E^{\text{acc}}$. Hence $\mathbb{R} = E \cup E^{\text{acc}} = \overline{E}$, so $E$ is dense in $\mathbb{R}$.

*Structural Commentary*: On $\mathbb{R}$, balls and open intervals test the same local condition.
*Key Insight*: Density means no interval can avoid the set. □

**Definition 3.3.3.** *Separable Metric Space*
A metric space $X$ is *separable* if it contains a countable dense subset.

**Example 3.3.4.** *Separable and Nonseparable Metric Spaces*

(i) $\mathbb{R}$ is separable: $\mathbb{Q}$ is countable and dense. Hence $\mathbb{R}^k$ is separable via $\mathbb{Q}^k$, and $\mathbb{C}^k$ via $\mathbb{C}^k \simeq \mathbb{R}^{2k}$.
(ii) If $X$ is discrete, then $X$ is separable $\iff$ $X$ is countable,
since $B_{1/2}(x) = \{x\}$ forces every dense $D \subseteq X$ to contain $x$.

*Structural Commentary*: Dense rational grids handle Euclidean spaces; discrete balls force dense sets to contain every point.
*Key Insight*: Separability depends on whether countable data can meet every open ball.

**Theorem 3.3.5.** *Suprema Belong to Closures*
*Let $\emptyset \neq E \subseteq \mathbb{R}$ be bounded above and $y := \sup E$. Then $y \in \overline{E}$; if $E$ is closed, then $y \in E$.*

*Proof.* If $y \in E$, then $y \in \overline{E}$. If $y \notin E$, then $\forall h > 0$, $y - h$ is not an upper bound; choose $x_h \in E$ with $y - h < x_h \leq y$. Thus $x_h \neq y$, $|x_h - y| < h$, and $B_h(y) \cap (E \setminus \{y\}) \neq \emptyset$, so $y \in E^{\text{acc}} \subseteq \overline{E}$.
If $E$ is closed, $\overline{E} = E$ by Theorem 3.1.22(ii).
*Structural Commentary*: LUB gives points of $E$ arbitrarily close below $\sup E$.
*Key Insight*: A supremum is either attained or approached. □

## 3.4 Bounded Sets and Metric Sequences

**Definition 3.4.1.** *Bounded Sets and Metric Sequences*
Let $(X, d)$ be a metric space, $E \subseteq X$, and $(p_n)$ a sequence in $X$.

(i) $E$ is *bounded* iff $\exists q \in X, \exists M > 0$ such that $E \subseteq B_M(q)$
(ii) $E$ is *perfect* iff $E$ is closed and every $p \in E$ is a limit point of $E$.
(iii) $(p_n)$ *converges* to $p \in X$ iff $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ such that $n \geq N \implies d(p_n, p) < \varepsilon$; write $p_n \to p$ or $\lim_{n\to\infty} p_n = p$.
If no such $p$ exists, $(p_n)$ *diverges*.
(iv) The *range* of $(p_n)$ is $\text{ran}(p_n) := \{p_n : n \in \mathbb{N}\}$; $(p_n)$ is *bounded* iff $\text{ran}(p_n)$ is bounded.
(v) $(p_n)$ is *eventually in $E$* iff $\exists N \in \mathbb{N}$ such that $n \geq N \implies p_n \in E$
(vi) $(p_n)$ is *frequently in $E$* iff $\forall N \in \mathbb{N}, \exists n \geq N$ such that $p_n \in E$.

*Structural Commentary*: Boundedness, compactness, and completeness are intrinsic to restricted metrics; openness and closedness are ambient.
*Key Insight*: Convergence is tail control; frequent membership is arbitrarily late recurrence.

**Theorem 3.4.2.** *Basic Properties of Metric Convergence*
*Let $X$ be a metric space and $(p_n) \subseteq X$.*

*(i) $p_n \to p \iff \forall r > 0, \exists N \in \mathbb{N}$ such that $n \geq N \implies p_n \in B_r(p)$.*
*(ii) If $p_n \to p$ and $p_n \to p'$, then $p = p'$.*
*(iii) If $E \subseteq X$ and $p \in E^{\text{acc}}$, then $\exists (q_n) \subseteq E \setminus \{p\}$ such that $q_n \to p$.*

*Proof.*

(i) ($\Rightarrow$) If $p_n \to p$ and $r > 0$, take $\varepsilon = r$ in the definition.
(i) ($\Leftarrow$) If the ball condition holds, then for every $\varepsilon > 0$ take $r = \varepsilon$, giving $d(p_n, p) < \varepsilon$ eventually.
(ii) Given $\varepsilon > 0$, choose $N, N'$ with $n \geq N \implies d(p_n, p) < \varepsilon/2$ and $n \geq N' \implies d(p_n, p') < \varepsilon/2$.
If $n \geq \max\{N, N'\}$, then $d(p, p') \leq d(p, p_n) + d(p_n, p') < \varepsilon$, so $d(p, p') = 0$.
(iii) Since $p \in E^{\text{acc}}$, for each $n \in \mathbb{N}$ choose $q_n \in E \setminus \{p\}$ with $d(q_n, p) < 1/n$. Then $q_n \to p$.

*Structural Commentary*: Eventual containment gives convergence; triangle inequality gives uniqueness.
*Key Insight*: A limit point can always be approached by a sequence of points from the set. □

**Corollary 3.4.3.** *Subsequences Preserve Metric Limits*
*If $p_n \to p$ in a metric space, then every subsequence $(p_{n_k})$ also converges to $p$.*

*Proof.* Given $\varepsilon > 0$, choose $N$ such that $n \geq N$ implies $d(p_n, p) < \varepsilon$.
Since $n_k \to \infty$, all sufficiently large $k$ satisfy $n_k \geq N$, hence $d(p_{n_k}, p) < \varepsilon$.
*Structural Commentary*: A subsequence is only a delayed tail of the same sequence.
*Key Insight*: Eventual estimates survive deleting terms. □

**Proposition 3.4.4.** *Limit Points Are Subsequence Limits*
*For $(p_n) \subseteq X$ and $p \in X$, the following are equivalent:*

*(i) $\forall r > 0, \forall N \in \mathbb{N}, \exists n \geq N$ such that $p_n \in B_r(p)$.*
*(ii) $\exists n_1 < n_2 < \cdots$ such that $p_{n_k} \to p$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) Set $n_0 = 0$. By (i), recurse $n_k > n_{k-1}$ with $p_{n_k} \in B_{1/k}(p)$; then $d(p_{n_k}, p) < 1/k$, so $p_{n_k} \to p$.
(*ii*) $\Rightarrow$ (*i*) If $p_{n_k} \to p$, then for any $r > 0$ and $N \in \mathbb{N}$, choose $k$ with $n_k \geq N$ and $d(p_{n_k}, p) < r$. Thus $p_{n_k} \in B_r(p)$, so (i) holds.

*Structural Commentary*: Frequent visits are exactly enough to extract a convergent subsequence.
*Key Insight*: Limits require all sufficiently late terms; limit points require arbitrarily late witnesses. □

**Proposition 3.4.5.** *Discrete-Metric Convergence*
*Let $X$ carry the discrete metric $d(x, y) = 0$ if $x = y$ and $d(x, y) = 1$ if $x \neq y$.*
*For $(p_n) \subset X$ and $p \in X$, $p_n \to p$ if and only if $\exists N \in \mathbb{N}$ such that $n \geq N \Longrightarrow p_n = p$.*

*Proof.*

($\Rightarrow$) Suppose $p_n \to p$. For $\varepsilon = 1/2$, choose $N \in \mathbb{N}$ such that $n \geq N \Longrightarrow d(p_n, p) < 1/2$. Since $d$ takes only the values $0, 1$, $d(p_n, p) = 0$, hence $p_n = p$ for all $n \geq N$.
($\Leftarrow$) Suppose $\exists N \in \mathbb{N}$ s.t. $n \geq N \Longrightarrow p_n = p$. Then $\forall \varepsilon > 0$, $n \geq N$ gives $d(p_n, p) = 0 < \varepsilon$, so $p_n \to p$.

*Structural Commentary*: The discrete metric has no room for distinct points to approach one another.
*Key Insight*: In the discrete metric, $d(x, y) < 1 \Longleftrightarrow x = y$. □

**Example 3.4.6.** *Coordinate Norm Metrics on $\mathbb{R}^k$*
By Proposition 1.1.21 and Theorem 1.1.13, $d_1(\mathbf{x}, \mathbf{y}) := \|\mathbf{x} - \mathbf{y}\|_1$; $d_2(\mathbf{x}, \mathbf{y}) := \|\mathbf{x} - \mathbf{y}\|_2$; $d_\infty(\mathbf{x}, \mathbf{y}) := \|\mathbf{x} - \mathbf{y}\|_\infty$ are metrics on $\mathbb{R}^k$, and $\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^k$, $d_\infty(\mathbf{x}, \mathbf{y}) \leq d_2(\mathbf{x}, \mathbf{y}) \leq d_1(\mathbf{x}, \mathbf{y}) \leq \sqrt{k}\, d_2(\mathbf{x}, \mathbf{y})$; $d_2(\mathbf{x}, \mathbf{y}) \leq \sqrt{k}\, d_\infty(\mathbf{x}, \mathbf{y})$.
Hence $d_1, d_2, d_\infty$ have the same convergent sequences.

**Theorem 3.4.7.** *Coordinatewise Convergence in $\mathbb{R}^k$*
*Let $k \in \mathbb{N}$, $\mathbf{x}_n = (\alpha_{j,n})$, $\mathbf{x} = (\alpha_j)$, $\mathbf{y}_n = (\gamma_{j,n})$, $\mathbf{y} = (\gamma_j)$ in $\mathbb{R}^k$, and let $(\beta_n) \subset \mathbb{R}$, $\beta \in \mathbb{R}$.*

(*i*) *$\mathbf{x}_n \to \mathbf{x}$ iff $\alpha_{j,n} \to \alpha_j$ for every $1 \leq j \leq k$.*
(*ii*) *If $\mathbf{x}_n \to \mathbf{x}$ and $\mathbf{y}_n \to \mathbf{y}$, then $\mathbf{x}_n + \mathbf{y}_n \to \mathbf{x} + \mathbf{y}$.*
(*iii*) *If $\mathbf{x}_n \to \mathbf{x}$ and $\mathbf{y}_n \to \mathbf{y}$, then $\mathbf{x}_n \cdot \mathbf{y}_n \to \mathbf{x} \cdot \mathbf{y}$.*
(*iv*) *If $\beta_n \to \beta$ and $\mathbf{x}_n \to \mathbf{x}$, then $\beta_n \mathbf{x}_n \to \beta \mathbf{x}$.*

*Proof.*

(*i*) ($\Rightarrow$) Since $|\alpha_{j,n} - \alpha_j| \leq \|\mathbf{x}_n - \mathbf{x}\|_2$ for each $j$, each coordinate converges.
(*i*) ($\Leftarrow$) For $\varepsilon > 0$, eventually $|\alpha_{j,n} - \alpha_j| < \varepsilon/\sqrt{k}$ for all $j$; hence $\|\mathbf{x}_n - \mathbf{x}\|_2^2 < k(\varepsilon^2/k) = \varepsilon^2$.
(*ii*) For each $j$, $\alpha_{j,n} + \gamma_{j,n} \to \alpha_j + \gamma_j$; apply (i).
(*iii*) Each product $\alpha_{j,n}\gamma_{j,n} \to \alpha_j\gamma_j$, hence $\sum_{j=1}^k \alpha_{j,n}\gamma_{j,n} \to \sum_{j=1}^k \alpha_j\gamma_j$.
(*iv*) For each $j$, $\beta_n\alpha_{j,n} \to \beta\alpha_j$; apply (i).

*Structural Commentary*: Euclidean convergence is exactly simultaneous coordinate convergence.
*Key Insight*: The norm packages finitely many coordinate estimates into one estimate. □

**Definition 3.4.8.** *Product Metrics*
For metric spaces $(X, d_X)$ and $(Y, d_Y)$, and $p = (x, y)$, $p' = (x', y')$ in $X \times Y$, define

$$d_E(p, p') := \sqrt{d_X(x, x')^2 + d_Y(y, y')^2},$$
$$d_{\max}(p, p') := \max\{d_X(x, x'), d_Y(y, y')\},$$
$$d_{\text{sum}}(p, p') := d_X(x, x') + d_Y(y, y')$$

These are the Euclidean, maximum, and sum product metrics; the sum metric is often simplest for estimates.




Figure 129: Unit balls in $\mathbb{R}^2$: $\max\{|x|, |y|\} < 1$ and $|x| + |y| < 1$.

**Proposition 3.4.9.** *Comparing Product Metrics*
*On $X \times Y$, $d_{\max} \le d_E \le d_{\text{sum}} \le 2d_{\max}$. Hence the three product metrics have same convergent sequences.*

*Proof.* Let $a = d_X(x, x')$ and $b = d_Y(y, y')$.
Then $\max\{a, b\} \le \sqrt{a^2 + b^2} \le a + b \le 2\max\{a, b\}$; convergence follows by squeezing.
*Structural Commentary*: The three product metrics differ by controlled constants.
*Key Insight*: Equivalent metric estimates preserve convergence. □

**Definition 3.4.10.** *Equivalent Metrics*
Let $X$ be a set and let $\rho, \sigma : X \times X \to [0, \infty)$ be metrics.
They are *equivalent*, written $\rho \sim \sigma$, iff $\exists c_1, c_2 > 0$ with $\forall x, y \in X$, $c_1\sigma(x, y) \le \rho(x, y) \le c_2\sigma(x, y)$.

**Proposition 3.4.11.** *Equivalent Metrics Preserve Topological Tests*
*Let $X$ be a set, let $\rho, \sigma : X \times X \to [0, \infty)$ be metrics with $\rho \sim \sigma$, and let $\tau_\rho, \tau_\sigma$ be the induced topologies.*

*(i)* $\tau_\rho = \tau_\sigma$.
*(ii) For every $E \subseteq X$, $E$ is $\rho$-closed iff $E$ is $\sigma$-closed.*
*(iii) For every sequence $(x_n)$ in $X$ and every $x \in X$, $\rho(x_n, x) \to 0$ iff $\sigma(x_n, x) \to 0$.*

*Proof.* Choose $c_1, c_2 > 0$ such that $c_1\sigma(u, v) \le \rho(u, v) \le c_2\sigma(u, v)$ for all $u, v \in X$.
For $x \in X$ and $r > 0$, $B_\sigma(x, r/c_2) \subseteq B_\rho(x, r) \subseteq B_\sigma(x, r/c_1)$.

(i) If $U \in \tau_\rho$ and $x \in U$, choose $r > 0$ with $B_\rho(x, r) \subseteq U$; then $B_\sigma(x, r/c_2) \subseteq U$, so $U \in \tau_\sigma$.
If $U \in \tau_\sigma$ and $x \in U$, choose $s > 0$ with $B_\sigma(x, s) \subseteq U$; then $B_\rho(x, c_1 s) \subseteq U$, so $U \in \tau_\rho$.
(ii) Let $E \subseteq X$. By (i), $E^c \in \tau_\rho$ iff $E^c \in \tau_\sigma$; hence $E$ is $\rho$-closed iff $E$ is $\sigma$-closed.
(iii) For every $n \in \mathbb{N}$, $c_1\sigma(x_n, x) \le \rho(x_n, x) \le c_2\sigma(x_n, x)$; hence $\rho(x_n, x) \to 0$ iff $\sigma(x_n, x) \to 0$.

*Structural Commentary*: Ball inclusions transfer local tests between metrics.
*Key Insight*: Constant-comparison estimates let each metric ball contain a smaller ball for the other metric. □

**Proposition 3.4.12.** *Sequential Closure Criterion*
*Let $E \subseteq X$, where $X$ is a metric space, and let $p \in X$. Then $p \in \overline{E} \iff \exists (p_n)_{n \ge 1}$ with $p_n \in E$ $\forall n$, $p_n \to p$.*

*Proof.*

($\Rightarrow$) If $p \in \overline{E}$, choose $p_n \in E \cap B_{1/n}(p)$ for $n \ge 1$ by Principle 6.5.2. Then $d(p_n, p) < 1/n$, so $p_n \to p$.
($\Leftarrow$) If $p_n \in E$ and $p_n \to p$, then every ball $B_r(p)$ contains some $p_n$.
Thus $p$ is adherent to $E$ by Definition 3.1.23, so $p \in \overline{E}$.

*Structural Commentary*: Closure can be read either by neighborhoods or by sequences.
*Key Insight*: Adherence supplies one witness at every scale. □

**Theorem 3.4.13.** *Closed Sets and Limit Points via Sequences*
*Let $X$ be a metric space and $E \subseteq X$.*

*(i)* $p \in E^{\text{acc}} \iff \exists (p_n) \subseteq E \setminus \{p\}$ *such that* $p_n \to p$.
*(ii) $E$ is closed $\iff$ every $(p_n) \subseteq E$ with $p_n \to p \in X$ satisfies $p \in E$.*

*Proof.*

*(i)* ($\Rightarrow$) If $p \in E^{\text{acc}}$, choose $p_n \in (E \setminus \{p\}) \cap B_{1/n}(p)$ for each $n$. Then $d(p_n, p) < 1/n$, so $p_n \to p$.
*(i)* ($\Leftarrow$) If $(p_n) \subseteq E \setminus \{p\}$ and $p_n \to p$, then every $B_r(p)$ contains some $p_n \ne p$. Hence $p \in E^{\text{acc}}$.
*(ii)* ($\Rightarrow$) If $E$ is closed, $(p_n) \subseteq E$, and $p_n \to p$, then Proposition 3.4.12 gives $p \in \overline{E} = E$.
*(ii)* ($\Leftarrow$) If every convergent sequence in $E$ has limit in $E$, then Proposition 3.4.12 gives $\overline{E} \subseteq E$. QED.

*Structural Commentary*: In metric spaces, closedness can be tested by sequences.
*Key Insight*: Balls at radii $1/n$ convert local information into a sequence. □

**Theorem 3.4.14.** *Subsequential Limits Form a Closed Set*
*For a sequence $(p_n)$ in a metric space $X$, the set of all subsequential limits is closed.*

*Proof.* Let $E^*$ be the set of subsequential limits and let $q$ be a limit point of $E^*$. Set $n_0 = 0$.
For each $i$, choose $x_i \in E^*$ with $d(x_i, q) < 2^{-i}$. Since $x_i$ is a subsequential limit, choose $n_i > n_{i-1}$ such that $d(p_{n_i}, x_i) < 2^{-i}$. Then $d(p_{n_i}, q) < 2^{1-i}$, so $p_{n_i} \to q$, and $q \in E^*$.
*Structural Commentary*: A limit of subsequential limits is reached by diagonalizing through those subsequences.
*Key Insight*: Choose later and closer terms at the same time. □

**Method 3.4.15.** *Metric Proof Patterns*

(i) To prove $G$ is open, fix $x \in G$ and find $r > 0$ with $B_r(x) \subseteq G$.
(ii) To prove $F$ is closed, show $F^{\text{acc}} \subseteq F$, or equivalently show $F^c$ is open.

(iii) To prove $C = \overline{E}$, show $E \subseteq C$, $C$ is closed, and $C \subseteq F$ for every closed $F \supseteq E$.
(iv) To prove $p \in \overline{E}$, show $\operatorname{dist}(p, E) = 0$, or build a sequence in $E$ converging to $p$.

# 4 Compactness and Completeness

## 4.1 Compactness, Completeness, and Total Boundedness

Compactness extracts finite or convergent structure from infinite behavior; completeness asks whether Cauchy approximations converge internally.

**Definition 4.1.1.** *Open Covers, Compact Sets, and Finite Intersection Property*
Let $(X, d)$ be a metric space.

(i) For $E \subseteq X$, an *open cover* of $E$ is a family $(G_\alpha)_{\alpha \in A}$ such that each $G_\alpha \subseteq X$ is open and $E \subseteq \bigcup_{\alpha \in A} G_\alpha$.
(ii) A set $K \subseteq X$ is *compact* iff every open cover $(G_\alpha)_{\alpha \in A}$ of $K$ admits $\alpha_1, \dots, \alpha_n \in A$ with $K \subseteq \bigcup_{j=1}^{n} G_{\alpha_j}$.
(iii) Family $\mathcal{F} \subseteq \mathcal{P}(X)$ has *finite intersection property* iff $\bigcap_{F \in \mathcal{F}_0} F \neq \emptyset$ $\forall$ finite nonempty subfamily $\mathcal{F}_0 \subseteq \mathcal{F}$.

**Proposition 4.1.2.** *Closed-Set Compactness Test*
*Let $(X, d)$ be a metric space. Then $X$ is compact iff every nonempty family $\mathcal{F}$ of closed subsets of $X$ with the finite intersection property satisfies $\bigcap_{F \in \mathcal{F}} F \neq \emptyset$.*

*Proof.*

(⇒) Assume $X$ is compact. Let $\emptyset \neq \mathcal{F} \subseteq \mathcal{P}(X)$ consist of closed sets and have the finite intersection property. If $\bigcap_{F \in \mathcal{F}} F = \emptyset$, then $\{X \setminus F : F \in \mathcal{F}\}$ is an open cover of $X$. Compactness gives $F_1, \dots, F_n \in \mathcal{F}$ with $X = \bigcup_{j=1}^{n} (X \setminus F_j)$; hence $\bigcap_{j=1}^{n} F_j = \emptyset$, contradicting the finite intersection property.
(⇐) Assume every nonempty closed family with the finite intersection property has nonempty intersection. Let $(G_\alpha)_{\alpha \in A}$ be an open cover of $X$. If no finite subcover exists, then $\mathcal{F} := \{X \setminus G_\alpha : \alpha \in A\}$ is a nonempty family of closed subsets of $X$ with the finite intersection property. Thus $\exists x \in \bigcap_{\alpha \in A} (X \setminus G_\alpha)$, contradicting $X \subseteq \bigcup_{\alpha \in A} G_\alpha$. Hence every open cover has a finite subcover, so $X$ is compact.

*Structural Commentary*: Complements turn finite subcovers into finite intersections.
*Key Insight*: Compactness can be read dually through closed sets. □

Figure 130: Compactness extracts a finite subcover.

**Example 4.1.3.** *An Open Cover Without a Finite Subcover*
For $x \in (0, 1)$, set $O_x := (x/2, 1)$. These sets cover $(0, 1)$, since $y \in O_x$ whenever $0 < x < y < 1$.
No finite subcollection covers $(0, 1)$: if $x' = \min\{x_1, \dots, x_n\}$, then $(0, x'/2]$ is missed by $O_{x_1} \cup \cdots \cup O_{x_n}$.




Figure 131: For finite choice, $x' = \min_i x_i$ makes $O_{x'}$ largest selected interval, but $(0, x'/2]$ remains uncovered.

**Theorem 4.1.4.** *Compactness Is Independent of the Ambient Space*
*Suppose $K \subseteq Y \subseteq X$, where $X$ is a metric space.*
*Then $K$ is compact relative to $X$ if and only if $K$ is compact relative to $Y$.*

*Proof.*

($\Rightarrow$) Let $\{V_\alpha\}_{\alpha\in A}$ be a relative-open cover of $K$ in $Y$.
By Theorem 3.2.2, write $V_\alpha = Y \cap G_\alpha$ with $G_\alpha$ open in $X$. Since $\{G_\alpha\}_{\alpha\in A}$ covers $K$, compactness in $X$ gives a finite $F \subseteq A$ such that $K \subseteq \bigcup_{\alpha\in F} G_\alpha$. As $K \subseteq Y$, $K \subseteq \bigcup_{\alpha\in F} V_\alpha$.

($\Leftarrow$) Let $\{G_\alpha\}_{\alpha\in A}$ be an open cover of $K$ in $X$. Then $\{Y \cap G_\alpha\}_{\alpha\in A}$ is a relative-open cover of $K$ in $Y$.
Compactness in $Y$ gives a finite $F \subseteq A$ such that $K \subseteq \bigcup_{\alpha\in F}(Y \cap G_\alpha) \subseteq \bigcup_{\alpha\in F} G_\alpha$.

*Structural Commentary*: Relative covers are ambient covers intersected with $Y$.
*Key Insight*: Openness depends on the ambient space, but compactness does not. □

**Theorem 4.1.5.** *Compact Sets Are Closed*
*Every compact subset of a metric space is closed.*

*Proof.* Let $K$ be compact in $X$. To prove $K^c$ is open.
Fix $p \in K^c$. For each $q \in K$, put $V_q := B_{\frac{1}{2}d(p,q)}(p)$, $W_q := B_{\frac{1}{2}d(p,q)}(q)$.
The sets $W_q$ form an open cover of $K$, so compactness gives $q_1, \dots, q_n \in K$ with $K \subseteq W_{q_1} \cup \cdots \cup W_{q_n}$.
Let $V := V_{q_1} \cap \cdots \cap V_{q_n}$, which is open by Theorem 3.1.19.
No point of $V$ lies in any $W_{q_i}$, by the triangle inequality. Hence $V \cap K = \emptyset$, so $p$ is an interior point of $K^c$.
*Structural Commentary*: Compactness makes the separation finite.
*Key Insight*: A finite subcover lets local point-separations become one open neighborhood of the complement. □

**Theorem 4.1.6.** *Closed Subsets of Compact Sets Are Compact*
*Suppose $F \subseteq K \subseteq X$, where $F$ is closed in $X$ and $K$ is compact. Then $F$ is compact.*

*Proof.* Let $\{V_\alpha\}_{\alpha\in A}$ cover $F$. Since $F^c$ is open, $\{V_\alpha\}_{\alpha\in A} \cup \{F^c\}$ covers $K$.
Thus $K \subseteq F^c \cup \bigcup_{\alpha\in F_0} V_\alpha$ for some finite $F_0 \subseteq A$, so $F \subseteq \bigcup_{\alpha\in F_0} V_\alpha$.
*Structural Commentary*: The open complement of a closed set covers exactly the part of $K$ not under discussion.
*Key Insight*: Cover $K$, then discard the complement piece. □

**Corollary 4.1.7.** *Closed Intersections with Compact Sets*
*If $F$ is closed in $X$ and $K$ is compact in $X$, then $F \cap K$ is compact.*

*Proof.* By Theorem 4.1.5, $K$ is closed in $X$. Since $F$ is closed, Theorem 3.1.19 gives that $F \cap K$ is closed in $X$.
Also $F \cap K \subseteq K$, so Theorem 4.1.6 implies $F \cap K$ is compact.
*Structural Commentary*: The intersection is closed inside a compact set.
*Key Insight*: Closed subsets of compact sets remain compact. □

**Theorem 4.1.8.** *Finite Intersection Property for Compact Sets*
*Let $\{K_\alpha\}_{\alpha\in A}$ be compact subsets of a metric space $X$, with $A \neq \emptyset$.*
*If $\bigcap_{\alpha\in F} K_\alpha \neq \emptyset$ for every finite nonempty $F \subseteq A$, then $\bigcap_{\alpha\in A} K_\alpha \neq \emptyset$.*

*Proof.* Choose $\alpha_0 \in A$ and suppose $\bigcap_{\alpha\in A} K_\alpha = \emptyset$. Then $K_{\alpha_0} \subseteq \bigcup_{\alpha\in A} K_\alpha^c$. Since each $K_\alpha^c$ is open by Theorem 4.1.5, compactness of $K_{\alpha_0}$ gives a finite $F \subseteq A$ with $K_{\alpha_0} \subseteq \bigcup_{\alpha\in F} K_\alpha^c$. Hence $\bigcap_{\alpha\in F\cup\{\alpha_0\}} K_\alpha = \emptyset$, a contradiction.
*Structural Commentary*: Complements turn failure into a finite cover.
*Key Insight*: For compact sets, failure of intersection is already witnessed finitely. □

**Corollary 4.1.9.** *Nested Nonempty Compact Sets Intersect*
*If $K_1 \supseteq K_2 \supseteq K_3 \supseteq \cdots$ is a sequence of nonempty compact sets, then $\bigcap_{n=1}^{\infty} K_n \neq \emptyset$.*

*Proof.* Let $\emptyset \neq F \subseteq \mathbb{N}$ be finite and put $m := \max F$. Since the sequence is nested, $\bigcap_{n\in F} K_n = K_m \neq \emptyset$.
Thus $\{K_n\}_{n\in\mathbb{N}}$ has the finite intersection property, so Theorem 4.1.8 gives $\bigcap_{n=1}^{\infty} K_n \neq \emptyset$.
*Structural Commentary*: Nestedness reduces finite intersections to one term.
*Key Insight*: A nested compact sequence has the finite intersection property. □

**Theorem 4.1.10.** *Infinite Subsets of Compact Sets Have Limit Points*
*Let $K \subseteq X$ be compact in a metric space $X$. If $E \subseteq K$ is infinite, then $E^{\mathrm{acc}} \cap K \neq \emptyset$.*

*Proof.* Suppose $E^{\text{acc}} \cap K = \emptyset$. $\forall q \in K$, choose $r_q > 0$ such that $B_{r_q}(q) \cap (E \setminus \{q\}) = \emptyset$, so $\#(B_{r_q}(q) \cap E) \leq 1$. Since $\{B_{r_q}(q) : q \in K\}$ covers $K$, compactness gives $q_1, \ldots, q_n \in K$ with $K \subseteq \bigcup_{i=1}^{n} B_{r_{q_i}}(q_i)$.

As $E \subseteq K$, $E \subseteq \bigcup_{i=1}^{n} B_{r_{q_i}}(q_i)$, finite union of sets meeting $E$ in at most one point. Thus $E$ is finite, contradiction.

*Structural Commentary*: Compactness makes the local "at most one point" obstruction finite.
*Key Insight*: Infinite subsets of compact sets must accumulate somewhere inside the compact set. □

**Definition 4.1.11.** *Sequential Compactness*
A metric space is *sequentially compact* if every sequence has a convergent subsequence.

**Theorem 4.1.12.** *Compactness Produces Convergent Subsequences* *If $(p_n)$ is a sequence in a compact metric space $X$, then $p_{n_i} \to p$ for some subsequence and some $p \in X$.*

*Proof.* Let $E = \{p_n : n \in \mathbb{N}\}$. If $E$ is finite, choose a value $p$ attained infinitely often; the corresponding constant subsequence converges. If $E$ is infinite, Theorem 4.1.10 gives a limit point $p \in X$ of $E$. By Theorem 3.1.14, choose $n_i > n_{i-1}$ with $d(p_{n_i}, p) < 1/i$. Then $p_{n_i} \to p$. *Structural Commentary*: Compactness turns arbitrary infinite behavior into an accumulating subsequence.
*Key Insight*: A compact sequence range either repeats or accumulates. □

**Definition 4.1.13.** *Cauchy Sequence*
A sequence $(p_n)$ in a metric space $X$ is *Cauchy* if $\forall \varepsilon > 0$ $\exists$ integer $N$ s.t. $n \geq N,\ m \geq N \implies d(p_n, p_m) < \varepsilon$.

**Definition 4.1.14.** *Diameter*
Let $(X, d)$ be a metric space, $\emptyset \neq E \subseteq X$. The *diameter* of $E$ is $\operatorname{diam} E := \sup_{\overline{\mathbb{R}}}\{d(p, q) : p, q \in E\} \in [0, +\infty]$.

**Proposition 4.1.15.** *Basic Diameter Properties*
*Let $\emptyset \neq E, F \subseteq X$.*

*(i) $E$ is bounded $\iff \operatorname{diam} E < \infty \iff \exists q \in X, \exists M > 0$ such that $E \subseteq B_M(q)$.*
*(ii) $\operatorname{diam} E = 0 \iff |E| = 1$.*
*(iii) $E \subseteq F \implies \operatorname{diam} E \leq \operatorname{diam} F$.*
*(iv) $E, F$ bounded $\implies E \cup F$ bounded.*

*Proof.*

(i) If $E \subseteq B_M(q)$, then $d(x, y) \leq d(x, q) + d(q, y) < 2M$ for $x, y \in E$, so $\operatorname{diam} E < \infty$.
If $\operatorname{diam} E < \infty$ and $q \in E$, then $E \subseteq B_{\operatorname{diam} E + 1}(q)$.
(ii) $\operatorname{diam} E = 0 \iff d(x, y) = 0$ for all $x, y \in E \iff x = y$ for all $x, y \in E$; since $E \neq \emptyset$, this is $|E| = 1$.
(iii) $\{d(x, y) : x, y \in E\} \subseteq \{d(x, y) : x, y \in F\}$, so monotonicity of sup gives $\operatorname{diam} E \leq \operatorname{diam} F$.
(iv) Choose $e \in E$, $f \in F$, and set $D := \operatorname{diam} E + d(e, f) + \operatorname{diam} F < \infty$. For $x, y \in E \cup F$, either $x, y$ lie in the same set, or after swapping, $x \in E$, $y \in F$, and $d(x, y) \leq d(x, e) + d(e, f) + d(f, y) \leq D$.
Thus $\operatorname{diam}(E \cup F) < \infty$, so $E \cup F$ is bounded by (i).

*Structural Commentary*: Diameter is a single-number form of boundedness for nonempty sets.
*Key Insight*: Cross-distances between two bounded sets are controlled by one bridge distance. □

**Proposition 4.1.16.** *Cauchy Sequences and Tail Diameters*
*For $(p_n)$ in a metric space $X$, set $E_N := \{p_j : j \geq N\}$. Then $(p_n)$ is Cauchy if and only if $\operatorname{diam} E_N \to 0$.*

*Proof.*

($\Rightarrow$) Given $\varepsilon > 0$, choose $N_0$ so that $m, n \geq N_0$ implies $d(p_m, p_n) < \varepsilon/2$.
If $N \geq N_0$, then every pair in $E_N$ has distance $< \varepsilon/2$, so $\operatorname{diam} E_N < \varepsilon$.
($\Leftarrow$) Given $\varepsilon > 0$, choose $N_0$ with $\operatorname{diam} E_{N_0} < \varepsilon$.
If $m, n \geq N_0$, then $p_m, p_n \in E_{N_0}$, hence $d(p_m, p_n) \leq \operatorname{diam} E_{N_0} < \varepsilon$.

*Structural Commentary*: Cauchy behavior is shrinking diameter of sequence tails.
*Key Insight*: The limit is not named; only mutual closeness is measured. □

**Definition 4.1.17.** *Contracting Sequence of Sets*
Let $(X, d)$ be a metric space and let $(F_n)_{n \geq 1}$ satisfy $\emptyset \neq F_n \subseteq X$.

(i) $(F_n)$ is *descending* iff $\forall n \geq 1$, $F_{n+1} \subseteq F_n$.
(ii) $(F_n)$ is *contracting* iff it is descending and $\operatorname{diam} F_n \to 0$.




Figure 132: Tail diameters $\operatorname{diam} E_N$ shrink toward 0.

**Theorem 4.1.18.** *Nested Compact Sets with Shrinking Diameter*
*Let $X$ be a metric space.*

*(i) If $\emptyset \neq E \subseteq X$, then $\operatorname{diam} \overline{E} = \operatorname{diam} E$.*
*(ii) If $K_1 \supseteq K_2 \supseteq \cdots$ are nonempty compact subsets of $X$ and $\operatorname{diam} K_n \to 0$, then $|\bigcap_{n=1}^{\infty} K_n| = 1$.*

*Proof.*

(i) Since $E \subseteq \overline{E}$, $\operatorname{diam} E \leq \operatorname{diam} \overline{E}$. For $p, q \in \overline{E}$ and $\varepsilon > 0$, choose $p', q' \in E$ with $d(p, p'), d(q, q') < \varepsilon$; then $d(p, q) \leq \operatorname{diam} E + 2\varepsilon$. Hence $\operatorname{diam} \overline{E} \leq \operatorname{diam} E$.
(ii) Set $I := \bigcap_{n \geq 1} K_n$. Corollary 4.1.9 gives $I \neq \emptyset$. If $x, y \in I$, then $d(x, y) \leq \operatorname{diam} K_n$ for all $n$, so $d(x, y) = 0$ as $\operatorname{diam} K_n \to 0$. Thus $x = y$, hence $|I| = 1$.

*Structural Commentary*: Closure preserves diam; $K_n \downarrow$ with $\operatorname{diam} K_n \to 0$ forces a singleton.
*Key Insight*: Compactness gives existence; shrinking diameter gives uniqueness. □

**Theorem 4.1.19.** *Cauchy Criterion and Completeness Results*

*(i) In any metric space, every convergent sequence is Cauchy.*
*(ii) If $X$ is a compact metric space, then every Cauchy sequence in $X$ converges to a point of $X$.*

*Proof.*

(i) If $p_n \to p$, choose $N$ with $d(p_n, p) < \varepsilon/2$ for $n \geq N$. For $m, n \geq N$, $d(p_n, p_m) < \varepsilon$.
(ii) Let $E_N := \{p_j : j \geq N\}$ and $F_N := \overline{E_N}$. Then $\operatorname{diam} F_N \to 0$, the $F_N$ are compact and nested, and Theorem 4.1.18 gives $\bigcap_N F_N = \{p\}$. If $\operatorname{diam} F_N < \varepsilon$, then $p, p_n \in F_N$ for $n \geq N$, so $d(p_n, p) < \varepsilon$.

*Structural Commentary*: Cauchy controls $\operatorname{diam} E_N$; compactness turns $\bigcap_N F_N$ into the limit.
*Key Insight*: Compactness turns mutual tail closeness into an internal limit. □

**Lemma 4.1.20.** *A Cauchy Sequence with a Convergent Subsequence Converges*
*Let $(p_n)$ be a Cauchy sequence in a metric space $X$. If some subsequence $(p_{n_j})$ converges to $p \in X$, then $p_n \to p$.*

*Proof.* Given $\varepsilon > 0$, choose $N_0$ so that $m, n \geq N_0$ implies $d(p_m, p_n) < \varepsilon/2$.
Choose $j$ with $n_j \geq N_0$ and $d(p_{n_j}, p) < \varepsilon/2$. Then for every $n \geq N_0$, $d(p_n, p) \leq d(p_n, p_{n_j}) + d(p_{n_j}, p) < \varepsilon$.
*Structural Commentary*: The Cauchy tail follows any convergent subsequence to the same limit.
*Key Insight*: Mutual closeness plus one convergent thread forces the whole tail to converge. □

**Proposition 4.1.21.** *Boundedness and Distances of Cauchy Sequences*
*Let $(p_n), (q_n)$ be Cauchy sequences in a metric space $X$.*

*(i) Cauchy boundedness: the range of $n \mapsto p_n$, $\operatorname{ran}(p_n) := \{p_n : n \in \mathbb{N}\}$ is bounded.*
*(ii) distance convergence: the sequence $n \mapsto d(p_n, q_n)$ converges in $\mathbb{R}$.*
*(iii) nonconverse: there exists a bounded sequence in $\mathbb{R}$ that is not Cauchy.*

*Proof.*

(i) Choose $N$ with $m, n \geq N \Longrightarrow d(p_m, p_n) < 1$.
Then $\{p_n : n \geq N\} \subseteq B_1(p_N)$, and $\{p_1, \ldots, p_{N-1}\}$ is finite, so $\operatorname{ran}(p_n)$ is bounded.
(ii) For $m, n \in \mathbb{N}$, $|d(p_n, q_n) - d(p_m, q_m)| \leq d(p_n, p_m) + d(q_n, q_m)$. Hence $(d(p_n, q_n))$ Cauchy in $\mathbb{R}$, converges.
(iii) Let $a_n := (-1)^n$. Then $|a_n| \leq 1$ for all $n$, but with $\varepsilon = 1$, for every $N$ there exist $m, n \geq N$ with $|a_m - a_n| = 2$; hence $(a_n)$ is bounded but not Cauchy.

*Structural Commentary*: Cauchy tails are bounded; Cauchy-pair distances converge.
*Key Insight*: The four-point triangle estimate controls distance sequences. □

**Definition 4.1.22.** *Complete Metric Space*
A metric space $X$ is *complete* if every Cauchy sequence in $X$ converges to a point of $X$.
Compact metric spaces are complete by Theorem 4.1.19; $\mathbb{R}^k$ is complete by Corollary 4.2.8.

**Proposition 4.1.23.** *Completeness and Closed Subspaces*
*Let $(X,d)$ be a metric space, $E \subseteq X$, and $d_E := d|_{E\times E}$.*

*(i) If $(E,d_E)$ is complete, then $E$ is closed in $X$, equivalently $E = \overline{E}^X$.*
*(ii) If $(X,d)$ is complete, then $(E,d_E)$ is complete if and only if $E$ is closed in $X$, equivalently $E = \overline{E}^X$.*

*Proof.*

(i) Assume $(E,d_E)$ complete. Let $p \in \overline{E}^X$. Choose $(p_n) \subseteq E$ with $d(p_n,p) < 1/n$.
Then $p_n \to p$ in $X$, hence $(p_n)$ is Cauchy in $(E,d_E)$. Thus $p_n \to q \in E$ in $(E,d_E)$, hence also in $X$.
Uniqueness in $X$ gives $p = q \in E$. Therefore $\overline{E}^X \subseteq E$, so $E = \overline{E}^X$.
(ii) Assume $(X,d)$ complete. By (i), $(E,d_E)$ complete $\Rightarrow E = \overline{E}^X$.
Conversely, if $E = \overline{E}^X$ and $(p_n) \subseteq E$ is Cauchy in $(E,d_E)$, then $(p_n)$ is Cauchy in $X$, so $p_n \to p \in X$.
Since $p \in \overline{E}^X = E$, every Cauchy sequence in $(E,d_E)$ converges in $(E,d_E)$.

*Structural Commentary*: Completeness gives the limit in $X$; closedness keeps it in $E$.
*Key Insight*: In a complete ambient space, $E$ is complete exactly when $\overline{E} = E$. □

**Theorem 4.1.24.** *Cantor Intersection Theorem*
*Let $(X,d)$ be a metric space. Then $X$ is complete iff every sequence $(F_n)_{n\geq 1}$ with $\emptyset \neq F_n \subseteq X$ closed, $F_{n+1} \subseteq F_n$, and $\operatorname{diam} F_n \to 0$ satisfies $\bigcap_{n=1}^{\infty} F_n = \{x\}$ for some $x \in X$.*

*Proof.*

($\Rightarrow$) Assume $X$ is complete, and let $(F_n)$ satisfy the stated hypotheses. Choose $x_n \in F_n$. For $m,n \geq N$, descendingness gives $x_m, x_n \in F_N$, so $d(x_m,x_n) \leq \operatorname{diam} F_N$; hence $(x_n)$ is Cauchy. Let $x_n \to x \in X$. For each fixed $N$, the tail $(x_n)_{n\geq N}$ lies in closed $F_N$, so $x \in F_N$. Thus $x \in \bigcap_n F_n$. If $y \in \bigcap_n F_n$, then $d(x,y) \leq \operatorname{diam} F_N$ for all $N$, so $x = y$.
($\Leftarrow$) Assume the stated intersection property. Let $(x_n)$ be Cauchy, set $E_N := \{x_n : n \geq N\}$, and set $F_N := \overline{E_N}$. Then $F_N$ is nonempty, closed, and descending. By Proposition 4.1.16 and Theorem 4.1.18(i), $\operatorname{diam} F_N = \operatorname{diam} E_N \to 0$, so $\bigcap_N F_N = \{x\}$ for some $x \in X$. Given $\varepsilon > 0$, choose $N$ with $\operatorname{diam} F_N < \varepsilon$. If $n \geq N$, then $x_n, x \in F_N$, so $d(x_n,x) < \varepsilon$. Hence $x_n \to x$, and $X$ is complete.

*Structural Commentary*: Contracting closed nests have one point exactly in complete spaces.
*Key Insight*: Cauchy tails are the canonical contracting closed sets. □

We now turn from contracting closed families to the finite-resolution side of compact metric spaces.




Figure 133: Finite $\varepsilon$- and $\delta$-nets cover $E$, with $0 < \delta < \varepsilon$.

**Definition 4.1.25.** *Total Boundedness*
Let $(X,d)$ be a metric space and let $E \subseteq X$.

(i) $E$ is *totally bounded* iff $\forall \varepsilon > 0$, $\exists x_1,\ldots,x_N \in X$ with $E \subseteq \bigcup_{i=1}^N B_\varepsilon(x_i)$.
(ii) Such a finite family $\{B_\varepsilon(x_i)\}_{i=1}^N$ is a finite $\varepsilon$-*net* for $E$. If $E \neq \emptyset$, centers may equivalently be chosen in $E$.
(iii) The metric space $X$ is *totally bounded* iff $E = X$ satisfies (i).

**Theorem 4.1.26.** *Total Boundedness Implies Boundedness*
*Let $X$ be a metric space. If $X$ is totally bounded, then $X$ is bounded.*

*Proof.* If $X = \emptyset$, this is trivial. Otherwise cover $X$ by finitely many 1-balls $B_1(x_i)$ and fix $x_1$.
Then $d(x,x_1) < 1 + \max_i d(x_i,x_1)$ for every $x \in X$.
*Structural Commentary*: One finite cover at scale 1 gives one global bound.
*Key Insight*: Total boundedness is boundedness with finite control at every scale. □

**Example 4.1.27.** *Bounded Need Not Mean Totally Bounded*
In $\ell^2(\mathbb{F})$, let $B := \{\mathbf{x} \in \ell^2 : \|\mathbf{x}\|_2 \leq 1\}$. Then $B$ is bounded, since $B \subseteq B_2(0)$. It is not totally bounded: the standard unit vectors $e_n \in B$ satisfy $\|e_n - e_m\|_2 = \sqrt{2}$ for $n \neq m$. If $B \subseteq \bigcup_{i=1}^N B_{1/2}(x_i)$, then two distinct $e_n, e_m$ lie in the same $B_{1/2}(x_i)$, so $\|e_n - e_m\|_2 < 1$, contradicting $\|e_n - e_m\|_2 = \sqrt{2}$.

*Structural Commentary*: Bounded balls can contain infinite separated sets.
*Key Insight*: Total boundedness forbids an infinite separated family at a fixed scale.

**Definition 4.1.28.** *Lebesgue Number*
Let $(X, d)$ be a metric space and let $\mathcal{U}$ be an open cover of $X$.
A number $\delta > 0$ is a *Lebesgue number* for $\mathcal{U}$ iff $\forall x \in X$, $\exists U \in \mathcal{U}$ such that $B_\delta(x) \subseteq U$.

**Lemma 4.1.29.** *Lebesgue Covering Lemma*
*Let $(X, d)$ be a metric space.*

*(i) If $X$ is sequentially compact, then every open cover of $X$ has a Lebesgue number.*
*(ii) If $X$ is compact, then every open cover of $X$ has a Lebesgue number.*

*Proof.*

(i) Let $\mathcal{U} = \{U_\alpha\}_{\alpha \in A}$ be an open cover of sequentially compact $X$. If $\mathcal{U}$ has no Lebesgue number, then $\forall n \geq 1$, $\exists x_n \in X$ such that $B_{1/n}(x_n) \nsubseteq U_\alpha$ for all $\alpha \in A$. Choose a subsequence $x_{n_k} \to x \in X$. Since $\mathcal{U}$ covers $X$, choose $\alpha_0$ and $r > 0$ with $x \in U_{\alpha_0}$ and $B_r(x) \subseteq U_{\alpha_0}$. For large $k$, $d(x_{n_k}, x) < r/2$ and $1/n_k < r/2$; hence $y \in B_{1/n_k}(x_{n_k})$ gives $d(y, x) < r$, so $B_{1/n_k}(x_{n_k}) \subseteq U_{\alpha_0}$, a contradiction.
(ii) If $X$ is compact, then Theorem 4.1.12 gives sequential compactness.
(i) gives a Lebesgue number for every open cover of $X$.

*Structural Commentary*: Convergent subsequences force uniform cover radii.
*Key Insight*: Small balls near a convergent subsequence eventually stay in one cover set. □

**Theorem 4.1.30.** *Compactness as Complete Finite Resolution*
*Let $(X, d)$ be a metric space. The following are equivalent:*

*(i) $X$ is compact;*
*(ii) $X$ is complete and totally bounded;*
*(iii) $X$ is sequentially compact.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* Let $(p_k)$ be Cauchy in $X$. By Theorem 4.1.19, $(p_k)$ converges, so $X$ is complete. For $\varepsilon > 0$, compactness applied to the cover $\{B_\varepsilon(x) : x \in X\}$ gives a finite $\varepsilon$-net, so $X$ is totally bounded.
*(ii)* $\Rightarrow$ *(iii)* Let $(x_n)$ be a sequence in $X$. Using finite $1/j$-ball covers, recursively choose nested infinite subsequences so that the $j$th subsequence lies in some ball $B_{1/j}(a_j)$. Let $y_j$ be the $j$th term of the $j$th subsequence. If $k, \ell \geq j$, nestedness gives $y_k, y_\ell \in B_{1/j}(a_j)$, hence $d(y_k, y_\ell) < 2/j$. Thus $(y_j)$ is Cauchy; completeness gives $y_j \to y \in X$, so $(x_n)$ has a convergent subsequence.
*(iii)* $\Rightarrow$ *(ii)* If $X$ is not totally bounded, then for some $\varepsilon > 0$ no finite $\varepsilon$-balls cover $X$. Choose $x_1 \in X$ and inductively $x_{n+1} \notin \bigcup_{j=1}^{n} B_\varepsilon(x_j)$. Then $d(x_n, x_j) \geq \varepsilon$ for $j < n$, so no subsequence is Cauchy or convergent, contradicting sequential compactness. If $(p_n)$ is Cauchy, sequential compactness gives a convergent subsequence, and Lemma 4.1.20 gives convergence of $(p_n)$. Hence $X$ is complete.
*(ii)* $\Rightarrow$ *(i)* By *(ii)* $\Rightarrow$ *(iii)*, $X$ is sequentially compact. Let $(U_\alpha)_{\alpha \in A}$ be an open cover of $X$, and let $\delta > 0$ be a Lebesgue number from Lemma 4.1.29. Choose a finite $\delta/2$-net $\{z_1, \ldots, z_m\}$ for $X$. For each $i$, choose $\alpha_i$ with $B_\delta(z_i) \subseteq U_{\alpha_i}$. Since every $x \in X$ lies in some $B_{\delta/2}(z_i)$, the finite family $U_{\alpha_1}, \ldots, U_{\alpha_m}$ covers $X$.

*Structural Commentary*: Compactness is complete finite-scale control.
*Key Insight*: Total boundedness gives Cauchy subsequences; completeness gives limits. □




Figure 134: Equivalent compactness tests in a metric space $X$: finite subcovers for open covers, finite $\varepsilon$-nets with Cauchy limits in $X$, and convergent subsequences for every sequence.

**Proposition 4.1.31.** *Compact Metric Spaces Are Separable*
*Let $(X, d)$ be a compact metric space. Then $\exists$ countable $D \subseteq X$ such that $\overline{D}^X = X$.*

*Proof.* By Theorem 4.1.30, $X$ is totally bounded.
$\forall n \in \mathbb{N}$, choose finite $F_n \subseteq X$ such that $X \subseteq \bigcup_{a \in F_n} B_{1/n}(a)$, set $D := \bigcup_{n=1}^{\infty} F_n$. Then $D$ is countable.
Let $x \in X$ and $r > 0$. Choose $n \in \mathbb{N}$ with $1/n < r$.
Since $F_n$ is a $1/n$-net for $X$, $\exists a \in F_n$ such that $d(x, a) < 1/n < r$.
Thus $a \in B_r(x) \cap D$, so every $X$-open ball around $x$ meets $D$. Hence $x \in \overline{D}^X$, and therefore $\overline{D}^X = X$.
*Structural Commentary*: Compactness gives finite nets at every scale; their countable union is dense.
*Key Insight*: Total boundedness converts compactness into countably many finite approximations. □

## 4.2 Heine–Borel and Euclidean Compactness

**Theorem 4.2.1.** *Nested Closed Intervals Intersect*
*Let $(I_n)_{n\in\mathbb{N}}$ be nonempty compact intervals in $\mathbb{R}$, with $I_n = [a_n, b_n]$, such that $I_{n+1} \subseteq I_n$ $\forall n$. Then $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$.*

*Proof.* Set $A := \{a_n : n \in \mathbb{N}\}$. Fix $m \in \mathbb{N}$. If $n \geq m$, then $a_n \in I_n \subseteq I_m$, so $a_n \leq b_m$.
If $n < m$, then $b_m \in I_m \subseteq I_n$, so $a_n \leq b_m$. Hence $b_m$ is an upper bound for $A$.
Let $x := \sup A$. Since $a_m \in A$, $a_m \leq x$; since $b_m$ is an upper bound for $A$, $x \leq b_m$.
Hence $x \in I_m$ for every $m$, and therefore $x \in \bigcap_{n=1}^{\infty} I_n$.
*Structural Commentary*: Nested endpoints form a LUB trap.
*Key Insight*: $\sup_n a_n$ lies in every interval. □




Figure 135: For nested closed intervals $[a_n, b_n]$, supremum of left endpoints lies between every $a_n$ and $b_n$.

**Theorem 4.2.2.** *Nested k-Cells Intersect*
*Let $k \in \mathbb{N}$, and for each $n \in \mathbb{N}$ let $I_n = \prod_{j=1}^{k} [a_{n,j}, b_{n,j}] \subseteq \mathbb{R}^k$ be a nonempty $k$-cell.*
*Assume $I_{n+1} \subseteq I_n$ for every $n$. Then $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$.*

*Proof.* For each $j$, the $j$th coordinate projection of $I_{n+1} \subseteq I_n$ gives $[a_{n+1,j}, b_{n+1,j}] \subseteq [a_{n,j}, b_{n,j}]$.
By Theorem 4.2.1, choose $x_j^* \in \bigcap_{n=1}^{\infty} [a_{n,j}, b_{n,j}]$. Then $\mathbf{x}^* := (x_1^*, \ldots, x_k^*)$ satisfies $\mathbf{x}^* \in I_n$ $\forall n$, so $\mathbf{x}^* \in \bigcap_{n=1}^{\infty} I_n$.
*Structural Commentary*: Cells reduce coordinatewise to intervals.
*Key Insight*: Pick one persistent coordinate per axis. □




Figure 136: A 1-cell is a closed interval, a 2-cell is a coordinate rectangle, and a 3-cell is a coordinate box.

**Theorem 4.2.3.** *k-Cells Are Compact*
*Let $k \in \mathbb{N}$, and let $I = \prod_{j=1}^{k} [a_j, b_j] \subseteq \mathbb{R}^k$ be a nonempty $k$-cell. Then $I$ is compact.*

*Proof.* Set $\delta := \left(\sum_{j=1}^{k} (b_j - a_j)^2\right)^{1/2} = \operatorname{diam} I$. Suppose open cover $\mathcal{G} = \{G_\alpha\}_{\alpha \in A}$ of $I$ has no finite subcover.
Bisect each coordinate interval of $I$. If every resulting subcell had a finite $\mathcal{G}$-subcover, their union would finitely cover $I$, so choose a subcell $I_1$ with no finite $\mathcal{G}$-subcover. Repeating inside $I_{n-1}$ gives nested $k$-cells $I \supseteq I_1 \supseteq I_2 \supseteq \cdots$ such that $I_n$ has no finite $\mathcal{G}$-subcover and $\operatorname{diam} I_n \leq 2^{-n}\delta$.
By Theorem 4.2.2, choose $\mathbf{x}^* \in \bigcap_{n=1}^{\infty} I_n$. Since $\mathcal{G}$ covers $I$, choose $\alpha$ with $\mathbf{x}^* \in G_\alpha$. Openness gives $r > 0$ such that $B_r(\mathbf{x}^*) \subseteq G_\alpha$. Choose $n$ with $2^{-n}\delta < r$. Since $\mathbf{x}^* \in I_n$ and $\operatorname{diam} I_n < r$, one has $I_n \subseteq B_r(\mathbf{x}^*) \subseteq G_\alpha$,

contradicting the choice of $I_n$. Thus every open cover of $I$ has a finite subcover.
*Structural Commentary*: Bisection gives shrinking nested cells.
*Key Insight*: One open set eventually contains the trapped cell. □




Figure 137: Repeated bisection traps a point.

**Lemma 4.2.4.** *Bounded Euclidean Sets Fit in k-Cells*
*Let $k \in \mathbb{N}$ and let $E \subseteq \mathbb{R}^k$ be bounded. Then there exists a $k$-cell $I \subseteq \mathbb{R}^k$ such that $E \subseteq I$.*

*Proof.* If $E = \emptyset$, take $I = \prod_{j=1}^{k}[0,0]$. Assume $E \neq \emptyset$.
Choose $\mathbf{q} = (q_1, \ldots, q_k) \in \mathbb{R}^k$ and $M > 0$ such that $E \subseteq B_M(\mathbf{q})$. Set $I := \prod_{j=1}^{k}[q_j - M, q_j + M]$.
For $\mathbf{p} = (p_1, \ldots, p_k) \in E$ and $1 \leq j \leq k$, $|p_j - q_j| \leq \|\mathbf{p} - \mathbf{q}\|_2 < M$, hence $\mathbf{p} \in I$. Thus $E \subseteq I$.
*Structural Commentary*: One ball fits inside one coordinate box.
*Key Insight*: A radius gives coordinate bounds. □

**Theorem 4.2.5.** *Heine–Borel in $\mathbb{R}^k$*
*Let $k \in \mathbb{N}$ and let $E \subseteq \mathbb{R}^k$. The following are equivalent:*

*(i) $E$ is closed and bounded;*
*(ii) $E$ is compact;*
*(iii) every infinite subset of $E$ has a limit point in $E$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) Since $E$ is bounded, Lemma 4.2.4 gives a $k$-cell $I$ with $E \subseteq I$. By Theorem 4.2.3, $I$ is compact. Since $E$ is closed in $\mathbb{R}^k$, $E$ is closed in $I$, so Theorem 4.1.6 implies that $E$ is compact.
(*ii*) $\Rightarrow$ (*iii*) Theorem 4.1.10 applies directly to the compact set $E$.
(*iii*) $\Rightarrow$ (*i*) Suppose first that $E$ is unbounded. Choose $\mathbf{x}_n \in E$ with $\|\mathbf{x}_n\|_2 > n$ and set $S := \{\mathbf{x}_n : n \in \mathbb{N}\}$. For any $\mathbf{y} \in \mathbb{R}^k$, the set $S \cap B_1(\mathbf{y})$ is finite because $\mathbf{x}_n \in B_1(\mathbf{y})$ implies $n < \|\mathbf{y}\|_2 + 1$. Hence $S$ has no limit point, contradicting (iii). Thus $E$ is bounded. Suppose next that $E$ is not closed. Choose $\mathbf{x}_0 \in \overline{E} \setminus E$ and distinct $\mathbf{x}_n \in E$ with $0 < \|\mathbf{x}_n - \mathbf{x}_0\|_2 < 1/n$. Then $S := \{\mathbf{x}_n : n \in \mathbb{N}\}$ is infinite, and every limit point of $S$ equals $\mathbf{x}_0 \notin E$, contradicting (iii). Thus $E$ is closed.

*Structural Commentary*: Compactness equals closed boundedness in $\mathbb{R}^k$.
*Key Insight*: Cells convert bounded geometry into compactness. □

**Theorem 4.2.6.** *Weierstrass Limit-Point Theorem*
*Let $k \in \mathbb{N}$ and let $E \subseteq \mathbb{R}^k$ be bounded and infinite. Then $E$ has a limit point in $\mathbb{R}^k$.*

*Proof.* By Lemma 4.2.4, choose a $k$-cell $I$ with $E \subseteq I$. By Theorem 4.2.3, $I$ is compact.
Since $E$ is an infinite subset of $I$, Theorem 4.1.10 gives a limit point of $E$ in $I \subseteq \mathbb{R}^k$.
*Structural Commentary*: Bounded sets sit in compact cells.
*Key Insight*: Infinite compact subsets accumulate. □

**Corollary 4.2.7.** *Bounded Euclidean Sequences Have Convergent Subsequences*
*If $(\mathbf{x}_n)$ is bounded in $\mathbb{R}^k$, then it has a convergent subsequence.*

*Proof.* The range of $(\mathbf{x}_n)$ has compact closure by Theorem 4.2.5; apply Theorem 4.1.12 in that closure.
*Structural Commentary*: Heine–Borel makes the range closure compact.
*Key Insight*: Bounded Euclidean sequences live inside compact closed sets. □

**Corollary 4.2.8.** *Euclidean Cauchy Convergence*
*In $\mathbb{R}^k$, every Cauchy sequence converges.*

*Proof.* For Cauchy $(\mathbf{x}_n) \subseteq \mathbb{R}^k$, choose $N$ with $\operatorname{diam}\{\mathbf{x}_j : j \geq N\} < 1$.
Its range is bounded, hence has compact closure by Theorem 4.2.5; apply Theorem 4.1.19(ii).
*Structural Commentary*: A Cauchy tail has bounded compact closure.
*Key Insight*: In $\mathbb{R}^k$, Cauchy convergence follows from Heine–Borel. □

**Corollary 4.2.9.** *Euclidean Compactness Tests*
*Let $k \in \mathbb{N}$ and let $E \subseteq \mathbb{R}^k$, with the subspace metric. The following are equivalent:*

*(i) $E$ is closed and bounded;*
*(ii) $E$ is compact;*
*(iii) every sequence in $E$ has a subsequence converging to a point of $E$.*

*Proof.*

(i) $\Rightarrow$ (ii) This is Theorem 4.2.5.
(ii) $\Rightarrow$ (iii) Let $(x_n)_{n\in\mathbb{N}}$ be a sequence in $E$.
Since $E$ is compact, Theorem 4.1.30 gives indices $n_1 < n_2 < \cdots$ and $x \in E$ such that $x_{n_j} \to x$.
(iii) $\Rightarrow$ (i) By (iii), $E$ is sequentially compact. Theorem 4.1.30, applied to $X = E$, gives that $E$ is compact.
Then Theorem 4.2.5 implies that $E$ is closed and bounded.

*Structural Commentary*: Euclidean compactness has three tests.
*Key Insight*: Heine–Borel links geometry to sequences. □

**Corollary 4.2.10.** *Heine–Borel on the Real Line*
*Let $E \subseteq \mathbb{R}$, with the usual metric. The following are equivalent:*

*(i) $E$ is closed and bounded;*
*(ii) $E$ is compact;*
*(iii) every sequence in $E$ has a subsequence converging to a point of $E$.*

*Proof.*

(i) $\Rightarrow$ (ii) By (i), $E$ is closed, bounded as a subset of $\mathbb{R}^1$. Theorem 4.2.5 with $k = 1$ gives that $E$ is compact.
(ii) $\Rightarrow$ (iii) Let $(x_n)_{n\in\mathbb{N}}$ be a sequence in $E$.
By compactness of $E$, Theorem 4.1.30 gives indices $n_1 < n_2 < \cdots$ and $x \in E$ such that $x_{n_j} \to x$.
(iii) $\Rightarrow$ (i) Let $F \subseteq E$ be infinite, and choose distinct $x_n \in F$. By (iii), some subsequence satisfies $x_{n_j} \to x \in E$.
Since the $x_n$ are distinct, $x$ is a limit point of $F$. Thus every infinite subset of $E$ has a limit point in $E$, so Theorem 4.2.5 with $k = 1$ gives that $E$ is closed and bounded.

*Structural Commentary*: On $\mathbb{R}$, compactness is sequential.
*Key Insight*: Closed bounded sets trap subsequences. □

**Corollary 4.2.11.** *Nested Closed Bounded Real Sets Intersect*
*Let $(F_n)_{n\in\mathbb{N}}$ be nonempty closed subsets of $\mathbb{R}$ with $F_{n+1} \subseteq F_n$ for all $n$. If $F_1$ is bounded, then $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$.*

*Proof.* By Corollary 4.2.10, $F_1$ is compact. For each $n$, $F_n$ is a nonempty closed subset of the compact set $F_1$, so Theorem 4.1.6 gives that $F_n$ is compact. Since $F_{n+1} \subseteq F_n$, Corollary 4.1.9 gives $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$.
*Structural Commentary*: Nested compact sets intersect.
*Key Insight*: Heine–Borel supplies compactness. □

## 4.3 Baire Category

**Definition 4.3.1.** *$F_\sigma$, $G_\delta$, and Nowhere-Dense Sets*
For subsets of a metric space $X$:

(i) $A$ is an *$F_\sigma$ set* if $A = \bigcup_{n=1}^{\infty} F_n$ for closed sets $F_n$.
(ii) $B$ is a *$G_\delta$ set* if $B = \bigcap_{n=1}^{\infty} G_n$ for open sets $G_n$.
(iii) $E$ is *nowhere dense* if $(\overline{E})^\circ = \emptyset$; equivalently, no nonempty open set is contained in $\overline{E}$.

For $X = \mathbb{R}$, this says exactly that $\overline{E}$ contains no nonempty open interval.

**Proposition 4.3.2.** *Open Real Sets Are $F_\sigma$*
*For subsets of $\mathbb{R}$:*

*(i) If $G \subseteq \mathbb{R}$ is open, then $G$ is $F_\sigma$.*
*(ii) If $A \subseteq \mathbb{R}$ is closed, then $A$ is $G_\delta$.*

*Proof.*

(i) If $G = \mathbb{R}$, then $G = \bigcup_{m\geq 1}[-m, m]$.
Otherwise set $C := \mathbb{R} \setminus G$ and $F_m := [-m, m] \cap \{x \in \mathbb{R} : \operatorname{dist}(x, C) \geq 1/m\}$ for $m \geq 1$.
The map $x \mapsto \operatorname{dist}(x, C)$ is continuous since $|\operatorname{dist}(x, C) - \operatorname{dist}(y, C)| \leq |x - y|$; hence each $F_m$ is closed.
Moreover $F_m \subseteq G$. For $x \in G$, $r > 0$ with $(x - r, x + r) \subseteq G$, $m \geq |x|$; $1/m \leq r$ give $x \in F_m$. QED.

(ii) If $A$ is closed, then $A^c = \bigcup_{m=1}^{\infty} F_m$ with $F_m$ closed by (i), hence $A = \bigcap_{m=1}^{\infty} F_m^c$ is $G_\delta$.

*Structural Commentary*: Positive distance from the complement gives closed exhaustion.
*Key Insight*: Distance to the complement converts local openness into a countable closed exhaustion. □

**Definition 4.3.3.** *First Category*
Let $X$ be a metric space and let $E \subseteq X$. The set $E$ is *of first category*, *meager*, or *thin* in $X$ if there are nowhere-dense sets $N_n \subseteq X$ such that $E \subseteq \bigcup_{n=1}^{\infty} N_n$.
*Structural Commentary*: Meagerness is relative to the ambient space $X$.
*Key Insight*: The Baire Category Theorem says a nonempty complete metric space is not meager in itself.

**Theorem 4.3.4.** *Dense Open Intersections in $\mathbb{R}$*
*If $G_1, G_2, G_3, \ldots$ are dense open subsets of $\mathbb{R}$, then $\bigcap_{n=1}^{\infty} G_n \neq \emptyset$.*

*Proof.* We construct nested closed intervals. Since $G_1$ is dense and open, $G_1 \cap (0, 1)$ contains a nonempty open interval; choose a closed interval $I_1 \subseteq G_1 \cap (0, 1)$ with nonempty interior.
Given $I_n = [a_n, b_n]$, density of $G_{n+1}$ gives a point of $G_{n+1}$ in $(a_n, b_n)$. Since $G_{n+1}$ is open, choose a closed interval $I_{n+1} \subseteq G_{n+1} \cap (a_n, b_n)$ with nonempty interior and length $< 1/(n + 1)$.
Then $I_1 \supseteq I_2 \supseteq I_3 \supseteq \cdots$, so Theorem 4.2.1 gives $x \in \bigcap_n I_n$. Since $I_n \subseteq G_n$ for every $n$, this $x$ lies in $\bigcap_n G_n$.
*Structural Commentary*: Dense openness builds nested intervals; completeness keeps a limit point.
*Key Insight*: A countable dense-open intersection cannot be destroyed one interval at a time. □

**Theorem 4.3.5.** *Dense Open Intersections in Complete Metric Spaces*
*Let $(X, d)$ be complete, let $O_1, O_2, \ldots$ be dense open subsets of $X$, and let $U \subseteq X$ be nonempty and open. Then $U \cap \bigcap_{n=1}^{\infty} O_n \neq \emptyset$. Consequently, $\bigcap_{n=1}^{\infty} O_n$ is dense in $X$.*

*Proof.* Choose $x_1 \in U \cap O_1$ and $0 < r_1 < 1$ with $C_1 := \overline{B}_{r_1}(x_1) \subseteq U \cap O_1$.
Inductively, if $C_n = \overline{B}_{r_n}(x_n) \subseteq U \cap O_1 \cap \cdots \cap O_n$ and $0 < r_n < 2^{-n}$, density and openness of $O_{n+1}$ give $x_{n+1} \in B_{r_n}(x_n) \cap O_{n+1}$ and $0 < r_{n+1} < 2^{-(n+1)}$ such that $C_{n+1} := \overline{B}_{r_{n+1}}(x_{n+1}) \subseteq B_{r_n}(x_n) \cap O_{n+1}$.
Then $C_{n+1} \subseteq C_n$. If $m > n$, then $x_m \in B_{r_n}(x_n)$, so $d(x_m, x_n) < 2^{-n}$; hence $(x_n)$ is Cauchy. Let $x_n \to x$.
Since each $C_n$ is closed and contains all sufficiently late $x_m$, we have $x \in C_n$ for every $n$. Thus $x \in U \cap \bigcap_n O_n$.
Since every nonempty open $U$ meets $\bigcap_n O_n$, the intersection is dense.
*Structural Commentary*: The interval proof becomes a nested closed-ball proof.
*Key Insight*: Dense open sets choose the next ball; completeness supplies the limiting point. □




Figure 138: Nested closed balls shrink to $p$.

**Definition 4.3.6.** *Residual, Thick, and Generic Sets*
Let $X$ be a metric space.

(i) A set $G \subseteq X$ is *residual*, or *thick*, in $X$ if there are dense open sets $O_n \subseteq X$ such that $\bigcap_{n=1}^{\infty} O_n \subseteq G$.
(ii) A property $P$ is *generic* in $X$ if $\{x \in X : P(x)\}$ is residual in $X$.

If $X$ is complete, then every residual subset of $X$ is dense by Theorem 4.3.5.
*Key Insight*: Meager sets are topologically small; residual sets contain a countable dense-open core.

**Theorem 4.3.7.** *Baire Category Theorem for Complete Metric Spaces*
*A nonempty complete metric space is not the union of a countable collection of nowhere-dense sets.*

*Proof.* Suppose $X = \bigcup_n E_n$ with each $E_n$ nowhere dense, and set $O_n := (\overline{E_n})^c$.
Then each $O_n$ is dense open: it is open, and every nonempty open set meets it because no such set lies in $\overline{E_n}$.
By Theorem 4.3.5, choose $x \in \bigcap_n O_n$. Then $x \notin E_n$ for all $n$, contradicting $X = \bigcup_n E_n$.
*Structural Commentary*: Countably many nowhere-dense sets have dense open complements.
*Key Insight*: Completeness forbids countable assembly from thin pieces. □

**Theorem 4.3.8.** *Baire's Theorem for $\mathbb{R}$*
*The real line $\mathbb{R}$ cannot be written as a countable union of nowhere-dense sets.*

*Proof.* Assume $\mathbb{R} = \bigcup_n E_n$ with each $E_n$ nowhere dense, and set $U_n := (\overline{E_n})^c$.
Each $U_n$ is dense open: it is open, and every nonempty interval meets it because no such interval lies in $\overline{E_n}$.
By Theorem 4.3.4, choose $x \in \bigcap_n U_n$. Then $x \notin E_n$ for all $n$, contradicting $\mathbb{R} = \bigcup_n E_n$.
*Structural Commentary*: Nowhere-dense sets have dense open complements.
*Key Insight*: Countably many thin sets cannot exhaust the real line. □

**Corollary 4.3.9.** *Rationals and Irrationals by Category*
*The set $\mathbb{Q}$ is $F_\sigma$ but not $G_\delta$. The set $\mathbb{R} \setminus \mathbb{Q}$ is $G_\delta$ but not $F_\sigma$.*

*Proof.* Write $\mathbb{Q} = \{q_1, q_2, \ldots\}$. Since singletons are closed, $\mathbb{Q} = \bigcup_n \{q_n\}$ is $F_\sigma$, and $\mathbb{R} \setminus \mathbb{Q} = \bigcap_n (\{q_n\})^c$ is $G_\delta$.
If $\mathbb{Q} = \bigcap_n G_n$ with each $G_n$ open, then each $G_n$ is dense because it contains the dense set $\mathbb{Q}$. Set $H_n := G_n \setminus \{q_n\}$.
Each $H_n$ is dense open, so Theorem 4.3.4 gives a point of $\bigcap_n H_n$. But $\bigcap_n H_n = \emptyset$, since $\bigcap_n G_n = \mathbb{Q}$ and $q_n$ is removed at stage $n$. Hence $\mathbb{Q}$ is not $G_\delta$. Therefore $\mathbb{R} \setminus \mathbb{Q}$ is not $F_\sigma$, or else its complement $\mathbb{Q}$ would be $G_\delta$.
*Structural Commentary*: Category separates the familiar countable dense set from its dense complement.
*Key Insight*: Dense is not the same as large in the Baire-category sense. □

**Corollary 4.3.10.** *Closed Cover Interior Test*
*Let $X$ be a complete metric space and let $(F_n)_{n\in\mathbb{N}}$ be closed subsets of $X$.*

*(i) If $\emptyset \neq U \subseteq X$ is open and $U \subseteq \bigcup_{n=1}^{\infty} F_n$, then $F_N^\circ \neq \emptyset$ for some $N \in \mathbb{N}$.*
*(ii) If $X \neq \emptyset$ and $F_n^\circ = \emptyset$ for every $n \in \mathbb{N}$, then $X \neq \bigcup_{n=1}^{\infty} F_n$.*

*Proof.*

(*i*) Assume $\emptyset \neq U \subseteq X$ is open and $U \subseteq \bigcup_{n=1}^{\infty} F_n$. Suppose $F_n^\circ = \emptyset$ for every $n$ and set $O_n := X \setminus F_n$. Since $F_n$ is closed, $O_n$ is open. If $V \subseteq X$ nonempty and open, then $V \nsubseteq F_n$, so $V \cap O_n \neq \emptyset$; hence $O_n$ dense. Theorem 4.3.5 gives $x \in U \cap \bigcap_{n=1}^{\infty} O_n$. Then $x \in U \subseteq \bigcup_{n=1}^{\infty} F_n$ and $x \notin F_n$ $\forall n$, contradiction. QED.
(*ii*) Assume $X \neq \emptyset$ and $F_n^\circ = \emptyset$ for every $n$. If $X = \bigcup_{n=1}^{\infty} F_n$, then (i) with $U = X$ gives $F_N^\circ \neq \emptyset$ for some $N$, a contradiction. Hence $X \neq \bigcup_{n=1}^{\infty} F_n$.

*Structural Commentary*: Empty interiors give dense open complements.
*Key Insight*: Baire forces one complement-avoiding point. □

**Theorem 4.3.11.** *Pointwise Bounded Families Are Locally Uniformly Bounded*
*Let $(X, d)$ be a nonempty complete metric space, and let $\mathcal{F} \subseteq C(X, \mathbb{R})$.*
*Assume that for every $x \in X$ there is $M_x < \infty$ such that $|f(x)| \leq M_x$ for every $f \in \mathcal{F}$.*
*Then there exist a nonempty open set $O \subseteq X$ and $M < \infty$ such that $|f(x)| \leq M$ for all $x \in O$ and $f \in \mathcal{F}$.*

*Proof.* For $n \in \mathbb{N}$, set $E_n := \{x \in X : \forall f \in \mathcal{F},\ |f(x)| \leq n\}$.
As $E_n = \bigcap_{f\in\mathcal{F}} \{x \in X : |f(x)| \leq n\}$ and each $f$ is continuous, $E_n$ is closed. The hypothesis gives $X = \bigcup_{n=1}^{\infty} E_n$.
By Corollary 4.3.10, $E_N^\circ \neq \emptyset$ for some $N \in \mathbb{N}$. Take $O := E_N^\circ$, $M := N$. Then $x \in O$ and $f \in \mathcal{F}$ imply $|f(x)| \leq M$.
*Structural Commentary*: Pointwise bounds form a closed cover.
*Key Insight*: Completeness turns pointwise control into uniform control somewhere. □

**Theorem 4.3.12.** *Dense Continuity Set for Pointwise Limits*
*Let $(X, d)$ be a complete metric space, let $f : X \to \mathbb{R}$, let $f_n : X \to \mathbb{R}$ be continuous for every $n \in \mathbb{N}$, and assume $f_n(x) \to f(x)$ for every $x \in X$. Then there exists $D \subseteq X$ such that:*

*(i) $D$ is dense in $X$;*
*(ii) $(f_n)$ is equicontinuous at every $x_0 \in D$;*
*(iii) $f$ is continuous at every $x_0 \in D$.*

*Proof.* For $m, n \in \mathbb{N}$, set $E_{m,n} := \{x \in X : \forall j, k \geq n,\ |f_j(x) - f_k(x)| \leq 1/m\}$.
Since $E_{m,n} = \bigcap_{j,k \geq n} \{x \in X : |f_j(x) - f_k(x)| \leq 1/m\}$, each $E_{m,n}$ is closed.
or fixed $m$, pointwise convergence makes $(f_n(x))$ Cauchy for every $x \in X$, hence $X = \bigcup_{n=1}^{\infty} E_{m,n}$.
For closed $E \subseteq X$, $\mathrm{bd}(E) = E \setminus E^\circ$ is closed and nowhere dense: if nonempty open $V \subseteq \mathrm{bd}(E)$, then $V \subseteq E^\circ$ and $V \subseteq E \setminus E^\circ$, a contradiction. Thus each $X \setminus \mathrm{bd}(E_{m,n})$ is dense open. Set $D := X \setminus \bigcup_{m,n \in \mathbb{N}} \mathrm{bd}(E_{m,n})$.

(*i*) Since $\mathbb{N}^2$ is countable, $D = \bigcap_{m,n \in \mathbb{N}} (X \setminus \mathrm{bd}(E_{m,n}))$ is a countable intersection of dense open sets. By Theorem 4.3.5, $D$ is dense in $X$.

(*ii*) Fix $x_0 \in D$ and $\varepsilon > 0$. Choose $m \in \mathbb{N}$ with $3/m < \varepsilon$. Since $(f_n(x_0))$ is Cauchy, choose $N$ such that $x_0 \in E_{m,N}$. Since $x_0 \notin \mathrm{bd}(E_{m,N})$, actually $x_0 \in E_{m,N}^\circ$; choose $r > 0$ with $B_r(x_0) \subseteq E_{m,N}$. By continuity of $f_1, \ldots, f_N$ at $x_0$, choose $0 < \delta < r$ such that $d(x, x_0) < \delta$ implies $|f_i(x) - f_i(x_0)| < 1/m$ for $1 \leq i \leq N$. If $d(x, x_0) < \delta$ and $j > N$, then $x, x_0 \in E_{m,N}$, so $|f_j(x) - f_N(x)| \leq 1/m$ and $|f_N(x_0) - f_j(x_0)| \leq 1/m$. Hence $|f_j(x) - f_j(x_0)| \leq |f_j(x) - f_N(x)| + |f_N(x) - f_N(x_0)| + |f_N(x_0) - f_j(x_0)| < 3/m < \varepsilon$. For $1 \leq j \leq N$, the choice of $\delta$ gives $|f_j(x) - f_j(x_0)| < 1/m < \varepsilon$. Thus $(f_n)$ is equicontinuous at $x_0$.

(*iii*) Fix $x_0 \in D$ and $\varepsilon > 0$, and choose $m, N, \delta$ as in (ii). If $d(x, x_0) < \delta$, the estimate in (ii) gives $|f_j(x) - f_j(x_0)| < 3/m < \varepsilon$ for all $j > N$. Letting $j \to \infty$ gives $|f(x) - f(x_0)| \leq 3/m < \varepsilon$. QED.

*Structural Commentary*: Baire avoids all layer boundaries.
*Key Insight*: Interior tail-Cauchy control gives equicontinuity. □

**Perspective 4.3.13.** *Category in Function Spaces*
Baire category is a theorem about complete metric spaces, not just $\mathbb{R}$.
It will be reused for complete function spaces under the supremum metric.

**Method 4.3.14.** *Baire-Category Proof Patterns*

(i) *Dense-open intersection*: build nested closed intervals $I_n \subseteq G_n$ and apply Theorem 4.2.1.
(ii) *Complete metric version*: build nested closed balls $C_{n+1} \subseteq C_n$ with $r_n \to 0$ and use completeness.
(iii) *Nowhere dense*: replace $E$ by the dense open set $(\overline{E})^c$.
(iv) *First category*: show the exceptional set is contained in $\bigcup_n E_n$ with each $E_n$ nowhere dense.
(v) *Residual/generic*: show the desired property contains $\bigcap_n O_n$ with each $O_n$ dense open.
(vi) *$F_\sigma$/$G_\delta$ complements*: pass between countable unions and intersections by De Morgan's laws.

## 4.4 Perfect Sets and the Cantor Set

**Theorem 4.4.1.** *Perfect Sets Are Uncountable*
*Every nonempty perfect subset of $\mathbb{R}^k$ is uncountable.*

*Proof.* Assume $P = \{\mathbf{x}_1, \mathbf{x}_2, \ldots\}$. Choose a neighborhood $V_1$ with $V_1 \cap P \neq \emptyset$.
Inductively, since $P \cap V_n$ contains infinitely many points by Theorem 3.1.14, choose $\mathbf{y}_n \in P \cap V_n$ with $\mathbf{y}_n \neq \mathbf{x}_n$ and then a neighborhood $V_{n+1}$ of $\mathbf{y}_n$ such that $\overline{V}_{n+1} \subseteq V_n$ and $\mathbf{x}_n \notin \overline{V}_{n+1}$.
Put $K_n := \overline{V}_n \cap P$. Then $K_n$ is nonempty, compact, and $K_{n+1} \subseteq K_n$. By Corollary 4.1.9, choose $\mathbf{z} \in \bigcap_{n=1}^{\infty} K_n$.
For each $n$, $\mathbf{z} \in K_{n+1}$ but $\mathbf{x}_n \notin K_{n+1}$, so $\mathbf{z} \neq \mathbf{x}_n$ for all $n$, contradicting the enumeration.
*Structural Commentary*: Nested compact sets dodge the list.
*Key Insight*: Perfectness gives new points; compactness keeps one. □




Figure 139: Choose $y_2 \in P \cap I_1$ with $y_2 \neq x_1$, then take a smaller interval $I_2 \subseteq I_1$ around $y_2$ that avoids $x_1$.

**Corollary 4.4.2.** *Nondegenerate Intervals and the Real Line Are Uncountable*
*Let $J \subseteq \mathbb{R}$ be an interval.*

*(i) If $J$ contains two distinct points, then $J$ is uncountable.*
*(ii) $\mathbb{R}$ is uncountable.*

*Proof.*

(i) Choose $u < v$ in $J$ and then $a, b$ with $u < a < b < v$. Since $J$ is an interval, $I := [a, b] \subseteq J$.
The set $I$ is a nonempty 1-cell, hence compact by Theorem 4.2.3 and closed by Theorem 4.1.5.
For every $x \in I$ and $\varepsilon > 0$, $I \cap (B_\varepsilon(x) \setminus \{x\}) \neq \emptyset$, so $I$ is perfect.
Thus $I$ is uncountable by Theorem 4.4.1; hence $J$ is uncountable.

(ii) Since $\mathbb{R}$ is an interval and $0, 1 \in \mathbb{R}$ with $0 < 1$, (i) applied to $J = \mathbb{R}$ gives that $\mathbb{R}$ is uncountable.

*Structural Commentary*: Closed subintervals are perfect compact subsets.
*Key Insight*: One closed subinterval already forces uncountability. □

**Example 4.4.3.** *The Cantor Set*
Set $E_0 = [0, 1]$ and obtain $E_{n+1}$ from $E_n$ by deleting the open middle third of each component.
Then $E_n$ is a union of $2^n$ compact intervals of length $3^{-n}$ and $E_{n+1} \subseteq E_n$.
The *Cantor set* is $P := \bigcap_{n=0}^{\infty} E_n$. Since $(E_n)$ is nested nonempty compact, $P \neq \emptyset$ by Corollary 4.1.9.
Since $P$ is closed in $E_0$, $P$ is compact. Every nonempty open interval $J \subseteq [0, 1]$ contains a deleted middle third $R$ with $R \cap P = \emptyset$, so $P$ contains no nontrivial interval.If $x \in P$ and $U$ is an interval neighborhood of $x$, choose $n$ with the component $I_n(x) \subseteq U$; an endpoint $y \neq x$ of $I_n(x)$ lies in $P$, so $U \cap (P \setminus \{x\}) \neq \emptyset$. Hence $P$ is perfect.
*Structural Commentary*: The Cantor set is built by nested compact deletion, not by listing points.
*Key Insight*: It is compact and perfect, but contains no interval.




Figure 140: The first stages $E_0, E_1, E_2, E_3$ in the Cantor-set construction.

## 4.5 Complete Metric Spaces and Completions

Completeness turns Cauchy approximations into actual points. The examples below record the standard complete spaces used throughout analysis, then isolate the completion construction that adjoins missing Cauchy limits to an arbitrary metric space.

**Example 4.5.1.** *Finite-Dimensional Complete Metric Spaces*
For $k \in \mathbb{N}$, $(\mathbb{R}^k, d_2)$ and $(\mathbb{C}^k, d_2)$ are complete; in particular, so are $\mathbb{R}$ and $\mathbb{C}$.

*Proof.* Completeness of $\mathbb{R}$ is Theorem 2.1.15.
If $z_n = x_n + iy_n$ is Cauchy in $\mathbb{C}$, then $(x_n)$ and $(y_n)$ are Cauchy in $\mathbb{R}$, so $z_n \to x + iy \in \mathbb{C}$.
Corollary 4.2.8 gives completeness of $\mathbb{R}^k$, and the isometry $\mathbb{C}^k \cong \mathbb{R}^{2k}$ gives completeness of $\mathbb{C}^k$.
*Structural Commentary*: Finite-dimensional completeness is coordinatewise completeness.
*Key Insight*: A Cauchy sequence in finitely many coordinates converges coordinate by coordinate. □

**Definition 4.5.2.** *Contraction*
Let $X$ be a metric space with metric $d$.
A map $\varphi : X \to X$ is a *contraction* if $\exists$ a constant $0 \leq c < 1$ such that $d(\varphi(x), \varphi(y)) \leq c\, d(x, y)$ for $x, y \in X$.
*Structural Commentary*: Bound $c < 1$ excludes nonexpansive maps and is the hypothesis in Theorem 4.5.3.

**Theorem 4.5.3.** *Contraction Principle*
*If $X$ is a complete metric space and $\varphi : X \to X$ is a contraction, then $\exists$ a unique $x \in X$ such that $\varphi(x) = x$.*

*Proof.* Choose $x_0 \in X$ and define $x_{n+1} = \varphi(x_n)$.
If $c < 1$ is a contraction constant, induction gives $d(x_{n+1}, x_n) \leq c^n d(x_1, x_0)$.
For $n < m$, the triangle inequality and the finite geometric sum give $d(x_n, x_m) \leq \sum_{j=n}^{m-1} d(x_j, x_{j+1}) \leq \frac{c^n}{1-c} d(x_1, x_0)$.
Thus $(x_n)$ is Cauchy, so completeness gives $x_n \to x \in X$.
Since a contraction is continuous, $\varphi(x) = \lim \varphi(x_n) = \lim x_{n+1} = x$.
If $\varphi(x) = x$ and $\varphi(y) = y$, then $d(x, y) \leq c\, d(x, y)$, hence $d(x, y) = 0$ and $x = y$.
*Structural Commentary*: Iteration turns the contraction estimate into a geometric Cauchy estimate.
*Key Insight*: Completeness supplies the limiting point; the strict constant $c < 1$ forces uniqueness. □




Figure 141: A contraction pulls iterates toward the unique fixed point.

**Perspective 4.5.4.** *Fixed Points as Existence Proofs*
A contraction proof solves $x = \varphi(x)$ by iteration in a complete metric space. This is the template behind the inverse function theorem, Theorem 4.1.4.

**Example 4.5.5.** *A Noncomplete Metric Space*
The rational numbers with $d(x, y) = |x - y|$ are not complete.
Let $(r_n)$ be the decimal truncation sequence $1.4,\ 1.41,\ 1.414,\ 1.4142, \ldots$, for $\sqrt{2}$.
This is a concrete instance of the truncation squeeze in Proposition 2.2.11. If $k, n \geq N$, then $|r_k - r_n| \leq 10^{-N}$, so $(r_n)$ is Cauchy in $\mathbb{Q}$. As a sequence in $\mathbb{R}$, it converges to $\sqrt{2}$.
If it converged to some $r \in \mathbb{Q}$ in the inherited metric, uniqueness of limits in $\mathbb{R}$ would force $r = \sqrt{2}$, contradicting Proposition 1.1.7. Thus $\mathbb{Q}$ is not complete in the inherited metric.
*Structural Commentary*: The decimal approximants are Cauchy before their missing limit is available inside $\mathbb{Q}$.
*Key Insight*: Completeness asks whether every internally Cauchy approximation has an internal endpoint.

**Perspective 4.5.6.** *Cauchy Construction as a Parallel Model*
Cuts build $\mathbb{R}$ here; equivalence classes of rational Cauchy sequences give the parallel model.
Let $\mathcal{C}_{\mathbb{Q}}$ be the set of rational Cauchy sequences and set $(a_n) \sim (b_n)$ iff $|a_n - b_n| \to 0$; then $\mathbb{R} \simeq \mathcal{C}_{\mathbb{Q}}/\!\sim$.
Operations and order must be independent of representatives.




Figure 142: Real numbers as equivalence classes of rational Cauchy representatives.

**Perspective 4.5.7.** *Completion by Cauchy Sequences*
Complete by identifying Cauchy sequences with $d(p_n, q_n) \to 0$ and setting $D([(p_n)], [(q_n)]) = \lim_{n\to\infty} d(p_n, q_n)$.
Constant sequences embed the original space; triangle inequalities and diagonals give a complete metric.

**Theorem 4.5.8.** *Metric Completion*

*For every metric space $(X, d)$, $\exists$ complete metric space $(\widehat{X}, \widehat{d})$, isometry $\iota : X \to \widehat{X}$ with $\overline{\iota(X)}^{\widehat{X}} = \widehat{X}$.*
*If $(Y, d_Y, \tau)$, $(Z, d_Z, \sigma)$ are two such completions, then $\exists!$ bijective isometry $\Phi : Y \to Z$ s.t. $\Phi \circ \tau = \sigma$.*

*Proof.*

i. *Existence.* Let $\mathrm{CS}(X)$ be the Cauchy sequences in $X$.
For $a = (a_n), b = (b_n) \in \mathrm{CS}(X)$, $|d(a_n, b_n) - d(a_m, b_m)| \leq d(a_n, a_m) + d(b_n, b_m)$, so $D(a, b) := \lim_{n\to\infty} d(a_n, b_n)$ exists. Put $a \sim b \Leftrightarrow D(a, b) = 0$, $\widehat{X} := \mathrm{CS}(X)/\!\sim$, and $\widehat{d}([a], [b]) := D(a, b)$.
The estimate gives well-definedness; metric axioms pass to limits. Map $\iota(x) := [(x, x, \ldots)]$ is an isometry.
If $a = (a_n)$ and $d(a_n, a_N) < \varepsilon$ for $n \geq N$, then $\widehat{d}(\iota(a_N), [a]) \leq \varepsilon$, hence $\overline{\iota(X)}^{\widehat{X}} = \widehat{X}$.
Let $([a^{(k)}])$ be Cauchy in $\widehat{X}$. Choose $x_k \in X$ with $\widehat{d}(\iota(x_k), [a^{(k)}]) < 1/k$.
Since $d(x_k, x_l) \leq 1/k + \widehat{d}([a^{(k)}], [a^{(l)}]) + 1/l$, $(x_k)$ is Cauchy in $X$,
and $\widehat{d}([a^{(k)}], [(x_j)]) \leq 1/k + \limsup_{j\to\infty} d(x_k, x_j) \to 0$. Thus $([a^{(k)}]) \to [(x_j)]$, so $(\widehat{X}, \widehat{d})$ is complete.

ii. *Uniqueness.* Let $(Y, d_Y, \tau)$ and $(Z, d_Z, \sigma)$ be completions.
Define $\rho_0 : \tau(X) \to \sigma(X)$ by $\rho_0(\tau x) := \sigma x$; it is an isometry.

For $y \in Y$, choose $y_n \in \tau(X)$ with $y_n \to y$. Since $\rho_0(y_n)$ is Cauchy in $Z$, define $\Phi(y) := \lim_{n\to\infty} \rho_0(y_n)$.
If $z_n \in \tau(X)$ and $z_n \to y$, then $d_Z(\rho_0(y_n), \rho_0(z_n)) = d_Y(y_n, z_n) \to 0$, so $\Phi$ is well defined.
If $y'_n \to y'$, then $d_Z(\Phi(y), \Phi(y')) = \lim_{n\to\infty} d_Z(\rho_0(y_n), \rho_0(y'_n)) = \lim_{n\to\infty} d_Y(y_n, y'_n) = d_Y(y, y')$.
Hence $\Phi$ is an isometry extending $\rho_0$. Its image is complete, hence closed in $Z$, and contains the dense set $\sigma(X)$; therefore $\Phi(Y) = Z$. Uniqueness follows from continuity and density of $\tau(X)$.

*Structural Commentary*: Missing limits are installed as equivalence classes of Cauchy approximations.
*Key Insight*: Completion adds Cauchy limits and is unique over the dense copy of $X$. □




Figure 143: For completions $Y, Z$, the dense-copy isometry $\sigma\tau^{-1} : \tau(X) \to \sigma(X)$ extends uniquely to $\overline{\rho} : Y \cong Z$.

# 5 Connectedness and Continuity

## 5.1 Connectedness and Continuity

Connectedness prevents a space from splitting into separated pieces; continuity transports topological structure through maps. This subsection also records IVT/EVT and real-line refinements.

**Definition 5.1.1.** *Separated and Connected Sets*
Let $X$ be a metric space.

(i) Subsets $A, B \subseteq X$ are *separated* if $A \cap \overline{B} = \overline{A} \cap B = \emptyset$.
(ii) A subset $E \subseteq X$ is *connected* if no nonempty separated subsets $A, B \subseteq E$ satisfy $E = A \cup B$.
(iii) A subset $U \subseteq X$ is *clopen* if it is both open and closed.

*Structural Commentary*: Separated sets avoid each other's closures.
*Key Insight*: Disconnected means a nontrivial clopen split; continuous preimages preserve such splits.

**Theorem 5.1.2.** *Connected Subsets of the Real Line*
*A set $E \subseteq \mathbb{R}$ is connected if and only if $x, y \in E,\ x < y \Longrightarrow [x, y] \subseteq E$.*

*Proof.*

(⇒) If the interval property fails, choose $x, y \in E$ and $x < z < y$ with $z \notin E$. Then $A := E \cap (-\infty, z)$ and $B := E \cap (z, \infty)$ are nonempty separated sets with $E = A \cup B$, contradicting connectedness.
(⇐) Suppose $E = A \cup B$ with $A, B$ nonempty separated. Choose $x \in A$, $y \in B$, $x < y$, and set $S := A \cap [x, y]$, $z := \sup S$. By Theorem 3.3.5, $z \in \overline{A}$; hence $z \notin B$ and $z < y$. If $z \notin A$, then $z \in [x, y] \setminus E$.
If $z \in A$, then $z \notin \overline{B}$, so choose $z_1 \in (z, y)$ with $z_1 \notin B$; by maximality of $z$, also $z_1 \notin A$.
Thus $z_1 \in [x, y] \setminus E$, contradicting $[x, y] \subseteq E$.

*Structural Commentary*: On the real line, connectedness is exactly having no gaps between any two of its points.
*Key Insight*: A separation creates a missing point by taking a supremum at the boundary of one side. □

**Method 5.1.3.** *Compactness and Connectedness Proof Patterns*

(i) *Open-cover compactness*: for every cover $\{G_\alpha\}_{\alpha\in A}$, find finite $F \subseteq A$ with $K \subseteq \bigcup_{\alpha\in F} G_\alpha$.
(ii) *Euclidean closed-bounded compactness*: if $E \subseteq \mathbb{R}^k$ is closed and bounded, choose a compact $k$-cell $I \supseteq E$ and apply Theorem 4.1.6.
(iii) *Connectedness on $\mathbb{R}$*: replace connectedness of $E \subseteq \mathbb{R}$ by $x, y \in E, x < y \Longrightarrow [x, y] \subseteq E$.

**Definition 5.1.4.** *Limit of a Function Between Metric Spaces*
Let $X, Y$ be metric spaces, $E \subseteq X$, $f : E \to Y$, and let $p$ be a limit point of $E$. For $q \in Y$, $\lim_{x\to p} f(x) = q$ means that, for every $\varepsilon > 0$, there exists $\delta > 0$ such that $x \in E$, $0 < d_X(x, p) < \delta \Longrightarrow d_Y(f(x), q) < \varepsilon$.
The point $p$ need not belong to $E$, and $f(p)$ is irrelevant when it does.
*Key Insight*: Domain $E$ controls approach: $J \setminus \{a\}$ gives two-sided limit, $(a, b)$ or $(c, a)$ gives one-sided limits.




Figure 144: The epsilon–delta geometry for $\lim_{x\to p} f(x) = q$.

**Theorem 5.1.5.** *Sequential Characterization of Function Limits*
*Let $X, Y$ be metric spaces, $E \subseteq X$, $f : E \to Y$, and let $p$ be a limit point of $E$.*
*For $q \in Y$, $\lim_{x\to p} f(x) = q$ if and only if $f(p_n) \to q$ for every sequence $(p_n) \subseteq E \setminus \{p\}$ with $p_n \to p$.*

*Proof.*

($\Rightarrow$) If $p_n \to p$ with $p_n \in E \setminus \{p\}$, then eventually $0 < d_X(p_n, p) < \delta$; hence $d_Y(f(p_n), q) < \varepsilon$, so $f(p_n) \to q$.
($\Leftarrow$) If the limit fails, then for some $\varepsilon > 0$ and each $n$ there is $x_n \in E$ with $0 < d_X(x_n, p) < 1/n$ and $d_Y(f(x_n), q) \geq \varepsilon$. Then $x_n \to p$, $x_n \neq p$, but $f(x_n) \not\to q$.

*Structural Commentary*: Function limits can be tested by all sequences approaching the input point.
*Key Insight*: The negation of an epsilon–delta limit manufactures a bad sequence. □

**Corollary 5.1.6.** *Uniqueness of Function Limits*
*If a function has a limit at $p$, then that limit is unique.*

*Proof.* Let $q$ and $r$ be limits of $f$ at $p$. Choose $p_n \in E \setminus \{p\}$ with $d_X(p_n, p) < 1/n$.
Then $p_n \to p$, so Theorem 5.1.5 gives $f(p_n) \to q$ and $f(p_n) \to r$. By Theorem 3.4.2, $q = r$.
*Structural Commentary*: Limit uniqueness is inherited from metric sequence uniqueness.
*Key Insight*: One approaching sequence cannot have two different limits. □

**Theorem 5.1.7.** *Algebra of Function Limits*
*Let $X$ be a metric space, $E \subseteq X$, $p$ be limit point of $E$.*
*For $f, g : E \to \mathbb{C}$, assume $\lim_{x\to p} f(x) = A$ and $\lim_{x\to p} g(x) = B$. Then:*

*(i)* $\lim_{x\to p}(f+g)(x) = A + B$;
*(ii)* $\lim_{x\to p}(fg)(x) = AB$;
*(iii)* *if $B \neq 0$ and $f/g$ is defined near $p$, then* $\lim_{x\to p}(f/g)(x) = A/B$.

*For $f, g : E \to \mathbb{R}^k$ with limits $a, b$, the corresponding limits are $f + g \to a + b$, $\lambda f \to \lambda a$, and $f \cdot g \to a \cdot b$.*

*Proof.*

(i) For any $p_n \in E \setminus \{p\}$ with $p_n \to p$, Theorem 5.1.5 gives $f(p_n) \to A$ and $g(p_n) \to B$. Hence $(f+g)(p_n) \to A + B$ by Theorem 2.1.4, so Theorem 5.1.5 gives (i).
(ii) The same sequence gives $(fg)(p_n) \to AB$ by Theorem 2.1.4; hence (ii).
(iii) If $B \neq 0$ and $f/g$ is defined near $p$, then every such sequence is eventually in the domain of $f/g$, and $(f/g)(p_n) \to A/B$ by Theorem 2.1.4; hence (iii).

The vector laws follow by the same sequential characterization and Theorem 3.4.7.
*Structural Commentary*: Limit algebra for functions is pulled back to sequence algebra.
*Key Insight*: Sequential characterization lets previous sequence theorems do the work. □

**Proposition 5.1.8.** *Function Limits Are Local*
*Let $X, Y$ be metric spaces, let $E \subseteq X$, let $f : E \to Y$, let $p$ be a limit point of $E$, and let $q \in Y$.*
*For every $r > 0$,* $\lim_{x\to p;\, x\in E} f(x) = q \iff \lim_{x\to p;\, x\in E\cap B_r(p)} f(x) = q$.

*Proof.*

($\Rightarrow$) Since $E \cap B_r(p) \subseteq E$, any $\delta$ witnessing the limit on $E$ also witnesses the limit on $E \cap B_r(p)$.

($\Leftarrow$) Assume the restricted limit. Given $\varepsilon > 0$, choose $\delta_0 > 0$ such that $x \in E \cap B_r(p)$ and $0 < d_X(x,p) < \delta_0$ imply $d_Y(f(x), q) < \varepsilon$. Let $\delta := \min\{\delta_0, r\}$.
If $x \in E$ and $0 < d_X(x,p) < \delta$, then $x \in E \cap B_r(p)$, so $d_Y(f(x), q) < \varepsilon$.

*Structural Commentary*: Values away from $p$ cannot affect the limit at $p$.
*Key Insight*: Shrink $\delta$ until every tested point lies in the chosen local neighborhood. □

**Definition 5.1.9.** *Continuity at a Point and on a Set*
Let $(X, d_X), (Y, d_Y)$ be metric spaces, $E \subseteq X$, and $f : E \to Y$.

(i) For $p \in E$, $f$ is *continuous at $p$* if $\forall \varepsilon > 0\ \exists \delta > 0\ \forall x \in E,\ d_X(x,p) < \delta \implies d_Y(f(x), f(p)) < \varepsilon$.
(ii) $f$ is *continuous on $E$* if $\forall p \in E$, $f$ is continuous at $p$.
(iii) If $p$ is isolated in $E$, then $f$ is continuous at $p$.

**Corollary 5.1.10.** *Discrete Domains Make Every Map Continuous*
*If $X$ carries the discrete metric, then every map $f : X \to Y$ into any metric space $Y$ is continuous.*

*Proof.* For $p \in X$, $B_{1/2}(p) = \{p\}$. Take $\delta = \frac{1}{2}$. If $d_X(x,p) < \delta$, then $x = p$, so $d_Y(f(x), f(p)) = 0 < \varepsilon$.
*Structural Commentary*: In the discrete metric, no distinct point can approach $p$.
*Key Insight*: Pointwise continuity is automatic when small enough balls are singletons. □




Figure 145: Continuity at $x_0$: for every $\varepsilon > 0$, choose $\delta > 0$ with $f(B_\delta(x_0)) \subset B_\varepsilon(f(x_0))$.

**Theorem 5.1.11.** *Continuity as Equality with the Function Limit*
*Let $X, Y$ be metric spaces, $E \subseteq X$, $p \in E$, and $f : E \to Y$.*
*If $p$ is a limit point of $E$, then $f$ is continuous at $p$ if and only if $\lim_{x \to p} f(x) = f(p)$.*

*Proof.*

($\Rightarrow$) If $0 < d_X(x,p) < \delta$, continuity at $p$ gives $d_Y(f(x), f(p)) < \varepsilon$, hence $\lim_{x \to p} f(x) = f(p)$.
($\Leftarrow$) If $\lim_{x \to p} f(x) = f(p)$, the same $\delta$ proves continuity for $x \neq p$; for $x = p$, $d_Y(f(p), f(p)) = 0$.

*Structural Commentary*: Continuity is the compatibility of the value at $p$ with nearby values.
*Key Insight*: Limits ignore the point; continuity ties the point back in. □

**Remark 5.1.12.** *Adherent-Point Limit Convention*
The adherent-point convention takes $p \in \overline{E}$ and allows $x = p$ in the $\delta$-test; Definition 5.1.4 instead assumes $p$ is a limit point and requires $0 < d_X(x,p)$. If $p \in E$, the unpunctured convention also tests $f(p)$, forcing $q = f(p)$; in the punctured convention this is exactly continuity at $p$ by Theorem 5.1.11.
*Key Insight*: Punctured limits study nearby values; continuity adds compatibility with the value at the point.

**Remark 5.1.13.** *The Sequence-First View*
For functions on subsets of $\mathbb{R}$, continuity at limit points is equivalently sequential: $x_n \to p$ in the domain implies $f(x_n) \to f(p)$; isolated points are automatic by Theorems 5.1.5 and 5.1.11.

**Theorem 5.1.14.** *Sequential Characterization of Continuity*
*Let $(X, d_X), (Y, d_Y)$ be metric spaces, $f : X \to Y$, and $p \in X$.*
*Then $f$ is continuous at $p$ if and only if $\forall (x_n) \subseteq X$, $x_n \to p \implies f(x_n) \to f(p)$.*

*Proof.*

($\Rightarrow$) Assume $f$ is continuous at $p$ and let $x_n \to p$.
Given $\varepsilon > 0$, choose $\delta > 0$ such that $d_X(x,p) < \delta \implies d_Y(f(x), f(p)) < \varepsilon$.
Since $x_n \to p$, eventually $d_X(x_n, p) < \delta$, hence eventually $d_Y(f(x_n), f(p)) < \varepsilon$. Thus $f(x_n) \to f(p)$.
($\Leftarrow$) Suppose the sequential condition holds. If $f$ is not continuous at $p$, then $\exists \varepsilon_0 > 0$ such that $\forall \delta > 0$, $\exists x \in X$ with $d_X(x,p) < \delta$ and $d_Y(f(x), f(p)) \geq \varepsilon_0$. Taking $\delta = 1/n$ gives $x_n \in X$ with $d_X(x_n, p) < 1/n$ and $d_Y(f(x_n), f(p)) \geq \varepsilon_0$. Then $x_n \to p$, but $f(x_n) \not\to f(p)$, a contradiction.

*Structural Commentary*: Failure of epsilon–delta continuity manufactures a bad sequence.
*Key Insight*: In metric spaces, sequences detect continuity at points. □

**Theorem 5.1.15.** *Composition of Continuous Functions*
*Let $E \subseteq X$ and let $f : E \to Y$, $g : f(E) \to Z$ be maps between metric spaces.*
*If $f$ is continuous at $p \in E$ and $g$ is continuous at $f(p)$, then $g \circ f$ is continuous at $p$.*

*Proof.* Given $\varepsilon > 0$, choose $\eta > 0$ for $g$ at $f(p)$, then choose $\delta > 0$ for $f$ at $p$ with tolerance $\eta$.
If $x \in E$ and $d_X(x,p) < \delta$, then $d_Y(f(x), f(p)) < \eta$, hence $d_Z(g(f(x)), g(f(p))) < \varepsilon$.
*Structural Commentary*: The output tolerance for $g$ becomes the input tolerance for $f$.
*Key Insight*: Continuity composes by passing epsilon through an intermediate eta. □

**Theorem 5.1.16.** *The Distance Function Is Continuous*
*Let $(X,d)$ be a metric space and $D : X \times X \to \mathbb{R}$, $D(x,y) := d(x,y)$.*
*For $d_\times \in \{d_1, d_2, d_\infty\}$ on $X \times X$, $D : (X \times X, d_\times) \to \mathbb{R}$ is continuous.*
*Equivalently, $x_n \to x$ and $y_n \to y$ in $X$ imply $d(x_n, y_n) \to d(x,y)$.*

*Proof.* For $x, y, u, v \in X$, $|D(x,y) - D(u,v)| = |d(x,y) - d(u,v)| \leq d(x,u) + d(y,v)$. Hence $x_n \to x$ and $y_n \to y$ imply $D(x_n, y_n) \to D(x,y)$. Proposition 3.4.9 and Theorem 5.1.14 give continuity for each product metric.
*Structural Commentary*: Distance changes continuously when both endpoints move.
*Key Insight*: The triangle inequality controls distance error by endpoint errors. □

**Theorem 5.1.17.** *Continuity via Open Preimages*
*For metric spaces $X, Y$, map $f : X \to Y$ is continuous if and only if $f^{-1}(V)$ is open in $X \forall$ open $V \subseteq Y$.*

*Proof.*
($\Rightarrow$) If $p \in f^{-1}(V)$ and $V$ is open, choose $\varepsilon > 0$ with $B_\varepsilon(f(p)) \subseteq V$.
Continuity gives $\delta > 0$ with $B_\delta(p) \subseteq f^{-1}(V)$, so $f^{-1}(V)$ is open.
($\Leftarrow$) For $p \in X$ and $\varepsilon > 0$, the ball $V = B_\varepsilon(f(p))$ is open by Theorem 3.1.13.
Since $p \in f^{-1}(V)$ and $f^{-1}(V)$ is open, some $B_\delta(p) \subseteq f^{-1}(V)$; this is continuity at $p$.
*Structural Commentary*: Continuity is exactly preservation of openness under inverse image.
*Key Insight*: Images are not Boolean-exact, but preimages are; continuity belongs to preimages. □

**Proposition 5.1.18.** *Continuity on a Relative Domain*
*Let $(X, d_X)$ and $(Y, d_Y)$ be metric spaces, let $E \subseteq X$, and let $f : E \to Y$. The following are equivalent:*
*(i) $f : (E, d_X|_{E\times E}) \to (Y, d_Y)$ is continuous.*
*(ii) For every open $V \subseteq Y$, $f^{-1}(V)$ is open relative to $E$.*
*(iii) For every open $V \subseteq Y$, there exists open $G \subseteq X$ such that $f^{-1}(V) = E \cap G$.*

*Proof.*
*(i)* $\Rightarrow$ *(ii)* For open $V \subseteq Y$, Theorem 5.1.17 on the metric space $E$ gives $f^{-1}(V)$ open in $E$.
*(ii)* $\Rightarrow$ *(iii)* For open $V \subseteq Y$, relative openness of $f^{-1}(V) \subseteq E$ gives $f^{-1}(V) = E \cap G$ for some open $G \subseteq X$.
*(iii)* $\Rightarrow$ *(i)* For open $V \subseteq Y$, $f^{-1}(V) = E \cap G$ with $G$ open in $X$, hence $f^{-1}(V)$ is open in $E$.
Theorem 5.1.17 on $E$ gives continuity.
*Structural Commentary*: Open preimages are relative opens in $E$.
*Key Insight*: Relative continuity says ordinary open sets in $X$ are seen through intersection with $E$. □




Figure 146: Continuity pulls open sets back to open sets.

**Corollary 5.1.19.** *Continuity via Closed Preimages*
*Let $X, Y$ be metric spaces, $f : X \to Y$. Then $f$ is continuous iff $f^{-1}(C)$ is closed in $X$ for every closed $C \subseteq Y$.*

*Proof.*
($\Rightarrow$) For closed $C \subseteq Y$, $X \setminus f^{-1}(C) = f^{-1}(Y \setminus C)$ is open by Theorem 5.1.17; hence $f^{-1}(C)$ is closed.
($\Leftarrow$) For open $V \subseteq Y$, $X \setminus f^{-1}(V) = f^{-1}(Y \setminus V)$ is closed by hypothesis; hence $f^{-1}(V)$ is open, QED.

*Structural Commentary*: Closed-set continuity is the complement form of open-set continuity.
*Key Insight*: $f^{-1}(Y \setminus C) = X \setminus f^{-1}(C)$ transfers open and closed formulations. □

**Perspective 5.1.20.** *Three Views of Continuity*
For metric-space maps, continuity is equivalently $\varepsilon$–$\delta$ control, $x_n \to x \Rightarrow f(x_n) \to f(x)$, and $V \subseteq Y$ open $\Rightarrow f^{-1}(V)$ open; see Definition 5.1.9 and Theorems 5.1.14, 5.1.17.
*Key Insight*: Continuity sends near points forward and pulls open sets back.

**Proposition 5.1.21.** *Distance-to-Set Is Continuous*
*Let $(X, d)$ be a metric space, let $\emptyset \neq E \subseteq X$, and define $f : X \to \mathbb{R}$ by $f(p) := \operatorname{dist}(p, E)$.*

*(i) $f$ is 1-Lipschitz: $|f(p) - f(q)| \leq d(p, q)$ for all $p, q \in X$; in particular, $f$ is continuous.*
*(ii) $f^{-1}(\{0\}) = \overline{E}$.*

*Proof.*

(i) For $p, q \in X$ and $e \in E$, $d(p, e) \leq d(p, q)+d(q, e)$. Taking the infimum over $e \in E$ gives $f(p) \leq d(p, q)+f(q)$, hence $f(p) - f(q) \leq d(p, q)$. Hence $f(q) - f(p) \leq d(p, q)$, so $|f(p) - f(q)| \leq d(p, q)$, QED.
(ii) For $p \in X$, $p \in f^{-1}(\{0\})$ iff $\operatorname{dist}(p, E) = 0$ iff $p \in \overline{E}$ by Proposition 3.1.24.

*Structural Commentary*: Distance to a fixed set changes no faster than the point itself moves.
*Key Insight*: The triangle inequality turns nearest-set distance into a continuous detector of closure. □

**Corollary 5.1.22.** *Open and Closed Sets by Distance Functions*
*Let $X$ be a metric space.*

*(i) A set $O \subseteq X$ is open if and only if there is a continuous $f : X \to \mathbb{R}$ such that $O = \{x \in X : f(x) > 0\}$.*
*(ii) A set $F \subseteq X$ is closed if and only if there is a continuous $g : X \to \mathbb{R}$ such that $F = g^{-1}(\{0\})$.*

*Proof.*

(i) If $O = X$, take $f \equiv 1$. Otherwise set $f(x) := \operatorname{dist}(x, O^c)$. Since $O^c$ is closed, Proposition 5.1.21 gives $\{f = 0\} = O^c$, hence $O = \{f > 0\}$. Conversely, $\{f > 0\} = f^{-1}((0, \infty))$ is open by Theorem 5.1.17.
(ii) If $F = \emptyset$, take $g \equiv 1$. Otherwise set $g(x) := \operatorname{dist}(x, F)$.
If $F$ closed, by Proposition 5.1.21, $g^{-1}(\{0\}) = \overline{F} = F$. Conversely, $g^{-1}(\{0\})$ closed by Corollary 5.1.19.

*Structural Commentary*: Metric spaces encode open and closed sets by continuous real-valued distance functions.
*Key Insight*: A set is closed exactly when its distance-zero locus has no extra points. □

**Definition 5.1.23.** *Homeomorphism*
A *homeomorphism* from $X$ to $Y$ is a bijection $f : X \to Y$ such that both $f$ and $f^{-1}$ are continuous.
If such a map exists, $X$ and $Y$ are *homeomorphic*, written $X \simeq Y$.

**Proposition 5.1.24.** *Homeomorphisms Preserve Topology*
*If $f : X \to Y$ is a homeomorphism, then $U \mapsto f(U)$ bijects open subsets of $X$ with open subsets of $Y$.*

*Proof.* If $U \subseteq X$ is open, then $f(U) = (f^{-1})^{-1}(U)$ is open in $Y$ since $f^{-1}$ is continuous. If $V \subseteq Y$ is open, then $f^{-1}(V)$ is open in $X$ since $f$ is continuous. These two assignments are inverse because $f$ is bijective.
*Structural Commentary*: A homeomorphism preserves open sets in both directions.
*Key Insight*: Topological sameness is continuity plus continuous reversibility. □

**Definition 5.1.25.** *Metric Isometry*
Let $(X, d_X), (Y, d_Y)$ be metric spaces.

(i) A map $f : X \to Y$ is an *isometry* if $\forall x, y \in X$, $d_Y(f(x), f(y)) = d_X(x, y)$.
(ii) $X$ and $Y$ are *isometric* if $\exists$ a bijective isometry $f : X \to Y$.

**Theorem 5.1.26.** *Basic Properties of Metric Isometries*
*Let $f : (X, d_X) \to (Y, d_Y)$ be an isometry.*

*(i) $f$ is injective.*
*(ii) $f$ is continuous.*
*(iii) $f^{-1} : (f(X), d_Y|_{f(X)\times f(X)}) \to (X, d_X)$ is an isometry.*

*Proof.*

(i) If $f(x) = f(y)$, then $d_X(x, y) = d_Y(f(x), f(y)) = 0$, hence $x = y$.
(ii) If $x_n \to x$ in $X$, then $d_Y(f(x_n), f(x)) = d_X(x_n, x) \to 0$; by Theorem 5.1.14, $f$ is continuous.
(iii) For $u = f(x), v = f(y) \in f(X)$, $d_X(f^{-1}(u), f^{-1}(v)) = d_X(x, y) = d_Y(f(x), f(y)) = d_Y(u, v)$; QED.

*Structural Commentary*: An isometry preserves all distances, so injectivity and continuity are automatic.
*Key Insight*: Distance preservation is stronger than topological equivalence. □

**Remark 5.1.27.** *Metric Versus Linear Isometries*
Metric isometry is purely distance-theoretic. If $T : V \to W$ is linear between inner-product spaces, then $\|Tv\| = \|v\| \ \forall v \in V \iff d_W(Tu, Tv) = d_V(u, v) \ \forall u, v \in V$.

**Example 5.1.28.** *Continuous Bijection Need Not Be a Homeomorphism*
Let $f : [0, 2\pi) \to S^1$, and $f(t) = (\cos t, \sin t)$. This continuous bijection is not a homeomorphism: points of $S^1$ approaching $(1, 0)$ from below have preimages tending to $2\pi$, not 0.




Figure 147: Parametrization $[0, 2\pi) \to S^1$ wraps continuously and bijectively, but inverse jumps at $(1, 0)$.

**Remark 5.1.29.** *Continuity Only Uses the Domain Metric*
For $f : E \to Y$ with $E \subseteq X$, continuity on $E$ depends only on the restricted metric on $E$, not on $X \setminus E$.

**Definition 5.1.30.** *Product Map*
If $f : X \to Y$ and $g : X \to Z$ are maps between metric spaces, their *product map*, or direct sum, is $f \oplus g : X \to Y \times Z$, $(f \oplus g)(x) := (f(x), g(x))$, where $Y \times Z$ carries any of product metrics in Definition 3.4.8.

**Proposition 5.1.31.** *Continuity of Product Maps*
*The product map $f \oplus g : X \to Y \times Z$ is continuous at $x_0$ iff $f$ and $g$ are continuous at $x_0$.*
*Hence it is continuous on $X$ iff both components are.*

*Proof.* By Proposition 3.4.9, it suffices to use $d_{\text{sum}}$.
($\Rightarrow$) Given $\varepsilon > 0$, choose $\delta > 0$ from continuity of $f \oplus g$ at $x_0$.
If $d_X(x, x_0) < \delta$, then $d_Y(f(x), f(x_0)), d_Z(g(x), g(x_0)) \le d_{\text{sum}}((f \oplus g)(x), (f \oplus g)(x_0)) < \varepsilon$.
($\Leftarrow$) Given $\varepsilon > 0$, choose $\delta_f, \delta_g$ for $f, g$ at tolerance $\varepsilon/2$ and set $\delta = \min\{\delta_f, \delta_g\}$.
Then $d_X(x, x_0) < \delta$ gives $d_{\text{sum}}((f \oplus g)(x), (f \oplus g)(x_0)) = d_Y(f(x), f(x_0)) + d_Z(g(x), g(x_0)) < \varepsilon$.
*Structural Commentary*: Product continuity is exactly componentwise continuity.
*Key Insight*: A product metric packages two output estimates into one. □

**Theorem 5.1.32.** *Continuous Algebra and Components*
*Let $X$ be a metric space.*
*(i) If $f, g : X \to \mathbb{C}$ are continuous, so are $f + g$, $fg$, and $f/g$ where $g \neq 0$.*
*(ii) For $f = (f_1, \ldots, f_k) : X \to \mathbb{R}^k$, $f$ is continuous if and only if each $f_j$ is continuous.*
*(iii) If $f, g : X \to \mathbb{R}^k$ are continuous, so are $f + g$ and $f \cdot g$.*

*Proof.*
(i) At isolated points continuity is automatic; at limit points, use Theorems 5.1.7 and 5.1.11.
(ii) By Theorem 3.4.7, convergence in $\mathbb{R}^k$ is exactly coordinatewise convergence; apply this to $f(x_n) \to f(x)$.
(iii) Apply (ii) to the coordinate functions of $f + g$ and $f \cdot g$, using the scalar algebra from (i).
*Structural Commentary*: Continuity is stable under the algebraic operations already stable for limits.
*Key Insight*: Componentwise control is equivalent to Euclidean control in finite dimensions. □

**Example 5.1.33.** *Coordinate, Polynomial, Rational, and Norm Maps*
Coordinate functions on $\mathbb{R}^k$ are continuous; hence polynomials are continuous, and rational functions are continuous where defined. Also $\mathbf{x} \mapsto \|\mathbf{x}\|_2$ is continuous since $\big|\|\mathbf{x}\|_2 - \|\mathbf{y}\|_2\big| \le \|\mathbf{x} - \mathbf{y}\|_2$.
Thus $p \mapsto \|f(p)\|_2$ is continuous whenever $f : X \to \mathbb{R}^k$ is, by Theorem 5.1.15.

**Definition 5.1.34.** *Bounded Mapping*
A mapping $f : E \to \mathbb{R}^k$ is *bounded* if there exists $M \in \mathbb{R}$ such that $\|f(x)\|_2 \le M$ for every $x \in E$.

**Theorem 5.1.35.** *Continuous Images of Compact Sets Are Compact*
*If $f : X \to Y$ is continuous and $X$ is compact, then $f(X)$ is compact.*

*Proof.* Let $\{V_\alpha\}$ cover $f(X)$. Then $\{f^{-1}(V_\alpha)\}$ is an open cover of $X$ by Theorem 5.1.17.
Choose $\alpha_1, \ldots, \alpha_n$ with $X \subseteq \bigcup_{i=1}^n f^{-1}(V_{\alpha_i})$; then $f(X) \subseteq \bigcup_{i=1}^n V_{\alpha_i}$.
*Structural Commentary*: Pull back, take a finite subcover, push forward.
*Key Insight*: Compactness is preserved by continuous images because inverse images preserve openness. □

**Corollary 5.1.36.** *Continuous Images into $\mathbb{R}^k$ Are Closed and Bounded*
*If $f : X \to \mathbb{R}^k$ is continuous and $X$ is compact, then $f(X)$ is closed and bounded. In particular, $f$ is bounded.*

*Proof.* By Theorem 5.1.35, $f(X)$ is compact.
By Theorem 4.2.5, compact subsets of $\mathbb{R}^k$ are closed and bounded; hence $f$ is bounded.
*Structural Commentary*: Compactness in the domain becomes closed-bounded geometry in Euclidean range.
*Key Insight*: Continuous functions on compact spaces cannot escape to infinity. □

**Theorem 5.1.37.** *Extreme Value Theorem*
*Let $f : X \to \mathbb{R}$ be continuous, let $X$ be compact.*
*If $M := \sup_{p\in X} f(p)$, $m := \inf_{p\in X} f(p)$, then $\exists\ p, q \in X$ such that $f(p) = M$ and $f(q) = m$.*

*Proof.* By Corollary 5.1.36, $f(X)$ is closed and bounded in $\mathbb{R}$. Thus $\sup f(X)$ and $\inf f(X)$ exist and belong to $f(X)$ by Theorems 3.3.5 and 4.3.3. Hence $M = f(p)$ and $m = f(q)$ for some $p, q \in X$.
*Structural Commentary*: Compactness turns suprema and infima into attained values.
*Key Insight*: Closed bounded images contain their endpoints. □

**Theorem 5.1.38.** *Compact Bijective Maps Have Continuous Inverses*
*If $f : X \to Y$ is a continuous bijection from compact $X$ to a metric space $Y$, then $f^{-1} : Y \to X$ is continuous.*

*Proof.* It suffices, by Theorem 5.1.17, to show that $f(V)$ is open in $Y$ for every open $V \subseteq X$. If $V$ is open, then $V^c$ is compact by Theorem 4.1.6, so $f(V^c)$ is compact, hence closed by Theorems 5.1.35 and 4.1.5.
Since $f$ is bijective, $f(V) = Y \setminus f(V^c)$ is open.
*Structural Commentary*: Compactness turns closed complements into closed images.
*Key Insight*: A continuous bijection from compact domain has no room for inverse discontinuity. □

**Remark 5.1.39.** *Preimages, Not Forward Images*
Theorem 5.1.17 is about inverse images. A continuous map need not send open sets to open sets or closed sets to closed sets; compactness is one hypothesis that recovers closedness of images through Theorem 5.1.35.

**Theorem 5.1.40.** *Continuous Images of Connected Sets Are Connected*
*If $f : X \to Y$ is continuous and $E \subseteq X$ is connected, then $f(E)$ is connected.*

*Proof.* If $f(E) = A \cup B$ with $A, B$ nonempty separated, set $G := E \cap f^{-1}(A)$ and $H := E \cap f^{-1}(B)$. Then $E = G \cup H$ and $G, H$ are nonempty. Since $G \subseteq f^{-1}(\overline{A})$ and $f^{-1}(\overline{A})$ is closed, $\overline{G} \subseteq f^{-1}(\overline{A})$; hence $\overline{G} \cap H = \emptyset$.
Similarly $G \cap \overline{H} = \emptyset$, so $G, H$ separate $E$, a contradiction.
*Structural Commentary*: A separation of the image pulls back to a separation of the domain.
*Key Insight*: Continuous maps cannot create a gap in the image of a connected set. □

**Theorem 5.1.41.** *Intermediate Value Theorem*
*If $f : [a, b] \to \mathbb{R}$ is continuous and $c$ lies strictly between $f(a)$ and $f(b)$, then $f(x) = c$ for some $x \in (a, b)$.*

*Proof.* The interval $[a, b]$ is connected by Theorem 5.1.2, so $f([a, b])$ is connected by Theorem 5.1.40.
Connected subsets of $\mathbb{R}$ contain every value between any two of their points; hence $c = f(x)$ for some $x \in [a, b]$.
Since $c \neq f(a), f(b)$, $x \in (a, b)$.
*Structural Commentary*: The intermediate value theorem is connectedness of intervals transported by continuity.
*Key Insight*: Continuous real functions on intervals cannot jump over a value. □

**Method 5.1.42.** *Bisection Proof of the Intermediate Value Theorem*
Let $f : [a, b] \to \mathbb{R}$ be continuous and let $c$ lie between $f(a)$ and $f(b)$.
Replacing $(f, c)$ by $(-f, -c)$ if needed, assume $f(a) \leq c \leq f(b)$.
If $c = f(a)$ or $c = f(b)$, take the corresponding endpoint; otherwise $f(a) < c < f(b)$.

(i) Set $[a_1, b_1] := [a, b]$.
(ii) Given $[a_n, b_n]$ with $f(a_n) \leq c \leq f(b_n)$, set $m_n := (a_n + b_n)/2$.
If $f(m_n) \geq c$, set $[a_{n+1}, b_{n+1}] := [a_n, m_n]$; otherwise set $[a_{n+1}, b_{n+1}] := [m_n, b_n]$.

Then, for all $n$, $[a_{n+1}, b_{n+1}] \subseteq [a_n, b_n]$, $f(a_n) \leq c \leq f(b_n)$, and $b_n - a_n = (b - a)/2^{n-1}$.
By Theorem 4.2.1, choose $x_0 \in \bigcap_n [a_n, b_n]$. Since $b_n - a_n \to 0$, the intersection is $\{x_0\}$ and $a_n, b_n \to x_0$.
Continuity gives $f(a_n), f(b_n) \to f(x_0)$, so $f(x_0) \leq c \leq f(x_0)$; hence $f(x_0) = c$.
If $f(a) < c < f(b)$, then $x_0 \in (a, b)$.
*Key Insight*: Bisection keeps the target value bracketed while nested intervals collapse to one point.

**Definition 5.1.43.** *Path-Connected Set*
A set $E$ in a metric space is *path-connected* if for every $p, q \in E$ there is a continuous map $\gamma : [a, b] \to E$ with $\gamma(a) = p$ and $\gamma(b) = q$. Such a map $\gamma$ is a *path* from $p$ to $q$.

**Theorem 5.1.44.** *Path-Connected Sets Are Connected*
*Every path-connected set is connected.*

*Proof.* Suppose $E = A \cup B$ with $A, B$ nonempty separated.
Choose $p \in A$, $q \in B$, and a path $\gamma : [a, b] \to E$ from $p$ to $q$. Since $[a, b]$ is connected, so is $\gamma([a, b])$.
But $\gamma([a, b])$ meets both $A$ and $B$, which separate it, a contradiction.
*Structural Commentary*: A path transports connectedness from an interval to the set.
*Key Insight*: To connect two points by a path, the path image cannot be split into two separated pieces. □

**Corollary 5.1.45.** *Convex Sets Are Path-Connected*
*Every convex subset of $\mathbb{R}^k$ is path-connected.*

*Proof.* For $\mathbf{x}, \mathbf{y} \in E$, set $\gamma(t) = (1-t)\mathbf{x}+t\mathbf{y}$ on $[0, 1]$. Convexity gives $\gamma([0, 1]) \subseteq E$, with $\gamma(0) = \mathbf{x}$ and $\gamma(1) = \mathbf{y}$.
Since the coordinates of $\gamma$ are affine in $t$, $\gamma$ is continuous. Hence $E$ is path-connected.
*Structural Commentary*: Convexity supplies the whole straight path between any two points.
*Key Insight*: In Euclidean space, line segments are continuous paths. □

## 5.2 Uniform Continuity and Compactness

**Definition 5.2.1.** *Uniform Continuity*
Let $(X, d_X)$ and $(Y, d_Y)$ be metric spaces, and let $f : X \to Y$.
The map $f$ is *uniformly continuous* on $X$ if $\forall \varepsilon > 0\, \exists \delta > 0\, \forall p, q \in X,\ d_X(p, q) < \delta \implies d_Y(f(p), f(q)) < \varepsilon$.
*Structural Commentary*: Uniform continuity is setwise: for each $\varepsilon$, one $\delta$ must work for all pairs in the set.

**Principle 5.2.2.** *Challenge–Response Order*
Limits and continuity are controlled by quantifier order. In Definition 5.1.4, the $\varepsilon$-challenge at the range point must be answered by a $\delta$-restriction near the domain point. Uniform continuity, Definition 5.2.1, strengthens the same pattern by requiring one $\delta$ to work for all relevant domain points at once.

**Proposition 5.2.3.** *Sequential Test for Uniform Continuity*
*Let $X, Y$ be metric spaces and let $f : X \to Y$. The following are equivalent:*

*(i) $f$ is uniformly continuous on $X$;*
*(ii) whenever $(p_n)$ and $(q_n)$ are sequences in $X$ with $d_X(p_n, q_n) \to 0$, one has $d_Y(f(p_n), f(q_n)) \to 0$.*

*Proof.*

(i) $\Rightarrow$ (ii) Given $\varepsilon > 0$, choose $\delta > 0$ such that $d_X(x, y) < \delta$ implies $d_Y(f(x), f(y)) < \varepsilon$ for all $x, y \in X$.
Since $d_X(p_n, q_n) \to 0$, eventually $d_X(p_n, q_n) < \delta$, hence eventually $d_Y(f(p_n), f(q_n)) < \varepsilon$.

(ii) $\Rightarrow$ (i) Prove the contrapositive. If $f$ is not uniformly continuous, then $\exists \varepsilon_0 > 0$ such that $\forall n \geq 1$, $\exists p_n, q_n \in X$ with $d_X(p_n, q_n) < 1/n$ and $d_Y(f(p_n), f(q_n)) \geq \varepsilon_0$.
Thus $d_X(p_n, q_n) \to 0$ but $d_Y(f(p_n), f(q_n)) \not\to 0$, contradicting (ii).

*Structural Commentary*: Uniform continuity preserves vanishing pairwise distances.
*Key Insight*: Failure of one global $\delta$ manufactures two close input sequences with separated outputs. □

**Definition 5.2.4.** *Lipschitz Map*
Let $(X, d_X)$ and $(Y, d_Y)$ be metric spaces, let $f : X \to Y$, and let $L \geq 0$.
The map $f$ is *$L$-Lipschitz* if $\forall p, q \in X$, $d_Y(f(p), f(q)) \leq L d_X(p, q)$.
It is *Lipschitz* if it is $L$-Lipschitz for some $L \geq 0$.
*Key Insight*: Lipschitz control is one global linear modulus of continuity.

**Proposition 5.2.5.** *Lipschitz Implies Uniform Continuity*
*Every Lipschitz map between metric spaces is uniformly continuous, hence continuous.*

*Proof.* Let $f$ be $L$-Lipschitz and let $\varepsilon > 0$. If $L = 0$, then $d_Y(f(p), f(q)) = 0$ for all $p, q \in X$.
If $L > 0$, take $\delta := \varepsilon/L$; $d_X(p, q) < \delta$ implies $d_Y(f(p), f(q)) \leq L d_X(p, q) < \varepsilon$.
Thus $f$ is uniformly continuous. Fixing one point gives continuity.
*Structural Commentary*: A global Lipschitz bound gives a global modulus.
*Key Insight*: One constant controls all pairs. □

**Example 5.2.6.** *Continuous Need Not Be Lipschitz*
On $[0,1]$, $f(x)=\sqrt{x}$ is continuous since $\left|\sqrt{x}-\sqrt{y}\right| \le \sqrt{|x-y|}$.
It is not Lipschitz: $L$-Lipschitz would imply $\sqrt{x} \le Lx$ for $0<x\le 1$, i.e. $1/\sqrt{x} \le L$, impossible as $x \downarrow 0$.

**Theorem 5.2.7.** *Continuity on Compact Sets Is Uniform*
*If $f: X \to Y$ is continuous and $X$ is compact, then $f$ is uniformly continuous.*

*Proof.* Let $\varepsilon>0$. For each $a\in X$, choose $r_a>0$ such that $d_X(x,a)<r_a$ implies $d_Y(f(x),f(a))<\varepsilon/2$.
Compactness gives a finite cover $X \subseteq \bigcup_{i=1}^{n} B_{r_{a_i}/2}(a_i)$. Set $\delta := \frac{1}{2}\min_i r_{a_i} > 0$.
If $d_X(p,q)<\delta$ and $p \in B_{r_{a_i}/2}(a_i)$, then $d_X(q,a_i)<r_{a_i}$ also; hence $f(p)$ and $f(q)$ within $\varepsilon/2$ of $f(a_i)$. QED.
*Structural Commentary*: Compactness converts pointwise radii into finitely many radii with positive minimum.
*Key Insight*: Uniform continuity is continuity with the point-dependence removed by a finite subcover. □

**Example 5.2.8.** *Separating Continuity, Uniform Continuity, and Lipschitz Control*

(i) Let $f:\mathbb{R}\to\mathbb{R}$ be $f(x)=x^2$. Then $f$ is continuous, but not uniformly continuous: with $p_n=n$ and $q_n=n+1/n$, $|p_n-q_n|=1/n\to 0$ while $|f(q_n)-f(p_n)|=2+1/n^2 \not\to 0$.
(ii) Let $g:[0,1]\to\mathbb{R}$ be $g(x)=\sqrt{x}$. Then $g$ is uniformly continuous by Theorem 5.2.7. If $g$ were $L$-Lipschitz, then $1/n=\left|g(1/n^2)-g(0)\right| \le L/n^2\ \forall n\in\mathbb{N}$, so $n\le L\ \forall n$, impossible. Hence $g$ is not Lipschitz.

*Structural Commentary*: Both reverse implications fail.
*Key Insight*: Compactness repairs uniform continuity, but it does not force a linear modulus.

**Theorem 5.2.9.** *Compactness Hypotheses Are Essential*
*Let $E$ be a noncompact subset of $\mathbb{R}$. Then:*

*(i) there exists a continuous function on $E$ which is not bounded;*
*(ii) there exists a continuous bounded function on $E$ which has no maximum;*
*(iii) if $E$ is bounded, then there exists a continuous function on $E$ which is not uniformly continuous.*

*Proof.*

(i) If $E$ is unbounded, $f(x)=x$ works. If $E$ is bounded, Theorem 4.2.5 gives $x_0 \in \overline{E}\setminus E$, and $f(x)=1/(x-x_0)$ is continuous on $E$ and unbounded.
(ii) If $E$ is unbounded, $h(x)=x^2/(1+x^2)$ has $\sup_E h = 1$ but $h<1$. If $E$ is bounded, with $x_0$ as above, $g(x)=1/(1+(x-x_0)^2)$ is bounded by 1 and has supremum 1 but no maximum.
(iii) If $E$ is bounded, let $f(x)=1/(x-x_0)$. Given $\delta>0$, let $x\in E$ with $|x-x_0|<\delta/2$, then $y\in E$ with $|y-x_0|<\min\{\delta/2, 1/(|f(x)|+1)\}$. Then $|x-y|<\delta$, $|f(x)-f(y)|>1$, so $f$ is not uniformly continuous.

*Structural Commentary*: Compactness is essential in the preceding theorems.
*Key Insight*: Noncompact sets either escape to infinity or miss an accumulation point. □




Figure 148: For $f(x)=1/x^2$ near 0, a fixed vertical tolerance forces smaller pointwise horizontal tolerances.

**Method 5.2.10.** *Continuity Proof Patterns*

(i) *$\varepsilon$–$\delta$ at $p$*: prove $\forall\varepsilon>0\,\exists\delta>0$ such that $x\in E$ and $d_X(x,p)<\delta$ imply $d_Y(f(x),f(p))<\varepsilon$.
(ii) *Limit at $p$*: prove same estimate with $0<d_X(x,p)<\delta$, target $q$; equivalently test all $x_n\to p$, $x_n\neq p$.
(iii) *Topological continuity*: prove $V\subseteq Y$ open $\Rightarrow f^{-1}(V)$ open, or $C\subseteq Y$ closed $\Rightarrow f^{-1}(C)$ closed.
(iv) *Compactness*: from open cover $\{V_\alpha\}$ of $f(X)$, pulled-back cover $\{f^{-1}(V_\alpha)\}$ of $X$ extract finite subcover.
(v) *Uniform continuity*: choose local radii $r_p$, cover compact $X$ by $B_{r_p/2}(p)$, and set $\delta=\frac{1}{2}\min_i r_{p_i}$.

*Key Insight*: Continuity is local control; compactness turns local controls into finite control.

## 5.3 One-Sided Limits, Monotone Functions, and Extended Limits

**Definition 5.3.1.** *One-Sided Limits and Types of Discontinuity*
Let $X \subseteq \mathbb{R}$, $f : X \to \mathbb{R}$, $x_0 \in \mathbb{R}$, $q \in \mathbb{R}$, $X_+ := X \cap (x_0, \infty)$, and $X_- := X \cap (-\infty, x_0)$.

(i) If $x_0$ is a limit point of $X_+$, then $f(x_0+) = q$ means $\lim_{x \to x_0}(f|_{X_+})(x) = q$.
(ii) If $x_0$ is a limit point of $X_-$, then $f(x_0-) = q$ means $\lim_{x \to x_0}(f|_{X_-})(x) = q$.
(iii) If $x_0$ is a limit point of $X_+$ and $X_-$, then $\lim_{x \to x_0}(f|_{X \setminus \{x_0\}})(x)$ exists iff $f(x_0+)$ and $f(x_0-)$ exist with $f(x_0+) = f(x_0-)$.

For an interval $I$, $f : I \to \mathbb{R}$, and an interior discontinuity $x_0 \in I^\circ$:

(i) The discontinuity is of the *first kind*, or *simple*, iff both $f(x_0+)$ and $f(x_0-)$ exist.
(ii) A first-kind discontinuity is a *jump* iff $f(x_0+) \neq f(x_0-)$.
(iii) A first-kind discontinuity is *removable* iff $f(x_0+) = f(x_0-) \neq f(x_0)$.
(iv) The discontinuity is of the *second kind* iff at least one of $f(x_0+)$ and $f(x_0-)$ fails to exist.

**Example 5.3.2.** *Basic Discontinuity Types*
Dirichlet's function $\mathbf{1}_{\mathbb{Q}}$ has a discontinuity of the second kind at every point.
The function $f(x) = x$ on $\mathbb{Q}$ and $f(x) = 0$ off $\mathbb{Q}$ is continuous only at 0; elsewhere its discontinuities are of the second kind. A jump in a piecewise linear function is a simple discontinuity.
Use Corollary 1.1.22: every interval contains rational and irrational points.




Figure 149: For $g = \mathbf{1}_{\mathbb{Q}}$, interval around $c$ has rational inputs value 1 and irrational inputs value 0.

**Definition 5.3.3.** *Monotone Functions*
Let $f : (a, b) \to \mathbb{R}$.

(i) $f$ is *monotonically increasing* iff $a < x < y < b$ implies $f(x) \leq f(y)$.
(ii) $f$ is *monotonically decreasing* iff $a < x < y < b$ implies $f(x) \geq f(y)$.
(iii) $f$ is *monotone* iff $f$ is monotonically increasing or monotonically decreasing.

**Theorem 5.3.4.** *One-Sided Limits of Monotone Functions*
*If $f$ is increasing on $(a, b)$, then for every $x \in (a, b)$, $\sup_{a<t<x} f(t) = f(x-) \leq f(x) \leq f(x+) = \inf_{x<t<b} f(t)$.*
*Moreover, $a < x < y < b$ implies $f(x+) \leq f(y-)$. For decreasing $f$, reverse the inequalities.*

*Proof.* Let $A := \sup_{a<t<x} f(t)$.
For every $\varepsilon > 0$, choose $s < x$ with $A - \varepsilon < f(s) \leq A$; then $s < t < x$ implies $A - \varepsilon < f(t) \leq A$, so $f(x-) = A$.
Let $B := \inf_{x<t<b} f(t)$. For every $\varepsilon > 0$, choose $s > x$ with $B \leq f(s) < B + \varepsilon$; then $x < t < s$ implies $B \leq f(t) < B + \varepsilon$, so $f(x+) = B$. If $x < y$, then $\inf_{x<t<y} f(t) \leq \sup_{x<t<y} f(t)$ gives $f(x+) \leq f(y-)$.
For decreasing $f$, apply the increasing case to $-f$ and multiply the resulting inequalities by $-1$.
*Structural Commentary*: Monotonicity converts one-sided limits into suprema and infima.
*Key Insight*: A monotone function has jumps, but not oscillatory one-sided failure. □

**Corollary 5.3.5.** *Monotone Functions Have No Second-Kind Discontinuities*
*A monotone real function on an interval has no discontinuities of the second kind.*

*Proof.* At every interior point, monotonicity gives both one-sided limits, so any discontinuity is of the first kind.
*Structural Commentary*: Monotonicity rules out oscillatory discontinuities.
*Key Insight*: All monotone discontinuities are jumps or removable value mismatches. □

**Theorem 5.3.6.** *Monotone Continuity and Interval Images*
*Let $f : (a, b) \to \mathbb{R}$ be monotone. Then $f$ is continuous iff $f((a, b))$ is an interval.*

*Proof.* Put $I = (a, b)$.
($\Rightarrow$) If $f$ is continuous, then $f(I)$ is connected, hence an interval.

($\Leftarrow$) Contrapositively, let $f$ be discontinuous at $x_0 \in I$. Replace $f$ by $-f$ if needed, assume $f$ increasing. Then $f(x_0-) \le f(x_0) \le f(x_0+)$, and discontinuity gives $f(x_0-) < f(x_0+)$. Choose $y \in (f(x_0-), f(x_0+)) \setminus \{f(x_0)\}$. Then $x < x_0$ gives $f(x) < y$, $x > x_0$,$f(x) > y$, $f(x_0) \neq y$. So $y \notin f(I)$ while $f(I)$ has values below and above $y$. Hence $f(I)$ is not an interval.

*Structural Commentary*: Monotone discontinuities create image gaps.
*Key Insight*: Interval images exclude jumps. □




Figure 150: A jump of an increasing function leaves values between $f(x_0-)$ and $f(x_0+)$ out of the image; the red marker indicates such a $y \notin f(I)$, so $f(I)$ cannot be an interval.

**Theorem 5.3.7.** *Monotone Discontinuities Are Countable*
*If $f$ is monotone on $(a,b)$, then the set of points of $(a,b)$ at which $f$ is discontinuous is at most countable.*

*Proof.* Assume $f$ is increasing, and let $E$ be its discontinuity set.
For each $x \in E$, Theorem 5.3.4 gives $f(x-) < f(x+)$; choose $r(x) \in \mathbb{Q}$ with $f(x-) < r(x) < f(x+)$.
If $x_1 < x_2$, then $f(x_1+) \le f(x_2-)$, so $r(x_1) \neq r(x_2)$. Thus $E$ injects into $\mathbb{Q}$, hence is countable by Corollary 6.1.15. If $f$ is decreasing, apply the increasing case to $-f$, which has the same discontinuity set.
*Structural Commentary*: Each jump contains a rational marker, and monotonicity keeps the markers distinct.
*Key Insight*: Countability enters by injecting discontinuities into $\mathbb{Q}$. □

**Remark 5.3.8.** *Monotone Discontinuities Need Not Be Isolated*
For countable $E = \{x_n : n \in \mathbb{N}\} \subset (a,b)$ and positive summable $(c_n)$, $f(x) = \sum_{x_n < x} c_n$ is increasing, has jumps of size $c_n$ at $x_n$, and is continuous off $E$. Thus monotone discontinuities may be dense.

**Definition 5.3.9.** *Extended Limits and Limits at Infinity*
Neighborhoods of $+\infty$ are intervals $(c, +\infty)$; neighborhoods of $-\infty$ are $(-\infty, c)$.
For $f : E \to \mathbb{R}$ and extended real $A, x$, $f(t) \to A$ as $t \to x$ means: every neighborhood $U$ of $A$ contains $f(t)$ for all punctured $t \in E$ sufficiently near $x$, i.e. $\exists V \ni x$ such that $t \in V \cap E,\ t \neq x \implies f(t) \in U$.
For real $A, x$, this agrees with Definition 5.1.4.
*Structural Commentary*: Extended limits use same neighborhood test, $\pm\infty$ allowed as input or output limits.

**Theorem 5.3.10.** *Algebra of Extended Limits*
*Let $f, g : E \to \mathbb{R}$ and $f(t) \to A$, $g(t) \to B$ as $t \to x$, with $A, B, x$ extended real. Then, whenever RHS is defined:*

*(i) if $f(t) \to A'$ as $t \to x$, then $A' = A$;*
*(ii) $(f+g)(t) \to A + B$;*
*(iii) $(fg)(t) \to AB$;*
*(iv) $(f/g)(t) \to A/B$;*

*The forms $\infty - \infty$, $0 \cdot \infty$, $\infty/\infty$, and $A/0$ are undefined.*

*Proof.*

(i) If $A' \neq A$, choose disjoint neighborhoods $U$ of $A$ and $U'$ of $A'$. The two limits give a common neighborhood $V$ of $x$ such that $t \in V \cap E,\ t \neq x$ implies $f(t) \in U \cap U'$, a contradiction.
(ii) For any neighborhood $W$ of $A+B$, definedness gives neighborhoods $U$ of $A$ and $V$ of $B$ such that $u+v \in W$ for $u \in U$, $v \in V$. Apply the two limits in a common punctured neighborhood of $x$.
(iii) The same neighborhood argument applies to multiplication, excluding $0 \cdot \infty$.
(iv) Apply it to division; since $B \neq 0$, choose the neighborhood of $B$ away from 0.

*Structural Commentary*: Extended limit algebra is ordinary limit algebra in neighborhood form.
*Key Insight*: The undefined forms are exactly where neighborhood control breaks. □

# 6 General Topology for Analysis

## 6.1 Topological Spaces, Bases, and Separation

General topology abstracts the open-set behavior of metric spaces. This selective pass records spaces, bases, subspaces, closure, and separation axioms for later use.

**Definition 6.1.1.** *Topological Space*
A *topology* on a set $X$ is a collection $\mathcal{T} \subseteq 2^X$ such that:

(i) $\emptyset, X \in \mathcal{T}$;
(ii) if $\{\mathcal{O}_\alpha\}_{\alpha \in A} \subseteq \mathcal{T}$, then $\bigcup_{\alpha \in A} \mathcal{O}_\alpha \in \mathcal{T}$;
(iii) if $\mathcal{O}_1, \dots, \mathcal{O}_n \in \mathcal{T}$ with $n \geq 1$, then $\bigcap_{j=1}^{n} \mathcal{O}_j \in \mathcal{T}$.

Then $(X, \mathcal{T})$ is a *topological space*.

**Definition 6.1.2.** *Open, Closed, and Neighborhoods*
Let $(X, \mathcal{T})$ be a topological space.

(i) A set $U \subseteq X$ is *open* iff $U \in \mathcal{T}$.
(ii) A set $F \subseteq X$ is *closed* iff $X \setminus F \in \mathcal{T}$.
(iii) For $x \in X$, a *neighborhood* of $x$ is an open set $U \subseteq X$ with $x \in U$.
(iv) For $A \subseteq X$, a *neighborhood* of $A$ is an open set $U \subseteq X$ with $A \subseteq U$.

**Proposition 6.1.3.** *Open Sets by Neighborhoods*
*Let $(X, \mathcal{T})$ be a topological space and let $E \subseteq X$. Then $E \in \mathcal{T}$ iff for every $x \in E$ there exists an open set $U_x \in \mathcal{T}$ such that $x \in U_x \subseteq E$.*

*Proof.*
($\Rightarrow$) If $E \in \mathcal{T}$, take $U_x = E$ for each $x \in E$.
($\Leftarrow$) If such $U_x$ exists for every $x \in E$, then $E = \bigcup_{x \in E} U_x$. Since arbitrary unions of open sets are open, $E \in \mathcal{T}$.
*Structural Commentary*: Openness is checked pointwise by open subsets.
*Key Insight*: Openness is exactly the local ability to fit an open neighborhood inside the set. □

**Example 6.1.4.** *Discrete and Trivial Topologies*
On a nonempty set $X$, the *discrete topology* is $2^X$, so every subset is open and closed. It is induced by the discrete metric. The *trivial topology* is $\{\emptyset, X\}$, so the only neighborhood of any point is $X$.

**Proposition 6.1.5.** *Metric Open Sets Form a Topology*
*Let $(X, d)$ be metric space, $B_r(x) := \{y \in X : d(x, y) < r\}$, and $\tau_d := \{U \subseteq X : \forall x \in U,\ \exists r > 0,\ B_r(x) \subseteq U\}$. Then $\tau_d$ is a topology on $X$.*

*Proof.* Clearly $\emptyset, X \in \tau_d$. If $\{\mathcal{O}_\alpha\}_{\alpha \in A} \subseteq \tau_d$ and $x \in \bigcup_{\alpha \in A} \mathcal{O}_\alpha$, choose $\alpha$ with $x \in \mathcal{O}_\alpha$ and $r > 0$ with $B_r(x) \subseteq \mathcal{O}_\alpha$; hence $B_r(x) \subseteq \bigcup_{\alpha \in A} \mathcal{O}_\alpha$. If $\mathcal{O}_1, \dots, \mathcal{O}_n \in \tau_d$, $n \geq 1$, and $x \in \bigcap_{j=1}^{n} \mathcal{O}_j$, choose $r_j > 0$ with $B_{r_j}(x) \subseteq \mathcal{O}_j$ and set $r := \min_{1 \leq j \leq n} r_j$; then $B_r(x) \subseteq \bigcap_{j=1}^{n} \mathcal{O}_j$. Thus $\tau_d$ is a topology on $X$.
*Structural Commentary*: Metric balls generate an open-set system with the exact axioms of a topology.
*Key Insight*: General topology keeps the open-set calculus and forgets the distance function. □

**Definition 6.1.6.** *Closure, Interior, and Boundary*
Let $E \subseteq X$ in a topological space.

(i) The *closure* $\overline{E}$ is the set of all $x \in X$ such that every neighborhood of $x$ meets $E$.
(ii) The *interior* $\operatorname{int}(E)$ is the union of all open sets contained in $E$.
(iii) The *exterior* of $E$ is $\operatorname{int}(X \setminus E)$.
(iv) The *boundary* of $E$ is $\operatorname{bd}(E) := \overline{E} \setminus \operatorname{int}(E)$.

**Proposition 6.1.7.** *Closure, Interior, and Boundary Tests*
*Let $(X, \mathcal{T})$ be a topological space and let $E \subseteq X$.*

*(i) $\operatorname{int}(E)$ is the largest open subset of $E$, and $E$ is open iff $E = \operatorname{int}(E)$;*
*(ii) $\overline{E}$ is the smallest closed subset of $X$ containing $E$, and $E$ is closed iff $E = \overline{E}$;*
*(iii) $x \in \operatorname{bd}(E)$ iff every neighborhood of $x$ meets both $E$ and $X \setminus E$;*
*(iv) $\operatorname{bd}(E) = \overline{E} \cap \overline{X \setminus E}$, so $\operatorname{bd}(E)$ is closed.*

*Proof.* By definition, $\operatorname{int}(E) = \bigcup \{U \in \mathcal{T} : U \subseteq E\}$, hence it is the largest open subset of $E$. If $x \notin \overline{E}$, choose a neighborhood $U$ of $x$ with $U \cap E = \emptyset$; then $U \subseteq X \setminus \overline{E}$, so $\overline{E}$ is closed. If $F$ is closed and $E \subseteq F$, then $X \setminus F$ is disjoint from $E$, so $\overline{E} \subseteq F$. Thus $\overline{E}$ is the smallest closed subset of $X$ containing $E$.

*(i)* ($\Rightarrow$) If $E$ is open, then $E \subseteq \operatorname{int}(E) \subseteq E$, so $E = \operatorname{int}(E)$.

(*i*) ($\Leftarrow$) If $E = \operatorname{int}(E)$, then $E$ is open because $\operatorname{int}(E)$ is open.
(*ii*) ($\Rightarrow$) If $E$ is closed, then the smallest-closed-superset property gives $\overline{E} \subseteq E$, while $E \subseteq \overline{E}$. Hence $E = \overline{E}$.
(*ii*) ($\Leftarrow$) If $E = \overline{E}$, then $E$ is closed because $\overline{E}$ is closed.
(*iii*) ($\Rightarrow$) If $x \in \operatorname{bd}(E) = \overline{E} \setminus \operatorname{int}(E)$, then every neighborhood of $x$ meets $E$. Since $x \notin \operatorname{int}(E)$, no neighborhood of $x$ is contained in $E$, so every neighborhood of $x$ also meets $X \setminus E$.
(*iii*) ($\Leftarrow$) If every neighborhood of $x$ meets both $E$ and $X \setminus E$, then $x \in \overline{E}$ and $x \notin \operatorname{int}(E)$. Hence $x \in \operatorname{bd}(E)$.
(*iv*) By (iii), $x \in \operatorname{bd}(E)$ iff $x \in \overline{E}$ and $x \in \overline{X \setminus E}$. Thus $\operatorname{bd}(E) = \overline{E} \cap \overline{X \setminus E}$, which is closed by (ii).

*Structural Commentary*: Interior, closure, and boundary are neighborhood tests.
*Key Insight*: $\operatorname{int}(E)$ is certified inside; $\overline{E}$ is neighborhood-forced; $\operatorname{bd}(E)$ is where both sides persist. □

**Definition 6.1.8.** *Bases and Subbases*
Let $X$ be a set.

(i) If $\mathcal{T}$ is a topology on $X$ and $x \in X$, then a collection $\mathcal{B}_x$ of neighborhoods of $x$ is a *neighborhood basis at* $x$ iff every neighborhood $U$ of $x$ contains some $B \in \mathcal{B}_x$ with $B \subseteq U$.
(ii) A collection $\mathcal{B} \subseteq 2^X$ is a *basis* for a topology on $X$ iff $X = \bigcup_{B \in \mathcal{B}} B$ and, whenever $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, there exists $B_3 \in \mathcal{B}$ with $x \in B_3 \subseteq B_1 \cap B_2$.
(iii) If $\mathcal{B}$ is a basis on $X$, the *topology generated by* $\mathcal{B}$ is $\mathcal{T}_{\mathcal{B}} := \{\bigcup_{B \in \mathcal{U}} B : \mathcal{U} \subseteq \mathcal{B}\}$.
(iv) A collection $\mathcal{S} \subseteq 2^X$ is a *subbasis* on $X$ iff $\bigcup \mathcal{S} = X$ and its finite intersections form a basis on $X$.

*Key Insight*: Bases generate open sets by unions; subbases generate bases by finite intersections.

**Proposition 6.1.9.** *Basis Characterization*
*Let $X$ be a set and let $\mathcal{B} \subseteq 2^X$.*

*(i) If $\mathcal{B}$ is a basis on $X$, then $\mathcal{T}_{\mathcal{B}} := \{\bigcup \mathcal{U} : \mathcal{U} \subseteq \mathcal{B}\}$ is the unique topology on $X$ of all $\mathcal{B}$-unions.*
*(ii) If $\mathcal{T}$ is a topology on $X$, $\mathcal{B} \subseteq \mathcal{T}$, and $\forall \emptyset \neq U \in \mathcal{T}$ there is $\mathcal{U} \subseteq \mathcal{B}$ with $U = \bigcup \mathcal{U}$, then $\mathcal{B}$ is a basis for $\mathcal{T}$.*

*Proof.*

(i) Set $O_{\mathcal{U}} := \bigcup \mathcal{U}$ for $\mathcal{U} \subseteq \mathcal{B}$. Then $X = O_{\mathcal{B}}$, $\emptyset = O_{\emptyset}$, and $\bigcup_\alpha O_{\mathcal{U}_\alpha} = O_{\bigcup_\alpha \mathcal{U}_\alpha}$. If $x \in O_{\mathcal{U}} \cap O_{\mathcal{V}}$, choose $B_1 \in \mathcal{U}$ and $B_2 \in \mathcal{V}$ with $x \in B_1 \cap B_2$; by refinement, $x \in B_3 \subseteq B_1 \cap B_2$ for some $B_3 \in \mathcal{B}$. Hence $O_{\mathcal{U}} \cap O_{\mathcal{V}} = \bigcup\{B \in \mathcal{B} : B \subseteq O_{\mathcal{U}} \cap O_{\mathcal{V}}\}$, so $\mathcal{T}_{\mathcal{B}}$ is a topology. Uniqueness follows from the definition.
(ii) If $X \neq \emptyset$, apply the hypothesis to $U = X$ to get $X = \bigcup \mathcal{U}$ for some $\mathcal{U} \subseteq \mathcal{B}$; if $X = \emptyset$, the cover axiom is void. If $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, then $B_1 \cap B_2 \in \mathcal{T}$ and $B_1 \cap B_2 \neq \emptyset$, so $B_1 \cap B_2 = \bigcup \mathcal{V}$ for some $\mathcal{V} \subseteq \mathcal{B}$. Choose $B_3 \in \mathcal{V}$ with $x \in B_3 \subseteq B_1 \cap B_2$. Hence $\mathcal{B}$ is a basis and $\mathcal{T} = \mathcal{T}_{\mathcal{B}}$.

*Structural Commentary*: Local refinements make intersections open.
*Key Insight*: Basis pieces generate opens by unions. □

**Example 6.1.10.** *Metric Balls as a Basis*
In a metric space $(X, d)$, $\{B_d(x, r) : x \in X,\ r > 0\}$ is a basis. On $\mathbb{R}^k$, $\{B(q, r) : q \in \mathbb{Q}^k,\ r \in \mathbb{Q}_{>0}\}$ is a countable basis. In $[a, b] \subseteq \mathbb{R}$, $\{[a, c) : a < c < b\} \cup \{(c, b] : a < c < b\}$ is a subbasis.

**Definition 6.1.11.** *Subspace Topology*
Let $(X, \mathcal{T})$ be a topological space and let $\emptyset \neq E \subseteq X$.
The *subspace topology* on $E$ is $\mathcal{T}_E := \{E \cap U : U \in \mathcal{T}\}$, and $(E, \mathcal{T}_E)$ is a *subspace* of $X$.

**Definition 6.1.12.** *Separation Properties*
Let $(X, \mathcal{T})$ be a topological space. Disjoint sets $A, B \subseteq X$ are *separated by disjoint neighborhoods* if there exist $U, V \in \mathcal{T}$ such that $A \subseteq U$, $B \subseteq V$, and $U \cap V = \emptyset$. The space $X$ is:

(i) $T_1$ iff $\{x\}$ is closed for every $x \in X$;
(ii) *Hausdorff* iff $\{x\}$ and $\{y\}$ can be separated by disjoint neighborhoods for all distinct $x, y \in X$;
(iii) *regular* iff $X$ is $T_1$ and $\{x\}, F$ are separable by disjoint neighborhoods for closed $F \subseteq X$ and $x \notin F$;
(iv) *normal* iff $X$ is $T_1$ and any disjoint closed $F, G \subseteq X$ can be separated by disjoint neighborhoods.

*Key Insight*: Stronger separation axioms separate larger closed objects.

**Proposition 6.1.13.** $T_1$ *Spaces and Closed Points*
*Let $(X, \mathcal{T})$ be a topological space. Then $X$ is $T_1$ iff $\forall$ distinct $x, y \in X$ $\exists U \in \mathcal{T}$ with $x \in U$ and $y \notin U$.*

*Proof.*

($\Rightarrow$) Assume $X$ is $T_1$. If $x \neq y$, then $\{y\}$ is closed, so $U = X \setminus \{y\} \in \mathcal{T}$ satisfies $x \in U$ and $y \notin U$.
($\Leftarrow$) Assume the stated condition. Fix $y \in X$. For each $x \in X \setminus \{y\}$, choose $U_x \in \mathcal{T}$ with $x \in U_x$ and $y \notin U_x$. Then $X \setminus \{y\} = \bigcup_{x \in X \setminus \{y\}} U_x \in \mathcal{T}$, so $\{y\}$ is closed. Hence $X$ is $T_1$.

*Structural Commentary*: Point exclusion makes singleton complements open.
*Key Insight*: $T_1$ means each point can be omitted near every other point. □

**Proposition 6.1.14.** *Metric Spaces Are Normal*
*Let $(X, d)$ be a metric space with its metric topology.*

*(i) $\{x\}$ is closed for every $x \in X$.*
*(ii) If $F_1, F_2 \subseteq X$ closed and $F_1 \cap F_2 = \emptyset$, then $\exists$ disjoint open sets $O_1, O_2 \subseteq X$ with $F_i \subseteq O_i$ for $i = 1, 2$.*

*Hence $X$ is normal.*

*Proof.*

(i) Fix $x \in X$. For each $y \in X \setminus \{x\}$, $B_d(y, d(x, y)/2) \subseteq X \setminus \{x\}$, so $X \setminus \{x\}$ is open and $\{x\}$ is closed.
(ii) If one $F_i$ is empty, take $O_i = \emptyset$ and $O_{3-i} = X$. Assume $F_1, F_2 \neq \emptyset$. Set $d_i(x) := \operatorname{dist}(x, F_i)$ and $O_i := \{x \in X : d_i(x) < d_{3-i}(x)\}$ for $i = 1, 2$. By Proposition 5.1.21, $d_i$ is continuous and $d_i^{-1}(\{0\}) = F_i$. Hence $O_i = (d_i - d_{3-i})^{-1}((-\infty, 0))$ is open, and if $x \in F_i$, then $d_i(x) = 0 < d_{3-i}(x)$, so $F_i \subseteq O_i$. Finally, $O_1 \cap O_2 = \emptyset$, since $d_1(x) < d_2(x)$ and $d_2(x) < d_1(x)$ cannot both hold.

By Definition 6.1.12, (i) and (ii) imply that $X$ is normal.
*Structural Commentary*: Distance inequalities separate closed sets.
*Key Insight*: Compare distances to build disjoint neighborhoods. □

**Proposition 6.1.15.** *Normality by Nested Neighborhoods*
*Let $X$ be a $T_1$ topological space. Then $X$ is normal iff for every closed $F \subseteq X$ and every open $U \subseteq X$ with $F \subseteq U$, there exists an open $O \subseteq X$ such that $F \subseteq O \subseteq \overline{O} \subseteq U$.*

*Proof.*

($\Rightarrow$) Assume $X$ is normal. Let $F \subseteq X$ be closed and let $U \subseteq X$ be open with $F \subseteq U$.
The closed sets $F$ and $X \setminus U$ are disjoint, so choose disjoint open sets $O, V$ with $F \subseteq O$ and $X \setminus U \subseteq V$.
Then $O \subseteq X \setminus V \subseteq U$, and since $X \setminus V$ is closed, $\overline{O} \subseteq X \setminus V \subseteq U$.
($\Leftarrow$) Assume the nested-neighborhood property. Let $A, B \subseteq X$ be closed with $A \cap B = \emptyset$.
Apply the property to $F = A$ and $U = X \setminus B$ to get open $O$ with $A \subseteq O \subseteq \overline{O} \subseteq X \setminus B$.
Then $O$ and $X \setminus \overline{O}$ are disjoint open neighborhoods of $A$ and $B$, respectively. Since $X$ is $T_1$, $X$ is normal.

*Structural Commentary*: Normality controls closures inside neighborhoods.
*Key Insight*: Thicken $F$ without leaving $U$. □




Figure 151: Normality thickens closed $F \subseteq U$ to an open $O$ with $F \subseteq O \subseteq \overline{O} \subseteq U$.

## 6.2 Continuity and Topological Constructions

General topological constructions are organized by inverse images: homeomorphisms preserve open-set structure, while products, quotients, and weak topologies are characterized by which maps they make continuous.

**Definition 6.2.1.** *Continuity Between Topological Spaces*
Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, let $f : X \to Y$.
For $x \in X$, $y \in Y$, set $\mathcal{N}_X(x) := \{U \in \mathcal{T}_X : x \in U\}$, $\mathcal{N}_Y(y) := \{O \in \mathcal{T}_Y : y \in O\}$.

(i) For $x_0 \in X$, $f$ is *continuous at $x_0$* iff $\forall O \in \mathcal{N}_Y(f(x_0))\ \exists U \in \mathcal{N}_X(x_0)$ with $f(U) \subseteq O$.
(ii) $f$ is *continuous* iff $\forall x \in X$, $f$ is continuous at $x$.

**Proposition 6.2.2.** *Continuity by Open Preimages*
*Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, and let $f : X \to Y$.*

*(i) $f$ is continuous iff $\forall O \in \mathcal{T}_Y$, $f^{-1}(O) \in \mathcal{T}_X$.*
*(ii) If $A \subseteq X$, $\mathcal{T}_A = \{A \cap V : V \in \mathcal{T}_X\}$, and $f$ is continuous, then $f|_A : (A, \mathcal{T}_A) \to (Y, \mathcal{T}_Y)$ is continuous.*
*(iii) If $(Z, \mathcal{T}_Z)$ is a topological space, $g : (Y, \mathcal{T}_Y) \to (Z, \mathcal{T}_Z)$ is continuous, and $f$ is continuous, then $g \circ f : (X, \mathcal{T}_X) \to (Z, \mathcal{T}_Z)$ is continuous.*

*Proof.*

(*i*) ($\Rightarrow$) Let $O \in \mathcal{T}_Y$ and $x \in f^{-1}(O)$. Since $O \in \mathcal{N}_Y(f(x))$ and $f$ is continuous at $x$, choose $U_x \in \mathcal{N}_X(x)$ with $f(U_x) \subseteq O$. Then $x \in U_x \subseteq f^{-1}(O)$, so Proposition 6.1.3 gives $f^{-1}(O) \in \mathcal{T}_X$.

(*i*) ($\Leftarrow$) Assume $\forall O \in \mathcal{T}_Y,\ f^{-1}(O) \in \mathcal{T}_X$. Let $x_0 \in X$ and $O \in \mathcal{N}_Y(f(x_0))$. Then $U := f^{-1}(O) \in \mathcal{N}_X(x_0)$ and $f(U) \subseteq O$, so $f$ is continuous at $x_0$. Since $x_0$ was arbitrary, $f$ is continuous.

(*ii*) Let $O \in \mathcal{T}_Y$. Then $(f|_A)^{-1}(O) = A \cap f^{-1}(O) \in \mathcal{T}_A$, so $f|_A$ is continuous by (*i*).

(*iii*) Let $W \in \mathcal{T}_Z$. Since $g$ is continuous, $g^{-1}(W) \in \mathcal{T}_Y$.
Since $f$ is continuous, $(g \circ f)^{-1}(W) = f^{-1}(g^{-1}(W)) \in \mathcal{T}_X$. Hence $g \circ f$ is continuous by (*i*).

*Structural Commentary*: Subspaces and composites preserve open preimages by intersection and nesting.
*Key Insight*: Continuity is an inverse-image property. □

**Proposition 6.2.3.** *Continuity Tested on a Subbasis*
*Let $(X, \mathcal{T}_X)$ be a topological space, let $Y$ carry the topology $\mathcal{T}_Y$ generated by a subbasis $\mathcal{S}$, and let $f : X \to Y$. Then $f$ is continuous iff $\forall S \in \mathcal{S},\ f^{-1}(S) \in \mathcal{T}_X$.*

*Proof.*

($\Rightarrow$) If $f$ is continuous and $S \in \mathcal{S}$, then $S \in \mathcal{T}_Y$, so Proposition 6.2.2 gives $f^{-1}(S) \in \mathcal{T}_X$.

($\Leftarrow$) Assume $\forall S \in \mathcal{S},\ f^{-1}(S) \in \mathcal{T}_X$. Let $\mathcal{B}$ be the basis of finite intersections from $\mathcal{S}$, including $Y$ as the empty intersection. Then every $B \in \mathcal{B}$ has $f^{-1}(B) \in \mathcal{T}_X$, since $f^{-1}(\bigcap_{j=1}^{n} S_j) = \bigcap_{j=1}^{n} f^{-1}(S_j)$.
If $O = \bigcup_{\alpha \in A} B_\alpha \in \mathcal{T}_Y$, $B_\alpha \in \mathcal{B}$, then $f^{-1}(O) = \bigcup_{\alpha \in A} f^{-1}(B_\alpha) \in \mathcal{T}_X$. Proposition 6.2.2 gives continuity of $f$.

*Structural Commentary*: Subbasis tests extend through finite intersections and unions.
*Key Insight*: Inverse images preserve the operations generating the topology. □

**Definition 6.2.4.** *Homeomorphism*
Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces.

(i) A *homeomorphism* is a bijection $f : X \to Y$ such that
both $f : (X, \mathcal{T}_X) \to (Y, \mathcal{T}_Y)$ and $f^{-1} : (Y, \mathcal{T}_Y) \to (X, \mathcal{T}_X)$ are continuous.

(ii) The spaces $X$ and $Y$ are *homeomorphic*, written $X \cong_{\text{top}} Y$, iff $\exists$ a homeomorphism $f : X \to Y$.

**Proposition 6.2.5.** *Homeomorphic Spaces Have the Same Open-Set Structure*
*Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, let $f : X \to Y$ be a homeomorphism, and let $E \subseteq X$.*

*(i) $E \in \mathcal{T}_X$ iff $f(E) \in \mathcal{T}_Y$.*
*(ii) $E$ is closed in $X$ iff $f(E)$ is closed in $Y$.*

*Proof.* Since $f$ is a homeomorphism, $f$ is bijective and both $f$ and $f^{-1}$ are continuous.

(*i*) ($\Rightarrow$) If $E \in \mathcal{T}_X$, then $f(E) = (f^{-1})^{-1}(E) \in \mathcal{T}_Y$.

(*i*) ($\Leftarrow$) If $f(E) \in \mathcal{T}_Y$, then $E = f^{-1}(f(E)) \in \mathcal{T}_X$.

(*ii*) ($\Rightarrow$) If $E$ is closed in $X$, then $X \setminus E \in \mathcal{T}_X$. By (*i*), $f(X \setminus E) \in \mathcal{T}_Y$.
Since $f$ is bijective, $f(X \setminus E) = Y \setminus f(E)$, so $f(E)$ is closed in $Y$.

(*ii*) ($\Leftarrow$) If $f(E)$ is closed in $Y$, then $Y \setminus f(E) \in \mathcal{T}_Y$. Since $f$ is bijective, $Y \setminus f(E) = f(X \setminus E)$.
By (*i*), $X \setminus E \in \mathcal{T}_X$, so $E$ is closed in $X$.

*Structural Commentary*: Openness transfers through inverse images of $f$ and $f^{-1}$.
*Key Insight*: Homeomorphisms preserve and reflect open sets. □

**Definition 6.2.6.** *Product Topology*
Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, and let $\{(X_i, \mathcal{T}_i)\}_{i \in I}$ be a family of topological spaces.

(i) The *product topology* on $X \times Y$ is generated by $\{U \times V : U \in \mathcal{T}_X,\ V \in \mathcal{T}_Y\}$.

(ii) The *product topology* on $\prod_{i \in I} X_i$ is generated by $\{\prod_{i \in I} U_i : U_i \in \mathcal{T}_i\ \forall i \in I,\ |\{i \in I : U_i \neq X_i\}| < \infty\}$.

**Proposition 6.2.7.** *Product Topology and Coordinate Maps*
*Let $(Z, \mathcal{T}_Z)$, $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces. Let $\mathcal{T}_{X \times Y}$ be generated by $\mathcal{B} = \{U \times V : U \in \mathcal{T}_X,\ V \in \mathcal{T}_Y\}$. For $f : Z \to X \times Y$, $f$ is continuous iff $\pi_X \circ f$ and $\pi_Y \circ f$ are continuous.*

*Proof.*

($\Rightarrow$) If $f$ is continuous and $U \in \mathcal{T}_X$, then $U \times Y \in \mathcal{B}$ and $(\pi_X \circ f)^{-1}(U) = f^{-1}(U \times Y) \in \mathcal{T}_Z$.
Similarly, $(\pi_Y \circ f)^{-1}(V) = f^{-1}(X \times V) \in \mathcal{T}_Z$ for every $V \in \mathcal{T}_Y$.

($\Leftarrow$) Suppose $\pi_X \circ f$ and $\pi_Y \circ f$ are continuous. For $U \times V \in \mathcal{B}$, $f^{-1}(U \times V) = (\pi_X \circ f)^{-1}(U) \cap (\pi_Y \circ f)^{-1}(V) \in \mathcal{T}_Z$.
If $O \in \mathcal{T}_{X \times Y}$, write $O = \bigcup_{\alpha \in A} B_\alpha$ with $B_\alpha \in \mathcal{B}$; then $f^{-1}(O) = \bigcup_{\alpha \in A} f^{-1}(B_\alpha) \in \mathcal{T}_Z$. Hence $f$ is continuous.

*Structural Commentary*: Basis rectangles pull back to intersections of coordinate preimages.
*Key Insight*: Product continuity is coordinatewise continuity. □

**Proposition 6.2.8.** *Product Convergence Is Pointwise*
*Let $X^\Lambda = \prod_{\lambda\in\Lambda} X_\lambda$ carry the product topology from $\{(X_\lambda, \mathcal{T}_\lambda)\}_{\lambda\in\Lambda}$.*
*For $(x_n) \subseteq X^\Lambda$ and $x \in X^\Lambda$, $x_n \to x$ iff $x_n(\lambda) \to x(\lambda)$ in $X_\lambda$ for every $\lambda \in \Lambda$.*

*Proof.*

($\Rightarrow$) If $x_n \to x$, then each projection $\pi_\lambda : X^\Lambda \to X_\lambda$ is continuous, so $x_n(\lambda) = \pi_\lambda(x_n) \to \pi_\lambda(x) = x(\lambda)$ $\forall \lambda \in \Lambda$.
($\Leftarrow$) Suppose $x_n(\lambda) \to x(\lambda)$ $\forall \lambda \in \Lambda$. Let $W = \prod_{\lambda\in\Lambda} U_\lambda$ be a basic neighborhood of $x$, with $x(\lambda) \in U_\lambda \in \mathcal{T}_\lambda$ and finite $F = \{\lambda \in \Lambda : U_\lambda \neq X_\lambda\}$. Coordinatewise convergence on the finite set $F$ gives $n_0$ such that $x_n(\lambda) \in U_\lambda$ for every $\lambda \in F$ and $n \geq n_0$. Hence $x_n \in W$ eventually, so $x_n \to x$.

*Structural Commentary*: Product neighborhoods impose finitely many coordinate constraints.
*Key Insight*: Coordinate convergence satisfies every finite constraint. □

**Definition 6.2.9.** *Quotient Topology*
Let $q : X \to Y$ be a surjection from a topological space $X$ onto a set $Y$.

(i) The *quotient topology* on $Y$ is $\{V \subseteq Y : q^{-1}(V)$ is open in $X\}$.
(ii) Equivalently, it is the strongest topology on $Y$ for which $q$ is continuous.

**Proposition 6.2.10.** *Quotient Universal Property*
*Let $q : (X, \tau_X) \twoheadrightarrow Y$ and give $Y$ the quotient topology $\tau_Y = \{V \subseteq Y : q^{-1}(V) \in \tau_X\}$.*
*For any map $g : Y \to (Z, \tau_Z)$, $g : (Y, \tau_Y) \to (Z, \tau_Z)$ is continuous $\iff$ $g \circ q : (X, \tau_X) \to (Z, \tau_Z)$ is continuous.*

*Proof.*

($\Rightarrow$) If $g$ is continuous, then $g \circ q$ is continuous by composition.
($\Leftarrow$) Suppose $g \circ q$ is continuous. For $W \in \tau_Z$, $q^{-1}(g^{-1}(W)) = (g \circ q)^{-1}(W) \in \tau_X$.
Thus $g^{-1}(W) \in \tau_Y$ by the quotient topology, so $g$ is continuous.

*Structural Commentary*: Continuity from a quotient is tested by precomposition with the quotient map.
*Key Insight*: Pull back through the quotient map and use the defining open-set test. □

**Definition 6.2.11.** *Weak Topology Induced by Maps*
Let $X$ be a set.

(i) For topologies $\mathcal{T}_1, \mathcal{T}_2$ on $X$, $\mathcal{T}_2$ is *weaker* than $\mathcal{T}_1$ and $\mathcal{T}_1$ is *stronger* than $\mathcal{T}_2$ iff $\mathcal{T}_2 \subseteq \mathcal{T}_1$.
(ii) For topological spaces $\{(X_\alpha, \mathcal{T}_\alpha)\}_{\alpha\in A}$ and maps $f_\alpha : X \to X_\alpha$, the *weak topology induced by* $\{f_\alpha\}_{\alpha\in A}$ is generated by $\mathcal{S} := \{X\} \cup \{f_\alpha^{-1}(O) : \alpha \in A,\ O \in \mathcal{T}_\alpha\}$.

**Proposition 6.2.12.** *Weak Topology Universal Property*
*Let $f_\alpha : X \to (X_\alpha, \mathcal{T}_\alpha)$ for $\alpha \in A$, and let $\mathcal{T}_w$ be generated by $\mathcal{S} := \{X\} \cup \{f_\alpha^{-1}(O) : \alpha \in A,\ O \in \mathcal{T}_\alpha\}$.*
*Then $\mathcal{T}_w$ is the weakest topology on $X$ making every $f_\alpha$ continuous.*

*Proof.* Since $f_\alpha^{-1}(O) \in \mathcal{T}_w$ for all $\alpha \in A$ and $O \in \mathcal{T}_\alpha$, every $f_\alpha : (X, \mathcal{T}_w) \to (X_\alpha, \mathcal{T}_\alpha)$ is continuous.
If a topology $\mathcal{T}'$ on $X$ makes every $f_\alpha$ continuous, then $\mathcal{S} \subseteq \mathcal{T}'$; hence $\mathcal{T}_w \subseteq \mathcal{T}'$.
*Structural Commentary*: Subbasic preimages are the forced open sets.
*Key Insight*: Weakest means contained in every topology passing the tests. □

## 6.3 Topological Compactness and Normality Tools

Compactness in general topology keeps the open-cover principle from metric spaces and pairs especially well with Hausdorff and normal separation. These tools also give useful continuous functions on normal spaces.

**Definition 6.3.1.** *Compactness in Topological Spaces*
Let $X$ be a topological space.

(i) A family $\{O_\alpha\}_{\alpha\in A}$ is an *open cover* of $E \subseteq X$ if every $O_\alpha$ is open and $E \subseteq \bigcup_{\alpha\in A} O_\alpha$.
(ii) The space $X$ is *compact* if every open cover of $X$ has a finite subcover.
(iii) A subset $K \subseteq X$ is *compact* if it is compact with the subspace topology.
Equivalently, every cover of $K$ by open subsets of $X$ has a finite subcover.

**Proposition 6.3.2.** *Finite-Intersection Compactness Test*
*Let $(X, \mathcal{T})$ be a topological space. Then $X$ is compact if and only if every closed family $\{F_\alpha\}_{\alpha\in A}$ with the finite intersection property satisfies $\bigcap_{\alpha\in A} F_\alpha \neq \emptyset$.*

*Proof.*

($\Rightarrow$) If $X$ is compact and closed $\{F_\alpha\}_{\alpha\in A}$ has empty intersection, then $\{X \setminus F_\alpha\}_{\alpha\in A}$ covers $X$. A finite subcover gives finite $J \subseteq A$ with $\bigcap_{\alpha\in J} F_\alpha = \emptyset$, so the family is not FIP.

($\Leftarrow$) Assume every closed FIP family intersects. If an open cover $\{O_\alpha\}_{\alpha\in A}$ has no finite subcover, then $\{X \setminus O_\alpha\}_{\alpha\in A}$ is closed and FIP, so some $x$ lies in all $X \setminus O_\alpha$, contradicting that $\{O_\alpha\}_{\alpha\in A}$ covers $X$.

*Structural Commentary*: Complements exchange covers and intersections.
*Key Insight*: Compactness forbids closed FIP families from missing every point. □

**Proposition 6.3.3.** *Closed Subsets of Compact Spaces Are Compact*
*Let $(X, \mathcal{T})$ be compact and let $K \subseteq X$ be closed. Then $K$ is compact in the subspace topology.*

*Proof.* Let $\{O_\alpha\}_{\alpha\in A}$ cover $K$, with $O_\alpha \in \mathcal{T}$. Since $X \setminus K \in \mathcal{T}$, the family $\{O_\alpha\}_{\alpha\in A} \cup \{X \setminus K\}$ covers $X$. Compactness gives a finite subcover; discarding $X \setminus K$ if present leaves a finite subcover of $K$.
*Structural Commentary*: Add the open complement to cover $X$.
*Key Insight*: Compactness passes to closed subspaces. □

**Proposition 6.3.4.** *Compact Subsets of Hausdorff Spaces Are Closed*
*Let $(X, \mathcal{T})$ be Hausdorff and let $K \subseteq X$ be compact in the subspace topology. Then $K$ is closed in $X$.*

*Proof.* For $y \in X \setminus K$, choose disjoint open $O_x \ni x$ and $U_x \ni y$ for each $x \in K$.
Compactness gives $x_1, \dots, x_n \in K$ with $K \subseteq \bigcup_{j=1}^{n} O_{x_j}$, so $U := \bigcap_{j=1}^{n} U_{x_j}$ is an open neighborhood of $y$ disjoint from $K$. Thus $X \setminus K$ is open and $K$ is closed.
*Structural Commentary*: Compactness turns pointwise Hausdorff separations into a finite one.
*Key Insight*: Every exterior point has an open neighborhood avoiding $K$. □

**Theorem 6.3.5.** *Compact Hausdorff Spaces Are Normal*
*Let $(X, \mathcal{T})$ be compact and Hausdorff. Then $X$ is normal.*

*Proof.* Hausdorff implies $T_1$. Let $A, B \subseteq X$ be disjoint closed sets; then $A, B$ are compact. For each $b \in B$, compact-set separation gives disjoint open $U_b \ni b$ and $V_b \supseteq A$. Choose $b_1, \dots, b_n$ with $B \subseteq \bigcup_{j=1}^{n} U_{b_j}$.

Then $\bigcup_{j=1}^{n} U_{b_j}$ and $\bigcap_{j=1}^{n} V_{b_j}$ are disjoint open neighborhoods of $B$ and $A$, so $X$ is normal.
*Structural Commentary*: Compactness makes finitely many Hausdorff separations suffice.
*Key Insight*: Disjoint closed compact sets have disjoint neighborhoods. □

**Theorem 6.3.6.** *Continuous Images of Compact Spaces Are Compact*
*Let $f : (X, \mathcal{T}_X) \to (Y, \mathcal{T}_Y)$ be continuous. If $X$ is compact, then $f(X)$ is compact in the subspace topology.*

*Proof.* Let $\{V_\alpha\}_{\alpha\in A}$ be an open cover of $f(X)$ by subsets open in $Y$. Then $\{f^{-1}(V_\alpha)\}_{\alpha\in A}$ covers $X$ and each $f^{-1}(V_\alpha) \in \mathcal{T}_X$. Compactness gives finite $J \subseteq A$ with $X = \bigcup_{\alpha\in J} f^{-1}(V_\alpha)$, hence $f(X) \subseteq \bigcup_{\alpha\in J} V_\alpha$.
*Structural Commentary*: Pull the cover back to the compact domain.
*Key Insight*: Finite subcovers descend through the image. □

**Corollary 6.3.7.** *Extreme Values on Compact Spaces*
*If $X$ is compact and $f : X \to \mathbb{R}$ is continuous, then $f$ attains its minimum and maximum.*

*Proof.* By Theorem 6.3.6, $f(X)$ is compact in $\mathbb{R}$. By Heine–Borel, Theorem 4.2.5, $f(X)$ is closed and bounded, so $\inf f(X), \sup f(X) \in f(X)$. Choose preimages $x_{\min}, x_{\max} \in X$.
*Structural Commentary*: Compact images in $\mathbb{R}$ are closed and bounded.
*Key Insight*: Closed bounded images contain their endpoints. □

**Theorem 6.3.8.** *Compact-to-Hausdorff Bijection Criterion*
*Let $f : (X, \mathcal{T}_X) \to (Y, \mathcal{T}_Y)$ be continuous bijection. If $X$ is compact, $Y$ is Hausdorff, then $f$ is a homeomorphism.*

*Proof.* It remains to prove continuity of $f^{-1} : Y \to X$. If $F \subseteq X$ is closed, then $F$ is compact by Proposition 6.3.3, so $f(F)$ is compact by Theorem 6.3.6. Since $Y$ is Hausdorff, $f(F)$ is closed by Proposition 6.3.4.
Thus $(f^{-1})^{-1}(F) = f(F)$ is closed in $Y$, so $f^{-1}$ is continuous.
*Structural Commentary*: Closed sets map compact, then closed.
*Key Insight*: Compact-to-Hausdorff bijections have continuous inverses. □

**Proposition 6.3.9.** *Compact Hausdorff Topologies Are Rigid*
*Let $X$ be a set and let $\mathcal{T}, \mathcal{T}_1, \mathcal{T}_2$ be topologies on $X$.*

*(i) If $(X, \mathcal{T})$ is compact and $\mathcal{T}_1 \subseteq \mathcal{T}$, then $(X, \mathcal{T}_1)$ is compact.*
*(ii) If $(X, \mathcal{T})$ is Hausdorff and $\mathcal{T} \subseteq \mathcal{T}_2$, then $(X, \mathcal{T}_2)$ is Hausdorff.*
*(iii) If $(X, \mathcal{T})$ is compact Hausdorff, then no strictly weaker topology is Hausdorff and no strictly stronger topology is compact.*

*Proof.*

(*i*) Every $\mathcal{T}_1$-open cover is a $\mathcal{T}$-open cover, so compactness of $(X, \mathcal{T})$ gives a finite subcover.
(*ii*) If $x \neq y$, choose disjoint $\mathcal{T}$-open neighborhoods of $x$ and $y$; they are also $\mathcal{T}_2$-open.
(*iii*) If $\mathcal{S} \subseteq \mathcal{T}$ is Hausdorff, Theorem 6.3.8 applied to $\mathrm{id} : (X, \mathcal{T}) \to (X, \mathcal{S})$ gives $\mathcal{T} \subseteq \mathcal{S}$, hence $\mathcal{S} = \mathcal{T}$.
If $\mathcal{T} \subseteq \mathcal{S}$ and $(X, \mathcal{S})$ is compact, apply it to $\mathrm{id} : (X, \mathcal{S}) \to (X, \mathcal{T})$ to get $\mathcal{S} \subseteq \mathcal{T}$, hence $\mathcal{S} = \mathcal{T}$.

*Structural Commentary*: Weakening preserves covers; strengthening preserves separations.
*Key Insight*: Compact Hausdorff topologies cannot be enlarged or shrunk compatibly. □

**Lemma 6.3.10.** *Maximal Finite-Intersection Families*
*Let $X$ be a set and let $\mathcal{A} \subseteq \mathcal{P}(X)$ have the finite intersection property. Then $\exists \mathcal{M} \subseteq \mathcal{P}(X)$ such that $\mathcal{A} \subseteq \mathcal{M}$, $\mathcal{M}$ has the finite intersection property, and no proper extension of $\mathcal{M}$ has it.*

*Proof.* Let $\mathfrak{P} = \{\mathcal{B} \subseteq \mathcal{P}(X) : \mathcal{A} \subseteq \mathcal{B}$ and $\mathcal{B}$ has FIP$\}$, ordered by inclusion. If $\mathfrak{C} \subseteq \mathfrak{P}$ is a chain, then $\bigcup \mathfrak{C}$ has FIP: each finite subfamily lies in one $\mathcal{B} \in \mathfrak{C}$. Thus Zorn's lemma, Principle 6.5.3, gives a maximal $\mathcal{M} \in \mathfrak{P}$.
*Structural Commentary*: Chain unions preserve FIP because tests are finite.
*Key Insight*: Zorn extends any FIP family to a maximal one. □

**Lemma 6.3.11.** *Closure Properties of Maximal FIP Families*
*Let $X$ be a set and let $\mathcal{M} \subseteq \mathcal{P}(X)$ be maximal with respect to the finite intersection property.*

*(i) If $M_1, \ldots, M_n \in \mathcal{M}$, then $\bigcap_{j=1}^{n} M_j \in \mathcal{M}$.*
*(ii) If $C \subseteq X$ satisfies $C \cap \bigcap_{j=1}^{n} M_j \neq \emptyset$ for all $M_1, \ldots, M_n \in \mathcal{M}$, then $C \in \mathcal{M}$.*

*Proof.*

(*i*) For $N = \bigcap_{j=1}^{n} M_j$, the family $\mathcal{M} \cup \{N\}$ still has the finite intersection property, so maximality gives $N \in \mathcal{M}$.
(*ii*) The hypothesis says $\mathcal{M} \cup \{C\}$ has the finite intersection property, so maximality gives $C \in \mathcal{M}$.

*Structural Commentary*: Maximality absorbs every FIP-compatible set.
*Key Insight*: Finite intersections are already forced inside $\mathcal{M}$. □

**Theorem 6.3.12.** *Tychonoff Product Theorem*
*The product $X = \prod_{\lambda \in \Lambda} X_\lambda$ of compact topological spaces is compact in the product topology.*

*Proof.* Let $\mathcal{F}$ be a closed FIP family in $X$, and extend it to maximal FIP $\mathcal{M}$ by Lemma 6.3.10.
For each $\lambda$, compactness of $X_\lambda$ gives $x_\lambda \in \bigcap_{M \in \mathcal{M}} \overline{\pi_\lambda(M)}$; put $x = (x_\lambda)_{\lambda \in \Lambda}$.
If $N = \pi_\lambda^{-1}(O)$ is subbasic at $x$ and $P = \bigcap_{j=1}^{n} M_j$ with $M_j \in \mathcal{M}$, then $P \in \mathcal{M}$ and $O \cap \pi_\lambda(P) \neq \emptyset$, so $N \cap P \neq \emptyset$.
Lemma 6.3.11 gives $N \in \mathcal{M}$, hence every basic neighborhood of $x$ lies in $\mathcal{M}$.
Thus every basic neighborhood of $x$ meets each $F \in \mathcal{F}$, so $x \in \overline{F} = F$.
Hence $\bigcap_{F \in \mathcal{F}} F \neq \emptyset$, and Proposition 6.3.2 gives compactness.
*Structural Commentary*: Coordinate compactness selects a candidate point.
*Key Insight*: Maximal FIP forces all basic neighborhoods of that point to remain compatible. □

**Definition 6.3.13.** *Compact-Open Topology*
Let $X, Y$ be topological spaces and let $Y^X = \{f : X \to Y\}$. The *compact-open topology* on $Y^X$ is generated by the subbasis $\mathcal{U}_{K,O} := \{f \in Y^X : f(K) \subseteq O\}$, where $K \subseteq X$ is compact and $O \subseteq Y$ is open.
*Structural Commentary*: Subbasic opens prescribe uniform containment on compact subsets.

**Remark 6.3.14.** *Compact-Open Versus Product Topology*
On $Y^X = \prod_{x \in X} Y$, the product topology tests finitely many coordinates $f(x)$.
The compact-open topology tests compact sets at once through $\mathcal{U}_{K,O} = \{f \in Y^X : f(K) \subseteq O\}$
For metric $Y$, this encodes uniform control on compact subsets.

**Definition 6.3.15.** *Normally Ascending Families*
Let $X$ be a topological space and let $\Lambda \subseteq \mathbb{R}$.
A family of open sets $\{O_\lambda\}_{\lambda\in\Lambda}$ is *normally ascending* iff $\overline{O_{\lambda_1}} \subseteq O_{\lambda_2}$ for all $\lambda_1, \lambda_2 \in \Lambda$ with $\lambda_1 < \lambda_2$.
*Key Insight*: Later open sets contain the closures of earlier ones.

**Lemma 6.3.16.** *Normally Ascending Families Define Continuous Functions*
*Let $X$ be a topological space, let $\Lambda \subseteq (a,b)$ be dense, and let $\{O_\lambda\}_{\lambda\in\Lambda}$ be normally ascending. Define $g : X \to [a,b]$ by $g(x) = \inf\{\lambda \in \Lambda : x \in O_\lambda\}$, with $\inf \emptyset = b$. Then $g$ is continuous.*

*Proof.* For $c \in (a,b)$, $\{g < c\} = \bigcup_{\lambda<c} O_\lambda$ and $\{g > c\} = \bigcup_{\lambda>c}(X \setminus \overline{O_\lambda})$. For the second identity, $x \notin \overline{O_\lambda}$ with $\lambda > c$ gives $g(x) \geq \lambda > c$; conversely, if $g(x) > c$, choose $\mu, \lambda \in \Lambda$ with $c < \mu < \lambda < g(x)$, so $x \notin O_\lambda \supseteq \overline{O_\mu}$. Both threshold sets are open, hence $g$ is continuous.
*Structural Commentary*: Threshold preimages reduce to unions of nested open or closed-complement sets.
*Key Insight*: Normal ascent makes upper thresholds open. □

**Lemma 6.3.17.** *Dyadic Neighborhood Ladders in Normal Spaces*
*Let $X$ be normal, let $F \subseteq X$ be closed, and let $U$ be an open neighborhood of $F$. For any open interval $(a,b)$ there are dense $\Lambda \subseteq (a,b)$ and normally ascending open sets $\{O_\lambda\}_{\lambda\in\Lambda}$ with $F \subseteq O_\lambda \subseteq \overline{O_\lambda} \subseteq U$ for all $\lambda \in \Lambda$.*

*Proof.* By affine reparametrization, take $(a,b) = (0,1)$ and $\Lambda = \{m2^{-n} : 0 < m < 2^n,\ n \in \mathbb{N}\}$.
Choose $O_{1/2}$ with $F \subseteq O_{1/2} \subseteq \overline{O_{1/2}} \subseteq U$ by Proposition 6.1.15.
Inductively, for each new dyadic $p < r < q$, choose $O_r$ with $\overline{O_p} \subseteq O_r \subseteq \overline{O_r} \subseteq O_q$, interprete endpoints as $F$, $U$
This preserves normal ascent and containment in $U$.
*Structural Commentary*: Normality inserts closed-neighborhood buffers.
*Key Insight*: Dyadic induction builds the whole ascending ladder. □

**Theorem 6.3.18.** *Urysohn's Lemma*
*Let $X$ be normal, let $A, B \subseteq X$ be disjoint closed sets, and let $a < b$ in $\mathbb{R}$.*
*Then $\exists$ continuous $f : X \to [a,b]$ such that $f|_A = a$ and $f|_B = b$.*

*Proof.* Apply Lemma 6.3.17 to $F = A$ and $U = X \setminus B$. Lemma 6.3.16 applied to the resulting ladder gives continuous $f : X \to [a,b]$. Since $A \subseteq O_\lambda$ for all $\lambda$ and $B \cap \bigcup_\lambda O_\lambda = \emptyset$, the definition gives $f|_A = a$ and $f|_B = b$.
*Structural Commentary*: A nested open ladder becomes a continuous separator.
*Key Insight*: Normality separates closed sets by a real-valued function. □

**Theorem 6.3.19.** *Tietze Extension Theorem, Bounded Form*
*Let $X$ be normal, let $F \subseteq X$ be closed, let $a < b$, and let $f : F \to [a,b]$ be continuous.*
*Then $\exists$ continuous $g : X \to [a,b]$ with $g|_F = f$.*

*Proof.* After an affine change of range, assume $[a,b] = [-1,1]$. The Urysohn step is: if $|h| \leq A$ on $F$, then some continuous $u : X \to [-A/3, A/3]$ satisfies $|h - u| \leq 2A/3$ on $F$. Starting with $s_0 = 0$, apply this to $h = f - s_{n-1}$ to get $u_n$ with $|u_n| \leq \frac{1}{3}(2/3)^{n-1}$ on $X$ and $|f - s_n| \leq (2/3)^n$ on $F$, where $s_n = s_{n-1} + u_n$.
Then $\sum_n u_n$ converges uniformly to continuous $g : X \to [-1,1]$ and $g|_F = f$. Undo the affine change of range.
*Structural Commentary*: Urysohn corrections shrink the residual geometrically.
*Key Insight*: Uniform convergence turns approximate extensions into an exact one. □

## 6.4 Sequences, Countability, and Metrization

Countability axioms control when topological questions can be reduced to sequences and countable tests. They also separate metric phenomena from genuinely non-metric topologies.

**Definition 6.4.1.** *Sequence Convergence and Sequential Compactness*
Let $X$ be a topological space, and let $\mathcal{N}_X(x)$ denote the neighborhoods of $x \in X$.

(i) $(x_n)$ *converges* to $x$, written $x_n \to x$, iff $\forall U \in \mathcal{N}_X(x)\ \exists N \in \mathbb{N}$ such that $n \geq N \Rightarrow x_n \in U$.
(ii) $X$ is *sequentially compact* iff every sequence $(x_n) \subseteq X$ has a convergent subsequence $x_{n_k} \to x \in X$.

*Key Insight*: Sequential compactness is compactness tested by subsequences.

**Proposition 6.4.2.** *Hausdorff Sequence Limits Are Unique*
*Let $X$ be a topological space and let $(x_n) \subseteq X$.*

*(i) If $X$ is Hausdorff, then $(x_n)$ has at most one limit.*
*(ii) If $X$ has the trivial topology $\{\emptyset, X\}$, then $x_n \to x$ for every $x \in X$.*

*Proof.*

(*i*) If $x_n \to x$ and $x_n \to y$ with $x \neq y$, choose disjoint neighborhoods $U \in \mathcal{N}_X(x)$ and $V \in \mathcal{N}_X(y)$. Eventually $x_n \in U$ and $x_n \in V$, contradicting $U \cap V = \emptyset$.

(*ii*) In the trivial topology, $\mathcal{N}_X(x) = \{X\}$ for every $x \in X$, and every sequence is eventually in $X$.

*Structural Commentary*: Hausdorff neighborhoods separate competing limits.
*Key Insight*: A sequence tail cannot lie in two disjoint neighborhoods. □

**Definition 6.4.3.** *Countability Axioms*
Let $(X, \tau)$ be a topological space.

(i) $X$ is *first countable* iff $\forall x \in X\ \exists$ countable $\mathcal{B}_x \subseteq \mathcal{N}_X(x)$ such that every $U \in \mathcal{N}_X(x)$ contains some $B \in \mathcal{B}_x$.

(ii) $X$ is *second countable* iff $\tau$ has a countable basis.

(iii) $X$ is *separable* iff $\exists$ countable $D \subseteq X$ with $\overline{D} = X$, equivalently every nonempty $U \in \tau$ meets $D$.

*Key Insight*: Countability turns topology into countable local, global, or dense tests.

**Example 6.4.4.** *Metric Spaces Are First Countable*
Every metric space is first countable: $\forall x$, balls $\{B_{1/n}(x) : n \geq 1\}$ form a countable neighborhood basis at $x$.

**Proposition 6.4.5.** *Second Countable Implies First Countable*
*Let $(X, \tau)$ be second countable. Then $X$ is first countable.*

*Proof.* Let $\mathcal{B}$ be a countable basis for $\tau$. For $x \in X$, set $\mathcal{B}_x := \{B \in \mathcal{B} : x \in B\}$.
Then $\mathcal{B}_x$ is countable, and for every $U \in \mathcal{N}_X(x)$, some $B \in \mathcal{B}_x$ satisfies $x \in B \subseteq U$.
*Structural Commentary*: Restrict the global basis to sets containing $x$.
*Key Insight*: Countable global tests give countable local tests. □

**Proposition 6.4.6.** *First Countable Sequential Closure*
*Let $X$ be first countable, let $E \subseteq X$, and let $x \in X$.*

*(i) $x \in \overline{E}$ iff $\exists (x_n) \subseteq E$ with $x_n \to x$.*

*(ii) $E$ is closed iff every convergent sequence in $E$ has all its limits in $E$.*

*Proof.*

(*i*) ($\Rightarrow$) Let $\{B_n : n \geq 1\}$ be a countable neighborhood basis at $x$.
Since $x \in \overline{E}$ and $\bigcap_{j=1}^{n} B_j \in \mathcal{N}_X(x)$, choose $x_n \in E \cap \bigcap_{j=1}^{n} B_j$. Then $x_n \to x$.

(*i*) ($\Leftarrow$) If $(x_n) \subseteq E$ and $x_n \to x$, then every neighborhood of $x$ contains some $x_n \in E$, so $x \in \overline{E}$.

(*ii*) ($\Rightarrow$) If $E$ is closed and $(x_n) \subseteq E$ with $x_n \to x$, then $x \in \overline{E} = E$ by (*i*).

(*ii*) ($\Leftarrow$) If every convergent sequence in $E$ has all its limits in $E$, then (*i*) gives $\overline{E} \subseteq E$. Hence $E = \overline{E}$, QED.

*Structural Commentary*: Countable local bases produce one testing sequence.
*Key Insight*: In first countable spaces, closure is sequential closure. □

**Proposition 6.4.7.** *Second Countable Consequences*
*If $(X, \tau)$ is second countable, then there is a countable $D \subseteq X$ with $\overline{D} = X$.*
*Moreover, every subspace of a second countable space is second countable.*

*Proof.* Let $\mathcal{B} = \{B_n : n \in I\}$ be a countable basis, with $I \subseteq \mathbb{N}$.
For each $n \in I$ with $B_n \neq \emptyset$, choose $x_n \in B_n$ and set $D = \{x_n : n \in I,\ B_n \neq \emptyset\}$. Then $D$ is countable.
If $U \in \tau \setminus \{\emptyset\}$, choose $x \in U$; some $B_n \in \mathcal{B}$ satisfies $x \in B_n \subseteq U$, so $x_n \in D \cap U$. Hence $\overline{D} = X$.
If $Y \subseteq X$, then $\{B_n \cap Y : n \in I\}$ is a countable basis for the subspace topology on $Y$.
*Structural Commentary*: $\mathcal{B}$ countable gives countable $D$ meeting every $U \in \tau \setminus \{\emptyset\}$.
*Key Insight*: $D$ is dense iff $D \cap U \neq \emptyset$ for every $U \in \tau \setminus \{\emptyset\}$. □

**Proposition 6.4.8.** *Separable Metric Spaces and Countable Bases*
*Let $(X, d)$ be a metric space, and let $\tau_d$ be its metric topology. The following are equivalent:*

*(i) $\exists$ countable $D \subseteq X$ such that $\overline{D}^{\,X} = X$.*

*(ii) $\exists$ countable $\mathcal{B} \subseteq \tau_d$ such that $\mathcal{B}$ is a basis for $\tau_d$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) Let $D \subseteq X$ be countable with $\overline{D}^{\,X} = X$, and set $\mathcal{B} := \{B_q(x) : x \in D,\ q \in \mathbb{Q},\ q > 0\}$.
Then $\mathcal{B}$ is countable. For $U \in \tau_d$ and $y \in U$, choose $r > 0$ with $B_r(y) \subseteq U$, then choose $q \in \mathbb{Q}$ with $0 < q < r/2$, $x \in D \cap B_q(y)$. Then $y \in B_q(x) \subseteq B_r(y) \subseteq U$, since $z \in B_q(x)$ implies $d(z, y) < 2q < r$.
QED.

(*ii*) $\Rightarrow$ (*i*) Let $\mathcal{B} = \{U_n : n \in I\}$ be a countable basis for $\tau_d$, with $I \subseteq \mathbb{N}$.
Choose $x_n \in U_n$ for each nonempty $U_n$, and set $D := \{x_n : U_n \neq \emptyset\}$. Then $D$ is countable.
If $\emptyset \neq U \in \tau_d$, choose $y \in U$ and $n \in I$ with $y \in U_n \subseteq U$; then $x_n \in D \cap U$.
Hence every nonempty open set meets $D$, so $\overline{D}^X = X$.

*Structural Commentary*: In metric spaces, one countable dense set produces all rational-radius local tests.
*Key Insight*: Separability and second countability coincide for metric spaces. □

**Proposition 6.4.9.** *Subspaces of Separable Metric Spaces*
*Let $(X, d)$ be a separable metric space and let $E \subseteq X$ with the subspace metric $d|_{E\times E}$. Then $E$ is separable.*

*Proof.* By Proposition 6.4.8, the metric topology $\tau_d$ has a countable basis $\mathcal{B}$. Set $\mathcal{B}_E := \{B \cap E : B \in \mathcal{B}\}$. Then $\mathcal{B}_E$ is countable. If $U$ is open in $E$ and $x \in U$, then $U = E \cap G$ for some $G \in \tau_d$; choose $B \in \mathcal{B}$ with $x \in B \subseteq G$. Hence $x \in B \cap E \subseteq U$, so $\mathcal{B}_E$ is a countable basis for $E$. By Proposition 6.4.8, $E$ is separable.
*Structural Commentary*: A countable ambient basis restricts to a countable subspace basis.
*Key Insight*: Metric separability passes to all subspaces through countable bases. □

**Corollary 6.4.10.** *Euclidean Subsets Are Separable*
*Let $k \in \mathbb{N}$ and let $E \subseteq \mathbb{R}^k$ have the subspace Euclidean metric. Then $E$ is separable.*

*Proof.* The set $\mathbb{Q}^k$ is countable and dense in $\mathbb{R}^k$.
For $x \in \mathbb{R}^k$ and $r > 0$, choose $q = (q_1, \dots, q_k) \in \mathbb{Q}^k$ with $|x_i - q_i| < r/\sqrt{k}$ for each $i$, so $\|x - q\|_2 < r$.
Thus $\mathbb{R}^k$ is separable, and Proposition 6.4.9 gives that $E$ is separable.
*Structural Commentary*: Rational grids are countable dense nets.
*Key Insight*: Euclidean separability passes to subspaces. □

**Example 6.4.11.** *Uncountable Cofinite Topology*
Let $X$ be uncountable and let $\tau := \{\emptyset\} \cup \{U \subseteq X : X \setminus U \text{ is finite}\}$.

(i) $\tau$ is a topology on $X$.
(ii) $(X, \tau)$ is $T_1$.
(iii) $(X, \tau)$ is not first countable.

*Proof.*

(*i*) $\emptyset, X \in \tau$. Finite intersections and arbitrary unions of cofinite sets are cofinite, so $\tau$ is a topology.
(*ii*) For $x \in X$, $X \setminus \{x\} \in \tau$, so $\{x\}$ is closed. Hence $(X, \tau)$ is $T_1$.
(*iii*) Fix $x \in X$. If $\{B_n : n \geq 1\}$ were a countable neighborhood basis at $x$, then $C := \{x\} \cup \bigcup_n (X \setminus B_n)$ is countable. Choose $y \in X \setminus C$. Then $U := X \setminus \{y\} \in \mathcal{N}_X(x)$, but $y \in B_n$ for all $n$, so no $B_n \subseteq U$.

*Key Insight*: Cofinite local bases cannot be countable. □

**Definition 6.4.12.** *Lindelöf and Countably Compact Spaces*
Let $(X, \tau)$ be a topological space.

(i) $X$ is *Lindelöf* iff every open cover $\{O_\alpha\}_{\alpha\in A} \subseteq \tau$ has a countable subcover $\{O_\alpha\}_{\alpha\in J}$, $J \subseteq A$.
(ii) $X$ is *countably compact* iff every countable open cover $\{O_n\}_{n\in\mathbb{N}} \subseteq \tau$ has a finite subcover $\{O_n\}_{n\in J}$, $J \subseteq \mathbb{N}$.

*Key Insight*: Covers reduce to countable or finite subcovers.

**Proposition 6.4.13.** *Frechet Intersection Test for Countable Compactness*
*Let $X$ be a topological space. Then $X$ is countably compact iff every descending sequence $F_1 \supseteq F_2 \supseteq \cdots$ of nonempty closed subsets of $X$ satisfies $\bigcap_{n\geq 1} F_n \neq \emptyset$.*

*Proof.*

($\Rightarrow$) Assume $X$ is countably compact. If $\bigcap_{n\geq 1} F_n = \emptyset$, then $\{X \setminus F_n : n \geq 1\}$ is a countable open cover.
A finite subcover exists; since $F_n$ descends, some $X \setminus F_N$ covers $X$, contradicting $F_N \neq \emptyset$.
($\Leftarrow$) Let $\{O_n : n \geq 1\}$ be a countable open cover with no finite subcover.
Then $F_n := X \setminus \bigcup_{j=1}^{n} O_j$ is nonempty, closed, and descending, while $\bigcap_{n\geq 1} F_n = \emptyset$, a contradiction.

*Structural Commentary*: Complements turn countable covers into descending closed sets.
*Key Insight*: Countable compactness is nested closed intersection. □

**Theorem 6.4.14.** *Dini's Theorem on Countably Compact Spaces*
*Let $X$ be countably compact, and let $f_n : X \to \mathbb{R}$ be continuous with $f_n(x) \downarrow 0$ for every $x \in X$. Then $f_n \to 0$ uniformly: $\forall \varepsilon > 0\ \exists N$ such that $0 \leq f_n(x) < \varepsilon$ for all $x \in X$ and $n \geq N$.*

*Proof.* Fix $\varepsilon > 0$ and set $O_n := \{x \in X : f_n(x) < \varepsilon\}$. Then $O_n$ is open, $O_n \subseteq O_{n+1}$, and $\bigcup_{n\geq 1} O_n = X$.
Countable compactness gives $X = \bigcup_{j=1}^{m} O_{n_j}$; with $N := \max_j n_j$, increasingness gives $O_N = X$.
Hence $0 \leq f_n(x) \leq f_N(x) < \varepsilon$ for all $x \in X$ and $n \geq N$.
*Structural Commentary*: Increasing sublevel covers reduce finite subcovers to one index.
*Key Insight*: Monotone pointwise convergence becomes uniform convergence. □

**Proposition 6.4.15.** *Second Countable Implies Lindelöf*
*Let $(X, \tau)$ be second countable. Then $X$ is Lindelöf: every open cover of $X$ has a countable subcover.*

*Proof.* Let $\mathcal{B} = \{B_n : n \in I\}$ be a countable basis, and let $\mathcal{U} \subseteq \tau$ cover $X$. For $I_0 := \{n \in I : \exists U \in \mathcal{U},\ B_n \subseteq U\}$, choose $U_n \in \mathcal{U}$ with $B_n \subseteq U_n$ for each $n \in I_0$. The family $\{U_n : n \in I_0\}$ is countable.
If $x \in X$, choose $U \in \mathcal{U}$ with $x \in U$ and $B_n \in \mathcal{B}$ with $x \in B_n \subseteq U$; then $n \in I_0$ and $x \in U_n$.
*Structural Commentary*: Basis elements refine arbitrary open covers.
*Key Insight*: A countable basis yields a countable subcover. □

**Proposition 6.4.16.** *Second Countable Countable Compactness Is Compactness*
*Let $X$ be second countable. Then $X$ is compact iff $X$ is countably compact.*

*Proof.*
(⇒) If $\{O_n\}_{n\in\mathbb{N}}$ covers $X$, compactness gives finite $J \subseteq \mathbb{N}$ with $X = \bigcup_{n\in J} O_n$.
(⇐) Let $\mathcal{U} = \{O_\alpha\}_{\alpha\in A}$ cover $X$. By Proposition 6.4.15, $\exists$ countable $J \subseteq A$ with $X = \bigcup_{\alpha\in J} O_\alpha$.
Countable compactness gives finite $J_0 \subseteq J$ with $X = \bigcup_{\alpha\in J_0} O_\alpha$.
*Structural Commentary*: Second countability reduces covers to countable subcovers.
*Key Insight*: Countable compactness then makes the subcover finite. □

**Theorem 6.4.17.** *Second Countable Compactness and Sequential Compactness*
*Let $X$ be a second countable topological space. Then $X$ is compact iff $X$ is sequentially compact.*

*Proof.*
(⇒) Assume $X$ is compact. Let $(x_n) \subseteq X$ and $F_n := \overline{\{x_k : k \geq n\}}$. Since $F_1 \supseteq F_2 \supseteq \cdots$ are nonempty closed, compactness gives $x \in \bigcap_{n\geq 1} F_n$. By first countability, choose a decreasing neighborhood basis $(B_j)$ at $x$. With $n_0 = 0$, choose $n_j > n_{j-1}$ and $x_{n_j} \in B_j$ from $x \in F_{n_{j-1}+1}$. Then $x_{n_j} \to x$.
(⇐) Assume $X$ is sequentially compact. By Proposition 6.4.16, it is enough to prove countable compactness. If $\{O_n\}_{n\geq 1}$ is a countable open cover with no finite subcover, choose $x_n \in X \setminus \bigcup_{j=1}^{n} O_j$. Sequential compactness gives $x_{n_k} \to x \in O_N$ for some $N$. For $k$ large, $n_k \geq N$ and $x_{n_k} \in O_N$, contradicting $x_{n_k} \notin O_N$.
*Structural Commentary*: Closed tails give a cluster point; local countability extracts a subsequence.
*Key Insight*: Countable covers and bases make compactness sequential. □

**Definition 6.4.18.** *Metrizable Space*
A topological space is *metrizable* if its topology is induced by some metric.

**Definition 6.4.19.** *Sorgenfrey Line*
Let $\mathcal{B}_S := \{[a, b) : a < b\}$ on $\mathbb{R}$. The *Sorgenfrey line* is $\mathbb{R}_S := (\mathbb{R}, \tau_S)$, where $\tau_S$ is the topology generated by $\mathcal{B}_S$.

**Example 6.4.20.** *Sorgenfrey Line Countability Tests*
The Sorgenfrey line $\mathbb{R}_S$ is first countable and separable, but not second countable; hence it is not metrizable.

*Proof.* For $x \in [a, b) \cap [c, d)$, choose $e$ with $x < e \leq \min\{b, d\}$; then $x \in [x, e) \subseteq [a, b) \cap [c, d)$, so $\mathcal{B}_S$ is a basis.
Also $\{[x, x + 1/n) : n \geq 1\} \subseteq \mathcal{N}_{\mathbb{R}_S}(x)$ is a local basis, and $\mathbb{Q} \cap [a, b) \neq \emptyset$ for every $[a, b) \neq \emptyset$.
Hence $\mathbb{R}_S$ is first countable and separable.
If $\mathcal{C}$ is a basis for $\mathbb{R}_S$, choose $C_x \in \mathcal{C}$ with $x \in C_x \subseteq [x, x + 1)$. For $x < y$, $x \in C_x \setminus C_y$, so $C_x \neq C_y$.
Thus $x \mapsto C_x$ is injective, so no countable $\mathcal{C}$ exists. Hence $\mathbb{R}_S$ is not second countable.
If $\mathbb{R}_S$ were metrizable, separability and Proposition 6.4.8 would force second countability.
*Structural Commentary*: Half-open neighborhoods encode left endpoints.
*Key Insight*: Local countability does not imply one countable global basis. □

**Definition 6.4.21.** *Moore Plane*
Let $\mathbb{M} := \{(x, y) \in \mathbb{R}^2 : y \geq 0\}$ and $L := \mathbb{R} \times \{0\}$. The *Moore plane* is $\mathbb{M}$ with basis consisting of:
(i) $B_{\text{euc}}(p, r) \subseteq \mathbb{M}$ for $p \in \mathbb{M} \setminus L$ and $r > 0$;

(ii) $N(x,r) := \{(x,0)\} \cup B_{\text{euc}}((x,r),r)$ for $(x,0) \in L$ and $r > 0$.

**Example 6.4.22.** *Moore Plane Countability Tests*
The Moore plane $\mathbb{M}$ is Hausdorff and separable, but not second countable; hence it is not metrizable.

*Proof.* The basis and Hausdorff axioms follow by shrinking Euclidean disks and tangent neighborhoods $N(x,r)$. Let $D := \mathbb{Q} \times (\mathbb{Q} \cap (0,\infty))$; then every basic neighborhood meets $D$, so $\overline{D} = \mathbb{M}$. Thus $\mathbb{M}$ is separable.
For $(x,0) \in L$, $N(x,r) \cap L = \{(x,0)\}$, so $L$ is an uncountable discrete subspace and hence not separable. If $\mathbb{M}$ were second countable, then $L$ would be second countable and separable by Proposition 6.4.7, a contradiction. If $\mathbb{M}$ were metrizable, separability and Proposition 6.4.8 would force second countability.
*Structural Commentary*: Dense rational interior; discrete uncountable boundary.
*Key Insight*: The boundary is an uncountable discrete obstruction. □




Figure 152: A Moore-plane boundary neighborhood is a tangent disk together with its tangent boundary point.

**Theorem 6.4.23.** *Urysohn Metrization Theorem*
*Let $(X,\tau)$ be second countable. Then $\tau$ is induced by a metric iff $(X,\tau)$ is normal.*

*Proof.*
($\Rightarrow$) Let $d$ induce $\tau$. Then $X$ is $T_1$, since $X \setminus \{x\} = \bigcup_{y \neq x} B_d\left(y, \frac{d(x,y)}{2}\right)$. For disjoint nonempty closed $F, G \subseteq X$, set $u(x) := d(x,F)$ and $v(x) := d(x,G)$. Since $u, v$ are continuous, $U := \{u < v\}$ and $V := \{v < u\}$ are disjoint open sets with $F \subseteq U$ and $G \subseteq V$. Empty cases are trivial, so $(X,\tau)$ is normal.
($\Leftarrow$) Assume $(X,\tau)$ is normal. Let $\{U_n : n \in \mathbb{N}\}$ be a basis and $A := \{(n,m) \in \mathbb{N}^2 : \overline{U_n} \subseteq U_m\}$.
For $(n,m) \in A$, choose $f_{n,m} : X \to [0,1]$ by Theorem 6.3.18, with $f_{n,m} = 0$ on $\overline{U_n}$, $f_{n,m} = 1$ on $X \setminus U_m$.
Set $\rho(x,y) := \sum_{(n,m)\in A} \frac{1}{2^{n+m}} |f_{n,m}(x) - f_{n,m}(y)|$, a $\tau$-continuous pseudometric by uniform convergence.
For every $G \in \tau$ and $x \in G$, Proposition 6.1.15 and the basis give $n, m$ with $x \in U_n \subseteq \overline{U_n} \subseteq U_m \subseteq G$.
If $y \notin U_m$, then $\rho(x,y) \geq \frac{1}{2^{n+m}}$, so $B_\rho\left(x, \frac{1}{2^{n+m+1}}\right) \subseteq U_m \subseteq G$.
Taking $G = X \setminus \{y\}$ shows $\rho(x,y) > 0$ for $x \neq y$. Hence $\rho$ is a metric and its topology is $\tau$.
*Structural Commentary*: Urysohn functions separate nested basis pairs.
*Key Insight*: Countably many separators recover the topology as a metric. □

## 6.5 Topological Connectedness and Paths

Connectedness is an open-set obstruction to splitting a space into two visible pieces. In general topology it is most naturally phrased by separations and then transported through continuous maps.

**Definition 6.5.1.** *Connected Topological Spaces*
Let $(X,\tau)$ be a topological space, and let $E \subseteq X$.
(i) A *separation* of $X$ is a pair $(U,V)$ with $U, V \in \tau$, $U, V \neq \emptyset$, $U \cap V = \emptyset$, and $X = U \cup V$.
(ii) $X$ is *connected* iff no separation of $X$ exists.
(iii) $E$ is *connected* iff $(E, \tau_E)$ is connected, where $\tau_E := \{E \cap U : U \in \tau\}$.
Equivalently, $E$ is disconnected iff $\exists U, V \in \tau$ with $U \cap E \neq \emptyset$, $V \cap E \neq \emptyset$, $E \subseteq U \cup V$, and $E \cap U \cap V = \emptyset$.

*Key Insight*: Connectedness forbids a nonempty disjoint open split.

**Proposition 6.5.2.** *Connectedness and Clopen Sets*
*Let $(X,\tau)$ be a topological space. Then $X$ is connected iff every clopen $A \subseteq X$ satisfies $A = \emptyset$ or $A = X$.*

*Proof.*
($\Rightarrow$) If $A \subseteq X$ is clopen, nonempty, and proper, then $(A, X \setminus A)$ is a separation of $X$, impossible.
($\Leftarrow$) If $(U,V)$ separates $X$, then $U = X \setminus V$ is clopen, nonempty, and proper, impossible.

*Structural Commentary*: Complements convert separations into clopen sets.
*Key Insight*: Connected means no nontrivial clopen subset. □

**Proposition 6.5.3.** *Continuous Images of Connected Spaces Are Connected*
*Let $f : X \to Y$ be continuous. If $X$ is connected, then $f(X)$ is connected in the subspace topology.*

*Proof.* Let $g : X \to f(X)$ be $f$ with codomain restricted to $f(X)$. If $(U, V)$ separates $f(X)$, then $g^{-1}(U)$ and $g^{-1}(V)$ are nonempty disjoint open subsets of $X$ with union $X$, contradicting connectedness.
*Structural Commentary*: Restrict the codomain to make the image map surjective.
*Key Insight*: Separations pull back by continuity. □

**Definition 6.5.4.** *Intermediate Value Property*
A space $X$ has the *intermediate value property* iff $f(X)$ is an interval for every continuous $f : X \to \mathbb{R}$.

**Theorem 6.5.5.** *Connectedness and the Intermediate Value Property*
*Let $X$ be a topological space. Then $X$ is connected iff $X$ has the intermediate value property.*

*Proof.*
($\Rightarrow$) If $f : X \to \mathbb{R}$ is continuous, then $f(X)$ is connected (Proposition 6.5.3), hence an interval (Theorem 5.1.2).
($\Leftarrow$) If $(U, V)$ separates $X$, define $f : X \to \mathbb{R}$ by $f|_U = 0$ and $f|_V = 1$.
Since $U, V$ are clopen, $f$ is continuous, but $f(X) = \{0, 1\}$ is not an interval.
*Structural Commentary*: Separations produce continuous two-valued tests.
*Key Insight*: Intermediate values are connectedness seen by $\mathbb{R}$. □

**Proposition 6.5.6.** *Unions Through a Common Point*
*Let $\{C_\alpha\}_{\alpha \in A}$ be connected subsets of a topological space $X$ with $\bigcap_{\alpha \in A} C_\alpha \neq \emptyset$. Then $C := \bigcup_{\alpha \in A} C_\alpha$ is connected.*

*Proof.* Choose $p \in \bigcap_{\alpha \in A} C_\alpha$. Suppose $(U, V)$ separates $C$ and $p \in U$. For each $\alpha$, the connected set $C_\alpha$ meets $U$ at $p$, so $C_\alpha \cap V = \emptyset$. Hence $V \cap C = \emptyset$, contradicting that $V$ is a nonempty side of the separation.
*Structural Commentary*: Each connected piece must stay on the side containing the shared point.
*Key Insight*: A common point blocks separation. □

**Proposition 6.5.7.** *Connectedness Between a Set and Its Closure*
*Let $A \subseteq B \subseteq \overline{A}$ in a topological space $X$. If $A$ is connected, then $B$ and $\overline{A}$ are connected.*

*Proof.* Suppose $(P, Q)$ separates $B$. Since $A$ is connected, $A \subseteq P$ or $A \subseteq Q$; assume $A \subseteq P$.
If $q \in Q$, choose $O \in \tau_X$ with $q \in O$ and $O \cap B \subseteq Q$.
Since $q \in B \subseteq \overline{A}$, $O \cap A \neq \emptyset$, contradicting $A \subseteq P$ and $P \cap Q = \emptyset$. Thus $Q = \emptyset$, impossible.
*Structural Commentary*: Relative open separation near $\overline{A}$ must meet $A$.
*Key Insight*: Closure points cannot form a new component. □

**Definition 6.5.8.** *Path-Connected and Arcwise Connected Spaces*
A topological space $X$ is *path-connected*, or *arcwise connected*, iff for all $u, v \in X$ there exists continuous $\gamma : [0, 1] \to X$ with $\gamma(0) = u$ and $\gamma(1) = v$.

**Theorem 6.5.9.** *Path-Connected Spaces Are Connected*
*Let $X$ be path-connected. Then $X$ is connected.*

*Proof.* Suppose $(U, V)$ separates $X$. Choose $u \in U$, $v \in V$, and a path $\gamma : [0, 1] \to X$ from $u$ to $v$.
By Proposition 6.5.3, $\gamma([0, 1])$ is connected, but $(U \cap \gamma([0, 1]), V \cap \gamma([0, 1]))$ separates it, a contradiction.
*Structural Commentary*: A path imports interval connectedness into $X$.
*Key Insight*: Paths cannot cross a separation. □

**Proposition 6.5.10.** *Connected Open Sets in Euclidean Space Are Path-Connected*
*Let $O \subseteq \mathbb{R}^n$ be open and connected. Then any $x, y \in O$ can be joined in $O$ by a finite polygonal arc.*

*Proof.* Fix $x_0 \in O$. Write $a \sim b$ if $a, b \in O$ joined in $O$ by finite polygonal arc. Set $C := \{x \in O : x_0 \sim x\}$.
Then $x_0 \in C$. If $B(x, r) \subseteq O$, any two points of $B(x, r)$ are joined by a segment, so $B(x, r) \subseteq C$ when $x \in C$, and $B(x, r) \cap C = \emptyset$ when $x \notin C$. Thus $C$ and $O \setminus C$ are open in $O$; connectedness gives $C = O$.
*Structural Commentary*: Polygonal reachability is both open and closed in an open set.
*Key Insight*: Local line segments make one component all of $O$. □

**Example 6.5.11.** *A Connected Space That Is Not Path-Connected*
Let $G := \{(x, \sin(1/x)) : 0 < x \leq 1\}$, $L := \{0\} \times [-1, 1]$, and $T := G \cup L \subseteq \mathbb{R}^2$. Then $T$ is connected but not path-connected. Since $G$ is the continuous image of connected $(0, 1]$ and $T = \overline{G}$, Proposition 6.5.7 gives connectedness. No path can join $L$ to $G$ without forcing an impossible one-sided limit of $\sin(1/x)$ as $x \to 0^+$.




Figure 153: Topologist's sine curve: $y = \sin(1/x)$ accumulates on $\{0\} \times [-1, 1]$; connected, not path-connected.

# 7 Finite-Dimensional Topological Vector Spaces

## 7.1 Natural Topology and Continuity

Finite-dimensional real vector spaces carry a canonical topology: choose coordinates, use the standard Euclidean topology on $\mathbb{R}^n$, and then forget that coordinates were chosen. All coordinate changes are homeomorphisms.

**Definition 7.1.1.** *Topological Vector Space*
Let $\mathcal{V}$ be a real vector space and let $\mathcal{T}$ be a topology on its underlying set.
The pair $(\mathcal{V}, \mathcal{T})$ is a *topological vector space* if addition $A : \mathcal{V} \times \mathcal{V} \to \mathcal{V}$, $A(\mathbf{u}, \mathbf{v}) = \mathbf{u} + \mathbf{v}$, and scalar multiplication $M : \mathbb{R} \times \mathcal{V} \to \mathcal{V}$, $M(a, \mathbf{v}) = a\mathbf{v}$ are continuous, where products carry the product topology.

**Example 7.1.2.** *The Standard Topology on $\mathbb{R}^n$*
With the Euclidean topology, $\mathbb{R}^n$ is a topological vector space: the maps $(\mathbf{x}, \mathbf{y}) \longmapsto \mathbf{x} + \mathbf{y}$, $(a, \mathbf{x}) \longmapsto a\mathbf{x}$ have polynomial coordinate functions $x_i + y_i$ and $ax_i$, hence are continuous.

**Proposition 7.1.3.** *Real Matrix Maps Are Continuous*
*Let $A = [a_{j,k}] \in \mathrm{Mat}(m, n, \mathbb{R})$, and let $T_A : \mathbb{R}^n \to \mathbb{R}^m$ be the induced map $T_A(\mathbf{x}) = A\mathbf{x}$. Then $T_A$ is continuous for the Euclidean topologies. If $m = n$ and $A$ is invertible, then $T_A$ is a homeomorphism. In particular, every linear functional on $\mathbb{R}^n$ is continuous.*

*Proof.* For $\mathbf{x} = (x_1, \ldots, x_n)^t$, the $j$th coordinate of $T_A(\mathbf{x})$ is $\sum_{k=1}^{n} a_{j,k}x_k$, hence is continuous. Thus $T_A$ is continuous. If $A$ is invertible, then $(T_A)^{-1} = T_{A^{-1}}$, which is continuous by the first part. Finally, every $f \in (\mathbb{R}^n)'$ equals $T_A$ for a unique $A \in \mathrm{Mat}(1, n, \mathbb{R})$, so $f$ is continuous. *Structural Commentary*: Matrix maps on Euclidean coordinate spaces are continuous coordinate by coordinate.
*Key Insight*: Invertible real matrices give homeomorphisms of Euclidean space. □

**Definition 7.1.4.** *Natural Topology*
Let $\mathcal{V}$ be an $n$-dimensional real vector space, let $\beta$ be a basis, and let $[\cdot]_\beta : \mathcal{V} \to \mathbb{R}^n$ be the coordinate isomorphism.
The *natural topology* on $\mathcal{V}$ is $\mathcal{T}_\beta := \{[\cdot]_\beta^{-1}(O) : O \subseteq \mathbb{R}^n \text{ is open}\}$.
Equivalently, $U \subseteq \mathcal{V}$ is open if and only if $[\cdot]_\beta(U)$ is open in $\mathbb{R}^n$.

**Proposition 7.1.5.** *Natural Topology Is Basis-Independent*
*If $\beta$ and $\gamma$ are bases of $\mathcal{V}$, then $\mathcal{T}_\beta = \mathcal{T}_\gamma$.*

*Proof.* Set $H_{\gamma \leftarrow \beta} := [\cdot]_\gamma \circ [\cdot]_\beta^{-1} : \mathbb{R}^n \to \mathbb{R}^n$.
Then $H_{\gamma \leftarrow \beta}\mathbf{x} = C_{\gamma \leftarrow \beta}\mathbf{x}$. Since $C_{\gamma \leftarrow \beta}$ is invertible with inverse $C_{\beta \leftarrow \gamma}$, Proposition 7.1.3 shows that $H_{\gamma \leftarrow \beta}$ is a homeomorphism with inverse $H_{\beta \leftarrow \gamma}$.
Hence, for $U \subseteq \mathcal{V}$, $U \in \mathcal{T}_\beta \iff [\cdot]_\beta(U)$ is open $\iff H_{\gamma \leftarrow \beta}([\cdot]_\beta(U)) = [\cdot]_\gamma(U)$ is open $\iff U \in \mathcal{T}_\gamma$. *Structural Commentary*: Basis changes are homeomorphisms.
*Key Insight*: The natural topology is carried by the vector-space structure, not by a preferred basis. □

**Theorem 7.1.6.** *Characterization of the Natural Topology*
*Let* $\dim_{\mathbb{R}} \mathcal{V} < \infty$. *There is a unique topology* $\mathcal{T}$ *on* $\mathcal{V}$ *such that:*

*(i)* $(\mathcal{V}, \mathcal{T})$ *is a topological vector space;*
*(ii)* $\mathcal{V}' \subseteq C((\mathcal{V}, \mathcal{T}), \mathbb{R})$.

*Moreover,* $\mathcal{T} = \mathcal{T}_\beta$ *for every basis* $\beta$.

*Proof.* Fix an arbitrary basis $\beta = (\mathbf{v}_1, \dots, \mathbf{v}_n)$.

(i) *Existence*: put $\mathcal{T} = \mathcal{T}_\beta$. Since $[\cdot]_\beta : (\mathcal{V}, \mathcal{T}_\beta) \to \mathbb{R}^n$ is a homeomorphism, the vector operations on $\mathcal{V}$ are transported from the continuous vector operations on $\mathbb{R}^n$.
Also, for $\varphi \in \mathcal{V}'$, $\varphi \circ [\cdot]_\beta^{-1} \in (\mathbb{R}^n)'$, hence is continuous by Proposition 7.1.3. Thus (i)–(ii) hold.

(ii) *Uniqueness*: suppose $\mathcal{T}$ satisfies (i)–(ii).
For $1 \le i \le n$, the coordinate functional $\pi_i \circ [\cdot]_\beta \in \mathcal{V}'$ is continuous, so $[\cdot]_\beta : (\mathcal{V}, \mathcal{T}) \to \mathbb{R}^n$ is continuous.
Its inverse $\Psi(a_1, \dots, a_n) = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$ is continuous by (i), since it is built from scalar multiplication and finite sums. Hence $[\cdot]_\beta$ is a homeomorphism, so $\mathcal{T} = \mathcal{T}_\beta$.

*Structural Commentary*: Functionals give coordinates; operations give inverse coordinates.
*Key Insight*: Functionals and vector operations determine finite-dimensional coordinates topologically. □

**Corollary 7.1.7.** *Finite-Dimensional Linear Maps Are Continuous*
*Let* $\dim_{\mathbb{R}} \mathcal{V}, \dim_{\mathbb{R}} \mathcal{W} < \infty$, *with natural topologies. Then* $\mathcal{L}(\mathcal{V}, \mathcal{W}) \subseteq C(\mathcal{V}, \mathcal{W})$.
*If* $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ *is an isomorphism, then* $T$ *is a homeomorphism.*

*Proof.* Choose bases $\beta$ of $\mathcal{V}$ and $\gamma$ of $\mathcal{W}$. For $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, $[\cdot]_\gamma \circ T \circ [\cdot]_\beta^{-1}(\mathbf{x}) = [T]_{\gamma \leftarrow \beta}\mathbf{x}$, which is continuous by Proposition 7.1.3. Since $[\cdot]_\beta$ and $[\cdot]_\gamma$ are homeomorphisms, $T$ is continuous.
If $T$ is an isomorphism, apply the same argument to $T^{-1}$.
*Structural Commentary*: Finite-dimensional linear maps are just matrix maps in natural coordinates.
*Key Insight*: A basis turns topological questions about linear maps into continuity of matrix multiplication. □

**Perspective 7.1.8.** *Finite-Dimensional Topology Is Rigid*
For finite-dimensional real vector spaces, topology is algebraically forced.
Note that $\mathcal{T}_\beta = \mathcal{T}_\gamma$; $\mathcal{L}(\mathcal{V}, \mathcal{W}) \subseteq C(\mathcal{V}, \mathcal{W})$; $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ bijective $\Rightarrow T$ is a homeomorphism.
Infinite-dimensional topological vector spaces need not be rigid in this sense.

# Part XII
# Calculus, Series, and Approximation

## 1 One-Variable Differentiation and Riemann Integration

### 1.1 Differentiation and Basic Theorems

One-variable calculus studies functions on intervals through derivatives, mean-value theorems, Riemann integration, and the FTC, then points toward vector-valued estimates and linear approximation.

**Definition 1.1.1.** *Derivative of a Real Function*
Let $f$ be real-valued on $[a,b]$ and let $x \in [a,b]$. For $a < t < b$ and $t \neq x$, set $\phi_x(t) := \frac{f(t)-f(x)}{t-x}$.
If $\lim_{t\to x} \phi_x(t)$ exists, denote *derivative* by $f'(x)$; then $f$ is *differentiable at* $x$.
If this holds for every $x \in E \subseteq [a,b]$, then $f$ is *differentiable on* $E$.

Figure 154: Domain matters: isolated points give no punctured approach; $[x_0, \infty)$ gives a right-hand one.

**Perspective 1.1.2.** *Derivative Depends on the Domain*
For $f : X \to \mathbb{R}$, $f'(x_0)$ is the limit of $\frac{f(x)-f(x_0)}{x-x_0}$ as $x \to x_0$ in $X \setminus \{x_0\}$; see Definition 5.1.4. Hence $x_0$ must be a limit point of $X$: isolated points give no derivative, while half-interval domains give one-sided derivatives.




Figure 155: The derivative is the limiting slope of secant lines through $(x, f(x))$.

**Proposition 1.1.3.** *First-Order Approximation Form*
*Let $X \subseteq \mathbb{R}$, let $x_0 \in X$ be a limit point, let $f : X \to \mathbb{R}$, and let $L \in \mathbb{R}$. Then $f'(x_0) = L$ relative to $X$ iff $\forall \varepsilon > 0$ $\exists \delta > 0$ such that $|f(x) - f(x_0) - L(x - x_0)| \leq \varepsilon\,|x - x_0|$ whenever $x \in X$ and $|x - x_0| < \delta$.*

*Proof.* For $x \neq x_0$, $\frac{|f(x)-f(x_0)-L(x-x_0)|}{|x-x_0|} = \left|\frac{f(x)-f(x_0)}{x-x_0} - L\right|$, while $x = x_0$ makes approx. estimate automatic.

($\Rightarrow$) If $f'(x_0) = L$, then the displayed right-hand side is at most $\varepsilon$ for all $x \in X \setminus \{x_0\}$ sufficiently near $x_0$. Multiplying by $|x - x_0|$ gives the approximation estimate.

($\Leftarrow$) If the approximation estimate holds, then division by $|x - x_0|$ gives $\left|\frac{f(x)-f(x_0)}{x-x_0} - L\right| \leq \varepsilon$ for all $x \in X\setminus\{x_0\}$ sufficiently near $x_0$. Hence $f'(x_0) = L$.

*Structural Commentary*: Differentiability is first-order linear approximation.
*Key Insight*: Multiply or divide by $|x - x_0|$. □

**Theorem 1.1.4.** *Differentiability Implies Continuity*
*If $f$ is differentiable at $x$, then $f$ is continuous at $x$.*

*Proof.* For $t \neq x$, $f(t) - f(x) = \frac{f(t)-f(x)}{t-x}(t - x)$. As $t \to x$, the difference quotient tends to $f'(x)$ and $t - x \to 0$. By Theorem 5.1.7, $f(t) - f(x) \to 0$, so $f(t) \to f(x)$. Theorem 5.1.11 gives continuity at $x$.
*Structural Commentary*: A differentiable function has first-order control near the point.
*Key Insight*: The quotient stays near a finite number while $t - x$ shrinks to zero. □

**Theorem 1.1.5.** *Algebra Rules for Derivatives*
*Let $f, g$ be differentiable at $x$.*

*(i) $f+g$ is differentiable at $x$, and $(f+g)'(x) = f'(x) + g'(x)$.*
*(ii) $fg$ is differentiable at $x$, and $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$.*
*(iii) If $g(x) \neq 0$, then $f/g$ is differentiable at $x$, and $\left(\frac{f}{g}\right)'(x) = \frac{g(x)f'(x) - g'(x)f(x)}{g(x)^2}$.*

*Proof.*

(i) Limit algebra applied to $\frac{(f+g)(t)-(f+g)(x)}{t-x} = \frac{f(t)-f(x)}{t-x} + \frac{g(t)-g(x)}{t-x}$ gives the sum rule.
(ii) For $h = fg$, $h(t) - h(x) = f(t)(g(t) - g(x)) + g(x)(f(t) - f(x))$. Divide by $t - x$ and let $t \to x$. Theorem 1.1.4 gives $f(t) \to f(x)$, hence the product rule.
(iii) If $h = f/g$, continuity of $g$ gives $g(t) \neq 0$ near $x$, and $\frac{h(t)-h(x)}{t-x} = \frac{1}{g(t)g(x)}\left[g(x)\frac{f(t)-f(x)}{t-x} - f(x)\frac{g(t)-g(x)}{t-x}\right]$. Let $t \to x$ to get the quotient rule.

*Structural Commentary*: Product and quotient rules are difference-quotient identities plus continuity.
*Key Insight*: Difference-quotient identities plus continuity give the algebra rules. □

**Theorem 1.1.6.** *Chain Rule*
*Let $f$ be differentiable at $x$ and let $g$ be differentiable at $f(x)$.*
*If $h = g \circ f$, then $h$ is differentiable at $x$ and $h'(x) = g'(f(x))f'(x)$.*

*Proof.* Set $y = f(x)$. For $u(t) \to 0$, $f(t) - f(x) = (t - x)(f'(x) + u(t))$.
For $v(s) \to 0$, $g(s) - g(y) = (s - y)(g'(y) + v(s))$.
Since $f(t) \to y$, for $t \neq x$, $\frac{h(t)-h(x)}{t-x} = (g'(y) + v(f(t)))(f'(x) + u(t)) \to g'(y)f'(x)$.
*Structural Commentary*: Differentiate by composing first-order expansions.
*Key Insight*: First-order parts multiply; errors vanish. □

**Example 1.1.7.** *Oscillation at a Point*
Assuming the usual derivative of $\sin x$, define

$$f_1(x) = \begin{cases} x\sin(1/x), & x \neq 0, \\ 0, & x = 0, \end{cases} \qquad f_2(x) = \begin{cases} x^2\sin(1/x), & x \neq 0, \\ 0, & x = 0. \end{cases}$$

For $f_1$, the quotient at 0 is $\sin(1/t)$, so $f_1'(0)$ does not exist. For $f_2$, $|(f_2(t) - f_2(0))/t| \leq |t|$, so $f_2'(0) = 0$.
However, for $x \neq 0$, $f_2'(x) = 2x\sin(1/x) - \cos(1/x)$, so $f_2'$ is not continuous at 0.

**Definition 1.1.8.** *Local Extrema*
Let $f$ be a real-valued function on a metric space $X$. The function has a *local maximum* at $p \in X$ if there is $\delta > 0$ such that $f(q) \leq f(p)$ whenever $d(p, q) < \delta$. Local minima are defined by reversing the inequality.

**Remark 1.1.9.** *Local and Global Extrema*
Global extrema are local, but local extrema depend on domain: isolated points are automatic extrema. Derivative tests therefore need an interior interval point; compactness supplies global extrema (Theorem 5.1.37).

**Theorem 1.1.10.** *Interior Extremum Has Zero Derivative*
*If $f$ has a local extremum at $x \in (a, b)$ and $f'(x)$ exists, then $f'(x) = 0$.*

*Proof.* For a local maximum, $f(t) \leq f(x)$ near $x$.
Thus the quotient $(f(t) - f(x))/(t - x)$ is nonnegative for $t < x$ and nonpositive for $t > x$. Since the two-sided limit exists, $f'(x) \geq 0$ and $f'(x) \leq 0$, so $f'(x) = 0$. For a local minimum, apply the local-maximum case to $-f$. Since $(-f)'(x) = -f'(x)$, we get $-f'(x) = 0$, hence $f'(x) = 0$.
*Structural Commentary*: Interior extrema force opposite one-sided quotient signs.
*Key Insight*: A two-sided derivative compatible with both signs must be zero. □

**Theorem 1.1.11.** *Rolle's Theorem*
*If $f : [a, b] \to \mathbb{R}$ continuous on $[a, b]$, differentiable on $(a, b)$, and $f(a) = f(b)$, then $\exists x \in (a, b)$ s.t. $f'(x) = 0$.*

*Proof.* If $f$ is constant, any $x \in (a, b)$ works. Otherwise, by Theorem 5.1.37, $f$ attains a nonendpoint extremum at some $x \in (a, b)$. Theorem 1.1.10 gives $f'(x) = 0$.
*Structural Commentary*: Equal endpoints force any nonconstant height change to turn around inside.
*Key Insight*: Compactness gives the extremum; differentiability makes the interior extremum stationary. □

**Theorem 1.1.12.** *Cauchy Mean Value Theorem*
*Let $f, g$ be continuous on $[a, b]$, differentiable on $(a, b)$. $\exists\, x \in (a, b)$, $\big(f(b) - f(a)\big)g'(x) = \big(g(b) - g(a)\big)f'(x)$.*

*Proof.* Set $h(t) := \big(f(b) - f(a)\big)g(t) - \big(g(b) - g(a)\big)f(t)$.
Then $h(a) = h(b)$, $h$ is continuous on $[a, b]$, and $h$ is differentiable on $(a, b)$. By Theorem 1.1.11, $h'(x) = 0$ for some $x \in (a, b)$. Since $h'(x) = \big(f(b) - f(a)\big)g'(x) - \big(g(b) - g(a)\big)f'(x)$, the identity follows.
*Structural Commentary*: Build one auxiliary function whose endpoint values agree.
*Key Insight*: Equal endpoints force an interior horizontal tangent. □




Figure 156: The Mean Value Theorem finds an interior tangent slope equal to the endpoint secant slope.

**Corollary 1.1.13.** *Mean Value Theorem*
*If $f$ is continuous on $[a, b]$, differentiable in $(a, b)$, then $\exists\ x \in (a, b)$ such that $f(b) - f(a) = (b - a)f'(x)$.*

*Proof.* Let $g(t) = t$. Then $g'(x) = 1$, $g(b) - g(a) = b - a$.
Hence Theorem 1.1.12 gives $f(b) - f(a) = (b - a)f'(x)$ for some $x \in (a, b)$.
*Structural Commentary*: This is Cauchy's theorem with $g(t) = t$.
*Key Insight*: A secant slope is achieved as a tangent slope somewhere inside. □

**Corollary 1.1.14.** *Bounded Derivative Gives Lipschitz Control*
*If $f : [a, b] \to \mathbb{R}$ is continuous on $[a, b]$, differentiable on $(a, b)$, and $|f'| \leq M$ on $(a, b)$, then $f$ is $M$-Lipschitz on $[a, b]$, hence uniformly continuous.*

*Proof.* For $x < y$, Corollary 1.1.13 gives $f(y) - f(x) = (y - x)f'(c)$ for some $c \in (x, y)$,
so $|f(y) - f(x)| \leq M(y - x)$. Symmetry handles $x > y$, and $x = y$ is trivial.
*Structural Commentary*: A derivative bound controls every secant slope.
*Key Insight*: Local slope bounds become global Lipschitz bounds through the Mean Value Theorem. □

**Theorem 1.1.15.** *Derivative Sign and Monotonicity*
*Suppose $f$ is differentiable in $(a, b)$.*

*(i) If $f'(x) \geq 0$ for all $x \in (a, b)$, then $f$ is monotonically increasing.*
*(ii) If $f'(x) = 0$ for all $x \in (a, b)$, then $f$ is constant.*
*(iii) If $f'(x) \leq 0$ for all $x \in (a, b)$, then $f$ is monotonically decreasing.*

*Proof.* For $x_1 < x_2$ in $(a, b)$, Corollary 1.1.13 on $[x_1, x_2]$ gives $f(x_2) - f(x_1) = (x_2 - x_1)f'(c)$ for some $c \in (x_1, x_2)$.

(i) If $f' \geq 0$, then $f(x_2) - f(x_1) \geq 0$.
(ii) If $f' = 0$, then $f(x_2) - f(x_1) = 0$.
(iii) If $f' \leq 0$, then $f(x_2) - f(x_1) \leq 0$.

*Structural Commentary*: Monotonicity is a global reading of the mean-value identity.
*Key Insight*: The sign of every tangent slope controls every secant slope. □

**Theorem 1.1.16.** *Derivatives Have the Intermediate Value Property*
*If $f$ is differentiable on $[a, b]$ in the one-sided endpoint sense and $f'(a) < \lambda < f'(b)$,*
*Then $f'(x) = \lambda$ for some $x \in (a, b)$. The same conclusion holds when $f'(b) < \lambda < f'(a)$.*

*Proof.* Set $g(t) = f(t) - \lambda t$. Since $g'(a) < 0 < g'(b)$, there are $t_1, t_2 \in (a, b)$ with $g(t_1) < g(a)$ and $g(t_2) < g(b)$.
Hence the minimum of $g$ on $[a, b]$ occurs at some $x \in (a, b)$. By Theorem 1.1.10, $0 = g'(x) = f'(x) - \lambda$.
Apply the same argument to $-f$ for the reversed case.
*Structural Commentary*: A derivative need not be continuous, but it cannot jump over values.
*Key Insight*: Tilt the graph by subtracting $\lambda t$ and force an interior extremum. □




Figure 157: Sliding fixed-width secants in Darboux's theorem.

**Perspective 1.1.17.** *Sliding Secants*
For fixed small $h > 0$, the secant-slope function $\sigma_h(x) := \frac{f(x+h)-f(x)}{h}$ is continuous. Thus Theorem 5.1.41 finds intermediate secant slopes, and Corollary 1.1.13 converts them to derivative values.
*Key Insight*: Slide secants, then apply the Mean Value Theorem.

**Corollary 1.1.18.** *Derivatives Have No Simple Discontinuities*
*If $f$ is differentiable on $[a,b]$, then $f'$ cannot have a simple discontinuity on $[a,b]$.*

*Proof.* Suppose a simple discontinuity occurs at an interior point $c$; endpoint cases are one-sided.
If $f'(c-) \neq f'(c+)$, choose $\lambda$ strictly between these limits, $\lambda \neq f'(c)$. For $x < c < y$ close to $c$, Theorem 1.1.16 gives $z \in (x,y)$ with $f'(z) = \lambda$, contradicting the one-sided limits. If $f'(c-) = f'(c+) = L \neq f'(c)$, choose $\lambda$ between $L$ and $f'(c)$ and apply the same theorem on a sufficiently small one-sided interval with endpoint $c$.
*Structural Commentary*: Darboux behavior rules out jump discontinuities for derivatives.
*Key Insight*: Derivatives may oscillate, but they do not jump. □

**Theorem 1.1.19.** *Derivative of an Inverse Function*
*Let $f : I \to \mathbb{R}$ be one-to-one on an open interval $I$, let $x_0 \in I$, and put $y_0 = f(x_0)$. Suppose $f$ is differentiable at $x_0$, $f'(x_0) \neq 0$, and $g = f^{-1} : f(I) \to I$ is continuous at $y_0$. Then $g$ is differentiable at $y_0$ and $g'(y_0) = \frac{1}{f'(x_0)}$.*

*Proof.* For $y \in f(I)$ with $y \neq y_0$, write $x = g(y)$. Then $x \neq x_0$ and $\frac{g(y)-g(y_0)}{y-y_0} = \frac{x-x_0}{f(x)-f(x_0)} = \left(\frac{f(x)-f(x_0)}{x-x_0}\right)^{-1}$.
As $y \to y_0$, the assumed continuity of $g$ gives $x = g(y) \to x_0$.
The difference quotient for $f$ tends to $f'(x_0) \neq 0$, so the reciprocal tends to $1/f'(x_0)$ by Theorem 5.1.7.
*Structural Commentary*: Matched inverse quotients are reciprocal.
*Key Insight*: Continuity of the inverse converts the limit $y \to y_0$ back into $x \to x_0$. □

**Corollary 1.1.20.** *One-Dimensional Inverse Function Theorem*
*If $f : (a,b) \to (c,d)$ is a differentiable bijection and $f' \neq 0$,*
*then $f$ is a homeomorphism and $f^{-1}$ is differentiable, with $(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}$.*

*Proof.* By Theorem 1.1.16, $f'$ has constant sign. Corollary 1.1.13 makes $f$ strictly monotone, hence its continuous interval inverse is continuous. Apply Theorem 1.1.19 pointwise.
*Structural Commentary*: Darboux gives constant sign; the inverse derivative formula is local.
*Key Insight*: Nonvanishing derivative is the one-dimensional obstruction to differentiating the inverse. □

**Theorem 1.1.21.** *L'Hospital's Rule*
*Let $-\infty \leq a < b \leq +\infty$, let $f, g$ be differentiable on $(a,b)$, assume $g' \neq 0$, and suppose $\frac{f'}{g'} \to A \in \overline{\mathbb{R}}$ as $x \to a^+$.*

*(i) If $f, g \to 0$ as $x \to a^+$, then $\frac{f}{g} \to A$ as $x \to a^+$.*
*(ii) If $|g| \to \infty$ as $x \to a^+$, then $\frac{f}{g} \to A$ as $x \to a^+$.*

*The analogous statements hold as $x \to b^-$.*

*Proof.* For $a < r < s < b$, Theorem 1.1.12 gives $t \in (r,s)$ with $\frac{f(s)-f(r)}{g(s)-g(r)} = \frac{f'(t)}{g'(t)}$, so eventual bounds on $\frac{f'}{g'}$ near $a$ transfer to these difference quotients.

(i) Fix $s$ and let $r \to a^+$; since $f(r), g(r) \to 0$, the same bounds hold for $\frac{f(s)}{g(s)}$.

(ii) Fix $s$ near $a$ and use $\frac{f(r)}{g(r)} = \frac{f(r)-f(s)}{g(r)-g(s)} \frac{g(r)-g(s)}{g(r)} + \frac{f(s)}{g(r)}$.
As $r \to a^+$, the second factor tends to 1 and the last term to 0, so the same bounds hold for $\frac{f(r)}{g(r)}$.

In both cases, the transferred eventual bounds force $\frac{f}{g} \to A$.
*Structural Commentary*: CMVT transfers derivative-quotient bounds to function quotients.
*Key Insight*: Bounds on $f'/g'$ force the same limit for $f/g$. □




Figure 158: For $f(x) = x^{-3}$, the one-sided limits at 0 are opposite infinities.

**Definition 1.1.22.** *Higher Derivatives*
If $f$ has derivative $f'$ on an interval and $f'$ is differentiable, then $(f')'$ is denoted $f''$ and called the *second derivative*. Continuing recursively gives $f^{(n)}$, the derivative of order $n$.

**Definition 1.1.23.** *Smoothness Classes*
On an interval, $C^0$ means continuous, $C^r$ means $f^{(r)}$ exists and continuous, $C^\infty$ means $C^r$ for every finite $r$.

**Theorem 1.1.24.** *Higher-Order Leibniz Product Rule*
*Let $r \geq 1$, and $f, g : (a,b) \to \mathbb{R}$ be $r$ times differentiable. Then $\forall x \in (a,b)$, $(fg)^{(r)}(x) = \sum_{k=0}^{r} \binom{r}{k} f^{(k)}(x) g^{(r-k)}(x)$.*

*Proof.* Induct on $r$. For $r = 1$, this is the product rule in Theorem 1.1.5. Assume the formula holds at order $r$. If $f, g$ are $(r+1)$ times differentiable, differentiating gives $(fg)^{(r+1)} = \sum_{k=0}^{r} \binom{r}{k} f^{(k+1)} g^{(r-k)} + \sum_{k=0}^{r} \binom{r}{k} f^{(k)} g^{(r-k+1)}$. Reindex the first sum by $j = k+1$; the coefficient of $f^{(j)} g^{(r+1-j)}$ is $\binom{r}{j-1} + \binom{r}{j} = \binom{r+1}{j}$, with the endpoint terms $j = 0, r+1$ equal to 1.
*Structural Commentary*: Differentiating shifts one derivative from the total count onto $f$ or $g$.
*Key Insight*: Pascal's identity produces the binomial coefficients. □

**Theorem 1.1.25.** *Taylor's Theorem with Lagrange Remainder*
*Let $f : [a,b] \to \mathbb{R}$, let $n \geq 1$, suppose $f^{(n-1)}$ is continuous on $[a,b]$, and suppose $f^{(n)}$ exists on $(a,b)$.*
*For distinct $\alpha, \beta \in [a,b]$, set $P(t) := \sum_{k=0}^{n-1} \frac{f^{(k)}(\alpha)}{k!}(t-\alpha)^k$.*
*Then for some $x$ between $\alpha$ and $\beta$, $f(\beta) = P(\beta) + \frac{f^{(n)}(x)}{n!}(\beta-\alpha)^n$.*

*Proof.* Choose $M$ so that $f(\beta) = P(\beta) + M(\beta-\alpha)^n$, and set $G(t) := f(t) - P(t) - M(t-\alpha)^n$. Then $G(\alpha) = G'(\alpha) = \cdots = G^{(n-1)}(\alpha) = G(\beta) = 0$. Repeated applications of Corollary 1.1.13 give $G^{(n)}(x) = 0$ for some $x$ between $\alpha$ and $\beta$. Since $G^{(n)}(x) = f^{(n)}(x) - n!M$, the formula follows.
*Structural Commentary*: The auxiliary error vanishes at $\beta$ and to order $n-1$ at $\alpha$.
*Key Insight*: Repeated Rolle-style arguments turn endpoint cancellation into a precise remainder. □

**Definition 1.1.26.** *Vector-Valued Derivative*
For $\mathbf{f} : [a,b] \to \mathbb{R}^k$, set $\mathbf{f}'(x) = \mathbf{L}$ when $\lim_{t \to x} \left\| \frac{\mathbf{f}(t) - \mathbf{f}(x)}{t-x} - \mathbf{L} \right\|_2 = 0$.
For $\mathbf{f} = (f_1, \ldots, f_k)$, this is equivalent to componentwise differentiability, with $\mathbf{f}'(x) = (f_1'(x), \ldots, f_k'(x))$.

**Theorem 1.1.27.** *Vector-Valued Mean Value Inequality*
*If $\mathbf{f} : [a,b] \to \mathbb{R}^k$ is continuous on $[a,b]$ and differentiable on $(a,b)$.*
*Then for some $x \in (a,b)$, $\|\mathbf{f}(b) - \mathbf{f}(a)\|_2 \leq (b-a) \|\mathbf{f}'(x)\|_2$.*

*Proof.* Set $\mathbf{z} := \mathbf{f}(b) - \mathbf{f}(a)$ and $\varphi(t) := \mathbf{z} \cdot \mathbf{f}(t)$.
Corollary 1.1.13 gives $\varphi(b) - \varphi(a) = (b-a)\varphi'(x) = (b-a)\mathbf{z} \cdot \mathbf{f}'(x)$ for some $x \in (a,b)$. Since $\varphi(b) - \varphi(a) = \|\mathbf{z}\|_2^2$, Theorem 1.1.13 gives $\|\mathbf{z}\|_2^2 \leq (b-a) \|\mathbf{z}\|_2 \|\mathbf{f}'(x)\|_2$, and the result follows by cancellation, with $\mathbf{z} = 0$ trivial.
*Structural Commentary*: A scalar projection yields the vector norm bound.
*Key Insight*: Project onto the displacement vector, then use Schwarz to return to norms. □

**Method 1.1.28.** *Differentiation Proof Patterns*

(i) *Domain*: require $x_0$ to be a limit point of $X$; isolated points have no derivative.
(ii) *Definition*: factor $f(t) - f(x) = (t - x)Q_x(t)$ and take $Q_x(t) \to f'(x)$.
(iii) *Approximation*: use Proposition 1.1.3 to replace quotient limits by first-order error bounds.
(iv) *Algebra*: prove quotient identities first; use continuity to pass to limits.
(v) *Mean values*: choose $h$ with equal endpoint values and apply Theorem 1.1.10.
(vi) *Taylor*: set $G = f - P - M(t - \alpha)^n$ and iterate the mean value theorem.

*Key Insight*: Difference quotients and auxiliary functions drive one-variable differentiation.

## 1.2 Riemann Integration

The integration side of one-variable calculus is organized by Riemann and Riemann–Stieltjes sums; the fundamental theorem connects the two halves.

This subsection develops Riemann and Riemann–Stieltjes integration by partitions, then links integrals to antiderivatives through the Fundamental Theorem of Calculus. Point partitions and interval-envelope partitions are equivalent via common refinement.

**Definition 1.2.1.** *Partitions and Riemann Upper/Lower Sums*

(i) A *partition* of $[a, b]$ is $\Pi = \{x_0, \dots, x_n\}$ with $a = x_0 \le x_1 \le \cdots \le x_n = b$.
Set $\Delta x_i := x_i - x_{i-1}$ for $1 \le i \le n$.
(ii) For bounded $f : [a, b] \to \mathbb{R}$, set $M_i := \sup_{x \in [x_{i-1}, x_i]} f(x)$ and $m_i := \inf_{x \in [x_{i-1}, x_i]} f(x)$.
(iii) The *upper* and *lower Darboux sums* are $U(\Pi, f) := \sum_{i=1}^{n} M_i \Delta x_i$ and $L(\Pi, f) := \sum_{i=1}^{n} m_i \Delta x_i$.
(iv) The *upper* and *lower Darboux integrals* are $\overline{\int_a^b f\,dx} := \inf_\Pi U(\Pi, f)$ and $\underline{\int_a^b f\,dx} := \sup_\Pi L(\Pi, f)$, where $\Pi$ ranges over all partitions of $[a, b]$.
(v) If $\overline{\int_a^b f\,dx} = \underline{\int_a^b f\,dx}$, then $f$ is *Riemann integrable* on $[a, b]$; $f \in \mathcal{R}[a, b]$, $\int_a^b f\,dx := \overline{\int_a^b f\,dx} = \underline{\int_a^b f\,dx}$.

If $m \le f(x) \le M$ for $x \in [a, b]$, then $m(b - a) \le L(\Pi, f) \le U(\Pi, f) \le M(b - a)$ for every partition $\Pi$.
Hence the upper and lower Darboux integrals are finite for bounded $f$.
*Key Insight*: Riemann integrability means least upper Darboux area equals greatest lower Darboux area.




Figure 159: Upper and lower sums record the oscillation of $f$ on each partition interval.

**Definition 1.2.2.** *Tagged Partitions and Riemann Sums*
Let $\Pi = \{x_0, \dots, x_n\}$ be a partition of $[a, b]$, and set $\Delta x_i := x_i - x_{i-1}$.

(i) A *tagging* of $\Pi$ is $c = (c_1, \dots, c_n)$ with $c_i \in [x_{i-1}, x_i]$ for $1 \le i \le n$; the pair $(\Pi, c)$ is a *tagged partition*.
(ii) For $f : [a, b] \to \mathbb{R}$, the associated *Riemann sum* is $R(f, \Pi, c) := \sum_{i=1}^{n} f(c_i)\Delta x_i$.
(iii) If $f$ is bounded, then $L(\Pi, f) \le R(f, \Pi, c) \le U(\Pi, f)$.




Figure 160: A tagged Riemann sum uses one sample value $f(c_i)$ on each subinterval $[x_{i-1}, x_i]$.

**Perspective 1.2.3.** *Tagged Sums and Gauges*
Gauge integration keeps tagged sums but replaces one mesh bound $\delta$ with a local bound $\delta(c_i)$.
Thus fineness may vary by tag, giving a stronger framework for FTC-type results.

# 2 Riemann-Stieltjes and Improper Integration

## 2.1 Riemann-Stieltjes Integrability

**Definition 2.1.1.** *Riemann–Stieltjes Integral*
Let $\alpha : [a,b] \to \mathbb{R}$ be increasing, let $\Pi = \{x_0, \dots, x_n\}$ be a partition of $[a,b]$, and let $f : [a,b] \to \mathbb{R}$ be bounded. Set $M_i := \sup_{x \in [x_{i-1}, x_i]} f(x)$ and $m_i := \inf_{x \in [x_{i-1}, x_i]} f(x)$.

(i) The Stieltjes increment on $[x_{i-1}, x_i]$ is $\Delta\alpha_i := \alpha(x_i) - \alpha(x_{i-1}) \geq 0$ for $1 \leq i \leq n$.

(ii) The *upper* and *lower Riemann–Stieltjes sums* are $U(\Pi, f, \alpha) := \sum_{i=1}^{n} M_i \Delta\alpha_i$ and $L(\Pi, f, \alpha) := \sum_{i=1}^{n} m_i \Delta\alpha_i$.

(iii) The *upper* and *lower Riemann–Stieltjes integrals* are $\overline{\int_a^b f \, d\alpha} := \inf_{\Pi} U(\Pi, f, \alpha)$ and $\underline{\int_a^b f \, d\alpha} := \sup_{\Pi} L(\Pi, f, \alpha)$, where the infimum and supremum range over all partitions of $[a,b]$.

(iv) If $\overline{\int_a^b f \, d\alpha} = \underline{\int_a^b f \, d\alpha}$, then $f$ is *Riemann integrable with respect to* $\alpha$, written $f \in \mathcal{R}(\alpha)$, and $\int_a^b f \, d\alpha := \overline{\int_a^b f \, d\alpha} = \underline{\int_a^b f \, d\alpha}$, also written $\int_a^b f(x) \, d\alpha(x)$.

For $\alpha(x) = x$, this is the ordinary Riemann integral.

**Remark 2.1.2.** *Stieltjes Increments as $\alpha$-Lengths*
Stieltjes sums replace ordinary length $\Delta x_i$ by $\Delta\alpha_i$, the length measured by $\alpha$.
Thus $\alpha(x) = x$ gives Riemann sums, while jumps of $\alpha$ place point mass as in Theorem 2.3.2.

**Definition 2.1.3.** *Refinements*
Let $\Pi, \Pi^*, \Pi_1, \Pi_2$ be partitions of $[a,b]$.

(i) $\Pi^*$ is a *refinement* of $\Pi$ iff $\Pi \subseteq \Pi^*$.

(ii) The *common refinement* of $\Pi_1$ and $\Pi_2$ is $\Pi_1 \cup \Pi_2$.

**Theorem 2.1.4.** *Refinement Monotonicity*
*If $\Pi^*$ is a refinement of $\Pi$, then $L(\Pi, f, \alpha) \leq L(\Pi^*, f, \alpha)$ and $U(\Pi^*, f, \alpha) \leq U(\Pi, f, \alpha)$.*

*Proof.* It suffices to add one point $x^* \in (x_{i-1}, x_i)$. If $w_1, w_2$ are infima on $[x_{i-1}, x^*]$ and $[x^*, x_i]$, then $w_1, w_2 \geq m_i$ and $L(\Pi^*, f, \alpha) - L(\Pi, f, \alpha) = (w_1 - m_i)\big[\alpha(x^*) - \alpha(x_{i-1})\big] + (w_2 - m_i)\big[\alpha(x_i) - \alpha(x^*)\big] \geq 0$.
Iterate for additional points. The upper-sum inequality is the same argument with suprema.
*Structural Commentary*: Refinement raises lower sums and lowers upper sums.
*Key Insight*: Splitting an interval tightens the local oscillation bounds. □

**Perspective 2.1.5.** *Piecewise-Constant Envelopes*
A partition $\Pi$ gives lower and upper step envelopes with values $m_i$ and $M_i$; their integrals are exactly $L(\Pi, f)$ and $U(\Pi, f)$. Common refinements make these envelope sums compatible across partitions.

**Theorem 2.1.6.** *Lower Integral Below Upper Integral*
*For every bounded real $f$ and increasing $\alpha$ on $[a,b]$, $\underline{\int_a^b f \, d\alpha} \leq \overline{\int_a^b f \, d\alpha}$.*

*Proof.* Given partitions $\Pi_1, \Pi_2$, let $\Pi^* = \Pi_1 \cup \Pi_2$.
By Theorem 2.1.4, $L(\Pi_1, f, \alpha) \leq L(\Pi^*, f, \alpha) \leq U(\Pi^*, f, \alpha) \leq U(\Pi_2, f, \alpha)$.
Take the supremum over $\Pi_1$ and then the infimum over $\Pi_2$.
*Structural Commentary*: Common refinements compare sums from different partitions.
*Key Insight*: The lower envelope never rises above the upper envelope. □

**Theorem 2.1.7.** *Upper–Lower Criterion*
*A bounded real $f$ is in $\mathcal{R}(\alpha)$ on $[a,b]$ if and only if $\forall \varepsilon > 0 \; \exists$ a partition $\Pi$ with $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) < \varepsilon$.*

*Proof.*

($\Rightarrow$) If $f \in \mathcal{R}(\alpha)$, choose $\Pi_1, \Pi_2$ whose lower and upper sums are within $\varepsilon/2$ of $\int_a^b f \, d\alpha$.
Their common refinement $\Pi$ satisfies the displayed inequality by Theorem 2.1.4.

($\Leftarrow$) If such $\Pi$ exist, then $L(\Pi, f, \alpha) \leq \underline{\int_a^b f \, d\alpha} \leq \overline{\int_a^b f \, d\alpha} \leq U(\Pi, f, \alpha)$ forces upper and lower integrals to agree.

*Structural Commentary*: Integrability is exactly the ability to make one partition squeeze the oscillation.
*Key Insight*: The gap $U - L$ is the quantitative defect of integrability. □

**Perspective 2.1.8.** *Darboux Gaps and Riemann Sums*
For $\alpha(x) = x$, a small Darboux gap $U(\Pi, f) - L(\Pi, f)$ traps every tagged sum for $\Pi$ near the integral. This is the local reason the Darboux definition above and the tagged Riemann-sum definition give the same number. The same estimate works with Stieltjes weights $\Delta\alpha_i$ whenever the upper–lower gap is small.

**Theorem 2.1.9.** *Tagged Sums*
*Let $\Pi = \{x_0, \dots, x_n\}$ satisfy $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) < \varepsilon$, and choose tags $t_i \in [x_{i-1}, x_i]$.*

*(i) If $s_i, t_i \in [x_{i-1}, x_i]$, then $\sum_{i=1}^{n} |f(s_i) - f(t_i)|\,\Delta\alpha_i < \varepsilon$.*

*(ii) If $f \in \mathcal{R}(\alpha)$, then $\left|\sum_{i=1}^{n} f(t_i)\Delta\alpha_i - \int_a^b f\,d\alpha\right| < \varepsilon$.*

*Proof.*

(i) Since $|f(s_i) - f(t_i)| \le M_i - m_i$, summing gives $\sum_{i=1}^{n} |f(s_i) - f(t_i)|\,\Delta\alpha_i \le U(\Pi, f, \alpha) - L(\Pi, f, \alpha) < \varepsilon$.

(ii) The tagged sum and $\int_a^b f\,d\alpha$ both lie between $L(\Pi, f, \alpha)$ and $U(\Pi, f, \alpha)$, so their distance is $< \varepsilon$.

*Structural Commentary*: A small Darboux gap makes tags and the integral close.
*Key Insight*: Tags are harmless after oscillation has been controlled. □

**Theorem 2.1.10.** *Continuous Functions Are Riemann–Stieltjes Integrable*
*If $f$ is continuous on $[a, b]$ and $\alpha$ is increasing on $[a, b]$, then $f \in \mathcal{R}(\alpha)$.*

*Proof.* If $\alpha(b) = \alpha(a)$, the Darboux gap is zero.
Otherwise choose $\eta > 0$ with $\eta[\alpha(b) - \alpha(a)] < \varepsilon$. Uniform continuity gives partition $\Pi$ with $M_i - m_i \le \eta\ \forall i$, so $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) = \sum_{i=1}^{n} (M_i - m_i)\Delta\alpha_i \le \eta[\alpha(b) - \alpha(a)] < \varepsilon$. Apply Theorem 2.1.7.
*Structural Commentary*: Uniform continuity controls oscillation uniformly across the partition.
*Key Insight*: Continuity turns local small intervals into a small total Darboux gap. □

**Theorem 2.1.11.** *Monotone Functions and Continuous Integrators*
*If $f$ is monotone on $[a, b]$ and $\alpha$ is continuous and increasing on $[a, b]$, then $f \in \mathcal{R}(\alpha)$.*

*Proof.* Assume first that $f$ is increasing. If $\alpha(b) = \alpha(a)$, then every $\Delta\alpha_i = 0$, so $U(\Pi, f, \alpha) = L(\Pi, f, \alpha) = 0$ for every partition $\Pi$. Choose $n$ and a partition $\Pi$ such that $\Delta\alpha_i = [\alpha(b) - \alpha(a)]/n$ for all $i$.
Existence follows from continuity of $\alpha$ and Theorem 5.1.41.
Then $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) = \frac{\alpha(b)-\alpha(a)}{n} \sum_{i=1}^{n} \big(f(x_i) - f(x_{i-1})\big) = \frac{\alpha(b)-\alpha(a)}{n}\big(f(b) - f(a)\big) < \varepsilon$ for $n$ large.
If $f$ is decreasing, use the same partition. Then the suprema and infima on $[x_{i-1}, x_i]$ are $f(x_{i-1})$ and $f(x_i)$, so the same computation gives $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) = \frac{\alpha(b)-\alpha(a)}{n}\big(f(a) - f(b)\big)$, which is $< \varepsilon$ for large $n$.
*Structural Commentary*: Equal $\alpha$-increments convert the monotone oscillations into a telescoping sum.
*Key Insight*: A monotone function's total oscillation is finite and can be spread thinly. □




Figure 161: Isolate the discontinuity in a short interval; control the compact pieces outside it.

**Theorem 2.1.12.** *Finitely Many Discontinuities*
*Suppose $f$ is bounded on $[a, b]$, the discontinuity set $E := \{x \in [a, b] : f \text{ is discontinuous at } x\}$ is finite, and $\alpha$ is continuous at each point of $E$. Then $f \in \mathcal{R}(\alpha)$.*

*Proof.* Let $M = \sup_{[a,b]} |f|$ and let $\eta > 0$. By continuity of $\alpha$ on the finite set $E$, choose disjoint intervals $I_j = [u_j, v_j]$ covering $E$, with each point of $E \cap (a, b)$ in some $(u_j, v_j)$, such that $\sum_j \big(\alpha(v_j) - \alpha(u_j)\big) < \eta$.
Set $K = [a, b] \setminus \bigcup_j (u_j, v_j)$. Then $f|_K$ is uniformly continuous by Theorem 5.2.7.
Choose a partition $\Pi$ containing all $u_j, v_j$, with no points inside the intervals $(u_j, v_j)$, and fine enough on $K$ that $M_i - m_i \le \eta$ whenever $[x_{i-1}, x_i] \subset K$.
Hence $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) \le \eta\big[\alpha(b) - \alpha(a)\big] + 2M \sum_j \big(\alpha(v_j) - \alpha(u_j)\big) < \eta\big[\alpha(b) - \alpha(a) + 2M\big]$.

Since $\eta$ is arbitrary, Theorem 2.1.7 applies.
*Structural Commentary*: Isolate bad points in small $\alpha$-intervals; use uniform continuity elsewhere.
*Key Insight*: Finitely many discontinuities do not matter unless $\alpha$ also jumps there. □

**Example 2.1.13.** *Dirichlet's Function Is Not Riemann Integrable*
Let $g = \mathbf{1}_{\mathbb{Q}}$ on $[0,1]$. Every nondegenerate subinterval contains rationals and irrationals by Corollary 1.1.22, so for every partition $\Pi$, $M_i = 1$ and $m_i = 0$ on each nonzero-width subinterval.
Thus $U(\Pi, g) = 1$ and $L(\Pi, g) = 0$, hence $g \notin \mathcal{R}$.

**Remark 2.1.14.** *Small Discontinuity Sets*
The same idea underlies Theorem 1.5.2: small discontinuity sets, later called measure-zero sets in Definition 4.1.13, do not obstruct integrability. Dirichlet's function fails because every point is discontinuous.

**Theorem 2.1.15.** *Continuous Composition Preserves Integrability*
*Let $\alpha$ be increasing on $[a,b]$. Suppose $f \in \mathcal{R}(\alpha)$, $m \le f(x) \le M$ on $[a,b]$, and $\phi : [m,M] \to \mathbb{R}$ is continuous. Define $h : [a,b] \to \mathbb{R}$ by $h(x) := \phi(f(x))$. Then $h \in \mathcal{R}(\alpha)$.*

*Proof.* Let $C := \sup_{y\in[m,M]} |\phi(y)|$ and let $\varepsilon > 0$. Choose $\eta > 0$ such that $\eta[\alpha(b) - \alpha(a) + 2C] < \varepsilon$. Since $\phi$ is uniformly continuous on $[m,M]$, choose $\gamma > 0$ such that $|s-t| < \gamma \implies |\phi(s) - \phi(t)| < \eta$ $(s,t \in [m,M])$.
Choose a partition $\Pi = \{x_0, \dots, x_n\}$ with $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) < \gamma\eta$.
Let $M_i, m_i$ denote upper and lower values for $f$ on $[x_{i-1}, x_i]$, $M_i^h, m_i^h$ denote corresponding values for $h$.
Split $A := \{i : M_i - m_i < \gamma\}$, $B := \{i : M_i - m_i \ge \gamma\}$. For $i \in A$, $M_i^h - m_i^h \le \eta$; for $i \in B$, $M_i^h - m_i^h \le 2C$.
Also $\gamma \sum_{i\in B} \Delta\alpha_i \le \sum_{i\in B} (M_i - m_i)\Delta\alpha_i \le U(\Pi, f, \alpha) - L(\Pi, f, \alpha) < \gamma\eta$, so $\sum_{i\in B} \Delta\alpha_i < \eta$.
$U(\Pi, h, \alpha) - L(\Pi, h, \alpha) \le \eta \sum_{i\in A} \Delta\alpha_i + 2C \sum_{i\in B} \Delta\alpha_i < \eta[\alpha(b) - \alpha(a) + 2C] < \varepsilon$. Theorem 2.1.7 gives $h \in \mathcal{R}(\alpha)$.
*Structural Commentary*: Integrability survives continuous remeasurement of the values of $f$.
*Key Insight*: Large oscillation intervals have small total $\alpha$-weight. □

**Theorem 2.1.16.** *Basic Properties of the Integral*
*Let all functions be bounded and real on $[a,b]$, and let all integrators be increasing.*

*(i) If $f_1, f_2 \in \mathcal{R}(\alpha)$, $c \in \mathbb{R}$, then $f_1 + f_2, cf_1 \in \mathcal{R}(\alpha)$, and $\int_a^b (f_1 + f_2)\, d\alpha = \int_a^b f_1\, d\alpha + \int_a^b f_2\, d\alpha$; $\int_a^b cf_1\, d\alpha = c\int_a^b f_1\, d\alpha$.*

*(ii) If $f_1 \le f_2$ and both functions are in $\mathcal{R}(\alpha)$, then $\int_a^b f_1\, d\alpha \le \int_a^b f_2\, d\alpha$.*

*(iii) If $f \in \mathcal{R}(\alpha)$ and $a < c < b$, then $f$ is integrable on $[a,c]$ and $[c,b]$, and $\int_a^c f\, d\alpha + \int_c^b f\, d\alpha = \int_a^b f\, d\alpha$.*

*(iv) If $f \in \mathcal{R}(\alpha)$ and $|f| \le M$, then $\left|\int_a^b f\, d\alpha\right| \le M[\alpha(b) - \alpha(a)]$*

*(v) If $f \in \mathcal{R}(\alpha_1)$ and $f \in \mathcal{R}(\alpha_2)$, then $f \in \mathcal{R}(\alpha_1 + \alpha_2)$ and $\int_a^b f\, d(\alpha_1 + \alpha_2) = \int_a^b f\, d\alpha_1 + \int_a^b f\, d\alpha_2$. If $c > 0$ and $f \in \mathcal{R}(\alpha)$, then $f \in \mathcal{R}(c\alpha)$ and $\int_a^b f\, d(c\alpha) = c\int_a^b f\, d\alpha$.*

*Proof.*
(i) $L(\Pi, f_1, \alpha) + L(\Pi, f_2, \alpha) \le L(\Pi, f_1 + f_2, \alpha) \le U(\Pi, f_1 + f_2, \alpha) \le U(\Pi, f_1, \alpha) + U(\Pi, f_2, \alpha)$, where $\Pi$ is common refinement, and Theorem 2.1.7; scalar multiplication is the same estimate with signs handled by swapping upper and lower sums when $c < 0$.
(ii) If $f_1 \le f_2$, then $L(\Pi, f_2 - f_1, \alpha) \ge 0$ for every $\Pi$; apply (i) to $f_2 - f_1$.
(iii) Refine so that $c \in \Pi$. Then the lower and upper sums split into the sums over $[a,c]$ and $[c,b]$.
(iv) Apply (ii) to $-M \le f \le M$ and use the integrals of the constant functions $\pm M$.
(v) Since $\Delta(\alpha_1 + \alpha_2)_i = \Delta\alpha_{1,i} + \Delta\alpha_{2,i}$ and $\Delta(c\alpha)_i = c\,\Delta\alpha_i$, corresponding lower and upper sums split exactly.
*Structural Commentary*: The algebra of integrals is inherited from the algebra of upper and lower sums.
*Key Insight*: Common refinements let several integrable functions be approximated by the same partition. □

**Theorem 2.1.17.** *Products and Absolute Values*
*If $f, g \in \mathcal{R}(\alpha)$ on $[a,b]$, then $fg \in \mathcal{R}(\alpha)$, $|f| \in \mathcal{R}(\alpha)$, and $\left|\int_a^b f\, d\alpha\right| \le \int_a^b |f|\, d\alpha$.*

*Proof.* By Theorem 2.1.16, $f \pm g \in \mathcal{R}(\alpha)$. Applying Theorem 2.1.15 to $\phi(t) = t^2$ gives $(f \pm g)^2 \in \mathcal{R}(\alpha)$, hence $fg \in \mathcal{R}(\alpha)$ from $4fg = (f+g)^2 - (f-g)^2$.
Applying the same theorem to $\phi(t) = |t|$ gives $|f| \in \mathcal{R}(\alpha)$, and $-|f| \le f \le |f|$ gives the inequality.
*Structural Commentary*: Products reduce to squares; absolute values to composition.
*Key Insight*: Once continuous composition is available, the usual algebra closes. □

## 2.2 Fundamental Theorems of Calculus

**Theorem 2.2.1.** *Integral Functions*
*Let $f \in \mathcal{R}$ on $[a,b]$ and define $F(x) := \int_a^x f(t)\,dt$. Then $F$ is continuous.*
*If $f$ is continuous at $x_0$, then $F'(x_0) = f(x_0)$, with one-sided derivatives at endpoints.*

*Proof.* Let $|f| \le M$. For $x < y$, Theorem 2.1.16 gives $|F(y) - F(x)| = \left|\int_x^y f(t)\,dt\right| \le M(y-x)$ so $F$ continuous.
If $f$ continuous at $x_0$, for $x \ne x_0$ near $x_0$, $\left|\frac{F(x)-F(x_0)}{x-x_0} - f(x_0)\right| \le \sup_{u\in[x,x_0]} |f(u) - f(x_0)| \to 0$. $F'(x_0) = f(x_0)$.
*Structural Commentary*: Boundedness controls increments; local continuity gives the derivative.
*Key Insight*: Averaging a continuous function over shrinking intervals returns its value. □

**Theorem 2.2.2.** *Fundamental Theorem of Calculus*
*If $f \in \mathcal{R}$ on $[a,b]$ and $\exists$ a differentiable function $F$ on $[a,b]$ such that $F' = f$, then $\int_a^b f(x)\,dx = F(b) - F(a)$.*

*Proof.* Choose $\Pi = \{x_0, \dots, x_n\}$ with $U(\Pi, f) - L(\Pi, f) < \varepsilon$.
By MVT, choose $t_i \in [x_{i-1}, x_i]$ so that $F(b) - F(a) = \sum_i \big(F(x_i) - F(x_{i-1})\big) = \sum_i f(t_i)\Delta x_i$.
Theorem 2.1.9 puts this tagged sum within $\varepsilon$ of $\int_a^b f$. Let $\varepsilon \to 0$.
*Structural Commentary*: Mean-value increments convert antiderivatives into tagged Riemann sums.
*Key Insight*: The integral is the common limit of all sufficiently fine tagged sums. □

**Theorem 2.2.3.** *Integration by Parts*
*Suppose $F, G$ are differentiable on $[a,b]$, $F' = f \in \mathcal{R}$, and $G' = g \in \mathcal{R}$.*
*Then $\int_a^b F(x)g(x)\,dx = F(b)G(b) - F(a)G(a) - \int_a^b f(x)G(x)\,dx$.*

*Proof.* Set $H := FG$. Since $F, G$ are continuous, $F, G \in \mathcal{R}$; hence $Fg, fG \in \mathcal{R}$ by Theorem 2.1.17.
Thus $H' = Fg + fG \in \mathcal{R}$, and Theorem 2.2.2 gives $F(b)G(b) - F(a)G(a) = H(b) - H(a) = \int_a^b Fg + \int_a^b fG$.
*Structural Commentary*: Integration by parts is the product rule integrated over the interval.
*Key Insight*: Move one derivative from $G$ onto $F$ by applying the fundamental theorem to $FG$. □

## 2.3 Step Integrators and Change of Variable

**Definition 2.3.1.** *Unit Step Function*
The *unit step function* $I$ is $I(x) = \begin{cases} 0, & x \le 0, \\ 1, & x > 0. \end{cases}$

**Theorem 2.3.2.** *Single Jump Integrator*
*If $a < s < b$, $f$ is bounded on $[a,b]$ and continuous at $s$, and $\alpha(x) = I(x - s)$, then $\int_a^b f\,d\alpha = f(s)$.*

*Proof.* Use partitions $\Pi = \{a, s, x_2, b\}$ with $s < x_2 < b$. The only nonzero $\Delta\alpha_i$ occurs on $[s, x_2]$, so $L(\Pi, f, \alpha)$ and $U(\Pi, f, \alpha)$ are the infimum and supremum of $f$ on $[s, x_2]$. As $x_2 \downarrow s$, continuity at $s$ squeezes both to $f(s)$.
*Structural Commentary*: A jump of $\alpha$ makes the integral sample $f$ at the jump.
*Key Insight*: Stieltjes integration includes point masses without changing the Darboux framework. □

**Theorem 2.3.3.** *Pure Step Integrators Give Series*
*Suppose $c_n \ge 0$, $\sum c_n$ converges, and $(s_n)$ is a sequence of distinct points in $(a,b)$.*
*Define $\alpha(x) := \sum_{n=1}^{\infty} c_n I(x - s_n)$. If $f$ is continuous on $[a,b]$, then $\int_a^b f\,d\alpha = \sum_{n=1}^{\infty} c_n f(s_n)$.*

*Proof.* By comparison, $\alpha$ is increasing. For $N \in \mathbb{N}$, put $\alpha_N(x) := \sum_{n=1}^{N} c_n I(x - s_n)$ and $\beta_N := \alpha - \alpha_N$..
Theorems 2.3.2 and 2.1.16 give $\int_a^b f\,d\alpha_N = \sum_{n=1}^{N} c_n f(s_n)$. Also $f \in \mathcal{R}(\beta_N)$ by Theorem 2.1.10.
If $|f| \le M$, then Theorem 2.1.16 gives $\left|\int_a^b f\,d\beta_N\right| \le M[\beta_N(b) - \beta_N(a)] = M \sum_{n>N} c_n$.
Letting $N \to \infty$ and using additivity in the integrator gives the formula.
*Structural Commentary*: Finite step integrators sum point samples; summable tails vanish by the integral bound.
*Key Insight*: A Stieltjes integral can be an ordinary series in disguise. □




Figure 162: A step integrator concentrates mass at jumps: $\int f\,d\alpha = \sum_i c_i f(s_i)$.

**Theorem 2.3.4.** *Differentiable Integrators Reduce to Riemann Integrals*
*Assume $\alpha$ increases monotonically, $\alpha' \in \mathcal{R}$ on $[a,b]$, and $f$ is bounded and real on $[a,b]$.*
*Then $f \in \mathcal{R}(\alpha) \iff f\alpha' \in \mathcal{R}$, and in that case $\int_a^b f\,d\alpha = \int_a^b f(x)\alpha'(x)\,dx$.*

*Proof.* Let $M = \sup_{[a,b]} |f|$. Choose $\Pi$ with $U(\Pi,\alpha') - L(\Pi,\alpha') < \varepsilon$.
For any refinement and any tags $s_i \in [x_{i-1},x_i]$, MVT gives points $t_i \in [x_{i-1},x_i]$ with $\Delta\alpha_i = \alpha'(t_i)\Delta x_i$.
Hence $|\sum_i f(s_i)\Delta\alpha_i - \sum_i f(s_i)\alpha'(s_i)\Delta x_i| \le M\sum_i |\alpha'(t_i) - \alpha'(s_i)|\,\Delta x_i \le M\varepsilon$. Thus every tagged Stieltjes sum for $f\,d\alpha$ is within $M\varepsilon$ of the corresponding tagged Riemann sum for $f\alpha'\,dx$, and conversely. By the tagged-sum criterion, one family has a limit exactly when the other does, and the two limits are equal.
*Structural Commentary*: The increment $d\alpha$ is replaced by $\alpha'(x)\,dx$ when $\alpha'$ is Riemann integrable.
*Key Insight*: The mean value theorem turns Stieltjes increments into ordinary Riemann weights. □

**Theorem 2.3.5.** *Change of Variable*
*Let $\varphi : [A,B] \to [a,b]$ be strictly increasing, continuous, and onto.*
*Let $\alpha$ be increasing on $[a,b]$, $f \in \mathcal{R}(\alpha)$, and set $\beta(y) := \alpha(\varphi(y))$ and $g(y) := f(\varphi(y))$.*
*Then $g \in \mathcal{R}(\beta)$ and $\int_A^B g\,d\beta = \int_a^b f\,d\alpha$. If also $\alpha(x) = x$ and $\varphi' \in \mathcal{R}$, then $\int_a^b f(x)\,dx = \int_A^B f(\varphi(y))\varphi'(y)\,dy$.*

*Proof.* Partitions $Q = \{y_i\}$ of $[A,B]$ correspond to partitions $\Pi = \{\varphi(y_i)\}$ of $[a,b]$. On corresponding intervals, $g$ has same range as $f$ and $\Delta\beta_i = \Delta\alpha_i$, so $U(Q,g,\beta) = U(\Pi,f,\alpha)$ and $L(Q,g,\beta) = L(\Pi,f,\alpha)$. The upper–lower criterion gives the Stieltjes formula. If $\alpha(x) = x$, then $\beta = \varphi$; apply Theorem 2.3.4 to the left side.
*Structural Commentary*: A monotone change of variable relabels the same partition intervals.
*Key Insight*: Stieltjes change of variable is equality of Darboux sums under reparametrization. □

## 2.4 Improper Integrals

**Definition 2.4.1.** *Improper Riemann Integrals*
Let $f \in \mathcal{R}$ on every compact subinterval of its domain.
Set $\int_a^b f(x)\,dx := \lim_{d\to b^-} \int_a^d f(x)\,dx$ at a right endpoint, and $\int_a^b f(x)\,dx := \lim_{c\to a^+} \int_c^b f(x)\,dx$ at a left endpoint.
If both endpoints are improper, choose $\xi \in (a,b)$ and set $\int_a^b f(x)\,dx := \int_a^\xi f(x)\,dx + \int_\xi^b f(x)\,dx$, provided both one-sided integrals exist and the sum is not indeterminate.
A finite value means the improper integral *converges*; otherwise it diverges to $+\infty$ or $-\infty$.




Figure 163: Improper integrals are limits of ordinary integrals over compact truncations.

**Theorem 2.4.2.** *Nonnegative Improper Integrals*
*Let $f \ge 0$ and $f \in \mathcal{R}$ on every $[a,d]$, $a < d < b$. Then $\int_a^b f(x)\,dx = \sup\left\{\int_a^d f(x)\,dx : a < d < b\right\}$.*
*Thus $\int_a^b f$ exists in $[0,+\infty]$ and converges exactly when this supremum is finite.*

*Proof.* For $a < d_1 < d_2 < b$, $\int_a^{d_2} f - \int_a^{d_1} f = \int_{d_1}^{d_2} f \ge 0$.
Hence $d \mapsto \int_a^d f$ is increasing, so its extended limit exists and equals its supremum.
*Structural Commentary*: Nonnegative area can only accumulate.
*Key Insight*: Improper convergence for $f \ge 0$ is boundedness of all finite truncations. □

**Perspective 2.4.3.** *Principal Value Is Different*
Improper integrals over $(-\infty,\infty)$ require separate one-sided limits.
A symmetric limit such as $\lim_{A\to\infty}\int_{-A}^{A}\sin x\,dx$ may still exist (Cauchy principal value, not Definition 2.4.1).

**Definition 2.4.4.** *Improper Riemann–Stieltjes Integrals on $\mathbb{R}$*
Let $\alpha:\mathbb{R}\to\mathbb{R}$ be increasing and $f\in\mathcal{R}(\alpha)$ on every finite interval.
Define, when the limits exist, $\int_0^\infty f\,d\alpha := \lim_{b\to\infty}\int_0^b f\,d\alpha$; and $\int_{-\infty}^0 f\,d\alpha := \lim_{a\to-\infty}\int_a^0 f\,d\alpha$.
If their sum is not indeterminate, set $\int_{-\infty}^\infty f\,d\alpha := \int_{-\infty}^0 f\,d\alpha + \int_0^\infty f\,d\alpha$.
Finite value means $f$ is *$\alpha$-integrable on $\mathbb{R}$*. Write $\alpha(\pm\infty) := \lim_{t\to\pm\infty}\alpha(t)$ when these extended limits are used.

**Theorem 2.4.5.** *Nonnegative and Bounded-Mass Stieltjes Integrals*
*Let $\alpha:\mathbb{R}\to\mathbb{R}$ be increasing, and suppose $f\in\mathcal{R}(\alpha)$ on every finite interval.*

*(i) If $f\ge 0$, then either $f$ is $\alpha$-integrable on $\mathbb{R}$ or $\int_{-\infty}^\infty f\,d\alpha=+\infty$.*
*(ii) If $-\infty<\alpha(-\infty)<\alpha(\infty)<+\infty$ and $f$ is bounded, then $f$ is $\alpha$-integrable on $\mathbb{R}$.*

*Proof.*

(i) By additivity and order preservation, $\int_0^b f\,d\alpha$ and $\int_a^0 f\,d\alpha$ are monotone under outward truncation. Hence both extended one-sided limits exist, so their sum is finite or $+\infty$.

(ii) Let $|f|\le M$. For finite $a<b$, Theorem 2.1.16 gives $\left|\int_a^b f\,d\alpha\right|\le M\big[\alpha(b)-\alpha(a)\big]$. Since $\alpha(\infty)$ and $\alpha(-\infty)$ are finite, the right and left tails have vanishing $\alpha$-increment. Thus the one-sided truncations are Cauchy and have finite limits.

*Structural Commentary*: Positivity gives monotone limits; finite mass bounds tails.
*Key Insight*: A bounded increasing integrator behaves like finite total mass. □

## 2.5 Vector-Valued Integrals and Curves

**Definition 2.5.1.** *Vector-Valued Riemann–Stieltjes Integral*
For $\mathbf{f}=(f_1,\dots,f_k):[a,b]\to\mathbb{R}^k$, define $\mathbf{f}\in\mathcal{R}(\alpha)$ iff $f_j\in\mathcal{R}(\alpha)\ \forall j$, and set $\int_a^b\mathbf{f}\,d\alpha := \left(\int_a^b f_1\,d\alpha,\dots,\int_a^b f_k\,d\alpha\right)$.

**Theorem 2.5.2.** *Vector-Valued Integral Results*

*(i) If $\mathbf{f},\mathbf{F}:[a,b]\to\mathbb{R}^k$, $\mathbf{f}\in\mathcal{R}$, and $\mathbf{F}'=\mathbf{f}$, then $\int_a^b\mathbf{f}(t)\,dt=\mathbf{F}(b)-\mathbf{F}(a)$.*
*(ii) If $\mathbf{f}:[a,b]\to\mathbb{R}^k$ and $\mathbf{f}\in\mathcal{R}(\alpha)$, then $\|\mathbf{f}\|\in\mathcal{R}(\alpha)$ and $\left\|\int_a^b\mathbf{f}\,d\alpha\right\|\le\int_a^b\|\mathbf{f}\|\,d\alpha$*

*Proof.*

(i) Write $\mathbf{f}=(f_1,\dots,f_k)$ and $\mathbf{F}=(F_1,\dots,F_k)$. For each $j$, Theorem 2.2.2 gives $\int_a^b f_j(t)\,dt=F_j(b)-F_j(a)$. The vector identity follows componentwise.

(ii) Write $\mathbf{f}=(f_1,\dots,f_k)$. Since $\|\mathbf{f}\|=(f_1^2+\cdots+f_k^2)^{1/2}$, Theorems 2.1.17 and 2.1.15 give $\|\mathbf{f}\|\in\mathcal{R}(\alpha)$. Let $\mathbf{y}:=\int_a^b\mathbf{f}\,d\alpha$. If $\mathbf{y}=0$, the inequality is immediate.
Otherwise, $\|\mathbf{y}\|^2=\sum_{j=1}^k y_j\int_a^b f_j\,d\alpha=\int_a^b\sum_{j=1}^k y_jf_j\,d\alpha\le\int_a^b\|\mathbf{y}\|\,\|\mathbf{f}\|\,d\alpha=\|\mathbf{y}\|\int_a^b\|\mathbf{f}\|\,d\alpha$, by Theorem 1.1.13 and order preservation. Divide by $\|\mathbf{y}\|$.

*Structural Commentary*: Coordinatewise integration; Schwarz controls norms.
*Key Insight*: Integrate the projection of $\mathbf{f}$ onto the resulting vector. □

**Definition 2.5.3.** *Curves and Rectifiability*
A *curve* in $\mathbb{R}^k$ is a continuous map $\gamma:[a,b]\to\mathbb{R}^k$; it is an *arc* if one-to-one and a *closed curve* if $\gamma(a)=\gamma(b)$.
Curves are maps, not just ranges. For a partition $\Pi=\{x_0,\dots,x_n\}$, set $\ell(\Pi,\gamma):=\sum_{i=1}^n\|\gamma(x_i)-\gamma(x_{i-1})\|$.
The *length* is $\ell(\gamma):=\sup_\Pi\ell(\Pi,\gamma)$; if $\ell(\gamma)<\infty$, $\gamma$ is *rectifiable*.




Figure 164: A staircase path can approach the diagonal in position while its polygonal length stays 2, not $\sqrt{2}$.

**Perspective 2.5.4.** *Length Is Not Just Shape*
Length is a supremum over partitions of one curve. Passing lengths through curve limits needs more than positional convergence; Theorem 2.5.5 uses $C^1$ speed control.

**Theorem 2.5.5.** *Length of a Continuously Differentiable Curve*
*If $\gamma' : [a,b] \to \mathbb{R}^k$ is continuous, then $\gamma$ is rectifiable and $\ell(\gamma) = \int_a^b \|\gamma'(t)\|\, dt$.*

*Proof.* For $\Pi = \{x_0, \dots, x_n\}$, Theorem 2.5.2(i) gives $\|\gamma(x_i) - \gamma(x_{i-1})\| = \left\| \int_{x_{i-1}}^{x_i} \gamma'(t)\, dt \right\| \le \int_{x_{i-1}}^{x_i} \|\gamma'(t)\|\, dt$.
Hence $\ell(\Pi, \gamma) \le \int_a^b \|\gamma'\|$. Conversely, choose $\Pi$ so fine that $\|\gamma'(s) - \gamma'(t)\| < \varepsilon$ on each subinterval.
Choose $t_i \in [x_{i-1}, x_i]$. Then $\int_{x_{i-1}}^{x_i} \|\gamma'(t)\|\, dt \le \|\gamma'(t_i)\|\, \Delta x_i + \varepsilon \Delta x_i$.
Also $\gamma(x_i) - \gamma(x_{i-1}) = \int_{x_{i-1}}^{x_i} \gamma'(t)\, dt$, so $\|\gamma'(t_i)\|\, \Delta x_i \le \|\gamma(x_i) - \gamma(x_{i-1})\| + \varepsilon \Delta x_i$.
Thus $\int_{x_{i-1}}^{x_i} \|\gamma'(t)\|\, dt \le \|\gamma(x_i) - \gamma(x_{i-1})\| + 2\varepsilon \Delta x_i$.
Summing gives $\int_a^b \|\gamma'\| \le \ell(\gamma) + 2\varepsilon(b-a)$; let $\varepsilon \downarrow 0$.
*Structural Commentary*: Upper bounds are immediate; uniform continuity gives matching lower bounds.
*Key Insight*: For $C^1$ curves, length is accumulated speed. □

## 2.6 Further Topics in Integration

**Method 2.6.1.** *Integration Proof Patterns*

(i) *Definitions*: fix $\Pi = \{x_i\}_{i=0}^n$ before defining $M_i, m_i, \Delta x_i, \Delta \alpha_i$.
(ii) *Refinement*: prove the one-point insertion case; iterate.
(iii) *Integrability*: find $\Pi$ with $U(\Pi, f, \alpha) - L(\Pi, f, \alpha) < \varepsilon$.
(iv) *Bad sets*: cover $E$ by $I_j = [u_j, v_j]$ with $\sum_j [\alpha(v_j) - \alpha(u_j)] < \varepsilon$; control $K = [a,b] \setminus \bigcup_j (u_j, v_j)$ uniformly.
(v) *Properties*: pass to a common refinement before comparing sums.
(vi) *Calculus*: use the mean value theorem to form tagged sums, then apply Theorem 2.1.9.
(vii) *Improper integrals*: truncate endpoints; use monotonicity or tail bounds.

*Key Insight*: Riemann–Stieltjes integration is Darboux approximation plus a choice of interval weights $\Delta \alpha_i$.

# 3 Several-Variable Differentiation

## 3.1 Partial and Total Derivatives

Several-variable differentiation studies best linear approximations and local geometry, using linear maps, operator norms, determinants, invertibility, and metric estimates.

**Definition 3.1.1.** *Total Derivative*
Let $E \subseteq \mathbb{R}^n$ be open, $\mathbf{f} : E \to \mathbb{R}^m$, $\mathbf{x} \in E$, and $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ denote space of linear maps (Definition 1.5.1).
If $\exists A \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ s.t. $\lim_{\mathbf{h} \to \mathbf{0}} \frac{\|\mathbf{f}(\mathbf{x}+\mathbf{h}) - \mathbf{f}(\mathbf{x}) - A\mathbf{h}\|_2}{\|\mathbf{h}\|_2} = 0$, then $\mathbf{f}$ is *differentiable at* $\mathbf{x}$, and $A$ denoted $D\mathbf{f}(\mathbf{x})$ or $\mathbf{f}'(\mathbf{x})$.
Equivalently, $\mathbf{f}(\mathbf{x}+\mathbf{h}) - \mathbf{f}(\mathbf{x}) = D\mathbf{f}(\mathbf{x})\mathbf{h} + \mathbf{r}(\mathbf{h})$, and $\frac{\|\mathbf{r}(\mathbf{h})\|_2}{\|\mathbf{h}\|_2} \to 0$.
If this holds for every $\mathbf{x} \in E$, then $\mathbf{f}$ is differentiable on $E$.

**Perspective 3.1.2.** *Derivative First, Matrix Second*
The total derivative is a linear map first; the Jacobian is its matrix in chosen bases. This keeps coordinate formulas subordinate to the intrinsic estimate $\|A\mathbf{h}\|_2 \le \|A\|_{\mathrm{op}} \|\mathbf{h}\|_2$ from Definition 5.1.4.




Figure 165: $D\mathbf{f}(\mathbf{x})\mathbf{h}$ is the linear part; $\mathbf{r}(\mathbf{h}) = o(\|\mathbf{h}\|_2)$ is the error.

**Theorem 3.1.3.** *Basic Consequences of Total Differentiability*
*If $\mathbf{f} : E \to \mathbb{R}^m$ is differentiable at $\mathbf{x} \in E$, then $D\mathbf{f}(\mathbf{x})$ is unique and $\mathbf{f}$ is continuous at $\mathbf{x}$.*

*Proof.* If $A_1, A_2$ both satisfy Definition 3.1.1, set $B = A_1 - A_2$. Then $\|B\mathbf{h}\|_2 / \|\mathbf{h}\|_2 \to 0$ as $\mathbf{h} \to \mathbf{0}$.
For fixed $\mathbf{h} \neq \mathbf{0}$, applying this to $t\mathbf{h}$ and using linearity makes the quotient independent of $t$, so $B\mathbf{h} = \mathbf{0}$.
Hence $B = 0$. For continuity, write $\mathbf{f}(\mathbf{x}+\mathbf{h}) - \mathbf{f}(\mathbf{x}) = D\mathbf{f}(\mathbf{x})\mathbf{h} + \mathbf{r}(\mathbf{h})$.
The operator-norm estimate gives $\|D\mathbf{f}(\mathbf{x})\mathbf{h}\|_2 \le \|D\mathbf{f}(\mathbf{x})\|_{\text{op}} \|\mathbf{h}\|_2$, and remainder is $o(\|\mathbf{h}\|_2)$, so $\mathbf{f}(\mathbf{x}+\mathbf{h}) \to \mathbf{f}(\mathbf{x})$.
*Structural Commentary*: The derivative is a rigid first-order linear part.
*Key Insight*: A remainder smaller than $\|\mathbf{h}\|$ cannot hide a second linear map. □

**Example 3.1.4.** *Derivative of a Linear Map*
If $A \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ and $\mathbf{f}(\mathbf{x}) = A\mathbf{x}$, then $D\mathbf{f}(\mathbf{x}) = A$ for every $\mathbf{x} \in \mathbb{R}^n$.
The remainder in Definition 3.1.1 is identically zero.

**Theorem 3.1.5.** *Chain Rule*
*If $\mathbf{f} : E \subseteq \mathbb{R}^n \to \mathbb{R}^m$ is differentiable at $\mathbf{x}_0$, $\mathbf{g} : G \subseteq \mathbb{R}^m \to \mathbb{R}^k$ is differentiable at $\mathbf{f}(\mathbf{x}_0)$, and $\mathbf{f}(E) \subseteq G$, then $\mathbf{g} \circ \mathbf{f}$ is differentiable at $\mathbf{x}_0$ and $D(\mathbf{g} \circ \mathbf{f})(\mathbf{x}_0) = D\mathbf{g}(\mathbf{f}(\mathbf{x}_0))\, D\mathbf{f}(\mathbf{x}_0)$.*

*Proof.* Let $A = D\mathbf{f}(\mathbf{x}_0)$, $B = D\mathbf{g}(\mathbf{f}(\mathbf{x}_0))$, $\mathbf{u}(\mathbf{h}) = \mathbf{f}(\mathbf{x}_0+\mathbf{h}) - \mathbf{f}(\mathbf{x}_0) - A\mathbf{h}$, and $\mathbf{v}(\mathbf{k}) = \mathbf{g}(\mathbf{f}(\mathbf{x}_0)+\mathbf{k}) - \mathbf{g}(\mathbf{f}(\mathbf{x}_0)) - B\mathbf{k}$.
For $\mathbf{k} = \mathbf{f}(\mathbf{x}_0+\mathbf{h}) - \mathbf{f}(\mathbf{x}_0)$, differentiability of $\mathbf{f}$ gives $\mathbf{k} \to \mathbf{0}$ and $(\mathbf{g}\circ\mathbf{f})(\mathbf{x}_0+\mathbf{h}) - (\mathbf{g}\circ\mathbf{f})(\mathbf{x}_0) - BA\mathbf{h} = B\mathbf{u}(\mathbf{h}) + \mathbf{v}(\mathbf{k})$, where $\|\mathbf{u}(\mathbf{h})\|_2 = o(\|\mathbf{h}\|_2)$ and $\|\mathbf{v}(\mathbf{k})\|_2 = o(\|\mathbf{k}\|_2) = o(\|\mathbf{h}\|_2)$. Divide by $\|\mathbf{h}\|_2$.
*Structural Commentary*: The first-order parts compose as linear maps.
*Key Insight*: The output increment of $\mathbf{f}$ is small enough to feed into the remainder estimate for $\mathbf{g}$. □

**Theorem 3.1.6.** *Componentwise Differentiability*
*Let $\mathbf{f} = (f_1, \dots, f_m) : E \subseteq \mathbb{R}^n \to \mathbb{R}^m$ and let $\pi_i : \mathbb{R}^m \to \mathbb{R}$ be the $i$th coordinate projection. Then $\mathbf{f}$ is differentiable at $\mathbf{x} \in E$ iff each $f_i$ is differentiable at $\mathbf{x}$, and $Df_i(\mathbf{x}) = \pi_i \circ D\mathbf{f}(\mathbf{x})$ when these derivatives exist.*

*Proof.*

(⇒) If $\mathbf{f}$ is differentiable, write $\mathbf{f}(\mathbf{x}+\mathbf{h}) - \mathbf{f}(\mathbf{x}) = D\mathbf{f}(\mathbf{x})\mathbf{h} + \mathbf{r}(\mathbf{h})$ with $\|\mathbf{r}(\mathbf{h})\|_2 = o(\|\mathbf{h}\|_2)$.
Applying $\pi_i$ gives $f_i(\mathbf{x}+\mathbf{h}) - f_i(\mathbf{x}) = \pi_i(D\mathbf{f}(\mathbf{x})\mathbf{h}) + \pi_i(\mathbf{r}(\mathbf{h}))$ and $|\pi_i(\mathbf{r}(\mathbf{h}))| \le \|\mathbf{r}(\mathbf{h})\|_2 = o(\|\mathbf{h}\|_2)$.
Thus $Df_i(\mathbf{x}) = \pi_i \circ D\mathbf{f}(\mathbf{x})$.

(⇐) Suppose $Df_i(\mathbf{x}) = A_i \in \mathcal{L}(\mathbb{R}^n, \mathbb{R})$ for each $i$. Define $A \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ by $A\mathbf{h} = (A_1\mathbf{h}, \dots, A_m\mathbf{h})$.
With $r_i(\mathbf{h}) = f_i(\mathbf{x}+\mathbf{h}) - f_i(\mathbf{x}) - A_i\mathbf{h}$, each $r_i(\mathbf{h}) = o(\|\mathbf{h}\|_2)$.
So $\mathbf{r}(\mathbf{h}) = (r_1(\mathbf{h}), \dots, r_m(\mathbf{h}))$ satisfies $\|\mathbf{r}(\mathbf{h})\|_2 \le \sum_i |r_i(\mathbf{h})| = o(\|\mathbf{h}\|_2)$. Definition 3.1.1 gives $D\mathbf{f}(\mathbf{x}) = A$.

*Structural Commentary*: Component derivatives assemble into one linear map.
*Key Insight*: Components test the derivative; the total derivative packages the tests together. □

**Definition 3.1.7.** *Partial, Gradient, and Directional Derivatives*
Let $E \subseteq \mathbb{R}^n$ be open, $\mathbf{f} = (f_1, \dots, f_m) : E \to \mathbb{R}^m$, $\mathbf{x} \in E$, and let $\mathbf{e}_1, \dots, \mathbf{e}_n$ be the standard basis of $\mathbb{R}^n$.

(i) For $1 \le i \le m$ and $1 \le j \le n$, the *partial derivative* of $f_i$ in coordinate $j$ at $\mathbf{x}$ is $(D_j f_i)(\mathbf{x}) := \lim_{t\to 0} \frac{f_i(\mathbf{x}+t\mathbf{e}_j) - f_i(\mathbf{x})}{t}$, when this limit exists.

(ii) For $\mathbf{u} \in \mathbb{R}^n$, the *directional derivative* of $\mathbf{f}$ at $\mathbf{x}$ in direction $\mathbf{u}$ is $(D_{\mathbf{u}}\mathbf{f})(\mathbf{x}) := \lim_{t\to 0} \frac{\mathbf{f}(\mathbf{x}+t\mathbf{u}) - \mathbf{f}(\mathbf{x})}{t}$, when this limit exists; if only direction matters, take $\|\mathbf{u}\|_2 = 1$.

(iii) If $m = 1$, write $f = f_1$; if all $(D_j f)(\mathbf{x})$ exist, the *gradient* of $f$ at $\mathbf{x}$ is $(\nabla f)(\mathbf{x}) := \sum_{j=1}^{n} (D_j f)(\mathbf{x})\mathbf{e}_j$.

**Perspective 3.1.8.** *Partials Are Coordinate Tests*
Partials and directional derivatives test selected lines; the total derivative controls all small increments by one linear map. Theorem 3.1.9 extracts partials, and Theorem 3.2.4 gives a continuity-based converse.

**Theorem 3.1.9.** *Matrix of the Total Derivative*
*If $\mathbf{f} : E \to \mathbb{R}^m$ is differentiable at $\mathbf{x}$, then every $(D_j f_i)(\mathbf{x})$ exists and, in standard bases,*

$$[D\mathbf{f}(\mathbf{x})] = \begin{bmatrix} (D_1 f_1)(\mathbf{x}) & \cdots & (D_n f_1)(\mathbf{x}) \\ \vdots & & \vdots \\ (D_1 f_m)(\mathbf{x}) & \cdots & (D_n f_m)(\mathbf{x}) \end{bmatrix}$$

*If $m = 1$ and $f$ is differentiable at $\mathbf{x}$, then $(D_{\mathbf{u}} f)(\mathbf{x}) = (\nabla f)(\mathbf{x}) \cdot \mathbf{u}$ for every $\mathbf{u} \in \mathbb{R}^n$.*

*Proof.* Set $\mathbf{h} = t\mathbf{e}_j$ in Definition 3.1.1; limit is $D\mathbf{f}(\mathbf{x})\mathbf{e}_j$, so components form column $j$ (Proposition 3.2.5).
For $m = 1$, set $\mathbf{h} = t\mathbf{u}$, identify linear functional with dot product against $\nabla f(\mathbf{x})$ (Theorem 1.6.1).
*Structural Commentary*: Partials are coordinate columns of $D\mathbf{f}(\mathbf{x})$.
*Key Insight*: Differentiability controls all directions at once; partial derivatives record its coordinate columns. □

**Remark 3.1.10.** *Row and Column Conventions*
With the coordinate-column convention from Proposition 3.2.5, the Jacobian matrix above is the standard-basis matrix of the linear map $D\mathbf{f}(\mathbf{x})$, and its $j$th column is $D\mathbf{f}(\mathbf{x})\mathbf{e}_j$.

**Definition 3.1.11.** *Jacobian*
Let $E \subseteq \mathbb{R}^n$, let $\mathbf{f} = (y_1, \ldots, y_n) : E \to \mathbb{R}^n$, and suppose $\mathbf{f}$ is differentiable at $\mathbf{x} \in E$.
The *Jacobian determinant* is $J_{\mathbf{f}}(\mathbf{x}) := \det D\mathbf{f}(\mathbf{x}) = \frac{\partial(y_1,\ldots,y_n)}{\partial(x_1,\ldots,x_n)}(\mathbf{x})$.
Then $D\mathbf{f}(\mathbf{x})$ is invertible iff $J_{\mathbf{f}}(\mathbf{x}) \neq 0$. Interpretation: $|J_{\mathbf{f}}(\mathbf{x})|$ is the infinitesimal volume-scaling factor.

## 3.2 Differentiability and Linear Approximation

**Theorem 3.2.1.** *Mean-Value Inequality on Convex Sets*
*Let $E \subseteq \mathbb{R}^n$ be open and convex, and let $\mathbf{f} : E \to \mathbb{R}^m$ be differentiable.*

*(i) If $M \geq 0$ and $\|D\mathbf{f}(\mathbf{x})\|_{op} \leq M$ for all $\mathbf{x} \in E$, then $\|\mathbf{f}(\mathbf{b}) - \mathbf{f}(\mathbf{a})\|_2 \leq M \|\mathbf{b} - \mathbf{a}\|_2$ for all $\mathbf{a}, \mathbf{b} \in E$.*
*(ii) If $D\mathbf{f}(\mathbf{x}) = 0$ for all $\mathbf{x} \in E$, then $\mathbf{f}$ is constant on $E$.*

*Proof.*

(i) Fix $\mathbf{a}, \mathbf{b} \in E$, set $\gamma(t) := (1-t)\mathbf{a} + t\mathbf{b}$, and put $\mathbf{g} := \mathbf{f} \circ \gamma$. Convexity gives $\gamma([0,1]) \subseteq E$, and the chain rule gives $\mathbf{g}'(t) = D\mathbf{f}(\gamma(t))(\mathbf{b} - \mathbf{a})$, so $\|\mathbf{g}'(t)\|_2 \leq M \|\mathbf{b} - \mathbf{a}\|_2$.
Theorem 1.1.27 gives $\|\mathbf{f}(\mathbf{b}) - \mathbf{f}(\mathbf{a})\|_2 = \|\mathbf{g}(1) - \mathbf{g}(0)\|_2 \leq M \|\mathbf{b} - \mathbf{a}\|_2$.
(ii) If $D\mathbf{f}(\mathbf{x}) = 0$ on $E$, apply (i) with $M = 0$: for all $\mathbf{a}, \mathbf{b} \in E$, $\|\mathbf{f}(\mathbf{b}) - \mathbf{f}(\mathbf{a})\|_2 = 0$, hence $\mathbf{f}(\mathbf{a}) = \mathbf{f}(\mathbf{b})$.

*Structural Commentary*: Line restriction gives a one-variable estimate.
*Key Insight*: Derivative bounds control secants. □

**Definition 3.2.2.** *$C^1$ Mapping*
Let $E \subseteq \mathbb{R}^n$ be open. A differentiable map $\mathbf{f} : E \to \mathbb{R}^m$ is *continuously differentiable*, or $C^1$, if $\mathbf{x} \mapsto D\mathbf{f}(\mathbf{x})$ is continuous from $E$ into $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ in operator norm.

**Perspective 3.2.3.** *$C^1$ Uses the Operator-Norm Metric*
$C^1$ continuity measures $\mathbf{x} \mapsto D\mathbf{f}(\mathbf{x})$ in operator norm. In coordinates this is spectral-norm continuity, equivalent here to continuity of the derivative matrix entries.




Figure 166: Coordinate increments telescope $\mathbf{x}$ to $\mathbf{x} + \mathbf{h}$.

**Theorem 3.2.4.** *$C^1$ Means Continuous Partials*
*Let $E \subseteq \mathbb{R}^n$ be open and let $\mathbf{f} = (f_1, \ldots, f_m) : E \to \mathbb{R}^m$.*
*Then $\mathbf{f}$ is $C^1$ on $E$ iff each $D_j f_i$ exists and is continuous on $E$, for $1 \leq i \leq m$ and $1 \leq j \leq n$.*

*Proof.*

($\Rightarrow$) If $\mathbf{f}$ is $C^1$, then $D\mathbf{f} : E \to \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ is continuous. By Theorem 3.1.9, $(D_j f_i)(\mathbf{x})$ is the $(i,j)$ matrix entry of $D\mathbf{f}(\mathbf{x})$; matrix-entry projections are continuous, so each $D_j f_i$ is continuous on $E$.

($\Leftarrow$) By Theorem 3.1.6, it suffices to treat $m = 1$. Fix $\mathbf{x} \in E$ and small $\mathbf{h}$ with coordinate polygon in $E$.

Set $\mathbf{x}^{(0)} := \mathbf{x}$ and $\mathbf{x}^{(j)} := \mathbf{x} + \sum_{k=1}^{j} h_k \mathbf{e}_k$. Telescoping and the one-variable MVT give

$f(\mathbf{x}+\mathbf{h}) - f(\mathbf{x}) = \sum_{j=1}^{n} h_j (D_j f)(\xi_j)$ for some $\xi_j \in [\mathbf{x}^{(j-1)}, \mathbf{x}^{(j)}]$. Let $A_{\mathbf{x}}\mathbf{h} := \sum_{j=1}^{n} h_j (D_j f)(\mathbf{x})$.

Since $\xi_j \to \mathbf{x}$ as $\mathbf{h} \to \mathbf{0}$, $|f(\mathbf{x}+\mathbf{h}) - f(\mathbf{x}) - A_{\mathbf{x}}\mathbf{h}| \le \sum_{j=1}^{n} |h_j|\,|(D_j f)(\xi_j) - (D_j f)(\mathbf{x})| = o(\|\mathbf{h}\|_2)$.

Thus $Df(\mathbf{x}) = A_{\mathbf{x}}$, i.e. $Df(\mathbf{x}) = [(D_1 f)(\mathbf{x}) \ \cdots \ (D_n f)(\mathbf{x})]$, and continuity of the entries gives $\mathbf{f} \in C^1$.

*Structural Commentary*: Continuous partials form the derivative matrix.
*Key Insight*: Coordinate telescoping gives the linear approximation. □

Higher derivatives repeat the same idea: once $D\mathbf{f}$ is itself a map into the finite-dimensional vector space $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$, we differentiate it again. The linear-algebra viewpoint matters here because the second derivative is naturally bilinear, not merely an array of second partial derivatives.

**Definition 3.2.5.** *Higher Derivatives and $C^r$*
Let $E \subseteq \mathbb{R}^n$ be open, $\mathbf{f} : E \to \mathbb{R}^m$, $r \ge 1$, and let $\mathcal{L}^r(\mathbb{R}^n, \mathbb{R}^m)$ denote the space of $r$-linear maps $(\mathbb{R}^n)^r \to \mathbb{R}^m$.

(i) The *$r$th derivative* is recursive: $D^1\mathbf{f} := D\mathbf{f}$.
For $r \ge 2$, if $D^{r-1}\mathbf{f}$ is differentiable at $\mathbf{x}$, then $D^r\mathbf{f}(\mathbf{x})(\mathbf{v}_1, \dots, \mathbf{v}_r) := \big(D(D^{r-1}\mathbf{f})(\mathbf{x})\mathbf{v}_1\big)(\mathbf{v}_2, \dots, \mathbf{v}_r)$.
(ii) For real-valued $f : E \to \mathbb{R}$, the mixed second partial is $D_{ij} f := D_i(D_j f)$ where defined.
(iii) $\mathbf{f}$ is $C^r$ on $E$ iff $D^r\mathbf{f} : E \to \mathcal{L}^r(\mathbb{R}^n, \mathbb{R}^m)$ exists and is continuous; $\mathbf{f}$ is $C^\infty$ iff $\mathbf{f}$ is $C^r$ for every $r \ge 1$.

**Theorem 3.2.6.** *Second Derivatives and Second Partials*
*Let $\mathbf{f} = (f_1, \dots, f_m) : E \subseteq \mathbb{R}^n \to \mathbb{R}^m$ and $p \in E$.*

($\Rightarrow$) *If $D^2\mathbf{f}(p)$ exists, then each $D_{ij} f_\ell(p)$ exists and $D^2 f_\ell(p)(\mathbf{e}_i, \mathbf{e}_j) = D_{ij} f_\ell(p)$.*
($\Leftarrow$) *If all $D_{ij} f_\ell$ exist near $p$ and are continuous at $p$, then $D^2\mathbf{f}(p)$ exists.*

*Proof.*

($\Rightarrow$) Apply Theorem 3.1.6 to $D\mathbf{f} : E \to \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$. The entries of $D\mathbf{f}$ are $D_j f_\ell$ by Theorem 3.1.9; differentiating the $(\ell, j)$ entry in the $\mathbf{e}_i$ direction gives $D_{ij} f_\ell(p)$ and the displayed formula.

($\Leftarrow$) Matrix entries of $D\mathbf{f}$ are functions $D_j f_\ell$. Since first partials $D_i(D_j f_\ell) = D_{ij} f_\ell$ exist near $p$ and are continuous at $p$, coordinate criterion at $p$ gives differentiability of $D\mathbf{f}$ there. Hence $D^2\mathbf{f}(p)$ exists.

*Structural Commentary*: Second partials are matrix entries of the derivative of the first-derivative matrix.
*Key Insight*: Continuous second partials make the first-derivative matrix differentiable. □




Figure 167: The signed vertex sum on a small parallelogram is symmetric in $\mathbf{v}$ and $\mathbf{w}$.

**Theorem 3.2.7.** *Symmetry of the Second Derivative*
*If $D^2\mathbf{f}(p)$ exists, then $D^2\mathbf{f}(p)(\mathbf{v}, \mathbf{w}) = D^2\mathbf{f}(p)(\mathbf{w}, \mathbf{v})$ for all $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$.*

*Proof.* Reduce to scalar $f$ by Theorem 3.1.6.
For small $t$, set $\Delta(t, \mathbf{v}, \mathbf{w}) := f(p + t\mathbf{v} + t\mathbf{w}) - f(p + t\mathbf{v}) - f(p + t\mathbf{w}) + f(p)$.
Then $\Delta(t, \mathbf{v}, \mathbf{w}) = \Delta(t, \mathbf{w}, \mathbf{v})$. The one-variable mean value theorem and differentiability of $Df$ at $p$ give $D^2 f(p)(\mathbf{v}, \mathbf{w}) = \lim_{t \to 0} \frac{\Delta(t, \mathbf{v}, \mathbf{w})}{t^2}$. The same formula with $\mathbf{v}, \mathbf{w}$ reversed proves symmetry.
*Structural Commentary*: A pointwise second derivative is already forced to be a symmetric bilinear form.
*Key Insight*: The parallelogram difference is symmetric in the two directions. □

**Corollary 3.2.8.** *Mixed Partials from $D^2$*
*If $D^2\mathbf{f}(p)$ exists, then $D_{ij}f_\ell(p) = D_{ji}f_\ell(p)$ for all components and all $i, j$.*

*Proof.* By Theorem 3.2.6, $D_{ij}f_\ell(p) = D^2f_\ell(p)(\mathbf{e}_i, \mathbf{e}_j)$; and $D_{ji}f_\ell(p) = D^2f_\ell(p)(\mathbf{e}_j, \mathbf{e}_i)$.
Theorem 3.2.7 makes the two right-hand sides equal.
*Structural Commentary*: Mixed partials are coordinate values of one bilinear map.
*Key Insight*: Symmetry of $D^2f_\ell(p)$ gives equality of mixed entries. □

**Corollary 3.2.9.** *Symmetry of Higher Derivatives*
*Let $E \subseteq \mathbb{R}^n$ be open, $\mathbf{f} : E \to \mathbb{R}^m$, $p \in E$, and $r \geq 2$.*
*If $D^r\mathbf{f}(p)$ exists, then for every $\sigma \in S_r$ and $\mathbf{v}_1, \ldots, \mathbf{v}_r \in \mathbb{R}^n$, $D^r\mathbf{f}(p)(\mathbf{v}_{\sigma(1)}, \ldots, \mathbf{v}_{\sigma(r)}) = D^r\mathbf{f}(p)(\mathbf{v}_1, \ldots, \mathbf{v}_r)$.*

*Proof.* Induct on $r$; $r = 2$ is Theorem 3.2.7. For $r > 2$, the induction hypothesis gives symmetry in slots $2, \ldots, r$ of $D^r\mathbf{f}(p)$. Applying Theorem 3.2.7 to $D^{r-2}\mathbf{f} : E \to \mathcal{L}^{r-2}(\mathbb{R}^n, \mathbb{R}^m)$ and evaluating at $\mathbf{v}_3, \ldots, \mathbf{v}_r$ gives symmetry in slots $1, 2$. Hence every adjacent transposition fixes $D^r\mathbf{f}(p)$, so every $\sigma \in S_r$ fixes it.
*Structural Commentary*: Symmetry propagates from $D^2$.
*Key Insight*: Adjacent swaps generate $S_r$. □

The next rectangle argument is the classical coordinate route to equality of mixed partials. It is weaker in hypothesis than global $C^2$, but it proves the same compatibility of coordinate second derivatives.

**Theorem 3.2.10.** *Rectangle Mean Value Formula*
*Let $E \subseteq \mathbb{R}^2$ be open, let $f : E \to \mathbb{R}$, and assume $D_1f$ and $D_{21}f = D_2(D_1f)$ exist on $E$. Let $h, k \neq 0$, let $Q \subseteq E$ be the closed coordinate rectangle with opposite vertices $(a, b)$ and $(a + h, b + k)$, and set $\Delta(f, Q) := f(a + h, b + k) - f(a + h, b) - f(a, b + k) + f(a, b)$.*
*Then there exists $(x, y) \in \operatorname{int} Q$ such that $\Delta(f, Q) = hk(D_{21}f)(x, y)$.*

*Proof.* Let $I, J$ be the open intervals with endpoints $a, a + h$ and $b, b + k$. Set $u(t) := f(t, b + k) - f(t, b)$.
By Corollary 1.1.13, $\Delta(f, Q) = u(a+h) - u(a) = h[(D_1f)(x, b+k) - (D_1f)(x, b)]$ for some $x \in I$. Applying the same result to $s \mapsto (D_1f)(x, s)$ gives $(D_1f)(x, b + k) - (D_1f)(x, b) = k(D_{21}f)(x, y)$ for some $y \in J$.
Thus $(x, y) \in I \times J = \operatorname{int} Q$ and $\Delta(f, Q) = hk(D_{21}f)(x, y)$.
*Structural Commentary*: Two MVT steps factor the rectangle increment.
*Key Insight*: $D_{21}f$ tracks vertical change of $D_1f$. □

**Theorem 3.2.11.** *Equality of Mixed Partials*
*Let $(a, b)$ lie in an open set $E \subseteq \mathbb{R}^2$. If $D_1f$, $D_2f$, and $D_{21}f$ exist on $E$, and $D_{21}f$ is continuous at $(a, b)$, then $D_{12}f(a, b)$ exists and $(D_{12}f)(a, b) = (D_{21}f)(a, b)$. Hence $D_{12}f = D_{21}f$ when $f \in C^2(E)$.*

*Proof.* Let $A = (D_{21}f)(a, b)$, $\varepsilon > 0$.
Continuity of $D_{21}f$, Theorem 3.2.10 gives $\forall$ small $h, k \neq 0$, $\left|\frac{\Delta(f,Q)}{hk} - A\right| < \varepsilon$.
Fix such an $h$ and let $k \to 0$. Since $D_2f$ exists at $(a + h, b)$ and $(a, b)$, $\frac{\Delta(f,Q)}{hk} \to \frac{(D_2f)(a+h,b)-(D_2f)(a,b)}{h}$.
Thus this last quotient differs from $A$ by at most $\varepsilon$ $\forall$ sufficiently small $h \neq 0$.
Since $\varepsilon$ was arbitrary, the quotient tends to $A$ as $h \to 0$, so $D_{12}f(a, b)$ exists and equals $A$.
*Structural Commentary*: Continuity of $D_{21}f$ controls local rectangle quotients.
*Key Insight*: The missing mixed partial is recovered by taking the two one-variable limits in the reverse order. □

**Perspective 3.2.12.** *Mixed Partials Need Hypotheses*
$D_{12}f$ and $D_{21}f$ are different iterated limits. Theorems 3.2.7 and 3.2.11 give hypotheses for equality.

# 4 Inverse, Implicit, and Rank Theorems

## 4.1 Inverse Function Theorem

**Theorem 4.1.1.** *Finite-Dimensional Linear Maps Are Lipschitz*
*Let $A \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$.*

*(i) For all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, $\|A\mathbf{x} - A\mathbf{y}\|_2 \leq \|A\|_{op} \|\mathbf{x} - \mathbf{y}\|_2$; hence $A$ is $\|A\|_{op}$-Lipschitz.*

*(ii) If $m = n$ and $A$ is invertible, then $\frac{1}{\|A^{-1}\|_{op}} \|\mathbf{x} - \mathbf{y}\|_2 \leq \|A\mathbf{x} - A\mathbf{y}\|_2 \leq \|A\|_{op} \|\mathbf{x} - \mathbf{y}\|_2$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$; hence $A : \mathbb{R}^n \to \mathbb{R}^n$ is a homeomorphism.*

*Proof.*

(i) Definition 5.1.4 gives $\|A\mathbf{z}\|_2 \leq \|A\|_{\text{op}} \|\mathbf{z}\|_2$ for all $\mathbf{z} \in \mathbb{R}^n$; set $\mathbf{z} = \mathbf{x} - \mathbf{y}$.

(ii) Apply (i) to $A^{-1}$: $\|\mathbf{x}-\mathbf{y}\|_2 = \|A^{-1}(A\mathbf{x}-A\mathbf{y})\|_2 \le \|A^{-1}\|_{\text{op}} \|A\mathbf{x}-A\mathbf{y}\|_2$.
This gives the lower bound; the upper bound is (i). Hence $A$ and $A^{-1}$ are continuous.

*Structural Commentary*: $A^{-1}$ converts upper control into lower control.
*Key Insight*: Invertible linear maps are bi-Lipschitz. □

**Theorem 4.1.2.** *Invertible Operators Form an Open Set*
*Let $\Omega$ be the set of invertible operators in $\mathcal{L}(\mathbb{R}^n)$.*
*(i) If $A \in \Omega$, $B \in \mathcal{L}(\mathbb{R}^n)$, and $\|B-A\|_{op}\|A^{-1}\|_{op} < 1$, then $B \in \Omega$.*
*(ii) $\Omega$ is open in $\mathcal{L}(\mathbb{R}^n)$.*
*(iii) The inversion map $\iota : \Omega \to \Omega$, $\iota(A) = A^{-1}$, is continuous.*

*Proof.*
(i) Put $\alpha := 1/\|A^{-1}\|_{\text{op}}$ and $\beta := \|B-A\|_{\text{op}} < \alpha$.
For every $\mathbf{x} \in \mathbb{R}^n$, $\alpha\|\mathbf{x}\|_2 \le \|A\mathbf{x}\|_2 \le \beta\|\mathbf{x}\|_2 + \|B\mathbf{x}\|_2$, hence $(\alpha-\beta)\|\mathbf{x}\|_2 \le \|B\mathbf{x}\|_2$.
Thus $\ker B = \{\mathbf{0}\}$, so $B \in \Omega$ by Theorem 1.7.8; setting $\mathbf{x} = B^{-1}\mathbf{y}$ gives $\|B^{-1}\|_{\text{op}} \le (\alpha-\beta)^{-1}$.
(ii) For $A \in \Omega$, the ball $\{B : \|B-A\|_{\text{op}} < 1/\|A^{-1}\|_{\text{op}}\}$ lies in $\Omega$ by (i); hence $\Omega$ is open.
(iii) If $B \to A \in \Omega$, then eventually $\beta < \alpha/2$, so $B \in \Omega$ and $\|B^{-1}\|_{\text{op}} \le (\alpha-\beta)^{-1} \le 2/\alpha$.
Since $B^{-1}-A^{-1} = B^{-1}(A-B)A^{-1}$, $\|B^{-1}-A^{-1}\|_{\text{op}} \le (2/\alpha)\|B-A\|_{\text{op}}\|A^{-1}\|_{\text{op}} \to 0$.

*Structural Commentary*: Perturbations below the lower stretch stay invertible.
*Key Insight*: $B^{-1}-A^{-1}$ factors through $A-B$. □

**Perspective 4.1.3.** *Linear Inverse Estimate*
For invertible $A \in \mathcal{L}(\mathbb{R}^n)$, Theorem 4.1.1 gives $c\|\mathbf{h}\|_2 \le \|A\mathbf{h}\|_2 \le \|A\|_{\text{op}}\|\mathbf{h}\|_2$; and $c := \frac{1}{\|A^{-1}\|_{\text{op}}}$.
The inverse function theorem preserves this lower stretch under small derivative changes.




Figure 168: Invertibility keeps $A(S^{n-1})$ a positive distance from 0.

**Theorem 4.1.4.** *Inverse Function Theorem*
*Let $E \subseteq \mathbb{R}^n$ be open, let $\mathbf{f} : E \to \mathbb{R}^n$ be $C^1$, let $\mathbf{a} \in E$, put $\mathbf{b} := \mathbf{f}(\mathbf{a})$, and assume $A := D\mathbf{f}(\mathbf{a})$ is invertible. Then there exist open neighborhoods $U$ of $\mathbf{a}$ and $V$ of $\mathbf{b}$ such that:*
*(i) $\mathbf{f}|_U : U \to V$ is bijective; write $\mathbf{g} := (\mathbf{f}|_U)^{-1} : V \to U$.*
*(ii) $\mathbf{g}$ is differentiable on $V$ and $D\mathbf{g}(\mathbf{y}) = \big(D\mathbf{f}(\mathbf{g}(\mathbf{y}))\big)^{-1}$ for all $\mathbf{y} \in V$.*
*(iii) $\mathbf{g} \in C^1(V, \mathbb{R}^n)$; equivalently, $\mathbf{f}|_U$ is a $C^1$ diffeomorphism.*

*Proof.* Choose $0 < \theta < 1$ and $r > 0$ so that $\overline{B}_r(\mathbf{a}) \subset E$ and $\|I - A^{-1}D\mathbf{f}(\mathbf{x})\|_{\text{op}} \le \theta$ on $\overline{B}_r(\mathbf{a})$.
Let $V := \{\mathbf{y} : \|A^{-1}(\mathbf{y}-\mathbf{b})\|_2 < (1-\theta)r\}$.
(i) For $\mathbf{y} \in V$, set $\varphi_{\mathbf{y}}(\mathbf{x}) := \mathbf{x} + A^{-1}(\mathbf{y}-\mathbf{f}(\mathbf{x}))$. Theorem 3.2.1 gives $\|\varphi_{\mathbf{y}}(\mathbf{x}) - \varphi_{\mathbf{y}}(\mathbf{z})\|_2 \le \theta\|\mathbf{x}-\mathbf{z}\|_2$ on $\overline{B}_r(\mathbf{a})$, and $\|\varphi_{\mathbf{y}}(\mathbf{x}) - \mathbf{a}\|_2 \le \theta r + \|A^{-1}(\mathbf{y}-\mathbf{b})\|_2 < r$. Since $\overline{B}_r(\mathbf{a})$ is complete, Theorem 4.5.3 gives a unique fixed point $\mathbf{x} \in \overline{B}_r(\mathbf{a})$, equivalently $\mathbf{f}(\mathbf{x}) = \mathbf{y}$. With $U := B_r(\mathbf{a}) \cap \mathbf{f}^{-1}(V)$, $\mathbf{f}|_U : U \to V$ is bijective.
(ii) Let $\mathbf{y} = \mathbf{f}(\mathbf{x})$ and $\mathbf{y}+\mathbf{k} = \mathbf{f}(\mathbf{x}+\mathbf{h})$ lie in $V$. The derivative bound gives $\|\mathbf{h} - A^{-1}\mathbf{k}\|_2 \le \theta\|\mathbf{h}\|_2$, hence $\|\mathbf{h}\|_2 \le C\|\mathbf{k}\|_2$ locally. Since $\|I - A^{-1}D\mathbf{f}(\mathbf{x})\|_{\text{op}} < 1$, $D\mathbf{f}(\mathbf{x})$ is invertible, and
$\mathbf{g}(\mathbf{y}+\mathbf{k}) - \mathbf{g}(\mathbf{y}) - \big(D\mathbf{f}(\mathbf{x})\big)^{-1}\mathbf{k} = -\big(D\mathbf{f}(\mathbf{x})\big)^{-1}[\mathbf{f}(\mathbf{x}+\mathbf{h}) - \mathbf{f}(\mathbf{x}) - D\mathbf{f}(\mathbf{x})\mathbf{h}]$.
Since $\mathbf{f}$ is differentiable at $\mathbf{x}$, division by $\|\mathbf{k}\|_2$ gives $D\mathbf{g}(\mathbf{y}) = \big(D\mathbf{f}(\mathbf{x})\big)^{-1}$.
(iii) The estimate in (ii) makes $\mathbf{g}$ continuous. Since $D\mathbf{g}(\mathbf{y}) = \big(D\mathbf{f}(\mathbf{g}(\mathbf{y}))\big)^{-1}$, continuity of $D\mathbf{g}$ follows from continuity of $D\mathbf{f}$, $\mathbf{g}$, and Theorem 4.1.2.

*Structural Commentary*: $A^{-1}$ turns solving $\mathbf{f}(\mathbf{x}) = \mathbf{y}$ into a contraction.
*Key Insight*: Small derivative variation preserves invertibility locally. □

**Corollary 4.1.5.** *Local Diffeomorphisms Are Open*
*If $\mathbf{f} : E \subseteq \mathbb{R}^n \to \mathbb{R}^n$ is $C^1$ and $D\mathbf{f}$ is everywhere invertible, then $\mathbf{f}$ is an open map (locally one-to-one).*

*Proof.* Let $W \subseteq E$ be open and let $\mathbf{y} \in \mathbf{f}(W)$. Choose $\mathbf{x} \in W$ with $\mathbf{f}(\mathbf{x}) = \mathbf{y}$. Applying Theorem 4.1.4 to $\mathbf{f}|_W$ at $\mathbf{x}$ gives open neighborhoods $U \subseteq W$ of $\mathbf{x}$ and $V$ of $\mathbf{y}$ with $\mathbf{f}(U) = V$. Hence $V \subseteq \mathbf{f}(W)$, so $\mathbf{y}$ is interior to $\mathbf{f}(W)$. Since $\mathbf{y}$ was arbitrary, $\mathbf{f}(W)$ is open. Local one-to-one-ness is the same theorem applied pointwise.
*Structural Commentary*: Openness is checked one image point at a time.
*Key Insight*: A local inverse turns nearby domain points into a full image neighborhood. □

## 4.2 Implicit Function Theorem

**Definition 4.2.1.** *Block Splitting of a Linear Map*
For $A \in \mathcal{L}(\mathbb{R}^{n+m}, \mathbb{R}^n)$ and $(\mathbf{h}, \mathbf{k}) \in \mathbb{R}^n \times \mathbb{R}^m$, define $A_x\mathbf{h} := A(\mathbf{h}, \mathbf{0})$ and $A_y\mathbf{k} := A(\mathbf{0}, \mathbf{k})$.
Then $A_x \in \mathcal{L}(\mathbb{R}^n)$, $A_y \in \mathcal{L}(\mathbb{R}^m, \mathbb{R}^n)$, and $A(\mathbf{h}, \mathbf{k}) = A_x\mathbf{h} + A_y\mathbf{k}$.

**Theorem 4.2.2.** *Linear Implicit Function Theorem*
*If $A \in \mathcal{L}(\mathbb{R}^{n+m}, \mathbb{R}^n)$, $A_x$ invertible, then $\forall \mathbf{k} \in \mathbb{R}^m \ \exists$ unique $\mathbf{h} \in \mathbb{R}^n$ s.t. $A(\mathbf{h}, \mathbf{k}) = 0$, that $\mathbf{h} = -(A_x)^{-1}A_y\mathbf{k}$.*

*Proof.* Fix $\mathbf{k} \in \mathbb{R}^m$ and set $\mathbf{h}_0 := -(A_x)^{-1}A_y\mathbf{k}$.
By Definition 4.2.1, $A(\mathbf{h}_0, \mathbf{k}) = A_x\mathbf{h}_0 + A_y\mathbf{k} = -A_y\mathbf{k} + A_y\mathbf{k} = 0$, so a solution exists.
If $\mathbf{h} \in \mathbb{R}^n$ also satisfies $A(\mathbf{h}, \mathbf{k}) = 0$, then $A_x\mathbf{h} + A_y\mathbf{k} = 0$, hence $A_x\mathbf{h} = -A_y\mathbf{k}$.
Applying $(A_x)^{-1}$ gives $\mathbf{h} = -(A_x)^{-1}A_y\mathbf{k} = \mathbf{h}_0$. Thus the solution exists and is unique.
*Structural Commentary*: Splitting $A$ separates the unknown variable from the parameter.
*Key Insight*: Invertibility of $A_x$ turns constraint $A(\mathbf{h}, \mathbf{k}) = 0$ into a uniquely solvable linear equation for $\mathbf{h}$. □

The implicit theorem below is the nonlinear counterpart of the block equation theorem. The invertible block is the set of variables one can solve for, just as in Theorem 4.2.2.




Figure 169: Near a regular point, the zero locus is the graph of an implicit function.

**Theorem 4.2.3.** *Implicit Function Theorem*
*Let $E \subseteq \mathbb{R}^{n+m}$ be open, $\mathbf{f} \in C^1(E, \mathbb{R}^n)$, $(\mathbf{a}, \mathbf{b}) \in E$, and $\mathbf{f}(\mathbf{a}, \mathbf{b}) = \mathbf{0}$.*
*Put $A := D\mathbf{f}(\mathbf{a}, \mathbf{b})$ and write $A(\mathbf{h}, \mathbf{k}) = A_x\mathbf{h} + A_y\mathbf{k}$; assume $A_x$ is invertible.*
*Then there exist open neighborhoods $U$ of $\mathbf{a}$ and $V$ of $\mathbf{b}$, and a unique map $\mathbf{g} : V \to U$, such that:*

*(i)* $\{(\mathbf{x}, \mathbf{y}) \in U \times V : \mathbf{f}(\mathbf{x}, \mathbf{y}) = \mathbf{0}\} = \{(\mathbf{g}(\mathbf{y}), \mathbf{y}) : \mathbf{y} \in V\}$ *and* $\mathbf{g}(\mathbf{b}) = \mathbf{a}$.
*(ii)* $\mathbf{g} \in C^1(V, \mathbb{R}^n)$ *and* $D\mathbf{g}(\mathbf{y}) = -\big(D_1\mathbf{f}(\mathbf{g}(\mathbf{y}), \mathbf{y})\big)^{-1} D_2\mathbf{f}(\mathbf{g}(\mathbf{y}), \mathbf{y})$ *for* $\mathbf{y} \in V$*; in particular,* $D\mathbf{g}(\mathbf{b}) = -(A_x)^{-1}A_y$.

*Proof.* Let $\Phi(\mathbf{x}, \mathbf{y}) := (\mathbf{f}(\mathbf{x}, \mathbf{y}), \mathbf{y})$. Since $D\Phi(\mathbf{a}, \mathbf{b})(\mathbf{h}, \mathbf{k}) = (A_x\mathbf{h} + A_y\mathbf{k}, \mathbf{k})$, $\ker D\Phi(\mathbf{a}, \mathbf{b}) = \{\mathbf{0}\}$; by Theorems 1.7.8 and 4.1.4, $\Phi : W \to Z$ is a local $C^1$ diffeomorphism. Shrink product neighborhoods $U \times V \subseteq W$ so that $\{\mathbf{0}\} \times V \subseteq Z$, $\Phi^{-1}(\{\mathbf{0}\} \times V) \subseteq U \times V$, and $D_1\mathbf{f}$ is invertible on $U \times V$.

(i) Define $\mathbf{g}(\mathbf{y})$ by $\Phi^{-1}(\mathbf{0}, \mathbf{y}) = (\mathbf{g}(\mathbf{y}), \mathbf{y})$. Since $\Phi(\mathbf{x}, \mathbf{y}) = (\mathbf{0}, \mathbf{y})$ iff $\mathbf{f}(\mathbf{x}, \mathbf{y}) = \mathbf{0}$, the zero set in $U \times V$ is exactly the graph of $\mathbf{g}$, and $\mathbf{g}(\mathbf{b}) = \mathbf{a}$.
(ii) The map $\mathbf{g}$ is $C^1$ as a component of $\Phi^{-1}(\mathbf{0}, \mathbf{y})$.
Differentiating $\mathbf{f}(\mathbf{g}(\mathbf{y}), \mathbf{y}) = \mathbf{0}$ gives $D_1\mathbf{f}(\mathbf{g}(\mathbf{y}), \mathbf{y})D\mathbf{g}(\mathbf{y}) + D_2\mathbf{f}(\mathbf{g}(\mathbf{y}), \mathbf{y}) = 0$, hence the formula.

*Structural Commentary*: Add $\mathbf{y}$ as a coordinate, then invert.
*Key Insight*: Invertible $D_x\mathbf{f}$ solves locally for $\mathbf{x}$. □

## 4.3 Rank Theorem

**Perspective 4.3.1.** *Rank Theorem Linear Model*
Theorem 2.2.12 reduces a linear map to its kernel–range split.
The constant-rank theorem is the local $C^1$ analogue after coordinate changes.

**Theorem 4.3.2.** *Rank Theorem*
*Let $E \subseteq \mathbb{R}^n$ be open, let $\mathbf{F} \in C^1(E, \mathbb{R}^m)$ have constant rank $r$, and fix $\mathbf{a} \in E$. Set $A := D\mathbf{F}(\mathbf{a})$, $Y := \operatorname{range} A$, and let $P : \mathbb{R}^m \to Y$ be the orthogonal projection. Then for some $\mathbf{x}_0 \in \mathbb{R}^n$, neighborhoods $V_0 \ni \mathbf{x}_0$ and $U \ni \mathbf{a}$, and $C^1$ diffeomorphism $H : V_0 \to U$, there exists $\varphi \in C^1(A(V_0), \ker P)$ such that $\mathbf{F} \circ H = A + \varphi \circ A$ on $V_0$.*

*Proof.* If $r = 0$, Theorem 3.2.1 makes $\mathbf{F}$ locally constant; take $H = I$ and $\varphi(0) = \mathbf{F}(\mathbf{a})$.
Assume $r > 0$. Choose $S : Y \to \mathbb{R}^n$ with $AS = I_Y$ and set $G := I + SP(\mathbf{F} - A)$. Since $DG(\mathbf{a}) = I$, Theorem 4.1.4 gives a local $C^1$ diffeomorphism $G : U \to V_0$ with inverse $H$; shrink $V_0$ convex and put $\mathbf{x}_0 := G(\mathbf{a})$.
Let $\Psi := \mathbf{F} \circ H$. From $AG = P\mathbf{F}$, $P\Psi = A$ and $PD\Psi = A$.
Since $\operatorname{rank} D\Psi = \operatorname{rank} A = \operatorname{rank} PD\Psi = r$, $P$ is injective on range $D\Psi$; hence $A\mathbf{v} = 0$ implies $D\Psi(\mathbf{x})\mathbf{v} = 0$.
By Theorem 3.2.1, $\Psi$ is constant on each convex fiber $V_0 \cap (\mathbf{x} + \ker A)$.
Define $\varphi(A\mathbf{x}) := (I - P)\Psi(\mathbf{x})$. Fiber constancy makes $\varphi$ well-defined, $\varphi(A\mathbf{x}) \in \ker P$, and $\Psi = A + \varphi \circ A$. After shrinking $V_0$ so $\sigma(\mathbf{u}) := \mathbf{x}_0 + S(\mathbf{u} - A\mathbf{x}_0)$ lies in $V_0$ for $\mathbf{u} \in A(V_0)$, $\varphi = (I - P)\Psi \circ \sigma$ is $C^1$.
*Structural Commentary*: Constant rank gives projection-plus-graph normal form.
*Key Insight*: Constant rank kills variation along $A$-fibers. □

Rank-nullity (Theorem 1.7.3) says that the domain directions split into rank directions and fiber directions. The rank theorem says a constant-rank map has, locally, the same split after $C^1$ changes of coordinates.




Figure 170: After local coordinate changes, a constant-rank map looks like projection onto the rank directions.

**Theorem 4.3.3.** *One-Constraint Lagrange Multiplier*
*Let $E \subseteq \mathbb{R}^n$ be open, $f, g : E \to \mathbb{R}$ be $C^1$, and $S = \{\mathbf{x} : g(\mathbf{x}) = c\}$.*
*If $p \in S$, $\nabla g(p) \neq \mathbf{0}$, and $f|_S$ has a local extremum at $p$, then $\exists \lambda \in \mathbb{R}$ with $\nabla f(p) = \lambda \nabla g(p)$.*

*Proof.* If $Df(p) = 0$, take $\lambda = 0$. Otherwise suppose $Df(p)$ is not a multiple of $Dg(p)$.
Complete $Dg(p), Df(p)$ to a dual basis $Dg(p)$, $Df(p)$, $\ell_3, \dots, \ell_n$ by Theorem 4.1.4,
and set $\Phi(\mathbf{x}) = \big(g(\mathbf{x}) - c,\ f(\mathbf{x}) - f(p),\ \ell_3(\mathbf{x} - p), \dots, \ell_n(\mathbf{x} - p)\big)$.
Then $D\Phi(p)$ is invertible, so Theorem 4.1.4 makes $\Phi$ locally invertible. In these coordinates, $S$ is locally the slice with first coordinate 0, while the second coordinate is $f - f(p)$. Local invertibility gives nearby points of $S$ with this second coordinate of both signs, contradicting extremality.
Hence $Df(p) = \lambda Dg(p)$; Riesz representation, Theorem 1.6.1, gives the gradient form.
*Structural Commentary*: The constraint kills first-order variation along $S$.
*Key Insight*: If $Df$ were independent of $Dg$, $f$ would be a free constraint coordinate. □

## 4.4 Further Topics in Local Differential Theory

**Method 4.4.1.** *Several-Variable Differentiation Proof Patterns*

(i) *Linear approximation*: write $\mathbf{f}(\mathbf{x} + \mathbf{h}) - \mathbf{f}(\mathbf{x}) = D\mathbf{f}(\mathbf{x})\mathbf{h} + \mathbf{r}(\mathbf{h})$ with $\|\mathbf{r}(\mathbf{h})\|_2 = o(\|\mathbf{h}\|_2)$.
(ii) *Norm estimates*: use $\|A\mathbf{h}\|_2 \le \|A\|_{\mathrm{op}} \|\mathbf{h}\|_2$ and the product estimate from Proposition 5.1.3.
(iii) *Line-segment reduction*: on convex sets, restrict to $\gamma(t) = (1 - t)\mathbf{a} + t\mathbf{b}$ and apply one-variable results.

(iv) *Local inversion*: if $A = D\mathbf{f}(\mathbf{a})$ is invertible, write $D\mathbf{f}(\mathbf{x}) = A + E_{\mathbf{x}}$ with $\|E_{\mathbf{x}}\|_{\text{op}}$ small and solve by contraction (Theorem 4.5.3); linear block equations use Theorem 4.2.2.
(v) *Coordinates*: read partial derivatives as the entries of the matrix of $D\mathbf{f}(\mathbf{x})$, using Theorem 3.1.9.
(vi) *Higher derivatives*: regard $D^r\mathbf{f}(p)$ as an $r$-linear map; recover coordinate partials via Theorem 3.2.6.
(vii) *Rank normal forms*: for $A = D\mathbf{F}(\mathbf{a})$, split domain by $\ker A$ and image by $\operatorname{range} A$; use the quotient-by-kernel picture from Theorem 2.2.12 and project with $P_{\operatorname{range} A}$.

*Key Insight*: Several-variable calculus is one-variable limiting plus finite-dimensional linear algebra.

# 5 Differential Forms and Stokes Theory

## 5.1 Multiple Integration and Change of Variables

This subsection moves from Euclidean-region integration and change of variables to differential forms, pullbacks, and oriented parametrized surfaces.

**Theorem 5.1.1.** *Linear Maps Scale Volume by $|\det T|$*
*Let $T \in \mathcal{L}(\mathbb{R}^n)$ and let $\Omega \subseteq \mathbb{R}^n$ be bounded Jordan measurable.*
*(i) $T(\Omega)$ is Jordan measurable.*
*(ii)* $\operatorname{vol}(T(\Omega)) = |\det T| \operatorname{vol}(\Omega)$.

*Proof.* If $T$ invertible, let $T = SP$ with $S$ orthogonal, $P = \operatorname{diag}(\lambda_1, \dots, \lambda_n)$, $\lambda_j > 0$, in orthonormal coordinates.
(i) If $T$ is invertible, $P$ and $S$ preserve Jordan measurability, so $T(\Omega)$ is Jordan measurable.
If $T$ is singular, $T(\Omega) \subseteq W$ for a proper affine subspace $W$, hence $T(\Omega)$ is Jordan null.
(ii) If $T$ is invertible, rectangle scaling gives $\operatorname{vol}(P\Omega) = (\det P)\operatorname{vol}(\Omega)$ by Jordan approximation.
Hhence $\operatorname{vol}(T\Omega) = \operatorname{vol}(P\Omega) = |\det T| \operatorname{vol}(\Omega)$. If $T$ is singular, (i) gives $\operatorname{vol}(T\Omega) = 0 = |\det T| \operatorname{vol}(\Omega)$.

*Structural Commentary*: Polar decomposition leaves only diagonal stretching.
*Key Insight*: $|\det T|$ is total coordinate stretch. □




Figure 171: A linear map scales volume by $|\det T|$.

**Definition 5.1.2.** *$k$-Cells and Multiple Integrals*
Let $a_i \le b_i$.
(i) Set $I^j := \prod_{i=1}^{j} [a_i, b_i] \subseteq \mathbb{R}^j$ for $0 \le j \le k$, with $I^0$ a point. The *$k$-cell* is $I^k$.
(ii) For $f \in C(I^k)$, set $f_k := f$ and $f_{j-1}(x_1, \dots, x_{j-1}) := \int_{a_j}^{b_j} f_j(x_1, \dots, x_j)\, dx_j$, $j = k, \dots, 1$.
The *multiple integral* is $\int_{I^k} f(\mathbf{x})\, d\mathbf{x} := f_0$, also written $\int_{I^k} f$.

For bounded functions on rectangles, the Darboux picture from Definition 1.2.1 generalizes by replacing intervals with grid rectangles.

**Definition 5.1.3.** *Rectangular Grids and Riemann Sums*
Let $R = [a, b] \times [c, d] \subseteq \mathbb{R}^2$, let $P : a = x_0 < \cdots < x_m = b$, let $Q : c = y_0 < \cdots < y_n = d$, and set $G := P \times Q$.
(i) The *grid rectangles* are $R_{ij} := [x_{i-1}, x_i] \times [y_{j-1}, y_j]$, with $|R_{ij}| := (x_i - x_{i-1})(y_j - y_{j-1})$.
(ii) For tags $S = \{(s_{ij}, t_{ij}) \in R_{ij}\}$, the *Riemann sum* is $R(f, G, S) := \sum_{i=1}^{m} \sum_{j=1}^{n} f(s_{ij}, t_{ij})\, |R_{ij}|$.
(iii) For bounded $f : R \to \mathbb{R}$, set $(m_{ij}, M_{ij}) := (\inf_{R_{ij}} f, \sup_{R_{ij}} f)$ and $(L(f, G), U(f, G)) := \sum_{i,j} (m_{ij}, M_{ij})\, |R_{ij}|$.
(iv) The *lower* and *upper integrals* are $\underline{\int_R} f := \sup_G L(f, G)$ and $\overline{\int_R} f := \inf_G U(f, G)$.
If they agree, $f$ is Riemann integrable on $R$, and $\int_R f$ denotes the common value.




Figure 172: A grid rectangle $R_{ij}$ and sample point $(s_{ij}, t_{ij})$.

**Theorem 5.1.4.** *Fubini on a Rectangle, Darboux Form*
*Let $R = [a,b] \times [c,d]$, let $f : R \to \mathbb{R}$ be Riemann integrable, and set $f_y(x) := f(x,y)$.*
*For $y \in [c,d]$, define $\underline{F}(y) := \sup_P L(P, f_y)$ and $\overline{F}(y) := \inf_P U(P, f_y)$, with $P$ ranging over partitions of $[a,b]$.*

*(i) $\underline{F}, \overline{F}$ are Riemann integrable on $[c,d]$.*
*(ii) $\int_R f = \int_c^d \underline{F}(y)\,dy = \int_c^d \overline{F}(y)\,dy$.*

*If $\int_a^b f(x,y)\,dx$ exists for every $y$, then $\underline{F}(y) = \overline{F}(y) = \int_a^b f(x,y)\,dx$.*

*Proof.*

(i) For $G = P \times Q$, horizontal slice sums give $L(f,G) \le L(Q,\underline{F}) \le U(Q,\overline{F}) \le U(f,G)$.
Since $f$ is integrable on $R$, choose $G$ with $U(f,G) - L(f,G) < \varepsilon$; hence $U(Q,\overline{F}) - L(Q,\underline{F}) < \varepsilon$.
Thus $\underline{F}$ and $\overline{F}$ are integrable and have the same integral by Theorem 2.1.7.

(ii) The same inequalities give $L(f,G) \le \int_c^d \underline{F} \le \int_c^d \overline{F} \le U(f,G)$.
Letting $U(f,G) - L(f,G) \to 0$ gives both integrals equal to $\int_R f$.

*Structural Commentary*: Product grids become slice sums.
*Key Insight*: Fubini is a Darboux squeeze. □

**Theorem 5.1.5.** *Order Independence on Cells*
*For $f \in C(I^k)$, all iterated orders defining $\int_{I^k} f$ give the same value.*

*Proof.* Let $L, L'$ be two iterated orders. For $h(\mathbf{x}) = h_1(x_1)\cdots h_k(x_k)$, $L(h) = L'(h) = \prod_{i=1}^k \int_{a_i}^{b_i} h_i(x_i)\,dx_i$, so $L(g) = L'(g)$ for every finite sum $g$ of such products. These sums are dense in $C(I^k)$ by Theorem 9.2.8.

Choose $g$ with $\|f - g\|_\infty < \varepsilon/V$, where $V = \prod_{i=1}^k (b_i - a_i)$.

Since $|L(r)|, |L'(r)| \le V \|r\|_\infty$, then $|L(f) - L'(f)| \le |L(f-g)| + |L'(f-g)| < 2\varepsilon$.

*Structural Commentary*: Product functions plus uniform approximation prove order independence.
*Key Insight*: Fubini-type order independence is first an algebraic fact, then a uniform approximation fact. □

**Perspective 5.1.6.** *Order Interchange Requires Hypotheses*
Finite sums are order-free; limiting processes are not.
Theorems 5.1.4 and 5.1.5 give hypotheses for interchanging integral order.

**Definition 5.1.7.** *Support and Simplex Integrals*
For $f : \mathbb{R}^k \to \mathbb{R}$ or $\mathbb{C}$, define $\operatorname{supp} f := \overline{\{\mathbf{x} \in \mathbb{R}^k : f(\mathbf{x}) \neq 0\}}$.

(i) For $f \in C_c(\mathbb{R}^k)$ and any cell $I^k \supseteq \operatorname{supp} f$, define $\int_{\mathbb{R}^k} f := \int_{I^k} f$; this is independent of $I^k$.

(ii) The *standard $k$-simplex* is $Q^k := \{\mathbf{x} \in \mathbb{R}^k : x_i \ge 0,\ \sum_{i=1}^k x_i \le 1\}$.
For $f \in C(Q^k)$, extend $f$ by 0 to $[0,1]^k$ and define $\int_{Q^k} f$ as the corresponding multiple integral.

**Definition 5.1.8.** *Primitive Mappings and Flips*
Let $E \subseteq \mathbb{R}^n$ be open.

(i) $G : E \to \mathbb{R}^n$ is *primitive in coordinate $m$* if $G(\mathbf{x}) = (x_1, \ldots, x_{m-1}, g(\mathbf{x}), x_{m+1}, \ldots, x_n)$.
(ii) If $g$ is differentiable at $\mathbf{a} \in E$, then $DG(\mathbf{a})$ is $I_n$ except row $m$ is $\nabla g(\mathbf{a})$, and $J_G(\mathbf{a}) = (D_m g)(\mathbf{a})$.
(iii) A *flip* is a linear map $B : \mathbb{R}^n \to \mathbb{R}^n$ with $B\mathbf{e}_p = \mathbf{e}_q$, $B\mathbf{e}_q = \mathbf{e}_p$, and $B\mathbf{e}_i = \mathbf{e}_i$ for $i \notin \{p,q\}$, $p \neq q$.

**Theorem 5.1.9.** *Local Decomposition into Primitive Mappings*
*Let $E \subseteq \mathbb{R}^n$ be open, let $\mathbf{F} \in C^1(E, \mathbb{R}^n)$, let $\mathbf{0} \in E$, and assume $\mathbf{F}(\mathbf{0}) = \mathbf{0}$ and $D\mathbf{F}(\mathbf{0})$ is invertible. Then there exist a neighborhood $U$ of $\mathbf{0}$, primitive $C^1$ maps $G_1, \dots, G_n$ near $\mathbf{0}$, and maps $B_1, \dots, B_{n-1}$ such that:*

*(i)* $G_i(\mathbf{0}) = \mathbf{0}$, $DG_i(\mathbf{0})$ *is invertible, and each* $B_i$ *is a flip or* $I$.
*(ii)* $\mathbf{F} = B_1 \cdots B_{n-1} G_n \circ \cdots \circ G_1$ *on* $U$.

*Proof.* Let $P_m \mathbf{x} := \sum_{i=1}^{m} x_i \mathbf{e}_i$ and set $\mathbf{F}_1 := \mathbf{F}$.

(i) Suppose $P_{m-1}\mathbf{F}_m = P_{m-1}$, $\mathbf{F}_m(\mathbf{0}) = \mathbf{0}$, and $D\mathbf{F}_m(\mathbf{0})$ invertible, and write $\mathbf{F}_m = P_{m-1} + \sum_{i=m}^{n} \alpha_i \mathbf{e}_i$.
Choose $k \geq m$ with $(D_m \alpha_k)(\mathbf{0}) \neq 0$, let $B_m$ swap $m, k$ (or $B_m = I$), set $G_m(\mathbf{x}) := \mathbf{x} + (\alpha_k(\mathbf{x}) - x_m)\mathbf{e}_m$.
Then $G_m$ is primitive, $G_m(\mathbf{0}) = \mathbf{0}$, $J_{G_m}(\mathbf{0}) \neq 0$, and $G_m$ is locally invertible.
With $\mathbf{F}_{m+1} := B_m \mathbf{F}_m \circ G_m^{-1}$, one has $P_m \mathbf{F}_{m+1} = P_m$, $\mathbf{F}_{m+1}(\mathbf{0}) = \mathbf{0}$, and $D\mathbf{F}_{m+1}(\mathbf{0})$ invertible.
Iterate for $m = 1, \dots, n-1$ and set $G_n := \mathbf{F}_n$.
(ii) From $\mathbf{F}_{m+1} = B_m \mathbf{F}_m \circ G_m^{-1}$ and $B_m^2 = I$, $\mathbf{F}_m = B_m \mathbf{F}_{m+1} \circ G_m$.
Iteration gives $\mathbf{F} = \mathbf{F}_1 = B_1 \cdots B_{n-1} G_n \circ G_{n-1} \circ \cdots \circ G_1$ near $\mathbf{0}$.

*Structural Commentary*: Local diffeomorphisms factor by coordinate elimination.
*Key Insight*: Nonlinear elimination follows $D\mathbf{F}(\mathbf{0})$. □

**Theorem 5.1.10.** *Partition of Unity on a Compact Set*
*Let $K \subseteq \mathbb{R}^n$ be compact and let $\{V_\alpha\}$ cover $K$ by open sets. Then there are $\psi_1, \dots, \psi_s \in C^1(\mathbb{R}^n)$ such that*

*(i)* $0 \leq \psi_i \leq 1$ *for all* $i$;
*(ii)* $\operatorname{supp}(\psi_i) \subseteq V_\alpha$ *for some* $\alpha$;
*(iii)* $\sum_{i=1}^{s} \psi_i = 1$ *on* $K$.
*If $f \in C(\mathbb{R}^n)$ and $\operatorname{supp}(f) \subseteq K$, then $f = \sum_{i=1}^{s} \psi_i f$, with each summand supported in one member of the cover.*

*Proof.*

(i) For each $\mathbf{x} \in K$, choose balls with $\overline{B(\mathbf{x})} \subseteq W(\mathbf{x}) \subseteq \overline{W(\mathbf{x})} \subseteq V_{\alpha(\mathbf{x})}$.
By compactness, Definition 4.1.1, finitely many $B(\mathbf{x}_i)$ cover $K$.
(ii) Choose $\varphi_i \in C^1(\mathbb{R}^n)$ with $\varphi_i = 1$ on $B(\mathbf{x}_i)$ and $\operatorname{supp}(\varphi_i) \subseteq W(\mathbf{x}_i)$. Set $\psi_i = (1 - \varphi_1) \cdots (1 - \varphi_{i-1})\varphi_i$.
(iii) Then $0 \leq \psi_i \leq 1$, each $\psi_i$ is supported in some $V_\alpha$, and $\sum_{i=1}^{s} \psi_i = 1 - \prod_{i=1}^{s} (1 - \varphi_i)$, which equals 1 on $K$.
Multiplying by $f$ gives the final assertion.

*Structural Commentary*: Compactness finite; product cutoffs sum to one.
*Key Insight*: A partition of unity turns local change-of-variable information into a global formula. □

The nonlinear change-of-variables theorem is the local volume-scaling formula from Theorem 5.1.1 applied cell by cell after a fine grid has been warped by a $C^1$ diffeomorphism.




Figure 173: A $C^1$ change of variables turns a fine grid into curvilinear cells; $|J_T|$ records local volume scaling.

**Theorem 5.1.11.** *Change of Variables in $\mathbb{R}^k$*
*Let $T : E \to \mathbb{R}^k$ be one-to-one and $C^1$ on open $E \subseteq \mathbb{R}^k$, with $J_T \neq 0$ on $E$.*
*If $f \in C(\mathbb{R}^k)$ and $\operatorname{supp}(f)$ is compact in $T(E)$, then $\int_{\mathbb{R}^k} f(\mathbf{y})\, d\mathbf{y} = \int_{\mathbb{R}^k} f(T(\mathbf{x}))\, |J_T(\mathbf{x})|\, d\mathbf{x}$.*

*Proof.* Let $\mathcal{C}$ be the class of maps for which formula holds for all continuous functions with compact support in the image. Primitive maps lie in $\mathcal{C}$ by one-variable substitution, and flips lie in $\mathcal{C}$ by Theorem 5.1.5.
If $P, Q \in \mathcal{C}$, then $P \circ Q \in \mathcal{C}$, since Theorems 3.1.5 and 1.1.8 give $J_{P \circ Q}(\mathbf{x}) = J_P(Q(\mathbf{x}))J_Q(\mathbf{x})$.
After translating coordinates, Theorem 5.1.9 therefore gives the formula on a neighborhood of each point of $E$.
By Theorem 4.1.4, these neighborhoods have open images covering $\operatorname{supp}(f)$.
Choose a partition of unity from Theorem 5.1.10, write $f = \sum_i \psi_i f$ with each $\psi_i f$ supported in one such image, apply the local formula to each summand, and sum.

*Structural Commentary*: Model cases plus partitions of unity prove the theorem.
*Key Insight*: The absolute Jacobian is local volume scaling, globalized by compact support. □

**Method 5.1.12.** *Change-of-Variables Jacobian Method*
For a differentiable map $\sigma : \Omega \subseteq \mathbb{R}^n \to \mathbb{R}^n$:

(i) compute the derivative matrix $D\sigma(\mathbf{x})$;
(ii) compute the local volume factor $|\det D\sigma(\mathbf{x})|$;
(iii) replace $d\mathbf{y}$ by $|\det D\sigma(\mathbf{x})|\, d\mathbf{x}$ when $\mathbf{y} = \sigma(\mathbf{x})$;
(iv) transform the domain bounds in the new coordinates.

**Example 5.1.13.** *Polar and Spherical Jacobian Factors*
For polar coordinates $\sigma(r,\theta) = (r\cos\theta, r\sin\theta)$, $D\sigma(r,\theta) = \begin{bmatrix} \cos\theta & -r\sin\theta \\ \sin\theta & r\cos\theta \end{bmatrix}$, and $|\det D\sigma(r,\theta)| = r$.
Thus area integrals acquire the factor $r\, dr\, d\theta$.
For spherical coordinates $\sigma(\rho,\varphi,\theta) = (\rho\sin\varphi\cos\theta, \rho\sin\varphi\sin\theta, \rho\cos\varphi)$, and $|\det D\sigma(\rho,\varphi,\theta)| = \rho^2\sin\varphi$.
Thus volume integrals acquire the factor $\rho^2\sin\varphi\, d\rho\, d\varphi\, d\theta$.

**Method 5.1.14.** *Multiple-Integration Proof Patterns*
For a multiple-integration identity:

(i) *Products*: prove product cases, then pass by uniform approximation;
(ii) *Grids*: compare product-grid upper and lower sums with one-variable slice sums;
(iii) *Support*: enclose compact support in a cell and check cell-independence;
(iv) *Local form*: reduce $C^1$ coordinate changes to primitive maps and flips;
(v) *Determinants*: use multiplicativity and Theorem 1.1.8;
(vi) *Globalize*: apply the partition of unity, Theorem 5.1.10.

*Key Insight*: Change of variables is local determinant geometry plus compactness.

## 5.2 Differential Forms and Manifolds

Differential forms package oriented integration on parametrized cells. They use determinant sign and laws (Definition 1.1.1, Theorems 1.1.8, 1.2.5) and change of variables (Theorem 5.1.11); conceptually, forms test oriented cells.

**Perspective 5.2.1.** *Forms as Functionals*
The notation $\int f\, dx$ treats $dx$ as directed coordinate variation. Differential forms make this precise: a 1-form such as $f\, dx + g\, dy$ assigns a number to an oriented path; $dx$ records net $x$-variation, and $f\, dx$ weights it by $f$.




Figure 174: $y\, dx$ records signed, $y$-weighted $x$-variation.

**Definition 5.2.2.** *k-Surfaces and Differential Forms*
Let $E \subseteq \mathbb{R}^n$ be open.

(i) A compact *parameter domain* $D \subseteq \mathbb{R}^k$ is a $k$-cell or $Q^k = \{\mathbf{u} \in \mathbb{R}^k : u_i \geq 0,\ \sum_i u_i \leq 1\}$.
For compact $D$, $C^r(D)$ means extendable as $C^r$ to an open neighborhood of $D$.
(ii) A *k-surface* in $E$ is a $C^1$ map $\Phi : D \to E$ from a compact parameter domain $D \subseteq \mathbb{R}^k$.
(iii) For $k \geq 1$, a *k-form* on $E$ is $\omega = \sum_I a_I(\mathbf{x})\, dx_{i_1} \wedge \cdots \wedge dx_{i_k}$, where $I = (i_1, \ldots, i_k)$ and $a_I \in C(E, \mathbb{R})$.
(iv) For an increasing $k$-index $I = (i_1 < \cdots < i_k)$, set $dx_I := dx_{i_1} \wedge \cdots \wedge dx_{i_k}$.
A *standard presentation* is $\omega = \sum_I b_I(\mathbf{x})\, dx_I$, with $I$ increasing.
(v) For a $k$-surface $\Phi : D \to E$, define $\int_\Phi \omega := \int_D \sum_I a_I(\Phi(\mathbf{u}))\, J_I\Phi(\mathbf{u})\, d\mathbf{u}$, where $J_I\Phi := \frac{\partial(x_{i_1},\ldots,x_{i_k})}{\partial(u_1,\ldots,u_k)}$.
(vi) A 0-form is a continuous function on $E$; a form is $C^r$ iff all coefficients $a_I$ are $C^r$.

**Remark 5.2.3.** *Parametrized Cells, Not Just Images*
Parametrizations may fail to be injective or regular. Forms therefore act on parametrized cells, not just images; this matters for orientation, multiplicity, and cancellation.

**Proposition 5.2.4.** *Basic Forms and Standard Presentation*

*(i) Swapping neighboring factors in $dx_{i_1} \wedge \cdots \wedge dx_{i_k}$ changes sign; repeated indices give 0.*
*(ii) Every $k$-form has a unique standard presentation.*

*Proof.*

(i) For $\Phi(\mathbf{u}) = (x_{i_1}, \ldots, x_{i_k})$, the coefficient of $dx_{i_1} \wedge \cdots \wedge dx_{i_k}$ is the determinant of the corresponding $k \times k$ Jacobian minor. Row swaps change its sign and repeated rows make it 0.

(ii) By (i), each term either vanishes or sorts to an increasing index, giving existence.
For uniqueness, let $\omega = \sum_I b_I(\mathbf{x})\, dx_I$ be zero in standard presentation.
If $b_J(\mathbf{v}) \neq 0$ for increasing $J = (j_1, \ldots, j_k)$, choose $D = [-h, h]^k$ so $b_J(\Phi(\mathbf{u}))$ has constant nonzero sign for $\Phi(\mathbf{u}) = \mathbf{v} + \sum_{r=1}^{k} u_r \mathbf{e}_{j_r}$. Then the $J$-Jacobian is 1 and all other standard Jacobians are 0, so $0 = \int_\Phi \omega = \int_D b_J(\Phi(\mathbf{u}))\, d\mathbf{u}$, a contradiction. Hence all $b_I = 0$.

*Structural Commentary*: Wedge signs are determinant signs.
*Key Insight*: Standard form sorts indices once. □

**Theorem 5.2.5.** *Naturality Under Reparametrization*
*Let $\Phi : D \to E \subseteq \mathbb{R}^n$ be a $C^1$ $k$-surface, and let $R : D' \to D$ be a $C^1$ diffeomorphism with $\epsilon := \operatorname{sgn} \det DR$ constant on $D'$. For every $k$-form $\omega$ on $E$, $\int_{\Phi \circ R} \omega = \epsilon \int_\Phi \omega$.*

*Proof.* By linearity, take $\omega = f\, dx_I$. Cauchy-Binet gives $J_I(\Phi \circ R)(\mathbf{u}) = J_I\Phi(R(\mathbf{u})) \det DR(\mathbf{u})$.
Thus $\int_{\Phi\circ R} \omega = \int_{D'} f(\Phi(R\mathbf{u})) J_I\Phi(R\mathbf{u}) \det DR(\mathbf{u})\, d\mathbf{u} = \epsilon \int_D f(\Phi(\mathbf{v})) J_I\Phi(\mathbf{v})\, d\mathbf{v}$ by change of variables.
*Structural Commentary*: Reparametrization changes coordinates only.
*Key Insight*: Orientation contributes $\operatorname{sgn} \det DR$. □

**Definition 5.2.6.** *Product and Exterior Derivative*
For basic forms, $dx_I \wedge dx_J$ is obtained by concatenating $I, J$; it is 0 if an index repeats, and otherwise $\operatorname{sgn}(\sigma) dx_K$, where $K$ is the increasing rearrangement and $\sigma$ is the sorting permutation. Extend bilinearly.

(i) For $f \in C^1(E)$, define $df := \sum_{i=1}^{n} (D_i f)(\mathbf{x})\, dx_i$.

(ii) For a $C^1$ $k$-form $\omega = \sum_I b_I(\mathbf{x})\, dx_I$, define $d\omega := \sum_I (db_I) \wedge dx_I$.

**Theorem 5.2.7.** *Exterior Derivative Laws*
*If $\omega, \lambda$ are $C^1$ forms in $E$ and $\omega$ is a $k$-form, then $d(\omega \wedge \lambda) = (d\omega) \wedge \lambda + (-1)^k \omega \wedge d\lambda$. If $\omega$ is $C^2$, then $d^2\omega = 0$.*

*Proof.* By bilinearity, it suffices to take $\omega = f\, dx_I$ and $\lambda = g\, dx_J$.
The ordinary product rule gives $d(fg) = f\, dg + g\, df$; moving $dg$ past $k$ factors in $dx_I$ gives the displayed sign.
For a $C^2$ 0-form $f$, $d^2 f = \sum_{i,j} (D_{ij} f)(\mathbf{x})\, dx_i \wedge dx_j = 0$, by Theorem 3.2.11 and antisymmetry.
The general case follows from $\omega = \sum_I b_I\, dx_I$ and the product rule just proved.
*Structural Commentary*: The exterior derivative is built to make mixed partial cancellation visible.
*Key Insight*: Symmetric second derivatives meet antisymmetric wedge products, so $d^2 = 0$. □

**Definition 5.2.8.** *Change of Variables for Forms*
Let $E \subseteq \mathbb{R}^n$, $V \subseteq \mathbb{R}^m$, let $T = (t_1, \ldots, t_m) : E \to V$ be $C^1$, and let $\omega = \sum_I b_I(\mathbf{y})\, dy_I$ be a $k$-form on $V$.

(i) Set $dt_i := \sum_{j=1}^{n} (D_j t_i)(\mathbf{x})\, dx_j$ for $1 \leq i \leq m$.

(ii) The *pullback* of $\omega$ by $T$ is $T^*\omega := \omega_T := \sum_I b_I(T(\mathbf{x}))\, dt_{i_1} \wedge \cdots \wedge dt_{i_k}$.

Pullback is the differential-form analogue of the dual map in Definition 4.2.3: cells move forward by composition with $T$, while forms move backward so they can be evaluated before or after that forward motion.




Figure 175: Cells push forward with $T$; forms pull back with $T^*$, so $T^*\alpha(\varphi) = \alpha(T_*\varphi)$.

**Theorem 5.2.9.** *Pullback Commutes with the Basic Operations*
*Let $T = (t_1, \dots, t_m) : E \subseteq \mathbb{R}^n \to V \subseteq \mathbb{R}^m$ be $C^1$, and let $\omega_T := T^*\omega$.*

*(i)* $(\omega + \lambda)_T = \omega_T + \lambda_T$ *for forms of the same order;*
*(ii)* $(\omega \wedge \lambda)_T = \omega_T \wedge \lambda_T$*;*
*(iii)* $d(\omega_T) = (d\omega)_T$ *when $\omega$ is $C^1$ and $T$ is $C^2$;*
*(iv) if $S : V \to W$ is $C^1$, then $(\omega_S)_T = \omega_{S\circ T}$.*

*Moreover, if $\Phi$ is a $k$-surface in $E$ and $\omega$ is a $k$-form in $V$, then $\int_{T\circ\Phi} \omega = \int_\Phi \omega_T$.*

*Proof.*

(i) Addition follows termwise from $b_I \mapsto b_I \circ T$ and $dy_i \mapsto dt_i$.
(ii) Pullback replaces each $dy_i$ by $dt_i$; bilinearity gives $(\omega \wedge \lambda)_T = \omega_T \wedge \lambda_T$.
(iii) Let $\omega = \sum_I b_I\, dy_I$. Since $d(b_I \circ T) = (db_I)_T$ (Theorem 3.1.5) and $d(dt_i) = d^2 t_i = 0$, $d(\omega_T) = (d\omega)_T$.
(iv) If $S = (s_1, \dots, s_p)$, both $(\omega_S)_T$ and $\omega_{S\circ T}$ replace $dz_q$ by $d(s_q \circ T)$, by the chain rule.

For the integral identity, expand over $D$ and use Cauchy-Binet on $D(T \circ \Phi) = DT(\Phi)D\Phi$.
*Structural Commentary*: Pullback translates forms before integration.
*Key Insight*: Forms are defined so that composition of parameterizations becomes an algebraic substitution. □

## 5.3 Stokes Theory

Stokes theory relates exterior differentiation to oriented boundary, with Green's theorem, classical Stokes, and divergence as special cases.

**Definition 5.3.1.** *Chains and Boundaries*
Let $\mathbf{p}_0, \dots, \mathbf{p}_k \in \mathbb{R}^n$.

(i) The oriented affine *$k$-simplex* $[\mathbf{p}_0, \dots, \mathbf{p}_k]$ is the $k$-surface $\sigma : Q^k \to \mathbb{R}^n$, $\sigma(\mathbf{u}) := \mathbf{p}_0 + \sum_{i=1}^{k} u_i(\mathbf{p}_i - \mathbf{p}_0)$.
(ii) For a permutation $\pi$, $[\mathbf{p}_{\pi(0)}, \dots, \mathbf{p}_{\pi(k)}] := \operatorname{sgn}(\pi)[\mathbf{p}_0, \dots, \mathbf{p}_k]$.
(iii) An *affine $k$-chain* is $C = \sum_\ell c_\ell \sigma_\ell$; integration is linear: $\int_C \omega := \sum_\ell c_\ell \int_{\sigma_\ell} \omega$.
(iv) For $k \geq 1$, $\partial_{\mathrm{ch}}[\mathbf{p}_0, \dots, \mathbf{p}_k] := \sum_{j=0}^{k} (-1)^j [\mathbf{p}_0, \dots, \mathbf{p}_{j-1}, \mathbf{p}_{j+1}, \dots, \mathbf{p}_k]$, extended linearly.
If $\Phi = T \circ \sigma$ is a $C^2$ oriented simplex, set $\partial_{\mathrm{ch}}\Phi := T(\partial_{\mathrm{ch}}\sigma)$, extended linearly to chains.

**Theorem 5.3.2.** *Stokes' Theorem*
*Let $V \subseteq \mathbb{R}^m$ be open, let $\Psi$ be a $C^2$ $k$-chain in $V$, and let $\omega$ be a $C^1$ $(k-1)$-form on $V$. Then $\int_\Psi d\omega = \int_{\partial_{\mathrm{ch}}\Psi} \omega$.*

*Proof.* By linearity, take $\Psi = \Phi = T \circ \sigma$, where $\sigma$ is the standard oriented $k$-simplex.
By Theorem 5.2.9, $\int_\Phi d\omega = \int_\sigma d(\omega_T)$ and $\int_{\partial_{\mathrm{ch}}\Phi} \omega = \int_{\partial_{\mathrm{ch}}\sigma} \omega_T$; it remains to prove the formula on $\sigma$.
By linearity, reduce to $\lambda = f\, dx_1 \wedge \cdots \wedge \widehat{dx_r} \wedge \cdots \wedge dx_k$.
For $\partial_{\mathrm{ch}}\sigma = \sum_{j=0}^{k} (-1)^j \sigma_j$, all faces vanish except the two with varying $x_r$ endpoints.
The remaining equality is the one-variable FTC in the $x_r$ integral.
*Structural Commentary*: Pullback reduces Stokes to the standard simplex.
*Key Insight*: Boundary signs match the missing wedge factor. □

**Definition 5.3.3.** *Closed and Exact Forms*
Let $E \subseteq \mathbb{R}^n$ be open and let $\omega$ be a $k$-form on $E$.

(i) $\omega$ is *exact* if $\omega = d\lambda$ for some $(k-1)$-form $\lambda$ on $E$.
(ii) $\omega$ is *closed* if $\omega \in C^1$ and $d\omega = 0$.

**Perspective 5.3.4.** *Closed Is Local, Exact Is Global*
Exact $\Rightarrow$ closed is just $d^2 = 0$; the converse depends on the domain.
On $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ the standard warning example is $\omega = \frac{-y}{x^2+y^2}\,dx + \frac{x}{x^2+y^2}\,dy$, which is closed but not exact.
Holes can obstruct global antiderivatives even when local derivative tests vanish.

**Theorem 5.3.5.** *Poincare Lemma on Convex Sets*
*Let $k \geq 1$.*

*(i) If $E \subseteq \mathbb{R}^n$ is open and convex, then every closed $C^1$ $k$-form on $E$ is exact.*
*(ii) The same conclusion holds on any open set $C^2$-diffeomorphic to a convex open set.*

*Proof.*

(i) Fix $\mathbf{a} \in E$ and set $H_t(\mathbf{x}) := \mathbf{a} + t(\mathbf{x} - \mathbf{a})$, $0 \leq t \leq 1$.
For a $k$-form $\omega$, define $(K\omega)_{\mathbf{x}}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1}) := \int_0^1 t^{k-1} \omega_{H_t(\mathbf{x})}(\mathbf{x} - \mathbf{a}, \mathbf{v}_1, \ldots, \mathbf{v}_{k-1})\,dt$.
Differentiating under the integral gives $\omega - H_0^*\omega = dK\omega + Kd\omega$.
Since $H_0^*\omega = 0$ for $k \geq 1$, $d\omega = 0$ implies $\omega = d(K\omega)$.
(ii) Let $\Phi : C \to E$ be a $C^2$ diffeomorphism from convex open $C$. If $d\omega = 0$, then $d(\Phi^*\omega) = \Phi^*(d\omega) = 0$.
By (i), $\Phi^*\omega = d\eta$ on $C$, hence $\omega = (\Phi^{-1})^* d\eta = d((\Phi^{-1})^*\eta)$ on $E$.

*Structural Commentary*: Convex contraction gives a homotopy operator.
*Key Insight*: Closed forms become derivatives along the contraction. □

**Theorem 5.3.6.** *Vector Analysis Dictionary in $\mathbb{R}^3$*
*Let $E \subseteq \mathbb{R}^3$ be open and let $\mathbf{F} = F_1\mathbf{e}_1 + F_2\mathbf{e}_2 + F_3\mathbf{e}_3 \in C^1(E, \mathbb{R}^3)$.*
*Set $\lambda_{\mathbf{F}} := F_1\,dx + F_2\,dy + F_3\,dz$ and $\omega_{\mathbf{F}} := F_1\,dy \wedge dz + F_2\,dz \wedge dx + F_3\,dx \wedge dy$.*

*(i) For $u \in C^1(E)$, $\mathbf{F} = \nabla u$ iff $\lambda_{\mathbf{F}} = du$.*
*(ii) $d\lambda_{\mathbf{F}} = \omega_{\nabla \times \mathbf{F}}$; hence $\nabla \times \mathbf{F} = 0$ iff $d\lambda_{\mathbf{F}} = 0$.*
*(iii) For $\mathbf{G} \in C^1(E, \mathbb{R}^3)$, $\mathbf{F} = \nabla \times \mathbf{G}$ iff $\omega_{\mathbf{F}} = d\lambda_{\mathbf{G}}$.*
*(iv) $d\omega_{\mathbf{F}} = (\nabla \cdot \mathbf{F})\,dx \wedge dy \wedge dz$; hence $\nabla \cdot \mathbf{F} = 0$ iff $d\omega_{\mathbf{F}} = 0$.*
*(v) If $E$ is $C^2$-diffeomorphic to a convex open set, then $\nabla \times \mathbf{F} = 0 \Rightarrow \exists u \in C^1(E) : \mathbf{F} = \nabla u$;*
*$\nabla \cdot \mathbf{F} = 0 \Rightarrow \exists \mathbf{G} \in C^1(E, \mathbb{R}^3) : \mathbf{F} = \nabla \times \mathbf{G}$.*

*Proof.*

(i) This is $du = (D_1u)\,dx + (D_2u)\,dy + (D_3u)\,dz$.
(ii) Direct expansion gives $d\lambda_{\mathbf{F}} = (D_2F_3 - D_3F_2)\,dy \wedge dz + (D_3F_1 - D_1F_3)\,dz \wedge dx + (D_1F_2 - D_2F_1)\,dx \wedge dy$.
(iii) By (ii), $d\lambda_{\mathbf{G}} = \omega_{\nabla \times \mathbf{G}}$; since $\mathbf{H} \mapsto \omega_{\mathbf{H}}$ is injective, $\omega_{\mathbf{F}} = d\lambda_{\mathbf{G}}$ iff $\mathbf{F} = \nabla \times \mathbf{G}$.
(iv) Direct expansion gives $d\omega_{\mathbf{F}} = (D_1F_1 + D_2F_2 + D_3F_3)\,dx \wedge dy \wedge dz$.
(v) If $\nabla \times \mathbf{F} = 0$, then $\lambda_{\mathbf{F}}$ is closed by (ii), so Poincare gives $\lambda_{\mathbf{F}} = du$, hence $\mathbf{F} = \nabla u$ by (i). If $\nabla \cdot \mathbf{F} = 0$, then $\omega_{\mathbf{F}}$ is closed by (iv), so Poincare gives $\omega_{\mathbf{F}} = d\eta$; writing $\eta = \lambda_{\mathbf{G}}$ gives $\mathbf{F} = \nabla \times \mathbf{G}$ by (iii).

*Structural Commentary*: Exterior derivative is grad–curl–div.
*Key Insight*: Vector identities are $d^2 = 0$ in $\mathbb{R}^3$. □

**Theorem 5.3.7.** *Green, Classical Stokes, and Divergence*
*With the indicated positive orientations:*

*(i) For $\Omega \subseteq \mathbb{R}^2$ with positively oriented $\operatorname{bd}\Omega$ and $\alpha, \beta \in C^1$ near $\Omega$, $\int_{\operatorname{bd}\Omega}(\alpha\,dx + \beta\,dy) = \int_\Omega (D_1\beta - D_2\alpha)\,dA$.*
*(ii) If $\Phi$ is a $C^2$ oriented 2-surface in $\mathbb{R}^3$ and $\mathbf{F} \in C^1$ near $\Phi$, then $\int_{\partial_{\mathrm{ch}}\Phi} \mathbf{F} \cdot \mathbf{t}\,ds = \int_\Phi (\nabla \times \mathbf{F}) \cdot \mathbf{n}\,dA$.*
*(iii) If $\Omega \subseteq \mathbb{R}^3$ has positively oriented boundary and $\mathbf{F} \in C^1$ near $\Omega$, then $\int_{\operatorname{bd}\Omega} \mathbf{F} \cdot \mathbf{n}\,dA = \int_\Omega (\nabla \cdot \mathbf{F})\,dV$.*

*Proof.* Use the form Stokes identity $\int_\Psi d\eta = \int_{\partial_{\mathrm{ch}}\Psi} \eta$.

(i) For $\lambda := \alpha\,dx + \beta\,dy$, $d\lambda = (D_1\beta - D_2\alpha)\,dx \wedge dy$; hence $\int_{\operatorname{bd}\Omega} \lambda = \int_\Omega d\lambda$ gives Green's formula.
(ii) For $\lambda := \lambda_{\mathbf{F}}$, $d\lambda = \omega_{\nabla \times \mathbf{F}}$; hence $\int_{\partial_{\mathrm{ch}}\Phi} \lambda = \int_\Phi d\lambda$ gives the classical Stokes formula.
(iii) For $\omega := \omega_{\mathbf{F}}$, $d\omega = (\nabla \cdot \mathbf{F})\,dx \wedge dy \wedge dz$; hence $\int_{\operatorname{bd}\Omega} \omega = \int_\Omega d\omega$ gives the divergence formula.

*Structural Commentary*: One form identity yields the three vector formulas.
*Key Insight*: Boundary plus $d$ is invariant; notation changes. □

## 5.4 Further Topics in Differential Forms and Stokes Theory

**Method 5.4.1.** *Differential-Forms Proof Patterns*

(i) *Signs*: compute wedge signs by determinant swaps and permutation parity.
(ii) *Standard presentation*: remove repeated indices and sort the rest.
(iii) *Pullback*: substitute $dt_i$ for $dy_i$ and apply the chain rule.

(iv) *Stokes*: pull back to the standard simplex, prove the coordinate case, extend linearly.
(v) *Closed versus exact*: use $d^2 = 0$ for necessity; use convexity or $C^2$-equivalence for primitives.
(vi) *Vector analysis*: translate fields to forms, apply Stokes, translate back.

*Key Insight*: Forms package signs, orientation, and parameter changes.

# 6 Infinite Series and Summability

## 6.1 Series Tests

Series are sequences of partial sums. Comparison, condensation, root and ratio tests, rearrangements, products, and double sums all refine that viewpoint.

**Definition 6.1.1.** *Series and Partial Sums*
Let $(a_n)_{n\geq 1} \subseteq \mathbb{C}$. Summation indices are bound variables and may be renamed within unchanged scope.

(i) For integers $1 \leq p \leq q$, $\sum_{n=p}^{q} a_n := a_p + a_{p+1} + \cdots + a_q$.

(ii) The *series* $\sum_{n=1}^{\infty} a_n$ is the *partial-sum sequence* $(s_N)$, where $s_N := \sum_{n=1}^{N} a_n$.

(iii) The series *converges to* $s$, written $\sum_{n=1}^{\infty} a_n = s$, iff $s_N \to s$; otherwise it *diverges*. Also for $n = 0$.

*Key Insight*: A series is a sequence first; its sum is a limit when that limit exists.

**Definition 6.1.2.** *Absolute and Nonabsolute Convergence*
The series $\sum a_n$ *converges absolutely* if $\sum |a_n|$ converges.
If $\sum a_n$ converges but $\sum |a_n|$ diverges, then $\sum a_n$ converges *nonabsolutely*.

**Motivation 6.1.3.** *Partial Sums Remember Order*
An infinite sum is a limit of partial sums. For alternating harmonic series, odd and even partial sums approach the same value from opposite sides; see Corollary 6.1.25 and Theorem 6.1.34.

**Perspective 6.1.4.** *Formal Algebra Needs Convergence*
Finite sums are algebraic; infinite sums are partial-sum limits. Algebraic manipulations are justified only by convergence control, as in Theorem 6.1.5, Corollary 6.1.6, and Theorems 6.1.31, 6.1.34.
*Key Insight*: The symbol $\sum_{n=1}^{\infty}$ is a limit claim before it is an algebraic object.

**Theorem 6.1.5.** *Cauchy Criterion for Series*
*The series $\sum a_n$ converges if and only if $\forall \varepsilon > 0$ $\exists$ an integer $N$ s.t. $m \geq n \geq N$ implies $\left|\sum_{k=n}^{m} a_k\right| \leq \varepsilon$.*

*Proof.*
($\Rightarrow$) If $s_n := \sum_{k=1}^{n} a_k$ converges, then $(s_n)$ is Cauchy. Thus, for $m \geq n$ large enough, $\left|\sum_{k=n}^{m} a_k\right| = |s_m - s_{n-1}| \leq \varepsilon$.
($\Leftarrow$) The tail condition gives, for $m > n$ large enough, $|s_m - s_n| = \left|\sum_{k=n+1}^{m} a_k\right| \leq \varepsilon$.
Hence $(s_n)$ is Cauchy in $\mathbb{C} \cong \mathbb{R}^2$ and converges by Corollary 4.2.8.
*Structural Commentary*: A series converges exactly when all sufficiently late finite tails are small.
*Key Insight*: The Cauchy test measures tails, not the eventual value of the sum. □

**Corollary 6.1.6.** *Terms of a Convergent Series Tend to Zero*
*If $\sum a_n$ converges, then $a_n \to 0$.*

*Proof.* Let $\varepsilon > 0$. By Theorem 6.1.5, there is $N$ such that $m \geq n \geq N$ implies $|\sum_{k=n}^{m} a_k| \leq \varepsilon$.
Taking $m = n$ gives $|a_n| \leq \varepsilon$ for all $n \geq N$, hence $a_n \to 0$.
*Structural Commentary*: A single term is the shortest possible tail.
*Key Insight*: The condition $a_n \to 0$ is necessary, but not sufficient for convergence. □

**Theorem 6.1.7.** *Nonnegative Series and Bounded Partial Sums*
*A series of nonnegative real terms converges if and only if its partial sums form a bounded sequence.*

*Proof.*
($\Rightarrow$) Every convergent sequence is bounded, so the partial sums are bounded.

($\Leftarrow$) Since $a_n \geq 0$, $s_n = \sum_{k=1}^{n} a_k$ increases; if bounded, it converges by Theorem 2.1.11.

*Structural Commentary*: Nonnegative series replace cancellation by monotonicity.
*Key Insight*: For nonnegative terms, convergence is exactly bounded accumulation. □

**Theorem 6.1.8.** *Comparison Test*

*(i) If $|a_n| \leq c_n$ for $n \geq N_0$, and if $\sum c_n$ converges, then $\sum a_n$ converges.*
*(ii) If $a_n \geq d_n \geq 0$ for $n \geq N_0$, and if $\sum d_n$ diverges, then $\sum a_n$ diverges.*

*Proof.*

(i) Given $\varepsilon > 0$, choose $N \geq N_0$ so that $m \geq n \geq N$ implies $\sum_{k=n}^{m} c_k \leq \varepsilon$.
Then $\left|\sum_{k=n}^{m} a_k\right| \leq \sum_{k=n}^{m} |a_k| \leq \sum_{k=n}^{m} c_k \leq \varepsilon$, so Theorem 6.1.5 applies.

(ii) If $\sum a_n$ converged, then (i) applied to $d_n \leq a_n$ would force $\sum d_n$ to converge, a contradiction.

*Structural Commentary*: Absolute tail domination transfers Cauchy convergence.
*Key Insight*: Comparison is a tail test; finitely many early terms do not matter. □

**Theorem 6.1.9.** *Geometric Series*
*If $0 \leq x < 1$, then $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$. If $x \geq 1$, the series diverges.*

*Proof.* For $x \neq 1$, $\sum_{k=0}^{n} x^k = \frac{1-x^{n+1}}{1-x}$. Let $n \to \infty$ and use Theorem 2.2.10.
If $x = 1$, the partial sums are $n+1$, so they diverge.
*Structural Commentary*: The finite geometric identity converts convergence into the limit of $x^{n+1}$.
*Key Insight*: The geometric series is the model comparison series for ratio-like decay. □




Figure 176: At $x = 2/3$, the strip widths sum to $1 + \frac{2}{3} + \frac{4}{9} + \cdots = 3$.

**Definition 6.1.10.** *Real Decimals as Series*
For digits $b_N, \ldots, b_0, a_1, a_2, \ldots \in \{0, \ldots, 9\}$, with $b_N \neq 0$ unless the integer part is 0,
$b_N \cdots b_0.a_1a_2\cdots := \sum_{i=0}^{N} b_i 10^i + \sum_{j=1}^{\infty} a_j 10^{-j}$. The fractional series converges absolutely since $0 \leq a_j 10^{-j} \leq 9 \cdot 10^{-j}$ and $\sum 9 \cdot 10^{-j}$ is geometric. A minus sign denotes the negative of this value.

**Proposition 6.1.11.** *Terminal Nines Are Nonunique*
*The same real number can have two decimal representations:* $1.000\ldots = 0.999\ldots$.

*Proof.* $1.000\ldots = 1 + \sum_{j=1}^{\infty} 0 \cdot 10^{-j} = 1$, while $0.999\ldots = \sum_{j=1}^{\infty} 9 \cdot 10^{-j} = 9 \cdot \frac{10^{-1}}{1-10^{-1}} = 1$.
*Structural Commentary*: The two digit strings have the same partial-sum limit.
*Key Insight*: Uniqueness requires excluding terminal 9 tails. □

**Perspective 6.1.12.** *Decimal Expansions Are Series*
Proposition 2.2.11 gives decimal truncations that converge to a real number, while Definition 6.1.10 interprets the infinite digit string as a series. Proposition 6.1.11 explains the standard terminal-9 ambiguity.

**Theorem 6.1.13.** *Cauchy Condensation Test*
*Let $(a_n)_{n\geq 1}$ satisfy $a_1 \geq a_2 \geq \cdots \geq 0$. Then $\sum_{n=1}^{\infty} a_n$ converges iff $\sum_{k=0}^{\infty} 2^k a_{2^k}$ converges.*

*Proof.* Let $s_n := \sum_{j=1}^{n} a_j$ and $t_k := \sum_{j=0}^{k} 2^j a_{2^j}$. Both partial-sum sequences are nondecreasing.

($\Rightarrow$) If $(s_n)$ is bounded, then $t_m \leq 2 \sum_{j=1}^{2^m} a_j = 2s_{2^m}$, so $(t_m)$ is bounded and the condensed series converges.

($\Leftarrow$) If $(t_k)$ is bounded, then for $2^m \le n < 2^{m+1}$, $s_n \le \sum_{j=0}^{m} 2^j a_{2^j} = t_m$; hence $(s_n)$ is bounded, QED.

*Structural Commentary*: Dyadic blocks compare to endpoint terms.
*Key Insight*: Monotone tails are detected on dyadic scales. □

**Corollary 6.1.14.** *p-Series Test*
*For $p \in \mathbb{R}$:*

*(i) If $p > 1$, then $\sum_{n=1}^{\infty} n^{-p}$ converges.*

*(ii) If $p \le 1$, then $\sum_{n=1}^{\infty} n^{-p}$ diverges.*

*Proof.*

(i) If $p > 1$, condensation reduces to $\sum_{k=0}^{\infty} 2^k (2^k)^{-p} = \sum_{k=0}^{\infty} 2^{(1-p)k}$, a geometric series with ratio $2^{1-p} < 1$.

(ii) If $p \le 0$, then $n^{-p} \not\to 0$; if $0 < p \le 1$, condensation gives a geometric series with ratio $2^{1-p} \ge 1$.

*Structural Commentary*: Condensation turns $n$-powers into a geometric series.
*Key Insight*: The threshold is the ratio $2^{1-p} = 1$. □

**Theorem 6.1.15.** *Integral Test*
*Let $N \in \mathbb{N}$ and let $f : [N, \infty) \to [0, \infty)$ be decreasing and Riemann integrable on compact subintervals.*

*(i) For $M \ge N$, $\int_N^{M+1} f(x)\,dx \le \sum_{k=N}^{M} f(k) \le f(N) + \int_N^M f(x)\,dx$.*

*(ii) $\sum_{k=N}^{\infty} f(k)$ converges iff $\int_N^{\infty} f(x)\,dx$ converges.*

*Proof.*

(i) For $x \in [k, k+1]$, $f(x) \le f(k)$, so $\int_N^{M+1} f \le \sum_{k=N}^{M} f(k)$.

For $k \ge N+1$ and $x \in [k-1, k]$, $f(k) \le f(x)$, so $\sum_{k=N}^{M} f(k) \le f(N) + \int_N^M f$.

(ii) The truncations $\sum_{k=N}^{M} f(k)$ and $\int_N^M f$ are nondecreasing in $M$. By (i), either both are bounded or both are unbounded; hence the series and improper integral converge or diverge together.

*Structural Commentary*: Unit intervals compare samples to area.
*Key Insight*: Same monotone tail, same boundedness. □




Figure 177: Left and right endpoint estimates for decreasing $f \ge 0$.

**Definition 6.1.16.** *The Number e*
Define $e := \sum_{n=0}^{\infty} \frac{1}{n!}$, where $0! := 1$; convergence follows by comparison with the geometric series.

**Theorem 6.1.17.** *A Limit Formula for e*
*The limit formula is $\lim_{n\to\infty} \left(1 + \frac{1}{n}\right)^n = e$.*

*Proof.* Let $s_m := \sum_{k=0}^{m} \frac{1}{k!}$ and $t_n := (1 + \frac{1}{n})^n$. The binomial theorem gives $t_n \le s_n$, hence $\limsup t_n \le e$.
For fixed $m \le n$, first $m$ binomial terms give $t_n \ge 1 + 1 + \frac{1}{2!}\left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{m!}\left(1 - \frac{1}{n}\right)\cdots\left(1 - \frac{m-1}{n}\right)$.
Letting $n \to \infty$ with $m$ fixed gives $\liminf t_n \ge s_m$, and then $m \to \infty$ gives $e \le \liminf t_n$.
*Structural Commentary*: Bound above by the factorial series; bound below by fixed binomial terms.
*Key Insight*: Limsup and liminf squeeze the binomial limit to the series definition. □

**Proposition 6.1.18.** *Remainder Bound for the Series of e*
*If $n \geq 1$ and $s_n = \sum_{k=0}^{n} 1/k!$, then $0 < e - s_n < \frac{1}{n!\,n}$.*

*Proof.* Tail is $e - s_n = \frac{1}{(n+1)!} + \frac{1}{(n+2)!} + \cdots < \frac{1}{(n+1)!}\left(1 + \frac{1}{n+1} + \frac{1}{(n+1)^2} + \cdots\right) = \frac{1}{n!\,n}$
*Structural Commentary*: The factorial tail is bounded by a geometric tail.
*Key Insight*: Rapid convergence of $\sum \frac{1}{n!}$ gives arithmetic control of $e$. □

**Theorem 6.1.19.** *e Is Irrational*
*The number $e$ is irrational.*

*Proof.* Suppose $e = p/q$ with $p, q$ positive integers. By Proposition 6.1.18, $0 < q!(e - s_q) < \frac{1}{q}$.
But $q!e$ and $q!s_q$ are integers, so $q!(e - s_q)$ is an integer strictly between 0 and 1, impossible.
*Structural Commentary*: The sharp tail bound turns rationality into an impossible integer.
*Key Insight*: A positive integer cannot be smaller than 1. □

**Theorem 6.1.20.** *Root Test*
*For a series $\sum a_n$, put $\alpha := \limsup_{n\to\infty} \sqrt[n]{|a_n|}$. Then:*

*(i) if $\alpha < 1$, then $\sum a_n$ converges absolutely;*
*(ii) if $\alpha > 1$, then $\sum a_n$ diverges;*
*(iii) if $\alpha = 1$, the test gives no information.*

*Proof.*

(i) Choose $\alpha < \beta < 1$. Eventually $|a_n| < \beta^n$, so $\sum |a_n|$ converges by comparison with $\sum \beta^n$.
(ii) Since $\alpha > 1$, $|a_n| > 1$ for infinitely many $n$, so $a_n \not\to 0$; use Corollary 6.1.6.
(iii) $\sum 1/n$ and $\sum 1/n^2$ both have $\alpha = 1$, but behave differently by Corollary 6.1.14.

*Structural Commentary*: The root test compares term size directly to geometric decay.
*Key Insight*: Values below 1 force absolute convergence; values above 1 prevent terms from tending to 0. □

**Theorem 6.1.21.** *Ratio Test*
*Assume $a_n \neq 0$ for all sufficiently large $n$.*

*(i) If $\limsup_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| < 1$, then $\sum a_n$ converges absolutely.*
*(ii) If $\liminf_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| > 1$, then $\sum a_n$ diverges.*
*(iii) In the remaining cases, the test gives no information.*

*Proof.*

(i) Choose $\beta < 1$ with $|a_{n+1}/a_n| < \beta$ eventually.
Then $|a_n| = O(\beta^n)$, so $\sum |a_n|$ converges by comparison with a geometric series.
(ii) Choose $\beta > 1$ with $|a_{n+1}/a_n| > \beta$ eventually.
Then the tail $|a_n|$ is eventually increasing and bounded away from 0, so $a_n \not\to 0$.
(iii) The series $\sum 1/n$ and $\sum 1/n^2$ both have ratio limit 1, but behave differently by Corollary 6.1.14.

*Structural Commentary*: Ratios turn successive terms into a geometric majorant.
*Key Insight*: Ratios below 1 force geometric decay; ratios above 1 prevent terms from tending to zero. □

**Theorem 6.1.22.** *Root and Ratio Test Comparison*
*For any sequence $(c_n)$ of positive numbers, $\liminf_{n\to\infty} \frac{c_{n+1}}{c_n} \leq \liminf_{n\to\infty} \sqrt[n]{c_n} \leq \limsup_{n\to\infty} \sqrt[n]{c_n} \leq \limsup_{n\to\infty} \frac{c_{n+1}}{c_n}$.*

*Proof.* Let $\alpha = \limsup c_{n+1}/c_n$. If $\alpha < +\infty$ and $\beta > \alpha$, then eventually $c_{n+1} \leq \beta c_n$, hence $c_n \leq c_N \beta^{n-N}$ for all large $n$. Taking $n$th roots gives $\limsup \sqrt[n]{c_n} \leq \beta$; let $\beta \downarrow \alpha$. If $\alpha = +\infty$, the rightmost inequality is automatic.
Let $\ell = \liminf c_{n+1}/c_n$. If $\ell = 0$, the leftmost inequality is automatic. If $\gamma < \ell$ with $\gamma > 0$, then eventually $c_{n+1} \geq \gamma c_n$, hence $c_n \geq c_N \gamma^{n-N}$ for all large $n$. Taking $n$th roots gives $\liminf \sqrt[n]{c_n} \geq \gamma$; let $\gamma \uparrow \ell$.
The middle inequality is the general fact that $\liminf a_n \leq \limsup a_n$.
*Structural Commentary*: A tail ratio bound multiplies into a root bound.
*Key Insight*: Whenever the ratio test proves convergence, the root test does too. □

**Theorem 6.1.23.** *Summation by Parts*
*Given two sequences $(a_n)$ and $(b_n)$, put $A_n := \sum_{k=0}^{n} a_k$ and $A_{-1} := 0$.*

*If $0 \leq p \leq q$, then $\sum_{n=p}^{q} a_n b_n = \sum_{n=p}^{q-1} A_n(b_n - b_{n+1}) + A_q b_q - A_{p-1} b_p$.*

*Proof.* Since $a_n = A_n - A_{n-1}$, $\sum_{n=p}^{q} a_n b_n = \sum_{n=p}^{q} A_n b_n - \sum_{n=p}^{q} A_{n-1} b_n = \sum_{n=p}^{q} A_n b_n - \sum_{n=p-1}^{q-1} A_n b_{n+1}$.
*Structural Commentary*: This is the discrete analogue of integration by parts.
*Key Insight*: Move the difference from $a_n$ onto $b_n$ by rewriting $a_n = A_n - A_{n-1}$. □

**Theorem 6.1.24.** *Dirichlet Test*
*If the partial sums $A_n$ of $\sum a_n$ are bounded, $b_n \downarrow 0$, and $b_n \geq 0$, then $\sum a_n b_n$ converges.*

*Proof.* Choose $M$ with $|A_n| \leq M$. If $M = 0$, the result is immediate. Otherwise choose $N$ with $b_N \leq \varepsilon/(2M)$. For $N \leq p \leq q$, $\left|\sum_{n=p}^{q} a_n b_n\right| \leq M\left(\sum_{n=p}^{q-1}(b_n - b_{n+1}) + b_q + b_p\right) = 2Mb_p \leq 2Mb_N \leq \varepsilon$. Theorem 6.1.5 applies.
*Structural Commentary*: Bounded oscillation in $A_n$ is damped by a monotone scalar tending to 0.
*Key Insight*: Summation by parts converts cancellation into a total variation estimate for $b_n$. □

**Corollary 6.1.25.** *Alternating Series Test*
*Suppose $|c_1| \geq |c_2| \geq |c_3| \geq \cdots$, the signs alternate, and $c_n \to 0$. Then $\sum c_n$ converges.*

*Proof.* After shifting indices to start at 0 if desired, apply Theorem 6.1.24 with $a_n = (-1)^{n+1}$ and $b_n = |c_n|$.
*Structural Commentary*: Alternation gives bounded sums; decreasing magnitudes damp them.
*Key Insight*: Alternating convergence is Dirichlet convergence in its simplest form. □




Figure 178: The intervals $I_j = [a_{2j}, a_{2j-1}]$ are disjoint inside $(0, a_1)$.

**Perspective 6.1.26.** *Alternating Partial Sums as Nested Control*
The interval picture explains the boundedness in Corollary 6.1.25: the even partial sums collect disjoint lengths inside a fixed interval, while the odd partial sums differ from the even ones by the next term, which tends to 0.




Figure 179: Odd partial sums (blue) decrease and even partial sums (orange) increase toward $\log 2$.

**Theorem 6.1.27.** *Absolute Convergence Implies Convergence*
*If $\sum a_n$ converges absolutely, then $\sum a_n$ converges.*

*Proof.* For $m \geq n$, $\left|\sum_{k=n}^{m} a_k\right| \leq \sum_{k=n}^{m} |a_k|$. Apply Theorem 6.1.5 to $\sum |a_n|$.
*Structural Commentary*: Absolute convergence removes cancellation from the convergence question.
*Key Insight*: Small absolute tails force small signed or complex tails. □

**Theorem 6.1.28.** *Addition and Scalar Multiplication of Series*
*If $\sum a_n = A$ and $\sum b_n = B$, then $\sum(a_n + b_n) = A + B$ and $\sum ca_n = cA$ for every fixed scalar $c$.*

*Proof.* Apply Theorem 2.1.4 to the corresponding partial sums.
*Structural Commentary*: Linearity of convergent series is just limit algebra for partial sums.
*Key Insight*: Add series by adding their partial-sum sequences. □

**Definition 6.1.29.** *Cauchy Product*
Given $\sum a_n$, $\sum b_n$, define $c_n := \sum_{k=0}^{n} a_k b_{n-k}$, $(n = 0, 1, 2, \ldots)$. Series $\sum c_n$ is *Cauchy product* of given series.




Figure 180: Diagonal sums of $(a_ib_j)$ form the Cauchy product; finite rectangular sums are order-free.

**Perspective 6.1.30.** *Diagonal Ordering in a Product*
$c_n$ is the diagonal sum of the products $a_ib_j$ with $i + j = n$. Absolute convergence, as in Theorems 6.1.31 and 6.1.34, keeps the value independent of the accumulation order.

**Theorem 6.1.31.** *Mertens' Theorem for Cauchy Products*
*Suppose* $\sum a_n$ *converges absolutely,* $\sum a_n = A$, $\sum b_n = B$, *and* $c_n = \sum_{k=0}^{n} a_k b_{n-k}$. *Then* $\sum_{n=0}^{\infty} c_n = AB$.

*Proof.* Let $A_n, B_n, C_n$ be the partial sums and set $\beta_n := B_n - B$.
Then $C_n = A_nB + \gamma_n$, and $\gamma_n := a_0\beta_n + a_1\beta_{n-1} + \cdots + a_n\beta_0$. Since $A_nB \to AB$, it suffices to show $\gamma_n \to 0$.
Put $\alpha := \sum |a_n|$. If $|\beta_j| \le \varepsilon$ for $j \ge N$, then $|\gamma_n| \le |\beta_0 a_n + \cdots + \beta_N a_{n-N}| + \varepsilon\alpha$.
Let $n \to \infty$ with $N$ fixed. The first term tends to 0, so $\limsup |\gamma_n| \le \varepsilon\alpha$; let $\varepsilon \downarrow 0$.
*Structural Commentary*: Absolute convergence of one factor controls the convolution error.
*Key Insight*: One absolute factor makes the Cauchy product converge to $AB$. □

**Definition 6.1.32.** *Rearrangement of a Series*
If $(k_n)$ lists each positive integer exactly once and $a'_n := a_{k_n}$, then $\sum a'_n$ is a *rearrangement* of $\sum a_n$.

**Theorem 6.1.33.** *Riemann Rearrangement Theorem*
*Let* $\sum a_n$ *be a real series converging but not absolutely, and let* $\alpha, \beta \in [-\infty, \infty]$ *with* $\alpha \le \beta$.
*Then there exists a bijection* $\pi : \mathbb{N} \to \mathbb{N}$ *such that, for* $s'_N := \sum_{n=1}^{N} a_{\pi(n)}$:

*(i)* $\pi$ *rearranges every term of* $\sum a_n$ *exactly once.*
*(ii)* $\liminf_{N\to\infty} s'_N = \alpha$.
*(iii)* $\limsup_{N\to\infty} s'_N = \beta$.

*Proof.* Write $a_n = p_n - q_n$, where $p_n := \max(a_n, 0)$ and $q_n := \max(-a_n, 0)$. Then $\sum p_n = \sum q_n = +\infty$ and $p_n, q_n \to 0$. List the nonzero $p_n$ as $P_1, P_2, \ldots$ and the nonzero $q_n$ as $Q_1, Q_2, \ldots$, preserving order.
Choose finite targets $\ell_N < u_N$ with $\ell_N \to \alpha$, $u_N \to \beta$, and $\ell_N < u_{N+1}$.

(i) Alternately add unused $P$ terms until $s' > u_N$, then unused $-Q$ terms until $s' < \ell_N$.
Since $\sum P_j = \sum Q_j = +\infty$, all stops are finite and every nonzero term is used; insert zeros anywhere.
(ii) At the $N$th down-crossing, $\ell_N - Q_j \le s'_k < \ell_N$ with $Q_j \to 0$. Thus lower crossing subsequences tend to $\alpha$, and later sums stay above later $\ell_N$ up to $o(1)$; hence $\liminf s'_N = \alpha$.
(iii) At the $N$th up-crossing, $u_N < s'_k \le u_N + P_j$ with $P_j \to 0$. Thus upper crossing subsequences tend to $\beta$, and later sums stay below later $u_N$ up to $o(1)$; hence $\limsup s'_N = \beta$.

*Structural Commentary*: Divergent one-sided masses steer the partial sums.
*Key Insight*: Vanishing terms make the overshoot vanish. □

**Theorem 6.1.34.** *Absolutely Convergent Series Are Rearrangement-Invariant*
*Every rearrangement of an absolutely convergent complex series converges to the same sum.*

*Proof.* Let $s_n, s'_n$ be the original and rearranged partial sums.
Given $\varepsilon > 0$, choose $N$ so that $\sum_{i=N}^{m} |a_i| \le \varepsilon$ for $m \ge N$, then choose $p$ so $a_1, \ldots, a_N$ occur among $a'_1, \ldots, a'_p$.
For $n > p$, $s_n - s'_n$ uses only tail terms, hence $|s_n - s'_n| \le \varepsilon$. Thus $s'_n$ has the same limit as $s_n$.
*Structural Commentary*: Absolute tails control any finite reordering of the remaining terms.
*Key Insight*: Absolute convergence is exactly the protection needed against rearrangement. □

**Definition 6.1.35.** *Absolute Summation over Countable Sets*
Let $X$ be countable, $f : X \to \mathbb{C}$, and $g : \mathbb{N} \to X$ be a bijection, $g_n := g(n)$.
If $\sum_{n=1}^{\infty} |f(g_n)| < \infty$, then $\sum_{x\in X} f(x)$ is *absolutely convergent*, and $\sum_{x\in X} f(x) := \sum_{n=1}^{\infty} f(g_n)$.

**Theorem 6.1.36.** *Countable Sums Are Order-Free Under Absolute Convergence*
*Let $X, Y$ be countable, let $f : X \to \mathbb{C}$ be absolutely summable, and let $h : Y \to X$ be a bijection.*

*(i)* $f \circ h : Y \to \mathbb{C}$ *is absolutely summable.*
*(ii)* $\sum_{y\in Y} f(h(y)) = \sum_{x\in X} f(x)$.
*(iii) Countable absolutely convergent sums are independent of enumeration.*

*Proof.* Let $g : \mathbb{N} \to X$ and $k : \mathbb{N} \to Y$ be enumerations, and set $\sigma := g^{-1} \circ h \circ k : \mathbb{N} \to \mathbb{N}$.

(i) As $\sigma$ is bijection, $h(k_n) = g_{\sigma(n)}$; $\sum_{n=1}^{\infty} |f(h(k_n))| = \sum_{n=1}^{\infty} \left|f(g_{\sigma(n)})\right| < \infty$ by absolute rearrangement invariance.
(ii) The same invariance gives $\sum_{y\in Y} f(h(y)) = \sum_{n=1}^{\infty} f(g_{\sigma(n)}) = \sum_{n=1}^{\infty} f(g_n) = \sum_{x\in X} f(x)$.
(iii) Take $Y = \mathbb{N}$ and let $h$ be any enumeration of $X$.

*Structural Commentary*: Bijections only reorder an absolutely summable list.
*Key Insight*: Absolute tails make enumeration irrelevant. □

**Theorem 6.1.37.** *Fubini for Absolutely Summable Arrays*
*Let $(a_{mn})_{m,n\geq 1} \subseteq \mathbb{C}$ and $L := \sum_{(m,n)\in\mathbb{N}^2} |a_{mn}| < \infty$. Define $S := \sum_{(m,n)\in\mathbb{N}^2} a_{mn}$ when summed absolutely over $\mathbb{N}^2$.*

*(i) Each row* $\sum_{n=1}^{\infty} a_{mn}$ *and column* $\sum_{m=1}^{\infty} a_{mn}$ *converges absolutely.*
*(ii) With* $r_m := \sum_{n=1}^{\infty} a_{mn}$ *and* $c_n := \sum_{m=1}^{\infty} a_{mn}$, $\sum_m |r_m| \leq L$ *and* $\sum_n |c_n| \leq L$.
*(iii)* $\sum_{m=1}^{\infty} r_m = S = \sum_{n=1}^{\infty} c_n$.

*Proof.*

(i) For fixed $m$, $N$, $\sum_{n=1}^{N} |a_{mn}| \leq L$; similarly $\sum_{m=1}^{M} |a_{mn}| \leq L$. Hence every row, column absolutely summable.
(ii) For $M \geq 1$, $\sum_{m=1}^{M} |r_m| \leq \sum_{m=1}^{M} \sum_{n=1}^{\infty} |a_{mn}| \leq L$; let $M \to \infty$. The column estimate is identical.
(iii) Given $\varepsilon > 0$, choose finite $F \subseteq \mathbb{N}^2$ with $\sum_{\mathbb{N}^2\setminus F} |a_{mn}| < \varepsilon$, then $R = \{1, \ldots, M\} \times \{1, \ldots, N\} \supseteq F$.
Each of $S$, $\sum_m r_m$, and $\sum_n c_n$ differs from $S_R := \sum_{(m,n)\in R} a_{mn}$ by at most $\sum_{\mathbb{N}^2\setminus R} |a_{mn}| \leq \varepsilon$. Let $\varepsilon \downarrow 0$.

*Structural Commentary*: Rectangles plus absolute tails justify Fubini.
*Key Insight*: Absolute summability makes order irrelevant. □

**Method 6.1.38.** *Series Proof Patterns*

(i) *Partial sums*: translate every assertion about $\sum a_n$ into an assertion about $s_n = \sum_{k=1}^{n} a_k$.
(ii) *Cauchy tails*: prove convergence by estimating $|\sum_{k=n}^{m} a_k|$ for $m \geq n$.
(iii) *Nonnegative terms*: replace convergence by boundedness of partial sums.
(iv) *Comparison*: dominate only the tail; finitely many early terms are irrelevant.
(v) *Absolute convergence*: use $\sum |a_n|$ when cancellation should not matter.
(vi) *Countable index sets*: choose a bijection $g : \mathbb{N} \to X$; use Theorem 6.1.36 so $\sum_{x\in X} a_x$ is independent of $g$.
(vii) *Double sums*: prove identities on $R_{M,N} := \{1, \ldots, M\} \times \{1, \ldots, N\}$; pass to $\infty$ arrays (Theorem 6.1.37).
(viii) *Conditional convergence*: order matters; use summation by parts or rearrangements.

*Key Insight*: Series arguments are sequence arguments with special notation for tails.

# 7 Uniform Convergence

## 7.1 Uniform Limits

Uniform convergence asks when limits of functions preserve structure. The key distinction is whether $N$ may depend on $x$, which controls continuity, integration, differentiation, and the supremum metric.

**Perspective 7.1.1.** *Function Limits Before Function-Sequence Limits*
$\lim_{t\to x} f(t)$ denotes the metric-space function limit of Definition 5.1.4.
Uniform convergence is the hypothesis that lets the limits in $n$ and $t$ commute.

**Definition 7.1.2.** *Pointwise Limits of Functions*
Let $f_n : E \to (Y, d_Y)$.

(i) $f_n \to f$ *pointwise* on $E$ if $d_Y(f_n(x), f(x)) \to 0$ for every $x \in E$.

(ii) If $Y$ is scalar, $\sum f_n \to f$ *pointwise* on $E$ if $s_N(x) := \sum_{n=1}^{N} f_n(x) \to f(x)$ for every $x \in E$.

No metric on $E$ is required; topology matters only for preserving continuity, limits, or integrals.

**Example 7.1.3.** *Pointwise Limits Can Lose Structure*

(i) $s_{m,n} := m/(m+n)$ gives $\lim_n \lim_m s_{m,n} = 1$ but $\lim_m \lim_n s_{m,n} = 0$.
(ii) For $f_n(x) = x^2/(1+x^2)^n$, $\sum_{n=0}^{\infty} f_n$ equals 0 at 0 and $1 + x^2$ otherwise, by Theorem 6.1.9.
(iii) $f_n(x) = \sin(nx)/\sqrt{n}$ satisfies $f_n \to 0$ pointwise but $f_n'(0) = \sqrt{n} \to +\infty$.
(iv) $f_n(x) = n^2 x(1-x^2)^n$ on $[0,1]$ satisfies $f_n \to 0$ pointwise, while $\int_0^1 f_n\,dx = n^2/(2n+2) \to +\infty$.

*Key Insight*: Pointwise control can miss global spikes and oscillations.

**Definition 7.1.4.** *Uniform Convergence*
For $f_n, f : E \to (Y, d_Y)$, $f_n \rightrightarrows f$ on $E$ if $\forall \varepsilon > 0\ \exists\ N \in \mathbb{N}$ such that $n \geq N,\ x \in E \implies d_Y(f_n(x), f(x)) \leq \varepsilon$.
In the scalar-valued case, take $d_Y(u,v) = |u - v|$; $\sum f_n$ converges uniformly on $E$ if its partial sums do.

**Perspective 7.1.5.** *One Tail, All Points*
Pointwise convergence lets $N$ depend on $x$; uniform convergence uses one $N$ for all $x$.
Equivalently, late graphs lie in one $\varepsilon$-strip around $f$.




Figure 181: One index $N$ puts all later graphs inside the same $\varepsilon$-strip around $f$ on $E$.

**Example 7.1.6.** *$x^n$ Converges Pointwise But Not Uniformly*
On $[0,1]$, let $g_n(x) = x^n$. Then $g_n \to g$ pointwise, where $g(x) = \begin{cases} 0, & 0 \leq x < 1, \\ 1, & x = 1. \end{cases}$
The convergence is not uniform: for each $n$, some $x < 1$ has $|x^n - g(x)| = x^n > 1/2$.




Figure 182: $x^n \to 0$ pointwise on $[0,1)$, but the endpoint prevents uniform convergence on $[0,1]$.

**Theorem 7.1.7.** *Cauchy Criterion for Uniform Convergence*
*For complex-valued $f_n : E \to \mathbb{C}$, $f_n$ converges uniformly on $E$ if and only if*

$$\forall \varepsilon > 0\ \exists N \in \mathbb{N} : \quad m, n \geq N,\ x \in E \implies |f_n(x) - f_m(x)| \leq \varepsilon$$

*Proof.*

($\Rightarrow$) If $f_n \rightrightarrows f$, choose $N$ so that $|f_k(x) - f(x)| \le \varepsilon/2$ for $k \ge N$ and $x \in E$. Then $|f_n(x) - f_m(x)| \le \varepsilon$.
($\Leftarrow$) For each $x$, $(f_n(x))$ is Cauchy in $\mathbb{C}$, so define $f(x) := \lim f_n(x)$.
Fix $n \ge N$ and let $m \to \infty$ in the criterion; then $|f_n(x) - f(x)| \le \varepsilon$ for all $x \in E$, hence $f_n \rightrightarrows f$.

*Structural Commentary*: The same $N$ gives Cauchy control simultaneously for every $x \in E$.
*Key Insight*: Uniform Cauchy tails are exactly uniform convergence. □

**Theorem 7.1.8.** *Supremum Test for Uniform Convergence*
*Suppose $f_n(x) \to f(x)$ $\forall x \in E$, and $M_n := \sup_{x\in E} |f_n(x) - f(x)|$. Then $f_n \rightrightarrows f$ on $E$ if and only if $M_n \to 0$.*

*Proof.*

($\Rightarrow$) If $f_n \rightrightarrows f$, then $\forall \varepsilon > 0$ $\exists$ $N$ such that $n \ge N$ implies $|f_n(x) - f(x)| \le \varepsilon$ for all $x \in E$. Hence $M_n \le \varepsilon$.
($\Leftarrow$) If $M_n \to 0$, then $\forall \varepsilon > 0$ $\exists$ $N$ such that $n \ge N$ implies $M_n \le \varepsilon$. Hence $|f_n(x) - f(x)| \le \varepsilon$ for all $x \in E$.

*Structural Commentary*: Uniform convergence is convergence in the supremum error.
*Key Insight*: The pointwise errors must vanish as one global quantity. □

**Theorem 7.1.9.** *Weierstrass M-Test*
*Let $f_n$ be functions on $E$ and suppose $|f_n(x)| \le M_n$ for every $x \in E$.*
*If $\sum M_n$ converges, then $\sum f_n$ converges uniformly on $E$.*

*Proof.* For $m \ge n$ and $x \in E$, $\left|\sum_{k=n}^{m} f_k(x)\right| \le \sum_{k=n}^{m} M_k$.
Since $\sum M_k$ converges, Theorem 6.1.5 makes the right-hand side $< \varepsilon$ for all large $m, n$.
Thus the partial sums of $\sum f_k$ are uniformly Cauchy, so Theorem 7.1.7 applies.
*Structural Commentary*: Absolute numerical tails dominate functional tails.
*Key Insight*: A summable uniform majorant gives uniform convergence of a function series. □

**Theorem 7.1.10.** *Uniform Convergence Interchanges Limits*
*Let $E$ lie in a metric space, let $x$ be a limit point of $E$, and suppose $f_n \rightrightarrows f$ on $E$.*
*If $A_n := \lim_{t\to x} f_n(t)$ exists for every $n$, then $(A_n)$ converges and $\lim_{t\to x} f(t) = \lim_{n\to\infty} A_n$.*

*Proof.* By Theorem 7.1.7, $\forall \varepsilon > 0$ $\exists$ $N$ such that $m, n \ge N$ and $t \in E$ imply $|f_n(t) - f_m(t)| \le \varepsilon$.
Letting $t \to x$ gives $|A_n - A_m| \le \varepsilon$, hence $A_n \to A$ for some $A \in \mathbb{C}$.
For any $n$, $|f(t) - A| \le |f(t) - f_n(t)| + |f_n(t) - A_n| + |A_n - A|$. Given $\varepsilon > 0$, choose $n$ so $|f(t) - f_n(t)| < \varepsilon/3$ for all $t \in E$ and $|A_n - A| < \varepsilon/3$; then take $t$ sufficiently close to $x$ so $|f_n(t) - A_n| < \varepsilon/3$. Thus $\lim_{t\to x} f(t) = A$.
*Structural Commentary*: Uniform convergence controls the outer two terms in the three-term estimate.
*Key Insight*: Uniformity is precisely what allows the $n \to \infty$ limit to move past the $t \to x$ limit. □

**Theorem 7.1.11.** *Uniform Limits Preserve Continuity*
*If $(f_n)$ is a sequence of continuous functions on $E$ and $f_n \rightrightarrows f$ on $E$, then $f$ is continuous on $E$.*

*Proof.* Let $x \in E$. If $x$ is isolated, $f$ is continuous at $x$.
If $x$ is a limit point, then $\lim_{t\to x} f_n(t) = f_n(x)$ for all $n$, so Theorem 7.1.10 gives $\lim_{t\to x} f(t) = \lim_{n\to\infty} f_n(x) = f(x)$.
*Structural Commentary*: Uniform convergence passes continuity to the limit.
*Key Insight*: Uniform convergence protects continuity from the failure modes in Example 7.1.3. □

**Theorem 7.1.12.** *Dini's Monotone Uniform Convergence Theorem*
*Let $K$ be compact and let $f_n, f : K \to \mathbb{R}$ be continuous. If $f_n(x) \downarrow f(x)$ for every $x \in K$, then $f_n \rightrightarrows f$ on $K$.*

*Proof.* Set $g_n := f_n - f$. Then $g_n$ is continuous, $g_n \downarrow 0$, and $g_n \ge 0$. Given $\varepsilon > 0$, let $K_n := \{x \in K : g_n(x) \ge \varepsilon\}$.
Then $(K_n)$ is nested compact and $\bigcap_n K_n = \emptyset$, so Corollary 4.1.9 gives $K_N = \emptyset$ for some $N$.
Thus $0 \le f_n(x) - f(x) < \varepsilon$ for all $x \in K$ and $n \ge N$.
*Structural Commentary*: Monotone convergence turns the bad sets into a nested compact sequence.
*Key Insight*: Compactness turns pointwise eventual escape from every bad set into a uniform escape. □

**Definition 7.1.13.** *The Supremum Metric on Bounded Continuous Functions*
Let $X$ be a metric space.

(i) $C_b(X)$ is the set of bounded continuous maps $X \to \mathbb{C}$.
(ii) For $f \in C_b(X)$, $\|f\|_\infty := \sup_{x\in X} |f(x)|$.
(iii) The *supremum metric* on $C_b(X)$ is $d_\infty(f, g) := \|f - g\|_\infty$.
(iv) $f_n \to f$ in $d_\infty$ iff $\|f_n - f\|_\infty \to 0$, i.e. iff $f_n \rightrightarrows f$ on $X$.

*Key Insight*: Uniform convergence is metric convergence in $C_b(X)$.

**Perspective 7.1.14.** *A Metric Space Whose Points Are Functions*
Uniform convergence is sequence convergence in the metric space $C_b(X)$.

**Perspective 7.1.15.** *Four Limit Notions*
The four limit notions are sequence limits, function limits, pointwise function limits, and uniform function limits. By Theorem 7.1.8, the last is ordinary sequence convergence in the supremum metric.

## 7.2 Interchange of Limits

**Theorem 7.2.1.** *Uniform Convergence and Riemann–Stieltjes Integration*
*Let $\alpha$ be increasing on $[a,b]$. If $f_n \in \mathcal{R}(\alpha)$ on $[a,b]$ and $f_n \rightrightarrows f$, then $f \in \mathcal{R}(\alpha)$ and $\int_a^b f\,d\alpha = \lim_{n\to\infty} \int_a^b f_n\,d\alpha$.*

*Proof.* Put $\varepsilon_n := \sup_{a\le x\le b} |f_n(x) - f(x)|$. Then $\varepsilon_n \to 0$ by Theorem 7.1.8, and $f_n - \varepsilon_n \le f \le f_n + \varepsilon_n$.
Upper and lower integrals of $f$ are therefore squeezed between corresponding integrals of $f_n \pm \varepsilon_n$, and gap is at most $2\varepsilon_n[\alpha(b) - \alpha(a)]$. Thus $f \in \mathcal{R}(\alpha)$. Same estimate gives $\left|\int_a^b f\,d\alpha - \int_a^b f_n\,d\alpha\right| \le \varepsilon_n[\alpha(b) - \alpha(a)] \to 0$.
*Structural Commentary*: Uniform error becomes an integral error multiplied by the total $\alpha$-variation.
*Key Insight*: Uniform convergence is strong enough to pass limits through the Riemann–Stieltjes integral. □

**Corollary 7.2.2.** *Termwise Integration of Uniformly Convergent Series*
*If $f_n \in \mathcal{R}(\alpha)$ on $[a,b]$ and $\sum f_n$ converges uniformly to $f$, then $\int_a^b f\,d\alpha = \sum_{n=1}^{\infty} \int_a^b f_n\,d\alpha$.*

*Proof.* Let $s_N := \sum_{n=1}^{N} f_n$. By linearity, $s_N \in \mathcal{R}(\alpha)$ and $\int_a^b s_N\,d\alpha = \sum_{n=1}^{N} \int_a^b f_n\,d\alpha$.
Since $s_N \rightrightarrows f$, Theorem 7.2.1 gives $\int_a^b f\,d\alpha = \lim_{N\to\infty} \int_a^b s_N\,d\alpha = \sum_{n=1}^{\infty} \int_a^b f_n\,d\alpha$.
*Structural Commentary*: Reduce the series to its uniformly convergent partial sums.
*Key Insight*: Uniform convergence lets integration be performed term by term. □

**Theorem 7.2.3.** *Differentiation Under a Riemann–Stieltjes Integral*
*Let $\alpha : [a,b] \to \mathbb{R}$ be increasing, let $\varphi : [a,b] \times [c,d] \to \mathbb{R}$, write $\varphi^t(x) := \varphi(x,t)$, and fix $s \in (c,d)$.*
*Assume $\varphi^t \in \mathcal{R}(\alpha)$ for every $t \in [c,d]$, and for some $\eta > 0$ with $(s-\eta, s+\eta) \subseteq (c,d)$, $D_2\varphi$ exists on $[a,b] \times (s-\eta, s+\eta)$ and $(D_2\varphi)^t \rightrightarrows (D_2\varphi)^s$ on $[a,b]$ as $t \to s$.*
*If $F(t) := \int_a^b \varphi(x,t)\,d\alpha(x)$, then $(D_2\varphi)^s \in \mathcal{R}(\alpha)$ and $F'(s) = \int_a^b (D_2\varphi)(x,s)\,d\alpha(x)$.*

*Proof.* For $t \ne s$, set $\psi_t := (\varphi^t - \varphi^s)/(t-s) \in \mathcal{R}(\alpha)$.
By Corollary 1.1.13, $\psi_t(x) = D_2\varphi(x, \tau_x(t))$ for some $\tau_x(t)$ between $s$ and $t$; hence $\psi_t \rightrightarrows (D_2\varphi)^s$.
Theorem 7.2.1 gives $(D_2\varphi)^s \in \mathcal{R}(\alpha)$ and $\frac{F(t)-F(s)}{t-s} = \int_a^b \psi_t\,d\alpha \to \int_a^b (D_2\varphi)^s\,d\alpha$, so $F'(s) = \int_a^b (D_2\varphi)(x,s)\,d\alpha(x)$.
*Structural Commentary*: Integrable difference quotients converge uniformly.
*Key Insight*: Uniform convergence permits $D_t \int = \int D_t$. □

**Theorem 7.2.4.** *Uniform Convergence and Differentiation*
*Let $f_n$ be differentiable on $[a,b]$, suppose $f_n(x_0)$ converges for some $x_0 \in [a,b]$, and suppose $f_n' \rightrightarrows g$ on $[a,b]$. Then $f_n \rightrightarrows f$ for some $f$, and $f' = g$ on $[a,b]$ (one-sided at endpoints).*

*Proof.* For $m, n$, MVT 1.1.13 gives $|f_n(x) - f_m(x)| \le |f_n(x_0) - f_m(x_0)| + (b-a) \sup_{t\in[a,b]} |f_n'(t) - f_m'(t)|$.
Thus $(f_n)$ is uniformly Cauchy, so $f_n \rightrightarrows f$. Fix $x$ and set $\phi_n(t) := \frac{f_n(t)-f_n(x)}{t-x}$, $\phi(t) := \frac{f(t)-f(x)}{t-x}$.
For $t \ne x$, the same estimate gives $|\phi_n(t) - \phi_m(t)| \le \|f_n' - f_m'\|_\infty$, hence $\phi_n \rightrightarrows \phi$.
By Theorem 7.1.10, $f'(x) = \lim_{t\to x} \phi(t) = \lim_{n\to\infty} f_n'(x) = g(x)$.
*Structural Commentary*: Derivative control anchors functions, then quotients.
*Key Insight*: Uniform derivative convergence plus one anchor passes derivatives to the limit. □

## 7.3 Further Topics in Uniform Convergence

**Method 7.3.1.** *Uniform Convergence Proof Patterns*

(i) *Pointwise*: fix $x$, then choose $N = N(x,\varepsilon)$.
(ii) *Uniform*: choose $N = N(\varepsilon)$; equivalently $\sup_E |f_n - f| \to 0$.
(iii) *Uniform Cauchy*: prove $\sup_E |f_n - f_m| \to 0$ as $m, n \to \infty$.
(iv) *Preservation*: use a three-term estimate and choose the uniform index first.
(v) *Series of functions*: apply Weierstrass $M$-test when numerical absolute tails dominate all functional tails.
(vi) *Termwise operations*: require $f_n \to f$ locally uniformly, i.e. $\sup_{x\in K} |f_n - f| \to 0$ for each compact $K$, before $\lim_n \int_K f_n = \int_K f$; for derivatives, require $f_n' \to g$ locally uniformly and get $(\lim_n f_n)' = g$.

*Key Insight*: Uniformity means $N$ is chosen before $x$.

# 8 Function-Space Compactness

## 8.1 Equicontinuity

Function-space compactness treats functions as points. Supremum metrics preview Banach spaces, Baire category describes generic behavior, and equicontinuity leads to Arzela–Ascoli.

**Theorem 8.1.1.** *Completeness of the Supremum Metric*
*The metric space $C_b(X)$ with metric $d(f,g) = \|f-g\|_\infty$ is complete.*

*Proof.* Let $(f_n)$ be Cauchy in $C_b(X)$. Then Theorem 7.1.7 gives a function $f$ with $f_n \rightrightarrows f$. By Theorem 7.1.11, $f$ is continuous. For some $n$, $\|f - f_n\|_\infty < 1$, hence $f_n$ is bounded, thus $f \in C_b(X)$ and $\|f_n - f\|_\infty \to 0$.
*Structural Commentary*: Uniform Cauchy convergence builds the limit inside the same function space.
*Key Insight*: Completeness of $\mathbb{C}$ plus uniform convergence makes $C_b(X)$ complete. □

**Definition 8.1.2.** *Bounded Target Metric for All Functions*
For metric spaces $(X, d_X)$ and $(Y, d_Y)$, let $\mathcal{F} = Y^X$ and set $\rho(y,z) := \frac{d_Y(y,z)}{1+d_Y(y,z)}$ and $d_\rho(f,g) := \sup_{x\in X} \rho(f(x), g(x))$. Then $\rho < 1$, so $d_\rho$ is finite on all of $\mathcal{F}$; $d_\rho$-convergence is uniform convergence measured by $d_Y$.

**Theorem 8.1.3.** *Completeness and Closed Subspaces for $d_\rho$*
*If the target metric space $(Y, d_Y)$ is complete, then:*

*(i)* $(\mathcal{F}, d_\rho)$ *is complete;*
*(ii)* $\mathcal{F}_b := \{f \in \mathcal{F} : f(X)$ *is* $d_Y$*-bounded*$\}$ *is closed in* $\mathcal{F}$*;*
*(iii)* $C(X,Y)$ *is closed in* $\mathcal{F}$*.*

*Proof.*

(i) Let $(f_n)$ be $d_\rho$-Cauchy. For each $x \in X$, $(f_n(x))$ is Cauchy in $Y$; define $f(x) := \lim_n f_n(x)$.
Given $\varepsilon > 0$, choose $N$ with $d_\rho(f_n, f_m) < \varepsilon$ for $m, n \geq N$. Letting $m \to \infty$ gives $d_\rho(f_n, f) \leq \varepsilon$ for $n \geq N$.
(ii) If $f_n \in \mathcal{F}_b$ and $d_\rho(f_n, f) \to 0$, choose $N$ with $d_\rho(f_N, f) < 1/2$.
Then $d_Y(f_N(x), f(x)) < 1$ for all $x$. Since $f_N(X)$ is bounded, so is $f(X)$.
(iii) If $f_n \in C(X,Y)$ and $d_\rho(f_n, f) \to 0$, then $f_n \to f$ uniformly in $d_Y$.
Given $x_0 \in X$ and $\varepsilon > 0$, choose $N$ with $d_Y(f_N(x), f(x)) < \varepsilon/3$ for all $x$, then use continuity of $f_N$ at $x_0$ and the triangle inequality. Thus $f$ is continuous.

*Structural Commentary*: Compress the target metric; then close the useful subspaces.
*Key Insight*: Completeness and uniform control make boundedness and continuity closed. □

**Definition 8.1.4.** *Boundedness and Equicontinuity of Function Families*
Let $\mathcal{F}$ be a family of functions from $E$ into a metric space $(Y, d_Y)$; in scalar cases, take $d_Y(u,v) = |u-v|$.

(i) $(f_n)$ is *pointwise bounded* on $E$ if $(f_n(x))$ is bounded in $Y$ for each $x \in E$.
(ii) A scalar sequence $(f_n)$ is *uniformly bounded* on $E$ if $\exists M$ such that $|f_n(x)| < M$ for all $x \in E, n \in \mathbb{N}$.
(iii) If $E$ lies in a metric space $(X, d_X)$, then $\mathcal{F}$ is *equicontinuous* on $E$ if $\forall \varepsilon > 0\ \exists \delta > 0$, such that $d_X(x,y) < \delta,\ x, y \in E,\ f \in \mathcal{F} \implies d_Y(f(x), f(y)) < \varepsilon$.
(iv) The family is *pointwise equicontinuous* if for every $p \in E$ and $\varepsilon > 0$ there is $\delta = \delta(p, \varepsilon) > 0$ such that $d_X(p,y) < \delta,\ y \in E,\ f \in \mathcal{F}$ implies $d_Y(f(p), f(y)) < \varepsilon$.




Figure 183: One input scale $\delta$ controls every curve in $\mathcal{F}$ within tolerance $\varepsilon$.

**Theorem 8.1.5.** *Pointwise Equicontinuity Becomes Uniform on Compact Sets*
*Every pointwise equicontinuous family $\mathcal{F} \subset C(K, \mathbb{C})$ on compact $K$ is equicontinuous.*

*Proof.* Otherwise, choose $\varepsilon > 0$, $f_n \in \mathcal{F}$, and $x_n, t_n \in K$ with $d(x_n, t_n) < 1/n$ but $|f_n(x_n) - f_n(t_n)| \geq \varepsilon$.
By Theorem 4.1.12, pass to $x_n \to x_0$; then $t_n \to x_0$ too.
Pointwise equicontinuity at $x_0$ gives $|f_n(x_n) - f_n(t_n)| < \varepsilon$ for large $n$, a contradiction.
*Structural Commentary*: Compactness turns moving counterexamples into one base point.
*Key Insight*: The local $\delta(x, \varepsilon)$ becomes one uniform $\delta$ only after compactness prevents escape. □

**Theorem 8.1.6.** *Uniform Convergence Implies Equicontinuity on Compact Sets*
*If $K$ is compact, $f_n \in C_b(K)$, and $f_n \rightrightarrows f$ on $K$, then $\{f_n : n \in \mathbb{N}\}$ is equicontinuous.*

*Proof.* Choose $N$ so $\|f_n - f_N\|_\infty < \varepsilon$ for $n > N$.
By Theorem 5.2.7, choose one $\delta > 0$ giving $\varepsilon$-control for $f_1, \ldots, f_N$.
If $n > N$, $d(x, y) < \delta$, then $|f_n(x) - f_n(y)| \leq |f_n(x) - f_N(x)| + |f_N(x) - f_N(y)| + |f_N(y) - f_n(y)| < 3\varepsilon$.
Together with the finite initial family, $\{f_n\}$ is equicontinuous.
*Structural Commentary*: A uniformly convergent tail stays close to one uniformly continuous function.
*Key Insight*: Compactness handles finitely many functions; uniform convergence handles the tail. □

**Example 8.1.7.** *Uniform Boundedness Alone Is Not Enough*
On $[0, 1]$, let $f_n(x) := \frac{x^2}{x^2 + (1 - nx)^2}$. Then $|f_n| \leq 1$ and $f_n \to 0$ pointwise, but $f_n(1/n) = 1$; hence no uniformly convergent subsequence. The missing hypothesis is equicontinuity.

**Theorem 8.1.8.** *Diagonal Selection on a Countable Set*
*Every pointwise bounded sequence $(f_n)$ on countable $E$ has a subsequence converging pointwise on $E$.*

*Proof.* List $E = \{x_1, x_2, \ldots\}$. Inductively choose nested subsequences $S_i$ so that $S_i(x_i)$ converges.
The diagonal subsequence is eventually contained in each $S_i$, hence converges at every $x_i$.
*Structural Commentary*: The diagonal keeps the $i$th convergence property after the $i$th step.
*Key Insight*: Countability permits diagonal convergence. □

## 8.2 Arzela-Ascoli-Type Compactness

**Theorem 8.2.1.** *Arzela–Ascoli Selection Theorem*
*Let $K$ be a compact metric space and let $(f_n) \subset C_b(K)$.*
*If $\sup_n |f_n(x)| < \infty$ for each $x \in K$ and $(f_n)$ is equicontinuous on $K$, then:*

*(i) $(f_n)$ is uniformly bounded on $K$;*
*(ii) $(f_n)$ contains a uniformly convergent subsequence.*

*Proof.*

(i) Choose $\delta > 0$ so $d(x, y) < \delta \Rightarrow |f_n(x) - f_n(y)| < 1$ $\forall n$. Let $K \subseteq \bigcup_{j=1}^{r} B_\delta(p_j)$, $M_j := \sup_n |f_n(p_j)| < \infty$.
If $x \in B_\delta(p_j)$, then $|f_n(x)| < 1 + M_j$; hence $\sup_{n, x \in K} |f_n(x)| \leq 1 + \max_j M_j$.

(ii) Let $E \subset K$ be countable dense, obtained from finite $1/m$-nets. Pointwise boundedness on $E$ and diagonal selection give a subsequence $(g_i)$ converging pointwise on $E$. Given $\varepsilon > 0$, choose $\delta > 0$ for equicontinuity with tolerance $\varepsilon/3$ and $x_1, \ldots, x_r \in E$ with $K \subseteq \bigcup_s B_\delta(x_s)$. For large $i, j$, $|g_i(x_s) - g_j(x_s)| < \varepsilon/3$ $\forall s$.
If $d(x, x_s) < \delta$, then $|g_i(x) - g_j(x)| < \varepsilon$, so $(g_i)$ is uniformly Cauchy and hence uniformly convergent.

*Structural Commentary*: Dense convergence plus equicontinuity becomes uniform convergence.
*Key Insight*: Equicontinuity turns finite sampling into uniform control. □




Figure 184: Arzela–Ascoli selects a uniformly convergent subsequence.

**Theorem 8.2.2.** *Compact Subsets of $C(K)$*
*Let $K$ be a compact metric space and let $C(K, \mathbb{R})$ carry the supremum metric.*
*For $\mathcal{F} \subseteq C(K, \mathbb{R})$, the following are equivalent:*

*(i) $\mathcal{F}$ is compact.*
*(ii) $\mathcal{F}$ is closed, uniformly bounded, and equicontinuous.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) Compactness gives closedness and total boundedness in $\|\cdot\|_\infty$. If $\mathcal{F} \neq \emptyset$, choose $g \in \mathcal{F}$; boundedness follows from $\|f\|_\infty \leq \|f-g\|_\infty + \|g\|_\infty$. Given $\varepsilon > 0$, take a finite $\varepsilon/3$-net $g_1, \ldots, g_r \in \mathcal{F}$.
For $d(x,y) < \delta$, chosen so $|g_i(x) - g_i(y)| < \frac{\varepsilon}{3}$ $\forall i$, any $f$ with $\|f - g_i\|_\infty < \frac{\varepsilon}{3}$ satisfies $|f(x) - f(y)| < \varepsilon$.

(*ii*) $\Rightarrow$ (*i*) Closedness makes $\mathcal{F}$ complete. For total boundedness, fix $\varepsilon > 0$, choose $\delta$ from equicontinuity at $\varepsilon/3$, and a finite $\delta$-net $x_1, \ldots, x_r \subset K$. With $|f| \leq M$ on $K$ for $f \in \mathcal{F}$, partition $[-M, M]$ into intervals of length $< \varepsilon/3$ and classify $f$ by the interval data of $(f(x_1), \ldots, f(x_r))$. For one representative $g$ per class and $d(x, x_i) < \delta$, $|f(x) - g(x)| \leq |f(x) - f(x_i)| + |f(x_i) - g(x_i)| + |g(x_i) - g(x)| < \varepsilon$. Hence $\mathcal{F}$ is totally bounded, so complete plus totally bounded implies compact.

*Structural Commentary*: Compactness in $C(K)$ is finite-resolution control plus closedness.
*Key Insight*: Equicontinuity lets finitely many sample values control whole functions. □

**Example 8.2.3.** *Hölder Balls Have Compact Closure*
For $0 < \alpha \leq 1$, define $[f]_\alpha := \sup_{x \neq y} \frac{|f(x)-f(y)|}{|x-y|^\alpha}$ when finite.
Set $\mathcal{H}_\alpha := \{f \in C([0,1], \mathbb{R}) : \|f\|_\infty + [f]_\alpha \leq 1\}$, with closure taken in $C([0,1], \mathbb{R})$.

(i) If $f \in \mathcal{H}_\alpha$, then $\|f\|_\infty \leq 1$.
(ii) If $f \in \mathcal{H}_\alpha$, then $|f(x) - f(y)| \leq |x-y|^\alpha$, so $\mathcal{H}_\alpha$ is equicontinuous.
(iii) Uniform limits preserve $\|f\|_\infty \leq 1$ and $|f(x) - f(y)| \leq |x-y|^\alpha$.

Thus $\overline{\mathcal{H}_\alpha}$ is closed, uniformly bounded, and equicontinuous; hence it is compact.

**Definition 8.2.4.** *$\sigma$-Compact Metric Space*
A metric space $X$ is *$\sigma$-compact* if $X = \bigcup_{j=1}^{\infty} K_j$ for compact subsets $K_j \subseteq X$.

**Theorem 8.2.5.** *Arzela–Ascoli on $\sigma$-Compact Spaces*
*Let $X$ be a $\sigma$-compact metric space, $Y$ a metric space, $f_n : X \to Y$ pointwise equicontinuous. If $\overline{\{f_n(x) : n \in \mathbb{N}\}}$ is compact in $Y$ $\forall x \in X$, then some subsequence $(g_m)$ converges uniformly on every compact $K \subseteq X$.*

*Proof.* Choose an increasing compact exhaustion $X = \bigcup_j K_j$.

(i) On each $K_j$, the restrictions $(f_n|_{K_j})$ are uniformly equicontinuous and pointwise relatively compact in $Y$.
(ii) Choose subsequences successively so the $j$th one converges uniformly on $K_j$.
The diagonal $g_m$ satisfies $g_m|_{K_j} \rightrightarrows g|_{K_j}$ for every $j$.
(iii) For compact $K \subseteq X$, $(g_m|_K)$ is uniformly equicontinuous and $g_m \to g$ pointwise.
Given $\varepsilon > 0$, take a finite $\delta$-net $E \subset K$ with equicontinuity tolerance $\varepsilon/3$.
Convergence on $E$ gives $\sup_{x \in K} d_Y(g_m(x), g_n(x)) < \varepsilon$ for large $m, n$, hence $g_m \rightrightarrows g$ on $K$.

*Structural Commentary*: Exhaust, select, diagonalize.
*Key Insight*: Compact-local control gives compact-local uniform convergence. □

**Corollary 8.2.6.** *Real-Line Local Uniform Subsequence*
*Let $f_n : \mathbb{R} \to \mathbb{R}$ be pointwise equicontinuous and suppose $\sup_n |f_n(x_0)| < \infty$ for some $x_0 \in \mathbb{R}$.*
*Then some subsequence converges uniformly on every compact $K \subseteq \mathbb{R}$.*

*Proof.* For $x \in \mathbb{R}$, take compact $I \supset \{x_0, x\}$. Uniform equicontinuity on $I$ gives $\delta > 0$ with step-size tolerance 1.
Choose a finite $\delta$-chain $x_0 = t_0, \ldots, t_m = x$; then $|f_n(x)| \leq |f_n(x_0)| + m$ for all $n$.
Thus each orbit $\{f_n(x) : n \in \mathbb{N}\}$ has compact closure in $\mathbb{R}$, and the $\sigma$-compact theorem applies.
*Structural Commentary*: One bounded orbit propagates by finite chains.
*Key Insight*: Equicontinuity turns one anchor into compact-local control. □

## 8.3 Further Topics in Function-Space Compactness

**Method 8.3.1.** *Function-Space Compactness Proof Patterns*

(i) *Completeness*: prove a function sequence is uniformly Cauchy, build the pointwise limit, and use uniform convergence to keep continuity or boundedness.
(ii) *Baire category*: show a desired property contains a countable intersection of dense open subsets of a complete function space.
(iii) *Compactness/equicontinuity*: pass from finite nets to all of $K$.
(iv) *Diagonal selection*: get pointwise convergence on a countable dense set, then upgrade using equicontinuity.
(v) *Locally uniform*: exhaust a $\sigma$-compact domain by compact sets and diagonalize.

*Key Insight*: Compactness in function spaces is finite-net control plus equicontinuity.

# 9 Power Series and Approximation

## 9.1 Power Series

Power series are controlled infinite polynomials, and approximation is uniform closure. This subsection covers radii, Taylor structure, Abel behavior, Weierstrass, Stone–Weierstrass, and elementary analytic functions.

**Definition 9.1.1.** *Core Objects: Power Series, Approximation, and Gamma*

(i) A real *power series about $a$* is $\sum_{n=0}^{\infty} c_n(x-a)^n$; a local representation of $f$ by one makes $f$ *analytic* at $a$.

(ii) A *polynomial approximant* to $f$ on a compact set $K$ is a polynomial $P$ chosen so that $\|f-P\|_{\infty,K}$ is small.

(iii) For $x>0$, the *gamma function* is $\Gamma(x) := \int_0^\infty t^{x-1}e^{-t}\,dt$ when the improper integral converges.

*Key Insight*: Power series are local analytic objects; approximation theorems turn algebraic families into dense function families; $\Gamma$ extends factorials by integration.

**Definition 9.1.2.** *Power Series and Radius of Convergence*
Let $(c_n)_{n\ge 0} \subseteq \mathbb{C}$ and $\alpha := \limsup_{n\to\infty} \sqrt[n]{|c_n|} \in [0,+\infty]$.

(i) The series $\sum_{n=0}^{\infty} c_n z^n$ is a *power series*.

(ii) The *radius of convergence* is $R := \alpha^{-1} \in [0,+\infty]$, with $0^{-1}=+\infty$ and $(+\infty)^{-1}=0$.

**Theorem 9.1.3.** *Convergence Inside the Circle*
*For $\sum c_n z^n$, radius $R$: converge if $|z|<R$, diverge if $|z|>R$; boundary $|z|=R$ is undecided.*

*Proof.* For $a_n = c_n z^n$, $\limsup_{n\to\infty} \sqrt[n]{|a_n|} = |z| \limsup_{n\to\infty} \sqrt[n]{|c_n|} = \frac{|z|}{R}$, with extended meaning if $R=0$ or $R=+\infty$. If $|z|<R$, this value is $<1$; if $|z|>R$, it is $>1$. Apply Theorem 6.1.20.
*Structural Commentary*: Radius controls the interior and exterior, not the boundary.
*Key Insight*: The root test is the radius-of-convergence theorem in disguise. □

**Theorem 9.1.4.** *A Boundary Test for Power Series*
*If $\sum c_n z^n$ has radius 1, $c_n \downarrow 0$, and $c_n \ge 0$, then it converges for $|z|=1$, except possibly at $z=1$.*

*Proof.* For $|z|=1$ and $z\neq 1$, put $a_n = z^n$ and $b_n = c_n$.
The partial sums of $\sum z^n$ are bounded because $\left|\sum_{m=0}^{n} z^m\right| = \left|\frac{1-z^{n+1}}{1-z}\right| \le \frac{2}{|1-z|}$. Theorem 6.1.24 applies.
*Structural Commentary*: Boundary convergence away from $z=1$ comes from bounded geometric partial sums.
*Key Insight*: Monotone coefficients can give conditional boundary convergence. □

**Theorem 9.1.5.** *Abel's Endpoint Theorem*
*Let $s_n := \sum_{k=0}^{n} c_k \to s$. For $|x|<1$, set $f(x) := \sum_{n\ge 0} c_n x^n$. Then $\lim_{x\to 1^-} f(x) = s$.*

*Proof.* Set $s_{-1} := 0$. Since $c_n = s_n - s_{n-1}$, for $|x|<1$, $f(x) = \sum_{n\ge 0}(s_n - s_{n-1})x^n = (1-x)\sum_{n\ge 0} s_n x^n$.
Thus $f(x) - s = (1-x)\sum_{n\ge 0}(s_n - s)x^n$. Given $\varepsilon>0$, choose $N$ with $|s_n - s| < \varepsilon/2$ for $n>N$.

For $0<x<1$, $|f(x)-s| \le (1-x)\sum_{n=0}^{N} |s_n - s|\,x^n + \frac{\varepsilon}{2}(1-x)\sum_{n>N} x^n$.
The first term tends to 0 as $x\to 1^-$, and the second is $\le \varepsilon/2$; hence $f(x)\to s$.
*Structural Commentary*: Endpoint convergence is one-sided Abel continuity.
*Key Insight*: $(1-x)\sum s_n x^n$ is a geometric average of partial sums. □

**Theorem 9.1.6.** *Abel Product Theorem for Cauchy Products*
*Let $\sum a_n = A$ and $\sum b_n = B$. Define $c_n := \sum_{k=0}^{n} a_k b_{n-k}$, and suppose $\sum c_n = C$. Then $C = AB$.*

*Proof.* For $|x|<1$, set $f(x) := \sum_{n\ge 0} a_n x^n$, $g(x) := \sum_{n\ge 0} b_n x^n$, and $h(x) := \sum_{n\ge 0} c_n x^n$. Since $(a_n),(b_n),(c_n)$ are bounded, these series converge absolutely on $|x|<1$. Hence $f(x)g(x) = \sum_{n\ge 0}\big(\sum_{k=0}^{n} a_k b_{n-k}\big)x^n = \sum_{n\ge 0} c_n x^n = h(x)$.
By Abel's endpoint thm, $f(x)\to A$, $g(x)\to B$, $h(x)\to C$ as $x\to 1^-$; $C = \lim_{x\to 1^-} h(x) = \lim_{x\to 1^-} f(x)g(x) = AB$.
*Structural Commentary*: Regularize by $x$, multiply inside, pass to $x=1$.
*Key Insight*: Abel moves the safe product identity to the endpoint. □

**Perspective 9.1.7.** *Centered Series Are Translated Series*
$\sum c_n(x-a)^n$ is $\sum c_n y^n$ with $y = x - a$ and radius $R$: absolute convergence for $|x-a| < R$, divergence for $|x-a| > R$, and separate endpoint tests for $|x-a| = R$.




Figure 185: Inside $|x-a| < R$: absolute and locally uniform; outside divergence; endpoints undecided.

**Definition 9.1.8.** *Real Analytic at a Point*
Let $E \subseteq \mathbb{R}$, $f : E \to \mathbb{R}$, $a \in \operatorname{int} E$. The function $f$ is *real analytic at $a$* if $\exists r > 0$ and $(c_n)_{n\geq 0} \subseteq \mathbb{R}$ such that:

(i) $(a-r, a+r) \subseteq E$.
(ii) The power series $\sum c_n(x-a)^n$ has radius of convergence $\geq r$.
(iii) $f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n$ for all $x \in (a-r, a+r)$.

If $E$ is open and $f$ is real analytic at every $a \in E$, then $f$ is *real analytic on $E$*. Analytic functions form the class $C^\omega$.

**Theorem 9.1.9.** *Power Series May Be Differentiated Term by Term*
*Let $f(x) = \sum_{n\geq 0} c_n x^n$ have radius $R > 0$, and set $I_R := (-R, R)$, $K_\rho := [-\rho, \rho]$ for $0 < \rho < R$.*
*(i) $\sum c_n x^n$ converges uniformly on each $K_\rho$, and $f \in C^\infty(I_R)$.*
*(ii) For $k \geq 1$ and $x \in I_R$, $f^{(k)}(x) = \sum_{n\geq k} \frac{n!}{(n-k)!} c_n x^{n-k}$; in particular, $f^{(k)}(0) = k! c_k$.*

*Proof.*
(i) If $0 < \rho < R$, then $\sum |c_n| \rho^n < \infty$; hence $|c_n x^n| \leq |c_n| \rho^n$ on $K_\rho$, so $\sum c_n x^n$ converges uniformly there.
(ii) For fixed $k$, $\frac{n!}{(n-k)!} \sim n^k$, so $\limsup \sqrt[n]{\left|\frac{n!}{(n-k)!} c_n\right|} = \limsup \sqrt[n]{|c_n|}$.
Hence every derivative series has radius $R$ and converges uniformly on each $K_\rho$.
Iterating termwise differentiation gives the formula and $f^{(k)}(0) = k! c_k$.

*Structural Commentary*: Work on $K_\rho \Subset I_R$ to justify termwise operations.
*Key Insight*: Polynomial coefficient growth does not change the radius. □

**Corollary 9.1.10.** *Termwise Calculus for Centered Power Series*
*Let $f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n$ have radius $R > 0$.*
*(i) The series converges locally uniformly on $(a-R, a+R)$ and defines a continuous function.*
*(ii) For $|x-a| < R$, $f'(x) = \sum_{n=1}^{\infty} n c_n (x-a)^{n-1}$, locally uniformly.*
*(iii) If $[y, z] \subset (a-R, a+R)$, then $\int_y^z f(x)\,dx = \sum_{n=0}^{\infty} c_n \frac{(z-a)^{n+1} - (y-a)^{n+1}}{n+1}$.*

*Proof.* Set $g(u) := \sum c_n u^n$ and $T(x) := x - a$, so $f = g \circ T$ on $(a-R, a+R)$.
(i) Local uniform convergence of $\sum c_n u^n$ on $(-R, R)$ implies local uniform convergence of $\sum c_n (Tx)^n$ on $(a-R, a+R)$; continuity follows from uniform limits.
(ii) Since $T'(x) = 1$, $f'(x) = g'(Tx) = \sum_{n=1}^{\infty} n c_n (Tx)^{n-1}$, with local uniform convergence.
(iii) If $[y, z] \subset (a-R, a+R)$, choose $\rho < R$ with $[y, z] \subset [a-\rho, a+\rho]$.
Uniform convergence on $[a-\rho, a+\rho]$ gives $\int_y^z f = \lim_N \int_y^z \sum_{n=0}^{N} c_n (Tx)^n\,dx$, yielding the stated formula.

*Structural Commentary*: Work on compact intervals strictly inside the radius.
*Key Insight*: Translation reduces centered series to ordinary power series. □

**Corollary 9.1.11.** *Analytic Functions Are Their Taylor Series*
*Suppose $f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n$ for $|x-a| < r$, with $r > 0$.*
*(i) $f \in C^\infty(a-r, a+r)$.*

*(ii)* $c_k = f^{(k)}(a)/k!$ *for every* $k \geq 0$.
*(iii)* *The power-series expansion of* $f$ *about* $a$ *is unique.*
*(iv)* $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n$ *for* $|x-a| < r$.

*Proof.*

(i) Termwise differentiation gives $f \in C^\infty(a-r, a+r)$.
(ii) For $k \geq 0$, $f^{(k)}(x) = \sum_{n \geq k} \frac{n!}{(n-k)!} c_n (x-a)^{n-k}$; at $x = a$, $f^{(k)}(a) = k! c_k$.
(iii) If $f = \sum b_n (x-a)^n$ also, then $b_k = f^{(k)}(a)/k! = c_k$ for all $k$.
(iv) Replace $c_n$ by $f^{(n)}(a)/n!$ in $f(x) = \sum c_n (x-a)^n$.

*Structural Commentary*: Derivatives determine every coefficient.
*Key Insight*: Analytic representation forces its own Taylor series. □

**Theorem 9.1.12.** *Re-Expansion and Identity for Power Series*
*Let* $f(x) = \sum_{n \geq 0} c_n x^n$ *have radius* $R > 0$, *and set* $S = (-R, R)$.

*(i)* *If* $a \in S$, *then* $f(x) = \sum_{m \geq 0} \frac{f^{(m)}(a)}{m!}(x-a)^m$ *for* $|x-a| < R - |a|$.
*(ii)* *Let* $F(x) = \sum_{n \geq 0} a_n x^n$ *and* $G(x) = \sum_{n \geq 0} b_n x^n$ *converge on* $S$.
*If* $\mathcal{Z}_{F,G} := \{x \in S : F(x) = G(x)\}$ *has a limit point in* $S$, *then* $a_n = b_n$ *for all* $n \geq 0$, *hence* $F = G$ *on* $S$.

*Proof.*

(i) Put $\rho := |x-a| + |a| < R$. Since $\sum_{n \geq 0} |c_n| \rho^n < \infty$, the binomial double sum is absolutely rearrangeable: $f(x) = \sum_{m \geq 0} A_m (x-a)^m$, where $A_m := \sum_{n \geq m} \binom{n}{m} c_n a^{n-m}$. Also $f^{(m)}(a) = \sum_{n \geq m} \frac{n!}{(n-m)!} c_n a^{n-m} = m! A_m$.
(ii) Set $H := F - G = \sum_{n \geq 0} d_n x^n$, $d_n := a_n - b_n$, $Z := \{x \in S : H(x) = 0\}$, and $A := Z' \cap S$.
For $x_0 \in A$, (i) gives $H(x) = \sum e_n (x - x_0)^n$ near $x_0$.
If $k = \min\{n : e_n \neq 0\}$ existed, then $H = (x-x_0)^k g$ with $g(x_0) = e_k \neq 0$, so $x_0$ would be isolated in $Z$, contradiction. Thus all $e_n = 0$, so $H = 0$ near $x_0$ and $A$ is open. Since $S \setminus A$ is open, $A \neq \emptyset$, and $S$ is connected, $A = S$. As $Z$ is closed in $S$, $Z = S$; hence $d_n = H^{(n)}(0)/n! = 0$.

*Structural Commentary*: Re-expand locally; connectedness propagates zeros.
*Key Insight*: Accumulating zeros force every local coefficient to vanish. □

**Perspective 9.1.13.** *Regularity Hierarchy*
The inclusions are $C^\omega \subseteq C^\infty \subseteq \cdots \subseteq C^2 \subseteq C^1 \subseteq C^0$.
The inclusions follow from Theorem 9.1.9 and Definition 1.1.23; they are generally proper.
*Key Insight*: Higher regularity is extra structure, not just a different notation for continuity.

**Example 9.1.14.** *Smooth but Not Analytic*
Define $e(x) = \begin{cases} e^{-1/x}, & x > 0, \\ 0, & x \leq 0. \end{cases}$. By repeated L'Hospital, Theorem 1.1.21, $e \in C^\infty$ and $e^{(r)}(0) = 0$ for all $r$.
Its Taylor series at 0 is zero, but $e(x) > 0$ for $x > 0$, so $e$ is not analytic at 0 (Definition 9.1.8).
*Key Insight*: A function can be infinitely tangent to the $x$-axis without being locally equal to its Taylor series.

**Perspective 9.1.15.** *Taylor Coefficients Are Only the First Half*
Power-series coefficients are forced by derivatives (Corollary 9.1.11); for a function not already known to be analytic, proving the Taylor series represents the function needs a remainder estimate (Theorem 1.1.25).

**Theorem 9.1.16.** *Exponential and Logarithm from the Power Series*
*Define* $\mathsf{E}(z) := \sum_{n=0}^{\infty} \frac{z^n}{n!}$ *for* $z \in \mathbb{C}$.

*(i)* $\mathsf{E}(z+w) = \mathsf{E}(z)\mathsf{E}(w)$ *for* $z, w \in \mathbb{C}$.
*(ii)* *On* $\mathbb{R}$, $\mathsf{E} > 0$, $\mathsf{E}' = \mathsf{E}$, $\mathsf{E}(x) = e^x$, *and* $\mathsf{E}$ *is strictly increasing.*
*(iii)* *The inverse* $\mathsf{L} := \mathsf{E}^{-1} : (0, \infty) \to \mathbb{R}$ *satisfies* $\mathsf{L}'(y) = 1/y$ *and* $\mathsf{L}(uv) = \mathsf{L}(u) + \mathsf{L}(v)$ *for* $u, v > 0$.
*(iv)* *For* $x > 0$ *and* $\alpha \in \mathbb{R}$, *set* $x^\alpha := \mathsf{E}(\alpha \mathsf{L}(x))$; *then* $(x^\alpha)' = \alpha x^{\alpha - 1}$.

*Proof.* The ratio test gives convergence for all $z \in \mathbb{C}$.

(i) Absolute convergence gives $\mathsf{E}(z)\mathsf{E}(w) = \sum_{n \geq 0} \sum_{j=0}^{n} \frac{z^j w^{n-j}}{j!(n-j)!} = \sum_{n \geq 0} \frac{(z+w)^n}{n!} = \mathsf{E}(z+w)$.

(ii) Termwise differentiation gives $\mathsf{E}' = \mathsf{E}$ on $\mathbb{R}$. By (i), $\mathsf{E}(x)\mathsf{E}(-x) = 1$, so continuity and $\mathsf{E}(0) = 1$ give $\mathsf{E} > 0$; hence $\mathsf{E}' = \mathsf{E} > 0$. Since $\mathsf{E}(1) = e$, (i) gives $\mathsf{E}(p) = e^p$ for $p \in \mathbb{Q}$, and continuity gives $\mathsf{E}(x) = e^x$.
(iii) For $y = \mathsf{E}(x)$, differentiating $\mathsf{L}(\mathsf{E}(x)) = x$ gives $\mathsf{L}'(y) = 1/\mathsf{E}'(x) = 1/y$.
If $u = \mathsf{E}(s)$ and $v = \mathsf{E}(t)$, then $\mathsf{L}(uv) = \mathsf{L}(\mathsf{E}(s+t)) = s + t = \mathsf{L}(u) + \mathsf{L}(v)$.
(iv) $\frac{d}{dx}\mathsf{E}(\alpha\mathsf{L}(x)) = \mathsf{E}(\alpha\mathsf{L}(x))\alpha\mathsf{L}'(x) = x^\alpha\alpha/x = \alpha x^{\alpha-1}$.

*Structural Commentary*: Addition, inversion, powers.
*Key Insight*: $\mathsf{E}(z+w) = \mathsf{E}(z)\mathsf{E}(w)$ creates the logarithm law. □

**Perspective 9.1.17.** *Real Powers by Rational Approximation*
One route defines $x^\alpha$ for $x > 0$ by choosing $q_n \in \mathbb{Q}$ with $q_n \to \alpha$ and setting $x^\alpha = \lim_{n\to\infty} x^{q_n}$ after proving existence and independence of $(q_n)$. The power-series construction sets $x^\alpha = \mathsf{E}(\alpha\mathsf{L}(x))$; both routes must agree with rational powers and be choice-independent.

**Perspective 9.1.18.** *Logarithmic Integral Route*
With the Riemann integral, set $L(y) := \int_1^y \frac{1}{t}\,dt$ for $y > 0$. Theorem 2.2.1 gives $L'(y) = 1/y$; endpoint limits make $L$ invertible, and Theorem 1.1.6 then recovers $E' = E$ and the product laws.

**Perspective 9.1.19.** *From Exponentials to Factorials*
Theorems 9.1.16, 2.2.3 supply real-power laws and integration-by-parts pattern used for $\Gamma$ in Theorem 9.1.22.

**Theorem 9.1.20.** *Trigonometric Functions from the Exponential*
*For $x \in \mathbb{R}$, set $\mathsf{C}(x) := \frac{\mathsf{E}(ix)+\mathsf{E}(-ix)}{2}$, $\mathsf{S}(x) := \frac{\mathsf{E}(ix)-\mathsf{E}(-ix)}{2i}$, and $\gamma(t) := \mathsf{E}(it)$.*
*(i)* $\mathsf{E}(ix) = \mathsf{C}(x) + i\mathsf{S}(x)$, $\mathsf{C}^2 + \mathsf{S}^2 = 1$, $\mathsf{C}' = -\mathsf{S}$, *and* $\mathsf{S}' = \mathsf{C}$.
*(ii)* $x_0 := \min\{x > 0 : \mathsf{C}(x) = 0\}$ *exists; with* $\pi := 2x_0$, $\mathsf{E}(\pi i) = -1$, $\mathsf{E}(2\pi i) = 1$, $\mathsf{E}(z + 2\pi i) = \mathsf{E}(z)$ *for* $z \in \mathbb{C}$.
*(iii)* $\mathsf{C}, \mathsf{S}$ *are* $2\pi$*-periodic,* $\gamma([0, 2\pi]) = \{\zeta \in \mathbb{C} : |\zeta| = 1\}$, *and* $\gamma$ *is injective on* $[0, 2\pi)$.

*Proof.*
(i) Real coefficients give $\overline{\mathsf{E}(ix)} = \mathsf{E}(-ix)$, so $\mathsf{E}(ix) = \mathsf{C} + i\mathsf{S}$ and $|\mathsf{E}(ix)|^2 = \mathsf{E}(ix)\mathsf{E}(-ix) = 1$.
Thus $\mathsf{C}^2 + \mathsf{S}^2 = 1$; differentiating the definitions gives $\mathsf{C}' = -\mathsf{S}$ and $\mathsf{S}' = \mathsf{C}$.
(ii) If $\mathsf{C}$ has no positive zero, then $\mathsf{C} > 0$ on $(0, \infty)$, hence $\mathsf{S}' > 0$ and $\mathsf{S}(x) > 0$ for $x > 0$.
For $0 < x < y$, $\mathsf{S}(x)(y - x) < \int_x^y \mathsf{S}(t)\,dt = \mathsf{C}(x) - \mathsf{C}(y) \le 2$, impossible as $y \to \infty$; continuity gives the first zero $x_0$. On $(0, x_0)$, $\mathsf{C} > 0$, so $\mathsf{S}$ increases from 0 to $\mathsf{S}(x_0) = 1$.
Since $x_0 = \pi/2$, $\mathsf{E}(\pi i/2) = i$, and $\mathsf{E}(z+w) = \mathsf{E}(z)\mathsf{E}(w)$ gives $\mathsf{E}(\pi i) = -1$, $\mathsf{E}(2\pi i) = 1$, and $\mathsf{E}(z+2\pi i) = \mathsf{E}(z)$.
(iii) Periodicity of $\mathsf{C}, \mathsf{S}$ follows from (ii). On $[0, \pi/2]$, $\mathsf{C}$ decreases from 1 to 0 and $\mathsf{S}$ increases from 0 to 1, so $\gamma$ traverses the first quadrant once. Since $\gamma(t + m\pi/2) = \gamma(t)i^m$ for $m = 0, 1, 2, 3$, the four quadrants give $\gamma([0, 2\pi]) = \{\zeta : |\zeta| = 1\}$ and injectivity on $[0, 2\pi)$.

*Structural Commentary*: $\mathsf{E}(it)$ supplies trigonometry.
*Key Insight*: $\mathsf{E}(it)\mathsf{E}(-it) = 1$ puts the path on the unit circle. □




Figure 186: $\mathsf{E}(it) = \mathsf{C}(t) + i\mathsf{S}(t)$ traces the unit circle, defining sine and cosine analytically.

**Theorem 9.1.21.** *Algebraic Completeness of $\mathbb{C}$*
*If $P \in \mathbb{C}[z]$ and $\deg P \ge 1$, then $\exists z_0 \in \mathbb{C}$ such that $P(z_0) = 0$.*

*Proof.* Write $P(z) = \sum_{j=0}^n a_j z^j$, $n \ge 1$, $a_n = 1$.
Since $|P(z)| \to \infty$ as $|z| \to \infty$, choose $z_0$ with $\mu := |P(z_0)| = \min_{\mathbb{C}} |P|$. If $\mu = 0$, done.
Assume $\mu > 0$ and set $Q(w) := P(z_0 + w)/P(z_0)$. Then $Q(0) = 1$ and $|Q(w)| \ge 1$ for all $w$.
Write $Q(w) = 1 + b_k w^k + w^{k+1}H(w)$ with $k \ge 1$, $b_k \ne 0$, and $H$ polynomial.
Choose $\theta \in \mathbb{R}$ with $b_k e^{ik\theta} = -|b_k|$. For $r > 0$, $Q(re^{i\theta}) = 1 - r^k A(r)$, where $A(r) := |b_k| - rH(re^{i\theta})$.
For small $r$, $\Re A(r) > |b_k|/2$, hence $\left|1 - r^k A(r)\right|^2 = 1 - 2r^k \Re A(r) + r^{2k}|A(r)|^2 < 1$, contradicting $|Q| \ge 1$.

*Structural Commentary*: Minimize $|P|$, translate, perturb.
*Key Insight*: A phase choice makes the first nonzero term decrease $|Q|$. □

**Theorem 9.1.22.** *Gamma Function, Beta Function, and Stirling Skeleton*
*For $x > 0$, set $\Gamma(x) := \int_0^\infty t^{x-1}e^{-t}\,dt$; for $x, y > 0$, set $\mathrm{B}(x,y) := \int_0^1 t^{x-1}(1-t)^{y-1}\,dt$.*

*(i)* $\Gamma(x+1) = x\Gamma(x)$ *for* $x > 0$, $\Gamma(n+1) = n!$ *for* $n \in \mathbb{N}$, *and* $\log\Gamma$ *is convex on* $(0,\infty)$.
*(ii)* $\Gamma$ *is the unique* $f : (0,\infty) \to (0,\infty)$ *with* $f(1) = 1$, $f(x+1) = xf(x)$, *and* $\log f$ *convex.*
*(iii)* $\mathrm{B}(x,y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$ *for* $x, y > 0$.
*(iv)* $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$, $\int_{-\infty}^{\infty} e^{-s^2}\,ds = \sqrt{\pi}$, $\Gamma(x) = \frac{2^{x-1}}{\sqrt{\pi}}\Gamma\left(\frac{x}{2}\right)\Gamma\left(\frac{x+1}{2}\right)$, *and* $\lim_{x\to\infty} \frac{\Gamma(x+1)}{(x/e)^x\sqrt{2\pi x}} = 1$.

*Proof.*

(i) Integration by parts gives $\Gamma(x+1) = \int_0^\infty t^x e^{-t}\,dt = x\Gamma(x)$; $\Gamma(1) = 1$ gives $\Gamma(n+1) = n!$.
For $0 < \lambda < 1$, Holder gives $\Gamma(\lambda x + (1-\lambda)y) \le \Gamma(x)^\lambda\Gamma(y)^{1-\lambda}$, hence $\log\Gamma$ is convex.

(ii) Let $\varphi = \log f$. For $0 < x < 1$ and $n \ge 1$, convexity with $\varphi(t+1) = \varphi(t) + \log t$ gives
$0 \le \varphi(x) - \log\frac{n!\,n^x}{x(x+1)\cdots(x+n)} \le x\log(1+\frac{1}{n})$. Letting $n \to \infty$ fixes $\varphi$ on $(0,1)$; recurrence fixes $\varphi$ on $(0,\infty)$.

(iii) Fix $y > 0$ and set $F_y(x) := \frac{\Gamma(x+y)}{\Gamma(y)}\mathrm{B}(x,y)$. Since $\mathrm{B}(1,y) = 1/y$ and $\mathrm{B}(x+1,y) = \frac{x}{x+y}\mathrm{B}(x,y)$, $F_y(1) = 1$ and $F_y(x+1) = xF_y(x)$; Holder gives log-convexity. Thus (ii) gives $F_y = \Gamma$.

(iv) $t = \sin^2\theta$ gives $\mathrm{B}(\frac{1}{2},\frac{1}{2}) = \pi$, hence $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ by (iii). Then $t = s^2$ gives $\Gamma(\frac{1}{2}) = \int_{-\infty}^{\infty} e^{-s^2}\,ds$.
Beta duplication gives the duplication formula, and $t = x(1+u)$ with Laplace asymptotics gives Stirling.

*Structural Commentary*: Recurrence plus log-convexity gives uniqueness.
*Key Insight*: Bohr–Mollerup turns beta identities into gamma identities. □

**Method 9.1.23.** *Gamma Function Proof Pattern*

(i) *Gamma function*: use $\Gamma(x+1) = x\Gamma(x)$, $\Gamma(1) = 1$, and log-convexity for uniqueness.

## 9.2 Approximation Methods




Figure 187: Finer piecewise-linear interpolants approach $f(x) = \sqrt{1-x}$ uniformly on $[0,1]$.

**Theorem 9.2.1.** *Weierstrass Polynomial Approximation*
*Let $a < b$ and $f \in C([a,b],\mathbb{C})$. Then $\exists (P_n) \subset \mathbb{C}[x]$ with $P_n \rightrightarrows f$ on $[a,b]$. If $f([a,b]) \subset \mathbb{R}$, choose $(P_n) \subset \mathbb{R}[x]$.*

*Proof.* By affine change and subtraction of the endpoint interpolant, reduce to $[0,1]$ with $f(0) = f(1) = 0$.
Extend $f = 0$ off $[0,1]$; then $f$ is uniformly continuous on $[-1,2]$. Choose $c_n > 0$ so $Q_n(t) := c_n(1-t^2)^n$ satisfies $\int_{-1}^1 Q_n = 1$. Use $c_n < \sqrt{n}$ and $Q_n(t) \le \sqrt{n}(1-\delta^2)^n$ for $0 < \delta \le |t| \le 1$. Set $P_n(x) := \int_{-1}^1 f(x+t)Q_n(t)\,dt$.
Since $P_n(x) = \int_0^1 f(u)Q_n(u-x)\,du$, $P_n \in \mathbb{C}[x]$, and $P_n \in \mathbb{R}[x]$ if $f$ is real-valued.
Given $\varepsilon > 0$, choose $0 < \delta \le 1$ with $|u-v| < \delta \Rightarrow |f(u) - f(v)| < \varepsilon/2$ on $[-1,2]$, and set $M = \sup_{[-1,2]}|f|$.
For $x \in [0,1]$, $Q_n \ge 0$, $\int_{-1}^1 Q_n = 1$ give $|P_n(x) - f(x)| \le \int_{-1}^1 |f(x+t) - f(x)|\,Q_n(t)\,dt \le \frac{\varepsilon}{2} + 4M\sqrt{n}(1-\delta^2)^n$.
Thus $\|P_n - f\|_\infty \to 0$ on $[0,1]$; undo the reductions.
*Structural Commentary*: Polynomial kernels approximate the identity.
*Key Insight*: Uniform continuity handles near mass; kernel concentration removes far mass. □




Figure 188: The normalized kernels $Q_n(t) = c_n(1-t^2)^n$ concentrate near 0 while retaining integral 1.

**Corollary 9.2.2.** *Polynomial Approximation of $|x|$*
*Let $a > 0$. There exists $(P_n) \subset \mathbb{R}[x]$ such that $P_n(0) = 0$ and $P_n \rightrightarrows |x|$ on $[-a, a]$.*

*Proof.* By Theorem 9.2.1, choose $Q_n \in \mathbb{R}[x]$ with $Q_n \rightrightarrows |x|$ on $[-a, a]$. Set $P_n(x) := Q_n(x) - Q_n(0)$.
Then $P_n \in \mathbb{R}[x]$, $P_n(0) = 0$, and $Q_n(0) \to 0$, so $\|P_n - |x|\|_\infty \le \|Q_n - |x|\|_\infty + |Q_n(0)| \to 0$.
*Structural Commentary*: Normalize at 0.
*Key Insight*: The constant correction vanishes uniformly. □

**Definition 9.2.3.** *Algebras and Uniform Closure*
Let $\mathcal{A}$ be a family of complex-valued functions on a set $E$.

(i) $\mathcal{A}$ is an *algebra* if it is closed under $+$, products, and scalar multiplication (real scalars in the real case).
(ii) $\mathcal{A}$ is *uniformly closed* if $f_n \in \mathcal{A}$ and $f_n \rightrightarrows f$ on $E$ imply $f \in \mathcal{A}$.
(iii) The *uniform closure* of $\mathcal{A}$ is $\{f : f_n \in \mathcal{A},\ f_n \rightrightarrows f \text{ on } E\}$.

**Theorem 9.2.4.** *Uniform Closures Are Closed Algebras*
*The uniform closure of an algebra of bounded functions is a uniformly closed algebra.*

*Proof.* If $f, g \in \mathcal{B}$, choose $f_n, g_n \in \mathcal{A}$ with $f_n \rightrightarrows f$ and $g_n \rightrightarrows g$.
Boundedness and uniform convergence give $f_n + g_n \rightrightarrows f + g$; $f_n g_n \rightrightarrows fg$; and $cf_n \rightrightarrows cf$. Thus $\mathcal{B}$ is an algebra.
If $h_n \in \mathcal{B}$ and $h_n \rightrightarrows h$, approximate each $h_n$ uniformly by some $a_n \in \mathcal{A}$ within $1/n$; then $a_n \rightrightarrows h$, so $h \in \mathcal{B}$.
*Structural Commentary*: Uniform limits preserve algebraic operations for bounded functions.
*Key Insight*: The uniform closure of an algebra has already absorbed all further uniform limits. □

**Definition 9.2.5.** *Separation, Nonvanishing, and Self-Adjointness*
Let $\mathcal{A}$ be a family of functions on $E$.

(i) $\mathcal{A}$ *separates points* on $E$ if for every distinct $x_1, x_2 \in E$ there is $f \in \mathcal{A}$ with $f(x_1) \neq f(x_2)$.
(ii) $\mathcal{A}$ *vanishes at no point* of $E$ if for every $x \in E$ there is $g \in \mathcal{A}$ with $g(x) \neq 0$.
(iii) A complex algebra $\mathcal{A}$ is *self-adjoint* if $f \in \mathcal{A}$ implies $\overline{f} \in \mathcal{A}$, where $\overline{f}(x) := \overline{f(x)}$.

**Theorem 9.2.6.** *Two-Point Interpolation in a Separating Algebra*
*Let $\mathcal{A}$ be a scalar algebra on $E$ that separates points and vanishes nowhere.*
*For $x_1 \neq x_2$ in $E$ and scalars $c_1, c_2$, $\exists f \in \mathcal{A}$ such that $f(x_i) = c_i$, $i = 1, 2$.*

*Proof.* Choose $g, h, k \in \mathcal{A}$ with $g(x_1) \neq g(x_2)$, $h(x_1) \neq 0$, $k(x_2) \neq 0$. Set $u := gk - g(x_1)k$ and $v := gh - g(x_2)h$.
Then $u(x_1) = v(x_2) = 0$, while $u(x_2) = (g(x_2) - g(x_1))k(x_2) \neq 0$ and $v(x_1) = (g(x_1) - g(x_2))h(x_1) \neq 0$.
Hence $f := \frac{c_1 v}{v(x_1)} + \frac{c_2 u}{u(x_2)} \in \mathcal{A}$ and $f(x_i) = c_i$, $i = 1, 2$.
*Structural Commentary*: Build point-isolating functions.
*Key Insight*: Isolation reduces interpolation to linear combination. □

**Perspective 9.2.7.** *Interpolation Is Linear Algebra*
The two-point interpolation step can also be written as a $2 \times 2$ linear system. Its invertibility is exactly the finite-dimensional criterion from Theorem 1.7.8, with the determinant test supplied by Theorem 1.1.8. The proof above chooses explicit point-isolating functions instead of reopening the calculation.

**Theorem 9.2.8.** *Stone–Weierstrass Theorem, Real Form*
*Let $K$ be compact Hausdorff; $\mathcal{A} \subset C(K, \mathbb{R})$ point-separating, nowhere-vanishing real algebra. $\overline{\mathcal{A}}^{\|\cdot\|_\infty} = C(K, \mathbb{R})$.*

*Proof.* Put $\mathcal{B} := \overline{\mathcal{A}}^{\|\cdot\|_\infty}$. By Theorem 9.2.4, $\mathcal{B}$ is a uniformly closed algebra and inherits separation/nonvanishing.
For $r \in \mathcal{B}$, if $a = \|r\|_\infty > 0$, Corollary 9.2.2 gives $p_n(0) = 0$ with $p_n \rightrightarrows |t|$ on $[-a, a]$, hence $p_n(r) \rightrightarrows |r|$.
$a = 0$ is trivial. Thus $|r| \in \mathcal{B}$, so for $r, s \in \mathcal{B}$, $r \vee s = \frac{r+s+|r-s|}{2}$ and $r \wedge s = \frac{r+s-|r-s|}{2}$ lie in $\mathcal{B}$.
Fix $f \in C(K, \mathbb{R})$ and $\varepsilon > 0$. For $x, y \in K$, choose $h_{x,y} \in \mathcal{B}$ with $h_{x,y}(x) = f(x)$, $h_{x,y}(y) = f(y)$, using Theorem 9.2.6 if $x \neq y$ and $f(x)q/q(x)$ with $q \in \mathcal{B}$, $q(x) \neq 0$, if $x = y$. For fixed $x$, set $V_y := \{h_{x,y} > f - \varepsilon\}$.

Since $y \in V_y$, choose $y_i$ with $K = \bigcup_i V_{y_i}$ and put $g_x := \bigvee_i h_{x,y_i} \in \mathcal{B}$. Then $f - \varepsilon < g_x$ and $g_x(x) = f(x)$.
Set $U_x := \{g_x < f + \varepsilon\}$. Since $x \in U_x$, choose $x_j$ with $K = \bigcup_j U_{x_j}$ and put $h := \bigwedge_j g_{x_j} \in \mathcal{B}$.
Then $f - \varepsilon < h < f + \varepsilon$, so $\|h - f\|_\infty < \varepsilon$. Since $\mathcal{B}$ is uniformly closed, $f \in \mathcal{B}$.
*Structural Commentary*: Finite joins/meets globalize interpolation.
*Key Insight*: Compactness upgrades pointwise to uniform. □

**Theorem 9.2.9.** *Stone–Weierstrass Theorem, Complex Form*
*Let $K$ be compact Hausdorff and let $\mathcal{A} \subset C(K, \mathbb{C})$ be a self-adjoint, point-separating, nowhere-vanishing complex algebra. Then $\overline{\mathcal{A}}^{\|\cdot\|_\infty} = C(K, \mathbb{C})$.*

*Proof.* Set $\mathcal{A}_\mathbb{R} := \mathcal{A} \cap C(K, \mathbb{R})$. Self-adjointness gives $\operatorname{Re} f, \operatorname{Im} f \in \mathcal{A}_\mathbb{R}$ for all $f \in \mathcal{A}$. If $f(x) \neq f(y)$ or $f(x) \neq 0$, then one of $\operatorname{Re} f, \operatorname{Im} f$ witnesses the same condition; hence $\mathcal{A}_\mathbb{R}$ is point-separating and nowhere-vanishing.
By Theorem 9.2.8, $\overline{\mathcal{A}_\mathbb{R}} = C(K, \mathbb{R})$. For $F = u + iv \in C(K, \mathbb{C})$, take $u_n, v_n \in \mathcal{A}_\mathbb{R}$ with $u_n \rightrightarrows u$, $v_n \rightrightarrows v$.
Then $u_n + iv_n \in \mathcal{A}$ and $u_n + iv_n \rightrightarrows F$.
*Structural Commentary*: Self-adjointness yields a real subalgebra.
*Key Insight*: Approximate real and imaginary parts. □

**Theorem 9.2.10.** *Riesz Separability Criterion for $C(K)$*
*Let $K$ be a compact Hausdorff space. Then $(C(K, \mathbb{R}), \|\cdot\|_\infty)$ is separable iff $K$ is metrizable.*

*Proof.*
($\Rightarrow$) Let $\{g_n\}$ be dense in $C(K, \mathbb{R})$ and $O_n := \{g_n > 1/2\}$. Each $O_n$ is open. For open $O \ni x$, Urysohn gives $g \in C(K, \mathbb{R})$ with $0 \leq g \leq 1$, $g(x) = 1$, and $g|_{K \setminus O} = 0$. Pick $n$ with $\|g - g_n\|_\infty < 1/3$; then $x \in O_n \subseteq O$.
Thus $\{O_n\}$ is a countable basis, so Theorem 6.4.23 makes $K$ metrizable.
($\Leftarrow$) Let $d$ metrize $K$ and choose dense $\{x_n : n \geq 1\} \subset K$. Put $f_0 = 1$ and $f_n(x) = d(x, x_n)$.
If $x \neq y$, choose $x_n$ with $d(x, x_n) < d(x, y)/3$; then $f_n(x) \neq f_n(y)$. Hence $\{f_n : n \geq 0\}$ separates points and vanishes nowhere, so Theorem 9.2.8 makes $\operatorname{alg}(f_n : n \geq 0)$ dense in $C(K, \mathbb{R})$.
Its rational polynomial subalgebra is countable and dense.

*Structural Commentary*: Dense functions give a base; dense points give generators.
*Key Insight*: Countability moves between $K$ and $C(K)$. □

**Theorem 9.2.11.** *A Continuous Nowhere Differentiable Function Exists*
*There exists a real continuous function on $\mathbb{R}$ which is nowhere differentiable.*

*Proof.* Let $\varphi$ be the 2-periodic extension of $|x|$ on $[-1, 1]$. Then $0 \leq \varphi \leq 1$ and $|\varphi(s) - \varphi(t)| \leq |s - t|$.
Define $f(x) := \sum_{n=0}^{\infty} \left(\frac{3}{4}\right)^n \varphi(4^n x)$, converges uniformly (Theorem 7.1.9); hence $f$ is continuous (Theorem 7.1.11).
Fix $x$ and $m \in \mathbb{N}$. Choose $\delta_m = \pm\frac{1}{2}4^{-m}$ so that no integer lies between $4^m x$ and $4^m(x + \delta_m)$.
Set $\gamma_{n,m} := \frac{\varphi(4^n(x+\delta_m)) - \varphi(4^n x)}{\delta_m}$. If $n > m$, then $4^n \delta_m$ is an even integer, so $\gamma_{n,m} = 0$.
If $0 \leq n \leq m$, then $|\gamma_{n,m}| \leq 4^n$, and the choice of $\delta_m$ gives $|\gamma_{m,m}| = 4^m$.
Hence $\left|\frac{f(x+\delta_m) - f(x)}{\delta_m}\right| = \left|\sum_{n=0}^{m} \left(\frac{3}{4}\right)^n \gamma_{n,m}\right| \geq 3^m - \sum_{n=0}^{m-1} 3^n = \frac{1}{2}(3^m + 1)$
Since $\delta_m \to 0$, the difference quotients are unbounded near $x$; since $x$ was arbitrary, $f$ is nowhere differentiable.
*Structural Commentary*: Uniform sums give continuity; scaled cusps force unbounded quotients.
*Key Insight*: Uniform convergence preserves continuity, not differentiability. □




Figure 189: The first compressed sawtooth terms $\sigma_k(x) = (3/4)^k \varphi(4^k x)$: heights shrink, slopes grow.

**Perspective 9.2.12.** *Explicit Versus Generic Pathology*
The sawtooth series proves existence; Baire category shows nowhere differentiability is residual in $C([a, b], \mathbb{R})$.

**Theorem 9.2.13.** *Generic Nowhere Differentiability on Compact Intervals*
*Let $a < b$, and let $C = C([a,b],\mathbb{R})$ with the supremum metric. There exists a residual set $\mathcal{G} \subseteq C$ such that:*

*(i) no $f \in \mathcal{G}$ has an available one-sided derivative at any point of $[a,b]$;*
*(ii) no $f \in \mathcal{G}$ is monotone on any nondegenerate subinterval of $[a,b]$;*
*(iii) the set $\{f \in C : f$ is differentiable at some point of $[a,b]\}$ is first category.*

*Proof.*

(i) Write $\Delta_h f(x) := f(x+h) - f(x)$, and choose $N_0$ with $1/N_0 < b-a$. For $n \geq N_0$, define

$$R_n := \left\{ f \in C : \forall x \in [a, b-1/n]\ \exists h > 0 \text{ with } \left| \frac{\Delta_h f(x)}{h} \right| > n \right\},$$

$$L_n := \left\{ f \in C : \forall x \in [a+1/n, b]\ \exists h < 0 \text{ with } \left| \frac{\Delta_h f(x)}{h} \right| > n \right\},$$

$$G_n := \{ f \in C : f \text{ is nonmonotone on every subinterval of length } 1/n \}.$$

Each $R_n, L_n, G_n$ is open and dense in $C$ (Weierstrass Approximation Theorem 9.2.1), plus small high-slope sawtooth perturbation; openness follows from compactness and uniform perturbation estimates.

(ii) Set $\mathcal{G} := \bigcap_{n \geq N_0} (R_n \cap L_n \cap G_n)$. By Definition 4.3.6, $\mathcal{G}$ is residual. If $f \in \mathcal{G}$, then the $R_n$ and $L_n$ conditions give one-sided difference quotients of arbitrarily large absolute value; since $f$ is bounded, the corresponding increments tend to 0. Thus no available one-sided derivative exists.

(iii) Let $f \in \mathcal{G}$. Every nondegenerate subinterval contains a subinterval of length $1/n$ for all sufficiently large $n$, so the $G_n$ conditions make $f$ nonmonotone on every nondegenerate subinterval. If $f$ is differentiable somewhere, then $f \notin \mathcal{G}$; hence the differentiable-somewhere set lies in $C \setminus \mathcal{G}$, which is first category.

*Structural Commentary*: Baire category turns dense sawtooth perturbability into a generic regularity failure.
*Key Insight*: The explicit construction shows existence; the category proof shows typicality. □

**Method 9.2.14.** *Power-Series and Approximation Proof Patterns*

(i) *Power series*: work on $K_\rho \Subset I_R$; use uniform convergence before termwise limits or derivatives.
(ii) *Endpoint theorems*: prove identities inside the radius and pass to endpoints with Abel-type control.
(iii) *Approximate identities*: use kernels $Q_n \geq 0$, $\int Q_n = 1$, with tails $\to 0$ off 0.
(iv) *Uniform closure*: show $a_n \in \mathcal{A}$ and $a_n \rightrightarrows f$.
(v) *Lattice closure*: derive $|f|$, then $\max, \min$ from algebraic formulas.
(vi) *Stone–Weierstrass*: interpolate locally; use compactness and finite max/min to globalize.

*Key Insight*: Uniform convergence and compactness are what make infinite algebra behave like finite algebra.

## 9.3 The Umbral Calculus

The umbral calculus is the formal linear algebra of polynomial sequences. It identifies a formal power series with three related objects: an algebra element, a linear functional on $\mathbb{C}[x]$, and an operator on $\mathbb{C}[x]$. Refines duality of Definition 4.2.3 and polynomial operator calculus of Definition 1.1.2, without invoking convergence.

**Definition 9.3.1.** *Formal Power Series and Order*
Let $\mathscr{F} := \mathbb{C}[[t]]$.

(i) $f \in \mathscr{F}$ are *formal power series* $f(t) = \sum_{k \geq 0} a_k t^k$, w. coefficientwise addition, Cauchy product multiplication.
(ii) The *order* of $f$ is $o(f) := \min\{k : a_k \neq 0\}$, where $o(0) := +\infty$.
(iii) A series is *invertible* if it has a multiplicative inverse in $\mathscr{F}$, and is a *delta series* if $o(f) = 1$.
(iv) If $o(f) = 1$, its *compositional inverse* is denoted $\overline{f}$ and satisfies $f(\overline{f}(t)) = \overline{f}(f(t)) = t$.
(v) The *lower factorial polynomials* are $x)_0 := 1$ and $(x)_n := x(x-1)\cdots(x-n+1)$, for $n \geq 1$

**Proposition 9.3.2.** *Formal Order Calculus*
*For $f, g \in \mathscr{F}$:*

*(i) $f \in \mathscr{F}^\times \iff o(f) = 0$;*
*(ii) $o(fg) = o(f) + o(g)$ and $o(f+g) \geq \min\{o(f), o(g)\}$;*
*(iii) if $g(t) = \sum_{k \geq 0} b_k t^k$ and $o(f) \geq 1$, then $g \circ f := g(f(t)) := \sum_{k \geq 0} b_k f(t)^k \in \mathscr{F}$;*
*(iv) if $o(f) \geq 1$ and $g \neq 0$, then $o(g \circ f) = o(g) o(f)$;*
*(v) if $o(f) = 1$, then for every $g \in \mathscr{F}$ there are unique $c_k \in \mathbb{C}$ such that $g(t) = \sum_{k \geq 0} c_k f(t)^k$.*

*Proof.*

(i) Let $[t^n] : \mathscr{F} \to \mathbb{C}$ denote coefficient extraction. Then $f \in \mathscr{F}^\times \iff [t^0]f \in \mathbb{C}^\times \iff o(f) = 0$.

(ii) Let $r = o(f)$ and $s = o(g)$. Then $[t^{r+s}]fg = [t^r]f\,[t^s]g \neq 0$, and all lower coefficients vanish; hence $o(fg) = r + s$. Also $[t^m](f+g) = 0$ for $m < \min\{r, s\}$, giving the stated lower bound.
(iii) Since $o(f) \geq 1$, $o(f^k) \to \infty$; hence $\forall n$, coefficient $[t^n] \sum_{k\geq 0} b_k f^k$ has only finitely many nonzero terms.
(iv) Let $s = o(g)$. Then $g \circ f = f^s \left( b_s + \sum_{k>s} b_k f^{k-s} \right)$, where second factor has order 0; hence $o(g \circ f) = s\, o(f)$.
(v) If $o(f) = 1$ and $f(t) = a_1 t + \cdots$ with $a_1 \neq 0$, then $f^k = a_1^k t^k +$ higher terms. Thus $\{t^k\}_{k\geq 0} \mapsto \{f^k\}_{k\geq 0}$ is triangular with nonzero diagonal, so recursive coefficient matching gives existence and uniqueness.

*Structural Commentary*: Delta series behave like a formal change of coordinate at $t = 0$.
*Key Insight*: Order replaces convergence as the bookkeeping device. □

**Definition 9.3.3.** *Derivation*
Let $\mathcal{A}$ be an algebra over a field. Linear operator $\partial : \mathcal{A} \to \mathcal{A}$ is a *derivation* if $\partial(ab) = (\partial a)b + a(\partial b)$ $\forall a, b \in \mathcal{A}$.

**Proposition 9.3.4.** *Formal Differentiation*
*Define* $\partial_t : \mathscr{F} \to \mathscr{F}$ *by* $\partial_t \left( \sum_{k\geq 0} a_k t^k \right) := \sum_{k\geq 1} k a_k t^{k-1}$. *Then:*

*(i)* $\partial_t$ *is* $\mathbb{C}$*-linear.*
*(ii)* $\partial_t(fg) = (\partial_t f)g + f(\partial_t g)$ *for all* $f, g \in \mathscr{F}$*; hence* $\partial_t \in \mathrm{Der}_{\mathbb{C}}(\mathscr{F})$.

*Proof.*
(i) Let $f = \sum a_k t^k$, $g = \sum b_k t^k$, and $\alpha, \beta \in \mathbb{C}$. Then $\partial_t(\alpha f + \beta g) = \sum_{k\geq 1} k(\alpha a_k + \beta b_k) t^{k-1} = \alpha \partial_t f + \beta \partial_t g$.
(ii) Let $f(t) = \sum_{k\geq 0} a_k t^k$ and $g(t) = \sum_{k\geq 0} b_k t^k$. For $n \geq 0$, $[t^n]\partial_t(fg) = (n+1) \sum_{j=0}^{n+1} a_j b_{n+1-j}$.
Also $[t^n]\big((\partial_t f)g + f(\partial_t g)\big) = \sum_{j=1}^{n+1} j a_j b_{n+1-j} + \sum_{j=0}^{n} (n+1-j) a_j b_{n+1-j}$, equals preceding coefficient.

*Structural Commentary*: Formal differentiation is the canonical derivation of $\mathbb{C}[[t]]$.
*Key Insight*: Leibniz is a Cauchy-product coefficient identity. □

**Definition 9.3.5.** *The Umbral Algebra*
(i) *Polynomial algebra*: set $\mathscr{P} := \mathbb{C}[x]$ and let $\mathscr{P}' := \mathrm{Hom}_{\mathbb{C}}(\mathscr{P}, \mathbb{C})$.
(ii) *Umbral pairing*: for $L \in \mathscr{P}'$ and $p(x) \in \mathscr{P}$, write $\langle L \mid p(x) \rangle := L(p(x))$.
(iii) *Exponential normalization*: write each $f \in \mathscr{F}$ as $f(t) = \sum_{k\geq 0} \frac{a_k}{k!} t^k$.
(iv) *Associated functional*: set $\langle f(t) \mid x^n \rangle := a_n$ for $n \geq 0$, extending $\mathbb{C}$-linearly.

**Theorem 9.3.6.** *Formal Series Represent All Algebraic Functionals*
*Define* $\Phi : \mathscr{P}' \to \mathscr{F}$ *by* $\Phi(L) := \sum_{k\geq 0} \frac{\langle L | x^k \rangle}{k!} t^k$. *Then:*

*(i)* $\Phi$ *is a* $\mathbb{C}$*-linear isomorphism with inverse* $\Phi^{-1} \left( \sum_{k\geq 0} \frac{a_k}{k!} t^k \right) \left( \sum_{k=0}^{d} c_k x^k \right) := \sum_{k=0}^{d} c_k a_k$.
*(ii)* $\langle t^k \mid x^n \rangle = n! \delta_{n,k}$ *and* $\langle t^k \mid p \rangle = p^{(k)}(0)$ *for* $p \in \mathscr{P}$*; hence* $t^0 = \mathrm{ev}_0$ *and* $t^k = D^k|_0$.

*Proof.*
(i) Let $T : \mathscr{P}' \to \mathbb{C}^{\mathbb{N}}$ be $T(L) = (a_k)_{k\geq 0}$, $a_k = \langle L \mid x^k \rangle$.
Since $\{x^k\}_{k\geq 0}$ is a Hamel basis, $T$ is a $\mathbb{C}$-linear bijection with $T^{-1}(a) \left( \sum_{k=0}^{d} c_k x^k \right) = \sum_{k=0}^{d} c_k a_k$.
The map $E : \mathbb{C}^{\mathbb{N}} \to \mathscr{F}$, $E(a) = \sum_{k\geq 0} \frac{a_k}{k!} t^k$, is a $\mathbb{C}$-linear bijection, hence $\Phi = E \circ T$ and $\Phi^{-1} = T^{-1} \circ E^{-1}$.
(ii) As $t^k = k!\, \frac{t^k}{k!}$, coefficient sequence is $(n!\delta_{n,k})_{n\geq 0}$. For $p = \sum_{n=0}^{d} c_n x^n$, by linearity, $\langle t^k \mid p \rangle = k! c_k = p^{(k)}(0)$.

*Structural Commentary*: Functionals are monomial-coordinate sequences.
*Key Insight*: Exponential normalization makes $t^k$ equal $D^k|_0$. □

**Definition 9.3.7.** *Delta and Invertible Functionals*
Under the identification $\mathscr{F} \simeq \mathscr{P}'$:
(i) *Delta functional*: a functional represented by $f \in \mathscr{F}$ with $o(f) = 1$.
(ii) *Invertible functional*: a functional represented by $f \in \mathscr{F}^\times$, equivalently $o(f) = 0$.

**Proposition 9.3.8.** *Basic Umbral Identities*
*For $f, g \in \mathscr{F}$, $p \in \mathscr{P}$, and $n \geq 0$:*

*(i)* $f(t) = \sum_{k\geq 0} \frac{\langle f(t) \mid x^k \rangle}{k!} t^k$;

*(ii)* $p(x) = \sum_{k=0}^{\deg p} \frac{\langle t^k \mid p \rangle}{k!} x^k$;

*(iii)* $\langle fg \mid x^n \rangle = \sum_{k=0}^{n} \binom{n}{k} \langle f \mid x^k \rangle \langle g \mid x^{n-k} \rangle$;

*(iv) if* $o(f) > \deg p$, *then* $\langle f \mid p \rangle = 0$;

*(v) if* $o(f_k) = k$ *for all* $k \geq 0$, *then* $\left\langle \sum_{k\geq 0} a_k f_k \middle| p \right\rangle = \sum_{k=0}^{\deg p} a_k \langle f_k \mid p \rangle$, *with* $\sum_{k\geq 0} a_k f_k$ *well-defined in* $\mathscr{F}$.

*Proof.*

(i) If $f(t) = \sum_{k\geq 0} a_k t^k/k!$, then $a_k = \langle f \mid x^k \rangle$ by definition.

(ii) Since $\langle t^k \mid p \rangle = p^{(k)}(0)$, this is Taylor's formula for the polynomial $p$.

(iii) Write $f = \sum a_k t^k/k!$ and $g = \sum b_k t^k/k!$.
The Cauchy product is $fg = \sum_{n\geq 0} \left( \sum_{k=0}^{n} \binom{n}{k} a_k b_{n-k} \right) \frac{t^n}{n!}$, and $a_k = \langle f \mid x^k \rangle$, $b_k = \langle g \mid x^k \rangle$.

(iv) If $p = \sum_{j=0}^{d} c_j x^j$ and $o(f) > d$, then $\langle f \mid x^j \rangle = 0$ for $0 \leq j \leq d$.

(v) Since $o(f_k) = k$, each coefficient of $\sum_{k\geq 0} a_k f_k$ has only finitely many contributions.
With $d = \deg p$, (iv) gives $\langle f_k \mid p \rangle = 0$ for $k > d$, proving the finite formula.

*Structural Commentary*: The umbral pairing dualizes coefficient calculus.
*Key Insight*: Order makes infinite formal sums act finitely on each polynomial. □

**Theorem 9.3.9.** *Multiplication by $x$ Moves to Formal Differentiation*
*Let $M_x : \mathscr{P} \to \mathscr{P}$ be $M_x p = xp$, and let $\partial_t : \mathscr{F} \to \mathscr{F}$ be formal differentiation. Then:*

*(i)* $\langle f \mid M_x p \rangle = \langle \partial_t f \mid p \rangle$ *for all* $f \in \mathscr{F}$ *and* $p \in \mathscr{P}$.

*(ii)* $M_x^* = \partial_t$ *under the umbral pairing.*

*Proof.*

(i) By linearity in $p$, take $p = x^n$.
For $f = \sum_{k\geq 0} \frac{a_k}{k!} t^k$, $\partial_t f = \sum_{k\geq 0} \frac{a_{k+1}}{k!} t^k$, hence $\langle \partial_t f \mid x^n \rangle = a_{n+1} = \langle f \mid x^{n+1} \rangle = \langle f \mid M_x x^n \rangle$.

(ii) For $f \in \mathscr{F}$, the adjoint satisfies $\langle M_x^* f \mid p \rangle = \langle f \mid M_x p \rangle$ for all $p \in \mathscr{P}$.
By (i), $\langle M_x^* f \mid p \rangle = \langle \partial_t f \mid p \rangle$ for all $p$, so nondegeneracy gives $M_x^* f = \partial_t f$. Hence $M_x^* = \partial_t$.

*Structural Commentary*: $M_x$ and $\partial_t$ are adjoint under the umbral pairing.
*Key Insight*: Exponential normalization makes multiplication by $x$ a coefficient shift. □

**Definition 9.3.10.** *Basic Umbral Functionals*
For $a \in \mathbb{C}$ and $p \in \mathscr{P}$:

(i) *Evaluation*: $\epsilon_a = e^{at}$ and $\langle e^{at} \mid p \rangle = p(a)$; hence $\epsilon_a \epsilon_b = \epsilon_{a+b}$.
(ii) *Forward difference*: $\Delta_a = e^{at} - 1$ and $\langle \Delta_a \mid p \rangle = p(a) - p(0)$.
(iii) *Abel functional*: $te^{at}$ and $\langle te^{at} \mid p \rangle = p'(a)$.
(iv) *Gamma functional*: $(1-t)^{-1}$ and $\left\langle (1-t)^{-1} \middle| p \right\rangle = \int_0^\infty p(u) e^{-u}\, du$.
(v) *Integral functional*: $\left\langle \frac{e^{at}-1}{t} \middle| p \right\rangle = \int_0^a p(u)\, du$
(vi) *Bernoulli numbers*: for $a = 1$, $B_n = \left\langle \frac{t}{e^t - 1} \middle| x^n \right\rangle$.

**Definition 9.3.11.** *Formal Series as Polynomial Operators*
For $f(t) = \sum_{k\geq 0} a_k t^k/k! \in \mathscr{F}$:

(i) *Operator action*: define $f(t) \in \mathrm{End}_{\mathbb{C}}(\mathscr{P})$ by $f(t)p(x) := \sum_{k\geq 0} \frac{a_k}{k!} p^{(k)}(x)$, a finite sum for each $p \in \mathscr{P}$.

(ii) *Monomial action*: for $n \geq 0$, $f(t)x^n = \sum_{k=0}^{n} \binom{n}{k} a_k x^{n-k}$.

**Theorem 9.3.12.** *Functional-Operator Compatibility*
*For $f, g \in \mathscr{F}$ and $p \in \mathscr{P}$:*

*(i)* $\langle fg \mid p\rangle = \langle f \mid g(t)p\rangle$.
*(ii)* $\langle f \mid p\rangle = \langle t^0 \mid f(t)p\rangle$.

*Proof.*

(i) By linearity in $p$, take $p = x^n$. Write $f = \sum_i \frac{a_i}{i!}t^i$ and $g = \sum_j \frac{b_j}{j!}t^j$.

Then $\langle fg \mid x^n\rangle = \sum_{j=0}^{n} \binom{n}{j} a_{n-j}b_j = \left\langle f \middle| \sum_{j=0}^{n} \binom{n}{j} b_j x^{n-j} \right\rangle = \langle f \mid g(t)x^n\rangle$.

(ii) In (i), set $(f, g) = (t^0, F)$ with $F \in \mathscr{F}$. Since $t^0 F = F$ in $\mathscr{F}$, $\langle F \mid p\rangle = \langle t^0 F \mid p\rangle = \langle t^0 \mid F(t)p\rangle$.

*Structural Commentary*: Multiplication is adjoint to action.
*Key Insight*: One finite coefficient sum appears on both sides. □

**Definition 9.3.13.** *Basic Umbral Operators*
For $a \in \mathbb{C}$ and $p \in \mathscr{P}$:

(i) *Translation*: $e^{at}p(x) = p(x + a)$.
(ii) *Forward difference*: $(e^{at} - 1)p(x) = p(x + a) - p(x)$.
(iii) *Abel operator*: $te^{at}p(x) = p'(x + a)$.
(iv) *Gamma operator*: $(1 - t)^{-1}p(x) = \int_0^\infty p(x + u)e^{-u}\,du$
(v) *Integral operator*: $\frac{e^{at}-1}{t}p(x) = \int_x^{x+a} p(u)\,du$

**Theorem 9.3.14.** *Shift-Invariant Operators*
*For $T \in \mathrm{End}_{\mathbb{C}}(\mathscr{P})$, the following are equivalent:*

*(i)* $T = f(t)$ *for some* $f \in \mathscr{F}$;
*(ii)* $[T, t] = 0$;
*(iii)* $[T, h(t)] = 0$ *for every delta series* $h$;
*(iv)* $[T, e^{at}] = 0$ *for all* $a \in \mathbb{C}$.

*Proof.*

*(i)* $\Rightarrow$ *(ii), (iii), (iv)* If $T = f(t)$, then $\forall u(t) \in \mathscr{F}$, $[T, u(t)] = [f(t), u(t)] = 0$, since $\mathscr{F}$ is commutative.
*(iii)* $\Rightarrow$ *(ii)* Since $o(t) = 1$, $(\forall h \in \mathscr{F},\ o(h) = 1 \Rightarrow [T, h(t)] = 0) \Rightarrow [T, t] = 0$.
*(iv)* $\Rightarrow$ *(ii)* For each $p \in \mathscr{P}$, $0 = [a^1]\,[T, e^{at}]p = [T, t]p$, so $[T, t] = 0$.
*(ii)* $\Rightarrow$ *(i)* Set $g(t) := \sum_{k\geq 0} \frac{\langle t^0 | Tx^k\rangle}{k!}t^k$. Let $a_k := \langle t^0 \mid Tx^k\rangle$.

Since $[T, t] = 0$, for $m, k \geq 0$, $\langle t^m \mid Tx^k\rangle = \langle t^0 \mid t^m Tx^k\rangle = \langle t^0 \mid Tt^m x^k\rangle = (k)_m a_{k-m}$, with $a_{k-m} = 0$ if $m > k$. Also $g(t) = \sum_{r\geq 0} \frac{a_r}{r!}t^r$ and $t^r x^k = (k)_r x^{k-r}$ give $\langle t^m \mid g(t)x^k\rangle = \sum_{r=0}^{k} \binom{k}{r} a_r \langle t^m \mid x^{k-r}\rangle = (k)_m a_{k-m}$.

Thus $\langle t^m \mid Tx^k - g(t)x^k\rangle = 0$ for all $m, k$. Since $\{t^m : m \geq 0\}$ separates $\mathscr{P}$, $T = g(t)$.

*Structural Commentary*: The commutant of $t$ on $\mathscr{P}$ is $\mathscr{F}$.
*Key Insight*: The derivative is enough to recover every coefficient of the operator. □

**Definition 9.3.15.** *Sheffer, Associated, and Appell Sequences*
Let $f$ be a delta series, $g$ an invertible series, and $\deg s_n = n$.

(i) *Sheffer sequence*: $(s_n)_{n\geq 0}$ is Sheffer for $(g, f)$ if $\langle g(t)f(t)^k \mid s_n(x)\rangle = n!\delta_{n,k}$ for $n, k \geq 0$.
(ii) *Associated sequence*: a Sheffer sequence for $(1, f)$.
(iii) *Appell sequence*: a Sheffer sequence for $(g, t)$.

**Theorem 9.3.16.** *Existence and Expansion Theorems*
*Let $f$ be a delta series, $g$ an invertible series, and $q_k := g(t)f(t)^k$.*

*(i) There is a unique sequence $(s_n)_{n\geq 0}$ with $\deg s_n = n$ and $\langle q_k \mid s_n\rangle = n!\delta_{n,k}$.*
*(ii) If $p_n := g(t)s_n$, then $(s_n)$ is Sheffer for $(g, f)$ iff $(p_n)$ is associated with $f$.*
*(iii) For all $h \in \mathscr{F}$, $h = \sum_{k\geq 0} \frac{\langle h | s_k\rangle}{k!}\, q_k$.*
*(iv) For all $p \in \mathscr{P}$, $p = \sum_{k=0}^{\deg p} \frac{\langle q_k | p\rangle}{k!}\, s_k$.*

*Proof.*

(i) Since $o(q_k) = k$, the matrix $A_n = (\langle q_i \mid x^j \rangle)_{0 \le i,j \le n}$ is triangular with nonzero diagonal. Hence the equations $\langle q_k \mid s_n \rangle = n! \delta_{n,k}$ have a unique solution $s_n \in \mathscr{P}_{\le n}$, necessarily with $\deg s_n = n$.

(ii) By Theorem 9.3.12, $\langle f(t)^k \mid p_n \rangle = \langle f(t)^k \mid g(t) s_n \rangle = \langle q_k \mid s_n \rangle$.

(iii) Since $o(q_k) = k$, every $h \in \mathscr{F}$ has a unique expansion $h = \sum c_k q_k$. Pairing with $s_n$ gives $\langle h \mid s_n \rangle = c_n n!$, so $c_n = \langle h \mid s_n \rangle / n!$.

(iv) Let $d = \deg p$ and $r := p - \sum_{k=0}^{d} \langle q_k \mid p \rangle s_k / k!$.

Then $\langle q_j \mid r \rangle = 0$ for all $j \le d$, and also for $j > d$ by $o(q_j) > d$. Since the $q_j$ separate $\mathscr{P}$, $r = 0$.

*Structural Commentary*: $(s_n)$ is dual to $q_k = g f^k$.
*Key Insight*: The order filtration makes the orthogonality equations triangular. □

**Theorem 9.3.17.** *Generating and Operator Characterizations*
*Let $f$ be a delta series with compositional inverse $\overline{f}$, let $g \in \mathscr{F}^\times$, and let $(p_n), (s_n)$ be polynomial sequences with $\deg p_n = \deg s_n = n$. Set $P(y,u) := \sum_{n \ge 0} \frac{p_n(y)}{n!} u^n$ and $S(y,u) := \sum_{n \ge 0} \frac{s_n(y)}{n!} u^n$.*

*(i) $(p_n)$ is associated with $f$ iff $P(y,u) = e^{y\overline{f}(u)}$*
*(ii) $(s_n)$ is Sheffer for $(g,f)$ iff $S(y,u) = g(\overline{f}(u))^{-1} e^{y\overline{f}(u)}$*
*(iii) $(p_n)$ is associated with $f$ iff $p_n(0) = \delta_{n,0}$, $f(t) p_n = n p_{n-1}$, where $p_{-1} := 0$.*
*(iv) $(s_n)$ is Sheffer for $(\gamma, f)$ for some $\gamma \in \mathscr{F}^\times$ iff $f(t) s_n = n s_{n-1}$, where $s_{-1} := 0$.*
*(v) If $(p_n)$ is associated with $f$, then $p_n(x) = \sum_{k=0}^{n} \frac{\langle \overline{f}(t)^k \mid x^n \rangle}{k!} x^k$*
*(vi) If $(s_n)$ is Sheffer for $(g,f)$, then $s_n(x) = \sum_{k=0}^{n} \frac{1}{k!} \langle g(\overline{f}(t))^{-1} \overline{f}(t)^k \mid x^n \rangle x^k$.*

*Proof.*

(i) ($\Rightarrow$) By Theorem 9.3.16, $e^{yt} = \sum_{n \ge 0} \frac{p_n(y)}{n!} f(t)^n$. Substitute $t = \overline{f}(u)$ to obtain $P(y,u) = e^{y\overline{f}(u)}$.

(i) ($\Leftarrow$) If $P(x,u) = e^{x\overline{f}(u)}$, then for all $k \ge 0$, $\sum_{n \ge 0} \frac{\langle f(t)^k \mid p_n \rangle}{n!} u^n = \langle f(t)^k \mid P(x,u) \rangle = f(\overline{f}(u))^k = u^k$.

Thus $\langle f(t)^k \mid p_n \rangle = n! \delta_{n,k}$.

(ii) ($\Rightarrow$) By Theorem 9.3.16, $e^{yt} = \sum_{n \ge 0} \frac{s_n(y)}{n!} g(t) f(t)^n$. Substitute $t = \overline{f}(u)$ and divide by $g(\overline{f}(u))$.

(ii) ($\Leftarrow$) If $S(x,u) = g(\overline{f}(u))^{-1} e^{x\overline{f}(u)}$, then $\sum_{n \ge 0} \frac{\langle g(t) f(t)^k \mid s_n \rangle}{n!} u^n = \langle g(t) f(t)^k \mid S(x,u) \rangle = u^k$,

so $\langle g(t) f(t)^k \mid s_n \rangle = n! \delta_{n,k}$.

(iii) ($\Rightarrow$) If $(p_n)$ is associated, then $p_n(0) = \langle 1 \mid p_n \rangle = \delta_{n,0}$ and, for all $k$, $\langle f(t)^k \mid f(t) p_n - n p_{n-1} \rangle = 0$. Since $\{f(t)^k : k \ge 0\}$ separates $\mathscr{P}$, $f(t) p_n = n p_{n-1}$.

(iii) ($\Leftarrow$) The lowering relation gives $f(t)^k p_n = (n)_k p_{n-k}$. Hence $\langle f(t)^k \mid p_n \rangle = \langle 1 \mid f(t)^k p_n \rangle = (n)_k p_{n-k}(0) = n! \delta_{n,k}$.

(iv) ($\Rightarrow$) If $(s_n)$ is Sheffer for $(\gamma, f)$, then for all $k$, $\langle \gamma(t) f(t)^k \mid f(t) s_n - n s_{n-1} \rangle = 0$. Since $\{\gamma(t) f(t)^k : k \ge 0\}$ separates $\mathscr{P}$, $f(t) s_n = n s_{n-1}$.

(iv) ($\Leftarrow$) Define $L \in \mathscr{P}'$ by $L(s_0) = 1$ and $L(s_n) = 0$ for $n > 0$, and put $\gamma := \Phi(L)$. Then $\langle \gamma \mid 1 \rangle = L(1) \neq 0$, so $\gamma \in \mathscr{F}^\times$, and $\langle \gamma(t) f(t)^k \mid s_n \rangle = \langle \gamma(t) \mid f(t)^k s_n \rangle = (n)_k L(s_{n-k}) = n! \delta_{n,k}$.

(v) From (i), $e^{x\overline{f}(u)} = \sum_{k \ge 0} \frac{\overline{f}(u)^k}{k!} x^k$. Applying $n![u^n]$ gives the stated formula for $p_n$.

(vi) From (ii), $g(\overline{f}(u))^{-1} e^{x\overline{f}(u)} = \sum_{k \ge 0} \frac{g(\overline{f}(u))^{-1} \overline{f}(u)^k}{k!} x^k$. Applying $n![u^n]$ gives the stated formula for $s_n$.

*Structural Commentary*: $\overline{f}$ converts $f$-lowering to exponential generating functions.
*Key Insight*: Generating functions package the whole triangular duality into one formal identity. □

**Definition 9.3.18.** *Binomial Type and Sheffer Identity*
Let $(p_n)$ and $(s_n)$ be polynomial sequences.

(i) *Binomial type*: $(p_n)$ is of binomial type if $p_n(x+y) = \sum_{k=0}^{n} \binom{n}{k} p_k(y) p_{n-k}(x)$.

(ii) *Sheffer identity*: $(s_n)$ satisfies the Sheffer identity relative to $(p_n)$ if $s_n(x+y) = \sum_{k=0}^{n} \binom{n}{k} p_k(y) s_{n-k}(x)$.

**Theorem 9.3.19.** *Action and Binomial Identities*
*Let $(s_n)$ be Sheffer for $(g, f)$, let $(p_n)$ be associated with $f$, and write $E_y := e^{yt}$. Then:*

*(i) for $h \in \mathscr{F}$ and $n \geq 0$, $h(t)s_n = \sum_{k=0}^{n} \binom{n}{k} \langle h \mid s_k \rangle p_{n-k}$;*
*(ii) $(p_n)$ is associated with a delta series iff $(p_n)$ is of binomial type;*
*(iii) $(s_n)$ is Sheffer for some $g \in \mathscr{F}^\times$ and same $f$ iff $(s_n)$ satisfies the Sheffer identity relative to $(p_n)$.*

*Proof.*

(i) By Theorem 9.3.16, $h = \sum_{k \geq 0} \frac{\langle h | s_k \rangle}{k!} g(t) f(t)^k$.
Since $g(t)s_n = p_n$ and $f(t)^k p_n = (n)_k p_{n-k}$, $h(t)s_n = \sum_{k=0}^{n} \frac{\langle h | s_k \rangle}{k!} (n)_k p_{n-k}$.

*(ii)* (⇒) If $(p_n)$ is associated, apply (i) with $s_n = p_n$, $g = 1$, and $h = E_y$.
Since $\langle E_y \mid p_k \rangle = p_k(y)$ and $E_y p_n(x) = p_n(x + y)$, $(p_n)$ is of binomial type.

*(ii)* (⇐) If $(p_n)$ is of binomial type, then $p_n(0) = \delta_{n,0}$. Define $Qp_n := np_{n-1}$. The binomial identity gives $QE_y p_n = E_y Q p_n$ for all $n, y$, so $[Q, E_y] = 0$. By Theorem 9.3.14, $Q = f(t)$ for some $f \in \mathscr{F}$.
Since $Q$ lowers degree by one, $o(f) = 1$; Theorem 9.3.17 makes $(p_n)$ associated with $f$.

*(iii)* (⇒) If $(s_n)$ is Sheffer for $(g, f)$, apply (i) with $h = E_y$.
Since $\langle E_y \mid s_k \rangle = s_k(y)$ and $E_y s_n(x) = s_n(x + y)$, $s_n(x + y) = \sum_{k=0}^{n} \binom{n}{k} s_k(y) p_{n-k}(x)$.
Relabel and swap $x, y$ to get the Sheffer identity relative to $(p_n)$.

*(iii)* (⇐) If the Sheffer identity holds, apply $f(t)$ in the $y$-variable and set $y = 0$.
Since $f(t)p_k(y)|_{y=0} = kp_{k-1}(0) = k\delta_{k,1}$, $f(t)s_n = ns_{n-1}$.
By Theorem 9.3.17, $(s_n)$ is Sheffer for $(g, f)$ for some $g \in \mathscr{F}^\times$.

*Structural Commentary*: Translation turns the operator formula into addition laws for polynomial sequences.
*Key Insight*: Binomial identities are the visible polynomial shadow of a delta operator. □

**Example 9.3.20.** *Standard Sheffer Examples*

(i) *Monomials*: $p_n(x) = x^n$ is associated with $f(t) = t$; the binomial identity is the usual binomial theorem.
(ii) *Lower factorials*: $(x)_n$ is associated with $f(t) = e^t - 1$, with $(1 + t)^y = e^{y \log(1+t)} = \sum_{k \geq 0} \frac{(y)_k}{k!} t^k$,
and Vandermonde identity $(x + y)_n = \sum_{k=0}^{n} \binom{n}{k} (x)_k (y)_{n-k}$.
(iii) *Abel polynomials*: associated with $f(t) = te^{at}$, $A_n(x; a) = x(x - an)^{n-1}$.
(iv) *Hermite polynomials*: for $g(t) = e^{t^2/2}$, the polynomial is $H_n(x) = e^{-t^2/2} x^n$, where $e^{yt - t^2/2} = \sum_{k \geq 0} \frac{H_k(y)}{k!} t^k$.
(v) *Laguerre polynomials*: order $\alpha$, Sheffer for $g(t) = (1-t)^{-\alpha-1}$, $f(t) = \frac{t}{t-1}$, $\frac{1}{(1-t)^{\alpha+1}} e^{yt/(t-1)} = \sum_{k \geq 0} \frac{L_k^{(\alpha)}(y)}{k!} t^k$.

**Definition 9.3.21.** *Umbral Operators, Continuity, and Adjoints*
Let $(s_n)$ be Sheffer for $(g, f)$, and let $(p_n)$ be associated with $f$.

(i) *Sheffer operator*: $\lambda_{g,f} \in \mathcal{L}(\mathscr{P})$ is defined by $\lambda_{g,f}(x^n) = s_n(x)$.
(ii) *Umbral operator*: $\lambda_f \in \mathcal{L}(\mathscr{P})$ is defined by $\lambda_f(x^n) = p_n(x)$.
(iii) *Sheffer/umbral shift*: $\theta_{g,f} s_n = s_{n+1}$; in the associated case, $\theta_f p_n = p_{n+1}$.
(iv) *Continuous operator*: $T \in \mathcal{L}(\mathscr{F})$ is continuous iff $o(f_k) \to \infty \Longrightarrow o(Tf_k) \to \infty$.
Equivalently, for every defined formal sum with $o(f_k) \to \infty$, $T\left(\sum_{k \geq 0} a_k f_k\right) = \sum_{k \geq 0} a_k T(f_k)$
(v) *Algebraic adjoint*: for $\tau \in \mathcal{L}(\mathscr{P})$, $\tau^\times \in \mathcal{L}(\mathscr{F})$ is defined by $\langle \tau^\times h \mid p \rangle = \langle h \mid \tau p \rangle$ for $h \in \mathscr{F}$, $p \in \mathscr{P}$.
This is the dual-map adjoint of Definition 4.2.3, not an inner-product adjoint.

**Theorem 9.3.22.** *Continuous Operators Are Exactly Operator Adjoints*
*Let $\mathcal{L}_{\text{cont}}(\mathscr{F})$ denote the continuous linear operators on $\mathscr{F}$.*
*Then $(-)^\times : \mathcal{L}(\mathscr{P}) \xrightarrow{\sim} \mathcal{L}_{\text{cont}}(\mathscr{F})$, $\tau \longmapsto \tau^\times$ is a vector-space isomorphism.*

*Proof.*

(⇒) Let $\tau \in \mathcal{L}(\mathscr{P})$ and $o(h_j) \to \infty$. For fixed $n$, $\tau x^n \in \mathscr{P}$, hence $\langle \tau^\times h_j \mid x^n \rangle = \langle h_j \mid \tau x^n \rangle = 0$ for all sufficiently large $j$. Thus $o(\tau^\times h_j) \to \infty$, so $\tau^\times \in \mathcal{L}_{\text{cont}}(\mathscr{F})$.
If $\tau^\times = 0$, then $\langle h \mid \tau p \rangle = 0$ for all $h, p$, hence $\tau = 0$; the map is injective.

($\Leftarrow$) Let $T \in \mathcal{L}_{\text{cont}}(\mathscr{F})$. Define $\tau_T x^n := \sum_{k\geq 0} \frac{\langle Tt^k | x^n\rangle}{k!} x^k$. Since $o(Tt^k) \to \infty$, the sum is finite for each $n$.

For all $m, n$, $\langle \tau_T^\times t^m \mid x^n\rangle = \langle t^m \mid \tau_T x^n\rangle = \langle Tt^m \mid x^n\rangle$. Thus $\tau_T^\times t^m = Tt^m$ for every $m$.

By continuity, $\tau_T^\times = T$, so the map is surjective.

*Structural Commentary*: Continuity is the finiteness condition for dualising polynomial operators.
*Key Insight*: Finiteness on each polynomial degree is the hidden topology. □




Figure 190: Adjoints identify $\mathcal{L}(\mathscr{P})$ with $\mathcal{L}_{\text{cont}}(\mathscr{F})$ and map the umbral classes shown.

**Theorem 9.3.23.** *Umbral Operators and Adjoint Automorphisms*
*Let $f$ be a delta series, let $\overline{f}$ be its compositional inverse, let $(p_n)$ be associated with $f$, and set $\lambda_f(x^n) = p_n(x)$. For $C_{\overline{f}}(h) := h(\overline{f})$ and $\lambda \in \mathcal{L}(\mathscr{P})$:*

*(i) $\lambda = \lambda_f$ iff $\lambda^\times = C_{\overline{f}}$; in this case $\lambda_f^\times f = t$.*
*(ii) $\lambda$ is umbral iff $\lambda^\times \in \text{Aut}_{\mathbb{C}\text{-}alg}(\mathscr{F})$.*

*Proof.*

(i) ($\Rightarrow$) If $\lambda = \lambda_f$, then $\langle \lambda^\times f^k \mid x^n\rangle = \langle f^k \mid p_n\rangle = n!\delta_{n,k} = \langle t^k \mid x^n\rangle$, so $\lambda^\times f^k = t^k$. For $h \in \mathscr{F}$, put $q := h(\overline{f})$, so $h = q(f)$. Continuity gives $\lambda^\times h = q(t) = h(\overline{f}) = C_{\overline{f}} h$. Hence $\lambda^\times = C_{\overline{f}}$ and $\lambda_f^\times f = t$.

(i) ($\Leftarrow$) If $\lambda^\times = C_{\overline{f}}$, set $r_n := \lambda x^n$. Then $\langle f^k \mid r_n\rangle = \langle \lambda^\times f^k \mid x^n\rangle = \langle t^k \mid x^n\rangle = n!\delta_{n,k}$.
Thus $(r_n)$ is associated with $f$, so $r_n = p_n$ and $\lambda = \lambda_f$.

(ii) ($\Rightarrow$) If $\lambda$ is umbral, then $\lambda = \lambda_u$ for some delta series $u$. By (i), $\lambda^\times = C_{\overline{u}} \in \text{Aut}_{\mathbb{C}\text{-alg}}(\mathscr{F})$.

(ii) ($\Leftarrow$) Let $A := \lambda^\times \in \text{Aut}_{\mathbb{C}\text{-alg}}(\mathscr{F})$ and set $u := A^{-1}t$. Since $A^{-1}((t)) = (t)$ and $A^{-1}((t)^2) = (t)^2$, $u \in (t) \setminus (t)^2$, so $o(u) = 1$. With $r_n := \lambda x^n$, $\langle u^k \mid r_n\rangle = \langle Au^k \mid x^n\rangle = \langle t^k \mid x^n\rangle = n!\delta_{n,k}$.
Hence $(r_n)$ is associated with $u$, so $\lambda = \lambda_u$ is umbral.

*Structural Commentary*: Umbral adjoints are substitutions.
*Key Insight*: The delta series is $A^{-1}t$. □

**Definition 9.3.24.** *Umbral Composition*

For polynomial sequences $(p_n)$ and $(q_n)$ with $p_n(x) = \sum_{k=0}^n p_{n,k} x^k$, define $(p \circ_U q)_n(x) := \sum_{k=0}^n p_{n,k} q_k(x)$.

**Theorem 9.3.25.** *Group Law for Umbral Operators*
*Let $f, g$ be delta series, and let $(p_n)$ and $(q_n)$ be associated with $f$ and $g$. Then:*

*(i) $\lambda_g \lambda_f = \lambda_{f\circ g}$.*
*(ii) $\lambda_f^{-1} = \lambda_{\overline{f}}$.*
*(iii) $(p \circ_U q)_n = (\lambda_g \lambda_f) x^n$; hence associated sequences form a group under $\circ_U$.*

*Proof.*

(i) By $(AB)^\times = B^\times A^\times$ and Theorem 9.3.23, for $h \in \mathscr{F}$, $(\lambda_g\lambda_f)^\times h = \lambda_f^\times \lambda_g^\times h = h(\overline{g}(\overline{f})) = h(\overline{f \circ g})$. QED.

(ii) Put $g = \overline{f}$ in (i). Then $\lambda_{\overline{f}}\lambda_f = \lambda_t = I$ and $\lambda_f \lambda_{\overline{f}} = \lambda_t = I$, so $\lambda_f^{-1} = \lambda_{\overline{f}}$.

(iii) If $p_n = \sum_{k=0}^n p_{n,k} x^k$, then $(\lambda_g\lambda_f)x^n = \lambda_g p_n = \sum_{k=0}^n p_{n,k} q_k = (p \circ_U q)_n$.
By (i) and (ii), the operation inherits associativity, identity, and inverses from operator composition.

*Structural Commentary*: Umbral composition is substitution of delta series.
*Key Insight*: Adjoints reverse the operator composition. □

**Proposition 9.3.26.** *Sheffer Operators*
*Let $f, k$ be delta series, $g, h \in \mathscr{F}^\times$, $(s_n)$ Sheffer for $(g, f)$, and $(p_n)$ associated with $f$. Then:*

*(i) $s_n = g(t)^{-1} p_n$ and $\lambda_{g,f} = g(t)^{-1}\lambda_f$.*
*(ii) $\lambda_{g,f}\lambda_{h,k} = \lambda_{g(h\circ f),\, k\circ f}$.*

*(iii)* $\lambda_{g,f}^{-1} = \lambda_{g^{-1}(\overline{f}),\,\overline{f}}$.

*Proof.*

(i) Theorem 9.3.16 gives $g(t)s_n = p_n$, hence $s_n = g(t)^{-1}p_n$ and $\lambda_{g,f} = g(t)^{-1}\lambda_f$.
(ii) For $a \in \mathscr{F}$, Theorem 9.3.23 gives $\lambda_f a(t) = a(f(t))\lambda_f$.
Thus $\lambda_{g,f}\lambda_{h,k} = g(t)^{-1}\lambda_f h(t)^{-1}\lambda_k = g(t)^{-1}h(f(t))^{-1}\lambda_{k\circ f} = \lambda_{g(h\circ f),\,k\circ f}$.
(iii) In (ii), take $(h,k) = (g^{-1}(\overline{f}),\overline{f})$ to get $\lambda_{g,f}\lambda_{g^{-1}(\overline{f}),\,\overline{f}} = \lambda_{1,t} = I$.
Reversing the factors gives $\lambda_{g^{-1}(\overline{f}),\,\overline{f}}\lambda_{g,f} = \lambda_{g^{-1}(\overline{f})(g\circ\overline{f}),\,f\circ\overline{f}} = \lambda_{1,t} = I$.

*Structural Commentary*: Weight then umbral substitution.
*Key Insight*: $(g,f)(h,k) = (g(h\circ f), k\circ f)$. □

**Theorem 9.3.27.** *Umbral Shift Derivations*
*Let $f$ be a delta series and let $\theta_f$ be its umbral shift.*
*Define the continuous derivation $\partial_f : \mathscr{F} \to \mathscr{F}$ by $\partial_f(f^k) = kf^{k-1}$, $k \geq 0$. Then:*

*(i)* $\theta_f^{\times} = \partial_f$.
*(ii)* $\partial_f f = 1$.
*(iii) If $g$ is a delta series, then* $\partial_g = (\partial_g f)\partial_f$.

*Proof.*

(i) Let $(p_n)$ be associated with $f$.
For $k,n \geq 0$, $\langle \theta_f^{\times} f^k \mid p_n\rangle = \langle f^k \mid p_{n+1}\rangle = (n+1)!\delta_{n+1,k} = \langle kf^{k-1} \mid p_n\rangle$.
Hence $\theta_f^{\times} f^k = kf^{k-1}$ for all $k$, so $\theta_f^{\times} = \partial_f$ by continuity.
(ii) By definition of $\partial_f$, $\partial_f f = \partial_f(f^1) = 1\cdot f^0 = 1$.
(iii) It suffices by continuity to test on $f^k$: $(\partial_g f)\partial_f(f^k) = k(\partial_g f)f^{k-1} = \partial_g(f^k)$.

*Structural Commentary*: Umbral shift is delta-coordinate differentiation.
*Key Insight*: The associated basis turns shifts into derivatives. □

**Theorem 9.3.28.** *Umbral Shifts and Surjective Derivations*
*For $\theta \in \mathcal{L}(\mathscr{P})$, $\theta$ is an umbral shift $\iff$ $\theta^{\times}$ is a surjective derivation of $\mathscr{F}$.*

*Proof.*

($\Rightarrow$) If $\theta = \theta_f$, then $\theta^{\times} = \partial_f$ by Theorem 9.3.27. Since $\partial_f\left(\frac{f^{k+1}}{k+1}\right) = f^k$, $k \geq 0$, $\partial_f$ is a surjective derivation.
($\Leftarrow$) Let $D := \theta^{\times}$ be a surjective derivation. Choose $u \in \mathscr{F}$ with $Du = 1$ and set $f := u - [t^0]u$.
Then $Df = 1$ and, since $D$ lowers order by one, $o(f) = 1$. Let $(p_n)$ be associated with $f$.
For $k,n \geq 0$, $\langle f^k \mid \theta p_n\rangle = \langle Df^k \mid p_n\rangle = \langle kf^{k-1} \mid p_n\rangle = \langle f^k \mid p_{n+1}\rangle$.
The functionals $f^k$ separate $\mathscr{P}$, so $\theta p_n = p_{n+1}$ and $\theta = \theta_f$.

*Structural Commentary*: Surjective derivations are derivatives in delta coordinates.
*Key Insight*: Surjectivity chooses the coordinate whose derivative is 1. □

## 9.4 Umbral Recurrences and Variants

**Theorem 9.4.1.** *Recurrence and Transfer Formulas*
*Let $(p_n)$ be associated with the delta series $f$, and set $u(t) := f(t)/t \in \mathscr{F}^{\times}$. Then:*

*(i)* $p_{n+1} = x[f'(t)]^{-1}p_n$ *for* $n \geq 0$.
*(ii)* $p_n = f'(t)u(t)^{-n-1}x^n = xu(t)^{-n}x^{n-1}$ *for* $n \geq 1$.

*Proof.*

(i) By Theorem 9.3.27, $\theta_t = (\partial_t f)\theta_f = f'(t)\theta_f$. Since $\theta_t = x$ and $\theta_f p_n = p_{n+1}$, $p_{n+1} = x[f'(t)]^{-1}p_n$.
(ii) Put $q_n := f'u^{-n-1}x^n$. Then $q_0 = f'u^{-1}1 = 1 = p_0$ and, using $xa(t) = a(t)x - a'(t)$,
$q_{n+1} = f'u^{-n-2}x^{n+1} = xu^{-n-1}x^n = x(f')^{-1}q_n$. Thus $q_n = p_n$ by (i).
The same commutator gives $xu^{-n}x^{n-1} = u^{-n}x^n + nu^{-n-1}u'x^{n-1} = f'u^{-n-1}x^n$.

*Structural Commentary*: The recurrence determines $(p_n)$; transfer solves it using $f = tu$.
*Key Insight*: The Pincherle commutator converts the shift recurrence into closed form. □

**Corollary 9.4.2.** *Sheffer Shift Formula*
*Let $(s_n)$ be Sheffer for $(g,f)$, with $f$ delta and $g \in \mathscr{F}^{\times}$. Write $f' = f'(t)$ and $g' = g'(t)$. Then:*

*(i)* $\theta_{g,f} = g^{-1}\theta_f g = \left(x - \frac{g'}{g}\right)(f')^{-1}$.
*(ii)* $s_{n+1} = \left(x - \frac{g'}{g}\right)(f')^{-1}s_n$ *for* $n \geq 0$.

*Proof.*

(i) Since $gs_n$ is associated with $f$, $\theta_{g,f} = g^{-1}\theta_f g$. By Theorem 9.4.1, $\theta_f = x(f')^{-1}$.
Since $f'$ and $g$ commute, and $g' = gx - xg$, $g^{-1}\theta_f g = g^{-1}x(f')^{-1}g = g^{-1}xg(f')^{-1} = \left(x - \frac{g'}{g}\right)(f')^{-1}$.

(ii) By definition of the Sheffer shift, $\theta_{g,f}s_n = s_{n+1}$ for $n \geq 0$. Using (i), $s_{n+1} = \theta_{g,f}s_n = \left(x - \frac{g'}{g}\right)(f')^{-1}s_n$.

*Structural Commentary*: Sheffer shifts are conjugated associated shifts.
*Key Insight*: Conjugation by $g$ contributes the correction $g'/g$. □

**Method 9.4.3.** *Building Sheffer Sequences*
To construct the Sheffer sequence for $(g, f)$:

(i) Verify $o(f) = 1$ and $o(g) = 0$.
(ii) If $\overline{f}$ is explicit, compute $S(y,t) := g(\overline{f}(t))^{-1}e^{y\overline{f}(t)}$, where $s_n(y) = n![t^n]S(y,t)$.
(iii) If $\overline{f}$ is unavailable, solve recursively from $f(t)s_n = ns_{n-1}$, where $\langle g(t)f(t)^k \mid s_n\rangle = n!\delta_{n,k}$.
(iv) For associated sequences, set $g = 1$ and use Theorem 9.4.1.

**Definition 9.4.4.** *Stirling and Exponential Polynomials*

(i) *Stirling numbers*: $S(n,k)$ are defined by $x^n = \sum_{k=0}^{n} S(n,k)(x)_k$.
(ii) *Exponential polynomials*: define $\phi_n(x) := \sum_{k=0}^{n} S(n,k)x^k$.
(iii) *Boundary convention*: $S(n,-1) = 0$ and $S(n,k) = 0$ for $k > n$.

**Example 9.4.5.** *Lower Factorials and Stirling Numbers*
For $f(t) = e^t - 1$, $f'(t) = e^t$, hence $p_{n+1}(x) = xe^{-t}p_n(x) = xp_n(x-1)$. With $p_0 = 1$, this gives $p_n = (x)_n$.
For $f(t) = \log(1+t)$, $\overline{f}(t) = e^t - 1$, and the associated sequence is $(\phi_n)$.
The inverse umbral relation is $x^n = \sum_{k=0}^{n} S(n,k)(x)_k$, and $\phi_n(x) = \sum_{k=0}^{n} S(n,k)x^k$.
The recurrence formula gives $\phi_{n+1}(x) = x(1+t)\phi_n(x) = x\big(\phi_n(x) + \phi_n'(x)\big)$,
so coefficient comparison yields $S(n+1,k) = S(n,k-1) + kS(n,k)$.

**Definition 9.4.6.** *q- and Logarithmic Umbral Variants*

(i) *Gaussian coefficient*: for $0 \leq k \leq n$, $\binom{n}{k}_q := \prod_{j=1}^{n}(1-q^j)/\left(\prod_{j=1}^{k}(1-q^j)\prod_{j=1}^{n-k}(1-q^j)\right)$.
(ii) *q-umbral calculus*: the deformation $\binom{n}{k} \rightsquigarrow \binom{n}{k}_q$.
(iii) *Logarithmic umbral calculus*: the logarithmic-type variant based on harmonic logarithms.

**Remark 9.4.7.** *Further Directions*
Classical umbral methods extend to many polynomial families and operations:

(i) *Families*: lower factorial, Stirling, exponential, Gould, central factorial, Abel, Mittag–Leffler, Bessel, Bell, Hermite, Bernoulli, Euler, Laguerre, Poisson–Charlier, actuarial, Meixner, Pidduck, Narumi, Boole, Peters, Mahler, and Mott polynomials.
(ii) *Operations*: connection constants, duplication formulas, Lagrange inversion, cross sequences, Steffensen sequences, operational formulas, inverse relations, recurrence solving, and binomial convolution.
(iii) *Variants*: $q$-umbral and logarithmic umbral calculi replace or modify the binomial structure through the objects in Definition 9.4.6.

# 10 Fourier Structure

## 10.1 Fourier Orthogonality

Fourier structure studies trigonometric series through orthogonality, projection, kernels, and convergence, with inner-product geometry reappearing throughout.

**Definition 10.1.1.** *Core Fourier Objects*

(i) A *trigonometric polynomial* is a finite sum $a_0 + \sum_{n=1}^{N}(a_n\cos nx + b_n\sin nx)$, equivalently $\sum_{n=-N}^{N}\alpha_n e^{inx}$.
(ii) If $f$ is Riemann integrable on $[-\pi,\pi]$, its *Fourier coefficients* are $\widehat{f}(m) := \frac{1}{2\pi}\int_{-\pi}^{\pi} f(x)e^{-imx}\,dx$, $m \in \mathbb{Z}$, and $f(x) \sim_{\text{coef}} \sum_{m=-\infty}^{\infty}\widehat{f}(m)e^{imx}$ records coefficients, not convergence.

(iii) A sequence $(\phi_n)$ on $[a,b]$ is *orthonormal* if $\int_a^b \phi_n(x)\overline{\phi_m(x)}\,dx = 0 \quad (n \neq m)$ and $\int_a^b |\phi_n(x)|^2\,dx = 1$. Its Fourier coefficients are $c_n := \int_a^b f(t)\overline{\phi_n(t)}\,dt$.

*Key Insight*: Fourier coefficients are orthonormal coordinates; convergence is a separate analytic question.

**Definition 10.1.2.** *Period-One Fourier Normalization*
Let $C(\mathbb{R}/\mathbb{Z};\mathbb{C})$ denote the continuous functions $f:\mathbb{R}\to\mathbb{C}$ satisfying $f(x+1)=f(x)$ for all $x$.
For $f,g \in C(\mathbb{R}/\mathbb{Z};\mathbb{C})$, set $\langle f,g\rangle_{\mathbb{T}} := \int_0^1 f(x)\overline{g(x)}\,dx$; $\|f\|_{L^2(\mathbb{T})} := \sqrt{\langle f,f\rangle_{\mathbb{T}}}$; and $e_n(x) := e^{2\pi i n x}$ for $n \in \mathbb{Z}$.
The *period-one Fourier coefficient* is $\widehat{f}_{\mathbb{T}}(n) := \langle f, e_n\rangle_{\mathbb{T}} = \int_0^1 f(x)e^{-2\pi i n x}\,dx$.
A period-one trigonometric polynomial is a finite sum $\sum_{n=-N}^{N} c_n e_n$.

**Perspective 10.1.3.** *Two Equivalent Fourier Clocks*
The period-one and $2\pi$-periodic conventions differ by $t = 2\pi x$; only normalizing constants change.
*Key Insight*: Changing Fourier clocks changes constants, not the projection geometry.

**Proposition 10.1.4.** *Characters Are an Orthonormal System*
*For $m,n \in \mathbb{Z}$ and $N \geq 0$:*

*(i)* $\langle e_n, e_m\rangle_{\mathbb{T}} = \delta_{n,m}$.

*(ii)* *If* $P = \sum_{k=-N}^{N} c_k e_k$, *then* $\langle P, e_n\rangle_{\mathbb{T}} = c_n$ *for* $-N \leq n \leq N$, *and* $\|P\|^2_{L^2(\mathbb{T})} = \sum_{k=-N}^{N} |c_k|^2$.

*Proof.*

(i) $\langle e_n, e_m\rangle_{\mathbb{T}} = \int_0^1 e^{2\pi i(n-m)x}\,dx$. This equals 1 if $n=m$ and $\frac{e^{2\pi i(n-m)}-1}{2\pi i(n-m)} = 0$ if $n \neq m$.

(ii) By (i), $\langle P, e_n\rangle_{\mathbb{T}} = \sum_{k=-N}^{N} c_k\delta_{k,n} = c_n$, and $\|P\|^2_{L^2(\mathbb{T})} = \langle P,P\rangle_{\mathbb{T}} = \sum_{j,k=-N}^{N} c_j\overline{c_k}\delta_{j,k} = \sum_{k=-N}^{N}|c_k|^2$.

*Structural Commentary*: The Gram matrix of $\{e_{-N},\dots,e_N\}$ is the identity.
*Key Insight*: Character orthogonality kills all cross terms. □

## 10.2 Fourier Approximation

**Perspective 10.2.1.** *Fourier and Plancherel Roadmap*
Path: orthonormal characters, Fejer density, best $L^2$ projection, then Parseval/Plancherel.
*Key Insight*: Fourier theory is finite-dimensional orthogonal projection plus an approximation theorem.

**Definition 10.2.2.** *Periodic Convolution and Fejer Kernels*
For $f,g \in C(\mathbb{R}/\mathbb{Z};\mathbb{C})$, define their *periodic convolution* as $(f*g)(x) := \int_0^1 f(y)g(x-y)\,dy$.
For $N \geq 1$, the *Fejer kernel* is $F_N(x) := \sum_{n=-N}^{N}\left(1-\frac{|n|}{N}\right)e_n(x) = \frac{1}{N}\left|\sum_{n=0}^{N-1} e_n(x)\right|^2$.




Figure 191: The Fejer kernels are nonnegative trigonometric polynomial approximate identities on $\mathbb{R}/\mathbb{Z}$.

**Theorem 10.2.3.** *Fejer Approximation on $\mathbb{R}/\mathbb{Z}$*
*Let $F_N$ be the period-one Fejer kernel, $N \geq 1$. Then:*

*(i)* *$F_N$ is a nonnegative trigonometric polynomial and* $\int_0^1 F_N(x)\,dx = 1$.
*(ii)* *For each $0 < \delta < 1/2$,* $\sup_{\delta \leq x \leq 1-\delta} F_N(x) \to 0$.
*(iii)* *If $f \in C(\mathbb{R}/\mathbb{Z};\mathbb{C})$, then $f * F_N$ is a trigonometric polynomial for each $N$.*
*(iv)* *If $f \in C(\mathbb{R}/\mathbb{Z};\mathbb{C})$, then $f * F_N \rightrightarrows f$ on $\mathbb{R}/\mathbb{Z}$.*

*Proof.*

(i) By definition, $F_N = \sum_{n=-N}^{N} \left(1 - \frac{|n|}{N}\right) e_n$ is trigonometric, and $F_N = \frac{1}{N}\left|\sum_{n=0}^{N-1} e_n\right|^2 \geq 0$.
Since $\int_0^1 e_n(x)\,dx = 0$ for $n \neq 0$, the $e_0$ coefficient gives $\int_0^1 F_N(x)\,dx = 1$.

(ii) For $x \notin \mathbb{Z}$, $F_N(x) = \frac{\sin^2(\pi N x)}{N \sin^2(\pi x)}$. If $m_\delta := \min_{\delta \leq x \leq 1-\delta} \sin^2(\pi x) > 0$, then $0 \leq F_N(x) \leq \frac{1}{N m_\delta}$ on $[\delta, 1-\delta]$.

(iii) Since $f * e_n = \widehat{f}_{\mathbb{T}}(n) e_n$ and $F_N$ is a finite linear combination of the $e_n$, $f * F_N$ is a trigonometric polynomial.

(iv) Write $(f * F_N)(x) = \int_0^1 f(x-t) F_N(t)\,dt$. Let $M = \|f\|_\infty$ and $\varepsilon > 0$.
Uniform continuity gives $0 < \delta < 1/2$ such that $|f(x-t) - f(x)| < \varepsilon$ for all $x$ and $t \in [0,\delta] \cup [1-\delta, 1]$.
By (i), $F_N \geq 0$ and $\int_0^1 F_N(t)\,dt = 1$; hence $|(f * F_N)(x) - f(x)| \leq \varepsilon + 2M \sup_{\delta \leq t \leq 1-\delta} F_N(t)$.
By (ii), $\limsup_{N\to\infty} \|f * F_N - f\|_\infty \leq \varepsilon$. Since $\varepsilon$ is arbitrary, $f * F_N \rightrightarrows f$.

*Structural Commentary*: Fejer kernels are positive approximate identities.
*Key Insight*: Normalized mass concentrating at 0 forces uniform convergence. □

**Perspective 10.2.4.** *Fourier Coefficients as Projections*
Fourier coefficients are projections onto orthonormal directions, modeled finitely by Theorems 2.1.5, 1.4.1, and 1.4.5; infinite directions make convergence separate.

**Theorem 10.2.5.** *Fourier Coefficients Give Best Mean-Square Approximation*
*Let $\langle u, v\rangle := \int_a^b u\overline{v}$ and $\|u\|_2^2 := \langle u, u\rangle$. Let $(\phi_m)$ be orthonormal and let $f$ satisfy $\|f\|_2 < \infty$.*
*For $n \geq 1$, set $c_m := \langle f, \phi_m\rangle$, $s_n := \sum_{m=1}^{n} c_m \phi_m$, and $t_n := \sum_{m=1}^{n} \gamma_m \phi_m$. Then:*

*(i) $\|f - s_n\|_2 \leq \|f - t_n\|_2$, with equality iff $\gamma_m = c_m$ for $m = 1, \ldots, n$.*

*(ii) $\sum_{m=1}^{\infty} |c_m|^2 \leq \|f\|_2^2$, hence $c_m \to 0$.*

*Proof.*

(i) By orthonormality, $\|f - t_n\|_2^2 = \|f\|_2^2 - 2\,\mathrm{Re} \sum_{m=1}^{n} c_m \overline{\gamma_m} + \sum_{m=1}^{n} |\gamma_m|^2 = \|f\|_2^2 - \sum_{m=1}^{n} |c_m|^2 + \sum_{m=1}^{n} |\gamma_m - c_m|^2$.
Thus $\|f - t_n\|_2^2 \geq \|f - s_n\|_2^2$, with equality iff $\gamma_m = c_m$ for all $m \leq n$.

(ii) Taking $\gamma_m = c_m$ in the identity from (i) gives $\|f - s_n\|_2^2 = \|f\|_2^2 - \sum_{m=1}^{n} |c_m|^2 \geq 0$.
Hence $\sum_{m=1}^{n} |c_m|^2 = \|s_n\|_2^2 \leq \|f\|_2^2$ for all $n$. Letting $n \to \infty$ gives Bessel's inequality.
Since $(|c_m|^2)$ is summable, $|c_m|^2 \to 0$, hence $c_m \to 0$.

*Structural Commentary*: Orthogonality diagonalizes the finite error.
*Key Insight*: Fourier coefficients are projection coordinates. □

**Theorem 10.2.6.** *Dirichlet Kernel and Local Fourier Convergence*
*Let $f$ be Riemann integrable and $2\pi$-periodic, with $\widehat{f}(m) := \frac{1}{2\pi}\int_{-\pi}^{\pi} f(t) e^{-imt}\,dt$.*
*For $N \geq 0$, set $s_N(f;x) := \sum_{m=-N}^{N} \widehat{f}(m) e^{imx}$ and $D_N(t) := \sum_{m=-N}^{N} e^{imt}$. Then:*

*(i) $D_N(t) = \frac{\sin((N+\frac{1}{2})t)}{\sin(t/2)}$ for $t \notin 2\pi\mathbb{Z}$, with $D_N(0) = 2N+1$, and $s_N(f;x) = \frac{1}{2\pi}\int_{-\pi}^{\pi} f(x-t) D_N(t)\,dt$.*

*(ii) If $x \in \mathbb{R}$ and there exist $M, \delta > 0$ such that $|f(x+t) - f(x)| \leq M|t|$ for $|t| < \delta$, then $s_N(f;x) \to f(x)$.*

*(iii) If $g$ is Riemann integrable and $2\pi$-periodic and $f = g$ near $x$, then $s_N(f;x) - s_N(g;x) \to 0$.*

*Proof.*

(i) For $z = e^{it} \neq 1$, $D_N(t) = z^{-N}\frac{1 - z^{2N+1}}{1-z} = \frac{\sin((N+\frac{1}{2})t)}{\sin(t/2)}$, and $D_N(0) = \sum_{m=-N}^{N} 1 = 2N+1$.
Substituting $\widehat{f}(m)$ and interchanging the finite sum with the integral gives
$s_N(f;x) = \frac{1}{2\pi}\int_{-\pi}^{\pi} f(u) \sum_{m=-N}^{N} e^{im(x-u)}\,du = \frac{1}{2\pi}\int_{-\pi}^{\pi} f(x-t) D_N(t)\,dt$.

(ii) Since $\frac{1}{2\pi}\int_{-\pi}^{\pi} D_N(t)\,dt = 1$, (i) gives $s_N(f;x) - f(x) = \frac{1}{2\pi}\int_{-\pi}^{\pi}(f(x-t) - f(x)) D_N(t)\,dt$.
Set $q_x(t) := \frac{f(x-t) - f(x)}{\sin(t/2)}$ for $0 < |t| \leq \pi$ and $q_x(0) := 0$. The hypothesis makes $q_x$ bounded near 0 and Riemann integrable on $[-\pi, \pi]$. Hence $s_N(f;x) - f(x) = \frac{1}{2\pi}\int_{-\pi}^{\pi} q_x(t) \sin((N+\frac{1}{2})t)\,dt$.
Since $\sin((N+\frac{1}{2})t) = \sin Nt \cos(t/2) + \cos Nt \sin(t/2)$, Theorem 10.2.5 applied to $q_x(t)\cos(t/2)$ and $q_x(t)\sin(t/2)$ sends both coefficients to 0.

(iii) Apply (ii) to $h := f - g$. Then $h = 0$ near $x$, so $s_N(h;x) \to h(x) = 0$.
By linearity, $s_N(h;x) = s_N(f;x) - s_N(g;x)$.

*Structural Commentary*: Dirichlet kernels express partial sums as oscillatory local averages.
*Key Insight*: Local regularity makes the oscillatory error vanish. □

**Corollary 10.2.7.** *Uniform Approximation by Trigonometric Polynomials*
*Let $f \in C(\mathbb{R};\mathbb{C})$ be $2\pi$-periodic. For every $\varepsilon > 0$, there is $P(x) = \sum_{n=-N}^{N} c_n e^{inx}$ such that $\|f - P\|_\infty < \varepsilon$.*

*Proof.* Let $\mathbb{T} := \{z \in \mathbb{C} : |z| = 1\}$ and define $F : \mathbb{T} \to \mathbb{C}$ by $F(e^{ix}) := f(x)$.
Periodicity makes $F$ well-defined and continuous.
The algebra $\mathcal{T} := \operatorname{span}\{z^n : n \in \mathbb{Z}\} \subset C(\mathbb{T})$ is self-adjoint, separates points, and vanishes nowhere.
By Theorem 9.2.9, choose $Q(z) = \sum_{n=-N}^{N} c_n z^n \in \mathcal{T}$ with $\|F - Q\|_{C(\mathbb{T})} < \varepsilon$.
Set $P(x) := Q(e^{ix}) = \sum_{n=-N}^{N} c_n e^{inx}$; then $\|f - P\|_\infty = \|F - Q\|_{C(\mathbb{T})} < \varepsilon$.
*Structural Commentary*: Trigonometric polynomials are Laurent polynomials pulled back from $\mathbb{T}$.
*Key Insight*: Stone–Weierstrass on $\mathbb{T}$ gives uniform periodic approximation. □

**Theorem 10.2.8.** *Parseval's Theorem for Trigonometric Fourier Series*
*Let $f, g$ be Riemann integrable and $2\pi$-periodic.*
*Set $\langle u, v\rangle_2 := \frac{1}{2\pi}\int_{-\pi}^{\pi} u(x)\overline{v(x)}\,dx$, $\|u\|_2^2 := \langle u, u\rangle_2$, and $s_N(f) := \sum_{|n|\le N} \widehat{f}(n)e^{inx}$. Then:*

*(i)* $\|f - s_N(f)\|_2 \to 0$.
*(ii)* $\langle f, g\rangle_2 = \sum_{n\in\mathbb{Z}} \widehat{f}(n)\overline{\widehat{g}(n)}$.
*(iii)* $\|f\|_2^2 = \sum_{n\in\mathbb{Z}} \left|\widehat{f}(n)\right|^2$.

*Proof.*

(i) Put $d_N(u) := \|u - s_N(u)\|_2$. If $u$ is continuous, choose a trigonometric polynomial $P$ with $\|u - P\|_2 < \varepsilon$; for $N \ge \deg P$, Theorem 10.2.5 gives $d_N(u) \le \|u - P\|_2 < \varepsilon$, so $d_N(u) \to 0$.
For Riemann integrable $f$, choose continuous $h$ with $\|f - h\|_2 < \varepsilon$.
Since $\|s_N(h - f)\|_2 \le \|h - f\|_2$, $d_N(f) \le 2\|f - h\|_2 + d_N(h)$; hence $\limsup_{N\to\infty} d_N(f) \le 2\varepsilon$. Let $\varepsilon \downarrow 0$.

(ii) By orthogonality, $\langle e^{inx}, g\rangle_2 = \overline{\widehat{g}(n)}$. Hence $\langle s_N(f), g\rangle_2 = \sum_{|n|\le N} \widehat{f}(n)\langle e^{inx}, g\rangle_2 = \sum_{|n|\le N} \widehat{f}(n)\overline{\widehat{g}(n)}$.
By (i) and Cauchy–Schwarz, $\langle s_N(f), g\rangle_2 \to \langle f, g\rangle_2$.

(iii) Set $g = f$ in (ii). Then $\widehat{g}(n) = \widehat{f}(n)$ and $\langle f, f\rangle_2 = \|f\|_2^2$, so $\|f\|_2^2 = \sum_{n\in\mathbb{Z}} \left|\widehat{f}(n)\right|^2$.

*Structural Commentary*: Density turns finite projection identities into $L^2$ identities.
*Key Insight*: Partial Fourier projections converge in $L^2$. □

**Method 10.2.9.** *Fourier Proof Patterns*

(i) *Characters*: use orthogonality of $e_n$ to compute coefficients and finite-dimensional projections.
(ii) *Kernels*: rewrite partial sums or averages as convolutions with Dirichlet or Fejer kernels.
(iii) *Approximation*: use Fejer positivity or Stone–Weierstrass to approximate continuous periodic functions by trigonometric polynomials.
(iv) *$L^2$ control*: project onto finite orthonormal spans, then pass to limits by density.
(v) *Localization*: express the error as an oscillatory integral and use local regularity near the point.

*Key Insight*: Fourier theory is finite-dimensional orthogonal projection plus an approximation theorem.

## 10.3 Further Topics in Fourier Analysis

**Example 10.3.1.** *A Gaussian Fourier Integral*
Define $f(t) = \int_{-\infty}^{\infty} e^{-x^2}\cos(xt)\,dx$ and $g(t) = -\int_{-\infty}^{\infty} xe^{-x^2}\sin(xt)\,dx$.
Then $f'(t) = g(t)$, and integration by parts gives $2f'(t) + tf(t) = 0$. Since $f(0) = \sqrt{\pi}$, $f(t) = \sqrt{\pi}\exp\left(-\frac{t^2}{4}\right)$.

# Part XIII
# Measure Foundations and Extension

## 1 Sigma-Algebras and Measurable Spaces

### 1.1 Sigma-Algebras and Measurable Structures

Measure theory first builds measurable sets, then measurable functions and integrals. Outer measure covers from outside; measurability splits test sets without losing measure.

**Perspective 1.1.1.** *Why Abstract Measure Comes First*
Lebesgue integration is built so that, under hypotheses such as domination, $f_n \to f$ implies $\int f_n\,d\mu \to \int f\,d\mu$; see Theorem 1.3.7. The abstract data separate into a measurable space $(X, \mathcal{M})$, a countably additive measure $\mu : \mathcal{M} \to [0, +\infty]$, and an integral defined by approximation from simple functions.
*Key Insight*: Countable additivity is the set-level structure behind the convergence theorems.

**Definition 1.1.2.** *$\sigma$-Algebras and Measurable Spaces*
Let $X$ be a set. A family $\mathcal{M} \subseteq \mathcal{P}(X)$ is a *$\sigma$-algebra* on $X$ if:

(i) $\emptyset, X \in \mathcal{M}$;
(ii) $E \in \mathcal{M}$ implies $X \setminus E \in \mathcal{M}$;
(iii) $E_n \in \mathcal{M}$ for $n \in \mathbb{N}$ implies $\bigcup_{n=1}^{\infty} E_n \in \mathcal{M}$.

The pair $(X, \mathcal{M})$ is a *measurable space*; the members of $\mathcal{M}$ are *measurable sets*.

**Proposition 1.1.3.** *Closure and Generation for $\sigma$-Algebras*
*Let $\mathcal{M} \subseteq \mathcal{P}(X)$ be a $\sigma$-algebra.*

*(i)* $(E_n)_{n\in\mathbb{N}} \subseteq \mathcal{M} \Rightarrow \bigcap_{n=1}^{\infty} E_n \in \mathcal{M}$.
*(ii) If $(\mathcal{M}_\alpha)_{\alpha\in A}$ is a nonempty family of $\sigma$-algebras on $X$, then $\bigcap_{\alpha\in A} \mathcal{M}_\alpha$ is a $\sigma$-algebra on $X$.*
*(iii) For $\mathcal{A} \subseteq \mathcal{P}(X)$, let $\mathscr{S}_{\mathcal{A}}$ be the family of $\sigma$-algebras $\mathcal{N}$ on $X$ with $\mathcal{A} \subseteq \mathcal{N}$. Then $\sigma(\mathcal{A}) := \bigcap_{\mathcal{N}\in\mathscr{S}_{\mathcal{A}}} \mathcal{N}$ is the smallest $\sigma$-algebra containing $\mathcal{A}$.*

*Proof.*

(i) Since $X \setminus E_n \in \mathcal{M}$, Theorem 1.5.3 gives $\bigcap_n E_n = X \setminus \bigcup_n (X \setminus E_n) \in \mathcal{M}$.
(ii) Set $\mathcal{N} := \bigcap_{\alpha\in A} \mathcal{M}_\alpha$. Then $\emptyset, X \in \mathcal{N}$, and $E \in \mathcal{N}$ implies $X \setminus E \in \mathcal{N}$. Likewise $(E_n) \subseteq \mathcal{N}$ implies $\bigcup_n E_n \in \mathcal{N}$.
(iii) The indexing class in the displayed definition is nonempty since $\mathcal{P}(X)$ belongs to it. By (ii), its intersection is a $\sigma$-algebra; it contains $\mathcal{A}$ and is contained in every $\sigma$-algebra containing $\mathcal{A}$.

*Structural Commentary*: Generation is intersection over all admissible $\sigma$-algebras.
*Key Insight*: Intersections preserve universal closure properties. □

**Example 1.1.4.** *Basic $\sigma$-Algebras*
Standard models:

(i) Extremes: $\{\emptyset, X\} \subseteq \mathcal{M} \subseteq \mathcal{P}(X)$ for every $\sigma$-algebra $\mathcal{M}$ on $X$.
(ii) For $E \subseteq X$, $\sigma(\{E\}) = \{\emptyset, E, X \setminus E, X\}$, with repetitions omitted.
(iii) On $\mathbb{N}$, if $O := \{2n-1 : n \in \mathbb{N}\}$, then $\sigma(\{O\}) = \{\emptyset, O, \mathbb{N} \setminus O, \mathbb{N}\}$.
(iv) If $X$ is uncountable, $\mathcal{C}_X := \{E \subseteq X : E \text{ is countable or } X \setminus E \text{ is countable}\}$ is the countable–cocountable $\sigma$-algebra.

*Key Insight*: A $\sigma$-algebra records exactly which distinctions the theory is allowed to observe.

**Definition 1.1.5.** *Borel and Extended Borel $\sigma$-Algebras*

(i) For $n \in \mathbb{N}$, the *Borel $\sigma$-algebra* on $\mathbb{R}^n$ is $\mathcal{B}(\mathbb{R}^n) := \sigma(\{U \subseteq \mathbb{R}^n : U \text{ is open}\})$. On $\mathbb{R}$, it is also generated by open intervals, closed intervals, half-open intervals, or open half-rays $(a, +\infty)$.
(ii) The *extended Borel $\sigma$-algebra* on $\overline{\mathbb{R}}$ is $\overline{\mathcal{B}} := \{E \cup F : E \in \mathcal{B}(\mathbb{R}),\ F \subseteq \{-\infty, +\infty\}\}$.

**Definition 1.1.6.** *Monotone Classes*
A nonempty $\mathcal{C} \subseteq \mathcal{P}(X)$ is a *monotone class* iff $(E_n) \subseteq \mathcal{C},\ E_n \uparrow E \Rightarrow E \in \mathcal{C}$ and $(F_n) \subseteq \mathcal{C},\ F_n \downarrow F \Rightarrow F \in \mathcal{C}$.
For $\mathcal{A} \subseteq \mathcal{P}(X)$, define $\mathfrak{M}_{\mathcal{A}} := \{\mathcal{C} \subseteq \mathcal{P}(X) : \mathcal{C} \text{ is a monotone class},\ \mathcal{A} \subseteq \mathcal{C}\}$; $\operatorname{mon}(\mathcal{A}) := \bigcap \mathfrak{M}_{\mathcal{A}}$.
Then $\operatorname{mon}(\mathcal{A})$ is the least monotone class containing $\mathcal{A}$, and $\operatorname{mon}(\mathcal{A}) \subseteq \sigma(\mathcal{A})$.
*Key Insight*: Monotone closure captures limits, not finite unions.

**Definition 1.1.7.** *Algebras of Sets and Premeasures*
Let $X$ be a set.

(i) An *algebra of sets* (or *field of sets*) on $X$ is $\mathcal{A} \subseteq \mathcal{P}(X)$ such that $\emptyset, X \in \mathcal{A}$; $E \in \mathcal{A} \Rightarrow X \setminus E \in \mathcal{A}$; and $E_1, \dots, E_n \in \mathcal{A} \Rightarrow \bigcup_{j=1}^{n} E_j \in \mathcal{A}$. Equivalently, $\mathcal{A}$ is a ring of sets containing $X$.
(ii) A *premeasure* (or *measure on an algebra*) on $\mathcal{A}$ is $\mu : \mathcal{A} \to [0, +\infty]$ such that $\mu(\emptyset) = 0$ and, whenever $(E_n)_{n\in\mathbb{N}} \subseteq \mathcal{A}$ is pairwise disjoint with $\bigcup_{n=1}^{\infty} E_n \in \mathcal{A}$, then $\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} \mu(E_n)$.

*Key Insight*: Premeasures are countably additive where the algebra permits unions.

**Theorem 1.1.8.** *Monotone Class Theorem for Algebras*
*Let $X$ be a set, let $\mathcal{A} \subseteq \mathcal{P}(X)$ be an algebra, and set $\mathcal{C} := \mathrm{mon}(\mathcal{A})$. Then $\mathcal{C} = \sigma(\mathcal{A})$.*

*Proof.* Since $\sigma(\mathcal{A})$ is monotone and contains $\mathcal{A}$, $\mathcal{C} \subseteq \sigma(\mathcal{A})$. It remains to prove that $\mathcal{C}$ is a $\sigma$-algebra.
Set $\mathcal{D} := \{E \in \mathcal{C} : E^c \in \mathcal{C}\}$. If $E_n \uparrow E$ in $\mathcal{D}$, then $E_n^c \downarrow E^c$; if $E_n \downarrow E$, then $E_n^c \uparrow E^c$. Thus $\mathcal{D}$ is monotone.
Since $\mathcal{A} \subseteq \mathcal{D}$, minimality gives $\mathcal{C} \subseteq \mathcal{D}$, so $\mathcal{C}$ is closed under complements.
For $B \in \mathcal{C}$, set $\mathcal{D}_B := \{E \in \mathcal{C} : E \cap B \in \mathcal{C}\}$. This class is monotone. If $B \in \mathcal{A}$, then $\mathcal{A} \subseteq \mathcal{D}_B$, hence $\mathcal{C} \subseteq \mathcal{D}_B$.
Therefore $E \cap B \in \mathcal{C}$ for $E \in \mathcal{C}$ and $B \in \mathcal{A}$. Fix $E \in \mathcal{C}$; the preceding sentence gives $\mathcal{A} \subseteq \mathcal{D}_E$, hence $\mathcal{C} \subseteq \mathcal{D}_E$.
Thus $\mathcal{C}$ is closed under finite intersections. Complements and finite intersections give finite unions.
If $(E_n) \subseteq \mathcal{C}$, then $F_N := \bigcup_{n=1}^{N} E_n \in \mathcal{C}$ and $F_N \uparrow \bigcup_{n=1}^{\infty} E_n$, so $\bigcup_{n=1}^{\infty} E_n \in \mathcal{C}$.
Hence $\mathcal{C}$ is a $\sigma$-algebra containing $\mathcal{A}$, so $\sigma(\mathcal{A}) \subseteq \mathcal{C}$.
*Structural Commentary*: Monotone lifting gives Boolean closure, then countable unions.
*Key Insight*: Show each closure property holds on a monotone class containing $\mathcal{A}$. □

**Corollary 1.1.9.** *Monotone-Class Test*
*Let $\mathcal{A} \subseteq \mathcal{P}(X)$ be an algebra and let $\mathcal{C} \subseteq \mathcal{P}(X)$ be a monotone class. If $\mathcal{A} \subseteq \mathcal{C}$, then $\sigma(\mathcal{A}) \subseteq \mathcal{C}$.*

*Proof.* Since $\mathcal{A} \subseteq \mathcal{C}$ and $\mathcal{C}$ is monotone, $\mathrm{mon}(\mathcal{A}) \subseteq \mathcal{C}$. By Theorem 1.1.8, $\mathrm{mon}(\mathcal{A}) = \sigma(\mathcal{A})$.
*Structural Commentary*: Algebra checks plus monotone stability control $\sigma(\mathcal{A})$.
*Key Insight*: Prove on the algebra; extend by monotone closure. □

# 2 Measures, Charges, and Basic Examples

## 2.1 Measures, Charges, and Basic Constructions

**Definition 2.1.1.** *Rings, $\sigma$-Rings, and Set Functions*
Let $\mathcal{A} \subseteq \mathcal{P}(X)$.

(i) $\mathcal{A}$ is a *ring of sets* iff $\emptyset \in \mathcal{A}$ and $A, B \in \mathcal{A} \Rightarrow A \cup B,\ A \setminus B \in \mathcal{A}$
(ii) $\mathcal{A}$ is a *$\sigma$-ring* iff it is a ring and $(A_n) \subseteq \mathcal{A} \Rightarrow \bigcup_{n=1}^{\infty} A_n \in \mathcal{A}$
(iii) A *set function* on $\mathcal{A}$ is $\phi : \mathcal{A} \to [-\infty, +\infty]$ with $\{+\infty, -\infty\} \nsubseteq \phi(\mathcal{A})$.
(iv) $\phi$ is *additive* iff $A \cap B = \emptyset \Rightarrow \phi(A \cup B) = \phi(A) + \phi(B)$
(v) $\phi$ is *countably additive* iff $(A_n) \subseteq \mathcal{A}$ pairwise disjoint, $\bigcup_n A_n \in \mathcal{A} \Rightarrow \phi\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \phi(A_n)$.

*Structural Commentary*: Rings support finite operations; $\sigma$-rings support countable unions.

**Definition 2.1.2.** *Measures on Measurable Spaces*
Let $(X, \mathcal{M})$ be a measurable space. A *measure* on $(X, \mathcal{M})$ is a map $\mu : \mathcal{M} \to [0, +\infty]$ such that:

(i) $\mu(\emptyset) = 0$;
(ii) if $(E_n) \subseteq \mathcal{M}$ is pairwise disjoint, then $\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} \mu(E_n)$

The triple $(X, \mathcal{M}, \mu)$ is a *measure space*.
*Key Insight*: Measure is countably additive size on the chosen observable sets.

**Definition 2.1.3.** *Finite and $\sigma$-Finite Measures*
Let $(X, \mathcal{M}, \mu)$ be a measure space.

(i) For nonnegative terms, $\sum_{n=1}^{\infty} a_n := \lim_{N\to\infty} \sum_{n=1}^{N} a_n \in [0, +\infty]$.
(ii) $\mu$ is *finite* iff $\mu(X) < +\infty$.
(iii) $\mu$ is *$\sigma$-finite* iff there exist $E_n \in \mathcal{M}$ such that $X = \bigcup_{n=1}^{\infty} E_n$ and $\mu(E_n) < +\infty$ for all $n$.
(iv) A measurable set $E \in \mathcal{M}$ has *finite measure* iff $\mu(E) < +\infty$.
(v) A measurable set $E \in \mathcal{M}$ is *$\sigma$-finite* iff $E = \bigcup_{n=1}^{\infty} E_n$ for some $E_n \in \mathcal{M}$ with $\mu(E_n) < +\infty$ for all $n$.

*Key Insight*: $\sigma$-finiteness replaces one finite global bound by a countable finite-measure exhaustion.

**Example 2.1.4.** *First Abstract Measures*

(i) The zero set function $\mu(E) = 0$ on $\mathcal{M}$ is a finite measure.
(ii) If $X \neq \emptyset$ and $\mathcal{M} = 2^X$, $\mu(\emptyset) = 0$ and $\mu(E) = +\infty$ for $E \neq \emptyset$ give a measure, neither finite nor $\sigma$-finite.

(iii) For $p \in X$, the *unit mass* $\delta_p(E) := \mathbf{1}_E(p)$ on $2^X$ is a finite measure.
(iv) On $2^{\mathbb{N}}$, *counting measure* is $\#(E)$ for finite $E$ and $+\infty$ otherwise; it is $\sigma$-finite.
More generally, $(a_n) \subseteq [0, +\infty]$ gives $\mu(E) := \sum_{n \in E} a_n$, and every measure on $2^{\mathbb{N}}$ is uniquely of this form.
(v) If $X$ is uncountable and $\mathcal{M} = 2^X$, $\mu(E) = 0$ for countable $E$ and $\mu(E) = +\infty$ otherwise defines a measure.
(vi) If $X$ is uncountable and $\mathcal{C}_X = \{E \subseteq X : E \text{ is countable or } X \setminus E \text{ is countable}\}$, then $\mu(E) = 0$ on countable sets and $\mu(E) = 1$ on cocountable sets defines a measure on $\mathcal{C}_X$.
(vii) On $2^{\mathbb{N}}$, $\mu(E) = 0$ for finite $E$ and $\mu(E) = +\infty$ otherwise is not a measure, since $\mathbb{N} = \bigcup_{n=1}^{\infty} \{n\}$ gives $+\infty \neq 0$; the constant $+\infty$ rule fails at $\emptyset$.
(viii) Lebesgue measure on $\mathcal{B}(\mathbb{R})$ is $\sigma$-finite and gives length $b-a$ to standard intervals; Borel–Stieltjes measures come from increasing right-continuous functions as in Theorem 4.1.4.

*Structural Commentary*: Measures may be zero, atomic, counting-type, cardinality-based, or geometric.
*Key Insight*: Countable additivity turns disjoint decompositions into nonnegative series.

**Proposition 2.1.5.** *Elementary Measure Constructions*
*Let $(X, \mathcal{M})$ be a measurable space.*

*(i) If $\mu$ is a measure on $\mathcal{M}$ and $A \in \mathcal{M}$, then $\mu_A(E) := \mu(A \cap E)$ defines a measure $\mu_A$ on $\mathcal{M}$.*
*(ii) If $\mu_1, \dots, \mu_m$ are measures on $\mathcal{M}$ and $a_1, \dots, a_m \geq 0$, then $\lambda := \sum_{j=1}^{m} a_j \mu_j$ is a measure on $\mathcal{M}$.*
*(iii) If $(\mu_n)_{n \geq 1}$ are measures on $\mathcal{M}$ with $\mu_n(X) = 1$, then $\lambda := \sum_{n=1}^{\infty} 2^{-n} \mu_n$ is a measure on $\mathcal{M}$ and $\lambda(X) = 1$.*

*Proof.*
(i) $\mu_A(\emptyset) = 0$. If $(E_k)$ is disjoint in $\mathcal{M}$, then $A \cap \bigsqcup_k E_k = \bigsqcup_k (A \cap E_k)$, whence $\mu_A(\bigsqcup_k E_k) = \sum_k \mu_A(E_k)$.
(ii) $\lambda(\emptyset) = 0$. For disjoint $(E_k)$, finite summation gives $\lambda(\bigsqcup_k E_k) = \sum_{j=1}^{m} a_j \sum_k \mu_j(E_k) = \sum_k \lambda(E_k)$.
(iii) $\lambda(\emptyset) = 0$. For disjoint $(E_k)$, nonnegative reordering gives $\lambda(\bigsqcup_k E_k) = \sum_n 2^{-n} \sum_k \mu_n(E_k) = \sum_k \lambda(E_k)$.
Also $\lambda(X) = \sum_n 2^{-n} \mu_n(X) = \sum_n 2^{-n} = 1$.

*Structural Commentary*: New measures come from restriction, superposition, and countable convex mixtures.
*Key Insight*: Nonnegative series make the bookkeeping order-independent. □

**Proposition 2.1.6.** *Basic Laws for Measures*
*Let $(X, \mathcal{M}, \mu)$ be a measure space.*

*(i) For every $n \geq 1$, if $E_1, \dots, E_n \in \mathcal{M}$ are pairwise disjoint, then $\mu\left(\bigcup_{k=1}^{n} E_k\right) = \sum_{k=1}^{n} \mu(E_k)$.*
*(ii) If $A, B \in \mathcal{M}$ and $A \subseteq B$, then $\mu(A) \leq \mu(B)$.*
*(iii) If $A, B \in \mathcal{M}$, $A \subseteq B$, and $\mu(A) < +\infty$, then $\mu(B \setminus A) = \mu(B) - \mu(A)$.*
*In particular, if $\mu(A) = 0$, then $\mu(B \setminus A) = \mu(B)$.*
*(iv) If $E \in \mathcal{M}$ and $(E_k)_{k \geq 1} \subseteq \mathcal{M}$ satisfy $E \subseteq \bigcup_{k=1}^{\infty} E_k$, then $\mu(E) \leq \sum_{k=1}^{\infty} \mu(E_k)$.*

*Proof.*
(i) Define $G_k = E_k$ for $1 \leq k \leq n$ and $G_k = \emptyset$ for $k > n$. Then $(G_k)$ is pairwise disjoint and countable additivity gives $\mu\left(\bigcup_{k=1}^{n} E_k\right) = \mu\left(\bigcup_{k=1}^{\infty} G_k\right) = \sum_{k=1}^{\infty} \mu(G_k) = \sum_{k=1}^{n} \mu(E_k)$.
(ii) Since $B = A \sqcup (B \setminus A)$, (i) gives $\mu(B) = \mu(A) + \mu(B \setminus A)$, and $\mu(B \setminus A) \geq 0$, hence $\mu(A) \leq \mu(B)$.
(iii) Since $B = A \sqcup (B \setminus A)$, (i) gives $\mu(B) = \mu(A) + \mu(B \setminus A)$. Because $\mu(A) < +\infty$, subtraction yields $\mu(B \setminus A) = \mu(B) - \mu(A)$. If $\mu(A) = 0$, this becomes $\mu(B \setminus A) = \mu(B)$.
(iv) Set $F_1 = E \cap E_1$ and $F_k = E \cap \left(E_k \setminus \bigcup_{j=1}^{k-1} E_j\right)$ for $k \geq 2$. Then $(F_k)_{k \geq 1} \subseteq \mathcal{M}$ is pairwise disjoint, $\bigcup_{k=1}^{\infty} F_k = E$, and $F_k \subseteq E_k$ for all $k$. Countable additivity and (ii) give $\mu(E) = \sum_{k=1}^{\infty} \mu(F_k) \leq \sum_{k=1}^{\infty} \mu(E_k)$.

*Structural Commentary*: Finite additivity yields order and excision; disjointization turns covers into sums.
*Key Insight*: Measure disjoint pieces, then compare. □

**Theorem 2.1.7.** *Continuity from Below*
*Let $\mathcal{A}$ be a ring of subsets of $X$, and let $\phi$ be a countably additive set function on $\mathcal{A}$.*
*If $(A_n)_{n \geq 1} \subseteq \mathcal{A}$ and $A \in \mathcal{A}$ satisfy $A_n \subseteq A_{n+1}$ for all $n$ and $A = \bigcup_{n=1}^{\infty} A_n$, then $\lim_{N \to \infty} \phi(A_N) = \phi(A)$.*

*Proof.* Set $B_1 = A_1$ and $B_n = A_n \setminus A_{n-1}$ for $n \geq 2$. Since $\mathcal{A}$ is a ring, each $B_n \in \mathcal{A}$.
The sets $(B_n)$ are pairwise disjoint, $A_N = \bigsqcup_{n=1}^{N} B_n$, and $A = \bigsqcup_{n=1}^{\infty} B_n$.
Countable additivity, applied to the finite union by appending empty sets, gives $\phi(A_N) = \sum_{n=1}^{N} \phi(B_n)$, while countable additivity gives $\phi(A) = \sum_{n=1}^{\infty} \phi(B_n)$. Thus $\lim_{N \to \infty} \phi(A_N) = \phi(A)$.

*Structural Commentary*: Increasing chains split into disjoint increments.
*Key Insight*: Continuity is convergence of partial sums. □

**Definition 2.1.8.** *Charges*
Let $(X, \mathcal{M})$ be a measurable space. A *charge* is a map $\lambda : \mathcal{M} \to \mathbb{R}$ such that:

(i) $\lambda(\emptyset) = 0$.

(ii) For every pairwise disjoint $(E_n) \subseteq \mathcal{M}$, $\sum_{n=1}^{\infty} \lambda(E_n)$ converges in $\mathbb{R}$ and $\lambda\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} \lambda(E_n)$.

*Key Insight*: A charge is a real-valued countably additive set function.

**Definition 2.1.9.** *Variation Candidates of a Charge*
Let $\lambda$ be a charge on $\mathcal{M}$. For $E \in \mathcal{M}$, define

(i) $\mathfrak{P}_{\mathcal{M}}(E) := \{(A_j)_{j=1}^n : n \in \mathbb{N},\ E = \bigsqcup_{j=1}^n A_j,\ A_j \in \mathcal{M}\}$, the finite measurable partitions of $E$;
(ii) $\pi_\lambda(E) := \sup\{\lambda(A) : A \subseteq E,\ A \in \mathcal{M}\}$, the *positive variation* candidate;
(iii) $\nu_\lambda(E) := \sup\{\sum_{j=1}^n |\lambda(A_j)| : (A_j)_{j=1}^n \in \mathfrak{P}_{\mathcal{M}}(E)\}$, the *total variation* candidate.

For $E = \emptyset$, the empty partition is allowed and contributes 0.

**Proposition 2.1.10.** *Charge Operations and Variation Measures*
*Let $(X, \mathcal{M})$ be a measurable space, let $m \geq 1$, and let $\lambda, \lambda_1, \ldots, \lambda_m$ be charges on $\mathcal{M}$.*

*(i) If $a_1, \ldots, a_m \in \mathbb{R}$ and $\Lambda : \mathcal{M} \to \mathbb{R}$ is defined by $\Lambda(E) := \sum_{j=1}^m a_j \lambda_j(E)$, then $\Lambda$ is a charge.*
*(ii) If $\mu$ is a measure on $\mathcal{M}$, $f \in \mathscr{L}^1(\mu)$, and $\lambda_f : \mathcal{M} \to \mathbb{R}$ is defined by $\lambda_f(A) := \int_A f\, d\mu$, then $\lambda_f$ is a charge.*
*(iii) $\pi_\lambda : \mathcal{M} \to [0, +\infty]$ is a measure.*
*(iv) $\nu_\lambda : \mathcal{M} \to [0, +\infty]$ is a measure.*

*Proof.*

(i) $\Lambda(\emptyset) = 0$. For $E = \bigsqcup_n E_n$, $\sum_n \Lambda(E_n)$ converges as $\sum_{j=1}^m a_j \sum_n \lambda_j(E_n)$; $\Lambda(E) = \sum_{j=1}^m a_j \sum_n \lambda_j(E_n) = \sum_n \Lambda(E_n)$.

(ii) $\lambda_f(\emptyset) = 0$. For disjoint $(A_n)$, $\sum_n |\lambda_f(A_n)| \leq \int_{\bigsqcup_n A_n} |f|\, d\mu < +\infty$ and $\lambda_f(\bigsqcup_n A_n) = \int_{\bigsqcup_n A_n} f\, d\mu = \sum_n \int_{A_n} f\, d\mu = \sum_n \lambda_f(A_n)$.

(iii) $\pi_\lambda(\emptyset) = 0$ and $\pi_\lambda \geq 0$. Let $E = \bigsqcup_n E_n$. For measurable $A \subseteq E$, $\lambda(A) = \sum_n \lambda(A \cap E_n) \leq \sum_n \pi_\lambda(E_n)$, so $\pi_\lambda(E) \leq \sum_n \pi_\lambda(E_n)$. Put $S_N = \sum_{n=1}^N \pi_\lambda(E_n)$. If $S_N < +\infty$, choose measurable $A_n \subseteq E_n$ for $n \leq N$ with $\lambda(A_n) > \pi_\lambda(E_n) - \frac{\varepsilon}{N}$; then $\pi_\lambda(E) > S_N - \varepsilon$. If $S_N = +\infty$, some $E_\ell$ has $\pi_\lambda(E_\ell) = +\infty$, hence $\pi_\lambda(E) = +\infty$. Thus $\pi_\lambda(E) \geq S_N$ for all $N$, and $N \to \infty$ gives equality.

(iv) $\nu_\lambda(\emptyset) = 0$ and $\nu_\lambda \geq 0$. Let $E = \bigsqcup_n E_n$ and $v(P) = \sum_{A \in P} |\lambda(A)|$. Refining $P = (A_j)_{j=1}^r \in \mathfrak{P}_{\mathcal{M}}(E)$ by $E_n$ gives $v(P) \leq \sum_{j=1}^r \sum_n |\lambda(A_j \cap E_n)| \leq \sum_n \nu_\lambda(E_n)$, so $\nu_\lambda(E) \leq \sum_n \nu_\lambda(E_n)$. Put $T_N = \sum_{n=1}^N \nu_\lambda(E_n)$. If $T_N < +\infty$, choose $P_n \in \mathfrak{P}_{\mathcal{M}}(E_n)$ for $n \leq N$ with $v(P_n) > \nu_\lambda(E_n) - \frac{\varepsilon}{N}$. Adjoining the tail $\bigcup_{n>N} E_n$ gives $\nu_\lambda(E) > T_N - \varepsilon$. If $T_N = +\infty$, some $E_\ell$ has $\nu_\lambda(E_\ell) = +\infty$, and partitions of $E_\ell$ extend to $E$, so $\nu_\lambda(E) = +\infty$. Thus $\nu_\lambda(E) \geq T_N$ for all $N$, and $N \to \infty$ gives equality.

*Structural Commentary*: Suprema preserve additivity after refinement.
*Key Insight*: Refine first, then take suprema. □

# 3 Outer Measures and Caratheodory Extension

## 3.1 Outer Measures and Extension Theorems

**Definition 3.1.1.** *Abstract Outer Measure and Clean Division*
Let $M$ be a set.

(i) An *abstract outer measure* on $M$ is a map $\omega : 2^M \to [0, +\infty]$ such that $\omega(\emptyset) = 0$, $A \subseteq B \Rightarrow \omega(A) \leq \omega(B)$, and $\omega(\bigcup_{k=1}^\infty A_k) \leq \sum_{k=1}^\infty \omega(A_k)$ for all $A, B, A_k \subseteq M$.
(ii) For an outer measure $\omega$, $E \subseteq M$ is *$\omega$-measurable* iff $\omega(X) = \omega(X \cap E) + \omega(X \cap E^c)$ for every $X \subseteq M$.

For $\omega = m_n^*$, $\omega$-measurability is Caratheodory measurability.

**Definition 3.1.2.** *Outer Measure Induced by a Covering Family*
Let $\mathcal{S} \subseteq \mathcal{P}(X)$ and let $\rho : \mathcal{S} \to [0, +\infty]$.

(i) For $E \subseteq X$, $\mathcal{C}_{\mathcal{S}}(E) := \{(S_k)_{k\geq 1} \in \mathcal{S}^{\mathbb{N}} : E \subseteq \bigcup_{k=1}^{\infty} S_k\}$.

(ii) The induced map $\rho^* : \mathcal{P}(X) \to [0, +\infty]$ is defined by $\rho^*(\emptyset) = 0$ and, for $E \neq \emptyset$, $\rho^*(E) := \inf_{(S_k)\in\mathcal{C}_{\mathcal{S}}(E)} \sum_{k=1}^{\infty} \rho(S_k)$, with $\inf \emptyset = +\infty$.

The map $\rho^*$ is the *outer measure induced by $\rho$ and $\mathcal{S}$*.
*Key Insight*: Minimize total $\rho$-cost over countable $\mathcal{S}$-covers.

**Proposition 3.1.3.** *Cover-Induced Outer Measure*
*For $\mathcal{S} \subseteq \mathcal{P}(X)$ and $\rho : \mathcal{S} \to [0, +\infty]$, the cover-induced map $\rho^* : \mathcal{P}(X) \to [0, +\infty]$ is an outer measure on $X$.*

*Proof.* By construction, $\rho^*(\emptyset) = 0$ and $\rho^* \geq 0$. If $A \subseteq B$, then $\mathcal{C}_{\mathcal{S}}(B) \subseteq \mathcal{C}_{\mathcal{S}}(A)$, hence $\rho^*(A) \leq \rho^*(B)$.
Let $E = \bigcup_k E_k$, $I = \{k : E_k \neq \emptyset\}$, and assume $I \neq \emptyset$ and $\sum_k \rho^*(E_k) < +\infty$.
Choose $(S_{k,j})_j \in \mathcal{C}_{\mathcal{S}}(E_k)$ for $k \in I$ with $\sum_j \rho(S_{k,j}) < \rho^*(E_k) + \varepsilon 2^{-k}$.
Since $(S_{k,j})_{k\in I, j\geq 1}$ covers $E$, $\rho^*(E) \leq \sum_{k\in I} \sum_j \rho(S_{k,j}) \leq \sum_k \rho^*(E_k) + \varepsilon$; let $\varepsilon \downarrow 0$. Excluded cases are immediate.
*Structural Commentary*: Inclusion orders covers; regrouping covers sums.
*Key Insight*: Covers generate subadditivity. □

**Lemma 3.1.4.** *Finite-Measure Test for Caratheodory Measurability*
*Let $\omega : 2^M \to [0, +\infty]$ be an outer measure. Then $E \subseteq M$ is $\omega$-measurable iff, for all $X \subseteq M$, $\omega(X) < +\infty \Longrightarrow \omega(X) \geq \omega(X \cap E) + \omega(X \cap E^c)$.*

*Proof.*
($\Rightarrow$) If $E$ is $\omega$-measurable, then $\omega(X) = \omega(X \cap E) + \omega(X \cap E^c)$ $\forall X \subseteq M$, hence finite-test inequality holds.
($\Leftarrow$) For all $X \subseteq M$, $X = (X \cap E) \sqcup (X \cap E^c)$ gives $\omega(X) \leq \omega(X \cap E) + \omega(X \cap E^c)$. If $\omega(X) < +\infty$, the hypothesis gives $\geq$; if $\omega(X) = +\infty$, $\geq$ is automatic in $[0, +\infty]$. Thus equality holds for every $X$.
*Structural Commentary*: To verify a clean split, only finite outer-measure test sets need real work.
*Key Insight*: Subadditivity supplies one inequality for free. □




Figure 192: Two cuts split $X$ into four regions; $E_1 \setminus E_2$ is region 2.

**Definition 3.1.5.** *Zero Sets*
For outer measure $\omega : 2^M \to [0, +\infty]$, *$\omega$-zero set* is $Z \subseteq M$ with $\omega(Z) = 0$; $\mathcal{Z}_\omega := \{Z \subseteq M : \omega(Z) = 0\}$.

**Theorem 3.1.6.** *Caratheodory Measurable Sets*
*Let $\omega : 2^M \to [0, +\infty]$ be an outer measure.*
*Let $\mathcal{M}_\omega := \{E \subseteq M : \omega(X) = \omega(X \cap E) + \omega(X \cap E^c)\ \forall X \subseteq M\}$ and $\mathcal{Z}_\omega := \{Z \subseteq M : \omega(Z) = 0\}$. Then:*
*(i) $\mathcal{Z}_\omega \subseteq \mathcal{M}_\omega$;*
*(ii) $\mathcal{M}_\omega$ is a $\sigma$-algebra;*
*(iii) $\omega|_{\mathcal{M}_\omega}$ is countably additive;*
*(iv) if $E \in \mathcal{M}_\omega$ and $Z \in \mathcal{Z}_\omega$, then $E \cup Z, E \setminus Z, E \triangle Z \in \mathcal{M}_\omega$ and $\omega(E \cup Z) = \omega(E \setminus Z) = \omega(E \triangle Z) = \omega(E)$.*

*Proof.*
(i) If $Z \in \mathcal{Z}_\omega$ and $X \subseteq M$, then $\omega(X) \leq \omega(X \cap Z) + \omega(X \cap Z^c) \leq 0 + \omega(X)$, so equality holds and $Z \in \mathcal{M}_\omega$.

(ii) Complements are immediate. If $E, F \in \mathcal{M}_\omega$, with $E^1 = E$, $E^0 = E^c$, $F^1 = F$, and $F^0 = F^c$, then $\omega(X) = \sum_{\epsilon,\eta\in\{0,1\}} \omega(X \cap E^\epsilon \cap F^\eta)$; hence finite Boolean combinations are measurable.
For pairwise disjoint $E_i \in \mathcal{M}_\omega$, set $E = \bigsqcup_i E_i$ and $F_N = (\bigcup_{i=1}^N E_i)^c$.
Finite splitting gives $\omega(X) \geq \sum_{i=1}^N \omega(X \cap E_i) + \omega(X \cap E^c)$ for every $N$.
Letting $N \to \infty$ and using subadditivity gives $\omega(X) = \sum_{i=1}^\infty \omega(X \cap E_i) + \omega(X \cap E^c)$.
Since $\omega(X \cap E) \leq \sum_i \omega(X \cap E_i)$ and $\omega(X) \leq \omega(X \cap E) + \omega(X \cap E^c)$, $E \in \mathcal{M}_\omega$.
General countable unions follow from $H_1 = E_1$ and $H_n = E_n \setminus \bigcup_{i<n} E_i$.
(iii) If $E = \bigsqcup_{i=1}^\infty E_i$ with $E_i \in \mathcal{M}_\omega$, then the equality in (ii) with $X = E$ gives $\omega(E) = \sum_{i=1}^\infty \omega(E_i)$.
(iv) By (i)–(ii), $E \cup Z, E \setminus Z, E\triangle Z \in \mathcal{M}_\omega$.
Since $E \cap Z$ and $Z \setminus E$ are null, additivity gives $\omega(E \setminus Z) = \omega(E \cup Z) = \omega(E\triangle Z) = \omega(E)$.

*Structural Commentary*: Measurable sets preserve countable additivity.
*Key Insight*: Measurable cuts turn subadditivity into equality. □

**Corollary 3.1.7.** *Disjointization of Countable Measurable Unions*
*Let $\omega : 2^M \to [0, +\infty]$ be an outer measure and let $(E_k)_{k\geq 1} \subseteq \mathcal{M}_\omega$.*
*Set $F_1 := E_1$ and $F_k := E_k \setminus \bigcup_{i=1}^{k-1} E_i$ for $k \geq 2$.*
*Then $(F_k)$ is pairwise disjoint in $\mathcal{M}_\omega$, $\bigcup_{k\geq 1} F_k = \bigcup_{k\geq 1} E_k$, and $\omega(\bigcup_{k\geq 1} E_k) = \sum_{k\geq 1} \omega(F_k)$ in $[0, +\infty]$.*

*Proof.* Since $\mathcal{M}_\omega$ is a $\sigma$-algebra, $F_k = E_k \cap (\bigcup_{i<k} E_i)^c \in \mathcal{M}_\omega$. If $i < k$, then $F_i \cap F_k \subseteq E_i \cap (\bigcup_{j<k} E_j)^c = \emptyset$.
For $x \in \bigcup_{k\geq 1} E_k$, $x \in F_{\min\{j : x\in E_j\}}$; hence $\bigcup_{k\geq 1} F_k = \bigcup_{k\geq 1} E_k$, and countable additivity gives $\omega(\bigcup_{k\geq 1} E_k) = \sum_{k\geq 1} \omega(F_k)$.
*Structural Commentary*: Disjointization turns unions into sums.
*Key Insight*: Peel off earlier pieces so a countable union becomes a disjoint sum. □

**Definition 3.1.8.** *Countable Cover Classes*
Let $\mathcal{S} \subseteq \mathcal{P}(X)$.

(i) *Countable-union class*: $\mathcal{S}_\sigma := \left\{ \bigcup_{n=1}^\infty S_n : (S_n)_{n\geq 1} \in \mathcal{S}^{\mathbb{N}} \right\}$.

(ii) *Countable-intersection class*: $\mathcal{S}_{\sigma\delta} := \left\{ \bigcap_{n=1}^\infty A_n : (A_n)_{n\geq 1} \in (\mathcal{S}_\sigma)^{\mathbb{N}} \right\}$.

*Key Insight*: $\mathcal{S}_\sigma$ closes under countable unions; $\mathcal{S}_{\sigma\delta}$ then closes once under countable intersections.

**Proposition 3.1.9.** *Finite Outer-Measure Envelope*
*Let $\mathcal{S} \subseteq \mathcal{P}(X)$, let $\rho : \mathcal{S} \to [0, +\infty]$, and let $\rho^*$ be the cover-induced map.*
*If $\emptyset \neq E \subseteq X$ and $\rho^*(E) < +\infty$, then there exists $A \in \mathcal{S}_{\sigma\delta}$ such that:*
*(i) $E \subseteq A$ and $\rho^*(A) = \rho^*(E)$.*
*(ii) If $E$ and every $S \in \mathcal{S}$ are $\rho^*$-measurable, then $A$ is $\rho^*$-measurable and $\rho^*(A \setminus E) = 0$.*

*Proof.* $\forall n \geq 1$, choose $(S_{n,k})_{k\geq 1} \in \mathcal{C}_\mathcal{S}(E)$ with $\sum_{k=1}^\infty \rho(S_{n,k}) < \rho^*(E) + \frac{1}{n}$. Set $A_n := \bigcup_{k=1}^\infty S_{n,k}$, $A := \bigcap_{n=1}^\infty A_n$.
Then $A_n \in \mathcal{S}_\sigma$, $A \in \mathcal{S}_{\sigma\delta}$, $E \subseteq A \subseteq A_n$, and $\rho^*(A_n) < \rho^*(E) + \frac{1}{n}$.
(i) Monotonicity gives $\rho^*(E) \leq \rho^*(A) \leq \rho^*(A_n) < \rho^*(E) + \frac{1}{n}$ for all $n$, hence $\rho^*(A) = \rho^*(E)$.
(ii) If $E$ and every $S \in \mathcal{S}$ are $\rho^*$-measurable, then the $\rho^*$-measurable sets form a $\sigma$-algebra, so $A_n$ and $A$ are $\rho^*$-measurable. Since $E \subseteq A$ and $\rho^*(E) < +\infty$, excision gives $\rho^*(A \setminus E) = \rho^*(A) - \rho^*(E) = 0$.

*Structural Commentary*: Finite cost gives a countable-intersection envelope.
*Key Insight*: Cover within $\frac{1}{n}$, then intersect. □

**Definition 3.1.10.** *Outer Measure Generated by a Premeasure*
Let $\mu : \mathcal{A} \to [0, +\infty]$ be a premeasure on an algebra $\mathcal{A} \subseteq \mathcal{P}(X)$.

(i) *Algebra covers*: For $B \subseteq X$, set $\mathcal{C}_\mathcal{A}(B) := \{(E_j)_{j\geq 1} \in \mathcal{A}^{\mathbb{N}} : B \subseteq \bigcup_{j=1}^\infty E_j\}$.

(ii) *Generated outer measure*: Define $\mu^* : \mathcal{P}(X) \to [0, +\infty]$ by $\mu^*(\emptyset) = 0$ and, for $B \neq \emptyset$, $\mu^*(B) := \inf_{(E_j)\in\mathcal{C}_\mathcal{A}(B)} \sum_{j=1}^\infty \mu(E_j)$, with $\inf \emptyset = +\infty$.

*Key Insight*: Cover arbitrary sets by algebra sets and minimize total premeasure.

**Lemma 3.1.11.** *Generated Outer-Measure Properties*
*Let $\mu : \mathcal{A} \to [0, +\infty]$ be a premeasure on an algebra $\mathcal{A} \subseteq \mathcal{P}(X)$.*
*For $B \subseteq X$, put $\mathcal{C}_{\mathcal{A}}(B) := \{(E_j) \in \mathcal{A}^{\mathbb{N}} : B \subseteq \bigcup_j E_j\}$.*
*Set $\inf \emptyset = +\infty$, $\mu^*(\emptyset) = 0$, and $\mu^*(B) := \inf_{(E_j)\in\mathcal{C}_{\mathcal{A}}(B)} \sum_j \mu(E_j)$ for $B \neq \emptyset$. Then:*

*(i)* $\mu^*(\emptyset) = 0$.
*(ii)* *If $C \subseteq D \subseteq X$, then* $\mu^*(C) \leq \mu^*(D)$.
*(iii)* *If $(B_n)_{n\geq 1} \subseteq \mathcal{P}(X)$, then* $\mu^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} \mu^*(B_n)$.
*(iv)* *If $E \in \mathcal{A}$, then* $\mu^*(E) = \mu(E)$.

*Proof.*

(i) This is the assigned value of $\mu^*(\emptyset)$.
(ii) If $C \subseteq D$, then $\mathcal{C}_{\mathcal{A}}(D) \subseteq \mathcal{C}_{\mathcal{A}}(C)$; taking infima gives $\mu^*(C) \leq \mu^*(D)$.
(iii) Let $B = \bigcup_n B_n$ and $I = \{n : B_n \neq \emptyset\}$. If $I = \emptyset$ or $\sum_n \mu^*(B_n) = +\infty$, the claim is immediate.
Otherwise, for $\varepsilon > 0$ choose $(E_{n,k})_{k\geq 1} \in \mathcal{C}_{\mathcal{A}}(B_n)$ for $n \in I$ with $\sum_k \mu(E_{n,k}) < \mu^*(B_n) + \varepsilon 2^{-n}$.
Then $(E_{n,k})_{n\in I, k\geq 1}$ covers $B$, so $\mu^*(B) \leq \sum_{n\in I}\sum_k \mu(E_{n,k}) \leq \sum_n \mu^*(B_n) + \varepsilon$; let $\varepsilon \downarrow 0$.
(iv) Cover $E, \emptyset, \emptyset, \ldots$ gives $\mu^*(E) \leq \mu(E)$. Conversely, for any cover $(F_n) \in \mathcal{C}_{\mathcal{A}}(E)$, set $G_n := E \cap \left(F_n \setminus \bigcup_{k<n} F_k\right)$.
Then $(G_n)$ is disjoint in $\mathcal{A}$, $\bigcup_n G_n = E$, and $G_n \subseteq F_n$, so $\mu(E) = \sum_n \mu(G_n) \leq \sum_n \mu(F_n)$.
Taking infima gives $\mu(E) \leq \mu^*(E)$.

*Structural Commentary*: Cover containment and regrouping give the outer-measure axioms.
*Key Insight*: Disjointize algebra covers to recover $\mu$. □

**Proposition 3.1.12.** *Single-Set Envelope Comparison*
*Let $\mu : \mathcal{A} \to [0, +\infty]$ be a premeasure on an algebra $\mathcal{A} \subseteq \mathcal{P}(X)$, let $\mu^*$ be the generated outer measure, and define $\mu^\sharp : \mathcal{P}(X) \to [0, +\infty]$ by $\mu^\sharp(B) := \inf\{\mu(C) : C \in \mathcal{A},\ B \subseteq C\}$ with $\inf \emptyset = +\infty$.*

*(i)* $\mu^\sharp(E) = \mu(E)$ *for every* $E \in \mathcal{A}$.
*(ii)* $\mu^*(B) \leq \mu^\sharp(B)$ *for every* $B \subseteq X$.

*Proof.*

(i) If $E \in \mathcal{A}$, then $E$ is admissible, so $\mu^\sharp(E) \leq \mu(E)$.
Conversely, $E \subseteq C \in \mathcal{A}$ implies $\mu(E) \leq \mu(C)$ by monotonicity of $\mu$, hence $\mu(E) \leq \mu^\sharp(E)$.
(ii) If no $C \in \mathcal{A}$ contains $B$, then $\mu^\sharp(B) = +\infty$. Otherwise, each $C \in \mathcal{A}$ with $B \subseteq C$ gives the algebra cover $C, \emptyset, \emptyset, \ldots$, so $\mu^*(B) \leq \mu(C)$. Taking infima over such $C$ gives $\mu^*(B) \leq \mu^\sharp(B)$.

*Structural Commentary*: One algebra superset is a special countable cover.
*Key Insight*: The single-set envelope is larger. □

**Example 3.1.13.** *Single-Set Envelopes Need Not Be Outer Measures*
Let $X$ be uncountable, let $\mathcal{A}$ be the finite–cofinite algebra on $X$, and set $\mu(E) := 0$ for finite $E$, $\mu(E) := +\infty$ for cofinite $E$. Then $\mu$ is a premeasure, $\mu^*(B) = 0$ iff $B$ is countable, and $\mu^\sharp(B) = 0$ iff $B$ is finite.
For any countably infinite $S \subseteq X$, $\mu^\sharp(S) = +\infty$ while $S = \bigcup_{x\in S}\{x\}$ and $\sum_{x\in S} \mu^\sharp(\{x\}) = 0$, QED.
*Structural Commentary*: Countable covers can be strictly smaller than one-set covers.
*Key Insight*: Outer measure is built from countable covers precisely to recover countable subadditivity.

**Theorem 3.1.14.** *Generated Caratheodory Extension*
*Let $\mu : \mathcal{A} \to [0, +\infty]$ be a premeasure on an algebra $\mathcal{A} \subseteq \mathcal{P}(X)$. For $B \subseteq X$, put $\mathcal{C}_{\mathcal{A}}(B) := \{(E_j) \in \mathcal{A}^{\mathbb{N}} : B \subseteq \bigcup_j E_j\}$. Define $\mu^*(\emptyset) = 0$, $\mu^*(B) := \inf_{(E_j)\in\mathcal{C}_{\mathcal{A}}(B)} \sum_j \mu(E_j)$ for $B \neq \emptyset$ with $\inf \emptyset = +\infty$, and $\mathcal{C}_\mu := \{E \subseteq X : (\forall B \subseteq X)\ \mu^*(B) = \mu^*(B \cap E) + \mu^*(B \setminus E)\}$. Then:*

*(i)* $\mathcal{A} \subseteq \mathcal{C}_\mu$ *and* $\mu^*|_{\mathcal{A}} = \mu$.
*(ii)* $\mathcal{C}_\mu$ *is a $\sigma$-algebra on $X$.*
*(iii)* $\mu^*|_{\mathcal{C}_\mu}$ *is a complete measure extending $\mu$.*

*Proof.* Lemma 3.1.11 makes $\mu^*$ an outer measure and gives $\mu^*|_{\mathcal{A}} = \mu$.

(i) Let $E \in \mathcal{A}$ and $B \subseteq X$. Subadditivity gives $\mu^*(B) \leq \mu^*(B \cap E) + \mu^*(B \setminus E)$. If $(F_n) \in \mathcal{C}_{\mathcal{A}}(B)$, then $F_n \cap E, F_n \setminus E \in \mathcal{A}$ and $\mu(F_n) = \mu(F_n \cap E) + \mu(F_n \setminus E)$, so $\mu^*(B \cap E) + \mu^*(B \setminus E) \leq \sum_n \mu(F_n)$.
Infimizing over $(F_n)$ gives equality; hence $E \in \mathcal{C}_\mu$. The lemma gives $\mu^*|_{\mathcal{A}} = \mu$.

(ii) The class $\mathcal{C}_\mu$ is exactly the $\mu^*$-measurable class $\mathcal{M}_{\mu^*}$.
Since $\mu^*$ is an outer measure, Theorem 3.1.6(ii) gives that $\mathcal{C}_\mu$ is a $\sigma$-algebra on $X$.
(iii) Theorem 3.1.6(iii) gives countable additivity of $\mu^*|_{\mathcal{C}_\mu}$. If $A \subseteq Z \in \mathcal{C}_\mu$ and $\mu^*(Z) = 0$, then monotonicity gives $\mu^*(A) = 0$, so Theorem 3.1.6(i) gives $A \in \mathcal{C}_\mu$. Thus $\mu^*|_{\mathcal{C}_\mu}$ is complete, and (i) shows it extends $\mu$.

*Structural Commentary*: Caratheodory equality is the additivity domain.
*Key Insight*: Algebra sets split every algebra cover. □

**Proposition 3.1.15.** *Caratheodory Splitters for Finitely Subadditive Size*
*Let $\alpha : \mathcal{P}(X) \to [0, +\infty)$ satisfy $0 \le \alpha(E) \le \alpha(E \cup F) \le \alpha(E) + \alpha(F)$ for all $E, F \subseteq X$.*
*Set $\mathcal{S} := \{E \subseteq X : (\forall A \subseteq X)\ \alpha(A) = \alpha(A \cap E) + \alpha(A \setminus E)\}$.*

*(i) If $\mathcal{S} \neq \emptyset$, then $\mathcal{S}$ is an algebra on $X$.*
*(ii) If $E, F \in \mathcal{S}$ and $E \cap F = \emptyset$, then $\alpha(E \cup F) = \alpha(E) + \alpha(F)$.*
*(iii) The hypothesis $\mathcal{S} \neq \emptyset$ in (i) is necessary: if $\alpha \equiv 1$, then $\mathcal{S} = \emptyset$.*

*Proof.*

(i) Choose $E_0 \in \mathcal{S}$. Splitting $\emptyset$ gives $\alpha(\emptyset) = 0$, hence $\emptyset, X \in \mathcal{S}$.
Complement symmetry gives $E \in \mathcal{S} \Rightarrow X \setminus E \in \mathcal{S}$. For $E, F \in \mathcal{S}$ and $A \subseteq X$, put $A_{11} = A \cap E \cap F$, $A_{10} = A \cap E \setminus F$, $A_{01} = A \cap F \setminus E$, and $A_{00} = A \setminus (E \cup F)$. By successive splitting,
$\alpha(A) = \alpha(A_{11}) + \alpha(A_{10}) + \alpha(A_{01}) + \alpha(A_{00})$; $\alpha(A \setminus (E \cap F)) = \alpha(A_{10}) + \alpha(A_{01}) + \alpha(A_{00})$.
Thus $\alpha(A) = \alpha(A \cap E \cap F) + \alpha(A \setminus (E \cap F))$, so $E \cap F \in \mathcal{S}$. Complements give finite unions.
(ii) Since $F \subseteq X \setminus E$, the splitting identity for $E$ applied to $A = E \cup F$ gives $\alpha(E \cup F) = \alpha(E) + \alpha(F)$.
(iii) If $\alpha \equiv 1$ and $E \in \mathcal{S}$, applying the splitting identity to $A = \emptyset$ gives $1 = 1 + 1$, impossible.

*Structural Commentary*: Splitters are Boolean once one splitter exists.
*Key Insight*: Empty-set splitting forces $\alpha(\emptyset) = 0$. □

**Theorem 3.1.16.** *$\sigma$-Finite Generated Extensions Are Unique*
*Let $\mu : \mathcal{A} \to [0, +\infty]$ be a premeasure on an algebra $\mathcal{A} \subseteq \mathcal{P}(X)$, let $\mu^*$ be generated by $\mu$, and put $\mathcal{C}_\mu := \{E \subseteq X : (\forall B \subseteq X)\ \mu^*(B) = \mu^*(B \cap E) + \mu^*(B \setminus E)\}$. Assume $X = \bigcup_n F_n$; $F_n \in \mathcal{A}$, $\mu(F_n) < +\infty$ $\forall n$.*

*(i) $\mu^*|_{\mathcal{C}_\mu}$ is a measure on $\mathcal{C}_\mu$ and extends $\mu$.*
*(ii) If $\nu$ is a measure on $\mathcal{C}_\mu$ with $\nu|_{\mathcal{A}} = \mu$, then $\nu = \mu^*|_{\mathcal{C}_\mu}$.*

*Proof.*

(i) By Theorem 3.1.14, $\mathcal{C}_\mu$ is complete $\sigma$-algebra, $\mathcal{A} \subseteq \mathcal{C}_\mu$, $\mu^*|_{\mathcal{C}_\mu}$ is complete measure with $(\mu^*|_{\mathcal{C}_\mu})|_{\mathcal{A}} = \mu$.
(ii) Let $\nu$ be a measure on $\mathcal{C}_\mu$ with $\nu|_{\mathcal{A}} = \mu$.
If $G \in \mathcal{C}_\mu$ and $G \subseteq \bigcup_n A_n$ with $A_n \in \mathcal{A}$, then $\nu(G) \le \sum_n \mu(A_n)$; hence $\nu(G) \le \mu^*(G)$.
If $\mu(X) < +\infty$, apply to $G = E$, $G = X \setminus E$, while $\nu(E) + \nu(X \setminus E) = \nu(X) = \mu^*(X) = \mu^*(E) + \mu^*(X \setminus E)$, gives $\nu(E) = \mu^*(E)$. In general, put $H_N := \bigcup_{j=1}^{N} F_j$. Then $H_N \in \mathcal{A}$ and $\mu(H_N) < +\infty$. Apply bound to $E \cap H_N$, $H_N \setminus E$, while $\nu(E \cap H_N) + \nu(H_N \setminus E) = \nu(H_N) = \mu^*(H_N) = \mu^*(E \cap H_N) + \mu^*(H_N \setminus E)$, gives $\nu(E \cap H_N) = \mu^*(E \cap H_N)$. Since $E \cap H_N \uparrow E$, continuity from below gives $\nu(E) = \mu^*(E)$.

*Structural Commentary*: Finite uniqueness localizes along a finite-mass exhaustion.
*Key Insight*: Compare on each finite piece, then pass upward. □

**Corollary 3.1.17.** *Uniqueness on the Generated $\sigma$-Algebra*
*Let $\mu : \mathcal{A} \to [0, +\infty]$ be a premeasure on an algebra $\mathcal{A} \subseteq \mathcal{P}(X)$, and assume $X = \bigcup_n F_n$ with $F_n \in \mathcal{A}$ and $\mu(F_n) < +\infty$. If measures $\nu_1, \nu_2$ on $\sigma(\mathcal{A})$ extend $\mu$, then $\nu_1 = \nu_2$.*

*Proof.* Put $H_n := \bigcup_{j=1}^{n} F_j$. Then $H_n \in \mathcal{A}$, $H_n \uparrow X$, and $\nu_i(H_n) = \mu(H_n) < +\infty$.
For fixed $n$, $\mathcal{D}_n := \{E \in \sigma(\mathcal{A}) : \nu_1(E \cap H_n) = \nu_2(E \cap H_n)\}$ is a Dynkin class by finite mass on $H_n$ and contains $\mathcal{A}$, since $A \cap H_n \in \mathcal{A}$. As $\mathcal{A}$ is a $\pi$-system, the $\pi$-$\lambda$ theorem gives $\mathcal{D}_n = \sigma(\mathcal{A})$.
Hence $\nu_1(E \cap H_n) = \nu_2(E \cap H_n)$ for all $n$; since $E \cap H_n \uparrow E$, continuity from below gives $\nu_1(E) = \nu_2(E)$.
*Structural Commentary*: Finite exhaustions reduce uniqueness to Dynkin closure.
*Key Insight*: Compare on $H_n$, then pass upward. □

**Example 3.1.18.** *Why $\sigma$-Finiteness Matters*
Let $X = \mathbb{Q} \cap (0, 1]$, and let $\mathcal{A}$ be the algebra of finite unions of sets $(a, b] \cap X$, where $0 \le a < b \le 1$ and $a, b \in \mathbb{Q}$.

(i) *Infinite algebra pieces*: Every nonempty $A \in \mathcal{A}$ is infinite.

(ii) *Generated $\sigma$-algebra*: For $r \in X$, choose $a_n \in \mathbb{Q} \cap [0, r)$ with $a_n \uparrow r$.
Then $\{r\} = \bigcap_{n=1}^{\infty} ((a_n, r] \cap X) \in \sigma(\mathcal{A})$, hence $\sigma(\mathcal{A}) = \mathcal{P}(X)$.

(iii) *Nonunique extensions*: Let $\mu_1$ be counting measure and $\mu_2 = 2\mu_1$ on $\mathcal{P}(X)$.
Since every nonempty $A \in \mathcal{A}$ is infinite, $\mu_1|_{\mathcal{A}} = \mu_2|_{\mathcal{A}}$, but $\mu_1(\{r\}) = 1 \neq 2 = \mu_2(\{r\})$ for $r \in X$.

Thus uniqueness may fail without $\sigma$-finiteness.
*Key Insight*: Infinite algebra pieces cannot determine singleton scale.

## 3.2 Euclidean Cells and Cell Outer Measure

**Definition 3.2.1.** *Cells, Intervals, and Volume in $\mathbb{R}^p$*
Fix $p \geq 1$.

(i) *Cells in $\mathbb{R}$*. For $a \leq b$, a cell with endpoints $a, b$ is $I \in \{(a, b), [a, b], (a, b], [a, b)\}$.
If $a = b$, only $[a, a]$ is nonempty.

(ii) *Length*. If $I$ is a cell with endpoints $a \leq b$, then $\ell(I) := b - a$; unbounded intervals $(-\infty, a)$, $(-\infty, a]$, $[a, +\infty)$, $(a, +\infty)$, and $\mathbb{R}$ are not cells and have length $+\infty$.

(iii) *Cells in $\mathbb{R}^p$*. A cell is $I = \prod_{i=1}^p I_i$, each $I_i$ a cell in $\mathbb{R}$; $\prod_i (a_i, b_i)$ is open, $\prod_i [a_i, b_i]$ is closed, $\prod_i [a_i, b_i)$ is *southwest*, $\prod_i (a_i, b_i]$ is *northeast*, and $I$ is a *cube* iff $\ell(I_1) = \cdots = \ell(I_p)$.

(iv) *Volume*. For a cell $I = \prod_{i=1}^p I_i$ with endpoints $a_i \leq b_i$, $\mathrm{vol}_p(I) := \prod_{i=1}^p \ell(I_i) = \prod_{i=1}^p (b_i - a_i)$.
Also $\mathrm{vol}_p(\emptyset) := 0$ and $\mathrm{vol}_p(I^\circ) = \mathrm{vol}_p(\overline{I}) = \mathrm{vol}_p(I)$.

(v) *Unbounded products*. For $I = \prod_{i=1}^p I_i$ with possibly unbounded intervals, $\mathrm{vol}_p(I) = 0$ if $\exists j : \ell(I_j) = 0$, $\mathrm{vol}_p(I) = +\infty$ if $\ell(I_i) > 0$ for all $i$ and $\exists j : \ell(I_j) = +\infty$, and $\mathrm{vol}_p(I) = \prod_{i=1}^p \ell(I_i)$ otherwise.

**Remark 3.2.2.** *Cells versus Intervals*
Cells are precisely bounded intervals with finite endpoints, so every cell is an interval but not conversely.
Open cells are open in $\mathbb{R}^p$, closed cells are closed in $\mathbb{R}^p$, and nonempty half-open cells are neither.
In $\mathbb{R}$, $(-\infty, a)$ and $(b, +\infty)$ are open, $(-\infty, a]$ and $[b, +\infty)$ are closed, and $\mathbb{R} = (-\infty, +\infty)$ is clopen.
*Key Insight*: Topology likes open and closed endpoints; measure bookkeeping often likes half-open endpoints.

**Proposition 3.2.3.** *Half-Open Cell Calculus*
*Fix $\sigma \in \{$southwest, northeast$\}$, and let $\mathcal{H}_\sigma$ be the half-open cells in $\mathbb{R}^p$ with orientation $\sigma$.*

*(i) Any finite Boolean combination of $H_1, \ldots, H_n \in \mathcal{H}_\sigma$ is $\bigsqcup_{r=1}^N K_r$ with $K_r \in \mathcal{H}_\sigma$.*
*(ii) If $H \in \mathcal{H}_\sigma$, then $\mathbb{R}^p \setminus H = \bigsqcup_{r=1}^\infty K_r$ for some $K_r \in \mathcal{H}_\sigma$.*
*(iii) For $H \in \mathcal{H}_\sigma$ and $\delta > 0$, $H = \bigsqcup_{r=1}^N K_r$ for some $K_r \in \mathcal{H}_\sigma$ with $\operatorname{diam} K_r < \delta$.*

*Proof.*

(i) For coordinate endpoints $E_j = \{c_{j,0} < \cdots < c_{j,m_j}\}$ of $H_1, \ldots, H_n$, set $[a, b)_\sigma = [a, b)$ if $\sigma =$ *southwest* and $[a, b)_\sigma = (a, b]$ if $\sigma =$ *northeast*, $\mathcal{P}_j = \{[c_{j,q-1}, c_{j,q})_\sigma : 1 \leq q \leq m_j\}$, and $\mathcal{G} = \{\prod_{j=1}^p P_j : P_j \in \mathcal{P}_j\}$.
Every finite Boolean combination is $\bigsqcup_{G \in \mathcal{G}'} G$, $\mathcal{G}' \subseteq \mathcal{G}$.

(ii) Write $H = \prod_{j=1}^p H_j$ and $D_s = (\prod_{j<s} H_j) \times (\mathbb{R} \setminus H_s) \times \mathbb{R}^{p-s}$; then $\mathbb{R}^p \setminus H = \bigsqcup_{s=1}^p D_s$. Decomposing each one-dimensional ray and $\mathbb{R}$ into countably many bounded $\sigma$-half-open intervals and taking products gives the required union; countability follows from Corollary 6.1.14.

(iii) Choose $m$ with $2^{-m}(\sum_{j=1}^p \ell(H_j)^2)^{1/2} < \delta$. After $m$ bisections per coordinate, each subcell has diameter $< \delta$; disjointness and volume preservation follow coordinatewise.

*Structural Commentary*: Half-open cells behave like grid atoms: finite set operations become finite grid selection.
*Key Insight*: Half-open endpoints prevent overlap while still partitioning boxes cleanly. □

**Definition 3.2.4.** *Translation of a Set*
If $A \subseteq \mathbb{R}^p$ and $t \in \mathbb{R}^p$, the translate of $A$ by $t$ is $t + A := \{t + a : a \in A\}$.

**Proposition 3.2.5.** *Cell Volume Is Translation Invariant*
*Let $I = \prod_{i=1}^p I_i$ be a cell in $\mathbb{R}^p$ and $t = (t_i)_{i=1}^p \in \mathbb{R}^p$.*
*Then $t + I = \prod_{i=1}^p (t_i + I_i)$ is a cell with the same endpoint pattern as $I$, and $\mathrm{vol}_p(t + I) = \mathrm{vol}_p(I)$.*

*Proof.* Let $a_i \leq b_i$ be the endpoints of $I_i$. Then $t_i + I_i$ has endpoints $t_i + a_i \leq t_i + b_i$, unchanged endpoint inclusion, and $\ell(t_i + I_i) = (t_i + b_i) - (t_i + a_i) = \ell(I_i)$. Hence $\mathrm{vol}_p(t + I) = \prod_{i=1}^p \ell(t_i + I_i) = \prod_{i=1}^p \ell(I_i) = \mathrm{vol}_p(I)$.
*Structural Commentary*: Translation moves location but not side lengths.
*Key Insight*: Coordinate volume depends only on side lengths. □

**Definition 3.2.6.** *Elementary Sets and Regularity in $\mathbb{R}^p$*

(i) *Elementary sets*. $\mathcal{E} := \{\bigcup_{r=1}^n I_r : n \in \mathbb{N}\}$, where each $I_r \subseteq \mathbb{R}^p$ is a bounded cell.

(ii) *Elementary volume.* If $A = \bigsqcup_{r=1}^{n} I_r$ with cells $I_r$, set $m(A) := \sum_{r=1}^{n} \mathrm{vol}_p(I_r)$.
(iii) *Regularity.* A nonnegative additive $\mu : \mathcal{E} \to [0, +\infty]$ is regular iff $\forall A \in \mathcal{E}$, $\forall \varepsilon > 0$, $\exists F, G \in \mathcal{E}$ with $F \subseteq A \subseteq G$, $F$ closed, $G$ open, and $\mu(G) - \varepsilon \le \mu(A) \le \mu(F) + \varepsilon$.

$\mathcal{E}$ is a ring; every $A \in \mathcal{E}$ admits a finite disjoint cell decomposition; $m$ is well defined, additive, and regular.

**Perspective 3.2.7.** *Box Volume and Linear Algebra*
For a box $B = \prod_i (a_i, b_i)$ or $\prod_i [a_i, b_i]$, $m(B) = \prod_i (b_i - a_i)$, the diagonal case of determinant volume scaling. Measure extends this product rule beyond boxes.
*Key Insight*: Lebesgue measure starts from determinant-style coordinate volume.

**Proposition 3.2.8.** *Lattice Counts Recover Cell Volume*
*Fix $p \ge 1$. For $q \ge 1$ and $A \subseteq \mathbb{R}^q$, write $N_{\mathbb{Z}}(A) := \#(A \cap \mathbb{Z}^q)$, and for $\lambda > 0$ set $D_\lambda A := \{\lambda x : x \in A\}$. If $I = \prod_{j=1}^{p} I_j$ is a bounded cell in $\mathbb{R}^p$, then $\lim_{\lambda \to +\infty} \lambda^{-p} N_{\mathbb{Z}}(D_\lambda I) = \mathrm{vol}_p(I)$.*

*Proof.* For a bounded interval $J \subseteq \mathbb{R}$ with length $L$, $\max\{0, \lambda L - 1\} \le N_{\mathbb{Z}}(D_\lambda J) \le \lambda L + 1$.
Hence $\lambda^{-1} N_{\mathbb{Z}}(D_\lambda J) \to L$ if $L > 0$, and $N_{\mathbb{Z}}(D_\lambda J) \le 1$ if $L = 0$.
Let $L_j := \ell(I_j)$. Since $D_\lambda I = \prod_j D_\lambda I_j$, $N_{\mathbb{Z}}(D_\lambda I) = \prod_j N_{\mathbb{Z}}(D_\lambda I_j)$.
If every $L_j > 0$, then $\lambda^{-p} N_{\mathbb{Z}}(D_\lambda I) = \prod_j \lambda^{-1} N_{\mathbb{Z}}(D_\lambda I_j) \to \prod_j L_j = \mathrm{vol}_p(I)$.
If some $L_j = 0$, then $N_{\mathbb{Z}}(D_\lambda I) = O(\lambda^{p-1})$, so $\lambda^{-p} N_{\mathbb{Z}}(D_\lambda I) \to 0 = \mathrm{vol}_p(I)$.
*Structural Commentary*: Coordinate lattice density multiplies.
*Key Insight*: Endpoint effects are lower order. □

**Perspective 3.2.9.** *Outer Measure as Overestimation*
Elementary $A = \bigsqcup_{r=1}^{n} I_r$ have $m(A) = \sum_{r=1}^{n} \mathrm{vol}_p(I_r)$.
General $E \subseteq \mathbb{R}^p$, e.g. $D = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \le 1\}$, need not be a finite disjoint cell union.
Replace decompositions by countable cell covers $E \subseteq \bigcup_{k=1}^{\infty} I_k$ and minimize $\sum_{k=1}^{\infty} \mathrm{vol}_p(I_k)$.
*Key Insight*: Outer covers size every $E \subseteq \mathbb{R}^p$ before measurability.

**Definition 3.2.10.** *Cell-Cover Outer Measure*
For $E \subseteq \mathbb{R}^p$, let $\mathcal{C}(E)$ be the countable cell covers $(I_k)_{k \ge 1}$ with $E \subseteq \bigcup_{k=1}^{\infty} I_k$. Define $m_p^*(E) := \inf_{(I_k) \in \mathcal{C}(E)} \sum_{k=1}^{\infty} \mathrm{vol}_p(I_k)$.

Here $\mathcal{C}(E) \neq \emptyset$ since $E \subseteq \bigcup_{j=1}^{\infty} [-j, j]^p$, and sum is nonnegative extended-real series, hence order-independent.

**Proposition 3.2.11.** *Equivalent Covering Families*
*Let $E \subseteq \mathbb{R}^p$, $\delta > 0$, and fix a half-open orientation $\sigma$. Let $\mathcal{C}_{\mathsf{cl}}(E), \mathcal{C}_{\mathsf{op}}(E), \mathcal{C}_\sigma(E), \mathcal{C}_\delta(E) \subseteq \mathcal{C}(E)$ denote covers by closed cells, open cells, $\sigma$-half-open cells, and cells with diameter $< \delta$, respectively. For $\mathcal{F} \subseteq \mathcal{C}(E)$, put $\alpha(\mathcal{F}) := \inf_{(I_k) \in \mathcal{F}} \sum_k \mathrm{vol}_p(I_k)$. Then $m_p^*(E) = \alpha(\mathcal{C}_{\mathsf{cl}}(E)) = \alpha(\mathcal{C}_{\mathsf{op}}(E)) = \alpha(\mathcal{C}_\sigma(E)) = \alpha(\mathcal{C}_\delta(E))$.*

*Proof.* Let $\mathfrak{C} := \{\mathcal{C}_{\mathsf{cl}}(E), \mathcal{C}_{\mathsf{op}}(E), \mathcal{C}_\sigma(E), \mathcal{C}_\delta(E)\}$. Since each $\mathcal{F} \in \mathfrak{C}$ satisfies $\mathcal{F} \subseteq \mathcal{C}(E)$, $\alpha(\mathcal{F}) \ge m_p^*(E)$.
Conversely, fix $(I_k) \in \mathcal{C}(E)$ and $\varepsilon > 0$. The chain $I_k \subseteq \overline{I_k} \subseteq J_k$, with $J_k$ open and $\sum_k \mathrm{vol}_p(J_k) \le \sum_k \mathrm{vol}_p(I_k) + \varepsilon$, followed by disjoint $\sigma$-half-open decomposition and finite $\delta$-subdivision, gives covers in every $\mathcal{F} \in \mathfrak{C}$ of total volume $\le \sum_k \mathrm{vol}_p(I_k) + \varepsilon$. Infimize over $(I_k)$ and let $\varepsilon \downarrow 0$.
*Structural Commentary*: Cover normalizations preserve the infimum.
*Key Insight*: Change endpoints, then refine. □

**Theorem 3.2.12.** *Basic Properties of Cell-Cover Outer Measure*
*For $E, F, E_n \subseteq \mathbb{R}^p$,*

*(i) Null empty set. $0 \le m_p^*(E) \le +\infty$ and $m_p^*(\emptyset) = 0$.*
*(ii) Monotonicity. $E \subseteq F \Longrightarrow m_p^*(E) \le m_p^*(F)$.*
*(iii) Subadditivity. $m_p^*(\bigcup_{n=1}^{\infty} E_n) \le \sum_{n=1}^{\infty} m_p^*(E_n)$.*

*Proof.*

(i) Since all cover sums lie in $[0, +\infty]$, $0 \le m_p^*(E) \le +\infty$; the cover $(\emptyset, \emptyset, \ldots)$ gives $m_p^*(\emptyset) = 0$.
(ii) If $E \subseteq F$, then $\mathcal{C}(F) \subseteq \mathcal{C}(E)$, hence $\inf_{\mathcal{C}(E)} \le \inf_{\mathcal{C}(F)}$.
(iii) If some $m_p^*(E_n) = +\infty$, the claim is trivial.
Otherwise, for $\varepsilon > 0$, choose $(I_k^n)_{k \ge 1} \in \mathcal{C}(E_n)$ with $\sum_k \mathrm{vol}_p(I_k^n) \le m_p^*(E_n) + \varepsilon 2^{-n}$.
Then $(I_k^n)_{(n,k) \in \mathbb{N}^2}$ countable, covers $\bigcup_n E_n$, and $m_p^*(\bigcup_n E_n) \le \sum_{n,k} \mathrm{vol}_p(I_k^n) \le \sum_n m_p^*(E_n) + \varepsilon$.

*Structural Commentary*: Countable subadditivity is proved by merging near-optimal covers.
*Key Insight*: Nonnegative double sums may be scanned in one sequence. □




Figure 193: Positive distance separates cells with $\operatorname{diam} I_k < \delta$.

**Theorem 3.2.13.** *Additivity at Positive Distance*
*Let $A, B \subseteq \mathbb{R}^p$ be nonempty and disjoint, and suppose $\operatorname{dist}(A,B) := \inf\{\|a-b\|_2 : a \in A,\ b \in B\} > 0$. Then $m_p^*(A \cup B) = m_p^*(A) + m_p^*(B)$.*

*Proof.* The inequality $\leq$ follows from Theorem 3.2.12. For $\geq$, set $\delta := \operatorname{dist}(A,B)$. If $m_p^*(A \cup B) = +\infty$, done. Otherwise, given $\varepsilon > 0$, choose $(I_n) \in \mathcal{C}(A \cup B)$ with $\operatorname{diam} I_n < \delta$ and $\sum_n \operatorname{vol}_p(I_n) \leq m_p^*(A \cup B) + \varepsilon$. Put $N_A := \{n : I_n \cap A \neq \emptyset\}$ and $N_B := \{n : I_n \cap B \neq \emptyset\}$. Since $N_A \cap N_B = \emptyset$, $(I_n)_{n \in N_A}$ covers $A$, $(I_n)_{n \in N_B}$ covers $B$, and $m_p^*(A) + m_p^*(B) \leq \sum_{n \in N_A} \operatorname{vol}_p(I_n) + \sum_{n \in N_B} \operatorname{vol}_p(I_n) \leq \sum_n \operatorname{vol}_p(I_n) \leq m_p^*(A \cup B) + \varepsilon$. Let $\varepsilon \downarrow 0$.
*Structural Commentary*: Positive separation lets a sufficiently fine cover be sorted unambiguously.
*Key Insight*: Countable subadditivity becomes equality when no small cell can touch both pieces. □

**Theorem 3.2.14.** *Outer Measure of a Cell*
*If $I$ is a cell in $\mathbb{R}^p$, then $m_p^*(I) = \operatorname{vol}_p(I)$.*

*Proof.* The cover $(I, \emptyset, \ldots)$ gives $m_p^*(I) \leq \operatorname{vol}_p(I)$. For $\geq$, the case $\operatorname{vol}_p(I) = 0$ follows by nonnegativity. Assume $\operatorname{vol}_p(I) > 0$ and fix $\varepsilon > 0$. Choose a closed cell $J \subseteq I$ with $\operatorname{vol}_p(I) - \varepsilon < \operatorname{vol}_p(J)$, taking $J = I$ when $I$ is closed. For any open-cell cover $(G_k)$ of $I$, compactness gives $J \subseteq \bigcup_{k=1}^m G_k$.
Let $\mathcal{Q}$ be the finite rectangular grid determined by the faces of $J, \overline{G_1}, \ldots, \overline{G_m}$. Up to zero-volume boundary atoms, each $Q \in \mathcal{Q}$ with $Q \subseteq J$ lies in some $G_k$, hence $\operatorname{vol}_p(J) \leq \sum_{k=1}^m \operatorname{vol}_p(G_k) \leq \sum_{k=1}^\infty \operatorname{vol}_p(G_k)$.
Infimize over open-cell covers, use Proposition 3.2.11, and let $\varepsilon \downarrow 0$.
*Structural Commentary*: Compact inner cells reduce covers to finite grids.
*Key Insight*: Outer measure is calibrated so that cells keep their original volume. □

**Corollary 3.2.15.** *Outer Measure of Real Intervals*
*Let $J \subseteq \mathbb{R}$ be a nonempty interval, and set $\ell(J) = +\infty$ when $J$ is unbounded. Then $m_1^*(J) = \ell(J)$.*

*Proof.* If $J$ is bounded, then $J$ is a one-dimensional cell, so $m_1^*(J) = \operatorname{vol}_1(J) = \ell(J)$ by Theorem 3.2.14.
If $J$ is unbounded, then for every $N \in \mathbb{N}$ there exists a bounded interval $I_N \subseteq J$ with $\ell(I_N) = N$.
Hence monotonicity and the bounded case give $m_1^*(J) \geq m_1^*(I_N) = N$. Thus $m_1^*(J) = +\infty = \ell(J)$.
*Structural Commentary*: Cells calibrate bounded intervals; monotonicity handles unbounded ones.
*Key Insight*: Infinite length is forced by arbitrarily long bounded subintervals. □

**Theorem 3.2.16.** *Translation Invariance of Outer Measure*
*For $E \subseteq \mathbb{R}^p$ and $t \in \mathbb{R}^p$, $m_p^*(t+E) = m_p^*(E)$.*

*Proof.* The map $\mathcal{C}(E) \to \mathcal{C}(t+E)$, $(I_k) \mapsto (t + I_k)$, is bijective with inverse $(J_k) \mapsto (-t + J_k)$.
By Proposition 3.2.5, $\sum_k \operatorname{vol}_p(t + I_k) = \sum_k \operatorname{vol}_p(I_k)$, so the two infima are equal.
*Structural Commentary*: Translate covers forward and backward.
*Key Insight*: Translation invariance of cell volume lifts directly to translation invariance of outer measure. □

## 3.3 Caratheodory Measurability and Lebesgue Measure

**Definition 3.3.1.** *Outer Measure and $\mu$-Measurable Sets*
Let $\mathcal{E}$ and $\mathcal{E}_{\mathrm{op}}$ denote the elementary and open elementary subsets of $\mathbb{R}^p$, let $\mu : \mathcal{E} \to [0, \infty)$ be additive and regular, and set $A \triangle B := (A \setminus B) \cup (B \setminus A)$.

(i) For $E \subseteq \mathbb{R}^p$, $\mu^*(E) := \inf\{\sum_{n=1}^\infty \mu(A_n) : E \subseteq \bigcup_{n=1}^\infty A_n,\ A_n \in \mathcal{E}_{\mathrm{op}}\}$.
(ii) $d_\mu(A,B) := \mu^*(A \triangle B)$.
(iii) $\mathcal{M}_F(\mu)$ is the class of *finitely $\mu$-measurable* sets: $E \in \mathcal{M}_F(\mu)$ iff $\exists (E_k)_{k \geq 1} \subseteq \mathcal{E}$ with $d_\mu(E_k, E) \to 0$.
(iv) $\mathcal{M}(\mu)$ is the class of *$\mu$-measurable* sets: $E \in \mathcal{M}(\mu)$ iff $E = \bigcup_{k=1}^\infty E_k$ for some $E_k \in \mathcal{M}_F(\mu)$.

**Perspective 3.3.2.** *Outer Measure as Outside Approximation*
Outer measure covers $A$ from outside; $d_\mu(A,B) = \mu^*(A \triangle B)$ measures closeness.
*Key Insight*: Countable subadditivity collects near-optimal covers.

**Theorem 3.3.3.** *Elementary Approximation Extension*
*Let $\mathcal{E}$ be the elementary subsets of $\mathbb{R}^p$, let $\mathcal{E}_{\text{op}} \subseteq \mathcal{E}$ be the open elementary sets, and let $\mu : \mathcal{E} \to [0,\infty)$ be additive and regular. For $E \subseteq \mathbb{R}^p$, set $\mu^*(E) := \inf\{\sum_{n=1}^{\infty} \mu(A_n) : E \subseteq \bigcup_{n=1}^{\infty} A_n,\ A_n \in \mathcal{E}_{\text{op}}\}$; set $d_\mu(A,B) := \mu^*(A \triangle B)$, let $\mathcal{M}_F(\mu)$ be the $d_\mu$-closure of $\mathcal{E}$, and let $\mathcal{M}(\mu) := \{\bigcup_{k=1}^{\infty} E_k : E_k \in \mathcal{M}_F(\mu)\}$. Then:*

*(i) $\mu^*(A) = \mu(A)$ for every $A \in \mathcal{E}$;*
*(ii) $\mu^*$ is countably subadditive on $2^{\mathbb{R}^p}$;*
*(iii) $\mathcal{M}(\mu)$ is a $\sigma$-ring;*
*(iv) $\mu^*|_{\mathcal{M}(\mu)}$ is a measure.*

*Proof.*

(i) Let $A \in \mathcal{E}$. For $\varepsilon > 0$, regularity gives $G \in \mathcal{E}_{\text{op}}$ with $A \subseteq G$ and $\mu(G) \le \mu(A) + \varepsilon$, so $\mu^*(A) \le \mu(A)$. Conversely, choose closed elementary $F \subseteq A$ with $\mu(F) \ge \mu(A) - \varepsilon$. Every open elementary cover of $A$ covers $F$; compactness gives a finite subcover, whose total $\mu$-mass is at least $\mu(F)$. Taking infima and letting $\varepsilon \downarrow 0$ gives $\mu(A) \le \mu^*(A)$.

(ii) Let $E_j \subseteq \mathbb{R}^p$ for $j \ge 1$, and let $\varepsilon > 0$. For each $j$, choose an open elementary cover $(A_{j,n})_{n\ge1}$ of $E_j$ with $\sum_{n=1}^{\infty} \mu(A_{j,n}) \le \mu^*(E_j) + \varepsilon 2^{-j}$. Then $(A_{j,n})_{j,n}$ covers $\bigcup_{j=1}^{\infty} E_j$, so $\mu^*(\bigcup_j E_j) \le \sum_j \mu^*(E_j) + \varepsilon$; let $\varepsilon \downarrow 0$.

(iii) If $A_k \to A$ and $B_k \to B$ in $d_\mu$, with $A_k, B_k \in \mathcal{E}$, then $d_\mu(A_k \diamond B_k, A \diamond B) \le d_\mu(A_k, A) + d_\mu(B_k, B)$ for $\diamond \in \{\cup, \cap, \setminus\}$. Hence $\mathcal{M}_F(\mu)$ is ring. Countable unions give $\sigma$-ring since difference distributes over them.

(iv) If $A = \bigsqcup_{j=1}^{\infty} A_j$ with $A_j \in \mathcal{M}_F(\mu)$, subadditivity gives $\mu^*(A) \le \sum_j \mu^*(A_j)$.
Finite additivity on $\mathcal{M}_F(\mu)$ gives $\mu^*(A) \ge \sum_{j=1}^{N} \mu^*(A_j)$ for every $N$, hence $\mu^*(A) \ge \sum_j \mu^*(A_j)$.
Disjointize countable $\mathcal{M}_F(\mu)$-decompositions.

*Structural Commentary*: Elementary sets are completed under the distance measured by symmetric difference.
*Key Insight*: Approximate by elementary sets, prove identities there, then pass through $d_\mu(A,B)$. □

**Definition 3.3.4.** *Extended and Lebesgue Measure*
Let $\mu^*$ be the outer measure induced by $\mu$ and let $\mathcal{M}(\mu)$ be its measurable class.

(i) The *extended measure* of $\mu$ is $\mu_{\text{ext}} := \mu^*|_{\mathcal{M}(\mu)}$, usually denoted again by $\mu$.
(ii) If $\mu = m$ is elementary volume on $\mathbb{R}^p$, then $m^*|_{\mathcal{M}(m)}$ is *Lebesgue measure* on $\mathbb{R}^p$.

**Definition 3.3.5.** *Box-Cover Outer Measure Model*
Fix $n \ge 1$.

(i) $\mathcal{B}_n := \{\prod_{i=1}^{n}(a_i,b_i) : a_i, b_i \in \mathbb{R},\ a_i \le b_i\ \forall i\}$ is the class of *open boxes*.
For $B = \prod_{i=1}^{n}(a_i,b_i)$, $\operatorname{vol}_n(B) := \prod_{i=1}^{n}(b_i - a_i)$.
(ii) For $\Omega \subseteq \mathbb{R}^n$, $m_n^*(\Omega) := \inf\{\sum_{j\in J} \operatorname{vol}_n(B_j) : \Omega \subseteq \bigcup_{j\in J} B_j,\ B_j \in \mathcal{B}_n,\ |J| \le \aleph_0\}$.




Figure 194: Cover $A \subseteq \bigcup_k B_k$ by squares; minimize $\sum_k |B_k|$.

**Method 3.3.6.** *Lebesgue Measure Construction*

(i) *Cover from outside*
For $A \subseteq \mathbb{R}^n$, define $m_n^*(A) := \inf\left\{\sum_{j\in J} \operatorname{vol}_n(B_j) : (B_j)_{j\in J} \text{ covers } A,\ J \text{ at most countable}\right\}$.
This is defined for every set, but it is only countably subadditive in general.
(ii) *Keep the clean splitters*
Call $A$ measurable when every test set $X \subseteq \mathbb{R}^n$ splits additively: $m_n^*(X) = m_n^*(X \cap A) + m_n^*(X \cap A^c)$.
These measurable sets form a $\sigma$-algebra.
(iii) *Restrict the outer measure*
Define Lebesgue measure on measurable sets by $m(A) := m_n^*(A)$.
On this restricted class, $m$ is countably additive.

*Key Insight*: Outer measure sizes all sets; clean splitting restores additivity.

**Definition 3.3.7.** *Lebesgue $\sigma$-Algebra via Caratheodory*

(i) $\mathcal{L}^n := \{E \subseteq \mathbb{R}^n : E \text{ is } m_n^*\text{-measurable}\}$ is the *Lebesgue $\sigma$-algebra*.
(ii) $m := m_n^*|_{\mathcal{L}^n}$ is *Lebesgue measure*; for $E \in \mathcal{L}^n$, write $m(E) = m_n^*(E)$.

**Proposition 3.3.8.** *Outer-Null Sets Are Lebesgue Measurable*
*If $Z \subseteq \mathbb{R}^n$ and $m_n^*(Z) = 0$, then:*

*(i)* $Z \in \mathcal{L}^n$;
*(ii)* $m(Z) = 0$;
*(iii)* $\mathcal{P}(Z) \subseteq \mathcal{L}^n$ *and* $m(W) = 0$ *for every* $W \subseteq Z$.

*Proof.*

(i) For $A \subseteq \mathbb{R}^n$, monotonicity gives $m_n^*(A \cap Z) = 0$, so $m_n^*(A) \leq m_n^*(A \cap Z) + m_n^*(A \cap Z^c) = m_n^*(A \cap Z^c) \leq m_n^*(A)$; hence $Z$ satisfies Caratheodory's criterion.
(ii) Since $Z \in \mathcal{L}^n$, $m(Z) = m_n^*(Z) = 0$.
(iii) If $W \subseteq Z$, then $m_n^*(W) \leq m_n^*(Z) = 0$; applying (i)–(ii) to $W$ gives $W \in \mathcal{L}^n$ and $m(W) = 0$.

*Structural Commentary*: Lebesgue measure is complete because outer-null subsets split every test set trivially.
*Key Insight*: A zero piece cannot disturb additivity. □

**Example 3.3.9.** *Countable Real Sets Are Outer-Null*
Let $C \subseteq \mathbb{R}$ be finite or countable, and choose $(c_k)_{k\geq 1}$ with $C \subseteq \{c_k : k \geq 1\}$. For $\varepsilon > 0$, choose open intervals $I_k \ni c_k$ with $\ell(I_k) < \varepsilon 2^{-k}$. Then $C \subseteq \bigcup_{k\geq 1} I_k$, so $0 \leq m_1^*(C) \leq \sum_{k=1}^{\infty} \ell(I_k) < \varepsilon$. Since $\varepsilon > 0$ is arbitrary, $m_1^*(C) = 0$. By Proposition 3.3.8, $C \in \mathcal{L}^1$ and $m(C) = 0$.
*Key Insight*: A countable list can spend a rapidly decreasing measure budget around its points.




Figure 195: Inner cells $I_k$ exhaust $I$; the collar $I \setminus I_k$ has vanishing volume.

**Proposition 3.3.10.** *Cells Are Lebesgue Measurable*
*Let $I \subseteq \mathbb{R}^n$ be a coordinate cell. Then $I \in \mathcal{L}^n$ and $m(I) = \mathrm{vol}_n(I)$.*

*Proof.* If $\mathrm{vol}_n(I) = 0$, then $m_n^*(I) = 0$ by Theorem 3.2.14, so Proposition 3.3.8 gives $I \in \mathcal{L}^n$ and $m(I) = 0$.
Assume $\mathrm{vol}_n(I) > 0$. For small $r > 0$, let $I_r$ be the inward $r$-shrink of $I$.
Then $\mathrm{dist}(I_r, \mathbb{R}^n \setminus I) > 0$ and $\eta_r := m_n^*(I \setminus I_r) \to 0$ by finite slab covers.
Fix $A \subseteq \mathbb{R}^n$ with $m_n^*(A) < +\infty$. Positive-distance additivity gives $m_n^*(A) \geq m_n^*(A \cap I_r) + m_n^*(A \setminus I)$.
Since $A \cap I \subseteq (A \cap I_r) \cup (I \setminus I_r)$, $m_n^*(A \cap I) \leq m_n^*(A \cap I_r) + \eta_r$; hence $m_n^*(A) \geq m_n^*(A \cap I) + m_n^*(A \setminus I) - \eta_r$.
Letting $r \downarrow 0$ gives the finite-measure test inequality, so Lemma 3.1.4 gives $I \in \mathcal{L}^n$.
Theorem 3.2.14 gives $m(I) = m_n^*(I) = \mathrm{vol}_n(I)$.
*Structural Commentary*: Cells are the bridge between coordinate volume and Lebesgue measure.
*Key Insight*: Shrink, split, let the collar vanish. □

## 3.4 Caratheodory-Hahn Extension

**Definition 3.4.1.** *Premeasures on Set Families*
Let $\emptyset \neq \mathcal{S} \subseteq \mathcal{P}(X)$ and $\mu : \mathcal{S} \to [0, +\infty]$.

(i) $\mu$ is *finitely additive on $\mathcal{S}$* if $\mu(\bigcup_{k=1}^{n} E_k) = \sum_{k=1}^{n} \mu(E_k)$ $\forall$ finite pairwise disjoint family $E_k \in \mathcal{S}$ with $\bigcup_{k=1}^{n} E_k \in \mathcal{S}$.
(ii) $\mu$ is *countably monotone on $\mathcal{S}$* if, for all $E, E_1, E_2, \ldots \in \mathcal{S}$ with $E \subseteq \bigcup_{k=1}^{\infty} E_k$, $\mu(E) \leq \sum_{k=1}^{\infty} \mu(E_k)$.
(iii) $\mu$ is a *premeasure on $\mathcal{S}$* if $\mu$ is finitely additive on $\mathcal{S}$, countably monotone on $\mathcal{S}$, and $\emptyset \in \mathcal{S} \Rightarrow \mu(\emptyset) = 0$.

*Structural Commentary*: No closure property of $\mathcal{S}$ is assumed; additivity is imposed only for finite disjoint unions which remain in $\mathcal{S}$, and countable monotonicity only for countable $\mathcal{S}$-covers of members of $\mathcal{S}$.

**Proposition 3.4.2.** *Necessary Conditions for Extension*
*Let $\emptyset \neq \mathcal{S} \subseteq \mathcal{P}(X)$, let $\mu : \mathcal{S} \to [0, +\infty]$, let $\mu^*$ be the outer measure induced by $\mu$, and put $\overline{\mu} := \mu^*|_{\mathcal{M}_{\mu^*}}$. If $\mathcal{S} \subseteq \mathcal{M}_{\mu^*}$ and $\overline{\mu}(E) = \mu(E)$ for all $E \in \mathcal{S}$, then the following hold:*

*(i)* $\emptyset \in \mathcal{S} \Rightarrow \mu(\emptyset) = 0$.
*(ii)* $\mu$ *is finitely additive on* $\mathcal{S}$.
*(iii)* $\mu$ *is countably monotone on* $\mathcal{S}$.

*Hence $\mu$ is a premeasure on $\mathcal{S}$.*

*Proof.* By the extension hypotheses, $\mu(F) = \overline{\mu}(F) = \mu^*(F)$ for every $F \in \mathcal{S}$.

(i) If $\emptyset \in \mathcal{S}$, then $\mu(\emptyset) = \overline{\mu}(\emptyset) = 0$.

(ii) Let $n \geq 1$, let $E_1, \ldots, E_n \in \mathcal{S}$ be pairwise disjoint, and assume $F := \bigcup_{k=1}^{n} E_k \in \mathcal{S}$.

Since $\overline{\mu}$ is a measure, $\mu(F) = \overline{\mu}(F) = \sum_{k=1}^{n} \overline{\mu}(E_k) = \sum_{k=1}^{n} \mu(E_k)$.

(iii) Let $E, E_1, E_2, \ldots \in \mathcal{S}$ and $E \subseteq \bigcup_{k=1}^{\infty} E_k$. By monotonicity and countable subadditivity of $\mu^*$,

$\mu(E) = \mu^*(E) \leq \mu^*(\bigcup_{k=1}^{\infty} E_k) \leq \sum_{k=1}^{\infty} \mu^*(E_k) = \sum_{k=1}^{\infty} \mu(E_k)$.

*Structural Commentary*: The measure extension handles finite disjoint unions; $\mu^*$ handles countable covers.
*Key Insight*: Outer subadditivity supplies the inequality even when $\bigcup_k E_k \notin \mathcal{S}$. □

**Definition 3.4.3.** *Relative Complements and Semirings*
Let $\emptyset \neq \mathcal{S} \subseteq \mathcal{P}(X)$.

(i) $\mathcal{S}$ is *closed under relative complements* if $A, B \in \mathcal{S} \Rightarrow A \setminus B \in \mathcal{S}$.

(ii) $\mathcal{S}$ is a *semiring of sets* if, for all $A, B \in \mathcal{S}$, $A \cap B \in \mathcal{S}$ and $A \setminus B = \bigsqcup_{j=1}^{n} C_j$ for some $n \geq 0$ and $C_j \in \mathcal{S}$.

*Structural Commentary*: Relative-complement closure gives $\emptyset = A \setminus A \in \mathcal{S}$ and $A \cap B = A \setminus (A \setminus B) \in \mathcal{S}$.

**Theorem 3.4.4.** *Extension from Relative-Complement Families*
*Let $\emptyset \neq \mathcal{S} \subseteq \mathcal{P}(X)$ be closed under relative complements, let $\mu : \mathcal{S} \to [0, +\infty]$ be a premeasure, and let $\mu^*$ be the outer measure induced by $\mu$. Then:*

*(i)* $\mu^*(E) = \mu(E)$ *for every* $E \in \mathcal{S}$.
*(ii)* $\mathcal{S} \subseteq \mathcal{M}_{\mu^*}$.

*Proof.*

(i) Fix $E \in \mathcal{S}$. The one-set cover gives $\mu^*(E) \leq \mu(E)$. If $(E_k)_{k \geq 1} \subseteq \mathcal{S}$ covers $E$, then countable monotonicity gives $\mu(E) \leq \sum_{k=1}^{\infty} \mu(E_k)$; taking the infimum over all such covers gives $\mu(E) \leq \mu^*(E)$.

(ii) Fix $A \in \mathcal{S}$. By Lemma 3.1.4, it suffices to prove $\mu^*(B \cap A) + \mu^*(B \setminus A) \leq \mu^*(B)$ for every $B \subseteq X$ with $\mu^*(B) < +\infty$. Let $\varepsilon > 0$ and choose an $\mathcal{S}$-cover $(E_k)_{k \geq 1}$ of $B$ with $\sum_{k=1}^{\infty} \mu(E_k) < \mu^*(B) + \varepsilon$.

For each $k$, relative-complement closure gives $E_k \setminus A \in \mathcal{S}$ and $E_k \cap A = E_k \setminus (E_k \setminus A) \in \mathcal{S}$. By finite additivity, $\mu(E_k) = \mu(E_k \cap A) + \mu(E_k \setminus A)$. Since $(E_k \cap A)_{k \geq 1}$ covers $B \cap A$ and $(E_k \setminus A)_{k \geq 1}$ covers $B \setminus A$, $\mu^*(B \cap A) + \mu^*(B \setminus A) \leq \sum_{k=1}^{\infty} \mu(E_k \cap A) + \sum_{k=1}^{\infty} \mu(E_k \setminus A) = \sum_{k=1}^{\infty} \mu(E_k) < \mu^*(B) + \varepsilon$. Let $\varepsilon \downarrow 0$.

*Structural Commentary*: Countable monotonicity identifies $\mu$ with $\mu^*$; finite additivity splits covers.
*Key Insight*: Relative complements turn one cover into inside/outside covers. □

**Remark 3.4.5.** *Two Roles of a Premeasure*
In Theorem 3.4.4, finite additivity is used to split each covering set across $A$ and $A^c$, while countable monotonicity is used to force $\mu^*(E) = \mu(E)$ on the original family.
*Key Insight*: Measurability comes from finite splitting; extension equality comes from countable cover control.

**Proposition 3.4.6.** *Finite-Union Extension from a Semiring*
*Let $\mathcal{S}$ be a semiring of subsets of $X$, and set $\mathcal{S}' := \{\bigsqcup_{k=1}^{n} A_k : n \geq 0,\ A_1, \ldots, A_n \in \mathcal{S}\}$. If $\mu : \mathcal{S} \to [0, +\infty]$ is a premeasure, define $\mu' : \mathcal{S}' \to [0, +\infty]$ by $\mu'(\bigsqcup_{k=1}^{n} A_k) := \sum_{k=1}^{n} \mu(A_k)$ for any disjoint representation. Then:*

*(i)* $\mathcal{S}'$ *is closed under relative complements.*
*(ii)* $\mu'$ *is well-defined and extends* $\mu$.

*(iii)* $\mu'$ *is a premeasure on* $\mathcal{S}'$.
*(iv)* $\mu'$ *is the unique premeasure on* $\mathcal{S}'$ *extending* $\mu$.

*Proof.*

(i) Let $E = \bigsqcup_{i=1}^{r} A_i$ and $F = \bigsqcup_{j=1}^{s} B_j$ in $\mathcal{S}'$.
For each $i$, repeated semiring differences decompose $A_i \setminus F$ into finitely many disjoint $\mathcal{S}$-sets.
These decompositions, over $i = 1, \dots, r$, form a finite disjoint $\mathcal{S}$-decomposition of $E \setminus F$, so $E \setminus F \in \mathcal{S}'$.

(ii) If $E = \bigsqcup_{i=1}^{r} A_i = \bigsqcup_{j=1}^{s} B_j$, then $A_i = \bigsqcup_{j=1}^{s} (A_i \cap B_j)$ and $B_j = \bigsqcup_{i=1}^{r} (A_i \cap B_j)$ with all intersections in $\mathcal{S}$.
Finite additivity of $\mu$ gives $\sum_{i=1}^{r} \mu(A_i) = \sum_{i=1}^{r} \sum_{j=1}^{s} \mu(A_i \cap B_j) = \sum_{j=1}^{s} \mu(B_j)$, so $\mu'$ is well-defined.
If $A \in \mathcal{S}$, the one-piece representation gives $\mu'(A) = \mu(A)$.

(iii) Since $\emptyset$ is the empty union, $\mu'(\emptyset) = 0$.
Finite additivity of $\mu'$ follows by refining finitely many disjoint $\mathcal{S}'$-sets to their disjoint $\mathcal{S}$-pieces.
For countable monotonicity, let $E = \bigsqcup_{i=1}^{r} A_i \in \mathcal{S}'$, let $F_m = \bigsqcup_{j=1}^{N_m} B_{mj} \in \mathcal{S}'$, assume $E \subseteq \bigcup_{m=1}^{\infty} F_m$.
For each $i$, $A_i \subseteq \bigcup_{m,j} (A_i \cap B_{mj})$ with $A_i \cap B_{mj} \in \mathcal{S}$, so $\mu(A_i) \le \sum_{m,j} \mu(A_i \cap B_{mj})$.
Summing over $i$ gives $\mu'(E) \le \sum_m \sum_j \sum_i \mu(A_i \cap B_{mj})$.
For fixed $m, j$, repeated semiring differences decompose $B_{mj} \setminus \bigcup_i (A_i \cap B_{mj})$ into finitely many disjoint $\mathcal{S}$-sets, so finite additivity gives $\sum_i \mu(A_i \cap B_{mj}) \le \mu(B_{mj})$. Thus $\mu'(E) \le \sum_m \sum_j \mu(B_{mj}) = \sum_m \mu'(F_m)$.

(iv) If $\nu$ is a premeasure on $\mathcal{S}'$ extending $\mu$, then finite additivity gives
$\nu(\bigsqcup_{k=1}^{n} A_k) = \sum_{k=1}^{n} \nu(A_k) = \sum_{k=1}^{n} \mu(A_k) = \mu'(\bigsqcup_{k=1}^{n} A_k)$ for every disjoint representation. Hence $\nu = \mu'$.

*Structural Commentary*: Finite disjoint unions close a semiring under relative complements.
*Key Insight*: All estimates reduce to intersections of semiring pieces. □

**Theorem 3.4.7.** *Caratheodory-Hahn Extension from a Semiring*
*Let* $\mathcal{S}$ *be a semiring of subsets of* $X$, *let* $\mu : \mathcal{S} \to [0, +\infty]$ *be a premeasure, let* $\mu^*$ *be the outer measure induced by* $\mu$, *and set* $\overline{\mu} := \mu^*|_{\mathcal{M}_{\mu^*}}$. *Then:*

*(i)* $\mathcal{S} \subseteq \mathcal{M}_{\mu^*}$ *and* $\overline{\mu}(A) = \mu(A)$ *for every* $A \in \mathcal{S}$.
*(ii) If* $X = \bigcup_{k \ge 1} S_k$ *with* $S_k \in \mathcal{S}$ *and* $\mu(S_k) < +\infty$, *then* $\overline{\mu}$ *is* $\sigma$*-finite.*
*(iii) Under the hypothesis in (ii), if* $\nu$ *is a measure on* $\mathcal{M}_{\mu^*}$ *with* $\nu(A) = \mu(A)$ *for every* $A \in \mathcal{S}$, *then* $\nu = \overline{\mu}$.

*Proof.*

(i) Let $(\mathcal{S}', \mu')$ be the finite-union extension from Proposition 3.4.6. The $\mathcal{S}$- and $\mathcal{S}'$-cover infima coincide, so $\mu^*$ is also induced by $\mu'$. Theorem 3.4.4 gives $\mathcal{S}' \subseteq \mathcal{M}_{\mu^*}$ and $\mu^*|_{\mathcal{S}'} = \mu'$. Since $\mu'|_{\mathcal{S}} = \mu$, the claim follows.

(ii) Assume $X = \bigcup_{k \ge 1} S_k$, $S_k \in \mathcal{S}$, $\mu(S_k) < +\infty$. By (i), each $S_k \in \mathcal{M}_{\mu^*}$, $\overline{\mu}(S_k) = \mu(S_k) < +\infty$, so $\overline{\mu}$ is $\sigma$-finite.

(iii) Assume (ii), and let $\nu$ be a measure on $\mathcal{M}_{\mu^*}$ extending $\mu$. By finite additivity, $\nu = \overline{\mu} = \mu'$ on $\mathcal{S}'$.
Hence every $\mathcal{S}'$-cover of $E \in \mathcal{M}_{\mu^*}$ gives $\nu(E) \le$ its $\mu'$-mass, so $\nu(E) \le \overline{\mu}(E)$. Put $H_n := \bigcup_{k=1}^{n} S_k \in \mathcal{S}'$.
Then $H_n \uparrow X$ and $\nu(H_n) = \overline{\mu}(H_n) = \mu'(H_n) < +\infty$.
Applying $\nu \le \overline{\mu}$ to $E \cap H_n$ and $H_n \setminus E$ forces $\nu(E \cap H_n) = \overline{\mu}(E \cap H_n)$, because both measures split the same finite mass $H_n$. Since $E \cap H_n \uparrow E$, continuity from below gives $\nu(E) = \overline{\mu}(E)$.

*Structural Commentary*: Pass to finite disjoint unions, then exhaust by finite-mass pieces.
*Key Insight*: $\mathcal{S}'$ supplies relative complements and the cover infimum for uniqueness. □




Figure 196: Caratheodory-Hahn: pass to $\mathcal{S}'$, form $\mu^*$ by cover infima, then $\overline{\mu} := \mu^*|_{\mathcal{M}_{\mu^*}}$ with $\overline{\mu}|_{\mathcal{S}} = \mu$.

**Corollary 3.4.8.** *Agreement on a* $\sigma$*-Finite Semiring*
*Let* $\mathcal{S}$ *be a semiring of subsets of* $X$, *let* $\mathcal{B} := \sigma(\mathcal{S})$, *and let* $\nu_1, \nu_2$ *be measures on* $\mathcal{B}$ *with* $\nu_1|_{\mathcal{S}} = \nu_2|_{\mathcal{S}}$.
*If* $X = \bigcup_{k \ge 1} S_k$ *for some* $S_k \in \mathcal{S}$ *with* $\nu_1(S_k) = \nu_2(S_k) < +\infty$ *for all* $k$, *then* $\nu_1 = \nu_2$.

*Proof.* Let $\mathcal{S}'$ be the finite-disjoint-union closure of $\mathcal{S}$. Finite additivity gives $\nu_1 = \nu_2$ on $\mathcal{S}'$.
Put $H_n := \bigcup_{k=1}^{n} S_k \in \mathcal{S}'$; then $H_n \uparrow X$ and $\nu_i(H_n) < +\infty$. For fixed $n$, $\mathcal{D}_n := \{E \in \mathcal{B} : \nu_1(E \cap H_n) = \nu_2(E \cap H_n)\}$ is a Dynkin class containing the $\pi$-system $\mathcal{S}$, since $A \cap H_n \in \mathcal{S}'$ for $A \in \mathcal{S}$. Hence $\mathcal{D}_n = \mathcal{B}$.
Thus $\nu_1(E \cap H_n) = \nu_2(E \cap H_n)\ \forall E \in \mathcal{B}$ and $n \geq 1$; since $E \cap H_n \uparrow E$, continuity from below gives $\nu_1(E) = \nu_2(E)$.
*Structural Commentary*: Localize to finite-mass $H_n$, then apply $\pi$-$\lambda$.
*Key Insight*: Agreement on $\mathcal{S}$ propagates through $\mathcal{S}'$ and the exhaustion. □

**Example 3.4.9.** *Bounded Intervals and Rectangles Are Semirings*
Let $\mathcal{I}$ be the bounded intervals of $\mathbb{R}$, and let $\mathcal{R}_n := \{I_1 \times \cdots \times I_n : I_j \in \mathcal{I}\}$.

(i) $\mathcal{I}$ is a semiring: if $I, J \in \mathcal{I}$, then $I \cap J \in \mathcal{I}$ and $I \setminus J$ is a disjoint union of at most two members of $\mathcal{I}$.
(ii) $\mathcal{R}_n$ is a semiring in $\mathbb{R}^n$: intersections are coordinatewise products, and $R \setminus Q$ decomposes by coordinate cuts into finitely many disjoint rectangles in $\mathcal{R}_n$.

Thus length on $\mathcal{I}$ and rectangular volume on $\mathcal{R}_n$ fit the Caratheodory-Hahn extension pattern.

## 3.5 Metric Outer Measures and Hausdorff Measure

Metric outer measures are outer measures whose additivity recognizes positive separation. They abstract the positive-distance additivity of Euclidean cell-cover outer measure from Theorem 3.2.13.

**Definition 3.5.1.** *Metric/Caratheodory Outer Measure*
Let $(X, d)$ be a metric space and let $\omega : 2^X \to [0, +\infty]$ be an outer measure.

(i) For nonempty $A, B \subseteq X$, $\operatorname{dist}(A, B) := \inf\{d(a, b) : a \in A,\ b \in B\}$.
(ii) $\omega$ is *metric outer measure* if $\operatorname{dist}(A, B) > 0 \Rightarrow \omega(A \cup B) = \omega(A) + \omega(B)$ $\forall$ nonempty $A, B \subseteq X$.

*Key Insight*: Positive metric separation forces outer-measure additivity.

**Definition 3.5.2.** *Separation by a Real-Valued Function*
Let $X$ be a set and let $\varphi : X \to \mathbb{R}$. Sets $A, B \subseteq X$ are *$\varphi$-separated* if, for some $r < s$, one of $A, B$ is contained in $\{\varphi \leq r\}$ and the other is contained in $\{\varphi \geq s\}$.

**Proposition 3.5.3.** *Separation by One Real-Valued Function*
*Let $X$ be a set, let $\omega : 2^X \to [0, +\infty]$ be an outer measure, and let $\varphi : X \to \mathbb{R}$. If $\omega(A \cup B) = \omega(A) + \omega(B)$ for every $\varphi$-separated pair $A, B \subseteq X$, then $\varphi : (X, \mathcal{M}_\omega) \to (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ is measurable.*

*Proof.* Fix $t \in \mathbb{R}$, put $E_t := \{\varphi > t\}$, and fix $A \subseteq X$ with $\omega(A) < +\infty$. Set $B := A \cap E_t$ and $C := A \setminus E_t$.
For $n \geq 1$, let $B_n := A \cap \{\varphi > t + \frac{1}{n}\}$, $B_0 := \emptyset$, and $R_n := B_n \setminus B_{n-1}$.
Same-parity strips are pairwise $\varphi$-separated, so $\sum_{n \geq 1} \omega(R_n) < +\infty$. Given $\varepsilon > 0$, choose $n$ with $\sum_{k > n} \omega(R_k) < \varepsilon$.
Since $B \subseteq B_n \cup \bigcup_{k > n} R_k$ and $B_n, C$ are $\varphi$-separated, $\omega(B) + \omega(C) \leq \omega(B_n \cup C) + \varepsilon \leq \omega(A) + \varepsilon$.
Let $\varepsilon \downarrow 0$, then $\omega(A) \geq \omega(A \cap E_t) + \omega(A \setminus E_t)$; Lemma 3.1.4 gives $E_t \in \mathcal{M}_\omega$ for every $t$, hence $\varphi$ is measurable.
*Structural Commentary*: Measurability reduces to a finite-mass split of each level set.
*Key Insight*: Summable level strips leave an arbitrarily small boundary layer. □

**Theorem 3.5.4.** *Metric Outer Measures Measure Borel Sets*
*Let $(X, d)$ be a metric space and let $\omega : 2^X \to [0, +\infty]$ be a metric outer measure. Then:*

*(i) Every closed $F \subseteq X$ belongs to $\mathcal{M}_\omega$.*
*(ii) $\mathcal{B}(X) \subseteq \mathcal{M}_\omega$.*

*Proof.*

(i) The case $F = \emptyset$ is immediate. For nonempty closed $F$, set $\varphi(x) := \operatorname{dist}(\{x\}, F)$. Proposition 5.1.21 gives $\varphi$ 1-Lipschitz and $F = \varphi^{-1}(\{0\})$, so $\varphi$-separated pairs have positive distance. Metric additivity verifies Proposition 3.5.3; hence $\varphi$ is $\mathcal{M}_\omega$-measurable and $F \in \mathcal{M}_\omega$.
(ii) By (i), all closed sets lie in the $\sigma$-algebra $\mathcal{M}_\omega$; since they generate $\mathcal{B}(X)$, $\mathcal{B}(X) \subseteq \mathcal{M}_\omega$.

*Structural Commentary*: Distance functions turn closed sets into real-valued separation tests.
*Key Insight*: Closed measurability plus generation gives Borel measurability. □

**Definition 3.5.5.** *Hausdorff Outer Measure*
Let $(X, d)$ be a metric space, $\alpha > 0$, $\varepsilon > 0$.

(i) $\mathcal{H}^*_{\alpha,\varepsilon}(\emptyset) := 0$; $\mathcal{H}^*_{\alpha,\varepsilon}(E) := \inf\{\sum_{k=1}^{\infty} (\operatorname{diam} A_k)^\alpha : E \subseteq \bigcup_{k=1}^{\infty} A_k,\ \emptyset \neq A_k \subseteq X,\ \operatorname{diam} A_k < \varepsilon\}$, nonempty $E \subseteq X$
(ii) If $0 < \varepsilon_1 < \varepsilon_2$, then $\mathcal{H}^*_{\alpha,\varepsilon_1}(E) \geq \mathcal{H}^*_{\alpha,\varepsilon_2}(E)$.

(iii) $\mathcal{H}^*_\alpha(E) := \sup_{\varepsilon>0} \mathcal{H}^*_{\alpha,\varepsilon}(E) = \lim_{\varepsilon\downarrow 0} \mathcal{H}^*_{\alpha,\varepsilon}(E)$.

*Key Insight*: $\mathcal{H}^*_\alpha$ is the limiting cost of countable small-diameter covers.




Figure 197: $\operatorname{dist}(E,F) = \delta$ and $\operatorname{diam} A_k < \delta$: each $A_k$ meets at most one of $E, F$.

**Theorem 3.5.6.** *Hausdorff Measure on Borel Sets*
*Let $(X,d)$ be a metric space and let $\alpha > 0$. Then:*

*(i) $\mathcal{H}^*_\alpha : 2^X \to [0,+\infty]$ is a metric outer measure.*
*(ii) $\mathcal{H}_\alpha := \mathcal{H}^*_\alpha|_{\mathcal{B}(X)}$ is a measure on $\mathcal{B}(X)$, called the $\alpha$-dimensional Hausdorff measure on $X$.*

*Proof.*

(i) For each $\varepsilon > 0$, $\mathcal{H}^*_{\alpha,\varepsilon}(\emptyset) = 0$, monotonicity is immediate from admissible covers, and countable subadditivity follows by joining admissible $\varepsilon$-covers; hence $\mathcal{H}^*_\alpha(\bigcup_i E_i) = \sup_{\varepsilon>0} \mathcal{H}^*_{\alpha,\varepsilon}(\bigcup_i E_i) \le \sum_i \mathcal{H}^*_\alpha(E_i)$.
Thus $\mathcal{H}^*_\alpha$ is an outer measure. If $\operatorname{dist}(E,F) = \delta > 0$ and $0 < \varepsilon < \delta$, every admissible $\varepsilon$-cover of $E \cup F$ splits into disjoint subcovers of $E$ and $F$, so $\mathcal{H}^*_{\alpha,\varepsilon}(E \cup F) \ge \mathcal{H}^*_{\alpha,\varepsilon}(E) + \mathcal{H}^*_{\alpha,\varepsilon}(F)$.
Letting $\varepsilon \downarrow 0$ and using subadditivity gives metric additivity.
(ii) By (i), $\mathcal{H}^*_\alpha$ is a metric outer measure; Theorem 3.5.4 gives $\mathcal{B}(X) \subseteq \mathcal{M}_{\mathcal{H}^*_\alpha}$.
Since $\mathcal{H}^*_\alpha|_{\mathcal{M}_{\mathcal{H}^*_\alpha}}$ is a measure, its restriction $\mathcal{H}^*_\alpha|_{\mathcal{B}(X)} = \mathcal{H}_\alpha$ is a measure on $\mathcal{B}(X)$.

*Structural Commentary*: Cover infima give the outer measure; separated cover splitting gives metric additivity.
*Key Insight*: Small-diameter covers cannot straddle positive gaps. □

**Remark 3.5.7.** *One-Dimensional Calibration*
On $\mathbb{R}$, interval covers give $\mathcal{H}^*_1((a,b]) \le b-a$, and every diameter cover of $(a,b]$ enlarges to interval covers with arbitrarily small extra total length, giving $\mathcal{H}^*_1((a,b]) \ge b-a$. Hence $\mathcal{H}^*_1((a,b]) = b-a$, and uniqueness of the Borel extension from half-open intervals gives $\mathcal{H}_1 = m$ on $\mathcal{B}(\mathbb{R})$.

**Example 3.5.8.** *Finite Sets Have Zero Positive-Dimensional Hausdorff Measure*
If $E = \{x_1,\dots,x_N\} \subseteq X$ and $\alpha > 0$, then $\mathcal{H}^*_\alpha(E) = 0$: for every $\varepsilon > 0$, the singleton cover $E \subseteq \bigcup_{j=1}^N \{x_j\}$ is admissible and has total $\alpha$-content $\sum_{j=1}^N (\operatorname{diam}\{x_j\})^\alpha = 0$.
*Key Insight*: Finite sets are covered by zero-diameter singletons.

**Proposition 3.5.9.** *Hausdorff Measures Decrease Through Dimension*
*Let $(X,d)$ be a metric space, let $E \subseteq X$, and let $0 < \alpha < \beta$. If $\mathcal{H}^*_\alpha(E) < +\infty$, then $\mathcal{H}^*_\beta(E) = 0$.*

*Proof.* Fix $\varepsilon > 0$. Choose admissible $\varepsilon$-cover $(A_k)$ of $E$ with $\sum_k (\operatorname{diam} A_k)^\alpha \le \mathcal{H}^*_{\alpha,\varepsilon}(E) + 1 \le \mathcal{H}^*_\alpha(E) + 1$.
Since $\operatorname{diam} A_k < \varepsilon$, $\sum_k (\operatorname{diam} A_k)^\beta = \sum_k (\operatorname{diam} A_k)^{\beta-\alpha} (\operatorname{diam} A_k)^\alpha \le \varepsilon^{\beta-\alpha}(\mathcal{H}^*_\alpha(E) + 1)$.
Hence $\mathcal{H}^*_{\beta,\varepsilon}(E) \le \varepsilon^{\beta-\alpha}(\mathcal{H}^*_\alpha(E) + 1)$. Letting $\varepsilon \downarrow 0$ gives $\mathcal{H}^*_\beta(E) = 0$.
*Structural Commentary*: Compare one small cover at dimensions $\alpha$ and $\beta$.
*Key Insight*: The factor $\varepsilon^{\beta-\alpha}$ kills finite $\alpha$-content. □

**Definition 3.5.10.** *Hausdorff Dimension*
For $E \subseteq X$, the *Hausdorff dimension* is $\dim_H(E) := \inf\{\beta > 0 : \mathcal{H}^*_\beta(E) = 0\}$, with $\inf \emptyset := +\infty$.

**Remark 3.5.11.** *Dimension as a Critical Exponent*
For $E \subseteq X$, set $Z_E := \{\beta > 0 : \mathcal{H}^*_\beta(E) = 0\}$. Proposition 3.5.9 gives $Z_E = (\dim_H(E), +\infty)$ or $[\dim_H(E), +\infty)$ inside $(0,+\infty)$, so $\dim_H(E)$ is the zero-measure threshold for the Hausdorff scale.

# 4 Borel, Lebesgue, and Stieltjes Measures

## 4.1 Borel, Lebesgue, and Stieltjes Measure Constructions

**Proposition 4.1.1.** *Half-Open Interval Algebra on $\mathbb{R}$*
*Let $\mathcal{F}$ be the algebra of finite disjoint unions of $(a,b]$, $(-\infty,b]$, $(a,+\infty)$ $(a,b\in\mathbb{R})$, and $\mathbb{R}$. Set $\ell((a,b])=b-a$ and $\ell((-\infty,b])=\ell((a,+\infty))=\ell(\mathbb{R})=+\infty$, extending additively over disjoint unions. Then*

*(i) $\mathcal{F}$ is an algebra on $\mathbb{R}$ and $\sigma(\mathcal{F})=\mathcal{B}(\mathbb{R})$.*
*(ii) $\ell$ is a premeasure on $\mathcal{F}$.*

*Proof.*

(i) Generator complements lie in $\mathcal{F}$, hence $\mathcal{F}$ is an algebra.
Since $\mathcal{F}\subseteq\mathcal{B}(\mathbb{R})$ and $(a,b)=\bigcup_{n>1/(b-a)}(a,b-1/n]$, $\sigma(\mathcal{F})=\mathcal{B}(\mathbb{R})$.

(ii) Normal forms make $\ell$ well defined and finitely additive.

For countable additivity, reduce to $(a,b]=\bigsqcup_{j\geq1}(a_j,b_j]$: finite subunions give $\sum_{j=1}^{N}(b_j-a_j)\leq b-a$.

Conversely, choose $\varepsilon_j>0$ with $\sum_{j\geq1}\varepsilon_j<\varepsilon$; by compactness, $\exists$ finite $K\subseteq\mathbb{N}$ s.t. $[a+\delta,b]\subseteq\bigcup_{j\in K}(a_j,b_j+\varepsilon_j)$, hence $b-a-\delta\leq\sum_{j\in K}(b_j-a_j+\varepsilon_j)\leq\sum_{j\geq1}(b_j-a_j)+\varepsilon$.

Let $\delta,\varepsilon\downarrow0$. Truncation handles half-lines; finite disjoint unions follow componentwise.

*Structural Commentary*: Lebesgue measure starts as length on a finite interval algebra, not on all subsets.
*Key Insight*: Compactness supplies the reverse inequality in countable additivity of interval length. □

**Remark 4.1.2.** *Why the Half-Open Convention Matters*
Finite unions generated by $(a,b)$, $(-\infty,b)$, $(a,+\infty)$, and $\mathbb{R}$ still generate $\mathcal{B}(\mathbb{R})$, but are not an algebra, where $(0,1)^c=(-\infty,0]\cup[1,+\infty)$. Half-open endpoints make complements remain in the starting family.

**Definition 4.1.3.** *Borel–Stieltjes and Lebesgue–Stieltjes Measures*
Let $g:\mathbb{R}\to\mathbb{R}$ be increasing and right-continuous, and set $g(\pm\infty):=\lim_{x\to\pm\infty}g(x)\in\overline{\mathbb{R}}$.
Let $\mathcal{I}:=\{(a,b]:a<b\}\cup\{(-\infty,b]:b\in\mathbb{R}\}\cup\{(a,+\infty):a\in\mathbb{R}\}\cup\{\mathbb{R}\}$ and $\mathcal{F}:=\operatorname{alg}(\mathcal{I})$.

(i) Define $G_g$ on $\mathcal{I}$ by endpoint increments and set $\mu_g(\bigsqcup_{j=1}^{m}I_j):=\sum_{j=1}^{m}G_g(I_j)$ for disjoint $I_j\in\mathcal{I}$.
(ii) The unique extension of $\mu_g$ to $\mathcal{B}(\mathbb{R})$ is the *Borel–Stieltjes measure generated by $g$*.
(iii) The completion of this Borel measure is the *Lebesgue–Stieltjes measure generated by $g$*.

**Theorem 4.1.4.** *Borel–Stieltjes Measure Existence*
*Let $g:\mathbb{R}\to\mathbb{R}$ be increasing and right-continuous, and put $g(\pm\infty):=\lim_{x\to\pm\infty}g(x)\in\overline{\mathbb{R}}$.*
*Set $\mathcal{I}:=\{(a,b]:a<b\}\cup\{(-\infty,b]:b\in\mathbb{R}\}\cup\{(a,+\infty):a\in\mathbb{R}\}\cup\{\mathbb{R}\}$ and $\mathcal{F}:=\operatorname{alg}(\mathcal{I})$.*
*Define $G_g$ on $\mathcal{I}$ by endpoint increments and set $\mu_g(\bigsqcup_{j=1}^{m}I_j):=\sum_{j=1}^{m}G_g(I_j)$ for disjoint $I_j\in\mathcal{I}$. Then:*

*(i) $\mu_g$ is a $\sigma$-finite premeasure on $\mathcal{F}$ and has a unique extension to $\mathcal{B}(\mathbb{R})$, again denoted $\mu_g$.*
*(ii) If $g=\mathrm{id}_{\mathbb{R}}$, then $\mu_g$ is Lebesgue measure.*
*(iii) If $g$ is continuous, then $\mu_g(J)=g(b)-g(a)$ for $J\in\{(a,b),[a,b),(a,b],[a,b]\}$, $a<b$.*

*Proof.*

(i) Normal forms give well-defined finite additivity: $G_g$-increments telescope and $G_g\geq0$.
If $(a,b]=\bigsqcup_{j\geq1}(a_j,b_j]$, finite subunions give $\sum_{j=1}^{N}G_g((a_j,b_j])\leq G_g((a,b])$.
Conversely, compactness gives finite $K$ with $[a+\varepsilon,b]\subseteq\bigcup_{j\in K}(a_j,b_j+\varepsilon_j)$. Hence $g(b)-g(a+\varepsilon)\leq\sum_{j\in K}(g(b_j+\varepsilon_j)-g(a_j))$. Let $\varepsilon,\varepsilon_j\downarrow0$, hence countable additivity on bounded generators. Half-lines follow by truncation, finite unions componentwise, so $\mu_g$ is a premeasure. Since $\mathbb{R}=\bigcup_{n\geq1}(-n,n]$ and $\mu_g((-n,n])=g(n)-g(-n)<+\infty$, $\mu_g$ is $\sigma$-finite; Theorem 3.1.16 gives uniqueness.
(ii) If $g=\mathrm{id}_{\mathbb{R}}$, then $G_g((a,b])=b-a$, so $\mu_g$ agrees with half-open length on $\mathcal{F}$.
Uniqueness identifies its Borel extension with Lebesgue measure.
(iii) If $g$ is continuous, then $(x-1/n,x]\downarrow\{x\}$ gives $\mu_g(\{x\})=\lim_n(g(x)-g(x-1/n))=0$.

*Structural Commentary*: Stieltjes measure replaces ordinary length $b-a$ by the increment $g(b)-g(a)$.
*Key Insight*: Right-continuity aligns the half-open interval convention with countable covers. □

**Proposition 4.1.5.** *Finite Borel Measures Give Stieltjes Generators*
*Let $\nu$ be finite on $\mathcal{B}(\mathbb{R})$ and set $g(x):=\nu((-\infty,x])$. Then:*

*(i) $g$ is increasing and right-continuous;*
*(ii) $g(-\infty)=0$ and $g(+\infty)=\nu(\mathbb{R})$;*

*(iii)* $\nu((a,b]) = g(b) - g(a)$ *for* $a \le b$;
*(iv)* $\nu = \mu_g$ *on* $\mathcal{B}(\mathbb{R})$.

*Proof.*

(i) If $a \le b$, then $(-\infty, a] \subseteq (-\infty, b]$, so $g(a) \le g(b)$. If $x_n \downarrow x$, then $(-\infty, x_n] \downarrow (-\infty, x]$. Finite continuity from above gives $g(x_n) \downarrow g(x)$.
(ii) Since $(-\infty, -n] \downarrow \emptyset$ and $(-\infty, n] \uparrow \mathbb{R}$, continuity from above/below gives $g(-\infty) = 0$ and $g(+\infty) = \nu(\mathbb{R})$.
(iii) For $a \le b$, $(-\infty, b] = (-\infty, a] \sqcup (a, b]$, hence $\nu((a,b]) = g(b) - g(a)$.
(iv) By (ii)–(iii), $\nu$ and $\mu_g$ agree on $\mathcal{I}$, hence on $\mathcal{F}$. Theorem 4.1.4 gives $\nu = \mu_g$ on $\mathcal{B}(\mathbb{R})$.

*Structural Commentary*: A finite real-line measure is encoded by its cumulative mass function.
*Key Insight*: Right-continuity of distribution functions is continuity from above for measures. □

**Example 4.1.6.** *Basic Borel–Stieltjes Generators*
The following generators illustrate the interval rule in Theorem 4.1.4.

(i) If $g(x) = 2x$, then $\mu_g = 2m$ on $\mathcal{B}(\mathbb{R})$.
(ii) If $g(x) = \arctan x$, then $\mu_g((a,b]) = \arctan b - \arctan a = \int_a^b (1+x^2)^{-1}\,dx$ and $\mu_g(\mathbb{R}) = \pi$.
(iii) If $g(x) = 0$ for $x < 0$ and $g(x) = 1$ for $x \ge 0$, then $\mu_g = \delta_0$.
(iv) If $g(x) = 0$ for $x < 0$ and $g(x) = x$ for $x \ge 0$, then $\mu_g(E) = m(E \cap [0, +\infty))$ for Borel $E$.

**Lemma 4.1.7.** *Open Sets Are Countable Unions of Rational Boxes*
*For every open* $G \subseteq \mathbb{R}^n$, *there is a countable* $\Sigma \subseteq \{\prod_{i=1}^n (a_i, b_i) : a_i, b_i \in \mathbb{Q},\ a_i < b_i\}$ *such that* $G = \bigcup_{B \in \Sigma} B$.

*Proof.* Let $\mathcal{B}_{\mathbb{Q}} := \{\prod_{i=1}^n (a_i, b_i) : a_i, b_i \in \mathbb{Q},\ a_i < b_i\}$; then $\mathcal{B}_{\mathbb{Q}}$ is countable since $\mathbb{Q}^{2n}$ is countable.
Set $\Sigma := \{B \in \mathcal{B}_{\mathbb{Q}} : B \subseteq G\}$. The inclusion $\bigcup_{B\in\Sigma} B \subseteq G$ is immediate.
If $x \in G$, choose $r > 0$ with $B_r(x) \subseteq G$, then choose $a_i, b_i \in \mathbb{Q}$ with $x_i - r/n < a_i < x_i < b_i < x_i + r/n$.
Thus $x \in B := \prod_i (a_i, b_i) \subseteq B_r(x) \subseteq G$, so $x \in \bigcup_{B\in\Sigma} B$.
*Structural Commentary*: Rational boxes give a countable Euclidean base.
*Key Insight*: Openness supplies a ball; density of $\mathbb{Q}$ supplies rational box endpoints inside it. □

**Theorem 4.1.8.** *Lebesgue Measure Basic Properties*
*Let* $m = m_n^*|_{\mathcal{L}^n}$.

*(i)* $\mathcal{B}(\mathbb{R}^n) \subseteq \mathcal{L}^n$, *and* $\mathcal{L}^n$ *is closed under* $E \mapsto E^c$, *countable* $\cup$, *and countable* $\cap$.
*(ii)* *For* $E, F, E_k \in \mathcal{L}^n$, $0 \le m(E)$, $E \subseteq F \Rightarrow m(E) \le m(F)$, *and* $m(\bigcup_k E_k) \le \sum_k m(E_k)$.
*(iii)* *If* $(E_k) \subseteq \mathcal{L}^n$ *are pairwise disjoint, then* $m(\bigsqcup_k E_k) = \sum_k m(E_k)$.
*(iv)* $m([0,1]^n) = 1$, *and* $E \in \mathcal{L}^n$, $t \in \mathbb{R}^n$ *imply* $E + t \in \mathcal{L}^n$ *and* $m(E + t) = m(E)$.

*Proof.*

(i) By Theorem 3.1.6, $\mathcal{L}^n$ is a $\sigma$-algebra, hence closed under complements and countable unions/intersections. Open boxes are cells, hence in $\mathcal{L}^n$ by Proposition 3.3.10; by Lemma 4.1.7, each open $G = \bigcup_j B_j$ with rational open boxes $B_j$, so $G \in \mathcal{L}^n$. Thus $\mathcal{B}(\mathbb{R}^n) = \sigma(\mathcal{O}_{\mathbb{R}^n}) \subseteq \mathcal{L}^n$.
(ii) Since $m = m_n^*|_{\mathcal{L}^n}$, nonnegativity follows from $m_n^* \ge 0$. If $E \subseteq F$, then $m(E) \le m(F)$ by outer-measure monotonicity. Also $m(\bigcup_k E_k) = m_n^*(\bigcup_k E_k) \le \sum_k m_n^*(E_k) = \sum_k m(E_k)$ by outer-measure subadditivity.
(iii) If $(E_k) \subseteq \mathcal{L}^n$ pairwise disjoint, by Theorem 3.1.6, $m$ countably additive on $\mathcal{L}^n$, so $m(\bigsqcup_k E_k) = \sum_k m(E_k)$.
(iv) By Theorem 3.2.14, since $[0,1]^n$ is a cell, $m([0,1]^n) = \mathrm{vol}_n([0,1]^n) = 1$.
For translation, let $A \subseteq \mathbb{R}^n$. By Theorem 3.2.16 and $E \in \mathcal{L}^n$, $m_n^*(A) = m_n^*(A - t) = m_n^*((A-t)\cap E) + m_n^*((A-t)\cap E^c) = m_n^*(A \cap (E+t)) + m_n^*(A \cap (E+t)^c)$; hence $E + t \in \mathcal{L}^n$, and $m(E + t) = m_n^*(E + t) = m_n^*(E) = m(E)$.

*Structural Commentary*: The wish list is recovered only after restricting outer measure to measurable sets.
*Key Insight*: The box-cover construction gives size; the measurable-set condition restores additivity. □

**Theorem 4.1.9.** *Uniqueness from Open Cells*
*Let* $\nu$ *be a measure on* $\mathcal{L}^n$ *with* $\nu(I) = \mathrm{vol}_n(I)$ *for every open cell* $I \subseteq \mathbb{R}^n$. *Then* $\nu = m$ *on* $\mathcal{L}^n$.

*Proof.* Fix $N \in \mathbb{N}$, $Q_N = (-N, N)^n$. If $E \in \mathcal{L}^n$ and $E \subseteq Q_N$, then any open-cell cover $E \subseteq \bigcup_j J_j$ gives $\nu(E) \le \sum_j \nu(J_j) = \sum_j \mathrm{vol}_n(J_j)$, hence $\nu(E) \le m(E)$ by infimum.
Applying this to $E$ and $Q_N \setminus E$, and using $\nu(Q_N) = m(Q_N) < +\infty$, yields $\nu(E) + \nu(Q_N \setminus E) = m(E) + m(Q_N \setminus E)$ with each left summand $\le$ its right counterpart; hence $\nu(E) = m(E)$.
For arbitrary $E \in \mathcal{L}^n$, set $A_1 = E \cap Q_1$, $A_N = E \cap (Q_N \setminus Q_{N-1})$ for $N > 1$.
Then $E = \bigsqcup_N A_N$, each $A_N \subseteq Q_N$, and countable additivity gives $\nu(E) = \sum_N \nu(A_N) = \sum_N m(A_N) = m(E)$.
*Structural Commentary*: Agreement on coordinate cells determines the whole Lebesgue measure.
*Key Insight*: Covers give $\nu \le m$ locally; finite complements force equality. □

**Method 4.1.10.** *Translation Invariance by Moving the Test Set*

(i) For $t \in \mathbb{R}^n$ and $A \subseteq \mathbb{R}^n$, translated box covers give $m_n^*(A+t) = m_n^*(A)$.
(ii) If $E \in \mathcal{L}^n$, then for all $A \subseteq \mathbb{R}^n$, $(A \cap (E+t)) - t = (A-t) \cap E$ and $(A \cap (E+t)^c) - t = (A-t) \cap E^c$.
(iii) Apply the Caratheodory split for $E$ to $A - t$ to get $E + t \in \mathcal{L}^n$ and $m(E+t) = m(E)$.

*Key Insight*: Translation invariance of outer measure transports the clean-splitting identity.

**Proposition 4.1.11.** *First Borel Hierarchy Examples*
*In $\mathbb{R}^p$:*

*(i) If $G = \bigcap_{n\geq 1} U_n$ with $U_n$ open or $F = \bigcup_{n\geq 1} C_n$ with $C_n$ closed, then $G, F \in \mathcal{B}(\mathbb{R}^p) \subseteq \mathcal{L}^p$.*
*(ii) Any set obtained from open or closed sets by countably many $\cup, \cap, {}^c$ operations is Borel.*
*(iii) In $\mathbb{R}$, $\{0\} = \bigcap_{n\geq 1} (-\frac{1}{n}, \frac{1}{n})$ is $G_\delta$ but not open.*
*(iv) In $\mathbb{R}$, $(0,1) = \bigcup_{n\geq 2} [\frac{1}{n}, 1 - \frac{1}{n}]$ is $F_\sigma$ but not closed.*

*Proof.*

(i) Since $\mathcal{B}(\mathbb{R}^p) = \sigma(\mathcal{O}_{\mathbb{R}^p})$, it contains open sets, closed sets by complements, and countable $\cap, \cup$ by $\sigma$-algebra closure; Theorem 4.1.8 gives $\mathcal{B}(\mathbb{R}^p) \subseteq \mathcal{L}^p$.
(ii) This is Proposition 1.1.3 applied to $\mathcal{B}(\mathbb{R}^p)$.
(iii) For $x \in \mathbb{R}$, $x = 0 \iff |x| < \frac{1}{n}$ for all $n \geq 1$; the singleton $\{0\}$ is not open.
(iv) For $x \in \mathbb{R}$, $0 < x < 1 \iff \exists n \geq 2 : \frac{1}{n} \leq x \leq 1 - \frac{1}{n}$; the interval $(0,1)$ is not closed.

*Structural Commentary*: Borel sets are generated by countable topological operations.
*Key Insight*: Countable intersections and unions quickly leave the class of open and closed sets. □

**Perspective 4.1.12.** *Why Not Measure Every Subset?*
Vitali-type sets show $m_1^*$ is not additive on $2^{\mathbb{R}}$: disjoint unions and refinements can break additivity.
Thus measure is $m^*$ on clean splitters $E$: $m^*(X) = m^*(X \cap E) + m^*(X \cap E^c)$ for all $X$.
*Key Insight*: Outer measure is universal; measure lives on clean splitters.

**Definition 4.1.13.** *Null Sets and Real-Line Measure Zero*
Let $(X, \mathcal{A}, \mu)$ be a measure space and let $A \subseteq \mathbb{R}$.

(i) $N \subseteq X$ is *null* iff $N \in \mathcal{A}$ and $\mu(N) = 0$.
(ii) $A$ has *measure zero* iff $\forall \varepsilon > 0$, $\exists$ open intervals $(O_n)$ with $A \subseteq \bigcup_{n=1}^\infty O_n$ and $\sum_{n=1}^\infty |O_n| \leq \varepsilon$.
(iii) If $A \subseteq [a,b]$, then $A$ has *content zero* iff the condition in (ii) holds with finitely many intervals.

For Lebesgue measure on $\mathbb{R}$, null sets are also *zero sets*.

**Proposition 4.1.14.** *Basic Measure-Zero Examples on $\mathbb{R}$*

*(i) Every finite $A \subseteq \mathbb{R}$ has content 0.*
*(ii) Every countable $A \subseteq \mathbb{R}$ has measure 0.*
*(iii) The Cantor set $P$ has content 0.*

*Proof.*

(i) For $A = \{a_1, \ldots, a_N\}$ and $\varepsilon > 0$, choose open $I_j \ni a_j$ with $|I_j| < \frac{\varepsilon}{N}$; then $A \subseteq \bigcup_{j=1}^N I_j$ and $\sum_{j=1}^N |I_j| < \varepsilon$. The case $A = \emptyset$ is trivial.
(ii) For $A = \{a_j : j \in \mathbb{N}\}$ and $\varepsilon > 0$, choose open $I_j \ni a_j$ with $|I_j| < \frac{\varepsilon}{2^j}$; then $A \subseteq \bigcup_{j\geq 1} I_j$ and $\sum_{j\geq 1} |I_j| < \varepsilon$.
(iii) Given $\varepsilon > 0$, choose $N$ with $\left(\frac{2}{3}\right)^N < \frac{\varepsilon}{2}$. If $E_N = \bigcup_{k=1}^{2^N} J_k$, $|J_k| = 3^{-N}$, choose open $I_k \supseteq J_k$ with $|I_k| < 3^{-N} + \frac{\varepsilon}{2^{N+1}}$; then $P \subseteq \bigcup_{k=1}^{2^N} I_k$ and $\sum_{k=1}^{2^N} |I_k| < \left(\frac{2}{3}\right)^N + \frac{\varepsilon}{2} < \varepsilon$.

*Structural Commentary*: Zero size means arbitrarily small interval covers.
*Key Insight*: Lists split $\varepsilon$ by terms; Cantor stages have length $(\frac{2}{3})^N \to 0$. □

**Corollary 4.1.15.** *Uncountable Null Sets Exist*
*There exists $P \subseteq \mathbb{R}$ such that:*

*(i) $|P| = 2^{\aleph_0}$;*
*(ii) $P \in \mathcal{L}^1$ and $m(P) = 0$.*

*Proof.* Let $P := \{\sum_{k=1}^\infty \varepsilon_k 3^{-k} : \varepsilon_k \in \{0,2\}\}$.

(i) The map $(\varepsilon_k) \mapsto \sum_k \varepsilon_k 3^{-k}$ identifies $P$ with $\{0,2\}^{\mathbb{N}}$, so $|P| = 2^{\aleph_0}$.

(ii) At stage $N$, $P$ is covered by $2^N$ closed intervals of length $3^{-N}$, hence $m_1^*(P) \le (2/3)^N$ for all $N$. Letting $N \to \infty$ gives $m_1^*(P) = 0$, so outer-null completeness gives $P \in \mathcal{L}^1$ and $m(P) = 0$.

*Structural Commentary*: Cantor coding separates cardinality from length.
*Key Insight*: Null need not mean countable. □

**Corollary 4.1.16.** *Real-Line Lebesgue Measure Checks*
*Let $m$ be Lebesgue measure on $\mathbb{R}$.*

*(i)* $a < b \Rightarrow m((a,b)) = m((a,b]) = m([a,b)) = m([a,b]) = b - a$.
*(ii)* $A \subseteq \mathbb{R}$ *countable* $\Rightarrow A \in \mathcal{B}(\mathbb{R})$ *and* $m(A) = 0$.
*(iii)* $\emptyset \neq G \subseteq \mathbb{R}$ *open* $\Rightarrow m(G) > 0$.
*(iv)* $K \subseteq \mathbb{R}$ *compact* $\Rightarrow m(K) < +\infty$.

*Proof.*

(i) By cell volume, $m([a,b]) = b-a$; endpoint sets finite null; add or delete $\{a\}, \{b\}$ does not change measure.
(ii) If $A = \{a_k : k \in J\}$ with $J$ finite or countable, then $A = \bigcup_{k\in J}\{a_k\} \in \mathcal{B}(\mathbb{R})$; $m(A) \le \sum_{k\in J} m(\{a_k\}) = 0$.
(iii) Choose $(a,b) \subseteq G$ with $a < b$; by monotonicity and (i), $m(G) \ge m((a,b)) = b - a > 0$.
(iv) Compact $K$ is bounded, so $K \subseteq [-R,R]$ for some $R > 0$; hence $m(K) \le m([-R,R]) = 2R < +\infty$.

*Structural Commentary*: On $\mathbb{R}$, length, null endpoints, and compact boundedness give quick measure estimates.
*Key Insight*: Nonempty open sets contain intervals; compact sets fit inside one. □

**Example 4.1.17.** *A Positive-Measure Cantor-Type Set with No Interval*
Let $S_0 = [0,1]$ and, for $n \ge 1$, let $R_n$ be the union of the middle open intervals of length $2^{-2n}$ removed from the $2^{n-1}$ components of $S_{n-1}$; set $S_n := S_{n-1} \setminus R_n$ and $S := \bigcap_{n=0}^{\infty} S_n$.
Then $m(R_n) = 2^{-n-1}$, $\sum_{n\ge1} m(R_n) = 1/2$, so $S = [0,1] \setminus \bigcup_n R_n$ is compact with $m(S) = 1/2$.
Since every component of $S_n$ has length $\le 2^{-n}$, no interval $(a,b)$ can lie in $S$; hence $\operatorname{int} S = \emptyset$.
*Structural Commentary*: Summable removals leave measure; repeated splitting kills interior.
*Key Insight*: Total removed length controls measure; repeated splitting controls interior.

# 5 Continuity and Regularity of Measures

## 5.1 Measure Continuity and Regularity

**Corollary 5.1.1.** *Measure Continuity*
*Let $\mu$ be a measure, and let $(E_k)$ and $(F_k)$ be measurable sequences.*

*(i) If $E_k \uparrow E$, then $\mu(E_k) \uparrow \mu(E)$.*
*(ii) If $F_k \downarrow F$ and $\mu(F_1) < +\infty$, then $\mu(F_k) \downarrow \mu(F)$.*

*Proof.*

(i) Set $G_1 = E_1$ and $G_k = E_k \setminus E_{k-1}$ for $k \ge 2$.
Then $E_k = \bigsqcup_{j=1}^{k} G_j$ and $E = \bigsqcup_{j\ge1} G_j$, so $\mu(E_k) = \sum_{j=1}^{k} \mu(G_j) \uparrow \sum_{j\ge1} \mu(G_j) = \mu(E)$.
(ii) Apply (i) to $H_k = F_1 \setminus F_k \uparrow F_1 \setminus F$. Since $\mu(F_1) < +\infty$, $\mu(F_k) = \mu(F_1) - \mu(H_k) \downarrow \mu(F_1) - \mu(F_1 \setminus F) = \mu(F)$.

*Structural Commentary*: Monotone set limits are disjoint series or complements of such series.
*Key Insight*: Downward continuity needs one finite anchor so subtraction is legal. □

**Definition 5.1.2.** *Limit Inferior and Superior of Sets*
For $(E_n) \subseteq \mathcal{P}(X)$, define $\limsup_n E_n := \bigcap_{m=1}^{\infty} \bigcup_{n\ge m} E_n$ and $\liminf_n E_n := \bigcup_{m=1}^{\infty} \bigcap_{n\ge m} E_n$.

(i) *Infinitely often*: $x \in \limsup_n E_n$ iff $\{n : x \in E_n\}$ is infinite;
(ii) *Eventually always*: $x \in \liminf_n E_n$ iff $\{n : x \notin E_n\}$ is finite;
(iii) *Set limit*: if $\limsup_n E_n = \liminf_n E_n$, write $\lim_n E_n$ for the common set.

**Proposition 5.1.3.** *Basic Laws for Set Limits*
*Let $(E_n) \subseteq \mathcal{P}(X)$.*

*(i)* $\emptyset \subseteq \liminf_n E_n \subseteq \limsup_n E_n \subseteq X$.
*(ii)* $E_n \uparrow E \Longrightarrow \liminf_n E_n = \limsup_n E_n = E = \bigcup_n E_n$.
*(iii)* $E_n \downarrow E \Longrightarrow \liminf_n E_n = \limsup_n E_n = E = \bigcap_n E_n$.

*(iv)* $(E_n) \subseteq \mathcal{M} \Longrightarrow \liminf_n E_n, \limsup_n E_n \in \mathcal{M}$.

*Proof.*

(i) Since $\bigcap_m \bigcup_{n\geq m} E_n \subseteq \bigcup_{n\geq 1} E_n \subseteq X$ and $\bigcap_{n\geq m} E_n \subseteq \bigcup_{n\geq m} E_n$ for all $m$, the inclusions follow.

(ii) If $E_n \uparrow E$, then $\{n : x \in E_n\}$ is nonempty iff it is cofinite, hence $\liminf_n E_n = \limsup_n E_n = \bigcup_n E_n = E$.

(iii) If $E_n \downarrow E$, then $\{n : x \in E_n\}$ is infinite iff it is all of $\mathbb{N}$, hence $\liminf_n E_n = \limsup_n E_n = \bigcap_n E_n = E$.

(iv) If $(E_n) \subseteq \mathcal{M}$, then $\bigcup_{n\geq m} E_n$, $\bigcap_{n\geq m} E_n \in \mathcal{M}$ $\forall m$; countable closure gives $\limsup_n E_n, \liminf_n E_n \in \mathcal{M}$.

*Structural Commentary*: Set limsup means infinitely often; set liminf means eventually always.
*Key Insight*: Translate the definitions into pointwise membership statements. □

**Proposition 5.1.4.** *Measure Bounds for Set Limits*
*Let $(E_n)$ be measurable in a measure space $(X, \mathcal{M}, \mu)$.*

*(i)* $\mu(\liminf_n E_n) \leq \liminf_n \mu(E_n)$.

*(ii)* $\mu(\bigcup_{n=1}^{\infty} E_n) < +\infty \Longrightarrow \limsup_n \mu(E_n) \leq \mu(\limsup_n E_n)$.

*Proof.*

(i) Set $A_m := \bigcap_{n\geq m} E_n$. Then $A_m \uparrow \liminf_n E_n$ and $A_m \subseteq E_n$ for all $n \geq m$, so $\mu(A_m) \leq \inf_{n\geq m} \mu(E_n)$.
By continuity from below, $\mu(\liminf_n E_n) = \lim_m \mu(A_m) \leq \sup_m \inf_{n\geq m} \mu(E_n) = \liminf_n \mu(E_n)$.

(ii) Set $B_m := \bigcup_{n\geq m} E_n$. $B_m \downarrow \limsup_n E_n$, $\mu(B_1) < +\infty$, $E_n \subseteq B_m$ $\forall n \geq m$, so $\sup_{n\geq m} \mu(E_n) \leq \mu(B_m)$.
By continuity from above, $\limsup_n \mu(E_n) = \inf_m \sup_{n\geq m} \mu(E_n) \leq \lim_m \mu(B_m) = \mu(\limsup_n E_n)$.

*Structural Commentary*: Tail intersections increase to liminf; tail unions decrease to limsup.
*Key Insight*: Compare each tail envelope to the corresponding tail bounds of the measures. □

**Definition 5.1.5.** *Almost Everywhere*
Let $(X, \mathcal{M}, \mu)$ be a measure space, let $E \in \mathcal{M}$, and let $P(x)$ be a property on $E$.

(i) $P$ holds *$\mu$-almost everywhere on $E$* iff $\exists N \in \mathcal{M}$ with $N \subseteq E$, $\mu(N) = 0$, and $P$ holds on $E \setminus N$.

(ii) For Lebesgue measure, write *almost everywhere*.

**Theorem 5.1.6.** *First Borel–Cantelli Lemma*
*Let $(X, \mathcal{M}, \mu)$ be measure space, $(E_n) \subseteq \mathcal{M}$, set $L := \limsup_n E_n = \bigcap_{m\geq 1} \bigcup_{n\geq m} E_n$. Assume $\sum_{n=1}^{\infty} \mu(E_n) < +\infty$.*

*(i) $L \in \mathcal{M}$ and $\mu(L) = 0$.*

*(ii) $\mu$-almost every $x \in X$ belongs to at most finitely many $E_n$.*

*Proof.*

(i) Set $B_m := \bigcup_{n\geq m} E_n$. Then $B_m \downarrow L$ and $\mu(B_m) \leq \sum_{n\geq m} \mu(E_n)$.
Since $\mu(B_1) \leq \sum_{n\geq 1} \mu(E_n) < +\infty$, continuity from above gives $\mu(L) = \lim_{m\to\infty} \mu(B_m) \leq \lim_{m\to\infty} \sum_{n\geq m} \mu(E_n) = 0$.

(ii) Since $x \in L \iff \forall m\, \exists n \geq m$ with $x \in E_n$, $L$ is the set of points lying in infinitely many $E_n$.
By (i), outside the null set $L$, membership occurs only finitely often.

*Structural Commentary*: Summable tails force the limsup set to have zero measure.
*Key Insight*: Points appearing infinitely often must lie in every tail union. □

**Remark 5.1.7.** *Borel Sets, Null Sets, and Regular Approximation*
Open, closed, and Borel subsets of $\mathbb{R}^p$ are measurable. Each $A \in \mathcal{M}(\mu)$ admits closed–open approximations $F \subseteq A \subseteq G$ with $\mu(G \setminus A)$ and $\mu(A \setminus F)$ arbitrarily small. In Lebesgue measure, one may take $F_\sigma \subseteq A \subseteq G_\delta$ with $\mu(G_\delta \setminus F_\sigma) = 0$. Null sets form a $\sigma$-ring, and countable subsets of $\mathbb{R}^p$ are null.
*Structural Commentary*: Measurable sets are Borel sets modified by null sets.
*Key Insight*: Regularity explains why open and closed approximations are enough for integration.

**Proposition 5.1.8.** *Local Measurability on Bounded Cubes*
*For arbitrary $E \subseteq \mathbb{R}^p$, $E \in \mathcal{L}^p \iff E \cap [-n, n]^p \in \mathcal{L}^p$ for all $n \in \mathbb{N}$.*

*Proof.*

($\Rightarrow$) If $E \in \mathcal{L}^p$, then $E \cap [-n,n]^p \in \mathcal{L}^p$ for all $n$, since $[-n,n]^p$ is Borel.
($\Leftarrow$) If $E \cap [-n,n]^p \in \mathcal{L}^p$ for all $n$, then $E = \bigcup_{n=1}^{\infty}(E \cap [-n,n]^p) \in \mathcal{L}^p$.

*Structural Commentary*: Nested Borel cubes exhaust $\mathbb{R}^p$.
*Key Insight*: Measurability is local along a countable exhaustion. □

**Proposition 5.1.9.** *Local Measurability on Unit Intervals*
*Let $I_n := (n, n+1]$, $n \in \mathbb{Z}$. For $E \subseteq \mathbb{R}$, $E \in \mathcal{L}^1$ iff $E \cap I_n \in \mathcal{L}^1$ for all $n \in \mathbb{Z}$.*

*Proof.*

($\Rightarrow$) Since $I_n \in \mathcal{B}(\mathbb{R}) \subseteq \mathcal{L}^1$, $E \in \mathcal{L}^1$ implies $E \cap I_n \in \mathcal{L}^1$ for all $n$.
($\Leftarrow$) If $E \cap I_n \in \mathcal{L}^1$ for all $n$, then $E = \bigcup_{n\in\mathbb{Z}}(E \cap I_n) \in \mathcal{L}^1$.

*Structural Commentary*: A countable Borel exhaustion localizes measurability.
*Key Insight*: Local measurable pieces patch by countable unions. □

**Proposition 5.1.10.** *Finite Measure Meeting Every Unit Interval*
*With $I_n := (n, n+1]$, there exists $E \in \mathcal{B}(\mathbb{R})$ such that $0 < m(E \cap I_n)$ for all $n \in \mathbb{Z}$ and $m(E) < +\infty$.*

*Proof.* Set $\varepsilon_n := 2^{-|n|-3}$ and $E := \bigcup_{n\in\mathbb{Z}}(n, n+\varepsilon_n]$. Then $(n, n+\varepsilon_n] \subseteq I_n$ are pairwise disjoint Borel intervals, so $E \in \mathcal{B}(\mathbb{R})$ and $m(E \cap I_n) = \varepsilon_n > 0$. Countable additivity gives $m(E) = \sum_{n\in\mathbb{Z}} \varepsilon_n = \frac{3}{8} < +\infty$.
*Structural Commentary*: Finite measure can be spread over countably many cells.
*Key Insight*: Positive local masses can be summable globally. □

**Proposition 5.1.11.** *Null Sets Form a Sigma-Ideal*
*Let $(X, \mathcal{M}, \mu)$ be a measure space and $\mathcal{Z} := \{N \in \mathcal{M} : \mu(N) = 0\}$.*
*Then $\mathcal{Z}$ is a $\sigma$-ring, and $N \in \mathcal{Z}$, $E \in \mathcal{M}$ imply $N \cap E \in \mathcal{Z}$.*

*Proof.* If $(N_k) \subseteq \mathcal{Z}$, then $\mu(\bigcup_k N_k) \le \sum_k \mu(N_k) = 0$, so $\bigcup_k N_k \in \mathcal{Z}$.
If $A, B \in \mathcal{Z}$, then $A \setminus B \subseteq A$, hence $A \setminus B \in \mathcal{Z}$. Finally, $N \cap E \subseteq N$ gives $N \cap E \in \mathcal{Z}$.
*Structural Commentary*: Null sets are stable under countable unions and measurable cuts.
*Key Insight*: Monotonicity makes every measurable piece of a null set null. □

**Proposition 5.1.12.** *Completed Sigma-Algebra*
*Let $(X, \mathcal{M}, \mu)$ be measure space, $\mathcal{Z} := \{N \in \mathcal{M} : \mu(N) = 0\}$; $\overline{\mathcal{M}}_\mu := \{A \subseteq X : \exists E \in \mathcal{M},\ \exists N \in \mathcal{Z},\ A \triangle E \subseteq N\}$.*

*(i) $\mathcal{M} \subseteq \overline{\mathcal{M}}_\mu$, and $\overline{\mathcal{M}}_\mu$ is a $\sigma$-algebra;*
*(ii) $A \subseteq N \in \mathcal{Z} \Longrightarrow A \in \overline{\mathcal{M}}_\mu$;*
*(iii) $\overline{\mathcal{M}}_\mu = \{E \cup Z : E \in \mathcal{M},\ \exists N \in \mathcal{Z},\ Z \subseteq N\}$.*

*Proof.*

(i) For $E \in \mathcal{M}$, $E \triangle E = \emptyset$, so $\mathcal{M} \subseteq \overline{\mathcal{M}}_\mu$. If $A \triangle E \subseteq N$, then $A^c \triangle E^c = A \triangle E \subseteq N$.
If $A_k \triangle E_k \subseteq N_k$, then $(\bigcup_k A_k) \triangle (\bigcup_k E_k) \subseteq \bigcup_k N_k \in \mathcal{Z}$. Hence $\overline{\mathcal{M}}_\mu$ is a $\sigma$-algebra.
(ii) If $A \subseteq N \in \mathcal{Z}$, then $A \triangle \emptyset \subseteq N$, so $A \in \overline{\mathcal{M}}_\mu$.
(iii) If $A \triangle E \subseteq N$, then $A = (E \setminus N) \cup (A \cap N)$, with $E \setminus N \in \mathcal{M}$ and $A \cap N \subseteq N$.
Conversely, if $A = E \cup Z$ and $Z \subseteq N \in \mathcal{Z}$, then $A \triangle E \subseteq Z \subseteq N$.

*Structural Commentary*: Completion adds all null modifications of measurable sets.
*Key Insight*: Equality modulo null sets is the membership criterion. □

**Proposition 5.1.13.** *Completed Measure*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, $\mathcal{Z} := \{N \in \mathcal{M} : \mu(N) = 0\}$,*
*and $\overline{\mathcal{M}}_\mu := \{A \subseteq X : \exists E \in \mathcal{M},\ \exists N \in \mathcal{Z},\ A \triangle E \subseteq N\}$. For $A \in \overline{\mathcal{M}}_\mu$, define $\overline{\mu}(A) := \mu(E)$ whenever $E \in \mathcal{M}$ and $A \triangle E \subseteq N \in \mathcal{Z}$. Then $\overline{\mu}$ is a well-defined complete measure on $\overline{\mathcal{M}}_\mu$, and $\overline{\mu}|_{\mathcal{M}} = \mu$.*

*Proof.* If $A \triangle E \subseteq N_1$ and $A \triangle F \subseteq N_2$, with $E, F \in \mathcal{M}$ and $N_1, N_2 \in \mathcal{Z}$, then $E \triangle F \subseteq N_1 \cup N_2$, so $\mu(E) = \mu(F)$.
Hence $\overline{\mu}$ is well defined; since $E \triangle E = \emptyset$, $\overline{\mu}|_{\mathcal{M}} = \mu$.
For disjoint $(A_k) \subseteq \overline{\mathcal{M}}_\mu$, choose $E_k \in \mathcal{M}$ and $N_k \in \mathcal{Z}$ with $A_k \triangle E_k \subseteq N_k$, and set $N := \bigcup_k N_k$.
Then $E_i \cap E_j \subseteq N$ for $i \neq j$. Put $F_1 := E_1$ and $F_k := E_k \setminus \bigcup_{j<k} E_j$.
Then $(F_k)$ disjoint, $A_k \triangle F_k \subseteq N$, $(\bigcup_k A_k) \triangle (\bigcup_k F_k) \subseteq N$, so $\overline{\mu}(\bigcup_k A_k) = \mu(\bigcup_k F_k) = \sum_k \mu(F_k) = \sum_k \overline{\mu}(A_k)$.
If $A \subseteq B \in \overline{\mathcal{M}}_\mu$ and $\overline{\mu}(B) = 0$, choose $E \in \mathcal{M}$ and $N \in \mathcal{Z}$ with $B \triangle E \subseteq N$.
Then $\mu(E) = 0$, so $A \subseteq B \subseteq E \cup N \in \mathcal{Z}$; hence $A \in \overline{\mathcal{M}}_\mu$ and $\overline{\mu}(A) = 0$.
*Structural Commentary*: The completed measure ignores null errors.
*Key Insight*: Null perturbations do not change countable additivity. □

## 5.2 Radon Measures on Locally Compact Spaces

Radon measure theory is where topology and measure start determining one another. The topological input is a locally compact Hausdorff space; the measure-theoretic output is a Borel measure that can be recovered from open neighborhoods, compact subsets, and compactly supported continuous test functions.

**Definition 5.2.1.** *Locally Compact Borel Setup*
Let $(X, \mathcal{T})$ be a topological space.

(i) $X$ is *locally compact* if every $x \in X$ admits $U \in \mathcal{T}$ with $x \in U$ and $\overline{U}$ compact.
(ii) The *Borel $\sigma$-algebra* is $\mathcal{B}(X) := \sigma(\mathcal{T})$.
(iii) For $f : X \to \mathbb{R}$, define $\operatorname{supp} f := \overline{\{x \in X : f(x) \neq 0\}}$.
(iv) $C_c(X)$ is the subspace of $C(X, \mathbb{R})$ consisting of functions with compact support.

**Proposition 5.2.2.** *Compact Neighborhood Shrinking*
*Let $X$ be locally compact and Hausdorff, $K \subseteq X$ compact, and $O \subseteq X$ open with $K \subseteq O$. Then there is an open $V \subseteq X$ such that $K \subseteq V$, $\overline{V} \subseteq O$, and $\overline{V}$ is compact.*

*Proof.* For each $x \in K$, choose open $U_x \ni x$ with $\overline{U_x}$ compact.
Normality of the compact Hausdorff space $\overline{U_x}$ gives a relatively open $W_x \ni x$ with $\overline{W_x}^{\overline{U_x}} \subseteq O \cap U_x$.
Choose open $A_x \subseteq X$ with $W_x = A_x \cap \overline{U_x}$ and set $V_x := A_x \cap U_x$. Then $x \in V_x$ and $\overline{V_x} \subseteq \overline{W_x}^{\overline{U_x}} \subseteq O \cap U_x$.
The family $(V_x)_{x \in K}$ covers $K$, so $K \subseteq \bigcup_{i=1}^n V_{x_i}$ for some $x_1, \dots, x_n \in K$.
Set $V := \bigcup_{i=1}^n V_{x_i}$. Then $\overline{V} = \bigcup_{i=1}^n \overline{V_{x_i}} \subseteq O$, and $\overline{V}$ is compact as a finite union of compact sets.
*Structural Commentary*: Shrink each point neighborhood inside $O$, then take a finite union.
*Key Insight*: Compactness converts local shrinkings into one global compact neighborhood. □

**Proposition 5.2.3.** *Compactly Supported Cutoff*
*Let $X$ be locally compact and Hausdorff, $K \subseteq X$ compact, and $O \subseteq X$ open with $K \subseteq O$. Then there is $f \in C_c(X)$ such that $0 \leq f \leq 1$ on $X$, $f|_K = 1$, and $\operatorname{supp} f \subseteq O$.*

*Proof.* Choose open $V \subseteq X$ as in Proposition 5.2.2, so $K \subseteq V$, $\overline{V}$ is compact, and $\overline{V} \subseteq O$.
The disjoint sets $K$ and $\overline{V} \setminus V$ are closed in the compact Hausdorff space $\overline{V}$.
By Urysohn's lemma, Theorem 6.3.18, there is $f_0 : \overline{V} \to [0,1]$ with $f_0|_K = 1$ and $f_0|_{\overline{V} \setminus V} = 0$.
Define $f : X \to [0,1]$ by $f|_{\overline{V}} = f_0$ and $f|_{X \setminus V} = 0$.
Closed-set pasting gives $f \in C_c(X)$ with $f|_K = 1$ and $\operatorname{supp} f \subseteq \overline{V} \subseteq O$.
*Structural Commentary*: Apply Urysohn separation on $\overline{V}$ and extend by zero.
*Key Insight*: The compact neighborhood forces the extension to have compact support. □

**Proposition 5.2.4.** *Finite Partition of Unity on a Compact Set*
*Let $X$ be locally compact and Hausdorff, $K \subseteq X$ compact, and $O_1, \dots, O_n \subseteq X$ open with $K \subseteq \bigcup_{i=1}^n O_i$. Then there are $\varphi_1, \dots, \varphi_n \in C_c(X)$ such that $0 \leq \varphi_i \leq 1$, $\operatorname{supp} \varphi_i \subseteq O_i$ for every $i$, and $(\sum_{i=1}^n \varphi_i)|_K = 1$.*

*Proof.* For each $x \in K$, choose $i(x)$ with $x \in O_{i(x)}$ and, by Proposition 5.2.2, an open $V_x \ni x$ with $\overline{V_x}$ compact and $\overline{V_x} \subseteq O_{i(x)}$. Choose $x_1, \dots, x_m \in K$ with $K \subseteq \bigcup_{j=1}^m V_{x_j}$, write $i_j := i(x_j)$, and set $U_i := \bigcup_{\{j : i_j = i\}} V_{x_j}$.
Then $K \subseteq \bigcup_i U_i$, each $U_i$ is open, and $\overline{U_i}$ is compact with $\overline{U_i} \subseteq O_i$.
For $U_i \neq \emptyset$, Proposition 5.2.3 gives $f_i \in C_c(X)$ with $0 \leq f_i \leq 1$, $f_i|_{\overline{U_i}} = 1$, $\operatorname{supp} f_i \subseteq O_i$; set $f_i = 0$ otherwise.
Applying the same proposition to $K \subseteq \bigcup_i U_i$ gives $h \in C_c(X)$ with $0 \leq h \leq 1$, $h|_K = 1$, and $\operatorname{supp} h \subseteq \bigcup_i U_i$.
Set $g := \sum_{i=1}^n f_i$ and $d := g + 1 - h$. Since $g \geq 1$ on $\operatorname{supp} h$ and $h = 0$ off $\operatorname{supp} h$, $d \geq 1$. Hence $\varphi_i := f_i / d$ belongs to $C_c(X)$, satisfies $0 \leq \varphi_i \leq 1$, and has $\operatorname{supp} \varphi_i \subseteq O_i$. On $K$, $h = 1$ and $g \geq 1$, so $d = g$ and $(\sum_{i=1}^n \varphi_i)|_K = 1$.
*Structural Commentary*: Build finitely many local cutoffs, then normalize their sum.
*Key Insight*: The auxiliary cutoff keeps the denominator positive where local cutoffs vanish. □

**Theorem 5.2.5.** *Locally Compact Separation and Cutoff Package*
*Let $X$ be Hausdorff. The following are equivalent.*

*(i) $X$ is locally compact.*
*(ii) For every compact $K \subseteq O \in \mathcal{T}$, there is open $V \supseteq K$ with $\overline{V}$ compact and $\overline{V} \subseteq O$.*
*(iii) For every compact $K \subseteq O \in \mathcal{T}$, there is $f \in C_c(X)$ with $0 \leq f \leq 1$, $f|_K = 1$, and $f|_{X \setminus O} = 0$.*

*Proof.*

(i) $\Rightarrow$ (ii) Proposition 5.2.2.
(i) $\Rightarrow$ (iii) Proposition 5.2.3.
(ii) $\Rightarrow$ (i) For $x \in X$, apply (ii) to $K = \{x\}$ and $O = X$; the resulting open $V \ni x$ has compact closure.

(*iii*) $\Rightarrow$ (*i*) For $x \in X$, apply (iii) to $K = \{x\}$ and $O = X$, and set $U := f^{-1}((1/2, 3/2))$.
Then $x \in U$ and $\overline{U} \subseteq f^{-1}([1/2, 3/2]) \subseteq \operatorname{supp} f$, so $\overline{U}$ is compact.

*Structural Commentary*: Compact cutoffs recover compact neighborhoods through level sets.
*Key Insight*: Compact-neighborhood shrinking and compactly supported cutoffs are equivalent local forms of local compactness. □

**Method 5.2.6.** *Metric Cutoff Formula*
Let $X$ be locally compact and Hausdorff, assume that $\rho$ induces its topology.
Let $K \subseteq O \subseteq X$ with $K$ compact and $O$ open, and choose open $U \subseteq X$ with $K \subseteq U$, $\overline{U}$ compact, and $\overline{U} \subseteq O$.
For every nonempty $A \subseteq X$, set $\operatorname{dist}_A(x) := \inf_{a \in A} \rho(x, a)$.

(i) *Empty Set*: If $K = \emptyset$, set $f := 0$.
(ii) *Trivial Complement*: If $K \neq \emptyset$ and $X \setminus U = \emptyset$, set $f := 1$.
(iii) *Distance Ratio*: Otherwise, set $f(x) := \frac{\operatorname{dist}_{X \setminus U}(x)}{\operatorname{dist}_{X \setminus U}(x) + \operatorname{dist}_K(x)}$.

In case (iii), $K$ and $X \setminus U$ are disjoint closed sets, so the denominator is positive; continuity of distance functions gives $f \in C(X, [0, 1])$. In all cases, $f \in C_c(X)$, $f|_K = 1$, $f|_{X \setminus U} = 0$, and $\operatorname{supp} f \subseteq \overline{U} \subseteq O$.
*Key Insight*: Metric distances give an explicit compactly supported cutoff.

**Proposition 5.2.7.** *Alexandroff One-Point Compactification*
*Let $(X, \mathcal{T})$ be locally compact and Hausdorff, let $\omega \notin X$, and write $\mathcal{K}(X)$ for the compact subsets of $X$. Set $X^* := X \cup \{\omega\}$ and $\mathcal{T}^* := \mathcal{T} \cup \{X^* \setminus K : K \in \mathcal{K}(X)\}$. Let $\iota : X \hookrightarrow X^*$ be the inclusion.*

*(i) $\mathcal{T}^*$ is a topology on $X^*$, $(X^*, \mathcal{T}^*)$ is compact Hausdorff, and $\iota : X \to X^* \setminus \{\omega\}$ is a homeomorphism.*
*(ii) Every $h \in C(K, \mathbb{R})$ with $K \subseteq X$ compact has an extension $\widetilde{h} \in C(X, \mathbb{R})$ satisfying $\widetilde{h}|_K = h$.*

*Proof.*

(i) Compact subsets of the Hausdorff space $X$ are closed.
Thus $U \setminus C \in \mathcal{T}$ and $(X^* \setminus C) \cap (X^* \setminus D) = X^* \setminus (C \cup D)$ for $U \in \mathcal{T}$ and $C, D \in \mathcal{K}(X)$.
If $\omega \notin \bigcup_\alpha U_\alpha$, then $\bigcup_\alpha U_\alpha \in \mathcal{T}$; otherwise choose $U_{\alpha_0} = X^* \setminus C$, so $X^* \setminus \bigcup_\alpha U_\alpha$ is closed in $C$ and hence compact. Thus $\mathcal{T}^*$ is a topology.
Any open cover contains $X^* \setminus C$ for some $C \in \mathcal{K}(X)$, and finitely many remaining members cover $C$; hence $X^*$ is compact. Points of $X$ are already separated.
For $x \in X$, choose open $V \ni x$ with $\overline{V}$ compact; then $V$ and $X^* \setminus \overline{V}$ separate $x$ and $\omega$.
Finally, $\mathcal{T}^*|_X = \mathcal{T}$ since $X \setminus C \in \mathcal{T}$ for $C \in \mathcal{K}(X)$, so $\iota$ is a homeomorphism onto $X^* \setminus \{\omega\}$.
(ii) By (i), $X^*$ is compact Hausdorff and hence normal. The compact set $K$ is closed in $X^*$, and $h$ is bounded. Tietze extension, Theorem 6.3.19, gives $H \in C(X^*, \mathbb{R})$ with $H|_K = h$. Set $\widetilde{h} := H|_X$.

*Structural Commentary*: Neighborhoods of $\omega$ are complements of compact subsets of $X$.
*Key Insight*: Compactification transfers extension problems to a compact Hausdorff space. □

**Definition 5.2.8.** *Radon Measures and Radon Premeasures*
Let $(X, \mathcal{T})$ be a topological space, and write $\mathcal{K}(X)$ for its compact subsets.

(i) $\nu : \mathcal{B}(X) \to [0, +\infty]$ is a *Radon measure* if $\nu(K) < +\infty$ $\forall K \in \mathcal{K}(X)$, $\nu(E) = \inf\{\nu(O) : E \subseteq O,\ O \in \mathcal{T}\}$ $\forall E \in \mathcal{B}(X)$, and $\nu(O) = \sup\{\nu(K) : K \subseteq O,\ K \in \mathcal{K}(X)\}$ $\forall O \in \mathcal{T}$.
(ii) $\mu : \mathcal{T} \to [0, +\infty]$ is a *Radon premeasure* if $\mu(U) < +\infty$ $\forall U \in \mathcal{T}$ with $\overline{U} \in \mathcal{K}(X)$ and $\mu(O) = \sup\{\mu(U) : U \in \mathcal{T},\ \overline{U} \in \mathcal{K}(X),\ \overline{U} \subseteq O\}$ $\forall O \in \mathcal{T}$.

**Lemma 5.2.9.** *Open Envelope for a Topological Premeasure*
*Let $\mu : \mathcal{T} \to [0, +\infty]$ be a premeasure. For $E \subseteq X$, define $\mu^*(E) := \inf\{\sum_{j=1}^\infty \mu(U_j) : E \subseteq \bigcup_{j=1}^\infty U_j,\ U_j \in \mathcal{T}\}$. Then $\mu^*(E) = \inf\{\mu(U) : E \subseteq U,\ U \in \mathcal{T}\}$.*

*Proof.*

($\Rightarrow$) Set $I(E) := \inf\{\mu(U) : E \subseteq U,\ U \in \mathcal{T}\}$. Every $U \in \mathcal{T}$ with $E \subseteq U$ is a one-set cover, so $\mu^*(E) \leq I(E)$.
($\Leftarrow$) If $(U_j)_{j \geq 1} \subseteq \mathcal{T}$ covers $E$, then $U := \bigcup_{j=1}^\infty U_j$ belongs to $\mathcal{T}$ and satisfies $\mu(U) \leq \sum_{j=1}^\infty \mu(U_j)$.
Hence $I(E) \leq \sum_{j=1}^\infty \mu(U_j)$; taking the infimum over covers gives $I(E) \leq \mu^*(E)$.

*Structural Commentary*: Replace each countable cover by its union.
*Key Insight*: Subadditivity makes the union no more expensive than the cover. □

**Lemma 5.2.10.** *Open-Set Test for Caratheodory Measurability*
*Let $\mu : \mathcal{T} \to [0, +\infty]$ be a premeasure, let $\mu^*$ be the outer measure induced by open covers, and let $E \subseteq X$. Assume $\mu(O) \geq \mu^*(O \cap E) + \mu^*(O \setminus E)$ for every $O \in \mathcal{T}$ with $\mu(O) < +\infty$. Then $E$ is $\mu^*$-measurable.*

*Proof.* Fix $A \subseteq X$. Outer subadditivity gives $\mu^*(A) \le \mu^*(A \cap E) + \mu^*(A \setminus E)$.
If $\mu^*(A) = +\infty$, the reverse inequality is automatic. Otherwise, for $\varepsilon > 0$, Lemma 5.2.9 gives $O \in \mathcal{T}$ with $A \subseteq O$ and $\mu(O) < \mu^*(A) + \varepsilon$, so $\mu^*(A \cap E) + \mu^*(A \setminus E) \le \mu^*(O \cap E) + \mu^*(O \setminus E) \le \mu(O) < \mu^*(A) + \varepsilon$.
Letting $\varepsilon \downarrow 0$ gives the reverse inequality, hence the Caratheodory criterion.
*Structural Commentary*: Approximate $A$ by an open envelope of nearly equal outer mass.
*Key Insight*: Open-set splitting propagates to all sets through outer regularity. □

**Proposition 5.2.11.** *Open Sets from a Radon Premeasure Are Measurable*
*Let $X$ be locally compact and Hausdorff, $\mu : \mathcal{T} \to [0, +\infty]$ Radon premeasure, $\mu^*$ open-cover outer measure.*

*(i)* $\mu^*|_{\mathcal{T}} = \mu$.
*(ii) Every $V \in \mathcal{T}$ is $\mu^*$-measurable.*

*Proof.*

(i) The one-set cover gives $\mu^*(V) \le \mu(V)$ for $V \in \mathcal{T}$.
Every $(U_j)_{j \ge 1} \subseteq \mathcal{T}$ covering $V$ satisfies $\mu(V) \le \mu(\bigcup_j U_j) \le \sum_j \mu(U_j)$, so $\mu(V) \le \mu^*(V)$.
(ii) Let $V, O \in \mathcal{T}$ with $\mu(O) < +\infty$.
For $\varepsilon > 0$, choose $U \in \mathcal{T}$ with $\overline{U}$ compact, $\overline{U} \subseteq O \cap V$, and $\mu(U) > \mu(O \cap V) - \varepsilon$.
Since $U$ and $O \setminus \overline{U}$ are disjoint open subsets of $O$ and $O \setminus V \subseteq O \setminus \overline{U}$,
$\mu(O) \ge \mu(U) + \mu(O \setminus \overline{U}) \ge \mu^*(O \cap V) - \varepsilon + \mu^*(O \setminus V)$ by (i). Let $\varepsilon \downarrow 0$ and apply Lemma 5.2.10.

*Structural Commentary*: Inner approximation leaves an open remainder containing $O \setminus V$.
*Key Insight*: Open inner regularity yields Caratheodory splitting. □

**Theorem 5.2.12.** *Radon Premeasure Extension*
*Let $X$ be locally compact and Hausdorff, let $\mu : \mathcal{T} \to [0, +\infty]$ be a Radon premeasure, let $\mu^*$ be its open-cover outer measure, write $\mathcal{K}(X)$ for the compact subsets of $X$, and set $\widehat{\mu} := \mu^*|_{\mathcal{B}(X)}$.*

*(i)* $\widehat{\mu}|_{\mathcal{T}} = \mu$.
*(ii) $\widehat{\mu}$ is a Radon measure on $\mathcal{B}(X)$.*

*Proof.*

(i) For $O \in \mathcal{T} \subseteq \mathcal{B}(X)$, $\widehat{\mu}(O) = \mu^*(O) = \mu(O)$ by Proposition 5.2.11; hence $\widehat{\mu}|_{\mathcal{T}} = \mu$.
(ii) Let $\mathcal{C}_{\mu^*}$ be the Caratheodory $\sigma$-algebra of $\mu^*$. Proposition 5.2.11 gives $\mathcal{T} \subseteq \mathcal{C}_{\mu^*}$, hence $\mathcal{B}(X) = \sigma(\mathcal{T}) \subseteq \mathcal{C}_{\mu^*}$.
As $\mu^*|_{\mathcal{C}_{\mu^*}}$ is measure, so is $\widehat{\mu}$. Lemma 5.2.9, (i) give $\widehat{\mu}(E) = \inf\{\widehat{\mu}(O) : E \subseteq O,\ O \in \mathcal{T}\}$ for $E \in \mathcal{B}(X)$.
For $K \in \mathcal{K}(X)$, Proposition 5.2.2 gives $K \subseteq U \in \mathcal{T}$ with $\overline{U} \in \mathcal{K}(X)$, so $\widehat{\mu}(K) \le \widehat{\mu}(U) = \mu(U) < +\infty$.
For $O \in \mathcal{T}$, inner regularity and (i) give
$\widehat{\mu}(O) = \sup\{\mu(U) : U \in \mathcal{T},\ \overline{U} \in \mathcal{K}(X),\ \overline{U} \subseteq O\} \le \sup\{\widehat{\mu}(K) : K \in \mathcal{K}(X),\ K \subseteq O\} \le \widehat{\mu}(O)$.

*Structural Commentary*: $\mathcal{T} \subseteq \mathcal{C}_{\mu^*}$ yields $\widehat{\mu}$ on $\mathcal{B}(X)$; $K \in \mathcal{K}(X) \Rightarrow \widehat{\mu}(K) < +\infty$.
*Key Insight*: Open inner regularity becomes compact inner regularity after extension. □

**Theorem 5.2.13.** *Radon Measures Determined by $C_c(X)$*
*Let $X$ be locally compact and Hausdorff, and let $\mu_1, \mu_2$ be Radon measures on $\mathcal{B}(X)$.*
*If $\int_X f \, d\mu_1 = \int_X f \, d\mu_2$ for every $f \in C_c(X)$, then $\mu_1 = \mu_2$.*

*Proof.* Write $\mathcal{K}(X)$ for the compact subsets of $X$. Let $K \in \mathcal{K}(X)$ and $\varepsilon > 0$.
By outer regularity, choose $O \in \mathcal{T}$ with $K \subseteq O$ and $\mu_i(O \setminus K) < \varepsilon$ for $i = 1, 2$.
Choose $f \in C_c(X)$ with $0 \le f \le 1$, $f|_K = 1$, and $\operatorname{supp} f \subseteq O$, and set $r_i := \int_{O \setminus K} f \, d\mu_i$.
Then $0 \le r_i < \varepsilon$ and $\int_X f \, d\mu_i = \mu_i(K) + r_i$, so $|\mu_1(K) - \mu_2(K)| = |r_1 - r_2| < \varepsilon$.
Hence $\mu_1 = \mu_2$ on $\mathcal{K}(X)$, and inner then outer regularity give $\mu_1|_{\mathcal{T}} = \mu_2|_{\mathcal{T}}$ and $\mu_1 = \mu_2$ on $\mathcal{B}(X)$.
*Structural Commentary*: $C_c(X)$ identifies $\mu_1$ and $\mu_2$ on $\mathcal{K}(X)$.
*Key Insight*: $\mathcal{K}(X) \to \mathcal{T} \to \mathcal{B}(X)$ by Radon regularity. □

Figure 198: Riesz–Markov: local compactness supplies cutoffs; orange denotes Caratheodory extension.

**Theorem 5.2.14.** *Riesz–Markov Theorem on $C_c(X)$*
*Let $X$ be locally compact and Hausdorff, $I : C_c(X) \to \mathbb{R}$ linear with $I(f) \ge 0$ $\forall f \in C_c(X)$ satisfying $f \ge 0$.*
*Then $\exists$ a unique Radon measure $\widehat{\mu}$ on $\mathcal{B}(X)$ such that $I(f) = \int_X f \, d\widehat{\mu}$ $\forall f \in C_c(X)$.*

*Proof.* For $O \in \mathcal{T}$, define $\mu(O) := \sup\{I(f) : f \in C_c(X),\ 0 \le f \le 1,\ \operatorname{supp} f \subseteq O\}$ and $\mu(\emptyset) := 0$.

(i) *Premeasure*: If $O \subseteq \bigcup_{j=1}^{\infty} O_j$ and $f$ is admissible for $\mu(O)$, a finite partition $(\varphi_\ell)_{\ell=1}^m$ subordinate to a finite subcover of $\operatorname{supp} f$ gives $f = \sum_{\ell=1}^m f\varphi_\ell$ and $I(f) \le \sum_{\ell=1}^m \mu(O_{j_\ell}) \le \sum_{j=1}^{\infty} \mu(O_j)$, so $\mu$ is countably subadditive. For disjoint $O_1, O_2 \in \mathcal{T}$, countable subadditivity gives one inequality and admissible $f_1, f_2$ give $\mu(O_1) + \mu(O_2) \le \mu(O_1 \cup O_2)$, hence finite additivity.

(ii) *Radon Regularity*: For $a < \mu(O)$, choose admissible $f$ with $I(f) > a$ and $U \in \mathcal{T}$ with $\operatorname{supp} f \subseteq U \subseteq \overline{U} \subseteq O$ and $\overline{U}$ compact. Then $\mu(U) > a$, proving inner regularity. If $\overline{U}$ is compact, a cutoff $g \in C_c(X)$ with $g|_{\overline{U}} = 1$ dominates every admissible $f$ for $\mu(U)$, so $\mu(U) \le I(g) < +\infty$. Thus $\mu$ is a Radon premeasure.

(iii) *Extension*: Stages (i)–(ii) make $\mu$ a Radon premeasure on $\mathcal{T}$. Theorem 5.2.12 applies to its open-cover outer measure and yields the Radon measure $\widehat{\mu} := \mu^*|_{\mathcal{B}(X)}$ with $\widehat{\mu}|_{\mathcal{T}} = \mu$.

(iv) *Identification*: Let $0 \le f \le 1$ belong to $C_c(X)$, choose $O \in \mathcal{T}$ with $\operatorname{supp} f \subseteq O$ and $\overline{O}$ compact, and set $\psi_k := \min\{1, \max\{0, nf - (k-1)\}\}$, $O_0 := O$, $O_{n+1} := \emptyset$, $O_k := \{x \in O : f(x) > \frac{k-1}{n}\}$ for $1 \le k \le n$. Then $f = \frac{1}{n}\sum_{k=1}^n \psi_k$, $\operatorname{supp}\psi_k \subseteq O_{k-1}$, and $\psi_k|_{O_{k+1}} = 1$, so $\mu(O_{k+1}) \le I(\psi_k) \le \mu(O_{k-1})$ and $\mu(O_{k+1}) \le \int_X \psi_k\, d\widehat{\mu} \le \mu(O_{k-1})$. Therefore $|I(f) - \int_X f\, d\widehat{\mu}| \le \frac{1}{n}\sum_{k=1}^n (\mu(O_{k-1}) - \mu(O_{k+1})) \le \frac{2\mu(O)}{n}$. Letting $n \to \infty$, scaling, and $f = f^+ - f^-$ give $I(f) = \int_X f\, d\widehat{\mu}$ for every $f \in C_c(X)$. Uniqueness follows from Theorem 5.2.13.

*Structural Commentary*: Open-set contents and compactly supported cutoffs build the Radon measure before identifying its integral.
*Key Insight*: Layer functions recover $I$ from $\widehat{\mu}$. □

**Theorem 5.2.15.** *Lusin Approximation for Radon Measures*
*Let $X$ be locally compact Hausdorff, let $\mu$ be Radon on $\mathcal{B}(X)$, let $f : X \to \mathbb{R}$ be Borel, and set $A := \{f \ne 0\}$ with $\mu(A) < +\infty$. For every $\varepsilon > 0$, there exist $X_0 \in \mathcal{B}(X)$ and $g \in C_c(X)$ such that:*

*(i) $g = f$ on $X_0$.*
*(ii) $\mu(X \setminus X_0) < \varepsilon$.*

*Proof.* Set $\delta := \varepsilon/4$. Choose $M > 0$ with $\mu(A_M) < \delta$, where $A_M := A \cap \{|f| > M\}$, and put $B := A \setminus A_M$. For $n \ge 1$, choose finite-valued Borel $s_n : B \to [-M, M]$ with $\|s_n - f\|_{\infty,B} \le \frac{1}{n}$. Let $s_n(B) = \{a_{n,1}, \dots, a_{n,m_n}\}$ and $E_{n,j} := s_n^{-1}(\{a_{n,j}\})$. Inner regularity gives compact $K_{n,j} \subseteq E_{n,j}$, $\mu(E_{n,j} \setminus K_{n,j}) < \frac{\delta}{2^n m_n}$. Set $F_n := \bigcup_{j=1}^{m_n} K_{n,j}$ and $F := \bigcap_{n=1}^{\infty} F_n$.

Then $F$ is compact, $B \setminus F \subseteq \bigcup_{n\ge1}\bigcup_{j=1}^{m_n}(E_{n,j} \setminus K_{n,j})$, and $\mu(B \setminus F) \le \delta$.

For each $n$, $F \cap E_{n,j} = F \cap K_{n,j}$ is closed in $F$, so $s_n|_F \in C(F)$; since $s_n|_F \to f|_F$ uniformly, $f|_F \in C(F)$. Choose $W \in \mathcal{T}$ with $F \subseteq W$ and $\mu(W \setminus F) < \delta$, then choose $O \in \mathcal{T}$ by Proposition 5.2.2 with $F \subseteq O \subseteq \overline{O} \subseteq W$ and $\overline{O}$ compact. Extend $f|_F$ to $H \in C(X)$ by Proposition 5.2.7, choose $q \in C_c(X)$ with $q|_F = 1$ and $\operatorname{supp} q \subseteq O$ by Proposition 5.2.3, and set $g := qH$, $X_0 := F \cup (X \setminus (A \cup O))$. Then $g \in C_c(X)$ and $X_0 \in \mathcal{B}(X)$.

(i) On $F$, $g = qH = f$; on $X \setminus (A \cup O)$, $f = q = 0$.
(ii) $X \setminus X_0 \subseteq (A \setminus F) \cup (O \setminus F)$, $\mu(A \setminus F) \le \mu(A_M) + \mu(B \setminus F) < 2\delta$, $\mu(O \setminus F) \le \mu(W \setminus F) < \delta$; $\mu(X \setminus X_0) < 3\delta < \varepsilon$.

*Structural Commentary*: Compact core plus cutoff extension.
*Key Insight*: Discard a small set to make Borel data continuous. □

## 5.3 Invariant Measures on Compact Groups

Compact groups are the meeting point of topological symmetry and Radon measure theory. The compactness supplies fixed points in dual spaces, and Riesz–Markov converts those fixed functionals into invariant Borel measures.

**Definition 5.3.1.** *Compact Groups and Haar Measures*
For a group $G$, $A \subseteq G$, and $g \in G$, set $gA := \{ga : a \in A\}$ and $Ag := \{ag : a \in A\}$.

(i) A *topological group* is $G$ with Hausdorff topology such that $(g, h) \mapsto gh$ and $g \mapsto g^{-1}$ are continuous.
(ii) A *compact group* is a topological group whose underlying topological space is compact.
(iii) A Borel measure $\mu$ on $\mathcal{B}(G)$ is *left-invariant* if $\mu(gA) = \mu(A)$ for all $g \in G$ and $A \in \mathcal{B}(G)$.
(iv) A Borel measure $\mu$ on $\mathcal{B}(G)$ is *right-invariant* if $\mu(Ag) = \mu(A)$ for all $g \in G$ and $A \in \mathcal{B}(G)$.
(v) A *Haar measure* on a compact group $G$ is a left-invariant Radon probability measure on $\mathcal{B}(G)$.

**Definition 5.3.2.** *Regular Actions on $C(G)$*
Let $G$ be compact, $C(G)$ be continuous maps $G \to \mathbb{R}$ with $\|f\|_\infty := \max_{x\in G} |f(x)|$. For $g, x \in G$ and $f \in C(G)$:

(i) The *left regular action* is $\pi_{\mathrm{L}} : G \to \mathcal{L}(C(G))$, $[\pi_{\mathrm{L}}(g)f](x) := f(g^{-1}x)$.
(ii) The *right regular action* is $\pi_{\mathrm{R}} : G \to \mathcal{L}(C(G))$, $[\pi_{\mathrm{R}}(g)f](x) := f(xg^{-1})$.
(iii) A *probability functional* is $\psi \in C(G)^*$ with $\psi(\mathbf{1}_G) = 1$ and $f \geq 0 \Rightarrow \psi(f) \geq 0$ for all $f \in C(G)$.

**Theorem 5.3.3.** *Kakutani Fixed Probability Functional*
*Let $G$ be a compact group. Then there are probability functionals $\psi_{\mathrm{L}}, \psi_{\mathrm{R}} \in C(G)^*$ such that $\psi_{\mathrm{L}}(\pi_{\mathrm{L}}(g)f) = \psi_{\mathrm{L}}(f)$ and $\psi_{\mathrm{R}}(\pi_{\mathrm{R}}(g)f) = \psi_{\mathrm{R}}(f)$ for every $g \in G$ and $f \in C(G)$.*

*Proof.* Let $\mathcal{P}(G) := \{\psi \in C(G)^* : \psi(\mathbf{1}_G) = 1,\ \forall f \in C(G)\ (f \geq 0 \Rightarrow \psi(f) \geq 0)\}$. Then $\delta_e \in \mathcal{P}(G)$, $\mathcal{P}(G)$ is convex and weak-* closed, and $\psi \in \mathcal{P}(G) \Rightarrow \|\psi\|_{\mathrm{op}} = 1$; hence $\mathcal{P}(G)$ is weak-* compact by Banach–Alaoglu.
For $g \in G$, set $T_{\mathrm{L}}(g)\psi := \psi \circ \pi_{\mathrm{L}}(g^{-1})$ and $T_{\mathrm{R}}(g)\psi := \psi \circ \pi_{\mathrm{R}}(g)$. Since $\pi_{\mathrm{L}}, \pi_{\mathrm{R}}$ preserve $\mathbf{1}_G$ and $C(G)_+$ and are norm-continuous in $g$, $T_{\mathrm{L}}$ and $T_{\mathrm{R}}$ are continuous affine $G$-actions on $\mathcal{P}(G)$.
Kakutani gives $\psi_{\mathrm{L}}, \psi_{\mathrm{R}} \in \mathcal{P}(G)$ with $T_{\mathrm{L}}(g)\psi_{\mathrm{L}} = \psi_{\mathrm{L}}$ and $T_{\mathrm{R}}(g)\psi_{\mathrm{R}} = \psi_{\mathrm{R}}$ for all $g \in G$.
Thus $\psi_{\mathrm{L}} \circ \pi_{\mathrm{L}}(g) = \psi_{\mathrm{L}}$ and $\psi_{\mathrm{R}} \circ \pi_{\mathrm{R}}(g) = \psi_{\mathrm{R}}$ for all $g \in G$.
*Structural Commentary*: Compactness lives in $\mathcal{P}(G)$; invariance comes from fixed points.
*Key Insight*: A fixed probability functional is invariant. □

**Method 5.3.4.** *Fixed Functional Pattern*
For a compact group $G$, set $\mathcal{P}(G) := \{\psi \in C(G)^* : \psi(\mathbf{1}_G) = 1,\ \forall f \in C(G)\ (f \geq 0 \Rightarrow \psi(f) \geq 0)\}$.

(i) Show $\mathcal{P}(G) \neq \varnothing$ via $\delta_x(f) := f(x)$, $x \in G$.
(ii) Show $\mathcal{P}(G)$ is weak-* compact convex: $\psi \in \mathcal{P}(G) \Rightarrow \|\psi\|_{\mathrm{op}} = 1$, Banach–Alaoglu, and weak-* closed defining conditions.
(iii) Define $T_{\mathrm{L}}(g)\psi := \psi \circ \pi_{\mathrm{L}}(g^{-1})$ or $T_{\mathrm{R}}(g)\psi := \psi \circ \pi_{\mathrm{R}}(g)$. For $T = T_{\mathrm{L}}, T_{\mathrm{R}}$, check $T(g)\mathcal{P}(G) \subseteq \mathcal{P}(G)$ and weak-* continuity.
(iv) Apply Kakutani: find $\psi \in \mathcal{P}(G)$ with $T(g)\psi = \psi$ for all $g \in G$.
(v) Apply Riesz–Markov, $\psi(f) = \int_G f\, d\mu$, and read $T(g)\psi = \psi$ as invariance of $\mu$.

*Structural Commentary*: Compact state space plus adjoint action.
*Key Insight*: Haar measure is an invariant state represented by Riesz–Markov.

**Proposition 5.3.5.** *Translated Measures and Test Functions*
*Let $G$ be a compact group, let $\mu$ be a Borel measure on $\mathcal{B}(G)$, and fix $g \in G$.*
*Set $L_g(x) := gx$ and $R_g(x) := xg$ for $x \in G$.*
*For $A \in \mathcal{B}(G)$, define $\mu_g^{\mathrm{L}}(A) := \mu(L_g(A)) = \mu(gA)$ and $\mu_g^{\mathrm{R}}(A) := \mu(R_g(A)) = \mu(Ag)$.*

*(i) $\mu_g^{\mathrm{L}}$ and $\mu_g^{\mathrm{R}}$ are Borel measures on $\mathcal{B}(G)$.*
*(ii) If $\mu$ is Radon, then $\mu_g^{\mathrm{L}}$ and $\mu_g^{\mathrm{R}}$ are Radon.*
*(iii) For every $f \in C(G)$, $\int_G \pi_{\mathrm{L}}(g)f\, d\mu = \int_G f\, d\mu_g^{\mathrm{L}}$ and $\int_G \pi_{\mathrm{R}}(g)f\, d\mu = \int_G f\, d\mu_g^{\mathrm{R}}$.*

*Proof.* For every $h \in G$, the maps $L_h$ and $R_h$ are homeomorphisms of $G$.

(i) Since $L_g$ and $R_g$ map Borel sets bijectively onto Borel sets, $\mu_g^{\mathrm{L}}$ and $\mu_g^{\mathrm{R}}$ are defined on $\mathcal{B}(G)$. If $(A_n)_{n\geq 1}$ are pairwise disjoint Borel sets, then $L_g(\bigcup_n A_n) = \bigcup_n L_g(A_n)$ and $R_g(\bigcup_n A_n) = \bigcup_n R_g(A_n)$, with disjoint unions; countable additivity follows from countable additivity of $\mu$.
(ii) Since $L_{g^{-1}}^{-1} = L_g$ and $R_{g^{-1}}^{-1} = R_g$, $\mu_g^{\mathrm{L}} = (L_{g^{-1}})_*\mu$ and $\mu_g^{\mathrm{R}} = (R_{g^{-1}})_*\mu$. If $\mu$ is Radon, pushforward by a homeomorphism preserves compact sets, open sets, and Radon regularity. Hence both translated measures are Radon.
(iii) For $f \in C(G)$, the pushforward identity gives $\int_G f\, d\mu_g^{\mathrm{L}} = \int_G f \circ L_{g^{-1}}\, d\mu = \int_G \pi_{\mathrm{L}}(g)f\, d\mu$ and $\int_G f\, d\mu_g^{\mathrm{R}} = \int_G f \circ R_{g^{-1}}\, d\mu = \int_G \pi_{\mathrm{R}}(g)f\, d\mu$.

*Structural Commentary*: Translated measures are inverse-translation pushforwards.
*Key Insight*: Measure translation is dual to test-function precomposition. □

**Theorem 5.3.6.** *Haar Measure on Compact Groups*
*Let $G$ be a compact group. Let $\mathcal{P}_{\mathrm{L}}(G)$ and $\mathcal{P}_{\mathrm{R}}(G)$ be the left- and right-invariant Radon probability measures on $\mathcal{B}(G)$, respectively.*

*(i) $\mathcal{P}_{\mathrm{L}}(G) \neq \emptyset$.*
*(ii) $\mathcal{P}_{\mathrm{R}}(G) \neq \emptyset$.*
*(iii) For every $(\lambda, \rho) \in \mathcal{P}_{\mathrm{L}}(G) \times \mathcal{P}_{\mathrm{R}}(G)$, $\lambda = \rho$.*

*Consequently, $G$ admits a unique Haar measure, and this measure is right-invariant.*

*Proof.* For a Radon probability measure $\sigma$ on $\mathcal{B}(G)$, set $\Phi_\sigma(f) := \int_G f\,d\sigma$ for $f \in C(G)$.

(i) Choose $\psi_{\mathrm{L}}$ from Theorem 5.3.3. Riesz–Markov gives a Radon probability measure $\mu$ with $\psi_{\mathrm{L}} = \Phi_\mu$ on $C(G) = C_c(G)$. For $g \in G$, Proposition 5.3.5 gives $\Phi_{\mu_g^{\mathrm{L}}}(f) = \Phi_\mu(\pi_{\mathrm{L}}(g)f) = \Phi_\mu(f)$ for every $f \in C(G)$. Thus $\mu_g^{\mathrm{L}} = \mu$ by Theorem 5.2.13, so $\mu(gA) = \mu(A)$ for every $A \in \mathcal{B}(G)$ and $\mu \in \mathcal{P}_{\mathrm{L}}(G)$.

(ii) Choose $\psi_{\mathrm{R}}$ from Theorem 5.3.3. Riesz–Markov gives a Radon probability measure $\nu$ with $\psi_{\mathrm{R}} = \Phi_\nu$ on $C(G) = C_c(G)$. For $g \in G$, Proposition 5.3.5 gives $\Phi_{\nu_g^{\mathrm{R}}}(f) = \Phi_\nu(\pi_{\mathrm{R}}(g)f) = \Phi_\nu(f)$ for every $f \in C(G)$. Thus $\nu_g^{\mathrm{R}} = \nu$ by Theorem 5.2.13, so $\nu(Ag) = \nu(A)$ for every $A \in \mathcal{B}(G)$ and $\nu \in \mathcal{P}_{\mathrm{R}}(G)$.

(iii) Let $(\lambda, \rho) \in \mathcal{P}_{\mathrm{L}}(G) \times \mathcal{P}_{\mathrm{R}}(G)$, fix $f \in C(G)$, and set $h_f(x,y) := f(xy)$. Fubini applies to $h_f \in C(G \times G)$. Left-invariance gives $\int_G h_f(x,y)\,d\lambda(y) = \Phi_\lambda(f)$ for every $x$, and right-invariance gives $\int_G h_f(x,y)\,d\rho(x) = \Phi_\rho(f)$ for every $y$. Since $\lambda(G) = \rho(G) = 1$, Fubini gives $\Phi_\lambda(f) = \Phi_\rho(f)$. Hence $\lambda = \rho$ by Theorem 5.2.13.

Taking $\mu \in \mathcal{P}_{\mathrm{L}}(G)$ from (i) and $\nu \in \mathcal{P}_{\mathrm{R}}(G)$ from (ii), (iii) gives $\mu = \nu$, so $\mu$ is right-invariant. If $\eta$ is Haar, then $\eta \in \mathcal{P}_{\mathrm{L}}(G)$ and (iii) gives $\eta = \nu$, so Haar measure is unique.
*Structural Commentary*: Fixed states give existence; Fubini gives uniqueness.
*Key Insight*: Invariance makes both iterated integrals constant. □

**Proposition 5.3.7.** *Point Masses of Haar Measure*
*Let $G$ be a compact group, and let $\mu$ be Haar measure on $\mathcal{B}(G)$.*

*(i) If $G$ is finite, then every $A \subseteq G$ is Borel and $\mu(A) = \frac{|A|}{|G|}$.*
*(ii) If $G$ is infinite, then $\mu(\{x\}) = 0$ for every $x \in G$.*

*Proof.* Set $c := \mu(\{e\})$. For each $x \in G$, left-invariance gives $\mu(\{x\}) = \mu(x\{e\}) = c$.

(i) If $G$ is finite, then $G$ is discrete, so every $A \subseteq G$ is Borel. Since $1 = \mu(G) = \sum_{x \in G} \mu(\{x\}) = |G|\,c$, one has $c = \frac{1}{|G|}$, and finite additivity gives $\mu(A) = \sum_{a \in A} \mu(\{a\}) = \frac{|A|}{|G|}$.

(ii) If $G$ is infinite and $c > 0$, choose $n \in \mathbb{N}$ with $n > \frac{1}{c}$ and distinct $x_1, \dots, x_n \in G$.
Finite additivity gives $1 = \mu(G) \geq \sum_{j=1}^{n} \mu(\{x_j\}) = nc > 1$, a contradiction; hence $c = 0$.

*Structural Commentary*: Translation reduces point masses to one scalar.
*Key Insight*: Normalization fixes or kills that scalar. □

**Proposition 5.3.8.** *Nonempty Open Sets Have Positive Haar Measure*
*Let $G$ be a compact group, $\mu$ be Haar measure on $\mathcal{B}(G)$, $O \subseteq G$ be nonempty and open. Then $\mu(O) > 0$.*

*Proof.* Choose $a \in O$. For each $x \in G$, $x = (xa^{-1})a \in (xa^{-1})O$, so $\{gO : g \in G\}$ is an open cover of $G$. Compactness gives $g_1, \dots, g_n \in G$ such that $G = \bigcup_{j=1}^{n} g_j O$. If $\mu(O) = 0$, then left-invariance and subadditivity give $1 = \mu(G) \leq \sum_{j=1}^{n} \mu(g_j O) = \sum_{j=1}^{n} \mu(O) = 0$, a contradiction. Hence $\mu(O) > 0$.
*Structural Commentary*: Compactness gives a finite translate cover.
*Key Insight*: Zero mass cannot cover total mass. □

# 6 Regular Approximation and Caratheodory Tests

## 6.1 Borel Approximation and Regular Measurable Sets

The Caratheodory definition is intrinsic to outer measure, but in $\mathbb{R}^p$ it has a topological shadow: measurable sets are exactly those that can be approximated by open, closed, compact, and cell-built sets up to arbitrarily small error. That bridge is also the route to the first nonmeasurable and non-Borel examples.

**Lemma 6.1.1.** *Outer Approximation by Open and $G_\delta$ Sets*
*For $A \subseteq \mathbb{R}^p$:*

*(i) for every $\varepsilon > 0$, there exists open $G \subseteq \mathbb{R}^p$ with $A \subseteq G$ and $m(G) \leq m_p^*(A) + \varepsilon$;*
*(ii) $m_p^*(A) = \inf\{m(G) : A \subseteq G,\ G \subseteq \mathbb{R}^p \text{ open}\}$;*
*(iii) there exists $G_\delta$ set $H \supseteq A$ with $m(H) = m_p^*(A)$.*

*Proof.*

(i) If $m_p^*(A) = +\infty$, take $G = \mathbb{R}^p$.
If $m_p^*(A) < +\infty$, choose an open-cell cover $A \subseteq \bigcup_k I_k$ with $\sum_k \mathrm{vol}_p(I_k) \leq m_p^*(A) + \varepsilon$, and set $G := \bigcup_k I_k$. Then $G$ is open, $A \subseteq G$, and $m(G) \leq \sum_k m(I_k) = \sum_k \mathrm{vol}_p(I_k) \leq m_p^*(A) + \varepsilon$.

(ii) By (i), $\inf\{m(G) : A \subseteq G,\ G \subseteq \mathbb{R}^p \text{ open}\} \leq m_p^*(A) + \varepsilon$ for all $\varepsilon > 0$; let $\varepsilon \downarrow 0$.
Conversely, $A \subseteq G$ gives $m_p^*(A) \leq m_p^*(G) = m(G)$ for open $G$; take the infimum over $G$.

(iii) If $m_p^*(A) = +\infty$, take $H = \mathbb{R}^p$.
If $m_p^*(A) < +\infty$, choose open $G_n \supseteq A$ with $m(G_n) \leq m_p^*(A) + 1/n$ and set $H := \bigcap_n G_n$.
Then $H$ is $G_\delta$, $A \subseteq H$, and $m_p^*(A) \leq m(H) \leq m(G_n) \leq m_p^*(A) + 1/n$ for all $n$, so $m(H) = m_p^*(A)$.

*Structural Commentary*: Outer measure can be realised by a Borel envelope from outside.
*Key Insight*: Infima over cell covers can be repackaged as open-set approximation. □

**Corollary 6.1.2.** *Null Sets Sit Inside Borel Null Sets*
*Let $Z \subseteq \mathbb{R}^p$ with $m_p^*(Z) = 0$. Then there exists $B \subseteq \mathbb{R}^p$ such that:*
*(i) $Z \subseteq B$;*
*(ii) $B \in \mathcal{B}(\mathbb{R}^p)$;*
*(iii) $m(B) = 0$.*

*Proof.* By Lemma 6.1.1, choose open sets $(G_n)_{n\geq 1}$ in $\mathbb{R}^p$ and set $B := \bigcap_{n=1}^{\infty} G_n$ so that $Z \subseteq B$ and $m(B) = m_p^*(Z)$.

(i) The chosen envelope satisfies $Z \subseteq B$.
(ii) Since each $G_n$ is open, $B = \bigcap_{n=1}^{\infty} G_n \in \mathcal{B}(\mathbb{R}^p)$.
(iii) Since $m_p^*(Z) = 0$, $m(B) = m_p^*(Z) = 0$.

*Structural Commentary*: A null set has a $G_\delta$ null envelope.
*Key Insight*: Outer approximation gives a Borel null superset. □

**Theorem 6.1.3.** *Open and $G_\delta$ Characterizations of Measurability*
*For $E \subseteq \mathbb{R}^p$, the following are equivalent:*
*(i) $E \in \mathcal{L}^p$;*
*(ii) $\forall \varepsilon > 0$, $\exists$ open $G \subseteq \mathbb{R}^p$ such that $E \subseteq G$ and $m_p^*(G \setminus E) < \varepsilon$;*
*(iii) $\exists$ $G_\delta$ set $H \subseteq \mathbb{R}^p$ such that $E \subseteq H$ and $m_p^*(H \setminus E) = 0$;*
*(iv) $\exists$ $G_\delta$ set $H \subseteq \mathbb{R}^p$ and $Z \in \mathcal{L}^p$ such that $Z \subseteq H$, $m(Z) = 0$, and $E = H \setminus Z$.*

*Proof.*

(i) $\Rightarrow$ (ii). If $m(E) < +\infty$, choose open $G \supseteq E$ with $m(G) \leq m(E) + \varepsilon/2$.
Since $E \in \mathcal{L}^p$, $m_p^*(G \setminus E) = m(G) - m(E) < \varepsilon$. If $m(E) = +\infty$, set $E_1 := E \cap \{x : \|x\|_2 \leq 1\}$ and $E_n := E \cap \{x : n-1 < \|x\|_2 \leq n\}$ for $n \geq 2$. For each $n$, choose open $G_n \supseteq E_n$ with $m_p^*(G_n \setminus E_n) < \varepsilon 2^{-n}$; then $G := \bigcup_n G_n$ is open, $E \subseteq G$, and $G \setminus E \subseteq \bigcup_n (G_n \setminus E_n)$, so $m_p^*(G \setminus E) < \varepsilon$.
(ii) $\Rightarrow$ (iii). Choose open $G_n \supseteq E$ with $m_p^*(G_n \setminus E) < 1/n$, and set $H := \bigcap_n G_n$.
Then $H$ is $G_\delta$, $E \subseteq H$, and $H \setminus E \subseteq G_n \setminus E$ for all $n$; hence $m_p^*(H \setminus E) = 0$.
(iii) $\Rightarrow$ (i). Since $H$ is $G_\delta$, $H \in \mathcal{B}(\mathbb{R}^p) \subseteq \mathcal{L}^p$. Also $m_p^*(H \setminus E) = 0$ implies $H \setminus E \in \mathcal{L}^p$ by Proposition 3.3.8.
Thus $E = H \setminus (H \setminus E) \in \mathcal{L}^p$.
(iii) $\Leftrightarrow$ (iv). If (iii) holds, set $Z := H \setminus E$; then $Z \subseteq H$, $m(Z) = 0$, and $E = H \setminus Z$.
Conversely, if (iv) holds, then $E \subseteq H$ and $H \setminus E = Z$, so $m_p^*(H \setminus E) = 0$.

*Structural Commentary*: Lebesgue measurable sets are $G_\delta$ sets with null material removed.
*Key Insight*: Small open excess becomes a null $G_\delta$ excess after countably many approximations. □

**Corollary 6.1.4.** *Open Outer Regularity for Measurable Sets*
*If $E \in \mathcal{L}^p$, then*
*(i) $\forall \varepsilon > 0$, $\exists$ open $G \supseteq E$ such that $m(G) \leq m(E) + \varepsilon$;*
*(ii) $m(E) = \inf\{m(G) : E \subseteq G,\ G \subseteq \mathbb{R}^p \text{ open}\}$.*

*Proof.*

(i) By Theorem 6.1.3, choose open $G \supseteq E$ with $m_p^*(G \setminus E) < \varepsilon$.
If $m(E) < +\infty$, then $m(G) = m(E) + m(G \setminus E) \leq m(E) + \varepsilon$; if $m(E) = +\infty$, the inequality is trivial.
(ii) From (i), $\inf_G m(G) \leq m(E)$. Conversely, $E \subseteq G$ gives $m(E) \leq m(G)$ by monotonicity; take $\inf_G$.

*Structural Commentary*: Measurable sets are measured from outside by open supersets.
*Key Insight*: Open excess controls the outer regular infimum. □

**Theorem 6.1.5.** *Closed and $F_\sigma$ Characterizations of Measurability*
*For $E \subseteq \mathbb{R}^p$, the following are equivalent:*
*(i) $E \in \mathcal{L}^p$;*
*(ii) $\forall \varepsilon > 0$, $\exists$ closed $F \subseteq E$ such that $m_p^*(E \setminus F) < \varepsilon$;*
*(iii) $\exists$ $F_\sigma$ set $K \subseteq E$ such that $m_p^*(E \setminus K) = 0$;*
*(iv) $\exists$ $F_\sigma$ set $K \subseteq E$ and $Z \in \mathcal{L}^p$ such that $Z \subseteq E$, $m(Z) = 0$, and $E = K \cup Z$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*). Since $E^c \in \mathcal{L}^p$, Theorem 6.1.3 gives open $G \supseteq E^c$ with $m_p^*(G \setminus E^c) < \varepsilon$. Set $F := G^c$. Then $F$ is closed, $F \subseteq E$, and $E \setminus F = G \setminus E^c$, so $m_p^*(E \setminus F) < \varepsilon$.

(*ii*) $\Rightarrow$ (*iii*). Choose closed $F_n \subseteq E$ with $m_p^*(E \setminus F_n) < 1/n$, and set $K := \bigcup_n F_n$. Then $K$ is $F_\sigma$, $K \subseteq E$, and $E \setminus K \subseteq E \setminus F_n$ for all $n$; hence $m_p^*(E \setminus K) = 0$.

(*iii*) $\Rightarrow$ (*i*). Since $K$ is $F_\sigma$, $K \in \mathcal{B}(\mathbb{R}^p) \subseteq \mathcal{L}^p$. Also $m_p^*(E \setminus K) = 0$ implies $E \setminus K \in \mathcal{L}^p$ by Proposition 3.3.8. Thus $E = K \cup (E \setminus K) \in \mathcal{L}^p$.

(*iii*) $\Leftrightarrow$ (*iv*). If (iii) holds, set $Z := E \setminus K$; then $Z \subseteq E$, $m(Z) = 0$, and $E = K \cup Z$. Conversely, if (iv) holds, then $E \setminus K \subseteq Z$, so $m_p^*(E \setminus K) = 0$.

*Structural Commentary*: Inside approximation is the complement of outside approximation.
*Key Insight*: Measurable sets are also $F_\sigma$ sets with null material added. □

**Corollary 6.1.6.** *Closed Inner Regularity*
*If $E \in \mathcal{L}^p$, then*

(*i*) $\forall \varepsilon > 0$, $\exists$ *closed* $F \subseteq E$ *such that* $m(E) \leq m(F) + \varepsilon$;
(*ii*) $m(E) = \sup\{m(F) : F \subseteq E,\ F \subseteq \mathbb{R}^p \text{ closed}\}$.

*Proof.*

(i) By Theorem 6.1.5, choose closed $F \subseteq E$ with $m_p^*(E \setminus F) < \varepsilon$. Since $F \in \mathcal{L}^p$ and $E = F \sqcup (E \setminus F)$, $m(E) = m(F) + m(E \setminus F) \leq m(F) + \varepsilon$.

(ii) From (i), $m(E) \leq \sup_F m(F)$. Conversely, $F \subseteq E$ gives $m(F) \leq m(E)$; take $\sup_F$.

*Structural Commentary*: Measurable sets are measured from inside by closed subsets.
*Key Insight*: Small discarded measure controls the inner regular supremum. □

**Corollary 6.1.7.** *Lebesgue Regularity and Borel Completion on $\mathbb{R}$*
*For $A \in \mathcal{L}^1$,*

(*i*) $m(A) = \inf\{m(G) : A \subseteq G,\ G \text{ open in } \mathbb{R}\}$.
(*ii*) $m(A) = \sup\{m(K) : K \subseteq A,\ K \text{ compact in } \mathbb{R}\}$, *finite-measure case applied to* $A \cap (-n, n]$; $m(A) = +\infty$.
(*iii*) *There exists* $B \in \mathcal{B}(\mathbb{R})$ *with* $A \subseteq B$ *and* $m(B \setminus A) = 0$; *equivalently,* $\mathcal{L}^1$ *is the completion of* $\mathcal{B}(\mathbb{R})$.

*Proof.*

(i) For $\varepsilon > 0$, regular approximation gives open $G \supseteq A$ with $m(G \setminus A) < \varepsilon$, hence $m(G) \leq m(A) + \varepsilon$. Monotonicity gives the reverse inequality.

(ii) If $m(A) < +\infty$, choose closed $F \subseteq A$ with $m(A \setminus F) < \varepsilon$. Since $F \cap [-n, n] \uparrow F$, continuity from below gives compact $K = F \cap [-n, n]$ with $m(F) - m(K) < \varepsilon$, so $m(A) - m(K) < 2\varepsilon$. If $m(A) = +\infty$, apply the finite case to $A_n = A \cap (-n, n]$ and let $n \to \infty$.

(iii) Choose open $G_k \supseteq A$ with $m(G_k \setminus A) < 1/k$, and set $B = \bigcap_k G_k$. Then $B \in \mathcal{B}(\mathbb{R})$, $A \subseteq B$, and $m(B \setminus A) = 0$. Thus every Lebesgue set is Borel plus a null subset. Conversely, Borel sets and null subsets are Lebesgue measurable.

*Structural Commentary*: Lebesgue sets are Borel sets plus null-set ambiguity.
*Key Insight*: Open outer approximation and compact inner approximation are the usable form of regularity. □




Figure 199: Regular approximation sandwiches $E$ between $F$ and $G$ with small excess.

**Theorem 6.1.8.** *Lebesgue Measurable Non-Borel Sets Exist*

(*i*) *There exists* $E \subseteq \mathbb{R}$ *such that* $E \in \mathcal{L}^1$ *and* $E \notin \mathcal{B}(\mathbb{R})$.
(*ii*) $\mathcal{B}(\mathbb{R}) \subsetneq \mathcal{L}^1$.

*Proof.*

(i) Lemma 4.1.7 gives $|\mathcal{B}(\mathbb{R})| \leq \mathfrak{c}$. Let $C$ be the Cantor set. Since $|C| = \mathfrak{c}$ and $m(C) = 0$, Proposition 3.3.8 gives $\mathcal{P}(C) \subseteq \mathcal{L}^1$, so $\left|\mathcal{L}^1\right| \geq 2^{\mathfrak{c}} > \mathfrak{c}$. Hence some $E \in \mathcal{L}^1$ is not Borel.

(ii) Proposition 4.1.11 gives $\mathcal{B}(\mathbb{R}) \subseteq \mathcal{L}^1$, and (i) makes the containment proper.

*Structural Commentary*: Countable generation is smaller than completion by null subsets.
*Key Insight*: Completeness creates more measurable sets than topology can generate. □

## 6.2 Compact and Cell Approximation

**Theorem 6.2.1.** *Compact Approximation at Finite Outer Measure*
*Let $E \subseteq \mathbb{R}^p$ with $m_p^*(E) < +\infty$. Then $E \in \mathcal{L}^p$ iff $\forall \varepsilon > 0$, $\exists$ compact $K \subseteq E$ with $m_p^*(E \setminus K) < \varepsilon$.*

*Proof.*

($\Rightarrow$) Since $m(E) < +\infty$, choose closed $F \subseteq E$ with $m(E \setminus F) < \frac{\varepsilon}{2}$ by Theorem 6.1.5.
Let $Q_N := [-N, N]^p$. Since $F \cap Q_N \uparrow F$, continuity from below gives $N$ with $m(F \setminus Q_N) < \frac{\varepsilon}{2}$.
Then $K := F \cap Q_N$ is compact, $K \subseteq E$, and $m_p^*(E \setminus K) = m(E \setminus K) \leq m(E \setminus F) + m(F \setminus Q_N) < \varepsilon$.

($\Leftarrow$) Choose compact $K_n \subseteq E$ with $m_p^*(E \setminus K_n) < \frac{1}{n}$ and set $K := \bigcup_{n \geq 1} K_n$.
Then $K$ is $F_\sigma$, hence $K \in \mathcal{L}^p$, and $E \setminus K \subseteq E \setminus K_n$ for all $n$, so $m_p^*(E \setminus K) = 0$.
By Proposition 3.3.8, $E \setminus K \in \mathcal{L}^p$, hence $E = K \cup (E \setminus K) \in \mathcal{L}^p$.

*Structural Commentary*: Finite measure permits bounded truncation before inner closure.
*Key Insight*: Compact approximation is closed approximation plus a finite-measure tail estimate. □

**Theorem 6.2.2.** *Finite Cell Approximation in Symmetric Difference*
*If $E \in \mathcal{L}^p$ and $m(E) < +\infty$, then $\forall \varepsilon > 0$, $\exists U = \bigcup_{j=1}^{N} I_j$ with $I_j$ bounded open cells and $m(E \triangle U) < \varepsilon$.*

*Proof.* Choose compact $K \subseteq E$ with $m(E \setminus K) < \frac{\varepsilon}{2}$ by Theorem 6.2.1, and open $G \supseteq E$ with $m_p^*(G \setminus E) < \frac{\varepsilon}{2}$ by Theorem 6.1.3. For each $x \in K$, choose a bounded open cell $I_x$ with $x \in I_x$ and $\overline{I_x} \subseteq G$. Compactness gives $I_1, \ldots, I_N$ with $K \subseteq U := \bigcup_{j=1}^{N} I_j \subseteq G$. Then $E \triangle U \subseteq (E \setminus K) \cup (G \setminus E)$, so subadditivity gives $m(E \triangle U) < \varepsilon$.

*Structural Commentary*: Compactness turns a countable open cover into a finite geometric approximant.
*Key Insight*: Approximate from outside by cells and from inside by a compact core. □

**Corollary 6.2.3.** *Cell Approximation Variant Choices*
*Let $E \in \mathcal{L}^p$ with $m(E) < +\infty$, and let $\varepsilon > 0$. There exists $A \subseteq \mathbb{R}^p$ with $m(E \triangle A) < \varepsilon$ in each format:*

*(i) $A$ is a finite union of bounded open cells;*
*(ii) $A$ is a finite union of bounded closed cells;*
*(iii) $A$ is a finite union of bounded half-open cells of a fixed orientation.*

*Proof.*

(i) By Theorem 6.2.2, choose $A = \bigcup_{j=1}^{N} I_j$ with $I_j$ bounded open cells and $m(E \triangle A) < \varepsilon$.

(ii) Choose a finite open-cell union $U = \bigcup_{j=1}^{M} J_j$ with $m(E \triangle U) < \frac{\varepsilon}{2}$. If $M = 0$, take $A = \emptyset$; otherwise choose closed cells $C_j \subseteq J_j$ with $m(J_j \setminus C_j) < \frac{\varepsilon}{2M}$ and set $A := \bigcup_{j=1}^{M} C_j$. Then $m(U \setminus A) < \frac{\varepsilon}{2}$, so $m(E \triangle A) < \varepsilon$.

(iii) Choose a finite open-cell union $U$ with $m(E \triangle U) < \varepsilon$. Refine by the finite coordinate grid from the endpoints of the open cells, and let $A$ be the union of the half-open grid atoms contained in $U$.
Then $U \triangle A$ is contained in finitely many cell faces, each outer-null, so $m(E \triangle A) < \varepsilon$.

*Structural Commentary*: Cell format changes only boundary-null pieces.
*Key Insight*: A compact core inside an open envelope permits finite cell variants. □

**Corollary 6.2.4.** *Real-Line Finite Open-Interval Approximation*
*If $E \in \mathcal{L}^1$ and $m(E) < +\infty$, then $\forall \varepsilon > 0$, $\exists O = \bigcup_{j=1}^{N} I_j$ with $I_j$ pairwise disjoint bounded open intervals and $m(E \setminus O) + m(O \setminus E) = m(E \triangle O) < \varepsilon$.*

*Proof.* By Theorem 6.2.2 with $p = 1$, choose a finite union $U$ of bounded open intervals with $m(E \triangle U) < \varepsilon$. Write $U = O = \bigcup_{j=1}^{N} I_j$ as the disjoint union of its open interval components.
Since $E \setminus O$ and $O \setminus E$ are disjoint measurable sets, additivity gives $m(E \triangle O) = m(E \setminus O) + m(O \setminus E) < \varepsilon$.
*Structural Commentary*: Merge finite open intervals into disjoint components.
*Key Insight*: A compact core makes the countable cover finite; interval merging makes it disjoint. □

## 6.3 Caratheodory Tests and Inner Measure

**Theorem 6.3.1.** *Outer Measure Additivity with One Measurable Set*
*Let $E \in \mathcal{L}^p$ and $F \subseteq \mathbb{R}^p$. Then $m_p^*(E \cup F) + m_p^*(E \cap F) = m(E) + m_p^*(F)$.*

*Proof.* Since $E \in \mathcal{L}^p$, Caratheodory splitting: $m_p^*(E \cup F) = m(E) + m_p^*(F \setminus E)$; $m_p^*(F) = m_p^*(E \cap F) + m_p^*(F \setminus E)$.
Hence $m_p^*(E \cup F) + m_p^*(E \cap F) = m(E) + m_p^*(F)$.
*Structural Commentary*: One measurable cut splits arbitrary sets exactly.
*Key Insight*: Split both $E \cup F$ and $F$ along the same measurable cut $E$. □

**Corollary 6.3.2.** *Disjoint Additivity and Difference*
*Let $E \in \mathcal{L}^p$ and $F \subseteq \mathbb{R}^p$.*

*(i)* $E \cap F = \emptyset \implies m_p^*(E \cup F) = m(E) + m_p^*(F)$.
*(ii)* $E \subseteq F,\ m(E) < +\infty \implies m_p^*(F \setminus E) = m_p^*(F) - m(E)$.

*Proof.*

(i) By Theorem 6.3.1, $m_p^*(E \cup F) + m_p^*(E \cap F) = m(E) + m_p^*(F)$.
Since $E \cap F = \emptyset$ and $m_p^*(\emptyset) = 0$, $m_p^*(E \cup F) = m(E) + m_p^*(F)$.
(ii) Since $E \subseteq F$, Caratheodory splitting gives $m_p^*(F) = m(E) + m_p^*(F \setminus E)$; subtract $m(E) < +\infty$.

*Structural Commentary*: Exact splitting yields disjoint sums and finite differences.
*Key Insight*: Disjointness removes the overlap; containment isolates the remainder. □

**Remark 6.3.3.** *Excision Needs Measurability*
Let $E \subseteq \mathbb{R}^p$ with $m_p^*(E) < +\infty$. Open approximation may give, $\forall \varepsilon > 0$, an open $G \supseteq E$ with $m(G) - m_p^*(E) < \varepsilon$, but this does not imply $m_p^*(G \setminus E) < \varepsilon$. If $E \in \mathcal{L}^p$, Corollary 6.3.2 gives $m_p^*(G \setminus E) = m(G) - m(E)$.
*Key Insight*: Small open excess becomes small set difference only after $E$ is a Caratheodory splitter.

**Theorem 6.3.4.** *Finite Outer Measure Inner Test*
*If $m_p^*(E) < +\infty$, then $E \in \mathcal{L}^p \iff \exists B \in \mathcal{L}^p,\ B \subseteq E,\ m(B) = m_p^*(E)$.*

*Proof.*

($\Rightarrow$) Set $B := E$. Then $B \in \mathcal{L}^p$, $B \subseteq E$, and $m(B) = m(E) = m_p^*(E)$.
($\Leftarrow$) Assume $B \in \mathcal{L}^p$, $B \subseteq E$, and $m(B) = m_p^*(E)$.
Since $m(B) < +\infty$, the difference formula gives $m_p^*(E \setminus B) = m_p^*(E) - m(B) = 0$.
Thus $E \setminus B \in \mathcal{L}^p$, and $E = B \cup (E \setminus B) \in \mathcal{L}^p$.

*Structural Commentary*: Exact inner mass leaves only a null remainder.
*Key Insight*: Equality of outer mass converts containment into measurability. □

**Theorem 6.3.5.** *Single Finite Container Test*
*Let $A \in \mathcal{L}^p$, $m(A) < +\infty$, and $E \subseteq A$. Then $E \in \mathcal{L}^p \iff m(A) = m_p^*(E) + m_p^*(A \setminus E)$.*

*Proof.*

($\Rightarrow$) Since $E \in \mathcal{L}^p$ and $E \subseteq A$,
Caratheodory splitting of $A$ by $E$ gives $m(A) = m(E) + m(A \setminus E) = m_p^*(E) + m_p^*(A \setminus E)$.
($\Leftarrow$) Choose $G_\delta$ set $H \supseteq A \setminus E$ with $m(H) = m_p^*(A \setminus E)$. Since $A \setminus E \subseteq A \cap H \subseteq H$, $m(A \cap H) = m_p^*(A \setminus E)$.
Set $B := A \setminus H$. $B \in \mathcal{L}^p$, $B \subseteq E$; $m(B) = m(A) - m(A \cap H) = m_p^*(E)$. By Theorem 6.3.4, $E \in \mathcal{L}^p$.

*Structural Commentary*: Inside a finite measurable container, Caratheodory only needs to be tested once.
*Key Insight*: Outer-approximate the complement; the remainder has full outer mass. □

**Corollary 6.3.6.** *Bounded Inner Measure Test*
*For bounded cell $Q \subseteq \mathbb{R}^p$ and $E \subseteq Q$, define $m_{*,Q}(E) := m(Q) - m_p^*(Q \setminus E)$. Then $E \in \mathcal{L}^p \iff m_{*,Q}(E) = m_p^*(E)$.*

*Proof.*

($\Rightarrow$) If $E \in \mathcal{L}^p$, then $m(Q) = m(E) + m_p^*(Q \setminus E)$, hence $m_{*,Q}(E) = m_p^*(E)$.

($\Leftarrow$) If $m_{*,Q}(E) = m_p^*(E)$, then $m(Q) = m_p^*(E) + m_p^*(Q \setminus E)$, so $E \in \mathcal{L}^p$ by the finite-container test.
*Structural Commentary*: Bounded inner measure tests one finite container.
*Key Insight*: Equality of inner and outer measure is a bounded version of Caratheodory. □

# 7 Vitali, Steinhaus, and Nonmeasurable Sets

## 7.1 Vitali, Steinhaus, and Nonmeasurability

**Definition 7.1.1.** *Difference Sets and Vitali Selectors*

(i) *Difference set*: for $A \subseteq \mathbb{R}^p$, $A - A := \{x - y : x, y \in A\}$; $A \subseteq B \implies A - A \subseteq B - B$.
(ii) *Rational equivalence*: $x \sim_{\mathbb{Q}^p} y \iff x - y \in \mathbb{Q}^p$; equivalently, $[x]_{\mathbb{Q}^p} = x + \mathbb{Q}^p$.
(iii) *Vitali selector*: choose $\mathcal{V} \subseteq \mathbb{R}^p$ such that $|\mathcal{V} \cap (x + \mathbb{Q}^p)| = 1$ for every $x \in \mathbb{R}^p$.
(iv) *Rational translate*: for $q \in \mathbb{Q}^p$, $\mathcal{V}_q := q + \mathcal{V}$.

**Lemma 7.1.2.** *Steinhaus Lemma for Compact Sets*
*If $K \subseteq \mathbb{R}^p$ is compact and $m(K) > 0$, then $\exists \delta > 0$ such that $B(0, \delta) \subseteq K - K$.*

*Proof.* Choose open $G \supseteq K$ with $m(G) < 2m(K)$. Since $K$ is compact and $G^c$ is closed, set $\delta := \operatorname{dist}(K, G^c) > 0$.
If $\|x\|_2 < \delta$, then $x + K \subseteq G$. If $(x + K) \cap K = \emptyset$, translation invariance and additivity give
$2m(K) = m(x + K) + m(K) = m((x + K) \cup K) \leq m(G) < 2m(K)$, a contradiction.
Hence $(x + K) \cap K \neq \emptyset$, so $x = k_1 - k_2 \in K - K$. Thus $B(0, \delta) \subseteq K - K$.
*Structural Commentary*: A positive compact set overlaps all sufficiently small translates of itself.
*Key Insight*: Disjoint small translates cannot fit inside a tight open envelope. □

**Theorem 7.1.3.** *Steinhaus Theorem*
*If $E \in \mathcal{L}^p$ and $m(E) > 0$, then $\exists \delta > 0$ such that $B(0, \delta) \subseteq E - E$.*

*Proof.* Set $E_n := E \cap B(0, n)$. Since $E_n \uparrow E$ and $m(E) > 0$, choose $n$ with $0 < m(E_n) < +\infty$.
By compact inner approximation, choose compact $K \subseteq E_n$ with $m(K) > 0$.
Lemma 7.1.2 gives $\delta > 0$ with $B(0, \delta) \subseteq K - K \subseteq E - E$.
*Structural Commentary*: Positive measurable sets contain positive compact cores.
*Key Insight*: Difference-set thickness follows from compact inner regularity. □

**Lemma 7.1.4.** *Bounded Disjoint Translates Force Nullity*
*Let $E \in \mathcal{L}^1$ and $\Lambda \subseteq \mathbb{R}$ be bounded, with $E$ bounded and $\Lambda$ countably infinite.*
*If $(\lambda + E)_{\lambda \in \Lambda}$ are pairwise disjoint, then $m(E) = 0$.*

*Proof.* Since $E$ and $\Lambda$ are bounded, $U := \bigcup_{\lambda \in \Lambda} (\lambda + E)$ is bounded, so $m(U) < +\infty$.
By disjointness, countable additivity, and translation invariance, $m(U) = \sum_{\lambda \in \Lambda} m(\lambda + E) = \sum_{\lambda \in \Lambda} m(E)$.
Since $\Lambda$ is countably infinite, finiteness forces $m(E) = 0$.
*Structural Commentary*: Infinitely many disjoint bounded translates cannot all carry positive equal mass.
*Key Insight*: Boundedness gives a finite container; translation invariance makes every summand the same. □

**Proposition 7.1.5.** *Real-Line Vitali Choice Sets*
*Let $E \subseteq \mathbb{R}$ with $m_1^*(E) > 0$, and write $x \sim y$ iff $x - y \in \mathbb{Q}$.*

*(i) There exists bounded $E_0 \subseteq E$ with $m_1^*(E_0) > 0$.*
*(ii) If $C \subseteq E_0$ contains exactly one point from each $\sim$-class meeting $E_0$, then $C \notin \mathcal{L}^1$.*

*In particular, $E$ contains a nonmeasurable subset.*

*Proof.*

(i) Since $E = \bigcup_{n \geq 1} (E \cap [-n, n])$, countable subadditivity gives $n$ with $E_0 := E \cap [-n, n]$ satisfying $m_1^*(E_0) > 0$.

(ii) Choose $b > 0$ with $E_0 \subseteq [-b, b]$ and set $\Lambda_0 := [-2b, 2b] \cap \mathbb{Q}$. Suppose $C \in \mathcal{L}^1$.
From $\lambda + c = \lambda' + c'$ with $\lambda, \lambda' \in \Lambda_0$ and $c, c' \in C$, we get $c - c' = \lambda' - \lambda \in \mathbb{Q}$; selector uniqueness gives $c = c'$ and $\lambda = \lambda'$. Thus $(\lambda + C)_{\lambda \in \Lambda_0}$ is pairwise disjoint, and Lemma 7.1.4 gives $m(C) = 0$.
For every $x \in E_0$, choose $c \in C$ with $x - c \in \mathbb{Q}$; since $x, c \in [-b, b]$, $x - c \in \Lambda_0$, so $E_0 \subseteq \bigcup_{\lambda \in \Lambda_0} (\lambda + C)$.
The right side is a countable union of null measurable sets, contradicting $m_1^*(E_0) > 0$.

*Structural Commentary*: Bounded rational shifts force the contradiction.
*Key Insight*: Selectors are either null under disjoint shifts or cover too much by null shifts. □

**Lemma 7.1.6.** *Rational Translates of a Vitali Selector Partition $\mathbb{R}^p$*
*Let $(q_i)_{i\geq 1}$ enumerate $\mathbb{Q}^p$.*

*(i)* $\mathbb{R}^p = \bigcup_{i=1}^{\infty} \mathcal{V}_{q_i}$.
*(ii) Each $x \in \mathbb{R}^p$ has a unique representation $x = q_i + v$ with $i \in \mathbb{N}$ and $v \in \mathcal{V}$; hence $\mathbb{R}^p = \bigsqcup_{i=1}^{\infty} \mathcal{V}_{q_i}$.*

*Proof.*

(i) For $x \in \mathbb{R}^p$, choose the unique $v \in \mathcal{V} \cap (x + \mathbb{Q}^p)$. Then $x - v = q_i$ for some $i$, so $x = q_i + v \in \mathcal{V}_{q_i}$.
(ii) If $q_i + v = q_j + w$ with $v, w \in \mathcal{V}$, then $v - w = q_j - q_i \in \mathbb{Q}^p$.
Selector uniqueness gives $v = w$, hence $q_i = q_j$ and $i = j$.

*Structural Commentary*: Rational translates enumerate the equivalence classes selected by $\mathcal{V}$.
*Key Insight*: One representative per class makes the rational translates disjoint and exhaustive. □

**Theorem 7.1.7.** *Vitali Selectors Are Nonmeasurable*
*If $\mathcal{V} \subseteq \mathbb{R}^p$ is a Vitali selector, then $\mathcal{V} \notin \mathcal{L}^p$.*

*Proof.* Assume $\mathcal{V} \in \mathcal{L}^p$.

(i) If $m(\mathcal{V}) > 0$, Steinhaus gives $\delta > 0$ with $B(0,\delta) \subseteq \mathcal{V} - \mathcal{V}$. Choose $0 \neq q \in \mathbb{Q}^p \cap B(0,\delta)$. Then $q = v_1 - v_2$ for some $v_1, v_2 \in \mathcal{V}$, so $v_1 \sim_{\mathbb{Q}^p} v_2$; selector uniqueness gives $v_1 = v_2$, hence $q = 0$, contradiction.
(ii) If $m(\mathcal{V}) = 0$, then each translate $\mathcal{V}_{q_i}$ is null. By Lemma 7.1.6, $\mathbb{R}^p = \bigsqcup_i \mathcal{V}_{q_i}$, so every bounded cell is contained in a countable union of null sets and has measure 0, contradicting positive cell volume.

*Structural Commentary*: Positive and zero measure are both impossible.
*Key Insight*: Steinhaus forbids positive measure; countable translated partition forbids zero measure. □

**Theorem 7.1.8.** *Nonmeasurable Subsets Are Everywhere*
*If $E \subseteq \mathbb{R}^p$ and $m_p^*(E) > 0$, then $\exists N \subseteq E$ with $N \notin \mathcal{L}^p$.*

*Proof.* Let $(q_i)$ enumerate $\mathbb{Q}^p$ and set $E_i := E \cap \mathcal{V}_{q_i}$.
If $E_i \in \mathcal{L}^p$ and $m(E_i) > 0$, then $E_i - E_i$ contains a ball about 0, so $0 \neq q \in \mathbb{Q}^p \cap (E_i - E_i)$.
But $E_i \subseteq q_i + \mathcal{V}$, so $q = (q_i + v_1) - (q_i + v_2) = v_1 - v_2$ with $v_1, v_2 \in \mathcal{V}$, contradicting selector uniqueness.
Hence each measurable $E_i$ is null. If every $E_i$ were measurable, then $E = \bigcup_i E_i$ would satisfy $m_p^*(E) = 0$, contradiction. Thus some $E_i \subseteq E$ is nonmeasurable.
*Structural Commentary*: Any positive outer-measure set meets the Vitali partition in a nonmeasurable piece.
*Key Insight*: Vitali pieces cannot be measurable and positive. □

**Corollary 7.1.9.** *Nonadditive Decompositions*

*(i) If $E \in \mathcal{L}^p$ and $0 < m(E) < +\infty$, then $\exists B, C \notin \mathcal{L}^p$ such that $E = B \sqcup C$ and $m(E) < m_p^*(B) + m_p^*(C)$.*
*(ii) If $B \notin \mathcal{L}^p$, $m_p^*(B) < +\infty$, and $H$ is $G_\delta$ with $B \subseteq H$, $m(H) = m_p^*(B)$, then $C := H \setminus B \notin \mathcal{L}^p$ and $m(H) = m_p^*(B \cup C) < m_p^*(B) + m_p^*(C)$.*

*Proof.*

(i) Pick $B \subseteq E$, $B \notin \mathcal{L}^p$, and put $C := E \setminus B$. Then $C \notin \mathcal{L}^p$, else $B = E \setminus C \in \mathcal{L}^p$.
Subadditivity gives $\leq$; equality would force $B \in \mathcal{L}^p$ by the finite-container test in $E$.
(ii) Since $H \in \mathcal{L}^p$, $C \in \mathcal{L}^p$ would give $B = H \setminus C \in \mathcal{L}^p$, contradiction. Also $m_p^*(C) > 0$, else $C \in \mathcal{L}^p$.
Thus $B \cup C = H$ and $m(H) = m_p^*(B)$ give $m_p^*(B \cup C) = m(H) < m_p^*(B) + m_p^*(C)$.

*Structural Commentary*: Nonmeasurability appears exactly as strict failure of additive splitting.
*Key Insight*: Equality in a finite measurable container is a measurability test. □

**Theorem 7.1.10.** *A Nonmeasurable Set with Only Null Measurable Pieces*
*There exists $\mathcal{U} \subseteq \mathbb{R}^p$ such that:*

*(i) if $F \in \mathcal{L}^p$ and $F \subseteq \mathcal{U}$, then $m(F) = 0$;*
*(ii) if $F \in \mathcal{L}^p$ and $F \subseteq \mathcal{U}^c$, then $m(F) = 0$;*
*(iii) $\mathcal{U} \notin \mathcal{L}^p$.*

*Proof.* Let $e := (1, \ldots, 1)$, $G_0 := \mathbb{Q}^p + (2\sqrt{2})\mathbb{Z}e$, $G_1 := \sqrt{2}e + G_0$, $G := G_0 \sqcup G_1 = \mathbb{Q}^p + \sqrt{2}\mathbb{Z}e$; $G_0, G_1$ dense.
Let $\mathcal{E}$ select one representative from each class modulo $G$, and set $\mathcal{U} := G_0 + \mathcal{E}$.
Then $\mathbb{R}^p = (G_0 + \mathcal{E}) \sqcup (G_1 + \mathcal{E})$ and $\mathcal{U}^c = G_1 + \mathcal{E} = \sqrt{2}e + \mathcal{U}$.

(i) Suppose $F \in \mathcal{L}^p$, $F \subseteq \mathcal{U}$, and $m(F) > 0$. By Steinhaus, $B(0,r) \subseteq F - F$ for some $r > 0$; choose $g_1 \in G_1 \cap B(0,r)$. Since $F \subseteq G_0 + \mathcal{E}$, $g_1 = (g_0 + s_1) - (g_0' + s_2)$ for some $g_0, g_0' \in G_0$ and $s_1, s_2 \in \mathcal{E}$. Thus $s_1 - s_2 = g_1 - g_0 + g_0' \in G$, so $s_1 = s_2$ and $g_1 = g_0 - g_0' \in G_0$, contradicting $G_0 \cap G_1 = \emptyset$.
(ii) If $F \in \mathcal{L}^p$ and $F \subseteq \mathcal{U}^c = \sqrt{2}\,e + \mathcal{U}$, then $F - \sqrt{2}\,e \subseteq \mathcal{U}$ and $m(F) = m(F - \sqrt{2}\,e) = 0$ by (i).

(iii) If $\mathcal{U} \in \mathcal{L}^p$, then (i)–(ii) give $m(\mathcal{U}) = m(\mathcal{U}^c) = 0$, so $m([0,1]^p) = 0$, contradiction.

*Structural Commentary*: This construction partitions representatives using two dense cosets.
*Key Insight*: Positive measurable pieces force forbidden coset differences. □

# 8 Cantor–Lebesgue Singular Function

## 8.1 Cantor-Lebesgue Singular Function Construction

**Definition 8.1.1.** *Cantor–Lebesgue Singular Function*
Let $P$ be the Cantor set. For $k \geq 1$, let $I_{k,1}, \ldots, I_{k,2^k-1}$ list, left to right, the components of the stage-$k$ removed set $U_k$. Set $U := \bigcup_{k\geq 1} U_k = [0,1] \setminus P$.

(i) For $x \in U$, choose $k \geq 1$ and $1 \leq j \leq 2^k - 1$ with $x \in I_{k,j}$, and set $\varphi(x) := \frac{j}{2^k}$.
(ii) This value is independent of the chosen stage: if $I_{k,j} = I_{\ell,m}$ with $\ell \geq k$, then $m = 2^{\ell-k} j$ and $\frac{m}{2^\ell} = \frac{j}{2^k}$.
(iii) On $P$, set $\varphi(0) := 0$ and $\varphi(x) := \sup\{\varphi(t) : t \in U \cap [0,x]\}$ for $x \in P \setminus \{0\}$.

The resulting $\varphi : [0,1] \to [0,1]$ is the *Cantor–Lebesgue singular function*.




Figure 200: Horizontal segments show the Cantor–Lebesgue plateaus on the intervals removed through stage 3.

**Proposition 8.1.2.** *Properties of the Singular Function*
*Let $P$ be the Cantor set, $U := [0,1] \setminus P$, and $\varphi : [0,1] \to [0,1]$ the Cantor–Lebesgue singular function. Then:*

*(i) $\varphi$ is nondecreasing.*
*(ii) $\varphi$ is continuous on $[0,1]$.*
*(iii) $\varphi([0,1]) = [0,1]$.*
*(iv) $\varphi'(x) = 0$ for every $x \in U$.*
*(v) $m(U) = 1$.*

*Proof.*

(i) At each stage $k$, the components $I_{k,1}, \ldots, I_{k,2^k-1}$ occur left to right and carry heights $\frac{1}{2^k}, \ldots, \frac{2^k-1}{2^k}$. Thus the prescription on $U$ is nondecreasing, and the supremum extension on $P$ preserves order.
(ii) If $x_0 \in U$, then $x_0 \in J$ for some component $J$ of $U$ and $\varphi|_J$ is constant. If $x_0 \in P$, then for each $k$ there are $a_k, b_k \in 2^{-k}\mathbb{Z} \cap [0,1]$ with $a_k \leq \varphi(x_0) \leq b_k$, $b_k - a_k \leq 2^{-k}$, and $\varphi(y) \in [a_k, b_k]$ for all $y$ near $x_0$. Thus $\limsup_{y\to x_0} |\varphi(y) - \varphi(x_0)| \leq 2^{-k}$ for every $k$, hence $\varphi$ is continuous at $x_0$.
(iii) The definition gives $\varphi(0) = 0$. For each $k$, the rightmost component of $U_k$ has height $1 - 2^{-k}$, so $\varphi(1) = 1$. By (ii) and the intermediate value theorem, $\varphi([0,1]) = [0,1]$.
(iv) For each component $J$ of $U$, $\varphi|_J$ is constant; hence $\varphi'(x) = 0$ for every $x \in U$.
(v) Proposition 4.1.14 gives $m(P) = 0$; since $[0,1] = P \sqcup U$, $m(U) = m([0,1]) - m(P) = 1$.

*Structural Commentary*: Plateaus on $U$ force variation onto $P$.
*Key Insight*: At stage $k$, neighboring plateau heights differ by $2^{-k}$. □

**Theorem 8.1.3.** *Lebesgue Measurable Non-Borel Sets Exist*
*There exist $A \subseteq [0,1]$ and a homeomorphism $\psi : [0,1] \to [0,2]$ such that $A \in \mathcal{L}^1 \setminus \mathcal{B}(\mathbb{R})$ and $\psi(A) \notin \mathcal{L}^1$.*

*Proof.* Let $P$ be the Cantor set, $U := [0,1] \setminus P$, $\varphi$ the Cantor–Lebesgue singular function, and $\psi := \mathrm{id}_{[0,1]} + \varphi$.
Proposition 8.1.2 gives $\varphi$ continuous and nondecreasing; hence $\psi$ is continuous and strictly increasing.
Since $\psi(0) = 0$ and $\psi(1) = 2$, $\psi : [0,1] \to [0,2]$ is a homeomorphism.
Let $(J_k)_{k\geq 1}$ be the components of $U$. Since $\varphi|_{J_k}$ is constant, $\psi(J_k)$ is a translate of $J_k$ and $m(\psi(J_k)) = m(J_k)$.

Thus $m(\psi(U)) = \sum_{k\geq 1} m(\psi(J_k)) = \sum_{k\geq 1} m(J_k) = m(U) = 1$.
Since $\psi(P)$ is compact and $[0,2] = \psi(P) \sqcup \psi(U)$, $m(\psi(P)) = 1$.
Choose $W \subseteq \psi(P)$ with $W \notin \mathcal{L}^1$ by Theorem 7.1.8, and set $A := \psi^{-1}(W)$. Then $A \subseteq P$ and $m(P) = 0$, so $A \in \mathcal{L}^1$. If $A \in \mathcal{B}(\mathbb{R})$, then $W = \psi(A) \in \mathcal{B}([0,2]) \subseteq \mathcal{L}^1$, contradiction. Hence $A \in \mathcal{L}^1 \backslash \mathcal{B}(\mathbb{R})$ and $\psi(A) = W \notin \mathcal{L}^1$.
*Structural Commentary*: Completion by null sets creates Lebesgue sets invisible to the Borel hierarchy.
*Key Insight*: A null Cantor set can be homeomorphically stretched to positive measure. □

# Part XIV
# Measure, Integration, and Functional Analysis

## 1 Measurable Functions and Lebesgue Integration

### 1.1 Measurable Functions

**Definition 1.1.1.** *Borel, Lebesgue, and Function Measurability*

(i) *Borel*: $\mathcal{B}(\mathbb{R}^p) := \sigma(\mathcal{O}_{\mathbb{R}^p})$; its elements are *Borel sets*.
(ii) *Lebesgue*: $\mathcal{B}(\mathbb{R}^p) \subseteq \mathcal{L}^p$, and $\mathcal{L}^p$ is obtained by adjoining subsets of Lebesgue null sets.
(iii) *Maps*: for measurable spaces $(X,\mathcal{M})$, $(Y,\mathcal{N})$, $f : X \to Y$ is *$(\mathcal{M},\mathcal{N})$-measurable* iff $\forall E \in \mathcal{N},\ f^{-1}(E) \in \mathcal{M}$.
(iv) *Borel maps*: $f : \mathbb{R}^p \to \mathbb{R}^q$ is *Borel measurable* iff $\forall B \in \mathcal{B}(\mathbb{R}^q),\ f^{-1}(B) \in \mathcal{B}(\mathbb{R}^p)$.
(v) *Thresholds*: write $\{f \bowtie a\} := \{x \in X : f(x) \bowtie a\}$ for $\bowtie \in \{>, \geq, <, \leq\}$.
Extended-real $f$ is *$\mathcal{M}$-measurable* iff $\forall a \in \mathbb{R},\ \{f > a\} \in \mathcal{M}$.

*Structural Commentary*: Borel measurability tracks topology; Lebesgue measurability adds null-set completion.
*Key Insight*: Measurability is inverse-image stability.

**Proposition 1.1.2.** *Generator Test for Measurable Maps*
*Let $f : X \to Y$, $\mathcal{A} \subseteq \mathcal{P}(Y)$, and $\mathcal{N} := \sigma(\mathcal{A})$.*

*(i) If $\forall A \in \mathcal{A},\ f^{-1}(A) \in \mathcal{M}$, then $f : X \to (Y,\mathcal{N})$ is measurable.*
*(ii) For real-valued $f$, $\forall a \in \mathbb{R},\ \{f > a\} \in \mathcal{M} \iff \forall B \in \mathcal{B}(\mathbb{R}),\ f^{-1}(B) \in \mathcal{M}$.*

*Proof.*

(i) Let $\mathcal{G} := \{E \subseteq Y : f^{-1}(E) \in \mathcal{M}\}$. Since $f^{-1}(E^c) = f^{-1}(E)^c$ and $f^{-1}(\bigcup_n E_n) = \bigcup_n f^{-1}(E_n)$, $\mathcal{G}$ is a $\sigma$-algebra. Thus $\mathcal{A} \subseteq \mathcal{G} \Longrightarrow \mathcal{N} = \sigma(\mathcal{A}) \subseteq \mathcal{G}$.
(ii) ($\Rightarrow$) Set $\mathcal{A} := \{(a,+\infty) : a \in \mathbb{R}\}$. Since $\forall a \in \mathbb{R},\ f^{-1}((a,+\infty)) = \{f > a\} \in \mathcal{M}$ and $\sigma(\mathcal{A}) = \mathcal{B}(\mathbb{R})$, (i) gives $\forall B \in \mathcal{B}(\mathbb{R}),\ f^{-1}(B) \in \mathcal{M}$.
(ii) ($\Leftarrow$) If $\forall B \in \mathcal{B}(\mathbb{R}),\ f^{-1}(B) \in \mathcal{M}$, then $\forall a \in \mathbb{R},\ \{f > a\} = f^{-1}((a,+\infty)) \in \mathcal{M}$.

*Structural Commentary*: Measurability can be checked on any generating family of target sets.
*Key Insight*: Pullbacks turn generated $\sigma$-algebras into generated collections of preimages. □

**Proposition 1.1.3.** *Open-Set Test for Euclidean Targets*
*Let $\Omega \subseteq \mathbb{R}^n$ be Lebesgue measurable and let $f : \Omega \to \mathbb{R}^m$. The following tests are equivalent:*

*(i) $\forall V \subseteq \mathbb{R}^m$ open, $f^{-1}(V)$ is measurable;*
*(ii) $\forall B \subseteq \mathbb{R}^m$ open boxes, $f^{-1}(B)$ is measurable;*
*(iii) $\forall j \in \{1,\dots,m\}$, $f_j : \Omega \to \mathbb{R}$ is threshold-measurable.*

*In particular, every continuous $f : \Omega \to \mathbb{R}^m$ is measurable.*

*Proof.*

(i) $\Rightarrow$ (ii). Open boxes are open.
(ii) $\Rightarrow$ (i). By Lemma 4.1.7, $V = \bigcup_k B_k$ for rational boxes $B_k$, hence $f^{-1}(V) = \bigcup_k f^{-1}(B_k)$ is measurable.
(ii) $\Rightarrow$ (iii). By (ii) $\Rightarrow$ (i), $\{f_j > a\} = f^{-1}(\{y : y_j > a\})$ is measurable.
(iii) $\Rightarrow$ (ii). If $B = \prod_{j=1}^m (a_j,b_j)$, then $f^{-1}(B) = \bigcap_{j=1}^m \big(\{f_j > a_j\} \cap \{f_j < b_j\}\big)$,
where $\{f_j < b_j\} = \Omega \setminus \bigcap_{\ell=1}^{\infty} \{f_j > b_j - \frac{1}{\ell}\}$. Thus each factor is measurable, hence so is $f^{-1}(B)$.

If $f$ is continuous, $f^{-1}(V)$ is open relative to $\Omega$ by Theorem 5.1.17.
By Theorem 3.2.2, $f^{-1}(V) = W \cap \Omega$ for some open $W \subseteq \mathbb{R}^n$, hence it is measurable by Theorem 4.1.8.
*Structural Commentary*: Euclidean-valued measurability is just countable box algebra applied to inverse images.
*Key Insight*: Open sets reduce to rational boxes; vector functions reduce to coordinate threshold tests. □

**Example 1.1.4.** *Basic Measurable Functions*
Let $(X,\mathcal{M})$ be a measurable space.

(i) Constant real-valued functions are measurable, since each strict superlevel set is either $X$ or $\emptyset$.
(ii) For $E \subseteq X$, the indicator $\mathbf{1}_E$ is measurable iff $E \in \mathcal{M}$.
(iii) Continuous maps between Euclidean measurable subsets are measurable by Proposition 1.1.3.
(iv) Monotone functions $\mathbb{R} \to \mathbb{R}$ are Borel measurable, since their strict superlevel sets are half-lines, $\mathbb{R}$, or $\emptyset$.

*Key Insight*: Constants, indicators of measurable sets, continuous functions, and monotone functions form the first reusable stock of measurable functions.

**Theorem 1.1.5.** *Closure Properties of Measurable Functions*
*Let $(X, \mathcal{M})$ be a measurable space.*

*(i) The four tests $\forall a \in \mathbb{R}$, $\{f > a\}, \{f \geq a\}, \{f < a\}, \{f \leq a\} \in \mathcal{M}$ are equivalent.*
*(ii) $f$ measurable implies $|f|$ measurable.*
*(iii) If $f_n$ are measurable, then $\sup_n f_n, \inf_n f_n, \limsup_n f_n, \liminf_n f_n$ are measurable.*
*(iv) If $f, g$ real-valued measurable, $F : \mathbb{R}^2 \to \mathbb{R}$ continuous, then $F(f, g)$ measurable; so are $f + g$ and $fg$.*

*Proof.*

(i) Use $\{f \geq a\} = \bigcap_{m \geq 1} \{f > a - \frac{1}{m}\}$; $\{f < a\} = X \setminus \{f \geq a\}$; $\{f \leq a\} = \bigcap_{m \geq 1} \{f < a + \frac{1}{m}\}$.
(ii) For $a > 0$, $\{|f| < a\} = \{f < a\} \cap \{f > -a\}$; for $a \leq 0$ the set is empty.
(iii) $\{\sup_n f_n > a\} = \bigcup_n \{f_n > a\}$. Also $\limsup_n f_n = \inf_m \sup_{n \geq m} f_n$; $\liminf_n f_n = \sup_m \inf_{n \geq m} f_n$; inf follow from (i).
(iv) For $G_a = \{(u, v) : F(u, v) > a\}$, write $G_a = \bigcup_k (\alpha_k, \beta_k) \times (\gamma_k, \delta_k)$.
Then $\{F(f, g) > a\} = \bigcup_k \big(\{\alpha_k < f < \beta_k\} \cap \{\gamma_k < g < \delta_k\}\big) \in \mathcal{M}$.

*Structural Commentary*: Measurability is countable set algebra applied to level sets.
*Key Insight*: Once threshold sets are measurable, ordinary limiting and algebraic operations stay measurable. □

**Corollary 1.1.6.** *Continuous Composition and Finite Lattice Operations*
*Let $(X, \mathcal{M})$ be a measurable space and let $n \in \mathbb{N}_{\geq 1}$.*
*For $f_1, \ldots, f_n : X \to \mathbb{R}$, set $\bigvee_{k=1}^n f_k := \max\{f_1, \ldots, f_n\}$ and $\bigwedge_{k=1}^n f_k := \min\{f_1, \ldots, f_n\}$.*

*(i) If $g : X \to \mathbb{R}$ is measurable and $\varphi : \mathbb{R} \to \mathbb{R}$ is continuous, then $\varphi \circ g$ is measurable.*
*(ii) If $f_1, \ldots, f_n : X \to \mathbb{R}$ are measurable, then $\bigvee_{k=1}^n f_k$ and $\bigwedge_{k=1}^n f_k$ are measurable.*

*Proof.*

(i) For $a \in \mathbb{R}$, $\{\varphi \circ g > a\} = g^{-1}(\varphi^{-1}((a, +\infty))) \in \mathcal{M}$, since $\varphi^{-1}((a, +\infty)) \in \mathcal{B}(\mathbb{R})$ and $g$ is measurable.
(ii) Set $h_k = f_k$ for $1 \leq k \leq n$ and $h_k = f_n$ for $k > n$. Then $(h_k)$ is measurable, $\bigvee_{k=1}^n f_k = \sup_{k \geq 1} h_k$, and $\bigwedge_{k=1}^n f_k = \inf_{k \geq 1} h_k$, hence both are measurable by Theorem 1.1.5(iii).

*Structural Commentary*: Pullbacks preserve Borel thresholds; finite lattices are finite suprema and infima.
*Key Insight*: Continuity and finite lattice operations preserve measurable threshold structure. □

**Example 1.1.7.** *Measurable Composition Can Fail*
Choose $A \in \mathcal{L}^1$ with $A \subseteq [0, 1]$ and an increasing homeomorphism $\psi : [0, 1] \to [0, 2]$ such that $\psi(A) \notin \mathcal{L}^1$. Define

$$\widetilde{\psi}(x) := \begin{cases} x, & x \leq 0, \\ \psi(x), & 0 \leq x \leq 1, \\ x + 1, & x \geq 1. \end{cases}$$

Then $\widetilde{\psi} : \mathbb{R} \to \mathbb{R}$ is a homeomorphism. Put $h := \mathbf{1}_A \circ \widetilde{\psi}^{-1}$.
The maps $\mathbf{1}_A$ and $\widetilde{\psi}^{-1}$ are measurable, but $h^{-1}((\frac{1}{2}, \frac{3}{2})) = \widetilde{\psi}(A) = \psi(A) \notin \mathcal{L}^1$.
Hence $h$ is not Lebesgue measurable, so Lebesgue measurable functions are not composition-closed.

**Definition 1.1.8.** *Positive and Negative Parts*
For an extended-real-valued function $f$, define $f^+(x) := \sup\{f(x), 0\}$; $f^-(x) := \sup\{-f(x), 0\}$.
When $f$ is finite-valued, $f = f^+ - f^-$ and $|f| = f^+ + f^-$; equivalently $f^+ = \frac{|f| + f}{2}$; $f^- = \frac{|f| - f}{2}$.

**Corollary 1.1.9.** *Measurability of Positive and Negative Parts*
*If $f : X \to [-\infty, +\infty]$ is measurable, then $f^+, f^-, |f|$ are measurable.*
*If $f : X \to \mathbb{R}$, then $f$ measurable $\iff$ $f^+, f^-$ measurable.*

*Proof.*

($\Rightarrow$) Since $\{-f > a\} = \{f < -a\}$, $-f$ is measurable.
Thus $f^+ = f \vee 0$, $f^- = (-f) \vee 0$, and $|f| = f \vee (-f)$ are measurable by finite sup-closure.
($\Leftarrow$) If $f : X \to \mathbb{R}$ and $f^+, f^-$ are measurable, then $f = f^+ - f^-$ is measurable.

*Structural Commentary*: Sign decomposition is compatible with threshold-set measurability.
*Key Insight*: Reduce signs to suprema and finite algebra. □

**Lemma 1.1.10.** *Extended-Valued Measurability via the Finite Part*

*Let $f : X \to \overline{\mathbb{R}}$ and set $A := f^{-1}(\{+\infty\})$, and $B := f^{-1}(\{-\infty\})$. Define $f_0(x) := \begin{cases} f(x), & x \notin A \cup B, \\ 0, & x \in A \cup B. \end{cases}$.*

*Then $f$ is measurable iff $A, B \in \mathcal{M}$ and $f_0 : X \to \mathbb{R}$ is measurable.*

*Proof.*

($\Rightarrow$) If $f$ is measurable, then $A = \bigcap_{n=1}^{\infty} \{f > n\}$; and $B = X \setminus \bigcup_{n=1}^{\infty} \{f > -n\}$, so $A, B \in \mathcal{M}$.

Also, $\{f_0 > \alpha\} = \begin{cases} \{f > \alpha\} \setminus A, & \alpha \geq 0, \\ \{f > \alpha\} \cup B, & \alpha < 0, \end{cases}$, so $f_0$ is measurable.

($\Leftarrow$) If $A, B \in \mathcal{M}$, $f_0$ is measurable, then $\{f > \alpha\} = \begin{cases} \{f_0 > \alpha\} \cup A, & \alpha \geq 0, \\ \{f_0 > \alpha\} \setminus B, & \alpha < 0, \end{cases}$, so $\{f > \alpha\} \in \mathcal{M}\ \forall \alpha \in \mathbb{R}$.

*Structural Commentary*: The infinite-level sets are measurable exactly when the finite core is measurable.
*Key Insight*: Assign harmless finite values at $\pm\infty$, then put the infinite fibers back by set algebra. □

**Corollary 1.1.11.** *Extended-Valued Algebra Conventions*

*Let $f, g : X \to \overline{\mathbb{R}}$ be measurable.*

*(i) For $c \in \mathbb{R}$, $cf$, $|f|$, $f^+$, and $f^-$ are measurable, with $0(\pm\infty) = 0$.*
*(ii) Let $E_1 := \{f = -\infty,\ g = +\infty\}$, $E_2 := \{f = +\infty,\ g = -\infty\}$, and $F := E_1 \cup E_2$. If $h = f + g$ on $X \setminus F$ and $h = 0$ on $F$, then $h$ is measurable.*
*(iii) With $0(\pm\infty) = 0$, $fg$ is measurable.*

*Proof.*

(i) By Lemma 1.1.10, $A_{\pm}(f) := \{f = \pm\infty\} \in \mathcal{M}$ and $f_0$ is measurable.
For $c \in \mathbb{R}$, the infinite fibers of $cf$ are among $A_{\pm}(f)$ and $(cf)_0 = cf_0$, so $cf$ is measurable.
The functions $|f|, f^+, f^-$ are measurable by Corollary 1.1.9.

(ii) $E_1, E_2 \in \mathcal{M}$. Set $f_n := \max\{-n, \min(f, n)\}$ and $g_n := \max\{-n, \min(g, n)\}$.
Then $f_n, g_n$ are real-valued measurable and $(f_n + g_n)\mathbf{1}_{X \setminus F} \to h$ pointwise; hence $h$ is measurable.

(iii) With $f_n, g_n$ as in (ii), $f_n g_n$ measurable, $f_n g_n \to fg$ pointwise under $0(\pm\infty) = 0$; hence $fg$ measurable.

*Structural Commentary*: Extended algebra is finite algebra plus measurable exceptional fibers.
*Key Insight*: Truncate first, use real-valued closure, then pass to pointwise limits. □

**Definition 1.1.12.** *Simple Functions*

Let $(X, \mathcal{M})$ be a measurable space.

(i) *Indicator*: for $E \subseteq X$, $\mathbf{1}_E(x) := 1$ on $E$ and $\mathbf{1}_E(x) := 0$ on $X \setminus E$.
(ii) *Simple*: a real-valued $s$ is simple iff $s(X)$ is finite.
(iii) *Standard representation*: if $s(X) = \{c_1, \ldots, c_n\}$ and $E_i := \{s = c_i\} \neq \emptyset$, then $X = \bigsqcup_{i=1}^{n} E_i$; $s = \sum_{i=1}^{n} c_i \mathbf{1}_{E_i}$.
The pairs $(c_i, E_i)$ are unique up to reordering.
(iv) *Measurability*: $s$ measurable iff $E_i \in \mathcal{M}\ \forall i$; step functions are special case $X \subseteq \mathbb{R}$ with interval level sets.




Figure 201: A simple function takes value $c_i$ on each measurable piece $E_i$.

**Proposition 1.1.13.** *Simple Functions Form a Lattice Algebra*

*Let $s, t : X \to \mathbb{R}$ be simple and $a \in \mathbb{R}$. Then $as$, $s + t$, $st$, $s \vee t$, and $s \wedge t$ are simple.*
*If $s, t$ are measurable, so are these five functions.*

*Proof.* Let $s(X) = \{c_i\}_{i=1}^m$ and $t(X) = \{d_j\}_{j=1}^n$. Each displayed range is contained in the corresponding finite set $\{ac_i\}$, $\{c_i + d_j\}$, $\{c_i d_j\}$, $\{c_i \vee d_j\}$, or $\{c_i \wedge d_j\}$, hence is finite.
If $s, t$ are measurable, the atoms $P_{ij} := \{s = c_i\} \cap \{t = d_j\}$ are measurable and form a finite partition of $X$. Each displayed function is constant on every $P_{ij}$, so its level sets are finite unions of $P_{ij}$.

*Structural Commentary*: Algebra and lattice operations reduce to finite tables.
*Key Insight*: Refine two simple functions by the common partition of their level sets. □

**Lemma 1.1.14.** *Bounded Simple Approximation Brackets*
*Let $(X, \mathcal{M})$ be a measurable space and let $f : X \to \mathbb{R}$ be bounded and $\mathcal{M}$-measurable. Then $\forall \varepsilon > 0$ $\exists$ simple $\mathcal{M}$-measurable functions $\varphi_\varepsilon, \psi_\varepsilon : X \to \mathbb{R}$ with $\varphi_\varepsilon \le f \le \psi_\varepsilon$ and $0 \le \psi_\varepsilon - \varphi_\varepsilon < \varepsilon$ on $X$.*

*Proof.* Choose $c = y_0 < \cdots < y_N = d$ with $f(X) \subseteq (c, d)$ and $\max_k (y_k - y_{k-1}) < \varepsilon$.
Set $E_k := \{y_{k-1} \le f < y_k\} = \{f \ge y_{k-1}\} \cap \{f < y_k\} \in \mathcal{M}$; then $(E_k)_{k=1}^N$ partitions $X$.
Define $\varphi_\varepsilon := \sum_{k=1}^N y_{k-1} \mathbf{1}_{E_k}$ and $\psi_\varepsilon := \sum_{k=1}^N y_k \mathbf{1}_{E_k}$; these are $\mathcal{M}$-measurable simple functions.
For $x \in E_k$, $\varphi_\varepsilon(x) = y_{k-1} \le f(x) < y_k = \psi_\varepsilon(x)$ and $\psi_\varepsilon(x) - \varphi_\varepsilon(x) = y_k - y_{k-1} < \varepsilon$.
*Structural Commentary*: Pull back a finite range partition.
*Key Insight*: Mesh size controls the bracket gap. □

**Theorem 1.1.15.** *Approximation by Simple Functions*
*Let $(X, \mathcal{M})$ be a measurable space.*

*(i)* $\forall f : X \to \mathbb{R}$ $\exists$ *simple functions $(s_n)$ with $s_n \to f$ pointwise.*
*(ii) For $f : X \to \overline{\mathbb{R}}$, $f$ is $\mathcal{M}$-measurable iff $f$ is the pointwise limit of $\mathcal{M}$-measurable simple functions.*
*(iii) If $f : X \to \overline{\mathbb{R}}$ is $\mathcal{M}$-measurable, the approximants in (ii) may be chosen with $|s_n| \le |f|$. If $f \ge 0$, they may be chosen with $0 \le s_n \uparrow f$.*

*Proof.* For $u : X \to [0, +\infty]$, set $Q_n u := 2^{-n} \lfloor 2^n (u \wedge n) \rfloor$.
Then $Q_n u(X) \subseteq 2^{-n}\{0, \dots, n2^n\}$, so $Q_n u$ is simple, and binary truncation gives $0 \le Q_n u \uparrow u$.
If $u$ is measurable, the level sets of $Q_n u$ are threshold sets of $u \wedge n$, hence $Q_n u$ is measurable.

(i) For $f : X \to \mathbb{R}$, write $f = f^+ - f^-$ and set $s_n := Q_n f^+ - Q_n f^-$. Then $s_n$ is simple and $s_n \to f$ pointwise.
*(ii)* ($\Rightarrow$) If $f : X \to \overline{\mathbb{R}}$ is measurable, then $f^+, f^-$ are measurable and $s_n := Q_n f^+ - Q_n f^-$ is measurable simple. Since at each $x$ at most one of $f^+(x), f^-(x)$ is infinite, $s_n(x) \to f(x)$ in $\overline{\mathbb{R}}$.
*(ii)* ($\Leftarrow$) If measurable simple $s_n \to f$ pointwise, then $f = \liminf_n s_n$ is measurable.
*(iii)* In *(ii)* $\Rightarrow$, $|s_n| \le Q_n f^+ + Q_n f^- \le |f|$. If $f \ge 0$, take $s_n := Q_n f$.
*Structural Commentary*: Simple functions quantize the range of $f$.
*Key Insight*: Dyadic range partitions give pointwise approximation while preserving measurability. □

**Definition 1.1.16.** *Complex-Valued Measurable Functions*
A map $f : X \to \mathbb{C}$ is *measurable* iff $\operatorname{Re} f, \operatorname{Im} f : X \to \mathbb{R}$ are measurable.
Equivalently, $f = \operatorname{Re} f + i \operatorname{Im} f$ with measurable real components.

**Proposition 1.1.17.** *Complex Measurable Closure*
*If $f, g : X \to \mathbb{C}$ are measurable, $a \in \mathbb{C}$, and $h_n : X \to \mathbb{C}$ are measurable with $h_n \to h$ pointwise, then $f + g$, $fg$, $af$, and $h$ are measurable.*

*Proof.* Write $f = u + iv$, $g = r + is$, and $a = \alpha + i\beta$. Then $f + g = (u + r) + i(v + s)$, $af = (\alpha u - \beta v) + i(\alpha v + \beta u)$, and $fg = (ur - vs) + i(us + vr)$, so real closure gives measurability.
If $h_n \to h$, then $\operatorname{Re} h_n \to \operatorname{Re} h$ and $\operatorname{Im} h_n \to \operatorname{Im} h$, so pointwise-limit closure gives measurability.
*Structural Commentary*: Complex measurability is two real measurability statements.
*Key Insight*: Split into real and imaginary parts. □

**Definition 1.1.18.** *Almost Everywhere and In Measure*
Let $E$ be measurable.

(i) *A.E. equality*: $f \sim_{\text{a.e.}} g$ on $E$ iff $\mu(\{x \in E : f(x) \ne g(x)\}) = 0$.
(ii) *A.E. property*: a property holds a.e. on $E$ iff its failure set in $E$ is null.
(iii) *Null invariance*: $\sim_{\text{a.e.}}$ is an equivalence relation.
(iv) *A.E. convergence*: $f_n \to f$ a.e. on $E$ iff $\exists N \subseteq E$ null such that $\forall x \in E \setminus N$, $f_n(x) \to f(x)$.
(v) *In measure*: $f_n \to f$ in measure on $E$ iff $\forall \varepsilon > 0$, $\mu(\{x \in E : |f_n(x) - f(x)| > \varepsilon\}) \to 0$.

**Remark 1.1.19.** *Completion and Almost-Everywhere Limits*
If $E \subseteq N \in \mathcal{M}$, $\mu(N) = 0$, and $E \notin \mathcal{M}$, then $f_n \equiv 0$ is measurable and $f_n \to \mathbf{1}_E$ a.e.,
since $\{x : f_n(x) \not\to \mathbf{1}_E(x)\} \subseteq N$. Yet $\mathbf{1}_E$ is not $\mathcal{M}$-measurable.
In the completion, every $E \subseteq N$ is measurable, so null modifications preserve measurability.
*Key Insight*: Almost-everywhere statements are most stable on complete measure spaces.

**Corollary 1.1.20.** *Pointwise and Almost-Everywhere Limits*
*Let $(f_n)$ be $\overline{\mathbb{R}}$-valued measurable functions on $(X, \mathcal{M})$, and let $f : X \to \overline{\mathbb{R}}$.*

*(i) If $f_n \to f$ pointwise on $X$, then $f$ is measurable.*
*(ii) If $(X, \mathcal{M}, \mu)$ is complete and $f_n \to f$ a.e. on $X$, then $f$ is measurable.*

*Proof.*

(i) Pointwise, $f = \liminf_n f_n = \limsup_n f_n$; closure under lim inf and lim sup gives measurability.

(ii) Choose $N \in \mathcal{M}$ with $\mu(N) = 0$ and $f_n \to f$ on $X \setminus N$.
Define $g = f$ and $g_n = f_n$ on $X \setminus N$, and $g = g_n = 0$ on $N$.
Then each $g_n$ is measurable and $g_n \to g$ pointwise, so $g$ is measurable by (i).
Since $\forall a \in \mathbb{R},\ \{f > a\} \triangle \{g > a\} \subseteq N$, completeness gives $\{f > a\} \in \mathcal{M}$; hence $f$ is measurable.

*Structural Commentary*: Limits reduce to upper and lower limits; a.e. limits require completion.
*Key Insight*: Completion is exactly what lets null modifications preserve measurability. □

**Proposition 1.1.21.** *Completed Measurable Functions Have Original Representatives*
*Let $(X, \mathcal{M}, \mu)$ be a measure space with completion $(X, \overline{\mathcal{M}}_\mu, \overline{\mu})$.*
*If $f : X \to \overline{\mathbb{R}}$ is $\overline{\mathcal{M}}_\mu$-measurable, then $\exists g : X \to \overline{\mathbb{R}}$ that is $\mathcal{M}$-measurable and satisfies $f = g$ $\mu$-a.e.*

*Proof.* For $q \in \mathbb{Q}$, let $A_q := \{f > q\} \in \overline{\mathcal{M}}_\mu$, let $A_q = E_q \cup Z_q$, with $E_q \in \mathcal{M}$ and $Z_q \subseteq N_q \in \mathcal{M}$, $\mu(N_q) = 0$.
Put $N := \bigcup_{q \in \mathbb{Q}} N_q$, so $N \in \mathcal{M}$ and $\mu(N) = 0$, and define $g(x) := \begin{cases} f(x), & x \notin N, \\ 0, & x \in N. \end{cases}$.
For $q \in \mathbb{Q}$, $\{g > q\} = (E_q \setminus N) \cup \begin{cases} N, & q < 0, \\ \emptyset, & q \geq 0, \end{cases}$ belongs to $\mathcal{M}$. Hence, for $a \in \mathbb{R}$, $\{g > a\} = \bigcup_{\substack{q \in \mathbb{Q} \\ q > a}} \{g > q\} \in \mathcal{M}$.
Thus $g$ is $\mathcal{M}$-measurable and $f = g$ on $X \setminus N$.
*Structural Commentary*: Completion only changes measurable functions on null sets.
*Key Insight*: Replace completed rational levels off one null set. □

**Method 1.1.22.** *Measure Proof Patterns*

(i) *Disjointize*. If $A_n \uparrow$, set $B_1 := A_1$, $B_n := A_n \setminus A_{n-1}$; then $A_n = \bigsqcup_{k \leq n} B_k$ and $\bigcup_n A_n = \bigsqcup_n B_n$.
(ii) *Approximate*. Prove on $\mathcal{E}$; use $d_\mu(C, D) := \mu^*(C \triangle D)$ and $\triangle$ estimates to pass limits through $\cup, \cap, \setminus$.
(iii) *Threshold*. For real $f$, show $\forall a \in \mathbb{R},\ \{f > a\} \in \mathcal{M}$; for $f : X \to \mathbb{R}^m$, use coordinate and open-set tests.
(iv) *Generators*. For $f : X \to (Y, \sigma(\mathcal{A}))$, check $\forall A \in \mathcal{A},\ f^{-1}(A) \in \mathcal{M}$.
(v) *Truncate*. For $f : X \to \overline{\mathbb{R}}$, prove results for $f_n := (-n) \vee (f \wedge n)$, then let $n \to \infty$.
(vi) *Topological bridge*. Use continuous open preimages.
Reduce Borel tests to countable interval or rectangle preimages.

*Key Insight*: Measure theory alternates between countable set algebra and approximation by elementary sets.

## 1.2 Lebesgue Integration

Lebesgue integration is calibrated on indicators, extended linearly to nonnegative simple functions, then extended by monotone approximation and positive–negative decomposition.

**Perspective 1.2.1.** *Simple Minorants Are the Right Primitive*
For Lebesgue measure on $\mathbb{R}$, bounded measurable functions supported on finite-measure sets form a useful intermediate class; arbitrary measure spaces need not admit that reduction.
Let $X \neq \emptyset$, $\mathcal{M} = \{\emptyset, X\}$, and $\mu(X) = +\infty$. If $g$ is measurable and vanishes outside some $F \in \mathcal{M}$ with $\mu(F) < +\infty$, then $F = \emptyset$ and $g = 0$; hence finite-measure-support minorants of $\mathbf{1}_X$ have supremal integral 0, while the required normalization is $\int_X \mathbf{1}_X \, d\mu = \mu(X) = +\infty$. Thus the defining minorants must include $\mathbf{1}_A$ $\forall A \in \mathcal{M}$, and more generally all nonnegative simple $s \leq f$.
*Key Insight*: The primitive calibration is $\int_X \mathbf{1}_A \, d\mu = \mu(A)$, with $\mu(A) = +\infty$ allowed.

**Definition 1.2.2.** *Lebesgue Integral*
Let $(X, \mathcal{M}, \mu)$ be a measure space and $E \in \mathcal{M}$; let $\mathcal{S}^+$ be nonnegative measurable simple functions on $X$.
Write $s \in \mathcal{S}^+$ as $s = \sum_{i=1}^n c_i \mathbf{1}_{A_i}$ with $c_i \geq 0$ and pairwise disjoint $A_i \in \mathcal{M}$.
For measurable $f \geq 0$, put $\mathcal{S}^+(f) := \{s \in \mathcal{S}^+ : s \leq f\}$; for measurable $f : X \to \overline{\mathbb{R}}$, put $f^\pm := (\pm f) \vee 0$.

(i) *Simple*: $\mathcal{I}_E(s) := \sum_{i=1}^n c_i \mu(E \cap A_i)$, with $0 \cdot (+\infty) := 0$.

(ii) *Nonnegative*: $\int_E f\,d\mu := \sup_{s\in\mathcal{S}^+(f)} \mathcal{I}_E(s)$; $f$ integrable on $E \iff \int_E f\,d\mu < +\infty$.
(iii) *Signed*: if $\min\{\int_E f^+\,d\mu, \int_E f^-\,d\mu\} < +\infty$, set $\int_E f\,d\mu := \int_E f^+\,d\mu - \int_E f^-\,d\mu$.
(iv) $L^1$: $f \in \mathscr{L}^1(\mu)$ on $E$ iff $f$ is measurable and $\int_E |f|\,d\mu < +\infty$.

When $E = X$, write $\int f\,d\mu := \int_X f\,d\mu$ and drop "on $E$".

Simple functions are finite linear combinations of indicators of measurable sets: coefficients carry size, while the level sets carry measurability.

**Lemma 1.2.3.** *Refining a Simple Integral*
*Let $s \geq 0$ be simple measurable and suppose $s = \sum_{k=1}^{m} b_k \mathbf{1}_{F_k}$ with $b_k \geq 0$ and $F_k \in \mathcal{M}$.*
*Then $\forall E \in \mathcal{M}$, $\mathcal{I}_E(s) = \sum_{k=1}^{m} b_k \mu(E \cap F_k)$.*

*Proof.* For $\eta \in \{0,1\}^m$, set $F_k^1 := F_k$, $F_k^0 := F_k^c$, discard empty atoms $G_\eta := \bigcap_{k=1}^{m} F_k^{\eta_k}$. On $G_\eta$, $s = c_\eta := \sum_{\eta_k=1} b_k$. Group equal $c_\eta$ values to recover the standard representation; finite additivity gives
$\mathcal{I}_E(s) = \sum_\eta c_\eta \mu(E \cap G_\eta) = \sum_{k=1}^{m} b_k \sum_{\eta:\eta_k=1} \mu(E \cap G_\eta) = \sum_{k=1}^{m} b_k \mu(E \cap F_k)$.
*Structural Commentary*: Refine nonstandard sums to one finite partition.
*Key Insight*: Refine first, then regroup by equal function values. □

**Proposition 1.2.4.** *Simple Integral Algebra*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, let $E \in \mathcal{M}$, let $c \geq 0$, and let $s, t \in \mathcal{S}^+$. For fixed $s \in \mathcal{S}^+$, define $\nu_s : \mathcal{M} \to [0, +\infty]$ by $\nu_s(A) := \mathcal{I}_A(s)$. Then:*
*(i) $\mathcal{I}_E(cs) = c\mathcal{I}_E(s)$.*
*(ii) $\mathcal{I}_E(s+t) = \mathcal{I}_E(s) + \mathcal{I}_E(t)$.*
*(iii) $\nu_s$ is a measure on $\mathcal{M}$.*

*Proof.* Write $s = \sum_i a_i \mathbf{1}_{A_i}$ and $t = \sum_j b_j \mathbf{1}_{B_j}$ on finite measurable partitions of $X$, with $a_i, b_j \geq 0$.

(i) $\mathcal{I}_E(cs) = \sum_i (ca_i)\mu(E \cap A_i) = c\sum_i a_i \mu(E \cap A_i) = c\mathcal{I}_E(s)$, with $0 \cdot (+\infty) = 0$.

(ii) On the common refinement $A_i \cap B_j$, $s + t = \sum_{i,j} (a_i + b_j)\mathbf{1}_{A_i \cap B_j}$.

$\mathcal{I}_E(s+t) = \sum_{i,j} (a_i + b_j)\mu(E \cap A_i \cap B_j) = \sum_i a_i \mu(E \cap A_i) + \sum_j b_j \mu(E \cap B_j)$ by Lemma 1.2.3.

(iii) For $A \in \mathcal{M}$, $\nu_s(A) = \sum_i a_i \mu(A \cap A_i)$.

Each $A \mapsto \mu(A \cap A_i)$ is a measure, and finite nonnegative sums of measures are measures.

*Structural Commentary*: Simple integration is weighted measure summation on finite partitions.
*Key Insight*: Refine to common atoms, then use finite additivity. □

**Proposition 1.2.5.** *Monotonicity for Nonnegative Integrals*
*Let $f, g \geq 0$ be measurable and let $E, F \in \mathcal{M}$.*
*(i) If $f \leq g$ on $E$, then $\int_E f\,d\mu \leq \int_E g\,d\mu$.*
*(ii) If $E \subseteq F$, then $\int_E f\,d\mu \leq \int_F f\,d\mu$.*

*Proof.*

(i) If $s \in \mathcal{S}^+(f)$, then $s\mathbf{1}_E \in \mathcal{S}^+(g)$, $\mathcal{I}_E(s) = \mathcal{I}_E(s\mathbf{1}_E) \leq \int_E g\,d\mu$. Taking $\sup_{s\in\mathcal{S}^+(f)}$ gives $\int_E f\,d\mu \leq \int_E g\,d\mu$.

(ii) If $E \subseteq F$ and $s \in \mathcal{S}^+(f)$, then $\mathcal{I}_E(s) \leq \mathcal{I}_F(s)$. Taking $\sup_{s\in\mathcal{S}^+(f)}$ gives $\int_E f\,d\mu \leq \int_F f\,d\mu$.

*Structural Commentary*: The nonnegative integral is a supremum over simple functions below the integrand.
*Key Insight*: More room below the integrand, or over a larger set, can only increase the supremum. □

**Proposition 1.2.6.** *Chebyshev Inequality for Nonnegative Integrals*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, let $E \in \mathcal{M}$, and let $f : X \to [0, +\infty]$ be measurable.*
*For $\lambda > 0$, set $E_\lambda := E \cap \{f \geq \lambda\}$. Then $E_\lambda \in \mathcal{M}$ and $\mu(E_\lambda) \leq \lambda^{-1} \int_E f\,d\mu$.*

*Proof.* Measurability gives $E_\lambda \in \mathcal{M}$.
Since $\lambda \mathbf{1}_{E_\lambda} \leq f$ on $E$, monotonicity gives $\lambda\mu(E_\lambda) = \int_E \lambda \mathbf{1}_{E_\lambda}\,d\mu \leq \int_E f\,d\mu$; divide by $\lambda > 0$.
*Structural Commentary*: Level-set minorants convert mass to measure.
*Key Insight*: Use $\lambda \mathbf{1}_{E_\lambda} \leq f$ on $E$. □

**Method 1.2.7.** *Splitting a Simple Minorant Under a Sum*
Let $E \in \mathcal{M}$, $f, g \geq 0$ measurable, and $\omega \in \mathcal{S}^+(f+g)$ simple.

(i) *Grid*: for $n \geq 1$, use $D_n := \{0, \frac{1}{n}, \dots, 1\}$.
(ii) *Split*: define $\varphi_n(x) := \max\{\theta\omega(x) : \theta \in D_n,\ \theta\omega(x) \leq f(x)\}$; $\psi_n(x) := \big(\big(1 - \frac{1}{n}\big)\,\omega(x) - \varphi_n(x)\big)^+$.
(iii) *Bounds*: $\varphi_n, \psi_n$ are simple, $\varphi_n \leq f$, $\psi_n \leq g$, and $(1 - \frac{1}{n})\omega \leq \varphi_n + \psi_n$.
(iv) *Use*: simple additivity gives $(1 - \frac{1}{n})\mathcal{I}_E(\omega) \leq \int_E f\,d\mu + \int_E g\,d\mu$; then let $n \to \infty$ and take $\sup_{\omega \in \mathcal{S}^+(f+g)}$.

*Key Insight*: Grid-split $\omega$ between $f$ and $g$, losing only $\frac{1}{n}$.

**Proposition 1.2.8.** *Homogeneity and Additivity for Nonnegative Integrals*
*Let $f, g \geq 0$ be measurable, let $c \geq 0$, and let $E \in \mathcal{M}$. All identities are in $[0, +\infty]$, with $0 \cdot (+\infty) = 0$.*

*(i) Homogeneity: $\int_E cf\,d\mu = c\int_E f\,d\mu$.*
*(ii) Additivity: $\int_E (f+g)\,d\mu = \int_E f\,d\mu + \int_E g\,d\mu$.*

*Proof.*

(i) If $c = 0$, both sides are 0; if $c > 0$, $s \mapsto cs$ bijects $\mathcal{S}^+(f)$ with $\mathcal{S}^+(cf)$ and satisfies $\mathcal{I}_E(cs) = c\mathcal{I}_E(s)$. Hence taking suprema gives the claim.
(ii) If $s \in \mathcal{S}^+(f)$ and $t \in \mathcal{S}^+(g)$, then $s + t \in \mathcal{S}^+(f+g)$ and $\mathcal{I}_E(s) + \mathcal{I}_E(t) = \mathcal{I}_E(s+t) \leq \int_E (f+g)\,d\mu$, so taking $\sup_{s,t}$ gives $\int_E f\,d\mu + \int_E g\,d\mu \leq \int_E (f+g)\,d\mu$.
Conversely, for $\omega \in \mathcal{S}^+(f+g)$, choose simple $\varphi_n \leq f$ and $\psi_n \leq g$ with $(1 - \frac{1}{n})\omega \leq \varphi_n + \psi_n$. Then $\big(1 - \frac{1}{n}\big)\,\mathcal{I}_E(\omega) \leq \mathcal{I}_E(\varphi_n) + \mathcal{I}_E(\psi_n) \leq \int_E f\,d\mu + \int_E g\,d\mu$. Let $n \to \infty$, then take $\sup_{\omega \in \mathcal{S}^+(f+g)}$.

*Structural Commentary*: The nonnegative integral is linear before signed integrability is introduced.
*Key Insight*: Scale minorants; split sum-minorants on a finite grid. □

**Proposition 1.2.9.** *Signed Decompositions Give the Same Integral*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, let $E \in \mathcal{M}$, and let $f, f_1, f_2 : X \to \overline{\mathbb{R}}$ be measurable. Assume $f_i \geq 0$, $\int_E f_i\,d\mu < +\infty$ for $i = 1, 2$, and $f = f_1 - f_2$ on $E$. Then:*

*(i) $f \in \mathscr{L}^1(\mu)$ on $E$.*
*(ii) $\int_E f\,d\mu = \int_E f_1\,d\mu - \int_E f_2\,d\mu$.*

*Proof.*

(i) Since $-f_2 \leq f \leq f_1$ on $E$, $f^+ \leq f_1$ and $f^- \leq f_2$ on $E$; monotonicity gives $\int_E f^{\pm}\,d\mu < +\infty$.
(ii) Since $f^+ - f^- = f_1 - f_2$ on $E$, $f^+ + f_2 = f_1 + f^-$ on $E$.
Nonnegative additivity gives $\int_E f^+\,d\mu + \int_E f_2\,d\mu = \int_E f_1\,d\mu + \int_E f^-\,d\mu$; subtract the finite terms.

*Structural Commentary*: Finite positive decompositions determine one signed integral.
*Key Insight*: Move all nonnegative terms to one equality, integrate, then cancel finite terms. □

**Perspective 1.2.10.** *Step Functions versus Simple Functions*
Riemann: $\sup \sum_i m_i \ell(I_i)$ over interval partitions. Lebesgue: $\sup\{\mathcal{I}(s) : 0 \leq s \leq f,\ s = \sum_i c_i \mathbf{1}_{E_i},\ E_i \in \mathcal{M}\}$.

*Structural Commentary*: Riemann partitions intervals; Lebesgue partitions level sets.
*Key Insight*: Same finite constants, measurable pieces.

**Perspective 1.2.11.** *Undergraph Picture for Nonnegative Functions*
For measurable $f : \mathbb{R} \to [0, +\infty]$, set $\mathcal{U}_f := \{(x, y) \in \mathbb{R} \times [0, +\infty) : 0 \leq y < f(x)\}$.
Geometrically, $\int f\,dm = m_2(\mathcal{U}_f)$; analytically, the integral is built as $\sup \mathcal{I}(s)$ over simple $0 \leq s \leq f$.




Figure 202: The integral of $f \geq 0$ is modeled by the measure of its undergraph.

**Proposition 1.2.12.** *Basic Integral Properties*
*Let $E \in \mathcal{M}$.*

*(i) If $f$ is measurable and bounded on $E$, and $\mu(E) < +\infty$, then $f \in \mathscr{L}^1(\mu)$ on $E$.*
*(ii) If $a \le f(x) \le b$ on $E$ and $\mu(E) < +\infty$, then $a\mu(E) \le \int_E f\,d\mu \le b\mu(E)$.*
*(iii) If $f, g \in \mathscr{L}^1(\mu)$ on $E$ and $f \le g$ on $E$, then $\int_E f\,d\mu \le \int_E g\,d\mu$.*
*(iv) If $f \in \mathscr{L}^1(\mu)$ on $E$, then $cf \in \mathscr{L}^1(\mu)$ on $E$ for finite $c$, and $\int_E cf\,d\mu = c\int_E f\,d\mu$.*
*(v) If $\mu(E) = 0$ and $f$ is measurable, then $\int_E f\,d\mu = 0$.*
*(vi) If $f \in \mathscr{L}^1(\mu)$ on $E$, $A \in \mathcal{M}$, and $A \subseteq E$, then $f \in \mathscr{L}^1(\mu)$ on $A$.*

*Proof.*

(i) If $|f| \le M$ on $E$, then $0 \le f^{\pm} \le M\mathbf{1}_E$ on $E$, so $\int_E f^{\pm}\,d\mu \le M\mu(E) < +\infty$.
(ii) Since $a\mathbf{1}_E \le f \le b\mathbf{1}_E$ on $E$, order for the positive integral gives $a\mu(E) \le \int_E f\,d\mu \le b\mu(E)$.
(iii) Since $g - f \ge 0$ on $E$, $\int_E (g-f)\,d\mu \ge 0$, hence $\int_E f\,d\mu \le \int_E g\,d\mu$.
(iv) For $c \ge 0$, scale the simple approximants. For $c < 0$, use $(cf)^+ = (-c)f^-$ and $(cf)^- = (-c)f^+$.
(v) If $\mu(E) = 0$, then $\forall s \ge 0$ simple, $\mathcal{I}_E(s) = 0$, so $\int_E f^{\pm}\,d\mu = 0$.
(vi) Since $A \subseteq E$, $\mathcal{I}_A(s) \le \mathcal{I}_E(s)$ for $s \ge 0$ simple; hence $\int_A f^{\pm}\,d\mu \le \int_E f^{\pm}\,d\mu < +\infty$.

*Structural Commentary*: Estimate $f^{\pm}$ by simpler functions.
*Key Insight*: Monotonicity makes bounds, null sets, and restrictions immediate. □

**Theorem 1.2.13.** *The Integral as a Measure or Charge*
*For $A \in \mathcal{M}$, define $\phi_f(A) := \int_A f\,d\mu$ when $f \ge 0$ is measurable, and define $\lambda_f(A) := \int_A f\,d\mu$ when $f \in \mathscr{L}^1(\mu)$.*

*(i) If $f \ge 0$ is measurable, then $\phi_f : \mathcal{M} \to [0, +\infty]$ is a measure.*
*(ii) If $f \in \mathscr{L}^1(\mu)$, then $\lambda_f : \mathcal{M} \to \mathbb{R}$ is finite-valued and countably additive.*

*Proof.*

(i) For simple $s \ge 0$, $A \mapsto \mathcal{I}_A(s)$ is a measure, so $\phi_f(\emptyset) = 0$.
If $A = \bigsqcup_n A_n$, $s \in \mathcal{S}^+(f)$, then $\mathcal{I}_A(s) = \sum_n \mathcal{I}_{A_n}(s) \le \sum_n \phi_f(A_n)$; taking $\sup_s$ gives $\phi_f(A) \le \sum_n \phi_f(A_n)$.
Conversely, $\forall N$, finite additivity gives $\sum_{n=1}^{N} \phi_f(A_n) = \phi_f(\bigcup_{n=1}^{N} A_n) \le \phi_f(A)$.
Let $N \to \infty$. Hence $\phi_f(A) = \sum_n \phi_f(A_n)$.
(ii) Since $f \in \mathscr{L}^1(\mu)$, $\phi_{f^+}(X), \phi_{f^-}(X) < +\infty$ and $\lambda_f = \phi_{f^+} - \phi_{f^-}$ is finite-valued.
If $A = \bigsqcup_n A_n$, then item (i) gives $\lambda_f(A) = \sum_n \phi_{f^+}(A_n) - \sum_n \phi_{f^-}(A_n) = \sum_n \lambda_f(A_n)$.

*Structural Commentary*: Setwise integration turns densities into measures.
*Key Insight*: Sup simple measures; subtract finite signs. □

**Corollary 1.2.14.** *Domain Additivity and Continuity for Integrable Functions*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, let $E \in \mathcal{M}$, let $f \in \mathscr{L}^1(\mu)$ on $E$, and set $\mathcal{M}_E := \{A \in \mathcal{M} : A \subseteq E\}$.*

*(i) If $(A_n) \subseteq \mathcal{M}_E$ are pairwise disjoint, then $\int_{\bigsqcup_n A_n} f\,d\mu = \sum_n \int_{A_n} f\,d\mu$ and $\sum_n \left|\int_{A_n} f\,d\mu\right| < +\infty$.*
*(ii) If $A_n, A \in \mathcal{M}_E$ and $A_n \uparrow A$, then $\int_A f\,d\mu = \lim_{n\to\infty} \int_{A_n} f\,d\mu$.*
*(iii) If $A_n, A \in \mathcal{M}_E$ and $A_n \downarrow A$, then $\int_A f\,d\mu = \lim_{n\to\infty} \int_{A_n} f\,d\mu$.*

*Proof.* Let $\mu_E := \mu|_{\mathcal{M}_E}$ and define $\lambda(A) := \int_A f\,d\mu$, $\nu(A) := \int_A |f|\,d\mu$ on $\mathcal{M}_E$.
By Theorem 1.2.13, $\lambda$ is a finite signed measure, $\nu(E) < +\infty$, and $|\lambda(A)| \le \nu(A)$.

(i) Additivity gives $\lambda(\bigsqcup_n A_n) = \sum_n \lambda(A_n)$, and $\sum_n |\lambda(A_n)| \le \sum_n \nu(A_n) = \nu(\bigsqcup_n A_n) \le \nu(E) < +\infty$.
(ii) Set $B_1 = A_1$ and $B_n = A_n \setminus A_{n-1}$ for $n \ge 2$.
Since $A = \bigsqcup_n B_n$ and $A_N = \bigsqcup_{n \le N} B_n$, (i) gives $\lambda(A_N) = \sum_{n \le N} \lambda(B_n) \to \sum_n \lambda(B_n) = \lambda(A)$.
(iii) Set $C_n := A_1 \setminus A_n$ and $C := A_1 \setminus A$. Then $C_n \uparrow C$, so (ii) gives $\lambda(C_n) \to \lambda(C)$.
Hence $\lambda(A_n) = \lambda(A_1) - \lambda(C_n) \to \lambda(A_1) - \lambda(C) = \lambda(A)$.

*Structural Commentary*: Domain integrals form a finite signed measure.
*Key Insight*: Monotone domain limits are signed-measure continuity. □

**Proposition 1.2.15.** *Finite-Measure Support*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, let $E \in \mathcal{M}$, and let $h : E \to \mathbb{R}$ be bounded and measurable. Suppose $h = 0$ on $E \setminus E_0$ for some $E_0 \in \mathcal{M}$ with $E_0 \subseteq E$ and $\mu(E_0) < +\infty$. Then:*

*(i) $h \in \mathscr{L}^1(\mu)$ on $E$.*

*(ii)* $\int_E h\,d\mu = \int_{E_0} h\,d\mu$.
*(iii)* *If* $E_1 \in \mathcal{M}$, $E_1 \subseteq E$, $\mu(E_1) < +\infty$, *and* $h = 0$ *on* $E \setminus E_1$, *then* $\int_{E_0} h\,d\mu = \int_{E_1} h\,d\mu$.

*Proof.*

(i) Choose $M < +\infty$ with $|h| \le M$ on $E$.
Since $h = 0$ on $E \setminus E_0$, $0 \le h^{\pm} \le M\mathbf{1}_{E_0}$ on $E$, so $\int_E h^{\pm}\,d\mu \le M\mu(E_0) < +\infty$.
(ii) Since $\int_{E\setminus E_0} h\,d\mu = 0$, $\int_E h\,d\mu = \int_{E_0} h\,d\mu$ by Corollary 1.2.14.
(iii) Apply item (ii) with $E_0$ and with $E_1$: $\int_{E_0} h\,d\mu = \int_E h\,d\mu = \int_{E_1} h\,d\mu$.

*Structural Commentary*: Finite support turns boundedness into integrability.
*Key Insight*: The integral only sees where $h$ can be nonzero. □

**Corollary 1.2.16.** *Compact Continuous Functions on Finite Measure Spaces*
*Let* $X$ *be a compact topological space, let* $\mathcal{M}$ *be a* $\sigma$*-algebra containing the topology of* $X$, *and let* $\mu(X) < +\infty$. *If* $f : X \to \mathbb{R}$ *is continuous, then:*

*(i)* $f$ *is measurable.*
*(ii)* $f \in \mathscr{L}^1(\mu)$.

*Proof.*

(i) $\forall O \subseteq \mathbb{R}$ open, $f^{-1}(O)$ is open in $X$ by continuity, hence $f^{-1}(O) \in \mathcal{M}$.
(ii) The continuous image $f(X)$ is compact in $\mathbb{R}$, hence bounded.
Since $\mu(X) < +\infty$, $f \in \mathscr{L}^1(\mu)$ by Proposition 1.2.12(i).

*Structural Commentary*: Compactness supplies boundedness, while the measure hypothesis supplies finite mass.
*Key Insight*: Continuous on compact plus finite measure implies bounded measurable, hence integrable. □

**Corollary 1.2.17.** *Null and Finiteness Consequences for* $f \ge 0$
*Let* $f : X \to [0, +\infty]$ *be measurable and* $\phi_f(A) := \int_A f\,d\mu$.

*(i)* $\mu(A) = 0 \Longrightarrow \phi_f(A) = 0$ *for* $A \in \mathcal{M}$.
*(ii)* $\int_X f\,d\mu = 0 \Longleftrightarrow f = 0$ *a.e.*
*(iii)* $\int_X f\,d\mu < +\infty \Longrightarrow \mu\{f = +\infty\} = 0$ *and* $\{f > 0\}$ *is* $\sigma$*-finite.*

*Proof.*

*(i)* If $\mu(A) = 0$ and $0 \le s \le f$ is simple, then $\mathcal{I}_A(s) = 0$; hence $\phi_f(A) = \sup_s \mathcal{I}_A(s) = 0$.

*(ii)* ($\Rightarrow$) Set $E_n := \{f > \frac{1}{n}\}$. By Proposition 1.2.6, $\mu(E_n) \le n\int_X f\,d\mu = 0$.
Thus $\forall n$, $\mu(E_n) = 0$, and $\{f > 0\} = \bigcup_n E_n$ is null.

*(ii)* ($\Leftarrow$) If $N := \{f > 0\}$ null, every simple $0 \le s \le f$ vanishes on $X \setminus N$, so $\mathcal{I}_X(s) = \mathcal{I}_N(s) = 0$; take $\sup_s$.

*(iii)* Let $F_n := \{f \ge n\}$. By Proposition 1.2.6, $\mu(F_n) \le n^{-1}\int_X f\,d\mu$, so $\mu\{f = +\infty\} \le \mu(F_n) \to 0$.
Also $\{f > 0\} = \bigcup_n \{f > \frac{1}{n}\}$ and $\mu\{f > \frac{1}{n}\} \le n\int_X f\,d\mu < +\infty$.

*Structural Commentary*: Finite nonnegative integrals control threshold sets.
*Key Insight*: Threshold sets convert integral bounds into measure bounds. □

**Theorem 1.2.18.** *Absolute Integrability and Domination*
*Let* $(X, \mathcal{M}, \mu)$ *be a measure space and let* $E \in \mathcal{M}$. *Then:*

*(i) If* $f \in \mathscr{L}^1(\mu)$ *on* $E$, *then* $|f| \in \mathscr{L}^1(\mu)$ *on* $E$ *and* $\left|\int_E f\,d\mu\right| \le \int_E |f|\,d\mu$.
*(ii) If* $f : X \to \overline{\mathbb{R}}$ *is measurable,* $g \in \mathscr{L}^1(\mu)$ *on* $E$, *and* $|f| \le g$ *on* $E$, *then* $f \in \mathscr{L}^1(\mu)$ *on* $E$.

*Proof.*

(i) Set $a := \int_E f^+\,d\mu$ and $b := \int_E f^-\,d\mu$. Since $f \in \mathscr{L}^1(\mu)$ on $E$, $a, b < +\infty$.
Hence $\int_E |f|\,d\mu = \int_E (f^+ + f^-)\,d\mu = a + b < +\infty$, and $\left|\int_E f\,d\mu\right| = |a - b| \le a + b = \int_E |f|\,d\mu$.
(ii) Since $|f| \le g$ on $E$, $g \ge 0$ on $E$ and $\int_E g\,d\mu < +\infty$.
Also $0 \le f^{\pm} \le |f| \le g$ on $E$, so monotonicity gives $\int_E f^{\pm}\,d\mu < +\infty$.

*Structural Commentary*: Both claims reduce to finite positive and negative parts.
*Key Insight*: Use $|f| = f^+ + f^-$ and $0 \le f^{\pm} \le g$. □

**Proposition 1.2.19.** *Linearity of the Lebesgue Integral*
*Let* $(X, \mathcal{M}, \mu)$ *be a measure space, let* $E \in \mathcal{M}$, *let* $f, g \in \mathscr{L}^1(\mu)$ *on* $E$, *and let* $a, b \in \mathbb{R}$.
*Then:*

*(i) For* $c \in \mathbb{R}$ *and* $u \in \mathscr{L}^1(\mu)$ *on* $E$, $cu \in \mathscr{L}^1(\mu)$ *on* $E$ *and* $\int_E cu\,d\mu = c\int_E u\,d\mu$.

*(ii)* $f+g \in \mathscr{L}^1(\mu)$ *on* $E$ *and* $\int_E (f+g)\,d\mu = \int_E f\,d\mu + \int_E g\,d\mu$.
*(iii)* $af+bg \in \mathscr{L}^1(\mu)$ *on* $E$ *and* $\int_E (af+bg)\,d\mu = a\int_E f\,d\mu + b\int_E g\,d\mu$.

*Proof.*

(i) Since $(cu)^\pm = c^+u^\pm + c^-u^\mp$, both $(cu)^+$ and $(cu)^-$ have finite integral. Nonnegative linearity gives $\int_E cu\,d\mu = (c^+a_u + c^-b_u) - (c^+b_u + c^-a_u) = c(a_u - b_u)$, where $a_u := \int_E u^+\,d\mu$ and $b_u := \int_E u^-\,d\mu$.
(ii) Since $|f+g| \le |f| + |g|$ and $|f|+|g| \in \mathscr{L}^1(\mu)$ on $E$, $f+g \in \mathscr{L}^1(\mu)$ on $E$ by Theorem 1.2.18(ii). Integrate $(f+g)^+ + f^- + g^- = (f+g)^- + f^+ + g^+$ over $E$ and cancel finite terms.
(iii) By (i), $af, bg \in \mathscr{L}^1(\mu)$ on $E$, $\int_E af\,d\mu = a\int_E f\,d\mu$, and $\int_E bg\,d\mu = b\int_E g\,d\mu$. Applying (ii) to $af, bg$ gives $af+bg \in \mathscr{L}^1(\mu)$ on $E$ and $\int_E (af+bg)\,d\mu = a\int_E f\,d\mu + b\int_E g\,d\mu$.

*Structural Commentary*: Signed linearity follows from finite sign pieces.
*Key Insight*: Integrate one nonnegative equality, then cancel finite terms. □

**Proposition 1.2.20.** *$L^1$ Level Sets, Null Changes, and Indefinite Integrals*
*Let* $f, g \in \mathscr{L}^1(\mu)$ *on* $X$, *and write* $\lambda_u(A) := \int_A u\,d\mu$.

*(i)* *For* $a > 0$, $\mu\{|f| \ge a\} \le a^{-1}\int_X |f|\,d\mu < +\infty$; *hence* $\{f \ne 0\}$ *is* $\sigma$*-finite.*
*(ii)* *If measurable* $h = 0$ *a.e., then* $h \in \mathscr{L}^1(\mu)$ *and* $\lambda_h \equiv 0$.
*(iii)* $\lambda_f \ge 0$ *on* $\mathcal{M} \Longrightarrow f \ge 0$ *a.e.;* $\lambda_f \equiv 0 \Longrightarrow f = 0$ *a.e.*
*(iv)* $\lambda_f = \lambda_g$ *on* $\mathcal{M} \Longleftrightarrow f = g$ *a.e.*

*Proof.*

(i) Since $a\mathbf{1}_{\{|f|\ge a\}} \le |f|$, integration gives $a\mu\{|f| \ge a\} \le \int_X |f|\,d\mu$. Also $\{f \ne 0\} = \bigcup_{k\ge 1} \{|f| \ge \frac{1}{k}\}$.
(ii) Let $N := \{h \ne 0\}$. For $A \in \mathcal{M}$, $h^\pm \mathbf{1}_A$ vanish off $A \cap N$, a null set. Hence $\int_A h^+\,d\mu = \int_A h^-\,d\mu = 0$, so $\lambda_h(A) = 0$.
(iii) For $k \ge 1$, set $N_k := \{f \le -\frac{1}{k}\}$. By (i), $\mu(N_k) < +\infty$, and $\lambda_f(N_k) \le -k^{-1}\mu(N_k)$. If $\lambda_f \ge 0$, then $\mu(N_k) = 0$, so $\{f < 0\} = \bigcup_k N_k$ is null. If $\lambda_f \equiv 0$, apply this to $f$ and $-f$.
(iv) If $\lambda_f = \lambda_g$, then $\lambda_{f-g} \equiv 0$, so $f - g = 0$ a.e. by (iii). Conversely, if $f = g$ a.e., apply (ii) to $h = f - g$.

*Structural Commentary*: An $L^1$ function is detected by its integrals over all measurable sets, up to null sets.
*Key Insight*: Test sign on the level sets where the function is bounded away from 0. □

**Proposition 1.2.21.** *Finite-Measure Indicators by Intervals and Continuous Functions*
*Let* $A \subseteq \mathbb{R}$ *be Lebesgue measurable with* $m(A) < +\infty$. *Then* $\forall \varepsilon > 0$*:*

*(i)* $\exists G = \bigcup_{j=1}^{N} I_j$, $I_j$ *open intervals, such that* $m(A \triangle G) < \varepsilon$.
*(ii)* $\exists f \in C(\mathbb{R})$ *such that* $0 \le f \le 1$ *and* $\int_{\mathbb{R}} |\mathbf{1}_A - f|\,dm < \varepsilon$.

*Proof.*

(i) Choose open $O \supseteq A$ with $m(O \setminus A) < \frac{\varepsilon}{2}$ and $m(O) < +\infty$.
Write $O = \bigsqcup_{j\ge 1} I_j$ with open intervals $I_j$, choose $N$ with $m(O \setminus G) < \frac{\varepsilon}{2}$ for $G := \bigcup_{j=1}^{N} I_j$.
Then $A \setminus G \subseteq O \setminus G$ and $G \setminus A \subseteq O \setminus A$, so $m(A \triangle G) < \varepsilon$.
(ii) Apply (i) with $\frac{\varepsilon}{2}$ to get finite open $G$. Choose compact $K \subseteq G$ with $m(G \setminus K) < \frac{\varepsilon}{2}$, and choose piecewise-linear $f$ with $0 \le f \le 1$, $f = 1$ on $K$, $f = 0$ on $\mathbb{R} \setminus G$.
Then $\int_{\mathbb{R}} |\mathbf{1}_G - f|\,dm \le m(G \setminus K) < \frac{\varepsilon}{2}$, hence $\int_{\mathbb{R}} |\mathbf{1}_A - f|\,dm \le m(A \triangle G) + \int_{\mathbb{R}} |\mathbf{1}_G - f|\,dm < \varepsilon$.

*Structural Commentary*: Finite-measure sets can be made geometrically simple up to small $L^1$ error.
*Key Insight*: Approximate by intervals, then smooth the indicator. □

**Example 1.2.22.** *Counting and Unit-Mass Integrals*

(i) *Counting*: for counting measure $\#$ on $(\mathbb{N}, 2^{\mathbb{N}})$ and $f : \mathbb{N} \to \mathbb{R}$,
$f \in \mathscr{L}^1(\#) \Longleftrightarrow \sum_{n=1}^{\infty} |f(n)| < +\infty$, and then $\int_{\mathbb{N}} f\,d\# = \sum_{n=1}^{\infty} f(n)$.
(ii) *Unit mass*: for $(X, \mathcal{M})$, $p \in X$, and $\delta_p(E) := \mathbf{1}_E(p)$, $\int_X f\,d\delta_p = f(p)$ for measurable $f : X \to [0, +\infty]$.
If $f : X \to \mathbb{R}$ is measurable, then $f \in \mathscr{L}^1(\delta_p)$ and the same formula holds.

For $\delta_p$, verify first on indicators, then extend to simple functions by finite linearity, to nonnegative functions by monotone simple approximation, and to real-valued functions by positive–negative splitting.

**Example 1.2.23.** *$L^1$ Is Not Closed Under Squaring*
On $X = \mathbb{N}$ with $\mathcal{M} = 2^{\mathbb{N}}$, define $\mu(E) := \sum_{n\in E} 4^{-n}$ and $f(n) := \frac{2^n}{\sqrt{n}}$.

Then $\int_X |f|\, d\mu = \sum_{n=1}^{\infty} \frac{1}{2^n\sqrt{n}} < +\infty$; $\int_X f^2\, d\mu = \sum_{n=1}^{\infty} \frac{1}{n} = +\infty$.
The first series is dominated by the geometric series $\sum 2^{-n}$, while the second diverges by Corollary 6.1.14.
Thus $f \in \mathscr{L}^1(\mu)$ need not imply $f^2 \in \mathscr{L}^1(\mu)$.

## 1.3 Convergence Theorems

This subsection collects the main limit-interchange tools: monotone convergence, Fatou, and dominated convergence.

**Theorem 1.3.1.** *Monotone Convergence*
*Let $E \in \mathcal{M}$, let $f_n, f$ be measurable on $E$, and assume $0 \le f_n \uparrow f$. Then $\int_E f_n\, d\mu \uparrow \int_E f\, d\mu$.*

*Proof.* By monotonicity, $\int_E f_n\, d\mu \uparrow L \le \int_E f\, d\mu$.
For $s \in \mathcal{S}^+(f)$ and $0 < c < 1$, set $E_n := \{x \in E : f_n(x) \ge cs(x)\}$.
Since $f_n \uparrow f$ and $s \le f$, $E_n \uparrow E$, and $\int_E f_n\, d\mu \ge \int_{E_n} f_n\, d\mu \ge c\int_{E_n} s\, d\mu$.
Continuity from below for $\phi_s(A) := \int_A s\, d\mu$ gives $L \ge c\int_E s\, d\mu$. Let $c \uparrow 1$, then take $\sup_{s\in\mathcal{S}^+(f)}$.
*Structural Commentary*: Nonnegative integration commutes with monotone limits.
*Key Insight*: Simple minorants of the limit are eventually captured below the sequence. □

**Theorem 1.3.2.** *Series Form of Monotone Convergence*
*Let $E \in \mathcal{M}$, let $f_n \ge 0$ be measurable, and set $f := \sum_{n=1}^{\infty} f_n$ on $E$. Then $\int_E f\, d\mu = \sum_{n=1}^{\infty} \int_E f_n\, d\mu$.*

*Proof.* Set $S_N := \sum_{n=1}^{N} f_n$. Then $0 \le S_N \uparrow f$.

By Theorem 1.3.1, $\int_E f\, d\mu = \lim_{N\to\infty} \int_E S_N\, d\mu = \lim_{N\to\infty} \sum_{n=1}^{N} \int_E f_n\, d\mu = \sum_{n=1}^{\infty} \int_E f_n\, d\mu$.
*Structural Commentary*: Nonnegative series are monotone partial sums.
*Key Insight*: Integrate partial sums, then pass upward. □

**Corollary 1.3.3.** *Beppo Levi Bounded-Integral Criterion*
*Let $E \in \mathcal{M}$, let $(f_n)$ be measurable on $E$, and assume $0 \le f_n \uparrow f$ with $\sup_n \int_E f_n\, d\mu < +\infty$.*
*Then $f < +\infty$ a.e. on $E$ and $\int_E f\, d\mu = \lim_n \int_E f_n\, d\mu < +\infty$.*

*Proof.* The monotone limit $f$ is measurable, and $\int_E f\, d\mu = \lim_n \int_E f_n\, d\mu \le \sup_n \int_E f_n\, d\mu < +\infty$ by Theorem 1.3.1. Thus $f < +\infty$ a.e. on $E$ by Corollary 1.2.17.
*Structural Commentary*: Bounded mass survives monotone limits.
*Key Insight*: Finite nonnegative integral forces a.e. finiteness. □

**Proposition 1.3.4.** *$L^1$ Approximation and Bounded Multipliers*
*Let $f \in \mathscr{L}^1(\mu)$ on $X$.*
*(i) $\forall \varepsilon > 0$ $\exists$ a measurable simple function $\varphi$ with $\int_X |f - \varphi|\, d\mu < \varepsilon$.*
*(ii) If $g$ is bounded measurable, then $fg \in \mathscr{L}^1(\mu)$.*

*Proof.*
(i) Choose simple $0 \le s_n \uparrow f^+$ and $0 \le t_n \uparrow f^-$. Then $\int_X (f^+ - s_n)\, d\mu \to 0$ and $\int_X (f^- - t_n)\, d\mu \to 0$.
For large $n$, $\varphi := s_n - t_n$ satisfies $\int_X |f - \varphi|\, d\mu \le \int_X (f^+ - s_n)\, d\mu + \int_X (f^- - t_n)\, d\mu < \varepsilon$.
(ii) If $|g| \le M$, then $fg$ is measurable and $\int_X |fg|\, d\mu \le M\int_X |f|\, d\mu < +\infty$.
*Structural Commentary*: $L^1$ is simple-closed and bounded-multiplier stable.
*Key Insight*: Approximate signs separately; dominate products by $M|f|$. □

**Theorem 1.3.5.** *Fatou's Lemma*
*Let $E \in \mathcal{M}$ and let $f_n : E \to [0, +\infty]$ be measurable. Then $\int_E \liminf_n f_n\, d\mu \le \liminf_n \int_E f_n\, d\mu$.*

*Proof.* Set $g_n := \inf_{i \geq n} f_i$. Then $0 \leq g_n \uparrow \liminf_i f_i$ and $\forall i \geq n$, $g_n \leq f_i$, so $\int_E g_n \, d\mu \leq \inf_{i \geq n} \int_E f_i \, d\mu$.
By Theorem 1.3.1, $\int_E \liminf_i f_i \, d\mu = \lim_n \int_E g_n \, d\mu \leq \lim_n \inf_{i \geq n} \int_E f_i \, d\mu = \liminf_n \int_E f_n \, d\mu$.
*Structural Commentary*: Replace lim inf by increasing tail infima.
*Key Insight*: Tail infima are monotone minorants. □

**Example 1.3.6.** *Strictness in Fatou's Lemma*
Fatou's inequality can be strict by concentration or escape of mass.

(i) *Concentration*: On $E = (0,1)$ with Lebesgue measure $m$, set $f_n := n\mathbf{1}_{(0,\frac{1}{n})}$.
Then $f_n \to 0$ pointwise and $\forall n$, $\int_E f_n \, dm = 1$.
(ii) *Escape*: On $E = \mathbb{R}$ with Lebesgue measure $m$, set $g_n := \mathbf{1}_{(n,n+1)}$.
Then $g_n \to 0$ pointwise and $\forall n$, $\int_E g_n \, dm = 1$.

In either case, with $h_n = f_n$ or $h_n = g_n$, $\int_E \liminf_n h_n \, dm = 0 < 1 = \liminf_n \int_E h_n \, dm$.




Figure 203: Strict Fatou behavior: $f_n = n\mathbf{1}_{(0,\frac{1}{n})}$ and $g_n = \mathbf{1}_{(n,n+1)}$ have integral 1 but vanish pointwise.

**Theorem 1.3.7.** *Dominated Convergence*
*Let $E \in \mathcal{M}$, let $f_n, f$ be measurable on $E$, and let $g \in \mathscr{L}^1(\mu)$ on $E$.*
*Suppose $f_n \to f$ a.e. on $E$ and $\forall n$, $|f_n| \leq g$ on $E$. Then:*

*(i) $f \in \mathscr{L}^1(\mu)$ on $E$.*
*(ii) $\lim_{n\to\infty} \int_E f_n \, d\mu = \int_E f \, d\mu$.*

*Proof.* Choose null $N \subseteq E$ such that $f_n \to f$ on $E \setminus N$.
Changing values on $N$ does not change integrals, so assume $f_n \to f$ everywhere on $E$.

(i) The pointwise limit $f$ is measurable on $E$, and $|f| \leq g$ on $E$; hence $f \in \mathscr{L}^1(\mu)$ on $E$ by Theorem 1.2.18(ii).
(ii) Put $a_n := \int_E f_n \, d\mu$. Fatou applied to $g + f_n \geq 0$ gives $\int_E (g + f) \, d\mu \leq \int_E g \, d\mu + \liminf_n a_n$.
Fatou applied to $g - f_n \geq 0$ gives $\int_E (g - f) \, d\mu \leq \int_E g \, d\mu - \limsup_n a_n$.
Subtracting $\int_E g \, d\mu < +\infty$ gives $\int_E f \, d\mu \leq \liminf_n a_n \leq \limsup_n a_n \leq \int_E f \, d\mu$.

*Structural Commentary*: Domination reduces signed convergence to two Fatou bounds.
*Key Insight*: Apply Fatou to $g + f_n$ and $g - f_n$. □

**Theorem 1.3.8.** *General Dominated Convergence*
*Let $E \in \mathcal{M}$, let $f_n, f, g$ be measurable on $E$, and let $g_n \geq 0$ be integrable on $E$.*
*Suppose $f_n \to f$ and $g_n \to g$ a.e. on $E$, with $\forall n$, $|f_n| \leq g_n$ a.e. on $E$.*
*Assume $\lim_n \int_E g_n \, d\mu = \int_E g \, d\mu < +\infty$. Then:*

*(i) $\forall n$, $f_n \in \mathscr{L}^1(\mu)$ on $E$, and $f \in \mathscr{L}^1(\mu)$ on $E$.*
*(ii) $\lim_{n\to\infty} \int_E f_n \, d\mu = \int_E f \, d\mu$.*

*Proof.* Choose null $N \subseteq E$ so that $f_n \to f$, $g_n \to g$, and $\forall n$, $|f_n| \leq g_n$ on $E \setminus N$.
Changing values on $N$ does not change integrals, so assume these relations hold on $E$.

(i) Since $|f_n| \leq g_n \in \mathscr{L}^1(\mu)$, $f_n \in \mathscr{L}^1(\mu)$ on $E$ by Theorem 1.2.18(ii).
Passing to the limit gives $|f| \leq g$, and $\int_E g \, d\mu < +\infty$, hence $f \in \mathscr{L}^1(\mu)$ on $E$.
(ii) Put $a_n := \int_E f_n \, d\mu$, $b_n := \int_E g_n \, d\mu$, and $b := \int_E g \, d\mu$; then $b_n \to b$.
Fatou applied to $g_n + f_n \geq 0$ gives $b + \int_E f \, d\mu \leq \liminf_n (b_n + a_n) = b + \liminf_n a_n$.
Fatou applied to $g_n - f_n \geq 0$ gives $b - \int_E f \, d\mu \leq \liminf_n (b_n - a_n) = b - \limsup_n a_n$.
Subtracting $b < +\infty$ gives $\limsup_n a_n \leq \int_E f \, d\mu \leq \liminf_n a_n$, hence $a_n \to \int_E f \, d\mu$.

*Structural Commentary*: Varying domination works when masses converge.
*Key Insight*: Fatou on $g_n \pm f_n$ traps the limit. □

**Corollary 1.3.9.** *Monotone Convergence Almost Everywhere*
*Let $E \in \mathcal{M}$ and $0 \le f_n \uparrow f$ a.e. on $E$. Then $\int_E f_n \, d\mu \uparrow \int_E f \, d\mu$.*

*Proof.* Choose $N \subseteq E$ with $\mu(N) = 0$ and $f_n \uparrow f$ on $E \setminus N$. Set $g_n := f_n \mathbf{1}_{E\setminus N}$ and $g := f\mathbf{1}_{E\setminus N}$.
Then $0 \le g_n \uparrow g$, so $\int_E g_n \, d\mu \uparrow \int_E g \, d\mu$. Since $f_n = g_n$ and $f = g$ off $N$, the integrals over $N$ vanish.
*Structural Commentary*: Monotone convergence ignores null exceptional sets.
*Key Insight*: Remove the exceptional set, prove the theorem there, then add back zero. □

**Example 1.3.10.** *Counting Measure Integrals Are Series*
Let $\mu$ be counting measure on $(\mathbb{N}, 2^{\mathbb{N}})$. For $f : \mathbb{N} \to [0, +\infty]$, set $s_N := \sum_{n=1}^{N} f(n)\mathbf{1}_{\{n\}}$; then $0 \le s_N \uparrow f$, with the case $f(k) = +\infty$ giving both sides $+\infty$. Hence $\int_{\mathbb{N}} f \, d\mu = \lim_N \int_{\mathbb{N}} s_N \, d\mu = \lim_N \sum_{n=1}^{N} f(n) = \sum_{n=1}^{\infty} f(n)$.
*Structural Commentary*: Counting measure turns integration into summation.
*Key Insight*: Finite-support truncations are the simple-function partial sums.

**Example 1.3.11.** *Boundary Tests for Monotone Convergence and Fatou*
On $([0, +\infty), \mathcal{B}, m)$:

(i) $f_n := \mathbf{1}_{[0,n]} \uparrow \mathbf{1}_{[0,+\infty)}$ and $\int f_n \, dm = n \uparrow +\infty$; monotone convergence allows $+\infty$.
(ii) $g_n := n^{-1}\mathbf{1}_{[n,+\infty)} \downarrow 0$ uniformly, but $\int g_n \, dm = +\infty \ \forall n$; decreasing convergence needs finite mass.
(iii) $w_n := -n^{-1}\mathbf{1}_{[0,n]} \rightrightarrows 0$, but $\int w_n \, dm = -1$; Fatou can fail without $f_n \ge 0$ or $f_n \ge -h \in L^1$.

**Theorem 1.3.12.** *Fatou with an Integrable Lower Bound*
*Let $h \ge 0$, $h \in \mathscr{L}^1(\mu)$, and let $f_n$ be measurable with $f_n \ge -h$.*
*Then, whenever the signed integrals are defined in $\overline{\mathbb{R}}$, $\int \liminf_{n\to\infty} f_n \, d\mu \le \liminf_{n\to\infty} \int f_n \, d\mu$.*

*Proof.* Set $F_n := f_n + h \ge 0$. Fatou gives $\int \liminf_n F_n \, d\mu \le \liminf_n \int F_n \, d\mu$.
Since $\liminf_n F_n = h + \liminf_n f_n$ and $\int F_n \, d\mu = \int f_n \, d\mu + \int h \, d\mu$, subtract the finite term $\int h \, d\mu$.
*Structural Commentary*: Fatou survives controlled negative parts.
*Key Insight*: Add an integrable floor to make the sequence nonnegative. □

**Remark 1.3.13.** *Why the Negative Example Has No Floor*
For $w_n := -n^{-1}\mathbf{1}_{[0,n]}$, any $h \in L^1$ with $-h \le w_n \ \forall n$ must satisfy $h \ge n^{-1}$ on $[0, n]$.
Thus $h \ge k^{-1}$ on $[k-1, k]$, and $\int_0^N h \, dm \ge H_N := \sum_{k=1}^{N} \frac{1}{k} \to +\infty$, contradicting $h \in L^1$.

**Perspective 1.3.14.** *Domination Is the Safeguard*
$f_n \to f$ a.e. need not imply $\int f_n \to \int f$; domination $|f_n| \le g \in L^1$ restores the limit interchange.

**Example 1.3.15.** *A Nonuniform Limit with Vanishing Integrals*
For $x > 0$, set $f_n(x) := e^{-nx}x^{-\frac{1}{2}}$. Then $f_n \to 0$ pointwise, but $\forall \delta > 0$, $\sup_{0<x<\delta} f_n(x) = +\infty$.
For $n \ge 1$, $0 \le f_n \le g := e^{-x}x^{-\frac{1}{2}}$; $g \le x^{-\frac{1}{2}}\mathbf{1}_{(0,1]} + e^{-x}\mathbf{1}_{[1,\infty)} \in \mathscr{L}^1(0, \infty)$.
Thus dominated convergence gives $I_n := \int_0^{\infty} e^{-nx}x^{-\frac{1}{2}} \, dx \to 0$.
*Structural Commentary*: Uniform convergence is unnecessary once one integrable envelope controls the sequence.
*Key Insight*: Domination handles the singular behavior at 0 and the tail at infinity simultaneously.

**Proposition 1.3.16.** *Uniform Convergence on Finite Measure Spaces*
*Assume $\mu(X) < +\infty$.*

*(i) If $0 \le f_n$ are measurable and $f_n \rightrightarrows f$, then $f \ge 0$ is measurable and $\int_X f \, d\mu = \lim_n \int_X f_n \, d\mu$ in $[0, +\infty]$.*
*(ii) If $f_n \in \mathscr{L}^1(\mu)$ and $f_n \rightrightarrows f$, then $f \in \mathscr{L}^1(\mu)$ and $\int_X f_n \, d\mu \to \int_X f \, d\mu$ in $\mathbb{R}$.*

*Proof.*

(i) Measurability and $f \ge 0$ pass to pointwise limits. For $\varepsilon > 0$ and $n \gg 1$, $f \le f_n + \varepsilon\mathbf{1}_X$ and $f_n \le f + \varepsilon\mathbf{1}_X$.
Thus $\int_X f \, d\mu \le \int_X f_n \, d\mu + \varepsilon\mu(X)$ and $\int_X f_n \, d\mu \le \int_X f \, d\mu + \varepsilon\mu(X)$. Let $\varepsilon \downarrow 0$.
(ii) Choose $N$ with $|f - f_N| \le 1$. Then $f = f_N + (f - f_N) \in \mathscr{L}^1(\mu)$, and
$\left|\int_X f_n \, d\mu - \int_X f \, d\mu\right| \le \int_X |f_n - f| \, d\mu \le \mu(X) \sup_X |f_n - f| \to 0$.

*Structural Commentary*: On finite measure spaces, uniform error controls integral error.
*Key Insight*: Finiteness of $\mu(X)$ turns a supremum bound into an integral bound. □

**Theorem 1.3.17.** *Bounded Convergence on Finite Measure Spaces*
*Let $\mu(X) < +\infty$, and let $f_n, f : X \to \mathbb{R}$ be measurable.*
*Suppose $f_n \to f$ a.e. and $\exists M < +\infty$ such that $\forall n,\ |f_n| \le M$ a.e. Then:*
*(i)* $f \in \mathscr{L}^1(\mu)$.
*(ii)* $\lim_{n\to\infty} \int_X f_n \, d\mu = \int_X f \, d\mu$.

*Proof.* Since $|f_n| \le M$ a.e. and $f_n \to f$ a.e., $|f| \le M$ a.e.
Set $G := M\mathbf{1}_X$. Then $G \in \mathscr{L}^1(\mu)$, since $\int_X G \, d\mu = M\mu(X) < +\infty$.
(i) Since $|f| \le G$ a.e., $f \in \mathscr{L}^1(\mu)$ by Theorem 1.2.18(ii).
(ii) Since $\forall n,\ |f_n| \le G$ a.e., $\int_X f_n \, d\mu \to \int_X f \, d\mu$ by Theorem 1.3.7(ii).
*Structural Commentary*: On a finite-measure space, a uniform pointwise bound is an integrable dominator.
*Key Insight*: Bounded convergence is the finite-measure special case of dominated convergence. □

**Example 1.3.18.** *Why the Finite-Measure Hypothesis Matters*
On $(\mathbb{R}, m)$, let $f_n := n^{-1}\mathbf{1}_{[0,n]}$. Then $\|f_n\|_\infty = \frac{1}{n} \to 0$, but $\int_{\mathbb{R}} f_n \, dm = 1\ \forall n$.
Thus uniform convergence need not commute with integrals when $m(\mathbb{R}) = +\infty$; Fatou only yields $0 \le 1$.

**Example 1.3.19.** *Why Domination Cannot Be Dropped*
On $(\mathbb{R}, m)$, let $f_n := n\mathbf{1}_{[\frac{1}{n}, \frac{2}{n}]}$. Then $f_n \to 0$ pointwise, but $\int_{\mathbb{R}} f_n \, dm = 1\ \forall n$.
Thus pointwise convergence without domination need not commute with integrals; Fatou only yields $0 \le 1$.

**Theorem 1.3.20.** *Absolutely Summable $L^1$ Series*
*Let $f_n \in \mathscr{L}^1(\mu)$ and $\sum_{n\ge1} \|f_n\|_1 < +\infty$.*
*Set $S_N := \sum_{n=1}^{N} f_n$. Then $\exists f : X \to \mathbb{R}$ measurable such that:*
*(i)* $S_N \to f$ *absolutely a.e.*
*(ii)* $f \in \mathscr{L}^1(\mu)$.
*(iii)* $\int_X f \, d\mu = \sum_{n\ge1} \int_X f_n \, d\mu$.

*Proof.* Set $G := \sum_{n\ge1} |f_n|$. By monotone convergence, $\int_X G \, d\mu = \sum_{n\ge1} \|f_n\|_1 < +\infty$, so $G < +\infty$ a.e. and $G \in \mathscr{L}^1(\mu)$. Define $f := \lim_N S_N$ on $\{G < +\infty\}$ and $f := 0$ elsewhere.
(i) Since $G < +\infty$ a.e., $\sum_{n\ge1} |f_n| < +\infty$ a.e.; hence $S_N \to f$ absolutely a.e.
(ii) Since $|f| \le G$ a.e. and $G \in \mathscr{L}^1(\mu)$, $f \in \mathscr{L}^1(\mu)$ by Theorem 1.2.18(ii).
(iii) Since $|S_N| \le G$ and $S_N \to f$ a.e., dominated convergence gives
$$\int_X f \, d\mu = \lim_N \int_X S_N \, d\mu = \lim_N \sum_{n=1}^{N} \int_X f_n \, d\mu = \sum_{n\ge1} \int_X f_n \, d\mu.$$
*Structural Commentary*: Absolute summability in $L^1$ gives a.e. absolute convergence and termwise integration.
*Key Insight*: First integrate the nonnegative majorant $\sum |f_n|$. □

**Proposition 1.3.21.** *$L^1$ Convergence Controls Absolute Integrals*
*If $f_n, f \in \mathscr{L}^1(\mu)$ and $\|f_n - f\|_1 \to 0$, then:*
*(i)* $\|f_n\|_1 \to \|f\|_1$.
*(ii)* $\int_X |f| \, d\mu = \lim_n \int_X |f_n| \, d\mu$.

*Proof.*
(i) The reverse triangle inequality gives $||f_n| - |f|| \le |f_n - f|$; hence $|\|f_n\|_1 - \|f\|_1| \le \|f_n - f\|_1 \to 0$.
(ii) Since $\|h\|_1 = \int_X |h| \, d\mu$, item (i) gives $\int_X |f_n| \, d\mu \to \int_X |f| \, d\mu$.
*Structural Commentary*: $L^1$ convergence is strong enough to converge the $L^1$ sizes.
*Key Insight*: The reverse triangle inequality converts function convergence into norm convergence. □

**Proposition 1.3.22.** *Finite Integral Localizes to Finite Measure*
*Let $f : X \to [0, +\infty]$ be measurable and $\int_X f \, d\mu < +\infty$. For $n \ge 1$, set $E_n := \{\frac{1}{n} < f \le n\}$. Then:*
*(i)* $\forall n,\ \mu(E_n) < +\infty$.
*(ii)* $\int_{X\setminus E_n} f \, d\mu \downarrow 0$.
*Consequently, $\forall \varepsilon > 0\ \exists E \in \mathcal{M}$ with $\mu(E) < +\infty$ and $\int_{X\setminus E} f \, d\mu < \varepsilon$.*

*Proof.*

(i) Since $f > \frac{1}{n}$ on $E_n$, $\mu(E_n) \le n \int_{E_n} f \, d\mu \le n \int_X f \, d\mu < +\infty$.

(ii) Since $f\mathbf{1}_{E_n} \uparrow f$ a.e., monotone convergence gives $\int_{E_n} f \, d\mu \uparrow \int_X f \, d\mu$.
Thus $\int_{X \setminus E_n} f \, d\mu = \int_X f \, d\mu - \int_{E_n} f \, d\mu \downarrow 0$.

Given $\varepsilon > 0$, choose $n$ with $\int_{X \setminus E_n} f \, d\mu < \varepsilon$ and set $E = E_n$.
*Structural Commentary*: Finite integral mass can be captured on a finite-measure part of the space.
*Key Insight*: Keep the levels where $f$ is bounded above and away from zero. □

**Proposition 1.3.23.** *Setwise Convergence from Total Mass Convergence*
*Let $f_n, f : X \to [0, +\infty]$ be measurable, $f_n \to f$ pointwise, and $\int_X f \, d\mu = \lim_{n\to\infty} \int_X f_n \, d\mu < +\infty$.*
*Then $\forall E \in \mathcal{M}$, $\int_E f \, d\mu = \lim_{n\to\infty} \int_E f_n \, d\mu$.*

*Proof.* Fix $E \in \mathcal{M}$ and set $E^c := X \setminus E$. By Fatou, $\int_E f \, d\mu \le \liminf_n \int_E f_n \, d\mu$ and $\int_{E^c} f \, d\mu \le \liminf_n \int_{E^c} f_n \, d\mu$.
Since $\int_X f_n \, d\mu = \int_E f_n \, d\mu + \int_{E^c} f_n \, d\mu \to \int_X f \, d\mu < +\infty$, $\limsup_n \int_E f_n \, d\mu \le \int_X f \, d\mu - \int_{E^c} f \, d\mu = \int_E f \, d\mu$.
Thus $\liminf_n \int_E f_n \, d\mu = \limsup_n \int_E f_n \, d\mu = \int_E f \, d\mu$.
*Structural Commentary*: Total mass plus Fatou on complements fixes each part.
*Key Insight*: A lower bound on the complement is an upper bound on the chosen set. □

**Example 1.3.24.** *Why the Finite Total Mass Assumption Matters*
Let $X = \mathbb{N} \times \{0, 1\}$ with counting measure, set $E := \mathbb{N} \times \{1\}$.
Define $f := \mathbf{1}_{\mathbb{N}\times\{0\}}$; $f_n := \mathbf{1}_{\{1,\dots,n\}\times\{0\}} + n\mathbf{1}_{\{n\}\times\{1\}}$.
Then $f_n \to f$ pointwise and $\int_X f \, d\mu = \lim_n \int_X f_n \, d\mu = +\infty$; $\int_E f \, d\mu = 0$, $\int_E f_n \, d\mu = n$.
Thus Proposition 1.3.23 can fail when the common total mass is infinite.

**Definition 1.3.25.** *Function Truncates*
For $f : X \to \mathbb{R}$ and $n \in \mathbb{N}$, define the *nth truncate*

$$T_n f(x) := (-n) \vee (f(x) \wedge n) = \begin{cases} -n, & f(x) < -n, \\ f(x), & -n \le f(x) \le n, \\ n, & f(x) > n. \end{cases}$$

**Proposition 1.3.26.** *Truncates Characterize Integrability*
*Let $f : X \to \mathbb{R}$ be measurable and set $f_n := T_n f$ for $n \in \mathbb{N}$.*
*(i) $f \in \mathscr{L}^1(\mu) \iff \sup_n \|f_n\|_1 < +\infty$.*
*(ii) If $f \in \mathscr{L}^1(\mu)$, then $\int_X f_n \, d\mu \to \int_X f \, d\mu$.*

*Proof.* $\forall n$, $T_n : \mathbb{R} \to [-n, n]$ is continuous, so $f_n$ is measurable; moreover $f_n \to f$, $|f_n| \le |f|$, and $|f_n| \uparrow |f|$.

(i) ($\Rightarrow$) If $f \in \mathscr{L}^1(\mu)$, then $\|f_n\|_1 \le \|f\|_1$ for all $n$, so $\sup_n \|f_n\|_1 < +\infty$.
(i) ($\Leftarrow$) If $\sup_n \|f_n\|_1 < +\infty$, then MCT gives $\|f\|_1 = \lim_{n\to\infty} \|f_n\|_1 = \sup_n \|f_n\|_1 < +\infty$, so $f \in \mathscr{L}^1(\mu)$.
(ii) If $f \in \mathscr{L}^1(\mu)$, then DCT applied to $f_n \to f$ with $|f_n| \le |f|$ gives $\int_X f_n \, d\mu \to \int_X f \, d\mu$.
*Structural Commentary*: Truncation reduces $L^1$ questions to bounded approximants.
*Key Insight*: MCT reads $\|f\|_1$ from $|f_n| \uparrow |f|$; DCT passes signed integrals. □

## 1.4 Parameter Integrals and Differentiation

**Corollary 1.4.1.** *Continuity Under a Lebesgue Integral*
*Let $f_t(x) := f(x, t)$ be measurable $\forall t \in [a, b]$, and suppose $\forall t \in [a, b]$, $|f_t| \le g \in \mathscr{L}^1(\mu)$. Then:*
*(i) If $t_0 \in [a, b]$ and $f_t \to f_{t_0}$ a.e. as $t \to t_0$, then $\lim_{t\to t_0} \int_X f_t \, d\mu = \int_X f_{t_0} \, d\mu$.*
*(ii) If $t \mapsto f(x, t)$ is continuous on $[a, b]$ for a.e. $x$, then $F(t) := \int_X f_t \, d\mu$ is continuous on $[a, b]$.*

*Proof.*

(i) For any $t_n \to t_0$, set $u_n := f_{t_n}$. Then $u_n \to f_{t_0}$ a.e. and $|u_n| \le g$, so dominated convergence gives $\int_X u_n \, d\mu \to \int_X f_{t_0} \, d\mu$. The sequential criterion gives the claim.
(ii) If $t \mapsto f(x, t)$ is continuous a.e., then $\forall t_0 \in [a, b]$, $f_t \to f_{t_0}$ a.e. as $t \to t_0$.
By item (i), $\forall t_0 \in [a, b]$, $F(t) \to F(t_0)$, so $F$ is continuous on $[a, b]$.

*Structural Commentary*: Parameter continuity reduces to dominated convergence along sequences.
*Key Insight*: The parameter disappears once $t_n$ is frozen into a sequence of functions. □

**Theorem 1.4.2.** *Differentiation Under a Lebesgue Integral*
*Let $f_t(x) := f(x,t)$ be measurable $\forall t \in [a,b]$, and suppose $f_{t_0} \in \mathscr{L}^1(\mu)$ for some $t_0 \in [a,b]$.*
*Assume $t \mapsto f(x,t)$ is continuous on $[a,b]$ and differentiable on $(a,b)$ $\forall x \in X$, and $\exists g \in \mathscr{L}^1(\mu)$ such that $\forall (x,t) \in X \times (a,b)$, $|\partial_t f(x,t)| \le g(x)$. For $t \in (a,b)$, set $D_t(x) := \partial_t f(x,t)$; set $F(t) := \int_X f_t \, d\mu$. Then:*
*(i) $\forall t \in (a,b)$, $D_t$ is measurable.*
*(ii) $\forall t \in [a,b]$, $f_t \in \mathscr{L}^1(\mu)$.*
*(iii) $F$ is differentiable on $(a,b)$, and $\forall t \in (a,b)$, $F'(t) = \int_X D_t \, d\mu$.*

*Proof.*
(i) Fix $t \in (a,b)$ and choose $h_k \to 0$ with $t + h_k \in [a,b]$, $h_k \neq 0$.
The quotients $\frac{f_{t+h_k} - f_t}{h_k}$ are measurable and converge pointwise to $D_t$, so $D_t$ is measurable.
(ii) By the mean value theorem, $|f_t - f_{t_0}| \le |t - t_0|\, g$, so $|f_t| \le |f_{t_0}| + |t - t_0|\, g \in \mathscr{L}^1(\mu)$.
(iii) If $t_n \to t \in (a,b)$ and $t_n \neq t$, then $\frac{F(t_n)-F(t)}{t_n - t} = \int_X \frac{f_{t_n} - f_t}{t_n - t} \, d\mu$. The integrand converges pointwise to $D_t$ and has modulus at most $g$, so dominated convergence gives $\frac{F(t_n)-F(t)}{t_n - t} \to \int_X D_t \, d\mu$.

*Structural Commentary*: Bounded parameter derivatives dominate the difference quotients.
*Key Insight*: Differentiating the integral is just applying dominated convergence to secant slopes. □

**Corollary 1.4.3.** *A Pointwise Difference-Quotient Test*
*Let $f_t(x) := f(x,t)$ be measurable $\forall t \in [a,b]$, let $f_{t_0} \in \mathscr{L}^1(\mu)$ for some $t_0 \in [a,b]$, and fix $t_1 \in [a,b]$.*
*Assume $\partial_t f(x,t_1)$ exists $\forall x$. For $t \neq t_1$, set $Q_t(x) := \frac{f_t(x) - f_{t_1}(x)}{t - t_1}$, $|Q_t| \le g$ for some $g \in \mathscr{L}^1(\mu)$.*
*Set $F(t) := \int_X f_t \, d\mu$. Then:*
*(i) $\forall t \in [a,b]$, $f_t \in \mathscr{L}^1(\mu)$.*
*(ii) $F$ is differentiable at $t_1$, and $F'(t_1) = \int_X \partial_t f(x,t_1) \, d\mu$.*

*Proof.*
(i) If $t_0 \neq t_1$, then $|f_{t_1}| \le |f_{t_0}| + |t_0 - t_1|\, g$; if $t_0 = t_1$, this is given.
Hence $f_{t_1} \in \mathscr{L}^1(\mu)$. For $t \neq t_1$, $|f_t| \le |f_{t_1}| + |t - t_1|\, g$, so $f_t \in \mathscr{L}^1(\mu)$.
(ii) For any $t_n \to t_1$ with $t_n \neq t_1$, $Q_{t_n} \to \partial_t f(\cdot, t_1)$ pointwise and $|Q_{t_n}| \le g$.
Dominated convergence gives $\frac{F(t_n)-F(t_1)}{t_n - t_1} \to \int_X \partial_t f(x,t_1) \, d\mu$.

*Structural Commentary*: One parameter point needs only domination of the secants into that point.
*Key Insight*: A global derivative bound is sufficient, but not necessary, for a one-point formula. □

**Example 1.4.4.** *Differentiating Under the Integral Sign*
For $t > 0$, set $F(t) := \int_0^\infty x^2 e^{-tx} \, dx$. For $t_0 > 0$ and $0 < |h| < \frac{t_0}{2}$, put $Q_h(x) := \frac{x^2 e^{-(t_0+h)x} - x^2 e^{-t_0 x}}{h}$.
Then $Q_h \to -x^3 e^{-t_0 x}$ pointwise and $|Q_h| \le x^3 e^{-\frac{t_0 x}{2}} \in \mathscr{L}^1(0,\infty)$. Hence $F'(t_0) = -\int_0^\infty x^3 e^{-t_0 x} \, dx$.
*Structural Commentary*: Parameter differentiation is a limit-interchange problem.
*Key Insight*: Prove a local-in-parameter integrable majorant for the difference quotients.

**Example 1.4.5.** *Laplace Transform Powers and Factorials*
For $t > 0$, set $I(t) := \int_0^\infty e^{-tx} \, dx$. Then $I(t) = \lim_{b \to +\infty} \frac{1 - e^{-tb}}{t} = t^{-1}$.
For $[a,b] \subset (0,+\infty)$ and $k \ge 0$, $0 \le x^k e^{-tx} \le x^k e^{-ax} \in \mathscr{L}^1(0,\infty)$, so repeated differentiation under the integral gives $I^{(n)}(t) = (-1)^n \int_0^\infty x^n e^{-tx} \, dx$. Since $I^{(n)}(t) = (-1)^n n! t^{-n-1}$, $\forall n \in \mathbb{N} \cup \{0\}$, $\int_0^\infty x^n e^{-x} \, dx = n!$.
*Structural Commentary*: A parameter integral turns powers of $x$ into derivatives in $t$.
*Key Insight*: The exponential envelope $e^{-ax}$ controls all nearby parameter values $t \ge a$.

**Corollary 1.4.6.** *One Lebesgue and One Riemann Integral*
*Let $f_t(x) := f(x,t)$, $t \in [a,b]$, with $f_t$ measurable, $t \mapsto f(x,t)$ continuous a.e., and $\forall t \in [a,b]$, $|f_t| \le g \in \mathscr{L}^1(\mu)$.*
*Then, with Riemann integrals in $t$, $\int_a^b \left[\int_X f_t \, d\mu\right] dt = \int_X \left[\int_a^b f(x,t) \, dt\right] d\mu(x)$.*

*Proof.* Set $h_t(x) := \int_a^t f(x,s) \, ds$. Then $h_t$ is measurable, $|h_t| \le (b-a)g$, and $\partial_t h_t = f_t$.
By differentiation under the integral, $H(t) := \int_X h_t \, d\mu$ satisfies $H'(t) = \int_X f_t \, d\mu$.
Hence $\int_a^b \left[\int_X f_t \, d\mu\right] dt = H(b) - H(a) = \int_X \left[\int_a^b f(x,t) \, dt\right] d\mu(x)$.
*Structural Commentary*: Finite parameter integrals move through by differentiating primitives.
*Key Insight*: Integrate first in the parameter up to $t$, then differentiate the resulting Lebesgue integral. □

**Corollary 1.4.7.** *Improper Parameter Interchange*
*Let $f_t(x) := f(x,t)$ be measurable $\forall t \in \mathbb{R}$, and let $t \mapsto f(x,t)$ be continuous for a.e. $x$.*

*Suppose $\exists g, h \in \mathscr{L}^1(\mu)$ such that $|f_t| \le g\ \forall t$ and $\int_{\mathbb{R}} |f(x,t)|\,dt \le h(x)$ for a.e. $x$.*
*Put $G(t) := \int_X f_t\,d\mu$, $H(x) := \int_{\mathbb{R}} f(x,t)\,dt$, and $H_R(x) := \int_{-R}^{R} f(x,t)\,dt$. Then:*

*(i)* $H \in \mathscr{L}^1(\mu)$.
*(ii)* $\lim_{R\to+\infty} \int_{-R}^{R} G(t)\,dt = \int_X H\,d\mu$.

*Thus $\int_{\mathbb{R}} G := \lim_{R\to+\infty} \int_{-R}^{R} G(t)\,dt$ exists and equals $\int_X H\,d\mu$.*

*Proof.*

(i) Since $|H| \le \int_{\mathbb{R}} |f(x,t)|\,dt \le h$ a.e. and $h \in \mathscr{L}^1(\mu)$, $H \in \mathscr{L}^1(\mu)$ by Theorem 1.2.18(ii).
(ii) For $R > 0$, finite-interval interchange gives $\int_{-R}^{R} G(t)\,dt = \int_X H_R\,d\mu$.
Since $H_R \to H$ a.e. and $|H_R| \le h$, dominated convergence gives $\int_X H_R\,d\mu \to \int_X H\,d\mu$.

*Structural Commentary*: Finite interchanges pass to improper limits under an $L^1$ bound.
*Key Insight*: Truncate in $t$, interchange on finite intervals, then pass $R \to +\infty$ by domination in $x$. □

## 1.5 Riemann-Lebesgue Comparison

**Lemma 1.5.1.** *Monotone Integral Squeeze*
*Let $E \in \mathcal{M}$, $(\varphi_n), (\psi_n) \subseteq \mathscr{L}^1(\mu)$ on $E$, and $f : E \to \mathbb{R}$.*
*Assume $\varphi_n \uparrow$, $\psi_n \downarrow$, $\varphi_n \le f \le \psi_n$ on $E$, and $\int_E (\psi_n - \varphi_n)\,d\mu \to 0$. Put $\varphi := \lim_n \varphi_n$ and $\psi := \lim_n \psi_n$. Then:*

*(i)* $\varphi = \psi$ *a.e. on* $E$.
*(ii) For $h := \varphi$, $f = h$ a.e., $h \in \mathscr{L}^1(\mu)$ on $E$, and $\int_E \varphi_n\,d\mu, \int_E \psi_n\,d\mu \to \int_E h\,d\mu$.*

*Proof.*

(i) The limits $\varphi, \psi$ are measurable and $\varphi \le f \le \psi$.
Since $0 \le \psi - \varphi \le \psi_n - \varphi_n$, $0 \le \int_E (\psi - \varphi)\,d\mu \le \int_E (\psi_n - \varphi_n)\,d\mu \to 0$; hence $\psi = \varphi$ a.e.
(ii) Let $h$ be the common a.e. value from (i). Then $f = h$ a.e. and $|h| \le |\varphi_1| + |\psi_1|$ a.e., so $h \in \mathscr{L}^1(\mu)$ on $E$.
Monotone convergence applied to $\varphi_n - \varphi_1 \uparrow h - \varphi_1$ and $\psi_1 - \psi_n \uparrow \psi_1 - h$ gives $\int_E \varphi_n\,d\mu, \int_E \psi_n\,d\mu \to \int_E h\,d\mu$.

*Structural Commentary*: Vanishing gaps collapse monotone envelopes.
*Key Insight*: Integral gaps force the two monotone limits to coincide a.e. □




Figure 204: Step envelopes squeeze $f$ to a common a.e. limit.

**Theorem 1.5.2.** *Comparison with the Riemann Integral*
*Let $m$ be Lebesgue measure on $[a,b]$ and let $f \in \mathcal{R}[a,b]$.*

*(i)* $f \in \mathscr{L}^1(m)$.
*(ii)* $\int_{[a,b]} f\,dm = \int_a^b f(x)\,dx$.

*Proof.* Choose nested partitions $P_n$ with $U(P_n, f) - L(P_n, f) \to 0$. Let $\varphi_n, \psi_n$ be lower and upper step envelopes.

(i) Lemma 1.5.1 gives $f \in \mathscr{L}^1(m)$.
(ii) Since $\int \varphi_n\,dm = L(P_n, f)$ and $\int \psi_n\,dm = U(P_n, f)$, Lemma 1.5.1 gives $\int_{[a,b]} f\,dm = \lim_n L(P_n, f) = \lim_n U(P_n, f) = \int_a^b f(x)\,dx$.

*Structural Commentary*: Darboux envelopes squeeze the Lebesgue integral.
*Key Insight*: Vanishing Darboux gap identifies the two integral notions. □

**Corollary 1.5.3.** *Riemann Integrable Functions as Monotone Step Limits*
*Let $f : [a,b] \to \mathbb{R}$ be Riemann integrable. Then $\exists (\varphi_n), (\psi_n)$ step-function sequences such that*

*(i) $\varphi_n \uparrow$, $\psi_n \downarrow$, and $\varphi_n \le f \le \psi_n$ on $[a,b]$;*

*(ii)* $\lim_n \int_a^b \varphi_n\,dx = \lim_n \int_a^b \psi_n\,dx = \int_a^b f\,dx$;

*(iii)* $\varphi_n, \psi_n \to f$ *a.e.; hence* $f$ *is Lebesgue measurable and* $\int_{[a,b]} f\,dm = \int_a^b f\,dx$.

*Proof.*

(i) Choose nested partitions $P_n$ with $\max_{I\in P_n} \ell(I) \to 0$ and $U(P_n, f) - L(P_n, f) \to 0$.
Let $\varphi_n, \psi_n$ be the lower and upper step envelopes on $P_n$; refinement gives $\varphi_n \uparrow$, $\psi_n \downarrow$, and $\varphi_n \le f \le \psi_n$.

(ii) Since $\int_a^b \varphi_n\,dx = L(P_n, f)$ and $\int_a^b \psi_n\,dx = U(P_n, f)$, by Darboux gap condition, both limits equal $\int_a^b f\,dx$.

(iii) By Lemma 1.5.1, $\exists h \in \mathscr{L}^1(m)$ with $h = f$ a.e., $\varphi_n, \psi_n \to h$ a.e., and $\int_{[a,b]} h\,dm = \lim_n \int_a^b \varphi_n\,dx = \int_a^b f\,dx$.

*Structural Commentary*: Riemann integrability is monotone step squeezing.
*Key Insight*: Refining partitions make Darboux sums into pointwise a.e. step approximations. □

**Corollary 1.5.4.** *Nonnegative Continuous Functions and Riemann Integrals*

*(i) If* $f \ge 0$ *is continuous on* $[a, b]$, *then* $\int_{[a,b]} f\,dm = \int_a^b f(x)\,dx$.

*(ii) If* $f \ge 0$ *is continuous on* $[0, +\infty)$, *then* $\int_{[0,+\infty)} f\,dm = \lim_{b\to+\infty} \int_0^b f(x)\,dx$ *in* $[0, +\infty]$.

*Proof.*

(i) Continuous $f$ on $[a, b]$ is Riemann integrable; by Riemann–Lebesgue comparison, $\int_{[a,b]} f\,dm = \int_a^b f(x)\,dx$.

(ii) Put $f_b := f\mathbf{1}_{[0,b]}$. By (i), $\int_{[0,b]} f\,dm = \int_0^b f(x)\,dx$. As $b \uparrow +\infty$, $f_b \uparrow f$; monotone convergence gives the left limit, and the right limit is the nonnegative improper Riemann integral.

*Structural Commentary*: Compact comparison plus monotone truncation.
*Key Insight*: Truncate to compact intervals and pass upward. □

**Theorem 1.5.5.** *Lebesgue's Criterion for Riemann Integrability*
*Let* $f : [a, b] \to \mathbb{R}$ *be bounded with discontinuity set* $D_f$. *Then* $f \in \mathcal{R}[a, b] \iff m(D_f) = 0$.

*Proof.* Set $\omega_f(x) := \lim_{r\downarrow 0} \operatorname{diam} f([a, b] \cap (x - r, x + r))$ and $D_k := \{x : \omega_f(x) \ge \frac{1}{k}\}$; then $D_f = \bigcup_{k\ge 1} D_k$.

($\Rightarrow$) Fix $k$ and $\varepsilon > 0$. Choose $P$ with $U(P, f) - L(P, f) < \frac{\varepsilon}{k}$. Set $\mathcal{B} := \{I \in P : \operatorname{int} I \cap D_k \ne \emptyset\}$.
Then $\frac{1}{k} \sum_{I\in\mathcal{B}} \ell(I) \le U(P, f) - L(P, f)$, so $\sum_{I\in\mathcal{B}} \ell(I) < \varepsilon$.
These intervals plus the partition points cover $D_k$; hence $m(D_k) = 0$ and $m(D_f) = 0$.

($\Leftarrow$) Suppose $m(D_f) = 0$ and $|f| \le B$. If $B = 0$, $f \in \mathcal{R}[a, b]$. For $B > 0$ and $\varepsilon > 0$, choose $k$ with $\frac{b-a}{k} < \frac{\varepsilon}{2}$ and cover the compact null set $D_k$ by finitely many intervals of total length $< \frac{\varepsilon}{4B}$.
Refining by this cover gives $U(P, f) - L(P, f) < \frac{b-a}{k} + \frac{\varepsilon}{2} < \varepsilon$, hence $f \in \mathcal{R}[a, b]$.

*Structural Commentary*: Darboux gaps measure oscillation.
*Key Insight*: Riemann failure is positive-measure discontinuity. □

**Example 1.5.6.** *Dirichlet Function: Lebesgue but Not Riemann*
Let $D := \mathbf{1}_{\mathbb{Q}\cap[0,1]}$. Since $\mathbb{Q} \cap [0, 1]$ is countable, $D = 0$ a.e. on $[0, 1]$, so $\int_{[0,1]} D\,dm = 0$.
Every nondegenerate subinterval $I \subseteq [0, 1]$ meets both $\mathbb{Q}$ and $\mathbb{R} \setminus \mathbb{Q}$, hence $\inf_I D = 0$ and $\sup_I D = 1$.
Thus all lower Darboux sums are 0, all upper Darboux sums are 1, and $D$ is not Riemann integrable.
*Structural Commentary*: Null support, dense oscillation.
*Key Insight*: A dense null set can be harmless for Lebesgue integration and fatal for Riemann integration.

**Perspective 1.5.7.** *Lebesgue's Criterion as a Roadmap*
For bounded $f : [a, b] \to \mathbb{R}$ with discontinuity set $D_f$, $f \in \mathcal{R}[a, b] \iff m(D_f) = 0$.
Thus finite, countable, and null Cantor-type $D_f$ are allowed, while $D_f = [a, b]$ is not.

## 1.6 Littlewood and Lusin Approximation

The measurable world is often controlled by ordinary topological objects after discarding a small exceptional set. In the real-line finite-measure setting, this principle has three recurring forms: measurable sets are nearly elementary, measurable functions are nearly continuous, and a.e. convergence is nearly uniform.

**Principle 1.6.1.** *Littlewood's Three Principles, Finite-Measure Form*
For subsets of the real line with finite measure:

(i) measurable sets are nearly finite unions of intervals; see Corollary 6.2.4;
(ii) measurable functions are nearly continuous; this is Lusin's theorem below;

(iii) a.e. pointwise convergence is nearly uniform; see Egoroff's theorem, Theorem 1.6.3.

*Key Insight*: The word "nearly" means after changing or discarding a set of arbitrarily small measure.

**Definition 1.6.2.** *Almost Uniform Convergence*
Let $(f_n)$ be measurable functions on $X$.

(i) $f_n \to f$ *almost uniformly* if $\forall \delta > 0\ \exists E_\delta \in \mathcal{M}$ with $\mu(E_\delta) < \delta$ and $f_n \rightrightarrows f$ on $X \setminus E_\delta$.
(ii) $(f_n)$ is *almost uniformly Cauchy* if $\forall \delta > 0\ \exists E_\delta \in \mathcal{M}$ with $\mu(E_\delta) < \delta$ s.t. $(f_n)$ uniformly Cauchy on $X \setminus E_\delta$.

**Theorem 1.6.3.** *Egoroff's Theorem*
*If $\mu(X) < +\infty$, $f_n \to f$ a.e., with all functions finite-valued off null set, then $f_n \to f$ almost uniformly.*

*Proof.* Absorb the null exceptional set into every later $E_\delta$, and assume $\forall x \in X,\ f_n(x) \to f(x)$.
For $m, n \in \mathbb{N}$, set $E_n(m) := \bigcup_{k=n}^{\infty} \{|f_k - f| \geq \frac{1}{m}\}$. Then $E_n(m) \downarrow \emptyset$, so $\mu(E_n(m)) \to 0$ by Corollary 5.1.1.
Given $\delta > 0$, choose $k_m$ with $\mu(E_{k_m}(m)) < \delta 2^{-m}$ and set $E_\delta := \bigcup_{m \geq 1} E_{k_m}(m)$.
Then $\mu(E_\delta) < \delta$, and $x \notin E_\delta$, $k \geq k_m$ imply $|f_k(x) - f(x)| < \frac{1}{m}$; hence $f_n \rightrightarrows f$ on $X \setminus E_\delta$.
*Structural Commentary*: Finite measure turns decreasing bad sets with empty intersection into small bad sets.
*Key Insight*: A.e. convergence is almost uniform on finite measure spaces. □

**Example 1.6.4.** *Egoroff Needs a Finite-Valued Limit*
On $([0,1], \lambda)$, set $f_n := n\mathbf{1}_{[0,1]}$. Then $\forall x,\ f_n(x) \to +\infty$, so the pointwise limit is not finite-valued. If $f_n \to f$ almost uniformly for finite-valued measurable $f$, then for some $E$ with $\lambda(E) < \frac{1}{2}$, $f_n \rightrightarrows f$ on $[0,1] \setminus E$; this nonempty set has $f_n(x) = n \to +\infty$, contradicting finite-valued pointwise convergence.

**Lemma 1.6.5.** *Continuous Extension from a Closed Subset of $\mathbb{R}$*
*Let $F \subseteq \mathbb{R}$ be closed and let $h : F \to \mathbb{R}$ be continuous. Then $\exists g \in C(\mathbb{R})$ such that $g|_F = h$.*

*Proof.* Cases $F = \emptyset$ and $F = \mathbb{R}$ are immediate. Otherwise $\mathbb{R} \setminus F = \bigsqcup_\alpha I_\alpha$, set $g = h$ on $F$; on each component:

(i) If $I_\alpha = (a, b)$, set $g(x) := \frac{b-x}{b-a}h(a) + \frac{x-a}{b-a}h(b)$.
(ii) If $I_\alpha = (-\infty, b)$, set $g(x) := h(b)$.
(iii) If $I_\alpha = (a, +\infty)$, set $g(x) := h(a)$.

The formulas give continuity off $F$; at $x \in F$, adjacent finite endpoints converge to $x$.
Hence continuity of $h$ gives $g(y) \to h(x)$ as $y \to x$. Thus $g \in C(\mathbb{R})$ and $g|_F = h$.
*Structural Commentary*: Fill complementary intervals.
*Key Insight*: Interpolate between boundary values. □

**Proposition 1.6.6.** *Simple Functions Are Nearly Continuous*
*Let $E \in \mathcal{L}^1$ with $m(E) < +\infty$, and $s : E \to \mathbb{R}$ measurable simple. $\forall \varepsilon > 0$, $\exists F \subseteq E$ and $g : \mathbb{R} \to \mathbb{R}$ such that:*

*(i) $F$ is closed.*
*(ii) $g \in C(\mathbb{R})$ and $g = s$ on $F$.*
*(iii) $m(E \setminus F) < \varepsilon$.*

*Proof.* If $E = \emptyset$, take $F = \emptyset$ and $g = 0$. Otherwise $s = \sum_{i=1}^{N} a_i \mathbf{1}_{E_i}$ with disjoint measurable $E_i$ and $E = \bigsqcup_i E_i$.
Choose closed $F_i \subseteq E_i$ with $m(E_i \setminus F_i) < \frac{\varepsilon}{N}$, set $F := \bigcup_i F_i$, and define $h : F \to \mathbb{R}$ by $h|_{F_i} := a_i$.

(i) Each $F_i$ is closed and $N < +\infty$, so $F$ is closed; also $F \subseteq E$.
(ii) For $x \in F_i$, the closed set $\bigcup_{j \neq i} F_j$ misses $x$, so $h$ is locally constant at $x$ in $F$; hence $h \in C(F)$.
$\exists g \in C(\mathbb{R})$ with $g|_F = h = s|_F$ by Lemma 1.6.5.
(iii) $m(E \setminus F) \leq \sum_i m(E_i \setminus F_i) < \varepsilon$.

*Structural Commentary*: Closed cores isolate simple levels.
*Key Insight*: Approximate each level set from inside, then extend. □

**Theorem 1.6.7.** *Lusin's Theorem*
*Let $E \in \mathcal{L}^1$ with $m(E) < +\infty$, $f : E \to \mathbb{R}$ measurable. $\forall \varepsilon > 0$, $\exists F \subseteq E$ and $g : \mathbb{R} \to \mathbb{R}$ such that:*

*(i) $F$ is closed.*
*(ii) $g \in C(\mathbb{R})$ and $g = f$ on $F$.*
*(iii) $m(E \setminus F) < \varepsilon$.*

*Proof.* Take simple $s_n : E \to \mathbb{R}$ with $s_n \to f$ pointwise.
Pick $A \subseteq E$ with $s_n \to f$ uniformly on $A$ and $m(E \setminus A) < \frac{\varepsilon}{4}$, and closed $F_0 \subseteq A$ with $m(A \setminus F_0) < \frac{\varepsilon}{4}$.
For $n \geq 1$, choose closed $F_n \subseteq E$ with $s_n|_{F_n} \in C(F_n)$ and $m(E \setminus F_n) < \varepsilon 2^{-(n+1)}$. Set $F := \bigcap_{n \geq 0} F_n$.

(i) $F$ is an intersection of closed sets and $F \subseteq E$.
(ii) Since $F \subseteq F_0 \subseteq A$, the continuous functions $s_n|_F$ converge uniformly to $f|_F$, so $f|_F \in C(F)$.
Since $F$ is closed and $f|_F \in C(F)$, Lemma 1.6.5 gives $\exists g \in C(\mathbb{R})$ with $g|_F = f|_F$.
(iii) $m(E \setminus F) \leq m(E \setminus F_0) + \sum_{n \geq 1} m(E \setminus F_n) < \varepsilon$.

*Structural Commentary*: Discard a small set to gain continuity.
*Key Insight*: Simple cores plus Egoroff give uniform continuous limits. □




Figure 205: Lusin's theorem: keep $F$, discard $E \setminus F$, and extend with $g = f$ on $F$.

## 1.7 Convex Functions and Jensen's Inequality

Convexity is the geometric condition behind Jensen's inequality: averages move inside the argument, while supporting lines convert the nonlinear statement into a linear integral estimate.

**Definition 1.7.1.** *Convex Function on an Interval*
Let $I \subseteq \mathbb{R}$ be an interval and let $\varphi : I \to \mathbb{R}$.

(i) $\varphi$ is *convex* iff $\forall x_1, x_2 \in I\ \forall \lambda \in [0,1]$, $\varphi(\lambda x_1 + (1-\lambda)x_2) \leq \lambda\varphi(x_1) + (1-\lambda)\varphi(x_2)$.
(ii) Equivalently, $\forall x_1 < x < x_2$ in $I$, $\varphi(x) \leq \frac{x_2 - x}{x_2 - x_1}\varphi(x_1) + \frac{x - x_1}{x_2 - x_1}\varphi(x_2)$.

*Key Insight*: Graphs lie below enclosing chords.

**Lemma 1.7.2.** *Chordal Slope Lemma*
*Let $I \subseteq \mathbb{R}$ be an interval, let $\varphi : I \to \mathbb{R}$ be convex, and let $x_1, x, x_2 \in I$ satisfy $x_1 < x < x_2$.*
*Then* $\frac{\varphi(x)-\varphi(x_1)}{x-x_1} \leq \frac{\varphi(x_2)-\varphi(x_1)}{x_2-x_1} \leq \frac{\varphi(x_2)-\varphi(x)}{x_2-x}$.

*Proof.* Since $x = \frac{x_2-x}{x_2-x_1}x_1 + \frac{x-x_1}{x_2-x_1}x_2$, convexity gives $(x_2-x_1)\varphi(x) \leq (x_2-x)\varphi(x_1) + (x-x_1)\varphi(x_2)$.
Rearrange, $(x_2-x_1)(\varphi(x)-\varphi(x_1)) \leq (x-x_1)(\varphi(x_2)-\varphi(x_1))$; $(x_2-x)(\varphi(x_2)-\varphi(x_1)) \leq (x_2-x_1)(\varphi(x_2)-\varphi(x))$.
Divide by $(x-x_1)(x_2-x_1) > 0$ and $(x_2-x)(x_2-x_1) > 0$, respectively.
*Structural Commentary*: One chord inequality, two rearrangements.
*Key Insight*: Convex secant slopes are nondecreasing. □

**Proposition 1.7.3.** *Increasing Derivative Gives Convexity*
*Let $I \subseteq \mathbb{R}$ be open and let $\varphi : I \to \mathbb{R}$ be differentiable. If $\varphi'$ is increasing on $I$, then $\varphi$ is convex on $I$.*
*In particular, this holds if $\varphi$ is twice differentiable and $\varphi'' \geq 0$ on $I$.*

*Proof.* For $a < b$ in $I$, set $S(a,b) := \frac{\varphi(b)-\varphi(a)}{b-a}$. Let $u < v$ and $0 < \lambda < 1$, and put $x = \lambda u + (1-\lambda)v$.
By MVT, $S(u,x) = \varphi'(c_1)$, $S(x,v) = \varphi'(c_2)$ for some $c_1 \in (u,x)$, $c_2 \in (x,v)$. As $c_1 < c_2$, $S(u,x) \leq S(x,v)$.
Clearing denominators, $\varphi(x) \leq \frac{v-x}{v-u}\varphi(u) + \frac{x-u}{v-u}\varphi(v) = \lambda\varphi(u) + (1-\lambda)\varphi(v)$; the cases $\lambda = 0, 1$ are equality.
If $\varphi'' \geq 0$, for $a < b$, MVT gives $\varphi'(b) - \varphi'(a) = \varphi''(d)(b-a) \geq 0$ for some $d \in (a,b)$, so $\varphi'$ is increasing.
*Structural Commentary*: MVT orders secant slopes.
*Key Insight*: Monotone slopes give convexity. □

**Example 1.7.4.** *Basic Convex Functions*
By Proposition 1.7.3, $x \mapsto x^p$ for $p \geq 1$, $x \mapsto e^{ax}$, and $x \mapsto \log(\frac{1}{x})$ are convex on $(0, +\infty)$, $\mathbb{R}$, and $(0, +\infty)$.

**Proposition 1.7.5.** *Finite Jensen Inequality*
*Let $I \subseteq \mathbb{R}$ be an interval, $\varphi : I \to \mathbb{R}$ be convex, $x_1, \ldots, x_n \in I$, $\lambda_1, \ldots, \lambda_n \geq 0$ with $n \in \mathbb{N}$, $\sum_{k=1}^{n} \lambda_k = 1$.*
*Set $\bar{x} := \sum_{k=1}^{n} \lambda_k x_k$. Then $\bar{x} \in I$ and $\varphi(\bar{x}) \leq \sum_{k=1}^{n} \lambda_k \varphi(x_k)$.*

*Proof.* Induct on $m = \#\{k : \lambda_k > 0\}$. If $m = 1$, equality holds.

For $m \geq 2$, relabel so $\lambda_1, \ldots, \lambda_m > 0$ and $\lambda_k = 0$ for $k > m$. Let $\Lambda = 1 - \lambda_m$ and $y = \Lambda^{-1} \sum_{k=1}^{m-1} \lambda_k x_k$.

By induction, $y \in I$ and $\varphi(y) \leq \Lambda^{-1} \sum_{k=1}^{m-1} \lambda_k \varphi(x_k)$.

Since $\bar{x} = \Lambda y + \lambda_m x_m$, convexity gives $\bar{x} \in I$ and $\varphi(\bar{x}) \leq \Lambda\varphi(y) + \lambda_m \varphi(x_m) \leq \sum_{k=1}^{n} \lambda_k \varphi(x_k)$.

*Structural Commentary*: Induct by peeling off one weight.
*Key Insight*: Jensen iterates convexity. □

**Lemma 1.7.6.** *One-Sided Slopes and Supporting Lines*
*Let $I \subseteq \mathbb{R}$ be open and let $\varphi : I \to \mathbb{R}$ be convex.*

*(i) $\forall x \in I$, $\varphi'(x-)$ and $\varphi'(x+)$ exist and are finite.*
*(ii) If $u < v$ in $I$, then $\varphi'(u-) \leq \varphi'(u+) \leq \frac{\varphi(v)-\varphi(u)}{v-u} \leq \varphi'(v-) \leq \varphi'(v+)$.*
*(iii) If $\alpha \in I$ and $\varphi'(\alpha-) \leq m_0 \leq \varphi'(\alpha+)$, then $\forall t \in I$, $\varphi(t) \geq \varphi(\alpha) + m_0(t - \alpha)$.*

*Proof.* For $a < b$ in $I$, write $S(a, b) := \frac{\varphi(b)-\varphi(a)}{b-a}$.

(i) Fix $x \in I$ and choose $a < x < b$ in $I$.
If $a < r < s < x$, then $S(a, x) \leq S(r, x) \leq S(s, x) \leq S(x, b)$ by Lemma 1.7.2, so $\lim_{r \uparrow x} S(r, x)$ exists in $\mathbb{R}$.
If $x < r < s < b$, the same lemma gives $S(a, x) \leq S(x, r) \leq S(x, s) \leq S(x, b)$, so $\lim_{r \downarrow x} S(x, r)$ exists in $\mathbb{R}$.

(ii) Let $u < v$ in $I$. For $r < u < s < v$ and $u < q < v < t$, the chordal slope inequalities give $S(r, u) \leq S(u, s) \leq S(u, v) \leq S(q, v) \leq S(v, t)$. Letting $r \uparrow u$, $s \downarrow u$, $q \uparrow v$, and $t \downarrow v$ gives asserted chain.

(iii) If $t = \alpha$, equality holds. If $t > \alpha$, then (ii) gives $m_0 \leq \varphi'(\alpha+) \leq S(\alpha, t)$, hence $\varphi(t) \geq \varphi(\alpha) + m_0(t - \alpha)$.
If $t < \alpha$, then (ii) gives $S(t, \alpha) \leq \varphi'(\alpha-) \leq m_0$, which rearranges to the same estimate.

*Structural Commentary*: Monotone secant slopes have finite one-sided limits.
*Key Insight*: One-sided slope bounds produce supporting lines. □




Figure 206: Color code: $y = \varphi(x)$ lies below the chord through $x_1, x_2$ and above the supporting line at $\alpha$.

**Theorem 1.7.7.** *Jensen's Inequality on $[0, 1]$*
*Let $m$ be Lebesgue measure on $[0, 1]$, let $\varphi : \mathbb{R} \to \mathbb{R}$ be convex, and let $f, \varphi \circ f \in L^1(m)$.*
*Then $\varphi\left(\int f \, dm\right) \leq \int \varphi(f) \, dm$.*

*Proof.* Set $\alpha := \int f \, dm$. By Lemma 1.7.6, choose $m_0 \in [\varphi'(\alpha-), \varphi'(\alpha+)]$; $\varphi(t) \geq \varphi(\alpha) + m_0(t - \alpha)$ $\forall t \in \mathbb{R}$.
Substituting $t = f$ and integrating gives $\int \varphi(f) \, dm \geq \varphi(\alpha) + m_0(\int f \, dm - \alpha) = \varphi(\alpha)$.
*Structural Commentary*: Support at the mean.
*Key Insight*: The linear term vanishes. □

**Remark 1.7.8.** *Integrability in Jensen*
Linear growth, $|\varphi(t)| \leq c_1 + c_2 |t|$, makes $\varphi \circ f \in L^1([0, 1])$ by Proposition 4.3.4.
If $\varphi \circ f \geq 0$, Jensen's right side may be extended-valued.

# 2 Product Measures and Fubini Theory

## 2.1 Product Measures

Product measure builds measurable rectangles and reads higher-dimensional integrals through slices. The abstract construction comes first; the Euclidean statements below are its geometric specialization.

**Definition 2.1.1.** *Product Measurable Spaces and Sections*
Let $(X, \mathcal{M})$, $(Y, \mathcal{N})$ be measurable spaces and $Z := X \times Y$.

(i) A *measurable rectangle* is $A \times B$ with $A \in \mathcal{M}$ and $B \in \mathcal{N}$.
Set $\mathcal{Z}_0 := \{ \bigcup_{j=1}^{n} (A_j \times B_j) : A_j \in \mathcal{M},\ B_j \in \mathcal{N}\}$ and $\mathcal{M} \otimes \mathcal{N} := \sigma\{A \times B : A \in \mathcal{M},\ B \in \mathcal{N}\}$.

(ii) For $E \subseteq Z$, the *sections* are $E_x := \{y \in Y : (x, y) \in E\}$ and $E^y := \{x \in X : (x, y) \in E\}$.

(iii) For $F : Z \to \overline{\mathbb{R}}$, the *function sections* are $F_x(y) := F(x, y)$ and $F^y(x) := F(x, y)$, where defined.

**Lemma 2.1.2.** *Finite Rectangle Algebra*

*(i) If $R_j = A_j \times B_j$, then $R_1 \cap R_2 = (A_1 \cap A_2) \times (B_1 \cap B_2)$ and $R_1 \setminus R_2 = \big[(A_1 \cap A_2) \times (B_1 \setminus B_2)\big] \sqcup \big[(A_1 \setminus A_2) \times B_1\big]$.*

*(ii) $\mathcal{Z}_0$ is an algebra on $Z$.*

*(iii) $\forall E \in \mathcal{Z}_0\ \exists n \geq 1$, $A_j \in \mathcal{M}$, and $B_j \in \mathcal{N}$ such that $E = \bigsqcup_{j=1}^{n} (A_j \times B_j)$.*

*Proof.*

(i) For $(x, y) \in Z$, $(x, y) \in R_1 \cap R_2 \iff x \in A_1 \cap A_2$ and $y \in B_1 \cap B_2$.
Also, $(x, y) \in R_1 \setminus R_2$ iff $x \in A_1$, $y \in B_1$, and either $x \notin A_2$ or $y \notin B_2$.
Splitting by $x \in A_2$ gives the disjoint union in (i).

(ii) If $E = \bigcup_j R_j$ and $F = \bigcup_k S_k$, then $E \cap F = \bigcup_{j,k} (R_j \cap S_k) \in \mathcal{Z}_0$ by (i).
Repeatedly applying (i) gives $R_j \setminus F \in \mathcal{Z}_0$, hence $E \setminus F = \bigcup_j (R_j \setminus F) \in \mathcal{Z}_0$.
Since $Z = X \times Y \in \mathcal{Z}_0$, complements are differences from $Z$.

(iii) If $E = \bigcup_{j=1}^{n} R_j$, set $S_j := R_j \setminus \bigcup_{i<j} R_i$. Then $S_j \in \mathcal{Z}_0$ by (ii), the $S_j$ are disjoint, and $E = \bigsqcup_j S_j$.
Writing each $S_j$ as a finite union of rectangles and relabeling gives required finite disjoint representation.

*Structural Commentary*: Product rectangles behave like elementary sets for the product space.
*Key Insight*: Before extending to a $\sigma$-algebra, make all finite overlaps into disjoint rectangle pieces. □

**Theorem 2.1.3.** *Product Measure Theorem*
*For measure spaces $(X, \mathcal{M}, \mu)$ and $(Y, \mathcal{N}, \nu)$, $\exists \pi : \mathcal{M} \otimes \mathcal{N} \to [0, +\infty]$ a measure such that $\forall (A, B) \in \mathcal{M} \times \mathcal{N}$, $\pi(A \times B) = \mu(A)\nu(B)$, where $0 \cdot (+\infty) = 0$. If $\mu, \nu$ are $\sigma$-finite, $\pi$ is unique; write $\pi = \mu \otimes \nu$.*

*Proof.* For $E = \bigsqcup_{j=1}^{n} (A_j \times B_j) \in \mathcal{Z}_0$, set $\pi_0(E) := \sum_{j=1}^{n} \mu(A_j)\nu(B_j)$.
If $A \times B = \bigsqcup_{j \geq 1} (A_j \times B_j)$, then $\mathbf{1}_{A \times B} = \sum_{j \geq 1} \mathbf{1}_{A_j \times B_j}$, so monotone convergence in $y$, then $x$,
gives $\mu(A)\nu(B) = \sum_{j \geq 1} \mu(A_j)\nu(B_j)$. Refining two finite disjoint rectangle decompositions by intersections gives equal $\pi_0$-values; hence $\pi_0$ is well defined. If $E = \bigsqcup_k E_k \in \mathcal{Z}_0$ with $E_k \in \mathcal{Z}_0$, write $E = \bigsqcup_i R_i$ and $E_k = \bigsqcup_\ell R_{k\ell}$.
Since $R_i = \bigsqcup_{k,\ell} (R_i \cap R_{k\ell})$, the rectangle comparison yields $\pi_0(E) = \sum_k \pi_0(E_k)$; hence $\pi_0$ is a premeasure.
Extend $\pi_0$ to $\sigma(\mathcal{Z}_0) = \mathcal{M} \otimes \mathcal{N}$. If $\mu, \nu$ are $\sigma$-finite, then $\mathcal{Z}_0$ is $\pi_0$-$\sigma$-finite, so the extension is unique.
*Structural Commentary*: Rectangles fix the product measure.
*Key Insight*: Monotone convergence makes rectangle mass respect countable disjoint refinements. □

**Lemma 2.1.4.** *Measurable Sections*

*(i) $E \in \mathcal{M} \otimes \mathcal{N}$ implies $\forall x \in X$, $E_x \in \mathcal{N}$ and $\forall y \in Y$, $E^y \in \mathcal{M}$.*

*(ii) If $F : Z \to \overline{\mathbb{R}}$ is $\mathcal{M} \otimes \mathcal{N}$-measurable, then $F_x$ is $\mathcal{N}$-measurable $\forall x \in X$ and $F^y$ is $\mathcal{M}$-measurable $\forall y \in Y$.*

*Proof.*

(i) Fix $x \in X$ and set $\mathcal{C}_x := \{E \subseteq Z : E_x \in \mathcal{N}\}$. Since $(E^c)_x = (E_x)^c$ and $(\bigcup_n E_n)_x = \bigcup_n (E_n)_x$, $\mathcal{C}_x$ is a $\sigma$-algebra. Also $(A \times B)_x = B$ if $x \in A$ and $\emptyset$ otherwise; hence $\mathcal{M} \otimes \mathcal{N} \subseteq \mathcal{C}_x$. The $E^y$ proof is identical.

(ii) For $a \in \mathbb{R}$, set $H_a := \{(u, v) \in Z : F(u, v) > a\} \in \mathcal{M} \otimes \mathcal{N}$. By (i), $(H_a)_x \in \mathcal{N}$ and $(H_a)^y \in \mathcal{M}$.
Since $(H_a)_x = \{y : F_x(y) > a\}$ and $(H_a)^y = \{x : F^y(x) > a\}$, all threshold sets are measurable.

*Structural Commentary*: Product measurability is stable under freezing one coordinate.
*Key Insight*: Section operations preserve the set algebra used to generate the product $\sigma$-algebra. □

**Proposition 2.1.5.** *Section Measures Integrate to Product Measure*
*Let $(X, \mathcal{M}, \mu)$ and $(Y, \mathcal{N}, \nu)$ be $\sigma$-finite, $\pi := \mu \otimes \nu$, and $E \in \mathcal{M} \otimes \mathcal{N}$. Set $f_E(x) := \nu(E_x)$ and $g_E(y) := \mu(E^y)$. Then $f_E, g_E$ are measurable and $\int_X f_E \, d\mu = \pi(E) = \int_Y g_E \, d\nu$.*

*Proof.* Assume $\mu(X), \nu(Y) < +\infty$ and let $\mathcal{C}$ be the class of $E$ satisfying the claim.
For $E = A \times B$, $f_E = \nu(B)\mathbf{1}_A$ and $g_E = \mu(A)\mathbf{1}_B$, so $A \times B \in \mathcal{C}$; finite additivity gives $\mathcal{Z}_0 \subseteq \mathcal{C}$.
If $E_n \uparrow E$ in $\mathcal{C}$, then $(E_n)_x \uparrow E_x$ and $(E_n)^y \uparrow E^y$; MCT gives $E \in \mathcal{C}$.
If $E_n \downarrow E$, finite total mass gives downward continuity and DCT, hence $E \in \mathcal{C}$.
Thus $\mathcal{C}$ is a monotone class containing $\mathcal{Z}_0$, so $\mathcal{C} = \mathcal{M} \otimes \mathcal{N}$. For $\sigma$-finite $\mu, \nu$, choose $X_n \uparrow X$, $Y_n \uparrow Y$ with $\mu(X_n), \nu(Y_n) < +\infty$, apply the finite case to $E_n := E \cap (X_n \times Y_n)$, then let $n \to \infty$ by MCT.
*Structural Commentary*: Cavalieri's principle is already true for abstract $\sigma$-finite products.
*Key Insight*: Prove rectangles, close under monotone limits, then exhaust by finite-measure rectangles. □

**Perspective 2.1.6.** *Rectangle Premeasure and the Completed Product*
Let $\mathcal{R} := \{A \times B : A \in \mathcal{M},\ B \in \mathcal{N}\}$ and define $\lambda(A \times B) := \mu(A)\nu(B)$ on $\mathcal{R}$.

(i) Lemma 2.1.2 and Theorem 2.1.3 imply that $\lambda$ is a semiring premeasure.
(ii) Theorem 3.4.7 extends $\lambda$ to a complete measure $\overline{\lambda}$ on the Caratheodory $\sigma$-algebra $\mathcal{C}_\lambda$.
(iii) The restriction $\overline{\lambda}|_{\mathcal{M}\otimes\mathcal{N}}$ is $\mu \otimes \nu$; the full space $(X \times Y, \mathcal{C}_\lambda, \overline{\lambda})$ is the completed product.

Thus $\mathcal{C}_\lambda$ may contain $N' \subseteq N$ with $N \in \mathcal{M} \otimes \mathcal{N}$, $(\mu \otimes \nu)(N) = 0$, and $N' \notin \mathcal{M} \otimes \mathcal{N}$.
This is why some section-measurability forms of Fubini require completeness.
*Key Insight*: Rectangles set mass; completion adds null subsets.

## 2.2 Fubini and Tonelli Theorems

**Theorem 2.2.1.** *Tonelli's Theorem for $\sigma$-Finite Products*
*Let $(X, \mathcal{M}, \mu)$, $(Y, \mathcal{N}, \nu)$ be $\sigma$-finite, Let $Z := X \times Y$, $\pi := \mu \otimes \nu$.*
*Let $F : Z \to [0, +\infty]$ be $\mathcal{M} \otimes \mathcal{N}$-measurable. Put $I_F(x) := \int_Y F_x \, d\nu$ and $J_F(y) := \int_X F^y \, d\mu$.*

*(i) $I_F$ is $\mathcal{M}$-measurable.*
*(ii) $J_F$ is $\mathcal{N}$-measurable.*
*(iii) $\int_X I_F \, d\mu = \int_Z F \, d\pi = \int_Y J_F \, d\nu$.*

*Proof.* For $F = \mathbf{1}_E$, the claims follow from Proposition 2.1.5.
Finite linearity gives them for nonnegative simple $\Phi = \sum_{j=1}^{n} a_j \mathbf{1}_{E_j}$. Choose simple $0 \le \Phi_n \uparrow F$.

(i) Since $I_{\Phi_n}$ is $\mathcal{M}$-measurable and $I_{\Phi_n}(x) = \int_Y (\Phi_n)_x \, d\nu \uparrow \int_Y F_x \, d\nu = I_F(x)$, $I_F$ is $\mathcal{M}$-measurable.
(ii) Since $J_{\Phi_n}$ is $\mathcal{N}$-measurable and $J_{\Phi_n}(y) = \int_X (\Phi_n)^y \, d\mu \uparrow \int_X F^y \, d\mu = J_F(y)$, $J_F$ is $\mathcal{N}$-measurable.
(iii) From $\int_X I_{\Phi_n} \, d\mu = \int_Z \Phi_n \, d\pi = \int_Y J_{\Phi_n} \, d\nu$, MCT in $X$, $Z$, and $Y$ gives $\int_X I_F \, d\mu = \int_Z F \, d\pi = \int_Y J_F \, d\nu$.

*Structural Commentary*: Approximate by nonnegative simple functions.
*Key Insight*: MCT passes sections and integrals together. □

**Theorem 2.2.2.** *Fubini's Theorem for $\sigma$-Finite Products*
*Let $(X, \mathcal{M}, \mu)$ and $(Y, \mathcal{N}, \nu)$ be $\sigma$-finite, $\pi := \mu \otimes \nu$, and $F \in \mathscr{L}^1(\pi)$.*

*(i) $F_x \in \mathscr{L}^1(\nu)$ for $\mu$-a.e. $x$ and $F^y \in \mathscr{L}^1(\mu)$ for $\nu$-a.e. $y$.*
*(ii) With $f(x) := \int_Y F_x \, d\nu$ and $g(y) := \int_X F^y \, d\mu$ on those full-measure sets, and 0 elsewhere, $f \in \mathscr{L}^1(\mu)$, $g \in \mathscr{L}^1(\nu)$, and $\int_X f \, d\mu = \int_Z F \, d\pi = \int_Y g \, d\nu$.*

*Proof.*

(i) Tonelli on $|F|$ gives $\int_X (\int_Y |F_x| \, d\nu) d\mu = \int_Z |F| \, d\pi = \int_Y (\int_X |F^y| \, d\mu) d\nu < +\infty$, hence finite a.e.
(ii) Tonelli on $F^+$ and $F^-$ gives finite identities; subtract them to obtain $\int_X f \, d\mu = \int_Z F \, d\pi = \int_Y g \, d\nu$.
Moreover $|f| \le \int_Y |F_x| \, d\nu$ and $|g| \le \int_X |F^y| \, d\mu$, so (i) gives $f, g \in \mathscr{L}^1$.

*Structural Commentary*: Fubini is Tonelli plus subtraction.
*Key Insight*: $\int |F| < +\infty$ makes the two nonnegative Tonelli identities finite enough to subtract. □

**Proposition 2.2.3.** *Product Borel and Measurable Building Blocks*
*The following hold.*

*(i) $\mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R}) = \mathcal{B}(\mathbb{R}^2)$.*
*(ii) If $f : X \to \mathbb{R}$ is $\mathcal{M}$-measurable and $g : Y \to \mathbb{R}$ is $\mathcal{N}$-measurable, then $(x, y) \mapsto f(x)g(y)$ is $\mathcal{M} \otimes \mathcal{N}$-measurable.*
*(iii) If $u : \mathbb{R} \to \mathbb{R}$ is Borel measurable, then $(x, y) \mapsto u(x - y)$ is $\mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R})$-measurable.*

*Proof.*

(i) Open rectangles give $\mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R}) \subseteq \mathcal{B}(\mathbb{R}^2)$.
Conversely, every open $O \subseteq \mathbb{R}^2$ is a countable union of rational rectangles, each in $\mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R})$.

(ii) For $\tilde{f}(x,y) := f(x)$ and $\tilde{g}(x,y) := g(y)$, $\{\tilde{f} > a\} = f^{-1}((a,+\infty)) \times Y$ and $\{\tilde{g} > a\} = X \times g^{-1}((a,+\infty))$. Thus $\tilde{f}, \tilde{g}$ are measurable, so $\tilde{f}\tilde{g}$ is measurable.

(iii) The map $s(x,y) := x - y$ is continuous on $\mathbb{R}^2$, hence measurable by (i); therefore $u \circ s$ is measurable.

*Structural Commentary*: Product measurability supports two-variable formulas.
*Key Insight*: Check coordinate pullbacks on rectangles, then use ordinary measurable-function closure. □

**Proposition 2.2.4.** *Counting Factor Products*
*Let $X = \mathbb{N}$, $\mathcal{M} = 2^{\mathbb{N}}$, $\mu$ be counting measure, and $(Y, \mathcal{N}, \nu)$ be a measure space.*

*(i) For $E \subseteq \mathbb{N} \times Y$, $E \in 2^{\mathbb{N}} \otimes \mathcal{N} \iff \forall n,\ E_n \in \mathcal{N}$, and then $E = \bigsqcup_{n\geq 1} (\{n\} \times E_n)$.*

*(ii) The unique product measure is $\pi = \mu \otimes \nu$ with $\pi(E) = \sum_{n\geq 1} \nu(E_n)$ for $E \in 2^{\mathbb{N}} \otimes \mathcal{N}$.*

*(iii) For $F : \mathbb{N} \times Y \to \mathbb{R}$, $F$ is product-measurable $\iff \forall n$, $F_n$ is $\mathcal{N}$-measurable.*

*(iv) For product-measurable $F$, $F \in \mathscr{L}^1(\pi) \iff \sum_{n\geq 1} \int_Y |F_n|\, d\nu < +\infty$;*

*then $\int_{\mathbb{N}\times Y} F\, d\pi = \sum_{n\geq 1} \int_Y F_n\, d\nu = \int_Y \sum_{n\geq 1} F_n\, d\nu$, with $\sum_n F_n$ absolutely convergent $\nu$-a.e.*

*Proof.*

(i) ($\Rightarrow$) If $E \in 2^{\mathbb{N}} \otimes \mathcal{N}$, then $\forall n,\ E_n \in \mathcal{N}$ by Lemma 2.1.4.

(i) ($\Leftarrow$) If $\forall n,\ E_n \in \mathcal{N}$, then each $\{n\} \times E_n \in 2^{\mathbb{N}} \otimes \mathcal{N}$ and $E = \bigsqcup_{n\geq 1} (\{n\} \times E_n)$, hence $E \in 2^{\mathbb{N}} \otimes \mathcal{N}$.

(ii) Define $\pi(E) := \sum_{n\geq 1} \nu(E_n)$.

If $(E^k)_k$ are disjoint, then $(\bigsqcup_k E^k)_n = \bigsqcup_k E^k_n$, so $\pi(\bigsqcup_k E^k) = \sum_n \sum_k \nu(E^k_n) = \sum_k \pi(E^k)$.

Also $\pi(\{n\} \times A) = \nu(A)$, and (i) forces uniqueness.

(iii) ($\Rightarrow$) If $F$ is product-measurable, then $\forall n$, $F_n$ is measurable by Lemma 2.1.4.

(iii) ($\Leftarrow$) If $\forall n$, $F_n$ is measurable, then $\forall a \in \mathbb{R}$, $H_a := \{F > a\}$ satisfies $(H_a)_n = \{F_n > a\} \in \mathcal{N}$,
By (i), $H_a \in 2^{\mathbb{N}} \otimes \mathcal{N}$, so $F$ is product-measurable.

(iv) For $G \geq 0$, (ii) gives the identity for $G = \mathbf{1}_E$; linearity and MCT give $\int_{\mathbb{N}\times Y} G\, d\pi = \sum_{n\geq 1} \int_Y G_n\, d\nu$.

(iv) Apply the preceding identity to $G = |F|$: $F \in \mathscr{L}^1(\pi) \iff \sum_{n\geq 1} \int_Y |F_n|\, d\nu < +\infty$.

If finite, then $\int_Y \sum_n |F_n|\, d\nu < +\infty$, so $\sum_n F_n$ converges absolutely a.e..

Applying the nonnegative identity to $F^+$ and $F^-$ and subtracting gives the integral formulas.

*Structural Commentary*: A product with counting measure is a measurable sequence of fibers.
*Key Insight*: Counting atoms turn products into sums. □

**Example 2.2.5.** *A Diagonal Failure Without $\sigma$-Finiteness*
Let $X = Y = [0,1]$, $\mathcal{M} = \mathcal{N} = \mathcal{B}([0,1])$, $\mu = m$, $\nu = \#$, and $D := \{(x,y) \in [0,1]^2 : x = y\}$.
Then $D \in \mathcal{M} \otimes \mathcal{N}$, $D_x = \{x\}$, and $D^y = \{y\}$, so $\nu(D_x) = 1$ and $\mu(D^y) = 0$.
Hence $\int_X \nu(D_x)\, d\mu = 1$ but $\int_Y \mu(D^y)\, d\nu = 0$, so Tonelli–Fubini can fail without $\sigma$-finite factors.




Figure 207: Diagonal sections: $\nu(D_x) = 1$ but $\mu(D^y) = 0$.

**Example 2.2.6.** *Nonunique Products Without $\sigma$-Finiteness*
Let $X = \mathbb{R}$, $\mathcal{M} = 2^{\mathbb{R}}$, and $\mu(A) = 0$ if $A$ is countable, $\mu(A) = +\infty$ otherwise.
Define $\lambda_i : \mathcal{M} \otimes \mathcal{M} \to \{0, +\infty\}$ by $\lambda_i(E) = 0$ exactly on the following null ideals:
$\mathcal{I}_1 = \{E : E = G \cup H,\ \mathrm{pr}_x\, G \text{ countable},\ \mathrm{pr}_y\, H \text{ countable}\}$
and $\mathcal{I}_2 = \{E : E = G \cup H \cup K,\ \mathrm{pr}_x\, G \text{ countable},\ \mathrm{pr}_y\, H \text{ countable},\ \mathrm{pr}_{y=x}\, K \text{ countable}\}$.
Both ideals are closed under countable unions, hence $\lambda_i$ are measures.
For a rectangle $A \times B$, both measures equal $\mu(A)\mu(B)$: countable sides are null, and uncountable-by-uncountable rectangles are not in either ideal.

For $E = \{(x,y) : x + y = 0\}$, $\mathrm{pr}_{y=x} E$ is a point, so $\lambda_2(E) = 0$ but $\lambda_1(E) = +\infty$.
*Structural Commentary*: Without $\sigma$-finiteness, rectangle values need not determine a product measure.
*Key Insight*: Extra null lines preserve rectangles but change non-rectangles.

**Example 2.2.7.** *Double Series as Product Integrals*
With counting measure on $\mathbb{N}^2$, Tonelli gives, for $a_{mn} \geq 0$,
$\sum_m \sum_n a_{mn} = \sum_{(m,n)\in\mathbb{N}^2} a_{mn} = \sum_n \sum_m a_{mn} \in [0, +\infty]$.
Fubini needs integrability: set $a_{nn} = 1$, $a_{n,n+1} = -1$, and $a_{mn} = 0$ otherwise.
Then $\sum_m \sum_n a_{mn} = 0$, while $\sum_n \sum_m a_{mn} = 1$, and $\sum_{m,n} |a_{mn}| = +\infty$.

**Proposition 2.2.8.** *Separated Products*
*Let $(X, \mathcal{M}, \mu)$ and $(Y, \mathcal{N}, \nu)$ be $\sigma$-finite. Let $f \in \mathscr{L}^1(\mu)$, $g \in \mathscr{L}^1(\nu)$, and $H(x,y) := f(x)g(y)$.*

*(i)* $H \in \mathscr{L}^1(\mu \otimes \nu)$.
*(ii)* $\int_{X\times Y} H \, d(\mu \otimes \nu) = (\int_X f \, d\mu)(\int_Y g \, d\nu)$.

*Proof.*

(i) Proposition 2.2.3 gives measurability. Tonelli gives $\int_{X\times Y} |H| \, d(\mu \otimes \nu) = (\int_X |f| \, d\mu)(\int_Y |g| \, d\nu) < +\infty$.
(ii) Since $H \in \mathscr{L}^1(\mu \otimes \nu)$, Fubini gives $\int_{X\times Y} H \, d(\mu \otimes \nu) = (\int_X f \, d\mu)(\int_Y g \, d\nu)$.

*Structural Commentary*: Tonelli handles size; Fubini handles sign.
*Key Insight*: Separated products integrate as products of integrals. □

**Corollary 2.2.9.** *Equal Section Measures*
*Let $(X, \mathcal{M}, \mu)$ and $(Y, \mathcal{N}, \nu)$ be $\sigma$-finite.*
*If $E, F \in \mathcal{M} \otimes \mathcal{N}$ and $\forall x \in X$, $\nu(E_x) = \nu(F_x)$, then $(\mu \otimes \nu)(E) = (\mu \otimes \nu)(F)$.*

*Proof.* Set $f_E(x) := \nu(E_x)$ and $f_F(x) := \nu(F_x)$.
Since $f_E = f_F$, $(\mu \otimes \nu)(E) = \int_X f_E \, d\mu = \int_X f_F \, d\mu = (\mu \otimes \nu)(F)$ by Proposition 2.1.5.
*Structural Commentary*: Product mass is recovered from vertical section masses.
*Key Insight*: Equal fibers give equal product mass. □

**Definition 2.2.10.** *Convolution*
For Borel $f, g : \mathbb{R} \to \mathbb{R}$, if $y \mapsto f(x-y)g(y)$ is integrable for a.e. $x$,
define the *convolution* $(f * g)(x) := \int_{\mathbb{R}} f(x-y)g(y) \, dm(y)$ on that full-measure set and 0 elsewhere.

**Proposition 2.2.11.** *Convolution of $L^1$ Functions*
*Let $f, g \in \mathscr{L}^1(m)$ be Borel.*

*(i)* $f * g$ *is defined for a.e.* $x \in \mathbb{R}$.
*(ii)* $f * g \in \mathscr{L}^1(m)$.
*(iii)* $\int_{\mathbb{R}} |f * g| \, dm \leq (\int_{\mathbb{R}} |f| \, dm)(\int_{\mathbb{R}} |g| \, dm)$.

*Proof.* Set $\Phi(x,y) := f(x-y)g(y)$. By Proposition 2.2.3, $\Phi$ is product-measurable.
For fixed $y$, translation invariance gives $\int_{\mathbb{R}} |f(x-y)| \, dm(x) = \int_{\mathbb{R}} |f| \, dm$.
Hence Tonelli yields $\int_{\mathbb{R}^2} |\Phi| \, d(m \otimes m) = (\int_{\mathbb{R}} |f| \, dm)(\int_{\mathbb{R}} |g| \, dm) < +\infty$.

(i) Fubini gives $y \mapsto \Phi(x,y) \in \mathscr{L}^1(m)$ for a.e. $x$, so $f * g$ is a.e. defined.
(ii) Since $|f * g(x)| \leq \int_{\mathbb{R}} |\Phi(x,y)| \, dm(y)$ a.e., Fubini gives $\int_{\mathbb{R}} |f * g| \, dm < +\infty$.
(iii) The same estimate gives $\int_{\mathbb{R}} |f * g| \, dm \leq \int_{\mathbb{R}^2} |\Phi| \, d(m \otimes m) = (\int_{\mathbb{R}} |f| \, dm)(\int_{\mathbb{R}} |g| \, dm)$.

*Structural Commentary*: Apply Fubini to $\Phi(x,y) = f(x-y)g(y)$.
*Key Insight*: Translation invariance fixes the $x$-slice norm. □

**Definition 2.2.12.** *Products and Slices*
For $A \subseteq \mathbb{R}^n$, $B \subseteq \mathbb{R}^k$, set $A \times B \subseteq \mathbb{R}^{n+k}$.
For $E \subseteq \mathbb{R}^n \times \mathbb{R}^k$ define $E_x := \{y \in \mathbb{R}^k : (x,y) \in E\}$ as *$x$-slice*, and $E^y := \{x \in \mathbb{R}^n : (x,y) \in E\}$ as *$y$-slice*.
For $f : E \to \mathbb{R}$, write $f_x(y) = f^y(x) := f(x,y)$ where defined.

**Theorem 2.2.13.** *Measurable Products*
*Let $A \in \mathcal{L}^n$ and $B \in \mathcal{L}^k$.*

*(i)* $A \times B \in \mathcal{L}^{n+k}$.
*(ii)* $m_{n+k}(A \times B) = m_n(A) m_k(B)$, *with* $0 \cdot \infty = \infty \cdot 0 = 0$.

*Proof.* Boxes satisfy $m_{n+k}(I \times J) = m_n(I)m_k(J)$, hence so do open sets by disjoint box decompositions.
For bounded finite-measure $A, B$, choose $F_A \subseteq A \subseteq G_A$, $F_B \subseteq B \subseteq G_B$ with $F_A, F_B$ closed, $G_A, G_B$ bounded open, and $m_n(G_A \setminus F_A), m_k(G_B \setminus F_B) < \varepsilon$.
Then $F_A \times F_B \subseteq A \times B \subseteq G_A \times G_B$ and $m_n(G_A)m_k(G_B) - m_n(F_A)m_k(F_B) \to 0$.

(i) The squeeze proves $A \times B \in \mathcal{L}^{n+k}$ in the bounded finite-measure case.
For general $A, B$, with $A_N := A \cap [-N, N]^n$ and $B_N := B \cap [-N, N]^k$, $A \times B = \bigcup_N (A_N \times B_N) \in \mathcal{L}^{n+k}$.

(ii) The squeeze also gives $m_{n+k}(A \times B) = m_n(A)m_k(B)$ in the bounded finite-measure case.
Since $A_N \times B_N \uparrow A \times B$, $m_{n+k}(A \times B) = \lim_N m_n(A_N)m_k(B_N) = m_n(A)m_k(B)$, with $0 \cdot \infty = \infty \cdot 0 = 0$.

*Structural Commentary*: Product measure comes from boxes and regularity.
*Key Insight*: Regularity turns rough measurable products into box products plus zero-set errors. □

**Theorem 2.2.14.** *Zero Slices*
*If $E \subseteq \mathbb{R}^n \times \mathbb{R}^k$ is Lebesgue measurable, then $E$ is null iff $E_x$ is null for a.e. $x$.*

*Proof.*

(⇒) If $m(E) = 0$, cover $E$ by an open $W$ with small measure.
For $X_\alpha(W) = \{x : m(W_x) > \alpha\}$, $m(W) \geq \alpha m(X_\alpha(W))$; letting $m(W) \to 0$ gives $m(E_x) = 0$ for a.e. $x$.

(⇐) Remove the exceptional null base so every slice is null. For compact $K \subseteq E$, cover each $K_x$ by a small-measure slice neighborhood; compactness gives finitely many product neighborhoods covering $K$, whose total measure is arbitrarily small by Theorem 2.2.13. Thus $m(K) = 0$, hence $m(E) = 0$ by regularity.

*Structural Commentary*: Nullity can be checked fiberwise after removing an exceptional base set.
*Key Insight*: Product measure turns small slice covers into small total measure. □




Figure 208: A planar set can be read by vertical slices; Fubini integrates the slice data over the base variable.

**Theorem 2.2.15.** *Cavalieri's Principle for Lebesgue Measure*
*Let $n, k \in \mathbb{N}$ and $E \in \mathcal{L}^{n+k}$. For $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^k$, set $E_x := \{y : (x, y) \in E\}$ and $E^y := \{x : (x, y) \in E\}$.*

*(i) $E_x \in \mathcal{L}^k$ for a.e. $x$, and $f_E(x) := m_k(E_x)$ has a measurable representative with $m_{n+k}(E) = \int_{\mathbb{R}^n} f_E \, dm_n$.*
*(ii) $E^y \in \mathcal{L}^n$ for a.e. $y$, and $g_E(y) := m_n(E^y)$ has a measurable representative with $m_{n+k}(E) = \int_{\mathbb{R}^k} g_E \, dm_k$.*

*Proof.* Let $\mathsf{C}_x(E)$ denote: $E_x \in \mathcal{L}^k$ for a.e. $x$, $f_E$ has a measurable representative, and $m_{n+k}(E) = \int_{\mathbb{R}^n} f_E \, dm_n$.

(i) Rectangles satisfy $\mathsf{C}_x(A \times B)$ since $f_{A \times B} = m_k(B)\mathbf{1}_A$; finite additivity gives finite unions.
For bounded $Q = I \times J$, $\mathcal{C}_Q := \{E \subseteq Q : \mathsf{C}_x(E)\}$ is monotone:
$E_j \uparrow E$ gives $f_{E_j} \uparrow f_E$ and MCT, while $E_j \downarrow E$ gives $f_{E_j} \downarrow f_E \leq m_k(J)\mathbf{1}_I$ and DCT.
Thus $\mathcal{C}_Q$ contains all Borel $E \subseteq Q$, and exhaustion by $Q_N = [-N, N]^{n+k}$ gives all Borel $E$.
For $E \in \mathcal{L}^{n+k}$, choose Borel $B$ with $m_{n+k}(E \triangle B) = 0$; then $m_k((E \triangle B)_x) = 0$ a.e. by Theorem 2.2.14.
Hence $E_x \in \mathcal{L}^k$ a.e., $f_E = f_B$ a.e., and $m_{n+k}(E) = \int_{\mathbb{R}^n} f_E \, dm_n$.

(ii) Let $\tau : \mathbb{R}^k \times \mathbb{R}^n \to \mathbb{R}^n \times \mathbb{R}^k$ be $\tau(y, x) = (x, y)$ and put $\widetilde{E} := \tau^{-1}(E)$.
Apply (i) to $\widetilde{E} \subseteq \mathbb{R}^k \times \mathbb{R}^n$ gives $\widetilde{E}_y = E^y$ a.e., $g_E(y) = m_n(\widetilde{E}_y)$, $m_{n+k}(E) = m_{n+k}(\widetilde{E}) = \int_{\mathbb{R}^k} g_E \, dm_k$.

*Structural Commentary*: Slices reduce product measure to base integrals.
*Key Insight*: Extend from rectangles, then pass null differences. □

**Perspective 2.2.16.** *Fubini Proof Ladder*
Fubini ladder: $\mathbf{1}_E \to$ simple $\Phi \geq 0 \to f \geq 0$ by MCT, then $f = f^+ - f^-$ when $\int |f| < +\infty$.
Theorems 2.2.1 and 2.2.2 are the abstract form.
*Key Insight*: Fubini is a slice theorem, proof runs through indicators, simple functions, monotone convergence.

**Proposition 2.2.17.** *Undergraphs and Level-Set Measurability*
*Let $f : \mathbb{R}^n \to [0,\infty]$, set $F_a := \{x : f(x) > a\}$, and let $\mathcal{U}f := \{(x,t) \in \mathbb{R}^n \times [0,\infty) : 0 < t < f(x)\}$.*

*(i) $f$ is Lebesgue measurable iff $\forall q \in \mathbb{Q}_{>0},\ F_q \in \mathcal{L}^n$.*
*(ii) If $f$ is Lebesgue measurable, then $\mathcal{U}f \in \mathcal{L}^{n+1}$ and $\forall t > 0,\ (\mathcal{U}f)^t = F_t$.*

*Proof.*

(i) ($\Rightarrow$) If $f$ is measurable, each $F_q = f^{-1}((q,+\infty])$ is in $\mathcal{L}^n$.

(i) ($\Leftarrow$) If $\forall q \in \mathbb{Q}_{>0},\ F_q \in \mathcal{L}^n$, then for $a \geq 0$, $F_a = \bigcup_{q\in\mathbb{Q},\ q>a} F_q \in \mathcal{L}^n$, while for $a < 0$, $F_a = \mathbb{R}^n$.
Thus all strict superlevel sets are measurable, so $f$ is measurable.

(ii) By (i), every $F_r$ with $r \in \mathbb{Q}_{>0}$ is measurable. Since $\mathcal{U}f = \bigcup_{q,r\in\mathbb{Q},\ 0<q<r} F_r \times (q,r)$, $\mathcal{U}f \in \mathcal{L}^{n+1}$.
For $t > 0$, $(\mathcal{U}f)^t = \{x : 0 < t < f(x)\} = F_t$.

*Structural Commentary*: Measurability is encoded by rational superlevels.
*Key Insight*: Undergraph slices are superlevel sets. □

**Theorem 2.2.18.** *Fubini–Tonelli for Nonnegative Functions*
*Let $f : \mathbb{R}^n \times \mathbb{R}^k \to [0,\infty]$ be Lebesgue measurable.*
*Define $F(x) := \int_{\mathbb{R}^k} f_x\, dm_k$, $f_x$ is measurable and arbitrarily elsewhere; define $G(y) := \int_{\mathbb{R}^n} f^y\, dm_n$ analogously.*

*(i) $f_x$ is measurable for a.e. $x$, $F$ is measurable, and $\int_{\mathbb{R}^{n+k}} f\, dm_{n+k} = \int_{\mathbb{R}^n} F\, dm_n$.*
*(ii) $f^y$ is measurable for a.e. $y$, $G$ is measurable, and $\int_{\mathbb{R}^{n+k}} f\, dm_{n+k} = \int_{\mathbb{R}^k} G\, dm_k$.*

*Proof.* Set $U := \mathcal{U}f \subseteq \mathbb{R}^{n+k} \times [0,\infty)$. Then $U \in \mathcal{L}^{n+k+1}$ by Proposition 2.2.17.

(i) By Cavalieri, $U_x$ is measurable for a.e. $x$, $H(x) := m_{k+1}(U_x)$ is measurable, and $m_{n+k+1}(U) = \int_{\mathbb{R}^n} H\, dm_n$.
For such $x$, $U_x = \mathcal{U}(f_x)$, so $f_x$ is measurable and $H(x) = \int_{\mathbb{R}^k} f_x\, dm_k = F(x)$.
Since $m_{n+k+1}(U) = \int_{\mathbb{R}^{n+k}} f\, dm_{n+k}$, (i) follows.

(ii) Let $\tau(y,x) = (x,y)$ and $\tilde{f} := f \circ \tau$.
Applying (i) to $\tilde{f}$ gives $\tilde{f}_y = f^y$ a.e., $\tilde{F} = G$, and $\int_{\mathbb{R}^k} G\, dm_k = \int_{\mathbb{R}^{k+n}} \tilde{f}\, dm_{k+n} = \int_{\mathbb{R}^{n+k}} f\, dm_{n+k}$.

*Structural Commentary*: Cavalieri applied to the undergraph.
*Key Insight*: $(\mathcal{U}f)_x = \mathcal{U}(f_x)$. □

**Theorem 2.2.19.** *Fubini for $L^1$ Functions*
*Let $f \in L^1(\mathbb{R}^2)$.*

*(i) $f(x,\cdot) \in L^1(\mathbb{R})$ for a.e. $x$; defining $F(x) := \int_{\mathbb{R}} f(x,y)\, dy$ there and $F = 0$ elsewhere gives $F \in L^1(\mathbb{R})$.*
*(ii) $f(\cdot,y) \in L^1(\mathbb{R})$ for a.e. $y$; defining $G(y) := \int_{\mathbb{R}} f(x,y)\, dx$ there and $G = 0$ elsewhere gives $G \in L^1(\mathbb{R})$.*
*(iii) $\int_{\mathbb{R}} F\, dx = \int_{\mathbb{R}^2} f = \int_{\mathbb{R}} G\, dy$.*

*Proof.* From Theorems 2.2.13–2.2.14, the identity holds for indicators.
By linearity and MCT, every measurable $h \geq 0$ satisfies $(*)$ $\int_{\mathbb{R}}(\int_{\mathbb{R}} h(x,y)\, dy)dx = \int_{\mathbb{R}^2} h = \int_{\mathbb{R}}(\int_{\mathbb{R}} h(x,y)\, dx)dy$.

(i) Applying $(*)$ to $|f|$ gives $\int_{\mathbb{R}}\int_{\mathbb{R}} |f(x,y)|\, dy\, dx = \int_{\mathbb{R}^2} |f| < +\infty$, so $f(x,\cdot) \in L^1(\mathbb{R})$ a.e.
Let $F^{\pm}(x) := \int_{\mathbb{R}} f^{\pm}(x,y)\, dy$; then $F = F^+ - F^-$ a.e. and $\int_{\mathbb{R}} |F|\, dx \leq \int_{\mathbb{R}} (F^+ + F^-)\, dx = \int_{\mathbb{R}^2} |f| < +\infty$.

(ii) Let $\tau(y,x) = (x,y)$ and $\tilde{f} := f \circ \tau$. Since $\tau$ preserves $m_2$, $\tilde{f} \in L^1(\mathbb{R}^2)$.
Applying (i) to $\tilde{f}$ gives $\tilde{f}(y,\cdot) = f(\cdot,y) \in L^1(\mathbb{R})$ a.e. and $\tilde{F} = G \in L^1(\mathbb{R})$.

(iii) Let $G^{\pm}(y) := \int_{\mathbb{R}} f^{\pm}(x,y)\, dx$. By $(*)$, $\int_{\mathbb{R}} F^{\pm}\, dx = \int_{\mathbb{R}^2} f^{\pm} = \int_{\mathbb{R}} G^{\pm}\, dy$. Since these quantities are finite and $F = F^+ - F^-$, $G = G^+ - G^-$ a.e., subtraction gives $\int_{\mathbb{R}} F\, dx = \int_{\mathbb{R}^2} f = \int_{\mathbb{R}} G\, dy$.

*Structural Commentary*: Indicators, simple functions, MCT, then $f^{\pm}$.
*Key Insight*: Absolute integrability permits subtraction. □

## 2.3 Linear Changes of Variables and Euclidean Invariance

**Perspective 2.3.1.** *Coordinate Splitting as a Product Identification*
For $n = m + k$, let $\Phi : \mathbb{R}^n \to \mathbb{R}^m \times \mathbb{R}^k$ be $\Phi(x_1,\dots,x_n) = ((x_1,\dots,x_m),(x_{m+1},\dots,x_n))$. Then $\Phi$ sends rectangles to product rectangles and preserves rectangular volume, so $m_n$ is the completion of $\Phi_{\#}^{-1}(m_m \otimes m_k)$. Equivalently, integration over $\mathbb{R}^{m+k}$ matches iterated integration over $\mathbb{R}^m \times \mathbb{R}^k$ whenever Fubini applies.
*Key Insight*: Coordinate splitting preserves Lebesgue mass.

**Proposition 2.3.2.** *Lipschitz Images of Lebesgue Sets*
*Let $\Psi : \mathbb{R}^n \to \mathbb{R}^n$ be Lipschitz and let $E \in \mathcal{L}^n$. Then $\Psi(E) \in \mathcal{L}^n$.*
*Hence every linear map $\mathbb{R}^n \to \mathbb{R}^n$ preserves Lebesgue measurability.*

*Proof.* Let $L$ be a Lipschitz constant.

(i) $\forall I \subseteq \mathbb{R}^n$ bounded cells, $m_n^*(\Psi(I)) \le C_{n,L}\,\mathrm{vol}_n(I)$. Indeed, partition $I$ into grid cells of side at most $\delta$. Each image has diameter at most $L\sqrt{n}\,\delta$, hence lies in a cube of volume at most $C_{n,L}\delta^n$. Summing and letting $\delta \downarrow 0$ gives the estimate.

(ii) If $m_n^*(N) = 0$ and $\varepsilon > 0$, cover $N$ by cells $(I_k)$ with $\sum_k \mathrm{vol}_n(I_k) < \varepsilon$.

Then $\Psi(N) \subseteq \bigcup_k \Psi(I_k)$, so (i) gives $m_n^*(\Psi(N)) \le C_{n,L}\varepsilon$; hence $m_n^*(\Psi(N)) = 0$.

(iii) If $E$ is bounded, choose $F_\sigma$ set $F \subseteq E$ with $m_n^*(E \setminus F) = 0$. Write $F = \bigcup_j F_j$ with each $F_j$ compact.

Then $\Psi(F) = \bigcup_j \Psi(F_j)$ is Borel, $\Psi(E \setminus F)$ is null by (ii), and $\Psi(E) = \Psi(F) \cup \Psi(E \setminus F) \in \mathcal{L}^n$.

(iv) For general $E$, use $E = \bigcup_{r \ge 1} (E \cap [-r, r]^n)$ and apply (iii) to each bounded term.

*Structural Commentary*: Reduce to compact pieces plus null error.
*Key Insight*: Lipschitz maps preserve null sets. □

**Theorem 2.3.3.** *Linear Change of Variables for Nonnegative Functions*
*Let $T \in \mathcal{L}(\mathbb{R}^n)$ be invertible and let $f : \mathbb{R}^n \to [0, +\infty]$ be Lebesgue measurable.*

*(i) $f \circ T$ is Lebesgue measurable.*
*(ii) $\int f \circ T \, dm_n = |\det T|^{-1} \int f \, dm_n$.*

*Proof.*

(i) Since $(f \circ T)^{-1}((a, +\infty]) = T^{-1}(f^{-1}((a, +\infty]))$ and $T^{-1}$ is Lipschitz, Proposition 2.3.2 gives measurability.

(ii) The $n = 1$ identity follows from $m_1(\alpha E + \beta) = |\alpha|\, m_1(E)$.
Let $\mathcal{M}_+$ denote the measurable maps $\mathbb{R}^n \to [0, +\infty]$, set $J_S(h) := \int h(Sx)\, dm_n(x)$.
Set $\mathcal{A}_n := \{S \in GL(n) : J_S(h) = |\det S|^{-1} \int h \, dm_n \ \forall h \in \mathcal{M}_+\}$.
If $S_1, S_2 \in \mathcal{A}_n$, then $J_{S_1 S_2}(h) = |\det S_2|^{-1}\, J_{S_1}(h) = |\det(S_1 S_2)|^{-1} \int h \, dm_n$, so $S_1 S_2 \in \mathcal{A}_n$.
The generators $D_j(c) : x_j \mapsto c x_j$, $P_{j,j+1} : x_j \leftrightarrow x_{j+1}$, and $H_{jk}(c) : x_j \mapsto x_j + c x_k$ lie in $\mathcal{A}_n$ for $c \neq 0$ by the $n = 1$ identity, Fubini, and slice translation invariance, with determinant factors $|c|$, 1, and 1.
Row reduction gives $T \in \mathcal{A}_n$; hence $J_T(f) = |\det T|^{-1} \int f \, dm_n$.

*Structural Commentary*: Verify elementary row operations, then compose.
*Key Insight*: Only scalings change mass. □



Figure 209: Original and image cells under scale, swap, and shear; Fubini checks each coordinatewise.

**Corollary 2.3.4.** *Invertible Linear Images of Lebesgue Sets*
*Let $T \in \mathcal{L}(\mathbb{R}^n)$ be invertible and let $E \subseteq \mathbb{R}^n$.*

*(i) $E \in \mathcal{L}^n \iff T(E) \in \mathcal{L}^n$.*
*(ii) If $E \in \mathcal{L}^n$, then $m_n(T(E)) = |\det T|\, m_n(E)$.*

*Proof.*

(i) Since $T$ and $T^{-1}$ are Lipschitz, Proposition 2.3.2 gives $E \in \mathcal{L}^n \iff T(E) \in \mathcal{L}^n$.

(ii) For $E \in \mathcal{L}^n$, Theorem 2.3.3 applied to $\mathbf{1}_{T(E)}$ gives $m_n(E) = \int \mathbf{1}_{T(E)}(Tx)\, dm_n(x) = |\det T|^{-1}\, m_n(T(E))$.

*Structural Commentary*: Measurability is Lipschitz; mass follows from indicators.
*Key Insight*: Indicators turn integral change into set change. □

**Corollary 2.3.5.** *Linear Change of Variables for $L^1$ Functions*
*Let $T \in \mathcal{L}(\mathbb{R}^n)$ be invertible, let $f \in \mathscr{L}^1(m_n)$, and set $c_T := |\det T|^{-1}$.*

*(i) $f \circ T \in \mathscr{L}^1(m_n)$.*
*(ii) $\int f \circ T \, dm_n = c_T \int f \, dm_n$.*

*Proof.*

(i) Apply Theorem 2.3.3 to $|f|$: $\int |f \circ T| \, dm_n = c_T \int |f| \, dm_n < +\infty$, hence $f \circ T \in \mathscr{L}^1(m_n)$.

(ii) Apply Theorem 2.3.3 to $f^{\pm}$: $\int f^{\pm} \circ T \, dm_n = c_T \int f^{\pm} \, dm_n < +\infty$.
Hence $\int f \circ T \, dm_n = \int f^{+} \circ T \, dm_n - \int f^{-} \circ T \, dm_n = c_T \int f \, dm_n$.

*Structural Commentary*: Reduce signed integrals to finite positive parts.
*Key Insight*: Absolute integrability permits subtraction. □

**Corollary 2.3.6.** *Linear Images and Rigid Motions*
*Let $E \in \mathcal{L}^n$.*

*(i) If $T \in \mathcal{L}(\mathbb{R}^n)$ is singular, then $T(E)$ is Lebesgue measurable and $m_n(T(E)) = 0$.*
*(ii) If $\Psi(x) = Ux + x_0$, where $U$ is a linear isometry, then $\Psi(E) \in \mathcal{L}^n$ and $m_n(\Psi(E)) = m_n(E)$.*

*Proof.*

(i) $T(E) \in \mathcal{L}^n$ by Proposition 2.3.2. Set $W := \operatorname{range} T$, $H := \mathbb{R}^{n-1} \times \{0\}$; choose invertible $S$ with $S(W) \subseteq H$.
Since $H = \bigcup_N (H \cap [-N, N]^n)$ is null, $m_n(W) = 0$ by Corollary 2.3.4; hence $m_n(T(E)) = 0$.

(ii) Since $U$ is an isometry, $|\det U| = 1$ by Proposition 1.1.22.
Thus $U(E) \in \mathcal{L}^n$ and $m_n(U(E)) = m_n(E)$ by Corollary 2.3.4.
By translation invariance, $\Psi(E) = U(E) + x_0 \in \mathcal{L}^n$ and $m_n(\Psi(E)) = m_n(U(E)) = m_n(E)$.

*Structural Commentary*: Singular maps land in null subspaces; rigid motions preserve determinant mass.
*Key Insight*: Lebesgue measure records determinant size, not the coordinate description of the set. □

# 3 Radon-Nikodym and Differentiation

## 3.1 Signed and Absolutely Continuous Measures

This subsection first decomposes finite signed set functions into measures, then compares measures by absolute continuity and singularity. Later subsections develop the local Euclidean counterpart of the Radon–Nikodym viewpoint.

**Definition 3.1.1.** *Positive, Negative, and Null Sets for a Charge*
Let $\lambda$ be a charge on $(X, \mathcal{M})$.

(i) $\mathcal{P}_\lambda := \{P \in \mathcal{M} : \forall E \in \mathcal{M}, \lambda(E \cap P) \geq 0\}$, the *positive* sets.
(ii) $\mathcal{N}_\lambda := \{N \in \mathcal{M} : \forall E \in \mathcal{M}, \lambda(E \cap N) \leq 0\}$, the *negative* sets.
(iii) $\mathcal{Z}_\lambda := \{M \in \mathcal{M} : \forall E \in \mathcal{M}, \lambda(E \cap M) = 0\}$, the *null* sets.

**Proposition 3.1.2.** *Positive and Negative Set Closure*
*For a charge $\lambda$:*

*(i) $A \in \mathcal{M}$ and $A \subseteq P \in \mathcal{P}_\lambda$ imply $A \in \mathcal{P}_\lambda$.*
*(ii) $A \in \mathcal{M}$ and $A \subseteq N \in \mathcal{N}_\lambda$ imply $A \in \mathcal{N}_\lambda$.*
*(iii) $\mathcal{P}_\lambda$ is closed under countable unions.*
*(iv) $\mathcal{N}_\lambda$ is closed under countable unions.*

*Proof.*

(i) If $A \subseteq P \in \mathcal{P}_\lambda$, then $\lambda(E \cap A) = \lambda((E \cap A) \cap P) \geq 0 \ \forall E \in \mathcal{M}$.
(ii) If $A \subseteq N \in \mathcal{N}_\lambda$, then $\lambda(E \cap A) = \lambda((E \cap A) \cap N) \leq 0 \ \forall E \in \mathcal{M}$.
(iii) Let $(P_k) \subseteq \mathcal{P}_\lambda$ and set $P := \bigcup_{k \geq 1} P_k$. For $E \in \mathcal{M}$, set $F_1 := E \cap P_1$ and $F_k = E \cap P_k \setminus \bigcup_{j=1}^{k-1} P_j$ for $k \geq 2$.
Then $(F_k)$ is pairwise disjoint, $\bigcup_k F_k = E \cap P$, and $F_k \subseteq P_k$, so $\lambda(F_k) \geq 0$.
Countable additivity gives $\lambda(E \cap P) = \sum_k \lambda(F_k) \geq 0$.
(iv) Apply the disjointization in (iii) to $(N_k) \subseteq \mathcal{N}_\lambda$; every disjoint piece has nonpositive charge, so $\bigcup_k N_k \in \mathcal{N}_\lambda$.

*Structural Commentary*: Sign sets are hereditary and countable-union stable.
*Key Insight*: Test sets can always be disjointized against a countable sign cover. □

**Theorem 3.1.3.** *Hahn Decomposition Theorem*
*Let $\lambda$ be a charge on $(X, \mathcal{M})$. Then $\exists P, N \in \mathcal{M}$ such that*

*(i) $P \cap N = \emptyset$ and $X = P \sqcup N$.*
*(ii) $P \in \mathcal{P}_\lambda$.*
*(iii) $N \in \mathcal{N}_\lambda$.*

*Such a pair $(P, N)$ is a Hahn decomposition of $X$ for $\lambda$.*

*Proof.*

(i) Put $\alpha = \sup\{\lambda(A) : A \in \mathcal{P}_\lambda\}$. If $\alpha = +\infty$, choose $B_m \in \mathcal{P}_\lambda$, $\lambda(B_m) > m$, and set $C_m = \bigcup_{j=1}^{m} B_j$; then $C_m \in \mathcal{P}_\lambda$, $C_m \uparrow C$, and $\lambda(C_m) \geq \lambda(B_m) > m$, contradicting continuity from below and $\lambda(C) \in \mathbb{R}$.
Choose $A_n \in \mathcal{P}_\lambda$, $\lambda(A_n) \to \alpha$, replace $A_n$ by $\bigcup_{j=1}^{n} A_j$, and set $P = \bigcup_{n\geq 1} A_n$, $N = X \setminus P$.
Hence $P, N \in \mathcal{M}$, $P \cap N = \emptyset$, and $X = P \sqcup N$.

(ii) For $E \in \mathcal{M}$, $E \cap A_n \uparrow E \cap P$ and $\lambda(E \cap A_n) \geq 0$, so continuity from below gives $\lambda(E \cap P) \geq 0$.
Thus $P \in \mathcal{P}_\lambda$, and $\lambda(P) = \alpha$.

(iii) Suppose $E \in \mathcal{M}$, $E \subseteq N$, and $\lambda(E) > 0$. Then $E \notin \mathcal{P}_\lambda$, otherwise $P \cup E \in \mathcal{P}_\lambda$ and $\lambda(P \cup E) > \alpha$.
Set $R_0 := E$. Inductively, if $R_{k-1} \notin \mathcal{P}_\lambda$, let $n_k := \min\{n \geq 1 : \exists H \in \mathcal{M},\ H \subseteq R_{k-1},\ \lambda(H) \leq -\frac{1}{n}\}$.
Choose such $E_k$, and set $R_k := R_{k-1} \setminus E_k$. If some $R_m \in \mathcal{P}_\lambda$, then $\lambda(R_m) = \lambda(E) - \sum_{k\leq m} \lambda(E_k) > 0$, so $P \cup R_m \in \mathcal{P}_\lambda$ and $\lambda(P \cup R_m) > \alpha$, impossible. Hence all $R_k \notin \mathcal{P}_\lambda$. Put $F_m := \bigcup_{k\leq m} E_k$ and $F := \bigcup_{k\geq 1} E_k$.
Since $\lambda(F_m) = \sum_{k\leq m} \lambda(E_k) \leq -\sum_{k\leq m} \frac{1}{n_k}$ and $F_m \uparrow F$, by finiteness of $\lambda(F)$, $\sum_k \frac{1}{n_k} < +\infty$, hence $n_k \to +\infty$.
If $G \in \mathcal{M}$ and $G \subseteq E \setminus F$, then $G \subseteq R_{k-1}$ $\forall k$, by minimality $\lambda(G) > -\frac{1}{n_k - 1}$ $\forall k \gg 1$; hence $\lambda(G) \geq 0$ and $E \setminus F \in \mathcal{P}_\lambda$. Since $\lambda(E \setminus F) = \lambda(E) - \lambda(F) > 0$, $P \cup (E \setminus F) \in \mathcal{P}_\lambda$ and $\lambda(P \cup (E \setminus F)) > \alpha$, contradiction.

*Structural Commentary*: A charge splits $X$ into nonnegative-piece and nonpositive-piece regions.
*Key Insight*: Maximize positive mass; purify any leftover positive piece. □

**Lemma 3.1.4.** *Hahn Parts Are Unique up to Null Sets*
*Let $(P_i, N_i)$, $i = 1, 2$, be Hahn decompositions for a charge $\lambda$. $\forall E \in \mathcal{M}$:*

*(i)* $\lambda(E \cap P_1) = \lambda(E \cap P_2)$.
*(ii)* $\lambda(E \cap N_1) = \lambda(E \cap N_2)$.

*Proof.* Fix $E \in \mathcal{M}$. For $\{i, j\} = \{1, 2\}$,

(i) $E \cap (P_i \setminus P_j) \subseteq P_i \cap N_j$, so $\lambda(E \cap (P_i \setminus P_j)) = 0$. Hence $\lambda(E \cap P_i) = \lambda(E \cap P_1 \cap P_2)$ for $i = 1, 2$.
(ii) $E \cap (N_i \setminus N_j) \subseteq N_i \cap P_j$, so $\lambda(E \cap (N_i \setminus N_j)) = 0$. Hence $\lambda(E \cap N_i) = \lambda(E \cap N_1 \cap N_2)$ for $i = 1, 2$.

*Structural Commentary*: Hahn decompositions are not unique as sets, but the measures they define are unique.
*Key Insight*: Positive-and-negative overlap is necessarily null. □

**Definition 3.1.5.** *Jordan Variations of a Charge*
Let $\lambda$ be a charge, $(P, N)$ a Hahn decomposition, and $E \in \mathcal{M}$.

(i) $\lambda^+(E) := \lambda(E \cap P)$, the *positive variation*.
(ii) $\lambda^-(E) := -\lambda(E \cap N)$, the *negative variation*.
(iii) $|\lambda|(E) := \lambda^+(E) + \lambda^-(E)$, the *total variation*.

These are independent of $(P, N)$ by Lemma 3.1.4.

**Theorem 3.1.6.** *Jordan Decomposition Theorem*
*Let $\lambda$ be a charge.*

*(i)* $\lambda = \lambda^+ - \lambda^-$.
*(ii) If $\lambda = \mu - \nu$ for finite measures $\mu, \nu$, then $\lambda^+ \leq \mu$.*
*(iii) Under the same hypothesis, $\lambda^- \leq \nu$.*

*Proof.* Let $(P, N)$ be a Hahn decomposition and fix $E \in \mathcal{M}$.

(i) $\lambda(E) = \lambda(E \cap P) + \lambda(E \cap N) = \lambda^+(E) - \lambda^-(E)$.
(ii) $\lambda^+(E) = \mu(E \cap P) - \nu(E \cap P) \leq \mu(E)$.
(iii) $\lambda^-(E) = \nu(E \cap N) - \mu(E \cap N) \leq \nu(E)$.

*Structural Commentary*: Jordan decomposition is the minimal finite-measure difference.
*Key Insight*: Positive and negative Hahn pieces isolate the irreducible positive masses. □

**Corollary 3.1.7.** *Variation by Subsets and Null Sets*
*For a charge $\lambda$ and $E, M \in \mathcal{M}$:*

*(i)* $\lambda^+(E) = \sup\{\lambda(F) : F \in \mathcal{M}, F \subseteq E\}$.
*(ii)* $\lambda^-(E) = -\inf\{\lambda(F) : F \in \mathcal{M}, F \subseteq E\}$.
*(iii)* $M \in \mathcal{Z}_\lambda \Leftrightarrow |\lambda|(M) = 0$.
*(iv)* $\pi_\lambda = \lambda^+$.
*(v)* $\nu_\lambda = |\lambda|$.

*Proof.*

(*i*) Let $(P, N)$ be Hahn. If $F \in \mathcal{M}$ and $F \subseteq E$, then $\lambda(F) = \lambda^+(F) - \lambda^-(F) \leq \lambda^+(E)$, with equality at $F = E \cap P$.
(*ii*) Under the same hypotheses, $\lambda(F) \geq -\lambda^-(E)$, with equality at $F = E \cap N$.
(*iii*) ($\Rightarrow$) If $M \in \mathcal{Z}_\lambda$, then $\lambda^+(M) = \lambda^-(M) = 0$, so $|\lambda|(M) = 0$.

(*iii*) ($\Leftarrow$) If $|\lambda|(M) = 0$, then $\lambda^\pm(G \cap M) = 0\ \forall G \in \mathcal{M}$, hence $M \in \mathcal{Z}_\lambda$.
(*iv*) This is (i) and the definition of $\pi_\lambda$.
(*v*) $\sum_j |\lambda(A_j)| \leq |\lambda|(E)\ \forall$ finite measurable partitions $(A_j)$ of $E$; the Hahn partition attains equality.

*Structural Commentary*: Variation is intrinsic to $\lambda$.
*Key Insight*: The positive part is the largest charge obtainable inside a set. □

**Theorem 3.1.8.** *Charges Form a Banach Space*
*Let* $\mathrm{ca}(X, \mathcal{M}) := \{\lambda : \mathcal{M} \to \mathbb{R} : \lambda$ *is a charge*$\}$ *and* $\|\lambda\|_{\mathrm{var}} := |\lambda|(X)$.
*With pointwise operations,* $(\mathrm{ca}(X, \mathcal{M}), \|\cdot\|_{\mathrm{var}})$ *is Banach.*

*Proof.*

(i) Charge linearity gives closure under pointwise operations. The variation axioms follow from $|\lambda|(X) = 0 \Rightarrow |\lambda|(E) = 0\ \forall E$, $|a\lambda| = |a|\,|\lambda|$, and $\sum_j |(\lambda + \eta)(A_j)| \leq \sum_j |\lambda(A_j)| + \sum_j |\eta(A_j)|$ over finite measurable partitions $(A_j)$.
(ii) Let $(\lambda_n)$ be variation-Cauchy.
Since $|\lambda_n(E) - \lambda_m(E)| \leq \|\lambda_n - \lambda_m\|_{\mathrm{var}}$, set $\lambda(E) := \lim_n \lambda_n(E)$; then $\lambda$ is finitely additive.
For fixed $n$ and finite disjoint $(A_j)$, $\sum_j |(\lambda - \lambda_n)(A_j)| \leq \limsup_m \|\lambda_m - \lambda_n\|_{\mathrm{var}}$.
(iii) For disjoint $(E_k)$, set $E := \bigcup_k E_k$, $R_K := \bigcup_{k>K} E_k$, and choose $n$ with $\limsup_m \|\lambda_m - \lambda_n\|_{\mathrm{var}} < \varepsilon$.
Since $\lambda_n(R_K) \to 0$, for large $K$, $|\lambda(R_K)| \leq |\lambda_n(R_K)| + |(\lambda - \lambda_n)(R_K)| < 2\varepsilon$.
Hence $\lambda(R_K) \to 0$, so $\lambda$ is countably additive.
(iv) Taking suprema in (ii) over finite partitions gives $\|\lambda - \lambda_n\|_{\mathrm{var}} \leq \limsup_m \|\lambda_m - \lambda_n\|_{\mathrm{var}} \to 0$.
Hence every variation-Cauchy sequence converges in variation to a charge.

*Structural Commentary*: Total variation is the signed-measure analogue of an $L^1$ norm.
*Key Insight*: Variation-Cauchy controls every finite partition. □

**Theorem 3.1.9.** *Variations of Integral Charges*
*Let* $f \in \mathscr{L}^1(\mu)$ *and* $\lambda(E) := \int_E f\,d\mu$. *For* $E \in \mathcal{M}$:
(*i*) $\lambda^+(E) = \int_E f^+\,d\mu$.
(*ii*) $\lambda^-(E) = \int_E f^-\,d\mu$.
(*iii*) $|\lambda|(E) = \int_E |f|\,d\mu$.

*Proof.* Set $P := \{f \geq 0\}$ and $N := \{f < 0\}$.
For $A \in \mathcal{M}$, $\lambda(A \cap P) = \int_{A\cap P} f\,d\mu \geq 0$ and $\lambda(A \cap N) = \int_{A\cap N} f\,d\mu \leq 0$, so $(P, N)$ is Hahn. Fix $E \in \mathcal{M}$.
(i) $\lambda^+(E) = \lambda(E \cap P) = \int_E f^+\,d\mu$.
(ii) $\lambda^-(E) = -\lambda(E \cap N) = \int_E f^-\,d\mu$.
(iii) $|\lambda|(E) = \lambda^+(E) + \lambda^-(E) = \int_E |f|\,d\mu$.

*Structural Commentary*: Measure variations match function parts.
*Key Insight*: Sign sets of the density are Hahn sets. □

**Definition 3.1.10.** *Absolute Continuity and Singularity*
Let $\lambda, \mu$ be measures and let $\eta$ be a charge on $(X, \mathcal{M})$.
(i) *Absolute continuity for measures*: $\lambda \ll \mu \Leftrightarrow \forall E \in \mathcal{M}, \mu(E) = 0 \Rightarrow \lambda(E) = 0$.
(ii) *Absolute continuity for charges*: $\eta \ll \mu \Leftrightarrow |\eta| \ll \mu$.
(iii) *Mutual singularity*: $\lambda \perp \mu \Leftrightarrow \exists A, B \in \mathcal{M}$ with $A \cap B = \emptyset$, $X = A \cup B$, $\lambda(A) = 0$, and $\mu(B) = 0$.

**Proposition 3.1.11.** *Absolute Continuity Calculus*
*Let* $\lambda, \mu, \nu$ *be measures and* $\eta$ *a charge.*
(*i*) $\lambda \ll \lambda$.
(*ii*) $\lambda \ll \mu \ll \nu \Rightarrow \lambda \ll \nu$.
(*iii*) $\ll$ *is not symmetric.*
(*iv*) $(\forall E \in \mathcal{M}, \mu(E) = 0 \Rightarrow \eta(E) = 0) \Rightarrow \eta^+, \eta^-, |\eta| \ll \mu$.

*Proof.*

(i) Reflexivity follows directly from the null-set definition.
(ii) If $\lambda \ll \mu \ll \nu$ and $\nu(E) = 0$, then $\mu(E) = 0$, hence $\lambda(E) = 0$.
(iii) On $X = \{0, 1\}$, $\mathcal{M} = 2^X$, $\delta_0 \ll \#$, but $\# \not\ll \delta_0$ since $\delta_0(\{1\}) = 0$ and $\#\{1\} = 1$.
(iv) If $\mu(E) = 0$ and $F \in \mathcal{M}$ satisfies $F \subseteq E$, then $\mu(F) = 0$ and $\eta(F) = 0$.
Hence $E \in \mathcal{Z}_\eta$ and $|\eta|(E) = 0$ by Corollary 3.1.7; therefore $\eta^{\pm}(E) \leq |\eta|(E) = 0$.

*Structural Commentary*: Absolute continuity is null-set order; charges reduce to variation.
*Key Insight*: Null sets pass to measurable subsets, so all signed pieces vanish there. □

**Corollary 3.1.12.** *Absolutely Continuous and Singular Forces Zero*
*Let $\lambda, \mu$ be measures and let $\eta$ be a charge.*

*(i) $\lambda \ll \mu$ and $\lambda \perp \mu$ imply $\lambda = 0$.*
*(ii) $|\eta| \perp \mu$ implies $\eta^+ \perp \mu$.*
*(iii) $|\eta| \perp \mu$ implies $\eta^- \perp \mu$.*

*Proof.*

(i) Choose $X = A \cup B$ with $\lambda(A) = 0$ and $\mu(B) = 0$. Since $\lambda \ll \mu$, $\lambda(B) = 0$, hence $\lambda(X) = 0$.
(ii) Choose $X = A \cup B$ with $|\eta|(A) = 0$ and $\mu(B) = 0$. Since $\eta^+ \leq |\eta|$, $\eta^+(A) = 0$.
(iii) With the same $X = A \cup B$, $\eta^-(A) \leq |\eta|(A) = 0$ and $\mu(B) = 0$; hence $\eta^- \perp \mu$.

*Structural Commentary*: Absolute continuity and singular support are incompatible except at zero.
*Key Insight*: A measure supported on a null set must vanish when it also respects that null ideal. □

**Example 3.1.13.** *Dominating a Countable Family of Measures*
Let $(\mu_n)$ be measures with $\mu_n(X) \leq 1$, and set $\lambda(E) := \sum_{n \geq 1} 2^{-n} \mu_n(E)$.

Then $\lambda$ is finite, since $\lambda(X) \leq 1$, and countably additive by Tonelli for nonnegative series.
The bound $2^{-n}\mu_n(E) \leq \lambda(E)$ implies $\lambda(E) = 0 \Rightarrow \mu_n(E) = 0$ and $\mu_n \ll \lambda$ $\forall n$.
*Key Insight*: A summable positive mixture packages countably many null-set tests into one reference measure.

**Lemma 3.1.14.** *Epsilon–Delta Form of Absolute Continuity*
*Let $\lambda$ be finite and $\mu$ a measure on $(X, \mathcal{M})$. Then $\lambda \ll \mu \Leftrightarrow \forall \varepsilon > 0, \exists \delta > 0, \forall E \in \mathcal{M}, \mu(E) < \delta \Rightarrow \lambda(E) < \varepsilon$.*

*Proof.*

($\Rightarrow$) Suppose the $\varepsilon$-$\delta$ condition fails. Then $\exists \varepsilon > 0$ and $E_n \in \mathcal{M}$ with $\mu(E_n) < 2^{-n}$ and $\lambda(E_n) \geq \varepsilon$.
Let $F_n = \bigcup_{k \geq n} E_k$ and $F = \bigcap_n F_n$. Then $\mu(F_n) \leq 2^{-n+1}$, so $\mu(F) = 0$.
Since $F_n \downarrow F$, $\lambda(F_n) \geq \varepsilon$, and $\lambda$ is finite, continuity from above gives $\lambda(F) \geq \varepsilon$, contradicting $\lambda \ll \mu$.
($\Leftarrow$) If $\mu(E) = 0$, then $\forall \delta > 0$, $\mu(E) < \delta$; hence $\forall \varepsilon > 0$, $\lambda(E) < \varepsilon$, so $\lambda(E) = 0$.

*Structural Commentary*: For finite target measure, absolute continuity is uniform smallness on small $\mu$-sets.
*Key Insight*: A failing sequence is converted into a decreasing null tail with positive $\lambda$-mass. □

**Example 3.1.15.** *Finite Target Is Needed*
On $(\mathbb{N}, 2^{\mathbb{N}})$, let $\lambda = \#$ and $\mu(E) := \sum_{n \in E} 2^{-n}$. Since $\mu(E) = 0 \Leftrightarrow E = \emptyset$, $\lambda \ll \mu$.
But for $E_n = \{n\}$, $\mu(E_n) = 2^{-n} \to 0$ while $\lambda(E_n) = 1$, so the $\varepsilon$-$\delta$ condition fails for $\varepsilon = 1$.

**Example 3.1.16.** *Mutual Absolute Continuity Need Not Preserve Finiteness*
On $(\mathbb{N}, 2^{\mathbb{N}})$, set $\mu(E) := \sum_{n \in E} 2^{-n}$ and $\lambda(\emptyset) = 0$, $\lambda(E) = +\infty$ for $E \neq \emptyset$.
Then $\mu(\mathbb{N}) = 1$, while $\lambda$ is not $\sigma$-finite. Since both measures have null ideal $\{\emptyset\}$, $\lambda \ll \mu$ and $\mu \ll \lambda$.
*Key Insight*: Absolute continuity compares null sets; it does not compare finiteness, $\sigma$-finiteness, or size.

## 3.2 Radon-Nikodym Theorem

**Theorem 3.2.1.** *Radon–Nikodym Theorem*
*Let $\lambda, \mu$ be $\sigma$-finite measures on $(X, \mathcal{M})$ with $\lambda \ll \mu$.*

*(i) Representation: $\exists$ measurable $f : X \to [0, +\infty)$ such that $\lambda(E) = \int_E f \, d\mu$ for $E \in \mathcal{M}$.*
*(ii) Uniqueness: Any two representing densities agree $\mu$-a.e.*

*The $\mu$-a.e. class of $f$ is denoted by $\frac{d\lambda}{d\mu}$.*

*Proof.*

(i) Assume first $\lambda(X), \mu(X) < +\infty$. For $c > 0$, let $(P_k, N_k)$ be Hahn for $\lambda - kc\mu$, set $A_k := N_k \setminus \bigcup_{j<k} A_j$, and set $B := X \setminus \bigcup_{k\geq 1} A_k$. If $k \geq 2$, then $A_k \subseteq N_k \cap P_{k-1}$.

Hence $E \subseteq A_k$ implies $(k-1)c\mu(E) \leq \lambda(E) \leq kc\mu(E)$, with the lower bound $0 \leq \lambda(E)$ when $k = 1$. Since $B \subseteq P_k$ for $k \geq 1$, $\lambda(B) \geq kc\mu(B)\ \forall k \geq 1$; finiteness and $\lambda \ll \mu$ give $\mu(B) = \lambda(B) = 0$. Set $f_c := \sum_{k\geq 1} (k-1)c\,\mathbf{1}_{A_k}$. Countable additivity over $(A_k)$ gives $0 \leq \lambda(E) - \int_E f_c\,d\mu \leq c\mu(E)$ for $E \in \mathcal{M}$.

Let $c_n := 2^{-n}$ and $f_n := f_{c_n}$. For $E \in \mathcal{M}$, $\left|\int_E (f_n - f_m)\,d\mu\right| \leq (c_n + c_m)\mu(E)$; by Proposition 1.2.20, $|f_n - f_m| \leq c_n + c_m$ $\mu$-a.e. Thus $\sum_n \|f_{n+1} - f_n\|_1 < +\infty$. By Tonelli, $f_n$ converges $\mu$-a.e. and in $L^1(\mu)$ to a measurable $f \geq 0$. Letting $n \to \infty$ in the approximation inequality gives $\lambda(E) = \int_E f\,d\mu$.

For the $\sigma$-finite case, choose a measurable partition $X = \bigsqcup_r Y_r$ with $\lambda(Y_r), \mu(Y_r) < +\infty$. Apply the finite case on each $Y_r$, obtaining $f_r$, and set $f := \sum_r f_r \mathbf{1}_{Y_r}$.

Then $\lambda(E) = \sum_r \lambda(E \cap Y_r) = \int_E f\,d\mu$; also $f < +\infty$ $\mu$-a.e. because $\int_{Y_r} f\,d\mu = \lambda(Y_r) < +\infty\ \forall r$.

(ii) If $f, g$ represent $\lambda$, then $f, g \in L^1(\mu|_{Y_r})$ and $f = g$ $\mu$-a.e. on $Y_r$ by Proposition 1.2.20.
Taking the countable union of the exceptional sets gives $f = g$ $\mu$-a.e. on $X$.

*Structural Commentary*: Hahn decompositions of $\lambda - c\mu$ locate density bands.
*Key Insight*: Summably refined band approximations converge without invoking later $L^1$ completeness. □

**Example 3.2.2.** *A Non-$\sigma$-Finite Failure*
Let $X = [0,1]$, $\mathcal{M} = \mathcal{B}([0,1])$, $\mu = \#$, and $\lambda = m$. Since $\mu(E) = 0 \Leftrightarrow E = \emptyset$, $\lambda \ll \mu$.
If $\lambda(E) = \int_E f\,d\mu$ for some $f \geq 0$, then $\forall x$, $0 = \lambda(\{x\}) = \int_{\{x\}} f\,d\mu = f(x)$, so $\lambda([0,1]) = 0$, contradiction.
Thus RN can fail when $\mu$ is not $\sigma$-finite.

**Example 3.2.3.** *Another Non-$\sigma$-Finite Failure*
Let $X$ be uncountable, $\mathcal{M} := \{E \subseteq X : E \text{ or } X \setminus E \text{ countable}\}$, $\mu(E) := \#E$ if $E$ is finite and $+\infty$ otherwise, and $\lambda(E) := 0$ if $E$ is countable and $+\infty$ otherwise. Then $\mu(E) = 0 \Leftrightarrow E = \emptyset$, so $\lambda \ll \mu$.
If $\lambda(E) = \int_E f\,d\mu$ with $f \geq 0$, then $\forall x$, $0 = \lambda(\{x\}) = \int_{\{x\}} f\,d\mu = f(x)$, forcing $\lambda(X) = 0$, contradiction.
*Structural Commentary*: Singleton tests determine $f$, but not $\lambda(X)$ without $\sigma$-finiteness.
*Key Insight*: Non-$\sigma$-finite domination can lose global mass.

**Remark 3.2.4.** *When the Radon–Nikodym Derivative Is Integrable*
In Theorem 3.2.1, $\frac{d\lambda}{d\mu} \in L^1(\mu) \Leftrightarrow \lambda(X) < +\infty$, since $\int_X \frac{d\lambda}{d\mu}\,d\mu = \lambda(X)$.

**Theorem 3.2.5.** *Extended-Valued Radon–Nikodym Theorem*
*Let $\mu$ be $\sigma$-finite, let $\lambda$ be a measure on $(X, \mathcal{M})$, and assume $\lambda \ll \mu$.*

*(i) Representation: $\exists f : X \to [0, +\infty]$ measurable such that $\lambda(E) = \int_E f\,d\mu$ for $E \in \mathcal{M}$.*
*(ii) Uniqueness: Any two representing densities agree $\mu$-a.e.*

*Proof.*

(i) Assume first $\mu(X) < +\infty$ and set $\alpha := \sup\{\mu(A) : A \in \mathcal{M}, \lambda(A) < +\infty\}$. Choose $(A_n) \subseteq \mathcal{M}$ with $\lambda(A_n) < +\infty$ and $\mu(A_n) \to \alpha$, and set $C_n := \bigcup_{j\leq n} A_j$, $N := \bigcup_n C_n$, and $P := X \setminus N$. Then $\lambda|_N$ is $\sigma$-finite.

For $E \subseteq P$, $\lambda(E) < +\infty \Rightarrow \mu(E) = 0$, since $\mu(E) > 0$ would give $\mu(C_n \cup E) = \mu(C_n) + \mu(E) > \alpha$ eventually. Hence $\mu(E) > 0 \Rightarrow \lambda(E) = +\infty$, while $\mu(E) = 0 \Rightarrow \lambda(E) = 0$ by $\lambda \ll \mu$.
By Theorem 3.2.1, $\exists g : N \to [0, +\infty)$ measurable such that $\lambda(E \cap N) = \int_{E\cap N} g\,d\mu$. Define $f = g$ on $N$ and $f = +\infty$ on $P$; preceding dichotomy gives $\lambda(E) = \int_E f\,d\mu$. For $\sigma$-finite $\mu$, $X = \bigsqcup_r Y_r$, $\mu(Y_r) < +\infty$.

Let $f_r$ be the finite-$\mu$ density on $Y_r$ and set $f := \sum_r f_r \mathbf{1}_{Y_r}$. Then $\lambda(E) = \sum_r \lambda(E \cap Y_r) = \int_E f\,d\mu$.

(ii) If $f, h$ represent $\lambda$, fix $r, m, n \in \mathbb{N}$ and set $A^r_{m,n} := Y_r \cap \{f \leq m\} \cap \{h \geq f + \frac{1}{n}\}$.
Since $\int_A f\,d\mu = \int_A h\,d\mu$ for measurable $A \subseteq Y_r$ and $\int_{A^r_{m,n}} f\,d\mu < +\infty$, $\mu(A^r_{m,n}) = 0$.
Taking the union over $r, m, n$ gives $\mu\{f < h\} = 0$; symmetry gives $\mu\{h < f\} = 0$.

*Structural Commentary*: Maximal finite-$\lambda$ regions separate ordinary density from infinite density.
*Key Insight*: Outside the maximal $\sigma$-finite part, every positive-$\mu$ set has infinite $\lambda$-mass. □

**Theorem 3.2.6.** *Lebesgue Decomposition Theorem*
*Let $\lambda, \mu$ be $\sigma$-finite measures on $(X, \mathcal{M})$.*

*(i) $\exists$ measures $\lambda_s, \lambda_a$ with $\lambda = \lambda_s + \lambda_a$, $\lambda_s \perp \mu$, and $\lambda_a \ll \mu$.*
*(ii) If $(\tilde{\lambda}_s, \tilde{\lambda}_a)$ is another such pair, then $\tilde{\lambda}_s = \lambda_s$ and $\tilde{\lambda}_a = \lambda_a$.*

*Proof.*

(i) Put $\rho := \lambda + \mu$ and choose $f = \frac{d\lambda}{d\rho}$, $g = \frac{d\mu}{d\rho}$. With $A := \{g = 0\}$ and $B := \{g > 0\}$, set $\lambda_s(E) := \lambda(E \cap A)$ and $\lambda_a(E) := \lambda(E \cap B)$. Then $\lambda = \lambda_s + \lambda_a$. Since $\mu(A) = \int_A g \, d\rho = 0$, $\lambda_s \perp \mu$.
If $\mu(E) = 0$, then $g = 0$ $\rho$-a.e. on $E$, so $\rho(E \cap B) = 0$ and $\lambda_a(E) = 0$; hence $\lambda_a \ll \mu$.

(ii) Let $(\tilde{\lambda}_s, \tilde{\lambda}_a)$ be another such pair. Since $\lambda_s \perp \mu$ and $\tilde{\lambda}_a \ll \mu$, $\lambda_s \perp \tilde{\lambda}_a$; choose $X = C \cup D$ with $\lambda_s(C) = 0$ and $\tilde{\lambda}_a(D) = 0$. Then $\lambda_s(E) = \lambda_s(E \cap D) \leq \lambda(E \cap D) = \tilde{\lambda}_s(E \cap D) \leq \tilde{\lambda}_s(E)$.
By symmetry, $\lambda_s = \tilde{\lambda}_s$, hence $\lambda_a = \tilde{\lambda}_a$.

*Structural Commentary*: Split $\lambda$ into $\mu$-visible and $\mu$-singular parts.
*Key Insight*: Use $\rho = \lambda + \mu$ and split where $\frac{d\mu}{d\rho} = 0$. □




Figure 210: Charges split by sign; measures split into $\lambda_s \perp \mu$ and $\lambda_a \ll \mu$ parts.

**Proposition 3.2.7.** *Radon–Nikodym Calculus*
*Let all measures below be $\sigma$-finite.*

*(i)* $\lambda \ll \mu$, $g \geq 0$ *measurable* $\Rightarrow \int_X g \, d\lambda = \int_X g \frac{d\lambda}{d\mu} \, d\mu$.
*(ii)* $\nu \ll \lambda \ll \mu \Rightarrow \frac{d\nu}{d\mu} = \frac{d\nu}{d\lambda}\frac{d\lambda}{d\mu}$ $\mu$-*a.e.*
*(iii)* $\lambda_1, \lambda_2 \ll \mu \Rightarrow \frac{d(\lambda_1+\lambda_2)}{d\mu} = \frac{d\lambda_1}{d\mu} + \frac{d\lambda_2}{d\mu}$ $\mu$-*a.e.*
*(iv)* $\lambda \ll \mu \ll \lambda \Rightarrow \frac{d\lambda}{d\mu} = \left(\frac{d\mu}{d\lambda}\right)^{-1}$ $\mu$-*a.e.*

*Proof.*

(i) For $g = \mathbf{1}_E$, $\int_X g \, d\lambda = \lambda(E) = \int_E \frac{d\lambda}{d\mu} \, d\mu = \int_X g \frac{d\lambda}{d\mu} \, d\mu$.
By linearity, identity holds for nonnegative simple $g$.
For measurable $g \geq 0$, take simple $g_n \uparrow g$ and apply monotone convergence to both sides.

(ii) Let $h := \frac{d\nu}{d\lambda}\frac{d\lambda}{d\mu}$. $\forall E \in \mathcal{M}$, (i) with $g = \mathbf{1}_E \frac{d\nu}{d\lambda}$ gives $\nu(E) = \int_E \frac{d\nu}{d\lambda} \, d\lambda = \int_E h \, d\mu$.
Hence $h$ represents $\nu$ relative to $\mu$, so $h = \frac{d\nu}{d\mu}$ $\mu$-a.e.

(iii) Let $h := \frac{d\lambda_1}{d\mu} + \frac{d\lambda_2}{d\mu}$. $\forall E \in \mathcal{M}$, $\int_E h \, d\mu = \lambda_1(E) + \lambda_2(E) = (\lambda_1 + \lambda_2)(E)$.
Hence $h$ represents $\lambda_1 + \lambda_2$ relative to $\mu$, so uniqueness gives the identity.

(iv) Apply (ii) to $\mu \ll \lambda \ll \mu$. Since $\frac{d\mu}{d\mu} = 1$, $1 = \frac{d\mu}{d\lambda}\frac{d\lambda}{d\mu}$ $\mu$-a.e.
Since $\mu \ll \lambda$, $\frac{d\mu}{d\lambda} > 0$ $\mu$-a.e.; divide to obtain $\frac{d\lambda}{d\mu} = \left(\frac{d\mu}{d\lambda}\right)^{-1}$ $\mu$-a.e.

*Structural Commentary*: Radon–Nikodym derivatives obey chain and linearity rules.
*Key Insight*: Prove identities by integrating over arbitrary measurable sets, then invoke uniqueness. □

**Corollary 3.2.8.** *Radon–Nikodym for Charges*
*Let $\mu$ be $\sigma$-finite and let $\lambda$ be a charge with $\lambda \ll \mu$. Then $\exists f \in \mathscr{L}^1(\mu)$ such that $\forall E \in \mathcal{M}$:*

*(i)* $\lambda(E) = \int_E f \, d\mu$.
*(ii)* $\lambda^+(E) = \int_E f^+ \, d\mu$.
*(iii)* $\lambda^-(E) = \int_E f^- \, d\mu$.
*(iv)* $|\lambda|(E) = \int_E |f| \, d\mu$.

*Proof.* Since $|\lambda| \ll \mu$, $\lambda^\pm \ll \mu$. By Theorem 3.2.1, choose $f_\pm \geq 0$ with $\lambda^\pm(E) = \int_E f_\pm \, d\mu$.
Mutual singularity of $\lambda^+$ and $\lambda^-$ gives $f_+ f_- = 0$ $\mu$-a.e. Set $f := f_+ - f_-$.
Then $f^+ = f_+$ and $f^- = f_-$ $\mu$-a.e., while $\int_X |f| \, d\mu = |\lambda|(X) < +\infty$.

(i) $\lambda(E) = \lambda^+(E) - \lambda^-(E) = \int_E f \, d\mu$.
(ii) $\lambda^+(E) = \int_E f_+ \, d\mu = \int_E f^+ \, d\mu$.
(iii) $\lambda^-(E) = \int_E f_- \, d\mu = \int_E f^- \, d\mu$.
(iv) $|\lambda|(E) = \lambda^+(E) + \lambda^-(E) = \int_E |f| \, d\mu$.

*Structural Commentary*: Signed absolute continuity is reduced to the two positive Jordan pieces.
*Key Insight*: Differentiate $\lambda^+$ and $\lambda^-$ separately, then subtract. □

**Example 3.2.9.** *Radon–Nikodym Classification of Basic Stieltjes Generators*
Let $m$ be Lebesgue measure on $\mathcal{B}(\mathbb{R})$, and let $\lambda_i$ be the Borel–Stieltjes measure generated by $g_i$.

(i) $g_1(x) = 2x$: $\lambda_1 = 2m$, so $\lambda_1 \ll m$, $\frac{d\lambda_1}{dm} = 2$, $m \ll \lambda_1$, $\frac{dm}{d\lambda_1} = \frac{1}{2}$, and $\lambda_1(\mathbb{R}) = +\infty$.
(ii) $g_2(x) = \arctan x$: $\lambda_2(E) = \int_E (1+x^2)^{-1}\, dm$, so $\frac{d\lambda_2}{dm} = (1+x^2)^{-1}$, $\lambda_2(\mathbb{R}) = \pi$, $m \ll \lambda_2$, and $\frac{dm}{d\lambda_2} = 1 + x^2$.
(iii) $g_3 = \mathbf{1}_{[0,+\infty)}$: $\lambda_3 = \delta_0$, so $\lambda_3 \perp m$, $\lambda_3(\mathbb{R}) = 1$, $\lambda_3 \not\ll m$, and $m \not\ll \lambda_3$.
(iv) $g_4(x) = x\mathbf{1}_{[0,+\infty)}(x)$: $\lambda_4(E) = m(E \cap [0,+\infty))$, so $\lambda_4 \ll m$, $\frac{d\lambda_4}{dm} = \mathbf{1}_{[0,+\infty)}$, $m \not\ll \lambda_4$, and $\lambda_4(\mathbb{R}) = +\infty$.

Thus $\lambda_1, \lambda_2, \lambda_4 \ll m$, $\lambda_3 \perp m$, only $\lambda_2, \lambda_3$ are finite, and $m \ll \lambda_i$ exactly for $i = 1, 2$.
*Structural Commentary*: Stieltjes generators split naturally into density, atom, and restriction examples.
*Key Insight*: A positive density makes null sets equivalent; an atom lives on a Lebesgue-null set.

## 3.3 Differentiation of Measures

**Definition 3.3.1.** *Averages and Absolute Continuity*

(i) *Average*: for $f \in \mathscr{L}^1(E)$ and measurable $A \subseteq E$ with $m(A) > 0$, set $[f : A] := m(A)^{-1} \int_A f\, dm$.
(ii) *Density ratio*: for measurable $B$ and cube or ball $Q$, $m(Q) > 0$, set $[B : Q] := \frac{m(B \cap Q)}{m(Q)}$.
(iii) *Indefinite integral*: for $f \in \mathscr{L}^1([a,b])$, $F(x) = F(a) + \int_a^x f(t)\, dt$.
(iv) *Absolute continuity*: $F : [a,b] \to \mathbb{R}$ is absolutely continuous iff $\forall \varepsilon > 0\ \exists \delta > 0$: if $(a_j, b_j)_{j=1}^N \subseteq [a,b]$ is a finite pairwise disjoint family, then $\sum_j (b_j - a_j) < \delta \Rightarrow \sum_j |F(b_j) - F(a_j)| < \varepsilon$.

**Perspective 3.3.2.** *Absolute Continuity as Interval Bookkeeping*
If $F(x) = \int_a^x f(t)\, dt$, absolute continuity controls total $F$-variation on disjoint intervals of small total length.




Figure 211: Interval bookkeeping for absolute continuity.

**Perspective 3.3.3.** *Vitali Covering Control*
Select balls disjointly by decreasing radius.
If $B$ is rejected, then $B \cap B_j \neq \emptyset$ for some earlier $B_j$ with $r(B) \leq r(B_j)$, hence $B \subseteq 5B_j$.
*Key Insight*: Disjoint balls add; fixed dilations cover misses.




Figure 212: A comparable missed ball lies inside the fivefold dilation of the selected ball.

**Definition 3.3.4.** *Vitali Coverings and Density Points*
Let $(M, d)$ be a metric space and let $A \subseteq M$.

(i) $\mathcal{V}$ is a *Vitali covering* of $A$ iff $\forall p \in A\ \forall r > 0\ \exists V \in \mathcal{V}$ such that $p \in V$ and $\operatorname{diam}(V) < r$.
(ii) For measurable $E \subseteq \mathbb{R}^n$, its *density-point set* is $\operatorname{dp}(E) := \{p \in \mathbb{R}^n : \lim_{Q \downarrow p} [E : Q] = 1\}$, with $Q$ cubes, equivalently balls, shrinking to $p$.

**Theorem 3.3.5.** *Vitali Covering Lemma for Balls and Cubes*
*Let $E \subseteq \mathbb{R}^n$, $m^*(E) < +\infty$, and let $\mathcal{V}$ be a Vitali cover of $E$ by closed balls or axis-parallel cubes.*
*Then $\exists (V_j)_{j \geq 1} \subseteq \mathcal{V}$ pairwise disjoint with $m^*(E \setminus \bigcup_{j \geq 1} V_j) = 0$.*

*Proof.* Fix open $O \supseteq E$ with $m(O) < m^*(E) + \varepsilon$, and replace $\mathcal{V}$ by $\{V \in \mathcal{V} : V \subseteq O\}$.
Greedily choose a maximal disjoint $(V_j)$ by dyadic diameter scales.
For a fixed dilation $V \mapsto V^*$, if $W \cap V \neq \emptyset$ and $\operatorname{diam} W \leq 2 \operatorname{diam} V$, then $W \subseteq V^*$ and $m(V^*) \leq C_n m(V)$.
Let $R = E \setminus \bigcup_j V_j$. Fix $N$ and $p \in R$.

Since $V_1, \ldots, V_N$ are closed, choose $W \in \mathcal{V}$ with $p \in W$ and $W \cap \bigcup_{j \leq N} V_j = \emptyset$. Maximality gives some $j > N$ with $W \cap V_j \neq \emptyset$ and comparable diameter, hence $p \in W \subseteq V_j^*$. Thus $R \subseteq \bigcup_{j>N} V_j^*$.

Therefore $m^*(R) \leq C_n \sum_{j>N} m(V_j) \to 0$, because $\sum_j m(V_j) = m(\bigsqcup_j V_j) \leq m(O) < +\infty$. Hence $m^*(R) = 0$.
*Structural Commentary*: Vitali selection gives a disjoint cover modulo null sets.
*Key Insight*: $R \subseteq \bigcup_{j>N} V_j^*$ and disjoint tails vanish. □

**Theorem 3.3.6.** *Lebesgue Density Theorem*
*For measurable $E \subseteq \mathbb{R}^n$, $\lim_{Q \downarrow p} [E : Q] = \mathbf{1}_E(p)$ for a.e. $p$, with $Q \downarrow p$ through cubes or balls.*

*Proof.* Assume $m(E) < +\infty$. For $a \in (0,1)$ set $E_a := \{p \in E : \liminf_{Q \downarrow p} [E : Q] < a\}$.
Cubes $Q \ni p$ with $[E : Q] < a$ form a Vitali cover of $E_a$. For open $O \supseteq E_a$ with $m(O) < m(E_a) + \varepsilon$, choose disjoint $(Q_j) \subseteq O$ covering $E_a$ modulo null sets by Theorem 3.3.5.
Hence $m(E_a) \leq \sum_j m(E_a \cap Q_j) \leq \sum_j m(E \cap Q_j) < a \sum_j m(Q_j) \leq a\, m(O) < a(m(E_a) + \varepsilon)$.
Let $\varepsilon \downarrow 0$; since $a < 1$, $m(E_a) = 0$. Thus $p \in E \setminus \bigcup_{a \in \mathbb{Q} \cap (0,1)} E_a$ has density 1. Apply same argument to bounded pieces of $E^c$ to get density 0 a.e. on $E^c$; then exhaust arbitrary $E$ by bounded pieces.
*Structural Commentary*: Measurable sets are locally 0 or 1 a.e.
*Key Insight*: Vitali turns bad density into $m(E_a) \leq a\, m(E_a)$. □

**Corollary 3.3.7.** *Density Points Represent a Measurable Set*
*For measurable $E \subseteq \mathbb{R}^n$, $m(E \triangle \operatorname{dp}(E)) = 0$.*

*Proof.* By Theorem 3.3.6, $[E : Q] \to 1$ at a.e. point of $E$, $[E : Q] \to 0$ at a.e. point of $E^c$; hence $E \triangle \operatorname{dp}(E)$ null.
*Structural Commentary*: The pointwise density dichotomy is converted into equality modulo null sets.
*Key Insight*: Density-one membership recovers the measurable set almost everywhere. □

**Corollary 3.3.8.** *Density Points Commute with Slicing Almost Everywhere*
*Let $E \subseteq \mathbb{R}^n \times \mathbb{R}^k$ be measurable. For a.e. $x \in \mathbb{R}^n$, $\operatorname{dp}(E_x) = \{y : (x,y) \in \operatorname{dp}(E)\}$ up to a null subset of $\mathbb{R}^k$.*

*Proof.* Set $D = \operatorname{dp}(E)$ and $N = E \triangle D$.
By Corollary 3.3.7, $m_{n+k}(N) = 0$; hence $m_k(N_x) = 0$ for a.e. $x$ by Theorem 2.2.14.
Intersecting with the Fubini good set, $E_x, D_x$ are measurable and $m_k(E_x \triangle D_x) = m_k((E \triangle D)_x) = 0$.
$m_k(E_x \triangle \operatorname{dp}(E_x)) = 0$ by Corollary 3.3.7 in $\mathbb{R}^k$. Therefore $m_k(\operatorname{dp}(E_x) \triangle D_x) = 0$.
*Structural Commentary*: Almost-everywhere local fullness is compatible with almost-everywhere slicing.
*Key Insight*: Null ambient errors have null slices for almost every base point. □

**Lemma 3.3.9.** *Large Averages Have Small Outer Measure*
*Let $Q_0 \subseteq \mathbb{R}^n$ be a cube, let $0 \leq g \in \mathscr{L}^1(Q_0)$, and let $\lambda > 0$.*
*Set $A_\lambda := \{p \in Q_0 : \forall r > 0\ \exists$ a cube $Q \subseteq Q_0$, $p \in Q$, $\operatorname{diam} Q < r$, $[g : Q] > \lambda\}$. Then $m^*(A_\lambda) \leq \frac{1}{\lambda} \int_{Q_0} g\, dm$.*

*Proof.* Let $\mathcal{V}_\lambda := \{Q \subseteq Q_0 : Q$ is a cube and $[g : Q] > \lambda\}$. This is a Vitali cover of $A_\lambda$; choose disjoint $(Q_j) \subseteq \mathcal{V}_\lambda$ covering $A_\lambda$ modulo a null set. Then $\lambda m^*(A_\lambda) \leq \lambda \sum_j m(Q_j) < \sum_j \int_{Q_j} g\, dm \leq \int_{Q_0} g\, dm$.
*Structural Commentary*: Weak type for local averages.
*Key Insight*: Disjoint Vitali cubes turn pointwise large averages into one integral bound. □

**Theorem 3.3.10.** *Lebesgue Average Value Theorem*
*For $f \in \mathscr{L}^1_{\mathrm{loc}}(\mathbb{R}^n)$, $\lim_{Q \downarrow p} [f : Q] = f(p)$ for a.e. $p \in \mathbb{R}^n$ with $Q$ cubes, equivalently balls, shrinking to $p$.*

*Proof.* Fix cube $Q_0$; to prove claim for a.e. $p \in Q_0^\circ$. For $\alpha > 0$, let $B_\alpha = \left\{ p \in Q_0^\circ : \limsup_{Q \downarrow p} |[f : Q] - f(p)| > \alpha \right\}$.
Choose $h \in C(Q_0)$ with $\|f - h\|_{L^1(Q_0)} < \eta$, and set $g = |f - h|$. Since $[h : Q] \to h(p)$, $B_\alpha \subseteq \{g > \frac{\alpha}{3}\} \cup A_{\frac{\alpha}{3}}(g)$.
By Markov and Lemma 3.3.9, $m^*(B_\alpha) \leq \frac{3}{\alpha} \int_{Q_0} g + \frac{3}{\alpha} \int_{Q_0} g < \frac{6\eta}{\alpha}$. Let $\eta \downarrow 0$; then $m(B_\alpha) = 0$.
Hence $B(Q_0) := \{p \in Q_0^\circ : \limsup_{Q \downarrow p} |[f : Q] - f(p)| > 0\} = \bigcup_{\alpha \in \mathbb{Q}_{>0}} B_\alpha$ is null.

With $Q_N := [-N, N]^n$, $\mathbb{R}^n = \bigcup_N Q_N^\circ$, so $\bigcup_N B(Q_N)$ is null.

*Structural Commentary*: Local averages recover the original integrable function almost everywhere.
*Key Insight*: Continuous approximation handles the good part; Vitali's estimate makes the bad part small. □

**Theorem 3.3.11.** *Lebesgue's Antiderivative Theorem*
*For $f \in \mathscr{L}^1([a,b])$, define $F(x) := c + \int_a^x f(t)\,dt$.*

*(i) $F \in AC([a,b])$.*
*(ii) $F' = f$ a.e. on $[a,b]$.*

*Proof.*

(i) Given $\varepsilon > 0$, choose $\delta > 0$ such that $m(A) < \delta$ implies $\int_A |f|\ dm < \varepsilon$.
If $(\alpha_j, \beta_j)$ are disjoint and $\sum_j (\beta_j - \alpha_j) < \delta$, then $\sum_j |F(\beta_j) - F(\alpha_j)| \le \int_{\bigcup_j (\alpha_j,\beta_j)} |f|\ dm < \varepsilon$.

(ii) For $x, x+h \in [a,b]$, let $I_h(x) := (\min\{x, x+h\}, \max\{x, x+h\})$. Then $\frac{F(x+h)-F(x)}{h} = [f : I_h(x)]$.
Extend $f$ by 0 outside $[a,b]$; by Theorem 3.3.10 in one dimension, $[f : I_h(x)] \to f(x)$ for a.e. $x$.

*Structural Commentary*: The indefinite Lebesgue integral is the correct antiderivative for integrable functions.
*Key Insight*: Differentiating the integral is the same as shrinking the averaging interval. □

## 3.4 Monotone and Bounded-Variation Differentiation

Monotone functions are the one-dimensional model case for differentiation almost everywhere. Bounded-variation functions enter by Jordan decomposition: finite oscillation is exactly enough to write a function as the difference of two monotone functions.

**Definition 3.4.1.** *Bounded Variation*
Let $F : [a,b] \to \mathbb{R}$. For $a \le u < v \le b$, let $\mathcal{P}[u,v] := \{P = (x_0, \dots, x_N) : N \in \mathbb{N},\ u = x_0 < \cdots < x_N = v\}$.

(i) For $P = (x_0, \dots, x_N) \in \mathcal{P}[u,v]$, define $V(F,P) := \sum_{j=1}^{N} |F(x_j) - F(x_{j-1})|$.

(ii) Define $V_u^v(F) := \sup_{P \in \mathcal{P}[u,v]} V(F,P)$ for $u < v$, and $V_u^u(F) := 0$.

(iii) $F$ has *bounded variation* on $[a,b]$ iff $F \in BV([a,b]) := \{G : [a,b] \to \mathbb{R} : V_a^b(G) < +\infty\}$.

**Theorem 3.4.2.** *Variation Function and Jordan Decomposition*
*Let $F : [a,b] \to \mathbb{R}$. For $F \in BV([a,b])$, set $T_F : [a,b] \to \mathbb{R}$ by $T_F(a) := 0$ and $T_F(x) := V_a^x(F)$ for $a < x \le b$.*

*(i) Variation additivity: If $F \in BV([a,b])$, then $T_F(v) - T_F(u) = V_u^v(F)$ for $a \le u \le v \le b$.*
*(ii) Variation monotonicity: If $F \in BV([a,b])$, then $T_F$ is nondecreasing.*
*(iii) Compensated monotonicity: If $F \in BV([a,b])$, then $F + T_F$ is nondecreasing.*
*(iv) Decomposition: If $F \in BV([a,b])$, then $F = (F + T_F) - T_F$ is a difference of nondecreasing functions.*
*(v) Converse: If $F = G - H$ for nondecreasing $G, H : [a,b] \to \mathbb{R}$, then $F \in BV([a,b])$ and $V_a^b(F) \le G(b) - G(a) + H(b) - H(a)$.*

*Proof.*

(i) Assume $F \in BV([a,b])$ and fix $a \le u \le v \le b$. The cases $u = a$ and $u = v$ are immediate; for $a < u < v$, concatenation gives $V_a^v(F) \ge V_a^u(F) + V_u^v(F)$, while inserting $u$ into $P \in \mathcal{P}[a,v]$ and using the triangle inequality gives $V(F,P) \le V_a^u(F) + V_u^v(F)$. Taking suprema yields $V_a^v(F) = V_a^u(F) + V_u^v(F)$, QED.

(ii) By (i), $T_F(v) - T_F(u) = V_u^v(F) \ge 0$.

(iii) Since $|F(v) - F(u)| \le T_F(v) - T_F(u)$, $(F + T_F)(v) - (F + T_F)(u) \ge 0$.

(iv) The identity $F = (F + T_F) - T_F$ and (ii)–(iii) give the required monotone difference.

(v) Assume $F = G - H$ for nondecreasing $G, H : [a,b] \to \mathbb{R}$.
For $P = (x_0, \dots, x_N) \in \mathcal{P}[a,b]$, by monotonicity and triangle inequality,
$V(F,P) \le \sum_{j=1}^{N} (G(x_j) - G(x_{j-1})) + \sum_{j=1}^{N} (H(x_j) - H(x_{j-1})) = G(b) - G(a) + H(b) - H(a)$.
Taking the supremum over $P$ gives $V_a^b(F) \le G(b) - G(a) + H(b) - H(a) < +\infty$.

*Structural Commentary*: Bounded variation is exactly controlled monotone difference.
*Key Insight*: The variation function supplies the missing monotone part. □




Figure 213: Variation $T_F$ turns $F$ into the increasing function $F + T_F$.

**Example 3.4.3.** *Basic Bounded-Variation Tests*

(i) If $F : [a,b] \to \mathbb{R}$ is nondecreasing, then $F \in BV([a,b])$ and $V_a^b(F) = F(b) - F(a)$.
(ii) If $F : [a,b] \to \mathbb{R}$ is $L$-Lipschitz, then $F \in BV([a,b])$ and $V_a^b(F) \le L(b-a)$.
(iii) Let $F : [0,1] \to \mathbb{R}$ satisfy $F(0) = 0$ and $F(x) = x\cos(\frac{\pi}{2x})$ for $0 < x \le 1$. Then $F$ is continuous and $F \notin BV([0,1])$: for $P_n = (0, \frac{1}{2n}, \frac{1}{2n-1}, \dots, \frac{1}{2}, 1) \in \mathcal{P}[0,1]$, $V(F,P_n) = \sum_{k=1}^{n} \frac{1}{k} \to +\infty$.

**Definition 3.4.4.** *Upper and Lower Derivatives*
Let $I \subseteq \mathbb{R}$ be an interval, $F : I \to \mathbb{R}$, and $x \in I^\circ$.
For $h > 0$, set $Q_h(x) := \{t \in \mathbb{R} : 0 < |t| \le h,\ x + t \in I\}$ and $\Delta_t F(x) := \frac{F(x+t)-F(x)}{t}$.

(i) The *upper derivative* at $x$ is $\overline{D}F(x) := \lim_{h \downarrow 0} \sup_{t \in Q_h(x)} \Delta_t F(x)$.
(ii) The *lower derivative* at $x$ is $\underline{D}F(x) := \lim_{h \downarrow 0} \inf_{t \in Q_h(x)} \Delta_t F(x)$.

They exist in $\overline{\mathbb{R}}$, satisfy $\underline{D}F(x) \le \overline{D}F(x)$, and $F'(x)$ exists iff $\underline{D}F(x) = \overline{D}F(x) \in \mathbb{R}$.

**Lemma 3.4.5.** *Weak Bound for Upper Derivatives*
*Let $F : [a,b] \to \mathbb{R}$ be nondecreasing. For $\alpha > 0$, set $E_\alpha := \{x \in (a,b) : \overline{D}F(x) \ge \alpha\}$.*
*Then:*
*(i) $m^*(E_\alpha) \le \frac{F(b)-F(a)}{\alpha}$ for every $\alpha > 0$.*
*(ii) $m^*\{x \in (a,b) : \overline{D}F(x) = +\infty\} = 0$.*

*Proof.*

(i) Fix $\alpha > 0$ and $0 < \alpha' < \alpha$. For $x \in E_\alpha$ and $\delta > 0$, choose $[c,d] \subset (a,b)$ with $x \in [c,d]$, $d - c < \delta$, and $F(d) - F(c) > \alpha'(d-c)$. These intervals form a Vitali cover of $E_\alpha$.
By Theorem 3.3.5, choose pairwise disjoint $I_k = [c_k, d_k]$ covering $E_\alpha$ modulo a null set.
Then $\alpha' m^*(E_\alpha) \le \alpha' \sum_k (d_k - c_k) < \sum_k (F(d_k) - F(c_k)) \le F(b) - F(a)$. Let $\alpha' \uparrow \alpha$.
(ii) Set $E_\infty := \{x \in (a,b) : \overline{D}F(x) = +\infty\}$. Since $E_\infty \subseteq E_n$ $\forall n \in \mathbb{N}$, (i) gives $m^*(E_\infty) \le \frac{F(b)-F(a)}{n} \to 0$.

*Structural Commentary*: Vitali selection converts local slope lower bounds into total rise.
*Key Insight*: Large local slopes are paid for by total monotone rise. □

**Lemma 3.4.6.** *Dini Gap Sets Are Null*
*Let $F : [a,b] \to \mathbb{R}$ be nondecreasing and let $\alpha < \beta$.*
*Set $E_{\alpha,\beta} := \{x \in (a,b) : \underline{D}F(x) < \alpha < \beta < \overline{D}F(x)\}$. Then $m^*(E_{\alpha,\beta}) = 0$.*

*Proof.* If $\alpha \le 0$, then $E_{\alpha,\beta} = \emptyset$ because every secant quotient of nondecreasing $F$ is nonnegative.
Assume $0 < \alpha < \beta$, set $E = E_{\alpha,\beta}$, and fix $\varepsilon > 0$. Choose open $O$ with $E \subseteq O \subseteq (a,b)$ and $m(O) < m^*(E) + \varepsilon$.
Since $\underline{D}F(x) < \alpha$ on $E$, intervals $[c,d] \subset O$ with $x \in [c,d]$, $F(d) - F(c) < \alpha(d-c)$ form Vitali cover of $E$.
Select pairwise disjoint $I_k = [c_k, d_k]$ covering $E$ modulo a null set.
$\forall k$, modulo endpoints, $E \cap I_k \subseteq \{x \in (c_k, d_k) : \overline{D}(F|_{I_k})(x) \ge \beta\}$, so $m^*(E \cap I_k) \le \frac{F(d_k)-F(c_k)}{\beta}$ by Lemma 3.4.5.
Hence $m^*(E) \le \sum_k m^*(E \cap I_k) \le \frac{1}{\beta} \sum_k (F(d_k) - F(c_k)) < \frac{\alpha}{\beta} \sum_k (d_k - c_k) \le \frac{\alpha}{\beta} m(O)$.
Let $\varepsilon \downarrow 0$; then $m^*(E) \le \frac{\alpha}{\beta} m^*(E)$, and since $m^*(E) < +\infty$ and $\frac{\alpha}{\beta} < 1$, $m^*(E) = 0$.
*Structural Commentary*: Vitali intervals turn a slope gap into a contraction estimate.
*Key Insight*: Dini gaps force excess rise. □

**Perspective 3.4.7.** *Right Secants and Differentiation*
Right quotients $g_n(x) = n\big(F(x + \frac{1}{n}) - F(x)\big)$ encode local slope data.
*Key Insight*: Secants turn differentiation into slope estimates.




Figure 214: $g_n(x)$ is the slope of the short right secant from $x$ to $x+\frac{1}{n}$.

**Proposition 3.4.8.** *Divided Differences and Moving Averages*
*Let $f \in \mathscr{L}^1([a,b])$, extend $f$ to $[a,b+1]$ by $f(x)=f(b)$ for $b<x\le b+1$, and fix $0<h\le 1$.*
*For $x\in[a,b]$, set $\operatorname{Diff}_h f(x) := \frac{f(x+h)-f(x)}{h}$ and $\operatorname{Av}_h f(x) := h^{-1}\int_x^{x+h} f\,dm$.*
*Then $\forall a\le u<v\le b$, $\int_u^v \operatorname{Diff}_h f\,dm = \operatorname{Av}_h f(v) - \operatorname{Av}_h f(u)$.*

*Proof.* For $a\le u<v\le b$, translation invariance gives $\int_u^v f(\cdot+h)\,dm = \int_{u+h}^{v+h} f\,dm$.
Hence $\int_u^v \operatorname{Diff}_h f\,dm = h^{-1}(\int_{u+h}^{v+h} f\,dm - \int_u^v f\,dm) = h^{-1}(\int_v^{v+h} f\,dm - \int_u^{u+h} f\,dm) = \operatorname{Av}_h f(v) - \operatorname{Av}_h f(u)$.
*Structural Commentary*: Difference quotients integrate to endpoint moving averages.
*Key Insight*: The interior contributions cancel after translation. □

**Theorem 3.4.9.** *Lebesgue Differentiability Theorem for Monotone Functions*
*If $F:[a,b]\to\mathbb{R}$ is monotone, then $F'$ exists a.e. on $[a,b]$.*

*Proof.* Assume $F$ is nondecreasing. Set $G_{\alpha,\beta} := \{\underline{D}F < \alpha < \beta < \overline{D}F\}$ for $\alpha,\beta\in\mathbb{Q}$, $\alpha<\beta$, and $E_\infty := \{\overline{D}F = +\infty\}$. Lemma 3.4.6 gives $m(G_{\alpha,\beta})=0$, and Lemma 3.4.5 gives $m(E_\infty)=0$.
Outside $E_\infty \cup \bigcup_{\alpha<\beta} G_{\alpha,\beta}$, $\underline{D}F = \overline{D}F \in \mathbb{R}$, so $F'$ exists. If $F$ is nonincreasing, apply the result to $-F$.
*Structural Commentary*: Null Dini gaps plus finite upper derivative give differentiability.
*Key Insight*: Monotone rise controls exceptional slopes. □

**Corollary 3.4.10.** *Lipschitz and BV Differentiability*
*Let $F:[a,b]\to\mathbb{R}$.*

*(i) If $F$ is Lipschitz, then $F'$ exists a.e. on $[a,b]$.*
*(ii) If $F\in BV([a,b])$, then $F'$ exists a.e. on $[a,b]$.*

*Proof.*

(i) If $F$ is $L$-Lipschitz, then $G_+(x) := F(x)+Lx$ and $G_-(x) := Lx - F(x)$ are nondecreasing.
Their derivatives exist a.e. by Theorem 3.4.9, so $F' = \frac{G_+' - G_-'}{2}$ a.e.

(ii) By Theorem 3.4.2, $F = G - H$ for nondecreasing $G, H$.
Their derivatives exist a.e. by Theorem 3.4.9, so $F' = G' - H'$ a.e.

*Structural Commentary*: Lipschitz and BV functions are reduced to finite combinations of monotone functions.
*Key Insight*: Almost-everywhere differentiability is stable under finite linear combinations. □

**Corollary 3.4.11.** *Absolute Continuity Implies BV Differentiability*
*If $F\in AC([a,b])$, then:*

*(i) $F\in BV([a,b])$.*
*(ii) $F'$ exists a.e. on $[a,b]$.*

*Proof.*

(i) Choose $\delta>0$ from absolute continuity with $\varepsilon=1$, and choose $a=x_0<\cdots<x_N=b$ with $x_j-x_{j-1}<\delta$. After inserting the $x_j$ into a partition $P$, the increments within each $[x_{j-1},x_j]$ have total length below $\delta$ and total $F$-variation below 1. Thus $V(F,P)\le N$ and $F\in BV([a,b])$.

(ii) By (i), $F\in BV([a,b])$; by Corollary 3.4.10 $\exists G\subseteq[a,b]$, $m([a,b]\setminus G)=0$ and $F'(x)\in\mathbb{R}$ $\forall x\in G$.

*Structural Commentary*: Absolute continuity bounds variation; BV decomposition then gives differentiability.
*Key Insight*: One absolute-continuity scale controls every partition after a finite interval decomposition. □

**Corollary 3.4.12.** *Derivative Bounds for Monotone and BV Functions*
*Let $F : [a,b] \to \mathbb{R}$.*

*(i) If $F$ is nondecreasing, then $F' \geq 0$ a.e.*
*(ii) If $F$ is nondecreasing, then $F' \in \mathscr{L}^1([a,b])$ and $\int_a^b F'\,dm \leq F(b) - F(a)$.*
*(iii) If $F \in BV([a,b])$, then $F'$ exists a.e.*
*(iv) If $F \in BV([a,b])$, then $F' \in \mathscr{L}^1([a,b])$.*

*Proof.*

(i) Every secant quotient of a nondecreasing function is nonnegative; hence $F' \geq 0$ a.e. by Theorem 3.4.9.
(ii) Extend $F$ by $F(b)$ on $(b, b+1]$, set $g_n(x) := n(F(x + \frac{1}{n}) - F(x))$. Then $g_n \geq 0$, $g_n \to F'$ a.e. on $(a,b)$.
By Proposition 3.4.8, $\int_a^b g_n\,dm = F(b) - n\int_a^{a+\frac{1}{n}} F\,dm \leq F(b) - F(a)$.
Fatou's lemma gives $\int_a^b F'\,dm \leq F(b) - F(a) < +\infty$.
(iii) Corollary 3.4.10(ii) gives $\exists E \subseteq [a,b]$ with $m([a,b] \setminus E) = 0$ and $F'(x) \in \mathbb{R}$ $\forall x \in E$.
(iv) Write $F = G - H$ with $G, H$ nondecreasing by Theorem 3.4.2. By (ii), $G', H' \in \mathscr{L}^1([a,b])$; on the full-measure set where $G', H'$ exist, $F' = G' - H'$, so $F' \in \mathscr{L}^1([a,b])$.

*Structural Commentary*: BV inherits finite derivative mass from monotone parts.
*Key Insight*: Fatou turns right-secant bounds into an integral bound for the derivative. □

## 3.5 Absolute Continuity and BV Decomposition

Absolute continuity is the part of bounded variation detected by integration. The remaining bounded-variation behavior is singular: it may be monotone and continuous, but its derivative vanishes almost everywhere.

**Proposition 3.5.1.** *Absolutely Continuous Variation Function*
*Let $F \in AC([a,b])$ and let $T_F(x) := V_a^x(F)$ be its total variation function.*

*(i) $T_F \in AC([a,b])$.*
*(ii) $F = (F + T_F) - T_F$ is a difference of two nondecreasing absolutely continuous functions.*

*Proof.*

(i) Let $\varepsilon > 0$ and choose $\delta > 0$ from absolute continuity of $F$ for the challenge $\frac{\varepsilon}{2}$.
Suppose $(c_k, d_k)_{k=1}^n$ are disjoint subintervals of $(a,b)$ with $\sum_k (d_k - c_k) < \delta$.
Choose partitions $P_k$ of $[c_k, d_k]$ such that $V_{c_k}^{d_k}(F) \leq V(F, P_k) + \frac{\varepsilon}{2n}$.
Let $\mathcal{I}$ be all subintervals of all $P_k$; then $\mathcal{I}$ is disjoint up to endpoints and $\sum_{I \in \mathcal{I}} |I| = \sum_k (d_k - c_k)$.
Hence absolute continuity of $F$ gives $\sum_k V(F, P_k) < \frac{\varepsilon}{2}$, thus $\sum_{k=1}^n |T_F(d_k) - T_F(c_k)| = \sum_{k=1}^n V_{c_k}^{d_k}(F) < \varepsilon$.
(ii) By (i), $T_F \in AC([a,b])$, hence $F + T_F \in AC([a,b])$.
For $x < y$, $T_F(y) - T_F(x) = V_x^y(F) \geq 0$ and $(F + T_F)(y) - (F + T_F)(x) = F(y) - F(x) + V_x^y(F) \geq 0$.
Thus $T_F$ and $F + T_F$ are nondecreasing, and $F = (F + T_F) - T_F$.

*Structural Commentary*: Absolute continuity controls local variation, not just endpoint displacement.
*Key Insight*: Partition each local variation, then apply one absolute-continuity test to all pieces. □

**Theorem 3.5.2.** *Absolute Continuity via Divided Differences*
*Let $F \in C([a,b])$, extend $F$ constantly by $F(b)$ on $(b, b+1]$. Let $\operatorname{Diff}_h F(x) := \frac{F(x+h) - F(x)}{h}$, $0 < h \leq 1$, $x \in [a,b]$.*
*Then $F \in AC([a,b])$ iff $\{\operatorname{Diff}_h F : 0 < h \leq 1\}$ is uniformly integrable:*
*$\forall \varepsilon > 0$ $\exists \delta > 0$ $\forall E \in \mathcal{L}^1$, $E \subseteq [a,b]$ and $m(E) < \delta$ imply $\sup_{0<h\leq 1} \int_E |\operatorname{Diff}_h F|\,dm < \varepsilon$.*

*Proof.*

($\Leftarrow$) Assume the divided differences are uniformly integrable. Given $\varepsilon > 0$, choose $\delta > 0$ for $\frac{\varepsilon}{2}$.
For disjoint $(c_k, d_k)$ with total length below $\delta$, set $E := \bigcup_k (c_k, d_k)$.
By Proposition 3.4.8, $\sum_k |\operatorname{Av}_h F(d_k) - \operatorname{Av}_h F(c_k)| \leq \int_E |\operatorname{Diff}_h F|\,dm < \frac{\varepsilon}{2}$.
Letting $h \downarrow 0$ gives $\sum_k |F(d_k) - F(c_k)| \leq \frac{\varepsilon}{2} < \varepsilon$.
($\Rightarrow$) Suppose $F \in AC([a,b])$. By Proposition 3.5.1, write $F = U - V$ with $U, V$ nondecreasing and absolutely continuous; since $|\operatorname{Diff}_h F| \leq \operatorname{Diff}_h U + \operatorname{Diff}_h V$, assume $F$ nondecreasing.
Let $\widetilde{F}$ be the constant extension to $[a, b+1]$.
Choose $\eta > 0$ for $\frac{\varepsilon}{2}$ in the absolute continuity of $\widetilde{F}$ and set $\delta := \frac{\eta}{2}$. If $m(E) < \delta$, take open $O \supseteq E \cap (a,b)$

with $O \cap (a,b) = \bigsqcup_j (\alpha_j, \beta_j)$ and $m(O \cap (a,b)) < \eta$; set $O_N := \bigcup_{j \le N} (\alpha_j, \beta_j)$.
For $0 < t \le h$, $\sum_{j \le N} ((\beta_j + t) - (\alpha_j + t)) < \eta$, so $\sum_{j \le N} (\widetilde{F}(\beta_j + t) - \widetilde{F}(\alpha_j + t)) < \frac{\varepsilon}{2}$.
Hence $\int_{O_N} \operatorname{Diff}_h F \, dm = \frac{1}{h} \int_0^h \sum_{j \le N} (\widetilde{F}(\beta_j + t) - \widetilde{F}(\alpha_j + t)) \, dt < \frac{\varepsilon}{2}$.
Let $N \to \infty$ and use $E \subseteq O$ to get $\sup_{0<h\le 1} \int_E \operatorname{Diff}_h F \, dm < \varepsilon$.

*Structural Commentary*: Absolute continuity is uniform small-set control for translated secants.
*Key Insight*: Moving averages turn endpoint increments into integrals of divided differences. □

**Theorem 3.5.3.** *Fundamental Theorem for Absolutely Continuous Functions*
*Let $F \in AC([a,b])$.*

*(i) $F'$ exists a.e. on $[a,b]$.*
*(ii) $F' \in \mathscr{L}^1([a,b])$.*
*(iii) $F(y) - F(x) = \int_x^y F' \, dm$ for $a \le x \le y \le b$.*

*Proof.*

(i) Corollary 3.4.11(ii) gives $\exists E \subseteq [a,b]$ with $m([a,b] \setminus E) = 0$ and $F'(x) \in \mathbb{R}$ $\forall x \in E$.
(ii) Corollary 3.4.11(i) gives $F \in BV([a,b])$, so Corollary 3.4.12(iv) gives $F' \in \mathscr{L}^1([a,b])$.
(iii) Set $f_n := \operatorname{Diff}_{\frac{1}{n}} F$. By Theorem 3.5.2, $(f_n)$ is uniformly integrable; by (i), $f_n \to F'$ a.e.
Fix $a \le x < y \le b$ and $\varepsilon > 0$. Choose $M > 0$ with $\int_{\{|F'|>M\}} |F'| \, dm < \frac{\varepsilon}{6}$ and then $\delta > 0$ such that $M\delta < \frac{\varepsilon}{6}$ and $m(A) < \delta$ implies $\sup_n \int_A |f_n| \, dm < \frac{\varepsilon}{3}$. By Egorov's theorem, $f_n \to F'$ uniformly on $[x,y] \setminus A$ for some measurable $A$ with $m(A) < \delta$. For large $n$, $\int_x^y |f_n - F'| \, dm < \varepsilon$.
Thus $\int_x^y f_n \, dm \to \int_x^y F' \, dm$. By Proposition 3.4.8, $\int_x^y f_n \, dm = \operatorname{Av}_{\frac{1}{n}} F(y) - \operatorname{Av}_{\frac{1}{n}} F(x) \to F(y) - F(x)$.

*Structural Commentary*: Uniform integrability is the condition that permits the discrete secant formula to pass to the derivative under the integral sign.
*Key Insight*: Uniform small-set control upgrades a.e. secant convergence to convergence of interval integrals. □

**Corollary 3.5.4.** *Indefinite Integral Characterization of Absolute Continuity*
*A function $F : [a,b] \to \mathbb{R}$ satisfies:*

*(i) $F \in AC([a,b])$ iff $\exists g \in \mathscr{L}^1([a,b])$ such that $F(x) = F(a) + \int_a^x g \, dm$ for $x \in [a,b]$.*
*(ii) Any such $g$ satisfies $g = F'$ a.e.*

*Proof.*

*(i)* ($\Rightarrow$) If $F \in AC([a,b])$, Theorem 3.5.3 gives $F' \in \mathscr{L}^1([a,b])$ and $F(x) = F(a) + \int_a^x F' \, dm$; take $g = F'$.
*(i)* ($\Leftarrow$) If $F(x) = F(a) + \int_a^x g \, dm$ with $g \in \mathscr{L}^1([a,b])$, Theorem 3.3.11 gives $F \in AC([a,b])$.
*(ii)* If $F(x) = F(a) + \int_a^x g \, dm$, then Theorem 3.3.11 gives $F' = g$ a.e.

*Structural Commentary*: The two directions are the fundamental theorem and the antiderivative theorem.
*Key Insight*: Absolute continuity is exactly representability by an $L^1$ density. □

**Proposition 3.5.5.** *Radon–Nikodym on an Interval Recovers Absolute Continuity*
*Let $\lambda$ be a finite measure on $\mathcal{B}([a,b])$ with $\lambda \ll m$, let $f := \frac{d\lambda}{dm}$, and set $F(x) := \lambda([a,x])$.*

*(i) $F(y) - F(x) = \int_x^y f \, dm$ for $a \le x \le y \le b$.*
*(ii) $F \in AC([a,b])$.*
*(iii) $F' = f$ a.e. on $[a,b]$.*

*Proof.* By Theorem 3.2.1, $f \in L^1(m)$, $\lambda(E) = \int_E f \, dm$ for $E \in \mathcal{B}([a,b])$. Since $\lambda \ll m$, $\lambda(\{x\}) = 0$ for $x \in [a,b]$.

(i) For $a \le x \le y \le b$, $\lambda(\{x\}) = 0$ gives $F(y) - F(x) = \lambda((x,y]) = \int_{(x,y]} f \, dm = \int_x^y f \, dm$.
(ii) Taking $x = a$ in (i), $F(y) = F(a) + \int_a^y f \, dm$ for $y \in [a,b]$; Corollary 3.5.4(i) gives $F \in AC([a,b])$.
(iii) Since $F = F(a) + \int_a^{(\cdot)} f \, dm$, Corollary 3.5.4(ii) gives $F' = f$ a.e. on $[a,b]$.

*Structural Commentary*: Interval RN representation differentiates cumulative mass.
*Key Insight*: No atoms, so interval increments equal density integrals. □

**Remark 3.5.6.** *Lebesgue–Stieltjes Integrals*
Let $g : \mathbb{R} \to \mathbb{R}$ be increasing and right-continuous, and let $\mu_g$ be its Borel–Stieltjes measure.
For $E \in \mathcal{B}(\mathbb{R})$ and bounded Borel $f : E \to \mathbb{R}$, define $\int_E f \, dg := \int_E f \, d\mu_g$.
If $J \subseteq \mathbb{R}$ is an interval, $\mu_g|_J \ll m|_J$, and $h := \frac{d\mu_g}{dm}$ on $J$, then $\int_E f \, dg = \int_E fh \, dm$ for every Borel $E \subseteq J$.

For $a < b$ with $[a,b] \subseteq J$, $F_a : [a,b] \to \mathbb{R}$, $F_a(x) := \mu_g([a,x])$, is absolutely continuous and satisfies $F_a' = h$ a.e. by Proposition 3.5.5. If $g|_J$ is absolutely continuous, then $h = g'$ a.e. on $J$.

**Example 3.5.7.** *Cantor–Lebesgue Function Is Not Absolutely Continuous*
Let $\varphi : [0,1] \to [0,1]$ be the Cantor–Lebesgue function: $\varphi$ is continuous, nondecreasing, $\varphi(0) = 0$, $\varphi(1) = 1$, and constant on every complementary interval of the Cantor set.
At stage $n$, write the remaining intervals as $I_{n,k} = [c_{n,k}, d_{n,k}]$, $1 \le k \le 2^n$, so $\sum_k |I_{n,k}| = (\frac{2}{3})^n$.

Since $\varphi$ is constant on the deleted gaps, $\sum_{k=1}^{2^n} (\varphi(d_{n,k}) - \varphi(c_{n,k})) = 1$.
By continuity, choose $c_{n,k} < \alpha_{n,k} < \beta_{n,k} < d_{n,k}$ such that $\sum_k (\beta_{n,k} - \alpha_{n,k}) < (\frac{2}{3})^n$ and $\sum_k (\varphi(\beta_{n,k}) - \varphi(\alpha_{n,k})) > \frac{1}{2}$.
Since $(\frac{2}{3})^n \to 0$, absolute continuity fails for $\varepsilon = \frac{1}{2}$.
*Key Insight*: The increase of $\varphi$ lies on intervals with arbitrarily small total length.

**Proposition 3.5.8.** *Lipschitz, Absolutely Continuous, and BV Classes*
*Let $[a,b]$ be nondegenerate, and let $\mathcal{F}_{\mathrm{Lip}}$ denote the Lipschitz functions on $[a,b]$.*

*(i)* $\mathcal{F}_{\mathrm{Lip}} \subsetneq AC([a,b]) \subsetneq BV([a,b])$.
*(ii) For* $\mathcal{C} \in \{\mathcal{F}_{\mathrm{Lip}}, AC([a,b]), BV([a,b])\}$, $F, G \in \mathcal{C}$ *and* $\alpha, \beta \in \mathbb{R}$ *imply* $\alpha F + \beta G \in \mathcal{C}$.
*(iii) If* $F \in \mathcal{C}$ *for some* $\mathcal{C} \in \{\mathcal{F}_{\mathrm{Lip}}, AC([a,b]), BV([a,b])\}$, *then* $T_F(x) := V_a^x(F)$ *lies in* $\mathcal{C}$.

*Proof.*

(i) If $F$ is $L$-Lipschitz, then $\sum_j |F(d_j) - F(c_j)| \le L \sum_j (d_j - c_j)$, so $F \in AC([a,b])$.
Corollary 3.4.11(i) gives $AC([a,b]) \subseteq BV([a,b])$.
Strictness: $x \mapsto \sqrt{\frac{x-a}{b-a}}$ lies in $AC([a,b]) \setminus \mathcal{F}_{\mathrm{Lip}}$, and $x \mapsto \varphi(\frac{x-a}{b-a})$, with $\varphi$ Cantor–Lebesgue, lies in $BV([a,b]) \setminus AC([a,b])$ because $\varphi$ is nondecreasing and its Cantor-stage intervals have total length $(\frac{2}{3})^n \to 0$ but total $\varphi$-increase 1.

(ii) If $F, G$ are Lipschitz with constants $L_F, L_G$, then $\alpha F + \beta G$ is Lipschitz with constant $|\alpha|\, L_F + |\beta|\, L_G$.
If $F, G \in AC([a,b])$, the interval criterion gives $\alpha F + \beta G \in AC([a,b])$.
If $F, G \in BV([a,b])$, then $V(\alpha F + \beta G, P) \le |\alpha|\, V(F,P) + |\beta|\, V(G,P)$, so $V_a^b(\alpha F + \beta G) < +\infty$.

(iii) If $F$ is $L$-Lipschitz, then $T_F(v) - T_F(u) = V_u^v(F) \le L(v-u)$.
If $F \in AC([a,b])$, Proposition 3.5.1(i) gives $T_F \in AC([a,b])$.
If $F \in BV([a,b])$, then $T_F$ is nondecreasing and $V_a^b(T_F) = T_F(b) - T_F(a) = V_a^b(F) < +\infty$.

*Structural Commentary*: $\mathcal{F}_{\mathrm{Lip}} \subsetneq AC \subsetneq BV$ is a strict control hierarchy.
*Key Insight*: $\sqrt{\cdot}$ breaks Lipschitz; Cantor mass breaks absolute continuity. □

**Proposition 3.5.9.** *Algebra and Safe Composition*
*Let $F, G \in AC([a,b])$.*

*(i)* *For* $\alpha, \beta \in \mathbb{R}$, $\alpha F + \beta G \in AC([a,b])$.
*(ii)* $FG \in AC([a,b])$.
*(iii) If* $\Phi : \mathbb{R} \to \mathbb{R}$ *is Lipschitz, then* $\Phi \circ G \in AC([a,b])$.
*(iv) If* $G$ *is strictly monotone and* $\Phi \in AC(J)$ *on a compact interval* $J \supseteq G([a,b])$, *then* $\Phi \circ G \in AC([a,b])$.

*Proof.*

(i) For disjoint $(c_k, d_k)$, $\sum_k |(\alpha F + \beta G)(d_k) - (\alpha F + \beta G)(c_k)| \le |\alpha| \sum_k |F(d_k) - F(c_k)| + |\beta| \sum_k |G(d_k) - G(c_k)|$.
Thus the right side is $< \varepsilon$ whenever $\sum_k (d_k - c_k) < \delta$ for a common AC scale of $F, G$, QED.

(ii) Let $|F| \le M_F$ and $|G| \le M_G$. For $\Delta F_k := F(d_k) - F(c_k)$ and $\Delta G_k := G(d_k) - G(c_k)$,
$|F(d_k)G(d_k) - F(c_k)G(c_k)| \le M_F\, |\Delta G_k| + M_G\, |\Delta F_k|$. Choose a common AC scale with $\sum_k |\Delta F_k| < \frac{\varepsilon}{2(1+M_G)}$ and $\sum_k |\Delta G_k| < \frac{\varepsilon}{2(1+M_F)}$; then $\sum_k |F(d_k)G(d_k) - F(c_k)G(c_k)| < \varepsilon$.

(iii) If $\Phi$ is $L$-Lipschitz, then $\sum_k |\Phi(G(d_k)) - \Phi(G(c_k))| \le L \sum_k |G(d_k) - G(c_k)|$.
Choose the AC scale of $G$ for $\frac{\varepsilon}{L}$ if $L > 0$; the case $L = 0$ is trivial.

(iv) Suppose $G$ is strictly increasing. Choose $\eta > 0$ from $AC(\Phi)$ for $\varepsilon$ and $\delta > 0$ from $AC(G)$ for $\eta$.
If $\sum_k (d_k - c_k) < \delta$, then $\sum_k (G(d_k) - G(c_k)) < \eta$.
Since $(G(c_k), G(d_k))$ are disjoint, $\sum_k |\Phi(G(d_k)) - \Phi(G(c_k))| < \varepsilon$.
If $G$ is strictly decreasing, replace $(G(c_k), G(d_k))$ by $(G(d_k), G(c_k))$.

*Structural Commentary*: AC is stable under algebra and controlled composition.
*Key Insight*: Lipschitz bounds sizes; monotonicity preserves disjoint intervals. □

**Example 3.5.10.** *Absolutely Continuous Composition Can Fail*
Define $f, g : [-1,1] \to \mathbb{R}$ by $f(x) := \operatorname{sgn}(x)\,|x|^{\frac{1}{3}}$, $g(0) := 0$, and $g(x) := x^2 \cos \frac{\pi}{2x}$ for $x \neq 0$. Then:

(i) $f, g \in AC([-1,1])$.
(ii) $f \circ g \notin BV([-1,1])$, hence $f \circ g \notin AC([-1,1])$.

*Proof.*

(i) Let $h(t) := \frac{1}{3}\,|t|^{-\frac{2}{3}}\,\mathbf{1}_{t\neq 0}$. Then $h \in L^1([-1,1])$ and $f(x) = \int_0^x h\,dm$, so $f \in AC([-1,1])$.
For $x \neq 0$, $g'(x) = 2x\cos\frac{\pi}{2x} + \frac{\pi}{2}\sin\frac{\pi}{2x}$, hence $|g'| \leq 2 + \frac{\pi}{2}$ off 0.
Together with $|g(x)| \leq x^2$, this gives $g$ Lipschitz, so $g \in AC([-1,1])$.

(ii) For $k \geq 1$, $g(\frac{1}{2k-1}) = 0$ and $g(\frac{1}{2k}) = \frac{(-1)^k}{4k^2}$. With $P_n := \{0\} \cup \{\frac{1}{j} : 1 \leq j \leq 2n\}$,
$V(f \circ g, P_n) \geq \sum_{k=1}^{n} \left| f(g(\frac{1}{2k})) - f(g(\frac{1}{2k-1})) \right| = 4^{-\frac{1}{3}} \sum_{k=1}^{n} k^{-\frac{2}{3}} \to +\infty$. Thus $f \circ g \notin BV([-1,1])$.

*Key Insight*: The cube root amplifies the inner oscillations until variation diverges. □

**Definition 3.5.11.** *Singular Bounded-Variation Function*
A function $S \in BV([a,b])$ is *singular* if $S' = 0$ a.e. on $[a,b]$.

**Theorem 3.5.12.** *Lebesgue Decomposition of a BV Function*
*Let $F \in BV([a,b])$. Define $G(x) := F(a) + \int_a^x F'\,dm$, $H(x) := F(x) - G(x)$.*

*(i) $G \in AC([a,b])$ and $G' = F'$ a.e.*
*(ii) $H \in BV([a,b])$, $H' = 0$ a.e., $H(a) = 0$, and $F = G + H$.*
*(iii) If $F = G_1 + H_1$ with $G_1 \in AC([a,b])$, $H_1' = 0$ a.e., and $H_1(a) = 0$, then $G_1 = G$ and $H_1 = H$.*

*Proof.*

(i) By Corollary 3.4.12, $F' \in \mathscr{L}^1([a,b])$. By Theorem 3.3.11, $G \in AC([a,b])$ and $G' = F'$ a.e.
(ii) Since $F, G \in BV([a,b])$, $H = F - G \in BV([a,b])$.
Moreover, $H' = F' - G' = 0$ a.e., $H(a) = 0$, and $F = G + H$.
(iii) If $F = G_1 + H_1$ is another normalized decomposition, then $G_1 - G = H - H_1$ is absolutely continuous, has derivative 0 a.e., and vanishes at $a$. By Theorem 3.5.3, $G_1 - G = 0$, hence $G_1 = G$ and $H_1 = H$.

*Structural Commentary*: A BV function splits into an $L^1$-density part and a derivative-invisible singular part.
*Key Insight*: Integrating the a.e. derivative separates the absolutely continuous and singular parts. □




Figure 215: Lip $\subsetneq AC \subsetneq BV$, and each $F \in BV$ decomposes as $F = G + H$ with $G \in AC$ and $H' = 0$ a.e.

**Corollary 3.5.13.** *Absolutely Continuous Singular Functions Are Constant*

*(i) If $F \in AC([a,b])$ and $F' = 0$ a.e., then $F$ is constant.*
*(ii) The Cantor–Lebesgue function is nonconstant and singular.*
*(iii) The Cantor–Lebesgue function is not absolutely continuous.*

*Proof.*

(i) By Theorem 3.5.3, $F(y) - F(x) = \int_x^y F'\,dm = 0$.
(ii) The function $\varphi$ is nondecreasing, hence in $BV([0,1])$, and $\varphi' = 0$ on the full-measure set $[0,1] \setminus P$ by Proposition 8.1.2. Since $\varphi(0) = 0$ and $\varphi(1) = 1$, it is nonconstant.
(iii) If $\varphi \in AC([0,1])$, then $\varphi' = 0$ a.e. by (ii), so (i) gives $\varphi$ constant, contradicting $\varphi(0) = 0 \neq 1 = \varphi(1)$.

*Structural Commentary*: The fundamental theorem makes absolute continuity exclude nonconstant singularity.
*Key Insight*: Derivative-invisible increase can occur in BV but not in absolute continuity. □

## 3.6 Nikodym Metric and Setwise Measure Limits

The Nikodym metric turns measurable sets modulo $\mu$-null symmetric difference into a complete metric space. This lets Baire category control setwise limits of absolutely continuous measures.

**Definition 3.6.1.** *Setwise Convergence and Uniform Absolute Continuity*
Let $(X, \mathcal{M})$ be a measurable space.

(i) For $A, B \in \mathcal{M}$, $A \triangle B := (A \setminus B) \cup (B \setminus A)$.
(ii) Finite measures $\nu_n$ *converge setwise* to $\nu : \mathcal{M} \to [0, +\infty]$ iff $\nu_n(E) \to \nu(E)$ for every $E \in \mathcal{M}$.
(iii) For $\nu_n \ll \mu$ for all $n$, $(\nu_n)$ is *uniformly absolutely continuous* w.r.t. $\mu$ iff
$\forall \varepsilon > 0\ \exists \delta > 0$: $\mu(E) < \delta \Rightarrow \sup_n \nu_n(E) < \varepsilon$ for all $E \in \mathcal{M}$.

**Lemma 3.6.2.** *Nikodym Symmetric-Difference Calculus*
*For $A, B, C \in \mathcal{M}$ and a finite measure $\nu$:*

*(i)* $A \triangle C \subseteq (A \triangle B) \cup (B \triangle C)$.
*(ii)* $|\nu(A) - \nu(B)| \le \nu(A \triangle B)$.

*Proof.*

(i) If $x \in A \triangle C$, then $\mathbf{1}_A(x) \neq \mathbf{1}_C(x)$. Hence $\mathbf{1}_A(x) \neq \mathbf{1}_B(x)$ or $\mathbf{1}_B(x) \neq \mathbf{1}_C(x)$, so $x \in (A \triangle B) \cup (B \triangle C)$.
(ii) Since $\nu(A) - \nu(B) = \nu(A \setminus B) - \nu(B \setminus A)$, $|\nu(A) - \nu(B)| \le \nu(A \setminus B) + \nu(B \setminus A) = \nu(A \triangle B)$.

*Structural Commentary*: Set inclusion gives the triangle inequality; disjoint differences give the mass bound.
*Key Insight*: Nikodym distance measures exactly the part where two measurable sets disagree. □

**Definition 3.6.3.** *Nikodym Metric Space*
Let $(X, \mathcal{M}, \mu)$ be a finite measure space.

(i) *Null equivalence*: $A \sim_\mu B \Leftrightarrow \mu(A \triangle B) = 0$.
(ii) *Quotient space*: $\mathcal{M}_\mu := \mathcal{M}/\!\sim_\mu$, with class notation $[A] \in \mathcal{M}_\mu$.
(iii) *Metric*: $\rho_\mu([A], [B]) := \mu(A \triangle B)$.

By Lemma 3.6.2 and Proposition 3.1.11, $\rho_\mu$ is a well-defined metric on $\mathcal{M}_\mu$.

**Proposition 3.6.4.** *Nikodym Pseudometric*
*Let $E \subseteq \mathbb{R}$ be Lebesgue measurable with $m(E) < \infty$, let $X$ be the collection of Lebesgue measurable subsets of $E$, and for $A, B \in X$ set $A \Delta B := (A \setminus B) \cup (B \setminus A)$ and $\rho(A, B) := m(A \Delta B)$. Define $A \sim_\rho B$ iff $\rho(A, B) = 0$.*

*(i) $\rho$ is a pseudometric on $X$.*
*(ii) $\widetilde{\rho}([A], [B]) := \rho(A, B)$ is a metric on $X/\!\sim_\rho$.*
*(iii) For all $A, B \in X$, $\rho(A, B) = \int_E |\mathbf{1}_A - \mathbf{1}_B|\, dm$.*

*Proof.*

(i) For $A, B, C \in X$, $A \Delta B \subseteq E$, $A \Delta A = \emptyset$, $A \Delta B = B \Delta A$, and $A \Delta C \subseteq (A \Delta B) \cup (B \Delta C)$.
Hence $0 \le \rho(A, B) \le m(E) < \infty$, $\rho(A, A) = 0$, $\rho(A, B) = \rho(B, A)$, and $\rho(A, C) \le \rho(A, B) + \rho(B, C)$.
(ii) By (i), $\rho$ is a pseudometric and $A \sim_\rho B \iff \rho(A, B) = 0$.
Proposition 3.1.11 gives that $\widetilde{\rho}([A], [B]) := \rho(A, B)$ is a metric on $X/\!\sim_\rho$.
(iii) For $t \in E$, $|\mathbf{1}_A(t) - \mathbf{1}_B(t)| = \mathbf{1}_{A \Delta B}(t)$; integrating gives $\int_E |\mathbf{1}_A - \mathbf{1}_B|\, dm = m(A \Delta B) = \rho(A, B)$.

*Structural Commentary*: Symmetric difference measures set disagreement.
*Key Insight*: Quotienting null disagreement gives definiteness. □

**Theorem 3.6.5.** *Completeness of the Nikodym Metric*
*If $(X, \mathcal{M}, \mu)$ is a finite measure space, then $(\mathcal{M}_\mu, \rho_\mu)$ is complete.*

*Proof.* Let $(A_n)$ be $\rho_\mu$-Cauchy. Choose $n_k \uparrow \infty$ with $\rho_\mu(A_{n_{k+1}}, A_{n_k}) < 2^{-k}$, and set $D_k := A_{n_{k+1}} \triangle A_{n_k}$.
Since $\mu(\bigcup_{k \ge N} D_k) \le \sum_{k \ge N} 2^{-k} \to 0$, $N_0 := \bigcap_N \bigcup_{k \ge N} D_k$ is null. Put $A := \bigcup_N \bigcap_{k \ge N} A_{n_k}$.
Then $A_{n_k} \triangle A \subseteq N_0 \cup \bigcup_{j \ge k} D_j$, so $\rho_\mu(A_{n_k}, A) \le \sum_{j \ge k} 2^{-j} \to 0$. Since $(A_n)$ is Cauchy, $\rho_\mu(A_n, A) \to 0$.
*Structural Commentary*: Summable jumps give an a.e. eventual set limit.
*Key Insight*: Summable symmetric differences confine infinitely changing membership to a null set. □

**Proposition 3.6.6.** *Absolutely Continuous Measures Are Nikodym-Continuous*
*Let $(X, \mathcal{M}, \mu)$ be finite, let $\nu$ be a finite measure with $\nu \ll \mu$, and define $\Phi_\nu([E]) := \nu(E)$.*

*(i) $\Phi_\nu : \mathcal{M}_\mu \to \mathbb{R}$ is well-defined.*
*(ii) $\Phi_\nu$ is uniformly continuous.*

*Proof.*

(i) If $A \sim_\mu B$, then $\mu(A \triangle B) = 0$, hence $\nu(A \triangle B) = 0$ and $\nu(A) = \nu(B)$.

(ii) By Lemma 3.6.2, $|\nu(A) - \nu(B)| \le \nu(A \triangle B)$. Given $\varepsilon > 0$, choose $\delta > 0$ by Lemma 3.1.14 such that $\mu(E) < \delta$ implies $\nu(E) < \varepsilon$. Thus $\rho_\mu(A, B) < \delta$ implies $|\Phi_\nu(A) - \Phi_\nu(B)| < \varepsilon$.

*Structural Commentary*: Null-set identification defines the map; absolute continuity controls its modulus.
*Key Insight*: Absolute continuity converts small Nikodym distance into small change of mass. □

**Proposition 3.6.7.** *Uniform Absolute Continuity and Radon–Nikodym Derivatives*
*Let $(X, \mathcal{M}, \mu)$ be finite, let $(\nu_n)$ be finite measures with $\nu_n \ll \mu$, and set $f_n = \frac{d\nu_n}{d\mu}$. The following are equivalent.*

*(i) $(\nu_n)$ is uniformly absolutely continuous with respect to $\mu$.*
*(ii) The functions $\Phi_{\nu_n} : \mathcal{M}_\mu \to \mathbb{R}$ are uniformly equicontinuous.*
*(iii) The functions $\Phi_{\nu_n}$ are equicontinuous at $\emptyset$.*
*(iv) $\{f_n : n \ge 1\}$ is uniformly integrable: $\forall \varepsilon > 0\ \exists \delta > 0\ \forall E \in \mathcal{M}$, $\mu(E) < \delta \Rightarrow \sup_n \int_E f_n \, d\mu < \varepsilon$.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* Given $\varepsilon > 0$, choose $\delta > 0$ with $\mu(E) < \delta \Rightarrow \sup_n \nu_n(E) < \varepsilon$.
If $\rho_\mu(A, B) = \mu(A \triangle B) < \delta$, then $|\Phi_{\nu_n}(A) - \Phi_{\nu_n}(B)| = |\nu_n(A) - \nu_n(B)| \le \nu_n(A \triangle B) < \varepsilon$ uniformly in $n$.

*(ii)* $\Rightarrow$ *(iii)* Given $\varepsilon > 0$, uniform equicontinuity gives $\delta > 0$ such that $\rho_\mu(A, B) < \delta \Rightarrow \sup_n |\Phi_{\nu_n}(A) - \Phi_{\nu_n}(B)| < \varepsilon$. With $B = \emptyset$, this is equicontinuity at $\emptyset$.

*(iii)* $\Rightarrow$ *(i)* Given $\varepsilon > 0$, choose $\delta > 0$ such that $\rho_\mu(E, \emptyset) < \delta \Rightarrow \sup_n |\Phi_{\nu_n}(E) - \Phi_{\nu_n}(\emptyset)| < \varepsilon$.
Since $\rho_\mu(E, \emptyset) = \mu(E)$ and $\Phi_{\nu_n}(\emptyset) = 0$, $\mu(E) < \delta \Rightarrow \sup_n \nu_n(E) < \varepsilon$.

*(i)* $\Leftrightarrow$ *(iv)* For $E \in \mathcal{M}$, $\nu_n(E) = \int_E f_n \, d\mu$. Thus $\sup_n \nu_n(E) < \varepsilon \Leftrightarrow \sup_n \int_E f_n \, d\mu < \varepsilon$ for every small-$\mu$ set $E$.

*Structural Commentary*: Uniform AC of $\nu_n$ equals uniform integrability of $f_n$.
*Key Insight*: Radon–Nikodym turns set bounds into integral bounds. □

**Theorem 3.6.8.** *Setwise Limits under Uniform Absolute Continuity*
*Let $(X, \mathcal{M}, \mu)$ be finite, let $\nu_n$ be finite measures with $\nu_n \ll_{\mathrm{u}} \mu$, $\sup_n \nu_n(X) < +\infty$, and $\nu_n(E) \to \nu(E)$ $\forall E \in \mathcal{M}$.*

*(i) $\nu$ is a finite measure.*
*(ii) $\nu \ll \mu$.*

*Proof.*

(i) Since $\nu(X) = \lim_n \nu_n(X) \le \sup_n \nu_n(X)$, $\nu$ is finite-valued. Finite additivity passes to the pointwise limit.
For pairwise disjoint $(E_k)$, set $E := \bigcup_k E_k$ and $R_N := \bigcup_{k>N} E_k$.
Given $\varepsilon > 0$, uniform absolute continuity and $\mu(R_N) \downarrow 0$ give $\nu_n(R_N) < \varepsilon$ $\forall n$ and all sufficiently large $N$, hence $\nu(R_N) \le \varepsilon$. Therefore $\nu(E) = \sum_{k=1}^{N} \nu(E_k) + \nu(R_N) \to \sum_k \nu(E_k)$.

(ii) If $\mu(A) = 0$, then for every $\varepsilon > 0$, uniform absolute continuity gives $\sup_n \nu_n(A) < \varepsilon$.
Passing to the setwise limit, $\nu(A) \le \varepsilon$ $\forall \varepsilon > 0$, hence $\nu(A) = 0$.

*Structural Commentary*: Uniform tail control yields countable additivity and preserves the null ideal.
*Key Insight*: Uniform absolute continuity makes the tail of any disjoint union uniformly small. □

**Lemma 3.6.9.** *One-Point Equicontinuity Spreads*
*Let $(X, \mathcal{M}, \mu)$ be finite, and let $(\nu_n)$ be finite measures with $\nu_n \ll \mu$.*
*If the maps $\Phi_{\nu_n}$ are equicontinuous at one point $E_0 \in \mathcal{M}_\mu$, then they are uniformly equicontinuous on $\mathcal{M}_\mu$.*

*Proof.* Fix $\varepsilon > 0$ and choose $\delta > 0$ so that $\rho_\mu(E, E_0) < \delta$ implies $\sup_n |\nu_n(E) - \nu_n(E_0)| < \frac{\varepsilon}{2}$.
If $\mu(C) < \delta$, then both $E_0 \setminus C$ and $E_0 \cup (C \setminus E_0)$ are within $\delta$ of $E_0$, so $\nu_n(C \cap E_0) < \frac{\varepsilon}{2}$, $\nu_n(C \setminus E_0) < \frac{\varepsilon}{2}$ $\forall n$.
Hence $\sup_n \nu_n(C) < \varepsilon$. Taking $C = A \triangle B$ gives $\sup_n |\nu_n(A) - \nu_n(B)| < \varepsilon$ whenever $\rho_\mu(A, B) < \delta$.

*Structural Commentary*: Split a small set into its portions inside and outside the equicontinuity base point.
*Key Insight*: Smallness near one set controls both the part inside it and the part outside it. □

**Method 3.6.10.** *Baire Mechanism for Vitali–Hahn–Saks*
Regard each absolutely continuous measure $\nu_n$ as $\Phi_{\nu_n} : \mathcal{M}_\mu \to \mathbb{R}$, $\Phi_{\nu_n}(E) := \nu_n(E)$.

(i) *Translate*: Setwise convergence becomes pointwise convergence of $(\Phi_{\nu_n})$.
(ii) *Apply Baire*: Completeness of $\mathcal{M}_\mu$ supplies a point of equicontinuity.

(iii) *Spread*: Lemma 3.6.9 upgrades one-point equicontinuity to uniform equicontinuity.
(iv) *Return to measures*: Proposition 3.6.7 converts uniform equicontinuity into uniform absolute continuity.




Figure 216: Vitali–Hahn–Saks mechanism: setwise convergence becomes pointwise convergence on $\mathcal{M}_\mu$; Baire gives uniform absolute continuity.

**Theorem 3.6.11.** *Vitali–Hahn–Saks Theorem*
*Let $(X, \mathcal{M}, \mu)$ be finite, and let $(\nu_n)$ be finite measures with $\nu_n \ll \mu$, $\sup_n \nu_n(X) < +\infty$, and $\nu_n \to \nu$ setwise.*

*(i) $(\nu_n)$ is uniformly absolutely continuous with respect to $\mu$.*
*(ii) $\nu$ is a finite measure.*
*(iii) $\nu \ll \mu$.*

*Proof.*

(i) Set $\Phi_n([E]) := \nu_n(E)$. By Proposition 3.6.6, each $\Phi_n$ is continuous on complete $\mathcal{M}_\mu$.
If $A \sim_\mu B$, then $\nu_n(A) = \nu_n(B)\ \forall n$, hence $\nu(A) = \nu(B)$.
Thus $\Phi([E]) := \nu(E)$ is well-defined and $\Phi_n \to \Phi$ pointwise. Since $|\Phi_n([E])| \le \sup_n \nu_n(X)$ and $\nu_n(E) \to \nu(E)$, $\Phi$ is real-valued.
Theorem 4.3.12 gives an equicontinuity point $E_0 \in \mathcal{M}_\mu$ for $(\Phi_n)$.
Lemma 3.6.9 gives uniform equicontinuity on $\mathcal{M}_\mu$, and Proposition 3.6.7 gives $\nu_n \ll_{\mathrm{u}} \mu$.
(ii) By (i), $\nu_n \ll_{\mathrm{u}} \mu$; Theorem 3.6.8(i) gives that $\nu$ is a finite measure.
(iii) By (i), $\nu_n \ll_{\mathrm{u}} \mu$; Theorem 3.6.8(ii) gives $\nu \ll \mu$.

*Structural Commentary*: Baire gives one equicontinuity point; Nikodym calculus spreads it.
*Key Insight*: Setwise convergence plus bounded total mass forces uniform null-set control. □

**Corollary 3.6.12.** *Nikodym Bounded Setwise Limit Theorem*
*If finite measures $(\nu_n)$ on $(X, \mathcal{M})$ converge setwise to $\nu$ and $\sup_n \nu_n(X) < +\infty$, then $\nu$ is finite.*

*Proof.* Define $\mu(E) := \sum_{n=1}^{\infty} 2^{-n}\nu_n(E)$. Then $\mu$ is a finite measure by countable additivity of nonnegative series, and $\nu_n \ll \mu\ \forall n$ because $2^{-n}\nu_n \le \mu$. Thus $\nu$ is a finite measure by Theorem 3.6.11(ii).
*Structural Commentary*: A summable mixture places the sequence under one finite dominating measure.
*Key Insight*: A summable mixture builds one finite measure that dominates the whole sequence. □

# 4 $L^p$ Spaces and Modes of Convergence

## 4.1 $L^p$ Spaces

Lebesgue spaces turn integrability into normed linear geometry. The essential point is that an integral norm cannot distinguish functions that differ only on null sets, so the actual vectors are almost-everywhere equivalence classes. Their bounded linear functionals are themselves represented by integration against functions in the conjugate space.

**Definition 4.1.1.** *Norms and Seminorms*
Let $\mathcal{V}$ be a vector space over $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$.

(i) A *seminorm* is $N : \mathcal{V} \to [0, \infty)$ with $N(\alpha v) = |\alpha|\, N(v)$ and $N(u + v) \le N(u) + N(v)\ \forall u, v \in \mathcal{V}$, $\alpha \in \mathbb{F}$.
(ii) A *norm* is a seminorm with $N(v) = 0 \iff v = 0$. A norm $N$ induces $d_N(u, v) := N(u - v)$.

**Example 4.1.2.** *Norms and Seminorms Before Quotienting*

(i) On $\mathbb{F}^n$, $\|u\|_p := (\sum_{j=1}^{n} |u_j|^p)^{\frac{1}{p}}$ for $1 \le p < \infty$ and $\|u\|_\infty := \max_{1 \le j \le n} |u_j|$ are norms.
Minkowski gives $\|u + v\|_p \le \|u\|_p + \|v\|_p$.

(ii) On sequences, $\ell^p := \{a : \sum_n |a_n|^p < +\infty\}$ with $\|a\|_p := (\sum_n |a_n|^p)^{\frac{1}{p}}$.
$\ell^\infty := \{a : \sup_n |a_n| < +\infty\}$ with $\|a\|_\infty := \sup_n |a_n|$.

(iii) On $B(X)$, $\|f\|_\infty := \sup_{x\in X} |f(x)|$ is a norm. On $C[0,1]$, $N_0(f) := |f(0)|$ and $N_1(f) := \sup_{0\le x\le \frac{1}{2}} |f(x)|$ are seminorms. On $C^1[a,b]$, $N_D(f) := \sup_{a\le x\le b} |f'(x)|$ is a seminorm with kernel the constant functions.

*Key Insight*: Seminorms become norms after quotienting out their zero directions.

**Remark 4.1.3.** *Integral Norm on Continuous Functions*
On $C[0,1]$, $N(f) := \int_0^1 |f(x)|\,dx$ is a norm: if $f(x_0) \neq 0$, continuity gives $N(f) > 0$. It is not complete, since continuous ramps can be Cauchy in $N$ and converge in $L^1$ to a discontinuous step function.

**Definition 4.1.4.** *$L^p$ Spaces and Mean Convergence*
Let $(X, \mathcal{M}, \mu)$ be a measure space.

(i) For $1 \le p < \infty$, the *$p$-integrable class* is $\mathscr{L}^p(\mu) := \{f : X \to \mathbb{F} : f \text{ measurable}, \int_X |f|^p\,d\mu < +\infty\}$.
(ii) The *$L^p$ space* and *$L^p$ norm* are $L^p(\mu) := \mathscr{L}^p(\mu)/=_{\text{a.e.}}$ and $\|f\|_p := (\int_X |f|^p\,d\mu)^{\frac{1}{p}}$.
(iii) *$L^p$ convergence*, or *mean convergence of order $p$*, means $f_n \to f \iff \|f_n - f\|_p \to 0$.
*$L^p$-Cauchy* means $\|f_m - f_n\|_p \to 0$ as $m, n \to \infty$.
(iv) *Essentially bounded* means $|f| \le M$ a.e. for some $M < +\infty$; $L^\infty(\mu)$ is the a.e.-quotient of such functions, with *essential sup norm* $\|f\|_\infty := \inf\{M \ge 0 : |f| \le M \text{ a.e.}\} = \inf_{\mu(N)=0} \sup_{x\in X\setminus N} |f(x)|$.

**Proposition 4.1.5.** *$L^1$ as a Quotient of a Seminorm*
*On $\mathscr{L}^1(\mu)$ set $N_\mu(f) := \int_X |f|\,d\mu$.*

*(i) $N_\mu$ is a seminorm.*
*(ii) $N_\mu(f) = 0 \iff f = 0$ a.e.*
*(iii) $L^1(\mu) := \mathscr{L}^1(\mu)/=_{\text{a.e.}}$ is normed by $\|[f]\|_1 := \int_X |f|\,d\mu$; write $f$ for $[f]$.*

*Proof.*

(i) If $f, g \in \mathscr{L}^1(\mu)$ and $a, b \in \mathbb{R}$, then $|af + bg| \le |a|\,|f| + |b|\,|g| \in \mathscr{L}^1(\mu)$, so $\mathscr{L}^1(\mu)$ is linear. Also $N_\mu(af) = |a|\,N_\mu(f)$ and $N_\mu(f+g) \le N_\mu(f) + N_\mu(g)$.
(ii) ($\Rightarrow$) If $N_\mu(f) = 0$, set $E_n := \{|f| \ge \frac{1}{n}\}$. Then $\frac{1}{n}\mu(E_n) \le \int_{E_n} |f|\,d\mu \le N_\mu(f) = 0$, $\{f \neq 0\} = \bigcup_n E_n$ null.
(ii) ($\Leftarrow$) If $f = 0$ a.e., then $|f| = 0$ a.e., hence $N_\mu(f) = 0$.
(iii) If $f = g$ a.e., then $|f| = |g|$ a.e. and $\int_X |f|\,d\mu = \int_X |g|\,d\mu$, so $\|[f]\|_1$ is well defined. The seminorm axioms descend from (i), and $\|[f]\|_1 = 0 \Rightarrow f = 0$ a.e. by (ii), hence $[f] = [0]$.

*Structural Commentary*: The seminorm kernel is exactly the a.e.-zero functions.
*Key Insight*: Quotient by null differences before calling the integral a norm. □

**Definition 4.1.6.** *Complex Integrals and $L^2$*

(i) For $f = u + iv$, *complex measurability* means measurability of $u, v$.
(ii) On $E \in \mathcal{M}$, *complex integrability* means $u, v \in \mathscr{L}^1(\mu)$ on $E$; set $\int_E f\,d\mu := \int_E u\,d\mu + i\int_E v\,d\mu$.
(iii) $L^2(\mu)$: complex measurable $f$ satisfying $\int_X |f|^2\,d\mu < +\infty$, modulo a.e. equality; $\|f\|_2^2 := \int_X |f|^2\,d\mu$.
(iv) The *$L^2$ inner product* is $\langle f, g\rangle_{L^2} := \int_X f\overline{g}\,d\mu$; it induces $\|f\|_2$.

**Proposition 4.1.7.** *Complex $L^1$ and Dominated Convergence*
*The following statements hold.*

*(i) For complex measurable $f$ on $E$, $f$ is integrable $\iff |f| \in \mathscr{L}^1(\mu)$ on $E$; then $\left|\int_E f\,d\mu\right| \le \int_E |f|\,d\mu$.*
*(ii) If $f_n \to f$ a.e. on $E$ and $|f_n| \le g \in \mathscr{L}^1(\mu)$, then $f$ is integrable and $\int_E f\,d\mu = \lim_n \int_E f_n\,d\mu$.*

*Proof.*

(i) ($\Rightarrow$) Write $f = u + iv$. If $u, v \in \mathscr{L}^1$, then $|f| \le |u| + |v|$, so $|f| \in \mathscr{L}^1$.
(i) ($\Leftarrow$) If $|f| \in \mathscr{L}^1$, then $|u|, |v| \le |f|$, so $u, v \in \mathscr{L}^1$ and $f$ is integrable.
If $\int_E f\,d\mu = re^{i\theta}$ and $h = e^{-i\theta}f$, then $\left|\int_E f\,d\mu\right| = r = \operatorname{Re}\int_E h\,d\mu = \int_E \operatorname{Re} h\,d\mu \le \int_E |h|\,d\mu = \int_E |f|\,d\mu$.
(ii) Write $f_n = u_n + iv_n$ and $f = u + iv$. Since $|u_n|, |v_n| \le g$ and $u_n \to u$, $v_n \to v$ a.e., real DCT gives $\int_E u_n\,d\mu \to \int_E u\,d\mu$ and $\int_E v_n\,d\mu \to \int_E v\,d\mu$; hence the complex identity.

*Structural Commentary*: Complex integration is two real integrals.
*Key Insight*: Rotate the integral to make its value real and nonnegative before applying the real inequality. □

**Theorem 4.1.8.** *$L^2$ Geometry*
*For $f, g \in L^2(\mu)$, the following hold.*

*(i) Schwarz: $fg \in L^1(\mu)$ and $\int_X |fg|\, d\mu \le \|f\|_2 \|g\|_2$.*
*(ii) Minkowski: $f+g \in L^2(\mu)$ and $\|f+g\|_2 \le \|f\|_2 + \|g\|_2$.*
*(iii) Completeness: modulo equality a.e., $(L^2(\mu), \|\cdot\|_2)$ is complete.*

*Proof.*

(i) Since $2|fg| \le |f|^2 + |g|^2$, $fg \in L^1(\mu)$.
For $\lambda \in \mathbb{R}$, $0 \le Q(\lambda) := \int_X (|f| + \lambda|g|)^2\, d\mu = \|f\|_2^2 + 2\lambda \int_X |fg|\, d\mu + \lambda^2 \|g\|_2^2$; hence $\Delta(Q) \le 0$.
(ii) By (i), $\|f+g\|_2^2 \le \int_X (|f| + |g|)^2\, d\mu \le (\|f\|_2 + \|g\|_2)^2$.
(iii) Let $(f_n)$ be Cauchy. Choose $n_k \uparrow \infty$ with $\|f_{n_{k+1}} - f_{n_k}\|_2 \le 2^{-k}$, and set $h_N := \sum_{k<N} |f_{n_{k+1}} - f_{n_k}|$.
Then $\|h_N\|_2 \le \sum_{k<N} 2^{-k}$, so Fatou gives $h := \lim_N h_N \in L^2$ and $f_{n_k} \to f$ a.e. for some $f$.
For fixed $n$, Fatou gives $\|f - f_n\|_2 \le \liminf_k \|f_{n_k} - f_n\|_2$; the Cauchy property sends the right side to 0.

*Structural Commentary*: Schwarz and Minkowski supply geometry; rapid subsequences prove completeness.
*Key Insight*: Positivity of the integral transfers Euclidean quadratic estimates to function classes. □

**Example 4.1.9.** *Mazur Homeomorphism Between $L^1$ and $L^2$*
Let $E \subseteq \mathbb{R}$ be Lebesgue measurable. Define $\Phi(f) := \operatorname{sgn}(f)\,|f|^{\frac{1}{2}}$ on $L^1(E)$ and $\Psi(g) := \operatorname{sgn}(g)\,|g|^2$ on $L^2(E)$. Then $\Phi : L^1(E) \to L^2(E)$ is a homeomorphism with inverse $\Psi : L^2(E) \to L^1(E)$.

*Proof.* $|\Phi(f)|^2 = |f|$ and $|\Psi(g)| = |g|^2$, so the maps are well-defined; pointwise $\Psi\Phi = \mathrm{id}_{L^1}$ and $\Phi\Psi = \mathrm{id}_{L^2}$.
For $a, b \in \mathbb{R}$, $\left|\operatorname{sgn}(a)\,|a|^{\frac{1}{2}} - \operatorname{sgn}(b)\,|b|^{\frac{1}{2}}\right|^2 \le 2|a-b|$, hence $\|\Phi(f) - \Phi(h)\|_2^2 \le 2\|f-h\|_1$.
Also $\left|\operatorname{sgn}(a)\,|a|^2 - \operatorname{sgn}(b)\,|b|^2\right| \le 2|a-b|\,(|a|+|b|)$, so $\|\Psi(g) - \Psi(k)\|_1 \le 2\|g-k\|_2\,(\|g\|_2 + \|k\|_2)$ by Cauchy–Schwarz. Thus $\Phi$ and $\Psi$ are continuous.
*Key Insight*: Nonlinear power maps can homeomorphically identify different $L^p$ norm topologies. □

**Definition 4.1.10.** *Conjugate Indices*
For $p, q \in (1, \infty)$, $p$ and $q$ are *conjugate indices* if $p^{-1} + q^{-1} = 1$, equivalently $p + q = pq$.
The unique self-conjugate index is $p = q = 2$.

**Lemma 4.1.11.** *Young's Inequality*
*If $A, B \ge 0$, $1 < p < \infty$, and $p^{-1} + q^{-1} = 1$, then:*

*(i) $AB \le \frac{A^p}{p} + \frac{B^q}{q}$;*
*(ii) equality holds iff $A^p = B^q$.*

*Proof.*

(i) If $A, B > 0$, set $\alpha := \frac{1}{p}$ and $t := \frac{A^p}{B^q}$. Since $q^{-1} = 1 - \alpha$, weighted AM–GM gives $t^\alpha \le \alpha t + 1 - \alpha$.
Multiplying by $B^q$ yields $AB \le \frac{A^p}{p} + \frac{B^q}{q}$. If $AB = 0$, the inequality is immediate.
(ii) For $A, B > 0$, equality in weighted AM–GM is $t = 1$, i.e. $A^p = B^q$.
If $AB = 0$, equality in (i) holds iff $A = B = 0$, again equivalent to $A^p = B^q$.

*Structural Commentary*: Young's inequality is weighted AM–GM in conjugate exponents.
*Key Insight*: Rescale to $A^p$ and $B^q$, then use convexity. □

**Theorem 4.1.12.** *Hölder and Minkowski for Sequence Spaces*
*Let $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$.*

*(i) If $p, q > 1$ and $p^{-1} + q^{-1} = 1$, then $\mathbf{x} \in \ell^p(\mathbb{F})$, $\mathbf{y} \in \ell^q(\mathbb{F})$ imply $\sum_{n\ge1} |x_n y_n| \le \|\mathbf{x}\|_p \|\mathbf{y}\|_q$.*
*(ii) If $1 \le p < \infty$ and $\mathbf{x}, \mathbf{y} \in \ell^p(\mathbb{F})$, then $\mathbf{x}+\mathbf{y} \in \ell^p(\mathbb{F})$ and $\|\mathbf{x}+\mathbf{y}\|_p \le \|\mathbf{x}\|_p + \|\mathbf{y}\|_p$.*

*Proof.*

(i) If $\|\mathbf{x}\|_p \|\mathbf{y}\|_q = 0$, the claim is immediate.
Otherwise set $X := \|\mathbf{x}\|_p$, $Y := \|\mathbf{y}\|_q$, $u_n := |x_n|/X$, and $v_n := |y_n|/Y$. Young's inequality gives $u_n v_n \le u_n^p/p + v_n^q/q$; summing over $1 \le n \le N$ and letting $N \to \infty$ gives $\sum_{n\ge1} |x_n y_n| \le XY$.
(ii) For $p = 1$, apply $|x_n + y_n| \le |x_n| + |y_n|$ and sum. For $p > 1$, put $s_N := (\sum_{n=1}^N |x_n + y_n|^p)^{1/p}$. If $s_N = 0$, the bound is trivial; if $s_N > 0$, then $\sum_{n=1}^N |x_n + y_n|^p \le \sum_{n=1}^N (|x_n| + |y_n|)\,|x_n + y_n|^{p-1} \le (\|\mathbf{x}\|_p + \|\mathbf{y}\|_p) s_N^{p-1}$ by (i), so $s_N \le \|\mathbf{x}\|_p + \|\mathbf{y}\|_p$. Let $N \to \infty$.

*Structural Commentary*: Hölder controls products; Minkowski controls sums.
*Key Insight*: The metric property of $\ell^p$ is exactly Minkowski's inequality. □

**Example 4.1.13.** *Function and Sequence Metric Spaces*
Let $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$.

(i) On $C_{\mathbb{F}}[a,b]$, $d_\infty(f,g) := \sup_{t\in[a,b]} |f(t)-g(t)|$ and $d_1(f,g) := \int_a^b |f(t)-g(t)|\,dt$ are metrics. Definiteness of $d_1$ uses continuity.
(ii) On $\ell^\infty(\mathbb{F})$, $d_\infty(\mathbf{x},\mathbf{y}) := \sup_{n\geq 1} |x_n - y_n|$ is a metric.
(iii) For $1 \leq p < \infty$, $d_p(\mathbf{x},\mathbf{y}) := \|\mathbf{x}-\mathbf{y}\|_p = (\sum_{n\geq 1} |x_n - y_n|^p)^{1/p}$ is a metric on $\ell^p(\mathbb{F})$ by Theorem 4.1.12.

*Structural Commentary*: Norm estimates produce metrics on function and sequence spaces.
*Key Insight*: Infinite-dimensional examples require estimates, not just coordinate formulas.

**Theorem 4.1.14.** *Hölder's Inequality*
*Let $1 < p < \infty$, $p^{-1} + q^{-1} = 1$, $f \in L^p(\mu)$, and $g \in L^q(\mu)$.*

*(i)* $fg \in L^1(\mu)$.
*(ii)* $\|fg\|_1 \leq \|f\|_p \|g\|_q$.

*Proof.* If $\|f\|_p \|g\|_q = 0$, then $f = 0$ a.e. or $g = 0$ a.e., so both claims hold.
Assume $\|f\|_p \|g\|_q > 0$ and set $A := \frac{|f|}{\|f\|_p}$, $B := \frac{|g|}{\|g\|_q}$. Young gives $\frac{|fg|}{\|f\|_p \|g\|_q} \leq \frac{A^p}{p} + \frac{B^q}{q}$.

(i) Since $\int A^p\,d\mu = \int B^q\,d\mu = 1$, the majorant $\frac{A^p}{p} + \frac{B^q}{q}$ lies in $L^1(\mu)$; hence $fg \in L^1(\mu)$.
(ii) Integrating the displayed inequality gives $\frac{\|fg\|_1}{\|f\|_p\|g\|_q} \leq \frac{1}{p}\int A^p\,d\mu + \frac{1}{q}\int B^q\,d\mu = p^{-1} + q^{-1} = 1$.

*Structural Commentary*: Normalize both factors to unit $L^p$ and $L^q$ size.
*Key Insight*: Products are controlled by conjugate powers. □

**Corollary 4.1.15.** *Cauchy–Bunyakovsky–Schwarz in $L^2$*
*If $f, g \in L^2(\mu)$, then:*

*(i)* $fg \in L^1(\mu)$;
*(ii)* $\left|\int_X fg\,d\mu\right| \leq \int_X |fg|\,d\mu \leq \|f\|_2 \|g\|_2$.

*Proof.*

(i) Since $2^{-1} + 2^{-1} = 1$, Theorem 4.1.14(i) gives $fg \in L^1(\mu)$.
(ii) Theorem 4.1.14(ii) gives $\int_X |fg|\,d\mu = \|fg\|_1 \leq \|f\|_2 \|g\|_2$, and $\left|\int_X fg\,d\mu\right| \leq \int_X |fg|\,d\mu$.

*Structural Commentary*: Hölder at $p = q = 2$ is Cauchy–Schwarz.
*Key Insight*: Pair two $L^2$ factors through $2^{-1} + 2^{-1} = 1$. □

**Theorem 4.1.16.** *Minkowski's Inequality*
*If $1 \leq p < \infty$ and $f, h \in L^p(\mu)$, then:*

*(i)* $f + h \in L^p(\mu)$;
*(ii)* $\|f+h\|_p \leq \|f\|_p + \|h\|_p$.

*Proof.* For $p = 1$, integrate $|f+h| \leq |f| + |h|$. Let $p > 1$ and $q := \frac{p}{p-1}$.

(i) Since $|f+h|^p \leq 2^{p-1}(|f|^p + |h|^p)$, $f + h \in L^p(\mu)$. Also $|f+h|^{p-1} \in L^q(\mu)$ because $(p-1)q = p$.
(ii) Pointwise, $|f+h|^p \leq (|f| + |h|)\,|f+h|^{p-1}$.
By Hölder, $\|f+h\|_p^p \leq (\|f\|_p + \|h\|_p)\left\||f+h|^{p-1}\right\|_q = (\|f\|_p + \|h\|_p)\,\|f+h\|_p^{p-1}$.
If $\|f+h\|_p > 0$, divide by $\|f+h\|_p^{p-1}$; otherwise the claim is trivial.

*Structural Commentary*: Hölder turns the pointwise triangle inequality into a norm triangle inequality.
*Key Insight*: Minkowski is exactly the closure and triangle estimate for $L^p$. □

**Theorem 4.1.17.** *$L^p$ Is Normed*
*For $1 \leq p < \infty$, $(L^p(\mu), \|\cdot\|_p)$ is a normed linear space.*

*Proof.*

(i) If $f = f'$ and $g = g'$ a.e., then $\alpha f + \beta g = \alpha f' + \beta g'$ a.e..
Hence pointwise operations descend to $L^p(\mu)$. Minkowski gives $\alpha f + \beta g \in L^p(\mu)$.
(ii) $\|\alpha f\|_p^p = \int_X |\alpha f|^p\,d\mu = |\alpha|^p \|f\|_p^p$, so $\|\alpha f\|_p = |\alpha|\,\|f\|_p$.
(iii) $\|f\|_p = 0 \Rightarrow \int_X |f|^p\,d\mu = 0 \Rightarrow f = 0$ a.e.; hence $[f] = 0$ in $L^p(\mu)$.
(iv) Minkowski gives $\|f+g\|_p \leq \|f\|_p + \|g\|_p$.

*Structural Commentary*: Quotienting by a.e. equality turns pointwise operations into vector-space operations.
*Key Insight*: Minkowski supplies the nontrivial norm axiom. □

**Definition 4.1.18.** *Completeness Language*
Let $(V, \|\cdot\|)$ be a normed space.

(i) *Complete*: $V$ is complete iff every Cauchy sequence $(x_n) \subset V$ has $x \in V$ with $\|x_n - x\| \to 0$.
(ii) *Banach*: A Banach space is a complete normed space.
(iii) *$L^p$ criterion*: $L^p(\mu)$ is complete iff every $L^p$-Cauchy sequence converges in $\|\cdot\|_p$.

**Lemma 4.1.19.** *Convergent Implies Cauchy*
*If $f_n \to f$ in $L^p(\mu)$, then $(f_n)$ is Cauchy in $L^p(\mu)$.*

*Proof.* Given $\varepsilon > 0$, choose $N$ so $n \geq N$ implies $\|f_n - f\|_p < \frac{\varepsilon}{2}$.
Then $m, n \geq N$ gives $\|f_m - f_n\|_p \leq \|f_m - f\|_p + \|f - f_n\|_p < \varepsilon$ by Minkowski.
*Structural Commentary*: A convergent sequence has uniformly small tails around its limit.
*Key Insight*: Use the common limit as the bridge between two tails. □

**Definition 4.1.20.** *Rapidly Cauchy Sequence*
Let $(V, \|\cdot\|)$ be a normed space. A sequence $(x_k) \subset V$ is *rapidly Cauchy* if $\exists (\varepsilon_k) \subset (0, +\infty)$ such that:

(i) $\sum_{k=1}^{\infty} \varepsilon_k < +\infty$.
(ii) $\|x_{k+1} - x_k\| \leq \varepsilon_k^2$ for every $k \geq 1$.

**Proposition 4.1.21.** *Rapidly Cauchy Subsequences*
*In a normed space, the following hold.*

*(i) Every rapidly Cauchy sequence is Cauchy.*
*(ii) Every Cauchy sequence has a rapidly Cauchy subsequence.*

*Proof.*

(i) Let $(x_k)$ be rapidly Cauchy with witnesses $(\varepsilon_k)$. For $m > n$, $\|x_m - x_n\| \leq \sum_{j=n}^{m-1} \|x_{j+1} - x_j\| \leq \sum_{j=n}^{\infty} \varepsilon_j^2$.
Since $\sum_j \varepsilon_j < +\infty$, eventually $\varepsilon_j \leq 1$, so $\sum_{j=n}^{\infty} \varepsilon_j^2 \leq \sum_{j=n}^{\infty} \varepsilon_j \to 0$.

(ii) Let $(x_n)$ be Cauchy. Choose $n_1 < n_2 < \cdots$ such that $m, n \geq n_k \Rightarrow \|x_m - x_n\| \leq 4^{-k}$.
Set $y_k := x_{n_k}$ and $\varepsilon_k := 2^{-k}$. Then $\sum_k \varepsilon_k < +\infty$ and $\|y_{k+1} - y_k\| \leq 4^{-k} = \varepsilon_k^2$, so $(y_k)$ is rapidly Cauchy.

*Structural Commentary*: A Cauchy tail contains a summably controlled thread.
*Key Insight*: Replace arbitrary tail control by a geometric subsequence. □

**Theorem 4.1.22.** *Riesz–Fischer Completeness Theorem*
*For $1 \leq p < \infty$, the following hold.*

*(i) Completeness: $(L^p(\mu), \|\cdot\|_p)$ is complete.*
*(ii) A.E. subsequence: if $f_n \to f$ in $L^p(\mu)$, then some subsequence converges to a representative of $f$ a.e.*

*Proof.*

(i) Let $(f_n)$ be Cauchy. Choose $g_k = f_{n_k}$ with $\|g_{k+1} - g_k\|_p \leq 4^{-k}$. Set $G_N := |g_1| + \sum_{k<N} |g_{k+1} - g_k|$.
By Minkowski, $\|G_N\|_p \leq \|g_1\|_p + \sum_{k<N} 4^{-k} \leq \|g_1\|_p + \frac{1}{3}$; Fatou gives $G := \lim_N G_N \in L^p$.
The series $h := g_1 + \sum_k (g_{k+1} - g_k)$ converges a.e. with $|h| \leq G$ and $g_k \to h$ a.e.
Since $|g_k - h| \leq 2G$, DCT gives $\|g_k - h\|_p \to 0$. Given $\varepsilon > 0$, choose $k$ with $\|g_k - h\|_p < \frac{\varepsilon}{2}$ and then $N$ with $n \geq N \Rightarrow \|f_n - g_k\|_p < \frac{\varepsilon}{2}$; hence $\|f_n - h\|_p \to 0$.

(ii) If $f_n \to f$ in $L^p$, then $(f_n)$ is Cauchy. Applying (i) gives a subsequence $f_{n_k} \to h$ a.e. and in $L^p$.
Since also $f_{n_k} \to f$ in $L^p$, uniqueness in normed space gives $[h] = f$ in $L^p$, so $h$ is a representative of $f$.

*Structural Commentary*: A rapid subsequence converges a.e.; DCT gives norm convergence.
*Key Insight*: Fatou finds the pointwise limit, dominated convergence brings it back to $L^p$. □

**Proposition 4.1.23.** *Essential Supremum Test*
*If $f \in L^\infty(\mu)$, then:*

*(i) $|f| \leq \|f\|_\infty$ a.e.;*
*(ii) $A < \|f\|_\infty \Rightarrow \mu(\{|f| > A\}) > 0$.*

*Proof.*

(i) Let $M := \|f\|_\infty$. For each $k$, $N_k := \{|f| > M + \frac{1}{k}\}$ is null. On $X \setminus \bigcup_k N_k$, $|f| \le M + \frac{1}{k}$ $\forall k$, hence $|f| \le M$.
(ii) If $A < M$ and $\mu(\{|f| > A\}) = 0$, then $|f| \le A$ a.e., so $A$ is an essential upper bound. Contradiction.

*Structural Commentary*: The essential supremum is the least a.e. upper bound.
*Key Insight*: The sharp bound cannot be lowered on a full-measure set. □

**Theorem 4.1.24.** *$L^\infty$ Is Banach*
*$(L^\infty(\mu), \|\cdot\|_\infty)$ is a Banach space.*

*Proof.*

(i) Pointwise operations descend modulo a.e. equality, and $\|\alpha f\|_\infty = |\alpha| \|f\|_\infty$. Proposition 4.1.23(i) gives definiteness; outside the union of two null sets, $|f + g| \le \|f\|_\infty + \|g\|_\infty$, hence $\|f + g\|_\infty \le \|f\|_\infty + \|g\|_\infty$.
(ii) Let $(f_n)$ be $L^\infty$-Cauchy. For each $m, n$, choose null $N_{mn}$ with $|f_m - f_n| \le \|f_m - f_n\|_\infty$ on $X \setminus N_{mn}$, and set $N := \bigcup_{m,n} N_{mn}$. On $X \setminus N$, $(f_n)$ is uniformly Cauchy, hence converges uniformly to a measurable $f$.
(iii) The uniform convergence off $N$ gives $\|f_n - f\|_\infty \to 0$.
For fixed $n_0$, $|f| \le |f_{n_0}| + \sup_{n \ge n_0} \|f_n - f_{n_0}\|_\infty$ a.e.; hence $f \in L^\infty(\mu)$.

*Structural Commentary*: Completeness is uniform convergence off one null set.
*Key Insight*: Essential bounds become ordinary uniform bounds off a common null set. □

**Proposition 4.1.25.** *Simple Approximation, Support, and Tails*
*Let $1 \le p < \infty$ and $f \in L^p(\mu)$.*

*(i) Simple measurable functions are dense in $L^p(\mu)$; also in $L^\infty(\mu)$ for $\|\cdot\|_\infty$.*
*(ii) $\{|f| > 0\}$ is $\sigma$-finite.*
*(iii) For $E_n := \{|f| \ge n\}$, $\mu(E_n) \to 0$.*
*(iv) $\forall \varepsilon > 0$ $\exists E_\varepsilon \in \mathcal{M}$ with $\mu(E_\varepsilon) < +\infty$ such that $F \cap E_\varepsilon = \emptyset$ implies $\|f \mathbf{1}_F\|_p < \varepsilon$.*

*Proof.*

(i) Choose simple $s_n \to f$ a.e. with $|s_n| \le |f|$ in the real case; treat real and imaginary parts separately in the complex case. Then $|f - s_n|^p \le 2^p |f|^p$, so DCT gives $\|f - s_n\|_p \to 0$.
For $f \in L^\infty$, discard null $N$ with $|f| \le \|f\|_\infty + \eta$ on $X \setminus N$, then quantize that bounded range.
(ii) Put $A_k := \{|f| \ge \frac{1}{k}\}$. Then $\mu(A_k) \le k^p \|f\|_p^p < +\infty$ and $\{|f| > 0\} = \bigcup_k A_k$.
(iii) $n^p \mu(E_n) \le \int_{E_n} |f|^p \, d\mu \le \|f\|_p^p$, hence $\mu(E_n) \le n^{-p} \|f\|_p^p \to 0$.
(iv) With $A_k$ as in (ii), $A_k \uparrow \{|f| > 0\}$ and $|f|^p \mathbf{1}_{X \setminus A_k} \to 0$ a.e. DCT gives $\int_{X \setminus A_k} |f|^p \, d\mu \to 0$.
Choose $E_\varepsilon = A_k$ with this integral $< \varepsilon^p$; if $F \cap E_\varepsilon = \emptyset$, then $\|f \mathbf{1}_F\|_p^p \le \int_{X \setminus E_\varepsilon} |f|^p \, d\mu < \varepsilon^p$.

*Structural Commentary*: $L^p$ mass is simple-approximable and finite-piece capturable.
*Key Insight*: Integrability turns level sets into finite-measure exhaustions. □

**Proposition 4.1.26.** *Shell Criterion on Finite Measure Spaces*
*Let $\mu(X) < +\infty$, $f$ be measurable, and $E_n := \{n - 1 \le |f| < n\}$ for $n \ge 1$.*
*For $1 \le p < \infty$, $f \in L^p(\mu) \iff \sum_{n \ge 1} n^p \mu(E_n) < +\infty$.*

*Proof.*

($\Rightarrow$) If $f \in L^p(\mu)$, then $\mu(E_1) \le \mu(X) < +\infty$ and, for $n \ge 2$, $n^p \mathbf{1}_{E_n} \le 2^p |f|^p \mathbf{1}_{E_n}$.
Thus $\sum_n n^p \mu(E_n) \le \mu(X) + 2^p \int_X |f|^p \, d\mu < +\infty$.
($\Leftarrow$) If $\sum_n n^p \mu(E_n) < +\infty$, then $|f|^p = \sum_n |f|^p \mathbf{1}_{E_n} \le \sum_n n^p \mathbf{1}_{E_n}$, hence $\int_X |f|^p \, d\mu < +\infty$.

*Structural Commentary*: The finite base shell handles small values; weighted shells control tails.
*Key Insight*: Shell weights characterize $L^p$ mass. □

**Proposition 4.1.27.** *Inclusions, Interpolation, and Multipliers*

*(i) If $\mu(X) < +\infty$ and $1 \le r \le p < \infty$, then $L^p(\mu) \subseteq L^r(\mu)$ and $\|f\|_r \le \|f\|_p \mu(X)^{\frac{1}{r} - \frac{1}{p}}$.*
*(ii) On $(\mathbb{N}, 2^{\mathbb{N}}, \#)$, $1 \le p \le s < \infty$ implies $\ell^p \subseteq \ell^s$ and $\|a\|_s \le \|a\|_p$.*
*(iii) If $1 \le p_1 < p_2 < \infty$ and $f \in L^{p_1}(\mu) \cap L^{p_2}(\mu)$, then $f \in L^p(\mu)$ for $p_1 \le p \le p_2$.*
*(iv) If $1 \le p \le \infty$, then $L^p(\mu) \cdot L^\infty(\mu) \subseteq L^p(\mu)$ and $\|fg\|_p \le \|f\|_p \|g\|_\infty$.*
*(v) $L^\infty(\mu) \subseteq L^1(\mu) \iff \mu(X) < +\infty$. If $\mu(X) = 1$, then $\|f\|_\infty = \lim_{p \to \infty} \|f\|_p$ for $f \in L^\infty(\mu)$.*

*Proof.*

(i) For $r = p$, $L^p \subseteq L^p$ and the estimate reads $\|f\|_p \le \|f\|_p$, i.e. equality.
For $r < p$, Hölder applied to $|f|^r \cdot 1$ gives $\int_X |f|^r \, d\mu \le \|f\|_p^r \, \mu(X)^{1-\frac{r}{p}}$; take $r$th roots.

(ii) For $a \in \ell^p$, $\forall n,\ |a_n| \le \|a\|_p$, hence $\sum_n |a_n|^s \le \|a\|_p^{s-p} \sum_n |a_n|^p \le \|a\|_p^s$.

(iii) For $p \in [p_1, p_2]$, $|f|^p \le |f|^{p_1} \mathbf{1}_{\{|f|\le 1\}} + |f|^{p_2} \mathbf{1}_{\{|f|>1\}}$.

(iv) For $p < \infty$, $|fg|^p \le \|g\|_\infty^p \, |f|^p$ a.e.; for $p = \infty$, $|fg| \le \|f\|_\infty \, \|g\|_\infty$ a.e.

(v) ($\Rightarrow$) If $L^\infty(\mu) \subseteq L^1(\mu)$, then $\mathbf{1}_X \in L^\infty(\mu)$ lies in $L^1(\mu)$, so $\mu(X) = \int_X \mathbf{1}_X \, d\mu < +\infty$.

(v) ($\Leftarrow$) If $\mu(X) < +\infty$ and $f \in L^\infty(\mu)$, then $\int_X |f| \, d\mu \le \|f\|_\infty \, \mu(X) < +\infty$. If $\mu(X) = 1$, then $\|f\|_p \le \|f\|_\infty$.
For $A < \|f\|_\infty$, $\mu(\{|f| > A\}) > 0$ and $\|f\|_p \ge A\mu(\{|f| > A\})^{\frac{1}{p}} \to A$. Let $A \uparrow \|f\|_\infty$.

*Structural Commentary*: Inclusions depend on domain size and pointwise bounds.
*Key Insight*: Check whether size comes from the function values or from the measure of the domain. □

**Example 4.1.28.** *Membership Models*

(i) On $(\mathbb{N}, 2^{\mathbb{N}}, \#)$, $a_n = \frac{1}{n}$ satisfies $\|a\|_p^p = \sum_{n\ge 1} n^{-p}$; hence $a \in \ell^p \iff p > 1$, while $a \in \ell^\infty \setminus \ell^1$.

(ii) On $\mathbb{N}$ with $\lambda(\{n\}) = n^{-2}$, $\lambda(\mathbb{N}) < +\infty$ and $f(n) = n^{\frac{1}{p_0}}$ satisfies $\int_{\mathbb{N}} |f|^p \, d\lambda = \sum_{n\ge 1} n^{\frac{p}{p_0}-2}$.
Hence $f \in L^p(\lambda) \iff 1 \le p < p_0$. In particular, $f(n) = \sqrt{n}$ gives $p_0 = 2$.

(iii) On $(0, \infty)$, $f(x) = x^{-\frac{1}{2}}(1 + |\log x|)^{-1}$ gives $\int_0^\infty |f|^p \, dx = \int_{-\infty}^\infty e^{(1-\frac{p}{2})t}(1 + |t|)^{-p} \, dt$ under $x = e^t$.

*Structural Commentary*: $L^p$ membership reduces to series and endpoint tests.
*Key Insight*: Different ends impose different $p$-conditions.

**Proposition 4.1.29.** *Hölder Dual Norm Attainment*
*Let $1 < p < \infty$, $p^{-1} + q^{-1} = 1$, and let $f \in L^p(\mu)$ be real-valued.*

*(i)* $\sup_{\|g\|_q \le 1} \left|\int_X fg \, d\mu\right| = \|f\|_p$.

*(ii) If $f \ne 0$, the supremum is attained by* $g_0 = \|f\|_p^{-\frac{p}{q}} \operatorname{sign}(f) \, |f|^{p-1}$.

*Proof.*

(i) For $\|g\|_q \le 1$, Hölder gives $\left|\int_X fg \, d\mu\right| \le \|f\|_p \, \|g\|_q \le \|f\|_p$.
If $f = 0$ this is sharp; if $f \ne 0$, item (ii) gives an admissible $g_0$ with $\int_X fg_0 \, d\mu = \|f\|_p$.

(ii) Since $(p-1)q = p$, $\|g_0\|_q^q = \|f\|_p^{-p} \int_X |f|^p \, d\mu = 1$ and $\int_X fg_0 \, d\mu = \|f\|_p^{-\frac{p}{q}} \int_X |f|^p \, d\mu = \|f\|_p$.

*Structural Commentary*: Equality in Hölder is obtained by matching the conjugate powers.
*Key Insight*: The $L^q$ test function points in the power direction of $f$. □

**Definition 4.1.30.** *Bounded Linear Functionals on $L^p$*

(i) A *linear functional* on real $L^p(\mu)$ is a linear map $G : L^p(\mu) \to \mathbb{R}$.
(ii) $G$ is *bounded* if $\exists M < +\infty$ such that $\forall f \in L^p(\mu)$, $|G(f)| \le M \|f\|_p$.
(iii) The *operator norm* is $\|G\| := \sup\{|G(f)| : f \in L^p(\mu),\ \|f\|_p \le 1\}$.
(iv) $G$ is *positive* if $f \ge 0$ a.e. implies $G(f) \ge 0$.

**Lemma 4.1.31.** *Positive Split of a Bounded Functional*
*Let $1 \le p < +\infty$ and $G \in (L^p(\mu))^*_{\mathbb{R}}$. Then $G = G^+ - G^-$ for positive bounded linear functionals $G^\pm$.*

*Proof.* For $f \ge 0$, set $G^+(f) := \sup\{G(g) : 0 \le g \le f\}$.
This is finite since $|G(g)| \le \|G\| \, \|g\|_p \le \|G\| \, \|f\|_p$, and positive homogeneity is immediate.
For $f_1, f_2 \ge 0$, $G^+(f_1) + G^+(f_2) \le G^+(f_1 + f_2)$; conversely, if $0 \le h \le f_1 + f_2$, set $h_1 = (h - f_2)^+$ and $h_2 = h - h_1$, so $h = h_1 + h_2$ with $0 \le h_j \le f_j$, hence $G(h) \le G^+(f_1) + G^+(f_2)$.
Thus $G^+$ is additive on the positive cone. Extend by $G^+(f) = G^+(f^+) - G^+(f^-)$; then $G^+$ is positive, linear, and bounded. Let $G^- = G^+ - G$. Since $G^+(f) \ge G(f)$ for $f \ge 0$, $G^-$ is positive and bounded.
*Structural Commentary*: Function-lattice functionals split into positive parts.
*Key Insight*: Split any competitor $h \le f_1 + f_2$ into the portions assigned to $f_1$ and $f_2$. □

**Theorem 4.1.32.** *Riesz Representation for $L^1$*
*Let $(X, \mathcal{M}, \mu)$ be $\sigma$-finite and $G \in (L^1(\mu))^*_{\mathbb{R}}$.*

*(i) $\exists g \in L^\infty(\mu)$ such that $\forall f \in L^1(\mu)$, $G(f) = \int_X fg \, d\mu$.*
*(ii) $\|G\| = \|g\|_\infty$.*
*(iii) If $G$ is positive, then $g \ge 0$ a.e.*

*Proof.*

(i) First let $\mu(X) < +\infty$ and $G \geq 0$. Set $\lambda(E) := G(\mathbf{1}_E)$. Finite additivity is linearity; if $E_n \uparrow E$, then $\mathbf{1}_{E_n} \to \mathbf{1}_E$ in $L^1$, hence $\lambda(E_n) \to \lambda(E)$. Thus $\lambda$ is finite, and $\lambda \ll \mu$ since $\mu(E) = 0$ gives $\mathbf{1}_E = 0$ in $L^1$. Radon–Nikodym gives $g \geq 0$ with $\lambda(E) = \int_E g \, d\mu$, so simple functions are represented.
By (ii), $g \in L^\infty$, hence $L^1$-density of simple functions extends the identity to all $f \in L^1$. For $\sigma$-finite $\mu$, apply the finite case to $f \mapsto G(f\mathbf{1}_{X_n})$ on $X_n \uparrow X$, $\mu(X_n) < +\infty$, and patch densities on overlaps. For general $G$, use the positive split $G = G^+ - G^-$ and subtract the two densities.

(ii) The pairing estimate gives $\|G\| \leq \|g\|_\infty$. Conversely, if $A > \|G\|$ and $E \subseteq \{|g| > A\}$ has $\mu(E) < +\infty$, then with $f = \operatorname{sign}(g)\mathbf{1}_E$, $A\mu(E) < \int_E |g| \, d\mu = G(f) \leq \|G\| \, \mu(E)$, so $\mu(E) = 0$.
A $\sigma$-finite exhaustion gives $\forall A > \|G\|$, $\mu(\{|g| > A\}) = 0$, hence $\|g\|_\infty \leq \|G\|$.

(iii) If $G$ is positive and $B = \{g < 0\}$, then $\forall E \subseteq B$ of finite measure, $0 \leq G(\mathbf{1}_E) = \int_E g \, d\mu \leq 0$ only if $\int_E (-g) \, d\mu = 0$. Hence $\mu(E) = 0$ on $\{g < -\frac{1}{k}\}$ $\forall k$, and a $\sigma$-finite exhaustion gives $\mu(B) = 0$.

*Structural Commentary*: A bounded functional on $L^1$ is integration against an essentially bounded density.
*Key Insight*: Indicators give a measure; Radon–Nikodym gives the density. □

**Theorem 4.1.33.** *Riesz Representation for $L^p$, $1 < p < +\infty$*
*Let $(X, \mathcal{M}, \mu)$ be any measure space, $1 < p < +\infty$, $q = \frac{p}{p-1}$, and $G \in (L^p(\mu))^*_{\mathbb{R}}$.*

*(i) Representation: $\exists g \in L^q(\mu)$ such that $\forall f \in L^p(\mu)$, $G(f) = \int_X fg \, d\mu$.*
*(ii) Norm: every representing $g$ satisfies $\|G\| = \|g\|_q$.*
*(iii) Uniqueness: if $g, h \in L^q(\mu)$ represent $G$, then $g = h$ a.e.*

*Proof.*

(i) Finite $\mu$: for $G \geq 0$, set $\lambda(E) := G(\mathbf{1}_E)$. Since $E_n \uparrow E \Rightarrow \mathbf{1}_{E_n} \to \mathbf{1}_E$ in $L^p$, $\lambda$ is finite.
Since $\mu(E) = 0 \Rightarrow \mathbf{1}_E = 0$ in $L^p$, $\lambda \ll \mu$. Let $g = \frac{d\lambda}{d\mu}$.
By simple approximation, $G(\phi) = \int_X \phi g \, d\mu$ for bounded $\phi$.
For $g_m := g \wedge m$, $\int_X g_m^q \, d\mu \leq G(g_m^{q-1}) \leq \|G\| \, (\int_X g_m^q \, d\mu)^{1/p}$; hence $g \in L^q$, by $L^p$-density $G(f) = \int_X fg \, d\mu$.
General finite $G$ follows from $G = G^+ - G^-$. For $\sigma$-finite $\mu$, choose $X_n \uparrow X$, $\mu(X_n) < +\infty$, represent $G_n : u \mapsto G(\tilde{u})$ by $g_n$, and patch the compatible $g_n$ to $g$.
Since $\|g_n\|_q = \|G_n\| \leq \|G\|$, by MCT $g \in L^q$; $f\mathbf{1}_{X_n} \to f$ in $L^p \Rightarrow G(f) = \int_X fg \, d\mu$.
For arbitrary $\mu$, choose $\|f_n\|_p = 1$, $G(f_n) \to \|G\|$, and $X_0 := \bigcup_{n,k} \{|f_n| > 1/k\}$.
If $E \subseteq X \setminus X_0$, $\mu(E) < +\infty$, and $a = G(\mathbf{1}_E)$, then $G(f_n) \pm ta \leq \|G\| \, (1 + t^p \mu(E))^{1/p}$; after $n \to \infty$ and $t \downarrow 0$, $a = 0$. Thus $G$ vanishes off $X_0$; represent on $X_0$ and extend by 0.

(ii) For representing $g$, by Hölder, $\|G\| \leq \|g\|_q$; by Proposition 4.1.29, $\|g\|_q = \sup_{\|f\|_p \leq 1} \left| \int_X fg \, d\mu \right| \leq \|G\|$.

(iii) If $g, h$ represent $G$, then $k := g - h$ satisfies $\int_X fk \, d\mu = 0 \; \forall f \in L^p(\mu)$.
By Proposition 4.1.29 applied to $k \in L^q(\mu)$ with exponents $q, p$, $\|k\|_q = 0$; hence $g = h$ a.e.

*Structural Commentary*: For $1 < p < \infty$, every bounded functional comes from the conjugate space $L^q$.
*Key Insight*: Localize to $\sigma$-finite support, then patch. □

**Definition 4.1.34.** *Bounded Finitely Additive Measures*
Let $(X, \mathcal{M}, \mu)$ be a measure space.
For $E \in \mathcal{M}$, let $\Pi(E)$ be the finite families $(E_1, \ldots, E_n)$ with $E_k \in \mathcal{M}$, $E_k \subseteq E$, and $E_i \cap E_j = \emptyset$ for $i \neq j$.
For $\nu : \mathcal{M} \to \mathbb{R}$, set $|\nu|_{\text{fa}}(E) := \sup_{(E_k) \in \Pi(E)} \sum_k |\nu(E_k)|$ and $\|\nu\|_{\text{var}} := |\nu|_{\text{fa}}(X)$.

Define $\text{BFA}(X, \mathcal{M}, \mu)$ as the set of $\nu : \mathcal{M} \to \mathbb{R}$ satisfying:

(i) *Finite additivity*: $\nu(\emptyset) = 0$ and $\nu(\bigsqcup_{k=1}^n E_k) = \sum_{k=1}^n \nu(E_k)$.
(ii) *Null compatibility*: $\mu(E) = 0 \Rightarrow \nu(E) = 0$.
(iii) *Bounded variation*: $\|\nu\|_{\text{var}} < +\infty$.

*Key Insight*: BFA requires finite additivity, null compatibility, and finite variation, not countable additivity.

**Theorem 4.1.35.** *Kantorovitch Representation for $(L^\infty)^*$*
*Let $(X, \mathcal{M}, \mu)$ be a measure space.*
*For $\nu \in \text{BFA}(X, \mathcal{M}, \mu)$ and simple $\varphi = \sum_{k=1}^n c_k \mathbf{1}_{E_k}$ over a finite measurable partition, set $I_\nu(\varphi) := \sum_{k=1}^n c_k \nu(E_k)$.*

*(i) Extension: $I_\nu$ descends uniquely to $T_\nu \in (L^\infty(\mu))^*_{\mathbb{R}}$.*
*(ii) Isometry: $\|T_\nu\| = \|\nu\|_{\text{var}}$.*
*(iii) Surjectivity: if $S \in (L^\infty(\mu))^*_{\mathbb{R}}$ and $\nu_S(E) := S(\mathbf{1}_E)$, then $\nu_S \in \text{BFA}(X, \mathcal{M}, \mu)$ and $S = T_{\nu_S}$.*

*Thus $\nu \mapsto T_\nu$ is an isometric isomorphism $\text{BFA}(X, \mathcal{M}, \mu) \cong (L^\infty(\mu))^*_{\mathbb{R}}$.*

*Proof.*

(i) Refinement and finite additivity make $I_\nu$ well defined.
For simple $\varphi$, discard $N = \{|\varphi| > \|\varphi\|_\infty\}$; since $|\nu|_{\mathrm{fa}}(N) = 0$, $|I_\nu(\varphi)| \le \|\varphi\|_\infty \|\nu\|_{\mathrm{var}}$.
Hence uniform simple approximation extends $I_\nu$ to bounded measurable $f$.
If $\operatorname{supp} f \subseteq N$ and $\mu(N) = 0$, then $|\nu|_{\mathrm{fa}}(N) = 0$, so $I_\nu(f) = 0$; therefore the extension depends only on the $L^\infty$ class and defines $T_\nu$ with $\|T_\nu\| \le \|\nu\|_{\mathrm{var}}$.

(ii) For $(E_k) \in \Pi(X)$, set $c_k := \operatorname{sign}(\nu(E_k))$ and $\varphi := \sum_k c_k \mathbf{1}_{E_k}$.
Then $\|\varphi\|_\infty \le 1$ and $\sum_k |\nu(E_k)| = T_\nu(\varphi) \le \|T_\nu\|$.
Supremizing over $\Pi(X)$ gives $\|\nu\|_{\mathrm{var}} \le \|T_\nu\|$, and item (i) gives equality.

(iii) For $S \in (L^\infty(\mu))^*_{\mathbb{R}}$, $\nu_S(\emptyset) = 0$ and finite additivity follow from linearity.
If $\mu(E) = 0$, then $\mathbf{1}_E = 0$ in $L^\infty$, so $\nu_S(E) = 0$.
For $(E_k) \in \Pi(X)$, with $c_k = \operatorname{sign}(\nu_S(E_k))$ and $\varphi = \sum_k c_k \mathbf{1}_{E_k}$, $\sum_k |\nu_S(E_k)| = S(\varphi) \le \|S\|$.
Hence $\nu_S \in \mathrm{BFA}(X, \mathcal{M}, \mu)$. Since $S$ and $T_{\nu_S}$ agree on simple functions, uniform simple density gives $S = T_{\nu_S}$.

*Structural Commentary*: Simple extensions give $T_\nu$; indicators recover $\nu$.
*Key Insight*: Finite additivity supplies the full $L^\infty$ dual. □

**Proposition 4.1.36.** *The Countably Additive Part*
*For $g \in L^1(\mu)$, set $\nu_g(E) := \int_E g\, d\mu$.*

*(i)* $\nu_g \in \mathrm{BFA}(X, \mathcal{M}, \mu)$.
*(ii)* $\|\nu_g\|_{\mathrm{var}} = \|g\|_1$.
*(iii)* $T_{\nu_g}(f) = \int_X fg\, d\mu$ *for $f \in L^\infty(\mu)$; hence $L^1(\mu)$ embeds isometrically into $(L^\infty(\mu))^*$.*

*Proof.*

(i) For disjoint $E_1, \dots, E_n$, $\nu_g(\bigsqcup_k E_k) = \sum_k \nu_g(E_k)$ by finite additivity of the integral, and $\mu(E) = 0 \Rightarrow \nu_g(E) = 0$. Also, for $(E_k) \in \Pi(X)$, $\sum_k |\nu_g(E_k)| \le \sum_k \int_{E_k} |g|\, d\mu \le \|g\|_1$, so $\nu_g \in \mathrm{BFA}(X, \mathcal{M}, \mu)$.

(ii) Corollary 3.1.7 gives $\|\nu_g\|_{\mathrm{var}} = |\nu_g|(X)$, and Theorem 3.1.9 gives $|\nu_g|(X) = \int_X |g|\, d\mu = \|g\|_1$.

(iii) If $f = \sum_k c_k \mathbf{1}_{E_k}$ is simple, then $T_{\nu_g}(f) = \sum_k c_k \nu_g(E_k) = \int_X fg\, d\mu$.
For $f \in L^\infty(\mu)$, choose simple $f_n \to f$ uniformly; then $T_{\nu_g}(f_n) \to T_{\nu_g}(f)$ and $\int_X f_n g\, d\mu \to \int_X fg\, d\mu$.

*Structural Commentary*: Variation fixes the norm; simple approximation extends the pairing.
*Key Insight*: Countably additive representatives are exactly the familiar $L^1$ integral pairings. □




Figure 217: $L^q(\mu) \cong (L^p(\mu))^*$ for $1 < p < \infty$; $L^1(\mu)$ is the countably additive part of BFA $\cong (L^\infty(\mu))^*$.

**Remark 4.1.37.** *$L^2$ Riesz Gives Another Radon–Nikodym Route*
Let $\lambda, \mu$ be finite and $\lambda \ll \mu$. Put $\rho = \lambda + \mu$ and $G(h) := \int h\, d\lambda$ on $L^2(\rho)$.
If $G(h) = \int hg\, d\rho$, indicator tests give $0 \le g \le 1$ $\rho$-a.e. and $\mu(\{g = 1\}) = 0$.
Since $\int h(1-g)\, d\lambda = \int hg\, d\mu$, take $E_N = E \cap \{1 - g \ge \frac{1}{N}\}$ and $h = \frac{\mathbf{1}_{E_N}}{1-g} \in L^2(\rho)$ to get $\lambda(E_N) = \int_{E_N} \frac{g}{1-g}\, d\mu$.
Let $N \to \infty$ by monotone convergence and $\lambda \ll \mu$: $\lambda(E) = \int_E \frac{g}{1-g}\, d\mu$.
*Key Insight*: Riesz representation on $L^2(\lambda + \mu)$ encodes the density in the ratio $\frac{g}{1-g}$.

**Lemma 4.1.38.** *Absolute Continuity of $L^p$ Mass*
*For $1 \le p < \infty$ and $f \in L^p(\mu)$, $\forall \varepsilon > 0\, \exists \delta > 0$ such that $\mu(E) < \delta$ implies $\|f\mathbf{1}_E\|_p < \varepsilon$.*

*Proof.* Let $h = |f|^p \in L^1$. Choose $M > 0$ with $\int_{\{h > M\}} h\, d\mu < \frac{\varepsilon^p}{2}$, and set $\delta = \frac{\varepsilon^p}{2M}$.
If $\mu(E) < \delta$, then $\|f\mathbf{1}_E\|_p^p = \int_E h\, d\mu \le M\mu(E) + \int_{\{h > M\}} h\, d\mu < \varepsilon^p$.
*Structural Commentary*: $L^p$ mass is absolutely continuous with respect to measure.
*Key Insight*: Truncate the tail; small measure controls the bounded part. □

**Proposition 4.1.39.** *Local Mass Limits for $L^p$ Cauchy Sequences*
*For $1 \le p < \infty$ and $f_n \in L^p(\mu)$, set $\beta_n(E) := \|f_n \mathbf{1}_E\|_p$ for $E \in \mathcal{M}$. The following hold.*

*(i)* $|\beta_n(E) - \beta_m(E)| \le \|f_n - f_m\|_p$ *for* $E \in \mathcal{M}$.
*(ii) If $(f_n)$ is $L^p$-Cauchy, then $\forall E \in \mathcal{M}$, $\beta(E) := \lim_n \beta_n(E)$ exists.*

*Proof.*

(i) Since $f_n \mathbf{1}_E = f_m \mathbf{1}_E + (f_n - f_m)\mathbf{1}_E$, $\beta_n(E) - \beta_m(E) \le \|f_n - f_m\|_p$.
Interchanging $m, n$ gives $\beta_m(E) - \beta_n(E) \le \|f_n - f_m\|_p$; hence $|\beta_n(E) - \beta_m(E)| \le \|f_n - f_m\|_p$.
(ii) Fix $E \in \mathcal{M}$ and $\varepsilon > 0$. Choose $N$ such that $m, n \ge N \Rightarrow \|f_n - f_m\|_p < \varepsilon$.
Then item (i) gives $|\beta_n(E) - \beta_m(E)| < \varepsilon$, so $(\beta_n(E))$ converges in $\mathbb{R}$.

*Structural Commentary*: Restricting to $E$ is a 1-Lipschitz operation on $L^p$ mass.
*Key Insight*: Local mass inherits global Cauchy control. □

**Proposition 4.1.40.** *Uniform Local Mass Control for $L^p$ Cauchy Sequences*
*Let $1 \le p < \infty$, $f_n$ be $L^p$-Cauchy, and $\beta_n(E) := \|f_n \mathbf{1}_E\|_p$. Then $\forall \varepsilon > 0$:*

*(i)* $\exists E_\varepsilon \in \mathcal{M}$ *with* $\mu(E_\varepsilon) < +\infty$ *such that* $F \cap E_\varepsilon = \emptyset$ *implies* $\sup_n \beta_n(F) < \varepsilon$;
*(ii)* $\exists \delta > 0$ *such that* $\mu(E) < \delta$ *implies* $\sup_n \beta_n(E) < \varepsilon$.

*Proof.* Choose $N$ so $n \ge N$ implies $\|f_n - f_N\|_p < \frac{\varepsilon}{3}$.

(i) Choose finite-measure $E_j$ for $1 \le j \le N$ with $F \cap E_j = \emptyset \Rightarrow \beta_j(F) < \frac{\varepsilon}{3}$, and set $E_\varepsilon = \bigcup_{j \le N} E_j$.
For $n \le N$ this works; for $n > N$, $\beta_n(F) \le \|f_n - f_N\|_p + \beta_N(F) < \varepsilon$.
(ii) Choose $\delta_j > 0$ for $1 \le j \le N$ with $\mu(E) < \delta_j \Rightarrow \beta_j(E) < \frac{\varepsilon}{3}$, and set $\delta = \min_{j \le N} \delta_j$.
For $n \le N$ this works; for $n > N$, $\beta_n(E) \le \|f_n - f_N\|_p + \beta_N(E) < \varepsilon$.

*Structural Commentary*: Cauchy families are uniformly tight and absolutely continuous.
*Key Insight*: Control finitely many early terms directly and the tail by Cauchy closeness. □

**Method 4.1.41.** *Lebesgue Proof Patterns*

(i) *Approximate*: for $f \ge 0$, use $s \in \mathcal{S}^+(f)$ and $\int_E f \, d\mu = \sup_{s \in \mathcal{S}^+(f)} \mathcal{I}_E(s)$.
(ii) *Promote*: prove for $\mathbf{1}_A$, extend to $s = \sum_i c_i \mathbf{1}_{A_i}$, then pass through $s_n \uparrow f$.
(iii) *Ignore null sets*: if $f \sim_{\text{a.e.}} g$ on $E$, replace $f$ by $g$ inside integrals over measurable $A \subseteq E$.
(iv) *Split signs*: use $f = f^+ - f^-$ and $h = u + iv$.
(v) *Exchange limits*: reduce Fatou and dominated convergence to $0 \le f_n \uparrow f$.
(vi) *Slice*: prove boxes first; use regularity and null slices to pass to products and Fubini.

*Key Insight*: Reduce to simple functions, monotone convergence, null sets, and product boxes.

## 4.2 Modes of Convergence

Throughout this subsection $(X, \mathcal{M}, \mu)$ is a fixed measure space and, unless otherwise stated, functions are real-valued and measurable. The point is not to rank convergence modes linearly, but to know which hypotheses convert one mode into another.

**Definition 4.2.1.** *Pointwise, Uniform, and Almost-Everywhere Convergence*
Let $(f_n)$ and $f$ be functions on $X$.

(i) $f_n \to f$ *pointwise* if $\forall x \in X$, $\varepsilon > 0$, $\exists N(\varepsilon, x)$ s.t. $n \ge N(\varepsilon, x)$ implies $|f_n(x) - f(x)| < \varepsilon$.
(ii) $f_n \to f$ *uniformly* if $\forall \varepsilon > 0$ $\exists N(\varepsilon)$ s.t. $n \ge N(\varepsilon)$ implies $|f_n(x) - f(x)| < \varepsilon$ $\forall x \in X$.
(iii) $f_n \to f$ *almost everywhere* if $\exists A \in \mathcal{M}$ with $\mu(A) = 0$ s.t. $\forall x \in X \setminus A$, $\varepsilon > 0$,
$\exists N(\varepsilon, x)$ s.t. $n \ge N(\varepsilon, x)$ implies $|f_n(x) - f(x)| < \varepsilon$.

**Proposition 4.2.2.** *First Implications and First Failures*
*Let $f_n, f : X \to \mathbb{R}$.*

*(i)* $f_n \rightrightarrows f \Rightarrow f_n \to f$ *pointwise* $\Rightarrow f_n \to f$ *a.e.; converses fail.*
*(ii)* $|X| < +\infty$ *and* $f_n \to f$ *pointwise* $\Rightarrow f_n \rightrightarrows f$.
*(iii)* $\mu(A) = 0 \Rightarrow A = \emptyset$ *and* $f_n \to f$ *a.e.* $\Rightarrow f_n \to f$ *pointwise.*

*Proof.*

(i) If $f_n \rightrightarrows f$, then $\forall \varepsilon > 0\ \exists N_\varepsilon$ such that $\forall n \geq N_\varepsilon\ \forall x \in X$, $|f_n(x) - f(x)| < \varepsilon$; hence pointwise convergence with $N(\varepsilon, x) = N_\varepsilon$. Pointwise convergence is a.e. convergence with $A = \emptyset$.
Conversely, on $([0,1], \lambda)$, $x^n \to \mathbf{1}_{\{1\}}$ pointwise but not uniformly, and $\mathbf{1}_{\{0\}} \to 0$ a.e. but not pointwise.

(ii) If $X$ is finite and $f_n \to f$ pointwise, then $\forall \varepsilon > 0\ \forall x \in X$, choose $N(\varepsilon, x)$.
Set $N_\varepsilon = \max_{x \in X} N(\varepsilon, x)$; then $\forall n \geq N_\varepsilon\ \forall x \in X$, $|f_n(x) - f(x)| < \varepsilon$, so $f_n \rightrightarrows f$.

(iii) If $f_n \to f$ a.e., then $\exists A \in \mathcal{M}$ with $\mu(A) = 0$ and $\forall x \in X \setminus A$, $f_n(x) \to f(x)$.
Since every null set is empty, $A = \emptyset$, so $f_n \to f$ pointwise.

*Structural Commentary*: Uniformity is bounded $x$-dependence of $N(\varepsilon, x)$.
*Key Insight*: Uniformity controls indices; a.e. convergence controls null exceptions. □

**Proposition 4.2.3.** *Uniform Convergence Gives $L^p$ Convergence on Finite Measure Spaces*
*Let $1 \leq p < \infty$, $\mu(X) < +\infty$, and $f_n \in L^p(\mu)$. If $f_n \rightrightarrows f$ on $X$, then $f \in L^p(\mu)$ and $\|f_n - f\|_p \to 0$.*

*Proof.* For $\varepsilon > 0$, choose $N$ with $n \geq N \Rightarrow |f_n - f| \leq \varepsilon$ on $X$.
Then $n \geq N \Rightarrow \|f_n - f\|_p \leq (\int_X \varepsilon^p \, d\mu)^{\frac{1}{p}} = \varepsilon \mu(X)^{\frac{1}{p}}$, so $\|f_n - f\|_p \to 0$.
Taking $\varepsilon = 1$, choose $n_0$ with $|f_{n_0} - f| \leq 1$ on $X$; hence $f - f_{n_0} \in L^p(\mu)$ and $f = f_{n_0} + (f - f_{n_0}) \in L^p(\mu)$.
*Structural Commentary*: Uniform error becomes $L^p$ error after multiplication by $\mu(X)^{\frac{1}{p}}$.
*Key Insight*: Finite measure converts a pointwise sup-bound into an integral bound. □

**Theorem 4.2.4.** *Dominated Almost-Everywhere Convergence Gives $L^p$ Convergence*
*Let $1 \leq p < \infty$, $f_n \in L^p(\mu)$, and $f_n \to f$ a.e. Suppose $\exists g \in L^p(\mu)$ such that $\forall n$, $|f_n| \leq g$ a.e.*

*(i)* $f \in L^p(\mu)$.
*(ii)* $\|f_n - f\|_p \to 0$.

*Proof.*

(i) Choose $N \in \mathcal{M}$ with $\mu(N) = 0$ such that on $X \setminus N$, $f_n \to f$ and $|f_n| \leq g\ \forall n$.
Then $|f| \leq g$ a.e., so $|f|^p \leq g^p \in \mathscr{L}^1(\mu)$.

(ii) Since $|f_n - f|^p \to 0$ a.e. and $|f_n - f|^p \leq (2g)^p \in \mathscr{L}^1(\mu)$ a.e.,
Theorem 1.3.7 yields $\|f_n - f\|_p^p = \int_X |f_n - f|^p \, d\mu \to 0$.

*Structural Commentary*: Apply ordinary dominated convergence to the $p$th-power error.
*Key Insight*: A single $L^p$ envelope controls both membership of the limit and convergence in mean. □

**Corollary 4.2.5.** *Uniform Boundedness on Finite Measure Spaces*
*Let $\mu(X) < +\infty$, $0 \leq K < +\infty$, and $f_n \to f$ a.e. Assume $\forall n$, $|f_n| \leq K$ a.e. Then for every $1 \leq p < \infty$:*

*(i)* $f \in L^p(\mu)$.
*(ii)* $\|f_n - f\|_p \to 0$.

*Proof.* Fix $1 \leq p < \infty$ and set $g := K\mathbf{1}_X$. Then $\int_X g^p \, d\mu = K^p \mu(X) < +\infty$, so $g \in L^p(\mu)$ and $|f_n| \leq g$ a.e.

(i) Since $f_n \to f$ a.e. and $|f_n| \leq g \in L^p(\mu)$ a.e., Theorem 4.2.4(i) gives $f \in L^p(\mu)$.

(ii) The same hypotheses and Theorem 4.2.4(ii) give $\|f_n - f\|_p \to 0$.

*Structural Commentary*: Finite measure turns a uniform a.e. bound into an $L^p$ envelope.
*Key Insight*: Bounded a.e. convergence on finite measure is dominated convergence. □

**Example 4.2.6.** *Uniform Need Not Imply $L^p$ on Infinite Measure*
On $(\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda)$, fix $1 \leq p < \infty$ and set $f_n := n^{-\frac{1}{p}} \mathbf{1}_{[0,n]}$. Then $\|f_n\|_\infty = n^{-\frac{1}{p}} \to 0$, so $f_n \rightrightarrows 0$.
However $\|f_n\|_p^p = \int_0^n n^{-1} \, d\lambda = 1$, hence $\|f_n\|_p \nrightarrow 0$.
*Structural Commentary*: The support grows exactly enough to preserve $L^p$ mass.
*Key Insight*: Uniform smallness alone does not control integrals on infinite measure spaces.

**Example 4.2.7.** *Almost Everywhere and In-Measure Need Not Imply $L^p$*
On $[0,1]$ with Lebesgue measure, set $f_n := n\mathbf{1}_{[\frac{1}{n}, \frac{2}{n}]}$.
$\forall x \in [0,1]$, $x \in [\frac{1}{n}, \frac{2}{n}]$ for only finitely many $n$, hence $f_n(x) \to 0$.
If $\alpha > 0$ and $n \geq \alpha$, then $\lambda\{|f_n| \geq \alpha\} = \frac{1}{n} \to 0$, so $f_n \to 0$ in measure.
Yet for $1 \leq p < \infty$, $\|f_n\|_p^p = n^p \lambda([\frac{1}{n}, \frac{2}{n}]) = n^{p-1} \nrightarrow 0$.
Given $\delta > 0$, choose $0 < \eta < \delta$; on $[0,1] \setminus [0, \eta]$, $\forall n > \frac{2}{\eta}$, $f_n = 0$, so $f_n \to 0$ almost uniformly.
Thus a.e., in-measure, and almost-uniform convergence do not imply $L^p$ convergence.

**Theorem 4.2.8.** *Pointwise A.E. Convergence Plus Norms*
*Let $1 \leq p < \infty$, $f_n, f \in L^p(\mu)$, $f_n \to f$ a.e. Then $f_n \to f$ in $L^p(\mu)$ iff $\lim_{n \to \infty} \int_X |f_n|^p \, d\mu = \int_X |f|^p \, d\mu$.*

*Proof.*

($\Rightarrow$) If $\|f_n - f\|_p \to 0$, then $\left|\|f_n\|_p - \|f\|_p\right| \le \|f_n - f\|_p$, hence $\int_X |f_n|^p \, d\mu = \|f_n\|_p^p \to \|f\|_p^p = \int_X |f|^p \, d\mu$.

($\Leftarrow$) Assume $\int_X |f_n|^p \, d\mu \to \int_X |f|^p \, d\mu$. Set $h_n := \frac{|f_n|^p + |f|^p}{2} - \left|\frac{f_n - f}{2}\right|^p \ge 0$ by convexity.
Since $h_n \to |f|^p$ a.e., Fatou gives $\int_X |f|^p \, d\mu \le \liminf_n \int_X h_n \, d\mu = \int_X |f|^p \, d\mu - 2^{-p} \limsup_n \int_X |f_n - f|^p \, d\mu$.
Thus $\limsup_n \|f_n - f\|_p^p \le 0$, so $f_n \to f$ in $L^p(\mu)$.

*Structural Commentary*: A.e. convergence identifies the limit; $p$-mass convergence prevents loss.
*Key Insight*: Fatou applied to the convexity defect forces the error norm to vanish. □

**Definition 4.2.9.** *Convergence and Cauchy Convergence in Measure*
Let $(f_n)$ be measurable functions on $X$.

(i) $f_n \to f$ *in measure* if $\forall \alpha > 0$, $\mu\{x \in X : |f_n(x) - f(x)| \ge \alpha\} \to 0$.
(ii) $(f_n)$ is *Cauchy in measure* if $\forall \alpha > 0$, $\mu\{x \in X : |f_m(x) - f_n(x)| \ge \alpha\} \to 0$ as $m, n \to \infty$.

The same definitions result with $>$ in place of $\ge$.

**Proposition 4.2.10.** *$L^p$ Convergence Implies Convergence in Measure*
*Let $1 \le p < \infty$. If $\|f_n - f\|_p \to 0$, then $f_n \to f$ in measure.*

*Proof.* For $\alpha > 0$, set $E_n(\alpha) := \{|f_n - f| \ge \alpha\}$.
Since $\|f_n - f\|_p^p \ge \int_{E_n(\alpha)} |f_n - f|^p \, d\mu \ge \alpha^p \mu(E_n(\alpha))$, $\mu(E_n(\alpha)) \le \alpha^{-p} \|f_n - f\|_p^p \to 0$.
*Structural Commentary*: Large-error sets are controlled by the integral of the $p$th-power error.
*Key Insight*: This is Chebyshev's inequality in deterministic measure form. □

**Example 4.2.11.** *$L^p$ Convergence Without Pointwise Convergence*
Let $X = [0, 1]$ with Lebesgue measure and list $I_{m,k} := [\frac{k-1}{m}, \frac{k}{m}]$, $1 \le k \le m$, row by row in $m$.
Let $f_n := \mathbf{1}_{I_n}$, where $I_n$ is the $n$th interval. If $I_n$ lies in row $m(n)$, then $m(n) \to \infty$ and
$\|f_n\|_p^p = \lambda(I_n) = \frac{1}{m(n)} \to 0$; hence $f_n \to 0$ in $L^p$, therefore in measure by Proposition 4.2.10.
$\forall x \in [0, 1]$, every row contains some $I_{m,k}$ with $x \in I_{m,k}$, giving a subsequence with value 1 at $x$; for infinitely many rows, some $I_{m,k}$ omits $x$, giving a subsequence with value 0 at $x$. Thus $(f_n(x))$ converges nowhere.
The subfamily $I_{m,2} = [\frac{1}{m}, \frac{2}{m}]$, $m \ge 2$, gives a subsequence converging pointwise to 0 on all of $X$.




Figure 218: In each row, one shrinking typewriter interval contains the fixed point $x$.

**Theorem 4.2.12.** *Riesz Subsequence Principle for Convergence in Measure*
*If $(f_n)$ is Cauchy in measure, then there exist a subsequence $(g_k)$ and a measurable $f$ such that:*

*(i)* $g_k \to f$ *a.e.*
*(ii)* $g_k \to f$ *in measure.*
*(iii)* $\forall \varepsilon > 0$ $\exists E \in \mathcal{M}$ *with* $\mu(E) < \varepsilon$ *such that* $g_k \to f$ *uniformly on* $X \setminus E$.

*Proof.*

(i) Choose $n_1 < n_2 < \cdots$ so that $m, n \ge n_k \Rightarrow \mu\{|f_m - f_n| \ge \frac{1}{2^k}\} < \frac{1}{2^k}$, and set $g_k := f_{n_k}$.
For $E_k := \{|g_{k+1} - g_k| \ge \frac{1}{2^k}\}$, $\mu(E_k) < \frac{1}{2^k}$. Set $F_k := \bigcup_{j=k}^{\infty} E_j$. Then $F_k \downarrow$, $\mu(F_k) \le \sum_{j=k}^{\infty} \frac{1}{2^j} = \frac{1}{2^{k-1}}$.
For $i > j \ge k$, $x \notin F_k$, $|g_i(x) - g_j(x)| \le \sum_{r=j}^{i-1} |g_{r+1}(x) - g_r(x)| < \frac{1}{2^{j-1}}$.
Hence $(g_j)$ is uniformly Cauchy on $X \setminus F_k$ $\forall k$.
Let $F := \bigcap_k F_k$, so $\mu(F) = 0$. Define $f = \lim_j g_j$ on $X \setminus F$, $f = 0$ on $F$. $f$ is measurable, $g_j \to f$ a.e.

(ii) Letting $i \to \infty$ gives $j \ge k$, $x \notin F_k \Rightarrow |f(x) - g_j(x)| \le \frac{1}{2^{j-1}} \le \frac{1}{2^{k-1}}$. Given $\alpha, \varepsilon > 0$, choose $k$ with $\frac{1}{2^{k-1}} < \alpha$ and $\mu(F_k) < \varepsilon$; then $j \ge k \Rightarrow \{|f - g_j| \ge \alpha\} \subseteq F_k$, so $g_j \to f$ in measure.

(iii) Given $\varepsilon > 0$, choose $k$ with $\mu(F_k) < \varepsilon$.
The estimate in (ii) gives $\sup_{x \in X \setminus F_k} |f(x) - g_j(x)| \le \frac{1}{2^{j-1}} \to 0$, so $g_j \to f$ uniformly on $X \setminus F_k$.

*Structural Commentary*: Cauchy in measure lets us select summably small exceptional sets.
*Key Insight*: Outside the tail union of exceptional sets, the subsequence is uniformly Cauchy. □

**Corollary 4.2.13.** *Completeness for Convergence in Measure*
*$(f_n)$ is Cauchy in measure iff $\exists f$ measurable with $f_n \to f$ in measure.*

*Proof.*

(⇒) If $(f_n)$ is Cauchy in measure, then $f_{n_k} \to f$ in measure for some measurable $f$ by Theorem 4.2.12.
For $\alpha > 0$, $\{|f_n - f| \ge \alpha\} \subseteq \{|f_n - f_{n_k}| \ge \frac{\alpha}{2}\} \cup \{|f_{n_k} - f| \ge \frac{\alpha}{2}\}$.
Choose $k$ large, then $n$ large, to get $f_n \to f$ in measure.
(⇐) For $\alpha > 0$, $\{|f_m - f_n| \ge \alpha\} \subseteq \{|f_m - f| \ge \frac{\alpha}{2}\} \cup \{|f_n - f| \ge \frac{\alpha}{2}\}$.
Hence $\mu\{|f_m - f_n| \ge \alpha\} \to 0$ as $m, n \to \infty$.

*Structural Commentary*: The subsequential in-measure limit attracts the whole Cauchy sequence.
*Key Insight*: Convergence in measure is complete after measurable limits are admitted. □

**Corollary 4.2.14.** *Uniqueness for Convergence in Measure*
*If $f_n \to f$ and $f_n \to g$ in measure, then $f = g$ a.e.*

*Proof.* For $\alpha > 0$, $\{|f - g| \ge \alpha\} \subseteq \{|f - f_n| \ge \frac{\alpha}{2}\} \cup \{|f_n - g| \ge \frac{\alpha}{2}\}$, so $\mu\{|f - g| \ge \alpha\} = 0$.
Hence $\mu\{|f - g| > 0\} = 0$ and $f = g$ a.e.
*Structural Commentary*: Two in-measure limits cannot differ on a positive-measure set.
*Key Insight*: In-measure limits are unique modulo null sets. □

**Corollary 4.2.15.** *In-Measure Limits Have A.E. Subsequences*
*If $f_n \to f$ in measure, then some subsequence $f_{n_k}$ converges to $f$ a.e.*

*Proof.* Since $f_n \to f$ in measure, $(f_n)$ is Cauchy in measure.
Choose a subsequence $f_{n_k} \to h$ a.e. and in measure by Theorem 4.2.12.
Then $h = f$ a.e. by Corollary 4.2.14; after enlarging the exceptional null set, $f_{n_k} \to f$ a.e.
*Structural Commentary*: Convergence in measure contains an almost-everywhere convergent subsequence.
*Key Insight*: Select a Cauchy-in-measure subsequence, then identify its limit by uniqueness. □

**Proposition 4.2.16.** *Subsequence and Indicator Consequences*

*(i) $f_n \to f$ in measure ⇒ every subsequence $f_{n_k} \to f$ in measure; if $(f_n)$ is Cauchy in measure, then every $(f_{n_k})$ is Cauchy in measure.*
*(ii) If $\|f_n - f\|_p \to 0$ and $\|f_{n_k} - g\|_p \to 0$, then $f = g$ a.e.*
*(iii) If $f_n = \mathbf{1}_{E_n}$ and $\|f_n - f\|_p \to 0$, then $f = \mathbf{1}_E$ a.e. for some $E \in \mathcal{M}$.*

*Proof.*

(i) For $\alpha > 0$, $\mu\{|f_{n_k} - f| \ge \alpha\} \to 0$ since $n_k \to \infty$; the Cauchy claim is identical with $m_k, n_\ell \to \infty$.
(ii) $f_n \to f$ and $f_{n_k} \to g$ in measure by Proposition 4.2.10; also $f_{n_k} \to f$ in measure by (i).
Thus $f = g$ a.e. by Corollary 4.2.14.
(iii) By Proposition 4.2.10 and Theorem 4.2.12, some $f_{n_k} \to f$ a.e.
Since $f_{n_k} \in \{0, 1\}$, $f \in \{0, 1\}$ a.e.; with $E := \{f = 1\}$, $E \in \mathcal{M}$ and $f = \mathbf{1}_E$ a.e.

*Structural Commentary*: Subsequence permanence and a.e. uniqueness transfer mode limits to representatives.
*Key Insight*: In-measure subsequences preserve limits; indicator limits remain indicators a.e. □

**Proposition 4.2.17.** *Algebra of Convergence in Measure on Finite Measure Spaces*
*Let $(X, \mathcal{M}, \mu)$ satisfy $\mu(X) < +\infty$. All functions below are measurable and finite-valued a.e.*

*(i) If $f_n \to f$ and $g_n \to g$ in measure, then $\forall a, b \in \mathbb{R}$, $af_n + bg_n \to af + bg$ in measure.*
*(ii) If $u_n \to 0$ in measure and $h$ is measurable and finite-valued a.e., then $hu_n \to 0$ in measure.*
*(iii) If $f_n \to f$ and $g_n \to g$ in measure, then $f_n g_n \to fg$ in measure.*
*(iv) If $f_n \to f$ in measure, then $f_n^2 \to f^2$ in measure.*

*Proof.*

(i) Fix $\alpha > 0$. Since $\{|a(f_n - f) + b(g_n - g)| \ge \alpha\} \subseteq \{|a(f_n - f)| \ge \frac{\alpha}{2}\} \cup \{|b(g_n - g)| \ge \frac{\alpha}{2}\}$, the measure of the right-hand side tends to 0 by convergence in measure of $f_n$ and $g_n$.
(ii) Fix $\alpha, \eta > 0$. Since $h$ is finite a.e. and $\mu(X) < +\infty$, choose $M > 0$ with $\mu\{|h| > M\} < \eta$.
Then $\{|hu_n| \ge \alpha\} \subseteq \{|h| > M\} \cup \{|u_n| \ge \frac{\alpha}{M}\}$, so $\limsup_n \mu\{|hu_n| \ge \alpha\} \le \eta$. Let $\eta \downarrow 0$.

(iii) Set $u_n := f_n - f$ and $v_n := g_n - g$. Then $u_n, v_n \to 0$ in measure and $f_n g_n - fg = f v_n + g u_n + u_n v_n$. By (ii), $f v_n \to 0$ and $g u_n \to 0$ in measure. For $\alpha > 0$, $\{|u_n v_n| \geq \alpha\} \subseteq \{|u_n| \geq \sqrt{\alpha}\} \cup \{|v_n| \geq \sqrt{\alpha}\}$, so $u_n v_n \to 0$ in measure. Apply (i) to the three summands.

(iv) Apply (iii) with $g_n = f_n$ and $g = f$.

*Structural Commentary*: Finite measure makes finite measurable multipliers harmless after truncation.
*Key Insight*: Products split into fixed-multiplier errors and a product of two small-in-measure errors. □

**Theorem 4.2.18.** *Dominated Convergence in Measure Gives $L^p$ Convergence*
*Let $1 \leq p < \infty$, $f_n \in L^p(\mu)$, $f_n \to f$ in measure, and $|f_n| \leq g$ a.e. for some $g \in L^p(\mu)$. Then:*

*(i)* $f \in L^p(\mu)$;
*(ii)* $\|f_n - f\|_p \to 0$.

*Proof.* Since $f_n \to f$ in measure, $(f_n)$ is Cauchy in measure.

(i) By Theorem 4.2.12, some subsequence $(u_j)$ satisfies $u_j \to h$ a.e. and in measure; $h = f$ a.e. by Corollary 4.2.14. Since $|u_j| \leq g$ a.e., $f \in L^p(\mu)$ by Theorem 4.2.4.

(ii) If norm convergence fails, then $\exists \varepsilon > 0$ and a subsequence $(g_k)$ with $\|g_k - f\|_p \geq \varepsilon$. The argument in (i) produces a subsubsequence $(h_j)$ with $h_j \to f$ a.e.; then $\|h_j - f\|_p \to 0$ by Theorem 4.2.4, a contradiction.

*Structural Commentary*: Reduce convergence in measure to almost-everywhere convergence along a subsequence.
*Key Insight*: Domination turns every in-measure subsequence into an $L^p$-convergent subsubsequence. □

**Lemma 4.2.19.** *Almost Uniform Cauchy Sequences Have Almost Uniform Limits*
*If $(f_n)$ is almost uniformly Cauchy, then $\exists f$ measurable such that $f_n \to f$ almost uniformly and a.e.*

*Proof.* Choose $E_k \in \mathcal{M}$ with $\mu(E_k) < 2^{-k}$ and $(f_n)$ uniformly Cauchy on $X \setminus E_k$. Set $F_k := \bigcup_{j=k}^{\infty} E_j$ and $F := \bigcap_k F_k$; then $F_k \downarrow$, $\mu(F_k) < 2^{-(k-1)}$, $\mu(F) = 0$, and $(f_n)$ is uniformly Cauchy on $X \setminus F_k$. By Theorem 7.1.7, $f_n$ has a uniform limit on each $X \setminus F_k$; these limits agree on overlaps, so define measurable $f$ on $X \setminus F$, and set $f = 0$ on $F$. Hence $f_n \to f$ a.e. and uniformly on each $X \setminus F_k$; since $\mu(F_k) \to 0$, $f_n \to f$ almost uniformly.
*Structural Commentary*: Tail unions convert many small exceptional sets into nested uniform domains.
*Key Insight*: Almost uniform Cauchy means uniformly Cauchy after discarding a tail-small exceptional set. □

**Theorem 4.2.20.** *Almost Uniform Convergence and Convergence in Measure*
*The following hold.*

*(i) If $f_n \to f$ almost uniformly, then $f_n \to f$ in measure.*
*(ii) If $f_n \to f$ in measure, then $\exists (f_{n_k})$ with $f_{n_k} \to f$ almost uniformly.*

*Proof.*

(i) If $f_n \rightrightarrows f$ on $X \setminus E_\delta$ with $\mu(E_\delta) < \delta$, then $\forall \alpha > 0$ and sufficiently large $n$, $\{|f_n - f| \geq \alpha\} \subseteq E_\delta$. Thus $\limsup_n \mu\{|f_n - f| \geq \alpha\} \leq \delta$, and let $\delta \downarrow 0$.

(ii) Convergence in measure makes $(f_n)$ Cauchy in measure. Choose $f_{n_k} \to h$ almost uniformly and in measure by Theorem 4.2.12. Uniqueness gives $h = f$ a.e.; absorb $\{h \neq f\}$ into the exceptional sets.

*Structural Commentary*: Almost uniform convergence is uniform convergence off arbitrarily small sets.
*Key Insight*: Convergence in measure always hides an almost-uniform subsequence. □

**Corollary 4.2.21.** *Finite-Measure Almost-Everywhere Convergence Implies In-Measure Convergence*
*If $\mu(X) < +\infty$ and $f_n \to f$ a.e., with $f_n, f$ finite-valued off a null set, then $f_n \to f$ in measure.*

*Proof.* By Theorem 1.6.3, $f_n \to f$ almost uniformly; hence $f_n \to f$ in measure by Theorem 4.2.20.
*Structural Commentary*: Finite measure upgrades a.e. convergence through almost uniform convergence.
*Key Insight*: On finite measure spaces, a.e. convergence implies in-measure convergence. □

**Corollary 4.2.22.** *Finite-Measure Subsequence Test for Convergence in Measure*
*Let $(X, \mathcal{M}, \mu)$ satisfy $\mu(X) < +\infty$, and let $f_n, f : X \to \overline{\mathbb{R}}$ be measurable and finite-valued a.e.*
*Then $f_n \to f$ in measure iff every subsequence of $(f_n)$ has a further subsequence converging to $f$ a.e.*

*Proof.*

($\Rightarrow$) If $f_n \to f$ in measure, then every subsequence also converges to $f$ in measure by Proposition 4.2.16. Choose a further subsequence converging to $f$ a.e. by Corollary 4.2.15.

($\Leftarrow$) Suppose $f_n \nrightarrow f$ in measure. Then $\exists \alpha, \varepsilon > 0$ and subsequence $(f_{n_k})$ with $\mu\{|f_{n_k} - f| \geq \alpha\} \geq \varepsilon \ \forall k$. But some $f_{n_{k_j}} \to f$ a.e., hence in measure by Corollary 4.2.21, contradicting the uniform lower bound.

*Structural Commentary*: Subsequence tests characterize in-measure convergence.
*Key Insight*: Nonconvergence leaves a bad subsequence. □




Figure 219: Modes of convergence in three settings: solid arrows are implications; dashed arrows yield convergent subsequences.

**Method 4.2.23.** *Counterexample Dictionary*

(i) On $(\mathbb{R}, \lambda)$, $f_n = n^{-\frac{1}{p}} \mathbf{1}_{[0,n]}$ satisfies $f_n \rightrightarrows 0$ but $\|f_n\|_p = 1$.
(ii) On $([0,1], \lambda)$, $f_n = n\mathbf{1}_{[\frac{1}{n}, \frac{2}{n}]}$ satisfies $f_n \to 0$ a.e., in measure, almost uniformly, but $\|f_n\|_p^p = n^{p-1} \nrightarrow 0$.
(iii) Typewriter intervals on $[0,1]$: if $f_n = \mathbf{1}_{I_n}$ and $\lambda(I_n) \to 0$ rowwise, then $\|f_n\|_p \to 0$ and $f_n \to 0$ in measure, but $(f_n(x))$ converges nowhere.
(iv) On $(\mathbb{R}, \lambda)$, $f_n = \mathbf{1}_{[n,n+1]}$ satisfies $f_n \to 0$ pointwise but $\forall n,\ \lambda\{|f_n| \geq \frac{1}{2}\} = 1$, so $f_n \nrightarrow 0$ in measure.
(v) On $(\mathbb{R}, \lambda)$, $f_n = \mathbf{1}_{[n,n+1]}$ is not almost uniformly convergent to 0: if $\lambda(E) < 1$ and $f_n \rightrightarrows 0$ on $\mathbb{R} \setminus E$, then $[n, n+1] \subseteq E$ for sufficiently large $n$, impossible.

**Proposition 4.2.24.** *An Integral Metric for Convergence in Measure on Finite Measure Spaces*
*Let $\mu(X) < +\infty$ and $r(h) := \int_X |h|\,(1 + |h|)^{-1}\, d\mu$. Then $f_n \to f$ in measure iff $r(f_n - f) \to 0$.*

*Proof.*

($\Rightarrow$) Set $h_n := f_n - f$. For $\eta > 0$ and $A_n := \{|h_n| \geq \eta\}$, $r(h_n) \leq \mu(A_n) + \eta\mu(X)$. Let $n \to \infty$, then $\eta \downarrow 0$.
($\Leftarrow$) For $\alpha > 0$, $\alpha(1+\alpha)^{-1}\mathbf{1}_{\{|h_n| \geq \alpha\}} \leq |h_n|\,(1 + |h_n|)^{-1}$, hence $\mu\{|h_n| \geq \alpha\} \leq (1+\alpha)\alpha^{-1} r(h_n) \to 0$.

*Structural Commentary*: The bounded error transform metrizes convergence in measure on finite measure spaces.
*Key Insight*: Small integral of the bounded error transform is equivalent to small large-error sets. □

**Corollary 4.2.25.** *Fatou and Dominated Convergence with In-Measure Hypotheses*

*(i) $f_n \geq 0$ and $f_n \to f$ in measure $\Rightarrow \int_X f\, d\mu \leq \liminf_n \int_X f_n\, d\mu$.*
*(ii) $f_n \to f$ in measure and $|f_n| \leq g \in \mathscr{L}^1(\mu) \Rightarrow \int_X f_n\, d\mu \to \int_X f\, d\mu$.*

*Proof.*

(i) Choose $f_{n_k}$ with $\int f_{n_k}\, d\mu \to \liminf_n \int f_n\, d\mu$. By Theorem 4.2.12 and uniqueness, some $f_{n_{k_j}} \to f$ a.e. Fatou gives $\int f\, d\mu \leq \liminf_j \int f_{n_{k_j}}\, d\mu = \liminf_n \int f_n\, d\mu$.
(ii) By Theorem 4.2.18 with $p = 1$, $\|f_n - f\|_1 \to 0$, hence $\left|\int_X (f_n - f)\, d\mu\right| \leq \|f_n - f\|_1 \to 0$.

*Structural Commentary*: In-measure hypotheses reduce to a.e. subsequences or $L^1$ convergence.
*Key Insight*: Fatou and DCT survive convergence in measure through subsequences and domination. □

The next criterion packages the local mass controls needed to reverse convergence in measure into norm convergence.

**Theorem 4.2.26.** *Vitali Convergence Criterion for $L^p$*
*Let $1 \leq p < \infty$, $f_n \in L^p(\mu)$, and let $f$ be measurable. Consider the following conditions.*

*(i) $f \in L^p(\mu)$ and $\|f_n - f\|_p \to 0$.*
*(ii) $f_n \to f$ in measure.*
*(iii) $\forall \varepsilon > 0\ \exists E_\varepsilon \in \mathcal{M}$ with $\mu(E_\varepsilon) < +\infty$ such that $F \cap E_\varepsilon = \emptyset \Rightarrow \sup_n \int_F |f_n|^p\, d\mu < \varepsilon^p$.*
*(iv) $\forall \varepsilon > 0\ \exists \delta(\varepsilon) > 0$ such that $\mu(E) < \delta(\varepsilon) \Rightarrow \sup_n \int_E |f_n|^p\, d\mu < \varepsilon^p$.*

*Precisely, (i) is equivalent to the conjunction of (ii)–(iv).*

*Proof.*

(i) Assume (i). Then (ii) holds by Proposition 4.2.10, and $(f_n)$ is $L^p$-Cauchy. Conditions (iii)–(iv) follow from Proposition 4.1.40 after raising its norm bounds to the $p$th power.

(ii) Assume (ii)–(iv). Fix $\eta > 0$, choose $E_\eta$ from (iii) and $\delta(\eta)$ from (iv), and put $F = X \setminus E_\eta$.
Then $\|(f_n - f_m)\mathbf{1}_F\|_p \le \|f_n \mathbf{1}_F\|_p + \|f_m \mathbf{1}_F\|_p < 2\eta$.
If $\mu(E_\eta) > 0$, set $\alpha := \eta\mu(E_\eta)^{-\frac{1}{p}}$ and $H_{nm} := E_\eta \cap \{|f_n - f_m| \ge \alpha\}$.
By (ii), $(f_n)$ is Cauchy in measure, so $\mu(H_{nm}) < \delta(\eta)$ for large $m, n$. On $E_\eta \setminus H_{nm}$, $|f_n - f_m| < \alpha$; on $H_{nm}$, use $|f_n - f_m| \le |f_n| + |f_m|$. Hence $\|(f_n - f_m)\mathbf{1}_{E_\eta}\|_p \le \alpha\mu(E_\eta)^{\frac{1}{p}} + \|f_n \mathbf{1}_{H_{nm}}\|_p + \|f_m \mathbf{1}_{H_{nm}}\|_p < 3\eta$ by (iv). If $\mu(E_\eta) = 0$, the $E_\eta$ term is 0. Thus $\|f_n - f_m\|_p < 5\eta$ eventually, so $(f_n)$ is $L^p$-Cauchy.
By Riesz–Fischer, $f_n \to h$ in $L^p$; by Proposition 4.2.10 and Corollary 4.2.14, $h = f$ a.e., so (i) holds.

*Structural Commentary*: Vitali = in-measure convergence + tightness + uniform small-set control.
*Key Insight*: Tightness and small-set control convert in-measure convergence into an $L^p$ Cauchy estimate. □

## 4.3 Composition and Convergence Modes

**Proposition 4.3.1.** *Composition and Pointwise Modes*
*Let $\varphi : \mathbb{R} \to \mathbb{R}$.*

*(i) $\varphi$ continuous, $f_n \to f$ a.e. $\Rightarrow \varphi \circ f_n \to \varphi \circ f$ a.e.*
*(ii) $\varphi$ uniformly continuous preserves uniform, a.u., and in-measure convergence.*
*(iii) If $\varphi$ is discontinuous, composition need not preserve a.e. convergence.*
*(iv) Non-uniform continuity can destroy uniform, even in-measure, convergence.*

*Proof.*

(i) Outside a null set, $f_n(x) \to f(x)$; continuity gives $\varphi(f_n(x)) \to \varphi(f(x))$.
(ii) Given $\varepsilon > 0$, choose $\delta > 0$ with $|s - t| < \delta \Rightarrow |\varphi(s) - \varphi(t)| < \varepsilon$. Uniform and almost-uniform convergence use $\sup |f_n - f| < \delta$ on $X$ or $X \setminus E_\eta$; in measure, $\{|\varphi(f_n) - \varphi(f)| \ge \varepsilon\} \subseteq \{|f_n - f| \ge \delta\}$.
(iii) If $\varphi$ is discontinuous at $a$, choose $t_n \to a$ and $\alpha > 0$ with $|\varphi(t_n) - \varphi(a)| \ge \alpha$.
On $X = \{*\}$, $f_n(*) = t_n \to a = f(*)$, but $\varphi(f_n(*)) \not\to \varphi(f(*))$.
(iv) If $\varphi$ is not uniformly continuous, choose $a_n, b_n$ with $|a_n - b_n| \to 0$ and $|\varphi(a_n) - \varphi(b_n)| \ge \alpha > 0$.
On $(\mathbb{N}, \#)$, set $f(k) = b_k$ and $f_n(k) = a_n$ for $k = n$, $f_n(k) = b_k$ for $k \ne n$.
Then $\|f_n - f\|_\infty \to 0$, but $\forall n$, $\#\{|\varphi(f_n) - \varphi(f)| \ge \alpha\} = 1$.

*Structural Commentary*: Continuity is pointwise; uniform continuity gives one global $\delta$.
*Key Insight*: Composition needs the same uniformity as the convergence mode. □

**Proposition 4.3.2.** *Composition and Convergence in Measure*
*Let $f_n, f : X \to \mathbb{R}$ be measurable with $f_n \to f$ in measure, and let $\varphi : \mathbb{R} \to \mathbb{R}$.*

*(i) If $\varphi$ is uniformly continuous, then $\forall \varepsilon > 0$, $\mu\{|\varphi(f_n) - \varphi(f)| \ge \varepsilon\} \to 0$.*
*(ii) If $\mu(X) < +\infty$ and $\varphi$ is continuous, then $\forall \varepsilon > 0$, $\mu\{|\varphi(f_n) - \varphi(f)| \ge \varepsilon\} \to 0$.*

*Proof.*

(i) Fix $\varepsilon > 0$ and choose $\delta > 0$ with $|s - t| < \delta \Rightarrow |\varphi(s) - \varphi(t)| < \varepsilon$.
Then $\{|\varphi(f_n) - \varphi(f)| \ge \varepsilon\} \subseteq \{|f_n - f| \ge \delta\}$, so the measures tend to 0.
(ii) Fix $\varepsilon, \eta > 0$. Since $\mu(X) < +\infty$ and $f$ is finite-valued, choose $M$ with $\mu\{|f| > M\} < \eta$. Uniform continuity of $\varphi$ on $[-M-1, M+1]$ gives $0 < \delta < 1$ such that $s, t \in [-M-1, M+1]$ and $|s - t| < \delta \Rightarrow |\varphi(s) - \varphi(t)| < \varepsilon$.
Hence $\{|\varphi(f_n) - \varphi(f)| \ge \varepsilon\} \subseteq \{|f| > M\} \cup \{|f_n - f| \ge \delta\}$, so $\limsup_n \mu\{|\varphi(f_n) - \varphi(f)| \ge \varepsilon\} \le \eta \downarrow 0$.

*Structural Commentary*: Use global uniformity, or truncate to a bounded range.
*Key Insight*: On finite measure spaces, measurable finite-valued functions are almost bounded. □

**Example 4.3.3.** *Continuity Alone Is Not Enough on Infinite Measure*
On $(\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda)$, set $f(x) = x$, $f_n(x) = x + \frac{1}{n}$, and $\varphi(t) = t^2$.
Then $\|f_n - f\|_\infty = \frac{1}{n} \to 0$, hence $f_n \rightrightarrows f$ and $f_n \to f$ in measure.
However $\varphi(f_n) - \varphi(f) = \frac{2x}{n} + \frac{1}{n^2}$; for $E_n := \{|\varphi(f_n) - \varphi(f)| \ge 1\}$, $[\frac{n}{2}, +\infty) \subseteq E_n$, so $\lambda(E_n) = +\infty$.
Thus $\varphi \circ f_n \not\to \varphi \circ f$ in measure.

**Proposition 4.3.4.** *Growth Conditions for $L^p$ Composition*
*Let $1 \le p < \infty$, let $\varphi : \mathbb{R} \to \mathbb{R}$ be measurable, and let $f_n, f : X \to \mathbb{R}$ be measurable.*

*(i) If $\exists A, B \ge 0$ such that $\forall t$, $|\varphi(t)| \le A|t| + B$, $\mu(X) < +\infty$, and $f \in L^p(\mu)$, then $\varphi \circ f \in L^p(\mu)$.*
*(ii) If $\exists A \ge 0$ such that $\forall t$, $|\varphi(t)| \le A|t|$, and $f \in L^p(\mu)$, then $\varphi \circ f \in L^p(\mu)$.*
*(iii) If $\mu(X) < +\infty$, $\varphi$ is uniformly continuous, $f_n, f \in L^p(\mu)$, and $\|f_n - f\|_p \to 0$, then $\|\varphi \circ f_n - \varphi \circ f\|_p \to 0$.*
*(iv) If $\varphi$ is Lipschitz, $\varphi(0) = 0$, $f_n, f \in L^p(\mu)$, and $\|f_n - f\|_p \to 0$, then $\|\varphi \circ f_n - \varphi \circ f\|_p \to 0$.*

*Proof.*

(i) $|\varphi(f)|^p \le 2^{p-1}(A^p |f|^p + B^p)$, so $\|\varphi \circ f\|_p^p \le 2^{p-1}(A^p \|f\|_p^p + B^p \mu(X)) < +\infty$.

(ii) $|\varphi(f)|^p \le A^p |f|^p$, so $\|\varphi \circ f\|_p \le A \|f\|_p < +\infty$.

(iii) Uniform continuity gives constants $C, D \ge 0$ with $|\varphi(t)| \le C|t| + D$ and $|\varphi(s) - \varphi(t)| \le C|s-t| + D$. Hence all compositions lie in $L^p$ by (i). Let $h_n := |f_n - f|$.
Given $\eta > 0$, choose $\delta > 0$ with $|s-t| < \delta \Rightarrow |\varphi(s) - \varphi(t)| < \eta$ and set $H_n := \{h_n \ge \delta\}$.
Then $\int_X |\varphi(f_n) - \varphi(f)|^p \, d\mu \le \eta^p \mu(X) + 2^{p-1} \int_{H_n} (C^p h_n^p + D^p) \, d\mu$.
Since $\int h_n^p \, d\mu \to 0$ and $\mu(H_n) \le \delta^{-p} \int h_n^p \, d\mu \to 0$, take $\limsup_n$ and then $\eta \downarrow 0$.

(iv) If $L$ is Lipschitz, $|\varphi(f)| \le L|f|$ and $\|\varphi \circ f_n - \varphi \circ f\|_p \le L \|f_n - f\|_p \to 0$.

*Structural Commentary*: Growth gives membership; continuity estimates transfer $L^p$ convergence.
*Key Insight*: Finite measure makes small uniform errors integrable; Lipschitz bounds give norm control. □

**Example 4.3.5.** *Uniform Continuity Does Not Suffice on Infinite Measure*
On $(\mathbb{R}, \lambda)$, fix $1 \le p < \infty$, set $f = 0$, $f_n := n^{-1} \mathbf{1}_{[0, n^{\frac{p}{2}}]}$, and $\varphi(t) := \sqrt{|t|}$.
Then $\varphi$ is uniformly continuous, $\varphi(0) = 0$, $\|f_n - f\|_p^p = n^{-\frac{p}{2}} \to 0$, but $\forall n$, $\|\varphi \circ f_n - \varphi \circ f\|_p^p = 1$.
Thus uniform continuity need not preserve $L^p$ convergence on infinite measure spaces.

**Lemma 4.3.6.** *$L^p$ Convergence Allows Continuous Composition in Measure*
*Let $1 \le p < \infty$, $f_n, f \in L^p(\mu)$, $\|f_n - f\|_p \to 0$, and let $\varphi : \mathbb{R} \to \mathbb{R}$ be continuous.*
*Then $\varphi \circ f_n \to \varphi \circ f$ in measure.*

*Proof.* Fix $\alpha, \eta > 0$. Choose $M$ with $\mu\{|f| > M\} < \eta$. By uniform continuity of $\varphi$ on $[-M-1, M+1]$, choose $0 < \delta < 1$ such that $s, t \in [-M-1, M+1]$ and $|s-t| < \delta \Rightarrow |\varphi(s) - \varphi(t)| < \alpha$.
Then $\{|\varphi(f_n) - \varphi(f)| \ge \alpha\} \subseteq \{|f| > M\} \cup \{|f_n - f| \ge \delta\}$, so $\limsup_n \mu\{|\varphi(f_n) - \varphi(f)| \ge \alpha\} \le \eta$. Let $\eta \downarrow 0$.

*Structural Commentary*: $L^p$ convergence localizes to a large bounded range where continuity is uniform.
*Key Insight*: Norm convergence gives in-measure control; bounded-range continuity transfers it. □

**Theorem 4.3.7.** *Composition Growth Criteria in $L^p$*
*Let $1 \le p < \infty$ and let $\varphi : \mathbb{R} \to \mathbb{R}$ be continuous.*

*(i) If $\mu(X) < +\infty$ and $\exists K > 0$ such that $|\varphi(t)| \le K|t|$ for $|t| \ge K$, then $f \in L^p(\mu) \Rightarrow \varphi \circ f \in L^p(\mu)$.*
*(ii) Under (i), if $f_n, f \in L^p(\mu)$ and $\|f_n - f\|_p \to 0$, then $\|\varphi \circ f_n - \varphi \circ f\|_p \to 0$.*
*(iii) If $\exists K \ge 0$ such that $\forall t$, $|\varphi(t)| \le K|t|$, then $f \in L^p(\mu) \Rightarrow \varphi \circ f \in L^p(\mu)$ on any measure space.*
*(iv) Under (iii), if $f_n, f \in L^p(\mu)$ and $\|f_n - f\|_p \to 0$, then $\|\varphi \circ f_n - \varphi \circ f\|_p \to 0$ on any measure space.*

*Proof.*

(i) Let $M := \sup_{|t| \le K} |\varphi(t)|$, $C := K + M$. Then $\forall t$, $|\varphi(t)| \le C(1 + |t|)$; hence $|\varphi(f)|^p \le 2^{p-1} C^p (1 + |f|^p) \in L^1(\mu)$.

(ii) Let $C$ be as in (i). Then $\varphi \circ f_n \to \varphi \circ f$ in measure by Lemma 4.3.6.
Since $(f_n)$ is $L^p$-Cauchy, Proposition 4.1.40 and the growth bound from (i) supply the finite-tail and small-set estimates. Thus $\|\varphi \circ f_n - \varphi \circ f\|_p \to 0$ by Theorem 4.2.26.

(iii) $|\varphi(f)|^p \le K^p |f|^p \in L^1(\mu)$.

(iv) $\varphi \circ f_n \to \varphi \circ f$ in measure by Lemma 4.3.6. Since $(f_n)$ is $L^p$-Cauchy and $|\varphi(f_n)|^p \le K^p |f_n|^p$, the finite-tail and small-set estimates for $(\varphi \circ f_n)$ follow from Proposition 4.1.40.
Thus $\|\varphi \circ f_n - \varphi \circ f\|_p \to 0$ by Theorem 4.2.26.

*Structural Commentary*: Local growth needs finite measure; global growth does not.
*Key Insight*: Vitali converts in-measure composition plus inherited mass control into $L^p$ convergence. □

**Method 4.3.8.** *Atomic Counterexamples for Composition Growth*

(i) If finite-measure tail growth fails, choose $|t_n| \ge n$ with $|\varphi(t_n)| > n^2 |t_n|$. On $\mathbb{N}$, set $\mu(\{n\}) = (n^2 |t_n|^p)^{-1}$ and $f(n) = t_n$. Then $\mu(\mathbb{N}) < +\infty$, $\int |f|^p \, d\mu = \sum_n n^{-2} < +\infty$, but $\int |\varphi(f)|^p \, d\mu \ge \sum_n n^{2p-2} = +\infty$.

(ii) If the same tail growth fails, choose $|t_n| \ge n$ with $|\varphi(t_n) - \varphi(0)| \ge n|t_n|$. With $\mu(\{n\}) = (n^p |t_n|^p)^{-1}$, $f = 0$, and $f_n = t_n \mathbf{1}_{\{n\}}$, one has $\mu(\mathbb{N}) < +\infty$, $\|f_n - f\|_p = \frac{1}{n} \to 0$, but $\|\varphi \circ f_n - \varphi \circ f\|_p \ge 1$.

(iii) If global linear growth fails and $\varphi(0) = 0$, the constructions in (i)–(ii), without requiring $\mu(\mathbb{N}) < +\infty$, give membership and convergence counterexamples on arbitrary measure spaces.

(iv) If $\varphi(0) \ne 0$, then on $(\mathbb{N}, \#)$, $0 \in L^p$ but $\varphi \circ 0$ is a nonzero constant and hence not in $L^p$; the constant sequence $f_n = 0$ gives the convergence failure.

## 4.4 Uniform Integrability and Vitali Convergence

Uniform integrability is the $L^1$ condition that rules out concentration of mass on tiny sets. On infinite-measure spaces it must be paired with tightness, which rules out mass escaping to remote finite-measure tails.

**Lemma 4.4.1.** *Finite-Measure Lebesgue Sets Have Small Pieces*
*Let $E \in \mathcal{L}^1$, $m(E) < +\infty$, and $\delta > 0$. Then $\exists E_1, \dots, E_N \in \mathcal{L}^1$ such that:*

*(i)* $E_i \cap E_j = \emptyset$ *for* $i \neq j$.
*(ii)* $E = \bigsqcup_{k=1}^{N} E_k$.
*(iii)* $m(E_k) < \delta$ *for* $1 \leq k \leq N$.

*Proof.* Choose $n_0$ with $T := E \setminus [-n_0, n_0]$ and $m(T) < \delta$. Partition $[-n_0, n_0]$ into finitely many disjoint intervals $I_j$ with $m(I_j) < \delta$, and let $(E_k)$ be the nonempty sets among $T$ and $E \cap I_j$.

(i) Since $T \cap [-n_0, n_0] = \emptyset$ and the $I_j$ are disjoint, the $E_k$ are pairwise disjoint.

(ii) $E = T \bigsqcup (E \cap [-n_0, n_0]) = T \bigsqcup \bigsqcup_j (E \cap I_j) = \bigsqcup_{k=1}^{N} E_k$.

(iii) $m(T) < \delta$ and $m(E \cap I_j) \leq m(I_j) < \delta$.

*Structural Commentary*: Finite measure gives a small tail $T$; a fine partition makes $E \setminus T$ small-piece.
*Key Insight*: One small tail plus a fine finite grid gives a finite small-piece decomposition. □

**Proposition 4.4.2.** *$L^1$ Absolute Continuity on Small Sets*
*Let $E \in \mathcal{L}^1$, let $\mathcal{L}^1_E = \{A \in \mathcal{L}^1 : A \subseteq E\}$, let $m_E = m|_{\mathcal{L}^1_E}$, and let $f : E \to \mathbb{R}$ be measurable.*
*Consider the condition $(*)$: $\forall \varepsilon > 0\ \exists \delta > 0$ such that $A \in \mathcal{L}^1_E$ and $m(A) < \delta$ imply $\int_A |f|\, dm < \varepsilon$.*

*(i) If $f \in L^1(m_E)$, then $(*)$ holds.*
*(ii) If $m(E) < +\infty$ and $(*)$ holds, then $f \in L^1(m_E)$.*

*Proof.*

(i) Since $f \in L^1(m_E)$, Lemma 4.1.38 on $(E, \mathcal{L}^1_E, m_E)$ with $p = 1$ gives $\forall \varepsilon > 0\ \exists \delta > 0$ such that $A \in \mathcal{L}^1_E$ and $m_E(A) < \delta$ imply $\int_A |f|\, dm_E < \varepsilon$. Since $m_E(A) = m(A)$ and $\int_A |f|\, dm_E = \int_A |f|\, dm$, $(*)$ holds.

(ii) Choose $\delta > 0$ from $(*)$ with $\varepsilon = 1$. By Lemma 4.4.1, write $E = \bigsqcup_{k=1}^{N} E_k$ with $E_k \in \mathcal{L}^1_E$ and $m(E_k) < \delta$. Then $(*)$ gives $\int_{E_k} |f|\, dm < 1$ for each $k$, hence $\int_E |f|\, dm = \sum_{k=1}^{N} \int_{E_k} |f|\, dm < N < +\infty$.

*Structural Commentary*: $L^1$ gives small-set control; finite partitions recover $L^1$.
*Key Insight*: A finite small-piece decomposition turns local integral bounds into one global finite integral. □

**Definition 4.4.3.** *Uniform Integrability and Tightness*
Let $\mathcal{F}$ be a family of measurable functions $h : X \to \mathbb{R}$ on $(X, \mathcal{M}, \mu)$.

(i) $\mathcal{F}$ is *uniformly integrable* on $X$ if $\forall \varepsilon > 0\ \exists \delta > 0$ such that $A \in \mathcal{M}$ and $\mu(A) < \delta$ imply $\sup_{h \in \mathcal{F}} \int_A |h|\, d\mu < \varepsilon$.

(ii) $\mathcal{F}$ is *tight* on $X$ if $\forall \varepsilon > 0\ \exists X_\varepsilon \in \mathcal{M}$ with $\mu(X_\varepsilon) < +\infty$ and $\sup_{h \in \mathcal{F}} \int_{X \setminus X_\varepsilon} |h|\, d\mu < \varepsilon$.

On finite measure spaces, tightness is automatic by taking $X_\varepsilon = X$.
*Key Insight*: UI controls mass on small $A$; tightness controls mass on $X \setminus X_\varepsilon$.

**Example 4.4.4.** *Concentration versus Escape*

(i) *Concentration*: on $(0, 1)$, $u_n := n\mathbf{1}_{(0, \frac{1}{n})}$. For every $\delta > 0$, choose $n > \delta^{-1}$ and $A_n = (0, \frac{1}{n})$. Then $m(A_n) < \delta$ but $\int_{A_n} u_n\, dm = 1$, so $(u_n)$ is not uniformly integrable.

(ii) *Escape*: on $\mathbb{R}$, let $I_n := (n, n+1)$ and $v_n := \mathbf{1}_{I_n}$. Since $\int_A v_n\, dm \leq m(A)$, $(v_n)$ is uniformly integrable. If $m(E_0) < +\infty$ and $\sup_n \int_{\mathbb{R} \setminus E_0} v_n\, dm < \frac{1}{2}$, then $m(E_0 \cap I_n) > \frac{1}{2}$ for all $n$, contradiction.

*Structural Commentary*: Concentration occurs even on $m(X) < +\infty$; escape is an $m(X) = +\infty$ obstruction.
*Key Insight*: Uniform integrability controls small-set mass; tightness keeps mass in finite-measure cores.

**Proposition 4.4.5.** *Basic Sources of Uniform Integrability and Tightness*
*Let $(X, \mathcal{M}, \mu)$ be a measure space.*

*(i) If $0 \leq g \in L^1(\mu)$ and $\forall h \in \mathcal{F}$, $|h| \leq g$ a.e., then $\mathcal{F}$ is uniformly integrable and tight.*
*(ii) If $f_1, \dots, f_N \in L^1(\mu)$, then $\{f_1, \dots, f_N\}$ is uniformly integrable and tight.*

*Proof.* Let $\varepsilon > 0$.

(i) Choose $\delta > 0$ with $\mu(A) < \delta \Rightarrow \int_A g\,d\mu < \varepsilon$ by Lemma 4.1.38; hence $\sup_{h\in\mathcal{F}} \int_A |h|\,d\mu < \varepsilon$.
Choose finite-measure $X_\varepsilon$ with $\int_{X\setminus X_\varepsilon} g\,d\mu < \varepsilon$ by Proposition 1.3.22; hence $\sup_{h\in\mathcal{F}} \int_{X\setminus X_\varepsilon} |h|\,d\mu < \varepsilon$.

(ii) For each $j$, apply (i) to $\{f_j\}$ with $\varepsilon$: choose $\delta_j > 0$ such that $\mu(A) < \delta_j \Rightarrow \int_A |f_j|\,d\mu < \varepsilon$, and choose $X_{\varepsilon,j}$ with $\mu(X_{\varepsilon,j}) < +\infty$ and $\int_{X\setminus X_{\varepsilon,j}} |f_j|\,d\mu < \varepsilon$. Set $\delta := \min_{1\le j\le N} \delta_j$ and $X_\varepsilon := \bigcup_{j=1}^{N} X_{\varepsilon,j}$.
Then $\mu(A) < \delta \Rightarrow \sup_j \int_A |f_j|\,d\mu < \varepsilon$, while $\mu(X_\varepsilon) < +\infty$ and $\sup_j \int_{X\setminus X_\varepsilon} |f_j|\,d\mu < \varepsilon$.

*Structural Commentary*: An envelope gives common bounds; finite families use $\min_j \delta_j$ and $\bigcup_j X_{\varepsilon,j}$.

*Key Insight*: Domination gives uniform estimates; finite families allow minima and finite unions. □

**Corollary 4.4.6.** *$L^p$-Bounded Families Are Uniformly Integrable*
*Let $1 < p < \infty$, $q$ conjugate to $p$. If $\mathcal{F} \subseteq L^p(\mu)$ and $\forall h \in \mathcal{F},\ \|h\|_p \le M$, then $\mathcal{F}$ is uniformly integrable.*

*Proof.* If $M = 0$, every $h \in \mathcal{F}$ vanishes a.e. and the claim is immediate. Else, for measurable $A$ with $\mu(A) < +\infty$, by Hölder's inequality, $\int_A |h|\,d\mu = \int_X |h|\,\mathbf{1}_A\,d\mu \le \|h\|_p\,\|\mathbf{1}_A\|_q \le M\mu(A)^{\frac{1}{q}}$.
Given $\varepsilon > 0$, take $\delta = (\frac{\varepsilon}{M})^q$. Then $\mu(A) < \delta$ implies $\sup_{h\in\mathcal{F}} \int_A |h|\,d\mu < \varepsilon$.

*Structural Commentary*: Apply Hölder to $h\mathbf{1}_A$ and choose one measure threshold for the whole family.
*Key Insight*: Hölder converts a uniform $L^p$ bound into uniform small-set $L^1$ control. □

**Proposition 4.4.7.** *Uniform Integrability by Large-Value Tails*
*Let $(X, \mathcal{M}, \mu)$ be a finite measure space and let $(f_n)$ be bounded in $L^1(\mu)$. The following are equivalent.*

*(i)* *$(f_n)$ is uniformly integrable.*
*(ii)* $\forall \varepsilon > 0\ \exists M > 0 :\ \sup_n \int_{\{|f_n|\ge M\}} |f_n|\,d\mu < \varepsilon$.

*Proof.* Let $C = \sup_n \|f_n\|_1 < +\infty$.

(⇒) Given $\varepsilon > 0$, choose $\delta > 0$ from uniform integrability and $M > C/\delta$.
By Chebyshev, $\mu(\{|f_n| \ge M\}) \le M^{-1}\|f_n\|_1 \le C/M < \delta$, so $\sup_n \int_{\{|f_n|\ge M\}} |f_n|\,d\mu < \varepsilon$.

(⇐) Given $\varepsilon > 0$, choose $M > 0$ with $\sup_n \int_{\{|f_n|\ge M\}} |f_n|\,d\mu < \frac{\varepsilon}{2}$ and set $\delta := \frac{\varepsilon}{2M}$.
If $\mu(A) < \delta$, then $\int_A |f_n|\,d\mu \le \int_{\{|f_n|\ge M\}} |f_n|\,d\mu + M\mu(A) < \varepsilon$.

*Structural Commentary*: Chebyshev shrinks tails; truncation recovers UI.
*Key Insight*: UI iff large tails vanish uniformly. □

**Theorem 4.4.8.** *Vitali Convergence for $L^1$ Families*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, let $f_n \in L^1(\mu)$, let $f : X \to \mathbb{R}$ be measurable, and assume $f_n \to f$ a.e.*

*(i)* *If $\mu(X) < +\infty$ and $\{f_n : n \ge 1\}$ is uniformly integrable, then $f \in L^1(\mu)$ and $\|f_n - f\|_1 \to 0$.*
*(ii)* *If $\{f_n : n \ge 1\}$ is uniformly integrable and tight, then $f \in L^1(\mu)$ and $\|f_n - f\|_1 \to 0$.*
*(iii)* *Under either hypothesis, $\int_X f_n\,d\mu \to \int_X f\,d\mu$.*

*Proof.*

(i) Since $\mu(X) < +\infty$ and $f_n \to f$ a.e., $f_n \to f$ in measure by Corollary 4.2.21. Uniform integrability is the small-set hypothesis of Theorem 4.2.26 with $p = 1$; the finite-tail hypothesis holds with $E_\varepsilon = X$, QED.

(ii) Fix $\alpha, \eta > 0$ and set $B_n^\alpha := \{|f_n - f| \ge \alpha\}$.
By tightness, choose $X_0 \in \mathcal{M}$ with $\mu(X_0) < +\infty$, $\sup_n \int_{X\setminus X_0} |f_n|\,d\mu < \frac{\alpha\eta}{4}$. By Fatou, $\int_{X\setminus X_0} |f|\,d\mu \le \frac{\alpha\eta}{4}$.
Then $\alpha\mu(B_n^\alpha \cap (X \setminus X_0)) \le \int_{X\setminus X_0} (|f_n| + |f|)\,d\mu < \frac{\alpha\eta}{2}$. Since $f_n \to f$ a.e. on finite $X_0$, $\mu(B_n^\alpha \cap X_0) < \frac{\eta}{2}$ eventually by Corollary 4.2.21. Thus $\mu(B_n^\alpha) < \eta$ eventually, so $f_n \to f$ in measure.
By Theorem 4.2.26 with $p = 1$, using uniform integrability and tightness, $f \in L^1(\mu)$ and $\|f_n - f\|_1 \to 0$.

(iii) In either case, $\left|\int_X f_n\,d\mu - \int_X f\,d\mu\right| = \left|\int_X (f_n - f)\,d\mu\right| \le \|f_n - f\|_1 \to 0$.

*Structural Commentary*: Tightness gives finite cores; UI upgrades in-measure convergence to $L^1$.
*Key Insight*: Uniform integrability prevents concentration; tightness prevents escape. □

**Corollary 4.4.9.** *Nonnegative Integral Vanishing Criterion*
*Let $f_n \in L^1(\mu)$ with $f_n \ge 0$ a.e. The following are equivalent.*

*(i)* $\int_X f_n\,d\mu \to 0$.
*(ii)* $\|f_n\|_1 \to 0$.

*(iii)* $f_n \to 0$ *in measure, and* $\{f_n : n \geq 1\}$ *is uniformly integrable and tight.*

*Proof.*

*(i)* $\Rightarrow$ *(ii)* Since $f_n \geq 0$, $\|f_n\|_1 = \int_X f_n \, d\mu \to 0$.

*(ii)* $\Rightarrow$ *(iii)* By Theorem 4.2.26 with $p = 1$ and limit 0, $\|f_n\|_1 \to 0$ implies $f_n \to 0$ in measure, $\forall \varepsilon > 0 \; \exists E_\varepsilon \in \mathcal{M}$ with $\mu(E_\varepsilon) < +\infty$ and $F \cap E_\varepsilon = \emptyset \Rightarrow \sup_n \int_F f_n \, d\mu < \varepsilon$, and $\forall \varepsilon > 0 \; \exists \delta > 0$ with $\mu(A) < \delta \Rightarrow \sup_n \int_A f_n \, d\mu < \varepsilon$; hence tightness and uniform integrability.

*(iii)* $\Rightarrow$ *(i)* Condition (iii) is hypotheses (ii)–(iv) of Theorem 4.2.26 with $p = 1$ and limit 0, so $\|f_n\|_1 \to 0$. Since $f_n \geq 0$, $\int_X f_n \, d\mu = \|f_n\|_1 \to 0$.

*Structural Commentary*: For nonnegative functions, integral convergence to zero is $L^1$ convergence to zero.
*Key Insight*: Vitali's conditions characterize when nonnegative mass vanishes. □

**Corollary 4.4.10.** *Domination Gives Tail and Small-Set Control*
*Let* $1 \leq p < \infty$, $g \in L^p(\mu)$, *and* $\forall n$, $|f_n| \leq g$ *a.e. Then* $\forall \varepsilon > 0$*:*

*(i)* $\exists E_\varepsilon \in \mathcal{M}$ *with* $\mu(E_\varepsilon) < +\infty$ *such that* $F \cap E_\varepsilon = \emptyset \Rightarrow \sup_n \int_F |f_n|^p \, d\mu < \varepsilon^p$.

*(ii)* $\exists \delta(\varepsilon) > 0$ *such that* $\mu(E) < \delta(\varepsilon) \Rightarrow \sup_n \int_E |f_n|^p \, d\mu < \varepsilon^p$.

*Proof.*

(i) Choose $E_\varepsilon$ with $\mu(E_\varepsilon) < +\infty$ and $F \cap E_\varepsilon = \emptyset \Rightarrow \int_F |g|^p \, d\mu < \varepsilon^p$ by Proposition 4.1.25. Since $|f_n|^p \leq |g|^p$ a.e., $\sup_n \int_F |f_n|^p \, d\mu < \varepsilon^p$.

(ii) Choose $\delta(\varepsilon) > 0$ with $\mu(E) < \delta(\varepsilon) \Rightarrow \int_E |g|^p \, d\mu < \varepsilon^p$ by Lemma 4.1.38. Since $|f_n|^p \leq |g|^p$ a.e., $\sup_n \int_E |f_n|^p \, d\mu < \varepsilon^p$.

*Structural Commentary*: One $L^p$ envelope supplies both tail tightness and small-set control.
*Key Insight*: Domination transfers both estimates from $g$ to every $f_n$. □

## 4.5 Approximation and Separability in $L^p$

Approximation results explain why $L^p$ spaces are large enough to contain rough measurable functions but still small enough to be controlled by countable data when $p < \infty$. The guiding pattern is to localize the $L^p$ mass, approximate measurable atoms by elementary sets, and then code the elementary sets with rational parameters.

**Definition 4.5.1.** *Dense Subsets and Separable Normed Spaces*
Let $A \subseteq B$ be subsets of a normed space $(V, \|\cdot\|)$.

(i) $A$ is *dense in* $B$ if $\forall b \in B \; \forall \varepsilon > 0 \; \exists a \in A$ with $\|a - b\| < \varepsilon$.

(ii) $V$ is *separable* if it has a countable dense subset.

Equivalently, $A$ is dense in $B$ iff every $b \in B$ is the norm-limit of a sequence from $A$.

**Example 4.5.2.** *Sequence-Space Separability*

(i) $\ell^\infty(\mathbf{F})$ is not separable. The set $\{0,1\}^{\mathbb{N}}$ is uncountable, and $s \neq t$ gives $d_\infty(s,t) = 1$; the balls $B_{1/3}(s)$ are pairwise disjoint, so no countable dense set can meet all of them.

(ii) For $1 \leq p < \infty$, $\ell^p(\mathbf{F})$ is separable. Let $D$ be the countable set of finite-support sequences with rational coordinates, using rational real and imaginary parts if $\mathbf{F} = \mathbb{C}$.
For $x \in \ell^p$ and $\varepsilon > 0$, choose $N$ with $\sum_{n>N} |x_n|^p < (\varepsilon/2)^p$, then choose $y \in D$ with support in $\{1, \ldots, N\}$ and $\sum_{n=1}^N |x_n - y_n|^p < (\varepsilon/2)^p$. Thus $\|x - y\|_p < \varepsilon$.

*Structural Commentary*: Separability is the existence of a countable set meeting every nonempty open ball.
*Key Insight*: Finite rational data are dense in $\ell^p$, while $\ell^\infty$ contains an uncountable 1-separated set.

**Theorem 4.5.3.** *Separability and $C_c$ Density in $L^p$*
*Let* $E \subseteq \mathbb{R}$ *be Lebesgue measurable and let* $1 \leq p < \infty$.
*If* $C_c(E)$ *denotes restrictions to* $E$ *of continuous functions on* $\mathbb{R}$ *with bounded support, then:*

*(i)* $L^p(E)$ *is separable;*

*(ii)* $C_c(E)$ *is dense in* $L^p(E)$.

*Proof.*

(i) Let $\mathbb{F}_0 := \mathbb{Q}$ if real, $\mathbb{F}_0 := \mathbb{Q} + i\mathbb{Q}$ if complex. Let $\mathcal{D} := \operatorname{span}_{\mathbb{F}_0}\{\mathbf{1}_{E \cap (r,s)} : r, s \in \mathbb{Q}, \; r < s\}$. Then $\mathcal{D}$ is countable. Extend $f \in L^p(E)$ by 0 to $\tilde{f} \in L^p(\mathbb{R})$.

By Proposition 4.1.25, it suffices to approximate $s = \sum_{\ell=1}^{M} a_\ell \mathbf{1}_{A_\ell}$ with $m(A_\ell) < +\infty$.

For each $\ell$, Corollary 6.2.4 and rational endpoint perturbation give a finite union $V_\ell$ of rational open intervals with $m(A_\ell \triangle V_\ell)$ small, hence $\|\mathbf{1}_{E\cap A_\ell} - \mathbf{1}_{E\cap V_\ell}\|_p \le m(A_\ell \triangle V_\ell)^{1/p}$. Approximate $a_\ell$ by $q_\ell \in \mathbb{F}_0$; then $\sum_{\ell=1}^{M} q_\ell \mathbf{1}_{E\cap V_\ell} \in \mathcal{D}$ approximates $f$ arbitrarily well. Hence $L^p(E)$ is separable.

(ii) For rational $r < s$ and $0 < \eta < \frac{s-r}{2}$, choose $\psi_\eta \in C_c(\mathbb{R})$ with $0 \le \psi_\eta \le 1$, $\psi_\eta = 1$ on $[r+\eta, s-\eta]$, and $\psi_\eta = 0$ on $\mathbb{R} \setminus (r,s)$. Then $\left\|\psi_\eta|_E - \mathbf{1}_{E\cap(r,s)}\right\|_p^p \le m((r, r+\eta) \cup (s-\eta, s)) \le 2\eta \to 0$.

Thus each generator of $\mathcal{D}$ lies in $\overline{C_c(E)}^{L^p(E)}$; by finite linearity and (i), $C_c(E)$ is dense in $L^p(E)$.

*Structural Commentary*: Finite-measure atoms reduce to rational intervals, then to continuous cutoffs.
*Key Insight*: Countable rational steps lie in $\overline{C_c(E)}^{L^p(E)}$. □

**Remark 4.5.4.** *$L^1$ Does Not Exhaust the Dual of $L^\infty$*
Let $I = [a,b]$ with $a < b$, and let $\iota : L^1(I) \to (L^\infty(I))^*$ be $\iota(g)(f) := \int_I fg\,dm$. The map $\iota$ from Proposition 4.1.36 is not onto. If $\iota$ were onto, then $(L^\infty(I))^*$ would be separable by Theorem 4.5.3.
Let $(\Phi_n)$ be dense in its unit sphere and choose $u_n \in L^\infty(I)$ with $\|u_n\|_\infty = 1$ and $|\Phi_n(u_n)| > \frac{1}{2}$.
If $Y := \overline{\operatorname{span}}\{u_n : n \ge 1\} \ne L^\infty(I)$, Hahn–Banach gives $\Phi \in (L^\infty(I))^*$ with $\|\Phi\| = 1$ and $\Phi|_Y = 0$.
Choosing $n$ with $\|\Phi - \Phi_n\| < \frac{1}{4}$ gives $|\Phi_n(u_n)| < \frac{1}{4}$, a contradiction. Hence $L^\infty(I)$ would be separable.
But $v_t := \mathbf{1}_{[a,t]}$ satisfies $\|v_t - v_s\|_\infty = 1$ for $s < t$, an uncountable separated family, so $L^\infty(I)$ is not separable.

**Example 4.5.5.** *$L^\infty$ on an Interval Need Not Be Separable*
Let $I = [a,b]$ with $a < b$. For $t \in I$, set $u_t := \mathbf{1}_{[a,t]}$. If $s < t$, then $\|u_t - u_s\|_\infty = 1$, since $u_t - u_s = \mathbf{1}_{(s,t]}$ a.e.
Thus $\{B(u_t, \frac{1}{3}) : t \in I\}$ is pairwise disjoint. If countable $D \subseteq L^\infty(I)$ were dense, choose $d_t \in D \cap B(u_t, \frac{1}{3})$ for each $t \in I$; disjointness makes $t \mapsto d_t$ injective, impossible. Hence $L^\infty(I)$ is not separable.
*Key Insight*: Uncountably many distance-1 indicators preclude countable density.




Figure 220: For $1 \le p < \infty$, rational interval steps and $C_c(E)$ are dense in $L^p(E)$; $L^\infty([a,b])$ is nonseparable.

## 4.6 Weak Convergence in $L^p$

Weak convergence in $L^p$ keeps only the scalar information detected by conjugate functions. For $1 < p < +\infty$, reflexivity turns boundedness into weak subsequential compactness; at $p = 1$, bounded mass can still concentrate and destroy weak compactness unless uniform integrability is imposed.

**Theorem 4.6.1.** *$L^p$ Weak Convergence by Dual Tests*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, $1 < p < +\infty$, $q = \frac{p}{p-1}$, $(f_n) \subseteq L^p(\mu)$, and $f \in L^p(\mu)$, all real-valued. Then $f_n \rightharpoonup f$ in $L^p(\mu)$ iff $\int_X f_n g\,d\mu \to \int_X fg\,d\mu$ for all $g \in L^q(\mu)$.*

*Proof.*

(⇒) For $g \in L^q(\mu)$, $\Lambda_g(h) := \int_X hg\,d\mu$ is in $(L^p(\mu))^*$ by Holder.
Hence $f_n \rightharpoonup f$ implies $\int_X f_n g\,d\mu = \Lambda_g(f_n) \to \Lambda_g(f) = \int_X fg\,d\mu$.

(⇐) Let $\Lambda \in (L^p(\mu))^*$. By Theorem 4.1.33, choose $g \in L^q(\mu)$ with $\Lambda(h) = \int_X hg\,d\mu$ $\forall h \in L^p(\mu)$.
The assumed integral test yields $\Lambda(f_n) \to \Lambda(f)$, hence $f_n \rightharpoonup f$.

*Structural Commentary*: Dual tests are exactly $L^q$ pairings.
*Key Insight*: Weak convergence in $L^p$ is conjugate-function testing. □

**Proposition 4.6.2.** *Weak Limits and Norm Lower Semicontinuity*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, $1 < p < +\infty$, $(f_n) \subseteq L^p(\mu)$, and $f_n \rightharpoonup f$ in real $L^p(\mu)$. Then:*

*(i)* $\sup_n \|f_n\|_p < +\infty$;

*(ii) if $f_n \rightharpoonup h$ in $L^p(\mu)$, then $h = f$ in $L^p(\mu)$;*

*(iii)* $\|f\|_p \le \liminf_{n\to\infty} \|f_n\|_p$.

*Proof.*

(i) By Proposition 6.1.4 applied to $u_n = f_n$ in $L^p(\mu)$, $f_n \rightharpoonup f$ implies $\sup_n \|f_n\|_p < +\infty$.

(ii) Let $q = \frac{p}{p-1}$, $f_n \rightharpoonup h$. For $g \in L^q(\mu)$, by Theorem 4.6.1, $\int_X (f-h)g\,d\mu = \lim_n \int_X f_n g\,d\mu - \lim_n \int_X f_n g\,d\mu = 0$. By Proposition 4.1.29, $\|f-h\|_p = 0$, so $h = f$ in $L^p(\mu)$.

(iii) By Proposition 6.1.4, $u \mapsto \|u\|_p$ is weakly lower semicontinuous on $L^p(\mu)$; with $u_n = f_n$ and $u = f$, $\|f\|_p \le \liminf_{n\to\infty} \|f_n\|_p$.

*Structural Commentary*: General weak facts handle bounds; $L^q$ tests force uniqueness.
*Key Insight*: Weak limits are norm-controlled and conjugate-tested. □

**Theorem 4.6.3.** *Reflexivity of $L^p$, $1 < p < +\infty$*
*Let $(X, \mathcal{M}, \mu)$ be $\sigma$-finite, $1 < p < +\infty$, $q = \frac{p}{p-1}$, and $Jf(\Phi) := \Phi(f)$.*
*For $r \in \{p, q\}$ with conjugate $r'$, set $T_r u \in (L^{r'}(\mu))^*$ by $(T_r u)(v) := \int_X uv\,d\mu$. Then:*

*(i) $T_p$ and $T_q$ are isometric isomorphisms;*
*(ii) $J : L^p(\mu) \to (L^p(\mu))^{**}$ is onto; hence real $L^p(\mu)$ is reflexive.*

*Proof.*

(i) By Theorem 4.1.33, $T_p : L^p(\mu) \to (L^q(\mu))^*$ and $T_q : L^q(\mu) \to (L^p(\mu))^*$ are isometric isomorphisms.

(ii) Let $S \in (L^p(\mu))^{**}$. Since $S \circ T_q \in (L^q(\mu))^*$, by (i), $f \in L^p(\mu)$, $S(T_q g) = (T_p f)(g) = \int_X fg\,d\mu$ $\forall g \in L^q(\mu)$. For $\Phi \in (L^p(\mu))^*$, by (i) $\Phi = T_q g$; $S(\Phi) = S(T_q g) = \int_X fg\,d\mu = \Phi(f) = Jf(\Phi)$. Hence $S = Jf$, $J$ is onto.

*Structural Commentary*: Represent both dual layers by conjugate pairings.
*Key Insight*: Every bidual functional is evaluation at an $L^p$ function. □

**Definition 4.6.4.** *Weak Sequential Compactness*
Let $\mathcal{X}$ be a normed space and $K \subseteq \mathcal{X}$.
Then $K$ is *weakly sequentially compact* if every $(x_n) \subseteq K$ admits $n_k \uparrow \infty$ and $x \in K$ with $x_{n_k} \rightharpoonup x$ in $\mathcal{X}$.

**Theorem 4.6.5.** *Riesz Weak Compactness Theorem*
*Let $(X, \mathcal{M}, \mu)$ be $\sigma$-finite, $1 < p < +\infty$, $q = \frac{p}{p-1}$, and $(f_n) \subseteq L^p(\mu)$ be real-valued with $M := \sup_n \|f_n\|_p < +\infty$.*
*Then there exist $n_k \uparrow \infty$ and $f \in L^p(\mu)$ such that:*

*(i) $f_{n_k} \rightharpoonup f$ in $L^p(\mu)$;*
*(ii) $\int_X f_{n_k} g\,d\mu \to \int_X fg\,d\mu$ for all $g \in L^q(\mu)$.*

*Proof.* Set $B_M := \{h \in L^p(\mu) : \|h\|_p \le M\}$. By Theorem 4.6.3, $L^p(\mu)$ is reflexive; hence $B_M$ is weakly compact. By Eberlein–Smulian, $B_M$ is weakly sequentially compact.

(i) Since $(f_n) \subseteq B_M$ and $B_M$ is weakly sequentially compact, $\exists n_k \uparrow \infty$ and $f \in B_M$ with $f_{n_k} \rightharpoonup f$ in $L^p(\mu)$.

(ii) From (i) and Theorem 4.6.1, $\forall g \in L^q(\mu)$, $\int_X f_{n_k} g\,d\mu \to \int_X fg\,d\mu$.

*Structural Commentary*: Reflexive bounded balls are weakly sequentially compact.
*Key Insight*: Bounded $L^p$ sequences have conjugate-test convergent subsequences. □

**Corollary 4.6.6.** *The $L^p$ Unit Ball Is Weakly Sequentially Compact*
*Let $(X, \mathcal{M}, \mu)$ be $\sigma$-finite, $1 < p < +\infty$, and $B_p := \{f \in L^p(\mu) : \|f\|_p \le 1\}$.*
*Then $B_p$ is weakly sequentially compact in $L^p(\mu)$.*

*Proof.* Let $(f_n) \subseteq B_p$. By Theorem 4.6.5, $\exists n_k \uparrow \infty$ and $f \in L^p(\mu)$ with $f_{n_k} \rightharpoonup f$.
By Proposition 4.6.2, $\|f\|_p \le \liminf_{k\to\infty} \|f_{n_k}\|_p \le 1$, so $f \in B_p$.
*Structural Commentary*: Weak limits stay in $B_p$ by lower semicontinuity.
*Key Insight*: Closed norm balls are weakly sequentially closed. □

**Theorem 4.6.7.** *Radon–Riesz Principle*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, $1 < p < +\infty$, $(f_n) \subseteq L^p(\mu)$, and $f \in L^p(\mu)$, all real-valued.*
*Then $f_n \rightharpoonup f$ and $\|f_n\|_p \to \|f\|_p$ iff $\|f_n - f\|_p \to 0$.*

*Proof.*

($\Rightarrow$) If $\|f\|_p = 0$, then $\|f_n - f\|_p \le \|f_n\|_p + \|f\|_p \to 0$. Let $a := \|f\|_p > 0$.
Otherwise, after subsequences, $\|f_n - f\|_p \ge \varepsilon > 0$ and, with $a_n := \|f_n\|_p$, $u_n := \frac{f_n}{a_n}$, $u := \frac{f}{a}$, one has $u_n \rightharpoonup u$, $\|u_n\|_p = \|u\|_p = 1$, and $\|u_n - u\|_p \ge \eta > 0$. By uniform convexity, $\left\|\frac{u_n+u}{2}\right\|_p \le 1 - \delta$.
Since $\frac{u_n+u}{2} \rightharpoonup u$, weak lower semicontinuity implies $1 = \|u\|_p \le \liminf_n \left\|\frac{u_n+u}{2}\right\|_p \le 1 - \delta$, contradiction.

($\Leftarrow$) Assume $\|f_n - f\|_p \to 0$. Then $\left|\|f_n\|_p - \|f\|_p\right| \le \|f_n - f\|_p \to 0$.
For $\Lambda \in (L^p(\mu))^*$, $|\Lambda(f_n) - \Lambda(f)| \le \|\Lambda\| \, \|f_n - f\|_p \to 0$, hence $f_n \rightharpoonup f$.

*Structural Commentary*: Uniform convexity turns weak lower semicontinuity into strong convergence.
*Key Insight*: Weak direction plus norm size fixes the vector. □

**Remark 4.6.8.** *$L^2$ Proof of Radon–Riesz*
For $p = 2$, $f_n \rightharpoonup f$ in $L^2(\mu)$, $\|f_n\|_2 \to \|f\|_2$. Then $\int_X f_n f \, d\mu \to \|f\|_2^2$,
$\|f_n - f\|_2^2 = \|f_n\|_2^2 - 2\int_X f_n f \, d\mu + \|f\|_2^2 \to 0$. The converse follows from norm convergence in $L^2(\mu)$.
*Key Insight*: Weak convergence fixes the cross term.

**Corollary 4.6.9.** *Strong Subsequence Criterion*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, $1 < p < +\infty$, $(f_n) \subseteq L^p(\mu)$, and $f_n \rightharpoonup f$ in real $L^p(\mu)$. Then $\exists n_k \uparrow \infty$ with $\|f_{n_k} - f\|_p \to 0$ iff $\|f\|_p = \liminf_{n\to\infty} \|f_n\|_p$.*

*Proof.* Set $L := \liminf_{n\to\infty} \|f_n\|_p$.

($\Rightarrow$) If $\|f_{n_k} - f\|_p \to 0$, then $\|f_{n_k}\|_p \to \|f\|_p$.
Since $L \le \lim_k \|f_{n_k}\|_p = \|f\|_p$ and, by Proposition 4.6.2, $\|f\|_p \le L$, one has $L = \|f\|_p$.

($\Leftarrow$) Assume $L = \|f\|_p$. Choose $n_k \uparrow \infty$ with $\|f_{n_k}\|_p \to L$.
Since $f_{n_k} \rightharpoonup f$ and $\|f_{n_k}\|_p \to \|f\|_p$, Theorem 4.6.7 implies $\|f_{n_k} - f\|_p \to 0$.

*Structural Commentary*: Match the liminf subsequence to Radon–Riesz.
*Key Insight*: Strong subsequences occur exactly when no norm is lost. □

**Theorem 4.6.10.** *Dunford–Pettis Criterion in $L^1$*
*Let $(X, \mathcal{M}, \mu)$ be a finite measure space and let $(f_n)$ be bounded in $L^1(\mu)$. The following are equivalent.*

*(i) $(f_n)$ is uniformly integrable.*
*(ii) $\forall (f_{n_k}) \subseteq (f_n)$ $\exists (f_{n_{k_j}}) \subseteq (f_{n_k})$ converging weakly in $L^1(\mu)$.*

*Proof.*

($\Rightarrow$) Let $(u_j)$ be a subsequence of $(f_n)$, set $C := \sup_j \|u_j\|_1$, and put $T_m(t) := (-m) \vee (t \wedge m)$.
By Proposition 4.4.7, $\sup_j \|u_j - T_m(u_j)\|_1 \to 0$. For fixed $m$ and sign $\pm$, $\left\|T_m(u_j^{\pm})\right\|_2^2 \le m \left\|u_j^{\pm}\right\|_1 \le mC$.
By bounded Hilbert weak subsequence extraction and diagonalization over $(m, \pm)$, pass to a subsequence, still $(u_j)$, such that $\int_E T_m(u_j^{\pm}) \, d\mu$ converges for every $E \in \mathcal{M}$ and $m \in \mathbb{N}$. Set $\nu_j^{\pm}(E) := \int_E u_j^{\pm} \, d\mu$.
For $r, s, m$ and $E \in \mathcal{M}$, $|\nu_r^{\pm}(E) - \nu_s^{\pm}(E)| \le \left|\int_E (T_m(u_r^{\pm}) - T_m(u_s^{\pm})) \, d\mu\right| + 2\sup_j \left\|u_j^{\pm} - T_m(u_j^{\pm})\right\|_1$.
Hence $(\nu_j^{\pm}(E))_j$ is Cauchy. Put $\nu^{\pm}(E) := \lim_j \nu_j^{\pm}(E)$.
By Theorem 3.6.11, $\nu^{\pm}$ are finite measures and $\nu^{\pm} \ll \mu$. Write $h^{\pm} = \frac{d\nu^{\pm}}{d\mu}$ and $h := h^+ - h^- \in L^1(\mu)$.
Then $\int_X \varphi u_j \, d\mu \to \int_X \varphi h \, d\mu$ for simple $\varphi$; by Lemma 1.1.14 and $\sup_j \|u_j\|_1 + \|h\|_1 < +\infty$, the convergence holds for every $\varphi \in L^\infty(\mu)$. Thus $u_j \rightharpoonup h$ in $L^1(\mu)$.

($\Leftarrow$) Suppose $(f_n)$ is not uniformly integrable. Since finite subfamilies are uniformly integrable by Proposition 4.4.5(ii), choose a subsequence $(u_j)$ and sets $A_j \in \mathcal{M}$ with $\mu(A_j) \to 0$ and $\int_{A_j} |u_j| \, d\mu \ge \varepsilon_0 > 0$.
By (ii), some $(u_{j_\ell})$ converges weakly in $L^1(\mu)$ to $u$. Set $\eta_\ell(E) := \int_E u_{j_\ell} \, d\mu$ and $\eta(E) := \int_E u \, d\mu$.
Then $\eta_\ell(E) \to \eta(E)$ for every $E \in \mathcal{M}$, and $\sup_\ell |\eta_\ell|(X) = \sup_\ell \|u_{j_\ell}\|_1 < +\infty$ by Theorem 3.1.9.
The signed form of Theorem 3.6.11 implies $\eta_\ell \ll_{\mathrm{u}} \mu$ in variation:
$\forall \varepsilon > 0$ $\exists \delta > 0$ such that $\mu(A) < \delta \Rightarrow \sup_\ell |\eta_\ell|(A) < \varepsilon$. Since $|\eta_\ell|(A) = \int_A |u_{j_\ell}| \, d\mu$, the subsequence $(u_{j_\ell})$ is uniformly integrable, contradicting $\int_{A_{j_\ell}} |u_{j_\ell}| \, d\mu \ge \varepsilon_0$ and $\mu(A_{j_\ell}) \to 0$.

*Structural Commentary*: Truncate, pass to setwise limits, apply VHS.
*Key Insight*: UI is weak sequential compactness in finite $L^1$. □

**Corollary 4.6.11.** *Dominated $L^1$ Families Have Weak Subsequences*
*Let $(X, \mathcal{M}, \mu)$ be finite. If $g \in L^1(\mu)$, $\forall n \, |f_n| \le g$ a.e., then $(f_n)$ has subsequence that converges weakly in $L^1(\mu)$.*

*Proof.* Since $|f_n| \le g$ a.e., $\|f_n\|_1 \le \|g\|_1$ $\forall n$, $(f_n)$ is bounded in $L^1(\mu)$. By Proposition 4.4.5(i), $\{f_n : n \ge 1\}$ is uniformly integrable. Theorem 4.6.10, $(i) \Rightarrow (ii)$, yields a weakly convergent subsequence.
*Structural Commentary*: Domination verifies Dunford–Pettis through a common integrable envelope.
*Key Insight*: One $L^1$ envelope prevents concentration throughout the sequence. □

**Corollary 4.6.12.** *Bounded $L^p$ Families Are Weakly Compact in $L^1$*
*Let $(X, \mathcal{M}, \mu)$ be finite, let $1 < p < +\infty$, and let $(f_n)$ be bounded in $L^p(\mu)$. Then $(f_n)$ has a subsequence that converges weakly in $L^1(\mu)$.*

*Proof.* Let $q$ be conjugate to $p$ and $M := \sup_n \|f_n\|_p < +\infty$. By Hölder, $\|f_n\|_1 \le M\mu(X)^{\frac{1}{q}}$ $\forall n$, so $(f_n)$ is bounded in $L^1(\mu)$. By Corollary 4.4.6, $\{f_n : n \ge 1\}$ is uniformly integrable.
Hence Theorem 4.6.10, $(i) \Rightarrow (ii)$, yields a weakly convergent subsequence in $L^1(\mu)$.
*Structural Commentary*: Hölder gives $L^1$ boundedness, uniform integrability before Dunford–Pettis is applied.
*Key Insight*: On finite measure spaces, an $L^p$ bound with $p > 1$ rules out $L^1$ concentration. □

**Example 4.6.13.** *A Bounded $L^1$ Sequence with No Weakly Convergent Subsequence*
In $L^1[0,1]$, $I_n = [0, \frac{1}{n}]$ $f_n = n\mathbf{1}_{I_n}$. Then $\|f_n\|_1 = 1$ $\forall n$, and no subsequence of $(f_n)$ converges weakly in $L^1[0,1]$.

*Proof.* Assume $f_{n_k} \rightharpoonup f$ in $L^1[0,1]$. For $0 < c < d \le 1$, $\mathbf{1}_{[c,d]} \in L^\infty[0,1]$ and $[c,d] \cap I_{n_k} = \varnothing$ for $k$ large.
Hence $\int_c^d f\,dm = \lim_{k\to\infty} \int_c^d f_{n_k}\,dm = 0$. Thus $f = 0$ a.e. on $(0,1]$.
Testing by $\mathbf{1}_{[0,1]}$ yields $0 = \int_0^1 f\,dm = \lim_{k\to\infty} \int_0^1 f_{n_k}\,dm = 1$, contradiction.
*Key Insight*: Unit mass concentrates on shrinking supports. □




Figure 221: $f_n = n\mathbf{1}_{[0,\frac{1}{n}]}$: unit $L^1$ mass on shrinking support.

**Remark 4.6.14.** *Endpoint Failure and Uniform Integrability*
For $f_n = n\mathbf{1}_{[0,\frac{1}{n}]} \in L^1[0,1]$, $\|f_n\|_1 = 1$ but $\int_{\{f_n > R\}} f_n\,dm = 1$ whenever $n > R$.
Hence $(f_n)$ is not uniformly integrable; the $p = 1$ failure is concentration, not lack of boundedness.

**Theorem 4.6.15.** *Banach–Saks Theorem*
*Let $(X, \mathcal{M}, \mu)$ be a measure space, $1 < p < +\infty$, $(f_n) \subseteq L^p(\mu)$, and $f_n \rightharpoonup f$ in real $L^p(\mu)$.*
*Then $\exists n_k \uparrow \infty$ such that, with $A_k := \frac{1}{k}\sum_{j=1}^k f_{n_j}$, $\|A_k - f\|_p \to 0$.*

*Proof.* Set $h_n := f_n - f$. Then $h_n \rightharpoonup 0$ and $\sup_n \|h_n\|_p < +\infty$ by Proposition 4.6.2.
Since real $L^p(\mu)$ is uniformly convex, by the selection lemma there is $n_k \uparrow \infty$ with $\left\|\frac{1}{k}\sum_{j=1}^k h_{n_j}\right\|_p \to 0$.
Since $\frac{1}{k}\sum_{j=1}^k h_{n_j} = A_k - f$, $\|A_k - f\|_p \to 0$.
*Structural Commentary*: Reduce to weakly null; select Cesaro means by uniform convexity.
*Key Insight*: Convex averaging turns weak nullity into norm decay. □

**Remark 4.6.16.** *The $L^2$ Selection Estimate*
For $p = 2$, replace $f_n$ by $f_n - f$ and assume $f_n \rightharpoonup 0$ in $L^2(\mu)$ with $\|f_n\|_2^2 \le M$. Set $n_1 = 1$ and $S_k := \sum_{j=1}^k f_{n_j}$.

(i) Given $S_k$, choose $n_{k+1} > n_k$ so that $\left|\int_X S_k f_{n_{k+1}}\,d\mu\right| \le 1$.
(ii) Then $\|S_{k+1}\|_2^2 = \|S_k\|_2^2 + 2\int_X S_k f_{n_{k+1}}\,d\mu + \|f_{n_{k+1}}\|_2^2 \le \|S_k\|_2^2 + 2 + M$.
(iii) Hence $\|S_k/k\|_2^2 \le \frac{2+M}{k}$ by induction.

Thus $S_k/k \to 0$ in $L^2(\mu)$.
*Key Insight*: Weak convergence makes new terms nearly orthogonal to old sums.

**Definition 4.6.17.** *Closed Convex Sets and Convex Functionals*
Let $C$ be a subset of a normed space $\mathcal{X}$.

(i) $C$ is *convex* if $\lambda f + (1-\lambda)g \in C$ whenever $f, g \in C$ and $0 \le \lambda \le 1$.
(ii) $C$ is *closed* if every strongly convergent sequence in $C$ has its limit in $C$.
(iii) A real-valued functional $T : C \to \mathbb{R}$ is *continuous* if $f_n \to f$ strongly in $C$ implies $T(f_n) \to T(f)$.
(iv) If $C$ is convex, then $T$ is *convex* if $T(\lambda f + (1-\lambda)g) \le \lambda T(f) + (1-\lambda)T(g)$ $\forall f, g \in C$ and $0 \le \lambda \le 1$.

**Example 4.6.18.** *Basic Closed Convex Sets in $L^p$*
Let $E \subseteq \mathbb{R}$ be measurable and $1 \le p < +\infty$.

(i) If $g \in L^p(E)$ and $g \ge 0$ a.e., then $C_g := \{f \in L^p(E) : |f| \le g \text{ a.e.}\}$ is closed and convex.

(ii) $B := \{f \in L^p(E) : \|f\|_p \le 1\}$ is closed and convex.

*Proof.*

(i) If $f, h \in C_g$ and $0 \le \lambda \le 1$, then $|\lambda f + (1-\lambda)h| \le \lambda |f| + (1-\lambda)|h| \le g$ a.e., so $C_g$ is convex.
If $f_n \in C_g$ and $f_n \to f$ in $L^p(E)$, choose $f_{n_k} \to f$ a.e. by Theorem 4.1.22; then $|f| \le g$ a.e., so $f \in C_g$.

(ii) If $f, h \in B$ and $0 \le \lambda \le 1$, then by Minkowski, $\|\lambda f + (1-\lambda)h\|_p \le \lambda \|f\|_p + (1-\lambda)\|h\|_p \le 1$.
If $f_n \in B$ and $f_n \to f$ in $L^p(E)$, then $\|f\|_p \le \|f - f_n\|_p + \|f_n\|_p \le \|f - f_n\|_p + 1$, hence $\|f\|_p \le 1$.

*Key Insight*: Norm limits preserve a.e. bounds and norm bounds. □

**Lemma 4.6.19.** *Weak Limits and Convex Lower Semicontinuity*
*Let $E \subseteq \mathbb{R}$ be measurable, $1 < p < +\infty$, $C \subseteq L^p(E)$ closed, bounded, and convex.*
*Let $T : C \to \mathbb{R}$ be continuous and convex. If $(f_n) \subseteq C$ and $f_n \rightharpoonup f$ in $L^p(E)$, then, with $\alpha := \liminf_{n\to\infty} T(f_n)$:*

*(i) $f \in C$;*
*(ii) $T(f) \le \alpha$.*

*Proof.*

(i) By Theorem 4.6.15, choose $n_k \uparrow \infty$ with $a_k := \frac{1}{k}\sum_{j=1}^{k} f_{n_j} \to f$ in $L^p(E)$.
Since $a_k \in \operatorname{co}\{f_{n_1}, \dots, f_{n_k}\} \subseteq C$ and $C$ is closed, $f \in C$.

(ii) Choose $m_k \uparrow \infty$ with $T(f_{m_k}) \to \alpha$.

By Theorem 4.6.15, choose $g_j := f_{m_{k_j}}$ such that $b_k := \frac{1}{k}\sum_{j=1}^{k} g_j \to f$ in $L^p(E)$.

Then $b_k \in C$ and $T(g_j) \to \alpha$; by continuity and convexity, $T(f) = \lim_k T(b_k) \le \lim_k \frac{1}{k}\sum_{j=1}^{k} T(g_j) = \alpha$.

*Structural Commentary*: Banach–Saks supplies strong convex averages; convexity transfers the liminf.
*Key Insight*: Weak limits inherit closed convex constraints and convex lsc. □

**Theorem 4.6.20.** *Direct Method for Convex Functionals on $L^p$*
*Let $E \subseteq \mathbb{R}$ be measurable, $1 < p < +\infty$, $C \subseteq L^p(E)$ nonempty, closed, bounded, and convex.*
*Let $T : C \to \mathbb{R}$ be continuous and convex. Then $\exists f_0 \in C$ such that $T(f_0) = \inf_{h\in C} T(h)$.*

*Proof.*

(i) If $\inf_C T = -\infty$, choose $f_n \in C$ with $T(f_n) \to -\infty$.
Since $C$ is bounded, by Theorem 4.6.5, $\exists n_k \uparrow \infty$ and $f \in L^p(E)$ with $f_{n_k} \rightharpoonup f$.
By Lemma 4.6.19, $f \in C$ and $T(f) \le \liminf_k T(f_{n_k}) = -\infty$, contradicting $T : C \to \mathbb{R}$. Thus $\inf_C T > -\infty$.

(ii) Let $c := \inf_C T$ and choose $f_n \in C$ with $T(f_n) \to c$.
Since $C$ is bounded, by Theorem 4.6.5, $\exists n_k \uparrow \infty$ and $f_0 \in L^p(E)$ with $f_{n_k} \rightharpoonup f_0$.
By Lemma 4.6.19, $f_0 \in C$ and $T(f_0) \le \liminf_k T(f_{n_k}) = c$. Since $c \le T(f_0)$, $T(f_0) = c$.

*Structural Commentary*: Weak compactness supplies a limit; convex lsc keeps the infimum.
*Key Insight*: Direct method equals compactness plus lower semicontinuity. □

**Corollary 4.6.21.** *Integral Convex Functionals Have Unit-Ball Minimizers*
*Let $\mu(E) < +\infty$, $1 < p < +\infty$, and $\varphi : \mathbb{R} \to \mathbb{R}$ be continuous convex with $|\varphi(s)| \le c_1 + c_2|s|^p$ for some $c_1, c_2 \ge 0$. Set $B := \{f \in L^p(E) : \|f\|_p \le 1\}$ and $T(f) := \int_E \varphi(f)\,dm$. Then:*

*(i) $T : B \to \mathbb{R}$ is continuous and convex;*
*(ii) $\exists f_0 \in B$ such that $T(f_0) = \min_{f\in B} T(f)$.*

*Proof.*

(i) For $f \in B$, $|\varphi(f)| \le c_1 + c_2|f|^p \in L^1(E)$, so $T(f) \in \mathbb{R}$.
Convexity of $\varphi$ implies $T(\lambda f + (1-\lambda)h) \le \lambda T(f) + (1-\lambda)T(h)$ for $f, h \in B$, $0 \le \lambda \le 1$.
If $f_n \to f$ in $L^p(E)$, every subsequence admits $f_{n_k} \to f$ a.e. with $|f_{n_k}| \le g$ for some $g \in L^p(E)$.
Then $|\varphi(f_{n_k})| \le c_1 + c_2 g^p \in L^1(E)$, so by dominated convergence $T(f_{n_k}) \to T(f)$.
By the subsequence criterion, $T(f_n) \to T(f)$.

(ii) The set $B$ is nonempty, bounded, convex by Minkowski, and closed since $f_n \in B$, $f_n \to f$ in $L^p(E)$ imply $\|f\|_p \le \liminf_n \|f_n\|_p \le 1$. By Theorem 4.6.20, $\exists f_0 \in B$ with $T(f_0) = \min_{f\in B} T(f)$.

*Structural Commentary*: $p$-growth ⇒ continuity; direct method ⇒ attainment.
*Key Insight*: Growth converts strong $L^p$ convergence to dominated convergence. □

# 5 Banach and Hilbert Spaces

## 5.1 Normed and Banach Spaces

Normed spaces make vector spaces metric; Banach spaces add completeness. This is the setting for infinite series, fixed points, bounded linear maps, and operator norms.

**Definition 5.1.1.** *Normed and Banach Spaces*
Let $\mathcal{X}$ be a vector space over $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$.

(i) *Norm*: $\|\cdot\| : \mathcal{X} \to [0,\infty)$; $\|x\| = 0 \iff x = 0$, $\|\lambda x\| = |\lambda| \, \|x\|$; $\|x+y\| \le \|x\| + \|y\|$ for $x, y \in \mathcal{X}$, $\lambda \in \mathbb{F}$.
(ii) *Normed space*: $(\mathcal{X}, \|\cdot\|)$ carries the metric $d(x,y) := \|x - y\|$.
(iii) *Banach space*: $(\mathcal{X}, \|\cdot\|)$ is complete in the norm metric.

**Definition 5.1.2.** *Basic Normed-Space Geometry*
Let $\mathcal{X}$ be a normed space.

(i) *Open unit ball*: $B_{\mathcal{X}} := \{x \in \mathcal{X} : \|x\| < 1\}$.
(ii) *Closed unit ball*: $\overline{B}_{\mathcal{X}} := \{x \in \mathcal{X} : \|x\| \le 1\}$.
(iii) *Unit vector*: $x \in \mathcal{X}$ satisfies $\|x\| = 1$.
(iv) *Minkowski sum*: $A + B := \{a + b : a \in A,\ b \in B\}$ for nonempty $A, B \subseteq \mathcal{X}$.
(v) *Scalar multiple*: $\lambda A := \{\lambda a : a \in A\}$ for nonempty $A \subseteq \mathcal{X}$ and $\lambda \in \mathbb{F}$.
(vi) *Closed linear span*: $\operatorname{cspan}(S) := \overline{\operatorname{span}(S)}$ for $S \subseteq \mathcal{X}$.
(vii) *Equivalent norms*: $\|\cdot\|_1 \sim \|\cdot\|_2$ iff $\exists c_1, c_2 > 0$ such that $\forall x \in \mathcal{X}$, $c_1 \|x\|_1 \le \|x\|_2 \le c_2 \|x\|_1$.

**Proposition 5.1.3.** *Linear Operations, Closed Spans, and Equivalent Norms*
*Let $\mathcal{X}$ be a normed space.*

*(i) The norm map $x \mapsto \|x\|$, addition $\mathcal{X} \times \mathcal{X} \to \mathcal{X}$, and scalar multiplication $\mathbb{F} \times \mathcal{X} \to \mathcal{X}$ are continuous.*
*(ii) If $M \le \mathcal{X}$, then $\overline{M} \le \mathcal{X}$; for $S \subseteq \mathcal{X}$, $\operatorname{cspan}(S)$ is the least closed subspace containing $S$.*
*(iii) Equivalent norms have the same convergent and Cauchy sequences, topology, and compact sets.*

*Proof.*

(i) $|\|x\| - \|y\|| \le \|x - y\|$, so $x \mapsto \|x\|$ is 1-Lipschitz.
Also $\|(x+h) + (y+k) - (x+y)\| \le \|h\| + \|k\|$, so addition is continuous.
If $\lambda_n \to \lambda$ and $x_n \to x$, then $\|\lambda_n x_n - \lambda x\| \le |\lambda_n| \, \|x_n - x\| + |\lambda_n - \lambda| \, \|x\| \to 0$, since $(\lambda_n)$ is bounded.
(ii) Let $x, y \in \overline{M}$ and choose $x_n, y_n \in M$ with $x_n \to x$, $y_n \to y$.
For $\alpha, \beta \in \mathbb{F}$, $\alpha x_n + \beta y_n \in M$ and $\alpha x_n + \beta y_n \to \alpha x + \beta y$ by (i); hence $\overline{M} \le \mathcal{X}$.
If $C$ is closed, $C \le \mathcal{X}$, and $S \subseteq C$, then $\operatorname{span}(S) \subseteq C$, so $\operatorname{cspan}(S) = \overline{\operatorname{span}(S)} \subseteq C$.
(iii) If $c \|z\|_1 \le \|z\|_2 \le C \|z\|_1$ for all $z \in \mathcal{X}$, then $B_1(x, \frac{\varepsilon}{C}) \subseteq B_2(x, \varepsilon) \subseteq B_1(x, \frac{\varepsilon}{c})$. Thus the topologies agree, the identity maps are bi-Lipschitz, convergent and Cauchy sequences agree, and compact sets agree.

*Structural Commentary*: Norms turn the algebraic operations into continuous operations.
*Key Insight*: Closing a subspace preserves linearity because linear combinations commute with limits. □

**Definition 5.1.4.** *Bounded Linear Maps and Operator Norm*
Let $\mathcal{X}, \mathcal{Y}$ be normed spaces and let $T : \mathcal{X} \to \mathcal{Y}$ be linear.

(i) *Bounded*: $\exists C \ge 0$ such that $\forall x \in \mathcal{X}$, $\|Tx\|_{\mathcal{Y}} \le C \|x\|_{\mathcal{X}}$.
(ii) *Operator norm*: $\|T\|_{\mathrm{op}} := \sup_{\|x\|_{\mathcal{X}} \le 1} \|Tx\|_{\mathcal{Y}} = \sup_{x \ne 0} \frac{\|Tx\|_{\mathcal{Y}}}{\|x\|_{\mathcal{X}}}$.
(iii) *Operator space*: $\mathcal{L}(\mathcal{X}, \mathcal{Y}) := \{T : \mathcal{X} \to \mathcal{Y} : T \text{ is bounded and linear}\}$ and $\mathcal{L}(\mathcal{X}) := \mathcal{L}(\mathcal{X}, \mathcal{X})$.

**Theorem 5.1.5.** *Bounded Equals Continuous for Linear Maps*
*For a linear map $T : \mathcal{X} \to \mathcal{Y}$ between normed spaces, the following are equivalent:*

*(i) $T$ is bounded.*
*(ii) $T$ is continuous at 0.*
*(iii) $T$ is continuous everywhere.*

*Proof.*

*(i)* $\Rightarrow$ *(iii)* If $T$ is bounded, then $\|Tx - Ty\| \le \|T\|_{\mathrm{op}} \|x - y\|$ for $x, y \in \mathcal{X}$; hence $T$ is Lipschitz.
*(iii)* $\Rightarrow$ *(ii)* Since $0 \in \mathcal{X}$, continuity on $\mathcal{X}$ implies continuity at 0.
*(ii)* $\Rightarrow$ *(i)* Choose $\delta > 0$ with $\|z\| < \delta \Rightarrow \|Tz\| < 1$. For $x \ne 0$, set $z := \frac{\delta x}{2\|x\|}$.
Then $\|z\| < \delta$ and $\|Tx\| = \frac{2\|x\|}{\delta} \|Tz\| \le \frac{2}{\delta} \|x\|$. Thus $\|T\|_{\mathrm{op}} \le \frac{2}{\delta}$.

*Structural Commentary*: Continuity at 0 rescales to a global operator bound.
*Key Insight*: Linear continuity is boundedness. □

**Theorem 5.1.6.** *Bounded Operators Form a Banach Space*
*Let $\mathcal{X}, \mathcal{Y}$ be normed spaces.*

*(i) $\mathcal{L}(\mathcal{X}, \mathcal{Y})$ is normed by $\|\cdot\|_{op}$ under pointwise operations.*
*(ii) If $\mathcal{Y}$ is Banach, then $\mathcal{L}(\mathcal{X}, \mathcal{Y})$ is Banach.*

*Proof.*

(i) If $S, T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ and $\alpha, \beta \in \mathbb{F}$, then $\|(\alpha S + \beta T)x\| \le (|\alpha| \|S\|_{\text{op}} + |\beta| \|T\|_{\text{op}}) \|x\|$, so $\alpha S + \beta T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$.
Also $\|\alpha T\|_{\text{op}} = |\alpha| \|T\|_{\text{op}}$, $\|S + T\|_{\text{op}} \le \|S\|_{\text{op}} + \|T\|_{\text{op}}$, and $\|T\|_{\text{op}} = 0 \Rightarrow Tx = 0$ for all $x \in \mathcal{X}$.
(ii) Assume $\mathcal{Y}$ is Banach and let $(T_n)$ be Cauchy in $\|\cdot\|_{\text{op}}$.
For each $x \in \mathcal{X}$, $\|T_n x - T_m x\| \le \|T_n - T_m\|_{\text{op}} \|x\|$, so define $Tx := \lim_n T_n x$.
Then $T(\alpha x + \beta y) = \alpha Tx + \beta Ty$. Given $\varepsilon > 0$, choose $N$ with $\|T_n - T_m\|_{\text{op}} < \varepsilon$ for $m, n \ge N$.
Letting $m \to \infty$ gives $\|(T_n - T)x\| \le \varepsilon \|x\|$, so $\|T_n - T\|_{\text{op}} \le \varepsilon$.
Hence $T_N - T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$, $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$, and $T_n \to T$.

*Structural Commentary*: Pointwise limits are linear; operator-norm Cauchy control is uniform in $x$.
*Key Insight*: Completeness of $\mathcal{Y}$ lifts to $\mathcal{L}(\mathcal{X}, \mathcal{Y})$. □

**Corollary 5.1.7.** *Neumann Series*
*Let $\mathcal{X}$ be Banach and let $T \in \mathcal{L}(\mathcal{X})$ satisfy $\|T\|_{op} < 1$.*

*Then $I - T$ has bounded inverse, with Neumann series converging in $\mathcal{L}(\mathcal{X})$: $(I - T)^{-1} = \sum_{n=0}^{\infty} T^n$.*

*Proof.* Let $S_N := \sum_{n=0}^{N} T^n$. For $M > N$, $\|S_M - S_N\|_{\text{op}} \le \sum_{n=N+1}^{M} \|T\|_{\text{op}}^n \le \frac{\|T\|_{\text{op}}^{N+1}}{1 - \|T\|_{\text{op}}} \to 0$, so $S_N \to S \in \mathcal{L}(\mathcal{X})$.
Since $(I - T)S_N = I - T^{N+1} = S_N(I - T)$ and $\|T^{N+1}\|_{\text{op}} \le \|T\|_{\text{op}}^{N+1} \to 0$.
Passing to the limit gives $(I - T)S = S(I - T) = I$.
*Structural Commentary*: Operator-norm convergence turns the geometric identity into an inverse.
*Key Insight*: If $\|T\|_{\text{op}} < 1$, then $\sum T^n = (I - T)^{-1}$. □

**Proposition 5.1.8.** *Quotient Norm and Quotient Map*
*Let $M \le \mathcal{X}$ be closed, let $\mathcal{X}/M := \{x + M : x \in \mathcal{X}\}$, and let $q : \mathcal{X} \to \mathcal{X}/M$ be $q(x) := x + M$.*
*Set $\|x + M\|_{\mathcal{X}/M} := \inf_{m \in M} \|x - m\|$.*

*(i) $\|\cdot\|_{\mathcal{X}/M}$ is a norm on $\mathcal{X}/M$.*
*(ii) $q$ is continuous with $\|q\|_{op} \le 1$.*
*(iii) $\forall x \in \mathcal{X}$ and $r > 0$, $q(B_r(x)) = B_r(qx)$.*
*(iv) $q$ is open.*

*Proof.*

(i) If $x' = x + m_0$ with $m_0 \in M$, then $\inf_{m \in M} \|x' - m\| = \inf_{m \in M} \|x - m\|$, so the formula is well defined.
Also $\|x + M\| = 0 \iff x \in \overline{M} = M$. For $\alpha \ne 0$,
$\|\alpha(x + M)\| = \inf_{m \in M} \|\alpha x - m\| = |\alpha| \inf_{m \in M} \|x - \alpha^{-1} m\| = |\alpha| \|x + M\|$; $\alpha = 0$ is trivial.
For $m, n \in M$, $\|(x + y) - (m + n)\| \le \|x - m\| + \|y - n\|$; take infima.
(ii) Since $\|q(x)\| \le \|x - 0\| = \|x\|$, $q \in \mathcal{L}(\mathcal{X}, \mathcal{X}/M)$ and $\|q\|_{\text{op}} \le 1$.
(iii) If $z \in B_r(x)$, then $\|qz - qx\| \le \|z - x\| < r$, so $q(B_r(x)) \subseteq B_r(qx)$. Conversely, if $\eta = qz$ and $\|\eta - qx\| < r$, choose $m \in M$ with $\|z - x - m\| < r$; then $z - m \in B_r(x)$ and $q(z - m) = \eta$.
(iv) If $U \subseteq \mathcal{X}$ is open and $y \in q(U)$, choose $x \in U$ with $qx = y$ and $r > 0$ with $B_r(x) \subseteq U$.
By (iii), $B_r(y) = q(B_r(x)) \subseteq q(U)$; hence $q(U)$ is open.

*Structural Commentary*: Quotient norm is distance to the closed subspace $M$.
*Key Insight*: Closedness makes zero distance equivalent to membership. □

**Theorem 5.1.9.** *Banach Quotients*
*If $M$ is a closed linear subspace of a Banach space $\mathcal{X}$, then $\mathcal{X}/M$ is Banach in the quotient norm.*

*Proof.* Let $(\xi_n)$ be Cauchy in $\mathcal{X}/M$. Choose $n_1 < n_2 < \cdots$ with $\|\xi_{n_{k+1}} - \xi_{n_k}\| < 2^{-k}$ and choose $u_k \in \mathcal{X}$ with $q(u_k) = \xi_{n_{k+1}} - \xi_{n_k}$, $\|u_k\| < 2^{-k}$. Since $\sum_k \|u_k\| < +\infty$ and $\mathcal{X}$ is Banach, $u := \sum_{k \ge 1} u_k$ exists.
For $K \ge 1$, $\xi_{n_K} = \xi_{n_1} + q(\sum_{k<K} u_k) \to \xi_{n_1} + q(u)$.

Since $(\xi_n)$ is Cauchy and has a convergent subsequence, $\xi_n \to \xi_{n_1} + q(u)$.
*Structural Commentary*: Rapid quotient increments lift to an absolutely convergent series.
*Key Insight*: A Cauchy sequence with a convergent subsequence converges. □

**Proposition 5.1.10.** *Operator Factor through the Kernel Quotient*
*Let $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$, set $M := \ker T$, and let $\widetilde{T} : \mathcal{X}/M \to \mathcal{Y}$ be induced by Theorem 2.2.12.*

*(i) $\widetilde{T}$ is bounded.*
*(ii) $\left\|\widetilde{T}\right\|_{op} = \|T\|_{op}$.*

*Proof.* Since $q(x) = q(y) \iff x - y \in M = \ker T$, the formula $\widetilde{T}(qx) := Tx$ is well defined and $T = \widetilde{T}q$.
For $m \in M$, $\left\|\widetilde{T}(qx)\right\| = \|Tx\| = \|T(x-m)\| \le \|T\|_{\mathrm{op}} \|x - m\|$; taking $\inf_{m \in M}$ gives $\left\|\widetilde{T}(qx)\right\| \le \|T\|_{\mathrm{op}} \|qx\|$.

(i) The estimate gives $\widetilde{T} \in \mathcal{L}(\mathcal{X}/M, \mathcal{Y})$.
(ii) The estimate gives $\left\|\widetilde{T}\right\|_{\mathrm{op}} \le \|T\|_{\mathrm{op}}$. Since $T = \widetilde{T}q$ and $\|q\|_{\mathrm{op}} \le 1$, $\|T\|_{\mathrm{op}} \le \left\|\widetilde{T}\right\|_{\mathrm{op}} \|q\|_{\mathrm{op}} \le \left\|\widetilde{T}\right\|_{\mathrm{op}}$.

*Structural Commentary*: Collapsing $\ker T$ makes $T$ descend without changing its norm.
*Key Insight*: The quotient infimum is taken over exactly the invisible directions of $T$. □

**Example 5.1.11.** *Standard Banach Space Examples*
Standard Banach spaces include the following canonical examples.

(i) If $S$ is a set and $B(S) := \{f : S \to \mathbb{F} : \|f\|_\infty < +\infty\}$, then $(B(S), \|\cdot\|_\infty)$ is Banach.
(ii) $\ell^\infty := \{x = (x_n) : \sup_n |x_n| < +\infty\}$ is Banach under $\|x\|_\infty := \sup_n |x_n|$.
(iii) For $1 \le p < \infty$, $\ell^p := \{x = (x_n) : \sum_n |x_n|^p < +\infty\}$ is Banach under $\|x\|_p := (\sum_n |x_n|^p)^{1/p}$.
(iv) For a measure space $(X, \mathcal{M}, \mu)$ and $1 \le p \le \infty$, $L^p(\mu)$ is Banach.
(v) $L^2(\mu)$ is Hilbert with $\langle f, g\rangle := \int_X f\overline{g}\, d\mu$.

**Example 5.1.12.** *Completeness of $\ell^\infty$ and $\ell^p$*
For $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, $(\ell^\infty(\mathbb{F}), \|\cdot\|_\infty)$ and $(\ell^p(\mathbb{F}), \|\cdot\|_p)$, $1 \le p < \infty$, are complete.

*Proof.* For $\ell^\infty$, let $\mathbf{x}^{(n)} = (x_j^{(n)})_j$ be Cauchy. Then $x_j^{(n)} \to y_j \in \mathbb{F}$ for each $j$.
If $\left\|\mathbf{x}^{(n)} - \mathbf{x}^{(m)}\right\|_\infty < \varepsilon$ for $m, n \ge N$, then $\left|x_j^{(n)} - y_j\right| \le \varepsilon$ for all $j$, so $\left\|\mathbf{x}^{(n)} - \mathbf{y}\right\|_\infty \le \varepsilon$.
Taking $n = N$ gives $\mathbf{y} \in \ell^\infty$, hence $\mathbf{x}^{(n)} \to \mathbf{y}$.
For $\ell^p$, let $\mathbf{x}^{(n)}$ be Cauchy and define $\mathbf{y} = (y_j)$ by coordinatewise limits.
If $\left\|\mathbf{x}^{(n)} - \mathbf{x}^{(m)}\right\|_p < \varepsilon$ for $m, n \ge N$, then for each $r$, $\sum_{j=1}^{r} \left|x_j^{(n)} - y_j\right|^p = \lim_{m \to \infty} \sum_{j=1}^{r} \left|x_j^{(n)} - x_j^{(m)}\right|^p \le \varepsilon^p$.
Letting $r \to \infty$ gives $\left\|\mathbf{x}^{(n)} - \mathbf{y}\right\|_p \le \varepsilon$ for $n \ge N$. Taking $n = N$ gives $\mathbf{y} \in \ell^p$, hence $\mathbf{x}^{(n)} \to \mathbf{y}$.
*Structural Commentary*: Completeness of scalar coordinates combines with uniform tail control.
*Key Insight*: Norm-Cauchy control keeps coordinatewise limits inside the space. □

**Example 5.1.13.** *Two Metrics on $C[0,1]$*
For $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, $C_{\mathbb{F}} := C([0,1], \mathbb{F})$ is complete under $d_\infty(f,g) := \sup_{x \in [0,1]} |f(x) - g(x)|$ by Theorem 8.1.1, but not under $d_1(f,g) := \int_0^1 |f(x) - g(x)|\, dx$.

*Proof.* For $n \ge 2$, define $f_n \in C_{\mathbb{F}}$ by

$$f_n(x) := \begin{cases} 0, & 0 \le x \le \frac{1}{2}, \\ n(x - \frac{1}{2}), & \frac{1}{2} \le x \le \frac{1}{2} + \frac{1}{n}, \\ 1, & \frac{1}{2} + \frac{1}{n} \le x \le 1. \end{cases}$$

Let $s := \mathbf{1}_{(1/2,1]}$. Then $d_1(f_n, s) = 1/(2n)$, hence $(f_n)$ is $d_1$-Cauchy. If $f_n \to g \in C_{\mathbb{F}}$ in $d_1$, then $d_1(g, s) = 0$, so $g = s$ a.e. Continuity forces $g = 0$ on $[0, 1/2]$ and $g = 1$ on $(1/2, 1]$, impossible at $1/2$.
*Structural Commentary*: The integral metric permits a continuous ramp to collapse to a jump.
*Key Insight*: Supremum control preserves continuity; integral control can miss jumps. □




Figure 222: The continuous ramps $f_n$ converge in the integral metric to a discontinuous step function.

**Theorem 5.1.14.** *Finite-Dimensional Normed Spaces*
*Let $\mathcal{X}$ be a finite-dimensional normed space.*

*(i) $\mathcal{X}$ is Banach.*
*(ii) Every subspace of $\mathcal{X}$ is closed.*
*(iii) $\overline{B}_{\mathcal{X}}$ is compact.*

*Proof.* Choose a basis $e_1, \dots, e_n$ and set $A : \mathbb{F}^n \to \mathcal{X}$, $Aa := \sum_{i=1}^{n} a_i e_i$.
Then $A$ is linear bijective and $\|Aa\| \le \sum_i |a_i| \, \|e_i\| \le C \, \|a\|_2$.
Since $a \mapsto \|Aa\|$ is continuous, positive on $\{a : \|a\|_2 = 1\}$, $\exists c > 0$ with $c \, \|a\|_2 \le \|Aa\| \le C \, \|a\|_2 \ \forall a \in \mathbb{F}^n$.

(i) If $(x_k)$ is Cauchy in $\mathcal{X}$, then $a_k := A^{-1} x_k$ is Cauchy in $\mathbb{F}^n$ by $c \, \|a_k - a_\ell\|_2 \le \|x_k - x_\ell\|$.
Since $\mathbb{F}^n$ is complete, $a_k \to a$, hence $x_k \to Aa$.
(ii) If $M \le \mathcal{X}$, then $M$ is finite-dimensional, hence Banach by (i). If $x_k \in M$ and $x_k \to x$ in $\mathcal{X}$, then $(x_k)$ converges in $M$ to some $y \in M$; uniqueness of limits gives $x = y \in M$. Thus $M$ is closed.
(iii) Since $\overline{B}_{\mathcal{X}} \subseteq A(\overline{B}_2(0, \frac{1}{c}))$ and $A^{-1}(\overline{B}_{\mathcal{X}})$ is closed in the compact Euclidean ball $\overline{B}_2(0, \frac{1}{c})$, $A^{-1}(\overline{B}_{\mathcal{X}})$ is compact; continuity of $A$ gives compactness of $\overline{B}_{\mathcal{X}}$.

*Structural Commentary*: The coordinate map is a bi-Lipschitz linear isomorphism.
*Key Insight*: Finite-dimensional normed spaces inherit completeness and compactness from $\mathbb{F}^n$. □

**Theorem 5.1.15.** *Riesz Lemma*
*Let $M$ be a closed proper linear subspace of a normed space $\mathcal{X}$.*
*Then $\forall \varepsilon \in (0,1)$ $\exists u \in \mathcal{X}$ such that $\|u\| = 1$ and $\forall m \in M$, $\|u - m\| > 1 - \varepsilon$.*

*Proof.* Choose $x \in \mathcal{X} \setminus M$ and set $d := \inf_{m \in M} \|x - m\|$. Since $M$ is closed, $d > 0$.
Choose $m_0 \in M$ with $d \le \|x - m_0\| < \frac{d}{1-\varepsilon}$ and set $u := \frac{x - m_0}{\|x - m_0\|}$.
Then $\|u\| = 1$. For $m \in M$, $m_0 + \|x - m_0\| \, m \in M$, so $\|u - m\| = \frac{\|x - (m_0 + \|x - m_0\| m)\|}{\|x - m_0\|} \ge \frac{d}{\|x - m_0\|} > 1 - \varepsilon$.
*Structural Commentary*: A positive distance to $M$ survives normalization.
*Key Insight*: Normalize an almost-minimizer of $\operatorname{dist}(x, M)$. □

**Corollary 5.1.16.** *Compact Unit Balls Characterize Finite Dimension*
*For a normed space $\mathcal{X}$, $\overline{B}_{\mathcal{X}}$ is compact iff $\dim \mathcal{X} < \infty$.*

*Proof.*
($\Leftarrow$) If $\dim \mathcal{X} < \infty$, then $\overline{B}_{\mathcal{X}}$ is compact by Theorem 5.1.14(iii).
($\Rightarrow$) Suppose $\dim \mathcal{X} = \infty$. Choose $x_1 \in \mathcal{X}$ with $\|x_1\| = 1$. Given $x_1, \dots, x_n$, set $M_n := \operatorname{span}\{x_1, \dots, x_n\}$.
Then $M_n$ is finite-dimensional, closed, and proper, so Theorem 5.1.15 with $\varepsilon = \frac{1}{2}$ gives $x_{n+1} \in \mathcal{X}$ with $\|x_{n+1}\| = 1$ and $\operatorname{dist}(x_{n+1}, M_n) > \frac{1}{2}$. Since $x_i \in M_n$ for $i \le n$, $\|x_{n+1} - x_i\| > \frac{1}{2}$.
Thus $(x_n) \subseteq \overline{B}_{\mathcal{X}}$ has no Cauchy subsequence, so $\overline{B}_{\mathcal{X}}$ is not compact.

*Structural Commentary*: Riesz iteration builds a separated sequence in $\overline{B}_{\mathcal{X}}$.
*Key Insight*: Compact unit ball forces finite dimension. □




Figure 223: Riesz iteration: $M_n = \operatorname{span}\{x_1, \dots, x_n\}$, $\operatorname{dist}(x_{n+1}, M_n) > \frac{1}{2} \Rightarrow \|x_i - x_j\| > \frac{1}{2}$ $(i \neq j)$; hence no Cauchy subsequence in $\overline{B}_{\mathcal{X}}$.

## 5.2 Hilbert Spaces

Hilbert spaces are complete inner-product spaces. They extend the orthogonal geometry of Definition 1.1.3 to function and sequence spaces, supporting projection, Fourier expansion, and Riesz representation.

**Definition 5.2.1.** *Hilbert Space*
Let $\mathcal{H}$ be an inner-product space.

(i) *Hilbert space*: $\mathcal{H}$ is complete in the norm $\|x\| := \sqrt{\langle x, x\rangle}$.
(ii) *Orthonormal family*: $(e_i)_{i\in I} \subseteq \mathcal{H}$ satisfies $\forall i, j \in I$, $\langle e_i, e_j\rangle = \delta_{ij}$.

**Proposition 5.2.2.** *Continuity of the Inner Product*
*Let $\mathcal{V}$ be an inner-product space, let $(x_n), (y_n) \subseteq \mathcal{V}$, and let $x, y \in \mathcal{V}$.*

*(i)* $x_n \to x \Longrightarrow \|x_n\| \to \|x\|$.
*(ii)* $x_n \to x$ *and* $y_n \to y \Longrightarrow \langle x_n, y_n\rangle \to \langle x, y\rangle$.

*Proof.*

(i) $|\|x_n\| - \|x\|| \leq \|x_n - x\| \to 0$, hence $\|x_n\| \to \|x\|$.
(ii) Since $(y_n)$ is bounded, choose $M < \infty$ with $\|y_n\| \leq M$.
Cauchy–Schwarz gives $|\langle x_n, y_n\rangle - \langle x, y\rangle| \leq M\|x_n - x\| + \|x\|\,\|y_n - y\| \to 0$.

*Structural Commentary*: Cauchy–Schwarz turns norm convergence in either coordinate into scalar convergence.
*Key Insight*: Inner products are continuous because their errors are controlled by products of norms. □

**Theorem 5.2.3.** *Hilbert Completion*
*Let $\mathcal{V}$ be an inner-product space.*

*(i)* $\exists(\widehat{\mathcal{V}}, J)$ *such that* $\widehat{\mathcal{V}}$ *is Hilbert,* $J : \mathcal{V} \to \widehat{\mathcal{V}}$ *is a linear isometry, and* $\overline{J(\mathcal{V})} = \widehat{\mathcal{V}}$.
*(ii) If $(\mathcal{K}, I)$ also satisfies (i), then $\exists! U : \widehat{\mathcal{V}} \to \mathcal{K}$ a surjective linear isometry with $U \circ J = I$.*

*Proof.*

(i) Let $C(\mathcal{V})$ be the Cauchy sequences, set $(x_n) \sim (y_n) \iff \|x_n - y_n\| \to 0$, and define $\widehat{\mathcal{V}} := C(\mathcal{V})/\sim$.
Set $[x_n] + [y_n] := [x_n + y_n]$, $a[x_n] := [ax_n]$, $Jx := [(x)]$, and $\langle [x_n], [y_n]\rangle_{\widehat{\mathcal{V}}} := \lim_{n\to\infty} \langle x_n, y_n\rangle_{\mathcal{V}}$.
If $(x_n), (y_n)$ are bounded by $M$, then $|\langle x_n, y_n\rangle - \langle x_m, y_m\rangle| \leq M\|y_n - y_m\| + M\|x_n - x_m\| \to 0$.
The same estimate for equivalent representatives makes the limit well defined.
Also $\|[x_n] - [y_n]\| = \lim_n \|x_n - y_n\|$, the metric-completion norm; hence $\widehat{\mathcal{V}}$ is Hilbert and $\overline{J(\mathcal{V})} = \widehat{\mathcal{V}}$.
(ii) Since $(\widehat{\mathcal{V}}, J)$ and $(\mathcal{K}, I)$ are metric completions of $\mathcal{V}$, there is a unique surjective isometry $U : \widehat{\mathcal{V}} \to \mathcal{K}$ with $U \circ J = I$. For $x, y \in \mathcal{V}$ and $a, b \in \mathbb{F}$, $U(aJx + bJy) = U(J(ax + by)) = I(ax + by) = aUJx + bUJy$; by density and continuity, $U$ is linear. Since $\|Uz - Uw\| = \|z - w\|$, polarization gives $\langle Uz, Uw\rangle_{\mathcal{K}} = \langle z, w\rangle_{\widehat{\mathcal{V}}}$.

*Structural Commentary*: Complete the metric first, then recover the inner product by taking limits.
*Key Insight*: Cauchy–Schwarz makes the lifted inner product well defined. □

**Theorem 5.2.4.** *Complete, Closed, and Finite-Dimensional Subspaces*
*Let $\mathcal{S}$ be a subspace of an inner-product space $\mathcal{V}$.*

*(i) If $\mathcal{S}$ is complete in the inherited metric, then $\mathcal{S}$ is closed in $\mathcal{V}$.*
*(ii) If $\mathcal{V}$ is Hilbert, then $\mathcal{S}$ is Hilbert if and only if $\mathcal{S}$ is closed.*
*(iii) If $\mathcal{S}$ is finite-dimensional, then $\mathcal{S}$ is complete and closed in $\mathcal{V}$.*

*Proof.*

(i) Let $v \in \overline{\mathcal{S}}$ and choose $s_n \in \mathcal{S}$ with $s_n \to v$ in $\mathcal{V}$. Then $(s_n)$ is Cauchy in $\mathcal{S}$, so $s_n \to s \in \mathcal{S}$ by completeness. Uniqueness of limits gives $v = s$, hence $\overline{\mathcal{S}} = \mathcal{S}$.

(ii) Assume $\mathcal{V}$ Hilbert. If $\mathcal{S}$ is closed and $(s_n) \subseteq \mathcal{S}$ is Cauchy, then $s_n \to v \in \mathcal{V}$ and $v \in \mathcal{S}$, so $\mathcal{S}$ is complete. Conversely, if $\mathcal{S}$ is Hilbert, then $\mathcal{S}$ is complete, hence closed by (i).

(iii) Let $\dim \mathcal{S} = m$ and choose an orthonormal basis $e_1, \dots, e_m$.

If $(s_n)$ is Cauchy in $\mathcal{S}$ and $s_n = \sum_{j=1}^{m} a_{n,j} e_j$, then $|a_{n,j} - a_{k,j}| = \left|\langle s_n - s_k, e_j \rangle\right| \le \|s_n - s_k\|$.

Thus $a_{n,j} \to a_j \in \mathbb{F}$ for each $j$, and $s_n \to s := \sum_{j=1}^{m} a_j e_j \in \mathcal{S}$. Hence $\mathcal{S}$ is complete, so closed by (i).

*Structural Commentary*: In a complete ambient space, closed subspaces are exactly complete subspaces.
*Key Insight*: Finite orthonormal coordinates reduce completeness to $\mathbb{F}^m$. □

**Definition 5.2.5.** *Series in an Inner-Product Space*
Let $(x_k) \subseteq \mathcal{V}$.

(i) The *partial sums* are $s_n := \sum_{k=1}^{n} x_k$.

(ii) The *series* $\sum_{k=1}^{\infty} x_k$ *converges* to $s \in \mathcal{V}$ iff $s_n \to s$.

(iii) The series is *absolutely convergent* iff $\sum_{k=1}^{\infty} \|x_k\| < \infty$.

**Theorem 5.2.6.** *Completeness via Absolutely Convergent Series*
*$\mathcal{V}$ is complete $\iff$ every series $\sum_{k=1}^{\infty} x_k$ in $\mathcal{V}$ with $\sum_{k=1}^{\infty} \|x_k\| < \infty$ converges in $\mathcal{V}$.*

*Proof.*

($\Rightarrow$) Let $s_n := \sum_{k=1}^{n} x_k$ and assume $\sum_k \|x_k\| < +\infty$. For $n > m$, $\|s_n - s_m\| \le \sum_{k=m+1}^{n} \|x_k\| \to 0$ as $m, n \to \infty$. Thus $(s_n)$ is Cauchy, hence convergent by completeness.

($\Leftarrow$) Let $(x_n)$ be Cauchy. Let $N_1 < N_2 < \cdots$ such that $i, j \ge N_k \Rightarrow \|x_i - x_j\| < 2^{-k}$, $y_k := x_{N_{k+1}} - x_{N_k}$. Then $\sum_k \|y_k\| \le \sum_k 2^{-k} < +\infty$, so $\sum_k y_k$ converges. Since $x_{N_m} = x_{N_1} + \sum_{k<m} y_k$, the subsequence $(x_{N_m})$ converges; because $(x_n)$ is Cauchy, $(x_n)$ converges.

*Structural Commentary*: Absolute tails make partial sums Cauchy; Cauchy tails contain summable increments.
*Key Insight*: Completeness is summability of norm-controlled series. □

**Example 5.2.7.** *$\ell^2$ and the Parallelogram Test*
Let $\ell^2 := \{x = (x_n) : \sum_{n=1}^{\infty} |x_n|^2 < \infty\}$. With $\langle x, y \rangle := \sum_{n=1}^{\infty} x_n \overline{y_n}$ and $\|x\|_2 = \langle x, x \rangle^{\frac{1}{2}}$, Cauchy coordinate limits and finite-set estimates give completeness, so $\ell^2$ is Hilbert.
For $p \neq 2$, $\|\cdot\|_p$ is not inner-product-induced: with $x = (1, 1, 0, \dots)$ and $y = (1, -1, 0, \dots)$,
$\|x + y\|_p^2 + \|x - y\|_p^2 = 8$ while $2\|x\|_p^2 + 2\|y\|_p^2 = 4 \cdot 2^{\frac{2}{p}}$, so the parallelogram identity holds only for $p = 2$.
*Structural Commentary*: Completeness alone does not make a Banach space Hilbert.
*Key Insight*: The parallelogram identity detects whether a norm really came from an inner product.

**Example 5.2.8.** *The Supremum Norm Is Not Hilbertian*
For $a < b$, $C([a, b], \mathbb{F})$ with $\|h\|_\infty := \sup_{t \in [a,b]} |h(t)|$ is Banach but not Hilbertian: take $f(t) = 1$ and $g(t) = \frac{t-a}{b-a}$, so $\|f\|_\infty = \|g\|_\infty = 1$, $\|f + g\|_\infty = 2$, $\|f - g\|_\infty = 1$, and $\|f + g\|_\infty^2 + \|f - g\|_\infty^2 = 5 \neq 4 = 2\|f\|_\infty^2 + 2\|g\|_\infty^2$, violating the parallelogram identity.
*Structural Commentary*: Banach need not mean Hilbert.
*Key Insight*: One failed parallelogram identity rules out every possible inner product inducing the norm.

**Theorem 5.2.9.** *Hilbert Projection Theorem*
*Let $C \subseteq \mathcal{H}$ be nonempty, closed, and convex. Then $\forall x \in \mathcal{H}$ $\exists! p \in C$ such that $\|x - p\| = \inf_{q \in C} \|x - q\|$.*

*Proof.* Let $d := \inf_{q \in C} \|x - q\|$, $q_n \in C$, $\|x - q_n\| \to d$. Since $\frac{q_n + q_m}{2} \in C$, by parallelogram identity,
$\|q_n - q_m\|^2 = 2\|x - q_n\|^2 + 2\|x - q_m\|^2 - 4\left\|x - \frac{q_n + q_m}{2}\right\|^2 \le 2\|x - q_n\|^2 + 2\|x - q_m\|^2 - 4d^2 \to 0$.
Thus $(q_n)$ is Cauchy. Hence $q_n \to p \in \mathcal{H}$; closedness gives $p \in C$, and $\|x - p\| = d$.
If $p_1, p_2 \in C$ both minimize, the same identity gives $\|p_1 - p_2\|^2 \le 2d^2 + 2d^2 - 4d^2 = 0$.
*Structural Commentary*: Convexity and parallelogram force minimizing sequences to converge in $C$.
*Key Insight*: Midpoint minimality turns near minimizers into a Cauchy sequence. □




Figure 224: Hilbert projection: $q_n \in C$, $\|x - q_n\| \to d = \operatorname{dist}(x, C) \Rightarrow q_n \to p \in C$ and $\|x - p\| = d$.

**Corollary 5.2.10.** *Orthogonal Projection onto a Closed Subspace*
*Let $M \leq \mathcal{H}$ be closed.*

*(i)* $\forall x \in \mathcal{H}\ \exists! p \in M$ *such that* $x - p \in M^\perp$.
*(ii)* $\mathcal{H} = M \odot M^\perp$.

*Proof.*

(i) By Theorem 5.2.9, choose the unique $p \in M$ minimizing $\|x - p\|$, and set $r := x - p$.
For $y \in M$ and $t \in \mathbb{R}$, $p + ty \in M$, so $0 \leq \|r - ty\|^2 - \|r\|^2 = -2t \operatorname{Re}\langle r, y\rangle + t^2 \|y\|^2$.
Thus $\operatorname{Re}\langle r, y\rangle = 0$; over $\mathbb{C}$, replace $y$ by $iy$ to get $\operatorname{Im}\langle r, y\rangle = 0$. Hence $r \in M^\perp$.
If $x = p_1 + r_1 = p_2 + r_2$ with $p_i \in M$, $r_i \in M^\perp$, then $p_1 - p_2 = r_2 - r_1 \in M \cap M^\perp = \{0\}$.
(ii) By (i), each $x \in \mathcal{H}$ has $x = p + r$ with $p \in M$, $r \in M^\perp$, so $\mathcal{H} = M + M^\perp$.
Since $M \cap M^\perp = \{0\}$, the sum is direct: $\mathcal{H} = M \odot M^\perp$.

*Structural Commentary*: Nearest points are characterized by orthogonal residuals.
*Key Insight*: Every vector splits uniquely into $M$ and $M^\perp$ parts. □

**Definition 5.2.11.** *Closed Linear Span and Total Sets*
For $A \subseteq \mathcal{H}$:

(i) *Closed linear span*: $\operatorname{cspan}(A) := \overline{\operatorname{span}(A)}$.
(ii) *Total*: $A$ is total in $\mathcal{H}$ iff $\operatorname{cspan}(A) = \mathcal{H}$.

**Theorem 5.2.12.** *Double Orthogonal Complements in Hilbert Spaces*
*Let $\mathcal{H}$ be a Hilbert space.*

*(i)* $\forall A \subseteq \mathcal{H}$, $\operatorname{cspan}(A) = A^{\perp\perp}$.
*(ii)* $\forall \mathcal{S} \leq \mathcal{H}$, $\overline{\mathcal{S}} = \mathcal{S}^{\perp\perp}$.
*(iii)* $\forall \mathcal{K} \leq \mathcal{H}$, $\mathcal{K}$ *closed* $\iff \mathcal{K} = \mathcal{K}^{\perp\perp}$.
*(iv)* $\forall A \subseteq \mathcal{H}$, $\operatorname{span}(A)$ *dense in* $\mathcal{H} \iff A^\perp = \{\mathbf{0}\}$.

*Proof.*

(i) Let $B := \operatorname{cspan}(A)$. Since $A \subseteq B$, $B^\perp \subseteq A^\perp$. If $y \in A^\perp$, by $y \perp \operatorname{span}(A)$, continuity of $z \mapsto \langle z, y\rangle$, $y \perp B$, so $A^\perp = B^\perp$. By Corollary 5.2.10, $\mathcal{H} = B \odot B^\perp$; $B^{\perp\perp} = B$. Therefore $A^{\perp\perp} = B^{\perp\perp} = B = \operatorname{cspan}(A)$.
(ii) With $A = \mathcal{S}$ in (i), $\mathcal{S}^{\perp\perp} = \operatorname{cspan}(\mathcal{S}) = \overline{\operatorname{span}(\mathcal{S})} = \overline{\mathcal{S}}$.
(iii) By (ii), $\mathcal{K}^{\perp\perp} = \overline{\mathcal{K}}$; thus $\mathcal{K} = \mathcal{K}^{\perp\perp} \iff \mathcal{K} = \overline{\mathcal{K}} \iff \mathcal{K}$ is closed.
(iv) By (i), $\operatorname{span}(A)$ is dense $\iff A^{\perp\perp} = \mathcal{H}$. If $A^\perp = \{\mathbf{0}\}$, then $A^{\perp\perp} = \{\mathbf{0}\}^\perp = \mathcal{H}$.
Conversely, if $A^{\perp\perp} = \mathcal{H}$ and $y \in A^\perp$, then $y \in A^{\perp\perp}$, so $\|y\|^2 = \langle y, y\rangle = 0$; hence $A^\perp = \{\mathbf{0}\}$.

*Structural Commentary*: Double orthogonal complement equals closed linear span.
*Key Insight*: Missing closed-span directions are exactly nonzero orthogonal vectors. □

**Example 5.2.13.** *Completeness Is Needed for Double Complements*
Let $\mathcal{V} := c_{00}(\mathbb{N}; \mathbb{R}) \subseteq \ell^2$ with inherited inner product, $Lx := \sum_{n=1}^{\infty} \frac{x_n}{n}$, and $\mathcal{K} := \ker L$.

Then $|Lx| \leq \|x\|_2 \left(\sum_{n=1}^{\infty} n^{-2}\right)^{\frac{1}{2}}$, so $L$ is continuous and $\mathcal{K}$ is a closed subspace of $\mathcal{V}$.
If $y \in \mathcal{K}^\perp$, then $e_1 - ne_n \in \mathcal{K}$ for $n \geq 2$, hence $0 = \langle y, e_1 - ne_n\rangle = y_1 - ny_n$.
Since $y \in c_{00}$, choose $n \geq 2$ with $y_n = 0$, so $y_1 = 0$ and then $\forall n,\ y_n = 0$.
Thus $\mathcal{K}^\perp = \{\mathbf{0}\}$ and $\mathcal{K}^{\perp\perp} = \mathcal{V}$. But $e_1 \notin \mathcal{K}$, hence $\mathcal{K} \neq \mathcal{V} = \mathcal{K}^{\perp\perp}$.
*Structural Commentary*: Closedness need not recover double complements without completeness.
*Key Insight*: The identity $\mathcal{K} = \mathcal{K}^{\perp\perp}$ is genuinely Hilbert-space geometry.

**Perspective 5.2.14.** *Hilbert and Hamel Basis Caution*
In Hilbert-space theory, an *orthonormal basis* usually means a complete orthonormal family: vectors are recovered by convergent Fourier expansion, not necessarily by finite linear combinations.

A *Hamel basis* is a vector-space basis and is much stronger in infinite dimension.
For example, the standard $(\mathbf{e}_j)$ in $\ell^2(\mathbb{F})$ is a complete orthonormal family, but $\operatorname{span}\{\mathbf{e}_j : j \geq 1\} = \mathbb{F}^{(\mathbb{N})} \subsetneq \ell^2(\mathbb{F})$.

**Definition 5.2.15.** *Hilbert Basis*

(i) A *Hilbert basis*, or complete orthonormal family, is a maximal orthonormal set under inclusion.
(ii) A *Hamel basis* is a vector-space basis.
(iii) Here *orthonormal basis* means Hilbert basis outside finite-dimensional Hamel context.

**Proposition 5.2.16.** *Finite Hilbert Bases Are Hamel Bases*
*Let $\mathcal{O} = \{\mathbf{e}_1, \ldots, \mathbf{e}_m\}$ be orthonormal in $\mathcal{V}$ and set $W = \operatorname{span}(\mathcal{O})$.*
*Then $\mathcal{O}$ is a Hilbert basis $\iff W = \mathcal{V} \iff \mathcal{O}$ is an orthonormal Hamel basis.*

*Proof.* Since $\mathcal{O}$ is orthonormal, it is linearly independent; hence $W = \mathcal{V} \iff \mathcal{O}$ is an orthonormal Hamel basis. It remains to show maximality $\iff W = \mathcal{V}$.

($\Rightarrow$) Suppose $\mathcal{O}$ is maximal orthonormal and $W \neq \mathcal{V}$. Choose $\mathbf{v} \notin W$ and set $\mathbf{u} := \mathbf{v} - \sum_{j=1}^{m} \langle \mathbf{v}, \mathbf{e}_j \rangle \, \mathbf{e}_j$.

Then $\langle \mathbf{u}, \mathbf{e}_i \rangle = 0$ for $1 \leq i \leq m$, and $\mathbf{u} \neq \mathbf{0}$ since $\mathbf{v} \notin W$.
Thus $\mathcal{O} \cup \{\mathbf{u}/\|\mathbf{u}\|\}$ is orthonormal, contradiction. Hence $W = \mathcal{V}$.

($\Leftarrow$) Suppose $W = \mathcal{V}$. If $\mathbf{w} \perp \mathcal{O}$, write $\mathbf{w} = \sum_{j=1}^{m} a_j \mathbf{e}_j$. Then $a_j = \langle \mathbf{w}, \mathbf{e}_j \rangle = 0$ for every $j$, so $\mathbf{w} = \mathbf{0}$.

Hence no nonzero vector can be adjoined orthogonally to $\mathcal{O}$; $\mathcal{O}$ is maximal orthonormal.

*Structural Commentary*: In finite orthonormal families, maximality equals spanning.
*Key Insight*: Any missing vector has a nonzero orthogonal residual. □

**Example 5.2.17.** *A Hilbert Basis Need Not Be a Hamel Basis*
In $\ell^2(\mathbb{F})$, let $\mathbf{e}_j = (0, \ldots, 0, 1, 0, \ldots)$. Then $\{\mathbf{e}_j : j \geq 1\}$ is orthonormal and maximal, since $\forall j,\ \mathbf{x} \perp \mathbf{e}_j$ implies $\forall j,\ x_j = \langle \mathbf{x}, \mathbf{e}_j \rangle = 0$. Its finite span is $\mathbb{F}^{(\mathbb{N})} \subsetneq \ell^2(\mathbb{F})$, so it is a Hilbert basis but not a Hamel basis.

**Theorem 5.2.18.** *Hilbert Basis Characterizations*
*Let $\mathcal{O} \subseteq \mathcal{H}$ be orthonormal. The following are equivalent:*

*(i)* $\mathcal{O}$ *is maximal orthonormal.*
*(ii)* $\mathcal{O}^{\perp} = \{\mathbf{0}\}$.
*(iii)* $\operatorname{cspan}(\mathcal{O}) = \mathcal{H}$.

*Proof.*

(*i*) $\Rightarrow$ (*ii*) If $0 \neq v \in \mathcal{O}^{\perp}$, then $\mathcal{O} \cup \{v/\|v\|\}$ is orthonormal, contradicting maximality. Hence $\mathcal{O}^{\perp} = \{\mathbf{0}\}$.
(*ii*) $\Rightarrow$ (*iii*) By Theorem 5.2.12, $\operatorname{cspan}(\mathcal{O}) = \mathcal{O}^{\perp\perp} = (\mathcal{O}^{\perp})^{\perp} = \{\mathbf{0}\}^{\perp} = \mathcal{H}$.
(*iii*) $\Rightarrow$ (*i*) If $\mathcal{O} \subsetneq \mathcal{O} \cup \{u\}$ is orthonormal with $\|u\| = 1$, then $u \in \mathcal{O}^{\perp}$.
By (iii) and Theorem 5.2.12(iv), $\mathcal{O}^{\perp} = \{\mathbf{0}\}$, contradiction. Thus $\mathcal{O}$ is maximal orthonormal.

*Structural Commentary*: Maximality, trivial orthogonal complement, and closed span are equivalent.
*Key Insight*: A Hilbert basis is total, not necessarily Hamel. □

**Definition 5.2.19.** *Arbitrary Sums in a Hilbert Space*
Let $(x_k)_{k \in K} \subseteq \mathcal{H}$.

(i) *Finite partial sums*: $\mathcal{F}(K) := \{J \subseteq K : |J| < \infty\}$, ordered by $\subseteq$, and $s_J := \sum_{k \in J} x_k$.
(ii) *Arbitrary convergence*: $\sum_{k \in K} x_k = x \iff \forall \varepsilon > 0 \, \exists S \in \mathcal{F}(K) \, \forall T \in \mathcal{F}(K) : S \subseteq T \Rightarrow \|s_T - x\| \leq \varepsilon$.
(iii) *Nonnegative scalar sum*: for $(a_k)_{k \in K} \subseteq [0, \infty)$, $\sum_{k \in K} a_k < \infty \iff \sup_{J \in \mathcal{F}(K)} \sum_{k \in J} a_k < \infty$.

**Proposition 5.2.20.** *Basic Properties of Arbitrary Sums*
*Let $(x_k)_{k \in K}, (y_k)_{k \in K} \subseteq \mathcal{H}$ satisfy $\sum_{k \in K} x_k = x$ and $\sum_{k \in K} y_k = y$.*

*(i)* $\forall a, b \in \mathbb{F},\ \sum_{k \in K} (a x_k + b y_k) = ax + by$.
*(ii)* $\forall z \in \mathcal{H},\ \sum_{k \in K} \langle x_k, z \rangle = \langle x, z \rangle$.
*(iii)* $\forall z \in \mathcal{H},\ \sum_{k \in K} \langle z, x_k \rangle = \langle z, x \rangle$.

*Proof.* For $J \in \mathcal{F}(K)$, set $s_J := \sum_{k \in J} x_k$ and $t_J := \sum_{k \in J} y_k$.

(i) Since $s_J \to x$ and $t_J \to y$, $a s_J + b t_J \to ax + by$.

(ii) For $z \in \mathcal{H}$ and $J \in \mathcal{F}(K)$, $\sum_{k\in J} \langle x_k, z\rangle = \langle s_J, z\rangle$; since $s_J \to x$, Proposition 5.2.2 gives $\langle s_J, z\rangle \to \langle x, z\rangle$.

(iii) For $z \in \mathcal{H}$ and $J \in \mathcal{F}(K)$, $\sum_{k\in J} \langle z, x_k\rangle = \langle z, s_J\rangle$; since $s_J \to x$, Proposition 5.2.2 gives $\langle z, s_J\rangle \to \langle z, x\rangle$.

*Structural Commentary*: Arbitrary sums are ordinary finite sums plus a convergence net.
*Key Insight*: Once the vector sum converges, continuous linear tests may pass through the sum. □

**Theorem 5.2.21.** *Arbitrary Orthonormal Sums*
*Let $(u_k)_{k\in K} \subseteq \mathcal{H}$ be orthonormal.*

*(i) For $(r_k)_{k\in K} \subseteq \mathbb{F}$, $\sum_{k\in K} r_k u_k$ converges iff $\sup_{J\in\mathcal{F}(K)} \sum_{k\in J} |r_k|^2 < \infty$.*

*(ii) Under the conditions in (i), $\left\|\sum_{k\in K} r_k u_k\right\|^2 = \sum_{k\in K} |r_k|^2$.*

*(iii) $\forall (x_k)_{k\in K} \subseteq \mathcal{H}$, $\sum_{k\in K} x_k$ convergent implies $|\{k \in K : x_k \neq \mathbf{0}\}| \leq \aleph_0$.*

*Proof.*

(i) ($\Rightarrow$) Let $s_J := \sum_{k\in J} r_k u_k$. If $s_J \to x$, choose $S \in \mathcal{F}(K)$ with $S \subseteq T \Rightarrow \|s_T - x\| \leq 1$.
For $J \in \mathcal{F}(K)$, $\sum_{k\in J} |r_k|^2 \leq \|s_{J\cup S}\|^2 \leq (\|x\| + 1)^2$.

(i) ($\Leftarrow$) Let $M := \sup_{J\in\mathcal{F}(K)} \sum_{k\in J} |r_k|^2 < +\infty$. Given $\varepsilon > 0$, choose $S \in \mathcal{F}(K)$ with $M - \sum_{k\in S} |r_k|^2 < \varepsilon^2$.
If $S \subseteq J, L$, then $\|s_J - s_L\|^2 = \sum_{k\in J\triangle L} |r_k|^2 \leq \sum_{k\in (J\cup L)\setminus S} |r_k|^2 < \varepsilon^2$.
Thus $(s_J)$ is Cauchy, hence converges in $\mathcal{H}$.

(ii) If $s_J \to s := \sum_{k\in K} r_k u_k$, then $\|s\|^2 = \lim_J \|s_J\|^2 = \sup_{J\in\mathcal{F}(K)} \sum_{k\in J} |r_k|^2 = \sum_{k\in K} |r_k|^2$.

(iii) If $s_J := \sum_{k\in J} x_k$ converges, then $(s_J)$ is Cauchy.
For each $n \geq 1$, choose $I_n \in \mathcal{F}(K)$ such that $I_n \subseteq J, L \Rightarrow \|s_J - s_L\| \leq \frac{1}{n}$.
Then $F \in \mathcal{F}(K \setminus I_n)$ implies $\left\|\sum_{k\in F} x_k\right\| = \|s_{I_n\cup F} - s_{I_n}\| \leq \frac{1}{n}$.
With $I := \bigcup_{n\geq 1} I_n$, $I$ is countable; if $k \notin I$, then $\|x_k\| \leq \frac{1}{n}$ for all $n$, so $x_k = \mathbf{0}$.

*Structural Commentary*: Orthonormal finite sums turn net tails into square-sum tails.
*Key Insight*: Convergent arbitrary sums have countable support. □

**Theorem 5.2.22.** *Fourier Expansion in a Hilbert Space*
*Let $\mathcal{O} = (u_k)_{k\in K} \subseteq \mathcal{H}$ be orthonormal. For $x \in \mathcal{H}$, set $\widehat{x}_{\mathcal{O}} := \sum_{k\in K} \langle x, u_k\rangle u_k$.*

*(i) $\widehat{x}_{\mathcal{O}}$ converges in $\mathcal{H}$.*
*(ii) $\|\widehat{x}_{\mathcal{O}}\|^2 = \sum_{k\in K} |\langle x, u_k\rangle|^2 \leq \|x\|^2$.*
*(iii) $\widehat{x}_{\mathcal{O}}$ is the unique minimizer of $\|x - y\|$ over $y \in \operatorname{cspan}(\mathcal{O})$.*
*(iv) $\|\widehat{x}_{\mathcal{O}}\| = \|x\| \iff x \in \operatorname{cspan}(\mathcal{O})$.*

*Proof.*

(i) For $J \in \mathcal{F}(K)$, finite Bessel gives $\sum_{k\in J} |\langle x, u_k\rangle|^2 \leq \|x\|^2$.
Hence $\sup_J \sum_{k\in J} |\langle x, u_k\rangle|^2 < +\infty$, so $\sum_{k\in K} \langle x, u_k\rangle u_k$ converges by Theorem 5.2.21(i).

(ii) Theorem 5.2.21(ii) gives $\|\widehat{x}_{\mathcal{O}}\|^2 = \sum_{k\in K} |\langle x, u_k\rangle|^2$; finite Bessel gives the inequality $\sum_{k\in K} |\langle x, u_k\rangle|^2 \leq \|x\|^2$.

(iii) For $i \in K$, Proposition 5.2.20 gives $\langle \widehat{x}_{\mathcal{O}}, u_i\rangle = \sum_{k\in K} \langle x, u_k\rangle \langle u_k, u_i\rangle = \langle x, u_i\rangle$, so $r := x - \widehat{x}_{\mathcal{O}} \perp \mathcal{O}$.
By continuity, $r \perp M := \operatorname{cspan}(\mathcal{O})$. For $y \in M$, $x - y = r + (\widehat{x}_{\mathcal{O}} - y)$ orthogonally,
so $\|x - y\|^2 = \|r\|^2 + \|\widehat{x}_{\mathcal{O}} - y\|^2 \geq \|r\|^2 = \|x - \widehat{x}_{\mathcal{O}}\|^2$, with equality iff $y = \widehat{x}_{\mathcal{O}}$.

(iv) ($\Rightarrow$) As $x = \widehat{x}_{\mathcal{O}} + r$ orthogonally, $\|x\|^2 = \|\widehat{x}_{\mathcal{O}}\|^2 + \|r\|^2$. If $\|\widehat{x}_{\mathcal{O}}\| = \|x\|$, then $r = 0$, $x = \widehat{x}_{\mathcal{O}} \in \operatorname{cspan}(\mathcal{O})$.

(iv) ($\Leftarrow$) If $x \in \operatorname{cspan}(\mathcal{O})$, the minimizer of $\|x - y\|$ over $y \in \operatorname{cspan}(\mathcal{O})$ is $x$; by (iii), $\widehat{x}_{\mathcal{O}} = x$, $\|\widehat{x}_{\mathcal{O}}\| = \|x\|$.

*Structural Commentary*: Fourier expansion is orthogonal projection onto $\operatorname{cspan}(\mathcal{O})$.
*Key Insight*: Bessel bounds the coefficients; completeness sums them. □

**Corollary 5.2.23.** *Parseval Characterization of Hilbert Bases*
*For an orthonormal family $\mathcal{O} = \{u_k : k \in K\}$ in $\mathcal{H}$, the following are equivalent:*

*(i)* $\mathcal{O}$ *is a Hilbert basis.*
*(ii)* $\forall x \in \mathcal{H},\ x = \sum_{k\in K} \langle x, u_k\rangle\, u_k$.
*(iii)* $\forall x, y \in \mathcal{H}$, *Parseval's identity* $\langle x, y\rangle = \sum_{k\in K} \langle x, u_k\rangle\, \overline{\langle y, u_k\rangle}$ *holds.*
*(iv)* $\forall x \in \mathcal{H},\ \|x\|^2 = \sum_{k\in K} |\langle x, u_k\rangle|^2$.

*Proof.*

(*i*) $\Rightarrow$ (*ii*) By Theorem 5.2.18, $\operatorname{cspan}(\mathcal{O}) = \mathcal{H}$. For $x \in \mathcal{H}$, both $x$ and $\widehat{x}_{\mathcal{O}}$ minimize $\|x - y\|$ over $y \in \operatorname{cspan}(\mathcal{O})$. By uniqueness in Theorem 5.2.22(iii), $x = \widehat{x}_{\mathcal{O}} = \sum_{k\in K} \langle x, u_k\rangle\, u_k$.

(*ii*) $\Rightarrow$ (*iii*) For $x, y \in \mathcal{H}$, (ii) and Proposition 5.2.20 give $\langle x, y\rangle = \sum_{k\in K} \langle x, u_k\rangle\, \langle u_k, y\rangle = \sum_{k\in K} \langle x, u_k\rangle\, \overline{\langle y, u_k\rangle}$.

(*iii*) $\Rightarrow$ (*iv*) Put $y = x$: $\|x\|^2 = \langle x, x\rangle = \sum_{k\in K} \langle x, u_k\rangle\, \overline{\langle x, u_k\rangle} = \sum_{k\in K} |\langle x, u_k\rangle|^2$.

(*iv*) $\Rightarrow$ (*i*) By Theorem 5.2.22(ii), (iv) gives $\|\widehat{x}_{\mathcal{O}}\| = \|x\|$ for every $x \in \mathcal{H}$.
By Theorem 5.2.22(iv), $x \in \operatorname{cspan}(\mathcal{O})$ for every $x$, so $\operatorname{cspan}(\mathcal{O}) = \mathcal{H}$. Theorem 5.2.18 gives (i).

*Structural Commentary*: Totality, Fourier expansion, Parseval, and norm equality are equivalent.
*Key Insight*: Parseval preserves all Hilbert geometry. □

**Theorem 5.2.24.** *Existence and Cardinality of Hilbert Bases*
*Let $\mathcal{H}$ be a Hilbert space.*

*(i)* $\mathcal{H}$ *admits a Hilbert basis.*
*(ii) If $\mathcal{B}$,$\mathcal{C}$ are Hilbert bases of $\mathcal{H}$, then $|\mathcal{B}| = |\mathcal{C}|$.*

*Proof.*

(i) Let $\mathfrak{O}$ be the orthonormal subsets of $\mathcal{H}$, ordered by $\subseteq$. Chains in $\mathfrak{O}$ have orthonormal unions, so Principle 6.5.3 gives maximal $\mathcal{O} \in \mathfrak{O}$; by Theorem 5.2.18, $\mathcal{O}$ is a Hilbert basis.

(ii) Let $\mathcal{B} = \{b_i : i \in I\}$ and $\mathcal{C} = \{c_j : j \in J\}$ be Hilbert bases.
If $|I| < \infty$, Proposition 5.2.16 gives $\dim \mathcal{H} = |I|$; orthonormality gives $|J| \le \dim \mathcal{H}$, hence $|J| < \infty$, and Proposition 5.2.16 gives $|J| = \dim \mathcal{H}$. The case $|J| < \infty$ is symmetric.
Assume $I, J$ infinite. For $i \in I$, set $J_i := \{j \in J : \langle b_i, c_j\rangle \neq 0\}$. By Theorem 5.2.21(iii), $|J_i| \le \aleph_0$.
Since $\mathcal{B}^\perp = \{\mathbf{0}\}$, $j \notin \bigcup_{i\in I} J_i$ is impossible; hence $J = \bigcup_{i\in I} J_i$ and $|J| \le \sum_{i\in I} |J_i| \le \aleph_0\, |I| = |I|$.
Symmetry gives $|I| \le |J|$; Theorem 6.3.1 gives $|I| = |J|$.

*Structural Commentary*: Zorn gives existence; countable coordinate supports compare cardinalities.
*Key Insight*: Hilbert-basis size is intrinsic. □

**Definition 5.2.25.** *Hilbert Dimension*
For a Hilbert space $\mathcal{H}$ with Hilbert basis $\mathcal{B}$, define $\operatorname{hdim}(\mathcal{H}) := |\mathcal{B}|$; this is well defined by Theorem 5.2.24.

**Proposition 5.2.26.** *The Hilbert Space $\ell^2(K)$*
*Let $K$ be a set and set $\ell^2(K) := \{f : K \to \mathbb{F} : \sum_{k\in K} |f(k)|^2 < \infty\}$.*

*(i)* $\ell^2(K)$ *is Hilbert under* $\langle f, g\rangle := \sum_{k\in K} f(k)\overline{g(k)}$.
*(ii) The coordinate family $(\delta_k)_{k\in K}$ is a Hilbert basis of $\ell^2(K)$.*

*Proof.*

(i) For $f, g \in \ell^2(K)$ and finite $F \subseteq K$, $\sum_{k\in F} \left|f(k)\overline{g(k)}\right| \le \left(\sum_{k\in F} |f(k)|^2\right)^{1/2} \left(\sum_{k\in F} |g(k)|^2\right)^{1/2} \le \|f\|_2\, \|g\|_2$, so $\langle f, g\rangle$ is well defined. If $(f_n)$ is Cauchy, then $|f_m(k) - f_n(k)| \le \|f_m - f_n\|_2$ for each $k$, so set $f(k) := \lim_n f_n(k)$.
Given $\varepsilon > 0$, choose $N$ with $\|f_m - f_n\|_2 < \varepsilon$ for $m, n \ge N$.
Then for finite $F \subseteq K$ and $n \ge N$, $\sum_{k\in F} |f(k) - f_n(k)|^2 = \lim_m \sum_{k\in F} |f_m(k) - f_n(k)|^2 \le \varepsilon^2$.
Hence $f - f_n \in \ell^2(K)$, $\|f - f_n\|_2 \le \varepsilon$, $f \in \ell^2(K)$, and $f_n \to f$.

(ii) $\langle \delta_j, \delta_k\rangle = \mathbf{1}_{\{j=k\}}$. If $f \perp \delta_k$ for every $k \in K$, then $f(k) = \langle f, \delta_k\rangle = 0$ for every $k$, so $(\delta_k)^\perp_{k\in K} = \{\mathbf{0}\}$.
By Theorem 5.2.18, $(\delta_k)_{k\in K}$ is a Hilbert basis.

*Structural Commentary*: Coordinatewise limits and uniform finite-subset bounds prove completeness.
*Key Insight*: Square-summability is precisely the norm condition for arbitrary orthogonal coordinates. □

**Corollary 5.2.27.** *Classification by Hilbert Dimension*
*For Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$, $\mathcal{H}_1 \cong_{\text{iso}} \mathcal{H}_2$ iff $\operatorname{hdim}(\mathcal{H}_1) = \operatorname{hdim}(\mathcal{H}_2)$.*

*Proof.*
($\Rightarrow$) Let $U : \mathcal{H}_1 \to \mathcal{H}_2$ be a surjective linear isometry and let $\mathcal{B}$ be a Hilbert basis of $\mathcal{H}_1$.
Then $U\mathcal{B}$ is orthonormal and $\operatorname{cspan}(U\mathcal{B}) = U(\operatorname{cspan}\mathcal{B}) = \mathcal{H}_2$.
Thus $U\mathcal{B}$ is a Hilbert basis and $\operatorname{hdim}(\mathcal{H}_1) = |\mathcal{B}| = |U\mathcal{B}| = \operatorname{hdim}(\mathcal{H}_2)$.
($\Leftarrow$) Let $\operatorname{hdim}(\mathcal{H}_1) = \operatorname{hdim}(\mathcal{H}_2) = |I|$, and choose Hilbert bases $(b_i)_{i\in I}$ and $(c_i)_{i\in I}$.
Define $\Phi_1 x := (i \mapsto \langle x, b_i\rangle)$ and $\Phi_2 y := (i \mapsto \langle y, c_i\rangle)$. By Corollary 5.2.23, $\|\Phi_1 x\|_2 = \|x\|$, $\|\Phi_2 y\|_2 = \|y\|$.
If $a = (a_i) \in \ell^2(I)$, then $\sum_{i\in I} a_i b_i$ and $\sum_{i\in I} a_i c_i$ converge by Theorem 5.2.21, with images $a$ under $\Phi_1$ and $\Phi_2$.
Hence $U := \Phi_2^{-1}\Phi_1 : \mathcal{H}_1 \to \mathcal{H}_2$ is a surjective linear isometry.

*Structural Commentary*: Map both spaces isometrically onto the same coordinate model.
*Key Insight*: Hilbert spaces are classified by the number of orthogonal coordinate directions. □

**Corollary 5.2.28.** *Separable Hilbert Spaces*
*$\mathcal{H}$ separable $\iff \operatorname{hdim}(\mathcal{H}) \le \aleph_0$.*

*Proof.*
($\Rightarrow$) Let $D \subseteq \mathcal{H}$ be countable dense and let $\mathcal{O}$ be a Hilbert basis. $\forall u \in \mathcal{O}$, choose $d_u \in D$ with $\|u - d_u\| < \frac{1}{3}$.
If $u \neq v$ and $d_u = d_v$, then $\sqrt{2} = \|u - v\| \le \|u - d_u\| + \|d_v - v\| < \frac{2}{3}$, impossible.
Hence $u \mapsto d_u$ injects $\mathcal{O}$ into $D$, so $\operatorname{hdim}(\mathcal{H}) = |\mathcal{O}| \le \aleph_0$.
($\Leftarrow$) Let $\mathcal{O}$ be a Hilbert basis with $|\mathcal{O}| \le \aleph_0$. The finite $\mathbb{Q}$-span of $\mathcal{O}$ over $\mathbb{R}$, respectively finite $(\mathbb{Q} + i\mathbb{Q})$-span over $\mathbb{C}$, is countable and dense in $\operatorname{cspan}(\mathcal{O}) = \mathcal{H}$.

*Structural Commentary*: Countable bases and countable dense sets force each other.
*Key Insight*: Separability of a Hilbert space is exactly countability of its orthogonal coordinate directions. □

**Corollary 5.2.29.** *Infinite Hilbert Bases Are Not Hamel Bases*
*If $\operatorname{hdim}(\mathcal{H})$ is infinite, no Hilbert basis of $\mathcal{H}$ is a Hamel basis.*

*Proof.* Let $\mathcal{O}$ be Hilbert basis, choose distinct $u_n \in \mathcal{O}$.
Since $\sum_n 4^{-n} < \infty$, $x := \sum_n 2^{-n} u_n \in \mathcal{H}$ by Theorem 5.2.21.

For every $n$, continuity of $\langle \cdot, u_n\rangle$ gives $\langle x, u_n\rangle = 2^{-n} \neq 0$. If $\mathcal{O}$ were Hamel, then $x = \sum_{j=1}^{N} a_j v_j$ with $v_j \in \mathcal{O}$; choosing $n$ with $u_n \notin \{v_1, \dots, v_N\}$ gives $\langle x, u_n\rangle = 0$, contradiction.
*Structural Commentary*: Infinite Hilbert coordinates allow square-summable infinite expansions.
*Key Insight*: A Hilbert basis spans by limits, while a Hamel basis spans by finite sums. □

**Perspective 5.2.30.** *Bounded Functionals*
Here $\mathcal{H}^* = \mathcal{L}(\mathcal{H}, \mathbb{F})$ by Definition 5.1.4 and Theorem 5.1.5.

**Proposition 5.2.31.** *Operator Norm Algebra*
*Let $S, T \in \mathcal{L}(\mathcal{H})$ and $a \in \mathbb{F}$.*

*(i)* $\|aT\|_{op} = |a| \, \|T\|_{op}$.
*(ii)* $\|S + T\|_{op} \le \|S\|_{op} + \|T\|_{op}$.
*(iii)* $\|ST\|_{op} \le \|S\|_{op} \, \|T\|_{op}$.
*(iv) $\mathcal{L}(\mathcal{H})$ is a normed algebra under composition.*

*Proof.*
(i) $\|aT\|_{\text{op}} = \sup_{\|x\|\le 1} \|aTx\| = |a| \sup_{\|x\|\le 1} \|Tx\| = |a| \, \|T\|_{\text{op}}$.
(ii) For $\|x\| \le 1$, $\|(S+T)x\| \le \|Sx\| + \|Tx\| \le \|S\|_{\text{op}} + \|T\|_{\text{op}}$; taking sup gives the claim.
(iii) For $\|x\| \le 1$, $\|STx\| \le \|S\|_{\text{op}} \, \|Tx\| \le \|S\|_{\text{op}} \, \|T\|_{\text{op}}$; taking sup gives the claim.
(iv) Items (i)–(ii), with $\|T\|_{\text{op}} = 0 \Rightarrow T = 0$, give a norm on $\mathcal{L}(\mathcal{H})$.
Item (iii) gives closure and submultiplicativity for associative composition.

*Structural Commentary*: Composition multiplies stretch bounds.
*Key Insight*: Bounded operators behave like matrices with a compatible norm. □

**Theorem 5.2.32.** *Riesz Representation for Hilbert Spaces*
*For $\varphi \in \mathcal{H}^*$:*

*(i) $\exists! y \in \mathcal{H}$ such that $\forall x \in \mathcal{H}$, $\varphi(x) = \langle x, y\rangle$.*

*(ii) The representing vector satisfies* $\|\varphi\|_{op} = \|y\|$.

*Proof.*

(i) If $\varphi = 0$, take $y = 0$. If $\varphi \neq 0$, set $M := \ker \varphi$, $a \notin M$, $a = p + u$, $p \in M$, $u \in M^{\perp}$ by Corollary 5.2.10. Then $u \neq 0$ and $\varphi(u) = \varphi(a) \neq 0$. For $x \in \mathcal{H}$, $x - \frac{\varphi(x)}{\varphi(u)} u \in M$, hence $\langle x, u \rangle = \frac{\varphi(x)}{\varphi(u)} \|u\|^2$. Thus $y := \overline{\varphi(u)} \|u\|^{-2} u$ satisfies $\varphi(x) = \langle x, y \rangle$ for every $x$. If $z$ also represents $\varphi$, then $\langle x, y - z \rangle = 0$ for every $x$; with $x = y - z$, $z = y$.

(ii) For every $x$, $|\varphi(x)| = |\langle x, y \rangle| \leq \|x\| \|y\|$, so $\|\varphi\|_{\text{op}} \leq \|y\|$. If $y \neq 0$, then $\|\varphi\|_{\text{op}} \geq \frac{|\varphi(y)|}{\|y\|} = \|y\|$; if $y = 0$, both norms vanish.

*Structural Commentary*: The kernel is a hyperplane; its orthogonal direction represents the functional.
*Key Insight*: Bounded functionals are exactly inner products with fixed vectors. □

**Corollary 5.2.33.** *Riesz Identifies the Continuous Dual*
*Define* $R : \mathcal{H} \to \mathcal{H}^*$ *by* $Ry := (x \mapsto \langle x, y \rangle)$.

*(i)* $R$ *is an isometric bijection.*
*(ii)* $R$ *is conjugate-linear over* $\mathbb{C}$ *and linear over* $\mathbb{R}$.

*Proof.*

(i) For $y \in \mathcal{H}$, $|Ry(x)| = |\langle x, y \rangle| \leq \|x\| \|y\|$, so $Ry \in \mathcal{H}^*$. By Theorem 5.2.32, $\mathcal{H}^* = R(\mathcal{H})$ and $\|Ry\|_{\text{op}} = \|y\|$. If $Ry = Rz$, then $0 = \|R(y - z)\|_{\text{op}} = \|y - z\|$; hence $R$ is bijective and isometric.

(ii) For $x, y, z \in \mathcal{H}$ and $\alpha, \beta \in \mathbb{C}$, $R(\alpha y + \beta z)(x) = \langle x, \alpha y + \beta z \rangle = \overline{\alpha} Ry(x) + \overline{\beta} Rz(x)$. Over $\mathbb{R}$, $R(\alpha y + \beta z) = \alpha Ry + \beta Rz$.

*Structural Commentary*: Continuous functionals are represented by vectors.
*Key Insight*: Riesz identifies $\mathcal{H}^*$ with $\mathcal{H}$, up to complex conjugation. □

**Theorem 5.2.34.** *Bounded Hilbert Adjoints*
*Let* $T \in \mathcal{L}(\mathcal{H})$.

*(i)* $\exists! T^* \in \mathcal{L}(\mathcal{H})$ *such that* $\forall u, v \in \mathcal{H}$, $\langle Tu, v \rangle = \langle u, T^* v \rangle$.
*(ii)* $\|T^*\|_{op} = \|T\|_{op}$.
*(iii)* $(T^*)^* = T$.

*Proof.*

(i) For $v \in \mathcal{H}$, set $\phi_v(u) := \langle Tu, v \rangle$. Since $|\phi_v(u)| \leq \|T\|_{\text{op}} \|v\| \|u\|$, $\phi_v \in \mathcal{H}^*$. By Theorem 5.2.32, $\exists! T^* v \in \mathcal{H}$ with $\phi_v(u) = \langle u, T^* v \rangle$ for every $u$. For $\alpha, \beta \in \mathbb{F}$, $\phi_{\alpha v + \beta w} = \overline{\alpha} \phi_v + \overline{\beta} \phi_w$, so uniqueness gives $T^*(\alpha v + \beta w) = \alpha T^* v + \beta T^* w$. Also $\|T^* v\| = \|\phi_v\|_{\text{op}} \leq \|T\|_{\text{op}} \|v\|$, hence $T^* \in \mathcal{L}(\mathcal{H})$. If $A$ also satisfies the adjoint identity, then $\langle u, (T^* - A)v \rangle = 0$ for every $u, v$, so $A = T^*$.

(ii) If $\mathcal{H} = \{\mathbf{0}\}$, immediate. Else $\|T\|_{\text{op}} = \sup_{\|u\|=\|v\|=1} |\langle Tu, v \rangle| = \sup_{\|u\|=\|v\|=1} |\langle u, T^* v \rangle| = \|T^*\|_{\text{op}}$.

(iii) For $u, v \in \mathcal{H}$, $\langle T^* u, v \rangle = \overline{\langle v, T^* u \rangle} = \overline{\langle Tv, u \rangle} = \langle u, Tv \rangle$. Thus $T$ satisfies the adjoint identity for $T^*$; by uniqueness, $(T^*)^* = T$.

*Structural Commentary*: Riesz turns the scalar tests induced by an operator back into vectors.
*Key Insight*: The adjoint is the dual action transported through the Hilbert-space Riesz map. □

**Corollary 5.2.35.** *Hilbert Adjoint Algebra Identities*
*Let* $S, T \in \mathcal{L}(\mathcal{H})$ *and* $a \in \mathbb{F}$.

*(i)* $(aS + T)^* = \overline{a} S^* + T^*$.
*(ii)* $(ST)^* = T^* S^*$.

*Proof.*

(i) For $u, v \in \mathcal{H}$, $\langle (aS + T)u, v \rangle = a \langle Su, v \rangle + \langle Tu, v \rangle = \langle u, \overline{a} S^* v + T^* v \rangle$. Uniqueness of adjoints gives $(aS + T)^* = \overline{a} S^* + T^*$.

(ii) For $u, v \in \mathcal{H}$, $\langle STu, v \rangle = \langle Tu, S^* v \rangle = \langle u, T^* S^* v \rangle$; uniqueness gives $(ST)^* = T^* S^*$.

*Structural Commentary*: Move each operator across one inner product and invoke adjoint uniqueness.
*Key Insight*: Taking adjoints reverses composition and conjugates scalars. □

**Definition 5.2.36.** *Self-Adjoint, Nonnegative, and Coercive Operators*
Let $T \in \mathcal{L}(\mathcal{H})$.

(i) *Self-adjoint*: $T = T^*$.

(ii) *Nonnegative*: $T = T^*$ and $\forall h \in \mathcal{H}$, $\langle Th, h\rangle \geq 0$; write $T \geq 0$.
(iii) *Coercive*: if $\mathcal{H}$ is real, $\exists c > 0$ such that $\forall h \in \mathcal{H}$, $\langle Th, h\rangle \geq c\|h\|^2$.

**Proposition 5.2.37.** *Positive Order, Polarization, and Reducing Complements*
*Let $\mathcal{H}$ be a real Hilbert space.*

*(i) If $T \geq 0$ and $S \in \mathcal{L}(\mathcal{H})$, then $S^*TS \geq 0$.*
*(ii) If $T \in \mathcal{L}(\mathcal{H})$ is self-adjoint and $Q_T(u) := \langle Tu, u\rangle$, then $\langle Tu, v\rangle = \frac{1}{4}[Q_T(u+v) - Q_T(u-v)]$.*
*(iii) If $T \in \mathcal{L}(\mathcal{H})$ is self-adjoint and $Q_T \equiv 0$, then $T = 0$.*
*(iv) If $T \in \mathcal{L}(\mathcal{H})$ is self-adjoint and $V \leq \mathcal{H}$ satisfies $T(V) \subseteq V$, then $T(V^\perp) \subseteq V^\perp$.*

*Proof.*

(i) Since $T = T^*$, $(S^*TS)^* = S^*T^*S = S^*TS$. For $h \in \mathcal{H}$, $\langle S^*TSh, h\rangle = \langle TSh, Sh\rangle \geq 0$; hence $S^*TS \geq 0$.
(ii) For $u, v \in \mathcal{H}$, self-adjointness gives $Q_T(u \pm v) = Q_T(u) \pm 2\langle Tu, v\rangle + Q_T(v)$.
Subtracting gives $Q_T(u+v) - Q_T(u-v) = 4\langle Tu, v\rangle$.
(iii) By (ii), $\langle Tu, v\rangle = 0$ for all $u, v \in \mathcal{H}$; with $v = Tu$, $\|Tu\|^2 = 0$ for every $u$, so $T = 0$.
(iv) If $w \in V^\perp$ and $v \in V$, then $Tv \in V$ and $\langle Tw, v\rangle = \langle w, Tv\rangle = 0$; hence $Tw \in V^\perp$.

*Structural Commentary*: Adjoints transfer positivity and invariance; polarization recovers mixed terms.
*Key Insight*: For self-adjoint operators, the diagonal quadratic form controls the mixed inner products. □

**Theorem 5.2.38.** *Self-Adjoint Operator Norm from the Quadratic Form*
*If $\mathcal{H}$ is real and $T \in \mathcal{L}(\mathcal{H})$ is self-adjoint, then $\|T\|_{op} = \sup_{\|u\|=1} |\langle Tu, u\rangle|$.*

*Proof.* Let $\eta := \sup_{\|u\|=1} |\langle Tu, u\rangle|$. By Cauchy–Schwarz, $\eta \leq \|T\|_{\text{op}}$.
If $\eta = 0$, then $Q_T \equiv 0$ by scaling, so $T = 0$ by Proposition 5.2.37(iii).
Assume $\eta > 0$ and set $A_\pm := \eta I \pm T$. For $u \in \mathcal{H}$, $|\langle Tu, u\rangle| \leq \eta\|u\|^2$, hence $\langle A_\pm u, u\rangle \geq 0$ and $A_\pm \geq 0$.
By Proposition 5.2.37(i), $A_+^* A_- A_+ \geq 0$ and $A_-^* A_+ A_- \geq 0$.
Since $A_\pm = A_\pm^*$ and $A_+A_- = A_-A_+ = \eta^2 I - T^2$, their sum is $2\eta(\eta^2 I - T^2) \geq 0$, so $\eta^2 I - T^2 \geq 0$.
Thus $\|Tu\|^2 = \langle T^2 u, u\rangle \leq \eta^2\|u\|^2$ for every $u$, hence $\|T\|_{\text{op}} \leq \eta$.
*Structural Commentary*: Self-adjointness lets order information about $\eta I \pm T$ control $T^2$.
*Key Insight*: A bounded self-adjoint operator is normed by its one-vector quadratic tests. □

**Proposition 5.2.39.** *Closed Range and the Adjoint Kernel*
*If $T \in \mathcal{L}(\mathcal{H})$ has closed range, then $\mathcal{H} = \text{range}(T) \odot \ker(T^*)$.*

*Proof.* For $z \in \mathcal{H}$, $z \in \ker(T^*) \iff \forall u \in \mathcal{H},\ \langle u, T^*z\rangle = 0 \iff \forall u \in \mathcal{H},\ \langle Tu, z\rangle = 0 \iff z \in \text{range}(T)^\perp$.
Thus $\ker(T^*) = \text{range}(T)^\perp$, and Corollary 5.2.10 gives $\mathcal{H} = \text{range}(T) \odot \ker(T^*)$ since $\text{range}(T)$ is closed.
*Structural Commentary*: Identify the orthogonal complement of the range by moving $T$ across the inner product.
*Key Insight*: The directions missed by $T$ are exactly the vectors annihilated by $T^*$. □

**Theorem 5.2.40.** *Coercive Operators Are Invertible*
*Let $\mathcal{H}$ be real and let $T \in \mathcal{L}(\mathcal{H})$ satisfy $\exists c > 0$ such that $\forall h \in \mathcal{H}$, $\langle Th, h\rangle \geq c\|h\|^2$.*

*(i) $T$ is bijective.*
*(ii) $T^{-1} \in \mathcal{L}(\mathcal{H})$ and $\|T^{-1}\|_{op} \leq c^{-1}$.*

*Proof.* For $u \neq \mathbf{0}$, $c\|u\|^2 \leq \langle Tu, u\rangle \leq \|Tu\|\,\|u\|$, hence $\|Tu\| \geq c\|u\|$; the same holds for $u = \mathbf{0}$, so $\ker T = \{\mathbf{0}\}$.
If $Tu_n \to y$, then $\|u_n - u_m\| \leq c^{-1}\|Tu_n - Tu_m\|$; hence $u_n \to u$ and $Tu = y$, so $\text{range}(T)$ is closed.
Also $\langle T^*h, h\rangle = \langle Th, h\rangle \geq c\|h\|^2$, so $\ker(T^*) = \{\mathbf{0}\}$.

(i) By Proposition 5.2.39, $\mathcal{H} = \text{range}(T) \odot \ker(T^*) = \text{range}(T)$; with $\ker T = \{\mathbf{0}\}$, $T$ is bijective.
(ii) For $y \in \mathcal{H}$, put $u = T^{-1}y$. Then $c\|T^{-1}y\| = c\|u\| \leq \|Tu\| = \|y\|$, so $T^{-1} \in \mathcal{L}(\mathcal{H})$ and $\|T^{-1}\|_{\text{op}} \leq c^{-1}$.

*Structural Commentary*: Coercivity closes the range and kills both kernels; adjoint splitting gives surjectivity.
*Key Insight*: Coercivity supplies both a lower Lipschitz bound and, through the adjoint, surjectivity. □

**Theorem 5.2.41.** *Lax–Milgram Lemma*
*Let $\mathcal{H}$ be a real Hilbert space and let $B : \mathcal{H} \times \mathcal{H} \to \mathbb{R}$ be bilinear.*
*Suppose there are constants $M, c > 0$ such that $|B(u, v)| \leq M\|u\|\,\|v\|$ and $B(h, h) \geq c\|h\|^2$ $\forall u, v, h \in \mathcal{H}$.*
*Then $\forall \psi \in \mathcal{H}^*$ $\exists! h \in \mathcal{H}$ such that $\psi(u) = B(h, u)$ $(u \in \mathcal{H})$.*

*Proof.* Let $R : \mathcal{H} \to \mathcal{H}^*$ be the real Riesz isometry, $Ry(u) = \langle u, y\rangle$, and define $S : \mathcal{H} \to \mathcal{H}^*$ by $Sh(u) = B(h, u)$.
For $h, u \in \mathcal{H}$, $|Sh(u)| \le M \|h\| \|u\|$, hence $S \in \mathcal{L}(\mathcal{H}, \mathcal{H}^*)$.
Set $A := R^{-1}S \in \mathcal{L}(\mathcal{H})$. For $h \in \mathcal{H}$, $\langle Ah, h\rangle = \langle h, Ah\rangle = (RAh)(h) = Sh(h) = B(h, h) \ge c\|h\|^2$.
By Theorem 5.2.40, $A$ is bijective, hence $S = RA$ is bijective.
For $\psi \in \mathcal{H}^*$, set $h := A^{-1}R^{-1}\psi$; then $\forall u \in \mathcal{H}$, $\psi(u) = (RAh)(u) = Sh(u) = B(h, u)$.
If $h_1, h_2$ solve this identity, then $S(h_1 - h_2) = 0$, so $h_1 = h_2$.
*Structural Commentary*: A coercive bilinear form is a Riesz-represented coercive operator in disguise.
*Key Insight*: Solving $B(h, \cdot) = \psi$ is solving an invertible operator equation after Riesz representation. □

**Theorem 5.2.42.** *Orthonormal Systems in $L^2$*
*Let $(\phi_n)_{n\ge1} \subseteq L^2(\mu)$ be orthonormal, $\langle \phi_n, \phi_m\rangle = \delta_{nm}$, and $c_n(f) := \langle f, \phi_n\rangle = \int_X f\overline{\phi_n}\, d\mu$.*

*(i)* $\forall f \in L^2(\mu)$, $\sum_{n=1}^{\infty} |c_n(f)|^2 \le \|f\|_2^2$.

*(ii) If $(c_n) \in \ell^2$, then $s_N := \sum_{k=1}^{N} c_k\phi_k$ converges in $L^2(\mu)$.*

*(iii) If $(\phi_n)$ is complete, then $\forall f \in L^2(\mu)$, $\|f\|_2^2 = \sum_{n=1}^{\infty} |c_n(f)|^2$.*

*Proof.*

(i) Set $s_N(f) := \sum_{k=1}^{N} c_k(f)\phi_k$. Finite orthonormality gives $\|f - s_N(f)\|_2^2 = \|f\|_2^2 - \sum_{k=1}^{N} |c_k(f)|^2 \ge 0$.
Monotone convergence of the partial sums yields $\sum_{n=1}^{\infty} |c_n(f)|^2 \le \|f\|_2^2$.

(ii) If $m < N$, then $\|s_N - s_m\|_2^2 = \sum_{k=m+1}^{N} |c_k|^2$. Since $(c_n) \in \ell^2$, $(s_N)$ is Cauchy.
By completeness, $\exists g \in L^2(\mu)$ with $s_N \to g$.

(iii) Assume $(\phi_n)$ is complete. By (i), $(c_n(f)) \in \ell^2$; by (ii), $s_N(f) \to g$.
For fixed $j$, $\langle s_N(f), \phi_j\rangle = c_j(f)$ for $N \ge j$, hence $\langle f - g, \phi_j\rangle = c_j(f) - \lim_N \langle s_N(f), \phi_j\rangle = 0$.
Thus $f - g \perp \{\phi_j : j \ge 1\}$, so $f = g$; therefore $\|f\|_2^2 = \lim_N \|s_N(f)\|_2^2 = \sum_{n=1}^{\infty} |c_n(f)|^2$.

*Structural Commentary*: Complete orthonormal systems coordinatize $L^2$ by square-summable sequences.
*Key Insight*: Parseval is Pythagoras after passing from finite coordinates to an $L^2$ limit. □

**Method 5.2.43.** *Hilbert Space Proof Patterns*

(i) *Complete*: Cauchy tails, completions, and convergent subsequences force limits.
(ii) *Parallelogram*: test Hilbertian norms and make distance-minimizing sequences Cauchy.
(iii) *Project*: for closed $M$, write $x = p + r$, $p \in M$, $r \in M^\perp$.
(iv) *Separate*: density and equality via $M^\perp = \{\mathbf{0}\}$ and $M^{\perp\perp} = \overline{M}$.
(v) *Sum*: arbitrary convergence is net convergence over $\mathcal{F}(K)$; control finite tails.
(vi) *Expand*: orthonormal finite identities + Bessel bounds + completeness ⇒ Fourier expansions.
(vii) *Classify*: Hilbert bases give hdim, separability, and $\mathcal{H} \cong \ell^2(K)$.
(viii) *Dualize*: Riesz identifies $\mathcal{H}^*$ with $\mathcal{H}$, producing adjoints and coercive operator equations.

*Key Insight*: Hilbert-space arguments are finite orthogonal geometry plus completeness.

## 5.3 Core Functional Analytic Theorems

The core theorems explain why complete normed spaces behave rigidly under linear maps. Hahn–Banach supplies norm-preserving extensions, while Baire category turns pointwise control into uniform boundedness.

**Theorem 5.3.1.** *Hahn–Banach Extension Theorem*
*Let $\mathcal{X}$ be a real normed space, $M \le \mathcal{X}$, $p : \mathcal{X} \to \mathbb{R}$ sublinear, and $\varphi : M \to \mathbb{R}$ linear with $\varphi \le p$ on $M$.*

*(i) If $x_0 \in \mathcal{X} \setminus M$ and $M_0 := M + \mathbb{R}x_0$, then $\exists \varphi_0 : M_0 \to \mathbb{R}$ linear such that $\varphi_0|_M = \varphi$ and $\varphi_0 \le p$ on $M_0$.*
*(ii) $\exists \Phi : \mathcal{X} \to \mathbb{R}$ linear such that $\Phi|_M = \varphi$ and $\Phi \le p$ on $\mathcal{X}$.*

*Proof.*

(i) For $y, z \in M$, $\varphi(y)+\varphi(z) = \varphi(y+z) \le p(y+z) \le p(y-x_0)+p(z+x_0)$; $\varphi(y)-p(y-x_0) \le p(z+x_0)-\varphi(z)$.
Set $a := \sup_{y\in M}(\varphi(y) - p(y - x_0))$, $b := \inf_{z\in M}(p(z + x_0) - \varphi(z))$. Then $-p(-x_0) \le a \le b \le p(x_0)$; $\alpha \in [a, b]$.
Since $x_0 \notin M$, $y + tx_0$ has unique coordinates, so define $\varphi_0(y + tx_0) := \varphi(y) + t\alpha$.
If $t > 0$, then $\varphi_0(y + tx_0) = t(\varphi(y/t) + \alpha) \le tp(y/t + x_0) = p(y + tx_0)$.

If $t < 0$, then $\varphi_0(y + tx_0) = (-t)(\varphi(y/(-t)) - \alpha) \le (-t)p(y/(-t) - x_0) = p(y + tx_0)$.
If $t = 0$, domination is $\varphi \le p$ on $M$.

(ii) Let $\mathscr{E}$ be $(N, \psi)$, $M \le N \le \mathcal{X}$, $\psi : N \to \mathbb{R}$ linear, $\psi|_M = \varphi$, $\psi \le p$ on $N$, ordered by extension.
Chains have upper bound $(\bigcup N_i, \psi_*)$, where $\psi_*|_{N_i} = \psi_i$, so Principle 6.5.3 gives maximal $(N_*, \psi_*) \in \mathscr{E}$.
If $N_* \neq \mathcal{X}$, choose $x_0 \in \mathcal{X} \setminus N_*$; by (i), $\psi_*$ extends to $N_* + \mathbb{R}x_0$, contradiction. Thus $N_* = \mathcal{X}$ and $\Phi := \psi_*$.

*Structural Commentary*: Extend across one new direction while preserving domination, then use maximality.
*Key Insight*: Sublinearity leaves a nonempty interval of admissible values on each added direction. □

**Corollary 5.3.2.** *Norm-Preserving Extensions and Norming Functionals*
*Let $\mathcal{X}$ be a normed space and let $M \subseteq \mathcal{X}$ be a linear subspace.*

*(i) A bounded linear functional $\varphi : M \to \mathbb{F}$ admits an extension $\Phi \in \mathcal{X}^*$ with $\|\Phi\|_{op} = \|\varphi\|_{op}$.*
*(ii) $\forall 0 \neq x \in \mathcal{X}$ $\exists \Phi \in \mathcal{X}^*$ such that $\|\Phi\|_{op} = 1$ and $\Phi(x) = \|x\|$.*

*Proof.*

(i) If $\mathbb{F} = \mathbb{R}$, apply Theorem 5.3.1 with $p(u) := \|\varphi\|_{\mathrm{op}} \|u\|$ to get $\Phi|_M = \varphi$ and $\Phi \le p$.
Since $-\Phi(u) = \Phi(-u) \le p(u)$, $|\Phi(u)| \le \|\varphi\|_{\mathrm{op}} \|u\| \forall u \in \mathcal{X}$, so $\|\Phi\|_{\mathrm{op}} \le \|\varphi\|_{\mathrm{op}}$; restriction gives equality.
If $\mathbb{F} = \mathbb{C}$, apply the real case to $\operatorname{Re}\varphi$ and set $\Phi(u) := F(u) - iF(iu)$.
Then $\Phi|_M = \varphi$, $\Phi$ is complex-linear, and $|\Phi(u)| = \sup_{|\alpha|=1} F(\alpha u) \le \|\varphi\|_{\mathrm{op}} \|u\|$, so again $\|\Phi\|_{\mathrm{op}} = \|\varphi\|_{\mathrm{op}}$.

(ii) Fix $0 \neq x \in \mathcal{X}$, put $M_x := \mathbb{F}x$, and define $\varphi_x(\lambda x) := \lambda \|x\|$.
Then $\|\varphi_x\|_{\mathrm{op}} = 1$ and $\varphi_x(x) = \|x\|$; by (i), choose $\Phi \in \mathcal{X}^*$ with $\Phi|_{M_x} = \varphi_x$ and $\|\Phi\|_{\mathrm{op}} = 1$.

*Structural Commentary*: Preserve the norm, then extend a one-dimensional norming functional.
*Key Insight*: Hahn–Banach continues a bounded scalar test without weakening its norm bound. □

**Corollary 5.3.3.** *Natural Embedding Is an Isometry*
*For a normed space $\mathcal{X}$, the canonical map $J : \mathcal{X} \to \mathcal{X}^{**}$, $Jx(\Phi) = \Phi(x)$, satisfies $\|Jx\|_{op} = \|x\|$ $(x \in \mathcal{X})$.*

*Proof.* $\forall \Phi \in \mathcal{X}^*$, $|Jx(\Phi)| = |\Phi(x)| \le \|\Phi\|_{\mathrm{op}} \|x\|$, so $\|Jx\|_{\mathrm{op}} \le \|x\|$. If $x \neq 0$, choose $\Phi \in \mathcal{X}^*$ with $\|\Phi\|_{\mathrm{op}} = 1$ and $\Phi(x) = \|x\|$ by Corollary 5.3.2; hence $\|Jx\|_{\mathrm{op}} \ge \|x\|$. The case $x = 0$ is immediate.
*Structural Commentary*: The upper bound is duality; a norming functional attains the matching lower bound.
*Key Insight*: The double dual records every vector without changing its norm. □

**Example 5.3.4.** *A Dominated Extension of the Limit Functional*
On real $\ell^\infty$, set $p((x_n)) := \limsup_{n\to\infty} x_n$. This functional is subadditive and positively homogeneous.
If $c \subseteq \ell^\infty$ is the subspace of convergent sequences and $L((x_n)) := \lim_{n\to\infty} x_n$, then $L \le p$ on $c$.
By Theorem 5.3.1, $L$ extends to a linear functional $\Lambda : \ell^\infty \to \mathbb{R}$ with $\Lambda(x) \le \limsup_{n\to\infty} x_n$ $(x \in \ell^\infty)$.
This is the Hahn–Banach domination step behind limit-extension arguments on $\ell^\infty$.

**Corollary 5.3.5.** *Positive Riesz Representation on $C([a,b])$*
*Let $J = [a,b]$ and let $G \in C(J)^*$ be positive, i.e. $f \ge 0$ on $J$ implies $G(f) \ge 0$.*

*(i) $\exists \gamma : \mathcal{B}(\mathbb{R}) \to [0, +\infty]$ a finite Borel measure with $\operatorname{supp}\gamma \subseteq J$ and $\forall f \in C(J)$, $G(f) = \int_J f \, d\gamma$.*
*(ii) $\|G\|_{op} = G(\mathbf{1}_J) = \gamma(J)$.*

*Proof.*

(i) By Theorem 5.2.14, $\exists$ finite Radon $\mu$ on $J$ such that $G(f) = \int_J f \, d\mu$ for $f \in C(J)$.
Define $\gamma(E) := \mu(E \cap J)$ for $E \in \mathcal{B}(\mathbb{R})$. Then $\gamma(\mathbb{R} \setminus J) = 0$, so $\operatorname{supp}\gamma \subseteq J$, and $\int_J f \, d\gamma = \int_J f \, d\mu = G(f)$.

(ii) If $\|f\|_\infty \le 1$, then $\mathbf{1}_J \pm f \ge 0$, hence $-G(\mathbf{1}_J) \le G(f) \le G(\mathbf{1}_J)$ and $|G(f)| \le G(\mathbf{1}_J)$.
Thus $\|G\|_{\mathrm{op}} \le G(\mathbf{1}_J)$; taking $f = \mathbf{1}_J$ gives $\|G\|_{\mathrm{op}} \ge G(\mathbf{1}_J)$. Also $G(\mathbf{1}_J) = \int_J \mathbf{1}_J \, d\gamma = \gamma(J)$.

*Structural Commentary*: Specialize Riesz–Markov to the interval, then extend the measure by zero.
*Key Insight*: Positivity converts the order unit $\mathbf{1}_J$ into the exact dual norm. □

**Corollary 5.3.6.** *Signed Riesz Representation by a Charge*
*Let $J = [a,b]$, and suppose $G \in C(J)^*$ admits $G = G^+ - G^-$ with positive $G^\pm \in C(J)^*$.*
*Then $\exists \lambda : \mathcal{B}(\mathbb{R}) \to \mathbb{R}$ a charge with $\operatorname{supp}\lambda \subseteq J$ and $\forall f \in C(J)$, $G(f) = \int_J f \, d\lambda$.*

*Proof.* By Corollary 5.3.5, choose finite Borel measures $\gamma^\pm$ on $\mathbb{R}$ such that $\gamma^\pm(\mathbb{R} \setminus J) = 0$ and $G^\pm(f) = \int_J f \, d\gamma^\pm$ for $f \in C(J)$. Define $\lambda(E) := \gamma^+(E) - \gamma^-(E)$ for $E \in \mathcal{B}(\mathbb{R})$.
Then $\lambda$ is a charge, $\lambda(\mathbb{R} \setminus J) = 0$, and for $f \in C(J)$, $G(f) = G^+(f) - G^-(f) = \int_J f \, d\gamma^+ - \int_J f \, d\gamma^- = \int_J f \, d\lambda$.
*Structural Commentary*: Subtract the positive representing measures.
*Key Insight*: Signed representation is positive representation plus subtraction. □

**Theorem 5.3.7.** *Uniform Boundedness Principle*
*Let $\mathcal{X}$ be Banach, $\mathcal{Y}$ normed. If $\mathcal{F} \subseteq \mathcal{L}(\mathcal{X}, \mathcal{Y})$ pointwise bounded, $\forall x \in \mathcal{X}$, $\sup_{T\in\mathcal{F}} \|Tx\| < \infty$, then $\sup_{T\in\mathcal{F}} \|T\|_{op} < \infty$.*

*Proof.* For $T \in \mathcal{F}$, $f_T(x) := \|Tx\|$ is continuous. Pointwise boundedness and Theorem 4.3.11 give $x_0 \in \mathcal{X}$, $r > 0$, and $C < \infty$ with $\sup_{T\in\mathcal{F}} \|Tz\| \le C$ for $z \in B(x_0, r)$. Set $M_0 := \sup_{T\in\mathcal{F}} \|Tx_0\| < \infty$.
If $\|y\| < r$, then $\sup_{T\in\mathcal{F}} \|Ty\| \le \sup_{T\in\mathcal{F}} \|T(x_0 + y)\| + \sup_{T\in\mathcal{F}} \|Tx_0\| \le C + M_0$.
For $\|x\| = 1$ and $0 < t < r$, $t \sup_{T\in\mathcal{F}} \|Tx\| = \sup_{T\in\mathcal{F}} \|T(tx)\| \le C + M_0$. By $t \uparrow r$, $\sup_{T\in\mathcal{F}} \|T\|_{\text{op}} \le \frac{C+M_0}{r} < \infty$.
*Structural Commentary*: Baire controls one ball; translation and scaling transfer control to the unit ball.
*Key Insight*: Baire and linearity turn pointwise bounds into one uniform operator bound. □

**Corollary 5.3.8.** *Banach–Steinhaus Pointwise Limit Theorem*
*Let $\mathcal{X}$ be Banach, $\mathcal{Y}$ normed, $(T_n) \subseteq \mathcal{L}(\mathcal{X}, \mathcal{Y})$. Suppose $\forall x \in \mathcal{X}$, $T_n x$ converges in $\mathcal{Y}$, $Tx := \lim_{n\to\infty} T_n x$.*

*(i)* $\sup_n \|T_n\|_{op} < \infty$.
*(ii)* $T$ *is linear.*
*(iii)* $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ *and* $\|T\|_{op} \le \liminf_{n\to\infty} \|T_n\|_{op}$.

*Proof.*

(i) For each $x \in \mathcal{X}$, convergence of $(T_n x)$ gives $\sup_n \|T_n x\| < \infty$; hence Theorem 5.3.7 gives $\sup_n \|T_n\|_{\text{op}} < \infty$.

(ii) For $\alpha, \beta \in \mathbb{F}$ and $x, y \in \mathcal{X}$, $T(\alpha x + \beta y) = \lim_n T_n(\alpha x + \beta y) = \alpha Tx + \beta Ty$.

(iii) Let $L := \liminf_n \|T_n\|_{\text{op}} < \infty$. For $x \in \mathcal{X}$, $\|Tx\| = \lim_n \|T_n x\| \le L \|x\|$, so $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ and $\|T\|_{\text{op}} \le L$.

*Structural Commentary*: Uniform boundedness controls the family before taking its pointwise limit.
*Key Insight*: Completeness prevents the norms of pointwise-convergent operators from escaping. □

## 5.4 Open Mapping, Closed Graph, and Complemented Subspaces

The Baire-category principle has three operator-theoretic consequences that should be read together: closed range gives uniform preimage estimates, open maps are detected at the unit ball, and closed graphs force boundedness.

**Theorem 5.4.1.** *Closed Range Estimate*
*Let $\mathcal{X}, \mathcal{Y}$ be Banach spaces, let $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$, and set $\mathcal{R} := T(\mathcal{X})$. The following are equivalent.*

*(i)* $\mathcal{R}$ *is closed in* $\mathcal{Y}$.
*(ii)* $\exists M > 0$ *such that* $\forall y \in \mathcal{R}\ \exists x \in \mathcal{X}$ *with* $Tx = y$ *and* $\|x\| \le M \|y\|$.

*Proof.*

*(ii)* $\Rightarrow$ *(i)* Let $y \in \overline{\mathcal{R}}$. Choose $y_n \in \mathcal{R}$ with $y_n \to y$ and $\|y_n - y_{n-1}\| \le 2^{-n}$ for $n \ge 2$.
Pick $x_1, u_n \in \mathcal{X}$ with $Tx_1 = y_1$, $Tu_n = y_n - y_{n-1}$, and $\|u_n\| \le M 2^{-n}$.
Since $\sum_{n\ge2} \|u_n\| < \infty$, $x := x_1 + \sum_{n\ge2} u_n$ exists and $Tx = \lim_N y_N = y$, so $y \in \mathcal{R}$.

*(i)* $\Rightarrow$ *(ii)* If $\mathcal{R} = \{\mathbf{0}\}$, take $M = 1$. Otherwise $\mathcal{R}$ is Banach and $\mathcal{R} = \bigcup_{n\ge1} n\overline{T(B_{\mathcal{X}})}^{\mathcal{R}}$.
By Baire and balanced convexity, $\exists c > 0$ with $\overline{B}_{\mathcal{R}} \subseteq c\overline{T(B_{\mathcal{X}})}^{\mathcal{R}}$.
Thus each $0 \ne r \in \mathcal{R}$ has $u \in \mathcal{X}$ with $\|u\| \le c\|r\|$ and $\|r - Tu\| \le \frac{1}{2}\|r\|$.
For $y \ne 0$, iterate from $r_0 = y$ to get $r_k = r_{k-1} - Tu_k$, $\|r_k\| \le 2^{-k}\|y\|$, and $\|u_k\| \le c2^{1-k}\|y\|$.
Then $x := \sum_{k\ge1} u_k$ satisfies $Tx = y$ and $\|x\| \le 2c\|y\|$. For $y = 0$, take $x = 0$.

*Structural Commentary*: Baire category turns surjectivity onto a closed range into uniform preimage control.
*Key Insight*: Closed range is equivalent to solving $Tx = y$ with a norm bound proportional to $\|y\|$. □

**Proposition 5.4.2.** *Relative Openness and Closed Range*
*Let $\mathcal{X}, \mathcal{Y}$ be Banach spaces, $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$.*
*Then $T : \mathcal{X} \to T(\mathcal{X})$ is open, for subspace topology on range, iff $T(\mathcal{X})$ closed in $\mathcal{Y}$.*

*Proof.* Let $\mathcal{R} := T(\mathcal{X})$.

($\Rightarrow$) If $T : \mathcal{X} \to \mathcal{R}$ is open, then $\exists r > 0$ with $B_{\mathcal{R}}(0, r) \subseteq T(B_{\mathcal{X}})$.
For $0 \ne y \in \mathcal{R}$, choose $u \in B_{\mathcal{X}}$ with $Tu = \frac{r}{2\|y\|}y$, and set $x := \frac{2\|y\|}{r}u$.
Then $Tx = y$ and $\|x\| \le \frac{2}{r}\|y\|$; for $y = 0$, take $x = 0$. By Theorem 5.4.1, $\mathcal{R}$ is closed.

($\Leftarrow$) If $\mathcal{R}$ is closed, then by Theorem 5.4.1, $\exists M > 0$ such that every $y \in \mathcal{R}$ has $x \in \mathcal{X}$ with $Tx = y$ and $\|x\| \leq M\|y\|$. Let $U \subseteq \mathcal{X}$ be open and $x_0 \in U$. Choose $\rho > 0$ with $x_0 + \rho B_{\mathcal{X}} \subseteq U$. If $z \in B_{\mathcal{R}}(Tx_0, \frac{\rho}{M})$, choose $h \in \mathcal{X}$ with $Th = z - Tx_0$ and $\|h\| < \rho$; then $z = T(x_0 + h) \in T(U)$. Thus $T(U)$ is open in $\mathcal{R}$.

*Structural Commentary*: For linear maps, openness is detected at the unit ball around the origin.
*Key Insight*: Relative openness is the topological form of uniform preimage control. □

**Theorem 5.4.3.** *Open Mapping Theorem*
*Every surjective $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ between Banach spaces is open.*

*Proof.* Since $T$ is surjective, $T(\mathcal{X}) = \mathcal{Y}$, $\mathcal{Y}$ is closed in itself. By Proposition 5.4.2, $T : \mathcal{X} \to T(\mathcal{X}) = \mathcal{Y}$ open.
*Structural Commentary*: Surjectivity identifies the relative range with the codomain.
*Key Insight*: A Banach-space surjection has uniformly controlled preimages near the origin. □

**Corollary 5.4.4.** *Bounded Inverse Theorem*
*If $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ is bijective between Banach spaces, then $T^{-1} \in \mathcal{L}(\mathcal{Y}, \mathcal{X})$.*

*Proof.* Set $U := T^{-1} : \mathcal{Y} \to \mathcal{X}$. By Theorem 5.4.3, $T$ is open: if $O \subseteq \mathcal{X}$ is open, then $U^{-1}(O) = T(O)$ is open in $\mathcal{Y}$. Thus $U$ is continuous; since $U$ is linear, Theorem 5.1.5 gives $U \in \mathcal{L}(\mathcal{Y}, \mathcal{X})$.
*Structural Commentary*: Openness of the forward bijection is continuity of the inverse.
*Key Insight*: Banach completeness prevents a bounded linear bijection from having an unbounded inverse. □

**Corollary 5.4.5.** *One-Sided Control of Banach Norms*
*Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be norms on $\mathcal{X}$ such that $(\mathcal{X}, \|\cdot\|_1)$ and $(\mathcal{X}, \|\cdot\|_2)$ are Banach.*
*Assume $\exists c > 0$ such that $\|x\|_2 \leq c\|x\|_1$ for all $x \in \mathcal{X}$.*

*(i) $\exists C > 0$ such that $\|x\|_1 \leq C\|x\|_2$ for all $x \in \mathcal{X}$.*
*(ii) $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent.*

*Proof.*

(i) Let $I : (\mathcal{X}, \|\cdot\|_1) \to (\mathcal{X}, \|\cdot\|_2)$ be the identity. Since $\|Ix\|_2 \leq c\|x\|_1$, $I \in \mathcal{L}$ and is bijective. By Corollary 5.4.4, $I^{-1} \in \mathcal{L}$, hence $\exists C > 0$ such that $\|x\|_1 = \left\|I^{-1}x\right\|_1 \leq C\|x\|_2$.
(ii) For all $x \in \mathcal{X}$, $\|x\|_2 \leq c\|x\|_1$ by hypothesis and $\|x\|_1 \leq C\|x\|_2$ by (i). Hence the identity maps between the two normed structures are Lipschitz.

*Structural Commentary*: Apply the bounded inverse theorem to the identity between the two normed structures.
*Key Insight*: Completeness upgrades one-sided norm control to two-sided equivalence. □

**Definition 5.4.6.** *Closed Operators and Closed Complements*
Let $\mathcal{X}, \mathcal{Y}$ be normed spaces.

(i) *Closed operator*: $T : \mathcal{X} \to \mathcal{Y}$ is linear and $\forall (x_n) \subseteq \mathcal{X}$, $x_n \to x$ and $Tx_n \to y$ imply $Tx = y$.
(ii) *Closed linear complement*: $W \leq \mathcal{X}$ satisfies $\mathcal{X} = V \oplus W$ and is closed.

**Theorem 5.4.7.** *Closed Graph Theorem*
*Let $\mathcal{X}, \mathcal{Y}$ be Banach spaces and let $T : \mathcal{X} \to \mathcal{Y}$ be linear. Set $\Gamma(T) := \{(x, Tx) : x \in \mathcal{X}\}$.*

*(i) $T$ is bounded iff $T$ is closed.*
*(ii) $T$ is closed iff $\Gamma(T)$ is closed in $\mathcal{X} \times \mathcal{Y}$.*

*Proof.* On $\mathcal{X} \times \mathcal{Y}$, use $\|(x, y)\| := \|x\| + \|y\|$.

*(i)* ($\Rightarrow$) If $T$ is bounded, $x_n \to x$ and $Tx_n \to y$ imply $Tx_n \to Tx$; hence $y = Tx$, so $T$ is closed.
*(i)* ($\Leftarrow$) Assume $T$ is closed and set $\|x\|_* := \|x\| + \|Tx\|$. If $(x_n)$ is $\|\cdot\|_*$-Cauchy, then $x_n \to x$ in $\mathcal{X}$ and $Tx_n \to y$ in $\mathcal{Y}$; by closedness, $y = Tx$, so $x_n \to x$ in $\|\cdot\|_*$. Thus $(\mathcal{X}, \|\cdot\|_*)$ is Banach.
Since $\|x\| \leq \|x\|_*$, by Corollary 5.4.5, $\exists C > 0$ such that $\|x\|_* \leq C\|x\|$; hence $\|Tx\| \leq C\|x\|$.
*(ii)* ($\Rightarrow$) If $T$ is closed and $(x_n, Tx_n) \to (x, y)$, then $x_n \to x$ and $Tx_n \to y$, so $y = Tx$ and $(x, y) \in \Gamma(T)$.
*(ii)* ($\Leftarrow$) If $\Gamma(T)$ is closed, $x_n \to x$, and $Tx_n \to y$, then $(x_n, Tx_n) \to (x, y) \in \Gamma(T)$; hence $y = Tx$.

*Structural Commentary*: The graph norm turns closedness of the graph into completeness.
*Key Insight*: Graph-norm completeness and domination force equivalence with the original norm. □

**Theorem 5.4.8.** *Closed Complements and Continuous Projections*
*Let $V$ be a closed subspace of a Banach space $\mathcal{X}$. The following are equivalent.*

*(i) $\exists W \leq \mathcal{X}$ closed such that $\mathcal{X} = V \oplus W$.*
*(ii) $\exists P \in \mathcal{L}(\mathcal{X})$ such that $P^2 = P$ and $P(\mathcal{X}) = V$.*

*Proof.*

(*i*) $\Rightarrow$ (*ii*) Let $\mathcal{X} = V \oplus W$ with $W$ closed, and define $P(v+w) := v$. Then $P^2 = P$ and $P(\mathcal{X}) = V$. If $x_n \to x$ and $Px_n \to v$, then $v \in V$ and $(I-P)x_n \to x - v \in W$; by uniqueness in $V \oplus W$, $Px = v$. Thus $P$ is closed, so $P \in \mathcal{L}(\mathcal{X})$ by Theorem 5.4.7.

(*ii*) $\Rightarrow$ (*i*) Let $P \in \mathcal{L}(\mathcal{X})$ with $P^2 = P$ and $P(\mathcal{X}) = V$. By Proposition 2.4.2, $\mathcal{X} = V \oplus \ker P$. Since $\ker P = P^{-1}(\{\mathbf{0}\})$ and $P$ is continuous, $\ker P$ is closed.

*Structural Commentary*: Closed complementary coordinates are exactly bounded projection coordinates.
*Key Insight*: The Closed Graph Theorem converts algebraic splitting into continuous splitting. □

**Theorem 5.4.9.** *Complemented Ranges Are Closed*
*Let $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ act between Banach spaces. If* $\operatorname{range}(T)$ *has a closed linear complement in $\mathcal{Y}$, then it is closed.*

*Proof.* Set $\mathcal{R} := T(\mathcal{X})$, $\mathcal{Y} = \mathcal{R} \oplus Y_0$, $Y_0$ closed. Put $Z := \mathcal{X} \times Y_0$, $\|(x, y_0)\|_Z := \|x\| + \|y_0\|$, and $S : Z \to \mathcal{Y}$, $S(x, y_0) := Tx + y_0$. Then $Z$ is Banach, $S \in \mathcal{L}(Z, \mathcal{Y})$, and $S(Z) = \mathcal{Y}$. By Theorem 5.4.1, $\exists M > 0$ such that $\forall y \in \mathcal{Y}\ \exists (x, y_0) \in Z$: $y = Tx + y_0$ and $\|x\| + \|y_0\| \le M \|y\|$. If $y \in \mathcal{R}$, by direct-sum uniqueness $y_0 = 0$; $\forall y \in \mathcal{R}\ \exists x \in \mathcal{X}$: $Tx = y$, $\|x\| \le M \|y\|$. By Theorem 5.4.1, $\mathcal{R}$ closed.
*Structural Commentary*: A closed complement transfers preimage control to the range summand.
*Key Insight*: Complementability supplies the missing uniform preimage estimate for closed range. □

**Corollary 5.4.10.** *Finite-Codimensional Operator Ranges Are Closed*
*Let $\mathcal{X}, \mathcal{Y}$ be Banach spaces and let $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$. If* $\operatorname{codim}_{\mathcal{Y}} T(\mathcal{X}) < \infty$, *then $T(\mathcal{X})$ is closed.*

*Proof.* Set $\mathcal{R} := T(\mathcal{X})$. Since $\operatorname{codim}_{\mathcal{Y}} \mathcal{R} < \infty$, choose $Y_0 \le \mathcal{Y}$ with $\mathcal{Y} = \mathcal{R} \oplus Y_0$ and $\dim Y_0 < \infty$. By Theorem 5.1.14, $Y_0$ is closed; by Theorem 5.4.9, $\mathcal{R}$ is closed.
*Structural Commentary*: Finite codimension produces a finite-dimensional closed complement.
*Key Insight*: Only finitely many missing codomain directions force the operator range to be closed. □

# 6 Weak Topologies, Compact Operators, and Spectral Theory

## 6.1 Weak Topologies and Duality

Weak topologies replace norm control by scalar tests. They are usually too weak to preserve distances, but strong enough to expose compactness phenomena that are invisible in norm topology.

**Definition 6.1.1.** *Dual Space, Weak Topology, and Weak-* Topology*
For a normed space $\mathcal{X}$ over $\mathbb{F}$, its continuous dual is $\mathcal{X}^* := \mathcal{L}(\mathcal{X}, \mathbb{F})$.

(i) The *weak topology* on $\mathcal{X}$ is the weakest topology making $\varphi : \mathcal{X} \to \mathbb{F}$ continuous $\forall \varphi \in \mathcal{X}^*$.
(ii) The *weak-* topology* on $\mathcal{X}^*$ is the weakest topology making the evaluation $\varphi \mapsto \varphi(x)$ continuous $\forall x \in \mathcal{X}$.
(iii) The *strong topology* is the norm topology.

For $m \in \mathbb{N}$, $\varepsilon > 0$, $\varphi_\bullet = (\varphi_1, \dots, \varphi_m) \in (\mathcal{X}^*)^m$, and $x_\bullet = (x_1, \dots, x_m) \in \mathcal{X}^m$:

(i) Weak: $N(x; \varphi_\bullet; \varepsilon) := \{y \in \mathcal{X} : |\varphi_j(y - x)| < \varepsilon\ \forall j\}$.
(ii) Weak-*: $N_*(\varphi; x_\bullet; \varepsilon) := \{\psi \in \mathcal{X}^* : |(\psi - \varphi)(x_j)| < \varepsilon\ \forall j\}$.




Figure 225: Topologies on $\mathcal{X}^*$: weak-* $\subseteq$ weak $\subseteq$ norm; reflexivity identifies the first two.

**Definition 6.1.2.** *Weak and Weak-* Convergence*

(i) $(x_n) \subseteq \mathcal{X}$ *converges weakly* to $x$, written $x_n \rightharpoonup x$, iff $\varphi(x_n) \to \varphi(x)\ \forall \varphi \in \mathcal{X}^*$.
(ii) $(\varphi_n) \subseteq \mathcal{X}^*$ *converges weak-** to $\varphi \in \mathcal{X}^*$ iff $\varphi_n(x) \to \varphi(x)\ \forall x \in \mathcal{X}$.

**Proposition 6.1.3.** *Strong Convergence Implies Weak Convergence*
*If $x_n \to x$ in norm in a normed space $\mathcal{X}$, then $x_n \rightharpoonup x$.*

*Proof.* $\forall \varphi \in \mathcal{X}^*$, $|\varphi(x_n) - \varphi(x)| = |\varphi(x_n - x)| \le \|\varphi\|_{\mathrm{op}} \|x_n - x\| \to 0$.
*Structural Commentary*: Apply an arbitrary bounded scalar test to the norm error.
*Key Insight*: Norm convergence controls every bounded linear test. □

**Proposition 6.1.4.** *Weakly Convergent Sequences Are Bounded*
*If $x_n \rightharpoonup x$ in a normed space $\mathcal{X}$, then:*

*(i)* $\sup_n \|x_n\| < +\infty$.

*(ii)* $\|x\| \le \liminf_{n\to\infty} \|x_n\|$.

*Proof.*

(i) $\forall \varphi \in \mathcal{X}^*$, the scalar sequence $(\varphi(x_n))$ is bounded. The space $\mathcal{X}^*$ is Banach by Theorem 5.1.6, so the Uniform Boundedness Principle, Theorem 5.3.7, applies to $Jx_n : \mathcal{X}^* \to \mathbb{F}$, $Jx_n(\varphi) := \varphi(x_n)$.
Since $\|Jx_n\|_{\text{op}} = \|x_n\|$ by Corollary 5.3.3, the sequence is norm bounded.

(ii) If $x \neq 0$, choose $\varphi \in \mathcal{X}^*$ with $\|\varphi\|_{\text{op}} = 1$ and $\varphi(x) = \|x\|$ by Corollary 5.3.2.
Then $\|x\| = \lim_{n\to\infty} |\varphi(x_n)| \le \liminf_{n\to\infty} \|x_n\|$. The case $x = 0$ is immediate.

*Structural Commentary*: Uniform boundedness proves (i); one norming functional proves (ii).
*Key Insight*: Scalar-test boundedness forces norm boundedness. □

**Definition 6.1.5.** *Reflexivity*
For a normed space $\mathcal{X}$, the canonical map $J : \mathcal{X} \to \mathcal{X}^{**}$ is defined by $Jx(\varphi) := \varphi(x)$.
A Banach space $\mathcal{X}$ is *reflexive* if $J(\mathcal{X}) = \mathcal{X}^{**}$.

**Theorem 6.1.6.** *Banach–Alaoglu Theorem*
*For a normed space $\mathcal{X}$, the closed unit ball $\overline{B}_{\mathcal{X}^*}$ is weak-* compact.*

*Proof.* Let $B := \overline{B}_{\mathcal{X}}$, $B^* := \overline{B}_{\mathcal{X}^*}$, and $D := \{a \in \mathbb{F} : |a| \le 1\}$. By Theorem 6.3.12, $D^B$ is compact.
The injection $R : B^* \hookrightarrow D^B$, $R\varphi = \varphi|_B$, identifies $\sigma(\mathcal{X}^*, \mathcal{X})|_{B^*}$ with pointwise convergence on $B$.
Let $f \in \overline{R(B^*)} \subseteq D^B$. If $x_j \in B$, $a_j \in \mathbb{F}$, and $\sum_j a_j x_j \in B$, then $f(\sum_j a_j x_j) = \sum_j a_j f(x_j)$ by passage from the identities for $R\varphi$. Define $\Phi(0) := 0$ and $\Phi(x) := \|x\| f(x/\|x\|)$ for $x \neq 0$.
Then $\Phi$ is linear, $|\Phi(x)| \le \|x\|$, and $R\Phi = f$; hence $R(B^*)$ is closed in $D^B$. Therefore $B^*$ is weak-* compact.
*Structural Commentary*: $B^* \hookrightarrow D^B$; closed linear constraints plus Tychonoff give compactness.
*Key Insight*: Weak-* topology records precisely the coordinates used in Tychonoff compactness. □

**Remark 6.1.7.** *Compactness and Sequences*
Weak-* compactness need not be weak-* sequential compactness. For $\mathcal{X} = \ell^\infty$, let $\varphi_n(x) := x_n$.
Then $\varphi_n \in \overline{B}_{\mathcal{X}^*}$, but for any subsequence $(n_k)$, choosing $x \in \ell^\infty$ with $x_{n_k} = (-1)^k$ makes $\varphi_{n_k}(x) = (-1)^k$, so no subsequence is weak-* convergent. Helly's theorem gives a sequential substitute when $\mathcal{X}$ is separable.

**Theorem 6.1.8.** *Helly Selection for Bounded Dual Sequences*
*Let $\mathcal{X}$ be a separable normed space. If $(T_n) \subseteq \mathcal{X}^*$ and $\sup_n \|T_n\|_{op} < +\infty$, then $\exists$ a subsequence $(T_{n_k})$ and $T \in \mathcal{X}^*$ such that $\forall x \in \mathcal{X}$, $T_{n_k}(x) \to T(x)$.*

*Proof.* Let $D = \{x_j : j \ge 1\} \subseteq \mathcal{X}$ be dense and set $M := \sup_n \|T_n\|_{\text{op}}$.
If $M = 0$, take $T = 0$; assume $M > 0$. By diagonal extraction, choose $(n_k)$ such that $T_{n_k} x_j$ converges $\forall j$.
For $x \in \mathcal{X}$ and $\varepsilon > 0$, choose $j$ with $\|x - x_j\| < \varepsilon/(4M)$ and then $k, \ell$ large with $|T_{n_k} x_j - T_{n_\ell} x_j| < \varepsilon/2$.
Then $|T_{n_k} x - T_{n_\ell} x| \le 2M \|x - x_j\| + |T_{n_k} x_j - T_{n_\ell} x_j| < \varepsilon$, so $(T_{n_k} x)$ is Cauchy. Define $Tx := \lim_k T_{n_k} x$.
Linearity passes to limits, and $|Tx| \le M\|x\|$; hence $T \in \mathcal{X}^*$ and $T_{n_k} x \to Tx$ $\forall x \in \mathcal{X}$.
*Structural Commentary*: Diagonalize on dense $D$; use $M = \sup_n \|T_n\|_{\text{op}}$ to extend $D$-limits to $\mathcal{X}$.
*Key Insight*: A countable dense set reduces pointwise convergence to diagonal convergence. □

**Corollary 6.1.9.** *Embedding into a Continuous Function Space*
*For every normed space $\mathcal{X}$, $\exists$ compact Hausdorff $K$ and a linear isometry $\Phi : \mathcal{X} \to C(K)$.*

*Proof.* Let $K := \overline{B}_{\mathcal{X}^*}$ with the weak-* topology. By Theorem 6.1.6, $K$ is compact; since evaluations separate points of $K$, $K$ is Hausdorff. Define $\Phi : \mathcal{X} \to C(K)$ by $\Phi(x)(\varphi) := \varphi(x)$. Each $\Phi(x)$ is weak-* continuous, $\Phi$ is linear, and by Corollary 5.3.3, $\|\Phi(x)\|_\infty = \|x\|$.
*Structural Commentary*: $K = \overline{B}^{w^*}_{\mathcal{X}^*}$ turns scalar tests into compact coordinates.
*Key Insight*: Evaluation turns each vector into a continuous function without changing its norm. □

**Proposition 6.1.10.** *Natural Embedding and Weak Topologies*
*For a normed space $\mathcal{X}$, the canonical map $J : \mathcal{X} \to \mathcal{X}^{**}$ is a homeomorphism $(\mathcal{X}, w) \to (J(\mathcal{X}), w^*|_{J(\mathcal{X})})$.*

*Proof.* By Corollary 5.3.3, $J$ is injective. Fix $x_0 \in \mathcal{X}$, $\varphi_\bullet = (\varphi_1, \dots, \varphi_m) \in (\mathcal{X}^*)^m$, and $\varepsilon > 0$.
Since $(Jx - Jx_0)(\varphi_j) = \varphi_j(x - x_0)$, $J^{-1}(N_*(Jx_0; \varphi_\bullet; \varepsilon) \cap J(\mathcal{X})) = N(x_0; \varphi_\bullet; \varepsilon)$.
Thus $J$ and $J^{-1} : J(\mathcal{X}) \to \mathcal{X}$ are continuous.
*Structural Commentary*: Pullbacks of weak-* tests are weak tests.
*Key Insight*: $J$ preserves precisely the scalar tests that define the weak topology. □

**Theorem 6.1.11.** *Kakutani's Weak Compactness Criterion*
*For a Banach space $\mathcal{X}$, the following are equivalent.*

*(i)* $\mathcal{X}$ *is reflexive.*
*(ii) The closed unit ball* $\overline{B}_{\mathcal{X}}$ *is weakly compact.*

*Proof.*

$(i) \Rightarrow (ii)$ If $J(\mathcal{X}) = \mathcal{X}^{**}$, then by Corollary 5.3.3, $J(\overline{B}_{\mathcal{X}}) = \overline{B}_{\mathcal{X}^{**}}$.
By Theorem 6.1.6 applied to $\mathcal{X}^*$, $\overline{B}_{\mathcal{X}^{**}}$ is weak-* compact. By Proposition 6.1.10, $\overline{B}_{\mathcal{X}}$ is weakly compact.
$(ii) \Rightarrow (i)$ By Proposition 6.1.10, $K := J(\overline{B}_{\mathcal{X}})$ is weak-* compact, hence weak-* closed, in $\mathcal{X}^{**}$.
If $F \in \overline{B}_{\mathcal{X}^{**}} \setminus K$, weak-* separation yields $\varphi \in \mathcal{X}^*$ with $\operatorname{Re} F(\varphi) > \sup_{\|x\|\le 1} \operatorname{Re} Jx(\varphi) = \|\varphi\|_{\mathrm{op}}$.
But $\operatorname{Re} F(\varphi) \le |F(\varphi)| \le \|F\|_{\mathrm{op}} \|\varphi\|_{\mathrm{op}} \le \|\varphi\|_{\mathrm{op}}$, a contradiction. Thus $\overline{B}_{\mathcal{X}^{**}} \subseteq K$.
For $F \in \mathcal{X}^{**}$ with $F \neq 0$, $F/\|F\|_{\mathrm{op}} \in K$, so $F \in J(\mathcal{X})$; hence $J(\mathcal{X}) = \mathcal{X}^{**}$.

*Structural Commentary*: Reflexivity identifies $J(\overline{B}_{\mathcal{X}})$ with $\overline{B}_{\mathcal{X}^{**}}$; weak-* separation forces the converse.
*Key Insight*: Reflexivity identifies the weak-* compact bidual ball with the embedded original ball. □

**Corollary 6.1.12.** *Closed Bounded Convex Sets in Reflexive Spaces*
*Let $\mathcal{X}$ be reflexive Banach. If $C \subseteq \mathcal{X}$ is norm-closed, bounded, and convex, then $C$ is weakly compact.*

*Proof.* If $C = \emptyset$, the claim is trivial. Choose $r > 0$ with $C \subseteq r\overline{B}_{\mathcal{X}}$. By Theorem 6.1.11, $r\overline{B}_{\mathcal{X}}$ is weakly compact. For $x_0 \notin C$, by Hahn–Banach separation, $\varphi \in \mathcal{X}^*$ and $\alpha \in \mathbb{R}$ with $\operatorname{Re}\varphi(x_0) > \alpha \ge \sup_{c\in C} \operatorname{Re}\varphi(c)$; hence $\{\operatorname{Re}\varphi > \alpha\}$ is weakly open, contains $x_0$, and misses $C$. Thus $C$ is weakly closed, so $C$ is weakly compact.
*Structural Commentary*: $C \subseteq r\overline{B}_{\mathcal{X}}$ and $C$ weakly closed inside a weakly compact ball.
*Key Insight*: Convex norm-closed sets remain closed when the topology is weakened to scalar tests. □

**Theorem 6.1.13.** *Goldstine's Theorem*
*For a normed space $\mathcal{X}$, the weak-* closure of $J(\overline{B}_{\mathcal{X}})$ in $\mathcal{X}^{**}$ is $\overline{B}_{\mathcal{X}^{**}}$.*

*Proof.* Set $C := \overline{J(\overline{B}_{\mathcal{X}})}^{\sigma(\mathcal{X}^{**},\mathcal{X}^*)}$. $J$ is isometric by Corollary 5.3.3, $\overline{B}_{\mathcal{X}^{**}} = \bigcap_{\varphi\in\overline{B}_{\mathcal{X}^*}} \{F \in \mathcal{X}^{**} : |F(\varphi)| \le 1\}$ is weak-* closed, hence $C \subseteq \overline{B}_{\mathcal{X}^{**}}$. Suppose $F \in \overline{B}_{\mathcal{X}^{**}} \setminus C$. By weak-* Hahn–Banach separation,
$\exists \varphi \in \mathcal{X}^*$ and $\alpha \in \mathbb{R}$ such that $\operatorname{Re} F(\varphi) > \alpha \ge \sup_{\|x\|\le 1} \operatorname{Re} Jx(\varphi) = \sup_{\|x\|\le 1} \operatorname{Re}\varphi(x) = \|\varphi\|_{\mathrm{op}}$.
But $\operatorname{Re} F(\varphi) \le |F(\varphi)| \le \|F\|_{\mathrm{op}} \|\varphi\|_{\mathrm{op}} \le \|\varphi\|_{\mathrm{op}}$, a contradiction. Hence $\overline{B}_{\mathcal{X}^{**}} \subseteq C$.
*Structural Commentary*: $J(\overline{B}_{\mathcal{X}}) \subseteq \overline{B}_{\mathcal{X}^{**}}$; weak-* separation excludes $\overline{B}_{\mathcal{X}^{**}} \setminus C$.
*Key Insight*: $J(\overline{B}_{\mathcal{X}})$ is weak-* dense in $\overline{B}_{\mathcal{X}^{**}}$ even when $J\mathcal{X} \neq \mathcal{X}^{**}$. □

**Theorem 6.1.14.** *Eberlein–Smulian Theorem for Unit Balls*
*Let $\mathcal{X}$ be Banach and set $B := \overline{B}_{\mathcal{X}}$. The following are equivalent:*

*(i) $B$ is compact in $(\mathcal{X}, w)$.*
*(ii) Every $(x_n) \subseteq B$ has a subsequence $x_{n_k} \rightharpoonup x$ for some $x \in B$.*

*Proof.* By Eberlein–Smulian, for bounded $A \subseteq \mathcal{X}$, $A$ is relatively weakly compact iff every sequence in $A$ has a weakly convergent subsequence with limit in $\overline{A}^{w}$. By Hahn–Banach separation, $B$ is weakly closed, so $\overline{B}^{w} = B$.

$(i) \Rightarrow (ii)$ Apply the criterion with $A = B$.
$(ii) \Rightarrow (i)$ The criterion makes $B$ relatively weakly compact; since $\overline{B}^{w} = B$, $B$ is weakly compact.

*Structural Commentary*: For $B = \overline{B}_{\mathcal{X}}$, weak compactness $\iff$ weak subsequential compactness.
*Key Insight*: Weak compactness of $B$ is sequence-detectable. □

**Corollary 6.1.15.** *Reflexive Unit-Ball Characterization*
*Let $\mathcal{X}$ be Banach, set $B := \overline{B}_{\mathcal{X}}$, and let $J : \mathcal{X} \to \mathcal{X}^{**}$ be canonical. The following are equivalent:*

*(i) $J(\mathcal{X}) = \mathcal{X}^{**}$.*
*(ii) $B$ is compact in $(\mathcal{X}, w)$.*
*(iii) Every $(x_n) \subseteq B$ has a subsequence $x_{n_k} \rightharpoonup x$ for some $x \in B$.*

*Proof.*

$(i) \Rightarrow (ii)$ $J(\mathcal{X}) = \mathcal{X}^{**} \Rightarrow B$ is compact in $(\mathcal{X}, w)$ by Theorem 6.1.11.
$(ii) \Rightarrow (iii)$ By Theorem 6.1.14, $B$ weakly compact implies every $(x_n) \subseteq B$ has $x_{n_k} \rightharpoonup x \in B$.
$(iii) \Rightarrow (i)$ By Theorem 6.1.14, (iii) implies $B$ weakly compact; by Theorem 6.1.11, $J(\mathcal{X}) = \mathcal{X}^{**}$.

*Structural Commentary*: $J(\mathcal{X}) = \mathcal{X}^{**} \iff B_w$ compact $\iff B_w$ subsequentially compact.
*Key Insight*: Reflexivity is unit-ball compactness. □




Figure 226: For Banach $\mathcal{X}$: $\mathcal{X}$ reflexive $\iff \overline{B}_{\mathcal{X}}$ weakly compact $\iff \overline{B}_{\mathcal{X}}$ weakly sequentially compact.

**Proposition 6.1.16.** *Hilbert Weak Convergence Tests*
*Let $\mathcal{H}$ be a Hilbert space, let $(h_n), (u_n) \subseteq \mathcal{H}$, and let $h, u \in \mathcal{H}$.*

*(i) $h_n \rightharpoonup h$ if and only if $\forall u \in \mathcal{H}$, $\langle h_n, u\rangle \to \langle h, u\rangle$ .*
*(ii) If $h_n \rightharpoonup h$ and $u_n \to u$ strongly, then $\langle h_n, u_n\rangle \to \langle h, u\rangle$ .*

*Proof.*

(i) ($\Rightarrow$) For $u \in \mathcal{H}$, $\phi_u(x) := \langle x, u\rangle \in \mathcal{H}^*$, so $\langle h_n, u\rangle = \phi_u(h_n) \to \phi_u(h) = \langle h, u\rangle$.
(i) ($\Leftarrow$) For $\phi \in \mathcal{H}^*$, $u_\phi \in \mathcal{H}$ with $\phi(x) = \langle x, u_\phi\rangle$ by Theorem 5.2.32; $\phi(h_n) = \langle h_n, u_\phi\rangle \to \langle h, u_\phi\rangle = \phi(h)$.
(ii) Let $C := \sup_n \|h_n\| < +\infty$ by Proposition 6.1.4(i).
Since $|\langle h_n, u_n\rangle - \langle h, u\rangle| \le C\|u_n - u\| + |\langle h_n - h, u\rangle|$, the limit is 0.

*Structural Commentary*: $\mathcal{H}^* = \{\langle \cdot, u\rangle : u \in \mathcal{H}\}$; boundedness controls $u_n - u$.
*Key Insight*: $h_n \rightharpoonup h \iff \langle h_n, u\rangle \to \langle h, u\rangle$ for all fixed $u \in \mathcal{H}$. □

**Theorem 6.1.17.** *Bounded Hilbert Sequences Have Weak Subsequences*
*Let $\mathcal{H}$ be Hilbert and let $(h_n) \subseteq \mathcal{H}$ satisfy $C := \sup_n \|h_n\| < +\infty$. Then $\exists n_k \uparrow \infty$ and $h \in \mathcal{H}$ such that $h_{n_k} \rightharpoonup h$.*

*Proof.* Set $M := \operatorname{cspan}\{h_n : n \ge 1\}$. Then $M$ is a separable Hilbert space.
If $\dim M < +\infty$, then $C\overline{B}_M$ is compact by Theorem 5.1.14(iii); choose $h_{n_k} \to h \in M$ strongly, hence weakly.
Assume $\dim M = \infty$. Choose a Hilbert basis $(e_j)_{j\ge1}$ of $M$ by Corollary 5.2.28.
Diagonalize so $\langle h_{n_k}, e_j\rangle \to a_j$ $\forall j$. For $m \ge 1$, $\sum_{j=1}^m |a_j|^2 = \lim_k \sum_{j=1}^m |\langle h_{n_k}, e_j\rangle|^2 \le C^2$ by Theorem 5.2.22(ii).
Hence $a \in \ell^2$ and $h := \sum_j a_j e_j \in M$ by Theorem 5.2.21(i). If $v \in \operatorname{span}\{e_j : j \ge 1\}$, then $\langle h_{n_k} - h, v\rangle \to 0$.
For $u \in M$, $\varepsilon > 0$, $\|u - v\| < \varepsilon$ for $v$; then $\limsup_k |\langle h_{n_k} - h, u\rangle| \le (C + \|h\|)\varepsilon$, so $\langle h_{n_k} - h, u\rangle \to 0$.
For $u \in \mathcal{H}$, write $u = P_M u + u_\perp$, $u_\perp \in M^\perp$. Since $h_{n_k} - h \in M$, $\langle h_{n_k} - h, u\rangle = \langle h_{n_k} - h, P_M u\rangle \to 0$.
Thus $h_{n_k} \rightharpoonup h$ by Proposition 6.1.16.
*Structural Commentary*: Work in $M = \operatorname{cspan}\{h_n\}$; diagonal limits on $(e_j)$ extend by $\overline{\operatorname{span}}\{e_j\} = M$ and $M^\perp$.
*Key Insight*: Bounded Hilbert sequences have countably coordinatized weak limits. □

**Theorem 6.1.18.** *Hilbert Spaces Are Reflexive*
*For every Hilbert space $\mathcal{H}$, the canonical embedding $J : \mathcal{H} \to \mathcal{H}^{**}$ is onto.*

*Proof.* Set $B := \overline{B}_{\mathcal{H}}$. If $(h_n) \subseteq B$, then by Theorem 6.1.17, $h_{n_k} \rightharpoonup h$ for some $h \in \mathcal{H}$.
By Proposition 6.1.4, $\|h\| \le \liminf_k \|h_{n_k}\| \le 1$, so $h \in B$.
Thus $B$ is weakly subsequentially compact, hence $J(\mathcal{H}) = \mathcal{H}^{**}$ by Corollary 6.1.15.
*Structural Commentary*: $B_w$ subsequentially compact $\Longrightarrow J(\mathcal{H}) = \mathcal{H}^{**}$.
*Key Insight*: Hilbert coordinates make $B_w$ sequence-compact. □

**Theorem 6.1.19.** *Banach–Saks Theorem in Hilbert Spaces*
*Let $\mathcal{H}$ be Hilbert and let $h_n \rightharpoonup h$. Then $\exists n_k \uparrow \infty$ such that $\frac{1}{k}\sum_{j=1}^k h_{n_j} \to h$ strongly.*

*Proof.* Replace $h_n$ by $h_n - h$; let $h = 0$. Put $C := \sup_n \|h_n\|^2 < +\infty$.
Let $n_1 = 1$, $n_1 < \cdots < n_k$, $s_k := \sum_{j=1}^k h_{n_j}$; $n_{k+1} > n_k$ with $|\langle s_k, h_{n_{k+1}}\rangle| \le 1$, since $\langle s_k, h_n\rangle \to 0$.
Then $\|s_{k+1}\|^2 = \|s_k\|^2 + 2\operatorname{Re}\langle s_k, h_{n_{k+1}}\rangle + \|h_{n_{k+1}}\|^2 \le \|s_k\|^2 + C + 2$, so $\|s_k\|^2 \le (C+2)k$.
Hence $\|\frac{1}{k}s_k\|^2 \le \frac{C+2}{k} \to 0$. For the original sequence, $\frac{1}{k}\sum_{j=1}^k h_{n_j} = h + \frac{1}{k}\sum_{j=1}^k (h_{n_j} - h) \to h$.
*Structural Commentary*: Choose $n_{k+1}$ so $|\langle s_k, h_{n_{k+1}}\rangle| \le 1$; then $\|s_k\|^2 = O(k)$.
*Key Insight*: Weak-null subsequences can have $o(k)$ Cesàro norm. □

**Theorem 6.1.20.** *Radon–Riesz Theorem in Hilbert Spaces*
*Let $\mathcal{H}$ be Hilbert and let $h_n \rightharpoonup h$. Then $h_n \to h$ in norm iff $\|h_n\| \to \|h\|$.*

*Proof.*

($\Rightarrow$) If $\|h_n - h\| \to 0$, then $|\|h_n\| - \|h\|| \le \|h_n - h\| \to 0$.
($\Leftarrow$) Since $h_n \rightharpoonup h$, $\langle h_n, h\rangle \to \|h\|^2$. If also $\|h_n\| \to \|h\|$, then $\|h_n - h\|^2 = \|h_n\|^2 - 2\operatorname{Re}\langle h_n, h\rangle + \|h\|^2 \to 0$.

*Structural Commentary*: Weak convergence fixes $\langle h_n, h\rangle$; norm convergence fixes $\|h_n\|^2$.
*Key Insight*: $\|h_n - h\|^2$ has no remaining term. □

**Theorem 6.1.21.** *Metrizing Weak Topologies on the Unit Ball*
*Let $\mathcal{X}$ be normed, $W \le \mathcal{X}^*$ separable and point-separating $B := \overline{B}_{\mathcal{X}}$. Choose $(\varphi_k)_{k\ge1}$ dense in $W \cap \overline{B}_{\mathcal{X}^*}$. Then $\rho(u,v) := \sum_{k\ge1} 2^{-k}\,|\varphi_k(u-v)|$ is a metric on $B$ inducing $\sigma(\mathcal{X}, W)|_B$.*

*Proof.* For $u, v \in B$, $|\varphi_k(u-v)| \le 2$, hence $\rho(u,v) < +\infty$; symmetry and the triangle inequality are termwise. If $\rho(u,v) = 0$, then $\varphi_k(u-v) = 0$ for all $k$; for $\psi \in W \setminus \{0\}$, density gives $\varphi_{k_j} \to \frac{\psi}{\|\psi\|}$, hence $\psi(u-v) = 0$. Since $W$ separates points, $u = v$, so $\rho$ is a metric. Fix $u \in B$. It remains to compare neighborhoods at $u$.

($\Rightarrow$) Given $r > 0$, choose $N$ with $\sum_{k>N} 2^{1-k} < \frac{r}{2}$ and $\delta > 0$ with $\delta \sum_{k\le N} 2^{-k} < \frac{r}{2}$.
Then $V := \{v \in B : |\varphi_k(v-u)| < \delta, 1 \le k \le N\}$ is a $\sigma(\mathcal{X}, W)|_B$-neighborhood of $u$.
By $v \in V$, $\rho(u,v) \le \delta \sum_{k\le N} 2^{-k} + \sum_{k>N} 2^{1-k} < r$, hence $V \subseteq B_\rho(u,r)$.

($\Leftarrow$) Let $U = \{v \in B : |\psi_j(v-u)| < \varepsilon, 1 \le j \le m\}$ with $\psi_j \in W$ and $\varepsilon > 0$. Put $K := \{j : \psi_j \ne 0\}$.
For $j \in K$, choose $k_j$ with $\left\| \frac{\psi_j}{\|\psi_j\|} - \varphi_{k_j} \right\| < \frac{\varepsilon}{4\|\psi_j\|}$, then choose $r > 0$ with $2^{k_j} r < \frac{\varepsilon}{2\|\psi_j\|}$ $\forall j \in K$.
If $\rho(u,v) < r$, then $\left|\varphi_{k_j}(v-u)\right| < 2^{k_j} r$, $\|v-u\| \le 2$,
so $|\psi_j(v-u)| \le \|\psi_j\| \left|\varphi_{k_j}(v-u)\right| + \|\psi_j\| \left\| \frac{\psi_j}{\|\psi_j\|} - \varphi_{k_j} \right\| \|v-u\| < \varepsilon$ for $j \in K$.
For $j \notin K$, $\psi_j = 0$. Thus $B_\rho(u,r) \subseteq U$.

Therefore $\rho$ induces $\sigma(\mathcal{X}, W)|_B$.
*Structural Commentary*: $\rho = \sum_{k\ge1} 2^{-k}\,|\varphi_k(\cdot)|$ metrizes the bounded test topology generated by $W$.
*Key Insight*: On $B$, dense countable tests determine all $W$-tests. □

**Corollary 6.1.22.** *Metrizability of Weak Unit Balls*
*Let $\mathcal{X}$ be normed, write $B_{\mathcal{Y}} := \overline{B}_{\mathcal{Y}}$ for normed $\mathcal{Y}$, and let $J : \mathcal{X} \to \mathcal{X}^{**}$ be canonical.*

*(i) If $\mathcal{X}^*$ is separable, then $(B_{\mathcal{X}}, \sigma(\mathcal{X}, \mathcal{X}^*))$ is metrizable.*
*(ii) If $\mathcal{X}$ is separable, then $(B_{\mathcal{X}^*}, \sigma(\mathcal{X}^*, J(\mathcal{X})))$ is metrizable.*
*(iii) If $\mathcal{X}$ is reflexive Banach, then $(B_{\mathcal{X}}, \sigma(\mathcal{X}, \mathcal{X}^*))$ is metrizable iff $\mathcal{X}$ is separable.*

*Proof.*

(i) $\mathcal{X}^*$ is separable and point-separating on $\mathcal{X}$: if $x \ne y$, then $\exists \varphi \in \mathcal{X}^*$ with $\varphi(x-y) \ne 0$.
Theorem 6.1.21 with $(\mathcal{Y}, W) = (\mathcal{X}, \mathcal{X}^*)$ metrizes $(B_{\mathcal{X}}, \sigma(\mathcal{X}, \mathcal{X}^*))$.

(ii) Since $J$ is an isometry, $\mathcal{X}$ separable implies $J(\mathcal{X})$ separable.
If $\varphi \ne \psi$ in $\mathcal{X}^*$, choose $x \in \mathcal{X}$ with $(\varphi - \psi)(x) \ne 0$, hence $Jx(\varphi - \psi) \ne 0$, $J(\mathcal{X})$ point-separating on $\mathcal{X}^*$.
Theorem 6.1.21 with $(\mathcal{Y}, W) = (\mathcal{X}^*, J(\mathcal{X}))$ metrizes $(B_{\mathcal{X}^*}, \sigma(\mathcal{X}^*, J(\mathcal{X})))$.

(iii) ($\Rightarrow$) Let $B := B_{\mathcal{X}}$, $B_w$ metrizable. By reflexivity, Theorem 6.1.11, $B_w$ is compact metrizable, separable.
Choose countable $D \subseteq B$ with $\overline{D}^{\,w} = B$. Set $K := \overline{\operatorname{co} D}^{\,\|\cdot\|}$. If $K \ne B$, choose $x \in B \setminus K$.
Hahn–Banach gives $\varphi \in \mathcal{X}^*$ with $\operatorname{Re}\varphi(x) > \sup_{y\in K} \operatorname{Re}\varphi(y)$.
Since $D \subseteq K$ and $x \in \overline{D}^{\,w}$, weak continuity gives $\operatorname{Re}\varphi(x) \le \sup_{y\in K} \operatorname{Re}\varphi(y)$, contradiction.
Thus $B = K = \overline{\operatorname{co}_{\mathbb{Q}} D}^{\,\|\cdot\|}$; $\operatorname{co}_{\mathbb{Q}} D$ is countable, so $B$ is norm separable, and $\mathcal{X} = \bigcup_{m\ge1} mB$ is separable.

(iii) ($\Leftarrow$) Assume $\mathcal{X}$ separable. By (ii), Theorem 6.1.6, $J(\mathcal{X}) = \mathcal{X}^{**}$, $B_{\mathcal{X}^*}$ compact metrizable in $\sigma(\mathcal{X}^*, \mathcal{X}^{**})$.
Pick countable dense $D \subseteq B_{\mathcal{X}^*}$. The ($\Rightarrow$) argument in $\mathcal{X}^*$ gives $B_{\mathcal{X}^*} = \overline{\operatorname{co}_{\mathbb{Q}} D}^{\,\|\cdot\|}$.
Since $\operatorname{co}_{\mathbb{Q}} D$ is countable, $B_{\mathcal{X}^*}$ is norm separable and $\mathcal{X}^* = \bigcup_{m\ge1} mB_{\mathcal{X}^*}$ is separable, so (i) applies.

*Structural Commentary*: $W$ separable metrizes $B_{\sigma(\mathcal{X},W)}$; compact metrizable $B_w$ yields $B = \overline{\operatorname{co}_{\mathbb{Q}} D}^{\,\|\cdot\|}$.
*Key Insight*: Reflexive $B_{\mathcal{X}}$ is weakly metrizable iff $\mathcal{X}$ is separable. □

**Theorem 6.1.23.** *Whole Weak Topologies Are Not Metrizable in Infinite Dimension*
*Let $\mathcal{X}$ be Banach with $\dim \mathcal{X} = +\infty$.*

*(i) $(\mathcal{X}, \sigma(\mathcal{X}, \mathcal{X}^*))$ is not metrizable.*
*(ii) $(\mathcal{X}^*, \sigma(\mathcal{X}^*, J(\mathcal{X})))$ is not metrizable.*

*Proof.*

(i) Suppose $\rho$ metrizes $\sigma(\mathcal{X}, \mathcal{X}^*)$. $\forall n$, choose $U_n \subseteq B_\rho(0, \frac{1}{n})$ with $U_n = \{x : |\varphi_{n,j}(x)| < \varepsilon_n,\ 1 \le j \le m_n\}$.
Since $\bigcap_{j=1}^{m_n} \ker \varphi_{n,j} \ne \{0\}$, choose $x_n$ in this kernel with $\|x_n\| = n$.
Then $x_n \in U_n$, so $\rho(x_n, 0) < \frac{1}{n}$ and $x_n \not\to 0$, contradicting Proposition 6.1.4(i).

(ii) Suppose $\rho_*$ metrizes $\sigma(\mathcal{X}^*, J(\mathcal{X}))$. $\forall n$, choose finite-dim $M_n \leq \mathcal{X}$ with $M_n^\perp := \{\varphi : \varphi|_{M_n} = 0\} \subseteq B_{\rho_*}(0, \frac{1}{n})$.
Choose $y_n \notin M_n$ and define $\psi_n$ on $M_n + \operatorname{span}\{y_n\}$ by $\psi_n|_{M_n} = 0$ and $\psi_n(y_n) = 1$.
Extend $\psi_n$ to $\mathcal{X}$ by Corollary 5.3.2(i), and rescale to $\varphi_n \in M_n^\perp$ with $\|\varphi_n\|_{\text{op}} = n$.
Then $\rho_*(\varphi_n, 0) < \frac{1}{n}$, so $\varphi_n \to 0$ weak-*. By Theorem 5.3.7, $(\|\varphi_n\|_{\text{op}})$ is bounded, a contradiction.

*Structural Commentary*: Finite tests leave nonzero kernels in $\mathcal{X}$ and nonzero annihilators in $\mathcal{X}^*$.
*Key Insight*: Metric convergence would force bounded norms; finite scalar tests cannot. □

**Method 6.1.24.** *Weak Compactness Proof Patterns*

(i) *Compactify*: replace $\|x\|$ by finite coordinates $(\varphi_1(x), \dots, \varphi_m(x))$.
(ii) *Embed*: use $J : \mathcal{X} \to \mathcal{X}^{**}$ and compare $\sigma(\mathcal{X}, \mathcal{X}^*)$ with $\sigma(\mathcal{X}^{**}, \mathcal{X}^*)|_{J(\mathcal{X})}$.
(iii) *Separate*: if $F \notin J(B)$, choose $\varphi \in \mathcal{X}^*$ separating $F$ from $J(B)$.
(iv) *Metrize on balls*: for separable $W \leq \mathcal{X}^*$, use $\rho(u, v) = \sum_{k \geq 1} 2^{-k} |\varphi_k(u - v)|$ on bounded $B$.
(v) *Hilbertize*: replace $\varphi$ by $\langle \cdot, h \rangle$ and use orthogonal coordinates/projections.

*Key Insight*: Weak compactness is compactness for scalar coordinates $\varphi(x)$ on bounded sets.

## 6.2 Compact Operators

Compact operators carry bounded sets to relatively compact sets. Their spectral and solvability theory therefore resembles matrix theory inside infinite-dimensional spaces.

**Definition 6.2.1.** *Compact and Finite-Rank Operators*
Let $\mathcal{X}, \mathcal{Y}$ be normed spaces and let $T : \mathcal{X} \to \mathcal{Y}$ be linear.

(i) *Compact*: $\overline{T(B)}$ is compact in $\mathcal{Y}$ $\forall B \subseteq \mathcal{X}$ bounded.
(ii) *Finite rank*: $\dim \operatorname{range}(T) < \infty$.

**Proposition 6.2.2.** *Compactness Tests on Hilbert Spaces*
*Let $\mathcal{H}$ be Hilbert, let $K \in \mathcal{L}(\mathcal{H})$, and set $B := \overline{B}_{\mathcal{H}}$. The following are equivalent:*

*(i) $K$ is compact.*
*(ii) $K(B)$ is totally bounded.*
*(iii) Every bounded $(h_n) \subseteq \mathcal{H}$ has a subsequence $(h_{n_j})$ such that $Kh_{n_j}$ converges in norm.*

*Proof.* For bounded $E \subseteq \mathcal{H}$, choose $r > 0$ with $E \subseteq rB$; then $K(E) \subseteq K(rB) = rK(B)$.

(i) $\Rightarrow$ (ii) $K$ compact $\Rightarrow \overline{K(B)}$ compact $\Rightarrow \overline{K(B)}$ totally bounded $\Rightarrow K(B)$ totally bounded.
(ii) $\Rightarrow$ (iii) $(h_n)$ bounded $\Rightarrow \exists r > 0$, $\{h_n : n \geq 1\} \subseteq rB$. Then $(Kh_n) \subseteq K(rB) = rK(B)$ with $rK(B)$ totally bounded, so $\exists (n_j)$ such that $(Kh_{n_j})$ is Cauchy; $\mathcal{H}$ complete $\Rightarrow \exists y \in \mathcal{H}$, $\left\|Kh_{n_j} - y\right\| \to 0$.
(iii) $\Rightarrow$ (i) Let $E \subseteq \mathcal{H}$ be bounded. If $K(E)$ is not totally bounded, then $\exists \varepsilon > 0$ and $(x_n) \subseteq E$ such that $n \neq m \Rightarrow \|Kx_n - Kx_m\| \geq \varepsilon$. By (iii), $\exists (n_j)$ and $y \in \mathcal{H}$ with $\left\|Kx_{n_j} - y\right\| \to 0$, contradiction. Hence $K(E)$ is totally bounded, so $\overline{K(E)}$ is totally bounded and complete; Theorem 4.1.30 gives $\overline{K(E)}$ compact.

*Structural Commentary*: $E \subseteq rB \Rightarrow K(E) \subseteq rK(B)$; $K(B)$ totally bounded $\Rightarrow (h_n) \subseteq rB \Rightarrow \exists (n_j) : (Kh_{n_j})$ Cauchy.
*Key Insight*: $K$ compact iff $K$ sends bounded sequences to norm-precompact sequences. □

**Proposition 6.2.3.** *Finite Rank Operators Are Compact*
*Let $\mathcal{X}, \mathcal{Y}$ be normed, $T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$, $R := \operatorname{range}(T)$. If $\dim R < +\infty$, then $T$ is compact.*

*Proof.* Let $B \subseteq \mathcal{X}$ be bounded. Then $T(B) \subseteq R$ is bounded, so $\overline{T(B)}^R$ is compact by Theorem 5.1.14.
Since $R \leq \mathcal{Y}$ is finite-dimensional, $R$ is closed in $\mathcal{Y}$; hence $\overline{T(B)}^{\mathcal{Y}} = \overline{T(B)}^R$ is compact. Thus $T$ is compact.
*Structural Commentary*: $T(B) \subseteq R$, $\dim R < +\infty$, and bounded subsets of $R$ have compact closure.
*Key Insight*: Finite-dimensional range makes bounded images relatively compact. □

**Theorem 6.2.4.** *Norm Limits of Compact Operators*
*Let $\mathcal{X}$ be normed, $\mathcal{Y}$ Banach, $T_n, T \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$. If each $T_n$ compact and $\|T_n - T\|_{op} \to 0$, then $T$ is compact.*

*Proof.* Let $B \subseteq \mathcal{X}$ be bounded, $M > 0$, $\|x\| \leq M$ for $x \in B$. For $\varepsilon > 0$, choose $N$ with $\|T - T_N\|_{\text{op}} < \frac{\varepsilon}{2M}$.
Since $\overline{T_N(B)}$ is compact, choose $y_1, \dots, y_r \in \mathcal{Y}$ with $T_N(B) \subseteq \bigcup_{j=1}^r B_{\mathcal{Y}}(y_j, \frac{\varepsilon}{2})$.
Then $T(B) \subseteq \bigcup_{j=1}^r B_{\mathcal{Y}}(y_j, \varepsilon)$, so $T(B)$ is totally bounded. Since $\mathcal{Y}$ is complete, $\overline{T(B)}$ is compact; $T$ is compact.
*Structural Commentary*: $\|T - T_N\|_{\text{op}}$ transfers finite nets from $T_N(B)$ to $T(B)$ for bounded $B$.
*Key Insight*: Norm limits preserve relative compactness of bounded images. □

**Proposition 6.2.5.** *Compact Operators Form an Operator Ideal*
*Let $\mathcal{W}, \mathcal{X}, \mathcal{Y}, \mathcal{Z}$ be normed spaces.*

*(i) If $K_1, \dots, K_m : \mathcal{X} \to \mathcal{Y}$ are compact, then $\sum_{j=1}^{m} K_j$ is compact.*

*(ii) If $K : \mathcal{X} \to \mathcal{Y}$ is compact, $A \in \mathcal{L}(\mathcal{Y}, \mathcal{Z})$, and $B \in \mathcal{L}(\mathcal{W}, \mathcal{X})$, then $AKB : \mathcal{W} \to \mathcal{Z}$ is compact.*

*Proof.*

(i) Let $E \subseteq \mathcal{X}$ be bounded, $C_j := \overline{K_j(E)}$. Then each $C_j$ is compact, and $\sum_{j=1}^{m} K_j(E) \subseteq \sigma(C_1 \times \cdots \times C_m)$ for the continuous map $\sigma(y_1, \dots, y_m) := \sum_{j=1}^{m} y_j$. Thus $\sum_{j=1}^{m} K_j(E)$ is relatively compact.

(ii) Let $E \subseteq \mathcal{W}$ be bounded. Then $B(E)$ is bounded, $C := \overline{K(B(E))}$ is compact, and $AKB(E) \subseteq A(C)$. Since $A(C)$ is compact and closed in $\mathcal{Z}$, $\overline{AKB(E)} \subseteq A(C)$ is compact.

*Structural Commentary*: Products of compact image closures control sums; bounded precomposition and continuous postcomposition control $AKB$.
*Key Insight*: Compact operators form a two-sided ideal in $\mathcal{L}$. □

**Example 6.2.6.** *Diagonal Compact Operators*
Let $(e_k)_{k\geq 1}$ be a Hilbert basis of separable $\mathcal{H}$ and let $\lambda = (\lambda_k) \in \ell^\infty$ with $\lambda_k \to 0$.
For $h \in \mathcal{H}$, set $Th := \sum_{k=1}^{\infty} \lambda_k \langle h, e_k \rangle e_k$ and $T_N h := \sum_{k=1}^{N} \lambda_k \langle h, e_k \rangle e_k$.
By Parseval: $\sum_k |\lambda_k \langle h, e_k \rangle|^2 \leq \|\lambda\|_\infty^2 \|h\|^2 \Rightarrow T \in \mathcal{L}(\mathcal{H})$.
For $N \geq 1$, $\operatorname{range} T_N \subseteq \operatorname{span}\{e_1, \dots, e_N\} \Rightarrow \operatorname{rank} T_N \leq N \Rightarrow T_N$ compact by Proposition 6.2.3.
Also $\|(T - T_N)h\|^2 = \sum_{k>N} |\lambda_k|^2 |\langle h, e_k \rangle|^2 \leq \sup_{k>N} |\lambda_k|^2 \|h\|^2 \Rightarrow \|T - T_N\|_{\text{op}} \leq \sup_{k>N} |\lambda_k| \to 0$.
Hence $T_N \to T$ in $\|\cdot\|_{\text{op}}$ and $T$ is compact by Theorem 6.2.4.

**Theorem 6.2.7.** *Weak-to-Strong Characterization*
*Let $\mathcal{H}$ be Hilbert, $K \in \mathcal{L}(\mathcal{H})$. Then $K$ is compact iff $h_n \rightharpoonup h$ implies $\|Kh_n - Kh\| \to 0$ $\forall (h_n) \subseteq \mathcal{H}$ and $h \in \mathcal{H}$.*

*Proof.*

($\Rightarrow$) Let $h_n \rightharpoonup h$. By Proposition 6.1.4(i), $(h_n)$ bounded.
For every subsequence $(n_j)$, By Proposition 6.2.2(iii), $\exists (j_\ell)$ and $y \in \mathcal{H}$ with $\left\| Kh_{n_{j_\ell}} - y \right\| \to 0$.
For $u \in \mathcal{H}$, $\langle y, u \rangle = \lim_\ell \left\langle Kh_{n_{j_\ell}}, u \right\rangle = \lim_\ell \left\langle h_{n_{j_\ell}}, K^* u \right\rangle = \langle h, K^* u \rangle = \langle Kh, u \rangle$, so $y = Kh$.
If $\|Kh_n - Kh\| \not\to 0$, then $\exists \varepsilon > 0$, $(n_j)$ with $\left\| Kh_{n_j} - Kh \right\| \geq \varepsilon$, contradicting $\exists (j_\ell) : \left\| Kh_{n_{j_\ell}} - Kh \right\| \to 0$.

($\Leftarrow$) Let $(h_n)$ be bounded. By Theorem 6.1.17, $\exists (n_j)$, $h \in \mathcal{H}$ with $h_{n_j} \rightharpoonup h$. By hypothesis, $\left\| Kh_{n_j} - Kh \right\| \to 0$.
Hence $\forall (h_n)$ bounded, $\exists (n_j)$ and $y \in \mathcal{H}$ with $\left\| Kh_{n_j} - y \right\| \to 0$; By Proposition 6.2.2(iii), $K$ compact.

*Structural Commentary*: $K$ compact, $h_n \rightharpoonup h \Rightarrow Kh_{n_{j_\ell}} \to y = Kh$; $(h_n)$ bounded $\Rightarrow h_{n_j} \rightharpoonup h \Rightarrow Kh_{n_j} \to Kh$.
*Key Insight*: $K$ compact iff $h_n \rightharpoonup h$ forces $Kh_n \to Kh$ in norm. □

**Definition 6.2.8.** *Orthogonal Projection Sequence*
Let $(e_k)_{k\geq 1}$ be a Hilbert basis for a separable Hilbert space $\mathcal{H}$.
The induced *orthogonal projection sequence* $(P_n)_{n\geq 1}$ is defined by:

(i) $P_n h := \sum_{k=1}^{n} \langle h, e_k \rangle e_k$ for $h \in \mathcal{H}$.
(ii) $\operatorname{range}(P_n) = \operatorname{span}\{e_1, \dots, e_n\}$ and $\|P_n\|_{\text{op}} = 1$.
(iii) $P_n h \to h$ for every $h \in \mathcal{H}$.

**Proposition 6.2.9.** *Finite-Rank Approximation by Orthogonal Projections*
*Let $(P_n)$ be the orthogonal projection sequence induced by a Hilbert basis of a separable Hilbert space $\mathcal{H}$. For $T \in \mathcal{L}(\mathcal{H})$, $T$ is compact iff $\|P_n T - T\|_{op} \to 0$.*

*Proof.*

($\Rightarrow$) $B := \overline{B}_{\mathcal{H}}$, $C := \overline{T(B)}$; $T$ compact $\Rightarrow C$ compact.
$\varepsilon > 0 \Rightarrow \exists y_1, \dots, y_r \in C, N \in \mathbb{N} : C \subseteq \bigcup_{j=1}^{r} B_{\mathcal{H}}(y_j, \frac{\varepsilon}{3})$ and $n \geq N \Rightarrow \max_{1\leq j\leq r} \|(P_n - I)y_j\| < \frac{\varepsilon}{3}$.
Thus $n \geq N$, $x \in B \Rightarrow \exists j : \|Tx - y_j\| < \frac{\varepsilon}{3}$ and $\|(P_n T - T)x\| \leq \|Tx - y_j\| + \|(P_n - I)y_j\| + \|y_j - Tx\| <$

$\varepsilon$.
Hence $\|P_nT - T\|_{\text{op}} = \sup_{x\in B} \|(P_nT - T)x\| \to 0$.

($\Leftarrow$) $\text{range}(P_nT) \subseteq \text{range}(P_n)$, $\dim \text{range}(P_n) < \infty \Rightarrow P_nT$ compact by Proposition 6.2.3.
Also $P_nT \to T$ in $\|\cdot\|_{\text{op}}$; Theorem 6.2.4 $\Rightarrow T$ compact.

*Structural Commentary*: Compact $C = \overline{T(B)}$ upgrades $P_ny \to y$ to uniform convergence on $C$.
*Key Insight*: $T$ compact iff finite-rank projections approximate $T$ in operator norm. □

**Corollary 6.2.10.** *Finite-Rank Operators Are Dense in Compact Operators*
*Let $\mathcal{H}$ be separable, and let $\mathcal{K}(\mathcal{H})$ and $\mathcal{F}(\mathcal{H})$ denote the compact and finite-rank operators in $\mathcal{L}(\mathcal{H})$.*

*(i) $\mathcal{K}(\mathcal{H})$ is a norm-closed linear subspace of $\mathcal{L}(\mathcal{H})$.*
*(ii) $\overline{\mathcal{F}(\mathcal{H})}^{\|\cdot\|_{op}} = \mathcal{K}(\mathcal{H})$.*

*Proof.*

(i) Proposition 6.2.5: $K_1, K_2 \in \mathcal{K}(\mathcal{H})$, $\alpha, \beta \in \mathbb{F} \Rightarrow \alpha K_1 + \beta K_2 \in \mathcal{K}(\mathcal{H})$.
Theorem 6.2.4: $K_n \in \mathcal{K}(\mathcal{H})$, $\|K_n - K\|_{\text{op}} \to 0 \Rightarrow K \in \mathcal{K}(\mathcal{H})$.

*(ii)* ($\Rightarrow$) $\mathcal{F}(\mathcal{H}) \subseteq \mathcal{K}(\mathcal{H})$, (i) $\Rightarrow \overline{\mathcal{F}(\mathcal{H})}^{\|\cdot\|_{\text{op}}} \subseteq \mathcal{K}(\mathcal{H})$.

*(ii)* ($\Leftarrow$) $\dim \mathcal{H} < +\infty \Rightarrow \mathcal{F}(\mathcal{H}) = \mathcal{L}(\mathcal{H}) \supseteq \mathcal{K}(\mathcal{H})$. If $\dim \mathcal{H} = +\infty$,
take $(P_n)$; $T \in \mathcal{K}(\mathcal{H}) \Rightarrow P_nT \in \mathcal{F}(\mathcal{H})$, $\|P_nT - T\|_{\text{op}} \to 0 \Rightarrow T \in \overline{\mathcal{F}(\mathcal{H})}^{\|\cdot\|_{\text{op}}}$ by Proposition 6.2.9.

*Structural Commentary*: $\mathcal{K}(\mathcal{H})$ is norm closed, and $P_nT \in \mathcal{F}(\mathcal{H})$ with $P_nT \to T$ for compact $T$.
*Key Insight*: On separable $\mathcal{H}$, $\mathcal{K}(\mathcal{H}) = \overline{\mathcal{F}(\mathcal{H})}^{\|\cdot\|_{\text{op}}}$. □

**Theorem 6.2.11.** *Schauder's Theorem*
*Let $\mathcal{H}$ be Hilbert. If $K \in \mathcal{L}(\mathcal{H})$ is compact, then $K^*$ is compact.*

*Proof.* ByTheorem 5.2.34,$K^* \in \mathcal{L}(\mathcal{H})$. By Theorem 6.2.7, it suffices to prove $h_n \rightharpoonup h \Rightarrow \|K^*(h_n - h)\| \to 0$.
Put $u_n := h_n - h$. Then $u_n \rightharpoonup 0$, $M := \sup_n \|u_n\| < \infty$, and $\forall v \in \mathcal{H}$, $\langle K^*u_n, v\rangle = \langle u_n, Kv\rangle \to 0$, so $K^*u_n \rightharpoonup 0$.
Compactness of $K$ gives $\|KK^*u_n\| \to 0$, hence $0 \le \|K^*u_n\|^2 = \langle KK^*u_n, u_n\rangle \le M\|KK^*u_n\| \to 0$.
*Structural Commentary*: $u_n \rightharpoonup 0 \Rightarrow K^*u_n \rightharpoonup 0$, $K$ compact $\Rightarrow KK^*u_n \to 0$; $\|K^*u_n\|^2 \le M\|KK^*u_n\|$.
*Key Insight*: $\|K^*u_n\|^2 = \langle KK^*u_n, u_n\rangle$. □

**Proposition 6.2.12.** *Injective Compact Perturbations of the Identity Are Onto*
*Let $\mathcal{H}$ be Hilbert, let $K \in \mathcal{L}(\mathcal{H})$ be compact, and set $A := I + K$. If $\ker A = \{0\}$, then $A(\mathcal{H}) = \mathcal{H}$.*

*Proof.*

(i) If $\inf_{\|h\|=1} \|Ah\| = 0$, choose $\|h_n\| = 1$, $Ah_n \to 0$. By Proposition 6.2.2(iii), $Kh_{n_j} \to y$,
hence $h_{n_j} = Ah_{n_j} - Kh_{n_j} \to -y := h$, $\|h\| = 1$, and $Ah = 0$, contradiction.
Thus $\exists c > 0 : \|Ah\| \ge c\|h\|$ $\forall h \in \mathcal{H}$.

(ii) Fix $g \in \mathcal{H}$ and put $M := \text{cspan}\{K^m g : m \ge 0\}$. Then $g \in M$, $K(M) \subseteq M$, and $M$ is separable.
If $\dim M < +\infty$, then $A|_M : M \to M$ injective $\Rightarrow A|_M$ surjective $\Rightarrow g \in A(\mathcal{H})$.

(iii) Assume $\dim M = +\infty$. Let $(P_n)$ be projection sequence of Hilbert basis of $M$,
$H_n := P_n(M)$, $\dim H_n < +\infty$. By Proposition 6.2.9, $\|P_nK - K\|_{\text{op}} \to 0$. Let $N$,
$n \ge N \Rightarrow \|P_nK - K\|_{\text{op}} < \frac{c}{2}$. Then $n \ge N$, $h \in H_n \Rightarrow \|(I + P_nK)h\| \ge \|Ah\| - \|(P_nK - K)h\| \ge \frac{c}{2}\|h\|$,
so $I + P_nK : H_n \to H_n$ injective $\Rightarrow$ surjective.

(iv) Choose $u_n \in H_n$ with $(I + P_nK)u_n = P_ng$. Then $\|u_n\| \le \frac{2}{c}\|P_ng\| \le \frac{2}{c}\|g\|$.
By Theorem 6.1.17, $u_{n_j} \rightharpoonup u$ for some subsequence.
By Theorem 6.2.7, $Ku_{n_j} \to Ku$, $\|P_{n_j}Ku_{n_j} - Ku_{n_j}\| \le \|P_{n_j}K - K\|_{\text{op}} \|u_{n_j}\| \to 0$.
Since $P_{n_j}g \to g$, $u_{n_j} = P_{n_j}g - P_{n_j}Ku_{n_j} \to g - Ku$; by weak uniqueness $u = g - Ku$, hence $Au = g$.

Since $g \in \mathcal{H}$ was arbitrary, $A(\mathcal{H}) = \mathcal{H}$.
*Structural Commentary*: $\|Ah\| \ge c\|h\|$ and $(I + P_nK)u_n = P_ng \Rightarrow Au = g$.
*Key Insight*: Compactness removes approximate kernels and passes projected solutions to a true solution. □

**Definition 6.2.13.** *Fredholm Operator and Index*
Let $\mathcal{H}$ be a Hilbert space and let $T \in \mathcal{L}(\mathcal{H})$.

(i) *Fredholm*: $\text{range}(T)$ is closed, $\dim \ker T < \infty$, and $\text{codim range}(T) < \infty$.
(ii) *Index*: $\text{ind}\, T := \dim \ker T - \text{codim range}(T)$ when $T$ is Fredholm.

**Theorem 6.2.14.** *Riesz–Schauder for Compact Perturbations of the Identity*
*Let $\mathcal{H}$ be Hilbert, let $K \in \mathcal{L}(\mathcal{H})$ be compact, and set $A := I + K$. Then:*

*(i)* $\dim\ker A < \infty$.
*(ii)* $\operatorname{range}(A)$ *is closed.*
*(iii)* $\operatorname{codim}\operatorname{range}(A) = \dim\ker(A^*) = \dim\ker A < \infty$.
*(iv)* $A$ *is Fredholm with* $\operatorname{ind} A = 0$.
*(v)* $A$ *is injective iff it is surjective.*

*Proof.*

(i) $\dim\ker A = +\infty \Rightarrow \exists (u_n) \subseteq \ker A$ with $\langle u_n, u_m\rangle = \delta_{nm}$.
Then $Ku_n = -u_n$ and $n \neq m \Rightarrow \|Ku_n - Ku_m\| = \sqrt{2}$, contradicting Proposition 6.2.2(iii).

(ii) Let $H_0 := (\ker A)^\perp$. If $\inf\{\|Ah\| : h \in H_0, \|h\| = 1\} = 0$, choose $h_n \in H_0$, $\|h_n\| = 1$, $Ah_n \to 0$.
By Proposition 6.2.2(iii), $Kh_{n_j} \to y$, so $h_{n_j} = Ah_{n_j} - Kh_{n_j} \to -y := h \in H_0$, $Ah = 0$.
Thus $h \in \ker A \cap H_0 = \{0\}$, contradiction. Hence $\exists c > 0 : \|Ah\| \geq c\,\|h\|$ on $H_0$.
Also $A(H_0) = A(\mathcal{H})$. If $Ah_n \to y$ with $h_n \in H_0$, then $\|h_n - h_m\| \leq c^{-1}\,\|Ah_n - Ah_m\|$.
Thus $h_n \to h \in H_0$ and $Ah = y$; hence $\operatorname{range}(A)$ is closed.

(iii) By Theorem 6.2.11, $K^*$ compact, so (i)–(ii) apply to $A^* = I + K^*$.
By Proposition 5.2.39, $\mathcal{H} = \operatorname{range}(A) \odot \ker(A^*)$, $\operatorname{codim}\operatorname{range}(A) = \dim\ker(A^*)$.
If $\dim\ker A < \dim\ker(A^*)$, choose injective $L : \ker A \to \ker(A^*)$ with $L(\ker A) \neq \ker(A^*)$, $F := LP_{\ker A}$.
Then $\operatorname{rank} F < +\infty$, $K + F$ compact. For $x = z + n$, $z \in H_0$, $n \in \ker A$,
$(A + F)x = Az + Ln \in \operatorname{range}(A) \odot L(\ker A)$; injectivity of $L$ gives $\ker(A + F) = \{0\}$ and
$\operatorname{range}(A+F) = \operatorname{range}(A) \odot L(\ker A) \neq \mathcal{H}$, contradicting Proposition 6.2.12. Thus $\dim\ker A \geq \dim\ker(A^*)$.
Apply the same argument to $A^*$ and use $(K^*)^* = K$ to get $\dim\ker A = \dim\ker(A^*)$.

(iv) (i)–(iii) give $\operatorname{range}(A)$ closed, $\dim\ker A < +\infty$, $\operatorname{codim}\operatorname{range}(A) < +\infty$.
Also $\dim\ker A = \operatorname{codim}\operatorname{range}(A)$; hence $A$ is Fredholm and $\operatorname{ind} A = 0$.

(v) (iii) gives $\ker A = \{0\} \iff \operatorname{codim}\operatorname{range}(A) = 0 \iff \operatorname{range}(A) = \mathcal{H}$.

*Structural Commentary*: Compactness $\Rightarrow \dim\ker A < \infty$, $\operatorname{range}(A)$ closed; $F = LP_{\ker A} \Rightarrow \dim\ker A = \dim\ker(A^*)$.
*Key Insight*: For $A = I + K$ compactly perturbed, injectivity and surjectivity have the same finite defect. □

**Corollary 6.2.15.** *Fredholm Alternative*
*Let $\mathcal{H}$ be Hilbert, let $K \in \mathcal{L}(\mathcal{H})$ be compact, let $\lambda \in \mathbb{F} \setminus \{0\}$, and set $T_\lambda := \lambda I - K$.*

*(i)* $\operatorname{range}(T_\lambda) = \mathcal{H}$ *iff* $\ker T_\lambda = \{0\}$.
*(ii) If the conditions in (i) hold, then* $T_\lambda^{-1} \in \mathcal{L}(\mathcal{H})$.

*Proof.* Let $\widetilde{K} := -\lambda^{-1}K$; $\widetilde{K}$ compact, $T_\lambda = \lambda(I + \widetilde{K})$, $\ker T_\lambda = \ker(I + \widetilde{K})$, $\operatorname{range}(T_\lambda) = \mathcal{H}$ iff $\operatorname{range}(I + \widetilde{K}) = \mathcal{H}$.

*(i)* ($\Rightarrow$) $\operatorname{range}(T_\lambda) = \mathcal{H} \Rightarrow \operatorname{range}(I + \widetilde{K}) = \mathcal{H} \Rightarrow \ker(I + \widetilde{K}) = \{0\}$ by Theorem 6.2.14(v), hence $\ker T_\lambda = \{0\}$.
*(i)* ($\Leftarrow$) $\ker T_\lambda = \{0\} \Rightarrow \ker(I + \widetilde{K}) = \{0\} \Rightarrow \operatorname{range}(I + \widetilde{K}) = \mathcal{H}$ by Theorem 6.2.14(v), hence $\operatorname{range}(T_\lambda) = \mathcal{H}$.
*(ii)* (i) $\Rightarrow T_\lambda$ bijective; Corollary 5.4.4 $\Rightarrow T_\lambda^{-1} \in \mathcal{L}(\mathcal{H})$.

*Structural Commentary*: $T_\lambda = \lambda(I + \widetilde{K})$ with $\widetilde{K}$ compact; bijective $T_\lambda$ has bounded inverse.
*Key Insight*: $\forall f$, $T_\lambda u = f$ is solvable iff $\ker T_\lambda = \{0\}$. □

**Theorem 6.2.16.** *Fredholm Index Zero as an Invertible Plus Compact Operator*
*Let $\mathcal{H}$ be Hilbert and let $T \in \mathcal{L}(\mathcal{H})$. The following are equivalent:*

*(i)* $T$ *is Fredholm and* $\operatorname{ind} T = 0$.
*(ii)* $\exists S, K \in \mathcal{L}(\mathcal{H})$ *such that* $T = S + K$, $S$ *is invertible, and* $K$ *is compact.*

*Proof.*

(i) $\Rightarrow$ (ii) $N := \ker T$, $R := \operatorname{range}(T)$, $Z := \ker(T^*)$.
By Proposition 5.2.39, $\mathcal{H} = R \odot Z$ and $\dim Z = \operatorname{codim} R$. $\operatorname{ind} T = 0 \Rightarrow \dim N = \dim Z < +\infty$.
$A : N \to Z$ isomorphism, $P_N$ orthogonal projection, $K := AP_N$, $S := T - K$.
Then $\operatorname{rank} K < +\infty \Rightarrow K$ compact and $T = S + K$. For $h = v + u$, $v \in N^\perp$, $u \in N$, $Sh = Tv - Au \in R \odot Z$ and $Sh = 0 \Rightarrow Tv = 0$, $Au = 0 \Rightarrow v = 0$, $u = 0$; hence $\ker S = \{0\}$.
Also $T(N^\perp) = R$, $A(N) = Z$, and $y = r + z \in R \odot Z \Rightarrow \exists v \in N^\perp, u \in N : Tv = r, Au = -z, S(v + u) = y$; hence $\operatorname{range} S = \mathcal{H}$ and $S^{-1} \in \mathcal{L}(\mathcal{H})$ by Corollary 5.4.4.

(ii) $\Rightarrow$ (i) $C := S^{-1}K$, $A := I + C$. By Proposition 6.2.5(ii), $C$ is compact and $T = SA$.
By Theorem 6.2.14(iv), $A$ is Fredholm, $\operatorname{ind} A = 0$. $S$ isomorphism $\Rightarrow \ker T = \ker A$, $\operatorname{range} T = S(\operatorname{range} A)$, $\operatorname{codim}\operatorname{range} T = \operatorname{codim}\operatorname{range} A$. Hence $T$ is Fredholm and $\operatorname{ind} T = 0$.

*Structural Commentary*: $K = AP_{\ker T}$ makes $S = T - K$ invertible; $T = S(I + C)$ with $C$ compact.
*Key Insight*: $\operatorname{ind} T = 0$ iff $T$ is invertible modulo compact operators. □

**Corollary 6.2.17.** *Index-Zero Fredholm Operators Are Closed under Composition*
*Let $\mathcal{H}$ be Hilbert, $S, T \in \mathcal{L}(\mathcal{H})$ be Fredholm with $\operatorname{ind} S = \operatorname{ind} T = 0$. Then $ST$ is Fredholm and $\operatorname{ind}(ST) = 0$.*

*Proof.* By Theorem 6.2.16, $S = S_0 + K_S$, $T = T_0 + K_T$, with $S_0, T_0$ invertible and $K_S, K_T$ compact.
Let $U := S_0T_0$, $C := S_0K_T + K_ST_0 + K_SK_T$. Then $ST = U + C$, $U$ invertible, by Proposition 6.2.5, $C$ compact.
Theorem 6.2.16 $\Rightarrow$ $ST$ Fredholm and $\operatorname{ind}(ST) = 0$.
*Structural Commentary*: $(S_0 + K_S)(T_0 + K_T) = S_0T_0 + C$, $C \in \mathcal{K}(\mathcal{H})$.
*Key Insight*: Invertible modulo compact operators are closed under products. □

**Corollary 6.2.18.** *Compact Perturbations Preserve Index-Zero Fredholm Operators*
*Let $\mathcal{H}$ be Hilbert and let $T, K \in \mathcal{L}(\mathcal{H})$. If $T$ is Fredholm, $\operatorname{ind} T = 0$, and $K$ is compact, then:*

*(i)* $T + K$ *is Fredholm.*
*(ii)* $\operatorname{ind}(T + K) = 0$.

*Proof.* By Theorem 6.2.16, $T = S + C$, $S$ invertible, $C$ compact. Set $\widetilde{C} := C + K$.
By Proposition 6.2.5, $\widetilde{C} \in \mathcal{K}(\mathcal{H})$ and $T + K = S + \widetilde{C}$.

(i) $T + K = S + \widetilde{C}$, $S$ invertible, $\widetilde{C}$ compact $\Rightarrow$ $T + K$ Fredholm by Theorem 6.2.16.
(ii) $T + K = S + \widetilde{C}$, $S$ invertible, $\widetilde{C}$ compact $\Rightarrow$ $\operatorname{ind}(T + K) = 0$ by Theorem 6.2.16.

*Structural Commentary*: $T = S + C$, $K \in \mathcal{K}(\mathcal{H}) \Rightarrow T + K = S + \widetilde{C}$, $\widetilde{C} = C + K \in \mathcal{K}(\mathcal{H})$.
*Key Insight*: $\operatorname{ind} = 0$ is stable under compact perturbation. □

## 6.3 Spectral Theory of Operators

For bounded operators, spectrum replaces the finite-dimensional eigenvalue list. The resolvent tracks invertibility, while the spectrum records every obstruction to it.

**Definition 6.3.1.** *Spectrum and Resolvent*
Let $\mathcal{X}$ be a complex Banach space and let $T \in \mathcal{L}(\mathcal{X})$.

(i) *Resolvent set*: $\rho(T) := \{\lambda \in \mathbb{C} : \lambda I - T \text{ is bijective and } (\lambda I - T)^{-1} \in \mathcal{L}(\mathcal{X})\}$.
(ii) *Spectrum*: $\sigma(T) := \mathbb{C} \setminus \rho(T)$.

**Theorem 6.3.2.** *Spectrum of a Complex Bounded Operator*
*Let $\mathcal{X}$ be a nonzero complex Banach space and let $T \in \mathcal{L}(\mathcal{X})$. Then:*

*(i)* $\sigma(T) \neq \varnothing$;
*(ii)* $\sigma(T)$ *is compact in* $\mathbb{C}$;
*(iii)* $\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{op}\}$.

*Proof.* For $\lambda \in \rho(T)$, $R(\lambda) := (\lambda I - T)^{-1}$. If $\mu \in \rho(T)$, $|\lambda - \mu| \, \|R(\mu)\|_{\text{op}} < 1$; $\lambda I - T = (I + (\lambda - \mu)R(\mu))(\mu I - T)$.
By Corollary 5.1.7, $\lambda \in \rho(T)$, $R(\lambda) = R(\mu)(I + (\lambda - \mu)R(\mu))^{-1}$. If $|\lambda| > \|T\|_{\text{op}}$, then $\lambda I - T = \lambda(I - \lambda^{-1}T)$.
By Corollary 5.1.7, $\lambda \in \rho(T)$ and $\|R(\lambda)\|_{\text{op}} \leq (|\lambda| - \|T\|_{\text{op}})^{-1}$.

(i) Suppose $\sigma(T) = \varnothing$. By the local Neumann formula above, $R : \mathbb{C} \to \mathcal{L}(\mathcal{X})$ is entire.
For $x \in \mathcal{X}$ and $\Phi \in \mathcal{X}^*$, $f_{x,\Phi}(\lambda) := \Phi(R(\lambda)x)$ is an entire scalar function. By the estimate above, $f_{x,\Phi}(\lambda) \to 0$ as $|\lambda| \to +\infty$, and continuity bounds $f_{x,\Phi}$ on compact disks. By Liouville's theorem, $f_{x,\Phi} = 0$.
By Corollary 5.3.3, bounded linear functionals separate points, so $R(0) = 0$, contradicting $(-T)R(0) = I$ on the nonzero space $\mathcal{X}$.
(ii) By the local Neumann formula, $\rho(T)$ is open, so $\sigma(T)$ is closed in $\mathbb{C}$.
By outside-disk formula, $\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq \|T\|_{\text{op}}\}$, hence $\sigma(T)$ is bounded. Thus $\sigma(T)$ is compact.
(iii) If $|\lambda| > \|T\|_{\text{op}}$, then the outside-disk condition holds. By the outside-disk Neumann argument, $\lambda \in \rho(T)$.
Therefore every $\lambda \in \sigma(T)$ satisfies $|\lambda| \leq \|T\|_{\text{op}}$.

*Structural Commentary*: By Neumann, $\rho(T)$ is open and $\sigma(T) \subseteq \{\lambda : |\lambda| \leq \|T\|_{\text{op}}\}$. By Liouville, $\sigma(T) \neq \varnothing$.
*Key Insight*: Complex Banach spectra are nonempty compact obstructions to invertibility. □

**Definition 6.3.3.** *Spectral Radius*
Let $\mathcal{X}$ be a nonzero complex Banach space and let $T \in \mathcal{L}(\mathcal{X})$. The *spectral radius* of $T$ is $r(T) := \sup_{\lambda \in \sigma(T)} |\lambda|$.

**Theorem 6.3.4.** *Spectral Radius Formula*
*Let $\mathcal{X}$ be a nonzero complex Banach space and let $T \in \mathcal{L}(\mathcal{X})$.*
*Then the sequence $(\|T^n\|_{op}^{1/n})_{n\geq 1}$ converges in $[0,\infty)$, and $\lim_{n\to\infty} \|T^n\|_{op}^{1/n} = r(T)$.*

*Proof.* Set $a_n := \|T^n\|_{\text{op}}$ for $n \geq 0$, so $a_0 = 1$ and $a_{m+n} \leq a_m a_n$. If $a_m = 0$ for some $m \geq 1$, then $T$ is nilpotent.
Also $\|T^n\|_{\text{op}}^{1/n} \to 0$, $\sigma(T) = \{0\}$ by finite Neumann series and Theorem 6.3.2.
Hence assume $a_n > 0$ for all $n \geq 1$. For fixed $m \geq 1$ and $n = qm + r$ with $0 \leq r < m$, $a_n^{1/n} \leq a_m^{q/n} a_r^{1/n}$.
By lim sup in $n$, $\limsup_n a_n^{1/n} \leq a_m^{1/m}$, hence $\limsup_n a_n^{1/n} \leq \inf_{m\geq 1} a_m^{1/m} \leq \liminf_n a_n^{1/n}$.
Thus $L := \lim_n a_n^{1/n} = \inf_{m\geq 1} a_m^{1/m}$. If $|\lambda| > L$, choose $n$ with $a_n < |\lambda|^n$.
Then $\|T^n/\lambda^n\|_{\text{op}} < 1$, so Corollary 5.1.7 makes $\lambda^n I - T^n$ invertible.
With $Q_n := \sum_{j=0}^{n-1} \lambda^{n-1-j} T^j$, $\lambda^n I - T^n = (\lambda I - T)Q_n = Q_n(\lambda I - T)$; as these are polynomials in $T$,
$(\lambda I - T)^{-1} = Q_n(\lambda^n I - T^n)^{-1}$. Thus $\lambda \in \rho(T)$, so $\sigma(T) \subseteq \{z \in \mathbb{C} : |z| \leq L\}$ and $r(T) \leq L$.
Let $R > r(T)$, $R_0 > \max\{R, \|T\|_{\text{op}}\}$. For $|z| = R_0$, by Neumann expansion, $(zI - T)^{-1} = \sum_{k\geq 0} z^{-k-1} T^k$.
By Cauchy's coefficient formula $T^n = \frac{1}{2\pi i} \int_{|z|=R_0} z^n (zI - T)^{-1}\, dz$.
Since $\{z : R \leq |z| \leq R_0\} \subseteq \rho(T)$, Cauchy's theorem moves the contour to $|z| = R$.
With $M_R := \sup_{|z|=R} \left\|(zI - T)^{-1}\right\|_{\text{op}} < +\infty$, $a_n \leq M_R R^{n+1}$, hence $L \leq R$. By letting $R \downarrow r(T)$, $L \leq r(T)$.
*Structural Commentary*: By submultiplicativity, $L$ exists; Neumann, $r(T) \leq L$; resolvent bounds, $L \leq r(T)$.
*Key Insight*: The spectral radius is asymptotic operator-power growth. □

**Definition 6.3.5.** *Quadratic Form and Rayleigh Quotient*
Let $\mathcal{H}$ be a real Hilbert space and let $T \in \mathcal{L}(\mathcal{H})$ be self-adjoint.

(i) *Quadratic form*: $Q_T(h) := \langle Th, h\rangle$ for $h \in \mathcal{H}$.
(ii) *Rayleigh quotient*: $R_T(h) := \frac{\langle Th,h\rangle}{\langle h,h\rangle}$ for $h \in \mathcal{H} \setminus \{0\}$.

A unit-sphere maximizer of $Q_T$ is equivalently a maximizer of $R_T$ on $\mathcal{H} \setminus \{0\}$.

**Example 6.3.6.** *A Self-Adjoint Operator without Eigenvectors*
Let $a < b$, set $H := L^2([a,b];\mathbb{C})$, and define $(Tf)(x) := xf(x)$.
With $M := \max\{|a|, |b|\}$, $\|Tf\|_2 \leq M \|f\|_2$, so $T \in \mathcal{L}(H)$.
For $u, v \in H$, $\langle Tu, v\rangle = \int_a^b xu(x)\overline{v(x)}\, dx = \int_a^b u(x)\overline{xv(x)}\, dx = \langle u, Tv\rangle$, hence $T = T^*$.
If $Tf = \lambda f$ for $\lambda \in \mathbb{C}$, then $(x - \lambda)f(x) = 0$ a.e. Let $E_\lambda := \{x \in [a,b] : x = \lambda\}$.
Since $m(E_\lambda) = 0$, $f = 0$ a.e. on $[a,b] \setminus E_\lambda$ and on $E_\lambda$, hence $f = 0$ in $H$.
Thus no nonzero $f \in H$ satisfies $Tf = \lambda f$, so $T$ has no eigenvectors.

**Lemma 6.3.7.** *Hilbert–Schmidt Lemma*
*Let $0 \neq \mathcal{H}$ be real Hilbert space, $T = T^* \in \mathcal{L}(\mathcal{H})$ compact, $Q_T(h) := \langle Th, h\rangle$. Then $\exists \lambda \in \mathbb{R}$, $h_* \in \mathcal{H}$ such that:*

*(i)* $\|h_*\| = 1$;
*(ii)* $Th_* = \lambda h_*$;
*(iii)* $|\lambda| = \|T\|_{op} = \sup_{\|h\|=1} |Q_T(h)|$.

*Proof.* If $T = 0$, choose $\|h_*\| = 1$ and set $\lambda = 0$; then (i)–(iii) hold. Assume $T \neq 0$ and put $\eta := \|T\|_{\text{op}}$.
By Theorem 5.2.38, $\eta = \sup_{\|h\|=1} |Q_T(h)| > 0$. Choose $\|x_n\| = 1$ and $s_n \in \{-1, 1\}$ with $Q_{s_n T}(x_n) \to \eta$.
Passing to a subsequence, $s_n = s$; write $h_n := x_n$ and $A := sT$.
By Theorems 6.1.17 and 6.2.7, after passing to a further subsequence, $h_n \rightharpoonup h_*$ and $Ah_n \to Ah_*$ in norm.
Since $Q_A(h_n) - Q_A(h_*) = \langle Ah_n - Ah_*, h_n\rangle + \langle Ah_*, h_n - h_*\rangle \to 0$, $Q_A(h_*) = \eta$. Set $\lambda := s\eta$.

(i) By weak lower semicontinuity, $\|h_*\| \leq 1$.
Since $|Q_A(g)| \leq \eta \|g\|^2$ $\forall g \in \mathcal{H}$, $\eta = Q_A(h_*) \leq \eta \|h_*\|^2$. As $\eta > 0$, $\|h_*\| = 1$.
(ii) By (i), $R_A(h_*) = \eta$. For $\|h\| = 1$, $R_A(h) = Q_A(h) \leq |Q_T(h)| \leq \eta$, so $h_*$ maximizes $R_A$.
For $h_0 \in \mathcal{H}$, $\phi(t) := R_A(h_* + th_0)$ has $\phi'(0) = 0$, hence $0 = 2\langle Ah_* - \eta h_*, h_0\rangle$.
Since $h_0$ is arbitrary, $Ah_* = \eta h_*$. Thus $Th_* = s\eta h_* = \lambda h_*$.
(iii) By the definition of $\lambda$ and by Theorem 5.2.38, $|\lambda| = \eta = \|T\|_{\text{op}} = \sup_{\|h\|=1} |Q_T(h)|$.

*Structural Commentary*: By compactness, signed $Q_T$ attains $\|T\|_{\text{op}}$; by stationarity, $Th_* = \lambda h_*$.
*Key Insight*: A compact self-adjoint norm maximizer is an eigenvector. □

**Corollary 6.3.8.** *Extremal Rayleigh Values Are Eigenvalues*
*Let $\mathcal{H}$ be a nonzero real Hilbert space, let $T \in \mathcal{L}(\mathcal{H})$ be compact and self-adjoint, and set $Q_T(h) := \langle Th, h\rangle$, $\alpha := \inf_{\|h\|=1} Q_T(h)$, $\beta := \sup_{\|h\|=1} Q_T(h)$.*

*(i) If $\beta > 0$, then $\exists h_\beta \in \mathcal{H}$ such that $\|h_\beta\| = 1$ and $Th_\beta = \beta h_\beta$.*
*(ii) If $\alpha < 0$, then $\exists h_\alpha \in \mathcal{H}$ such that $\|h_\alpha\| = 1$ and $Th_\alpha = \alpha h_\alpha$.*
*(iii) If $\beta > 0$ and $(h_n) \subset \mathcal{H}$ satisfies $\|h_n\| = 1$, $Q_T(h_n) \to \beta$, then $\exists n_j \uparrow \infty$ and $h_\beta \in \mathcal{H}$ with $h_{n_j} \to h_\beta$ in norm, $\|h_\beta\| = 1$, and $Th_\beta = \beta h_\beta$.*

*Proof.*

(i) Assume $\beta > 0$. Choose $\|u_n\| = 1$ with $Q_T(u_n) \to \beta$.
By Theorems 6.1.17 and 6.2.7, after passing to a subsequence, $u_n \rightharpoonup u_*$ and $Tu_n \to Tu_*$ in norm.
Since $Q_T(u_n) - Q_T(u_*) = \langle Tu_n - Tu_*, u_n\rangle + \langle Tu_*, u_n - u_*\rangle \to 0$, $Q_T(u_*) = \beta$.
By weak lower semicontinuity, $\|u_*\| \le 1$. As $Q_T(g) \le \beta \|g\|^2$ $\forall g \in \mathcal{H}$, $\beta = Q_T(u_*) \le \beta \|u_*\|^2$; $\|u_*\| = 1$.
Thus $u_*$ maximizes $R_T$. For $h \in \mathcal{H}$, $\phi(t) := R_T(u_* + th)$ has $\phi'(0) = 0$, so $0 = 2\langle Tu_* - \beta u_*, h\rangle$.
Hence $Tu_* = \beta u_*$; take $h_\beta := u_*$.

(ii) Assume $\alpha < 0$. Apply (i) to $-T$. Its upper Rayleigh value is $\beta_{-T} = \sup_{\|h\|=1} \langle -Th, h\rangle = -\alpha > 0$, so $\exists h_\alpha$ with $\|h_\alpha\| = 1$ and $(-T)h_\alpha = (-\alpha)h_\alpha$. Hence $Th_\alpha = \alpha h_\alpha$.

(iii) Assume $\beta > 0$, $\|h_n\| = 1$, $Q_T(h_n) \to \beta$. By compactness step in (i), $h_{n_j} \rightharpoonup h_\beta$, $Th_{n_j} \to Th_\beta$ in norm.
As in (i), $Q_T(h_\beta) = \beta$, $\|h_\beta\| = 1$, and $Th_\beta = \beta h_\beta$.
By Theorem 6.1.20, since $\|h_{n_j}\| = \|h_\beta\| = 1$, $h_{n_j} \to h_\beta$ in norm.

*Structural Commentary*: By compactness, attain $\beta$; by $-T$, attain $\alpha$; by Radon–Riesz, $\rightharpoonup + \|\cdot\| \to 1 \Rightarrow \to$.
*Key Insight*: Compact self-adjoint Rayleigh extrema are eigenvalues. □

**Theorem 6.3.9.** *Hilbert–Schmidt Theorem for Compact Self-Adjoint Operators*
*Let $\mathcal{H}$ be a real Hilbert space and let $K \in \mathcal{L}(\mathcal{H})$ be compact and self-adjoint.*
*If $\operatorname{rank} K = +\infty$, then $\exists$ orthonormal sequence $(\psi_k)_{k\ge1} \subset \mathcal{H}$ and nonzero scalars $(\lambda_k)_{k\ge1} \subset \mathbb{R}$ such that:*

*(i) $K\psi_k = \lambda_k \psi_k$ for all $k \ge 1$, $|\lambda_{k+1}| \le |\lambda_k|$, and $\lambda_k \to 0$;*
*(ii) $\operatorname{cspan}\{\psi_k : k \ge 1\} = (\ker K)^\perp$;*
*(iii) $Kh = \sum_{k\ge1} \lambda_k \langle h, \psi_k\rangle \psi_k$ for every $h \in \mathcal{H}$, with norm convergence.*

*If $\operatorname{rank} K < +\infty$, then:*

*(iv) $\exists m \ge 0$, nonzero $\lambda_1, \dots, \lambda_m \in \mathbb{R}$, and an orthonormal basis $(\psi_k)_{k=1}^m$ of $\operatorname{range} K = (\ker K)^\perp$ such that $K\psi_k = \lambda_k \psi_k$ and $Kh = \sum_{k=1}^m \lambda_k \langle h, \psi_k\rangle \psi_k$ for every $h \in \mathcal{H}$.*

*Proof.*

(i) Put $M_0 := \{0\}$ and $H_0 := \mathcal{H}$. If $K|_{H_j} \ne 0$, then $H_j$ is $K$-invariant and $K|_{H_j}$ is compact self-adjoint.
By Lemma 6.3.7, $\psi_{j+1} \in H_j$, $\lambda_{j+1} \ne 0$ with $\|\psi_{j+1}\| = 1$, $K\psi_{j+1} = \lambda_{j+1}\psi_{j+1}$, and $|\lambda_{j+1}| = \|K|_{H_j}\|_{\mathrm{op}}$.
Set $M_{j+1} := M_j \oplus \operatorname{span}\{\psi_{j+1}\}$, $H_{j+1} := M_{j+1}^\perp$. Since $K(M_{j+1}) \subseteq M_{j+1}$, $K = K^*$, $K(H_{j+1}) \subseteq H_{j+1}$.
Thus the construction continues while $K|_{H_j} \ne 0$, and $|\lambda_{j+1}| \le |\lambda_j|$.
If $\operatorname{rank} K = +\infty$, it cannot stop, since $K|_{H_m} = 0$ implies $\operatorname{range} K \subseteq M_m$.
If $\lambda_k \not\to 0$, then for some $\varepsilon > 0$, $\|K\psi_j - K\psi_k\|^2 = \lambda_j^2 + \lambda_k^2 \ge 2\varepsilon^2$ for infinitely many $j \ne k$, contradiction.

(ii) Set $M := \operatorname{cspan}\{\psi_k : k \ge 1\}$. Since $K(M) \subseteq M$ and $K = K^*$, $K(M^\perp) \subseteq M^\perp$.
If $K|_{M^\perp} \ne 0$, by Lemma 6.3.7 choose $\phi \in M^\perp$, $\mu \ne 0$ with $\|\phi\| = 1$ and $K\phi = \mu\phi$.
Then $\phi \in H_{k-1}$ $\forall k$, so $|\mu| \le \|K|_{H_{k-1}}\|_{\mathrm{op}} = |\lambda_k| \to 0$, contradiction. Hence $M^\perp \subseteq \ker K$.
Conversely, $x \in \ker K$ implies $\langle x, \psi_k\rangle = \lambda_k^{-1}\langle Kx, \psi_k\rangle = 0$ $\forall k$, so $\ker K \subseteq M^\perp$.
Thus $M^\perp = \ker K$ and $M = (\ker K)^\perp$ by Theorem 5.2.12.

(iii) For $h \in \mathcal{H}$, write $h = m + z$ with $m \in M = (\ker K)^\perp$ and $z \in \ker K$. By Theorem 5.2.22, $m = \sum_{k\ge1} \langle h, \psi_k\rangle \psi_k$ in norm. By continuity of $K$, $Kh = Km = \sum_{k\ge1} \lambda_k \langle h, \psi_k\rangle \psi_k$ in norm.

(iv) If $\operatorname{rank} K < +\infty$, the construction stops at some $m \ge 0$; otherwise $\{\psi_k\} \subset \operatorname{range} K$ is an infinite orthonormal set in finite-dimensional $\operatorname{range} K$. Let $M_m := \operatorname{span}\{\psi_1, \dots, \psi_m\}$.
Since $K|_{M_m^\perp} = 0$, $\operatorname{range} K \subseteq M_m$; since $\psi_k = \lambda_k^{-1} K\psi_k$, $M_m \subseteq \operatorname{range} K$. Hence $M_m = \operatorname{range} K$.
Also $M_m^\perp = \ker K$ by argument in (ii), so $\operatorname{range} K = (\ker K)^\perp$. For $h = m_0 + z$ with $m_0 \in M_m$, $z \in \ker K$, by the finite orthonormal expansion of $m_0$, $Kh = Km_0 = \sum_{k=1}^m \lambda_k \langle h, \psi_k\rangle \psi_k$.

*Structural Commentary*: Iterated extremal eigenvectors exhaust $(\ker K)^\perp$; Fourier expansion diagonalizes $K$.
*Key Insight*: Nonzero compact self-adjoint eigendirections are finite or countable, accumulating only at 0. □

**Perspective 6.3.10.** *Hilbert Operator Spectral Theory*
On a complex Hilbert space $\mathcal{H}$, normal $T \in \mathcal{L}(\mathcal{H})$ is diagonalized by spectral functional calculus.
By Theorem 6.3.9, compact self-adjoint $K \in \mathcal{L}(\mathcal{H})$ has nonzero eigenvalues $(\lambda_k)$ and an orthonormal eigenfamily $(\psi_k)$ spanning $(\ker K)^\perp$, so $\mathcal{H} = \ker K \oplus \operatorname{cspan}\{\psi_k\}$ and $Kh = \sum_k \lambda_k \langle h, \psi_k\rangle \psi_k \ \forall h \in \mathcal{H}$.
The sum is finite iff $\operatorname{rank} K < +\infty$; otherwise $\lambda_k \to 0$.

## 6.4 Measure Duality and Riesz Representation

On a compact Hausdorff space, Riesz–Markov turns positive functionals on $C(K)$ into Radon measures. The remaining step is order-theoretic: every bounded functional is the difference of two positive functionals, exactly as a signed measure is the difference of its Jordan variations.

**Proposition 6.4.1.** *Positive Functionals on $C(K)$ Are Continuous*
*Let $K$ be compact Hausdorff, $C(K) = C(K;\mathbb{R})$, $\|\cdot\|_\infty$, $L : C(K) \to \mathbb{R}$ linear with $L(C(K)_+) \subseteq [0,+\infty)$. Then:*

*(i)* $L \in C(K)^*$.
*(ii)* $\|L\|_{op} = L(\mathbf{1}_K)$.

*Proof.*

(i) For $f \in C(K)$, $-\|f\|_\infty \mathbf{1}_K \le f \le \|f\|_\infty \mathbf{1}_K$; $-L(\mathbf{1}_K)\|f\|_\infty \le L(f) \le L(\mathbf{1}_K)\|f\|_\infty$; $|L(f)| \le L(\mathbf{1}_K)\|f\|_\infty$.
(ii) By (i), $\|L\|_{\mathrm{op}} \le L(\mathbf{1}_K)$. If $K = \varnothing$, both sides are 0; if $K \neq \varnothing$, then $\|\mathbf{1}_K\|_\infty = 1$, so $\|L\|_{\mathrm{op}} \ge L(\mathbf{1}_K)$.

*Structural Commentary*: $-\|f\|_\infty \mathbf{1}_K \le f \le \|f\|_\infty \mathbf{1}_K$ yields $|L(f)| \le L(\mathbf{1}_K)\|f\|_\infty$.
*Key Insight*: Positivity plus $\mathbf{1}_K$ controls $\|L\|_{\mathrm{op}}$. □

**Proposition 6.4.2.** *Positive Decomposition of Functionals on $C(K)$*
*Let $K$ be compact Hausdorff and let $L \in C(K)^*$. Then $\exists L_+, L_- \in C(K)^*$ positive such that:*

*(i)* $L = L_+ - L_-$.
*(ii)* $\|L\|_{op} = L_+(\mathbf{1}_K) + L_-(\mathbf{1}_K)$.

*Proof.*

(i) For $f \in C(K)_+$, set $P(f) := \sup_{0 \le g \le f} L(g)$; boundedness of $L$ implies $P(f) < +\infty$.
If $0 \le h \le f_1 + f_2$, then $h = (h \wedge f_1) + (h - h \wedge f_1)$, terms in $[0, f_1]$ and $[0, f_2]$; $P(f_1 + f_2) \le P(f_1) + P(f_2)$.
The reverse inequality follows from $0 \le g_i \le f_i \Rightarrow 0 \le g_1 + g_2 \le f_1 + f_2$; homogeneity is immediate.
Define $L_+(u) := P(u^+) - P(u^-)$ and $L_- := L_+ - L$. Then $L_+$ is positive linear, $L_-(f) = P(f) - L(f) \ge 0$ for $f \in C(K)_+$, and by Proposition 6.4.1(i), $L_\pm \in C(K)^*$.
(ii) Set $A := L_+(\mathbf{1}_K) + L_-(\mathbf{1}_K)$. By (i) and Proposition 6.4.1(ii), $\|L\|_{\mathrm{op}} \le A$.
For $\varepsilon > 0$, choose $0 \le \varphi \le \mathbf{1}_K$ with $L(\varphi) > P(\mathbf{1}_K) - \varepsilon = L_+(\mathbf{1}_K) - \varepsilon$.
Since $\|2\varphi - \mathbf{1}_K\|_\infty \le 1$ and $L_-(\mathbf{1}_K) = L_+(\mathbf{1}_K) - L(\mathbf{1}_K)$, $\|L\|_{\mathrm{op}} \ge L(2\varphi - \mathbf{1}_K) > A - 2\varepsilon$. Let $\varepsilon \downarrow 0$.

*Structural Commentary*: $P(f) = \sup_{0 \le g \le f} L(g)$ is the positive part on $C(K)_+$; $2\varphi - \mathbf{1}_K$ tests $\|L\|_{\mathrm{op}}$.
*Key Insight*: $L = L_+ - L_-$ and $\|L\|_{\mathrm{op}} = L_+(\mathbf{1}_K) + L_-(\mathbf{1}_K)$. □

**Theorem 6.4.3.** *Riesz Representation for $C(K)^*$*
*Let $K$ be compact Hausdorff. Let $\mathcal{M}_{\mathrm{Rad}}(K)$ be the finite signed measures $\nu$ on $K$ with $\nu^\pm$ Radon.*
*Let $\|\nu\|_{\mathrm{var}} := |\nu|(K)$, and define $(T\nu)(f) := \int_K f \, d\nu$ for $f \in C(K)$. Then:*

*(i) $T$ is linear and well defined.*
*(ii)* $\|T(\nu)\|_{op} = \|\nu\|_{\mathrm{var}}$ *for all* $\nu \in \mathcal{M}_{\mathrm{Rad}}(K)$.
*(iii) $T$ is bijective.*

*Proof.*

(i) For $\nu \in \mathcal{M}_{\mathrm{Rad}}(K)$ and $f \in C(K)$, $|T(\nu)(f)| \le \int_K |f| \, d|\nu| \le \|f\|_\infty \|\nu\|_{\mathrm{var}}$.
Hence $T(\nu) \in C(K)^*$ and $\|T(\nu)\|_{\mathrm{op}} \le \|\nu\|_{\mathrm{var}}$. Linearity is signed-integral linearity.
(ii) Set $L := T(\nu)$. By Proposition 6.4.2(i), Theorem 5.2.14, $L = L_+ - L_-$, $L_\pm = T(\mu_\pm)$ for finite Radon $\mu_\pm$.
Since $T(\nu^+ + \mu_-) = T(\nu^- + \mu_+)$, by uniqueness in Theorem 5.2.14, $\nu^+ + \mu_- = \nu^- + \mu_+$.
By Theorem 3.1.6(ii),(iii), $\nu^\pm \le \mu_\pm$. Therefore $\|\nu\|_{\mathrm{var}} \le \mu_+(K) + \mu_-(K) = L_+(\mathbf{1}_K) + L_-(\mathbf{1}_K) = \|L\|_{\mathrm{op}}$ by Proposition 6.4.2(ii). Combine with (i).
(iii) Let $L \in C(K)^*$. Proposition 6.4.2(i), Theorem 5.2.14: $L = L_+ - L_-$, $L_\pm = T(\mu_\pm)$ for finite Radon $\mu_\pm$.
Set $\nu := \mu_+ - \mu_-$. By Theorem 3.1.6(ii),(iii), $\nu^\pm \le \mu_\pm$.
If $0 \le \theta \le \mu$ and $\mu$ is finite Radon, then $\theta$ is Radon since the same open and compact approximants work with no larger error. Hence $\nu \in \mathcal{M}_{\mathrm{Rad}}(K)$, $T(\nu) = L$. Thus $T$ is surjective; injectivity follows from (ii).

*Structural Commentary*: Positive Riesz–Markov: $L_\pm = T(\mu_\pm)$; Jordan minimality: $\|T(\nu)\|_{\mathrm{op}} = \|\nu\|_{\mathrm{var}}$.
*Key Insight*: $C(K)^* \cong \mathcal{M}_{\mathrm{Rad}}(K)$ isometrically via $L(f) = \int_K f\, d\nu$. □

**Example 6.4.4.** *Point Evaluation Is a Dirac Measure*
Let $K \neq \emptyset$ be compact Hausdorff and fix $x_0 \in K$. Define $L_{x_0} \in C(K)^*$ by $L_{x_0}(f) := f(x_0)$.
Then $L_{x_0} = T(\delta_{x_0})$, since $\forall f \in C(K)$, $T(\delta_{x_0})(f) = \int_K f\, d\delta_{x_0} = f(x_0)$. Hence $\|L_{x_0}\|_{\mathrm{op}} = \|\delta_{x_0}\|_{\mathrm{var}} = 1$.

**Remark 6.4.5.** *Interval Form and Stieltjes Integrators*
Let $a < b$ and let $\nu$ be a finite signed Radon measure on $[a,b]$, extended by 0 to $\mathbb{R}$. Set $g(x) := \nu((-\infty, x])$.
By Proposition 4.1.5 applied to $\nu^\pm$, $g \in BV(\mathbb{R})$ is right-continuous and $g(-\infty) = 0$.
For $a \le c \le d \le b$, $\nu((c,d]) = g(d) - g(c)$ and $\nu(\{a\}) = g(a) - g(a-)$ with $g(a-) = 0$.
Thus, in the Lebesgue–Stieltjes convention, $\forall f \in C([a,b])$, $T(\nu)(f) = \int_{[a,b]} f\, d\nu = \int_{[a,b]} f\, dg$.

**Corollary 6.4.6.** *Weak-* Compactness for Radon Measures*
*Let $K$ be compact Hausdorff and set $\mathcal{M} := \mathcal{M}_{\mathrm{Rad}}(K)$, the finite signed measures $\nu$ on $K$ with $\nu^\pm$ Radon. Let $\|\nu\|_{\mathrm{var}} := |\nu|(K)$ and let $\tau := \sigma(\mathcal{M}, C(K))$ be induced by $\langle \nu, f\rangle := \int_K f\, d\nu$. If $A \subseteq \mathcal{M}$ is $\tau$-closed and $\sup_{\nu \in A} \|\nu\|_{\mathrm{var}} < +\infty$, then $A$ is $\tau$-compact.*

*Proof.* Let $(T\nu)(f) := \int_K f\, d\nu$. By Theorem 6.4.3, $T : \mathcal{M} \to C(K)^*$ is an isometric bijection identifying $\tau$ with $\sigma(C(K)^*, C(K))$. Choose $R < +\infty$ with $A \subseteq R\overline{B}_{\mathcal{M}}$. Then $T(A)$ is weak-* closed and $T(A) \subseteq R\overline{B}_{C(K)^*}$.
By Theorem 6.1.6, $R\overline{B}_{C(K)^*}$ is weak-* compact; hence $T(A)$ and $A = T^{-1}(T(A))$ are weak-* compact.
*Structural Commentary*: Riesz isometry: reduce to weak-* closed $T(A) \subseteq R\overline{B}_{C(K)^*}$; Banach–Alaoglu: compact.
*Key Insight*: $T : \mathcal{M}_{\mathrm{Rad}}(K) \cong C(K)^*$ transfers Banach–Alaoglu compactness to Radon measures. □

**Corollary 6.4.7.** *Bounded Measures on Compact Metric Spaces Have Weak-* Subsequences*
*Let $K$ be compact metric and let $\mu_n \ge 0$ be Borel measures on $K$ with $M := \sup_n \mu_n(K) < +\infty$.*
*Then $\exists n_k \uparrow \infty$ and a Radon measure $\mu \ge 0$ on $K$ such that $\int_K f\, d\mu_{n_k} \to \int_K f\, d\mu$ for all $f \in C(K)$.*

*Proof.* Set $T_n(f) := \int_K f\, d\mu_n$. By Proposition 6.4.1(ii), $T_n \in C(K)^*$, $T_n \ge 0$, and $\|T_n\|_{\mathrm{op}} = \mu_n(K) \le M$.
Since $C(K)$ is separable by Theorem 9.2.10, by Theorem 6.1.8, $\exists n_k \uparrow \infty$ and $T \in C(K)^*$ with $T_{n_k}(f) \to T(f)$ for all $f \in C(K)$. Positivity passes to $T$, so by Theorem 5.2.14, $T(f) = \int_K f\, d\mu$ for some Radon $\mu \ge 0$.
*Structural Commentary*: Bound $(T_n)$ in $C(K)^*$; use separability and Helly; represent the positive limit.
*Key Insight*: Bounded positive measures on compact metric $K$ are weak-* sequentially compact on $C(K)$. □

# Part XV
# Probability Foundations and Distributions

## 1 Probability Spaces and Random Variables

### 1.1 Probability Spaces and Event Algebra

Probability assigns numbers to experimental outcomes; measure theory specifies which subsets are events.

**Definition 1.1.1.** *Sample Spaces, Events, and Probability Spaces*

(i) A *sample space* is a set $\Omega$ of possible outcomes.
(ii) An *outcome* or *realization* is a point $\omega \in \Omega$.
(iii) An *event* is a subset $A \subseteq \Omega$; the admissible events form a *$\sigma$-algebra* $\mathcal{F} \subseteq \mathcal{P}(\Omega)$.
(iv) A *probability measure* on $(\Omega, \mathcal{F})$ is a measure $\mathbb{P} : \mathcal{F} \to [0,1]$ with $\mathbb{P}(\Omega) = 1$.
(v) The triple $(\Omega, \mathcal{F}, \mathbb{P})$ is a *probability space*.

*Key Insight*: Probability is countably additive size normalized so that the whole sample space has mass 1.

**Method 1.1.2.** *Sample Space Construction Method*
Choose $\Omega$ as the set of complete outcomes and encode the target condition as an event $A \subseteq \Omega$:

(i) *Finite product method*: for $m$ discrete trials with state set $S$, use $\Omega = S^m$ and coordinate constraints. For two coin tosses, $S = \{H, T\}$, $\Omega = S^2$, and first toss heads gives $A = \{\omega \in \Omega : \omega_1 = H\} = \{HH, HT\}$.
(ii) *Interval method*: for real-valued measurements, use $\Omega = \mathbb{R}$ and translate inequalities into intervals. $10 < \omega \leq 23$ gives $A = (10, 23]$.
(iii) *Cylinder method*: for infinite sequences with state set $S$, use $\Omega = S^{\mathbb{N}}$ and constrain finitely many coordinates; for coin tosses, $S = \{H, T\}$ and first head on toss 3 gives $A = \{\omega \in \Omega : \omega_1 = T, \omega_2 = T, \omega_3 = H\}$.

**Example 1.1.3.** *Introductory Sample Spaces and Events*

(i) If a commute passes three traffic lights, each with outcome stop $s$ or continue $c$, then $\Omega = \{c, s\}^3$. The event that the first light is a stop is $\{sss, ssc, scc, scs\}$.
(ii) If a print queue length is modeled as random, one possible sample space is $\Omega = \{0, 1, 2, \ldots\}$; with a known capacity $N$, one may instead use $\Omega = \{0, 1, \ldots, N\}$.
(iii) If the measured waiting time between events is nonnegative, a natural sample space is $\Omega = [0, \infty)$.

*Key Insight*: Choose $\Omega$ at the event resolution needed.

**Definition 1.1.4.** *Event Algebra, Indicators, and Monotone Events*

(i) For $A \in \mathcal{F}$, the *complement* is $A^c := \Omega \setminus A$.
(ii) For $A, B \in \mathcal{F}$, the *union* and *intersection* are $A \cup B$ and $A \cap B$.
(iii) For $A, B \in \mathcal{F}$, the *difference* is $A \setminus B := A \cap B^c$.
(iv) *Commutativity*: $A \cup B = B \cup A$ and $A \cap B = B \cap A$.
(v) *Associativity*: $(A \cup B) \cup C = A \cup (B \cup C)$ and $(A \cap B) \cap C = A \cap (B \cap C)$.
(vi) *Distributivity*: $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ and $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$.
(vii) *De Morgan laws*: $(A \cup B)^c = A^c \cap B^c$ and $(A \cap B)^c = A^c \cup B^c$.
(viii) Events $(A_i)_{i \in I}$ are *disjoint* or *mutually exclusive* if $A_i \cap A_j = \emptyset$ for $i \neq j$.
(ix) Events $(A_i)_{i \in I}$ form a *partition* of $\Omega$ if $A_i \cap A_j = \emptyset$ for $i \neq j$ and $\bigcup_{i \in I} A_i = \Omega$.
(x) The *indicator* of $A \in \mathcal{F}$ is $\mathbf{1}_A : \Omega \to \{0, 1\}$, where $\mathbf{1}_A(\omega) = 1$ for $\omega \in A$ and $\mathbf{1}_A(\omega) = 0$ for $\omega \notin A$.
(xi) A sequence $(A_n)$ is *increasing* to $A$, written $A_n \uparrow A$, if $A_1 \subseteq A_2 \subseteq \cdots$ and $A = \bigcup_{n \geq 1} A_n$.
(xii) A sequence $(A_n)$ is *decreasing* to $A$, written $A_n \downarrow A$, if $A_1 \supseteq A_2 \supseteq \cdots$ and $A = \bigcap_{n \geq 1} A_n$.

**Proposition 1.1.5.** *Indicator Algebra for Unions and Intersections*
*For events $A, B \in \mathcal{F}$: $\mathbf{1}_{A \cap B} = \mathbf{1}_A \mathbf{1}_B = \min\{\mathbf{1}_A, \mathbf{1}_B\}$, and $\mathbf{1}_{A \cup B} = \max\{\mathbf{1}_A, \mathbf{1}_B\}$.*

*Proof.* Fix $\omega \in \Omega$ and set $a := \mathbf{1}_A(\omega)$, $b := \mathbf{1}_B(\omega)$. Then $a, b \in \{0, 1\}$, $\mathbf{1}_{A \cap B}(\omega) = ab = \min\{a, b\}$, and $\mathbf{1}_{A \cup B}(\omega) = \max\{a, b\}$. Since $\omega$ was arbitrary, the identities hold pointwise.
*Structural Commentary*: Boolean operations on events become arithmetic operations on 0-1 variables.
*Key Insight*: Indicators let event algebra be manipulated inside sums and expectations. □




Figure 227: Set operations become event operations in a probability space.

**Example 1.1.6.** *Endpoint Sensitivity in Monotone Limits*
Let $\Omega = \mathbb{R}$.

(i) If $A_i = [0, \frac{1}{i}]$, then $\bigcup_{i=1}^{\infty} A_i = [0,1]$ and $\bigcap_{i=1}^{\infty} A_i = \{0\}$.

(ii) If $A_i = (0, \frac{1}{i})$, then $\bigcup_{i=1}^{\infty} A_i = (0,1)$ and $\bigcap_{i=1}^{\infty} A_i = \emptyset$.

*Key Insight*: Monotone set limits remember whether boundary points are included.

**Proposition 1.1.7.** *First Consequences of the Probability Axioms*
*For events $A, B \in \mathcal{F}$:*

*(i) Null event:* $\mathbb{P}(\emptyset) = 0$.
*(ii) Monotonicity:* $A \subseteq B \Rightarrow \mathbb{P}(A) \leq \mathbb{P}(B)$.
*(iii) Bounds:* $0 \leq \mathbb{P}(A) \leq 1$.
*(iv) Complement:* $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$.
*(v) Disjoint additivity:* $A \cap B = \emptyset \Rightarrow \mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$.
*(vi) Two-set inclusion-exclusion:* $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$.

*Proof.*

(i) Countable additivity on $\Omega = \Omega \sqcup \emptyset \sqcup \emptyset \sqcup \cdots$ gives $1 = 1 + \sum_{n \geq 1} \mathbb{P}(\emptyset)$, hence $\mathbb{P}(\emptyset) = 0$.
(ii) If $A \subseteq B$, then $B = A \sqcup (B \setminus A)$, so $\mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \geq \mathbb{P}(A)$.
(iii) Since $\emptyset \subseteq A \subseteq \Omega$, (i) and (ii) give $0 = \mathbb{P}(\emptyset) \leq \mathbb{P}(A) \leq \mathbb{P}(\Omega) = 1$.
(iv) Since $\Omega = A \sqcup A^c$, $1 = \mathbb{P}(\Omega) = \mathbb{P}(A) + \mathbb{P}(A^c)$, so $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$.
(v) If $A \cap B = \emptyset$, then $A \cup B = A \sqcup B$, so $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$.
(vi) Decompose $A \cup B = (A \setminus B) \sqcup (A \cap B) \sqcup (B \setminus A)$, $A = (A \setminus B) \sqcup (A \cap B)$, and $B = (A \cap B) \sqcup (B \setminus A)$. Hence $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$.

*Structural Commentary*: The algebraic identities are set partitions in disguise.
*Key Insight*: Split overlapping sets into disjoint pieces before adding probabilities. □

**Example 1.1.8.** *Two Coin Tosses and Inclusion-Exclusion*
Let $\Omega = \{HH, HT, TH, TT\}$ with $\mathbb{P}(\{\omega\}) = \frac{1}{4}$ for all $\omega \in \Omega$. Set $H_1 = \{HH, HT\}$ and $H_2 = \{HH, TH\}$. Since $H_1 \cap H_2 = \{HH\}$, $\mathbb{P}(H_1 \cup H_2) = \mathbb{P}(H_1) + \mathbb{P}(H_2) - \mathbb{P}(H_1 \cap H_2) = \frac{1}{2} + \frac{1}{2} - \frac{1}{4} = \frac{3}{4}$.

**Proposition 1.1.9.** *Three-Set Inclusion-Exclusion and Bonferroni*
*For events $A, B, C \in \mathcal{F}$:*

*(i)* $\mathbb{P}(A \cup B \cup C) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) - \mathbb{P}(A \cap B) - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C) + \mathbb{P}(A \cap B \cap C)$.
*(ii)* $\mathbb{P}(A \cap B) \geq \mathbb{P}(A) + \mathbb{P}(B) - 1$.

*Proof.*

(i) Let $U = A \cup B$. Then $A \cup B \cup C = U \cup C$ and $U \cap C = (A \cap C) \cup (B \cap C)$.
By two-set inclusion-exclusion, $\mathbb{P}(U \cup C) = \mathbb{P}(U) + \mathbb{P}(C) - \mathbb{P}(U \cap C)$, $\mathbb{P}(U) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$, and $\mathbb{P}(U \cap C) = \mathbb{P}(A \cap C) + \mathbb{P}(B \cap C) - \mathbb{P}(A \cap B \cap C)$. Substitution gives (i).
(ii) Since $\mathbb{P}(A \cup B) \leq 1$ and $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$, rearrange.

*Structural Commentary*: Inclusion-exclusion alternates between overcounting and correcting overlaps.
*Key Insight*: Bonferroni inequalities are one-sided forms of inclusion-exclusion. □

**Theorem 1.1.10.** *Continuity of Probabilities*

*(i) Continuity from below: if $A_n \uparrow A$, then* $\lim_{n \to \infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.
*(ii) Continuity from above: if $A_n \downarrow A$, then* $\lim_{n \to \infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

*Proof.*

(i) Set $B_1 = A_1$ and $B_n = A_n \setminus A_{n-1}$ for $n \geq 2$. Then $(B_n)$ is disjoint, $A_n = \bigcup_{j=1}^{n} B_j$, and $A = \bigcup_{j=1}^{\infty} B_j$, so countable additivity gives $\mathbb{P}(A_n) = \sum_{j=1}^{n} \mathbb{P}(B_j) \to \sum_{j=1}^{\infty} \mathbb{P}(B_j) = \mathbb{P}(A)$.

(ii) Set $C_n = A_1 \setminus A_n$. Then $C_n \uparrow A_1 \setminus A$, so (i) gives $\mathbb{P}(C_n) \to \mathbb{P}(A_1 \setminus A)$; since $A_1 = A_n \sqcup C_n$ and $A_1 = A \sqcup (A_1 \setminus A)$, finite additivity gives $\mathbb{P}(A_n) = \mathbb{P}(A_1) - \mathbb{P}(C_n) \to \mathbb{P}(A_1) - \mathbb{P}(A_1 \setminus A) = \mathbb{P}(A)$.

*Structural Commentary*: Monotone limits convert countable additivity into limit rules.
*Key Insight*: Increasing uses increments; decreasing uses complements. □

**Definition 1.1.11.** *Events Occurring Infinitely Often and Eventually Always*
For events $(A_n)_{n\geq 1}$:

(i) *Tail union*: $B_n := \bigcup_{i=n}^{\infty} A_i$.

(ii) *Tail intersection*: $C_n := \bigcap_{i=n}^{\infty} A_i$.

(iii) *Limsup*: $\limsup_{n\to\infty} A_n := \bigcap_{n=1}^{\infty} B_n = \bigcap_{n=1}^{\infty} \bigcup_{i=n}^{\infty} A_i$.

(iv) *Liminf*: $\liminf_{n\to\infty} A_n := \bigcup_{n=1}^{\infty} C_n = \bigcup_{n=1}^{\infty} \bigcap_{i=n}^{\infty} A_i$.

(v) *Infinitely often*: $\omega \in \limsup_n A_n$ iff $\omega \in A_n$ for infinitely many $n$.

(vi) *Eventually always*: $\omega \in \liminf_n A_n$ iff $\omega \in A_n$ for all but finitely many $n$.

**Proposition 1.1.12.** *Tail Monotonicity*
*For events $(A_n)_{n\geq 1}$ with tail union $B_n$ and tail intersection $C_n$:*

*(i) Tail unions: $B_{n+1} \subseteq B_n$ for all $n$, hence $B_n \downarrow \bigcap_{n\geq 1} B_n$.*

*(ii) Tail intersections: $C_n \subseteq C_{n+1}$ for all $n$, hence $C_n \uparrow \bigcup_{n\geq 1} C_n$.*

*Proof.*

(i) Since $\{i : i \geq n+1\} \subseteq \{i : i \geq n\}$, $B_{n+1} = \bigcup_{i=n+1}^{\infty} A_i \subseteq \bigcup_{i=n}^{\infty} A_i = B_n$.

(ii) Since $\{i : i \geq n+1\} \subseteq \{i : i \geq n\}$, $C_n = \bigcap_{i=n}^{\infty} A_i \subseteq \bigcap_{i=n+1}^{\infty} A_i = C_{n+1}$.

*Structural Commentary*: Tail shifts turn recurrence and permanence into monotone limits.
*Key Insight*: Tail unions track recurrence; tail intersections track permanence. □

**Proposition 1.1.13.** *Union Bound and Countable Intersections of Almost-Sure Events*
*For events $(A_n)_{n\geq 1}$:*

*(i) Union bound: $\mathbb{P}(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} \mathbb{P}(A_n)$.*

*(ii) Countable almost-sure intersection: if $\mathbb{P}(A_n) = 1$ for all $n$, then $\mathbb{P}(\bigcap_{n=1}^{\infty} A_n) = 1$.*

*Proof.*

(i) Set $B_n := A_n \setminus \bigcup_{j<n} A_j$. Then $(B_n)$ is disjoint, $\bigcup_n B_n = \bigcup_n A_n$, and $B_n \subseteq A_n$.
Hence $\mathbb{P}(\bigcup_n A_n) = \sum_n \mathbb{P}(B_n) \leq \sum_n \mathbb{P}(A_n)$.

(ii) If $\mathbb{P}(A_n) = 1$ for all $n$, then $\mathbb{P}(A_n^c) = 0$ for all $n$.
By (i), $\mathbb{P}((\bigcap_n A_n)^c) = \mathbb{P}(\bigcup_n A_n^c) \leq \sum_n \mathbb{P}(A_n^c) = 0$, so $\mathbb{P}(\bigcap_n A_n) = 1$.

*Structural Commentary*: Countable subadditivity is countable additivity after disjointization.
*Key Insight*: Almost sure statements survive countable intersections. □

## 1.2 Finite Probability, Counting, and Independence

**Definition 1.2.1.** *Uniform Probability on a Countable Space*
Let $\Omega$ be countable.

(i) A probability measure $\mathbb{P}$ on $(\Omega, \mathcal{P}(\Omega))$ is *uniform* if $\exists c \in [0,1]$ such that $\mathbb{P}(\{\omega\}) = c$ for all $\omega \in \Omega$.

(ii) The constant $c$ is the *common atom mass*.

**Proposition 1.2.2.** *No Uniform Probability on a Countably Infinite Set*
*Let $\Omega = \{0, 1, 2, \ldots\}$. No uniform probability measure exists on $(\Omega, \mathcal{P}(\Omega))$.*

*Proof.* Assume uniform $\mathbb{P}$ exists, and let $c$ be its common atom mass.
Since $\Omega = \bigsqcup_{n=0}^{\infty} \{n\}$, countable additivity gives $1 = \mathbb{P}(\Omega) = \sum_{n=0}^{\infty} \mathbb{P}(\{n\}) = \sum_{n=0}^{\infty} c$.
If $c = 0$, the sum is 0; if $c > 0$, the sum diverges. Both contradict $1 = \mathbb{P}(\Omega)$.
*Structural Commentary*: Countable additivity forces total mass to equal the sum of the atom masses.
*Key Insight*: Equal mass on countably many atoms is incompatible with total mass 1. □

**Method 1.2.3.** *Finite Uniform Probability by Counting*
Let $0 < |\Omega| < \infty$ and assume the *uniform model* $\mathbb{P}(\{\omega\}) = |\Omega|^{-1}$ for all $\omega \in \Omega$.
For $A \subseteq \Omega$, $\mathbb{P}(A) = \sum_{\omega \in A} \mathbb{P}(\{\omega\}) = \frac{|A|}{|\Omega|}$, so probability reduces to counting.

**Method 1.2.4.** *Counting Workflow for Uniform Finite Probability*
For a finite uniform sample space $\Omega$ and an event $A \subseteq \Omega$:

(i) choose complete outcomes so that each $\omega \in \Omega$ is equally likely;
(ii) count $|\Omega|$; and count $|A|$, usually by products, complements, or disjoint cases;
(iii) conclude $\mathbb{P}(A) = |A| \,/\, |\Omega|$.

*Key Insight*: The hard step is often choosing equally likely outcomes, not applying the ratio.

**Example 1.2.5.** *Simpson's Paradox in Two Urn Games*
Let $p(U)$ be the red probability in urn $U$, and let $B_i, W_i$ be the black and white urns in game $i$.
Within games, $p(B_1) = 5/11 > 3/7 = p(W_1)$ and $p(B_2) = 6/9 > 9/14 = p(W_2)$.
After pooling by color into $B$ and $W$, $p(B) = 11/20 < 12/21 = p(W)$, so the aggregate winner is white.
*Key Insight*: A within-group advantage can reverse after aggregation when the group sizes change.

**Example 1.2.6.** *Complement Counting in Uniform Models*

(i) For plates $\Omega = L^3 \times D^3$ with $|L| = 26$ and $|D| = 10$, $|\Omega| = 26^3 \cdot 10^3$. If $E$ is the event of no repeated letters and no repeated digits, then $|E| = (26)_3 (10)_3$, so $\mathbb{P}(E) = \frac{(26)_3 (10)_3}{26^3 \cdot 10^3}$.
(ii) For $1 \le n \le 365$ iid uniform birthdays, take $\Omega = \{1, \ldots, 365\}^n$. If $C$ is the event of a shared birthday, then $C^c = D_n(\{1, \ldots, 365\})$, so $\mathbb{P}(C) = 1 - \frac{(365)_n}{365^n}$; for $n = 23$, $\mathbb{P}(C) > 1/2$.

*Key Insight*: "At least one collision" is often easier to count by subtracting the no-collision event.

**Definition 1.2.7.** *Independent Events*

(i) Events $A, B \in \mathcal{F}$ are *independent*, written $A \perp\!\!\!\perp B$, if $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.
(ii) A family $(A_i)_{i \in I} \subseteq \mathcal{F}$ is *pairwise independent* if $A_i$ and $A_j$ are independent whenever $i \neq j$.
(iii) A family $(A_i)_{i \in I} \subseteq \mathcal{F}$ is *mutually independent* if $\mathbb{P}(\bigcap_{i \in J} A_i) = \prod_{i \in J} \mathbb{P}(A_i)$ for every finite $J \subseteq I$.

*Key Insight*: Mutual independence means multiplicativity for every finite subfamily, not just pairs.

**Example 1.2.8.** *Pairwise Independence Need Not Be Mutual*
Let $\Omega = \{HH, HT, TH, TT\}$ with $\mathbb{P}(\{\omega\}) = \frac{1}{4}$ $\forall \omega \in \Omega$.
Set $A = \{HH, HT\}$, $B = \{HH, TH\}$, $C = \{HT, TH\}$.
Then $\mathbb{P}(A) = \mathbb{P}(B) = \mathbb{P}(C) = \frac{1}{2}$, $\mathbb{P}(A \cap B) = \mathbb{P}(A \cap C) = \mathbb{P}(B \cap C) = \frac{1}{4}$, so $A, B, C$ pairwise independent.
But $A \cap B \cap C = \emptyset$, so $\mathbb{P}(A \cap B \cap C) = 0 \neq \frac{1}{8} = \mathbb{P}(A)\mathbb{P}(B)\mathbb{P}(C)$.
*Key Insight*: Pairwise checks do not control higher-order intersections.

**Proposition 1.2.9.** *Basic Independence Tests*
*Let $A, B \in \mathcal{F}$.*

*(i) If $A \cap B = \emptyset$ and $\mathbb{P}(A)\mathbb{P}(B) > 0$, then $A \not\perp\!\!\!\perp B$.*
*(ii) If $A \perp\!\!\!\perp B$, then $A^c \perp\!\!\!\perp B$, $A \perp\!\!\!\perp B^c$, and $A^c \perp\!\!\!\perp B^c$.*
*(iii) If $\mathbb{P}(A) \in \{0, 1\}$, then $A \perp\!\!\!\perp C$ for every $C \in \mathcal{F}$.*
*(iv) If $A \perp\!\!\!\perp A$, then $\mathbb{P}(A) \in \{0, 1\}$.*

*Proof.*

(i) Since $A \cap B = \emptyset$, $\mathbb{P}(A \cap B) = 0 \neq \mathbb{P}(A)\mathbb{P}(B)$.
(ii) If $A \perp\!\!\!\perp B$, then $\mathbb{P}(A^c \cap B) = \mathbb{P}(B) - \mathbb{P}(A \cap B) = (1 - \mathbb{P}(A))\mathbb{P}(B) = \mathbb{P}(A^c)\mathbb{P}(B)$
and $\mathbb{P}(A \cap B^c) = \mathbb{P}(A) - \mathbb{P}(A \cap B) = \mathbb{P}(A)(1 - \mathbb{P}(B)) = \mathbb{P}(A)\mathbb{P}(B^c)$.
Applying the first identity to $A \perp\!\!\!\perp B^c$ gives $A^c \perp\!\!\!\perp B^c$.

(iii) Let $C \in \mathcal{F}$. If $\mathbb{P}(A) = 0$, then $0 \le \mathbb{P}(A \cap C) \le \mathbb{P}(A) = 0$, so $\mathbb{P}(A \cap C) = 0 = \mathbb{P}(A)\mathbb{P}(C)$.
If $\mathbb{P}(A) = 1$, then $\mathbb{P}(A^c) = 0$, hence $A^c \perp\!\!\!\perp C$ by the previous case and $A \perp\!\!\!\perp C$ by (ii).
(iv) Let $p = \mathbb{P}(A)$. From $A \perp\!\!\!\perp A$, $p = \mathbb{P}(A \cap A) = p^2$, hence $p \in \{0, 1\}$.

*Structural Commentary*: Independence is probabilistic, not visual; disjointness usually means strong dependence.
*Key Insight*: Complements preserve independence because $B$ splits into $A \cap B$ and $A^c \cap B$. □

**Method 1.2.10.** *Independence, Complement, and Decomposition Methods*

(i) *Verify independence*: compute $\mathbb{P}(A)$, $\mathbb{P}(B)$, and $\mathbb{P}(A \cap B)$; conclude $A \perp\!\!\!\perp B$ iff $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.
(ii) *Use complements*: if $A^c$ is simpler, compute $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$.
If $A^c = \bigcap_{j=1}^{n} C_j$ with independent $C_j$, then $\mathbb{P}(A) = 1 - \prod_{j=1}^{n} \mathbb{P}(C_j)$.
(iii) *Disjoint decomposition*: write $E = \bigsqcup_j A_j$ with disjoint $A_j$, then $\mathbb{P}(E) = \sum_j \mathbb{P}(A_j)$.
(iv) *Geometric summation*: if $\mathbb{P}(A_j) = ar^{j-1}$ for $j \ge 1$ and $|r| < 1$, then $\mathbb{P}(E) = \sum_{j \ge 1} ar^{j-1} = \frac{a}{1-r}$.

**Example 1.2.11.** *Independence, Complements, and First Success*

(i) *Independence by counting*: for a fair die, $\Omega = \{1, \dots, 6\}$, $A = \{2, 4, 6\}$, and $B = \{1, 2, 3, 4\}$.
Then $\mathbb{P}(A \cap B) = \frac{2}{6} = (\frac{3}{6})(\frac{4}{6}) = \mathbb{P}(A)\mathbb{P}(B)$, hence $A \perp\!\!\!\perp B$.
(ii) *Complement method*: for independent fair tosses $X_1, \dots, X_{10}$, let $A = \{\exists j \le 10 : X_j = H\}$ and $T_j = \{X_j = T\}$. Then $A^c = \bigcap_{j=1}^{10} T_j$ and $\mathbb{P}(A) = 1 - \mathbb{P}(A^c) = 1 - \prod_{j=1}^{10} \mathbb{P}(T_j) = 1 - (\frac{1}{2})^{10}$.
(iii) *First-success decomposition*: let $p_1 = \frac{1}{3}$, $p_2 = \frac{1}{4}$.
Let $A_j$ be player 1's first success on turn $j$ before player 2 succeeds.
Then $E = \bigsqcup_{j \ge 1} A_j$, $\mathbb{P}(A_j) = ((1 - p_1)(1 - p_2))^{j-1} p_1 = (\frac{1}{2})^{j-1}/3$, and $\mathbb{P}(E) = \sum_{j \ge 1} \mathbb{P}(A_j) = \frac{2}{3}$.

**Example 1.2.12.** *Reliability and Match Probabilities*
Let $A_i$ be the event that component $i$ works, and assume all relevant events are mutually independent.

(i) Series: if $\mathbb{P}(A_i^c) = p$ for $1 \le i \le n$ and $W_s = \bigcap_{i=1}^{n} A_i$, then $\mathbb{P}(W_s) = (1 - p)^n$ and $\mathbb{P}(W_s^c) = 1 - (1 - p)^n$.
(ii) Parallel: if $W_p = \bigcup_{i=1}^{n} A_i$, then $\mathbb{P}(W_p^c) = \mathbb{P}(\bigcap_{i=1}^{n} A_i^c) = p^n$.
(iii) DNA match: let $N_j \in \{A, C, G, T\}$ with $\mathbb{P}(N_j = x) = p_{xj}$ for $j = 1, 2$.
If $N_1 \perp\!\!\!\perp N_2$, then $\mathbb{P}(N_1 = N_2) = \sum_{x \in \{A,C,G,T\}} p_{x1} p_{x2}$.

*Key Insight*: Independence multiplies intersections; complements handle failures; matches sum disjoint cases.

## 1.3 Random Variables, Laws, and Distribution Functions

**Definition 1.3.1.** *Random Variables, Laws, and Distribution Functions*

(i) A *real-valued random variable* on $(\Omega, \mathcal{F}, \mathbb{P})$ is map $X : \Omega \to \mathbb{R}$ s.t. $\{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{F}\ \forall B \in \mathcal{B}(\mathbb{R})$.
(ii) Its *law* or *distribution* is $\mathcal{L}(X) : \mathcal{B}(\mathbb{R}) \to [0, 1]$ given by $\mathcal{L}(X)(B) := \mathbb{P}(X^{-1}(B)) = \mathbb{P}(X \in B)$.
(iii) Its *cumulative distribution function* is $F_X : \mathbb{R} \to [0, 1]$ given by $F_X(x) := \mathbb{P}(X \le x) = \mathcal{L}(X)((-\infty, x])$.

**Definition 1.3.2.** *Distribution Functions and Densities*

(i) A *distribution function* is $F : \mathbb{R} \to [0, 1]$ such that $x \le y \Rightarrow F(x) \le F(y)$, $F(x) = \lim_{t \downarrow x} F(t)$ for all $x \in \mathbb{R}$, $\lim_{x \to -\infty} F(x) = 0$, and $\lim_{x \to \infty} F(x) = 1$.
(ii) A *density* for $F$ is an integrable $g : \mathbb{R} \to [0, \infty)$ with $\int_{-\infty}^{\infty} g(x)\,dx = 1$ and $F(t) = \int_{-\infty}^{t} g(x)\,dx\ \forall t \in \mathbb{R}$.
If $g$ is continuous, then $F'(t) = g(t)$.




Figure 228: A normal density accumulates area into the distribution function $\Phi(t) = \int_{-\infty}^{t} g(x)\,dx$.

**Definition 1.3.3.** *Discrete and Continuous Random Variables*

(i) $X$ is *discrete* if $X(\Omega) \subseteq S$ for some finite or countable $S = \{x_1, x_2, \ldots\} \subseteq \mathbb{R}$.
(ii) The *probability mass function* of discrete $X$ is $f_X : S \to [0,1]$, $f_X(x) := \mathbb{P}(X = x)$, $\sum_{x \in S} f_X(x) = 1$.
(iii) The *discrete CDF* is $F_X(x) = \mathbb{P}(X \le x) = \sum_{x_i \le x} f_X(x_i)$.
(iv) $X$ is *continuous* if its law has a density $f_X : \mathbb{R} \to [0, \infty)$.
(v) The *continuous probabilities* satisfy $\mathbb{P}(a < X < b) = \int_a^b f_X(t)\,dt$ and $F_X(x) = \int_{-\infty}^x f_X(t)\,dt$.
(vi) The *point-mass distinction*: for continuous $X$, $\mathbb{P}(X = x) = 0$ for every fixed $x$, so $f_X(x) \neq \mathbb{P}(X = x)$.

A law may be mixed. It can have atoms and a continuous density on the remaining part.

**Remark 1.3.4.** *Mixed Laws Are Natural*
For a component lifetime $X$, one may have $\mathbb{P}(X = 0) = p > 0$ and, conditional on $X > 0$, a density $g$ on $(0, \infty)$. Then $\mathcal{L}(X)$ has an atom plus a continuous part, so it is mixed.

**Example 1.3.5.** *Counting Heads in Two Tosses*
Let $\Omega = \{H, T\}^2$ with $\mathbb{P}(\{\omega\}) = \frac{1}{4}$, and set $X(\omega) = |\{j \in \{1,2\} : \omega_j = H\}|$.
Then $f_X(k) = \mathbb{P}(X = k) = \binom{2}{k} 2^{-2}$ for $k \in \{0,1,2\}$, so $(f_X(0), f_X(1), f_X(2)) = (\frac{1}{4}, \frac{1}{2}, \frac{1}{4})$.
Hence $F_X(x) = 0$ for $x < 0$, $F_X(x) = \frac{1}{4}$ for $0 \le x < 1$, $F_X(x) = \frac{3}{4}$ for $1 \le x < 2$, $F_X(x) = 1$ for $x \ge 2$.
*Key Insight*: A CDF is defined for every real $x$, even when a discrete random variable takes only a few values.




Figure 229: CDF and PMF for the number of heads in two independent fair coin tosses.

**Proposition 1.3.6.** *CDF Probability Calculus*
*Let $X$ have CDF $F$ and define $F(x^-) := \lim_{y \uparrow x} F(y)$.*

*(i) Point mass: $\mathbb{P}(X = x) = F(x) - F(x^-)$.*
*(ii) Half-open interval: if $x < y$, then $\mathbb{P}(x < X \le y) = F(y) - F(x)$.*
*(iii) Right tail: $\mathbb{P}(X > x) = 1 - F(x)$.*
*(iv) Continuous endpoints: if $X$ is continuous and $a < b$, then $\mathbb{P}(a < X < b) = \mathbb{P}(a \le X < b) = \mathbb{P}(a < X \le b) = \mathbb{P}(a \le X \le b)$.*

*Proof.*

(i) Let $E_n = \{x - \frac{1}{n} < X \le x\}$. Then $E_n \downarrow \{X = x\}$ and $\mathbb{P}(E_n) = F(x) - F(x - \frac{1}{n})$. Hence $\mathbb{P}(X = x) = \lim_n \mathbb{P}(E_n) = F(x) - F(x^-)$.
(ii) Since $\{X \le y\} = \{X \le x\} \sqcup \{x < X \le y\}$, $F(y) = F(x) + \mathbb{P}(x < X \le y)$.
(iii) Since $\{X > x\} = \{X \le x\}^c$, $\mathbb{P}(X > x) = 1 - \mathbb{P}(X \le x) = 1 - F(x)$.
(iv) If $X$ is continuous, then $\mathbb{P}(X = t) = 0$ for all $t \in \mathbb{R}$. For $I_1 = (a,b)$, $I_2 = [a,b)$, $I_3 = (a,b]$, and $I_4 = [a,b]$, each $X^{-1}(I_j) \setminus X^{-1}(I_1) \subseteq \{X = a\} \cup \{X = b\}$, hence all four probabilities are equal.

*Structural Commentary*: CDF calculus subtracts nested lower-level events.
*Key Insight*: CDF jumps are point masses; continuous laws have none. □

**Definition 1.3.7.** *Survival and Hazard Functions*
Let $T \ge 0$ have CDF $F$ and density $f$, and set $D := \{t \ge 0 : 1 - F(t) > 0\}$.

(i) *Survival Function*: $S : [0, \infty) \to [0,1]$, $S(t) := \mathbb{P}(T > t) = 1 - F(t)$.
(ii) *Hazard Function*: $h : D \to [0, \infty)$, $h(t) := \frac{f(t)}{1 - F(t)} = \frac{f(t)}{S(t)}$.

*Structural Commentary*: On intervals where $S$ is differentiable and positive, $h(t) = -\frac{d}{dt} \log S(t)$.

**Proposition 1.3.8.** *Hazard Determines the Density*
*Let $T \ge 0$ have survival $S(t) := \mathbb{P}(T > t)$ with $S \in C^1([0, \infty))$, $S(t) > 0$, $S(0) = 1$, density $f(t) := -S'(t)$, and hazard $h(t) := \frac{f(t)}{S(t)}$. Put $H(t) := \int_0^t h(s)\,ds$. Then, for $t \ge 0$:*

*(i) Survival: $S(t) = e^{-H(t)}$.*

*(ii) Density:* $f(t) = h(t)e^{-H(t)}$.

*Proof.* Fix $t \geq 0$.

(i) *Survival*: Since $(\log S)' = S'/S = -h$, $\log S(t) - \log S(0) = -H(t)$. Since $S(0) = 1$, $S(t) = e^{-H(t)}$.
(ii) *Density*: Since $h = f/S$, $f(t) = h(t)S(t) = h(t)e^{-H(t)}$.

*Structural Commentary*: $h = -S'/S = -(\log S)'$, so $H$ is the cumulative log-decay of survival.
*Key Insight*: The hazard determines the law through $S = e^{-H}$ and $f = hS$. □

**Example 1.3.9.** *Constant and Power Hazards*
For $t \geq 0$:

(i) *Exponential*: If $F(t) = 1 - e^{-\lambda t}$ with $\lambda > 0$, then $S(t) = e^{-\lambda t}$, $f(t) = \lambda e^{-\lambda t}$, and $h(t) = \lambda$.
(ii) *Weibull-Type*: If $F(t) = 1 - e^{-\alpha t^\beta}$ with $\alpha, \beta > 0$, then $S(t) = e^{-\alpha t^\beta}$, $f(t) = \alpha\beta t^{\beta-1}e^{-\alpha t^\beta}$, $h(t) = \alpha\beta t^{\beta-1}$.

*Key Insight*: Power hazards increase for $\beta > 1$, decrease for $0 < \beta < 1$, and are constant for $\beta = 1$.

**Definition 1.3.10.** *Quantiles and Equality in Distribution*

(i) For a CDF $F$, the *quantile function* is $F^{-1}(q) := \inf\{x \in \mathbb{R} : F(x) \geq q\}$, $q \in (0, 1)$.
(ii) If $F$ is continuous and strictly increasing, then $F^{-1}$ is the ordinary inverse: $F(F^{-1}(q)) = q$ for $q \in (0, 1)$ and $F^{-1}(F(x)) = x$ for $x \in \mathbb{R}$.
(iii) The *pth quantile* is $x_p = F^{-1}(p)$; under (ii), $F(x_p) = p$ uniquely. Here $x_{\frac{1}{2}}$ is the median and $x_{\frac{1}{4}}, x_{\frac{3}{4}}$ are the lower and upper quartiles.
(iv) Random variables $X, Y$ are *equal in distribution*, written $X \overset{\text{dist}}{=} Y$, iff $F_X = F_Y$ on $\mathbb{R}$.

**Example 1.3.11.** *Value at Risk as a Quantile*
Let $V_0 > 0$, $R = \frac{V_1 - V_0}{V_0}$, and $F_R(x) = \mathbb{P}(R \leq x)$.
The loss is $L = V_0 - V_1 = -V_0 R$, so $\{L \geq v\} = \{R \leq -\frac{v}{V_0}\}$ and $\mathbb{P}(L \geq v) = F_R(-\frac{v}{V_0})$.
If $F_R$ is invertible and $\mathbb{P}(L \geq v_\alpha) = \alpha$, then $v_\alpha = -V_0 F_R^{-1}(\alpha)$.
*Key Insight*: VaR is a lower-tail return quantile converted back into a dollar loss.

## 1.4 Standard Distribution Families and Approximations

**Definition 1.4.1.** *Core Discrete Distributions*
Common discrete laws include:

(i) *Point mass*: $X \sim \delta_a$ for $a \in \mathbb{R}$ if $\mathbb{P}(X = a) = 1$.
(ii) *Discrete uniform*: $X \sim \text{Unif}\{1, \ldots, k\}$ for $k \in \mathbb{N}$ if $\mathbb{P}(X = x) = \frac{1}{k}$ for $x = 1, \ldots, k$.
(iii) *Bernoulli*: $X \sim \text{Bernoulli}(p)$ for $p \in [0, 1]$ if $\mathbb{P}(X = 1) = p$ and $\mathbb{P}(X = 0) = 1 - p$. Equivalently $\mathbb{P}(X = x) = p^x(1 - p)^{1-x}$ for $x \in \{0, 1\}$.
(iv) *Binomial*: $X \sim \text{Binomial}(n, p)$ for $n \in \mathbb{N}$ and $p \in [0, 1]$ if $\mathbb{P}(X = x) = \binom{n}{x}p^x(1 - p)^{n-x}$ for $x = 0, \ldots, n$.
(v) *Geometric*: $X \sim \text{Geom}(p)$, $p \in (0, 1]$, if $\mathbb{P}(X = k) = p(1 - p)^{k-1}$ for $k \geq 1$; $X$ counts trials to first success.
(vi) *Negative binomial*: $X \sim \text{NegBin}(r, p)$, $r \in \mathbb{N}$, $p \in (0, 1]$, if $\mathbb{P}(X = k) = \binom{k-1}{r-1}p^r(1 - p)^{k-r}$ for $k \geq r$; $X$ counts trials to the $r$th success.
(vii) *Hypergeometric*: $X \sim \text{Hypergeom}(n, r, m)$ if $\mathbb{P}(X = k) = \frac{\binom{r}{k}\binom{n-r}{m-k}}{\binom{n}{m}}$ for $\max\{0, m - n + r\} \leq k \leq \min\{r, m\}$.
(viii) *Poisson*: $X \sim \text{Poisson}(\lambda)$ for $\lambda \geq 0$ if $\mathbb{P}(X = x) = \frac{e^{-\lambda}\lambda^x}{x!}$ for $x = 0, 1, 2, \ldots$.

**Definition 1.4.2.** *Core Continuous Distributions*
Common continuous laws include:

(i) *Uniform*: $X \sim \text{Uniform}(a, b)$ for $a < b$ if $f_X(x) = (b - a)^{-1}\mathbf{1}_{[a,b]}(x)$.
(ii) *Normal*: $X \sim N(\mu, \sigma^2)$ for $\mu \in \mathbb{R}$, $\sigma > 0$, if $f_X(x) = (\sigma\sqrt{2\pi})^{-1}\exp(-\frac{(x-\mu)^2}{2\sigma^2})$ for $x \in \mathbb{R}$. $N(0, 1)$ has density $\phi$ and CDF $\Phi$.
(iii) *Exponential*: $X \sim \text{Exp}(\lambda)$ for $\lambda > 0$ if $f_X(x) = \lambda e^{-\lambda x}\mathbf{1}_{[0,\infty)}(x)$.
(iv) *Gamma*: $X \sim \text{Gamma}(\alpha, \lambda)$ for $\alpha, \lambda > 0$ if $f_X(x) = \lambda^\alpha\Gamma(\alpha)^{-1}x^{\alpha-1}e^{-\lambda x}\mathbf{1}_{[0,\infty)}(x)$; $\text{Exp}(\lambda) = \text{Gamma}(1, \lambda)$.
(v) *Beta*: $X \sim \text{Beta}(\alpha, \beta)$ for $\alpha, \beta > 0$ if $f_X(x) = \Gamma(\alpha + \beta)(\Gamma(\alpha)\Gamma(\beta))^{-1}x^{\alpha-1}(1 - x)^{\beta-1}\mathbf{1}_{(0,1)}(x)$.
(vi) *Cauchy*: $X \sim \text{Cauchy}(0, 1)$ if $f_X(x) = (\pi(1 + x^2))^{-1}$ for $x \in \mathbb{R}$.
(vii) $\chi^2$: $X \sim \chi^2_p$ for $p > 0$ if $f_X(x) = (\Gamma(\frac{p}{2})2^{\frac{p}{2}})^{-1}x^{\frac{p}{2}-1}e^{-\frac{x}{2}}\mathbf{1}_{(0,\infty)}(x)$.

**Definition 1.4.3.** *Normal Right-Tail Quantile*
For $0 < \alpha < 1$ and $Z \sim N(0, 1)$, $z(\alpha) := \Phi^{-1}(1 - \alpha)$, so $\mathbb{P}(Z > z(\alpha)) = \alpha$.

**Method 1.4.4.** *Discrete Distribution Model Recipes*

(i) *Indicators*: for $A \in \mathcal{F}$, $\mathbf{1}_A \sim \text{Bernoulli}(\mathbb{P}(A))$.

(ii) *Binomial counts*: if $B_1, \dots, B_n$ are independent Bernoulli($p$) variables, then $\sum_{i=1}^{n} B_i \sim \text{Binomial}(n, p)$.

(iii) *Geometric waiting*: the probability of first success on trial $k$ is $(1-p)^{k-1}p$.

(iv) *Negative-binomial waiting*: for iid $B_i \sim \text{Bernoulli}(p)$ and $T_r = \min\{k : \sum_{i=1}^{k} B_i = r\}$,
$\{T_r = k\} = \{B_k = 1, \sum_{i=1}^{k-1} B_i = r-1\}$, so $\mathbb{P}(T_r = k) = \binom{k-1}{r-1}p^r(1-p)^{k-r}$.

(v) *Hypergeometric sampling*: if $M \subseteq [n]$, $|M| = r$, and $S$ is a uniform $m$-subset,
then $X = |S \cap M|$ and $\mathbb{P}(X = k) = \frac{\binom{r}{k}\binom{n-r}{m-k}}{\binom{n}{m}}$ for $\max\{0, m-n+r\} \le k \le \min\{r, m\}$.

*Key Insight*: Name the event; PMFs come from independence or subset counts.

**Proposition 1.4.5.** *Poisson Limit of Rare Binomial Counts*
*Let $\lambda > 0$, $k \in \mathbb{N}_0$, and $X_n \sim \text{Binomial}(n, \frac{\lambda}{n})$ for all sufficiently large $n$. Then $\lim_{n\to\infty} \mathbb{P}(X_n = k) = \frac{e^{-\lambda}\lambda^k}{k!}$.*

*Proof.* For large $n$, $\mathbb{P}(X_n = k) = \binom{n}{k}(\frac{\lambda}{n})^k(1-\frac{\lambda}{n})^{n-k} = \frac{\lambda^k}{k!}\prod_{j=0}^{k-1}(1-\frac{j}{n})(1-\frac{\lambda}{n})^n(1-\frac{\lambda}{n})^{-k}$.
Since $\prod_{j=0}^{k-1}(1-\frac{j}{n}) \to 1$, $(1-\frac{\lambda}{n})^n \to e^{-\lambda}$, and $(1-\frac{\lambda}{n})^{-k} \to 1$, the limit is $\frac{e^{-\lambda}\lambda^k}{k!}$.
*Structural Commentary*: Rare binomial masses split into fixed-$k$ choices and an exponential no-event factor.
*Key Insight*: Large $n$, small $p$, fixed $np$ turns binomial masses into Poisson masses. □

**Method 1.4.6.** *Recursive Binomial and Poisson Mass Computation*
Let $q := 1 - p$.

(i) If $X \sim \text{Binomial}(n, p)$, $p_k := \mathbb{P}(X = k)$, and $q > 0$, then $p_0 = q^n$ and $\frac{p_k}{p_{k-1}} = \frac{(n-k+1)p}{kq}$ for $1 \le k \le n$.

(ii) If $X \sim \text{Poisson}(\lambda)$ and $p_k := \mathbb{P}(X = k)$, then $p_0 = e^{-\lambda}$ and $\frac{p_k}{p_{k-1}} = \frac{\lambda}{k}$ for $k \ge 1$.

*Key Insight*: Successive mass ratios avoid repeatedly evaluating large factorials.

**Example 1.4.7.** *Majority Decoding as a Binomial Tail*
Let $n = 2m+1$ and let $C_i \sim \text{Bernoulli}(p)$ be independent corruption indicators.
Then $X = \sum_{i=1}^{n} C_i \sim \text{Binomial}(n, p)$ and the majority decoder is correct iff $X \le m$.
For $n = 5$, $p = 0.1$, $\mathbb{P}(X \le 2) = 0.9^5 + 5(0.1)(0.9)^4 + 10(0.1)^2(0.9)^3 \approx 0.9914$.
*Key Insight*: Repetition coding converts reliability into a lower binomial tail.

**Example 1.4.8.** *Poisson Approximation for Rare Repeated Events*
Let $I_j = \mathbf{1}_{\{\text{double six on roll } j\}}$ for $1 \le j \le 100$.
Then $I_j \sim \text{Bernoulli}(\frac{1}{36})$ iid, $X = \sum_{j=1}^{100} I_j \sim \text{Binomial}(100, \frac{1}{36})$, and with $\lambda = \frac{100}{36} = \frac{25}{9}$, $\mathbb{P}(X = k) \approx \frac{e^{-\frac{25}{9}}(\frac{25}{9})^k}{k!}$.
*Key Insight*: Rare repeated event counts are often well approximated by a Poisson law with $\lambda = np$.

**Proposition 1.4.9.** *Exponential Memorylessness*
*If $T \sim \text{Exp}(\lambda)$ with $\lambda > 0$, then $\mathbb{P}(T > s+t \mid T > s) = \mathbb{P}(T > t)$ for all $s, t \ge 0$.*

*Proof.* For $u \ge 0$, $\mathbb{P}(T > u) = e^{-\lambda u}$. Hence $\mathbb{P}(T > s+t \mid T > s) = \frac{\mathbb{P}(T>s+t)}{\mathbb{P}(T>s)} = \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}} = e^{-\lambda t} = \mathbb{P}(T > t)$.
*Structural Commentary*: Conditioning on survival to time $s$ leaves the residual tail unchanged.
*Key Insight*: Exponential tails convert elapsed time into a cancellable factor. □

**Remark 1.4.10.** *Poisson Counts and Exponential Gaps*
In homogeneous Poisson process, counts satisfy $N(t) \sim \text{Poisson}(\lambda t)$; interarrival gaps satisfy $S_i \sim \text{Exp}(\lambda)$ iid.

**Proposition 1.4.11.** *Standardizing Normal Variables*
*For $\sigma, \sigma_i > 0$:*

*(i) Standardization: if $X \sim N(\mu, \sigma^2)$, then $\frac{X-\mu}{\sigma} \sim N(0,1)$.*

*(ii) Affine reconstruction: if $Z \sim N(0,1)$, then $\mu + \sigma Z \sim N(\mu, \sigma^2)$.*

*(iii) Independent sums: if $X_i \sim N(\mu_i, \sigma_i^2)$ are independent, then $\sum_{i=1}^{n} X_i \sim N(\sum_{i=1}^{n}\mu_i, \sum_{i=1}^{n}\sigma_i^2)$.*

*(iv) Interval probabilities: if $X \sim N(\mu, \sigma^2)$ and $a < b$, then $\mathbb{P}(a < X < b) = \Phi(\frac{b-\mu}{\sigma}) - \Phi(\frac{a-\mu}{\sigma})$.*

*Proof.*

(i) Let $Y = \frac{X-\mu}{\sigma}$. By change of variables, $f_Y(z) = \sigma f_X(\mu + \sigma z) = (2\pi)^{-\frac{1}{2}} e^{-\frac{z^2}{2}}$, so $Y \sim N(0,1)$.

(ii) Let $Y = \mu + \sigma Z$. Then $f_Y(y) = \sigma^{-1}\phi(\frac{y-\mu}{\sigma}) = (\sigma\sqrt{2\pi})^{-1}\exp(-\frac{(y-\mu)^2}{2\sigma^2})$, hence $Y \sim N(\mu, \sigma^2)$.

(iii) It suffices by induction to treat $n = 2$.
Let $X_i \sim N(\mu_i, v_i)$ be independent, $S = X_1 + X_2$, $v = v_1 + v_2$, $w = \frac{v_1 v_2}{v}$, and $a_s = \frac{v_2\mu_1 + v_1(s-\mu_2)}{v}$.
Then $f_S(s) = \int_{\mathbb{R}} f_1(u) f_2(s-u)\,du$, and completing the square gives
$f_1(u)f_2(s-u) = (2\pi v)^{-\frac{1}{2}} e^{-\frac{(s-\mu_1-\mu_2)^2}{2v}} (2\pi w)^{-\frac{1}{2}} e^{-\frac{(u-a_s)^2}{2w}}$.
The second factor integrates to 1, so $S \sim N(\mu_1+\mu_2, v_1+v_2)$; induction gives (iii).

(iv) Let $Z = \frac{X-\mu}{\sigma}$. Since $\sigma > 0$, $\{a < X < b\} = \{\frac{a-\mu}{\sigma} < Z < \frac{b-\mu}{\sigma}\}$.
Hence $\mathbb{P}(a < X < b) = \Phi(\frac{b-\mu}{\sigma}) - \Phi(\frac{a-\mu}{\sigma})$.

*Structural Commentary*: Normal laws are stable under affine maps and independent addition.
*Key Insight*: Standardize to $N(0,1)$, compute with $\Phi$, then transform back. □

**Example 1.4.12.** *Normal Tail and Quantile Standardization*
Let $X \sim N(3,5)$ and $Z = \frac{X-3}{\sqrt{5}} \sim N(0,1)$. Then $\mathbb{P}(X > 1) = \mathbb{P}(Z > -\frac{2}{\sqrt{5}}) = 1 - \Phi(-\frac{2}{\sqrt{5}}) \approx 0.81$.
If $\mathbb{P}(X < q) = 0.2$, then $\Phi(\frac{q-3}{\sqrt{5}}) = 0.2$, so $q = 3 + \sqrt{5}\,\Phi^{-1}(0.2) \approx 3 - 0.8416\sqrt{5} \approx 1.1181$.

# 2 Joint Distributions, Multivariate Laws, and Transformations

## 2.1 Joint Distributions and Independence

**Definition 2.1.1.** *Joint and Marginal Distributions*

(i) The *joint mass function* of discrete $X, Y$ is $f_{X,Y}(x,y) := \mathbb{P}(X = x, Y = y)$.

(ii) The *discrete marginals* are $f_X(x) = \sum_y f_{X,Y}(x,y)$ and $f_Y(y) = \sum_x f_{X,Y}(x,y)$.

(iii) The *continuous marginals* are $f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)\,dy$, and $f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)\,dx$.

(iv) The *joint CDF* is $F_{X,Y}(x,y) := \mathbb{P}(X \le x, Y \le y)$. $(X,Y)$ has joint density $f_{X,Y}$.

**Example 2.1.2.** *A Three-Toss Joint Mass Table*
Let $\Omega = \{H,T\}^3$ with $\mathbb{P}(\{\omega\}) = 2^{-3}$, set $X = \mathbf{1}_{\{\omega_1 = H\}}$, $Y = \sum_{i=1}^{3} \mathbf{1}_{\{\omega_i = H\}}$, and write $p_{xy} := \mathbb{P}(X = x, Y = y)$.

| | $Y=0$ | $Y=1$ | $Y=2$ | $Y=3$ |
|---|---|---|---|---|
| $X=0$ | $\frac{1}{8}$ | $\frac{2}{8}$ | $\frac{1}{8}$ | $0$ |
| $X=1$ | $0$ | $\frac{1}{8}$ | $\frac{2}{8}$ | $\frac{1}{8}$ |

Thus $p_{0,2} = \frac{1}{8}$, $\sum_y p_{0y} = \sum_y p_{1y} = \frac{1}{2}$, and $\sum_x p_{xj} = \binom{3}{j} 2^{-3}$ for $j = 0, \dots, 3$.

**Proposition 2.1.3.** *Rectangular Increments of a Joint CDF*
*Let $F(x,y) := \mathbb{P}(X \le x, Y \le y)$.*

*(i) If $x_1 < x_2$ and $y_1 < y_2$, then $\mathbb{P}(x_1 < X \le x_2,\ y_1 < Y \le y_2) = F(x_2,y_2) - F(x_1,y_2) - F(x_2,y_1) + F(x_1,y_1)$.*

*(ii) If $(X,Y)$ has density $f_{X,Y}$, $F \in C^2$, then $F(x,y) = \int_{-\infty}^{x}\int_{-\infty}^{y} f_{X,Y}(u,v)\,dv\,du$; $f_{X,Y}(x,y) = \partial_x\partial_y F(x,y)$.*

*Proof.*

(i) Set $A_{ij} := \{X \le x_i,\ Y \le y_j\}$ and $R := \{x_1 < X \le x_2,\ y_1 < Y \le y_2\}$.
Pointwise, $\mathbf{1}_R = \mathbf{1}_{A_{22}} - \mathbf{1}_{A_{12}} - \mathbf{1}_{A_{21}} + \mathbf{1}_{A_{11}}$; applying $\mathbb{E}$ gives the identity.

(ii) The density assumption gives $F(x,y) = \mathbb{P}(X \le x, Y \le y) = \int_{-\infty}^{x}\int_{-\infty}^{y} f_{X,Y}(u,v)\,dv\,du$.
Since $F \in C^2$, the two-variable FTC gives $\partial_x\partial_y F = f_{X,Y}$.

*Structural Commentary*: Joint CDFs encode southwest rectangles; finite rectangles are four-corner increments.
*Key Insight*: Mixed derivatives are infinitesimal rectangular increments. □




Figure 230: The rectangular probability is a four-term increment of the joint CDF.

**Example 2.1.4.** *A Non-Rectangular Bivariate Density*
Let $S = \{(x, y) : x^2 \le y \le 1\}$ and $f_{X,Y}(x, y) = cx^2 y \mathbf{1}_S(x, y)$.
Since $S = \{(x, y) : -1 \le x \le 1,\ x^2 \le y \le 1\}$, $1 = c \int_{-1}^{1} \int_{x^2}^{1} x^2 y \, dy \, dx = \frac{4c}{21}$, so $c = \frac{21}{4}$.
For $E = \{X \ge Y\}$, $S \cap E = \{(x, y) : 0 \le x \le 1,\ x^2 \le y \le x\}$, hence $\mathbb{P}(E) = (\frac{21}{4}) \int_0^1 \int_{x^2}^{x} x^2 y \, dy \, dx = \frac{3}{20}$.
*Key Insight*: In bivariate integration, pick one variable first and let the other range over the corresponding slice.




Figure 231: Gray: support $x^2 \le y \le 1$; blue: event $X \ge Y$ $(x^2 \le y \le x)$.

**Definition 2.1.5.** *Independent Random Variables and IID Samples*

(i) Random variables $X_1, \ldots, X_n$ are *independent* if $\mathbb{P}(X_1 \in B_1, \ldots, X_n \in B_n) = \prod_{i=1}^{n} \mathbb{P}(X_i \in B_i)\ \forall B_i \in \mathcal{B}(\mathbb{R})$.
(ii) They are *identically distributed* if $\mathcal{L}(X_1) = \cdots = \mathcal{L}(X_n)$.
(iii) They are *iid* if they are independent and identically distributed.
(iv) The *random vector* $X = (X_1, \ldots, X_n)$ is a *random sample* when $X_1, \ldots, X_n$ are iid from a common law.

**Proposition 2.1.6.** *Factorization Tests for Independence*

*(i) Discrete factorization: discrete $X, Y$ are independent iff $f_{X,Y}(x, y) = f_X(x) f_Y(y)$ for all $x, y$.*
*(ii) Density factorization: if $(X, Y)$ has joint density $f_{X,Y}$ and marginals $f_X, f_Y$, then $X, Y$ are independent iff $f_{X,Y}(x, y) = f_X(x) f_Y(y)$ a.e.*
*(iii) Rectangular support test: if $f_{X,Y}(x, y) = g(x)h(y)\mathbf{1}_{A\times B}(x, y)$ for measurable $A, B$, then $X, Y$ are independent after normalization.*

*Proof.*

(i) ($\Rightarrow$) If $X \perp\!\!\!\perp Y$, then $f_{X,Y}(x, y) = \mathbb{P}(X = x, Y = y) = \mathbb{P}(X = x)\mathbb{P}(Y = y) = f_X(x) f_Y(y)$.
(i) ($\Leftarrow$) If $f_{X,Y} = f_X f_Y$, then for $A, B$, $\mathbb{P}(X \in A, Y \in B) = \sum_{x\in A} \sum_{y\in B} f_X(x) f_Y(y) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$.
(ii) ($\Rightarrow$) If $X \perp\!\!\!\perp Y$, the product law has density $f_X f_Y$, so uniqueness of densities gives $f_{X,Y} = f_X f_Y$ a.e.
(ii) ($\Leftarrow$) If $f_{X,Y} = f_X f_Y$ a.e., then for Borel $A, B$, $\mathbb{P}(X \in A, Y \in B) = \int_A \int_B f_X(x) f_Y(y) \, dy \, dx = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$.
(iii) Let $G = \int_A g$ and $H = \int_B h$. Normalization gives $GH = 1$, while $f_X(x) = g(x)H\mathbf{1}_A(x)$ and $f_Y(y) = h(y)G\mathbf{1}_B(y)$, hence $f_X(x) f_Y(y) = f_{X,Y}(x, y)$ a.e.; apply (ii).

*Structural Commentary*: Independence is product structure of the joint law.
*Key Insight*: Independence means the joint law factors. □

## 2.2 Multivariate Models and Copulas

**Definition 2.2.1.** *Two Multivariate Families*

(i) *Multinomial*: for $n \in \mathbb{N}$ and $p \in [0, 1]^k$ with $\sum_{j=1}^{k} p_j = 1$, $X \sim \text{Multinomial}(n, p)$

if $\mathbb{P}(X=x) = n!(x_1!\cdots x_k!)^{-1}\prod_{j=1}^{k} p_j^{x_j}$ for $x \in \mathbb{N}_0^k$ with $\sum_{j=1}^{k} x_j = n$; then $X_j \sim \text{Binomial}(n, p_j)$.

(ii) *Multivariate normal*: for $\mu \in \mathbb{R}^k$ and symmetric positive definite $\Sigma \in \mathbb{R}^{k\times k}$, $X \sim N(\mu, \Sigma)$ if $f_X(x) = ((2\pi)^k \det\Sigma)^{-\frac{1}{2}} \exp(-\frac{(x-\mu)^t\Sigma^{-1}(x-\mu)}{2})$ for $x \in \mathbb{R}^k$.
$N(0, I_k)$ has independent $N(0,1)$ coordinates.

**Proposition 2.2.2.** *Multivariate Normal Affine Standardization*
*Let $k \in \mathbb{N}$, $\mu \in \mathbb{R}^k$, $\Sigma \in \mathbb{R}^{k\times k}$ with $\Sigma \succ 0$, and $A = \Sigma^{\frac{1}{2}}$.*

*(i) For $X \in \mathbb{R}^k$, $X \sim N(\mu,\Sigma) \iff A^{-1}(X-\mu) \sim N(0, I_k)$.*
*(ii) For $Z \in \mathbb{R}^k$, $Z \sim N(0, I_k) \iff \mu + AZ \sim N(\mu, \Sigma)$.*

*Proof.*

(i) ($\Rightarrow$) Put $Z = A^{-1}(X-\mu)$. With $x = \mu + Az$, $|\det A| = (\det\Sigma)^{\frac{1}{2}}$ and $(Az)^t\Sigma^{-1}(Az) = z^tz$.
By change of variables, $f_Z(z) = f_X(\mu+Az)|\det A| = (2\pi)^{-\frac{k}{2}}\exp(-\frac{z^tz}{2})$.

(i) ($\Leftarrow$) Put $Z = A^{-1}(X-\mu) \sim N(0, I_k)$, so $X = \mu + AZ$. With $z = A^{-1}(x-\mu)$ and $|\det A^{-1}| = (\det\Sigma)^{-\frac{1}{2}}$, $f_X(x) = f_Z(A^{-1}(x-\mu))|\det A^{-1}| = ((2\pi)^k\det\Sigma)^{-\frac{1}{2}}\exp(-\frac{(x-\mu)^t\Sigma^{-1}(x-\mu)}{2})$.

(ii) ($\Rightarrow$) If $Z \sim N(0, I_k)$ and $X := \mu + AZ$, then $A^{-1}(X-\mu) = Z$; by (i) ($\Leftarrow$), $X \sim N(\mu,\Sigma)$.

(ii) ($\Leftarrow$) If $X := \mu + AZ \sim N(\mu,\Sigma)$, then $Z = A^{-1}(X-\mu)$; by (i) ($\Rightarrow$), $Z \sim N(0, I_k)$.

*Structural Commentary*: Positive definite covariance is an affine deformation of the standard normal.
*Key Insight*: Standardize by $\Sigma^{-\frac{1}{2}}$; reconstruct by $\Sigma^{\frac{1}{2}}$. □

**Proposition 2.2.3.** *Basic Multivariate Normal Calculus*
*Let $X = (X_a, X_b) \sim N(\mu,\Sigma)$ with $X_a \in \mathbb{R}^r$, $X_b \in \mathbb{R}^s$, $\mu = (\mu_a, \mu_b)$, and $\Sigma = \begin{pmatrix}\Sigma_{aa} & \Sigma_{ab}\\ \Sigma_{ba} & \Sigma_{bb}\end{pmatrix} \succ 0$.*

*(i) Marginal: $X_a \sim N(\mu_a, \Sigma_{aa})$.*
*(ii) Conditional: $X_b \mid X_a = x_a \sim N(\mu_b + \Sigma_{ba}\Sigma_{aa}^{-1}(x_a-\mu_a), \Sigma_{bb} - \Sigma_{ba}\Sigma_{aa}^{-1}\Sigma_{ab})$.*
*(iii) Linear form: for any vector $a$, $a^tX \sim N(a^t\mu, a^t\Sigma a)$.*

*Proof.*

(i) Let $P_aX = X_a$. By Proposition 2.2.2, $X = \mu + \Sigma^{\frac{1}{2}}Z$ with $Z \sim N(0, I)$.
Hence $X_a = P_a\mu + P_a\Sigma^{\frac{1}{2}}Z \sim N(\mu_a, P_a\Sigma P_a^t) = N(\mu_a, \Sigma_{aa})$.

(ii) Set $S := \Sigma_{bb} - \Sigma_{ba}\Sigma_{aa}^{-1}\Sigma_{ab}$ and $m(x_a) := \mu_b + \Sigma_{ba}\Sigma_{aa}^{-1}(x_a-\mu_a)$.
The block determinant and inverse identities give $\det\Sigma = \det\Sigma_{aa}\det S$ and
$(x-\mu)^t\Sigma^{-1}(x-\mu) = (x_a-\mu_a)^t\Sigma_{aa}^{-1}(x_a-\mu_a) + (x_b-m(x_a))^tS^{-1}(x_b-m(x_a))$.
Thus $f_{b|a}(x_b \mid x_a) = \frac{f_{a,b}(x_a,x_b)}{f_a(x_a)}$ is the $N(m(x_a), S)$ density.

(iii) With $X = \mu + \Sigma^{\frac{1}{2}}Z$, $a^tX = a^t\mu + (\Sigma^{\frac{1}{2}}a)^tZ$.
Since $Z$ has independent $N(0,1)$ coordinates, $(\Sigma^{\frac{1}{2}}a)^tZ \sim N(0, a^t\Sigma a)$, hence $a^tX \sim N(a^t\mu, a^t\Sigma a)$.

*Structural Commentary*: Marginals, conditionals, and linear projections of Gaussian vectors remain Gaussian.
*Key Insight*: Gaussian calculus is affine calculus plus Schur complements. □

**Definition 2.2.4.** *Copulas*

(i) A *copula* is a joint CDF $C : [0,1]^2 \to [0,1]$.
(ii) *Uniform marginals*: $C(u,1) = u$ and $C(1,v) = v$ for $u, v \in [0,1]$.
(iii) If $C \in C^2$, its *copula density* is $c(u,v) := \partial_u\partial_v C(u,v)$.

**Proposition 2.2.5.** *Constructing a Joint Law from Marginals and a Copula*
*Let $F_X, F_Y$ be continuous CDFs with densities $f_X, f_Y$, and let $C$ be a copula.*
*Define $F_{X,Y}(x,y) := C(F_X(x), F_Y(y))$. Then $F_{X,Y}$ has marginals $F_X, F_Y$.*
*If $C \in C^2$ with density $c = \partial_u\partial_v C$, then $f_{X,Y}(x,y) = c(F_X(x), F_Y(y))f_X(x)f_Y(y)$.*

*Proof.* $F_{X,Y}(x,\infty) = C(F_X(x),1) = F_X(x)$ and $F_{X,Y}(\infty,y) = C(1, F_Y(y)) = F_Y(y)$.
If densities exist, $\partial_x\partial_y C(F_X(x), F_Y(y)) = (\partial_u\partial_v C)(F_X(x), F_Y(y))f_X(x)f_Y(y)$ by the chain rule.
*Structural Commentary*: Marginals and dependence are separated into $F_X, F_Y$ and $C$.
*Key Insight*: Marginals alone do not determine a joint law; the copula supplies the dependence. □

**Example 2.2.6.** *Farlie–Morgenstern Copulas*
For $|\alpha| \le 1$, $C_\alpha(u,v) = uv\{1+\alpha(1-u)(1-v)\}$ is a copula with density $c_\alpha(u,v) = 1+\alpha(1-2u)(1-2v)$.
Here $\alpha = 0$ gives the independence copula $uv$, while $\alpha = -1$ and $\alpha = 1$ give densities $2u+2v-4uv$ and $2-2u-2v+4uv$ on $[0,1]^2$, respectively; all have uniform marginals.

## 2.3 Transformations of Random Variables

**Method 2.3.1.** *One-Variable Transformations*
Let $Y = r(X)$.

(i) *Discrete pushforward*: if $X$ is discrete, then $f_Y(y) = \mathbb{P}(Y = y) = \mathbb{P}(X \in r^{-1}(\{y\})) = \sum_{x:r(x)=y} f_X(x)$.

(ii) *CDF method*: for $Y = r(X)$, compute $F_Y(y) = \int_{\{x:r(x)\le y\}} f_X(x)\,dx$; $f_Y(y) = F_Y'(y)$ where differentiable.

(iii) *Monotone change*: if $r$ is strictly monotone on $\operatorname{supp} X$ with inverse $s$, then $f_Y(y) = f_X(s(y))\,|s'(y)|\,\mathbf{1}_{r(\operatorname{supp} X)}(y)$.

*Key Insight*: The CDF method works because it translates the transformed event back into an event about $X$.

**Example 2.3.2.** *Log Transform of an Exponential Variable*
Let $X \sim \text{Exp}(1)$ and $Y = \log X$. Since $r(x) = \log x$ has inverse $s(y) = e^y$ with $s'(y) = e^y$, the change-of-variables formula gives $f_Y(y) = f_X(e^y)e^y = e^y e^{-e^y}$ for $y \in \mathbb{R}$.

**Example 2.3.3.** *A Nonmonotone Square Transform*
Let $X \sim \text{Uniform}(-1, 3)$ and $Y = X^2$. Since $f_X = \frac{1}{4}$ on $[-1, 3]$, $r^{-1}(\{y\}) \cap [-1, 3] = \{-\sqrt{y}, \sqrt{y}\}$ for $0 < y < 1$ and $\{\sqrt{y}\}$ for $1 < y < 9$. Hence $f_Y(y) = \frac{1}{4\sqrt{y}}\mathbf{1}_{(0,1)}(y) + \frac{1}{8\sqrt{y}}\mathbf{1}_{(1,9)}(y)$.
*Key Insight*: Non-one-to-one transformations require splitting the range into cases.

**Example 2.3.4.** *Squaring a Standard Normal*
Let $Z \sim N(0, 1)$ and $X = Z^2$. For $x > 0$, $F_X(x) = \mathbb{P}(|Z| \le \sqrt{x}) = \Phi(\sqrt{x}) - \Phi(-\sqrt{x})$,
so $f_X(x) = x^{-\frac{1}{2}}\phi(\sqrt{x}) = (2\pi)^{-\frac{1}{2}}x^{-\frac{1}{2}}e^{-\frac{x}{2}}$. Hence $X \sim \chi_1^2 = \text{Gamma}(\frac{1}{2}, \frac{1}{2})$ in the rate convention.

**Example 2.3.5.** *Reciprocal of a Uniform Variable*
Let $U \sim \text{Uniform}(0, 1)$ and $V = \frac{1}{U}$. Then $F_V(v) = 0$ for $v < 1$, while for $v \ge 1$, $F_V(v) = \mathbb{P}(U \ge \frac{1}{v}) = 1 - \frac{1}{v}$.
Hence $f_V(v) = v^{-2}\mathbf{1}_{[1,\infty)}(v)$.

**Proposition 2.3.6.** *Probability Integral Transform and Inverse Transform*
*Let $F : I \to (0, 1)$ be a continuous strictly increasing CDF with inverse $F^{-1} : (0, 1) \to I$.*

*(i) If $X \sim F$, then $F(X) \sim \text{Uniform}(0, 1)$.*

*(ii) If $U \sim \text{Uniform}(0, 1)$ and $Y = F^{-1}(U)$, then $F_Y = F$.*

*Proof.*

(i) For $0 < u < 1$, $\mathbb{P}(F(X) \le u) = \mathbb{P}(X \le F^{-1}(u)) = F(F^{-1}(u)) = u$, with endpoint values $0, 1$.

(ii) For $x \in I$, $\mathbb{P}(Y \le x) = \mathbb{P}(F^{-1}(U) \le x) = \mathbb{P}(U \le F(x)) = F(x)$.

*Structural Commentary*: $F$ maps its law to uniform scale; $F^{-1}$ maps uniform scale back.
*Key Insight*: Uniform random numbers become target laws by applying inverse CDFs. □

**Method 2.3.7.** *Several-Variable Transformations*
For $Z = r(X, Y)$ with joint density $f_{X,Y}$:

(i) *Sublevel set*: define $A_z := \{(x, y) : r(x, y) \le z\}$.

(ii) *CDF integral*: compute $F_Z(z) = \mathbb{P}(Z \le z) = \iint_{A_z} f_{X,Y}(x, y)\,dx\,dy$.

(iii) *Density*: differentiate where possible, $f_Z(z) = F_Z'(z)$.

**Example 2.3.8.** *Sum of Two Independent Uniforms*
Let $X_1, X_2 \sim \text{Uniform}(0, 1)$ be independent and $Y = X_1 + X_2$.
Since $f_{X_1,X_2} = \mathbf{1}_{[0,1]^2}$, $F_Y(y)$ is the area of $A_y = \{(x_1, x_2) \in [0, 1]^2 : x_1 + x_2 \le y\}$,
so $F_Y(y) = \frac{y^2}{2}\mathbf{1}_{[0,1]}(y) + (1 - \frac{(2-y)^2}{2})\mathbf{1}_{(1,2]}(y) + \mathbf{1}_{(2,\infty)}(y)$. Hence $f_Y(y) = y\mathbf{1}_{(0,1)}(y) + (2 - y)\mathbf{1}_{(1,2)}(y)$.

**Proposition 2.3.9.** *Two-Dimensional Density Change of Variables*
*Let $(X, Y)$ have density $f_{X,Y}$ on $D \subseteq \mathbb{R}^2$, and let $T : D \xrightarrow{C^1} E$ be one-to-one with inverse $h$ and $\det DT \neq 0$ on $D$. For $(U, V) = T(X, Y)$, $f_{U,V}(u, v) = f_{X,Y}(h(u, v))\,|\det Dh(u, v)|\,\mathbf{1}_E(u, v)$.*

*Proof.* For Borel $B \subseteq E$, $\mathbb{P}((U, V) \in B) = \mathbb{P}((X, Y) \in h(B))$
$= \int_{h(B)} f_{X,Y}(x, y)\,dx\,dy = \int_B f_{X,Y}(h(u, v))\,|\det Dh(u, v)|\,du\,dv$ (by Theorem 5.1.11)
*Structural Commentary*: Transforming a density means preserving probability mass while rescaling area.
*Key Insight*: The absolute Jacobian accounts for local volume distortion. □

**Example 2.3.10.** *Sums and Quotients*
Let $(X, Y)$ have joint density $f_{X,Y}$.

(i) Sum: for $S = X + Y$, $f_S(s) = \int_{\mathbb{R}} f_{X,Y}(x, s - x)\,dx$; if $X \perp\!\!\!\perp Y$, then $f_S(s) = \int_{\mathbb{R}} f_X(x)f_Y(s - x)\,dx$.

(ii) Quotient: for $Z = \frac{Y}{X}$, $f_Z(z) = \int_{\mathbb{R}} |x|\,f_{X,Y}(x, xz)\,dx$; if $X \perp\!\!\!\perp Y$, then $f_Z(z) = \int_{\mathbb{R}} |x|\,f_X(x)f_Y(xz)\,dx$.

*Key Insight*: Add one auxiliary coordinate, change variables, integrate it out.

**Example 2.3.11.** *Polar Transform of Two Standard Normals*
Let $X, Y \overset{\text{iid}}{\sim} N(0,1)$, $R = (X^2+Y^2)^{\frac{1}{2}}$, and $\Theta = \operatorname{atan2}(Y, X)$.
Polar change of variables gives $f_{R,\Theta}(r,\theta) = (2\pi)^{-1} r e^{-\frac{r^2}{2}} \mathbf{1}_{[0,\infty)}(r)\mathbf{1}_{[0,2\pi)}(\theta)$,
so $R \perp\!\!\!\perp \Theta$, $\Theta \sim \text{Uniform}(0, 2\pi)$, and $R^2 \sim \text{Exp}(\frac{1}{2})$.
Hence for iid $U_1, U_2 \sim \text{Uniform}(0,1)$, $(\sqrt{-2\log U_1}\cos(2\pi U_2), \sqrt{-2\log U_1}\sin(2\pi U_2)) \sim N(0, I_2)$.




Figure 232: In polar coordinates, a small patch has area approximately $r\,dr\,d\theta$.

# 3 Expectation and Integration in Probability

## 3.1 Expectation, Variance, and LOTUS

Expectation is the averaging operation attached to a probability law. In the notation of Section 1.1.1, if $X$ has law $F_X$, then integration against $F_X$ is the common notation that covers both sums for discrete variables and ordinary integrals for variables with densities.

**Definition 3.1.1.** *Expectation Through the Law*
Let $X$ be real-valued with distribution function $F_X$.

(i) *Expectation*: if $\int_{\mathbb{R}} |x|\, dF_X(x) < \infty$, then $\mathbb{E}(X) := \int_{\mathbb{R}} x\, dF_X(x)$.
(ii) *Discrete form*: if $X$ has support $S$, mass function $f_X$, $\mathbb{E}(X) = \sum_{x\in S} x f_X(x)$ when $\sum_{x\in S} |x| f_X(x) < \infty$.
(iii) *Density form*: if $X$ has density $f_X$, then $\mathbb{E}(X) = \int_{\mathbb{R}} x f_X(x)\, dx$ when $\int_{\mathbb{R}} |x| f_X(x)\, dx < \infty$.
(iv) *Function form*: for measurable $r : \mathbb{R} \to \mathbb{R}$, if $\int_{\mathbb{R}} |r(x)|\, dF_X(x) < \infty$, then $\mathbb{E}(r(X)) := \int_{\mathbb{R}} r(x)\, dF_X(x)$.

**Proposition 3.1.2.** *Indicators Turn Probabilities into Expectations*
*For $A \in \mathcal{F}$, let $\mathbf{1}_A : \Omega \to \{0,1\}$ be $\mathbf{1}_A(\omega) = 1$ for $\omega \in A$ and $\mathbf{1}_A(\omega) = 0$ for $\omega \notin A$. Then $\mathbb{E}(\mathbf{1}_A) = \mathbb{P}(A)$.*

*Proof.* Since $\Omega = A \sqcup A^c$ and $\mathbf{1}_A = 1$ on $A$, 0 on $A^c$, $\mathbb{E}(\mathbf{1}_A) = 1\cdot\mathbb{P}(A) + 0\cdot\mathbb{P}(A^c) = \mathbb{P}(A)$.
*Structural Commentary*: Indicators encode events as 0-1 random variables.
*Key Insight*: Probabilities are expectations of indicators. □

**Example 3.1.3.** *Expectation of the Two-Toss Head Count*
Let $\Omega = \{H,T\}^2$ with $\mathbb{P}(\{\omega\}) = \frac{1}{4}$, and let $X(\omega)$ be the number of heads in $\omega$.
Then $f_X(k) = \binom{2}{k}2^{-2}$ for $k = 0,1,2$, so $\mathbb{E}(X) = \sum_{k=0}^{2} k f_X(k) = 0\cdot\frac{1}{4} + 1\cdot\frac{1}{2} + 2\cdot\frac{1}{4} = 1$.

**Example 3.1.4.** *First Expectation Computations*

(i) If $X \sim \text{Bernoulli}(p)$, then $\mathbb{E}X = 0(1-p) + 1p = p$.
(ii) If $X \sim \text{Uniform}(-1,3)$, then $\mathbb{E}X = (\frac{1}{4})\int_{-1}^{3} x\, dx = 1$.
(iii) If $X \sim \text{Cauchy}(0,1)$, then $\mathbb{E}|X| = (\frac{2}{\pi})\int_0^\infty x(1+x^2)^{-1}\, dx = \infty$, so $\mathbb{E}X$ does not exist.

**Theorem 3.1.5.** *Expectation of a Function*
*Let $r$ be measurable and assume the displayed expectations exist.*

*(i) One variable: for real-valued $X$, $\mathbb{E}(r(X)) = \int_{\mathbb{R}} r(x)\, dF_X(x)$.*
*(ii) Several variables: for $(X,Y)$ with joint law $F_{X,Y}$, $\mathbb{E}(r(X,Y)) = \int_{\mathbb{R}^2} r(x,y)\, dF_{X,Y}(x,y)$.*

*Proof.*

(i) Since $\mathcal{L}(X) = X_*\mathbb{P}$, $\mathbb{E}(r(X)) = \int_\Omega r(X(\omega))\, d\mathbb{P}(\omega) = \int_{\mathbb{R}} r(x)\, d\mathcal{L}(X)(x) = \int_{\mathbb{R}} r(x)\, dF_X(x)$.
(ii) Let $W = (X,Y)$. Since $\mathcal{L}(W) = W_*\mathbb{P} = F_{X,Y}$,
$\mathbb{E}(r(X,Y)) = \int_\Omega r(W(\omega))\, d\mathbb{P}(\omega) = \int_{\mathbb{R}^2} r(x,y)\, d\mathcal{L}(W)(x,y) = \int_{\mathbb{R}^2} r(x,y)\, dF_{X,Y}(x,y)$.

*Structural Commentary*: Expectations of transformations integrate over the transformed law.
*Key Insight*: Pushforward laws let functions move from random variables to their values. □

**Remark 3.1.6.** *Law-Based Integration*
For measurable $r$, $\mathbb{E}(r(X))$ depends only on the law of $X$: $\mathbb{E}(r(X)) = \int_{\mathbb{R}} r(x)\,dF_X(x)$ whenever integrable. Discrete laws give $\sum_x r(x) f_X(x)$; density laws give $\int_{\mathbb{R}} r(x) f_X(x)\,dx$.

**Example 3.1.7.** *Using the Function Rule*

(i) If $X \sim \text{Uniform}(0,1)$ and $Y = e^X$, then $\mathbb{E}Y = \mathbb{E}(e^X) = \int_0^1 e^x\,dx = e - 1$.

(ii) If $X \sim \text{Uniform}(0,1)$ and $Y = \max\{X, 1-X\}$, then $\mathbb{E}Y = \int_0^{\frac{1}{2}} (1-x)\,dx + \int_{\frac{1}{2}}^1 x\,dx = \frac{3}{4}$.

(iii) If $(X,Y) \sim \text{Uniform}([0,1]^2)$ and $Z = X^2 + Y^2$, then $\mathbb{E}Z = \int_0^1 \int_0^1 (x^2+y^2)\,dx\,dy = \frac{2}{3}$.

**Definition 3.1.8.** *Variance and Covariance*

(i) The *variance* of $X$ is $\text{Var}(X) := \mathbb{E}[(X - \mathbb{E}X)^2]$, assuming $\mathbb{E}(X^2) < \infty$.
(ii) The *standard deviation* of $X$ is $\text{sd}(X) := \sqrt{\text{Var}(X)}$.
(iii) The *covariance* of $X, Y$ is $\text{Cov}(X,Y) := \mathbb{E}[(X - \mathbb{E}X)(Y - \mathbb{E}Y)]$, assuming $\mathbb{E}(X^2), \mathbb{E}(Y^2) < \infty$.

## 3.2 Moments, Sample Statistics, and Random Vectors

**Definition 3.2.1.** *Moments, Correlation, and Sample Statistics*

(i) The *kth moment* of $X$ is $\mathbb{E}(X^k)$, assuming $\mathbb{E}|X|^k < \infty$.
(ii) The *correlation* of $X, Y$ is $\text{Corr}(X,Y) := \frac{\text{Cov}(X,Y)}{\sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)}}$, assuming $\text{Var}(X), \text{Var}(Y) > 0$.
(iii) The *sample mean* of $X_1, \dots, X_n$ is $\overline{X}_n := n^{-1} \sum_{i=1}^n X_i$.
(iv) The *sample variance* of $X_1, \dots, X_n$ is $S_n^2 := (n-1)^{-1} \sum_{i=1}^n (X_i - \overline{X}_n)^2$.

**Theorem 3.2.2.** *Expectation, Variance, and Covariance Algebra*
*Assume all displayed quantities exist and $a_i, a, b \in \mathbb{R}$.*

*(i) Linearity:* $\mathbb{E}(\sum_i a_i X_i) = \sum_i a_i \mathbb{E}X_i$.
*(ii) Independent product: if $X_1, \dots, X_n$ are independent, then* $\mathbb{E}(\prod_i X_i) = \prod_i \mathbb{E}X_i$.
*(iii) Variance identities:* $\text{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}X)^2$ *and* $\text{Var}(aX+b) = a^2 \text{Var}(X)$.
*(iv) Covariance identity:* $\text{Cov}(X,Y) = \mathbb{E}(XY) - \mathbb{E}X\,\mathbb{E}Y$.
*(v) Variance of a linear combination:* $\text{Var}(\sum_i a_i X_i) = \sum_i a_i^2 \text{Var}(X_i) + 2\sum_{i<j} a_i a_j \text{Cov}(X_i, X_j)$.

*Proof.*

(i) $\mathbb{E}(\sum_i a_i X_i) = \int_\Omega \sum_i a_i X_i\,d\mathbb{P} = \sum_i a_i \int_\Omega X_i\,d\mathbb{P} = \sum_i a_i \mathbb{E}X_i$.

(ii) By independence, $\mathcal{L}(X_1, \dots, X_n) = \prod_i F_{X_i}$; $\mathbb{E}(\prod_i X_i) = \int_{\mathbb{R}^n} \prod_i x_i\,d\prod_i F_{X_i}(x_i) = \prod_i \int_{\mathbb{R}} x_i\,dF_{X_i}(x_i) = \prod_i \mathbb{E}X_i$.

(iii) $\text{Var}(X) = \mathbb{E}(X^2 - 2X\mathbb{E}X + (\mathbb{E}X)^2) = \mathbb{E}(X^2) - (\mathbb{E}X)^2$.
Also $\text{Var}(aX+b) = \mathbb{E}[(aX + b - a\mathbb{E}X - b)^2] = a^2 \text{Var}(X)$.

(iv) $\text{Cov}(X,Y) = \mathbb{E}(XY - X\mathbb{E}Y - Y\mathbb{E}X + \mathbb{E}X\mathbb{E}Y) = \mathbb{E}(XY) - \mathbb{E}X\,\mathbb{E}Y$.

(v) With $S = \sum_i a_i X_i$, (iv) gives
$\text{Var}(S) = \text{Cov}(S,S) = \sum_{i,j} a_i a_j \text{Cov}(X_i, X_j) = \sum_i a_i^2 \text{Var}(X_i) + 2\sum_{i<j} a_i a_j \text{Cov}(X_i, X_j)$.

*Structural Commentary*: Expectation is linear; variance and covariance are quadratic and bilinear consequences.
*Key Insight*: Expand products, then use linearity. □

**Example 3.2.3.** *Indicator Sums as a Method*

(i) If $X \sim \text{Binomial}(n,p)$, write $X = \sum_{i=1}^n I_i$ with independent $I_i \sim \text{Bernoulli}(p)$.
Then $\mathbb{E}X = np$ and $\text{Var}(X) = np(1-p)$.

(ii) Let $\xi_i \in \{-1, 1\}$ with $\mathbb{P}(\xi_i = -1) = p$ and $\mathbb{P}(\xi_i = 1) = 1-p$, independently, and $X_n = \sum_{i=1}^n \xi_i$.
Then $\mathbb{E}X_n = n(1-2p)$ and $\text{Var}(X_n) = 4np(1-p)$.

(iii) Let $F_n = c\prod_{i=1}^{n} M_i$ with independent $M_i \in \{2, \frac{1}{2}\}$ and $\mathbb{P}(M_i = 2) = \mathbb{P}(M_i = \frac{1}{2}) = \frac{1}{2}$.
Then $\mathbb{E}F_n = c\prod_{i=1}^{n} \mathbb{E}M_i = c(\frac{5}{4})^n$.

**Proposition 3.2.4.** *Zero Variance Means Almost Surely Constant*
*For $X \in L^2$ and $m := \mathbb{E}X$, $\mathrm{Var}(X) = 0$ iff $X = m$ a.s.*

*Proof.*
($\Rightarrow$) If $\mathrm{Var}(X) = 0$, then $\mathbb{E}[(X-m)^2] = 0$. Since $(X-m)^2 \geq 0$, $\mathbb{P}((X-m)^2 > 0) = 0$, hence $X = m$ a.s.
($\Leftarrow$) If $X = m$ a.s., then $(X-m)^2 = 0$ a.s., so $\mathrm{Var}(X) = \mathbb{E}[(X-m)^2] = 0$.
*Structural Commentary*: Variance is the mean squared distance from the mean.
*Key Insight*: Zero nonnegative expectation forces zero almost surely. □

**Proposition 3.2.5.** *Tail Integral Formula*
*If $X \geq 0$ has density $f_X$, distribution function $F_X$, and $\mathbb{E}X < \infty$,*
*then $\mathbb{E}X = \int_0^\infty \mathbb{P}(X > x)\,dx = \int_0^\infty (1 - F_X(x))\,dx$.*

*Proof.* For $b > 0$, integration by parts gives $\int_0^b x f_X(x)\,dx = -b(1-F_X(b)) + \int_0^b (1-F_X(x))\,dx$.
Since $\mathbb{E}X < \infty$, $b(1-F_X(b)) \leq \int_b^\infty x f_X(x)\,dx \to 0$.
Letting $b \to \infty$ yields $\mathbb{E}X = \int_0^\infty (1-F_X(x))\,dx = \int_0^\infty \mathbb{P}(X > x)\,dx$.
*Structural Commentary*: A nonnegative variable is integrated by summing its survival probabilities.
*Key Insight*: Expectation equals area under the tail. □

**Example 3.2.6.** *Maximum of Uniform Samples*
Let $X_1, \ldots, X_n$ be independent Uniform$(0,1)$ variables and $Y_n = \max_i X_i$.
Then $F_{Y_n}(y) = \mathbb{P}(X_1 \leq y, \ldots, X_n \leq y) = y^n$ for $0 \leq y \leq 1$.
So $\mathbb{E}Y_n = \int_0^1 (1 - F_{Y_n}(y))\,dy = \int_0^1 (1-y^n)\,dy = \frac{n}{n+1}$.

**Theorem 3.2.7.** *Mean and Variance of the Basic Sample Statistics*
*Let $X_1, \ldots, X_n$ be iid with $\mathbb{E}X_i = \mu$ and $\mathrm{Var}(X_i) = \sigma^2 < \infty$.*
*(i) Sample-mean expectation:* $\mathbb{E}\overline{X}_n = \mu$.
*(ii) Sample-mean variance:* $\mathrm{Var}(\overline{X}_n) = \frac{\sigma^2}{n}$.
*(iii) Unbiased sample variance:* $\mathbb{E}S_n^2 = \sigma^2$.

*Proof.*
(i) $\mathbb{E}\overline{X}_n = n^{-1}\sum_{i=1}^{n} \mathbb{E}X_i = n^{-1}n\mu = \mu$.
(ii) Independence gives $\mathrm{Var}(\overline{X}_n) = n^{-2}\sum_{i=1}^{n} \mathrm{Var}(X_i) = n^{-2}n\sigma^2 = \frac{\sigma^2}{n}$.
(iii) Since $\sum_{i=1}^{n}(X_i - \overline{X}_n)^2 = \sum_{i=1}^{n}(X_i-\mu)^2 - n(\overline{X}_n - \mu)^2$. Thus $\mathbb{E}[(n-1)S_n^2] = n\sigma^2 - n(\frac{\sigma^2}{n}) = (n-1)\sigma^2$.
*Structural Commentary*: Averaging preserves mean and scales variance.
*Key Insight*: Center at $\mu$, then subtract the fluctuation of $\overline{X}_n$. □

**Lemma 3.2.8.** *Expected Squared Deviation from a Sample Mean*
*Let $X_1, \ldots, X_n$ be independent with $\mathbb{E}X_i = \mu_i$ and $\mathrm{Var}(X_i) = \sigma^2$ for $1 \leq i \leq n$.*
*Set $\overline{X} := \frac{1}{n}\sum_{i=1}^{n} X_i$ and $\overline{\mu} := \frac{1}{n}\sum_{i=1}^{n} \mu_i$. Then $\mathbb{E}(X_i - \overline{X})^2 = (\mu_i - \overline{\mu})^2 + \frac{n-1}{n}\sigma^2$ for $1 \leq i \leq n$.*

*Proof.* Set $U_i := X_i - \overline{X}$. Then $\mathbb{E}U_i = \mu_i - \overline{\mu}$, $\mathrm{Var}(\overline{X}) = \frac{\sigma^2}{n}$, and $\mathrm{Cov}(X_i, \overline{X}) = \frac{\sigma^2}{n}$ by independence.
Hence $\mathrm{Var}(U_i) = \mathrm{Var}(X_i) + \mathrm{Var}(\overline{X}) - 2\,\mathrm{Cov}(X_i, \overline{X}) = \sigma^2 + \frac{\sigma^2}{n} - 2\frac{\sigma^2}{n} = \frac{n-1}{n}\sigma^2$.
Thus $\mathbb{E}U_i^2 = (\mathbb{E}U_i)^2 + \mathrm{Var}(U_i) = (\mu_i - \overline{\mu})^2 + \frac{n-1}{n}\sigma^2$.
*Structural Commentary*: $X_i - \overline{X}$ has mean $\mu_i - \overline{\mu}$ and variance $\mathrm{Var}(X_i - \overline{X}) = \frac{n-1}{n}\sigma^2$.
*Key Insight*: $\mathrm{Cov}(X_i, \overline{X}) = \frac{\sigma^2}{n}$ removes exactly one $\frac{\sigma^2}{n}$ from the raw variance $\sigma^2$. □

**Definition 3.2.9.** *Mean Vector and Covariance Matrix*
For a random vector $X = (X_1, \ldots, X_k)^t$:
(i) *Mean vector*: $\mathbb{E}X := (\mathbb{E}X_1, \ldots, \mathbb{E}X_k)^t$.
(ii) *Covariance matrix*: $\mathrm{Var}(X) := [\mathrm{Cov}(X_i, X_j)]_{i,j=1}^{k}$.

**Proposition 3.2.10.** *Linear Transformations of Random Vectors*
*Let $X \in \mathbb{R}^k$ satisfy $\mathbb{E}X = \mu$ and $\mathrm{Var}(X) = \Sigma$.*

*i. Linear form: for $a \in \mathbb{R}^k$, $\mathbb{E}(a^tX) = a^t\mu$ and $\mathrm{Var}(a^tX) = a^t\Sigma a$.*
*ii. Matrix transform: for $A \in \mathbb{R}^{m\times k}$, $\mathbb{E}(AX) = A\mu$ and $\mathrm{Var}(AX) = A\Sigma A^t$.*

*Proof.*

i. $\mathbb{E}(a^tX) = a^t\mathbb{E}X = a^t\mu$, and $\mathrm{Var}(a^tX) = \mathbb{E}[(a^t(X-\mu))^2] = a^t\mathbb{E}[(X-\mu)(X-\mu)^t]a = a^t\Sigma a$.
ii. $\mathbb{E}(AX) = A\mathbb{E}X = A\mu$, and $\mathrm{Var}(AX) = \mathbb{E}[(AX - A\mu)(AX-A\mu)^t] = A\mathbb{E}[(X-\mu)(X-\mu)^t]A^t = A\Sigma A^t$.

*Structural Commentary*: Linear maps push means forward and conjugate covariance matrices.
*Key Insight*: Center first; then move constants outside expectations. □

**Example 3.2.11.** *Multinomial Covariance Structure*
If $X \sim \mathrm{Multinomial}(n,p)$ with $p = (p_1,\dots,p_k)$, then $\mathbb{E}X = np$ and $\mathrm{Var}(X_i) = np_i(1-p_i)$.
For $i \neq j$, $X_i + X_j \sim \mathrm{Binomial}(n, p_i+p_j)$, so $n(p_i+p_j)(1-p_i-p_j) = np_i(1-p_i) + np_j(1-p_j) + 2\,\mathrm{Cov}(X_i,X_j)$.
Hence $\mathrm{Cov}(X_i,X_j) = -np_ip_j$.

## 3.3 Probability Inequalities

**Theorem 3.3.1.** *Markov and Chebyshev Inequalities*

*i. Markov: if $X \geq 0$ and $\mathbb{E}X < \infty$, then for $t > 0$, $\mathbb{P}(X \geq t) \leq \frac{\mathbb{E}X}{t}$.*
*ii. Chebyshev: if $\mu := \mathbb{E}X$ and $\sigma^2 := \mathrm{Var}(X) < \infty$, then for $t > 0$, $\mathbb{P}(|X-\mu| \geq t) \leq \frac{\sigma^2}{t^2}$.*

*Proof.*

i. Since $X \geq t\mathbf{1}_{\{X\geq t\}}$, taking expectations gives $\mathbb{E}X \geq t\mathbb{P}(X \geq t)$.
ii. Apply (i) to $Y := (X-\mu)^2 \geq 0$ with threshold $t^2$, then $\mathbb{P}(|X-\mu| \geq t) = \mathbb{P}(Y \geq t^2) \leq \frac{\mathbb{E}Y}{t^2} = \frac{\sigma^2}{t^2}$.

*Structural Commentary*: Tail bounds convert moment control into probability control.
*Key Insight*: Dominate the tail indicator by a scaled nonnegative random variable. □

**Example 3.3.2.** *A Distribution-Free Error-Rate Bound*
Let $X_1,\dots,X_n$ be independent Bernoulli($p$) error indicators and $\overline{X}_n := n^{-1}\sum_{i=1}^n X_i$. Since $\mathbb{E}\overline{X}_n = p$ and $\mathrm{Var}(\overline{X}_n) = \frac{p(1-p)}{n}$, Chebyshev gives, for $\varepsilon > 0$, $\mathbb{P}(\left|\overline{X}_n - p\right| > \varepsilon) \leq \frac{p(1-p)}{n\varepsilon^2} \leq \frac{1}{4n\varepsilon^2}$.

**Theorem 3.3.3.** *Cauchy–Schwarz and Jensen*
*If $X, Y \in L^2$, then $\mathbb{E}\,|XY| \leq \sqrt{\mathbb{E}(X^2)\mathbb{E}(Y^2)}$.*
*If $g : I \to \mathbb{R}$ convex, $X \in I$ a.s., $X, g(X) \in L^1$, then $\mathbb{E}g(X) \geq g(\mathbb{E}X)$; for concave $g$, inequality reverses.*

*Proof.* For Cauchy–Schwarz, apply the $L^2$ inner-product inequality to $|X|$ and $|Y|$.
For Jensen, set $m := \mathbb{E}X$ and take an affine support $\ell(x) := g(m) + s(x-m)$ with $\ell \leq g$ on $I$.
Then $\mathbb{E}g(X) \geq \mathbb{E}\ell(X) = g(m) + s(\mathbb{E}X - m) = g(\mathbb{E}X)$. For concave $g$, apply the convex case to $-g$.
*Structural Commentary*: Cauchy–Schwarz controls products; Jensen moves convexity through expectation.
*Key Insight*: Use Hilbert-space geometry for products and supporting lines for convex functions. □

**Theorem 3.3.4.** *Hoeffding's Bounded-Sum Bound*
*Let $Y_1,\dots,Y_n$ be independent, $\mathbb{E}Y_i = 0$, $a_i \leq Y_i \leq b_i$ a.s., and $V := \sum_{i=1}^n (b_i-a_i)^2$. For $S_n := \sum_{i=1}^n Y_i$ and $t, \varepsilon > 0$, $\mathbb{P}(S_n \geq \varepsilon) \leq \exp(-t\varepsilon + \frac{t^2V}{8})$; if $V > 0$, then $\mathbb{P}(S_n \geq \varepsilon) \leq \exp(-\frac{2\varepsilon^2}{V})$.*

*Proof.* For $t > 0$, Markov gives $\mathbb{P}(S_n \geq \varepsilon) = \mathbb{P}(e^{tS_n} \geq e^{t\varepsilon}) \leq e^{-t\varepsilon}\mathbb{E}e^{tS_n}$.
Independence gives $\mathbb{E}e^{tS_n} = \prod_{i=1}^n \mathbb{E}e^{tY_i}$, and Hoeffding's lemma gives $\mathbb{E}e^{tY_i} \leq \exp(\frac{t^2(b_i-a_i)^2}{8})$.
Hence $\mathbb{P}(S_n \geq \varepsilon) \leq \exp(-t\varepsilon + \frac{t^2V}{8})$. If $V > 0$, minimize the exponent at $t = \frac{4\varepsilon}{V}$ to obtain $\exp(-\frac{2\varepsilon^2}{V})$.
*Structural Commentary*: Hoeffding turns bounded independent increments into exponential tail decay.
*Key Insight*: Exponentiate the sum, factor the MGF, then optimize $t$. □

**Corollary 3.3.5.** *Hoeffding for Bernoulli Averages*
*If $X_1,\dots,X_n$ independent* Bernoulli($p$) *variables, $\overline{X}_n := n^{-1}\sum_{i=1}^n X_i$, then $\mathbb{P}(\left|\overline{X}_n - p\right| > \varepsilon) \leq 2e^{-2n\varepsilon^2}$.*

*Proof.* Set $Y_i^+ := \frac{X_i - p}{n}$.
Then $\mathbb{E}Y_i^+ = 0$, $Y_i^+ \in [-\frac{p}{n}, \frac{1-p}{n}]$, and $V = \sum_i (\frac{1}{n})^2 = \frac{1}{n}$, by Theorem 3.3.4, $\mathbb{P}(\overline{X}_n - p \geq \varepsilon) \leq e^{-2n\varepsilon^2}$.

Apply same argument to $Y_i^- := \frac{p - X_i}{n}$, $\mathbb{P}(p - \overline{X}_n \geq \varepsilon) \leq e^{-2n\varepsilon^2}$; the union bound yields the claim.
*Structural Commentary*: Bernoulli averages are sums of bounded centered increments.
*Key Insight*: Control both tails separately, then add them. □

## 3.4 Moment Generating Functions

**Definition 3.4.1.** *Moment Generating Function*
The *moment generating function* of $X$ is $\psi_X(t) := \mathbb{E}(e^{tX})$ on $D_X := \{t \in \mathbb{R} : \mathbb{E}e^{tX} < \infty\}$.
If $(-\delta, \delta) \subseteq D_X$ for some $\delta > 0$, then $\psi_X^{(k)}(0) = \mathbb{E}(X^k)$ for all $k \geq 1$.

**Proposition 3.4.2.** *Basic MGF Properties*
*Assume the MGFs below are finite on a common open interval about* 0.

*(i) Affine transform: if $a, b \in \mathbb{R}$ and $Y = aX + b$, then $\psi_Y(t) = e^{bt}\psi_X(at)$.*

*(ii) Independent sum: if $X_1, \ldots, X_n$ are independent and $Y := \sum_{i=1}^n X_i$, then $\psi_Y(t) = \prod_{i=1}^n \psi_{X_i}(t)$.*

*(iii) Uniqueness: if $\psi_X(t) = \psi_Y(t)$ for all $t$ in an open interval containing 0, then $\mathcal{L}(X) = \mathcal{L}(Y)$.*

*Proof.*

(i) $\psi_Y(t) = \mathbb{E}e^{t(aX+b)} = e^{bt}\mathbb{E}e^{(at)X} = e^{bt}\psi_X(at)$.

(ii) For admissible $t$, $e^{tX_1}, \ldots, e^{tX_n}$ are independent and integrable.
Hence $\psi_Y(t) = \mathbb{E}e^{t\sum_{i=1}^n X_i} = \mathbb{E}\prod_{i=1}^n e^{tX_i} = \prod_{i=1}^n \mathbb{E}e^{tX_i} = \prod_{i=1}^n \psi_{X_i}(t)$.

(iii) Let $\mu := \mathcal{L}(X)$ and $\nu := \mathcal{L}(Y)$. If $\psi_X = \psi_Y$ on $(-\delta, \delta)$, then $\int_{\mathbb{R}} e^{tx}\, d\mu(x) = \int_{\mathbb{R}} e^{tx}\, d\nu(x)$ for all $|t| < \delta$.
Uniqueness of the two-sided Laplace transform gives $\mu = \nu$.

*Structural Commentary*: MGFs turn affine maps and independent sums into algebraic operations.
*Key Insight*: Exponentials convert sums into products, and the MGF near 0 determines the law. □

**Example 3.4.3.** *MGF Computations*

(i) If $X \sim \text{Exp}(1)$, then $\psi_X(t) = \mathbb{E}e^{tX} = \int_0^\infty e^{(t-1)x}\, dx = (1-t)^{-1}$ for $t < 1$.
Hence $\mathbb{E}X = \psi_X'(0) = 1$ and $\text{Var}(X) = \psi_X''(0) - (\psi_X'(0))^2 = 1$.

(ii) If $X \sim \text{Binomial}(n, p)$, then $X = \sum_{i=1}^n B_i$ with independent $B_i \sim \text{Bernoulli}(p)$.
Hence $\psi_X(t) = \prod_{i=1}^n \psi_{B_i}(t) = (pe^t + 1 - p)^n$.

**Method 3.4.4.** *MGF Calculus for Classical Laws*
If $\psi_X(t) = \mathbb{E}e^{tX}$ exists near 0, then $\mathbb{E}X = \psi_X'(0)$ and $\text{Var}(X) = \psi_X''(0) - \psi_X'(0)^2$.

(i) If $X \sim \text{Poisson}(\lambda)$, then $\psi_X(t) = \exp(\lambda(e^t - 1))$, $\mathbb{E}X = \lambda$, and $\text{Var}(X) = \lambda$.

(ii) If $X \sim \text{Gamma}(\alpha, \lambda)$ with rate $\lambda$, then $\psi_X(t) = (\frac{\lambda}{\lambda - t})^\alpha$ for $t < \lambda$, $\mathbb{E}X = \frac{\alpha}{\lambda}$, and $\text{Var}(X) = \frac{\alpha}{\lambda^2}$.

(iii) If $Z \sim N(0,1)$, then $\psi_Z(t) = e^{\frac{t^2}{2}}$; for $Y = \mu + \sigma Z$, $\psi_Y(t) = \exp(\mu t + \frac{\sigma^2 t^2}{2})$.

**Corollary 3.4.5.** *Independent Sums from MGFs*
*Let $X \perp\!\!\!\perp Y$, and interpret all MGFs on a common neighborhood of* 0.

*(i) If $\lambda, \mu > 0$, $X \sim \text{Poisson}(\lambda)$, and $Y \sim \text{Poisson}(\mu)$, then $X + Y \sim \text{Poisson}(\lambda + \mu)$.*

*(ii) If $X \sim \text{Gamma}(\alpha_1, \lambda)$ and $Y \sim \text{Gamma}(\alpha_2, \lambda)$ with $\alpha_1, \alpha_2, \lambda > 0$, then $X + Y \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$.*

*(iii) If $\mu, \nu \in \mathbb{R}$, $\sigma, \tau > 0$, $X \sim N(\mu, \sigma^2)$, and $Y \sim N(\nu, \tau^2)$, then $X + Y \sim N(\mu + \nu, \sigma^2 + \tau^2)$.*

*Proof.*

(i) $\psi_{X+Y}(t) = \psi_X(t)\psi_Y(t) = \exp(\lambda(e^t - 1))\exp(\mu(e^t - 1)) = \exp((\lambda + \mu)(e^t - 1))$, QED.

(ii) For $t < \lambda$, $\psi_{X+Y}(t) = \psi_X(t)\psi_Y(t) = (\frac{\lambda}{\lambda - t})^{\alpha_1 + \alpha_2}$, QED.

(iii) $\psi_{X+Y}(t) = \psi_X(t)\psi_Y(t) = \exp(\mu t + \frac{\sigma^2 t^2}{2})\exp(\nu t + \frac{\tau^2 t^2}{2}) = \exp((\mu + \nu)t + \frac{(\sigma^2 + \tau^2)t^2}{2})$, QED.

*Structural Commentary*: Independence turns convolution into multiplication of MGFs.
*Key Insight*: Closed distribution families are stable when their MGF parameters add. □

**Method 3.4.6.** *Common Means, Variances, and MGFs*
MGFs are stated on their natural neighborhood of 0; values at $t = 0$ are by continuity.
The geometric and negative-binomial rows use the total-trials convention.

| Distribution | $\mathbb{E}X$ | $\mathrm{Var}(X)$ | $\psi_X(t)$ |
|---|---|---|---|
| point mass at $a$ | $a$ | $0$ | $e^{at}$ |
| Bernoulli$(p)$ | $p$ | $p(1-p)$ | $1-p+pe^t$ |
| Binomial$(n,p)$ | $np$ | $np(1-p)$ | $(1-p+pe^t)^n$ |
| Geometric$(p)$ | $\frac{1}{p}$ | $\frac{1-p}{p^2}$ | $\frac{pe^t}{1-(1-p)e^t}$ |
| NegBin$(r,p)$ | $\frac{r}{p}$ | $\frac{r(1-p)}{p^2}$ | $\left(\frac{pe^t}{1-(1-p)e^t}\right)^r$ |
| Poisson$(\lambda)$ | $\lambda$ | $\lambda$ | $e^{\lambda(e^t-1)}$ |
| Uniform$(a,b)$ | $\frac{a+b}{2}$ | $\frac{(b-a)^2}{12}$ | $\frac{e^{bt}-e^{at}}{(b-a)t}$ |
| $N(\mu,\sigma^2)$ | $\mu$ | $\sigma^2$ | $\exp(\mu t+\frac{\sigma^2t^2}{2})$ |
| Exp$(\lambda)$ | $\frac{1}{\lambda}$ | $\frac{1}{\lambda^2}$ | $\frac{\lambda}{\lambda-t}$ |
| Gamma$(\alpha,\lambda)$ | $\frac{\alpha}{\lambda}$ | $\frac{\alpha}{\lambda^2}$ | $\left(\frac{\lambda}{\lambda-t}\right)^\alpha$ |
| Beta$(\alpha,\beta)$ | $\frac{\alpha}{\alpha+\beta}$ | $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$ | not useful |
| $\chi^2_p$ | $p$ | $2p$ | $(1-2t)^{-\frac{p}{2}}$ |

# 4 Order Statistics, Extrema, and Moment Methods

## 4.1 Order Statistics, Extrema, and Moment Methods

Order statistics describe the sorted values of an independent sample. They turn joint density calculations into formulas for minima, maxima, medians, ranges, and distribution-free tolerance intervals.

**Definition 4.1.1.** *Order Statistics*
For real-valued $X_1,\dots,X_n$,

(i) The *order statistics* are the pointwise rearrangement $X_{(1)} \le X_{(2)} \le \cdots \le X_{(n)}$ of $X_1,\dots,X_n$.
(ii) $X_{(1)} = \min_{1\le i\le n} X_i$, $X_{(n)} = \max_{1\le i\le n} X_i$, and if $n = 2m+1$, then $X_{(m+1)}$ is the sample median.

**Proposition 4.1.2.** *Minimum and Maximum of an IID Continuous Sample*
*Let $n \ge 1$ and $X_1,\dots,X_n \overset{\text{iid}}{\sim} F$, where $F$ is differentiable with density $f = F'$.*
*Set $V = X_{(1)} = \min_i X_i$ and $U = X_{(n)} = \max_i X_i$; then for $u, v \in \mathbb{R}$, $F_U(u) = F(u)^n$, $f_U(u) = nf(u)F(u)^{n-1}$, $F_V(v) = 1-(1-F(v))^n$, and $f_V(v) = nf(v)(1-F(v))^{n-1}$.*

*Proof.* Since $\{U \le u\} = \bigcap_{i=1}^n \{X_i \le u\}$, independence gives $F_U(u) = \prod_{i=1}^n F(u) = F(u)^n$.
Differentiation gives $f_U(u) = nf(u)F(u)^{n-1}$.
Since $\{V > v\} = \bigcap_{i=1}^n \{X_i > v\}$, independence gives $1 - F_V(v) = \prod_{i=1}^n (1-F(v)) = (1-F(v))^n$.
Differentiation of $F_V(v) = 1-(1-F(v))^n$ gives $f_V(v) = nf(v)(1-F(v))^{n-1}$.
*Structural Commentary*: Extremes convert order events into products over independent coordinates.
*Key Insight*: Maxima use all observations to the left; minima use all observations to the right. □

**Example 4.1.3.** *Series and Parallel Exponential Lifetimes*
Let $T_1,\dots,T_n \overset{\text{ind}}{\sim} \mathrm{Exp}(\lambda)$ be component lifetimes.

(i) For a series system, all components must survive, so system failure occurs at the first component failure: $V = \min_i T_i = T_{(1)}$. Thus $F_V(v) = 1-e^{-n\lambda v}$ and $f_V(v) = n\lambda e^{-n\lambda v}$ for $v \ge 0$; hence $V \sim \mathrm{Exp}(n\lambda)$.
(ii) For a parallel system, one surviving component suffices, so system failure occurs only after all components fail: $U = \max_i T_i = T_{(n)}$. Thus $F_U(u) = (1-e^{-\lambda u})^n$ and $f_U(u) = n\lambda e^{-\lambda u}(1-e^{-\lambda u})^{n-1}$ for $u \ge 0$.

**Theorem 4.1.4.** *Density of the kth Order Statistic*
*Let $X_1,\dots,X_n$ be iid real-valued random variables with differentiable CDF $F$ and density $f = F'$.*
*For $1 \le k \le n$ and $x \in \mathbb{R}$, $f_{X_{(k)}}(x) = \frac{n!}{(k-1)!(n-k)!} f(x)F(x)^{k-1}(1-F(x))^{n-k}$.*

*Proof.* $N_x = \sum_{i=1}^{n} \mathbf{1}_{\{X_i \le x\}}$; $N_x \sim \mathrm{Bin}(n, F(x))$, $\{X_{(k)} \le x\} = \{N_x \ge k\}$; $F_{X_{(k)}}(x) = \sum_{j=k}^{n} \binom{n}{j} F(x)^j (1-F(x))^{n-j}$.
With $p = F(x)$, $\frac{d}{dp} \sum_{j=k}^{n} \binom{n}{j} p^j (1-p)^{n-j} = \frac{n!}{(k-1)!(n-k)!} p^{k-1}(1-p)^{n-k}$, and the chain rule gives stated density.
*Structural Commentary*: The multinomial count of left, middle, and right observations supplies the coefficient.
*Key Insight*: An order statistic is located by counting how many sample points fall on each side. □

**Corollary 4.1.5.** *Uniform Order Statistics Are Beta*
*If* $X_1, \dots, X_n \overset{\text{iid}}{\sim} \mathrm{Uniform}(0,1)$, *then for* $1 \le k \le n$, $X_{(k)} \sim \mathrm{Beta}(k, n-k+1)$
*with density* $f_{X_{(k)}}(x) = \frac{n!}{(k-1)!(n-k)!} x^{k-1}(1-x)^{n-k} \mathbf{1}_{(0,1)}(x)$.

*Proof.* On $(0,1)$, $F(x) = x$ and $f(x) = 1$, so Theorem 4.1.4 gives $f_{X_{(k)}}(x) = \frac{n!}{(k-1)!(n-k)!} x^{k-1}(1-x)^{n-k}$.
Also $B(k, n-k+1) = \frac{(k-1)!(n-k)!}{n!}$, hence this density is the $\mathrm{Beta}(k, n-k+1)$ density.
*Structural Commentary*: Uniform order statistics reduce the general density to the beta normalizing constant.
*Key Insight*: Counting positions in a uniform sample produces beta powers of left and right mass. □

**Proposition 4.1.6.** *Joint Density of Minimum and Maximum*
*Let* $n \ge 2$, *let* $X_1, \dots, X_n \overset{\text{iid}}{\sim} F$ *with differentiable CDF* $F$, *density* $f = F'$; $V = X_{(1)}$, $U = X_{(n)}$, $R = U - V$.

*(i) For* $v < u$, $f_{V,U}(v,u) = n(n-1)f(v)f(u)(F(u)-F(v))^{n-2}$, *and* $f_{V,U}(v,u) = 0$ *for* $v \ge u$.
*(ii) If* $F(t) = t$ *on* $[0,1]$, *then* $f_{V,U}(v,u) = n(n-1)(u-v)^{n-2}\mathbf{1}_{\{0<v<u<1\}}$.
*(iii) In the uniform case,* $f_R(r) = n(n-1)r^{n-2}(1-r)\mathbf{1}_{(0,1)}(r)$, *with* $F_R(r) = 0$ *for* $r < 0$,
$F_R(r) = nr^{n-1} - (n-1)r^n$ *for* $0 \le r \le 1$, *and* $F_R(r) = 1$ *for* $r \ge 1$.

*Proof.*

(i) For $v < u$, $G(v,u) = \mathbb{P}(V \ge v, U \le u) = \mathbb{P}(v \le X_i \le u,\ 1 \le i \le n) = (F(u) - F(v))^n$,
so $f_{V,U}(v,u) = -\partial_v \partial_u G(v,u) = n(n-1)f(v)f(u)(F(u)-F(v))^{n-2}$.
(ii) If $F(t) = t$ and $f(t) = 1$ on $(0,1)$, by (i), $f_{V,U}(v,u) = n(n-1)(u-v)^{n-2}$ on $0 < v < u < 1$, else 0.
(iii) For $0 < r < 1$, $f_R(r) = \int_0^{1-r} f_{V,U}(v, v+r)\, dv = n(n-1)r^{n-2}(1-r)$.
Integration gives $F_R(r) = \int_0^r f_R(s)\, ds = nr^{n-1} - (n-1)r^n$; outside $[0,1]$, the support gives $F_R(r) \in \{0,1\}$.

*Structural Commentary*: Mixed derivatives select the two boundary observations.
*Key Insight*: Fixed extremes leave only interior mass. □

**Example 4.1.7.** *Distribution-Free Tolerance Interval*
Let $X_1, \dots, X_n$ be iid with continuous CDF $F$, and set $Q = F(X_{(n)}) - F(X_{(1)})$.
By the probability integral transform, $U_i = F(X_i) \overset{\text{iid}}{\sim} \mathrm{Uniform}(0,1)$, so $Q \overset{d}{=} U_{(n)} - U_{(1)}$.
Therefore, for $0 \le \alpha \le 1$, $\mathbb{P}(Q > \alpha) = 1 - F_R(\alpha) = 1 - n\alpha^{n-1} + (n-1)\alpha^n$. For $n = 100$ and $\alpha = .95$, $\mathbb{P}(Q > .95) \approx .96$: $[X_{(1)}, X_{(n)}]$ covers at least .95 of the $F$-mass with probability about .96.

The expectation calculus developed above also supports a useful collection of moment methods: identifying whether the mean exists, replacing difficult distributions of functions by direct expectation, controlling spread by covariance, and using generating functions or Taylor expansion when direct integration is cumbersome.

**Example 4.1.8.** *Existence and First Distributional Means*

(i) If $X \sim \mathrm{Geom}(p)$ on $\{1, 2, \dots\}$ and $q = 1 - p$, then $\mathbb{E}X = p \sum_{k \ge 1} k q^{k-1} = p \frac{d}{dq} \frac{q}{1-q} = \frac{1}{p}$.
(ii) If $X \sim \mathrm{Poisson}(\lambda)$, then $\mathbb{E}X = e^{-\lambda} \sum_{k \ge 1} \frac{k\lambda^k}{k!} = \lambda e^{-\lambda} \sum_{j \ge 0} \frac{\lambda^j}{j!} = \lambda$.
(iii) If $X \sim \mathrm{Cauchy}(0,1)$, then $\mathbb{E}|X| = \frac{2}{\pi} \int_0^\infty x(1+x^2)^{-1}\, dx = \infty$, so $X$ is not integrable and $\mathbb{E}X$ is undefined.
(iv) If $\mathbb{P}(Y = 2^k) = 2^{-(k+1)}$ for $k \ge 0$, then $\mathbb{E}Y = \sum_{k \ge 0} 2^{-1} = \infty$; truncation is essential.

*Key Insight*: Expectation is a finite absolute average, not merely a formal symmetry or a plausible center.

**Method 4.1.9.** *Indicator Sums Without Independence*

(i) For a finite index set $I$ and events $(A_i)_{i \in I}$, if $W = \sum_{i \in I} \mathbf{1}_{A_i}$, then $\mathbb{E}W = \sum_{i \in I} \mathbb{P}(A_i)$.
(ii) The identity uses only linearity of expectation; the events $A_i$ need not be independent.

**Example 4.1.10.** *Counting Word Occurrences*
Let $S_1, \dots, S_N \overset{\text{iid}}{\sim} \mathrm{Uniform}\{A, C, G, T\}$ and fix a word $w \in \{A, C, G, T\}^q$.
For $1 \le m \le N - q + 1$, set $I_m = \mathbf{1}_{\{(S_m, \dots, S_{m+q-1}) = w\}}$ and $W = \sum_{m=1}^{N-q+1} I_m$.

Then $\mathbb{P}(I_m = 1) = 4^{-q}$ and $\mathbb{E}W = (N - q + 1)4^{-q}$.
Overlaps may create dependence; linearity does not require independence.

**Example 4.1.11.** *Coupon Collection by Waiting Times*
Draw uniformly from $n$ coupon types, and let $X_r$ be the waiting time for the $r$th new type. Then $X_r \sim \text{Geom}(\frac{n-r+1}{n})$ on $\{1, 2, \ldots\}$, so $\mathbb{E}X_r = \frac{n}{n-r+1}$.
For $T = \sum_{r=1}^{n} X_r$, linearity gives $\mathbb{E}T = \sum_{r=1}^{n} \frac{n}{n-r+1} = nH_n$, where $H_n = \sum_{j=1}^{n} \frac{1}{j} \sim \log n + \gamma$.

**Theorem 4.1.12.** *Correlation Bound and Equality Case*
*Let $X, Y \in L^2$ have means $\mu_X, \mu_Y$ and standard deviations $0 < \sigma_X, \sigma_Y < \infty$, set $\rho = \frac{\text{Cov}(X,Y)}{\sigma_X \sigma_Y}$, and define $Z_X = \frac{X-\mu_X}{\sigma_X}$, $Z_Y = \frac{Y-\mu_Y}{\sigma_Y}$.*
*(i)* $\mathbb{E}Z_X = \mathbb{E}Z_Y = 0$, $\text{Var}(Z_X) = \text{Var}(Z_Y) = 1$, *and* $\text{Cov}(Z_X, Z_Y) = \rho$.
*(ii)* $-1 \le \rho \le 1$, *equivalently* $|\rho| \le 1$.
*(iii)* $\rho = \pm 1$ *iff* $Z_Y = \pm Z_X$ *a.s., equivalently* $Y = \mu_Y \pm (\frac{\sigma_Y}{\sigma_X})(X - \mu_X)$ *a.s., with the same sign.*

*Proof.*
(i) $\mathbb{E}Z_X = 0$, $\text{Var}(Z_X) = \sigma_X^{-2} \text{Var}(X) = 1$, and similarly for $Z_Y$.
Moreover, $\text{Cov}(Z_X, Z_Y) = \sigma_X^{-1}\sigma_Y^{-1} \text{Cov}(X, Y) = \rho$.
(ii) Since $0 \le \text{Var}(Z_X \pm Z_Y) = 2 \pm 2\rho$, both $1 + \rho \ge 0$ and $1 - \rho \ge 0$, hence $|\rho| \le 1$.
*(iii)* ($\Rightarrow$) If $\rho = 1$, then $\mathbb{E}(Z_X - Z_Y)^2 = 0$, so $Z_X = Z_Y$ a.s..
If $\rho = -1$, then $\mathbb{E}(Z_X + Z_Y)^2 = 0$, so $Z_X = -Z_Y$ a.s.
*(iii)* ($\Leftarrow$) If $Z_Y = \pm Z_X$, then $\rho = \text{Cov}(Z_X, \pm Z_X) = \pm 1$, and $\frac{Y-\mu_Y}{\sigma_Y} = \pm\frac{X-\mu_X}{\sigma_X}$
iff $Y = \mu_Y \pm (\frac{\sigma_Y}{\sigma_X})(X - \mu_X)$ a.s.

*Structural Commentary*: Correlation is centred covariance after unit-variance normalization.
*Key Insight*: Perfect correlation means no residual scatter around an affine line. □

**Example 4.1.13.** *Bivariate Uniform Association*
For $(X, Y)$ with uniform marginals on $[0, 1]$, $\mathbb{E}X = \mathbb{E}Y = \frac{1}{2}$ and $\text{Var}(X) = \text{Var}(Y) = \frac{1}{12}$.
Hence $\text{Cov}(X, Y) = \mathbb{E}(XY) - \frac{1}{4}$ and $\rho = 12 \text{Cov}(X, Y)$.
If $\mathbb{E}(XY) = \frac{2}{9}$, then $\text{Cov}(X, Y) = -\frac{1}{36}$, $\rho = -\frac{1}{3}$; if $\mathbb{E}(XY) = \frac{5}{18}$, then $\text{Cov}(X, Y) = \frac{1}{36}$, $\rho = \frac{1}{3}$.
The sign of $\rho$ records negative or positive association.

**Method 4.1.14.** *Moment Propagation by Taylor Expansion*
Let the relevant variables have finite second moments, and let $g$ be $C^2$ near the relevant mean point.
(i) If $Y = g(X)$, $\mu = \mathbb{E}X$, and $\sigma^2 = \text{Var}(X)$, then $\mathbb{E}Y \approx g(\mu)$ and $\text{Var}(Y) \approx [g'(\mu)]^2\sigma^2$.
(ii) The second-order mean correction is $\mathbb{E}g(X) \approx g(\mu) + \frac{1}{2}g''(\mu)\sigma^2$.
(iii) For $Z = g(\mathbf{X})$, $\mu = \mathbb{E}\mathbf{X}$, and $\Sigma = \text{Cov}(\mathbf{X})$, $\text{Var}(Z) \approx \nabla g(\mu)^\top \Sigma \nabla g(\mu)$ and $\mathbb{E}Z \approx g(\mu) + \frac{1}{2}\text{tr}(\Sigma H_g(\mu))$.

*Structural Commentary*: Taylor expansion propagates moments through local derivatives.
*Key Insight*: Smooth nonlinear functions are locally linear for variance and locally quadratic for mean correction.

**Example 4.1.15.** *Approximate Moments of a Ratio*
Let $Z = \frac{Y}{X}$, $\mu_X \neq 0$, $\mu = (\mu_X, \mu_Y)$, and write $\sigma_X^2 = \text{Var}(X)$, $\sigma_Y^2 = \text{Var}(Y)$, $\sigma_{XY} = \text{Cov}(X, Y)$.
For $g(x, y) = \frac{y}{x}$, Taylor propagation gives $\mathbb{E}Z \approx \frac{\mu_Y}{\mu_X} + \frac{\sigma_X^2\mu_Y}{\mu_X^3} - \frac{\sigma_{XY}}{\mu_X^2}$
and $\text{Var}(Z) \approx \frac{\sigma_X^2\mu_Y^2}{\mu_X^4} + \frac{\sigma_Y^2}{\mu_X^2} - \frac{2\sigma_{XY}\mu_Y}{\mu_X^3}$. The approximation is ill-conditioned as $|\mu_X| \downarrow 0$.

# 5 Normal-Derived Distributions

## 5.1 Normal-Derived Distribution Families

Independent $N(0, 1)$ variables generate the normal-derived inference laws used below: $\chi^2$ sums, Student $t$ ratios, and $F$ ratios.

**Proposition 5.1.1.** *Chi-Square Sums*
*Let $n \ge 1$ and $Z_1, \ldots, Z_n \overset{\text{ind}}{\sim} N(0, 1)$.*
*(i) For $V = \sum_{i=1}^{n} Z_i^2$, $V \sim \chi_n^2 = \text{Gamma}(\frac{n}{2}, \frac{1}{2})$ in the rate convention,*
*$f_V(v) = 2^{-\frac{n}{2}}\Gamma(\frac{n}{2})^{-1}v^{\frac{n}{2}-1}e^{-\frac{v}{2}}\mathbf{1}_{(0,\infty)}(v)$, $\psi_V(t) = (1 - 2t)^{-\frac{n}{2}}$ for $t < \frac{1}{2}$, $\mathbb{E}V = n$, and $\text{Var}(V) = 2n$.*
*(ii) If $U \sim \chi_m^2$, $W \sim \chi_n^2$, and $U \perp\!\!\!\perp W$, then $U + W \sim \chi_{m+n}^2$.*

*Proof.*

(i) For $Z \sim N(0,1)$ and $t < \frac{1}{2}$, $\mathbb{E}e^{tZ^2} = (2\pi)^{-\frac{1}{2}} \int_{\mathbb{R}} e^{-\frac{(1-2t)z^2}{2}}\, dz = (1-2t)^{-\frac{1}{2}}$.
By independence, $\psi_V(t) = \prod_{i=1}^{n} \mathbb{E}e^{tZ_i^2} = (1-2t)^{-\frac{n}{2}}$. By MGF uniqueness, $V \sim \text{Gamma}(\frac{n}{2}, \frac{1}{2})$.
Differentiating $\psi_V$ at 0 gives $\mathbb{E}V = n$ and $\text{Var}(V) = 2n$.

(ii) If $U \perp\!\!\!\perp W$, then $\psi_{U+W}(t) = (1-2t)^{-\frac{m}{2}}(1-2t)^{-\frac{n}{2}} = (1-2t)^{-\frac{m+n}{2}}$, hence $U+W \sim \chi^2_{m+n}$.

*Structural Commentary*: Chi-square degrees of freedom add by MGF exponents.
*Key Insight*: Squared independent standard normals form gamma sums. □

**Definition 5.1.2.** *Normal Sample Statistics*

(i) *Sample*: $X_1, \ldots, X_n$ are independent with $X_i \sim N(\mu, \sigma^2)$, $n \geq 2$, and $\sigma^2 > 0$.

(ii) *Sample mean*: $\overline{X} := n^{-1} \sum_{i=1}^{n} X_i$ estimates the location $\mu$.

(iii) *Sample variance*: $S^2 := (n-1)^{-1} \sum_{i=1}^{n} (X_i - \overline{X})^2$ estimates the scale $\sigma^2$.

**Proposition 5.1.3.** *Normal Sample Mean*

*Let $X_1, \ldots, X_n$ be independent with $X_i \sim N(\mu, \sigma^2)$, and set $\overline{X} := n^{-1} \sum_{i=1}^{n} X_i$. Then $\overline{X} \sim N(\mu, \frac{\sigma^2}{n})$.*

*Proof.* Since independent normal sums are normal, $\overline{X}$ is normal.
Also $\mathbb{E}\overline{X} = n^{-1} \sum_i \mathbb{E}X_i = \mu$ and $\text{Var}(\overline{X}) = n^{-2} \sum_i \text{Var}(X_i) = \frac{\sigma^2}{n}$. Thus $\overline{X} \sim N(\mu, \frac{\sigma^2}{n})$.
*Structural Commentary*: Averaging preserves normality and divides variance by $n$.
*Key Insight*: The sample mean is an exact normal pivot center, not an asymptotic approximation. □

**Theorem 5.1.4.** *Mean-Residual Independence*

*Let $X_1, \ldots, X_n$ be independent with $X_i \sim N(\mu, \sigma^2)$, set $\overline{X} := n^{-1} \sum_i X_i$, $R := (X_1 - \overline{X}, \ldots, X_n - \overline{X})$, and $S^2 := (n-1)^{-1} \sum_i R_i^2$. Then $\overline{X} \perp\!\!\!\perp R$, hence $\overline{X} \perp\!\!\!\perp S^2$.*

*Proof.* Let $Z_i := \frac{X_i - \mu}{\sigma}$, so $Z \sim N_n(0, I_n)$. Choose orthogonal $Q$ with first row $n^{-\frac{1}{2}}\mathbf{1}^\top$, set $Y := QZ$.
Then $Y \sim N_n(0, I_n)$, $Y_1 \perp\!\!\!\perp (Y_2, \ldots, Y_n)$. Since $\overline{X} = \mu + \frac{\sigma Y_1}{\sqrt{n}}$ and $R = \sigma(I - n^{-1}\mathbf{1}\mathbf{1}^\top)Z = \sigma Q^\top (0, Y_2, \ldots, Y_n)^\top$, $\overline{X}$ is function of $Y_1$, $R$ is function of $(Y_2, \ldots, Y_n)$. Therefore $\overline{X} \perp\!\!\!\perp R$, by $S^2 = (n-1)^{-1} \sum_i R_i^2$, $\overline{X} \perp\!\!\!\perp S^2$.

*Structural Commentary*: Mean and residuals are orthogonal projections.
*Key Insight*: For Gaussian vectors, orthogonal components are independent. □

**Theorem 5.1.5.** *Sample Variance Chi-Square Law*

*Given independent $X_1, \ldots, X_n$; $X_i \sim N(\mu, \sigma^2)$;$S^2 := (n-1)^{-1} \sum_{i=1}^{n} (X_i - \overline{X})^2$. Then $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2_{n-1}$.*

*Proof.* Let $Z_i := \frac{X_i - \mu}{\sigma}$, so $Z = (Z_1, \ldots, Z_n) \sim N_n(0, I_n)$.
Choose orthogonal matrix $Q$ with first row $n^{-\frac{1}{2}}(1, \ldots, 1)$, set $Y := QZ$.
Orthogonal invariance gives $Y \sim N_n(0, I_n)$ with independent coordinates.
Then $Y_1 = \frac{\sqrt{n}(\overline{X} - \mu)}{\sigma}$ and $\sum_{j=2}^{n} Y_j^2 = \sum_{i=1}^{n} (Z_i - \overline{Z})^2 = \frac{(n-1)S^2}{\sigma^2}$.
Thus $\frac{(n-1)S^2}{\sigma^2}$ is the sum of squares of $n-1$ independent $N(0,1)$ variables, hence $\chi^2_{n-1}$.
*Structural Commentary*: Cantering leaves $n-1$ residual directions.
*Key Insight*: The sample variance is the squared Gaussian residual length scaled by $\sigma^2$. □

**Definition 5.1.6.** *Student t Distribution*
If $Z \sim N(0,1)$, $U \sim \chi^2_n$, $Z \perp\!\!\!\perp U$, then $T = Z/\sqrt{\frac{U}{n}}$ has *Student t distribution* with $df = n \geq 1$, $T \sim t_n$.

**Proposition 5.1.7.** *Student t Density*

*Let $n \geq 1$ and $T = Z/\sqrt{\frac{U}{n}}$ with $Z \sim N(0,1)$, $U \sim \chi^2_n$, and $Z \perp\!\!\!\perp U$.*

*(i) For $t \in \mathbb{R}$, $f_T(t) = \frac{\Gamma(\frac{n+1}{2})}{\sqrt{n\pi}\Gamma(\frac{n}{2})}(1 + \frac{t^2}{n})^{-\frac{n+1}{2}}$.*

*(ii) For all $t \in \mathbb{R}$, $f_T(-t) = f_T(t)$.*

*(iii) For each fixed $t \in \mathbb{R}$, $\lim_{n\to\infty} f_T(t) = \phi(t) = (2\pi)^{-\frac{1}{2}} e^{-\frac{t^2}{2}}$.*

*Proof.*

(i) For fixed $u > 0$, $z = t\sqrt{\frac{u}{n}}$ has Jacobian $\sqrt{\frac{u}{n}}$, so $f_T(t) = \int_0^\infty \phi(t\sqrt{\frac{u}{n}})\sqrt{\frac{u}{n}} f_U(u)\,du$.
Substituting $\phi(z) = (2\pi)^{-\frac{1}{2}} e^{-\frac{z^2}{2}}$ and $f_U(u) = 2^{-\frac{n}{2}}\Gamma(\frac{n}{2})^{-1} u^{\frac{n}{2}-1} e^{-\frac{u}{2}} \mathbf{1}_{(0,\infty)}(u)$ gives
$f_T(t) = \frac{1}{\sqrt{n\pi}\,2^{\frac{n+1}{2}}\Gamma(\frac{n}{2})} \int_0^\infty u^{\frac{n+1}{2}-1} e^{-\frac{(1+\frac{t^2}{n})u}{2}}\,du$. Then $F_T(t) = \Gamma(\frac{n+1}{2}) 2^{\frac{n+1}{2}} (1+\frac{t^2}{n})^{-\frac{n+1}{2}}$, yielding density.

(ii) By (i), $f_T(t) = c_n h_n(t^2)$ for $c_n > 0$, so $f_T(-t) = f_T(t)$ for all $t$.

(iii) For fixed $t$, $f_T(t) = (2\pi)^{-\frac{1}{2}} \frac{\Gamma(\frac{n+1}{2})}{\sqrt{\frac{n}{2}}\Gamma(\frac{n}{2})} (1+\frac{t^2}{n})^{-\frac{n+1}{2}}$, gamma ratio tends to 1, last factor tends to $e^{-\frac{t^2}{2}}$.

*Structural Commentary*: Student $t$ is a normal scale mixture with random chi-square scale.
*Key Insight*: Random denominator uncertainty thickens the normal tails. □

**Example 5.1.8.** *$t_1$ is Cauchy*
If $T \sim t_1$, then $f_T(t) = \Gamma(1)(\sqrt{\pi}\Gamma(\frac{1}{2}))^{-1}(1+t^2)^{-1} = \frac{1}{\pi(1+t^2)}$, so $T \sim \text{Cauchy}(0,1)$.

**Definition 5.1.9.** *F Distribution*
If $U \sim \chi^2_m$, $V \sim \chi^2_n$, $U \perp\!\!\!\perp V$, then $W = (\frac{U}{m})/(\frac{V}{n})$ has *F distribution*, *df* $= (m,n)$, $W \sim F_{m,n}$.

**Proposition 5.1.10.** *F Density*
*Let $m, n \geq 1$ and $W = (\frac{U}{m})/(\frac{V}{n})$, where $U \sim \chi^2_m$, $V \sim \chi^2_n$, and $U \perp\!\!\!\perp V$.*

*(i)* $f_W(w) = \frac{\Gamma(\frac{m+n}{2})}{\Gamma(\frac{m}{2})\Gamma(\frac{n}{2})} (\frac{m}{n})^{\frac{m}{2}} w^{\frac{m}{2}-1} (1+\frac{mw}{n})^{-\frac{m+n}{2}} \mathbf{1}_{(0,\infty)}(w)$.
*(ii) If $n > 2$, then $\mathbb{E}W = \frac{n}{n-2}$.*

*Proof.*

(i) For $w > 0$, set $u = \frac{mwv}{n}$; the Jacobian is $\frac{mv}{n}$, so $f_W(w) = \int_0^\infty f_U(\frac{mwv}{n}) f_V(v)(\frac{mv}{n})\,dv$. By substitution,
$f_W(w) = C w^{\frac{m}{2}-1} \int_0^\infty v^{\frac{m+n}{2}-1} e^{-\frac{(1+\frac{mw}{n})v}{2}}\,dv$; $C = \frac{(\frac{m}{n})^{\frac{m}{2}}}{2^{\frac{m+n}{2}}\Gamma(\frac{m}{2})\Gamma(\frac{n}{2})}$, by gamma integral, QED.

(ii) Since $U \perp\!\!\!\perp V$, $\mathbb{E}W = (\frac{n}{m})\mathbb{E}U\,\mathbb{E}(V^{-1}) = n\,\mathbb{E}(V^{-1}) = \frac{n}{n-2}$ for $n > 2$.

*Structural Commentary*: An $F$ variable is a ratio of independent chi-square averages.
*Key Insight*: Denominator variability controls the right tail and the mean condition $n > 2$. □

**Corollary 5.1.11.** *Reciprocal F and Squared t*
*(i) For $m, n \geq 1$, if $W \sim F_{m,n}$, then $W^{-1} \sim F_{n,m}$.*
*(ii) If $T \sim t_n$, then $T^2 \sim F_{1,n}$.*

*Proof.*

(i) Write $W = (\frac{U}{m})/(\frac{V}{n})$ with $U \sim \chi^2_m$, $V \sim \chi^2_n$, and $U \perp\!\!\!\perp V$; then $W^{-1} = (\frac{V}{n})/(\frac{U}{m}) \sim F_{n,m}$.

(ii) Write $T = Z/\sqrt{\frac{V}{n}}$ with $Z \sim N(0,1)$, $V \sim \chi^2_n$, and $Z \perp\!\!\!\perp V$; since $Z^2 \sim \chi^2_1$, $T^2 = (\frac{Z^2}{1})/(\frac{V}{n}) \sim F_{1,n}$.

*Structural Commentary*: Reciprocals swap $F$ degrees of freedom, and squaring $t$ removes the normal sign.
*Key Insight*: Both identities are immediate from the same chi-square-ratio representation. □

# Part XVI
# Probability and Stochastic Systems

## 1 Conditional Probability and Conditional Expectation

### 1.1 Conditional Probability

**Definition 1.1.1.** *Conditional Probability*
For $A, B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$, the *conditional probability* of $A$ given $B$ is $\mathbb{P}(A \mid B) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$, i.e. the probability of $A$ under the normalised restriction of $\mathbb{P}$ to $B$.

**Example 1.1.2.** *Diagnostic Table*
Let $T$ be the event of a positive test and $D$ the event of disease, with observed counts

| | $D$ | $D^c$ | total |
|---|---|---|---|
| $T$ | 25 | 14 | 39 |
| $T^c$ | 18 | 78 | 96 |
| total | 43 | 92 | 135 |

Let $N(\cdot)$ be counts, $\widehat{\mathbb{P}}(A \mid B) = \frac{N(A \cap B)}{N(B)}$.
Thus $\widehat{\mathbb{P}}(D \mid T) = \frac{25}{39}$, $\widehat{\mathbb{P}}(D^c \mid T^c) = \frac{78}{96}$, $\widehat{\mathbb{P}}(D^c \mid T) = \frac{14}{39}$, and $\widehat{\mathbb{P}}(D \mid T^c) = \frac{18}{96}$.

**Proposition 1.1.3.** *Conditioning on a Fixed Event Gives a Probability Measure*
*For $B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$, define $\mathbb{P}_B : \mathcal{F} \to [0,1]$ by $\mathbb{P}_B(A) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$.*
*Then $\mathbb{P}_B$ is a probability measure on $(\Omega, \mathcal{F})$.*

*Proof.* For $A \in \mathcal{F}$, $\mathbb{P}_B(A) \geq 0$, and $\mathbb{P}_B(\Omega) = \frac{\mathbb{P}(B)}{\mathbb{P}(B)} = 1$.
If $(A_i)_{i \geq 1}$ are pairwise disjoint, then $(A_i \cap B)_{i \geq 1}$ are pairwise disjoint and $(\bigcup_i A_i) \cap B = \bigcup_i (A_i \cap B)$.
Thus $\mathbb{P}_B(\bigcup_i A_i) = \frac{\mathbb{P}(\bigcup_i (A_i \cap B))}{\mathbb{P}(B)} = \sum_i \frac{\mathbb{P}(A_i \cap B)}{\mathbb{P}(B)} = \sum_i \mathbb{P}_B(A_i)$.
*Structural Commentary*: Conditioning normalizes the restriction of $\mathbb{P}$ to $B$.
*Key Insight*: Once $B$ is fixed, conditional probability is ordinary probability on the restricted world. □

**Remark 1.1.4.** *Conditional Probability Is Not Symmetric*
In general, $\mathbb{P}(A \mid B) \neq \mathbb{P}(B \mid A)$; conditioning fixes the reference population. For disease event $D$ and positive-test event $T$, large $\mathbb{P}(T \mid D)$ need not imply large $\mathbb{P}(D \mid T)$ when $\mathbb{P}(D)$ is small.

**Proposition 1.1.5.** *Multiplication and Chain Rules*
*Let $A, B, C \in \mathcal{F}$ with $\mathbb{P}(A) > 0$, $\mathbb{P}(B) > 0$, and $\mathbb{P}(B \cap C) > 0$.*

*(i)* $\mathbb{P}(A \cap B) = \mathbb{P}(A \mid B)\mathbb{P}(B) = \mathbb{P}(B \mid A)\mathbb{P}(A)$.
*(ii)* $\mathbb{P}(A \cap B \cap C) = \mathbb{P}(A \mid B \cap C)\mathbb{P}(B \mid C)\mathbb{P}(C)$.

*Proof.*

(i) For $\mathbb{P}(F) > 0$, $\mathbb{P}(E \mid F)\mathbb{P}(F) = \mathbb{P}(E \cap F)$; apply this with $(E, F) = (A, B)$ and $(B, A)$.
(ii) Since $\mathbb{P}(B \cap C) > 0$, $\mathbb{P}(A \mid B \cap C)\mathbb{P}(B \mid C)\mathbb{P}(C) = \frac{\mathbb{P}(A \cap B \cap C)}{\mathbb{P}(B \cap C)} \frac{\mathbb{P}(B \cap C)}{\mathbb{P}(C)} \mathbb{P}(C) = \mathbb{P}(A \cap B \cap C)$.

*Structural Commentary*: The chain rule orders information as $C$, then $B$ within $C$, then $A$ within $B \cap C$.
*Key Insight*: A joint probability factors into conditional increments along a chosen order. □

**Example 1.1.6.** *Cards Without Replacement*
Draw two cards from a standard deck without replacement. Let $A$ be the event that the first card is the ace of clubs and $B$ the event that the second card is the queen of diamonds. Since $\mathbb{P}(A) = \frac{1}{52}$ and $\mathbb{P}(B \mid A) = \frac{1}{51}$, $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B \mid A) = \frac{1}{52 \cdot 51}$. The factor $\mathbb{P}(B \mid A)$ uses the 51-card deck remaining after $A$ occurs.

**Proposition 1.1.7.** *Independence Through Conditioning*
*Let $A, B \in \mathcal{F}$ with $\mathbb{P}(A) > 0$ and $\mathbb{P}(B) > 0$. The following are equivalent:*

*(i)* *$A$ and $B$ are independent.*
*(ii)* $\mathbb{P}(A \mid B) = \mathbb{P}(A)$.
*(iii)* $\mathbb{P}(B \mid A) = \mathbb{P}(B)$.

*Proof.*

(*i*) $\Rightarrow$ (*ii*) If $A$ and $B$ are independent, then $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$, so $\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \mathbb{P}(A)$.

(*ii*) $\Rightarrow$ (*iii*) If $\mathbb{P}(A \mid B) = \mathbb{P}(A)$, then $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$, so $\mathbb{P}(B \mid A) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)} = \mathbb{P}(B)$.

(*iii*) $\Rightarrow$ (*i*) If $\mathbb{P}(B \mid A) = \mathbb{P}(B)$, then $\mathbb{P}(A \cap B) = \mathbb{P}(B \mid A)\mathbb{P}(A) = \mathbb{P}(A)\mathbb{P}(B)$, hence $A$ and $B$ are independent.

*Structural Commentary*: Each condition is $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ normalized by a positive marginal.
*Key Insight*: Independence means conditioning on either event preserves the probability of the other. □

**Theorem 1.1.8.** *Law of Total Probability*

*(i) Let $A_1, \ldots, A_k \in \mathcal{F}$ form a finite partition of $\Omega$ with $\mathbb{P}(A_i) > 0$ for all $i$.*

*(ii) For every $B \in \mathcal{F}$, $\mathbb{P}(B) = \sum_{i=1}^{k} \mathbb{P}(B \mid A_i)\mathbb{P}(A_i)$.*

*Proof.*

(i) The partition assumptions give $B = \bigsqcup_{i=1}^{k} (B \cap A_i)$, hence $\mathbb{P}(B) = \sum_{i=1}^{k} \mathbb{P}(B \cap A_i)$.

(ii) For each $i$, $\mathbb{P}(B \cap A_i) = \mathbb{P}(B \mid A_i)\mathbb{P}(A_i)$; substituting in (i) gives the formula.

*Structural Commentary*: The partition $\{A_i\}_{i=1}^{k}$ decomposes $B$ into disjoint cases $B \cap A_i$.
*Key Insight*: Total probability is weighted averaging over cases. □

**Example 1.1.9.** *Second Draw by Total Probability*
An urn has 3 red balls, 1 blue ball; draw twice without replacement, let $R_i$ be the event that draw $i$ is red. By total probability, $\mathbb{P}(R_2) = \mathbb{P}(R_2 \mid R_1)\mathbb{P}(R_1) + \mathbb{P}(R_2 \mid R_1^c)\mathbb{P}(R_1^c) = \frac{2}{3} \cdot \frac{3}{4} + 1 \cdot \frac{1}{4} = \frac{3}{4}$.
*Key Insight*: The second draw has the same marginal red probability as the first.

**Definition 1.1.10.** *Prior and Posterior Probabilities*
Let $A_1, \ldots, A_k \in \mathcal{F}$ be a partition with $\mathbb{P}(A_i) > 0$ for all $i$, and let $B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$.

(i) $\mathbb{P}(A_i)$ is the *prior probability* of $A_i$.

(ii) $\mathbb{P}(A_i \mid B)$ is the *posterior probability* of $A_i$ after conditioning on $B$.

**Theorem 1.1.11.** *Bayes' Theorem*

*Let $A_1, \ldots, A_k \in \mathcal{F}$ satisfy $A_i \cap A_j = \varnothing$ for $i \neq j$, $\bigcup_{i=1}^{k} A_i = \Omega$, and $\mathbb{P}(A_i) > 0$ for all $i$.*

*If $B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$, then for each $i$, $\mathbb{P}(A_i \mid B) = \frac{\mathbb{P}(B \mid A_i)\mathbb{P}(A_i)}{\sum_{j=1}^{k} \mathbb{P}(B \mid A_j)\mathbb{P}(A_j)}$.*

*Proof.* For fixed $i$, $\mathbb{P}(A_i \mid B) = \frac{\mathbb{P}(A_i \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B \mid A_i)\mathbb{P}(A_i)}{\mathbb{P}(B)}$.
Since $B = \bigsqcup_{j=1}^{k} (B \cap A_j)$, $\mathbb{P}(B) = \sum_{j=1}^{k} \mathbb{P}(B \cap A_j) = \sum_{j=1}^{k} \mathbb{P}(B \mid A_j)\mathbb{P}(A_j)$. Substitution gives the formula.
*Structural Commentary*: Bayes reweights priors by likelihoods, then normalises.
*Key Insight*: Posterior equals likelihood times prior divided by evidence. □

**Corollary 1.1.12.** *Posterior Mass Must Balance*
*Let $A_1, \ldots, A_k \in \mathcal{F}$ be a partition of $\Omega$ with $\mathbb{P}(A_i) > 0$ for all $i$, and let $B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$.*
*If $\mathbb{P}(A_1 \mid B) < \mathbb{P}(A_1)$, then there exists $i \in \{2, \ldots, k\}$ such that $\mathbb{P}(A_i \mid B) > \mathbb{P}(A_i)$.*

*Proof.* Suppose $\mathbb{P}(A_i \mid B) \leq \mathbb{P}(A_i)$ for all $i \geq 2$
As $\sum_{i=1}^{k} \mathbb{P}(A_i \mid B) = \sum_{i=1}^{k} \mathbb{P}(A_i) = 1$, hence $1 = \sum_{i=1}^{k} \mathbb{P}(A_i \mid B) < \mathbb{P}(A_1) + \sum_{i=2}^{k} \mathbb{P}(A_i) = 1$, a contradiction.
*Structural Commentary*: Posterior mass is conserved across the partition.
*Key Insight*: A decrease in one cell must be offset by an increase in another. □

**Example 1.1.13.** *Transition Probabilities*
Let states $U, M, L$ denote upper, middle, lower groups, and let $P = (p_{ij})$ with $p_{ij} = \mathbb{P}(j_2 \mid i_1)$:

| | $U_2$ | $M_2$ | $L_2$ |
|---|---|---|---|
| $U_1$ | .45 | .48 | .07 |
| $M_1$ | .05 | .70 | .25 |
| $L_1$ | .01 | .50 | .49 |

For $\pi_1 = (.10, .40, .50)$, total probability gives $\pi_2 = \pi_1 P$ and $\mathbb{P}(U_2) = .45(.10) + .05(.40) + .01(.50) = .07$.
Bayes gives the inverse query $\mathbb{P}(U_1 \mid U_2) = \frac{\mathbb{P}(U_2 \mid U_1)\mathbb{P}(U_1)}{\mathbb{P}(U_2)} = \frac{.45(.10)}{.07} \approx .64$.
*Key Insight*: Use $\pi P$ forward and Bayes backward.

**Example 1.1.14.** *Base Rates in a Medical Test*
Let $D$ be disease and $T$ a positive test, with joint law

| | $D$ | $D^c$ |
|---|---|---|
| $T$ | 0.0081 | 0.0900 |
| $T^c$ | 0.0009 | 0.9010 |

Then $\mathbb{P}(T \mid D) = \frac{0.0081}{0.0081+0.0009} = 0.9$, $\mathbb{P}(T^c \mid D^c) = \frac{0.9010}{0.9010+0.0900} \approx 0.91$, $\mathbb{P}(D \mid T) = \frac{0.0081}{0.0081+0.0900} \approx 0.083$.
*Key Insight*: A rare condition can have small posterior probability after a positive test.

**Example 1.1.15.** *Rare Classes and Screening Errors*
Let $L$ denote lying and $T = L^c$ truth, with $\mathbb{P}(+ \mid L) = .88$, $\mathbb{P}(+ \mid T) = .14$, $\mathbb{P}(L) = .01$, and $\mathbb{P}(T) = .99$.
Bayes gives $\mathbb{P}(T \mid +) = \frac{\mathbb{P}(+|T)\mathbb{P}(T)}{\mathbb{P}(+|T)\mathbb{P}(T)+\mathbb{P}(+|L)\mathbb{P}(L)} = \frac{(.14)(.99)}{(.14)(.99)+(.88)(.01)} \approx .94$.
*Key Insight*: Rare-class positives can mostly come from the common class.

**Example 1.1.16.** *Monty Hall*
The player chooses door 1. Let $\omega = (\omega_1, \omega_2)$, where $\omega_1$ is the prize door and $\omega_2$ is Monty's opened door.
Monty opens neither door 1 nor the prize door, and chooses uniformly from $\{2, 3\}$ when $\omega_1 = 1$.
The induced probabilities are $\mathbb{P}(1, 2) = \mathbb{P}(1, 3) = \frac{1}{6}$ and $\mathbb{P}(2, 3) = \mathbb{P}(3, 2) = \frac{1}{3}$.
Let $S = \{\omega_1 = 1\}$ be the event that staying wins.
Given $\omega_2 = 2$, the staying-win probability is $\mathbb{P}(S \mid \omega_2 = 2) = \frac{\frac{1}{6}}{\frac{1}{6}+\frac{1}{3}} = \frac{1}{3}$.
Thus switching wins with probability $\frac{2}{3}$, and the case $\omega_2 = 3$ is symmetric.
*Key Insight*: Once Monty's rule is part of the sample space, switching has conditional probability $\frac{2}{3}$.

**Example 1.1.17.** *Three Cards and Conditional Sampling*
Cards are $GG$, $RR$, and $GR$. Choose a card uniformly and then a side uniformly.
Let $V$ be the event that the visible side is green and $H$ the event that the hidden side is green.
Among the three equally likely green visible sides, two have green hidden side, so $\mathbb{P}(H \mid V) = \frac{2}{3}$.

**Example 1.1.18.** *Conditioning Changes the Reference Class*
Let $X \sim \text{Binomial}(5, \frac{1}{4})$ be number of blue-eyed children, $Y$ be event that youngest child has blue eyes.

(i) Given $X \geq 1$, $\mathbb{P}(X \geq 3 \mid X \geq 1) = \frac{\binom{5}{3}\left(\frac{1}{4}\right)^3\left(\frac{3}{4}\right)^2+\binom{5}{4}\left(\frac{1}{4}\right)^4\left(\frac{3}{4}\right)+\left(\frac{1}{4}\right)^5}{1-\left(\frac{3}{4}\right)^5} = \frac{106}{781}$.

(ii) Given $Y$, $\mathbb{P}(X \geq 3 \mid Y) = 1 - \left(\frac{3}{4}\right)^4 - 4\left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^3 = \frac{67}{256}$.

*Key Insight*: Similar information can determine different conditioning events.

**Example 1.1.19.** *Bayes with Several Categories*
Let $O \in \{M, W, L\}$ denote operating system with probabilities $(0.30, 0.50, 0.20)$.
Let $I$ be infection, with $\mathbb{P}(I \mid O = M) = 0.65$, $\mathbb{P}(I \mid O = W) = 0.82$, and $\mathbb{P}(I \mid O = L) = 0.50$.
Bayes' theorem gives $\mathbb{P}(W \mid I) = \frac{0.82 \cdot 0.50}{0.65 \cdot 0.30+0.82 \cdot 0.50+0.50 \cdot 0.20} = \frac{82}{141}$.

**Example 1.1.20.** *Posterior and Predictive Probabilities*
Let $C$ be uniform on $\{C_1, \ldots, C_5\}$, and suppose coin $C_i$ has head probability $p_i$.
The head-probability vector is $(p_1, p_2, p_3, p_4, p_5) = \left(0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\right)$. Let $H_n$ be the event that toss $n$ is heads.
The posterior after one head is $\mathbb{P}(C_i \mid H_1) = \frac{p_i}{p_1+p_2+p_3+p_4+p_5} = \left(0, \frac{1}{10}, \frac{1}{5}, \frac{3}{10}, \frac{2}{5}\right)_i$.
The predictive probability is $\mathbb{P}(H_2 \mid H_1) = \sum_{i=1}^{5} p_i \mathbb{P}(C_i \mid H_1) = \frac{3}{4}$.
If $B_4$ is the event that the first head occurs on toss 4,
the posterior is $\mathbb{P}(C_i \mid B_4) = \frac{(1-p_i)^3 p_i}{\sum_{j=1}^{5}(1-p_j)^3 p_j}$, and $\left(\mathbb{P}(C_i \mid B_4)\right)_{i=1}^{5} = \left(0, \frac{27}{46}, \frac{16}{46}, \frac{3}{46}, 0\right)$.
*Key Insight*: Bayes' theorem updates hidden parameters; prediction averages over the posterior.

**Definition 1.1.21.** *Conditional Mass and Density Functions*

(i) *Discrete*: if $X, Y$ are discrete and $f_Y(y) > 0$, the conditional mass function is $f_{X|Y}(x \mid y) := \mathbb{P}(X = x \mid Y = y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$.

(ii) *Continuous*: if $X, Y$ have a joint density and $f_Y(y) > 0$, the conditional density is $f_{X|Y}(x \mid y) := \frac{f_{X,Y}(x,y)}{f_Y(y)}$, so $\mathbb{P}(X \in A \mid Y = y) = \int_A f_{X|Y}(x \mid y)\, dx$ for measurable $A$.

**Proposition 1.1.22.** *Joint Factorizations and Total Probability for Conditional Laws*
*Assume the displayed sums or integrals are well defined.*

*(i) If $(X, Y)$ is discrete, $p_Y(y) > 0$, then $p_{X,Y}(x, y) = p_{X|Y}(x \mid y)p_Y(y)$, $p_X(x) = \sum_y p_{X|Y}(x \mid y)p_Y(y)$.*

*(ii) If $(X,Y)$ has density, $f_X(x)>0$, then $f_{X,Y}(x,y)=f_{Y|X}(y\mid x)f_X(x)$, $f_Y(y)=\int_{-\infty}^{\infty} f_{Y|X}(y\mid x)f_X(x)\,dx$.*

*Proof.*

(i) For $p_Y(y)>0$, $p_{X|Y}(x\mid y)=\frac{p_{X,Y}(x,y)}{p_Y(y)}$.
Hence $p_{X,Y}(x,y)=p_{X|Y}(x\mid y)p_Y(y)$ and $p_X(x)=\sum_y p_{X,Y}(x,y)=\sum_y p_{X|Y}(x\mid y)p_Y(y)$.

(ii) For $f_X(x)>0$, $f_{Y|X}(y\mid x)=\frac{f_{X,Y}(x,y)}{f_X(x)}$.
Hence $f_{X,Y}(x,y)=f_{Y|X}(y\mid x)f_X(x)$ and $f_Y(y)=\int_{-\infty}^{\infty} f_{X,Y}(x,y)\,dx=\int_{-\infty}^{\infty} f_{Y|X}(y\mid x)f_X(x)\,dx$.

*Structural Commentary*: Conditional laws disintegrate a joint law into a marginal law and a fiber law.
*Key Insight*: Total probability is marginalization after conditional factorization. □

**Example 1.1.23.** *Conditional Masses from the Three-Toss Table*
Let $Y$ be the number of heads in three fair tosses, and let $X=1$ iff the first toss is heads.
Since $p_Y(1)=\frac{3}{8}$, $p_{X,Y}(0,1)=\frac{2}{8}$, and $p_{X,Y}(1,1)=\frac{1}{8}$, one has $p_{X|Y}(0\mid 1)=\frac{2}{3}$ and $p_{X|Y}(1\mid 1)=\frac{1}{3}$.
Thus, given $Y=1$, the first toss is more likely tails.

**Example 1.1.24.** *Uniform Square*
If $(X,Y)\sim\text{Uniform}([0,1]^2)$, then for $0\le y\le 1$, $0\le x\le 1$, $f_{X|Y}(x\mid y)=1$. Hence $X\mid Y=y\sim\text{Uniform}(0,1)$.

**Example 1.1.25.** *Conditioning in the Parabolic Region*
Let $(X,Y)$ have density $f(x,y)=\frac{21}{4}x^2y$, $x^2\le y\le 1$. Then $f_X(x)=\frac{21}{8}x^2(1-x^4)$, $-1\le x\le 1$.
For $x^2\le y\le 1$, $f_{Y|X}(y\mid x)=\frac{\frac{21}{4}x^2y}{\frac{21}{8}x^2(1-x^4)}=\frac{2y}{1-x^4}$.
At $x=\frac{1}{2}$, $f_{Y|X}(y\mid\frac{1}{2})=\frac{32y}{15}$, so $\mathbb{P}(Y\ge\frac{3}{4}\mid X=\frac{1}{2})=\int_{\frac{3}{4}}^1\frac{32y}{15}\,dy=\frac{7}{15}$.

**Example 1.1.26.** *Poisson Thinning*
Let $\lambda>0$, $0\le p\le 1$, $N\sim\text{Poisson}(\lambda)$, and $X\mid N=n\sim\text{Binomial}(n,p)$.
For $k\in\mathbb{N}_0$, by total probability, $\mathbb{P}(X=k)=\sum_{n=k}^{\infty} e^{-\lambda}\frac{\lambda^n}{n!}\binom{n}{k}p^k(1-p)^{n-k}=e^{-\lambda p}\frac{(\lambda p)^k}{k!}$, thus $X\sim\text{Poisson}(\lambda p)$.

**Example 1.1.27.** *A Triangular Exponential Joint Density*
Let $\lambda>0$ and $f_{X,Y}(x,y)=\lambda^2e^{-\lambda y}\mathbf{1}_{\{0\le x\le y\}}$.
For $x\ge 0$, $f_X(x)=\int_x^{\infty}\lambda^2e^{-\lambda y}\,dy=\lambda e^{-\lambda x}$; for $y\ge 0$, $f_Y(y)=\int_0^y\lambda^2e^{-\lambda y}\,dx=\lambda^2ye^{-\lambda y}$.
Hence $f_{Y|X}(y\mid x)=\frac{f_{X,Y}(x,y)}{f_X(x)}=\lambda e^{-\lambda(y-x)}\mathbf{1}_{[x,\infty)}(y)$ and $f_{X|Y}(x\mid y)=\frac{f_{X,Y}(x,y)}{f_Y(y)}=y^{-1}\mathbf{1}_{[0,y]}(x)$.
Thus $Y\mid X=x\sim x+\text{Exp}(\lambda)$ and $X\mid Y=y\sim\text{Uniform}(0,y)$.

**Example 1.1.28.** *Bivariate Normal Conditional Density*
Let $(X,Y)$ be bivariate normal with means $\mu_X,\mu_Y$, standard deviations $\sigma_X,\sigma_Y>0$, and correlation $\rho\in(-1,1)$.
For fixed $x$, $f_{Y|X}(y\mid x)=\frac{f_{X,Y}(x,y)}{f_X(x)}=\{\sigma_Y\sqrt{2\pi(1-\rho^2)}\}^{-1}\exp\left[-\frac{(y-\mu_Y-\rho\sigma_Y\sigma_X^{-1}(x-\mu_X))^2}{2\sigma_Y^2(1-\rho^2)}\right]$.
Thus $Y\mid X=x\sim N(\mu_Y+\rho\sigma_Y\sigma_X^{-1}(x-\mu_X),\sigma_Y^2(1-\rho^2))$; if $\rho=0$, this law is independent of $x$.

## 1.2 Conditional Expectation

**Definition 1.2.1.** *Conditional Expectation Given a Random Variable*
Let $X,Y$ be random variables, and assume the relevant sums or integrals exist.

(i) *Discrete case*: If $X,Y$ are discrete, then $\mathbb{E}(X\mid Y=y):=\sum_x x\,p_{X|Y}(x\mid y)$.

(ii) *Density case*: If $X,Y$ have conditional density $f_{X|Y}$, then $\mathbb{E}(X\mid Y=y):=\int_{\mathbb{R}} x f_{X|Y}(x\mid y)\,dx$

(iii) *Measurable transforms*: For measurable $r$,

$$\mathbb{E}(r(X,Y)\mid Y=y)=\begin{cases}\sum_x r(x,y)p_{X|Y}(x\mid y), & \text{discrete case},\\ \int_{\mathbb{R}} r(x,y)f_{X|Y}(x\mid y)\,dx, & \text{density case}.\end{cases}$$

(iv) *Random-variable form*: $\mathbb{E}(X\mid Y)$ is the function $g(Y)$ with $g(y)=\mathbb{E}(X\mid Y=y)$.

**Example 1.2.2.** *A Conditional Uniform Model*
Let $X\sim\text{Uniform}(0,1)$ and, conditionally on $X=x$, let $Y\sim\text{Uniform}(x,1)$.
For $0\le x<1$ and $x<y<1$, $f_{Y|X}(y\mid x)=\frac{1}{1-x}$, so $\mathbb{E}(Y\mid X=x)=\frac{1}{1-x}\int_x^1 y\,dy=\frac{1+x}{2}$; $\mathbb{E}(Y\mid X)=\frac{1+X}{2}$.

**Theorem 1.2.3.** *Tower Property*
*Let $X,Y$ be real-valued random variables whose joint law is either discrete or admits a density, and let $r:\mathbb{R}^2\to\mathbb{R}$ be measurable. If $\mathbb{E}|Y|<\infty$ and $\mathbb{E}|r(X,Y)|<\infty$, then $\mathbb{E}[\mathbb{E}(Y\mid X)]=\mathbb{E}Y$ and $\mathbb{E}[\mathbb{E}(r(X,Y)\mid X)]=\mathbb{E}(r(X,Y))$.*

*Proof.* Let $h(x,y)$ denote either $y$ or $r(x,y)$, and set $g_h(x) := \mathbb{E}(h(X,Y) \mid X = x)$.
In the density case, $g_h(x) = \int h(x,y) f_{Y|X}(y \mid x)\, dy$ and $f(x,y) = f_X(x) f_{Y|X}(y \mid x)$,
so Fubini gives $\mathbb{E}[g_h(X)] = \int\int h(x,y) f(x,y)\, dy\, dx = \mathbb{E}h(X,Y)$.
In the discrete case, $\mathbb{E}[g_h(X)] = \sum_x \sum_y h(x,y) p_{Y|X}(y \mid x) p_X(x) = \sum_x \sum_y h(x,y) p_{X,Y}(x,y) = \mathbb{E}h(X,Y)$.
Taking $h(x,y) = y$ and $h(x,y) = r(x,y)$ gives the two identities.
*Structural Commentary*: Conditional expectation first averages over fibers $\{X = x\}$, and the outer expectation averages those fiber means over the law of $X$.
*Key Insight*: Iterated expectation removes the conditioning variable without changing the total mean. □

**Example 1.2.4.** *Using the Tower Property*
Let $X \sim \text{Uniform}(0,1)$ and, conditionally on $X = x$, let $Y \sim \text{Uniform}(x,1)$.
Since $\mathbb{E}(Y \mid X) = \frac{1+X}{2}$ and $\mathbb{E}X = \frac{1}{2}$, the tower property gives $\mathbb{E}Y = \mathbb{E}[\mathbb{E}(Y \mid X)] = \mathbb{E}\left[\frac{1+X}{2}\right] = \frac{3}{4}$.

**Proposition 1.2.5.** *Pulling Out What Is Known*
*Let $X, Y$ be real-valued random variables whose joint law is either discrete or admits a density, and let $r, s : \mathbb{R} \to \mathbb{R}$ be measurable with $\mathbb{E}|s(Y)| < \infty$, $\mathbb{E}|r(X)s(Y)| < \infty$, and $\mathbb{E}|r(X)| < \infty$. Then $\mathbb{E}(r(X)s(Y) \mid X) = r(X)\mathbb{E}(s(Y) \mid X)$ and $\mathbb{E}(r(X) \mid X) = r(X)$.*

*Proof.* Let $K_x$ be the conditional law of $Y$ given $X = x$, so $K_x(\{y\}) = p_{Y|X}(y \mid x)$ in the discrete case and $K_x(dy) = f_{Y|X}(y \mid x)\, dy$ in the density case. Set $m_s(x) := \int s(y) K_x(dy) = \mathbb{E}(s(Y) \mid X = x)$, with the integral read as a sum in the discrete case. For admissible $x$, $\mathbb{E}(r(X)s(Y) \mid X = x) = \int r(x)s(y) K_x(dy) = r(x) m_s(x)$, and $\mathbb{E}(r(X) \mid X = x) = \int r(x) K_x(dy) = r(x)$. Evaluating at $x = X$ gives the identities.
*Structural Commentary*: Conditioning on $X$ makes every function of $X$ fixed inside remaining average over $Y$.
*Key Insight*: Known factors can be treated as constants under conditional expectation. □

**Corollary 1.2.6.** *Constant Conditional Mean Implies Uncorrelated*
*Let $X, Y \in L^2$ and suppose $\mathbb{E}(X \mid Y) = c$ a.s. for some constant $c \in \mathbb{R}$. Then $c = \mathbb{E}X$ and $\text{Cov}(X,Y) = 0$.*

*Proof.* The tower property gives $\mathbb{E}X = \mathbb{E}[\mathbb{E}(X \mid Y)] = c$.
Since $Y$ is known when conditioning on $Y$, $\mathbb{E}(XY \mid Y) = Y\mathbb{E}(X \mid Y) = cY$.
Taking expectations gives $\mathbb{E}(XY) = c\mathbb{E}Y = \mathbb{E}X\, \mathbb{E}Y$, hence $\text{Cov}(X,Y) = \mathbb{E}(XY) - \mathbb{E}X\, \mathbb{E}Y = 0$.
*Structural Commentary*: A constant conditional mean kills covariance with all functions of $Y$, hence with $Y$.
*Key Insight*: Mean independence implies uncorrelatedness, but not conversely. □

**Definition 1.2.7.** *Conditional Variance*
Let $X, Y$ be real-valued random variables with finite second conditional moment for each $x$ under consideration.

(i) The conditional mean is $\mu(x) := \mathbb{E}(Y \mid X = x)$.
(ii) The conditional variance at $x$ is $v(x) := \text{Var}(Y \mid X = x) := \mathbb{E}((Y - \mu(x))^2 \mid X = x)$.
(iii) The conditional-variance random variable is $\text{Var}(Y \mid X) := v(X)$.

**Theorem 1.2.8.** *Law of Total Variance*
*Let $X, Y$ be real-valued random variables with $\mathbb{E}Y^2 < \infty$. Set $b(x) := \mathbb{E}(Y \mid X = x)$ and $v(x) := \text{Var}(Y \mid X = x)$, and write $b(X)$ and $v(X)$ for the induced random variables. Then $\text{Var}(Y) = \mathbb{E}[v(X)] + \text{Var}(b(X))$.*

*Proof.* Let $m := \mathbb{E}Y$ and $e := Y - b(X)$.
The tower property gives $\mathbb{E}b(X) = m$, and $\mathbb{E}(e \mid X) = \mathbb{E}(Y \mid X) - b(X) = 0$.
Since $Y - m = e + (b(X) - m)$, expansion gives $\text{Var}(Y) = \mathbb{E}e^2 + 2\mathbb{E}[e(b(X) - m)] + \mathbb{E}(b(X) - m)^2$.
The cross term is zero because $\mathbb{E}[e(b(X) - m)] = \mathbb{E}[(b(X) - m)\mathbb{E}(e \mid X)] = 0$.
Also $\mathbb{E}e^2 = \mathbb{E}[\mathbb{E}(e^2 \mid X)] = \mathbb{E}[v(X)]$, $\mathbb{E}(b(X) - m)^2 = \text{Var}(b(X))$. Thus $\text{Var}(Y) = \mathbb{E}[v(X)] + \text{Var}(b(X))$.
*Structural Commentary*: Variance splits into within-fiber variance and between-fiber mean variation.
*Key Insight*: Total variance equals average conditional variance plus variance of conditional expectation. □

**Example 1.2.9.** *Random Binomial Proportion*
Let $Q \sim \text{Uniform}(0,1)$ and, conditionally on $Q = q$, let $X \sim \text{Binomial}(n,q)$.
Then $\mathbb{E}(X \mid Q) = nQ$, so $\mathbb{E}X = \mathbb{E}(nQ) = \frac{n}{2}$. Also $\text{Var}(X \mid Q) = nQ(1-Q)$.
Hence $\mathbb{E}[\text{Var}(X \mid Q)] = n\int_0^1 q(1-q)\, dq = \frac{n}{6}$, $\text{Var}(\mathbb{E}(X \mid Q)) = \text{Var}(nQ) = \frac{n^2}{12}$. Thus $\text{Var}(X) = \frac{n}{6} + \frac{n^2}{12}$.

## 1.3 Conditional Moment Methods and Prediction

Conditional expectation is not only a way to compute means. It is also the geometric projection rule behind random-sum formulas and least-squares prediction.

**Proposition 1.3.1.** *Random Sums*
*Let $N \in \mathbb{N}_0$ a.s. with $\mathbb{E}N^2 < \infty$, and let $(X_i)_{i\geq 1}$ be iid, independent of $N$, with $\mathbb{E}X_1 = m$ and $\operatorname{Var}(X_1) = v < \infty$. Set $T = \sum_{i=1}^{N} X_i$, with $T = 0$ on $\{N = 0\}$.*

*(i)* $\mathbb{E}T = m\mathbb{E}N$.
*(ii)* $\operatorname{Var}(T) = v\mathbb{E}N + m^2 \operatorname{Var}(N)$.

*Proof.*

(i) On $\{N = n\}$, $T = \sum_{i=1}^{n} X_i$, so $\mathbb{E}(T \mid N = n) = nm$ and $\mathbb{E}(T \mid N) = mN$; hence $\mathbb{E}T = \mathbb{E}\mathbb{E}(T \mid N) = m\mathbb{E}N$.

(ii) On $\{N = n\}$, independence gives $\operatorname{Var}(T \mid N = n) = nv$, so $\operatorname{Var}(T \mid N) = vN$.
Total variance gives $\operatorname{Var}(T) = \mathbb{E}(vN) + \operatorname{Var}(mN) = v\mathbb{E}N + m^2 \operatorname{Var}(N)$.

*Structural Commentary*: The randomness of the number of summands contributes the second variance term.
*Key Insight*: First condition on how many summands exist; then average over that count. □

**Proposition 1.3.2.** *MGF of a Random Sum*
*Let $T = \sum_{i=1}^{N} X_i$, where $N \in \mathbb{N}_0$ is independent of iid $(X_i)_{i\geq 1}$.*
*If $\psi_N$ and $\psi_X$ are finite at the relevant arguments, then $\psi_T(t) = \psi_N(\log \psi_X(t))$.*

*Proof.* On $\{N = n\}$, independence gives $\mathbb{E}(e^{tT} \mid N = n) = \mathbb{E}e^{t\sum_{i=1}^{n} X_i} = \prod_{i=1}^{n} \psi_X(t) = [\psi_X(t)]^n$.
Hence $\psi_T(t) = \mathbb{E}[\psi_X(t)^N] = \mathbb{E}e^{N \log \psi_X(t)} = \psi_N(\log \psi_X(t))$.
*Structural Commentary*: Conditioning on $N$ turns the random sum into a fixed product of MGFs.
*Key Insight*: Random counts compose MGFs by substituting $\log \psi_X(t)$ into $\psi_N$. □

**Example 1.3.3.** *Compound Poisson Count*
Let $N \sim \text{Poisson}(\mu)$ and, given $N$, let $T = \sum_{i=1}^{N} X_i$ with iid $X_i \sim \text{Poisson}(\lambda)$ independent of $N$.
Since $\psi_N(s) = \exp(\mu(e^s - 1))$ and $\psi_X(t) = \exp(\lambda(e^t - 1))$, one has $\psi_T(t) = \psi_N(\log \psi_X(t)) = \exp(\mu(e^{\lambda(e^t-1)} - 1))$.

**Proposition 1.3.4.** *Best Constant Predictor*
*For $Y \in L^2$, $\arg\min_{c\in\mathbb{R}} \mathbb{E}(Y - c)^2 = \mathbb{E}Y$, and $\min_{c\in\mathbb{R}} \mathbb{E}(Y - c)^2 = \operatorname{Var}(Y)$.*

*Proof.* Let $\mu = \mathbb{E}Y$. Since $\mathbb{E}(Y - \mu) = 0$, $\mathbb{E}(Y - c)^2 = \mathbb{E}[(Y - \mu) + (\mu - c)]^2 = \operatorname{Var}(Y) + (\mu - c)^2$.
Since $(\mu - c)^2 \geq 0$ with equality iff $c = \mu$, the unique minimizer is $c = \mu$.
*Structural Commentary*: The mean is the orthogonal projection of $Y$ onto the constants.
*Key Insight*: Best constant prediction equals the average. □

**Theorem 1.3.5.** *Conditional Expectation as Best Predictor*
*Let $X, Y \in L^2$ and $\mathcal{G} := \{h(X) : h(X) \in L^2\}$. Let $M := \mathbb{E}(Y \mid X)$, then $M \in \mathcal{G}$, $\mathbb{E}(Y - M)^2 \leq \mathbb{E}(Y - Z)^2 \forall Z \in \mathcal{G}$.*

*Proof.* For $Z \in \mathcal{G}$, both $M$ and $Z$ are $\sigma(X)$-measurable.
Since $\mathbb{E}(Y - M \mid X) = 0$, $\mathbb{E}[(Y - M)(M - Z)] = \mathbb{E}[(M - Z)\mathbb{E}(Y - M \mid X)] = 0$.
Hence $\mathbb{E}(Y - Z)^2 = \mathbb{E}(Y - M)^2 + \mathbb{E}(M - Z)^2 \geq \mathbb{E}(Y - M)^2$.
*Structural Commentary*: Prediction from $X$ is performed fiber by fiber over the conditional laws of $Y$.
*Key Insight*: The optimal nonlinear least-squares predictor is the conditional mean. □

**Definition 1.3.6.** *Best Linear Predictor*
For $X, Y \in L^2$, set $\mathcal{A}_X := \{\alpha + \beta X : \alpha, \beta \in \mathbb{R}\}$.
A *best linear predictor* of $Y$ from $X$ is any $\widehat{Y}^* \in \arg\min_{Z\in\mathcal{A}_X} \mathbb{E}(Y - Z)^2$.

**Theorem 1.3.7.** *Best Linear Predictor*
*Let $X, Y \in L^2$, $\sigma_X^2 := \operatorname{Var}(X) > 0$, $\mu_X := \mathbb{E}X$, and $\mu_Y := \mathbb{E}Y$.*

*(i)* $\arg\min_{Z\in\mathcal{A}_X} \mathbb{E}(Y - Z)^2 = \{\alpha^* + \beta^* X\}$, *where* $\beta^* = \frac{\operatorname{Cov}(X,Y)}{\sigma_X^2}$ *and* $\alpha^* = \mu_Y - \beta^*\mu_X$.

*(ii)* $\min_{Z\in\mathcal{A}_X} \mathbb{E}(Y - Z)^2 = \operatorname{Var}(Y) - \frac{\operatorname{Cov}(X,Y)^2}{\sigma_X^2}$.

*(iii) If* $\mathrm{Var}(Y) > 0$ *and* $\rho = \mathrm{Corr}(X, Y)$*, then the minimum error is* $\mathrm{Var}(Y)(1-\rho^2)$.

*Proof.*

(i) Let $q(\alpha, \beta) := \mathbb{E}(Y - \alpha - \beta X)^2$. For fixed $\beta$, $q(\alpha, \beta) = \mathrm{Var}(Y - \beta X) + (\mu_Y - \alpha - \beta\mu_X)^2$, so $\alpha(\beta) = \mu_Y - \beta\mu_X$ uniquely minimizes over $\alpha$.

(ii) Then $q(\alpha(\beta), \beta) = \mathrm{Var}(Y - \beta X) = \mathrm{Var}(Y) + \beta^2\sigma_X^2 - 2\beta\,\mathrm{Cov}(X, Y)$. Since $\sigma_X^2 > 0$, this quadratic has unique minimizer $\beta^* = \frac{\mathrm{Cov}(X,Y)}{\sigma_X^2}$.

(iii) Substituting $\beta^*$ gives $\mathrm{Var}(Y) - \frac{\mathrm{Cov}(X,Y)^2}{\sigma_X^2}$. If $\rho = \frac{\mathrm{Cov}(X,Y)}{\sigma_X\sigma_Y}$, then this equals $\mathrm{Var}(Y)(1-\rho^2)$.

*Structural Commentary*: Linear prediction projects $Y$ onto $\mathcal{A}_X$.
*Key Insight*: The optimal slope is normalized covariance. □

**Example 1.3.8.** *Bivariate Normal Prediction*
If $(X, Y)$ is bivariate normal with means $\mu_X, \mu_Y$, standard deviations $\sigma_X, \sigma_Y > 0$, and correlation $\rho \in (-1, 1)$, then $Y \mid X \sim N(\mu_Y + \rho\sigma_Y\sigma_X^{-1}(X - \mu_X), \sigma_Y^2(1-\rho^2))$.
Thus $\mathbb{E}(Y \mid X) = \mu_Y + \rho\sigma_Y\sigma_X^{-1}(X - \mu_X)$ is linear, with conditional variance $\sigma_Y^2(1-\rho^2)$.

**Example 1.3.9.** *Regression to the Mean*
Let $\mathbb{E}X = \mathbb{E}Y = \mu$, $\mathrm{Var}(X) = \mathrm{Var}(Y) = \sigma^2 > 0$, and $\mathrm{Corr}(X, Y) = \rho$. Best linear predictor is $\widehat{Y} = \mu + \rho(X - \mu)$.
For $0 < \rho < 1$, $|\widehat{Y} - \mu| = \rho|X - \mu| < |X - \mu|$, so predicted scores regress toward $\mu$.

# 2 Probabilistic Convergence

## 2.1 Modes of Probabilistic Convergence

Pointwise convergence is too strict for most random sequences. The useful replacement is to ask whether probabilities, laws, or moments of the errors become small.

**Definition 2.1.1.** *Modes of Probabilistic Convergence*
Let $(X_n)_{n\geq 1}$ and $X$ be real-valued random variables with distribution functions $F_n$ and $F$.

(i) *Convergence in probability*: $X_n \xrightarrow{\mathbb{P}} X$ iff $\mathbb{P}(|X_n - X| > \varepsilon) \to 0$ for every $\varepsilon > 0$.
(ii) *Convergence in distribution*: $X_n \rightsquigarrow X$ iff $F_n(t) \to F(t)$ at every continuity point $t$ of $F$.
(iii) *Convergence in $L^p$*: for $p \geq 1$, $X_n \xrightarrow{L^p} X$ iff $\mathbb{E}\,|X_n - X|^p \to 0$.
(iv) *Convergence in quadratic mean*: $X_n \xrightarrow{qm} X$ iff $X_n \xrightarrow{L^2} X$.
(v) *Almost-sure convergence*: $X_n \xrightarrow{\text{a.s.}} X$ iff $\mathbb{P}(\{\omega \in \Omega : X_n(\omega) \to X(\omega)\}) = 1$.
(vi) *Constant limits*: if $X = c$ almost surely for a constant $c$, then $X_n \xrightarrow{\mathbb{P}} c$, $X_n \rightsquigarrow c$, $X_n \xrightarrow{L^p} c$, $X_n \xrightarrow{qm} c$, and $X_n \xrightarrow{\text{a.s.}} c$ denote convergence to that constant random variable in the corresponding modes.

**Example 2.1.2.** *A Shrinking Normal Law*
Let $X_n \sim N(0, \frac{1}{n})$, and let $F$ be the distribution function of $\delta_0$.
Since $\sqrt{n}X_n \sim N(0, 1)$, $F_n(t) = \mathbb{P}(X_n \leq t) = \Phi(\sqrt{n}t)$, so $F_n(t) \to 0$ for $t < 0$ and $F_n(t) \to 1$ for $t > 0$.
Hence $F_n(t) \to F(t)$ at every continuity point $t \neq 0$ of $F$, and therefore $X_n \rightsquigarrow 0$.

**Theorem 2.1.3.** *Basic Implications Between Convergence Modes*
*Let $X_n$ and $X$ be real-valued random variables with distribution functions $F_n$ and $F$, and let $c \in \mathbb{R}$.*

*(i)* $X_n \xrightarrow{qm} X \Rightarrow X_n \xrightarrow{\mathbb{P}} X$.
*(ii)* $X_n \xrightarrow{\mathbb{P}} X \Rightarrow X_n \rightsquigarrow X$.
*(iii)* $X_n \rightsquigarrow c \Rightarrow X_n \xrightarrow{\mathbb{P}} c$.
*(iv)* $X_n \xrightarrow{\text{a.s.}} X \Rightarrow X_n \xrightarrow{\mathbb{P}} X$.
*(v)* $X_n \xrightarrow{L^1} X \Rightarrow X_n \xrightarrow{\mathbb{P}} X$, *and* $X_n \xrightarrow{qm} X \Rightarrow X_n \xrightarrow{L^1} X$.

*Proof.*

(i) If $X_n \xrightarrow{qm} X$, then for each $\varepsilon > 0$, Markov gives $\mathbb{P}(|X_n - X| > \varepsilon) \leq \varepsilon^{-2}\mathbb{E}\,|X_n - X|^2 \to 0$.

(ii) Let $x$ be a continuity point of $F$.
For each $\varepsilon > 0$, $F(x - \varepsilon) - \mathbb{P}(|X_n - X| > \varepsilon) \leq F_n(x) \leq F(x + \varepsilon) + \mathbb{P}(|X_n - X| > \varepsilon)$,
so $F(x - \varepsilon) \leq \liminf_n F_n(x) \leq \limsup_n F_n(x) \leq F(x + \varepsilon)$; letting $\varepsilon \downarrow 0$ gives $F_n(x) \to F(x)$.

(iii) For $\varepsilon > 0$, the distribution function of $c$ is continuous at $c - \varepsilon$ and $c + \varepsilon$.
Hence $\mathbb{P}(|X_n - c| > \varepsilon) \leq F_n(c - \varepsilon) + 1 - F_n(c + \varepsilon) \to 0$.

(iv) Fix $\varepsilon > 0$ and set $Y_n = \mathbf{1}_{\{|X_n - X| > \varepsilon\}}$. If $X_n \to X$ a.s., then $Y_n \to 0$ a.s. and $0 \le Y_n \le 1$.
Hence dominated convergence gives $\mathbb{P}(|X_n - X| > \varepsilon) = \mathbb{E}Y_n \to 0$.

(v) If $X_n \xrightarrow{L^1} X$, then $\mathbb{P}(|X_n - X| > \varepsilon) \le \varepsilon^{-1}\mathbb{E}\,|X_n - X| \to 0$ by Markov.
If $X_n \xrightarrow{qm} X$, then $\mathbb{E}\,|X_n - X| \le (\mathbb{E}\,|X_n - X|^2)^{\frac{1}{2}} \to 0$ by Cauchy–Schwarz.

*Structural Commentary*: Strong modes give tail bounds; distribution convergence gives c.d.f. control.
*Key Insight*: Convert to tail bounds; use c.d.f. squeezes when laws enter. □




Figure 233: Solid: always; dotted: point-mass limit.

**Example 2.1.4.** *Why the Reverse Arrows Fail*

(i) Let $U \sim \text{Uniform}(0,1)$ and $X_n = \sqrt{n}\mathbf{1}_{(0,\frac{1}{n})}(U)$. Then for each $\varepsilon > 0$, $\mathbb{P}(|X_n| > \varepsilon) \le \frac{1}{n} \to 0$,
so $X_n \xrightarrow{\mathbb{P}} 0$, but $\mathbb{E}X_n^2 = 1$ for all $n$; hence $X_n \not\to 0$ in quadratic mean.

(ii) Let $X \sim N(0,1)$ and set $X_n = -X$. Then $X_n \sim X$ for all $n$, so $X_n \rightsquigarrow X$,
but for every $\varepsilon > 0$, $\mathbb{P}(|X_n - X| > \varepsilon) = \mathbb{P}(2\,|X| > \varepsilon) > 0$ for all $n$; hence $X_n \not\to X$ in probability.

**Remark 2.1.5.** *A Warning About Expectations*
Convergence in probability does not imply convergence of expectations.
If $\mathbb{P}(X_n = n^2) = \frac{1}{n}$, $\mathbb{P}(X_n = 0) = 1 - \frac{1}{n}$, then $\forall \varepsilon > 0$, $\mathbb{P}(|X_n| > \varepsilon) = \frac{1}{n} \to 0$, so $X_n \xrightarrow{\mathbb{P}} 0$, $\mathbb{E}X_n = n \to +\infty$.

**Definition 2.1.6.** *Asymptotic Uniform Integrability*
A sequence $(X_n)$ is *asymptotically uniformly integrable* if $\lim_{M\to\infty} \limsup_{n\to\infty} \mathbb{E}(|X_n|\,\mathbf{1}_{\{|X_n|>M\}}) = 0$.

**Proposition 2.1.7.** *Recovering Expectations from Probability Convergence*
*Let $b \in \mathbb{R}$. If $X_n \xrightarrow{\mathbb{P}} b$ and $(X_n)$ is asymptotically uniformly integrable, then $\mathbb{E}X_n \to b$.*

*Proof.* For $M > |b|$, set $\tau_M(x) = (-M) \vee (x \wedge M)$. As $\tau_M$ bounded, continuous; $\tau_M(b) = b$, $\tau_M(X_n) \xrightarrow{\mathbb{P}} b$.
For $\eta > 0$, $\mathbb{E}\,|\tau_M(X_n) - b| \le \eta + 2M\mathbb{P}(|\tau_M(X_n) - b| > \eta)$, so $\mathbb{E}\tau_M(X_n) \to b$ after $n \to \infty$ and $\eta \downarrow 0$.
Moreover, $|X_n - \tau_M(X_n)| \le |X_n|\,\mathbf{1}_{\{|X_n|>M\}}$, hence $\limsup_n |\mathbb{E}X_n - b| \le \limsup_n \mathbb{E}(|X_n|\,\mathbf{1}_{\{|X_n|>M\}})$.
Letting $M \to \infty$ and using asymptotic uniform integrability gives $\mathbb{E}X_n \to b$.
*Structural Commentary*: Truncation makes convergence bounded; asymptotic uniform integrability kills tails.
*Key Insight*: Probability convergence handles the truncations; tail control restores expectations. □

**Theorem 2.1.8.** *Continuous Mapping and Slutsky Rules*
*Let $g : \mathbb{R} \to \mathbb{R}$ be continuous, and let $c \in \mathbb{R}$.*

*(i)* $X_n \xrightarrow{\mathbb{P}} X \Rightarrow g(X_n) \xrightarrow{\mathbb{P}} g(X)$.
*(ii)* $X_n \rightsquigarrow X \Rightarrow g(X_n) \rightsquigarrow g(X)$.
*(iii)* $X_n \rightsquigarrow X$ *and* $Y_n \xrightarrow{\mathbb{P}} c \Rightarrow X_n + Y_n \rightsquigarrow X + c$ *and* $X_nY_n \rightsquigarrow cX$.
*(iv)* $X_n \xrightarrow{\mathbb{P}} X$ *and* $Y_n \xrightarrow{\mathbb{P}} Y \Rightarrow X_n + Y_n \xrightarrow{\mathbb{P}} X + Y$ *and* $X_nY_n \xrightarrow{\mathbb{P}} XY$.

*Proof.*

(i) Fix $\varepsilon, \eta > 0$. Choose $K$, $\mathbb{P}(|X| > K) < \eta$, $\delta \le 1$ s.t. $|x - y| < \delta$, $|x|, |y| \le K + 1$; $|g(x) - g(y)| < \varepsilon$.
Then $\mathbb{P}(|g(X_n) - g(X)| > \varepsilon) \le \mathbb{P}(|X| > K) + \mathbb{P}(|X_n - X| > \delta)$, so lim sup at most $\eta$.

(ii) Let $t$ be a continuity point of the law of $g(X)$ and set $A_t = \{x : g(x) \le t\}$.
Since $\partial A_t \subseteq \{x : g(x) = t\}$ and $\mathbb{P}(g(X) = t) = 0$, the continuity-set form of weak convergence gives
$\mathbb{P}(g(X_n) \le t) = \mathbb{P}(X_n \in A_t) \to \mathbb{P}(X \in A_t) = \mathbb{P}(g(X) \le t)$.

(iii) If $Z_n \rightsquigarrow Z$ and $R_n \xrightarrow{\mathbb{P}} 0$, then $Z_n + R_n \rightsquigarrow Z$ by the c.d.f. squeeze
$F_{Z_n}(t - \eta) - \mathbb{P}(|R_n| > \eta) \le F_{Z_n+R_n}(t) \le F_{Z_n}(t + \eta) + \mathbb{P}(|R_n| > \eta)$,
followed by $n \to \infty$ and $\eta \downarrow 0$ through continuity points of $F_Z$.
Since $X_n + c \rightsquigarrow X + c$ by (ii), applying this with $Z_n = X_n + c$ and $R_n = Y_n - c$ gives $X_n + Y_n \rightsquigarrow X + c$.
Also $X_n \rightsquigarrow X$ implies $(X_n)$ is tight, so $\mathbb{P}(|X_n(Y_n - c)| > \varepsilon) \le \mathbb{P}(|X_n| > K) + \mathbb{P}(|Y_n - c| > \frac{\varepsilon}{K})$.
Letting $n \to \infty$, then $K \uparrow \infty$ gives $X_n(Y_n - c) \xrightarrow{\mathbb{P}} 0$.
Since $cX_n \rightsquigarrow cX$ by (ii), the additive perturbation just proved gives $X_nY_n = cX_n + X_n(Y_n - c) \rightsquigarrow cX$.

(iv) For sums, $\mathbb{P}(|(X_n+Y_n)-(X+Y)|>\varepsilon)\le\mathbb{P}(|X_n-X|>\frac{\varepsilon}{2})+\mathbb{P}(|Y_n-Y|>\frac{\varepsilon}{2})\to 0$.
For products, $(X_n)$ and $Y$ are tight, $\mathbb{P}(|X_n(Y_n-Y)|>\frac{\varepsilon}{2})\le\mathbb{P}(|X_n|>K)+\mathbb{P}(|Y_n-Y|>\frac{\varepsilon}{2K})$, while $\mathbb{P}(|Y(X_n-X)|>\frac{\varepsilon}{2})\le\mathbb{P}(|Y|>K)+\mathbb{P}(|X_n-X|>\frac{\varepsilon}{2K})$; let $n\to\infty$ and then $K\uparrow\infty$.

*Structural Commentary*: Continuity transports convergence; Slutsky absorbs probability-small errors.
*Key Insight*: Control the bounded high-probability part; let the tails vanish. □

## 2.2 Probability on Metric Spaces

# 3 Limit Theorems

## 3.1 Laws of Large Numbers

The law of large numbers formalizes the idea that a large sample average is close to the mean of the distribution.

**Perspective 3.1.1.** *Law of the Sample Mean*

If $X_1,\dots,X_n$ are IID, mean $\mu$, variance $\sigma^2$, then $\overline{X}_n:=n^{-1}\sum_{i=1}^n X_i$ is a random variable with law $\mathcal{L}(\overline{X}_n)$.

By Theorem 3.2.7, $\mathbb{E}\overline{X}_n=\mu$ and $\mathrm{Var}(\overline{X}_n)=\frac{\sigma^2}{n}$, so $\mathcal{L}(\overline{X}_n)$ concentrates around $\mu$ as $n\to\infty$.

**Theorem 3.1.2.** *Weak Law of Large Numbers*

*Let $(X_i)_{i\ge1}$ be IID real-valued random variables with $\mathbb{E}X_1=\mu$, $\mathrm{Var}(X_1)=\sigma^2<+\infty$. Set $\overline{X}_n:=n^{-1}\sum_{i=1}^n X_i$.*

*Then $\overline{X}_n\xrightarrow{\mathbb{P}}\mu$, equivalently $\mathbb{P}(|\overline{X}_n-\mu|>\varepsilon)\to0$ for every $\varepsilon>0$.*

*Proof.* For $n\ge1$, $\mathbb{E}\overline{X}_n=n^{-1}\sum_{i=1}^n\mathbb{E}X_i=\mu$ and, by independence, $\mathrm{Var}(\overline{X}_n)=n^{-2}\sum_{i=1}^n\mathrm{Var}(X_i)=\frac{\sigma^2}{n}$.
For every $\varepsilon>0$, Chebyshev gives $\mathbb{P}(|\overline{X}_n-\mu|>\varepsilon)\le\frac{\mathrm{Var}(\overline{X}_n)}{\varepsilon^2}=\frac{\sigma^2}{n\varepsilon^2}\to0$, hence $\overline{X}_n\xrightarrow{\mathbb{P}}\mu$.
*Structural Commentary*: Averaging keeps the mean fixed and divides variance by $n$.
*Key Insight*: Chebyshev turns vanishing variance into convergence in probability. □

**Proposition 3.1.3.** *Weak LLN with Vanishing Average Variance*

*Let $(X_i)_{i\ge1}$ be independent, $X_i\in L^2$, with $\mu_i:=\mathbb{E}X_i$ and $\sigma_i^2:=\mathrm{Var}(X_i)$.*
*Set $\overline{X}_n:=n^{-1}\sum_{i=1}^n X_i$ and $m_n:=n^{-1}\sum_{i=1}^n\mu_i$. If $m_n\to\mu$ and $n^{-2}\sum_{i=1}^n\sigma_i^2\to0$, then $\overline{X}_n\xrightarrow{\mathbb{P}}\mu$.*

*Proof.* $\mathbb{E}\overline{X}_n=m_n$. By independence, $\mathrm{Var}(\overline{X}_n)=n^{-2}\sum_{i=1}^n\sigma_i^2\to0$. Fix $\varepsilon>0$. $\forall$ large $n$, $|m_n-\mu|<\frac{\varepsilon}{2}$.
Hence $\{|\overline{X}_n-\mu|>\varepsilon\}\subseteq\{|\overline{X}_n-m_n|>\frac{\varepsilon}{2}\}$. By Chebyshev, $\mathbb{P}(|\overline{X}_n-\mu|>\varepsilon)\le\frac{4\,\mathrm{Var}(\overline{X}_n)}{\varepsilon^2}\to0$.
*Structural Commentary*: Identical distribution is not essential for a weak law.
*Key Insight*: It is enough that the mean target stabilises and the average variance vanishes. □

**Example 3.1.4.** *Coin Proportions and Chebyshev Scale*

Let $X_i\sim\mathrm{Bernoulli}(p)$ be IID head indicators, $\overline{X}_n:=n^{-1}\sum_{i=1}^n X_i$. Then $\mathbb{E}\overline{X}_n=p$, $\mathrm{Var}(\overline{X}_n)=\frac{p(1-p)}{n}$, $\overline{X}_n\xrightarrow{\mathbb{P}}p$.
For $p=\frac12$, Chebyshev gives $\mathbb{P}(0.4\le\overline{X}_n\le0.6)\ge1-\frac{1}{4n(0.1)^2}=1-\frac{25}{n}$, at least 0.7 for $n\ge84$.
If $n=10000$ fair tosses produce 5067 heads, then $\overline{X}_n=0.5067$, while $\mathrm{Var}(\overline{X}_n)=\frac{1}{4n}=2.5\cdot10^{-5}$ and $\mathrm{sd}(\overline{X}_n)=0.005$. The observed deviation is about 1.34 standard deviations from $\frac12$.

**Theorem 3.1.5.** *Strong Law of Large Numbers*

*Let $(X_i)_{i\ge1}$ be IID real-valued random variables with $X_1\in L^1$, $\mu:=\mathbb{E}X_1$, and $\overline{X}_n:=n^{-1}\sum_{i=1}^n X_i$.*

*Then $\overline{X}_n\xrightarrow{\text{a.s.}}\mu$, equivalently $\mathbb{P}(\lim_{n\to\infty}\overline{X}_n=\mu)=1$.*

*Proof.* After replacing $X_i$ by $X_i-\mu$, assume $\mu=0$. Set $Y_i:=X_i\mathbf{1}_{\{|X_i|\le i\}}$.
Since $\sum_{i=1}^\infty\mathbb{P}(X_i\ne Y_i)=\sum_{i=1}^\infty\mathbb{P}(|X_1|>i)<+\infty$, Borel–Cantelli gives $X_i=Y_i$ eventually a.s.
Hence $n^{-1}\sum_{i=1}^n(X_i-Y_i)\to0$ a.s. For $Z_i:=Y_i-\mathbb{E}Y_i$, independence gives
$\sum_{i=1}^\infty\frac{\mathrm{Var}(Z_i)}{i^2}\le\sum_{i=1}^\infty\frac{\mathbb{E}(X_1^2\mathbf{1}_{\{|X_1|\le i\}})}{i^2}<+\infty$, by Kolmogorov's variance criterion, $n^{-1}\sum_{i=1}^n Z_i\to0$ a.s.
By dominated convergence, $\mathbb{E}Y_i\to0$, hence Cesaro averaging gives $n^{-1}\sum_{i=1}^n\mathbb{E}Y_i\to0$.

Combining $X_i - Y_i$, $Z_i$, and $\mathbb{E}Y_i$ gives $\overline{X}_n \to 0$ a.s.; adding back $\mu$ completes the proof.
*Structural Commentary*: Truncate tails; centered truncations have summable variance.
*Key Insight*: Replace by an eventually equal truncated sequence, then average the centered part. □

**Corollary 3.1.6.** *Strong Implies Weak*
*Let $(X_i)_{i\geq 1}$ be IID real-valued random variables, $X_1 \in L^1$, $\mu := \mathbb{E}X_1$, $\overline{X}_n := n^{-1}\sum_{i=1}^{n} X_i$. Then $\overline{X}_n \xrightarrow{\mathbb{P}} \mu$.*

*Proof.* By Theorem 3.1.5, $\overline{X}_n \to \mu$ a.s. For $\varepsilon > 0$, set $Y_n := \mathbf{1}_{\{|\overline{X}_n - \mu| > \varepsilon\}}$.
Then $Y_n \to 0$ a.s. and $0 \leq Y_n \leq 1$, so dominated convergence gives $\mathbb{P}(|\overline{X}_n - \mu| > \varepsilon) = \mathbb{E}Y_n \to 0$.
*Structural Commentary*: Almost-sure convergence gives pointwise tail indicators tending to zero.
*Key Insight*: Dominate the indicators by 1 and integrate. □

**Proposition 3.1.7.** *Consistency of Empirical Moments and Variance*
*Let $(X_i)_{i\geq 1}$ be iid, $X_1 \in L^2$, $\mu := \mathbb{E}X_1$, and $\sigma^2 := \mathrm{Var}(X_1)$.*
*For $n \geq 2$, set $\overline{X}_n := n^{-1}\sum_{i=1}^{n} X_i$ and $S_n^2 := (n-1)^{-1}\sum_{i=1}^{n}(X_i - \overline{X}_n)^2$. Then $S_n^2 \xrightarrow{\mathbb{P}} \sigma^2$.*
*More generally, if $X_1 \in L^r$ for an integer $r \geq 1$, then $n^{-1}\sum_{i=1}^{n} X_i^r \xrightarrow{\mathbb{P}} \mathbb{E}X_1^r$.*

*Proof.* Since $(X_i^r)_{i\geq 1}$ is iid and integrable, the strong law gives $n^{-1}\sum_{i=1}^{n} X_i^r \to \mathbb{E}X_1^r$ a.s., hence in probability.
Let $m_2 := \mathbb{E}X_1^2 = \sigma^2 + \mu^2$. The cases $r = 1, 2$ give $\overline{X}_n \to \mu$ and $n^{-1}\sum_{i=1}^{n} X_i^2 \to m_2$ a.s.
Since $S_n^2 = \frac{n}{n-1}(n^{-1}\sum_{i=1}^{n} X_i^2 - \overline{X}_n^2)$, continuous mapping gives $S_n^2 \to m_2 - \mu^2 = \sigma^2$ a.s., hence in probability.
*Structural Commentary*: Sample moments are averages of transformed observations.
*Key Insight*: Apply the strong law to $X_i^r$, then pass through the continuous variance formula. □

**Remark 3.1.8.** *Why Integrability Matters*
If $X_1 \notin L^1$, the mean $\mathbb{E}X_1$ need not exist and sample averages need not converge to any finite deterministic limit; integrability is the condition that makes a law of large numbers meaningful.

## 3.2 Central Limit Theory

The law of large numbers says that $\overline{X}_n$ is close to $\mu$ for large $n$. The central limit theorem describes the scale and shape of the remaining fluctuation.

**Theorem 3.2.1.** *Central Limit Theorem*
*Let $(X_i)_{i\geq 1}$ be IID real-valued random variables, $\mathbb{E}X_1 = \mu$, $\mathrm{Var}(X_1) = \sigma^2 \in (0, +\infty)$.*
*Let $\overline{X}_n := n^{-1}\sum_{i=1}^{n} X_i$, $Z_n := \sigma^{-1}\sqrt{n}(\overline{X}_n - \mu)$.*
*Then $Z_n \rightsquigarrow N(0,1)$, equivalently $\mathbb{P}(Z_n \leq z) \to \Phi(z) := \int_{-\infty}^{z}(2\pi)^{-\frac{1}{2}}e^{-\frac{x^2}{2}}\,dx$ $\forall z \in \mathbb{R}$.*

*Proof.* Set $Y_i := \frac{X_i - \mu}{\sigma}$, so $\mathbb{E}Y_i = 0$, $\mathrm{Var}(Y_i) = 1$, and $Z_n = n^{-\frac{1}{2}}\sum_{i=1}^{n} Y_i$.
For $\varphi(t) := \mathbb{E}e^{itY_1}$, the finite second moment expansion gives $\varphi(u) = 1 - \frac{u^2}{2} + o(u^2)$ as $u \to 0$.
By independence, $\varphi_{Z_n}(t) = [\varphi(\frac{t}{\sqrt{n}})]^n = [1 - \frac{t^2}{2n} + o(\frac{1}{n})]^n \to e^{-\frac{t^2}{2}}$ for every $t \in \mathbb{R}$.
Since $e^{-\frac{t^2}{2}}$ is the characteristic function of $N(0,1)$, Levy's continuity theorem gives $Z_n \rightsquigarrow N(0,1)$; continuity of $\Phi$ gives the displayed c.d.f. convergence.
*Structural Commentary*: Centering and scaling turn the sum into unit-variance IID increments.
*Key Insight*: Characteristic functions turn normalized sums into powers, whose limit is Gaussian. □

**Method 3.2.2.** *Equivalent CLT Approximation Forms*
Let $\overline{X}_n := n^{-1}\sum_{i=1}^{n} X_i$ with mean $\mu$ and variance $\sigma^2 \in (0, +\infty)$, and set $Z_n := \sigma^{-1}\sqrt{n}(\overline{X}_n - \mu)$.

(i) Standardized form: $Z_n \approx N(0,1)$.
(ii) Centered form: $\sqrt{n}(\overline{X}_n - \mu) \approx N(0, \sigma^2)$.
(iii) Sample-mean form: $\overline{X}_n \approx N(\mu, \frac{\sigma^2}{n})$.

Use any equivalent form to approximate probabilities involving $\overline{X}_n$.

**Example 3.2.3.** *Measurement Error and Sample Size*
Let $X_1, \dots, X_n$ be independent measurements with $\mathbb{E}X_i = \mu$ and $\mathrm{Var}(X_i) = \sigma^2$.
For $c > 0$, the CLT gives $\mathbb{P}(|\overline{X}_n - \mu| < c) \approx \Phi(\frac{c\sqrt{n}}{\sigma}) - \Phi(\frac{-c\sqrt{n}}{\sigma})$.
For $(n, \sigma, c) = (16, 1, 0.5)$, this is $\Phi(2) - \Phi(-2) \approx 0.954$.

**Lemma 3.2.4.** *MGF Continuity Theorem*
*Let $Z_n, Z$ be real-valued with MGFs $\psi_n(t) := \mathbb{E}e^{tZ_n}$ and $\psi(t) := \mathbb{E}e^{tZ}$ finite on $(-a, a)$ for some $a > 0$. If $\psi_n(t) \to \psi(t)$ for all $t \in (-a, a)$, then $Z_n \rightsquigarrow Z$.*

*Proof.* For each $s \in \mathbb{R}$, the characteristic functions satisfy $\varphi_n(s) = \mathbb{E}e^{\mathrm{i}sZ_n} = \psi_n(\mathrm{i}s)$ by analytic continuation on the strip $\{z : |\mathrm{Re}\, z| < a\}$. Pointwise MGF convergence near 0 gives $\varphi_n(s) \to \varphi(s)$, the characteristic function of $Z$; Levy's continuity theorem yields $Z_n \rightsquigarrow Z$.
*Structural Commentary*: Local MGF convergence determines local characteristic-function convergence.
*Key Insight*: MGFs near 0 identify the weak limit. □

**Proposition 3.2.5.** *MGF Proof of the IID CLT*
*Let $(X_i)_{i\geq 1}$ be iid with $\mathbb{E}X_1 = 0$, $\mathrm{Var}(X_1) = \sigma^2 \in (0, +\infty)$, and MGF $M(t) := \mathbb{E}e^{tX_1}$ finite near 0.*
*Set $S_n := \sum_{i=1}^{n} X_i$ and $Z_n := \frac{S_n}{\sigma\sqrt{n}}$. Then $Z_n \rightsquigarrow N(0,1)$, equivalently $\mathbb{P}(Z_n \leq x) \to \Phi(x)$ for all $x \in \mathbb{R}$.*

*Proof.* For fixed $t$ near 0, independence gives $M_{Z_n}(t) = \mathbb{E}e^{\frac{tS_n}{\sigma\sqrt{n}}} = \prod_{i=1}^{n} \mathbb{E}e^{\frac{tX_i}{\sigma\sqrt{n}}} = [M(\frac{t}{\sigma\sqrt{n}})]^n$.
Since $M(0) = 1$, $M'(0) = 0$, and $M''(0) = \sigma^2$, Taylor expansion gives $M(s) = 1 + \frac{\sigma^2 s^2}{2} + o(s^2)$ as $s \to 0$.
Hence $M_{Z_n}(t) = (1 + \frac{t^2}{2n} + o(\frac{1}{n}))^n \to e^{\frac{t^2}{2}}$, the MGF of $N(0,1)$.
Lemma 3.2.4 gives $Z_n \rightsquigarrow N(0,1)$, and continuity of $\Phi$ gives the c.d.f. convergence.
*Structural Commentary*: The MGF proof needs the stronger near-zero MGF assumption.
*Key Insight*: Standardization gives $\frac{t^2}{2n}$; independence exponentiates. □

**Example 3.2.6.** *A Poisson Error-Count Approximation*
Let $X_1, \dots, X_{125} \sim \mathrm{Poisson}(5)$ be IID and set $\overline{X}_{125} = 125^{-1} \sum_{i=1}^{125} X_i$.
Since $\mu = 5$ and $\sigma^2 = 5$, the CLT gives $\mathbb{P}(\overline{X}_{125} < 5.5) \approx \Phi(\frac{\sqrt{125}(5.5-5)}{\sqrt{5}}) = \Phi(2.5) \approx 0.9938$.

**Example 3.2.7.** *Normal Approximation to a Poisson Count*
Let $X_\lambda \sim \mathrm{Poisson}(\lambda)$, $Z_\lambda := \frac{X_\lambda - \lambda}{\sqrt{\lambda}}$. Then $M_{Z_\lambda}(t) = \exp(-t\sqrt{\lambda} + \lambda(e^{\frac{t}{\sqrt{\lambda}}} - 1))$ and $\log M_{Z_\lambda}(t) = \frac{t^2}{2} + o(1)$, so $Z_\lambda \rightsquigarrow N(0,1)$. If $X \sim \mathrm{Poisson}(900)$, then $\mathbb{P}(X > 950) = \mathbb{P}(\frac{X-900}{30} > \frac{5}{3}) \approx 1 - \Phi(\frac{5}{3}) \approx 0.04779$, close to 0.04712.

**Example 3.2.8.** *Normal Approximation to the Binomial*
If $X \sim \mathrm{Binomial}(n, p)$, $np > 5$, $n(1-p) > 5$, then $X \approx N(np, np(1-p))$; $\frac{X-np}{\sqrt{np(1-p)}} \approx N(0,1)$.
For $X \sim \mathrm{Binomial}(100, \frac{1}{2})$, $\mathbb{P}(X \geq 60) \approx 1 - \Phi(\frac{60-50}{5}) = 1 - \Phi(2) \approx 0.0228$.

**Example 3.2.9.** *Skewed Summands Need More Averaging*
Let $E_1, \dots, E_n$ be iid Exp(1) and $G_n := \sum_{i=1}^{n} E_i \sim \mathrm{Gamma}(n, 1)$. Then $\frac{G_n - n}{\sqrt{n}} \rightsquigarrow N(0,1)$.
Skewness makes the normal approximation slower for small $n$.

**Example 3.2.10.** *Products and Lognormal Shape*
Let $Y_n = y_0 \prod_{i=1}^{n} X_i$ with iid $X_i > 0$, and suppose $\mu_\ell := \mathbb{E}\log X_1$ and $\sigma_\ell^2 := \mathrm{Var}(\log X_1) \in (0, +\infty)$.
Since $\log Y_n = \log y_0 + \sum_{i=1}^{n} \log X_i$, the CLT gives $\log Y_n \approx N(\log y_0 + n\mu_\ell, n\sigma_\ell^2)$, so $Y_n$ is approximately lognormal.
Likewise $V_n = v_0 \prod_{i=1}^{n} R_i$ when iid log-returns $\log R_i$ have finite positive variance.

**Theorem 3.2.11.** *Studentized Central Limit Theorem*
*Let $(X_i)_{i\geq 1}$ be IID real-valued random variables with $\mathbb{E}X_1 = \mu$ and $\mathrm{Var}(X_1) = \sigma^2 \in (0, +\infty)$.*
*Set $\overline{X}_n := n^{-1} \sum_{i=1}^{n} X_i$, $S_n^2 := (n-1)^{-1} \sum_{i=1}^{n} (X_i - \overline{X}_n)^2$, and $T_n := S_n^{-1}\sqrt{n}(\overline{X}_n - \mu)$. Then $T_n \rightsquigarrow N(0,1)$.*

*Proof.* The CLT gives $Z_n := \sigma^{-1}\sqrt{n}(\overline{X}_n - \mu) \rightsquigarrow N(0,1)$, and Proposition 3.1.7 gives $S_n^2 \xrightarrow{\mathbb{P}} \sigma^2$, hence $S_n \xrightarrow{\mathbb{P}} \sigma$.
Thus $\frac{\sigma}{S_n} \xrightarrow{\mathbb{P}} 1$, and Slutsky's rule from Theorem 2.1.8 gives $T_n = Z_n(\frac{\sigma}{S_n}) \rightsquigarrow N(0,1)$.
*Structural Commentary*: Studentization replaces the unknown $\sigma$ by the consistent estimate $S_n$.
*Key Insight*: Slutsky permits multiplying the CLT statistic by $\frac{\sigma}{S_n} \to 1$. □

**Theorem 3.2.12.** *Berry–Esseen Rate*
*Let $(X_i)_{i\geq 1}$ be IID real-valued random variables, $\mathbb{E}X_1 = \mu$, $\mathrm{Var}(X_1) = \sigma^2 \in (0, +\infty)$, $\rho_3 := \mathbb{E}\,|X_1 - \mu|^3 < +\infty$. Let $\overline{X}_n := n^{-1}\sum_{i=1}^{n} X_i$, $Z_n := \sigma^{-1}\sqrt{n}(\overline{X}_n - \mu)$, $\Phi(z) := \int_{-\infty}^{z}(2\pi)^{-\frac{1}{2}}e^{-\frac{x^2}{2}}\,dx$; $\sup_{z\in\mathbb{R}}|\mathbb{P}(Z_n \leq z) - \Phi(z)| \leq \frac{33}{4}\frac{\rho_3}{\sigma^3\sqrt{n}}$.*

*Proof.* Set $Y_i := \frac{X_i - \mu}{\sigma}$, $\rho := \mathbb{E}\,|Y_1|^3 = \frac{\rho_3}{\sigma^3}$, and $W_n := n^{-\frac{1}{2}}\sum_{i=1}^{n} Y_i = Z_n$.
For $\varphi(t) := \mathbb{E}e^{itY_1}$, Taylor's formula with $\mathbb{E}Y_1 = 0$, $\mathbb{E}Y_1^2 = 1$, $\mathbb{E}\,|Y_1|^3 = \rho$ gives $\varphi(t) = 1 - \frac{t^2}{2} + O(\rho\,|t|^3)$ near 0.
Thus $\varphi_{W_n}(t) = [\varphi(\frac{t}{\sqrt{n}})]^n$ differs from $e^{-\frac{t^2}{2}}$ by $O(\rho n^{-\frac{1}{2}})$ in Esseen's smoothing integral.
Standard optimised smoothing bound gives $\sup_z |\mathbb{P}(W_n \leq z) - \Phi(z)| \leq \frac{33}{4}\frac{\rho}{\sqrt{n}}$. Substitute $\rho = \frac{\rho_3}{\sigma^3}$, QED.
*Structural Commentary*: Berry–Esseen bounds the CLT error by standardised third absolute moment.
*Key Insight*: Normalise, compare characteristic functions, then apply Esseen smoothing. □

**Theorem 3.2.13.** *Multivariate Central Limit Theorem*
*Let $(X_i)_{i\geq 1}$ be IID random vectors in $\mathbb{R}^k$ with $\mathbb{E}X_1 = \mu \in \mathbb{R}^k$ and $\mathrm{Cov}(X_1) = \Sigma$, and set $\overline{X}_n := n^{-1}\sum_{i=1}^{n} X_i$.*
*Then $\sqrt{n}(\overline{X}_n - \mu) \rightsquigarrow N_k(0, \Sigma)$.*

*Proof.* For $a \in \mathbb{R}^k$, set $Y_{i,a} := a^t(X_i - \mu)$ and $s_a^2 := a^t\Sigma a$.
Then $(Y_{i,a})_{i\geq 1}$ are IID, $\mathbb{E}Y_{1,a} = 0$, $\mathrm{Var}(Y_{1,a}) = s_a^2$, and $a^t\sqrt{n}(\overline{X}_n - \mu) = n^{-\frac{1}{2}}\sum_{i=1}^{n} Y_{i,a}$.
If $s_a^2 > 0$, the scalar CLT gives $a^t\sqrt{n}(\overline{X}_n - \mu) \rightsquigarrow N(0, s_a^2)$; if $s_a^2 = 0$, then $\mathbb{E}Y_{1,a}^2 = 0$, so the same convergence holds with degenerate limit $N(0,0)$. Since $a^tZ \sim N(0, s_a^2)$ for $Z \sim N_k(0, \Sigma)$, by Cramer–Wold, QED.
*Structural Commentary*: Multivariate convergence is checked through one-dimensional projections.
*Key Insight*: Project first, apply the scalar CLT, then use Cramer–Wold. □

**Theorem 3.2.14.** *Delta Method*
*Let $(Y_n)_{n\geq 1}$ be real-valued random variables, let $\mu \in \mathbb{R}$ and $\sigma > 0$, and assume $\sigma^{-1}\sqrt{n}(Y_n - \mu) \rightsquigarrow N(0,1)$.*
*If $g : \mathbb{R} \to \mathbb{R}$ is differentiable at $\mu$ and $g'(\mu) \neq 0$, then $(|g'(\mu)|\,\sigma)^{-1}\sqrt{n}(g(Y_n) - g(\mu)) \rightsquigarrow N(0,1)$.*
*Equivalently $g(Y_n) \approx N(g(\mu), \frac{(g'(\mu))^2\sigma^2}{n})$.*

*Proof.* Set $A_n := \sigma^{-1}\sqrt{n}(Y_n - \mu)$ and define $\rho(\mu) := 0$, $\rho(y) := \frac{g(y)-g(\mu)}{y-\mu} - g'(\mu)$ for $y \neq \mu$.
Then $A_n \rightsquigarrow Z \sim N(0,1)$, $A_n = O_{\mathbb{P}}(1)$, and $Y_n - \mu = \frac{\sigma A_n}{\sqrt{n}} \xrightarrow{\mathbb{P}} 0$.
Differentiability gives $\rho(y) \to 0$ as $y \to \mu$, hence $\rho(Y_n) \xrightarrow{\mathbb{P}} 0$.
For all $n$, $g(Y_n) - g(\mu) = (g'(\mu) + \rho(Y_n))(Y_n - \mu)$. Thus, with $T_n := (|g'(\mu)|\,\sigma)^{-1}\sqrt{n}(g(Y_n) - g(\mu))$,
$T_n = \mathrm{sgn}(g'(\mu))A_n(1 + \frac{\rho(Y_n)}{g'(\mu)}) \rightsquigarrow \mathrm{sgn}(g'(\mu))Z \sim N(0,1)$ by Theorem 2.1.8.
*Structural Commentary*: The delta method is first-order Taylor expansion at the limit point.
*Key Insight*: The derivative rescales the fluctuation; the remainder is negligible in probability. □

**Theorem 3.2.15.** *Multivariate Delta Method*
*Let $(Y_n)_{n\geq 1}$ be random vectors in $\mathbb{R}^k$ with $\sqrt{n}(Y_n - \mu) \rightsquigarrow N_k(0, \Sigma)$, and let $g : \mathbb{R}^k \to \mathbb{R}$ be differentiable at $\mu$ with gradient $\nabla g(\mu)$. Then $\sqrt{n}(g(Y_n) - g(\mu)) \rightsquigarrow N(0, \nabla g(\mu)^t\Sigma\nabla g(\mu))$.*

*Proof.* Set $A_n := \sqrt{n}(Y_n - \mu)$, so $A_n \rightsquigarrow N_k(0, \Sigma)$ and $Y_n \xrightarrow{\mathbb{P}} \mu$.
Differentiability gives $g(Y_n) - g(\mu) = \nabla g(\mu)^t(Y_n - \mu) + r(Y_n)\,\|Y_n - \mu\|$ with $r(Y_n) \xrightarrow{\mathbb{P}} 0$.
Since $A_n = O_{\mathbb{P}}(1)$, $\sqrt{n}r(Y_n)\,\|Y_n - \mu\| = r(Y_n)\,\|A_n\| \xrightarrow{\mathbb{P}} 0$.
Thus $\sqrt{n}(g(Y_n) - g(\mu)) = \nabla g(\mu)^t A_n + o_{\mathbb{P}}(1) \rightsquigarrow N(0, \nabla g(\mu)^t\Sigma\nabla g(\mu))$ by Theorem 2.1.8.
*Structural Commentary*: Multivariate delta method linearizes $g$ by its gradient.
*Key Insight*: The gradient projection carries the Gaussian limit; the differentiability remainder vanishes. □

**Proposition 3.2.16.** *Delta Method*
*If $\sqrt{n}(T_n - \theta) \Rightarrow N(0, \Sigma)$ and $g : \mathbb{R}^d \to \mathbb{R}^k$ is differentiable at $\theta$ with derivative $Dg(\theta)$, then $\sqrt{n}(g(T_n) - g(\theta)) \Rightarrow N(0, Dg(\theta)\Sigma Dg(\theta)^\top)$.*

*Proof.* Differentiability gives $g(T_n) - g(\theta) = Dg(\theta)(T_n - \theta) + r_n$ with $\frac{\|r_n\|}{\|T_n - \theta\|} \to 0$ in probability.
Since $\sqrt{n}(T_n - \theta) = O_P(1)$, $\sqrt{n}r_n = o_P(1)$; $\sqrt{n}(g(T_n) - g(\theta)) = Dg(\theta)\sqrt{n}(T_n - \theta) + o_P(1)$, by Slutsky QED.
*Structural Commentary*: The derivative transports the limiting normal fluctuation through $g$.
*Key Insight*: First-order linearisation determines asymptotic variance. □

**Example 3.2.17.** *Exponentiating a Sample Mean*
Let $(X_i)_{i=1}^n$ be IID with mean $\mu$ and variance $\sigma^2 < +\infty$, and set $W_n := e^{\overline{X}_n}$.
For $g(s) = e^s$, $g'(\mu) = e^\mu$, so the delta method gives $W_n \approx N(e^\mu, \frac{e^{2\mu}\sigma^2}{n})$.

**Example 3.2.18.** *Product and Ratio of Sample Means*
Let $X_i = (X_{1i}, X_{2i})^t$ be IID with mean $\mu = (\mu_1, \mu_2)^t$ and covariance $\Sigma = (\sigma_{ij})$, and set $\overline{X}_{jn} := n^{-1} \sum_{i=1}^n X_{ji}$.
For $g(x_1, x_2) = x_1 x_2$, $\nabla g(\mu) = (\mu_2, \mu_1)^t$, so $\sqrt{n}(\overline{X}_{1n}\overline{X}_{2n} - \mu_1\mu_2) \rightsquigarrow N(0, \mu_2^2\sigma_{11} + 2\mu_1\mu_2\sigma_{12} + \mu_1^2\sigma_{22})$.
If $\mu_2 \neq 0$, then for $h(x_1, x_2) = \frac{x_1}{x_2}$ and $a := \nabla h(\mu) = (\frac{1}{\mu_2}, -\frac{\mu_1}{\mu_2^2})^t$, $\sqrt{n}(\frac{\overline{X}_{1n}}{\overline{X}_{2n}} - \frac{\mu_1}{\mu_2}) \rightsquigarrow N(0, a^t\Sigma a)$.

# 4 Stochastic Processes

## 4.1 Process-Level Probability

**Definition 4.1.1.** *Stochastic Process, State Space, and Index Set*

(i) A *stochastic process* is a family $(X_t)_{t\in T}$ of random variables.
(ii) The set $T$ is the *index set*, often representing time.
(iii) Each $X_t$ takes values in a *state space* $\mathcal{X}$.

Typical choices are discrete time $T = \{0, 1, 2, \ldots\}$ and continuous time $T = [0, \infty)$.

**Example 4.1.2.** *IID Sequences as Processes*
An IID sequence $(X_n)_{n\geq 1}$ is a stochastic process indexed by $T = \{1, 2, \ldots\}$.

**Example 4.1.3.** *Discrete States and Discrete Time*
If $\mathcal{X} = \{\text{sunny}, \text{cloudy}\}$ and $X_n$ is the weather on day $n$, then $(X_n)_{n\geq 1}$ is a stochastic process with discrete state space and discrete index set.

**Example 4.1.4.** *The Empirical CDF as a Process*
Let $X_1, \ldots, X_n$ be IID with CDF $F$ on $[0, 1]$, and define $\widehat{F}_n(t) = n^{-1} \sum_{i=1}^n \mathbf{1}\{X_i \leq t\}$.
Then $(\widehat{F}_n(t))_{t\in[0,1]}$ is a stochastic process with continuous index set $[0, 1]$ and state space $[0, 1]$.

**Remark 4.1.5.** *Sample Paths*
A *sample path* is one realized function $t \mapsto X_t(\omega)$ of a stochastic process.
The state space and index set may each be discrete or continuous.

**Proposition 4.1.6.** *Finite-Dimensional Chain Rule*
*Let $X_{1:n} = (X_1, \ldots, X_n)$ have joint mass function or density $f_{1:n}$.*
*At points where the needed marginals are positive, $f_{1:n}(x_{1:n}) = f_1(x_1) \prod_{i=2}^n f_{i|1:i-1}(x_i \mid x_{1:i-1})$.*

*Proof.* For $i \geq 2$, the definition of conditional density gives $f_{1:i}(x_{1:i}) = f_{i|1:i-1}(x_i \mid x_{1:i-1}) f_{1:i-1}(x_{1:i-1})$.
Iterating this identity from $i = n$ down to 2 yields the product formula.
*Structural Commentary*: The joint law factors by revealing one coordinate at a time.
*Key Insight*: Conditional densities convert successive joint marginals into a telescoping product. □

## 4.2 Filtrations and Adapted Processes

## 4.3 Martingales in Discrete and Continuous Time

## 4.4 Markov Chains and Markov Processes

**Definition 4.4.1.** *Probability Vectors and Markov Matrices*
A vector $\mathbf{x} \in \mathbb{R}^n$ is a *probability vector* if its entries are nonnegative and sum to 1. Let $\mathbf{e} := (1, \ldots, 1)^t$. A matrix $A = [a_{ij}] \in \mathrm{Mat}(n, n, \mathbb{R})$ is a *Markov matrix* if $a_{ij} \geq 0$ for all $i, j$ and $A\mathbf{e} = \mathbf{e}$; equivalently, every row sum is 1.

**Lemma 4.4.2.** *Basic Properties of Markov Matrices*
*If $A \in \mathrm{Mat}(n, n, \mathbb{R})$ is Markov, then $1 \in \mathrm{spec}(A)$, and each $A^p$ is Markov with entries in $[0, 1]$.*

*Proof.* Since $A\mathbf{e} = \mathbf{e}$, 1 is an eigenvalue. Inductively, $A^p$ has nonnegative entries and $A^p\mathbf{e} = \mathbf{e}$, so $A^p$ is Markov.
Each nonnegative row sums to 1, so all entries lie in $[0, 1]$.
*Structural Commentary*: Markov structure is closed under powers.
*Key Insight*: Nonnegative row sums of 1 bound every entry. □

**Theorem 4.4.3.** *Unit-Circle Spectrum of Markov Matrices*
*If $A \in \mathrm{Mat}(n,n,\mathbb{R})$ is Markov, then $\rho(A) = 1$, and $\dim E_\lambda(A) = m_A(\lambda)$ for every $\lambda \in \mathrm{spec}(A)$ with $|\lambda| = 1$.*

*Proof.* Lemma 4.4.2 makes $A$ power bounded. Theorem 3.3.7 gives $\rho(A) \le 1$ and $\dim E_\lambda(A) = m_A(\lambda)$ on $|\lambda| = 1$. Since $A\mathbf{e} = \mathbf{e}$, $1 \in \mathrm{spec}(A)$, so $\rho(A) \ge 1$. Thus $\rho(A) = 1$.
*Structural Commentary*: Bounded Markov powers rule out unit-circle Jordan growth.
*Key Insight*: $A\mathbf{e} = \mathbf{e}$ forces the eigenvalue 1. □

**Theorem 4.4.4.** *Positive Markov Matrices Converge to a Stationary Distribution*
*Let $n \ge 2$, $\mathbf{e} := (1,\dots,1)^t$, and $A = (a_{ij}) \in \mathrm{Mat}(n,n,\mathbb{R})$ satisfy $a_{ij} > 0$ and $A\mathbf{e} = \mathbf{e}$.*
*Let $\Delta_{n-1} := \{\mathbf{u} \in \mathbb{R}^n : \mathbf{u} \ge 0,\ \mathbf{e}^t\mathbf{u} = 1\}$. Then $\exists!\, \mathbf{x} \in \Delta_{n-1}$ s.t. $A^t\mathbf{x} = \mathbf{x}$, $\mathbf{x} > 0$, $\lim_{p\to\infty} A^p = \mathbf{e}\mathbf{x}^t$.*

*Proof.* Let $A\mathbf{y} = \lambda\mathbf{y}$ with $\mathbf{y} \ne 0$ and $|\lambda| = 1$, and choose $k$ with $|y_k| = \|\mathbf{y}\|_\infty$.
Since $\sum_j a_{kj} = 1$, then $\|\mathbf{y}\|_\infty = |\lambda y_k| = \left|\sum_{j=1}^n a_{kj} y_j\right| \le \sum_{j=1}^n a_{kj}\,|y_j| \le \|\mathbf{y}\|_\infty$. Thus all inequalities are equalities.
Since $a_{kj} > 0$, $|y_j| = \|\mathbf{y}\|_\infty\ \forall j$; equality in the triangle inequality gives $\mathbf{y} = c\mathbf{e}$, $c \ne 0$.
Hence $\lambda = 1$, $E_1(A) = \mathrm{span}\{\mathbf{e}\}$, and $\mathrm{spec}(A) \cap \{z \in \mathbb{C} : |z| = 1\} = \{1\}$.
Theorem 4.4.3 makes 1 semisimple, so $A = S([1] \oplus J)S^{-1}$ with $S = [\mathbf{e}\ Y]$ and $\rho(J) < 1$.
First row of $S^{-1}$ is $\mathbf{x}^t$. Then $\mathbf{x}^t A = \mathbf{x}^t$, $\mathbf{x}^t\mathbf{e} = 1$, and $A^p = S([1] \oplus J^p)S^{-1} \longrightarrow S([1] \oplus 0)S^{-1} = \mathbf{e}\mathbf{x}^t$.
Since $A^p \ge 0$, the limit gives $\mathbf{x} \ge 0$; since $A^t\mathbf{x} = \mathbf{x}$, $\mathbf{x}^t\mathbf{e} = 1$, and $A^t > 0$, we get $\mathbf{x} > 0$.
Finally, $\dim E_1(A^t) = n - \mathrm{rank}(A^t - I_n) = n - \mathrm{rank}(A - I_n) = \dim E_1(A) = 1$, so any $\mathbf{u} \in \Delta_{n-1}$ with $A^t\mathbf{u} = \mathbf{u}$ satisfies $\mathbf{u} = c\mathbf{x}$ and $\mathbf{e}^t\mathbf{u} = 1 = \mathbf{e}^t\mathbf{x}$, hence $\mathbf{u} = \mathbf{x}$.
*Structural Commentary*: Positivity leaves one simple unit-circle block; all other blocks decay.
*Key Insight*: The limit is the rank-one projection $\mathbf{e}\mathbf{x}^t$. □




Figure 234: The line span($\mathbf{e}$) remains; all other Jordan directions decay, so $A^p \to \mathbf{e}\mathbf{x}^t$.

The preceding results are matrix facts. In probability language, a row-stochastic matrix becomes a transition matrix for a discrete-time chain.

**Definition 4.4.5.** *Markov Chain and Transition Matrix*
(i) Let $\mathcal{X}$ be countable and let $(X_n)_{n\ge0}$ take values in $\mathcal{X}$.
(ii) Process is *Markov chain* if $\mathbb{P}(X_n = x \mid X_0,\dots,X_{n-1}) = \mathbb{P}(X_n = x \mid X_{n-1})$ $\forall$ admissible $n$ and $x \in \mathcal{X}$.
(iii) It is *homogeneous* if $p_{ij} := \mathbb{P}(X_{n+1} = j \mid X_n = i)$ is independent of $n$.
(iv) The matrix $P = [p_{ij}]$ is the *transition matrix*; $p_{ij} \ge 0$ and $\sum_{j\in\mathcal{X}} p_{ij} = 1$ for each $i$.




Figure 235: A Markov chain remembers the current state, not the full past.

**Example 4.4.6.** *Random Walk with Absorbing Barriers*
Let $\mathcal{X} = \{1,\dots,N\}$ and $q = 1 - p$. The absorbing random walk has $p_{11} = p_{NN} = 1$, $p_{i,i+1} = p$ and $p_{i,i-1} = q$ for $2 \le i \le N-1$, and all other transition probabilities equal to 0.

**Definition 4.4.7.** *n-Step Transition Probabilities*
For a homogeneous Markov chain, the *n-step transition probability* from $i$ to $j$ is $p_{ij}(n) := \mathbb{P}(X_{m+n} = j \mid X_m = i)$; set $P_n = [p_{ij}(n)]$.

**Theorem 4.4.8.** *Chapman-Kolmogorov Equations*
*Let $(X_r)_{r\ge0}$ be homogeneous Markov chain on countable $\mathcal{X}$. Let $p_{ij}(r) := \mathbb{P}_i(X_r = j)$, $P_r = (p_{ij}(r))_{i,j\in\mathcal{X}}$.*
*$\forall i,j \in \mathcal{X}$ and $m,n \in \mathbb{N}_0$, $p_{ij}(m+n) = \sum_{k\in\mathcal{X}} p_{ik}(m)p_{kj}(n)$; equivalently, $P_{m+n} = P_mP_n$ and $P_n = P^n$ for $P = P_1$.*

*Proof.* Fix $i, j \in \mathcal{X}$ and $m, n \in \mathbb{N}_0$. By total probability, the Markov property, and homogeneity, $p_{ij}(m+n) = \mathbb{P}_i(X_{m+n} = j) = \sum_{k \in \mathcal{X}} \mathbb{P}_i(X_{m+n} = j \mid X_m = k)\mathbb{P}_i(X_m = k) = \sum_{k \in \mathcal{X}} p_{kj}(n)p_{ik}(m)$.
Thus $(P_m P_n)_{ij} = \sum_{k \in \mathcal{X}} p_{ik}(m)p_{kj}(n) = p_{ij}(m+n)$ for all $i, j$, so $P_{m+n} = P_m P_n$.
With $P_0 = I$ and $P = P_1$, induction gives $P_n = P^n$.
*Structural Commentary*: Transition over $m+n$ steps factors through the intermediate state after $m$ steps.
*Key Insight*: Conditioning over intermediate states is matrix multiplication. □

**Lemma 4.4.9.** *Marginal Evolution*
*Let $(X_n)_{n \geq 0}$ be a homogeneous Markov chain on countable $\mathcal{X}$ with $P = P_1 = (p_{ij})$, where $p_{ij}(n) := \mathbb{P}_i(X_n = j)$ and $P_n = (p_{ij}(n))$. Define row laws $\mu_n(j) := \mathbb{P}(X_n = j)$. Then $\mu_n = \mu_0 P^n$ for all $n \in \mathbb{N}_0$.*

*Proof.* $\forall j \in \mathcal{X}$, by total probability $\mu_n(j) = \sum_{i \in \mathcal{X}} \mathbb{P}(X_n = j \mid X_0 = i)\mathbb{P}(X_0 = i) = \sum_{i \in \mathcal{X}} \mu_0(i)p_{ij}(n) = (\mu_0 P_n)_j$.
By Theorem 4.4.8, $P_n = P^n$, so $\mu_n(j) = (\mu_0 P^n)_j$ for every $j \in \mathcal{X}$; hence $\mu_n = \mu_0 P^n$.
*Structural Commentary*: Marginals evolve by multiplying the initial law by the transition semigroup.
*Key Insight*: Conditioning on $X_0$ turns distribution evolution into row-vector matrix multiplication. □

**Definition 4.4.10.** *Communication Classes*

(i) State $i$ *reaches* $j$, written $i \to j$, if $p_{ij}(n) > 0$ for some $n \geq 0$.
(ii) States $i, j$ *communicate*, written $i \leftrightarrow j$, if $i \to j$ and $j \to i$.
(iii) A *communication class* is an equivalence class under $\leftrightarrow$.
(iv) The chain is *irreducible* if all states lie in one communication class.
(v) A set $C$ is *closed* if $p_{ij} = 0$ for all $i \in C$ and $j \notin C$; a singleton closed class is an *absorbing state*.

**Proposition 4.4.11.** *Communication Classes Partition the State Space*
*Let $(X_n)_{n \geq 0}$ be a homogeneous Markov chain on countable $\mathcal{X}$ with $n$-step probabilities $p_{ij}(n)$.*
*Define $i \to j$ iff $\exists n \in \mathbb{N}_0$ such that $p_{ij}(n) > 0$, and $i \leftrightarrow j$ iff $i \to j$ and $j \to i$.*
*Then $\leftrightarrow$ is equivalence relation, and $\mathcal{X} = \bigsqcup_{C \in \mathcal{C}} C$ where $\mathcal{C} := \{[i]_{\leftrightarrow} : i \in \mathcal{X}\}$ is set of communication classes.*

*Proof.* Since $p_{ii}(0) = 1$, $i \leftrightarrow i$ for all $i \in \mathcal{X}$; symmetry is immediate from the definition.
If $i \leftrightarrow j$ and $j \leftrightarrow k$, choose $m, n, r, s \in \mathbb{N}_0$ with $p_{ij}(m) > 0$, $p_{jk}(n) > 0$, $p_{kj}(r) > 0$, and $p_{ji}(s) > 0$.
By Theorem 4.4.8, $p_{ik}(m+n) \geq p_{ij}(m)p_{jk}(n) > 0$ and $p_{ki}(r+s) \geq p_{kj}(r)p_{ji}(s) > 0$, hence $i \leftrightarrow k$.
Thus $\leftrightarrow$ is reflexive, symmetric, and transitive.
Its equivalence classes are pairwise disjoint, cover $\mathcal{X}$, and are exactly the communication classes.
*Structural Commentary*: Communication classes are the equivalence classes generated by mutual reachability.
*Key Insight*: Positive transition paths concatenate by Chapman-Kolmogorov. □

**Definition 4.4.12.** *Recurrent and Transient States*

(i) Let $\tau_i^+ := \inf\{n \geq 1 : X_n = i\}$ be the first return time to $i$.
(ii) State $i$ is *recurrent* or *persistent* if $\mathbb{P}_i(\tau_i^+ < \infty) = 1$.
(iii) State $i$ is *transient* if $\mathbb{P}_i(\tau_i^+ < \infty) < 1$.

**Theorem 4.4.13.** *Recurrence Criterion*

*(i) State $i$ is recurrent iff $S_i := \sum_{n=0}^{\infty} p_{ii}(n) = \infty$.*
*(ii) State $i$ is transient iff $S_i < \infty$.*

*Proof.* Let $N_i := \sum_{n=0}^{\infty} \mathbf{1}\{X_n = i\}$ and $r_i := \mathbb{P}_i(\tau_i^+ < \infty)$. By Tonelli, $\mathbb{E}_i N_i = \sum_{n=0}^{\infty} p_{ii}(n) = S_i$.
By the Markov property after returns to $i$, $\mathbb{P}_i(N_i \geq k) = r_i^{k-1}$ for $k \geq 1$, hence $\mathbb{E}_i N_i = \sum_{k=1}^{\infty} r_i^{k-1}$.

*(i)* ($\Rightarrow$) If $i$ is recurrent, then $r_i = 1$, so $S_i = \mathbb{E}_i N_i = \infty$.
*(i)* ($\Leftarrow$) If $S_i = \infty$, then $r_i < 1$ would give $\mathbb{E}_i N_i = (1 - r_i)^{-1} < \infty$, a contradiction; hence $r_i = 1$.
*(ii)* ($\Rightarrow$) If $i$ is transient, then $r_i < 1$, so $S_i = \mathbb{E}_i N_i = (1 - r_i)^{-1} < \infty$.
*(ii)* ($\Leftarrow$) If $S_i < \infty$, then $r_i = 1$ would give $\mathbb{E}_i N_i = \infty$, a contradiction; hence $r_i < 1$.

*Structural Commentary*: Recurrence is detected by the expected total number of returns.
*Key Insight*: Visits to $i$ repeat with geometric return probability $r_i$. □

**Definition 4.4.14.** *Period and Ergodicity*

(i) For recurrent $i$, it is *positive recurrent* if $\mathbb{E}_i \tau_i^+ < \infty$ and *null recurrent* otherwise.
(ii) The *period* of $i$ is $d(i) := \gcd\{n \geq 1 : p_{ii}(n) > 0\}$.

(iii) State $i$ is *aperiodic* if $d(i) = 1$.
(iv) State $i$ is *ergodic* if it is recurrent, positive recurrent, and aperiodic.
(v) The chain is *ergodic* if every state is ergodic.

**Definition 4.4.15.** *Stationary and Limiting Distributions*

(i) A probability row vector $\pi$ is *stationary* or *invariant* if $\pi = \pi P$.
(ii) A finite chain has *limiting distribution* $\pi$ if $p_{ij}(n) \to \pi_j$ for all states $i, j$.

Equivalently, every row of $P^n$ converges to $\pi$.

**Remark 4.4.16.** *Stationary Need Not Mean Convergent*
For the deterministic three-cycle on $\{1, 2, 3\}$, set $p_{12} = p_{23} = p_{31} = 1$ and all other $p_{ij} = 0$.
Then $\pi = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ satisfies $\pi P = \pi$, but $P^{n+3} = P^n$, so $P^n$ does not converge.

**Theorem 4.4.17.** *Finite Ergodic Chain Convergence*
*Let $P$ be the transition matrix of a finite irreducible aperiodic Markov chain on $\mathcal{X}$.*
*There is a unique $\pi \in \Delta(\mathcal{X})$ with $\pi P = \pi$, and $P^n \to \mathbf{e}\pi$.*
*For every bounded $g : \mathcal{X} \to \mathbb{R}$, $N^{-1} \sum_{n=1}^{N} g(X_n) \to \pi(g) := \sum_{j \in \mathcal{X}} g(j)\pi_j$ almost surely.*

*Proof.* Finite irreducibility and aperiodicity imply $P$ is primitive: $P^m > 0$ for some $m \geq 1$.
Row-vector form of Theorem 4.4.4 on $A = P^m$, gives unique $\pi \in \Delta(\mathcal{X})$ with $\pi P^m = \pi$, $(P^m)^k \to \mathbf{e}\pi$.
Since $(\pi P)P^m = \pi P$ and $\pi P \in \Delta(\mathcal{X})$, uniqueness gives $\pi P = \pi$.
For $n = mk + r$ with $0 \leq r < m$, $P^n = (P^m)^k P^r \to \mathbf{e}\pi P^r = \mathbf{e}\pi$.
The finite-state Markov-chain strong law yields $N^{-1} \sum_{n=1}^{N} g(X_n) \to \pi(g)$ almost surely for bounded $g$.
*Structural Commentary*: Irreducibility gives one communicating class; aperiodicity removes cyclic obstruction.
*Key Insight*: Primitivity makes $P^n$ converge to the rank-one projection $\mathbf{e}\pi$. □

**Definition 4.4.18.** *Detailed Balance*
Let $P = [p_{ij}]$ be a transition matrix on $\mathcal{X}$.
A probability row vector $\pi$ satisfies *detailed balance* for $P$ if $\pi_i p_{ij} = \pi_j p_{ji}$ for all $i, j \in \mathcal{X}$.

**Theorem 4.4.19.** *Detailed Balance Implies Stationarity*
*Let $P = [p_{ij}]$ be row-stochastic on countable $\mathcal{X}$, and let $\pi \in \Delta(\mathcal{X})$. If $\pi_i p_{ij} = \pi_j p_{ji}$ $\forall i, j \in \mathcal{X}$, then $\pi P = \pi$.*

*Proof.* $\forall j \in \mathcal{X}$, $(\pi P)_j = \sum_{i \in \mathcal{X}} \pi_i p_{ij} = \sum_{i \in \mathcal{X}} \pi_j p_{ji} = \pi_j \sum_{i \in \mathcal{X}} p_{ji} = \pi_j$, since row $j$ of $P$ sums to 1. QED.
*Structural Commentary*: Detailed balance gives pairwise cancellation of probability fluxes.
*Key Insight*: Reversibility is stronger than stationarity. □

**Proposition 4.4.20.** *Detailed Balance Implies Stationarity*
*Let $\pi(dx) = f(x)\,dx$ and let $P$ be a Markov kernel on state space $\mathsf{X}$.*
*If $\pi(dx)P(x, dy) = \pi(dy)P(y, dx)$ as measures on $\mathsf{X} \times \mathsf{X}$, then $\pi P = \pi$.*

*Proof.* For measurable $A \subseteq \mathsf{X}$, $\pi P(A) = \int P(x, A)\,\pi(dx) = \int \mathbf{1}_A(y)\,\pi(dx)P(x, dy)$.
By detailed balance, this equals $\int \mathbf{1}_A(y)\,\pi(dy)P(y, dx) = \int_A P(y, \mathsf{X})\,\pi(dy) = \pi(A)$, proving $\pi P = \pi$.
*Structural Commentary*: Detailed balance makes the joint law of consecutive states symmetric.
*Key Insight*: Marginalising the symmetric joint measure gives the same one-step density. □

**Proposition 4.4.21.** *Two-State Chain Limit*
*Let $0 < a, b < 1$, $P = \begin{bmatrix} 1-a & a \\ b & 1-b \end{bmatrix}$. Set $\pi = \left(\frac{b}{a+b}, \frac{a}{a+b}\right)$. Then $\pi P = \pi$, $\pi$ unique, $P^n \to \mathbf{e}\pi = \begin{bmatrix} \frac{b}{a+b} & \frac{a}{a+b} \\ \frac{b}{a+b} & \frac{a}{a+b} \end{bmatrix}$.*

*Proof.* Solving $\pi P = \pi$ with $\pi \mathbf{e} = 1$ gives $\pi = \left(\frac{b}{a+b}, \frac{a}{a+b}\right)$.
The characteristic polynomial is $(\lambda - 1)(\lambda - (1 - a - b))$, so $\operatorname{spec}(P) = \{1, 1 - a - b\}$ and $|1 - a - b| < 1$.
Thus stationary component is projected onto $\mathbf{e}\pi$, transverse component decays geometrically, giving $P^n \to \mathbf{e}\pi$.
*Structural Commentary*: The dynamics split into the stationary line and one transverse eigendirection.
*Key Insight*: The non-unit eigenvalue $1 - a - b$ controls convergence. □

## 4.5 Stationary Processes and Ergodicity

**Definition 4.5.1.** *Strict and Weak Stationarity*
Let $(X_t)_{t\in\mathbb{Z}}$ be a real-valued stochastic process.

(i) *Strict stationarity*: $\forall k \in \mathbb{N},\ t_1,\dots,t_k,h \in \mathbb{Z}$, $(X_{t_1},\dots,X_{t_k}) \overset{\text{dist}}{=} (X_{t_1+h},\dots,X_{t_k+h})$.
(ii) *Weak stationarity*: $\exists \mu \in \mathbb{R}$ and $\gamma : \mathbb{Z} \to \mathbb{R}$ with $\forall t,h \in \mathbb{Z}$, $X_t \in L^2$, $\mathbb{E}X_t = \mu$, and $\operatorname{Cov}(X_t, X_{t-h}) = \gamma(h)$.

*Key Insight*: (i) plus $L^2$ implies (ii); (ii) fixes only $\mu$ and $\gamma$, not finite-dimensional laws.

## 4.6 Gaussian Processes

## 4.7 Point Processes and Renewal Structure

Counting processes model random event arrivals over time. The Poisson process is the basic model in which disjoint time intervals contribute independent counts and short intervals usually contain either zero or one event.

**Definition 4.7.1.** *Counting Process*
A *counting process* is a stochastic process $\{X(t) : t \geq 0\}$ with state space $\{0,1,2,\dots\}$ such that $X(t)$ records the number of events observed up to and including time $t$.

**Definition 4.7.2.** *Poisson Process with Intensity Function*
Let $\lambda : [0,\infty) \to [0,\infty)$ be locally integrable.
A counting process $X = \{X(t) : t \geq 0\}$ is a *Poisson process* with *intensity function* $\lambda$ if:

(i) $X(0) = 0$.
(ii) For $0 = t_0 < t_1 < \cdots < t_n$, the increments $\Delta_k X := X(t_k) - X(t_{k-1})$, $1 \leq k \leq n$, are independent.
(iii) For each $t \geq 0$, as $h \downarrow 0$, $\mathbb{P}(X(t+h) - X(t) = 1) = \lambda(t)h + o(h)$ and $\mathbb{P}(X(t+h) - X(t) \geq 2) = o(h)$.

**Theorem 4.7.3.** *Increment Distribution*
*Let $X$ be a Poisson process with intensity $\lambda$, and set $\Lambda(s,t) := \int_s^t \lambda(u)\,du$ for $0 \leq s \leq t$.*
*Then $X(t) - X(s) \sim \text{Poisson}(\Lambda(s,t))$. In particular, $X(t) \sim \text{Poisson}(\Lambda(0,t))$, so $\mathbb{E}X(t) = \operatorname{Var}(X(t)) = \Lambda(0,t)$.*

*Proof.* Fix $s \geq 0$ and let $p_k(u) := \mathbb{P}(X(s+u) - X(s) = k)$ with $p_{-1} := 0$. By small-time conditions and independent increments, $p_k(u+h) = p_k(u)(1 - \lambda(s+u)h + o(h)) + p_{k-1}(u)\lambda(s+u)h + o(h)$.
Hence $p_k'(u) = -\lambda(s+u)p_k(u) + \lambda(s+u)p_{k-1}(u)$, with $p_0(0) = 1$ and $p_k(0) = 0$ for $k \geq 1$.
Writing $M(u) := \Lambda(s, s+u)$, the unique solution is $p_k(u) = e^{-M(u)}\frac{M(u)^k}{k!}$ by induction on $k$.
Taking $u = t - s$ yields $X(t) - X(s) \sim \text{Poisson}(\Lambda(s,t))$. Mean-variance identities follow from Poisson law.
*Structural Commentary*: The infinitesimal jump rules form a triangular system for the increment probabilities.
*Key Insight*: Integrating the local rate $\lambda$ gives the Poisson mean over the interval. □

**Definition 4.7.4.** *Homogeneous Poisson Process*
If $\lambda(t) \equiv \lambda > 0$, then $X(t)$ is a *homogeneous Poisson process* with *rate* $\lambda$. In this case, $X(t) \sim \text{Poisson}(\lambda t)$.

**Definition 4.7.5.** *Waiting and Interarrival Times*
For a homogeneous Poisson process:

(i) Set $W_0 = 0$ and $W_n := \inf\{t \geq 0 : X(t) \geq n\}$ for $n \geq 1$; $W_1, W_2, \dots$ are *waiting times*.
(ii) Set $S_n := W_{n+1} - W_n$ for $n \geq 0$; $S_0, S_1, \dots$ are *sojourn* or *interarrival times*.

**Theorem 4.7.6.** *Poisson Interarrival Times*
*Let $X$ be a homogeneous Poisson process with rate $\lambda > 0$.*
*Then $S_0, S_1, \dots \overset{\text{iid}}{\sim} \text{Exp}(\lambda)$, with density $f(s) = \lambda e^{-\lambda s}$ for $s \geq 0$.*
*For $n \geq 1$, $W_n = \sum_{r=0}^{n-1} S_r \sim \text{Gamma}(n,\lambda)$, so $\mathbb{E}W_n = \frac{n}{\lambda}$ and $\operatorname{Var}(W_n) = \frac{n}{\lambda^2}$.*

*Proof.* For $t \geq 0$, $\mathbb{P}(S_0 > t) = \mathbb{P}(X(t) = 0) = e^{-\lambda t}$, so $S_0 \sim \text{Exp}(\lambda)$.
By independent stationary increments, $\mathbb{P}(S_n > t \mid \mathcal{F}_{W_n}) = \mathbb{P}(X(W_n + t) - X(W_n) = 0 \mid \mathcal{F}_{W_n}) = e^{-\lambda t}$.
Thus each $S_n \sim \text{Exp}(\lambda)$ and is independent of the past, so the $S_n$ are IID.
Since $W_n = \sum_{r=0}^{n-1} S_r$, the Gamma law and its mean and variance follow.
*Structural Commentary*: Arrival times are cumulative sums of IID interarrival gaps.
*Key Insight*: No events in a length-$t$ interval has probability $e^{-\lambda t}$. □

**Proposition 4.7.7.** *Conditional Binomial Count*
*Let $X$ be a homogeneous Poisson process with rate $\lambda > 0$.*
*For $s,t > 0$ and $n \in \mathbb{N}_0$, $X(t) \mid X(t+s) = n \sim \text{Binomial}\left(n, \frac{t}{t+s}\right)$.*

*Proof.* Let $Y = X(t)$ and $Z = X(t+s) - X(t)$. Then $Y \sim \text{Poisson}(\lambda t)$, $Z \sim \text{Poisson}(\lambda s)$, and $Y \perp Z$.
For $0 \le k \le n$, $\mathbb{P}(Y = k \mid Y + Z = n) = \frac{\frac{(\lambda t)^k (\lambda s)^{n-k}}{k!(n-k)!}}{\frac{(\lambda(t+s))^n}{n!}} = \binom{n}{k} \left(\frac{t}{t+s}\right)^k \left(\frac{s}{t+s}\right)^{n-k}$; QED.
*Structural Commentary*: Conditioning on the total allocates independent Poisson counts.
*Key Insight*: Independent Poisson counts become multinomial after conditioning on their sum. □

**Remark 4.7.8.** *Uniform Event-Time Diagnostic*
For a homogeneous Poisson process on $[0, T]$, conditional on $X(T) = n$, the event times $(W_1, \ldots, W_n)$ have the law of the order statistics of $n$ IID Uniform$(0, T)$ variables.
Marked nonuniformity in the observed event times is evidence against homogeneity.

**Proposition 4.7.9.** *Odd Poisson Count*
*Let $X$ be a homogeneous Poisson process with rate $\lambda > 0$. For $t \ge 0$, $\mathbb{P}(X(t) \in 2\mathbb{N}_0 + 1) = \frac{1 - e^{-2\lambda t}}{2}$.*

*Proof.* Let $\mu = \lambda t$. Since $X(t) \sim \text{Poisson}(\mu)$, $\mathbb{P}(X(t) \in 2\mathbb{N}_0 + 1) = e^{-\mu} \sum_{k=0}^{\infty} \frac{\mu^{2k+1}}{(2k+1)!} = e^{-\mu} \sinh \mu = \frac{1 - e^{-2\mu}}{2}$.
*Structural Commentary*: Split the Poisson mass by parity.
*Key Insight*: The odd terms of $e^{\mu}$ sum to $\sinh \mu$. □

**Proposition 4.7.10.** *Nearest Neighbor in a Planar Poisson Process*
*Let $\Phi$ be a planar Poisson point process with intensity $\lambda > 0$, fix $x_0 \in \mathbb{R}^2$, and set $R := \inf_{Y \in \Phi} \|Y - x_0\|$.*
*Then for $t \ge 0$, $\mathbb{P}(R > t) = e^{-\lambda \pi t^2}$, and $\mathbb{E}R = \frac{1}{2\sqrt{\lambda}}$.*

*Proof.* Let $B_t := B(x_0, t)$, $N_t := \#(\Phi \cap B_t)$. Then $N_t \sim \text{Poisson}(\lambda \pi t^2)$, $\{R > t\} = \{N_t = 0\}$,
so $\mathbb{P}(R > t) = e^{-\lambda \pi t^2}$. Since $R \ge 0$, $\mathbb{E}R = \int_0^{\infty} \mathbb{P}(R > t)\, dt = \int_0^{\infty} e^{-\lambda \pi t^2}\, dt = \frac{1}{2\sqrt{\lambda}}$.
*Structural Commentary*: Nearest-neighbor distance is encoded by empty disks around $x_0$.
*Key Insight*: The tail of $R$ is the zero-count probability of a Poisson disk count. □

# 5 Stochastic Calculus and SDEs

## 5.1 Brownian Motion

## 5.2 Stochastic Integration

## 5.3 Itô Calculus

## 5.4 Stochastic Differential Equations

## 5.5 Change of Measure and Girsanov Theory

## 5.6 Diffusions and PDE Connections

## 5.7 Stochastic Flows and Advanced Dynamics

# Part XVII
# Dynamical and PDE Systems

## 1 Dynamical Systems and ODEs

### 1.1 ODE Foundations and Flows

Ordinary differential equations prescribe the evolution of an unknown curve through relations among the curve and its derivatives. Their order, dependence on time, and linearity determine the natural state variables and solution methods.

**Definition 1.1.1.** *Implicit Scalar ODE and Regular Point*
Let $k \in \mathbb{N}$, $k \geq 1$, $U \subseteq \mathbb{R}^{k+2}$ be open, and $F \in C(U, \mathbb{R})$. For $I \subseteq \mathbb{R}$, $x \in C^k(I, \mathbb{R})$, set $J_x^k(t) := (t, x(t), x^{(1)}(t), \dots, x^{(k)}(t))$.

(i) $F(t, y_0, \dots, y_k) = 0$ is an *implicit scalar ODE written through order $k$*; its $k$-jet form is $F(J_x^k(t)) = 0$.
(ii) $x$ is a *solution* on $I$ iff $J_x^k(t) \in U$ and $F(J_x^k(t)) = 0\ \forall t \in I$.
(iii) $F \in C^1(U, \mathbb{R})$; $p = (t_*, y_0^*, \dots, y_k^*) \in F^{-1}(0)$ is *regular w.r.t. highest derivative* iff $\partial_{y_k} F(p) \neq 0$.

*Key Insight*: $\partial_{y_k} F(p) \neq 0$ locally solves the relation for $x^{(k)}$.

**Proposition 1.1.2.** *Local Explicit Form of a Regular Implicit ODE*
*Let $k \in \mathbb{N}$, $k \geq 1$, $U \subseteq \mathbb{R}^{k+2}$ be open, $F \in C^1(U, \mathbb{R})$, $u_* = (t_*, y_0^*, \dots, y_{k-1}^*)$, and $p = (u_*, y_k^*) \in U$ with $F(p) = 0$ and $\partial_{y_k} F(p) \neq 0$. Then $\exists$ open $V \subseteq \mathbb{R}^{k+1}$, $W \subseteq \mathbb{R}$ with $u_* \in V$, $y_k^* \in W$, and a unique $f \in C^1(V, \mathbb{R})$ such that $f(V) \subseteq W$, $f(u_*) = y_k^*$, $V \times W \subseteq U$, and:*

*(i) $F(u, v) = 0$ iff $v = f(u)\ \forall (u, v) \in V \times W$.*
*(ii) If $x \in C^k(I, \mathbb{R})$ and $J_x^k(I) \subseteq V \times W$, then $F(J_x^k(t)) = 0\ \forall t \in I$ iff $x^{(k)}(t) = f(t, x(t), \dots, x^{(k-1)}(t))\ \forall t \in I$.*

*Proof.* Let $u = (t, y_0, \dots, y_{k-1})$, $v = y_k$, $\widehat{U} := \{(v, u) : (u, v) \in U\}$, and $\widehat{F}(v, u) := F(u, v)$.
Since $\partial_v \widehat{F}(y_k^*, u_*) = \partial_{y_k} F(p) \neq 0$, by Theorem 4.2.3, there are $W, V$ and a unique $f \in C^1(V, \mathbb{R})$ with $W \times V \subseteq \widehat{U}$, $f(V) \subseteq W$, $f(u_*) = y_k^*$, and $\widehat{F}(v, u) = 0$ iff $v = f(u)$.

(i) For $(u, v) \in V \times W$, $F(u, v) = \widehat{F}(v, u)$, so $F(u, v) = 0$ iff $v = f(u)$.
(ii) Set $u_x(t) := (t, x(t), \dots, x^{(k-1)}(t))$ and $v_x(t) := x^{(k)}(t)$. Since $(u_x(t), v_x(t)) \in V \times W\ \forall t \in I$, (i) gives $F(J_x^k(t)) = 0$ iff $v_x(t) = f(u_x(t))\ \forall t \in I$.

*Structural Commentary*: IFT solves $F(u, y_k) = 0$ for $y_k$.
*Key Insight*: $\partial_{y_k} F(p) \neq 0$ turns $F = 0$ into $y_k = f(u)$ locally. □

**Definition 1.1.3.** *Explicit ODE Systems and Classes*
Let $k, n \in \mathbb{N}$, $k \geq 1$, $U \subseteq \mathbb{R} \times (\mathbb{R}^n)^k$ open, and $f : U \to \mathbb{R}^n$.
Use $X_x(t) := (x(t), x^{(1)}(t), \dots, x^{(k-1)}(t))$, $\xi = (\xi_0, \dots, \xi_{k-1})$, and $\pi(U) := \{\xi \in (\mathbb{R}^n)^k : \exists t \in \mathbb{R},\ (t, \xi) \in U\}$.

(i) $x^{(k)}(t) = f(t, X_x(t))$ is an *explicit ODE system of order $k$*.
(ii) For an interval $I \subseteq \mathbb{R}$, $x \in C^k(I, \mathbb{R}^n)$ is a *solution* iff $\forall t \in I$, $(t, X_x(t)) \in U$ and $x^{(k)}(t) = f(t, X_x(t))$.
(iii) The system is *autonomous* iff $\exists F : \pi(U) \to \mathbb{R}^n$ with $f(t, \xi) = F(\xi)\ \forall (t, \xi) \in U$.
(iv) On an interval $I$, the system is *linear* iff $\exists g \in C(I, \mathbb{R}^n)$ and $A_j \in C(I, \mathbb{R}^{n \times n})$, $0 \leq j < k$, with $f(t, \xi) = g(t) + \sum_{j=0}^{k-1} A_j(t)\xi_j\ \forall (t, \xi) \in U \cap (I \times (\mathbb{R}^n)^k)$.
(v) A linear system on $I$ is *homogeneous* iff (iv) holds with $g = 0$.
(vi) A solution $x : I \to \mathbb{R}^n$ is *global* iff $I = \mathbb{R}$.
(vii) A solution $x : I \to \mathbb{R}^n$ is *maximal* iff $\nexists$ solution $\bar{x} : \bar{I} \to \mathbb{R}^n$ with $I \subsetneq \bar{I}$ and $\bar{x}|_I = x$.

*Key Insight*: The state lift $X_x$ has dimension $kn$.

**Proposition 1.1.4.** *Reduction to First Order*
*Let $k, n \in \mathbb{N}$, $k \geq 1$, $U \subseteq \mathbb{R} \times (\mathbb{R}^n)^k$ open, and $f : U \to \mathbb{R}^n$.*
*For $y = (y_1, \dots, y_k)$, define $G(t, y) := (y_2, \dots, y_k, f(t, y))$.*
*For an interval $I \subseteq \mathbb{R}$ and $x \in C^k(I, \mathbb{R}^n)$, set $y := (x, x^{(1)}, \dots, x^{(k-1)})$ and assume $(t, y(t)) \in U\ \forall t \in I$.*
*Then $y \in C^1(I, (\mathbb{R}^n)^k)$, and $x^{(k)}(t) = f(t, y(t))\ \forall t \in I$ iff $y'(t) = G(t, y(t))\ \forall t \in I$.*

*Proof.*

(⇒) Since $x \in C^k(I, \mathbb{R}^n)$, $y \in C^1(I, (\mathbb{R}^n)^k)$. The identities $y_j' = x^{(j)} = y_{j+1}$, $1 \leq j < k$, and $y_k' = x^{(k)} = f(t, y(t))$ imply $y' = G(\cdot, y)$ on $I$.
(⇐) Since $y_k = x^{(k-1)}$, the $k$th coordinate gives $x^{(k)}(t) = y_k'(t) = G_k(t, y(t)) = f(t, y(t))\ \forall t \in I$.

*Structural Commentary*: Coordinates shift; the last coordinate carries $f$.
*Key Insight*: Adjoin derivatives to get first order. □

**Proposition 1.1.5.** *Autonomization by Adjoining Time*
*Let $m \in \mathbb{N}$, $U \subseteq \mathbb{R} \times \mathbb{R}^m$ be open, $f : U \to \mathbb{R}^m$, and $\widetilde{f}(s,\xi) := (1, f(s,\xi))$. For an interval $I \subseteq \mathbb{R}$:*

*(i) If $y \in C^1(I, \mathbb{R}^m)$, $(t, y(t)) \in U$, and $y' = f(t,y)$, then $z(t) := (t, y(t))$ solves $z' = \widetilde{f}(z)$.*
*(ii) If $z = (s,y) \in C^1(I, \mathbb{R}^{m+1})$, $z(I) \subseteq U$, $z' = \widetilde{f}(z)$, and $s(t_0) = t_0$ for some $t_0 \in I$, then $s = \mathrm{id}_I$ and $y'(t) = f(t, y(t))$ on $I$.*

*Proof.*
(i) For $z = (t,y)$, $z' = (1, y') = (1, f(t,y)) = \widetilde{f}(t,y) = \widetilde{f}(z)$.
(ii) Since $(s', y') = z' = \widetilde{f}(s,y) = (1, f(s,y))$, $s' = 1$ and $y' = f(s,y)$.
With $s(t_0) = t_0$, $s = \mathrm{id}_I$; hence $y'(t) = f(t, y(t))$.

*Structural Commentary*: The clock coordinate has $s' = 1$.
*Key Insight*: $y' = f(t,y)$ becomes $z' = \widetilde{f}(z)$ with $z = (t,y)$. □

**Example 1.1.6.** *Newtonian Motion as a First-Order System*
Let $q : \mathbb{R} \to \mathbb{R}^3$ be the position of a particle of mass $m > 0$, let $v := q'$, and let $F : \mathbb{R}^3 \to \mathbb{R}^3$ be a position-dependent force. Newton's equation $mq''(t) = F(q(t))$ becomes $(q,v)' = (v, m^{-1}F(q))$ by Proposition 1.1.4. For uniform gravity $F(q) = -mge_3$, where $e_3 := (0,0,1)^{\mathsf{T}}$ and $g > 0$, the initial state $(q(0), v(0)) = (q_0, v_0)$ gives $q(t) = q_0 + v_0 t - \frac{g}{2} e_3 t^2$. Thus position and velocity together determine the mechanical state.

**Proposition 1.1.7.** *Scalar Autonomous Separation*
*Let $D \subseteq \mathbb{R}$ be open, $f \in C(D, \mathbb{R})$, $D_f := \{x \in D : f(x) \neq 0\}$, and $x_0 \in D_f$.*
*Let $C$ be the component of $D_f$ containing $x_0$, and set $H(x) := \int_{x_0}^{x} \frac{du}{f(u)}$, $K := H(C)$. Then:*

*(i) $K$ is an open interval and $H : C \to K$ is a differentiable bijection.*
*(ii) $\phi := H^{-1} : K \to C$ is the unique $C$-valued solution of $x' = f(x)$ with $x(0) = x_0$.*
*(iii) $K$ is maximal among intervals of $C$-valued solutions through $x_0$, and $\phi$ is global iff $K = \mathbb{R}$.*

*Proof.*
(i) Since $D_f$ is open, $C$ is an open interval. By Theorem 2.2.1, $H' = 1/f \neq 0$ on $C$.
Since $f$ has constant sign on $C$ by Theorem 5.1.41, $H$ is strictly monotone.
Hence $H : C \to K = H(C)$ is bijective and $K$ is an open interval.
(ii) By Corollary 1.1.20, $\phi \in C^1(K, C)$ and $\phi' = 1/(H' \circ \phi) = f \circ \phi$. Since $H(x_0) = 0$, $\phi(0) = x_0$.
If $u : L \to C$ solves $u' = f(u)$ with $0 \in L$ and $u(0) = x_0$, then $(H \circ u)' = H'(u)u' = 1$ and $(H \circ u)(0) = 0$.
Hence $H(u(t)) = t$, so $L \subseteq K$ and $u = \phi|_L$.
(iii) If $\bar{\phi} : \bar{K} \to C$ solves $x' = f(x)$, $K \subseteq \bar{K}$, and $\bar{\phi}|_K = \phi$, then (ii) gives $\bar{K} \subseteq K$; hence $\bar{K} = K$.
Thus $K$ is maximal. Since $\operatorname{dom}\phi = K$, $\phi$ is global iff $K = \mathbb{R}$.

*Structural Commentary*: Monotone coordinate, inverse trajectory.
*Key Insight*: Time is accumulated reciprocal speed. □

**Method 1.1.8.** *Separation of Variables*
Let $D \subseteq \mathbb{R}$ be open, $J \subseteq \mathbb{R}$ open interval, $f \in C(D, \mathbb{R})$, $g \in C(J, \mathbb{R})$, $t_0 \in J$, $x_0 \in D_f := \{x \in D : f(x) \neq 0\}$.
Let $C \ni x_0$ be the component of $D_f$ and $H(x) := \int_{x_0}^{x} \frac{du}{f(u)}$. Since $f$ has constant sign on $C$ and $H' = 1/f$, $H : C \to H(C)$ is a differentiable bijection.

(i) Set $G(t) := \int_{t_0}^{t} g(s)\,ds$ and let $L$ be the component of $\{t \in J : G(t) \in H(C)\}$ containing $t_0$.
(ii) Output $x := H^{-1} \circ G$ on $L$; then $x' = f(x)g$ on $L$ and $x(t_0) = x_0$.

**Example 1.1.9.** *Global Growth, Finite-Time Blow-Up, and Nonuniqueness*
The separation coordinate distinguishes three basic behaviors.

(i) For $x' = x$ and $x(0) = x_0 > 0$, $H(x) = \log\left(\frac{x}{x_0}\right)$ and $x(t) = x_0 e^t$ on $\mathbb{R}$.
(ii) For $x' = x^2$ and $x(0) = x_0 > 0$, $H(x) = x_0^{-1} - x^{-1}$ and $x(t) = \frac{x_0}{1 - x_0 t}$ on $(-\infty, x_0^{-1})$; $x(t) \to +\infty$ as $t \uparrow x_0^{-1}$.
(iii) For $x' = \sqrt{|x|}$ and $x(0) = 0$, $a \in [0, +\infty)$ determines $x_a(t) := \begin{cases} 0, & t \le a, \\ \frac{(t-a)^2}{4}, & t \ge a. \end{cases}$ Then $x_a \in C^1(\mathbb{R})$, $x_a' = \sqrt{|x_a|}$, $x_a(0) = 0$, and if $0 \le a < b$, then $x_a = x_b$ on $(-\infty, a]$ while $x_a > x_b$ on $(a, b]$; hence the IVP is nonunique.

First-order equations can also be rewritten as fixed-point problems on complete function spaces.

**Definition 1.1.10.** *Scalar Initial-Value Problem*
Let $O \subseteq \mathbb{R}^2$ be open, $(x_0, y_0) \in O$, $g : O \to \mathbb{R}$, and $\mathrm{IVP}(O, g, x_0, y_0) := \{y' = g(x, y),\ y(x_0) = y_0\}$.
For $J \subseteq \mathbb{R}$ with $x_0 \in J$, $f : J \to \mathbb{R}$ solves $\mathrm{IVP}(O, g, x_0, y_0)$ iff

(i) $J$ is open and $f$ is differentiable.
(ii) $\Gamma_f := \{(x, f(x)) : x \in J\} \subseteq O$.
(iii) $f'(x) = g(x, f(x))\ \forall x \in J$, and $f(x_0) = y_0$.

*Structural Commentary*: $\Gamma_f \subseteq O$ makes $g(\,\cdot\,, f)$ defined; $f(x_0) = y_0$ selects the curve.

**Proposition 1.1.11.** *Differential and Integral Forms*
*Let $O \subseteq \mathbb{R}^2$ be open, $(x_0, y_0) \in O$, $g \in C(O, \mathbb{R})$, $J \subseteq \mathbb{R}$ open with $x_0 \in J$, and $f \in C(J, \mathbb{R})$.*
*Let $\Gamma_f := \{(x, f(x)) : x \in J\} \subseteq O$. Then $f$ solves $\mathrm{IVP}(O, g, x_0, y_0)$ iff $f(x) = y_0 + \int_{x_0}^{x} g(t, f(t))\, dt\ \forall x \in J$.*

*Proof.*

($\Rightarrow$) If $f'(x) = g(x, f(x))$ on $J$, $f(x_0) = y_0$, by Theorem 2.2.2, $f(x) - y_0 = \int_{x_0}^{x} f'(t)\, dt = \int_{x_0}^{x} g(t, f(t))\, dt\ \forall x \in J$.
($\Leftarrow$) If $f(x) = y_0 + \int_{x_0}^{x} g(t, f(t))\, dt\ \forall x \in J$, then $g(\,\cdot\,, f(\,\cdot\,))$ is continuous on $J$.
By Theorem 2.2.1, $f'(x) = g(x, f(x))$ on $J$. Evaluating at $x = x_0$ gives $f(x_0) = y_0$.

*Structural Commentary*: Differentiation and integration are inverse along the continuous graph.
*Key Insight*: The IVP is equivalent to its Volterra integral equation. □

**Theorem 1.1.12.** *Picard Local Existence Theorem*
*Let $O \subseteq \mathbb{R}^2$ be open, $(x_0, y_0) \in O$, and $g \in C(O, \mathbb{R})$.*
*Assume $|g(x, y_1) - g(x, y_2)| \le M\,|y_1 - y_2|$ for some $M > 0$ whenever $(x, y_1), (x, y_2) \in O$.*
*Then $\exists a, b, \ell > 0$ with $Q := [x_0 - a, x_0 + a] \times [y_0 - b, y_0 + b] \subseteq O$ and $J_\ell := (x_0 - \ell, x_0 + \ell)$ such that:*

*(i) The IVP $y' = g(x, y)$, $y(x_0) = y_0$, has a differentiable solution $f : J_\ell \to \mathbb{R}$ with $\Gamma_f \subseteq Q$.*
*(ii) If $h : J_\ell \to \mathbb{R}$ is a differentiable solution with $\Gamma_h \subseteq Q$, then $h = f$.*

*Proof.*

(i) Choose $a, b > 0$ with $Q := [x_0 - a, x_0 + a] \times [y_0 - b, y_0 + b] \subseteq O$.
Let $K := \sup_Q |g| < +\infty$, and choose $0 < \ell \le a$ with $\ell K \le b$ and $\ell M < 1$.
Put $I_\ell := [x_0 - \ell, x_0 + \ell]$, $\mathcal{X}_\ell := \{u \in C(I_\ell, \mathbb{R}) : \|u - y_0\|_\infty \le b\}$, and $(Tu)(x) := y_0 + \int_{x_0}^{x} g(t, u(t))\, dt$.
The space $\mathcal{X}_\ell$ is complete, and for $u, v \in \mathcal{X}_\ell$, $\|Tu - y_0\|_\infty \le \ell K \le b$ and $\|Tu - Tv\|_\infty \le \ell M\, \|u - v\|_\infty$.
Thus $T : \mathcal{X}_\ell \to \mathcal{X}_\ell$ is a contraction; by Theorem 4.5.3, $\exists! u_* \in \mathcal{X}_\ell$ with $Tu_* = u_*$.
For $f := u_*|_{J_\ell}$, $\Gamma_f \subseteq Q$; by Proposition 1.1.11, $f' = g(\,\cdot\,, f)$ and $f(x_0) = y_0$.
(ii) Let $h : J_\ell \to \mathbb{R}$ solve the IVP with $\Gamma_h \subseteq Q$. For $0 < r < \ell$, put $I_r := [x_0 - r, x_0 + r]$,
$\mathcal{X}_r := \{u \in C(I_r, \mathbb{R}) : \|u - y_0\|_\infty \le b\}$, and $(T_r u)(x) := y_0 + \int_{x_0}^{x} g(t, u(t))\, dt$.
Since $rM < 1$, $T_r$ is a contraction. By Proposition 1.1.11, $T_r(h|_{I_r}) = h|_{I_r}$ and $T_r(u_*|_{I_r}) = u_*|_{I_r}$.
Uniqueness gives $h = f$ on $I_r$. Hence $h = f$ on $\bigcup_{0<r<\ell} I_r = J_\ell$.

*Structural Commentary*: Box control plus short-time contraction.
*Key Insight*: Picard iteration is Banach iteration. □

**Method 1.1.13.** *Picard Iteration*
Let $Q := [x_0 - a, x_0 + a] \times [y_0 - b, y_0 + b] \subseteq O$, assume $|g(x, y)| \le K$ for $(x, y) \in Q$,
$|g(x, y_1) - g(x, y_2)| \le M\,|y_1 - y_2|$ whenever $(x, y_1), (x, y_2) \in Q$. Let $\ell > 0$ with $\ell \le a$, $\ell K \le b$, and $\ell M < 1$.
Set $I_\ell := [x_0 - \ell, x_0 + \ell]$, $J_\ell := (x_0 - \ell, x_0 + \ell)$, $\mathcal{X}_\ell := \{u \in C(I_\ell, \mathbb{R}) : \|u - y_0\|_\infty \le b\}$,
$(Tu)(x) := y_0 + \int_{x_0}^{x} g(t, u(t))\, dt$. Then

(i) Initialize $f_0(x) \equiv y_0$ on $I_\ell$.
(ii) Iterate $f_{n+1} := T f_n$, i.e. $f_{n+1}(x) = y_0 + \int_{x_0}^{x} g(t, f_n(t))\, dt$ for $x \in I_\ell$.
(iii) Let $f_\infty := \lim_{n \to \infty} f_n$ uniformly on $I_\ell$, and output $f := f_\infty|_{J_\ell}$.

By Theorem 4.5.3, the limit exists and is the unique fixed point of $T$; Proposition 1.1.11 identifies its restriction as the unique local solution with graph in $Q$.




Figure 236: Picard setup: choose $\ell$ with $\ell K \le b$ so $T(\mathcal{X}_\ell) \subseteq \mathcal{X}_\ell$ and graphs stay in $Q$.

The same fixed-point formulation applies to vector fields and supports quantitative iteration estimates.

**Definition 1.1.14.** *Vector IVP and Local State Lipschitz Condition*
Let $n \in \mathbb{N}$, $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f : U \to \mathbb{R}^n$, and $(t_0, x_0) \in U$.

(i) The *initial-value problem* is $x' = f(t, x)$, $x(t_0) = x_0$; a solution on an interval $I \ni t_0$ is $x \in C^1(I, \mathbb{R}^n)$ with $(t, x(t)) \in U$ and $x'(t) = f(t, x(t))$ $\forall t \in I$.

(ii) $f$ is *locally Lipschitz in the state, uniformly in time* iff $\forall$ compact $V \subseteq U$, $\exists L_V < +\infty$ such that $\|f(t, x) - f(t, y)\| \le L_V \|x - y\|$ whenever $(t, x), (t, y) \in V$.

*Key Insight*: State-Lipschitz control makes the integral operator asymptotically contractive under iteration.

**Lemma 1.1.15.** *$C^1$ Vector Fields Are Locally State-Lipschitz*
*Let $n \in \mathbb{N}$, $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, and $f \in C^1(U, \mathbb{R}^n)$.*
*$\forall$ compact $K \subseteq U$, $\exists L_K < +\infty$ such that $\|f(t, x) - f(t, y)\| \le L_K \|x - y\|$ whenever $(t, x), (t, y) \in K$.*

*Proof.* Fix compact $K \subseteq U$. If $K = \emptyset$, take $L_K = 0$. Otherwise choose $r > 0$, $K_r := \{z : \operatorname{dist}(z, K) \le r\} \subseteq U$.
Set $A := \max_{K_r} \|D_x f\|$, $M := \max_K \|f\|$, and $L_K := \max\{A, \frac{2M}{r}\}$. For $(t, x), (t, y) \in K$, put $\rho := \|x - y\|$.
If $\rho < r$, then $\sigma(s) := (t, x + s(y - x)) \in K_r$ for $s \in [0, 1]$; by Theorem 3.2.1, $\|f(t, x) - f(t, y)\| \le A\rho$.
If $\rho \ge r$, then $\|f(t, x) - f(t, y)\| \le 2M \le \frac{2M}{r}\rho$. Thus $\|f(t, x) - f(t, y)\| \le L_K \rho$.
*Structural Commentary*: Nearby pairs use $D_x f$; separated pairs use boundedness.
*Key Insight*: Compactness yields one state-Lipschitz constant. □

**Theorem 1.1.16.** *Quantitative Vector Picard–Lindelöf Theorem*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C(U, \mathbb{R}^n)$, and $(t_0, x_0) \in U$. Choose $T, \delta > 0$ and set $V := [t_0, t_0 + T] \times \overline{B}_\delta(x_0) \subseteq U$.*
*Assume $L \ge 0$ and $\|f(t, x) - f(t, y)\| \le L \|x - y\|$ $\forall (t, x), (t, y) \in V$.*
*Set $M := \max_V \|f\|$, $T_0 := \min\{T, \delta/M\}$ with $\delta/0 := +\infty$, and $I := [t_0, t_0 + T_0]$.*
*Let $X := C(I, \mathbb{R}^n)$, $\|u\|_\infty := \max_{t \in I} \|u(t)\|$, and $\mathcal{C} := \{u \in X : \|u - x_0\|_\infty \le \delta\}$.*
*Define $Ku := x_0 + \int_{t_0}^{(\cdot)} f(s, u(s))\, ds$, $u_0 \equiv x_0$, and $u_{m+1} := Ku_m$.*

*(i) $(u_m) \subseteq \mathcal{C}$ and $u_m \to \phi$ uniformly for some $\phi \in \mathcal{C}$ satisfying $K\phi = \phi$; hence $\phi \in C^1(I, \mathbb{R}^n)$ solves the IVP.*
*(ii) $\phi$ is the unique solution on $I$ whose graph lies in $V$.*
*(iii) If $M_0 := \max_{t \in I} \|f(t, x_0)\|$, then $\|\phi - u_m\|_\infty \le T_0 M_0 e^{LT_0} \frac{(LT_0)^m}{m!}$ $\forall m \ge 1$.*

*Proof.*

(i) For $u \in \mathcal{C}$, $\|Ku - x_0\|_\infty \le MT_0 \le \delta$; hence $K(\mathcal{C}) \subseteq \mathcal{C}$.
For $u, v \in \mathcal{C}$, induction on $m \ge 1$ yields $\|K^m u(t) - K^m v(t)\| \le \frac{L^m (t - t_0)^m}{m!} \|u - v\|_\infty$ for $t \in I$.
Hence $\|u_{m+1} - u_m\|_\infty \le \frac{(LT_0)^m}{m!} \|Ku_0 - u_0\|_\infty$, so $(u_m)$ is uniformly Cauchy.
By Theorem 8.1.1, $X$ is complete; since $\mathcal{C}$ is closed, $u_m \to \phi \in \mathcal{C}$.
The case $m = 1$ implies $K$ is continuous on $\mathcal{C}$, so $K\phi = \lim_{m \to \infty} Ku_m = \lim_{m \to \infty} u_{m+1} = \phi$.
Thus $\phi(t) = x_0 + \int_{t_0}^t f(s, \phi(s))\, ds$, so $\phi \in C^1(I, \mathbb{R}^n)$, $\phi(t_0) = x_0$, and $\phi' = f(\cdot, \phi)$.

(ii) If $Ku = u$ and $Kv = v$ in $\mathcal{C}$, then $\|u - v\|_\infty = \|K^m u - K^m v\|_\infty \le \frac{(LT_0)^m}{m!} \|u - v\|_\infty$ $\forall m \ge 1$.
Choose $m$ with $(LT_0)^m / m! < 1$; hence $u = v$.
If $y$ solves the IVP on $I$ with graph in $V$, then $y \in \mathcal{C}$ and integration implies $Ky = y$; hence $y = \phi$.

(iii) Since $\|Ku_0 - u_0\|_\infty \le T_0 M_0$, (i) implies $\|\phi - u_m\|_\infty \le \sum_{j=m}^{\infty} \|u_{j+1} - u_j\|_\infty \le T_0 M_0 \sum_{j=m}^{\infty} \frac{(LT_0)^j}{j!}$.
For $a \ge 0$, $(m + k)! \ge m! k!$ implies $\sum_{j=m}^{\infty} \frac{a^j}{j!} \le \frac{a^m}{m!} e^a$; set $a = LT_0$.

*Structural Commentary*: Factorial iterate bounds replace the short-time contraction restriction.
*Key Insight*: Iterates contract after sufficiently many integrations. □

**Theorem 1.1.17.** *Picard–Lindelöf with an Integrable Lipschitz Bound*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C(U, \mathbb{R}^n)$, $(t_0, x_0) \in U$, and choose $T, \delta > 0$ with $V := [t_0, t_0 + T] \times \overline{B}_\delta(x_0) \subseteq U$. Set $b(t) := \max_{x \in \overline{B}_\delta(x_0)} \|f(t,x)\|$. Let $L : [t_0, t_0 + T] \to [0, +\infty)$ be integrable,*

*satisfy $\|f(t,x) - f(t,y)\| \le L(t)\,\|x - y\|$ for $(t,x), (t,y) \in V$. Choose $0 < T_0 \le T$ with $\int_{t_0}^{t_0+T_0} b(s)\,ds \le \delta$.*
*Put $I := [t_0, t_0 + T_0]$, $A(t) := \int_{t_0}^t L(s)\,ds$, $A := A(t_0 + T_0)$, and $B_0 := \int_{t_0}^{t_0+T_0} \|f(s, x_0)\|\,ds$.*
*Let $\mathcal{C} := \{u \in C(I, \mathbb{R}^n) : \|u - x_0\|_\infty \le \delta\}$, $(Ku)(t) := x_0 + \int_{t_0}^t f(s, u(s))\,ds$, $u_0 \equiv x_0$, and $u_{m+1} := Ku_m$. Then:*

*(i) $K(\mathcal{C}) \subseteq \mathcal{C}$ and $u_m \to \phi \in \mathcal{C}$ uniformly, where $K\phi = \phi$.*
*(ii) $\phi \in C^1(I, \mathbb{R}^n)$ is the unique solution of $x' = f(t,x)$, $x(t_0) = x_0$, whose graph lies in $V$.*
*(iii) $\|\phi - u_m\|_\infty \le B_0 e^A \frac{A^m}{m!}$ for $m \ge 0$.*

*Proof.* For $u \in \mathcal{C}$ and $t \in I$, $\|Ku(t) - x_0\| \le \int_{t_0}^t b(s)\,ds \le \delta$; hence $K(\mathcal{C}) \subseteq \mathcal{C}$.
Induction and $dA = L\,ds$ give $\|K^m u(t) - K^m v(t)\| \le \frac{A(t)^m}{m!}\,\|u - v\|_\infty$ for $u, v \in \mathcal{C}$ and $m \ge 1$.
With $u = Ku_0$ and $v = u_0$, $\|u_{j+1} - u_j\|_\infty \le B_0 \frac{A^j}{j!}$ for $j \ge 0$. Thus $(u_m)$ is uniformly Cauchy.

(i) By Theorem 8.1.1, $C(I, \mathbb{R}^n)$ is complete; closedness of $\mathcal{C}$ gives $u_m \to \phi \in \mathcal{C}$.
The case $m = 1$ makes $K$ continuous, so $K\phi = \lim_{m\to\infty} Ku_m = \lim_{m\to\infty} u_{m+1} = \phi$.

(ii) By Theorem 2.2.1, the fixed-point identity implies $\phi \in C^1(I, \mathbb{R}^n)$, $\phi(t_0) = x_0$, and $\phi' = f(\cdot, \phi)$.
If $y$ is a solution with graph in $V$, then $y \in \mathcal{C}$ and $Ky = y$.
If $Ku = u$ and $Kv = v$, then $\|u - v\|_\infty \le \frac{A^m}{m!}\,\|u - v\|_\infty$ for $m \ge 1$; choose $m$ with $\frac{A^m}{m!} < 1$, hence $u = v$.

(iii) By telescoping, $\|\phi - u_m\|_\infty \le B_0 \sum_{j=m}^{\infty} \frac{A^j}{j!}$. Since $(m + r)! \ge m!r!$, $\sum_{j=m}^{\infty} \frac{A^j}{j!} \le e^A \frac{A^m}{m!}$.

*Structural Commentary*: Integrated Lipschitz mass gives factorial iterate bounds.
*Key Insight*: Repeated integration contracts eventually. □

**Remark 1.1.18.** *Carathéodory Solutions*
Let $I \subseteq \mathbb{R}$, $t_0 \in I$, $x_0 \in \mathbb{R}^n$, $\delta > 0$, $V := I \times \overline{B}_\delta(x_0)$, and $f : V \to \mathbb{R}^n$.
Suppose $f(\cdot, x)$ is measurable, $f(t, \cdot)$ is continuous for a.e. $t$, and $\exists b, L \in L^1(I, [0, +\infty))$ with $\|f(t,x)\| \le b(t)$ and $\|f(t,x) - f(t,y)\| \le L(t)\,\|x - y\|$ for a.e. $t$ and $x, y \in \overline{B}_\delta(x_0)$.
Then a fixed point of $(Kx)(t) := x_0 + \int_{t_0}^t f(s, x(s))\,ds$ lies in $AC(I, \mathbb{R}^n)$ and satisfies $x' = f(\cdot, x)$ a.e.
Generally $AC$ cannot be replaced by $C^1$. For $f(t,x) = \operatorname{sign}(t)x$, $x(t) := e^{|t|}$ satisfies $x(0) = 1$,
$x(t) = 1 + \int_0^t \operatorname{sign}(s)x(s)\,ds$, and $x' = \operatorname{sign}(t)x$ for $t \ne 0$, but $x \notin C^1((-1, 1))$.

**Lemma 1.1.19.** *Time Regularity of ODE Solutions*
*Let $I \subseteq \mathbb{R}$ be an interval, $k, n \in \mathbb{N}$, $k \ge 1$, $U \subseteq \mathbb{R} \times \mathbb{R}^n$ open, and $f \in C^k(U, \mathbb{R}^n)$.*
*Let $x \in C^1(I, \mathbb{R}^n)$ and $\gamma := (\mathrm{id}_I, x)$. If $\gamma(I) \subseteq U$ and $x' = f \circ \gamma$, then $x \in C^{k+1}(I, \mathbb{R}^n)$.*

*Proof.* Let $P_j$ mean $x \in C^j(I, \mathbb{R}^n)$. Since $x \in C^1$, $P_1$ holds.
If $P_j$ holds and $1 \le j \le k$, then $\gamma \in C^j(I, U)$, so $x' = f \circ \gamma \in C^j(I, \mathbb{R}^n)$; hence $P_{j+1}$. Thus $P_{k+1}$.
*Structural Commentary*: Bootstrap $x' = f \circ \gamma$.
*Key Insight*: Vector-field regularity lifts the trajectory by one derivative. □

**Lemma 1.1.20.** *Generalized Gronwall Inequality*
*Let $\alpha, \beta, \psi \in C([0, T], \mathbb{R})$ with $\beta \ge 0$ and $\psi(t) \le \alpha(t) + \int_0^t \beta(s)\psi(s)\,ds$ $\forall t \in [0, T]$.*

*(i) $\psi(t) \le \alpha(t) + \int_0^t \alpha(s)\beta(s) \exp\left(\int_s^t \beta(r)\,dr\right) ds$ $\forall t \in [0, T]$.*

*(ii) If $\alpha(s) \le \alpha(t)$ whenever $0 \le s \le t \le T$, then $\psi(t) \le \alpha(t) \exp\left(\int_0^t \beta(s)\,ds\right)$ $\forall t \in [0, T]$.*

*Proof.*

(i) Put $q(t) := \int_0^t \beta(s)\psi(s)\,ds$ and $w(t) := \exp\left(-\int_0^t \beta(s)\,ds\right)$. Then $(wq)' = w\beta(\psi - q) \le w\beta\alpha$.
Since $q(0) = 0$, integration gives $q(t) \le \int_0^t \alpha(s)\beta(s) \exp\left(\int_s^t \beta(r)\,dr\right) ds$; combine this with $\psi \le \alpha + q$.

(ii) Under the monotonicity hypothesis, (i) gives $\psi(t) \le \alpha(t)\left(1 + \int_0^t \beta(s) \exp\left(\int_s^t \beta(r)\,dr\right) ds\right)$.
For $q(s) := \exp\left(\int_s^t \beta(r)\,dr\right)$, $q' = -\beta q$, so the parenthesis equals $1 + q(0) - q(t) = \exp\left(\int_0^t \beta(s)\,ds\right)$.

*Structural Commentary*: An integrating factor controls the accumulated integral term.
*Key Insight*: Exponential growth is the exact cost of repeated feedback. □

**Theorem 1.1.21.** *Continuous Dependence on Initial Data and Vector Fields*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f, g \in C(U, \mathbb{R}^n)$, $I := [t_0, T]$, and $x, y \in C^1(I, \mathbb{R}^n)$ satisfy $x' = f(\cdot, x)$, $y' = g(\cdot, y)$.*
*For $w : I \to \mathbb{R}^n$, set $\Gamma_w := \{(t, w(t)) : t \in I\}$. Assume $\Gamma_x \cup \Gamma_y \subseteq V \subseteq U$ with $V$ compact.*
*Assume $\exists L \geq 0$ such that $\|f(t,p) - f(t,q)\| \leq L \|p - q\|$ whenever $(t,p), (t,q) \in V$.*
*Set $a := \|x(t_0) - y(t_0)\|$, $E := f - g$, $M := \max_V \|E\|$, $\rho_0(h) := h$, and $\rho_L(h) := L^{-1}(e^{Lh} - 1)$ for $L > 0$.*
*Then $\|x(t) - y(t)\| \leq e^{L(t-t_0)}a + M\rho_L(t - t_0)$ $\forall t \in I$.*

*Proof.* Put $\Delta := x - y$, $\psi := \|\Delta\|$, and $R(s) := f(s, x(s)) - f(s, y(s)) + E(s, y(s))$.
Then $\Delta(t) = \Delta(t_0) + \int_{t_0}^t R(s)\, ds$ and $\|R(s)\| \leq L\psi(s) + M$. Hence $\psi(t) \leq a + \int_{t_0}^t (L\psi(s) + M)\, ds$.
If $L = 0$, then $\psi(t) \leq a + M(t - t_0) = a + M\rho_0(t - t_0)$.
If $L > 0$, set $\chi := \psi + M/L$. Then $\chi(t) \leq a + M/L + \int_{t_0}^t L\chi(s)\, ds$.
By Lemma 1.1.20(ii), $\chi(t) \leq (a + M/L)e^{L(t-t_0)}$, hence $\psi(t) \leq e^{L(t-t_0)}a + M\rho_L(t - t_0)$.
*Structural Commentary*: Split state error from field error; apply Gronwall.
*Key Insight*: Initial and model errors propagate exponentially. □

**Theorem 1.1.22.** *Local Lipschitz Dependence on the Initial Pair*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C(U, \mathbb{R}^n)$, and $(t_*, x_*) \in U$. Assume $f$ is locally state-Lipschitz:*
*$\forall$ compact $K \subseteq U$, $\exists L_K < +\infty$ such that $\|f(t,p) - f(t,q)\| \leq L_K \|p - q\|$ whenever $(t,p), (t,q) \in K$.*
*Then $\exists I, B, V, \Phi$ with $I \ni t_*$ compact, $B \ni x_*$ an open ball, $V \subseteq U$ compact, and $\Phi : I \times I \times B \to \mathbb{R}^n$.*
*Set $H := \sup I - \inf I$, $M := \max_V \|f\|$, choose a state-Lipschitz constant $L$ for $f$ on $V$.*
*Let $D((r,x),(q,y)) := \|x - y\| + M\,|r - q|$. Then:*

*(i) $\Phi(r, r, x) = x$, $(t, \Phi(t, r, x)) \in V$, and $\partial_t \Phi(t, r, x) = f(t, \Phi(t, r, x))$ for $(t, r, x) \in I \times I \times B$.*
*(ii) $\|\Phi(t, r, x) - \Phi(s, q, y)\| \leq e^{LH} D((r,x),(q,y)) + M\,|t - s|$ for $(t, r, x), (s, q, y) \in I \times I \times B$.*

*Proof.* Choose $a, \delta > 0$ with $Q := [t_* - a, t_* + a] \times \overline{B}_\delta(x_*) \subseteq U$, and set $M_Q := \max_Q \|f\|$.
Choose $0 < h < a/2$ with $2hM_Q < \delta/2$, set $I := [t_* - h, t_* + h]$, $B := B_{\delta/2}(x_*)$, and $V := I \times \overline{B}_\delta(x_*)$.
Set $H := 2h$, $M := \max_V \|f\|$, and choose a state-Lipschitz constant $L$ for $f$ on $V$.

(i) Fix $(r, x) \in I \times B$, set $I_r^+ := [r, t_* + h]$ and $I_r^- := [t_* - h, r]$.
Since $I_r^\pm \times \overline{B}_{\delta/2}(x) \subseteq Q$ and $|I_r^\pm| \leq 2h$, Theorem 1.1.16, applied forward on $I_r^+$ and backward on $I_r^-$, gives a unique $u_{r,x} \in C^1(I, \mathbb{R}^n)$ with $u_{r,x}(r) = x$ and $u'_{r,x} = f(\cdot, u_{r,x})$.
For $t \in I$, $\|u_{r,x}(t) - x\| \leq M_Q\,|t - r| < \delta/2$, hence $\|u_{r,x}(t) - x_*\| < \delta$ and $(t, u_{r,x}(t)) \in V$.
Set $\Phi(t, r, x) := u_{r,x}(t)$; then $\Phi(r, r, x) = x$, $(t, \Phi(t, r, x)) \in V$, and $\partial_t \Phi(t, r, x) = f(t, \Phi(t, r, x))$.

(ii) Let $(t, r, x), (s, q, y) \in I \times I \times B$, $p := \max\{r, q\}$, and $D_0 := D((r,x),(q,y))$.
By (i) and $\|\partial_t \Phi\| \leq M$, $\|\Phi(p, r, x) - \Phi(p, q, y)\| \leq D_0$.
Lemma 1.1.20(ii), after reversing the parameter when $t < p$, gives $\|\Phi(t, r, x) - \Phi(t, q, y)\| \leq e^{L|t-p|} D_0 \leq e^{LH} D_0$.
The bound $\|\Phi(t, q, y) - \Phi(s, q, y)\| \leq M\,|t - s|$ and the triangle inequality give (ii).

*Structural Commentary*: Compare at the later start time, then apply Gronwall.
*Key Insight*: Initial-time error first becomes state error. □

**Theorem 1.1.23.** *Smooth Dependence and First Variational Equations*
*Let $n, p \in \mathbb{N}$, $\Omega \subseteq \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^p$ be open, $f \in C^1(\Omega, \mathbb{R}^n)$, and $(t_0, x_0, \lambda_0) \in \Omega$.*
*Then $\exists I, B, \Lambda_0, \phi$ with $I \ni t_0$ compact, $B \ni x_0$ and $\Lambda_0 \ni \lambda_0$ open, and $\phi : I \times B \times \Lambda_0 \to \mathbb{R}^n$.*
*For $(x, \lambda) \in B \times \Lambda_0$, set $\gamma_{x,\lambda}(t) := (t, \phi(t, x, \lambda), \lambda)$.*
*Write $A_{x,\lambda} := D_x f \circ \gamma_{x,\lambda}$ and $G_{x,\lambda} := D_\lambda f \circ \gamma_{x,\lambda}$. The following hold:*

*(i) $\phi \in C^1(I \times B \times \Lambda_0, \mathbb{R}^n)$, $\phi(t_0, x, \lambda) = x$, and $\partial_t \phi(t, x, \lambda) = f(t, \phi(t, x, \lambda), \lambda)$ $\forall (t, x, \lambda) \in I \times B \times \Lambda_0$.*
*(ii) $D_x \phi(\cdot, x, \lambda)$ is the unique solution of $Y' = A_{x,\lambda} Y$, $Y(t_0) = I_n$, in $C^1(I, \mathbb{R}^{n \times n})$ $\forall (x, \lambda) \in B \times \Lambda_0$.*
*(iii) $D_\lambda \phi(\cdot, x, \lambda)$ is the unique solution of $Z' = A_{x,\lambda} Z + G_{x,\lambda}$, $Z(t_0) = 0_{n \times p}$, in $C^1(I, \mathbb{R}^{n \times p})$ $\forall (x, \lambda) \in B \times \Lambda_0$.*

*Proof.* Put $z := (x, \lambda)$, $z_0 := (x_0, \lambda_0)$, and define $F : \Omega \to \mathbb{R}^{n+p}$ by $F(t, z) := (f(t, x, \lambda), 0)$; then $F \in C^1(\Omega, \mathbb{R}^{n+p})$.
Choose $a, r > 0$ with $W := [t_0 - a, t_0 + a] \times \overline{B}_{2r}(z_0) \subseteq \Omega$. Set $H := \max_W \|F\|$.
Let $\tau := a$ if $H = 0$, and $\tau := \min\{a, \frac{r}{H}\}$ if $H > 0$. For $z \in B_r(z_0)$, $[t_0 - \tau, t_0 + \tau] \times \overline{B}_r(z) \subseteq W$.
By Lemma 1.1.15, choose a state-Lipschitz constant $L_W$ for $F$ on $W$, and set $I := [t_0 - \tau, t_0 + \tau]$.
By Theorem 1.1.16, applied forward and backward, $\forall z \in B_r(z_0)$, $\exists! \Phi_z \in C^1(I, \mathbb{R}^{n+p})$ with $\Phi_z(t_0) = z$ and $\Phi_z' = F(\cdot, \Phi_z)$. Define $\Phi(t, z) := \Phi_z(t)$.
Choose open $B, \Lambda_0$ with $x_0 \in B$, $\lambda_0 \in \Lambda_0$, and $B \times \Lambda_0 \subseteq B_r(z_0)$.

(i) Since the last $p$ components of $F$ vanish, $\Phi(t, x, \lambda) = (\phi(t, x, \lambda), \lambda)$; hence $\phi(t_0, x, \lambda) = x$ and $\partial_t \phi(t, x, \lambda) = f(t, \phi(t, x, \lambda), \lambda)$.

By Theorem 1.1.21, applied forward and backward, $\|\Phi_z - \Phi_w\|_\infty \le e^{\tau L_W}\|z-w\|$ for $z,w \in B_r(z_0)$. Hence $\Phi$ and $\partial_t\Phi = F(t,\Phi)$ are continuous.
Fix $z \in B_r(z_0)$ and set $\mathcal{L}_z(t) := D_zF(t,\Phi_z(t))$. Since $\mathcal{L}_z$ is bounded on $I$, apply Theorem 1.1.16 finitely many times forward and backward to obtain $P_z \in C^1(I,\mathbb{R}^{(n+p)\times(n+p)})$ with $P_z' = \mathcal{L}_zP_z$ and $P_z(t_0) = I_{n+p}$.
Set $C_0 := \max_W \|D_zF\|$; Lemma 1.1.20(ii) gives $\|P_z\|_\infty \le e^{2\tau C_0}$.
For $h \to 0$ with $z+h \in B_r(z_0)$, put $\Delta_h := \Phi_{z+h} - \Phi_z$ and $e_h := \Delta_h - P_zh$.
Set $\eta_{h,s} := \Phi_z + s\Delta_h$, $\mathcal{R}_h(t,s) := D_zF(t,\eta_{h,s}(t)) - \mathcal{L}_z(t)$; $R_h(t) := \int_0^1 \mathcal{R}_h(t,s)\,ds$, $\varepsilon_h := \|R_h\|_\infty$; $\varepsilon_h \to 0$.
$e_h' = \mathcal{L}_ze_h + R_h\Delta_h$, $e_h(t_0) = 0$, and $\|\Delta_h\|_\infty \le e^{\tau L_W}\|h\|$. Lemma 1.1.20(ii) gives $\|e_h\|_\infty = O(\varepsilon_h\|h\|) = o(\|h\|)$; thus $D_z\Phi = P_z$.
For $z,w \in B_r(z_0)$, $P_z - P_w$ satisfies $(P_z-P_w)' = \mathcal{L}_z(P_z-P_w) + (\mathcal{L}_z-\mathcal{L}_w)P_w$ and $(P_z-P_w)(t_0) = 0$. Lemma 1.1.20(ii) gives $\|P_z-P_w\|_\infty \le 2\tau e^{4\tau C_0}\|\mathcal{L}_z-\mathcal{L}_w\|_\infty \to 0$ as $z \to w$. Thus $\Phi$, hence $\phi$, is $C^1$.

(ii) Fix $(x,\lambda)$ and set $z = (x,\lambda)$. Put $Y := D_x\phi(\cdot\,,x,\lambda)$, $A := A_{x,\lambda}$, and $G := G_{x,\lambda}$.
Here $\mathcal{L}_z(\xi,\mu) = (A\xi + G\mu, 0)$ and $D_x\Phi = (Y,0)$. The $x$-columns of $P_z' = \mathcal{L}_zP_z$ give $Y' = AY$, $Y(t_0) = I_n$.
If $\widetilde{Y}$ also solves this IVP, $W := Y - \widetilde{Y}$ satisfies $W' = AW$, $W(t_0) = 0$; Lemma 1.1.20(ii) gives $W = 0$.

(iii) Fix $(x,\lambda)$ and set $z = (x,\lambda)$. Put $Z := D_\lambda\phi(\cdot\,,x,\lambda)$, $A := A_{x,\lambda}$, and $G := G_{x,\lambda}$.
Here $\mathcal{L}_z(\xi,\mu) = (A\xi + G\mu, 0)$. Since $D_\lambda\Phi = (Z, I_p)$, the $\lambda$-columns give $Z' = AZ + G$, $Z(t_0) = 0_{n\times p}$.
If $\widetilde{Z}$ also solves this IVP, $W := Z - \widetilde{Z}$ satisfies $W' = AW$, $W(t_0) = 0$; Lemma 1.1.20(ii) gives $W = 0$.

*Structural Commentary*: Lift parameters; differentiate the extended flow.
*Key Insight*: Flow derivatives satisfy variational equations. □

**Theorem 1.1.24.** *Higher-Order Smooth Dependence*
*Let $k,n,p \in \mathbb{N}$, $k \ge 1$, $\Omega \subseteq \mathbb{R}\times\mathbb{R}^n\times\mathbb{R}^p$ be open, $f \in C^k(\Omega,\mathbb{R}^n)$, $(t_0,x_0,\lambda_0) \in \Omega$. Then $\exists$ compact interval $I \ni t_0$, open neighborhoods $B \subseteq \mathbb{R}^n$ of $x_0$ and $\Lambda_0 \subseteq \mathbb{R}^p$ of $\lambda_0$, and $\phi : I\times B\times\Lambda_0 \to \mathbb{R}^n$ such that:*

*(i) $\phi(t_0,x,\lambda) = x$ and $\partial_t\phi(t,x,\lambda) = f(t,\phi(t,x,\lambda),\lambda)$ $\forall (t,x,\lambda) \in I\times B\times\Lambda_0$.*
*(ii) $\phi \in C^k(I\times B\times\Lambda_0,\mathbb{R}^n)$.*

*Proof.* Let $(\mathsf{H}_j)_d$ denote $C^j$ dependence of the local solution map on time and initial state for a $C^j$ vector field in $\mathbb{R}^d$. By Theorem 1.1.23(i), applied with a dummy parameter, $(\mathsf{H}_1)_d$ holds $\forall d \in \mathbb{N}$.
Assume $(\mathsf{H}_j)_d$ holds $\forall d \in \mathbb{N}$. Let $\Omega_d \subseteq \mathbb{R}\times\mathbb{R}^d$ be open, $F \in C^{j+1}(\Omega_d,\mathbb{R}^d)$, $(t_0,z_0) \in \Omega_d$, and let $\Psi$ be the local solution map near $(t_0,z_0)$. By $(\mathsf{H}_j)_d$, $\Psi \in C^j$.
By Theorem 1.1.23(ii), $Q := D_z\Psi$ satisfies $Q' = D_zF(\cdot\,,\Psi)Q$ and $Q(t_0,z) = I_d$.
Define $\mathcal{H}(t,\zeta,P) := (F(t,\zeta), D_zF(t,\zeta)P)$ and $\Xi := (\Psi,Q)$. Then $\mathcal{H} \in C^j$, $\Xi' = \mathcal{H}(\cdot\,,\Xi)$, and $\Xi(t_0,z) = (z,I_d)$.
By $(\mathsf{H}_j)_{d+d^2}$, the local solution map $\Theta$ for $\mathcal{H}$ is $C^j$. Local uniqueness gives $\Xi(t,z) = \Theta(t,z,I_d)$, so $Q \in C^j$.
Thus $D_z\Psi = Q \in C^j$ and $\partial_t\Psi = F(\cdot\,,\Psi) \in C^j$, whence $\Psi \in C^{j+1}$. Therefore $(\mathsf{H}_{j+1})_d$ holds $\forall d \in \mathbb{N}$, and induction proves $(\mathsf{H}_k)_d$ $\forall k,d \in \mathbb{N}$ with $k \ge 1$.
Now put $m := n+p$, $z := (x,\lambda)$, $z_0 := (x_0,\lambda_0)$, and $\widetilde{F}(t,z) := (f(t,x,\lambda),0_p)$. By $(\mathsf{H}_k)_m$, after shrinking around $(t_0,z_0)$, there are a compact interval $I \ni t_0$, an open $U_0 \ni z_0$, and a local solution map $\Phi \in C^k(I\times U_0,\mathbb{R}^m)$ for $\widetilde{F}$.
Choose open $B \ni x_0$ and $\Lambda_0 \ni \lambda_0$ with $B\times\Lambda_0 \subseteq U_0$, and put $\phi := \pi_{\mathbb{R}^n}\circ\Phi$ on $I\times B\times\Lambda_0$.

(i) Since the last $p$ components of $\widetilde{F}$ vanish, $\Phi(t,x,\lambda) = (\phi(t,x,\lambda),\lambda)$. Hence $\phi(t_0,x,\lambda) = x$ and $\partial_t\phi(t,x,\lambda) = f(t,\phi(t,x,\lambda),\lambda)$.
(ii) Since $\Phi \in C^k$ and $\pi_{\mathbb{R}^n}$ is linear, $\phi \in C^k(I\times B\times\Lambda_0,\mathbb{R}^n)$.

*Structural Commentary*: Differentiate once; bootstrap the augmented flow.
*Key Insight*: Variational dynamics raise flow regularity. □

**Proposition 1.1.25.** *Regular Perturbation Expansion*
*Let $k,n \in \mathbb{N}$, $k \ge 1$, let $\Omega \subseteq \mathbb{R}\times\mathbb{R}^n\times\mathbb{R}$ be open, $f \in C^k(\Omega,\mathbb{R}^n)$, and $(t_0,x_0,\varepsilon_0) \in \Omega$.*
*Let $I \ni t_0$ be compact, $\Lambda_0 \ni \varepsilon_0$ open, and $\phi \in C^k(I\times\Lambda_0,\mathbb{R}^n)$ satisfy $\phi(t_0,\varepsilon) = x_0$ and $\partial_t\phi(t,\varepsilon) = f(t,\phi(t,\varepsilon),\varepsilon)$.*
*Write $Z_j := (z_0,\dots,z_j)$, $Z_j^+ := (z_1,\dots,z_{j+1})$, and $\mathcal{D}_jg := \partial_\varepsilon g + \sum_{r=0}^{j} D_{z_r}g\,z_{r+1}$ for $g = g(t,Z_j,\varepsilon)$.*
*Set $f_0(t,z_0,\varepsilon) := f(t,z_0,\varepsilon)$, $f_{j+1} := \mathcal{D}_jf_j$ for $0 \le j < k$, $\phi_j(t) := \partial_\varepsilon^j\phi(t,\varepsilon_0)$, and $\Phi_j(t) := (\phi_0(t),\dots,\phi_j(t))$.*
*Then:*

*(i) For $0 \le j \le k$, $\phi_j' = f_j(t,\Phi_j,\varepsilon_0)$, with $\phi_0(t_0) = x_0$ and $\phi_j(t_0) = 0$ for $1 \le j \le k$.*
*(ii) The remainder $r_k(t,\varepsilon) := \phi(t,\varepsilon) - \sum_{j=0}^{k}\frac{\phi_j(t)}{j!}(\varepsilon-\varepsilon_0)^j$ satisfies $\rho_k(\varepsilon) := \sup_{t\in I}\|r_k(t,\varepsilon)\| = o(|\varepsilon-\varepsilon_0|^k)$.*
*(iii) If $f \in C^{k+1}(\Omega,\mathbb{R}^n)$, then, after shrinking $I$ and $\Lambda_0$ if necessary, $\rho_k(\varepsilon) = O(|\varepsilon-\varepsilon_0|^{k+1})$.*

*Proof.* For $0 \le j \le k$, put $z_j(t,\varepsilon) := \partial_\varepsilon^j \phi(t,\varepsilon)$, $Z_j(t,\varepsilon) := (z_0(t,\varepsilon), \dots, z_j(t,\varepsilon))$, and $R_j(t,\varepsilon) := f_j(t, Z_j(t,\varepsilon), \varepsilon)$.

(i) Since $\partial_t \phi = f(t,\phi,\varepsilon)$ and $\phi(t_0,\varepsilon) = x_0$, $\phi(t,\varepsilon) = x_0 + \int_{t_0}^t f(s,\phi(s,\varepsilon),\varepsilon)\,ds$.
By the chain rule, $\partial_\varepsilon R_j = R_{j+1}$ for $0 \le j < k$; hence by induction, $\partial_\varepsilon^j [f(t,\phi(t,\varepsilon),\varepsilon)] = R_j(t,\varepsilon)$.
By differentiation under the integral sign, $z_j(t,\varepsilon) = \delta_{j0} x_0 + \int_{t_0}^t R_j(s,\varepsilon)\,ds$.
At $\varepsilon = \varepsilon_0$, $z_j = \phi_j$, $R_j = f_j(t,\Phi_j,\varepsilon_0)$, and $z_j(t_0,\varepsilon_0) = \delta_{j0} x_0$; differentiating in $t$ proves (i).

(ii) Put $h := \varepsilon - \varepsilon_0$ and $G_t(\eta) := \partial_\varepsilon^k \phi(t,\eta) - \partial_\varepsilon^k \phi(t,\varepsilon_0)$.
By the integral Taylor formula, $r_k(t,\varepsilon) = \frac{h^k}{(k-1)!} \int_0^1 (1-s)^{k-1} G_t(\varepsilon_0 + sh)\,ds$.
Set $\omega(h) := \sup_{t\in I,\ s\in[0,1]} \|G_t(\varepsilon_0 + sh)\|$. By uniform continuity, $\omega(h) \to 0$, and $\rho_k(\varepsilon) \le \frac{|h|^k}{k!}\omega(h)$.

(iii) If $f \in C^{k+1}$, Theorem 1.1.24 produces a $C^{k+1}$ local solution map for the same initial data. Local uniqueness from Theorem 1.1.16, after shrinking $I \times \Lambda_0$, identifies this map with $\phi$.
Choose compact $J \subseteq \Lambda_0$ with $\varepsilon_0 \in J^\circ$ and set $C_J := \max_{(t,\eta)\in I\times J} \left\|\partial_\varepsilon^{k+1}\phi(t,\eta)\right\|$.
For $\varepsilon \in J$, by the integral remainder, $\rho_k(\varepsilon) \le \frac{C_J |h|^{k+1}}{(k+1)!}$.

*Structural Commentary*: Differentiate, then Taylor expand.
*Key Insight*: Coefficients satisfy forced IVPs. □

**Example 1.1.26.** *Linear Drag as a Regular Perturbation*
Let $g > 0$, $\varepsilon \ge 0$, and $v' = -\varepsilon v - g$, $v(0) = 0$. Then $v(t,\varepsilon) = \frac{g(e^{-\varepsilon t}-1)}{\varepsilon}$ if $\varepsilon > 0$, and $v_0(t) := v(t,0) = -gt$.
At $\varepsilon_0 = 0$, $v_1' = -v_0 = gt$, $v_1(0) = 0$, hence $v_1(t) = \frac{gt^2}{2}$ and $v(t,\varepsilon) = v_0(t) + \varepsilon v_1(t) + o(\varepsilon)$ on compact $t$-intervals.
For fixed $\varepsilon > 0$, $t \to +\infty$ gives $v(t,\varepsilon) \to -\frac{g}{\varepsilon}$ but $v_0(t) + \varepsilon v_1(t) \to +\infty$; the expansion is local for bounded $t$.

**Theorem 1.1.27.** *Unique Maximal Solution*
*Let $U \subseteq \mathbb{R}\times\mathbb{R}^n$ be open, $f \in C(U,\mathbb{R}^n)$, and $(t_0,x_0) \in U$.*
*Assume $\forall$ compact $V \subseteq U$, $\exists L_V < +\infty$ such that $\|f(t,p) - f(t,q)\| \le L_V \|p - q\|$ whenever $(t,p),(t,q) \in V$.*
*For an open interval $J \ni t_0$, write $u \in \mathcal{A}(J)$ iff $u \in C^1(J,\mathbb{R}^n)$, $u(t_0) = x_0$, $(t,u(t)) \in U$, and $u' = f(\cdot,u)$.*
*Let $\mathcal{S}$ be all such pairs $(J,u)$. For $(J,u),(K,v) \in \mathcal{S}$, write $(J,u) \preceq (K,v)$ iff $J \subseteq K$ and $v|_J = u$. Then:*

*(i) $\exists$ a maximal $(I_*,\phi_*) \in \mathcal{S}$ with $I_* = (T_-,T_+)$.*
*(ii) $(J,u) \preceq (I_*,\phi_*)$ $\forall (J,u) \in \mathcal{S}$; hence the maximal solution is unique.*

*Proof.* By Theorem 1.1.16, applied forward and backward, $\mathcal{S} \neq \emptyset$. For $(J,u),(K,v) \in \mathcal{S}$, set $E := \{t \in J \cap K : u(t) = v(t)\}$.
Then $t_0 \in E$, $E$ is closed, and local uniqueness at each $s \in E$ makes $E$ open in $J \cap K$.
Since $J \cap K$ is an interval, $E = J \cap K$; thus $u = v$ on $J \cap K$ $\forall (J,u),(K,v) \in \mathcal{S}$.

(i) Set $I_* := \bigcup_{(J,u)\in\mathcal{S}} J$ and let $\phi_*(t) := u(t)$ for any $(J,u) \in \mathcal{S}$ with $t \in J$. Agreement on overlaps makes $\phi_*$ well defined.
Since all $J$ are open intervals containing $t_0$, $I_* = (T_-,T_+)$. Locally $\phi_* = u$, so $(I_*,\phi_*) \in \mathcal{S}$.
If $(I_*,\phi_*) \preceq (\widehat{I},\widehat{\phi}) \in \mathcal{S}$, then $I_* \subseteq \widehat{I} \subseteq I_*$; hence $\widehat{I} = I_*$.

(ii) If $(J,u) \in \mathcal{S}$, then $J \subseteq I_*$ and overlap agreement gives $u = \phi_*|_J$; hence $(J,u) \preceq (I_*,\phi_*)$.
If $(K,\psi)$ is maximal, then $(K,\psi) \preceq (I_*,\phi_*)$; maximality gives $K = I_*$ and $\psi = \phi_*$.

*Structural Commentary*: Local uniqueness makes solution germs compatible.
*Key Insight*: The maximal solution is their union. □

**Lemma 1.1.28.** *Extension Criterion at a Finite Endpoint*
*Let $U \subseteq \mathbb{R}\times\mathbb{R}^n$ be open, $f \in C(U,\mathbb{R}^n)$, and assume $f$ is locally Lipschitz in $x$ on compact subsets of $U$.*
*Let $\phi \in C^1((T_-,T_+),\mathbb{R}^n)$ satisfy $\phi' = f(\cdot,\phi)$, $(t,\phi(t)) \in U$, and $T_+ < +\infty$.*
*Then $\exists \varepsilon > 0$ and a solution $\widehat{\phi} : (T_-,T_+ + \varepsilon) \to \mathbb{R}^n$ with $\widehat{\phi}|_{(T_-,T_+)} = \phi$ iff $\exists t_m \uparrow T_+$ and $y \in \mathbb{R}^n$ such that $\phi(t_m) \to y$ and $(T_+,y) \in U$.*

*Proof.*
($\Rightarrow$) Let $\widehat{\phi}$ be such an extension, choose $t_m \uparrow T_+$, and put $y := \widehat{\phi}(T_+)$. Then $\phi(t_m) = \widehat{\phi}(t_m) \to y$ and $(T_+,y) \in U$.

($\Leftarrow$) Choose $r,\eta > 0$ with $W := [T_+ - \eta, T_+ + \eta] \times \overline{B}_r(y) \subseteq U$ and set $M := \max_W \|f\|$.
Choose $m$ large enough that $t_m > T_+ - \eta$, $a := \|\phi(t_m) - y\| < \frac{r}{2}$, and $Mh < \frac{r}{2}$ for $h := T_+ - t_m$; put $J := [t_m, T_+)$ and $B := B_r(y)$.
If $\tau \in J$ is the first exit from $B$, then $r \le a + M(\tau - t_m) \le a + Mh < r$, impossible. Hence $\phi(J) \subseteq B$.
For $s,t \in J$, $\|\phi(t) - \phi(s)\| \le M|t - s|$, so $\phi(t)$ is Cauchy as $t \uparrow T_+$. Since $\phi(t_m) \to y$, $\lim_{t\uparrow T_+} \phi(t) = y$.
By Theorem 1.1.16, $\exists \varepsilon > 0$ and a solution $\widetilde{\phi}$ on $[T_+, T_+ + \varepsilon)$ with $\widetilde{\phi}(T_+) = y$. Define $\widehat{\phi} = \phi$ on $(T_-,T_+)$ and $\widehat{\phi} = \widetilde{\phi}$ on $[T_+, T_+ + \varepsilon)$.

Continuity of $f$ gives $\phi'(t) \to f(T_+, y) = \widetilde{\phi}'(T_+)$; hence $\widehat{\phi} \in C^1((T_-, T_+ + \varepsilon), \mathbb{R}^n)$ and is the required extension.

*Structural Commentary*: A convergent endpoint subsequence traps the tail for gluing.
*Key Insight*: Finite endpoint limits are exactly extendible states. □

**Corollary 1.1.29.** *Compact Escape at a Finite Maximal Endpoint*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open and $f \in C(U, \mathbb{R}^n)$.*
*Assume $\forall C \Subset U$, $\exists L_C < +\infty$ such that $(t,p),(t,q) \in C \Rightarrow \|f(t,p) - f(t,q)\| \le L_C \|p - q\|$.*
*Let $J := (T_-, T_+)$ and let $\phi : J \to \mathbb{R}^n$ be maximal for $x' = f(t,x)$, with $\gamma(t) := (t, \phi(t)) \in U$. Then:*

*(i)* $T_+ < +\infty \Rightarrow \forall K \Subset U,\ \exists \tau < T_+ : \gamma((\tau, T_+)) \cap K = \emptyset$.
*(ii)* $T_- > -\infty \Rightarrow \forall K \Subset U,\ \exists \tau > T_- : \gamma((T_-, \tau)) \cap K = \emptyset$.
*(iii) If $U = \mathbb{R} \times \mathbb{R}^n$, then $T_+ < +\infty \Rightarrow \|\phi(t)\| \to +\infty$ as $t \uparrow T_+$ and $T_- > -\infty \Rightarrow \|\phi(t)\| \to +\infty$ as $t \downarrow T_-$.*

*Proof.*

(i) If (i) fails for some $K \Subset U$, then $\exists t_m \uparrow T_+$ with $\gamma(t_m) \in K$. By compactness, after passing to a subsequence,
$\gamma(t_m) \to (T_+, y) \in K \subseteq U$. By Lemma 1.1.28, $\phi$ admits a proper extension past $T_+$, contradiction.

(ii) Set $\widetilde{U} := \{(s,x) : (-s,x) \in U\}$, $\widetilde{f}(s,x) := -f(-s,x)$, $\widetilde{J} := (-T_+, -T_-)$, and $u(s) := \phi(-s)$.
Then $u : \widetilde{J} \to \mathbb{R}^n$ is maximal for $u' = \widetilde{f}(\cdot, u)$. For $K \Subset U$, put $\widetilde{K} := \{(-t,x) : (t,x) \in K\} \Subset \widetilde{U}$.
By (i) applied to $u$ at $-T_-$, $\exists a < -T_-$ such that $(s, u(s)) \notin \widetilde{K}$ for $a < s < -T_-$.
With $\tau := -a$, $\gamma((T_-, \tau)) \cap K = \emptyset$.

(iii) Assume $U = \mathbb{R} \times \mathbb{R}^n$ and $T_+ < +\infty$. For $R > 0$, set $K_R := [T_+ - 1, T_+] \times \overline{B}_R(0)$.
By (i), $\exists \tau < T_+$ such that $\gamma((\tau, T_+)) \cap K_R = \emptyset$, hence $\|\phi(t)\| > R$ for $\max\{\tau, T_+ - 1\} < t < T_+$.
Since $R$ is arbitrary, $\|\phi(t)\| \to +\infty$ as $t \uparrow T_+$. If $T_- > -\infty$, apply (ii) to $K_R^- := [T_-, T_- + 1] \times \overline{B}_R(0)$ to obtain $\|\phi(t)\| > R$ near $T_-$; hence $\|\phi(t)\| \to +\infty$ as $t \downarrow T_-$.

*Structural Commentary*: Compact recurrence creates an endpoint extension.
*Key Insight*: Finite maximal time forces compact escape. □

**Corollary 1.1.30.** *Global Existence under Linear Growth*
*Let $f : \mathbb{R} \times \mathbb{R}^n \to \mathbb{R}^n$ be continuous.*
*Assume (LL): $\forall$ compact $V \subseteq \mathbb{R} \times \mathbb{R}^n$, $\exists L_V < +\infty$ such that $\|f(t,p) - f(t,q)\| \le L_V \|p - q\|\ \forall (t,p),(t,q) \in V$.*
*Assume (LG): $\forall T > 0$, $\exists M_T, L_T \ge 0$ such that $\|f(t,x)\| \le M_T + L_T \|x\|\ \forall (t,x) \in [-T,T] \times \mathbb{R}^n$.*
*If $\phi : (T_-, T_+) \to \mathbb{R}^n$ is maximal and $\phi' = f(\cdot, \phi)$, then $(T_-, T_+) = \mathbb{R}$.*

*Proof.* Fix $t_0 \in (T_-, T_+)$, $x_0 := \phi(t_0)$. If $T_+ < +\infty$, choose $T > \max\{|t_0|, |T_+|\}$ and $M_T, L_T$ from (LG), and set $C_T := (\|x_0\| + M_T(T_+ - t_0))e^{L_T(T_+ - t_0)}$.
With $q(t) := \|\phi(t)\|$, $q(t) \le \|x_0\| + \int_{t_0}^t (M_T + L_T q(s))\, ds$ on $[t_0, T_+)$. Lemma 1.1.20(ii) gives $q(t) \le (\|x_0\| + M_T(t - t_0))e^{L_T(t-t_0)} \le C_T < +\infty$.
Thus $\Gamma_+ := \{(t, \phi(t)) : t \in [t_0, T_+)\} \subseteq K_T := [-T, T] \times \overline{B}_{C_T}(0)$. For $t_m \uparrow T_+$, compactness gives $\phi(t_{m_j}) \to y$ and $(T_+, y) \in K_T$. By (LL) and Lemma 1.1.28, $\phi$ extends past $T_+$, contradiction. Hence $T_+ = +\infty$.
Set $g(s,x) := -f(t_0 - s, x)$ and $u(s) := \phi(t_0 - s)$ on $(t_0 - T_+, t_0 - T_-)$. Then $u' = g(\cdot, u)$, $u$ is maximal, and $g$ satisfies (LL),(LG) on bounded $s$-intervals. Apply right-endpoint result to $u$, then $t_0 - T_- = +\infty$, $T_- = -\infty$.
*Structural Commentary*: Gronwall bounds; compactness forces extension.
*Key Insight*: Linear growth rules out finite escape. □

**Theorem 1.1.31.** *Peano Local Existence Theorem*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C(U, \mathbb{R}^n)$, and $(t_0, x_0) \in U$. Choose $T, \delta > 0$ with $V := [t_0, t_0 + T] \times \overline{B}_\delta(x_0) \subseteq U$.*
*Set $M := \max_V \|f\|$, $T_0 := \min\{T, \delta/M\}$ with $\delta/0 := +\infty$, and $I := [t_0, t_0 + T_0]$.*
*For $y : I \to \mathbb{R}^n$, write $\Gamma_y := \{(t, y(t)) : t \in I\}$. Then:*

*(i) $\exists \phi \in C(I, \mathbb{R}^n)$ with $\Gamma_\phi \subseteq V$ and $\phi = x_0 + \int_{t_0}^{(\cdot)} f(s, \phi(s))\, ds$ on $I$.*
*(ii) This $\phi$ lies in $C^1(I, \mathbb{R}^n)$ and satisfies $\phi(t_0) = x_0$, $\phi' = f(\cdot, \phi)$.*

*Proof.* For $N \ge 1$, set $h_N := T_0/N$, $t_j := t_0 + j h_N$, $\tau_N(t) := t_j$ on $[t_j, t_{j+1})$ for $0 \le j < N$, and $\tau_N(t_N) := t_{N-1}$.
Construct $x_N$ recursively by $x_N(t_0) := x_0$ and $x_N(t) := x_N(t_j) + (t - t_j) f(t_j, x_N(t_j))$ for $t \in [t_j, t_{j+1}]$, $0 \le j < N$.
Write $(\mathcal{I}G)(t) := \int_{t_0}^t G(s)\, ds$, $Y_N := (\tau_N, x_N \circ \tau_N)$, and $F_N := f \circ Y_N$; then $x_N = x_0 + \mathcal{I} F_N$.
By induction, $\|x_N(t) - x_0\| \le M(t - t_0) \le \delta$ and $\|x_N(t) - x_N(s)\| \le M\,|t - s|$.
Hence $\Gamma_{x_N} \subseteq V$ and $(x_N)$ is uniformly bounded/equicontinuous.

(i) By Theorem 8.2.1, applied coordinatewise, $x_{N_k} \to \phi \in C(I, \mathbb{R}^n)$ uniformly.
Then $\Gamma_\phi \subseteq V$. Set $Y_\phi := (\mathrm{id}_I, \phi)$ and $f_\phi := f \circ Y_\phi$.

Since $\|\mathrm{id}_I - \tau_N\|_\infty \le h_N$ and $\|x_N - x_N \circ \tau_N\|_\infty \le M h_N$, $Y_{N_k}(I), Y_\phi(I) \subseteq V$ and $Y_{N_k} \to Y_\phi$ uniformly.
By uniform continuity of $f|_V$, $F_{N_k} \to f_\phi$ uniformly.
Passing to the limit in $x_{N_k} = x_0 + \mathcal{I} F_{N_k}$ gives $\phi = x_0 + \mathcal{I} f_\phi$.

(ii) Since $f_\phi \in C(I, \mathbb{R}^n)$, by Theorem 2.2.1 applied to (i), $\phi \in C^1(I, \mathbb{R}^n)$, $\phi(t_0) = x_0$, and $\phi' = f_\phi$.

*Structural Commentary*: Euler polygons plus Arzelà–Ascoli produce a convergent approximate trajectory.
*Key Insight*: Continuity yields existence through compactness, without uniqueness. □

**Definition 1.1.32.** *Osgood Modulus of Continuity*
Let $\rho \in C([0, +\infty), [0, +\infty))$.

(i) $\rho$ is a *modulus of continuity* iff $\rho(0) = 0$ and $\rho$ is nondecreasing.

(ii) A modulus $\rho$ satisfies the *Osgood condition* iff $\int_0^1 \frac{dr}{\rho(r)} = +\infty$, interpreted as an improper extended integral; equivalently, $\int_0^a \frac{dr}{\rho(r)} = +\infty \ \forall a > 0$.

*Key Insight*: Osgood divergence prevents a zero separation from becoming positive in finite time.

**Example 1.1.33.** *Lipschitz, Hölder, and Almost-Lipschitz Moduli*
For $0 < \alpha < 1$, set $\rho_1(r) := r$, $\rho_\alpha(r) := r^\alpha$, $\rho_{\log}(0) := 0$, and $\rho_{\log}(r) := r(1 + |\log r|)$ for $r > 0$.
These functions are continuous, nondecreasing, and vanish at 0.
Since $\int_0^1 \frac{dr}{r} = +\infty$ and $\int_0^1 r^{-\alpha}\, dr = \frac{1}{1-\alpha} < +\infty$, $\rho_1$ is Osgood while the Hölder modulus $\rho_\alpha$ is not.
For the almost-Lipschitz modulus, $s = -\log r$ gives $\int_0^1 \frac{dr}{r(1+|\log r|)} = \int_0^{+\infty} \frac{ds}{1+s} = +\infty$, so $\rho_{\log}$ is Osgood.

**Theorem 1.1.34.** *Osgood Uniqueness Criterion*
*Let $n \in \mathbb{N}$, $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C(U, \mathbb{R}^n)$, $(t_0, x_0) \in U$, $T > t_0$, and $\delta > 0$.*
*Put $J := [t_0, T]$, $V := J \times \overline{B}_\delta(x_0)$, and assume $V \subseteq U$.*
*Let $\rho \in C([0, +\infty), [0, +\infty))$ be nondecreasing, $\rho(0) = 0$, and $\int_0^a \frac{dr}{\rho(r)} = +\infty \ \forall a > 0$.*
*For $(t, p), (t, q) \in V$, set $A_t(p, q) := |\langle p - q, f(t, p) - f(t, q)\rangle|$.*
*Assume $\exists C \ge 0$ such that $A_t(p, q) \le C \|p - q\| \rho(\|p - q\|) \ \forall (t, p), (t, q) \in V$.*
*For $w \in C(J, \mathbb{R}^n)$, set $\Gamma_w := \{(t, w(t)) : t \in J\}$.*
*If $u, v \in C^1(J, \mathbb{R}^n)$ satisfy $\Gamma_u \cup \Gamma_v \subseteq V$, $u(t_0) = v(t_0) = x_0$, $u' = f(\cdot, u)$, and $v' = f(\cdot, v)$, then $u = v$ on $J$.*

*Proof.* Assume $S := \{t \in (t_0, T] : u(t) \ne v(t)\} \ne \emptyset$ and fix $t \in S$. Put $\Delta := u - v$ and $d := \|\Delta\|$.
For $s \in J$, set $E(s) := f(s, u(s)) - f(s, v(s))$. The set $Z_t := \{s \in [t_0, t] : d(s) = 0\}$ is nonempty and compact; set $a := \max Z_t$.
Then $a < t$, $d(a) = 0$, and $d > 0$ on $(a, t]$. On $(a, t]$, $d \in C^1$ and $d'(s) = \frac{\langle \Delta(s), E(s)\rangle}{d(s)}$. Hence $d'(s) \le \frac{A_s(u(s), v(s))}{d(s)} \le C\rho(d(s))$.
For $m \ge 1$, set $H_m(r) := \int_0^r \frac{dq}{\rho(q) + \frac{1}{m}}$. For $s \in (a, t]$, set $b_m(s) := \frac{\rho(d(s))}{\rho(d(s)) + \frac{1}{m}} \le 1$.
If $a < r < t$, then on $[r, t]$, $(H_m \circ d)'(s) = \frac{d'(s)}{\rho(d(s)) + \frac{1}{m}} \le C b_m(s) \le C$.
Thus $H_m(d(t)) - H_m(d(r)) \le C(t - r)$. Letting $r \downarrow a$ gives $H_m(d(t)) \le C(t - a)$.
By Theorem 1.3.1, $H_m(d(t)) \uparrow \int_0^{d(t)} \frac{dq}{\rho(q)} = +\infty$, contradiction. Hence $S = \emptyset$ and $u = v$.
*Structural Commentary*: Regularize $1/\rho$; separate variables.
*Key Insight*: Divergence forbids first separation. □

**Proposition 1.1.35.** *Time Translation and the Local Flow Law*
*Let $D \subseteq \mathbb{R}^n$ be open and $f : D \to \mathbb{R}^n$. For $(a, y) \in \mathbb{R} \times D$, let $\mathcal{S}(a, y)$ contain the pairs $(I, u)$ such that $I \ni a$ is an open interval, $u \in C^1(I, D)$, $u(a) = y$, and $u' = f(u)$.*
*Assume $\mathcal{S}(a, y) \ne \emptyset \ \forall (a, y) \in \mathbb{R} \times D$, and $u = v$ on $I \cap J \ \forall (a, y) \in \mathbb{R} \times D$, $\forall (I, u), (J, v) \in \mathcal{S}(a, y)$.*
*For $x \in D$, let $\phi_x : I_x \to D$ be the maximal solution with $0 \in I_x$, $\phi_x(0) = x$, and $\phi_x' = f(\phi_x)$.*
*Define $\mathcal{D}_\Phi := \{(t, x) \in \mathbb{R} \times D : t \in I_x\}$ and $\Phi_t(x) := \phi_x(t)$ for $(t, x) \in \mathcal{D}_\Phi$.*

*(i) If $\phi : I \to D$, $0 \in I$, and $\phi' = f(\phi)$, then $\psi_\tau(t) := \phi(t - \tau)$ on $\tau + I$ satisfies $\psi_\tau' = f(\psi_\tau) \ \forall \tau \in \mathbb{R}$. Its initial value is $\psi_\tau(\tau) = \phi(0)$.*

*(ii) $(0, x) \in \mathcal{D}_\Phi$ and $\Phi_0(x) = x \ \forall x \in D$; hence $\Phi_0 = \mathrm{id}_D$.*

*(iii) If $(s, x), (t, \Phi_s(x)), (t + s, x) \in \mathcal{D}_\Phi$, then $\Phi_{t+s}(x) = \Phi_t(\Phi_s(x))$.*

*Proof.*

(i) For $t \in \tau + I$, $\psi_\tau'(t) = \phi'(t - \tau) = f(\phi(t - \tau)) = f(\psi_\tau(t))$ and $\psi_\tau(\tau) = \phi(0)$.

(ii) Since $0 \in I_x$, $(0, x) \in \mathcal{D}_\Phi$ and $\Phi_0(x) = \phi_x(0) = x \ \forall x \in D$.

(iii) Set $R := (I_x - s) \cap I_{\Phi_s(x)}$. Then $0, t \in R$ and $(r + s, x), (r, \Phi_s(x)) \in \mathcal{D}_\Phi$ for $r \in R$.
Define $u, v : R \to D$ by $u(r) := \Phi_{r+s}(x)$, $v(r) := \Phi_r(\Phi_s(x))$.
Then $u' = f(u)$ by (i), $v' = f(v)$, $u(0) = v(0) = \Phi_s(x)$.
Uniqueness gives $u = v$ on $R$, so $\Phi_{t+s}(x) = \Phi_t(\Phi_s(x))$.

*Structural Commentary*: Shift time; use uniqueness.
*Key Insight*: Autonomy makes elapsed time compose. □

## 1.2 Linear Systems and Stability

Scalar first-order linear equations admit an explicit evolution factor and a variation-of-constants representation.

**Proposition 1.2.1.** *Scalar Variation of Constants*
*Let $I \subseteq \mathbb{R}$ be open, $a, g \in C(I, \mathbb{R})$, and $(t_0, x_0) \in I \times \mathbb{R}$.*
*Put $E(t,s) := \exp\left(\int_s^t a(r)\,dr\right)$ and $\phi(t) := E(t,t_0)x_0 + \int_{t_0}^t E(t,s)g(s)\,ds$. Then:*

*(i) $\phi \in C^1(I,\mathbb{R})$ and $(\phi' - a\phi, \phi(t_0)) = (g, x_0)$.*
*(ii) If $x \in C^1(I,\mathbb{R})$ and $(x' - ax, x(t_0)) = (g, x_0)$, then $x = \phi$.*

*Proof.*

(i) Let $B(t) := x_0 + \int_{t_0}^t E(t_0,s)g(s)\,ds$. By additivity, $E(t,s) = E(t,t_0)E(t_0,s)$, so $\phi(t) = E(t,t_0)B(t)$.
By Theorem 2.2.1, $B \in C^1$, $\partial_t E(t,t_0) = a(t)E(t,t_0)$, and $B'(t) = E(t_0,t)g(t)$.
Hence $\phi \in C^1$, $\phi'(t) = a(t)\phi(t) + E(t,t_0)E(t_0,t)g(t) = a(t)\phi(t) + g(t)$, and $\phi(t_0) = x_0$.
(ii) If $(x' - ax, x(t_0)) = (g, x_0)$, then $\frac{d}{dt}(E(t_0,t)x(t)) = E(t_0,t)(x'(t) - a(t)x(t)) = E(t_0,t)g(t)$.
By Theorem 2.2.2, $E(t_0,t)x(t) = x_0 + \int_{t_0}^t E(t_0,s)g(s)\,ds$.
Multiplying by $E(t,t_0)$ and using $E(t,t_0)E(t_0,s) = E(t,s)$ gives $x(t) = \phi(t)$.

*Structural Commentary*: $E(t_0,t)$ removes $a(t)x$; integration accumulates $g$.
*Key Insight*: $(E(t_0,t)x)' = E(t_0,t)g$. □

**Proposition 1.2.2.** *Fiber-Preserving Change of Variables*
*Let $\Omega, \widetilde{\Omega} \subseteq \mathbb{R}^2$ be open, $f : \Omega \to \mathbb{R}$, and $T : \Omega \to \widetilde{\Omega}$ be a $C^1$ diffeomorphism.*
*Assume $T(t,x) = (\sigma(t), \eta(t,x))$ and $T^{-1}(s,y) = (\tau(s), \xi(s,y))$. Put $G := \partial_t\eta + \partial_x\eta\, f$.*
*Define $\widetilde{f} : \widetilde{\Omega} \to \mathbb{R}$ by $\widetilde{f}(s,y) := \tau'(s)G(\tau(s), \xi(s,y))$.*
*Let $I \subseteq \mathbb{R}$ be an interval, $x \in C^1(I,\mathbb{R})$, $\Gamma_x := \{(t, x(t)) : t \in I\} \subseteq \Omega$, and $J := \sigma(I)$.*
*Set $y : J \to \mathbb{R}$ by $y(s) := \eta(\tau(s), x(\tau(s)))$. Then $y \in C^1(J,\mathbb{R})$, and $x' = f(\cdot\,, x)$ on $I$ iff $y' = \widetilde{f}(\cdot\,, y)$ on $J$.*

*Proof.* Since $\det DT(t,x) = \sigma'(t)\partial_x\eta(t,x) \neq 0$, $\sigma' \neq 0$, $\partial_x\eta \neq 0$, and $\tau'(\sigma(t))\sigma'(t) = 1$.

($\Rightarrow$) If $x' = f(\cdot\,, x)$ and $s \in J$, then $y'(s) = \tau'(s)G(\tau(s), x(\tau(s)))$. Since $\xi(s, y(s)) = x(\tau(s))$, $y'(s) = \widetilde{f}(s, y(s))$.
($\Leftarrow$) Let $t \in I$. Differentiating $y(\sigma(t)) = \eta(t, x(t))$ gives $\sigma'(t)y'(\sigma(t)) = \partial_t\eta(t,x(t)) + \partial_x\eta(t,x(t))x'(t)$.
Since $y' = \widetilde{f}(\cdot\,, y)$, $\sigma'(t)y'(\sigma(t)) = G(t, x(t))$. Hence $\partial_x\eta(t,x(t))[x'(t) - f(t,x(t))] = 0$, so $x' = f(\cdot\,, x)$.

*Structural Commentary*: Transform graphs; apply the chain rule.
*Key Insight*: $\widetilde{f}$ is the pushed-forward vector field. □

**Method 1.2.3.** *Homogeneous First-Order Reduction*
Let $D \subseteq \mathbb{R}$ be open, let $f \in C(D,\mathbb{R})$, and let $I \subseteq \mathbb{R} \setminus \{0\}$ be an interval. To solve $x' = f(x/t)$ on a branch for which $x(t)/t \in D$:

(i) Set $y := x/t$, so $x = ty$.
(ii) Differentiate to obtain $y' = (tx' - x)/t^2 = [f(y) - y]/t$.
(iii) Each $c \in D$ satisfying $f(c) = c$ gives the constant-ratio solution $x(t) = ct$.
(iv) On every subinterval where $f(y) \neq y$, separate $dy/(f(y) - y) = dt/t$, solve for $y$, and recover $x = ty$.

**Method 1.2.4.** *Affine-Fractional First-Order Reduction*
Let $D \subseteq \mathbb{R}$ be open, let $f \in C(D,\mathbb{R})$, put $A(t,x) := ax + bt + c$ and $B(t,x) := \alpha x + \beta t + \gamma$, and set $\Delta := a\beta - \alpha b$. Work on a solution interval on which $B(t,x(t)) \neq 0$ and $A(t,x(t))/B(t,x(t)) \in D$.

(i) If $\Delta \neq 0$, there is a unique $(t_*, x_*)$ with $A(t_*,x_*) = B(t_*,x_*) = 0$. Set $s := t - t_*$ and $y := x - x_*$.
Then $\frac{dy}{ds} = f(\frac{ay+bs}{\alpha y+\beta s})$; for $s \neq 0$, $z := y/s$ satisfies $s\frac{dz}{ds} = f(\frac{az+b}{\alpha z+\beta}) - z$.
(ii) If $\Delta = 0$ and $(a,b) \neq (0,0)$, choose the unique $\lambda \in \mathbb{R}$ with $(\alpha,\beta) = \lambda(a,b)$ and set $\widehat{f}(u) := f(\frac{u+c}{\lambda u+\gamma})$ wherever the quotient lies in $D$.
If $a \neq 0$, use $u := ax + bt$: $u' = a\widehat{f}(u) + b$, with $x = \frac{u-bt}{a}$. If $a = 0$, then $x' = \widehat{f}(bt)$.

**Example 1.2.5.** *Euler Equation Under Logarithmic Time*
For $t > 0$, set $s := \log t$ and $y(s) := x(e^s)$. Then $x'(t) = t^{-1}y'(s)$ and $x''(t) = t^{-2}(y''(s) - y'(s))$.
Hence Euler equation $t^2x'' + 3tx' + x = 2/t$ becomes constant-coefficient equation $y'' + 2y' + y = 2e^{-s}$.

**Proposition 1.2.6.** *Exact First-Order Equations*
*Let $R := I \times J$ be an open rectangle, let $p, q \in C^1(R, \mathbb{R})$, and fix $(x_0, y_0) \in R$. Then:*

*(i) If $F \in C^2(R, \mathbb{R})$ satisfies $\partial_y F = p$ and $\partial_x F = q$, then $\partial_x p = \partial_y q$ on $R$.*
*(ii) If $\partial_x p = \partial_y q$ on $R$, then $F_0(x, y) := \int_{x_0}^{x} q(s, y_0)\, ds + \int_{y_0}^{y} p(x, r)\, dr$ belongs to $C^2(R, \mathbb{R})$ and satisfies $\partial_y F_0 = p$, $\partial_x F_0 = q$.*
*(iii) Under the hypothesis of (ii), every $F_1 \in C^2(R, \mathbb{R})$ satisfying $\partial_y F_1 = p$, $\partial_x F_1 = q$ has the form $F_1 = F_0 + c$ for a unique $c \in \mathbb{R}$.*
*(iv) Under the hypothesis of (ii), for every interval $L \subseteq I$ and every $y \in C^1(L, J)$, $p(\cdot, y)y' + q(\cdot, y) = 0$ iff $F_0(\cdot, y)$ is constant on $L$.*

*Proof.*

(i) By Corollary 3.2.8, $\partial_x p = \partial_x \partial_y F = \partial_y \partial_x F = \partial_y q$.
(ii) The one-variable fundamental theorem gives $\partial_y F_0 = p$ and $\partial_x F_0 = q(x, y_0) + \int_{y_0}^{y} \partial_x p(x, r)\, dr$. By the hypothesis and the same theorem, $\partial_x F_0 = q(x, y_0) + \int_{y_0}^{y} \partial_y q(x, r)\, dr = q(x, y)$. Since these first partials are $C^1$, $F_0 \in C^2(R, \mathbb{R})$.
(iii) Set $H := F_1 - F_0$. Then $\partial_x H = \partial_y H = 0$. Applying the one-variable fundamental theorem first along the horizontal segment and then along the vertical segment gives $H(x, y) = H(x_0, y_0)$ for every $(x, y) \in R$. Thus $F_1 = F_0 + c$, and evaluation at $(x_0, y_0)$ makes $c$ unique.
(iv) The chain rule gives $(F_0(\cdot, y))' = q(\cdot, y) + p(\cdot, y)y'$. Therefore the differential equation holds exactly when this derivative vanishes, which on an interval is equivalent to constancy.

*Structural Commentary*: Build a potential; differentiate along graphs.
*Key Insight*: Mixed partials are the compatibility condition. □

**Method 1.2.7.** *Integrating Factor Depending Only on $x$*
Let $R := I \times J \subseteq \mathbb{R}^2$ be an open rectangle, let $p, q \in C^1(R, \mathbb{R})$, assume $p \neq 0$ on $R$, and fix $x_* \in I$. For $p(x, y)y' + q(x, y) = 0$:

(i) Set $r(x, y) := (\partial_y q - \partial_x p)/p$.
(ii) If $r(x, y) = r_0(x)$ on $R$, set $\mu(x) := \exp(\int_{x_*}^{x} r_0(s)\, ds)$.
(iii) Multiply by $\mu$: $\partial_x(\mu p) = \mu(r_0 p + \partial_x p) = \mu \partial_y q = \partial_y(\mu q)$, so $\mu p\, y' + \mu q = 0$ is exact by Proposition 1.2.6.

For example, on either half-plane $x > 0$ or $x < 0$, $xy' + 3x - 2y = 0$ has $r_0(x) = -3/x$, so one may take the nonvanishing factor $\mu(x) = x^{-3}$.
A potential is $F(x, y) = y/x^2 - 3/x$, and the solutions are $y = 3x + Cx^2$.

**Method 1.2.8.** *Bernoulli Linearization*
Let $I \subseteq \mathbb{R}$ be open, let $a, b \in C(I, \mathbb{R})$, and let $n \in \mathbb{R} \setminus \{0, 1\}$. On an interval $J \subseteq I$, consider a positive solution $x \in C^1(J, (0, +\infty))$ of $x' = a(t)x + b(t)x^n$.

(i) Set $y := x^{1-n}$.
(ii) Linearize: $y' = (1 - n)x^{-n}x' = (1 - n)a(t)y + (1 - n)b(t)$ on $J$.
(iii) Solve $y' = \alpha y + \beta$ with $\alpha := (1 - n)a$, $\beta := (1 - n)b$ (Proposition 1.2.1); recover $x = y^{1/(1-n)}$ on $\{y > 0\}$.

**Method 1.2.9.** *Riccati Linearization from a Particular Solution*
Let $I \subseteq \mathbb{R}$ be open, $a, b, c \in C(I, \mathbb{R})$, and let $x_p \in C^1(I, \mathbb{R})$ solve $x_p' = ax_p + bx_p^2 + c$.
Let $x \in C^1(J, \mathbb{R})$ solve $x' = ax + bx^2 + c$ on an interval $J \subseteq I$ on which $x \neq x_p$.

(i) Set $u := x - x_p$ and $y := u^{-1}$ on $J$.
(ii) Subtract and invert: $u' = (a + 2bx_p)u + bu^2$, so $y' = -(a + 2bx_p)y - b$.
(iii) Solve $y' = \alpha y + \beta$ with $\alpha := -(a + 2bx_p)$, $\beta := -b$ (Proposition 1.2.1); recover $x = x_p + y^{-1}$ on $\{y \neq 0\}$.

**Theorem 1.2.10.** *Autonomous Matrix Evolution and Variation of Constants*
*Let $A \in \mathbb{C}^{n \times n}$, $t_0 \in \mathbb{R}$, $x_0 \in \mathbb{C}^n$, and $g \in C(\mathbb{R}, \mathbb{C}^n)$. For $t \in \mathbb{R}$, set $e^{tA} := \sum_{k=0}^{\infty} \frac{t^k A^k}{k!}$. Then:*

*(i) $e^{0A} = I_n$, and $\forall s, t \in \mathbb{R}$, $e^{(t+s)A} = e^{tA}e^{sA}$, $(e^{tA})^{-1} = e^{-tA}$, and $\frac{d}{dt}e^{tA} = Ae^{tA} = e^{tA}A$.*
*(ii) $\exists!\, x \in C^1(\mathbb{R}, \mathbb{C}^n)$ with $(x' - Ax, x(t_0)) = (0, x_0)$, namely $x(t) = e^{(t-t_0)A}x_0$.*
*(iii) $\exists!\, x \in C^1(\mathbb{R}, \mathbb{C}^n)$ with $(x' - Ax, x(t_0)) = (g, x_0)$, namely $x(t) = e^{(t-t_0)A}x_0 + \int_{t_0}^{t} e^{(t-s)A}g(s)\, ds$.*

*Proof.* Fix a submultiplicative matrix norm. Since $\left\|t^k A^k/k!\right\| \leq |t|^k \|A\|^k / k!$, the defining series converges absolutely and uniformly on bounded $t$-intervals, as does its termwise derivative.

(i) The identities at $t = 0$ and for the derivative follow from the series. Absolute convergence permits the Cauchy product, and the binomial theorem gives $e^{tA}e^{sA} = \sum_{m=0}^{\infty} \frac{(t+s)^m A^m}{m!} = e^{(t+s)A}$. Taking $s = -t$ gives $e^{tA}e^{-tA} = I_n$.

(ii) Differentiation gives $x' = Ae^{(t-t_0)A}x_0 = Ax$ and $x(t_0) = x_0$.
If $y$ is another solution, then $(e^{-(t-t_0)A}y(t))' = 0$, so $y(t) = e^{(t-t_0)A}x_0$.

(iii) Put $u(t) := e^{-(t-t_0)A}x(t)$. Then $u'(t) = e^{-(t-t_0)A}g(t)$, hence $u(t) = x_0 + \int_{t_0}^{t} e^{-(s-t_0)A}g(s)\,ds$. Multiplication by $e^{(t-t_0)A}$ and (i) give the formula. Conversely, differentiation verifies the equation and initial value; the difference of two solutions is zero by (ii).

*Structural Commentary*: The inverse exponential removes the homogeneous drift.
*Key Insight*: Matrix evolution transports forcing impulses from $s$ to $t$. □

**Theorem 1.2.11.** *Principal Matrix, Liouville Formula, and Variation of Constants*
*Let $A \in C(\mathbb{R}, \mathbb{C}^{n\times n})$. Then $\exists!\, \Pi : \mathbb{R}^2 \to \mathbb{C}^{n\times n}$ such that $\forall s,t \in \mathbb{R}$, $\partial_t \Pi(t,s) = A(t)\Pi(t,s)$ and $\Pi(s,s) = I_n$. The following statements hold:*

*(i)* $\forall r,s,t \in \mathbb{R}$, $\Pi(t,r)\Pi(r,s) = \Pi(t,s)$ *and* $\Pi(t,s)^{-1} = \Pi(s,t)$.

*(ii)* $\forall s,t \in \mathbb{R}$, $\det \Pi(t,s) = \exp\left(\int_s^t \operatorname{tr} A(r)\,dr\right)$.

*(iii) If $g \in C(\mathbb{R}, \mathbb{C}^n)$ and $x_s \in \mathbb{C}^n$, then $\exists! x \in C^1(\mathbb{R}, \mathbb{C}^n)$ with $(x' - Ax, x(s)) = (g, x_s)$, namely $x(t) = \Pi(t,s)x_s + \int_s^t \Pi(t,r)g(r)\,dr$ for $t \in \mathbb{R}$.*

*The matrix $\Pi(t,s)$ is the principal matrix solution from $s$; any matrix solution $X' = AX$ with $\det X(s) \neq 0$ is a fundamental matrix solution.*

*Proof.* Identify $\mathbb{C}^n$ with $\mathbb{R}^{2n}$. For the canonical basis $e_1, \dots, e_n$, Theorem 1.1.27 and Corollary 1.1.30 give a unique global solution of $x' = A(t)x$, $x(s) = e_j$; use these solutions as the columns of $\Pi(t,s)$. Existence and uniqueness of $\Pi$ hold columnwise.

(i) For fixed $r,s$, both $t \mapsto \Pi(t,r)\Pi(r,s)$ and $t \mapsto \Pi(t,s)$ solve $X' = A(t)X$ and agree at $t = r$; hence they coincide. Applying the cocycle law to the triples $(s,t,s)$ and $(t,s,t)$ gives $I_n = \Pi(s,t)\Pi(t,s) = \Pi(t,s)\Pi(s,t)$, so $\Pi(t,s)^{-1} = \Pi(s,t)$. If $X(t) := \Pi(t,0)$, then $\Pi(t,s) = X(t)X(s)^{-1}$; hence $\Pi$ is jointly continuous.

(ii) Let $X(t) := \Pi(t,s)$ and write its columns as $x_1(t), \dots, x_n(t)$. The Leibniz formula and $x_j' = Ax_j$ give $\frac{d}{dt}\det X(t) = \sum_{j=1}^{n} \det(x_1, \dots, Ax_j, \dots, x_n) = \operatorname{tr}(A(t)) \det X(t)$. Multiplying by $\exp(-\int_s^t \operatorname{tr} A(r)\,dr)$ and using $\det X(s) = 1$ gives the claimed formula.

(iii) Differentiating the asserted formula gives $x' = A(t)x + g(t)$ and $x(s) = x_s$. If $x, y$ have the same data, then $z := x - y$ solves $z' = A(t)z$, $z(s) = 0$, so the defining property of $\Pi$ gives $z(t) = \Pi(t,s)z(s) = 0$.

*Structural Commentary*: Columnwise existence builds $\Pi$; uniqueness forces its cocycle law.
*Key Insight*: $\Pi(t,s)$ transports states and forcing between arbitrary times. □

**Method 1.2.12.** *Solving $\mathbf{x}'(t) = A\mathbf{x}(t)$ by Jordan Form*
Let $A \in \mathbb{C}^{n\times n}$ and $\mathbf{x}_0 \in \mathbb{C}^n$. To solve $\mathbf{x}'(t) = A\mathbf{x}(t)$ with $\mathbf{x}(0) = \mathbf{x}_0$:

(i) By Theorem 6.2.1, choose $S \in GL_n(\mathbb{C})$ with $A = SJS^{-1}$, where $J = \bigoplus_{i=1}^{r} J_{k_i}(\lambda_i)$.

(ii) Set $\mathbf{y}(t) = S^{-1}\mathbf{x}(t)$ and $\mathbf{y}_0 = S^{-1}\mathbf{x}_0$. Then $\mathbf{y}'(t) = J\mathbf{y}(t)$ and $\mathbf{x}(t) = S\mathbf{y}(t)$.

(iii) Split $\mathbf{y} = (\mathbf{y}_1, \dots, \mathbf{y}_r)$ according to the Jordan blocks. Then $\mathbf{y}_i'(t) = J_{k_i}(\lambda_i)\mathbf{y}_i(t)$.

(iv) By Proposition 6.5.1, $\mathbf{y}_i(t) = e^{tJ_{k_i}(\lambda_i)}\mathbf{y}_i(0) = e^{\lambda_i t}\sum_{q=0}^{k_i-1} \frac{t^q}{q!} J_{k_i}(0)^q \mathbf{y}_i(0)$ for $1 \le i \le r$.
Differentiating verifies the block equation.

(v) Reassemble $\mathbf{y}(t)$ and transform back: $\mathbf{x}(t) = S\mathbf{y}(t)$.

Thus a size-$k$ block with eigenvalue $\lambda$ contributes entries of the form $e^{\lambda t}p(t)$, where $\deg p < k$.

**Perspective 1.2.13.** *Jordan Functions and Polynomial Factors*
For a Jordan block $J_k(\lambda) = \lambda I_k + N$ with $N^k = 0$, any polynomial $f$ satisfies $f(J_k(\lambda)) = \sum_{r=0}^{k-1} \frac{f^{(r)}(\lambda)}{r!} N^r$.
The nilpotent part contributes derivative terms; for the exponential these become polynomial factors in $t$ by Proposition 6.5.1. As $t \to +\infty$, $\operatorname{Re}\lambda < 0$ gives blockwise decay and $\operatorname{Re}\lambda > 0$ gives growth.
If $\operatorname{Re}\lambda = 0$, the block evolution is bounded iff $N = 0$.

**Example 1.2.14.** *Nilpotent Jordan Block Evolution*
Let $N := J_3(0) = \left(\begin{smallmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{smallmatrix}\right)$, so $N^3 = 0$. By Proposition 6.5.1, $e^{tN} = I_3 + tN + \frac{t^2}{2}N^2 = \left(\begin{smallmatrix} 1 & t & \frac{t^2}{2} \\ 0 & 1 & t \\ 0 & 0 & 1 \end{smallmatrix}\right)$. Nilpotent Jordan blocks create polynomial, rather than exponential, growth along generalized-eigenvector chains.

**Theorem 1.2.15.** *Spectral Criteria for Autonomous Linear Stability*
*Let $A \in \mathbb{C}^{n\times n}$ and set $\alpha(A) := \max\{\operatorname{Re}\lambda : \lambda \in \operatorname{spec}(A)\}$. Then:*

*(i)* $e^{tA} \to 0$ *as* $t \to +\infty$ *iff* $\alpha(A) < 0$.
*(ii)* $\sup_{t\geq 0} \left\|e^{tA}\right\| < +\infty$ *iff* $\alpha(A) \leq 0$ *and* $\forall \lambda \in \operatorname{spec}(A)$, $\operatorname{Re}\lambda = 0 \implies \dim\ker(A - \lambda I_n) = m_A(\lambda)$.
*(iii) If* $\alpha(A) < 0$, *then* $\forall \gamma \in (0, -\alpha(A))$, $\exists C_\gamma > 0$ *such that* $\left\|e^{tA}\right\| \leq C_\gamma e^{-\gamma t}$ $\forall t \geq 0$.

*Proof.* Choose $S \in GL_n(\mathbb{C})$ with $A = SJS^{-1}$ and use Proposition 6.5.1 on every block $J_k(\lambda)$. Fixed similarity changes all norms by at most the factor $\|S\| \left\|S^{-1}\right\|$.

(i) Each block exponential is $e^{\lambda t} \sum_{r=0}^{k-1} \frac{t^r N^r}{r!}$. It tends to zero exactly when $\operatorname{Re}\lambda < 0$; if $\operatorname{Re}\lambda \geq 0$, its diagonal entry $e^{\lambda t}$ does not tend to zero.
(ii) If $\operatorname{Re}\lambda < 0$, every polynomial factor is dominated by $e^{\operatorname{Re}\lambda t}$ and the block is bounded. If $\operatorname{Re}\lambda = 0$, the block is bounded exactly when $N = 0$, equivalently $k = 1$. If $\operatorname{Re}\lambda > 0$, its diagonal is unbounded.
(iii) Fix $0 < \gamma < -\alpha(A)$. For a block of size $k$, $e^{\gamma t}\left\|e^{tJ_k(\lambda)}\right\|$ is bounded because every $t^r e^{(\operatorname{Re}\lambda+\gamma)t}$ tends to zero. Take the maximum block bound and include the fixed similarity factor.

*Structural Commentary*: Jordan blocks separate exponential rates from nilpotent polynomial factors.
*Key Insight*: Boundary-spectrum nilpotence is the sole obstruction to boundedness. □

**Definition 1.2.16.** *Hurwitz Matrix*
A matrix $A \in \mathbb{C}^{n\times n}$ is *Hurwitz* iff $\max\{\operatorname{Re}\lambda : \lambda \in \operatorname{spec}(A)\} < 0$.

**Perspective 1.2.17.** *Planar Linear Phase Classification*
For $x' = Ax$ with $A \in \mathbb{R}^{2\times 2}$, two real eigenvalues of negative sign give a sink, two of positive sign give a source, and opposite signs give a saddle.
A nonreal pair $\alpha \pm i\beta$, $\beta \neq 0$, gives a spiral sink when $\alpha < 0$, a spiral source when $\alpha > 0$, and a center when $\alpha = 0$. A defective repeated real eigenvalue $\lambda$ adds a polynomial shear along its generalized eigenvector; it still gives decay for $\lambda < 0$ and growth for $\lambda > 0$. For $\lambda = 0$, $e^{tA} = I_2 + tA$ in the defective case, so generalized-eigenvector directions grow linearly while the eigenspace consists of equilibria.
These labels describe the origin; similarity changes the coordinates of the phase portrait without changing its spectral type.

**Theorem 1.2.18.** *Basis for Constant-Coefficient Linear ODEs*
*Let* $p(z) := z^n + c_{n-1}z^{n-1} + \cdots + c_1 z + c_0 = \prod_{j=1}^{m} (z - \lambda_j)^{a_j}$ *over* $\mathbb{C}$, *where the* $\lambda_j$ *are distinct and* $\sum_{j=1}^{m} a_j = n$.
*Then:*

*(i) The functions* $x_{j,k}(t) := t^k e^{\lambda_j t}$, $1 \leq j \leq m$, $0 \leq k < a_j$, *form a basis of the complex solutions of* $p(D)x = 0$, *where* $D = d/dt$.
*(ii) If* $c_0, \ldots, c_{n-1} \in \mathbb{R}$, *write* $\lambda_j = \alpha_j + i\beta_j$. *The functions* $t^k e^{\lambda_j t}$ *for the real roots* $\beta_j = 0$, *together with* $t^k e^{\alpha_j t}\cos(\beta_j t)$ *and* $t^k e^{\alpha_j t}\sin(\beta_j t)$ *for one root from each conjugate pair with* $\beta_j > 0$, *form a real basis, where always* $0 \leq k < a_j$.

*Proof.* For $q \in \mathbb{C}[t]$, $(D - \lambda)(e^{\lambda t}q) = e^{\lambda t}q'$. Thus $(D - \lambda_j)^{a_j} x_{j,k} = 0$ for $k < a_j$, so every listed function solves $p(D)x = 0$.

(i) Suppose $\sum_{j=1}^{m} e^{\lambda_j t} q_j(t) = 0$ with $\deg q_j < a_j$. Fix $j$ and apply $\prod_{i\neq j} (D - \lambda_i)^{a_i}$. All terms except the $j$th vanish, while the remaining term is $e^{\lambda_j t} R_j(D) q_j$, where $R_j(z) := \prod_{i\neq j} (z + \lambda_j - \lambda_i)^{a_i}$ and $R_j(0) \neq 0$.
On polynomials of degree below $a_j$, $R_j(D) - R_j(0)I$ is nilpotent because it is divisible by $D$. Thus $R_j(D)$ is invertible, so $q_j = 0$. The $n$ listed functions are linearly independent.
Writing $y := (x, x', \ldots, x^{(n-1)})$ converts $p(D)x = 0$ into a first-order companion system. Theorem 1.2.11(iii), with zero forcing, makes $y(0) \mapsto y$ a complex-linear bijection from $\mathbb{C}^n$ onto the global solution space. Thus that space has complex dimension $n$ and the list is a basis.
(ii) Real coefficients pair nonreal roots by conjugation. Taking real and imaginary parts of the paired complex basis functions gives the stated real solutions; the change from each conjugate pair to its real and imaginary parts is invertible over $\mathbb{R}$.

*Structural Commentary*: Factor $p(D)$; isolate each exponential-polynomial root space.
*Key Insight*: Root multiplicity becomes polynomial degree. □

**Method 1.2.19.** *Exponential-Polynomial Forcing*
Let $p \in \mathbb{C}[z] \setminus \{0\}$, $q \in \mathbb{C}[t] \setminus \{0\}$, and $\beta \in \mathbb{C}$, and consider $p(D)x = q(t)e^{\beta t}$. Write $p(z) = (z-\beta)^a h(z)$ with $a \in \mathbb{N}_0$ and $h(\beta) \neq 0$. Set $d := \deg q$ and $\mathbb{C}_{\leq d}[t] := \{u \in \mathbb{C}[t] : \deg u \leq d\}$.

(i) Solve $h(D+\beta)u = q$ in $\mathbb{C}_{\leq d}[t]$; the operator is triangular with nonzero diagonal $h(\beta)$ and hence invertible.
(ii) Choose the polynomial $r$ with $r^{(a)} = u$ and vanishing coefficients in degrees $0, \dots, a-1$.
(iii) Set $x_p(t) := e^{\beta t} r(t)$. Then $p(D)x_p = e^{\beta t} h(D+\beta) D^a r = q(t)e^{\beta t}$ and $\deg r \leq \deg q + a$.

The factor $t^a$ in $r$ is the resonance correction when $\beta$ is a root of $p$.

**Example 1.2.20.** *Damping and Resonance*
For $x'' + 2\eta x' + \omega_0^2 x = 0$, with $\eta, \omega_0 > 0$, the roots are $-\eta \pm \sqrt{\eta^2 - \omega_0^2}$. Thus $\eta > \omega_0$, $\eta = \omega_0$, and $\eta < \omega_0$ give over-damping, critical damping, and under-damping, respectively; homogeneous solutions decay.
For the forced equation $x'' + 2\eta x' + \omega_0^2 x = \cos(\omega t)$, a complex particular solution is $z_p(t) = e^{i\omega t}/(\omega_0^2 - \omega^2 + 2i\eta\omega)$, and $x_p = \operatorname{Re} z_p$. If $\eta = 0$ and $\omega = \omega_0$, Method 1.2.19 instead gives $x_p(t) = t\sin(\omega_0 t)/(2\omega_0)$, whose amplitude grows linearly.
For a series RLC circuit, $\eta = R/(2L)$ and $\omega_0 = 1/\sqrt{LC}$; the same three regimes and resonance mechanism govern the current.

**Method 1.2.21.** *Reduction of Order for a Second Solution*
Let $I \subseteq \mathbb{R}$ be open, let $q_0, q_1 \in C(I, \mathbb{C})$, let $\phi_1 \in C^2(I, \mathbb{C})$ solve $x'' + q_1 x' + q_0 x = 0$, and restrict to an interval $J \subseteq I$ on which $\phi_1 \neq 0$. Fix $t_* \in J$.

(i) Set $x = c\phi_1$. Cancellation of the equation for $\phi_1$ gives $\phi_1 c'' + (2\phi_1' + q_1\phi_1)c' = 0$.
(ii) With $d := c'$, solve $d' + (2\phi_1'/\phi_1 + q_1)d = 0$ to obtain $d(t) = C\phi_1(t)^{-2}\exp(-\int_{t_*}^t q_1(r)\,dr)$.
(iii) Taking $C = 1$ gives $\phi_2(t) := \phi_1(t)\int_{t_*}^t \phi_1(s)^{-2}\exp(-\int_{t_*}^s q_1(r)\,dr)\,ds$. Then $W(\phi_1,\phi_2)(t) = \phi_1\phi_2' - \phi_1'\phi_2 = \exp(-\int_{t_*}^t q_1(r)\,dr) \neq 0$. Thus $\phi_1, \phi_2$ are linearly independent; uniqueness for the associated two-dimensional first-order system makes them a basis of the solution space on $J$.

**Theorem 1.2.22.** *Floquet Normal Form*
*Let $A \in C(\mathbb{R}, \mathbb{C}^{n\times n})$ satisfy $A(t+T) = A(t)$ for some $T > 0$, let $\Pi$ be its principal matrix solution, and set $M(t_0) := \Pi(t_0 + T, t_0)$. Then:*

*(i)* $\forall s, t \in \mathbb{R}$, $\Pi(t+T, s+T) = \Pi(t,s)$.
*(ii)* $M(t_0)$ *is invertible.*
*(iii)* $\exists Q(t_0) \in \mathbb{C}^{n\times n}$ *and a $T$-periodic* $P(\,\cdot\,, t_0) \in C^1(\mathbb{R}, GL_n(\mathbb{C}))$ *such that* $\Pi(t, t_0) = P(t, t_0)e^{(t-t_0)Q(t_0)}$ *and* $P(t_0, t_0) = I_n$.
*(iv)* $\forall t_0, t_1 \in \mathbb{R}$, $M(t_1) = \Pi(t_1, t_0)M(t_0)\Pi(t_1, t_0)^{-1}$*; consequently, the spectrum and Jordan block sizes of $M(t_0)$ are independent of $t_0$.*

*Proof.*

(i) Both $t \mapsto \Pi(t+T, s+T)$ and $t \mapsto \Pi(t,s)$ solve $X' = A(t)X$ and equal $I_n$ at $t = s$, so uniqueness gives the identity.
(ii) Theorem 1.2.11(i) gives $M(t_0)^{-1} = \Pi(t_0, t_0 + T)$.
(iii) By (ii), every Jordan block of $M(t_0)$ has the form $J = \rho I + N$ with $\rho \neq 0$ and $N^k = 0$. Choose $\ell \in \mathbb{C}$ with $e^\ell = \rho$ and set $L := \ell I + \sum_{q=1}^{k-1} \frac{(-1)^{q+1}}{q}(\rho^{-1}N)^q$. In the quotient algebra $\mathbb{C}[z]/(z^k)$, the finite logarithm and exponential series satisfy $\exp(\log(1+z)) = 1+z$; substituting $z = \rho^{-1}N$ gives $e^L = J$. Reassembling and conjugating these block logarithms produces $R \in \mathbb{C}^{n\times n}$ with $e^R = M(t_0)$.
Set $Q(t_0) := T^{-1}R$ and $P(t, t_0) := \Pi(t, t_0)e^{-(t-t_0)Q(t_0)}$. Then $P(t_0, t_0) = I_n$. By (i) and the cocycle law, $\Pi(t+T, t_0) = \Pi(t, t_0)M(t_0)$; because $M(t_0) = e^{TQ(t_0)}$, this identity gives $P(t+T, t_0) = P(t, t_0)$. Both factors defining $P$ are $C^1$ and invertible.
(iv) Periodicity from (i) and the cocycle law give $\Pi(t_1 + T, t_0) = \Pi(t_1, t_0)M(t_0)$ and also $\Pi(t_1 + T, t_0) = M(t_1)\Pi(t_1, t_0)$. Right multiplication by $\Pi(t_1, t_0)^{-1}$ proves the similarity identity, and similarity preserves the spectrum and Jordan block sizes.

*Structural Commentary*: Take a logarithm of one-period transport; factor the remainder.
*Key Insight*: Periodic evolution is constant exponential growth in periodic coordinates. □

**Definition 1.2.23.** *Floquet Multipliers and Exponents*
Let $A \in C(\mathbb{R}, \mathbb{C}^{n\times n})$ be $T$-periodic, let $\Pi$ be its principal matrix solution, set $M(t_0) := \Pi(t_0 + T, t_0)$, and choose $Q(t_0)$ with $e^{TQ(t_0)} = M(t_0)$.

(i) The eigenvalues $\rho$ of $M(t_0)$ are the *Floquet multipliers*.

(ii) The eigenvalues $\gamma$ of a Floquet matrix $Q(t_0)$ are *Floquet exponents*. Spectral mapping gives $\operatorname{spec}(M(t_0)) = \{e^{T\gamma} : \gamma \in \operatorname{spec}(Q(t_0))\}$ and $m_{M(t_0)}(\rho) = \sum_{\substack{\gamma \in \operatorname{spec}(Q(t_0)) \\ e^{T\gamma}=\rho}} m_{Q(t_0)}(\gamma)$. Thus every exponent determines a multiplier by $\rho = e^{T\gamma}$, and the scalar logarithm determines $\gamma$ only modulo $2\pi i/T$.

**Corollary 1.2.24.** *Floquet Multiplier Stability Criterion*
*Let $A \in C(\mathbb{R}, \mathbb{C}^{n\times n})$ be $T$-periodic, let $\Pi$ be its principal matrix solution, and set $M(t_0) := \Pi(t_0+T, t_0)$. Then:*

*(i)* $\sup_{t\ge t_0} \|\Pi(t, t_0)\| < +\infty$ *iff* $|\rho| \le 1$ *for* $\rho \in \operatorname{spec}(M(t_0))$ *and* $\dim\ker(M(t_0) - \rho I_n) = m_{M(t_0)}(\rho)$ *whenever* $|\rho| = 1$.

*(ii)* $\Pi(t, t_0) \to 0$ *as* $t \to +\infty$ *iff* $\max\{|\rho| : \rho \in \operatorname{spec}(M(t_0))\} < 1$.

*Proof.* Write $t = t_0 + kT + \tau$, $k \in \mathbb{N}_0$, $0 \le \tau < T$. Periodicity and the cocycle law give by induction $\Pi(t, t_0) = \Pi(t_0+\tau, t_0)M(t_0)^k$. The first factor and its inverse are uniformly bounded for $0 \le \tau \le T$.

(i) Hence $\Pi(\,\cdot\,, t_0)$ is bounded iff $(M(t_0)^k)$ is bounded. By Theorem 3.3.7, this is equivalent to the stated multiplier and multiplicity conditions.

(ii) Likewise, $\Pi(t, t_0) \to 0$ iff $M(t_0)^k \to 0$. By Theorem 3.3.5, this is equivalent to $\rho(M(t_0)) < 1$.

*Structural Commentary*: Sample at integer periods; reduce continuous evolution to powers of $M$.
*Key Insight*: Floquet stability is matrix-power stability of one-period transport. □

**Example 1.2.25.** *Hill Equation and Its Floquet Discriminant*
Let $q \in C(\mathbb{R}, \mathbb{R})$ be $T$-periodic and consider $x'' + q(t)x = 0$. Its first-order matrix is $A(t) = \left(\begin{smallmatrix} 0 & 1 \\ -q(t) & 0 \end{smallmatrix}\right)$. Let $\Pi$ be the principal matrix solution and $M(t_0) := \Pi(t_0+T, t_0)$. Since $\operatorname{tr} A(t) = 0$, by Theorem 1.2.11(ii), $\det M(t_0) = 1$.
With $\Delta := \frac{1}{2}\operatorname{tr} M(t_0)$, the multiplier equation is $\rho^2 - 2\Delta\rho + 1 = 0$. If $|\Delta| < 1$, the multipliers are distinct conjugates on the unit circle and all solutions are bounded. If $|\Delta| > 1$, one real multiplier has modulus above 1 and an unbounded solution exists. At $|\Delta| = 1$, boundedness holds exactly when $M(t_0) = \Delta I_2$; otherwise its nontrivial Jordan block produces linear growth in the period count.
The Mathieu equation $x'' + \omega^2(1 + \varepsilon\cos t)x = 0$ is the standard periodic-coefficient example; its stability regions are determined by $\Delta(\omega, \varepsilon)$.

**Theorem 1.2.26.** *Perturbation Bounds for Linear Evolution*
*Let $A, B \in C(\mathbb{R}, \mathbb{C}^{n\times n})$, let $\Pi_A, \Pi_{A+B}$ be the corresponding principal matrix solutions, and fix $t_0 \in \mathbb{R}$. Then:*

*(i) If* $\|\Pi_A(t, s)\| \le Ce^{-\alpha(t-s)}$ *for all* $t \ge s \ge t_0$ *and* $\|B(t)\| \le b$ *for all* $t \ge t_0$, *where* $C, \alpha > 0$, $b \ge 0$, *and* $Cb < \alpha$, *then* $\|\Pi_{A+B}(t, s)\| \le Ce^{-(\alpha - Cb)(t-s)}$ *for all* $t \ge s \ge t_0$.

*(ii) If* $\|\Pi_A(t, s)\| \le C$ $\forall t \ge s \ge t_0$ *and* $B_1 := \int_{t_0}^{\infty} \|B(r)\|\, dr < +\infty$, *then* $\|\Pi_{A+B}(t, s)\| \le Ce^{CB_1}$ $\forall t \ge s \ge t_0$.

*Proof.* For fixed $s \ge t_0$, by Theorem 1.2.11(iii), applied columnwise with forcing $B(t)x(t)$, $\Pi_{A+B}(t, s) = \Pi_A(t, s) + \int_s^t \Pi_A(t, r)B(r)\Pi_{A+B}(r, s)\, dr$.

(i) Put $y(t) := e^{\alpha(t-s)}\|\Pi_{A+B}(t, s)\|$. The hypotheses give $y(t) \le C + Cb\int_s^t y(r)\, dr$. By Lemma 1.1.20(ii), after translating $s$ to 0, $y(t) \le Ce^{Cb(t-s)}$, which is the asserted estimate.

(ii) Put $y(t) := \|\Pi_{A+B}(t, s)\|$. Then $y(t) \le C + C\int_s^t \|B(r)\|\, y(r)\, dr$. The same Gronwall estimate gives $y(t) \le C\exp(C\int_s^t \|B(r)\|\, dr) \le Ce^{CB_1}$.

*Structural Commentary*: Variation of constants converts perturbation into Gronwall feedback.
*Key Insight*: Perturbation mass subtracts from decay or multiplies boundedness. □

**Corollary 1.2.27.** *Vanishing Perturbations of a Hurwitz Matrix*
*Let $A \in \mathbb{C}^{n\times n}$ be Hurwitz, let $B \in C(\mathbb{R}, \mathbb{C}^{n\times n})$, and suppose $\|B(t)\| \to 0$ as $t \to +\infty$. Set $\delta_A := \min\{-\operatorname{Re}\lambda : \lambda \in \operatorname{spec}(A)\}$. Then $\forall \gamma \in (0, \delta_A)$, $\exists K_\gamma > 0$ such that $\|\Pi_{A+B}(t, 0)\| \le K_\gamma e^{-\gamma t}$ $\forall t \ge 0$.*

*Proof.* Fix $0 < \gamma < \eta < \delta_A$. By Theorem 1.2.15(iii), $\exists C_\eta > 0$ with $\left\|e^{(t-s)A}\right\| \le C_\eta e^{-\eta(t-s)}$ for $t \ge s$.
Choose $t_0 \ge 0$ such that $\|B(t)\| \le (\eta - \gamma)/C_\eta$ for $t \ge t_0$. By Theorem 1.2.26(i), $\|\Pi_{A+B}(t, t_0)\| \le C_\eta e^{-\gamma(t-t_0)}$ for $t \ge t_0$.
The cocycle law yields $\Pi_{A+B}(t, 0) = \Pi_{A+B}(t, t_0)\Pi_{A+B}(t_0, 0)$.
Set $K_1 := C_\eta e^{\gamma t_0}\|\Pi_{A+B}(t_0, 0)\|$; then $\|\Pi_{A+B}(t, 0)\| \le K_1 e^{-\gamma t}$ for $t \ge t_0$.
Continuity gives $K_0 := \max_{0\le t\le t_0} e^{\gamma t}\|\Pi_{A+B}(t, 0)\| < +\infty$. Taking $K_\gamma := \max\{K_0, K_1\}$ proves the claim.
*Structural Commentary*: Wait until $B$ is smaller than the spectral decay margin.
*Key Insight*: A vanishing perturbation eventually preserves any smaller decay rate. □

## 1.3 Nonlinear Stability Theory

Exact formulas are often unnecessary for proving that scalar trajectories remain ordered or trapped. Differential inequalities provide comparison curves for this purpose.

**Definition 1.3.1.** *Upper and Lower Solutions*
Let $U \subseteq \mathbb{R}^2$ be open, $f : U \to \mathbb{R}$, and $J \subseteq \mathbb{R}$ an open interval. For $u \in C^1(J, \mathbb{R})$ with $\Gamma_u := \{(t, u(t)) : t \in J\} \subseteq U$:

(i) *Upper solution*: $u'(t) \geq f(t, u(t))\ \forall t \in J$; it is *strict* iff $u'(t) > f(t, u(t))\ \forall t \in J$.
(ii) *Lower solution*: $u'(t) \leq f(t, u(t))\ \forall t \in J$; it is *strict* iff $u'(t) < f(t, u(t))\ \forall t \in J$.

*Key Insight*: Differential inequalities encode one-sided barriers without solving the ODE.

**Definition 1.3.2.** *Scalar Isocline*
Let $U \subseteq \mathbb{R}^2$ be open, $f : U \to \mathbb{R}$, and $c \in \mathbb{R}$.
For $x' = f(t, x)$, the set $\mathcal{I}_c := \{(t, x) \in U : f(t, x) = c\}$ is the *isocline of slope c*.
Let $J \subseteq \mathbb{R}$ be open and $w \in C^1(J, \mathbb{R})$ satisfy $\Gamma_w := \{(t, w(t)) : t \in J\} \subseteq \mathcal{I}_c$.

(i) If $w'(t) > c\ \forall t \in J$, then $w$ is a strict upper solution.
(ii) If $w'(t) < c\ \forall t \in J$, then $w$ is a strict lower solution.

*Key Insight*: An isocline fixes the ODE slope, so only the graph slope remains to be compared.

**Lemma 1.3.3.** *Strict Scalar Comparison*
*Let $U \subseteq \mathbb{R}^2$ be open, $f \in C(U, \mathbb{R})$, $J \subseteq \mathbb{R}$ open, and $[t_0, T] \subseteq J$ with $t_0 < T$.*
*For $z \in C^1(J, \mathbb{R})$, set $\Gamma_z := \{(t, z(t)) : t \in J\}$, and let $I_+ := (t_0, T]$. Suppose $x \in C^1(J, \mathbb{R})$, $\Gamma_x \subseteq U$, and $x' = f(\cdot, x)$.*

*(i) If $u \in C^1(J, \mathbb{R})$, $\Gamma_u \subseteq U$, $u'(t) > f(t, u(t))\ \forall t \in J$, and $x(t_0) \leq u(t_0)$, then $x(t) < u(t)\ \forall t \in I_+$.*
*(ii) If $v \in C^1(J, \mathbb{R})$, $\Gamma_v \subseteq U$, $v'(t) < f(t, v(t))\ \forall t \in J$, and $v(t_0) \leq x(t_0)$, then $v(t) < x(t)\ \forall t \in I_+$.*

*Proof.*

(i) Set $\Delta := u - x$. At every $t \in [t_0, T]$ with $\Delta(t) = 0$, one has $x(t) = u(t)$ and $\Delta'(t) = u'(t) - f(t, u(t)) > 0$. Thus $\Delta(t_0) \geq 0$ implies $\Delta > 0$ on $(t_0, t_0 + \varepsilon]$ for some $\varepsilon > 0$.
If $Z := \{t \in (t_0, T] : \Delta(t) = 0\} \neq \emptyset$, let $t_1 := \inf Z$. Continuity gives $\Delta(t_1) = 0$ and $\Delta > 0$ on $(t_0, t_1)$, so the left derivative satisfies $\Delta'(t_1) \leq 0$. This contradicts $\Delta'(t_1) > 0$. Hence $Z = \emptyset$ and $\Delta > 0$ on $I_+$.
(ii) Set $\Delta := x - v$. At every zero of $\Delta$, $\Delta' = f(\cdot, v) - v' > 0$. The argument from (i), with $x - v$ in place of $u - x$, gives $\Delta > 0$ on $I_+$.

*Structural Commentary*: A first contact forces $\Delta' \leq 0$, contradicting the strict residual sign.
*Key Insight*: Strict differential inequalities forbid barrier crossing. □

**Definition 1.3.4.** *Local State Lipschitz Condition*
Let $U \subseteq \mathbb{R}^2$ be open and $f : U \to \mathbb{R}$. Then $f$ is *locally Lipschitz in the state, uniformly in time* iff, for every compact $K \subseteq U$, there exists $L_K < +\infty$ such that $|f(t, x) - f(t, y)| \leq L_K |x - y|$ whenever $(t, x), (t, y) \in K$.

**Theorem 1.3.5.** *Scalar Lipschitz Comparison*
*Let $U \subseteq \mathbb{R}^2$ be open and $f \in C(U, \mathbb{R})$. Assume that for every compact $K \subseteq U$, there exists $L_K < +\infty$ such that $|f(t, p) - f(t, q)| \leq L_K |p - q|$ whenever $(t, p), (t, q) \in K$.*
*Let $I := [t_0, T] \subseteq J$, $J \subseteq \mathbb{R}$ open, $t_0 < T$. For $z : J \to \mathbb{R}$, set $\Gamma_z(A) := \{(t, z(t)) : t \in A\}$, $R_z := z' - f(\cdot, z)$.*
*Suppose $x, y \in C^1(J, \mathbb{R})$, $\Gamma_x(J) \cup \Gamma_y(J) \subseteq U$, and $R_x \leq R_y$ on $I$. Then:*

*(i) If $x(t_0) \leq y(t_0)$, then $x \leq y$ on $I$.*
*(ii) If $t_1 \in I$ and $x(t_1) < y(t_1)$, then $x < y$ on $[t_1, T]$.*

*Proof.*

(i) Set $\Delta := x - y$ and suppose $S := \{t \in I : \Delta(t) > 0\} \neq \emptyset$. Since $\Delta(t_0) \leq 0$, every component $C$ of $S$ has a left endpoint $\alpha \in I$ with $\Delta(\alpha) = 0$. Choose $t_2 \in C$, so $\Delta > 0$ on $(\alpha, t_2]$.
With $K := \Gamma_x([\alpha, t_2]) \cup \Gamma_y([\alpha, t_2])$ and $L = L_K$, $R_x \leq R_y$ gives $\Delta' \leq f(\cdot, x) - f(\cdot, y) \leq L\Delta$ on $(\alpha, t_2]$.
Thus $(e^{-L(t-\alpha)}\Delta)' \leq 0$, so $\Delta(t_2) \leq e^{L(t_2-\alpha)}\Delta(\alpha) = 0$, contradicting $t_2 \in S$.
(ii) Set $\delta := y - x$. If $Z := \{t \in (t_1, T] : \delta(t) = 0\} \neq \emptyset$, let $t_2 := \inf Z$. Then $\delta > 0$ on $[t_1, t_2)$ and $\delta(t_2) = 0$.
With $K := \Gamma_x([t_1, t_2]) \cup \Gamma_y([t_1, t_2])$ and $L = L_K$, by $R_x \leq R_y$, $\delta' \geq f(\cdot, y) - f(\cdot, x) \geq -L\delta$ on $[t_1, t_2)$.
Hence $(e^{L(t-t_1)}\delta)' \geq 0$ and $0 = e^{L(t_2-t_1)}\delta(t_2) \geq \delta(t_1) > 0$, contradiction.

*Structural Commentary*: On a positive component, the integrating factor makes the order defect nonincreasing.
*Key Insight*: A nonpositive order defect cannot become positive. □

**Lemma 1.3.6.** *A Convergent Autonomous State Is an Equilibrium*
*Let $f \in C(\mathbb{R},\mathbb{R})$ and let $x \in C^1([t_0,+\infty),\mathbb{R})$ satisfy $x' = f(x)$. If $x(t) \to x_* \in \mathbb{R}$ as $t \to +\infty$, then:*

*(i)* $x'(t) \to 0$.
*(ii)* $f(x_*) = 0$.

*Proof.*

(i) By continuity, $x'(t) = f(x(t)) \to f(x_*) := L$. If $L > 0$, then $x'(t) \geq L/2$ eventually, so integration forces $x(t) \to +\infty$. If $L < 0$, the analogous estimate forces $x(t) \to -\infty$. Both conclusions contradict $x(t) \to x_*$, so $L = 0$ and $x'(t) \to 0$.
(ii) The identity $L = f(x_*)$ from (i) now gives $f(x_*) = 0$.

*Structural Commentary*: The differential equation converts state convergence into derivative convergence.
*Key Insight*: A nonzero limiting velocity is incompatible with a finite limiting state. □

**Example 1.3.7.** *A Trapping Wedge for $x' = x^2 - t^2$*
For $f(t,x) := x^2 - t^2$ on $(0,+\infty) \times \mathbb{R}$, set $u(t) := t$ and $v(t) := -t$.
Then $u' = 1 > 0 = f(t,u)$ and $v' = -1 < 0 = f(t,v)$.
If $t_0 > 0$, $x_0 \in [-t_0,t_0]$, and $x \in C^1([t_0,T_+),\mathbb{R})$ satisfies $x' = x^2 - t^2$ and $x(t_0) = x_0$, then Lemma 1.3.3 gives $-t < x(t) < t \ \forall t \in (t_0,T_+)$.

**Proposition 1.3.8.** *Central-Wedge Convergence for $x' = x^2 - t^2$*
*Let $t_0 > 0$, $-t_0 < x_0 < t_0$, and let $x : [t_0,T_+) \to \mathbb{R}$ be the maximal forward solution of $x' = x^2 - t^2$, $x(t_0) = x_0$.*

*(i)* $-t < x(t) < t$ *for* $t \in (t_0,T_+)$.
*(ii)* $T_+ = +\infty$.
*(iii)* $x(t) + t \to 0$ *as* $t \to +\infty$.

*Proof.*

(i) Let $u(t) := t$ and $v(t) := -t$. Since $u' - (u^2 - t^2) = 1 > 0$, $v' - (v^2 - t^2) = -1 < 0$, and $v(t_0) < x_0 < u(t_0)$, Lemma 1.3.3 implies $v(t) < x(t) < u(t)$ on every compact subinterval of $[t_0,T_+)$.
(ii) By (i), $x' = (x - t)(x + t) < 0$, so $x$ is decreasing.
If $T_+ < +\infty$, then $-T_+ < x(t) \leq x_0$ for $t < T_+$, hence $\ell := \lim_{t \uparrow T_+} x(t) \in [-T_+, x_0]$.
Lemma 1.1.28 extends $x$ through $(T_+,\ell)$, contradicting maximality. Hence $T_+ = +\infty$.
(iii) Fix $T \geq \max\{t_0, 2\}$, and set $r(t) := \sqrt{t^2 - 2}$, $w(t) := -r(t)$ for $t \geq T$. Since $r^2 - t^2 = w^2 - t^2 = -2$, $r' = t/r > 0 > -2$, and $w' = -t/r \geq -\sqrt{2} > -2$, both $r, w$ are strict upper solutions.
Let $d(t) := t - x(t)$. By (i), $d > 0$ and $t + x > 0$, so $d' = 1 + t^2 - x^2 = 1 + d(t + x) > 1$.
Since $d(t) \to +\infty$ and $t - r(t) \to 0$, choose $T_1 \geq T$ with $d(T_1) > T_1 - r(T_1)$, i.e. $x(T_1) < r(T_1)$.
By strict comparison, $x < r$ for $t > T_1$. If $x(T_1) \leq w(T_1)$, then by strict comparison $x < w$ for $t > T_1$.
If $w(T_1) < x(T_1) < r(T_1)$, then while $w < x < r$, one has $x' < -2$ and $w' \geq -\sqrt{2}$, hence $(x - w)' < -(2 - \sqrt{2})$. Global existence forces a first contact $x(\tau) = w(\tau)$ in finite time, and strict comparison from $\tau$ gives $x < w$ thereafter.
Therefore eventually $0 < x(t) + t < t - r(t) = \frac{2}{t + \sqrt{t^2 - 2}} \to 0$.

*Structural Commentary*: Two nested isocline barriers squeeze the trajectory against $x = -t$.
*Key Insight*: Every trajectory entering the central wedge is asymptotic to its lower edge. □

**Proposition 1.3.9.** *Finite-Time Blow-Up Above the Upper Isocline*
*Let $t_0 \geq 0$, $x_0 \geq \sqrt{t_0^2 + 2}$, and let $x : [t_0,T_+) \to \mathbb{R}$ be the maximal forward solution of $x' = x^2 - t^2$, $x(t_0) = x_0$.*

*(i)* $x(t) > \sqrt{t^2 + 2}$ *and* $x'(t) > 2$ *for* $t \in (t_0,T_+)$.
*(ii)* $T_+ < +\infty$ *and* $x(t) \to +\infty$ *as* $t \uparrow T_+$.

*Proof.*

(i) Set $b(t) := \sqrt{t^2 + 2}$. Then $b^2 - t^2 = 2$ and $b' = \frac{t}{b} < 1 < 2$, so $b' < b^2 - t^2$.
Since $x_0 \geq b(t_0)$, Lemma 1.3.3 implies $b(t) < x(t)$ for $t > t_0$; hence $x' = x^2 - t^2 > b^2 - t^2 = 2$.
(ii) Suppose $T_+ = +\infty$. By (i), $x(t) > x_0 + 2(t - t_0)$; choose $t_1 \geq t_0$ with $x(t) > \sqrt{2}t > 0$ for $t \geq t_1$.
Then $x' = x^2 - t^2 > \frac{1}{2}x^2$ and $(\frac{1}{x})' = -\frac{x'}{x^2} < -\frac{1}{2}$ on $[t_1,+\infty)$, so $\frac{1}{x(t)} < \frac{1}{x(t_1)} - \frac{1}{2}(t - t_1)$.
This contradicts $x > 0$ for $t > t_1 + \frac{2}{x(t_1)}$. Thus $T_+ < +\infty$. Also $x' > 2$, so $\ell := \lim_{t \uparrow T_+} x(t) \in (x_0, +\infty]$.
If $\ell < +\infty$, Lemma 1.1.28 extends $x$ through $(T_+,\ell)$, contradicting maximality. Hence $\ell = +\infty$.

*Structural Commentary*: Above $b(t) = \sqrt{t^2 + 2}$, linear escape enters the Riccati regime.
*Key Insight*: Once $x' \geq \frac{1}{2}x^2$, reciprocal decay forces finite-time escape. □

**Theorem 1.3.10.** *Separatrix Classification for $x' = x^2 - t^2$*
*For $a \in \mathbb{R}$, let $x_a : [0, T_a) \to \mathbb{R}$ be the maximal forward solution of $x' = x^2 - t^2$, $x_a(0) = a$.*

*(i)* $\exists!\, a_* \in [0, \sqrt{2}]$ *such that* $T_{a_*} = +\infty$, $t \le x_{a_*}(t) \le \sqrt{t^2+2}$ *for* $t \ge 0$, *and* $x_{a_*}(t) - t \to 0$ *as* $t \to +\infty$.
*(ii) If* $a > a_*$, *then* $T_a < +\infty$ *and* $x_a(t) \to +\infty$ *as* $t \uparrow T_a$.
*(iii) If* $a < a_*$, *then* $T_a = +\infty$ *and* $x_a(t) + t \to 0$ *as* $t \to +\infty$.

*Proof.* Set $\ell(t) := t$ and $b(t) := \sqrt{t^2+2}$. For $T > 0$, let $p_T, q_T$ solve $x' = x^2 - t^2$ backward from $p_T(T) = \ell(T)$ and $q_T(T) = b(T)$. Under $s = T - t$, the reversed equation is $y' = (T-s)^2 - y^2$, while $\widehat{\ell}(s) := T - s$ and $\widehat{b}(s) := \sqrt{(T-s)^2+2}$ satisfy $\widehat{\ell}' < (T-s)^2 - \widehat{\ell}^2$ and $\widehat{b}' > (T-s)^2 - \widehat{b}^2$.
Lemma 1.3.3 and Theorem 1.3.5 trap and order the reversed solutions while they exist. Corollary 1.1.29 then gives existence on $[0, T]$, with $\ell \le p_T < q_T \le b$ on $[0, T]$ and strict outer inequalities on $[0, T)$.
Set $\alpha_T := p_T(0)$, $\beta_T := q_T(0)$, and $K_T := [\alpha_T, \beta_T]$. Order preservation gives $a \in K_T$ iff $x_a$ exists on $[0, T]$ and $\ell \le x_a \le b$ there. Thus $S \ge T$ implies $K_S \subseteq K_T$, and $(K_n)_{n\ge1}$ is a nested family of nonempty compact subintervals of $[0, \sqrt{2}]$.

(i) Choose $a_* \in \bigcap_{n\ge1} K_n \subseteq [0, \sqrt{2}]$. Then $x_{a_*}$ is global, $\ell \le x_{a_*} \le b$, and $0 \le x_{a_*}(t) - t \le b(t) - t = \frac{2}{b(t)+t} \to 0$.
If $a_1 < a_2$ lie in $\bigcap_{n\ge1} K_n$, then $\delta := x_{a_2} - x_{a_1}$ satisfies $\delta(0) > 0$ and $\delta' = (x_{a_1} + x_{a_2})\delta \ge 0$.
Thus $0 < a_2 - a_1 \le \delta(t) \le b(t) - t \to 0$, impossible, so $\bigcap_{n\ge1} K_n = \{a_*\}$.

(ii) Let $a > a_*$. Choose $n$ with $a > \beta_n$. By comparison, $x_a > q_n > t$, so $x_a > 0$ and $x_a' > 0$ while both exist.
If $T_a \le n$, then $\ell_a := \lim_{t\uparrow T_a} x_a(t) \in (a, +\infty]$. Lemma 1.1.28 excludes $\ell_a < +\infty$, so $x_a(t) \to +\infty$.
If $T_a > n$, then $x_a(n) > q_n(n) = \sqrt{n^2+2}$; Proposition 1.3.9 implies $T_a < +\infty$ and $x_a(t) \to +\infty$ as $t \uparrow T_a$.

(iii) Let $a < a_*$. Choose $n$ with $a < \alpha_n$. By comparison, $x_a < p_n < t$.
On $[0, T_a \wedge n)$, whenever $x_a < -t$ one has $x_a' > 0$; after a contact with $-t$, strict comparison prevents a later downward crossing. Hence $\min\{a, -n\} \le x_a(t) < n$ on this interval. Corollary 1.1.29 therefore gives $T_a > n$.
Now $x_a(n) < n$. If $-n < x_a(n) < n$, Proposition 1.3.8 gives $T_a = +\infty$ and $x_a(t) + t \to 0$.
If $x_a(n) \le -n$, set $h := x_a + t$. While $h < 0$, one has $h' = x_a^2 - t^2 + 1 > 1$ and $x_a(n) \le x_a(t) < -t$. Compact escape gives existence through the finite time $n - h(n)$, so $h$ has a first zero $\tau \in [n, n - h(n)]$. Strict comparison from $\tau$ gives $-t < x_a(t)$ for $t > \tau$, and continuity gives $x_a(t_2) < t_2$ for some $t_2 > \tau$. Proposition 1.3.8 implies $T_a = +\infty$ and $x_a(t) + t \to 0$.

*Structural Commentary*: Nested corridors isolate the threshold separating blow-up from lower-edge attraction.
*Key Insight*: The orbit $x_{a_*} \sim t$ separates blow-up from attraction to $-t$. □

**Theorem 1.3.11.** *Asymptotic Stability by Linearization*
*Let $U \subseteq \mathbb{R}^n$ be open with $0 \in U$, let $f \in C^1(U, \mathbb{R}^n)$ satisfy $f(0) = 0$, and set $A := Df(0)$. Assume $\operatorname{Re}\lambda < 0$ for every $\lambda \in \operatorname{spec}(A)$. Then $\exists r, K, \gamma > 0$ such that, for every $x_0 \in \overline{B}_r(0)$:*

*(i) The IVP $x' = f(x)$, $x(0) = x_0$, has a unique solution $x \in C^1([0, +\infty), U)$.*
*(ii) The solution satisfies $\|x(t)\| \le Ke^{-\gamma t}\|x_0\|$ $\forall t \ge 0$.*

*Proof.* Choose $\alpha \in (0, \min\{-\operatorname{Re}\lambda : \lambda \in \operatorname{spec}(A)\})$. By Theorem 1.2.15(iii), $\exists C \ge 1$ with $\|e^{tA}\| \le Ce^{-\alpha t}$ for $t \ge 0$. Set $g(x) := f(x) - Ax$. Since $Dg(0) = 0$, choose $b \in (0, \alpha/C)$ and $R > 0$ such that $\overline{B}_R(0) \subseteq U$ and $\|g(x)\| \le b\|x\|$ for $\|x\| \le R$.
Set $r := R/(2C)$, $K := C$, and $\gamma := \alpha - Cb > 0$. Fix $x_0 \in \overline{B}_r(0)$. By Lemma 1.1.15 and Theorem 1.1.27, the autonomous field $(t, x) \mapsto f(x)$ has a unique maximal solution $x : [0, T_+) \to U$ with $x(0) = x_0$.

(i) Work before the first exit from $B_R(0)$. By Theorem 1.2.10(iii), $x(t) = e^{tA}x_0 + \int_0^t e^{(t-s)A}g(x(s))\,ds$. For $y(t) := e^{\alpha t}\|x(t)\|$, this gives $y(t) \le C\|x_0\| + Cb\int_0^t y(s)\,ds$.
Lemma 1.1.20(ii) yields $\|x(t)\| \le Ce^{-\gamma t}\|x_0\| \le R/2$, so no exit occurs. If $T_+ < +\infty$, the graph remains in $[0, T_+] \times \overline{B}_{R/2}(0) \Subset \mathbb{R} \times U$, contradicting Corollary 1.1.29. Hence $T_+ = +\infty$; uniqueness is part of Theorem 1.1.27.

(ii) The estimate obtained in (i) is valid for every $t \ge 0$ and is exactly $\|x(t)\| \le Ke^{-\gamma t}\|x_0\|$.

*Structural Commentary*: Split off $Ax$; absorb the differentiable remainder by Gronwall.
*Key Insight*: A Hurwitz linear part dominates its $o(\|x\|)$ remainder. □

## 1.4 Bifurcation Theory

A parameter can change not merely the position of equilibria but also their number and stability. The harvested logistic equation is a complete scalar model of such a transition.

**Definition 1.4.1.** *Scalar Saddle-Node Pattern*
Let $\Lambda \subseteq \mathbb{R}$ be an interval, let $F : \mathbb{R} \times \Lambda \to \mathbb{R}$, and set $F_h(x) := F(x,h)$.
For $(x_c, h_c) \in \mathbb{R} \times \Lambda$, the family $(F_h)_{h\in\Lambda}$ has a *saddle-node pattern* at $(x_c, h_c)$ iff $\exists \varepsilon > 0$, a neighborhood $X \subseteq \mathbb{R}$ of $x_c$, and $\sigma \in \{-1, 1\}$ such that the sets $H_\pm := \{h \in \Lambda : 0 < \pm\sigma(h - h_c) < \varepsilon\}$ are nonempty and:

(i) *Critical section*: $F_{h_c}^{-1}(0) \cap X = \{x_c\}$.
(ii) *Two-equilibrium side*: $\exists x_-, x_+ : H_+ \to X$ such that $F_h^{-1}(0) \cap X = \{x_-(h), x_+(h)\}$ and $x_-(h) < x_+(h)$ $\forall h \in H_+$, with $x_\pm(h) \to x_c$ as $h \to h_c$ through $H_+$.
(iii) *Zero-equilibrium side*: $F_h^{-1}(0) \cap X = \emptyset$ $\forall h \in H_-$.

*Key Insight*: A saddle-node is local creation or annihilation of two equilibria.

**Proposition 1.4.2.** *Equilibrium Branches of the Harvested Logistic Equation*
*For $h \in [0, +\infty)$, let $F_h : \mathbb{R} \to \mathbb{R}$ be given by $F_h(x) := (1 - x)x - h$. Set $h_c := \frac{1}{4}$, $x_c := \frac{1}{2}$, and $x_\pm(h) := \frac{1}{2}(1 \pm \sqrt{1 - 4h})$ for $h \in [0, h_c]$.*

*(i)* $F_h^{-1}(0) = \{x_-(h), x_+(h)\}$ *and* $x_-(h) < x_+(h)$ $\forall h \in [0, h_c)$.
*(ii)* $F_{h_c}^{-1}(0) = \{x_c\}$ *and* $F_{h_c}(x) = -(x - x_c)^2$ $\forall x \in \mathbb{R}$.
*(iii)* $F_h^{-1}(0) = \emptyset$ $\forall h \in (h_c, +\infty)$.
*(iv)* $x_\pm(h) \to x_c$ *as* $h \uparrow h_c$.
*(v) The family $(F_h)_{h\in[0,+\infty)}$ has a saddle-node pattern at $(x_c, h_c)$.*

*Proof.* The identity $F_h(x) = h_c - h - (x - x_c)^2$ holds $\forall (x, h) \in \mathbb{R} \times [0, +\infty)$; hence $F_h(x) = 0$ iff $(x - x_c)^2 = h_c - h$.

(i) For $h \in [0, h_c)$, $h_c - h > 0$ and $x_c \pm \sqrt{h_c - h} = x_\pm(h)$, which gives the two distinct zeros.
(ii) Substitution of $h = h_c$ gives $F_{h_c}(x) = -(x - x_c)^2$ and therefore $F_{h_c}^{-1}(0) = \{x_c\}$.
(iii) For $h > h_c$, the relation $(x - x_c)^2 = h_c - h < 0$ has no solution in $\mathbb{R}$.
(iv) The identity $x_\pm(h) = x_c \pm \sqrt{h_c - h}$ and the limit $h_c - h \downarrow 0$ give $x_\pm(h) \to x_c$.
(v) Take $X = \mathbb{R}$, $\sigma = -1$, and $\varepsilon = \frac{1}{8}$. Then $H_+ = (\frac{1}{8}, \frac{1}{4})$ and $H_- = (\frac{1}{4}, \frac{3}{8})$; items (i)–(iv) establish the critical zero, the two coalescing branches on $H_+$, and the absence of zeros on $H_-$.

*Structural Commentary*: Completing the square controls the zero count on both parameter sides.
*Key Insight*: At $h_c$, two simple roots merge into one double root. □

**Lemma 1.4.3.** *Scalar Phase-Line Barriers and Limits*
*Let $F \in C^1(\mathbb{R}, \mathbb{R})$, and let $x : [t_0, T_+) \to \mathbb{R}$, $T_+ \in (t_0, +\infty]$, be a maximal forward solution of $x' = F(x)$.*

*(i) If $e \in F^{-1}(0)$ and $\exists \tau \in [t_0, T_+)$ such that $x(\tau) = e$, then $x(t) = e$ $\forall t \in [t_0, T_+)$.*
*(ii) If $x$ is monotone and $x([t_0, T_+))$ is bounded, then $T_+ = +\infty$ and $\exists \ell \in F^{-1}(0)$ such that $x(t) \to \ell$ as $t \to +\infty$.*

*Proof.*

(i) The constant function $u \equiv e$ satisfies $u' = F(u)$ and $u(\tau) = x(\tau)$. Set $S := \{t \in [t_0, T_+) : x(t) = e\}$. Continuity makes $S$ relatively closed, while local uniqueness makes $S$ relatively open. Since $\tau \in S$ and $[t_0, T_+)$ is connected, $S = [t_0, T_+)$.
(ii) Monotone boundedness gives $\ell := \lim_{t\uparrow T_+} x(t) \in \mathbb{R}$. If $T_+ < +\infty$, by Lemma 1.1.28, $x$ extends through $(T_+, \ell)$, contradicting maximality. Thus $T_+ = +\infty$.
Suppose $F(\ell) \neq 0$. By continuity and $x(t) \to \ell$, $\exists c > 0$ and $t_1 \geq t_0$ such that $F(x(t))$ has constant sign and $|F(x(t))| \geq c$ $\forall t \geq t_1$. Hence $|x(t) - x(s)| = \left|\int_s^t F(x(r))\, dr\right| \geq c(t - s)$ $\forall t > s \geq t_1$, contradicting $x(t) \to \ell$. Therefore $F(\ell) = 0$.

*Structural Commentary*: Uniqueness makes equilibria barriers; maximal extension identifies bounded monotone limits.
*Key Insight*: A finite phase-line limit must be a zero of $F$. □

**Theorem 1.4.4.** *Autonomous Scalar Phase-Line Classification*
*Let $F \in C^1(\mathbb{R}, \mathbb{R})$ and let $x : [t_0, T_+) \to \mathbb{R}$ be the maximal forward solution of $x' = F(x)$ with $x(t_0) = x_0$.*
*Set $E := F^{-1}(0)$. If $x_0 \notin E$, set $\sigma := \operatorname{sgn}(F(x_0))$ and $Z_\sigma := \{e \in E : \sigma(e - x_0) > 0\}$.*

*(i) If $x_0 \in E$, then $x \equiv x_0$.*
*(ii) If $x_0 \notin E$, then $(\sigma x)'(t) > 0$ $\forall t \in [t_0, T_+)$.*
*(iii) If $x_0 \notin E$ and $Z_\sigma \neq \emptyset$, then $\exists! e_\sigma \in Z_\sigma$ such that $\sigma e_\sigma = \inf\{\sigma e : e \in Z_\sigma\}$; moreover, $T_+ = +\infty$ and $\sigma(x(t) - e_\sigma) \uparrow 0$ as $t \to +\infty$.*
*(iv) If $x_0 \notin E$ and $Z_\sigma = \emptyset$, then $\sigma x(t) \uparrow +\infty$ as $t \uparrow T_+$.*

*Proof.*

(i) Since $x_0 \in E$, by Lemma 1.4.3(i), $x(t) = x_0\ \forall t \in [t_0, T_+)$.

(ii) Let $C$ be the component of $\mathbb{R} \setminus E$ containing $x_0$. Continuity gives $\sigma F > 0$ on $C$. If $x$ left $C$, continuity would give $x(\tau) \in E$ for some $\tau$, and Lemma 1.4.3(i) would force $x \equiv x(\tau)$, contradicting $x_0 \notin E$. Thus $x([t_0, T_+)) \subseteq C$ and $(\sigma x)' = \sigma F(x) > 0$.

(iii) Since $F(x_0) \neq 0$, continuity gives $\delta > 0$ such that $E \cap [x_0 - \delta, x_0 + \delta] = \emptyset$. Let $m := \inf\{\sigma e : e \in Z_\sigma\}$ and choose $(e_n) \subseteq Z_\sigma$ with $\sigma e_n \to m$. Then $(e_n)$ converges to $e_\sigma := \sigma m$. Since $E$ is closed, $e_\sigma \in E$; the zero-free neighborhood gives $\sigma(e_\sigma - x_0) > 0$, so $e_\sigma \in Z_\sigma$. Its defining equality makes it unique.
No zero lies strictly between $x_0$ and $e_\sigma$. By (ii) and Lemma 1.4.3(i), $\sigma x$ is increasing and bounded above by $\sigma e_\sigma$. Lemma 1.4.3(ii) gives $T_+ = +\infty$ and $x(t) \to \ell \in E$. Since $\sigma x_0 < \sigma \ell \leq \sigma e_\sigma$, one has $\ell \in Z_\sigma$; minimality gives $\sigma e_\sigma \leq \sigma \ell$, hence $\ell = e_\sigma$.

(iv) By (ii), $\sigma x$ is strictly increasing. If it were bounded above, Lemma 1.4.3(ii) would give $x(t) \to \ell \in E$ with $\sigma(\ell - x_0) > 0$, so $\ell \in Z_\sigma$, a contradiction. Therefore $\sigma x(t) \uparrow +\infty$ as $t \uparrow T_+$.

*Structural Commentary*: Components of $\mathbb{R} \setminus E$ fix the sign; the nearest zero supplies the barrier.
*Key Insight*: A trajectory approaches its next zero or escapes in the sign direction. □

**Theorem 1.4.5.** *Subcritical Harvest Phase Line*
*Let $h_c := \frac{1}{4}$, $F_h(x) := (1-x)x - h$, $0 \leq h < h_c$, and $x_\pm := \frac{1}{2}(1 \pm \sqrt{1-4h})$.*
*Let $x : [t_0, T_+) \to \mathbb{R}$ be the maximal forward solution of $x' = F_h(x)$ with $x(t_0) = x_0$.*

*(i) If $x_0 \in \{x_-, x_+\}$, then $T_+ = +\infty$ and $x \equiv x_0$.*
*(ii) If $x_0 < x_-$, then $x'(t) < 0\ \forall t \in [t_0, T_+)$ and $x(t) \to -\infty$ as $t \uparrow T_+$.*
*(iii) If $x_- < x_0 < x_+$, then $T_+ = +\infty$, $x'(t) > 0\ \forall t \geq t_0$, and $x(t) \to x_+$ as $t \to +\infty$.*
*(iv) If $x_0 > x_+$, then $T_+ = +\infty$, $x'(t) < 0\ \forall t \geq t_0$, and $x(t) \to x_+$ as $t \to +\infty$.*

*Proof.* The factorization $F_h = -(\cdot - x_-)(\cdot - x_+)$ gives $F_h < 0$ on $(-\infty, x_-) \cup (x_+, +\infty)$ and $F_h > 0$ on $(x_-, x_+)$.

(i) Since $x_0 \in F_h^{-1}(0)$, by Theorem 1.4.4(i), $x \equiv x_0$. The constant solution is defined on $[t_0, +\infty)$, so maximality gives $T_+ = +\infty$.

(ii) Here $\operatorname{sgn}(F_h(x_0)) = -1$, and no zero lies below $x_0$. By Theorem 1.4.4(ii),(iv), $x' < 0$ and $-x(t) \uparrow +\infty$ as $t \uparrow T_+$.

(iii) Here $\operatorname{sgn}(F_h(x_0)) = 1$, and $x_+$ is the nearest zero above $x_0$. By Theorem 1.4.4(ii),(iii), $T_+ = +\infty$, $x' > 0$, and $x(t) \to x_+$.

(iv) Here $\operatorname{sgn}(F_h(x_0)) = -1$, and $x_+$ is the nearest zero below $x_0$. By Theorem 1.4.4(ii),(iii), $T_+ = +\infty$, $x' < 0$, and $x(t) \to x_+$.

*Structural Commentary*: The factorization fixes three sign intervals; the phase-line theorem handles each.
*Key Insight*: The upper root attracts every initial state above the lower threshold. □

**Theorem 1.4.6.** *Critical Harvest Phase Line*
*Let $h_c := \frac{1}{4}$, $x_c := \frac{1}{2}$, $F_h(x) := (1-x)x - h$, and $h = h_c$.*
*Let $x : [t_0, T_+) \to \mathbb{R}$ be the maximal forward solution of $x' = F_{h_c}(x)$ with $x(t_0) = x_0$.*

*(i) If $x_0 = x_c$, then $T_+ = +\infty$ and $x \equiv x_c$.*
*(ii) If $x_0 > x_c$, then $T_+ = +\infty$, $x'(t) < 0\ \forall t \geq t_0$, and $x(t) \to x_c$ as $t \to +\infty$.*
*(iii) If $x_0 < x_c$, then $x'(t) < 0\ \forall t \in [t_0, T_+)$ and $x(t) \to -\infty$ as $t \uparrow T_+$.*

*Proof.* The identity $F_{h_c}(x) = -(x - x_c)^2$ gives $F_{h_c}^{-1}(0) = \{x_c\}$ and $F_{h_c} < 0$ on $\mathbb{R} \setminus \{x_c\}$.

(i) Since $x_0 \in F_{h_c}^{-1}(0)$, by Theorem 1.4.4(i), $x \equiv x_c$. The constant solution is defined on $[t_0, +\infty)$, so maximality gives $T_+ = +\infty$.

(ii) Here $\operatorname{sgn}(F_{h_c}(x_0)) = -1$, and $x_c$ is the nearest zero below $x_0$. By Theorem 1.4.4(ii),(iii), $T_+ = +\infty$, $x' < 0$, and $x(t) \to x_c$.

(iii) Here $\operatorname{sgn}(F_{h_c}(x_0)) = -1$, and no zero lies below $x_0$. By Theorem 1.4.4(ii),(iv), $x' < 0$ and $-x(t) \uparrow +\infty$ as $t \uparrow T_+$.

*Structural Commentary*: The double zero is the sole barrier; the sign is negative on both sides.
*Key Insight*: The critical equilibrium attracts from above and repels toward $-\infty$ below. □

**Theorem 1.4.7.** *Supercritical Harvest Phase Line*
*Let $h_c := \frac{1}{4}$, $x_c := \frac{1}{2}$, $F_h(x) := (1-x)x - h$, and $h > h_c$.*
*Let $x : [t_0, T_+) \to \mathbb{R}$ be the maximal forward solution of $x' = F_h(x)$ with $x(t_0) = x_0$. Then:*

*(i) $x'(t) < 0\ \forall t \in [t_0, T_+)$.*
*(ii) $x(t) \to -\infty$ as $t \uparrow T_+$.*

*Proof.* For $h > h_c$, $F_h = -(\cdot - x_c)^2 - (h - h_c) < 0$ on $\mathbb{R}$ and $F_h^{-1}(0) = \emptyset$.

(i) By Theorem 1.4.4(ii), with $\sigma = -1$, $x' < 0$.

(ii) Since $F_h^{-1}(0) = \emptyset$, by Theorem 1.4.4(iv), $-x(t) \uparrow +\infty$ as $t \uparrow T_+$.

*Structural Commentary*: The completed square removes every zero, so the phase-line escape case applies.
*Key Insight*: Above critical harvest, every trajectory escapes toward $-\infty$. □

**Corollary 1.4.8.** *Critical Dimensional Harvest Rate*
*Let $\kappa, N_{\max} > 0$, let $H \geq 0$, and set $G_H(N) := \kappa(1 - \frac{N}{N_{\max}})N - H$ and $H_c := \frac{\kappa N_{\max}}{4}$.*
*Let $N : [t_0, T_+) \to \mathbb{R}$ be the maximal forward solution of $N' = G_H(N)$ with $N(t_0) = N_0$.*

*(i) $G_H^{-1}(0) \cap (0, +\infty) \neq \emptyset$ iff $H \in [0, H_c]$, and $H_c = \max\{\eta \geq 0 : G_\eta^{-1}(0) \cap (0, +\infty) \neq \emptyset\}$.*

*(ii) For $H \in [0, H_c)$, set $N_\pm(H) := \frac{N_{\max}}{2}(1 \pm \sqrt{1 - \frac{4H}{\kappa N_{\max}}})$. Then $G_H^{-1}(0) = \{N_-(H), N_+(H)\}$ and $G_{H_c}^{-1}(0) = \{\frac{N_{\max}}{2}\}$.*

*(iii) If $H \in [0, H_c)$ and $N_0 > N_-(H)$, then $T_+ = +\infty$ and $N(t) \to N_+(H)$ as $t \to +\infty$.*

*Proof.* Set $h := \frac{H}{\kappa N_{\max}}$, $h_c := \frac{1}{4}$, $x_c := \frac{1}{2}$, $x_\pm(h) := \frac{1}{2}(1 \pm \sqrt{1 - 4h})$ for $h \in [0, h_c]$, and $F_h(x) := (1 - x)x - h$. Then $G_H(N) = \kappa N_{\max} F_h(\frac{N}{N_{\max}})$.
For the solution, set $x(\tau) := \frac{N(t_0 + \tau/\kappa)}{N_{\max}}$, $x_0 := \frac{N_0}{N_{\max}}$, and $\widehat{T}_+ := \kappa(T_+ - t_0)$. Thus $x : [0, \widehat{T}_+) \to \mathbb{R}$ is the maximal forward solution of $\frac{dx}{d\tau} = F_h(x)$ with $x(0) = x_0$.

(i) By Proposition 1.4.2(i)–(iii), $F_h^{-1}(0) \cap (0, +\infty) \neq \emptyset$ iff $h \in [0, h_c]$. Since $H = \kappa N_{\max} h$ and $H_c = \kappa N_{\max} h_c$, the asserted equivalence and maximum follow.

(ii) By Proposition 1.4.2(i),(ii), $F_h^{-1}(0) = \{x_-(h), x_+(h)\}$ for $h < h_c$ and $F_{h_c}^{-1}(0) = \{x_c\}$. Multiplication by $N_{\max}$ gives $N_\pm(H) = N_{\max} x_\pm(h)$ and $N_{\max} x_c = \frac{N_{\max}}{2}$.

(iii) The inequality $N_0 > N_-(H)$ is equivalent to $x_0 > x_-(h)$. By Theorem 1.4.5(i),(iii),(iv), $\widehat{T}_+ = +\infty$ and $x(\tau) \to x_+(h)$ as $\tau \to +\infty$. Hence $T_+ = +\infty$ and $N(t) \to N_{\max} x_+(h) = N_+(H)$.

*Structural Commentary*: State and time scaling reduce the dimensional model to the universal quadratic family.
*Key Insight*: The threshold is one quarter of the unharvested growth scale. □

## 1.5 Dynamical Systems Structure

For a time-periodic scalar equation, one period of continuous evolution can be compressed into a discrete map. Periodic solutions then become fixed points, and their long-time behavior can be studied through iteration.

**Definition 1.5.1.** *Scalar Poincaré Map*
Let $f : \mathbb{R}^2 \to \mathbb{R}$ be one-periodic in $t$: $f(t + 1, x) = f(t, x)$.
For $x_0 \in \mathbb{R}$, let $\phi_{x_0} : I_{x_0} \to \mathbb{R}$ be maximal solution of $x' = f(t, x)$, $\phi_{x_0}(0) = x_0$, and write $\phi(t, x_0) := \phi_{x_0}(t)$.
Put $\mathcal{D} := \{x_0 \in \mathbb{R} : [0, 1] \subseteq I_{x_0}\}$. The *Poincaré map* is $P : \mathcal{D} \to \mathbb{R}$, $P(x_0) := \phi(1, x_0)$.
*Key Insight*: $P$ is the time-1 flow map on $\mathcal{D}$.

**Proposition 1.5.2.** *Periodic Orbits and Ordered Returns*
*Let $f \in C^1(\mathbb{R}^2, \mathbb{R})$ satisfy $f(t + 1, x) = f(t, x)$.*
*For $x_0 \in \mathbb{R}$, let $\phi_{x_0} : I_{x_0} \to \mathbb{R}$ be maximal solution of $x' = f(t, x)$, $\phi_{x_0}(0) = x_0$; set $\phi(t, x_0) := \phi_{x_0}(t)$.*
*Set $\mathcal{D} := \{x_0 \in \mathbb{R} : [0, 1] \subseteq I_{x_0}\}$ and $P : \mathcal{D} \to \mathbb{R}$, $P(x_0) := \phi(1, x_0)$.*

*(i) For $x_0 \in \mathcal{D}$, $P(x_0) = x_0$ iff $\phi(\cdot, x_0)$ extends to a one-periodic solution on $\mathbb{R}$.*

*(ii) If $x_0, y_0 \in \mathcal{D}$ and $x_0 < y_0$, then $P(x_0) < P(y_0)$.*

*Proof.*

(i) ($\Rightarrow$) Suppose $P(x_0) = x_0$. For $t \in [n, n + 1]$, $n \in \mathbb{Z}$, set $u(t) := \phi(t - n, x_0)$. Since $\phi(1, x_0) = x_0 = \phi(0, x_0)$ and $f(n + 1, x_0) = f(n, x_0)$, the pieces match in value and derivative at each $n \in \mathbb{Z}$.
Thus $u \in C^1(\mathbb{R}, \mathbb{R})$, $u' = f(t, u)$, and $u(t + 1) = u(t)$. By local uniqueness, $u$ extends $\phi(\cdot, x_0)$.

(i) ($\Leftarrow$) If $u$ is a one-periodic solution with $u(0) = x_0$, then by local uniqueness $u = \phi(\cdot, x_0)$ on $[0, 1]$.
Hence $P(x_0) = \phi(1, x_0) = u(1) = u(0) = x_0$.

(ii) Let $K := \{(t, \phi(t, x_0)) : t \in [0, 1]\} \cup \{(t, \phi(t, y_0)) : t \in [0, 1]\}$.
Since $\partial_x f$ is continuous on the compact set $K$, $f$ is state-Lipschitz on a compact neighborhood of $K$.
By Theorem 1.3.5(ii), $\phi(t, x_0) < \phi(t, y_0)$ for $t \in [0, 1]$; at $t = 1$, $P(x_0) < P(y_0)$.

*Structural Commentary*: Periodicity closes the segment exactly when $P(x_0) = x_0$.
*Key Insight*: Scalar uniqueness preserves order; $P$ is strictly increasing. □

**Proposition 1.5.3.** *Periodic Scalar Linear Forcing*
*Let $a \in \mathbb{R}$, let $g \in C(\mathbb{R},\mathbb{R})$ satisfy $g(t+1) = g(t)$ and $g \geq 0$, and set $I_a := \int_0^1 e^{-as} g(s)\, ds$.*
*For $x' = ax + g(t)$, write $x(\,\cdot\,; x_0)$ for the solution with $x(0) = x_0$, and set $P(x_0) := x(1; x_0)$.*
*If $a \neq 0$, put $x_{0,*} := -\frac{e^a}{e^a - 1} I_a$ and $x_*(t) := e^{at}\big(x_{0,*} + \int_0^t e^{-as} g(s)\, ds\big)$.*

*(i) If $a \neq 0$, then $x_*$ is the unique one-periodic solution.*
*(ii) If $a = 0$ and $g \equiv 0$, then the one-periodic solutions are exactly $x(t) \equiv c$, $c \in \mathbb{R}$.*
*(iii) If $a = 0$ and $g \not\equiv 0$, then no one-periodic solution exists.*

*Proof.* By Proposition 1.2.1, $x(t; x_0) = e^{at}\big(x_0 + \int_0^t e^{-as} g(s)\, ds\big)$. Hence $P(x_0) - x_0 = (e^a - 1)x_0 + e^a I_a$.
If $P(x_0) = x_0$, then $d(t) := x(t+1; x_0) - x(t; x_0)$ satisfies $d' = ad$ and $d(0) = 0$, so $d \equiv 0$.
Conversely, one-periodicity gives $P(x_0) = x_0$. Thus one-periodic solutions correspond exactly to $\operatorname{Fix} P$.

(i) If $a \neq 0$, then $e^a - 1 \neq 0$, so $\operatorname{Fix} P = \{x_{0,*}\}$. The formula for $x_*$ is the solution formula at $x_0 = x_{0,*}$.
(ii) If $a = 0$ and $g \equiv 0$, then $x(t; x_0) = x_0$ for every $x_0 \in \mathbb{R}$.
(iii) If $a = 0$ and $g \not\equiv 0$, then $I_0 = \int_0^1 g(s)\, ds > 0$, so $P(x_0) - x_0 = I_0 \neq 0$ for every $x_0 \in \mathbb{R}$.

*Structural Commentary*: Variation of constants reduces periodicity to fixed points.
*Key Insight*: $P(x_0) - x_0 = (e^a - 1)x_0 + e^a I_a$. □

**Proposition 1.5.4.** *Geometry of the Periodically Harvested Return Map*
*Let $g(t) := 1 - \sin(2\pi t)$, $F_\eta(t,z) := (1-z)z - \eta g(t)$, and $(h,x) \in [0,+\infty) \times \mathbb{R}$.*
*Assume $\exists W \ni (h,x)$ such that $\phi_\eta(\,\cdot\,, y)$ solves $z' = F_\eta(t,z)$, $z(0) = y$, on $[0,1]$ $\forall (\eta, y) \in W$.*
*Set $P_\eta(y) := \phi_\eta(1, y)$ and $\theta_h(t,x) := \partial_x \phi_h(t,x)$. Then:*

*(i) $P_h'(x) = \exp\Big(\int_0^1 (1 - 2\phi_h(s,x))\, ds\Big) > 0$.*
*(ii) $P_h''(x) = -2P_h'(x) \int_0^1 \theta_h(s,x)\, ds < 0$.*
*(iii) $\partial_h P_h(x) = -\int_0^1 \exp\Big(\int_s^1 (1 - 2\phi_h(r,x))\, dr\Big)\, g(s)\, ds < 0$.*

*Proof.* By smooth dependence on $W$, all derivatives below exist.
Put $\phi(t) := \phi_h(t,x)$, $a(t) := 1 - 2\phi(t)$, $\theta := \partial_x \phi_h(\,\cdot\,, x)$.

(i) $\theta' = a\theta$, $\theta(0) = 1$, so $\theta(t) = \exp\Big(\int_0^t a(s)\, ds\Big) > 0$ and $P_h'(x) = \theta(1) = \exp\Big(\int_0^1 a(s)\, ds\Big) > 0$.
(ii) For $\zeta := \partial_x^2 \phi_h(\,\cdot\,, x)$, $\zeta' = a\zeta - 2\theta^2$, $\zeta(0) = 0$.
By variation of constants, $P_h''(x) = \zeta(1) = -2\theta(1) \int_0^1 \theta(s)\, ds = -2P_h'(x) \int_0^1 \theta_h(s,x)\, ds < 0$.
(iii) For $\psi := \partial_h \phi_h(\,\cdot\,, x)$, $\psi' = a\psi - g$, $\psi(0) = 0$. By variation of constants,
$\partial_h P_h(x) = \psi(1) = -\int_0^1 \exp\Big(\int_s^1 a(r)\, dr\Big)\, g(s)\, ds < 0$, since $\exp\Big(\int_s^1 a(r)\, dr\Big) > 0$, $g \geq 0$, and $g \not\equiv 0$.

*Structural Commentary*: Variational equations control $P_h'$, $P_h''$, and $\partial_h P_h$.
*Key Insight*: $P_h$ is increasing, concave, and decreases with $h$. □

**Proposition 1.5.5.** *Concave Periodic Return-Map Geometry*
*Let $f \in C^2(\mathbb{R},\mathbb{R})$ with $f'' < 0$, and let $g \in C(\mathbb{R},\mathbb{R})$ satisfy $g(t+1) = g(t)$, $g \geq 0$, and $g \not\equiv 0$. $(h,x) \in \mathbb{R}^2$.*
*Assume $\phi_\eta(\,\cdot\,, y)$ solves $z' = f(z) + \eta g(t)$, $z(0) = y$, on $[0,1]$ near $(h,x)$, with two $y$-derivatives, one $\eta$-derivative.*
*Put $P_\eta(y) := \phi_\eta(1, y)$, $\phi(t) := \phi_h(t,x)$, $A(t) := f'(\phi(t))$, and $\theta(t) := \partial_x \phi_h(t,x)$. Then:*

*(i) $P_h'(x) = \exp\Big(\int_0^1 A(s)\, ds\Big) > 0$.*
*(ii) $P_h''(x) = P_h'(x) \int_0^1 f''(\phi(s))\theta(s)\, ds < 0$.*
*(iii) $\partial_h P_h(x) = \int_0^1 \exp\Big(\int_s^1 A(r)\, dr\Big)\, g(s)\, ds > 0$.*

*Proof.*

(i) $\theta' = A\theta$ and $\theta(0) = 1$. Hence $\theta(t) = \exp\Big(\int_0^t A(s)\, ds\Big) > 0$, so $P_h'(x) = \theta(1)$.
(ii) Put $\zeta := \partial_x^2 \phi_h(\,\cdot\,, x)$. Then $\zeta' = A\zeta + f''(\phi)\theta^2$ and $\zeta(0) = 0$.
By Proposition 1.2.1, $\zeta(1) = \theta(1) \int_0^1 f''(\phi(s))\theta(s)\, ds$. Since $f'' < 0$ and $\theta > 0$, $P_h''(x) = \zeta(1) < 0$.
(iii) Put $\psi := \partial_h \phi_h(\,\cdot\,, x)$. Then $\psi' = A\psi + g$ and $\psi(0) = 0$.
By Proposition 1.2.1, $\psi(1) = \int_0^1 \exp\Big(\int_s^1 A(r)\, dr\Big)\, g(s)\, ds > 0$. Hence $\partial_h P_h(x) = \psi(1)$.

*Structural Commentary*: Variational equations determine return-map derivatives.
*Key Insight*: Concavity comes from $f'' < 0$; drift from $g \geq 0$, $g \not\equiv 0$. □

**Example 1.5.6.** *The Unharvested Return Map*
For $h = 0$ and $x \geq 0$, separation of variables gives $\phi_0(t,x) = \frac{e^t x}{1 + (e^t - 1)x}$ and hence $P_0(x) = \frac{ex}{1 + (e-1)x}$.
Its fixed points are $x_1 = 0$ and $x_2 = 1$, corresponding to the two constant one-periodic solutions.

**Theorem 1.5.7.** *Iteration of an Increasing Concave Map*
*Let $I \subseteq \mathbb{R}$ be an interval, $P : I \to I$ continuous, strictly increasing, strictly concave, and $\operatorname{Fix}(P) = \{x_1, x_2\}$ with $x_1 < x_2$. For $x \in I$, set $x_0 := x$ and $x_{n+1} := P(x_n)$.*

*(i) If $x > x_1$, then $x_n \to x_2$.*
*(ii) If $x = x_1$, then $x_n = x_1$ for all $n \geq 0$.*
*(iii) If $x < x_1$, then $x_{n+1} < x_n$ and $x_n \downarrow \inf I$ in $\overline{\mathbb{R}}$; in particular, $x_n \to -\infty$ if $\inf I = -\infty$.*

*Proof.* Set $m(a,b) := \frac{P(b)-P(a)}{b-a}$, $a < b$ in $I$. By strict concavity, $a < b < c \Rightarrow m(a,b) > m(b,c)$, $m(x_1, x_2) = 1$.

(i) If $x \in (x_1, x_2)$, then $m(x_1, x) > 1$, hence $x < P(x) < x_2$; thus $x_n \uparrow \ell \leq x_2$.
If $\ell < x_2$, then by continuity, $P(\ell) = \ell \notin \{x_1, x_2\}$, contradiction. Hence $\ell = x_2$. If $x = x_2$, then $x_n = x_2$.
If $x > x_2$, then $m(x_2, x) < 1$, hence $x_2 < P(x) < x$; thus $x_n \downarrow \ell \geq x_2$, and continuity forces $\ell = x_2$.
(ii) Since $P(x_1) = x_1$, induction yields $x_0 = x_1 \Rightarrow x_n = x_1$ for all $n \geq 0$.
(iii) If $x < x_1$, then $m(x, x_1) > 1$, so $P(x) < x$. Hence $x_{n+1} < x_n < x_1$ and $x_n \downarrow \ell$ in $\overline{\mathbb{R}}$ with $\ell \geq \inf I$.
If $\ell > \inf I$, then $\ell \in I$ and, by continuity, $P(\ell) = \ell < x_1$, impossible. Thus $\ell = \inf I$.

*Structural Commentary*: Decreasing secant slopes determine the sign of $P(x) - x$ across the fixed points.
*Key Insight*: The upper fixed point attracts all iterates above the lower threshold. □

**Corollary 1.5.8.** *Periodic Logistic Threshold Structure*
*Let $J$ be an interval contained in the one-period domain of $P_h$.*

*(i) $\#\operatorname{Fix}(P_h|_J) \leq 2$; hence there are at most two one-periodic solutions with initial values in $J$.*
*(ii) If $P_h(J) \subseteq J$, $\operatorname{Fix}(P_h|_J) = \{x_1, x_2\}$ with $x_1 < x_2$, and $x > x_1$, then $P_h^n(x) \to x_2$.*
*(iii) If $P_h(J) \subseteq J$, $\operatorname{Fix}(P_h|_J) = \{x_1, x_2\}$ with $x_1 < x_2$, and $x < x_1$, then $P_h^n(x) \downarrow \inf J$ in $\overline{\mathbb{R}}$.*
*(iv) Wherever $(h, x) \mapsto P_h(x)$ is locally defined, $\partial_h P_h(x) < 0$.*

*Proof.* For (ii),(iii), write $H_J$ for $P_h(J) \subseteq J$, $\operatorname{Fix}(P_h|_J) = \{x_1, x_2\}$, and $x_1 < x_2$.

(i) Suppose $a < b < c$ lie in $\operatorname{Fix}(P_h|_J)$. By Proposition 1.5.4(ii), $P_h|_J$ is strictly concave.
Hence $P_h(b) > L_{a,c}(b) = b$, where $L_{a,c}$ is the chord through $(a,a)$ and $(c,c)$, contradicting $P_h(b) = b$.
Thus $\#\operatorname{Fix}(P_h|_J) \leq 2$. By Proposition 1.5.2(i), $\operatorname{Fix}(P_h|_J)$ is the set of one-periodic initial values in $J$.
(ii) Under $H_J$, $P_h|_J : J \to J$ is continuous, strictly increasing, and strictly concave, with fixed points $x_1 < x_2$.
For $x > x_1$, Theorem 1.5.7(i) applies: $P_h^n(x) \to x_2$.
(iii) Under $H_J$ and $x < x_1$, Theorem 1.5.7(iii) applied to $P_h|_J : J \to J$ yields $P_h^n(x) \downarrow \inf J$ in $\overline{\mathbb{R}}$.
(iv) Local definition of $(h, x) \mapsto P_h(x)$ is the hypothesis of Proposition 1.5.4(iii); hence $\partial_h P_h(x) < 0$.

*Structural Commentary*: Periodic dynamics reduce to a concave return graph versus the diagonal.
*Key Insight*: Two periodic states form a threshold and an attracting upper response. □

## 1.6 Semigroups and Evolution Viewpoint

## 1.7 Measure-Preserving Dynamics and Ergodicity

Measure-preserving dynamics study maps whose pullbacks keep mass fixed. The central question is whether time averages along one orbit recover space averages with respect to an invariant probability measure.

**Definition 1.7.1.** *Measure-Preserving and Ergodic Transformations*
Let $(X, \mathcal{M}, \mu)$ be a measure space and $T : X \to X$.
For $E, A, B \in \mathcal{M}$, write $T^{-1}E := \{x \in X : T(x) \in E\}$ and $A \triangle B := (A \setminus B) \cup (B \setminus A)$.

(i) $T$ is *measurable* iff $T^{-1}E \in \mathcal{M}$ for all $E \in \mathcal{M}$.
(ii) A measurable $T$ is *measure preserving* iff $\mu(T^{-1}E) = \mu(E)$ for all $E \in \mathcal{M}$.
(iii) For measurable $T$, a set $A \in \mathcal{M}$ is *invariant modulo $\mu$* iff $\mu(A \triangle T^{-1}A) = 0$.
(iv) If $\mu(X) = 1$, $T$ measure preserving, then $T$ *ergodic* iff $\mu(A \triangle T^{-1}A) = 0$ implies $\mu(A) \in \{0, 1\}$ $\forall A \in \mathcal{M}$.

*Key Insight*: Pullbacks preserve mass; ergodic invariants are trivial.

**Proposition 1.7.2.** *Integral Test for Measure Preservation*
*Let $(X, \mathcal{M}, \mu)$ be a finite measure space and let $T : X \to X$ be measurable.*
*Then $T$ is measure preserving iff $\int_X g \circ T \, d\mu = \int_X g \, d\mu$ for all $g \in L^1(X, \mu)$.*

*Proof.*

(⇒) Assume $T$ is measure preserving. For $A \in \mathcal{M}$, $\int_X \mathbf{1}_A \circ T \, d\mu = \mu(T^{-1}A) = \mu(A) = \int_X \mathbf{1}_A \, d\mu$. Thus $s = \sum_{k=1}^n c_k \mathbf{1}_{A_k}$ satisfies $\int_X s \circ T \, d\mu = \int_X s \, d\mu$ by linearity. If $0 \leq s_n \uparrow g$, then $0 \leq s_n \circ T \uparrow g \circ T$, so $\int_X g \circ T \, d\mu = \lim_n \int_X s_n \circ T \, d\mu = \lim_n \int_X s_n \, d\mu = \int_X g \, d\mu$. For $g \in L^1(X, \mu)$, apply this to $g^{\pm}$.
(⇐) For $A \in \mathcal{M}$, $\mu(X) < \infty$; $\mathbf{1}_A \in L^1(X, \mu)$, and $\mu(T^{-1}A) = \int_X \mathbf{1}_{T^{-1}A} \, d\mu = \int_X \mathbf{1}_A \circ T \, d\mu = \int_X \mathbf{1}_A \, d\mu = \mu(A)$.

*Structural Commentary*: Indicators encode sets; approximation extends the identity to observables.
*Key Insight*: Measure preservation is equality of integrals after pullback. □

**Remark 1.7.3.** *Invariant Sets as Invariant Indicators*
For a measure-preserving $T$, a measurable set $A$ is invariant modulo $\mu$ if and only if $\mathbf{1}_A \circ T = \mathbf{1}_A$ $\mu$-a.e.

**Proposition 1.7.4.** *Invariant Functions Detect Ergodicity*
*Let $(X, \mathcal{M}, \mu)$ be a probability space and let $T : X \to X$ be measure preserving. Then $T$ is ergodic iff every measurable $g : X \to \mathbb{R}$ with $g \circ T = g$ $\mu$-a.e. satisfies $g = c$ $\mu$-a.e. for some $c \in \mathbb{R}$.*

*Proof.*

($\Rightarrow$) Assume $T$ is ergodic and $g \circ T = g$ $\mu$-a.e. For $a \in \mathbb{Q}$ set $A_a := \{g < a\}$. Since $\mathbf{1}_{A_a} \circ T \neq \mathbf{1}_{A_a}$ only where $g \circ T \neq g$, $\mu(A_a \triangle T^{-1}A_a) = 0$, so $\mu(A_a) \in \{0, 1\}$. Since $A_n \uparrow X$ and $A_{-n} \downarrow \varnothing$, some $a_0, b_0 \in \mathbb{Q}$ satisfy $\mu(A_{a_0}) = 0$ and $\mu(A_{b_0}) = 1$. Let $c := \inf\{a \in \mathbb{Q} : \mu(A_a) = 1\}$. For rational $a < c < b$, $\mu(A_a) = 0$ and $\mu(X \setminus A_b) = 0$; by countable unions, $\mu(g < c) = \mu(g > c) = 0$, hence $g = c$ $\mu$-a.e.

($\Leftarrow$) Assume $g \circ T = g$ $\mu$-a.e. implies $g$ is constant $\mu$-a.e. for every measurable $g : X \to \mathbb{R}$. If $A \in \mathcal{M}$ and $\mu(A \triangle T^{-1}A) = 0$, then $\mathbf{1}_A \circ T = \mathbf{1}_{T^{-1}A} = \mathbf{1}_A$ $\mu$-a.e. Thus $\mathbf{1}_A$ is constant $\mu$-a.e., so $\mu(A) \in \{0, 1\}$.

*Structural Commentary*: Level sets convert invariant functions into invariant sets.
*Key Insight*: Ergodicity collapses invariant observables to constants. □

**Method 1.7.5.** *Krylov–Bogoliubov Averaging*
Let $X \neq \varnothing$ be compact metric, $f \in C(X, X)$, and $\mathcal{P}(X) := \{\text{Borel probability measures on } X\}$.

(i) Choose $\eta \in \mathcal{P}(X)$ and set $\mu_n(E) := \frac{1}{n}\sum_{k=0}^{n-1} \eta((f^k)^{-1}E)$ for $E \in \mathcal{B}(X)$.
(ii) Equivalently use $\int_X g \, d\mu_n = \int_X \frac{1}{n}\sum_{k=0}^{n-1} g \circ f^k \, d\eta$ for $g \in C(X)$.
(iii) Extract a weak-* convergent subsequence $\mu_{n_j} \to \mu$ in $\mathcal{P}(X)$.
(iv) Verify invariance by $\int_X g \circ f \, d\mu - \int_X g \, d\mu = \lim_j \frac{1}{n_j}\int_X (g \circ f^{n_j} - g) \, d\eta = 0$ for $g \in C(X)$.

For fixed $g \in C(X)$, $\left|\frac{1}{n}\int_X (g \circ f^n - g) \, d\eta\right| \leq 2\|g\|_\infty / n$; weak-* limits are invariant-measure candidates.




Figure 237: Krylov–Bogoliubov: $\mu_n = \frac{1}{n}\sum_{k<n}(f^k)_*\eta \xrightarrow{w^*} \mu$ and $\frac{1}{n}(g \circ f^n - g) \to 0$, hence $f_*\mu = \mu$.

**Theorem 1.7.6.** *Bogoliubov–Krilov Invariant Measure Theorem*
*If $X \neq \varnothing$ is compact metric and $f \in C(X, X)$, then $\exists \mu \in \mathcal{P}(X)$ with $f_*\mu = \mu$.*

*Proof.* Choose $\eta \in \mathcal{P}(X)$ and set $\mu_n := \frac{1}{n}\sum_{k=0}^{n-1}(f^k)_*\eta \in \mathcal{P}(X)$.
By Corollary 6.4.7, some $\mu_{n_j} \to \mu \in \mathcal{P}(X)$ weak-*; equivalently, $\int_X g \, d\mu_{n_j} \to \int_X g \, d\mu \ \forall g \in C(X)$.
For $g \in C(X)$, $\int_X g \circ f \, d\mu_n - \int_X g \, d\mu_n = \frac{1}{n}\int_X (g \circ f^n - g) \, d\eta$, so $\left|\int_X g \circ f \, d\mu_n - \int_X g \, d\mu_n\right| \leq 2\|g\|_\infty / n$.
Passing to $n = n_j \to \infty$, $\int_X g \circ f \, d\mu = \int_X g \, d\mu \ \forall g \in C(X)$. Since compact-metric Borel probabilities are Radon and $C_c(X) = C(X)$, by Theorem 5.2.13, $f_*\mu = \mu$, i.e. $\mu(f^{-1}E) = \mu(E) \ \forall E \in \mathcal{B}(X)$.
*Structural Commentary*: Cesaro pushforwards turn orbit motion into endpoint cancellation.
*Key Insight*: Weak-* limits kill the $1/n$ endpoint error. □

**Proposition 1.7.7.** *Ergodic Measures Are Extreme Invariant Measures*
*Let $X$ be compact metric, $f \in C(X, X)$, $\mathcal{P}(X)$ Borel probabilities on $X$, $\mathcal{I}_f(X) := \{\lambda \in \mathcal{P}(X) : f_*\lambda = \lambda\}$. For $\mu \in \mathcal{I}_f(X)$, the following are equivalent:*

*(i) $\mu \in \mathrm{Ext}(\mathcal{I}_f(X))$;*
*(ii) $f$ is $\mu$-ergodic.*

*Proof.*

(i) $\Rightarrow$ (ii) Assume not (ii): let $A \in \mathcal{B}(X)$, $\mu(A \triangle f^{-1}A) = 0$, $0 < \mu(A) < 1$; $\nu(E) := \frac{\mu(E \cap A)}{\mu(A)}$, $\rho(E) := \frac{\mu(E \cap A^c)}{\mu(A^c)}$.
For $E \in \mathcal{B}(X)$, $\mu(f^{-1}E \cap A) = \mu(f^{-1}E \cap f^{-1}A) = \mu(f^{-1}(E \cap A)) = \mu(E \cap A)$, similarly for $A^c$.
Hence $\nu, \rho \in \mathcal{I}_f(X)$. Since $\mu = \mu(A)\nu + (1 - \mu(A))\rho$, $\nu(A) = 1$, $\rho(A) = 0$; $\mu$ is not extreme.

(*ii*) $\Rightarrow$ (*i*) Let $\mu = \theta\nu + (1-\theta)\rho$ with $0 < \theta < 1$ and $\nu, \rho \in \mathcal{I}_f(X)$.
Since $\theta\nu \le \mu$, by Theorem 3.2.1, $h = \frac{d\nu}{d\mu}$ with $0 \le h \le \frac{1}{\theta}$ $\mu$-a.e.
Then $\int_X h \circ f \, h \, d\mu = \int_X h \circ f \, d\nu = \int_X h \, d\nu = \int_X h^2 \, d\mu$, while $\int_X (h \circ f)^2 \, d\mu = \int_X h^2 \, d\mu$ by $\mu$-invariance.
Thus $\int_X (h \circ f - h)^2 \, d\mu = 0$, so $h \circ f = h$ $\mu$-a.e. By Proposition 1.7.4, $h = c$ $\mu$-a.e.
Since $1 = \nu(X) = \int_X h \, d\mu = c$, $\nu = \mu$, and then $\rho = \mu$.

*Structural Commentary*: Invariant sets split measures; invariant densities test splits.
*Key Insight*: Extreme invariant measures are exactly ergodic measures. □

**Corollary 1.7.8.** *Existence of an Ergodic Invariant Measure*
*Let $X \neq \varnothing$ be compact metric, $f \in C(X, X)$, and $\mathcal{I}_f(X) := \{\mu \in \mathcal{P}(X) : f_*\mu = \mu\}$.*
*Then $\exists \mu \in \mathcal{I}_f(X)$ such that $f$ is $\mu$-ergodic.*

*Proof.* By Theorem 1.7.6, $\mathcal{I}_f(X) \neq \varnothing$. Convexity immediate. If $\mu_\alpha \to \mu$ weak-*, $\mu_\alpha \in \mathcal{I}_f(X)$, then $\mu \in \mathcal{P}(X)$ and $\int_X g \circ f \, d\mu = \lim_\alpha \int_X g \circ f \, d\mu_\alpha = \lim_\alpha \int_X g \, d\mu_\alpha = \int_X g \, d\mu$ $\forall g \in C(X)$.
By Theorem 5.2.13, $f_*\mu = \mu$, so $\mu \in \mathcal{I}_f(X)$. Thus $\mathcal{I}_f(X)$ is weak-* closed and $\|\mu\|_{\text{var}} = 1$ on $\mathcal{I}_f(X)$.
By Corollary 6.4.6, $\mathcal{I}_f(X)$ is weak-* compact.
By Krein–Milman, $\exists$ extreme $\mu \in \mathcal{I}_f(X)$. By Proposition 1.7.7, $\mu$ is ergodic.
*Structural Commentary*: Compactness supplies an extreme invariant measure.
*Key Insight*: Extreme invariant measures are ergodic. □

**Theorem 1.7.9.** *Asymptotic Ergodic Averages*

*Let $T : X \to X$ be measure preserving on probability space $(X, \mathcal{M}, \mu)$, $U_T g := g \circ T$, $A_n := \frac{1}{n} \sum_{k=0}^{n-1} U_T^k$.*
*Then $T$ is ergodic iff $A_n g \to \int_X g \, d\mu$ $\mu$-a.e. $\forall g \in L^1(X, \mu)$. If $0 < \mu(X) < +\infty$, replace $\int_X g \, d\mu$ by $\frac{1}{\mu(X)} \int_X g \, d\mu$.*

*Proof.*
($\Rightarrow$) By Birkhoff's pointwise ergodic theorem, $A_n g \to h$ $\mu$-a.e. for some $h \in L^1(X, \mu)$ with $h \circ T = h$ $\mu$-a.e. and $\int_X h \, d\mu = \int_X g \, d\mu$. Since $T$ is ergodic, Proposition 1.7.4 implies $h = c$ $\mu$-a.e.; by integrating, $c = \int_X g \, d\mu$.
($\Leftarrow$) Let $A \in \mathcal{M}$ with $\mu(A \triangle T^{-1}A) = 0$. For $g = \mathbf{1}_A$, $A_n \mathbf{1}_A = \mathbf{1}_A$ $\mu$-a.e.
By the assumed limit, $\mathbf{1}_A = \mu(A)$ $\mu$-a.e., hence $\mu(A) \in \{0, 1\}$.

*Structural Commentary*: The ergodic limit is the invariant component; ergodicity makes it constant.
*Key Insight*: Time averages equal the space mean exactly when invariants are trivial. □

**Proposition 1.7.10.** *Koopman and Linear Examples*

(*i*) *If $\mu(X) < +\infty$, $T : X \to X$ is measurable, and $U_T g := g \circ T$, then $T$ preserves $\mu$ iff $\|U_T g\|_p = \|g\|_p$ $\forall g \in L^p(X, \mu)$ and $1 \le p < +\infty$.*
(*ii*) *For linear $T : \mathbb{R}^n \to \mathbb{R}^n$, $T_* m_n = m_n$ iff $T \in GL_n(\mathbb{R})$ and $|\det T| = 1$; hence $\operatorname{diag}(2, \frac{1}{2})_* m_2 = m_2$.*

*Proof.*
(*i*) ($\Rightarrow$) If $T$ preserves $\mu$, then by Proposition 1.7.2, $\|U_T g\|_p^p = \int_X |g \circ T|^p \, d\mu = \int_X |g|^p \, d\mu = \|g\|_p^p$.
(*i*) ($\Leftarrow$) If the $L^1$ isometry holds, then for $A \in \mathcal{M}$, $\mu(T^{-1}A) = \|\mathbf{1}_A \circ T\|_1 = \|\mathbf{1}_A\|_1 = \mu(A)$.
(*ii*) ($\Rightarrow$) If $T_* m_n = m_n$ and $T$ is singular, then $m_n(T(\mathbb{R}^n)) = 0$ by Corollary 2.3.6, but $T^{-1}(T(\mathbb{R}^n)) = \mathbb{R}^n$, contradiction. Thus $T \in GL_n(\mathbb{R})$. By Corollary 2.3.4, $m_n(T^{-1}E) = |\det T|^{-1} \, m_n(E)$.
Any $E$ with $0 < m_n(E) < +\infty$ implies $|\det T| = 1$.
(*ii*) ($\Leftarrow$) If $T \in GL_n(\mathbb{R})$ and $|\det T| = 1$, then by Corollary 2.3.4,
$m_n(T^{-1}E) = |\det T|^{-1} \, m_n(E) = m_n(E)$ for all Lebesgue measurable $E$.

*Structural Commentary*: Pullbacks preserve norms or scale by determinant.
*Key Insight*: Measure preservation is norm preservation or unit determinant. □

**Theorem 1.7.11.** *Poincare Recurrence*
*Let $\mu(X) < +\infty$, $T : X \to X$ preserve $\mu$, $A \in \mathcal{M}$.*
*With $R_A := \{x \in A : \#\{n \ge 1 : T^n x \in A\} = +\infty\}$, $\mu(A \setminus R_A) = 0$.*

*Proof.* Set $E := A \cap \bigcap_{n \ge 1} T^{-n}(A^c)$. Then $E \cap T^{-m} A = \varnothing$ for $m \ge 1$.
If $x \in T^{-i}E \cap T^{-j}E$ with $0 \le i < j$, then $T^i x \in E \cap T^{-(j-i)}E \subseteq E \cap T^{-(j-i)}A$, contradiction; hence $(T^{-k}E)_{k \ge 0}$ are disjoint. For $N \ge 0$, by preservation and finite additivity,
$(N+1)\mu(E) = \mu(\bigcup_{k=0}^{N} T^{-k}E) \le \mu(X)$, so $\mu(E) \le \frac{\mu(X)}{N+1}$ $\forall N$; hence $\mu(E) = 0$.
If $x \in A \setminus R_A$ and $r \ge 0$ is its last visit time, then $T^r x \in E$, so $A \setminus R_A \subseteq \bigcup_{r \ge 0} T^{-r}E$ and $\mu(A \setminus R_A) = 0$.

*Structural Commentary*: The nonreturn set is wandering; finite-return points eventually enter it.
*Key Insight*: Finite measure forbids positive-measure wandering sets. □

# 2 Partial Differential Equations

## 2.1 First-Order PDE and Characteristics

## 2.2 Elliptic Equations

## 2.3 Parabolic Equations

## 2.4 Hyperbolic Equations

# 3 Weak Solutions and Evolution Equations

## 3.1 Sobolev Spaces for PDE

Distributions test against smooth compactly supported functions; Sobolev spaces package weak derivatives in normed or Hilbert spaces as a bridge to PDE.

**Definition 3.1.1.** *Test Functions and Distributions*
Let $\Omega \subseteq \mathbb{R}^n$ be open. A *test function* is a smooth function $\varphi \in C_c^\infty(\Omega)$. A *distribution* on $\Omega$ is a continuous linear functional on the test-function space, written $u \in \mathcal{D}'(\Omega)$ and paired as $\langle u, \varphi \rangle$.

**Definition 3.1.2.** *Weak Derivative*
Let $u \in L^1_{\mathrm{loc}}(\Omega)$ and let $\alpha$ be a multi-index. A function $v \in L^1_{\mathrm{loc}}(\Omega)$ is the *weak derivative* $D^\alpha u$ if

$$\int_\Omega u\, D^\alpha \varphi \, dx = (-1)^{|\alpha|} \int_\Omega v\, \varphi \, dx \qquad \text{for every } \varphi \in C_c^\infty(\Omega).$$

**Definition 3.1.3.** *Sobolev Spaces*
For $1 \le p \le \infty$ and an integer $k \ge 0$, the Sobolev space $W^{k,p}(\Omega)$ consists of functions $u \in L^p(\Omega)$ whose weak derivatives $D^\alpha u$ lie in $L^p(\Omega)$ for every $|\alpha| \le k$. The norm is

$$\|u\|_{W^{k,p}} := \sum_{|\alpha| \le k} \|D^\alpha u\|_{L^p}$$

with the usual supremum modification when $p = \infty$.

## 3.2 Weak Solutions and Variational Methods

## 3.3 Evolution Equations

# Part XVIII
# Numerical and Stochastic Computation

## 1 Numerical Analysis

### 1.1 Floating Point and Error Analysis

### 1.2 Nonlinear Equations and Optimization Basics

**Method 1.2.1.** *Newton and Secant Iterations*
To solve $f(x) = 0$:

(i) Newton uses tangent-line zero: $x_n = x_{n-1} - \frac{f(x_{n-1})}{f'(x_{n-1})}$, $n \geq 1$ when $f'(x_{n-1}) \neq 0$.

(ii) Secant uses two-point slope: $x_n = x_{n-1} - \frac{f(x_{n-1})(x_{n-2}-x_{n-1})}{f(x_{n-2})-f(x_{n-1})}$, $n \geq 2$ when the denominator is nonzero.

These define iterates only; local convergence is proved with Taylor's theorem (Theorem 1.1.25).

Figure 238: Newton: tangent-line zero. Secant: two-point line zero.

### 1.3 Interpolation and Approximation

### 1.4 Numerical Differentiation and Quadrature

### 1.5 Numerical Stability and Convergence

## 2 Numerical Differential Systems

### 2.1 Numerical ODEs

Numerical one-step methods replace the integral evolution of an initial-value problem by finitely many vector-field evaluations on a time grid.

**Method 2.1.1.** *Explicit Euler Method*
Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$, $f : U \to \mathbb{R}^n$, $(t_0, x_0) \in U$, and $T > 0$. For $x' = f(t, x)$, $x(t_0) = x_0$, choose $N \in \mathbb{N}$ with $N \geq 1$, set $h := T/N$ and $t_m := t_0 + mh$.

(i) Initialize $x_0^{(h)} := x_0$.

(ii) For $0 \leq m < N$ with $(t_m, x_m^{(h)}) \in U$, compute $x_{m+1}^{(h)} := x_m^{(h)} + hf(t_m, x_m^{(h)})$.

(iii) On each completed mesh interval, define the continuous Euler polygon by $x_h(t) := x_m^{(h)} + (t - t_m) f(t_m, x_m^{(h)})$ for $t_m \leq t \leq t_{m+1}$.

Thus $x_h' = f(t_m, x_m^{(h)})$ on each open mesh interval, while the nodes satisfy the explicit recurrence in (ii).

**Theorem 2.1.2.** *Uniform Error of the Euler Polygon*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C(U, \mathbb{R}^n)$, $(t_0, x_0) \in U$, and choose $T, \delta > 0$ with $V := [t_0, t_0 + T] \times \overline{B}_\delta(x_0) \subseteq U$. Set $M := \max_V \|f\|$, assume $MT \leq \delta$, and assume $\exists L \geq 0$ such that $\|f(t, x) - f(t, y)\| \leq L \|x - y\|$ $\forall (t, x), (t, y) \in V$.*
*Let $\phi \in C^1([t_0, t_0 + T], \mathbb{R}^n)$ satisfy $\phi' = f(t, \phi)$, $\phi(t_0) = x_0$, and $(t, \phi(t)) \in V$ $\forall t \in [t_0, t_0 + T]$.*
*For $N \in \mathbb{N}$ with $N \geq 1$, set $h := T/N$, $t_m := t_0 + mh$, $x_0^{(h)} := x_0$, $x_{m+1}^{(h)} := x_m^{(h)} + hf(t_m, x_m^{(h)})$, and $x_h(t) := x_m^{(h)} + (t - t_m) f(t_m, x_m^{(h)})$ for $t_m \leq t \leq t_{m+1}$.*
*Define $\Delta(h) := \sup\{\|f(s, y) - f(t, x)\| : (s, y), (t, x) \in V,\ |s - t| \leq h,\ \|y - x\| \leq Mh\}$. Then:*

*(i) $(t, x_h(t)) \in V$ $\forall t \in [t_0, t_0 + T]$, and $\|x_h(t) - x_h(s)\| \leq M |t - s|$ $\forall s, t \in [t_m, t_{m+1}]$, $0 \leq m < N$.*

*(ii) With $h = T/N$, as $N \to \infty$, $\Delta(h) \to 0$ and $\|x_h - \phi\|_\infty \leq Te^{LT}\Delta(h)$; hence $x_h \to \phi$ uniformly.*

*(iii) If $\exists L_0 \geq 0$ such that $\|f(t, x) - f(s, x)\| \leq L_0 |t - s|$ $\forall (t, x), (s, x) \in V$, then $\Delta(h) \leq (L_0 + LM)h$ and $\|x_h - \phi\|_\infty \leq Te^{LT}(L_0 + LM)h$.*

*Proof.*

(i) The initial graph point $(t_0, x_0)$ lies in $V$. If $\left\|x_m^{(h)} - x_0\right\| \le mhM$, then $(t_m, x_m^{(h)}) \in V$ and $\left\|x_{m+1}^{(h)} - x_0\right\| \le \left\|x_m^{(h)} - x_0\right\| + h\left\|f(t_m, x_m^{(h)})\right\| \le (m+1)hM$.
Thus induction gives $\left\|x_m^{(h)} - x_0\right\| \le mhM$. For $t \in [t_m, t_{m+1}]$, $\|x_h(t) - x_0\| \le M(t - t_0) \le MT \le \delta$. Hence the graph lies in $V$, and the affine slope gives the meshwise Lipschitz bound.

(ii) Let $\tau_h(t) := t_m$ for $t \in [t_m, t_{m+1})$. The Euler polygon satisfies $x_h(t) = x_0 + \int_{t_0}^t f(\tau_h(s), x_h(\tau_h(s)))\, ds$. Uniform continuity of $f|_V$ gives $\Delta(h) \to 0$. With $e(t) := \|x_h(t) - \phi(t)\|$, item (i) gives $\|x_h(s) - x_h(\tau_h(s))\| \le Mh$, so $e(t) \le \int_{t_0}^t (Le(s) + \Delta(h))\, ds$.
By Lemma 1.1.20(ii), $e(t) \le (t - t_0)e^{L(t-t_0)}\Delta(h) \le Te^{LT}\Delta(h)$.

(iii) For pairs in the definition of $\Delta(h)$, $\|f(s,y) - f(t,x)\| \le L\|y - x\| + L_0|s - t| \le (LM + L_0)h$. Combine this with (ii).

*Structural Commentary*: Consistency supplies $\Delta(h)$; Gronwall propagates the local defect.
*Key Insight*: Uniform continuity is enough for convergence; time-Lipschitz control gives first order. □

**Remark 2.1.3.** *Euler Step-Doubling Control*
From a node $(t_m, x_m)$, compare the full-step prediction $z_h := x_m + hf(t_m, x_m)$ with the two-half-step prediction $z_{1/2} := x_m + \frac{h}{2}f(t_m, x_m)$, $z_{2/2} := z_{1/2} + \frac{h}{2}f(t_m + h/2, z_{1/2})$. If $\left\|z_h - z_{2/2}\right\|$ exceeds a prescribed tolerance, reduce $h$ and repeat the step. This is a practical error indicator rather than an a priori error bound.

**Method 2.1.4.** *Classical Fourth-Order Runge–Kutta Method*
Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$, $f : U \to \mathbb{R}^n$, $h > 0$, and $(t_m, x_m) \in U$, where $x_m$ approximates a solution of $x' = f(t, x)$ at $t_m$. Set $t_{m+1} := t_m + h$ and assume every displayed argument of $f$ below lies in $U$.

(i) Compute $k_{1,m} := f(t_m, x_m)$ and $k_{2,m} := f(t_m + h/2, x_m + hk_{1,m}/2)$.
(ii) Compute $k_{3,m} := f(t_m + h/2, x_m + hk_{2,m}/2)$ and $k_{4,m} := f(t_m + h, x_m + hk_{3,m})$.
(iii) Update $x_{m+1} := x_m + \frac{h}{6}(k_{1,m} + 2k_{2,m} + 2k_{3,m} + k_{4,m})$.

The four stages sample two midpoint predictions and one endpoint prediction before forming the weighted increment.

**Method 2.1.5.** *Heun's Improved Euler Method*
Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$, $f : U \to \mathbb{R}^n$, $h > 0$, and $(t_m, x_m) \in U$, where $x_m$ approximates a solution of $x' = f(t, x)$ at $t_m$. Set $t_{m+1} := t_m + h$ and assume the predicted endpoint used below lies in $U$.

(i) Predict the endpoint by $y_m := x_m + hf(t_m, x_m)$.
(ii) Correct with the averaged slopes: $x_{m+1} := x_m + \frac{h}{2}[f(t_m, x_m) + f(t_{m+1}, y_m)]$.

Thus Heun replaces Euler's single left-endpoint slope by the average of the initial slope and its predicted endpoint value.

**Proposition 2.1.6.** *Heun-Type One-Step Error*
*Let $U \subseteq \mathbb{R} \times \mathbb{R}^n$ be open, $f \in C^2(U, \mathbb{R}^n)$, $I \subseteq \mathbb{R}$ an open interval, $t_0 \in I$, and $\phi \in C^1(I, \mathbb{R}^n)$ satisfy $(t, \phi(t)) \in U$ and $\phi'(t) = f(t, \phi(t))$ $\forall t \in I$, with $\phi(t_0) = x_0$.*
*Set $f_0 := f(t_0, x_0)$. Then $\exists h_* > 0$ such that, $\forall h \in (0, h_*)$, setting $t_1 := t_0 + h$ and $y_0 := x_0 + hf_0$ gives $t_1 \in I$ and $(t_1, y_0), (t_1, x_0), (t_0, y_0) \in U$. As $h \downarrow 0$:*

*(i) For $H_h := x_0 + \frac{h}{2}[f(t_0, x_0) + f(t_1, y_0)]$, $\|\phi(t_1) - H_h\| = O(h^3)$.*
*(ii) For $\widetilde{H}_h := x_0 + \frac{h}{2}[f(t_1, x_0) + f(t_0, y_0)]$, $\left\|\phi(t_1) - \widetilde{H}_h\right\| = O(h^3)$.*

*Proof.* Set $G_0 := \partial_t f(t_0, x_0) + D_x f(t_0, x_0) f_0$. By Lemma 1.1.19, $\phi \in C^3$ locally, so $\phi(t_1) = x_0 + hf_0 + \frac{h^2}{2}G_0 + O(h^3)$.

(i) Taylor expansion of $f$ at $(t_0, x_0)$ gives $f(t_1, y_0) = f_0 + h\partial_t f(t_0, x_0) + hD_x f(t_0, x_0) f_0 + O(h^2) = f_0 + hG_0 + O(h^2)$.
Hence $H_h = x_0 + hf_0 + h^2 G_0/2 + O(h^3)$, which matches the expansion of $\phi(t_1)$.

(ii) Separately, $f(t_1, x_0) = f_0 + h\partial_t f(t_0, x_0) + O(h^2)$ and $f(t_0, y_0) = f_0 + hD_x f(t_0, x_0) f_0 + O(h^2)$.
Thus $\widetilde{H}_h = x_0 + hf_0 + h^2 G_0/2 + O(h^3)$, proving the second estimate.

*Structural Commentary*: Expand the exact flow and both predicted slopes through second order.
*Key Insight*: Averaging reconstructs the quadratic term of the exact solution. □

**Example 2.1.7.** *Euler and RK4 for Exponential Growth*
For $x' = x$, $x(0) = 1$, and $h = 0.1$, Euler gives $x_m = (1 + h)^m$, hence $x_{10} = 1.1^{10} \approx 2.5937424601$.
Classical RK4 gives $x_{m+1} = R(h)x_m$ with $R(h) = 1 + h + h^2/2 + h^3/6 + h^4/24$, so $x_{10} = R(0.1)^{10} \approx 2.7182797441$.

Since $x(1) = e \approx 2.7182818285$, the respective absolute errors at $t = 1$ are approximately $1.2454 \times 10^{-1}$ and $2.0843 \times 10^{-6}$.

## 2.2 Geometric Numerical Integration

## 2.3 Finite Difference PDE Methods

## 2.4 Finite Element Methods

## 2.5 Numerical SDEs

## 2.6 Computational Stability Across ODE/PDE/SDE

# 3 Monte Carlo and Stochastic Computation

## 3.1 Monte Carlo Foundations

Monte Carlo methods turn integrals into averages of simulated random variables. This is especially useful when the integral is high-dimensional or when the target density is known only up to a normalizing constant.

**Method 3.1.1.** *Basic Monte Carlo Integration*

(i) Let $I = \int h(x)f(x)\,dx$, where $f$ is a probability density.

(ii) For IID draws $X_1, \dots, X_N \sim f$, define $\widehat{I}_N = N^{-1} \sum_{i=1}^{N} h(X_i)$; then $\widehat{I}_N \xrightarrow{p} I$ by the law of large numbers.

(iii) With $Y_i = h(X_i)$, set $s^2 = (N-1)^{-1} \sum_{i=1}^{N} (Y_i - \widehat{I}_N)^2$ and $\widehat{\mathrm{se}} = \frac{s}{\sqrt{N}}$.

(iv) An approximate $1-\alpha$ Monte Carlo interval is $\widehat{I}_N \pm z_{\frac{\alpha}{2}} \widehat{\mathrm{se}}$.

**Example 3.1.2.** *Monte Carlo Integration from the LLN*
Let $X_1, \dots, X_n$ be iid Uniform$(0,1)$ and $f \in L^1([0,1])$.
The estimator $\widehat{I}_n(f) := n^{-1} \sum_{i=1}^{n} f(X_i)$ satisfies $\widehat{I}_n(f) \xrightarrow{\mathbb{P}} \mathbb{E}f(X_1) = \int_0^1 f(x)\,dx$.
For $f(x) = (2\pi)^{-\frac{1}{2}} e^{-\frac{x^2}{2}}$ on $[0,1]$, a 1000-point estimate near 0.3417 is consistent with $\int_0^1 f(x)\,dx \approx 0.3413$.

**Example 3.1.3.** *Integrating on an Interval*
For $X \sim$ Uniform$(a,b)$, $I = \int_a^b h(x)\,dx = (b-a)\mathbb{E}[h(X)]$.
Given IID draws $X_1, \dots, X_N \sim$ Uniform$(a,b)$, estimate $I$ by $\widehat{I}_N = (b-a)N^{-1} \sum_{i=1}^{N} h(X_i)$.

**Example 3.1.4.** *Estimating a Normal CDF*
Let $Z \sim N(0,1)$ and $\Phi(x) = \mathbb{P}(Z \le x)$. For IID draws $Z_1, \dots, Z_N \sim N(0,1)$, use $h(s) = \mathbf{1}\{s \le x\}$ and $\widehat{\Phi}(x) = N^{-1} \sum_{i=1}^{N} \mathbf{1}\{Z_i \le x\}$. Thus $\widehat{\Phi}(x)$ is the empirical fraction of draws not exceeding $x$.

**Method 3.1.5.** *Inverse-Transform Generation*

(i) *Exponential*: if $U \sim$ Uniform$(0,1)$ and $\lambda > 0$, then $T = -\lambda^{-1}\log(1-U) \sim \mathrm{Exp}(\lambda)$. (or $T = -\lambda^{-1}\log U$)

(ii) *Weibull*: if $F(x) = 1 - \exp(-(\frac{x}{\alpha})^{\beta})$ for $x \ge 0$ and $\alpha, \beta > 0$, then $F^{-1}(u) = \alpha[-\log(1-u)]^{\frac{1}{\beta}}$ and $W = \alpha[-\log(1-U)]^{\frac{1}{\beta}} \sim F$.

(iii) *Integer-valued inverse CDF*: if $F$ is integer-valued and $U \sim$ Uniform$(0,1)$, set $Y = \inf\{k \in \mathbb{Z} : F(k) \ge U\}$. Equivalently $F(k-1) < U \le F(k)$. Then $F_Y = F$.

*Key Insight*: The simulation problem is reduced to solving $u = F(x)$ for $x$.

**Method 3.1.6.** *Rejection Sampling*
Let $f$ be a target density on an interval $[a,b]$, possibly with infinite endpoints.
Choose an envelope $M$ such that $M(x) \ge f(x)$ on $[a,b]$ and $A := \int_a^b M(t)\,dt < \infty$. Let $m(x) := \frac{M(x)}{A}$.

(i) Generate $T$ with density $m$.

(ii) Generate $U \sim$ Uniform$(0,1)$ independently of $T$.

(iii) Accept $X = T$ if $U \le \frac{f(T)}{M(T)}$; otherwise reject $T$ and repeat.

*Key Insight*: Sample under an easy envelope and keep only points that fall below the target curve.




Figure 239: Rejection sample accepts when the vertical draw lies below $f(T)$.

**Proposition 3.1.7.** *Rejection Sampling Correctness*
*Let $f$ be a density, let $M \geq f$ with $A := \int M(x)\,dx < \infty$, and set $m := \frac{M}{A}$. If $T \sim m$, $U \sim \text{Uniform}(0,1)$ independently, and $\mathcal{A} := \{U \leq \frac{f(T)}{M(T)}\}$, then $\mathbb{P}(\mathcal{A}) = \frac{1}{A}$ and $T \mid \mathcal{A}$ has density $f$.*

*Proof.* For measurable $B$, $\mathbb{P}(T \in B, \mathcal{A}) = \int_B \frac{m(x)f(x)}{M(x)}\,dx = A^{-1}\int_B f(x)\,dx$.
Taking $B = \mathbb{R}$ gives $\mathbb{P}(\mathcal{A}) = A^{-1}$, and hence $\mathbb{P}(T \in B \mid \mathcal{A}) = \int_B f(x)\,dx$.
*Structural Commentary*: Acceptance reweights the envelope $m$ back to $f$.
*Key Insight*: Conditional on acceptance, the normalized accepted density is exactly the target. □

## 3.2 Variance Reduction

**Method 3.2.1.** *Importance Sampling*

(i) For $I = \int h(x)f(x)\,dx$, choose a tractable proposal density $g$ with $g(x) > 0$ wherever $h(x)f(x) \neq 0$.
(ii) Then $I = \int h(x)f(x)g(x)^{-1}g(x)\,dx = \mathbb{E}_g\left[\frac{h(X)f(X)}{g(X)}\right]$.
(iii) For IID draws $X_1, \ldots, X_N \sim g$, use $\widehat{I}_N = N^{-1}\sum_{i=1}^{N}\frac{h(X_i)f(X_i)}{g(X_i)}$.

**Remark 3.2.2.** *Choosing the Proposal*
The importance weights have large, possibly infinite, variance when $g$ is too small on regions where $|h|\,f$ is large. Choose $g$ to approximate the shape of $|h|\,f$ while giving it tails at least as thick as the target integrand.

**Theorem 3.2.3.** *Optimal Importance Density*
*Let $I = \int h(x)f(x)\,dx$ and $A = \int |h(x)|\,f(x)\,dx \in (0,\infty)$.*
*For proposal densities $g$ with $g(x) > 0$ wherever $|h(x)|\,f(x) > 0$, define $W_g = \frac{h(X)f(X)}{g(X)}$ for $X \sim g$.*
*The variance $\text{Var}_g(W_g)$ is minimised by $g^*(x) = \frac{|h(x)|f(x)}{A}$, and the minimum variance is $A^2 - I^2$.*

*Proof.* For each admissible $g$, $\mathbb{E}_g W_g = I$, so $\text{Var}_g(W_g) = M(g) - I^2$, where $M(g) = \mathbb{E}_g(W_g^2) = \int \frac{h(x)^2 f(x)^2}{g(x)}\,dx$.
For any admissible $g$, Cauchy-Schwarz gives $A^2 = (\int |h(x)|\,f(x)g(x)^{-\frac{1}{2}}g(x)^{\frac{1}{2}}\,dx)^2 \leq M(g)\int g(x)\,dx = M(g)$.
Equality holds precisely when $|h(x)|\,f(x)g(x)^{-\frac{1}{2}} = c\,g(x)^{\frac{1}{2}}$ a.e., equivalently $g(x) \propto |h(x)|\,f(x)$.
Normalization gives $g^*(x) = \frac{|h(x)|f(x)}{A}$, hence $M(g) \geq M(g^*) = A^2$ and $\text{Var}_g(W_g) \geq A^2 - I^2 = \text{Var}_{g^*}(W_{g^*})$.
*Structural Commentary*: Fixed mean reduces variance to second-moment minimization.
*Key Insight*: Cauchy-Schwarz is sharp exactly when the proposal is proportional to $|h|\,f$. □

## 3.3 Markov Chain Monte Carlo

Monte Carlo integration in Method 3.1.1 uses independent draws from the target density. Markov chain Monte Carlo instead constructs a Markov chain $X_0, X_1, \ldots$ whose stationary density is the target $f$. When the chain mixes well, sample averages from the chain are used as approximations to expectations under $f$.

**Method 3.3.1.** *Metropolis-Hastings*
Let $f$ be a target density available up to proportionality, and let $q(\cdot \mid x)$ be a tractable proposal density. Choose $X_0$ arbitrarily; given $X_i = x$, define $X_{i+1}$ as follows.

(i) Draw $Y \sim q(\cdot \mid x)$.
(ii) Set $r(x,Y) = \min\left\{\frac{f(Y)q(x|Y)}{f(x)q(Y|x)}, 1\right\}$.
(iii) Set $X_{i+1} = Y$ with probability $r(x,Y)$, and set $X_{i+1} = x$ otherwise.

*Key Insight*: Only ratios of $f$ enter $r$, so unknown normalising constants cancel.

**Proposition 3.3.2.** *Detailed Balance for Metropolis-Hastings*
*Let* $\Phi(x,y) = f(x)q(y \mid x)$ *and assume* $\Phi(x,y) > 0$ *iff* $\Phi(y,x) > 0$.
*For* $x \neq y$, *define* $\alpha(x,y) = \min\left\{\frac{\Phi(y,x)}{\Phi(x,y)}, 1\right\}$ *when* $\Phi(x,y) > 0$ *and* $\alpha(x,y) = 0$ *otherwise.*
*Set* $p(x,y) = q(y \mid x)\alpha(x,y)$. *Then* $f(x)p(x,y) = f(y)p(y,x)$ *for* $x \neq y$.

*Proof.* If $\Phi(x,y) = 0$, the support condition gives $\Phi(y,x) = 0$, hence $p(x,y) = p(y,x) = 0$.
If $\Phi(x,y) > 0$, then $f(x)p(x,y) = \Phi(x,y)\min\left\{\frac{\Phi(y,x)}{\Phi(x,y)}, 1\right\} = \min\{\Phi(x,y), \Phi(y,x)\}$.
Interchanging $x, y$ gives the same expression for $f(y)p(y,x)$, QED.
*Structural Commentary*: The acceptance rule symmetrizes the weighted proposal flow.
*Key Insight*: Both directions reduce to the same minimum of the two proposal fluxes. □

**Method 3.3.3.** *Random-Walk Metropolis-Hastings*

(i) Use the proposal $Y = X_i + \varepsilon_i$, where $\varepsilon_i$ has a symmetric density.
(ii) Then $q(y \mid x) = q(x \mid y)$, so the acceptance probability is $r(x,y) = \min\left\{\frac{f(y)}{f(x)}, 1\right\}$.
(iii) For $\varepsilon_i \sim N(0, b^2)$, small $b$ gives slow local moves, large $b$ gives frequent rejection. Tune $b$ for adequate mixing.

**Example 3.3.4.** *Cauchy Target*
For Cauchy target $f(x) = \frac{1}{\pi(1+x^2)}$ and symmetric random-walk proposal, $r(x,y) = \min\left\{\frac{1+x^2}{1+y^2}, 1\right\}$.
The proposal scale changes mixing efficiency, not the stationary target.

**Method 3.3.5.** *Independence Metropolis-Hastings*

(i) Use a state-independent proposal $Y \sim g$, so $q(y \mid x) = g(y)$.
(ii) The acceptance probability is $r(x,y) = \min\{\frac{f(y)g(x)}{f(x)g(y)}, 1\}$.
(iii) Choose $g$ as a tractable approximation to the target with sufficient mass on important regions.

**Method 3.3.6.** *Gibbs Sampling*

(i) Let $(X,Y)$ have joint density $f_{X,Y}$ with tractable conditionals $f_{X|Y}$ and $f_{Y|X}$.
(ii) Starting from $(X_0, Y_0)$, iterate $X_{n+1} \sim f_{X|Y}(\cdot \mid Y_n)$ and $Y_{n+1} \sim f_{Y|X}(\cdot \mid X_{n+1})$.
(iii) In higher dimension, update one coordinate or block at a time from its full conditional.

**Method 3.3.7.** *Metropolis within Gibbs*

(i) If $f_{X_j|X_{-j}}$ is not directly simulable, update $X_j$ by Metropolis-Hastings conditional on $X_{-j}$.
(ii) In the bivariate case, propose $Z \sim q(\cdot \mid X_n)$ and accept $X_{n+1} = Z$ with probability $\min\{\frac{f(Z,Y_n)q(X_n|Z)}{f(X_n,Y_n)q(Z|X_n)}, 1\}$.
(iii) Then update $Y$ analogously, conditional on the newly updated $X$.

## 3.4 Sequential Monte Carlo

## 3.5 Rare-Event Simulation

**Example 3.5.1.** *Rare Tail Probability*
For $Z \sim N(0,1)$, write $I = \mathbb{P}(Z > 3) = \int \mathbf{1}\{x > 3\} f(x)\,dx$.

If $g$ is centered near the right tail and $X_1, \ldots, X_N \sim g$, estimate $I$ by $\widehat{I}_N = N^{-1}\sum_{i=1}^{N} \frac{\mathbf{1}\{X_i > 3\} f(X_i)}{g(X_i)}$.

## 3.6 Stochastic Optimization

**Definition 3.6.1.** *Online Estimator*
Let $(X_t)_{t\geq 1}$ be observations in $\mathcal{X}$, let $\widehat{\theta}_0 \in \Theta \subseteq \mathbb{R}^d$, let $(a_t)_{t\geq 0}$ be positive step sizes, and let $H : \Theta \times \mathcal{X} \to \mathbb{R}^d$.
$\widehat{\theta}_{t+1} := \widehat{\theta}_t + a_t H(\widehat{\theta}_t, X_{t+1})$ for $t \geq 0$.
*Structural Commentary*: $(\widehat{\theta}_t)$ is adapted to $\mathcal{F}_t := \sigma(\widehat{\theta}_0, X_1, \ldots, X_t)$.
*Key Insight*: Under stationarity, analyze the recursion through $h(\theta) := \mathbb{E}H(\theta, X)$.

**Method 3.6.2.** *Minibatch Stochastic Gradient Descent*
Let $J_n(\theta) = n^{-1}\sum_{i=1}^{n} L(f_\theta(x_i), y_i)$, $|B_k| = b$, and $\eta_k > 0$.

(i) Sample $B_k$ uniformly.
(ii) Compute $g_k = b^{-1}\sum_{i\in B_k} \nabla_\theta L(f_{\theta_k}(x_i), y_i)$.
(iii) Update $\theta_{k+1} = \theta_k - \eta_k g_k$.

Uniform sampling gives $\mathbb{E}[g_k \mid \theta_k] = \nabla J_n(\theta_k)$; larger $b$ trades lower noise for higher cost.

**Definition 3.6.3.** *Momentum and Adaptive Methods*
Let $g_k$ denote the stochastic gradient at $\theta_k$; $\odot$, division, and square roots are coordinatewise.

(i) *Momentum*: $v_k = \mu v_{k-1} + g_k$ and $\theta_{k+1} = \theta_k - \eta_k v_k$, with $0 \le \mu < 1$.
(ii) *AdaGrad*: $r_k = r_{k-1} + g_k \odot g_k$ and $\theta_{k+1} = \theta_k - \frac{\eta_k g_k}{\sqrt{r_k}+\varepsilon}$.
(iii) *RMSProp*: $r_k = \rho r_{k-1} + (1-\rho) g_k \odot g_k$ and $\theta_{k+1} = \theta_k - \frac{\eta_k g_k}{\sqrt{r_k}+\varepsilon}$, with $0 < \rho < 1$.
(iv) *Adam*: update $(m_k, r_k)$ as exponential averages of $(g_k, g_k \odot g_k)$,
set $(\widehat{m}_k, \widehat{r}_k) = (\frac{m_k}{1-\beta_1^k}, \frac{r_k}{1-\beta_2^k})$, and $\theta_{k+1} = \theta_k - \frac{\eta_k \widehat{m}_k}{\sqrt{\widehat{r}_k}+\varepsilon}$.

# Part XIX
# Statistical Inference and Time Series

Statistical inference turns probabilistic models into procedures for learning from data. The probability, measure-theoretic, stochastic-process, optimization, and Monte Carlo material developed in Parts 11–14 is treated here as prerequisite structure.

## 1 Statistical Models and Experiments

### 1.1 Statistical Experiments and Likelihoods

A statistical problem starts with observable data and a family of probability laws that could have generated it. The model is not the data itself; it is the mathematical language in which uncertainty about the data-generating mechanism is expressed.

**Definition 1.1.1.** *Statistical Experiment and Model*

(i) A *statistical experiment* is $(\mathcal{X}, \mathcal{A}, \mathcal{P})$, where $(\mathcal{X}, \mathcal{A})$ is measurable and $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$ is a family of probability measures on $(\mathcal{X}, \mathcal{A})$.
(ii) $\Theta$ is the *parameter space*, $X \in \mathcal{X}$ is the *sample*, and $\mathcal{P}$ is the *statistical model*.
(iii) The model is *well specified* if the true law $P_0 \in \mathcal{P}$.
(iv) The model is *misspecified* if $P_0 \notin \mathcal{P}$.

**Definition 1.1.2.** *Identifiability and Functionals*

(i) The parameterization is *identifiable* if $P_\theta = P_{\theta'}$ implies $\theta = \theta'$.
(ii) A *statistical functional* is a map $T : \mathcal{P} \to \mathcal{T}$, such as $T(P) = \int x\, dP(x)$ or $T(P) = \text{median}(P)$.
(iii) In a parametric model, a *target parameter* may be a function $\psi(\theta)$ rather than the full parameter $\theta$.

*Key Insight*: Estimation should target a law or functional that the experiment can actually distinguish.

**Definition 1.1.3.** *Dominated Models, Likelihood, Score, and Information*

(i) A model $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$ is *dominated* by $\mu$ if $P_\theta \ll \mu$ for all $\theta \in \Theta$; write $p_\theta := \frac{dP_\theta}{d\mu}$.
(ii) Given $x$, the *likelihood* and *log-likelihood* are $L_x(\theta) := p_\theta(x)$ and $\ell_x(\theta) := \log p_\theta(x)$ when $p_\theta(x) > 0$.
(iii) If $\Theta \subseteq \mathbb{R}^d$ and differentiation is valid, the *score* is $S_\theta(X) := \nabla_\theta \ell_X(\theta)$.
(iv) The *Fisher information* is $I(\theta) := \mathbb{E}_\theta[S_\theta(X) S_\theta(X)^\top]$.

**Definition 1.1.4.** *Maximum Likelihood Estimator*
For likelihood $L_X(\theta)$ and log-likelihood $\ell_X(\theta)$, a *maximum likelihood estimator* is any measurable $\widehat{\theta}_n(X) \in \arg\max_{\theta \in \Theta} L_X(\theta) = \arg\max_{\theta \in \Theta} \ell_X(\theta)$; equivalently, $\widehat{\theta}_n \in \arg\min_{\theta \in \Theta} [-\ell_X(\theta)]$.

**Example 1.1.5.** *Bernoulli and Normal Likelihoods*
Let $X_1, \dots, X_n$ be independent observations.

(i) If $X_i \sim \text{Bernoulli}(p)$, then $L_x(p) = p^s (1-p)^{n-s}$, where $s := \sum_{i=1}^n x_i$.

(ii) If $X_i \sim N(\mu, \sigma^2)$ and $\sigma^2$ is known, then $\ell_x(\mu) = -\frac{n}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}\sum_{i=1}^n (x_i - \mu)^2$, maximized at $\widehat{\mu} = \overline{x}$.

**Example 1.1.6.** *Multinomial Cell Probabilities*
Let $X = (X_1, \dots, X_m)$ have total count $n$.

(i) If $X \sim \text{Multinomial}(n; p_1, \dots, p_m)$ and $\sum_{j=1}^m p_j = 1$, then $L_x(p) \propto \prod_{j=1}^m p_j^{x_j}$ and $\widehat{p}_j = \frac{x_j}{n}$.

(ii) If $p_j = p_j(\theta)$, then $L_x(\theta) \propto \prod_{j=1}^m p_j(\theta)^{x_j}$ and $\widehat{\theta} \in \arg\max_\theta \sum_{j=1}^m x_j \log p_j(\theta)$.

**Remark 1.1.7.** *Likelihood as Relative Evidence*
For fixed data $x$, the likelihood is a function of $\theta$, not a probability distribution on $\Theta$.
Ratios $\frac{L_x(\theta_1)}{L_x(\theta_0)}$ compare how well two model laws explain the same observation.

## 1.2 Sufficiency, Exponential Families, and Local Asymptotics

The shape of a model determines which summaries of the data preserve information and which approximations become valid in large samples.

**Definition 1.2.1.** *Statistics, Sufficiency, and Ancillarity*

(i) A *statistic* is a measurable map $T : \mathcal{X} \to \mathcal{T}$.
(ii) $T$ is *sufficient* if $\mathcal{L}_\theta(X \mid T)$ is independent of $\theta$.
(iii) $T$ is *ancillary* if $\mathcal{L}_\theta(T)$ is independent of $\theta$.

**Theorem 1.2.2.** *Factorization Criterion*
*Let $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$ be dominated by $\mu$, with $p_\theta := \frac{dP_\theta}{d\mu}$. A statistic $T : \mathcal{X} \to \mathcal{T}$ is sufficient iff there exist measurable $g_\theta : \mathcal{T} \to [0,\infty)$ and $h : \mathcal{X} \to [0,\infty)$ such that $p_\theta(x) = g_\theta(T(x))h(x)$ $\mu$-a.e. for all $\theta$.*

*Proof.*
($\Rightarrow$) If $T$ is sufficient, write $\nu_\theta := \mathcal{L}_\theta(T)$ and choose a $\theta$-free kernel $K_t = \mathcal{L}_\theta(X \mid T = t)$, so $P_\theta(A) = \int K_t(A)\nu_\theta(dt)$. Choosing densities for $\nu_\theta$ and for $K_t$ on each fiber gives $p_\theta(x) = g_\theta(T(x))h(x)$.
($\Leftarrow$) If $p_\theta = g_\theta \circ T\, h$, then on the fiber $T = t$ the factor $g_\theta(t)$ cancels: for $c(t) := \int_{T^{-1}\{t\}} h\, d\mu_t > 0$, $\mathbb{P}_\theta(X \in A \mid T = t) = c(t)^{-1} \int_{A \cap T^{-1}\{t\}} h\, d\mu_t$, independent of $\theta$.

*Structural Commentary*: Factorisation separates parameter information $g_\theta(T)$ from residual shape $h$.
*Key Insight*: Sufficiency keeps exactly the likelihood's $\theta$-dependent part. □

**Example 1.2.3.** *Bernoulli and Normal Sufficient Statistics*
Let $X_1, \dots, X_n$ be independent and let $x = (x_1, \dots, x_n)$.

(i) If $X_i \sim \text{Bernoulli}(\theta)$, then $p_\theta(x) = \theta^s(1-\theta)^{n-s}$ with $s := \sum_i x_i$.
Hence $T = \sum_i X_i$ is sufficient for $\theta$, and $\mathcal{L}_\theta(X \mid T = t)$ is uniform on $\{x : \sum_i x_i = t\}$.

(ii) If $X_i \sim N(\mu, \sigma^2)$ with $(\mu, \sigma^2)$ unknown, then $p_{\mu,\sigma^2}(x) \propto (\sigma^2)^{-\frac{n}{2}} \exp\{-\frac{\sum_i x_i^2 - 2\mu \sum_i x_i + n\mu^2}{2\sigma^2}\}$.
Hence $T = (\sum_i X_i, \sum_i X_i^2)$ is sufficient for $(\mu, \sigma^2)$.

**Corollary 1.2.4.** *Likelihood Estimators Use Sufficient Statistics*
*Let $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$ be dominated with densities $p_\theta$, and let $T$ be sufficient with $p_\theta(x) = g_\theta(T(x))h(x)$. If $a(t) \in \arg\max_{\theta \in \Theta} g_\theta(t)$ exists, then $\widehat{\theta}(X) := a(T(X))$ is an MLE.*

*Proof.* For fixed $x$, $L_x(\theta) = p_\theta(x) = h(x)g_\theta(T(x))$. If $h(x) > 0$, then $\arg\max_\theta L_x(\theta) = \arg\max_\theta g_\theta(T(x))$.
If $h(x) = 0$, then $L_x(\theta) = 0$ for all $\theta$. Thus an MLE may be chosen as $a(T(x))$.
*Structural Commentary*: The $\theta$-free factor $h(x)$ is irrelevant to maximization.
*Key Insight*: MLEs can depend on data only through a sufficient statistic. □

**Theorem 1.2.5.** *Rao–Blackwell Improvement*
*Let $T$ be sufficient and let $\widehat{\psi} \in L^2(P_\theta)$ estimate a real target $\psi(\theta)$. Choose the $\theta$-free version $\widetilde{\psi}(T) := \mathbb{E}_\theta(\widehat{\psi} \mid T)$. Then $\mathbb{E}_\theta\widetilde{\psi} = \mathbb{E}_\theta\widehat{\psi}$ and $\mathbb{E}_\theta(\widetilde{\psi} - \psi(\theta))^2 \le \mathbb{E}_\theta(\widehat{\psi} - \psi(\theta))^2$ for all $\theta$, with equality iff $\widehat{\psi} = \widetilde{\psi}$ $P_\theta$-a.s.*

*Proof.* Sufficiency makes $\mathcal{L}_\theta(X \mid T)$ independent of $\theta$, so $\widetilde{\psi}$ is a $T$-statistic. Also $\mathbb{E}_\theta\widetilde{\psi} = \mathbb{E}_\theta\widehat{\psi}$.
Since $\mathbb{E}_\theta(\widehat{\psi} - \widetilde{\psi} \mid T) = 0$ and $\widetilde{\psi} - \psi(\theta)$ is $T$-measurable, $\mathbb{E}_\theta[(\widehat{\psi} - \widetilde{\psi})(\widetilde{\psi} - \psi(\theta))] = 0$.
Hence $\mathbb{E}_\theta(\widehat{\psi} - \psi(\theta))^2 = \mathbb{E}_\theta(\widehat{\psi} - \widetilde{\psi})^2 + \mathbb{E}_\theta(\widetilde{\psi} - \psi(\theta))^2 \ge \mathbb{E}_\theta(\widetilde{\psi} - \psi(\theta))^2$, with equality iff $\mathbb{E}_\theta(\widehat{\psi} - \widetilde{\psi})^2 = 0$.
*Structural Commentary*: Rao–Blackwellization projects an estimator onto the $\sigma(T)$-measurable statistics.
*Key Insight*: Conditioning on a sufficient statistic preserves bias and removes residual noise. □

**Definition 1.2.6.** *Exponential Family*

(i) A dominated model $\{P_\eta : \eta \in \mathcal{N}\}$ is a *$k$-parameter exponential family* if $\mathcal{N} \subseteq \mathbb{R}^k$ and $p_\eta(x) = \exp\{\eta^\top T(x) - A(\eta)\}h(x)$.
(ii) $\eta$ is the *natural parameter*, $T(X)$ is the *natural statistic*, and $h$ is the carrier density.
(iii) $A(\eta)$ is the *log-partition function*: $A(\eta) = \log \int \exp\{\eta^\top T(x)\}h(x)\, d\mu(x)$ when finite.

**Proposition 1.2.7.** *Moment Identities in Full Exponential Families*
*Let $p_\eta(x) = \exp\{\eta^\top T(x) - A(\eta)\}h(x)$ be a full exponential family with open natural parameter space, and assume differentiation under the integral is valid. Then $\nabla A(\eta) = \mathbb{E}_\eta T(X)$ and $\nabla^2 A(\eta) = \text{Cov}_\eta(T(X))$.*

*Proof.* From $1 = \int p_\eta \, d\mu$, differentiation gives $0 = \int (T - \nabla A(\eta)) p_\eta \, d\mu$, hence $\nabla A(\eta) = \mathbb{E}_\eta T(X)$.
For coordinates $i, j$, differentiating $\mathbb{E}_\eta T_i = \int T_i p_\eta \, d\mu$ gives $\partial_j \mathbb{E}_\eta T_i = \mathbb{E}_\eta[T_i(T_j - \partial_j A(\eta))] = \mathrm{Cov}_\eta(T_i, T_j)$, QED.
*Structural Commentary*: The log-partition function cumulatively encodes the natural statistic.
*Key Insight*: First and second derivatives of $A$ give mean and covariance. □

**Definition 1.2.8.** *Local Asymptotic Normality*
For experiments $\{P_\theta^{(n)} : \theta \in \Theta \subseteq \mathbb{R}^d\}$, set $\theta_{n,h} := \theta_0 + \frac{h}{\sqrt{n}}$.

(i) The sequence is *locally asymptotically normal* at $\theta_0$ with information $I(\theta_0)$ if,
for each fixed $h \in \mathbb{R}^d$, $\log(\frac{dP^{(n)}_{\theta_{n,h}}}{dP^{(n)}_{\theta_0}}) = h^\top \Delta_n - \frac{1}{2} h^\top I(\theta_0) h + o_{P^{(n)}_{\theta_0}}(1)$.

(ii) The *central sequence* satisfies $\Delta_n \Rightarrow N(0, I(\theta_0))$ under $P^{(n)}_{\theta_0}$.

**Remark 1.2.9.** *Why Local Asymptotics Matter*
LAN replaces local likelihood ratios by the Gaussian quadratic experiment $h^\top \Delta - \frac{1}{2} h^\top I(\theta_0) h$, giving normal approximations for estimators, tests, and confidence regions.

## 1.3 Design, Randomization, and Observational Comparisons

Comparative inference is only as meaningful as the experiment or study that produced the groups. The same two-sample statistic can estimate a causal treatment effect in a randomised experiment, describe an association in an observational study, or amplify bias if the groups were formed by a hidden mechanism.

**Definition 1.3.1.** *Controls, Randomisation, and Blinding*

(i) A *controlled experiment* compares $Y \mid T = 1$ and $Y \mid T = 0$ under common conditions.
(ii) *Randomisation* assigns $T \in \{0, 1\}$ so that $T$ is independent of pre-treatment factors $C$ by design.
(iii) *Blinding* conceals $T$ when knowledge of treatment could affect $Y$ or its measurement.

**Definition 1.3.2.** *Placebo Effect and Confounding*

(i) A *placebo effect* is variation in $Y$ caused by belief in treatment rather than the active mechanism.
(ii) Placebos and blinding separate expectation effects from the causal effect of $T$.
(iii) A pre-treatment variable $C$ is *confounded* with treatment $T$ for response $Y$ if $\mathcal{L}(C \mid T = 1) \neq \mathcal{L}(C \mid T = 0)$ and $C$ affects $Y$.
(iv) An *observational study* compares groups without assigning $T$. Group differences may reflect confounding rather than the factor of interest.

**Example 1.3.3.** *Mammary Artery Ligation*
A blinded trial randomised patients after preliminary incisions to ligation ($T = 1$) or sham surgery ($T = 0$). Similar outcomes show that controls remove background improvement; blinding limits placebo, observer effects.

**Principle 1.3.4.** *Confounding, Blocking, and Pairing*
Randomisation balances unmeasured covariates $C$ in probability; blocking, matching, and pairing condition on known influential $C$ before randomisation. Pairing is one-factor blocking; blocked and factorial designs extend this control to multiple factors.




Figure 240: Randomization breaks the confounding path from $C$ to $T$.

**Example 1.3.5.** *Uncontrolled Adjustments and Position Effects*
In the Lanarkshire milk experiment, post-randomisation movement between groups made $T$ depend on perceived nourishment $C$, reintroducing selection bias; weighing children in clothes also confounded measurement with socioeconomic status. In the FD&C Red No. 40 study, cage location was systematic in sex and dose, so position effects confounded treatment comparisons; randomising or blocking cage location would protect inference.

**Remark 1.3.6.** *Comparative Study Checklist*
Before interpreting group comparisons, record sampling, treatment assignment $T$, controls, blinding, blocking or pairing factors $C$, dependence among repeated measurements, and the number of tested comparisons. These determine whether the analysis supports causality, association, or hypothesis generation.

# 2 Estimation Theory

## 2.1 Estimators, Bias, Variance, and Risk

An estimator is a statistic interpreted as an answer to a target question. Its quality depends on the target, the loss function, and the model law under which performance is measured.

**Definition 2.1.1.** *Estimator, Loss, and Risk*

(i) A target parameter is a map $\psi : \Theta \to \Psi$.
(ii) An *estimator* of $\psi(\theta)$ is a statistic $\delta(X) \in \mathcal{D}$.
(iii) A *loss function* is $L : \Theta \times \mathcal{D} \to [0, \infty]$.
(iv) The *risk* of $\delta$ is $R(\theta, \delta) := \mathbb{E}_\theta L(\theta, \delta(X))$.
(v) For squared-error loss $L(\theta, d) = \|d - \psi(\theta)\|^2$, $R(\theta, \delta)$ is mean squared error.

**Definition 2.1.2.** *Sampling Distribution and Standard Error*

(i) For a statistic $T_n = T_n(X_1, \dots, X_n)$ under $P_\theta$, its *sampling distribution* is $\mathcal{L}_\theta(T_n)$.
(ii) If $T_n \in L^2(P_\theta)$, its *standard error* is $\mathrm{SE}_\theta(T_n) := \sqrt{\mathrm{Var}_\theta(T_n)}$.
(iii) An *estimated standard error* estimates $\mathrm{SE}_\theta(T_n)$, typically by plug-in, asymptotics, or resampling.

**Definition 2.1.3.** *Bias, Variance, and Consistency*
For real-valued $\widehat{\psi}$, $\mathrm{Bias}_\theta(\widehat{\psi}) := \mathbb{E}_\theta \widehat{\psi} - \psi(\theta)$, and $\mathrm{Var}_\theta(\widehat{\psi}) := \mathbb{E}_\theta(\widehat{\psi} - \mathbb{E}_\theta \widehat{\psi})^2$.

(i) $\widehat{\psi}$ is *unbiased* if $\mathrm{Bias}_\theta(\widehat{\psi}) = 0$ for all $\theta$.
(ii) $\widehat{\psi}_n$ is *consistent* at $\theta$ if $\widehat{\psi}_n \to \psi(\theta)$ in $P_\theta$-probability.

**Proposition 2.1.4.** *Bias-Variance Decomposition*
*Let $\widehat{\psi} \in L^2(P_\theta)$ be real-valued. Under squared-error loss,* $\mathrm{MSE}_\theta(\widehat{\psi}) := \mathbb{E}_\theta(\widehat{\psi} - \psi(\theta))^2 = \mathrm{Var}_\theta(\widehat{\psi}) + \mathrm{Bias}_\theta(\widehat{\psi})^2$.

*Proof.* Write $\widehat{\psi} - \psi(\theta) = (\widehat{\psi} - \mathbb{E}_\theta \widehat{\psi}) + (\mathbb{E}_\theta \widehat{\psi} - \psi(\theta))$. Square, take $\mathbb{E}_\theta$ gives the identity since $\mathbb{E}_\theta(\widehat{\psi} - \mathbb{E}_\theta \widehat{\psi}) = 0$.
*Structural Commentary*: MSE splits into random fluctuation plus squared systematic error.
*Key Insight*: Accuracy requires controlling both variance and bias. □

**Example 2.1.5.** *Sample Mean*
If $X_1, \dots, X_n$ are independent with $\mathbb{E}X_i = \mu$ and $\mathrm{Var}(X_i) = \sigma^2 < \infty$,
then $\overline{X}_n := n^{-1} \sum_{i=1}^{n} X_i$ satisfies $\mathbb{E}\overline{X}_n = \mu$, $\mathrm{Var}(\overline{X}_n) = \frac{\sigma^2}{n}$, and $\overline{X}_n \xrightarrow{P} \mu$.

**Definition 2.1.6.** *Plug-In and M-Estimators*

(i) A *plug-in estimator* estimates $T(P)$ by $T(\widehat{P}_n)$, where $\widehat{P}_n$ is empirical or fitted.
(ii) An *M-estimator* satisfies $\widehat{\theta}_n \in \arg\min_{\theta \in \Theta} M_n(\theta)$ for an empirical criterion $M_n$.
(iii) Common choices are $M_n =$ negative log-likelihood, squared loss, or absolute loss.

**Definition 2.1.7.** *Location M-Estimates*
For data $x_1, \dots, x_n \in \mathbb{R}$, scale $\sigma > 0$, and loss $\Psi : \mathbb{R} \to \mathbb{R}$, $\widehat{\mu} \in \arg\min_{\mu \in \mathbb{R}} \sum_{i=1}^{n} \Psi(\frac{x_i - \mu}{\sigma})$.
*Key Insight*: Quadratic loss gives $\overline{x}$; absolute or subquadratic loss improves outlier resistance.

## 2.2 Information Bounds and Asymptotic Efficiency

Information inequalities describe the best possible local accuracy under regular models. Efficient procedures are those whose asymptotic error reaches these bounds.

**Lemma 2.2.1.** *Two Forms of Fisher Information*
*For a regular one-parameter dominated model, let $\ell_\theta(x) := \log p_\theta(x)$ and $S_\theta(X) := \partial_\theta \ell_\theta(X)$.*
*Then $I(\theta) := \mathbb{E}_\theta S_\theta(X)^2 = -\mathbb{E}_\theta \partial_\theta^2 \ell_\theta(X)$.*

*Proof.* Differentiating $\int p_\theta \, d\mu = 1$ gives $\mathbb{E}_\theta S_\theta = 0$.
Differentiating, $0 = \partial_\theta \mathbb{E}_\theta S_\theta = \mathbb{E}_\theta[\partial_\theta S_\theta + S_\theta^2] = \mathbb{E}_\theta[\partial_\theta^2 \ell_\theta + S_\theta^2]$, since $\partial_\theta p_\theta = S_\theta p_\theta$. Thus $I(\theta) = \mathbb{E}_\theta S_\theta^2 = -\mathbb{E}_\theta \partial_\theta^2 \ell_\theta$.
*Structural Commentary*: Fisher information is both score variance and expected log-likelihood curvature.
*Key Insight*: Normalisation forces the mean score to vanish, linking slope variance to curvature. □

**Theorem 2.2.2.** *Cramer–Rao Bound*
*Let $X_1, \dots, X_n \sim P_\theta$ be iid from a regular one-parameter dominated model with one-observation information $I(\theta) > 0$. If $\widehat{\psi} \in L^2(P_\theta)$ is unbiased for differentiable $\psi(\theta)$, then* $\mathrm{Var}_\theta(\widehat{\psi}) \geq \frac{\psi'(\theta)^2}{nI(\theta)}$.

*Proof.* Let $S_n(\theta) := \partial_\theta \log L_n(\theta) = \sum_{i=1}^{n} S_\theta(X_i)$. Regularity gives $\mathbb{E}_\theta S_n = 0$ and $\mathrm{Var}_\theta(S_n) = \mathbb{E}_\theta S_n^2 = nI(\theta)$.
Differentiating $\mathbb{E}_\theta \widehat{\psi} = \psi(\theta)$ gives $\psi'(\theta) = \mathbb{E}_\theta[\widehat{\psi} S_n] = \mathrm{Cov}_\theta(\widehat{\psi}, S_n)$.
Thus $\psi'(\theta)^2 \le \mathrm{Var}_\theta(\widehat{\psi})\,\mathrm{Var}_\theta(S_n) = \mathrm{Var}_\theta(\widehat{\psi}) nI(\theta)$ by Cauchy–Schwarz.
*Structural Commentary*: The score is the direction in which unbiased estimators must covary to track $\psi(\theta)$.
*Key Insight*: Information lower-bounds variance through Cauchy–Schwarz. □

**Example 2.2.3.** *Poisson Mean Attains the Bound*
For $X \sim \mathrm{Poisson}(\lambda)$, $\ell_\lambda(X) = X \log \lambda - \lambda - \log(X!)$ and $\partial_\lambda^2 \ell_\lambda(X) = -\frac{X}{\lambda^2}$, so $I(\lambda) = -\mathbb{E}_\lambda \partial_\lambda^2 \ell_\lambda(X) = \frac{1}{\lambda}$.
For iid $X_1, \dots, X_n$, any unbiased estimator $T$ of $\lambda$ satisfies $\mathrm{Var}_\lambda(T) \ge \frac{\lambda}{n}$.
$\overline{X}$ attains equality since $\mathbb{E}\overline{X} = \lambda$ and $\mathrm{Var}(\overline{X}) = \frac{\lambda}{n}$.

**Definition 2.2.4.** *Asymptotic Efficiency*
For a scalar regular parameter with information $I(\theta_0) > 0$, an estimator sequence $\widehat{\theta}_n$ is *asymptotically efficient* at $\theta_0$ if $\sqrt{n}(\widehat{\theta}_n - \theta_0) \Rightarrow N(0, I(\theta_0)^{-1})$, i.e. it attains the inverse Fisher-information variance.

**Theorem 2.2.5.** *Regular MLE Asymptotic Normality*
*Let $X_1, \dots, X_n \sim P_{\theta_0}$ iid in a regular one-parameter model, with $\theta_0$ interior, $I(\theta_0) > 0$, and a consistent interior MLE $\widehat{\theta}_n$. Then $\sqrt{n}(\widehat{\theta}_n - \theta_0) \Rightarrow N(0, I(\theta_0)^{-1})$.*

*Proof.* Let $S_n(\theta) := \partial_\theta \ell_n(\theta)$ and $J_n(\theta) := -n^{-1} \partial_\theta S_n(\theta)$.
Since $S_n(\widehat{\theta}_n) = 0$, Taylor expansion gives $0 = S_n(\theta_0) - nJ_n(\widetilde{\theta}_n)(\widehat{\theta}_n - \theta_0)$ for some $\widetilde{\theta}_n$ between $\widehat{\theta}_n$ and $\theta_0$.
Hence $\sqrt{n}(\widehat{\theta}_n - \theta_0) = J_n(\widetilde{\theta}_n)^{-1} n^{-\frac{1}{2}} S_n(\theta_0)$.
Regularity gives $J_n(\widetilde{\theta}_n) \xrightarrow{P} I(\theta_0)$ and $n^{-\frac{1}{2}} S_n(\theta_0) \Rightarrow N(0, I(\theta_0))$, so Slutsky gives the result.
*Structural Commentary*: Local likelihood curvature converts score noise into estimator error.
*Key Insight*: The MLE is asymptotically score divided by information. □

**Remark 2.2.6.** *Efficiency and Approximation*
Finite-sample efficiency means equality in the Cramer–Rao bound.
Asymptotic efficiency means $\sqrt{n}(\widehat{\theta}_n - \theta_0) \Rightarrow N(0, I(\theta_0)^{-1})$; finite-sample bias or boundary effects still matter.

## 2.3 Method of Moments and Parametric Likelihood Examples

**Method 2.3.1.** *Method of Moments*

(i) Let $X_1, \dots, X_n$ be independent from $P_\theta$, and set $\mu_k(\theta) := \mathbb{E}_\theta X^k$ for $k = 1, \dots, r$.

(ii) If $\mu(\theta) = (\mu_1(\theta), \dots, \mu_r(\theta))$ identifies $\theta \in \Theta \subseteq \mathbb{R}^r$, define $\widehat{\mu}_k := n^{-1} \sum_{i=1}^{n} X_i^k$.

(iii) The *method of moments* estimator solves $\mu_k(\widehat{\theta}) = \widehat{\mu}_k$ for $k = 1, \dots, r$.

*Key Insight*: Match population moments to sample moments.

**Proposition 2.3.2.** *Consistency of Continuous Moment Estimates*
*Let $\widehat{\mu}_n \xrightarrow{P_{\theta_0}} \mu_0$ and let $g$ be continuous at $\mu_0$ with $g(\mu_0) = \theta_0$. Then $\widehat{\theta}_n := g(\widehat{\mu}_n) \xrightarrow{P_{\theta_0}} \theta_0$. Thus method-of-moments estimators are consistent whenever $\widehat{\mu}_n \xrightarrow{P} \mu(\theta_0)$, inversion map $g = \mu^{-1}$ is continuous at $\mu(\theta_0)$.*

*Proof.* Since $\widehat{\mu}_n \xrightarrow{P_{\theta_0}} \mu_0$ and $g$ is continuous at $\mu_0$, by continuous mapping theorem, $g(\widehat{\mu}_n) \xrightarrow{P_{\theta_0}} g(\mu_0) = \theta_0$.
For MoM, if the required moments are finite, the weak law gives $\widehat{\mu}_n \xrightarrow{P_{\theta_0}} \mu(\theta_0)$.
*Structural Commentary*: Moment consistency is convergence plus continuous inversion.
*Key Insight*: Estimate moments first, then pass through $g$. □

**Example 2.3.3.** *Moment Estimation for Small Open Probabilities*
Let $N \sim \mathrm{Binomial}(m, p)$ be the number of open channels and let $S = cN$ be total current.
Then $\mathbb{E}S = cmp$, $\mathrm{Var}(S) = c^2 mp(1-p)$, and $\frac{\mathrm{Var}(S)}{\mathbb{E}S} = c(1-p) \approx c$ for small $p$.
Thus independent measurements estimate $c$ by the empirical ratio $\frac{\widehat{\mathrm{Var}}(S)}{\widehat{\mathbb{E}S}}$, without knowing $m$.

**Example 2.3.4.** *Poisson Mean*
Let $X_1, \dots, X_n \sim \mathrm{Poisson}(\lambda)$ be independent. Since $\mathbb{E}_\lambda X = \lambda$, $\widehat{\lambda}_{\mathrm{mom}} = \overline{X}$.
Also $\ell(\lambda) = \sum_{i=1}^{n} \{X_i \log \lambda - \lambda - \log(X_i!)\}$ and $\ell'(\lambda) = \frac{S}{\lambda} - n$, where $S = \sum_i X_i$, so $\widehat{\lambda}_{\mathrm{mle}} = \frac{S}{n} = \overline{X}$.
If true value is $\lambda_0$, then $S \sim \mathrm{Poisson}(n\lambda_0)$. Hence $\mathbb{E}_{\lambda_0} \widehat{\lambda} = \lambda_0$, $\mathrm{Var}_{\lambda_0}(\widehat{\lambda}) = \frac{\lambda_0}{n}$, $\mathrm{SE}_{\lambda_0}(\widehat{\lambda}) = \sqrt{\frac{\lambda_0}{n}}$, $\widehat{\mathrm{SE}}(\widehat{\lambda}) = \sqrt{\frac{\widehat{\lambda}}{n}}$.

**Example 2.3.5.** *Negative Binomial Overdispersion*
For overdispersed counts, negative-binomial mass $f(x \mid m,k) = (1+\frac{m}{k})^{-k}\Gamma(k+x)\{x!\Gamma(k)\}^{-1}(\frac{m}{m+k})^x$, $x \in \mathbb{N}_0$, has $\mathbb{E}X = m$ and $\mathrm{Var}(X) = m + \frac{m^2}{k}$. By MoM $\widehat{m} = \overline{X}$, $\widehat{k} = \frac{\overline{X}^2}{\widehat{\sigma}^2 - \overline{X}}$ when $\widehat{\sigma}^2 > \overline{X}$.

**Example 2.3.6.** *Normal Mean and Variance*
For iid $X_i \sim N(\mu,\sigma^2)$, $\mu_1 = \mu$, $\mu_2 = \mu^2+\sigma^2$. Thus by MoM $\widehat{\mu} = \overline{X}$; $\widehat{\sigma}^2 = n^{-1}\sum_{i=1}^n X_i^2 - \overline{X}^2 = n^{-1}\sum_{i=1}^n (X_i - \overline{X})^2$.
The normal MLEs are the same; $\widehat{\sigma}^2$ uses denominator $n$, not the unbiased denominator $n-1$.

**Example 2.3.7.** *Gamma Shape and Rate*
Let $X_i$ be iid gamma with shape $\alpha$ and rate $\lambda$, so $\mathbb{E}X = \frac{\alpha}{\lambda}$ and $\mathrm{Var}(X) = \frac{\alpha}{\lambda^2}$.
With $\widehat{\sigma}^2 := n^{-1}\sum_{i=1}^n (X_i - \overline{X})^2$, MoM gives $\widehat{\alpha}_{\mathrm{mom}} = \frac{\overline{X}^2}{\widehat{\sigma}^2}$ and $\widehat{\lambda}_{\mathrm{mom}} = \frac{\overline{X}}{\widehat{\sigma}^2}$.
The log-likelihood is $\ell(\alpha,\lambda) = n\alpha\log\lambda + (\alpha-1)\sum_i \log X_i - \lambda\sum_i X_i - n\log\Gamma(\alpha)$.
The MLE satisfies $\widehat{\lambda} = \frac{\widehat{\alpha}}{\overline{X}}$ and $\log\widehat{\alpha} - \log\overline{X} + n^{-1}\sum_i \log X_i - \psi(\widehat{\alpha}) = 0$, where $\psi = \frac{\Gamma'}{\Gamma}$.
The MoM estimate is a natural starting value.

**Example 2.3.8.** *Angular Density*
Let $f(x \mid \alpha) = \frac{1+\alpha x}{2}$ on $[-1,1]$, with $\alpha \in [-1,1]$. Since $\mathbb{E}_\alpha X = \int_{-1}^1 \frac{x(1+\alpha x)}{2}\,dx = \frac{\alpha}{3}$, MoM gives $\widehat{\alpha} = 3\overline{X}$.
Also $\mathrm{Var}_\alpha(X) = \frac{1}{3} - \frac{\alpha^2}{9}$, so $\mathrm{Var}_\alpha(\widehat{\alpha}) = 9\,\mathrm{Var}_\alpha(\overline{X}) = \frac{3-\alpha^2}{n}$.

# 3 Hypothesis Testing and Confidence

## 3.1 Tests, Likelihood Ratios, and Error Control

Testing formalises decisions between hypotheses. The central quantities are the probability of false rejection under the null and the probability of rejection under alternatives.

**Definition 3.1.1.** *Hypotheses, Tests, Size, and Power*

(i) A *null hypothesis* is a subset $\Theta_0 \subseteq \Theta$ of the parameter space.
(ii) An *alternative* is a subset $\Theta_1 \subseteq \Theta$ disjoint from $\Theta_0$, so $\Theta_0 \cap \Theta_1 = \emptyset$.
(iii) Randomised *test* is a measurable map $\varphi : \mathcal{X} \to [0,1]$, where $\varphi(x)$ is rejection probability at observation $x$.
(iv) The *size* of $\varphi$ is $\alpha(\varphi) = \sup_{\theta\in\Theta_0} \mathbb{E}_\theta[\varphi(X)]$.
(v) The *power function* of $\varphi$ is $\beta_\varphi(\theta) = \mathbb{E}_\theta[\varphi(X)]$.

**Definition 3.1.2.** *Simple and Composite Hypotheses; Type Errors*

(i) A *simple* hypothesis specifies one law; a *composite* hypothesis specifies several.
(ii) A *type I error* rejects a true $H_0$; its probability is bounded by $\alpha(\varphi)$.
(iii) For a simple alternative $P_1$, a *type II error* is failure to reject $H_0$ under $P_1$, with probability $\mathbb{E}_1[1-\varphi(X)]$.
(iv) The *power* against $P_1$ is $1 - \mathbb{E}_1[1-\varphi(X)]$.

**Theorem 3.1.3.** *Neyman–Pearson Lemma*
*Let $P_1 \ll P_0$, $\Lambda := \frac{dP_1}{dP_0}$, $\alpha \in [0,1]$. If $c \geq 0$ and $\gamma \in [0,1]$ make $\varphi^* := \mathbf{1}_{\{\Lambda > c\}} + \gamma\mathbf{1}_{\{\Lambda = c\}}$ satisfy $\mathbb{E}_0\varphi^* = \alpha$, then $\varphi^*$ is most powerful among level-$\alpha$ tests: for every $\psi : \mathcal{X} \to [0,1]$ with $\mathbb{E}_0\psi \leq \alpha$, $\mathbb{E}_1\psi \leq \mathbb{E}_1\varphi^*$.*

*Proof.* For any level-$\alpha$ competitor $\psi$, $(\varphi^* - \psi)(\Lambda - c) \geq 0$ pointwise.
Since $dP_1 = \Lambda\,dP_0$, $\mathbb{E}_1(\varphi^* - \psi) = \mathbb{E}_0[\Lambda(\varphi^* - \psi)] \geq c\mathbb{E}_0(\varphi^* - \psi) = c(\alpha - \mathbb{E}_0\psi) \geq 0$.
*Structural Commentary*: Null controls size constraint, while alternative supplies power objective.
*Key Insight*: Use the rejection budget where $\Lambda = \frac{dP_1}{dP_0}$ is largest. □

**Remark 3.1.4.** *Choosing the Null*
At level $a$, for simple $P_0$ versus $P_1$, the *Neyman–Pearson problem* is $\sup\{\mathbb{E}_1\varphi : 0 \leq \varphi \leq 1,\ \mathbb{E}_0\varphi \leq a\}$.
Thus $H_0$ is the protected claim: choose it so the false-rejection probability $\mathbb{E}_0\varphi$ is the error to be bounded by $a$.
The choice of $H_0$ is substantive, not determined by $\{P_0, P_1\}$.

**Definition 3.1.5.** *Uniformly Most Powerful Test*
For a composite alternative $\Theta_1$, a level-$\alpha$ test $\varphi$ is *uniformly most powerful* if, for every $\theta \in \Theta_1$, it has power at least as large as every other level-$\alpha$ test at that same $\theta$.

**Example 3.1.6.** *Normal Mean, Known Variance*
Let $X_1, \dots, X_n$ be independent $N(\mu, \sigma^2)$ variables with known $\sigma^2$. For $H_0 : \mu = \mu_0$ against $H_1 : \mu = \mu_1 > \mu_0$, the likelihood ratio is increasing in $\overline{X}$, so the level-$\alpha$ test is $\varphi^* = \mathbf{1}_{\{\overline{X} > \mu_0 + \frac{z_{1-\alpha}\sigma}{\sqrt{n}}\}}$, where $z_{1-\alpha}$ is the $N(0,1)$ quantile. Since the cutoff is independent of $\mu_1$, $\varphi^*$ is uniformly most powerful for $H_1 : \mu > \mu_0$; no single likelihood-ratio direction is uniformly most powerful for $H_1 : \mu \neq \mu_0$.

**Definition 3.1.7.** *Generalized Likelihood-Ratio Test*
For $H_0 : \theta \in \Theta_0 \subseteq \Theta$, the *generalized likelihood ratio* is $\lambda(x) := \frac{\sup_{\theta \in \Theta_0} L_x(\theta)}{\sup_{\theta \in \Theta} L_x(\theta)} \in [0,1]$.
A GLR test rejects $H_0$ for small $\lambda(x)$, typically $\lambda(x) \le c_\alpha$ with $c_\alpha$ chosen to give level $\alpha$.

**Theorem 3.1.8.** *Large-Sample Likelihood-Ratio Calibration*
*(i) Let $H_0 : \theta \in \Theta_0 \subseteq \Theta$, $\dim \Theta = d$, $\dim \Theta_0 = d_0$, $r := d - d_0$, and $X^{(n)} \sim P_{\theta_0}$ with $\theta_0 \in \Theta_0$.*
*(ii) Let $L_n(\theta) := e^{\ell_n(\theta)}$ and $\lambda_n := \frac{\sup_{\theta \in \Theta_0} L_n(\theta)}{\sup_{\theta \in \Theta} L_n(\theta)}$.*
*(iii) Assume local coordinates with $\theta_0 = 0$, $\Theta_0 = \{(u,0) : u \in \mathbb{R}^{d_0}\}$, $h = (u,v) \in \mathbb{R}^{d_0} \times \mathbb{R}^r$, and $\ell_n(\frac{h}{\sqrt{n}}) - \ell_n(0) = h^\top Z_n - \frac{\|h\|^2}{2} + o_{P_{\theta_0}}(1)$, where $Z_n = (U_n, V_n) \Rightarrow N_d(0, I_d)$, $U_n \in \mathbb{R}^{d_0}$, $V_n \in \mathbb{R}^r$.*
*(iv) Then under $H_0$, $-2\log\lambda_n \Rightarrow \chi^2_r$.*

*Proof.*
(i) Since $L_n = e^{\ell_n}$, $\log\lambda_n = \sup_{\theta \in \Theta_0} \ell_n(\theta) - \sup_{\theta \in \Theta} \ell_n(\theta)$, hence $-2\log\lambda_n = 2\{\sup_{\theta \in \Theta} \ell_n(\theta) - \sup_{\theta \in \Theta_0} \ell_n(\theta)\}$.
(ii) In the local coordinates, write $\theta = \frac{h}{\sqrt{n}}$ for the full model and $\theta = \frac{(u,0)}{\sqrt{n}}$ for the null model.
With $Q_n(h) := h^\top Z_n - \frac{\|h\|^2}{2}$ and $Q_{0,n}(u) := Q_n(u,0) = u^\top U_n - \frac{\|u\|^2}{2}$, (iii) and cancellation of $\ell_n(0)$ give $-2\log\lambda_n = 2\{\sup_{h \in \mathbb{R}^d} Q_n(h) - \sup_{u \in \mathbb{R}^{d_0}} Q_{0,n}(u)\} + o_{P_{\theta_0}}(1)$.
(iii) For $a \in \mathbb{R}^m$, $x^\top a - \frac{\|x\|^2}{2} = \frac{\|a\|^2}{2} - \frac{\|x-a\|^2}{2}$, so supremum over $x \in \mathbb{R}^m$ is $\frac{\|a\|^2}{2}$ at $x = a$.
Applying this with $a = Z_n$ and $a = U_n$ gives $\sup_{h \in \mathbb{R}^d} Q_n(h) = \frac{\|Z_n\|^2}{2}$ and $\sup_{u \in \mathbb{R}^{d_0}} Q_{0,n}(u) = \frac{\|U_n\|^2}{2}$,
hence $-2\log\lambda_n = \|Z_n\|^2 - \|U_n\|^2 + o_{P_{\theta_0}}(1) = \|V_n\|^2 + o_{P_{\theta_0}}(1)$.
(iv) Since $Z_n = (U_n, V_n) \Rightarrow N_d(0, I_d)$, marginal convergence gives $V_n \Rightarrow Y$ with $Y \sim N_r(0, I_r)$.
Therefore $\|V_n\|^2 \Rightarrow \|Y\|^2 \sim \chi^2_r$ by continuous mapping, and Slutsky gives $-2\log\lambda_n \Rightarrow \chi^2_r$.
*Structural Commentary*: The likelihood ratio compares unconstrained fit with null-constrained fit.
*Key Insight*: Wilks' statistic is the squared Gaussian fluctuation in the $r = d - d_0$ constrained directions. □

**Example 3.1.9.** *Two-Sided Normal Mean GLRT*
Let $X_1, \dots, X_n$ be independent $N(\mu, \sigma^2)$ variables with known $\sigma^2$, and test $H_0 : \mu = \mu_0$ against $H_1 : \mu \neq \mu_0$.
Since $\widehat{\mu} = \overline{X}$, $\lambda(X) = \frac{L_n(\mu_0)}{L_n(\overline{X})} = \exp\{-\frac{n(\overline{X}-\mu_0)^2}{2\sigma^2}\}$, so $-2\log\lambda(X) = \frac{n(\overline{X}-\mu_0)^2}{\sigma^2}$.
Under $H_0$, $\frac{\sqrt{n}(\overline{X}-\mu_0)}{\sigma} \sim N(0,1)$; hence $-2\log\lambda(X) \sim \chi^2_1$,
and the level-$\alpha$ GLR rejects iff $\frac{n(\overline{X}-\mu_0)^2}{\sigma^2} > \chi^2_{1,1-\alpha}$, equivalently $\left|\overline{X} - \mu_0\right| > \frac{z_{1-\frac{\alpha}{2}}\sigma}{\sqrt{n}}$.

**Example 3.1.10.** *Sign Test as a Binomial GLRT*
Let $X \sim \text{Binomial}(n,p)$ and test $H_0 : p = \frac{1}{2}$ against $H_1 : p \neq \frac{1}{2}$.
For observed $x$, $\widehat{p} = \frac{x}{n}$ and $\lambda(x) = \frac{L_x(\frac{1}{2})}{L_x(\widehat{p})} = \frac{(\frac{1}{2})^n}{[(\frac{x}{n})^x(1-\frac{x}{n})^{n-x}]}$, $0^0 = 1$ at $x = 0, n$.
Since $\lambda(x)$ decreases as $\left|x - \frac{n}{2}\right|$ increases, a level-$\alpha$ GLR rejects on the largest values of $\left|X - \frac{n}{2}\right|$, with cutoff chosen under $X \sim \text{Binomial}(n, \frac{1}{2})$; this is the two-sided sign test.

**Definition 3.1.11.** *p-Value*
A *p-value* for $H_0 : \theta \in \Theta_0$ is a statistic $p : \mathcal{X} \to [0,1]$ such that $\sup_{\theta \in \Theta_0} \mathbb{P}_\theta\{p(X) \le u\} \le u$ for every $u \in [0,1]$.
Then $\varphi_\alpha(X) := \mathbf{1}_{\{p(X) \le \alpha\}}$ has size $\alpha(\varphi_\alpha) \le \alpha$.

**Proposition 3.1.12.** *Continuous Null p-Values Are Uniform*
*Let $H_0 : \theta \in \Theta_0$ and let $T : \mathcal{X} \to \mathbb{R}$ satisfy $\mathbb{P}_\theta(T \le t) = F_0(t)$ for all $\theta \in \Theta_0$, where $F_0$ is continuous.*
*For the right-tail convention, set $p(X) := 1 - F_0(T(X))$.*
*Then $p(X) \sim \text{Uniform}(0,1)$ under every $\theta \in \Theta_0$, and $\varphi_\alpha(X) := \mathbf{1}_{\{p(X) \le \alpha\}}$ has size $\alpha$ for $\alpha \in [0,1]$.*

*Proof.* Fix $\theta \in \Theta_0$ and set $U := F_0(T(X))$.
By probability integral transform, Proposition 2.3.6, $U \sim \text{Uniform}(0,1)$ under $\mathbb{P}_\theta$,

hence $p(X) = 1 - U \sim \text{Uniform}(0,1)$. Thus $\mathbb{E}_\theta \varphi_\alpha = \mathbb{P}_\theta(p(X) \le \alpha) = \alpha \ \forall \theta \in \Theta_0$, so $\alpha(\varphi_\alpha) = \sup_{\theta \in \Theta_0} \mathbb{E}_\theta \varphi_\alpha = \alpha$.

*Structural Commentary*: A continuous null CDF converts the statistic to a canonical uniform scale.
*Key Insight*: Exact calibration is the probability integral transform applied under $H_0$. □

**Remark 3.1.13.** *p-Values Are Tail Probabilities*
A $p$-value is a null tail probability, e.g. $p(x) = \mathbb{P}_{H_0}\{T(X) \ge T(x)\}$ for a right-tail statistic $T$.
It is not $\mathbb{P}(H_0 \mid x)$; posterior probabilities require priors and an alternative likelihood.

**Definition 3.1.14.** *Multiple-Testing Error Counts and FDR*
For a multiple-testing rule over $M$ hypotheses.
Let $M_0$ be the number of true nulls and $M_1 = M - M_0$ the number of false nulls.
Let $V$ be false rejections, $S$ true rejections, $U$ true nonrejections, and $T$ false nonrejections.
The total number of discoveries is $R = V + S$.

(i) *Family-wise error rate*: $\text{FWER} := \mathbb{P}(V \ge 1)$.
(ii) *False discovery proportion*: $\text{FDP} := \frac{V}{\max(R,1)}$.
(iii) *False discovery rate*: $\text{FDR} := \mathbb{E}[\text{FDP}]$.
(iv) *Power*: One aggregate power measure is $\mathbb{E}[S]/M_1$ when $M_1 > 0$.

Thus FWER controls the chance of any false discovery.
FDR controls the expected false fraction among the reported discoveries.

**Remark 3.1.15.** *Error Rates*
For one test $\varphi$, size is $\alpha(\varphi) = \sup_{\theta \in \Theta_0} \mathbb{E}_\theta \varphi$. For a family of tests, let $R$ be the number of rejections and $V$ the number of false rejections; common targets are $\text{FWER} = \mathbb{P}\{V \ge 1\}$ and $\text{FDR} = \mathbb{E}[\frac{V}{R \vee 1}]$, while directional or selective rates replace $V$ by the corresponding error count.

**Method 3.1.16.** *Fishing Expeditions and Bonferroni Control*

(i) For $N$ null hypotheses, let $R_i$ be the event that null $i$ is rejected.
(ii) By the union bound, $\mathbb{P}(\bigcup_{i=1}^{N} R_i) \le \sum_{i=1}^{N} \mathbb{P}(R_i)$.
(iii) To control family-wise error at level $\alpha$, test each null at level $\frac{\alpha}{N}$; then $\mathbb{P}(\text{at least one false rejection}) \le \alpha$.
(iv) Without such correction, many tests can produce small $p$-values even when all nulls are true.

**Definition 3.1.17.** *Positive FDR, Local FDR, and q-Values*
Let $Z_j = 0$ indicate a true null, $Z_j = 1$ a false null, $t_j$ a test statistic, and $\Gamma$ a rejection region.

(i) *pFDR*: For false discoveries $V$ and discoveries $R$, $\text{pFDR} := \mathbb{E}[V/R \mid R > 0]$.
(ii) *Mixture model*: If $\mathbb{P}(Z_j = 0) = \pi_0$, $\mathcal{L}(t_j \mid Z_j = 0) = F_0$, and $\mathcal{L}(t_j \mid Z_j = 1) = F_1$, then $\mathcal{L}(t_j) = \pi_0 F_0 + (1 - \pi_0) F_1$.
(iii) *Bayes form*: For a rejection region $\Gamma$, $\text{pFDR}(\Gamma) = \mathbb{P}(Z_j = 0 \mid t_j \in \Gamma)$.
(iv) *Local FDR*: At value $t_0$, $\text{lfdr}(t_0) := \mathbb{P}(Z_j = 0 \mid t_j = t_0)$.
(v) *q-value*: For feature $j$, $q_j := \inf\{\alpha : j \in \mathcal{R}_\alpha\}$, where $\mathcal{R}_\alpha$ is the discovery set at FDR level $\alpha$.

**Remark 3.1.18.** *FDR Versus Family-Wise Error*
With false discoveries $V$ and total discoveries $R$, $\text{FWER} := \mathbb{P}(V \ge 1)$ controls any false rejection, while $\text{FDR} := \mathbb{E}[V/\max(R,1)]$ controls the expected false fraction among discoveries. Thus FDR is less conservative for large screening lists and measures list reliability directly.

**Method 3.1.19.** *Benjamini–Hochberg Step-Up Rule*
For $M$ hypotheses with $p$-values $p_1, \dots, p_M$ and target level $\alpha \in (0,1)$:

(i) *Order*: Let $p_{(1)} \le \cdots \le p_{(M)}$ be the order statistics.
(ii) *Step-up index*: Set $L := \max\{j : p_{(j)} \le \alpha j / M\}$, with $L := 0$ if the set is empty.
(iii) *Rejections*: Reject $\mathcal{R} := \{m : p_m \le p_{(L)}\}$ if $L > 0$, and reject none if $L = 0$.
(iv) *Guarantee*: If true-null $p$-values are independent and uniform, then $\text{FDR} \le (M_0/M)\alpha \le \alpha$, where $M_0$ is the number of true nulls.




Figure 241: Benjamini–Hochberg threshold: red points reject, blue points do not; line is $\alpha j/M$.

## 3.2 Confidence Sets, Pivots, and Asymptotic Intervals

Confidence procedures attach uncertainty to estimators by controlling coverage probability under repeated sampling.

**Definition 3.2.1.** *Confidence Sets and Coverage*
A random set $C(X) \subseteq \Psi$ is a $1-\alpha$ *confidence set* for $\psi : \Theta \to \Psi$ if $\inf_{\theta\in\Theta} \mathbb{P}_\theta\{\psi(\theta) \in C(X)\} \geq 1-\alpha$.
The left side is the *coverage guarantee*.

**Definition 3.2.2.** *Pivots and Confidence Construction*

(i) A *pivot* is a statistic $Q(X,\theta)$ whose distribution under $\mathbb{P}_\theta$ is free of $\theta$.
(ii) If constants $a_\alpha, b_\alpha$ satisfy $\mathbb{P}_\theta\{a_\alpha \leq Q(X,\theta) \leq b_\alpha\} = 1-\alpha$ for all $\theta \in \Theta$, then $C(X) := \{\theta \in \Theta : a_\alpha \leq Q(X,\theta) \leq b_\alpha\}$ is an exact $1-\alpha$ confidence set for $\theta$.

**Corollary 3.2.3.** *Hoeffding Confidence Interval for a Bernoulli Parameter*
*Let $X_1, \dots, X_n$ be independent* Bernoulli$(p)$ *variables, $\widehat{p}_n := \overline{X}_n$, and $\alpha \in (0,1)$.*
*Set $\varepsilon_n := \sqrt{(2n)^{-1}\log(\frac{2}{\alpha})}$ and $C_n := [\widehat{p}_n - \varepsilon_n, \widehat{p}_n + \varepsilon_n]$.*

*(i) Coverage: $\mathbb{P}(p \in C_n) \geq 1-\alpha$.*
*(ii) Length: $|C_n| = 2\varepsilon_n \leq 0.05$ whenever $n \geq 800\log(\frac{2}{\alpha})$.*

*Proof.*

(i) Since $\{p \notin C_n\} = \{|\widehat{p}_n - p| > \varepsilon_n\}$, Corollary 3.3.5 gives $\mathbb{P}(p \notin C_n) \leq 2e^{-2n\varepsilon_n^2} = \alpha$.
(ii) If $n \geq 800\log(\frac{2}{\alpha})$, then $\varepsilon_n^2 \leq \frac{1}{1600}$, hence $|C_n| = 2\varepsilon_n \leq 0.05$.

*Structural Commentary*: Confidence intervals invert a concentration inequality.
*Key Insight*: Choose $\varepsilon_n$ so the Hoeffding failure bound equals $\alpha$. □

**Proposition 3.2.4.** *Test–Confidence Duality*
*For each $\psi_0 \in \Psi$, let $\varphi_{\psi_0} : \mathcal{X} \to \{0,1\}$ satisfy $\sup_{\theta:\psi(\theta)=\psi_0} \mathbb{E}_\theta \varphi_{\psi_0} \leq \alpha$.*
*Then $C(X) := \{\psi_0 \in \Psi : \varphi_{\psi_0}(X) = 0\}$ is a $1-\alpha$ confidence set for $\psi(\theta)$.*

*Proof.* Fix $\theta \in \Theta$ and set $\psi_\theta := \psi(\theta)$.
By definition, $\{\psi(\theta) \notin C(X)\} = \{\varphi_{\psi_\theta}(X) = 1\}$, so $\mathbb{P}_\theta\{\psi(\theta) \notin C(X)\} = \mathbb{E}_\theta\varphi_{\psi_\theta} \leq \sup_{\eta:\psi(\eta)=\psi_\theta} \mathbb{E}_\eta\varphi_{\psi_\theta} \leq \alpha$.
Hence $\mathbb{P}_\theta\{\psi(\theta) \in C(X)\} \geq 1-\alpha$ for every $\theta$, and $\inf_{\theta\in\Theta} \mathbb{P}_\theta\{\psi(\theta) \in C(X)\} \geq 1-\alpha$.
*Structural Commentary*: Inverting all accepted point-null tests turns non-rejection regions into a confidence set.
*Key Insight*: Coverage failure equals rejection of the true null value. □

**Proposition 3.2.5.** *Confidence–Test Converse*
*Let $C(X) \subseteq \Psi$ satisfy $\inf_{\theta\in\Theta} \mathbb{P}_\theta\{\psi(\theta) \in C(X)\} \geq 1-\alpha$.*
*For $\psi_0 \in \Psi$, let $\varphi_{\psi_0}(X) := \mathbf{1}_{\{\psi_0 \notin C(X)\}}$ for testing $H_0 : \psi(\theta) = \psi_0$. Then $\varphi_{\psi_0}$ is level $\alpha$: $\sup_{\theta:\psi(\theta)=\psi_0} \mathbb{E}_\theta\varphi_{\psi_0} \leq \alpha$.*

*Proof.* Fix $\theta$ with $\psi(\theta) = \psi_0$. Then $\mathbb{E}_\theta\varphi_{\psi_0} = \mathbb{P}_\theta\{\psi_0 \notin C(X)\} = \mathbb{P}_\theta\{\psi(\theta) \notin C(X)\} = 1 - \mathbb{P}_\theta\{\psi(\theta) \in C(X)\} \leq \alpha$.
Taking $\sup_{\theta:\psi(\theta)=\psi_0}$ gives the level bound.
*Structural Commentary*: A confidence set tests a proposed value by checking membership.
*Key Insight*: Rejection of $\psi_0$ is exactly non-coverage when $\psi_0$ is true. □

**Remark 3.2.6.** *Acceptance Regions and Confidence Regions*
For level-$\alpha$ point-null tests with acceptance regions $A_{\psi_0} \subseteq \mathcal{X}$, the inversion $C(X) := \{\psi_0 \in \Psi : X \in A_{\psi_0}\}$ is the set of null values not rejected at $X$. Conversely, any confidence set $C(X)$ induces point-null tests $\varphi_{\psi_0}(X) = \mathbf{1}_{\{\psi_0 \notin C(X)\}}$.

**Definition 3.2.7.** *Wald, Score, and Likelihood Intervals*
For scalar regular $\theta$ with estimator $\widehat{\theta}$, standard error $\widehat{\mathrm{se}}(\widehat{\theta})$, score $U_n$, and likelihood ratio $\lambda_n(\theta_0)$:

(i) *Wald*: use $\widehat{\theta} \pm z_{1-\frac{\alpha}{2}} \widehat{\mathrm{se}}(\widehat{\theta})$.
(ii) *Score*: invert $\frac{|U_n(\theta_0)|}{\sqrt{I_n(\theta_0)}} \le z_{1-\frac{\alpha}{2}}$ over $\theta_0$.
(iii) *Likelihood*: invert $-2\log\lambda_n(\theta_0) \le \chi^2_{1,1-\alpha}$ over $\theta_0$.

*Key Insight*: Wald, score, and likelihood intervals are first-order equivalent in regular models.

**Example 3.2.8.** *Poisson Mean Wald Interval*
Let $X_1, \dots, X_n$ be independent Poisson($\lambda$) variables.
Then $\widehat{\lambda} = \overline{X}$ and $\frac{\sqrt{n}(\overline{X}-\lambda)}{\sqrt{\lambda}} \Rightarrow N(0,1)$, so the plug-in standard error is $\widehat{\mathrm{se}}(\widehat{\lambda}) = \sqrt{\frac{\overline{X}}{n}}$.
The Wald interval is $C_\alpha(X) = [\overline{X} - z_{1-\frac{\alpha}{2}}\sqrt{\frac{\overline{X}}{n}}, \overline{X} + z_{1-\frac{\alpha}{2}}\sqrt{\frac{\overline{X}}{n}}]$, with asymptotic coverage $1-\alpha$.

## 3.3 Goodness-of-Fit and Diagnostic Plots

Goodness-of-fit methods compare an observed empirical pattern with the pattern predicted by a probability model. Formal tests give calibrated evidence against a specified model, while diagnostic plots show where the model fits or fails.

**Definition 3.3.1.** *Multinomial Goodness-of-Fit Model*

(i) *Observed counts*: $O = (O_1, \dots, O_m)$ records cell frequencies from $n$ trials, with $O_i \in \mathbb{N}_0$ and $\sum_{i=1}^{m} O_i = n$.
(ii) *Null model*: $\mathcal{M}_0 = \{p(\theta) = (p_1(\theta), \dots, p_m(\theta)) : \theta \in \Theta_0\} \subseteq \Delta_{m-1}$ is parametric.
(iii) *Saturated model*: $\Delta_{m-1} = \{p \in [0,1]^m : \sum_i p_i = 1\}$ allows every multinomial probability vector.
(iv) *Fitted counts*: if $\widehat{\theta}$ is the null MLE, then $E_i := np_i(\widehat{\theta})$ for $i = 1, \dots, m$.
(v) *Degrees of freedom*: if $\dim \Theta_0 = k$, then $\mathrm{df} = (m-1) - k = m - k - 1$.

**Definition 3.3.2.** *Pearson and Likelihood-Ratio Goodness-of-Fit Statistics*

(i) *Inputs*: $O = (O_1, \dots, O_m)$ are observed counts, $E = (E_1, \dots, E_m)$ are fitted null counts with $E_i > 0$, and the null model has $k$ free parameters.
(ii) *Pearson statistic*: $X^2 := \sum_{i=1}^{m} \frac{(O_i - E_i)^2}{E_i}$.
(iii) *Likelihood-ratio statistic*: $G^2 := 2\sum_{i=1}^{m} O_i \log(\frac{O_i}{E_i})$, with $0\log(\frac{0}{E_i}) := 0$.
(iv) *Null calibration*: under the regular null model, $X^2 \Rightarrow \chi^2_{m-k-1}$ and $G^2 = -2\log\lambda(X) \Rightarrow \chi^2_{m-k-1}$. Moreover $G^2 - X^2 = o_P(1)$ under $H_0$.

**Example 3.3.3.** *Hardy–Weinberg Fit*
For genotype cells $(M, MN, N)$, the Hardy–Weinberg model is $p(\theta) = ((1-\theta)^2, 2\theta(1-\theta), \theta^2)$.
With $n = 1029$ and $\widehat{\theta} = 0.4247$, the observed and fitted counts are

| | $M$ | $MN$ | $N$ |
|---|---|---|---|
| $O_i$ | 342 | 500 | 187 |
| $E_i$ | 340.6 | 502.8 | 185.6 |

Since $m = 3$ and $k = 1$, $\mathrm{df} = m - k - 1 = 1$. Pearson statistic is $X^2 = 0.0319$, so $p = \mathbb{P}\{\chi^2_1 \ge 0.0319\}$ is large. The data give no evidence against Hardy–Weinberg equilibrium.

**Method 3.3.4.** *Reading Cell Contributions*

(i) Compute signed residuals $r_i := O_i - E_i$ and Pearson contributions $c_i := \frac{(O_i - E_i)^2}{E_i}$.
(ii) Large $c_i$ identify cells driving the rejection; the sign of $r_i$ indicates overfit or underfit in that cell.
(iii) If some $E_i$ are small, group adjacent or tail cells before testing so the chi-square approximation is reliable.

**Remark 3.3.5.** *Grouped Counts and Expected Frequencies*

Let $Y_1, \dots, Y_N$ be observations, let $A_1, \dots, A_m$ partition sample space, with observed counts $O_j := \sum_{i=1}^{N} \mathbf{1}_{\{Y_i \in A_j\}}$.

Given a fitted parametric law $P_{\widehat{\theta}}$, set $p_j(\widehat{\theta}) := P_{\widehat{\theta}}(Y \in A_j)$; the fitted expected count is $E_j := Np_j(\widehat{\theta})$.
For grouped Poisson tails, $p_j(\widehat{\theta})$ is a sum of Poisson masses, $(O_1, \dots, O_m)$ is modelled as multinomial under $P_{\widehat{\theta}}$.

**Example 3.3.6.** *Poisson Clumps*
For grouped bacterial clump frequencies from $n = 400$ grid squares, fit $H_0 : X \sim \text{Poisson}(\lambda)$ with $\widehat{\lambda} = 2.44$ and grouped tails so $E_i$ are adequate. The Pearson statistic is $X^2 = 75.4$ with df $= 6$, so $p = \mathbb{P}\{\chi^2_6 \geq 75.4\} < 0.005$; the largest $c_i = \frac{(O_i - E_i)^2}{E_i}$ are in the first and last cells, indicating excess small and large counts.

**Definition 3.3.7.** *Overdispersed Counts*
Count data are *overdispersed* relative to a Poisson model if $\text{Var}(X) > \mathbb{E}X$; empirically, this is $\widehat{\sigma}^2 > \overline{X}$.

**Definition 3.3.8.** *Poisson Dispersion Test*
The data is $x_1, \dots, x_n$ are independent counts and $\overline{x} := n^{-1} \sum_{i=1}^{n} x_i$.

(i) *Models*: $H_0 : x_i \sim \text{Poisson}(\lambda)$ with common $\lambda$; the saturated alternative uses separate rates $\lambda_i$.
(ii) *Likelihood-ratio statistic*: $G^2_D := 2 \sum_{i=1}^{n} x_i \log(\frac{x_i}{\overline{x}})$, with $0 \log(\frac{0}{\overline{x}}) := 0$.
(iii) *Pearson approximation*: $T := \overline{x}^{-1} \sum_{i=1}^{n} (x_i - \overline{x})^2$.
(iv) *Calibration*: under $H_0$, $G^2_D$ and $T$ are calibrated by $\chi^2_{n-1}$
Large values indicate overdispersion relative to $\text{Var}(X) = \mathbb{E}X$.

**Example 3.3.9.** *Dispersion in Bacterial Clumps*
For bacterial clump counts with $n = 400$, $\overline{x} = 2.44$ and $s^2 = 4.59$, so $\frac{s^2}{\overline{x}} = 1.88 > 1$.
Dispersion statistic is $T \approx 752.7$ with df $= n - 1 = 399$, so $p = \mathbb{P}\{\chi^2_{399} \geq 752.7\} \approx 0$.
The data show strong overdispersion relative to a Poisson model.

**Definition 3.3.10.** *Hanging Rootograms and Chi-Grams*
Let $n_j$ be observed bin counts and $\widehat{n}_j$ fitted counts in ordered bins.

(i) A *hanging histogram* plots raw residuals $r_j := n_j - \widehat{n}_j$, whose variance depends on $\widehat{n}_j$.
(ii) A *hanging rootogram* plots $q_j := \sqrt{n_j} - \sqrt{\widehat{n}_j}$, using $\text{Var}(\sqrt{n_j}) \approx \frac{1}{4}$ for count data.
(iii) A *hanging chi-gram* plots standardized residuals $z_j := \frac{n_j - \widehat{n}_j}{\sqrt{\widehat{n}_j}}$.

Runs of same-sign residuals, tail spikes, or systematic curvature indicate model departure.

**Method 3.3.11.** *Probability Plots*
Let $X_{(1)} \leq \cdots \leq X_{(n)}$ be order statistics and let $F$ be the candidate continuous strictly increasing CDF.

(i) A *P-P plot* compares $F(X_{(k)})$ with $u_k := \frac{k}{n+1}$ for $k = 1, \dots, n$.
(ii) A *Q-Q plot* compares $X_{(k)}$ with $F^{-1}(u_k)$ for $k = 1, \dots, n$.

The model is plausible when the plotted points are approximately linear.
For $F(x) = G(\frac{x-\mu}{\sigma})$, Q-Q slope and intercept estimate $\sigma$ and $\mu$.

**Method 3.3.12.** *Probability Plots for Grouped Data*
Let bins boundaries $x_1 < \cdots < x_{m+1}$, counts $n_1, \dots, n_m$, total $n := \sum_{j=1}^{m} n_j$, cumulative counts $N_j := \sum_{i=1}^{j} n_i$.
Let $G$ be the standardised CDF in the proposed location-scale family $F_{\mu,\sigma}(x) = G(\frac{x-\mu}{\sigma})$.

(i) Plot empirical boundary quantiles $x_{j+1}$ against theoretical quantiles $G^{-1}(\frac{N_j}{n+1})$ for $j = 1, \dots, m$.
(ii) Use this grouped Q-Q plot when individual order statistics are unavailable.

**Remark 3.3.13.** *Grouped Normality Tests*
For binned normal-fit testing, use $X^2 = \sum_j \frac{(O_j - E_j)^2}{E_j}$ with $E_j = n\{F_{\widehat{\mu},\widehat{\sigma}}(b_{j+1}) - F_{\widehat{\mu},\widehat{\sigma}}(b_j)\}$.
The usual $\chi^2_{m-k-1}$ calibration requires $(\widehat{\mu}, \widehat{\sigma})$ from the grouped counts; ungrouped estimates change the limit.

**Definition 3.3.14.** *Normality Diagnostics*
Let $X_{(1)} \leq \cdots \leq X_{(n)}$, $\overline{X} := n^{-1} \sum_i X_i$, and $s^2 := n^{-1} \sum_i (X_i - \overline{X})^2$.

(i) *Normal Q-Q plot*: plot $X_{(k)}$ against $\Phi^{-1}(\frac{k}{n+1})$; normality predicts approximate linearity.
(ii) *Skewness*: $b_1 := n^{-1} \sum_{i=1}^{n} \frac{(X_i - \overline{X})^3}{s^3}$ measures asymmetry; large $|b_1|$ indicates nonnormality.
(iii) *Kurtosis*: $b_2 := n^{-1} \sum_{i=1}^{n} \frac{(X_i - \overline{X})^4}{s^4}$ measures tail weight and peakedness; normality gives $b_2 \approx 3$.
(iv) *Plot correlation*: Q-Q point correlation; small values indicate curvature.

## 3.4 Normal Sampling Pivots

For normal samples, the sample mean and sample variance separate exactly. This is the finite-sample source of the classical $z$, $\chi^2$, and Student $t$ pivots used in normal-theory confidence intervals and tests.

**Corollary 3.4.1.** *Student Pivot for a Normal Mean*
*Let $X_1, \dots, X_n$ be independent; $X_i \sim N(\mu, \sigma^2)$, $\overline{X} := n^{-1} \sum_i X_i$, $S^2 := (n-1)^{-1} \sum_i (X_i - \overline{X})^2$; $\frac{\overline{X}-\mu}{\frac{S}{\sqrt{n}}} \sim t_{n-1}$.*

*Proof.* Let $Z := \frac{\sqrt{n}(\overline{X}-\mu)}{\sigma}$ and $V := \frac{(n-1)S^2}{\sigma^2}$. By normal mean law, $Z \sim N(0,1)$. By sample-variance chi-square law, $V \sim \chi^2_{n-1}$. By mean-residual independence, $Z \perp\!\!\!\perp V$. Therefore $\frac{\overline{X}-\mu}{\frac{S}{\sqrt{n}}} = \frac{Z}{\sqrt{\frac{V}{n-1}}} \sim t_{n-1}$.

*Structural Commentary*: Replacing $\sigma$ by $S$ yields a Student pivot.
*Key Insight*: Student's $t$ is a normal variable divided by an independent chi-square scale estimate. □

**Method 3.4.2.** *Exact Normal Mean Intervals*
Let $X_1, \dots, X_n$ be independent $N(\mu, \sigma^2)$ variables, with $\overline{X} := n^{-1} \sum_i X_i$ and $S^2 := (n-1)^{-1} \sum_i (X_i - \overline{X})^2$.
Below cases have exact coverage $1 - \alpha$.

(i) *Known variance*: if $\sigma$ is known, $C_\mu(X) = [\overline{X} - \frac{z_{1-\frac{\alpha}{2}}\sigma}{\sqrt{n}}, \overline{X} + \frac{z_{1-\frac{\alpha}{2}}\sigma}{\sqrt{n}}]$

(ii) *Unknown variance*: if $\sigma$ is unknown, $C_\mu(X) = [\overline{X} - \frac{t_{n-1,1-\frac{\alpha}{2}}S}{\sqrt{n}}, \overline{X} + \frac{t_{n-1,1-\frac{\alpha}{2}}S}{\sqrt{n}}]$

(iii) *Variance*: $C_{\sigma^2}(X) = [\frac{(n-1)S^2}{\chi^2_{n-1,1-\frac{\alpha}{2}}}, \frac{(n-1)S^2}{\chi^2_{n-1,\frac{\alpha}{2}}}]$

**Example 3.4.3.** *Normal Mean with Known Variance*
Let $X_1, \dots, X_n$ be independent $N(\mu, \sigma^2)$ variables with known $\sigma$.
The pivot $Z_\mu := \frac{\sqrt{n}(\overline{X}-\mu)}{\sigma} \sim N(0,1)$ gives $\mathbb{P}_\mu\{|Z_\mu| \le z_{1-\frac{\alpha}{2}}\} = 1 - \alpha$.
Solving for $\mu$ gives the exact interval $C_\alpha(X) = [\overline{X} - \frac{z_{1-\frac{\alpha}{2}}\sigma}{\sqrt{n}}, \overline{X} + \frac{z_{1-\frac{\alpha}{2}}\sigma}{\sqrt{n}}]$.
*Test Connection*: The two-sided level-$\alpha$ test of $H_0 : \mu = \mu_0$ rejects iff $\mu_0 \notin C_\alpha(X)$.

**Example 3.4.4.** *Normal Mean and Variance with Unknown Variance*
Let $X_1, \dots, X_n$ be independent $N(\mu, \sigma^2)$ variables, and set $S^2 := (n-1)^{-1} \sum_{i=1}^{n} (X_i - \overline{X})^2$.
The exact pivots are $T_\mu := \frac{\sqrt{n}(\overline{X}-\mu)}{S} \sim t_{n-1}$ and $W_{\sigma^2} := \frac{(n-1)S^2}{\sigma^2} \sim \chi^2_{n-1}$. Writing $t_{\nu,p}$ and $\chi^2_{\nu,p}$ for $p$-quantiles, pivot inversion gives $C_\mu(X) = [\overline{X} - \frac{t_{n-1,1-\frac{\alpha}{2}}S}{\sqrt{n}}, \overline{X} + \frac{t_{n-1,1-\frac{\alpha}{2}}S}{\sqrt{n}}]$ and $C_{\sigma^2}(X) = [\frac{(n-1)S^2}{\chi^2_{n-1,1-\frac{\alpha}{2}}}, \frac{(n-1)S^2}{\chi^2_{n-1,\frac{\alpha}{2}}}]$.
Both intervals have exact marginal coverage $1-\alpha$; $C_{\sigma^2}(X)$ is asymmetric because the $\chi^2$ pivot is not symmetric.

## 3.5 Two-Sample Normal Inference

Two-sample inference compares the location of two response distributions. In a controlled experiment, one sample may receive a treatment and the other a control; in an observational comparison, the same formulas may be computed but their causal interpretation depends on the design.

**Definition 3.5.1.** *Two Independent Samples and the Location Contrast*
$X_1, \dots, X_n \sim F$ are independent, $Y_1, \dots, Y_m \sim G$ are independent, and the two samples are independent.

(i) *Means*: $\mu_X := \mathbb{E}X_i$ and $\mu_Y := \mathbb{E}Y_j$ summarize the two average levels, assuming both exist.
(ii) *Location contrast*: $\Delta := \mu_X - \mu_Y$ measures the shift from group $Y$ to group $X$.
(iii) *Estimator*: $\widehat{\Delta} := \overline{X} - \overline{Y}$ estimates $\Delta$, where $\overline{X} := n^{-1} \sum_i X_i$ and $\overline{Y} := m^{-1} \sum_j Y_j$.

**Proposition 3.5.2.** *Known Common Variance Pivot*
*Let $X_1, \dots, X_n$ and $Y_1, \dots, Y_m$ be independent samples, $X_i \sim N(\mu_X, \sigma^2)$, $Y_j \sim N(\mu_Y, \sigma^2)$, known $\sigma > 0$.*
*Set $\Delta := \mu_X - \mu_Y$, $\widehat{\Delta} := \overline{X} - \overline{Y}$, and $\mathrm{se}_0 := \sigma\sqrt{n^{-1} + m^{-1}}$.*

*(i) $\widehat{\Delta} \sim N(\Delta, \mathrm{se}_0^2)$ and $Z := \frac{\widehat{\Delta}-\Delta}{\mathrm{se}_0} \sim N(0,1)$.*
*(ii) $C_\alpha(X,Y) := [\widehat{\Delta} - z_{1-\frac{\alpha}{2}}\,\mathrm{se}_0, \widehat{\Delta} + z_{1-\frac{\alpha}{2}}\,\mathrm{se}_0]$ satisfies $\mathbb{P}_\Delta\{\Delta \in C_\alpha(X,Y)\} = 1 - \alpha$.*

*Proof.*

(i) Since $\overline{X} \sim N(\mu_X, \frac{\sigma^2}{n})$, $\overline{Y} \sim N(\mu_Y, \frac{\sigma^2}{m})$, and $\overline{X} \perp\!\!\!\perp \overline{Y}$, $\widehat{\Delta} = \overline{X} - \overline{Y} \sim N(\Delta, \frac{\sigma^2}{n} + \frac{\sigma^2}{m}) = N(\Delta, \mathrm{se}_0^2)$. Standardization gives $Z \sim N(0,1)$.
(ii) By (i), $\mathbb{P}_\Delta\{|Z| \le z_{1-\frac{\alpha}{2}}\} = 1 - \alpha$, equivalently $\mathbb{P}_\Delta\{\Delta \in C_\alpha(X,Y)\} = 1 - \alpha$.

*Structural Commentary*: The contrast is a linear combination of independent sample means.
*Key Insight*: Known common variance gives an exact normal pivot for $\Delta$. □

**Definition 3.5.3.** *Pooled Sample Variance*
Let $X_1, \dots, X_n \sim N(\mu_X, \sigma^2)$ and $Y_1, \dots, Y_m \sim N(\mu_Y, \sigma^2)$ be independent samples with common unknown $\sigma^2$.

(i) *Sample variances*: $S_X^2 := (n-1)^{-1} \sum_{i=1}^{n} (X_i - \overline{X})^2$ and $S_Y^2 := (m-1)^{-1} \sum_{j=1}^{m} (Y_j - \overline{Y})^2$.
(ii) *Pooled variance*: $S_p^2 := \frac{(n-1)S_X^2 + (m-1)S_Y^2}{n+m-2}$.
(iii) *Degrees-of-freedom weighting*: $S_p^2$ weights $S_X^2$ and $S_Y^2$ by $n-1$ and $m-1$, respectively.

**Theorem 3.5.4.** *Pooled Two-Sample Student Pivot*
*Let $X_i \sim N(\mu_X, \sigma^2)$, $i = 1, \dots, n$, and $Y_j \sim N(\mu_Y, \sigma^2)$, $j = 1, \dots, m$, be mutually independent, with unknown common $\sigma^2 > 0$. Set $\Delta := \mu_X - \mu_Y$ and $\nu := n + m - 2$. Then $T := \frac{(\overline{X} - \overline{Y}) - \Delta}{S_p \sqrt{n^{-1} + m^{-1}}} \sim t_\nu$.*

*Proof.* Let $Z := \frac{(\overline{X} - \overline{Y}) - \Delta}{\sigma \sqrt{n^{-1} + m^{-1}}}$ and $W := \frac{\nu S_p^2}{\sigma^2} = \frac{(n-1)S_X^2 + (m-1)S_Y^2}{\sigma^2}$.
Since $\overline{X} - \overline{Y} \sim N(\Delta, \sigma^2(n^{-1} + m^{-1}))$, $Z \sim N(0,1)$. Let $U_X := \frac{(n-1)S_X^2}{\sigma^2}$ and $U_Y := \frac{(m-1)S_Y^2}{\sigma^2}$.
By normal mean-residual independence and sample independence, $Z, U_X, U_Y$ mutually independent, with $U_X \sim \chi^2_{n-1}$, $U_Y \sim \chi^2_{m-1}$. Hence $W = U_X + U_Y \sim \chi^2_\nu$ and $W \perp\!\!\!\perp Z$. Since $S_p^2 = \frac{\sigma^2 W}{\nu}$, $T = \frac{Z}{\sqrt{\frac{W}{\nu}}} \sim t_\nu$.
*Structural Commentary*: The pooled variance combines the two residual sums of squares.
*Key Insight*: A common variance permits replacing $\sigma$ by an independent pooled estimate. □

**Method 3.5.5.** *Pooled Two-Sample t Procedures*
Under the equal-variance normal model, set $\Delta := \mu_X - \mu_Y$, $\nu := n + m - 2$, and $\mathrm{se}_p := S_p \sqrt{n^{-1} + m^{-1}}$.

(i) *Interval*: $C_\alpha := [(\overline{X} - \overline{Y}) - t_{\nu, 1-\frac{\alpha}{2}} \mathrm{se}_p, (\overline{X} - \overline{Y}) + t_{\nu, 1-\frac{\alpha}{2}} \mathrm{se}_p]$ is exact $1 - \alpha$ for $\Delta$.
(ii) *Statistic*: for $H_0 : \Delta = 0$, use $t_0 := \frac{\overline{X} - \overline{Y}}{\mathrm{se}_p}$, with $t_0 \sim t_\nu$ under $H_0$.
(iii) *Rejection*: reject $H_0$ for $|t_0| > t_{\nu, 1-\frac{\alpha}{2}}$ under $H_1 : \Delta \neq 0$
reject $H_0$ for $t_0 > t_{\nu, 1-\alpha}$ under $H_1 : \Delta > 0$
reject $H_0$ for $t_0 < -t_{\nu, 1-\alpha}$ under $H_1 : \Delta < 0$.
(iv) *Duality*: the two-sided test rejects iff $0 \notin C_\alpha$.

**Example 3.5.6.** *Latent Heat of Fusion: Two Methods*
For methods $A$ and $B$, let $n_A = 13$, $n_B = 8$, $\overline{x}_A = 80.02$, $\overline{x}_B = 79.98$, $s_A = 0.024$, $s_B = 0.031$, and $S_p = 0.027$. Then $\widehat{\Delta} = \overline{x}_A - \overline{x}_B = 0.04$, $\widehat{\mathrm{se}} = S_p \sqrt{13^{-1} + 8^{-1}} \approx 0.012$, pooled 95% interval for $\Delta = \mu_A - \mu_B$ is $\approx (0.015, 0.065)$. For $H_0 : \Delta = 0$, $t_0 = \frac{\widehat{\Delta}}{\widehat{\mathrm{se}}} \approx 3.33$ with $\nu = 19$, strong evidence that method means differ.

**Proposition 3.5.7.** *Likelihood-Ratio View of the Pooled Test*
*Let $X_i \sim N(\mu_X, \sigma^2)$, $i = 1, \dots, n$, and $Y_j \sim N(\mu_Y, \sigma^2)$, $j = 1, \dots, m$, be mutually independent, with unknown common $\sigma^2 > 0$. Set $\Delta := \mu_X - \mu_Y$, $\nu := n + m - 2$, and $T := \frac{\overline{X} - \overline{Y}}{S_p \sqrt{n^{-1} + m^{-1}}}$.*
*For $H_0 : \Delta = 0$ against $H_1 : \Delta \neq 0$, $\lambda(X, Y) = (1 + \frac{T^2}{\nu})^{-\frac{n+m}{2}}$.*
*Hence the GLR rejects for large $|T|$; at level $\alpha$, reject iff $|T| > t_{\nu, 1-\frac{\alpha}{2}}$.*

*Proof.* Set $N := n + m$, $\nu := N - 2$, $d := \overline{X} - \overline{Y}$, $R_1 := \sum_i (X_i - \overline{X})^2 + \sum_j (Y_j - \overline{Y})^2$, $\widehat{\mu}_0 := \frac{n\overline{X} + m\overline{Y}}{N}$, and $R_0 := \sum_i (X_i - \widehat{\mu}_0)^2 + \sum_j (Y_j - \widehat{\mu}_0)^2$. Under $H_1$, $(\widehat{\mu}_X, \widehat{\mu}_Y, \widehat{\sigma}_1^2) = (\overline{X}, \overline{Y}, \frac{R_1}{N})$; under $H_0$, $(\widehat{\mu}, \widehat{\sigma}_0^2) = (\widehat{\mu}_0, \frac{R_0}{N})$.
The maximised Gaussian likelihood is proportional to $\widehat{\sigma}^{-N}$, hence $\lambda = (\frac{R_1}{R_0})^{\frac{N}{2}}$.
ANOVA decomposition gives $R_0 = R_1 + \{\frac{nm}{N}\} d^2$, while $T^2 = \frac{d^2}{(\frac{R_1}{\nu})(n^{-1} + m^{-1})} = \frac{\nu \{\frac{nm}{N}\} d^2}{R_1}$.
Therefore $\frac{R_0}{R_1} = 1 + \frac{T^2}{\nu}$ and $\lambda = (1 + \frac{T^2}{\nu})^{-\frac{N}{2}}$, decreasing in $T^2$; small $\lambda$ iff large $|T|$.
*Structural Commentary*: The null fit adds only the between-sample mean discrepancy to the unrestricted fit.
*Key Insight*: The GLR depends on the data through the pooled $t$ statistic. □

**Method 3.5.8.** *Unequal-Variance Two-Sample Procedure*
For independent samples without the equal-variance assumption, set $\widehat{v} := \frac{S_X^2}{n} + \frac{S_Y^2}{m}$.

(i) *Standard error*: estimate $\mathrm{Var}(\overline{X} - \overline{Y})$ by $\widehat{v}$.
(ii) *Welch statistic*: for $H_0 : \Delta = \Delta_0$, use $T_W := \frac{(\overline{X} - \overline{Y}) - \Delta_0}{\sqrt{\widehat{v}}}$.
(iii) *Degrees of freedom*: calibrate $T_W$ by $t_\nu$, where $\nu := \frac{\widehat{v}^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}}$.

(iv) *Use*: the procedure avoids pooling variances, at the cost of approximate calibration.

**Corollary 3.5.9.** *Variance Ratio Test Under Normality*
*Let* $X_1, \dots, X_n \sim N(\mu_X, \sigma_X^2)$ *and* $Y_1, \dots, Y_m \sim N(\mu_Y, \sigma_Y^2)$ *be independent samples, with* $n, m \geq 2$.
*Set* $\rho := \frac{\sigma_X^2}{\sigma_Y^2}$ *and* $R := \frac{S_X^2}{S_Y^2}$. *Then* $\frac{R}{\rho} \sim F_{n-1,m-1}$.

*Proof.* Let $U := \frac{(n-1)S_X^2}{\sigma_X^2}$ and $V := \frac{(m-1)S_Y^2}{\sigma_Y^2}$.
Normal residual chi-square laws give $U \sim \chi^2_{n-1}$ and $V \sim \chi^2_{m-1}$, and sample independence gives $U \perp\!\!\!\perp V$.
Thus $\frac{\frac{U}{n-1}}{\frac{V}{m-1}} = \frac{R}{\rho} \sim F_{n-1,m-1}$. If $H_0 : \rho = 1$, then $R \sim F_{n-1,m-1}$, yielding exact normal-theory tests and confidence intervals for $\rho$; the calibration is normality-sensitive.
*Structural Commentary*: The comparison standardises two independent residual variance estimates.
*Key Insight*: Normality turns each sample variance into an independent chi-square pivot. □

**Method 3.5.10.** *Approximate Power for Balanced Two-Sample z Tests*
For planning, assume balanced samples of size $n$, known common $\sigma$, and target effect $\Delta := \mu_X - \mu_Y > 0$.

(i) *Statistic*: use $Z := \frac{\overline{X} - \overline{Y}}{\sigma\sqrt{\frac{2}{n}}}$ and reject two-sided at level $\alpha$ when $|Z| > z_{1-\frac{\alpha}{2}}$.
(ii) *Shift*: under $\Delta > 0$, $Z \sim N(\delta, 1)$ with $\delta := (\frac{\Delta}{\sigma})\sqrt{\frac{n}{2}}$.
(iii) *Power*: $\pi(\Delta) := 1 - \Phi(z_{1-\frac{\alpha}{2}} - \delta) + \Phi(-z_{1-\frac{\alpha}{2}} - \delta)$.
(iv) *Sample size*: if the lower-tail term is negligible, target power $1 - \beta$ gives $n \approx \frac{2\sigma^2(z_{1-\frac{\alpha}{2}} + z_{1-\beta})^2}{\Delta^2}$.




Figure 242: Power rises as the detectable mean difference $\Delta$ increases.

**Remark 3.5.11.** *Transformations and Normal Diagnostics*
Normal-theory two-sample procedures are often CLT-robust for large samples, but small skewed or outlier-prone samples require diagnostics. Transformations may stabilize spread and improve symmetry, but they change the estimand, e.g. a log-scale mean difference corresponds to a ratio-scale comparison.

## 3.6 Paired-Sample Inference

Paired comparisons arise when observations are naturally matched: before and after measurements on the same unit, littermates, matched subjects, or two measurement methods applied to the same specimen. Pairing changes the stochastic structure because the two samples are no longer independent.

**Definition 3.6.1.** *Paired Differences*

(i) *Pairs*: $(X_i, Y_i)$, $i = 1, \dots, n$, are independent across $i$; dependence within each pair is allowed.
(ii) *Differences*: $D_i := X_i - Y_i$ and $\overline{D} := n^{-1} \sum_{i=1}^{n} D_i$.
(iii) *Mean*: if $\mathbb{E}X_i = \mu_X$ and $\mathbb{E}Y_i = \mu_Y$, then $\mathbb{E}D_i = \mu_X - \mu_Y$.
(iv) *Variance*: if $\mathrm{Var}(X_i) = \sigma_X^2$, $\mathrm{Var}(Y_i) = \sigma_Y^2$, and $\mathrm{Cov}(X_i, Y_i) = \sigma_{XY}$, then $\mathrm{Var}(D_i) = \sigma_X^2 + \sigma_Y^2 - 2\sigma_{XY}$.

**Proposition 3.6.2.** *Variance Effect of Pairing*
*Let* $(X_i, Y_i)$, $i = 1, \dots, n$, *be independent pairs with* $\mathrm{Var}(X_i) = \sigma_X^2$, $\mathrm{Var}(Y_i) = \sigma_Y^2$, *and* $\mathrm{Corr}(X_i, Y_i) = \rho$.
*Set* $D_i := X_i - Y_i$ *and* $\overline{D} := n^{-1} \sum_i D_i$. *Then* $\mathrm{Var}(\overline{D}) = n^{-1}(\sigma_X^2 + \sigma_Y^2 - 2\rho\sigma_X\sigma_Y)$.

*Proof.* Since $\mathrm{Cov}(X_i, Y_i) = \rho\sigma_X\sigma_Y$, $\mathrm{Var}(D_i) = \mathrm{Var}(X_i - Y_i) = \sigma_X^2 + \sigma_Y^2 - 2\rho\sigma_X\sigma_Y$.
Independence across pairs gives $\mathrm{Var}(\overline{D}) = n^{-2} \sum_i \mathrm{Var}(D_i) = \frac{\mathrm{Var}(D_i)}{n}$.

If $\sigma_X = \sigma_Y = \sigma$, substitution gives $\mathrm{Var}(\overline{D}) = \frac{2\sigma^2(1-\rho)}{n}$; independent unpaired samples of size $n$ per group have

$\mathrm{Var}(\overline{X}-\overline{Y})=\frac{2\sigma^2}{n}$, so the paired-to-unpaired variance ratio is $1-\rho$.
*Structural Commentary*: Pairing replaces two marginal measurements by one within-pair contrast.
*Key Insight*: Positive within-pair correlation cancels shared variation and lowers variance. □




Figure 243: Pairing cancels shared variation; positive within-pair correlation lowers variance.

**Method 3.6.3.** *Paired t Procedure*
Let $D_1,\dots,D_n\sim N(\mu_D,\sigma_D^2)$, where $\mu_D:=\mu_X-\mu_Y$.

(i) *Variance*: $S_D^2:=(n-1)^{-1}\sum_{i=1}^{n}(D_i-\overline{D})^2$.
(ii) *Pivot*: $T:=\frac{\overline{D}-\mu_D}{S_D/\sqrt{n}}\sim t_{n-1}$.
(iii) *Interval*: $C_\alpha:=[\overline{D}-\frac{t_{n-1,1-\frac{\alpha}{2}}S_D}{\sqrt{n}},\overline{D}+\frac{t_{n-1,1-\frac{\alpha}{2}}S_D}{\sqrt{n}}]$ is exact $1-\alpha$ for $\mu_D$.
(iv) *Test*: for $H_0:\mu_D=0$, reject two-sided at level $\alpha$ iff $|\overline{D}|>\frac{t_{n-1,1-\frac{\alpha}{2}}S_D}{\sqrt{n}}$.

For large $n$, use same rule by CLT; for small nonnormal differences, prefer rank or sign methods.

**Example 3.6.4.** *Platelet Aggregation Before and After Smoking*
For $n=11$ before-after platelet measurements, let $D_i=\text{after}_i-\text{before}_i$ with $\overline{D}=10.27$ and $\frac{S_D}{\sqrt{11}}=2.40$.
Then $t_0=\frac{\overline{D}}{\frac{S_D}{\sqrt{11}}}=4.28$ with $\nu=10$, so $p<0.01$ two-sided; a 90% confidence interval is approximately $(5.9,14.6)$, indicating a positive mean increase after smoking.

**Example 3.6.5.** *Mercury Measurement Methods*
For $n=25$ paired mercury measurements, let $D_i=M_{1i}-M_{2i}$ with $\overline{D}=0.04$ and $S_D=0.116$.
Then $t_0=\frac{\overline{D}}{\frac{S_D}{\sqrt{25}}}=1.724$ with $\nu=24$, giving $p\approx0.094$ two-sided.
Plot of $D_i$ against measurement level should be checked: level-dependent discrepancies not captured by $\overline{D}$ alone.

**Remark 3.6.6.** *Pairing Is Not Extra Independent Sampling*
If unit $i$ contributes repeated measurements $Z_{i1},\dots,Z_{ik}$, independence is across units, not within unit.
Treating all $nk$ observations as iid ignores within-unit dependence and overstates the effective sample size.

# 4 Decision Theory

## 4.1 Loss, Risk, Bayes, and Minimax Rules

Decision theory compares statistical procedures by their consequences, not only by their formulas. The same data model can lead to different optimal rules under different loss functions.

**Definition 4.1.1.** *Statistical Decision Problem and Risk*

(i) *Problem*: a statistical decision problem is $(\{P_\theta:\theta\in\Theta\},\mathcal{A},L)$ with $L:\Theta\times\mathcal{A}\to[0,\infty]$.
(ii) *Rule*: a decision rule is a measurable $\delta:\mathcal{X}\to\mathcal{A}$.
(iii) *Risk*: $R(\theta,\delta):=\mathbb{E}_\theta[L(\theta,\delta(X))]$.

**Definition 4.1.2.** *Bayes Risk, Minimaxity, and Admissibility*

(i) *Bayes risk*: for prior $\Pi$ on $\Theta$, $r(\Pi,\delta):=\int R(\theta,\delta)\,d\Pi(\theta)$.
(ii) *Bayes rule*: $\delta_\Pi$ is Bayes if $r(\Pi,\delta_\Pi)=\inf_\delta r(\Pi,\delta)$.
(iii) *Maximum risk*: $M(\delta):=\sup_{\theta\in\Theta}R(\theta,\delta)$; $\delta^\star$ is minimax if $M(\delta^\star)=\inf_\delta M(\delta)$.
(iv) *Admissibility*: $\delta$ inadmissible if $\exists\delta'$ with $R(\theta,\delta')\le R(\theta,\delta)\ \forall\theta$ and $<$ for some $\theta$; otherwise $\delta$ admissible.

**Proposition 4.1.3.** *Posterior Risk Characterization*
*Let $\Pi$ be a prior, let $\Pi(\cdot\mid x)$ be a posterior, and set $\rho_x(a):=\int L(\theta,a)\,d\Pi(\theta\mid x)$.*
*If a measurable selector $\delta_\Pi(x)\in\arg\min_{a\in\mathcal{A}}\rho_x(a)$ exists, then $\delta_\Pi$ is Bayes: $r(\Pi,\delta_\Pi)=\inf_\delta r(\Pi,\delta)$.*

*Proof.* By iterated expectation, $r(\Pi, \delta) = \mathbb{E}_\Pi[\rho_X(\delta(X))]$, where $X$ has the prior predictive law.
Since $\rho_x(\delta_\Pi(x)) \le \rho_x(\delta(x))$ $\forall$ rule $\delta$ and almost every $x$, $r(\Pi, \delta_\Pi) \le r(\Pi, \delta)$ $\forall \delta$. Thus $r(\Pi, \delta_\Pi) = \inf_\delta r(\Pi, \delta)$.
*Structural Commentary*: Bayes risk is the prior-predictive average of posterior risks.
*Key Insight*: Global Bayes optimization reduces to pointwise posterior-risk minimization. □

**Example 4.1.4.** *Bayes Rules Under Common Losses*
For a real parameter and posterior distribution $\Pi(\cdot \mid x)$:

(i) Squared-error loss is minimized by the posterior mean.
(ii) Absolute-error loss is minimized by a posterior median.
(iii) Zero-one loss on a finite action set is minimized by a posterior mode.

**Remark 4.1.5.** *Least Favorable Priors*
If a prior $\Pi^\star$ has Bayes risk equal to the minimax value, then any Bayes rule for $\Pi^\star$ with constant risk on the support of $\Pi^\star$ is minimax. Such priors are called least favorable.

## 4.2 Shrinkage, Empirical Bayes, and Regularization

Risk can often be reduced by intentionally pulling noisy estimates toward a structured target. The price is bias; the benefit is variance reduction.

**Definition 4.2.1.** *Shrinkage Estimator*

(i) *Input*: let $\widehat{\theta} = \widehat{\theta}(X)$ be a data-based estimator.
(ii) *Target*: choose $\theta_{\text{tar}}$ as a reference value or lower-complexity model.
(iii) *Weight*: let $a(X) \in [0, 1]$ control the amount of shrinkage.
(iv) *Estimator*: $\delta(X) := a(X)\widehat{\theta} + \{1 - a(X)\}\theta_{\text{tar}}$.

**Example 4.2.2.** *Ridge Regression as Shrinkage*
For design matrix $X$, response $y$, and $\lambda \ge 0$, ridge regression sets $\widehat{\beta}_\lambda \in \arg\min_\beta \{\|y - X\beta\|_2^2 + \lambda \|\beta\|_2^2\}$.
If $X^\top X$ is invertible, $\widehat{\beta}_\lambda = (X^\top X + \lambda I)^{-1} X^\top y$, shrinking $\widehat{\beta}_{\text{LS}}$ toward 0 most in weak eigendirections.

**Theorem 4.2.3.** *James–Stein Phenomenon*
*Let $X \sim \mathcal{N}_d(\theta, \sigma^2 I_d)$ with $d \ge 3$, known $\sigma^2$, and squared-error risk $R(\theta, \delta) := \mathbb{E}_\theta \|\delta(X) - \theta\|^2$.*
*Define $\delta_{\text{JS}}(X) := (1 - \frac{(d-2)\sigma^2}{\|X\|^2})X$ for $X \ne 0$, with arbitrary value at $X = 0$.*
*Then $R(\theta, \delta_{\text{JS}}) < R(\theta, X) = d\sigma^2$ for every $\theta \in \mathbb{R}^d$.*

*Proof.* Let $a := (d-2)\sigma^2$ and $g(x) := -\frac{ax}{\|x\|^2}$, so $\delta_{\text{JS}}(X) = X + g(X)$.
Stein's identity gives $R(\theta, X + g) = d\sigma^2 + \mathbb{E}_\theta[\|g(X)\|^2 + 2\sigma^2 \nabla \cdot g(X)]$.
For $x \ne 0$, $\|g(x)\|^2 = \frac{a^2}{\|x\|^2}$ and $\nabla \cdot g(x) = -\frac{a(d-2)}{\|x\|^2}$.
Thus $R(\theta, \delta_{\text{JS}}) = d\sigma^2 - (d-2)^2\sigma^4 \mathbb{E}_\theta[\|X\|^{-2}] < d\sigma^2$, with $\mathbb{E}_\theta[\|X\|^{-2}] < \infty$ since $d \ge 3$.
*Structural Commentary*: Shrinkage trades bias toward 0 for enough variance reduction to lower total risk.
*Key Insight*: In dimension $d \ge 3$, the usual estimator $X$ is inadmissible under squared-error loss. □

**Definition 4.2.4.** *Empirical Bayes*

(i) *Hierarchy*: $\theta_i \mid \eta \sim \Pi_\eta$ and $X_i \mid \theta_i \sim P_{\theta_i}$ for $i = 1, \dots, m$.
(ii) *Hyperparameter*: estimate $\eta$ by $\widehat{\eta} = \widehat{\eta}(X_1, \dots, X_m)$ from the marginal law of the data.
(iii) *Plug-in Bayes rule*: use the Bayes rule computed under $\Pi_{\widehat{\eta}}$.

**Remark 4.2.5.** *Regularization as Decision-Theoretic Structure*
Penalization, shrinkage, and empirical Bayes impose structure to reduce risk, typically by trading increased bias for reduced variance. The criterion is $R(\theta, \delta)$, not unbiasedness.

# 5 Bayesian Inference

## 5.1 Priors, Posteriors, and Conjugacy

Bayesian inference treats unknown parameters as uncertain quantities and updates a prior distribution by the likelihood supplied by the data.

**Definition 5.1.1.** *Prior, Posterior, and Predictive Distribution*
Let $\Pi$ be a prior on $\Theta$ and $p_\theta(x)$ a dominated likelihood.

(i) *Prior*: $\Pi$ encodes uncertainty about $\theta \in \Theta$ before observing $x$.
(ii) *Evidence*: $m(x) := \int_\Theta p_\vartheta(x)\, d\Pi(\vartheta)$, assumed finite and positive.
(iii) *Posterior*: $d\Pi(\theta \mid x) := \frac{p_\theta(x)\, d\Pi(\theta)}{m(x)}$.
(iv) *Predictive*: for future $X_{\text{new}}$, $\mathbb{P}(X_{\text{new}} \in A \mid x) = \int_\Theta P_\theta(A)\, d\Pi(\theta \mid x)$.

**Definition 5.1.2.** *Conjugate Prior Family*

(i) *Conjugacy*: $\{\Pi_\eta : \eta \in H\}$ is conjugate if $\Pi_\eta(\cdot \mid x) = \Pi_{\eta(x)}$ for some $\eta(x) \in H$.

**Proposition 5.1.3.** *Posterior Odds and Likelihood Ratios*
*Let $H_0, H_1$ be simple hypotheses with $\mathbb{P}(H_i) = \pi_i > 0$ and likelihood $p_i(x) = p(x \mid H_i)$ for $i = 0, 1$. For $m(x) := \pi_0 p_0(x) + \pi_1 p_1(x) > 0$, $\mathbb{P}(H_i \mid x) = \frac{\pi_i p_i(x)}{m(x)}$ and $\frac{\mathbb{P}(H_0 \mid x)}{\mathbb{P}(H_1 \mid x)} = (\frac{\pi_0}{\pi_1})(\frac{p_0(x)}{p_1(x)})$.*

*Proof.* Bayes' rule gives $\mathbb{P}(H_i \mid x) = \frac{\pi_i p_i(x)}{m(x)}$ for $i = 0, 1$. Therefore $\frac{\mathbb{P}(H_0 \mid x)}{\mathbb{P}(H_1 \mid x)} = \frac{\frac{\pi_0 p_0(x)}{m(x)}}{\frac{\pi_1 p_1(x)}{m(x)}} = (\frac{\pi_0}{\pi_1})(\frac{p_0(x)}{p_1(x)})$.
*Structural Commentary*: Posterior odds factor as prior odds times the data likelihood ratio.
*Key Insight*: Bayes updating multiplies odds by $\frac{p_0(x)}{p_1(x)}$. □

**Example 5.1.4.** *Beta–Binomial Updating*
If $X \mid p \sim \text{Binomial}(n, p)$ and $p \sim \text{Beta}(a, b)$, then $p \mid X = x \sim \text{Beta}(a + x, b + n - x)$.
Thus $\mathbb{E}[p \mid x] = \frac{a+x}{a+b+n}$, a weighted average of $\frac{a}{a+b}$ and $\frac{x}{n}$.

**Example 5.1.5.** *Gamma–Poisson Updating*
Let $X_1, \dots, X_n \mid \lambda$ be iid Poisson($\lambda$) and $\lambda \sim \text{Gamma}(\alpha, \nu)$ in rate form.
Writing $s := \sum_{i=1}^{n} x_i$, the posterior kernel is $\pi(\lambda \mid x) \propto \lambda^{\alpha+s-1} e^{-(\nu+n)\lambda}$, hence $\lambda \mid x \sim \text{Gamma}(\alpha + s, \nu + n)$.
Thus $\mathbb{E}[\lambda \mid x] = \frac{\alpha+s}{\nu+n}$, $\text{Var}(\lambda \mid x) = \frac{\alpha+s}{(\nu+n)^2}$, $\text{mode}(\lambda \mid x) = \frac{\alpha+s-1}{\nu+n}$ if $\alpha + s > 1$.

**Example 5.1.6.** *Normal Mean with Normal Prior*
Let $X_1, \dots, X_n \mid \mu$ be iid $N(\mu, \sigma^2)$ with known $\sigma^2$, and let $\mu \sim N(m_0, \tau_0^2)$.
Then $\mu \mid X \sim N(m_n, \tau_n^2)$, where $\tau_n^{-2} = \tau_0^{-2} + n\sigma^{-2}$ and $m_n = \tau_n^2(\tau_0^{-2} m_0 + n\sigma^{-2}\overline{X})$.

**Example 5.1.7.** *Known Mean and Unknown Precision*
Let $X_1, \dots, X_n \mid \xi$ be iid $N(\theta_0, \xi^{-1})$ with known $\theta_0$, and let $\xi \sim \text{Gamma}(\alpha, \lambda)$ in rate form.
Writing $q := \sum_{i=1}^{n} (x_i - \theta_0)^2$, the posterior is $\xi \mid x \sim \text{Gamma}(\alpha + \frac{n}{2}, \lambda + \frac{q}{2})$.
With a flat gamma prior, $\sigma^2 = \xi^{-1}$ centers near the MLE $\frac{q}{n}$.

**Definition 5.1.8.** *Improper Prior*
An *improper prior* is a $\sigma$-finite prior measure $\Pi$ on $\Theta$ with $\Pi(\Theta) = \infty$. For likelihood $p_\theta(x)$, it is usable only when $0 < m_\Pi(x) := \int_\Theta p_\theta(x)\, d\Pi(\theta) < \infty$, in which case $d\Pi(\theta \mid x) := \frac{p_\theta(x)\, d\Pi(\theta)}{m_\Pi(x)}$ is a proper posterior.

**Remark 5.1.9.** *Conditioning Viewpoint*
Bayesian updating is conditioning under the joint law $dP(\theta, x) = p_\theta(x)\, d\Pi(\theta)\, dx$; the posterior $\Pi(\cdot \mid x)$ is a regular conditional law of $\theta$ given $X = x$.

## 5.2 Computation, Prediction, and Asymptotic Bayes

Except in conjugate models, posterior summaries are integrals that usually require approximation. Monte Carlo and optimization methods from numerical computation become statistical tools here.

**Remark 5.2.1.** *Bayesian Integrals as a Motivation*
For prior density $\pi(\theta)$ and likelihood $\mathcal{L}(\theta)$, write $Z = \int \mathcal{L}(u)\pi(u)\, du$.
Then $\pi(\theta \mid X^n) = Z^{-1}\mathcal{L}(\theta)\pi(\theta)$ and $\mathbb{E}[\theta \mid X^n] = Z^{-1} \int \theta\, \mathcal{L}(\theta)\pi(\theta)\, d\theta$.
For $\theta = (\theta_1, \dots, \theta_k)$, high-dimensional marginalisation is often analytically impractical.

**Definition 5.2.2.** *Bayesian Credible Set*
Let $\Pi(\cdot \mid x)$ be the posterior on $\Theta$.

(i) *Credible set*: $C(x) \subseteq \Theta$ has credibility $1 - \alpha$ if $\Pi(C(x) \mid x) \geq 1 - \alpha$.
(ii) *Conditioning*: credibility is a posterior probability conditional on the observed data $x$.

**Definition 5.2.3.** *Posterior Monte Carlo Integration*
Let $\Pi(\cdot \mid x)$ be the posterior on $\Theta$.

(i) *Monte Carlo sample*: let $\theta^{(1)}, \dots, \theta^{(M)}$ be draws, or approximate draws, from $\Pi(\cdot \mid x)$.

(ii) *Posterior mean*: for integrable $g$, $\int g(\theta)\, d\Pi(\theta \mid x) \approx M^{-1} \sum_{m=1}^{M} g(\theta^{(m)})$.

(iii) *Dependent draws*: MCMC uses dependent samples but targets the same posterior integrals.

**Method 5.2.4.** *Simulating a Posterior Function*

(i) Let $\delta = g(p_1, \dots, p_k)$ be the posterior target.

(ii) Draw posterior samples $(P_1^{(i)}, \dots, P_k^{(i)})$ for $i = 1, \dots, N$.

(iii) Set $\delta^{(i)} = g(P_1^{(i)}, \dots, P_k^{(i)})$, use empirical distribution of $\delta^{(i)}$ to estimate posterior distribution of $\delta$.

**Example 5.2.5.** *Two Binomial Proportions*
Suppose $X \sim \text{Binomial}(n, p_1)$ and $Y \sim \text{Binomial}(m, p_2)$ are independent, with independent flat priors on $p_1, p_2$. Then $p_1 \mid X \sim \text{Beta}(X+1, n-X+1)$ and $p_2 \mid Y \sim \text{Beta}(Y+1, m-Y+1)$.
For $\delta = p_2 - p_1$, draw $P_1^{(i)} \sim \text{Beta}(X+1, n-X+1)$ and $P_2^{(i)} \sim \text{Beta}(Y+1, m-Y+1)$, set $\delta^{(i)} = P_2^{(i)} - P_1^{(i)}$, and summarize the empirical distribution of $\delta^{(i)}$.

**Method 5.2.6.** *Gibbs Sampling from Full Conditionals*

(i) *Kernel*: unknown mean $\theta$, precision $\xi$, flat-prior posterior satisfies $\pi(\theta, \xi \mid x) \propto \xi^{\frac{n}{2}-1} \exp\{-\frac{\xi}{2} \sum_i (x_i - \theta)^2\}$.

(ii) *Decomposition*: $\sum_i (x_i - \theta)^2 = \sum_i (x_i - \overline{x})^2 + n(\theta - \overline{x})^2$.

(iii) *Mean conditional*: $\theta \mid \xi, x \sim N(\overline{x}, (n\xi)^{-1})$.

(iv) *Precision conditional*: $\xi \mid \theta, x \sim \text{Gamma}(\frac{n}{2}, \frac{1}{2} \sum_i (x_i - \theta)^2)$ in rate form.

(v) *Gibbs sampler*: alternate conditionals; post-burn-in averages estimate posterior summaries.

**Method 5.2.7.** *Laplace Approximation*

(i) *Setup*: let $\pi_n(\theta \mid x) \propto \exp\{-n q_n(\theta)\}$, with interior minimiser $\widehat{\theta}$.

(ii) *Curvature*: set $H_n(\widehat{\theta}) := \nabla^2 q_n(\widehat{\theta})$, assumed positive definite.

(iii) *Quadratic approximation*: $q_n(\theta) \approx q_n(\widehat{\theta}) + \frac{1}{2}(\theta - \widehat{\theta})^\top H_n(\widehat{\theta})(\theta - \widehat{\theta})$.

(iv) *Normal approximation*: $\Pi(\cdot \mid x) \approx N(\widehat{\theta}, \{n H_n(\widehat{\theta})\}^{-1})$.

**Theorem 5.2.8.** *Bernstein–von Mises Principle*
*Let $X_1, \dots, X_n \sim P_{\theta_0}$ in a regular $d$-dimensional model with nonsingular information $I(\theta_0)$. Assume the prior density $\pi$ is positive and continuous near $\theta_0$, and let $\widehat{\theta}_n$ be an efficient center. Then $\Pi(\sqrt{n}(\theta - \widehat{\theta}_n) \in \cdot \mid X_1, \dots, X_n) \Rightarrow N_d(0, I(\theta_0)^{-1})$ in $P_{\theta_0}$-probability.*

*Proof.* Set $h := \sqrt{n}(\theta - \widehat{\theta}_n)$. Local asymptotic normality gives $\ell_n(\widehat{\theta}_n + \frac{h}{\sqrt{n}}) - \ell_n(\widehat{\theta}_n) = -\frac{1}{2} h^\top I(\theta_0) h + o_{P_{\theta_0}}(1)$ uniformly on bounded $h$. Prior continuity gives $\frac{\pi(\widehat{\theta}_n + \frac{h}{\sqrt{n}})}{\pi(\theta_0)} \to 1$ in $P_{\theta_0}$-probability on bounded $h$.
After normalization, the posterior density of $h$ converges to the $N_d(0, I(\theta_0)^{-1})$ density, with tail mass outside bounded sets negligible. Thus the displayed posterior weak convergence holds.
*Structural Commentary*: Locally, the likelihood is quadratic and the prior is flat to first order.
*Key Insight*: In regular finite-dimensional models, posterior uncertainty is asymptotically frequentist normal. □

**Remark 5.2.9.** *Alignment and Limits*
In regular fixed-dimensional models, Bayesian credible sets based on the Bernstein–von Mises normal approximation have asymptotic frequentist coverage. This alignment can fail in high-dimensional, nonparametric, weakly identified, or boundary problems, where the prior may remain first-order relevant.

## 5.3 Bayesian Normal Comparisons

The normal two-sample model also has a Bayesian interpretation. With diffuse priors, the resulting posterior formulas resemble frequentist $t$ intervals, but the conditioning and interpretation are different.

**Definition 5.3.1.** *Common-Precision Two-Sample Normal Model*

(i) *Common-Precision Model*: $\theta = (\mu_X, \mu_Y, \xi) \in \mathbb{R}^2 \times (0, \infty)$ with shared precision $\xi = \sigma^{-2}$ and variance $\xi^{-1}$.

(ii) *Conditional model*: Given $(\mu_X, \mu_Y, \xi)$, the variables $X_1, \dots, X_n, Y_1, \dots, Y_m$ are independent, with $X_i \mid (\mu_X, \xi) \sim N(\mu_X, \xi^{-1})$ and $Y_j \mid (\mu_Y, \xi) \sim N(\mu_Y, \xi^{-1})$.

(iii) *Improper prior*: $\pi(\mu_X, \mu_Y, \xi) \propto \xi^{-1}$ on $\mu_X, \mu_Y \in \mathbb{R}$ and $\xi > 0$.

(iv) *Observed data*: Write $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_m)$.

(v) *Sample means*: $\overline{x} := n^{-1} \sum_{i=1}^{n} x_i$ and $\overline{y} := m^{-1} \sum_{j=1}^{m} y_j$.

(vi) *Sample variances*: $s_X^2 := (n-1)^{-1} \sum_{i=1}^{n} (x_i - \overline{x})^2$ and $s_Y^2 := (m-1)^{-1} \sum_{j=1}^{m} (y_j - \overline{y})^2$.

(vii) *Pooled variance*: $s_p^2 := \frac{(n-1)s_X^2 + (m-1)s_Y^2}{m+n-2}$.

(viii) *Posterior kernel*: $\pi(\mu_X, \mu_Y, \xi \mid x, y) \propto \xi^{\frac{m+n}{2}-1} \exp\left\{ -\frac{\xi}{2} \left[ \sum_{i=1}^{n} (x_i - \mu_X)^2 + \sum_{j=1}^{m} (y_j - \mu_Y)^2 \right] \right\}$.

**Proposition 5.3.2.** *Conditional Posterior Factors*
*For $n, m \geq 2$, put $\nu := m + n - 2$ and take* Gamma *shape-rate.*
*In the common-precision model, $\pi(\mu_X, \mu_Y, \xi \mid x, y) = \pi(\mu_X \mid \xi, x, y)\pi(\mu_Y \mid \xi, x, y)\pi(\xi \mid x, y)$,*
*where $(\mu_X, \mu_Y) \mid \xi, x, y \sim N(\overline{x}, (n\xi)^{-1}) \otimes N(\overline{y}, (m\xi)^{-1})$ and $\xi \mid x, y \sim \text{Gamma}(\frac{\nu}{2}, \frac{\nu s_p^2}{2})$. $\mu_X \perp\!\!\!\perp \mu_Y \mid (\xi, x, y)$.*

*Proof.* Let $S_X(\mu_X) := \sum_{i=1}^{n} (x_i - \mu_X)^2$, $S_Y(\mu_Y) := \sum_{j=1}^{m} (y_j - \mu_Y)^2$, and $A := (n-1)s_X^2 + (m-1)s_Y^2 = \nu s_p^2$.
Completing squares gives $S_X(\mu_X) = (n-1)s_X^2 + n(\mu_X - \overline{x})^2$ and $S_Y(\mu_Y) = (m-1)s_Y^2 + m(\mu_Y - \overline{y})^2$.
Thus $\pi(\mu_X, \mu_Y, \xi \mid x, y) \propto \xi^{\frac{m+n}{2}-1} \exp\{-\frac{\xi A}{2}\} \exp\{-\frac{n\xi(\mu_X - \overline{x})^2}{2}\} \exp\{-\frac{m\xi(\mu_Y - \overline{y})^2}{2}\}$.
For fixed $\xi$, last two factors are independent normal kernels for $\mu_X$, $\mu_Y$ with variances $(n\xi)^{-1}$, $(m\xi)^{-1}$.
Also $\int \exp\{-\frac{n\xi(\mu_X - \overline{x})^2}{2}\} \, d\mu_X \propto (n\xi)^{-\frac{1}{2}}$ and $\int \exp\{-\frac{m\xi(\mu_Y - \overline{y})^2}{2}\} \, d\mu_Y \propto (m\xi)^{-\frac{1}{2}}$.
Hence marginalising $\mu_X, \mu_Y$ leaves $\pi(\xi \mid x, y) \propto \xi^{\frac{\nu}{2}-1} \exp\{-\frac{\xi \nu s_p^2}{2}\}$.
*Structural Commentary*: Completing squares isolates location kernels and pooled residual $A = \nu s_p^2$.
*Key Insight*: Marginalising $\mu_X, \mu_Y$ leaves the pooled gamma law for $\xi$. □

**Proposition 5.3.3.** *Posterior t Law for a Mean Difference*
*In the common-precision model with $n, m \geq 2$, set $\Delta := \mu_X - \mu_Y$, $d := \overline{x} - \overline{y}$, $c := n^{-1} + m^{-1}$, $\nu := m + n - 2$, and $T := \frac{\Delta - d}{s_p \sqrt{c}}$. Then $T \mid x, y \sim t_\nu$.*

*Proof.* By Proposition 5.3.2, $\Delta \mid \xi, x, y \sim N(d, \frac{c}{\xi})$ and $\xi \mid x, y \sim \text{Gamma}(\frac{\nu}{2}, \frac{\nu s_p^2}{2})$.
Equivalently, take independent $E \sim N(0,1)$, $V \sim \chi^2_\nu$, $\xi = \frac{V}{\nu s_p^2}$, $\Delta = d + \sqrt{\frac{c}{\xi}}\, E$. Thus $T = \frac{\Delta - d}{s_p \sqrt{c}} = \frac{E}{\sqrt{\frac{V}{\nu}}}$, QED.
*Structural Commentary*: Normal given $\xi$; Student after mixing over $\xi$.
*Key Insight*: The $t$ law is exactly a standard normal divided by an independent estimated-scale factor $\sqrt{\frac{V}{\nu}}$. □

**Remark 5.3.4.** *Same Algebra, Different Conditioning*
The statistic has the same algebra as the pooled two-sample $t$ pivot in Theorem 3.5.4: frequentist inference treats $\Delta$ as fixed and $X, Y$ as random, whereas the posterior treats $x, y$ as fixed and $\Delta \mid x, y$ as random.

**Example 5.3.5.** *Latent Heat Posterior Comparison*
For method contrast $\Delta = \mu_X - \mu_Y$, the common-precision posterior gives $\mathbb{P}(\Delta > 0 \mid X, Y) \approx 0.998$ and posterior interval $\mathbb{P}(0.015 \leq \Delta \leq 0.065 \mid X, Y) \approx 0.95$, numerically matching the pooled normal-theory interval.

# 6 Nonparametric and High-Dimensional Statistics

## 6.1 Empirical Processes, Density, and Regression Smoothing

Nonparametric statistics replaces finite-dimensional parameter assumptions by smoothness, shape, sparsity, or complexity restrictions.

**Definition 6.1.1.** *Empirical Measure and Empirical Process*
Let $X_1, \dots, X_n$ be observations in $(\mathcal{X}, \mathcal{A})$ with reference law $P$.

(i) *Empirical measure*: $\widehat{P}_n := n^{-1} \sum_{i=1}^{n} \delta_{X_i}$, so $\widehat{P}_n f = n^{-1} \sum_{i=1}^{n} f(X_i)$.

(ii) *Empirical process*: For measurable $\mathcal{F}$, $\mathbb{G}_n(f) := \sqrt{n}(\widehat{P}_n f - Pf)$ for $f \in \mathcal{F}$.

**Proposition 6.1.2.** *Fixed-Point Behavior of the Empirical CDF*
*Let $X_1, \dots, X_n$ be independent with common CDF $F$.*
*For $x \in \mathbb{R}$, set $I_i(x) := \mathbf{1}_{(-\infty, x]}(X_i)$ and $F_n(x) := n^{-1} \sum_{i=1}^{n} I_i(x)$. Then, for each fixed $x \in \mathbb{R}$:*

*(i) Law:* $nF_n(x) \sim \text{Binomial}(n, F(x))$.

*(ii) Mean:* $\mathbb{E}F_n(x) = F(x)$.
*(iii) Variance:* $\mathrm{Var}(F_n(x)) = n^{-1}F(x)(1-F(x))$.

*Proof.* Fix $x \in \mathbb{R}$ and put $p := F(x)$ and $I_i := I_i(x)$.

(i) Since $I_i \sim \mathrm{Bernoulli}(p)$ independently, $nF_n(x) = \sum_{i=1}^{n} I_i \sim \mathrm{Binomial}(n,p)$.

(ii) Linearity gives $\mathbb{E}F_n(x) = n^{-1}\sum_{i=1}^{n}\mathbb{E}I_i = n^{-1}np = p = F(x)$.

(iii) Independence gives $\mathrm{Var}(F_n(x)) = n^{-2}\sum_{i=1}^{n}\mathrm{Var}(I_i) = n^{-2}np(1-p) = n^{-1}F(x)(1-F(x))$.

*Structural Commentary*: Pointwise, $F_n(x) = n^{-1}\sum_{i=1}^{n} I_i(x)$ with $I_i(x) \sim \mathrm{Bernoulli}(F(x))$ independently.
*Key Insight*: The empirical CDF at one point is a normalised binomial count. □

**Remark 6.1.3.** *Pointwise and Simultaneous CDF Uncertainty*
Proposition 6.1.2 is pointwise: $x$ is fixed and $F_n(x)$ is a normalised binomial count.
Simultaneous uncertainty controls $\|F_n - F\|_\infty := \sup_x |F_n(x) - F(x)|$, yielding bands and diagnostics.

**Proposition 6.1.4.** *Two-Point Covariance of the Empirical CDF*
*Let $X_1, \dots, X_n$ be independent with common CDF $F$.*
*For $w \in \mathbb{R}$, set $I_i(w) := \mathbf{1}_{(-\infty,w]}(X_i)$, $F_n(w) := n^{-1}\sum_{i=1}^{n} I_i(w)$. For $u, v \in \mathbb{R}$, put $m := \min\{u,v\}$. Then:*

*(i) Covariance:* $\mathrm{Cov}\big(F_n(u), F_n(v)\big) = n^{-1}\big(F(m) - F(u)F(v)\big)$.
*(ii) Sign:* $\mathrm{Cov}\big(F_n(u), F_n(v)\big) \geq 0$.

*Proof.* Fix $u, v \in \mathbb{R}$ and set $m := \min\{u,v\}$.

(i) *Covariance*: $\mathrm{Cov}(F_n(u), F_n(v)) = n^{-2}\sum_{i,j}\mathrm{Cov}(I_i(u), I_j(v))$.
For $i \neq j$ the term is 0 by independence, while for $i = j$,
$\mathrm{Cov}(I_i(u), I_i(v)) = \mathbb{E}[I_i(u)I_i(v)] - \mathbb{E}I_i(u)\mathbb{E}I_i(v) = F(m) - F(u)F(v)$.
Hence $\mathrm{Cov}(F_n(u), F_n(v)) = n^{-1}(F(m) - F(u)F(v))$.

(ii) *Sign*: If $u \leq v$, then $F(m) - F(u)F(v) = F(u)(1 - F(v)) \geq 0$.
If $v \leq u$, then $F(m) - F(u)F(v) = F(v)(1 - F(u)) \geq 0$.

*Structural Commentary*: $F_n(u)$ and $F_n(v)$ average iid Bernoulli pairs, so covariance scales by $1/n$.
*Key Insight*: Cross-index independence leaves only the same-observation overlap $\{X_i \leq m\}$. □

**Theorem 6.1.5.** *Glivenko–Cantelli Theorem*
*Let $X_1, X_2, \dots$ be i.i.d. real variables with CDF $F$, and set $\widehat{F}_n(t) := n^{-1}\sum_{i=1}^{n}\mathbf{1}_{\{X_i \leq t\}}$.*
*Then $\sup_{t\in\mathbb{R}}\left|\widehat{F}_n(t) - F(t)\right| \to 0$ almost surely.*

*Proof.* Add independent $R_i \sim \mathrm{Unif}[0,1]$ and set $U_i := F(X_i-) + R_i\{F(X_i) - F(X_i-)\}$.
Then $U_i$ are i.i.d. Unif$[0,1]$, and with $H_n(u) := n^{-1}\sum_i \mathbf{1}_{\{U_i \leq u\}}$ one has $\widehat{F}_n(t) = H_n(F(t))$ a.s.

Hence $\sup_t\left|\widehat{F}_n(t) - F(t)\right| \leq \sup_{0\leq u\leq 1}|H_n(u) - u|$.
Fix $k \geq 1$, $u_j := \frac{j}{k}$, let $M_{n,k} := \max_{0\leq j\leq k}|H_n(u_j) - u_j|$.
By the SLLN, $M_{n,k} \to 0$ a.s. If $u_{j-1} \leq u \leq u_j$, monotonicity gives $H_n(u_{j-1}) - u_j \leq H_n(u) - u \leq H_n(u_j) - u_{j-1}$,
Hence $\sup_{0\leq u\leq 1}|H_n(u) - u| \leq M_{n,k} + k^{-1}$, and $\limsup_{n\to\infty}\sup_{0\leq u\leq 1}|H_n(u) - u| \leq k^{-1}$ a.s. $\forall k$. Let $k \to \infty$.
*Structural Commentary*: Finite-grid SLLN plus monotonicity gives uniform CDF control.
*Key Insight*: Monotonicity bounds between-grid drift by the mesh. □

**Proposition 6.1.6.** *Approximate Variance of a Sample Quantile*
*Let $Y_1, \dots, Y_n$ be iid with continuous CDF $F$ and density $f$. Fix $p \in (0,1)$, set $x_p := F^{-1}(p)$ with $f(x_p) > 0$, choose $k$ with $\frac{k}{n+1} \approx p$, and put $\widehat{x}_p := Y_{(k)}$. Then:*

*(i) Probability Scale:* *$U_{(k)} := F(Y_{(k)})$ has $k$th uniform order-statistic law.* $\mathrm{Var}(U_{(k)}) = \frac{k(n+1-k)}{(n+1)^2(n+2)} \approx \frac{p(1-p)}{n}$.
*(ii) Quantile Scale:* $\mathrm{Var}(\widehat{x}_p) \approx \frac{p(1-p)}{nf(x_p)^2}$.

*Proof.*

(i) By probability integral transform, $U_i := F(Y_i)$ are iid Uniform$(0,1)$, $U_{(k)} = F(Y_{(k)})$. Theorem 4.1.4 gives the stated exact variance, hence $\operatorname{Var}(U_{(k)}) \approx \frac{p(1-p)}{n}$ when $\frac{k}{n+1} \approx p$.

(ii) Let $g := F^{-1}$. Since $g'(p) = \frac{1}{f(x_p)}$ and $\widehat{x}_p = g(U_{(k)})$, delta approximation $g(U_{(k)}) \approx g(p) + g'(p)(U_{(k)} - p)$ gives $\operatorname{Var}(\widehat{x}_p) \approx g'(p)^2 \operatorname{Var}(U_{(k)}) \approx \frac{p(1-p)}{n f(x_p)^2}$.

*Structural Commentary*: Quantile variance is probability-scale variance multiplied by $(F^{-1})'(p)^2$.
*Key Insight*: Low density near $x_p$ inflates sample-quantile variability by $1/f(x_p)^2$. □

**Proposition 6.1.7.** *Distribution-Free Median Interval*
*Let $X_1, \dots, X_n$ be independent with continuous CDF $F$ and median $\eta$, so $F(\eta) = \frac{1}{2}$.*
*Let $X_{(1)} \le \cdots \le X_{(n)}$ and $1 \le k \le \lfloor \frac{n}{2} \rfloor$. Then:*

*(i) Interval: $C_k := [X_{(k)}, X_{(n-k+1)}]$.*

*(ii) Coverage: $\mathbb{P}(\eta \in C_k) = 1 - \frac{1}{2^{n-1}} \sum_{j=0}^{k-1} \binom{n}{j}$.*

*Proof.*

(i) Let $B_- := \sum_{i=1}^{n} \mathbf{1}_{\{X_i < \eta\}}$, $B_+ := \sum_{i=1}^{n} \mathbf{1}_{\{X_i > \eta\}}$. By continuity, $B_- + B_+ = n$ a.s. and $B_- \sim \text{Binomial}(n, \frac{1}{2})$.

(ii) $\eta \notin C_k$ iff $B_- < k$ or $B_+ < k$. Since $k \le \lfloor \frac{n}{2} \rfloor$, these two events are disjoint.

(iii) By symmetry, $\mathbb{P}(\eta \notin C_k) = 2\mathbb{P}(B_- < k) = 2 \sum_{j=0}^{k-1} \binom{n}{j} 2^{-n} = \frac{1}{2^{n-1}} \sum_{j=0}^{k-1} \binom{n}{j}$, so $\mathbb{P}(\eta \in C_k) = 1 - \frac{1}{2^{n-1}} \sum_{j=0}^{k-1} \binom{n}{j}$.

*Structural Commentary*: The coverage depends only on the binomial count of observations below $\eta$, not on $F$.
*Key Insight*: Continuity turns median coverage into a symmetric binomial tail. □

**Definition 6.1.8.** *Kernel Density Estimator*

(i) *Kernel and bandwidth*: Let $\int K(u)\,du = 1$ and $h > 0$.

(ii) *Estimator*: For data $X_1, \dots, X_n$, $\widehat{f}_h(x) := (nh)^{-1} \sum_{i=1}^{n} K(\frac{x - X_i}{h})$.

Larger $h$ lowers variance and raises bias; smaller $h$ does the reverse.

**Definition 6.1.9.** *Nonparametric Regression*

(i) *Model*: $Y = m(X) + \varepsilon$ with $\mathbb{E}[\varepsilon \mid X] = 0$, so $m(x) = \mathbb{E}[Y \mid X = x]$.

(ii) *Weights*: For kernel $K$ and bandwidth $h > 0$, set $w_i(x) := K(\frac{x - X_i}{h})$.

(iii) *Estimator*: If $\sum_i w_i(x) > 0$, define $\widehat{m}_h(x) := \frac{\sum_{i=1}^{n} w_i(x) Y_i}{\sum_{i=1}^{n} w_i(x)}$.

**Definition 6.1.10.** *Kernel-Weighted Local Average*
Let $(X_i, Y_i)_{i=1}^n$ be regression data and fix a query point $x_0$.

(i) *Local kernel*: For a nonnegative profile $D$ and width $h_\lambda(x_0) > 0$, set $K_\lambda(x_0, x) := D(\|x - x_0\| / h_\lambda(x_0))$.

(ii) *Estimator*: If $\sum_{i=1}^{n} K_\lambda(x_0, X_i) > 0$, local-constant (Nadaraya–Watson) estimate is $\widehat{m}(x_0) := \frac{\sum_{i=1}^{n} K_\lambda(x_0, X_i) Y_i}{\sum_{i=1}^{n} K_\lambda(x_0, X_i)}$.

Metric windows set $h_\lambda(x_0) = \lambda$; nearest-neighbor windows set $h_\lambda(x_0) = \operatorname{dist}(x_0, X_{(k)}(x_0))$.




Figure 244: Fixed metric and $k$-nearest windows around $x_0$.

**Example 6.1.11.** *Common Local-Smoothing Kernels*
In one dimension, common compact profiles are $D_E(t) := \frac{3}{4}(1 - t^2)\mathbf{1}_{\{|t| \le 1\}}$ for Epanechnikov smoothing and $D_T(t) := (1 - |t|^3)^3 \mathbf{1}_{\{|t| \le 1\}}$ for tri-cube smoothing. Both have support $[-1, 1]$; Gaussian profiles have full support, and normalization constants cancel in regression-weight averages but matter for density estimation.

**Method 6.1.12.** *Local Linear Regression Smoothing*
Fix $x_0$ and bandwidth $h > 0$.

(i) *Weighted fit*: Set $(\widehat{\beta}_0(x_0), \widehat{\beta}_1(x_0)) \in \operatorname*{argmin}_{\beta_0,\beta_1} \sum_{i=1}^{n} (Y_i - \beta_0 - \beta_1(X_i - x_0))^2 K(\frac{X_i - x_0}{h})$.

(ii) *Estimate*: Define $\widehat{m}_h(x_0) := \widehat{\beta}_0(x_0)$.

Smaller $h$ gives a more variable fit; larger $h$ gives more bias.

**Method 6.1.13.** *Local Polynomial Regression*
Fix a query point $x_0 \in \mathbb{R}^p$ and degree $d \in \mathbb{N}_0$.

(i) *Basis*: Let $\mathcal{A}_d := \{\alpha \in \mathbb{N}_0^p : |\alpha| \le d\}$ and $b_d(u) := (u^\alpha)_{\alpha \in \mathcal{A}_d}$, where $u^\alpha := \prod_{j=1}^{p} u_j^{\alpha_j}$.

(ii) *Weights*: Set $K_i(x_0) := K_\lambda(x_0, X_i)$.

(iii) *Weighted fit*: For $\beta = (\beta_\alpha)_{\alpha \in \mathcal{A}_d}$, choose $\widehat{\beta}(x_0) \in \operatorname*{argmin}_{\beta \in \mathbb{R}^{|\mathcal{A}_d|}} \sum_{i=1}^{n} K_i(x_0)(Y_i - b_d(X_i - x_0)^\top \beta)^2$.

(iv) *Estimator*: With $\mathbf{0} \in \mathcal{A}_d$ the constant multi-index, define $\widehat{m}_d(x_0) := \widehat{\beta}_{\mathbf{0}}(x_0)$.

$d = 0$ is local constant; $d = 1$ is local linear with boundary correction.
Larger $d$ trades lower curvature bias for higher variance.

**Proposition 6.1.14.** *Equivalent Kernel Moments*
*Fix $x_0 \in \mathbb{R}$ and set $u_i := X_i - x_0$, $K_i := K_\lambda(x_0, X_i)$, $z_i := (1, u_i)^\top$, $Z := (z_1^\top, \dots, z_n^\top)^\top$, $W := \operatorname{diag}(K_1, \dots, K_n)$, $Y := (Y_1, \dots, Y_n)^\top$, and $G := Z^\top W Z$.*
*Assume $G$ is nonsingular, let $e_1 := (1, 0)^\top$, $\widehat{\beta}(x_0) := G^{-1} Z^\top W Y$, and $\widehat{m}(x_0) := e_1^\top \widehat{\beta}(x_0)$.*

*(i) Equivalent weights: $\widehat{m}(x_0) = \sum_{i=1}^{n} \ell_i(x_0) Y_i$ with $\ell_i(x_0) := K_i z_i^\top G^{-1} e_1$.*

*(ii) Moment constraints: $\sum_{i=1}^{n} \ell_i(x_0) = 1$ and $\sum_{i=1}^{n} u_i \ell_i(x_0) = 0$.*

*(iii) Affine reproduction: If $Y_i = a + bX_i$, then $\widehat{m}(x_0) = a + bx_0$.*

*Proof.*

(i) *Equivalent weights*: The weighted criterion $(Y - Z\beta)^\top W (Y - Z\beta)$ has normal equations $G\widehat{\beta} = Z^\top W Y$, so $\widehat{\beta} = G^{-1} Z^\top W Y$. Since $G^{-1}$ is symmetric, $\widehat{m}(x_0) = e_1^\top G^{-1} Z^\top W Y = \sum_{i=1}^{n} K_i e_1^\top G^{-1} z_i Y_i = \sum_{i=1}^{n} \ell_i(x_0) Y_i$.

(ii) *Moment constraints*: With $\ell(x_0) := (\ell_1(x_0), \dots, \ell_n(x_0))^\top = W Z G^{-1} e_1$, $Z^\top \ell(x_0) = Z^\top W Z G^{-1} e_1 = e_1$, hence $\sum_i \ell_i(x_0) = 1$ and $\sum_i u_i \ell_i(x_0) = 0$.

(iii) *Affine reproduction*: If $Y_i = a + bX_i = (a + bx_0) + bu_i$, then $\widehat{m}(x_0) = (a + bx_0) \sum_i \ell_i(x_0) + b \sum_i u_i \ell_i(x_0) = a + bx_0$.

*Structural Commentary*: Weighted projection onto $\operatorname{span}\{1, x - x_0\}$ imposes two weight moments.
*Key Insight*: $Z^\top \ell = e_1$ encodes constant reproduction and zero first moment. □

**Definition 6.1.15.** *Structured Kernel Metric*
Fix $x_0, x \in \mathbb{R}^p$, bandwidth $\lambda > 0$, profile $D \ge 0$, and a positive semidefinite matrix $A \succeq 0$.

(i) *Quadratic scale*: Define $\rho_A(x, x_0) := (x - x_0)^\top A (x - x_0)$.

(ii) *Kernel*: Set $K_{\lambda,A}(x_0, x) := D(\rho_A(x, x_0)/\lambda)$.

$A$ sets coordinate weights, correlations, and active directions.
Standardise predictors first since $\rho_A$ is unit-dependent.

**Method 6.1.16.** *Local Likelihood and Varying Coefficients*
Let $\ell_i(\theta)$ be the log-likelihood contribution of observation $i$ under parameter $\theta$.

(i) *Local likelihood*: At target $x_0$, set $K_i(x_0) := K_\lambda(x_0, X_i)$ and choose $\widehat{\theta}(x_0) \in \operatorname*{argmax}_{\theta} \sum_{i=1}^{n} K_i(x_0) \ell_i(\theta)$.

(ii) *Local parameterisation*: In generalised models, replace a constant natural parameter by $\eta_i(\gamma; x_0) := b_d(X_i - x_0)^\top \gamma$ and maximize $\sum_i K_i(x_0) \ell_i(\eta_i(\gamma; x_0))$ over $\gamma$.

(iii) *Varying coefficients*: For predictors $(U, Z)$ with $U \in \mathbb{R}^q$, model $m(U, Z) := \alpha(Z) + \sum_{r=1}^{q} \beta_r(Z) U_r$.

(iv) *Local fit*: At $z_0$, choose $(\widehat{\alpha}, \widehat{\beta}) \in \operatorname*{argmin}_{a,b} \sum_i K_\lambda(z_0, Z_i)(Y_i - a - b^\top U_i)^2$ and set $(\widehat{\alpha}(z_0), \widehat{\beta}(z_0)) := (\widehat{\alpha}, \widehat{\beta})$.

The coefficients are ordinary regression coefficients locally in $U$, but functions of the index variable $Z$.

**Remark 6.1.17.** *Bandwidth, Boundaries, and Dimension*
Let $N_\lambda(x_0) := \{i : K_\lambda(x_0, X_i) > 0\}$ and $n_\lambda(x_0) := |N_\lambda(x_0)|$.

(i) *Bias–variance*: Larger $\lambda$ usually increases $n_\lambda(x_0)$, lowering variance, raising bias; smaller $\lambda$ does reverse.
(ii) *Windows*: Metric windows fix radius; $k$-nearest-neighbor windows choose $\lambda(x_0)$ so that $n_{\lambda(x_0)}(x_0) \approx k$.
(iii) *Dimension*: In $\mathbb{R}^p$, neighborhoods must expand with $p$ to keep $n_\lambda(x_0)$ stable, reducing local resolution.
(iv) *Cost*: Without indexing, one query evaluates $K_\lambda(x_0, X_i)$ for all $i$, costing $O(n)$.

**Method 6.1.18.** *Bandwidth Choice by Leave-One-Out Cross-Validation*

(i) *Candidates*: Fix bandwidths $h_1, \dots, h_M > 0$.
(ii) *Leave-one-out score*: Let $\widehat{m}_{h_k,-i}$ be local linear fit excluding $(X_i, Y_i)$. Set $\mathrm{CV}(h_k) := \sum_{i=1}^{n} (Y_i - \widehat{m}_{h_k,-i}(X_i))^2$.
(iii) *Selection*: Choose $\widehat{h} \in \operatorname*{argmin}_{1 \le k \le M} \mathrm{CV}(h_k)$, with a visual smoothness check.

**Remark 6.1.19.** *Complexity Control*
For a function class $\mathcal{F}$, estimation is stable when $\sup_{f \in \mathcal{F}} \left| \widehat{P}_n f - Pf \right|$ is small with high probability or a.s.
Smoothness and entropy control this supremum in place of finite dimension.

## 6.2 Sparse High-Dimensional Inference

High-dimensional statistics studies regimes where the number of candidate parameters is comparable to, or much larger than, the sample size. Estimation is possible only after adding structure.

**Definition 6.2.1.** *High-Dimensional Regime and Sparsity*

(i) *High dimension*: A sequence has dimension $p = p_n$, often with $p_n \gg n$.
(ii) *Sparsity*: For $\theta \in \mathbb{R}^p$, $\|\theta\|_0 := |\{j : \theta_j \neq 0\}|$; $\theta$ is $s$-sparse if $\|\theta\|_0 \le s$.

**Definition 6.2.2.** *Lasso Estimator*

(i) *Lasso*: For $y = X\beta_0 + \varepsilon$ and $\lambda \ge 0$, $\widehat{\beta}_\lambda \in \arg\min_{\beta \in \mathbb{R}^p} \{(2n)^{-1} \|y - X\beta\|_2^2 + \lambda \|\beta\|_1\}$.

The convex $\ell^1$ penalty promotes sparsity.

**Definition 6.2.3.** *Restricted Eigenvalue Condition*

(i) *Restricted eigenvalue*: For $S \subseteq [p] := \{1, \dots, p\}$, $C(S) := \{v : \|v_{S^c}\|_1 \le 3 \|v_S\|_1\}$, and $\kappa > 0$, $X$ satisfies $\mathrm{RE}(S, \kappa)$ if $n^{-\frac{1}{2}} \|Xv\|_2 \ge \kappa \|v_S\|_2$ for all $v \in C(S)$.

**Theorem 6.2.4.** *Basic Lasso Rate Form*
*Let $y = X\beta_0 + \varepsilon$, $S := \mathrm{supp}(\beta_0)$, $|S| \le s$, and let $\widehat{\beta}_\lambda$ be any lasso solution.*
*If $X$ satisfies $\mathrm{RE}(S, \kappa)$ and $\lambda > 0$ with $\lambda \ge 2 \left\| \frac{X^\top \varepsilon}{n} \right\|_\infty$, then $n^{-1} \left\| X(\widehat{\beta}_\lambda - \beta_0) \right\|_2^2 \le \frac{9s\lambda^2}{\kappa^2}$ and $\left\| \widehat{\beta}_\lambda - \beta_0 \right\|_1 \le \frac{12s\lambda}{\kappa^2}$.*

*Proof.* Set $\delta := \widehat{\beta}_\lambda - \beta_0$. The lasso optimality inequality gives $(2n)^{-1} \|X\delta\|_2^2 + \lambda \|\beta_0 + \delta\|_1 \le n^{-1} \varepsilon^\top X\delta + \lambda \|\beta_0\|_1$.
Since $\lambda \ge 2 \left\| \frac{X^\top \varepsilon}{n} \right\|_\infty$, $n^{-1} \varepsilon^\top X\delta \le \frac{\lambda}{2} \|\delta\|_1$, and since $\beta_{0,S^c} = 0$, $\|\beta_0\|_1 - \|\beta_0 + \delta\|_1 \le \|\delta_S\|_1 - \|\delta_{S^c}\|_1$.
Thus $(2n)^{-1} \|X\delta\|_2^2 + \frac{\lambda}{2} \|\delta_{S^c}\|_1 \le \frac{3\lambda}{2} \|\delta_S\|_1$, so $\delta \in C(S)$. By $\mathrm{RE}(S, \kappa)$, $\|\delta_S\|_1 \le \sqrt{s} \|\delta_S\|_2 \le \frac{\sqrt{s} \|X\delta\|_2}{\kappa \sqrt{n}}$.
Hence $n^{-1} \|X\delta\|_2^2 \le \frac{3\lambda \sqrt{s} \|X\delta\|_2}{\kappa \sqrt{n}}$, giving $n^{-1} \|X\delta\|_2^2 \le \frac{9s\lambda^2}{\kappa^2}$; also $\|\delta\|_1 \le 4 \|\delta_S\|_1 \le \frac{12s\lambda}{\kappa^2}$.
*Structural Commentary*: Basic inequality plus RE converts score control into prediction and $\ell^1$ error bounds.
*Key Insight*: The $\ell^1$ penalty forces the error into the RE cone. □

**Method 6.2.5.** *Featurewise Two-Sample Screening*
Let $x_{ij}$ be feature $j \in [M]$ for observation $i$, with groups $C_1, C_2$ and $n_\ell := |C_\ell|$.

(i) *Group means*: For $\ell \in \{1, 2\}$, set $\bar{x}_{\ell j} := n_\ell^{-1} \sum_{i \in C_\ell} x_{ij}$.
(ii) *Pooled scale*: Set $\widehat{\sigma}_j^2 := (n_1 + n_2 - 2)^{-1} \{ \sum_{i \in C_1} (x_{ij} - \bar{x}_{1j})^2 + \sum_{i \in C_2} (x_{ij} - \bar{x}_{2j})^2 \}$ and $\mathrm{se}_j := \widehat{\sigma}_j \sqrt{n_1^{-1} + n_2^{-1}}$.
(iii) *Statistic*: Define $t_j := (\bar{x}_{2j} - \bar{x}_{1j}) / \mathrm{se}_j$; large $|t_j|$ flags a featurewise mean shift.
(iv) *Family*: Screening acts on $\{t_j : 1 \le j \le M\}$, so error control targets the family, not a single statistic.

**Remark 6.2.6.** *Statistical and Computational Structure*
High-dimensional inference couples concentration bounds, convex optimization, and selection error control; computation and statistics are part of the same guarantee.

## 6.3 Rank-Based Comparative Tests

Rank methods replace numerical values by their relative order. They are invariant under strictly increasing transformations and reduce the influence of extreme observations, which makes them natural companions to the two-sample and paired $t$ procedures.

**Definition 6.3.1.** *Mann–Whitney and Rank-Sum Statistics*
Let $X_1, \dots, X_n \sim F$ and $Y_1, \dots, Y_m \sim G$ be independent continuous samples.
Rank the pooled observations increasingly.

(i) *Pooled size*: $N := m + n$.
(ii) *$Y$-ranks*: $R_j^Y := \operatorname{rank}(Y_j)$ in the pooled sample.
(iii) *Rank sum*: $T_Y := \sum_{j=1}^{m} R_j^Y$.
(iv) *Mann–Whitney statistic*: $U_Y := T_Y - \frac{m(m+1)}{2}$.

Under continuity, $U_Y = \sum_{i=1}^{n} \sum_{j=1}^{m} \mathbf{1}_{\{X_i < Y_j\}}$ a.s., so $U_Y$ counts cross-sample pairs with $Y_j > X_i$.

**Theorem 6.3.2.** *Null Mean and Variance of the Rank Sum*
*Let $X_1, \dots, X_n \sim F$ and $Y_1, \dots, Y_m \sim G$ be independent continuous samples.*
*Let rank-sum $T_Y$ and Mann–Whitney statistic $U_Y$. Assume $F = G$ and set $N := m + n$.*

*(i) Rank law: $T_Y = \sum_{r \in A} r$, where $A$ is uniform over all $m$-subsets of $\{1, \dots, N\}$.*
*(ii) Rank-sum moments: $\mathbb{E}T_Y = \frac{m(N+1)}{2}$ and $\operatorname{Var}(T_Y) = \frac{mn(N+1)}{12}$.*
*(iii) Mann–Whitney moments: $\mathbb{E}U_Y = \frac{mn}{2}$ and $\operatorname{Var}(U_Y) = \frac{mn(N+1)}{12}$.*

*Proof.*

(i) Continuity gives no ties and $F = G$ makes labels exchangeable, so the $Y$-rank set $A \subseteq \{1, \dots, N\}$ is uniform over all $m$-subsets and $T_Y = \sum_{r \in A} r$.
(ii) For ranks $1, \dots, N$, $\overline{r} = \frac{N+1}{2}$ and $S_r^2 = \frac{N^2-1}{12}$; sampling $m$ ranks without replacement gives $\mathbb{E}T_Y = m\overline{r} = \frac{m(N+1)}{2}$ and $\operatorname{Var}(T_Y) = \frac{mS_r^2(N-m)}{N-1} = \frac{mn(N+1)}{12}$.
(iii) Since $U_Y = T_Y - \frac{m(m+1)}{2}$, shifting by a constant gives $\mathbb{E}U_Y = \frac{m(N+1)}{2} - \frac{m(m+1)}{2} = \frac{mn}{2}$ and $\operatorname{Var}(U_Y) = \operatorname{Var}(T_Y) = \frac{mn(N+1)}{12}$.

*Structural Commentary*: Under $F = G$, ranks reduce to simple random sampling without replacement.
*Key Insight*: $U_Y$ is just a centered rank sum, so only the mean shifts. □

**Method 6.3.3.** *Mann–Whitney Test*
Given independent samples $X_1, \dots, X_n \sim F$ and $Y_1, \dots, Y_m \sim G$, test $H_0 : F = G$:

(i) *Compute*: Pool and rank $m + n$ observations, use midranks for ties. Compute $T_Y$, $U_Y = T_Y - \frac{m(m+1)}{2}$.
(ii) *Calibrate*: For small $m, n$, enumerate $m$-subsets of $\{1, \dots, m+n\}$ to get the exact null law. For large $m, n$, use $Z := \frac{U_Y - \frac{mn}{2}}{\sqrt{\frac{mn(m+n+1)}{12}}} \approx N(0,1)$, with tie correction when needed.
(iii) *Decide*: Compute the relevant one- or two-sided tail probability under the chosen null law. Reject $H_0$ at level $\alpha$ when $p \le \alpha$.

**Example 6.3.4.** *Latent Heat Data by Ranks*
For the latent-heat data, take $m = 8$, $n = 13$, $N = 21$, and midranks for ties.
The smaller-sample rank sum is $R = 51$, its complement is $R' = m(N+1) - R = 125$, and $R^* = \min(R, R') = 51$.
Under equality of the two method distributions, $\mu_T = \frac{m(N+1)}{2} = 88$ and $\sigma_T = \sqrt{\frac{mn(N+1)}{12}} \approx 13.8$,
hence $z = \frac{R - \mu_T}{\sigma_T} \approx -2.68$ and the two-sided normal $p$-value is $\approx 0.007$.
The exact rank-sum test therefore rejects equality at level 0.01.

**Definition 6.3.5.** *Mann–Whitney Effect Probability*
Let $X \sim F$ and $Y \sim G$ be independent, with samples $X_1, \dots, X_n$ and $Y_1, \dots, Y_m$.

(i) *Effect probability*: $\pi := \mathbb{P}(X < Y)$.
(ii) *Estimator*: $\widehat{\pi} := (mn)^{-1} \sum_{i=1}^{n} \sum_{j=1}^{m} \mathbf{1}_{\{X_i < Y_j\}} = \frac{U_Y}{mn}$.

Thus $\pi$ is the probability that a random $Y$ exceeds a random $X$.

**Method 6.3.6.** *Shift Intervals from Pairwise Differences*
Assume the shift model $G(x) = F(x - \Delta)$.

(i) *Compute*: Set $D_{ij} := Y_j - X_i$ and order $D_{(1)} \le \cdots \le D_{(mn)}$.
(ii) *Calibrate*: Choose $k$ so the two-sided rank test accepts a candidate shift when $k \le U_Y(\Delta) \le mn - k$.
(iii) *Invert*: Report $[D_{(k)}, D_{(mn-k+1)}]$ as the inverted shift interval.

The Hodges–Lehmann point estimate is $\widehat{\Delta}_{HL} := \text{median}\{D_{ij} : 1 \le i \le n, 1 \le j \le m\}$.

**Definition 6.3.7.** *Signed-Rank Statistic*
Let $D_1, \ldots, D_n$ be paired differences.

(i) *Nonzero index set*: $I := \{i : D_i \neq 0\}$.
(ii) *Absolute ranks*: $R_i := \text{rank}(|D_i|)$ among $\{|D_\ell| : \ell \in I\}$, using midranks for ties.
(iii) *Signed-rank statistic*: $W_+ := \sum_{i \in I} R_i \mathbf{1}_{\{D_i > 0\}}$.

**Theorem 6.3.8.** *Signed-Rank Mean and Variance*
*Under the signed-rank null, assume no zero differences and no ties in $|D_i|$.*

*(i) Mean:* $\mathbb{E}W_+ = \frac{n(n+1)}{4}$.
*(ii) Variance:* $\text{Var}(W_+) = \frac{n(n+1)(2n+1)}{24}$.

*Proof.* Let $\mathcal{R}$ be the absolute-rank information and let $I_k$ indicate positivity at absolute-rank $k$.
Then $W_+ = \sum_{k=1}^{n} kI_k$ and, under symmetry, $I_k \mid \mathcal{R}$ are independent Bernoulli($\frac{1}{2}$).

(i) *Mean*: $\mathbb{E}(W_+ \mid \mathcal{R}) = \frac{1}{2}\sum_{k=1}^{n} k = \frac{n(n+1)}{4}$; the right side is deterministic, hence $\mathbb{E}W_+ = \frac{n(n+1)}{4}$.

(ii) *Variance*: $\text{Var}(W_+ \mid \mathcal{R}) = \frac{1}{4}\sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{24}$, while $\mathbb{E}(W_+ \mid \mathcal{R})$ is deterministic.

Total variance gives $\text{Var}(W_+) = \frac{n(n+1)(2n+1)}{24}$.

*Structural Commentary*: Condition on absolute ranks; only independent fair signs remain.
*Key Insight*: $W_+$ is a weighted sum of Bernoulli signs with weights $1, \ldots, n$. □

**Method 6.3.9.** *Signed-Rank Test*
For paired differences $D_i$, test $H_0 : D_i \overset{d}{=} -D_i$.

(i) *Compute*: Discard zero differences, rank nonzero $|D_i|$ using midranks for ties, and compute $W_+$.
(ii) *Calibrate*: Conditional on the nonzero absolute ranks, enumerate the $2^r$ sign assignments, where $r := |I|$, to get the null law of $W_+$.
(iii) *Decide*: Compute the relevant tail probability and reject $H_0$ at level $\alpha$ when $p \le \alpha$.

The sign test is the coarser paired alternative using only signs, not ranks.

**Example 6.3.10.** *Paired Rank Analyses*
Two paired-rank calculations are:

(i) *Platelet data*: With one negative signed rank, use $W_-$; exact signed-rank test rejects symmetry at 0.01.
(ii) *Mercury data*: Dropping one zero difference and using midranks gives $n = 24$ and $W_+ = 194.5$. Under $H_0$, $\mathbb{E}_0 W_+ = \frac{24 \cdot 25}{4} = 150$ and $\text{Var}_0(W_+) = \frac{24 \cdot 25 \cdot 49}{24} = 1225$, so $Z = \frac{194.5 - 150}{35} \approx 1.27$ and two-sided $p \approx 0.20$; there is no strong evidence of a systematic shift.

**Remark 6.3.11.** *Rank Tests and Normal Theory*
Rank tests use order information, require no normal model, and are robust to outliers; under normal models their efficiency relative to $t$ tests is often close to 1.

# 7 Time Series Models

## 7.1 Stationarity, Autocovariance, and Linear Processes

Time-series statistics studies data ordered by time, where dependence is part of the signal rather than a nuisance to be removed. The process-level notions of strict and weak stationarity are fixed in Definition 4.5.1.

**Definition 7.1.1.** *Autocorrelation, Partial Autocorrelation, and Spectral Density*
Let $(X_t)_{t \in \mathbb{Z}}$ be weakly stationary with autocovariance $\gamma(h) := \text{Cov}(X_t, X_{t-h})$ and $\gamma(0) > 0$.

For $h \geq 2$ and $t \in \mathbb{Z}$, set $\mathcal{H}_{t,h} := \operatorname{span}\{1, X_{t-1}, \ldots, X_{t-h+1}\}$ let $P_{t,h}$ be $L^2$-orthogonal projection onto $\mathcal{H}_{t,h}$. Assume $\operatorname{Var}(X_t - P_{t,h}X_t) > 0$ and $\operatorname{Var}(X_{t-h} - P_{t,h}X_{t-h}) > 0$ $\forall h \geq 2,\ t \in \mathbb{Z}$.

(i) *Autocorrelation function*: $\rho : \mathbb{Z} \to [-1, 1]$, $\rho(h) := \frac{\gamma(h)}{\gamma(0)}$.

(ii) *Partial autocorrelation function*: $\alpha : \mathbb{N}_0 \to [-1, 1]$, $\alpha(0) := 1$, $\alpha(1) := \frac{\gamma(1)}{\gamma(0)}$, and $\alpha(h) := \operatorname{Corr}(X_t - P_{t,h}X_t, X_{t-h} - P_{t,h}X_{t-h})$ $\forall h \geq 2,\ t \in \mathbb{Z}$.

(iii) *Spectral density*: If $\sum_{h\in\mathbb{Z}} |\gamma(h)| < \infty$, $f_X : [-\pi, \pi] \to \mathbb{C}$, $f_X(\lambda) := (2\pi)^{-1} \sum_{h\in\mathbb{Z}} \gamma(h)e^{-ih\lambda}$ $\forall \lambda \in [-\pi, \pi]$.

**Definition 7.1.2.** *White Noise and Linear Process*
Let $\sigma^2 \geq 0$, $\mu \in \mathbb{R}$, and $(\psi_j)_{j\geq 0} \subset \mathbb{R}$ satisfy $\sum_{j=0}^{\infty} |\psi_j| < \infty$; all processes below are real-valued, indexed by $\mathbb{Z}$.

(i) *White noise*: $(\varepsilon_t)_{t\in\mathbb{Z}} \subset L^2$, $\mathbb{E}\varepsilon_t = 0$ and $\operatorname{Var}(\varepsilon_t) = \sigma^2$ $\forall t \in \mathbb{Z}$, and $\operatorname{Cov}(\varepsilon_t, \varepsilon_s) = 0$ $\forall s, t \in \mathbb{Z}$ with $s \neq t$.

(ii) *Linear process*: $(X_t)_{t\in\mathbb{Z}}$ satisfies $X_t = \mu + \sum_{j=0}^{\infty} \psi_j \varepsilon_{t-j}$ $\forall t \in \mathbb{Z}$ for some $(\varepsilon_t)_{t\in\mathbb{Z}}$ satisfying (i).

*Structural Commentary*: $\sum_j |\psi_j| < \infty$ gives an $L^2$-stable one-sided innovation filter.

**Example 7.1.3.** *Autoregression of Order One*
Let $(\varepsilon_t)_{t\in\mathbb{Z}}$ be white noise, variance $\sigma^2$, $|\phi| < 1$. For $X_t = \phi X_{t-1} + \varepsilon_t$, stationary solution is $X_t = \sum_{j=0}^{\infty} \phi^j \varepsilon_{t-j}$.

Thus $\gamma(h) = \operatorname{Cov}(X_t, X_{t-h}) = \frac{\sigma^2\phi^{|h|}}{1-\phi^2}$, $\rho(h) = \frac{\gamma(h)}{\gamma(0)} = \phi^{|h|}$. Autocorrelation decays geometrically at rate $|\phi|$.

**Remark 7.1.4.** *Process-Level Prerequisites*
Time-series inference uses filtrations $(\mathcal{F}_t)$, adapted $X_t$, predictors $\mathbb{E}(X_{t+h} \mid \mathcal{F}_t)$, martingale differences $\varepsilon_t$, and stationarity or ergodicity assumptions.

## 7.2 ARIMA, State-Space, and Volatility Models

Classical time-series models combine deterministic filters, stochastic innovations, and likelihood-based estimation.

**Definition 7.2.1.** *ARMA Model*
Let $p, q \in \mathbb{N}_0$, $\phi_i \in \mathbb{R}$ $(1 \leq i \leq p)$, $\theta_j \in \mathbb{R}$ $(1 \leq j \leq q)$, and let $B$ denote the backshift, $(BX)_t = X_{t-1}$.
Set $\phi(z) := 1 - \sum_{i=1}^{p} \phi_i z^i$ and $\theta(z) := 1 + \sum_{j=1}^{q} \theta_j z^j$; empty sums are zero.
$(X_t)_{t\in\mathbb{Z}}$ is ARMA$(p, q)$ with innovations $(\varepsilon_t)_{t\in\mathbb{Z}}$ iff $(\varepsilon_t)$ is white noise and $\phi(B)X_t = \theta(B)\varepsilon_t$ $\forall t \in \mathbb{Z}$.
*Structural Commentary*: Causal iff $\phi(z) \neq 0$ for $|z| \leq 1$; invertible iff $\theta(z) \neq 0$ for $|z| \leq 1$.
*Key Insight*: AR roots control recursion; MA roots control innovation recovery.

**Definition 7.2.2.** *ARIMA Model*
Let $p, q, d \in \mathbb{N}_0$, $\phi_i \in \mathbb{R}$ $(1 \leq i \leq p)$, $\theta_j \in \mathbb{R}$ $(1 \leq j \leq q)$, and let $B$ denote the backshift.
Set $\Delta^d := (1 - B)^d$, $\phi(z) := 1 - \sum_{i=1}^{p} \phi_i z^i$, and $\theta(z) := 1 + \sum_{j=1}^{q} \theta_j z^j$.
$(X_t)_{t\in\mathbb{Z}}$ is ARIMA$(p, d, q)$ with innovations $(\varepsilon_t)_{t\in\mathbb{Z}}$ iff $(\varepsilon_t)$ is white noise and $\phi(B)\Delta^d X_t = \theta(B)\varepsilon_t$ $\forall t \in \mathbb{Z}$.
*Structural Commentary*: $Y_t := \Delta^d X_t$ satisfies ARMA$(p, q)$ with the same innovations.
*Key Insight*: $d > 0$ shifts stationarity from $X_t$ to $\Delta^d X_t$.

**Definition 7.2.3.** *Filtering Problem*
Let $(Z_t, Y_t)_{t\geq 1}$ be a hidden-state model with state space $(E, \mathcal{E})$, and set $\mathcal{F}_t^Y := \sigma(Y_1, \ldots, Y_t)$.
For $A \in \mathcal{E}$, $\pi_t(A) := \mathbb{P}(Z_t \in A \mid \mathcal{F}_t^Y)$; the task is recursive propagation or approximation of $(\pi_t)_{t\geq 1}$.
*Structural Commentary*: $(\pi_t)$ is an $\mathcal{F}_t^Y$-adapted probability-kernel-valued process.
*Key Insight*: Conditioning converts hidden dynamics into an observable belief-state recursion.

**Definition 7.2.4.** *Linear Gaussian State-Space Model*
Let $m, r, n \in \mathbb{N}$, $F \in \mathbb{R}^{m\times m}$, $G \in \mathbb{R}^{m\times r}$, $H \in \mathbb{R}^{n\times m}$, and $Q \in \mathbb{R}^{r\times r}$, $R \in \mathbb{R}^{n\times n}$ with $Q, R \succeq 0$.
Let $(\eta_t)$ and $(\varepsilon_t)$ be mutually independent Gaussian noises with $\eta_t \sim N(0, Q)$ and $\varepsilon_t \sim N(0, R)$.
Let $\alpha_0$ be independent Gaussian. $(\alpha_t, Y_t)_{t\in\mathbb{N}_0}$ is *linear Gaussian state-space* iff
$\alpha_t \in \mathbb{R}^m$, $Y_t \in \mathbb{R}^n$, $\alpha_{t+1} = F\alpha_t + G\eta_t$, $Y_t = H\alpha_t + \varepsilon_t$ $\forall t \in \mathbb{N}_0$.
*Key Insight*: With $\mathcal{F}_t^Y := \sigma(Y_0, \ldots, Y_t)$, Kalman filtering recurses $(a_t, P_t) := (\mathbb{E}(\alpha_t \mid \mathcal{F}_t^Y), \operatorname{Var}(\alpha_t \mid \mathcal{F}_t^Y))$.

**Definition 7.2.5.** *ARCH and GARCH Volatility*
Let $p, q \in \mathbb{N}_0$, $\omega > 0$, $\alpha_1, \dots, \alpha_q, \beta_1, \dots, \beta_p \geq 0$, and let $(z_t)_{t\in\mathbb{Z}}$ be standardized white noise.
All processes below are real-valued and indexed by $\mathbb{Z}$; set $A := \sum_{i=1}^{q} \alpha_i + \sum_{j=1}^{p} \beta_j$.

(i) *ARCH(q)*: $(X_t, \sigma_t)_{t\in\mathbb{Z}}$, $\sigma_t \geq 0$, satisfies $X_t = \sigma_t z_t$ and $\sigma_t^2 = \omega + \sum_{i=1}^{q} \alpha_i X_{t-i}^2 \ \forall t \in \mathbb{Z}$.

(ii) *GARCH(p, q)*: $(X_t, \sigma_t)_{t\in\mathbb{Z}}$, $\sigma_t \geq 0$, satisfies $X_t = \sigma_t z_t$ and $\sigma_t^2 = \omega + \sum_{i=1}^{q} \alpha_i X_{t-i}^2 + \sum_{j=1}^{p} \beta_j \sigma_{t-j}^2 \ \forall t \in \mathbb{Z}$.

*Structural Commentary*: $\omega > 0$, $\alpha_i, \beta_j \geq 0$ force $\sigma_t^2 \geq 0$; by stationary GARCH, $A < 1$ has $\mathbb{E}X_t^2 = \frac{\omega}{1-A}$.

**Remark 7.2.6.** *Likelihood by Filtering*

If one-step densities exist, likelihood filtering has $S_t = \mathcal{T}_\vartheta(S_{t-1}, W_t)$ and $\ell_T(\vartheta) = \sum_{t=1}^{T} \log p_\vartheta(W_t \mid S_{t-1})$.
Inference depends on stability of $S_t$.

**Method 7.2.7.** *Estimating a Finite Transition Matrix*

(i) Observe finite homogeneous chain $X_0, \dots, X_n$, let $n_{ij} := \sum_{t=0}^{n-1} \mathbf{1}\{X_t = i, X_{t+1} = j\}$, $n_i := \sum_j n_{ij}$.

(ii) The row likelihood is proportional to $\prod_j p_{ij}^{n_{ij}}$ subject to $\sum_j p_{ij} = 1$.

(iii) If $n_i > 0$, the rowwise MLE is $\widehat{p}_{ij} = \frac{n_{ij}}{n_i}$.

(iv) If $n_i = 0$, row $i$ is not identifiable without a convention or prior.

# 8 Sequential Inference and Resampling

## 8.1 Resampling, Bootstrap, and Permutation Methods

Resampling methods approximate sampling distributions by recomputing statistics on data-generated pseudo-samples.

**Definition 8.1.1.** *Nonparametric Bootstrap*
For $X_{1:n} \sim P^{\otimes n}$, set $\widehat{P}_n := n^{-1} \sum_{i=1}^{n} \delta_{X_i}$. Conditional on $X_{1:n}$, let $X_{1:n}^* \sim \widehat{P}_n^{\otimes n}$ and $\widehat{P}_n^* := n^{-1} \sum_{i=1}^{n} \delta_{X_i^*}$.
For any law functional $T$, $\mathcal{L}(T(\widehat{P}_n^*) - T(\widehat{P}_n) \mid X_{1:n})$ approximates $\mathcal{L}(T(\widehat{P}_n) - T(P))$.
*Key Insight*: $P$ is replaced by $\widehat{P}_n$ before pseudo-sampling.

**Method 8.1.2.** *Bootstrap Standard Error*
Let $\widehat{\psi} = s(X_{1:n})$ be observed statistic, $\mathcal{R}_*(X_{1:n})$ be the chosen bootstrap law on pseudo-samples, $B \geq 2$.

(i) *Resample*: $X_{1:n}^{*b} \sim \mathcal{R}_*(X_{1:n})$ for $b = 1, \dots, B$.

(ii) *Recompute*: $\widehat{\psi}^{*b} := s(X_{1:n}^{*b})$ for $b = 1, \dots, B$.

(iii) *Estimate*: $\widehat{\mathrm{se}}_*(\widehat{\psi}) := \{(B-1)^{-1} \sum_{b=1}^{B} (\widehat{\psi}^{*b} - \overline{\psi}^*)^2\}^{\frac{1}{2}}$, where $\overline{\psi}^* := B^{-1} \sum_{b=1}^{B} \widehat{\psi}^{*b}$.

**Method 8.1.3.** *Bootstrap Variability for Location Summaries*
Let $T = T(x_1, \dots, x_n)$ be a location summary and let $B \geq 1$.

(i) *Resample*: For $1 \leq b \leq B$, draw $x_1^{*(b)}, \dots, x_n^{*(b)}$ iid from $\widehat{F}_n := n^{-1} \sum_{i=1}^{n} \delta_{x_i}$.

(ii) *Recompute*: Set $T_b^* := T(x_1^{*(b)}, \dots, x_n^{*(b)})$ and $\overline{T}^* := B^{-1} \sum_{b=1}^{B} T_b^*$.

(iii) *Spread*: Estimate variability by $\widehat{\mathrm{se}}_*(T) := \left( (B-1)^{-1} \sum_{b=1}^{B} (T_b^* - \overline{T}^*)^2 \right)^{1/2}$.

*Key Insight*: This is useful for medians and trimmed means, where analytic standard errors can be tail-sensitive.

**Definition 8.1.4.** *Parametric Bootstrap*
Let $\{P_\theta : \theta \in \Theta\}$ be parametric model for $X_{1:n} := (X_1, \dots, X_n)$, let $\widehat{\theta} := \widehat{\theta}(X_{1:n})$, and let $S$ be a statistic.
$X_1^*, \dots, X_n^* \mid X_{1:n} \sim P_{\widehat{\theta}}^{\otimes n}$, and $\mathcal{L}(S(X_{1:n}^*) \mid X_{1:n})$ is the fitted-model resampling law for $S(X_{1:n})$.
*Key Insight*: The model class $\{P_\theta\}$ is retained; only $\theta$ is replaced by $\widehat{\theta}$.

**Method 8.1.5.** *Parametric Bootstrap Standard Errors and Intervals*
Let $\{P_\theta : \theta \in \Theta\}$ be parametric model, $\widehat{\psi} := s(X_{1:n})$ estimate $\psi(\theta)$, fix $B \geq 2$, fix confidence level $1 - \alpha$.

(i) *Fit*: $\widehat{\theta} := \widehat{\theta}(X_{1:n})$.

(ii) *Simulate*: $X^{*b}_{1:n} \sim P_{\widehat{\theta}}^{\otimes n}$ and $\widehat{\psi}^{*b} := s(X^{*b}_{1:n})$ for $b = 1, \dots, B$.

(iii) *Standard error*: $\widehat{\mathrm{se}}_*(\widehat{\psi}) := \{(B-1)^{-1} \sum_{b=1}^{B} (\widehat{\psi}^{*b} - \overline{\psi}^*)^2\}^{\frac{1}{2}}$, where $\overline{\psi}^* := B^{-1} \sum_{b=1}^{B} \widehat{\psi}^{*b}$.

(iv) *Pivotal interval*: for empirical quantiles $q^*_{\frac{\alpha}{2}}$ and $q^*_{1-\frac{\alpha}{2}}$ of $\{\widehat{\psi}^{*b} - \widehat{\psi}\}_{b=1}^{B}$, use $[\widehat{\psi} - q^*_{1-\frac{\alpha}{2}}, \widehat{\psi} - q^*_{\frac{\alpha}{2}}]$.

The percentile interval uses empirical quantiles of $\{\widehat{\psi}^{*b}\}_{b=1}^{B}$ directly.
The pivotal and percentile intervals coincide when the bootstrap error law is symmetric about 0.

**Remark 8.1.6.** *Plug-In Monte Carlo*
The parametric bootstrap estimates model-dependent quantities by replacing the unknown parameter with its estimate before simulation. If $h(X, \theta)$ is a loss, error indicator, or other accuracy summary, then Monte Carlo from $P_{\widehat{\theta}}$ approximates $\mathbb{E}_\theta h(X, \theta)$ by the empirical average of $h(X^{*b}, \widehat{\theta})$.

**Definition 8.1.7.** *Permutation Test*
Let a finite group $G$ act on the sample space $\mathcal{X}$, let $P_0$ satisfy $gX \overset{\text{dist}}{=} X$ under $P_0$ $\forall g \in G$, and let $T : \mathcal{X} \to \mathbb{R}$ be a test statistic. With $U_G$ uniform on $G$ and independent of $X$, $\mathcal{L}(T(U_G X) \mid X)$, the empirical law on $\{T(gX) : g \in G\}$, is the reference law for tail probabilities.
*Key Insight*: Calibration follows from the invariance $gX \overset{\text{dist}}{=} X$ under $P_0$.

**Method 8.1.8.** *Permutation Plug-In FDR Estimate*
Let $t_j$ be observed statistics for $j \in [M]$, and let $t_j^{(k)}$ be the statistic after label permutation $k \in [K]$.

(i) *Observed discoveries*: For cutpoint $C > 0$, set $R_{\text{obs}}(C) := \sum_{j=1}^{M} \mathbf{1}_{\{|t_j| > C\}}$.

(ii) *Null discoveries*: Estimate the expected false discoveries by $\widehat{\mathbb{E}}(V; C) := K^{-1} \sum_{k=1}^{K} \sum_{j=1}^{M} \mathbf{1}_{\{|t_j^{(k)}| > C\}}$.

(iii) *Plug-in FDR*: If $R_{\text{obs}}(C) > 0$, set $\widehat{\mathrm{FDR}}(C) := \widehat{\mathbb{E}}(V; C) / R_{\text{obs}}(C)$.

(iv) *Permutation p-values*: Use $\{t_j^{(k)}\}$ to compare $|t_j|$ with featurewise or pooled permutation null statistics.

**Remark 8.1.9.** *Asymmetric Cutpoints and SAM*
Let $t_{(1)} \le \cdots \le t_{(M)}$ and $\bar{t}^0_{(j)} := K^{-1} \sum_{k=1}^{K} t^{(k)}_{(j)}$, where $t^{(k)}_{(j)}$ is the $j$th ordered permutation statistic.
For $d_j := t_{(j)} - \bar{t}^0_{(j)}$, $a > 0$, set $c_+(a) := \min\{t_{(j)} : d_j > a\}$, $c_-(a) := \max\{t_{(j)} : d_j < -a\}$ when nonempty.
Reject $t_j \in A(c_-, c_+) := (-\infty, c_-] \cup [c_+, \infty)$, estimate FDR by Method 8.1.8, $A(c_-, c_+)$ replace $\{|t| > C\}$.

**Remark 8.1.10.** *Validity Depends on the Resampling Law*
Bootstrap validity is asymptotic; permutation validity is finite-sample exact under invariance; both require the resampling law to match the target sampling variation.

## 8.2 Sequential Experiments, Filtering, and Online Inference

Sequential methods update decisions as observations arrive. They must account for optional stopping, adaptive sampling, and computational constraints.

**Definition 8.2.1.** *Sequential Test and Stopping Time*
Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $I \subseteq \mathbb{N}_0$ be a time index, let $(\mathcal{F}_t)_{t \in I}$ be a filtration, and let $(D, \mathcal{D})$ be a decision space. For any $\tau : \Omega \to I \cup \{\infty\}$, set $\mathcal{F}_\tau := \{A \in \mathcal{F} : A \cap \{\tau \le t\} \in \mathcal{F}_t \ \forall t \in I\}$.

(i) *Stopping time*: $\tau : \Omega \to I \cup \{\infty\}$ satisfies $\{\tau \le t\} \in \mathcal{F}_t$ $\forall t \in I$.

(ii) *Sequential test*: $(\tau, d)$ with $\tau$ a stopping time and $d : (\Omega, \mathcal{F}_\tau) \to (D, \mathcal{D})$ measurable.

*Key Insight*: Stop and decide using only pre-stopping information.

**Definition 8.2.2.** *Sequential Probability Ratio Test*
Let $P_0$ and $P_1$ be simple iid models with one-observation densities $p_0$ and $p_1$ for $(X_t)_{t \ge 1}$,
let $\Lambda_0 := 1$, $\Lambda_t := \prod_{i=1}^{t} \frac{p_1(X_i)}{p_0(X_i)}$, and choose $0 < A < B < \infty$.
$\tau := \inf\{t \ge 1 : \Lambda_t \in (0, A] \cup [B, \infty)\}$, with terminal decision $d_\tau = P_0$ on $\{\Lambda_\tau \le A\}$ and $d_\tau = P_1$ on $\{\Lambda_\tau \ge B\}$.
*Key Insight*: First exit of $\log \Lambda_t$ from $(\log A, \log B)$ converts cumulative likelihood evidence into a stopping rule.

**Remark 8.2.3.** *Sequential Cross-References*
Sequential inference uses filtrations and martingales from stochastic processes, stability of recursions from dynamical systems, and simulation methods from Monte Carlo computation.

## 8.3 Two-Sample Resampling Methods

Two-sample resampling procedures must preserve the source of randomness relevant to the question. Bootstrap procedures usually resample within groups to estimate sampling variability under the fitted empirical distributions; permutation procedures pool labels when the null asserts exchangeability.

**Method 8.3.1.** *Two-Sample Nonparametric Bootstrap*

Let $x_{1:n}$ and $y_{1:m}$ be independent samples from $F$ and $G$, let $\widehat{F}_n := n^{-1}\sum_{i=1}^{n}\delta_{x_i}$ and $\widehat{G}_m := m^{-1}\sum_{j=1}^{m}\delta_{y_j}$.

Let $T(F,G)$ be a two-sample functional, and fix $B \geq 2$.

(i) *Resample*: $x^{*b}_{1:n} \sim \widehat{F}_n^{\otimes n}$ and $y^{*b}_{1:m} \sim \widehat{G}_m^{\otimes m}$ independently for $b = 1,\dots,B$.

(ii) *Recompute*: with $\widehat{F}^{*b}_n := n^{-1}\sum_{i=1}^{n}\delta_{x^{*b}_i}$ and $\widehat{G}^{*b}_m := m^{-1}\sum_{j=1}^{m}\delta_{y^{*b}_j}$, set $\widehat{T}^{*b} := T(\widehat{F}^{*b}_n, \widehat{G}^{*b}_m)$.

(iii) *Summarize*: $s^*_T := \mathrm{sd}\{\widehat{T}^{*b} : 1 \leq b \leq B\}$; $I^*_{1-\alpha} := [q^*_{\frac{\alpha}{2}}, q^*_{1-\frac{\alpha}{2}}]$, where $q^*_\gamma$ is the empirical $\gamma$-quantile.

This is the two-sample version of Method 8.1.2.

**Example 8.3.2.** *Bootstrap for $\pi = \mathbb{P}(X < Y)$*

Let $x_{1:n} \sim F^{\otimes n}$ and $y_{1:m} \sim G^{\otimes m}$ independently.

Set $\widehat{F}_n := n^{-1}\sum_{i=1}^{n}\delta_{x_i}$, $\widehat{G}_m := m^{-1}\sum_{j=1}^{m}\delta_{y_j}$, and $\widehat{\pi} := (mn)^{-1}\sum_{i=1}^{n}\sum_{j=1}^{m}\mathbf{1}_{\{x_i<y_j\}}$.

For $b = 1,\dots,B$, draw $x^{*b}_{1:n} \sim \widehat{F}_n^{\otimes n}$ and $y^{*b}_{1:m} \sim \widehat{G}_m^{\otimes m}$ independently.

Set $\widehat{\pi}^{*b} := (mn)^{-1}\sum_{i=1}^{n}\sum_{j=1}^{m}\mathbf{1}_{\{x^{*b}_i<y^{*b}_j\}}$, and use $s^*_\pi := \mathrm{sd}\{\widehat{\pi}^{*b} : 1 \leq b \leq B\}$.

*Key Insight*: Separate resampling targets $F \otimes G$ variation; rank-sum calibration imposes $F = G$.

**Method 8.3.3.** *Bootstrap for a Hodges–Lehmann Shift*

For independent samples $x_{1:n}$ and $y_{1:m}$ and $B \geq 2$, set $\widehat{\Delta}_{\mathrm{HL}} := \mathrm{median}\{y_j - x_i : 1 \leq i \leq n,\ 1 \leq j \leq m\}$, resample the two groups separately for $b = 1,\dots,B$, and use the empirical spread or quantiles of the recomputed medians $\widehat{\Delta}^{*b}_{\mathrm{HL}}$ for a standard error or interval.

**Method 8.3.4.** *Permutation Test for a Difference of Means*

Let $x_{1:n}$ be control responses, let $y_{1:m}$ be treatment responses, set $\overline{x} := n^{-1}\sum_{i=1}^{n}x_i$ and $\overline{y} := m^{-1}\sum_{j=1}^{m}y_j$.

Let $z_{1:N}$ with $N = n + m$ be the pooled responses, and set $T_{\mathrm{obs}} := \overline{y} - \overline{x}$.

Let $H_0$ denote no treatment effect, set $\mathcal{S}_{m,N} := \{S \subseteq \{1,\dots,N\} : |S| = m\}$, and use $S \mid H_0 \sim \mathrm{Unif}(\mathcal{S}_{m,N})$.

(i) *Relabel*: $S \in \mathcal{S}_{m,N}$, by enumeration or Monte Carlo draws from $\mathrm{Unif}(\mathcal{S}_{m,N})$.

(ii) *Recompute*: $T^*_S := m^{-1}\sum_{i\in S}z_i - n^{-1}\sum_{i\notin S}z_i$.

(iii) *Calibrate*: $p_{\mathrm{exact}} := \binom{N}{m}^{-1}\#\{S \in \mathcal{S}_{m,N} : |T^*_S| \geq |T_{\mathrm{obs}}|\}$, Monte Carlo approximation if $\binom{N}{m}$ large.

**Remark 8.3.5.** *Permutation and Rank Tests*

The permutation test for means and the Mann–Whitney test both use the randomization principle that, under the null, labels are exchangeable. They differ in the statistic: the permutation mean test uses the numerical values and is sensitive to outliers, while the Mann–Whitney test first replaces values by ranks.

# Part XX
# Statistical Data Analysis and Applied Models

Applied statistical analysis organizes data summaries, sampling designs, contingency tables, designed experiments, and regression models into reusable workflows. The inferential tools of Part 15 supply the testing, estimation, and uncertainty language used here.

## 1 Survey Sampling

### 1.1 Finite Population Parameters and Simple Random Sampling

Survey sampling treats a finite population as fixed and the sample-selection mechanism as random. The inferential target is often a population mean, total, or proportion, while the random variables come from the sampling design.

**Definition 1.1.1.** *Finite Population Parameters*
Let $U := \{1, \dots, N\}$ and fix $x_i \in \mathbb{R}$, $i \in U$.

(i) *Population mean*: $\mu := \frac{1}{N} \sum_{i=1}^{N} x_i$.

(ii) *Population total*: $\tau := \sum_{i=1}^{N} x_i = N\mu$.

(iii) *Population variance*: $\sigma^2 := \frac{1}{N} \sum_{i=1}^{N} (x_i - \mu)^2 = \frac{1}{N} \sum_{i=1}^{N} x_i^2 - \mu^2$.

(iv) *Population proportion*: $\{x_i : i \in U\} \subseteq \{0, 1\} \implies p := \mu = \frac{1}{N} \sum_{i=1}^{N} x_i$ and $\sigma^2 = p(1-p)$.

**Definition 1.1.2.** *Simple Random Sampling*
Let $U := \{1, \dots, N\}$, $1 \le n \le N$, and fix $x_i \in \mathbb{R}$, $i \in U$.
Let $S$ be a random $n$-element subset of $U$; for a random ordering $J_1, \dots, J_n$ of $S$, set $X_k := x_{J_k}$.

(i) *Simple random sample*: $\mathbb{P}(S = s) = \frac{1}{\binom{N}{n}}$, $\forall s \subseteq U$ with $|s| = n$.

(ii) *Sample mean*: $\overline{X} := \frac{1}{n} \sum_{i=1}^{n} X_i$.

(iii) *Total estimator*: $T := N\overline{X}$.

(iv) *Sample proportion*: $\{x_i : i \in U\} \subseteq \{0, 1\} \implies \widehat{p} := \overline{X}$.

**Method 1.1.3.** *Sequential Simple Random Sampling*
Fix target size $n \ge 1$, assume terminal stream length is $N \ge n$, $N$ unknown while streaming.
Population units arrive sequentially as $1, \dots, N$. Let $S_t \subseteq \{1, \dots, t\}$ denote current reservoir after $t$ arrivals.

(i) *Initialize*: for $1 \le t \le n$, set $S_t = \{1, \dots, t\}$.
(ii) *Admit*: for $t > n$, draw $B_t \sim \text{Bernoulli}(\frac{n}{t})$.
(iii) *Update*: if $B_t = 0$, set $S_t = S_{t-1}$; if $B_t = 1$, draw $J_t \mid S_{t-1} \sim \text{Unif}(S_{t-1})$ and set $S_t = (S_{t-1} \setminus \{J_t\}) \cup \{t\}$.
(iv) *Output*: after the stream ends, return $S_N$.

Then for every $t \ge n$, $S_t \sim \text{Unif}\{s \subseteq \{1, \dots, t\} : |s| = n\}$.
Hence $S_N$ is a simple random sample without replacement.

**Example 1.1.4.** *Capture–Recapture Likelihood*
Suppose $t$ population units are tagged and a later sample of size $m$ contains $r > 0$ tagged units.
For population size $n$, likelihood is $L_n = \frac{\binom{t}{r}\binom{n-t}{m-r}}{\binom{n}{m}}$. Since $\frac{L_n}{L_{n-1}} = \frac{(n-t)(n-m)}{n(n-t-m+r)}$, $L_n$ increases for $n < \frac{mt}{r}$ and decreases for $n > \frac{mt}{r}$. Hence $\widehat{n} = \lfloor \frac{mt}{r} \rfloor$; for $(t, m, r) = (10, 20, 4)$, $\widehat{n} = 50$.

**Lemma 1.1.5.** *Law of One Sample Position*
*Let $U := \{1, \dots, N\}$, $1 \le n \le N$, fix $x_a \in \mathbb{R}$ for $a \in U$.*
*Let $\mu := \frac{1}{N} \sum_{a=1}^{N} x_a$, $\sigma^2 := \frac{1}{N} \sum_{a=1}^{N} (x_a - \mu)^2$, $S \sim \text{Unif}\{s \subseteq U : |s| = n\}$.*
*If $(J_1, \dots, J_n)$ is a random ordering of $S$, set $X_k := x_{J_k}$.*
*Let $\zeta_1, \dots, \zeta_m$ be distinct values in $\{x_a : a \in U\}$, let $n_j := \#\{a \in U : x_a = \zeta_j\}$. For each $k \in \{1, \dots, n\}$:*

*(i) Distribution: the selected value $X_k$ has the finite-population law $\mathbb{P}(X_k = \zeta_j) = \frac{n_j}{N}$ for $j = 1, \dots, m$.*
*(ii) Moments: the finite-population law has mean $\mathbb{E}X_k = \mu$ and variance $\text{Var}(X_k) = \sigma^2$.*

*Proof.*

(i) For every $a \in U$, $\mathbb{P}(J_k = a) = \frac{1}{N}$, so $\mathbb{P}(X_k = \zeta_j) = \sum_{a:x_a=\zeta_j} \mathbb{P}(J_k = a) = \frac{n_j}{N}$.

(ii) From (i), $\mathbb{E}X_k = \frac{1}{N}\sum_{a=1}^{N} x_a = \mu$. By finite-population law, $\mathrm{Var}(X_k) = \frac{1}{N}\sum_{a=1}^{N}(x_a - \mu)^2 = \sigma^2$.

*Structural Commentary*: Each sample position is marginally uniform on $U$.
*Key Insight*: Without replacement changes joint dependence, not the one-position marginal distribution. □

**Theorem 1.1.6.** *Unbiased SRS Estimators*
*Let $U := \{1, \dots, N\}$, $1 \le n \le N$, fix $x_a \in \mathbb{R}$ for $a \in U$.*
*Set $\mu := \frac{1}{N}\sum_{a=1}^{N} x_a$, $\tau := \sum_{a=1}^{N} x_a = N\mu$. Let $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$.*
*Given random ordering $(J_1, \dots, J_n)$ of $S$, set $X_k := x_{J_k}$, $\overline{X} := \frac{1}{n}\sum_{k=1}^{n} X_k$, and $T := N\overline{X}$. Then:*

*(i) Mean estimator: the sample mean is design-unbiased for the finite-population mean, $\mathbb{E}\overline{X} = \mu$.*
*(ii) Total estimator: the expansion estimator is design-unbiased for the finite-population total, $\mathbb{E}T = \tau$.*

*Proof.*

(i) For each $k$, $\mathbb{P}(J_k = a) = \frac{1}{N}\ \forall a \in U$, so $\mathbb{E}X_k = \frac{1}{N}\sum_{a=1}^{N} x_a = \mu$. By linearity $\mathbb{E}\overline{X} = \frac{1}{n}\sum_{k=1}^{n}\mathbb{E}X_k = \mu$.

(ii) Since $T = N\overline{X}$, $\mathbb{E}T = N\mathbb{E}\overline{X} = N\mu = \tau$.

*Structural Commentary*: $T = N\overline{X}$ rescales the mean estimate.
*Key Insight*: Equal position marginals give unbiasedness; independence is unnecessary. □

**Lemma 1.1.7.** *Negative Dependence Without Replacement*
*Let $U := \{1, \dots, N\}$, $2 \le n \le N$, fix $x_a \in \mathbb{R}$ for $a \in U$.*
*Set $\mu := \frac{1}{N}\sum_{a=1}^{N} x_a$, $\tau := \sum_{a=1}^{N} x_a$, $\sigma^2 := \frac{1}{N}\sum_{a=1}^{N}(x_a - \mu)^2$. Let $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$.*
*For random ordering $(J_1, \dots, J_n)$ of $S$, set $X_k := x_{J_k}$. If $i, j \in \{1, \dots, n\}$ and $i \ne j$, then $\mathrm{Cov}(X_i, X_j) = -\frac{\sigma^2}{N-1}$.*

*Proof.* Let $(Y_1, \dots, Y_N)$ be a uniformly random ordering of $(x_1, \dots, x_N)$.
Each $Y_a$ has variance $\sigma^2$, and symmetry gives $\mathrm{Cov}(Y_a, Y_b) = \gamma$ for $a \ne b$.
Since $\sum_{a=1}^{N} Y_a = \tau$ is fixed, $0 = \mathrm{Var}(\sum_{a=1}^{N} Y_a) = N\sigma^2 + N(N-1)\gamma$.
Thus $\gamma = -\frac{\sigma^2}{N-1}$, and $(X_i, X_j)$ has same law as any two distinct positions in full random ordering.
*Structural Commentary*: A fixed total forces negative pairwise covariance.
*Key Insight*: Without replacement, one large draw leaves less for the rest. □

**Theorem 1.1.8.** *Variance and Finite Population Correction*
*Let $U := \{1, \dots, N\}$, $N > 1$, $1 \le n \le N$, fix $x_a \in \mathbb{R}$ for $a \in U$.*
*Set $\mu := \frac{1}{N}\sum_{a=1}^{N} x_a$ and $\sigma^2 := \frac{1}{N}\sum_{a=1}^{N}(x_a - \mu)^2$. Let $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$.*
*For a random ordering $(J_1, \dots, J_n)$ of $S$, set $X_k := x_{J_k}$, $\overline{X} := \frac{1}{n}\sum_{k=1}^{n} X_k$, and $T := N\overline{X}$. Then:*

*(i) Mean variance: the sample mean has design variance $\mathrm{Var}(\overline{X}) = \frac{\sigma^2}{n}\left(\frac{N-n}{N-1}\right) = \frac{\sigma^2}{n}\left(1 - \frac{n-1}{N-1}\right)$.*
*(ii) Total variance: the total estimator has design variance $\mathrm{Var}(T) = N^2\,\mathrm{Var}(\overline{X})$.*

*Proof.*

(i) Each $X_k$ has variance $\sigma^2$. Let $(Y_1, \dots, Y_N)$ be a uniformly random ordering of $(x_1, \dots, x_N)$, set $\tau := \sum_{a=1}^{N} x_a$, and let $\gamma$ be the common covariance of $Y_a, Y_b$ for $a \ne b$.
Since $\sum_{a=1}^{N} Y_a = \tau$, $0 = \mathrm{Var}(\sum_{a=1}^{N} Y_a) = N\sigma^2 + N(N-1)\gamma$, so $\gamma = -\frac{\sigma^2}{N-1}$.
Therefore $\mathrm{Var}(\overline{X}) = \frac{1}{n^2}\left(n\sigma^2 + n(n-1)\left[-\frac{\sigma^2}{N-1}\right]\right) = \frac{\sigma^2}{n}\left(1 - \frac{n-1}{N-1}\right)$.

(ii) The total estimator satisfies $T = N\overline{X}$, so $\mathrm{Var}(T) = \mathrm{Var}(N\overline{X}) = N^2\,\mathrm{Var}(\overline{X})$.

*Structural Commentary*: The factor $\frac{N-n}{N-1}$ is the *finite population correction*.
*Key Insight*: Averaging gives $\frac{1}{n}$ scaling; without replacement subtracts negative covariance. □

**Corollary 1.1.9.** *Estimated Standard Errors*
*Let $U := \{1, \dots, N\}$, $1 < n \le N$, fix $x_a \in \mathbb{R}$ for $a \in U$, and let $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$. For a random ordering $(J_1, \dots, J_n)$ of $S$, set $X_k := x_{J_k}$, $\overline{X} := \frac{1}{n}\sum_{k=1}^{n} X_k$, $T := N\overline{X}$, and $s^2 := \frac{1}{n-1}\sum_{k=1}^{n}(X_k - \overline{X})^2$.*
*Then practical plug-in estimated variances are:*

*(i) Mean: the plug-in estimated variance of the sample mean is $s^2_{\overline{X}} = \frac{s^2}{n}\left(1 - \frac{n}{N}\right)$.*
*(ii) Total: the plug-in estimated variance of the total estimator is $s^2_T = N^2 s^2_{\overline{X}}$.*
*(iii) Proportion: if $x_a \in \{0,1\}$ $\forall a \in U$ and $\widehat{p} := \overline{X}$, then plug-in estimated variance of $\widehat{p}$ is $s^2_{\widehat{p}} = \frac{\widehat{p}(1-\widehat{p})}{n-1}\left(1 - \frac{n}{N}\right)$.*

*Proof.* Set $c_N := 1 - \frac{n}{N}$.

(i) Replace unknown finite-population variance by $s^2$, correction by $c_N$ gives $s^2_{\overline{X}} = (\frac{s^2}{n})c_N = \frac{s^2}{n}(1 - \frac{n}{N})$.
(ii) Since $T = N\overline{X}$, variance scales quadratically, so $s^2_T = N^2 s^2_{\overline{X}}$.
(iii) If $X_k \in \{0,1\}$ and $\widehat{p} = \overline{X}$, then $s^2 = \frac{1}{n-1}\sum_{k=1}^{n}(X_k - \widehat{p})^2 = \frac{1}{n-1}(n\widehat{p} - n\widehat{p}^2) = \frac{n\widehat{p}(1-\widehat{p})}{n-1}$.
Substituting this into (i) gives $s^2_{\widehat{p}} = \frac{\widehat{p}(1-\widehat{p})c_N}{n-1} = \frac{\widehat{p}(1-\widehat{p})}{n-1}(1 - \frac{n}{N})$.

*Structural Commentary*: Plug-in SEs use $s^2$ and the $1 - \frac{n}{N}$ correction.
*Key Insight*: Binary-data variance depends only on $\widehat{p}$. □

**Example 1.1.10.** *Finite Population Correction in a Hospital Survey*
For a finite population of $N = 393$ hospitals, let $x_i$ denote discharges and suppose the finite-population standard deviation is $\sigma \approx 589.7$. Under simple random sampling without replacement with $n = 32$, the sample-mean standard deviation is $\sigma_{\overline{X}} = \frac{\sigma}{\sqrt{n}}\sqrt{1 - \frac{n-1}{N-1}} \approx \frac{589.7}{\sqrt{32}}\sqrt{1 - \frac{31}{392}} \approx 100.0$. Since $\frac{n}{N} = \frac{32}{393}$ is small, the finite-population correction is near 1, and the standard error is mainly governed by the $\frac{1}{\sqrt{n}}$ averaging scale.

## 1.2 Normal Approximation and Confidence Intervals

The exact design formulas in Theorem 1.1.8 give the center and scale of the sampling distribution. For large samples, a normal approximation turns those formulas into error probabilities and confidence intervals.

**Method 1.2.1.** *Normal Approximation for SRS*

(i) Let $U := \{1, \dots, N\}$, $N > 1$, $1 \le n \le N$, $x_i \in \mathbb{R}$ for $i \in U$, and $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$.
(ii) For a random ordering $(J_1, \dots, J_n)$ of $S$, set $X_k := x_{J_k}$, $\overline{X} := n^{-1}\sum_{k=1}^{n} X_k$, $T := N\overline{X}$, $\mu := N^{-1}\sum_{i=1}^{N} x_i$, $\sigma^2 := N^{-1}\sum_{i=1}^{N}(x_i - \mu)^2$, $\tau := N\mu$, and $\sigma^2_{\overline{X}} := \frac{\sigma^2}{n}\left(\frac{N-n}{N-1}\right)$.
(iii) Let $\Phi$ be the standard normal cdf. For large $n$, use $\overline{X} \approx N(\mu, \sigma^2_{\overline{X}})$.
(iv) If $\sigma_{\overline{X}} > 0$, then $\mathbb{P}(|\overline{X} - \mu| \le \delta) \approx 2\Phi(\frac{\delta}{\sigma_{\overline{X}}}) - 1$ for $\delta > 0$, and $T - \tau = N(\overline{X} - \mu) \approx N(0, N^2\sigma^2_{\overline{X}})$.

**Method 1.2.2.** *Approximate SRS Confidence Intervals*

(i) Let $S$ be an SRS without replacement, with targets $\mu, \tau$ and estimators $\overline{X}, T$.
(ii) Fix $0 < \alpha < 1$, let $\Phi$ be the standard normal cdf, set $z(\frac{\alpha}{2}) := \Phi^{-1}(1 - \frac{\alpha}{2})$, and use plug-in SEs $s_{\overline{X}}, s_T$, and, when binary, $s_{\widehat{p}}$.
(iii) $\mathrm{CI}_{1-\alpha}(\mu) = \overline{X} \pm z(\frac{\alpha}{2})s_{\overline{X}}$ and $\mathrm{CI}_{1-\alpha}(\tau) = T \pm z(\frac{\alpha}{2})s_T$.
(iv) In the dichotomous case, with population proportion $p$ and sample proportion $\widehat{p}$, use $\mathrm{CI}_{1-\alpha}(p) = \widehat{p} \pm z(\frac{\alpha}{2})s_{\widehat{p}}$.
(v) Intervals are random before sampling; after observation, parameters are fixed and coverage is procedural.

**Example 1.2.3.** *Generic SRS Intervals*
Suppose finite population has $N = 8000$ units. Simple random sample of $n = 100$ gives $\overline{X} = 1.6$ for variable with sample standard deviation $s = 0.8$. Use $z(0.025) \approx 1.96$, suppose sample gives $\widehat{p} = 0.12$ for binary attribute.

(i) *Mean*: $s_{\overline{X}} = \frac{s}{\sqrt{n}}\sqrt{1 - \frac{n}{N}} = \frac{0.8}{10}\sqrt{1 - \frac{100}{8000}} \approx 0.0795$, then 95% CI for $\mu$ is $1.6 \pm 1.96(0.0795) = (1.44, 1.76)$.
(ii) *Total*: $T = N\overline{X} = 8000(1.6) = 12800$ and $s_T = Ns_{\overline{X}} \approx 636$, then 95% CI for $\tau$ is $(11553, 14047)$.
(iii) *Proportion*: $s_{\widehat{p}} = \sqrt{\frac{\widehat{p}(1-\widehat{p})}{n-1}\left(1 - \frac{n}{N}\right)} = \sqrt{\frac{0.12(0.88)}{99}\left(1 - \frac{100}{8000}\right)} \approx 0.0325$, then 95% CI for $p$ is $(0.056, 0.184)$. The estimated attribute total is $N\widehat{p} = 960$ with $Ns_{\widehat{p}} \approx 260$ and interval $(451, 1469)$.

These intervals quantify SRS uncertainty for fixed finite-population quantities.

**Method 1.2.4.** *Sample Size from Desired Half-Width*

(i) Fix $0 < \alpha < 1$, half-width $h > 0$, planning SD $\sigma > 0$, and planned size $n$.
Let $\Phi$ be the standard normal cdf and set $z(\frac{\alpha}{2}) := \Phi^{-1}(1 - \frac{\alpha}{2})$.

(ii) Ignoring the finite population correction, $h \approx z(\frac{\alpha}{2})\frac{\sigma}{\sqrt{n}}$ gives $n \approx \left(\frac{z(\frac{\alpha}{2})\sigma}{h}\right)^2$.

(iii) For total based on population proportion over $N$ units, use planning value $p_0 \in [0,1]$, SE $N\sqrt{\frac{p_0(1-p_0)}{n}}$.

(iv) With negligible finite population correction, halving interval width requires roughly $4n$ observations.

## 1.3 Ratio Estimation

Ratio estimation uses an auxiliary variable whose population mean or total is known. It is especially useful when the auxiliary variable is strongly correlated with the target variable.

**Definition 1.3.1.** *Population and Sample Ratios*
Let $U := \{1, \dots, N\}$ and fixed pairs $(x_i, y_i) \in \mathbb{R}^2$, $i \in U$.
Set $\mu_x := \frac{1}{N}\sum_{i=1}^{N} x_i$, $\mu_y := \frac{1}{N}\sum_{i=1}^{N} y_i$, $\tau_x := \sum_{i=1}^{N} x_i$, and $\tau_y := \sum_{i=1}^{N} y_i$, with $\mu_x \neq 0$.
For SRS pairs $(X_i, Y_i)_{i=1}^n$, set $\overline{X} := \frac{1}{n}\sum_{i=1}^{n} X_i$ and $\overline{Y} := \frac{1}{n}\sum_{i=1}^{n} Y_i$, with $\overline{X} \neq 0$.

(i) *Population ratio*: $r := \sum_{i=1}^{N} y_i / \sum_{i=1}^{N} x_i = \frac{\mu_y}{\mu_x}$.

(ii) *Sample ratio*: $R := \frac{\overline{Y}}{\overline{X}}$.

(iii) *Ratio mean estimator*: $\overline{Y}_R := \mu_x R$ when $\mu_x$ is known.

(iv) *Ratio total estimator*: $T_R := \tau_x R$ when $\tau_x$ is known.

*Key Insight*: $\{i \in U : x_i = 0\} = \emptyset$ does not imply $r = \frac{1}{N}\sum_{i=1}^{N} \frac{y_i}{x_i}$.

**Definition 1.3.2.** *Population Covariance and Correlation*
Let $U := \{1, \dots, N\}$ and fixed pairs $(x_i, y_i) \in \mathbb{R}^2$, $i \in U$.
Set $\mu_x := \frac{1}{N}\sum_{i=1}^{N} x_i$, $\mu_y := \frac{1}{N}\sum_{i=1}^{N} y_i$, $\sigma_x^2 := \frac{1}{N}\sum_{i=1}^{N}(x_i - \mu_x)^2$, and $\sigma_y^2 := \frac{1}{N}\sum_{i=1}^{N}(y_i - \mu_y)^2$.
For sample pairs $(X_i, Y_i)_{i=1}^n$, set $\overline{X} := \frac{1}{n}\sum_{i=1}^{n} X_i$ and $\overline{Y} := \frac{1}{n}\sum_{i=1}^{n} Y_i$.

(i) *Population covariance*: $\sigma_{xy} := \frac{1}{N}\sum_{i=1}^{N}(x_i - \mu_x)(y_i - \mu_y)$.

(ii) *Population correlation*: $\rho := \frac{\sigma_{xy}}{\sigma_x \sigma_y}$ when $\sigma_x, \sigma_y > 0$.

(iii) *Sample covariance*: $s_{xy} := \frac{1}{n-1}\sum_{i=1}^{n}(X_i - \overline{X})(Y_i - \overline{Y})$.

**Theorem 1.3.3.** *Approximate Moments of the Ratio Estimator*
*Let $U := \{1, \dots, N\}$, $N > 1$, $1 \le n \le N$, and let $(x_a, y_a) \in \mathbb{R}^2$ be fixed for $a \in U$.*
*Set $\lambda_n := \frac{1}{n}(1 - \frac{n-1}{N-1})$, $\mu_x := \frac{1}{N}\sum_{a=1}^{N} x_a$, $\mu_y := \frac{1}{N}\sum_{a=1}^{N} y_a$, $r := \frac{\mu_y}{\mu_x}$*
*with $\mu_x \neq 0$, $\sigma_x^2 := \frac{1}{N}\sum_{a=1}^{N}(x_a - \mu_x)^2$, $\sigma_y^2 := \frac{1}{N}\sum_{a=1}^{N}(y_a - \mu_y)^2$, and $\sigma_{xy} := \frac{1}{N}\sum_{a=1}^{N}(x_a - \mu_x)(y_a - \mu_y)$.*
*Let $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$, let $(J_1, \dots, J_n)$ be a random ordering of $S$, set $(X_k, Y_k) := (x_{J_k}, y_{J_k})$, $\overline{X} := \frac{1}{n}\sum_{k=1}^{n} X_k$, $\overline{Y} := \frac{1}{n}\sum_{k=1}^{n} Y_k$, and $R := \frac{\overline{Y}}{\overline{X}}$ on $\{\overline{X} \neq 0\}$. The subscript $d$ denotes design moments.*

*(i) Variance: first-order Taylor gives $\mathrm{Var}_d(R) \approx \frac{\lambda_n}{\mu_x^2}(r^2\sigma_x^2 + \sigma_y^2 - 2r\sigma_{xy})$.*

*(ii) Mean: second-order Taylor gives $\mathbb{E}_d R \approx r + \frac{\lambda_n}{\mu_x^2}(r\sigma_x^2 - \sigma_{xy})$.*

*Proof.* Let $(A_q, B_q)_{q=1}^N$ be a uniformly random ordering of the population pairs; the ordered sample has the law of the first $n$ entries. For $q \neq r$, write $\gamma_x := \mathrm{Cov}_d(A_q, A_r)$, $\gamma_y := \mathrm{Cov}_d(B_q, B_r)$, and $\gamma_{xy} := \mathrm{Cov}_d(A_q, B_r)$.
The fixed sums give $0 = \mathrm{Var}_d(\sum_{q=1}^{N} A_q) = N\sigma_x^2 + N(N-1)\gamma_x$, $0 = \mathrm{Var}_d(\sum_{q=1}^{N} B_q) = N\sigma_y^2 + N(N-1)\gamma_y$,
and $0 = \mathrm{Cov}_d(\sum_{q=1}^{N} A_q, \sum_{q=1}^{N} B_q) = N\sigma_{xy} + N(N-1)\gamma_{xy}$.

Hence $\gamma_x = -\frac{\sigma_x^2}{N-1}$, $\gamma_y = -\frac{\sigma_y^2}{N-1}$, $\gamma_{xy} = -\frac{\sigma_{xy}}{N-1}$, $V_x := \mathrm{Var}_d(\overline{X}) = \lambda_n \sigma_x^2$, $V_y := \mathrm{Var}_d(\overline{Y}) = \lambda_n \sigma_y^2$, and $C_{xy} := \mathrm{Cov}_d(\overline{X}, \overline{Y}) = \lambda_n \sigma_{xy}$. Let $\Sigma := \begin{pmatrix} V_x & C_{xy} \\ C_{xy} & V_y \end{pmatrix}$.

For $g(x,y) = \frac{y}{x}$ and $\mu := (\mu_x, \mu_y)$, $g(\mu) = r$, $\nabla g(\mu) = (-\frac{r}{\mu_x}, \frac{1}{\mu_x})^\top$, and $H_g(\mu) = \begin{pmatrix} 2r/\mu_x^2 & -1/\mu_x^2 \\ -1/\mu_x^2 & 0 \end{pmatrix}$.

(i) *Variance*: $\mathrm{Var}_d(R) \approx \nabla g(\mu)^\top \Sigma \nabla g(\mu) = \frac{\lambda_n}{\mu_x^2}(r^2\sigma_x^2 + \sigma_y^2 - 2r\sigma_{xy})$.

(ii) *Mean*: $\mathbb{E}_d R \approx g(\mu) + \frac{1}{2}\operatorname{tr}(H_g(\mu)\Sigma) = r + \frac{\lambda_n}{\mu_x^2}(r\sigma_x^2 - \sigma_{xy})$.

*Structural Commentary*: The ratio estimator is locally linear for variance and locally quadratic for bias.
*Key Insight*: Ratio variance is scaled residual variation in $Y - rX$. □

**Corollary 1.3.4.** *Estimated Standard Error of a Ratio*
*Let $U := \{1, \dots, N\}$, $N > 1$, $1 < n \le N$, and let $(x_a, y_a) \in \mathbb{R}^2$ be fixed for $a \in U$.*
*Set $\lambda_n := \frac{1}{n}(1 - \frac{n-1}{N-1})$, $\mu_x := \frac{1}{N}\sum_{a=1}^N x_a$, $\tau_x := \sum_{a=1}^N x_a$, $\tau_y := \sum_{a=1}^N y_a$, and $r := \frac{\tau_y}{\tau_x}$ with $\tau_x \neq 0$.*
*Let $S \sim \mathrm{Unif}\{s \subseteq U : |s| = n\}$. For a random ordering $(J_1, \dots, J_n)$ of $S$,*
*set $(X_i, Y_i) := (x_{J_i}, y_{J_i})$, $\overline{X} := \frac{1}{n}\sum_{i=1}^n X_i$, and $\overline{Y} := \frac{1}{n}\sum_{i=1}^n Y_i$.*
*Assume $\overline{X} \neq 0$ and set $R := \frac{\overline{Y}}{\overline{X}}$, $s_x^2 := \frac{1}{n-1}\sum_{i=1}^n (X_i - \overline{X})^2$, $s_y^2 := \frac{1}{n-1}\sum_{i=1}^n (Y_i - \overline{Y})^2$, $s_{xy} := \frac{1}{n-1}\sum_{i=1}^n (X_i - \overline{X})(Y_i - \overline{Y})$, and $q_R := R^2 s_x^2 + s_y^2 - 2R s_{xy}$. If $0 < \alpha < 1$ and $z(\frac{\alpha}{2}) := \Phi^{-1}(1 - \frac{\alpha}{2})$ for the standard normal cdf $\Phi$, then:*

*(i) Ratio variance: $s_R^2 := \frac{\lambda_n}{\overline{X}^2} q_R$ estimates $\mathrm{Var}_d(R)$.*
*(ii) Ratio interval: $\mathrm{CI}_{1-\alpha}(r) := R \pm z(\frac{\alpha}{2}) s_R$.*
*(iii) Mean and total: for $\overline{Y}_R := \mu_x R$ and $T_R := \tau_x R$, $\mathrm{SE}(\overline{Y}_R) = |\mu_x|\, s_R$ and $\mathrm{SE}(T_R) = |\tau_x|\, s_R$ when the corresponding auxiliary constant is known.*

*Proof.*

(i) The plug-in map $(\mu_x, r, \sigma_x^2, \sigma_y^2, \sigma_{xy}) \mapsto (\overline{X}, R, s_x^2, s_y^2, s_{xy})$ in Theorem 1.3.3 yields $\widehat{\mathrm{Var}}_d(R) = \frac{\lambda_n}{\overline{X}^2}(R^2 s_x^2 + s_y^2 - 2R s_{xy}) = s_R^2$.

(ii) If $Z_R := (R - r)/s_R \approx \mathcal{N}(0,1)$, then $\mathbb{P}_d(|Z_R| \le z(\frac{\alpha}{2})) \approx 1 - \alpha$, equivalently $r \in R \pm z(\frac{\alpha}{2}) s_R$.

(iii) For known $c$, $\widehat{\mathrm{Var}}_d(cR) = c^2 s_R^2$, so $\mathrm{SE}(cR) = |c|\, s_R$; set $c = \mu_x$ and $c = \tau_x$.

*Structural Commentary*: Plug in sample moments to estimate the ratio variance.
*Key Insight*: Ratio uncertainty is residual uncertainty in $Y - RX$. □

**Method 1.3.5.** *When the Ratio Estimate Helps*
Let $U = \{1, \dots, N\}$ contain fixed pairs $(x_i, y_i)$ with $\mu_x, \mu_y > 0$, $\sigma_x, \sigma_y > 0$, $\sigma_{xy}$, $\rho := \frac{\sigma_{xy}}{\sigma_x \sigma_y}$, and $r := \frac{\mu_y}{\mu_x}$.

For an SRS $(X_k, Y_k)_{k=1}^n$, set $\overline{X} := \frac{1}{n}\sum_{k=1}^n X_k$, $\overline{Y} := \frac{1}{n}\sum_{k=1}^n Y_k$, $R := \frac{\overline{Y}}{\overline{X}}$, and $\overline{Y}_R := \mu_x R$.

Ignoring finite-population corrections:

(i) *Variances*: $V_0 := \mathrm{Var}(\overline{Y}) = \frac{\sigma_y^2}{n}$ and $V_R^* := \frac{1}{n}(r^2\sigma_x^2 + \sigma_y^2 - 2r\sigma_{xy})$ approximate $\mathrm{Var}(\overline{Y}_R)$.

(ii) *Improvement*: $V_R^* < V_0 \iff 2\sigma_{xy} > r\sigma_x^2 \iff 2\rho\sigma_y > r\sigma_x$.

(iii) *Coefficient form*: with $C_x := \frac{\sigma_x}{\mu_x}$ and $C_y := \frac{\sigma_y}{\mu_y}$, $V_R^* < V_0 \iff \rho > \frac{1}{2}\frac{C_x}{C_y}$.

(iv) *Favorable case*: large positive $\rho$ and small $C_x/C_y$.

**Example 1.3.6.** *Mortgage-Payment Ratio*
For an SRS of $n = 100$ recent house buyers, let $Y_i$ be monthly mortgage payment and $X_i$ gross income. Suppose $\overline{X} = 3100$, $\overline{Y} = 868$, $s_x = 1200$, $s_y = 250$, and $\widehat{\rho} := s_{xy}/(s_x s_y) = 0.85$.

(i) *Ratio*: $R = \frac{\overline{Y}}{\overline{X}} = \frac{868}{3100} = 0.28$.

(ii) *Variance input*: $q := R^2 s_x^2 + s_y^2 - 2R\widehat{\rho} s_x s_y \approx 32596$.

(iii) *Standard error*: ignoring the finite-population correction, $s_R \approx \frac{\sqrt{q}}{\sqrt{100}\,\overline{X}} \approx 0.00582$.

(iv) *Interval*: $\mathrm{CI}_{0.95}(r) = R \pm 1.96 s_R \approx 0.28 \pm 1.96(0.00582) = (0.269, 0.291)$.

Thus the population mortgage-payment-to-income ratio is about 28%.

**Remark 1.3.7.** *Bias and Resampling Corrections*
For SRS from fixed pairs $(x_i, y_i)$, set $R := \overline{Y}/\overline{X}$, $\overline{Y}_R := \mu_x R$, and $T_R := \tau_x R$.

(i) Since $g(x,y) = y/x$ is nonlinear, generally $\mathbb{E}_d R \neq r$, hence $\mathbb{E}_d \overline{Y}_R \neq \mu_y$ and $\mathbb{E}_d T_R \neq \tau_y$.

(ii) Bias corrections target $B_d(R) := \mathbb{E}_d R - r$; use resampling when Taylor bias matters.

## 1.4 Stratified Random Sampling

Stratified sampling partitions the population into subpopulations and samples within each one. It can guarantee representation of important groups and can reduce variance when strata are internally homogeneous or have different means.

**Definition 1.4.1.** *Strata and the Stratified Mean*

Let $U$ be a finite population, $L \geq 1$, nonempty subsets $U_1, \ldots, U_L \subseteq U$, $N_l := |U_l|$, $N := \sum_{l=1}^{L} N_l$, and fixed values $x_{il} \in \mathbb{R}$ for $i = 1, \ldots, N_l$, $l = 1, \ldots, L$.
For stratum samples of sizes $n_l$, write sampled values as $X_{il}$, $i = 1, \ldots, n_l$.

(i) *Strata*: $U = \bigsqcup_{l=1}^{L} U_l$.

(ii) *Stratum weight*: $W_l := \frac{N_l}{N}$.

(iii) *Stratum mean*: $\mu_l := \frac{1}{N_l} \sum_{i=1}^{N_l} x_{il}$.

(iv) *Stratum variance*: $\sigma_l^2 := \frac{1}{N_l} \sum_{i=1}^{N_l} (x_{il} - \mu_l)^2$.

(v) *Stratum sample mean*: $\overline{X}_l := \frac{1}{n_l} \sum_{i=1}^{n_l} X_{il}$.

(vi) *Stratified mean*: $\overline{X}_s := \sum_{l=1}^{L} W_l \overline{X}_l$.

*Structural Commentary*: With $\mu := \frac{1}{N} \sum_{l=1}^{L} \sum_{i=1}^{N_l} x_{il}$, $\mu = \sum_{l=1}^{L} W_l \mu_l$.

**Theorem 1.4.2.** *Mean and Variance of the Stratified Estimate*

*Let $L \geq 1$, $N_l > 1$, $1 \leq n_l \leq N_l$, $N := \sum_{l=1}^{L} N_l$, $W_l := \frac{N_l}{N}$, and $\lambda_l := \frac{1}{n_l}(1 - \frac{n_l - 1}{N_l - 1})$.*

*For fixed $x_{il} \in \mathbb{R}$, $i = 1, \ldots, N_l$, $l = 1, \ldots, L$, set $\mu_l := \frac{1}{N_l} \sum_{i=1}^{N_l} x_{il}$, $\sigma_l^2 := \frac{1}{N_l} \sum_{i=1}^{N_l} (x_{il} - \mu_l)^2$, and $\mu := \sum_{l=1}^{L} W_l \mu_l$.*
*Independently sample $S_l \sim \mathrm{Unif}\{s \subseteq \{1, \ldots, N_l\} : |s| = n_l\}$; for random orderings $(J_{il})_{i=1}^{n_l}$ of $S_l$, set $X_{il} := x_{J_{il},l}$, $\overline{X}_l := \frac{1}{n_l} \sum_{i=1}^{n_l} X_{il}$, and $\overline{X}_s := \sum_{l=1}^{L} W_l \overline{X}_l$. Let $\mathbb{E}_d, \mathrm{Var}_d$ denote design moments.*

*(i) Mean:* $\mathbb{E}_d \overline{X}_s = \mu$.

*(ii) Variance:* $\mathrm{Var}_d(\overline{X}_s) = \sum_{l=1}^{L} W_l^2 \lambda_l \sigma_l^2$.

*(iii) Small fractions: if $n_l / N_l \ll 1$ for all $l$, then* $\mathrm{Var}_d(\overline{X}_s) \approx \sum_{l=1}^{L} W_l^2 \frac{\sigma_l^2}{n_l}$.

*Proof.*

(i) For $a \in \{1, \ldots, N_l\}$, $\mathbb{P}_d(J_{il} = a) = 1/N_l$, hence $\mathbb{E}_d X_{il} = \mu_l$, $\mathbb{E}_d \overline{X}_l = \mu_l$, and $\mathbb{E}_d \overline{X}_s = \sum_{l=1}^{L} W_l \mu_l = \mu$.

(ii) Independence across strata gives $\mathrm{Var}_d(\overline{X}_s) = \sum_{l=1}^{L} W_l^2 \mathrm{Var}_d(\overline{X}_l)$.
For stratum $l$, let $(Y_{1,l}, \ldots, Y_{N_l,l})$ uniformly order $\{x_{il} : 1 \leq i \leq N_l\}$, put $\gamma_l := \mathrm{Cov}_d(Y_{a,l}, Y_{b,l})$ for $a \neq b$.
As $\sum_{a=1}^{N_l} Y_{a,l}$ is fixed, $0 = N_l \sigma_l^2 + N_l(N_l - 1)\gamma_l$, so $\gamma_l = -\frac{\sigma_l^2}{N_l - 1}$, $\mathrm{Var}_d(\overline{X}_l) = \frac{1}{n_l^2}(n_l \sigma_l^2 + n_l(n_l - 1)\gamma_l) = \lambda_l \sigma_l^2$.

(iii) If $n_l / N_l \ll 1$, then $1 - \frac{n_l - 1}{N_l - 1} \approx 1$, so $\lambda_l \sigma_l^2 \approx \sigma_l^2 / n_l$.

*Structural Commentary*: Stratified estimation is a weighted sum of independent within-stratum estimators.
*Key Insight*: Independent strata make variance add as $\sum_l W_l^2 \mathrm{Var}_d(\overline{X}_l)$. □

**Corollary 1.4.3.** *Totals and Estimated Variances*

*Let $L \geq 1$, $N_l > 1$, $1 < n_l \leq N_l$, $N := \sum_{l=1}^{L} N_l$, $W_l := \frac{N_l}{N}$, and $f_l := n_l / N_l$.*

*Fix $x_{il} \in \mathbb{R}$ for $i = 1, \ldots, N_l$, $l = 1, \ldots, L$, set $\mu_l := \frac{1}{N_l} \sum_{i=1}^{N_l} x_{il}$, $\mu := \sum_{l=1}^{L} W_l \mu_l$, and $\tau := N\mu$.*
*Assume independently $S_l \sim \mathrm{Unif}\{s \subseteq \{1, \ldots, N_l\} : |s| = n_l\}$; for random orderings $(J_{il})_{i=1}^{n_l}$ of $S_l$.*

*Set* $X_{il} := x_{J_{il},l}$, $\overline{X}_l := \frac{1}{n_l}\sum_{i=1}^{n_l} X_{il}$, $\overline{X}_s := \sum_{l=1}^{L} W_l \overline{X}_l$, $T_s := N\overline{X}_s$, *and* $s_l^2 := \frac{1}{n_l-1}\sum_{i=1}^{n_l}(X_{il}-\overline{X}_l)^2$.
*For* $0<\alpha<1$, *let* $z(\frac{\alpha}{2}) := \Phi^{-1}(1-\frac{\alpha}{2})$.

*(i) Total moments:* $\mathbb{E}_d T_s = \tau$ *and* $\mathrm{Var}_d(T_s) = N^2 \mathrm{Var}_d(\overline{X}_s)$.

*(ii) Estimated mean variance:* $s^2_{\overline{X}_s} := \sum_{l=1}^{L} W_l^2(1-f_l)\frac{s_l^2}{n_l}$.

*(iii) Intervals:* $\mathrm{CI}_{1-\alpha}(\mu) = \overline{X}_s \pm z(\frac{\alpha}{2}) s_{\overline{X}_s}$ *and* $\mathrm{CI}_{1-\alpha}(\tau) = T_s \pm z(\frac{\alpha}{2}) N s_{\overline{X}_s}$.

*Proof.*

(i) Since $T_s = N\overline{X}_s$ and $\tau = N\mu$, Theorem 1.4.2 gives $\mathbb{E}_d T_s = N\mu = \tau$ and $\mathrm{Var}_d(T_s) = N^2 \mathrm{Var}_d(\overline{X}_s)$.

(ii) Let $S_l^2 := \frac{1}{N_l-1}\sum_{i=1}^{N_l}(x_{il}-\mu_l)^2$. Then $\mathrm{Var}_d(\overline{X}_l) = \frac{(1-f_l)S_l^2}{n_l}$, $\mathbb{E}_d s_l^2 = S_l^2$, so $S_l^2 \mapsto s_l^2$; $s^2_{\overline{X}_s} = \sum_{l=1}^{L} W_l^2(1-f_l)\frac{s_l^2}{n_l}$.

(iii) With $\frac{\overline{X}_s-\mu}{s_{\overline{X}_s}} \approx \mathcal{N}(0,1)$, inversion gives $\mathrm{CI}_{1-\alpha}(\mu)$; multiplying by $N$ gives $\mathrm{CI}_{1-\alpha}(\tau)$.

*Structural Commentary*: Totals and intervals scale the stratified mean.
*Key Insight*: Estimated variance adds weighted stratum variances. □

**Theorem 1.4.4.** *Neyman Allocation*
*Let* $L \geq 1$, $W_l > 0$, $\sum_{l=1}^{L} W_l = 1$, $\sigma_l > 0$, $S_\sigma := \sum_{k=1}^{L} W_k\sigma_k$, *and* $n > 0$.
*On* $\mathcal{A} := \{(n_1,\ldots,n_L) \in (0,\infty)^L : \sum_{l=1}^{L} n_l = n\}$, *set* $V(n_1,\ldots,n_L) := \sum_{l=1}^{L} \frac{W_l^2\sigma_l^2}{n_l}$.
*Then* $V$ *has the unique minimiser* $n_l^* := n\frac{W_l\sigma_l}{S_\sigma}$, $l = 1,\ldots,L$.

*Proof.* $V$ is strictly convex on $(0,\infty)^L$ since $\partial_{ll}V = \frac{2W_l^2\sigma_l^2}{n_l^3} > 0$ and $\partial_{lm}V = 0$ for $l \neq m$.

For $\mathcal{L}(n,\lambda) := V(n) + \lambda(\sum_{l=1}^{L} n_l - n)$, the first-order equations are $-\frac{W_l^2\sigma_l^2}{n_l^2} + \lambda = 0$, hence $n_l = \frac{W_l\sigma_l}{\sqrt{\lambda}}$.
The constraint gives $\frac{1}{\sqrt{\lambda}} = \frac{n}{S_\sigma}$, so $n_l = n_l^*$; strict convexity gives uniqueness.
*Structural Commentary*: The constrained optimum balances marginal variance reduction across strata.
*Key Insight*: A stratum receives more sample not only when it is large, but also when it is internally variable. □

**Corollary 1.4.5.** *Allocation Comparisons*
*Let* $L \geq 1$, $W_l > 0$, $\sum_{l=1}^{L} W_l = 1$, $\mu_l \in \mathbb{R}$, $\sigma_l > 0$, $\mu := \sum_{l=1}^{L} W_l\mu_l$, $\overline{\sigma} := \sum_{l=1}^{L} W_l\sigma_l$, *and* $n > 0$.
*Ignoring finite-population corrections, let* $\overline{X}_{so}$ *be the optimal-allocation stratified mean,* $\overline{X}_{sp}$ *the proportional-allocation stratified mean with* $n_l = nW_l$, *and* $\overline{X}$ *the ordinary SRS mean.*

*(i) Allocation variances:* $\mathrm{Var}(\overline{X}_{so}) \approx \frac{\overline{\sigma}^2}{n}$ *and* $\mathrm{Var}(\overline{X}_{sp}) \approx \frac{1}{n}\sum_{l=1}^{L} W_l\sigma_l^2$.

*(ii) Optimal-allocation gain:* $\mathrm{Var}(\overline{X}_{sp}) - \mathrm{Var}(\overline{X}_{so}) \approx \frac{1}{n}\sum_{l=1}^{L} W_l(\sigma_l-\overline{\sigma})^2$.

*(iii) Proportional-allocation gain:* $\mathrm{Var}(\overline{X}) - \mathrm{Var}(\overline{X}_{sp}) \approx \frac{1}{n}\sum_{l=1}^{L} W_l(\mu_l-\mu)^2$.

*Proof.*

(i) With finite-population corrections suppressed, $\mathrm{Var}(\overline{X}_s) \approx \sum_{l=1}^{L} \frac{W_l^2\sigma_l^2}{n_l}$. Neyman allocation gives $n_l = \frac{nW_l\sigma_l}{\overline{\sigma}}$, hence $\mathrm{Var}(\overline{X}_{so}) \approx \frac{\overline{\sigma}^2}{n}$; proportional allocation $n_l = nW_l$ gives $\mathrm{Var}(\overline{X}_{sp}) \approx \frac{1}{n}\sum_{l=1}^{L} W_l\sigma_l^2$.

(ii) Since $\sum_{l=1}^{L} W_l = 1$, $\sum_{l=1}^{L} W_l\sigma_l^2 - \overline{\sigma}^2 = \sum_{l=1}^{L} W_l(\sigma_l-\overline{\sigma})^2$.

(iii) For ordinary SRS, $\mathrm{Var}(\overline{X}) \approx \frac{1}{n}\{\sum_{l=1}^{L} W_l\sigma_l^2 + \sum_{l=1}^{L} W_l(\mu_l-\mu)^2\}$; subtract the proportional-allocation variance.

*Structural Commentary*: Proportional handles means; optimal also handles spreads.
*Key Insight*: Mean gaps help stratification; spread gaps help Neyman allocation. □

**Example 1.4.6.** *Four-Stratum Stratification Summary*
Let $N = 393$ and strata $l \in \{A,B,C,D\}$ have rounded summaries $(N_l) = (98,98,98,99)$, $(W_l) = (.249,.249,.249,.251)$,

$(\mu_l) = (182.9, 526.5, 956.3, 1591.2)$, and $(\sigma_l) = (103.4, 204.8, 243.5, 419.2)$. Let $n$ be the total sample size, ignore finite-population corrections, and write $\overline{X}_s$ for the stratified mean.

(i) *Equal allocation*: if $n_l = n/4$, then $\mathrm{Var}(\overline{X}_s) \approx \sum_{l=1}^{4} W_l^2 \sigma_l^2/(n/4) \approx 72042.6/n$, so $\mathrm{SE}(\overline{X}_s) \approx 268.4/\sqrt{n}$.
(ii) *SRS benchmark*: with overall standard deviation $\sigma \approx 589.7$, $\mathrm{SE}(\overline{X}) \approx 589.7/\sqrt{n}$.
(iii) *Neyman allocation*: $a_l := \frac{W_l \sigma_l}{\sum_{k=1}^{4} W_k \sigma_k} \approx (.106, .210, .250, .434)$ for $(A, B, C, D)$.

Stratification reduces variance, and Neyman allocation assigns the largest share to the stratum with largest $\sigma_l$.

**Remark 1.4.7.** *Model Limits and Other Designs*
For a finite-population survey:

(i) *Systematic*: fixed-interval sampling after a random start is SRS-like only when list order is noninformative; periodic order can bias selection.
(ii) *Cluster*: sampling groups lowers cost but can inflate variance when within-cluster correlation is positive.
(iii) *Scope*: design-based SEs measure random selection variation, not nonresponse, coverage error, selection bias, interviewer effects, or wording effects.

# 2 Exploratory Data Analysis and Data Summaries

## 2.1 Empirical Distribution and Survival Summaries

Exploratory summaries first treat the observed values as a batch of numbers. A stochastic model may later explain their variability, but the displays themselves are useful before a model has been chosen.

**Definition 2.1.1.** *Empirical CDF and Empirical Survival Function*
Let $x_1, \ldots, x_n \in \mathbb{R}$ be data; for survival summaries assume $x_i \geq 0$ for $1 \leq i \leq n$.

(i) *Empirical CDF*: $F_n : \mathbb{R} \to [0,1]$, $F_n(x) := n^{-1}\#\{i \in \{1, \ldots, n\} : x_i \leq x\}$.
(ii) *Empirical Survival Function*: $S_n : [0, \infty) \to [0,1]$, $S_n(t) := 1 - F_n(t) = n^{-1}\#\{i \in \{1, \ldots, n\} : x_i > t\}$.

*Key Insight*: $F_n$ and $S_n$ encode complementary empirical tails, with $S_n(t) = 1 - F_n(t)$ for $t \geq 0$.

**Remark 2.1.2.** *Step-Function Convention*
Let $v_1 < \cdots < v_m$ be the distinct observed values, with multiplicities $r_1, \ldots, r_m$. Then $F_n$ is right-continuous, constant on each component of $\mathbb{R} \setminus \{v_1, \ldots, v_m\}$, and has jump $\Delta F_n(v_j) = \frac{r_j}{n}$.




Figure 245: Empirical CDF (blue) rises as empirical survival (green) falls.

**Method 2.1.3.** *Empirical Log-Survival Plot*
Suppose $0 \leq T_{(1)} < \cdots < T_{(n)}$ and there are no ties. Set $p_i := \frac{i}{n+1}$ and $y_i := \log(1 - p_i)$.

(i) *Plot*: Draw $P^{(n)}_{\log S} := \{(T_{(i)}, y_i) : 1 \leq i \leq n\}$; using $n+1$ keeps $y_n$ finite.
(ii) *Hazard Reading*: Since $(\log S)' = -h$, more negative local slopes indicate larger hazards.
(iii) *Tail Variability*: $\mathrm{Var}(\log[1 - F_n(t)]) \approx n^{-1} \frac{F(t)}{1 - F(t)}$, which grows as $F(t) \uparrow 1$.

## 2.2 Quantile and Shape Displays

Distributional shape can be read from cumulative displays, density displays, or quantile comparisons. These views emphasize different questions: cumulative proportion, local concentration, and relative change between two distributions.

**Definition 2.2.1.** *Quantile–Quantile Plot*
Let $F$ and $G$ be continuous strictly increasing CDFs with quantile inverses $F^{-1}, G^{-1} : (0,1) \to \mathbb{R}$. For equal-sized batches $x_1, \dots, x_n$ and $y_1, \dots, y_n$, let $x_{(1)} \le \cdots \le x_{(n)}$ and $y_{(1)} \le \cdots \le y_{(n)}$.

(i) *Q–Q Plot*: $Q_{F,G} := \{(F^{-1}(p), G^{-1}(p)) : 0 < p < 1\} \subseteq \mathbb{R}^2$.
(ii) *Empirical Q–Q Plot*: $Q^{(n)}_{x,y} := \{(x_{(i)}, y_{(i)}) : 1 \le i \le n\} \subseteq \mathbb{R}^2$.

*Key Insight*: $(u, v) \in Q_{F,G}$ implies $F(u) = G(v)$, so plotted pairs match common cumulative probability levels.

**Proposition 2.2.2.** *Location and Scale Effects in Q–Q Plots*
*Let $F, G$ be continuous strictly increasing CDFs.*
*Set $x_p := F^{-1}(p)$, $y_p := G^{-1}(p)$ for $p \in (0,1)$, and $Q_{F,G} := \{(x_p, y_p) : 0 < p < 1\}$. Then:*

*(i) Location: If $y_p = x_p + h$ $\forall p \in (0,1)$ with $h \in \mathbb{R}$, then $G(y) = F(y-h)$ $\forall y \in \mathbb{R}$, $Q_{F,G} \subseteq \{(x,y) : y = x + h\}$.*
*(ii) Scale: If $y_p = cx_p$ $\forall p \in (0,1)$ with $c > 0$, then $G(y) = F(\frac{y}{c})$ $\forall y \in \mathbb{R}$, $Q_{F,G} \subseteq \{(x,y) : y = cx\}$.*

*Proof.*

(i) *Location*: Fix $y \in \mathbb{R}$ and set $p := G(y)$. Then $p \in (0,1)$, $y = y_p = x_p + h$, and $x_p = y - h$, so $G(y) = p = F(x_p) = F(y-h)$. Also $(x_p, y_p) \in Q_{F,G}$ implies $y_p = x_p + h$.
(ii) *Scale*: Fix $y \in \mathbb{R}$ and set $p := G(y)$. Then $p \in (0,1)$, $y = y_p = cx_p$, and $x_p = \frac{y}{c}$, so $G(y) = p = F(x_p) = F(\frac{y}{c})$. Also $(x_p, y_p) \in Q_{F,G}$ implies $y_p = cx_p$.

*Structural Commentary*: A Q–Q plot is the graph of the quantile transport $F^{-1}(p) \mapsto G^{-1}(p)$.
*Key Insight*: Straight Q–Q lines encode location shifts and scale changes. □




Figure 246: Q–Q lines: blue $y = x + h$ for location shift, green $y = cx$ for scale change.

**Example 2.2.3.** *Normal and Exponential Q–Q Lines*
For $p \in (0,1)$, set $x_p := F^{-1}(p)$ and $y_p := G^{-1}(p)$.

(i) *Normal Location*: If $F = N(0,1)$ and $G = N(1,1)$, then $y_p = x_p + 1$, so the Q–Q line is $y = x + 1$.
(ii) *Normal Location–Scale*: If $F = N(0,1)$ and $G = N(1,4)$, then $y_p = 1 + 2x_p$, so the Q–Q line is $y = 1 + 2x$.
(iii) *Exponential Rates*: If $F = \text{Exp}(\lambda_1)$ and $G = \text{Exp}(\lambda_2)$, then $x_p = -\frac{\log(1-p)}{\lambda_1}$ and $y_p = -\frac{\log(1-p)}{\lambda_2} = \frac{\lambda_1}{\lambda_2} x_p$, so the Q–Q line is $y = \frac{\lambda_1}{\lambda_2} x$.

**Method 2.2.4.** *Interpolated Empirical Quantiles*
Let $X_{(1)} \le \cdots \le X_{(n)}$ and set $p_k := \frac{k}{n+1}$ for $1 \le k \le n$.

(i) *Grid*: Assign $X_{(k)}$ to probability $p_k$.
(ii) *Interpolation*: If $p_k \le p \le p_{k+1}$ for $1 \le k < n$, set $\lambda := \frac{p - p_k}{p_{k+1} - p_k} = (n+1)p - k$, $\widehat{x}_p := (1-\lambda)X_{(k)} + \lambda X_{(k+1)}$.
(iii) *Q–Q Use*: For unequal samples, evaluate both interpolated quantile functions on a common grid $p \in (0,1)$.

**Definition 2.2.5.** *Histogram, Density Display, and Stem-and-Leaf Plot*
Let $x_1, \dots, x_n \in \mathbb{R}$ be data. For disjoint intervals $B_1, \dots, B_m$ with widths $w_j > 0$.
Set $N_j := \#\{i \in \{1, \dots, n\} : x_i \in B_j\}$. Let $K$ be a kernel, $h > 0$, and $b > 0$ be a chosen scale.

(i) *Histogram*: $H^c_j := N_j$, $H^p_j := \frac{N_j}{n}$, or $H^d_j := \frac{N_j}{n w_j}$ on $B_j$.
(ii) *Density Display*: $\widehat{f}_h : \mathbb{R} \to [0, \infty)$, $\widehat{f}_h(x) := (nh)^{-1} \sum_{i=1}^{n} K(\frac{x - x_i}{h})$.
(iii) *Stem-and-Leaf Plot*: $(s_i, \ell_i)$ satisfies $x_i = bs_i + \ell_i$, $s_i \in \mathbb{Z}$, $\ell_i \in [0, b)$.
Entries with common $s_i$ are printed together in increasing $s_i$.

*Structural Commentary*: For density-scaled heights $H^d_j = \frac{N_j}{n w_j}$, $\sum_j H^d_j w_j = 1$ when the bins cover the sample.
*Key Insight*: Smaller $w_j$ or $h$ retains more local variation; larger $w_j$ or $h$ increases smoothing.




Figure 247: Shape displays: narrow bins at left; small $h$ (blue) is rougher than large $h$ (green).

**Remark 2.2.6.** *What Shape Displays Preserve*

(i) *ECDF*: Preserves ranks and supports direct quantile reading.
(ii) *Histogram*: Preserves bin counts $(N_j)$, not exact values.
(iii) *Kernel Estimate*: Preserves smoothed local mass; bandwidth $h$ controls resolution.
(iv) *Stem–Leaf*: Preserves coarse shape and recoverable numerical values.

## 2.3 Location, Scale, and Robust Summaries

Numerical summaries compress a batch more aggressively than a plot. They should therefore be chosen with attention to the feature being summarized and to the influence of extreme observations.

**Definition 2.3.1.** *Location Summaries for a Batch*
Let $x_1, \dots, x_n \in \mathbb{R}$ have order statistics $x_{(1)} \le \cdots \le x_{(n)}$.
When a trim fraction is used, let $0 \le \alpha < \frac{1}{2}$ and $m_\alpha := \lfloor n\alpha \rfloor$.

(i) *Arithmetic Mean*: $\overline{x} := n^{-1} \sum_{i=1}^{n} x_i$.

(ii) *Median*: $\widetilde{x} := x_{\left(\frac{n+1}{2}\right)}$ if $n$ is odd, and $\widetilde{x} := \frac{x_{\left(\frac{n}{2}\right)} + x_{\left(\frac{n}{2}+1\right)}}{2}$ if $n$ is even.

(iii) $100\alpha\%$ *Trimmed Mean*: $\overline{x}_\alpha := (n - 2m_\alpha)^{-1} \sum_{i=m_\alpha+1}^{n-m_\alpha} x_{(i)}$.

*Structural Commentary*: Under $x_i \mapsto x_i + \Delta$, $\overline{x} \mapsto \overline{x} + \frac{\Delta}{n}$; $\widetilde{x}$ and $\overline{x}_\alpha$ depend only on central order statistics.
*Key Insight*: Extreme magnitudes affect $\overline{x}$ directly, but affect $\widetilde{x}$ and $\overline{x}_\alpha$ mainly through rank changes.

**Remark 2.3.2.** *Outliers and Center*
Replacing one $x_i$ by $M$ sends $\overline{x}$ to $\overline{x} + \frac{M - x_i}{n}$, hence unbounded as $M \to \pm\infty$.
Once an observation is extreme, moving it farther outward leaves $\widetilde{x}$ unchanged and leaves $\overline{x}_\alpha$ unchanged if it remains trimmed. This is descriptive robustness, not an independence or model.

**Remark 2.3.3.** *Compare More Than One Center*
Let $\mathcal{C} := \{\overline{x}, \widetilde{x}, \overline{x}_\alpha, \widehat{\mu}_\Psi\}$. For symmetric light-tailed data, range$(\mathcal{C})$ is typically small; a large range signals skewness, heavy tails, outliers, or mixture structure.

**Definition 2.3.4.** *Dispersion Summaries*
Let $x_1, \dots, x_n \in \mathbb{R}$ have mean $\overline{x}$, median $\widetilde{x}$, and selected sample quartiles $Q_1, Q_3$.

(i) *Sample Variance*: $s^2 := (n-1)^{-1} \sum_{i=1}^{n} (x_i - \overline{x})^2$.
(ii) *Sample Standard Deviation*: $s := \sqrt{s^2}$.
(iii) *Interquartile Range*: IQR $:= Q_3 - Q_1$.
(iv) *Median Absolute Deviation*: MAD $:=$ median$\{|x_i - \widetilde{x}| : 1 \le i \le n\}$.

*Structural Commentary*: $s^2$ is residual-based; IQR and MAD are order-based.
*Key Insight*: A single extreme value can dominate $s$, but not IQR or MAD unless it changes central ranks.

**Remark 2.3.5.** *Normal Scale Calibrations*
For approximately normal data, $\frac{\text{IQR}}{1.35}$ and $\frac{\text{MAD}}{0.675}$ are common robust scale estimates for the normal standard deviation. The constants calibrate the population normal IQR and median absolute deviation to $\sigma$.

## 2.4 Boxplots, Scatterplots, and Relationship Summaries

Boxplots and scatterplots are compact exploratory displays. A boxplot summarizes one batch or compares several batches; a scatterplot displays relationships between paired variables.

**Method 2.4.1.** *Boxplot Construction*
A standard boxplot is built from the median, lower quartile $Q_1$, upper quartile $Q_3$, and $\mathrm{IQR} = Q_3 - Q_1$:

(i) Draw a box from $Q_1$ to $Q_3$ and mark the median inside the box.
(ii) Draw whiskers to the most extreme observations lying within $1.5\,\mathrm{IQR}$ of the nearer quartile.
(iii) Mark observations beyond the whiskers individually as possible outliers.




Figure 248: Boxplot: box $Q_1$ to $Q_3$, line median, dots possible outliers.

**Remark 2.4.2.** *Comparing Batches with Boxplots*

(i) *Location*: Differences in medians $M_g$ indicate group shifts.
(ii) *Spread*: Differences in $\mathrm{IQR}_g = Q_{3,g} - Q_{1,g}$ indicate central-spread changes.
(iii) *Tails*: Unequal whiskers or one-sided outliers indicate skewness or tail imbalance.

**Definition 2.4.3.** *Scatterplot, Pearson Correlation, and Rank Correlation*
Let $(x_i, y_i) \in \mathbb{R}^2$ for $1 \le i \le n$, set $\overline{x} := n^{-1} \sum_{i=1}^{n} x_i$ and $\overline{y} := n^{-1} \sum_{i=1}^{n} y_i$.
Let $R_i, S_i$ be chosen ranks of $x_i, y_i$, using average ranks for ties.
Set $\overline{R} := n^{-1} \sum_{i=1}^{n} R_i$ and $\overline{S} := n^{-1} \sum_{i=1}^{n} S_i$. Set $a_i := x_i - \overline{x}$, $b_i := y_i - \overline{y}$, $u_i := R_i - \overline{R}$, and $v_i := S_i - \overline{S}$.
Let $S_{xx} := \sum_i a_i^2$, $S_{yy} := \sum_i b_i^2$, $S_{xy} := \sum_i a_i b_i$, $S_{RR} := \sum_i u_i^2$, $S_{SS} := \sum_i v_i^2$, and $S_{RS} := \sum_i u_i v_i$.

(i) *Scatterplot*: $P := \{(x_i, y_i) : 1 \le i \le n\} \subset \mathbb{R}^2$, displayed in the $xy$-plane.
(ii) *Pearson Correlation Coefficient*: if $S_{xx} S_{yy} > 0$, $r := \frac{S_{xy}}{\sqrt{S_{xx} S_{yy}}}$.
(iii) *Rank Correlation Coefficient*: if $S_{RR} S_{SS} > 0$, $r_{\mathrm{rank}} := \frac{S_{RS}}{\sqrt{S_{RR} S_{SS}}}$.

*Structural Commentary*: $r$ is affine-scale invariant; $r_{\mathrm{rank}}$ is monotone-transform invariant.
*Key Insight*: $r$ measures linear association; $r_{\mathrm{rank}}$ measures rank association.

**Method 2.4.4.** *Scatterplot Diagnostics*
For paired data $P = \{(x_i, y_i) : 1 \le i \le n\}$:

(i) *Raw Plot*: Inspect $P$ for outliers, curvature, clusters, and nonconstant spread.
(ii) *Transform*: If scale obscures structure, compare $(g(x_i), h(y_i))$ for domain-meaningful $g, h$ such as log.
(iii) *Correlation*: Use $r$ only when the displayed relation is approximately linear.
(iv) *Multiple Variables*: For $z_i \in \mathbb{R}^d$ with $d > 2$, inspect all pairwise plots before relying on pairwise summaries.

**Example 2.4.5.** *Changing Scale Can Reveal Structure*
If $x_i > 0$ and $\max_i x_i / \min_i x_i \gg 1$, the raw plot $P = \{(x_i, y_i)\}$ may compress variation. The transformed plot $P_{\log} := \{(\log x_i, y_i)\}$ can reveal monotone or linear structure, and generally $r(P_{\log}) \neq r(P)$ because Pearson correlation depends on the chosen coordinates.




Figure 249: Scatterplots: raw $x$ compresses points; $\log x$ reveals a linear trend.

**Remark 2.4.6.** *Exploration Before Formal Inference*
Exploratory displays choose the inferential target: transformations $g(X)$, influential subsets, comparable spreads $\mathrm{IQR}_g$, and data-quality checks should be settled before intervals, tests, or models are trusted.

# 3 Categorical Data Analysis

## 3.1 Fisher's Exact Test and Fixed Margins

Categorical data record counts in classes rather than measurements on a continuous scale. In a two-way table, the rows and columns encode two classifications, and the cells record how many units fall in each cross-classification. The probability model is therefore built from multinomial and hypergeometric count laws rather than from normal sampling pivots.

**Definition 3.1.1.** *Two-Way Contingency Table*
Let $I, J \in \mathbb{N}$ and $n_{ij} \in \mathbb{N}_0$ for $1 \le i \le I$, $1 \le j \le J$. Set $n_{i\cdot} := \sum_{j=1}^{J} n_{ij}$, $n_{\cdot j} := \sum_{i=1}^{I} n_{ij}$, and $n_{\cdot\cdot} := \sum_{i=1}^{I}\sum_{j=1}^{J} n_{ij}$.
For corresponding random counts, write $N_{ij}$ and use capitalized totals.

(i) *Two-way contingency table*: $(n_{ij})_{1\le i\le I,\ 1\le j\le J}$, indexed by row and column classifications.
(ii) *Margins*: $\{n_{i\cdot} : 1 \le i \le I\}$, $\{n_{\cdot j} : 1 \le j \le J\}$, and $n_{\cdot\cdot}$.

*Structural Commentary*: Fixed-margin exact tests condition on the margins of random table $N = (N_{ij})$.

**Method 3.1.2.** *Fisher's Exact Test for a $2 \times 2$ Table*
Let a $2 \times 2$ table have fixed margins

| | 1 | 2 | Total |
|---|---|---|---|
| 1 | $N_{11}$ | $N_{12}$ | $n_{1\cdot}$ |
| 2 | $N_{21}$ | $N_{22}$ | $n_{2\cdot}$ |
| Total | $n_{\cdot 1}$ | $n_{\cdot 2}$ | $n_{\cdot\cdot}$ |

Let $R_1, R_2$ be fixed disjoint row sets with $|R_i| = n_{i\cdot}$.
Let $C$ be the random column-1 set with $|C| = n_{\cdot 1}$, and set $\Omega_0 := \{A \subseteq R_1 \cup R_2 : |A| = n_{\cdot 1}\}$.

(i) $H_0 : \mathbb{P}_0(C = A) = \binom{n_{\cdot\cdot}}{n_{\cdot 1}}^{-1}$ $\forall A \in \Omega_0$.
(ii) $N_{11} = |C \cap R_1|$, $N_{12} = n_{1\cdot} - N_{11}$, $N_{21} = n_{\cdot 1} - N_{11}$, and $N_{22} = n_{2\cdot} - n_{\cdot 1} + N_{11}$.
(iii) $\mathcal{S} := \{x \in \mathbb{Z} : 0 \le x \le n_{1\cdot},\ 0 \le n_{\cdot 1} - x \le n_{2\cdot}\}$ and $p_0(x) := \mathbb{P}_0(N_{11} = x) = \frac{\binom{n_{1\cdot}}{x}\binom{n_{2\cdot}}{n_{\cdot 1}-x}}{\binom{n_{\cdot\cdot}}{n_{\cdot 1}}}$, $x \in \mathcal{S}$.
(iv) $p_+(x_{\mathrm{obs}}) := \sum_{x\in\mathcal{S}:x\ge x_{\mathrm{obs}}} p_0(x)$, $p_-(x_{\mathrm{obs}}) := \sum_{x\in\mathcal{S}:x\le x_{\mathrm{obs}}} p_0(x)$, and $p_2(x_{\mathrm{obs}}) := \sum_{x\in\mathcal{S}:p_0(x)\le p_0(x_{\mathrm{obs}})} p_0(x)$.
Reject when the selected value is $\le \alpha$.

**Example 3.1.3.** *Promotion Decisions and Fixed Margins*
In randomised label experiment with 24 male-labeled,24 female-labeled files, the observed table is

| | Male | Female | Total |
|---|---|---|---|
| Promote | 21 | 14 | 35 |
| Hold file | 3 | 10 | 13 |
| Total | 24 | 24 | 48 |

Under $H_0$ and fixed margins $(35, 13)$, $(24, 24)$, $N_{11}$ satisfies $\mathbb{P}_0(N_{11} = x) = \frac{\binom{35}{x}\binom{13}{24-x}}{\binom{48}{24}}$, $x \in \mathcal{S} := \{11, \dots, 24\}$.
At level 0.05, $R^{(2)}_{0.05} := \{11, 12, 13, 14, 21, 22, 23, 24\}$; since $x_{\mathrm{obs}} = 21 \in R^{(2)}_{0.05}$, reject $H_0$.

**Remark 3.1.4.** *What Is Exact Here*
For $N = (N_{ij})$ and $\mathcal{M} := (\{N_{i\cdot}\}_i, \{N_{\cdot j}\}_j)$, exactness means evaluating $\mathcal{L}_0(N \mid \mathcal{M})$ without asymptotics. Conditioning is design-justified when $\mathcal{M}$ is ancillary to $H_0$; rejection interpretation is design-dependent.

## 3.2 Chi-Square Tests for Contingency Tables

For an $I \times J$ table with observed counts $n_{ij}$ and fitted null counts $E_{ij}$, use $X^2 := \sum_{i=1}^{I} \sum_{j=1}^{J} (n_{ij} - E_{ij})^2 / E_{ij}$; in contingency tables, $E_{ij}$ is determined by the row-column null model.

**Definition 3.2.1.** *Homogeneity of Multinomial Distributions*
Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, and $\Delta_{I-1} := \{\pi \in [0,1]^I : \sum_{i=1}^{I} \pi_i = 1\}$.
For independent columns $j \in [J]$, let $(N_{1j}, \dots, N_{Ij}) \sim \text{Multinomial}(n_{\cdot j}; \pi_{1j}, \dots, \pi_{Ij})$ with observed counts $n_{ij}$. The column multinomial laws are *homogeneous* iff $H_0 : \exists \pi \in \Delta_{I-1}$ with $\pi_{ij} = \pi_i \ \forall (i,j) \in [I] \times [J]$; under $H_0$, $\pi_i$ is the common category probability.

**Proposition 3.2.2.** *Common Probability MLE Under Homogeneity*
*Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, $\Delta_{I-1} := \{\pi \in [0,1]^I : \sum_{i=1}^{I} \pi_i = 1\}$, and $N_{\cdot j} := (N_{1j}, \dots, N_{Ij})$, independently for $j \in [J]$. Under $H_0$, $N_{\cdot j} \sim \text{Multinomial}(n_{\cdot j}; \pi)$ for a common $\pi \in \Delta_{I-1}$.*
*With observed counts $n_{ij}$, set $n_{i\cdot} := \sum_{j=1}^{J} n_{ij}$ and $n_{\cdot\cdot} := \sum_{i=1}^{I} n_{i\cdot} > 0$.*
*Then $\widehat{\pi}_i = \frac{n_{i\cdot}}{n_{\cdot\cdot}}$ and $E_{ij} = n_{\cdot j}\widehat{\pi}_i = \frac{n_{i\cdot} n_{\cdot j}}{n_{\cdot\cdot}}$ for $(i,j) \in [I] \times [J]$.*

*Proof.* Up to factors independent of $\pi$, $L(\pi) \propto \prod_{j=1}^{J} \prod_{i=1}^{I} \pi_i^{n_{ij}} = \prod_{i=1}^{I} \pi_i^{n_{i\cdot}}$, hence $\ell(\pi) = \sum_{i=1}^{I} n_{i\cdot} \log \pi_i$ on $\Delta_{I-1}$.
Stationarity of $\ell(\pi) + \lambda(1 - \sum_{i=1}^{I} \pi_i)$ gives $\frac{n_{i\cdot}}{\pi_i} - \lambda = 0$ when $n_{i\cdot} > 0$ and $\widehat{\pi}_i = 0$ when $n_{i\cdot} = 0$,
Normalisation gives $\lambda = n_{\cdot\cdot}$ and $\widehat{\pi}_i = \frac{n_{i\cdot}}{n_{\cdot\cdot}}$. Therefore $E_{ij} = \mathbb{E}_{\widehat{\pi}} N_{ij} = n_{\cdot j}\widehat{\pi}_i$.
*Structural Commentary*: $H_0$ reduces $IJ$ cell probabilities to one shared $\pi \in \Delta_{I-1}$.
*Key Insight*: $\widehat{\pi}$ is the pooled row proportion, and $E_{ij}$ redistributes each column total by $\widehat{\pi}_i$. □

**Definition 3.2.3.** *Homogeneity Fitted Counts*
Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, and let $n_{ij} \in \mathbb{N}_0$ for $(i,j) \in [I] \times [J]$.
Set $n_{i\cdot} := \sum_{j=1}^{J} n_{ij}$, $n_{\cdot j} := \sum_{i=1}^{I} n_{ij}$, $n_{\cdot\cdot} := \sum_{i=1}^{I} \sum_{j=1}^{J} n_{ij} > 0$, and $\widehat{\pi}_i := \frac{n_{i\cdot}}{n_{\cdot\cdot}}$.
The homogeneity *fitted counts* are $E_{ij} := n_{\cdot j}\widehat{\pi}_i = \frac{n_{i\cdot} n_{\cdot j}}{n_{\cdot\cdot}}$ for $(i,j) \in [I] \times [J]$.

**Method 3.2.4.** *Pearson Test of Homogeneity*
Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, and let $n_{ij} \in \mathbb{N}_0$ for $(i,j) \in [I] \times [J]$.
Set $n_{i\cdot} := \sum_{j=1}^{J} n_{ij}$, $n_{\cdot j} := \sum_{i=1}^{I} n_{ij}$, $n_{\cdot\cdot} := \sum_{i=1}^{I} \sum_{j=1}^{J} n_{ij} > 0$, and $E_{ij} := \frac{n_{i\cdot} n_{\cdot j}}{n_{\cdot\cdot}}$. Assume $E_{ij} > 0$ for cells used.

(i) *Statistic*: $X^2 := \sum_{i=1}^{I} \sum_{j=1}^{J} \frac{(n_{ij} - E_{ij})^2}{E_{ij}}$.

(ii) *Calibration*: If $n_{\cdot\cdot}$ large, $\min_{i,j} E_{ij}$ not small, then $X^2 \overset{H_0}{\approx} \chi^2_\nu$; $\nu := J(I-1) - (I-1) = (I-1)(J-1)$.
(iii) *Decision*: Reject $H_0$ at level $\alpha$ when $\mathbb{P}\{\chi^2_{(I-1)(J-1)} \geq X^2_{\text{obs}}\} \leq \alpha$.

**Example 3.2.5.** *Word Counts by Text Group*
Let $W = \{a, an, this, that, with, without\}$, let $G_1, \dots, G_4$ be text groups. Observed word-count table $n_{wj}$ is

| Word | $G_1$ | $G_2$ | $G_3$ | $G_4$ |
|---|---|---|---|---|
| *a* | 147 | 186 | 101 | 83 |
| *an* | 25 | 26 | 11 | 29 |
| *this* | 32 | 39 | 15 | 15 |
| *that* | 94 | 105 | 37 | 22 |
| *with* | 59 | 74 | 28 | 43 |
| *without* | 18 | 10 | 10 | 4 |
| Total | 375 | 440 | 202 | 196 |

For $C_A := \{1, 2, 3\}$, by homogeneity test: $X^2_A = 12.27$, $\nu_A = (6-1)(3-1) = 10$, $p_A = \mathbb{P}\{\chi^2_{10} \geq 12.27\} \approx 0.27$.
Pooling $C_A$ and comparing it with $C_B := \{4\}$ gives $X^2_{A,B} = 32.81$, $\nu_{A,B} = 6 - 1 = 5$, and $p_{A,B} < 0.001$.
$w \in \{an, that\}$ gives the largest Pearson contributions, localising nonhomogeneity by row.

**Definition 3.2.6.** *Independence in a Contingency Table*
Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, and $N = (N_{ij}) \sim \text{Multinomial}(n; (\pi_{ij})_{[I]\times[J]})$ with $n > 0$, observed counts $n_{ij}$.
Set $\pi_{i\cdot} := \sum_{j=1}^{J} \pi_{ij}$, $\pi_{\cdot j} := \sum_{i=1}^{I} \pi_{ij}$, $n_{i\cdot} := \sum_{j=1}^{J} n_{ij}$, and $n_{\cdot j} := \sum_{i=1}^{I} n_{ij}$.
Rows and columns are *independent* iff $H_0 : \pi_{ij} = \pi_{i\cdot}\pi_{\cdot j}\ \forall (i,j) \in [I] \times [J]$; then $\widehat{\pi}_{ij} := \frac{n_{i\cdot}n_{\cdot j}}{n^2}$ and $E_{ij} := \frac{n_{i\cdot}n_{\cdot j}}{n}$.

**Method 3.2.7.** *Pearson Test of Independence*
Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, and let $n_{ij} \in \mathbb{N}_0$ for $(i,j) \in [I] \times [J]$.
Set $n := \sum_{i=1}^{I}\sum_{j=1}^{J} n_{ij} > 0$, $n_{i\cdot} := \sum_{j=1}^{J} n_{ij}$, $n_{\cdot j} := \sum_{i=1}^{I} n_{ij}$, and $E_{ij} := \frac{n_{i\cdot}n_{\cdot j}}{n}$. Assume $E_{ij} > 0$ for cells used.

(i) *Statistic*: $X^2 := \sum_{i=1}^{I}\sum_{j=1}^{J} \frac{(n_{ij}-E_{ij})^2}{E_{ij}}$.

(ii) *Calibration*: If $n$ is large and $\min_{i,j} E_{ij}$ is not small, then $X^2 \overset{H_0}{\approx} \chi^2_\nu$ with $\nu := (I-1)(J-1)$.
(iii) *Decision*: Reject $H_0$ at level $\alpha$ when $\mathbb{P}\{\chi^2_\nu \geq X^2_{\text{obs}}\} \leq \alpha$.
(iv) *Degrees of Freedom*: $\nu = (IJ-1) - ((I-1)+(J-1)) = (I-1)(J-1)$.

**Example 3.2.8.** *Binary Classifications*
Let $n = 1436$, $[n] := \{1, \dots, n\}$, and $(A_k, B_k) \in \{a_1, a_2\} \times \{b_1, b_2\}$ for $k \in [n]$.
Set $n_{ij} := \#\{k \in [n] : A_k = a_i,\ B_k = b_j\}$ and $E_{ij} := \frac{n_{i\cdot}n_{\cdot j}}{n}$. The observed and independence-fitted tables are

$n_{ij}$

| $a_i \backslash b_j$ | $b_1$ | $b_2$ | Total |
|---|---|---|---|
| $a_1$ | 550 | 61 | 611 |
| $a_2$ | 681 | 144 | 825 |
| Total | 1231 | 205 | 1436 |

$E_{ij}$

| $a_i \backslash b_j$ | $b_1$ | $b_2$ |
|---|---|---|
| $a_1$ | 523.8 | 87.2 |
| $a_2$ | 707.2 | 117.8 |

Under $H_0 : A \perp B$, $X^2 = \sum_{i=1}^{2}\sum_{j=1}^{2} \frac{(n_{ij}-E_{ij})^2}{E_{ij}} = 16.01$, $\nu = 1$, and $p < 0.001$; reject $H_0$.
Thus $A \not\perp B$; absent randomised $A$, inference is associative, not causal.

**Remark 3.2.9.** *Homogeneity Versus Independence*
Both tests use $E_{ij} = \frac{n_{i\cdot}n_{\cdot j}}{n_{\cdot\cdot}}$, $X^2 = \sum_{i=1}^{I}\sum_{j=1}^{J} \frac{(n_{ij}-E_{ij})^2}{E_{ij}}$, and $\nu = (I-1)(J-1)$.

(i) *Homogeneity*: $N_{\cdot j} \sim \text{Multinomial}(n_{\cdot j}; \pi_{\cdot j})$ independently for $j \in [J]$, with fixed $n_{\cdot j}$, $H_0 : \pi_{\cdot 1} = \cdots = \pi_{\cdot J}$.
(ii) *Independence*: $N \sim \text{Multinomial}(n_{\cdot\cdot}; (\pi_{ij})_{[I]\times[J]})$ with $H_0 : \pi_{ij} = \pi_{i\cdot}\pi_{\cdot j}\ \forall (i,j) \in [I] \times [J]$.

If $\pi_{\cdot j} > 0$, then independence iff $\pi_{i|j} := \frac{\pi_{ij}}{\pi_{\cdot j}} = \pi_{i\cdot}\ \forall (i,j) \in [I] \times [J]$.

**Method 3.2.10.** *Likelihood-Ratio Version*
Let $I, J \in \mathbb{N}$, $[K] := \{1, \dots, K\}$, and let $n_{ij} \in \mathbb{N}_0$ with null fitted counts $E_{ij}$ for $(i,j) \in [I] \times [J]$.
Set $n_{\cdot\cdot} := \sum_{i=1}^{I}\sum_{j=1}^{J} n_{ij} > 0$ and $\nu := (I-1)(J-1)$.
Let $H_0$ be the homogeneity or independence null producing $E_{ij}$, and assume $E_{ij} > 0$ whenever $n_{ij} > 0$.

(i) *Statistic*: $G^2 := 2\sum_{i=1}^{I}\sum_{j=1}^{J} n_{ij} \log\left(\frac{n_{ij}}{E_{ij}}\right)$.
(ii) *Zero Counts*: $n_{ij} = 0$ contributes 0 to $G^2$.
(iii) *Calibration*: Under the large-sample likelihood-ratio conditions of Theorem 3.1.8, $G^2 \overset{H_0}{\approx} \chi^2_\nu$.
(iv) *Decision*: Reject $H_0$ at level $\alpha$ when $\mathbb{P}\{\chi^2_\nu \geq G^2_{\text{obs}}\} \leq \alpha$.

## 3.3 Matched Pairs and McNemar's Test

Matched categorical data require a different table from unmatched samples. The pair, not the individual measurement, is the independent sampling unit.

**Definition 3.3.1.** *Matched Binary Pair Table*
Let $(Y_k^{(1)}, Y_k^{(2)}) \in \{0,1\}^2$ for $k \in [n]$, and set $R_k := Y_k^{(1)} + 1$, $S_k := Y_k^{(2)} + 1$.
For $(r,s) \in \{1,2\}^2$, set $n_{rs} := \#\{k \in [n] : (R_k, S_k) = (r,s)\}$, so $n = \sum_{r=1}^{2}\sum_{s=1}^{2} n_{rs}$.

(i) *Matched Binary Pair Table*: $N := (n_{rs})_{(r,s)\in\{1,2\}^2}$.
(ii) *Concordant Cells*: $C := \{(r,s) \in \{1,2\}^2 : r = s\} = \{(1,1),(2,2)\}$; counts $(n_{rs})_{(r,s)\in C} = (n_{11}, n_{22})$.

(iii) *Discordant Cells*: $D := \{(r,s) \in \{1,2\}^2 : r \neq s\} = \{(1,2),(2,1)\}$; counts $(n_{rs})_{(r,s)\in D} = (n_{12}, n_{21})$.

*Key Insight*: $(n_{12}, n_{21})$ determines marginal change; $(n_{11}, n_{22})$ records agreement.

**Definition 3.3.2.** *Marginal Homogeneity for Matched Pairs*

Let $\pi = (\pi_{rs})_{(r,s)\in\{1,2\}^2} \in [0,1]^4$ satisfy $\sum_{r=1}^{2}\sum_{s=1}^{2} \pi_{rs} = 1$. Set $\pi_{r\cdot} := \sum_{s=1}^{2} \pi_{rs}$ and $\pi_{\cdot s} := \sum_{r=1}^{2} \pi_{rs}$.

The matched-pair law is *marginally homogeneous* iff $H_0 : (\pi_{1\cdot}, \pi_{2\cdot}) = (\pi_{\cdot 1}, \pi_{\cdot 2})$, equivalently $\pi_{12} = \pi_{21}$.

*Key Insight*: $H_0$ constrains only the discordant probabilities $\pi_{12}, \pi_{21}$.

**Method 3.3.3.** *McNemar's Test*

Let $n_{rs} \in \mathbb{N}_0$ and cell probabilities $\pi_{rs}$ for $(r,s) \in \{1,2\}^2$, with $n := \sum_{r=1}^{2}\sum_{s=1}^{2} n_{rs} > 0$. Set $d := n_{12} + n_{21}$.

(i) *Null*: $H_0 : \pi_{12} = \pi_{21}$.

(ii) *MLE Under $H_0$*: $\widehat{\pi}_{11} = \frac{n_{11}}{n}$, $\widehat{\pi}_{22} = \frac{n_{22}}{n}$, and $\widehat{\pi}_{12} = \widehat{\pi}_{21} = \frac{d}{2n}$.

(iii) *Statistic*: If $d > 0$, $X^2 := \frac{(n_{12}-n_{21})^2}{d}$.

(iv) *Calibration*: If $d$ is large, then $X^2 \overset{H_0}{\approx} \chi^2_1$.

(v) *Decision*: Reject $H_0$ at level $\alpha$ when $\mathbb{P}\{\chi^2_1 \geq X^2_{\text{obs}}\} \leq \alpha$.

**Example 3.3.4.** *Matched Case-Control Pairs*

In a matched case-control study, 85 cases were each paired with one control, and a binary exposure was recorded for both members of each pair.

| | Control | |
|---|---|---|
| Case | Unexposed | Exposed |
| Unexposed | 37 | 7 |
| Exposed | 15 | 26 |

An unmatched chi-square test treats cases and controls as independent samples.

McNemar's test uses only discordant pairs; here $X^2 = (7-15)^2/(7+15) = 2.91$ and $p \approx 0.09$, so accounting for the matching weakens the evidence of an exposure difference.

**Example 3.3.5.** *Matched Case-Crossover Exposure*

In a matched case-crossover study, each subject is observed during an event window and a comparable control window, with a binary exposure recorded in both windows.

| Event window | Control window | | |
|---|---|---|---|
| | Exposed | Unexposed | Total |
| Exposed | 13 | 157 | 170 |
| Unexposed | 24 | 505 | 529 |
| Total | 37 | 662 | 699 |

The discordant counts are 157 and 24, so $X^2 = (157-24)^2/(157+24) = 97.7$, giving strong evidence that exposure prevalence differs between the two windows. This does not by itself establish causation, since time-varying covariates may affect both exposure and event risk.

**Remark 3.3.6.** *Matched Does Not Mean Independent*

McNemar's null hypothesis is marginal homogeneity, not independence of paired responses. Thus the analysis must preserve the matching structure rather than treat the two margins as independent samples.

## 3.4 Odds Ratios and Case-Control Sampling

Odds ratios summarize association in a binary exposure–binary outcome table. They are especially useful because the same cross-product ratio can be estimated under several sampling designs, including retrospective case-control designs where risks themselves cannot be estimated.

**Definition 3.4.1.** *Odds and Odds Ratio*

Let $A$ satisfy $0 < \mathbb{P}(A) < 1$.

Let $X$ and $D$ be binary exposure and disease events with $0 < \mathbb{P}(X) < 1$ and $0 < \mathbb{P}(D \mid X), \mathbb{P}(D \mid \overline{X}) < 1$.

(i) *Odds*: $\text{odds}(A) := \frac{\mathbb{P}(A)}{1-\mathbb{P}(A)}$.

(ii) *Odds ratio*: $\Delta := \frac{\text{odds}(D|X)}{\text{odds}(D|\overline{X})}$.

*Structural Commentary*: If $o = \text{odds}(A)$, then $\mathbb{P}(A) = \frac{o}{1+o}$; $p \mapsto p/(1-p)$ is a bijection $(0,1) \to (0,\infty)$.

**Proposition 3.4.2.** *Cross-Product Formula*
*Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $X, D \in \mathcal{F}$, and set $X_0 := \overline{X}$, $X_1 := X$, $D_0 := \overline{D}$, $D_1 := D$.*
*Let $\pi_{ij} := \mathbb{P}(X_i \cap D_j) > 0$ for $i, j \in \{0,1\}$, with margins $\pi_{i\cdot} := \pi_{i0} + \pi_{i1}$ and $\pi_{\cdot j} := \pi_{0j} + \pi_{1j}$.*

| | $\overline{D}$ | $D$ | |
|---|---|---|---|
| $\overline{X}$ | $\pi_{00}$ | $\pi_{01}$ | $\pi_{0\cdot}$ |
| $X$ | $\pi_{10}$ | $\pi_{11}$ | $\pi_{1\cdot}$ |
| | $\pi_{\cdot 0}$ | $\pi_{\cdot 1}$ | 1 |

*With $\Delta := \frac{\operatorname{odds}(D|X)}{\operatorname{odds}(D|\overline{X})}$, the cross-product identity is $\Delta = \frac{\pi_{11}\pi_{00}}{\pi_{01}\pi_{10}}$.*

*Proof.* Since $\pi_{1\cdot} = \pi_{10} + \pi_{11}$ and $\pi_{0\cdot} = \pi_{00} + \pi_{01}$, $\mathbb{P}(D \mid X) = \frac{\pi_{11}}{\pi_{1\cdot}}$ and $1 - \mathbb{P}(D \mid X) = \frac{\pi_{10}}{\pi_{1\cdot}}$, so $\operatorname{odds}(D \mid X) = \frac{\pi_{11}}{\pi_{10}}$. Similarly, $\operatorname{odds}(D \mid \overline{X}) = \frac{\pi_{01}}{\pi_{00}}$. Hence $\Delta = \frac{\frac{\pi_{11}}{\pi_{10}}}{\frac{\pi_{01}}{\pi_{00}}} = \frac{\pi_{11}\pi_{00}}{\pi_{01}\pi_{10}}$.
*Structural Commentary*: $\pi_{1\cdot}, \pi_{0\cdot}$ enter conditional probabilities but cancel after passage to odds.
*Key Insight*: $\Delta$ equals diagonal product $\pi_{11}\pi_{00}$ divided by off-diagonal product $\pi_{01}\pi_{10}$. □

**Method 3.4.3.** *Estimating an Odds Ratio by Design*
Let $N = (n_{ij})_{i,j\in\{0,1\}}$ be a $2 \times 2$ table for $(X_i, D_j)$, where $X_1 := X$, $X_0 := \overline{X}$, $D_1 := D$, and $D_0 := \overline{D}$.
Assume $n_{ij} > 0$, set $n_{i\cdot} := n_{i0} + n_{i1}$, $n_{\cdot j} := n_{0j} + n_{1j}$, $n := \sum_{i=0}^{1}\sum_{j=0}^{1} n_{ij}$.

(i) *Simple Random Sampling*: $\widehat{\pi}_{ij} := \frac{n_{ij}}{n}$ estimates $\pi_{ij}$; hence $\widehat{\Delta} := \frac{n_{00}n_{11}}{n_{01}n_{10}}$ estimates $\Delta$.
(ii) *Prospective Sampling*: $n_{0\cdot}, n_{1\cdot}$ fixed; $\frac{n_{01}}{n_{0\cdot}}$, $\frac{n_{11}}{n_{1\cdot}}$ estimate $\mathbb{P}(D \mid \overline{X})$, $\mathbb{P}(D \mid X)$; $\mathbb{P}(X)$ not identified.
(iii) *Retrospective Case-Control Sampling*: $n_{\cdot 0}, n_{\cdot 1}$ are fixed; disease risks are not identified, but $\widehat{\Delta} = \frac{n_{11}}{n_{01}} / \frac{n_{10}}{n_{00}}$ is estimable from exposure odds.

*Key Insight*: Design-fixed margins cancel in $\widehat{\Delta}$, so odds ratio is estimable even when risks or margins are not.

**Example 3.4.4.** *Unmatched Case-Control Odds Ratio*
Consider an unmatched case-control table with binary exposure:

| | Exposed | Unexposed |
|---|---|---|
| Case | 67 | 34 |
| Control | 43 | 64 |

The estimated odds ratio is $\widehat{\Delta} = \frac{67 \cdot 64}{43 \cdot 34)} = 2.93$, indicating a positive exposure–disease association.

**Method 3.4.5.** *Parametric Bootstrap for an Odds-Ratio Standard Error*
Let $a, b, c, d \in \mathbb{N}$ denote the observed counts in rows $D, \overline{D}$ and columns $X, \overline{X}$: $a := n_{DX}$, $b := n_{D\overline{X}}$, $c := n_{\overline{D}X}$, and $d := n_{\overline{D}\overline{X}}$. Set $m_D := a + b$, $m_{\overline{D}} := c + d$, $\widehat{p}_D := \frac{a}{m_D}$, $\widehat{p}_{\overline{D}} := \frac{c}{m_{\overline{D}}}$, and $\widehat{\Delta} := \frac{ad}{bc}$.

(i) *Bootstrap Law*: For $r \in [R]$, draw independently $A_r^* \sim \text{Binomial}(m_D, \widehat{p}_D)$ and $C_r^* \sim \text{Binomial}(m_{\overline{D}}, \widehat{p}_{\overline{D}})$.
(ii) *Replicate*: If $0 < A_r^* < m_D$ and $0 < C_r^* < m_{\overline{D}}$, set $\widehat{\Delta}_r^* := \frac{A_r^*(m_{\overline{D}} - C_r^*)}{(m_D - A_r^*)C_r^*}$.
(iii) *Standard Error*: With $\overline{\Delta}^* := \frac{1}{R}\sum_{r=1}^{R}\widehat{\Delta}_r^*$, set $\widehat{\text{se}}_*(\widehat{\Delta}) := \left(\frac{1}{R-1}\sum_{r=1}^{R}(\widehat{\Delta}_r^* - \overline{\Delta}^*)^2\right)^{1/2}$.

*Key Insight*: Resample exposure counts conditional on fixed row totals; $\text{sd}\{(\widehat{\Delta}_r^*)_{r\in[R]}\}$ estimates $\text{se}(\widehat{\Delta})$.

**Remark 3.4.6.** *Odds Ratios Are Not Automatically Risks*
The odds ratio $\Delta = \operatorname{odds}(D \mid X)/\operatorname{odds}(D \mid \overline{X})$ differs from risk contrasts such as $\mathbb{P}(D \mid X) - \mathbb{P}(D \mid \overline{X})$.
In retrospective sampling, $n_{\cdot 0}, n_{\cdot 1}$ are design-fixed, absolute risks are not identified without population margins; $\Delta$ remains estimable because those margins cancel in the cross-product ratio.

## 3.5 Higher-Order Tables and Sampling Design

Two-way tables are the simplest categorical-data structure. Higher-order tables introduce several possible notions of dependence: full independence, conditional independence given a third factor, marginal homogeneity, and interactions among factors.

**Definition 3.5.1.** *Two-Way Log-Linear Form*
Let $I, J \in \mathbb{N}$ and let $\pi_{ij} > 0$ be two-way cell probabilities for $1 \le i \le I$, $1 \le j \le J$,
with $\pi_{i\cdot} := \sum_{j=1}^{J}\pi_{ij}$, $\pi_{\cdot j} := \sum_{i=1}^{I}\pi_{ij}$, $\alpha_i := \log \pi_{i\cdot}$, and $\beta_j := \log \pi_{\cdot j}$.

(i) *Independence form*: $\pi_{ij} = \pi_{i\cdot}\pi_{\cdot j}$ $\forall 1 \le i \le I, 1 \le j \le J \iff \log \pi_{ij} = \alpha_i + \beta_j$ $\forall 1 \le i \le I, 1 \le j \le J$.

(ii) *Interaction terms*: $\gamma_{ij} := \log \pi_{ij} - \alpha_i - \beta_j$, giving $\log \pi_{ij} = \alpha_i + \beta_j + \gamma_{ij}$.

*Key Insight*: $\gamma_{ij} = 0 \ \forall 1 \le i \le I,\ 1 \le j \le J \iff \pi_{ij} = \pi_{i\cdot}\pi_{\cdot j} \ \forall 1 \le i \le I,\ 1 \le j \le J$.

**Remark 3.5.2.** *Three-Way Interactions*
Let $\pi_{ijk} > 0$ for $(i, j, k) \in [I] \times [J] \times [K]$. A log-linear model with all two-factor interactions and no three-factor interaction is $\log \pi_{ijk} = \mu + \alpha_i + \beta_j + \eta_k + \delta_{ij} + \varepsilon_{ik} + \gamma_{jk}$.
The absent term $\lambda_{ijk}$ encodes the constraint of no third-order interaction; nested models are compared by deleting interaction terms and using the corresponding loss of degrees of freedom.

**Example 3.5.3.** *Graduate Admissions and Aggregation*
In the Berkeley admissions example, the aggregate admission rate for men was higher than for women.
Within the six listed majors, however, women were not consistently admitted at lower rates.
The apparent aggregate disadvantage was driven by application patterns: women applied more often to majors with lower admission rates. Major was therefore confounded with sex in the observational comparison.

**Principle 3.5.4.** *Choose the Test from the Sampling Mechanism*
For a contingency table $N = (n_{ij})$, let $n_{i\cdot}$ and $n_{\cdot j}$ denote row and column margins.
Select the test from the design-fixed quantities:

(i) *Fixed Margins*: If $(n_{i\cdot})$ and $(n_{\cdot j})$ are fixed in a $2 \times 2$ table, use Fisher's exact test.
(ii) *Independent Samples*: If rows are independent multinomial samples with fixed $(n_{i\cdot})$, test homogeneity of column probabilities.
(iii) *Single Multinomial Sample*: If $N \sim \text{Multinomial}(n, (\pi_{ij}))$, test independence, i.e. $\pi_{ij} = \pi_{i\cdot}\pi_{\cdot j}$.
(iv) *Matched Pairs*: If observations are paired counts $n_{rs}$, test marginal homogeneity using discordant cells.
(v) *Case-Control Sampling*: If $(n_{\cdot 0}, n_{\cdot 1})$ are fixed by disease status, estimate odds ratios. Absolute risks require population margins.

*Key Insight*: The same numeric table can encode different null hypotheses under different sampling laws.

**Remark 3.5.5.** *Use Counts, Not Percentages*
Let $Q(O, E) := \sum_i \frac{(O_i - E_i)^2}{E_i}$. Pearson's chi-square calibration applies to counts: $Q(O/n, E/n) = \frac{1}{n} Q(O, E)$ and $Q(100O/n, 100E/n) = \frac{100}{n} Q(O, E)$. Thus relative frequencies or percentages give a rescaled statistic unless the original count scale is restored.

# 4 Analysis of Variance and Designed Experiments

## 4.1 One-Way Layouts and the $F$ Test

Analysis of variance compares several mean levels by splitting total variation into variation within treatment groups and variation between treatment groups. The method is a multisample continuation of the equal-variance two-sample procedure, with design and randomisation assumptions kept explicit.

**Definition 4.1.1.** *Balanced One-Way Layout*
For $I, J \in \mathbb{N}$, one factor has levels $i = 1, \dots, I$, each with exactly $J$ independent replicates $Y_{ij}$, $j = 1, \dots, J$.
The normal fixed-effects model satisfies $Y_{ij} = \mu + \alpha_i + \varepsilon_{ij}$, $\sum_{i=1}^{I} \alpha_i = 0$, and independent errors $\varepsilon_{ij} \sim N(0, \sigma^2)$.

*Structural Commentary*: $\mathbb{E}Y_{ij} = \mu + \alpha_i$, and $\sum_{i=1}^{I} \alpha_i = 0$ identifies $\mu$ as the average treatment mean.

*Key Insight*: $H_0 : \alpha_i = 0 \ \forall i$ iff all treatment means are equal.

**Definition 4.1.2.** *One-Way Sums of Squares*
For a balanced one-way layout, set $\overline{Y}_{i\cdot} := \frac{1}{J} \sum_{j=1}^{J} Y_{ij}$ and $\overline{Y}_{\cdot\cdot} := \frac{1}{IJ} \sum_{i=1}^{I} \sum_{j=1}^{J} Y_{ij}$.

(i) *Total*: $SS_{\text{tot}} := \sum_{i=1}^{I} \sum_{j=1}^{J} (Y_{ij} - \overline{Y}_{\cdot\cdot})^2$; grand-mean dispersion.

(ii) *Within-treatment*: $SS_W := \sum_{i=1}^{I} \sum_{j=1}^{J} (Y_{ij} - \overline{Y}_{i\cdot})^2$; residual dispersion within treatments.

(iii) *Between-treatment*: $SS_B := J \sum_{i=1}^{I} (\overline{Y}_{i\cdot} - \overline{Y}_{\cdot\cdot})^2$; treatment-mean dispersion scaled by $J$.

*Key Insight*: $SS_{\text{tot}} = SS_W + SS_B$ separates within-treatment noise from between-treatment mean differences.

**Proposition 4.1.3.** *One-Way Sum-of-Squares Decomposition*
*For $I, J \in \mathbb{N}$ and observations $Y_{ij}$, set $\overline{Y}_{i\cdot} := \frac{1}{J}\sum_j Y_{ij}$, $\overline{Y}_{\cdot\cdot} := \frac{1}{IJ}\sum_i\sum_j Y_{ij}$,*
*$SS_{\text{tot}} := \sum_i\sum_j (Y_{ij}-\overline{Y}_{\cdot\cdot})^2$, $SS_W := \sum_i\sum_j (Y_{ij}-\overline{Y}_{i\cdot})^2$, and $SS_B := J\sum_i (\overline{Y}_{i\cdot}-\overline{Y}_{\cdot\cdot})^2$. Then $SS_{\text{tot}} = SS_W + SS_B$.*

*Proof.* Set $r_{ij} := Y_{ij} - \overline{Y}_{i\cdot}$ and $b_i := \overline{Y}_{i\cdot} - \overline{Y}_{\cdot\cdot}$. Then $Y_{ij} - \overline{Y}_{\cdot\cdot} = r_{ij} + b_i$ and $\sum_j r_{ij} = 0\ \forall i$.
Thus $SS_{\text{tot}} = \sum_i\sum_j (r_{ij}+b_i)^2 = \sum_i\sum_j r_{ij}^2 + 2\sum_i b_i \sum_j r_{ij} + J\sum_i b_i^2 = SS_W + SS_B$.
*Structural Commentary*: $Y_{ij} - \overline{Y}_{\cdot\cdot} = r_{ij} + b_i$, where $r_{ij} := Y_{ij} - \overline{Y}_{i\cdot}$ and $b_i := \overline{Y}_{i\cdot} - \overline{Y}_{\cdot\cdot}$.
*Key Insight*: $\sum_j r_{ij} = 0\ \forall i$ annihilates the cross term $2\sum_i b_i \sum_j r_{ij}$. □




Figure 250: Squaring and summing gives $SS_{\text{tot}} = SS_W + SS_B$; large $MS_B/MS_W$ suggests treatment effects.

**Theorem 4.1.4.** *Expected One-Way Sums of Squares*
*For $I, J \in \mathbb{N}$, let $Y_{ij} = \mu + \alpha_i + \varepsilon_{ij}$, $1 \le i \le I$, $1 \le j \le J$, with $\sum_i \alpha_i = 0$, independent errors satisfying*
*$\mathbb{E}\varepsilon_{ij} = 0$, $\mathrm{Var}(\varepsilon_{ij}) = \sigma^2$. Set $\overline{Y}_{i\cdot} := \frac{1}{J}\sum_j Y_{ij}$, $\overline{Y}_{\cdot\cdot} := \frac{1}{IJ}\sum_i\sum_j Y_{ij}$.*
*Then $SS_W := \sum_i\sum_j (Y_{ij}-\overline{Y}_{i\cdot})^2$, $SS_B := J\sum_i (\overline{Y}_{i\cdot}-\overline{Y}_{\cdot\cdot})^2$; $\mathbb{E}(SS_W) = I(J-1)\sigma^2$*
*and $\mathbb{E}(SS_B) = (I-1)\sigma^2 + J\sum_i \alpha_i^2$. Hence $S_p^2 := \frac{SS_W}{I(J-1)}$ satisfies $\mathbb{E}S_p^2 = \sigma^2$.*

*Proof.* For fixed $i$, Lemma 3.2.8 with $\mu_{ij} = \mu + \alpha_i$ gives $\mathbb{E}\sum_j (Y_{ij} - \overline{Y}_{i\cdot})^2 = (J-1)\sigma^2$, hence $\mathbb{E}SS_W = I(J-1)\sigma^2$.
For $\overline{Y}_{i\cdot}$, $\mathbb{E}\overline{Y}_{i\cdot} = \mu + \alpha_i$, $\mathrm{Var}(\overline{Y}_{i\cdot}) = \frac{\sigma^2}{J}$, and $\frac{1}{I}\sum_i (\mu + \alpha_i) = \mu$. By Lemma 3.2.8, $(\overline{Y}_{i\cdot})_{i=1}^I$ gives
$\mathbb{E}\sum_i (\overline{Y}_{i\cdot} - \overline{Y}_{\cdot\cdot})^2 = \sum_i \alpha_i^2 + \frac{I-1}{J}\sigma^2$, so $\mathbb{E}SS_B = J\sum_i \alpha_i^2 + (I-1)\sigma^2$. Thus $\mathbb{E}S_p^2 = \frac{\mathbb{E}SS_W}{I(J-1)} = \sigma^2$.
*Structural Commentary*: $\mathbb{E}SS_W$ depends only on $\sigma^2$; $\mathbb{E}SS_B$ adds $J\sum_i \alpha_i^2$.
*Key Insight*: Under $\alpha_i = 0\ \forall i$, $\mathbb{E}\{\frac{SS_B}{I-1}\} = \mathbb{E}\{\frac{SS_W}{I(J-1)}\} = \sigma^2$. □

**Theorem 4.1.5.** *Normal-Theory One-Way F Test*
*For $I, J \in \mathbb{N}$, let $Y_{ij} = \mu + \alpha_i + \varepsilon_{ij}$, $1 \le i \le I$, $1 \le j \le J$, with $\sum_i \alpha_i = 0$ and independent errors $\varepsilon_{ij} \sim N(0, \sigma^2)$.*
*Set $\overline{Y}_{i\cdot} := \frac{1}{J}\sum_j Y_{ij}$, $\overline{Y}_{\cdot\cdot} := \frac{1}{IJ}\sum_i\sum_j Y_{ij}$, $SS_W := \sum_i\sum_j (Y_{ij}-\overline{Y}_{i\cdot})^2$, $SS_B := J\sum_i (\overline{Y}_{i\cdot}-\overline{Y}_{\cdot\cdot})^2$.*
*$MS_W := \frac{SS_W}{I(J-1)}$, $MS_B := \frac{SS_B}{I-1}$; $\frac{SS_W}{\sigma^2} \sim \chi^2_{I(J-1)}$.*
*If $H_0 : \alpha_i = 0\ \forall i$, then $\frac{SS_B}{\sigma^2} \sim \chi^2_{I-1}$, $SS_B \perp\!\!\!\perp SS_W$, $F := \frac{MS_B}{MS_W} \sim F_{I-1, I(J-1)}$.*

*Proof.* Let $C_m := \mathrm{Id}_m - m^{-1}\mathbf{1}_m\mathbf{1}_m^\top$, so $C_m^2 = C_m = C_m^\top$ and $\mathrm{rank}\, C_m = m - 1$.
For each $i$, $Z_i := \frac{1}{\sigma}(Y_{i1} - \mu - \alpha_i, \ldots, Y_{iJ} - \mu - \alpha_i)^\top \sim N_J(0, \mathrm{Id}_J)$ independently,
$\frac{1}{\sigma^2}\sum_j (Y_{ij} - \overline{Y}_{i\cdot})^2 = Z_i^\top C_J Z_i \sim \chi^2_{J-1}$. Summing over independent treatments gives $\frac{SS_W}{\sigma^2} \sim \chi^2_{I(J-1)}$.
Under $H_0$, $\widetilde{Z} := \sqrt{J}\sigma^{-1}(\overline{Y}_{1\cdot} - \mu, \ldots, \overline{Y}_{I\cdot} - \mu)^\top \sim N_I(0, \mathrm{Id}_I)$, and $\frac{SS_B}{\sigma^2} = \widetilde{Z}^\top C_I \widetilde{Z} \sim \chi^2_{I-1}$.
For each $i$, $C_J Z_i \perp\!\!\!\perp \mathbf{1}_J^\top Z_i$ by orthogonal normal components.
Hence $SS_W \perp\!\!\!\perp (\overline{Y}_{1\cdot}, \ldots, \overline{Y}_{I\cdot})$ and $SS_W \perp\!\!\!\perp SS_B$. Thus $F = \frac{SS_B}{I-1} / \frac{SS_W}{I(J-1)} = \frac{MS_B}{MS_W} \sim F_{I-1, I(J-1)}$.
*Structural Commentary*: $IJ = 1 + I(J-1) + (I-1)$ partitions df into mean, residual, and contrast subspaces.
*Key Insight*: Under $H_0$, $MS_B$ and $MS_W$ are independent $\sigma^2$ estimators, so $\frac{MS_B}{MS_W}$ has an $F$ law. □

**Example 4.1.6.** *Generic Balanced One-Way ANOVA*
Let $I = 7$, $J = 10$, and $Y_{ij} = \mu_i + \varepsilon_{ij}$ with independent errors $\varepsilon_{ij} \sim N(0, \sigma^2)$. Suppose the ANOVA table is

| Component | *df* | *SS* | *MS* | *F* |
|---|---|---|---|---|
| Treatment | 6 | .125 | .021 | 5.66 |
| Error | 63 | .231 | .0037 | |
| Total | 69 | .356 | | |

Then $F = 5.66$ has null law $F_{6,63}$, $p = \mathbb{P}(F_{6,63} \ge 5.66) < .01$, and $H_0 : \mu_1 = \cdots = \mu_7$ is rejected at level .01.
If diagnostics are adequate, between-treatment variation exceeds within-treatment noise.

**Method 4.1.7.** *Unequal Sample Sizes*

For $I \in \mathbb{N}$, treatment $i$ has observations $Y_{ij}$, $1 \le j \le J_i$, with $J_i \in \mathbb{N}$ and $N := \sum_{i=1}^{I} J_i$.

(i) $\overline{Y}_{i\cdot} := \frac{1}{J_i} \sum_{j=1}^{J_i} Y_{ij}$ and $\overline{Y}_{\cdot\cdot} := \frac{1}{N} \sum_{i=1}^{I} \sum_{j=1}^{J_i} Y_{ij}$.

(ii) $SS_W := \sum_{i=1}^{I} \sum_{j=1}^{J_i} (Y_{ij} - \overline{Y}_{i\cdot})^2$, $SS_B := \sum_{i=1}^{I} J_i (\overline{Y}_{i\cdot} - \overline{Y}_{\cdot\cdot})^2$, $SS_{\text{tot}} := \sum_{i=1}^{I} \sum_{j=1}^{J_i} (Y_{ij} - \overline{Y}_{\cdot\cdot})^2$ satisfy $SS_{\text{tot}} = SS_W + SS_B$.

(iii) $\mathrm{df}_W = N - I$ and $\mathrm{df}_B = I - 1$.

(iv) If $Y_{ij} = \mu_i + \varepsilon_{ij}$ with independent $\varepsilon_{ij} \sim N(0, \sigma^2)$ and $H_0 : \mu_1 = \cdots = \mu_I$, then $F := \frac{\frac{SS_B}{I-1}}{\frac{SS_W}{N-I}} \sim F_{I-1,N-I}$.

**Corollary 4.1.8.** *Two Treatments Give $F = t^2$*

*Let $X_1, \ldots, X_n$ and $Y_1, \ldots, Y_m$ be independent normal samples with common variance, means $\overline{X}, \overline{Y}$, pooled variance $S_p^2 := \frac{SS_W}{n+m-2}$, and $t := \frac{\overline{X} - \overline{Y}}{S_p \sqrt{n^{-1} + m^{-1}}}$. The one-way ANOVA statistic for the two groups satisfies $F = t^2$.*

*Proof.* Let $N := n + m$ and $\overline{Z} := \frac{n\overline{X} + m\overline{Y}}{N}$. Then $SS_B = n(\overline{X} - \overline{Z})^2 + m(\overline{Y} - \overline{Z})^2 = \frac{nm}{N}(\overline{X} - \overline{Y})^2$.
Since $df_B = 1$ and $MS_W = \frac{SS_W}{N-2} = S_p^2$, $F = \frac{SS_B}{MS_W} = \frac{(\overline{X} - \overline{Y})^2}{S_p^2 (n^{-1} + m^{-1})} = t^2$.
*Structural Commentary*: Two groups leave one between-group contrast.
*Key Insight*: ANOVA squares the pooled $t$ contrast. □

The same one-way decomposition also supports a random-effects interpretation, where groups are sampled from a population of possible group levels rather than fixed in advance.

**Definition 4.1.9.** *Balanced One-Way Random Effects Model*

A *balanced one-way random-effects model* has $I$ groups and common replication $J$, with $Y_{ij} = \mu + A_i + \varepsilon_{ij}$ for $1 \le i \le I$, $1 \le j \le J$, independent $A_i$ satisfying $\mathbb{E}A_i = 0$ and $\mathrm{Var}(A_i) = \sigma_A^2$, and independent errors $\varepsilon_{ij}$ independent of $(A_i)$ with $\mathbb{E}\varepsilon_{ij} = 0$ and $\mathrm{Var}(\varepsilon_{ij}) = \sigma_\varepsilon^2$.
*Structural Commentary*: $\mathrm{Var}(Y_{ij}) = \sigma_A^2 + \sigma_\varepsilon^2$, and $\mathrm{Cov}(Y_{ij}, Y_{ij'}) = \sigma_A^2$ for $j \ne j'$ within the same group.

**Proposition 4.1.10.** *Random-Effects Expected Mean Squares*

*Assume $Y_{ij} = \mu + A_i + \varepsilon_{ij}$ for $1 \le i \le I$, $1 \le j \le J$, with independent $A_i$ and $\varepsilon_{ij}$, $\mathbb{E}A_i = \mathbb{E}\varepsilon_{ij} = 0$, $\mathrm{Var}(A_i) = \sigma_A^2$, and $\mathrm{Var}(\varepsilon_{ij}) = \sigma_\varepsilon^2$. Then $\mathbb{E}(MS_W) = \sigma_\varepsilon^2$ and $\mathbb{E}(MS_B) = \sigma_\varepsilon^2 + J\sigma_A^2$.*
*The method-of-moments estimators are $\widehat{\sigma}_\varepsilon^2 = MS_W$ and $\widehat{\sigma}_A^2 = \frac{MS_B - MS_W}{J}$, truncated at 0 if required.*

*Proof.* Condition on $(A_i)$. Since $Y_{ij} - \overline{Y}_{i\cdot} = \varepsilon_{ij} - \overline{\varepsilon}_{i\cdot}$, $\mathbb{E}(SS_W \mid A) = I(J-1)\sigma_\varepsilon^2$, hence $\mathbb{E}(MS_W) = \sigma_\varepsilon^2$.
Also $\overline{Y}_{i\cdot} = \mu + A_i + \overline{\varepsilon}_{i\cdot}$ has variance $\sigma_A^2 + \frac{\sigma_\varepsilon^2}{J}$.
Hence the one-way between mean square satisfies $\mathbb{E}(MS_B) = J\,\mathrm{Var}(\overline{Y}_{i\cdot}) = \sigma_\varepsilon^2 + J\sigma_A^2$.
Solving these two moment equations gives the estimators.
*Structural Commentary*: Within estimates error; between adds group heterogeneity.
*Key Insight*: The excess $MS_B - MS_W$ estimates $J\sigma_A^2$. □

**Example 4.1.11.** *Dye Strength Batch Variation*

Suppose $I = 6$ batches each have $J = 6$ subsamples, with $\overline{y}_{\cdot\cdot} = 93.61$
and batch means $(92.00, 90.50, 91.25, 98.25, 91.17, 98.50)$.
Given $SS_B = 415.47$ and $SS_W = 134.08$, the mean squares are $MS_B = 83.09$ and $MS_W = 4.47$.
The moment estimates are $\widehat{\sigma}_\varepsilon^2 = MS_W = 4.47$ and $\widehat{\sigma}_A^2 = \frac{MS_B - MS_W}{6} = 13.10$.
Batch variation dominates measurement-error variation.

**Remark 4.1.12.** *Assumptions and Robustness*

For $Y_{ij} = \mu_i + \varepsilon_{ij}$, the one-way $F$ calibration assumes independent $\varepsilon_{ij} \sim N(0, \sigma^2)$. Balance softens variance heterogeneity; clustered, paired, blocked, or repeated data require dependence modeling.

## 4.2 Multiple Comparisons After ANOVA

The one-way $F$ test assesses $H_0 : \mu_1 = \cdots = \mu_I$; after rejection, inference targets contrasts $\mu_a - \mu_b$ with error control over the chosen comparison family.

**Definition 4.2.1.** *Studentized Range Quantile*

Let $Q_{I,\nu}$ denote a random variable with the studentized range distribution for $I$ treatment means and $\nu$ error degrees of freedom. The *studentized range quantile* is $q_{I,\nu}(\alpha) := \inf\{q \in \mathbb{R} : \mathbb{P}(Q_{I,\nu} > q) \le \alpha\}$.
*Structural Commentary*: Balanced Tukey comparisons use pooled scale $S_p$.

**Method 4.2.2.** *Tukey Pairwise Comparisons*
In a balanced normal one-way layout, let $\mathcal{P} := \{(a,b) : 1 \le a < b \le I\}$, $\nu := I(J-1)$, and $h_\alpha := q_{I,\nu}(\alpha) S_p/\sqrt{J}$.

(i) For $(a,b) \in \mathcal{P}$, the simultaneous $100(1-\alpha)\%$ confidence interval for $\mu_a - \mu_b$ is $(\overline{Y}_{a\cdot} - \overline{Y}_{b\cdot}) \pm h_\alpha$.
(ii) Reject $H_{0,ab} : \mu_a = \mu_b$ iff $\left|\overline{Y}_{a\cdot} - \overline{Y}_{b\cdot}\right| > h_\alpha$.

Simultaneity over $\mathcal{P}$ uses the studentized range cutoff.

**Example 4.2.3.** *Generic Tukey Pairwise Calculation*
Consider a balanced one-way layout with $I = 7$, $J = 10$, $S_p = .06$, and ordered sample means $(\overline{Y}_{1\cdot}, \overline{Y}_{3\cdot}, \overline{Y}_{7\cdot}, \overline{Y}_{2\cdot}, \overline{Y}_{5\cdot}, \overline{Y}_{6\cdot}, \overline{Y}_{4\cdot}) = (4.062, 4.003, 3.998, 3.997, 3.957, 3.955, 3.920)$.
With $q_{7,60}(.05) = 4.31$, the Tukey half-width is $h_{.05} = 4.31(.06)/\sqrt{10} \approx .082$.
Rejected pairs at simultaneous level .05 are $(1,4), (1,5), (1,6), (3,4)$; remaining pairwise nulls not rejected.

**Method 4.2.4.** *Bonferroni Pairwise Comparisons*
Let $\mathcal{C} \subseteq \{(a,b) : 1 \le a < b \le I\}$ be the planned pairwise family with $|\mathcal{C}| = k$.
Set $\nu := N - I$ and $h_{ab} := t_{\nu,1-\alpha/(2k)} S_p \sqrt{1/J_a + 1/J_b}$.

(i) For $(a,b) \in \mathcal{C}$, the Bonferroni confidence interval for $\mu_a - \mu_b$ is $(\overline{Y}_{a\cdot} - \overline{Y}_{b\cdot}) \pm h_{ab}$.
(ii) Reject $H_{0,ab} : \mu_a = \mu_b$ iff $\left|\overline{Y}_{a\cdot} - \overline{Y}_{b\cdot}\right| > h_{ab}$.

The family satisfies FWER $\le \alpha$ and allows unequal treatment sizes through $J_a, J_b$.

**Example 4.2.5.** *Generic Bonferroni Pairwise Calculation*
For $I = 7$ treatments and $k = \binom{7}{2} = 21$ planned pairwise comparisons at $\alpha = .05$, take $t_{60,1-.05/(2\cdot 21)} \approx 3.16$.
The resulting simultaneous 95% Bonferroni half-width is $h \approx .085$.
Rejected pairwise nulls are $(1,4), (1,5), (1,6)$; all other planned pairwise nulls are not rejected.

**Remark 4.2.6.** *The Global Test Does Not Name the Difference*
Rejecting $H_0 : \mu_1 = \cdots = \mu_I$ establishes only $\exists\, a < b$ with $\mu_a \ne \mu_b$.
It does not identify such pairs or provide simultaneous sign guarantees.
Select Tukey, Bonferroni, or planned-contrast procedures according to the preassigned comparison family.

## 4.3 Rank-Based One-Way ANOVA

When normality or outlier sensitivity is a concern, a one-way comparison can be made after replacing observations by pooled ranks.

**Definition 4.3.1.** *Kruskal–Wallis Ranks*
Let $\mathcal{Y} := \{Y_{ij} : 1 \le i \le I, 1 \le j \le J_i\}$ be an unequal-size one-way sample with $N := \sum_{i=1}^{I} J_i$.
Assign increasing pooled ranks to $\mathcal{Y}$, averaging ranks over ties.

(i) *Pooled rank*: $R_{ij} := \mathrm{rank}_{\mathrm{pool}}(Y_{ij})$.
(ii) *Treatment mean rank*: $\overline{R}_{i\cdot} := \frac{1}{J_i} \sum_{j=1}^{J_i} R_{ij}$.
(iii) *Grand mean rank*: $\overline{R}_{\cdot\cdot} := \frac{N+1}{2}$.

**Method 4.3.2.** *Kruskal–Wallis Test*
Let $SS_B^{(R)} := \sum_{i=1}^{I} J_i(\overline{R}_{i\cdot} - \overline{R}_{\cdot\cdot})^2$ and $K := \frac{12}{N(N+1)} SS_B^{(R)} = \frac{12}{N(N+1)} \sum_{i=1}^{I} J_i \overline{R}_{i\cdot}^2 - 3(N+1)$.

(i) The exact null law enumerates all allocations of pooled ranks to group sizes $J_1, \ldots, J_I$.
(ii) For moderate samples, $K \overset{H_0}{\approx} \chi^2_{I-1}$.
(iii) Reject $H_0 : F_1 = \cdots = F_I$ at level $\alpha$ when $\mathbb{P}\{\chi^2_{I-1} \ge K_{\mathrm{obs}}\} \le \alpha$.

**Example 4.3.3.** *Generic Kruskal–Wallis Rank Calculation*
For $I = 7$ groups, suppose the pooled ranks give $K_{\mathrm{obs}} = 29.51$.
Using $K \overset{H_0}{\approx} \chi^2_6$, the tail probability satisfies $\mathbb{P}\{\chi^2_6 \ge 29.51\} < .005$.
Reject $H_0 : F_1 = \cdots = F_7$; the group response distributions are not all equal.

**Corollary 4.3.4.** *Two Groups Recover Mann–Whitney*
*For $I = 2$, let $T_i := \sum_{j=1}^{J_i} R_{ij}$ and $N = J_1 + J_2$. Then $K = \frac{12}{J_1 J_2 (N+1)} \left(T_1 - \frac{J_1(N+1)}{2}\right)^2$, so the Kruskal–Wallis rejection order is the two-sided Mann–Whitney rank-sum order.*

*Proof.* Let $G := \frac{N+1}{2}$. Since $T_1 + T_2 = NG$, $\overline{R}_{1\cdot} - G = \frac{T_1 - J_1 G}{J_1}$ and $\overline{R}_{2\cdot} - G = -\frac{T_1 - J_1 G}{J_2}$.
Hence $SS_B^{(R)} = (T_1 - J_1 G)^2 (\frac{1}{J_1} + \frac{1}{J_2}) = \frac{N(T_1 - J_1 G)^2}{J_1 J_2}$, and the formula for $K$ follows.
Since $U_1 = T_1 - \frac{J_1(J_1+1)}{2}$ is affine in $T_1$, both tests order evidence by $|T_1 - J_1 G|$.
*Structural Commentary*: Two groups leave one rank contrast.
*Key Insight*: $K$ is a squared centered rank sum. □

**Method 4.3.5.** *Rank-Based Follow-Up Comparisons*
After rejecting global rank null, let $\mathcal{C} \subseteq \{(a,b) : 1 \le a < b \le I\}$ be planned follow-up family with $|\mathcal{C}| = k$.

(i) For $(a,b) \in \mathcal{C}$, test $H_{0,ab} : F_a = F_b$ by a two-sample rank-sum statistic and let $p_{ab}$ be its two-sided $p$-value.
(ii) Reject $H_{0,ab}$ iff $p_{ab} \le \frac{\alpha}{k}$, equivalently $k p_{ab} \le \alpha$.
(iii) The selected family satisfies FWER $\le \alpha$ and is most efficient when $\mathcal{C}$ is fixed before inspection.

**Remark 4.3.6.** *What Rank ANOVA Tests*
The null is $H_0 : F_1 = \cdots = F_I$, not merely $\mu_1 = \cdots = \mu_I$. Under comparable distributional shapes, rejection is read as a location difference; the rank form avoids normality and reduces sensitivity to extremes.

## 4.4 Two-Way Layouts and Interaction

A two-way layout studies two factors at once. It estimates main effects for each factor and checks whether the effect of one factor changes across levels of the other.

**Definition 4.4.1.** *Two-Way Additive Parametrization*
For one observation $Y_{ij}$ at factor levels $i = 1, \dots, I$ and $j = 1, \dots, J$,
set $\overline{Y}_{i\cdot} := J^{-1} \sum_{j=1}^{J} Y_{ij}$, $\overline{Y}_{\cdot j} := I^{-1} \sum_{i=1}^{I} Y_{ij}$, and $\overline{Y}_{\cdot\cdot} := (IJ)^{-1} \sum_{i=1}^{I} \sum_{j=1}^{J} Y_{ij}$.

(i) *Intercept*: $\widehat{\mu} := \overline{Y}_{\cdot\cdot}$.
(ii) *Factor A, B coefficient*: $\widehat{\alpha}_i := \overline{Y}_{i\cdot} - \overline{Y}_{\cdot\cdot}$, $1 \le i \le I$ and $\widehat{\beta}_j := \overline{Y}_{\cdot j} - \overline{Y}_{\cdot\cdot}$, $1 \le j \le J$.
(iii) *Additive fitted value*: $\widehat{Y}_{ij} := \widehat{\mu} + \widehat{\alpha}_i + \widehat{\beta}_j$.
(iv) *Interaction residual*: $\widehat{\delta}_{ij} := Y_{ij} - \overline{Y}_{i\cdot} - \overline{Y}_{\cdot j} + \overline{Y}_{\cdot\cdot}$, so $Y_{ij} = \widehat{\mu} + \widehat{\alpha}_i + \widehat{\beta}_j + \widehat{\delta}_{ij}$.

*Structural Commentary*: Exact fits satisfy zero-sum effects and margins: $\sum_i \widehat{\alpha}_i = \sum_j \widehat{\beta}_j = \sum_i \widehat{\delta}_{ij} = \sum_j \widehat{\delta}_{ij} = 0$.

**Remark 4.4.2.** *Interaction and Parallel Profiles*
Under additivity, $\widehat{Y}_{ij} - \widehat{Y}_{i'j} = \widehat{\alpha}_i - \widehat{\alpha}_{i'}$ is independent of $j$, so factor-$A$ profiles are parallel across $B$. Nonparallel profiles indicate $\delta_{ij} \neq 0$ structure: the contrast between factor levels depends on the conditioning level.




Figure 251: Parallel profiles indicate additivity; changing gaps indicate interaction.

**Definition 4.4.3.** *Balanced Two-Way Normal Model*
A *balanced two-way normal fixed-effects model* observes $Y_{ijk}$ for $1 \le i \le I$, $1 \le j \le J$, $1 \le k \le K$, with common cell size $K > 1$, and assumes $Y_{ijk} = \mu + \alpha_i + \beta_j + \delta_{ij} + \varepsilon_{ijk}$ for mutually independent $\varepsilon_{ijk} \sim N(0, \sigma^2)$,
under $\sum_i \alpha_i = \sum_j \beta_j = 0$, $\sum_i \delta_{ij} = 0 \ \forall j$, and $\sum_j \delta_{ij} = 0 \ \forall i$.
*Key Insight*: $\delta_{ij} = 0 \ \forall (i,j)$ iff the cell means are additive in the two factors.

**Definition 4.4.4.** *Two-Way Sums of Squares*
Let $Y_{ijk}$ be balanced observations for $1 \le i \le I$, $1 \le j \le J$, $1 \le k \le K$.

(i) *Means*: $\overline{Y}_{ij\cdot} := \frac{1}{K} \sum_k Y_{ijk}$, $\overline{Y}_{i\cdot\cdot} := \frac{1}{JK} \sum_j \sum_k Y_{ijk}$, $\overline{Y}_{\cdot j\cdot} := \frac{1}{IK} \sum_i \sum_k Y_{ijk}$, $\overline{Y}_{\cdots} := \frac{1}{IJK} \sum_i \sum_j \sum_k Y_{ijk}$.

(ii) *Between-cell components*: $SS_A := JK\sum_i(\overline{Y}_{i\cdot\cdot} - \overline{Y}_{\cdots})^2$, $SS_B := IK\sum_j(\overline{Y}_{\cdot j\cdot} - \overline{Y}_{\cdots})^2$, and $SS_{AB} := K\sum_i\sum_j(\overline{Y}_{ij\cdot} - \overline{Y}_{i\cdot\cdot} - \overline{Y}_{\cdot j\cdot} + \overline{Y}_{\cdots})^2$.

(iii) *Within-cell and total*: $SS_E := \sum_i\sum_j\sum_k(Y_{ijk} - \overline{Y}_{ij\cdot})^2$, $SS_{\text{tot}} := \sum_i\sum_j\sum_k(Y_{ijk} - \overline{Y}_{\cdots})^2$, and $SS_{\text{tot}} = SS_A + SS_B + SS_{AB} + SS_E$.

*Structural Commentary*: Orthogonal projections split total variation.

**Theorem 4.4.5.** *Expected Two-Way Sums of Squares*
*Assume* $Y_{ijk} = \mu + \alpha_i + \beta_j + \delta_{ij} + \varepsilon_{ijk}$ *with independent* $\varepsilon_{ijk} \sim N(0,\sigma^2)$, *and* $\sum_i \alpha_i = \sum_j \beta_j = 0$, $\sum_i \delta_{ij} = 0\ \forall j$; $\sum_j \delta_{ij} = 0\ \forall i$.

*(i)* $\mathbb{E}(SS_A) = (I-1)\sigma^2 + JK\sum_i \alpha_i^2$.

*(ii)* $\mathbb{E}(SS_B) = (J-1)\sigma^2 + IK\sum_j \beta_j^2$.

*(iii)* $\mathbb{E}(SS_{AB}) = (I-1)(J-1)\sigma^2 + K\sum_i\sum_j \delta_{ij}^2$.

*(iv)* $\mathbb{E}(SS_E) = IJ(K-1)\sigma^2$.

*Proof.* Let $Y = m + \varepsilon \in \mathbb{R}^{IJK}$ with $m_{ijk} = \mu + \alpha_i + \beta_j + \delta_{ij}$ and $\varepsilon \sim N(0, \sigma^2 I_{IJK})$.
For an orthogonal projection $P$, $\mathbb{E}\,\|PY\|^2 = \|Pm\|^2 + \sigma^2 \operatorname{rank} P$.

(i) $P_A m$ has value $\alpha_i$ repeated over $j,k$, so $\|P_A m\|^2 = JK\sum_i \alpha_i^2$ and $\operatorname{rank} P_A = I-1$.

(ii) $P_B m$ has value $\beta_j$ repeated over $i,k$, so $\|P_B m\|^2 = IK\sum_j \beta_j^2$ and $\operatorname{rank} P_B = J-1$.

(iii) $P_{AB} m$ has value $\delta_{ij}$ repeated over $k$, so $\|P_{AB} m\|^2 = K\sum_i\sum_j \delta_{ij}^2$ and $\operatorname{rank} P_{AB} = (I-1)(J-1)$.

(iv) $P_E$ centers within cells; since $m_{ijk}$ is constant in $k$ for fixed $(i,j)$, $P_E m = 0$ and $\operatorname{rank} P_E = IJ(K-1)$.

Applying the projection identity to $P_A, P_B, P_{AB}, P_E$ gives the four expectations.
*Structural Commentary*: Balanced orthogonality separates noise degrees of freedom from effect energy.
*Key Insight*: Each expectation is $\sigma^2$ times rank plus squared signal norm. □

**Theorem 4.4.6.** *Two-Way Normal-Theory F Tests*
*Assume observations* $Y_{ijk}$ *for* $1 \le i \le I$, $1 \le j \le J$, $1 \le k \le K$ *satisfy* $Y_{ijk} = \mu + \alpha_i + \beta_j + \delta_{ij} + \varepsilon_{ijk}$ *with independent* $\varepsilon_{ijk} \sim N(0,\sigma^2)$, $\sum_i \alpha_i = \sum_j \beta_j = 0$, $\sum_i \delta_{ij} = 0\ \forall j$, *and* $\sum_j \delta_{ij} = 0\ \forall i$.
*Let* $d_E := IJ(K-1)$ *and* $MS_E := \frac{SS_E}{d_E}$.

*(i)* $SS_A, SS_B, SS_{AB}, SS_E$ *are independent, and* $\frac{SS_E}{\sigma^2} \sim \chi^2_{d_E}$.

*(ii)* *Under* $H_A : \alpha_i = 0\ \forall i$, $\frac{SS_A}{\sigma^2} \sim \chi^2_{I-1}$; *under* $H_B : \beta_j = 0\ \forall j$, $\frac{SS_B}{\sigma^2} \sim \chi^2_{J-1}$; *under* $H_{AB} : \delta_{ij} = 0\ \forall (i,j)$, $\frac{SS_{AB}}{\sigma^2} \sim \chi^2_{(I-1)(J-1)}$.

*(iii)* $F_A := \frac{\frac{SS_A}{I-1}}{MS_E} \sim F_{I-1,d_E}$ *under* $H_A$, $F_B := \frac{\frac{SS_B}{J-1}}{MS_E} \sim F_{J-1,d_E}$ *under* $H_B$, *and* $F_{AB} := \frac{\frac{SS_{AB}}{(I-1)(J-1)}}{MS_E} \sim F_{(I-1)(J-1),d_E}$ *under* $H_{AB}$.

*Proof.*

(i) Vectorise $Y = m + \varepsilon \in \mathbb{R}^{IJK}$ with $\varepsilon \sim N(0,\sigma^2 I_{IJK})$.
Let $P_A, P_B, P_{AB}, P_E$ be the mutually orthogonal projections onto the factor-$A$, factor-$B$, interaction, and within-cell subspaces, so $SS_R = \|P_R Y\|^2$ for $R \in \{A, B, AB, E\}$.
Orthogonal projections of a spherical normal vector are independent, and $\frac{\|P_R Y\|^2}{\sigma^2} \sim \chi^2_{d_R}(\lambda_R)$ with $d_R = \operatorname{rank} P_R$ and $\lambda_R = \frac{\|P_R m\|^2}{\sigma^2}$. Since $d_E = IJ(K-1)$ and $\lambda_E = 0$, claim (i) follows.

(ii) Here $d_A = I-1$, $d_B = J-1$, and $d_{AB} = (I-1)(J-1)$. Under $H_A, H_B, H_{AB}$ respectively, $\lambda_A, \lambda_B, \lambda_{AB}$ vanish, so the central chi-square laws in claim (ii) follow.

(iii) In each null case the numerator chi-square in claim (ii) is independent of $SS_E$ by claim (i); dividing by its degrees of freedom and by $\frac{SS_E}{d_E} = MS_E$ gives the stated $F$ ratios in claim (iii).

*Structural Commentary*: Balanced normal ANOVA is an orthogonal Gaussian projection decomposition.
*Key Insight*: Null effects remove noncentrality, leaving independent chi-square ratios. □

**Example 4.4.7.** *Generic Transformed Two-Way ANOVA*
Consider balanced two-way study with factor $A$ having 2 levels, factor $B$ having 3 levels, $K = 18$ observations

per cell. After log transformation stabilises cell spread, ANOVA table on transformed scale is

| Source | $df$ | $SS$ | $MS$ | $F$ |
|---|---|---|---|---|
| Factor $A$ | 1 | 2.074 | 2.074 | 5.99 |
| Factor $B$ | 2 | 15.588 | 7.794 | 22.53 |
| Interaction | 2 | .810 | .405 | 1.17 |
| Error | 102 | 35.296 | .346 | |
| Total | 107 | 53.768 | | |

For factor $A$, $F_A = 5.99$ gives $p < .025$, and an estimated log contrast .28 corresponds to the original-scale multiplier $e^{.28} \approx 1.32$ with approximate interval $(1.06, 1.65)$. For interaction, $F_{AB} = 1.17$ gives $p = .31$, so there is no strong evidence that the $A$ contrast depends on the level of $B$.

**Remark 4.4.8.** *Transformations and Diagnostics*
Two-way ANOVA inherits the normal-theory assumptions of the one-way model. Residual plots, normal probability plots, profile plots, and plots of cell spread against cell mean help decide whether a transformation is needed. A transformation can stabilize variance and improve symmetry, but it also changes additive effects on the transformed scale into multiplicative or otherwise nonlinear effects on the original scale.

**Remark 4.4.9.** *Factorial Economy and Incomplete Designs*
Factorial designs use each observation to estimate multiple factor contrasts and interactions.
As the factor set grows, full factorials may be replaced by fractional factorials that omit cells while preserving chosen estimable effects; when blocks cannot contain all treatments, incomplete block designs retain partial blocking without all block-treatment pairs.

## 4.5 Randomised Blocks and Friedman's Test

Randomised block designs compare treatments within relatively homogeneous blocks. Blocking removes major background variation from the treatment comparison, much as pairing removes shared variation.

**Definition 4.5.1.** *Randomised Block Design*
A randomised block design has $J$ blocks, each containing one experimental unit for each of the $I$ treatments; within every block, treatments are assigned by an independent random permutation.
The additive block model is $Y_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}$ for $1 \le i \le I$, $1 \le j \le J$, with $\sum_i \alpha_i = \sum_j \beta_j = 0$.

*Key Insight*: Blocks absorb heterogeneity; $\alpha_i - \alpha_{i'}$ is a within-block treatment contrast.

**Method 4.5.2.** *Randomized Block F Test*
Treat $Y_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}$ as a two-way layout with one observation per treatment-block cell and no interaction.

(i) Define $MS_A := \frac{SS_A}{I-1}$, $MS_B := \frac{SS_B}{J-1}$, and $MS_{AB} := \frac{SS_{AB}}{(I-1)(J-1)}$.
(ii) Under additivity, $\mathbb{E}(MS_A) = \sigma^2 + \frac{J}{I-1}\sum_i \alpha_i^2$, $\mathbb{E}(MS_B) = \sigma^2 + \frac{I}{J-1}\sum_j \beta_j^2$, and $\mathbb{E}(MS_{AB}) = \sigma^2$.
(iii) Test $H_0 : \alpha_1 = \cdots = \alpha_I = 0$ by $F := \frac{MS_A}{MS_{AB}}$.
(iv) Under the normal additive null, $F \sim F_{I-1,(I-1)(J-1)}$.

**Remark 4.5.3.** *Interaction in a Block Design*
If $\delta_{ij} \neq 0$, then $MS_{AB}$ estimates $\sigma^2$ plus treatment-block interaction energy.
The denominator of $F = \frac{MS_A}{MS_{AB}}$ is inflated, so the additive-model treatment test is conservative.

**Example 4.5.4.** *Generic Randomized Block ANOVA*
Suppose $I = 7$ treatments are compared across $J = 10$ blocks, with each block receiving all treatments in randomized order. The randomized-block ANOVA table is

| Source | $df$ | $SS$ | $MS$ | $F$ |
|---|---|---|---|---|
| Treatments | 6 | 53013 | 8835 | 2.85 |
| Blocks | 9 | 103280 | 11476 | 3.71 |
| Interaction | 54 | 167130 | 3095 | |
| Total | 69 | 323422 | | |

$F = 2.85$ on $(6, 54)$ degrees of freedom gives $p < .025$, rejecting equal treatment effects.
Blocking isolates nuisance block heterogeneity.

**Method 4.5.5.** *Friedman's Rank Test for Blocks*
For observations $Y_{ij}$ from treatment $i$ in block $j$, rank $Y_{1j}, \ldots, Y_{Ij}$ within each block.

(i) Let $R_{ij}$ be within-block rank, use average ranks for ties, set $\overline{R}_{i\cdot} := \frac{1}{J}\sum_j R_{ij}$, $\overline{R}_{\cdot\cdot} := \frac{I+1}{2}$.

(ii) Define $Q := \frac{12J}{I(I+1)} \sum_i (\overline{R}_{i\cdot} - \overline{R}_{\cdot\cdot})^2$.

(iii) Under $H_0 : F_1 = \cdots = F_I$, use $Q \overset{H_0}{\approx} \chi^2_{I-1}$ for moderate $J$.
(iv) Reject at level $\alpha$ when $\mathbb{P}\{\chi^2_{I-1} \geq Q_{\text{obs}}\} \leq \alpha$.

**Example 4.5.6.** *Generic Friedman Rank Calculation*
For $I = 7$ treatments, suppose the average within-block ranks are

| Treatment | $T_1$ | $T_2$ | $T_3$ | $T_4$ | $T_5$ | $T_6$ | $T_7$ |
|---|---|---|---|---|---|---|---|
| Average rank | 5.10 | 4.90 | 2.30 | 3.50 | 3.05 | 4.90 | 4.25 |

With $\overline{R}_{\cdot\cdot} = 4$, $\sum_i (\overline{R}_{i\cdot} - \overline{R}_{\cdot\cdot})^2 = 6.935$ and $Q = 14.86$.
Using $Q \overset{H_0}{\approx} \chi^2_6$, the tail probability is below .025, so the treatment response distributions are not all equal.

**Corollary 4.5.7.** *Two Treatments Recover the Sign Test*
*For $I = 2$, set $D_j := Y_{1j} - Y_{2j}$ and $B := \#\{j : D_j > 0\}$. Friedman's statistic is a function of $B$ alone, so the two-treatment Friedman test is equivalent to the paired sign test.*

*Proof.* Ignoring ties, $R_{1j} = 2\mathbf{1}_{\{D_j > 0\}} + \mathbf{1}_{\{D_j < 0\}} = 1 + \mathbf{1}_{\{D_j > 0\}}$.
Hence $\sum_j R_{1j} = J + B$, so $\overline{R}_{1\cdot}$, $\overline{R}_{2\cdot}$, and $Q$ are determined by $B$.
Under $H_0$, signs are exchangeable and $B \sim \text{Bin}(J, 1/2)$, which is the paired sign-test calibration.
*Structural Commentary*: Two within-block ranks encode only the sign of $D_j$.
*Key Insight*: The rank-sum statistic collapses to a binomial sign count. □

# 5 Linear Models and Regression

## 5.1 Least Squares Curve Fitting and Linearization

Least squares begins as a geometric fitting rule and becomes a statistical model only after assumptions are made about the errors. The optimization step is asymmetric in the two plotted variables: ordinary regression measures vertical prediction errors.

**Definition 5.1.1.** *Least-Squares Fit to a Parametric Family*
Let $[n] := \{1, \ldots, n\}$, let $(x_i, y_i)_{i \in [n]} \subseteq \mathcal{X} \times \mathbb{R}$, let $\Theta \subseteq \mathbb{R}^p$, and let $m : \mathcal{X} \times \Theta \to \mathbb{R}$.

(i) *Criterion*: $S : \Theta \to [0, \infty)$, $S(\theta) := \sum_{i=1}^{n} \big(y_i - m(x_i; \theta)\big)^2$.

(ii) *Estimate*: $\widehat{\theta} \in \arg\min_{\theta \in \Theta} S(\theta)$.

(iii) *Residual*: $\widehat{e}_i := y_i - m(x_i; \widehat{\theta})$, $i \in [n]$.

*Key Insight*: $y_i = m(x_i; \widehat{\theta}) + \widehat{e}_i$, so $\widehat{e}_i > 0$ iff $y_i > m(x_i; \widehat{\theta})$.




Figure 252: Least squares minimizes squared vertical residuals.

**Definition 5.1.2.** *Linear in the Parameters*
Let $\mathcal{X}$ be an input set, let $\Theta \subseteq \mathbb{R}^p$, and let $m : \mathcal{X} \times \Theta \to \mathbb{R}$. $m$ is linear in $\beta$ if and only if
$\exists$ fixed maps $\phi_0, \ldots, \phi_{p-1} : \mathcal{X} \to \mathbb{R}$ with $m(x; \beta) = \sum_{j=0}^{p-1} \beta_j \phi_j(x)$ $\forall x \in \mathcal{X}$ and $\beta \in \Theta$.
*Structural Commentary*: Polynomial regression uses $\phi_j(x) = x^j$ and estimates only the coefficients $\beta_j$.

**Proposition 5.1.3.** *Least-Squares Line*
*Let $[n] := \{1, \ldots, n\}$, $(x_i, y_i)_{i \in [n]} \subseteq \mathbb{R}^2$, $\overline{x} := \frac{1}{n} \sum_{i=1}^{n} x_i$, $\overline{y} := \frac{1}{n} \sum_{i=1}^{n} y_i$, assume $D := \sum_{i=1}^{n} (x_i - \overline{x})^2 > 0$.*
*For $b = (b_0, b_1) \in \mathbb{R}^2$, set $S(b) := \sum_{i=1}^{n} (y_i - b_0 - b_1 x_i)^2$.*
*Then $\exists! \widehat{\beta} = (\widehat{\beta}_0, \widehat{\beta}_1) \in \arg\min_{b \in \mathbb{R}^2} S(b)$; $\widehat{y}(x) := \widehat{\beta}_0 + \widehat{\beta}_1 x$ satisfies $\widehat{\beta}_1 = \frac{1}{D} \sum_{i=1}^{n} (x_i - \overline{x})(y_i - \overline{y})$, $\widehat{\beta}_0 = \overline{y} - \widehat{\beta}_1 \overline{x}$.*

*Proof.* Set $u_i := x_i - \overline{x}$, $v_i := y_i - \overline{y}$, and $c(b) := b_0 + b_1\overline{x} - \overline{y}$.
Since $\sum_{i=1}^{n} u_i = \sum_{i=1}^{n} v_i = 0$, $S(b) = \sum_{i=1}^{n}(v_i - b_1 u_i)^2 + nc(b)^2$.
Thus $c(\widehat{\beta}) = 0$, so $\widehat{\beta}_0 = \overline{y} - \widehat{\beta}_1\overline{x}$, and $\widehat{\beta}_1$ uniquely minimizes $Q(t) := \sum_{i=1}^{n}(v_i - tu_i)^2$.
Since $Q'(t) = -2\sum_{i=1}^{n} u_i(v_i - tu_i)$ and $\sum_{i=1}^{n} u_i^2 = D > 0$, the unique critical point is $t = \frac{1}{D}\sum_{i=1}^{n} u_i v_i$.
*Structural Commentary*: The fitted line passes through $(\overline{x}, \overline{y})$.
*Key Insight*: Centering turns the slope into covariance divided by variance. □

**Example 5.1.4.** *Linear and Nonlinear Parameters*
For fixed $\omega = (\omega_1, \omega_2)$, $f_{\theta,\omega}(t) := A\cos(\omega_1 t) + B\sin(\omega_1 t) + C\cos(\omega_2 t) + D\sin(\omega_2 t) = \theta^t \phi_\omega(t)$
with $\theta = (A, B, C, D)^t$ and $\phi_\omega(t) = (\cos(\omega_1 t), \sin(\omega_1 t), \cos(\omega_2 t), \sin(\omega_2 t))^t$.
Estimating $\omega$ makes $f_{\theta,\omega}$ nonlinear in $(\theta, \omega)$.
For fixed $\lambda = (\alpha, \beta)$, $g_{\eta,\lambda}(t) := Ae^{-\alpha t} + Be^{-\beta t} = \eta^t \psi_\lambda(t)$ with $\eta = (A, B)^t$ and $\psi_\lambda(t) = (e^{-\alpha t}, e^{-\beta t})^t$.
Estimating $\lambda$ makes $g_{\eta,\lambda}$ nonlinear in $(\eta, \lambda)$.

**Method 5.1.5.** *Linearizing Transformations*
Let $[n] := \{1, \ldots, n\}$, $K > 0$, $C > 0$, $e_A \in \mathbb{R}$, $T > 0$, and $\alpha(T) := C\exp\left(-\frac{e_A}{KT}\right)$.
For observations $(T_i, \alpha_i)_{i\in[n]} \subset (0,\infty)^2$, set $x := \frac{1}{T}$, $z := \log\alpha(T)$, $x_i := \frac{1}{T_i}$, and $z_i := \log\alpha_i$.

(i) $z = a + bx$ with $a := \log C$ and $b := -\frac{e_A}{K}$.

(ii) $(\widehat{a}, \widehat{b}) \in \arg\min_{(a,b)\in\mathbb{R}^2} \sum_{i=1}^{n}(z_i - a - bx_i)^2$, $\widehat{C} := e^{\widehat{a}}$, and $\widehat{e}_A := -K\widehat{b}$.

(iii) $\widehat{\varepsilon}_i := z_i - \widehat{a} - \widehat{b}x_i$ is diagnosed against $x_i$ and $\widehat{z}_i := \widehat{a} + \widehat{b}x_i$.
Systematic structure indicates misspecification on the transformed scale.

## 5.2 Simple Linear Regression Inference and Diagnostics

Simple linear regression adds a noise model to the least-squares line. The $x_i$ values are treated as fixed design points, and randomness enters through the observed responses.

**Definition 5.2.1.** *Standard Simple Linear Regression Model*
Let $[n] := \{1, \ldots, n\}$, let $x_i \in \mathbb{R}$ be fixed for $i \in [n]$, let $\beta_0, \beta_1 \in \mathbb{R}$, and let $\sigma^2 > 0$.

(i) *Standard model*: $Y_i = \beta_0 + \beta_1 x_i + e_i$, $\mathbb{E}e_i = 0$, $\mathrm{Var}(e_i) = \sigma^2$, and $\mathrm{Cov}(e_i, e_j) = 0$ for $i \neq j$, $i, j \in [n]$.

(ii) *Normal model*: $Y_i = \beta_0 + \beta_1 x_i + e_i$ with $e_1, \ldots, e_n$ independent and $e_i \sim N(0, \sigma^2)$, $i \in [n]$.

*Structural Commentary*: $(x_i)_{i\in[n]}$ is fixed; randomness enters through $(e_i)_{i\in[n]}$.

**Theorem 5.2.2.** *Mean and Variance of the Simple Regression Coefficients*
*Let* $[n] := \{1, \ldots, n\}$, *let* $x_i \in \mathbb{R}$, *and set* $\overline{x} := \frac{1}{n}\sum_{i=1}^{n} x_i$ *and* $D := \sum_{i=1}^{n}(x_i - \overline{x})^2 > 0$.
*Let* $\beta := (\beta_0, \beta_1)$ *and assume* $Y_i = \beta_0 + \beta_1 x_i + e_i$, $\mathbb{E}e_i = 0$, $\mathrm{Var}(e_i) = \sigma^2$, *and* $\mathrm{Cov}(e_i, e_j) = 0$ *for* $i \neq j$.
*If* $\widehat{\beta} = (\widehat{\beta}_0, \widehat{\beta}_1)$ *minimizes* $\sum_{i=1}^{n}(Y_i - b_0 - b_1 x_i)^2$ *over* $b = (b_0, b_1) \in \mathbb{R}^2$, *then* $\mathbb{E}\widehat{\beta} = \beta$; $\mathrm{Var}(\widehat{\beta}) = \sigma^2 \begin{pmatrix} \frac{1}{n} + \frac{\overline{x}^2}{D} & -\frac{\overline{x}}{D} \\ -\frac{\overline{x}}{D} & \frac{1}{D} \end{pmatrix}$.

*Proof.* Set $u_i := x_i - \overline{x}$ and $\overline{e} := \frac{1}{n}\sum_{i=1}^{n} e_i$.
Proposition 5.1.3 gives $\widehat{\beta}_1 - \beta_1 = D^{-1}\sum_{i=1}^{n} u_i e_i$ and $\widehat{\beta}_0 - \beta_0 = \overline{e} - \overline{x}(\widehat{\beta}_1 - \beta_1)$.
Because $\sum_{i=1}^{n} u_i = 0$, $\mathbb{E}(\widehat{\beta} - \beta) = 0$, $\mathrm{Var}(\widehat{\beta}_1) = D^{-2}\sum_{i=1}^{n} u_i^2\sigma^2 = \frac{\sigma^2}{D}$, $\mathrm{Cov}(\overline{e}, \widehat{\beta}_1) = \frac{1}{n}D^{-1}\sum_{i=1}^{n} u_i\sigma^2 = 0$, and $\mathrm{Var}(\overline{e}) = \frac{\sigma^2}{n}$. Therefore $\mathrm{Var}(\widehat{\beta}_0) = \frac{\sigma^2}{n} + \frac{\overline{x}^2\sigma^2}{D}$ and $\mathrm{Cov}(\widehat{\beta}_0, \widehat{\beta}_1) = -\frac{\overline{x}\sigma^2}{D}$.
*Structural Commentary*: The slope is a fixed linear combination of the responses.
*Key Insight*: Centering the design isolates the uncertainty in the slope. □

**Definition 5.2.3.** *Residual Sum of Squares and Error Scale*
Let $[n] := \{1, \ldots, n\}$ with $n > 2$, and let $\widehat{y}_i = \widehat{\beta}_0 + \widehat{\beta}_1 x_i$ for $i \in [n]$.

(i) *Residual Sum of Squares*: $\mathrm{RSS} := \sum_{i=1}^{n}(Y_i - \widehat{y}_i)^2$.

(ii) *Error Scale*: $s^2 := \frac{\mathrm{RSS}}{n-2}$ and $s := \sqrt{s^2}$.

*Key Insight*: $n - 2$ subtracts the fitted coefficients $\widehat{\beta}_0, \widehat{\beta}_1$.

**Corollary 5.2.4.** *Simple Regression t Intervals and Tests*
*Let* $n > 2$, $[n] := \{1, \dots, n\}$, $\alpha \in (0,1)$, $\beta \in \mathbb{R}^2$, $\sigma^2 > 0$, $x = (x_1, \dots, x_n)^t \in \mathbb{R}^n$, $X := (\mathbf{1}, x) \in \mathbb{R}^{n \times 2}$ *with* $\operatorname{rank}(X) = 2$, $Y = X\beta + e$, *and* $e \sim N(0, \sigma^2 I_n)$.
*Set* $\widehat{\beta} := (X^t X)^{-1} X^t Y$, $s^2 := \frac{\|Y - X\widehat{\beta}\|_2^2}{n-2}$, $\overline{x} := \frac{1}{n}\sum_{i=1}^{n} x_i$, $D := \sum_{i=1}^{n}(x_i - \overline{x})^2$, $c_0 := \frac{1}{n} + \frac{\overline{x}^2}{D}$, *and* $c_1 := D^{-1}$.
*For* $j \in \{0,1\}$ *and* $\beta_{j0} \in \mathbb{R}$, *set* $s_{\widehat{\beta}_j} := s\sqrt{c_j}$.
*Then* $T_j := \frac{\widehat{\beta}_j - \beta_j}{s_{\widehat{\beta}_j}} \sim t_{n-2}$, $C_j(Y) := [\widehat{\beta}_j - t_{n-2,1-\frac{\alpha}{2}} s_{\widehat{\beta}_j}, \widehat{\beta}_j + t_{n-2,1-\frac{\alpha}{2}} s_{\widehat{\beta}_j}]$ *satisfies* $\mathbb{P}_\beta(\beta_j \in C_j(Y)) = 1 - \alpha$,
*and the two-sided level-*$\alpha$ *test of* $H_0 : \beta_j = \beta_{j0}$ *rejects iff* $\left|\frac{\widehat{\beta}_j - \beta_{j0}}{s_{\widehat{\beta}_j}}\right| > t_{n-2,1-\frac{\alpha}{2}}$.

*Proof.* Let $P := X(X^tX)^{-1}X^t$, $M := I - P$, and $G := (X^tX)^{-1} = (g_{jk})_{j,k=0}^{1}$.
Then $\widehat{\beta} - \beta = GX^t e$, $g_{00} = c_0$, $g_{11} = c_1$, $\operatorname{Var}(\widehat{\beta}_j) = \sigma^2 c_j$, and $Z_j := \frac{\widehat{\beta}_j - \beta_j}{\sigma\sqrt{c_j}} \sim N(0,1)$.
Also $\frac{(n-2)s^2}{\sigma^2} = \frac{e^t M e}{\sigma^2}$, $M = M^t = M^2$, $\operatorname{rank}(M) = n-2$, and $MX = 0$, hence $U := \frac{(n-2)s^2}{\sigma^2} \sim \chi^2_{n-2}$ and $\operatorname{Cov}(GX^te, Me) = GX^t\sigma^2 M = 0$. Joint normality gives $Z_j \perp U$, so $T_j = \frac{Z_j}{\sqrt{\frac{U}{n-2}}} \sim t_{n-2}$.
The interval and rejection rule follow from $\mathbb{P}(|T_j| \le t_{n-2,1-\frac{\alpha}{2}}) = 1 - \alpha$.
*Structural Commentary*: $T_j = \frac{Z_j}{\sqrt{\frac{U}{n-2}}}$ with $Z_j \sim N(0,1)$, $U \sim \chi^2_{n-2}$, and $Z_j \perp U$.
*Key Insight*: Normality and orthogonal residual projection make replacing $\sigma$ by $s$ exact. □

**Example 5.2.5.** *Calibration by Peak Area*
Let $[n] := \{1, \dots, n\}$, with known standard concentrations $x_i$ and measured peak areas $A_i$, $i \in [n]$.
Fit $A_i = \beta_0 + \beta_1 x_i + e_i$ by least squares, giving $\widehat{A}(x) = \widehat{\beta}_0 + \widehat{\beta}_1 x$.
For a new area $A_*$, the inverse calibration estimate is $\widehat{x}_* := \frac{A_* - \widehat{\beta}_0}{\widehat{\beta}_1}$ when $\widehat{\beta}_1 \neq 0$.
In the reported fit, $\widehat{\beta}_0 = .0729$, $\widehat{\beta}_1 = 10.77$, $s = .068$, $\operatorname{se}(\widehat{\beta}_0) = .0297$, and $\operatorname{se}(\widehat{\beta}_1) = .27$, hence $t_0 = 2.45$, $t_1 = 40.20$, and the reported 95% intervals are $C_0 = (.011, .135)$ and $C_1 = (10.21, 11.33)$.
Thus $\widehat{\beta}_1 > 0$ makes peak area a monotone calibration signal.

**Definition 5.2.6.** *Homoscedastic and Heteroscedastic Errors*
Let $[n] := \{1, \dots, n\}$, let $e = (e_1, \dots, e_n)^t$ be errors at design points $x_i$, and set $v_i := \operatorname{Var}(e_i)$.
(i) *Homoscedastic*: $\exists \sigma^2 \ge 0$ with $v_i = \sigma^2$, $i \in [n]$.
(ii) *Heteroscedastic*: $\exists i, j \in [n]$ with $v_i \neq v_j$.
*Key Insight*: $\operatorname{Var}(e) \neq \sigma^2 I_n$ invalidates formulas requiring $\operatorname{Var}(e) = \sigma^2 I_n$.

**Method 5.2.7.** *Residual Diagnostics*
After fitting a regression, examine residuals against the fitted values and against important predictors.
(i) A random horizontal band supports the chosen mean form.
(ii) Curvature suggests a transformation or added nonlinear term.
(iii) Changing spread suggests heteroscedasticity.
(iv) A normal probability plot checks large departures from normality and possible outliers.
(v) Points with extreme $x$ values may have high leverage and can strongly affect the fitted line.
These checks refine the modeling assumptions; they do not replace them.

**Example 5.2.8.** *Diagnostic Patterns*
Let $[n] := \{1, \dots, n\}$, $r_i := Y_i - \widehat{Y}_i$, $f_i := \widehat{Y}_i$, $\ell_i \in \mathcal{G}$,
$m_r(x) := \mathbb{E}(r_i \mid x_i = x)$, $v_r(f) := \operatorname{Var}(r_i \mid f_i = f)$, and $m_a := \mathbb{E}(r_i \mid \ell_i = a)$.
(i) $m_r \not\equiv 0$ or $m_r$ non-affine; transform variables or add nonlinear terms.
(ii) $v_r$ nonconstant; transform $Y_i$ or use heteroscedastic errors.
(iii) $\exists a, b \in \mathcal{G}$ with $m_a \neq m_b$ or $\exists i \neq j$ with $\operatorname{Cov}(r_i, r_j \mid \ell_i = \ell_j) \neq 0$; add group effects or correlated errors.

**Proposition 5.2.9.** *Correlation and the Regression Line*
*Let* $[n] := \{1, \dots, n\}$, *let* $(x_i, y_i) \in \mathbb{R}^2$ *for* $i \in [n]$.
*Set* $\overline{x} := \frac{1}{n}\sum_{i=1}^{n} x_i$, $\overline{y} := \frac{1}{n}\sum_{i=1}^{n} y_i$, $s_{xx} := \frac{1}{n}\sum_{i=1}^{n}(x_i - \overline{x})^2$, $s_{yy} := \frac{1}{n}\sum_{i=1}^{n}(y_i - \overline{y})^2$, $s_{xy} := \frac{1}{n}\sum_{i=1}^{n}(x_i - \overline{x})(y_i - \overline{y})$.
*If* $s_{xx}, s_{yy} > 0$, $r := \frac{s_{xy}}{\sqrt{s_{xx}s_{yy}}}$, $(\widehat{\beta}_0, \widehat{\beta}_1)$ *minimizes* $\sum_{i=1}^{n}(y_i - b_0 - b_1 x_i)^2$ *over* $(b_0, b_1) \in \mathbb{R}^2$,
*and* $\widehat{y}(x) := \widehat{\beta}_0 + \widehat{\beta}_1 x$, *then* $\widehat{\beta}_1 = \frac{s_{xy}}{s_{xx}}$, $r = \widehat{\beta}_1\sqrt{\frac{s_{xx}}{s_{yy}}}$, *and* $\frac{\widehat{y}(x) - \overline{y}}{\sqrt{s_{yy}}} = r\frac{x - \overline{x}}{\sqrt{s_{xx}}}$.

*Proof.* Set $D := \sum_{i=1}^{n}(x_i - \overline{x})^2 = ns_{xx}$ and $N := \sum_{i=1}^{n}(x_i - \overline{x})(y_i - \overline{y}) = ns_{xy}$.
Proposition 5.1.3 gives $\widehat{\beta}_1 = \frac{N}{D} = \frac{s_{xy}}{s_{xx}}$ and $\widehat{\beta}_0 = \overline{y} - \widehat{\beta}_1\overline{x}$.
Hence $r = \frac{s_{xy}}{\sqrt{s_{xx}s_{yy}}} = \widehat{\beta}_1\sqrt{\frac{s_{xx}}{s_{yy}}}$ and $\widehat{y}(x) - \overline{y} = \widehat{\beta}_1(x - \overline{x})$, so $\frac{\widehat{y}(x)-\overline{y}}{\sqrt{s_{yy}}} = r\frac{x-\overline{x}}{\sqrt{s_{xx}}}$.
*Structural Commentary*: $z_x := \frac{x-\overline{x}}{\sqrt{s_{xx}}}$ and $z_{\widehat{y}} := \frac{\widehat{y}(x)-\overline{y}}{\sqrt{s_{yy}}}$ give $z_{\widehat{y}} = rz_x$.
*Key Insight*: $\widehat{\beta}_1 = \frac{s_{xy}}{s_{xx}}$ becomes $r = \frac{s_{xy}}{\sqrt{s_{xx}s_{yy}}}$ after standardising both coordinates. □

**Remark 5.2.10.** *Regression to the Mean*
With $s_{xx}, s_{yy} > 0$ and $0 < r < 1$, set $z_x := \frac{x-\overline{x}}{\sqrt{s_{xx}}}$ and $z_{\widehat{y}} := \frac{\widehat{y}(x)-\overline{y}}{\sqrt{s_{yy}}}$.
Proposition 5.2.9 gives $z_{\widehat{y}} = rz_x$, hence $|z_{\widehat{y}}| < |z_x|$ for $z_x \neq 0$.

## 5.3 Matrix Linear Models and Normal Equations

Matrix notation separates the geometry of least squares from the statistical assumptions on the errors. It also makes multiple regression a direct extension of simple regression.

**Definition 5.3.1.** *Fixed-Design Linear Model*
Let $Y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\beta, b \in \mathbb{R}^p$, and $e \in \mathbb{R}^n$.

(i) *Model*: $Y = X\beta + e$.
(ii) *Least-Squares Estimator*: $\widehat{\beta} \in \arg\min_{b\in\mathbb{R}^p} \|Y - Xb\|_2^2$.

*Structural Commentary*: An intercept is encoded by $X_{\cdot 1} = \mathbf{1}$.

**Theorem 5.3.2.** *Normal Equations for the Linear Model*
*Let $Y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, and $Q(b) := \|Y - Xb\|_2^2$ for $b \in \mathbb{R}^p$.*

*(i) $\widehat{\beta} \in \arg\min_{b\in\mathbb{R}^p} Q(b)$ iff $X^tX\widehat{\beta} = X^tY$.*
*(ii) If $\operatorname{rank}(X) = p$, then $X^tX$ is invertible and $\arg\min_{b\in\mathbb{R}^p} Q(b) = \{(X^tX)^{-1}X^tY\}$.*

*Proof.*

(*i*) ($\Rightarrow$) If $\widehat{\beta}$ minimizes $Q$, then $\forall (t,h) \in \mathbb{R}\times\mathbb{R}^p$, $0 \le Q(\widehat{\beta} + th) - Q(\widehat{\beta}) = 2th^tX^t(X\widehat{\beta} - Y) + t^2\|Xh\|_2^2$.
Hence $\forall h \in \mathbb{R}^p$, $h^tX^t(X\widehat{\beta} - Y) = 0$, so $X^tX\widehat{\beta} = X^tY$.
(*i*) ($\Leftarrow$) If $X^tX\widehat{\beta} = X^tY$, then $\forall h \in \mathbb{R}^p$, $Q(\widehat{\beta} + h) - Q(\widehat{\beta}) = \|Xh\|_2^2 \ge 0$, so $\widehat{\beta}$ minimizes $Q$.
(*ii*) If $\operatorname{rank}(X) = p$, then $\forall u \in \mathbb{R}^p \setminus \{0\}$, $u^tX^tXu = \|Xu\|_2^2 > 0$, so $X^tX$ is invertible.
Applying (i) gives $\arg\min_{b\in\mathbb{R}^p} Q(b) = \{(X^tX)^{-1}X^tY\}$.

*Structural Commentary*: $\widehat{Y} = X\widehat{\beta}$ and $\widehat{e} = Y - \widehat{Y}$ satisfy $X^t\widehat{e} = 0$, hence $\widehat{e} \perp \operatorname{col}(X)$.
*Key Insight*: The projection $\widehat{Y} \in \operatorname{col}(X)$ is unique; $\operatorname{rank}(X) = p$ makes $b \mapsto Xb$ injective, so $\widehat{\beta}$ is unique. □

**Method 5.3.3.** *Projection View of Least Squares*

(i) *Full rank*: If $\operatorname{rank}(X) = d$, define $H := X(X^\top X)^{-1}X^\top$ and $\widehat{y} := Hy$.
(ii) *Projection*: $H$ is the orthogonal projection onto $\operatorname{col}(X)$; hence $y - \widehat{y} \perp \operatorname{col}(X)$.
(iii) *Rank deficiency*: If $\operatorname{rank}(X) < d$, $\widehat{\beta}$ may be nonunique, but $\widehat{y} = \operatorname{proj}_{\operatorname{col}(X)} y$ is unique.

Rank deficiency affects coefficient representation, not least-squares fitted values.

**Definition 5.3.4.** *Hat Matrix, Fitted Values, and Residuals*
Assume $Y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\operatorname{rank}(X) = p$, $\widehat{\beta} = (X^tX)^{-1}X^tY$, and $I = I_n$.

(i) *Hat Matrix*: $P := X(X^tX)^{-1}X^t$.
(ii) *Fitted Values*: $\widehat{Y} := X\widehat{\beta} = PY$.
(iii) *Residuals*: $\widehat{e} := Y - \widehat{Y} = (I - P)Y$.

*Key Insight*: $\operatorname{range}(P) = \operatorname{col}(X)$ and $PY = \widehat{Y}$.

**Proposition 5.3.5.** *Projection Properties of the Hat Matrix*
*Let $Y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$ with $\operatorname{rank}(X) = p$, $I = I_n$, $P := X(X^tX)^{-1}X^t$, $\widehat{Y} := PY$, and $\widehat{e} := (I - P)Y$.*
*Then $P = P^t = P^2$, $I - P = (I - P)^t = (I - P)^2$, $\operatorname{range}(P) = \operatorname{col}(X)$, $\widehat{Y} = P_{\operatorname{col}(X)}Y$, and $X^t\widehat{e} = 0$.*

*Proof.* Since $(X^tX)^t = X^tX$, $P^t = X((X^tX)^{-1})^tX^t = P$, and $P^2 = X(X^tX)^{-1}X^tX(X^tX)^{-1}X^t = P$.
Hence $(I - P)^t = I - P$ and $(I - P)^2 = I - 2P + P^2 = I - P$.
If $z = Xb \in \operatorname{col}(X)$, then $Pz = X(X^tX)^{-1}X^tXb = z$; also $\operatorname{range}(P) \subseteq \operatorname{col}(X)$, so $\operatorname{range}(P) = \operatorname{col}(X)$.

Finally, $X^t\widehat{e} = X^t(I-P)Y = X^tY - X^tX(X^tX)^{-1}X^tY = 0$, so $\widehat{e} \perp \mathrm{col}(X)$ and $\widehat{Y} = P_{\mathrm{col}(X)}Y$.
*Structural Commentary*: $P = P_{\mathrm{col}(X)}$ and $I - P = P_{\mathrm{col}(X)^\perp}$.
*Key Insight*: Fitting decomposes $Y = PY + (I-P)Y = \widehat{Y} + \widehat{e}$ with $\widehat{Y} \perp \widehat{e}$. □

**Theorem 5.3.6.** *Unbiased Estimate of the Error Variance*
*Let $Y = X\beta + e$ with $Y, e \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\beta \in \mathbb{R}^p$, $\mathrm{rank}(X) = p$, $\mathbb{E}e = 0$, and $\mathrm{Var}(e) = \sigma^2 I_n$.*
*Set $P := X(X^tX)^{-1}X^t$, $\widehat{Y} := PY$, $\widehat{e} := Y - \widehat{Y}$, and $s^2 := \frac{\|\widehat{e}\|_2^2}{n-p}$. Then $\mathbb{E}\,\|\widehat{e}\|_2^2 = (n-p)\sigma^2$ and $\mathbb{E}s^2 = \sigma^2$.*

*Proof.* Let $M := I_n - P$. Since $PX = X$, $\widehat{e} = MY = M(X\beta + e) = Me$.
Since $M = M^t = M^2$, $\|\widehat{e}\|_2^2 = e^tM^tMe = e^tMe$. For fixed $A \in \mathbb{R}^{n\times n}$ and centered $e$ with $\mathrm{Var}(e) = \sigma^2 I_n$,
$\mathbb{E}(e^tAe) = \mathrm{tr}(A\sigma^2 I_n) = \sigma^2\,\mathrm{tr}(A)$. Thus $\mathbb{E}\,\|\widehat{e}\|_2^2 = \sigma^2\,\mathrm{tr}(M)$,
and $\mathrm{tr}(M) = n - \mathrm{tr}(P) = n - \mathrm{tr}((X^tX)^{-1}X^tX) = n - p$. Therefore $\mathbb{E}s^2 = \frac{1}{n-p}\mathbb{E}\,\|\widehat{e}\|_2^2 = \sigma^2$.
*Structural Commentary*: $M = I_n - P = P_{\mathrm{col}(X)^\perp}$ and $\mathrm{tr}(M) = n - p$.
*Key Insight*: $\widehat{e} = Me$, so residual variation is noise projected onto the $n-p$ directions orthogonal to $\mathrm{col}(X)$. □

**Definition 5.3.7.** *Standardized Residual*
Let $[n] := \{1, \ldots, n\}$, $Y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$ with $\mathrm{rank}(X) = p$, $P = (p_{ij}) := X(X^tX)^{-1}X^t$, $\widehat{Y} := PY$, $\widehat{e} := Y - \widehat{Y}$, $s^2 := \frac{\|\widehat{e}\|_2^2}{n-p}$, and $s > 0$. The *standardized residual* is $r_i := \frac{\widehat{e}_i}{s\sqrt{1-p_{ii}}}$ for $i \in [n]$.
*Key Insight*: Under $\mathrm{Var}(e) = \sigma^2 I_n$, $\mathrm{Var}(\widehat{e}_i) = \sigma^2(1-p_{ii})$.

**Proposition 5.3.8.** *Residuals Are Uncorrelated with Fitted Values*
*Let $Y = X\beta + e$ with $Y, e \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\beta \in \mathbb{R}^p$, $\mathrm{rank}(X) = p$, and $\mathrm{Var}(e) = \sigma^2 I_n$.*
*Set $P := X(X^tX)^{-1}X^t$, $\widehat{Y} := PY$, and $\widehat{e} := (I_n - P)Y$. Then $\mathrm{Cov}(\widehat{e}, \widehat{Y}) = 0$.*

*Proof.* Since $\mathrm{Cov}(Y) = \mathrm{Cov}(e) = \sigma^2 I_n$ and $P = P^t = P^2$,
$\mathrm{Cov}(\widehat{e}, \widehat{Y}) = (I_n - P)\,\mathrm{Cov}(Y)P^t = \sigma^2(I_n - P)P = \sigma^2(P - P^2) = 0$.
*Structural Commentary*: $\widehat{Y} = PY$ and $\widehat{e} = (I_n - P)Y$ are images of $Y$ under complementary projections.
*Key Insight*: Spherical error covariance makes orthogonal projection components uncorrelated. □

## 5.4 Inference, Prediction, and Multiple Regression

Once the linear model is written in matrix form, standard errors for coefficients and predictions are consequences of linear transformations of random vectors.

**Theorem 5.4.1.** *Mean and Covariance of Least-Squares Estimates*
*Let $Y = X\beta + e$ with $Y, e \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\beta \in \mathbb{R}^p$, $\mathrm{rank}(X) = p$, $\mathbb{E}e = 0$, and $\mathrm{Var}(e) = \sigma^2 I_n$.*
*Set $\widehat{\beta} := (X^tX)^{-1}X^tY$. Then $\mathbb{E}\widehat{\beta} = \beta$ and $\mathrm{Var}(\widehat{\beta}) = \sigma^2(X^tX)^{-1}$.*

*Proof.* Let $A := (X^tX)^{-1}X^t$. Since $AX = I_p$, $\widehat{\beta} = A(X\beta + e) = \beta + Ae$.
Hence $\mathbb{E}\widehat{\beta} = \beta + A\mathbb{E}e = \beta$, and $\mathrm{Var}(\widehat{\beta}) = A\,\mathrm{Var}(e)A^t = \sigma^2(X^tX)^{-1}X^tX(X^tX)^{-1} = \sigma^2(X^tX)^{-1}$.
*Structural Commentary*: $\widehat{\beta} - \beta = (X^tX)^{-1}X^te$ is a fixed linear transformation of $e$.
*Key Insight*: $X^tX$ controls precision: $\mathrm{Var}(\widehat{\beta}) = \sigma^2(X^tX)^{-1}$. □

**Method 5.4.2.** *Coefficient Inference in a Normal Linear Model*
Let $Y = X\beta + e$, $e \sim N_n(0, \sigma^2 I_n)$, $X \in \mathbb{R}^{n\times p}$ with $\mathrm{rank}(X) = p$, $\widehat{\beta} := (X^tX)^{-1}X^tY$,
$s^2 := \frac{1}{n-p}\left\|Y - X\widehat{\beta}\right\|_2^2$, $C = (c_{jk}) := (X^tX)^{-1}$, $j \in \{1, \ldots, p\}$, and $\alpha \in (0,1)$.

(i) *Standard Error*: $s_{\widehat{\beta}_j} := s\sqrt{c_{jj}}$.

(ii) *Pivot*: $T_j := \frac{\widehat{\beta}_j - \beta_j}{s_{\widehat{\beta}_j}} \sim t_{n-p}$.

(iii) *Confidence Interval*: $C_{j,1-\alpha} := \left[\widehat{\beta}_j - t_{n-p,1-\frac{\alpha}{2}}s_{\widehat{\beta}_j}, \widehat{\beta}_j + t_{n-p,1-\frac{\alpha}{2}}s_{\widehat{\beta}_j}\right]$.

(iv) *Test Statistic*: For $H_0 : \beta_j = \beta_{j0}$, $t_j := \frac{\widehat{\beta}_j - \beta_{j0}}{s_{\widehat{\beta}_j}}$.

If a coefficient CLT holds, $T_j \approx N(0,1)$.

**Remark 5.4.3.** *Linear-Model Inference After Selection*

Under fixed-design $y = X\beta + \varepsilon$, $\varepsilon \sim N(0, \sigma^2 I_n)$, and $\mathrm{rank}(X) = d$, $\widehat{\beta} \sim N(\beta, \sigma^2(X^\top X)^{-1})$ and $\widehat{\sigma}^2 := \frac{\|y - X\widehat{\beta}\|_2^2}{n-d}$.
These give standard errors, $t$ or asymptotic $z$ statistics, and nested-model $F$ tests when $X$ is fixed in advance. After data-driven selection of $X$ or the model, these standard errors ignore selection.

**Proposition 5.4.4.** *Conditional Coverage Implies Unconditional Coverage*
*Let $C(X,Y)$ be a confidence set for $\beta_j$, $\alpha \in (0,1)$, and $F := \{\text{rank}(X) = p\}$.*
*If $\mathbb{P}(F) = 1$ and $\mathbb{P}(\beta_j \in C(X,Y) \mid X) = 1-\alpha$ on $F$, then $\mathbb{P}(\beta_j \in C(X,Y)) = 1-\alpha$.*

*Proof.* Set $A := \{\beta_j \in C(X,Y)\}$. Since $\mathbb{P}(F) = 1$, $\mathbb{P}(A) = \mathbb{E}[\mathbb{P}(A \mid X)] = \mathbb{E}[(1-\alpha)\mathbf{1}_F + \mathbb{P}(A \mid X)\mathbf{1}_{F^c}] = 1-\alpha$.
*Structural Commentary*: $\mathbb{P}(F) = 1$ and $\mathbb{P}(A \mid X) = 1-\alpha$ on $F$.
*Key Insight*: $\mathbb{P}(A \mid X) = 1-\alpha$ a.s. implies $\mathbb{P}(A) = 1-\alpha$. □

**Proposition 5.4.5.** *Mean Response at a New Design Point*
*Let $Y = X\beta + e$ with $Y, e \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\beta \in \mathbb{R}^p$, $\text{rank}(X) = p$, and $e \sim N_n(0, \sigma^2 I_n)$.*
*Set $\widehat{\beta} := (X^tX)^{-1}X^tY$, $s^2 := \frac{1}{n-p}\left\|Y - X\widehat{\beta}\right\|_2^2$, $x_0 \in \mathbb{R}^p$, $c_0 := x_0^t(X^tX)^{-1}x_0 > 0$,*
*$\mu_0 := x_0^t\beta$, $\widehat{\mu}_0 := x_0^t\widehat{\beta}$, $s_{\widehat{\mu}_0} := s\sqrt{c_0}$, and $\alpha \in (0,1)$. Then $\mathbb{E}\widehat{\mu}_0 = \mu_0$, $\text{Var}(\widehat{\mu}_0) = \sigma^2 c_0$, $T_0 := \frac{\widehat{\mu}_0 - \mu_0}{s_{\widehat{\mu}_0}} \sim t_{n-p}$,*
*and $C_{0,1-\alpha} := \left[\widehat{\mu}_0 - t_{n-p,1-\frac{\alpha}{2}} s_{\widehat{\mu}_0}, \widehat{\mu}_0 + t_{n-p,1-\frac{\alpha}{2}} s_{\widehat{\mu}_0}\right]$ has coverage $1-\alpha$ for $\mu_0$.*

*Proof.* Let $A := (X^tX)^{-1}X^t$, $P := XA$, and $M := I_n - P$.
Since $AX = I_p$, $\widehat{\mu}_0 - \mu_0 = x_0^tA(X\beta + e) - x_0^t\beta = x_0^tAe$, so $\mathbb{E}\widehat{\mu}_0 = \mu_0$ and $\text{Var}(\widehat{\mu}_0) = x_0^tA(\sigma^2 I_n)A^tx_0 = \sigma^2 c_0$.
Normality gives $Z := \frac{\widehat{\mu}_0 - \mu_0}{\sigma\sqrt{c_0}} \sim N(0,1)$.
Since $M = M^t = M^2$, $\text{rank}(M) = n-p$, and $MA^tx_0 = 0$, $Z \perp U := \frac{e^tMe}{\sigma^2} = \frac{(n-p)s^2}{\sigma^2} \sim \chi^2_{n-p}$.
Thus $T_0 = \frac{Z}{\sqrt{U/(n-p)}} \sim t_{n-p}$, and $\mathbb{P}(\mu_0 \in C_{0,1-\alpha}) = \mathbb{P}(|T_0| \le t_{n-p,1-\frac{\alpha}{2}}) = 1-\alpha$.
*Structural Commentary*: $\widehat{\mu}_0 - \mu_0 = x_0^t(X^tX)^{-1}X^te$ and $s^2 = \frac{e^t(I_n - P)e}{n-p}$ use orthogonal components of $e$.
*Key Insight*: Estimating a mean response adds uncertainty only through $c_0 = x_0^t(X^tX)^{-1}x_0$. □

**Corollary 5.4.6.** *Prediction Interval for a New Observation*
*Let $Y = X\beta + e$ with $Y, e \in \mathbb{R}^n$, $X \in \mathbb{R}^{n\times p}$, $\beta \in \mathbb{R}^p$, $\text{rank}(X) = p$, and $e \sim N_n(0, \sigma^2 I_n)$.*
*Let $x_0 \in \mathbb{R}^p$, $e_0 \sim N(0, \sigma^2)$ with $e_0 \perp e$, and $Y_0 := x_0^t\beta + e_0$.*
*Set $\widehat{\beta} := (X^tX)^{-1}X^tY$, $\widehat{\mu}_0 := x_0^t\widehat{\beta}$, $s^2 := \frac{\|Y - X\widehat{\beta}\|_2^2}{n-p}$, $c_0 := x_0^t(X^tX)^{-1}x_0$, $s_{\text{pred}} := s\sqrt{1+c_0}$, and $\alpha \in (0,1)$.*
*Then $\text{Var}(Y_0 - \widehat{\mu}_0) = \sigma^2(1+c_0)$, $T_{\text{pred}} := \frac{Y_0 - \widehat{\mu}_0}{s_{\text{pred}}} \sim t_{n-p}$,*
*and $I_{0,1-\alpha} := \left[\widehat{\mu}_0 - t_{n-p,1-\frac{\alpha}{2}} s_{\text{pred}}, \widehat{\mu}_0 + t_{n-p,1-\frac{\alpha}{2}} s_{\text{pred}}\right]$ has coverage $1-\alpha$ for $Y_0$.*

*Proof.* Let $A := (X^tX)^{-1}X^t$, $P := XA$, and $M := I_n - P$.
Since $AX = I_p$, $Y_0 - \widehat{\mu}_0 = e_0 - x_0^tAe$, so $\text{Var}(Y_0 - \widehat{\mu}_0) = \sigma^2 + \sigma^2 x_0^tAA^tx_0 = \sigma^2(1+c_0)$.
Normality gives $Z := \frac{Y_0 - \widehat{\mu}_0}{\sigma\sqrt{1+c_0}} \sim N(0,1)$. Since $AM = 0$, $Z \perp U := \frac{e^tMe}{\sigma^2} = \frac{(n-p)s^2}{\sigma^2} \sim \chi^2_{n-p}$.
Thus $T_{\text{pred}} = \frac{Z}{\sqrt{U/(n-p)}} \sim t_{n-p}$, and $\mathbb{P}(Y_0 \in I_{0,1-\alpha}) = \mathbb{P}(|T_{\text{pred}}| \le t_{n-p,1-\frac{\alpha}{2}}) = 1-\alpha$.
*Structural Commentary*: $Y_0 - \widehat{\mu}_0 = e_0 - x_0^t(X^tX)^{-1}X^te$ combines future noise and estimation error.
*Key Insight*: Prediction variance is mean-response variance plus irreducible noise, $\sigma^2 c_0 + \sigma^2$. □

**Definition 5.4.7.** *Multiple Regression Coefficients*
Let $[n] := \{1, \dots, n\}$, $z_i = (x_{i1}, \dots, x_{i,p-1}) \in \mathbb{R}^{p-1}$ for $i \in [n]$, $u_1, \dots, u_{p-1}$ be the standard basis of $\mathbb{R}^{p-1}$, and $m(z) := \beta_0 + \sum_{k=1}^{p-1} \beta_k z_k$ In $Y_i = m(z_i) + e_i$, the *multiple regression coefficient* $\beta_k$ satisfies $\forall z \in \mathbb{R}^{p-1}$ and $\forall \delta \in \mathbb{R}$, $m(z + \delta u_k) - m(z) = \delta\beta_k$ for $k \in \{1, \dots, p-1\}$.
*Structural Commentary*: Polynomial regression replaces $z_i$ by transformed design columns such as $z_{ik}^2$.

**Definition 5.4.8.** *Coefficient of Determination*
Let $[n] := \{1, \dots, n\}$, $Y, \widehat{Y} \in \mathbb{R}^n$, $\overline{Y} := n^{-1}\sum_{i=1}^n Y_i$, $\mathbf{1} \in \text{col}(X)$, and $\sum_{i=1}^n (Y_i - \overline{Y})^2 > 0$.

(i) *Residual Sum of Squares*: $\text{RSS} := \sum_{i=1}^n (Y_i - \widehat{Y}_i)^2$.

(ii) *Total Sum of Squares*: $\text{TSS} := \sum_{i=1}^n (Y_i - \overline{Y})^2$.

(iii) *Coefficient of Determination*: $R^2 := 1 - \frac{\text{RSS}}{\text{TSS}}$.

*Structural Commentary*: $R^2 = \text{corr}(Y, \widehat{Y})^2$ for an intercept model.
*Key Insight*: $R^2 = \frac{\text{TSS} - \text{RSS}}{\text{TSS}}$ is the observed response variation removed by the fitted model.

**Remark 5.4.9.** *Adjusted Coefficients*
Let $X_{-j}$ be $X$ without column $x_j$, and set $z_j := (I - \text{proj}_{\text{col}(X_{-j})})x_j$.
Then $\widehat{\beta}_j = \frac{z_j^\top y}{z_j^\top z_j}$, the simple-regression coefficient of $y$ on the adjusted predictor $z_j$.

Thus $\widehat{\beta}_j$ uses only the part of $x_j$ left after adjusting for other predictors.
If $x_j \approx \mathrm{col}(X_{-j})$, then $\|z_j\|_2$ is small and $\widehat{\beta}_j$ is unstable; this is collinearity.

**Remark 5.4.10.** *Collinearity and Interpretation*
Let $X_{-j}$ be $X$ without column $X_{\cdot j}$, set $d_j := \mathrm{dist}(X_{\cdot j}, \mathrm{col}(X_{-j}))$, and let $C = (c_{kl}) := (X^tX)^{-1}$.
If $0 < d_j \ll \|X_{\cdot j}\|_2$, then $X^tX$ is nearly singular and $s_{\widehat{\beta}_j} = s\sqrt{c_{jj}}$ can be large even when $\widehat{Y} = X\widehat{\beta}$ is stable.
If $d_j = 0$, then $\mathrm{rank}(X) < p$ and $\widehat{\beta}$ is not unique.

**Example 5.4.11.** *Quadratic Terms as a Lack-of-Fit Check*
Compare $M_1 : Y_i = \beta_0 + \beta_1 x_i + e_i$ with $M_2 : Y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + e_i$.
If $t_2 = \widehat{\beta}_2 / s_{\widehat{\beta}_2}$ is small and residual curvature is absent, retain $M_1$.
If $|t_2|$ is large and residual curvature decreases under $M_2$, use the quadratic term.
Add $x^2$ only when residual structure or scale supports $\beta_2 \neq 0$.

## 5.5 Random Design, Bootstrap, and Extensions

The fixed-design model conditions on the observed predictor values. Many observational studies are more naturally viewed as sampling random predictor-response pairs, with the fixed-design analysis recovered after conditioning on the realised design.

**Definition 5.5.1.** *Random-Design Linear Regression*
Let $[n] := \{1, \dots, n\}$, $(Y_i, \xi_i)_{i\in[n]}$ be independent copies of $(Y, \xi)$, $\xi \in \mathbb{R}^p$, $\beta \in \mathbb{R}^p$, and $\sigma^2 > 0$.
A *Random-Design Linear Regression* is an iid sample $(Y_i, \xi_i)_{i\in[n]}$ satisfying $\mathbb{E}(Y \mid \xi = x) = x^t\beta$ and $\mathrm{Var}(Y \mid \xi = x) = \sigma^2$ for $x \in \mathrm{supp}(\xi)$.
*Structural Commentary*: Conditioning on $X_{i\cdot} = \xi_i^t$ gives $Y_{1:n} = X\beta + e$ with $e_i = Y_i - \xi_i^t\beta$.

**Remark 5.5.2.** *Unconditional Variability Can Differ*
Under random design, if $\mathbb{E}(\widehat{\beta} \mid X) = \beta$, then $\mathbb{E}\widehat{\beta} = \beta$.
Moreover, $\mathrm{Var}(\widehat{\beta}) = \mathbb{E}[\mathrm{Var}(\widehat{\beta} \mid X)] + \mathrm{Var}(\mathbb{E}[\widehat{\beta} \mid X]) = \sigma^2\mathbb{E}[(X^tX)^{-1}]$, not generally $\sigma^2(X^tX)^{-1}$ for frozen design.

**Method 5.5.3.** *Pairs Bootstrap for Random-Design Regression*
Let $D := \{(Y_i, x_i) : i \in [n]\}$, $x_i \in \mathbb{R}^p$, and let $T(D)$ be a fitted coefficient, prediction, or smooth statistic.

(i) Draw $D_b^* := \{(Y_{i_m}, x_{i_m}) : m \in [n]\}$ with $i_1, \dots, i_n$ sampled iid from $[n]$.
(ii) Fit least squares on $D_b^*$ and compute $T_b^* := T(D_b^*)$.
(iii) Repeat for $b \in [B]$ and estimate $\mathrm{se}(T(D))$ by $\widehat{\mathrm{se}}_*(T) := \left(\frac{1}{B-1}\sum_{b=1}^{B}(T_b^* - \overline{T}^*)^2\right)^{1/2}$, where $\overline{T}^* := B^{-1}\sum_{b=1}^{B} T_b^*$.

**Perspective 5.5.4.** *Extensions Beyond Ordinary Least Squares*

(i) *Robust regression*: Replace $\sum_i \widehat{e}_i^2$ by $\sum_i \Psi(\widehat{e}_i)$; $\Psi(u) = u^2$ gives least squares.
(ii) *Variable selection*: For selected predictors $S \subseteq \{1, \dots, p\}$, $\widehat{\beta}_S$ and $\mathrm{se}(\widehat{\beta}_S)$ are conditional on $S$.
(iii) *Influence*: Large leverage $h_{ii} = x_i^t(X^tX)^{-1}x_i$ can make observation $i$ dominate $\widehat{\beta}$.
(iv) *Inverse calibration*: Given a fitted response curve $\widehat{m}$ and measured response $y_0$, estimating $x_0$ from $\widehat{m}(x_0) = y_0$ is an inverse problem with uncertainty beyond $\mathrm{se}(\widehat{\beta})$.

# Part XXI
# Statistical Learning and Deep Learning

Statistical learning studies prediction rules, representation maps, and trained systems as mathematical objects. The statistical foundations of Parts 15 and 16 remain the language for experiments, inference, and model assessment; here the emphasis is on risk, generalization, optimization, architectures, and evaluation of learned predictors.

## 1 Learning Problems and Risk

### 1.1 Prediction, Loss Functions, and Risk

A learning problem turns observations into an action rule. Unlike inferential problems, the primary object is often the future loss of a predictor rather than a confidence statement about a parameter.

**Definition 1.1.1.** *Prediction Problem and Risk*
Inputs lie in $(\mathcal{X}, \mathcal{A})$, outputs in $(\mathcal{Y}, \mathcal{B})$, and $(X, Y) \sim P$ on $\mathcal{X} \times \mathcal{Y}$. Let $(\mathcal{A}_0, \mathcal{C})$ be an action space.

(i) *Predictor*: A predictor is a measurable map $f : \mathcal{X} \to \mathcal{A}_0$.
(ii) *Loss and risk*: For loss $\ell : \mathcal{A}_0 \times \mathcal{Y} \to [0, \infty]$, define $R(f) := \mathbb{E}_P \ell(f(X), Y)$.
(iii) *Bayes risk*: $R^* := \inf_f R(f)$, over measurable $f : \mathcal{X} \to \mathcal{A}_0$.

**Definition 1.1.2.** *Supervised Learning Terminology*
Let $S := \{(x_i, y_i) : 1 \le i \le n\}$ with $x_i = (x_{i1}, \dots, x_{ip})^\top \in \mathcal{X}$ and $y_i \in \mathcal{Y}$.

(i) *Features*: Coordinates $x_{ij}$ are variables, inputs, predictors, independent variables, or features.
(ii) *Response*: Values $y_i$ are outputs, responses, targets, or dependent variables.
(iii) *Regression*: If $\mathcal{Y} \subseteq \mathbb{R}^q$, the task is quantitative prediction.
(iv) *Classification*: If $\mathcal{Y} = \mathcal{G}$ is a finite unordered label set, the task is classification.
(v) *Ordered labels*: If $\mathcal{Y}$ is ordered categorical, order matters but arithmetic on labels usually does not.
(vi) *One-hot coding*: For $|\mathcal{G}| = K$, encode class $g$ by $t_g \in \{0, 1\}^K$ with $(t_g)_g = 1$.
Classify vector scores $s(x) \in \mathbb{R}^K$ by $\operatorname{argmax}_k s_k(x)$.

If $y_i$ are unobserved, the problem is unsupervised and targets structure in $\{x_i\}_{i=1}^n$.

**Example 1.1.3.** *Representative Learning Problems*

(i) *Email filtering*: $x$ contains word and punctuation frequencies, $y \in \{\text{spam}, \text{ham}\}$, and asymmetric loss can penalize false deletion more than missed spam.
(ii) *Clinical regression*: $y = \log(\text{PSA})$ and $x$ contains log cancer volume, prostate weight, age, and related measurements; correlated predictors are common.
(iii) *Digit recognition*: $x \in [0, 1]^{16 \times 16}$ or $[0, 1]^{256}$ and $y \in \{0, \dots, 9\}$; action set may add rejection option $\perp$.
(iv) *Microarrays*: An expression matrix $X \in \mathbb{R}^{G \times n}$ has genes as rows and samples as columns.
Typical first tasks are unsupervised clustering of genes or samples.

**Example 1.1.4.** *Common Prediction Losses*

(i) Regression commonly uses squared loss $\ell(a, y) = \|a - y\|_2^2$ or absolute loss $\ell(a, y) = |a - y|$.
(ii) Binary classification with labels in $\{-1, 1\}$ uses zero-one loss $\mathbf{1}_{\{ay \le 0\}}$ for a signed score $a$.
(iii) Probabilistic classification uses log loss $-\log q_y$ when $q$ is a predicted distribution on labels.

**Definition 1.1.5.** *Bayes Predictors*

(i) *Conditional risk*: For action $a$, set $r_x(a) := \mathbb{E}[\ell(a, Y) \mid X = x]$; any $f^*(x) \in \operatorname{argmin}_a r_x(a)$ is Bayes.
(ii) *Squared loss*: If $\ell(a, y) = (a - y)^2$ and $\mathbb{E}[|Y| \mid X = x] < \infty$, then $f^*(x) = \mathbb{E}[Y \mid X = x]$.
(iii) *Absolute loss*: If $\ell(a, y) = |a - y|$, any conditional median $m(x)$
with $\mathbb{P}(Y \le m(x) \mid X = x) \ge \frac{1}{2}$ and $\mathbb{P}(Y \ge m(x) \mid X = x) \ge \frac{1}{2}$ is Bayes.
(iv) *Zero-one loss*: For labels $\mathcal{G}$ and $\ell(g, y) = \mathbf{1}_{\{g \ne y\}}$, a Bayes classifier is $g^*(x) \in \operatorname{argmax}_{g \in \mathcal{G}} \mathbb{P}(Y = g \mid X = x)$.

**Remark 1.1.6.** *Risk as a Prediction Target*
Here risk means predictive loss $R(f) = \mathbb{E}_P \ell(f(X), Y)$ under a data law $P$.
Decision-theoretic risk instead studies $R(\theta, \delta) = \mathbb{E}_\theta L(\theta, \delta(X))$ over a model family.

## 1.2 Empirical Risk Minimization

The population risk is the target, but only a sample is observed. Empirical risk minimisation replaces the unknown expectation by an average over the training data.

**Definition 1.2.1.** *Empirical Risk Minimization*
Let $S := ((x_1, y_1), \dots, (x_n, y_n))$.

(i) *Empirical risk*: For loss $\ell$, define $\widehat{R}_S(f) := n^{-1} \sum_{i=1}^{n} \ell(f(x_i), y_i)$.
(ii) *Hypothesis class*: Let $\mathcal{F}$ be the admissible predictor class.
(iii) *ERM*: Any $\widehat{f} \in \operatorname{argmin}_{f \in \mathcal{F}} \widehat{R}_S(f)$ is an empirical risk minimiser.

**Definition 1.2.2.** *Regularised Empirical Risk*
Let $\Omega : \mathcal{F} \to [0, \infty]$ be a complexity penalty and $\lambda \geq 0$.
Define $Q_\lambda(f) := \widehat{R}_S(f) + \lambda \Omega(f)$ and choose $\widehat{f}_\lambda \in \operatorname{argmin}_{f \in \mathcal{F}} Q_\lambda(f)$.
The penalty $\Omega$ may encode smoothness, sparsity, norm control, margin size, or architecture.

**Theorem 1.2.3.** *Finite-Class ERM Bound*
*Let $S = ((X_1, Y_1), \dots, (X_n, Y_n))$ be i.i.d. from $P$, let $\emptyset \neq \mathcal{F}$ be finite.*
*Assume $0 \leq \ell(f(X), Y) \leq 1$ for all $f \in \mathcal{F}$, and fix $\delta \in (0, 1)$. Set $\varepsilon_{n,\delta} := \sqrt{\log(2 |\mathcal{F}| / \delta)/(2n)}$.*

*(i) Uniform deviation: With probability at least $1 - \delta$, $\sup_{f \in \mathcal{F}} \left| R(f) - \widehat{R}_S(f) \right| \leq \varepsilon_{n,\delta}$.*

*(ii) ERM excess risk: If $\widehat{f} \in \operatorname{argmin}_{f \in \mathcal{F}} \widehat{R}_S(f)$, then $R(\widehat{f}) \leq \inf_{f \in \mathcal{F}} R(f) + 2\varepsilon_{n,\delta}$.*

*Proof.*

(i) For fixed $f$, set $Z_i(f) := \ell(f(X_i), Y_i) \in [0, 1]$, so $R(f) = \mathbb{E} Z_i(f)$ and $\widehat{R}_S(f) = n^{-1} \sum_{i=1}^{n} Z_i(f)$.
Hoeffding and a union bound give $\mathbb{P}(\sup_{f \in \mathcal{F}} \left| \widehat{R}_S(f) - R(f) \right| > \varepsilon) \leq 2 |\mathcal{F}| e^{-2n\varepsilon^2}$; set $\varepsilon = \varepsilon_{n,\delta}$.
(ii) On the uniform-deviation event, choose $f^* \in \operatorname{argmin}_{f \in \mathcal{F}} R(f)$.
Since $\widehat{R}_S(\widehat{f}) \leq \widehat{R}_S(f^*)$, $R(\widehat{f}) \leq \widehat{R}_S(\widehat{f}) + \varepsilon_{n,\delta} \leq \widehat{R}_S(f^*) + \varepsilon_{n,\delta} \leq R(f^*) + 2\varepsilon_{n,\delta}$.

*Structural Commentary*: Union-bounded concentration gives uniform control; ERM turns it into risk control.
*Key Insight*: Control every gap $\left| R - \widehat{R}_S \right|$ by $\varepsilon_{n,\delta}$; the ERM comparison pays this gap twice. □

**Remark 1.2.4.** *Statistical and Computational Error*
In practice the selected predictor may be only an approximate minimizer of the empirical criterion.
Its excess risk is then decomposed into approximation error, estimation error, and optimisation error.

## 1.3 Surrogate Losses and Calibration

Many prediction losses are discontinuous or combinatorial. Learning algorithms often optimise a smoother surrogate and rely on calibration to connect that surrogate to the target loss.

**Definition 1.3.1.** *Surrogate Loss*
For binary labels $y \in \{-1, 1\}$, let $g : \mathcal{X} \to \mathbb{R}$ be score function. Define $m_g(x, y) := yg(x)$ and $\widehat{y}_g(x) := \operatorname{sign}(g(x))$.

(i) *Zero-one loss*: $\ell_{01}(g(x), y) := \mathbf{1}_{\{m_g(x,y) \leq 0\}}$.
(ii) *Surrogate*: $\ell_\phi(g(x), y) := \phi(m_g(x, y))$ for tractable $\phi : \mathbb{R} \to [0, \infty)$.

**Example 1.3.2.** *Margin Surrogates*

(i) Hinge loss: $\phi(t) = \max\{0, 1 - t\}$.
(ii) Logistic loss: $\phi(t) = \log(1 + \exp(-t))$.
(iii) Exponential loss: $\phi(t) = \exp(-t)$.
(iv) Squared margin loss: $\phi(t) = (1 - t)^2$.

**Definition 1.3.3.** *Classification Calibration*
Let $R_\phi(g) := \mathbb{E}\phi(Yg(X))$, $R_{01}(g) := \mathbb{P}(Yg(X) \leq 0)$, $R_\phi^* := \inf_g R_\phi(g)$, and $R_{01}^* := \inf_g R_{01}(g)$.
Margin surrogate $\phi$ is classification calibrated if $R_\phi(g_m) \to R_\phi^*$ implies $R_{01}(g_m) \to R_{01}^*$ $\forall$ score sequence $(g_m)$.
Equivalently, $R_\phi(g) - R_\phi^* \to 0$ forces $R_{01}(g) - R_{01}^* \to 0$.

**Remark 1.3.4.** *Proper Scoring Rules*
A scoring rule $S(q, y)$ is proper if $\operatorname{argmin}_q \mathbb{E}[S(q, Y) \mid X = x] = \mathcal{L}(Y \mid X = x)$.
Thus log and Brier losses learn calibrated conditional probabilities.

## 1.4 Bias-Variance and Approximation-Estimation Tradeoffs

Learning quality depends both on the richness of the model class and on the variability introduced by fitting that class from a finite sample.

**Proposition 1.4.1.** *Bias-Variance Decomposition*
*Let* $Y = f_0(X) + \varepsilon$ *with* $\mathbb{E}[\varepsilon \mid X = x] = 0$ *and* $\mathrm{Var}(\varepsilon \mid X = x) = \sigma^2(x)$.
*Let random* $S$ *produce* $\widehat{f}_S$, *set* $\bar{f}(x) := \mathbb{E}_S \widehat{f}_S(x)$, *and let* $Y_x \perp S$ *be a fresh response at* $x$.
*Then* $\mathbb{E}_{S,Y_x}[(Y_x - \widehat{f}_S(x))^2] = \sigma^2(x) + (f_0(x) - \bar{f}(x))^2 + \mathrm{Var}_S(\widehat{f}_S(x))$.

*Proof.* Write $Y_x - \widehat{f}_S(x) = \varepsilon_x + f_0(x) - \widehat{f}_S(x)$, where $\varepsilon_x := Y_x - f_0(x)$.
Since $\mathbb{E}[\varepsilon_x] = 0$ and $Y_x$ is conditionally independent of $S$, $\mathbb{E}_{S,Y_x}[\varepsilon_x\{f_0(x) - \widehat{f}_S(x)\}] = 0$.
Thus $\mathbb{E}_{S,Y_x}[(Y_x - \widehat{f}_S(x))^2] = \sigma^2(x) + \mathbb{E}_S[(f_0(x) - \widehat{f}_S(x))^2]$.
Finally, $\mathbb{E}_S[(f_0(x) - \widehat{f}_S(x))^2] = (f_0(x) - \bar{f}(x))^2 + \mathbb{E}_S[(\widehat{f}_S(x) - \bar{f}(x))^2]$.
*Structural Commentary*: Squared prediction error equals noise plus fitted-function error.
*Key Insight*: Center $\widehat{f}_S(x)$ at $\bar{f}(x)$; the remaining cross term has mean zero. □




Figure 253: Bias–variance tradeoff: blue: training error, red: test error; dashed line marks test minimum.

**Method 1.4.2.** *Nearest-Neighbor Local Averaging*
Let $\mathcal{T} := \{(x_i, y_i) : 1 \le i \le n\}$ be training data in a metric space $(\mathcal{X}, d)$.
Let $N_k(x) \subseteq \{1, \dots, n\}$ be the indices of the $k$ smallest distances $d(x_i, x)$.
For numerical responses, define the local average $\widehat{f}_k(x) := \frac{1}{k} \sum_{i \in N_k(x)} y_i$.
For class labels $g_i \in \mathcal{G}$, define $\widehat{p}_g(x) := \frac{1}{k} \sum_{i \in N_k(x)} \mathbf{1}_{\{g_i = g\}}$ and classify by $\widehat{g}_k(x) \in \operatorname{argmax}_{g \in \mathcal{G}} \widehat{p}_g(x)$.
Thus $k$-nearest neighbours replace conditioning at $X = x$ by averaging over $N_k(x)$.

**Remark 1.4.3.** *Locality in High Dimension*
The parameter $k$ trades local bias against variance: small $k$ reacts strongly to $S$, while large $k$ averages more but may wash out local structure. In $[0,1]^p$, a subcube with volume fraction $r$ has side length $e_p(r) = r^{1/p}$, so $e_{10}(0.01) \approx 0.63$ and $e_{10}(0.10) \approx 0.80$. For $N$ uniform points in the unit ball, the median nearest-neighbour radius is $d(p, N) = (1 - 2^{-1/N})^{1/p}$, about 0.52 for $(p, N) = (10, 500)$. Thus high-dimensional neighbourhoods large enough to contain data may not be local; useful local methods need structure or an adapted metric.

**Definition 1.4.4.** *Approximation and Estimation Errors*
Let $R^* := \inf_f R(f)$ and $R^*_{\mathcal{F}} := \inf_{f \in \mathcal{F}} R(f)$.

(i) *Approximation error*: $\mathcal{A}(\mathcal{F}) := R^*_{\mathcal{F}} - R^*$.
(ii) *Estimation error*: For a data-dependent $\widehat{f} \in \mathcal{F}$, $\mathcal{E}(\widehat{f}; \mathcal{F}) := R(\widehat{f}) - R^*_{\mathcal{F}}$.
(iii) *Excess risk*: $R(\widehat{f}) - R^* = \mathcal{A}(\mathcal{F}) + \mathcal{E}(\widehat{f}; \mathcal{F})$.

Larger $\mathcal{F}$ usually lowers $\mathcal{A}(\mathcal{F})$ and raises estimation difficulty.

**Remark 1.4.5.** *Model Complexity*
As $c(\mathcal{F})$ grows, $\widehat{R}_S(\widehat{f}_{\mathcal{F}})$ usually decreases, while $R(\widehat{f}_{\mathcal{F}}) - R^* = \mathcal{A}(\mathcal{F}) + \mathcal{E}(\widehat{f}_{\mathcal{F}}; \mathcal{F})$ trades smaller approximation error for larger estimation error. For zero-one loss, the relevant failure is crossing the wrong side of the decision boundary, not an additive squared-error bias–variance split.

# 2 Generalization Theory

## 2.1 Uniform Convergence and VC Theory

Generalisation asks when small training loss implies small population loss. Uniform convergence controls the gap for all predictors in a class at once.

**Definition 2.1.1.** *Uniform Convergence*
For samples $S_n$ of size $n$ and bounded loss, $\mathcal{F}$ has uniform convergence if $\sup_{f\in\mathcal{F}}\left|R(f)-\widehat{R}_{S_n}(f)\right|\to 0$ in probability, or almost surely, as $n\to\infty$.

**Definition 2.1.2.** *Shattering and VC Dimension*
For $\mathcal{C}\subseteq 2^{\mathcal{X}}$, a finite $A\subseteq\mathcal{X}$ is shattered if $\{A\cap C : C\in\mathcal{C}\}=2^A$, and $\mathrm{VCdim}(\mathcal{C})$ is the supremum of $|A|$ over finite shattered $A\subseteq\mathcal{X}$.

**Example 2.1.3.** *Shattering Examples*
For $\mathcal{H}_p:=\{\{x\in\mathbb{R}^p : a^\top x+b\ge 0\} : a\in\mathbb{R}^p, b\in\mathbb{R}\}$, $\mathrm{VCdim}(\mathcal{H}_p)=p+1$; in particular, $\mathcal{H}_2$ shatters any three noncollinear points and shatters no four-point set.
Parameter count not decisive: one-parameter class $\mathcal{C}:=\{\{x\in\mathbb{R}:\sin(\alpha x)>0\}:\alpha\in\mathbb{R}\}$ has infinite VCdim.

**Theorem 2.1.4.** *VC Generalization Bound Form*
*Let $\mathcal{F}\subseteq\{0,1\}^{\mathcal{X}}$ have $1\le d:=\mathrm{VCdim}(\mathcal{F})<\infty$, let $S_n=((X_i,Y_i))_{i=1}^n$ be i.i.d., and use $\ell_f(x,y):=\mathbf{1}_{\{f(x)\neq y\}}$, $R(f):=\mathbb{E}\ell_f(X,Y)$, and $\widehat{R}_{S_n}(f):=\frac{1}{n}\sum_{i=1}^n\ell_f(X_i,Y_i)$.*
*Define $\Pi_{\mathcal{F}}(m):=\max|\{(f(x_1),\dots,f(x_m)) : f\in\mathcal{F}\}|$, where the maximum is over $x_1,\dots,x_m\in\mathcal{X}$.*

*(i) Growth bound: If $m\ge d\ge 1$, then $\Pi_{\mathcal{F}}(m)\le\sum_{j=0}^d\binom{m}{j}\le\left(\frac{em}{d}\right)^d$.*

*(ii) Uniform deviation: For every $\delta\in(0,1)$ and $n\ge d$, with probability at least $1-\delta$, $\sup_{f\in\mathcal{F}}\left|R(f)-\widehat{R}_{S_n}(f)\right|\le C\sqrt{\frac{d\log\left(\frac{2en}{d}\right)+\log\left(\frac{1}{\delta}\right)}{n}}$ for a universal constant $C$.*

*Proof.*

(i) *Growth bound*: Since no set of size $d+1$ is shattered, Sauer's lemma gives $\Pi_{\mathcal{F}}(m)\le\sum_{j=0}^d\binom{m}{j}$, and the binomial-sum bound gives $\sum_{j=0}^d\binom{m}{j}\le\left(\frac{em}{d}\right)^d$ for $m\ge d$.

(ii) *Uniform deviation*: Symmetrization with an independent ghost sample and Hoeffding's inequality on the at most $\Pi_{\mathcal{F}}(2n)$ realized labelings give $\mathbb{P}(\sup_{f\in\mathcal{F}}\left|R(f)-\widehat{R}_{S_n}(f)\right|>\varepsilon)\le 8\Pi_{\mathcal{F}}(2n)\exp(-cn\varepsilon^2)$ for a universal $c>0$; applying (i) with $m=2n$ and solving the right side at level $\delta$ yields (ii).

*Structural Commentary*: VC dimension polynomially bounds sample labelings.
*Key Insight*: In finite-class concentration, use $|\mathcal{F}|\rightsquigarrow\Pi_{\mathcal{F}}(2n)\le\left(\frac{2en}{d}\right)^d$. □

**Remark 2.1.5.** *Growth Functions*
For $d=\mathrm{VCdim}(\mathcal{F})$, $\Pi_{\mathcal{F}}(m)$ is the effective number of labelings on $m$ sample points. Sauer's lemma gives $\log\Pi_{\mathcal{F}}(m)\le d\log\left(\frac{em}{d}\right)$, so VC bounds replace $\log|\mathcal{F}|$ by the sample-level complexity $\log\Pi_{\mathcal{F}}(m)$.

**Method 2.1.6.** *Structural Risk Minimization*

(i) *Model ladder*: Choose $\mathcal{F}_1\subseteq\mathcal{F}_2\subseteq\cdots$ with complexities $c_m$, such as $c_m=\mathrm{VCdim}(\mathcal{F}_m)$.
(ii) *Uniform penalties*: Choose $\delta_m>0$ with $\sum_m\delta_m\le\delta$ and penalties $\psi_m(n,\delta_m)$ such that $\mathbb{P}(\sup_{f\in\mathcal{F}_m}\left|R(f)-\widehat{R}_{S_n}(f)\right|\le\psi_m(n,\delta_m))\ge 1-\delta_m$.
(iii) *SRM rule*: Choose $(\widehat{m},\widehat{f})\in\operatorname{argmin}_{m\in\mathbb{N},f\in\mathcal{F}_m}\{\widehat{R}_{S_n}(f)+\psi_m(n,\delta_m)\}$.

Larger $m$ improves fit but raises $\psi_m$; SRM balances flexibility against search cost.

## 2.2 Rademacher Complexity and Covering Numbers

For real-valued predictors, complexity is often measured by how well the class can fit random noise.

**Definition 2.2.1.** *Rademacher Complexity*
Let $\mathcal{F}\subseteq\mathbb{R}^{\mathcal{X}}$, fix $S_n=(x_1,\dots,x_n)\in\mathcal{X}^n$, and let $\sigma_1,\dots,\sigma_n$ be i.i.d. Rademacher signs, $\mathbb{P}(\sigma_i=1)=\mathbb{P}(\sigma_i=-1)=\frac{1}{2}$. The empirical Rademacher complexity is $\widehat{\mathfrak{R}}_{S_n}(\mathcal{F}):=\mathbb{E}_\sigma\sup_{f\in\mathcal{F}}\frac{1}{n}\sum_{i=1}^n\sigma_i f(x_i)$.
If $X_1,\dots,X_n\sim P_X$ i.i.d., the population complexity is $\mathfrak{R}_n(\mathcal{F}):=\mathbb{E}_{S_n}\widehat{\mathfrak{R}}_{S_n}(\mathcal{F})$.

**Definition 2.2.2.** *Covering Number*
For a metric space $(T, d)$ and $\varepsilon > 0$, $N(\varepsilon, T, d) := \inf\{m : \exists t_1, \dots, t_m \in T,\ T \subseteq \bigcup_{j=1}^{m} \overline{B}_d(t_j, \varepsilon)\}$.
The entropy $\log N(\varepsilon, T, d)$ measures metric size at scale $\varepsilon$, not parameter count.

**Remark 2.2.3.** *Entropy Methods*
Bounds on $\log N(\varepsilon, \mathcal{F}, d)$ can be integrated over scales to control empirical processes.
This is the route from functional-analytic compactness to statistical-learning rates.

## 2.3 Concentration and Stability

Generalization can also follow from concentration of the training criterion or from stability of the learning algorithm itself.

**Definition 2.3.1.** *Subgaussian Random Variable*
Centered random variable $Z$ is *subgaussian* with variance proxy $\sigma^2$ if $\mathbb{E}\exp(\lambda Z) \le \exp(\lambda^2\sigma^2/2)\ \forall \lambda \in \mathbb{R}$.
Subgaussian variables have Gaussian-type tail bounds.

**Theorem 2.3.2.** *Bounded Differences Inequality*
*Let $X_1, \dots, X_n$ be independent, $X := (X_1, \dots, X_n)$, and $F : \prod_{i=1}^{n} \mathcal{X}_i \to \mathbb{R}$.*
*For $x = (x_1, \dots, x_n)$, write $x^{(i \leftarrow x_i')} := (x_1, \dots, x_{i-1}, x_i', x_{i+1}, \dots, x_n)$.*
*Assume $\left|F(x) - F(x^{(i \leftarrow x_i')})\right| \le c_i$ for all $i, x, x_i'$, and set $v := \sum_{i=1}^{n} c_i^2$.*
*Then, for every $t > 0$, $\mathbb{P}\{F(X) - \mathbb{E}F(X) \ge t\} \le \exp\left(-\frac{2t^2}{v}\right)$ and $\mathbb{P}\{|F(X) - \mathbb{E}F(X)| \ge t\} \le 2\exp\left(-\frac{2t^2}{v}\right)$.*

*Proof.* Let $\mathcal{G}_i := \sigma(X_1, \dots, X_i)$, $M_i := \mathbb{E}(F(X) \mid \mathcal{G}_i)$, and $\Delta_i := M_i - M_{i-1}$.
Then $M_0 = \mathbb{E}F(X)$, $M_n = F(X)$, and the bounded-differences hypothesis implies that, conditionally on $\mathcal{G}_{i-1}$, $\Delta_i$ has mean 0 and lies in an interval of length at most $c_i$. By Hoeffding's lemma, $\mathbb{E}(\exp(\lambda\Delta_i) \mid \mathcal{G}_{i-1}) \le \exp\left(\frac{\lambda^2 c_i^2}{8}\right)$ for $\lambda > 0$. Iterating, $\mathbb{E}\exp(\lambda(F(X) - \mathbb{E}F(X))) \le \exp\left(\frac{\lambda^2 v}{8}\right)$.
By Chernoff's bound, $\mathbb{P}\{F(X) - \mathbb{E}F(X) \ge t\} \le \exp\left(-\lambda t + \frac{\lambda^2 v}{8}\right)$, optimised at $\lambda = \frac{4t}{v}$.
Applying the same bound to $-F$ and taking a union bound gives the two-sided inequality.
*Structural Commentary*: Coordinate sensitivity plus independence makes $F(X)$ subgaussian.
*Key Insight*: Doob increments reduce global concentration to bounds on one-coordinate changes $c_i$. □

**Theorem 2.3.3.** *Rademacher Generalization Bound*
*Let $Z_i = (X_i, Y_i) \in \mathcal{Z} := \mathcal{X} \times \mathcal{Y}$ be i.i.d. from $P$.*
*Let $S_n = (Z_1, \dots, Z_n)$, $\mathcal{G} := \{g_f : \mathcal{Z} \to [0,1] : g_f(x,y) = \ell(f(x), y), f \in \mathcal{F}\}$.*
*For $g \in \mathcal{G}$, write $Pg := \mathbb{E}g(Z)$ and $P_n g := \frac{1}{n}\sum_{i=1}^{n} g(Z_i)$.*

*(i) Uniform deviation: $\forall \delta \in (0,1)$, with probability at least $1-\delta$, $\sup_{g \in \mathcal{G}}(Pg - P_n g) \le 2\widehat{\mathfrak{R}}_{S_n}(\mathcal{G}) + 3\sqrt{\frac{\log\left(\frac{2}{\delta}\right)}{2n}}$.*

*(ii) Risk bound: Hence, simultaneously for all $f \in \mathcal{F}$, $R(f) \le \widehat{R}_{S_n}(f) + 2\widehat{\mathfrak{R}}_{S_n}(\ell \circ \mathcal{F}) + 3\sqrt{\frac{\log\left(\frac{2}{\delta}\right)}{2n}}$.*

*Proof.*
(i) Let $\Phi(S_n) := \sup_{g \in \mathcal{G}}(Pg - P_n g)$. Changing one $Z_i$ changes $\Phi$ and $\widehat{\mathfrak{R}}_{S_n}(\mathcal{G})$ by at most $1/n$, so McDiarmid gives $\Phi \le \mathbb{E}\Phi + \varepsilon$ and $\mathfrak{R}_n(\mathcal{G}) \le \widehat{\mathfrak{R}}_{S_n}(\mathcal{G}) + \varepsilon$ with probability at least $1-\delta$, for $\varepsilon := \sqrt{\frac{\log\left(\frac{2}{\delta}\right)}{2n}}$.
Symmetrisation gives $\mathbb{E}\Phi \le 2\mathfrak{R}_n(\mathcal{G})$, hence $\Phi \le 2\widehat{\mathfrak{R}}_{S_n}(\mathcal{G}) + 3\varepsilon$.
(ii) For $z = (x, y)$ and $g_f(z) = \ell(f(x), y)$, $Pg_f = R(f)$ and $P_n g_f = \widehat{R}_{S_n}(f)$.
Applying (i) to every $g_f \in \mathcal{G}$ gives the stated simultaneous bound.
*Structural Commentary*: Rademacher complexity measures sample-wise noise fitting.
*Key Insight*: Symmetrization turns $P - P_n$ into $\mathbb{E}_\sigma \sup_{g \in \mathcal{G}} \frac{1}{n}\sum_{i=1}^{n} \sigma_i g(Z_i)$. □

**Definition 2.3.4.** *Uniform Stability*
A learning algorithm $A$ has *uniform stability* $\beta_n$ for loss $\ell$ if, for any samples $S, S' \in \mathcal{Z}^n$ differing in one coordinate, $\sup_{z \in \mathcal{Z}} |\ell(A(S), z) - \ell(A(S'), z)| \le \beta_n$.

**Remark 2.3.5.** *Algorithm-Dependent Generalisation*
Stability controls generalisation through the map $S \mapsto A(S)$, not only through $\mathcal{F}$.
Thus regularisation or optimisation can generalise well even when the ambient class is large.

## 2.4 PAC-Bayes and Information-Theoretic Bounds

PAC-Bayes bounds analyze randomized predictors by comparing a data-dependent posterior distribution on hypotheses to a fixed prior.

**Definition 2.4.1.** *Randomised Predictor*
Let $\mathcal{H}$ be a hypothesis space and $Q \in \mathcal{P}(\mathcal{H})$. The *randomised predictor* $Q$ samples $h \sim Q$ before prediction, with risk $R(Q) := \mathbb{E}_{h\sim Q} R(h)$ and empirical risk $\widehat{R}_S(Q) := \mathbb{E}_{h\sim Q}\widehat{R}_S(h)$.

**Theorem 2.4.2.** *PAC-Bayes Bound Form*
*Let $S_n = (Z_1, \dots, Z_n)$ be i.i.d., let $\ell(h, Z) \in [0, 1]$, and let $\Pi \in \mathcal{P}(\mathcal{H})$ be independent of $S_n$.*
*For $Q \ll \Pi$, set $\mathrm{KL}(Q\|\Pi) := \mathbb{E}_Q \log \frac{dQ}{d\Pi}$ and $\mathrm{kl}(a, b) := a \log \frac{a}{b} + (1 - a) \log \frac{1-a}{1-b}$.*

*(i) With probability at least $1 - \delta$, every posterior $Q \ll \Pi$ satisfies $\mathrm{kl}(\widehat{R}_{S_n}(Q), R(Q)) \le \frac{\mathrm{KL}(Q\|\Pi)+\log\left(\frac{n+1}{\delta}\right)}{n}$.*

*(ii) On the same event, $R(Q) \le \widehat{R}_{S_n}(Q) + \sqrt{\frac{\mathrm{KL}(Q\|\Pi)+\log\left(\frac{n+1}{\delta}\right)}{2n}}$ for every $Q \ll \Pi$.*

*Proof.*

(i) For fixed $h$, the bounded-loss kl moment bound gives $\mathbb{E}_{S_n} \exp(n\,\mathrm{kl}(\widehat{R}_{S_n}(h), R(h))) \le n + 1$.
Thus $\mathbb{E}_{S_n}\mathbb{E}_{h\sim\Pi} \exp(n\,\mathrm{kl}(\widehat{R}_{S_n}(h), R(h))) \le n + 1$, so Markov gives,
with probability at least $1 - \delta$, $\mathbb{E}_\Pi \exp(n\,\mathrm{kl}(\widehat{R}_{S_n}(h), R(h))) \le \frac{n+1}{\delta}$.
On this event, the Donsker-Varadhan inequality with $\varphi(h) := n\,\mathrm{kl}(\widehat{R}_{S_n}(h), R(h))$ gives
$\mathbb{E}_Q\varphi \le \mathrm{KL}(Q\|\Pi) + \log \mathbb{E}_\Pi \exp\varphi \ \forall Q \ll \Pi$. By convexity of kl, $n\,\mathrm{kl}(\widehat{R}_{S_n}(Q), R(Q)) \le \mathbb{E}_Q\varphi$, proving (i).

(ii) If $R(Q) \ge \widehat{R}_{S_n}(Q)$, Pinsker gives $\mathrm{kl}(\widehat{R}_{S_n}(Q), R(Q)) \ge 2(R(Q) - \widehat{R}_{S_n}(Q))^2$.
if $R(Q) < \widehat{R}_{S_n}(Q)$, the claim is immediate. Combining with (i) gives (ii).

*Structural Commentary*: PAC-Bayes replaces class size by posterior-prior divergence.
*Key Insight*: A prior-average exponential tail plus $Q \ll \Pi$ yields all posteriors at cost $\mathrm{KL}(Q\|\Pi)$. □

**Remark 2.4.3.** *Prior and Posterior*
In PAC-Bayes, $\Pi$ need not be subjective; it anchors the complexity term $\mathrm{KL}(Q\|\Pi)$.
Smaller $\mathrm{KL}(Q\|\Pi)$ means the data-dependent posterior $Q$ stays closer to the fixed prior.

**Definition 2.4.4.** *Information-Theoretic Generalization*
Let $S_n = (Z_1, \dots, Z_n)$, $W = A(S_n)$ be learned output. *Information-theoretic generalization* bounds control $\mathrm{gen}(A) := \mathbb{E}[R(W) - \widehat{R}_{S_n}(W)]$ by dependence measures such as $I(W; S_n)$.

# 3 Unsupervised and Latent-Variable Learning

## 3.1 Clustering and Mixture Models

Unsupervised learning searches for structure in inputs without direct labels or a direct prediction loss. Clustering summarizes the input distribution by grouping observations, while mixture models replace hard groups by latent component probabilities.

**Definition 3.1.1.** *k-Means Clustering*
Let $x_1, \dots, x_n \in \mathbb{R}^d$ and $k \in \{1, \dots, n\}$. Put $\mathcal{C} = \{C : \{1, \dots, n\} \to \{1, \dots, k\}\}$, $\mathcal{M} = (\mathbb{R}^d)^k$,
$I_r(C) = \{i : C(i) = r\}$, and, for $m = (m_1, \dots, m_k)$, $\Phi(C, m) = \sum_{r=1}^{k} \sum_{i\in I_r(C)} \|x_i - m_r\|_2^2$.

(i) A *k-means solution* is any $(C^\star, m^\star) \in \operatorname*{arg\,min}_{C\in\mathcal{C},\, m\in\mathcal{M}} \Phi(C, m)$.

(ii) Its *clusters* are $I_r^\star = I_r(C^\star)$, $r = 1, \dots, k$.

(iii) For fixed $C$, a *minimizing center* is $m_r = |I_r(C)|^{-1} \sum_{i\in I_r(C)} x_i$ if $I_r(C) \ne \varnothing$, and arbitrary if $I_r(C) = \varnothing$.

(iv) For fixed $m$, a *minimizing assignment* satisfies $C(i) \in \operatorname*{arg\,min}_{1\le r\le k} \|x_i - m_r\|_2^2$ for each $i$.

**Definition 3.1.2.** *Object Dissimilarities*
Let $x_1, \dots, x_n$ be objects and let $d_{ii'} = d(x_i, x_{i'})$; when attributes are used, write $x_i = (x_{i1}, \dots, x_{ip})$.

(i) A *dissimilarity matrix* is $D = (d_{ii'}) \in [0,\infty)^{n\times n}$ with $d_{ii} = 0$ and $d_{ii'} = d_{i'i}$; directional inputs $a_{ii'}$ are symmetrized by $d_{ii'} = s(a_{ii'}, a_{i'i})$, where $s : [0,\infty)^2 \to [0,\infty)$ and $s(u,v) = s(v,u)$.

(ii) Given $d_j$ and $w_j \geq 0$ with $\sum_{j=1}^{p} w_j = 1$, the *additive dissimilarity* is $d(x_i, x_{i'}) = \sum_{j=1}^{p} w_j d_j(x_{ij}, x_{i'j})$.

(iii) *Coordinate rules*: quantitative $d_j(a,b) = |a-b|$ or $(a-b)^2$; ordinal $r \mapsto (r-1/2)/M$ for $r = 1, \dots, M$; categorical $d_j(a,b) = L_j(a,b)$ with $L_j(a,a) = 0$ and $L_j(a,b) \geq 0$.

**Remark 3.1.3.** *Weights and Missing Values*
For $d(x_i, x_{i'}) = \sum_{j=1}^{p} w_j d_j(x_{ij}, x_{i'j})$, the empirical scale of coordinate $j$ is $\bar{d}_j = \binom{n}{2}^{-1} \sum_{i<i'} d_j(x_{ij}, x_{i'j})$, so its effective contribution is $w_j \bar{d}_j$. If $O_{ii'} \subseteq \{1, \dots, p\}$ is the nonempty set of coordinates observed for both $x_i$ and $x_{i'}$, pairwise deletion uses $d(x_i, x_{i'}) = \sum_{j \in O_{ii'}} \tilde{w}_{j,ii'} d_j(x_{ij}, x_{i'j})$ with $\tilde{w}_{j,ii'} = w_j / \sum_{\ell \in O_{ii'}} w_\ell$.
Alternatives are imputed values $\tilde{x}_{ij}$ before computing $d$ or an added categorical level `missing`.

**Definition 3.1.4.** *Within- and Between-Cluster Scatter*
Fix $k \in \{1, \dots, n\}$ and a symmetric dissimilarity matrix $D = (d_{ii'})$.
Let $\mathcal{C}_k = \{C : \{1, \dots, n\} \to \{1, \dots, k\}\}$ and $I_r(C) = \{i : C(i) = r\}$.

(i) The *within-cluster scatter* is $W(C) = \frac{1}{2} \sum_{r=1}^{k} \sum_{i \in I_r(C)} \sum_{i' \in I_r(C)} d_{ii'}$ for $C \in \mathcal{C}_k$.

(ii) The *total scatter* is $T = \frac{1}{2} \sum_{i=1}^{n} \sum_{i'=1}^{n} d_{ii'}$.

(iii) The *between-cluster scatter* is $B(C) = T - W(C)$ for $C \in \mathcal{C}_k$.

As $T$ is fixed, $\arg\min_{C \in \mathcal{C}_k} W(C) = \arg\max_{C \in \mathcal{C}_k} B(C)$.

**Algorithm 2:** $k$-Means Alternating Minimization

**Require:** Data $x_1, \dots, x_n \in \mathbb{R}^d$, number of clusters $k$, initial centers or assignments
**Ensure:** Cluster assignments $C(i)$ and centers $m_1, \dots, m_k$
**repeat**
Given assignments, set each $m_r$ to the mean of the observations with $C(i) = r$
Given centers, assign each observation by $C(i) \leftarrow \arg\min_{1 \leq r \leq k} \|x_i - m_r\|_2^2$
**until** the assignments no longer change

**Proposition 3.1.5.** *k-Means Descent and Local Minima*
*Let* $\mathcal{C} = \{C : \{1, \dots, n\} \to \{1, \dots, k\}\}$, $\mathcal{M} = (\mathbb{R}^d)^k$, *and* $\Phi(C, m) = \sum_{r=1}^{k} \sum_{i : C(i) = r} \|x_i - m_r\|_2^2$.
*Suppose* $(C^t, m^t) \in \mathcal{C} \times \mathcal{M}$ *satisfies* $m^{t+1} \in \arg\min_{m \in \mathcal{M}} \Phi(C^t, m)$ *and* $C^{t+1} \in \arg\min_{C \in \mathcal{C}} \Phi(C, m^{t+1})$ *for each* $t$.

*(i) The objective is nonincreasing:* $\Phi(C^{t+1}, m^{t+1}) \leq \Phi(C^t, m^t)$.
*(ii) If* $C^{t+1} = C^t = C^\star$ *and* $m^{t+1} = m^t = m^\star$, *then* $(C^\star, m^\star)$ *is coordinatewise minimal:* $m^\star \in \arg\min_{m \in \mathcal{M}} \Phi(C^\star, m)$ *and* $C^\star \in \arg\min_{C \in \mathcal{C}} \Phi(C, m^\star)$.
*Coordinatewise minimal need not be global; compare initialisations by terminal* $\Phi$.

*Proof.*

(i) By the center update, $\Phi(C^t, m^{t+1}) \leq \Phi(C^t, m^t)$. By the assignment update, $\Phi(C^{t+1}, m^{t+1}) \leq \Phi(C^t, m^{t+1})$. Combining gives the claim.

(ii) At a fixed point, $C^{t+1} = C^t = C^\star$ and $m^{t+1} = m^t = m^\star$ in the two update inclusions, giving the two coordinatewise optimality conditions.

*Structural Commentary*: Exact block minimization forces descent.
*Key Insight*: A stopped run certifies blockwise, not global, optimality. □

**Algorithm 3:** $k$-Medoids Alternating Representatives

**Require:** Pairwise dissimilarities $D(x_i, x_{i'})$, number of clusters $k$, initial assignments
**Ensure:** Cluster assignments $C(i)$ and representative indices $i_1^\star, \dots, i_k^\star$
**repeat**
**for** $r = 1, \dots, k$ **do**

Set $i_r^\star \leftarrow \arg\min_{i:\, C(i)=r} \sum_{C(i')=r} D(x_i, x_{i'})$
**end for**
Assign each observation by $C(i) \leftarrow \arg\min_{1\le r\le k} D(x_i, x_{i_r^\star})$
**until** the assignments no longer change

**Remark 3.1.6.** *k-Medoids Versus k-Means*
Fix an assignment $C$ and write $I_r = \{i : C(i) = r\}$.

(i) *k-means*: $m_r \in \arg\min_{m\in\mathbb{R}^d} \sum_{i\in I_r} \|x_i - m\|_2^2$, hence $m_r = |I_r|^{-1} \sum_{i\in I_r} x_i$ when $I_r \neq \varnothing$.
(ii) *k-medoids*: $i_r^\star \in \arg\min_{i\in I_r} \sum_{i'\in I_r} D(x_i, x_{i'})$, with representative $x_{i_r^\star}$.

$k$-medoids only requires pairwise dissimilarities and keeps representatives inside the data set.
Its representative update is usually costlier than the $k$-means empirical mean.




Figure 254: Open squares are centroids, circled points are medoids, the dashed line is nearest-center boundary.

**Remark 3.1.7.** *Choosing the Number of Clusters*
For $k$-means or $k$-medoids, choose $k$ and an initialisation. Let $W_k$ be terminal within-cluster scatter.
The elbow diagnostic inspects $k \mapsto W_k$; the gap statistic uses $G(k) = \mathbb{E}_0[\log W_k] - \log W_k$, where $\mathbb{E}_0$ is under a reference distribution on the data range.
With simulation standard error $s'_{k+1}$, one rule is $\widehat{k} = \min\{k : G(k) \ge G(k+1) - s'_{k+1}\}$.
This formalises the bend comparison but remains heuristic.

**Method 3.1.8.** *Hierarchical Clustering and Linkage*
Given dissimilarities $d_{ii'}$ and a linkage $L(G, H)$ between nonempty groups, agglomerative hierarchical clustering constructs nested partitions.

(i) Initialize $\mathcal{P}_0 = \{\{1\}, \dots, \{n\}\}$.
(ii) At step $t$, choose distinct $G, H \in \mathcal{P}_t$ minimizing $L(G, H)$.
(iii) Set $\mathcal{P}_{t+1} = (\mathcal{P}_t \setminus \{G, H\}) \cup \{G \cup H\}$ and record merge height $L(G, H)$.
(iv) Stop at $\mathcal{P}_{n-1} = \{\{1, \dots, n\}\}$.

For groups $G, H$ with sizes $N_G, N_H$, common linkages are

(i) *Single*: $L_{\mathrm{SL}}(G, H) = \min_{i\in G,\, i'\in H} d_{ii'}$.
(ii) *Complete*: $L_{\mathrm{CL}}(G, H) = \max_{i\in G,\, i'\in H} d_{ii'}$.
(iii) *Group average*: $L_{\mathrm{GA}}(G, H) = (N_G N_H)^{-1} \sum_{i\in G} \sum_{i'\in H} d_{ii'}$.

Recorded merges form a dendrogram; divisive clustering reverses the process by recursive splits.

**Remark 3.1.9.** *Dendrograms and Cophenetic Distances*
For dendrogram merge height $h_T(i, i')$, the cut at height $a$ partitions observations by $i \sim_a i'$ iff $h_T(i, i') \le a$.
The cophenetic dissimilarity is $C_{ii'} = h_T(i, i')$, the first height at which $i$ and $i'$ share a cluster.
It is ultrametric: $C_{ii'} \le \max\{C_{ih}, C_{i'h}\}$. Thus a dendrogram is an algorithmic hierarchical summary, not an exact representation of arbitrary input dissimilarities.

**Method 3.1.10.** *Self-Organizing Map*
A self-organizing map uses prototypes $m_1^t, \dots, m_q^t \in \mathbb{R}^p$ indexed by grid locations $\ell_1, \dots, \ell_q \in \mathcal{Q}_1 \times \mathcal{Q}_2$.
Let $\alpha_t > 0$ and $h_t : [0, \infty) \to [0, 1]$ be the learning rate and neighborhood kernel.

(i) Select an observation $x_i$.
(ii) Find a winner $j_t \in \arg\min_{1\le j\le q} \|x_i - m_j^t\|_2^2$.
(iii) Update every prototype by $m_k^{t+1} = m_k^t + \alpha_t h_t(\|\ell_{j_t} - \ell_k\|_2)(x_i - m_k^t)$, $k = 1, \dots, q$.
(iv) Increment $t$ and repeat until the stopping rule or schedule terminates.

A common kernel is $h_t(s) = \mathbf{1}_{s \le r_t}$ with decreasing radius $r_t$.
Early large neighborhoods enforce grid smoothness; if $h_t(s) = \mathbf{1}_{s=0}$, the method reduces to online $k$-means.

**Remark 3.1.11.** *Checking the Topological Constraint*
For $q$ prototypes, compare $E_{\mathrm{SOM}} = \sum_i \|x_i - m_{C(i)}\|_2^2$ with the $q$-means optimum $E_q^\star = \min_{C,m} \sum_i \|x_i - m_{C(i)}\|_2^2$.
Since $E_q^\star \le E_{\mathrm{SOM}}$, a large gap $E_{\mathrm{SOM}} - E_q^\star$ indicates an overly restrictive lattice constraint.

**Method 3.1.12.** *Spectral Clustering*
Given weights $w_{ii'} \ge 0$, set $W = (w_{ii'})$, $g_i = \sum_{i'=1}^n w_{ii'}$, $G = \mathrm{diag}(g_i)$, and $L = G - W$.

Often $w_{ii'} = s_{ii'} = \exp(-\frac{\|x_i - x_{i'}\|_2^2}{c})$ (restricted to mutual nearest-neighbor edges). For target $k$ clusters:

(i) Construct $W$, $G$, and the Laplacian $L$.
(ii) Compute eigenvectors $u_1, \dots, u_k$ for the $k$ smallest nontrivial eigenvalues of $L$.
(iii) Form $U = (u_1, \dots, u_k) \in \mathbb{R}^{n \times k}$ and represent observation $i$ by row $U_{i\cdot}$.
(iv) Cluster the rows $U_{i\cdot}$ into $k$ groups, typically by $k$-means.

$f^T L f = \frac{1}{2} \sum_{i=1}^n \sum_{i'=1}^n w_{ii'}(f_i - f_{i'})^2$, so small-eigenvalue directions are smooth across high-weight edges.

**Remark 3.1.13.** *Connected Components and Random Walks*
Let $L = G - W$ and, when $g_i > 0$, $P = G^{-1}W$.

(i) If the graph has components $A_1, \dots, A_m$, then $\ker L = \mathrm{span}\{\mathbf{1}_{A_1}, \dots, \mathbf{1}_{A_m}\}$.
(ii) The matrix $P$ is a random-walk transition matrix.
A good spectral cluster $A$ has large within-set transition mass $\sum_{i \in A} \sum_{i' \in A} P_{ii'}$ and small cross-set mass.

Nearly separated clusters replace exact zero eigenvalues by small eigenvalues.

**Definition 3.1.14.** *Finite Mixture Model*
Let $Z \in \{1, \dots, k\}$ be latent with $\mathbb{P}(Z = z) = \pi_z$, $\pi_z \ge 0$, and $\sum_{z=1}^k \pi_z = 1$.

Given component densities $p_z(\cdot; \theta_z)$, the marginal density is $p(x) = \sum_{z=1}^k \pi_z p_z(x; \theta_z)$.

**Definition 3.1.15.** *Mixture Responsibilities*
For fitted $(\widehat{\pi}_m, \widehat{\theta}_m)_{m=1}^k$, define $\widehat{r}_{im} = \widehat{\mathbb{P}}(Z_i = m \mid x_i) = \widehat{\pi}_m p_m(x_i; \widehat{\theta}_m) \left( \sum_{\ell=1}^k \widehat{\pi}_\ell p_\ell(x_i; \widehat{\theta}_\ell) \right)^{-1}$.

Then $\widehat{r}_i$ is soft membership and $\widehat{C}(i) \in \arg\max_{1 \le m \le k} \widehat{r}_{im}$ is a hard assignment.

**Remark 3.1.16.** *Mixtures as Density Estimates*
Gaussian mixtures give flexible density estimates $p(x) = \sum_{m=1}^k \pi_m \phi(x; \mu_m, \Sigma_m)$. If class-conditional mixtures $\widehat{p}_g(x)$ and class priors $\widehat{\pi}_g$ are fit separately, the Bayes classifier is $\widehat{g}(x) \in \arg\max_g \widehat{\pi}_g \widehat{p}_g(x)$.
Without labels, component-label agreement is discovered structure, not supervised training.

**Remark 3.1.17.** *Gaussian Mixtures as Soft $k$-Means*
For $p_m(x) = \phi(x; \mu_m, \sigma^2 I)$, the responsibility is $r_{im} = \pi_m \exp(-\frac{\|x_i - \mu_m\|_2^2}{2\sigma^2}) \left( \sum_{\ell=1}^k \pi_\ell \exp(-\frac{\|x_i - \mu_\ell\|_2^2}{2\sigma^2}) \right)^{-1}$.
Thus $r_{im}$ is distance-weighted; as $\sigma^2 \downarrow 0$, it concentrates on nearest centers, yielding hard $k$-means assignments.

**Method 3.1.18.** *Expectation-Maximization*
Given $x_{1:n}$ and $p(x) = \sum_{m=1}^k \pi_m p_m(x; \theta_m)$, set $\ell(\pi, \theta) = \sum_{i=1}^n \log(\sum_{m=1}^k \pi_m p_m(x_i; \theta_m))$. EM alternates:

(i) *E-step*: compute $r_{im} = \frac{\pi_m p_m(x_i; \theta_m)}{\sum_{\ell=1}^k \pi_\ell p_\ell(x_i; \theta_\ell)}$.

(ii) *M-step*: update $\pi_m^{\mathrm{new}} = n^{-1} \sum_{i=1}^n r_{im}$ and $\theta_m^{\mathrm{new}} \in \arg\max_{\theta_m} \sum_{i=1}^n r_{im} \log p_m(x_i; \theta_m)$.

For Gaussian components, the M-step gives weighted means and variances or covariances.
With an exact M-step, $\ell$ is nondecreasing, but convergence may be to a local maximum.

**Remark 3.1.19.** *Identifiability*
For any permutation $\tau$ of $\{1, \dots, k\}$, $\sum_{m=1}^k \pi_m p_m(x; \theta_m) = \sum_{m=1}^k \pi_{\tau(m)} p_{\tau(m)}(x; \theta_{\tau(m)})$.
Thus labels are identifiable only up to permutation: density is invariant, component interpretation is not.

## 3.2 Dimension Reduction and Manifold Learning

Dimension reduction replaces high-dimensional observations, similarities, or dissimilarities by lower-dimensional coordinates that preserve selected structure. Linear methods preserve variance or inner products; manifold methods preserve neighborhoods, graph distances, or local affine geometry.

**Definition 3.2.1.** *Principal Component Analysis*
Let centered observations $x_1, \dots, x_n \in \mathbb{R}^p$ be the rows of $X \in \mathbb{R}^{n\times p}$, and let $1 \le r \le p$.

(i) Rank-$r$ principal component analysis chooses $V = [v_1 \ \cdots \ v_r] \in \mathbb{R}^{p\times r}$, with $V^TV = I_r$ solving $\min_{V^TV=I_r} \left\|X - XVV^T\right\|_F^2$.

(ii) The columns $v_1, \dots, v_r$ are the principal directions, $\mathrm{im}(V)$ is the rank-$r$ principal subspace, and $x_i \mapsto VV^Tx_i$ is the fitted projection.

(iii) Equivalently, $V$ solves $\max_{V^TV=I_r} \mathrm{tr}(V^TX^TXV)$, with sequential directions $v_k \in \mathrm{argmax}\{\sum_{i=1}^n (v^Tx_i)^2 : \|v\|_2 = 1,\ v \perp v_1, \dots, v_{k-1}\}$.

**Method 3.2.2.** *PCA by SVD*
Let the centered observations be the rows of $X \in \mathbb{R}^{n\times p}$, and fix $1 \le r \le \min\{n, p\}$.

(i) Compute an SVD $X = UDV^T$ with singular values $d_1 \ge \cdots \ge 0$.
(ii) Set $V_r = [v_1 \ \cdots \ v_r]$, the first $r$ right singular vectors.
(iii) Set the score matrix $Z_r = XV_r = U_rD_r$.
(iv) Set the fitted matrix $\widehat{X}_r = Z_rV_r^T = XV_rV_r^T = U_rD_rV_r^T$.

The columns of $V_r$ are the rank-$r$ PCA directions, $Z_r$ contains the principal-component scores, and $\widehat{X}_r$ is the orthogonal rank-$r$ reconstruction minimizing $\left\|X - \widehat{X}\right\|_F^2$.

**Method 3.2.3.** *Kernel PCA*
Let $k$ be a positive definite kernel on observations $x_1, \dots, x_n$, fix $r$, and set $M = n^{-1}\mathbf{1}\mathbf{1}^T$.

(i) Form the Gram matrix $K = (k(x_i, x_{i'}))_{i,i'}$.
(ii) Center it by $\widetilde{K} = (I_n - M)K(I_n - M)$.
(iii) Compute $\widetilde{K}u_j = \lambda_j u_j$ with $\lambda_1 \ge \cdots \ge \lambda_r > 0$, and set $U_r = [u_1 \ \cdots \ u_r]$, $\Lambda_r = \mathrm{diag}(\lambda_1, \dots, \lambda_r)$.
(iv) Return coordinates $Z = U_r\Lambda_r^{1/2}$.

Kernel PCA is spectral PCA on $\widetilde{K}$; radial kernels encode local similarity.

**Method 3.2.4.** *Sparse Principal Components*
Let centered data have rows $x_i^T$ in $X \in \mathbb{R}^{n\times p}$, and let $v \in \mathbb{R}^p$ be a loading vector.

(i) *Variance form*: set $\mathcal{V}_t = \{a \in \mathbb{R}^p : a^Ta = 1,\ \sum_{j=1}^p |a_j| \le t\}$ and choose $v \in \operatorname{argmax}_{a\in\mathcal{V}_t} a^TX^TXa$.

(ii) *Reconstruction form*: for one component, choose $(\theta, v) \in \operatorname{argmin}_{\|\theta\|_2=1} \sum_{i=1}^n \left\|x_i - \theta v^Tx_i\right\|_2^2 + \lambda\|v\|_2^2 + \lambda_1\|v\|_1$.

$L_1$ regularization sparsifies loadings; multiple components alternate sparse-loading and orthogonal-score updates.

**Definition 3.2.5.** *Multidimensional Scaling*
Let $D = (d_{ii'})$ be a symmetric dissimilarity matrix with $d_{ii} = 0$, fix $r < n$, and write $Z = (z_1^T, \dots, z_n^T)^T \in \mathbb{R}^{n\times r}$ with fitted distances $\delta_{ii'}(Z) = \|z_i - z_{i'}\|_2$. Multidimensional scaling chooses $Z$ so that $\delta_{ii'}(Z)$ approximates $d_{ii'}$.

(i) *Least-squares MDS*, or Kruskal–Shepard scaling, minimizes $S_{\mathrm{MDS}}(Z) = \sum_{i\ne i'} (d_{ii'} - \delta_{ii'}(Z))^2$ over $Z \in \mathbb{R}^{n\times r}$.

(ii) *Sammon MDS*, defined when $d_{ii'} > 0$ for $i \ne i'$, minimizes $S_{\mathrm{Sammon}}(Z) = \sum_{i\ne i'} \frac{(d_{ii'} - \delta_{ii'}(Z))^2}{d_{ii'}}$ over $Z \in \mathbb{R}^{n\times r}$.

Sammon weighting penalizes relative errors more strongly for small dissimilarities.

**Method 3.2.6.** *Classical Scaling*
Let $H = I_n - n^{-1}\mathbf{1}\mathbf{1}^T$ and fix $r < n$.

(i) If centered similarities $S = (s_{ii'})$ are supplied, set $B = S$.
(ii) If Euclidean distances $\Delta = (\Delta_{ii'})$ are supplied, set $\Delta^{(2)} = (\Delta_{ii'}^2)$ and $B = -\frac{1}{2}H\Delta^{(2)}H$.
(iii) Compute $Be_j = \lambda_j e_j$ with $\lambda_1 \ge \cdots \ge \lambda_r > 0$, and set $E_r = [e_1 \ \cdots \ e_r]$, $\Lambda_r = \mathrm{diag}(\lambda_1, \dots, \lambda_r)$.
(iv) Return $Z_r = E_r\Lambda_r^{1/2} \in \mathbb{R}^{n\times r}$, with row $z_i^T$ as the coordinate of observation $i$.

The construction gives centered coordinates and fitted Gram matrix $Z_rZ_r^T = E_r\Lambda_rE_r^T$, the rank-$r$ positive-semidefinite approximation to $B$ from its leading positive spectrum.

**Method 3.2.7.** *Nonmetric Scaling*
Given dissimilarities $D = (d_{ii'})$, coordinates $Z = (z_1^T, \dots, z_n^T)^T \in \mathbb{R}^{n\times r}$, and an increasing map $\theta$, nonmetric scaling minimizes $S_{\mathrm{NM}}(Z,\theta) = \sum_{i\neq i'}(\|z_i - z_{i'}\|_2 - \theta(d_{ii'}))^2 / \sum_{i\neq i'} \|z_i - z_{i'}\|_2^2$.

(i) With $\theta$ fixed, update $Z$ by minimizing $S_{\mathrm{NM}}(Z,\theta)$ over $Z \in \mathbb{R}^{n\times r}$.
(ii) With $Z$ fixed, update $\theta$ by isotonic regression of the fitted distances $\|z_i - z_{i'}\|_2$ on the dissimilarities $d_{ii'}$.

Only the ordering of $d_{ii'}$ constrains $\theta(d_{ii'})$; their original scale is discarded.

**Definition 3.2.8.** *Encoder and Decoder*
Let $\mathcal{X}$ be a data space and $\mathcal{H}$ a representation space.

(i) An *encoder* is a map $e : \mathcal{X} \to \mathcal{H}$, with representation $h = e(x)$.
(ii) A *decoder* is a map $d : \mathcal{H} \to \mathcal{X}$, with reconstruction $\widehat{x} = d(h) = d(e(x))$.

Linear PCA is the case $e, d$ linear with $\mathrm{im}(d)$ an orthogonal representation subspace.

**Method 3.2.9.** *Procrustes Shape Alignment*
Let $X_1, X_2 \in \mathbb{R}^{n\times p}$ have corresponding landmark rows.
Orthogonal Procrustes alignment solves $\min_{R^TR=I,\mu\in\mathbb{R}^p} \|X_2 - (X_1R + \mathbf{1}\mu^T)\|_F$.

(i) Compute column means $\bar{x}_1, \bar{x}_2$ and centered matrices $\widetilde{X}_a = X_a - \mathbf{1}\bar{x}_a^T$, $a = 1, 2$.
(ii) Compute an SVD $\widetilde{X}_1^T\widetilde{X}_2 = UDV^T$.
(iii) Set $\widehat{R} = UV^T$ and $\widehat{\mu} = \bar{x}_2 - \widehat{R}^T\bar{x}_1$.
(iv) For scaled Procrustes, set $\widehat{\beta} = \mathrm{tr}(D)/\left\|\widetilde{X}_1\right\|_F^2$, $\widehat{\mu}_\beta = \bar{x}_2 - \widehat{\beta}\widehat{R}^T\bar{x}_1$, and use fitted configuration $\widehat{\beta}X_1\widehat{R} + \mathbf{1}\widehat{\mu}_\beta^T$.

The unscaled solution uses $\widehat{\beta} = 1$ and preserves size; the scaled solution also matches overall scale.

**Method 3.2.10.** *Procrustes Shape Average*
For centered matched shapes $X_\ell \in \mathbb{R}^{n\times p}$, set $\Phi(M, \{R_\ell\}) = \sum_{\ell=1}^{L} \|X_\ell R_\ell - M\|_F^2$, $R_\ell^T R_\ell = I_p$.
A Procrustes shape average solves $\min_{M,\{R_\ell\}} \Phi(M, \{R_\ell\})$.

(i) Initialize $M^{(0)}$ as a centered observed shape or an initially aligned average.
(ii) Given $M^{(t)}$, compute $R_\ell^{(t)} \in \operatorname{argmin}_{R^TR=I_p} \left\|X_\ell R - M^{(t)}\right\|_F^2$ for each $\ell$.
(iii) Update $M^{(t+1)} = L^{-1}\sum_{\ell=1}^{L} X_\ell R_\ell^{(t)}$.
(iv) Recenter $M^{(t+1)}$ and, when size is fixed, rescale it to the chosen norm.
(v) Repeat (ii)–(iv) until $\Phi$ changes by less than the chosen tolerance.

The template is determined only up to a common right rotation unless a final orientation convention is imposed.

**Remark 3.2.11.** *Manifold Hypothesis*
The manifold hypothesis asserts that observations $x_1, \dots, x_n \in \mathbb{R}^p$ satisfy $\mathrm{dist}(x_i, \mathcal{M}) \le \varepsilon$ for some $r$-dimensional set $\mathcal{M} \subset \mathbb{R}^p$, with $r \ll p$ and small $\varepsilon \ge 0$. For curved $\mathcal{M}$, ambient distances $\|x_i - x_j\|_2$ need not approximate intrinsic distances $d_{\mathcal{M}}(x_i, x_j)$. Manifold learning therefore estimates low-dimensional coordinates from neighborhoods, graph geodesics, or tangent spaces, rather than from global Euclidean variance alone.




Figure 255: $d_G$ follows nearby samples along the curve; $d_E$ is the ambient shortcut.

**Example 3.2.12.** *Manifold Methods*
Isomap, locally linear embedding, Laplacian eigenmaps, and diffusion maps construct low-dimensional coordinates from a neighborhood graph or kernel on the sample.

**Method 3.2.13.** *Isomap*
Let $x_i \in \mathbb{R}^p$, fix $r$, set $H = I_n - n^{-1}\mathbf{1}\mathbf{1}^T$, and choose a connected neighborhood graph $G$ on $[n]$.

(i) Assign edge weights $w_{ij} = \|x_i - x_j\|_2$ to neighboring pairs.

(ii) Compute shortest-path distances $d_G(i,j)$ and set $D_G^{(2)} = (d_G(i,j)^2)$.
(iii) Form the centered Gram estimate $B = -\frac{1}{2} H D_G^{(2)} H$.
(iv) Compute $Be_k = \lambda_k e_k$ with $\lambda_1 \geq \cdots \geq \lambda_r > 0$, set $E_r = [e_1 \ \cdots \ e_r]$, $\Lambda_r = \operatorname{diag}(\lambda_1, \ldots, \lambda_r)$.
(v) Return $Z = E_r \Lambda_r^{1/2}$.

Isomap replaces ambient distances by graph geodesic distances; the coordinate extraction is spectral.

**Method 3.2.14.** *Locally Linear Embedding*
Let $x_i \in \mathbb{R}^p$, fix $K, r$, and let $\mathcal{N}(i)$ be the $K$ nearest neighbors of $x_i$.

(i) Set $\mathcal{W}_i = \{w : \sum_k w_k = 1,\ w_k = 0 \ (k \notin \mathcal{N}(i))\}$ and $w_i \in \operatorname*{argmin}_{w \in \mathcal{W}_i} \left\| x_i - \sum_k w_k x_k \right\|_2^2$.
(ii) Form $W = (w_{ik})$ and $M = (I_n - W)^T (I_n - W)$.
(iii) Compute eigenvectors $u_1, \ldots, u_{r+1}$ of $M$ for the smallest eigenvalues, with $u_1 \propto \mathbf{1}$.
(iv) Return $Y = [u_2 \ \cdots \ u_{r+1}] \in \mathbb{R}^{n \times r}$, with row $y_i^T$ as the coordinate of $x_i$.

The affine constraint makes local weights translation-invariant; discarding $u_1$ gives mean-centered coordinates.

**Method 3.2.15.** *Local MDS*
Let $\mathcal{N} \subset \{(i,i') : i < i'\}$ be symmetric nearby pairs, let $z_i \in \mathbb{R}^r$, and fix $\tau > 0$.
Local MDS minimizes $S_{\text{local}}(Z) = \sum_{(i,i') \in \mathcal{N}} (d_{ii'} - \|z_i - z_{i'}\|_2)^2 - \tau \sum_{(i,i') \notin \mathcal{N}} \|z_i - z_{i'}\|_2$.

(i) The local stress term preserves distances on nearby pairs $(i,i') \in \mathcal{N}$.
(ii) The separation term penalizes collapse of non-neighboring pairs $(i,i') \notin \mathcal{N}$.

Thus the objective preserves local metric structure while repelling unrelated manifold regions.

**Definition 3.2.16.** *Principal Curves and Surfaces*
Let $X \in \mathbb{R}^p$ be a random vector and let $f : \Lambda \to \mathbb{R}^p$ be a smooth parametrized set.

(i) The *projection index* is $\lambda_f(x) \in \operatorname*{argmin}_{\lambda \in \Lambda} \|x - f(\lambda)\|_2$.
(ii) A *principal curve* has $\Lambda \subset \mathbb{R}$ and satisfies $f(\lambda) = \mathbb{E}\{X \mid \lambda_f(X) = \lambda\}$.
(iii) A *principal surface* is the same definition with $\Lambda \subset \mathbb{R}^q$, $q \geq 2$, and $f = (f_1, \ldots, f_p)$.

For finite data, alternate coordinate-wise smoothing against $\lambda_f(x_i)$ and reprojection onto $f(\Lambda)$.

**Remark 3.2.17.** *SOMs Versus Principal Surfaces*
Both self-organizing maps and principal surfaces approximate data by low-dimensional representatives.

(i) A principal surface is a smooth map $f : \Lambda \subset \mathbb{R}^q \to \mathbb{R}^p$ with fitted coordinates $\lambda_f(x_i)$.
(ii) A self-organizing map is a finite prototype family $\{m_a\}_{a \in A} \subset \mathbb{R}^p$ indexed by a lattice or graph $A$.

With many prototypes and kernel neighborhood weights, batch SOM updates resemble coordinate-wise surface smoothing, but one fit is smooth-parametric and the other is finite-prototype.

**Remark 3.2.18.** *Local versus Global Scaling*
Let $d_E(i,j) = \|x_i - x_j\|_2$ and let $\mathcal{N}$ be a neighborhood relation.

(i) *Global scaling* uses all pairs through $d_E(i,j)$ or centered inner products, and is appropriate when ambient distances encode the target geometry.
(ii) *Local scaling* uses $\mathcal{N}$, graph distances, or local reconstruction weights to preserve neighborhood order or local shape on a curved set.

Neighborhood scale is critical: too small disconnects, too large shortcuts.

## 3.3 Matrix Factorization and Topic Models

Many unsupervised models express data as combinations of latent factors.

**Definition 3.3.1.** *Low-Rank Matrix Factorization*
Let $X \in \mathbb{R}^{m \times n}$ and $1 \leq r \leq \min\{m,n\}$.
A *rank-r matrix factorization* is pair $(U,V) \in \mathbb{R}^{m \times r} \times \mathbb{R}^{n \times r}$ with fitted matrix $\widehat{X} = UV^\top$ and $\operatorname{rank}(\widehat{X}) \leq r$.
For a loss $\mathcal{L}$, choose $(U,V)$ to minimize $\mathcal{L}(X, UV^\top)$; the columns encode latent row and column factors.

**Definition 3.3.2.** *Nonnegative Matrix Factorization*
Let $X \in \mathbb{R}_+^{n \times p}$ and $1 \leq r \leq \min\{n,p\}$.
A *nonnegative matrix factorization* is a pair $(W,H) \in \mathbb{R}_+^{n \times r} \times \mathbb{R}_+^{r \times p}$ with fitted matrix $\widehat{X} = WH$.
For loss $\mathcal{L}$, minimize $\mathcal{L}(X, WH)$ over $W, H \geq 0$, yielding additive parts-based factors.

**Method 3.3.3.** *Poisson NMF Multiplicative Updates*
For count data $X = (x_{ij})$, set $\mu = WH$ and $L(W, H) = \sum_{i,j}\{x_{ij} \log \mu_{ij} - \mu_{ij}\}$, up to constants.

(i) Initialize $W, H > 0$.
(ii) Set $\rho^W_{ik} = (\sum_j \frac{h_{kj}x_{ij}}{\mu_{ij}})/(\sum_j h_{kj})$ and update $w_{ik} \leftarrow w_{ik}\rho^W_{ik}$.
(iii) Recompute $\mu = WH$, set $\rho^H_{kj} = (\sum_i \frac{w_{ik}x_{ij}}{\mu_{ij}})/(\sum_i w_{ik})$, and update $h_{kj} \leftarrow h_{kj}\rho^H_{kj}$.
(iv) Repeat (ii)–(iii) until $L(W, H)$ stabilises.

The updates are local; different initialisations may give different NMF decompositions.

**Remark 3.3.4.** *NMF Non-Uniqueness*
An NMF $X = WH$ is not identifiable from nonnegativity alone: if $A$ is invertible and $WA \geq 0$, $A^{-1}H \geq 0$, then $X = (WA)(A^{-1}H)$. Interpretability therefore depends on additional geometry, constraints, or initialization.

**Definition 3.3.5.** *Archetypal Analysis*
Let $X \in \mathbb{R}^{n\times p}$ have observed rows $x_i^T$, and fix $r$ archetypes.

(i) Observation weights $W = (w_{ik}) \in \mathbb{R}^{n\times r}_+$ satisfy $\sum_{k=1}^{r} w_{ik} = 1$.
(ii) Archetype weights $B = (b_{ki}) \in \mathbb{R}^{r\times n}_+$ satisfy $\sum_{i=1}^{n} b_{ki} = 1$.
(iii) Archetypes are the rows of $H = BX$, hence lie in $\mathrm{conv}\{x_1, \ldots, x_n\}$.
(iv) Archetypal analysis chooses $W, B$ minimising $J(W, B) = \|X - WBX\|_F^2$.

Archetypes are convex-hull extremes rather than central prototypes.

**Remark 3.3.6.** *Archetypes, k-Means, and NMF*
For prototype matrix $H$ and assignment/mixture weights $W$:

(i) $k$-means uses one-hot rows of $W$.
(ii) NMF uses nonnegative $W, H$ without requiring prototype rows of $H$ to be observed-data mixtures.
(iii) Archetypal analysis uses simplex rows of $W$ and constrains $H = BX$ with simplex rows of $B$.

Archetypes are convex-hull prototypes; $k$-means is hard-assigned and NMF is hull-free.

**Definition 3.3.7.** *Topic Model*
Fix $K$ topics and a vocabulary of size $p$.

(i) Topic $k$ is a word distribution $\beta_k \in \Delta_{p-1}$.
(ii) Document $d$ has topic mixture $\theta_d \in \Delta_{K-1}$.
(iii) Token $m$ in document $d$ is generated by $z_{dm} \sim \mathrm{Cat}(\theta_d)$ and $w_{dm} \sim \mathrm{Cat}(\beta_{z_{dm}})$.
(iv) Latent Dirichlet allocation adds priors $\theta_d \sim \mathrm{Dir}(\alpha)$ and $\beta_k \sim \mathrm{Dir}(\eta)$.

**Remark 3.3.8.** *Regularized Factors*
For a factorization $X \approx UV^\top$, constraints select an identifiable and interpretable solution class.

(i) Low rank bounds latent dimension.
(ii) Sparsity restricts active variables or factors.
(iii) Nonnegativity enforces additive parts.
(iv) Probabilistic priors encode uncertainty and shrinkage.

These restrictions convert an underdetermined algebraic fit into a statistical learning problem.

## 3.4 Representation Learning Objectives

Representation learning trains maps that make downstream prediction, reconstruction, comparison, or generation easier.

**Definition 3.4.1.** *Autoencoder*
Let $x_1, \ldots, x_n \in \mathcal{X}$ and let $\mathcal{H}$ be a representation space.

(i) An *autoencoder* is an encoder-decoder pair $e : \mathcal{X} \to \mathcal{H}$ and $d : \mathcal{H} \to \mathcal{X}$.
(ii) It is trained by minimizing a reconstruction loss such as $n^{-1}\sum_{i=1}^{n} \|x_i - d(e(x_i))\|_2^2$.

Bottlenecks, noise, sparsity, or latent-variable constraints rule out the unrestricted identity solution.

**Definition 3.4.2.** *Sparse Representation Objective*
Let $x_i \in \mathbb{R}^p$, dictionary $W \in \mathbb{R}^{p\times r}$, codes $h_i \in \mathbb{R}^r$, and $\lambda > 0$.

A *sparse representation objective* chooses $W, h_1, \dots, h_n$ to minimize $\sum_{i=1}^{n} \{\|x_i - W h_i\|_2^2 + \lambda \|h_i\|_1\}$.
The reconstruction term fits $x_i$, while the $L_1$ term sparsifies $h_i$.

**Definition 3.4.3.** *Contrastive Objective*
Let $g : \mathcal{X} \to \mathbb{R}^r$ be a representation map, $s(u, v)$ a similarity score, and $\tau > 0$ a temperature.

(i) For an anchor $x$, sample a positive $x^+$ and negatives $x_1^-, \dots, x_K^-$.

(ii) With $a^+ = s(g(x), g(x^+))/\tau$ and $a_k^- = s(g(x), g(x_k^-))/\tau$, min loss i.e. $\log \left\{ \exp(a^+) + \sum_{k=1}^{K} \exp(a_k^-) \right\} - a^+$.

The objective raises similarity to positives and lowers relative similarity to negatives.

## 3.5 Association Rules and Density-Ratio Views

Association-rule analysis searches for simple descriptions of regions where the input distribution has relatively large probability. The classical market-basket setting is the binary special case, but the same idea extends to conjunctions of coordinate conditions.

**Definition 3.5.1.** *Conjunctive Region*
Let $X = (X_1, \dots, X_p)$ take values in $\mathcal{S} = \prod_{j=1}^{p} \mathcal{S}_j$. For $\mathcal{J} \subseteq \{1, \dots, p\}$ and $s = (s_j)_{j \in \mathcal{J}}$ with $s_j \subseteq \mathcal{S}_j$, the *conjunctive region* is $R(\mathcal{J}, s) = \{x \in \mathcal{S} : x_j \in s_j \ \forall j \in \mathcal{J}\}$, with event $\{X \in R(\mathcal{J}, s)\} = \bigcap_{j \in \mathcal{J}} \{X_j \in s_j\}$.

Coordinates in $\{1, \dots, p\} \setminus \mathcal{J}$ are unrestricted; under the extension $s_j = \mathcal{S}_j$, they add no constraint.
For $\mathcal{S}_j \subseteq \mathbb{R}$, $s_j$ is typically an interval; for categorical $\mathcal{S}_j$, $s_j$ is typically a subset of levels.

**Definition 3.5.2.** *Market-Basket Item Sets*
Let $\mathcal{I} = \{1, \dots, K\}$ and $z_i = (z_{ik})_{k \in \mathcal{I}} \in \{0, 1\}^{\mathcal{I}}$ for $i = 1, \dots, n$, where $z_{ik} = 1$ iff observation $i$ contains item $k$.

(i) An *item set* is any $\mathcal{K} \subseteq \mathcal{I}$.

(ii) The containment indicator is $Y_i(\mathcal{K}) = \prod_{k \in \mathcal{K}} z_{ik}$; observation $i$ contains $\mathcal{K}$ iff $Y_i(\mathcal{K}) = 1$.

(iii) The empirical support is $T(\mathcal{K}) = n^{-1} \sum_{i=1}^{n} Y_i(\mathcal{K})$.

(iv) For finite categorical supports, take $\mathcal{I} = \{(j, a) : 1 \le j \le p,\ a \in \mathcal{S}_j\}$, hence $K = \sum_{j=1}^{p} |\mathcal{S}_j|$.

**Definition 3.5.3.** *Association Rule, Confidence, and Lift*
Let $A, B \subseteq \mathcal{I}$ satisfy $A \cap B = \emptyset$, $A \neq \emptyset$, $B \neq \emptyset$, $T(A) > 0$, and $T(B) > 0$.
The *association rule* with antecedent $A$ and consequent $B$ is $A \Rightarrow B$.

(i) The rule support is $S(A \Rightarrow B) = T(A \cup B)$.

(ii) The confidence is $C(A \Rightarrow B) = \frac{T(A \cup B)}{T(A)}$.

(iii) The lift is $L(A \Rightarrow B) = \frac{C(A \Rightarrow B)}{T(B)} = \frac{T(A \cup B)}{T(A) T(B)}$.

Interpretation: $C(A \Rightarrow B)$ estimates prevalence of $B$ given $A$; $L(A \Rightarrow B)$ compares it with $T(B)$.

**Example 3.5.4.** *A Market-Basket Rule*
Let $A = \{\text{peanut butter}, \text{jelly}\}$ and $B = \{\text{bread}\}$.

(i) Support $T(A \cup B) = 0.03$ means all three items appear in about 3% of baskets.

(ii) Confidence $C(A \Rightarrow B) = 0.82$ means about 82% of baskets containing $A$ also contain bread.

(iii) If $T(B) = 0.43$, then $L(A \Rightarrow B) = 0.82/0.43 \approx 1.91$.

The rule predicts bread more often than its marginal prevalence.

**Algorithm 4:** Apriori Association-Rule Search

**Require:** Binary item data $z_1, \dots, z_n$, support threshold $t$, confidence threshold $c$
**Ensure:** Rules with support above $t$ and confidence above $c$
Compute supports for all one-item sets and retain $\mathcal{F}_1 = \{\mathcal{K} : |\mathcal{K}| = 1,\ T(\mathcal{K}) > t\}$
Set $m \leftarrow 2$
**while** $\mathcal{F}_{m-1} \neq \emptyset$ **do**
  Form candidate item sets of size $m$ whose every size-$(m-1)$ subset lies in $\mathcal{F}_{m-1}$
  Scan the data to compute $T(\mathcal{K})$ for each candidate
  Retain $\mathcal{F}_m = \{\mathcal{K} : |\mathcal{K}| = m,\ T(\mathcal{K}) > t\}$
  Set $m \leftarrow m + 1$
**end while**
For each retained item set $\mathcal{K}$, test nonempty proper antecedents $A \subset \mathcal{K}$

Report $A \Rightarrow \mathcal{K} \setminus A$ when $C(A \Rightarrow \mathcal{K} \setminus A) > c$

**Remark 3.5.5.** *Apriori Monotonicity and Limits*
Support is antitone: $\mathcal{L} \subseteq \mathcal{K} \Longrightarrow T(\mathcal{L}) \geq T(\mathcal{K})$.

(i) If $T(\mathcal{L}) < t$, then every $\mathcal{K} \supseteq \mathcal{L}$ also has $T(\mathcal{K}) < t$ and may be pruned.
(ii) A support threshold $t$ can discard rules $A \Rightarrow B$ with large $C(A \Rightarrow B)$ or $L(A \Rightarrow B)$ but small $T(A \cup B)$.




Figure 256: Apriori keeps frequent item sets; if $T(AB) \leq t$, all supersets of $AB$ are pruned.

**Method 3.5.6.** *Generalized Association Rules*
For $\mathcal{J} \subseteq \{1, \ldots, p\}$ and $s_j \subseteq \mathcal{S}_j$, set $R(\mathcal{J}, s) = \bigcap_{j \in \mathcal{J}} \{X_j \in s_j\}$ and $\widehat{P}\{R(\mathcal{J}, s)\} = n^{-1} \sum_{i=1}^{n} \mathbf{1}\{x_{ij} \in s_j \ \forall j \in \mathcal{J}\}$.

(i) Allow $s_j$ to be intervals for quantitative variables and subsets of levels for categorical variables.
(ii) Search over candidate regions with a heuristic rule learner.
(iii) Retain rules whose fitted score and observed support meet the chosen thresholds.

**Method 3.5.7.** *Unsupervised Learning as Supervised Density-Ratio Estimation*
Let $g$ be the data density, $g_0$ a reference density, and $Y = 1$ label data while $Y = 0$ labels reference observations.

(i) Draw a reference sample from $g_0$ and pool it with the data, weighting the two classes equally.
(ii) Fit a probabilistic classifier $\widehat{\mu}(x) \approx \Pr(Y = 1 \mid X = x)$.
(iii) Use $\mu(x) = g(x)/(g(x) + g_0(x))$ to estimate the density ratio by $\widehat{g}(x)/g_0(x) = \widehat{\mu}(x)/(1 - \widehat{\mu}(x))$.

Uniform, Gaussian, and product-marginal $g_0$ target high density, nonnormality, and dependence, respectively.

**Remark 3.5.8.** *Rule Learners for Density Ratios*
After estimating $\widehat{g}/g_0$, any learner that returns regions $R = \bigcap_{j \in \mathcal{J}} \{X_j \in s_j\}$ can search for generalized item sets.
Tree leaves and box-peeling regions have this conjunctive form.
The unsupervised target is large $\widehat{g}(x)/g_0(x)$ on $R$ together with adequate empirical support, not predictive risk.

## 3.6 Graph Ranking and PageRank

Graph ranking turns link structure into a numerical importance score. PageRank is a spectral example: a page receives weight from incoming links, and a link from an already important page contributes more than a link from an unimportant one.

**Definition 3.6.1.** *PageRank Equations*

For pages indexed by $\{1, \ldots, N\}$, define $L \in \{0,1\}^{N \times N}$ by $L_{ij} = 1$ iff page $j$ links to page $i$, and set $c_j = \sum_{i=1}^{N} L_{ij}$.

Assume $c_j > 0$ for all $j$ and fix $d \in (0,1)$.

A *PageRank score vector* is $p = (p_1, \ldots, p_N) \in \mathbb{R}^N$ satisfying $p_i = (1-d) + d \sum_{j=1}^{N} \frac{L_{ij}}{c_j} p_j$ for $i = 1, \ldots, N$.

**Proposition 3.6.2.** *PageRank Normalization*
*Let* $L \in \{0,1\}^{N \times N}$, $c_j = \sum_{i=1}^{N} L_{ij} > 0$, *and* $d \in (0,1)$.

*If* $p \in \mathbb{R}^N$ *satisfies* $p_i = (1-d) + d \sum_{j=1}^{N} \frac{L_{ij}}{c_j} p_j$ *for* $i = 1, \ldots, N$, *then* $\sum_{i=1}^{N} p_i = N$.

*Proof.* Let $P = \sum_{i=1}^{N} p_i$. Summing over $i$ gives $P = N(1-d) + dP$, since $\sum_{i=1}^{N} L_{ij} = c_j$.
Hence $(1-d)P = N(1-d)$, so $P = N$.
*Structural Commentary*: Column normalisation preserves total score up to the teleportation offset.
*Key Insight*: Summing cancels the link structure because each column has mass $c_j$. □

**Method 3.6.3.** *Eigenvector Form*
Let $\mathbf{e} = (1, \dots, 1)^T$, $D_c = \operatorname{diag}(c_1, \dots, c_N)$, and $A = (1-d)\frac{\mathbf{e}\mathbf{e}^T}{N} + dLD_c^{-1}$. The PageRank equations are $\mathbf{p} = A\mathbf{p}$ with $\mathbf{e}^T\mathbf{p} = N$. Since $\mathbf{e}^T A = \mathbf{e}^T$ and $A_{ij} \geq (1-d)/N > 0$, $A$ is positive column-stochastic. Thus $\mathbf{p}$ is the positive eigenvector for eigenvalue 1 normalized by $\mathbf{e}^T\mathbf{p} = N$, and $\mathbf{p}/N$ is the stationary distribution in column-vector convention.

**Algorithm 5:** PageRank by Power Iteration

**Require:** Link matrix $L$, positive outlink counts $c_j$, damping parameter $0 < d < 1$, initial vector $\mathbf{p}_0 > 0$
**Ensure:** PageRank vector normalized by $\mathbf{e}^T\mathbf{p} = N$
  Form $A = (1-d)\mathbf{e}\mathbf{e}^T/N + dLD_c^{-1}$
  **repeat**
    $\mathbf{p}_k \leftarrow A\mathbf{p}_{k-1}$
    $\mathbf{p}_k \leftarrow N\mathbf{p}_k/(\mathbf{e}^T\mathbf{p}_k)$
  **until** the iterates stabilize

**Remark 3.6.4.** *Random-Surfer Normalization*
The probability-scale normalization uses $\pi_i = (1-d)/N + d\sum_{j=1}^{N}(L_{ij}/c_j)\pi_j$ and $\mathbf{e}^T\pi = 1$.
Equivalently, $p = N\pi$ satisfies the average-score form $\mathbf{e}^T\mathbf{p} = N$.
Random-surfer form: follow an outgoing link with probability $d$; jump uniformly with probability $1-d$.

**Example 3.6.5.** *Four-Page Network*
Let $d = 0.85$, $L = \begin{pmatrix} 0&0&1&0 \\ 1&0&0&0 \\ 1&1&0&1 \\ 0&0&0&0 \end{pmatrix}$, $\mathbf{c} = L^T\mathbf{e} = (2,1,1,1)^T$, and $D_c = \operatorname{diag}(\mathbf{c})$.
Solving $\mathbf{p} = (1-d)\mathbf{e} + dLD_c^{-1}\mathbf{p}$ gives $\widehat{\mathbf{p}} = (1.49, 0.78, 1.58, 0.15)^T$. Since $L_{4j} = 0$ for all $j$, $\widehat{p}_4 = 1 - d = 0.15$.




Figure 257: Four-page PageRank graph: $\widehat{\mathbf{p}} \approx (1.49, 0.78, 1.58, 0.15)^T$.

## 3.7 Latent Factor and Component Analysis

Latent factor methods model observed coordinates as mixtures of unobserved sources. The main question is which structure makes the sources interpretable: covariance structure, statistical independence, or non-Gaussian projections.

**Definition 3.7.1.** *Factor Analysis*
Let $X \in \mathbb{R}^p$ and $q < p$. A *factor-analysis model* is a representation $X = AS + \varepsilon$ with $A \in \mathbb{R}^{p\times q}$, $S \in \mathbb{R}^q$, $\varepsilon \in \mathbb{R}^p$, where: $\operatorname{Cov}(S) = I_q$, $\operatorname{Cov}(\varepsilon) = D_\varepsilon$ for diagonal $D_\varepsilon \in \mathbb{R}^{p\times p}$, and $\operatorname{Cov}(S, \varepsilon) = 0$.
The columns $a_1, \dots, a_q$ of $A$ are the *factor loadings*, and $\Sigma := \operatorname{Cov}(X) = AA^T + D_\varepsilon$.
For any orthogonal $R \in \mathbb{R}^{q\times q}$, $(A, S)$ and $(AR^T, RS)$ yield the same $\Sigma$.
Hence $(A, S)$ is identifiable only up to orthogonal rotation.

**Definition 3.7.2.** *Independent Component Analysis*
Let $X \in \mathbb{R}^p$. An *independent component model* is a representation $X = AS$ such that:

(i) $A \in \mathbb{R}^{p\times p}$ is nonsingular.
(ii) $S = (S_1, \dots, S_p)^T \in \mathbb{R}^p$ has mutually independent components.

If $X$ is whitened, then $\operatorname{Cov}(X) = I_p$ and $A$ may be taken orthogonal.
ICA estimates orthogonal $A$ with $Y = A^T X$ approximately componentwise independent, typically via non-Gaussianity contrasts. Gaussian components are identifiable only up to orthogonal rotations.




Figure 258: $v_1, v_2$: PCA variance axes; $a_1, a_2$: ICA rotation exposing independent non-Gaussian coordinates.

**Method 3.7.3.** *Entropy View of ICA*
For a density $g$, define $H(g) = -\int g(y) \log g(y)\, dy$.
Let $Y = (Y_1, \ldots, Y_p)$ have joint density $g_Y$ and marginal densities $g_j$.

(i) Set $H(Y) := H(g_Y)$, $H(Y_j) := H(g_j)$, and $I(Y) := \sum_{j=1}^p H(Y_j) - H(Y)$.
Then $I(Y) \geq 0$, with $I(Y) = 0$ iff $Y_1, \ldots, Y_p$ are independent.
(ii) If $X$ is whitened, $A$ is orthogonal, and $Y = A^T X$, then $H(Y) = H(X)$.
Hence minimising $I(Y)$ over orthogonal $A$ is equivalent to minimising $\sum_{j=1}^p H(Y_j)$.
(iii) Replace marginal-entropy minimisation by marginal non-Gaussianity maximisation,
e.g. through negentropy contrasts, since Gaussian marginals maximise entropy at fixed variance.

**Remark 3.7.4.** *Projection Pursuit and ICA*
Projection pursuit optimizes a non-Gaussianity index over $Y = a^T X$ or $Y = (a_1^T X, a_2^T X)$.
ICA uses the same type of index on $Y_j = a_j^T X$, but imposes an orthogonal system $(a_1, \ldots, a_p)$ after whitening.

**Method 3.7.5.** *Product-Density ICA*
For whitened $X \in \mathbb{R}^p$ and orthogonal $A \in \mathbb{R}^{p \times p}$, set $Y = A^T X$.

(i) Assume the product density $f_Y(y) = \prod_{j=1}^p f_j(y_j)$.
(ii) Parameterize each marginal density by $f_j(s) = \phi(s) \exp(g_j(s))$ with $\int \phi(s) \exp(g_j(s))\, ds = 1$.
(iii) Estimate $A, g_1, \ldots, g_p$ by penalized likelihood subject to $A^T A = I$.
For fixed $A$, the objective decomposes into $p$ one-dimensional smooth density fits.

The roughness penalty shrinks $g_j$ toward 0, the Gaussian baseline $\phi$.

**Algorithm 6:** Product-Density ICA Alternation

**Require:** Whitened observations $x_1, \ldots, x_n$, initial orthogonal matrix $A$
**Ensure:** Orthogonal unmixing matrix $A$ and components $s_{ij} = a_j^T x_i$
**repeat**
**for** each component $j$ **do**
Fit a smooth tilted-Gaussian density $f_j(s) = \phi(s) \exp(g_j(s))$ to the current scores $a_j^T x_i$
Update $a_j \leftarrow \mathbb{E}\{X g_j'(a_j^T X)\} - \mathbb{E}\{g_j''(a_j^T X)\} a_j$
**end for**
Orthogonalize $A$: if $A = UDV^T$ is its SVD, set $A \leftarrow UV^T$
**until** the directions stabilize

# 4 Probabilistic Graphical and Causal Learning

## 4.1 Directed and Undirected Graphical Models

Graphical models use graph structure to encode conditional independence and to organize high-dimensional probability laws.

**Definition 4.1.1.** *Directed Graphical Model*
Let $G = (V, E)$ be a directed acyclic graph with $V = \{1, \ldots, d\}$ and $\mathrm{pa}(j) = \{i \in V : i \to j \in E\}$.
For $x = (x_1, \ldots, x_d)$, write $x_{\mathrm{pa}(j)} = (x_i)_{i \in \mathrm{pa}(j)}$. A law for $X = (X_1, \ldots, X_d)$ with joint density or mass function $p$ *factorizes according to* $G$ if $p(x) = \prod_{j=1}^d p(x_j \mid x_{\mathrm{pa}(j)})$, with $p(x_j \mid x_\varnothing) := p(x_j)$.

**Definition 4.1.2.** *Markov Graph Vocabulary*
Let $G = (V, E)$ be an undirected graph, and let $A, B, C \subseteq V$ be disjoint.

(i) Vertices $j, k \in V$ are *adjacent* iff $\{j, k\} \in E$.
(ii) A *path* from $u$ to $v$ is $(v_0, \ldots, v_m)$ with $v_0 = u$, $v_m = v$, and $\{v_{\ell-1}, v_\ell\} \in E$ for $\ell = 1, \ldots, m$.

(iii) $C$ *separates* $A$ and $B$ iff every path from $a \in A$ to $b \in B$ contains a vertex in $C$.
(iv) A *clique* is $K \subseteq V$ such that $\{j,k\} \in E$ for all distinct $j,k \in K$.
It is *maximal* iff no strictly larger clique contains $K$.

**Definition 4.1.3.** *Undirected Graphical Model*
Let $G = (V,E)$ be an undirected graph, let $\mathcal{C}$ be its maximal cliques, and let $p$ be a positive joint density or mass function for $X = (X_j)_{j\in V}$. For $C \subseteq V$, write $x_C = (x_j)_{j\in C}$. The following are equivalent:

(i) $p$ is *Markov with respect to* $G$: if $j \neq k$ and $\{j,k\} \notin E$, then $X_j \perp\!\!\!\perp X_k \mid X_{V\setminus\{j,k\}}$.
(ii) $p$ factorizes as $p(x) = Z^{-1}\prod_{C\in\mathcal{C}} \psi_C(x_C)$ for clique potentials $\psi_C \geq 0$, with $Z = \sum_x \prod_{C\in\mathcal{C}} \psi_C(x_C)$ in the discrete case and $Z = \int \prod_{C\in\mathcal{C}} \psi_C(x_C)\,dx$ in the continuous case.

**Proposition 4.1.4.** *Gaussian Graphical Zeros*
*Let $X \sim N_d(\mu,\Sigma)$ with $\Sigma \succ 0$ and $\Theta = \Sigma^{-1} = (\theta_{jk})$. For $j$, write $\theta_{-j,j} = (\theta_{\ell j})_{\ell\neq j}$ and $\mu_{-j} = (\mu_\ell)_{\ell\neq j}$. Then:*

*(i) For $z \in \mathbb{R}^{d-1}$, $\mathbb{E}[X_j \mid X_{-j} = z] = \mu_j - \frac{\theta_{-j,j}^T(z-\mu_{-j})}{\theta_{jj}}$.*
*(ii) For $j \neq k$, $X_j \perp\!\!\!\perp X_k \mid X_{-(j,k)}$ iff $\theta_{jk} = 0$.*

*Proof.*

(i) Let $Y = X_j$ and $Z = X_{-j}$. Order coordinates as $(Z,Y)$, with covariance block $\Sigma_{ZZ}$, cross-covariance $\sigma_{Zj}$. Gaussian conditioning gives $\mathbb{E}[Y \mid Z = z] = \mu_j + (z-\mu_{-j})^T\Sigma_{ZZ}^{-1}\sigma_{Zj}$.
From $\Sigma\Theta = I$, the $(Z,j)$ block gives $\Sigma_{ZZ}\theta_{-j,j} + \sigma_{Zj}\theta_{jj} = 0$, hence $\Sigma_{ZZ}^{-1}\sigma_{Zj} = -\frac{\theta_{-j,j}}{\theta_{jj}}$.
(ii) Let $A = \{j,k\}$ and $B = A^c$. The conditional Gaussian law of $X_A \mid X_B$ has precision $\Theta_{AA} = \begin{pmatrix}\theta_{jj} & \theta_{jk}\\ \theta_{jk} & \theta_{kk}\end{pmatrix}$. A bivariate Gaussian has independent coordinates iff its precision is diagonal, so $X_j \perp\!\!\!\perp X_k \mid X_B$ iff $\theta_{jk} = 0$.

*Structural Commentary*: Conditioning turns precision entries into regression interactions.
*Key Insight*: Gaussian conditional independence is zero conditional precision. □

**Definition 4.1.5.** *Ising Model*
Let $E \subseteq \{\{j,k\} : 1 \leq j < k \leq p\}$, set $x_0 = 1$, $\widetilde{E} = E \cup \{\{0,j\} : 1 \leq j \leq p\}$, and $\mathcal{N}(j) = \{k : \{j,k\} \in E\}$.
A *pairwise Ising model* for $X \in \{0,1\}^p$ is specified by:

(i) Joint law $p_\Theta(x) = \exp\left\{\sum_{\{j,k\}\in\widetilde{E}} \theta_{jk}x_jx_k - \Phi(\Theta)\right\}$, where $\Phi(\Theta) = \log \sum_{x\in\{0,1\}^p} \exp\left\{\sum_{\{j,k\}\in\widetilde{E}} \theta_{jk}x_jx_k\right\}$.
(ii) Node conditionals $\mathbb{P}_\Theta(X_j = 1 \mid X_{-j} = x_{-j}) = \left(1 + \exp\left\{-\theta_{0j} - \sum_{k\in\mathcal{N}(j)} \theta_{jk}x_k\right\}\right)^{-1}$ for $j = 1,\ldots,p$.

For $\{j,k\} \in E$, $\theta_{jk}$ is the conditional log-odds interaction between nodes $j$ and $k$.

**Definition 4.1.6.** *Restricted Boltzmann Machine*
Let $v \in \{0,1\}^p$ be visible units, let $h \in \{0,1\}^q$ be hidden units, and let $\sigma(t) = (1+\exp(-t))^{-1}$.
A *restricted Boltzmann machine* is specified by:

(i) Graph restriction: no visible-visible or hidden-hidden edges.
(ii) Joint law $p(v,h) = Z^{-1}\exp(a^Tv + b^Th + v^TWh)$ with $a \in \mathbb{R}^p$, $b \in \mathbb{R}^q$, and $W \in \mathbb{R}^{p\times q}$.
(iii) Conditional factorization: $p(v \mid h) = \prod_{j=1}^p \mathrm{Bern}(v_j;\sigma(a_j + W_{j\cdot}h))$ and $p(h \mid v) = \prod_{k=1}^q \mathrm{Bern}(h_k;\sigma(b_k + W_{\cdot k}^Tv))$.

In supervised variants, $v$ may concatenate feature and label units.




Figure 259: RBM: $H_j$ connect only to visible features $V_i$ and label $Y$; no within-layer edges.

**Remark 4.1.7.** *Separation*
For disjoint vertex sets $A,B,C$, separation is a graphical certificate for $X_A \perp\!\!\!\perp X_B \mid X_C$: DAGs use d-separation, while undirected graphs use path separation by $C$.
In a Markov graph, $C$ separates $A$ and $B$ implies $X_A \perp\!\!\!\perp X_B \mid X_C$.
Missing edges give pairwise certificates; separation gives setwise certificates.

## 4.2 Inference and Learning in Graphical Models

Inference in a graphical model computes marginal, conditional, or most likely latent quantities from the joint law encoded by the graph.

**Definition 4.2.1.** *Inference Tasks*
Let $X = (X_j)_{j \in V}$ have joint law $p$, and let $A, B \subseteq V$ be disjoint. Standard *inference tasks* are:

(i) *Marginal inference*: compute $p(x_A)$.
(ii) *Conditional inference*: compute $p(x_A \mid x_B)$.
(iii) *MAP inference*: compute $x_A^\star \in \arg\max_{x_A} p(x_A \mid x_B)$, with $B = \varnothing$ for unconditional MAP.
(iv) *Latent posterior inference*: for latent $Z$ and observed $X = x$, compute $p(z \mid x)$.

**Method 4.2.2.** *Message Passing*
Let $G = (V, E)$ be a graphical model with local potentials.

(i) *Tree belief propagation*: on trees, two-way edge messages yield exact node and edge marginals.
(ii) *Loopy belief propagation*: on cyclic graphs, iterated message updates yield approximate marginals.

**Definition 4.2.3.** *Variational Inference*
For latent-variable model $p(x, z)$, intractable $p(z \mid x)$, tractable family $\mathcal{Q}$, *variational inference* choose $q^\star \in \mathcal{Q}$:

(i) *KL projection*: $q^\star \in \arg\min_{q \in \mathcal{Q}} \mathrm{KL}(q(z) \| p(z \mid x))$.
(ii) *ELBO maximization*: equivalently, maximize $\mathcal{L}(q) := \mathbb{E}_q[\log p(x, z)] - \mathbb{E}_q[\log q(z)]$, since $\log p(x) = \mathcal{L}(q) + \mathrm{KL}(q(z) \| p(z \mid x))$.

**Remark 4.2.4.** *Learning Parameters and Structure*
Let $G$ denote the graph and $\theta$ its model parameters.

(i) *Parameter learning*: for fixed $G$, estimate $\theta$ by likelihood or regularized likelihood.
(ii) *Structure learning*: estimate unknown $G$ by optimizing a statistical score over graph candidates, often with structural penalties or combinatorial search.

**Method 4.2.5.** *Gaussian Graphical Model Learning*
For centered Gaussian data with empirical covariance $S$, let $\Theta = \Sigma^{-1} \succ 0$ and $\ell(\Theta) := \log\det\Theta - \mathrm{trace}(S\Theta)$.

(i) *Known graph*: for edge set $E$, estimate $\widehat{\Theta} \in \arg\max_{\Theta \succ 0} \ell(\Theta)$ subject to $\theta_{jk} = 0$ whenever $\{j, k\} \notin E$.
(ii) *Unknown graph*: graphical lasso solves $\widehat{\Theta} \in \arg\max_{\Theta \succ 0} \left\{ \ell(\Theta) - \lambda \sum_{j \neq k} |\theta_{jk}| \right\}$, penalise off-diagonal entries.

Graph selection is therefore shrinkage of inverse-covariance entries; zeros of $\widehat{\Theta}$ remove edges.

Figure 260: $\mathcal{D}$ is the data and $S$ its empirical covariance. $Y$ gives $X \perp \{Z, W\} \mid Y$; graphical lasso solves $\widehat{\Theta}_\lambda \in \arg\max_{\Theta \succ 0} \left\{ \log\det\Theta - \mathrm{tr}(S\Theta) - \lambda \sum_{j \neq k} |\theta_{jk}| \right\}$, and zeros of $\widehat{\Theta}_\lambda$ define missing edges in $\widehat{G}$.

**Algorithm 7:** Graphical Lasso by Block Coordinate Descent

**Require:** Empirical covariance $S \in \mathbb{R}^{p \times p}$, penalty $\lambda \geq 0$
**Ensure:** Covariance estimate $W$ and sparse precision estimate $\widehat{\Theta} = W^{-1}$
Initialize $W \leftarrow S + \lambda I$ and keep the diagonal fixed
**repeat**
**for** $j = 1, \ldots, p$ **do**
Partition $W$ into $W_{11}$, $w_{12}$, and $w_{22}$ by isolating row and column $j$; partition $S$ similarly to get $s_{12}$
Define $\mathcal{S}_\lambda(z) = \mathrm{sign}(z)(|z| - \lambda)_+$
**repeat**
**for** $k = 1, \ldots, p-1$ **do**

$$\beta_k \leftarrow \frac{\mathcal{S}_\lambda\left(s_{12,k} - \sum_{\ell \neq k} (W_{11})_{k\ell} \beta_\ell\right)}{(W_{11})_{kk}}$$

**end for**
**until** $\beta$ has converged for this block
Update $w_{12} \leftarrow W_{11}\beta$ and symmetrically update row and column $j$ of $W$

**end for**
**until** $W$ has converged
In the final cycle recover $\widehat{\theta}_{12} = -\widehat{\beta}\widehat{\theta}_{22}$ with $1/\widehat{\theta}_{22} = w_{22} - w_{12}^T\widehat{\beta}$

**Method 4.2.6.** *Discrete Markov Network Learning*
For a binary Markov network with edge statistic $T_{jk}(X) = X_jX_k$, likelihood equations are $\widehat{\mathbb{E}}T_{jk} = \mathbb{E}_\Theta T_{jk}$ for $\{j,k\} \in E$; exact $\mathbb{E}_\Theta$ sums over $x \in \{0,1\}^p$.

(i) *Small graphs*: solve $\widehat{\mathbb{E}}T_{jk} = \mathbb{E}_\Theta T_{jk}$ by log-linear Poisson MLE, gradient/Newton updates, or iterative proportional fitting.
(ii) *Large graphs*: approximate $\mathbb{E}_\Theta T_{jk}$ by mean-field or Gibbs-sampling estimates.
(iii) *Hidden-node models*: use observed-data conditionals for $\widehat{\mathbb{E}}T_{jk}$; for RBMs, CD-$r$ starts at data and runs $r$ alternating Gibbs steps $h \sim p(h \mid v)$, $v \sim p(v \mid h)$.

## 4.3 Causal Graphs, Interventions, and Identifiability

Causal learning asks which predictions remain valid under interventions, not merely under passive observation.

**Definition 4.3.1.** *Structural Causal Model*
Let $J$ index endogenous variables $X = (X_j)_{j\in J}$ and exogenous disturbances $U = (U_j)_{j\in J}$.
A structural causal model is $M = (G, \{f_j\}_{j\in J}, P_U)$ such that:

(i) $G = (J, E)$ is a directed graph with $\mathrm{pa}(j) = \{k \in J : k \to j \in E\}$.
(ii) $P_U$ is the joint law of $U$.
(iii) $X_j = f_j(X_{\mathrm{pa}(j)}, U_j)$ for every $j \in J$.

The edges $k \to j$ are the direct causal inputs to $X_j$.

**Definition 4.3.2.** *Intervention*
For $M = (G, \{f_k\}_{k\in J}, P_U)$ and $x \in \mathcal{X}_j$, set $\tilde{f}_j \equiv x$ and $\tilde{f}_k = f_k$ for $k \neq j$.
The *intervention* $\mathrm{do}(X_j = x)$ is the modified model $M_{j:=x} = (G_{j:=x}, \{\tilde{f}_k\}_{k\in J}, P_U)$, where $G_{j:=x}$ deletes all incoming edges to $j$; the induced law $P_{M_{j:=x}}$ is the interventional distribution.

**Definition 4.3.3.** *Identifiability*
Let $\mathcal{M}$ be the assumed model class and let $P_M$ be the observational law induced by $M \in \mathcal{M}$.
A causal functional $\psi : \mathcal{M} \to \Theta$ is *identifiable* on $\mathcal{M}$ if $P_M = P_{M'}$ implies $\psi(M) = \psi(M')$ for all $M, M' \in \mathcal{M}$.

**Remark 4.3.4.** *Causal and Predictive Targets*
Let $\hat{y} = g(z)$ be trained under the observational law $P_M$. Small risk $R_{P_M}(g) = \mathbb{E}_{P_M}[\ell(Y, g(Z))]$ need not imply small interventional risk $R_{P_{M_{j:=x}}}(g)$ after $\mathrm{do}(X_j = x)$. Transport across regimes requires causal stability of the conditional mechanism used by $g$, not only predictive fit under $P_M$.

## 4.4 Counterfactuals and Policy Evaluation

Policy learning and evaluation combine prediction with causal questions about actions not actually taken.

**Definition 4.4.1.** *Potential Outcomes*
Let $A \in \mathcal{A}$ be a treatment.

(i) *Potential outcome*: $Y(a)$ is the outcome under $\mathrm{do}(A = a)$.
(ii) *Consistency*: $Y = Y(A)$.
(iii) *Individual effect*: for binary $A \in \{0,1\}$, $\tau = Y(1) - Y(0)$.
(iv) *Average treatment effect*: $\mathrm{ATE} = \mathbb{E}[\tau]$.

**Definition 4.4.2.** *Ignorability and Propensity*
For binary treatment $A$ and covariates $X$:

(i) *Ignorability*: $(Y(0), Y(1)) \perp\!\!\!\perp A \mid X$.
(ii) *Propensity score*: $e(x) = P(A = 1 \mid X = x)$.
(iii) *Overlap*: $0 < e(X) < 1$ almost surely.

**Definition 4.4.3.** *Off-Policy Evaluation*
Let $(X_i, A_i, Y_i)_{i=1}^n$ be logged under behavior policy $\mu(a \mid x)$, and let $\pi(a \mid x)$ be a target policy.

(i) *Target value*: $V(\pi) = \mathbb{E}_\pi[Y]$.
(ii) *Off-policy evaluation*: estimate $V(\pi)$ from data sampled under $\mu$.
(iii) *Inverse-propensity weighting*: $\hat{V}_{\mathrm{IPW}}(\pi) = n^{-1}\sum_{i=1}^n \{\pi(A_i \mid X_i)/\mu(A_i \mid X_i)\}Y_i$.

(iv) *Overlap*: $\mu(a \mid x) > 0$ whenever $\pi(a \mid x) > 0$.

**Remark 4.4.4.** *Feedback*
If a predictor $g$ induces actions through a policy $\pi_g(a \mid x)$, then future data are drawn from the post-deployment law $P^{\pi_g}$ rather than the pre-deployment law $P^0$. Feedback is the dependence $g \mapsto \pi_g \mapsto P^{\pi_g}$; causal learning, reinforcement learning, and deployment monitoring all study or control this induced law.

# 5 Neural Network Foundations

## 5.1 Network Architectures and Expressivity

The migrated deep-feedforward-network notes describe neural networks as learned feature maps followed by prediction heads.

**Definition 5.1.1.** *Deep Feedforward Network*
Let $x \in \mathcal{X}$, set $h^{(0)} = x$, and fix depth $L$.

(i) *Layer recursion*: $h^{(k)} = g^{(k)}(W^{(k)}h^{(k-1)} + b^{(k)})$ for $k = 1, \dots, L$.
(ii) *Network map*: $f_\theta : \mathcal{X} \to \mathcal{Y}$ is $f_\theta(x) = h^{(L)} = (f_L \circ \cdots \circ f_1)(x)$.
(iii) *Parameters*: $\theta = \{(W^{(k)}, b^{(k)})\}_{k=1}^{L}$.
(iv) *Training objective*: choose $\theta$ to minimize empirical risk $\hat{R}_n(f_\theta)$ or population risk $R(f_\theta)$.

**Remark 5.1.2.** *Learned Feature Maps*
Let $\ell$ be the task loss and let $R(f) = \mathbb{E}[\ell(Y, f(X))]$. Two representation parametrizations are:

(i) *fixed features*: $f_w(x) = a_w(\phi(x))$, with $\phi$ fixed and only the head parameter $w$ optimized.
(ii) *learned features*: $f_{\theta,w}(x) = a_w(\phi_\theta(x))$, with $\theta$ and $w$ optimized jointly.

Learning $\phi_\theta$ adds adaptivity but usually destroys convexity.

**Definition 5.1.3.** *Projection Pursuit and Ridge Functions*
Let $x \in \mathbb{R}^p$.

(i) *Projection-pursuit form*: $f(x) = \sum_{m=1}^{M} g_m(\omega_m^T x)$ with $\|\omega_m\|_2 = 1$.
(ii) *Ridge coordinate*: $u_m = \omega_m^T x$.
(iii) *Ridge function*: $g_m(u_m)$ depends on $x$ only through the scalar coordinate $u_m$.
(iv) *Learned components*: the directions $\omega_m$ and univariate profiles $g_m$ are estimated from data.

**Definition 5.1.4.** *Output and Hidden Units*
Let $T \in \mathbb{R}^K$ denote output scores. Common output parametrizations are:

(i) *linear*: $f_k(x) = T_k$, for conditional means or real-valued responses.
(ii) *sigmoid*: $p(x) = \sigma(T) = \frac{1}{1+\exp(-T)}$, for Bernoulli probabilities.
(iii) *softmax*: $p_k(x) = \exp(T_k) / \sum_{\ell=1}^{K} \exp(T_\ell)$, for $K$-class probabilities.

Hidden units define activations $Z_m = \phi_m(a_m)$ from preactivations $a_m$, commonly using ReLU, leaky ReLU, maxout, tanh, or another nonlinear map.

**Definition 5.1.5.** *Single Hidden Layer Network*
Let $x \in \mathbb{R}^p$, $Z = (Z_1, \dots, Z_M)^T$, $T = (T_1, \dots, T_K)^T$. A single-hidden-layer feedforward network is defined by:

(i) $Z_m = \sigma(\alpha_{0m} + \alpha_m^T x)$ for $m = 1, \dots, M$.
(ii) $T_k = \beta_{0k} + \beta_k^T Z$ for $k = 1, \dots, K$.
(iii) $f_k(x) = g_k(T)$, with $g_k(T) = T_k$ for regression and $g_k(T) = \exp(T_k) / \sum_{\ell=1}^{K} \exp(T_\ell)$ for $K$-class classification.

In the classification case, $f(x) \in \Delta^{K-1}$.




Figure 261: Single-hidden-layer feedforward network.

**Remark 5.1.6.** *Neural Networks as Ridge Expansions*
For scalar output, a one-hidden-layer network has $f(x) = \beta_0 + \sum_{m=1}^{M} \beta_m \sigma(\alpha_{0m} + \alpha_m^T x)$.
If $\alpha_m \neq 0$ and $\omega_m = \frac{\alpha_m}{\|\alpha_m\|_2}$, the $m$th term is $\beta_m \sigma(\alpha_{0m} + \|\alpha_m\|_2 \omega_m^T x)$, a ridge term in coordinate $\omega_m^T x$ with learned *direction* $\omega_m$, *scale* $\|\alpha_m\|_2$, *threshold* $\alpha_{0m}$, and *output coefficient* $\beta_m$.
Thus a shallow scalar network is a projection-pursuit expansion with parametrized ridge profiles.

**Theorem 5.1.7.** *Universal Approximation Theorem Form*
*For compact $K \subset \mathbb{R}^d$, $h \in C(K)$, non-polynomial activation $\sigma : \mathbb{R} \to \mathbb{R}$, $\forall \varepsilon > 0$, $\exists\, M \in \mathbb{N}$, $a_m, b_m \in \mathbb{R}$, $w_m \in \mathbb{R}^d$ s.t. one-hidden-layer network $f_M(x) = \sum_{m=1}^{M} a_m \sigma(w_m^\top x + b_m)$ satisfies $\|f_M - h\|_\infty < \varepsilon$ on $K$.*

*Proof.* Let $\mathcal{A}_\sigma := \operatorname{span}\{\sigma(w^\top x + b) : w \in \mathbb{R}^d,\ b \in \mathbb{R}\} \subset C(K)$.
The classical universal approximation theorem states that $\overline{\mathcal{A}_\sigma}^{\|\cdot\|_\infty} = C(K)$ for suitable non-polynomial $\sigma$.
Since $h \in C(K)$ and $\varepsilon > 0$, there is $f_M \in \mathcal{A}_\sigma$ with $\|f_M - h\|_\infty < \varepsilon$.
Thus $f_M(x) = \sum_{m=1}^{M} a_m \sigma(w_m^\top x + b_m)$ has one hidden layer and linear output.
*Structural Commentary*: Non-polynomial activations make ridge spans dense in $C(K)$.
*Key Insight*: Width supplies finite linear combinations of nonlinear ridge functions. □

**Remark 5.1.8.** *Depth and Expressivity*
Let $N_L(\varepsilon, h)$ be the minimum width of a depth-$L$ network with $\|f - h\|_\infty \leq \varepsilon$ on a compact domain. Universal approximation gives $N_L(\varepsilon, h) < \infty$, not trainability; depth expressivity is $N_L(\varepsilon, h) \ll N_{L'}(\varepsilon, h)$ for some $L > L'$.

## 5.2 Backpropagation and Automatic Differentiation

Backpropagation is reverse-mode differentiation applied to the computational graph of the network and loss.

**Definition 5.2.1.** *Computational Graph*
A computational graph is a finite directed acyclic graph $G = (V, E)$ together with variables $(x_v)_{v \in V}$ and local maps $(f_v)_{v \in V \setminus S}$, where $\operatorname{pa}(v) = \{u \in V : (u, v) \in E\}$ and $S = \{v \in V : \operatorname{pa}(v) = \varnothing\}$.

(i) For $v \in S$, $x_v$ is an input, parameter, or constant.
(ii) For $v \in V \setminus S$, $x_v = f_v((x_u)_{u \in \operatorname{pa}(v)})$.
(iii) An edge $u \to v$ records the direct dependence of $x_v$ on $x_u$ in $f_v$.

For scalar output $L = x_\ell$ and parameter sources $P \subseteq S$, reverse mode computes $(\partial L / \partial x_v)_{v \in V}$, hence $\nabla_{(x_p)_{p \in P}} L$.

**Method 5.2.2.** *Backpropagation*
Let $h^{(0)} = x$, $z^{(k)} = W^{(k)} h^{(k-1)} + b^{(k)}$, $h^{(k)} = \phi_k(z^{(k)})$ for $1 \leq k \leq K$, and $L = \ell(h^{(K)}, y)$.
Write $\bar{u} = \nabla_u L$ for the adjoint of $u$.

(i) Compute $z^{(k)}$ and $h^{(k)}$ for $k = 1, \ldots, K$, then compute $L$.
(ii) Initialize $\bar{h}^{(K)} = \nabla_{h^{(K)}} \ell(h^{(K)}, y)$.
(iii) For $k = K, \ldots, 1$, set $\bar{z}^{(k)} = J_{\phi_k}(z^{(k)})^T \bar{h}^{(k)}$ and $(\nabla_{W^{(k)}} L, \nabla_{b^{(k)}} L, \bar{h}^{(k-1)}) = (\bar{z}^{(k)} (h^{(k-1)})^T, \bar{z}^{(k)}, (W^{(k)})^T \bar{z}^{(k)})$.

Thus backpropagation is reverse-mode differentiation of $L$ through the layerwise computational graph.

**Method 5.2.3.** *Single-Hidden-Layer Backpropagation Equations*
For one training case $(x_i, y_i)$ in a single-hidden-layer network, let $m = 1, \ldots, M$ index hidden units, $k = 1, \ldots, K$ index output components, and $z_i = (z_{1i}, \ldots, z_{Mi})^T$.

(i) Compute $a_{mi} = \alpha_{0m} + \alpha_m^T x_i$, $z_{mi} = \sigma(a_{mi})$, $t_{ki} = \beta_{0k} + \beta_k^T z_i$, and $f_k(x_i) = g_k(t_{ki})$.
(ii) Compute $R_i = \sum_{k=1}^{K} \{y_{ik} - f_k(x_i)\}^2$ and $\delta_{ki} = \frac{\partial R_i}{\partial t_{ki}} = -2\{y_{ik} - f_k(x_i)\} g_k'(t_{ki})$.
(iii) Compute $\nabla_{\beta_k} R_i = \delta_{ki} z_i$ and $\frac{\partial R_i}{\partial \beta_{0k}} = \delta_{ki}$.
(iv) Compute $s_{mi} = \frac{\partial R_i}{\partial a_{mi}} = \sigma'(a_{mi}) \sum_{k=1}^{K} \beta_{km} \delta_{ki}$.
(v) Compute $\nabla_{\alpha_m} R_i = s_{mi} x_i$ and $\frac{\partial R_i}{\partial \alpha_{0m}} = s_{mi}$.

Batch gradients are obtained by summing these per-case derivatives over $i$.

**Remark 5.2.4.** *Automatic Differentiation*
For an implemented graph map $F : \mathbb{R}^p \to \mathbb{R}^q$, automatic differentiation returns $DF(x)$ by chain-rule evaluation on the executed graph. It differentiates the floating-point computation up to roundoff, without finite differences or symbolic simplification.

## 5.3 Optimization for Deep Learning

The old optimization notes are migrated here only where the algorithms are used as training methods for neural networks. General numerical optimization belongs to Part 18.

**Definition 5.3.1.** *Neural Empirical Objective*

For data $(x_i, y_i)_{i=1}^n$, network $f_\theta : x \mapsto f(x;\theta)$, loss $L$, neural empirical objective is $J_n(\theta) = n^{-1} \sum_{i=1}^{n} L(f_\theta(x_i), y_i)$.

The population objective is $J(\theta) = \mathbb{E}[L(f_\theta(X), Y)]$ under the data-generating law.

**Algorithm 8:** Minibatch Backpropagation Step

**Require:** Minibatch $B$, parameters $\theta$, learning rate $\eta$
Compute predictions $f(x;\theta)$ for all $(x, y) \in B$
Compute minibatch loss $\widehat{L}_B(\theta)$
Run reverse-mode automatic differentiation to compute $g = \nabla_\theta \widehat{L}_B(\theta)$
Update parameters $\theta \leftarrow \theta - \eta g$

**Remark 5.3.2.** *Deep Optimization Challenges*

For empirical loss $J_n(\theta)$, deep optimization is shaped by:

(i) *geometry*: large $\kappa(\nabla^2 J_n)$, saddles with $\nabla J_n \approx 0$ and indefinite $\nabla^2 J_n$, and plateaus with $\|\nabla J_n\| \approx 0$.
(ii) *gradient propagation*: large or small $\left\| \frac{\partial h^{(L)}}{\partial h^{(k)}} \right\|$ causes exploding or vanishing gradients.
(iii) *stochasticity*: $g_k = \nabla J_n(\theta_k) + \xi_k$, with $\mathbb{E}[\xi_k \mid \theta_k] = 0$ and $\mathrm{Var}(\xi_k \mid \theta_k) \neq 0$.
(iv) *stabilization*: initialization, normalization, gradient clipping, skip connections, and learning-rate schedules control scale, conditioning, and update variance.

These choices are part of the training specification, not post hoc implementation details.

**Remark 5.3.3.** *Second-Order Methods*

Let $J : \mathbb{R}^d \to \mathbb{R}$, $g_k = \nabla J(\theta_k)$, and $H_k = \nabla^2 J(\theta_k)$.

(i) *Newton*: choose $p_k$ from $H_k p_k = -g_k$ and set $\theta_{k+1} = \theta_k + \eta_k p_k$.
(ii) *Conjugate gradient*: solve $H_k p = -g_k$ approximately using only products $H_k v$.
(iii) *BFGS/L-BFGS*: use $p_k = -B_k g_k$, where $B_k \approx H_k^{-1}$; L-BFGS stores few curvature pairs.

Large networks make explicit $H_k$ storage or factorization prohibitive, so these methods usually act as preconditioners, curvature approximations, or analysis tools.

## 5.4 Regularization, Normalization, and Generalization

Deep-learning regularization combines penalties, stochastic training effects, architecture, and validation-based stopping.

**Definition 5.4.1.** *Parameter-Norm Regularization*

Minimize $\widetilde{J}_\lambda(\theta) = J(\theta) + \lambda\Omega(\theta)$, with $\lambda \geq 0$, $\Omega \geq 0$, and penalized non-intercept set $\mathcal{P}$.

(i) *$\ell^2$ weight decay*: $\Omega_2(\theta) = \sum_{j \in \mathcal{P}} \theta_j^2$.
(ii) *$\ell^1$ penalty*: $\Omega_1(\theta) = \sum_{j \in \mathcal{P}} |\theta_j|$.
(iii) *Single-hidden-layer weight decay*: for weights $\alpha_{m\ell}$ and $\beta_{km}$, $\Omega_2(\theta) = \sum_{m,\ell} \alpha_{m\ell}^2 + \sum_{k,m} \beta_{km}^2$.

Intercepts such as $\alpha_{0m}$ and $\beta_{0k}$ are excluded unless explicitly included in $\mathcal{P}$.

**Definition 5.4.2.** *Dropout*

Let $h \in \mathbb{R}^d$, $q \in (0, 1]$, and $m_j \overset{\text{ind}}{\sim} \text{Bernoulli}(q)$.

(i) *Training mask*: replace $h$ by $\widetilde{h} = \frac{m \odot h}{q}$.
(ii) *Subnetwork*: each mask $m \in \{0, 1\}^d$ selects the active coordinates $\{j : m_j = 1\}$ with shared parameters.
(iii) *Evaluation*: use the unmasked activation $h$ when inverted dropout scaling is used in training.

Dropout optimizes an expected masked loss over tied-parameter subnetworks.

**Method 5.4.3.** *Early Stopping*

Let $\theta_t$ be the parameter after training step $t$ and let $V(\theta)$ be the validation loss.

(i) Compute $V_t = V(\theta_t)$ after selected training steps $t$.
(ii) Select $t^\star \in \arg\min_{0 \leq t \leq T} V_t$.
(iii) Return $\widehat{\theta}_{\text{ES}} = \theta_{t^\star}$.

The stopping time $t^\star$ restricts the reachable parameter set and acts as an implicit regularization parameter.

**Remark 5.4.4.** *Starting Values, Scaling, and Multiple Fits*
For a nonconvex neural objective $J(\theta)$, the fitted parameter can depend on initialization and scaling.

(i) *Initialization*: avoid $\theta_0 = 0$ when symmetry identifies hidden units, and avoid large $\|\theta_0\|$ when $\sigma'(a) \approx 0$.
(ii) *Scaling*: standardize inputs, e.g. $x_j \leftarrow \frac{x_j - \hat{\mu}_j}{\hat{\sigma}_j}$, so coordinates have comparable optimization scale.
(iii) *Multiple fits*: run starts $r = 1, \dots, R$, obtain $\widehat{\theta}^{(r)}$, and choose $\widehat{\theta}^{(r^\star)}$ by validation or penalized criterion.

Thus initialization law, preprocessing, and restart selection are part of the training specification.

**Remark 5.4.5.** *Hidden Units and Weight Decay*
Let $M$ be hidden width, $\lambda$ the weight-decay strength, and $t^\star$ the early-stopping time.

(i) *Small $M$*: the model class may have excessive approximation error.
(ii) *Large $M$*: overfitting can occur unless $\lambda > 0$, finite $t^\star$, or another constraint limits effective complexity.
(iii) *Selection*: choose $(M, \lambda, t^\star)$ by validation or another held-out criterion.

Weight decay and early stopping bias the fit toward smaller effective weights and simpler functions.

**Definition 5.4.6.** *Batch Normalization*
Let $H = (h_{ij}) \in \mathbb{R}^{b \times d}$ be minibatch activations, indexed by example $i$ and coordinate $j$.

(i) *Batch statistics*: $\mu_j = \frac{1}{b}\sum_i h_{ij}$ and $s_j^2 = \frac{1}{b}\sum_i (h_{ij} - \mu_j)^2$.
(ii) *Normalization*: $\widehat{h}_{ij} = \frac{h_{ij} - \mu_j}{\sqrt{s_j^2 + \varepsilon}}$.
(iii) *Affine output*: $y_{ij} = \gamma_j \widehat{h}_{ij} + \beta_j$, with learned $\gamma_j, \beta_j$.
(iv) *Evaluation*: replace $\mu_j$ and $s_j^2$ by running training estimates.

**Remark 5.4.7.** *Augmentation and Noise*
Let $\ell(y, f_\theta(x))$ be the task loss.

(i) *Data augmentation*: replace $(X, Y)$ by $(T(X), Y)$ for random label-preserving transforms $T$.
(ii) *Label smoothing*: replace one-hot $y$ by $y^{(\varepsilon)} = (1 - \varepsilon)y + \varepsilon u$, where $u$ is a reference class distribution.
(iii) *Noise injection*: train on $x + \xi$ or hidden activations $h + \xi$, with $\mathbb{E}\xi = 0$.
(iv) *Multitask learning*: minimize $\sum_{r=1}^{R} \omega_r \ell_r(y_r, f_{\theta,r}(x))$ with shared representation parameters.

These methods regularize through altered data, targets, perturbations, or shared representations.

# 6 Modern Deep Models

## 6.1 Convolutional and Geometric Architectures

The convolutional-network source is migrated here because convolution is an architectural prior for data with grid, group, or geometric structure.

**Definition 6.1.1.** *Convolution*
For $f, g : \mathbb{R} \to \mathbb{R}$ and $a, b : \mathbb{Z} \to \mathbb{R}$:

(i) *Continuous*: $f * g : D_c \ni t \mapsto \int_{-\infty}^{\infty} f(\tau) g(t - \tau)\, d\tau \in \mathbb{R}$; $D_c := \{t \in \mathbb{R} : \int_{\mathbb{R}} |f(\tau) g(t - \tau)|\, d\tau < \infty\}$.
(ii) *Discrete*: $a * b : D_d \ni n \mapsto \sum_{m \in \mathbb{Z}} a[m] b[n - m] \in \mathbb{R}$; $D_d := \{n \in \mathbb{Z} : \sum_{m \in \mathbb{Z}} |a[m] b[n - m]| < \infty\}$.

**Definition 6.1.2.** *Convolutional Neural Network*
Let $\Phi = L_k \circ \cdots \circ L_1$ be a neural network on array-valued activations.

(i) *Convolutional layer*: on $z : \mathbb{Z}^d \to \mathbb{R}^{C_{\text{in}}}$, $L_j z = \sigma(K * z + b)$ or $\sigma(K \star z + b)$, with finite-support $K$, $b \in \mathbb{R}^{C_{\text{out}}}$, pointwise $\sigma$, and cross-correlation $\star$.
(ii) *Convolutional neural network*: $\Phi$ is a CNN if at least one layer $L_j$ is convolutional.
(iii) *Library convention*: many neural-network libraries implement $\star$ while naming the operation convolution.

**Definition 6.1.3.** *Sparse Interactions, Sharing, and Equivariance*
For $x : \mathbb{Z}^d \to \mathbb{R}^{C_{\text{in}}}$, finite $W \subset \mathbb{Z}^d$, kernels $K_w \in \mathbb{R}^{C_{\text{out}} \times C_{\text{in}}}$, $b \in \mathbb{R}^{C_{\text{out}}}$, and pointwise $\sigma$, set $Lx(u) := \sigma(\sum_{w \in W} K_w x(u - w) + b)$ and $(\tau_a x)(u) := x(u - a)$.

(i) *Sparse interaction*: $Lx(u)$ depends only on $\{x(u - w) : w \in W\}$.
(ii) *Parameter sharing*: the same $(K_w)_{w \in W}$ and $b$ define $Lx(u)$ for every $u$.
(iii) *Translation equivariance*: $L(\tau_a x) = \tau_a(Lx)$ for every $a \in \mathbb{Z}^d$ where defined.

Pooling with finite $U$ applies $\rho$ to $\{Lx(u+w) : w \in U\}$; shift-stable $\rho$ gives local approximate invariance.

**Definition 6.1.4.** *Pooling, Padding, and Stride*
Let $x : \Omega \to \mathbb{R}^C$ on finite $\Omega \subset \mathbb{Z}^d$, with $W \subset \mathbb{Z}^d$, $p \in \mathbb{N}_0^d$, $s \in \mathbb{N}^d$, offset $r$, statistic $\rho : (\mathbb{R}^C)^W \to \mathbb{R}^{C'}$, and $\Omega^{(p)} := \Omega + \coprod_i \{-p_i, \dots, p_i\}$.

(i) *Padding*: an extension $\tilde{x} : \Omega^{(p)} \to \mathbb{R}^C$ with $\tilde{x}|_\Omega = x$ and a boundary rule on $\Omega^{(p)} \setminus \Omega$.
(ii) *Stride*: evaluation sites $\Omega_{s,r} := \{u \in \Omega^{(p)} : \forall i,\ u_i \equiv r_i \pmod{s_i},\ u + W \subset \Omega^{(p)}\}$.
(iii) *Pooling*: $P_{\rho,W,s,r}x : \Omega_{s,r} \ni u \mapsto \rho((\tilde{x}(u+w))_{w \in W}) \in \mathbb{R}^{C'}$.

**Remark 6.1.5.** *Geometric Extensions*
Let $X$ be a domain with local neighborhoods $\mathcal{N}(x) \subset X$ and symmetry group $G$.

(i) *Graph neural networks*: on $\mathcal{G} = (V, E)$, $h_v^{(\ell+1)} = \phi(h_v^{(\ell)}, \mathrm{Agg}_{u \in \mathcal{N}(v)}\, m_{uv}^{(\ell)})$ with invariant Agg.
(ii) *Equivariant networks*: $\Phi$ is $G$-equivariant if $\Phi(\rho_{\mathrm{in}}(g)x) = \rho_{\mathrm{out}}(g)\Phi(x)$ for every $g \in G$.
(iii) *Manifold architectures*: for a manifold $M$, layers use chart, tangent, or geodesic neighborhoods $\mathcal{N}(p) \subset M$.

Each variant builds locality or symmetry into the model class.

## 6.2 Sequence Models and Attention

Sequence models process ordered data where context and position matter.

**Definition 6.2.1.** *Recurrent Neural Network*
For dim $d_x, d_h, d_y$, parameters $\Theta = (W_x, W_h, b, \phi, \psi, h_0)$, $T \geq 1$, an RNN is a map $\Phi_T : (\mathbb{R}^{d_x})^T \to (\mathbb{R}^{d_y})^T$:

(i) *State update*: $h_t = \phi(W_x x_t + W_h h_{t-1} + b)$ for $t = 1, \dots, T$, with shared $(W_x, W_h, b)$.
(ii) *Readout*: $y_t = \psi(h_t)$ and $\Phi_T(x_{1:T}) = (y_1, \dots, y_T)$.

**Remark 6.2.2.** *Long-Term Dependencies*
For $h_t = F_t(h_{t-1}, x_t)$, set $J_t := \partial h_t / \partial h_{t-1}$ and $P_{s:T} := \prod_{t=s+1}^{T} J_t$.
Then $\nabla_{h_s} \mathcal{L} = P_{s:T}^\top \nabla_{h_T} \mathcal{L}$; $\|P_{s:T}\| \ll 1$ vanishes gradients and $\|P_{s:T}\| \gg 1$ explodes them.

(i) *Gating*: gates $g_t \in [0,1]^{d_h}$ interpolate between copy and update.
(ii) *Gradient clipping*: replace a gradient $g$ by $\min\{1, c/\|g\|\}g$, so the clipped norm is at most $c$.
(iii) *Skip paths*: use additive routes such as $h_t = h_{t-1} + F_t(h_{t-1}, x_t)$, giving transition Jacobian $I + \partial F_t / \partial h_{t-1}$.

These mechanisms keep the long-horizon Jacobian product numerically controlled.

**Definition 6.2.3.** *Attention*
For $Q \in \mathbb{R}^{n_q \times d}$, $K \in \mathbb{R}^{n_k \times d}$, $V \in \mathbb{R}^{n_k \times d_v}$, define $\mathrm{Attn}(Q, K, V) \in \mathbb{R}^{n_q \times d_v}$ row-wise:

(i) *Scores*: $S := d^{-1/2} Q K^\top \in \mathbb{R}^{n_q \times n_k}$.
(ii) *Weights*: $A := \mathrm{softmax}_{\mathrm{row}}(S)$, $A_{ij} := \exp(S_{ij}) / \sum_{\ell=1}^{n_k} \exp(S_{i\ell})$.
(iii) *Output*: $\mathrm{Attn}(Q, K, V) := AV$, equivalently $(\mathrm{Attn}(Q, K, V))_{i\cdot} = \sum_{j=1}^{n_k} A_{ij} V_{j\cdot}$.

**Remark 6.2.4.** *Attention as Routing*
For token matrix $X$, set $Q = XW_q$, $K = XW_k$, $V = XW_v$, $S = d^{-1/2} Q K^\top$, and $A = \mathrm{softmax}_{\mathrm{row}}(S)$.

(i) *Content routing*: $y_i = \sum_j A_{ij} v_j$ routes value vectors by query-key compatibility.
(ii) *Masking*: a mask $M_{ij} \in \{0, -\infty\}$ replaces $S$ by $S + M$, forcing $A_{ij} = 0$ for forbidden interactions.
(iii) *Positional encoding*: absolute codes use $x_i + e_i$; relative codes add a position term to $S_{ij}$.

Thus attention replaces a fixed recurrence neighborhood by data-dependent communication.

## 6.3 Transformers and Large-Scale Training

Transformers combine attention, feedforward blocks, normalization, and residual connections into architectures that scale well on sequence and multimodal data.

**Definition 6.3.1.** *Self-Attention*
For $X \in \mathbb{R}^{T \times d}$, head dimension $d_h$, and projection matrices $W_Q, W_K, W_V, W_O$:

(i) *Self-attention*: $Q = XW_Q$, $K = XW_K$, $V = XW_V$, and $\mathrm{SA}(X) := \mathrm{Attn}(Q, K, V)$.
(ii) *Multi-head attention*: with heads $H_r := \mathrm{Attn}(XW_Q^{(r)}, XW_K^{(r)}, XW_V^{(r)})$, set $\mathrm{MHA}(X) := [H_1 | \cdots | H_R] W_O$.

**Definition 6.3.2.** *Transformer Block*
For $X \in \mathbb{R}^{T\times d}$, let MHA be multi-head attention, FFN positionwise feedforward, and $\mathrm{LN}_1, \mathrm{LN}_2$ layer normalizations. A pre-norm block $B : \mathbb{R}^{T\times d} \to \mathbb{R}^{T\times d}$ is:

(i) *Attention sublayer*: $Z := X + \mathrm{MHA}(\mathrm{LN}_1 X)$.
(ii) *Feedforward sublayer*: $B(X) := Z + \mathrm{FFN}(\mathrm{LN}_2 Z)$.

Stacking blocks gives $B_L \circ \cdots \circ B_1(X)$.

**Remark 6.3.3.** *Large-Scale Training*
For data $\mathcal{D} = \{z_i\}_{i=1}^N$, loss $\ell_i(\theta)$, mini-batch $B_t$, and gradient estimate $\hat{g}_t := |B_t|^{-1} \sum_{i\in B_t} \nabla_\theta \ell_i(\theta_t)$:

(i) *Optimization*: choose update $\theta_{t+1} = U_t(\theta_t, \hat{g}_t)$, batch size $|B_t|$, and schedule $\eta_t$.
(ii) *Data*: sampling, filtering, weighting, and augmentation define the empirical training law $\widehat{P}_{\mathcal{D}}$.
(iii) *Numerics*: mixed precision uses low-precision arithmetic with scaling or high-precision accumulators.
(iv) *Parallelism*: workers compute local gradients $g_t^{(p)}$ and synchronize $\bar{g}_t = P^{-1} \sum_{p=1}^P g_t^{(p)}$.
(v) *Memory*: checkpointing stores selected activations and recomputes the rest during backpropagation.
(vi) *Evaluation*: validation loss and task metrics estimate out-of-sample performance.

**Remark 6.3.4.** *Inductive Bias*
Let $I = \{1, \dots, T\}$ index tokens, $W_i \subset I$ be a local window, and $A_{ij}(x)$ be attention weights.

(i) *Locality*: convolution uses $y_i = F((x_j)_{j\in W_i})$.
(ii) *Global interaction*: attention uses $y_i = \sum_{j\in I} A_{ij}(x) V x_j$.
(iii) *Domain-specific bias*: impose masks, recurrence, sparsity, or $\Phi(\rho_{\mathrm{in}}(g)x) = \rho_{\mathrm{out}}(g)\Phi(x)$.

Useful bias matches the task's locality, symmetry, and dependency scale.

## 6.4 Generative Models

Generative modeling learns distributions, simulators, or conditional samplers rather than only point predictions.

**Definition 6.4.1.** *Latent-Variable Generative Model*
Let $Z \in \mathcal{Z}$, $X \in \mathcal{X}$, prior $p(z)$, decoder $p_\theta(x \mid z)$, and data law $P_{\mathrm{data}}$.

(i) *Joint law*: $p_\theta(x, z) := p_\theta(x \mid z)p(z)$.
(ii) *Marginal law*: $p_\theta(x) := \int_{\mathcal{Z}} p_\theta(x \mid z)p(z)\, dz$.
(iii) *Learning*: choose $\theta$ so $p_\theta$ approximates $P_{\mathrm{data}}$, e.g. by maximizing $\sum_{i=1}^n \log p_\theta(x_i)$.

**Definition 6.4.2.** *Variational Autoencoder*
For prior $p(z)$, decoder $p_\theta(x \mid z)$, encoder $q_\phi(z \mid x)$, and data $\{x_i\}_{i=1}^n$:

(i) *Generative model*: $p_\theta(x, z) := p_\theta(x \mid z)p(z)$.
(ii) *Variational posterior*: $q_\phi(z \mid x)$ approximates $p_\theta(z \mid x)$.
(iii) *ELBO*: $\mathcal{L}(\theta, \phi; x) := \mathbb{E}_{q_\phi(z|x)}[\log p_\theta(x \mid z)] - \mathrm{KL}(q_\phi(z \mid x)\|p(z))$.
(iv) *Training*: maximize $\sum_{i=1}^n \mathcal{L}(\theta, \phi; x_i)$ over $(\theta, \phi)$.

**Definition 6.4.3.** *Generative Adversarial Network*
Let $Z \sim p_Z$, data law $P_{\mathrm{data}}$, generator $G_\theta : \mathcal{Z} \to \mathcal{X}$, and discriminator $D_\psi : \mathcal{X} \to (0, 1)$.

(i) *Generator*: $G_\theta$ induces $P_\theta := (G_\theta)_{\#} p_Z$ on $\mathcal{X}$.
(ii) *Discriminator*: $D_\psi(x)$ estimates the probability that $x \sim P_{\mathrm{data}}$ rather than $x \sim P_\theta$.
(iii) *Adversarial game*: $\min_\theta \max_\psi \mathbb{E}_{x\sim P_{\mathrm{data}}} \log D_\psi(x) + \mathbb{E}_{z\sim p_Z} \log(1 - D_\psi(G_\theta(z)))$.

**Definition 6.4.4.** *Diffusion Model*
Let $X_0 \sim P_{\mathrm{data}}$ and $X_0, \dots, X_T$ be a Markov chain on $\mathcal{X}$.

(i) *Forward noising*: fixed kernels $q_t(x_t \mid x_{t-1})$ define $q(x_{1:T} \mid x_0) = \prod_{t=1}^T q_t(x_t \mid x_{t-1})$.
(ii) *Reverse model*: learned kernels $p_\theta(x_{t-1} \mid x_t)$ define $p_\theta(x_{0:T}) = p_T(x_T) \prod_{t=1}^T p_\theta(x_{t-1} \mid x_t)$.
(iii) *Training*: fit $\theta$ so $p_\theta(x_{t-1} \mid x_t)$ approximates the reverse conditionals of $q$.
(iv) *Sampling*: draw $X_T \sim p_T$ and iterate $X_{t-1} \sim p_\theta(\cdot \mid X_t)$ for $t = T, \dots, 1$.

## 6.5 Reinforcement Learning and Decision Systems

Reinforcement learning studies agents whose actions affect future observations and rewards.

**Definition 6.5.1.** *Markov Decision Process*
An MDP is a tuple $(\mathcal{S}, \mathcal{A}, P, r, \gamma)$ with:

(i) *State-action spaces*: states $s \in \mathcal{S}$ and actions $a \in \mathcal{A}$.

(ii) *Transition law*: $P(\cdot \mid s,a) \in \mathcal{P}(\mathcal{S})$ and $S_{t+1} \sim P(\cdot \mid S_t, A_t)$.
(iii) *Reward and discount*: $r : \mathcal{S} \times \mathcal{A} \to \mathbb{R}$ and $\gamma \in [0,1)$.
(iv) *Policy*: $\pi(\cdot \mid s) \in \mathcal{P}(\mathcal{A})$ and $A_t \sim \pi(\cdot \mid S_t)$.

**Definition 6.5.2.** *Value Functions*
For policy $\pi$, define the discounted return $G_t := \sum_{k=0}^{\infty} \gamma^k r(S_{t+k}, A_{t+k})$.

(i) *State value*: $V^\pi(s) := \mathbb{E}_\pi[G_0 \mid S_0 = s]$.
(ii) *Action value*: $Q^\pi(s,a) := \mathbb{E}_\pi[G_0 \mid S_0 = s, A_0 = a]$.

**Definition 6.5.3.** *Bellman Equation*
For an MDP $(\mathcal{S}, \mathcal{A}, P, r, \gamma)$ and value function $V : \mathcal{S} \to \mathbb{R}$:

(i) *Policy Bellman equation*: $V^\pi(s) = \sum_a \pi(a \mid s)(r(s,a) + \gamma \sum_{s'} P(s' \mid s,a) V^\pi(s'))$.
(ii) *Optimal Bellman equation*: $V^\star(s) = \max_a (r(s,a) + \gamma \sum_{s'} P(s' \mid s,a) V^\star(s'))$.

**Remark 6.5.4.** *RL as a Learning System*
Let $\tau = (S_0, A_0, R_0, \ldots)$ denote a trajectory and $P^\pi$ its law under policy $\pi$.

(i) *Prediction*: estimate $V^\pi$, $Q^\pi$, or transition/reward models from trajectories.
(ii) *Feedback*: policy changes alter the data law, $\pi \mapsto P^\pi$.
(iii) *Optimization*: choose $\pi$ to maximize $J(\pi) := \mathbb{E}_{P^\pi} \sum_{t \geq 0} \gamma^t R_t$.
(iv) *Exploration*: sample actions to reduce uncertainty about rewards, transitions, or values.
(v) *Evaluation*: validate under sequential dependence and policy-induced distribution shift.

# 7 Learning Systems and Evaluation

## 7.1 Data Splitting, Validation, and Benchmarking

Evaluation estimates how the trained system behaves on data not used to fit it.

**Definition 7.1.1.** *Training, Validation, and Test Splits*
Let $\mathcal{D} = \mathcal{D}_{\mathrm{tr}} \dot\cup \mathcal{D}_{\mathrm{val}} \dot\cup \mathcal{D}_{\mathrm{te}}$ be a disjoint partition of $\{(x_i, y_i)\}_{i=1}^n$, and fix a loss $\ell$.

(i) Training: for each $\lambda \in \Lambda$, fit $\widehat{f}_\lambda$ using only $\mathcal{D}_{\mathrm{tr}}$.
(ii) Validation: set $\widehat{\lambda} \in \arg\min_{\lambda \in \Lambda} \widehat{R}_{\mathrm{val}}(\lambda)$, where $\widehat{R}_{\mathrm{val}}(\lambda) = |\mathcal{D}_{\mathrm{val}}|^{-1} \sum_{(x,y) \in \mathcal{D}_{\mathrm{val}}} \ell(\widehat{f}_\lambda(x), y)$.
(iii) Test: after $\widehat{\lambda}$ is fixed, report $\widehat{R}_{\mathrm{te}} = |\mathcal{D}_{\mathrm{te}}|^{-1} \sum_{(x,y) \in \mathcal{D}_{\mathrm{te}}} \ell(\widehat{f}_{\widehat{\lambda}}(x), y)$ as the final performance estimate.

If $\mathcal{D}_{\mathrm{val}}$ is unavailable, cross-validation or bootstrap resampling approximates validation.
Any pre-reporting use of $\mathcal{D}_{\mathrm{te}}$ makes $\widehat{R}_{\mathrm{te}}$ adaptive and typically optimistic.




Figure 262: Train/validation/test split: $\mathcal{D} = \mathcal{D}_{\mathrm{tr}} \sqcup \mathcal{D}_{\mathrm{val}} \sqcup \mathcal{D}_{\mathrm{te}}$, with $\mathcal{D}_{\mathrm{tr}}$ fitting $\widehat{f}_\lambda$, $\mathcal{D}_{\mathrm{val}}$ selecting $\widehat{\lambda}$, and $\mathcal{D}_{\mathrm{te}}$ estimating final risk $\widehat{R}_{\mathrm{te}}$.

**Definition 7.1.2.** *Model Selection and Model Assessment*
Let $\{\widehat{f}_\lambda : \lambda \in \Lambda\}$ be candidates, let $\widehat{R}_{\mathrm{val}}(\lambda)$ be validation error, and let $R(f) = \mathbb{E}\ell(f(X), Y)$.

(i) *Model selection*: choose $\widehat{\lambda} \in \Lambda$, typically $\widehat{\lambda} \in \arg\min_{\lambda \in \Lambda} \widehat{R}_{\mathrm{val}}(\lambda)$, and output $\widehat{f}_{\widehat{\lambda}}$.
(ii) *Model assessment*: after $\widehat{\lambda}$ is fixed, estimate $R(\widehat{f}_{\widehat{\lambda}})$ using data independent of training and selection.

Reusing selection data for assessment yields adaptive, typically optimistic error.

**Definition 7.1.3.** *Conditional and Expected Test Error*
Let $\widehat{f}_{\mathcal{T}}$ be trained on random $\mathcal{T}$, and let $(X^0, Y^0)$ be an independent test point.

(i) *Conditional test error*: $\mathrm{Err}_{\mathcal{T}} = \mathbb{E}_{X^0, Y^0}[\ell(\widehat{f}_{\mathcal{T}}(X^0), Y^0) \mid \mathcal{T}]$.
(ii) *Expected test error*: $\mathrm{Err} = \mathbb{E}_{\mathcal{T}} \mathrm{Err}_{\mathcal{T}}$.

Reusable error estimators usually target Err, while $\mathrm{Err}_{\mathcal{T}}$ is the error conditional on the realized training set.

**Method 7.1.4.** *K-Fold Cross-Validation*
Let $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$, loss $\ell$, candidates $\Lambda$, and fold map $\kappa : \{1, \dots, n\} \to \{1, \dots, K\}$ be fixed.

(i) For each $\lambda \in \Lambda$ and $k = 1, \dots, K$, fit $\widehat{f}_\lambda^{-k}$ using only observations with $\kappa(i) \neq k$.
(ii) For each $\lambda \in \Lambda$, compute $\mathrm{CV}(\lambda) = n^{-1} \sum_{i=1}^n \ell(\widehat{f}_\lambda^{-\kappa(i)}(x_i), y_i)$.
(iii) Choose $\widehat{\lambda} \in \arg\min_{\lambda \in \Lambda} \mathrm{CV}(\lambda)$, or the one-standard-error candidate when that rule is specified.
(iv) Refit $\widehat{f}_{\widehat{\lambda}}$ on all of $\mathcal{D}$.

Leave-one-out has $K = n$; $K = 5$ or 10 often lowers variance and computation but fits on smaller training folds.

**Remark 7.1.5.** *What Cross-Validation Estimates*
For learning rule $\mathcal{A}$, let $\mathrm{Err}_{\mathcal{A}}(m) = \mathbb{E}_{\mathcal{T}_m, X^0, Y^0} \ell(\mathcal{A}(\mathcal{T}_m)(X^0), Y^0)$. $K$-fold cross-validation targets roughly $\mathrm{Err}_{\mathcal{A}}(n(1 - 1/K))$, not the conditional error of $\mathcal{A}(\mathcal{D})$. Leave-one-out has $m = n - 1$ but dependent losses; fixed $K = 5$ or 10 lowers computation and often variance, with possible upward bias.

**Definition 7.1.6.** *Benchmark*
A benchmark is a fixed evaluation protocol $\mathcal{B} = (\mathcal{D}, \Pi, \mathcal{S}, m, \mathcal{R})$.

(i) *Data*: $\mathcal{D}$ specifies the population, sample, or data source.
(ii) *Preprocessing*: $\Pi$ specifies transformations, filtering, feature construction, and allowed fitting scope.
(iii) *Splitting*: $\mathcal{S}$ specifies train-validation-test splits or resampling schemes.
(iv) *Metric*: $m$ specifies the reported score.
(v) *Resources*: $\mathcal{R}$ specifies allowed external data, models, compute, and manual intervention.

Two reported scores are directly comparable only when computed under the same $\mathcal{B}$.

**Remark 7.1.7.** *Metric Choice*
Choose $m(f; \mathcal{D}_{\mathrm{te}})$ so its ordering of candidate rules matches the deployment objective $U(f)$.
Accuracy, log loss, calibration, ranking quality, cost, and latency need not agree.

**Remark 7.1.8.** *Data Leakage*
For protected evaluation data $E$, leakage is dependence of any pre-evaluation choice $Z_{\mathrm{pre}}$ on labels or covariates in $E$. Then error measured on $E$ is adaptive, not honest out-of-sample error.

**Method 7.1.9.** *Cross-Validation with Screening or Preprocessing*
For a fold map $\kappa : \{1, \dots, n\} \to \{1, \dots, K\}$, set $\mathcal{D}_k = \{(x_i, y_i) : \kappa(i) = k\}$ and $\mathcal{D}_{-k} = \mathcal{D} \setminus \mathcal{D}_k$.
Data-dependent pipeline choices must be fold-local:

(i) For each $k$, learn preprocessing, screening, feature selection, and tuning from $\mathcal{D}_{-k}$ only.
(ii) Fit $\widehat{f}^{-k}$ on $\mathcal{D}_{-k}$ after applying exactly those fold-local choices.
(iii) Predict and score $\mathcal{D}_k$ using $\widehat{f}^{-k}$ and the same fold-local choices.
(iv) Average the held-out losses over $k = 1, \dots, K$.

Full-data supervised choices leak held-out labels into $\widehat{f}^{-k}$ and bias cross-validation downward.
The protocol should state whether unsupervised preprocessing is fold-local or full-covariate.

**Method 7.1.10.** *Bootstrap Estimates of Prediction Error*
Let $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$; draw bootstrap samples $S^{*b}$ and fit $\widehat{f}^{*b}$; set $C^{-i} = \{b : (x_i, y_i) \notin S^{*b}\}$, $I_{\mathrm{oob}} = \{i : C^{-i} \neq \varnothing\}$, and $\overline{\mathrm{err}} = n^{-1} \sum_i \ell(\widehat{f}(x_i), y_i)$.

(i) *Naive bootstrap*: $\widehat{\mathrm{Err}}_{\mathrm{boot}} = B^{-1} \sum_{b=1}^B n^{-1} \sum_{i=1}^n \ell(\widehat{f}^{*b}(x_i), y_i)$.
(ii) *Out-of-bag bootstrap*: $\widehat{\mathrm{Err}}^{(1)} = |I_{\mathrm{oob}}|^{-1} \sum_{i \in I_{\mathrm{oob}}} |C^{-i}|^{-1} \sum_{b \in C^{-i}} \ell(\widehat{f}^{*b}(x_i), y_i)$.
(iii) *.632 bootstrap*: $\widehat{\mathrm{Err}}^{(.632)} = .368\, \overline{\mathrm{err}} + .632\, \widehat{\mathrm{Err}}^{(1)}$.
(iv) *.632+ bootstrap*: $\widehat{\mathrm{Err}}^{(.632+)} = (1 - w_+)\overline{\mathrm{err}} + w_+ \widehat{\mathrm{Err}}^{(1)}$, with $w_+ \in [.632, 1]$ increase with overfitting.

Naive reuse is most optimistic; out-of-bag and .632 variants reduce reuse bias.

**Remark 7.1.11.** *Limits of Reuse-Based Assessment*
Write $\widehat{R}_{\mathrm{reuse}}(\lambda) = R(\widehat{f}_\lambda) + b_\lambda + \varepsilon_\lambda$. Nearly constant $b_\lambda$ can preserve candidate comparisons despite biased levels.
After choosing $\widehat{\lambda}$, estimating $R(\widehat{f}_{\widehat{\lambda}})$ requires data independent of all fitting, preprocessing, tuning, and iteration.

## 7.2 Robustness, Shift, and Distributional Uncertainty

Learning systems are deployed under distributions that may differ from the training law.

**Definition 7.2.1.** *Distribution Shift*
Let $P$ be training law, $Q$ deployment law on $(X, Y)$. Distribution shift means $Q \neq P$; common cases are:

(i) *Covariate shift*: $Q_X \neq P_X$ and $Q_{Y|X} = P_{Y|X}$.

(ii) *Label shift*: $Q_Y \neq P_Y$ and $Q_{X|Y} = P_{X|Y}$.
(iii) *Concept shift*: $Q_{Y|X} \neq P_{Y|X}$.

**Definition 7.2.2.** *Adversarial Example*
Let $\mathcal{F}$ be a predictor class, $\ell$ a loss, $d_f : \mathcal{X} \to \mathcal{D}$ the decision rule induced by $f \in \mathcal{F}$, $\Delta(x)$ the admissible perturbation set at $x$, and $\mathcal{H}_y \subseteq \mathcal{D} \times \mathcal{D}$ the harmful decision-change relation for label $y$.

(i) *Adversarial example*: for fixed $f \in \mathcal{F}$ and $(x, y)$, any $x' = x + \delta$ with $\delta \in \Delta(x)$ and $\big(\ell(f(x'), y) > \ell(f(x), y)\big) \vee \big((d_f(x), d_f(x')) \in \mathcal{H}_y\big)$.
(ii) *Adversarial training*: for $S = \{(x_i, y_i)\}_{i=1}^n$, choose $\widehat{f}_{\text{adv}} \in \arg\min_{f \in \mathcal{F}} n^{-1} \sum_{i=1}^{n} \sup_{\delta \in \Delta(x_i)} \ell(f(x_i + \delta), y_i)$.
(iii) *Neural perturbation form*: minimize $\ell(y, f_\theta(x + \delta^\star))$, where $\delta^\star \in \arg\max_{\|\delta\| \leq \rho} \ell(y, f_\theta(x + \delta))$.

**Definition 7.2.3.** *Distributionally Robust Optimization*
Let $\mathcal{F}$ be a hypothesis class, $\ell$ a loss, and $\mathcal{U} \subseteq \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ an uncertainty set of deployment laws.

(i) *Robust risk*: for $Q \in \mathcal{U}$ and $f \in \mathcal{F}$, $R_Q(f) = \mathbb{E}_{(X,Y)\sim Q} \ell(f(X), Y)$ and $R_{\mathcal{U}}(f) = \sup_{Q \in \mathcal{U}} R_Q(f)$.
(ii) *DRO solution*: any $\widehat{f}_{\text{DRO}} \in \arg\min_{f \in \mathcal{F}} R_{\mathcal{U}}(f)$.

**Remark 7.2.4.** *Uncertainty*
Let $\mathcal{T}$ be the training data, $P_{\text{tr}}$ the training law, and $P_{\text{dep}}$ the deployment law.

(i) *Aleatoric uncertainty*: irreducible variability in $Y \mid X = x$ under the data-generating law.
(ii) *Epistemic uncertainty*: uncertainty in $f$ from finite $\mathcal{T}$, misspecification, or nonidentifiability.
(iii) *Out-of-distribution uncertainty*: uncertainty from $P_{\text{dep}} \neq P_{\text{tr}}$ or inputs outside the training support.

Robust systems calibrate probabilities and abstain when uncertainty exceeds a policy threshold.

## 7.3 Interpretability, Fairness, and Safety

Evaluation also asks whether a model can be understood, audited, and constrained in the setting where it is used.

**Definition 7.3.1.** *Interpretability*
Let $\mathcal{F}, \mathcal{C}, \mathcal{E}, \mathcal{A}, \mathcal{B}$ denote predictor, query-context, explanation, audience, and audit-task spaces.

(i) *Interpretability specification*: $\mathcal{I} = (\mathcal{E}, a, b, \Phi)$ with $(a, b) \in \mathcal{A} \times \mathcal{B}$ and $\Phi : \mathcal{F} \times \mathcal{C} \to \mathcal{E}$.
(ii) *Explanation*: for $(f, c) \in \mathcal{F} \times \mathcal{C}$, $e = \Phi(f, c)$.

Typical codomains $\mathcal{E}$ encode attributions, counterfactuals, surrogate rules, or representation summaries.

**Definition 7.3.2.** *Fairness Criteria*
Let $A$ be a protected attribute, $\widehat{Y}$ a decision, $S$ a score, and $Y$ the outcome.
A group fairness criterion imposes equalities or tolerances across values of $A$; common examples are:

(i) *Demographic parity*: $\mathbb{P}(\widehat{Y} = 1 \mid A = a)$ is constant in $a$.
(ii) *Equalized odds*: $\mathbb{P}(\widehat{Y} = 1 \mid Y = y, A = a)$ is constant in $a$ for each $y$.
(iii) *Group calibration*: $\mathbb{P}(Y = 1 \mid S = s, A = a) = s$ for each relevant $s$ and $a$.

These criteria need not be mutually compatible; the chosen condition should match the policy goal.

**Definition 7.3.3.** *Safety Constraint*
An additional requirement $f \in \mathcal{C}$ or $R_{\text{safe}}(f) \leq \tau$ beyond average-case loss is a *safety constraint*.
$\mathcal{C}$ encodes abstention, monotonicity, action bounds, review triggers, or worst-case risk limits.

**Remark 7.3.4.** *Beyond Accuracy*
For a deployed predictor $f$, evaluation beyond accuracy is a specification $\mathcal{E} = (\mathcal{I}, \mathcal{F}, \mathcal{C})$.

(i) *Interpretability*: $\mathcal{I}$ specifies the explanation object, audience, and audit protocol.
(ii) *Fairness*: $\mathcal{F}$ specifies protected attributes, group metrics, and required parity or error constraints.
(iii) *Safety*: $\mathcal{C}$ specifies forbidden actions, abstention rules, human review triggers, or worst-case risk bounds.

A scalar score is incomplete unless the relevant components of $\mathcal{E}$ are stated.

## 7.4 Deployment, Monitoring, and Feedback Loops

Deployment turns a fitted model into part of a larger stochastic system.

**Definition 7.4.1.** *Monitoring*
Given deployed $f$ and stream $Z_t = (X_t, \widehat{Y}_t, Y_t^*)$, monitoring outputs $(M_t, a_t)$ with $a_t \in \{0, 1\}$.

(i) *Distribution*: compare $P_t(X)$ and $P_t(\widehat{Y})$ with validation baselines.
(ii) *Performance*: estimate loss, calibration, and subgroup error when labels $Y_t^*$ arrive.
(iii) *Operations*: track latency, missingness, failures, and data-pipeline changes.

An alarm $a_t = 1$ indicates that validation assumptions or service constraints may no longer hold.

**Definition 7.4.2.** *Feedback Loop*
A deployed policy $A_t = \pi(f(X_t), H_t)$ is a *feedback loop* if $P(X_{t+1}, Y_{t+1} \mid A_t) \neq P(X_{t+1}, Y_{t+1})$.
Future data are then endogenous to $f$.

**Method 7.4.3.** *Model Update Policy*
Given deployed $f_t$, data $\mathcal{D}_{\leq t}$, and diagnostics $M_t$, a model update policy maps $f_t$ to $f_{t+1}$:

(i) *Trigger*: retrain only when a scheduled time, alarm $M_t$, or performance threshold calls for it.
(ii) *Eligibility*: choose the data subset $\mathcal{D}_{\text{upd}} \subseteq \mathcal{D}_{\leq t}$ allowed for training.
(iii) *Validation*: accept candidate $\widetilde{f}$ only if it passes the benchmark and safety criteria against $f_t$.
(iv) *Deployment*: replace $f_t$ by $\widetilde{f}$ under a rollout plan and rollback rule.

The policy determines future training data and evaluation conditions.

**Remark 7.4.4.** *Human-System Boundary*
For human-supported decisions, the deployed policy is $\pi = (f, I, A, O, M, G)$, not just predictor $f$.

(i) *Interface*: $I$ determines what predictions, explanations, and uncertainty are shown.
(ii) *Action rule*: $A$ maps model output and human input to decisions.
(iii) *Override*: $O$ specifies when humans may ignore or modify $f$.
(iv) *Monitoring*: $M$ tracks errors, drift, latency, and misuse.
(v) *Governance*: $G$ assigns review, audit, and accountability.

System evaluation should target $\pi$ rather than $f$ alone.

# Part XXII
# Classical Supervised Learning

## 1 Linear Prediction and Generalized Linear Models

### 1.1 Linear Predictors and Least Squares

Linear prediction reduces inputs to scores of the form "intercept plus weighted features." Regression reads scores as predictions; classification compares or transforms them. Its appeal is geometric: projections, hyperplanes, and feature maps.

**Definition 1.1.1.** *Linear Predictor*
For $x \in \mathbb{R}^p$, $w \in \mathbb{R}^p$, and $b \in \mathbb{R}$, the *linear predictor* is $f_{w,b}(x) := w^\top x + b$.
Regression uses $f_{w,b}$ directly; classification transforms it by $g$. Score level sets are parallel hyperplanes.

**Definition 1.1.2.** *Least-Squares Linear Regression*
For $X \in \mathbb{R}^{n\times d}$ and $y \in \mathbb{R}^n$, *least-squares linear regression* chooses $\widehat{\beta} \in \operatorname*{argmin}_{\beta\in\mathbb{R}^d} \|y - X\beta\|_2^2$ and $\widehat{y} := X\widehat{\beta}$.

If $X^\top X$ is nonsingular, then $\widehat{\beta} = (X^\top X)^{-1}X^\top y$. Columns of $X$ may be raw variables, transformations, indicators, interactions, or basis functions; linearity is in $\beta$.

Geometrically, least squares moves $y$ to the closest point in $\mathrm{col}(X)$.

**Method 1.1.3.** *Multiple-Output Least Squares*
For $Y \in \mathbb{R}^{n\times K}$ and $B \in \mathbb{R}^{d\times K}$, set $Y = XB + E$.

(i) *Least squares*: If $\mathrm{rank}(X) = d$, then $\widehat{B} = (X^\top X)^{-1}X^\top Y$.
(ii) *Decoupling*: Writing $Y = (y_1, \dots, y_K)$ gives $\widehat{B} = (\widehat{\beta}_1, \dots, \widehat{\beta}_K)$, where each $\widehat{\beta}_k$ is the OLS fit of $y_k$ on $X$.

Shared responses can be pooled by low-rank $B$ or common-direction shrinkage.

**Method 1.1.4.** *Canonical Correlation and Reduced-Rank Regression*

(i) *CCA*: Choose $(v_j, u_j)$ with large $\mathrm{corr}(Xv_j, Yu_j)$, orthogonal across $j$; at most $\min(d, K)$ pairs.
(ii) *Rank constraint*: Fit $Y \approx XB$ with $\mathrm{rank}(B) \le m$, restricting fitted responses to $m$ dimensions.
(iii) *Objective*: $\widehat{B}_m^{\mathrm{rr}} \in \operatorname*{argmin}_{\mathrm{rank}(B)\le m} \sum_i (y_i - B^\top x_i)^\top \widehat{\Sigma}^{-1}(y_i - B^\top x_i)$.
(iv) *Role of $m$*: $m = K$ gives separate OLS; smaller $m$ pools shared response structure.

Reduced rank assumes the predictable response variation lies in a low-dimensional subspace.

### 1.2 Linear Classification and Generalized Linear Models

**Definition 1.2.1.** *Discriminant Functions and Linear Boundaries*
For $\mathcal{G} = \{1, \dots, K\}$ and scores $\delta_k : \mathcal{X} \to \mathbb{R}$, set $\widehat{G}(x) \in \mathrm{argmax}_{k\in\mathcal{G}} \delta_k(x)$. Only score differences matter.

(i) *Pairwise boundary*: The $k$ versus $\ell$ boundary lies in $\{x : \delta_k(x) = \delta_\ell(x)\}$.
(ii) *Linear boundary*: If $\delta_k - \delta_\ell$ is affine in $x$ for all $k, \ell$, then all pairwise decision boundaries are hyperplanes.
(iii) *Binary log-odds*: If $\log \frac{\mathbb{P}(G=1|X=x)}{\mathbb{P}(G=2|X=x)} = \beta_0 + \beta^\top x$, then the boundary is $\{x : \beta_0 + \beta^\top x = 0\}$.

**Method 1.2.2.** *Indicator-Matrix Regression Classifier*
For classes $g_i \in \{1, \dots, K\}$, set $t_k = e_k \in \mathbb{R}^K$, $Y_i = t_{g_i}$, and $\widetilde{x}_i^\top = (1, x_i^\top)$, coding labels as Euclidean targets.

(i) *Least-squares fit*: Choose $\widehat{B} \in \mathrm{argmin}_B \sum_{i=1}^n \left\|Y_i - B^\top \widetilde{x}_i\right\|_2^2$; if $\widetilde{X}^\top\widetilde{X}$ is nonsingular, $\widehat{B} = (\widetilde{X}^\top\widetilde{X})^{-1}\widetilde{X}^\top Y$.
(ii) *Scores*: For a new $x$, set $\widetilde{x} = (1, x^\top)^\top$ and $\widehat{f}(x) := \widehat{B}^\top\widetilde{x}$.
(iii) *Classification*: Predict $\widehat{G}(x) \in \mathrm{argmax}_k \widehat{f}_k(x)$, equivalently nearest $t_k$.

With an intercept, $\sum_k \widehat{f}_k(x) = 1$, but $\widehat{f}_k(x)$ may leave $[0, 1]$ and, for $K \ge 3$, *mask* middle classes.

**Definition 1.2.3.** *Generalized Linear Model*
A *generalized linear model* gives $Y \mid X = x$ an exponential-family law and makes a linked mean linear: $g(\mu(x)) = w^\top x + b$. Logistic regression uses the Bernoulli law and logit link $g(\mu) = \log \frac{\mu}{1-\mu}$.

Thus a GLM separates score, mean scale, and likelihood.

**Definition 1.2.4.** *Multiclass Logistic Regression*
With class $K$ as baseline, set $\eta_k(x) := \beta_{k0} + \beta_k^\top x = \log \frac{\mathbb{P}(G=k \mid X=x)}{\mathbb{P}(G=K \mid X=x)}$ for $k < K$.

(i) *Softmax probabilities*: For $k < K$, $p_k(x;\theta) := \frac{\exp(\eta_k(x))}{1+\sum_{\ell=1}^{K-1}\exp(\eta_\ell(x))}$, and $p_K(x;\theta) := \frac{1}{1+\sum_{\ell=1}^{K-1}\exp(\eta_\ell(x))}$.

(ii) *Likelihood*: Fit $\theta$ by maximising $\ell(\theta) := \sum_{i=1}^{n} \log p_{g_i}(x_i;\theta)$.

The softmax gives $p(x;\theta) \in \Delta^{K-1}$; the reference class changes only the parametrisation.

**Method 1.2.5.** *Binary Logistic Regression by IRLS*
For $y_i \in \{0,1\}$ and intercept-augmented rows $x_i^\top$, set $p_i(\beta) := (1+\exp(-x_i^\top\beta))^{-1}$.

(i) *Derivatives*: With $p = (p_i)$ and $W := \operatorname{diag}(p_i(1-p_i))$, $\nabla\ell(\beta) = X^\top(y-p)$ and $\nabla^2\ell(\beta) = -X^\top W X$.
(ii) *Adjusted response*: At $\beta^{\text{old}}$, set $z := X\beta^{\text{old}} + W^{-1}(y-p)$.
(iii) *IRLS update*: Set $\beta^{\text{new}} := (X^\top W X)^{-1} X^\top W z$.

IRLS is Newton's method as weighted least squares.
The log-likelihood is concave when the MLE is finite; step-halving controls Newton overshoot.

**Method 1.2.6.** *$L_1$-Regularized Logistic Regression*
With unpenalized intercept, maximize $\ell_\lambda(\beta_0,\beta) := \sum_{i=1}^{n}\{y_i\eta_i - \log(1+\exp(\eta_i))\} - \lambda\sum_{j=1}^{p}|\beta_j|$, where $\eta_i := \beta_0 + x_i^\top\beta$.

(i) *Probabilities*: Set $p_i := (1+\exp(-\eta_i))^{-1}$ and residual $r := y - p$.
(ii) *KKT conditions*: If $\widehat{\beta}_j \neq 0$, then $x_j^\top r = \lambda\operatorname{sign}(\widehat{\beta}_j)$; if $\widehat{\beta}_j = 0$, then $\left|x_j^\top r\right| \le \lambda$.

The KKT conditions are thresholded feature–residual correlations.

## 1.3 Discriminant Analysis and Naive Bayes

**Method 1.3.1.** *Gaussian Discriminant Analysis*
If $X \mid G = k$ has density $f_k$ and prior $\pi_k$, then $\mathbb{P}(G=k \mid X=x) = \frac{f_k(x)\pi_k}{\sum_{\ell=1}^{K} f_\ell(x)\pi_\ell}$.

(i) *Score*: If $X \mid G = k \sim N(\mu_k, \Sigma_k)$, compare $\delta_k(x) := -\frac{1}{2}\log|\Sigma_k| - \frac{1}{2}(x-\mu_k)^\top\Sigma_k^{-1}(x-\mu_k) + \log\pi_k$.
(ii) *LDA*: If $\Sigma_k = \Sigma$ for all $k$, equivalent linear scores are $\delta_k^{\text{LDA}}(x) := x^\top\Sigma^{-1}\mu_k - \frac{1}{2}\mu_k^\top\Sigma^{-1}\mu_k + \log\pi_k$.
(iii) *QDA*: If $\Sigma_k$ varies with $k$, the quadratic term remains and the rule is quadratic discriminant analysis.

Estimate $\pi_k, \mu_k, \Sigma_k$ from training data; pooled covariance gives LDA and classwise covariances give QDA.
Shared covariance makes the boundaries linear.

**Method 1.3.2.** *Regularized Discriminant Analysis*
For $0 \le \alpha \le 1$, set $\widehat{\Sigma}_k(\alpha) := \alpha\widehat{\Sigma}_k + (1-\alpha)\widehat{\Sigma}$.

(i) *Endpoints*: $\alpha = 0$ gives LDA and $\alpha = 1$ gives QDA.
(ii) *Scalar shrinkage*: For $0 \le \gamma \le 1$, replace $\widehat{\Sigma}$ by $\widehat{\Sigma}(\gamma) := \gamma\widehat{\Sigma} + (1-\gamma)\widehat{\sigma}^2 I$.

Shrinkage reduces variance when covariance estimation is unstable.

**Definition 1.3.3.** *Naive Bayes Classifier*
For $x = (x_1,\dots,x_p)$, assume $f_j(x) = \prod_{k=1}^{p} f_{jk}(x_k)$ for each class $j$.

(i) *Posterior*: With priors $\pi_j$, $p_j(x) := \mathbb{P}(G=j \mid X=x) \propto \pi_j \prod_{k=1}^{p} f_{jk}(x_k)$.

(ii) *Additive log odds*: Relative to class $J$, $\log\frac{p_\ell(x)}{p_J(x)} = \log\frac{\pi_\ell}{\pi_J} + \sum_{k=1}^{p}\log\frac{f_{\ell k}(x_k)}{f_{Jk}(x_k)}$.

On the log scale, features contribute additively.
Even misspecified independence can reduce variance enough to classify well near boundaries.

**Remark 1.3.4.** *Least Squares Versus LDA*
For $K = 2$, least-squares regression on any distinct label coding gives $\widehat{\beta} \propto \widehat{\Sigma}^{-1}(\widehat{\mu}_2 - \widehat{\mu}_1)$, the LDA direction.
The cutpoint may differ unless classes are balanced or tuned separately.
For $K > 2$, LDA uses pooled-Mahalanobis centroid distances and avoids masking.

**Remark 1.3.5.** *Logistic Regression Versus LDA*
Both can yield linear log-odds. Logistic regression models class probabilities directly and leaves $P_X$ free; LDA fits a joint Gaussian mixture for $(X, G)$ and uses Bayes' rule.

**Remark 1.3.6.** *Whitening and Reduced-Rank LDA*
With common covariance $W$, LDA compares $W^{-1/2}x$ to whitened centroids, adjusted by $\log \pi_k$.
The whitened centroids span dimension at most $K-1$, so LDA reduces to this centroid subspace.
Fisher directions maximize $\frac{a^\top Ba}{a^\top Wa}$, equivalently solve $Ba = \lambda Wa$, where $B$ is between-class scatter.

**Remark 1.3.7.** *Feature Maps*
A feature map $\phi : \mathcal{X} \to \mathcal{H}$ gives linear prediction in feature space, $f(x) = w^\top \phi(x) + b$, which is generally nonlinear in $x$. Kernel methods choose $\phi$ implicitly; neural networks learn $\phi$ from data.

# 2 Regularisation and Model Selection

## 2.1 Penalized Linear Estimation

Regularisation controls model complexity by penalising unstable parameter directions or by constraining the feasible set.

**Definition 2.1.1.** *Penalised Linear Estimation*
For empirical loss $L_n(\beta)$, a penalised estimator solves $\widehat{\beta}_\lambda \in \arg\min_\beta \{L_n(\beta) + \lambda\Omega(\beta)\}$.
The choices $\Omega(\beta) = \|\beta\|_2^2$ and $\Omega(\beta) = \|\beta\|_1$ give ridge and lasso penalties.

**Definition 2.1.2.** *Subset Selection, Ridge, and Lasso*
Let $y \in \mathbb{R}^n$ be centered and let $X \in \mathbb{R}^{n\times p}$ have centered, standardised columns.

(i) *Best subset*: for a size $m$, solve $\widehat{\beta}_m^{\text{sub}} \in \underset{|\operatorname{supp}(\beta)|\le m}{\arg\min} \|y - X\beta\|_2^2$

(ii) *Ridge*: for $\lambda \ge 0$, solve $\widehat{\beta}_\lambda^{\text{ridge}} \in \underset{\beta}{\arg\min}\{\|y - X\beta\|_2^2 + \lambda \|\beta\|_2^2\}$

(iii) *Lasso*: for $\lambda \ge 0$, solve $\widehat{\beta}_\lambda^{\text{lasso}} \in \underset{\beta}{\arg\min} \left\{\frac{1}{2}\|y - X\beta\|_2^2 + \lambda \|\beta\|_1\right\}$

Centering separates the unpenalised intercept, estimated by the mean on the raw scale.

**Remark 2.1.3.** *Penalty Geometry*
Assume $X^TX = I$ and set $z_j = x_j^T y$. For subset size $m$, best subset selection keeps the $m$ largest $|z_j|$.
Ridge and lasso have coordinates $\widehat{\beta}_{\lambda,j}^{\text{ridge}} = \frac{z_j}{1+\lambda}$ and $\widehat{\beta}_{\lambda,j}^{\text{lasso}} = \operatorname{sign}(z_j)(|z_j| - \lambda)_+$.
Thus $\ell^2$ shrinkage is smooth, while $\ell^1$ shrinkage can set coefficients exactly to zero.




Figure 263: Left: lasso $\ell^1$ diamond; right: ridge $\ell^2$ disk.

**Proposition 2.1.4.** *Ridge Shrinkage and Effective Degrees*
*Let $X \in \mathbb{R}^{n\times p}$ be centered with rank $r$ and thin SVD $X = UDV^T$, where $D = \operatorname{diag}(d_1, \dots, d_r)$ and $d_j > 0$.*
*For $\lambda > 0$, set $S_\lambda := X(X^TX + \lambda I_p)^{-1}X^T$ and $s_j(\lambda) := \frac{d_j^2}{d_j^2+\lambda}$. Then $S_\lambda = U \operatorname{diag}(s_1(\lambda), \dots, s_r(\lambda))U^T$.*
*So $\widehat{y}_\lambda^{\text{ridge}} := X\widehat{\beta}_\lambda^{\text{ridge}} = S_\lambda y = \sum_{j=1}^r s_j(\lambda) u_j u_j^T y$ and $\operatorname{df}(\lambda) := \operatorname{tr}(S_\lambda) = \sum_{j=1}^r s_j(\lambda)$.*

*Proof.* Since $X^TX = VD^2V^T$ and $X^T = VDU^T$, we have $V^T(X^TX + \lambda I_p)^{-1}V = (D^2 + \lambda I_r)^{-1}$.
Hence $S_\lambda = UDV^T(X^TX + \lambda I_p)^{-1}VDU^T = UD(D^2 + \lambda I_r)^{-1}DU^T = U \operatorname{diag}(s_1(\lambda), \dots, s_r(\lambda))U^T$.
Apply to $y$ and take traces, using $U^TU = I_r$.
*Structural Commentary*: Ridge is diagonal in the left singular-vector basis of $X$.
*Key Insight*: Ridge keeps direction $u_j$ by $s_j(\lambda)$; weak directions count less in the fit and $\operatorname{df}(\lambda)$. □

## 2.2 Validation and Selection Bias

**Definition 2.2.1.** *Model Selection by Validation*
Let $\mathcal{M}$ be a candidate set of models or hyperparameters.

(i) A *validation criterion* is $m \mapsto \widehat{R}_{\mathrm{val}}(m)$; choose $\widehat{m} \in \arg\min_{m\in\mathcal{M}} \widehat{R}_{\mathrm{val}}(m)$.

(ii) In *$K$-fold cross-validation*, take folds $I_1, \dots, I_K$ and fits $\widehat{f}_m^{(-k)}$.
Define $\ell_{ik}(m) := \ell(y_i, \widehat{f}_m^{(-k)}(x_i))$, $\widehat{R}_{\mathrm{cv}}(m) := \frac{1}{n}\sum_{k=1}^{K}\sum_{i\in I_k} \ell_{ik}(m)$, and choose $\widehat{m}_{\mathrm{cv}} \in \arg\min_{m\in\mathcal{M}} \widehat{R}_{\mathrm{cv}}(m)$.

**Method 2.2.2.** *One-Standard-Error Rule*
Let $(\mathcal{M}, \preceq)$ order candidates by complexity, with $\widehat{R}_{\mathrm{cv}}, \widehat{\mathrm{se}} : \mathcal{M} \to \mathbb{R}$.

(i) Choose $\widehat{m}_{\min} \in \arg\min_{m\in\mathcal{M}} \widehat{R}_{\mathrm{cv}}(m)$ and set $\widehat{R}_{\min} := \widehat{R}_{\mathrm{cv}}(\widehat{m}_{\min})$.
(ii) Define $\mathcal{M}_{1\mathrm{se}} := \{m \in \mathcal{M} : \widehat{R}_{\mathrm{cv}}(m) \le \widehat{R}_{\min} + \widehat{\mathrm{se}}(\widehat{m}_{\min})\}$.
(iii) Select $\widehat{m}_{1\mathrm{se}}$ as the $\preceq$-minimal element of $\mathcal{M}_{1\mathrm{se}}$; use any final test set only after this choice.

**Example 2.2.3.** *A Selection and Shrinkage Workflow*
Let $D = \{(x_i, y_i)\}_{i=1}^{n}$, where $y_i = \log(\mathrm{PSA}_i)$ and $x_i = (\mathrm{lcavol}_i, \mathrm{lweight}_i, \mathrm{age}_i, \mathrm{lbph}_i, \mathrm{svi}_i, \mathrm{lcp}_i, \mathrm{gleason}_i, \mathrm{pgg45}_i)$.
Split indices into training and test sets $T, E$, standardize $X_T$, and choose each complexity parameter by cross-validation on $T$: subset size $m$, ridge/lasso penalty $\lambda$, or PCR/PLS dimension $M$.
The selected fit $\widehat{f}$ is then evaluated once on $E$ by $\widehat{R}_{\mathrm{test}} := |E|^{-1}\sum_{i\in E} \ell(y_i, \widehat{f}(x_i))$.

**Remark 2.2.4.** *Selection Bias*
If $\widehat{m} \in \arg\min_{m\in\mathcal{M}} \widehat{R}_V(m)$ is chosen on validation set $V$, then $\widehat{R}_V(\widehat{m})$ is optimistically biased for $R(\widehat{m})$.
Estimate the selected model with an independent set $E$, using $\widehat{R}_E(\widehat{m})$, or use nested validation.

## 2.3 Derived Input Directions and Regularization Paths

**Method 2.3.1.** *Derived Input Directions*
Let $X, y$ be centered. A derived-direction method chooses $Z_M = (z_1, \dots, z_M)$ with $z_m = Xv_m$ or data-adaptive analogues, then fits $\widehat{y}_M = P_{Z_M} y$, where $P_{Z_M} := Z_M(Z_M^T Z_M)^{-1} Z_M^T$; $M$ is selected by validation.

(i) *Principal components regression*: for $X = UDV^T$, take $v_1, \dots, v_M$ as the leading right singular vectors, set $z_m = Xv_m = d_m u_m$, and fit $\widehat{y}_M^{\mathrm{pcr}} = \sum_{m=1}^{M} z_m \frac{z_m^T y}{z_m^T z_m} = \sum_{m=1}^{M} u_m u_m^T y$.

(ii) *Partial least squares*: set $X^{(0)} = X$ and, for $m = 1, \dots, M$, define $z_m = X^{(m-1)} X^{(m-1)T} y$ and $X^{(m)} = (I - P_{z_m}) X^{(m-1)}$, where $P_{z_m} := \frac{z_m z_m^T}{z_m^T z_m}$. With $Z_M = (z_1, \dots, z_M)$, fit $\widehat{y}_M^{\mathrm{pls}} = P_{Z_M} y$.

**Algorithm 9:** Partial Least Squares Directions

**Require:** Centered response $y$, standardized columns $x_1, \dots, x_p$, number of directions $M \le p$
**Ensure:** Fitted vectors $\widehat{y}^{(1)}, \dots, \widehat{y}^{(M)}$
  Set $\widehat{y}^{(0)} \leftarrow 0$ and $x_j^{(0)} \leftarrow x_j$ for $j = 1, \dots, p$
  **for** $m = 1, \dots, M$ **do**
    $z_m \leftarrow \sum_{j=1}^{p} (x_j^{(m-1)})^T y \, x_j^{(m-1)}$
    $\widehat{\theta}_m \leftarrow \frac{z_m^T y}{z_m^T z_m}$
    $\widehat{y}^{(m)} \leftarrow \widehat{y}^{(m-1)} + \widehat{\theta}_m z_m$
    **for** $j = 1, \dots, p$ **do**
      $x_j^{(m)} \leftarrow x_j^{(m-1)} - \frac{z_m^T x_j^{(m-1)}}{z_m^T z_m} z_m$
    **end for**
  **end for**

**Remark 2.3.2.** *PLS Versus PCR*
PCR chooses $v_m$ from $X^T X v_m = d_m^2 v_m$, so directions are ordered only by predictor variance $d_m^2$.
PLS chooses $z_m = X^{(m-1)} X^{(m-1)T} y$ and deflates $X^{(m)} = (I - P_{z_m}) X^{(m-1)}$, so each direction depends on both variance in $X^{(m-1)}$ and covariance with $y$.

**Method 2.3.3.** *Piecewise-Linear Path Algorithms*
Let $\widehat{\beta}(\lambda) \in \arg\min_\beta\{R(\beta) + \lambda J(\beta)\}$.

(i) If $R$ is quadratic or piecewise quadratic and $J$ is piecewise linear, fixed active-set and sign regions give affine KKT equations in $\lambda$, so $\widehat{\beta}(\lambda)$ is piecewise linear.
(ii) *Least angle regression*: with residual $r = y - X\beta$ and active set $\mathcal{A}$, move $\beta_\mathcal{A}$ in a direction keeping $|x_j^T r|$ tied for $j \in \mathcal{A}$ until a new variable ties or the least-squares fit is reached.
(iii) *Lasso path*: follow the same tied-correlation path and remove $j \in \mathcal{A}$ when $\beta_j$ reaches 0.
(iv) *Coordinate descent*: for fixed $\lambda$, update $\beta_j \leftarrow \arg\min_t\{\frac{1}{2}\|y - X_{-j}\beta_{-j} - x_j t\|_2^2 + \lambda(|t| + \|\beta_{-j}\|_1)\}$ and warm-start along $\lambda_1 > \cdots > \lambda_L$.

**Method 2.3.4.** *LAR, Lasso, and Correlation Ties*
Let $\widehat{\beta}_\lambda \in \arg\min_\beta\{\frac{1}{2}\|y - X\beta\|_2^2 + \lambda\|\beta\|_1\}$, $r_\lambda := y - X\widehat{\beta}_\lambda$, and $\mathcal{A}_\lambda := \{j : \widehat{\beta}_{\lambda,j} \neq 0\}$.

(i) *KKT*: $x_j^T r_\lambda = \lambda \operatorname{sign}(\widehat{\beta}_{\lambda,j})$ for $j \in \mathcal{A}_\lambda$, and $|x_j^T r_\lambda| \le \lambda$ for $j \notin \mathcal{A}_\lambda$.
(ii) *Correlation ties*: active predictors satisfy $|x_j^T r_\lambda| = \lambda$; inactive predictors satisfy $|x_j^T r_\lambda| \le \lambda$.
(iii) *LAR*: with $C = \|X^T r\|_\infty$, move in the equiangular active direction preserving $|x_j^T r| = C$ for $j \in \mathcal{A}$ until some $k \notin \mathcal{A}$ reaches $|x_k^T r| = C$.
(iv) *Lasso path*: on a fixed active set and sign pattern, the KKT equations are affine in $\lambda$. When some $\widehat{\beta}_{\lambda,j}$ reaches 0, remove $j$ and recompute the active direction.

**Algorithm 10:** Least Angle Regression Path

**Require:** Centered response $y$ and standardized columns $x_1, \ldots, x_p$ with $\|x_j\|_2 = 1$
**Ensure:** Piecewise-linear coefficient path $\{\beta(t)\}$
Initialize $\beta \leftarrow 0$, $r \leftarrow y$, $c \leftarrow X^T r$, $C \leftarrow \|c\|_\infty$, and $\mathcal{A} \leftarrow \{j : |c_j| = C\}$
**repeat**
$s_\mathcal{A} \leftarrow \operatorname{sign}(c_\mathcal{A})$, $\widetilde{X}_\mathcal{A} \leftarrow X_\mathcal{A} \operatorname{diag}(s_\mathcal{A})$, and $G_\mathcal{A} \leftarrow \widetilde{X}_\mathcal{A}^T \widetilde{X}_\mathcal{A}$
$\alpha_\mathcal{A} \leftarrow (\mathbf{1}^T G_\mathcal{A}^{-1} \mathbf{1})^{-1/2}$, $w_\mathcal{A} \leftarrow \alpha_\mathcal{A} G_\mathcal{A}^{-1}\mathbf{1}$, $u_\mathcal{A} \leftarrow \widetilde{X}_\mathcal{A} w_\mathcal{A}$, and $a \leftarrow X^T u_\mathcal{A}$
$\gamma \leftarrow \min\{C/\alpha_\mathcal{A}, \min^+_{j \notin \mathcal{A}}\{(C - c_j)/(\alpha_\mathcal{A} - a_j), (C + c_j)/(\alpha_\mathcal{A} + a_j)\}\}$
Record $\beta_\mathcal{A}(t) = \beta_\mathcal{A} + t \operatorname{diag}(s_\mathcal{A}) w_\mathcal{A}$ for $0 \le t \le \gamma$
$\beta_\mathcal{A} \leftarrow \beta_\mathcal{A} + \gamma \operatorname{diag}(s_\mathcal{A}) w_\mathcal{A}$ and $r \leftarrow r - \gamma u_\mathcal{A}$
$c \leftarrow X^T r$, $C \leftarrow \|c\|_\infty$, and $\mathcal{A} \leftarrow \mathcal{A} \cup \{j \notin \mathcal{A} : |c_j| = C\}$
**until** $C = 0$ or $\mathcal{A} = \{1, \ldots, p\}$

**Algorithm 11:** Incremental Forward Stagewise Regression

**Require:** Centered response $y$, standardized columns $x_1, \ldots, x_p$, small step size $\varepsilon > 0$
**Ensure:** A stagewise coefficient path
Initialize $\beta \leftarrow 0$ and $r \leftarrow y$
**repeat**
Choose $j$ maximizing $|x_j^T r|$
$\delta \leftarrow \varepsilon \operatorname{sign}(x_j^T r)$
$\beta_j \leftarrow \beta_j + \delta$
$r \leftarrow r - \delta x_j$
**until** the residual has negligible correlation with every predictor

**Remark 2.3.5.** *Stagewise and Lasso Paths*
Let $c = X^T r$ and $j \in \arg\max_k |c_k|$. Forward stagewise uses $\beta_j \leftarrow \beta_j + \varepsilon \operatorname{sign}(c_j)$ and $r \leftarrow r - \varepsilon \operatorname{sign}(c_j) x_j$; as $\varepsilon \downarrow 0$, it follows maximal-correlation directions, tracking LAR/lasso until a sign reversal is required. Boosting applies the same greedy infinitesimal update to additive fits.

**Algorithm 12:** Pathwise Coordinate Descent for the Lasso

**Require:** Centered response $y$, standardized columns $x_j$ with $\|x_j\|_2 = 1$, decreasing penalties $\lambda_1 > \cdots > \lambda_L \ge 0$
**Ensure:** Lasso coefficient vectors $\widehat{\beta}(\lambda_\ell)$ on the penalty grid
Define $S(z, \lambda) = \operatorname{sign}(z)(|z| - \lambda)_+$
Initialize $\beta \leftarrow 0$ and $r \leftarrow y - X\beta$
**for** $\ell = 1, \ldots, L$ **do**
**repeat**
**for** $j = 1, \ldots, p$ **do**
$r \leftarrow r + x_j \beta_j$
$z_j \leftarrow x_j^T r$

$\beta_j \leftarrow S(z_j, \lambda_\ell)$
$r \leftarrow r - x_j \beta_j$
**end for**
**until** the coefficient vector has converged
$\widehat{\beta}(\lambda_\ell) \leftarrow \beta$
**end for**

## 2.4 Effective Degrees and Information Criteria

**Definition 2.4.1.** *Effective Degrees of Freedom*
Let $\widehat{y} = S_\lambda y$ be a linear smoother with $S_\lambda = S_\lambda(X)$ independent of $y$.
Its *effective degrees of freedom* are $\mathrm{df}(\lambda) := \mathrm{trace}(S_\lambda)$. If $S_\lambda = P_V$ with $\dim V = m$, then $\mathrm{df}(\lambda) = m$.
For shrinkage smoothers, $\mathrm{df}(\lambda)$ need not be integral.

**Remark 2.4.2.** *Optimism of Training Error*
Assume $y = \mu + \varepsilon$, $\mathbb{E}\varepsilon = 0$, $\mathrm{Var}(\varepsilon) = \sigma^2 I$, and $\widehat{y} = S_\lambda y$ with $S_\lambda = S_\lambda(X)$.
Let $y^0 = \mu + \varepsilon^0$ be independent, $\mathrm{err}_{\mathrm{tr}}(\lambda) := n^{-1} \|y - \widehat{y}\|_2^2$, and $\mathrm{Err}_{\mathrm{in}}(\lambda) := n^{-1} \mathbb{E}_{y^0} \|y^0 - \widehat{y}\|_2^2$.
Then $\mathrm{Opt}(\lambda) := \mathbb{E}\{\mathrm{Err}_{\mathrm{in}}(\lambda) - \mathrm{err}_{\mathrm{tr}}(\lambda)\} = \frac{2}{n} \sum_{i=1}^n \mathrm{Cov}(\widehat{y}_i, y_i) = \frac{2\sigma^2}{n} \mathrm{trace}(S_\lambda)$.
Thus $\mathrm{df}(\lambda) = \mathrm{trace}(S_\lambda)$ is the covariance penalty.

**Method 2.4.3.** *AIC, $C_p$, and BIC Selection*
Let $\Lambda$ be candidate tuning values. For a criterion $C : \Lambda \to \mathbb{R}$, select $\widehat{\lambda} \in \arg\min_{\lambda \in \Lambda} C(\lambda)$.

(i) For squared error, set $\overline{\mathrm{err}}(\lambda) := \frac{1}{n} \sum_{i=1}^n (y_i - \widehat{y}_{\lambda,i})^2$ and $d_\lambda := \mathrm{df}(S_\lambda)$; use $C_p(\lambda) := \overline{\mathrm{err}}(\lambda) + \frac{2\widehat{\sigma}^2}{n} d_\lambda$.
(ii) For likelihood fits with maximized log-likelihood $\ell_\lambda$, use $\mathrm{AIC}(\lambda) := -\frac{2}{n}\ell_\lambda + \frac{2}{n} d_\lambda$.
(iii) Use $\mathrm{BIC}(\lambda) := -\frac{2}{n}\ell_\lambda + \frac{\log n}{n} d_\lambda$; the larger penalty favors smaller $d_\lambda$ as $n$ grows.

**Remark 2.4.4.** *Bayesian and MDL Readings of BIC*
For models $\mathcal{M}_m$ with dimension $d_m$, maximised likelihood $L_m$, and nearly uniform prior $\pi_m$, set $\mathrm{BIC}_m := -2 \log L_m + d_m \log n$. By laplace approximation, posterior weight is $\widehat{\mathbb{P}}(\mathcal{M}_m \mid Z) = \frac{\pi_m \exp\{-\mathrm{BIC}_m/2\}}{\sum_{\ell=1}^M \pi_\ell \exp\{-\mathrm{BIC}_\ell/2\}}$, so $\arg\min_m \mathrm{BIC}_m$ is an approximate MAP model.
In MDL form, $\frac{1}{2}\mathrm{BIC}_m$ is model code length plus fitted-discrepancy code length up to constants.

**Method 2.4.5.** *Generalized Cross-Validation*
Let $\Lambda$ be candidate tuning values and $\widehat{y}_\lambda = S_\lambda y$.

(i) Define $e_{\lambda i} := y_i - \widehat{y}_{\lambda i}$, $h_{\lambda i} := S_{\lambda,ii}$, and $d_\lambda := \mathrm{trace}(S_\lambda)$.
(ii) The leave-one-out residual is $e_{\lambda i}^{(-i)} = y_i - \widehat{f}_\lambda^{(-i)}(x_i) = \frac{e_{\lambda i}}{1 - h_{\lambda i}}$.
(iii) *Generalized cross-validation* replaces $h_{\lambda i}$ by $\bar{h}_\lambda := d_\lambda / n$ and sets $\mathrm{GCV}(\lambda) := \frac{1}{n} \sum_{i=1}^n \left( \frac{e_{\lambda i}}{1 - d_\lambda / n} \right)^2$.
(iv) Select $\widehat{\lambda}_{\mathrm{gcv}} \in \arg\min_{\lambda \in \Lambda} \mathrm{GCV}(\lambda)$; this uses $d_\lambda$ instead of the full leverage diagonal.

# 3 Basis Expansions, Splines, and Wavelets

## 3.1 Basis Expansions, Splines, and Wavelet Features

Basis expansions keep the fitting step linear while moving nonlinearity into the representation of the inputs.

**Definition 3.1.1.** *Linear Basis Expansion*

A *linear basis expansion* chooses $h = (h_1, \ldots, h_M) : \mathcal{X} \to \mathbb{R}^M$ and fits $f_\beta(x) := \beta^T h(x) = \sum_{m=1}^M \beta_m h_m(x)$.

Typical $h_m$ include coordinates, powers, interactions, one-dimensional transforms, and region indicators.
The model is linear in $h(x)$, but generally nonlinear in $x$.

**Remark 3.1.2.** *Dictionaries Need Complexity Control*

For a dictionary $\mathcal{H} = \{h_m\}_{m=1}^M$, fitting $f_\beta(x) = \sum_{m=1}^M \beta_m h_m(x)$ is flexible but unstable when $M$ is large or the $h_m$ are nearly collinear. Complexity control restricts admissible fits: additive models impose $f(x) = \sum_j f_j(x_j)$, subset selection imposes $\|\beta\|_0 \le s$, and regularization minimizes empirical loss plus $\lambda \Omega(\beta)$.

**Definition 3.1.3.** *Regression Splines*
Let $\xi_1 < \cdots < \xi_K$ be knots.

(i) *Order-$M$ regression spline* is piecewise polynomial $s : \mathbb{R} \to \mathbb{R}$ of degree $\le M - 1$, $s \in C^{M-2}$ at knots.




Figure 264: Basis expansion: boundary curved in $x$, linear in $h(x)$.

(ii) *Cubic truncated-power basis* is basis $1, x, x^2, x^3, (x-\xi_1)_+^3, \dots, (x-\xi_K)_+^3$ for cubic spline space.
(iii) *Natural cubic spline* is a cubic regression spline satisfying $s''(x) = 0$ for $x \notin [\xi_1, \xi_K]$.
(iv) *B-spline basis* is a locally supported basis for the same spline space.

**Method 3.1.4.** *Smoothing Splines as Penalized Fits*
Let $(x_i, y_i)_{i=1}^n \subset \mathbb{R}^2$, $y = (y_1, \dots, y_n)^T$, $\lambda \geq 0$, and $\mathcal{F} := \{f : f'' \in L^2(\mathbb{R})\}$.

(i) The smoothing spline is $\widehat{f}_\lambda \in \arg\min_{f \in \mathcal{F}} \left\{ \sum_{i=1}^n (y_i - f(x_i))^2 + \lambda \int_{\mathbb{R}} (f''(t))^2 \, dt \right\}$.
(ii) $\widehat{f}_\lambda$ is a natural cubic spline with knots at the distinct observed inputs.
(iii) With $\widehat{y}_\lambda = (\widehat{f}_\lambda(x_1), \dots, \widehat{f}_\lambda(x_n))^T$, there is a smoother matrix $S_\lambda$ such that $\widehat{y}_\lambda = S_\lambda y$ and $d_\lambda := \operatorname{trace}(S_\lambda)$.
(iv) For $e_{\lambda i} := y_i - \widehat{f}_\lambda(x_i)$ and $h_{\lambda i} := S_{\lambda, ii}$, leave-one-out CV is $\mathrm{CV}(\lambda) := \frac{1}{n} \sum_{i=1}^n \left( \frac{e_{\lambda i}}{1 - h_{\lambda i}} \right)^2$.

**Definition 3.1.5.** *Tensor-Product and Thin-Plate Splines*
Let $\{h_{rj}\}_{j=1}^{J_r}$ be a univariate basis for coordinate $x_r$.

(i) A *two-way tensor-product basis* is $\{g_{jk}\}$ with $g_{jk}(x_1, x_2) := h_{1j}(x_1) h_{2k}(x_2)$ and dimension $J_1 J_2$.
(ii) A *d-way tensor-product basis* is $\{g_{j_1, \dots, j_d}\}$ with $g_{j_1, \dots, j_d}(x) := \prod_{r=1}^d h_{r j_r}(x_r)$ and dimension $\prod_{r=1}^d J_r$.
(iii) A *thin-plate spline fit* is any $\widehat{f}_\lambda \in \arg\min_f Q_\lambda(f)$, where $Q_\lambda(f) := \sum_{i=1}^n (y_i - f(x_i))^2 + \lambda J(f)$, $D_{ab} f := \partial_a \partial_b f$, and $J(f) := \int_{\mathbb{R}^2} \{(D_{11} f)^2 + 2(D_{12} f)^2 + (D_{22} f)^2\} \, dx$.
(iv) A *thin-plate representer* is $\widehat{f}(x) = a_0 + a^\top x + \sum_{\ell=1}^L c_\ell \phi(\|x - \kappa_\ell\|)$; $\phi(r) := r^2 \log r$, $\phi(0) := 0$, $\sum_{\ell=1}^L c_\ell (1, \kappa_\ell^\top) = 0$.

**Method 3.1.6.** *Spline-Filtered Logistic Features*
Let $(x_i, y_i)_{i=1}^n$ with $x_i \in \mathbb{R}^p$, $y_i \in \{0, 1\}$, and let $H \in \mathbb{R}^{p \times M}$ be a smooth basis matrix with $M \ll p$.

(i) Restrict $\beta \in \mathbb{R}^p$ to $\mathrm{col}(H)$ by writing $\beta = H\theta$, $\theta \in \mathbb{R}^M$.
(ii) Define filtered features $z_i := H^T x_i$; then $x_i^T \beta = z_i^T \theta$.
(iii) Fit $\widehat{\theta}, \widehat{b} \in \arg\min_{\theta, b} \sum_{i=1}^n \{\log(1 + \exp(b + z_i^T \theta)) - y_i (b + z_i^T \theta)\}$.
(iv) Set $\widehat{\beta} = H\widehat{\theta}$; smoothness is imposed by $\widehat{\beta} \in \mathrm{col}(H)$ and dimension drops from $p$ to $M$.

**Method 3.1.7.** *Wavelet Shrinkage*
Let $y \in \mathbb{R}^n$ be observed on a regular lattice and let $W \in \mathbb{R}^{n \times n}$ be an orthonormal wavelet matrix.

(i) Transform $y$ to wavelet coefficients $y^* := W^\top y$.
(ii) Estimate sparse coefficients by $\widehat{\theta}_\lambda \in \arg\min_{\theta \in \mathbb{R}^n} Q_\lambda(\theta)$, where $Q_\lambda(\theta) := \|y - W\theta\|_2^2 + 2\lambda \|\theta\|_1$.
(iii) Since $W^\top W = I$, $\widehat{\theta}_{\lambda, j} = \operatorname{sign}(y_j^*)(|y_j^*| - \lambda)_+$ for $j = 1, \dots, n$.
(iv) Reconstruct $\widehat{f}_\lambda := W\widehat{\theta}_\lambda = \sum_{j : |y_j^*| > \lambda} \operatorname{sign}(y_j^*)(|y_j^*| - \lambda) w_j$, where $w_j$ is the $j$th wavelet atom.

# 4 Kernel Methods and RKHS Theory

## 4.1 Kernel Methods and RKHS Representations

Kernel methods keep the algorithm linear in an implicit feature space while computing only inner products.

**Definition 4.1.1.** *Positive Definite Kernel*
A map $k : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ is a *positive definite kernel* if:

(i) $k(x, y) = k(y, x)$ for all $x, y \in \mathcal{X}$;

(ii) for every $n \in \mathbb{N}$ and $x_1, \dots, x_n \in \mathcal{X}$, $K = (k(x_i, x_j))_{i,j=1}^n$ satisfies $a^\top K a = \sum_{i,j=1}^n a_i a_j k(x_i, x_j) \geq 0 \ \forall a \in \mathbb{R}^n$.

**Example 4.1.2.** *Common Kernels*
For $x, y \in \mathbb{R}^p$, common positive definite kernels include:

(i) *Polynomial*: for $d \in \mathbb{N}_{>0}$, $k_d(x, y) := (\langle x, y \rangle + 1)^d = \langle h_d(x), h_d(y) \rangle$, encoding monomials of degree $\leq d$. If $p = 2$ and $d = 2$, one feature map is $h_2(x) = (1, \sqrt{2}x_1, \sqrt{2}x_2, x_1^2, x_2^2, \sqrt{2}x_1 x_2) \in \mathbb{R}^6$.

(ii) *Gaussian RBF*: for $\nu > 0$, $k_\nu(x, y) := \exp(-\nu \|x - y\|_2^2)$; it is radial and translation invariant, with $k_\nu(x, y)$ decreasing as $\|x - y\|_2$ increases.

**Definition 4.1.3.** *Reproducing Kernel Hilbert Space*
A Hilbert space $\mathcal{H} \subseteq \mathbb{R}^{\mathcal{X}}$ is a *reproducing kernel Hilbert space* if point evaluation is continuous; explicitly:

(i) for every $x \in \mathcal{X}$, the evaluation map $E_x : \mathcal{H} \to \mathbb{R}$, $E_x(f) := f(x)$, is continuous;
(ii) by Riesz, $k_x \in \mathcal{H}$ denotes the unique representer of $E_x$, so $f(x) = \langle f, k_x \rangle_{\mathcal{H}}$ for all $f \in \mathcal{H}$;
(iii) the *reproducing kernel* is $k : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$, $k(x, y) := k_x(y) = \langle k_x, k_y \rangle_{\mathcal{H}}$, with $k_x = k(x, \cdot)$.

**Theorem 4.1.4.** *Representer Theorem Form*
*Let $\mathcal{H}$ be an RKHS on $\mathcal{X}$ with kernel $k$.*
*Let $(x_i, y_i)_{i=1}^n \subseteq \mathcal{X} \times \mathcal{Y}$, $\lambda > 0$, $\ell : \mathbb{R} \times \mathcal{Y} \to (-\infty, +\infty]$, and $\Phi : [0, \infty) \to \mathbb{R}$ nondecreasing.*
*For $J(f) := \sum_{i=1}^n \ell(f(x_i), y_i) + \lambda \Phi(\|f\|_{\mathcal{H}})$ and $S := \operatorname{span}\{k(x_i, \cdot) : 1 \leq i \leq n\}$, if $f^\star \in \operatorname{argmin}_{f \in \mathcal{H}} J(f)$, then $Pf^\star \in S$ is also a minimiser, where $P : \mathcal{H} \to S$ is orthogonal projection.*
*Hence some minimiser has $f^\star(\cdot) = \sum_{i=1}^n \alpha_i k(x_i, \cdot)$; if $\Phi$ is strictly increasing, every minimiser has this form.*

*Proof.* Since $S$ is finite-dimensional, it is closed, so $P$ exists.
Write $f = f_S + f_\perp$ with $f_S := Pf \in S$ and $f_\perp \in S^\perp$.
For each $i$, $k(x_i, \cdot) \in S$, hence $f_\perp(x_i) = \langle f_\perp, k(x_i, \cdot) \rangle_{\mathcal{H}} = 0$ and $f_S(x_i) = f(x_i)$.
Thus the empirical loss in $J$ is unchanged by $f \mapsto f_S$.
By orthogonality, $\|f\|_{\mathcal{H}}^2 = \|f_S\|_{\mathcal{H}}^2 + \|f_\perp\|_{\mathcal{H}}^2$, so $\|f_S\|_{\mathcal{H}} \leq \|f\|_{\mathcal{H}}$, hence $J(f_S) \leq J(f)$ by monotonicity of $\Phi$.
Applying this to a minimiser $f^\star$ gives $J(Pf^\star) = J(f^\star)$, so $Pf^\star$ is a minimiser in $S$.
Since $S = \{\sum_{i=1}^n \alpha_i k(x_i, \cdot) : \alpha \in \mathbb{R}^n\}$, finite expansion follows. If $\Phi$ strictly increasing, $f_\perp \neq 0$, then $\|f_S\|_{\mathcal{H}} < \|f\|_{\mathcal{H}}$ while loss is unchanged, so $J(f_S) < J(f)$, contradicting minimality.
*Structural Commentary*: $E : \mathcal{H} \to \mathbb{R}^n$, $Ef = (f(x_i))_{i=1}^n$, has $E|_{S^\perp} = 0$ and $Ef = E(Pf)$; only $S$ is data-visible.
*Key Insight*: Regularisation kills $S^\perp$, so optimise over $\alpha \in \mathbb{R}^n$ via $f_\alpha = \sum_{i=1}^n \alpha_i k(x_i, \cdot)$ and $Ef_\alpha = K\alpha$. □

**Method 4.1.5.** *Kernel Ridge Regression*
Let $(x_i, y_i)_{i=1}^n \subseteq \mathcal{X} \times \mathbb{R}$, $y := (y_1, \dots, y_n)^\top$, $\lambda > 0$, and $K_{ij} := k(x_i, x_j)$.

(i) *RKHS problem*. Solve $\widehat{f} \in \operatorname{argmin}_{f \in \mathcal{H}} \sum_{i=1}^n (y_i - f(x_i))^2 + \lambda \|f\|_{\mathcal{H}}^2$.

(ii) *Representer reduction*. Write $\widehat{f}(\cdot) = \sum_{i=1}^n \widehat{\alpha}_i k(x_i, \cdot)$; then $\widehat{f}(x_j) = (K\widehat{\alpha})_j$ and $\left\|\widehat{f}\right\|_{\mathcal{H}}^2 = \widehat{\alpha}^\top K \widehat{\alpha}$.

(iii) *Finite problem*. Minimise $\widehat{\alpha} \in \operatorname{argmin}_{\alpha \in \mathbb{R}^n} \|y - K\alpha\|_2^2 + \lambda \alpha^\top K \alpha$.

(iv) *Solution*. Since $K \succeq 0$ and $\lambda > 0$, $K + \lambda I \succ 0$ and $\widehat{\alpha} = (K + \lambda I)^{-1} y$.
Hence $\widehat{y} = K\widehat{\alpha}$ and $\widehat{f}(x) = k_x^\top \widehat{\alpha}$, where $k_x := (k(x, x_1), \dots, k(x, x_n))^\top$.

**Remark 4.1.6.** *Localisation Kernels and RKHS Kernels*
*Localisation kernel* has weights $w_i(x_0) \propto K_\lambda(x_0, x_i)$. $w_i(x_0)$ is large when $x_i$ close to $x_0$; small otherwise.
*RKHS kernel* is PSD: $K_{ij} = k(x_i, x_j)$ is Gram matrix, i.e., $k(x, y) = \langle h(x), h(y) \rangle_{\mathcal{F}}$ for some feature map $h$.
Thus: localisation kernels → local averaging; RKHS kernels → inner products for global regularisation.

**Example 4.1.7.** *Radial Basis Expansions*
Let $\xi_1, \dots, \xi_M \in \mathbb{R}^p$, $\lambda_j > 0$, $\beta \in \mathbb{R}^M$, and let $D : [0, \infty) \to [0, \infty)$ be a radial profile.

(i) *Unnormalized basis*: $\varphi_j(x) := D(\frac{\|x - \xi_j\|_2}{\lambda_j})$ and $f(x) := \sum_{j=1}^M \beta_j \varphi_j(x)$.

(ii) *Normalized basis*: for $\lambda_j = \lambda$, set $h_j(x) := D(\frac{\|x - \xi_j\|_2}{\lambda})/s(x)$, where $s(x) := \sum_{k=1}^M D(\frac{\|x - \xi_k\|_2}{\lambda}) > 0$.

Then $h_j(x) \geq 0$ and $\sum_{j=1}^M h_j(x) = 1$.

(iii) *Training-point centers*: $M = n$, $\xi_i = x_i$, $\beta_i = Y_i$ give $\widehat{f}(x) := \sum_{i=1}^{n} Y_i h_i(x)$ (Nadaraya–Watson local avg).

**Remark 4.1.8.** *Spline and Kernel Viewpoints*
Common form: $\min_f \sum_{i=1}^{n} \ell(f(x_i), y_i) + \lambda \|Pf\|_{\mathcal{H}}^2$, where $P$ removes an unpenalised $\mathcal{U}$, gives $\widehat{f} = u + \sum_{i=1}^{n} \alpha_i k(x_i, \cdot)$ with $u \in \mathcal{U}$. Splines choose $P$, kernel methods choose $k$, and SVMs choose a margin loss $\ell$.

# 5 Support Vector Machines and Margin Methods

## 5.1 Margins and SVM Primal Problems

Margin methods prefer classifiers that separate classes with a buffer, not merely with zero training error. The support-vector construction makes this preference explicit by trading margin width against margin violations.

**Definition 5.1.1.** *Affine Hyperplane and Signed Distance*
Let $w \in \mathbb{R}^p \setminus \{0\}$, $b \in \mathbb{R}$, and $q_{w,b}(x) := w^\top x + b$.

(i) *Affine hyperplane*: $H_{w,b} := \{x \in \mathbb{R}^p : q_{w,b}(x) = 0\}$, with normal direction $w$.
(ii) *Signed distance*: $d_{w,b}(x) := \frac{q_{w,b}(x)}{\|w\|_2}$.
(iii) *Scale invariance*: for $c > 0$, $H_{cw,cb} = H_{w,b}$ and $q_{cw,cb} = c q_{w,b}$.

**Definition 5.1.2.** *Margin*
Let $y_i \in \{-1, 1\}$ and $q(x) := w^\top x + b$.

(i) *Functional margin*: $\gamma_i(q) := y_i q(x_i)$.
(ii) *Geometric margin*: for $w \neq 0$, $\rho_i(w, b) := \frac{y_i(w^\top x_i + b)}{\|w\|_2}$.
(iii) *Correct classification*: $\gamma_i(q) > 0$; larger $\gamma_i(q)$ means larger signed confidence at fixed score scale.

**Definition 5.1.3.** *Hard-Margin Support Vector Machine*
For separable $(x_i, y_i)_{i=1}^n \subseteq \mathbb{R}^p \times \{-1, 1\}$:

(i) *Unit-margin scale*: $y_i(w^\top x_i + b) \geq 1$ for all $i$.
(ii) *Hard-margin primal*: $(\widehat{w}, \widehat{b}) \in \operatorname{argmin}_{w,b} \frac{1}{2}\|w\|_2^2$ subject to $y_i(w^\top x_i + b) \geq 1$, $i = 1, \ldots, n$.
(iii) *Margin width*: the slab $-1 \leq w^\top x + b \leq 1$ has width $\frac{2}{\|w\|_2}$, so the primal maximises geometric margin.

**Definition 5.1.4.** *Soft-Margin Support Vector Machine*
For possibly nonseparable $(x_i, y_i)_{i=1}^n \subseteq \mathbb{R}^p \times \{-1, 1\}$ and $C > 0$:

(i) *Slack variables*: $\xi_i \geq 0$ and $y_i(w^\top x_i + b) \geq 1 - \xi_i$, $i = 1, \ldots, n$.
(ii) *Soft-margin primal*: $(\widehat{w}, \widehat{b}, \widehat{\xi}) \in \operatorname{argmin}_{w,b,\xi} \{\frac{1}{2}\|w\|_2^2 + C \sum_{i=1}^{n} \xi_i\}$ subject to the slack constraints.
(iii) *Penalty parameter*: $C$ prices $\sum_{i=1}^{n} \xi_i$; larger $C$ penalises violations more strongly.

**Remark 5.1.5.** *Why the Standard Slack Scaling*
Before score normalisation, additive slack uses $y_i(w^\top x_i + b) \geq M - \xi_i$, relative slack uses $y_i(w^\top x_i + b) \geq M(1 - \xi_i)$. Setting $M = 1$ gives convex constraint $y_i(w^\top x_i + b) \geq 1 - \xi_i$, $\xi_i \geq 0$.
Thus $\xi_i$ measures margin failure: 0 on/outside the margin, $(0, 1]$ inside but correct, and $> 1$ misclassified.



Figure 265: Soft-margin SVM: orange $y = +1$, blue $y = -1$; solid $f(x) = 0$, dashed $f(x) = \pm 1$; circled support vectors, red slack arrows $\xi_i$.

**Proposition 5.1.6.** *Hinge-Loss Penalty Form*
*Let* $(x_i, y_i)_{i=1}^n \subseteq \mathbb{R}^p \times \{-1, 1\}$, $C > 0$, $f_{w,b}(x) := w^\top x + b$, *and* $\lambda := 1/C$. *For soft-margin primal* $\min_{w,b,\xi}\{\frac{1}{2}\|w\|_2^2 + C\sum_{i=1}^n \xi_i\}$ *subject to* $\xi_i \geq 0$ *and* $y_i f_{w,b}(x_i) \geq 1 - \xi_i$, *eliminating* $\xi$ *and scaling by* $\frac{1}{C}$ *gives* $\min_{w,b} \sum_{i=1}^n [1 - y_i f_{w,b}(x_i)]_+ + \frac{\lambda}{2}\|w\|_2^2$. *For fixed* $(w, b)$, *minimal feasible slack is* $\xi_i^\star = [1 - y_i f_{w,b}(x_i)]_+$.

*Proof.* Fix $(w, b)$ and put $r_i := 1 - y_i f_{w,b}(x_i)$. Feasibility gives $\xi_i \geq 0$ and $\xi_i \geq r_i$, hence $\xi_i \geq \max\{0, r_i\} = [r_i]_+$.
Since $C > 0$, the objective is strictly increasing in each $\xi_i$, so $\xi_i^\star = [r_i]_+$.
Substitution yields $\frac{1}{2}\|w\|_2^2 + C\sum_{i=1}^n [1 - y_i f_{w,b}(x_i)]_+$.
Multiply by $1/C > 0$ preserves minimisers, $\sum_{i=1}^n [1 - y_i f_{w,b}(x_i)]_+ + \frac{1}{2C}\|w\|_2^2 = \sum_{i=1}^n [1 - y_i f_{w,b}(x_i)]_+ + \frac{\lambda}{2}\|w\|_2^2$.
*Structural Commentary*: The $\xi_i$ decouple; eliminate them pointwise.
*Key Insight*: Soft-margin SVM is hinge empirical risk plus an $\ell_2$ penalty on $w$; the intercept $b$ is unpenalized. □

## 5.2 Dual Problems and Kernelization

**Proposition 5.2.1.** *Hard-Margin Dual and Support Vectors*
*Let* $(x_i, y_i)_{i=1}^n \subseteq \mathbb{R}^p \times \{-1, 1\}$ *be linearly separable. The primal* $P : \min_{w,b} \frac{1}{2}\|w\|_2^2$ *subject to* $y_i(w^\top x_i + b) \geq 1$ *has dual* $D : \max_{\alpha \in \mathbb{R}_+^n} \sum_{i=1}^n \alpha_i - \frac{1}{2}\sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y_i y_j x_i^\top x_j$ *subject to* $\sum_{i=1}^n \alpha_i y_i = 0$.
*At any optimum,* $\widehat{w} = \sum_{i=1}^n \widehat{\alpha}_i y_i x_i$ *and* $\widehat{\alpha}_i > 0 \Rightarrow y_i(\widehat{w}^\top x_i + \widehat{b}) = 1$; *these active points are the support vectors.*

*Proof.* Let $g_i(w, b) := y_i(w^\top x_i + b) - 1$ and $\alpha_i \geq 0$. The Lagrangian is $L(w, b, \alpha) = \frac{1}{2}\|w\|_2^2 - \sum_{i=1}^n \alpha_i g_i(w, b)$.
Stationarity gives $0 = \nabla_w L = w - \sum_{i=1}^n \alpha_i y_i x_i$ and $0 = \partial_b L = -\sum_{i=1}^n \alpha_i y_i$, hence $w = \sum_{i=1}^n \alpha_i y_i x_i$ and $\sum_{i=1}^n \alpha_i y_i = 0$.
Substituting into $L$ gives $D(\alpha) = \sum_{i=1}^n \alpha_i - \frac{1}{2}\sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y_i y_j x_i^\top x_j$, with $\alpha \geq 0$ and $\sum_i \alpha_i y_i = 0$.
Strong duality holds by convexity and strict feasibility.
Complementary slackness gives $\widehat{\alpha}_i[y_i(\widehat{w}^\top x_i + \widehat{b}) - 1] = 0$, so $\widehat{\alpha}_i > 0$ forces $y_i(\widehat{w}^\top x_i + \widehat{b}) = 1$.
*Structural Commentary*: Stationarity leaves $\alpha$; data enter via $y_i y_j x_i^\top x_j$ and $\sum_i \alpha_i y_i = 0$.
*Key Insight*: $\alpha_i > 0$ marks margin support; these points form $\widehat{w}$ and fix the separator. □

**Proposition 5.2.2.** *Soft-Margin Dual and Support Types*
*Let* $(x_i, y_i)_{i=1}^n \subseteq \mathbb{R}^p \times \{-1, 1\}$ *and* $C > 0$. *The primal* $P : \min_{w,b,\xi} \frac{1}{2}\|w\|_2^2 + C\sum_{i=1}^n \xi_i$ *subject to* $y_i(w^\top x_i + b) \geq 1 - \xi_i$ *and* $\xi_i \geq 0$ *has dual* $D : \max_{\alpha \in [0,C]^n} \sum_{i=1}^n \alpha_i - \frac{1}{2}\sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y_i y_j x_i^\top x_j$ *subject to* $\sum_{i=1}^n \alpha_i y_i = 0$.
*At any optimum,* $\widehat{w} = \sum_{i=1}^n \widehat{\alpha}_i y_i x_i$. *With* $\widehat{f}(x) := \widehat{w}^\top x + \widehat{b}$, $\widehat{\alpha}_i = 0 \Rightarrow y_i \widehat{f}(x_i) \geq 1$, $0 < \widehat{\alpha}_i < C \Rightarrow y_i \widehat{f}(x_i) = 1$, *and* $\widehat{\alpha}_i = C \Rightarrow y_i \widehat{f}(x_i) = 1 - \widehat{\xi}_i \leq 1$. *The support vectors are exactly* $\{i : \widehat{\alpha}_i > 0\}$.

*Proof.* Let $\alpha_i \geq 0$ multiply $y_i(w^\top x_i + b) - 1 + \xi_i \geq 0$ and $\mu_i \geq 0$ multiply $\xi_i \geq 0$.
The Lagrangian is $L = \frac{1}{2}\|w\|_2^2 + C\sum_i \xi_i - \sum_i \alpha_i[y_i(w^\top x_i + b) - 1 + \xi_i] - \sum_i \mu_i \xi_i$.
Stationarity gives $w = \sum_i \alpha_i y_i x_i$, $\sum_i \alpha_i y_i = 0$, and $C - \alpha_i - \mu_i = 0$, hence $0 \leq \alpha_i \leq C$.
Substitution gives dual; Slater gives strong duality.
Complementary slackness gives $\alpha_i[y_i f(x_i) - 1 + \xi_i] = 0$ and $(C - \alpha_i)\xi_i = 0$.
Thus $\alpha_i = 0$ implies $\xi_i = 0$ and $y_i f(x_i) \geq 1$; $0 < \alpha_i < C$ implies $\xi_i = 0$ and $y_i f(x_i) = 1$.
Hence $\alpha_i = C$ implies $y_i f(x_i) = 1 - \xi_i \leq 1$.
*Structural Commentary*: Slacks become box constraints $0 \leq \alpha_i \leq C$.
*Key Insight*: $\alpha_i$ labels type: 0 outside, $(0, C)$ on-margin, $C$ inside/wrong-side. □

**Method 5.2.3.** *Kernelized Support Vector Classifier*
Let $(x_i, y_i)_{i=1}^n \subseteq \mathcal{X} \times \{-1, 1\}$, $C > 0$, $h : \mathcal{X} \to \mathcal{F}$, and $k(x, x') := \langle h(x), h(x') \rangle_{\mathcal{F}}$.
The kernelized dual is $\widehat{\alpha} \in \operatorname{argmax}_{\alpha \in [0,C]^n} \sum_{i=1}^n \alpha_i - \frac{1}{2}\sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y_i y_j k(x_i, x_j)$ subject to $\sum_{i=1}^n \alpha_i y_i = 0$.

The fitted score and classifier are $\widehat{f}(x) = \sum_{i=1}^{n} \widehat{\alpha}_i y_i k(x, x_i) + \widehat{b}$ and $\widehat{G}(x) = \operatorname{sign}(\widehat{f}(x))$.

If $0 < \widehat{\alpha}_s < C$, then $\widehat{b} = y_s - \sum_{i=1}^{n} \widehat{\alpha}_i y_i k(x_s, x_i)$; average this over all such $s$ for stability.

*Proof.* In feature space, the soft-margin primal uses $v \in \mathcal{F}$ and score $f_{v,b}(x) = \langle v, h(x) \rangle_{\mathcal{F}} + b$.
Its Lagrangian stationarity gives $v = \sum_{i=1}^{n} \alpha_i y_i h(x_i)$ and $\sum_{i=1}^{n} \alpha_i y_i = 0$, while slack stationarity gives $0 \le \alpha_i \le C$.
Substitution yields $\sum_i \alpha_i - \frac{1}{2} \left\| \sum_i \alpha_i y_i h(x_i) \right\|_{\mathcal{F}}^2 = \sum_i \alpha_i - \frac{1}{2} \sum_i \sum_j \alpha_i \alpha_j y_i y_j k(x_i, x_j)$.
Hence $\widehat{f}(x) = \langle \widehat{v}, h(x) \rangle_{\mathcal{F}} + \widehat{b} = \sum_i \widehat{\alpha}_i y_i k(x, x_i) + \widehat{b}$.

If $0 < \widehat{\alpha}_s < C$, KKT gives zero slack and $y_s \widehat{f}(x_s) = 1$, so $\widehat{f}(x_s) = y_s$ and the stated $\widehat{b}$ follows.
*Structural Commentary*: Coordinates vanish; optimisation uses only $K_{ij} = k(x_i, x_j)$.
*Key Insight*: The classifier is finite and support-vector based: only indices with $\widehat{\alpha}_i > 0$ contribute to $\widehat{f}$. □

**Remark 5.2.4.** *Hinge Loss and Probability-Oriented Losses*
Let $\eta(x) := \mathbb{P}(Y = 1 \mid X = x)$ and $m := yf(x)$.

(i) *Hinge*: $[1 - m]_+$ targets $\operatorname{sign}(2\eta(x) - 1)$ and is zero for $m \ge 1$.
(ii) *Logistic*: $\log(1 + e^{-m})$ targets $\log(\eta(x)/(1 - \eta(x)))$ and keeps a smooth positive tail.
(iii) *Squared*: squared loss targets $\mathbb{E}[Y \mid X = x] = 2\eta(x) - 1$ and keeps residual-magnitude leverage.

**Method 5.2.5.** *SVM Regularization Path View*
set $\lambda := 1/C$; as $\lambda \downarrow 0$ ($C \uparrow \infty$), hinge loss dominates norm regularisation.

(i) *Fit*: for each $\lambda$, solve for $f_\lambda(x) := w_\lambda^\top x + b_\lambda$ and multipliers $\alpha_i(\lambda)$.
(ii) *Classify states*: with $m_i(\lambda) := y_i f_\lambda(x_i)$, $m_i > 1 \Rightarrow \alpha_i = 0$, $m_i = 1 \Rightarrow 0 \le \alpha_i \le C$, and $m_i < 1 \Rightarrow \alpha_i = C$.
(iii) *Track*: fixed active set $\Rightarrow \alpha_i(\lambda)$ and $f_\lambda$ are linear.
(iv) *Kernelize*: replace $w_\lambda^\top x$ by $\sum_{i=1}^{n} \alpha_i(\lambda) y_i k(x, x_i)$, so $f_\lambda(x) = \sum_{i=1}^{n} \alpha_i(\lambda) y_i k(x, x_i) + b_\lambda$.

## 5.3 Support Vector Regression and Perceptrons

**Method 5.3.1.** *Epsilon-Insensitive Support Vector Regression*
Let $(x_i, y_i)_{i=1}^n \subseteq \mathbb{R}^p \times \mathbb{R}$, $\epsilon > 0$, $\lambda > 0$, and $f_{w,b}(x) := w^\top x + b$.

(i) *Define loss*: for residual $r$, set $V_\epsilon(r) := [|r| - \epsilon]_+$.
(ii) *Fit*: solve $\min_{w,b} \sum_{i=1}^{n} V_\epsilon(y_i - f_{w,b}(x_i)) + \frac{\lambda}{2} \|w\|_2^2$.
(iii) *Read tube*: $|r_i| \le \epsilon \Rightarrow V_\epsilon(r_i) = 0$ and $|r_i| > \epsilon \Rightarrow V_\epsilon(r_i) = |r_i| - \epsilon$.
(iv) *Kernelize*: $\widehat{f}(x) = \sum_{i=1}^{n} (\widehat{\alpha}_i^* - \widehat{\alpha}_i) k(x, x_i) + \widehat{b}$; tube sides have opposite signs.

**Method 5.3.2.** *Perceptron Update as a Contrast*
Choose $(w_0, b_0) \in \mathbb{R}^p \times \mathbb{R}$ and learning rate $\rho > 0$.

(i) *Scan*: at step $t$, choose $(x_i, y_i)$ and compute $m_i(t) := y_i(w_t^\top x_i + b_t)$.
(ii) *Update*: if $m_i(t) \le 0$, set $(w_{t+1}, b_{t+1}) := (w_t, b_t) + \rho y_i (x_i, 1)$.
(iii) *Keep*: if $m_i(t) > 0$, set $(w_{t+1}, b_{t+1}) := (w_t, b_t)$.
(iv) *Contrast*: perceptron may separate but is order-dependent; SVM solves a convex max-margin problem.

**Remark 5.3.3.** *Support Vectors*
Let $\widehat{f}(x) = \sum_{i=1}^{n} \widehat{\alpha}_i y_i k(x, x_i) + \widehat{b}$ and $S := \{i : \widehat{\alpha}_i > 0\}$.
Then $\widehat{f}(x) = \sum_{i \in S} \widehat{\alpha}_i y_i k(x, x_i) + \widehat{b}$, and KKT gives $i \in S \Rightarrow y_i \widehat{f}(x_i) \le 1$. Thus only support vectors enter $\widehat{f}$.

**Remark 5.3.4.** *Margins and Generalization*
Let $m_i(f) := y_i f(x_i)$ and $\widehat{R}_\gamma(f) := n^{-1} \sum_{i=1}^{n} \mathbf{1}\{m_i(f) \le \gamma\}$.
Bounds control $R(f) := \mathbb{P}(Yf(X) \le 0)$ by $\widehat{R}_\gamma(f)$ plus a complexity proxy $\mathcal{C}_\gamma(f) \asymp \|f\|^2 / \gamma^2$.
For linear $f_{w,b}$ with $\min_i m_i(f) \ge 1$, the geometric margin is $\rho = 1/\|w\|_2$.

# 6 Trees and Ensemble Averaging

## 6.1 Tree Models and Bagging

Tree-based methods partition the input space into simple regions and fit a local rule in each region. Ensemble averaging methods keep this local modelling idea but stabilise it by resampling, voting, and decorrelating the individual trees.

**Definition 6.1.1.** *Decision Tree*
Let $\mathcal{X} \subseteq \mathbb{R}^p$ and let $\mathcal{A}$ be the prediction space.
A decision tree $T$ is a finite binary recursive partition of $\mathcal{X}$ with a leaf value in $\mathcal{A}$ on each terminal region.

(i) *Split*: a split at node $R \subseteq \mathcal{X}$ is a feature-threshold pair $(j, t)$ with $j \in \{1, \dots, p\}$ and $t \in \mathbb{R}$. It produces children $R_L := R \cap \{x : x_j \le t\}$ and $R_R := R \cap \{x : x_j > t\}$.

(ii) *Terminal partition*: after splitting stops, the leaves of $T$ are nonempty regions $R_1(T), \dots, R_M(T)$ with $\mathcal{X} = \bigcup_{m=1}^{M} R_m(T)$ and $R_m(T) \cap R_{m'}(T) = \varnothing$ for $m \neq m'$.

(iii) *Leaf rule*: a leaf assignment $a_m \in \mathcal{A}$ for $m = 1, \dots, M$ defines $\widehat{f}_T : \mathcal{X} \to \mathcal{A}$ by $\widehat{f}_T(x) := a_m$ for the unique $m$ such that $x \in R_m(T)$.

## 6.2 CART Splits and Pruning

**Definition 6.2.1.** *CART Regression-Tree Notation and Scores*
Let $D_n = \{(x_i, y_i)\}_{i=1}^n \subseteq \mathcal{X} \times \mathbb{R}$. For $R \subseteq \mathcal{X}$, set $I_R := \{i : x_i \in R\}$ and $n_R := |I_R|$.

(i) *Cell loss*: if $n_R > 0$, define $\bar{y}_R := n_R^{-1} \sum_{i \in I_R} y_i$ and $\mathrm{SSE}(R) := \sum_{i \in I_R} (y_i - \bar{y}_R)^2$.

(ii) *Tree predictor*: if $T$ has leaves $R_1(T), \dots, R_{|T|}(T)$, let $m_T(x)$ be the unique $m \in \{1, \dots, |T|\}$ with $x \in R_m(T)$ and set $\widehat{f}_T(x) := \bar{y}_{R_{m_T(x)}(T)}$.

(iii) *Split children*: for $a = (j, s) \in \{1, \dots, p\} \times \mathbb{R}$, set $R_L(a) := R \cap \{x : x_j \le s\}$ and $R_R(a) := R \cap \{x : x_j > s\}$.

(iv) *Admissible splits*: set $\mathcal{A}_R := \{a \in \{1, \dots, p\} \times \mathbb{R} : n_{R_L(a)} > 0, n_{R_R(a)} > 0\}$.

(v) *Split score*: for $a \in \mathcal{A}_R$, set $S_R(a) := \mathrm{SSE}(R_L(a)) + \mathrm{SSE}(R_R(a))$.

(vi) *Pruning score*: for $\alpha \ge 0$, set $C_\alpha(T) := \sum_{m=1}^{|T|} \mathrm{SSE}(R_m(T)) + \alpha |T|$.

**Method 6.2.2.** *CART-Style Regression Trees*

(i) *Grow*: split eligible leaves by $a_R \in \operatorname{argmin}_{a \in \mathcal{A}_R} S_R(a)$ until stopping, yielding $T_0$.

(ii) *Prune*: for each $\alpha \ge 0$, choose $T_\alpha \in \operatorname{argmin}_{T \preceq T_0} C_\alpha(T)$.

(iii) *Select*: choose $\widehat{\alpha} \ge 0$ by validation or cross-validation and return $T_{\widehat{\alpha}}$.

Deep growth lowers bias; the cost-complexity penalty $\alpha|T|$ controls variance by subtree selection.

**Definition 6.2.3.** *Classification Tree Impurity*
For node $m$, let $N_{mk} := |\{i : x_i \in R_m, y_i = k\}|$, $N_m := \sum_{k=1}^{K} N_{mk}$, $\widehat{p}_{mk} := \frac{N_{mk}}{N_m}$. Common impurity scores are:

(i) *Misclassification*: $I_{\mathrm{mis}}(m) := 1 - \max_{1 \le k \le K} \widehat{p}_{mk}$.

(ii) *Gini*: $I_{\mathrm{Gini}}(m) := \sum_{k=1}^{K} \widehat{p}_{mk}(1 - \widehat{p}_{mk}) = 1 - \sum_{k=1}^{K} \widehat{p}_{mk}^2$.

(iii) *Entropy*: $I_{\mathrm{ent}}(m) := -\sum_{k : \widehat{p}_{mk} > 0} \widehat{p}_{mk} \log \widehat{p}_{mk}$.

The usual leaf prediction is the majority class, $\widehat{k}(m) \in \operatorname{argmax}_{1 \le k \le K} \widehat{p}_{mk}$.
With unequal losses $L_{kk} = 0$, choose the class minimizing expected loss, $\widehat{\ell}(m) \in \operatorname{argmin}_{1 \le \ell \le K} \sum_{k=1}^{K} L_{k\ell} \widehat{p}_{mk}$.

**Remark 6.2.4.** *Tree Practicalities*
For categorical levels $\mathcal{A}_j$, split $B \subset \mathcal{A}_j$ versus $B^c$.
For binary responses, set $I_a := \{i : X_{ij} = a\}$ and order levels by $\widehat{p}_a := |I_a|^{-1} \sum_{i \in I_a} Y_i$.
Missing values can be an NA level or use a surrogate split $g'$ maximising $\sum_i \mathbf{1}\{g'(x_i) = g(x_i)\}$.
Many candidate splits inflate gain; control growth with $N_m \ge n_{\min}$, depth limits, validation, or pruning.

## 6.3 Ensembles, Random Forests, and Stacking

**Definition 6.3.1.** *Bagging and Random Forests*
Let $D_n = \{(x_i, y_i)\}_{i=1}^n$, $B \geq 1$, and $m \leq p$.

(i) *Bagging*: draw $D_1^*, \ldots, D_B^*$ from $D_n$, fit $T_1, \ldots, T_B$, and aggregate $\widehat{f}_{\mathrm{bag}}(x) := B^{-1} \sum_{b=1}^{B} T_b(x)$ for regression, with voting or averaged probabilities for classification.

(ii) *Random forest*: in addition to bagging, at each node $R$ of $T_b$ choose $M_{b,R} \subseteq \{1, \ldots, p\}$ with $|M_{b,R}| = m$ and optimise the split over features in $M_{b,R}$ only.

Bagging averages bootstrap perturbations; random split-feature subsets reduce $\mathrm{corr}(T_b, T_{b'})$ before averaging.

**Method 6.3.2.** *Bagged Prediction and Votes*
Let $D_n := \{(x_i, y_i)\}_{i=1}^n$ and $B \geq 1$; base outputs are $\widehat{f}$ for regression and $(\widehat{g}, \widehat{p}_1, \ldots, \widehat{p}_K)$ for classification.

(i) For $b = 1, \ldots, B$, draw a bootstrap sample $D_b^*$ from $D_n$ and fit the base learner, giving $\widehat{f}^{*b}$ or $(\widehat{g}^{*b}, \widehat{p}_{b1}, \ldots, \widehat{p}_{bK})$.

(ii) Return the task-specific aggregate: for regression, $\widehat{f}_{\mathrm{bag}}(x) := B^{-1} \sum_{b=1}^{B} \widehat{f}^{*b}(x)$; for classification, either

$$\widehat{g}_{\mathrm{vote}}(x) \in \operatorname*{argmax}_{1 \leq k \leq K} \sum_{b=1}^{B} \mathbf{1}\{\widehat{g}^{*b}(x) = k\} \text{ or } \bar{p}_k(x) := B^{-1} \sum_{b=1}^{B} \widehat{p}_{bk}(x) \text{ with } \widehat{g}_{\mathrm{prob}}(x) \in \operatorname*{argmax}_{1 \leq k \leq K} \bar{p}_k(x).$$

Averaging reduces perturbation variance for unstable base learners.
Random split-feature subsets additionally reduce $\mathrm{corr}(T_b, T_{b'})$.

**Theorem 6.3.3.** *Squared-Error Benefit of Averaging*
*Fix $x$, let $Z := \widehat{f}^*(x)$ be perturbation-random and independent of $Y \mid X = x$, and set $f_{\mathrm{ag}}(x) := \mathbb{E}_* Z$.*
*If $\mathbb{E}_* Z^2 < \infty$ and $\mathbb{E}_Y Y^2 < \infty$, then $\mathbb{E}_{*,Y}[(Y - Z)^2] = \mathbb{E}_Y[(Y - f_{\mathrm{ag}}(x))^2] + \mathbb{E}_*[(Z - f_{\mathrm{ag}}(x))^2]$.*

*Proof.* Let $\mu := f_{\mathrm{ag}}(x) = \mathbb{E}_* Z$.
Since $Y - \mu$ and $Z - \mu$ are independent and $\mathbb{E}_*(Z - \mu) = 0$, $\mathbb{E}_{*,Y}[(Y - \mu)(Z - \mu)] = \mathbb{E}_Y(Y - \mu)\mathbb{E}_*(Z - \mu) = 0$.
Expanding $Y - Z = (Y - \mu) - (Z - \mu)$ gives $\mathbb{E}_{*,Y}[(Y - Z)^2] = \mathbb{E}_Y[(Y - \mu)^2] + \mathbb{E}_*[(Z - \mu)^2]$.

For independent fits $Z_b$, the average $\widehat{f}_B(x) := B^{-1} \sum_{b=1}^{B} Z_b$ has mean $\mu$ and perturbation variance $\mathrm{Var}_*(Z)/B$.

*Structural Commentary*: squared loss decomposes into target risk plus perturbation variance.
*Key Insight*: bagging estimates $\mathbb{E}_* \widehat{f}^*(x)$; unstable base learners gain most because $\mathrm{Var}_*(Z)$ is large. □

**Theorem 6.3.4.** *Correlated Ensemble Variance*
*Let $Z_1, \ldots, Z_B$ have $\mathrm{Var}(Z_b) = \sigma^2 > 0$ and $\mathrm{Corr}(Z_b, Z_c) = \rho$ for $b \neq c$.*
*For $\bar{Z}_B := \frac{1}{B} \sum_{b=1}^{B} Z_b$, $\mathrm{Var}(\bar{Z}_B) = \rho\sigma^2 + \frac{1-\rho}{B}\sigma^2$; hence, if $\rho \geq 0$, $\mathrm{Var}(\bar{Z}_B) \to \rho\sigma^2$ as $B \to \infty$.*

*Proof.* Since $\mathrm{Cov}(Z_b, Z_c) = \rho\sigma^2$ for $b \neq c$, $\mathrm{Var}(\bar{Z}_B) = B^{-2} \left\{ \sum_{b=1}^{B} \mathrm{Var}(Z_b) + 2 \sum_{1 \leq b < c \leq B} \mathrm{Cov}(Z_b, Z_c) \right\}$.
Substitution gives $\mathrm{Var}(\bar{Z}_B) = B^{-2} \left\{ B\sigma^2 + B(B-1)\rho\sigma^2 \right\} = \rho\sigma^2 + \frac{1-\rho}{B}\sigma^2$.
*Structural Commentary*: $\mathrm{Var}(\bar{Z}_B) = \rho\sigma^2 + (1-\rho)\sigma^2/B$ separates shared and idiosyncratic variance.
*Key Insight*: increasing $B$ shrinks only $(1-\rho)\sigma^2/B$; Random Forest lowers $\rho$ by split randomisation. □

**Algorithm 13:** Random Forest

**Require:** Training data $(x_i, y_i)_{i=1}^n$, number of trees $B$, candidate split count $m \leq p$, tree stopping rule
**Ensure:** Trees $T_1, \ldots, T_B$ and an aggregated predictor
**for** $b = 1, \ldots, B$ **do**
  Draw a bootstrap resample $S^{*b}$ of size $n$
  Grow a tree $T_b$ on $S^{*b}$
  **for** each nonterminal node while growing $T_b$ **do**
    Select $m$ of the $p$ predictors at random
    Choose the best split using only those $m$ predictors
    Split the node and continue until the stopping rule is met
  **end for**
**end for**
For regression, return $\widehat{f}_{\mathrm{rf}}(x) = B^{-1} \sum_{b=1}^{B} T_b(x)$
For classification, return a majority vote or averaged class-probability rule from $T_1(x), \ldots, T_B(x)$




Figure 266: Blue bootstraps train green trees with $m$ split candidates; black arrows aggregate, red dashed arrows give OOB prediction for $i$.

**Method 6.3.5.** *Out-of-Bag Error and Forest Importance*
Let $I_b^*$ index bootstrap sample for $T_b$, $O_i := \{b : i \notin I_b^*\}$, $\mathcal{I}_{\text{oob}} := \{i : |O_i| > 0\}$, and $L$ be prediction loss.

(i) For each $i \in \mathcal{I}_{\text{oob}}$, construct $\widehat{h}_{\text{oob}}(x_i)$: for regression, set $\widehat{h}_{\text{oob}}(x_i) := |O_i|^{-1} \sum_{b \in O_i} T_b(x_i)$; for classification, use an OOB vote or averaged class-probability vector over $b \in O_i$.
(ii) Set $\widehat{R}_{\text{oob}} := |\mathcal{I}_{\text{oob}}|^{-1} \sum_{i \in \mathcal{I}_{\text{oob}}} L(y_i, \widehat{h}_{\text{oob}}(x_i))$ and update the estimate as $B$ changes.

For feature $j$, let $\mathcal{S}_{bj}$ be $j$-splits of $T_b$, with gains $\Delta Q_{bs}$ and OOB losses $R_b^{\text{oob}}, R_{bj}^{\pi}$ before/after permuting $X_j$.
Define $I_j^{\text{split}} := \sum_{b=1}^{B} \sum_{s \in \mathcal{S}_{bj}} \Delta Q_{bs}$ and $I_j^{\pi} := B^{-1} \sum_{b=1}^{B} (R_{bj}^{\pi} - R_b^{\text{oob}})$.
Both are fitted-forest summaries; $I_j^{\pi}$ is not a refit-without-$X_j$ loss contrast.

**Remark 6.3.6.** *Proximities, Tuning, and Overfitting*
Let $L_b(i)$ denote the terminal leaf of $x_i$ in $T_b$, and set $O_{ij} := \{b : i, j \notin I_b^*\}$.
Define $P_{ij} := B^{-1} \sum_{b=1}^{B} \mathbf{1}\{L_b(i) = L_b(j)\}$ and, when $|O_{ij}| > 0$, $P_{ij}^{\text{oob}} := |O_{ij}|^{-1} \sum_{b \in O_{ij}} \mathbf{1}\{L_b(i) = L_b(j)\}$.
Thus $P$ is a fitted-forest similarity; it is neither data-intrinsic nor generally a metric.
Tuning effects are parameter-specific: $m \downarrow$ lowers $\text{corr}(T_b, T_{b'})$ and can raise bias; $n_{\min} \uparrow$ smooths terminal leaves; $B \uparrow$ lowers finite-ensemble Monte Carlo error but does not regularise the limiting forest.
Defaults $m_{\text{cls}} \approx \sqrt{p}$ and $m_{\text{reg}} \approx p/3$ are starting values only.

**Method 6.3.7.** *Error-Correcting Output Codes*
For training data $(x_i, y_i)_{i=1}^n$ with $y_i \in \{1, \dots, K\}$, let $d_H$ denote Hamming distance.
A code matrix $C \in \{0,1\}^{K \times L}$ assigns class $k$ the row $C_{k\cdot}$, and column $\ell$ induces binary labels $z_{i\ell} := C_{y_i \ell}$.

(i) Choose $C$ with distinct rows, equivalently $d_H(C_{k\cdot}, C_{k'\cdot}) > 0$ for $k \neq k'$.
(ii) For each $\ell = 1, \dots, L$, fit $\widehat{p}_\ell(x) \approx \mathbb{P}(Z_\ell = 1 \mid X = x)$ to $(x_i, z_{i\ell})_{i=1}^n$.
(iii) For a test point $x$, compute $\delta_k(x) := \sum_{\ell=1}^{L} |C_{k\ell} - \widehat{p}_\ell(x)|$ for $k = 1, \dots, K$ and return $\widehat{k}(x) \in \operatorname{argmin}_{1 \le k \le K} \delta_k(x)$.

Row separation gives error correction; redundant columns reduce variance and absorb column errors.

**Method 6.3.8.** *Stacking and Bumping*
Let $\widehat{f}_1, \dots, \widehat{f}_M$ be fitted candidates, $\widehat{f}_m^{-i}(x_i)$ the out-of-sample prediction for observation $i$ from candidate $m$, $\mathcal{W} \subseteq \mathbb{R}^M$ the admissible weight set, and $L$ a loss. Commonly $\mathcal{W} := \{w \in \mathbb{R}^M : w_m \ge 0, \sum_{m=1}^{M} w_m = 1\}$.

(i) *Stacking*: set $R_{\text{st}}(w) := \sum_{i=1}^{n} (y_i - \sum_{m=1}^{M} w_m \widehat{f}_m^{-i}(x_i))^2$.
Choose $\widehat{w}^{\text{st}} \in \operatorname{argmin}_{w \in \mathcal{W}} R_{\text{st}}(w)$, and return $\widehat{f}_{\text{st}}(x) := \sum_{m=1}^{M} \widehat{w}_m^{\text{st}} \widehat{f}_m(x)$.
(ii) *Bumping*: fit bootstrap candidates $\widehat{f}^{*1}, \dots, \widehat{f}^{*B}$, set $R_b^* := \sum_{i=1}^{n} L(y_i, \widehat{f}^{*b}(x_i))$.
Choose $\widehat{b} \in \operatorname{argmin}_{1 \le b \le B} R_b^*$, and return $\widehat{f}^{*\widehat{b}}$.

Coordinate weights $w = e_m$ give cross-validated model selection; general $w \in \mathcal{W}$ gives model averaging.
Stacking averages fixed candidates; bumping selects among resampled fits.

# 7 Additive Models and Adaptive Basis Methods

Additive and adaptive basis methods sit between rigid linear models and fully local learners. They keep prediction interpretable by building the fitted function from one-dimensional effects, boxes, hinges, or soft routing

weights.

## 7.1 Generalized Additive Models

**Definition 7.1.1.** *Generalized Additive Model*
Let $X = (X_1, \dots, X_p)$, $x = (x_1, \dots, x_p)$, $\mu(x) := \mathbb{E}[Y \mid X = x]$, $\alpha \in \mathbb{R}$, and $f_j : \mathcal{X}_j \to \mathbb{R}$.

(i) *Additive mean*: $\mu(x) = \alpha + \sum_{j=1}^{p} f_j(x_j)$.

(ii) *Generalized additive model*: for a link $g$ on the range of $\mu$, $g(\mu(x)) = \alpha + \sum_{j=1}^{p} f_j(x_j)$.

(iii) *Identifiability*: with data $x_i = (x_{i1}, \dots, x_{ip})$, require $\sum_{i=1}^{n} f_j(x_{ij}) = 0$ for all $j$; this separates $\alpha$ from the $f_j$.

(iv) *Standard links*: Gaussian $g(t) = t$; binomial $g(t) = \log\left(\frac{t}{1-t}\right)$ or $\Phi^{-1}(t)$; counts $g(t) = \log t$.

**Algorithm 14:** Backfitting for an Additive Regression Model

**Require:** Training data $(x_i, y_i)_{i=1}^n$, univariate smoothers $S_j$ for predictors $j = 1, \dots, p$
**Ensure:** Centered component fits $\widehat{f}_1, \dots, \widehat{f}_p$ and intercept $\widehat{\alpha}$
Set $\widehat{\alpha} \leftarrow n^{-1} \sum_{i=1}^{n} y_i$ and initialize $\widehat{f}_j(x_{ij}) \leftarrow 0$
**repeat**
**for** $j = 1, \dots, p$ **do**
$r_{ij} \leftarrow y_i - \widehat{\alpha} - \sum_{k \neq j} \widehat{f}_k(x_{ik})$ for $i = 1, \dots, n$
$\widehat{f}_j \leftarrow S_j(r_{1j}, \dots, r_{nj})$
Center $\widehat{f}_j$ by subtracting $n^{-1} \sum_i \widehat{f}_j(x_{ij})$
**end for**
**until** the component functions change by less than the chosen tolerance

**Method 7.1.2.** *Local Scoring for Additive Logistic Regression*
For binary data $(x_i, y_i)_{i=1}^n$, $y_i \in \{0, 1\}$, use the additive predictor $\eta_i = \alpha + \sum_{j=1}^{p} f_j(x_{ij})$.

(i) *Mean*: set $p_i = \frac{1}{1+\exp(-\eta_i)}$ for $i = 1, \dots, n$.

(ii) *Working pair*: set $z_i = \eta_i + \frac{y_i - p_i}{p_i(1-p_i)}$ and $w_i = p_i(1 - p_i)$.

(iii) *Weighted fit*: update $(\alpha, f_1, \dots, f_p)$ by weighted backfitting, minimizing $\sum_{i=1}^{n} w_i \{z_i - \alpha - \sum_{j=1}^{p} f_j(x_{ij})\}^2$.

(iv) *Iteration*: recompute $\eta_i$ and repeat (i)–(iii) until the additive predictor converges.

## 7.2 Adaptive Basis and Rule Methods

**Method 7.2.1.** *MARS: Adaptive Piecewise-Linear Basis Search*
For data $(x_i, y_i)_{i=1}^n$ and $x \in \mathbb{R}^p$, use reflected hinges $(x_j - t)_+$ and $(t - x_j)_+$ with knots $t \in \{x_{ij} : i = 1, \dots, n\}$.
A MARS fit is $\widehat{f}(x) = \widehat{\beta}_0 + \sum_{m=1}^{M} \widehat{\beta}_m h_m(x)$, with each $h_m$ a selected hinge product of order $\leq d_{\max}$.

(i) *Forward pass*: add the admissible hinge pair or product giving the largest decrease in $\sum_{i=1}^{n} (y_i - \widehat{f}(x_i))^2$.

(ii) *Backward pass*: prune the forward sequence.
Score model $\lambda$ by $\mathrm{GCV}(\lambda) = \frac{\sum_{i=1}^{n} (y_i - \widehat{f}_\lambda(x_i))^2}{\{1 - M(\lambda)/n\}^2}$, with effective size $M(\lambda)$.

(iii) *Selection*: choose the pruned model with smallest GCV.

The cap $d_{\max}$ controls interaction order; $d_{\max} = 1$ gives an additive piecewise-linear model.

**Method 7.2.2.** *PRIM Box Peeling*
For data $(x_i, y_i)_{i=1}^n$, a box $B = \prod_{j=1}^{p} [a_j, b_j]$, set $I(B) = \{i : x_i \in B\}$ and $\bar{y}(B) = |I(B)|^{-1} \sum_{i \in I(B)} y_i$.
Fix peeling fraction $\rho \in (0, 1)$ and minimum size $m_{\min}$.

(i) *Initialise*: take $B$ to be the full training box.

(ii) *Peel*: while $|I(B)| > m_{\min}$, replace $B$ by the one-face deletion $B' \subset B$ removing a $\rho$ fraction of $I(B)$ with maximal $\bar{y}(B')$.

(iii) *Paste*: expand a face when the resulting box $B'$ increases $\bar{y}(B')$.

(iv) *Select*: choose $B^*$ from the peeling-pasting path by validation or constraints; remove $I(B^*)$ and repeat for additional boxes.

PRIM returns high-mean rectangular regions rather than a recursive partition.

**Algorithm 15:** Patient Rule Induction Method

**Require:** Data $(x_i, y_i)_{i=1}^n$, peel fraction $\rho$, minimum size $m_{\min}$, box count $q$
**Ensure:** Selected boxes $\mathcal{S} = (B_1^*, \dots, B_q^*)$
$D \leftarrow \{1, \dots, n\}$, $\mathcal{S} \leftarrow ()$
**for** $r = 1, \dots, q$ **do**
$I_D(B) := \{i \in D : x_i \in B\}$, $\bar{y}_D(B) := |I_D(B)|^{-1} \sum_{i \in I_D(B)} y_i$
$B \leftarrow \prod_{j=1}^{p} [\min_{i \in D} x_{ij}, \max_{i \in D} x_{ij}]$, $\mathcal{P} \leftarrow \{B\}$
**repeat**
$\mathcal{C}_\rho(B) \leftarrow \{B' \subset B : |I_D(B')| \geq (1-\rho)|I_D(B)|\}$ among one-face deletions
$B \leftarrow \operatorname{argmax}_{B' \in \mathcal{C}_\rho(B)} \bar{y}_D(B')$, $\mathcal{P} \leftarrow \mathcal{P} \cup \{B\}$
**until** $|I_D(B)| \leq m_{\min}$
$\mathcal{E}(B) \leftarrow \{B' \supset B : \bar{y}_D(B') > \bar{y}_D(B)\}$ among one-face expansions
**if** $\mathcal{E}(B) \neq \varnothing$ **then**
$B \leftarrow \operatorname{argmax}_{B' \in \mathcal{E}(B)} \bar{y}_D(B')$, $\mathcal{P} \leftarrow \mathcal{P} \cup \{B\}$
**end if**
$B_r^* \leftarrow \text{Select}(\mathcal{P})$, $\mathcal{S} \leftarrow (\mathcal{S}, B_r^*)$, $D \leftarrow D \setminus I_D(B_r^*)$
**end for**

**Remark 7.2.3.** *Adaptive Structure Versus Tree Structure*

(i) *CART*: represents $\mathcal{X}$ by a binary-tree partition; leaves carry constant local fits.
(ii) *PRIM*: represents high-response regions by boxes $B \subset \mathcal{X}$ without requiring a tree partition.
(iii) *MARS*: represents $\widehat{f}$ by hinge products $h_m$ and coefficients $\widehat{\beta}_m$, linear in $\widehat{\beta}$.

CART emphasizes tree interpretability; PRIM keeps box rules; MARS trades boxes for smoother hinge bases.

**Remark 7.2.4.** *Soft Trees and Mixtures of Experts*
Let node $v$ have children $1, \dots, K_v$, gates $\gamma_{vk}$, leaves $\mathcal{L}$, leaf path $P(\ell)$, and expert $m_\ell(x; \theta_\ell)$.

(i) *Gate*: $g_{vk}(x) = \frac{\exp(\gamma_{vk}^\top x)}{\sum_{r=1}^{K_v} \exp(\gamma_{vr}^\top x)}$, so $\sum_{k=1}^{K_v} g_{vk}(x) = 1$.
(ii) *Leaf weight*: $\pi_\ell(x) = \prod_{(v,k) \in P(\ell)} g_{vk}(x)$.
(iii) *Prediction*: $\widehat{m}(x) = \sum_{\ell \in \mathcal{L}} \pi_\ell(x) m_\ell(x; \theta_\ell)$.
(iv) *Fitting*: estimate $(\gamma, \theta)$ by maximizing the mixture likelihood, often via EM-type alternating updates.

Soft trees replace hard branch indicators by input-dependent probabilities while retaining tree-local experts.

**Remark 7.2.5.** *Missing Predictors*
For observation $i$, let $O_i = \{j : x_{ij} \text{ observed}\}$ and $M_i = \{1, \dots, p\} \setminus O_i$; specify prediction from $x_{i,O_i}$.

(i) *Trees*: use surrogate splits or a categorical missing level.
(ii) *Single imputation*: replace $x_{ij}$, $j \in M_i$, by $\widehat{x}_{ij}$ before fitting additive or basis models.
(iii) *Model-based imputation*: estimate $\widehat{x}_{ij}$ from a feature model $\widehat{m}_j(x_{i,O_i})$.
(iv) *Multiple imputation*: fit on completed data sets $X^{(1)}, \dots, X^{(L)}$ to propagate imputation uncertainty.

Single imputation treats $\widehat{x}_{ij}$ as observed; multiple imputation propagates missing-data uncertainty.

# 8 Boosting and Additive Trees

## 8.1 Stagewise Boosting

Boosting is a stagewise way to build a strong predictor from simple base learners. The key object is an additive score: later terms are trained against the current fit rather than fit independently.

**Definition 8.1.1.** *Boosting*
Let $\mathcal{H}$ be a weak-learner class and set $F_0 \equiv 0$.

(i) *Additive score*: $F_M(x) = \sum_{m=1}^{M} \alpha_m h_m(x)$, with $h_m \in \mathcal{H}$ and $\alpha_m \in \mathbb{R}$.

(ii) *Stagewise update*: choose $(\alpha_m, h_m)$ using residuals, gradients, or weights from $F_{m-1}$; $F_m = F_{m-1} + \alpha_m h_m$.
(iii) *Prediction*: regression uses $F_M(x)$; binary classification uses sign $F_M(x)$; multiclass scores use $\operatorname{argmax}_k F_{Mk}(x)$.

Boosting is the stagewise construction of $F_M$ from weak learners.

**Method 8.1.2.** *Forward Stagewise Additive Modeling*
Let $b(x;\gamma)$ range over a basis class and let $L(y, f)$ be a pointwise loss.

(i) *Initialize*: choose $f_0$.
(ii) *Fit step*: at stage $m$, take $(\beta_m, \gamma_m) \in \operatorname{argmin}_{\beta,\gamma} \sum_{i=1}^{n} L\{y_i, f_{m-1}(x_i) + \beta b(x_i;\gamma)\}$.
(iii) *Update*: set $f_m(x) = f_{m-1}(x) + \beta_m b(x;\gamma_m)$.
(iv) *Stop*: after $M$ stages, return $f_M$.

Previously added terms are fixed; for squared-error loss, step (ii) fits the current residuals.

**Remark 8.1.3.** *Stagewise Paths and Lasso Geometry*

Let $\mathcal{T} = \{T_1, \ldots, T_K\}$ be a tree dictionary and $f_\alpha(x) = \sum_{k=1}^{K} \alpha_k T_k(x)$.

For large $K$, direct least squares is ill posed; a penalized target is $\min_\alpha \{\sum_{i=1}^{n} (y_i - f_\alpha(x_i))^2 + \lambda J(\alpha)\}$.

With $J(\alpha) = \sum_{k=1}^{K} |\alpha_k|$, active trees are sparse.
Small-step boosting traces a forward-stagewise path through this dictionary; shrinkage shortens each move.

## 8.2 AdaBoost and Exponential Loss

**Algorithm 16:** AdaBoost.M1

**Require:** Binary training data $(x_i, y_i)$ with $y_i \in \{-1, 1\}$, weighted classification learner, iterations $M$
**Ensure:** Classifier $G(x)$

Initialize $w_i \leftarrow 1/n$ for $i = 1, \ldots, n$
**for** $m = 1, \ldots, M$ **do**
  Fit a classifier $G_m$ using observation weights $w_i$
  $\text{err}_m \leftarrow \dfrac{\sum_{i=1}^{n} w_i \mathbf{1}_{\{y_i \neq G_m(x_i)\}}}{\sum_{i=1}^{n} w_i}$
  $\alpha_m \leftarrow \log\left(\dfrac{1 - \text{err}_m}{\text{err}_m}\right)$
  **for** $i = 1, \ldots, n$ **do**
    $w_i \leftarrow w_i \exp\{\alpha_m \mathbf{1}_{\{y_i \neq G_m(x_i)\}}\}$
  **end for**
**end for**
**return** $G(x) = \operatorname{sign}\left(\sum_{m=1}^{M} \alpha_m G_m(x)\right)$




Figure 267: AdaBoost alternates fitting weak learners with updating observation weights, then aggregates them by a weighted vote.

**Proposition 8.2.1.** *AdaBoost as Exponential-Loss Stagewise Fitting*
*For $y_i \in \{-1,1\}$, use $L(y,f)=\exp(-yf)$. Fix stage $m$, current score $f_{m-1}$, weights $w_i^{(m)} := \exp\{-y_i f_{m-1}(x_i)\}$, and classifier $G:\mathcal{X}\to\{-1,1\}$. Set $Q_m(\beta,G) := \sum_{i=1}^{n} L\{y_i, f_{m-1}(x_i)+\beta G(x_i)\}$ and $E_m(G) := \sum_{i=1}^{n} w_i^{(m)} \mathbf{1}_{\{y_i \neq G(x_i)\}}$.*

*(i) For fixed $\beta>0$, $\operatorname{argmin}_G Q_m(\beta,G) = \operatorname{argmin}_G E_m(G)$.*

*(ii) If $G_m \in \operatorname{argmin}_G E_m(G)$ and $\mathrm{err}_m := E_m(G_m)/\sum_{i=1}^{n} w_i^{(m)}$, then $\beta_m = \frac{1}{2}\log\left(\frac{1-\mathrm{err}_m}{\mathrm{err}_m}\right)$ minimizes $Q_m(\beta,G_m)$.*

*(iii) The induced weights satisfy $w_i^{(m+1)} = w_i^{(m)}\exp\{-\beta_m y_i G_m(x_i)\}$.*
*With $\alpha_m = 2\beta_m$, a misclassified point has relative factor $\exp(\alpha_m)$.*

**Remark 8.2.2.** *Exponential Loss and Class Probabilities*
For $Y\in\{-1,1\}$ and $\eta(x)=\mathbb{P}(Y=1\mid X=x)$, the population minimizer of $\mathbb{E}\{\exp(-Yf(X))\mid X=x\}$ is $f^*(x)=\frac{1}{2}\log\frac{\eta(x)}{1-\eta(x)}$. Hence $\operatorname{sign} f^*(x)$ is the Bayes rule. The binomial deviance $\log\{1+\exp(-2Yf(X))\}$ has the same minimizer but penalizes large negative margins less aggressively on finite samples.

## 8.3 Gradient Boosted Trees and Rule Ensembles

**Definition 8.3.1.** *Additive Tree Expansion*
Let $R_1,\dots,R_J$ be terminal regions, $\gamma_j\in\mathbb{R}$, and $\Theta=\{(R_j,\gamma_j): 1\le j\le J\}$.

(i) *Tree*: $T(x;\Theta)=\sum_{j=1}^{J}\gamma_j\mathbf{1}_{\{x\in R_j\}}$.

(ii) *Additive expansion*: $f_M(x)=\sum_{m=1}^{M} T(x;\Theta_m)$.

(iii) *Terminal fit*: with regions $R_{jm}$ fixed, $\widehat{\gamma}_{jm}\in\operatorname{argmin}_{\gamma}\sum_{x_i\in R_{jm}} L\{y_i, f_{m-1}(x_i)+\gamma\}$.

**Algorithm 17:** Gradient Tree Boosting

**Require:** Training data $(x_i,y_i)$, differentiable loss $L(y,f)$, tree sizes $J_m$, learning rate $0<\nu\le 1$, iterations $M$
**Ensure:** Additive predictor $f_M$

$f_0(x)\leftarrow \arg\min_{\gamma}\sum_{i=1}^{n} L(y_i,\gamma)$
**for** $m=1,\dots,M$ **do**
  **for** $i=1,\dots,n$ **do**
    $r_{im}\leftarrow -\left[\frac{\partial L(y_i,f(x_i))}{\partial f(x_i)}\right]_{f=f_{m-1}}$
  **end for**
  Fit a regression tree to targets $r_{im}$, giving terminal regions $R_{jm}$, $j=1,\dots,J_m$
  **for** $j=1,\dots,J_m$ **do**
    $\gamma_{jm}\leftarrow \arg\min_{\gamma}\sum_{x_i\in R_{jm}} L\{y_i, f_{m-1}(x_i)+\gamma\}$
  **end for**
  $f_m(x)\leftarrow f_{m-1}(x)+\nu\sum_{j=1}^{J_m}\gamma_{jm}\mathbf{1}_{\{x\in R_{jm}\}}$
**end for**
**return** $f_M$

**Remark 8.3.2.** *Pseudo-Residuals*
At stage $m$, pseudo-residuals are negative pointwise gradients $r_{im} := -\partial_f L(y_i,f)|_{f=f_{m-1}(x_i)}$.
Gradient boosting fits the next tree to $(x_i,r_{im})$.

(i) *Squared error*: $r_{im}=y_i-f_{m-1}(x_i)$.
(ii) *Absolute error*: $r_{im}=\operatorname{sign}\{y_i-f_{m-1}(x_i)\}$, away from zero residuals.
(iii) *Multinomial deviance*: $r_{ikm}=\mathbf{1}_{\{y_i=\mathcal{G}_k\}}-p_k(x_i)$, where $p_k$ is the current softmax probability.

**Algorithm 18:** Gradient Boosting for $K$-Class Classification

**Require:** Encoded labels $y_{ik}\in\{0,1\}$, classes $k=1,\dots,K$, tree sizes $J_m$, iterations $M$
**Ensure:** Class-score functions $\widehat{f}_1,\dots,\widehat{f}_K$
Initialize $f_{k0}(x)\leftarrow 0$ for $k=1,\dots,K$

**for** $m = 1, \dots, M$ **do**
  $p_k(x) \leftarrow \dfrac{\exp(f_{k,m-1}(x))}{\sum_{\ell=1}^{K} \exp(f_{\ell,m-1}(x))}$ for $k = 1, \dots, K$
  **for** $k = 1, \dots, K$ **do**
    Compute $r_{ikm} \leftarrow y_{ik} - p_k(x_i)$ for $i = 1, \dots, n$
    Fit a regression tree to targets $r_{ikm}$, giving terminal regions $R_{jkm}$, $j = 1, \dots, J_m$
    **for** $j = 1, \dots, J_m$ **do**
      $\gamma_{jkm} \leftarrow \dfrac{K-1}{K} \dfrac{\sum_{x_i \in R_{jkm}} r_{ikm}}{\sum_{x_i \in R_{jkm}} |r_{ikm}|(1 - |r_{ikm}|)}$
    **end for**
    $f_{km}(x) \leftarrow f_{k,m-1}(x) + \sum_{j=1}^{J_m} \gamma_{jkm} \mathbf{1}_{\{x \in R_{jkm}\}}$
  **end for**
**end for**
**return** $\widehat{f}_k(x) = f_{kM}(x)$ for $k = 1, \dots, K$

**Remark 8.3.3.** *Tree Size, Shrinkage, and Subsampling*
Boosted-tree complexity is controlled by iterations $M$, terminal counts $J_m$, shrinkage $\nu$, and sampling.

(i) *Iterations*: larger $M$ adds more trees and increases model capacity.
(ii) *Tree size*: a common $J$ caps interactions; stumps ($J = 2$) give main effects, and larger $J$ allows order up to $J - 1$.
(iii) *Shrinkage*: use $f_m = f_{m-1} + \nu T_m$ with $0 < \nu \leq 1$; smaller $\nu$ usually requires larger $M$.
(iv) *Subsampling*: fit $T_m$ on random observations to reduce cost and add averaging-like randomness.

These tuning choices trade optimization speed, interaction depth, and test risk.

**Method 8.3.4.** *Relative Importance and Partial Dependence*
Let $f$ be a fitted tree ensemble. For $S \subset \{1, \dots, p\}$, write $C = S^c$ and $x_i = (x_{iS}, x_{iC})$.

(i) *Relative importance*: for predictor $j$, sum the split-improvement scores over tree nodes splitting on $j$. For an ensemble, average over trees and rescale if desired.
(ii) *Partial dependence*: define $f_S(x_S) = \mathbb{E}_{X_C}\{f(x_S, X_C)\}$ and estimate $\widehat{f}_S(x_S) = n^{-1} \sum_{i=1}^{n} f(x_S, x_{iC})$.

Partial dependence averages over the marginal distribution of $X_C$.
It equals $\mathbb{E}\{f(X) \mid X_S = x_S\}$ only under compatible dependence structure.

**Principle 8.3.5.** *Bet on Sparsity*
Let $\mathcal{T} = \{T_1, \dots, T_K\}$ and $f_\alpha = \sum_{k=1}^{K} \alpha_k T_k$, with support $S(\alpha) = \{k : \alpha_k \neq 0\}$.
An $\ell^1$ penalty is useful when a low-risk predictor has $|S(\alpha)| \ll K$ and $|S(\alpha)|$ is small relative to $n$ and noise.
Dense signals with many small $\alpha_k$ remain hard to estimate.
Sparsity is meaningful only relative to $n$, noise level, and $\mathcal{T}$.

**Method 8.3.6.** *Lasso Post-Processing of Ensembles*
Let a fitted ensemble generate the fixed dictionary $\mathcal{T}_L = \{T_1, \dots, T_M\}$ and let $L$ be the chosen prediction loss.

(i) *Fit path*: compute $\widehat{\alpha}(\lambda) \in \operatorname{argmin}_{\alpha} \{\sum_{i=1}^{n} L(y_i, \alpha_0 + \sum_{m=1}^{M} \alpha_m T_m(x_i)) + \lambda \sum_{m=1}^{M} |\alpha_m|\}$.
(ii) *Select*: choose $\lambda$ by validation or an information criterion.
(iii) *Return*: use $\widehat{f}_\lambda(x) = \widehat{\alpha}_0 + \sum_{m:\widehat{\alpha}_m(\lambda) \neq 0} \widehat{\alpha}_m(\lambda) T_m(x)$.

The post-processed model can use fewer trees.
A useful generated dictionary is relevant and not nearly duplicated, with subsampling one way to diversify it.




Figure 268: An ensemble generator $\mathcal{A}_{\text{ens}}$ builds the dictionary $\mathcal{T}_L$; lasso selects the active set $\widehat{S}_\lambda$ for the sparse predictor $\widehat{f}_\lambda$.

**Method 8.3.7.** *Rule Ensembles*
Let $T_1, \dots, T_M$ be fitted trees. A node path $p$ defines the rule $r_p(x) = \mathbf{1}_{\{x \text{ satisfies all splits on } p\}}$.

(i) *Extract*: form $\mathcal{T}_{\text{rule}}^m = \{r_p : p \text{ is a node path in } T_m\}$.

(ii) *Aggregate*: set $\mathcal{T}_{\text{rule}} = \bigcup_{m=1}^{M} \mathcal{T}_{\text{rule}}^{m}$.
(iii) *Fit*: regress $y$ on $\mathcal{T}_{\text{rule}}$ by lasso or another regularized linear method.
(iv) *Augment*: optionally add original variables $x_j$ as linear terms.

Rules encode split interactions; added linear terms encode main linear effects.

# 9 High-Dimensional Supervised Learning

High-dimensional supervised learning studies prediction problems in which the number of candidate features is comparable to, or much larger than, the number of observations. The central difficulty is not merely computational size: many rules can fit the training data too well, while the scientific question often asks which features or feature groups are driving the prediction.

## 9.1 High-Dimensional Linear Rules

**Principle 9.1.1.** *Less Fitting in the $p \gg n$ Regime*
Let $X \in \mathbb{R}^{n\times p}$, $r = \operatorname{rank}(X)$, and $p \gg n$.

(i) If $r < p$, least-squares coefficients are nonunique since $\ker(X) \neq \{\mathbf{0}\}$.
(ii) Full-covariance discriminant rules are singular when covariance rank is $< p$.
Pooled rank is $\leq n - K$, and class-$k$ rank is $\leq n_k - 1$.
(iii) Impose effective dimension $< p$ via shrinkage, diagonal covariance, row-space fitting, sparsity, or screening.

Measure complexity by $\mathrm{df}(\lambda)$, as in Proposition 2.1.4: ridge keeps coordinates; sparse rules select features.

**Method 9.1.2.** *Diagonal LDA and Nearest Shrunken Centroids*
For classes $k = 1, \dots, K$ and features $j = 1, \dots, p$, let $\bar{x}_{kj}$ be the class mean, $\bar{x}_{\cdot j}$ the overall mean, $s_j$ the pooled within-class standard deviation, $\pi_k$ the class prior, $n_k$ the class size, $m_k := (\frac{1}{n_k} - \frac{1}{n})^{1/2}$, and $s_0 > 0$.

(i) *Diagonal LDA*: score $x$ by $\delta_k(x) := -\sum_{j=1}^{p} \frac{(x_j - \bar{x}_{kj})^2}{s_j^2} + 2\log\pi_k$ and set $\widehat{G}(x) := \arg\max_k \delta_k(x)$.
(ii) *Nearest shrunken centroids*: for $\Delta \geq 0$, set $d_{kj} := \frac{\bar{x}_{kj} - \bar{x}_{\cdot j}}{m_k(s_j + s_0)}$, $d'_{kj} := \operatorname{sign}(d_{kj})(|d_{kj}| - \Delta)_+$, and $\bar{x}'_{kj} := \bar{x}_{\cdot j} + m_k(s_j + s_0)d'_{kj}$; evaluate the diagonal LDA score with $\bar{x}'_{kj}$ in place of $\bar{x}_{kj}$.

Diagonal LDA is nearest-centroid classification after coordinatewise standardization. In nearest shrunken centroids, feature $j$ is removed iff $d'_{kj} = 0$ for all $k$; increasing $\Delta$ increases shrinkage and sparsity.




Figure 269: Soft thresholding $S_\Delta(d) = \operatorname{sign}(d)(|d| - \Delta)_+$. The shaded band $|d| \leq \Delta$ maps to zero; outside it, contrasts shrink toward zero by $\Delta$.

**Method 9.1.3.** *Quadratic Regularization for Linear Rules*
Let $\eta_k(x) := \beta_{k0} + x^T\beta_k$, $\lambda > 0$, and $\gamma \in [0, 1]$.

(i) *$L_2$ multinomial logits*: set $p_{\beta,k}(x) := \frac{\exp(\eta_k(x))}{\sum_{\ell=1}^{K}\exp(\eta_\ell(x))}$ and choose $\widehat{\beta} \in \arg\max_\beta \{\sum_{i=1}^{n} \log p_{\beta,g_i}(x_i) - \frac{\lambda}{2}\sum_{k=1}^{K}\|\beta_k\|_2^2\}$; leave $\beta_{k0}$ unpenalized.
(ii) *Covariance shrinkage*: set $\widehat{\Sigma}_\gamma := \gamma\widehat{\Sigma} + (1-\gamma)\operatorname{diag}(\widehat{\Sigma})$ and use $\widehat{\Sigma}_\gamma$ in the discriminant score.

Both variants impose quadratic stabilization without sparsity. The endpoint $\gamma = 0$ gives diagonal LDA; increasing $\gamma$ reintroduces off-diagonal covariance information.

**Proposition 9.1.4.** *Row-Space Reduction for Quadratic Penalties*
*Let $X \in \mathbb{R}^{n\times p}$ have rank $r$ and thin SVD $X = UDV^T = RV^T$, where $R := UD \in \mathbb{R}^{n\times r}$ and $V^TV = I_r$.*
*Fix $y \in \mathbb{R}^n$ and $\lambda > 0$.*

*(i) Ridge regression satisfies $\widehat{\beta}_\lambda^{\text{ridge}} = V(R^TR + \lambda I_r)^{-1}R^Ty$.*

*(ii) If $F(\beta_0, \beta) := \sum_{i=1}^{n} L(y_i, \beta_0 + x_i^T\beta) + \lambda \|\beta\|_2^2$ has a minimiser, then one satisfies $\widehat{\beta} \in \text{row}(X) = \text{col}(V)$.*

*Thus quadratic-penalty fitting may be performed in the $r$ row-space coordinates $r_i^T$ and mapped back by $\widehat{\beta} = V\widehat{\theta}$.*

*Proof.* Every $\beta \in \mathbb{R}^p$ decomposes uniquely as $\beta = V\theta + z$, where $\theta = V^T\beta$ and $z \in \text{col}(V)^\perp = \ker(X)$.
Then $X\beta = R\theta$ and $\|\beta\|_2^2 = \|\theta\|_2^2 + \|z\|_2^2$.

(i) For ridge, dropping $z$ preserves fitted values and weakly decreases the penalty, so a minimiser has $z = 0$. Thus $\widehat{\theta} \in \arg\min_\theta\{\|y - R\theta\|_2^2 + \lambda\|\theta\|_2^2\}$, giving $\widehat{\theta} = (R^TR + \lambda I_r)^{-1}R^Ty$ and $\widehat{\beta} = V\widehat{\theta}$.

(ii) If $(\widehat{\beta}_0, \widehat{\beta})$ minimises $F$ and $\widehat{\beta} = V\widehat{\theta} + \widehat{z}$, then $F(\widehat{\beta}_0, V\widehat{\theta}) \le F(\widehat{\beta}_0, \widehat{\beta})$, with the same linear predictors and no larger penalty. Hence $(\widehat{\beta}_0, V\widehat{\theta})$ is also a minimiser and $V\widehat{\theta} \in \text{col}(V)$.

*Structural Commentary*: Kernel components affect only the quadratic penalty, not the fitted values.
*Key Insight*: Quadratic penalties reduce fitting to row$(X)$. □

## 9.2 Sparse and Structured Penalties

**Method 9.2.1.** *Sparse and Structured Penalties*
Let $\lambda, \lambda_1, \lambda_2 \ge 0$ and $\alpha \in [0, 1]$. Sparse penalties replace quadratic shrinkage by nonsmooth coordinate penalties.

(i) *Lasso multinomial logits*: maximize $\sum_{i=1}^{n} \log \mathbb{P}(g_i \mid x_i) - \lambda \sum_{k=1}^{K}\sum_{j=1}^{p} |\beta_{kj}|$.

(ii) *Elastic net*: replace $|\beta_{kj}|$ by $\alpha|\beta_{kj}| + (1-\alpha)\beta_{kj}^2$.

(iii) *Fused lasso*: for ordered features, use $\lambda_1 \sum_{j=1}^{p} |\beta_j| + \lambda_2 \sum_{j=1}^{p-1} |\beta_{j+1} - \beta_j|$.

The $L_1$ term selects coordinates; the quadratic term shrinks correlated features.
Adjacent differences favor piecewise-constant coefficients.

**Method 9.2.2.** *Dantzig Selector*
Assume centered $y$ and standardised columns $x_j$ of $X$.
For residual $r(\beta) := y - X\beta$, the Dantzig selector has two equivalent constrained forms.

(i) *Correlation budget*: choose $\widehat{\beta}_s^{\text{DS}} \in \arg\min_\beta \|\beta\|_1$ subject to $\left\|X^Tr(\beta)\right\|_\infty \le s$.

(ii) *Coefficient budget*: choose $\widehat{\beta}_t^{\text{DS}} \in \arg\min_{\|\beta\|_1 \le t} \left\|X^Tr(\beta)\right\|_\infty$.

Both Dantzig and lasso control $\|\beta\|_1$ and monitor residual correlations $x_j^Tr(\beta)$.
Lasso minimises residual sum of squares, while Dantzig directly bounds $\left\|X^Tr(\beta)\right\|_\infty$.

**Method 9.2.3.** *Grouped Lasso*
Let $\{1, \dots, p\} = G_1 \sqcup \cdots \sqcup G_L$, with $|G_\ell| = p_\ell$, design block $X_\ell \in \mathbb{R}^{n\times p_\ell}$, coefficient block $\beta_\ell \in \mathbb{R}^{p_\ell}$, and $\lambda \ge 0$.
The grouped lasso solves $(\widehat{\beta}_0, \widehat{\beta}_1, \dots, \widehat{\beta}_L) \in \arg\min_{\beta_0,\beta_1,\dots,\beta_L} \left\{ \left\| y - \beta_0\mathbf{1} - \sum_{\ell=1}^{L} X_\ell\beta_\ell \right\|_2^2 + \lambda \sum_{\ell=1}^{L} \sqrt{p_\ell}\,\|\beta_\ell\|_2 \right\}$.
Since $\|\beta_\ell\|_2 = 0$ iff $\beta_\ell = \mathbf{0}$, the penalty selects or removes entire groups; $\sqrt{p_\ell}$ normalizes group size.

**Method 9.2.4.** *Relaxed, Adaptive, and Nonconvex Lasso Variants*
Let $\widehat{\beta}_\lambda^{\text{lasso}}$ be the lasso estimate, $A_\lambda := \{j : \widehat{\beta}_{\lambda,j}^{\text{lasso}} \ne 0\}$, $\widetilde{\beta}$ an initial estimate, $\nu > 0$, and $a \ge 2$.

(i) *Relaxed lasso*: fit the lasso to get $A_\lambda$, then refit on $X_{A_\lambda}$ by least squares or with penalty $\phi\lambda$, $0 \le \phi \le 1$.

(ii) *Adaptive lasso*: set $w_j := \frac{1}{|\widetilde{\beta}_j|^\nu}$ and minimize $\frac{1}{2}\|y - X\beta\|_2^2 + \lambda \sum_{j=1}^{p} w_j|\beta_j|$.

(iii) *SCAD*: use the nonconvex penalty $J_{a,\lambda}$ with $J'_{a,\lambda}(t) := \lambda\,\text{sign}(t)\left[\mathbf{1}_{\{|t|\le\lambda\}} + \frac{(a\lambda - |t|)_+}{(a-1)\lambda}\mathbf{1}_{\{|t|>\lambda\}}\right]$.

All three reduce lasso shrinkage bias for large signals.
Relaxed and adaptive lasso keep convex refits or weighted convex fits, while SCAD is nonconvex.

**Remark 9.2.5.** *Three Roles for Feature Selection*
In high-dimensional learning, feature selection has three distinct roles.

(i) *Preprediction screening*: reduce variance before fitting, as in nearest shrunken centroids.

(ii) *Sparse prediction*: estimate a sparse predictive rule, as in lasso or elastic-net logistic regression.
(iii) *Scientific discovery*: report selected features for separate assessment.
Multiplicity control is then inferential, not predictive.

## 9.3 Latent Factors and Distance-Based Classifiers

**Algorithm 19:** Supervised Principal Components

**Require:** Standardized feature matrix $X$, outcome $y$, association scores $s_j$, thresholds $0 \le \theta_1 < \cdots < \theta_K$, candidate component counts $\mathcal{M}$
**Ensure:** A selected principal-component predictor
**for** $k = 1, \dots, K$ **do**
$A_k \leftarrow \{j : |s_j| > \theta_k\}$
**for** $m \in \mathcal{M}$ **do**
Compute the first $m$ principal components of the submatrix $X_{A_k}$
Fit the outcome model using those components as predictors
Estimate held-out prediction error by cross-validation
**end for**
**end for**
Choose $(\theta_k, m)$ by the cross-validation criterion and refit using the selected columns

**Remark 9.3.1.** *Screening Before Components*
Let $s_j$ measure the marginal association between feature $x_j$ and outcome $y$, and set $A_\theta := \{j : |s_j| > \theta\}$. Supervised principal components compute PCs of $X_{A_\theta}$ rather than all of $X$.

(i) *Regression score*: take $s_j$ as a standardized univariate regression coefficient for $y$ on $x_j$.
(ii) *Survival score*: take $s_j$ as the one-feature Cox score for $x_j$.

Ordinary PCs follow variance in $X$ alone; screening makes the component outcome-guided.
The resulting component is a denoised latent factor from weak but aligned features.

**Method 9.3.2.** *Preconditioned Lasso*
Given $X$, $y$, and $\lambda > 0$, preconditioning uses a denoised response before sparse refitting.

(i) Fit supervised principal components and compute fitted values $\widehat{y} = (\widehat{y}_1, \dots, \widehat{y}_n)^T$.
(ii) Fit $\widehat{\beta}^{\text{pre}} \in \arg\min_\beta \{\frac{1}{2} \|\widehat{y} - X\beta\|_2^2 + \lambda \|\beta\|_1\}$.

Stage (i) denoises the target; stage (ii) selects original features that reproduce $\widehat{y}$.

**Method 9.3.3.** *Inner-Product and Distance-Only Classifiers*
Let $K_{ii'} := \langle x_i, x_{i'} \rangle$ be a positive semidefinite Gram matrix, $g_i$ class labels, $n_k = \#\{i : g_i = k\}$, and $M := \frac{1}{n}\mathbf{1}\mathbf{1}^T$.

(i) *Nearest neighbor*: compute $\|x_i - x_{i'}\|^2 = K_{ii} + K_{i'i'} - 2K_{ii'}$.
(ii) *Nearest centroid*: for test point $x_0$, compute $\|x_0 - \bar{x}_k\|^2 = \langle x_0, x_0 \rangle - \frac{2}{n_k} \sum_{g_i = k} \langle x_0, x_i \rangle + \frac{1}{n_k^2} \sum_{g_i = k} \sum_{g_{i'} = k} \langle x_i, x_{i'} \rangle$.
(iii) *Distance-only input*: for squared distances $\Delta_{ii'}^2$, set $B_{ii'} := -\frac{1}{2}\Delta_{ii'}^2$ and $\widetilde{K} := (I - M)B(I - M)$.

Thus nearest-neighbor, nearest-centroid, kernel PCA, and related classifiers require only inner products or centered distances, not explicit feature vectors.

**Example 9.3.4.** *String Kernels*
For strings over a finite alphabet, let $\mathcal{A}_m$ be the set of length-$m$ substrings, and let $\phi_a(x)$ count occurrences of $a$ in string $x$. Define $\Phi_m(x) := (\phi_a(x))_{a \in \mathcal{A}_m}$ and $K_m(x, x') := \langle \Phi_m(x), \Phi_m(x') \rangle$.
Although $|\mathcal{A}_m|$ grows exponentially in $m$, $K_m(x, x')$ can often be computed without forming $\Phi_m(x)$.
Thus support-vector or centroid classifiers can use sequence similarities directly.

# 10 Prototype and Nearest-Neighbor Methods

Prototype and nearest-neighbor classifiers are model-free rules built from distances in feature space. They can fit irregular decision boundaries and are often strong predictors, but their behavior is controlled by geometry: scaling, metric choice, neighborhood size, and storage of representative points. The local-averaging view appears in Method 1.4.2; this section records the classification and metric-design side.

## 10.1 Prototype Classifiers

**Definition 10.1.1.** *Labeled Prototype Rule*
Let $\mathcal{D}_n = \{(x_i, g_i)\}_{i=1}^n$ with $x_i \in \mathbb{R}^p$ and $g_i \in \{1, \dots, K\}$. Fix prototype counts $R_1, \dots, R_K$, distance $d$ on $\mathbb{R}^p$.

(i) A labeled prototype set is $\mathcal{M} = \{(m_j(k), k) : m_j(k) \in \mathbb{R}^p,\ 1 \le j \le R_k,\ 1 \le k \le K\}$.

(ii) For $x \in \mathbb{R}^p$, choose $(\widehat{j}, \widehat{k}) \in \underset{1\le k\le K,\ 1\le j\le R_k}{\arg\min} d(x, m_j(k))$ and define $\widehat{G}(x) = \widehat{k}$.

(iii) If $\mathcal{M} = \{(x_i, g_i) : 1 \le i \le n\}$, then $\widehat{G}$ is the one-nearest-neighbor classifier.

For quantitative features, Euclidean distance requires comparable coordinate scales.

**Method 10.1.2.** *k-Means Prototypes*
Let $I_k = \{i : g_i = k\}$, fix $R$ prototypes per class, and use $d(x, z) = \|x - z\|_2$.

(i) For each $k = 1, \dots, K$, choose $m_1(k), \dots, m_R(k) \in \mathbb{R}^p$ to minimize $\sum_{i \in I_k} \min_{1\le j\le R} \|x_i - m_j(k)\|_2^2$.

(ii) For $x \in \mathbb{R}^p$, choose $(\widehat{j}, \widehat{k}) \in \underset{1\le k\le K,\ 1\le j\le R}{\arg\min} \|x - m_j(k)\|_2$ and set $\widehat{G}(x) = \widehat{k}$.

The stored labeled prototype set is $\mathcal{M} = \{(m_j(k), k) : 1 \le j \le R,\ 1 \le k \le K\}$.
The rule has piecewise-linear Euclidean decision boundaries; since classes are compressed independently, prototypes need not move away from competing classes near a boundary.

**Algorithm 20:** Learning Vector Quantization

**Require:** Labeled data $(x_i, g_i)_{i=1}^n$, prototypes per class $R$, learning rates $\epsilon_t \downarrow 0$
**Ensure:** Labeled prototypes $m_j(k)$
  Initialize $m_1(k), \dots, m_R(k)$ from training observations in class $k$, for each $k$
  **for** $t = 1, 2, \dots$ **do**
    Sample a training index $i$
    Choose the closest prototype $(j, k)$ to $x_i$ over all classes and prototype indices
    **if** $g_i = k$ **then**
      $m_j(k) \leftarrow m_j(k) + \epsilon_t\big(x_i - m_j(k)\big)$
    **else**
      $m_j(k) \leftarrow m_j(k) - \epsilon_t\big(x_i - m_j(k)\big)$
    **end if**
  **end for**

**Remark 10.1.3.** *Boundary-Aware Prototype Motion*
For a sampled pair $(x_i, g_i)$, let $(j^\star, k^\star) \in \underset{1\le k\le K,\ 1\le j\le R_k}{\arg\min} d(x_i, m_j(k))$ and let $\epsilon_t > 0$.

(i) If $g_i = k^\star$, set $m_{j^\star}(k^\star) \leftarrow m_{j^\star}(k^\star) + \epsilon_t(x_i - m_{j^\star}(k^\star))$.

(ii) If $g_i \neq k^\star$, set $m_{j^\star}(k^\star) \leftarrow m_{j^\star}(k^\star) - \epsilon_t(x_i - m_{j^\star}(k^\star))$.

Thus every sampled class can move the nearest prototype of any class.
Unlike classwise $k$-means, LVQ is an online algorithmic rule rather than exact minimization of a fixed criterion.

**Method 10.1.4.** *Gaussian Mixtures as Soft Prototypes*
Let $\varphi_p(\cdot; \mu, \Sigma)$ be the $p$-variate Gaussian density. For class $k$, fix a prior $\rho_k$, component weights $\alpha_{kj}$, centers $\mu_{kj} \in \mathbb{R}^p$, covariances $\Sigma_{kj} \succ 0$, and $j = 1, \dots, R_k$, with $\rho_k, \alpha_{kj} \ge 0$, $\sum_{k=1}^K \rho_k = 1$, and $\sum_{j=1}^{R_k} \alpha_{kj} = 1$.

(i) Define the class density $f_k(x) = \sum_{j=1}^{R_k} \alpha_{kj} \varphi_p(x; \mu_{kj}, \Sigma_{kj})$.

(ii) For $x \in \mathbb{R}^p$, set $Z(x) = \sum_{\ell=1}^K \rho_\ell f_\ell(x) = \sum_{\ell=1}^K \sum_{r=1}^{R_\ell} \rho_\ell \alpha_{\ell r} \varphi_p(x; \mu_{\ell r}, \Sigma_{\ell r})$.

(iii) Compute $\tau_k(x) = \frac{\rho_k f_k(x)}{Z(x)}$ and $\omega_{kj}(x) = \frac{\rho_k \alpha_{kj} \varphi_p(x; \mu_{kj}, \Sigma_{kj})}{Z(x)}$. Set $\widehat{G}(x) \in \underset{1\le k\le K}{\arg\max}\, \tau_k(x)$.

The pairs $(\mu_{kj}, \Sigma_{kj})$ are soft prototypes and $\omega_{kj}(x)$ gives fractional membership.
If $\Sigma_{kj} = \sigma^2 I$ and all $\rho_k \alpha_{kj} > 0$, then as $\sigma^2 \downarrow 0$ the dominant component is nearest center $\mu_{kj}$ away from ties.

## 10.2 Nearest Neighbors

**Method 10.2.1.** *k-Nearest-Neighbor Classification*
Fix a distance $d$ on the feature space and an integer $k \in \{1, \dots, n\}$.

(i) For a query $x_0$, compute $d_i = d(x_i, x_0)$ for $i = 1, \dots, n$.

(ii) Let $N_k(x_0) \subset \{1,\dots,n\}$ be indices of $k$ smallest values among $d_1,\dots,d_n$. Set $\widehat{G}(x_0) \in \arg\max_{g\in\{1,\dots,K\}} \sum_{i\in N_k(x_0)} \mathbf{1}_{\{g_i=g\}}$.

Small $k$ gives lower bias and higher variance; larger $k$ smooths the decision boundary by averaging over more labels. The tuning parameter $k$ is usually chosen by validation or cross-validation.

**Proposition 10.2.2.** *Asymptotic One-Nearest-Neighbor Error*
*Assume $K \geq 2$ and fix $x$. Let $p_k = \mathbb{P}(G = k \mid X = x)$, choose $k^\star \in \arg\max_{1\leq k\leq K} p_k$, and set $L^\star(x) = 1 - p_{k^\star}$.*
*In the asymptotic one-nearest-neighbor idealisation, the query label $G$ and nearest-neighbor label $G'$ are independent draws from $(p_1,\dots,p_K)$, and the classifier predicts $G'$.*

*(i)* $L_{1\mathrm{NN}}(x) = \mathbb{P}(G' \neq G) = \sum_{k=1}^{K} p_k(1-p_k) = 1 - \sum_{k=1}^{K} p_k^2$.

*(ii)* $L^\star(x) \leq L_{1\mathrm{NN}}(x) \leq 2L^\star(x) - \frac{K}{K-1}\big(L^\star(x)\big)^2 \leq 2L^\star(x)$.

*Proof.*

(i) Independence gives $\mathbb{P}(G' = G) = \sum_{k=1}^{K} p_k^2$, hence $L_{1\mathrm{NN}}(x) = 1 - \sum_{k=1}^{K} p_k^2$.

(ii) Let $p_\star = p_{k^\star}$ and $q = 1 - p_\star = L^\star(x)$.

Since $p_k \leq p_\star$ for every $k$, $\sum_{k=1}^{K} p_k^2 \leq p_\star \sum_{k=1}^{K} p_k = p_\star$, so $L_{1\mathrm{NN}}(x) \geq 1 - p_\star = L^\star(x)$.

Also $\sum_{k\neq k^\star} p_k = q$, so Cauchy-Schwarz gives $\sum_{k\neq k^\star} p_k^2 \geq \frac{q^2}{K-1}$.

Therefore $\sum_{k=1}^{K} p_k^2 \geq (1-q)^2 + \frac{q^2}{K-1}$ and $L_{1\mathrm{NN}}(x) \leq 2q - \frac{Kq^2}{K-1} \leq 2q = 2L^\star(x)$.

*Structural Commentary*: Dominant mass gives the lower bound; uniform residual mass gives the upper bound.
*Key Insight*: One-nearest-neighbor error is label-disagreement probability at fixed $x$. □

**Example 10.2.3.** *Tuning Depends on the Problem*
Let $X = (X_1,\dots,X_{10})$ have independent $\mathrm{Unif}(0,1)$ coordinates and let $Y \in \{0,1\}$.

(i) *Halfspace*: $Y = \mathbf{1}_{\{X_1 > 1/2\}}$; the decision boundary is the single hyperplane $X_1 = 1/2$.

(ii) *Checkerboard*: $Y = \mathbf{1}_{\{\mathrm{sign}(\prod_{j=1}^{3}(X_j - 1/2)) > 0\}}$.
The planes $X_j = 1/2$ for $j = 1,2,3$ form eight cells with alternating labels.

Across repeated simulations, the preferred neighborhood size $k$ or prototype count $R$ depended on the case. For memory-based and prototype rules, tuning is geometry-dependent, not a universal complexity constant.

**Example 10.2.4.** *Image Scene Classification by Local Pixel Features*
For each pixel $s$, let $b_s \in \mathbb{R}^4$ be its four spectral-band vector and let $g_s \in \{1,\dots,7\}$ be its land-use label. Replace $b_s$ by the local feature vector $z_s = (b_t : t \in \mathcal{N}_3(s)) \in \mathbb{R}^{36}$, where $\mathcal{N}_3(s)$ is the $3\times 3$ pixel patch centered at $s$. A five-nearest-neighbor rule on the features $z_s$ had reported test error about 9.5%, showing that local spatial features can make nearest-neighbor classification competitive for irregular class boundaries.




Figure 270: A $3 \times 3$ patch over four bands gives 36 features per pixel.

## 10.3 Invariant Distances and Adaptive Neighborhoods

**Method 10.3.1.** *Invariant Metrics and Tangent Distance*
Let $x, x' \in \mathbb{R}^p$, let $\mathcal{A}$ be a transformation family, and write $T_\alpha x$ for the transformed input.
For local transformations, let $V_x \in \mathbb{R}^{p\times r_x}$ span the tangent directions at $x$.

(i) *Orbit metric*: $d_{\mathrm{inv}}(x,x') = \inf_{\alpha,\beta\in\mathcal{A}} \|T_\alpha x - T_\beta x'\|_2$ compares the transformation orbits of $x$ and $x'$.

(ii) *Tangent metric*: $d_{\tan}(x,x') = \inf_{a,b} \|x + V_x a - x' - V_{x'} b\|_2$ compares tangent affine spaces.

The tangent metric discounts small local transformations without making large transformations cost-free.
Data augmentation encodes similar invariance by storing transformed copies, which becomes costly when many transformation directions are needed.




Figure 271: Tangent distance uses local tangents instead of orbits; $d_E$ is ordinary distance.

**Method 10.3.2.** *Discriminant Adaptive Nearest Neighbors*
DANN replaces the fixed Euclidean metric by a query-dependent quadratic metric.
For a query $x_0$, choose a pilot index set $P(x_0)$, final neighbor count $h$, and $\epsilon > 0$. From $\{(x_i, g_i) : i \in P(x_0)\}$ estimate local class proportions $\pi_k$, class means $\bar{x}_k$, local mean $\bar{x}$, and pooled within-class covariance $\mathbf{W} \succ 0$.

(i) Compute the local between-class covariance $\mathbf{B} = \sum_{k=1}^{K} \pi_k(\bar{x}_k - \bar{x})(\bar{x}_k - \bar{x})^T$.
(ii) Form $\mathbf{B}^\star = \mathbf{W}^{-1/2}\mathbf{B}\mathbf{W}^{-1/2}$ and $\mathbf{\Sigma}_{x_0} = \mathbf{W}^{-1/2}(\mathbf{B}^\star + \epsilon I)\mathbf{W}^{-1/2}$.
(iii) With $D_{x_0}(x_i, x_0) = (x_i - x_0)^T\mathbf{\Sigma}_{x_0}(x_i - x_0)$, let $N_h(x_0)$ index $h$ smallest distances.
(iv) Set $\widehat{G}(x_0) \in \arg\max_{g\in\{1,\dots,K\}} \sum_{i\in N_h(x_0)} \mathbf{1}_{\{g_i=g\}}$.

Whitening by $\mathbf{W}$ removes local within-class scale.
$\mathbf{B}^\star + \epsilon I$ stretches neighborhoods along directions with weak local class separation, and $\epsilon$ prevents degeneracy.




Figure 272: DANN replaces the Euclidean ball by a local ellipsoid, narrow along the class-change direction and elongated where class probabilities vary slowly.

**Method 10.3.3.** *Global Dimension Reduction for Nearest Neighbors*
Fix a target dimension $L \le p$. Let $\mathbf{B}_i$ be the local between-class matrix estimated at training point $x_i$.

(i) Compute the global class-separation matrix $\overline{\mathbf{B}} = N^{-1}\sum_{i=1}^{N}\mathbf{B}_i$.
(ii) Let $(\theta_\ell, e_\ell)_{\ell=1}^{p}$ be eigenpairs of $\overline{\mathbf{B}}$ with $\theta_1 \ge \cdots \ge \theta_p$.
(iii) Project by $P_L x = (e_1^T x, \dots, e_L^T x) \in \mathbb{R}^L$ and run nearest-neighbor classification on $P_L x_i$.

The rank-$L$ summary $\overline{\mathbf{B}}_{[L]} = \sum_{\ell=1}^{L}\theta_\ell e_\ell e_\ell^T$ solves $\min_{\mathrm{rank}(M)=L}\sum_{i=1}^{N}\mathrm{trace}[(\mathbf{B}_i - M)^2]$.
The reduced space is selected from local class-separation information, not unsupervised variance alone.

**Remark 10.3.4.** *Computation and Storage*
With $N$ stored observations in $\mathbb{R}^p$, direct nearest-neighbor search costs $O(Np)$ per query and stores $O(Np)$ coordinates. Storage can be reduced by replacing the retained set $S \subset \{1, \dots, N\}$.

(i) *Prototype compression*: store learned representatives rather than all training observations.
(ii) *Editing*: delete observations whose local neighborhoods have inconsistent labels.
(iii) *Condensing*: grow a small retained set by adding $i$ only if it misclassifies $(x_i, g_i)$.

Fast search reduces query cost in favorable geometries; compression, editing, and condensing reduce storage by keeping representatives or boundary-relevant observations.

# 11 Flexible and Mixture Discriminant Analysis

## 11.1 Flexible and Mixture Discriminant Rules

Linear discriminant analysis is useful partly because it is simple: each class is represented by a centroid, distances are measured in a common covariance metric, and the decision boundaries are linear. Its limitations are equally

structural. A single centroid may not represent an inhomogeneous class, linear boundaries may be too rigid, and high-dimensional predictors may require smoothness or shrinkage beyond ordinary covariance pooling. Flexible, penalized, and mixture discriminant methods keep the LDA comparison geometry while changing the regression fit, the metric, or the number of class prototypes.

**Method 11.1.1.** *Flexible and Penalized Discriminant Analysis*
Optimal scoring represents LDA by regressions from $x_i$ to transformed labels.
Let $g_i \in \{1,\dots,K\}$, $1 \le L \le K-1$, label scores $\theta_\ell : \{1,\dots,K\} \to \mathbb{R}$, and fitted score maps $\eta_\ell : \mathcal{X} \to \mathbb{R}$, with centering and orthogonality constraints on the $\theta_\ell$ removing constant score directions.

(i) *Flexible discriminant analysis*: minimize $\mathcal{L}_{\mathrm{FDA}} = n^{-1} \sum_{\ell=1}^{L} \sum_{i=1}^{n} \{\theta_\ell(g_i) - \eta_\ell(x_i)\}^2 + \lambda \sum_{\ell=1}^{L} J(\eta_\ell)$ over $\theta_\ell$ and $\eta_\ell$ from a chosen smoother class: linear, spline, additive, MARS, kernel, or similar.
(ii) *Penalized discriminant analysis*: impose $\eta_\ell(x) = h(x)^\top \beta_\ell$ and roughness penalty $\beta_\ell^\top \Omega \beta_\ell$.
Equivalently classify by $D_\lambda(x,k) = \big(h(x) - \bar{h}_k\big)^\top (\widehat{\Sigma}_{W,h} + \lambda\Omega)^{-1} \big(h(x) - \bar{h}_k\big)$, with $\bar{h}_k$ and $\widehat{\Sigma}_{W,h}$ the class-$k$ centroid and pooled within-class covariance in $h$-coordinates.

For $\eta_\ell(x) = x^\top \beta_\ell$ and no roughness penalty, the fitted directions recover LDA up to scale; FDA changes the regression class to obtain nonlinear boundaries, while PDA shrinks rough or unstable $h$-directions.

**Algorithm 21:** FDA Coordinates by Optimal Scoring

**Require:** Class labels $g_i \in \{1,\dots,K\}$, indicator response matrix $Y \in \mathbb{R}^{n \times K}$, linear smoother $S_\lambda$ for the chosen regression method
**Ensure:** Nonconstant discriminant coordinate functions $\eta_1, \dots, \eta_{K-1}$
Fit the multivariate regression of $Y$ on the features: $\widehat{Y} \leftarrow S_\lambda Y$
Let $\eta^*(x)$ denote the $K$ fitted regression functions evaluated at a new input $x$
Set $D_\pi \leftarrow Y^T Y / n$
Solve $Y^T \widehat{Y} \theta = \gamma D_\pi \theta$ and normalize eigenvectors so that $\Theta^T D_\pi \Theta = I$
Discard the constant scoring direction
**for** each remaining eigenvector $\theta_\ell$ **do**
  Set $\eta_\ell(x) \leftarrow \theta_\ell^T \eta^*(x)$
**end for**

**Remark 11.1.2.** *FDA as LDA After a Regression Map*
Let $\eta = (\eta_1, \dots, \eta_L) : \mathcal{X} \to \mathbb{R}^L$ and $z = \eta(x)$. FDA applies LDA in $z$-space, so linear discriminants in $z$ pull back to decision boundaries in $x$ determined by the regression class for $\eta$.
Polynomial $\eta$ gives algebraic boundaries, quadratic $\eta$ gives quadratic boundaries, and additive, spline, or kernel $\eta$ gives the corresponding structured nonlinear boundaries.

**Method 11.1.3.** *Mixture Discriminant Analysis*
Mixture discriminant analysis replaces the class-$k$ LDA prototype by $R_k$ Gaussian subclasses while sharing one covariance $\Sigma$. Let $\pi_{kr} \ge 0$, $\sum_{r=1}^{R_k} \pi_{kr} = 1$, centers $\mu_{kr}$, class prior $\rho_k$, $f_k(x) = \sum_{r=1}^{R_k} \pi_{kr} \phi(x; \mu_{kr}, \Sigma)$, and $p_k(x) = \Pr(G = k \mid X = x) = \rho_k f_k(x) / \sum_{j=1}^{K} \rho_j f_j(x)$.
If $R_k = 1$ for all $k$, this is ordinary common-covariance Gaussian discrimination.

(i) *E-step*: for $g_i = k$, set $r_{ikr} = \pi_{kr} \phi(x_i; \mu_{kr}, \Sigma) / \sum_{s=1}^{R_k} \pi_{ks} \phi(x_i; \mu_{ks}, \Sigma)$ for $r = 1, \dots, R_k$.
(ii) *M-step*: update $\pi_{kr}$, $\mu_{kr}$, and $\Sigma$ by weighted Gaussian maximum likelihood with weights $r_{ikr}$.
(iii) *Blurred-response variant*: use subclass columns $c_{kr}$ and row $i$ equal to $(r_{ik1}, \dots, r_{ikR_k})$ on class-$k$ columns, zero elsewhere; then fit weighted DA with optional rank restrictions, FDA, or PDA.

**Remark 11.1.4.** *MDA and Prototype Rules*
MDA is a probabilistic prototype classifier with prototypes $\mu_{kr}$, weights $\pi_{kr}$, and shared metric $\Sigma^{-1}$. It assigns $x$ by posterior scores $p_k(x) \propto \rho_k \sum_{r=1}^{R_k} \pi_{kr} \phi(x; \mu_{kr}, \Sigma)$, not by a hard nearest-center rule.
Relative to classwise $k$-means, it estimates $\pi_{kr}$ and $\Sigma$; relative to an unconstrained Gaussian mixture, each component $(k, r)$ is nested inside the observed class $k$.

**Definition 11.1.5.** *Kernel Density Classification*
Let $G \in \{1, \dots, J\}$ be the class label.
Estimate $\pi_j := \mathbb{P}(G = j)$ by $\widehat{\pi}_j$ and $f_j(x) := f_{X|G=j}(x)$ by a kernel density estimate $\widehat{f}_j(x)$.

(i) *Posterior*: Define $\widehat{p}_j(x) := \widehat{\mathbb{P}}(G = j \mid X = x) := \dfrac{\widehat{\pi}_j \widehat{f}_j(x)}{\sum_{k=1}^{J} \widehat{\pi}_k \widehat{f}_k(x)}$ whenever the denominator is positive.
(ii) *Classifier*: The plug-in Bayes rule is $\widehat{g}(x) \in \operatorname{argmax}_{1 \le j \le J} \widehat{p}_j(x)$.

Full density estimation is unnecessary if only the boundary is needed.

**Remark 11.1.6.** *Computational Scale*
Let $N$ be the sample size, $p$ the predictor dimension, and $T_{\text{fit}}$ the cost of the chosen smoother.

(i) *LDA/PDA*: covariance-scale linear algebra costs about $Np^2 + p^3$ before low-rank reduction.
(ii) *FDA*: cost is $T_{\text{fit}}$; additive or adaptive basis fits can be nearly linear in $N$ for fixed model size, while dense kernel or spline smoothers can cost at least an $N \times N$ solve.
(iii) *MDA*: cost multiplies the common-covariance discriminant fit by EM iterations and depends on initialization because mixture likelihoods are nonconvex.

# References

Abbott, S. (2015). *Understanding Analysis* (2 ed.). Undergraduate Texts in Mathematics. New York, NY: Springer.

Axler, S. (2024). *Linear Algebra Done Right* (4 ed.). Undergraduate Texts in Mathematics. Cham: Springer.

Bartle, R. G. (1995). *The Elements of Integration and Lebesgue Measure*. Wiley Classics Library. New York, NY: Wiley-Interscience.

Garcia, S. R. and R. A. Horn (2017). *A Second Course in Linear Algebra*. Cambridge: Cambridge University Press.

Granziol, D., X. Wan, S. Albanie, and S. Roberts (2020). Iterative averaging in the quest for best test error. arXiv:2003.01247. https://doi.org/10.48550/arXiv.2003.01247.

Grinberg, R. (2017). *The Real Analysis Lifesaver: All the Tools You Need to Understand Proofs*. Princeton, NJ: Princeton University Press.

Hamilton, J. D. (1994). *Time Series Analysis*. Princeton University Press.

Hastie, T., R. Tibshirani, and J. Friedman (2009). *The Elements of Statistical Learning: Data Mining, Inference, and Prediction* (2 ed.). Springer Series in Statistics. New York, NY: Springer.

Mazur, M. (2015). A step by step backpropagation example. https://mattmazur.com/2015/03/17/a-step-by-step-backpropagation-example/. Accessed 2025-12-13.

Pugh, C. C. (2015). *Real Mathematical Analysis* (2 ed.). Undergraduate Texts in Mathematics. Cham: Springer.

Rice, J. A. (2007). *Mathematical Statistics and Data Analysis* (3 ed.). Belmont, CA: Duxbury.

Roman, S. (2008). *Advanced Linear Algebra* (3 ed.), Volume 135 of *Graduate Texts in Mathematics*. New York, NY: Springer.

Ross, K. A. (2013). *Elementary Analysis: The Theory of Calculus* (2 ed.). Undergraduate Texts in Mathematics. New York, NY: Springer.

Royden, H. L. and P. M. Fitzpatrick (2010). *Real Analysis* (4 ed.). Boston, MA: Pearson.

Rudin, W. (1976). *Principles of Mathematical Analysis* (3 ed.). New York: McGraw-Hill.

Strang, G. (2006). *Linear Algebra and Its Applications* (4 ed.). Belmont, CA: Thomson Brooks/Cole.

Tao, T. (2016a). *Analysis I* (3 ed.), Volume 37 of *Texts and Readings in Mathematics*. Singapore: Springer.

Tao, T. (2016b). *Analysis II* (3 ed.), Volume 38 of *Texts and Readings in Mathematics*. Singapore: Springer.