Mathematics

Arthur Li

October 6, 2025

Contents

1 Fundamentals of Higher Mathematics **1**
1.1 Single Variable Calculus 1
1.2 Linear Algebra 4
1.2.1 Vector Spaces 4
1.2.2 Linear Maps 8
1.2.3 Eigenvalues and Eigenvectors 12
1.2.4 Inner-Product Spaces 17
1.2.5 Operators on Inner-Product Spaces 23
1.2.6 Operators on Complex Vector Spaces 31

1 Fundamentals of Higher Mathematics

Generally covers four areas: single-variable calculus, linear algebra, multivariable calculus, and differential equations. As this is fundamentals, the focus is generally on computation instead of proofs.

1.1 Single Variable Calculus

Calculus is the branch of mathematics that studies change and accumulation. It has two main branches:

i. Differential Calculus: Concerned with the rate at which things change (e.g. velocity).

ii. Integral Calculus: Concerned with accumulation (e.g. area under a curve).

Calculus underpins physics, engineering, economics, biology, and more.

Definition 1.1.1. The *derivative* is defined as:

$$\lim_{\Delta x \to 0} \frac{\Delta f}{\Delta x} = \lim_{\Delta x \to 0} \frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x} = f'(x_0)$$

Definition 1.1.2. A function $f(x)$ is continuous at x_0 when

$$\lim_{x \to x_0} f(x) = f(x_0)$$

Definition 1.1.3. The *right-hand* and *left-hand* limits are as follows:

i. *Right-Hand Limit*: $\lim_{x \to x_0^+} f(x)$ means $\lim_{x \to x_0} f(x)$ for $x > x_0$

ii. *Left-Hand Limit*: $\lim_{x \to x_0^-} f(x)$ means $\lim_{x \to x_0} f(x)$ for $x < x_0$

Definition 1.1.4. A *removable discontinuity* is where

$$\lim_{x \to x_0^+} f(x) = \lim_{x \to x_0^-} f(x)$$

but not at $f(x_0)$, or if $f(x_0)$ is undefined.

Definition 1.1.5. A *jump discontinuity* is where $\lim_{x \to x_0^+} f(x)$ and $\lim_{x \to x_0^-} f(x)$ exists, but not equal.

Definition 1.1.6. A *infinite discontinuity* is where $\lim_{x \to x_0^+} f(x) \neq \lim_{x \to x_0^-} f(x) \in \{+\infty, -\infty\}$

Theorem 1.1.7. *Differentiability Implies Continuity*
If f is differentiable at x_0, then f is continuous at x_0.

Proof.

$$\lim_{x \to x_0} (f(x) - f(x_0)) = \lim_{x \to x_0} \left[\frac{f(x) - f(x_0)}{x - x_0} \right] (x - x_0) = f'(x_0) \cdot 0 = 0$$

□

Definition 1.1.8. *Product Formula*
The derivative of product of two functions.

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

Definition 1.1.9. *Quotient Formula*
The derivate of ratio of two differentiable functions. Let $v(x) \neq 0$, then

$$\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$$

Definition 1.1.10. *Chain Rule*
The derivative of composition of two functions.

$$\frac{d}{dt} f(g(t)) = f'(g(t))g'(t)$$

May be used to differentiate implicitly defined functions.

Theorem 1.1.11. *Taylor's Theorem*
Let $k \geq 1$ be an integer and let function $f : \mathbb{R} \to \mathbb{R}$ be k-times differentiable at point $a \in \mathbb{R}$. Then there exists a function $h_k : \mathbb{R} \to \mathbb{R}$ such that

$$f(x) = \sum_{i=0}^{k} \frac{f^{(i)}(a)}{i!}(x-a)^i + h_k(x)(x-a)^k$$

where $\lim_{x \to a} h_k(x) = 0$ is the Peano form of the remainder.

Theorem 1.1.12. *Mean Value Theorem*
Let $f : [a, b] \to \mathbb{R}$ be a continuous function on closed interval $[a, b]$ and differentiable on open interval (a, b) where $a < b$. Then $\exists c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Theorem 1.1.13. *First Fundamental Theorem of Calculus*
Let f be continuous real-valued function on closed interval $[a, b]$ Let F be function defined for all $x \in [a, b]$, by

$$F(x) = \int_a^x f(t)dt$$

Then F is uniformly continuous on $[a, b]$ and differentiable on open interval (a, b), and

$$F'(x) = f(x)$$

for all x in (a, b) so F is an antiderivative of f.

Remark 1.1.14. *Change of Variable*
If $f(x) = g(u(x))$, then $du = u'(x)dx$, hence

$$\int g(u)du = \int g(u(x))u'(x)dx = \int f(x)u'(x)dx$$

For definite integrals,

$$\int_{x_1}^{x_2} f(x)u'(x)dx = \int_{u_1}^{u_2} g(u)d(u)$$

where $u_1 = u(x_1), u_2 = u(x_2)$.

Theorem 1.1.15. *Second Fundamental Theorem of Calculus*
Let $f : \mathbb{R} \to \mathbb{R}$ on $[a, b]$ and F a continuous function on $[a, b]$ which is an antiderivative of f in (a, b),

$$F'(x) = f(x)$$

If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(x)dx = F(b) - F(a)$$

Remark 1.1.16. *Integration by Trigonometric Substitution*
Evaluate trigonometric integral, work backwards to original variable by using $\text{trig}(\text{trig}^{-1})$ with right-triangle.

i. $a^2 - x^2$ use $x = a \sin u$ and $dx = a \cos u \; du$
ii. $a^2 + x^2$ use $x = a \tan u$ and $dx = a \sec^2 u \; du$
iii. $x^2 - a^2$ use $x = a \sec u$ and $dx = a \sec u \tan u \; du$

Remark 1.1.17. *Systematic Integration of Rational Functions*
For a rational function P/Q (ratio of polynomials), by method of partial fractions, factor Q completely into linear and quadratic factors, and write P/Q as a sum of simpler terms.

i. If $Q(x) = (x-a)^m(x-b)^n$, then the expression is

$$\frac{A_1}{x-a} + \frac{A_2}{(x-a)^2} + \cdots + \frac{A_m}{(x-a)^m} + \frac{B_1}{x-b} + \frac{B_2}{(x-b)^2} + \cdots + \frac{B_n}{(x-b)^n}$$

ii. If there are quadratic factor such as $(Ax^2+Bx+C)^p$, then there are terms

$$\frac{a_1x+b_1}{Ax^2+Bx+C}+\frac{a_2x+b_2}{(Ax^2+Bx+C)^2}+\cdots+\frac{a_px+b_p}{(Ax^2+Bx+C)^p}$$

for each factor. To integrate, complete the square and make trigonometric substitution.

Remark 1.1.18. *Integration by Parts*

$$\int_b^a uv'\,dx = uv|_a^b - \int_a^b u'v\,dx$$

This is used when $u'v$ is simpler than uv' (often the case when u' is simpler than u).

Remark 1.1.19. *Arc-Length*
Given $ds=\sqrt{dx^2+dy^2}$, depending on whether to integrate with respect to x,t,y, this is

$$ds=\sqrt{1+\left(\frac{dy}{dx}\right)^2}\,dx;\quad ds=\sqrt{\left(\frac{dx}{dt}\right)^2+\left(\frac{dy}{dt}\right)^2}\,dt;\quad ds=\sqrt{\left(\frac{dx}{dy}\right)^2+1}\,dy$$

Remark 1.1.20. *Surface Area for Surface of Revolution*

i. Around x-axis: (requires formula for $y=y(x)$)

$$2\pi y\,ds = 2\pi y\sqrt{1+\left(\frac{dy}{dx}\right)^2}\,dx$$

ii. Around y-axis: (requires formula for $x=x(y)$)

$$2\pi x\,ds = 2\pi x\sqrt{\left(\frac{dx}{dy}\right)^2+1}\,dy$$

Remark 1.1.21. *Polar Coordinates*
Given $x=r\cos\theta$, $y=r\sin\theta$ (or more rarely $r=\sqrt{x^2+y^2}\theta$, $\theta=\tan^{-1}(y/x)$)

i. Find the polar equation for a curve from its equation in (x,y) variables by substitution.
ii. Sketch curves given in polar coordinates and find range of θ for integration.
iii. Integration area formula is then

$$\int_{\theta_1}^{\theta_2}\frac{1}{2}r^2\,d\theta$$

Theorem 1.1.22. *L'Hopital Rule*
For functions f and g defined on open interval I and differentiable on $I\backslash\{c\}$ for a accumulation point c of I, if

$$\lim_{x\to c}f(x)=\lim_{x\to c}g(x)=0\quad or\quad \pm\infty$$

and $g'(x)\neq 0$ for all $x\in I\backslash\{c\}$, and $\lim_{x\to c}\frac{f'(x)}{g'(x)}$ exists, then

$$\lim_{x\to c}\frac{f(x)}{g(x)}=\lim_{x\to c}\frac{f'(x)}{g'(x)}$$

1.2 Linear Algebra

Linear algebra is the branch of mathematics that studies vectors, matrices, and linear transformations. It provides the foundation for solving systems of equations and analysing multidimensional data. Its applications span across machine learning, computer graphics, physics, finance, and engineering.
References from Garcia and Horn (2017) and Axler (2024).

1.2.1 Vector Spaces

Definition 1.2.1. *Vector Space*
Let $\mathbb{F} = \mathbb{R}$ or $\mathbb{C}$. Then $\mathcal{V}$ is a vector space over a field $\mathbb{F}$ if the following are satisfied:

i. Identities: $\exists\ \mathbf{0} \in \mathcal{V}$ such that $\mathbf{0} + \mathbf{u} = \mathbf{u}\ \forall\ \mathbf{u} \in \mathcal{V}$, and $\exists\ \mathbf{1} \in \mathcal{V}$ such that $\mathbf{1u} = \mathbf{u}\ \forall\ \mathbf{u} \in \mathcal{V}$
ii. Commutativity: $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}\ \forall\ \mathbf{u}, \mathbf{v} \in \mathcal{V}$
iii. Associativity: $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$ and $(ab)\mathbf{u} = a(b\mathbf{u})\ \forall\ a, b \in \mathbb{F}$ and $\forall\ \mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$
iv. Distributive: $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$ and $(a + b)\mathbf{u} = a\mathbf{u} + b\mathbf{u}\ \forall\ a, b \in \mathbb{F}$ and $\forall\ \mathbf{u}, \mathbf{v} \in \mathcal{V}$
v. Additive Inverse: $\forall\ \mathbf{u} \in \mathcal{V}\ \exists$ unique vector $-\mathbf{u} \in \mathcal{V}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
vi. Multiplicative Inverse: $\forall\ \mathbf{u} \in \mathcal{V}$ with $\mathbf{u} \neq \mathbf{0}\ \exists$ unique $\mathbf{w} \in \mathcal{V}$ such that $\mathbf{uw} = 1$

Remark 1.2.2. *Properties of Vector Spaces*
Let $\mathcal{V}$ be a vector space over a field $\mathbb{F}$.

i. A vector space has a unique additive identity
ii. Every element in a vector space has a unique additive inverse
iii. $\mathbf{0v} = \mathbf{v}$ for every $\mathbf{v} \in \mathcal{V}$
iv. $a\mathbf{0} = \mathbf{0}$ for every $a \in \mathbb{F}$
v. $(-1)\mathbf{v} = -\mathbf{v}$ for every $\mathbf{v} \in \mathcal{V}$

Definition 1.2.3. *Subspace*
A subspace of an $\mathbb{F}$-vector space $\mathcal{V}$ is a subset $\mathcal{U}$ of $\mathcal{V}$ that is an $\mathbb{F}$-vector space with the same vector addition and scalar multiplication operations as in $\mathcal{V}$. To check $\mathcal{U}$ satisfies the following:

i. Additive Identity: $\mathbf{0} \in \mathcal{U}$
ii. Closure Under Addition: $\mathbf{u}, \mathbf{v} \in \mathcal{U}$ implies $\mathbf{u} + \mathbf{v} \in \mathcal{U}$
iii. Closed Under Scalar Multiplication: $a \in \mathbb{F}$ and $\mathbf{u} \in \mathcal{U}$ implies $a\mathbf{u} \in \mathcal{U}$.

Definition 1.2.4. *Sum of Subspaces*
Suppose $\mathcal{U}_1, \ldots, \mathcal{U}_m$ are subspaces of $\mathcal{V}$. The sum of $\mathcal{U}_1, \ldots, \mathcal{U}_m$ as denoted by $\mathcal{U}_1 + \cdots + \mathcal{U}_m$ is defined to be set of all possible sums of elements of $\mathcal{U}_1, \ldots, \mathcal{U}_m$:

$$\mathcal{U}_1 + \cdots + \mathcal{U}_m \coloneqq \{\mathbf{u}_1 + \cdots + \mathbf{u}_m : \mathbf{u}_1 \in \mathcal{U}_1, \ldots, \mathbf{u}_m \in \mathcal{U}_m\}$$

Definition 1.2.5. *Direct Sum of Subspaces*
If $\mathcal{U}_1, \ldots, \mathcal{U}_m$ are subspaces of $\mathcal{V}$ such that $\mathcal{V} = \mathcal{U}_1 + \cdots + \mathcal{U}_m$, then every element of $\mathcal{V}$ may be written as

$$\mathbf{u}_1 + \cdots + \mathbf{u}_m$$

where each $\mathbf{u}_j \in \mathcal{U}_j$. Then $\mathcal{V}$ is the direct sum of subspaces $\mathcal{U}_1, \ldots, \mathcal{U}_m$, written as $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$

Definition 1.2.6. *Linear Combination*
Let $\mathcal{U}$ be non-empty subset of $\mathbb{F}$-vector space $\mathcal{V}$. A linear combination of elements of $\mathcal{U}$ is expressed as

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_r\mathbf{v}_r$$

where r is a positive integer, and $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_r \in \mathcal{U}$, $c_1, c_2, \ldots, c_r \in \mathbb{F}$.
A linear combination is *trivial* if $c_1 = c_2 = \cdots = c_r = 0$.

Definition 1.2.7. *Span*
Let $\mathcal{U}$ be subset of $\mathbb{F}$-vector space $\mathcal{V}$. If $\mathcal{U} \neq \emptyset$, then $\text{span}(\mathcal{U})$ is the set of linear combinations of elements of $\mathcal{U}$. Define $\text{span}(\emptyset) = \{\mathbf{0}\}$. If $\beta = \mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_r$ is a list of $r \geq 1$ vectors in $\mathcal{V}$, define $\text{span}(\beta) = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_r\}$.

Lemma 1.2.8. *Linear Dependence Lemma*
If $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly dependent in $\mathcal{V}$ and $\mathbf{v}_1 \neq 0$, then $\exists\ j \in \{2, \ldots, m\}$ such that the following hold:

i. $\mathbf{v}_j \in \mathit{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$

ii. if the jth term is removed from $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$*, the span of the remaining list equals* $\mathit{span}(\mathbf{v}_1, \ldots, \mathbf{v}_m)$.

Proof. i. Suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly dependent in $\mathcal{V}$, and $\mathbf{v}_1 \neq 0$. Then $\exists\ a_1, \ldots, a_m \in \mathbb{F}$, not all 0, s.t.

$$a_1\mathbf{v}_1 + \cdots + a_m\mathbf{v}_m = 0$$

Not all of $a_2, \ldots, a_m$ can be 0 as $\mathbf{v}_1 \neq 0$. Let j be largest element of $2, \ldots, m$ s.t. $a_j \neq 0$. Then

$$\mathbf{v}_j = -\frac{a_1}{a_j}\mathbf{v}_1 - \cdots - \frac{a_{j-1}}{a_j}\mathbf{v}_{j-1}$$

ii. Suppose $\boldsymbol{u} \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_m)$. Then $\exists\ c_1, \ldots, c_m \in \mathbb{F}$ such that

$$\mathbf{u} = c_1\mathbf{v}_1 + \cdots + c_m\mathbf{v}_m$$

Replace $\mathbf{v}_j$ with equation from (i), hence (ii) holds.

□

Theorem 1.2.9. *In a finite-dimensional vector space, length of every linearly independent list of vectors is less than or equal to length of every spanning list of vectors.*

Proof. Suppose $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ is linearly independent in $\mathcal{V}$, and $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$ spans $\mathcal{V}$. Need to prove $m \leq n$.

i. List $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$ spans $\mathcal{V}$, hence adjoining any vector to it produces linearly dependent list. By Lemma 1.2.8, can remove one of $\mathbf{w}$ such that the list containing $\mathbf{u}_1$ and remaining $\mathbf{w}$'s spans $\mathcal{V}$.

ii. List of length n from previous step $j-1$ spans $\mathcal{V}$, hence adjoining any vector to it produces a linearly dependent list. The list of length $(n+1)$ obtained by adjoining $\mathbf{u}_j$ to the list, placing it just after $\mathbf{u}_1, \ldots, \mathbf{u}_{j-1}$ is linearly dependent. By Lemma 1.2.8, one of the vectors in this list is span of previous ones, because $(\mathbf{u}_1, \ldots, \mathbf{u}_j)$ is linearly independent, hence this vector must be one of $\mathbf{w}$ not one of $\mathbf{u}$. Hence can remove $\mathbf{w}$ from the list so that the new list consists of $\mathbf{u}_1, \ldots, \mathbf{u}_j$ and remaining $\mathbf{w}$ spans $\mathcal{V}$.

iii. At step m, all $\mathbf{u}$ has been added, and process tops. If at any step $\mathbf{u}$ is added and no more $\mathbf{w}$ could be removed, there is a contradiction. hence there must be at least as many $\mathbf{w}$ as $\mathbf{u}$.

□

Proposition 1.2.10. *Every subspace of finite-dimensional vector space is finite dimensional.*

Proof. Suppose $\mathcal{V}$ is finite dimensional, and $\mathcal{U}$ is subspace of $\mathcal{V}$.

i. If $\mathcal{U} = \{0\}$, then $\mathcal{U}$ is finite dimensional. If $\mathcal{U} \neq \{0\}$, choose nonzero vector $\mathbf{v}_1 \in \mathcal{U}$.

ii. If $\mathcal{U} = \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$, then U is finite dimensional. If $\mathcal{U} \neq \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$, choose vector $\mathbf{v}_j \in \mathcal{U}$ such that $\mathbf{v}_j \neq \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$

After each step, a linearly independent list is constructed by Lemma 1.2.8. This linearly independent list cannot be longer than any spanning list of $\mathcal{V}$, hence process must terminate, which means $\mathcal{U}$ is finite dimensional. □

Definition 1.2.11. A *basis* of $\mathcal{V}$ is a list of vectors in $\mathcal{V}$ that is linearly independent and spans $\mathcal{V}$.

Proposition 1.2.12. *A list* $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ *of vectors in* $\mathcal{V}$ *is a basis of* $\mathcal{V}$ *if and only if every* $\mathbf{v} \in \mathcal{V}$ *can be written uniquely in form* $\mathbf{v} = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$*, where* $a_1, \ldots, a_n \in \mathbb{F}$.

Proof. First suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is basis of $\mathcal{V}$. Let $\mathbf{v} \in \mathcal{V}$. As $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ spans $\mathcal{V}$, $\exists\ a_1, \ldots, a_n \in \mathbb{F}$ such that proposition holds. To show the representation is unique, suppose $b_1, \ldots, b_n$ are scalars, hence

$$\mathbf{v} = b_1\mathbf{v}_1 + \cdots + b_n\mathbf{v}_n$$

Subtracting this from proposition equation,

$$0 = (a_1 - b_1)\mathbf{v}_1 + \cdots + (a_n - b_n)\mathbf{v}_n$$

This implies each $a_j - b_j = 0$ as $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is linearly independent, hence $a_1 = b_1, \ldots, a_n = b_n$.
For other direction, suppose every $\mathbf{v} \in \mathcal{V}$ can be written uniquely in proposition form. To show $(\mathbf{v}_1, \ldots \mathbf{v}_n)$ is linearly independent, suppose $a_1, \ldots a_n \in \mathbb{F}$ are such that

$$0 = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$$

The uniqueness of representation from proposition with $\mathbf{v} = 0$ implies $a_1 = \cdots = a_n = 0$. Hence $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is linearly independent and hence is a basis of $\mathcal{V}$. □

Theorem 1.2.13. *Every spanning list in a vector space can be reduced to a basis of vector space.*

Proof. Suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ spans $\mathcal{V}$. To remove some vectors from $(\mathbf{v}_1, \ldots \mathbf{v}_n)$ so that remaining vectors form a basis of $\mathcal{V}$. Start with $\mathcal{B} = (\mathbf{v}_1, \ldots, \mathbf{v}_n)$.

i. If $\mathbf{v}_1 = 0$, delete $\mathbf{v}_1$ from B, If $\mathbf{v}_1 \neq 0$, leave $\mathcal{B}$ unchanged.

ii. If $\mathbf{v}_j$ is in span$(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$, delete $\mathbf{v}_j$ from $\mathcal{B}$. If $\mathbf{v}_j$ is not in span$(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$, leave $\mathcal{B}$ unchanged.

Stop process after step n, getting a list $\mathcal{B}$ that spans $\mathcal{V}$, and discarded only vectors that were already in span of previous vectors. Hence $\mathcal{B}$ is linearly independent by Lemma 1.2.8. Hence $\mathcal{B}$ is a basis of $\mathcal{V}$. □

Corollary 1.2.14. *Every finite-dimensional vector space has a basis.*

Proof. By definition, a finite-dimensional vector space has a spanning list. By Theorem 1.2.13, any spanning list can be reduced to a basis. □

Theorem 1.2.15. *Every linearly independent list of vectors in a finite-dimensional vector space can be extended to a basis of the vector space.*

Proof. Suppose $\mathcal{V}$ is finite dimensional and $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly independent in $\mathcal{V}$. To extend $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ to a basis of $\mathcal{V}$. Let $(w_1, \ldots, w_n)$ be any list of vectors in $\mathcal{V}$ that spans $\mathcal{V}$.

i. If $\mathbf{w}_1 \in$ span$(\mathbf{v}_1, \ldots, \mathbf{v}_m)$, let $\mathcal{B} = (\mathbf{v}_1, \ldots, \mathbf{v}_m)$. If $\mathbf{w}_1$ is not in span$(\mathbf{v}_1, \ldots, \mathbf{v}_m)$, let $\mathcal{B} = (\mathbf{v}_1, \ldots, \mathbf{v}_m, \mathbf{w}_1)$.

ii. If $\mathbf{w}_j \in$ span(B), leave $\mathcal{B}$ unchanged. If $\mathbf{w}_j$ not in span(B), extend $\mathcal{B}$ by adjoining $\mathbf{w}_j$ to it.

After each step, $\mathcal{B}$ is still linearly independent as otherwise Lemma 1.2.8 would give a contradiction. After step n, the span of $\mathcal{B}$ includes all the $\mathbf{w}$. Hence the $\mathcal{B}$ obtained after step n spans $\mathcal{V}$ and hence is a basis of $\mathcal{V}$. □

This theorem can be applied to show that every subspace of a finite-dimensional vector space can be paired with another subspace to form a direct sum of the whole space.

Proposition 1.2.16. *Suppose $\mathcal{V}$ is finite dimensional and $\mathcal{U}$ is a subspace of $\mathcal{V}$. Then there is a subspace $\mathcal{W}$ of $\mathcal{V}$ such that $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.*

Proof. As $\mathcal{V}$ is finite dimensional, so is $\mathcal{U}$, hence $\exists$ basis$(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ of $\mathcal{U}$. As $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ is a linearly independent list of vectors of $\mathcal{V}$, it can be extended to a basis $(\mathbf{u}_1, \ldots \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_m)$ of $\mathcal{V}$. Let $\mathcal{W} =$ span$(\mathbf{w}_1, \ldots, \mathbf{w}_n)$. To prove $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, need to show that $\mathcal{V} = \mathcal{U} + \mathcal{W}$ and $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$.
Suppose $\mathbf{v} \in \mathcal{V}$. As $\mathbf{u}_1, \ldots, \mathbf{u}_m$ spans $\mathcal{V}$, there exists scalars $a_1, \ldots, a_m, b_1, \ldots, b_n \in \mathbb{F}$ such that

$$\mathbf{v} = a_1\mathbf{u}_1 + \cdots + a_m\mathbf{u}_m + b_1\mathbf{w}_1 + \cdots + b_n\mathbf{w}_n = \mathbf{u} + \mathbf{w}$$

where $\mathbf{u} \in \mathcal{U}$, $\mathbf{w} \in \mathcal{W}$ are defined as above. Hence $\mathbf{v} \in \mathcal{U} + \mathcal{W}$, thus $\mathcal{V} = \mathcal{U} + \mathcal{W}$.
To show $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$, suppose $\mathbf{v} \in \mathcal{U} \cap \mathcal{W}$. Then $\exists$ scalars $a_1, \ldots, a_m, b_1, \ldots, b_n \in \mathbb{F}$. such that

$$\mathbf{v} = a_1\mathbf{u}_1 + \cdots + a_m\mathbf{u}_m = b_1\mathbf{w}_1 + \cdots + b_n\mathbf{w}_n$$

Hence

$$a_1\mathbf{u}_1 + \cdots + a_m\mathbf{u}_m - b_1\mathbf{w}_1 - \cdots - b_n\mathbf{w}_n = \mathbf{0}$$

As $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_n)$ is linearly independent, this implies that $a_1 = \cdots = a_m = b_1 = \cdots = b_n = 0$, hence $\mathbf{v} = \mathbf{0}$, completing proof that $\mathcal{U} \cap \mathcal{W} = \{\mathbf{0}\}$. □

Theorem 1.2.17. *Any two bases of a finite-dimensional vector space have the same length.*

Proof. Suppose $\mathcal{V}$ is finite dimensional, and $\mathcal{B}_1, \mathcal{B}_2$ be any two bases of $\mathcal{V}$. Then $\mathcal{B}_1$ is linearly independent in $\mathcal{V}$, and $\mathcal{B}_2$ spans $\mathcal{V}$. Hence length of $\mathcal{B}_1$ is at most the length of $\mathcal{B}_2$. Interchanging roles of $\mathcal{B}_1$ and $\mathcal{B}_2$, the length of $\mathcal{B}_2$ is at most the length of $\mathcal{B}_1$. Hence length of $\mathcal{B}_1$ must equal length of $\mathcal{B}_2$ as desired. □

Definition 1.2.18. The *dimension* of a finite-dimensional vector space is defined to be the length of any basis of the vector space, denoted as dim$(\mathcal{V})$.

Proposition 1.2.19. *If $\mathcal{V}$ is finite dimensional and $\mathcal{U}$ is a subspace of $\mathcal{V}$, then dim$(\mathcal{U}) \leq$ dim$(\mathcal{V})$.*

Proof. Suppose $\mathcal{V}$ is finite dimensional and $\mathcal{U}$ is a subspace of $\mathcal{V}$. Any basis of $\mathcal{U}$ is a linearly independent list of vectors in $\mathcal{V}$, thus can be extended to a basis of $\mathcal{V}$. Hence length of a basis of $\mathcal{U} \leq$ length of a basis of $\mathcal{V}$. □

Proposition 1.2.20. *If $\mathcal{V}$ is finite dimensional, then every spanning list of vectors in $\mathcal{V}$ with length $dim(\mathcal{V})$ is a basis of $\mathcal{V}$.*

Proof. Suppose $\dim(\mathcal{V}) = n$ and $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ spans $\mathcal{V}$. The list $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ can be reduced to a basis of $\mathcal{V}$. However, every basis of $\mathcal{V}$ has length n, hence reduction must be trivial, i.e. no elements are deleted from $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$. Hence, $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$ as desired. □

Proposition 1.2.21. *If $\mathcal{V}$ is finite dimensional, then every linearly independent list of vectors in $\mathcal{V}$ with length $dim(\mathcal{V})$ is a basis of $\mathcal{V}$.*

Proof. Suppose $\dim(\mathcal{V}) = n$ and $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is linearly independent in $\mathcal{V}$. The list $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ can be extended to a basis of $\mathcal{V}$. However, every basis of $\mathcal{V}$ has length n, hence the extension must be trivial, i.e., no new elements are adjoined to $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$. Hence, $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$ as desired. □

The next theorem gives formula for dimension of sum of two subspaces of a finite-dimensional vector space.

Theorem 1.2.22. *If $\mathcal{U}_1$ and $\mathcal{U}_2$ are subspaces of a finite-dimensional vector space, then*

$$dim(\mathcal{U}_1 + \mathcal{U}_2) = dim(\mathcal{U}_1) + dim(\mathcal{U}_2) - dim(\mathcal{U}_1 \cap \mathcal{U}_2)$$

Proof. Let $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ be basis of $\mathcal{U}_1 \cap \mathcal{U}_2$, hence $\dim(\mathcal{U}_1 \cap \mathcal{U}_2) = m$.
As $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ is a basis of $\mathcal{U}_1 \cap \mathcal{U}_2$, it is linearly independent in $\mathcal{U}_1$, hence can be extended to a basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_j)$ of $\mathcal{U}_1$. Hence $\dim(\mathcal{U}_1) = m+j$. Extend $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ to a basis of $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_k)$ of $\mathcal{U}_2$, hence $\dim(\mathcal{U}_2) = m + k$.
Note that $\text{span}(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_j, \mathbf{w}_1, \ldots, \mathbf{w}_k)$ contains $\mathcal{U}_1$ and $\mathcal{U}_2$, hence contains $\mathcal{U}_1 + \mathcal{U}_2$. To show that this list is a basis of $\mathcal{U}_1 + \mathcal{U}_2$, need to show that it is linearly independent. Suppose

$$a_1\mathbf{u}_1 + \cdots + a_m\mathbf{u}_m + b_1\mathbf{v}_1 + \cdots + b_j\mathbf{v}_j + c_1\mathbf{w}_1 + \cdots + c_k\mathbf{w}_k = \mathbf{0}$$

where $a_i, b_i, c_i \in \mathbb{F}$. Rewriting the equation,

$$c_1\mathbf{w}_1 + \cdots + c_k\mathbf{w}_k = -a_1\mathbf{u}_1 - \cdots - a_m\mathbf{u}_m - b_1\mathbf{v}_1 - \cdots - b_j\mathbf{v}_j$$

which shows $c_1\mathbf{w}_1 + \cdots + c_k\mathbf{w}_k \in \mathcal{U}_1$. All $\mathbf{w}$ are in $\mathcal{U}_2$, hence this implies $c_1\mathbf{w}_1 + \cdots + c_k\mathbf{w}_k \in \mathcal{U}_1 \cap \mathcal{U}_2$. As $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ is a basis of $\mathcal{U}_1 \cap \mathcal{U}_2$, write this as

$$c_1\mathbf{w}_1 + \cdots + c_k\mathbf{w}_k = d_1\mathbf{u}_1 + \cdots + d_m\mathbf{u}_m$$

for some $d_i \in \mathbb{F}$. As $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_k)$ is linearly independent, the last equation implies all c_i and d_i equal 0. Hence the original equation becomes

$$a_1\mathbf{u}_1 + \cdots + a_m\mathbf{u}_m + b_1\mathbf{v}_1 + \cdots + b_j\mathbf{v}_j = \mathbf{0}$$

This implies all a_i and b_i are 0 as the list $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_k)$ is linearly independent.
Hence $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_j, \mathbf{w}_1, \ldots, \mathbf{w}_k)$ is a basis of $\mathcal{U}_1 + \mathcal{U}_2$, and thus

$$\begin{aligned}\dim(\mathcal{U}_1 + \mathcal{U}_2) &= m + j + k \\ &= (m + j) + (m + k) - m \\ &= \dim(\mathcal{U}_1) + \dim(\mathcal{U}_2) - \dim(\mathcal{U}_1 \cap \mathcal{U}_2)\end{aligned}$$

□

Note that dimension meshes well with direct sums, which is analogous to disjoint union.

Proposition 1.2.23. *Suppose $\mathcal{V}$ is finite dimensional and $\mathcal{U}_1, \ldots, \mathcal{U}_m$ are subspaces of $\mathcal{V}$ such that*

$$\begin{aligned}\mathcal{V} &= \mathcal{U}_1 + \cdots + \mathcal{U}_m \\ dim(\mathcal{V}) &= dim(\mathcal{U}_1) + \cdots + dim(\mathcal{U}_m)\end{aligned}$$

Then $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$.

Proof. Choose a basis for each $\mathcal{U}_j$. Put these bases together in a list which spans $\mathcal{V}$ and has length $\dim(\mathcal{V})$. Hence this list is a basis of $\mathcal{V}$, and it is linearly independent. Suppose $\mathbf{u}_1 \in \mathcal{U}_1, \ldots, \mathbf{u}_m \in \mathcal{U}_m$ such that

$$\mathbf{0} = \mathbf{u}_1 + \cdots + \mathbf{u}_m$$

Write each $\mathbf{u}_j$ as a linear combination of basis vectors of $\mathcal{U}_j$. Substitute these linear combinations into equation above, hence $\mathbf{0}$ is written as a linear combination of basis of $\mathcal{V}$ constructed above. Hence all scalars used in this linear combination must be 0. Thus each $\mathbf{u}_j = \mathbf{0}$, which proves $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$. □

1.2.2 Linear Maps

Definition 1.2.24. A *Linear Map* from $\mathcal{V}$ to $\mathcal{W}$ is a function $T : \mathcal{V} \to \mathcal{W}$ with following properties:

i. *Additivity*: $T(\mathbf{u} + \mathbf{v}) = T\mathbf{u} + T\mathbf{v}$ for all $\mathbf{u}, \mathbf{v} \in \mathcal{V}$

ii. *Homogeneity*: $T(a\mathbf{v}) = a(T\mathbf{v})$ for all $a \in \mathbb{F}$ and all $\mathbf{v} \in \mathcal{V}$.

The set of all linear maps from $\mathcal{V}$ to $\mathcal{W}$ is denoted $\mathcal{L}(\mathcal{V}, \mathcal{W})$.
The linear map is *injective* if whenever $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ and $T\mathbf{u} = T\mathbf{v}$, that $\mathbf{u} = \mathbf{v}$.
The linear map is *surjective* if its range equals $\mathcal{W}$.

Example 1.2.25. Examples of linear maps include

i. Zero Map: $0 \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ is defined by $0\mathbf{v} = 0$.

ii. Identity Map: $I \in \mathcal{L}(\mathcal{V}, \mathcal{V})$ is defined by $I\mathbf{v} = \mathbf{v}$

iii. Differentiation: $T \in \mathcal{L}(\mathcal{P}(\mathbf{R}), \mathcal{P}(\mathbf{R}))$ is defined by $T\mathbf{p} = \mathbf{p}'$

iv. Integration: $T \in \mathcal{L}(\mathcal{P}(\mathbf{R}), \mathbf{R})$ is defined by $T\mathbf{p} = \int_0^1 \mathbf{p}(\mathbf{x})\, d\mathbf{x}$

v. Backward Shift: Let $\mathbb{F}^\infty$ be vector space of all sequences of elements of $\mathbb{F}$. Let $T \in \mathcal{L}(\mathbb{F}^\infty, \mathbb{F}^\infty)$. The backward shift map is then $T(x_1, x_2, x_3, \ldots) = (x_2, x_3, \ldots)$

vi. From $\mathbb{F}^n$ to $\mathbb{F}^m$: $T \in \mathcal{L}(\mathbb{F}^n, \mathbb{F}^m)$ defined by $T(x_1, \ldots, x_n) = (a_{1,1}x_1 + \cdots + a_{1,n}x_n, \ldots, a_{m,1}x_1 + \cdots + a_{m,n}x_n)$, where $a_{j,k} \in \mathbb{F}$, $j = 1, \ldots, m$ and $k = 1, \ldots, n$.

Remark 1.2.26. *Properties of Product of Linear Maps*
Let $T \in \mathcal{L}(\mathcal{U}, \mathcal{V})$, and $S \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. The *product* of S and T is ST.

i. Associativity: $(T_1T_2)T_3 = T_1(T_2T_3)$

ii. Identity: $TI = T$ (identity map on $\mathcal{V}$) and $IT = T$ (identity map on $\mathcal{W}$)

iii. Distributive: $(S_1 + S_2)T = S_1T + S_2T$ and $S(T_1 + T_2) = ST_1 + ST_2$ whenever $T, T_1, T_2 \in \mathcal{L}(\mathcal{U}, \mathcal{V})$ and $S, S_1, S_2 \in \mathcal{L}(\mathcal{V}, \mathcal{W})$

Multiplication of linear maps is not commutative.

Definition 1.2.27. For $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, the *null space* of T, denoted $\text{null}(T)$, is the subset of $\mathcal{V}$ consisting of those vectors that maps T to 0, $\text{null}(T) = \{\mathbf{v} \in \mathcal{V} : T\mathbf{v} = \mathbf{0}\}$.

Proposition 1.2.28. *If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then null$(T)$ is a subspace of $\mathcal{V}$.*

Proof. Suppose $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. By additivity,

$$T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0})$$

which implies $T(\mathbf{0}) = \mathbf{0}$. Hence $\mathbf{0} \in \text{null}(T)$.
If $\mathbf{u}, \mathbf{v} \in \text{null}(T)$, then

$$T(\mathbf{u} + \mathbf{v}) = T\mathbf{u} + T\mathbf{v} = \mathbf{0} + \mathbf{0} = \mathbf{0}$$

Hence $\mathbf{u} + \mathbf{v} \in \text{null}(T)$, and $\text{null}(T)$ is closed under addition.
If $\mathbf{u} \in \text{null}(T)$ and $a \in \mathbb{F}$, then

$$T(a\mathbf{u}) = aT\mathbf{u} = a\mathbf{0} = \mathbf{0}$$

Hence $a\mathbf{u} \in \text{null}(T)$, and $\text{null}(T)$ is closed under scalar multiplication.
As $\text{null}(T)$ contains $\mathbf{0}$ and is closed under addition and scalar multiplication, $\text{null}(T)$ is subspace of $\mathcal{V}$. □

Proposition 1.2.29. *Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. Then T is injective if and only if null$(T) = \{\mathbf{0}\}$.*

Proof. ($\Rightarrow$) Suppose T is injective. By Proposition 1.2.28, $\{\mathbf{0}\} \subset \text{null}(T)$. Suppose $\mathbf{v} \in \text{null}(T)$. Then $T(\mathbf{v}) = \mathbf{0} = T(\mathbf{0})$. As T is injective, equation above implies $\mathbf{v} = \mathbf{0}$. Thus $\text{null}(T) = \{\mathbf{0}\}$.
($\Leftarrow$) Suppose $\text{null}(T) = \{\mathbf{0}\}$, $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ and $T\mathbf{u} = T\mathbf{v}$. Then $\mathbf{0} = T\mathbf{u} - T\mathbf{v} = T(\mathbf{u} - \mathbf{v})$. Hence $\mathbf{u} - \mathbf{v}$ is in $\text{null}(T)$, which equals $\{\mathbf{0}\}$. Thus $\mathbf{u} - \mathbf{v} = \mathbf{0}$, which implies $\mathbf{u} = \mathbf{v}$. Hence T is injective. □

Definition 1.2.30. For $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, the *range* of T, denoted $\text{range}(T)$, is subset of $\mathcal{W}$ consisting of vectors that are of from $T\mathbf{v}$ for some $\mathbf{v} \in \mathcal{V}$, where $\text{range}(T) = \{T\mathbf{v} : \mathbf{v} \in \mathcal{V}\}$.

Proposition 1.2.31. *If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then range$(T)$ is a subspace of $\mathcal{W}$.*

Proof. Suppose $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then $T(\mathbf{0}) = \mathbf{0}$ by Proposition 1.2.28, implying $\mathbf{0} \in \text{range}(T)$.
If $\mathbf{w}_1, \mathbf{w}_2 \in \text{range}(T)$, then $\exists\ \mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}$ such that $T\mathbf{v}_1 = \mathbf{w}_1$ and $T\mathbf{v}_2 = \mathbf{w}_2$. Hence

$$T(\mathbf{v}_1 + \mathbf{v}_2) = T\mathbf{v}_1 + T\mathbf{v}_2 = \mathbf{w}_1 + \mathbf{w}_2$$

thus $\mathbf{w}_1 + \mathbf{w}_2 \in \text{range}(T)$. Hence $\text{range}(T)$ is closed under addition.
If $\mathbf{w} \in \text{range}(T)$ and $a \in \mathbb{F}$, then $\exists\ \mathbf{v} \in \mathcal{V}$ such that $T\mathbf{v} = \mathbf{w}$. Hence $T(a\mathbf{v}) = aT\mathbf{v} = a\mathbf{w}$, and $a\mathbf{w} \in \text{range}(T)$. Thus $\text{range}(T)$ is closed under scalar multiplication.
As $\text{range}(T)$ contains $\mathbf{0}$ and is closed under addition and scalar multiplication, $\text{range}(T)$ is a subspace of $\mathcal{W}$. □

The next theorem states that the dimension of the null space plus dimension of range of linear map on finite-dimensional vector space equals dimension of the domain.

Theorem 1.2.32. *If $\mathcal{V}$ is finite-dimensional and $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then range$(T)$ is a finite dimensional subspace of $\mathcal{W}$, and dim$(\mathcal{V})$ = dim(null(T)) + dim(range(T)).*

Proof. Suppose $\mathcal{V}$ is a finite-dimensional vector space and $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.
Let $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ be a basis of $\text{null}(T)$, thus $\dim(\text{null}(T)) = m$. The linearly independent list $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ can be extended to a basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_n)$ of $\mathcal{V}$. Hence $\dim(\mathcal{V}) = m + n$. Need to show $\text{range}(T)$ is finite dimensional and $\dim(\text{range}(T)) = n$, by proving that $(T\mathbf{w}_1, \ldots, T\mathbf{w}_n)$ is a basis of $\text{range}(T)$.
Let $\mathbf{v} \in \mathcal{V}$. As $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_n)$ spans $\mathcal{V}$, then

$$\mathbf{v} = a_1\mathbf{u}_1 + \cdots + a_m\mathbf{u}_m + b_1\mathbf{w}_1 + \cdots + b_n\mathbf{w}_n$$

where $a, b \in \mathbb{F}$. Applying T to both sides of equation,

$$T\mathbf{v} = b_1T\mathbf{w}_1 + \cdots + b_nT\mathbf{w}_n$$

Note $T\mathbf{u}_j$ disappeared as each $\mathbf{u}_j \in \text{null}(T)$. Last equation implies $(T\mathbf{w}_1, \ldots T\mathbf{w}_n)$ spans $\text{range}(T)$, hence that $\text{range}(T)$ is finite dimensional.
To show $(T\mathbf{w}_1, \ldots T\mathbf{w}_n)$ is linearly independent, suppose $c_1, \ldots, c_n \in \mathbb{F}$, and

$$\begin{aligned} c_1T\mathbf{w}_1 + \cdots + c_nT\mathbf{w}_n &= \mathbf{0} \\ T(c_1\mathbf{w}_1 + \cdots + c_n\mathbf{w}_n) &= \mathbf{0} \\ c_1\mathbf{w}_1 + \cdots + c_n\mathbf{w}_n &\in \text{null}(T) \end{aligned}$$

As $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ spans $\text{null}(T)$, then

$$c_1\mathbf{w}_1 + \cdots + c_n\mathbf{w}_n = d_1\mathbf{u}_1 + \cdots + d_m\mathbf{u}_m$$

where $d_i \in \mathbb{F}$. This implies all c and d are 0, as $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{w}_1, \ldots, \mathbf{w}_n)$ are linearly independent.
Hence $(T\mathbf{w}_1, \ldots, T\mathbf{w}_n)$ is linearly independent and is a basis for $\text{range}(T)$. □

No linear map from finite-dimensional vector space to a smaller vector space can be injective.

Corollary 1.2.33. *If $\mathcal{V}$ and $\mathcal{W}$ are finite-dimensional vector spaces such that dim$(\mathcal{V})$ > dim$(\mathcal{W})$, then no linear map from $\mathcal{V}$ to $\mathcal{W}$ is injective.*

Proof. Suppose $\mathcal{V}$ and $\mathcal{W}$ are finite-dimensional vector spaces such that $\dim(\mathcal{V}) > \dim(\mathcal{W})$. Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$,

$$\begin{aligned} \dim(\text{null}(T)) &= \dim(\mathcal{V}) - \dim(\text{range}(T)) \\ &\geq \dim(\mathcal{V}) - \dim(\mathcal{W}) \geq 0 \end{aligned}$$

As $\dim(\text{null}(T)) \geq 0$, $\text{null}(T)$ must contain vectors other than 0, hence T is not injective. □

No linear map from finite-dimensional vector space to a bigger vector space can be surjective.

Corollary 1.2.34. *If $\mathcal{V}$ and $\mathcal{W}$ are finite-dimensional vector spaces such that $\dim(\mathcal{V}) < \dim(\mathcal{W})$, then no linear map from $\mathcal{V}$ to $\mathcal{W}$ is surjective.*

Proof. Suppose $\mathcal{V}$ and $\mathcal{W}$ are finite-dimensional vector spaces such that $\dim(\mathcal{V}) < \dim(\mathcal{W})$. Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$,

$$\begin{aligned}\dim(\text{range}(T)) &= \dim(\mathcal{V}) - \dim(\text{null}(T)) \\ &\leq \dim(\mathcal{V}) < \dim(\mathcal{W})\end{aligned}$$

As $\dim(\text{range}(T)) < \dim(\mathcal{W})$, $\text{range}(T)$ cannot equal $\mathcal{W}$, hence T is not surjective. □

Definition 1.2.35. A $m \times n$ *matrix* $\mathcal{M}$ has m rows and n columns,

$$\begin{bmatrix} a_{1,1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots \\ a_{m,1} & \cdots & a_{m,n} \end{bmatrix}$$

Proposition 1.2.36. *Suppose $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, and $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$, and $(\mathbf{w}_1, \ldots, \mathbf{w}_m)$ is a basis of $\mathcal{W}$. Then $\mathcal{M}(T\mathbf{v}) = \mathcal{M}(T)\mathcal{M}(\mathbf{v})$ for every $\mathbf{v} \in \mathcal{V}$.*

Proof. Let

$$\begin{bmatrix} a_{1,1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots \\ a_{m,1} & \cdots & a_{m,n} \end{bmatrix}$$

Note that

$$T\mathbf{v}_k = \sum_{j=1}^{m} a_{j,k}\mathbf{w}_j$$

for each k. Let $\mathbf{v}$ be an arbitrary vector in $\mathcal{V}$, then $\mathcal{M}(\mathbf{v})$ is given by

$$\begin{aligned} T\mathbf{v} &= b_1 T\mathbf{v}_1 + \cdots + b_n T\mathbf{v}_n \\ &= b_1 \sum_{j=1}^{m} a_{j,1}\mathbf{w}_j + \cdots + b_n \sum_{j=1}^{m} a_{j,n}\mathbf{w}_j \\ &= \sum_{j=1} (a_{j,1}b_1 + \cdots + a_{j,n}b_n)\mathbf{w}_j \end{aligned}$$

The equation shows that $\mathcal{M}(T\mathbf{v})$ is a $m \times 1$ matrix of vector $T\mathbf{v}$ with respect to basis $(\mathbf{w}_1, \ldots, \mathbf{w}_m)$, given by

$$\mathcal{M}(T\mathbf{v}) = \begin{bmatrix} a_{1,1}b_1 + \cdots + a_{1,n}b_n \\ \vdots \\ a_{m,1}b_1 + \cdots + a_{m,n}b_n \end{bmatrix}$$

Hence $\mathcal{M}(T\mathbf{v}) = \mathcal{M}(T)\mathcal{M}(\mathbf{v})$. □

Definition 1.2.37. A linear map $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ is *invertible* if $\exists$ linear map $S \in \mathcal{L}(\mathcal{W}, \mathcal{V})$ such that ST equals identity map on $\mathcal{V}$ and TS equals identity map on $\mathcal{W}$.
A linear map $S \in \mathcal{L}(\mathcal{W}, \mathcal{V})$ satisfying $ST = I$ and $TS = I$ is the *inverse* of T.

Proposition 1.2.38. *A linear map is invertible if and only if it is injective and surjective.*

Proof. Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$.
($\Rightarrow$) Suppose T is invertible, $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, and $T\mathbf{u} = T\mathbf{v}$. Then

$$\mathbf{u} = T^{-1}(T\mathbf{u}) = T^{-1}T(\mathbf{v}) = \mathbf{v}$$

hence $\mathbf{u} = \mathbf{v}$. Thus T is injective.
To show T is surjective, let $\mathbf{w} \in \mathcal{W}$, then $\mathbf{w} = T(T^{-1}\mathbf{w})$, thus $\mathbf{w}$ is in range of T. Hence $\text{range}(T) = W$.
Suppose T is injective and surjective, to prove T is invertible. For each $\mathbf{w} \in \mathcal{W}$, define $S\mathbf{w}$ to be unique element

of $\mathcal{V}$ such that $T(S\mathbf{w}) = \mathbf{w}$, which shows TS equals identity map on $\mathcal{W}$. To prove ST equals identity map on $\mathcal{V}$, let $\mathbf{v} \in \mathcal{V}$, then

$$T(ST\mathbf{v}) = (TS)(T\mathbf{v}) = I(T\mathbf{v}) = T\mathbf{v}$$

which implies $ST\mathbf{v} = \mathbf{v}$ (as T is injective), thus ST equals identity map on $\mathcal{V}$. Let $\mathbf{w}_1, \mathbf{w}_2 \in \mathcal{W}$, then

$$T(S\mathbf{w}_1 + S\mathbf{w}_2) = T(S\mathbf{w}_1) + T(S\mathbf{w}_2) = \mathbf{w}_1 + \mathbf{w}_2$$

Thus $S\mathbf{w}_1 + S\mathbf{w}_2$ is unique element of $\mathcal{V}$ that maps T to $\mathbf{w}_1 + \mathbf{w}_2$. This implies S satisfies the additive property requires for linearity.
For proof of homogeneity, let $\mathbf{w} \in \mathcal{W}$ and $a \in \mathbb{F}$, then

$$T(aS\mathbf{w}) = aT(S\mathbf{w}) = a\mathbf{w}$$

Thus $aS\mathbf{w}$ is the unique element of $\mathcal{V}$ that T maps to $a\mathbf{w}$. By definition of S, this implies $S(a\mathbf{w}) = aS\mathbf{w}$, hence S is linear, as desired. □

Definition 1.2.39. Two vector spaces are *isomorphic* if there is an invertible linear map from one vector space onto the other. Two isomorphic spaces have the same properties.

Theorem 1.2.40. *Two finite-dimensional vector spaces are isomorphic if and only if they have the same dim.*

Proof. ($\Rightarrow$) Suppose $\mathcal{V}$ and $\mathcal{W}$ are isomorphic finite-dimensional vector spaces.
$\exists$ invertible linear map $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. As T is invertible, $\text{null}(T) = \{\mathbf{0}\}$, and $\text{range}(T) = \mathcal{W}$, thus $\dim(\text{null}(T)) = 0$ and $\dim(\text{range}(T)) = \dim(\mathcal{W})$.

$$\begin{aligned}\dim(\mathcal{V}) &= \dim(\text{null}(T)) + \dim(\text{range}(T)) \\ &= \dim(\mathcal{W})\end{aligned}$$

($\Leftarrow$) Suppose $\mathcal{V}$ and $\mathcal{W}$ are finite-dimensional vector spaces with the same dimension. let $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be basis of $\mathcal{V}$ and $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$ be basis of $\mathcal{W}$. Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ defined by

$$T(a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n) = a_1\mathbf{w}_1 + \cdots + a_n\mathbf{w}_n$$

Then T is surjective as $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$ spans $\mathcal{W}$, and T is injective as $(\mathbf{w}_1, \ldots, \mathbf{w}_n)$ is linearly independent.
As T is injective and surjective, it is invertible, hence $\mathcal{V}$ and $\mathcal{W}$ are isomorphic. □

The next theorem implies that every finite-dimensional vector space is isomorphic to some $\mathbb{F}^n$.

Theorem 1.2.41. *Suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$ and $(\mathbf{w}_1, \cdots, \mathbf{w}_m)$ is a basis of $\mathcal{W}$. Then $\mathcal{M}$ is an invertible linear map between $\mathcal{L}(\mathcal{V}, \mathcal{W})$ and Mat$(m, n, \mathbb{F})$,*

Proof. Note that $\mathcal{M}$ is linear, hence to prove that $\mathcal{M}$ is injective and surjective.
If $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ and $\mathcal{M}(T) = 0$, then $T\mathbf{v}_k = \mathbf{0}$ for $k = 1, \ldots, n$, As $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$, this implies $T = 0$, hence $\mathcal{M}$ is injective.
To show $\mathcal{M}$ is surjective, let

$$A = \begin{bmatrix} a_{1,1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots \\ a_{m,1} & \cdots & a_{m,n} \end{bmatrix}$$

be a matrix in $\text{Mat}(m, n, \mathbb{F})$. Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ such that

$$T\mathbf{v}_k = \sum_{j=1}^{m} a_{j,k}\mathbf{w}_k$$

for $k = 1, \ldots, n$. Note $\mathcal{M}(T) = A$, so range of $\mathcal{M}$ equals $\text{Mat}(m, n, \mathbb{F})$ as desired. □

Proposition 1.2.42. *If $\mathcal{V}$ and $\mathcal{W}$ are finite-dimensional, then $\mathcal{L}(\mathcal{V}, \mathcal{W})$ is finite-dimensional, and*

$$dim\mathcal{L}(\mathcal{V}, \mathcal{W}) = (dim(\mathcal{V}))(dim(\mathcal{W}))$$

Proof. This follows from $\dim(\text{Mat}(m, n, \mathbb{F})) = mn$ and Theorem 1.2.40 and Theorem 1.2.41. □

Definition 1.2.43. A linear map from a vector space to itself is an *operator*, denoted $\mathcal{L}(\mathcal{V})$.

Theorem 1.2.44. *Suppose $\mathcal{V}$ is finite-dimensional. If $T \in \mathcal{L}(\mathcal{V})$, then the following are equivalent:*

i. T is invertible.

ii. T is injective.

iii. T is surjective.

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$, then $(i) \Rightarrow (ii)$.
Suppose (ii) holds so that T is injective. Thus $\text{null}(T) = \{\mathbf{0}\}$. Then

$$\begin{aligned}\dim(\text{range}(T)) &= \dim(\mathcal{V}) - \dim(\text{null}(T)) \\ &= \dim(\mathcal{V})\end{aligned}$$

which implies range of T equals $\mathcal{V}$, hence T is surjective, and $(ii) \Rightarrow (iii)$.
Suppose (iii) holds, so T is surjective. Thus $\text{range}(T) = \mathcal{V}$. Then

$$\dim(\text{null}(T)) = \dim(\mathcal{V}) - \dim(\text{range}(T)) = 0$$

which implies $\text{null}(T)$ equals $\{\mathbf{0}\}$, hence T is injective and thus invertible. Hence $(iii) \Rightarrow (i)$. □

1.2.3 Eigenvalues and Eigenvectors

Definition 1.2.45. Let $T \in \mathcal{L}(\mathcal{V})$, and $\exists$ direct sum decomposition $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$ where each $\mathcal{U}_j$ is proper subspace of $\mathcal{V}$. The *restriction* of T to smaller domain $\mathcal{U}_j$ is denoted $T|_{\mathcal{U}_j}$

Definition 1.2.46. For $T \in \mathcal{L}(\mathcal{V})$ and $\mathcal{U}$ a subspace of $\mathcal{V}$, $\mathcal{U}$ is invariant under T if $\mathbf{u} \in \mathcal{U}$ implies $T\mathbf{u} \in \mathcal{U}$ ($\mathcal{U}$ is invariant under T if $T|_{\mathcal{U}}$ is an operator under $\mathcal{U}$.

Definition 1.2.47. A scalar $\lambda \in \mathbb{F}$ is an *eigenvalue* of $T \in \mathcal{L}(\mathcal{V})$ if $\exists$ a nonzero vector $\mathbf{u} \in \mathcal{V}$ such that $T\mathbf{u} = \lambda\mathbf{u}$. A vector $\mathbf{u} \in \mathcal{V}$ is an *eigenvector* of T if $T\mathbf{u} = \lambda\mathbf{u}$.

Note that the set of eigenvectors of T corresponding to λ equals $\text{null}(T - \lambda I)$.

Theorem 1.2.48. *Let $T \in \mathcal{L}(\mathcal{V})$, and $\lambda_1, \ldots, \lambda_m$ distinct eigenvalues of T, $\mathbf{v}_1, \ldots, \mathbf{v}_m$ are corresponding nonzero eigenvectors, then $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly independent.*

Proof. Suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly dependent. Let k be smallest positive integer such that

$$\mathbf{v}_k \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$$

The existence of k with this property follows from Lemma 1.2.8. Hence $\exists\ a_1, \ldots, a_{k-1} \in \mathbb{F}$ such that

$$\mathbf{v}_k = a_1\mathbf{v}_1 + \cdots + a_{k-1}\mathbf{v}_{k-1}$$

Applying T to both sides of equation gives

$$\lambda_k\mathbf{v}_k = a_1\lambda_1\mathbf{v}_1 + \cdots + a_{k-1}\lambda_{k-1}\mathbf{v}_{k-1}$$

Multiply both sides of second equation by λ_k then subtracting above gives

$$0 = a_1(\lambda_k - \lambda_1)\mathbf{v}_1 + \cdots + a_{k-1}(\lambda_k - \lambda_{k-1})\mathbf{v}_{k-1}$$

As k is chosen to be smallest positive integer satisfying first equation, $(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$ is linearly dependent, hence equation above implies that all a are 0, which also means $\mathbf{v}_k = \mathbf{0}$, contracting hypothesis that all $\mathbf{v}$ are nonzero. Hence assumption that $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly dependent must be false. □

Corollary 1.2.49. *Each operator on $\mathcal{V}$ has at most $\dim(\mathcal{V})$ distinct eigenvalues.*

Proof. Let $T \in \mathcal{L}(\mathcal{V})$, and $\lambda_1, \ldots, \lambda_m$ distinct eigenvalues of T. Let $\mathbf{v}_1, \ldots, \mathbf{v}_m$ be corresponding nonzero eigenvectors. Theorem 1.2.48 implies $(\mathbf{v}_1, \ldots, \mathbf{v}_m)$ is linearly independent, hence $m \leq \dim(\mathcal{V})$. □

Theorem 1.2.50. *Every operator on a finite-dimensional, nonzero complex vector space has an eigenvalue.*

Proof. Suppose $\mathcal{V}$ is a complex vector space with $\dim(\mathcal{V}) = n > 0$ and $T \in \mathcal{L}(\mathcal{V})$. Choose $\mathbf{v} \in \mathcal{V}$ with $\mathbf{v} \neq \mathbf{0}$, then $(\mathbf{v}, T\mathbf{v}, T^2\mathbf{v}, \ldots, T^n\mathbf{v})$ cannot be linearly independent as $\mathcal{V}$ has dimension n and $n+1$ vectors. Thus $\exists\ a_0, \ldots, a_n \in \mathbb{C}$ not all 0 such that

$$0 = a_0\mathbf{v} + a_1T\mathbf{v} + \cdots + a_nT^n\mathbf{v}$$

Let m be largest index such that $a_m \neq 0$. As $\mathbf{v} \neq \mathbf{0}$, the coefficients $a_1, \ldots, a_m$ cannot all be 0, hence $0 < m \leq n$. Let a e coefficients of a polynomial in factored form

$$a_0 + a_1z + \cdots + a_nz^n = c(z - \lambda_1)\cdots(z - \lambda_m)$$

where c is nonzero complex number, each $\lambda_j \in \mathbb{C}$, and equation holds for all $z \in \mathbb{C}$, then

$$\begin{aligned}\mathbf{0} &= a_0\mathbf{v} + a_1T\mathbf{v} + \cdots + a_nT^n\mathbf{v} \\ &= (a_0I + a_1T + \cdots + a_nT^n)\mathbf{v} \\ &= c(T - \lambda_1I)\cdots(T - \lambda_mI)\mathbf{v}\end{aligned}$$

which means $T - \lambda_jI$ is not injective for at least one j. Hence T has an eigenvalue. □

Definition 1.2.51. *Upper triangular matrices* are of the form

$$\begin{bmatrix}\lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n\end{bmatrix}$$

Next proposition shows useful connection between upper-triangular matrices and invariant subspaces.

Proposition 1.2.52. *Suppose $T \in \mathcal{L}(\mathcal{V})$, and $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$. Following are equivalent:*

i. The matrix of T with respect to $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is upper triangular.

ii. $T\mathbf{v}_k \in$ span$(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ for each $k = 1, \ldots, n$

iii. span$(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is invariant under T for each $k = 1, \ldots, n$

Proof. $(i) \Rightarrow (ii)$ follows from definitions. Also, $(iii) \Rightarrow (ii)$ as (iii) implies (ii).
$(ii) \Rightarrow (iii)$, suppose (ii) holds. Fix $k \in \{1, \ldots, n\}$. From (ii), we know that

$$\begin{aligned}T\mathbf{v}_1 &\in \text{span}(\mathbf{v}_1) \subset \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k) \\ T\mathbf{v}_2 &\in \text{span}(\mathbf{v}_1, \mathbf{v}_2) \subset \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k) \\ &\vdots \\ T\mathbf{v}_k &\in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)\end{aligned}$$

Hence if $\mathbf{v}$ is a linear combination of $(\mathbf{v}_1, \ldots, \mathbf{v}_k)$, then

$$T\mathbf{v} \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$$

Hence, span$(\mathbf{v}_1, \ldots, \mathbf{v}_k)$ is invariant under T. □

Next theorem shows that for each operator on a complex vector space, there is a basis of the vector space with respect to which the matrix of operator only has 0 below the diagonal.

Theorem 1.2.53. *Suppose $\mathcal{V}$ is a complex vector space and $T \in \mathcal{L}(\mathcal{V})$, then T has an upper-triangular matrix with respect to some basis of $\mathcal{V}$.*

Proof. Note that result holds if $\dim(\mathcal{V}) = 1$.
Suppose $\dim(\mathcal{V}) > 1$ and the result holds for all complex vector spaces with dimension less than $\dim(\mathcal{V})$. Let λ be any eigenvalue of T, and let $\mathcal{U} = \text{range}(T - \lambda I)$. As $T - \lambda I$ is not surjective, $\dim(\mathcal{U}) < \dim(\mathcal{V})$, hence $\mathcal{U}$ is invariant under T. To show this, let $\mathbf{u} \in \mathcal{U}$, then

$$T\mathbf{u} = (T - \lambda I)\mathbf{u} + \lambda\mathbf{u}$$

$(T - \lambda I)\mathbf{u} \in \mathcal{U}$ by definition, and $\lambda\mathbf{u} \in \mathcal{U}$. Hence, equation shows $T\mathbf{u} \in \mathcal{U}$, and $\mathcal{U}$ is invariant under T.
Hence $T|_{\mathcal{U}}$ is an operator on $\mathcal{U}$. By induction, $\exists$ basis $(\mathbf{u}_1, \ldots \mathbf{u}_m)$ of $\mathcal{U}$ with respect to which $T|_{\mathcal{U}}$ has an

upper-triangular matrix. Hence for each j,

$$T\mathbf{u}_j = (T|_{\mathcal{U}})(\mathbf{u}_j) \in \text{span}(\mathbf{u}_1, \ldots, \mathbf{u}_j)$$

Extend $(\mathbf{u}_1, \ldots, \mathbf{u}_m)$ to basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_n)$ of $\mathcal{V}$. For each k,

$$T\mathbf{v}_k = (T - \lambda I)\mathbf{v}_k + \lambda \mathbf{v}_k$$

Definition of $\mathcal{U}$ shows that $(T - \lambda I)\mathbf{v}_k \in \mathcal{U} = \text{span}(\mathbf{u}_1, \ldots, \mathbf{u}_m)$. Hence,

$$T\mathbf{v}_k \in \text{span}(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_k)$$

Hence T has an upper triangular matrix with respect to basis $(\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_k)$. □

Proposition 1.2.54. *Suppose $T \in \mathcal{L}(\mathcal{V})$ has an upper-triangular matrix with respect to some basis of $\mathcal{V}$. Then T is invertible if and only if all entries of the diagonal of upper-triangular matrix are non-zero.*

Proof. Let $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ be basis of $\mathcal{V}$ with respect to T which has upper-triangular matrix

$$\mathcal{M}(T, (\mathbf{v}_1, \ldots, \mathbf{v}_n)) = \begin{bmatrix} \lambda_1 & & * \\ & \lambda_2 & \\ & & \ddots \\ & & & \lambda_n \end{bmatrix}$$

($\Rightarrow$) If $\lambda_1 = 0$, then $T\mathbf{v}_1 = \mathbf{0}$, hence T is not invertible
Suppose $1 < k \leq n$ and $\lambda_k = 0$, then T maps each of the vectors $\mathbf{v}_1, \ldots, \mathbf{v}_{k-1}$ into $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$. As $\lambda_k = 0$, the matrix representation implies $T\mathbf{v}_k \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$. Hence, define a linear map

$$S : \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k) \to \text{span}(\mathbf{v}_1, \mathbf{v}_{k-1})$$

by $S\mathbf{v} = T\mathbf{v}$ for $\mathbf{v} \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$. Then S is T restricted to $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$.
Note $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$ has dimension k and $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$ has dimension $k-1$ (as $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is linearly independent). As $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$ has larger dimension than $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$, no linear map from $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$ to $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$ is injective, hence there exists nonzero vector $\mathbf{v} \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_k)$ such that $S\mathbf{v} = \mathbf{0}$. Hence $T\mathbf{v} = \mathbf{0}$, and hence T is not invertible.
($\Leftarrow$) Suppose T is not invertible. Thus T is not injective, and hence there exists a nonzero vector $\mathbf{v} \in \mathcal{V}$ such that $T\mathbf{v} = \mathbf{0}$. As $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$, then $\mathbf{v} = a_1\mathbf{v}_1 + \cdots + a_k\mathbf{v}_k$ where $a_1, \ldots, a_k \in \mathbb{F}$ and $a_k \neq 0$ (represent $\mathbf{v}$ as linear combination of $(\mathbf{v}_1, \ldots, \mathbf{v}_k)$ then choose k to be largest index with nonzero coefficient).

$$\begin{aligned} \mathbf{0} = T\mathbf{v} &= T(a_1\mathbf{v}_1 + \cdots + a_k\mathbf{v}_k) \\ &= (a_1T\mathbf{v}_1 + \cdots + a_{k-1}T\mathbf{v}_{k-1}) + a_kT\mathbf{v}_k \end{aligned}$$

The last term in parentheses is in $\text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$ (due to upper-triangular form).
Hence this shows $a_kT\mathbf{v}_k \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$. Multiply by $1/a_k$, which is allowed as $a_k \neq 0$, conclude that $T\mathbf{v}_k \in \text{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{k-1})$. Thus when $T\mathbf{v}_k$ is written as linear combination of the basis $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$, the coefficient of $\mathbf{v}_k$ will be 0, hence λ_k must be 0, completing the proof. □

Proposition 1.2.55. *Suppose $T \in \mathcal{L}(\mathcal{V})$ has upper-triangular matrix with respect to some basis of $\mathcal{V}$, then eigenvalues of T consists of precisely the entries of the diagonal of the upper-triangular matrix.*

Proof. Suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$ with respect to T which T has an upper-triangular matrix

$$\mathcal{M}(T, (\mathbf{v}_1, \ldots, \mathbf{v}_n)) = \begin{bmatrix} \lambda_1 & & * \\ & \lambda_2 & \\ & & \ddots \\ 0 & & & \lambda_n \end{bmatrix}$$

Let $\lambda \in \mathbb{F}$, then

$$\mathcal{M}(T - \lambda I, (\mathbf{v}_1, \ldots, \mathbf{v}_n)) = \begin{bmatrix} \lambda_1 - \lambda & & & * \\ & \lambda_2 - \lambda & & \\ & & \ddots & \\ 0 & & & \lambda_n - \lambda \end{bmatrix}$$

Hence $T - \lambda I$ is not invertible if and only if λ equals one of the λ_j.
Hence λ is an eigenvalue of T if and only if λ equals one of the λ_j. □

Definition 1.2.56. A *diagonal matrix* is a square matrix that is 0 everywhere except possibly along diagonal.

Operator $T \in \mathcal{L}(\mathcal{V})$ has diagonal matrix $\text{diag}(\lambda_1, \ldots, \lambda_n)$ w.r.t. basis $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ of $\mathcal{V}$ if and only if

$$\begin{aligned} T\mathbf{v}_1 &= \lambda_1 \mathbf{v}_1 \\ &\vdots \\ T\mathbf{v}_n &= \lambda_n \mathbf{v}_n \end{aligned}$$

This follows from definition of matrix of operator with respect to a basis,

If an operator has a diagonal matrix w.r.t. some basis, the entires along diagonal are precisely eigenvalues.

Proposition 1.2.57. *If $T \in \mathcal{L}(\mathcal{V})$ has $\dim(\mathcal{V})$ distinct eigenvalues, then T has a diagonal matrix with respect to some basis of $\mathcal{V}$.*

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$ has $\dim(\mathcal{V})$ distinct eigenvalues $\lambda_1, \ldots, \lambda_{\dim(\mathcal{V})}$. For each j, let $\mathbf{v} \in \mathcal{V}$ be nonzero eigenvector corresponding to the eigenvalue λ_j. As nonzero eigenvectors corresponding to distinct eigenvalues are linearly independent, $(\mathbf{v}_1, \ldots, \mathbf{v}_{\dim(\mathcal{V})})$ is linearly independent. A linearly independent list of $\dim(\mathcal{V})$ vectors in $\mathcal{V}$ is a basis of $\mathcal{V}$, hence $(\mathbf{v}_1, \ldots, \mathbf{v}_{\dim(\mathcal{V})})$ is a basis of $\mathcal{V}$. With respect to this basis, T is a diagonal matrix. □

Proposition 1.2.58. *Suppose $T \in \mathcal{L}(\mathcal{V})$. Let $\lambda_1, \ldots, \lambda_m$ denote distinct eigenvalues of T. Then the following are equivalent:*

i. T has a diagonal matrix with respect to some basis of $\mathcal{V}$;

ii. $\mathcal{V}$ has a basis consisting of eigenvectors of T;

iii. $\exists$ one-dimensional subspaces $\mathcal{U}_1, \ldots, \mathcal{U}_n$ of $\mathcal{V}$, each invariant under T, such that

$$\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_n$$

iv. $\mathcal{V} = \text{null}(T - \lambda_1 I) \oplus \cdots \oplus \text{null}(T - \lambda_m I)$

v. $\dim(\mathcal{V}) = \dim(\text{null}(T - \lambda_1 I)) + \cdots + \dim(\text{null}(T - \lambda_m I))$

Proof. Already shown earlier that (i) and (ii) are equivalent.
$(ii) \Rightarrow (iii)$. Suppose (ii) holds, thus $\mathcal{V}$ has a basis $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ consisting of eigenvectors of T. For each j, let $\mathcal{U}_j = \text{span}(\mathbf{v}_j)$, and hence each $\mathcal{U}_j$ is a one-dimensional subspace of $\mathcal{V}$ that is invariant under T (as each $\mathbf{v}_j$ is an eigenvector of T). As $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$, each vector in $\mathcal{V}$ can be written uniquely as a linearly combination of $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$. Hence each vector in $\mathcal{V}$ can be written uniquely as a sum $\mathbf{u}_1 + \cdots + \mathbf{u}_n$, $\mathbf{u}_j \in \mathcal{U}_j$. Thus $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_n$, and (ii) implies (iii).
$(iii) \Rightarrow (ii)$. Suppose (iii) holds, hence there are one-dimensional subspaces $\mathcal{U}_1, \ldots, \mathcal{U}_n$ of $\mathcal{V}$, each invariant under T, such that $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_n$. For each j, let $\mathbf{v}_j$ be nonzero vector in $\mathcal{U}_j$. Then each $\mathbf{v}_j$ is an eigenvector of T. As each vector in $\mathcal{V}$ can be written uniquely as $\mathbf{u}_1 + \cdots + \mathbf{u}_n, \mathbf{u}_j \in \mathcal{U}_j$, $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a basis of $\mathcal{V}$.
Hence this shows so far $(i) \Leftrightarrow (ii) \Leftrightarrow (iii)$.
$(ii) \Rightarrow (iv)$. Suppose (ii) holds, then $\mathcal{V}$ has a basis consisting of eigenvectors of T, and hence every vector in $\mathcal{V}$ is a linear combination of eigenvectors of T. Thus

$$\mathcal{V} = \text{null}(T - \lambda_1 I) + \cdots + \text{null}(T - \lambda_m I)$$

Suppose $\mathbf{0} = \mathbf{u}_1 + \cdots + \mathbf{u}_m, \mathbf{u}_j \in \text{null}(T - \lambda_j I)$. As nonzero eigenvectors corresponding to distinct eigenvalues are linearly independent, this implies each $\mathbf{u}_j = \mathbf{0}$. This implies the sum is a direct sum.
$(iv) \Rightarrow (v)$. Suppose (iv) holds. By definition, each $\text{null}(T - \lambda_j I)$ is a subspace of $\mathcal{V}$. By definition of direct sum, each $\mathbf{V} \in \mathcal{V}$ can be uniquely expressed as $\mathbf{v} = \mathbf{u}_1 + \cdots + \mathbf{u}_m, \mathbf{u}_j \in \text{null}(T - \lambda_j I)$.
The only way for each $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$ with each $\mathbf{u}_j \in \text{null}(T - \lambda_j I)$ is if each $\mathbf{u}_j = \mathbf{0}$. Hence these properties together imply $(iv) \Rightarrow (v)$.
$(v) \Rightarrow (ii)$. Suppose (v) holds. Choose a basis $\mathbf{v}_j \in \text{null}(T - \lambda_j I)$, and form a list $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ of eigenvectors of T, where $n = \dim(\mathcal{V})$. Suppose $a_1 \mathbf{v}_1 + \cdots + a_n \mathbf{v}_n = \mathbf{0}$, where $a_1, \ldots, a_n \in \mathbb{F}$. For each $j = 1, \ldots, m$, let $\mathbf{u}_j$ be sum of all terms $a_k \mathbf{v}_k$ such that $\mathbf{v}_k \in \text{null}(T - \lambda_j I)$. Thus each $\mathbf{u}_j$ is an eigenvector of T with eigenvalue λ_j, and $\mathbf{u}_1 + \cdots + \mathbf{u}_m = \mathbf{0}$. As nonzero eigenvectors corresponding to distinct eigenvalues are linearly independent, this implies each $\mathbf{u}_j = \mathbf{0}$. As each $\mathbf{u}_j$ is sum of $a_k \mathbf{v}_k$, where $\mathbf{v}_k$ were chosen to be a basis of $\text{null}(T - \lambda_j I)$, this implies all $a_k = 0$. Hence $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is linearly independent and hence is a basis of $\mathcal{V}$. □

Next theorem shows that an invariant subspace of dimension 1 or 2 always exists.

Theorem 1.2.59. *Every operator on a finite-dimensional, nonzero, real vector space has an invariant subspace of dimension* 1 *or* 2.

Proof. Let $\mathcal{V}$ be a real vector space, $\dim(\mathcal{V}) = n > 0$, and $T \in \mathcal{L}(\mathcal{V})$. Choose $\mathbf{v} \in \mathcal{V}$ with $\mathbf{v} \neq \mathbf{0}$, then

$$(\mathbf{v}, T\mathbf{v}, T^2\mathbf{v}, \dots, T^n\mathbf{v})$$

cannot be linearly independent as $\mathcal{V}$ has dimension n, and there are $n+1$ vectors.
Hence $\exists\ a_0, \dots a_n \in \mathbb{R}$ not all 0, such that

$$\mathbf{0} = a_0\mathbf{v} + a_1T\mathbf{v} + \cdots + a_nT^n\mathbf{v}$$

Let a be coefficients of a polynomial, then the factored form is

$$a_0 + a_1x + \cdots + a_nx^n = c(x - \lambda_1)\cdots(x - \lambda_m)(x^2\alpha_1x + \beta_1)\cdots(x^2 + \alpha_Mx + \beta_M)$$

where $c \neq 0 \in \mathbb{R}$, $\lambda_j, \alpha_j, \beta_j \in \mathbb{R}$, $m + M \geq 1$, and equation holds for all $x \in \mathbb{R}$. Then

$$\begin{aligned}\mathbf{0} &= a_0\mathbf{v} + a_1T\mathbf{v} + \cdots + a_nT^n\mathbf{v}\\ &= (a_0I + a_1T + \cdots + a_nT^n)\mathbf{v}\\ &= c(T - \lambda_1I)\cdots(T - \lambda_mI)(T^2 + \alpha_1T + \beta_1I)\cdots(T^2 + \alpha_MT + \beta_MI)\mathbf{v}\end{aligned}$$

This means $T - \lambda_jI$ is not injective for at least one j, or $(T^2 + \alpha_jT + \beta_jI)$ is not injective for at least one j. If $T - \lambda_jI$ is not injective for at least one j, then T has an eigenvalue, and hence a one-dimensional subspace.
If $(T^2 + \alpha_jT + \beta_jT)$ is not injective for some j, then there exists nonzero vector $\mathbf{u} \in \mathcal{V}$ such that

$$T^2\mathbf{u} + \alpha_jT\mathbf{u} + \beta_j\mathbf{u} = \mathbf{0}$$

Consider typical element of $\operatorname{span}(\mathbf{u}, T\mathbf{u})$ of form $a\mathbf{u} + bT\mathbf{u}$, where $a, b \in \mathbb{R}$, then

$$\begin{aligned}T(a\mathbf{u} + bT\mathbf{u}) &= aT\mathbf{u} + bT^2\mathbf{u}\\ &= aT\mathbf{u} - b\alpha_jT\mathbf{u} - b\beta_j\mathbf{u}\end{aligned}$$

last equality comes from solving the equality for $T^2\mathbf{u}$. This shows $T(a\mathbf{u}+bT\mathbf{u}) \in \operatorname{span}(\mathbf{u}, T\mathbf{u})$. hence $\operatorname{span}(\mathbf{u}, T\mathbf{u})$ is invariant under T. □

Definition 1.2.60. Let $\mathcal{U}$, $\mathcal{W}$ be subspaces of $\mathcal{V}$, where $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$.
Let $\mathbf{u} \in \mathcal{U}, \mathbf{w} \in \mathcal{W}, \mathbf{v} \in \mathcal{V}$, and $\mathbf{v} = \mathbf{u} + \mathbf{w}$. Define *projection* $P_{\mathcal{U},\mathcal{W}} \in \mathcal{L}(\mathcal{V})$ by $P_{\mathcal{U},\mathcal{W}}\mathbf{v} = \mathbf{u}$. Note that

$$\begin{aligned}P_{\mathcal{W},\mathcal{U}}\mathbf{v} = \mathbf{w}, &\quad \mathbf{v} = P_{\mathcal{U},\mathcal{W}}\mathbf{v} + P_{\mathcal{W},\mathcal{U}}\mathbf{v}\\ \operatorname{range}(P_{\mathcal{U},\mathcal{W}}) = \mathcal{U}, &\quad \operatorname{null}(P_{\mathcal{U},\mathcal{W}}) = \mathcal{W}\end{aligned}$$

Theorem 1.2.61. *Every operator on an odd-dimensional real vector space has an eigenvalue.*

Proof. Suppose $\mathcal{V}$ is a real vector space with odd dimension.
Note that result holds for $\dim(\mathcal{V}) = 1$.
Suppose $\dim(\mathcal{V})$ is an odd number greater than 1. Suppose $T \in \mathcal{L}(\mathcal{V})$, to prove T has an eigenvalue. By Theorem 1.2.59, there is a two-dimensional subspace $\mathcal{U}$ of $\mathcal{V}$ that is invariant under T. Let $\mathcal{W}$ be any subspace of $\mathcal{V}$ such that $\mathcal{V} = \mathcal{U} \oplus \mathcal{W}$, then Theorem 1.2.59 guarantees that such a $\mathcal{W}$ exists.
As $\mathcal{W}$ has dimension 2 less than $\dim(\mathcal{V})$, apply induction hypothesis to $T|_{\mathcal{W}}$. However, $\mathcal{W}$ might not be invariant under T, which means $T_{\mathcal{W}}$ might not be an operator on $\mathcal{W}$.
Define $S \in \mathcal{L}(\mathcal{W})$ by $S\mathbf{w} = P_{\mathcal{W},\mathcal{U}}(T\mathbf{w})$ for $\mathbf{w} \in \mathcal{W}$. By induction, S has an eigenvalue λ.
Let $\mathbf{w} \in \mathcal{W}$ be nonzero eigenvector for S corresponding to the eigenvalue λ, thus $(S - \lambda I)\mathbf{w} = 0$. To look for an eigenvector of T in $\mathcal{U} + \operatorname{span}(\mathbf{w})$. Consider typical vector $\mathbf{u} + a\mathbf{w}$ in $\mathcal{U} = \operatorname{span}(\mathbf{w})$, where $\mathbf{u} \in \mathcal{U}$ and $a \in \mathbb{R}$.

$$\begin{aligned}(T - \lambda I)(\mathbf{u} + a\mathbf{w}) &= T\mathbf{u} - \lambda\mathbf{u} + a(T\mathbf{w} - \lambda\mathbf{w})\\ &= T\mathbf{u} - \lambda\mathbf{u} + a(P_{\mathcal{U},\mathcal{W}}(T\mathbf{w}) + P_{\mathcal{W},\mathcal{U}}(T\mathbf{w}) - \lambda\mathbf{w})\\ &= T\mathbf{u} - \lambda\mathbf{u} + a(P_{\mathcal{U},\mathcal{W}}(T\mathbf{w}) + S\mathbf{w} - \lambda\mathbf{w})\\ &= T\mathbf{u} - \lambda\mathbf{u} + aP_{\mathcal{U},\mathcal{W}}(T\mathbf{w})\end{aligned}$$

Note $T\mathbf{u} \in \mathcal{U}$ (as $\mathcal{U}$ is invariant under T), $\lambda\mathbf{u} \in \mathcal{U}$ (as $\mathbf{u} \in \mathcal{U}$), and $aP_{\mathcal{U},\mathcal{W}}(T\mathbf{w}) \in \mathcal{U}$ (by definition). Hence $T - \lambda I$ maps $\mathcal{U} + \text{span}(\mathbf{w})$ into $\mathcal{U}$. As $\mathcal{U} + \text{span}(\mathbf{w})$ has larger dimension than $\mathcal{U}$ this means $(T - \lambda I)|_{\mathcal{U}+\text{span}(\mathbf{w})}$ is not injective, and there exists a nonzero vector $\mathbf{v} \in \mathcal{U} + \text{span}(\mathbf{w})$ such that $(T - \lambda I)\mathbf{v} = \mathbf{0}$. Hence T has an eigenvalue as desired. □

1.2.4 Inner-Product Spaces

Definition 1.2.62. The *norm* is defined as $\|\mathbf{z}\| = \sqrt{|z_1|^2 + \cdots + |z_n|^2} = \sqrt{\langle \mathbf{z}, \mathbf{z} \rangle}$ for $\mathbf{z} = (z_1, \ldots, z_n) \in \mathbb{C}^n$.
The norm should be think of as the *inner product* of $\mathbf{z}$ with itself, $\|z\|^2 = z_1\overline{z_1} + \cdots + z_n\overline{z_n}$.
If $\mathbf{p} \in \mathcal{P}_m(\mathbb{F})$ (polynomial space0, then $\|\mathbf{p}\| = \sqrt{\int_0^1 |\mathbf{p}(x)|^2 \, dx}$

Remark 1.2.63. An inner product on $\mathcal{V}$ that is a function that takes each ordered pair $(\mathbf{u}, \mathbf{v})$ of elements in $\mathcal{V}$ to a number $\langle \mathbf{u}, \mathbf{v} \rangle \in \mathbb{F}$ has following properties:

i. Positivity: $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0 \; \forall \; \mathbf{v} \in \mathcal{V}$
ii. Definiteness: $\langle \mathbf{v}, \mathbf{v} \rangle = 0$ if and only if $\mathbf{v} = \mathbf{0}$
iii. Additivity in First Slot: $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$ for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$
iv. Homogeneity in First Slot: $\langle a\mathbf{v}, \mathbf{w} \rangle = a \langle \mathbf{v}, \mathbf{w} \rangle$ for all $a \in \mathbb{F}$ and $\mathbf{v}, \mathbf{w} \in \mathcal{V}$
v. Conjugate Symmetry: $\langle \mathbf{v}, \mathbf{w} \rangle = \overline{\langle \mathbf{w}, \mathbf{v} \rangle}$ for all $\mathbf{v}, \mathbf{w} \in \mathcal{V}$

Definition 1.2.64. An *inner-product space* is a vector space $\mathcal{V}$ with an inner product on $\mathcal{V}$.

Definition 1.2.65. Two vectors $\mathbf{u}$, $\mathbf{v}$ are *orthogonal* if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

Theorem 1.2.66. *Pythagorean Theorem*
If $\mathbf{u}, \mathbf{v}$ are orthogonal vectors in $\mathcal{V}$, then $\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$.

Proof. Suppose $\mathbf{u}, \mathbf{v}$ are orthogonal vectors in $\mathcal{V}$, then

$$\begin{aligned} \|\mathbf{u} + \mathbf{v}\|^2 &= \langle \mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{v} \rangle \\ &= \|\mathbf{u}\|^2 + \|\mathbf{u}\|^2 + \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{v}, \mathbf{u} \rangle \\ &= \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 \end{aligned}$$

□




Figure 1: Orthogonal decomposition of $\mathbf{u}$ as a scalar multiple of $\mathbf{v}$ plus vector $\mathbf{w}$ orthogonal to $\mathbf{v}$

Definition 1.2.67. To compute the *orthogonal decomposition* of vector $\mathbf{u}$, let $a \in \mathbb{F}$, then

$$\mathbf{u} = a\mathbf{v} + (\mathbf{u} - a\mathbf{v})$$

To choose a so that $\mathbf{v}$ is orthogonal to $(\mathbf{u} - a\mathbf{v})$, which is

$$0 = \langle \mathbf{u} - a\mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle - a \|\mathbf{v}\|^2$$

Need to choose a to be $\langle \mathbf{u}, \mathbf{v} \rangle / \|\mathbf{v}\|^2$, where $\mathbf{v} \neq \mathbf{0}$, then

$$\mathbf{u} = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{v}\|^2}\mathbf{v} + \left(\mathbf{u} - \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\|^2}\mathbf{v} \right)$$

Theorem 1.2.68. *Cauchy-Schwarz Inequality*
If $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, *then* $|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \, \|\mathbf{u}\|$.
This inequality is an equality if and only if one of $\mathbf{u}, \mathbf{v}$ *is a scalar multiple of the other.*

Proof. Let $\mathbf{u}, \mathbf{v} \in \mathcal{V}$. If $\mathbf{v} = \mathbf{0}$, then inequality holds.
Assume $\mathbf{u} \neq \mathbf{0}$. Consider orthogonal decomposition

$$\mathbf{u} = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{v}\|^2} \mathbf{v} + \mathbf{w}$$

where $\mathbf{w}$ is orthogonal to $\mathbf{v}$. By Pythagorean Theorem 1.2.66,

$$\begin{aligned} \|\mathbf{u}\|^2 &= \left\| \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{v}\|^2} \mathbf{v} \right\|^2 + \|\mathbf{w}\|^2 \\ &= \frac{|\langle \mathbf{u}, \mathbf{v} \rangle|^2}{\|\mathbf{v}\|^2} + \|\mathbf{w}\|^2 \\ &\geq \frac{|\langle \mathbf{u}, \mathbf{v} \rangle|^2}{\|\mathbf{v}\|^2} \end{aligned}$$

Multiplying both sides by $\|\mathbf{v}\|^2$ then taking square roots give the desired inequality.
Note that this is an equality if and only if $\mathbf{w} = \mathbf{0}$, but $\mathbf{w} = \mathbf{0}$ if and only if $\mathbf{u}$ is a multiple of $\mathbf{v}$. □

The next result is on triangle inequality, which states the length of any side of a triangle is less than the sum of length of the other two sides.

Theorem 1.2.69. *Triangle Inequality*
If $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, *then* $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$.
This inequality is an equality if and only if one of $\mathbf{u}, \mathbf{v}$ *is a nonnegative multiple of the other.*

Proof. Let $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, then

$$\begin{aligned} \|\mathbf{u} + \mathbf{v}\|^2 &= \langle \mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{v} \rangle \\ &= \langle \mathbf{u}, \mathbf{u} \rangle + \langle \mathbf{v}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{v}, \mathbf{u} \rangle \\ &= \langle \mathbf{u}, \mathbf{u} \rangle + \langle \mathbf{v}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{v} \rangle + \overline{\langle \mathbf{u}, \mathbf{v} \rangle} \\ &= \|\mathbf{u}\|^2 + \|\mathbf{u}\|^2 + 2\mathrm{Re} \, \langle \mathbf{u}, \mathbf{v} \rangle \\ &\leq \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2 \, |\langle \mathbf{u}, \mathbf{v} \rangle| \\ &\leq \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2 \, \|\mathbf{u}\| \, \|\mathbf{v}\| \qquad \text{by Cauchy-Schwarz Inequality Theorem 1.2.68} \\ &= (\|\mathbf{u}\| + \|\mathbf{v}\|)^2 \end{aligned}$$

Taking square roots on both sides gives the triangle inequality.
Note this is an equality if and only if $\langle \mathbf{u}, \mathbf{v} \rangle = \|\mathbf{u}\| \, \|\mathbf{v}\|$. □

The next result is on parallelogram inequality, which states that in any parallelogram, the sum of squares of length of diagonals equals the sum of lengths of the four sides.

Theorem 1.2.70. *Parallelogram Inequality*
If $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, *then* $\|\mathbf{u} + \mathbf{v}\|^2 + \|\mathbf{u} - \mathbf{v}\|^2 = 2(\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2)$.

Proof. Let $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, then

$$\begin{aligned} \|\mathbf{u} + \mathbf{v}\|^2 + \|\mathbf{u} - \mathbf{v}\|^2 &= \langle \mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{v} \rangle + \langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle \\ &= 2 \, \|\mathbf{u}\|^2 + 2 \, \|\mathbf{v}\|^2 + \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{v}, \mathbf{u} \rangle - \langle \mathbf{u}, \mathbf{v} \rangle - \langle \mathbf{v}, \mathbf{u} \rangle \\ &= 2(\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2) \end{aligned}$$

□

Definition 1.2.71. A list of vectors $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ in $\mathcal{V}$ is *orthonormal* if

$$\langle \mathbf{e}_j, \mathbf{e}_k \rangle = \begin{cases} \langle \mathbf{e}_j, \mathbf{e}_k \rangle = 0 & \text{if } j \neq k \\ \langle \mathbf{e}_j, \mathbf{e}_k \rangle = 1 & \text{if } j = k \end{cases}$$

Proposition 1.2.72. *If $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ is an orthonormal list of vectors in $\mathcal{V}$, then*

$$\|a_1\mathbf{e}_1 + \cdots + a_m\mathbf{e}_m\|^2 = |a_1|^2 + \cdots + |a_m|^2$$

for all $a_1, \dots, a_m \in \mathbb{F}$.

Proof. As each $\mathbf{e}_j$ has norm 1, by Pythagorean Theorem 1.2.66, this holds. □

Corollary 1.2.73. *Every orthonormal list of vectors is linearly independent.*

Proof. Suppose $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ is an orthonormal list of vectors in $\mathcal{V}$ and $a_1, \dots, a_m \in \mathbb{F}$ are such that

$$a_1\mathbf{e}_1 + \cdots + a_m\mathbf{e}_m = \mathbf{0}$$

then $|a_1|^2 + \cdots + |a_m|^2 = 0$ by Proposition 1.2.72, which means all a_j are 0 as desired. □

Definition 1.2.74. An *orthonormal basis* of $\mathcal{V}$ is an orthonormal list of vectors in $\mathcal{V}$ that is also a basis.

Theorem 1.2.75. *Suppose $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ is an orthonormal basis of $\mathcal{V}$, then*

$$\mathbf{v} = \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_n \rangle \mathbf{e}_n$$
$$\|\mathbf{v}\|^2 = |\langle \mathbf{v}, \mathbf{e}_1 \rangle|^2 + \cdots + |\langle \mathbf{v}, \mathbf{e}_n \rangle|^2$$

for every $\mathbf{e} \in \mathcal{V}$.

Proof. Let $\mathbf{v} \in \mathcal{V}$. As $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ is a basis of $\mathcal{V}$, there exists scalars $a_1, \dots, a_n$ such that

$$\mathbf{v} = a_1\mathbf{e}_1 + \cdots + a_n\mathbf{e}_n$$

Take inner product of both sides of equation with $\mathbf{e}_j$ to get $\langle \mathbf{v}, \mathbf{e}_j \rangle = a_j$, hence proving. □

Theorem 1.2.76. *If $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is a linearly independent list of vectors in $\mathcal{V}$, then there exists an orthonormal list $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ of vectors in $\mathcal{V}$ such that*

$$\mathit{span}(\mathbf{v}_1, \dots, \mathbf{v}_j) = \mathit{span}(\mathbf{e}_1, \dots, \mathbf{e}_j)$$

for $j = 1, \dots, m$.

Proof. Suppose $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is linearly independent list of vectors in $\mathcal{V}$. To construct $\mathbf{e}$, let $\mathbf{e} = \mathbf{v}_1 / \|\mathbf{v}_1\|$, which satisfies condition for $j = 1$.
Suppose $j > 1$, and orthonormal list $(\mathbf{e}_1, \dots, \mathbf{e}_{j-1})$ has been chosen so that

$$\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_{j-1}) = \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_{j-1})$$

Let

$$\mathbf{e}_j = \frac{\mathbf{v}_j - \langle \mathbf{v}_j, \mathbf{e}_1 \rangle \mathbf{e}_1 - \cdots - \langle \mathbf{v}_j, \mathbf{e}_{j-1} \rangle \mathbf{e}_{j-1}}{\left\| \mathbf{v}_j - \langle \mathbf{v}_j, \mathbf{e}_1 \rangle \mathbf{e}_1 - \cdots - \langle \mathbf{v}_j, \mathbf{e}_{j-1} \rangle \mathbf{e}_{j-1} \right\|}$$

Note that $\mathbf{v}_j \notin \operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_{j-1})$ (as $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is linearly independent), hence $\mathbf{v}_j \notin \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_{j-1})$. Hence above equation is not dividing by $\mathbf{0}$, and $\mathbf{e}_j$ is well defined. Note $\|\mathbf{e}_j\| = 1$. Let $1 \le k < j$, then

$$\begin{aligned}
\langle \mathbf{e}_j, \mathbf{e}_k \rangle &= \left\langle \frac{\mathbf{v}_j - \langle \mathbf{v}_j, \mathbf{e}_1 \rangle \mathbf{e}_1 - \cdots - \langle \mathbf{v}_j, \mathbf{e}_{j-1} \rangle \mathbf{e}_{j-1}}{\left\| \mathbf{v}_j - \langle \mathbf{v}_j, \mathbf{e}_1 \rangle \mathbf{e}_1 - \cdots - \langle \mathbf{v}_j, \mathbf{e}_{j-1} \rangle \mathbf{e}_{j-1} \right\|}, \mathbf{e}_k \right\rangle \\
&= \frac{\langle \mathbf{v}_j, \mathbf{e}_k \rangle - \langle \mathbf{v}_j, \mathbf{e}_k \rangle}{\left\| \mathbf{v}_j - \langle \mathbf{v}_j, \mathbf{e}_1 \rangle \mathbf{e}_1 - \cdots - \langle \mathbf{v}_j, \mathbf{e}_{j-1} \rangle \mathbf{e}_{j-1} \right\|} \\
&= 0
\end{aligned}$$

Hence $(\mathbf{e}_1, \dots, \mathbf{e}_j)$ is an orthonormal list.
Note that $\mathbf{v}_j \in \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_j)$, and hence

$$\operatorname{span}(\mathbf{v}_1, \dots, \mathbf{v}_j) \subset \operatorname{span}(\mathbf{e}_1, \dots, \mathbf{e}_j)$$

Both lists are linearly independent, hence both spaces have dimension j, thus must be equal. □

Corollary 1.2.77. *Every finite-dimensional inner-product space has an orthonormal basis.*

Proof. Choose a basis of $\mathcal{V}$. Apply Gram-Schmidt 1.2.76 to it to produce an orthonormal list. This orthonormal list is linearly independent by Corollary 1.2.73 and its span equals $\mathcal{V}$, hence it is an orthonormal basis of $\mathcal{V}$. □

Corollary 1.2.78. *Every orthonormal list of vectors in $\mathcal{V}$ can be extended to an orthonormal basis of $\mathcal{V}$.*

Proof. Suppose $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ is an orthonormal list of vectors in $\mathcal{V}$. Then $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ is linearly independent by Corollary 1.2.73, hence it can be extended to a basis $(\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{v}_1, \ldots, \mathbf{v}_n)$ of $\mathcal{V}$. Apply Gram-Schmidt procedure 1.2.76 to $(\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{v}_1, \ldots, \mathbf{v}_n)$ of $\mathcal{V}$ to produce an orthonormal list

$$(\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{f}_1, \ldots, \mathbf{f}_n)$$

Note Gram-Schmidt leaves first m vectors unchanged as they are already orthonormal. Note that above is an orthonormal basis of $\mathcal{V}$ as it is linearly independent, and its span equals $\mathcal{V}$. Hence this gives the extension of $(\mathbf{e}_1, \ldots, \mathbf{e}_m,)$ to an orthonormal basis of $\mathcal{V}$. □

Corollary 1.2.79. *Suppose $T \in \mathcal{L}(\mathcal{V})$. If T has an upper-triangular matrix with respect to some basis of $\mathcal{V}$, then T has an upper-triangular matrix with respect to some orthonormal basis of $\mathcal{V}$.*

Proof. Suppose T has an upper-triangular matrix with respect to some basis $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ of $\mathcal{V}$. Thus $\operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$ is invariant under T for each $j = 1, \ldots, n$ by Proposition 1.2.52.
Apply Gram-Schmidt procedure 1.2.76 to $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ to produce an orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ of $\mathcal{V}$. Note that $\operatorname{span}(\mathbf{e}_1, \ldots, \mathbf{e}_j) = \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_j)$ for each j, hence $\operatorname{span}(\mathbf{e}_1, \ldots, \mathbf{e}_j)$ is invariant under T for each $j = 1, \ldots, n$. Hence by Proposition 1.2.52, T has an upper-triangular matrix with respect to some orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$. □

Corollary 1.2.80. *Suppose $\mathcal{V}$ is a complex vector space and $T \in \mathcal{L}(\mathcal{V})$, then T has an upper-triangular matrix with respect to some orthonormal basis of $\mathcal{V}$.*

Proof. This follows from Theorem 1.2.53 and Corollary 1.2.79. □

Definition 1.2.81. If $\mathcal{U}$ is a subset of $\mathcal{V}$, then the *orthogonal complement* of $\mathcal{U}$, denoted $\mathcal{U}^\perp$, is the set of all vectors in $\mathcal{V}$ that are orthogonal to every vector in $\mathcal{U}$:

$$\mathcal{U}^\perp = \{\mathbf{v} \in \mathcal{V} : \langle \mathbf{v}, \mathbf{u} \rangle = 0 \ \ \forall\, \mathbf{u} \in \mathcal{U}\}$$

Note that $\mathcal{U}^\perp \subset \mathcal{V}$, $\mathcal{V}^\perp = \{\mathbf{0}\}$, $\{\mathbf{0}\}^\perp = \mathcal{V}$. Also note that if $\mathcal{U}_1 \subset \mathcal{U}_2$, then $\mathcal{U}_1^\perp \subset \mathcal{U}_2^\perp$.

Theorem 1.2.82. *If $\mathcal{U}$ is a subspace of $\mathcal{V}$, then $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$.*

Proof. Suppose $\mathcal{U}$ is a subspace of $\mathcal{V}$.
First show that $\mathcal{V} = \mathcal{U} + \mathcal{U}^\perp$. Suppose $\mathbf{v} \in \mathcal{V}$, let $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ be an orthonormal basis of $\mathcal{U}$. Note that

$$\begin{aligned} \mathbf{v} &= \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_m \rangle \mathbf{e}_m + \mathbf{v} - \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{e}_1 - \cdots - \langle \mathbf{v}, \mathbf{e}_m \rangle \mathbf{e}_m \\ &= \mathbf{u} + \mathbf{w} \end{aligned}$$

Clearly $\mathbf{u} \in \mathcal{U}$. As $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ is an orthonormal list, for each j, note that

$$\langle \mathbf{w}, \mathbf{e}_j \rangle = \langle v, \mathbf{e}_j \rangle - \langle \mathbf{v}, \mathbf{e}_j \rangle = 0$$

Hence $\mathbf{w}$ is orthogonal to every vector in $\operatorname{span}(\mathbf{e}_1, \ldots, \mathbf{e}_m)$. Hence $\mathbf{w} \in \mathcal{U}^\perp$, hence $\mathbf{v} = \mathbf{u} + \mathbf{w}$, where $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{U}^\perp$, completing the proof for equation $\mathcal{V} = \mathcal{U} + \mathcal{U}^\perp$.
If $\mathbf{v} \in \mathcal{U} \cap \mathcal{U}^\perp$, then $\mathbf{v}$ is orthogonal to every vector in $\mathcal{U}$, which implies $(\mathbf{v}, \mathbf{v}) = 0$, which implies $\mathbf{v} = \mathbf{0}$. Hence $\mathcal{U} \cap \mathcal{U}^\perp = \{\mathbf{0}\}$. Note that $\mathcal{V} = \mathcal{U} + \mathcal{U}^\perp$ and above equation implies $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$. □

Corollary 1.2.83. *If $\mathcal{U} \subset \mathcal{V}$, then $\mathcal{U} = (\mathcal{U}^\perp)^\perp$.*

Proof. Suppose $\mathcal{U} \subset \mathcal{V}$.
First to show $\mathcal{U} \subset (\mathcal{U}^\perp)^\perp$. Let $\mathbf{u} \in \mathcal{U}$. Then $\langle \mathbf{u}, \mathbf{v} \rangle = 0$ for every $\mathbf{v} \in \mathcal{U}^\perp$ by definition of $\mathcal{U}^\perp$. As $\mathbf{u}$ is orthogonal to every vector in $\mathcal{U}^\perp$, then we have $\mathbf{u} \in (\mathcal{U}^\perp)^\perp$, completing the proof.
Next to show $(\mathcal{U}^\perp)^\perp \subset \mathcal{U}$, suppose $\mathbf{v} \in (\mathcal{U}^\perp)^\perp$. By Theorem 1.2.82, $\mathbf{v} = \mathbf{u} + \mathbf{w}$, where $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{U}^\perp$. Then $\mathbf{v} - \mathbf{u} = \mathbf{w} \in \mathcal{U}^\perp$. As $\mathbf{v} \in (\mathcal{U}^\perp)^\perp$ and $\mathbf{u} \in (\mathcal{U}^\perp)^\perp$, then $\mathbf{v} - \mathbf{u} \in (\mathcal{U}^\perp)^\perp$, Thus $\mathbf{v} - \mathbf{u} \in \mathcal{U}^\perp \cap (\mathcal{U}^\perp)^\perp$, which implies $\mathbf{v} - \mathbf{u}$ is orthogonal to itself, which implies $\mathbf{v} - \mathbf{u} = \mathbf{0}$, hence $\mathbf{v} = \mathbf{u} \in \mathcal{U}$. Hence $(\mathcal{U}^\perp)^\perp \subset \mathcal{U}$. □

Definition 1.2.84. Let $\mathcal{U} \subset \mathcal{V}$. Decomposition $\mathcal{V} = \mathcal{U} \oplus \mathcal{U}^\perp$ means each vector $\mathbf{v} \in \mathbf{V}$ can be written as $\mathbf{v} = \mathbf{u} + \mathbf{w}$, where $\mathbf{u} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{U}^\perp$.
The *orthogonal projection* of $\mathcal{V}$ onto $\mathcal{U}$ is an operator on $\mathcal{V}$, denoted $P_\mathcal{U} = P_{\mathcal{U},\mathcal{U}^\perp}$.
Note that for $\mathbf{v} \in \mathcal{V}$, denote $P_\mathcal{U}\mathbf{v}$ to be vector $\mathbf{u}$ in decomposition $\mathbf{v} = \mathbf{u} + \mathbf{w}$.

Remark 1.2.85. *Properties of Orthogonal Projection* of $P_\mathcal{U} \in \mathcal{L}(\mathcal{V})$:

i. $\text{range} P_\mathcal{U} = \mathcal{U}$
ii. $\text{null} P_\mathcal{U} = \mathcal{U}^\perp$
iii. $\mathbf{v} - P_\mathcal{U}\mathbf{v} \in \mathcal{U}^\perp$ for every $\mathbf{v} \in \mathcal{V}$
iv. $P_\mathcal{U}^2 = P_\mathcal{U}$
v. $\|P_\mathcal{U}\mathbf{v}\| \leq \|\mathbf{v}\|$ for every $\mathbf{v} \in \mathcal{V}$
vi. If $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ is an orthonormal basis of $\mathcal{U}$, then $P_\mathcal{U}\mathbf{v} = \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_m \rangle \mathbf{v}_m$ for every $\mathbf{v} \in \mathcal{V}$

Proposition 1.2.86. *Suppose $\mathcal{U} \subset \mathcal{V}$ and $\mathbf{v} \in \mathcal{V}$. Then*

$$\|\mathbf{v} - P_\mathcal{U}\mathbf{v}\| \leq \|\mathbf{v} - \mathbf{u}\|$$

for every $\mathbf{v} \in \mathcal{U}$. Furthermore, if $\mathbf{v} \in \mathcal{U}$ and the inequality above is an equality, then $\mathbf{v} = P_\mathcal{U}\mathbf{v}$.

Proof. Suppose $\mathbf{u} \in \mathcal{U}$, then

$$\begin{aligned} \|\mathbf{v} - P_\mathcal{U}\mathbf{v}\| &\leq \|\mathbf{v} - P_\mathcal{U}\mathbf{v}\|^2 + \|P_\mathcal{U}\mathbf{v} - \mathbf{u}\|^2 \\ &= \|(\mathbf{v} - P_\mathcal{U}\mathbf{v}) + (P_\mathcal{U}\mathbf{v} - \mathbf{u})\|^2 \quad \text{by Pythagorean Theorem 1.2.66} \\ &= \|\mathbf{v} - \mathbf{u}\|^2 \end{aligned}$$

Note that for equality case, this happens if and only if $\|P_\mathcal{U}\mathbf{v} - \mathbf{u}\| = 0$, hence $\mathbf{u} = P_\mathcal{U}\mathbf{v}$. □

Example 1.2.87. Approximate $\sin(x)$ on $[-\pi, \pi]$ with basis $(1, x, x^2, x^3, x^4, x^5)$ of $\mathcal{U}$.

Proof. Let

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f(x)g(x)\, dx, \qquad U_{\text{odd}} = \text{span}\{x, x^3, x^5\}$$

Then

$$\begin{aligned} p_1 &= x, & p_2 &= x^3 - \frac{3\pi^2}{5}x, & p_3 &= x^5 - \frac{10\pi^2}{9}x^3 + \frac{5\pi^4}{21}x \\ \|p_1\|^2 &= \frac{2\pi^3}{3}, & \|p_2\|^2 &= \frac{8\pi^7}{175}, & \|p_3\|^2 &= \frac{128\pi^{11}}{43659} \\ \langle \sin x, p_1 \rangle &= 2\pi, & \langle \sin x, p_2 \rangle &= \frac{4\pi}{5}(\pi^2 - 15), & \langle \sin x, p_3 \rangle &= \frac{16\pi}{63}(\pi^4 - 105\pi^2 + 945) \end{aligned}$$

Hence

$$u = \sum_{k=1}^{3} \frac{\langle \sin x, p_k \rangle}{\langle p_k, p_k \rangle} p_k = Ax + Bx^3 + Cx^5,$$

$$A = \frac{105(-153\pi^2 + \pi^4 + 1485)}{8\pi^6}, \quad B = \frac{315(-1155 - \pi^4 + 125\pi^2)}{4\pi^8}, \quad C = \frac{693(-105\pi^2 + \pi^4 + 945)}{8\pi^{10}}.$$

□

Definition 1.2.88. A *linear functional* on $\mathcal{V}$ is a linear map from $\mathcal{V}$ to the scalars $\mathbb{F}$.

Theorem 1.2.89. *Suppose φ is a linear functional on $\mathcal{V}$. Then there is a unique vector $\mathbf{v} \in \mathcal{V}$ such that*

$$\varphi(\mathbf{u}) = \langle \mathbf{u}, \mathbf{v} \rangle$$

for every $\mathbf{u} \in \mathcal{V}$.

Proof. First, to show $\exists$ a vector $\mathbf{v} \in \mathcal{V}$ such that $\varphi(\mathbf{u}) = \langle \mathbf{u}, \mathbf{v} \rangle$ for every $\mathbf{u} \in \mathcal{V}$.
Let $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be an orthonormal basis of $\mathcal{V}$, then

$$\begin{aligned} \varphi\mathbf{u} &= \varphi(\langle \mathbf{u}, \mathbf{e}_1 \rangle \, \mathbf{e}_1 + \cdots + \langle \mathbf{u}, \mathbf{e}_n \rangle \, \mathbf{e}_n) \quad \text{by Theorem 1.2.75} \\ &= \langle \mathbf{u}, \mathbf{e}_1 \rangle \, \varphi(\mathbf{e}_1) + \cdots + \langle \mathbf{u}, \mathbf{e}_n \rangle \, \varphi(\mathbf{e}_n) \\ &= \left\langle \mathbf{u}, \overline{\varphi(\mathbf{e}_1)}\mathbf{e}_1 + \cdots + \overline{\varphi(\mathbf{e}_n)}\mathbf{e}_n \right\rangle \end{aligned}$$

for every $\mathbf{u} \in \mathcal{V}$. Setting $\mathbf{v} = \overline{\varphi(\mathbf{e}_1)}\mathbf{e}_1 + \cdots + \overline{\varphi(\mathbf{e}_n)}\mathbf{e}_n$, then $\varphi(\mathbf{u}) = \langle \mathbf{u}, \mathbf{v} \rangle$ for every $\mathbf{v} \in \mathcal{V}$ as desired.
Next, to prove that only one vector $\mathbf{v} \in \mathcal{V}$ has the desired behaviour. Suppose $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}$, such that

$$\varphi(\mathbf{u}) = \langle \mathbf{u}, \mathbf{v}_1 \rangle = \langle \mathbf{u}, \mathbf{v}_2 \rangle$$

for every $\mathbf{u} \in \mathcal{V}$. Taking $\mathbf{u} = \mathbf{v}_1 - \mathbf{v}_2$ shows $\mathbf{v}_1 - \mathbf{v}_2 = \mathbf{0}$, hence $\mathbf{v}_1 = \mathbf{v}_2$, proving uniqueness. □

Definition 1.2.90. Let $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, where $\mathcal{W}, \mathcal{V}$ are finite-dimensional nonzero, inner-product space over $\mathbb{F}$. Fix $\mathbf{w} \in \mathcal{W}$, and consider the linear functional $f_{\mathbf{w}} : \mathcal{V} \to \mathbb{F}$ given by $f_{\mathbf{w}}(\mathbf{v}) = \langle T\mathbf{v}, \mathbf{w} \rangle$. There is a unique vector $\mathbf{y} \in \mathcal{V}$ such that $f_{\mathbf{w}}(\mathbf{v}) = \langle \mathbf{v}, \mathbf{y} \rangle \ \forall \ \mathbf{v} \in \mathcal{V}$, and let this unique vector $\mathbf{y}$ be $T^*\mathbf{w}$. Hence

$$\langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, T^*\mathbf{w} \rangle$$

for all $\mathbf{v} \in \mathcal{V}$. Doing this for every $\mathbf{w} \in \mathcal{W}$ gives a map $T^* : \mathcal{W} \to \mathcal{V}$, which is the *adjoint* of T.

Remark 1.2.91. *Property of Adjoints*

i. Additivity: $(S + T)^* = S^* + T^*$ for all $S, T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$
ii. Conjugate Homogeneity: $(aT)^* = \overline{a}T^*$ for all $a \in \mathbb{F}$ and $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$
iii. Adjoint of Adjoint: $(T^*)^* = T$ for all $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$
iv. Identity: $I^* = I$, where I is the identity operator on $\mathcal{V}$
v. Products: $(ST^*) = T^*S^*$ for all $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$ and $S \in \mathcal{L}(\mathcal{W}, \mathcal{U})$ ($\mathcal{U}$ is inner-product space over $\mathbb{F}$)

Next result shows relationship between null space and range of a linear map and its adjoint.

Proposition 1.2.92. *Suppose $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$, then*

i. $null(T^*) = (range(T))^{\perp}$
ii. $range(T^*) = (null()T)^{\perp}$
iii. $null(T) = (range(T^*))^{\perp}$
iv. $range(T) = (null(T^*))^{\perp}$

Proof. For (i), let $\mathbf{w} \in \mathcal{W}$, then

$$\mathbf{w} \in \text{null}(T^*) \Leftrightarrow T^*\mathbf{w} = \mathbf{0} \Leftrightarrow \langle \mathbf{v}, T^*\mathbf{w} \rangle = \mathbf{0} \ \forall \ \mathbf{v} \in \mathcal{V} \Leftrightarrow \langle T\mathbf{v}, \mathbf{w} \rangle = \mathbf{0} \ \forall \ \mathbf{v} \in \mathcal{V} \Leftrightarrow \mathbf{w} \in \text{range(T)}^{\perp}$$

Take orthogonal complement of both sides of (i) to get (iv).
Replace T with T^* in (i) and (iv) to get (iii) and (ii). □

Definition 1.2.93. The *conjugate transpose* of an $m \times n$ matrix is obtained by interchanging rows and columns, then taking the complex conjugate of each entry.

Proposition 1.2.94. *Suppose $T \in \mathcal{L}(\mathcal{V}, \mathcal{W})$. If $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ is an orthonormal basis of $\mathcal{V}$ and $(\mathbf{f}_1, \ldots, \mathbf{f}_m)$ is an orthonormal basis of $\mathcal{W}$, then*

$$\mathcal{M}(T^*, (\mathbf{f}_1, \ldots, \mathbf{f}_m), (\mathbf{e}_1, \ldots, \mathbf{e}_n))$$

is the conjugate transpose of

$$\mathcal{M}(T, (\mathbf{e}_1, \ldots, \mathbf{e}_n), (\mathbf{f}_1, \ldots, \mathbf{f}_m))$$

Proof. Let $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be orthonormal basis of $\mathcal{V}$ and $(\mathbf{f}_1, \ldots, \mathbf{f}_m)$ be orthonormal basis of $\mathcal{W}$. The k column of $\mathcal{M}(T)$ is obtained by writing $T\mathbf{e}_k$ as a linear combination of $\mathbf{f}_j$, and the scalars used in this linear combination

becomes the k column of $\mathcal{M}(T)$. As $(\mathbf{f}_1, \ldots, \mathbf{f}_m)$ is an orthonormal basis of $\mathcal{W}$, $T\mathbf{e}_k$ can be written as a linear combination of the $\mathbf{f}_j$.

$$T\mathbf{e}_k = \langle T\mathbf{e}_k, \mathbf{f}_1 \rangle \, \mathbf{f}_1 + \cdots + \langle T\mathbf{e}_k, \mathbf{f}_m \rangle \, \mathbf{f}_m$$

Hence the entry in row j column k of $\mathcal{M}(T)$ is $\langle T\mathbf{e}_k, \mathbf{f}_j \rangle$. Replace T with T^* and interchanging the roles played by $\mathbf{e}_j$ and $\mathbf{f}_j$, then the entry in row j column k of $\mathcal{K}(T^*)$ is $\langle T^*\mathbf{f}_k, \mathbf{e}_j \rangle = \langle \mathbf{f}_k, T\mathbf{e}_j \rangle = \overline{\langle T\mathbf{e}_j, \mathbf{f}_k \rangle}$, which equals the complex conjugate of entry in row k column j of $\mathcal{M}(T)$. Hence $\mathcal{M}(T^*)$ equals conjugate transpose of $\mathcal{M}(T)$. □

1.2.5 Operators on Inner-Product Spaces

Definition 1.2.95. An operator $T \in \mathcal{L}(\mathcal{V})$ is *self-adjoint* or *Hermitian* if $T = T^*$.

A self-adjoint operator is analogous to a real number. It plays similar role to complex conjugation on $\mathbb{C}$.

Proposition 1.2.96. *Every eigenvalue of a self-adjoint operator is real.*

Proof. Let $T \in \mathcal{L}(\mathcal{V})$. Let λ be an eigenvalue of T, $\mathbf{v}$ be nonzero vector in $\mathcal{V}$ such that $T\mathbf{v} = \lambda\mathbf{v}$. Then

$$\lambda \|\mathbf{v}\|^2 = \langle \lambda\mathbf{v}, \mathbf{v} \rangle = \langle T\mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{v}, T\mathbf{v} \rangle = \langle \mathbf{v}, \lambda, \mathbf{v} \rangle = \overline{\lambda} \|\mathbf{v}\|^2$$

Note that $\lambda = \overline{\lambda}$, hence λ is real. □

Proposition 1.2.97. *If $\mathcal{V}$ is a complex inner-product space and T is an operator on $\mathcal{V}$ such that $\langle T\mathbf{v}, \mathbf{v} \rangle = 0$ for all $\mathbf{v} \in \mathcal{V}$, then $T = 0$.*

Proof. Suppose $\mathcal{V}$ is a complex inner-product space and $T \in \mathcal{L}(\mathcal{V})$, then

$$\langle T\mathbf{u}, \mathbf{w} \rangle = \frac{\langle T(\mathbf{u}+\mathbf{w}), \mathbf{u}+\mathbf{w} \rangle - \langle T(\mathbf{u}-\mathbf{w}), \mathbf{u}-\mathbf{w} \rangle}{4} + \frac{\langle T(\mathbf{u}+i\mathbf{w}), \mathbf{u}+i\mathbf{w} \rangle - \langle T(\mathbf{u}-i\mathbf{w}), \mathbf{u}-i\mathbf{w} \rangle}{4} i$$

for all $\mathbf{u}, \mathbf{w} \in \mathcal{V}$. Note each term on right side is of form $\langle T\mathbf{v}, \mathbf{v} \rangle$ for appropriate $\mathbf{v} \in \mathcal{V}$. If $\langle T\mathbf{v}, \mathbf{v} \rangle = 0$ for all $\mathbf{v} \in \mathcal{V}$, then equation above implies $\langle T\mathbf{u}, \mathbf{w} \rangle = 0$ for all $\mathbf{u}, \mathbf{w} \in \mathcal{V}$. This implies $T = 0$ (take $\mathbf{w} = T\mathbf{u}$). □

Corollary 1.2.98. *Let $\mathcal{V}$ be complex inner-product space and let $T \in \mathcal{L}(\mathcal{V})$. Then T is self-adjoint if and only if $\langle T\mathbf{v}, \mathbf{v} \rangle \in \mathbb{R}$ for every $\mathbf{v} \in \mathcal{V}$.*

Proof. Let $\mathbf{v} \in \mathcal{V}$. Then

$$\langle T\mathbf{v}, \mathbf{v} \rangle - \overline{\langle T\mathbf{v}, \mathbf{v} \rangle} = \langle T\mathbf{v}, \mathbf{v} \rangle - \langle \mathbf{v}, T\mathbf{v} \rangle = \langle T\mathbf{v}, \mathbf{v} \rangle - \langle T^*\mathbf{v}, \mathbf{v} \rangle = \langle (T - T^*)\mathbf{v}, \mathbf{v} \rangle$$

If $\langle T\mathbf{v}, \mathbf{v} \rangle \in \mathbb{R}$ for every $\mathbf{v} \in \mathcal{V}$, then the left-side of equation equals 0, hence $\langle (T - T^*)\mathbf{v}, \mathbf{v} \rangle = 0$ for every $\mathbf{v} \in \mathcal{V}$. This implies $T - T^* = 0$ by Proposition 1.2.97, hence T is self-adjoint.
Conversely, if T is self-adjoint, then the right-side of the equation above equals 0, hence $\langle T\mathbf{v}, \mathbf{v} \rangle = \overline{\langle T\mathbf{v}, \mathbf{v} \rangle}$ for every $\mathbf{v} \in \mathcal{V}$. This implies $\langle T\mathbf{v}, \mathbf{v} \rangle \in \mathbb{R}$ for every $\mathbf{v} \in \mathcal{V}$ as desired. □

Proposition 1.2.99. *If T is a self-adjoint operator on $\mathcal{V}$ such that $\langle T\mathbf{v}, \mathbf{v} \rangle = 0$ for all $\mathbf{v} \in \mathcal{V}$, then $T = 0$.*

Proof. Let $\mathcal{V}$ be a real inner-product space, and T a self-adjoint operator on $\mathcal{V}$. For $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, we have

$$\langle T\mathbf{u}, \mathbf{v} \rangle = \frac{\langle T(\mathbf{u}+\mathbf{w}), \mathbf{u}+\mathbf{w} \rangle - \langle T(\mathbf{u}-\mathbf{w}), \mathbf{u}-\mathbf{w} \rangle}{4}$$

Note that

$$\begin{aligned} \langle T\mathbf{w}, \mathbf{u} \rangle &= \langle \mathbf{w}, T\mathbf{u} \rangle && \text{as } T \text{ is self-adjoint} \\ &= \langle T\mathbf{u}, \mathbf{w} \rangle && \text{as this is real inner-product space} \end{aligned}$$

If $\langle T\mathbf{v}, \mathbf{v} \rangle = 0$ for all $\mathbf{v} \in \mathcal{V}$, then first equation implies $\langle T\mathbf{u}, \mathbf{w} \rangle = 0$ for all $\mathbf{u}, \mathbf{w} \in \mathcal{V}$.
This implies $T = 0$ (taking $\mathbf{w} = T\mathbf{u}$). □

Definition 1.2.100. An operator on an inner-product space is *normal* if it commutes with its adjoint, i.e., $T \in \mathcal{L}(\mathcal{V})$ is normal if $TT^* = T^*T$. Every self-adjoint operator is normal.

The next proposition implies $\text{null}(T) = \text{null}(T^*)$ for every normal operator T.

Proposition 1.2.101. *An operator $T \in \mathcal{L}(\mathcal{V})$ is normal if and only if $\|T\mathbf{v}\| = \|T^*\mathbf{v}\|$ for all $\mathbf{v} \in \mathcal{V}$.*

Proof. Let $T \in \mathcal{L}(\mathcal{V})$. Note that

$$\begin{aligned} T \text{ is normal} &\Leftrightarrow T^*T - TT^* = 0 \\ &\Leftrightarrow \langle (T^*T - TT^*)\mathbf{v}, \mathbf{v}\rangle = 0 \quad \text{for all } \mathbf{v} \in \mathcal{V} \quad \text{(by Proposition 1.2.99)} \\ &\Leftrightarrow \langle T^*T\mathbf{v}, \mathbf{v}\rangle = \langle TT^*\mathbf{v}, \mathbf{v}\rangle \quad \text{for all } \mathbf{v} \in \mathcal{V} \\ &\Leftrightarrow \|T\mathbf{v}\|^2 = \|T^*\mathbf{v}\|^2 \quad \text{for all } \mathbf{v} \in \mathcal{V} \end{aligned}$$

hence proving the proposition. □

The next corollary implies that a normal operator and its adjoint have the same eigenvectors.

Corollary 1.2.102. *Suppose $T \in \mathcal{L}(\mathcal{V})$ is normal. If $\mathbf{v} \in \mathcal{V}$ is an eigenvector of T with eigenvalue $\lambda \in \mathbb{F}$, then $\mathbf{v}$ is also an eigenvector of T^* with eigenvalue $\overline{\lambda}$.*

Proof. Suppose $\mathbf{v} \in \mathcal{V}$ is an eigenvector of T with eigenvalue λ, hence $(T - \lambda I)\mathbf{v} = \mathbf{0}$. As T is normal, so is $T - \lambda I$. By Proposition 1.2.101,

$$0 = \|(T - \lambda I)\mathbf{v}\| = \|(T - \lambda I)^*\mathbf{v}\| = \left\|(T^* - \overline{\lambda} I)\mathbf{v}\right\|$$

hence $\mathbf{v}$ is an eigenvector of T^* with eigenvalue $\overline{\lambda}$ as desired. □

Corollary 1.2.103. *If $T \in \mathcal{L}(\mathcal{V})$ is normal, eigenvectors of T with unique eigenvalues are orthogonal.*

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$ is normal, and α, β are distinct eigenvalues of T, with corresponding eigenvectors $\mathbf{u}, \mathbf{v}$. Thus $T\mathbf{u} = \alpha\mathbf{u}$ and $T\mathbf{v} = \beta\mathbf{v}$. By Corollary 1.2.102, $T^*\mathbf{v} = \overline{\beta}\mathbf{v}$, hence

$$\begin{aligned} (\alpha - \beta)\langle \mathbf{u}, \mathbf{v}\rangle &= \langle \alpha\mathbf{u}, \mathbf{v}\rangle - \langle \mathbf{u}, \overline{\beta}\mathbf{v}\rangle \\ &= \langle T\mathbf{u}, \mathbf{v}\rangle - \langle \mathbf{u}, T^*\mathbf{v}\rangle = 0 \end{aligned}$$

As $\alpha \neq \beta$, equation implies $\langle \mathbf{u}, \mathbf{v}\rangle = 0$, hence $\mathbf{u}$ and $\mathbf{v}$ are orthogonal as desired. □

The complex spectral theorem implies that every self-adjoint operator on a finite dimensional complex inner-product space has a diagonal matrix with respect to some orthonormal basis.

Theorem 1.2.104. *Complex Spectral Theorem*
Suppose $\mathcal{V}$ is a complex inner-product space and $T \in \mathcal{L}(\mathcal{V})$. Then $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of T if and only if T is normal.

Proof. ($\Rightarrow$) First, suppose $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of T, and T has a diagonal matrix with respect to this basis. The matrix of T^* is from taking the conjugate transpose of T, hence T^* also has a diagonal matrix. Any two diagonal matrices commute, hence T commutes with T^*, and T is normal.
($\Leftarrow$) Suppose T is normal. There is an orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ of $\mathcal{V}$ with respect to which T has an upper-triangular matrix by Corollary 1.2.80, hence

$$\mathcal{M}(T, (\mathbf{e}_1, \ldots, \mathbf{e}_n)) = \begin{bmatrix} a_{1,1} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots \\ 0 & \cdots & a_{n,n} \end{bmatrix}$$

Note that from the matrix, $\|T\mathbf{e}_1\|^2 = |a_{1,1}|^2$, and $\|T^*\mathbf{e}_1\|^2 = |a_{1,1}|^2 + |a_{1,2}|^2 + \cdots + |a_{1,n}|^2$.
As T is normal, $\|T\mathbf{e}_1\| = \|T^*\mathbf{e}_1\|$ by Proposition 1.2.101, hence above equations imply all entires in first row of matrix except possibly first entry $a_{1,1}$ is equal 0.
From matrix, note that $\|T\mathbf{e}_2\|^2 = |a_{2,2}|^2$ (as $a_{1,2} = 0$ is shown above), $\|T^*\mathbf{e}_2\|^2 = |a_{2,2}|^2 + |a_{2,3}|^2 + \cdots + |a_{2,n}|^2$.
As T is normal, $\|T\mathbf{e}_2\| = \|T^*\mathbf{e}_2\|$, hence above equations imply all entries in second row of matrix except possibly the diagonal entry $a_{2,2}$ is equal 0.
Doing this for all non-diagonal entries show that these equal 0 as desired. □

Note that for $\alpha, \beta \in \mathbb{R}$, and $\alpha^2 < 4\beta$, let $x \in \mathbb{R}$, then $x^2 + \alpha x + \beta = \left(x + \frac{\alpha}{2}\right)^2 + \left(\beta - \frac{\alpha^2}{4}\right) > 0$, hence $x^2 + \alpha x + \beta$ is an invertible real number. Replacing x with a self-adjoint operator leads to the lemma below.

Lemma 1.2.105. *Suppose $T \in \mathcal{L}(\mathcal{V})$ is self-adjoint. If $\alpha, \beta \in \mathbb{R}$ are such that $\alpha^2 < 4\beta$, then*

$$T^2 + \alpha T + \beta I$$

is invertible.

Proof. Suppose $\alpha, \beta \in \mathbb{R}$ are such that $\alpha^2 < 4\beta$. Let $\mathbf{v}$ be nonzero vector in $\mathcal{V}$, then

$$\begin{aligned}\left\langle (T^2 + \alpha T + \beta I)\mathbf{v}, \mathbf{v} \right\rangle &= \left\langle T^2\mathbf{v}, \mathbf{v} \right\rangle + \alpha \left\langle T\mathbf{v}, \mathbf{v} \right\rangle + \beta \left\langle \mathbf{v}, \mathbf{v} \right\rangle \\ &= \left\langle T\mathbf{v}, T\mathbf{v} \right\rangle + \alpha \left\langle T\mathbf{v}, \mathbf{v} \right\rangle + \beta \left\| \mathbf{v} \right\|^2 \\ &\geq \left\| T\mathbf{v} \right\|^2 - |\alpha| \left\| T\mathbf{v} \right\| \left\| \mathbf{v} \right\| + \beta \left\| \mathbf{v} \right\|^2 \quad \text{by Cauchy-Schwarz Inequality} \\ &= \left(\left\| T\mathbf{v} \right\| - \frac{|\alpha| \left\| \mathbf{v} \right\|}{2} \right)^2 + \left(\beta - \frac{\alpha^2}{4} \right) \left\| \mathbf{v} \right\|^2 > 0\end{aligned}$$

Note last inequality implies $(T^2+\alpha T+\beta I)\mathbf{v} \neq 0$, hence $T^2+\alpha T+\beta I$ is injective, which implies it is invertible. □

Lemma 1.2.106. *Suppose $T \in \mathcal{L}(\mathcal{V})$ is self-adjoint, then T has an eigenvalue.*

Proof. Assume $\mathcal{V}$ is a real inner-product space. Let $\dim(\mathcal{V}) = n$, $\mathbf{v} \in \mathcal{V}$ with $\mathbf{v} \neq \mathbf{0}$, then

$$(\mathbf{v}, T\mathbf{v}, T^2\mathbf{v}, \ldots, T^n\mathbf{v})$$

is not linearly independent as $\dim(\mathcal{V}) = n$ and there are $n+1$ vectors. Hence $\exists\ a_0, \ldots, a_n \in \mathbb{R}$ not all 0 s.t.

$$0 = a_0\mathbf{v} + a_1 T\mathbf{v} + \cdots + a_n T^n\mathbf{v}$$

Let a be coefficients of a polynomial, written in factored form as

$$a_0 + a_1 x + \cdots + a_n x^n = c(x^2 + \alpha_1 x + \beta_1) \cdots (x^2 + \alpha_M x + \beta_M)(x - \lambda_1) \cdots (x - \lambda_m)$$

where $c \neq 0 \in \mathbb{R}, \alpha_j, \beta_j, \lambda_j \in \mathbb{R}$, each $\alpha_j^2 < 4\beta_j$, $m + M \geq 1$, and equation holds for all $x \in \mathbb{R}$. Then

$$\begin{aligned}0 &= a_0\mathbf{v} + a_1 T\mathbf{v} + \cdots + a_n T^n\mathbf{v} \\ &= (a_0 I + a_1 T + \cdots + a_n T^n)\mathbf{v} \\ &= c(T^2 + \alpha_1 T + \beta_1 I) \cdots (T^2 + \alpha_M T + \beta_M I)(T - \lambda_1 I) \cdots (T - \lambda_m I)\mathbf{v}\end{aligned}$$

Note each $T^2 + \alpha_j T + \beta_j I$ is invertible as T is self-adjoint and each $\alpha_j^2 < 4\beta_j$ as by Lemma 1.2.105.
Also $c \neq 0$, hence equation implies

$$0 = (T - \lambda_1 T) \cdots (T - \lambda_m I)\mathbf{v}$$

Hence $T - \lambda_j I$ is not injective for at least one j, thus T has an eigenvalue. □

Theorem 1.2.107. *Real Spectral Theorem*
Suppose $\mathcal{V}$ is a real inner-product space and $T \in \mathcal{L}(\mathcal{V})$, then $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of T if and only if T is self-adjoint.

Proof. ($\Rightarrow$) Suppose $\mathcal{V}$ has an orthonormal basis consisting of eigenvectors of T. With respect to this basis, T has a diagonal matrix which equals its conjugate transpose, $T = T^*$, hence T is self-adjoint.
($\Leftarrow$) Suppose T is self-adjoint. Note $\mathcal{V}$ has an orthonormal basis clearly if $\dim(\mathcal{V}) = 1$.
Assume $\dim(\mathcal{V}) > 1$, and λ be any eigenvalue of T, let $\mathbf{u} \in \mathcal{V}$ denote corresponding eigenvector with $\|\mathbf{u}\| = 1$. Let $\mathcal{U}$ be one-dimensional subspace of $\mathcal{V}$ consisting of all scalar multiples of $\mathbf{u}$. Note $\mathbf{v} \in \mathcal{V}$ is in $\mathcal{U}^\perp$ if and only if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$. Suppose $\mathbf{v} \in \mathcal{U}^\perp$, then T is self-adjoint, and

$$\langle \mathbf{u}, T\mathbf{v} \rangle = \langle T\mathbf{u}, \mathbf{v} \rangle = \langle \lambda\mathbf{u}, \mathbf{v} \rangle = \lambda \langle \mathbf{u}, \mathbf{v} \rangle = 0$$

hence $T\mathbf{v} \in \mathcal{U}^\perp$. Thus $T\mathbf{v} \in \mathcal{U}^\perp$ whenever $\mathbf{v} \in \mathcal{U}^\perp$, i.e., $\mathcal{U}^\perp$ is invariant under T. Hence define an operator $S \in \mathcal{L}(\mathcal{U}^\perp)$ by $S = T|_{\mathcal{U}^\perp}$. If $\mathbf{v}, \mathbf{w} \in \mathcal{U}^\perp$, then

$$\langle S\mathbf{v}, \mathbf{w} \rangle = \langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, T\mathbf{w} \rangle = \langle \mathbf{v}, S\mathbf{w} \rangle$$

which shows S is self-adjoint. By induction hypothesis, there is an orthonormal basis of $\mathcal{U}^\perp$ consisting of eigenvectors of S. As $S\mathbf{v} = T\mathbf{v}$ for every $\mathbf{v} \in \mathcal{U}^\perp$, clearly every eigenvector of S is an eigenvector of T. Hence adjoining $\mathbf{u}$ to an orthonormal basis of $\mathcal{U}^\perp$ consisting of eigenvectors of S gives an orthonormal basis of $\mathcal{V}$ consisting of eigenvectors of T, as desired. □

For $T \in \mathcal{L}(\mathcal{V})$ self-adjoint, next corollary provides nicest decomposition of $\mathcal{V}$ into subspaces invariant in T.

Corollary 1.2.108. *Suppose $T \in \mathcal{L}(\mathcal{V})$ is self-adjoint, let $\lambda_1, \ldots, \lambda_m$ be distinct eigenvalues of T, then*

$$\mathcal{V} = \textit{null}(T - \lambda_1 I) \oplus \cdots \oplus \textit{null}(T - \lambda_m I)$$

Each vector in each $\textit{null}(T - \lambda_j I)$ is orthogonal to all vectors in other subspaces of this decomposition.

Proof. The Spectral Theorems 1.2.104 and 1.2.107 implies $\mathcal{V}$ has a basis consisting of eigenvectors of T. The desired decomposition of $\mathcal{V}$ follows from Proposition 1.2.52.
The orthogonality statement follows from Corollary 1.2.103. □

Lemma 1.2.109. *Suppose $\mathcal{V}$ is a real inner-product space, $\dim(\mathcal{V})$ and $T \in \mathcal{L}(\mathcal{V})$, these are equivalent:*

i. T is normal but not self-adjoint

ii. the matrix of T with respect to every orthonormal basis of $\mathcal{V}$ has the form

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

with $b \neq 0$.

iii. the matrix of T with respect to some orthonormal basis of $\mathcal{V}$ has the form

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

with $b > 0$.

Proof. $(i) \Rightarrow (ii)$. Suppose (i) holds. Let $(\mathbf{e}_1, \mathbf{e}_2)$ be an orthonormal basis of $\mathcal{V}$. Let

$$\mathcal{M}(T, (\mathbf{e}_1, \mathbf{e}_2)) = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$$

Then $\|T\mathbf{e}_1\|^2 = a^2 + b^2$ and $\|T^*\mathbf{e}_1\|^2 = a^2 + c^2$. As T is normal, $\|T\mathbf{e}_1\| = \|T^*\mathbf{e}_1\|$, then these equations imply $b^2 = c^2$, hence $c = b$ or $c = -b$. Note that $c \neq b$ (otherwise T will be self-adjoint), hence $c = -b$, thus

$$\mathcal{M}(T, (\mathbf{e}_1, \mathbf{e}_2)) = \begin{bmatrix} a & -b \\ b & d \end{bmatrix}$$

Hence T^* is the transpose of the matrix above. As T is normal, $TT^* = T^*T$, hence these two matrices must be equal, and $bd = ab$. Also, $b \neq 0$ otherwise T would be self-adjoint, hence $d = a$.
$(ii) \Rightarrow (iii)$. Suppose (ii) holds. Let $(\mathbf{e}_1, \mathbf{e}_2)$ be any orthonormal basis of $\mathcal{V}$. The matrix of T with respect to this basis has form given in (ii), with $b \neq 0$. If $b > 0$, then (iii) holds, hence (ii) implies (iii). If $b < 0$, then

$$\mathcal{M}(T, (\mathbf{e}_1, -\mathbf{e}_2)) = \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$$

where $-b > 0$, hence (ii) implies (iii).
$(iii) \Rightarrow (i)$. Suppose (iii) holds, so matrix of T with respect to some orthonormal basis has form given in (iii) with $b > 0$. Clearly matrix of T is not equal to its transpose as $b \neq 0$, hence T is not self-adjoint. Note that $TT^* = T^*T$, and hence T is normal. Thus (iii) implies (i). □

The next proposition play key role in characterisation of normal operators on real inner-product space.

Proposition 1.2.110. *Suppose $T \in \mathcal{L}(\mathcal{V})$ is normal and $\mathcal{U}$ is a subspace of $\mathcal{V}$ that is invariant under T. Then*

i. $\mathcal{U}^\perp$ is invariant under T

ii. $\mathcal{U}$ is invariant under T^*

iii. $(T|_{\mathcal{U}})^* = (T^*)_{\mathcal{U}}$

iv. $T|_{\mathcal{U}}$ is a normal operator on $\mathcal{U}$

v. $T_{\mathcal{U}^\perp}$ is a normal operator on $\mathcal{U}^\perp$

Proof. (i). Let $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ be an orthonormal basis of $\mathcal{U}$. Extend to an orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{f}_1, \ldots, \mathbf{f}_n)$ of $\mathcal{V}$ (by Corollary 1.2.78). As $\mathcal{U}$ is invariant under T, each $T\mathbf{e}_j$ is a linear combination of $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$. Hence

matrix of T with respect to basis $(\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{f}_1, \ldots, \mathbf{f}_n)$ is of the form

$$\mathcal{M}(T, (\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{f}_1, \ldots, \mathbf{f}_n)) = \begin{bmatrix} A & B \\ 0 & C \end{bmatrix}$$

where A is a $m \times m$ matrix, 0 is a $m \times n$ matrix of all 0s, B is a $m \times n$ matrix, C is a $n \times n$ matrix.
For each $j \in \{1, \ldots, m\}$, $\|T\mathbf{e}_j\|^2$ equals sum of squares of absolute values of entries in jth column of A, hence $\sum_{j=1}^{m} \|T\mathbf{e}_j\|^2$ is sum of squares of absolute values of entries of A.
For each $j \in \{1, \ldots, m\}$, $\|T^*\mathbf{e}_j\|^2$ equals sum of squares of absolute values of entires in jth row of A and B, hence $\sum_{j=1}^{m} \|T^*\mathbf{e}_j\|^2$ is sum of squares of the absolute values of entries of A and B.
As T is normal, $\|T\mathbf{e}_j\| = \|T^*\mathbf{e}_j\|$ for each j, hence $\sum_{j=1}^{m} \|T\mathbf{e}_j\|^2 = \sum_{j=1}^{m} \|T^*\mathbf{e}_j\|^2$ This implies the sum of squares of absolute values of entries of B must equal 0, hence B is matrix consisting of all 0s, hence

$$\mathcal{M}(T, (\mathbf{e}_1, \ldots, \mathbf{e}_m, \mathbf{f}_1, \ldots, \mathbf{f}_n)) = \begin{bmatrix} A & 0 \\ 0 & C \end{bmatrix}$$

This representation shows that $T\mathbf{f}_k$ is the span of $(\mathbf{f}_1, \ldots, \mathbf{f}_n)$ for each k. As $(\mathbf{f}_1, \ldots, \mathbf{f}_n)$ is a basis of $\mathcal{U}^\perp$, this implies $T\mathbf{v} \in \mathcal{U}^\perp$ whenever $\mathbf{v} \in \mathcal{U}^\perp$. In other words, $\mathcal{U}^\perp$. is invariant under T.
(ii). Note that $\mathcal{T}^*$ has block of 0 is lower left, hence each $T^*\mathbf{e}_j$ can be written as linear combination of $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$, thus $\mathcal{U}$ is invariant under T^*.
(iii). Let $S = T|_{\mathcal{U}}$. Fix $\mathbf{v} \in \mathcal{U}$, then

$$\langle S\mathbf{u}, \mathbf{v} \rangle = \langle T\mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, T^*\mathbf{v} \rangle$$

for all $\mathbf{u} \in \mathcal{U}$. As $T^*\mathbf{v} \in \mathcal{U}$ by (ii), equation above shows $S^*\mathbf{v} = T^*\mathbf{v}$. Hence $(T|_{\mathcal{U}})^* = (T^*)_{\mathcal{U}}$.
(iv). Note T commutes with T^* as T is normal, and $(T|_{\mathcal{U}})^* = (T^*)_{\mathcal{U}}$ by (iii), hence $T|_{\mathcal{U}}$ commutes with its adjoint and hence is normal.
(v). Note that in (iv), it is shown that restriction of T to any invariant subspace is normal. However, $\mathcal{U}^\perp$ is invariant under T by (i), hence $T|_{\mathcal{U}}$ is normal. □

Definition 1.2.111. A *block diagonal matrix* is a square matrix of form

$$\begin{bmatrix} A_1 & & 0 \\ & \ddots & \\ 0 & & A_m \end{bmatrix}$$

where $A_1, \ldots, A_m$ are square matrices along the diagonal, and all other entries of matrix equal 0.

Theorem 1.2.112. *Suppose $\mathcal{V}$ is a real inner-product space and $T \in \mathcal{L}(\mathcal{V})$. Then T is normal if and only if there is an orthonormal basis of $\mathcal{V}$ with respect to which T has a block diagonal matrix where each block is a 1×1 or 2×2 matrix of form*

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

where $b > 0$.

Proof. ($\Rightarrow$) Suppose $\exists$ orthonormal basis of $\mathcal{V}$ such that T is a block-diagonal matrix where each block is 1×1 or 2×2 matrix of form shown earlier. Note $TT^* = T^*T$, hence T is normal.
($\Leftarrow$) Suppose T is normal. Note result is trivial if $\dim(\mathcal{V}) = 1$. If $\dim(\mathcal{V}) = 2$, if T is self-adjoint, result holds by Real Spectral Theorem 1.2.107; if not, then result holds by Lemma 1.2.109.
Assume $\dim(\mathcal{V}) > 2$, and desired result holds on vector spaces of smaller dimension. Let $\mathcal{U} \subset \mathcal{V}$, $\dim(\mathcal{U}) = 1$ that is invariant under T if the subspace exists (if T has nonzero eigenvector, let $\mathcal{U}$ be span of this eigenvector). If no such subspace exists, let $\mathcal{U} \subset \mathcal{V}$, $\dim(\mathcal{U}) = 2$ that is invariant under T.
If $\dim(\mathcal{U}) = 1$, choose vector in $\mathcal{U}$ with norm 1, and this vector will b an orthonormal basis of $\mathcal{U}$, and $T|_{\mathcal{U}}$ is 1×1 matrix. If $\dim(\mathcal{U}) = 2$, then $T_{\mathcal{U}}$ is normal but not self adjoint (else $T|_{\mathcal{U}}$ have nonzero eigenvector), hence choose an orthonormal basis of $\mathcal{U}$ with respect to which the matrix of $T|_{\mathcal{U}}$ has form in statement.
Now $\mathcal{U}^\perp$ is invariant under T and $T|_{\mathcal{U}^\perp}$ is a normal operator on $\mathcal{U}^\perp$. By induction, there is an orthonormal basis of $\mathcal{U}^\perp$ with respect to which matrix of $T|_{\mathcal{U}^\perp}$ has the desired form. Adjoining this basis to basis of $\mathcal{U}$ gives an orthonormal basis of $\mathcal{V}$ with respect to which matrix of T has the desireed form. □

Definition 1.2.113. An operator $T \in \mathcal{L}(\mathcal{V})$ is *positive semi-definite* if T is self-adjoint and $\langle T\mathbf{v}, \mathbf{v}\rangle \geq 0 \; \forall \mathbf{v} \in \mathcal{V}$.

Definition 1.2.114. An operator S is a *square root* of an operator T if $S^2 = T$.

The next theorem characterises positive operators that correspond to characterisations of nonnegative numbers among $\mathbb{C}$.

Theorem 1.2.115. *Let $T \in \mathcal{L}(\mathcal{V})$, then the following are equivalent:*

i. T is positive

ii. T is self-adjoint and all eigenvalues of T are nonnegative

iii. T has a positive square root

iv. T has a self-adjoint square root

v. there exists an operator $S \in \mathcal{L}(\mathcal{V})$ such that $T = S^*S$

Proof. $(i) \Rightarrow (ii)$. Suppose (i) holds, so T is positive. By definition of positive operator, T is self-adjoint. Suppose λ is an eigenvalue of T. Let $\mathbf{v}$ be a nonzero eigenvector of T corresponding to λ, then

$$0 \leq \langle T\mathbf{v}, \mathbf{v}\rangle = \langle \lambda\mathbf{v}, \mathbf{v}\rangle = \lambda \langle \mathbf{v}, \mathbf{v}\rangle$$

hence λ is a nonnegative number.
$(ii) \Rightarrow (iii)$. Suppose (ii) holds, so T is self-adjoint and all the eigenvalues of T are nonnegative. By Spectral Theorem 1.2.107 and 1.2.104, there is an orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ of $\mathcal{V}$ consisting of eigenvectors of T. Let $\lambda_1, \ldots, \lambda_n$ be eigenvalues of T corresponding to $\mathbf{e}_1, \ldots, \mathbf{e}_n$ so that each λ_j is a nonnegative number. Let $S \in \mathcal{L}(\mathcal{V})$ such that $S\mathbf{e}_j = \sqrt{\lambda_j}\mathbf{e}_j$ for $j = 1, \ldots, n$, then S is a positive operator. Also, $S^2\mathbf{e}_j = \lambda_j\mathbf{e}_j = T\mathbf{e}_j$ for each j, which implies $S^2 = T$. Hence S is a positive square root of T, and (iii) holds.
$(iii) \Rightarrow (iv)$. Note that by definition, every positive operator is self-adjoint hence this holds.
$(iv) \Rightarrow (v)$. Let $S \in \mathcal{L}(\mathcal{V})$, $T = S^*S$, then $T^* = (S^*S)^* = S^*(S^*)^* = S^*S = T$, hence T is self-adjoint. Then

$$\langle T\mathbf{v}, \mathbf{v}\rangle = \langle S^*S\mathbf{v}, \mathbf{v}\rangle = \langle S\mathbf{v}, S\mathbf{v}\rangle \geq 0$$

for every $\mathbf{v} \in \mathcal{V}$, hence T is positive. □

A positive operator can have infinitely many square roots, but only one can be positive.

Proposition 1.2.116. *Every positive operator on $\mathcal{V}$ has a unique positive square root.*

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$ is positive. Let $\lambda_1, \ldots, \lambda_m$ denote distinct eigenvalues of T. As T is positive, all these are nonnegative by Theorem 1.2.115. As T is self-adjoint, by Corollary 1.2.108,

$$\mathcal{V} = \text{null}(T - \lambda_1 I) \oplus \cdots \oplus \text{null}(T - \lambda_m I)$$

Suppose $S \in \mathcal{L}(\mathcal{V})$ is a positive square root of T, and α is an eigenvalue of S. If $\mathbf{v} \in \text{null}(S - \alpha I)$, then $S\mathbf{v} = \alpha\mathbf{v}$, implying $T\mathbf{v} = S^2\mathbf{v} = \alpha^2\mathbf{v}$ so that $\mathbf{v} \in \text{null}(T - \alpha^2 I)$. Thus α^2 is an eigenvalue of T, which means α^2 must equal some λ_j. Hence $\alpha = \sqrt{\lambda_j}$ for some j. Also,

$$\text{null}(S - \sqrt{\lambda_j} I) \subset \text{null}(T - \lambda_j I)$$

From above, the only possible eigenvalues for S are $\sqrt{\lambda_1}, \ldots, \sqrt{\lambda_m}$. As S is self-adjoint, this implies

$$\mathcal{V} = \text{null}(S - \sqrt{\lambda_1} I) \oplus \cdots \oplus \text{null}(S - \sqrt{\lambda_m} I)$$

Hence these implies $\text{null}(S - \sqrt{\lambda_j} I) = \text{null}(T - \lambda_j I)$ for each j. On $\text{null}(T - \lambda_j I)$, the operator S is just multiplication by $\sqrt{\lambda_j}$. Hence S is a uniquely determined positive square root of T. □

Definition 1.2.117. An operator $S \in \mathcal{L}(\mathcal{V})$ is an *isometry* if $\|S\mathbf{v}\| = \|\mathbf{v}\|$ for all $\mathbf{v} \in \mathcal{V}$.
An isometry on a real inner-product space is an *orthogonal operator*.
An isometry on a complex inner-product space is an *unitary operator*.
If $S \in \mathcal{L}(\mathcal{V})$ is an isometry, then S is injective, hence every isometry is invertible.

Next theorem provides several conditions that are equivalent to being an isometry.

Theorem 1.2.118. *Suppose $S \in \mathcal{L}(\mathcal{V})$, then the following are equivalent:*

i. S is an isometry

ii. $\langle S\mathbf{u}, S\mathbf{v}\rangle = \langle \mathbf{u}, \mathbf{v}\rangle$ *for all* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$ *(inner product is preserved)*

iii. $S^*S = I$

iv. $(S\mathbf{e}_1, \ldots, S\mathbf{e}_n)$ *is orthonormal whenever* $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ *is an orthonormal list of vectors in* $\mathcal{V}$

v. $\exists$ *an orthonormal basis* $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ *of* $\mathcal{V}$ *such that* $(S\mathbf{e}_1, \ldots, S\mathbf{e}_n)$ *is orthonormal*

vi. S^* *is an isometry*

vii. $\langle S^*\mathbf{u}, S^*\mathbf{v}\rangle = \langle \mathbf{u}, \mathbf{v}\rangle$ *for all* $\mathbf{u}, \mathbf{v} \in \mathcal{V}$

viii. $SS^* = I$

ix. $(S^*\mathbf{e}_1, \ldots, S^*\mathbf{e}_n)$ *is orthonormal whenever* $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ *is an orthonormal list of vectors in* $\mathcal{V}$

x. $\exists$ *an orthonormal basis* $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ *of* $\mathcal{V}$ *such that* $(S^*\mathbf{e}_1, \ldots, S^*\mathbf{e}_n)$ *is orthonormal*

Proof. (i). Suppose (i) holds, and $\mathcal{V}$ is a real-inner product space, then for every $\mathbf{u}, \mathbf{v} \in \mathcal{V}$, we have

$$\begin{aligned}\langle S\mathbf{u}, S\mathbf{v}\rangle &= (\|S\mathbf{u} + S\mathbf{v}\|^2 - \|S\mathbf{u} - S\mathbf{v}\|^2)/4 \\ &= (\|S(\mathbf{u} + \mathbf{v})\|^2 - \|S(\mathbf{u} - \mathbf{v})\|^2)/4 \\ &= (\|\mathbf{u} + \mathbf{v}\|^2 - \|\mathbf{u} - \mathbf{v}\|^2)/4 \\ &= \langle \mathbf{u}, \mathbf{v}\rangle\end{aligned}$$

$(ii) \Rightarrow (iii)$. Suppose (ii) holds, then

$$\langle (S^*S - I)\mathbf{u}, \mathbf{v}\rangle = \langle S\mathbf{u}, S\mathbf{v}\rangle - \langle \mathbf{u}, \mathbf{v}\rangle = 0$$

for every $\mathbf{u}, \mathbf{v} \in \mathcal{V}$. Let $\mathbf{v} = (S^*S - I)\mathbf{u}$, then $S^*S - I = 0$, hence $S^*S = I$.
$(iii) \Rightarrow (iv)$. Suppose (iii) holds, and $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ is an orthonormal list of vectors in $\mathcal{V}$, then

$$\langle S\mathbf{e}_j, S\mathbf{e}_k\rangle = \langle S^*S\mathbf{e}_j, \mathbf{e}_k\rangle = \langle \mathbf{e}_j, \mathbf{e}_k\rangle$$

hence $(S\mathbf{e}_1, \ldots, S\mathbf{e}_n)$ is orthonormal.
$(iv) \Rightarrow (v)$ is quite obvious by implication.
$(v) \Rightarrow (i)$. Suppose (v) holds, let $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ be an orthonormal basis of $\mathcal{V}$ such that $(S\mathbf{e}_1, \ldots, S\mathbf{e}_n)$ is orthonormal. If $\mathbf{v} \in \mathcal{V}$, then

$$\begin{aligned}\|S\mathbf{v}\|^2 &= \|S(\langle \mathbf{v}, \mathbf{e}_1\rangle \mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_n\rangle \mathbf{e}_n)\|^2 \\ &= \|\langle \mathbf{v}, \mathbf{e}_1\rangle S\mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e_n}\rangle S\mathbf{e}_n\|^2 \\ &= |\langle \mathbf{v}, \mathbf{e}_1\rangle|^2 + \cdots + |\langle \mathbf{v}, \mathbf{e}_n\rangle|^2 = \|\mathbf{v}\|^2\end{aligned}$$

Taking square roots, S is shown to be an isometry.
Replacing S with S^*, (vi) to (x) are all equivalent to each other.
To connect the two groups of conditions, show $(iii) \to (viii)$. In general, S does not commute with S^*. However, $S^*S = I$ if and only if $SS^* = I$. Hence this completes the proof. □

Theorem 1.2.119. *Suppose* $\mathcal{V}$ *is a complex inner-product space and* $S \in \mathcal{L}(\mathcal{V})$*, then* S *is an isometry if and only if there is an orthonormal basis of* $\mathcal{V}$ *consisting of eigenvectors of* S *all of whose corresponding eigenvalues have absolute value* 1*.*

Proof. $(\Rightarrow)$. Note that λI is an isometry whenever $\lambda \in \mathbb{F}$ satisfies $|\lambda| = 1$.
Let $\lambda_1, \ldots, \lambda_n$ be scalars with absolute value 1 and $S \in \mathcal{L}(\mathcal{V})$ satisfies $S(\mathbf{e}_j) = \lambda_j \mathbf{e}_j$ for some orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ of $\mathcal{V}$. Suppose $\mathbf{v} \in \mathcal{V}$, then

$$\begin{aligned}\mathbf{v} &= \langle \mathbf{v}, \mathbf{e}_1\rangle \mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_n\rangle \mathbf{e}_n \\ \|\mathbf{v}\|^2 &= |\langle \mathbf{v}, \mathbf{e}_1\rangle|^2 + \cdots + |\langle \mathbf{v}, \mathbf{e}_n\rangle|^2\end{aligned}$$

Applying S to both sides gives

$$\begin{aligned}S\mathbf{v} &= \langle \mathbf{v}, \mathbf{e}_1\rangle S\mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_n\rangle S\mathbf{e}_n \\ &= \lambda_1 \langle \mathbf{v}, \mathbf{e}_1\rangle \mathbf{e}_1 + \cdots + \lambda_n \langle \mathbf{v}, \mathbf{e}_n\rangle \mathbf{e}_n \\ \|S\mathbf{v}\|^2 &= |\langle \mathbf{v}, \mathbf{e}_1\rangle|^2 + \cdots + |\langle \mathbf{v}, \mathbf{e}_n\rangle|^2\end{aligned}$$

Hence this shows $\|\mathbf{v}\| = \|S\mathbf{v}\|$, thus S is an isometry.
($\Leftarrow$). Suppose S is an isometry. By Complex Spectral Theorem 1.2.104, there is an orthonormal basis $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ of $\mathcal{V}$ consisting of eigenvectors of S. For $j \in \{1, \ldots, n\}$, let λ_j be eigenvalue corresponding to $\mathbf{e}_j$, then

$$|\lambda_j| = \|\lambda_j \mathbf{e}_j\| = \|S\mathbf{e}_j\| = \|\mathbf{e}_j\| = 1$$

Hence each eigenvalue of S has absolute value 1. □

The next theorem implies that an isometry on an odd-dimensional real inner-product space must have 1 or -1 as an eigenvalue.

Theorem 1.2.120. *Suppose $\mathcal{V}$ is a real inner-product space with $S \in \mathcal{L}(\mathcal{V})$. Then S is an isometry if and only if there is an orthonormal basis of $\mathcal{V}$ with respect to which S has a block diagonal matrix where each block on the diagonal is a 1×1 matrix containing 1 or -1, or a 2×2 matrix of form*

$$\begin{bmatrix} \sin\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$

with $\theta \in (0, \pi)$.

Proof. ($\Rightarrow$). Suppose S is an isometry. As S is normal, $\exists$ an orthonormal basis of V such that with respect to this basis S has a block diagonal matrix, where each block is a 1×1 matrix or 2×2 matrix of form

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

with $b > 0$.
If λ is an entry in a 1×1 along the diagonal of matrix of S with respect to basis above, then $\exists$ a basis vector $\mathbf{e}_j$ such that $S\mathbf{e}_j = \lambda \mathbf{e}_j$. As S is an isometry, this implies $|\lambda| = 1$, hence $\lambda = 1$ or -1.
Consider 2×2 matrix from earlier along the diagonal of the matrix of S. $\exists$ basis vectors $\mathbf{e}_j, \mathbf{e}_{j+1}$ such that

$$S\mathbf{e}_j = a\mathbf{e}_j + b\mathbf{e}_{j+1}$$

Hence $1 = \|\mathbf{e}_j\|^2 = \|S\mathbf{e}_j\|^2 = a^2 + b^2$. Along with condition $b > 0$, this implies $\exists$ number $\theta \in (0, \pi)$ such that $a = \cos\theta$ and $b = \sin\theta$, hence the matrix has the required form.
($\Leftarrow$). Suppose $\exists$ an orthonormal basis of $\mathcal{V}$ with respect to which matrix S has form in theorem. Then there is a direct sum decomposition $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$ where each $\mathcal{U}_j$ is a subspace of $\mathcal{V}$ of dimension 1 or 2. Furthermore any two vectors belonging to distinct $\mathcal{U}_j$ are orthogonal, and each $S|_{\mathcal{U}_j}$ is an isometry mapping $\mathcal{U}_j$ to $\mathcal{U}_j$. If $\mathbf{v} \in \mathcal{V}$, then $\mathbf{v} = \mathbf{u}_1 + \cdots + \mathbf{u}_m$ where each $\mathbf{u}_j \in \mathcal{U}_j$. Applying S to above equation and taking norms gives

$$\begin{aligned} \|S\mathbf{v}\|^2 &= \|S\mathbf{u}_1 + \cdots + S\mathbf{u}_m\|^2 \\ &= \|S\mathbf{u}_1\|^2 + \cdots + \|S\mathbf{u}_m\|^2 \\ &= \|\mathbf{u}_1\|^2 + \cdots + \|\mathbf{u}_m\|^2 = \|\mathbf{v}\|^2 \end{aligned}$$

Hence S is an isometry as desired. □

Theorem 1.2.121. *Polar Decomposition*
If $T \in \mathcal{L}(\mathcal{V})$, then there exists an isometry $S \in \mathcal{L}(\mathcal{V})$ such that $T = S\sqrt{T^*T}$.

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$. If $\mathbf{v} \in \mathcal{V}$, then

$$\|T\mathbf{v}\|^2 = \langle T\mathbf{v}, T\mathbf{v}\rangle = \langle T^*T\mathbf{v}, \mathbf{v}\rangle = \left\langle \sqrt{T^*T}\sqrt{T^*T}\mathbf{v}, \mathbf{v} \right\rangle = \left\langle \sqrt{T^*T}\mathbf{v}, \sqrt{T^*T}\mathbf{v} \right\rangle = \left\| \sqrt{T^*T}\mathbf{v} \right\|^2$$

Hence $\|T\mathbf{v}\| = \left\| \sqrt{T^*T}\mathbf{v} \right\|$ for all $\mathbf{v} \in \mathcal{V}$.
Define linear map $S_1 : \text{range}(\sqrt{T^*T}) \to \text{range}(T)$ by $S_1(\sqrt{T^*T}\mathbf{v}) = T\mathbf{v}$. To extend S_1 to an isometry $S \in \mathcal{L}(\mathcal{V})$ such that $T = S\sqrt{T^*T}$.
First to check S_1 is well defined. Suppose $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}$ such that $\sqrt{T^*T}\mathbf{v}_1 = \sqrt{T^*T}\mathbf{v}_2$, to show $T\mathbf{v}_1 = T\mathbf{v}_2$,

$$\|T\mathbf{v}_1 - T\mathbf{v}_2\| = \|T(\mathbf{v}_1 - \mathbf{v}_2)\| = \left\| \sqrt{T^*T}(\mathbf{v}_1 - \mathbf{v}_2) \right\| = \left\| \sqrt{T^*T}\mathbf{v}_1 - \sqrt{T^*T}\mathbf{v}_2 \right\| = 0$$

As $T\mathbf{v}_1 = T\mathbf{v}_2$, S_1 is well defined. Next, note that

$$S_1(\sqrt{T^*T}\mathbf{v}_1 + \sqrt{T^*T}\mathbf{v}_2) = S_1(\sqrt{T^*T}(\mathbf{v}_1 + \mathbf{v}_2)) = T(\mathbf{v}_1 + \mathbf{v}_2) = T\mathbf{v}_1 + T\mathbf{v}_2 = S_1(\sqrt{T^*T}\mathbf{v}_1) + S_1(\sqrt{T^*T}\mathbf{v}_2)$$

Hence S_1 is additive. In addition, for any scalar $\alpha in \mathbb{F}$,

$$S_1(\alpha\sqrt{T^*T}\mathbf{v}_1) = S_1(\sqrt{T^*T}(\alpha\mathbf{v}_1)) = T(\alpha\mathbf{v}) = \alpha T\mathbf{v} = \alpha S_1(\sqrt{T^*T}\mathbf{v}_1)$$

hence scalar multiplication is preserved. Since S_1 satisfies additivity and homogeneity, it is a linear map.
By definition, S_1 maps range$(\sqrt{T^*T})$ onto range(T), hence this imply $\|S_1\mathbf{u}\| = \|\mathbf{u}\|$ for all $\mathbf{u} \in$ range$(\sqrt{T^*T})$, hence S_1 is injective. Thus dim(range$(\sqrt{T^*T})$) = dim(range(T)), and thus dim(range$(\sqrt{T^*T}))^\perp$ = dim(range$(T))^\perp$.
The orthonormal bases $(\mathbf{e}_1, \ldots, \mathbf{e}_m)$ of (range$(\sqrt{T^*T}))^\perp$ and $(\mathbf{f}_1, \ldots, \mathbf{f}_m)$ of (range$(T))^\perp$ can be chosen, and these two bases have the same length. Define linear map $S_2 : (\text{range}(\sqrt{T^*T}))^\perp \to (\text{range}(T))^\perp$ by

$$S_2(a_1\mathbf{e}_1 + \cdots + a_m\mathbf{e}_m) = a_1\mathbf{f}_1 + \cdots + a_m\mathbf{f}_1$$

Obviously $\|S_2\mathbf{w}\| = \|\mathbf{w}\|$ for all $\mathbf{w} \in (\text{range}(\sqrt{T^*T}))^\perp$.
Let S be the operator on $\mathcal{V}$ that equals S_1 on range$(\sqrt{T^*T})$ and equals S_2 on (range$(\sqrt{T^*T}))^\perp$. Note each $\mathbf{v} \in \mathcal{V}$ can be written uniquely in the form $\mathbf{v} = \mathbf{u} + \mathbf{w}$ where $\mathbf{u} \in$ range$(\sqrt{T^*T})$ and $\mathbf{w} \in$ (range$(\sqrt{T^*T}))^\perp$. Then define $S\mathbf{v}$ by $S\mathbf{v} = S_1\mathbf{u} + S_2\mathbf{w}$. For each $\mathbf{v} \in \mathcal{V}$, then $S(\sqrt{T^*T}\mathbf{v}) = S_1(\sqrt{T^*T}\mathbf{v}) = T\mathbf{v}$, hence $T = S\sqrt{T^*T}$.
To show S is an isometry, note that

$$\|S\mathbf{v}\|^2 = \|S_1\mathbf{u} + S_2\mathbf{w}\|^2 = \|S_1\mathbf{u}\|^2 + \|S_2\mathbf{w}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v}\|^2$$

where the second equality above holds as $S_1\mathbf{u} \in$ range(T) and $S_2\mathbf{u} \in$ (range$(T))^\perp$. □

Polar decomposition states that each operator on $\mathcal{V}$ is the product of an isometry and a positive operator.

Definition 1.2.122. Suppose $T \in \mathcal{L}(\mathcal{V})$. The *singular values* of T are eigenvalues of $\sqrt{T^*T}$, with each eigenvalue λ repeated dim(null$(\sqrt{T^*T} - \lambda I)$) times, all nonnegative as these are eigenvalues of positive operator $\sqrt{T^*T}$.

Theorem 1.2.123. *Singular Value Decomposition*
If $T \in \mathcal{L}(\mathcal{V})$ has singular values $s_1, \ldots, s_n$, then $\exists$ orthonormal bases $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ and $(\mathbf{f}_1, \ldots, \mathbf{f}_n)$ of $\mathcal{V}$ s.t.

$$T\mathbf{v} = s_1 \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{f}_1 + \cdots + s_n \langle \mathbf{v}, \mathbf{e}_n \rangle \mathbf{f}_n$$

for every $\mathbf{v} \in \mathcal{V}$.

Proof. By Spectral Theorem 1.2.104 applied to $\sqrt{T^*T}$, $\exists$ an orthonormal bases $(\mathbf{e}_1, \ldots, \mathbf{e}_n)$ of $\mathcal{V}$ such that $\sqrt{T^*T}\mathbf{e}_j = s_j\mathbf{e}_j$ for $j = 1, \ldots, n$. Then $\mathbf{v} = \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{e}_1 + \cdots + \langle \mathbf{v}, \mathbf{e}_n \rangle \mathbf{e}_n$ for every $\mathbf{v} \in \mathcal{V}$. Apply $\sqrt{T^*T}$ to both sides of this equation to get $\sqrt{T^*T}\mathbf{v} = s_1 \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{e}_1 + \cdots + s_n \langle \mathbf{v}, \mathbf{e}_n \rangle \mathbf{e}_n$ for every $\mathbf{v} \in \mathcal{V}$.
By Polar Decomposition Theorem 1.2.121, $\exists$ an isometry $S \in \mathcal{L}(\mathcal{V})$ such that $T = S\sqrt{T^*T}$. Apply S to both sides of the equation to get $T\mathbf{v} = s_1 \langle \mathbf{v}, \mathbf{e}_1 \rangle S\mathbf{e}_1 + \cdots + s_n \langle \mathbf{v}, \mathbf{e}_n \rangle S\mathbf{e}_n$ for every $\mathbf{v} \in \mathcal{V}$.
For each j, let $\mathbf{f}_j = S\mathbf{e}_j$. As S is an isometry, $(\mathbf{f}_1, \ldots, \mathbf{f}_n)$ is an orthonormal basis of $\mathcal{V}$. Then

$$T\mathbf{v} = s_1 \langle \mathbf{v}, \mathbf{e}_1 \rangle \mathbf{f}_1 + \cdots + s_n \langle \mathbf{v}, \mathbf{e}_n \rangle \mathbf{f}_n$$

for every $\mathbf{v} \in \mathcal{V}$, completing the proof. □

1.2.6 Operators on Complex Vector Spaces

Definition 1.2.124. Given $T \in \mathcal{L}(\mathcal{V})$, and λ an eigenvalue of T, a vector $\mathbf{v} \in \mathcal{V}$ is a *generalised eigenvector* of T corresponding to λ if $(T - \lambda I)^j\mathbf{v} = \mathbf{0}$ for some positive integer j.

Suppose $T \in \mathcal{L}(\mathcal{V})$ and $k \geq 0$, $k \in \mathbb{Z}$. If $T^k\mathbf{v} = \mathbf{0}$, then $T^{k+1}\mathbf{v} = T(T^k\mathbf{v}) = T(\mathbf{0}) = \mathbf{0}$, hence,

$$\{\mathbf{0}\} = \text{null}(T^0) \subset \text{null}(T^1) \subset \cdots \subset \text{null}(T^k) \subset \cdots$$

Next proposition states that once two consecutive terms in this sequence of subspaces are equal, then later terms in the sequence are equal.

Proposition 1.2.125. *If $T \in \mathcal{L}(\mathcal{V})$ and $m \geq 0$, $m \in \mathbb{Z}$ such that null(T^m) = null(T^{m+1}), then*

$$null(T^0) \subset null(T^1) \subset \cdots \subset null(T^m) = null(T^{m+1}) = null(T^{m+2}) = \cdots$$

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$ and $m \geq 0$, $m \in \mathbb{Z}$ such that $\text{null}(T^m) = \text{null}(T^{m+1})$. Let k be a positive integer. Note that $\text{null}(T^{m+k}) \subset \text{null}(T^{m+k+1})$. To prove inclusion in other direction, suppose $\mathbf{v} \in \text{null}(T^{m+k+1})$, then

$$\mathbf{0} = T^{m+k+1}\mathbf{v} = T^{m+1}(T^k\mathbf{v})$$
$$T^k\mathbf{v} \in \text{null}(T^{m+1}) = \text{null}(T^m)$$

Hence $\mathbf{0} = T^m(T^k\mathbf{v}) = T^{m+k}\mathbf{v}$, thus $\mathbf{v} \in \text{null}(T^{m+k})$. Hence this implies $\text{null}(T^{m+k+1}) \subset \text{null}(T^{m+k})$. □

Proposition 1.2.126. *If $T \in \mathcal{L}(\mathcal{V})$, then*

$$\textit{null}(T^{\dim(\mathcal{V})}) = \textit{null}(T^{\dim(\mathcal{V}+1)}) = \textit{null}(T^{\dim(\mathcal{V}+2)}) = \cdots$$

Proof. Suppose $T \in \mathcal{L}(\mathcal{V})$, and $\text{null}(T^{\dim(\mathcal{V})}) \neq \text{null}(T^{\dim(\mathcal{V}+1)})$, then by Proposition 1.2.125,

$$\{\mathbf{0}\} = \text{null}(T^0) \subsetneq \text{null}(T^1) \subsetneq \cdots \subsetneq \text{null}(T^{\dim(\mathcal{V})}) \subsetneq \text{null}(T^{\dim(\mathcal{V}+1)})$$

By strict inclusion, dimension must increase by at least 1, hence $\dim(\text{null}(T^{\dim(\mathcal{V}+1)})) \geq \dim(\mathcal{V} = 1)$, a contraction as a subspace of $\mathcal{V}$ cannot have a larger dimension than $\dim(\mathcal{V})$. □

The next corollary implies that the generalised eigenvectors of $T \in \mathcal{L}(\mathcal{V})$ corresponding to an eigenvalue λ is a subspace of $\mathcal{V}$.

Corollary 1.2.127. *Suppose $T \in \mathcal{L}(\mathcal{V})$ and λ is an eigenvalue of T, then the set of generalised eigenvectors of T corresponding to λ equals $\textit{null}(T - \lambda I)^{\dim(\mathcal{V})}$.*

Proof. If $\mathbf{v} \in \text{null}(T - \lambda I)^{\dim(\mathcal{V})}$, then $\mathbf{v}$ is a generalised eigenvector of T corresponding to λ.
Conversely, suppose $\mathbf{v} \in \mathcal{V}$ is a generalised eigenvector of T corresponding to λ, thus there exists $j \geq 0 \in \mathbb{Z}$ such that $\mathbf{v} \in \text{null}(T - \lambda I)^j$. By Proposition 1.2.125 and Proposition 1.2.126, replacing T with $T - \lambda I$, then $\mathbf{v} \in \text{null}(T - \lambda I)^{\dim(\mathcal{V})}$ as desired. □

Definition 1.2.128. An operator is *nilpotent* if some power of it equals 0.

Corollary 1.2.129. *Suppose $N \in \mathcal{L}(\mathcal{V})$ is nilpotent, then $N^{\dim(\mathcal{V})} = 0$.*

Proof. As N is nilpotent, every eigenvector in $\mathcal{V}$ is a generalised eigenvector corresponding to eigenvalue 0. Hence from Corollary 1.2.127, $\text{null}(N^{\dim(\mathcal{V})}) = \mathcal{V}$ as desired. □

Suppose $T \in \mathcal{L}(\mathcal{V})$ and $k \geq 0, k \in \mathbb{Z}$. If $\mathbf{w} \in \text{range}(T^{k+1})$, then $\exists\ \mathbf{v} \in \mathcal{V}$ with $\mathbf{w} = T^{k+1}\mathbf{v} = T^k(T\mathbf{v}) \in \text{range}(T^k)$, hence $\text{range}(T^{k+1}) \subset \text{range}(T^k)$, thus

$$\mathcal{V} = \text{range}(T^0) \supset \text{range}(T^1) \supset \cdots \supset \text{range}(T^k) \subset \text{range}(T^{k+1}) \subset \cdots$$

Next proportion shows that inclusions become equalities once power reaches dimension $\mathcal{V}$.

Proposition 1.2.130. *If $T \in \mathcal{L}(\mathcal{V})$, then $\textit{range}(T^{\dim(\mathcal{V})}) = \textit{range}(T^{\dim(\mathcal{V}+1)}) = \textit{range}(T^{\dim(\mathcal{V})+2}) = \cdots$*

Proof. Suppose $m > \dim(\mathcal{V})$, then

$$\dim(\text{range}(T^m)) = \dim(\mathcal{V}) - \dim(\text{null}(T^m)) = \dim(\mathcal{V}) - \dim(\text{null}(T^{\dim(\mathcal{V})})) = \dim(\text{range}(T^{\dim(\mathcal{V})}))$$

by Proposition 1.2.126. As $\text{range}(T^{\dim(\mathcal{V})}) \subset \text{range}(T^m)$, and $\dim(\text{range}(T^{\dim(\mathcal{V})})) = \dim(\text{range}(T^m))$ from above, this implies $\text{range}(T^{\dim(\mathcal{V})}) = \text{range}(T^m)$ as desired. □

Next theorem implies that the number of times an eigenvalue is repeated on diagonal of upper-triangular matrix of T is independent of the choice of basis.

Theorem 1.2.131. *Let $T \in \mathcal{L}(\mathcal{V})$ and $\lambda \in \mathbb{F}$, then for every basis of $\mathcal{V}$ with respect to which T has an upper-triangular matrix, λ appears on diagonal of matrix of T precisely $\dim(\textit{null}(T - \lambda I))^{\dim(\mathcal{V})}$ times.*

Proof. If T is upper-triangular with some basis $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ and diagonal $\lambda_1, \ldots, \lambda_n$, then the number of times a given scalar λ_j appears on diagonal equals $\dim(\text{null}(T - \lambda I)^n)$. As replacing T by $T - \lambda I$ shifts diagonal by

$-\lambda$, it suffes to prove the case $\lambda = 0$. Let $n = \dim(\mathcal{V})$.
Assume $n > 1$, and statement is true in dimension $n-1$. Pick basis $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ which makes T upper-triangular.

$$\mathcal{M}(T) = \begin{bmatrix} \lambda_1 & & & * \\ & \ddots & & \\ & & \lambda_{n-1} & \\ 0 & & & \lambda_n \end{bmatrix}$$

Let $\mathcal{U} = \text{span}(\mathbf{v}_1, \dots \mathbf{v}_{n-1})$. By Proposition 1.2.52, $\mathcal{U}$ is invariant under T, and

$$\mathcal{M}(T|_{\mathcal{U}}) = \begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_{n-1} \end{bmatrix}$$

By induction hypothesis applied to $T|_{\mathcal{U}}$ (space dimension $n-1$), the number of zeros on diagonal is $\dim(\text{null}(T|_{\mathcal{U}}))^{n-1}$.
The nullspaces in power sequence stabilises by dimension $n-1$, hence $\text{null}(T|_{\mathcal{U}})^{n-1} = \text{null}(T|_{\mathcal{U}})^n$, thus 0 appears on diagonal exactly $\text{null}(T|_{\mathcal{U}})^n$ times. To finish proof, relate $\text{null}(T^n)$ to $\text{null}(K|_{\mathcal{U}})^n$ times.
For case $\lambda_n \neq 0$, claim $\text{null}(T^n) \subset \mathcal{U}$. Note that powers of upper-triangular matrix are still upper triangular with diagonal $\lambda_1^n, \dots \lambda_n^n$. In particular

$$\mathcal{M}(T^n) = (T)^n \Rightarrow T^n\mathbf{v}_n = \mathbf{u} + \lambda_n^n\mathbf{v}_n$$

for some $\mathbf{u} \in \mathcal{U}$. Take any $\mathbf{v} \in \text{null}(T^n)$ and write $\mathbf{v} = \tilde{\mathbf{u}} + a\mathbf{v}_n$, where $\tilde{\mathbf{u}} \in \mathcal{U}$ and $a \in \mathbb{F}$, then

$$\mathbf{0} = T^n\mathbf{v} = T^n\tilde{\mathbf{u}} + aT^n\mathbf{v}_n = T^n\tilde{\mathbf{u}} + a\mathbf{u} + a\lambda_n^n\mathbf{v}_n$$

Note that $T^n\tilde{\mathbf{u}}, a\mathbf{u} \in \mathcal{U}$ and $\mathbf{v}_n \neq \mathcal{U}$, which implies $a\lambda_n^n = 0$. However $\lambda_n \neq 0$, so $a = 0$, hence $\mathbf{v} = \tilde{\mathbf{u}} \in \mathcal{U}$. Thus $\text{null}(T^n) = \text{null}(K|_{\mathcal{U}})^n$, hence total number of zeros on diagonal of $\mathcal{M}(T)$ equals $\mathcal{M}(T|_{\mathcal{U}})$, namely $\dim(\text{null}(T^n))$.
For case $\lambda_n = 0$, to get $\dim(\text{null}(T^n)) = \dim(\text{null}(K|_{\mathcal{U}}))^n + 1$. By dimension for sum of subspaces,

$$\dim(\text{null}(T^n)) = \dim z\mathcal{U} \cap \text{null}(T^n)) + \dim(\mathcal{U} + \text{null}(T^n)) - \dim(\mathcal{U})$$

Note that $\mathcal{U} \cap \text{null}(T^n) = \text{null}(K|_{\mathcal{U}})^n$, hence result will follow if $\mathcal{U} + \text{null}(T^n) = \mathcal{V}$, i.e., there is a vector in $\text{null}(T^n)$ not in $\mathcal{U}$. The vector might be of form $\mathbf{u} - \mathbf{v}_n$ with $\mathbf{u} \in \mathcal{U}$. Want $T^n(\mathbf{u} - \mathbf{v}_n) = \mathbf{0}$, i.e., $T^n\mathbf{u} = T^n\mathbf{v}_n$. This is solvable if $T^n\mathbf{v}_n \in \text{range}(T|_{\mathcal{U}})^n$. When $\lambda_n = 0$, the last diagonal entry is 0, hence the last coordinate of $T\mathbf{v}_n$ is 0, hence $T\mathbf{v}_n \in \mathcal{U}$, therefore

$$T^n\mathbf{v}_n = T^{n-1}(T\mathbf{v}_n) \in \text{range}(T|_{\mathcal{U}})^{n-1}$$

As range stabilises, thus $\text{range}(T|_{\mathcal{U}})^{n-1} = \text{range}(T|_{\mathcal{U}})^n$, hence there exists $\mathbf{u} \in \mathcal{U}$ with $T^n\mathbf{u} = T^n\mathbf{v}_n$, making $\mathbf{u} - \mathbf{v}_n \in \text{null}(T^n)$. Since $\mathbf{v}_n \notin \mathcal{U}$, this vector is not in $\mathcal{U}$. As $\mathcal{U} + \text{null}(T^n) = \mathcal{V}$, and with first equation in this case, the $\lambda_n = 0$ case is done.
In either case the number of zeroes on diagonal of $\mathcal{M}(T)$ equals $\dim(\text{null}(T^n))$. Replacing T with $T - \lambda I$ gives the general statement: the number of times λ appears on diagonal equals $\dim(\text{null}(T - \lambda I)^n)$. □

Definition 1.2.132. Suppose $T \in \mathcal{L}(\mathcal{V})$. The *multiplicity* of eigenvalue λ in T is defined to be dimension of subspace of generalised eigenvectors corresponding to λ, which is $\dim(\text{null}(T - \lambda I)^{\dim(\mathcal{V})})$.

Proposition 1.2.133. *If $\mathcal{V}$ is a complex vector space and $T \in \mathcal{L}(\mathcal{V})$, then the sum of multiplicities of all eigenvalues of T equals $\dim(\mathcal{V})$.*

Proof. Over $\mathbb{C}$, every complex operator T has at least one eigenvalue, ensuring that T can be represented in upper-triangular form. There exists a basis of $\mathcal{V}$ such that the matrix of T is upper-triangular

$$\begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

where $n = \dim(\mathcal{V})$. The eigenvalues of T are exactly diagonal entries $\lambda_1, \dots, \lambda_n$, where each eigenvalue may repeat. By Theorem 1.2.131, the multiplicity of λ is number of times λ appears on diagonal, hence listing all diagonal entries counting repeats is listing all eigenvalues with their multiplicities.
The diagonal of triangular matrix has length $n = \dim(\mathcal{V})$, hence proving this. □

Definition 1.2.134. Let $\mathcal{V}$ be complex vector space, and $T \in \mathcal{L}(\mathcal{V})$. Let $\lambda_1, \ldots, \lambda_m$ be distinct eigenvalues of T, d_j be multiplicity of λ_j as an eigenvalue of T. The *characteristic polynomial* of T is $(z-\lambda_1)^{d_1} \cdots (z-\lambda_m)^{d_m}$.

Theorem 1.2.135. *Cayley-Hamilton Theorem*
Suppose $\mathcal{V}$ is complex vector space and $T \in \mathcal{L}(\mathcal{V})$, and q is characteristic polynomial of T. Then $q(T) = 0$.

Proof. Suppose $(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is basis of $\mathcal{V}$ with respect to which matrix of T has upper-triangular form

$$\mathcal{M}(T) = \begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

To show $q(T)\mathbf{v}_j = 0$ for $j = 1, \ldots, n$. As $q(x) = (x-\lambda_1)\cdots(x-\lambda_n)$, this reduces to showing

$$(T-\lambda_1 I)(T-\lambda_2 I)\cdots(T-\lambda_j I)\mathbf{v}_j = 0$$

for each j. For base case $j = 1$, since $T\mathbf{v}_1 = \lambda_1\mathbf{v}_1$, then $(T-\lambda_1 I)\mathbf{v}_1 = 0$.
Suppose claim holds for all $\mathbf{v}_1, \ldots, \mathbf{v}_{j-1}$. As T is upper-triangular in this basis, then

$$(T-\lambda_j I)\mathbf{v}_j \in \operatorname{span}(\mathbf{v}_1, \ldots, \mathbf{v}_{j-1})$$

By induction hypothesis, $(T-\lambda_1 I)\cdots(T-\lambda_{j-1}I)$ applied $(T-\lambda_j I)\mathbf{v}_j$ gives 0, hence completing the proof. □

Next proposition shows that the null space of any polynomial of $T \in \mathcal{L}(\mathcal{V})$ is invariant under T.

Proposition 1.2.136. *If $T \in \mathcal{L}(\mathcal{V})$ and $p \in \mathcal{P}(\mathbb{F})$, then null$(p(T))$ is invariant under T.*

Proof. Let $\mathbf{v} \in \operatorname{null}(p(T))$, then $p(T)\mathbf{v} = 0$, thus

$$(p(T))(T\mathbf{v}) = T(p(T)\mathbf{v}) = T(0) = 0$$

hence $T\mathbf{v} \in \operatorname{null}(p(T))$, thus $\operatorname{null}(p(T))$ is invariant under T as desired. □

Next theorem shows every operator on complex vector space can be composed of nilpotent operator plus scalar multiple of the identity.

Theorem 1.2.137. *Major Structure Theorem / Primary Decomposition Theorem*
Suppose $\mathcal{V}$ is a complex vector space and $T \in \mathcal{L}(\mathcal{V})$. Let $\lambda_1, \ldots, \lambda_m$ be distinct eigenvalues of T, and let $\mathcal{U}_1, \ldots, \mathcal{U}_m$ be corresponding subspaces of generalised eigenvalues. Then

i. $\mathcal{V} = \mathcal{U}_1 \oplus \cdots \oplus \mathcal{U}_m$
ii. each $\mathcal{U}_j$ is invariant under T
iii. each $(T-\lambda_j I)|_{\mathcal{U}_j}$ is nilpotent

Proof. (i) The multiplicity of λ_j is $\dim(\mathcal{U}_j)$. The sum of these multiplicities equal $\dim(\mathcal{U}_j)$, hence

$$\dim(\mathcal{V}) = \dim(\mathcal{U}_1) + \cdots + \dim(\mathcal{U}_m)$$

Let $\mathcal{U} = \mathcal{U}_1 + \cdots + \mathcal{U}_m$, clearly $\mathcal{U}$ is invariant under T. Define $S \in \mathcal{L}(\mathcal{U})$ by $S = T_{\mathcal{U}}$. Note that S has same eigenvalues with same multiplicities as T, as all generalised eigenvectors of T are in $\mathcal{U}$ (domain of S). Applying Proposition 1.2.133, then

$$\dim(\mathcal{U}) = \dim(\mathcal{U}_1) + \cdots + \dim(\mathcal{U}_m)$$

Hence $\dim(\mathcal{U}) = \dim(\mathcal{V})$. As $\mathcal{U}$ is a subspace of $\mathcal{V}$, this implies $\mathcal{U} = \mathcal{V}$, i.e., $\mathcal{V} = \mathcal{U}_1 + \cdots + \mathcal{U}_m$.
(ii). Note that $\mathcal{U}_j = \operatorname{null}(T-\lambda_j I)^{\dim(\mathcal{V})}$ for each j (by 1.2.127). By 1.2.136 (with $p(z) = (z-\lambda_j)^{\dim(\mathcal{V})}$).
(iii). This is obvious from the definitions. □

The following corollary shows that on a complex vector space there are enough generalised eigenvectors to form a basis for the domain of an operator.

Corollary 1.2.138. *Suppose $\mathcal{V}$ is a complex vector space and $T \in \mathcal{L}(\mathcal{V})$. Then there is a basis of $\mathcal{V}$ consisting of generalised eigenvectors of T.*

Proof. Choose a basis for each $\mathcal{U}_j$ in Theorem 1.2.137. Put all these bases together to form a basis of $\mathcal{V}$ consisting of generalised eigenvectors of T. □

Lemma 1.2.139. *Suppose N is nilpotent operator on $\mathcal{V}$, then $\exists$ a basis of $\mathcal{V}$ w.r.t. which matrix of N has form*

$$\begin{bmatrix} 0 & & * \\ & \ddots & \\ 0 & & 0 \end{bmatrix}$$

where all entries on and below diagonal are 0s.

Proof. Since N is nilpotent, $\exists$ some integer $m \geq 1$ such that $N^m = 0$. Hence $\exists$ ascending chain of subspaces

$$\{0\} \subseteq \text{null}(N) \subseteq \text{null}(N^2) \subseteq \cdots \subseteq \text{null}(N^m) = \mathcal{V}$$

Build a basis of $\mathcal{V}$ inductively along this chain. First choose a basis of null(N), then extend this to a basis of null(N^2), then to null(N^3) and so on. After finitely many steps, a basis of $\mathcal{V}$ is constructed that is adapted to the chain of subspaces

$$\text{null}(N) \subseteq \text{null}(N^2) \subseteq \cdots \subseteq \text{null}(N^m)$$

Examine matrix of N with respect to this basis.
The first set of basis vectors (in null(N)) are sent to 0 under N, hence the corresponding columns of the matrix are zero columns. The next set of basis vectors (those in null(N^2) but not in null(N)) are mapped by N into null(N), hence the image of these vectors are linear combinations of the earlier basis vectors. Thus the nonzero entries in these columns occur only in rows corresponding to earlier basis vectors (i.e., above the diagonal). Similarly, the next set of basis vectors (in null(N^3) but not in null(N^2)) are mapped to null(N^2), hence their images are linear combinations of basis vectors chosen earlier, hence nonzero entries in their columns lie above the diagonal. Continuing inductively, for each k, the basis vectors from null(N^k) are mapped by N into null(N^{k-1}), hence each column has nonzero entries only in rows corresponding to earlier basis vectors.
Hence, all diagonal entries of N are zero (since N is nilpotent, no vector is mapped to a nonzero multiple of itself). All entries below the diagonal are zero (since images of later basis vectors lie in the span of earlier ones). All possible nonzero entries occur strictly above diagonal. Hence matrix of N is strictly upper triangular. □

Theorem 1.2.140. *Triangular Form Theorem for Linear Operators over $\mathbb{C}$*
Suppose $\mathcal{V}$ is a complex vector space and $T \in \mathcal{L}(\mathcal{V})$. Let $\lambda_1, \ldots, \lambda_m$ be distinct eigenvalues of T. Then there exists a basis of $\mathcal{V}$ with respect to which the matrix of T is block diagonal of form

$$\begin{bmatrix} A_1 & & 0 \\ & \ddots & \\ 0 & & A_m \end{bmatrix}$$

where each A_j is an upper triangular matrix of form

$$A_j = \begin{bmatrix} \lambda_j & & * \\ & \ddots & \\ 0 & & \lambda_j \end{bmatrix}$$

Proof. For $j = 1, \ldots, m$, let $\mathcal{U})_j$ be subspace of generalised eigenvectors of T corresponding to λ_j. By Major Structure Theorem 1.2.137(c), $(T - \lambda_j I)|_{\mathcal{U}_j}$ is nilpotent. For each j, choose basis of $\mathcal{U}_j$ such that matrix of $(T - \lambda_j I)|_{\mathcal{U}_j}$ with respect to this basis is as seen in Lemma 1.2.139. Hence the matrix of $T|_{\mathcal{U}_j}$ will be as seen in A_j above. Putting together the basis for $\mathcal{U}_j$ gives a basis for $\mathcal{V}$, by Major Structure Theorem 1.2.137(a). The matrix of T with respect to this basis has the desired form. □

Lemma 1.2.141. *Suppose $N \in \mathcal{L}(\mathcal{V})$ is nilpotent, then $I + N$ has a square root.*

Proof. Let $m \geq 1$ with $N^m = 0$. Seek square rot of $I + N$ in form

$$S = I + a_1 N + a_2 N^2 + \cdots + a_{m-1} N^{m-1}$$

where $a_j \in \mathbb{C}$ are to be chosen (ignore higher powers as $N^m = 0$.
Let $a_0 = 1$. Using $N^i N^j = N^{i+j}$, expand

$$S^2 = \sum_{i=0}^{m-1} \sum_{j=0}^{m-1} a_i a_j N^{i+j} = \sum_{k=0}^{m-1} \left(\sum_{r=0}^{k} a_r a_{k-r} \right) N^k$$

as all terms with exponent $\geq m$ vanish. For each k, let $c_k = \sum_{r=0}^{k}$ be coefficient of N^k in S^2. To have

$$c_0 = 1, \quad c_1 = 1, \quad c_k = 0 \ (2 \leq k \leq m-1)$$

Note that $c_0 = a_0^2 = 1$ automatically. For $k \geq 1$, split off the terms with $r = 0, k$:

$$c_k = a_0 a_k + a_k a_0 + \sum_{r=1}^{k-1} a_r a_{k-r} = 2a_k + \sum_{r=1}^{k-1} a_r a_{k-r}$$

Hence to solve triangular system

$$2a_1 = 1, \quad 2a_k + \sum_{r=1}^{k-1} a_r a_{k-r} = 0 \ (2 \leq k \leq m-1)$$

Recursively define $a_1 = \frac{1}{2}, a_k = -\frac{1}{2}\sum_{r=1}^{k-1} a_r a_{k-r}$. As the right-hand side for a_k involves only $a_1, \ldots, a_{k-1}$, this uniquely determines a_k for all k. With these choices, then $c_1 = 1$, $c_k = 0$ for $2 \leq k \leq m-1$, therefore $S^2 = 1+N$. Hence S is a square root of $1 + N$. □

Theorem 1.2.142. *Let $\mathcal{V}$ be a complex vector space and $T \in \mathcal{L}(\mathcal{V})$. If T is invertible, then T has a square root.*

Proof. Let $\lambda_1, \ldots, \lambda_m$ be distinct eigenvalues of T, and $\mathcal{U}_j$ be the generalised eigenspace corresponding to λ_j. By Major Structure Theorem 1.2.137, for each j, there exists a nilpotent operator $N_j \in \mathcal{L}(\mathcal{U}_j)$ such that $T|_{\mathcal{U}_j} = \lambda_j I + N_j$. As T is invertible, 0 is not an eigenvalue, hence each $\lambda_j \neq 0$, thus

$$T|_{\mathcal{U}_j} = \lambda_j \left(I + \frac{N_j}{\lambda_j} \right)$$

Since N_j is nilpotent, so is N_j/λ_j, hence by Lemma 1.2.141, there exists operator $R_j \in \mathcal{L}(\mathcal{U}_j)$ with

$$R^2 = I + \frac{N_j}{\lambda_j}$$

on $\mathcal{U}_j$. Choose $\alpha_j \in \mathbb{C}$ with $\alpha_j^2 = \lambda_j$ (possible as the field is $\mathbb{C}$). Define $S_j = \alpha_j R_j \in \mathcal{L}(\mathcal{U}_j)$, then

$$S_j^2 = \alpha_j^2 R_j^2 = \lambda_j \left(I + \frac{N_j}{\lambda_j} \right) = T|_{\mathcal{U}_j}$$

Using direct sum decomposition $\mathcal{V} = \oplus_j \mathcal{U}_j$, each $\mathbf{v} \in \mathcal{V}$ can be written as $\mathbf{v} = \mathbf{u}_1 + \cdots + \mathbf{u}_m$ with $\mathbf{u} \in \mathcal{U}_j$. Define $S \in \mathcal{L}(\mathcal{V})$ by

$$S(\mathbf{v}) = S(\mathbf{u}_1 + \cdots + \mathbf{u}_m) = S_1\mathbf{u}_1 + \cdots + S_m\mathbf{u}_m$$

This is well-defined (the decomposition is unique) and linear (blockwise linearity).
As each $S_j(\mathcal{U}_j) \subseteq \mathcal{U}_j$, then $\mathbf{v} = \mathbf{u}_1 + \cdots + \mathbf{u}_m$,

$$S^2\mathbf{v} = S(S_1\mathbf{u}_1 + \cdots + S_m\mathbf{u}_m) = S_1^2\mathbf{u}_1 + \cdots + S_m^2\mathbf{u}_m = T|_{\mathcal{U}_1}\mathbf{u}_1 + \cdots + T|_{\mathcal{U}_m}\mathbf{u}_m = T(\mathbf{u}_1 + \cdots + \mathbf{u}_m) = T\mathbf{v}$$

Hence $S^2 = T$, and thus S is a square root of T. □

Definition 1.2.143. A *monic polynomial* is a non-zero univariate polynomial in which the leading coefficient equals 1, i.e., $x^n + c_{n-1}x^{n-1} + \cdots + c_2x^2 + c_1x + x_0$ with $n \geq 0$.

Definition 1.2.144. A *minimal polynomial* of T is monic polynomial $\mathbf{p} \in \mathcal{P}(\mathbb{F})$ of smallest degree s.t. $\mathbf{p}(T) = 0$.

Theorem 1.2.145. *Let $T \in \mathcal{L}(\mathcal{V})$ and let $\mathbf{q} \in \mathcal{P}(\mathbb{F})$, then $\mathbf{q}(T) = 0$ if and only if minimal polynomial of T divides $\mathbf{q}$.*

Proof. ($\Leftarrow$) Let $\mathbf{q}$ be minimal polynomial of T. Suppose $\mathbf{p}$ divides $\mathbf{q}$, hence there exists polynomial $\mathbf{s} \in \mathcal{P}(\mathbb{F})$ such that $\mathbf{q} = \mathbf{sp}$. Then

$$\mathbf{q}(T) = \mathbf{s}(T)\mathbf{p}(T) = \mathbf{s}(T)0 = 0$$

($\Rightarrow$) Suppose $\mathbf{q}(T) = 0$. By division algorithm, there exists polynomials $\mathbf{s}, \mathbf{r} \in \mathcal{P}(\mathbb{F})$ such that $\mathbf{q} = \mathbf{sp} + \mathbf{r}$ and $\deg(\mathbf{r}) < \deg(\mathbf{p})$, then

$$0 = \mathbf{q}(T) = \mathbf{s}(T)\mathbf{p}(T) + \mathbf{r}(T) = \mathbf{r}(T)$$

As $\mathbf{p}$ is minimal polynomial of T and $\deg(\mathbf{r}) < \deg(\mathbf{p})$, the equation above implies $\mathbf{r} = 0$. Hence equation becomes $\mathbf{q} = \mathbf{sp}$, hence $\mathbf{p}$ divides $\mathbf{q}$ as desired. □

Theorem 1.2.146. *Let $T \in \mathcal{L}(\mathcal{V})$, then the roots of minimal polynomial of T are precisely eigenvalues of T.*

Proof. ($\Rightarrow$) Let

$$\mathbf{p} = a_0 + a_1 z + a_2 z^2 + \cdots + a_{m-1} z^{m-1} + z^m$$

be minimal polynomial of T. Suppose $\lambda \in \mathbb{F}$ is a root of $\mathbf{p}$, then the polynomial of T is $\mathbf{p}(z) = (z - \lambda)\mathbf{q}(z)$, where $\mathbf{q}$ is a monic polynomial with coefficients in $\mathbb{F}$. As $\mathbf{p}(T) = 0$, then

$$0 = (T - \lambda I)(\mathbf{q}(T)\mathbf{v})$$

for all $\mathbf{v} \in \mathcal{V}$. As $\deg(\mathbf{q}) < \deg(\mathbf{q})$, there must exist at least one vector $\mathbf{v} \in \mathcal{V}$ such that $\mathbf{q}(T)\mathbf{v} \neq 0$. This implies λ is an eigenvalue of T as desired.
($\Leftarrow$) Suppose $\lambda \in \mathbb{F}$ is an eigenvalue of T. Let $\mathbf{v}$ be nonzero vector in $\mathcal{V}$ such that $T\mathbf{v} = \lambda\mathbf{v}$, Repeated application of T to both sides of equation show $T^j\mathbf{v} = \lambda^j\mathbf{v}$ for every nonnegative integer j. Hence

$$\begin{aligned} 0 = \mathbf{p}(T)\mathbf{v} &= (a_0 + a_1 T + a_2 T^2 + \cdots + a_{m-1} T^{m-1} + T^m)\mathbf{v} \\ &= (a_0 + a_1 \lambda + a_2 \lambda^2 + \cdots + a_{m-1} \lambda^{m-1} + \lambda^m)\mathbf{v} \\ &= \mathbf{p}(\lambda)\mathbf{v} \end{aligned}$$

As $\mathbf{v} \neq \mathbf{0}$, this implies $\mathbf{p}(\lambda) = 0$ as desired. □

References

Axler, S. (2024). *Linear Algebra Done Right* (4 ed.). Undergraduate Texts in Mathematics. Cham: Springer.

Garcia, S. R. and R. A. Horn (2017). *A Second Course in Linear Algebra*. Cambridge: Cambridge University Press.