

## Accepted Manuscript

Dissipativity and passivity analysis for memristor-based neural networks with leakage and two additive time-varying delays


Qianhua Fu, Jingye Cai, Shouming Zhong, Yongbin Yu

PII: S0925-2312(17)31509-6  
DOI: [10.1016/j.neucom.2017.09.014](https://doi.org/10.1016/j.neucom.2017.09.014)  
Reference: NEUCOM 18872

To appear in: *Neurocomputing*

Received date: 19 May 2017  
Revised date: 8 August 2017  
Accepted date: 4 September 2017

Please cite this article as: Qianhua Fu, Jingye Cai, Shouming Zhong, Yongbin Yu, Dissipativity and passivity analysis for memristor-based neural networks with leakage and two additive time-varying delays, *Neurocomputing* (2017), doi: [10.1016/j.neucom.2017.09.014](https://doi.org/10.1016/j.neucom.2017.09.014)



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Dissipativity and passivity analysis for memristor-based neural networks with leakage and two additive time-varying delays<sup>☆</sup>

Qianhua Fu<sup>a,b,\*</sup>, Jingye Cai<sup>a</sup>, Shouming Zhong<sup>c</sup>, Yongbin Yu<sup>a</sup>

<sup>a</sup>*School of Information and Software Engineering, University of Electronic Science and Technology of China, Chengdu 610054, P.R. China*

<sup>b</sup>*School of Electrical Engineering and Electronic Information, Xihua University, Chengdu 610039, P.R. China*

<sup>c</sup>*School of Mathematical Sciences, University of Electronic Science and Technology of China, Chengdu 611731, P.R. China*

## Abstract

In this paper, the problems of dissipativity and passivity analysis for memristor-based neural networks (MNNs) with both time-varying leakage delay and two additive time-varying delays are studied. By introducing an improved Lyapunov-Krasovskii functional (LKF) with triple integral terms, and combining the reciprocally convex combination technique, Wirtinger-based integral inequality with free-weighting matrices technique, some less conservative delay-dependent dissipativity and passivity criteria are obtained. The proposed criteria that depend on the upper bounds of the leakage and additive time-varying delays are given in terms of linear matrix inequalities (LMI), which can be solved by MATLAB LMI Control Toolbox. Meanwhile, the criteria for the system with a single time-varying delay are also provided. Finally, some examples are given to illustrate the effectiveness and superiority of the obtained results.

**Keywords:** memristor-based neural networks, leakage delays, additive time-varying delays, dissipativity, passivity

<sup>☆</sup>This work was supported in part by the National Natural Science Foundation of China under Grants 61533006, in part by the Research Fund for International Young Scientists of National Natural Science Foundation of China under Grant 61550110248.

\*Corresponding author

Email addresses: qhfu8@mail.xhu.edu.cn (Qianhua Fu), jycai@uestc.edu.cn (Jingye Cai), zhongsm@uestc.edu.cn (Shouming Zhong), ybyu@uestc.edu.cn (Yongbin Yu)

## 1. Introduction

Memristor was postulated first by Chua in 1971 as the fourth basic element of electrical circuits [1]. About 37 years later, HP researchers showed that memristance arises naturally in nanoscale systems in which solid-state electronic and ionic transport are coupled under an external bias voltage [2]. Compared with ordinary resistor, the value of memristor is not unique and it hinges on the voltage applied to the corresponding history and current state [3]. Memristor owns a lot of characteristics, such as good scalability, a high density and low power [4]. Therefore, the nonvolatile temperament of memristor makes it a noticeable candidate for the next-generation memory technology [5, 6, 7]. By substituting resistors in artificial neural networks, a class of memristor-based neural networks (MNNs) can be constructed. This kinds of system can exhibit intelligent behaviors [8], such as cognition and simulate forgetting mechanism of neuromorphic system [9].

Recently most of the researchers have paid attention to know about the dynamical behaviors of MNNs [10, 11, 12, 13, 14]. In some practical applications, owing to the restricted speed of information processing, the reality of time delays frequently causes oscillation, divergence, or instability in neural networks [15]. Signals transmitted from one point to another may experience a few segments of networks, which can induce successive delays with different properties due to variable network transmission conditions [16]. A typical time delay called leakage delay may exist in the negative feedback terms of the system and these terms are variously known as forgetting or leakage terms. Since leakage delays have a destabilizing influence on the dynamical behaviors of MNNs, it is necessary and important to consider the leakage delay effects on the study of state estimation of MNNs, however to the best of our knowledge, there are only a few relative works on MNNs with leakage delay [17, 18, 19]. Thus, the stability problem of MNNs with leakage and additive time-varying delays has become a topic of great theoretical and practical importance [20].

30 The dissipativity and passivity problems for a spread of sensible systems are attracting researchers' attention for several years [21, 22, 23]. It is well-known that dissipativity analysis contains other dynamic analysis such as passivity analysis,  $H_\infty$  analysis, and so on [24, 25]. The problem of strict  $(Q, S, R) - \gamma -$  dissipativity analysis for memristive neural networks with a constant leakage delay and a single time-varying delay is studied in [26]. The quadratical stability and extended dissipative conditions for the memristive neural networks with two additive time-varying delays have been proposed in [27]. The problem of passivity analysis of stochastic neural networks with leakage delay and Markovian jumping parameters is considered in [28]. X. Li and R. Rakkiyappan et al. [29] have investigated extensively the problem of dissipativity analysis for memristor-based complex-valued neural networks with time-varying delays. S. Ding and Z. Wang et al. [30] have investigated the dissipativity problem for a new array of discrete-time memristive neural networks with time-varying delays by defining a set of logical switched signals and utilizing robust analysis method.

45 This paper further investigates the dissipativity and passivity analysis for MNNs. The main contributions of this paper are summarized as follows:

1. The considered MNNs in this paper include not only additive time-varying delays, but also time-varying leakage delay, so it is firstly studied in the present paper, the model of MNNs is more general than ever.
- 50 2. By constructing a new Lypunov-Krasovskii functional (LKF) and using the second-order reciprocally convex combination technique, Wirtinger-based integral inequality with free-weighting matrices technique, some less conservative delay-dependent dissipativity and passivity criteria are obtained.
- 55 3. Our result is more general and it is valid for the usual neural networks model. Many relative works are included in this paper.

The remaining part of this paper is organized as follows. In Section 2, the model of MNNs is formulated by applying the theory of set-valued maps and functional differential inclusions, some predictable assumptions on the bound-

edness and Lipschitz continuity of activation functions are formulated base on the knowledge of memristor. In Section 3, some less conservative dissipativity and passivity criteria are established. Two numerical simulation examples are provided to validate all of the theoretical results in Section 4. Finally, the conclusion is drawn in Section 5.

**Notation:** The notations used throughout this paper are standard.  $\mathbb{R}^n$  and  $\mathbb{R}^{m \times n}$ , respectively, denote the  $n$ -dimensional Euclidean space and the set of all  $m \times n$  real matrices.  $\|\cdot\|$  refers to the Euclidean vector norm.  $\text{diag}\{\dots\}$  is the block diagonal matrix. The superscript  $T$  is the transposition and the notation  $P > 0$  ( $P \geq 0$ ) means that  $P$  is the real symmetric matrix and positive definite (semidefinite).  $\mathbf{I}$  and  $\mathbf{O}$  are the identity matrix and zero matrix with appropriate dimensions, respectively. The notation  $\star$  is the symmetric block in one symmetric matrix, and  $\mathbb{C}$  is the set of continuous functions.  $\text{co}\{\Pi_1, \Pi_2\}$  represents closure of the convex hull generated by real matrices  $\Pi_1$  and  $\Pi_2$  or real numbers  $\Pi_1$  and  $\Pi_2$ .  $\mathcal{L}_2^n$  is the space of square integrable functions on  $\mathbb{R}^+$  with values in  $\mathbb{R}^n$ .  $\mathcal{L}_{2e}^n$  is the extended  $\mathcal{L}_2^n$  space defined by  $\mathcal{L}_{2e}^n = \{f : f \text{ is a measurable function on } \mathbb{R}^+, P_T f \in \mathcal{L}_2^n, \forall T \in \mathbb{R}^+\}$ , where  $(P_T f)(t) = f(t)$  if  $t \leq T$ , and 0 if  $t > T$ . For any function  $x = \{x(t)\}, y = \{y(t)\} \in \mathcal{L}_{2e}^n$ , matrix  $Q$ , we define  $\langle x, Qy \rangle_T = \int_0^T x^T(t) Q y(t) dt$ .

## 2. Problem Description Preliminaries

Consider the following class of MNNs with leakage and two additive time-varying delays:

$$\begin{cases} \frac{dx_i(t)}{dt} = -d_i(t)x_i(t - \eta_i(t)) + \sum_{j=1}^n a_{ij}(t)f_j(x_j(t)) \\ \quad + \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_{j1}(t) - \tau_{j2}(t))) + u_i(t) \\ y_i(t) = f_i(x_i(t)), i = 1, 2, \dots, n. \end{cases} \quad (1)$$

where  $n$  is the amount of neuronal cell in a neural networks;  $x_i(t) \in \mathbb{R}$  is the  $i$ th neuronal cell state;  $y_i(t) \in \mathbb{R}$  is the output of the  $i$ th neuronal cell;

$d_i(t) > 0$  represents the rate which the  $i$ th neuronal cell will reset its potential to the resting state in isolation when disconnected from the network and external inputs at time  $t$ ;  $\tau_{j1}(t)$  and  $\tau_{j2}(t)$  are the two additive time varying delays that are assumed to satisfy the conditions  $0 \leq \tau_{j1}(t) \leq \tau_1, 0 \leq \tau_{j2}(t) \leq \tau_2, |\dot{\tau}_{j1}(t)| \leq \mu_1, |\dot{\tau}_{j2}(t)| \leq \mu_2$ ;  $\eta_i(t)$  denotes the leakage delay satisfying  $0 \leq \eta_i(t) \leq \eta, |\dot{\eta}_i(t)| \leq \mu = \mu_1 + \mu_2$ ;  $\eta, \tau_1, \tau_2, \mu_1, \mu_2, \mu$  are nonnegative constants;  $f_j(\cdot)$  and  $g_j(\cdot)$  are activation functions;  $u_i(t)$  is the controller to be designed;  $a_{ij}(t)$  and  $b_{ij}(t)$  represent the memristive synaptic weights, denote the strengths of the  $j$ th neuronal cell on the  $i$ th neuronal cell at time  $t$ , which are defined as follows:

$$a_{ij}(t) = \begin{cases} \hat{a}_{ij}, & \dot{\kappa}_{ij}^1(t) < 0 \\ \text{unchanged}, & \dot{\kappa}_{ij}^1(t) = 0 \\ \check{a}_{ij}, & \dot{\kappa}_{ij}^1(t) > 0 \end{cases}$$

$$b_{ij}(t) = \begin{cases} \hat{b}_{ij}, & \dot{\kappa}_{ij}^2(t) < 0 \\ \text{unchanged}, & \dot{\kappa}_{ij}^2(t) = 0 \\ \check{b}_{ij}, & \dot{\kappa}_{ij}^2(t) > 0 \end{cases}$$

80 where  $\kappa_{ij}^1(t) = f_{ij}(x_j(t)) - x_i(t)$ ,  $\kappa_{ij}^2(t) = g_{ij}(x_j(t - \tau_{j1}(t) - \tau_{j2}(t))) - x_i(t)$ ,  $\kappa_{ij}^1(t)$  and  $\kappa_{ij}^2(t)$  represent the voltage of the corresponding memresistance.  $\hat{a}_{ij}$ ,  $\check{a}_{ij}$ ,  $\hat{b}_{ij}$ ,  $\check{b}_{ij}$  are constants for  $i, j = 1, 2, \dots, n$ . The initial values associated with system (1) are  $x_i(t) = \varphi_i(t) \in \mathbb{C}([-\tau^*, 0], \mathbb{R})$ ,  $\tau^* = \max\{\eta, \tau_1 + \tau_2\}$  for  $i = 1, 2, \dots, n$ .

85 **Remark 1.** In some existing literatures [18, 31, 32, 33, 34, 35], the memristor is just looked as a simple switch, its state switching depends on the potential  $x_i(t)$  which is connected to one port of the memristor, if  $x_i(t)$  is more or less than the switching jumps  $T_i$ , the memristive synaptic weights are switched. But the memristor states depend on the voltage of its two ports and the voltage variation trend [3, 36]. So we use the derivative of  $\kappa_{ij}^1(t)$  and  $\kappa_{ij}^2(t)$  to switch the  $a_{ij}(t)$  and  $b_{ij}(t)$ , the model of MNNs is more general than ever.

90 **Remark 2.** The dissipativity and passivity problems of MNNs have received much of the focus in recent years. So far, a great many important results on

dissipativity or passivity have been obtained for MNNs. Unfortunately, The  
 95 references [29, 30] just considered a single time-varying delay. The references  
 [37, 38] considered additive time-varying delays, but didn't include leakage delay.  
 The references [39, 40] considered a leakage delay, but the leakage delay is a  
 constant. Practically, the leakage delay and multiple signal transmission delays  
 coexist in the system of MNNs.

100 For convenience, we introduce the following Definitions about set-valued map  
 and differential inclusion.

**Definition 1 [41].** Let  $E \subset \mathbb{R}^n$ ,  $x \mapsto F(x)$  is called a set-valued map from  
 $E \rightarrow \mathbb{R}^n$ , if to each point  $x$  of a set  $E \subset \mathbb{R}^n$ , there corresponds a nonempty set  
 $F(x) \subset \mathbb{R}^n$ .

105 **Definition 2 [42].** A set-valued map  $F$  with nonempty values is said to be  
 upper semi-continuous at  $x_0 \in E \subset \mathbb{R}^n$ , if for any open set  $N$  containing  $F(x_0)$ ,  
 there exists a neighborhood  $M$  of  $x_0$  such that  $F(M) \subset N$ .  $F(x)$  is said to have  
 a closed (convex, compact) image if for each  $x \in E$ ,  $F(x)$  is closed (convex,  
 compact).

By applying the above theories of set-valued maps and differential inclusions,  
 the MNNs (1) can be written as the following differential inclusion:

$$\begin{cases} \frac{dx_i(t)}{dt} \in -\underline{d_i(t)}x_i(t - \eta_i(t)) + \sum_{j=1}^n co[a_{ij}^-, a_{ij}^+]f_j(x_j(t)) \\ \quad + \sum_{j=1}^n co[b_{ij}^-, b_{ij}^+]g_j(x_j(t - \tau_{j1}(t) - \tau_{j2}(t))) + u_i(t) \\ y_i(t) = f_i(x_i(t)), i = 1, 2, \dots, n. \end{cases} \quad (2)$$

where  $a_{ij}^- = \min\{\hat{a}_{ij}, \check{a}_{ij}\}$ ,  $a_{ij}^+ = \max\{\hat{a}_{ij}, \check{a}_{ij}\}$ ,  $b_{ij}^- = \min\{\hat{b}_{ij}, \check{b}_{ij}\}$ ,  $b_{ij}^+ = \max\{\hat{b}_{ij}, \check{b}_{ij}\}$ . There exist  $\bar{a}_{ij}(t) \in co[a_{ij}^-, a_{ij}^+]$ ,  $\bar{b}_{ij}(t) \in co[b_{ij}^-, b_{ij}^+]$  satisfy

$$\begin{cases} \frac{dx_i(t)}{dt} = -d_i(t)x_i(t - \eta_i(t)) + \sum_{j=1}^n \bar{a}_{ij}(t)f_j(x_j(t)) \\ \quad + \sum_{j=1}^n \bar{b}_{ij}(t)g_j(x_j(t - \tau_{j1}(t) - \tau_{j2}(t))) + u_i(t) \\ y_i(t) = f_i(x_i(t)) \end{cases} \quad (3)$$

for a.e.  $t \in [0, \infty)$ ,  $i = 1, 2, \dots, n$ .

Now the (3) can be rewritten as the following matrix form:

$$\begin{cases} \dot{x}(t) = -Dx(t - \eta(t)) + Af(x(t)) + Bg(x(t - \tau_1(t) - \tau_2(t))) + u(t) \\ \underline{y(t) = f(x(t))} \end{cases} \quad (4)$$

where  $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T$ ,  $D = \text{diag}\{d_1(t), d_2(t), \dots, d_n(t)\}$ ,  $A = [\bar{a}_{ij}(t)]_{n \times n}$ ,  $B = [\bar{b}_{ij}(t)]_{n \times n}$ ,  $f(x(t)) = (f_1(x_1(t)), f_2(x_2(t)), \dots, f_n(x_n(t)))^T$ ,  $x(t - \eta(t)) = (x_1(t - \eta_1(t)), x_2(t - \eta_2(t)), \dots, x_n(t - \eta_n(t)))^T$ ,  $g(x(t - \tau_1(t) - \tau_2(t))) = (g_1(x_1(t - \tau_{11}(t) - \tau_{12}(t))), g_2(x_2(t - \tau_{21}(t) - \tau_{22}(t))), \dots, g_n(x_n(t - \tau_{n1}(t) - \tau_{n2}(t))))^T$ ,  $u(t) = (u_1(t), u_2(t), \dots, u_n(t))^T$ .

The time-varying delay  $\tau(t) = \tau_1(t) + \tau_2(t)$  satisfying the conditions:

$$0 \leq \tau_1(t) \leq \tau_1; 0 \leq \tau_2(t) \leq \tau_2;$$

$$|\dot{\tau}_1(t)| \leq \mu_1; |\dot{\tau}_2(t)| \leq \mu_2;$$

where  $\tau_1, \tau_2, \mu_1, \mu_2$  are nonnegative constants, we denote  $\tau = \tau_1 + \tau_2$ ,  $\mu = \mu_1 + \mu_2$ .

To derive the main results, we introduce the following assumptions, definitions and lemmas.

**Assumption 1 [43].** For all  $\alpha, \beta \in \mathbb{R}$  and  $\alpha \neq \beta$ ,  $i \in \{1, 2, \dots, n\}$ , the activation functions  $f, g$  are bounded and there exist four constant matrices  $L^- = \text{diag}\{l_1^-, l_2^-, \dots, l_n^-\}$ ,  $L^+ = \text{diag}\{l_1^+, l_2^+, \dots, l_n^+\}$ ,  $H^- = \text{diag}\{h_1^-, h_2^-, \dots, h_n^-\}$ ,  $H^+ = \text{diag}\{h_1^+, h_2^+, \dots, h_n^+\}$  such that

$$l_i^- \leq \frac{f_i(\alpha) - f_i(\beta)}{\alpha - \beta} \leq l_i^+ \quad \text{这个条件比Lipschitz条件弱吗?}$$

$$h_i^- \leq \frac{g_i(\alpha) - g_i(\beta)}{\alpha - \beta} \leq h_i^+$$

where  $l_i^-, l_i^+, h_i^-$  and  $h_i^+$  are constants,  $f_i(0) = 0, g_i(0) = 0$ .

**Remark 3.** Assumption 1 limits the lower and upper bounds of the activation functions slope, it is weaker than the well-known Lipschitz-type conditions (i.e.  $|f_i(\alpha) - f_i(\beta)| \leq l_i|\alpha - \beta|$ ,  $|g_i(\alpha) - g_i(\beta)| \leq h_i|\alpha - \beta|$ ,  $l_i = \max\{|l_i^-|, |l_i^+|\}$ ,  $h_i = \max\{|h_i^-|, |h_i^+|\}$ ), which is used in [25, 29].



**Definition 3 [39].** Given real symmetric matrices  $Q$ ,  $R$  and  $S$  with appropriate dimensions, the system (4) is called strictly  $(Q, S, R) - \gamma - \text{dissipative}$ , if there exists a scalar  $\gamma > 0$  such that the inequality

$$\langle y, Qy \rangle_T + 2\langle y, Su \rangle_T + \langle u, Ru \rangle_T \geq \gamma \langle u, u \rangle_T, \forall u \in \mathcal{L}_{2e}^n \quad (5)$$

hold for all  $t \geq 0$  and under the zero initial condition.

**Definition 4 [37].** The system (4) is said to be passive, if for all solutions of (4), there exists a scalar  $\gamma > 0$  such that the inequality

$$2 \int_0^T y^T(s)u(s)ds \geq -\gamma \int_0^T u^T(s)u(s)ds \quad (6)$$

is satisfied under the zero initial condition.

**Lemma 1 [44].** Consider a given matrix  $R > 0$ . Then, for all continuous function  $\omega$  in  $[a, b] \rightarrow \mathbb{R}^n$  the following inequality holds:

$$\begin{aligned} \int_a^b \omega^T(u)R\omega(u)du &\geq \frac{1}{b-a} \left( \int_a^b \omega(u)du \right)^T R \int_a^b \omega(u)du + \frac{3}{b-a} \Omega^T R \Omega \\ &\geq \frac{1}{b-a} \left( \int_a^b \omega(u)du \right)^T R \int_a^b \omega(u)du \end{aligned} \quad (7)$$

where  $\Omega = \int_a^b \omega(s)ds - \frac{2}{b-a} \int_a^b \int_a^s \omega(r)drds$ .

**Lemma 2 [45].** For any constant matrix  $W \in \mathbb{R}^{n \times n}$ , two scalars  $b \geq a \geq 0$  and vector function  $x(s) : [a, b] \rightarrow \mathbb{R}^n$ , such that the following integration is well defined, then:

$$\begin{aligned} &-\frac{(b^2 - a^2)}{2} \int_{-b}^{-a} \int_{t+\theta}^t x^T(s)Wx(s)dsd\theta \\ &\leq -\left( \int_{-b}^{-a} \int_{t+\theta}^t x(s)dsd\theta \right)^T W \int_{-b}^{-a} \int_{t+\theta}^t x(s)dsd\theta \end{aligned} \quad (8)$$

**Lemma 3 [46].** Let  $f_1, f_2, \dots, f_N : \mathbb{R}^m \rightarrow \mathbb{R}$  have positive values in an open subset  $\mathbf{D}$  of  $\mathbb{R}^m$ . Then, the reciprocally convex combination of  $f_i$  over  $\mathbf{D}$  satisfies

$$\min_{\{a_i | a_i > 0, \sum_i a_i = 1\}} \frac{1}{a_i} f_i(t) = \sum_i f_i(t) + \max_{g_{i,j}(t)} \sum_{i \neq j} g_{i,j}(t) \quad (9)$$

Subject to

$$\{g_{i,j} : \mathbb{R}^m \rightarrow \mathbb{R}, g_{j,i} \triangleq g_{i,j}(t), \begin{bmatrix} f_i(t) & g_{i,j}(t) \\ g_{i,j}(t) & f_j(t) \end{bmatrix} \geq 0\}$$

**Remark 4.** Lemma 1 shows the Wirtinger-based integral inequality and Jensen inequality. The Wirtinger-based integral inequality delivers a more accurate lower bound of  $\int_a^b w^T(u)Rw(u)du$  so it reduces the conservatism of the Jensen inequality. Lemma 2 gets from Jensen inequality and Schur Complements, which can be used to deal with multiple integral terms. Lemma 3 (The reciprocally convex combination technique) achieves performance behavior identical to approaches based on the integral inequality but also decreases the number of decision variables dramatically up to those based on the Jensen inequality.

### 3. Main results

In this section, by using the Lyapunov functional method and LMI technique, we will derive the delay-dependent dissipativity criterion for the MNNs (4) in the following theorem.

**Theorem 1:** Given real symmetric matrices  $Q$ ,  $R$  and  $S$  with appropriate dimensions, under Assumption 1, the MNNs (4) is dissipative in the sense of Definition 3, if there exist symmetric matrices  $P > 0$ ,  $Q_i > 0$ ,  $R_i > 0$ ,  $T_i > 0$ , ( $i = 1, 2, 3, 4$ ),  $U_j > 0$  ( $j = 1, 2, 3$ ),  $Z > 0$ , diagonal matrices  $M = \text{diag}\{m_1, m_2, \dots, m_n\} > 0$ ,  $W = \text{diag}\{w_1, w_2, \dots, w_n\} > 0$ ,  $\Lambda_1 = \text{diag}\{\lambda_{11}, \lambda_{12}, \dots, \lambda_{1n}\} > 0$ ,  $\Lambda_2 = \text{diag}\{\lambda_{21}, \lambda_{22}, \dots, \lambda_{2n}\} > 0$ , and matrices  $G_i$  ( $i = 1, 2, 3, 4$ ),  $V_j$  ( $j = 1, 2, 3$ ) and scalar  $\gamma > 0$ , such that the following LMIs hold:

$$\Phi_l = \Xi - \Upsilon_l^T \begin{bmatrix} U_1 & V_1 & 0 & 0 & 0 & 0 \\ * & U_1 & 0 & 0 & 0 & 0 \\ * & * & U_2 & V_2 & 0 & 0 \\ * & * & * & U_2 & 0 & 0 \\ * & * & * & * & U_3 & V_3 \\ * & * & * & * & * & U_3 \end{bmatrix} \Upsilon_l \leq 0, \quad (10)$$

$l = (1, 2, 3, 4).$

$$\begin{bmatrix} T_i & G_i \\ \star & T_i \end{bmatrix} \geq 0, (i = 1, 2, 3, 4). \quad (11)$$

$$\begin{bmatrix} U_j & V_j \\ \star & U_j \end{bmatrix} \geq 0, (j = 1, 2, 3). \quad (12)$$

where

140

$$\Xi = \begin{bmatrix} F_{11} & F_{12} & F_{13} & F_{14} \\ \star & F_{22} & F_{23} & F_{24} \\ \star & \star & F_{33} & F_{34} \\ \star & \star & \star & F_{44} \end{bmatrix},$$

$$F_{11} = \begin{bmatrix} \Psi_{1,1} & \Psi_{1,2} & \Psi_{1,3} & \Psi_{1,4} & -2G_1^T & \Psi_{1,6} \\ \star & \Psi_{2,2} & \Psi_{2,3} & 0 & 0 & 0 \\ \star & \star & \Psi_{3,3} & 0 & 0 & 0 \\ \star & \star & \star & \Psi_{4,4} & \Psi_{4,5} & 0 \\ \star & \star & \star & \star & \Psi_{5,5} & 0 \\ \star & \star & \star & \star & \star & \Psi_{6,6} \end{bmatrix},$$

145

$$F_{12} = \begin{bmatrix} -2G_2^T & \Psi_{1,8} & -2G_4^T & \Psi_{1,10} & PB \\ 0 & 0 & 0 & \Psi_{2,10} & \Psi_{2,11} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \Psi_{6,7} & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$F_{13} = \begin{bmatrix} 6T_3 & 6G_3^T & 6T_1 & 6G_1^T & 6T_2 \\ \Psi_{2,12} & \Psi_{2,13} & 0 & 0 & 0 \\ 6G_3 & 6T_3 & 0 & 0 & 0 \\ 0 & 0 & \Psi_{4,14} & \Psi_{4,15} & 0 \\ 0 & 0 & 6G_1 & 6T_1 & 0 \\ 0 & 0 & 0 & 0 & \Psi_{6,16} \end{bmatrix},$$

$$F_{14} = \begin{bmatrix} 6G_2^T & 6T_4 & 6G_4^T & MPM & P \\ 0 & 0 & 0 & DPM & \Psi_{2,21} \\ 0 & 0 & 0 & -MPM & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \Psi_{6,17} & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$F_{22} = \begin{bmatrix} \Psi_{7,7} & 0 & 0 & 0 & 0 \\ * & \Psi_{8,8} & \Psi_{8,9} & 0 & \Psi_{8,11} \\ * & * & \Psi_{9,9} & 0 & 0 \\ * & * & * & \Psi_{10,10} & \Psi_{10,11} \\ * & * & * & * & \Psi_{11,11} \end{bmatrix},$$

$$F_{23} = \begin{bmatrix} 0 & 0 & 0 & 0 & 6G_2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$F_{24} = \begin{bmatrix} 6T_2 & 0 & 0 & 0 & 0 \\ 0 & \Psi_{8,18} & \Psi_{8,19} & 0 & 0 \\ 0 & 6G_4 & 6T_4 & 0 & 0 \\ 0 & 0 & 0 & \Psi_{10,20} & \Psi_{10,21} \\ 0 & 0 & 0 & \Psi_{11,20} & \Psi_{11,21} \end{bmatrix},$$

$$F_{33} = \begin{bmatrix} -12T_3 & -12G_3^T & 0 & 0 & 0 \\ * & -12T_3 & 0 & 0 & 0 \\ * & * & -12T_1 & -12G_1^T & 0 \\ * & * & * & -12T_1 & 0 \\ * & * & * & * & -12T_2 \end{bmatrix},$$

$$F_{34} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -12G_2^T & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$F_{44} = \begin{bmatrix} -12T_2 & 0 & 0 & 0 & 0 \\ * & -12T_4 & -12G_4^T & 0 & 0 \\ * & * & -12T_4 & 0 & 0 \\ * & * & * & -Z & -MP \\ * & * & * & * & \Psi_{21,21} \end{bmatrix},$$

$$\begin{aligned} \Psi_{1,1} &= +PM - MP + Q_1 + Q_2 + Q_3 + Q_4 + R_1 + R_2 + R_3 + R_4 + \eta^2 Z - 4T_1 - \\ &4T_2 - 4T_3 - 4T_4 - 2L^- \Lambda_1 L^+, \Psi_{1,2} = -PD - 2T_3 - 4G_3^T, \Psi_{1,3} = PM - 2G_3^T, \\ \Psi_{1,4} &= -2T_1 - 4G_1^T, \Psi_{1,6} = -2T_2 - 4G_2^T, \Psi_{1,8} = -2T_4 - 4G_4^T, \Psi_{1,10} = \\ &PA + (L^- + L^+) \Lambda_1, \Psi_{2,2} = -(1 - \mu)R_4 + \tau_1^2 DT_1 D + \tau_2^2 DT_2 D + \eta^2 DT_3 D + \\ &\tau^2 DT_4 D + \frac{\tau_1^4}{4} DU_1 D + \frac{\tau_2^4}{4} DU_2 D + \frac{\tau_4^4}{4} DU_3 D - 8T_3 - 2(G_3 + G_3^T), \Psi_{2,3} = -2T_3 - \\ &4G_3^T, \Psi_{2,10} = -DW - \tau_1^2 DT_1 A - \tau_2^2 DT_2 A - \eta^2 DT_3 A - \tau^2 DT_4 A - \frac{\tau_1^4}{4} DU_1 A - \\ &\frac{\tau_2^4}{4} DU_2 A - \frac{\tau_4^4}{4} DU_3 A, \Psi_{2,11} = -\tau_1^2 DT_1 B - \tau_2^2 DT_2 B - \eta^2 DT_3 B - \tau^2 DT_4 B - \end{aligned}$$

$$\begin{aligned}
 & \frac{\tau_1^4}{4} DU_1 B - \frac{\tau_2^4}{4} DU_2 B - \frac{\tau_4^4}{4} DU_3 B, \Psi_{2,12} = 6T_3 + 6G_3, \Psi_{2,13} = 6T_3 + 6G_3^T, \\
 & \Psi_{2,21} = -\tau_1^2 DT_1 - \tau_2^2 DT_2 - \eta^2 DT_3 - \tau^2 DT_4 - \frac{\tau_1^4}{4} DU_1 - \frac{\tau_2^4}{4} DU_2 - \frac{\tau_4^4}{4} DU_3, \\
 & \Psi_{3,3} = -Q_4 - 4T_3, \Psi_{4,4} = -(1 - \mu_1)R_1 - 8T_1 - 2(G_1 + G_1^T), \Psi_{4,5} = -2T_1 - 4G_1^T, \\
 & \Psi_{4,14} = 6T_1 + 6G_1, \Psi_{4,15} = 6T_1 + 6G_1^T, \Psi_{5,5} = -Q_1 - 4T_1, \Psi_{6,6} = -(1 - \mu_2)R_2 - \\
 & 8T_2 - 2(G_2 + G_2^T), \Psi_{6,7} = -2T_2 - 4G_2^T, \Psi_{6,16} = 6T_2 + 6G_2, \Psi_{6,17} = 6T_2 + 6G_2^T, \\
 & \Psi_{7,7} = -Q_2 - 4T_2, \Psi_{8,8} = -(1 - \mu)R_3 - 2H^- \Lambda_2 H^+ - 8T_4 - 2(G_4 + G_4^T), \\
 & \Psi_{8,9} = -2T_4 - 4G_4^T, \Psi_{8,11} = (H^- + H^+) \Lambda_2, \Psi_{8,18} = 6T_4 + 6G_4, \Psi_{8,19} = \\
 & 6T_4 + 6G_4^T, \Psi_{9,9} = -Q_3 - 4T_4, \Psi_{10,10} = AW + WA^T + \tau_1^2 A^T T_1 A + \tau_2^2 A^T T_2 A + \\
 & \eta^2 A^T T_3 A + \tau^2 A^T T_4 A + \frac{\tau_1^4}{4} A^T U_1 A + \frac{\tau_2^4}{4} A^T U_2 A + \frac{\tau_4^4}{4} A^T U_3 A - 2\Lambda_1 - Q, \Psi_{10,11} = \\
 & WB^T + \tau_1^2 A^T T_1 B + \tau_2^2 A^T T_2 B + \eta^2 A^T T_3 B + \tau^2 A^T T_4 B + \frac{\tau_1^4}{4} A^T U_1 B + \frac{\tau_2^4}{4} A^T U_2 B + \\
 & \frac{\tau_4^4}{4} A^T U_3 B, \Psi_{10,20} = -A^T PM, \Psi_{10,21} = W + \tau_1^2 A^T T_1 + \tau_2^2 A^T T_2 + \eta^2 A^T T_3 + \\
 & \tau^2 A^T T_4 + \frac{\tau_1^4}{4} A^T U_1 + \frac{\tau_2^4}{4} A^T U_2 + \frac{\tau_4^4}{4} A^T U_3 - S, \Psi_{11,11} = \tau_1^2 B^T T_1 B + \tau_2^2 B^T T_2 B + \\
 & \eta^2 B^T T_3 B + \tau^2 B^T T_4 B + \frac{\tau_1^4}{4} B^T U_1 B + \frac{\tau_2^4}{4} B^T U_2 B + \frac{\tau_4^4}{4} B^T U_3 B - 2\Lambda_2, \Psi_{11,20} = \\
 & -B^T PM, \Psi_{11,21} = \tau_1^2 B^T T_1 + \tau_2^2 B^T T_2 + \eta^2 B^T T_3 + \tau^2 B^T T_4 + \frac{\tau_1^4}{4} B^T U_1 + \frac{\tau_2^4}{4} B^T U_2 + \\
 & \frac{\tau_4^4}{4} B^T U_3, \Psi_{21,21} = \tau_1^2 T_1 + \tau_2^2 T_2 + \eta^2 T_3 + \tau^2 T_4 + \frac{\tau_1^4}{4} U_1 + \frac{\tau_2^4}{4} U_2 + \frac{\tau_4^4}{4} U_3 - (R - \gamma I), \\
 & \Upsilon_l^T = [\Gamma_{1l}^T, \Gamma_{2l}^T, \Gamma_{3l}^T, \Gamma_{4l}^T, \Gamma_{5l}^T, \Gamma_{6l}^T], (l = 1, 2, 3, 4), \\
 & \Gamma_{11}^T = \Gamma_{12}^T = \tau_1(e_1 - e_{15}), \Gamma_{13}^T = \Gamma_{14}^T = \mathbf{0}, \\
 & \Gamma_{21}^T = \Gamma_{22}^T = \mathbf{0}, \Gamma_{23}^T = \Gamma_{24}^T = \tau_1(e_1 - e_{14}), \\
 & \Gamma_{31}^T = \Gamma_{33}^T = \tau_2(e_1 - e_{17}), \Gamma_{32}^T = \Gamma_{34}^T = \mathbf{0}, \\
 & \Gamma_{41}^T = \Gamma_{43}^T = \mathbf{0}, \Gamma_{42}^T = \Gamma_{44}^T = \tau_2(e_1 - e_{16}), \\
 & \Gamma_{51}^T = \tau(e_1 - e_{19}), \Gamma_{52}^T = \tau_1(e_1 - e_{19}), \Gamma_{53}^T = \tau_2(e_1 - e_{19}), \Gamma_{54}^T = \mathbf{0}, \\
 & \Gamma_{61}^T = \mathbf{0}, \Gamma_{62}^T = \tau_2(e_1 - e_{18}), \Gamma_{63}^T = \tau_1(e_1 - e_{18}), \Gamma_{64}^T = \tau(e_1 - e_{19}), \\
 & e_i = [\mathbf{0}_{n \times (i-1)n}, \mathbf{I}_{n \times n}, \mathbf{0}_{n \times (21-i)n}], (i = 1, 2, \dots, n).
 \end{aligned}$$

**Proof.** We construct the Lyapunov-Krasovskii functional as follows:

$$V(t) = \sum_{l=1}^7 V_l(t) \quad (13)$$

where

$$\begin{aligned}
 V_1(t) &= [x(t) - M \int_{t-\eta}^t x(s) ds]^T P [x(t) - M \int_{t-\eta}^t x(s) ds], \\
 V_2(t) &= 2 \sum_{i=1}^n w_i \int_0^{x_i(t)} f_i(s) ds,
 \end{aligned}$$

$$\begin{aligned}
 V_3(t) &= \int_{t-\tau_1}^t x^T(s)Q_1x(s)ds + \int_{t-\tau_2}^t x^T(s)Q_2x(s)ds + \\
 &\quad \int_{t-\tau}^t x^T(s)Q_3x(s)ds + \int_{t-\eta}^t x^T(s)Q_4x(s)ds, \\
 V_4(t) &= \int_{t-\tau_1(t)}^t x^T(s)R_1x(s)ds + \int_{t-\tau_2(t)}^t x^T(s)R_2x(s)ds + \\
 &\quad \int_{t-\tau(t)}^t x^T(s)R_3x(s)ds + \int_{t-\eta(t)}^t x^T(s)R_4x(s)ds, \\
 V_5(t) &= \tau_1 \int_{-\tau_1}^0 \int_{t+\theta}^t \dot{x}^T(s)T_1\dot{x}(s)dsd\theta + \tau_2 \int_{-\tau_2}^0 \int_{t+\theta}^t \dot{x}^T(s)T_2\dot{x}(s)dsd\theta + \\
 &\quad \eta \int_{-\eta}^0 \int_{t+\theta}^t \dot{x}^T(s)T_3\dot{x}(s)dsd\theta + \tau \int_{-\tau}^0 \int_{t+\theta}^t \dot{x}^T(s)T_4\dot{x}(s)dsd\theta, \\
 V_6(t) &= \frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{\theta}^0 \int_{t+\lambda}^t \dot{x}^T(s)U_1\dot{x}(s)dsd\lambda d\theta + \\
 &\quad \frac{\tau_2^2}{2} \int_{-\tau_2}^0 \int_{\theta}^0 \int_{t+\lambda}^t \dot{x}^T(s)U_2\dot{x}(s)dsd\lambda d\theta + \\
 &\quad \frac{\tau^2}{2} \int_{-\tau}^0 \int_{\theta}^0 \int_{t+\lambda}^t \dot{x}^T(s)U_3\dot{x}(s)dsd\lambda d\theta, \\
 V_7(t) &= \eta \int_{-\eta}^0 \int_{t+\theta}^t x^T(s)Zx(s)dsd\theta.
 \end{aligned}$$

Let

$$\begin{aligned}
 \zeta^T(t) &= [x^T(t), x^T(t-\eta(t)), x^T(t-\eta), x^T(t-\tau_1(t)), x^T(t-\tau_1), x^T(t-\tau_2(t)), \\
 &\quad x^T(t-\tau_2), x^T(t-\tau(t)), x^T(t-\tau), f^T(x(t)), g^T(x(t-\tau(t))), \\
 &\quad \frac{1}{\eta(t)} \int_{t-\eta(t)}^t x^T(s)ds, \frac{1}{\eta-\eta(t)} \int_{t-\eta}^{t-\eta(t)} x^T(s)ds, \frac{1}{\tau_1(t)} \int_{t-\tau_1(t)}^t x^T(s)ds, \\
 &\quad \frac{1}{\tau_1-\tau_1(t)} \int_{t-\tau_1}^{t-\tau_1(t)} x^T(s)ds, \frac{1}{\tau_2(t)} \int_{t-\tau_2(t)}^t x^T(s)ds, \frac{1}{\tau_2-\tau_2(t)} \int_{t-\tau_2}^{t-\tau_2(t)} x^T(s)ds, \\
 &\quad \frac{1}{\tau(t)} \int_{t-\tau(t)}^t x^T(s)ds, \frac{1}{\tau-\tau(t)} \int_{t-\tau}^{t-\tau(t)} x^T(s)ds, \int_{t-\eta}^t x^T(s)ds, u^T(t)]
 \end{aligned}$$

Calculating the upper right Dini derivative of  $V(t)$  along the trajectory of system (4), we have

$$\begin{aligned}
 D^+V_1(t) &= 2[x^T(t) - \int_{t-\eta}^t x^T(s)ds \times M]P[-Dx(t-\eta(t)) + \\
 &\quad Af(x(t)) + Bg(x(t-\tau(t))) + u(t) - Mx(t) + Mx(t-\eta)],
 \end{aligned} \tag{14}$$

$$D^+V_2(t) = 2[-Dx(t - \eta(t)) + Af(x(t)) + Bg(x(t - \tau(t))) + u(t)]^T Wf(x(t)), \quad (15)$$

$$D^+V_3(t) = x^T(t)(Q_1 + Q_2 + Q_3 + Q_4)x(t) - x^T(t - \tau_1)Q_1x(t - \tau_1) - x^T(t - \tau_2)Q_2x(t - \tau_2) - x^T(t - \tau)Q_3x(t - \tau) - x^T(t - \eta)Q_4x(t - \eta) \quad (16)$$

$$D^+V_4(t) \leq x^T(t)(R_1 + R_2 + R_3 + R_4)x(t) - x^T(t - \tau_1(t))R_1x(t - \tau_1(t))(1 - \mu_1) - x^T(t - \tau_2(t))R_2x(t - \tau_2(t))(1 - \mu_2) - x^T(t - \tau(t))R_3x(t - \tau(t))(1 - \mu) - x^T(t - \eta(t))R_4x(t - \eta(t))(1 - \mu) \quad (17)$$

$$D^+V_5(t) = \tau_1^2 \dot{x}^T(t)T_1\dot{x}(t) - \tau_1 \int_{t-\tau_1}^t \dot{x}^T(s)T_1\dot{x}(s)ds + \tau_2^2 \dot{x}^T(t)T_2\dot{x}(t) - \tau_2 \int_{t-\tau_2}^t \dot{x}^T(s)T_2\dot{x}(s)ds + \eta^2 \dot{x}^T(t)T_3\dot{x}(t) - \eta \int_{t-\eta}^t \dot{x}^T(s)T_3\dot{x}(s)ds + \tau^2 \dot{x}^T(t)T_4\dot{x}(t) - \tau \int_{t-\tau}^t \dot{x}^T(s)T_4\dot{x}(s)ds \quad (18)$$

$$D^+V_6(t) = \frac{\tau_1^4}{4} \dot{x}^T(t)U_1\dot{x}(t) - \frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{t+\theta}^t \dot{x}^T(s)U_1\dot{x}(s)dsd\theta + \frac{\tau_2^4}{4} \dot{x}^T(t)U_2\dot{x}(t) - \frac{\tau_2^2}{2} \int_{-\tau_2}^0 \int_{t+\theta}^t \dot{x}^T(s)U_2\dot{x}(s)dsd\theta + \frac{\tau^4}{4} \dot{x}^T(t)U_3\dot{x}(t) - \frac{\tau^2}{2} \int_{-\tau}^0 \int_{t+\theta}^t \dot{x}^T(s)U_3\dot{x}(s)dsd\theta \quad (19)$$

$$D^+V_7(t) \leq \eta^2 x^T(t)Zx(t) - \int_{t-\eta}^t x^T(s)dsZ \int_{t-\eta}^t x(s)ds \quad (20)$$



For any matrix  $T_1$  with  $\begin{bmatrix} T_1 & G_1 \\ \star & T_1 \end{bmatrix} \geq 0$ , by using Lemma 1 and Lemma 3, we can get:

$$\begin{aligned}
 & -\tau_1 \int_{t-\tau_1}^t \dot{x}^T(s) T_1 \dot{x}(s) ds \\
 & = -\tau_1 \left[ \int_{t-\tau_1}^{t-\tau_1(t)} \dot{x}^T(s) T_1 \dot{x}(s) ds + \int_{t-\tau_1(t)}^t \dot{x}^T(s) T_1 \dot{x}(s) ds \right] \\
 & \leq -\frac{\tau_1}{\tau_1 - \tau_1(t)} [\alpha_1^T(t) T_1 \alpha_1(t) + 3\alpha_2^T(t) T_1 \alpha_2(t)] \\
 & \quad - \frac{\tau_1}{\tau_1(t)} [\alpha_3^T(t) T_1 \alpha_3(t) + 3\alpha_4^T(t) T_1 \alpha_4(t)] \\
 & \leq -\alpha_1^T(t) T_1 \alpha_1(t) - 3\alpha_2^T(t) T_1 \alpha_2(t) - \alpha_3^T(t) T_1 \alpha_3(t) - 3\alpha_4^T(t) T_1 \alpha_4(t) \\
 & \quad - 2\alpha_1^T(t) G_1 \alpha_3(t) - 6\alpha_2^T(t) G_1 \alpha_4(t)
 \end{aligned} \tag{21}$$

where

$$\begin{aligned}
 \alpha_1(t) &= x(t - \tau_1(t)) - x(t - \tau_1); \\
 \alpha_2(t) &= x(t - \tau_1(t)) + x(t - \tau_1) - \frac{2}{\tau_1 - \tau_1(t)} \int_{t-\tau_1}^{t-\tau_1(t)} x(s) ds; \\
 \alpha_3(t) &= x(t) - x(t - \tau_1(t)); \\
 \alpha_4(t) &= x(t) + x(t - \tau_1(t)) - \frac{2}{\tau_1(t)} \int_{t-\tau_1(t)}^t x(s) ds.
 \end{aligned}$$

Similarly, applying Lemma 1 and Lemma 3 in fourth, sixth and eighth term of Equ.(18), we can obtain the following:

$$\begin{aligned}
 & -\tau_2 \int_{t-\tau_2}^t \dot{x}^T(s) T_2 \dot{x}(s) ds \\
 & \leq -\alpha_5^T(t) T_2 \alpha_5(t) - 3\alpha_6^T(t) T_2 \alpha_6(t) - \alpha_7^T(t) T_2 \alpha_7(t) \\
 & \quad - 3\alpha_8^T(t) T_2 \alpha_8(t) - 2\alpha_5^T(t) G_2 \alpha_7(t) - 6\alpha_6^T(t) G_2 \alpha_8(t)
 \end{aligned} \tag{22}$$

$$\begin{aligned}
 & -\eta \int_{t-\eta}^t \dot{x}^T(s) T_3 \dot{x}(s) ds \\
 & \leq -\alpha_9^T(t) T_3 \alpha_9(t) - 3\alpha_{10}^T(t) T_3 \alpha_{10}(t) - \alpha_{11}^T(t) T_3 \alpha_{11}(t) \\
 & \quad - 3\alpha_{12}^T(t) T_3 \alpha_{12}(t) - 2\alpha_9^T(t) G_3 \alpha_{11}(t) - 6\alpha_{10}^T(t) G_3 \alpha_{12}(t)
 \end{aligned} \tag{23}$$

$$\begin{aligned}
 & -\tau \int_{t-\tau}^t \dot{x}^T(s) T_4 \dot{x}(s) ds \\
 & \leq -\alpha_{13}^T(t) T_4 \alpha_{13}(t) - 3\alpha_{14}^T(t) T_4 \alpha_{14}(t) - \alpha_{15}^T(t) T_4 \alpha_{15}(t) \\
 & \quad - 3\alpha_{16}^T(t) T_4 \alpha_{16}(t) - 2\alpha_{13}^T(t) G_4 \alpha_{15}(t) - 6\alpha_{14}^T(t) G_4 \alpha_{16}(t)
 \end{aligned} \tag{24}$$

where

$$\begin{aligned}
 \alpha_5(t) &= x(t - \tau_2(t)) - x(t - \tau_2); \\
 \alpha_6(t) &= x(t - \tau_2(t)) + x(t - \tau_2) - \frac{2}{\tau_2 - \tau_2(t)} \int_{t-\tau_2}^{t-\tau_2(t)} x(s) ds; \\
 \alpha_7(t) &= x(t) - x(t - \tau_2(t)); \\
 \alpha_8(t) &= x(t) + x(t - \tau_2(t)) - \frac{2}{\tau_2(t)} \int_{t-\tau_2(t)}^t x(s) ds; \\
 \alpha_9(t) &= x(t - \eta(t)) - x(t - \eta); \\
 \alpha_{10}(t) &= x(t - \eta(t)) + x(t - \eta) - \frac{2}{\eta - \eta(t)} \int_{t-\eta}^{t-\eta(t)} x(s) ds; \\
 \alpha_{11}(t) &= x(t) - x(t - \eta(t)); \\
 \alpha_{12}(t) &= x(t) + x(t - \eta(t)) - \frac{2}{\eta(t)} \int_{t-\eta(t)}^t x(s) ds; \\
 \alpha_{13}(t) &= x(t - \tau(t)) - x(t - \tau); \\
 \alpha_{14}(t) &= x(t - \tau(t)) + x(t - \tau) - \frac{2}{\tau - \tau(t)} \int_{t-\tau}^{t-\tau(t)} x(s) ds; \\
 \alpha_{15}(t) &= x(t) - x(t - \tau(t)); \\
 \alpha_{16}(t) &= x(t) + x(t - \tau(t)) - \frac{2}{\tau(t)} \int_{t-\tau(t)}^t x(s) ds;
 \end{aligned}$$

Now, the second term of Equ.(19) can be written as

$$\begin{aligned}
 & -\frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{t+\theta}^t \dot{x}^T(s) U_1 \dot{x}(s) ds d\theta \\
 & = -\frac{\tau_1^2}{2} \int_{-\tau_1}^{-\tau_1(t)} \int_{t+\theta}^t \dot{x}^T(s) U_1 \dot{x}(s) ds d\theta - \frac{\tau_1^2}{2} \int_{-\tau_1(t)}^0 \int_{t+\theta}^t \dot{x}^T(s) U_1 \dot{x}(s) ds d\theta
 \end{aligned} \tag{25}$$

Further, using Lemma 2, the above equation becomes

$$\begin{aligned}
 & +\frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{t+\theta}^t \dot{x}^T(s) U_1 \dot{x}(s) ds d\theta \\
 & \leq -\frac{\tau_1^2}{\tau_1^2 - \tau_1^2(t)} \left[ \int_{-\tau_1}^{-\tau_1(t)} \int_{t+\theta}^t \dot{x}(s) ds d\theta \right]^T U_1 \left[ \int_{-\tau_1}^{-\tau_1(t)} \int_{t+\theta}^t \dot{x}(s) ds d\theta \right] \\
 & \quad -\frac{\tau_1^2}{\tau_1^2(t)} \left[ \int_{-\tau_1(t)}^0 \int_{t+\theta}^t \dot{x}(s) ds d\theta \right]^T U_1 \left[ \int_{-\tau_1(t)}^0 \int_{t+\theta}^t \dot{x}(s) ds d\theta \right]
 \end{aligned} \tag{26}$$

For any matrix  $V_1$  with  $\begin{bmatrix} U_1 & V_1 \\ \star & U_1 \end{bmatrix} \geq 0$ , from Lemma 3, the above inequality

becomes

$$\begin{aligned}
 & -\frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{t+\theta}^t \dot{x}^T(s) U_1 \dot{x}(s) ds d\theta \\
 & \leq - \left[ \int_{-\tau_1}^{-\tau_1(t)} \int_{t+\theta}^t \dot{x}(s) ds d\theta \right]^T U_1 \left[ \int_{-\tau_1}^{-\tau_1(t)} \int_{t+\theta}^t \dot{x}(s) ds d\theta \right] \\
 & \quad - \left[ \int_{-\tau_1(t)}^0 \int_{t+\theta}^t \dot{x}(s) ds d\theta \right]^T 2V_1 \left[ \int_{-\tau_1}^{-\tau_1(t)} \int_{t+\theta}^t \dot{x}(s) ds d\theta \right] \\
 & \quad - \left[ \int_{-\tau_1(t)}^0 \int_{t+\theta}^t \dot{x}(s) ds d\theta \right]^T U_1 \left[ \int_{-\tau_1(t)}^0 \int_{t+\theta}^t \dot{x}(s) ds d\theta \right] \\
 & = -\Delta_1^T U_1 \Delta_1 - 2\Delta_1^T V_1 \Delta_2 - \Delta_2^T U_1 \Delta_2 \\
 & = \zeta^T(t) [-\Gamma_1^T(t) U_1 \Gamma_1(t) - 2\Gamma_1^T(t) V_1 \Gamma_2(t) - \Gamma_2^T(t) U_1 \Gamma_2(t)] \zeta(t)
 \end{aligned} \tag{27}$$

where  $\Delta_1 = (\tau_1 - \tau_1(t))x(t) - \int_{t-\tau_1}^{t-\tau_1(t)} x(s) ds$ ,  $\Delta_2 = \tau_1(t)x(t) - \int_{t-\tau_1(t)}^t x(s) ds$ ,  $\Gamma_1(t) = (\tau_1 - \tau_1(t))(e_1 - e_{15})$ ,  $\Gamma_2(t) = \tau_1(t)(e_1 - e_{14})$ .

Similarly, applying Lemma 3 in fourth and sixth term of Equ.(19), we can obtain the following:

$$\begin{aligned}
 & -\frac{\tau_2^2}{2} \int_{-\tau_2}^0 \int_{t+\theta}^t \dot{x}^T(s) U_2 \dot{x}(s) ds d\theta \\
 & \leq -\Delta_3^T U_2 \Delta_3 - 2\Delta_3^T V_2 \Delta_4 - \Delta_4^T U_2 \Delta_4 \\
 & = \zeta^T(t) [-\Gamma_3^T(t) U_2 \Gamma_3(t) - 2\Gamma_3^T(t) V_2 \Gamma_4(t) - \Gamma_4^T(t) U_2 \Gamma_4(t)] \zeta(t)
 \end{aligned} \tag{28}$$

$$\begin{aligned}
 & -\frac{\tau^2}{2} \int_{-\tau}^0 \int_{t+\theta}^t \dot{x}^T(s) U_3 \dot{x}(s) ds d\theta \\
 & \leq -\Delta_5^T U_3 \Delta_5 - 2\Delta_5^T V_3 \Delta_6 - \Delta_6^T U_3 \Delta_6 \\
 & = \zeta^T(t) [-\Gamma_5^T(t) U_3 \Gamma_5(t) - 2\Gamma_5^T(t) V_3 \Gamma_6(t) - \Gamma_6^T(t) U_3 \Gamma_6(t)] \zeta(t)
 \end{aligned} \tag{29}$$

where  $\Delta_3 = (\tau_2 - \tau_2(t))x(t) - \int_{t-\tau_2}^{t-\tau_2(t)} x(s) ds$ ,  $\Delta_4 = \tau_2(t)x(t) - \int_{t-\tau_2(t)}^t x(s) ds$ ,  $\Delta_5 = (\tau - \tau(t))x(t) - \int_{t-\tau}^{t-\tau(t)} x(s) ds$ ,  $\Delta_6 = \tau(t)x(t) - \int_{t-\tau(t)}^t x(s) ds$ ,  $\Gamma_3(t) = (\tau_2 - \tau_2(t))(e_1 - e_{17})$ ,  $\Gamma_4(t) = \tau_2(t)(e_1 - e_{16})$ ,  $\Gamma_5(t) = (\tau - \tau(t))(e_1 - e_{19})$ ,  $\Gamma_6(t) = \tau(t)(e_1 - e_{18})$ .

According to Assumption 1, we can get

$$\sum_{i=1}^n 2\lambda_{1i} [l_i^- x_i(t) - f_i(x_i(t))] [f_i(x_i(t)) - l_i^+ x_i(t)] \geq 0 \tag{30}$$

$$\sum_{i=1}^n 2\lambda_{2i} [h_i^- x_i(t - \tau_{i1}(t) - \tau_{i2}(t)) - g_i(x_i(t - \tau_{i1}(t) - \tau_{i2}(t)))] \times$$

$$[g_i(x_i(t - \tau_{i1}(t) - \tau_{i2}(t))) - h_i^+ x_i(t - \tau_{i1}(t) - \tau_{i2}(t))] \geq 0 \quad (31)$$

Then Equations (30) - (31) can be written in the following vector form:

$$-2x^T(t)L^-\Lambda_1L^+x(t) - 2f^T(x(t))\Lambda_1f(x(t))$$

$$+ 2x^T(t)(L^- + L^+)\Lambda_1f(x(t)) \geq 0 \quad (32)$$

$$-2x^T(t - \tau(t))H^-\Lambda_2H^+x(t - \tau(t)) - 2g^T(x(t - \tau(t))) \times$$

$$\Lambda_2g(x(t - \tau(t))) + 2x^T(t - \tau(t))(H^- + H^+)\Lambda_2g(x(t - \tau(t))) \geq 0 \quad (33)$$

From (14) - (29) and adding (32) and (33), we have

$$D^+V(t) - f^T(x(t))Qf(x(t)) - 2f^T(x(t))Su(t) - u^T(t)(R - \gamma I)u(t)$$

$$\leq \zeta^T(t)\Phi(\tau_1(t), \tau_2(t))\zeta(t) \quad (34)$$

where

$$\Phi(\tau_1(t), \tau_2(t)) = \Xi - \Gamma_1^T(t)U_1\Gamma_1(t) - 2\Gamma_1^T(t)V_1\Gamma_2(t) - \Gamma_2^T(t)U_1\Gamma_2(t) - \Gamma_3^T(t)U_2\Gamma_3(t) -$$

$$2\Gamma_3^T(t)V_2\Gamma_4(t) - \Gamma_4^T(t)U_2\Gamma_4(t) - \Gamma_5^T(t)U_3\Gamma_5(t) - 2\Gamma_5^T(t)V_3\Gamma_6(t) - \Gamma_6^T(t)U_3\Gamma_6(t).$$

Letting  $\tau_1(t) = 0$ ,  $\tau_1(t) = \tau_1$  and  $\tau_2(t) = 0$ ,  $\tau_2(t) = \tau_2$  respectively, we can get

$$\begin{cases} \Phi_1 = \Phi(0, 0), \\ \Phi_2 = \Phi(0, \tau_2), \\ \Phi_3 = \Phi(\tau_1, 0), \\ \Phi_4 = \Phi(\tau_1, \tau_2) \end{cases} \quad (35)$$

According to the LMIs (10), we have

$$D^+V(t) - f^T(x(t))Qf(x(t)) - 2f^T(x(t))Su(t) - u^T(t)(R - \gamma I)u(t) \leq 0 \quad (36)$$

Under the zero initial conditon,  $V(0) = 0$ . For any  $T > 0$ ,  $V(T) \geq 0$ . Integrating Equ.(36) with respect to  $t$  over the time period from 0 to  $T$ , we have

$$\int_0^T [-f^T(x(t))Qf(x(t)) - 2f^T(x(t))Su(t) - u^T(t)(R - \gamma I)u(t)]dt \leq -V(T) \leq 0 \quad (37)$$

So Equ. (5) holds, which implies MNNs (4) is strictly  $(Q, S, R) - \gamma -$  dissipative. This completes the proof.

**Remark 5.** In Theorem 1, not only free-weighting matrices technique but also the reciprocally convex combination technique, Wirtinger-based integral inequality are employed. Also, double integral form in  $V_5(t)$  and novel triple integral form in  $V_6(t)$  are introduced by considering both time-varying leakage delay and two additive time-varying delays, which has not been addressed yet in the previous literature [37, 39, 47]. Construction this form of double and triple integral terms in the LKF is a recent tool for getting less conservative results.

**Remark 6.** According to Definition 3 and Definition 4, we can obtain the passivity conditions for the system (4) by substituting  $Q = 0$ ,  $S = I$  and  $R = 2\gamma I$  in Theorem 1. In this case, we can get Theorem 2 in the same way as Theorem 1.

**Theorem 2.** Under Assumption 1, the MNNs (4) is passive in the sense of Definition 4, if there exist symmetric matrices  $P > 0$ ,  $Q_i > 0$ ,  $R_i > 0$ ,  $T_i > 0$ , ( $i = 1, 2, 3, 4$ ),  $U_j > 0$  ( $j = 1, 2, 3$ ),  $Z > 0$ , diagonal matrices  $M = \text{diag}\{m_1, m_2, \dots, m_n\} > 0$ ,  $W = \text{diag}\{w_1, w_2, \dots, w_n\} > 0$ ,  $\Lambda_1 = \text{diag}\{\lambda_{11}, \lambda_{12}, \dots, \lambda_{1n}\} > 0$ ,  $\Lambda_2 = \text{diag}\{\lambda_{21}, \lambda_{22}, \dots, \lambda_{2n}\} > 0$ , and matrices  $G_i$  ( $i = 1, 2, 3, 4$ ),  $V_j$  ( $j = 1, 2, 3$ ) and scalar  $\gamma > 0$ , such that the following LMIs

hold:

$$\tilde{\Phi}_l = \tilde{\Xi} - \Upsilon_l^T \begin{bmatrix} U_1 & V_1 & 0 & 0 & 0 & 0 \\ \star & U_1 & 0 & 0 & 0 & 0 \\ \star & \star & U_2 & V_2 & 0 & 0 \\ \star & \star & \star & U_2 & 0 & 0 \\ \star & \star & \star & \star & U_3 & V_3 \\ \star & \star & \star & \star & \star & U_3 \end{bmatrix} \Upsilon_l \leq 0, \quad (38)$$

$l = (1, 2, 3, 4).$

$$\begin{bmatrix} T_i & G_i \\ \star & T_i \end{bmatrix} \geq 0, (i = 1, 2, 3, 4). \quad (39)$$

$$\begin{bmatrix} U_j & V_j \\ \star & U_j \end{bmatrix} \geq 0, (j = 1, 2, 3). \quad (40)$$

where  $\tilde{\Xi} = (\tilde{\Psi}_{p,q})_{21 \times 21}$ ,  $\tilde{\Psi}_{p,q} = \Psi_{p,q}$ ,  $\forall p, q = 1, 2, \dots, 21$ , except

$$\begin{aligned} \tilde{\Psi}_{10,10} &= AW + WA^T + \tau_1^2 A^T T_1 A + \tau_2^2 A^T T_2 A + \eta^2 A^T T_3 A + \tau^2 A^T T_4 A + \frac{\tau_1^4}{4} A^T U_1 A + \\ &\frac{\tau_2^4}{4} A^T U_2 A + \frac{\tau^4}{4} A^T U_3 A - 2\Lambda_1, \quad \Psi_{10,21} = W + \tau_1^2 A^T T_1 + \tau_2^2 A^T T_2 + \eta^2 A^T T_3 + \\ &\tau^2 A^T T_4 + \frac{\tau_1^4}{4} A^T U_1 + \frac{\tau_2^4}{4} A^T U_2 + \frac{\tau^4}{4} A^T U_3 - I, \quad \Psi_{21,21} = \tau_1^2 T_1 + \tau_2^2 T_2 + \eta^2 T_3 + \\ &\tau^2 T_4 + \frac{\tau_1^4}{4} U_1 + \frac{\tau_2^4}{4} U_2 + \frac{\tau^4}{4} U_3 - \gamma I. \end{aligned}$$

The remaining coefficients are same as in Theorem 1.

**Proof.** The proof is same as that of Theorem 1 and hence it is omitted.

**Remark 7.** The less conservative of the proposed delay-dependent dissipativity and passivity criterion over [22, 29] relies on how to handle the terms  $-\tau_1 \int_{t-\tau_1}^t \dot{x}^T(s) T_1 \dot{x}(s) ds$ ,  $-\tau_2 \int_{t-\tau_2}^t \dot{x}^T(s) T_2 \dot{x}(s) ds$ ,  $-\eta \int_{t-\eta}^t \dot{x}^T(s) T_3 \dot{x}(s) ds$  and  $-\tau \int_{t-\tau}^t \dot{x}^T(s) T_4 \dot{x}(s) ds$ . Lemma 1 and Lemma 3 is employed to handle these terms instead of using Jensen Inequality in [22, 29].

**Remark 8.** When  $u(t) = 0$  and  $Q = 0$ , it can be seen from Equ.(36) that  $D^+V(t) \leq 0$ . So it implies the asymptotic stability of the system (4). So the dissipativity and passivity analysis for MMNs is more general and applicable.

#### 4. Illustrative examples

In this section, two numerical examples are given to illustrate the effectiveness of the obtained results.

**Example 1.** Consider the following MNNs with leakage and two additive time-varying delays:

$$\begin{cases} \frac{dx_i(t)}{dt} = -d_i x_i(t - \eta_i(t)) + \sum_{j=1}^2 a_{ij}(t) f_j(x_j(t)) \\ \quad + \sum_{j=1}^2 b_{ij}(t) g_j(x_j(t - \bar{\tau}_j(t))) + u_i(t) \\ y_i(t) = f_i(x_i(t)), i = 1, 2. \end{cases} \quad (41)$$

where

$$\bar{\tau}_j(t) = \tau_{j1}(t) + \tau_{j2}(t), d_1 = 1.10, d_2 = 1.20, u_1(t) = \sin(0.5t), u_2(t) = \cos(2t),$$

$$\begin{aligned} a_{11}(t) &= \begin{cases} 0.60, & \dot{\kappa}_{11}^1 < 0 \\ \text{unchanged}, & \dot{\kappa}_{11}^1 = 0, \\ 0.80, & \dot{\kappa}_{11}^1 > 0 \end{cases}, a_{12}(t) = \begin{cases} 0.10, & \dot{\kappa}_{12}^1 < 0 \\ \text{unchanged}, & \dot{\kappa}_{12}^1 = 0, \\ 0.15, & \dot{\kappa}_{12}^1 > 0 \end{cases}, \\ a_{21}(t) &= \begin{cases} 0.12, & \dot{\kappa}_{21}^1 < 0 \\ \text{unchanged}, & \dot{\kappa}_{21}^1 = 0, \\ 0.16, & \dot{\kappa}_{21}^1 > 0 \end{cases}, a_{22}(t) = \begin{cases} 0.70, & \dot{\kappa}_{22}^1 < 0 \\ \text{unchanged}, & \dot{\kappa}_{22}^1 = 0, \\ 0.80, & \dot{\kappa}_{22}^1 > 0 \end{cases}, \\ b_{11}(t) &= \begin{cases} 0.30, & \dot{\kappa}_{11}^2 < 0 \\ \text{unchanged}, & \dot{\kappa}_{11}^2 = 0, \\ 0.35, & \dot{\kappa}_{11}^2 > 0 \end{cases}, b_{12}(t) = \begin{cases} 0.08, & \dot{\kappa}_{12}^2 < 0 \\ \text{unchanged}, & \dot{\kappa}_{12}^2 = 0, \\ 0.10, & \dot{\kappa}_{12}^2 > 0 \end{cases}, \\ b_{21}(t) &= \begin{cases} 0.07, & \dot{\kappa}_{21}^2 < 0 \\ \text{unchanged}, & \dot{\kappa}_{21}^2 = 0, \\ 0.12, & \dot{\kappa}_{21}^2 > 0 \end{cases}, b_{22}(t) = \begin{cases} 0.40, & \dot{\kappa}_{22}^2 < 0 \\ \text{unchanged}, & \dot{\kappa}_{22}^2 = 0, \\ 0.42, & \dot{\kappa}_{22}^2 > 0 \end{cases}, \end{aligned}$$

$$\kappa_{ij}^1 = f_{ij}(x_j(t)) - x_i(t), \kappa_{ij}^2 = g_{ij}(x_j(t - \bar{\tau}_j(t))) - x_i(t), i = 1 \text{ or } 2, j = 1 \text{ or } 2.$$

Let us consider the nonlinear activation functions as  $f(x(t)) = g(x(t)) = \sin(x(t))$ , it can be verified that the activation functions satisfy Assumption

1 with  $l_i^- = h_i^- = -1$ ,  $l_i^+ = h_i^+ = 1$ , ( $i = 1, 2$ ). We shall assume that the leakage delay to be  $\eta(t) = 0.2 + 0.2\sin(2t)$  and the time-varying delays to be  $\tau_1(t) = 0.1 + 0.1\sin(t)$ ,  $\tau_2(t) = 0.1 + 0.1\cos(t)$ . So  $\eta = 0.4$ ,  $\tau_1 = 0.2$ ,  $\tau_2 = 0.2$ ,  $\mu_1 = 0.1$ ,  $\mu_2 = 0.1$ ,  $\mu = 0.2$ , Then a set of corresponding feasible solution to  
 285 the LMIs in Theorem 1 can be obtained by the Matlab LMI control toolbox as following:

$$\begin{aligned} P &= \begin{bmatrix} 0.0475 & -0.0291 \\ -0.0291 & 0.0287 \end{bmatrix}, Q_1 = \begin{bmatrix} 2.3090 & -1.3697 \\ -1.3697 & 1.4282 \end{bmatrix}, \\ 290 \quad Q_2 &= \begin{bmatrix} 2.2678 & -1.3111 \\ -1.3111 & 1.4202 \end{bmatrix}, R_1 = \begin{bmatrix} 3.2735 & -1.9551 \\ -1.9551 & 2.0149 \end{bmatrix}, \\ T_1 &= \begin{bmatrix} 0.0746 & -0.0445 \\ -0.0445 & 0.0459 \end{bmatrix}, U_1 = \begin{bmatrix} 15.4356 & -9.0026 \\ -9.0026 & 9.6499 \end{bmatrix}, \end{aligned}$$

Due to the page limit, we omit the remaining feasible matrices. Fig.1 shows  
 295 phase trajectories of MNNs (41) under the zero initial condition. Fig.2 shows trajectories of neuron state  $x_1(t)$  and  $x_2(t)$  of MNNs (41). Fig.3 shows 3-D trajectories of neuron state  $x_1(t)$  and  $x_2(t)$  of MNNs (41). It can be seen that neuron state  $x_1(t)$  and  $x_2(t)$  are tending to periodic when the outputs of MNNs (41) controllers are designed to periodic signals. According to Theorem 1 and  
 300 Definition 3, the system (41) is dissipative.

**Example 2.** Consider a two-neuron MNNs (4) with the following parameters:

$$\begin{aligned} D &= \begin{bmatrix} 1.50 & 0 \\ 0 & 1.80 \end{bmatrix}, u(t) = \begin{bmatrix} \cos t \\ \sin(2t) \end{bmatrix}, \\ 305 \quad A &= \begin{bmatrix} 1.8 & 0.10 \\ 0.12 & 2.00 \end{bmatrix}, B = \begin{bmatrix} 0.80 & 0.07 \\ 0.05 & 1.20 \end{bmatrix}, \end{aligned}$$

Let us consider the nonlinear activation functions as  $f(x(t)) = g(x(t)) = 0.01(|x + 1| - |x - 1|)$ . We shall assume that the leakage delay to be  $\eta(t) = 0.1 + 0.1\sin(2t)$  and the time-varying delays to be  $\tau_1(t) = 0.05 + 0.05\sin(t)$ ,







Figure 1: Phase trajectories of MNNs (41).


 Figure 2: Trajectories of neuron state  $x_1(t)$  and  $x_2(t)$  of MNNs (41).

<sup>310</sup>  $\tau_2(t) = 0.05 + 0.05\cos(t)$ . So  $\eta = 0.2$ ,  $\tau_1 = 0.1$ ,  $\tau_2 = 0.1$ ,  $\mu_1 = 0.05$ ,  $\mu_2 = 0.05$ ,  $\mu = 0.1$ , Then a set of corresponding feasible solution to the LMIs in Theorem 2 can be obtained by the Matlab LMI control toolbox as following:

$$P = \begin{bmatrix} 0.0026 & -0.0001 \\ -0.0001 & 0.0016 \end{bmatrix}, Q_1 = \begin{bmatrix} 1.1953 & -0.0439 \\ -0.0439 & 0.7844 \end{bmatrix},$$




Figure 3: 3-D trajectories of neuron state  $x_1(t)$  and  $x_2(t)$  of MNNs (41).

$$Q_2 = \begin{bmatrix} 1.1574 & -0.0423 \\ -0.0423 & 0.7586 \end{bmatrix}, R_1 = \begin{bmatrix} 1.6760 & -0.0614 \\ -0.0614 & 1.1008 \end{bmatrix},$$

$$T_1 = \begin{bmatrix} 0.0396 & -0.0014 \\ -0.0014 & 0.0260 \end{bmatrix}, U_1 = \begin{bmatrix} 29.7285 & -1.0527 \\ -1.0527 & 19.8825 \end{bmatrix},$$

Due to the page limit, we omit the remaining feasible matrices. Fig.4 shows the time respond trajectories of MNNs (4) with 20 random initial conditions and without external input respectively, it is clear that  $x_1(t)$  and  $x_2(t)$  are asymptotic stability. Therefore, we conclude that system (4) is passive.

In order to demonstrate the improvement of our results, we compare our results with those in [26, 48, 49]. Table 1 shows the comparison results for allowable upper bounds of  $\tau$  for different  $\mu$ . From the table, the results calculated based on the criteria given in this paper are less conservative than the ones reported in the existing literatures.

## 5. Conclusion

In this paper, the problem of dissipativity and passivity has been analysed for MNNs in the case of both leakage and two additive time-varying delays. MNNs




Figure 4: Time response trajectories of MNNs (4). ( $u_1(t) = u_2(t) = 0$ )

Table 1: Allowable upper bounds of  $\tau$  for different  $\mu$ .

Conditions	[48]	[49]	[26]	Theorem 2
$\mu = 0$	0.4528	0.7340	Unknown	Large
$\mu = 0.5$	0.3638	0.6834	1.0760	1.2803
$\mu = 0.7$	0.3575	0.6355	1.0704	1.2207

have been converted into the conventional neural networks by applying differential inclusions and set-valued maps. Wirtinger-based integral inequality, reciprocally convex combination technique and free-weighting matrices technique  
335 have been used to obtain less conservative dissipativity and passivity criteria, which is based on proper Lyapunov functional approach. The effectiveness of the corresponding criteria have been shown via two numerical examples. The employed methods can be extended to other problems of MNNs with leakage  
340 MNNs is a new and challenging topic, looking for more complex and practical MNNs model and more advanced mathematical method is our further work. In future, we shall consider the dissipativity and passivity for stochastic memristive

complex-values neural networks with two delay components based on dynamic delay interval method.

## References

- [1] L. Chua, Memristor-the missing circuit element, *IEEE Transactions on circuit theory* 18 (5) (1971) 507–519.
- [2] D. B. Strukov, G. S. Snider, D. R. Stewart, R. S. Williams, The missing memristor found, *nature* 453 (7191) (2008) 80–83.
- [3] L. Chua, Resistance switching memories are memristors, *Applied Physics A* 102 (4) (2011) 765–783.
- [4] D. Liu, S. Zhu, W. Chang, Input-to-state stability of memristor-based complex-valued neural networks with time delays, *Neurocomputing* 221 (2017) 159–167.
- [5] L. Wang, H. Li, S. Duan, T. Huang, H. Wang, Pavlov associative memory in a memristive neural network and its circuit implementation, *Neurocomputing* 171 (2016) 23–29.
- [6] S. Hu, Y. Liu, Z. Liu, T. Chen, Q. Yu, L. Deng, Y. Yin, S. Hosaka, Synaptic long-term potentiation realized in pavlov’s dog model based on a niox-based memristor, *Journal of Applied Physics* 116 (21) (2014) 214502.
- [7] H. Kim, M. P. Sah, C. Yang, T. Roska, L. O. Chua, Neural synaptic weighting with a pulse-based memristor circuit, *IEEE Transactions on Circuits and Systems I: Regular Papers* 59 (1) (2012) 148–158.
- [8] X. Hu, S. Duan, G. Chen, L. Chen, Modeling affections with memristor-based associative memory neural networks, *Neurocomputing* 223 (2017) 129–137.
- [9] P. Zhang, C. Li, T. Huang, L. Chen, Y. Chen, Forgetting memristor based neuromorphic system for pattern training and recognition, *Neurocomputing* 222 (2016) 47–53.

- [10] M. Itoh, L. O. Chua, Memristor cellular automata and memristor discrete-time cellular neural networks, *International Journal of Bifurcation and Chaos* 19 (11) (2009) 3605–3656.
- [11] R. Sakthivel, R. Anbuviya, K. Mathiyalagan, Y.-K. Ma, P. Prakash, Reliable anti-synchronization conditions for bam memristive neural networks with different memductance functions, *Applied Mathematics and Computation* 275 (2016) 213–228.
- [12] M. S. Ali, Stability of markovian jumping recurrent neural networks with discrete and distributed time-varying delays, *Neurocomputing* 149 (2015) 1280–1285.
- [13] A. Chandrasekar, R. Rakkiyappan, Impulsive controller design for exponential synchronization of delayed stochastic memristor-based recurrent neural networks, *Neurocomputing* 173 (2016) 1348–1355.
- [14] G. Jie, P. Zhu, A. Alsaedi, F. E. Alsaadi, T. Hayat, A new switching control for finite-time synchronization of memristor-based recurrent neural networks, *Neural Networks* 86 (2017) 1–9.
- [15] Y. Liu, B. Z. Guo, J. H. Park, S. M. Lee, Nonfragile exponential synchronization of delayed complex dynamical networks with memory sampled-data control, *IEEE Transactions on Neural Networks and Learning Systems* PP (99) (2016) 1–11.
- [16] H. Zhang, Q. Shan, Z. Wang, Stability analysis of neural networks with two delay components based on dynamic delay interval method, *IEEE Transactions on Neural networks and learning systems* 28 (2) (2017) 259–267.
- [17] A. Chandrasekar, R. Rakkiyappan, X. Li, Effects of bounded and unbounded leakage time-varying delays in memristor-based recurrent neural networks with different memductance functions, *Neurocomputing* 202 (2016) 67–83.

- [18] P. Jiang, Z. Zeng, J. Chen, On the periodic dynamics of memristor-based neural networks with leakage and time-varying delays, *Neurocomputing* 219 (2017) 163–173.
- 400 [19] C. Xu, P. Li, Y. Pang, Exponential stability of almost periodic solutions for memristor-based neural networks with distributed leakage delays, *Neural Computation* 28 (12) (2016) 2726–2756.
- [20] Y. Liu, S. M. Lee, O. M. Kwon, H. P. Ju, New approach to stability criteria for generalized neural networks with interval time-varying delays, *Neuro-*  
405 *computing* 149 (2015) 1544–1551.
- [21] S. Wen, Z. Zeng, T. Huang, Y. Chen, Passivity analysis of memristor-based recurrent neural networks with time-varying delays, *Journal of the Franklin Institute* 350 (8) (2013) 2354–2370.
- [22] R. Rakkiyappan, K. Sivarajani, G. Velmurugan, Passivity and passifica-  
410 *tion of memristor-based complex-valued recurrent neural networks with interval time-varying delays, Neurocomputing* 144 (1) (2014) 391–407.
- [23] J. Liu, R. Xu, Passivity analysis and state estimation for a class of memristor-based neural networks with multiple proportional delays, *Advances in Difference Equations* 2017 (1) (2017) 34.
- 415 [24] M. Syed Ali, R. Saravanakumar, J. Cao, New passivity criteria for memristor-based neutral-type stochastic bam neural networks with mixed time-varying delays, *Neurocomputing* 171 (C) (2016) 1533–1547.
- [25] Y. Huang, S. Ren, Passivity and passivity-based synchronization of switched coupled reaction-diffusion neural networks with state and spatial  
420 *diffusion couplings, Neural Processing Letters* (2017) 1–17.
- [26] J. Xiao, S. Zhong, Y. Li, Relaxed dissipativity criteria for memristive neural networks with leakage and time-varying delays, *Neurocomputing* 171 (C) (2016) 708–718.

- [27] H. Wei, R. Li, C. Chen, Z. Tu, Extended dissipative analysis for memristive neural networks with two additive time-varying delay components , Neurocomputing 216 (2016) 429–438.
- [28] M. S. Ali, S. Arik, M. E. Rani, Passivity analysis of stochastic neural networks with leakage delay and markovian jumping parameters , Neurocomputing 218 (2016) 139–145.
- [29] X. Li, R. Rakkiyappan, G. Velmurugan, Dissipativity analysis of memristor-based complex-valued neural networks with time-varying delays, Information Sciences 294 (2015) 645 – 665.
- [30] S. Ding, Z. Wang, H. Zhang, Dissipativity analysis for stochastic memristive neural networks with time-varying delays: A discrete-time case, IEEE Transactions on Neural Networks and Learning Systems PP (99) (2016) 1–13.
- [31] A. Wu, S. Wen, Z. Zeng, Synchronization control of a class of memristor-based recurrent neural networks, Information Sciences 183 (1) (2012) 106–116.
- [32] Y. Song, S. Wen, Synchronization control of stochastic memristor-based neural networks with mixed delays, Neurocomputing 156 (2015) 121–128.
- [33] N. Li, J. Cao, Lag synchronization of memristor-based coupled neural networks via-measure, IEEE transactions on neural networks and learning systems 27 (3) (2016) 686–697.
- [34] H. Li, H. Jiang, C. Hu, Existence and global exponential stability of periodic solution of memristor-based bam neural networks with time-varying delays, Neural Networks 75 (2016) 97–109.
- [35] H. Bao, J. H. Park, J. Cao, Exponential synchronization of coupled stochastic memristor-based neural networks with time-varying probabilistic delay coupling and impulsive delay, IEEE transactions on neural networks and learning systems 27 (1) (2016) 190–201.

- [36] S. P. Adhikari, M. P. Sah, H. Kim, L. O. Chua, Three fingerprints of memristor, *IEEE Transactions on Circuits and Systems I: Regular Papers* 60 (11) (2013) 3008–3021.
- 455 [37] G. Nagamani, S. Ramasamy, Dissipativity and passivity analysis for uncertain discrete-time stochastic markovian jump neural networks with additive time-varying delays, *Neurocomputing* 174 (2016) 795–805.
- [38] R. Rakkiyappan, A. Chandrasekar, J. Cao, Passivity and passification of memristor-based recurrent neural networks with additive time-varying delays, *IEEE transactions on neural networks and learning systems* 26 (9) 460 (2015) 2043–2057.
- [39] J. Xiao, S. Zhong, Y. Li, Relaxed dissipativity criteria for memristive neural networks with leakage and time-varying delays, *Neurocomputing* 171 (C) (2015) 708–718.
- 465 [40] J. Liu, R. Xu, Delay-dependent passivity and stability analysis for a class of memristor-based neural networks with time delay in the leakage term, *Neural Processing Letters* (2017) 1–19.
- [41] J. P. Aubin, A. Cellina, *Differential Inclusions*, Springer Berlin Heidelberg, 1984.
- 470 [42] J. P. Aubin, *Set-Valued Analysis*, Birkhäuser Boston, 1999.
- [43] S. Wen, T. Huang, Z. Zeng, Y. Chen, P. Li, Circuit design and exponential stabilization of memristive neural networks, *Neural Networks* 63 (2015) 48–56.
- 475 [44] A. Seuret, F. Gouaisbaut, Wirtinger-based integral inequality: Application to time-delay systems, *Automatica* 49 (9) (2013) 2860–2866.
- [45] O. M. Kwon, S. M. Lee, H. P. Ju, E. J. Cha, New approaches on stability criteria for neural networks with interval time-varying delays, *Applied Mathematics and Computation* 218 (19) (2012) 9953–9964.



- [46] P. Park, J. W. Ko, C. Jeong, Reciprocally convex approach to stability of  
480 systems with time-varying delays, *Automatica* 47 (1) (2011) 235–238.
- [47] L. Duan, L. Huang, Periodicity and dissipativity for memristor-based mixed  
time-varying delayed neural networks via differential inclusions, *Neural  
Networks* 57 (9) (2014) 12–22.
- [48] H. B. Zeng, Y. He, M. Wu, S. P. Xiao, Passivity analysis for neural networks  
485 with a time-varying delay, *Neurocomputing* 74 (5) (2011) 730–734.
- [49] B. Zhang, S. Xu, J. Lam, Relaxed passivity conditions for neural networks  
with time-varying delays, *Neurocomputing* 142 (142) (2014) 299–306.



**Qianhua Fu** received the B.S. degree in electronic information engineering from Chongqing University of Technology, China, in 2003, and received the M.S. degree in communication and information systems from University of Electronic Science and Technology of China (UESTC) in 2010. He was a R&D engineer in HUAWEI company from 2010 to 2014. He is currently pursuing the Ph.D. degree in information and communication Engineering, UESTC and working as an engineer at Xihua University. His main research interests are memristor neural network, RF circuits and systems for wireless communications, and signal processing in modern communication.



**Jingye Cai** received the B.S. degree from Sichuan University in 1983, and the M.S. degree from the University of Electronic Science and Technology of China (UESTC), Chengdu, in 1990. He is currently a professor with the School of Software and Information Engineering, UESTC. His research interests include nonlinear circuits and systems (memristor), communication signal processing, frequency synthesis, RF and wireless systems.



**Shouming Zhong** was born on November 5, 1955. He graduated from University of Electronic Science and Technology of China, majoring Applied Mathematics on Differential Equation. He is a professor of School of Mathematical Sciences, University of Electronic Science and Technology of China, since June 1997-present. He is the director of Chinese Mathematical Biology Society, the chair of Biomathematics in Sichuan, and editor of Journal of Biomathematics. He has reviewed for many journals, such as Journal of Theory and Application on Control, Journal of Automation, Journal of Electronics, and Journal of Electronics Science. His research interest is stability theorem and its application research of the differential system, the robustness control, neural network and biomathematics.



**Yongbin Yu** received his M.S. degree and Ph.D. degree in circuits and systems from University of Electronic Science and Technology of China (UESTC) in 2004 and 2008. He is currently an associate professor with the School of Software and Information Engineering, UESTC. His current research interest covers nonlinear circuits and systems (memristor), artificial intelligence (neural networks, genetic algorithm), VLSI physical design, modern control theory and its application.