


DART Tutorial Section 9: More on Dealing with Error: Inflation






©UCAR 2014




The National Center for Atmospheric Research is sponsored by the National Science Foundation. Any opinions, findings and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.



Some Error Sources in Ensemble Filters



Dealing with Ensemble Filter Errors



Fix 1, 2, 3 independently, HARD but ongoing.


Often, ensemble filters...

1-4: Variance inflation, Increase prior uncertainty to give obs more impact.


5. 'Localization': only let obs. impact a set of 'nearby' state variables.

Often smoothly decrease impact to 0 as function of distance.


1. History of observations and physical system => 'true' distribution.



- 1. History of observations and physical system => 'true' distribution.
- 2. Sampling error, some model errors lead to insufficient prior variance.
- 3. Can lead to 'filter divergence': prior is too confident, obs. Ignored.




- 1. History of observations and physical system => 'true' distribution.
- 2. Sampling error, some model errors lead to insufficient prior variance.
- 3. Can lead to 'filter divergence': prior is too confident, obs. Ignored.




Naïve solution is variance inflation: just increase spread of prior. For ensemble member i, $inflate(x_i) = \sqrt{\lambda}(x_i - \overline{x}) + \overline{x}$


1. History of observations and physical system => 'true' distribution.



- 1. History of observations and physical system => 'true' distribution.
- 2. Most model errors also lead to erroneous shift in entire distribution.
- 3. Again, prior can be viewed as being TOO CERTAIN.



- 1. History of observations and physical system => 'true' distribution.
- 2. Most model errors also lead to erroneous shift in entire distribution.
- 3. Again, prior can be viewed as being TOO CERTAIN.



Inflating can ameliorate this.

Obviously, if we knew E(error), we'd correct for it directly.

Physical Space Variance Inflation

Inflate all state variables by same amount before assimilation.


Capabilities:

- 1. Can be effective for a variety of models.
- Can maintain linear balances.
- 3. Stays on local flat manifolds.
- 4. Simple and cheap.


Liabilities:

- State variables not constrained by observations can 'blow up'.
 For instance unobserved regions near the top of AGCMs.
- 2. Magnitude of λ normally selected by trial and error.


Observation outside prior: danger of filter divergence.



After inflating, observation is in prior cloud: filter divergence avoided.




Prior distribution is significantly 'curved'.



Observation in red.

Prior ensemble in green.

Inflated prior outside attractor. Posterior will also be off attractor.



Basic control of inflation in DART is in filter_nml

```
After
                       Before
                       Assimilation
                                                  Assimilation
inf flavor
                            = 0,
                                                        0,
                                                                   Flavor:
                                                                              1 => obs. space
                           = .false.,
                                                        .false.,
inf initial from restart
                                                                              2,3 => physical space
inf sd initial from restart = .false.,
                                                        .false.,
                                                                              0 \Rightarrow NONE
inf output restart
                           = .true.,
                                                        .true.,
inf deterministic
                           = .true.,
                                                        .true.,
inf in file name
                           = 'prior inflate ics',
                                                        'post inflate ics',
inf out file name
                           = 'prior inflate restart',
                                                        'post inflate restart',
inf diag file name
                           = 'prior inflate diag',
                                                        'post inflate diag',
inf initial
                           = 1.0,
                                                        1.0,
                                                                             Inflation
inf sd initial
                           = 0.0,
                                                        0.0,
                                                                             Value
inf damping
                           = 1.0,
                                                        1.0,
inf lower bound
                           = 1.0.
                                                        1.0,
inf upper bound
                           = 1000000.0,
                                                        1000000.0,
inf sd lower bound
                            = 0.0,
                                                        0.0,
```


Initially, we'll change inf_flavor and inf_initial in first column.

Set *inf_flavor=3*, state space inflation, in the first column.

Try some values and see what happens to L96 assimilation. Set *inf_initial* to values like 1.05, 1.08, 1.10 in the first column.


Make sure that *cutoff=10000000* and *ens_size=20* (These were settings that diverged without inflation)

Variance inflation in observation space



1. For observed variable, have estimate of prior-observed inconsistency.


Variance inflation in observation space



1. For observed variable, have estimate of prior-observed inconsistency.

2. Expected (prior_mean – observation) = $\sqrt{\sigma_{prior}^2 + \sigma_{obs}^2}$ Assumes that prior and observation are supposed to be unbiased. Is it model error or random chance?

Variance inflation in observation space



- 1. For observed variable, have estimate of prior-observed inconsistency.
- 2. Expected (prior_mean observation) = $\sqrt{\sigma_{prior}^2 + \sigma_{obs}^2}$
- 3. Inflating increases expected separation. Increases 'apparent' consistency between prior and observation.

Variance inflation in observation space: Lorenz 96

Try some values and see what happens to L96 assimilation. Set *inf_flavor=1*, observation space inflation in first column.

Try some values and see what happens to L96 assimilation. Set *inf_initial* to values like 1.05, 1.08, 1.10 in first column.

Make sure that *cutoff=10000000* and *ens_size=20*. (These were settings that diverged without inflation)

DART Tutorial Index to Sections

- 1. Filtering For a One Variable System
- 2. The DART Directory Tree
- 3. DART Runtime Control and Documentation
- 4. How should observations of a state variable impact an unobserved state variable? Multivariate assimilation.
- 5. Comprehensive Filtering Theory: Non-Identity Observations and the Joint Phase Space
- 6. Other Updates for An Observed Variable
- 7. Some Additional Low-Order Models
- 8. Dealing with Sampling Error
- 9. More on Dealing with Error; Inflation
- 10. Regression and Nonlinear Effects
- 11. Creating DART Executables
- 12. Adaptive Inflation
- 13. Hierarchical Group Filters and Localization
- 14. Quality control
- 15. DART Experiments: Control and Design
- 16. Diagnostic Output
- 17. Creating Observation Sequences
- 18. Lost in Phase Space: The Challenge of Not Knowing the Truth
- 19. DART-Compliant Models and Making Models Compliant
- 20. Model Parameter Estimation
- 21. Observation Types and Observing System Design
- 22. Parallel Algorithm Implementation
- 23. Location module design (not available)
- 24. Fixed lag smoother (not available)