Appendix

Compliance in regulatory gray areas: The case of the organic seed standard

Survey sample

1. Data collection method

- 2010: Data were collected by convenience sample only, shared by 23 organizations, including non-profits and foundations, organic certifiers, and departments of agriculture. Surveys were distributed primarily over email, but paper, phone, and in-person surveys were conducted at the request of the grower. We received a total of 1,027 responses (712 web-based, 315 paper-based), 542 of which included complete enough data for our analysis.
- 2015: Data were collected by both random and convenience sampling. For the random sample, 800 certified organic producers were randomly selected from the approximately 12,000 organic producers that use seed, recorded in the NOP database. To encourage responsiveness, three follow-up emails were sent following an initial email, paper copies were mailed on request, and individual phone calls were made to follow-up. From these 800, 197 (25%) responded (114 web-based, 83 paper-based) and 128 (16%) were complete enough to use in our analyses. For the convenience sample, surveys were shared by 41 organizations, primarily over email. We received a total of 1,148 responses from the convenience sample, 656 of which included complete enough data for our analysis. In total we had 784 usable responses.
- 2020: Data were collected by both random and convenience sampling using methods similar to 2015. For the random sample, 2,000 certified organic producers were randomly selected from the the NOP database and given the option to complete the survey over email or paper. After an initial email and postcard invitation, growers received follow-up contacts by email, postcard, and phone, and hard copies of the survey were sent. Survey distribution was managed by Washington State University's Social and Economic Sciences Research Center (SESRC) and distributed alongside the Organic Farming Research Foundation's (OFRF) National Organic Research Agenda. From these 2,000 surveys, 448 (22%) responded (206 web-based and 242 paper-based) and 185 (10%) were complete enough to use in our analyses. For the convenience sample, surveys were distributed by SESRC to all remaining certified organic operations over email, with postcard reminders. The survey was also advertised by Organic Seed Alliance (OSA) and OFRF. We received a total of 611 responses from the convenience sample, 240 of which included complete enough data for our analysis. In total we had 425 usable responses.

The breakdown between random and convenience sampling used in our analysis is summarized in Table S1. We consider the effect of sampling method, as well as survey distributor, through models in Section 2.

Table S1: Convenience and random samples, usable responses

Year	Total complete responses	Convenience responses (%)	Random responses (%)
2010	542	542 (100 %)	0 (0 %)
2015	784	656 (84 %)	128 (16 %)
2020	425	240 (56 %)	185 (44 %)

2. Checking for biases induced by data collection methods

In order to test how our data collection methods may bias the sample, we consider the effect of the sampling approach and survey distributor in our model. First we model the effect of sampling method with our 2015 and 2020 survey data (the only survey periods where we had both sampling approaches) by including a term for responses that were gathered by random sampling compared to convenience sampling (Table 2, Model 1). We see a small and insignificant effect for sampling method, and negligible changes in the results between the models run with and without the sampling consideration (Table 2, Model 2), suggesting no significant effect on results based on sampling approach.

Regarding concerns of survey distributor (i.e. that the distribution by OSA elicited responses from a select type of organic farmer), we model this effect with our 2010 and 2015 survey data (the only survey periods that identified specific distributors) by including a term for responses that were gathered through OSA channels compared to all others (Table 2, Model 3). We see a small and insignificant effect for distributor, and negligible changes in the results between the models run with and without the distributor consideration (Table 2, Model 4), suggesting no significant effect on results based on survey distributor.

Table S2: Beta regression model results including sampling and distributor effects

	Proportion of organic land planted with organic seed			
	Including sampling method (2015, 2020)	Without sampling method (2015, 2020)	Including survey distributor (2010, 2015)	Without survey distributor (2010, 2015)
	(1)	(2)	(3)	(4)
Capability	0.690***	0.692***	0.518***	0.509***
	(0.147)	(0.147)	(0.150)	(0.150)
Norms	0.314***	0.313***	0.286***	0.282***
	(0.053)	(0.053)	(0.046)	(0.046)
Certifier request	−0.016	−0.015	−0.141*	−0.139*
	(0.078)	(0.078)	(0.074)	(0.074)
Farm size (log acres)	−0.128***	−0.128***	−0.128***	−0.126***
	(0.024)	(0.024)	(0.023)	(0.023)
Crop diversity	0.118*	0.118*	0.133**	0.134**
	(0.069)	(0.069)	(0.063)	(0.063)
Seed saved or traded (%)	0.059	0.058	0.092***	0.090***
	(0.038)	(0.038)	(0.035)	(0.035)
Years certified organic	0.014	0.014	0.036	0.036
	(0.036)	(0.036)	(0.038)	(0.038)
Forage crops (vs. Field)	−0.948***	−0.947***	−1.017***	−1.021***
	(0.116)	(0.116)	(0.107)	(0.107)
Vegetable crops (vs. Field)	−0.565***	−0.567***	−0.613***	−0.622***
	(0.125)	(0.124)	(0.118)	(0.118)
Western region (vs. North Central)	−0.280***	−0.279***	−0.347***	−0.387***
	(0.102)	(0.102)	(0.102)	(0.098)
Northeastern region (vs. North Central)	−0.287***	−0.283***	−0.212*	−0.261**
	(0.107)	(0.106)	(0.119)	(0.114)
Southern region (vs. North Central)	−0.038	−0.038	−0.194	−0.221*
	(0.139)	(0.139)	(0.132)	(0.131)
Year: 2020 (vs. 2015)	0.058	0.066		
	(0.083)	(0.079)		
Random data collection (vs. convenience)	0.028			
	(0.090)			
Year: 2010 (vs. 2015)			−0.265***	−0.259***
			(0.079)	(0.079)
Distributor: OSA (vs. Other)			−0.118	
			(0.082)	
Constant	−0.124***	−0.124***	−0.096**	−0.093**
	(0.041)	(0.041)	(0.042)	(0.042)
Constant	−0.188	−0.182	0.114	0.099
	(0.204)	(0.202)	(0.181)	(0.181)
Observations	1,186	1,186	1,310	1,310
R^2	0.245	0.244	0.249	0.248
Log Likelihood	1,712.499	1,712.451	1,633.746	1,632.704

Note: *p<0.1; **p<0.05; ***p<0.01

3. Sample representativeness

While the mix of sampling methods creates a challenge for determining the sample's representativeness of the population, we aim to demonstrate that our sample is representative by comparing the geographic distribution

of operations (Table S3) and farm size distribution (Figure S1) between our sample and USDA Census data.

We first use USDA Census data of organic farmers from 2011, 2016, and 2021 to align with our 2010, 2015, and 2020 survey periods. Across the three survey periods, the number of respondents represented approximately 6.3%, 5.9%, and 2.5% of the total number of organic farmers identified in the USDA Census in the related years. In Table S3 we compare the number of operations in our sample (n) to the population (N) by year and region. For each census period we calculated the standard deviation of the percentage of operations represented, which we used to identify regions that are over or under-represented. Over-represented year-region combinations are signaled by italic text formatting, while under-represented year-region combinations are signaled by bold text formatting. Only in 2010 do we see significant variation in representation across regions, with the North Central and Southern regions over-represented and the Northeast under-represented.

Table S3: Sample representativeness compared to USDA Censuses (2011, 2016, 2021)

Year	Region	Operations in sample (n)	Operations in USDA census (N)	Percent of operations represented (n/N)
2010	North Central	305	2986	*10.2*
2010	Northeast	51	1897	**2.7**
2010	South	51	540	*9.4*
2010	West	139	3296	4.2
2015	North Central	386	4689	8.2
2015	Northeast	141	3246	4.3
2015	South	75	1096	6.8
2015	West	189	4529	4.2
2020	North Central	174	5824	3
2020	Northeast	98	3888	2.5
2020	South	38	1209	3.1
2020	West	98	5273	1.9

Note:
Italic formatting sigals over-representation and bold formatting signals under-representation

Next, to understand farm size representativeness, we look at USDA Census organic tabulations, reported for as the 2012, 2017, and 2022 census tabulations.

In Figure S1 we compare these USDA Census data against our survey sample. These comparisons show a consistent over-representation of the smallest farms (1-9 acres), especially in the North Central and South, and under-representation of the largest farms. Still, our sample captures a distribution of organic farm sizes that includes medium and large farms, allowing us to understand the effect of farm size in our analysis. Additionally, an important consideration is that we calculate farm size as reported acreage planted to three major crop categories, but this approach likely underestimates acreage generally compared to USDA Census data, which reports all certified land. Furthermore these data are not crop specific and so USDA data also includes non-seed crop operations (i.e. orchards).




Figure S1: Sample farm size distribution compared to USDA Census data

Model checks

4. Dissagregating allowable and non-allowable factors (Table S4)

Table S4: Beta regression model results disaggregating allowable and non-allowable ability factors

	Proportion of production acreage planted with organic seed			
Ability: Allowable	0.192***	0.386***	0.192***	0.297**
	(0.031)	(0.122)	(0.031)	(0.131)
Ability: Non-allowable	0.108***	0.105***	0.311***	0.277***
	(0.026)	(0.026)	(0.093)	(0.101)
Norms	0.200***	0.235***	0.287***	0.292***
	(0.038)	(0.043)	(0.053)	(0.053)
Certifier request	−0.048	−0.050	−0.051	−0.052
	(0.065)	(0.065)	(0.065)	(0.065)
Farm size (log acres)	−0.118***	−0.116***	−0.116***	−0.115***
	(0.021)	(0.021)	(0.021)	(0.021)
Crop diversity	0.177***	0.180***	0.179***	0.180***
	(0.056)	(0.056)	(0.056)	(0.056)
Seed saved or traded (%)	0.104***	0.104***	0.101***	0.102***
	(0.032)	(0.032)	(0.032)	(0.032)
Years certified organic	0.034	0.034	0.035	0.035
	(0.032)	(0.032)	(0.032)	(0.032)
Forage crops (vs. Field)	−0.899***	−0.901***	−0.900***	−0.900***
	(0.093)	(0.093)	(0.093)	(0.093)
Vegetable crops (vs. Field)	−0.496***	−0.485***	−0.480***	−0.476***
	(0.103)	(0.103)	(0.103)	(0.103)
Western region (vs. North Central)	−0.380***	−0.379***	−0.386***	−0.385***
	(0.085)	(0.085)	(0.085)	(0.085)
Northeastern region (vs. North Central)	−0.290***	−0.291***	−0.296***	−0.296***
	(0.094)	(0.094)	(0.094)	(0.094)
Southern region (vs. North Central)	−0.174	−0.170	−0.178	−0.176
	(0.115)	(0.115)	(0.115)	(0.115)
Year: 2010 (vs. 2015)	−0.215***	−0.213***	−0.217***	−0.215***
	(0.078)	(0.078)	(0.078)	(0.078)
Year: 2020 (vs. 2015)	0.101	0.098	0.098	0.097
	(0.079)	(0.079)	(0.079)	(0.079)
Ability: Allowable x Duty		−0.057*		−0.031
		(0.034)		(0.037)
Ability: Non-allowable x Duty			−0.062**	−0.052*
			(0.027)	(0.030)
Constant	0.002	−0.113	−0.269	−0.288
	(0.154)	(0.167)	(0.192)	(0.193)
Observations	1,725	1,725	1,725	1,725
R^2	0.237	0.238	0.239	0.239
Log Likelihood	2,239.216	2,240.591	2,241.742	2,242.084

Note: *p<0.1; **p<0.05; ***p<0.01

5. Correlations of all ability factors (Table S5)

Table S5: Correlations among organic seed sourcing ability factors

Barrier	Availability	Quantity	Quality	Form	Price
Quantity	0.50	NA	NA	NA	NA
Quality	0.64	0.42	NA	NA	NA
Form	0.20	0.22	0.23	NA	NA
Price	0.36	0.31	0.39	0.34	NA
Buyer requirement	0.41	0.38	0.27	0.21	0.22