

Malaria in Madagascar

Benjamin Rice

November 29, 2017

Supplementary Figure 2: Map of Madagascar showing the regions sampled and some climatic characteristics





Figure 1: Schematic of nested study design where individuals were sampled within households at 6 sites per region for 4 regions



01	02	03	04	05	06	07	08
09	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	00	01	02	03	04

05	06	07	08	09	10	11	12
13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28
29	30	31	32	33	34	35	36
37	38	39	40	41	42	43	44
45	46	47	48	49	50	51	52
53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68
69	70	71	72	73	74	75	76
77	78	79	80	81	82	83	84
85	86	87	88	89	90	91	92
93	94	95	96	97	98	99	00

01
 02
 03
 04
 05
 06
 07
 08
 09
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 00




[illegible]

Summary of data set analyzed									
Regions	4								
Sites	24								
Households	1129								
Individuals	6293								
Mean household size (number of individuals)	5.57								
Household size range	(2-19)								
Individuals with valid malaria result	5544								
Percent of individuals with a malaria result (%)	88.1								
Number of malaria positive individuals	776								
Malaria prevalence (overall) (%)	13.99								
Mean age of individuals (years)	17.34								

Spatial Clustering Example: Site R2V4



**Southeast:
Vatovavy Fitovinany
(Mananjary)**



R2V4

Spatial Clustering Example: Site R2V4



Spatial Clustering Example: Site R2V4



Spatial Clustering Example: Site R2V4





Figure 3: Density plots of age for all individuals, those infected with malaria, and uninfected with malaria (n = 5533)




Figure 2: Variation in the percentage of individuals infected (prevalence) between sites within regions and between regions




Table 2: Using a simple generalized linear model of four site-specific predictors of malaria prevalence to demonstrate the need for accounting for inter-regional differences

Table 2A: All regions combined ($n = 24$ sites)

Predictor	Coefficient	Confidence interval
Travel time by foot to nearest doctor (hours) (continuous)	-0.73	[-2.31, 0.85]
Road access (binary)	-10.64	[-23.15, 1.87]
Coastal versus inland (binary)	15.56	[2.25, 28.87]
Distance to nearest hospital (km) (integer)	0.31	[-0.099, 0.73]

Table 2B: Excluding Region 5 ($n = 18$ sites)

Predictor	Coefficient	Confidence interval
Travel time by foot to nearest doctor (hours) (continuous)	-0.93	[-3.0684, 1.2084]
Road access (binary)	-13.73	[-29.2334, 1.7734]
Coastal versus inland (binary)	11.72	[-5.506, 28.946]
Distance to nearest hospital (km) (integer)	0.28	[-0.215, 0.775]

Supplementary Table 2: Parameter values: Empirically derived, chosen simulation inputs or simulation outputs for the intercepts, coefficients, and residual standard deviations show

	Empirical estimate	Simulation (1X)* Inputs	Simulation (1X) Outputs (mean)	Simulation (2X)* Inputs	Simulation (2X) Outputs (mean)
Intercept	14.09	14	5.64	14	18.72
Travel time by foot to nearest doctor (hours)	-0.72	-0.70	-0.71	-1.50	-1.49
Distance to nearest hospital (km)	0.14	0.13	0.13	0.26	0.26
Residual <u>sd</u>	15.61	15	14.84	7.30	7.18

*In the "2X" simulation scenario the coefficients (i.e. slopes) were increased approximately by a factor of two and the residual sd was decreased by approximately a fact of two (intercept unchanged). For both simulation scenarios, the proportion of repetitions in which the simulation returned parameter estimates with confidence intervals that covered the inputted parameter values were recorded. For both coefficients in both simulation scenarios, the proportions were above 0.93.

Summary of data set analyzed

Regions

4

Sites

24

Households

1129

Individuals

6293

Mean household size (number of individuals)

5.57

Household size range

(2-19)

Individuals with valid malaria result

5544

Percent of individuals with a malaria result (%)

88.1

Number of malaria positive individuals

776

Malaria prevalence (overall) (%)

13.99

Mean age of individuals (years)

17.34

Table 3: Mean household size as a site predictor – not enough sites to have sufficient statistical power

Model: $Pr(\text{infection}) = \alpha + \beta * (\text{mean household size}) + \text{error}$

Model parameter estimates	All regions included*	Region 5 excluded*
α	38.089 [1.20, 74.97]	7.44 [-76.01, 90.89]
β	-4.301 [-10.749, 2.14]	2.04 [-13.58, 17.66]
Residual sd	15.03	15.65

*95% confidence intervals shown in brackets next to parameter estimates

Figure 2: Variation in the percentage of individuals infected (prevalence) between sites within regions and between regions



Table 3: Mean household size as a site predictor – not enough sites to have sufficient statistical power

$$\text{Model: } \text{Pr}(\text{infection}) = \alpha + \beta * (\text{mean household size}) + \text{error}$$

Model parameter estimates	All regions included*	Region 5 excluded*
α	38.089 [1.20, 74.97]	7.44 [-76.01, 90.89]
β	-4.301 [-10.749, 2.14]	2.04 [-13.58, 17.66]
Residual sd	15.03	15.65

*95% confidence intervals shown in brackets next to parameter estimates