The problem and promise of modeling plant leafout under warmer winters

E.M. Wolkovich[1*], Justin Ngo[1], Victor Van der Meersch[1], Robert D. Guy[1] & Jonathan Auerbach[2]

November 7, 2026

[1] Forest & Conservation Sciences, Faculty of Forestry & Environmental Stewardship, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4, Canada

[2] Department of Statistics, George Mason University, 4511 Patriot Circle, Fairfax, VA 22030, United States

* e.wolkovich@ubc.ca

Abstract

How warmer winters affect plant leafout has major implications for carbon storage and ecosystem stability. Recent evidence that the advance of spring budburst has slowed with anthropogenic climate change has reignited a decades-old debate over winter chilling—the concept that woody temperate plants require sufficient winter cool temperatures to budburst each spring. The problem with chilling is that it does not correspond with an observable event. This makes models of it rarely falsifiable and of limited use for advancing scientific understanding. In this paper, we review how current research built on the concept of chilling combined with limited data has led to the proliferation of untestable models that are consistent with a wide range of forecasts and provide little insight into actual plant biology. We argue that the concept of chilling should be reevaluated like past theoretical constructs of limited scientific value (such as 'dormin' or 'anthesin') to help accelerate our understanding of plant development. This can be done by leveraging new insights on the biochemistry and physiology of dormancy to build better models that fit to both experimental and observational data, and by designing better experiments to more rigorously test theory.

Keywords: Chilling, spring phenology, dormancy, endodormancy, process-based modeling, forests

Plant leafout in the spring has shifted weeks earlier in many regions due to warming from anthropogenic climate change with consequences for a wide range of ecosystem services (Keenan *et al.*, 2014; Pörtner *et al.*, 2022). Which underlying processes drive this trend, however, is debated, as recent research suggests winter warming may slow or stall further advances (Fu *et al.*, 2015; Piao *et al.*, 2017). Such reports focus on a two-step model of leafout where plants first require cool winter temperatures, often called 'chilling,' before they can accumulate enough warm temperatures—'forcing'—to leafout each year (Figure 1). This model of chilling is only one of many models proposed since the concept was first introduced (Basler, 2016; Dennis, 2003; Spafford *et al.*, 2025). But, while forcing ends in an observable event such as leafout, the day plants achieve their chilling requirements coincides with no known measurable event.

Despite chilling relating to an unobserved event, decades of work suggest it impacts budburst. Experimental studies, which use controlled temperatures to vary the hypothesized amount of chilling, almost always find evidence that cool temperatures affect the timing and/or success of plant budburst (Erez & Lavee, 1971; Charrier *et al.*, 2015; Baumgarten *et al.*, 2021). Observational studies, which estimate chilling requirements or effects of chilling using observed budburst and temperature data from plants (usually woody plants) in natural settings, report similar effects. Models generally build on both types of data to predict how winter temperatures impact leafout (Charrier *et al.*, 2015; Baumgarten *et al.*, 2021), but new research has highlighted major potential flaws these models.

Current models of chilling predict wildly different leafout times for the same study system and temperature regime (Guy, 2014; Chuine *et al.*, 2016). Even simple models yield radically different predictions for the exact same conditions depending on, for example, the calibration data (Olsson & Jönsson, 2014) or which parameters are treated as 'known' (Figure 2). Such extreme variability of spring phenology predicted from current models highlights fundamental gaps in our understanding. At the same time, multiple papers now suggest estimates of 'chilling' could easily be artifacts of poor models or correlated observational climate data (Wolkovich *et al.*, 2021; Gao *et al.*, 2024).

Now appears an opportune time to address the problems in our concept of chilling in models of leafout. Because accurate forecasts of spring phenology are critical for carbon sequestration (Green & Keenan, 2022; Keenan *et al.*, 2014), crop yields and relating food security (Lee *et al.*, 2024; Leisner, 2020), there is widespread interest in improving chilling models (Luedeling *et al.*, 2015; Chuine *et al.*, 2016). At the same time, recent results from molecular and cellular studies of

dormancy are providing insights into when and how chilling works (Pan *et al.*, 2023; Zhu *et al.*, 2021), which could rule in—or out—some of the myriad current models. Here, we review the theory of chilling, its origins and potential problems, as well as new opportunities for major advances and how shifting current practices could accelerate progress. We argue that revisiting the current concept of chilling—and its utility—could help build better models and design better experiments.

What is chilling?

How plants in temperature-limited systems avoid leafout during warm spells in the winter has long been debated by plant biologists (e.g., Lamb, 1948; Weinberger, 1950). Most work to date has focused on the idea that plants enter some form of dormancy in the fall, which is then released before warm temperatures in the spring begin. This idea hypothesizes that the slow accumulation of cool temperatures—or chilling—over the winter extends through periods of short warm spells in the winter and thus prevents leafout before spring.

The term 'chilling' is now used across numerous fields in plant biology to refer to a process where dormant embryonic tissues exposed to cool temperatures accelerate a developmental event—usually budburst—that later occurs after warm temperatures (Fig. 1). The term relates to vernalization—cool temperatures required for flowering (Kim *et al.*, 2009; Kim, 2020)—with much work on the herbaceous species *Arabidopsis thaliana*. Today the term 'chilling' is most often used for dormant buds that later transition to leafout or flowering, but also used for seeds (though 'cold stratification,' which indicates an additional requirement of moist conditions, is more often used for seeds, Baskin & Baskin, 2000; Considine & Considine, 2016). While we also focus on chilling in regards to bud dormancy, the mechanisms underlying chilling requirements of seeds likely provide important parallels to bud dormancy (Leida *et al.*, 2012; Penfield & Springthorpe, 2012; Sato & Yamane, 2024); evolution is more likely to co-opt an existing mechanism than to start de novo.

Much of our fundamental understanding of chilling for buds comes from studies on temperate woody fruit crops where chilling can be critical to yield. Peach trees planted into warmer climates well outside their range often have extremely low fruitset because most flower buds do not burst (Weinberger, 1950; Overcash & Campbell, 1955; Erez *et al.*, 1971). Early studies of this phenomenon with related experiments found cuttings (cut ends of dormant branches from trees) exposed to cool temperatures in controlled chambers (with a standard of 0-7.2°C for many fruit tree studies Weinberger, 1950) burst more fully and more quickly (Vitasse & Basler, 2014; Ettinger *et al.*, 2020).

These studies underlie most of the models of chilling used today for crops and wild tree species (Dennis, 2003).

Focused on how chilling can accelerate observable events, researchers have calculated how much 'chilling' is required for leafout of forest trees from cutting experiments similar to those used for peaches (reviewed in Ettinger *et al.*, 2020), ground observations of budburst (Luedeling *et al.*, 2009b), and satellite measures of greenup (Kaduk & Los, 2011). These estimates rely on phenological models that have become critical across a suite of fields. In climatology and global change biology, modeling how vegetation on the land surface responds to anthropogenic climate change affects spring leafout and carbon storage (Schwartz, 1994; Cleland *et al.*, 2007; Chuine *et al.*, 2013). In crop biology 'chill units' guide growers in which specific cultivar to plant (Figure **??**). Alongside these more macro-scale studies of chilling, molecular and cellular studies have examined variation in hormones, transcripts and other cellular compounds that coincide or precede budburst, generally using controlled temperatures to vary the hypothesized amount of chilling (e.g., Azeez *et al.*, 2021; Cai *et al.*, 2024, and see Fig. 4).

Today these studies have led to over 30 basic models where accumulated chilling releases plants from dormancy (Basler, 2016; Hufkens *et al.*, 2018). Though early debates considered whether plants were truly dormant or only growing slow ('dormancy' or 'rest' versus 'quiescence'; Considine & Considine, 2016). Today most research assumes a model with two phases of dormancy: (1) endodormany, the period when chilling occurs, and (2) a period between endodormancy but before the observed event, called ecodormancy (Figure 4).

In most models, chilling only accumulates under certain temperatures—traditionally above zero but below 10°C—with certain temperatures being optimal for the most rapid accumulation of 'chill units,' where some unknown sum of chill units breaks endodormancy (Figure **??**). Early models introduced 'chill hours' where chilling only accumulated during hours where the temperature was 0-7.2°C (corresponding to the range of 32-45°F developed for peach trees, Weinberger, 1950). After this, the Utah model expanded the range of temperatures to 0-16°C and allowed higher temperatures (>16°C) to negate previous accumulated chilling (Richardson, 1974); such 'chill units' from the Utah model is still widely used, though the term 'chill units' is often used more broadly (as we do here). Since then, more complex models have continually been developed including the Dynamic model, which is broadly similar to the Utah model in its range (0-18°C) and negation but adds in a response that varies with duration of exposure and can be enhanced depending on how

the temperatures fluctuate Erez *et al.* (1987); Erez & Fishman (1997); Fishman *et al.* (1987). The dynamic model introduced the term 'chill portions' and has been widely promoted for fruit tree chilling estimates (Luedeling *et al.*, 2009a, 2021). For many forest trees, which temperatures are most effective at providing chilling is a common question continually addressed in experiments, with different experiments providing different answers (Vitasse *et al.*, 2013; Baumgarten *et al.*, 2021).

Variation in responses to chilling across and within species yields hundreds of variants of the basic models (Basler, 2016; Hufkens *et al.*, 2018). Complexity comes from the hypothesized diversity of chilling temperature thresholds and sums across species and populations. Most assume different species require different sums of chill units, and may have different lower, upper, and optimal chill temperatures. Within species, populations may require different sums of chill units, with populations in more mild climates—where warm interruptions in the winter are more common—requiring more chill units than those in areas with cold winters, where temperatures rarely rise above zero before spring (Campbell & Sugano, 1979; Leinonen, 1996).

The problem with chilling

An unobserved process

Chilling is a latent, unobserved process. Typically, 'chilling' describes the physiological phase (endodormancy) in which a plant experiences environmental temperatures that induce progress towards the next physiological phase (ecodormancy) that ends in budburst (Figure 1). The problem is that the transition from endo- to ecodormancy currently corresponds with no clearly measurable phenomenon, and thus the properties of each period cannot be determined from the observed budburst, even under experimental conditions in which the temperature can be manipulated.

The parameters of a model matching this hypothesized dormancy process are said to be underdetermined or unidentified. For example, consider a simple model in which one would need to estimate the minimum and maximum temperatures that allow chilling to accumulate (two parameters) and the total sum of those temperature units needed to trigger a shift into the next physiological phase (often called 'endodormancy break') for one additional parameter. Models then need to estimate when plants start and stop accumulating (two more parameters). An experiment that raises temperatures and observes an earlier leafout could be explained by more rapid chilling accumulation leading to earlier endodormancy break, more rapid ecodormancy break, or an almost limitless mix

of the two (Figures 1-2).

To address this, models often assign the start date of endodormancy as known (e.g., starting 1 September in the northern hemisphere) and rely on assumptions to set the end date (Figure 2). A common and widely-used assumption, developed by early work on peaches, is that high and rapid budburst (leaf or flower) is evidence that chilling has been met (i.e. endodormancy has ended, Erez & Lavee, 1971, see also Figure 1), but this assumption has rarely been evaluated for other taxa or populations. The approach of assigning some unknown parameters as known has the benefit of avoiding adding more unknown model parameters, but has led researchers to be overly confident in a model of chilling where more is actually unobserved and unknown than acknowledged.

Hidden assumptions in complicated models

More complex models may seem to solve the identification problem, but often come with their own untestable assumptions. Even if we assume that high and rapid percent budburst signals sufficient chilling, most models today include parameters that cannot be uniquely identified with current data, which explains how one model can often predict a wide variety of leafout possibilities (Figure 2). The current tacit approach to these major issues likely has contributed to the expansion of models over the last few decades without any clear advances. While various models have added complexity via the shape of the optimal temperature range for chilling, allowing accumulated chilling to be reduced, shifting the start date of chilling, and/or allowing chilling and forcing to act simultaneously (Hänninen, 1990; Kramer, 1994; Luedeling *et al.*, 2009a; Pope *et al.*, 2014; Gusewell *et al.*, 2017), none of these have swept through the field.

As experiments and models have become more complicated, data have remained relatively uninformative to try to estimate all the complexities the models include. Further, recent models have often relied on even less data (Hänninen *et al.*, 2019). Many current methods use only observational data of the timing of leafout (or flowering) to attempt to estimate a model of chilling for different species or locations and project it forward to understand effects of anthropogenic climate change (Luedeling *et al.*, 2011; Luedeling, 2012; Gao *et al.*, 2024; Fernandez *et al.*, 2022). Perhaps not surprisingly then, which model is deemed best varies strongly by method, underlying data and approach (Caffarra *et al.*, 2011; Olsson & Jönsson, 2014; Basler, 2016; Hufkens *et al.*, 2018), with no clear pattern.

Unclear aims: Prediction versus mechanistic understanding

The flood of current models of budburst with similar levels of predictive accuracy is not necessarily a problem, depending on the aim. Many models of chilling are used to guide crop and tree planting—preventing planting too far outside the range of climates with sufficient chilling for a species, population, or cultivar. For this aim, models need to translate climate into relevant chilling and forcing values and define appropriate thresholds, but how exactly each model defines chilling and forcing is immaterial if the predictions are accurate enough (many models of 'chill units' are thus based simply on average winter conditions). In such cases moving towards machine learning models—which emphasize predictive accuracy often through black box approaches—may provide more practical models that also clearly acknowledge that the underlying process is obscure (Khodadadzadeh *et al.*, 2024; Bauer *et al.*, 2026).

In contrast, if the aim is biological understanding, then the number of highly contrasting but similarly predictive models suggests a problem with the current approach. Models to help elucidate the process that underlies the concept of chilling should represent the current biological understanding. Thus the specification of the model—what parameters it includes, and how well they can be estimated from the data—becomes more important, as does how to evaluate models. Tests that compare models for how accurate they represent our understanding of plant biology, and not only their predictive accuracies across a range of climate regimes, should guide model development. For this aim, new models are unlikely to be enough without new data and approaches that target the major problems facing models of chilling.

Overcoming the chilling problem

Richer, more informative data from molecular biology studies and other approaches (Fouché *et al.*, 2023; Walde *et al.*, 2024) hold the promise to elevate chilling from an unobserved complex process to something we understand and can robustly forecast. Taking full advantage of this opportunity, however, would benefit from re-examining the concept of chilling across different disciplines to combine observational and experimental evidence, while providing the opportunity to question the future utility of chilling.

Leverage new molecular insights to overcome problems in the simplest models

New molecular research provides hope that the transition from endo- to ecodormancy is measurable. Molecular insights have long contributed to crop and forest tree models of chilling (Chuine & Regniere, 2017). Decades of work on vernalization have outlined the pathways—and genes—that lead to flowering only after winter's cool temperatures in annual/biennial (herbaceous) populations of *Arabidopsis thaliana* (Figure 4, Wilczek *et al.*, 2009; Kim, 2020). Research has linked some of these pathways to similar ones in woody species and have also highlighted the polysaccharide callose (1,3-β-D-glucan) as potentially pivotal for chilling (van der Schoot *et al.*, 2014; Pan *et al.*, 2023).

Multiple studies across multiple species have now shown that (1) lower temperatures appear to degrade callose and (2) the release/loss of callose appears to re-start cell-to-cell communication before budburst (van der Schoot *et al.*, 2014; Pandey *et al.*, 2026). Taken together, these results suggest the loss of callose may be an indicator of endodormancy release.

If callose is a major controller of endodormancy and its release, chilling models could be limited to those that match the idea of callose degradation. This would include models that have a temperature range over which the enzyme(s) degrading callose is active (Figure 4). In contrast, models using simple temperature thresholds (e.g., all hours below -5°C equally allow chilling) would appear less biologically accurate, as enzymes generally do not work over such a wide range of temperatures. Models have begun to introduce the idea of factors or enzymes that may control dormancy or frost hardiness, with many now focusing on how starch to sugar dynamics may mediate budbreak (Charrier *et al.*, 2018; Sperling *et al.*, 2019). To date, however, most only add these parts as additional complexity onto of existing models (Caspersen *et al.*, 2025; Charrier *et al.*, 2018), which may limit progress (see 'Hidden assumptions in complicated models').

Callose removal itself may also represent only one downstream event of a larger process aligned to chilling. Many other factors, such as phytohormones (e.g., GA and ABA), also often change at the same time (Figure 4, Rinne *et al.*, 2018; Tylewicz *et al.*, 2018; Pan *et al.*, 2023; André *et al.*, 2022). Yet many of these compounds are also integral to other plant development and defense pathways, making it difficult to tie them directly to bud dormancy and budbreak. Increasing work, however, suggests promise through measuring DAMs (dormancy associated MADS-box genes), which are master regulators of various pathways involved in dormancy for both buds and seeds (Leida

et al., 2012; Tuan *et al.*, 2017). For example, Zhang *et al.* (2023) recently showed (PpDAM6 and PpMAPK6) may be useful markers for endodormancy maintenance through their effects on ABA—and thus transitions to ecodormancy. Similarly upregulation of 1,3-β-glucanases responsible for plasmodesmata reopening (Rinne *et al.*, 2011; Singh *et al.*, 2018) suggests the 1,3--glucanase family could be yet another potential marker for modelling dormancy progression that eventually allows water reactivation into cells and thus growth (Faust *et al.*, 1991; Kalcsits *et al.*, 2009; Walde *et al.*, 2024)—the event currently used in models. Making these new findings most useful to improving models of chilling will require tying these genes more fully to the process and then finding optimum temperature for their expression and bioactivity (see also supplement, 'The search for better indicators').

Other recent insights similarly suggest that most simplified temperature metrics used in many chilling models may not map to molecular realities. For example, new work on how slow growth itself may act a 'long-term thermosensor' (Zhao *et al.*, 2020) adds to an increasing number of molecular studies that suggest plants integrate long-term thermo-sensing in the winter alongside responses to short-term temperatures (Antoniou-Kourounioti *et al.*, 2021; Satake *et al.*, 2022). The best models of leafout may thus need to integrate across multiple timescales. While this could easily complicate models already burdened with complexity, insights from molecular biology could rule out models by focusing on new experiments and modeling approaches.

Model experimental and observational data together to test assumptions

Research on chilling could accelerate by working across what today are three fairly separated silos: 1) model building for crops and other forecasting needs (usually called process-based models), 2) experiments at the branch or whole-plant level, and 3) molecular research. Currently, crop biologists, phenological process-based modelers of forest trees, molecular biologists, and modelers of related processes (e.g., hardiness) all develop unique and rarely compared models of dormancy and budburst (but see Kovaleski, 2022), highlighting a major problem, but also a major opportunity. Synthesizing and benchmarking models—and their underlying biological understanding of chilling—across the many research fields developing chilling models today would help identify models that are equivalent and/or perform especially poorly, and could begin to discard some models.

Comparing models more directly would hopefully drive new approaches that estimate chilling by fitting experimental and observational data together in one model. This is rarely done, in part be-

cause of how differently they may be observed, including the challenging diversity of environmental conditions across these two data types (for example, many experiments apply cold temperatures in the dark and include extreme temperature differences, while in observational data photoperiod shifts each day and temperatures are more similar) but also because of separate modeling approaches. Researchers almost never fit process-based models to new empirical data; instead they use so-called 'statistical models' that often follow canonical treatment designs (e.g., ANOVA). Statistical models are usually far simpler, and make a suite of unstated assumptions that contradict the current understanding of chilling; for example, most statistical models used assume a linear response to chilling temperatures and durations, whereas all process-based model include an accumulation function, which is inherently non-linear (Figure fig:multimodelexps Wolkovich *et al.*, 2021). Many of the original studies that led to the concept of chilling, however, bridged across observational and experimental studies more often and leveraged datasets that created greater extremes in observational data—by focusing on crops planted well outside their natural range (e.g., peaches in Florida and Israel, Erez & Lavee, 1971; Richardson, 1974). These approaches did this in part by more explicitly mapping from biological processes to statistical models, making it easier to use the models to define new tests to challenge the biological theory.

Experiments bridging across methods may have the greatest opportunity to provide data that would truly challenge current models of chilling. Testing models across large environmental gradients in the field is one of the best ways to find out where models work—and where they fail. For example, Wilczek *et al.* (2009) tested vernalization models by comparing predicted to observed flowering times in a common garden design across Europe and supported the temperature-dependent growth model by testing its predictions of what happens when growth is altered but temperature is held constant (Zhao *et al.*, 2020). Similar examples for challenging other models of chilling date back over 40 years to when many of the models used today were developed (Richardson, 1974; Chuine *et al.*, 2016; Ettinger *et al.*, 2020), but could take place now. Models of chilling can make predictions under lower field chilling then test them using individuals planted beyond the range (either by planting those individuals now or identifying such cases in forestry provenance trials or similar). Process-based modelers could also challenge their models through more dramatic variation in biology; new experiments that include genetic mutants or similar variants (e.g., Azeez *et al.*, 2021; Pan *et al.*, 2023) could help test the effects of genes that putatively influence endo/ecodormancy under a spectrum of natural conditions (Satake *et al.*, 2022).

Is the concept of chilling necessary for scientific progress?

Results from new molecular experiments designed to identify novel ways to directly observe and measure chilling could upend the current concept of chilling. As increasing evidence suggests the concept aligns with multiple shifts at the cellular and molecular level (Figure 4), researchers across disciplines will need to consider how to continue using the term 'chilling.' One approach is to try to align previous and current definitions through new studies, for example, by testing for callose loss (Yu *et al.*, 2024) alongside previously used markers of dormancy shifts—including the often-used bioassay of high and rapid budburst at higher temperatures (indicating endodormancy release) and additional methods, such as weighing flower buds (Chuine *et al.*, 2016) or tracing water reactivation into cells (Faust *et al.*, 1991; Kalcsits *et al.*, 2009; Walde *et al.*, 2024). Another approach is to work towards new terms that more clearly align with new measurements and insights.

Because chilling is an unobserved process, it may link eventually to one measurable compound or regulator (similar to florigen or auxin) or to a more complicated biology, making the term less useful (Aksenova *et al.*, 2006). Current results suggest the latter, and already the term 'chilling' is used to mean different things. For example, chilling sometimes refers to an accumulation process during an unobserved physiological phase (endodormancy), or to the environmental conditions that cause that plants then respond to with an underlying process. In many experiments, the term 'chilling' refers to a treatment where researchers do not know the physiological phase (for example, cuttings or buds from woody plants are often chilled at 5°C for 6 weeks in the dark in a 'chilling' treatment, then transferred to warm 'forcing' conditions, without any tests that could confirm the phase, such as recording percent budburst at high temperatures, or tracking cellular processes Flynn & Wolkovich, 2018; Ettinger *et al.*, 2020).

Diverging definitions suggest that the best path for chilling may be to follow that of other terms modern plant biologists may not remember, such as dormin or anthesin (hypothesized compounds to induce dormancy and floral anthesis, respectively, Aksenova *et al.*, 2006; Dörffling, 2015), and let the term fade into obsolescence. Fundamental research on the physiological and molecular biological mechanisms of bud dormancy is primed to discover new measurable processes and observable events related to what we call 'chilling' today. With this, new terms and concepts that better match our improved understanding could emerge to aid research across the disciplines studying chilling. Models for forecasting and new experiments at the branch and whole plant level could focus on predicting and measuring these new events and processes. 'Chill units' for widely planted forest trees

and specific varieties of crops, which are already less linked to the mechanism of chilling, would likely hold on longer. They too, however, may shift to new estimates or units based on better measures, especially as increasing anthropogenic climate change makes getting the mechanistic biology of plant development right all the more important.

Acknowledgements: We thank two anonymous reviewers for feedback that improved the manuscript. We did not use any any Artificial Intelligence Generated Content tools in the development of this manuscript beyond several brief queries after writing the manuscript to confirm that we were not missing any relevant literature and separately verified all information ourselves (e.g., we learned of 'dormin' from one query, but this result also gave a list of hypothetical compounds that do not ever appear to have existed).

Author contributions: All authors contributed to developing the research and editing the manuscript. In addition EMW and JN led the literature review, EMW, VVdM, JN and JA developed the figures. EMW wrote the manuscript.

1 References

Aksenova, N., Milyaeva, E. & Romanov, G. (2006) Florigen goes molecular: seventy years of the hormonal theory of flowering regulation. *Russian Journal of Plant Physiology* **53**, 401–406.

André, D., Zambrano, J.A., Zhang, B., Lee, K.C., Rühl, M., Marcon, A. & Nilsson, O. (2022) Populus svl acts in leaves to modulate the timing of growth cessation and bud set. *Frontiers in Plant Science* **13**, 823019.

Antoniou-Kourounioti, R.L., Zhao, Y., Dean, C. & Howard, M. (2021) Feeling every bit of winter–distributed temperature sensitivity in vernalization. *Frontiers in Plant Science* **12**, 628726.

Azeez, A., Zhao, Y.C., Singh, R.K., Yordanov, Y.S., Dash, M., Miskolczi, P., Stojkovič, K., Strauss, S.H., Bhalerao, R.P. & Busov, V.B. (2021) EARLY BUD-BREAK 1 and EARLY BUD-BREAK 3 control resumption of poplar growth after winter dormancy. *Nature communications* **12**, 1123.

Baskin, C.C. & Baskin, J.M. (2000) *Seeds: ecology, biogeography, and, evolution of dormancy and germination*. Academic press.

Basler, D. (2016) Evaluating phenological models for the prediction of leaf-out dates in six temperate tree species across central europe. *Agricultural and Forest Meteorology* **217**, 10–21.

Bauer, J.N., Schiffers, K., Caspersen, L., Yamane, H. & Luedeling, E. (2026) Rethinking model transferability: Validity domains as a new approach to delineate the limits of bloom date projections. *Global Change Biology* **32**, e70776.

Baumgarten, F., Zohner, C.M., Gessler, A. & Vitasse, Y. (2021) Chilled to be forced: the best dose to wake up buds from winter dormancy. *New Phytologist* **230**, 1366–1377.

Caffarra, A., Donnelly, A. & Chuine, I. (2011) Modelling the timing of *Betula pubescens* budburst. II. Integrating complex effects of photoperiod into process-based models. *Climate Research* **46**, 159–170.

Cai, F., Jin, X., Tian, Y., Huang, Z., Wang, X., Zhang, Y., Sun, Y. & Shao, C. (2024) Molecular regulation of bud dormancy in perennial plants. *Plant Growth Regulation* **102**, 1–11.

Campbell, R.K. & Sugano, A.I. (1979) Genecology of bud-burst phenology in Douglas-fir - response to flushing temperature and chilling. *Botanical Gazette* **140**, 223–231.

Caspersen, L., Schiffers, K., Picornell, A., Egea, J.A., Delgado, A., El Yaacoubi, A., Benmoussa, H., Rodrigo, J., Fadón, E., Mimoun, M.B. *et al.* (2025) Contrasting responses to climate change–predicting bloom of major temperate fruit tree species in the mediterranean region and central europe. *Agricultural and Forest Meteorology* **375**, 110859.

Charrier, G., Lacointe, A. & Améglio, T. (2018) Dynamic modeling of carbon metabolism during the dormant period accurately predicts the changes in frost hardiness in walnut trees juglans regia l. *Frontiers in Plant Science* **9**, 1746.

Charrier, G., Ngao, J., Saudreau, M. & Ameglio, T. (2015) Effects of environmental factors and management practices on microclimate, winter physiology, and frost resistance in trees. *Frontiers in Plant Science* **6**.

Chuine, I., Bonhomme, M., Legave, J.M., de Cortazar-Atauri, I.G., Charrier, G., Lacointe, A. & Ameglio, T. (2016) Can phenological models predict tree phenology accurately in the future? the unrevealed hurdle of endodormancy break. *Global Change Biology* **22**, 3444–3460.

Chuine, I., Garcia de Cortazar Atauri, I., Hanninen, H. & Kramer, K. (2013) *Plant development models*, pp. 275–293. Kluwer, Dordrecht, the Netherlands.

Chuine, I. & Regniere, J. (2017) Process-based models of phenology for plants and animals. *Annual Review of Ecology, Evolution, and Systematics* **48**, 159–182.

Cleland, E.E., Chuine, I., Menzel, A., Mooney, H.A. & Schwartz, M.D. (2007) Shifting plant phenology in response to global change. *Trends in Ecology & Evolution* **22**, 357–365.

Considine, M.J. & Considine, J.A. (2016) On the language and physiology of dormancy and quiescence in plants. *Journal of Experimental Botany* **67**, 3189–3203.

Dennis, F. (2003) Problems in standardizing methods for evaluating the chilling requirements for the breaking of dormancy in buds of woody plants. *HortScience* **38**, 347–350.

Dörffling, K. (2015) The discovery of abscisic acid: a retrospect. *Journal of Plant Growth Regulation* **34**, 795–808.

Erez, A. & Fishman, S. (1997) The dynamic model for chilling evaluation in peach buds. *IV International Peach Symposium 465*, pp. 507–510.

Erez, A., Fishman, S., Gat, Z. & Couvillon, G. (1987) Evaluation of winter climate for breaking bud rest using the dynamic model. *International Workshop on Apple Culture in the Tropics and Subtropics 232*, pp. 76–89.

Erez, A. & Lavee, S. (1971) Effect of climatic conditions on dormancy development of peach buds. i. temperature. *Journal of the American Society for Horticultural Science* .

Erez, A., Lavee, S. & Samish, R.M. (1971) Improved methods for breaking rest in the peach and other deciduous fruit species. .

Ettinger, A.K., Chamberlain, C.J., Morales-Castilla, I., Buonaiuto, D.M., Flynn, D.F.B., Savas, T., Samaha, J.A. & Wolkovich, E.M. (2020) Winter temperatures predominate in spring phenological responses to warming. *Nature Climate Change* **10**, 1137–U119.

Faust, M., Liu, D., Millard, M.M. & Stutte, G. (1991) Bound versus free water in dormant apple buds—a theory for endodormancy. *HortScience* **26**, 887–890.

Fernandez, E., Schiffers, K., Urbach, C. & Luedeling, E. (2022) Unusually warm winter seasons may compromise the performance of current phenology models–predicting bloom dates in young apple trees with phenoflex. *Agricultural and forest meteorology* **322**, 109020.

Fishman, S., Erez, A. & Couvillon, G. (1987) The temperature dependence of dormancy breaking in plants: mathematical analysis of a two-step model involving a cooperative transition. *Journal of Theoretical Biology* **124**, 473–483.

Flynn, D.F.B. & Wolkovich, E.M. (2018) Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytologist* **29**, 1135–1492.

Fouché, M., Bonnet, H., Bonnet, D.M. & Wenden, B. (2023) Transport capacity is uncoupled with endodormancy breaking in sweet cherry buds: physiological and molecular insights. *Frontiers in Plant Science* **14**, 1240642.

Fu, Y.S.H., Zhao, H.F., Piao, S.L., Peaucelle, M., Peng, S.S., Zhou, G.Y., Ciais, P., Huang, M.T., Menzel, A., Uelas, J.P., Song, Y., Vitasse, Y., Zeng, Z.Z. & Janssens, I.A. (2015) Declining global warming effects on the phenology of spring leaf unfolding. *Nature* **526**, 104–107.

Gao, X., Richardson, A.D., Friedl, M.A., Moon, M. & Gray, J.M. (2024) Thermal forcing versus chilling? misspecification of temperature controls in spring phenology models. *Global Ecology and Biogeography* **33**, e13932.

Green, J.K. & Keenan, T.F. (2022) The limits of forest carbon sequestration. *Science* **376**, 692–693.

Gusewell, S., Furrer, R., Gehrig, R. & Pietragalla, B. (2017) Changes in temperature sensitivity of spring phenology with recent climate warming in switzerland are related to shifts of the preseason. *Global Change Biology* **23**, 5189–5202.

Guy, R.D. (2014) The early bud gets to warm. *New Phytologist* **202**, 7–9.

Hänninen, H. (1990) *Modelling bud dormancy release in trees from cool and temperate regions.* 213.

Hänninen, H., Kramer, K., Tanino, K., Zhang, R., Wu, J. & Fu, Y.H. (2019) Experiments are necessary in process-based tree phenology modelling. *Trends in Plant Science* **24**, 199–209.

Hufkens, K., Basler, D., Milliman, T., Melaas, E.K. & Richardson, A.D. (2018) An integrated phenology modelling framework in r. *Methods in Ecology and Evolution* **9**, 1276–1285.

Kaduk, J.D. & Los, S.O. (2011) Predicting the time of green up in temperate and boreal biomes. *Climatic Change* **107**, 277–304.

Kalcsits, L., Kendall, E., Silim, S. & Tanino, K. (2009) Magnetic resonance microimaging indicates water diffusion correlates with dormancy induction in cultured hybrid poplar (populus spp.) buds. *Tree Physiology* **29**, 1269–1277.

Keenan, T.F., Gray, J., Friedl, M.A., Toomey, M., Bohrer, G., Hollinger, D.Y., Munger, J.W., O'Keefe, J., Schmid, H.P., Wing, I.S. *et al.* (2014) Net carbon uptake has increased through warming-induced changes in temperate forest phenology. *Nature Climate Change* **4**, 598–604.

Khodadadzadeh, M., Kalverla, P. & Zurita-Milla, R. (2024) Harmonizing machine learning based phenological modeling: A unified workflow for comparative analyses. *IGARSS 2024 - 2024 IEEE International Geoscience and Remote Sensing Symposium*, pp. 5333–5336.

Kim, D.H. (2020) Current understanding of flowering pathways in plants: focusing on the vernalization pathway in arabidopsis and several vegetable crop plants. *Horticulture, Environment, and Biotechnology* **61**, 209–227.

Kim, D.H., Doyle, M.R., Sung, S. & Amasino, R.M. (2009) Vernalization: winter and the timing of flowering in plants. *Annual Review of Cell and Developmental* **25**, 277–299.

Kovaleski, A. (2022) Woody species do not differ in dormancy progression: differences in time to budbreak due to forcing and cold hardiness. *Proceedings of the National Academy of Sciences* **119**, e2112250119.

Kramer, K. (1994) Selelecting a model to predict the onset of growth of *Fagus sylvatica*. *Journal of Applied Ecology* **31**, 172–181.

Lamb, R.C. (1948) Effect of temperatures above and below freezing on the breaking of rest in the Latham raspberry. *Proceedings of the American Society for Horticultural Science* **51**, 313–315.

Lee, Z., Lim, J.A., Harikrishna, J.A., Islam, T., Abd Rahim, M.H. & Yaacob, J.S. (2024) Regulation of plant responses to temperature stress: a key factor in food security and for mitigating effects of climate change. *International Journal of Plant Production* **18**, 141–159.

Leida, C., Conejero, A., Arbona, V., Gómez-Cadenas, A., Llácer, G., Badenes, M.L. & Ríos, G. (2012) Chilling-dependent release of seed and bud dormancy in peach associates to common changes in gene expression. *PLoS One* **7**, e35777.

Leinonen, I. (1996) Dependence of dormancy release on temperature in different origins of pinus sylvestris and betula pendula seedlings. *Scandinavian Journal of Forest Research* **11**, 122–128.

Leisner, C.P. (2020) Climate change impacts on food security-focus on perennial cropping systems and nutritional value. *Plant Science* **293**, 110412.

Luedeling, E. (2012) Climate change impacts on winter chill for temperate fruit and nut production: a review. *Scientia horticulturae* **144**, 218–229.

Luedeling, E., Blanke, M. & Gebauer, J. (2015) *Chilling Challenges in a Warming World*, vol. 1099 of *Acta Horticulturae*, pp. 901–907.

Luedeling, E., Girvetz, E.H., Semenov, M.A. & Brown, P.H. (2011) Climate change affects winter chill for temperate fruit and nut trees. *Plos One* **6**.

Luedeling, E., Schiffers, K., Fohrmann, T. & Urbach, C. (2021) Phenoflex-an integrated model to predict spring phenology in temperate fruit trees. *Agricultural and Forest Meteorology* **307**, 108491.

Luedeling, E., Zhang, M.H. & Girvetz, E.H. (2009a) Climatic changes lead to declining winter chill for fruit and nut trees in California during 1950-2099. *Plos One* **4**.

Luedeling, E., Zhang, M.H., McGranahan, G. & Leslie, C. (2009b) Validation of winter chill models using historic records of walnut phenology. *Agricultural and Forest Meteorology* **149**, 1854–1864.

Olsson, C. & Jönsson, A.M. (2014) Process-based models not always better than empirical models for simulating budburst of norway spruce and birch in europe. *Global Change Biology* **20**, 3492–3507.

Overcash, J. & Campbell, J. (1955) The effects of intermittent warm and cold periods on breaking the rest period of peach leaf buds. .

Pan, W., Li, J., Du, Y., Zhao, Y., Xin, Y., Wang, S., Liu, C., Lin, Z., Fang, S., Yang, Y. *et al.* (2023) Epigenetic silencing of callose synthase by VIL1 promotes bud-growth transition in lily bulbs. *Nature Plants* **9**, 1451–1467.

Pandey, S.K., Moraes, T.S., Nair, A., Aryal, B., Azeez, A., Miskolczi, P., Maucort, G., Cordelières, F.P., Brocard, L., Davis, G.V. *et al.* (2026) Variable temperature processing by plasmodesmata regulates robust bud dormancy release. *Nature Communications* **17**, 348.

Penfield, S. & Springthorpe, V. (2012) Understanding chilling responses in *Arabidopsis* seeds and their contribution to life history. *Philosophical Transactions of the Royal Society B-Biological Sciences* **367**, 291–297.

Piao, S.L., Liu, Z., Wang, T., Peng, S.S., Ciais, P., Huang, M.T., Ahlstrom, A., Burkhart, J.F., Chevallier, F., Janssens, I.A., Jeong, S.J., Lin, X., Mao, J.F., Miller, J., Mohammat, A., Myneni, R.B., Penuelas, J., Shi, X.Y., Stohl, A., Yao, Y.T., Zhu, Z.C. & Tans, P.P. (2017) Weakening temperature control on the interannual variations of spring carbon uptake across northern lands. *Nature Climate Change* **7**, 359–+.

Pope, K., Da Silva, D., Brown, P. & DeJong, T. (2014) A biologically based approach to modeling spring phenology in temperate deciduous trees. *Agricultural and Forest Meteorology* **198**, 15–23.

Pörtner, H.O., Roberts, D.C., Tignor, M., Poloczanska, E.S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A. & Rama, B. (2022) *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.

494 Richardson, E. (1974) A model for estimating the completion of rest for 'Redhaven' and 'Elberta'
495 peach trees. *HortScience* **9**, 331–332.

496 Rinne, P.L.H., Paul, L.K. & van der Schoot, C. (2018) Decoupling photo- and thermoperiod by
497 projected climate change perturbs bud development, dormancy establishment and vernalization
498 in the model tree populus. *BMC Plant Biology* **18**.

499 Rinne, P.L.H., Welling, A., Vahala, J., Ripel, L., Ruonala, R., Kangasjarvi, J. & van der Schoot,
500 C. (2011) Chilling of dormant buds hyperinduces FLOWERING LOCUS T and recruits GA-
501 Inducible 1,3-beta-Glucanases to reopen signal conduits and release dormancy in *Populus*. *Plant*
502 *Cell* **23**, 130–146.

503 Satake, A., Nagahama, A. & Sasaki, E. (2022) A cross-scale approach to unravel the molecular
504 basis of plant phenology in temperate and tropical climates. *New Phytologist* **233**, 2340–2353.

505 Sato, H. & Yamane, H. (2024) Histone modifications affecting plant dormancy and dormancy
506 release: common regulatory effects on hormone metabolism. *Journal of Experimental Botany*
507 **75**, 6142–6158.

508 Schwartz, M.D. (1994) Monitoring global change with phenology – the case of the spring green
509 wave. *International Journal of Biometeorology* **38**, 18–22.

510 Singh, R.K., Maurya, J.P., Azeez, A., Miskolczi, P., Tylewicz, S., Stojkovič, K., Delhomme, N.,
511 Busov, V. & Bhalerao, R.P. (2018) A genetic network mediating the control of bud break in
512 hybrid aspen. *Nature communications* **9**, 4173.

513 Spafford, L., Lever, J.J., Gessler, A., Zweifel, R., Pietragalla, B., Wegner, J.D., Garnot, V.S.F.,
514 Sigg, C. & Vitasse, Y. (2025) Stability and transferability of broadly trained phenology models
515 in a changing climate. *Agricultural and Forest Meteorology* **372**, 110685.

516 Sperling, O., Kamai, T., Tixier, A., Davidson, A., Jarvis-Shean, K., Raveh, E., DeJong, T.M. &
517 Zwieniecki, M.A. (2019) Predicting bloom dates by temperature mediated kinetics of carbohy-
518 drate metabolism in deciduous trees. *Agricultural and Forest Meteorology* **276**, 107643.

519 Tuan, P.A., Bai, S., Saito, T., Ito, A. & Moriguchi, T. (2017) Dormancy-associated mads-box (dam)
520 and the abscisic acid pathway regulate pear endodormancy through a feedback mechanism. *Plant*
521 *and Cell Physiology* **58**, 1378–1390.

Tylewicz, S., Petterle, A., Marttila, S., Miskolczi, P., Azeez, A., Singh, R.K., Immanen, J., Mähler, N., Hvidsten, T.R., Eklund, D.M. *et al.* (2018) Photoperiodic control of seasonal growth is mediated by aba acting on cell-cell communication. *Science* **360**, 212–215.

van der Schoot, C., Paul, L.K. & Rinne, P.L.H. (2014) The embryonic shoot: a lifeline through winter. *Journal of Experimental Botany* **65**, 1699–1712.

Vitasse, Y. & Basler, D. (2014) Is the use of cuttings a good proxy to explore phenological responses of temperate forests in warming and photoperiod experiments? *Tree physiology* **34**, 174–183.

Vitasse, Y., Hoch, G., Randin, C.F., Lenz, A., Kollas, C., Scheepens, J.F. & Koerner, C. (2013) Elevational adaptation and plasticity in seedling phenology of temperate deciduous tree species. *Oecologia* **171**, 663–678.

Walde, M.G., Wenden, B., Chuine, I., Gessler, A., Saurer, M. & Vitasse, Y. (2024) Stable water isotopes reveal the onset of bud dormancy in temperate trees, whereas water content is a better proxy for dormancy release. *Tree Physiology* **44**, tpae028.

Walde, M.G., Wu, Z., Fox, T., Baumgarten, F., Fu, Y.H., Wang, S. & Vitasse, Y. (2022) Higher spring phenological sensitivity to forcing temperatures of asian compared to european tree species under low and high pre-chilling conditions. *Frontiers in Forests and Global Change* **5**, 1063127.

Weinberger, J.H. (1950) Chilling requirements of peach varieties. *Proceedings of the American Society for Horticultural Science* **56**, 122–128.

Wilczek, A.M., Roe, J.L., Knapp, M.C., Cooper, M.D., Lopez-Gallego, C., Martin, L.J., Muir, C.D., Sim, S., Walker, A., Anderson, J., Egan, J.F., Moyers, B.T., Petipas, R., Giakountis, A., Charbit, E., Coupland, G., Welch, S.M. & Schmitt, J. (2009) Effects of genetic perturbation on seasonal life history plasticity. *Science* **323**, 930–934.

Wolkovich, E.M., Auerbach, J., Chamberlain, C.J., Buonaiuto, D.M., Ettinger, A.K., Morales-Castilla, I. & Gelman, A. (2021) A simple explanation for declining temperature sensitivity with warming. *Global Change Biology* **27**, 4947–4949.

Yu, S., Wenden, B., Ferguson, L. & Tian, L. (2024) Building climate resilient deciduous tree crops by deciphering winter dormancy.

549 Zhang, Y.z., Chen, X., Ning, W., Ling, L. *et al.* (2023) PpMAPK6 regulates peach bud endodor-
550 mancy release through interactions with PpDAM6. *Journal of Integrative Agriculture* **22**, 139–
551 148.

552 Zhao, Y., Antoniou-Kourounioti, R.L., Calder, G., Dean, C. & Howard, M. (2020) Temperature-
553 dependent growth contributes to long-term cold sensing. *Nature* **583**, 825–829.

554 Zhu, P., Lister, C. & Dean, C. (2021) Cold-induced arabidopsis frigida nuclear condensates for flc
555 repression. *Nature* **599**, 657–661.

2 Figures




Figure 1: The term chilling generally refers to a process where dormant embryonic tissues exposed to cool temperatures accelerate a plant event that later occurs. While chilling cannot be directly observed currently, experiments for over a hundred years have tried to identify how much chilling plants need by finding the conditions that lead to rapid and full budburst (see right side where chilling is shown as the modeled unit of 'chill days'). Many studies have measured this via experiments that take dormant tissue (e.g., cut ends of tree branches taken from the field during the winter), which they then expose to sequential cool temperature treatments (generally called 'chilling' treatments or conditions) followed by warmer temperature treatments (called 'forcing' treatments or conditions), and observe budburst (left). Required chilling is then defined from these experiments by when additional cool temperatures do not yield faster and more complete budburst (right).




Figure 2: **Current models of chilling cannot uniquely estimate the most important aspects of chilling.** We illustrate this using a simple chilling model with a symmetrical thermal response curve, assuming the end of the chilling phase (often called endodormancy break) is observed (dashed vertical lines in right panes). In both cases, multiple parameter sets reproduce the same observation equally well. **(A)** Estimating the thermal response curve parameters (min. and max. temperatures) and the chilling threshold jointly (with a fixed start date) leads to many potential solutions. For example, a wider temperature range requires a higher threshold to reach the observed date (yellow), while a narrower range requires a lower threshold (blue). **(B)** The same issue happens when the response curve is fixed but the start date and threshold are estimated jointly. An earlier start date with a higher threshold (orange) leads to the same observed date as a later start date with a lower threshold (blue). These two simple examples illustrate that the parameter space contains multiple indistinguishable solutions, even when the end of the chilling phase is known (which is rare, more often models are built with only data on the end of the next phase—observed via budbreak). This issue would be even worse if the thermal response curve, the threshold, and the start date were all estimated jointly—and when the end of the phase is not directly observed. This reality is present in every model of chilling, but rarely discussed and often not even mentioned.




Figure 3: There are a lot of chilling models ... .

??




Figure 4: Aligning phenological models and molecular findings. Current phenological models of budbreak (top panel) assume two underlying states that lead to the observed process of budbreak: (1) endodormancy during which plants accumulate sufficient chilling to break endodormancy and transition (shown in blue) to (2) ecodormancy, a period during which plants accumulate sufficient heat or 'forcing' (warm temperatures, shown in orange) to break bud (flower or leaf), phases which may be sequential (darker shading) or occur in parallel (dark and light shading) yielding many variants of this model (second panel, and see Figure **??**). Cellular and molecular studies have found a suite of gene expression patterns than may underlie this transition, including shifts in callose (third panel), phytohormones (fourth panel) and those related to flowering time and master regulators (bottom). We show the glucanase family as dotted, given how many there are and that different glucanases are upregulated depending on the dormancy phase (Rinne *et al.*, 2011). We provide information on the species and citations underlying this figure in the Supplement (see 'Additional details on Figure 4')




Figure 5: Many experiments studying chilling alter 'chilling' and 'forcing' by varying the duration that cool temperatures are applied ('chilling') followed by different warm temperatures ('forcing'). Fitting simple linear models to these data (e.g., ANOVA models, with main effects of 'chilling' and 'forcing' treatments and their interaction—'chilling' × 'forcing'—to find a sub-additive effect of the two) usually does not match the current biological understanding of chilling. Fitting models that match the proposed accumulation process is possible, however, as we show here using data from Walde *et al.* (2022) for *Quercus robur*, and considering two alternative biologies: (A) cool ('chilling') and warm treatments (x axis) interact, such that longer cool temperatures mean more warm temperatures are required for leafout, versus (B) longer cool temperatures change the start date of the accumulation of warm ('forcing') temperatures (also known as endodormancy break).