

# The illusion of declining temperature sensitivity with warming OR A simple explanation for declining temperature sensitivity with warming

E. M. Wolkovich<sup>1,a</sup>, C. J. Chamberlain<sup>2</sup>, D. M. Buonaiuto<sup>2</sup>,  
A. K. Ettinger<sup>3</sup>, I. Morales-Castilla<sup>4</sup>, A. Gelman<sup>5</sup> & J. Auerbach<sup>5</sup>

March 31, 2020

\*\*Authorship order is not set.

<sup>1</sup>Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, Vancouver, British Columbia, Canada

<sup>2</sup>Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA

<sup>3</sup>TNC, USA

<sup>4</sup>Department of Life Sciences, University of Alcalà CTRA N-II, KM., 33,600, 28802, Alcalà de Henares, Spain

<sup>5</sup>Department of Statistics, Columbia University, New York, NY 10027, USA

<sup>a</sup>Corresponding author.

## Abstract

The concept of temperature sensitivity is fundamental across scientific disciplines, especially in biology, where temperature determines the rate of many plant, animal and ecosystem processes. Recently, a growing body of literature has found declining temperature sensitivities with global warming (Fu et al., 2015; Güsewell et al., 2017; Dai et al., 2019). Such declines are predicted if warming causes fundamental shifts in underlying biological processes, but researchers have not conclusively documented changes in the underlying biology. Here we present a simple explanation for observed declining sensitivities: the use of linear regression to estimate non-linear temperature responses. Simple corrections for the non-linearity of temperature response in simulated data, and long-term phenological data from Europe, remove the apparent decline. By accelerating biological time, climate change makes methods that may suffice in stationary systems problematic for inferring mechanism from measurements today.

# 1 Main text

Climate change has reshaped biological processes around the globe, with shifts in the timing of major life history events (phenology), in carbon dynamics and other ecosystem processes (IPCC, 2014). With increasing warming, a growing body of literature has documented changes in temperature sensitivity—the magnitude of a response scaled per °C—including declining responses to temperature over time (Fu et al., 2015; Güsewell et al., 2017; Piao et al., 2017; Dai et al., 2019), and more uniform sensitivities across elevation (Vitasse et al., 2018).

Researchers often suggest these changes in temperature sensitivity are driven by shifts in underlying biological processes. For example, fundamental science suggests that warm spring temperatures are the main controller of many temperate phenological events (e.g., leafout, insect emergence), but winter temperatures and photoperiod can also determine the timing of events. Thus, an increased role of winter temperatures or photoperiod could explain observed declines in the temperature sensitivity of plant phenology with warming. Yet, providing strong evidence of this mechanistic link has proven difficult (e.g., Fu et al., 2015; Gauzere et al., 2019), in part because of uncertainty in how these other factors control phenological events (Chuine et al., 2016), and because these factors are often correlated in nature (e.g., Fu et al., 2015).

Alongside efforts to identify the biology that may underlie shifting temperature sensitivities, a small but increasing number of studies have focused on potential statistical issues in common metrics of temperature sensitivity. These studies have highlighted important complexities that can influence estimated sensitivities (Clark et al., 2014; Güsewell et al., 2017; Keenan et al., 2020). They have generally not, however, explained observed shifts. And all have examined sensitivities through methods based on assumptions of linearity, generally relying on some form of linear regression to compute a change in a quantity—days to leafout or carbon sequestered, for example—per °C—thus ignoring that many biological responses to temperature are non-linear.

Many observed biological events are the result of continuous processes that depend on temperature, which are discretized into temporal units for measurement. Leafout, for example, is generally observable only after a certain thermal sum is reached, and plants will reach this threshold more quickly—in calendar time—when average daily temperatures are warmer. Biologically, however, the plants require the exact same temperature sum and have not shifted their sensitivity to temperature. Indeed any process observed or measured as the time until reaching a threshold is inversely proportional to the speed at which that threshold is approached. Thus, at very low temperatures plants would never leaf out and at higher temperatures they could leaf out in only a matter of days—yet sensitivities estimated from simple linear regression at these higher temperatures would appear much lower than those observed at lower temperatures (given the low variance of the response variable). Warming acts to step on the biological accelerator, and makes the use of classic calendar time precarious.

Simple simulations of biological events observed after a certain thermal sum show that sensitivities estimated from simple linear regression will always appear to decline with warming (Fig. 1, ??, code link). Examining the same responses using proportional change or logged variables re-

moves the apparent decline, and yields a constant sensitivity of -1 (the expected slope given that these sensitivities effectively include temperature as both the predictor and response variable, see ‘Model of leafout timing’ in SI and Nee et al., 2005). Using alternative simulations where warming increases the required thermal sum for a biological event—a common hypothesis for declining sensitivities in spring phenological events—yields declining sensitivities that remain after adjusting response variables using proportions or logs (Fig. ??).

Comparing these simulation results with long-term leafout data from Europe, we find little evidence for declining sensitivities with warming (Figs. 1, ??, ??). An apparent decline in sensitivity for silver birch (*Betula pendula*) from -4.3 days/°C to -3.6 days/°C from 1950-1960 compared to 2000-2010 disappears using a log-log regression (-0.17 versus -0.22). We see similar corrections using 20-year windows, and a potential increase in sensitivity for European beech (*Fagus sylvatica*, see Tables ??-??). Across both species, there are large declines in the variance of leafout dates—(declines of roughly 50%, see Tables ??-??), as expected if warming accelerates towards a thermal threshold that triggers leafout (and in contrast to predictions from shifting biology, see Ford et al., 2016).

Inferring biological processes from statistical artifacts is not a new problem (e.g., Nee et al., 2005), but climate change provides a new challenge in discerning mechanism from measurements because it accelerates biological time over years and reshapes the spatial landscape of temperature. Before anthropogenic climate change, the use of sensitivities calculated from linear models may have been less prone to yielding notable temporal patterns. With warming declining sensitivities should be the null model for analyses using simple linear regressions, and highlights how the nonstationarity of climate change upends methods and approaches that may work in stationary systems (Milly et al., 2008; Wolkovich et al., 2014). Attempts to use sensitivities to identify shifting biological process across space has always required caution (e.g., Tansey et al., 2017), but climate change adds further complexity.

Research inferring biological processes from differing temperature sensitivities across space and time must look beyond shifting sensitivities with warming as strong evidence. Other fields, focused on temperature sensitivity generally use approaches that avoid the issue we have outlined here (e.g., Yuste et al., 2004). Researchers have called for more use of process-based models (Keenan et al., 2020), which is also beneficial. But many fields, still lack the underlying mechanistic understanding to robustly develop and fit process-based models, with many parameters and aspects of the model specification unknown (Chuine et al., 2016). Thus we expect more exploratory methods, such as regression, will continue to inform science, but findings from such methods must be interrogated—confronted with multiple diverse methods of calculating similar metrics, tested for logical outcomes, and compared against null models based on biological time.

*Acknowledgements:* Thanks to TMG, TJD and others.

## References

- Chuine, I., M. Bonhomme, J.-M. Legave, I. García de Cortázar-Atauri, G. Charrier, A. Lacointe, and T. Améglio. 2016. Can phenological models predict tree phenology accurately in the future? The unrevealed hurdle of endodormancy break. *Global Change Biology* 22:3444–3460.
- Clark, J. S., J. Melillo, J. Mohan, and C. Salk. 2014. The seasonal timing of warming that controls onset of the growing season. *Global Change Biology* 20:1136–1145.
- Dai, W. J., H. Y. Jin, Y. H. Zhang, T. Liu, and Z. Q. Zhou. 2019. Detecting temporal changes in the temperature sensitivity of spring phenology with global warming: Application of machine learning in phenological model. *Agricultural and Forest Meteorology* 279.
- Ford, K. R., C. A. Harrington, S. Bansal, J. Gould, Peter, and J. B. St. Clair. 2016. Will changes in phenology track climate change? A study of growth initiation timing in coast Douglas–fir. *Global Change Biology* 22:3712–3723.
- Fu, Y. S. H., H. F. Zhao, S. L. Piao, M. Peaucelle, S. S. Peng, G. Y. Zhou, P. Ciais, M. T. Huang, A. Menzel, J. P. Uelas, Y. Song, Y. Vitasse, Z. Z. Zeng, and I. A. Janssens. 2015. Declining global warming effects on the phenology of spring leaf unfolding. *Nature* 526:104–107.
- Gauzere, J., C. Lucas, O. Ronce, H. Davi, and I. Chuine. 2019. Sensitivity analysis of tree phenology models reveals increasing sensitivity of their predictions to winter chilling temperature and photoperiod with warming climate. *Ecological Modelling* 441:108805.
- Güsewell, S., R. Furrer, R. Gehrig, and B. Pietragalla. 2017. Changes in temperature sensitivity of spring phenology with recent climate warming in switzerland are related to shifts of the preseason. *Global Change Biology* 23:5189–5202.
- IPCC. 2014. *Climate Change 2014: Impacts, Adaptation, and Vulnerability*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Keenan, T. F., A. D. Richardson, and K. Hufkens. 2020. On quantifying the apparent temperature sensitivity of plant phenology. *New Phytologist* 225:1033–1040.
- Milly, P. C. D., J. Betancourt, M. Falkenmark, R. M. Hirsch, Z. W. Kundzewicz, D. P. Lettenmaier, and R. J. Stouffer. 2008. Climate change - stationarity is dead: Whither water management? *Science* 319:573–574.
- Nee, S., N. Colegrave, S. A. West, and A. Grafen. 2005. The illusion of invariant quantities in life histories. *Science* 309:1236–1239.
- Piao, S., Z. Liu, T. Wang, S. Peng, P. Ciais, M. Huang, A. Ahlstrom, J. F. Burkhart, F. Chevallier, I. A. Janssens, et al. 2017. Weakening temperature control on the interannual variations of spring carbon uptake across northern lands. *Nature climate change* 7:359.
- Tansey, C. J., J. D. Hadfield, and A. B. Phillimore. 2017. Estimating the ability of plants to plastically track temperature-mediated shifts in the spring phenological optimum. *Global Change Biology* 23:3321–3334.

- Vitasse, Y., C. Signarbieux, and Y. H. Fu. 2018. Global warming leads to more uniform spring phenology across elevations. *Proceedings of the National Academy of Sciences* 115:1004–1008.
- Wolkovich, E. M., B. I. Cook, K. K. McLauchlan, and T. J. Davies. 2014. Temporal ecology in the anthropocene. *Ecology Letters* 17:1365–1379. Cook, Benjamin/H-2265-2012.
- Yuste, J., I. A. Janssens, A. Carrara, and R. Ceulemans. 2004. Annual  $Q_{10}$  of soil respiration reflects plant phenological patterns as well as temperature sensitivity. *Global Change Biology* 10:161–169.

## **2 Tasks, milestones etc.**

- Turn this into Sweave and add in-text refs for the sensitivities.
- Dan, please help check refs (did I use the correct Dai one? They seem very different!)

## Figures




Figure 1: **Shifts in temperature sensitivities with warming occur when using linear regression for a non-linear process.** We found estimated sensitivities declined with warming from simulations with no underlying change in the biological process when sensitivities were estimated with simple linear regression (‘Sims: untransformed,’ estimated across 45 sites with a base temperature of  $\mathcal{N}(6, 4)$ ). This decline disappears when performing the regression on logged predictor and response variables (‘Sims: logged’). Such issues may underlie declining sensitivities calculated from long-term phenological data, such as from PEP725 observations. ‘PEP: untransformed’ shows a declining sensitivity with warming, while ‘PEP725: logged’ shows a similar sensitivity with warming. We used sites with leafout data for *Betula pendula* from PEP725 from 1950-1960 versus 2000-2010 (45 sites). We show means  $\pm$  standard deviations of regressions across sites for all points.