2424 Main Mall Vancouver, BC Canada V6T 1Z4

Ph: 604.827.5246



Dear Dr. Sudgen:

Please consider our manuscript, entitled "A simple explanation for declining temperature sensitivity with warming," for publication as a Report in *Science*.

Climate change has shifted many biological events (1), and recent observations that organisms' sensitivity to temperature is declining has raised concerns that fundamental biological processes are now also changing. Studies suggest warming has altered the main drivers of leafout in temperate plants (2; 3; 4; 5) and altered carbon uptake in the tundra possibly due to increased light limitation or shifts in photosynthesis (6; 7). Such possible changes are alarming, but they are rarely reported with strong evidence beyond declines in temperature sensitivity.

Here we provide a simpler alternative explanation for declining temperature sensitivity with warming: the use of linear models to model non-linear temperature responses. We show theoretically, then through simulations and empirical data (using the same dataset used in 2), that observations of declining sensitivities are the default given current methods. This is because many biological events are the outcome of threshold processes and thus will occur more quickly with warming. This in turn will lead to statistically lower estimated responses per degree C when using linear models, without any changes in the underlying biology. Effectively, warming acts to step on the biological accelerator, making the use of classic calendar time combined with linear models problematic.

We believe this issue is currently rampant throughout the phenological (e.g., 2; 4; 5; 8) and related literature (e.g., 6) and may apply more broadly to many fields using linear methods and fixed time periods with warming. Understanding this problem only came through close collaboration between biologists (Wolkovich group) and statisticians (J. Auerbach and A. Gelman).

We have suggested three potential reviewers through the online submission site (Theresa Crimmins, Alison Donnelly, Mark Schwartz). Prior to submission Jonathan Davies, David Lipson and Christy Rollinson reviewed the manuscript. This manuscript is not under consideration elsewhere, nor has it been previously submitted. Empirical data are already publicly available through the PEP 725 portal, and we provide a link in the main text to our simulation code. All authors approved of this version for submission.

Sincerely,

Elizabeth M Wolkovich

Salimuelle-

Associate Professor of Forest & Conservation Sciences

## References

- [1] IPCC. Climate Change 2014: Impacts, Adaptation, and Vulnerability (Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014).
- [2] Fu, Y. S. H. *et al.* Declining global warming effects on the phenology of spring leaf unfolding. *Nature* **526**, 104–107 (2015).
- [3] Güsewell, S., Furrer, R., Gehrig, R. & Pietragalla, B. Changes in temperature sensitivity of spring phenology with recent climate warming in switzerland are related to shifts of the preseason. *Global Change Biology* **23**, 5189–5202 (2017).
- [4] Samplonius, J. M. et al. Phenological sensitivity to climate change is higher in resident than in migrant bird populations among european cavity breeders. Global Change Biology 24, 3780–3790 (2018).
- [5] Vitasse, Y., Signarbieux, C. & Fu, Y. H. Global warming leads to more uniform spring phenology across elevations. *Proceedings of the National Academy of Sciences* **115**, 1004–1008 (2018).
- [6] Piao, S. et al. Weakening temperature control on the interannual variations of spring carbon uptake across northern lands. Nature Climate Change 7, 359 (2017).
- [7] Zhu, P. et al. Recent warming has resulted in smaller gains in net carbon uptake in northern high latitudes. *Journal of Climate* **32**, 5849–5863 (2019).
- [8] Meng, L. et al. Urban warming advances spring phenology but reduces the response of phenology to temperature in the conterminous united states. *Proceedings of the National Academy of Sciences* 117, 4228 (2020).