

Scattered spring: How climate change disrupts the synchrony of biological events

Jonathan Auerbach*
Department of Statistics
George Mason University
Fairfax, VA USA

Andrew Gelman[†]
Department of Statistics
Department of Political Science
Columbia University
New York, NY USA

E. M. Wolkovich[‡]
Forest and Conservation Sciences
University of British Columbia
Vancouver, BC Canada

(Dated: June 18, 2026)

Many biological processes, including plant leafout and flowering, occur once cumulative temperatures reach a threshold, the ‘thermal-sum model’. In this way, temperature is thought to coordinate the timing of biological events. But growing evidence suggests that as climates warm, both the advancement of spring has slowed (declining sensitivity) and the variance in the timing of spring events has increased (declining synchrony)—raising questions about the resilience of temperature-based coordination to anthropogenic climate change. To answer these questions, researchers have modified the thermal-sum model by introducing additional factors and mechanisms, such as chilling and photoperiod. We show such complexity is not necessary to explain current trends of sensitivity and synchrony. Using experimental and real-world data, we find these trends are exactly as predicted by the thermal-sum model. In particular, the model predicts that as temperatures continue to increase and springtime events shift from the equinox toward the winter solstice, those events will become less synchronized and more variable, a phenomenon we refer to as a scattered spring. By 2100, our results predict much of the American South will experience a scattered spring under a high-warming scenario, with the average time between spring flowering events in many locations increasing by several weeks.

Introduction

Many biological events are triggered not by temperature on a single day, but by temperature accumulated over many days (Larcher, 1980). The long-standing and widespread ‘thermal-sum’ model states that the bud of a flowering plant, for example, first blooms in the spring once cumulative temperatures exceed a plant-specific threshold (Quetelet, 1849; Chuine and Regnier, 2017). This mechanism is thought to syn-

chronize a wide variety of seasonal events within an ecosystem—for instance, ensuring that pollinators emerge when flowers are in bloom (Schwartz, 2024). It also supports an array of forecasting tools in which scientists track cumulative temperatures or ‘growing degree days’ to predict harvest dates, manage pest emergence, and assess how changing temperature patterns reshape ecological communities (Schwartz, 2024).

Shifts in the timing of biological events have been among the clearest fingerprints of anthropogenic climate change across both natural and managed systems

*jauerba@gmu.edu

[†]ag389@columbia.edu

[‡]wolkovic@mail.ubc.ca

(Pörtner et al., 2022). But researchers have struggled to fully predict or explain these shifts in recent years. One puzzle is that while the timing of many biological events has moved earlier (i.e., advanced) as climates warm, the rate of advancement has slowed (Fu et al., 2015). We refer to this pattern in which events appear less sensitive to additional warming as ‘declining sensitivity.’

A second puzzle is that the variance in the timing of biological events has in some cases declined and in others increased (Vitasse et al., 2018; Stemkovski et al., 2023). Higher variance has also manifested as previously unknown or rare events becoming more common (Chuine et al., 2025). We refer to this pattern in which events appear less coordinated as ‘declining synchrony.’

Declining sensitivity and synchrony are almost always interpreted as evidence that the long-standing and widespread thermal-sum model is now wrong or incomplete. Researchers have suggested myriad complexities or alternatives (Fu et al., 2015; Wolkovich et al., 2021; Gao et al., 2024). Yet no work has examined whether these puzzles are explained by the thermal-sum model, or how sensitivity and synchrony will continue to evolve under the thermal-sum model as climates continue to warm.

Here we show that the thermal-sum model is an example of a stopped random walk, and we build a simple and transparent framework for the analysis of biological event times using statistical tools developed for stopped random walks. Applying the framework to observational and experimental data, we find that the thermal-sum model explains both declining sensitivity and synchrony. Thus modifications and alternative approaches often suggested (Fu et al., 2015; Vitasse et al., 2018) may be unnecessary and yield inaccurate predictions. In contrast to predictions from current models of biological events, our framework implies that as springtime events advance with anthropogenic warming, they enter a new temperature regime in which the typical time between events in the same ecosystem can increase from a few days to a month or more, a phenomenon we refer to as a ‘scattered spring.’

Results & Discussion

The thermal-sum model states that a biological event occurs when cumulative temperature reaches a threshold. This model is an example of a stopped random walk where the cumulative temperature is the position of the walk, and the biological event occurs when the

walk arrives at the threshold.

Consider the timing of a specific biological event, such as the bloom date of a bud. Let X_i denote the effective temperature experienced by that bud on day i . We model X_i under two temperature regimes (Figure 1).

The first regime is the ‘winter regime’ in which the average daily temperature is approximately constant at some level $\alpha > 0$. This regime is often observed in the winter months following the solstice, and we refer to α as the ‘winter temperature.’ The second regime is the ‘spring regime’ in which the average daily temperature rises at a constant rate $\beta > 0$. This regime is often observed in the spring months closer to the equinox, and we refer to β as ‘spring warming.’

In either regime, effective temperature is modeled as average temperature plus noise ϵ_i . That is, $X_i = \alpha + \beta i + \epsilon_i$, where $\beta = 0$ in the winter regime and $\beta > 0$ in the spring regime. We assume the ϵ_i are independent with mean 0 and variance σ^2 .

Noise in this formulation can represent the influence of various biological factors, depending on the scale of analysis and on our evolving understanding of what triggers springtime events. At the bud level, it represents idiosyncratic variation specific to that bud, which could arise from microclimate differences (Schwartz et al., 2014), bud color variation (Peaucelle et al., 2022), or internal biological differences that affect when the bud begins accumulating temperatures (van der Schoot et al., 2014; Pan et al., 2023).

This two regime model in which average temperatures switch from constant to increasing well-describes both historic springtime temperature data (Figure S1 in SI.3) and projected springtime temperature data (Figure S4 in SI.3). Here we assume that warming occurs entirely in one of the two regimes, although similar results hold when warming occurs under both regimes (since in such cases the spring regime dominates, see SI.1 for details.)

The bloom date $\nu(\tau)$ is the day n on which the cumulative temperature $\sum_{i=1}^n X_i$ first reaches a plant-specific threshold $\tau > 0$. This formulation is standard in growing degree day models and, more generally, the ‘forcing’ component of many process-based models of plant phenology (Schwartz, 2024). This formulation also allows us to use standard results for stopped random walks, which yield asymptotically normal approximations for the bloom date when the threshold is sufficiently large that accumulation occurs over many days.

In the winter regime, average temperatures are con-




Figure 1: We model the daily temperature (solid line) as trend (dashed line) plus noise. The timing of springtime events in temperate climates depends on whether temperature is accumulated during one of two regimes, 1. the winter regime (relatively flat average daily temperature near the winter solstice, which we refer to as the ‘winter temperature’ or α), or 2. the spring regime (relatively linear average daily temperatures near the spring equinox, the slope of which we refer to as ‘spring warming’ or β). We assume the effective temperature experienced by the bud is positive since biological limit prevents plants from accumulating very low temperatures, taken here to be 0°C (Larcher, 1980) although this is easily adjusted.

stant during accumulation ($\beta = 0$), and cumulative temperature grow at an approximately linear rate. The average bloom date advances with climate change in step with the winter temperature ($\mathbb{E}[\nu(\tau)] \approx \tau/\alpha$) as is often described in the literature (Vitasse et al., 2022). In the spring regime, average temperatures increase during accumulation ($\beta > 0$), and cumulative temperatures grow quadratically. In this regime, the average bloom date squared advances in step with spring warming ($\mathbb{E}[\nu(\tau)] \approx \sqrt{2\tau/\beta}$).

Within either regime, the relationship between bloom dates and temperature is non-linear as observed in recent papers (Fu et al., 2015; Wolkovich et al., 2021). Indeed, both regimes predict declining sensitivity in that the average bloom date advances at a diminishing rate when warming increases as measured by the winter temperature (α) or spring warming (β).

Importantly, the effects of warming within the two regimes make very different predictions regarding the synchrony of events—that is, the variance in the timing of events within the same year and location due to idiosyncratic differences described above such as microclimate. Both predict decreased variance with warming ($\text{Var}(\nu(\tau)) \approx \sigma^2 \tau / \alpha^3$ in the winter regime

and $\text{Var}(\nu(\tau)) \approx \sigma^2 / \sqrt{2\beta^3 \tau}$ in the spring regime). But the overall size of the variance depends on whether the threshold (τ) is in the numerator or the denominator.

When the threshold is in the denominator as in the spring regime, the variance is small, reflecting the fact that noise is quickly overwhelmed by the quadratic increase in cumulative temperatures that occurs during spring warming each year. This is how temperatures near the equinox synchronize plants (and buds within plants) with the same threshold—by making idiosyncratic differences such as microclimate negligible. In contrast, when the threshold is in the numerator as in the winter regime, the variance is large, reflecting the fact that noise remains relatively influential in that idiosyncratic differences can translate into larger differences in threshold-crossing times.

In this way, our results reveal the key mechanism that determines whether synchrony declines: the temperature regime under which accumulation occurs. When average temperatures increase day over day, as they often do near the equinox, springtime events tend to be tightly synchronized. But when warming shifts those events closer to the solstice, where average tempera-

tures are comparatively constant, event times become more variable. The shift in regimes is observable as a ‘scattered spring’ in which, as springtime events advance, they become less synchronized.

Our results also show that the two most important aspects of temperature are the levels of winter temperature (α) and spring warming (β). This simplification yields a simple and transparent framework for translating spatially varying climate change into spatially varying predictions of both the advancement of spring and its variability.

To demonstrate the framework, we analyze bloom dates from the lilac–honeysuckle phenology network, compiled by Rosemartin et al. (2015) from sites across the continental United States between 1956–2014. We limit our analysis to the purple common lilac (*Syringa vulgaris*) and the phenophase full bloom, and we update the dataset using USA National Phenology Network records from the same sites and species until 2025 (Switzer et al., 2025). For each record, we estimate the winter temperature (α) and spring warming (β). We then estimate the conditional mean and standard deviation functions of the bloom date using a generalized additive model (Hastie and Tibshirani, 1986; Wood, 2017).

We find that increasing either the winter temperature, spring warming, or both has shifted lilac flowering times exactly as predicted by the thermal-sum model (Figures 2). Warming in either regime (increasing α or β) leads to earlier mean bloom dates, which advance at a decreasing rate (declining sensitivity). Warming in the winter regime only (increasing α and holding β fixed) returns greater variance (higher standard deviation, declining synchrony), while warming in the spring regime only (increasing β while holding α fixed) returns lower variance (lower standard deviation, increasing synchrony). This is the ‘scattered spring’ predicted by the model: in the first case, plants are more likely to accumulate their thermal sum in the winter regime, which results in the larger variance.

In this way, the thermal-sum model explains a wide variety of trends in the mean and variance of biological event times, challenging a growing body of work claiming such trends either invalidate the model or justify additional complexity (e.g., Fu et al., 2015; Kovaleski, 2021). For example, researchers have suggested that climate change may influence photoperiod or chilling (cool temperatures required to break endodormancy), which in turn change the threshold of the thermal sum.


We show that these additional factors are unneces-

sary by applying our framework to rare experimental data in which photoperiod and chilling are held fixed but forcing temperatures vary. We use results from Charrier et al. (2011), who tested the effect of different constant temperatures on walnut (*Juglans regia* and *Juglans regia x nigra* hybrids) tree budburst after the plants had experienced sufficient chilling to break endodormancy (385 buds were observed on stems sampled from 15 trees, chilled at 4°C), and we fit the winter regime approximation to the thermal-sum model (since temperatures are held constant, this experiment replicates this regime) using weighted least squares. The data exhibit the exact nonlinear relationship in the mean and variance as predicted by the thermal-sum model (Figure 3), establishing the sufficiency of the model even under an extreme temperature range and controlled chilling and photoperiod conditions.


Our results show that the long-standing and widely-used thermal-sum model (Quetelet, 1849; Chuine and Regniere, 2017) explains observed shifts in the timing of biological events including declining sensitivity and synchrony. This finding has two important consequences. First, it challenges an extensive body of work that has attempted to revise the thermal-sum model by adding additional factors or mechanisms (Fu et al., 2015). These factors can influence the timing of biological events, but our work suggests they do not themselves explain current trends in sensitivity and synchrony. Further, they can produce inaccurate forecasts if the factors are modeled incorrectly.

To understand the role played by any additional factors, the basic relationship between event time and temperature must be modeled correctly. For example, the literature often assumes a linear relationship between temperature and event time with constant variance, which we have shown is inconsistent with the thermal-sum model. Our framework provides a simple and transparent way to encode the thermal-sum model, on top of which complexity may be added and better studied.

The second consequence is that the thermal-sum model predicts a ‘scattered spring’ or general desynchronization of springtime events as their timing moves closer to the winter solstice. We find that the synchrony of lilac bloom dates, a common indicator of spring, will decline in much of the Southern United States by the year 2100 under a high-warming scenario (SSP5-8.5): For example, were lilac planted across the South in 2050, the average time between blooms would be less than two weeks in 76% of locations. By 2100, however, the average time will be less than two weeks in only 18% of locations (Figure



(a) Bivariate plots showing mean (left) and log standard deviation (right) of lilac bloom dates as a function of winter temperature (α) and spring warming (β).



(b) Cross-section plots showing mean (left) and log standard deviation (right) of lilac bloom dates as a function of winter temperature (α) holding spring warming (β) constant (top) and as a function of spring warming (β) holding winter temperature (α) constant (bottom).

Figure 2: Mean and log standard deviation of lilac bloom dates ($n = 10,613$ from Rosemartin et al. 2015, updated to 2025 using USA National Phenology Network data) by winter temperature (α) and spring warming (β) as estimated by a generalized additive model. The estimated functions are shown using bivariate plots (a, top) and cross-section plots (b, bottom). The bivariate plots show the mean decreases as α and β increase (left), while the standard deviation can increase or decrease depending on the relative values of α and β (right). The cross-section plots show the mean decreases both as α increases holding β constant (top left) and as β increases holding α constant (top right), while the standard deviation can increase or decrease depending on whether α increases with β held constant (bottom left) or β increases with α held constant (bottom right).




Figure 3: Mean time until budburst as a function of the average daily temperature (α) in a controlled experiment conducted by Charrier et al. (2011) in which 385 buds were observed on stems sampled from 15 walnut trees (*Juglans regia* and *Juglans regia* $\times$ *nigra* hybrids) trees, chilled at 4°C, and then warmed in different temperatures environments (x-axis) with the same photoperiod conditions across temperatures until budburst. For each tree (panel), we show the observed dates (points), fitted mean (solid line), 95% confidence interval (inner grey region, approximately two standard errors), and 95% prediction interval (outer grey region, approximately two standard deviations). The figure shows that the mean and variance are nonlinear functions of temperature even when biological factors such as chilling and photoperiod are held fixed. Both decrease with increasing temperatures as predicted by the thermal-sum model in the winter regime.

4, note that the synchrony of species with lower threshold (τ) may decline sooner). In many locations, the average time will increase by several weeks.

Absent some compensating mechanism, the long-standing and widespread thermal-sum model predicts that the timing of biological events across the ecosystem—from crops to forests, and from plants to insects—will in many cases become more variable, with potential cascading consequences for industry and natural ecosystems.

Methods

Our observational data analysis considers all sites in the historical lilac–honeysuckle phenology network (Rosemartin et al., 2015) that recorded the phenophase full bloom and were within 10 miles of a Global Historical Climatology Network daily (GHCNd) site with sufficiently complete temperature records. We limit our analysis to common lilac (*Syringa vulgaris*), including all observations made from 1955 to 2025 as recorded by the USA National Phenology Network.

The data were retrieved using the R package `rnnpn`, resulting in 10,613 observations.

For each observation, we estimate the winter temperature (α) using the average daily temperature between January and February, and we estimate spring warming (β) using the slope of a linear regression model fit by regressing daily temperature on day index (i.e., $i = 1$ for January 1, $i = 2$ for January 2, etc.) over March and April (Figure S1, S2, and S3 in SI.3). Temperatures below 0 are set to 0, a common base temperature in growing degree day models.

We then fit a two-parameter Gaussian location-scale model using the `gaulss` function from the R package `mgcv` in which we define

$$\mu(\alpha, \beta) = \mathbb{E}[\nu(\tau) \mid \alpha, \beta]$$

$$\sigma(\alpha, \beta) = \text{SD}(\nu(\tau) \mid \alpha, \beta).$$

Both μ and σ are modeled with tensor-product splines with basis dimension $k = 10$ (Figure 2).

Finally, we calculate winter temperatures ($\tilde{\alpha}$) and




Figure 4: Predicted mean (top) and standard deviation (bottom, log scale) of lilac bloom dates in years 2050 and 2100 across the Southern United States under temperatures projected in the NEX-GDDP-CMIP6 high-warming scenario SSP5-8.5 (Thrasher et al., 2022). Lighter areas indicate earlier blooms (top) or more variable blooms (bottom) due to higher temperatures. The mean bloom date advances moderately between 2050 and 2100, while the standard deviation increases by several weeks in many locations.

spring warming ($\tilde{\beta}$) for the years 2025, 2050, 2075, and 2100 using the NEX-GDDP-CMIP6 temperature projections for two scenarios: SSP2-4.5 and SSP5-8.5 (Thrasher et al., 2022) (Figure S4 in SI.3). We also calculate the projected mean bloom date $\mu(\tilde{\alpha}, \tilde{\beta})$ and standard deviation $\sigma(\tilde{\alpha}, \tilde{\beta})$ (Figure 4 and Figures S5 and S6 in SI.3).

References

- Charrier, G., Bonhomme, M., Lacointe, A., and Améglio, T. (2011). Are budburst dates, dormancy and cold acclimation in walnut trees (*Juglans regia* L.) under mainly genotypic or environmental control? *International journal of biometeorology*, 55(6):763–774.
- Chuine, I., Garcia de Cortazar-Atauri, I., Jean, F., and Van Reeth, C. (2025). Living things are showing increasing anomalies in their seasonal activity, which could disrupt the dynamics of biodiversity and ecosystems. *Scientific Reports*, 15(1):32860.
- Chuine, I. and Regniere, J. (2017). Process-based models of phenology for plants and animals. *Annual Review of Ecology, Evolution, and Systematics*, 48:159–182.
- Fu, Y. H., Zhao, H., Piao, S., Peaucelle, M., Peng, S., Zhou, G., Ciais, P., Huang, M., Menzel, A., Peñuelas, J., et al. (2015). Declining global warming effects on the phenology of spring leaf unfolding. *Nature*, 526(7571):104–107.
- Gao, X., Richardson, A. D., Friedl, M. A., Moon, M., and Gray, J. M. (2024). Thermal forcing versus chilling? misspecification of temperature controls in spring phenology models. *Global Ecology and Biogeography*, 33(12):e13932.
- Guillaume, C., Isabelle, C., Marc, B., and Thierry, A. (2018). Assessing frost damages using dynamic models in walnut trees: exposure rather than vulnerability controls frost risks. *Plant, Cell & Environment*, 41(5):1008–1021.
- Gut, A. (2009). *Stopped random walks*. Springer.
- Hastie, T. and Tibshirani, R. (1986). Generalized additive models. *Statistical science*, 1(3):297–310.
- Kovaleski, A. (2021). Woody species do not differ in dormancy progression: differences in time to bud-break due to forcing and cold hardiness. *bioRxiv*.
- Lai, T. L. and Siegmund, D. (1977). A nonlinear renewal theory with applications to sequential analysis I. *The Annals of Statistics*, pages 946–954.
- Larcher, W. (1980). *Plant Physiological Ecology*. Springer-Verlag.
- Pan, W., Li, J., Du, Y., Zhao, Y., Xin, Y., Wang, S., Liu, C., Lin, Z., Fang, S., Yang, Y., et al. (2023). Epigenetic silencing of callose synthase by VIL1 promotes bud-growth transition in lily bulbs. *Nature Plants*, 9(9):1451–1467.
- Peaucelle, M., Penuelas, J., and Verbeeck, H. (2022). Accurate phenology analyses require bud traits and energy budgets. *Nature Plants*, 8(8):915–922.
- Pörtner, H. O., Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., and Rama, B. (2022). *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Quetelet, A. (1849). *Letters addressed to HRH the Grand Duke of Saxe Coburg and Gotha: On the theory of probabilities, as applied to the moral and political sciences*. C. & E. Layton.
- Rosemartin, A. H., Denny, E. G., Weltzin, J. F., Lee Marsh, R., Wilson, B. E., Mehdipour, H., Zurita-Milla, R., and Schwartz, M. D. (2015). Lilac and honeysuckle phenology data 1956–2014. *Scientific Data*, 2(1):1–8.
- Schwartz, M. D., editor (2024). *Phenology: An Integrative Environmental Science*. Springer, New York.
- Schwartz, M. D., Hanes, J. M., and Liang, L. (2014). Separating temperature from other factors in phenological measurements. *International journal of biometeorology*, 58(7):1699–1704.
- Stemkovski, M., Bell, J. R., Ellwood, E. R., Inouye, B. D., Kobori, H., Lee, S. D., Lloyd-Evans, T., Primack, R. B., Templ, B., and Pearse, W. D. (2023). Disorder or a new order: how climate change affects phenological variability. *Ecology*, 104(1):e3846.
- Switzer, J., Chamberlain, S., Marsh, L., Wong, K., and Scott, E. R. (2025). *rnnpn: Interface to the national 'phenology' network 'api'*.
- Thrasher, B., Wang, W., Michaelis, A., Melton, F., Lee, T., and Nemani, R. (2022). Nasa global daily downscaled projections, cmip6. *Scientific data*, 9(1):262.
- van der Schoot, C., Paul, L. K., and Rinne, P. L. H. (2014). The embryonic shoot: a lifeline through win-

- ter. Journal of Experimental Botany, 65(7):1699–1712.
- Vitasse, Y., Baumgarten, F., Zohner, C., Rutishauser, T., Pietragalla, B., Gehrig, R., Dai, J., Wang, H., Aono, Y., and Sparks, T. (2022). The great acceleration of plant phenological shifts. Nature Climate Change, 12(4):300–302.
- Vitasse, Y., Signarbieux, C., and Fu, Y. H. (2018). Global warming leads to more uniform spring phenology across elevations. Proceedings of the National Academy of Sciences, 115(5):1004–1008.
- Wolkovich, E., Auerbach, J., Chamberlain, C., Buonaiuto, D., Ettinger, A., Morales-Castilla, I., and Gelman, A. (2021). A simple explanation for declining temperature sensitivity with warming. Global Change Biology, 27(20):4947–4949.
- Wood, S. N. (2017). Generalized additive models: an introduction with R. chapman and hall/CRC.

S. Supporting Information

SI.1 Sketch of Theoretical Results

We follow the notation in Gut (2009). Let $X_i = \alpha + \beta i + \epsilon_i$ denote the temperature on day i , where $\alpha > 0$, $\beta \geq 0$, and the $\{\epsilon_i\}$ are independent and identically distributed with $\mathbb{E}[\epsilon_i] = 0$ and $\text{Var}(\epsilon_i) = \sigma^2$. Let $\gamma = \frac{\alpha}{\beta} + \frac{1}{2}$, and define the deterministic cumulative sum

$$\xi_n = \sum_{i=1}^n \mathbb{E}[X_i] = \alpha n + \frac{\beta}{2} n(n+1) = \frac{\beta}{2} n^2 + \beta \gamma n$$

when $\beta > 0$ and $\xi_n = \alpha n$ when $\beta = 0$.

Denote the sum of the stochastic component $S_n = \sum_{i=1}^n \epsilon_i$. The object of interest is the first-passage time of $Z_n = S_n + \xi_n$,

$$\nu(\tau) = \min\{n \geq 1 : Z_n > \tau\}.$$

In the language of Lai and Siegmund (1977), $\{Z_n\}$ is a perturbed random walk. That is, the random walk $\{S_n\}$ is perturbed by the deterministic term $\{\xi_n\}$. This setup yields two asymptotic regimes, corresponding to whether $\beta \approx 0$ as in the winter regime or $\beta > 0$ as in the spring regime.

When $\beta = 0$, we apply the usual asymptotic argument for the hitting time of a random walk with constant positive drift as $\tau \rightarrow \infty$,

$$\nu(\tau) \sim \text{Normal}\left(\frac{\tau}{\alpha}, \frac{\sigma^2 \tau}{\alpha^3}\right),$$

see Gut (2009).

When $\beta > 0$, the deterministic component $\xi_n = \frac{\beta}{2} n^2 + \beta \gamma n$ grows quadratically so that the threshold τ is reached near the deterministic crossing time $m(\tau)$ satisfying $\xi_{m(\tau)} \approx \tau$. The solution is

$$m(\tau) = \frac{-\beta \gamma + \sqrt{\beta^2 \gamma^2 + 2\beta \tau}}{\beta} = \sqrt{\frac{2\tau}{\beta}} - \gamma + o(1).$$

By the functional central limit theorem, the deviation of $\nu(\tau)$ about $m(\tau)$ is approximately normal. Linearization of $Z_{\nu(\tau)} \approx \tau$ around $m(\tau)$ yields

$$\nu(\tau) \sim \text{Normal}\left(m(\tau), \frac{\sigma^2 m(\tau)}{(\alpha + \beta m(\tau))^2}\right) \approx \text{Normal}\left(\sqrt{\frac{2\tau}{\beta}} - \gamma, \frac{\sigma^2}{\beta^{3/2} \sqrt{2\tau}}\right),$$

where we have used the fact that $\alpha + \beta m(\tau) \approx \sqrt{2\beta \tau}$ when τ is large.

We make two observations about this asymptotic approximation. First, we have written the cumulative sum as beginning on a known day $i = 1$. However, the approximation does not actually depend on the day the accumulation begins. This is because when τ is sufficiently large, the first finite number of terms of the cumulative sum are negligible by comparison. For example, for any start day $a > 1$, incorrectly starting at day $i = 1$ yields error

$$\sum_{i=1}^{\nu(\tau)} X_i - \sum_{i=a}^{\nu(\tau)} X_i = \sum_{i=1}^{a-1} X_i = O_p(1).$$

Second, we have described accumulation as occurring entirely within one of two regimes. In practice, accumulation may begin near the end of the winter regime ($\beta = 0$) and continue into the spring regime ($\beta > 0$). In such cases, the event time will behave as occurring in the spring regime when τ is large as long as a nontrivial amount of warming happens under the spring regime (i.e., the percent of time in the spring regime does not go to 0 as τ goes to infinity). This is because the quadratic accumulation in the spring regime dominates the contribution from the winter regime. Moreover, the linearization argument used to compute the variance shows that the asymptotic variance depends only on the accumulation that occurs near the event time. That is, the variance only depends on the regime active when the event occurs.

Table S1: Summary data from the Walnut forcing experiment (Charrier et al. 2011).

α	n	mean	sd
5	69	172	34.6
10	86	88	17.1
15	76	42	13.2
20	77	26	8.4
25	77	21	7.4

Table S2: Mean bloom date (day-of-year) of lilac (Rosemartin et al. 2015, updated to 2025 using USA NPN records) by winter temperatures (α , rows) and spring warming (β , columns).

$\alpha \setminus \beta$	$[-0.28, 0.07]$	$(0.07, 0.11]$	$(0.11, 0.15]$	$(0.15, 0.68]$
$(0, 0.7]$	157.70	152.07	147.60	142.20
$(0.7, 1.9]$	151.12	146.88	141.64	136.15
$(1.9, 4.7]$	136.30	131.92	127.58	124.94
$(4.7, 16.4]$	109.56	109.77	107.36	105.71

Table S3: Standard deviation of lilac bloom date (days, Rosemartin et al. 2015, updated to 2025 using USA NPN records) by winter temperatures (α , rows) and spring warming (β , columns).

$\alpha \setminus \beta$	$[-0.28, 0.07]$	$(0.07, 0.11]$	$(0.11, 0.15]$	$(0.15, 0.68]$
$(0, 0.7]$	12.80	11.94	10.90	10.95
$(0.7, 1.9]$	12.76	11.91	12.67	12.99
$(1.9, 4.7]$	16.68	14.97	14.45	12.55
$(4.7, 16.4]$	19.39	17.06	15.31	12.35

SI.2 Additional Tables

In addition to the model-based approach in the main text, we also examine the data using a binning approach. Table S1 shows data from a controlled experiment on walnut trees (*Juglans regia* and *Juglans regia* $\times$ *nigra* hybrids) conducted by Charrier et al. (2011). See also Guillaume et al. (2018). The data reflect 385 buds from 15 trees comprising 6 genotypes at 2 locations. In November of the study year, stems were sampled from each tree and cut into 7 centimeter segments containing a single bud. All segments were chilled at 4°C and then transferred to constant ‘forcing’ environments with the air temperature set at one of 5, 10, 15, 20, or 25°C, denoted α . For each temperature, the outcome is the response time, the time (in days) until budburst determined from the date at which the buds reached stage 10 of the BBCH scale.

This experiment isolates the winter regime because the forcing temperature is held constant at level α over time. To show consistency with this regime, Table S1 reports the mean and standard deviation of the response times across the 30 stems at each forcing temperature. Higher forcing temperatures advance budburst at a decreasing rate, and the standard deviation of budburst timing is smaller at warmer temperatures. Thus, in an experimental setting, the thermal-sum model captures the relationship between constant-temperature forcing and the level and variability of budburst times. This pattern cannot be explained by factors such as chilling or photoperiod, which are held constant.

Tables S2-S3 show data from the historical lilac-honeysuckle phenology network described in the main text. We bin sites and years within a grid of winter temperature (α , rows) and spring warming (β , columns) and calculate the mean and standard deviation of the observed bloom date, as the number of days since January 1. The results are consistent with Figure 2 in the main text. The mean and standard deviation are nonlinear functions of α and β . In particular, the standard deviation generally increases when α increases (holding β

fixed) and decreases when β decreases (holding α fixed).

SI.3 Additional Figures

The winter temperature (α) and spring warming (β) are shown for nine sites in Figures S1-S3, chosen from the historical lilac–honeysuckle phenology network. Figure S1 shows nine randomly selected site-years and has the same interpretation as Figure 1. Figures S2 and S3 show α and β for the nine sites with the longest running records. Figure S4 shows α and β for two random sites in the years 2050, 2075, and 2100 under temperature projections from NEX-GDDP-CMIP6 scenarios of moderate warming (SSP2-4.5) and high warming (SSP5-8.5) (Thrasher et al., 2022). Figures S5 and S6 show the projected mean and standard deviation of lilac bloom dates using the α and β calculated from the NEX-GDDP-CMIP6 data and the generalized additive model described in the main text.




Figure S1: Springtime daily midrange temperatures (points) with α (mean from Jan to Feb) and β (slope from Mar to Apr) for nine randomly selected sites from lilac-honeysuckle dataset (Rosemartin et al. 2015) as measured by GHCND stations nearby. Temperatures below 0 are set to 0.




Figure S2: Changes in α for the nine sites from lilac-honeysuckle data (Rosemartin et al. 2015) with longest running records as as measured by GHCND stations nearby. Trend line calculated using linear regression.




Figure S3: Changes in β for the nine sites from lilac-honeysuckle data (Rosemartin et al. 2015) with longest running records as as measured by GHCND stations nearby. Trend line calculated using linear regression.




Figure S4: Springtime daily midrange temperatures (points) with α (mean from Jan to Feb) and β (slope from Mar to Apr) for two randomly selected sites under two scenarios for the years 2050, 2075, and 2100 from the NEX-GDDP-CMIP6 dataset (Thrasher et al. 2022). Temperatures below 0 are set to 0.




Figure S5: Predicted mean of lilac bloom dates in years 2025, 2050, 2075, and 2100 (columns) across the continental United States under temperatures projected under two scenarios (rows) from NEX-GDDP-CMIP6 (Thrasher et al., 2022) in which additional warming is moderate (top) or high (bottom). Lighter areas indicate earlier blooms due to higher temperatures. Under SSP2-4.5, the mean bloom date is largely constant between 2050 and 2100, while under SSP5-8.5, the mean bloom date advances moderately.




Figure S6: Predicted standard deviation of lilac bloom dates in years 2025, 2050, 2075, and 2100 (columns) across the continental United States under temperatures projected under two scenarios (rows) from NEX-GDDP-CMIP6 (Thrasher et al., 2022) in which additional warming is moderate (top) or high (bottom). Light areas indicate more variation (less synchrony) due to higher temperatures. Under SSP2-4.5, the standard deviation is largely constant between 2050 and 2100, while under SSP5-8.5, the standard deviation increases dramatically, particularly in the Southeast and Southwest. These dramatic increase is due to a regime change, which we refer to as a 'scattered spring.'

SI.4 Simulations

We run two simulations to verify the interpretation of our findings. The first verifies the asymptotic approximation of the thermal-sum model derived in Section A.1. The second verifies the interpretation of the data in Section A.2.

Simulation 1

We verify the accuracy of the two asymptotic normal approximations for the hitting time

$$\nu(\tau) = \min \left\{ n \geq 1 : \sum_{i=1}^n X_i > \tau \right\}$$

under the model

$$X_i = \alpha + \beta i + \varepsilon_i, \quad \varepsilon_i \sim \text{Normal}(0, \sigma^2).$$

We set $\sigma = 20$ and generate daily temperatures until the threshold τ is reached, recording the corresponding hitting time $\nu(\tau)$. We repeat this procedure $R = 10,000$ times for thresholds $\tau = \{1000, 2000\}$, $\alpha = \{2, 4\}$ and $\beta = \{0, 0.1\}$. When $\beta = 0$, the simulation matches the winter regime, and when $\beta = 0.1$, it matches the spring regime.

We then compare these simulations with the corresponding asymptotic distribution. When $\beta = 0$, we use the distribution

$$\nu(\tau) \sim \text{Normal}\left(\frac{\tau}{\alpha}, \frac{\sigma^2 \tau}{\alpha^3}\right).$$

When $\beta > 0$, we use the distribution

$$\nu(\tau) \sim \text{Normal}\left(\sqrt{\frac{2\tau}{\beta}} - \frac{\alpha}{\beta} + \frac{1}{2}, \frac{\sigma^2}{\beta^{3/2}\sqrt{2\tau}}\right).$$

The simulated distribution of $\nu(\tau)$ is represented in Figure S7 below by the histograms, where we have standardized by subtracting the mean and dividing by the standard deviation of the corresponding asymptotic distribution. Overlaid is the standard normal density. Across both regimes, the standardized histograms align closely with the $\text{Normal}(0, 1)$ curve, and the agreement improves as τ increases, providing a visual confirmation of the derived asymptotic distributions.

Simulation 2

To verify our interpretation of the data, we simulate daily temperatures

$$X_i = \mu_i + \varepsilon_i, \quad \varepsilon_i \sim \text{Normal}(0, \sigma^2),$$

with $\sigma = 20$, where the deterministic component follows

$$\mu_i = \begin{cases} \alpha, & i = 1, \dots, 90 \quad (\text{January–February}), \\ \alpha + \beta(i - 90), & i = 91, \dots, 180 \quad (\text{March–April}). \end{cases}$$

Bloom occurs when cumulative temperature first exceeds a threshold τ , i.e.

$$\nu(\tau) = \min \left\{ n \geq 1 : \sum_{i=1}^n X_i > \tau \right\}.$$

We simulated $R = 10,000$ independent realizations of $\nu(\tau)$ for each combination of $\alpha \in \{4, 8, 10\}$, $\beta \in \{0.2, 0.4, 0.8\}$, and $\tau \in \{1000, 2000\}$. We report the mean (left) and standard deviation (right) of $\nu(\tau)$ in Tables S4a–S4d, arranged with rows indexing α and columns indexing β .




Figure S7: Simulations comparing the distribution of bloom dates (histogram) with an asymptotic approximation (solid line) under two regimes: winter (top two rows) and spring (bottom two rows).

Table S4: Simulated mean and standard deviation of the bloom date $\nu(t)$ across $R = 10,000$ replicates with thresholds $\tau \in \{1000, 2000\}$. Rows denote the winter temperature (α), and columns denote spring warming (β).

(a) mean, $\tau = 1000$				(c) sd, $\tau = 1000$			
$\alpha \setminus \beta$	0.2	0.4	0.8	$\alpha \setminus \beta$	0.2	0.4	0.8
4	151.66	136.83	124.86	4	15.48	10.42	7.21
8	114.87	111.27	106.85	8	16.53	13.27	10.50
10	98.45	97.20	95.72	10	15.94	14.45	12.71

(b) mean, $\tau = 2000$				(d) sd, $\tau = 2000$			
$\alpha \setminus \beta$	0.2	0.4	0.8	$\alpha \setminus \beta$	0.2	0.4	0.8
4	199.54	171.15	149.16	4	10.96	7.22	4.84
8	169.81	152.43	137.37	8	10.93	7.50	5.11
10	156.22	143.40	131.34	10	10.69	7.66	5.36

The results reproduce the trends observed in the lilac data. Mean bloom dates decrease as either α increases or β increases (Tables S4a and S4b). Within a fixed α row, increasing β reduces the standard deviation of bloom dates (Tables S4c and S4d), consistent with the prediction that increasing average temperatures diminish the influence of noise on the threshold-crossing time. Holding β fixed, the standard deviation is often larger at higher α (particularly for $\beta \in \{0.4, 0.8\}$), consistent with the prediction that earlier crossings occur in a flatter portion of the seasonal cycle where microclimate variability has greater influence.