

# Temperature responses of developmental processes have not been affected by breeding in different ecological areas for 17 crop species

Boris Parent<sup>1</sup> and François Tardieu<sup>2</sup>

<sup>1</sup>Australian Centre for Plant Functional Genomics, PMB1, Glen Osmond, SA 5064, Australia; <sup>2</sup>INRA, UMR759 Laboratoire d'Ecophysiologie des Plantes sous Stress Environnementaux. Place Viala, F-34060 Montpellier, France

Author for correspondence:  
François Tardieu  
Tel: +33 04 99 61 26 32  
Email: francois.tardieu@supagro.inra.fr

Received: 30 November 2011  
Accepted: 19 January 2012

New Phytologist (2012) **194**: 760–774  
doi: 10.1111/j.1469-8137.2012.04086.x

**Key words:** development, growth, genetic variability, maize (*Zea mays*), response, rice (*Oryza* spp.), temperature, wheat (*Triticum aestivum*).

## Summary

- Rates of tissue expansion, cell division and progression in the plant cycle are driven by temperature, following common Arrhenius-type response curves.
- We analysed the genetic variability of this response in the range 6–37°C in seven to nine lines of maize (*Zea mays*), rice (*Oryza* spp.) and wheat (*Triticum aestivum*) and in 18 species (17 crop species, different genotypes) via the meta-analysis of 72 literature references.
- Lines with tropical or north-temperate origins had common response curves over the whole range of temperature. Conversely, appreciable differences in response curves, including optimum temperatures, were observed between species growing in temperate and tropical areas.
- Therefore, centuries of crop breeding have not impacted on the response of development to short-term changes in temperature, whereas evolution over millions of years has. This slow evolution may be a result of the need for a synchronous shift in the temperature response of all developmental processes, otherwise plants will not be viable. Other possibilities are discussed. This result has important consequences for the breeding and modelling of temperature effects associated with global changes.

## Introduction

Lines and species of crop plants have evolved and been selected in different latitudes and altitudes, thereby generating differences in temperature adaptation via a large range of mechanisms (Atkin *et al.*, 2006). Indeed, gradients of latitude are accompanied by gradients of traits, such as the duration of the plant cycle, specific leaf area and survival from heat or cold stresses (Kratsch & Wise, 2000; Atkin *et al.*, 2006; Barnabas *et al.*, 2008; Penfield, 2008). The analysis of the genetic variability of the response of crop plants to temperature is therefore essential in a context in which climate changes increase the risk of yield loss because of high temperatures, without necessarily reducing the risk of exposure to low temperatures (Battisti & Naylor, 2009; Lobell *et al.*, 2011).

The analysis of short-term response curves differs from the study of the irreversible effects at high or low temperature, which require several weeks to become evident (Wise *et al.*, 1983; Bukovnik *et al.*, 2009; Gorsuch *et al.*, 2010). It involves the quantitative analysis of the short-term effects of changes in temperature over a large range, not limited to its low or high ends, in which effects are reproducible for a given line and a given range of conditions, are reversible and not dependent on the temperature history of the plant, and are independent of the intrinsic value of the studied variable measured at a standard temperature. When these

conditions are fulfilled, meta-analyses of a series of experiments result in response curves to temperature or other environmental variables that allow the comparison of life kingdoms (Gillooly *et al.*, 2001; Dell *et al.*, 2011) or fitness between species in ecological studies (Poorter *et al.*, 2010). They also allow the study of quantitative genetics based on response parameters (Reymond *et al.*, 2003; Yin *et al.*, 2004) or analysis of the coordination between different mechanisms under fluctuating conditions (Parent *et al.*, 2010b; Tardieu *et al.*, 2011).

Following pioneering studies that compared the temperature responses of phyllochron and leaf elongation rate (LER) (Ong, 1983a,b; Warrington & Kanemasu, 1983a,b), we have shown recently that the germination rate, cell division rate, leaf initiation rate, LER and the reciprocal of the duration of phenological phases follow a common response to temperature after normalization by their absolute rates at 20°C (Parent *et al.*, 2010b). This implies that these processes have similar temperature responses. Conversely, the responses of *in vitro* enzyme activities and of net photosynthesis differ from those mentioned above (Parent *et al.*, 2010b). It is therefore relevant to analyse jointly the temperature responses of rates involved in the progression of plant development or in expansive growth (called 'developmental rates' hereafter), independent of those involved in plant metabolism and biomass production.

The response to temperature of enzyme activities has been described by the equation of Johnson *et al.* (1942), adequately extended to processes as diverse as the growth of bacteria (Johnson & Lewin, 1946), parameters of photosynthesis (Farquhar *et al.*, 1980), insect development (Sharpe & Demichele, 1977), plant development (Feng *et al.*, 1990; Parent *et al.*, 2010b) and 1072 physiological and ecological traits in microbes, animals and plants (Dell *et al.*, 2011).

$$F(T) = \frac{A T e^{\left(\frac{-\Delta H_A^\ddagger}{RT}\right)}}{1 + e^{\left[\frac{\Delta S_D}{R} - \frac{\Delta H_D}{RT}\right]}} \quad \text{Eqn 1}$$

(Johnson *et al.*, 1942) ( $F(T)$  is the considered rate,  $T$  is the temperature (K),  $\Delta H_A^\ddagger$  (J mol<sup>-1</sup>) is the enthalpy of activation of the reaction and determines the curvature at low temperature (Supporting Information Fig. S1a),  $\Delta H_D$  (J mol<sup>-1</sup>) and  $\Delta S_D$  (J mol<sup>-1</sup> K<sup>-1</sup>) are the enthalpy and entropy of enzyme inactivation at high temperature and  $A$  is the trait scaling coefficient). When describing integrated variables instead of enzyme activities, the parameters lose their biochemical nature, and so a more intuitive modified version was preferred:

$$F(T) = \frac{A T e^{\left(\frac{-\Delta H_A^\ddagger}{RT}\right)}}{1 + \left[ e^{\left(\frac{-\Delta H_A^\ddagger}{RT}\right)} \right]^{\alpha \left(1 - \frac{T}{T_0}\right)}} \quad \text{Eqn 2}$$

where  $\alpha$  (dimensionless) is the ratio  $\Delta H_D/\Delta H_A^\ddagger$  in Eqn 1 and determines how sharp is the decrease in rate at high temperature (Fig. S1c).  $T_0$  (K) equals the ratio  $\Delta H_D/\Delta S_D$  in Eqn 1. It is the temperature at which half of the system is in an inactive state (Fig. S1b) and determines the temperature at which the rate is at a maximum ( $T_{opt}$ , Methods S1). After normalization by the rate at 20°C (to scale different processes and species and to compare them independently of their absolute values; Parent *et al.*, 2010b), parameter  $A$  is not free and depends on  $\Delta H_A^\ddagger$ ,  $T_0$  and  $\alpha$ .

The analysis of the genetic variability of the temperature response of the developmental rates will provide essential elements to understand the thermal adaptation of contrasting genotypes grown in common ranges of temperature. A very low or nonexistent genetic variability between species or lines originating from cold or warm areas may suggest that this relationship is based on an immutable law (Eqn 2) with parameters based on physical constraints. The existence of genetic variability would indicate that mutations on key genes can substantially alter the regulatory pathways that govern the temperature response of developmental rates (McClung & Davis, 2010). Although large-scale comparisons between the temperature responses of different species and life kingdoms have been studied for enzyme activities (Campbell *et al.*, 2007) and metabolic rates (Gillooly *et al.*, 2001), such comparisons are not available for developmental processes. In this study, we performed the following investigations:

- We tested the working hypotheses for the analysis of temperature responses, that is, that these responses are reproducible and reversible without after-effects of previous temperature, and that different developmental processes have similar temperature responses after normalization by their absolute rate at a standard temperature.
- We tested whether lines originating from cold or warm areas have different temperature responses; this was performed for lines of maize (*Zea mays*), wheat (*Triticum aestivum*) and rice (*Oryza*) with diverse breeding histories and ecological origins, from temperate to tropical, by testing whether the parameters of Eqn 2 were common or differed between the lines of one species.
- We extended our analysis to the diversity of responses in 18 species (17 major crop species comprising the 14 most cultivated species in the world (<http://faostat.fao.org>; Table 1) and *Arabidopsis thaliana*) by compiling literature data. We tested the commonality of response curves by asking whether one or more parameter(s) of Eqn 2 could be considered as common to several species and, within each species, to different genotypes.

## Description

### Genetic material

Eight maize (*Zea mays* L.) lines were used in the experiments, originating from the Caribbean zone (latitudes 10–20°), tropical areas (lines CML69 and CML444; [http://apps.cimmyt.org/english/wps/obtain\\_seed/cimmytCMLS.htm](http://apps.cimmyt.org/english/wps/obtain_seed/cimmytCMLS.htm)) or tropical highlands (line F331); the US corn belt (latitudes 30–40°; dent lines B73, A188 and Io); and Canada, grown since then in Europe in latitudes 40–50° (flint lines UH007 and F2). These lines are genetically very distant (Camus-Kulandaivelu *et al.*, 2006; A. Charcosset, pers. comm.) and were bred in different thermal environments (average temperature of 19°C in temperate environments, for example, north USA, or 24°C in tropical environments; Lobell *et al.*, 2011). The hybrid Io × F2 (Dea) was considered in the same analysis.

The seven studied rice lines belonged to two species, *Oryza glaberrima* Steud. (CG14) and *Oryza sativa* L. (other cultivars), and three subfamilies of *Oryza sativa*: temperate *japonica* (Nipponbare), tropical *japonica* (Azucena, Moroberekan) and *indica* (Apo, Vandana and IR64). They originated from temperate (Nipponbare; Koumura, 1972) or tropical (others) regions. They therefore experienced different thermal environments during their breeding history, from nearly constant high temperatures in tropical areas (23–31°C in the Philippines; Peng *et al.*, 2004) to large temperature amplitudes (5–30°C in Japan; <http://www.knowledgebank.irri.org>). These lines also differ in their year of release, from traditional cultivars (Azucena), cultivars developed during the green revolution (IR64) in the 1980s (Moroberekan) and recently released cultivars (Vandana, Apo).

Seven wheat (*Triticum aestivum* L.) lines were analysed (Drysdale, Gladius, Kukri, Excalibur, RAC875, Chinese Spring and Bob White). Chinese Spring is a Chinese landrace, selected in a cool climate (average temperature *c.* 10°C; Lobell *et al.*, 2011), whereas South Australian lines were selected under high thermal

**Table 1** Origin of the data for the different processes in 18 species

Species	Origin of data	Process	Species	Origin of data	Process
Maize	New data	Leaf elongation	Sorghum	Lafarge <i>et al.</i> (1998)	Leaf elongation
	Database	Leaf elongation		Lawlor <i>et al.</i> (1990)	Germination
	Parent <i>et al.</i> (2010b)	Germination		Wade <i>et al.</i> (1993)	Germination
	Barlow & Boersma (1972)	Seedling elongation	Millet	Garciahuidobro <i>et al.</i> (1982)	Germination
	Ben-Haj-Salah & Tardieu (1995)	Cell division		Mohamed <i>et al.</i> (1988a)	Germination
	Blacklow (1972)	Shoot initiation		Mohamed <i>et al.</i> (1988b)	Leaf elongation
		Shoot elongation		Ong & Monteith (1985)	Root elongation
		Radicle elongation		Ong (1983a)	Leaf appearance
		Radicle initiation		Ong (1983b)	Leaf elongation
		Development (primordium–tip)		Pearson (1975)	Leaf appearance
	Hesketh & Warrington (1989)	Development (tip–ligule)	Cotton	Arndt (1945)	Cotyledon emergence
		Growth duration <sup>-1</sup>			Hypocotyl elongation
		Leaf elongation			Primary root elongation
		Leaf appearance		Hesketh <i>et al.</i> (1972)	Development (flower–boll)
	Lehenbauer (1914)	Seedling elongation			Development (square–flower)
	Tollenaar <i>et al.</i> (1979)	Leaf appearance			Relative leaf growth
	Warrington & Kanemasu (1983a)	Development (sowing–anthesis)			Plastochrons (flower, square, leaf)
		Development (sowing–tassel initiation)		Mutsaers (1983)	Relative growth
		Emergence			
	Warrington & Kanemasu (1983b)	Leaf appearance		Reddy <i>et al.</i> (1997)	Relative internode expansion
Rice	New data	Leaf elongation			Relative leaf expansion
	Database	Leaf elongation			Leaf unfolding
	Parent <i>et al.</i> (2010b)	Germination	Canola	Marshall & Squire (1996)	Germination
	Summerfield <i>et al.</i> (1992)	Development (sowing–flowering)		Morrison <i>et al.</i> (1989)	Development (sowing–maturity)
		Leaf appearance (f5–f6)		Singh <i>et al.</i> (2008)	Pollen tube elongation
	Yin & Kropff (1996)	Development (sowing–flowering)		Vigil <i>et al.</i> (1997)	Pollen germination
	Yin <i>et al.</i> (1996)				Emergence
Wheat	New data	Emergence	Sunflower	Chimenti <i>et al.</i> (2001)	Embryo growth
		Seedling elongation			
	Addae & Pearson (1992)	Germination	Peanut	Villalobos & Ritchie (1992)	Leaf appearance
		Emergence		Kakani <i>et al.</i> (2002)	Pollen tube elongation
	Cutforth <i>et al.</i> (1992)	Germination		Mohamed <i>et al.</i> (1988a)	Germination
	Dejong & Best (1979)	Emergence		Prasad <i>et al.</i> (2006)	Development (sowing–V2 stage)
		Emergence	Sugar cane		Emergence
		Coleoptyl elongation		Campbell <i>et al.</i> (1998)	Node production
		Internode elongation		Inmanbamber (1994)	Plant expansion
	Gallagher <i>et al.</i> (1979)a	Leaf elongation		Mongelar & Mimura (1972)	Spindel elongation
	Kemp & Blacklow (1982)	Leaf elongation	Cassava	Akparobi <i>et al.</i> (2000)	Growth duration <sup>-1</sup>
	Lafond & Baker (1986)	Germination			Leaf initiation
	Lafond & Fowler (1989)	Germination			Leaf opening
<i>Arabidopsis thaliana</i>	Lindstrom <i>et al.</i> (1976)	Emergence		Keating & Evenson (1979)	Emergence
	Parent <i>et al.</i> (2010b)	Germination	Potato	Benoit <i>et al.</i> (1986)	Stem elongation
	Granier <i>et al.</i> (2002)	Growth duration <sup>-1</sup>		Fleisher & Timlin (2006)	Cell expansion and division
		Leaf expansion			Growth
Soybean	Orbovic & Poff (2007)	Leaf initiation		Fleisher <i>et al.</i> (2006)	Leaf appearance
		Seedling elongation		Kirk & Marshall (1992)	Leaf elongation
	Covell <i>et al.</i> (1986)	Germination			Leaf appearance
	Hesketh <i>et al.</i> (1973)	Development (3rd trifoliolate–flower)	Cowpea	Covell <i>et al.</i> (1986)	Germination
		Development (planting–trifoliolate)		Craufurd <i>et al.</i> (1996a)	Emergence
		Development (unifoliolate–trifoliolate)			Germination
		Trifoliolate production		Craufurd <i>et al.</i> (1996b)	Development (sowing–flowering)
		Node production		Craufurd <i>et al.</i> (1996c)	Development (sowing–flowering)
	Lawn & Hume (1985)	Pod production		Craufurd <i>et al.</i> (1997)	Leaf appearance
				Hadley <i>et al.</i> (1983)	Development (sowing–flowering)

Table 1 (Continued)

Species	Origin of data	Process	Species	Origin of data	Process
Barley	New data	Emergence	Common bean	Moss & Mullett (1982)	Development (sowing–ripe pods)
		Seedling elongation		Scully & Waines (1987)	Development (sowing–visible bud)
	Ellis <i>et al.</i> (1987)	Germination		White & Montes (1993)	Seedling elongation
	Gallagher & Biscoe (1979)	Germination	Cauliflower	Olesen & Grevsen (1997)	Germination
		Leaf elongation		Wurr <i>et al.</i> (1990)	Leaf area expansion
					Curd growth


Information on exact processes, materials and methods, and genotypes is available in the original papers.

amplitude (RAC875, Excalibur, Gladius, temperature from 3 to 40°C, average temperature *c.* 16°C, records from ACPFG weather stations). Excalibur and RAC875 are considered as heat tolerant and Kukri as heat sensitive (Izanloo *et al.*, 2008; Bukovnik *et al.*, 2009). Lines differ in their year of selection, from the old cultivars Chinese Spring (1940) and Bob White (1970) to recent varieties Drysdale (2002) and Gladius (2008). A barley line (Golden Promise) was added to this series to analyse the behaviour of another species from the Triticeae tribe.

### LER in maize and rice subjected to short-term temperature changes

All measurements of LER were performed during the night, under well-watered conditions with wet air (air vapour pressure deficit maintained under 0.8 kPa) to avoid confusion between the effects of temperature and of evaporative demand (Ben-Haj-Salah & Tardieu, 1997). Data presented in this study are the result of a meta-analysis of several independent experiments with similar protocols.

Maize and rice plants were grown in polyvinylchloride (PVC) columns in a glasshouse with temperature fluctuating in the range 18–32°C, light in the range 10–20 mol d<sup>-1</sup> m<sup>-2</sup> and air vapour pressure deficit in the range 1–2.5 kPa. Soil was maintained at a water potential above –0.05 MPa via irrigation with a 1/10 Hoagland-type nutrient solution controlled by individual pot weight (<http://bioweb.supagro.inra.fr/phenodyn/>). Temperatures taken into account in this article are the meristem temperatures, measured with thermocouples inserted near the meristem of at least five plants (<http://bioweb.supagro.inra.fr/phenodyn/>). Meristem temperatures were averaged every 15 min and stored in the database PhenodynDB (<http://bioweb.supagro.inra.fr/phenodyn/>), together with air temperature, irradiance, air vapour pressure deficit and soil water content. At the six-leaf stage, plants were transferred to the growth chamber under different temperature conditions. Environmental conditions were measured using the same methods as in the glasshouse. Typical temperature scenarios are presented in Fig. 1(c). They were intentionally varied between days and plants to avoid any confounding effect of sequences of temperature (e.g. the step at 16°C that would systematically follow a step at 28°C). In 10 plants of lines B73, CML444, A188, F-2 and UH007, LER was measured at 20°C after periods of 4 h at 6 or 35°C in order to test the possible after-effect of low or high temperature.



**Fig. 1** Time courses of meristem temperature (*T*) and leaf elongation rate (LER) of the maize (*Zea mays*) line CML444. Grey panels indicate night periods. (a,b) Plants were placed in the growth chamber under a controlled environment, transferred to the glasshouse for a cold night (second night) and then transferred back to the growth chamber (third night) under a similar temperature scenario. (c,d) Plants were placed in the growth chamber under a wide range of temperatures from 6 to 37°C. It should be noted that temperature scenarios changed every day.

The temperature control of the growth chamber was based on the meristem temperature. Relative humidity, light intensity and air temperature were measured as in Parent *et al.* (2010b). The night temperature varied in steps of 3–4 h (Fig. 1c). LER was measured in leaf 6 with rotating displacement transducers (<http://bioweb.supagro.inra.fr/phenodyn/>) during the period with linear elongation (i.e. 2 d after leaf appearance in rice, Parent *et al.*, 2009; 5–6 d in maize, Sadok *et al.*, 2007), and after stabilization of temperature and LER (usually the last 2 h of each temperature step). A cloud of points of LER vs temperature, corresponding to one line in one experiment, can be seen in Fig. S2(a).



In addition, glasshouse experiments with maize or rice were included in the dataset, in particular that presented in Fig. 1(a) in maize. During the latter experiment, plants were transferred between the growth chamber and the glasshouse in order to test the after-effect of a cold night with continuously decreasing temperature. Each data point for regression is a coupled value of the meristem temperature and LER for 3 h. A cloud of points corresponding to one line in several experiments can be seen in Fig. S2b. A total of 2630 coupled values of LER and temperature were collected for maize and 598 for rice in growth chamber and glasshouse experiments. Finally, a dataset was added, for hybrid Dea only, comprising measurements of the cell division rate in leaves (Ben-Haj-Salah & Tardieu, 1995).

### Germination rate at constant temperature in wheat, barley, maize and rice

The germination rate was measured under constant temperature in the seven above-mentioned wheat lines, one barley line (Golden Promise), one maize hybrid (Dea) and two rice lines (Azucena and IR64). Seeds were sown in Petri dishes (100 seeds per Petri dish) lined with saturated Whatman filter paper and placed under different temperatures (4, 9, 15.5, 20, 24.5, 28 and 35°C). Germination was scored every 8 h (6 h in rice and maize) for 2 d, every 12 h for the next 3 d, and every 1 d after that. The appearance of a radicle was scored as a germinated seed. The time for 50% cumulative germination was determined by regression on the progression curves. Data points were not taken into account when fewer than 50% of seeds germinated.

### Emergence rate and seedling elongation rate at constant temperature in wheat and barley

Seedling elongation rate was measured in the wheat and barley lines presented above. Twelve seeds per line and per tested temperature were sown at a depth of 1 cm in pots filled with coco peat. Pots were placed in trays with 1 cm of free water and kept in the dark, with saturated soil and air. The appearance of coleoptiles at the soil surface was scored, and the length of the emerged coleoptiles was measured every 12 h during 5 d, and every 1 d thereafter. The time for 50% cumulative emergence was determined by regression on the progression curves. Data points were not taken into account when fewer than 50% of the initial number of seeds emerged. The coleoptile elongation rate was calculated by linear regression between 10 and 100 mm length for wheat (common linear phase in all lines) and between 10 and 60 mm length for barley.

### Meta-analysis of lines originating from our database

**Normalization and pooling of data** Each process rate (e.g. LER, germination rate, etc.) was normalized by its mean value at 20°C (Parent *et al.*, 2010b), thereby allowing joint analysis for different processes, regardless of dimensions, units or absolute values. The normalization at 20°C was chosen because data at 20°C are

available in most publications, species and processes. The rate at 20°C was determined via a linear regression performed in the vicinity of 20°C (e.g. from 18 to 22°C). Because information criteria for model comparison are sensitive to the number of data points, analyses were performed after pooling data points in 15–16 groups (example in Fig. S2c).

**Test of differences in model parameters** The model presented in Eqn 2 has three parameters (because  $A$  depends on  $\Delta H_A^\ddagger$ ,  $T_0$  and  $\alpha$ , Methods S2). In order to determine which parameters are common/differ between lines, the model with three parameters (called M.3 hereafter) was compared with seven models in which one, two or three parameters were common to the studied lines (Table 2). Three models had two free parameters (M.2a, M.2b and M.2c) and one parameter common to the studied lines. Three models (M.1a, M.1b and M.1c) had one free parameter and two parameters common to the studied lines. Finally, one model (M.0) had no free parameter, thereby resulting in a common temperature response for all lines.

**Calculation of model parameters** Parameters of individual response curves were calculated using the nonlinear regression function *nls* of the R language (R Development Core Team, 2005). Depending on the number of free/fixed parameters, different procedures were used to ensure convergence in the fitting procedure (Fig. S3).

**Model comparison** We used several indicators to compare nonlinear models, including the root-mean-square error (RMSE) and the coefficient of variation of the RMSE ( $CV_{RMSE}$ ). We also used indicators that add a penalty to the number of estimated parameters, thereby avoiding overfitting, namely the Akaike information criterion (AIC; Akaike, 1974; function *AIC()* in R software) and the Bayesian information criterion (BIC; Schwarz, 1978; function *BIC()*). The best model according to these indicators is that with the lowest values of BIC and AIC. The Akaike weights (Akaike, 1978) indicate the probability for one model to be the best model given the dataset. Here, we calculated these weights for BIC values ( $w_{BIC}$ ).

**Table 2** Number of free/fixed parameters in each of the eight studied models

Model	No. of free parameters	$\Delta H_A^\ddagger$	$T_0$	$\alpha$
M.3 (Eqn 2)	3	Free	Free	Free
M.2a	2	Fixed	Free	Free
M.2b	2	Free	Free	Fixed
M.2c	2	Free	Fixed	Free
M.1a	1	Fixed	Free	Fixed
M.1b	1	Fixed	Fixed	Free
M.1c	1	Free	Fixed	Fixed
M.0	0	Fixed	Fixed	Fixed

The three parameters are those of Eqn 2. In each model, a free parameter is calculated individually for each genotype or species. A fixed parameter is common to all considered genotypes or species.

For an easier interpretation of these criteria, we compared model indicators with that of model M.3.  $\Delta CV$  is the difference between  $CV_{RMSE}$  of one model and that of model M.3, and is interpreted as the additional error when fixing parameters. With regard to experimental errors and the difference targeted between lines, an additional error of 5% was accepted for models with fewer parameters.  $\Delta AIC$  and  $\Delta BIC$  are the differences in values between the considered model and model M.3.  $\Delta AIC$  and  $\Delta BIC$  of  $< 5$  were accepted for models with fewer parameters compared with model M.3.

**Sensitivity analysis on model comparison** The sensitivity of statistical indicators to changes in the temperature response was tested using the procedure described in Fig. S4. New datasets were created by modifying that of maize. To test a shift in the whole response curve, including  $T_{opt}$ , all temperatures in the original dataset were shifted by a step of 1°C between  $-2$  to  $+2$ °C in relation to the original values (for example, the temperature was lowered by  $-1$ °C for each coupled value of rate/temperature; Fig. S4c,d). To test a difference in the sensitivity of rates to high temperature, rates above the optimum temperature were lowered or increased by a step of 5% from  $-20\%$  to  $+20\%$  in relation to the original values (e.g. for  $+10\%$ , all data points above  $30.8$ °C were multiplied by 1.1; Fig. S4a,b). Each new dataset was compared with the original by calculating  $\Delta BIC$  and  $\Delta CV$  between model M.3 and model M.0.

**Comparison of four species from our database** Normalized data corresponding to different lines were pooled within each of the four tested species (maize, rice, wheat and *Arabidopsis thaliana*). We used a similar procedure to that explained above to compare lines, taking into account 13–20 groups of data per species. Parameter calculation and model comparison were carried out as described above for lines. To confirm the results of model comparison obtained with the Bayesian approach, a cascade of *F*-tests was carried out on the nested models (R function `anova.nls()`). The simplest model was accepted unless the test was significant ( $P < 0.05$ ).

#### Comparison of 18 species from our database or from the literature

All other data were collected from the literature (Table 1), from tables or figures of the original papers. The positions of data points were recorded in figures by image analysis (software ImageJ (<http://rsbweb.nih.gov/ij/>)). In Covell *et al.* (1986) and Ellis *et al.* (1987), only the 50% germination rates were retained in the analysis. For the germination rates taken from Scully & Waines (1987), only the crop species were analysed. In Hadley *et al.* (1983), only the photoperiod insensitive lines were analysed. In Weaich *et al.* (1996), the coleoptile elongation rate was calculated as the reciprocal of the duration to reach a length of 15 mm. The internode elongation rate was calculated as the reciprocal of the duration to reach a length of 25 mm. In Summerfield *et al.* (1992), Yin & Kropff (1996) and Yin *et al.* (1996), only data obtained under constant temperature were used in the analysis.

Each dataset was normalized by the corresponding value at 20°C using the procedure described above. A pooling of the data resulted in 13–20 groups per species. For each model, the fixed parameter values were those of the four species analysed. Parameter values and model criteria were calculated as in the line analysis.

The optimum temperature  $T_{opt}$ , at which the rate is at a maximum, was calculated from model parameters (Methods S1):

$$\frac{1}{T_{opt}} = \frac{1}{T_0} + \frac{R}{\alpha \Delta H_A^\ddagger} \log \left( \frac{(\alpha - 1) \Delta H_A^\ddagger - R * 303}{\Delta H_A^\ddagger + R * 303} \right) \quad \text{Eqn 3}$$

Confidence intervals of model parameters and  $T_{opt}$  were calculated by bootstrapping (1000 replicates).


## Results

The working hypotheses for the analysis of temperature responses were fulfilled

The first condition mentioned in the Introduction is that a trait has a unique response to temperature over a large range of conditions for a given line of the studied species. This was the case when either similar or different temperature scenarios were applied on several days (Fig. 1a,b, days 1, 3, 4, 5, and Fig. S2a). Similar results were found in rice (Parent *et al.*, 2010b).


Second, the establishment of response curves requires that the temperature effect is reversible, that is, the rate at a given temperature is independent of previous temperatures during the experiment. This was the case for durations of exposure ranging from hours to days. Exposures to 6 or 34°C for 4 h (the duration of a typical peak of temperature in natural conditions) had no significant effect on LER at 20°C ( $P = 0.6$ ; Figs 1c,d, S5a,b). In the same way, LER was unaffected by a previous long night at low temperature (Fig. 1a,b, night 2). This resulted in a unique response curve per line, regardless of plant, experiment and thermal history, even though absolute rates at a given temperature differed between lines (Fig. S2). Lines of rice showed a similar behaviour in this respect (data not shown).

The domain of validity of the response covered a large range of developmental processes. After normalization by the rate at 20°C, which allows one to scale different processes that have different units, the response to temperature in one line was common between several developmental processes in maize (hybrid Dea; Fig. 2a), rice (line Azucena; Fig. 2r) and wheat (variety Drysdale; Fig. 2j). A meta-analysis of 21 sources of information (Table 1) extended this conclusion to traits referring to leaves, roots or seedlings, growth rates, cell division rate, germination rate or reciprocal of duration of phenological phases. This led to a common response curve for all of these processes, exemplified in Fig. 3(b) for maize. It is noteworthy that processes associated with response times of a few days (e.g. phyllochron) showed similar response curves to those having a response time of minutes (e.g. elongation rates; Figs 2a, 3b), indicating that the responses of the processes studied here were essentially time independent in the range 6–37°C.



**Fig. 2** Response curves to temperature of several rates involved in plant development in lines of maize (a–i), wheat (j–p), barley (q) and rice (r–x). Rates are normalized by their mean value at 20°C, set to unity. For each species, the red line corresponds to a common model for all genotypes (model M.0, Table 2). A common model applies to barley (q) and wheat (j–p). (a,j,r,q) The normalized responses to temperature of several developmental processes presented on the same graph. Green circles, leaf (a,r) or seedling (j,q) elongation rate; red squares, reciprocal of the duration from sowing to emergence; blue triangles, germination rate; orange circles, cell division rate. In all panels, blue circles represent the average of all studied processes. Other panels present the response of leaf elongation rate (maize and rice) or germination and seedling elongation (wheat). Error bars indicate standard deviation (SD).

Hence, response curves were established hereafter by considering data points originating from different thermal scenarios, different developmental processes and from our own experiments or from the literature. In the experiments presented below, most comparisons between lines were performed on the temperature response of LER.




**Fig. 3** Response curve to temperature of leaf elongation rate for nine maize (*Zea mays*) lines (a) and of several processes in a meta-analysis of literature data (b). Each process is normalized by its corresponding value at 20°C. The red line represents a common model for all processes and genotypes (M.0, Table 2). Regression parameters are summarized in Table 3. (a) Leaf elongation rate for genotypes: CML444, purple; CML69, red; B73, orange; A188, black; F2, cyan; F331, green; Io, brown; Dea, deep blue. (b) Temperature dependence of several developmental processes. Circles represent expansion processes: leaf elongation, new data (blue); coleoptile elongation from Weaich *et al.* (1996) (red); seedling elongation from Lehenbauer (1914) (cyan) and Barlow & Boersma (1972) (purple); inter-node elongation from Weaich *et al.* (1996) (black); shoot elongation from Blacklow (1972) (green); radicle elongation from Blacklow (1972) (yellow). Triangles represent organ initiation and appearance: radicle initiation from Blacklow (1972) (black); shoot initiation from Blacklow (1972) (blue); leaf initiation from Warrington & Kanemasu (1983b) (red); leaf appearance from Hesketh & Warrington (1989) (purple); leaf appearance from Warrington & Kanemasu (1983b) (yellow); leaf appearance from Tollenaar *et al.* (1979) (cyan). Diamonds represent development rate between two events: leaf primordium to leaf tip appearance from Hesketh & Warrington (1989) (purple); leaf tip appearance to ligulation from Hesketh & Warrington (1989) (red); growth duration from Hesketh & Warrington (1989) (blue); sowing to tassel initiation from Warrington & Kanemasu (1983a) (cyan); sowing to anthesis from Warrington & Kanemasu (1983a) (yellow). Squares represent germination and emergence: germination from Parent *et al.* (2010b) (blue); emergence from Warrington & Kanemasu (1983a) (turquoise). Stars represent cell division from Ben-Haj-Salah & Tardieu (1995).

The genetic variability was very low between lines originating from tropical or temperate regions, from different breeding generations and with different heat tolerances

In maize, the nine studied lines originating from various regions and climates (Caribbean zone, tropical highlands, US corn belt and temperate regions of Canada and Europe) were associated

with similar response curves (Fig. 2), almost indistinguishable over the whole range of temperature when plotted together (Fig. 3a). To identify putative differences between these curves, we tested whether one or more parameters of Eqn 2 could be considered as common to all lines. A model with three free parameters (model M.3, three parameters differing between lines) was compared with seven models with zero, one or two free parameters (Table 2). Statistical criteria indicated that the best model was model M.0, which considers that all three parameters are common to the nine studied lines (Fig. 4a,b and Table S1). Furthermore, CVs of errors increased by 2% only in model M.0 compared with the others, far from the 5% error that was considered as the limit in view of experimental errors (Fig. 4b). A sensitivity analysis showed that this procedure was able to detect a 1°C shift in the whole response curve towards higher or lower temperature (Fig. 4c,d). It also showed that we could detect a difference in rate of  $\pm 10\%$  in the range above  $T_{opt}$ , where precision was lower than in the range below  $T_{opt}$  (Fig. 4a,b). It can therefore be concluded that a unique response curve applied to maize lines with very different origins, to hybrid as well as to inbred lines. This conclusion can be extended to 23 lines described in the literature, with different origins and dates of release, including one line released in 1914 (Table 1, Fig. 3b). All lines had a response curve with common parameters to those calculated with our data (Fig. 3b).



**Fig. 4** Comparison of models presented in Table 2. (a,b) Comparison of models assuming that zero to three parameters differ between genotypes of each species. (a) Box-plot of the differences in the Bayesian information criterion ( $\Delta BIC$ ) between model M.3 and the seven other models for genotypes of maize (white), rice (grey) and wheat (black). The lowest values of  $\Delta BIC$  indicate the best models. (b) Difference in coefficient of variation of the error ( $\Delta CV$ , in % of CV of model M3) between model M.3 and the seven other models for genotypes of the three species. Lowest values of  $\Delta CV$  indicate the best models. The horizontal line represents the threshold of tolerance of 5% of additional error compared with model M.3. (c,d) Comparison of models assuming that zero to three parameters differ between four species (maize, wheat, rice and *Arabidopsis thaliana*). (c) Box-plot of  $\Delta BIC$  between model M.3 and the seven other models for the four species. (d)  $\Delta CV$  between model M.3 and the seven other models for the four species. Line in (d) as in (b). Other model selection criteria are presented in Supporting Information Table S1.

In rice, the seven studied lines belonged to different species and subspecies (*O. glaberrima*, *O. sativa* ssp. *indica* and *japonica*) from tropical or temperate regions of Asia or Africa. As in maize, the comparison of the different statistical criteria led to the conclusion that the model with common parameters for all lines was acceptable (Fig. 4a and Table S1), with an added error lower than the 5% threshold compared with models specific to each line (2.0% on average; Fig. 4b and Table S1). This indicated that differences between rice lines with regard to the response curve and  $T_{opt}$  corresponded, if they existed, to a shift lower than 1°C (Fig. S4). Lines originating from different ecological areas and from different species or subspecies therefore had a common temperature–rate relationship. There was no trend for an effect of breeding between lines that were traditional (Azucena, Moroberkan), developed during the green revolution (IR64) or recently released (Vandana, Apo).

In wheat, we compared several lines selected in cool or warm climates, tolerant or sensitive to heat stress and released from the 1940s until recently. As above, a common model applied to the whole dataset (Fig. 2), with no significant differences between lines, as shown by the statistical criteria comparing models (Fig. 4a), as well as by the low additional error obtained when considering that the three parameters were common to all lines ( $\Delta CV = 1.4\%$  on average; Fig. 4b and Table S1). It can be concluded that any difference between lines would involve a shift of  $< 1^\circ C$  (Fig. S4c,d). The response was also common for a barley line (*Hordeum vulgare* cv. Golden Promise), which, like wheat, belongs to the Triticeae tribe (Fig. 2q), with an increase in error of  $< 2\%$  when considering a unique model compared with separate models for each species (not shown).

#### The responses of developmental processes differed in 18 species with a trend between cool-/warm-adapted species

The variability of the temperature response was explored in 18 species (17 crop species comprising the 14 most cultivated species worldwide and *Arabidopsis thaliana*). Results were collected from either our own database or the literature, with 42 variables, 73 references and 122 experiments (Table 1). The variables analysed in this study involved different genotypes for each species and developmental processes as different as the expansion rates of several organs (pollen tube, roots, leaves and embryo), the reciprocal of durations of phases, the cell division rate and germination rate.

In order to select the model that best suited the comparison between species, the analysis was first restricted to four species in which we had access to well-controlled datasets obtained by our groups (dataset of Fig. 2 in maize, rice and wheat, and the *Arabidopsis thaliana* dataset presented in Parent *et al.*, 2010b). When comparing the eight models of Table 2, statistical indicators showed that the response curves differed between species, because a unique model for different species (M.0) was not acceptable ( $\Delta BIC = +40$  and  $\Delta CV = 25\%$ , far from the thresholds of  $+5$  and  $+5\%$ , respectively; Fig. 4c,d and Table S1). An *F*-test procedure resulted in a similar conclusion (Fig. S6). The model M.2b, with a common  $\alpha$  for all species, was the best model ( $\Delta BIC = -2.2$ ; Fig. 4c, Table S1, Fig. S4), with 0.1% of




additional error compared with the model with three parameters. The other models, in particular those assuming that  $T_0$  was common between species, were rejected. We therefore compared the 18 species using model M.2b, in which species differ by two parameters of Eqn 2, namely the characteristic temperature  $T_0$  and activation enthalpy  $\Delta H_A^\ddagger$ .

Response curves corresponding to the 18 studied species followed similar patterns with different parameters (Fig. 5). The model M.2b applied to all species, with  $r^2$  ranging from 0.92 to 0.98 (Table 3). Its two parameters,  $\Delta H_A^\ddagger$  and  $T_0$ , are presented for each species in Table 3 and in spreadsheet format in Table S2. They were independent (Fig. S7a), confirming the two degrees of freedom of this model. In the following paragraph, we have transformed them into two more intuitive characteristics of response curves for each species, namely the optimum

temperature  $T_{opt}$  for which rates are at a maximum (related to  $T_0$ ; Eqn 3; Figs S1b, S7b), and the range of temperature in which the considered rate is higher than 50% of the maximum rate (range<sub>50%</sub>), related to  $\Delta H_A^\ddagger$  (Figs 5, 6 and Table 3).

Species originating from cool areas, such as canola (*Brassica napus* L.) and cauliflower (*Brassica oleracea*), showed the lowest values for  $T_{opt}$  ( $23.1 \pm 0.7$  and  $21.6 \pm 1.2^\circ\text{C}$ ; Table 3 and Fig. 6). Species adapted to warm climates, such as pearl millet (*Pennisetum glaucum*), peanut (*Arachis hypogaea*), cotton (*Gossypium*), sorghum (*Sorghum bicolor*) and cowpea (*Vigna unguiculata*), showed high values of  $T_{opt}$  (from  $32.4 \pm 0.5$  to  $34.1 \pm 1.5^\circ\text{C}$ ). This climatic effect applied within families of species, for instance in Poaceae species, in which  $T_{opt}$  ranged from relatively low to high values in wheat and sorghum ( $27.7 \pm 0.2$  and  $34.1 \pm 1.5^\circ\text{C}$ , respectively).




**Fig. 5** Response curve to temperature of developmental processes in 18 species. Each variable is normalized by its corresponding value at  $20^\circ\text{C}$ . Lines represent models for the considered species (model 2b). The origin of the data is given in Table 1. Model parameters are shown in Table 3 and Supporting Information Table S2.

**Table 3** Parameter values and quality of regression for the 18 studied species (model M.2b)

Species	Model M.2b					
	$R^2$	$\Delta H_A^\ddagger$ (kJ mol <sup>-1</sup> )	$T_0$ (°C)	$T_{opt}$ (°C)	Range <sub>50%</sub> (°C)	Rate <sub>max</sub>
<i>Arabidopsis thaliana</i>	0.98	65.1 ± 3.5	29.3 ± 0.4	26.5 ± 0.5	15.6–34.7	1.35
Barley	0.95	65.1 ± 3.5	29.4 ± 0.9	26.5 ± 0.9	15.6–34.7	1.36
Canola	0.94	80 ± 6	25.4 ± 0.8	23.1 ± 0.7	15.0–29.6	1.13
Cassava	0.95	89.6 ± 5.5	32 ± 0.6	29.9 ± 0.6	21.6–35.9	2.43
Cauliflower	0.98	63.6 ± 4.3	24.4 ± 1.2	21.6 ± 1.2	15.0–29.7	1.02
Common bean	0.96	61.7 ± 3.8	33.8 ± 1.2	30.7 ± 1	18.9–39.6	1.81
Cotton	0.95	70.3 ± 3.7	35.4 ± 0.9	32.6 ± 0.7	22.1–40.5	2.44
Cowpea	0.93	60 ± 3.5	36.3 ± 1.1	33.1 ± 0.9	20.8–42.4	2.13
Maize	0.98	73.9 ± 2.1	33.4 ± 0.2	30.8 ± 0.2	20.8–38.2	2.17
Millet	0.92	73.6 ± 4.3	35.1 ± 0.5	32.4 ± 0.5	22.3–39.9	2.53
Peanut	0.97	73.9 ± 2.9	35.3 ± 0.4	32.7 ± 0.3	22.6–40.2	2.61
Potato	0.97	81.7 ± 4	33 ± 1.4	30.6 ± 0.9	21.6–37.3	2.38
Rice	0.99	78.9 ± 3.1	33.4 ± 0.5	30.9 ± 0.5	21.6–37.8	2.36
Sorghum	0.98	60.6 ± 6	37.3 ± 1.7	34.1 ± 1.5	21.8–43.3	2.32
Soybean	0.97	52.3 ± 4.7	35.8 ± 1.2	32.2 ± 1	18.3–42.9	1.76
Sugarcane	0.98	75.2 ± 6	34.2 ± 1.2	31.6 ± 1.3	21.7–38.9	2.39
Sunflower	0.94	60.4 ± 5.8	32.4 ± 1.1	29.3 ± 0.9	17.3–38.3	1.60
Wheat	0.97	63.3 ± 3.6	30.6 ± 0.3	27.7 ± 0.2	16.4–36.2	1.46

$R^2$ , coefficient of determination of the regression between fitted and observed values;  $\Delta H_A^\ddagger$ ,  $T_0$  and  $\alpha$  are the three parameters of Eqn 2. In model M.2b,  $\alpha$  was common to all species ( $\alpha = 3.5$ ).  $T_{opt}$  (optimum temperature) is the temperature at which the development rate is at a maximum. Range<sub>50%</sub> is the temperature range within which the development rate reaches 50% of its maximum value. Rate<sub>max</sub> is the maximum rate in proportion to the value at 20°C. Parameter values are given ± SE calculated by bootstrapping.



**Fig. 6** Temperature at which rates are at a maximum ( $T_{opt}$ , black dots) and range of temperature for which the rate of development is at least 50% of its maximum (horizontal bars) in 18 species. Species are ranked by values of  $T_{opt}$ , also representing a gradient from temperate to tropical species.

## Discussion

### Coordination between developmental processes and absence of acclimation of temperature responses

The results presented here confirm and extend to a larger range of processes and species those observations recorded earlier on the coordination of developmental processes via a common

response curve to temperature (Ong, 1983a,b; Warrington & Kanemasu, 1983a,b; Parent *et al.*, 2010b). By contrast, photosynthesis and enzyme activities have response curves that differ markedly from those presented here (Campbell *et al.*, 2007; Sage *et al.*, 2008; Parent *et al.*, 2010b). Furthermore, they have nonunique temperature responses because they acclimate to changes in temperature within a few days (Sage & Kubien, 2007).

The absence of acclimation of the temperature response of developmental processes is not in contradiction with plant adaptation to low or high temperatures. Marked differences in the duration of the plant cycle (e.g. number of days after sowing at a given temperature) are observed following selection over several generations in plants, but without modification of the temperature response of this duration (Craufurd *et al.*, 1999). We show here that the response to temperature of these durations was common to all lines within a cultivated species. Morphological or anatomical adaptations to low or high temperature occur over periods of weeks (Gorsuch *et al.*, 2010). For example, changes in anatomy are still small after 6 d at 5°C, are noticeable after 30 d and become dramatic in new leaves developing under cold temperatures (Gorsuch *et al.*, 2010). The same conclusion applies to trees grown in different environments (Rehfeldt *et al.*, 2002).

Morphological changes linked to temperature may be an emerging property generated by the difference in behaviour between metabolic processes, with changes in response to temperature after acclimation, and developmental processes which do not acclimate. This would cause changes in carbon accumulation and specific leaf area (Tardieu *et al.*, 1999), observed experimentally by several groups (Loveys *et al.*, 2002; Shahba & Bauerle, 2009; Atkinson *et al.*, 2010; Fajardo & Piper, 2011).

A similar view has been proposed recently for the response to water deficit (Tardieu *et al.*, 2011).

### An absence of genetic variability of responses within the most common species

A different choice of lines within each species is unlikely to have yielded a different conclusion with regard to the genetic variability within species. In maize, the set of studied lines included tropical lines at very large genetic distances from the temperate lines (Camus-Kulandaivelu *et al.*, 2006). Data originating from the literature confirm this conclusion (Fig. 3b). Furthermore, no quantitative trait loci were detected for the temperature response of LER over a shorter range of temperature in three mapping populations (Reymond *et al.*, 2003; Sadok *et al.*, 2007; Welcker *et al.*, 2007), confirming the absence of a structured genetic variability. In the same way, the diversity tested in rice was large, with different species or subspecies, origins and generations of selection. The same result applied to wheat lines with different stress tolerances, suggesting that differences in heat tolerance involve the lowest and highest temperatures for which response curves can be established, but not the response curves themselves. It is noteworthy that, in the panels of lines tested here or in mapping populations derived from them, other traits showed a large genetic variability and clear quantitative trait loci (Reymond *et al.*, 2003; Chardon *et al.*, 2004; Izanloo *et al.*, 2008; Parent *et al.*, 2010a; Welcker *et al.*, 2011). Hence, the lack of genetic variability on the temperature response observed here was not a consequence of a poor choice of lines. It was probably not the result of a low power of statistical tests either, because our sensitivity analysis showed that we could identify a shift of  $< 1^{\circ}\text{C}$  when comparing response curves of lines or species. Finally, the conclusion with regard to the absence of genetic variability can be extended to the other crop species presented in Table 1, which were analysed on at least three genotypes each (except barley, two genotypes).

### A shift in temperature response probably requires millennia under very contrasting climates

The above paragraph shows that breeding in climates involving either high or low temperature has not affected the response of developmental processes to temperature, in particular  $T_{\text{opt}}$ . Centuries of mutations and selection are therefore not sufficient to affect this response within each species, contrary to most other traits that have responded to breeding. We therefore propose that a shift in temperature response requires two conditions. First, it requires millennia, i.e. at least the period during which most species have been bred. The second proposed condition is a marked contrast of climate. *Triticum* and *Hordeum* genera that genetically diverged more than 10 million yr (My) ago (Feuillet *et al.*, 2008) still have response curves with common parameters, probably because they grew under similar climatic conditions. This may apply to rice species and subspecies that have diverged for a longer period than that of agriculture (*c.* 11 000 years). No significant difference was observed between the subspecies *indica*

and *japonica*, which diverged 0.4 My ago, or between the species *O. sativa* and *O. glaberrima*, which diverged 0.6 My ago (Ma & Bennetzen, 2004). This is probably also the case for sorghum and sugarcane, which diverged 5 My ago (Ming *et al.*, 1998), and sorghum and maize, which diverged 10 My ago (Ramakrishna *et al.*, 2002). Differences were only observed in the case of very ancient divergence and very different climates, as in the contrast between Triticeae species and tropical grasses, such as sorghum, maize, millet or sugarcane, which diverged 50 My ago (Gale & Devos, 1998) and grew for a long period in different climates.

### Hypotheses for very slow evolution

A first hypothesis is that the evolution of temperature responses has been slowed down by the necessity for a synchronous shift in the responses of several processes. What would happen if a mutation changed the temperature response of one developmental process but not of the others? Simple computer simulations (not shown) suggest that a plant which loses the coordination between developmental processes because of a mutation would have highly disturbed performances, with, for example, a reproductive development occurring too early in comparison with vegetative development, or a disturbed aspect because of the lack of coordination between processes involved in the control of growth. We can therefore hypothesize that natural and breeder's selection has eliminated mutations that affect temperature responses of one process only, thereby requiring a very long time to obtain the response of all processes to shift simultaneously.

A second hypothesis is based on the fact that even the diversity found in the 18 tested species is low when compared with the six-fold differences in activation enthalpy found in different traits and life kingdoms (Dell *et al.*, 2011). The theory of 'thermodynamic constraint' (or 'warmer is better'), developed in animal studies (Huey & Kingsolver, 1989; Frazier *et al.*, 2006), argues that low temperatures slow down all biochemical reactions and that adaptation cannot change this rule. Because maximum rates occur at low temperature in cold-adapted plant species, such species would have lower maximum rates and performances than warm-adapted species, for which maximum rates occur at higher temperature, with higher rates of all biochemical reactions. By focusing mainly on plant performance, breeding may have selected lines and species with high optimum temperatures regardless of the ecological area. In our results,  $T_{\text{opt}}$  was generally higher than the mean temperature during the crop cycle. Eleven of the 17 studied crop species had optimum temperatures above  $30^{\circ}\text{C}$ , higher than those commonly observed in their growing regions. In these species, the temperature response may have reached a final equilibrium between a broad adaptation to a range of climate and maximum growth.

### Consequences for breeding and modelling

Our study suggests that there is little chance that breeding can substantially affect the temperature responses of the developmental rates analysed here, because of a nonexistent source of genetic

variability in crop species. This does not rule out several possibilities related to genetic adaptations of other traits related to temperature responses. First, the duration of the plant cycle has a considerable genetic variability, in particular between tropical and temperate materials. We therefore argue here that, although durations or rates have a large genetic variability, the change with temperature in these durations or rates does not vary between tropical and temperate lines. One way to avoid a decrease in yield with increasing temperature is to use lines with a longer crop cycle (Lobell *et al.*, 2008). Manipulating the respective durations of crop phases would also improve the yield potential in cereals (Borràs *et al.*, 2010). Second, a genetic gain has been obtained by breeding for the range of temperatures in which plant metabolism is not damaged under prolonged high or low temperature. For instance, the wheat line Excalibur used here tolerates high temperatures that cannot be tolerated by Kukri (Bukovnik *et al.*, 2009). This implies that the responses observed here, which apply to nonstressed ranges, are largely independent of tolerance to cold or heat stress. Third, high temperatures usually occur together with high evaporative demand, which has a considerable effect on gas exchange and growth, with a large genetic variability of this effect (Sadok *et al.*, 2007; Welcker *et al.*, 2007).

Our results also simplify the modelling of the effects of global changes, because lines grown in 50 or 100 yr will probably have a response to temperature similar to that of current lines. Existing crop models (Brisson *et al.*, 2003; Hammer *et al.*, 2010) already implicitly assume a common response of developmental processes, which is confirmed here. The response to temperature of a large range of species can be characterized by two traits that are relatively easy to measure, namely  $T_{\text{opt}}$  and the ratio between the rate at  $T_{\text{opt}}$  and that at 20°C. The parameters of Eqn 2 can be derived from these two traits (Eqn 3). Furthermore, because of the similar values of species in a given ecological area or not too distant genetically, modelling can be performed using the values of related species.

## Acknowledgements

This work has been funded in part by the project DROPS (DROught tolerant yielding PlantS; <http://www.drops-project.eu>), which received funding from the European Community's Seventh Framework Programme under the grant agreement no FP7-244374.

## References

- Addae PC, Pearson CJ. 1992. Thermal requirements for germination and seedling growth of wheat. *Australian Journal of Agricultural Research* 43: 585–594.
- Akaike H. 1974. New look at statistical-model identification. *Ieee Transactions on Automatic Control* AC19: 716–723.
- Akaike H. 1978. Likelihood of a time-series model. *Statistician* 27: 217–235.
- Akparobi SO, Togun AO, Ekanayake IJ. 2000. Temperature effects on leaf growth of cassava (*Manihot esculenta* Crantz) in controlled environments. *African Journal of Root and Tuber Crops* 4: 1–4.
- Arndt CH. 1945. Temperature–growth relations of the roots and hypocotyls of cotton seedlings. *Plant Physiology* 20: 200–220.
- Atkin OK, Loveys BR, Atkinson LJ, Pons TL. 2006. Phenotypic plasticity and growth temperature: understanding interspecific variability. *Journal of Experimental Botany* 57: 267–281.
- Atkinson LJ, Campbell D, Zaragoza-Castells J, Hurry V, Atkin OK. 2010. Impact of growth temperature on scaling relationships linking photosynthetic metabolism to leaf functional traits. *Functional Ecology* 2010: 1181–1191.
- Barlow EWR, Boersma L. 1972. Growth response of corn to changes in root temperature and soil-water suction measured with an LVDT. *Crop Science* 12: 251.
- Barnabas B, Jager K, Feher A. 2008. The effect of drought and heat stress on reproductive processes in cereals. *Plant, Cell & Environment* 31: 11–38.
- Battisti DS, Naylor RL. 2009. Historical warnings of future food insecurity with unprecedented seasonal heat. *Science* 323: 240–244.
- Ben-Haj-Salah H, Tardieu F. 1995. Temperature affects expansion rate of maize leaves without change in spatial distribution of cell length: analysis of the coordination between cell division and cell expansion. *Plant Physiology* 109: 870.
- Ben-Haj-Salah H, Tardieu F. 1997. Control of leaf expansion rate of droughted maize plants under fluctuating evaporative demand: a superposition of hydraulic and chemical messages? *Plant Physiology* 114: 893–900.
- Benoit GR, Grant WJ, Devine OJ. 1986. Potato top growth as influenced by day–night temperature differences. *Agronomy Journal* 78: 264–269.
- Blacklow WM. 1972. Influence of temperature on germination and elongation of radicle and shoot of corn (*Zea mays* L.). *Crop Science* 12: 647.
- Borràs G, Slafer GA, Casas AM, van Eeuwijk F, Romagosa I. 2010. Genetic control of pre-heading phases and other traits related to development in a double haploid barley (*Hordeum vulgare* L.) population. *Field Crops Research* 119: 36–47.
- Brisson N, Gary C, Justes E, Roche R, Mary B, Ripoche D, Zimmer D, Sierra J, Bertuzzi P, Burger P *et al.* 2003. An overview of the crop model STICS. *European Journal of Agronomy* 18: 309–332.
- Bukovnik U, Fu JM, Bennett M, Prasad PVV, Ristic Z. 2009. Heat tolerance and expression of protein synthesis elongation factors, EF-Tu and EF-1 alpha, in spring wheat. *Functional Plant Biology* 36: 234–241.
- Campbell C, Atkinson L, Zaragoza-Castells J, Lundmark M, Atkin O, Hurry V. 2007. Acclimation of photosynthesis and respiration is asynchronous in response to changes in temperature regardless of plant functional group. *New Phytologist* 176: 375–389.
- Campbell JA, Robertson MJ, Grof CPL. 1998. Temperature effects on node appearance in sugarcane. *Australian Journal of Plant Physiology* 25: 815–818.
- Camus-Kulandaivelu L, Veyrieras JB, Madur D, Combes V, Fourmann M, Barraud S, Dubreuil P, Gouesnard B, Manicacci D, Charcosset A. 2006. Maize adaptation to temperate climate: relationship between population structure and polymorphism in the *Dwarf8* gene. *Genetics* 172: 2449–2463.
- Chardon F, Virlon B, Moreau L, Falque M, Joets J, Decousset L, Murigneux A, Charcosset A. 2004. Genetic architecture of flowering time in maize as inferred from quantitative trait loci meta-analysis and synteny conservation with the rice genome. *Genetics* 168: 2169–2185.
- Chimenti CA, Hall AJ, Lopez MS. 2001. Embryo-growth rate and duration in sunflower as affected by temperature. *Field Crops Research* 69: 81–88.
- Covell S, Ellis RH, Roberts EH, Summerfield RJ. 1986. The influence of temperature on seed-germination rate in grain legumes. 1. A comparison of chickpea, lentil, soybean and cowpea at constant temperatures. *Journal of Experimental Botany* 37: 705–715.
- Craufurd PQ, Ellis RH, Summerfield RJ, Menin L. 1996a. Development in cowpea (*Vigna unguiculata*). 1. The influence of temperature on seed germination and seedling emergence. *Experimental Agriculture* 32: 1–12.
- Craufurd PQ, Mahalakshmi V, Bidinger FR, Mukuru SZ, Chantreau J, Omanga PA, Qi A, Roberts EH, Ellis RH, Summerfield RJ *et al.* 1999. Adaptation of sorghum: characterisation of genotypic flowering responses to temperature and photoperiod. *Theoretical and Applied Genetics* 99: 900–911.
- Craufurd PQ, Qi A, Ellis RH, Summerfield RJ, Roberts EH. 1996b. Development in cowpea (*Vigna unguiculata*). 2. Effect of temperature and saturation deficit on time to flowering in photoperiod-insensitive genotypes. *Experimental Agriculture* 32: 13–28.
- Craufurd PQ, Qi A, Summerfield RJ, Ellis RH, Roberts EH. 1996c. Development in cowpea (*Vigna unguiculata*). 3. Effects of temperature and



- photoperiod on time to flowering in photoperiod-sensitive genotypes and screening for photothermal responses. *Experimental Agriculture* 32: 29–40.
- Craufurd PQ, Subedi M, Summerfield RJ. 1997. Leaf appearance in cowpea: effects of temperature and photoperiod. *Crop Science* 37: 167–172.
- Cutforth HW, Jame YW, Jefferson PG. 1992. Effect of temperature, vernalisation, and water-stress on phyllochron and final main-stem leaf number of HY320 and Neepawa spring wheats. *Canadian Journal of Plant Science* 72: 1141–1151.
- Dejong R, Best KF. 1979. Effects of soil water potential, temperature and seeding depth on seedlings emergence of wheat. *Canadian Journal of Soil Science* 59: 259–264.
- Dell AI, Pawar S, Savage VM. 2011. Systematic variation in the temperature dependence of physiological and ecological traits. *Proceedings of the National Academy of Sciences, USA* 108: 10591–10596.
- Ellis RH, Hong TD, Roberts EH. 1987. Comparison of cumulative germination and rate of germination of dormant and aged barley seed lots at different constant temperatures. *Seed Science and Technology* 15: 717–727.
- Fajardo A, Piper FI. 2011. Intraspecific trait variation and covariation in a widespread tree species (*Nothofagus pumilio*) in southern Chile. *New Phytologist* 189: 259–271.
- Farquhar GD, Caemmerer Sv, Berry JA. 1980. A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C3 species. *Planta* 149: 78–90.
- Feng YS, Li XM, Boersma L. 1990. The Arrhenius equation as a model for explaining plant-responses to temperature and water stresses. *Annals of Botany* 66: 237–244.
- Feuillet C, Langridge P, Waugh R. 2008. Cereal breeding takes a walk on the wild side. *Trends in Genetics* 24: 24–32.
- Fleisher DH, Shillito RM, Timlin DJ, Kim SH, Reddy VR. 2006. Approaches to modeling potato leaf appearance rate. *Agronomy Journal* 98: 522–528.
- Fleisher DH, Timlin D. 2006. Modeling expansion of individual leaves in the potato canopy. *Agricultural and Forest Meteorology* 139: 84–93.
- Frazier MR, Huey RB, Berrigan D. 2006. Thermodynamics constrains the evolution of insect population growth rates: “warmer is better”. *The American Naturalist* 168: 512–520.
- Gale MD, Devos KM. 1998. Comparative genetics in the grasses. *Proceedings of the National Academy of Sciences, USA* 95: 1971–1974.
- Gallagher JN, Biscoe PV. 1979. Field studies of cereal leaf growth. 3. Barley leaf extension in relation to temperature, irradiance and water potential. *Journal of Experimental Botany* 30: 645–655.
- Gallagher JN, Biscoe PV, Wallace JS. 1979. Field studies of cereal leaf growth. 4. Winter-wheat leaf extension in relation to temperature and leaf water status. *Journal of Experimental Botany* 30: 657–668.
- Garciahuidobro J, Monteith JL, Squire GR. 1982. Time, temperature and germination of pearl-millet (*Pennisetum-typoides* S. and H.). 1. Constant temperature. *Journal of Experimental Botany* 33: 288–296.
- Gay C, Corbineau F, Come D. 1991. Effects of temperature and oxygen on seed-germination and seedling growth in sunflower (*Helianthus-annuus* L.). *Environmental and Experimental Botany* 31: 193–200.
- Gillooly JF, Brown JH, West GB, Savage VM, Charnov EL. 2001. Effects of size and temperature on metabolic rate. *Science* 293: 2248–2251.
- Gorsuch PA, Pandey S, Atkin OK. 2010. Temporal heterogeneity of cold acclimation phenotypes in *Arabidopsis* leaves. *Plant, Cell & Environment* 33: 244–258.
- Granier C, Massonnet C, Turc O, Muller B, Chenu K, Tardieu F. 2002. Individual leaf development in *Arabidopsis thaliana*: a stable thermal-time-based programme. *Annals of Botany* 89: 595–604.
- Hadley P, Roberts EH, Summerfield RJ, Minchin FR. 1983. A quantitative model of reproductive development in cowpea (*Vigna-unguiculata* (L) Walp.) in relation to photoperiod and temperature, and implications for screening germplasm. *Annals of Botany* 51: 531–543.
- Hammer GL, van Oosterom E, McLean G, Chapman SC, Broad I, Harland P, Muchow RC. 2010. Adapting APSIM to model the physiology and genetics of complex adaptive traits in field crops. *Journal of Experimental Botany* 61: 2185–2202.
- Hesketh JD, Duncan WG, Baker DN. 1972. Simulation of growth and yield in cotton. 2. Environmental conditions of morphogenesis. *Crop Science* 12: 436–439.
- Hesketh JD, Myhre DL, Willey CR. 1973. Temperature control of time intervals between vegetative and reproductive events in soybeans. *Crop Science* 13: 250–254.
- Hesketh JD, Warrington IJ. 1989. Corn growth-response to temperature: rate and duration of leaf emergence. *Agronomy Journal* 81: 696–701.
- Huey RB, Kingsolver JG. 1989. Evolution of thermal sensitivity of exotherm performance. *Trends in Ecology and Evolution* 4: 131–135.
- Inmanbamber NG. 1994. Temperature and seasonal effects on canopy development and light interception of sugarcane. *Field Crops Research* 36: 41–51.
- Izanloo A, Condon AG, Langridge P, Tester M, Schnurbusch T. 2008. Different mechanisms of adaptation to cyclic water stress in two South Australian bread wheat cultivars. *Journal of Experimental Botany* 59: 3327–3346.
- Johnson FH, Eyring H, Williams RW. 1942. The nature of enzyme inhibitions in bacterial luminescence: sulfanilamide, urethane, temperature and pressure. *Journal of Cellular and Comparative Physiology* 20: 247–268.
- Johnson FH, Lewin I. 1946. The growth rate of *E. coli* in relation to temperature, quinine, and coenzyme. *Journal of Cellular and Comparative Physiology* 28: 47–75.
- Kakani VG, Prasad PVV, Craufurd PQ, Wheeler TR. 2002. Response of in vitro pollen germination and pollen tube growth of groundnut (*Arachis hypogaea* L.) genotypes to temperature. *Plant, Cell & Environment* 25: 1651–1661.
- Keating BA, Evenson JP. 1979. Effect of soil-temperature on sprouting and sprout elongation of stem cuttings of cassava (*Manihot esculenta* Crantz.). *Field Crops Research* 2: 241–251.
- Kemp DR, Blacklow WM. 1982. The responsiveness to temperature of the extension rates of leaves of wheat growing in the field under different levels of nitrogen fertiliser. *Journal of Experimental Botany* 33: 29–36.
- Kirk WW, Marshall B. 1992. The influence of temperature on leaf development and growth in potatoes in controlled environments. *Annals of Applied Biology* 120: 511–525.
- Koumura T. 1972. Breeding of new rice variety ‘Nipponbare’. *Agricultural Technology* 27: 112–116.
- Kratsch HA, Wise RR. 2000. The ultrastructure of chilling stress. *Plant, Cell & Environment* 23: 337–350.
- Lafarge T, de Raissac M, Tardieu F. 1998. Elongation rate of sorghum leaves has a common response to meristem temperature in diverse African and European environmental conditions. *Field Crops Research* 58: 69–79.
- Lafond GP, Baker RJ. 1986. Effects of temperature, moisture stress, and seed size on germination of 9 spring wheat cultivars. *Crop Science* 26: 563–567.
- Lafond GP, Fowler DB. 1989. Soil-temperature and moisture stress effects on kernel water-uptake and germination of winter-wheat. *Agronomy Journal* 81: 447–450.
- Lawlor DJ, Kanemasu ET, Albrecht WC, Johnson DE. 1990. Seed production environment influence on the base temperature for growth of sorghum genotypes. *Agronomy Journal* 82: 643–647.
- Lawn RJ, Hume DJ. 1985. Response of tropical and temperate soybean genotypes to temperature during early reproductive growth. *Crop Science* 25: 137–142.
- Lehenbauer PA. 1914. Growth of maize seedlings in relation to temperature. *Physiological Researches* 1: 247.
- Lindstrom MJ, Papendick RI, Koehler FE. 1976. Model to predict winter-wheat emergence as affected by soil temperature, water potential and depth of planting. *Agronomy Journal* 68: 137–141.
- Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL. 2008. Prioritizing climate change adaptation needs for food security in 2030. *Science* 319: 607–610.
- Lobell DB, Schlenker W, Costa-Roberts J. 2011. Climate trends and global crop production since 1980. *Science* 333: 616–620.
- Loveys BR, Scurwater T, Pons TL, Fitter AH, Atkin OK. 2002. Growth temperature influences the underlying components of relative growth rate: an investigation using inherently fast- and slow-growing plant species. *Plant, Cell & Environment* 25: 975–987.

- Ma JX, Bennetzen JL. 2004. Rapid recent growth and divergence of rice nuclear genomes. *Proceedings of the National Academy of Sciences, USA* 101: 12404–12410.
- Marshall B, Squire GR. 1996. Non-linearity in rate–temperature relations of germination in oilseed rape. *Journal of Experimental Botany* 47: 1369–1375.
- McClung CR, Davis SJ. 2010. Ambient thermometers in plants: from physiological outputs towards mechanisms of thermal sensing. *Current Biology* 20: 1086–1092.
- Ming R, Liu SC, Lin YR, da Silva J, Wilson W, Braga D, van Deynze A, Wenslaff TF, Wu KK, Moore PH *et al.* 1998. Detailed alignment of *Saccharum* and *Sorghum* chromosomes: comparative organization of closely related diploid and polyploid genomes. *Genetics* 150: 1663–1682.
- Mohamed HA, Clark JA, Ong CK. 1988a. Genotypic differences in the temperature responses of tropical crops. 1. Germination characteristics of groundnut (*Arachis hypogaea* L.) and pearl-millet (*Pennisetum typhoides* S. & H.). *Journal of Experimental Botany* 39: 1121–1128.
- Mohamed HA, Clark JA, Ong CK. 1988b. Genotypic differences in the temperature responses of tropical crops. 2. Seedling emergence and leaf growth of groundnut (*Arachis hypogaea* L.) and pearl-millet (*Pennisetum typhoides* S. & H.). *Journal of Experimental Botany* 39: 1129–1135.
- Mongelar JC, Mimura L. 1972. Growth studies of sugarcane plant. 2. Some effects of root temperature and gibberellic-acid and their interactions on growth. *Crop Science* 12: 52.
- Morrison MJ, McVetty PBE, Shaykewich CF. 1989. The determination and verification of a baseline temperature for the growth of Westar summer rape. *Canadian Journal of Plant Science* 69: 455–464.
- Moss GI, Mullett JH. 1982. Potassium release and seed vigor in germinating bean (*Phaseolus vulgaris* L.) seed as influenced by temperature over the previous 5 generations. *Journal of Experimental Botany* 33: 1147–1160.
- Mutsaers HJW. 1983. Leaf growth in cotton (*Gossypium hirsutum* L.). 2. The influence of temperature, light, water-stress and root restriction on the growth and initiation of leaves. *Annals of Botany* 51: 521–529.
- Olesen JE, Grevsen K. 1997. Effects of temperature and irradiance on vegetative growth of cauliflower (*Brassica oleracea* L. botrytis) and broccoli (*Brassica oleracea* L. italica). *Journal of Experimental Botany* 48: 1591–1598.
- Ong CK. 1983a. Responses to temperature in a stand of pearl-millet (*Pennisetum typhoides* S. & H.). 1. Vegetative development. *Journal of Experimental Botany* 34: 322–336.
- Ong CK. 1983b. Responses to temperature in a stand of pearl-millet (*Pennisetum typhoides* S. & H.). 1. Extension of individual leaves. *Journal of Experimental Botany* 34: 1731–1739.
- Ong CK, Monteith JL. 1985. Response of pearl-millet to light and temperature. *Field Crops Research* 11: 141–160.
- Orbovic V, Poff K. 2007. Effect of temperature on growth and phototropism of *Arabidopsis thaliana* seedlings. *Journal of Plant Growth Regulation* 26: 222–228.
- Parent B, Conejero G, Tardieu F. 2009. Spatial and temporal analysis of non-steady elongation of rice leaves. *Plant, Cell & Environment* 32: 1561–1572.
- Parent B, Suard B, Serraj R, Tardieu F. 2010a. Rice leaf growth and water potential are resilient to evaporative demand and soil water deficit once the effects of root system are neutralized. *Plant, Cell & Environment* 33: 1256–1267.
- Parent B, Turc O, Gibon Y, Stitt M, Tardieu F. 2010b. Modelling temperature-compensated physiological rates, based on the co-ordination of responses to temperature of developmental processes. *Journal of Experimental Botany* 61: 2057–2069.
- Pearson CJ. 1975. Thermal adaptation of *Pennisetum*: seedling development. *Australian Journal of Plant Physiology* 2: 413–424.
- Penfield S. 2008. Temperature perception and signal transduction in plants. *New Phytologist* 179: 615–628.
- Peng S, Jianliang H, John ES, Rebecca CL, Romeo MV, Xuhua Z, Grace SC, Gurdev SK, Kenneth GC. 2004. Rice yields decline with higher night temperature from global warming. *Proceedings of the National Academy of Sciences, USA* 101: 9971–9975.
- Poorter H, Niinemets U, Walter A, Fiorani F, Schurr U. 2010. A method to construct dose–response curves for a wide range of environmental factors and plant traits by means of a meta-analysis of phenotypic data. *Journal of Experimental Botany* 61: 2043–2055.
- Prasad PVV, Boote KJ, Thomas JMG, Allen LH, Gorbett DW. 2006. Influence of soil temperature on seedling emergence and early growth of peanut cultivars in field conditions. *Journal of Agronomy and Crop Science* 192: 168–177.
- R Development Core Team. 2005. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. ISBN 3-900051-07-0, <http://www.R-project.org>.
- Ramakrishna W, Dubcovsky J, Park YJ, Busso C, Emberton J, SanMiguel P, Bennetzen JL. 2002. Different types and rates of genome evolution detected by comparative sequence analysis of orthologous segments from four cereal genomes. *Genetics* 162: 1389–1400.
- Reddy KR, Hodges HF, McKinion JM. 1997. Modeling temperature effects on cotton internode and leaf growth. *Crop Science* 37: 503–509.
- Rehfeldt GE, Tchebakova JM, Parfenova YI, Wykoff WR, Kuzmina NA, Milyutin LI. 2002. Intraspecific response to climate in *Pinus sylvestris*. *Global Change Biology* 8: 1–18.
- Reymond M, Muller B, Leonardi A, Charcosset A, Tardieu F. 2003. Combining quantitative trait loci analysis and an ecophysiological model to analyze the genetic variability of the responses of maize leaf growth to temperature and water deficit. *Plant Physiology* 131: 664–675.
- Sadok W, Naudin P, Boussuge B, Muller B, Welcker C, Tardieu F. 2007. Leaf growth rate per unit thermal time follows QTL-dependent daily patterns in hundreds of maize lines under naturally fluctuating conditions. *Plant, Cell & Environment* 30: 135–146.
- Sage RF, Kubien DS. 2007. The temperature response of C3 and C4 photosynthesis. *Plant, Cell & Environment* 30: 1086–1106.
- Sage RF, Way DA, Kubien DS. 2008. Rubisco, Rubisco activase, and global climate change. *Journal of Experimental Botany* 59: 1581–1595.
- Schwarz G. 1978. Estimating the dimension of a model. *Annals of Statistics* 6: 461–464.
- Scully B, Waines JG. 1987. Germination and emergence response of common and tepary beans to controlled temperature. *Agronomy Journal* 79: 287–291.
- Shahba MA, Bauerle WL. 2009. Growth temperature modulates the spatial variability of leaf morphology and chemical elements within crowns of climatically divergent *Acer rubrum* genotypes. *Tree Physiology* 29: 869–877.
- Sharpe PJH, Demichele DW. 1977. Reaction-kinetics of poikilotherm development. *Journal of Theoretical Biology* 64: 649–670.
- Singh SK, Kakani VG, Brand D, Baldwin B, Reddy KR. 2008. Assessment of cold and heat tolerance of winter-grown canola (*Brassica napus* L.) cultivars by pollen-based parameters. *Journal of Agronomy and Crop Science* 194: 225–236.
- Summerfield RJ, Collinson ST, Ellis RH, Roberts EH, Devries F. 1992. Photothermal responses of flowering in rice (*Oryza sativa*). *Annals of Botany* 69: 101–112.
- Tardieu F, Granier C, Muller B. 1999. Modelling leaf expansion in a fluctuating environment: are changes in specific leaf area a consequence of changes in expansion rate? *New Phytologist* 143: 33–43.
- Tardieu F, Granier C, Muller B. 2011. Water deficit and growth. Co-ordinating processes without an orchestrator? *Current Opinion in Plant Biology* 14: 283–289.
- Tollenaar M, Daynard TB, Hunter RB. 1979. Effect of temperature on rate of leaf appearance and flowering date in maize. *Crop Science* 19: 363–366.
- Vigil MF, Anderson RL, Beard WE. 1997. Base temperature and growing-degree-hour requirements for the emergence of canola. *Crop Science* 37: 844–849.
- Villablobos FJ, Ritchie JT. 1992. The effect of temperature on leaf emergence rates of sunflower genotypes. *Field Crops Research* 29: 37–46.
- Wade LJ, Hammer GL, Davey MA. 1993. Response of germination to temperature amongst diverse sorghum hybrids. *Field Crops Research* 31: 295–308.
- Warrington IJ, Kanemasu ET. 1983a. Corn growth response to temperature and photoperiod. 1. Seedling emergence, tassel initiation, and anthesis. *Agronomy Journal* 75: 749–754.
- Warrington IJ, Kanemasu ET. 1983b. Corn growth response to temperature and photoperiod. 2. Leaf-initiation and leaf-appearance rates. *Agronomy Journal* 75: 755–761.

- Weaich K, Bristow KL, Cass A. 1996. Modeling preemergent maize shoot growth. 1. Physiological temperature conditions. *Agronomy Journal* **88**: 391–397.
- Welcker C, Bousuge B, Bencivenni C, Ribaut JM, Tardieu F. 2007. Are source and sink strengths genetically linked in maize plants subjected to water deficit? A QTL study of the responses of leaf growth and of anthesis–silking interval to water deficit. *Journal of Experimental Botany* **58**: 339–349.
- Welcker C, Sadok W, Dignat G, Renault M, Salvi S, Charcosset A, Tardieu F. 2011. A common genetic determinism for sensitivities to soil water deficit and evaporative demand: meta-analysis of quantitative trait loci and introgression lines of maize. *Plant Physiology* **157**: 718–729.
- White JW, Montes C. 1993. The influence of temperature on seed-germination in cultivars of common bean. *Journal of Experimental Botany* **44**: 1795–1800.
- Wise RR, McWilliam JR, Naylor AW. 1983. A comparative study of low-temperature-induced ultrastructural alteration of 3 species with different chilling sensitivities. *Plant, Cell & Environment* **6**: 525–535.
- Wurr DCE, Fellows JR, Sutherland RA, Elphinstone ED. 1990. A model of cauliflower curd growth to predict when curds reach a specified size. *Journal of Horticultural Science* **65**: 555–564.
- Yin XY, Kropff MJ. 1996. The effect of temperature on leaf appearance in rice. *Annals of Botany* **77**: 215–221.
- Yin XY, Kropff MJ, Goudriaan J. 1996. Differential effects of day and night temperature on development to flowering in rice. *Annals of Botany* **77**: 203–213.
- Yin XY, Struik PC, Kropff MJ. 2004. Role of crop physiology in predicting gene-to-phenotype relationships. *Trends in Plant Science* **9**: 426–432.

## Supporting Information

Additional supporting information may be found in the online version of this article.

**Fig. S1** Sensitivity analysis of the model M.3.

**Fig. S2** Data handling, example of genotype CML444.

**Fig. S3** Calculation of parameter values for models M.3, M.2b and M.0 in the maize hybrid ‘Dea’.

**Fig. S4** Analysis of the sensitivity of the information criterion ( $\Delta\text{BIC}$ , a,c) and supplemental error ( $\Delta\text{CV}$ , b,d) to changes in the temperature response. BIC, Bayesian information criterion; CV, coefficient of variation.

**Fig. S5** Effect of previous high or low temperature on leaf elongation rate at 20°C.

**Fig. S6** *F*-test between the different nested models in four species.

**Fig. S7** Correlation between model parameters and the optimum temperature in 18 species.

**Table S1** Goodness-of-fit evaluation of the different models considering one species (maize, rice or wheat) or four species

**Table S2** Spreadsheet with parameters and temperature response functions for the 17 crop species and *Arabidopsis thaliana*

**Methods S1** Calculation of optimum temperature from parameters of Eqn 2.

**Methods S2** Fitting version of Eqn 2 after normalization at 20°C.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.