

# Why longer seasons with climate change may not increase tree growth

E.M. Wolkovich<sup>1</sup>, Ailene K. Ettinger<sup>2</sup>, Alana Chin<sup>3</sup>, Catherine J. Chamberlain<sup>4</sup>,  
Frederik Baumgarten<sup>1</sup>, Kavya Pradhan<sup>5,6</sup>, Rubén D. Manzanedo<sup>7-9</sup> &  
Janneke Hille Ris Lambers<sup>7</sup>

February 2, 2025

<sup>1</sup> Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4, Canada

<sup>2</sup> The Nature Conservancy of Washington, 74 Wall Street, Seattle, WA USA

<sup>3</sup> California State Polytechnic University, Humboldt. Department of Biological Sciences. Arcata, CA, USA

<sup>4</sup> The Nature Conservancy, 334 Blackwell St Ste 300, Durham, NC, USA

<sup>5</sup> Department of Biology, University of Washington, Seattle, WA, 98195

<sup>6</sup> Moore Center for Science, Conservation International, Arlington, VA, 22202

<sup>7</sup> Institute for Integrative Biology, ETH Zürich, 8092 Zürich, Switzerland

<sup>8</sup> Institute of Plant Sciences, University of Bern, Bern, Switzerland

<sup>9</sup> Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland

*Author contributions:* All authors conceived of the idea, prepared the literature review, edited the manuscript and contributed to designing some of the figures; in addition EMW wrote the manuscript, AKE, AC, CJC and EMW synthesized the literature review, RD led developing Fig. 1, CJC developed Fig 2, EMW developed Fig. 4, and AC, RD and EMW led developing the Box figure and Fig. 3.

## Abstract

A number of recent studies have challenged the assumption that longer growing seasons lead to increased tree growth, raising concerns that forecasts of future climate change—which include increased carbon storage through this assumption—may be overly optimistic. In a review of recent literature, we found that 58% of studies supported the assumption of increased growth with longer seasons, while 36% of studies did not. Diverging results remained when holding methodology constant, which suggests the current major challenge is to understand what underlies this widespread variation. Studies have proposed a suite of hypotheses for why longer growing seasons may not always increase tree growth, including drought-related constraints and internal limits. These hypotheses and their underlying mechanisms, however, were generally tested in different ways by different fields on different species, making comparisons difficult. We outline how bridging these current divides while simultaneously integrating evolutionary history and ecological theory could yield new advances and build a unified model across species for when longer seasons will—or will not—lead to greater tree growth, with major forecasting implications.

## 1 Introduction

2 The idea that longer growing seasons lead to increased plant growth is an intuitive tenet across  
3 multiple fields of biology, including physiology, dendrochronology and ecosystem ecology (Nobel  
4 *et al.*, 1983; Frank *et al.*, 2022). It is also a foundational assumption of many global carbon  
5 cycle models (e.g. Ito *et al.*, 2020; Friedlingstein *et al.*, 2022). These models project that contin-  
6 ued anthropogenic warming will be partly offset by increased carbon sequestration as warming  
7 lengthens growing seasons in many forests (Friedlingstein *et al.*, 2022), an assumption supported  
8 by ecosystem-scale studies (Chen *et al.*, 1999; Keenan *et al.*, 2014; Finzi *et al.*, 2020).

9 Yet recent work has questioned this longstanding assumption (e.g. Dow *et al.*, 2022; Green  
10 & Keenan, 2022; Silvestro *et al.*, 2023), with potentially large implications for future climate  
11 change. These recent studies challenge decades of research reporting increased growth with  
12 longer seasons, from observations along elevational and latitudinal gradients (Myneni *et al.*, 1997;  
13 Berdanier & Klein, 2011; King *et al.*, 2013; Cuapio-Hernández *et al.*, 2022), classic experiments  
14 in lab settings (Went, 1957), to trends in ecosystem fluxes with warming (Chen *et al.*, 1999;  
15 Keenan *et al.*, 2014; Finzi *et al.*, 2020). Proposed mechanisms for the apparent disconnect are  
16 diverse (Fig. 1), including the complex nature of climate change (e.g. drought or heat stress,  
17 Dow *et al.*, 2022) and internal limits on plant growth (Zohner *et al.*, 2023).

18 Here we review how different fields have studied the relationship between growing season length  
19 and tree growth to identify the potential mechanisms that unite—and could disconnect—these  
20 processes. Working across multiple definitions (see Box: Growth, growing season length and the  
21 challenge of standardized metrics), we find results that suggest substantial variation in growth  
22  $\times$  season length relationships. We also find a pervasive disciplinary split between studies, which  
23 often test different mechanisms on different species. Current work often implicitly ignores the

importance of shared evolutionary history and community ecology to plant growth (e.g. Grime, 1977; Webb *et al.*, 2002; Ávila-Lovera *et al.*, 2023), which could aid the search for a universal model when studying different species. We show how increased cross-disciplinary efforts to build a model across species would allow the field to rapidly develop a framework to predict when, where and how climate change may increase tree growth.

## Evidence that longer seasons increase plant growth, or not

The idea that time limits growth is a fundamental principle across biology. Many biological processes—including photosynthesis and aspects of growth—are rate-limited, making time a crucial commodity (Nobel *et al.*, 1983; Cosgrove, 2005; Hilty *et al.*, 2021). Thus, the hypothesis that longer growing seasons should increase growth is intuitive—and pervasive.

Foundational evidence comes from spatial clines across elevation and latitude, with growth decreasing alongside growing season length at higher elevations and latitudes (Fig. S1). Experimentally, this assumption is supported by small-scale field warming studies that find that phenologically advancing species also grow more with warming (Cleland *et al.*, 2012), while observationally, ecosystem-scale studies have reported a similar positive relationship between season length and carbon fluxes across decades with global warming (Keenan *et al.*, 2014) or in years with warm, early springs (Chen *et al.*, 1999). However, some recent high-profile studies find no support for this relationship (Dow *et al.*, 2022). These studies, which often focus on inter-annual correlations with metrics of standardized individual tree growth (Dow *et al.*, 2022; Silvestro *et al.*, 2023), have generated debate about whether future carbon storage forecasts are overestimated and which metrics of growth (Green & Keenan, 2022), or growing season length (Körner *et al.*, 2023), are relevant (see Box).

Despite this recent debate, we found that longer seasons lead to increased growth in a slight majority of papers spanning 25 years. Though the number of papers directly addressing this topic is small (36 papers), we found studies have variously found evidence for—or against—the relationship, with no clear pattern by method or year (Fig. 2 and see ‘Literature review methods’ in Supplement). For example, carbon assimilation studies were evenly split in finding evidence for or against the relationship (or simply not testing it, Fig. 2). Diverging results were consistently found within methods, suggesting the drivers of this variation are likely due to biological mechanisms, not solely inconsistent definitions of growth or growing season length (as some have recently suggested, e.g. Green & Keenan, 2022; Körner *et al.*, 2023, and see Box).

Most studies tested the hypothesis that longer seasons with climate change increase growth via either increased time to grow (10 of 36 papers) or because longer seasons are usually warmer (8 papers), although many also considered hypotheses that could disconnect growth from season length. Studies from dendrochronology (the study of tree rings and their dating) and physiology have readily offered explanations for findings that increased growth may not be a universal

outcome of longer seasons (Fig. 1). External climatic drivers that offset the positive growth effects of longer seasons were often reported in tree ring studies (Kolář *et al.*, 2016; de Sauvage *et al.*, 2022; Camarero *et al.*, 2022). In particular, the hypothesis that higher temperatures paired with lower precipitation produce negative correlations of season length with growth appeared in 58% of tree ring studies we reviewed (and was only mentioned once outside of these studies, see also Fig. 1). In contrast, 43% of lab experimental and wood phenology (xylogenesis) studies suggested fundamental internal constraints that prevent trees from responding to longer seasons (Fig. S2, Cuny *et al.*, 2012; Michelot *et al.*, 2012; Zohner *et al.*, 2023). Yet we found that these hypotheses have been tested in radically different ways on different species, rarely together, and ignore a suite of relevant research from other disciplines.

## Controllers on growth $\times$ season length relationships

Major mechanisms that could limit or disrupt the positive effects of longer growing seasons generally fall into two categories: (1) external factors, such as drought, which should impact ecosystem-level trends at regional scales, and (2) internal physiological constraints, which some research suggests are either universal across plants (e.g. Zohner *et al.*, 2023), or species- and population-specific (e.g. Soolanayakanahally *et al.*, 2013). While we address each in turn, these drivers can clearly operate together. Further, the importance of internal versus external drivers likely varies by species, highlighting the need to integrate perspectives from community and phylogenetic ecology.

### External drivers

Temperature limits many biological processes. Temperatures that are too cool (below 5°C for temperate trees) and too warm (an area of active research, but likely between 35-45°C; Martinez-Meier *et al.*, 2008; Cabon *et al.*, 2022) slow down biological processes and eventually can lead to tissue death (see Fig. 3a, Larcher, 1980; Kramer, 2012). Between these upper and lower limits, biological processes underpinning growth generally accelerate such that warming can have a direct effect, by accelerating biological time, up until the maximum rate for that particular process. Assuming a common growth response curve to temperature, possible increased growth should be predictable based on the current seasonal temperatures and the amount of warming (Fig. 3b).

How much or whether growth increases at all depends on the non-linear effect of temperature on biological processes (Fig. 3a). At very cool temperatures—such as in early spring—a small increase in temperature may have limited effect (or even increase frost risk through early budburst, Fig. 1e, Chamberlain *et al.*, 2021), while an increase at warmer temperatures—such as those more common in the summer (e.g. 16 to 18°C)—could have a larger physiological impact. However, warming that pushes plants beyond their optima, where many biological rates crash, could have large negative impacts (Nobel *et al.*, 1983; Leuning, 2002). Thus, some studies

hypothesize that longer seasons effectively only extend the very cool early-season periods and may have no discernible effect on growth (with varying definitions of growth, see Box), while other studies—based on tree rings—suggest that any increases in growth due to longer seasons can be offset by reduced growth due to high summer temperatures (Fig. 1, Gantois, 2022; Dow *et al.*, 2022). In contrast, other researchers argue that warmer temperatures have not yet pushed trees above their optima (Schaber & Badeck, 2002), and instead have driven increases in growth through accelerated rates, rather than longer seasons (e.g. Ren *et al.*, 2019), or through a combination of both.

Positive effects of longer—or warmer—seasons on growth predicted from temperature responses alone, however, could be counteracted by other external drivers. Moisture deficits from reduced precipitation or higher evaporative demand (commonly invoked in tree ring studies, Fig. 1) can slow or stall growth. Support for this hypothesis comes from negative correlations between growth and precipitation (or other metrics related to plant access to water in tree ring studies, Kolář *et al.*, 2016; Etzold *et al.*, 2022), and is well supported by physiological observations that tree water status can be a biophysical limit to growth (i.e., cells cannot expand without sufficient turgor, Peters *et al.*, 2021; Cosgrove, 2023), though we found few physiological studies on season length that considered this effect (Fig. 2). External biotic factors are also shifting with longer seasons—including herbivory, disease and competition (Mitton & Ferrenberg, 2012; Lange *et al.*, 2006; Cleland & Wolkovich, 2024)—and can limit productivity (Sturrock *et al.*, 2011; La Porta *et al.*, 2008; Senf *et al.*, 2017), though they are missing from the current debate on the impacts of longer seasons on growth (we found no mention of them, Fig. 1e).

## Internal constraints

When and how growth is initiated and ceases is under genetic and developmental control, and thus plants’ internal programming could limit growth responses to longer seasons (Howe *et al.*, 2003). Research has repeatedly shown that populations vary in their growth and its responses to extended seasons (Fig. 1d), reflecting differences in genetic and developmental controls that likely evolved to limit tissue loss to rare early or late-season events (Mitton & Ferrenberg, 2012; Lange *et al.*, 2006; Cleland & Wolkovich, 2024). Populations often vary predictably in their end-of-season phenology, with more poleward populations tending to stop height growth (budset) earlier using locally adapted photoperiod cues (Soolanayakanahally *et al.*, 2013; Aitken & Bemmels, 2016). This means longer seasons are generally driven by spring phenology, which appears far more flexible, and has advanced more rapidly than fall events (Aitken & Bemmels, 2016). Some recent studies suggest novel roles for the summer solstice (Zohner *et al.*, 2023) in setting a fixed universal developmental switch between when warming temperatures hasten or delay leaf senescence, and in determining when warmer temperatures trigger greater reproduction (Journé *et al.*, 2024).

Trade-offs between vegetative and reproductive investments may produce important growth response differences across years within individuals, as well as between species. Years of high

reproductive output can reduce growth (Thomas, 2011; Hacket-Pain *et al.*, 2016a). For species that mast—producing abundant cones or fruits in only some years—high reproduction could especially impact measures of wood growth. Higher summer temperatures may trigger masting in the following year (Hacket-Pain *et al.*, 2016a,b); if true, then reduced growth in years following warm summers may not indicate temperatures too high for growth, as recent studies have suggested (e.g. Gantois, 2022; Dow *et al.*, 2022), but instead shifting investment to reproduction.

## Species-level variation

The effects of these external and internal drivers are likely to vary across species, with species identity strongly predicting variation in growth  $\times$  season length relationships (e.g. Cuny *et al.*, 2012; Michelot *et al.*, 2012). Though this reality was rarely acknowledged in studies we reviewed (Fig. 1c), research in dendrochronology, physiology and in phenology often mentions important differences between certain species groups that should affect how longer seasons affect growth.

The distinct strategies of deciduous versus evergreen species, including in how and when they invest in leaf and shoot elongation versus cambial growth, can affect how they respond to longer seasons. While evergreen species generally leaf out later than deciduous species they can more immediately photosynthesize with earlier springs, though both types of species generally invest in buds (for new leaves, shoots and flowers) in the preceding year. This means neither can rapidly change their investment in leaf area in response to an earlier spring, but both can have multiple flushes of leaves (Day & Greenwood, 2011; Soolanayakanahally *et al.*, 2013). Wood growth in evergreen species is generally thought to come from current season photosynthate, while deciduous species may more often use stored carbon resources (Gordon & Larson, 1968; Monson *et al.*, 2018). These differences would suggest season length by growth relationships may be most apparent via lagged effects in deciduous species, but this is rarely studied (and not clearly supported to date, see Coulthard *et al.*, 2020; Klesse *et al.*, 2023).

This division between evergreen and deciduous species hints at a larger suite of traits that predict growth by growing season length relationships among species. Species that budburst earlier and more readily produce additional leaves (e.g. leaf flushes after budset, and other characteristics more common to ‘indeterminate’ species, Kikuzawa, 1982; Lechowicz, 1984) may grow more with longer seasons (though potentially with a lag, see Box) versus those that budburst later and flush new primary growth only once. Similarly, species adapted to cold, dry or high latitude conditions across their range may have different thresholds for when these external drivers limit or promote growth (e.g. some *Populus* and *Quercus* species, Soolanayakanahally *et al.*, 2013; McKown *et al.*, 2016; Delpierre *et al.*, 2017; de Sauvage *et al.*, 2022, and see Fig.3). Such differences could easily obscure any overall relationship between growth and growing season length. Supporting this possibility, current studies finding divergent results (Fig. S3) span a wide range of species (we found 57 species from 26 genera across 36 papers). While this diversity may appear to make identifying a common relationship between growth and growing season length more difficult, it may instead offer the path to an improved framework.

## Building a new framework for growth $\times$ season length relationships

Useful models of tree growth for climate change forecasting must include a diversity of species, while overcoming the challenges of uneven sampling across species and their contrasting responses. Leveraging the diversity of responses observed across species is possible by integrating ecological advances in how species traits and evolutionary history shape responses to climate change (Cornwell & Nakagawa, 2017; Harmon, 2024). In particular, advances in phylogenetic comparative methods (Webb *et al.*, 2002) have moved research away from treating species identity as a simple grouping factor where each species is unique (e.g. *Fagus sylvatica* is different from *Quercus robur* and *Pinus sylvestris*) or fits into a limited set of groups (e.g. deciduous versus evergreen) and towards species as suites of correlated observations, separated by their evolutionary distance (e.g. *Fagus sylvatica* is much more closely related to *Quercus robur* compared to *Pinus sylvestris*). New models built from these advances can fit data from all species at once and layer in species-level information. Traits can capture differences in growth strategies while phylogeny can capture additional species differences, which likely capture unmeasured ‘latent’ traits (Pearse *et al.*, 2019; Morales-Castilla *et al.*, 2024).

In step with these advances, trait ecology has documented leaf and wood economic spectra that suggest major traits to include in these models (with related databases of these traits often available, Chave *et al.*, 2009; Diaz *et al.*, 2016). These ‘economics’ define a common trade-off along an acquisitive to conservative axis, where some species grow rapidly and more flexibly to take advantage of resources, but are less defended against herbivores and compete poorly at low resource levels, whereas other species compete well at low resource levels, but at the expense of growing slower (Grime, 1977; Chave *et al.*, 2009; Diaz *et al.*, 2016). While these traits likely miss critical components for understanding how growing season length shapes growth, such as when different species invest in shoot and leaf versus wood growth, they provide a baseline from which to build, and a powerful approach to combine data usefully across species. This approach has already been used to identify that early-leafout species often show faster-growing more acquisitive strategies compared to later-leafout species (reviewed in Cleland & Wolkovich, 2024)—differences that may also impact how they respond to longer seasons.

In addition to naturally organizing species differences, a trait-based phylogenetic comparative approach can help build a more testable and predictable framework. Because this approach can flexibly fit evolutionary history and traits together, it allows clades or species groupings that respond similarly to emerge from the data and models (Davies *et al.*, 2019), versus being a priori grouped or defined. Similarly traits that co-vary with different responses can be more quickly identified (e.g. Willis *et al.*, 2008; Davies *et al.*, 2019, see Fig. 4). Both of these benefits could highlight which species or traits to focus additional studies on to gain the most insights, while similarly suggesting areas that should be less studied (e.g. traits that may be too confounded with evolutionary history, Cornwell *et al.*, 2014; Westoby *et al.*, 2023) or outlier species that may not represent most species (Morales-Castilla *et al.*, 2024). This approach may thus redefine

debates over which metrics of growth or growing season length are relevant into debates over which metrics are most relevant for which clades and/or traits.

Importantly, trait-mediated phylogenetic models aid the search a universal model that can be useful for global forecasts of how growing season length influences growth. By effectively assuming one model, which is then shaped by evolutionary history and trait differences to produce the divergent responses observed across species today (Fig. 4), this approach yields an overall estimated growth response alongside a response for each species, which can then be variously combined to scale up. For example, species-level estimates combined with data on species abundance across forests (e.g. Forest Service - United States Department of Agriculture; Fischer *et al.*, 2019) could predict larger-scale metrics, such as satellite observations of phenology and productivity. This framework also provides an organizing method to re-approach the fundamental question of how external and internal drivers shape growth responses to longer seasons, and a way to tackle what we are argue the three most critical questions in this area.

## **What is the scale of variation in season length x growth relationships across space and time?**

The idea that growing season length influences plant growth is fundamental to plant biology, but we found it is rarely tested in ways relevant to the current debate (see ‘Growth  $\times$  elevation relationships’ in Supplement), providing a major gap that limits progress. While multiple papers report a lack of relationship between growth and growing season length (Figs. 1, 2), we have no fundamental understanding of what the effect size of this relationship should be, and thus no way to know if we have sufficient power in current studies to detect it.

Identifying the macro-scale pattern of how growth and growing season length relate should be a primary goal, and one that seems broadly tractable. Tree ring studies designed to leverage latitudinal and elevational gradients in climate could quickly provide the raw data. Research will then need to develop models that tease out the effects of warmer temperatures across the season—likely affecting important biological rates (Fig. 3)—versus longer seasons. Disentangling these may require focused efforts to understand xylogenesis across species and climates, but doing so across major climatic gradients could make differences more obvious. Wood growth provides an obvious and tractable baseline from which to set expectations of how much growth can vary across space, and links to existing major datasets (Fig. S4). Research will also then need to integrate beyond wood growth, including methods to better characterize changes across the leaf, shoot and wood architecture of different species (e.g. Puletti *et al.*, 2020; Sillett *et al.*, 2024) and also extending to the complexity of roots (McKown *et al.*, 2016; Radville *et al.*, 2016). These data can provide a baseline to compare to the scale of shifts over time, which studies of growth  $\times$  growing season length to date have focused on (Fig. 2), since the same tree rings measured for understanding spatial variation will also capture inter-annual variation.



## How prevalent are internal constraints across species and populations?

New evidence suggests inter-annual variation in growth may be limited because of internal constraints that prevent plants from fully using longer seasons (Zohner *et al.*, 2023). If true, this would have major ramifications for how much we expect growth to shift with warming. All plants are limited by internal constraints and how quickly they can build new tissues (Marchand *et al.*, 2021; Luo *et al.*, 2024), but selection towards different growth strategies (e.g. acquisitive versus conservative) should drive variation in these constraints across species. Selection should also drive local adaptation in these constraints at the population-level (McKown *et al.*, 2016; Soolanayakanahally *et al.*, 2013), by favoring individuals that match to local environmental optima (Colautti *et al.*, 2010; McKown *et al.*, 2014). This appears to be the case for budset—which indicates the end of height growth, though we currently have data on only a few species (Aitken & Bemmels, 2016; Zeng & Wolkovich, 2024). New studies could rapidly test for constraints across species and populations, to work towards a predictive framework using phylogeny and traits to predict these constraints. This approach has already yielded useful insights in spring phenology, highlighting which environmental factors consistently drive budburst across species while also showing widely-cited results may not extend beyond one well-studied species (Morales-Castilla *et al.*, 2024).

## How do external drivers and internal constraints act together?

The external and internal factors that affect how longer seasons impact growth are inherently interconnected (Nobel *et al.*, 1983). While research often acknowledges this, modeling these together will require both experiments and observational studies, ideally designed to integrate into trait-mediated phylogenetic models. Studies across space could provide some inference by studying how growing seasons measured by vegetative versus wood phenology vary—and attributing variation through models that nest species within populations and include traits while also testing for how climate drives growth.

The complexity of climate change and plant growth in response to longer, warmer seasons makes experiments vital to building useful mechanistic models for forecasting. Observational data—used mainly to date to tackle this question (Figs. 2, S2)—generally confounds multiple external drivers, including season length, temperature and precipitation regimes (Ren *et al.*, 2019; Canadell *et al.*, 2021; Camarero *et al.*, 2022), making it impossible to tease out actual drivers behind observed trends. Experiments, in contrast, can provide more robust tests. While juvenile stages of trees are often more flexible than their adult forms, they usually provide predictable inference in differences across species and populations, and so should be integrated far more into studies of how season length affects growth. Using saplings and controlled environments could quickly test how much growth can shift with longer seasons—providing a potentially standardized way to compare constraints across species and populations—and then layer on shifts in external drivers.


In particular, mechanistic models will need to tease out changes in season length from warming that affects rates; a challenge best addressed by new experiments that decouple these two factors. Such experiments could start on juvenile trees to help inform the underlying model, select representative species to focus on, and develop predictions for large-scale studies. Experiments could also inform a better model of lag effects across species, with small-scale studies sampling saplings multiple years after manipulations (versus the common practice of destructive sampling at the end of the treatment growing season) and large-scale studies following existing efforts to test for ecological ‘memory’ (e.g. Flinker *et al.*, 2021; Schweiger *et al.*, 2022; Chin *et al.*, 2025). These efforts should help bridge across the contrasting timescales of current physiological and dendrochronological studies of growing season length: we found most physiological studies of growth  $\times$  growing season length relationships studied 1-2 years of dynamics, usually of juvenile trees, while tree ring studies focused on synthesizing across decades of adult tree growth.

Expanding studies across more species will be critical for useful models that can forecast at relevant scales and to help design large-scale experiments. While experimenting on adult trees is difficult, previous challenges in climate change research have led to large-scale experiments to understand other complex drivers (e.g. SPRUCE, DroughtNet, Pfywald, Norby & Zak, 2011; Hanson *et al.*, 2017; Smith *et al.*, 2016). We expect similar experiments will be critical here. Preparing for these large experiments using trait-mediated phylogenetic models to understand responses across species, however, could advance new experiments well beyond past efforts. By informing which species or clades to study, new experiments could span enough phylogenetic and trait diversity to forecast to species beyond the experiment and maximize the information gained (Cadotte *et al.*, 2017).

*Conclusions:* Anthropogenic climate change has often been described as an unfortunate and unplanned experiment. Like many experiments, it has highlighted important biology we do not know well. Understanding when, how and why longer seasons lead to increased tree growth requires an interdisciplinary reckoning with how temperature, time and a suite of external and internal drivers affect plant growth across species. A mechanistic understanding of how these drivers integrate over species diverse growth strategies and the imprints of evolutionary history to affect growth today is possible, but will require new approaches. Starting now to leverage data across species to inform and design new large-scale studies and experiments be critical for accurate models of future forest and related carbon dynamics, with implications for projections of carbon sequestration and carbon markets.

*Acknowledgements:* B. Wu for extracting growth  $\times$  elevation data; R. Zäch for logistical support; N. Pederson for discussion, J. Davies and three reviewers for comments that improved the manuscript.

## Box. Growth, growing season length and the challenge of standardized metrics



318

319 A major challenge in determining how growth responds to longer growing seasons is the complex-  
 320 ity of each, which means that neither can have one simple definition. Here we show the simplified  
 321 climate of one year (a), which drives variation (b) in primary growth (root and shoot elongation  
 322 and leaf production from meristems) and secondary growth (radial wood and bark growth from  
 323 cambia), both of which often depend on conditions determining non-structural carbohydrates  
 324 (NSC, which are sugars and starch needed for growth and an important area of study, for more  
 325 details see Luo *et al.*, 2024) and storage from previous seasons. Each of these types of growth  
 326 could define the growing season length (GSL, c) but GSL can also be defined meteorologically  
 327 (shown here as time,  $t$ , above some minimum  $X^\circ\text{C}$  and below above some maximum  $X^\circ\text{C}$ , with  
 328 sufficient soil moisture) or by large-scale measures of plant productivity (Körner *et al.*, 2023).  
 329 Lagged effects (shown in gray in b) are lasting impacts of previous time periods either in the  
 330 form of NSC stores or structural legacies influencing productivity (e.g. vessel diameter)

331 Of studies in our literature review, the largest proportion used metrics related to secondary  
 332 growth, quantifying growth by measuring radial growth (e.g. through increment cores or den-  
 333 drometers,  $n = 28$ ), but a number also looked at metrics related to primary growth, including C  
 334 assimilation (e.g. net ecosystem productivity or gross primary productivity,  $n = 20$ ). A smaller  
 335 number used metrics that reflect combined primary and secondary growth, including biomass,

height, or number of stems ( $n=9$ ), and root:shoot ratio ( $n=1$ ), while others used various modeled estimates potentially related to growth. For growing season length, the largest number of studies used vegetative (e.g. budburst to leaf senescence in our figure above, 26 studies) or wood phenology (11 studies) as their definition, while a smaller number used a meteorological definitions or fixed dates (7 studies). We found 14 studies that did not directly measure GSL (e.g. Zhu *et al.*, 2021; Dow *et al.*, 2022; Zohner *et al.*, 2023). Further, these definitions of GSL are simplified and could not be easily aligned, as we found 14 different metrics of start of season and 16 metrics of end of season, resulting in 25 metrics of growing season length. See also ‘The challenge of metrics: Measuring growth and growing season length’ in the Supplement.

## 1 References

- Aitken, S.N. & Bemmels, J.B. (2016) Time to get moving: assisted gene flow of forest trees. *Evol Appl* **9**, 271–90.
- Ávila-Lovera, E., Winter, K. & Goldsmith, G.R. (2023) Evidence for phylogenetic signal and correlated evolution in plant–water relation traits. *New Phytologist* **237**, 392–407.
- Berdanier, A.B. & Klein, J.A. (2011) Growing season length and soil moisture interactively constrain high elevation aboveground net primary production. *Ecosystems* **14**, 963–974.
- Cabon, A., Kannenberg, S.A., Arain, A., Babst, F., Baldocchi, D., Belmecheri, S., Delpierre, N., Guerrieri, R., Maxwell, J.T., McKenzie, S. *et al.* (2022) Cross-biome synthesis of source versus sink limits to tree growth. *Science* **376**, 758–761.
- Cadotte, M.W., Davies, T.J. & Peres-Neto, P.R. (2017) Why phylogenies do not always predict ecological differences. *Ecological Monographs* **87**, 535–551.
- Camarero, J.J., Campelo, F., Colangelo, M., Valeriano, C., Knorre, A., Solé, G. & Rubio-Cuadrado, Á. (2022) Decoupled leaf-wood phenology in two pine species from contrasting climates: Longer growing seasons do not mean more radial growth. *Agricultural and Forest Meteorology* **327**, 109223.
- Canadell, J., Monteiro, P., Costa, M., Cotrim da Cunha, L., Cox, P., Eliseev, A., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S. & Zickfeld, K. (2021) *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, New York, NY.
- Chamberlain, C.J., Cook, B.I., Morales-Castilla, I. & Wolkovich, E.M. (2021) Climate change reshapes the drivers of false spring risk across european trees. *New Phytologist* **229**, 323–334.
- Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A.E. (2009) Towards a worldwide wood economics spectrum. *Ecology Letters* **12**, 351–366.
- Chen, W., Black, T., Yang, P., Barr, A., Neumann, H., Nesic, Z., Blanken, P., Novak, M., Eley, J., Ketler, R. *et al.* (1999) Effects of climatic variability on the annual carbon sequestration by a boreal aspen forest. *Global Change Biology* **5**, 41–53.
- Chin, A.R., Gessler, A., Laín, O., Østerlund, I., Schaub, M., Thérroux-Rancourt, G., Voggeneder, K. & Lambers, J.H.R. (2025) The memory of past water abundance shapes trees 7 years later. *American Journal of Botany* p. e16452.

378 Cleland, E.E., Allen, J.M., Crimmins, T.M., Dunne, J.A., Pau, S., Travers, S.E., Zavaleta, E.S.  
379 & Wolkovich, E.M. (2012) Phenological tracking enables positive species responses to climate  
380 change. *Ecology* **93**, 1765–1771.

381 Cleland, E.E. & Wolkovich, E. (2024) Effects of phenology on plant community assembly and  
382 structure. *Annual Review of Ecology, Evolution, and Systematics* **55**, 471–492.

383 Colautti, R.I., Eckert, C.G. & Barrett, S.C.H. (2010) Evolutionary constraints on adaptive  
384 evolution during range expansion in an invasive plant. *Proceedings of the Royal Society B-  
385 Biological Sciences* **277**, 1799–1806, colautti, Robert I. Eckert, Christopher G. Barrett, Spencer  
386 C. H.

387 Cornwell, W. & Nakagawa, S. (2017) Phylogenetic comparative methods. *Current Biology* **27**,  
388 R333–R336.

389 Cornwell, W.K., Westoby, M., Falster, D.S., FitzJohn, R.G., O’Meara, B.C., Pennell, M.W.,  
390 McGlinn, D.J., Eastman, J.M., Moles, A.T., Reich, P.B. *et al.* (2014) Functional distinctive-  
391 ness of major plant lineages. *Journal of Ecology* **102**, 345–356.

392 Cosgrove, D.J. (2005) Growth of the plant cell wall. *Nature reviews molecular cell biology* **6**,  
393 850–861.

394 Cosgrove, D.J. (2023) Structure and growth of plant cell walls. *Nature Reviews Molecular Cell  
395 Biology* pp. 1–19.

396 Coulthard, B.L., George, S.S. & Meko, D.M. (2020) The limits of freely-available tree-ring  
397 chronologies. *Quaternary Science Reviews* **234**, 106264.

398 Cuapio-Hernández, L., Reyes-Ortiz, J.L., De La Rosa, A.B., Pavón, N.P., López-Herrera, M.,  
399 Villanueva-Díaz, J. & Sánchez-González, A. (2022) Is there a response pattern between radial  
400 growth of trees and elevation gradient? *Tree-Ring Research* .

401 Cuny, H.E., Rathgeber, C.B., Lebourgeois, F., Fortin, M. & Fournier, M. (2012) Life strategies  
402 in intra-annual dynamics of wood formation: example of three conifer species in a temperate  
403 forest in north-east france. *Tree physiology* **32**, 612–625.

404 Davies, T.J., Regetz, J., Wolkovich, E.M. & McGill, B.J. (2019) Phylogenetically weighted  
405 regression: A method for modelling non-stationarity on evolutionary trees. *Global ecology and  
406 biogeography* **28**, 275–285.

407 Day, M.E. & Greenwood, M.S. (2011) Regulation of ontogeny in temperate conifers. *Size-and  
408 age-related changes in tree structure and function*, pp. 91–119, Springer.

409 de Sauvage, J.C., Vitasse, Y., Meier, M., Delzon, S. & Bigler, C. (2022) Temperature rather  
410 than individual growing period length determines radial growth of sessile oak in the pyrenees.  
411 *Agricultural and Forest Meteorology* **317**, 108885.

412 Delpierre, N., Guillemot, J., Dufrêne, E., Cecchini, S. & Nicolas, M. (2017) Tree phenological  
413 ranks repeat from year to year and correlate with growth in temperate deciduous forests.  
414 *Agricultural and Forest Meteorology* **234**, 1–10.

415 Diaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., Reu, B., Kleyer, M.,  
416 Wirth, C., Prentice, I.C., Garnier, E., Bonisch, G., Westoby, M., Poorter, H., Reich, P.B.,  
417 Moles, A.T., Dickie, J., Gillison, A.N., Zanne, A.E., Chave, J., Wright, S.J., Sheremet'ev,  
418 S.N., Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup, D., Casanoves,  
419 F., Joswig, J.S., Gunther, A., Falczuk, V., Ruger, N., Mahecha, M.D. & Gorne, L.D. (2016)  
420 The global spectrum of plant form and function. *Nature* **529**, 167–U73.

421 Dow, C., Kim, A.Y., D'Orangeville, L., Gonzalez-Akre, E.B., Helcoski, R., Herrmann, V., Harley,  
422 G.L., Maxwell, J.T., McGregor, I.R., McShea, W.J. *et al.* (2022) Warm springs alter timing  
423 but not total growth of temperate deciduous trees. *Nature* **608**, 552–557.

424 Etzold, S., Sterck, F., Bose, A.K., Braun, S., Buchmann, N., Eugster, W., Gessler, A., Kahmen,  
425 A., Peters, R.L., Vitasse, Y. *et al.* (2022) Number of growth days and not length of the growth  
426 period determines radial stem growth of temperate trees. *Ecology Letters* **25**, 427–439.

427 Finzi, A.C., Giasson, M.A., Plotkin, A.A.B., Aber, J.D., Boose, E.R., Davidson, E.A., Dietze,  
428 M.C., Ellison, A.M., Frey, S.D., Goldman, E., Keenan, T.F., Melillo, J.M., Munger, J.W.,  
429 Nadelhoffer, K.J., Ollinger, V., S., Orwig, D.A., Pederson, N., Richardson, A.D., Savage, K.,  
430 Tang, J., Thompson, J.R., Williams, C.A., Wofsy, S.C., Zhou, Z. & Foster, D.R. (2020) Carbon  
431 budget of the harvard forest long-term ecological research site: pattern, process, and response  
432 to global change. *Ecological Monographs* **90**.

433 Fischer, C., Traub, B. *et al.* (2019) *Swiss National Forest Inventory-methods and models of the*  
434 *fourth assessment*. Springer.

435 Flinker, R.H., Cardenas, M.B., Caldwell, T.G., Flerchinger, G.N., Roy, R. & Reich, P.B. (2021)  
436 Promise and pitfalls of modeling grassland soil moisture in a free-air co2 enrichment experi-  
437 ment (biocon) using the shaw model. *Pedosphere* **31**, 783–795.

438 Forest Service - United States Department of Agriculture (????) Forest Inventory and Analysis  
439 (FIA) database.

440 Frank, D., Fang, K. & Fonti, P. (2022) Dendrochronology: Fundamentals and innovations. *Stable*  
441 *Isotopes in Tree Rings: Inferring Physiological, Climatic and Environmental Responses*, pp.  
442 21–59, Springer International Publishing Cham.

443 Friedlingstein, P., O'sullivan, M., Jones, M.W., Andrew, R.M., Gregor, L., Hauck, J., Le Quéré,  
444 C., Luijkx, I.T., Olsen, A., Peters, G.P. *et al.* (2022) Global carbon budget 2022. *Earth System*  
445 *Science Data Discussions* **2022**, 1–159.

446 Gantois, J. (2022) New tree-level temperature response curves document sensitivity of tree  
447 growth to high temperatures across a us-wide climatic gradient. *Global Change Biology* **28**,  
448 6002–6020.

449 Gordon, J.C. & Larson, P.R. (1968) Seasonal course of photosynthesis, respiration, and distri-  
450 bution of 14c in young pinus resinosa trees as related to wood formation. *Plant Physiology* **43**,  
451 1617–1624.

452 Green, J.K. & Keenan, T.F. (2022) The limits of forest carbon sequestration. *Science* **376**,  
453 692–693.

454 Grime, J.P. (1977) Evidence for existence of 3 primary strategies in plants and its relevance to  
455 ecological and evolutionary theory. *American Naturalist* **111**, 1169–1194.

456 Hacket-Pain, A., Friend, A., Lageard, J. & Thomas, P. (2016a) Tree rings and masting: con-  
457 sidering reproductive phenomena when interpreting tree rings? *Tree Rings in Archaeology,*  
458 *Climatology and Ecology* **14**, 78–85.

459 Hacket-Pain, A.J., Cavin, L., Friend, A.D. & Jump, A. (2016b) Consistent limitation of growth  
460 by high temperature and low precipitation from range core to southern edge of european beech  
461 indicates widespread vulnerability to changing climate. *European Journal of Forest Research*  
462 **135**, 897–909.

463 Hanson, P.J., Riggs, J.S., Nettles, W.R., Phillips, J.R., Krassovski, M.B., Hook, L.A., Gu, L.,  
464 Richardson, A.D., Aubrecht, D.M., Ricciuto, D.M. *et al.* (2017) Attaining whole-ecosystem  
465 warming using air and deep-soil heating methods with an elevated co 2 atmosphere. *Biogeo-*  
466 *sciences* **14**, 861–883.

467 Harmon, L. (2024) *Phylogenetic Comparative Methods*. University of Idaho.

468 Hilty, J., Muller, B., Pantin, F. & Leuzinger, S. (2021) Plant growth: The what, the how, and  
469 the why. *New Phytologist* **232**, 25–41.

470 Howe, G.T., Aitken, S.N., Neale, D.B., Jermstad, K.D., Wheeler, N.C. & Chen, T.H. (2003)  
471 From genotype to phenotype: unraveling the complexities of cold adaptation in forest trees.  
472 *Canadian Journal of Botany* **81**, 1247–1266.

473 Ito, G., Romanou, A., Kiang, N.Y., Faluvegi, G., Aleinov, I., Ruedy, R., Russell, G., Lerner,  
474 P., Kelley, M. & Lo, K. (2020) Global carbon cycle and climate feedbacks in the nasa giss  
475 modele2. 1. *Journal of Advances in Modeling Earth Systems* **12**, e2019MS002030.

476 Journé, V., Szymkowiak, J., Foest, J., Hacket-Pain, A., Kelly, D. & Bogdziewicz, M. (2024) Sum-  
477 mer solstice orchestrates the subcontinental-scale synchrony of mast seeding. *Nature Plants*  
478 **10**, 367–373.



479 Keenan, T.F., Gray, J., Friedl, M.A., Toomey, M., Bohrer, G., Hollinger, D.Y., Munger, J.W.,  
480 O’Keefe, J., Schmid, H.P., Wing, I.S. *et al.* (2014) Net carbon uptake has increased through  
481 warming-induced changes in temperate forest phenology. *Nature Climate Change* **4**, 598–604.

482 Kikuzawa, K. (1982) Leaf survival and evolution in betulaceae. *Annals of Botany* **50**, 345–353.

483 King, G.M., Gugerli, F., Fonti, P. & Frank, D.C. (2013) Tree growth response along an eleva-  
484 tional gradient: climate or genetics? *Oecologia* **173**, 1587–1600.

485 Klesse, S., Babst, F., Evans, M.E., Hurley, A., Pappas, C. & Peters, R.L. (2023) Legacy effects  
486 in radial tree growth are rarely significant after accounting for biological memory. *Journal of*  
487 *Ecology* **111**, 1188–1202.

488 Kolář, T., Giagli, K., Trnka, M., Bednářová, E., Vavřík, H. & Rybníček, M. (2016) Response of  
489 the leaf phenology and tree-ring width of european beech to climate variability. *Silva Fennica*  
490 **50**.

491 Körner, C., Möhl, P. & Hiltbrunner, E. (2023) Four ways to define the growing season. *Ecology*  
492 *Letters* .

493 Kramer, P. (2012) *Physiology of woody plants*. Elsevier, New York.

494 La Porta, N., Capretti, P., Thomsen, I., Kasanen, R., Hietala, A. & Von Weissenberg, K. (2008)  
495 Forest pathogens with higher damage potential due to climate change in europe. *Canadian*  
496 *Journal of Plant Pathology* **30**, 177–195.

497 Lange, H., Økland, B. & Krokene, P. (2006) Thresholds in the life cycle of the spruce bark beetle  
498 under climate change. *Interjournal for Complex Systems* **1648**, 1–10.

499 Larcher, W. (1980) *Plant Physiological Ecology*. Springer-Verlag.

500 Lechowicz, M.J. (1984) Why do temperate deciduous trees leaf out at different times - adaptation  
501 and ecology of forest communities. *American Naturalist* **124**, 821–842.

502 Leuning, R. (2002) Temperature dependence of two parameters in a photosynthesis model. *Plant,*  
503 *Cell & Environment* **25**, 1205–1210.

504 Luo, Y., Zohner, C., Crowther, T.W., Feng, J., Hoch, G., Li, P., Richardson, A.D., Vitasse, Y. &  
505 Gessler, A. (2024) Internal physiological drivers of leaf development in trees: Understanding  
506 the relationship between non-structural carbohydrates and leaf phenology. *Functional Ecology*  
507 .

508 Marchand, L.J., Dox, I., Gričar, J., Prislan, P., Van den Bulcke, J., Fonti, P. & Campioli, M.  
509 (2021) Timing of spring xylogenesis in temperate deciduous tree species relates to tree growth  
510 characteristics and previous autumn phenology. *Tree Physiology* **41**, 1161–1170.

511 Martinez-Meier, A., Sanchez, L., Pastorino, M., Gallo, L. & Rozenberg, P. (2008) What is hot  
512 in tree rings? the wood density of surviving douglas-firs to the 2003 drought and heat wave.  
513 *Forest Ecology and Management* **256**, 837–843.

514 McKown, A.D., Guy, R.D., Klapste, J., Geraldles, A., Friedmann, M., Cronk, Q.C.B., El-  
515 Kassaby, Y.A., Mansfield, S.D. & Douglas, C.J. (2014) Geographical and environmental gra-  
516 dients shape phenotypic trait variation and genetic structure in populus trichocarpa. *New*  
517 *Phytologist* **201**, 1263–1276, McKown, Athena D. Guy, Robert D. Klapste, Jaroslav Geraldles,  
518 Armando Friedmann, Michael Cronk, Quentin C. B. El-Kassaby, Yousry A. Mansfield, Shawn  
519 D. Douglas, Carl J.

520 McKown, A.D., Guy, R.D. & Quamme, L.K. (2016) Impacts of bud set and lammas phenology on  
521 root: shoot biomass partitioning and carbon gain physiology in poplar. *Trees* **30**, 2131–2141.

522 Michelot, A., Simard, S., Rathgeber, C., Dufrêne, E. & Damesin, C. (2012) Comparing the  
523 intra-annual wood formation of three European species (*Fagus sylvatica*, *Quercus petraea* and  
524 *Pinus sylvestris*) as related to leaf phenology and non-structural carbohydrate dynamics. *Tree*  
525 *physiology* **32**, 1033–1045.

526 Mitton, J.B. & Ferrenberg, S.M. (2012) Mountain pine beetle develops an unprecedented summer  
527 generation in response to climate warming. *The American Naturalist* **179**, E163–E171.

528 Monson, R.K., Szejner, P., Belmecheri, S., Morino, K.A. & Wright, W.E. (2018) Finding the  
529 seasons in tree ring stable isotope ratios. *American journal of botany* **105**, 819–821.

530 Morales-Castilla, I., Davies, T., Legault, G., Buonaiuto, D., Chamberlain, C.J., Ettinger, A.K.,  
531 Garner, M., Jones, F.A., Loughnan, D., Pearse, W.D. *et al.* (2024) Phylogenetic estimates of  
532 species-level phenology improve ecological forecasting. *Nature Climate Change* pp. 1–7.

533 Myneni, R.B., Keeling, C., Tucker, C.J., Asrar, G. & Nemani, R.R. (1997) Increased plant  
534 growth in the northern high latitudes from 1981 to 1991. *Nature* **386**, 698–702.

535 Nobel, P.S. *et al.* (1983) *Biophysical plant physiology and ecology*. WH Freeman and company.

536 Norby, R.J. & Zak, D.R. (2011) Ecological lessons from free-air co2 enrichment (face) experi-  
537 ments. *Annual review of ecology, evolution, and systematics* **42**, 181–203.

538 Pearse, W.D., Legendre, P., Peres-Neto, P.R. & Davies, T.J. (2019) The interaction of phylogeny  
539 and community structure: Linking the community composition and trait evolution of clades.  
540 *Global Ecology and Biogeography* **28**, 1499–1511.

541 Peters, R.L., Steppe, K., Cuny, H.E., De Pauw, D.J., Frank, D.C., Schaub, M., Rathgeber, C.B.,  
542 Cabon, A. & Fonti, P. (2021) Turgor—a limiting factor for radial growth in mature conifers  
543 along an elevational gradient. *New Phytologist* **229**, 213–229.

544 Puletti, N., Grotti, M., Ferrara, C. & Chianucci, F. (2020) Lidar-based estimates of aboveground  
545 biomass through ground, aerial, and satellite observation: a case study in a mediterranean  
546 forest. *Journal of Applied Remote Sensing* **14**, 044501–044501.

547 Radville, L., McCormack, M.L., Post, E. & Eissenstat, D.M. (2016) Root phenology in a chang-  
548 ing climate. *Journal of Experimental Botany* **67**, 3617–3628.

549 Ren, P., Ziaco, E., Rossi, S., Biondi, F., Prislán, P. & Liang, E. (2019) Growth rate rather than  
550 growing season length determines wood biomass in dry environments. *Agricultural and Forest*  
551 *Meteorology* **271**, 46–53.

552 Rezende, E.L. & Bozinovic, F. (2019) Thermal performance across levels of biological organiza-  
553 tion. *Philosophical Transactions of the Royal Society B* **374**, 20180549.

554 Schaber, J. & Badeck, F.W. (2002) Evaluation of methods for the combination of phenological  
555 time series and outlier detection. *Tree Physiology* **22**, 973–982.

556 Schweiger, A., Awdamet, F.T., Drenckhan, W. & Berauer, B.J. (2022) Transgenerational effects  
557 of elevated co<sub>2</sub>: Downregulation of photosynthetic efficiency and stomatal sensitivity to  
558 drought .

559 Senf, C., Seidl, R. & Hostert, P. (2017) Remote sensing of forest insect disturbances: Current  
560 state and future directions. *International journal of applied earth observation and geoinfor-*  
561 *mation* **60**, 49–60.

562 Sillett, S.C., Graham, M.E., Montague, J.P., Antoine, M.E. & Koch, G.W. (2024) Ground-based  
563 calibration for remote sensing of biomass in the tallest forests. *Forest Ecology and Management*  
564 **561**, 121879.

565 Silvestro, R., Zeng, Q., Buttò, V., Sylvain, J.D., Drolet, G., Mencuccini, M., Thiffault, N.,  
566 Yuan, S. & Rossi, S. (2023) A longer wood growing season does not lead to higher carbon  
567 sequestration. *Scientific reports* **13**, 4059.

568 Smith, M.D., Wilcox, K., Sala, O., Phillips, R., Luo, Y., Knapp, A. & Lemoine, N.P. (2016)  
569 Drought-net: A global network merging observations, experiments, and modeling to forecast  
570 terrestrial ecosystem sensitivity to drought. *AGU Fall Meeting Abstracts*, vol. 2016, pp. B11J–  
571 06.

572 Soolanayakanahally, R.Y., Guy, R.D., Silim, S.N. & Song, M. (2013) Timing of photoperiodic  
573 competency causes phenological mismatch in balsam poplar (*populus balsamifera* l.). *Plant,*  
574 *cell & environment* **36**, 116–127.

575 Sturrock, R., Frankel, S., Brown, A., Hennon, P., Kliejunas, J., Lewis, K., Worrall, J. & Woods,  
576 A. (2011) Climate change and forest diseases. *Plant pathology* **60**, 133–149.

- 577 Thomas, S.C. (2011) Age-related changes in tree growth and functional biology: The role  
578 of reproduction. *Size- and Age-Related Changes in Tree Structure and Function* (eds. F.C.  
579 MEINZER, B. Lachenbruch & T.E. Dawson), chap. 2, pp. 33–64, Springer.
- 580 Waring, R.H. & Franklin, J.F. (1979) Evergreen coniferous forests of the pacific northwest:  
581 Massive long-lived conifers dominating these forests are adapted to a winter-wet, summer-dry  
582 environment. *Science* **204**, 1380–1386.
- 583 Webb, C.O., Ackerly, D.D., McPeck, M. & Donoghue, M.J. (2002) Phylogenies and community  
584 ecology. *Annual Review of Ecology and Systematics* **33**, 475–505.
- 585 Went, F.W. (1957) The experimental control of plant growth. *The experimental control of plant*  
586 *growth*. **17**.
- 587 Westoby, M., Yates, L., Holland, B. & Halliwell, B. (2023) Phylogenetically conservative trait  
588 correlation: quantification and interpretation. *Journal of Ecology* **111**, 2105–2117.
- 589 Willis, C.G., Ruhfel, B., Primack, R.B., Miller-Rushing, A.J. & Davis, C.C. (2008) Phylogenetic  
590 patterns of species loss in thoreau’s woods are driven by climate change. *Proceedings of the*  
591 *National Academy of Sciences* **105**, 17029–17033.
- 592 Zeng, Z.A. & Wolkovich, E.M. (2024) Weak evidence of provenance effects in spring phenology  
593 across europe and north america. *New Phytologist* .
- 594 Zhu, L., Liu, S., Arzac, A., Cooper, D.J., Jin, Y., Yuan, D., Zhu, Y., Zhang, X., Li, Z., Zhang, Y.  
595 *et al.* (2021) Different response of earlywood vessel features of fraxinus mandshurica to rapid  
596 warming in warm-dry and cold-wet areas. *Agricultural and Forest Meteorology* **307**, 108523.
- 597 Zohner, C.M., Mirzaghali, L., Renner, S.S., Mo, L., Rebindaine, D., Bucher, R., Palouš, D.,  
598 Vitasse, Y., Fu, Y.H., Stocker, B.D. *et al.* (2023) Effect of climate warming on the timing of  
599 autumn leaf senescence reverses after the summer solstice. *Science* **381**, eadf5098.

## 2 Figures

GS: Growing season  
GSL: Growing season length




Figure 1: Climate change may alter growing season length (GSL), which can then affect growth through diverse pathways. We review hypotheses for these pathways showing the number of papers (from a review of papers studying growth  $\times$  growing season length) that mentioned each hypothesis (width of the shaded areas of left column is proportional to the number of papers with the number also given, right column shows the expected growth response for each hypothesis). We group hypotheses as focused on mechanisms moderated by the environment ('external') versus those focused on internal physiological constraints, which span both source (photosynthesis-limited) and sink limitation, and could act together. For more details, see Supplement.




Figure 2: A review of growth  $\times$  growing season length relationship studies spanned a diversity of methods, but there was no coherency in which methods did or did not find a positive relationship. A number of studies tested relationships possibly related to growth  $\times$  growing season length (e.g. they tested how spring temperatures related to growth) but never directly growth  $\times$  growing season length, thus ‘not tested’ was surprisingly common across methods. See Supplement for review details.




Figure 3: Understanding how longer seasons with climate change affect growth requires teasing out effects of longer seasons versus warmer seasons, which generally co-vary in observational data. Here we show two examples of this complexity. In (a) we show a general net photosynthesis response curve (top panel), which has a non-linear response to temperature (blue curve, adapted from meta-analysis of Rezende & Bozinovic, 2019), though it is often modeled as linear (red). This non-linearity means that increases in lower temperatures—such as those in the spring when much of growing season extensions may happen—have lower absolute increases in photosynthesis compared to increases in later-season (e.g. summer) warmer temperatures, while a linear response assumes a constant scale of effect across low to high temperatures (bottom). In (b) we show conceptual growth responses to temperature for two different species with different growth rate responses to temperature (top, inset), which impacts their growth across the season, leading to small absolute differences in accumulated growth at a conceptual high elevation site (top) versus larger differences in accumulated growth low elevation site (bottom). Testing how growth varies across larger spatial gradients of growing season length, as we conceptualize here (b) could help establish a baseline expectation of the scale of temporal—especially inter-annual variation—and force a greater reckoning with drivers that shift alongside growing season length.




Figure 4: A trait-based phylogenetic model provides a way to naturally organize species (and, not shown, populations, which nest within species) responses to predict how they respond to longer seasons. This approach estimates a universal model that is then shaped by species evolutionary history (shown at left via a phylogenetic tree) and traits to estimate how each species should uniquely respond. We argue this framework can organize and guide experiments that separate out changes in temperature from changes in growing season length ( $^{\circ}\text{C}$  and GSL in see middle panels) to better integrate observational data and identify different responses by species that can help forecast (see ‘Building a new framework for growth  $\times$  season length relationships’ section for more details). In this example, we show how this approach can identify one clade (top) with a common response to longer seasons that also shares a suite of similar traits, and can identify a unique response by one species in a clade where that species also has a unique trait compared to other species with the same common ancestor (lower clade). Estimating responses across species through a phylogenetic model has the additional benefit of robustly estimating responses given uneven sampling and missing data (the dashed red lines represent that the model will predict a response for each species, informed to various degrees—with the degree determined by the model—by the traits and phylogeny).