

Shifting tree growth in response to shifting seasons

Background – Forests across the globe provide critical resources for people.^{1,2} Economically their timber underpins the economies of many states, provinces and countries, with their economic value increasing with climate change through their role in the carbon cycle. Temperate forests in particular currently sequester substantial fossil fuel emissions, offsetting planetary warming.³⁻⁵

Yet how long forests can continue to offset fossil fuel emissions depends on how they shift their growth in response to the longer, warmer seasons that come with anthropogenic climate change. Fundamentally, longer seasons should allow trees more time to grow and thus increase their growth, an assumption that was previously supported by several studies.^{6,7} Recently, however, new work suggests trees do not increase growth with longer seasons,⁸ but why and when this happens is a major open question.⁹

My approach – My focus on temporal ecology and shifting seasons with climate change (see *Career narrative*) make me well positioned to address this major open question. Further, my current progress on this question, alongside new datasets^{10,11} and advances in modeling approaches,^{12,13} positions me well to make major progress on it now through a Guggenheim Fellowship.

Over the past three years I have begun to work in-depth on tree growth, including outlining the major potential reasons we may see variability in tree growth with climate change.⁹ It seems clear that the likely reasons tree do not grow more with longer seasons are drought stress (not enough water), heat stress (too high temperatures), or species-specific constraints (some species are simply conservative in their responses, while others are not). But testing these hypotheses has proved much harder because of the complexity of tree growth.

Individual-level tree growth is often highly variable and currently difficult to predict (Fig. 1). For these reasons, most work on radial tree growth to date has worked from stand-level averages, after correcting for age-trends (tree-level




Figure 1: An example of the complexity of tree growth from one *Pinus ponderosa* tree sampled in Utah (USA). Dots show the observed radial growth each year, while the coloring represents estimates from a hierarchical model; shown here is a hierarchical Gaussian process model I may use.¹⁴

allometry).^{15,16} Yet this approach is far less useful for understanding the relative importance of longer seasons, which requires understanding the actual drivers of variability. It thus means modeling the time-series of individual trees (Fig. 1) and then building up to understand how different species and stands (sites) respond to longer seasons alongside climate. It effectively requires an entirely new approach to modeling tree growth.

Methods – For my Guggenheim fellowship I propose to develop new models of tree growth that can tease apart how longer seasons, alongside shifting climate, have impacted diverse forests across North America. I have already begun preliminary work on these models (Fig. 1) and am well positioned to complete them for trees across North America during my fellowship.

I propose to work with a leader in hierarchical modeling (Andrew Gelman, Columbia University) and an expert in tree growth and drought (Dr. Benjamin Cook, NASA-GISS/Monash University) to understand how tree growth and forests have shifted in recent decades with longer seasons, and project shifts in the future. I will use tree ring data from the International Tree Ring Data Bank¹⁰, and climate data from PRISM and NASA to model tree growth across North America since 1896 (start date of of robust gridded climate data). We will estimate the start and end of the growing season using newly available gridded data.¹¹ For climate, I plan to begin by focusing on summer warmth through growing degree days (GDD), moisture (and thus drought) estimated through precipitation sums and stress effects of both heat and drought possible through high vapor pressure deficit (VPD). Given the different climate drivers in western versus eastern North America, I plan a multi-step approach: I will fit data from western North America first where winter precipitation and VPD often dominate as drivers, then I will fit data from eastern North America where GDD is likely more important. Multiple aspects of my proposed work make it highly feasible, including the scientific backgrounds, interests and skills of myself and proposed collaborators (Gelman and Cook), as well as extensive work to compile data and design a plan that can be rapidly achieved.

Impact – My focus on climate will be critical for understanding how longer seasons affect tree growth outside of the shifting climate that drives longer seasons. Further, it will allow me to forecast from my model to predict how forests will respond to continuing anthropogenic climate change. These forecasts will be especially unique in providing guidance on how forest health may shift depending on species and age composition (both of which can be managed through forest policy).

My approach will yield a multi-level and multi-risk view, highlighting areas where management and policy could affect outcomes, yielding more resilient forests.

This project will shift our fundamental scientific understanding of how trees grow, yielding our best estimates of radial growth responses to climate—and how growth shifts with longer seasons. The work during my fellowship will lead to 1-2 papers in high profile scientific journals (e.g., *Science*, *Nature*) and likely high media coverage, which I have often received for my work. Beyond, I expect this new outlook on tree growth will lead to new experiments, studies and drive new knowledge. Additionally, multiple aspects of my approach are designed to extend beyond. My multi-scale approach can be extended both outward—to forests in other regions, countries and continents—and temporally downscaled, moving beyond annual resolution to potentially drawing on weekly or daily data (increasingly available through dendrometer networks). Both areas I hope to expand to after my fellowship period.

Bibliography

- [1] Oldekop, J. A., Rasmussen, L. V., Agrawal, A., Bebbington, A. J., Meyfroidt, P., Bengston, D. N., Blackman, A., Brooks, S., Davidson-Hunt, I., Davies, P., et al. (2020). Forest-linked livelihoods in a globalized world. *Nature Plants* *6*, 1400–1407.
- [2] Costanza, R., Fioramonti, L., and Kubiszewski, I. (2016). The un sustainable development goals and the dynamics of well-being. *Frontiers in Ecology & the Environment* *14*, 59.
- [3] Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., et al. (2011). A large and persistent carbon sink in the world’s forests. *science* *333*, 988–993.
- [4] Griscom, B. W., Adams, J., Ellis, P. W., Houghton, R. A., Lomax, G., Miteva, D. A., Schlesinger, W. H., Shoch, D., Siikamäki, J. V., Smith, P., et al. (2017). Natural climate solutions. *Proceedings of the national Academy of sciences* *114*, 11645–11650.
- [5] Friedlingstein, P., O’sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T., Olsen, A., Peters, G. P., et al. (2022). Global carbon budget 2022. *Earth System Science Data Discussions* *2022*, 1–159.
- [6] Keenan, T. F., Gray, J., Friedl, M. A., Toomey, M., Bohrer, G., Hollinger, D. Y., Munger, J. W., O’Keefe, J., Schmid, H. P., Wing, I. S., et al. (2014). Net carbon uptake has increased through warming-induced changes in temperate forest phenology. *Nature Climate Change* *4*, 598–604.
- [7] Finzi, A. C., Giasson, M.-A., Plotkin, A. A. B., Aber, J. D., Boose, E. R., Davidson, E. A., Dietze, M. C., Ellison, A. M., Frey, S. D., Goldman, E., et al. (2020). Carbon budget of the harvard forest long-term ecological research site: pattern, process, and response to global change. *Ecological Monographs* *90*.
- [8] Dow, C., Kim, A. Y., D’Orangeville, L., Gonzalez-Akre, E. B., Helcoski, R., Herrmann, V., Harley, G. L., Maxwell, J. T., McGregor, I. R., McShea, W. J., et al. (2022). Warm springs alter timing but not total growth of temperate deciduous trees. *Nature* *608*, 552–557.
- [9] Wolkovich, E., Ettinger, A. K., Chin, A. R., Chamberlain, C. J., Baumgarten, F., Pradhan, K., Manzanedo, R. D., and Hille Ris Lambers, J. (2025). Why longer seasons with climate change may not increase tree growth. *Nature Climate Change* *15*, 1283–1292.
- [10] Zhao, S., Pederson, N., D’Orangeville, L., HilleRisLambers, J., Boose, E., Penone, C., Bauer, B., Jiang, Y., and Manzanedo, R. D. (2019). The international tree-ring data bank (itrd) revisited: data availability and global ecological representativity. *Journal of Biogeography* *46*, 355–368.
- [11] Nelson, J. A., Walther, S., Gans, F., Kraft, B., Weber, U., Novick, K., Buchmann, N., Migliavacca, M., Wohlfahrt, G., Šigut, L., et al. (2024). X-base: the first terrestrial carbon and water flux products from an extended data-driven scaling framework, fluxcom-x. *Biogeosciences* *21*, 5079–5115.
- [12] Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M. A., Guo, J., Li, P., and Allen, R. (2017). Stan: A probabilistic programming language. *Journal of Statistical Software* *76*, 10.18637/jss.v076.i01.
- [13] Gelman, A., Vehtari, A., McElreath, R., Simpson, D., Margossian, C. C., Yao, Y., Kennedy, L., Gabry, J., Bürkner, P.-C., Modrák, M., et al. (2026). *Bayesian workflow*. (CRC Press).
- [14] Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B. (2014). *Bayesian Data Analysis*. (New York: CRC Press) 3rd edition.
- [15] Cook, E. R. (1985). A time series analysis approach to tree ring standardization (dendrochronology, forestry, dendroclimatology, autoregressive process). (The University of Arizona).
- [16] Schofield, M. R., Barker, R. J., Gelman, A., Cook, E. R., and Briffa, K. R. (2016). A model-based approach to climate reconstruction using tree-ring data. *Journal of the American Statistical Association* *111*, 93–106.