

¹ Woody plant phenological responses are strongly associated
² with key functional traits

³ Deirdre Loughnan¹, Faith A M Jones^{1,2}, Geoffrey Legault¹, Daniel Buonaiuto^{3,4,5},
Catherine Chamberlain^{3,4,6}, Ailene Ettinger⁷, Mira Garner¹, Ignacio Morales-Castilla^{8,9},
Darwin Sodhi¹ and E M Wolkovich^{1,3,4}

⁴ September 3, 2024

⁵ ¹ Department of Forest and Conservation Sciences, Faculty of Forestry, University of British Columbia,
⁶ 2424 Main Mall Vancouver, BC Canada V6T 1Z4.

⁷ ² Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sci-
⁸ ences, 901 83 Umeå, Sweden

⁹ ³ Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

¹⁰ ⁴ Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts,
¹¹ USA;

¹² ⁵ Department of Environmental Conservation, University of Massachusetts, Amherst, 160 Holdsworth
¹³ Way, Amherst, MA, USA

¹⁴ ⁶ The Nature Conservancy, 334 Blackwell St, #300, Durham, NC, USA

¹⁵ ⁷ The Nature Conservancy of Washington, 74 Wall Street, Seattle, WA USA

¹⁶ ⁸ GloCEE—Global Change Ecology and Evolution Group, Department of Life Sciences, University of
¹⁷ Alcalá, Alcalá de Henares, Spain

¹⁸ ⁹ Edificio Ciencias, Campus Universitario 28805 Alcalá de Henares, Madrid, Spain

²⁰ Corresponding Author: Deirdre Loughnan deirdre.loughnan@ubc.ca

²¹ Running title: Budburst responses associated with traits

³² **Summary**

³³ Species-level variation in phenology—the timing of recurring life history events—can vary seasonally
³⁴ with changes in climatic risk, light, and nutrients. This favours acquisitive growth strategies early in
³⁵ the spring season and conservative growth strategies under the more climatically benign, but compet-
³⁶ itive, environment later in the season. This framework infers suites of traits that may co-vary with
³⁷ phenologies, but high variability across environments makes this challenging to test. Here, we combine
³⁸ a new joint modeling approach to accommodate this variability with global data on plant traits and
³⁹ budburst responses in controlled environment experiments. We find that earlier species—which are
⁴⁰ generally most responsive to anthropogenic warming—are generally shorter with denser, lower nitro-
⁴¹ gen leaves. These results suggest warming may reshape the trait structure of plant communities, and
⁴² could help improve predictions of how growth strategies and phenologies together shift with continued
⁴³ climate change.

⁴⁴

⁴⁵ Key Words: Budburst, spring phenology, functional traits, trees, climate change, forest communities

⁴⁶ **Introduction**

⁴⁷ The timing of life history events—phenology—can shape both ecosystem services and community
⁴⁸ dynamics. Spring phenology, for example, defines the start and overall length of the growing season—
⁴⁹ shaping forest carbon storage and species interactions^{3,9,18}. Shifts in phenology with climate change
⁵⁰ across systems^{31,45} have thus led to growing concerns over their possible impacts.

⁵¹

⁵² Predicting these changes requires understanding the drivers of phenology both at a proximate scale—
⁵³ the environmental triggers of phenology each year, such as temperature and daylength—and at an
⁵⁴ ultimate scale, where long-term environmental pressures may select for different phenologies across
⁵⁵ species, e.g., certain species are early or late relative to other species each year^{33,51}. At the proximate
⁵⁶ level, environmental conditions throughout the winter and spring cause species to start growth at dif-
⁵⁷ ferent times. Similar environmental conditions appear to trigger spring phenological events across taxa,
⁵⁸ including the start of growth in woody plants each year^{15,24}, the timing of egg laying in birds^{11,12},
⁵⁹ and the advance of spawning in amphibians^{23,43}, but current work provides limited insights into the
⁶⁰ drivers of species differences^{6,15,24}.

⁶¹

⁶² At the ultimate level, species phenologies may vary due to changing pressures across the growing season.
⁶³ Species that start growth early often risk high tissue loss—due to frost damage^{2,37} or high herbivore
⁶⁴ apparency⁴⁶—but benefit from higher resource availability^{21,34}. In contrast, later species face greater
⁶⁵ biotic pressures, especially from high competition for resources^{27,52}. For plants, this variation in early
⁶⁶ to late season growth may mirror the stressors from early to late successional communities, and may
⁶⁷ similarly shape phenology²⁴.

⁶⁸

⁶⁹ Different pressures could shape a number of species attributes related to their growth strategies, includ-
⁷⁰ ing phenology. Species with earlier phenology may produce cheaper tissues that are easily replaced if
⁷¹ damaged³⁶, while species with later phenology may benefit from investing in tissues that infer greater
⁷² resource retention¹⁶. Differences in traits, and trade-offs in allocation of resources to growth and tissue
⁷³ quality, can be related to a broader framework of species growth strategies and functional traits⁵² (Fig
⁷⁴ 1), where species range from acquisitive (fast) to more conservative (slow) growth^{5,53}.

⁷⁵

⁷⁶ Globally, gradients from acquisitive to conservative strategies appear common, and form the foun-
⁷⁷ dation of the leaf economic and the wood economic spectra^{5,13,47,48,53}, but they can make limited
⁷⁸ predictions of trait variability⁴⁴. As a result, highly variable traits like phenology are often excluded
⁷⁹ from trait studies, leaving the relationships between broader trait syndromes and phenology largely

80 unknown. Understanding these relationships is critical to forecasting community dynamics and re-
81 sponses to climate change. To date, studies have generally only examined the relationship between
82 traits and phenology within a single site (as reviewed by Wolkovich and Cleland⁵⁰ and Wolkovich and
83 Donahue⁵¹) where the problem of proximate drivers causing phenological variation can be more easily
84 ignored. Adding phenology to broader trait frameworks becomes more complex when high levels of
85 variation occurs across large spatial and temporal ranges.

86

87 Consistently defining early to late phenology is possible using the underlying cues that predict gradients
88 in phenology, which do not generally vary strongly across space and time^{7,15,20}. For many plants, early
89 species generally have responses that are small in magnitude to all three major cues of spring leafout:
90 warm spring temperatures (forcing), cool winter temperatures (chilling) and daylength (photoperiod).
91 In contrast, later species have larger responses to chilling and/or photoperiod^{15,24}, and likely larger
92 forcing responses.

93

94 Studies of spring phenology in temperate forests may provide the best opportunity to integrate phe-
95 nology into functional trait research. In addition to how well we understand the environmental cues
96 that trigger early versus late leafout, spring in many forests includes strong gradients in potential se-
97 lective environments (Fig 1). Based on trade-offs between early and late spring phenologies, we predict
98 acquisitive species to be shorter, with leaf traits favourable to higher light availability and tolerance
99 to late spring frost (high specific leaf area, SLA, and leaf nitrogen content, LNC; Fig 1). Such species
100 should exhibit early phenology, with small cue responses. Canopy species that budburst later via larger
101 cue responses, when competition for soil resources is greater, would then have traits associated with
102 conservative growth—taller with denser wood²⁵—with leaf traits suited for more variable light (low
103 SLA and LNC, Fig 1). Seed size may similarly be predicted from this acquisitive to conservative con-
104 tinuum, as acquisitive species produce smaller seeds and conservative species produce larger—better
105 provisioned—seeds (Fig 1).

106

107 To test our predicted relationships between budburst responses to environmental cues and common
108 functional traits (height, SLA, seed mass, and LNC), we merged available data from trait databases—
109 BIEN²⁹ and TRY²²—with budburst data from the OSPREE database of controlled environment stud-
110 ies¹⁴. We developed a hierarchical Bayesian joint model that predicts phenological responses to forcing,
111 chilling and photoperiod treatments based on species-level trait values, while allowing additional varia-
112 tion due to species. This approach takes a step towards predicting variation via species traits instead of
113 species identity (when traits explain a significant portion of the variation, species identity will explain
114 only a small amount), which could help forecast species phenological responses based on trait values
115 alone.

116

117 Methods

118 We merged three major databases for our analysis. We gathered phenological data from the OSPREE
119 database¹⁴, which contains budburst data for woody, species from experiments of forcing, chilling and
120 photoperiod. We updated this database since its initial publication, the methods of which are dis-
121 cussed by Morales-Castilla et al.³². We gathered trait data from TRY and BIEN (v. 4.0)^{22,29}, both
122 of which are large trait databases that include plant trait data across many individuals, species, and
123 studies (Table S1). We obtained data from both databases on 5 December, 2018, with an updated
124 version of the TRY data obtained 10 April, 2019. We focused our search for trait data on the subset
125 of 234 OSPREE species used in Morales-Castilla et al.³². Using the BIEN R package (v. 1.2.5)²⁹, we
126 downloaded trait data for the 94 species available, for which there were 13 traits. The TRY database
127 included data for 10 traits for 96 of our focal species²². Given our focus on the phenology of adult
128 trees, we included trait data from adult individuals with a minimum height of 1.38 m. We further

removed all data from experiments or from plants growing in non-natural habitats. We also grouped trait values where appropriate, for example categorizing trait values for “SLA”, “SLA with petioles”, and “SLA without petioles” as simply “SLA” in our analysis (see Table S1). Duplicated data in both the TRY and BIEN datasets were also removed ($n = 434905$). Based on our selection criteria, our final dataset included data for 11 traits from 91 of the species also represented in the OSPREE database, with each species differing in the number and types of traits measured.

135

For our analysis, we only include species for which we had a complete trait profile (i.e., all traits measured for all species). We initially considered six commonly measured traits—SLA, leaf dry matter content (LDMC), height, seed mass, stem specific density (SSD), and LNC—for which 26 species had at least one trait measurement for each trait. We then used a principle component analysis (PCA) to understand trait correlations and adjusted which traits we included. A PCA of our six initial traits identified high correlations between SLA and LDMC, and between height and SSD. The first principal component explained 32% of variation while the second explained 24.2% of the variation (Fig. S1). By excluding one trait from each of these highly correlated trait pairs (specifically LDMC and SSD) we increased the number of species in our dataset from the 26 species with six traits, to 37 species for which we had complete datasets for four traits. The data for these 37 species were from 24 unique studies (samples sizes: height $n = 47781$, seed mass $n = 281$, LNC $n = 3853$, SLA $n = 7656$). We subsampled height measurements to reduce the influence of the 13 most frequently measured tree species on our height model. Since these 13 species were measured 19 times more frequently than other species, for each species, we randomly sampled 3000 height measurements.

150

151 Joint model of trait and phenology

To understand connections between phenology and species traits, we developed and then parameterized a joint model for each trait: height, SLA, LNC, and seed mass. Our model is a joint model insofar as it involves two types of data—trait observations and phenological observations—that arise from shared latent processes. In particular, we assume that species “true” trait values determine observed trait values across different studies (trait sub-model), and separately, that the same “true” trait values interact with phenological cues (forcing, chilling, and photoperiod) to determine observed phenology, specifically the day of year of budburst (phenology sub-model). Below we describe the two sub-models, noting which parameters are shared across sub-models and which are independent.

160 Trait sub-model

The trait sub-model describes the processes that determine trait observations for 1 to n species across each of the 1 to m studies in our trait dataset (TRY and BIEN data). We use hierarchical modeling to partition trait variation by measurement error, species identity, and study identity. In particular, we assume that a trait observation for species i from study j , $Y_{\text{trait}_{i,j}}$, has the following normal distribution:

$$Y_{\text{trait}_{i,j}} \sim \mathcal{N}(\mu_{i,j}, \sigma_m^2) \quad (1)$$

with

$$\mu_{i,j} = \alpha_{\text{grand trait}} + \alpha_{\text{trait}_i} + \alpha_{\text{study}_j} \quad (2)$$

where α_{trait_i} and α_{study_j} are elements of the normal random vectors:

$$\begin{aligned} \boldsymbol{\alpha}_{\text{trait}} &= \{\alpha_{\text{trait}_1}, \dots, \alpha_{\text{trait}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{trait}} \sim \mathcal{N}(0, \sigma_{\text{trait}}^2) \\ \boldsymbol{\alpha}_{\text{study}} &= \{\alpha_{\text{study}_1}, \dots, \alpha_{\text{study}_m}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{study}} \sim \mathcal{N}(0, \sigma_{\text{study}}^2) \end{aligned} \quad (3)$$

166 The latent parameter $\alpha_{\text{grand trait}}$ represents a trait value that is independent of study and species,
 167 $\alpha_{\text{species},i}$ and $\alpha_{\text{study},j}$ are species and study-level offsets from that trait value, σ_m^2 is measurement error,
 168 and σ_{trait}^2 and σ_{study}^2 represent species and study-level variances in trait values. Of these parameters,
 169 α_{trait} are shared by the phenology sub-model.

170 Phenology sub-model

171 The phenology sub-model describes the processes that determine phenological observations for 1 to
 172 n species, specifically the timing (day of year) of budburst from the updated OSPREE dataset. We
 173 assume that an observation of budburst day for species k under set g of chilling, forcing, and photoper-
 174 iod treatments (c_g, f_g, p_g), $Y_{\text{pheno}_{k,g}}$ (which we z -scored to allow direct comparison of cues) has the
 175 following normal distribution:

$$Y_{\text{pheno}_{k,g}} \sim \mathcal{N}(\mu_{k,g}, \sigma_d^2) \quad (4)$$

with

$$\mu_{k,g} = \alpha_{\text{pheno},k} + \beta_{\text{chill}_k} \cdot c_g + \beta_{\text{force}_k} \cdot f_g + \beta_{\text{photo}_k} \cdot p_g \quad (5)$$

and

$$\begin{aligned} \beta_{\text{chill}_k} &= \alpha_{\text{chill},k} + \beta_{\text{trait.chill}} \cdot \alpha_{\text{trait},k} \\ \beta_{\text{force}_k} &= \alpha_{\text{force},k} + \beta_{\text{trait.force}} \cdot \alpha_{\text{trait},k} \\ \beta_{\text{photo}_k} &= \alpha_{\text{photo},k} + \beta_{\text{trait.photo}} \cdot \alpha_{\text{trait},k} \end{aligned} \quad (6)$$

176 where α_{pheno_k} , α_{chill_k} , α_{force_k} , and α_{photo_k} are elements of the normal random vectors:

$$\begin{aligned} \boldsymbol{\alpha}_{\text{pheno}} &= \{\alpha_{\text{pheno}_1}, \dots, \alpha_{\text{pheno}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{pheno}} \sim \mathcal{N}(\mu_{\text{pheno}}, \sigma_{\text{pheno}}^2) \\ \boldsymbol{\alpha}_{\text{chill}} &= \{\alpha_{\text{chill}_1}, \dots, \alpha_{\text{chill}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{chill}} \sim \mathcal{N}(\mu_{\text{chill}}, \sigma_{\text{chill}}^2) \\ \boldsymbol{\alpha}_{\text{force}} &= \{\alpha_{\text{force}_1}, \dots, \alpha_{\text{force}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{force}} \sim \mathcal{N}(\mu_{\text{force}}, \sigma_{\text{force}}^2) \\ \boldsymbol{\alpha}_{\text{photo}} &= \{\alpha_{\text{photo}_1}, \dots, \alpha_{\text{photo}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{photo}} \sim \mathcal{N}(\mu_{\text{photo}}, \sigma_{\text{photo}}^2) \end{aligned} \quad (7)$$

177 Parameter $\alpha_{\text{pheno},k}$ represents the day of budburst for species k without the influence of chilling, forc-
 178 ing, or photoperiod treatments. Average day of budburst (independent of treatments) is μ_{pheno} , and
 179 σ_{pheno}^2 is the variance across species. The parameters $\alpha_{\text{chill},k}$, $\alpha_{\text{force},k}$, and $\alpha_{\text{photo},k}$ represent the trait-
 180 independent responses of species k to chilling, forcing, and photoperiod treatments respectively, each
 181 with an associated mean (μ_{chill} , μ_{force} , μ_{photo}) and variance (σ_{chill}^2 , σ_{force}^2 , σ_{photo}^2) across species. The
 182 effect of the species trait value, $\alpha_{\text{trait},k}$ (parameter shared with trait sub-model above), on its responses
 183 to chilling, forcing, and photoperiod are described by parameters $\beta_{\text{trait.chill}}$, $\beta_{\text{trait.force}}$, $\beta_{\text{trait.photo}}$. Fi-
 184 nally, σ_d^2 is the variance arising from measurement error.

185
 186 We chose weakly informative priors, and validated them using a series of prior predictive checks. The
 187 model was coded in the Stan programming language, fit using the rstan package (version 3.3.6)⁴¹,
 188 with 1,000 iterations per chain across 4 chains (4,000 total sampling iterations), and all models met
 189 basic diagnostic checks, including no divergences, high effective sample size (n_{eff}), and \hat{R} close to 1,
 190 fitting the data well (Fig S2). Here we present our model estimates as the means and 90% posterior
 191 uncertainty intervals.

192 **Results**

193 Across traits, height, SLA, and LNC strongly related to chilling (β_{chill_k}), forcing (β_{force_k}), and photoperiod (β_{photo_k}) treatments, Fig 2 a-f & j-l), but the direction of these relationships only showed
194 consistent trends for LNC (Fig. 2 j-l). As we predicted, height was negatively related to chilling (β_{chill_k})
195 and photoperiod (β_{photo_k}), with taller species having larger responses to cues (-0.5 m per standard-
196 ized chilling; 90% uncertainty interval (UI): -1.0, -0.1 and -0.2 m per standardized photoperiod; 90%
197 UI: -0.5, 0.0, Fig 2 a-c, Table S2). As illustrated for one characteristically acquisitive species, *Alnus*
198 *incana*, and one characteristically conservative species, *Quercus rubra* (Fig S3), the cue relationships
199 with height led to generally later budburst relative to estimates without trait effects ($\mu_{k,g}$; Fig. 3). In
200 contrast, seed mass had the smallest responses, with no relationship between seed mass and any cue
201 (Fig. 2 g-i, Fig 3 d-f, & Table S3).

202
203 Of our leaf traits, we found that species SLA related to photoperiod (β_{photo_k} , -0.2 mm²/mg per stan-
204 dardized photoperiod; 90% UI: -0.4, 0.0, Fig. 2 f, Table S4), but did not strongly predict responses
205 to chilling (β_{chill_k}) or forcing treatments (β_{force_k} , Fig. 2 d and e). Thus, species with more acquisitive
206 growth strategies (thin leaves and a lower investment in leaf mass that leads to large SLA values),
207 had larger responses to photoperiod, contrary to our predictions (Fig. 2 f). For LNC, we found that
208 species that produce leaves with high nitrogen content, which relates generally to high photosynthetic
209 rates and acquisitive growth, show smaller responses to cues (Fig. 2 j-l). These findings are in line
210 with our predictions that high LNC species (acquisitive) would be less responsive to chilling (0.7 mg/g
211 per standardized chilling; 90% UI: 0.3, 1.2, Table S5), but we also found high LNC species to be less
212 responsive to photoperiod (0.3 mg/g per standardized photoperiod; 90% UI: 0.0, 0.6) and to forcing
213 (0.5 mg/g per standardized forcing; 90% UI: 0.1, 0.9, Fig 2 j-l & Fig S3 d-f).

214
215 We found species-level variation across traits (σ^2_{trait}) were comparable to or greater than variation
216 across studies (σ^2_{study} , Fig 4). The magnitude of study-level variation (σ^2_{study}) that we found, however,
217 suggests that models using large trait databases that fail to separate out study from species-level vari-
218 ation (σ^2_{trait}) may poorly estimate species traits. Variation across studies was greatest for height (with
219 σ^2_{study} of 7.5 m compared to 5.9 m for σ^2_{trait} , Fig 4a). For seed mass and LNC, study-level variation
220 was less than that of the species-level variation, with estimates of 1 mg for study-level variation versus
221 1.6 mg for species-level variation in seed mass and estimates of 3.6 mg g⁻¹ for study-level variation
222 and 5.1 mg g⁻¹ for the species-level variation in LNC (Fig 4c and d). At the lowest end, study-level
223 variation in SLA was approximately half the value of the species-level variation (3.3 mm² mg⁻¹ versus
224 7.8 mm² mg⁻¹ for σ^2_{study} and σ^2_{trait} , respectively, Fig 4b).

225
226

227 **Discussion**

228 We found species traits influenced the timing of budburst in response to the three primary cues of
229 spring phenology: chilling, forcing and photoperiod. These trait effects were associated with earlier
230 or later phenology following well-established gradients in growth strategies predicted by functional
231 trait frameworks^{5,13,47,48,53}: early species tended to have traits associated with fast and acquisitive
232 strategies while later species had traits associated with conservative, slower strategies. We found the
233 largest budburst responses occurred for traits related to resource acquisition and structure, with SLA,
234 LNC, and height all showing large responses across our three cues. In contrast, our one reproductive
235 trait—seed mass—showed a smaller response. Our results provide a major step forward in integrating
236 phenology into broader trait syndromes that shape species growth strategies, and support previous
237 findings from more local scales that found strong relationships between height and species phenol-
238 ogy^{39,40,42}. Our findings also suggest other traits—such as seed mass—show no relationship with
239 phenology in our more global analysis.

241 Effects of phenology-trait relationships on community assembly

242 Our findings suggest the changing pressures across the early growing season may affect the temporal
 243 assembly of communities. Strong abiotic pressures alongside weak competition early in the season were
 244 associated with early-budbursting species with acquisitive traits (shorter heights and low LNC) that
 245 allow faster return on resource investments^{5,17,47}. These traits should allow early species to more easily
 246 replace tissue if lost to frost or other abiotic disturbances, and benefit from greater light availability in
 247 the open canopy of many temperate forests in the early spring. In contrast, later-budbursting species
 248 had traits associated with greater competitive abilities and slower growth^{5,17,47} (Fig 2), which may
 249 help them compete for soil and light resources when most other species are already growing. These
 250 traits can be linked to other ecological processes and species characteristics, such as species succes-
 251 sional position, as illustrated by the differences between early and late successional species (e.g., *Alnus*
 252 *incana* and *Quercus rubra*; Fig 2).

253

254 The traits with cue responses that deviated from our expectations also offer novel insights into the
 255 tradeoffs between traits and environmental cues. All of our traits are associated with multiple aspects
 256 of species growth, and may be adaptive for reasons other than those we predicted. Contrary to our
 257 predictions, we found large responses to forcing for short trees, which could prevent frost damage or
 258 xylem cavitation under a late spring frost^{10,30} and influence annual cambial meristem growth²⁶. Simi-
 259 larly, the lack of a response to chilling or forcing by high SLA individuals could be driven by other trait
 260 attributes and environmental cues—selecting for species relative growth rates or leaf longevity—and
 261 not photosynthetic potential^{35,47}. These findings highlight the complexity of determining the drivers
 262 of species trait profiles, and further our understanding of how traits affect community dynamics under
 263 variable environments.

264

265 Phenology-trait relationships under future climates

266 Incorporating phenology within broader trait syndromes could aid forecasting of species and commu-
 267 nity responses to climate change. While decades of research have documented phenological shifts with
 268 anthropogenic climate change, increasing research suggests a potential connection between phenolog-
 269 ical responses to warming and performance with warming, where species that shift their phenology
 270 more also perform better^{8,28}.

271

272 Our results suggest this phenology-performance relationship could be driven in part by a suite of traits
 273 that covary with phenological cues to determine how responsive species are to warming. Species with
 274 smaller responses to all cues, especially chilling and photoperiod, would tend to advance more with
 275 warming, which our results suggest would allow these species to also grow more quickly. These results
 276 could further aid in predicting the potential for invasion, as communities with similar phenologies
 277 and suites of traits, appear more susceptible to fast growing, phenologically more responsive invasive
 278 species^{1,38,49}.

279

280 Our analytical approach and results may be especially useful to help forecast changes in forest dy-
 281 namics. Identifying the varying trait syndromes of forest communities over a spring season can aid
 282 predictions of how climate change will alter species growth and productivity. For example, our results
 283 suggest that, by favoring more phenologically responsive species (i.e., with small chilling and pho-
 284 toperiod responses), warming may also favor species with acquisitive growth strategies. In contrast,
 285 conservative species, which appear less phenologically responsive to changes in temperature (due to
 286 larger chilling and photoperiod responses) could face greater abiotic and biotic stress¹⁹.

287

288 Our results could further help identify which species are most likely to be negatively impacted under
289 future climates, and develop better strategies for climate change mitigation and conservation. Species
290 that fail to advance phenologically with warming might experience more competition^{1,4}, as species that
291 begin growth increasingly earlier with warming have more time to deplete resources. In addition to
292 altering the timing and interactions between species within a season, species trait syndromes have the
293 potential to further redefine the environmental conditions under which growth occurs, and as a result,
294 shape community assembly and productivity within ecological communities. By identifying the species
295 most vulnerable to climate change impacts, we can develop more effective management practices that
296 prevent the loss of critical ecosystem services and preserve community diversity under future conditions.

297

298 **References**

- 299 [1] Alexander, J. M., and J. M. Levine. 2019. Earlier phenology of a nonnative plant increases impacts
300 on native competitors. *PNAS* 116:6199–6204.
- 301 [2] Augspurger, C. K. 2009. Spring 2007 warmth and frost : phenology , damage and refoliation in a
302 temperate deciduous forest. *Functional Ecology* 23:1031–1039.
- 303 [3] Beard, K. H., K. C. Kelsey, A. J. Leffler, and J. M. Welker. 2019. The Missing Angle : Ecosystem
304 Consequences of Phenological Mismatch. *Trends in Ecology and Evolution* 34:885–888.
- 305 [4] Carter, S. K., D. Saenz, and V. H. Rudolf. 2018. Shifts in phenological distributions reshape
306 interaction potential in natural communities. *Ecology Letters* 21:1143–1151.
- 307 [5] Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a
308 worldwide wood economics spectrum. *Ecology Letters* 12:351–366.
- 309 [6] Chuine, I., M. Bonhomme, J. M. Legave, I. García de Cortázar-Atauri, G. Charrier, A. Lacointe,
310 and T. Améglio. 2016. Can phenological models predict tree phenology accurately in the future?
311 The unrevealed hurdle of endodormancy break. *Global change biology* 22:3444–3460.
- 312 [7] Chuine, I., and P. Cour. 1999. Climatic determinants of budburst seasonality in four temperate-
313 zone tree species. *New Phytologist* 143:339–349.
- 314 [8] Cleland, E. E., J. M. Allen, T. M. Crimmins, J. A. Dunne, S. Pau, S. E. Travers, E. S. Zavaleta,
315 and E. M. Wolkovich. 2012. Phenological tracking enables positive species responses to climate
316 change.pdf. *Ecology* 93:1765–1771.
- 317 [9] Cleland, E. E., I. Chuine, A. Menzel, H. A. Mooney, and M. D. Schwartz. 2007. Shifting plant
318 phenology in response to global change. *Trends in Ecology and Evolution* 22:357–365.
- 319 [10] Clements, J. R., W. Fraser, J, and C. W. Yeatman. 1972. Frost Damage to White Spruce Buds.
320 *Canadian Journal of Forest Research* 2:62–63.
- 321 [11] Crick, H., C. Dudley, D. E. Glue, and D. L. Thomson. 1997. UK birds are laying eggs earlier.
322 *Nature* 388:526–527.
- 323 [12] D'Alba, L., P. Monaghan, and R. G. Nager. 2010. Advances in laying date and increasing popu-
324 lation size suggest positive responses to climate change in Common Eiders Somateria mollissima
325 in Iceland. *International Journal of Avian Science* 152:19–28.
- 326 [13] Díaz, S., J. Kattge, J. H. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer,
327 C. Wirth, I. Colin Prentice, E. Garnier, G. Bönisch, M. Westoby, H. Poorter, P. B. Reich, A. T.
328 Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright, S. N. Sheremet Ev,
329 H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig,
330 A. Günther, V. Falcuk, N. Rüger, M. D. Mahecha, and L. D. Gorné. 2016. The global spectrum
331 of plant form and function. *Nature* 529:167–171.
- 332 [14] Ettinger, A. K., C. J. Chamberlain, I. Morales-Castilla, D. M. Buonaiuto, D. F. Flynn, T. Savas,
333 J. A. Samaha, and E. M. Wolkovich. 2020. Winter temperatures predominate in spring phenolog-
334 ical responses to warming. *Nature Climate Change* 10:1137–1142.
- 335 [15] Flynn, D. F. B., and E. M. Wolkovich. 2018. Temperature and photoperiod drive spring phenology
336 across all species in a temperate forest community. *New Phytologist* 219:1353–1362.
- 337 [16] Gorné, L. D., S. Díaz, V. Minden, Y. Onoda, K. Kramer, C. Muir, S. T. Michaletz, S. Lavorel,
338 J. Sharpe, S. Jansen, M. Slot, E. Chacon, and G. Boenisch. 2020. The acquisitive–conservative
339 axis of leaf trait variation emerges even in homogeneous environments. *Annals of Botany* .

- 340 [17] Grime, J. P. 1977. Evidence for the Existence of Three Primary Strategies in Plants and Its
341 Relevance to Ecological and Evolutionary Theory Author (s): J . P . Grime Source : The
342 American Naturalist , Vol . 111 , No . 982 (Nov . - Dec ., 1977), pp . 1169-1194 Published. The
343 American Naturalist 111:1169–1194.
- 344 [18] Gu, H., Y. Qiao, Z. Xi, S. Rossi, N. G. Smith, J. Liu, and L. Chen. 2022. Warming-induced
345 increase in carbon uptake is linked to earlier spring phenology in temperate and boreal forests.
346 Nature Communications 13:1–8.
- 347 [19] Guy, R. D. 2014. The early bud gets to warm. New Phytologist 202:7–9.
- 348 [20] Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying
349 dormancy requirements for Pacific Northwest tree species. Frontiers in Plant Science 6:1–12.
- 350 [21] Hufkens, K., M. A. Friedl, T. F. Keenan, O. Sonnentag, A. Bailey, J. O'keefe, and A. D. Richardson.
351 2012. Ecological impacts of a widespread frost event following early spring leaf-out. Global
352 Change Biology 18:2365–2377.
- 353 [22] Kattge, J., G. Bönisch, S. Díaz, S. Lavorel, I. C. Prentice, and et al. 2020. TRY plant trait
354 database – enhanced coverage and open access. Global Change Biology 26:119–188.
- 355 [23] Kusano, T., and M. Inoue. 2008. Long-Term Trends toward Earlier Breeding of Japanese Am-
356 phibians. Journal of Herpetology 42:608–614.
- 357 [24] Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling
358 outweighs photoperiod in preventing precocious spring development. Global Change Biology
359 20:170–182.
- 360 [25] Laughlin, D. C., J. J. Leppert, M. M. Moore, and C. H. Sieg. 2010. A multi-trait test of the leaf-
361 height-seed plant strategy scheme with 133 species from a pine forest flora. Functional Ecology
362 24:493–501.
- 363 [26] Lenz, A., G. Hoch, C. Körner, and Y. Vitasse. 2016. Convergence of leaf-out towards minimum
364 risk of freezing damage in temperate trees. Functional Ecology 30:1480–1490.
- 365 [27] Lopez, O. R., K. Farris-Lopez, R. A. Montgomery, and T. J. Givnish. 2008. Leaf phenology in
366 relation to canopy closure in southern Appalachian trees. American Journal of Botany 95:1395–
367 1407.
- 368 [28] Macgregor, C. J., C. D. Thomas, D. B. Roy, M. A. Beaumont, J. R. Bell, T. Brereton, J. R. Bridle,
369 C. Dytham, R. Fox, K. Gotthard, A. A. Hoffmann, G. Martin, I. Middlebrook, S. Nylin, P. J.
370 Platts, R. Rasteiro, I. J. Saccheri, R. Villoutreix, C. W. Wheat, and J. K. Hill. 2019. Climate-
371 induced phenology shifts linked to range expansions in species with multiple reproductive cycles
372 per year. Nature Communications 10:1–10.
- 373 [29] Maitner, B. S., B. Boyle, N. Casler, R. Condit, J. Donoghue, S. M. Durán, D. Guaderrama, C. E.
374 Hinchliff, P. M. Jørgensen, N. J. Kraft, B. McGill, C. Merow, N. Morueta-Holme, R. K. Peet,
375 B. Sandel, M. Schildhauer, S. A. Smith, J. C. Svenning, B. Thiers, C. Violette, S. Wiser, and B. J.
376 Enquist. 2018. The bien r package: A tool to access the Botanical Information and Ecology
377 Network (BIEN) database. Methods in Ecology and Evolution 9:373–379.
- 378 [30] Marquis, B., Y. Bergeron, M. Simard, and F. Tremblay. 2020. Growing-season frost is a better
379 predictor of tree growth than mean annual temperature in boreal mixedwood forest plantations.
380 Global Change Biology 26:6537–6554.

- 381 [31] Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kübler, P. Bissolli,
 382 O. Braslavská, A. Briede, F. M. Chmielewski, Z. Crepinsek, Y. Curnel, Å. Dahl, C. Defila, A. Don-
 383 nnelly, Y. Filella, K. Jatczak, F. Måge, A. Mestre, Ø. Nordli, J. Peñuelas, P. Pirinen, V. Remišová,
 384 H. Scheifinger, M. Striz, A. Susnik, A. J. Van Vliet, F. E. Wielgolaski, S. Zach, and A. Zust. 2006.
 385 European phenological response to climate change matches the warming pattern. *Global Change
 386 Biology* 12:1969–1976.
- 387 [32] Morales-Castilla, I., T. J. Davies, G. Legault, D. M. Buonaiuto, C. J. Chamberlain, A. K. Ettinger,
 388 M. Garner, F. A. M. Jones, D. Loughnan, W. D. Pearse, D. Sodhi, and E. M. Wolkovich. 2024.
 389 Phylogenetic estimates of species-level phenology improve ecological forecasting. *Nature Climate
 390 Change* .
- 391 [33] Ovaskainen, O., S. Skorokhodova, M. Yakovleva, A. Sukhov, A. Kutenkov, N. Kutenkova,
 392 A. Shcherbakov, E. Meyke, and M. del Mar Delgado. 2013. Community-level phenological re-
 393 sponse to climate change. *PNAS* 110:13434–13439.
- 394 [34] Rathcke, B., and E. P. Lacey. 1985. Phenological patterns of terrestrial plants. *Annual Review of
 395 Ecology and Systematics* 16:179–214.
- 396 [35] Reich, P. B. 2014. The world-wide ‘ fast – slow ’ plant economics spectrum : a traits manifesto.
 397 *Journal of Ecology* 102:275–301.
- 398 [36] Reich, P. B., D. S. Ellsworth, M. B. Walters, J. M. Vose, C. Gresham, J. C. Volin, and W. D.
 399 Bowman. 1999. Generality of leaf trait relationships: A test across six biomes. *Ecology* 80:1955–
 400 1969.
- 401 [37] Sakai, A., and W. Larcher. 1987. Frost Survival of Plants: Responses and adaptation to freezing
 402 stress. Springer-Verlag, Berlin, Heidelberg.
- 403 [38] Schuster, M. J., P. D. Wragg, and P. B. Reich. 2021. Phenological niche overlap between invasive
 404 buckthorn (*Rhamnus cathartica*) and native woody species. *Forest Ecology and Management*
 405 498:119568.
- 406 [39] Segrestin, J., M.-l. Navas, and E. Garnier. 2020. Reproductive phenology as a dimension of the
 407 phenotypic space in 139 plant species from the Mediterranean. *New Phytologist* 225:740–753.
- 408 [40] Sporbert, M., D. Jakubka, S. F. Bucher, I. Hensen, M. Freiberg, K. Heubach, A. Konig, B. Nordt,
 409 C. Plos, I. Blinova, A. Bonn, B. Knickmann, T. Koubek, A. Linstadter, T. Maskova, R. Primack,
 410 C. Rosche, M. A. Shah, A.-D. Stevens, K. Teilborger, S. Trager, C. Wirth, and C. Romermann.
 411 2022. Functional traits influence patterns in vegetative and reproductive plant phenology – a
 412 multi-botanical garden study. *New Phytologist* 235:2199–2210.
- 413 [41] Stan Development Team. 2018. RStan: the R interface to Stan. R package version 2.17.3.
- 414 [42] Sun, S., and L. E. Frelich. 2011. Flowering phenology and height growth pattern are associated
 415 with maximum plant height, relative growth rate and stem tissue mass density in herbaceous
 416 grassland species. *Journal of Ecology* 99:991–1000.
- 417 [43] Tryjanowski, P., M. Rybacki, and T. Sparks. 2003. Changes in the first spawning dates of common
 418 frogs and common toads in western Poland in 1978-2002. *Annales Zoologici Fennici* 40:459–464.
- 419 [44] Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung, and J. Messier.
 420 2012. The return of the variance: Intraspecific variability in community ecology. *Trends in Ecology
 421 and Evolution* 27:244–252.

- 422 [45] Vitasse, Y., S. Ursenbacher, G. Klein, T. Bohnenstengel, Y. Chittaro, A. Delestrade, C. Monnerat,
423 M. Rebetez, C. Rixen, N. Strelbel, B. R. Schmidt, S. Wipf, T. Wohlgemuth, N. G. Yoccoz, and
424 J. Lenoir. 2021. Phenological and elevational shifts of plants , animals and fungi under climate
425 change in the European Alps. *Biological Reviews* 96:1816–1835.
- 426 [46] Wainwright, C. E., E. M. Wolkovich, and E. E. Cleland. 2012. Seasonal priority effects : implications
427 for invasion and restoration in a semi-arid system. *Journal of Applied Ecology* 49:234–241.
- 428 [47] Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*
429 199:213–227.
- 430 [48] Westoby, M., and I. J. Wright. 2006. Land-plant ecology on the basis of functional traits. *Trends
431 in Ecology and Evolution* 21:261–268.
- 432 [49] Wolkovich, E. M., and E. E. Cleland. 2011. The phenology of plant invasions: A community
433 ecology perspective. *Frontiers in Ecology and the Environment* 9:287–294.
- 434 [50] ———. 2014. Phenological niches and the future of invaded ecosystems with climate change. *AoB
435 PLANTS* 6:1–16.
- 436 [51] Wolkovich, E. M., and M. J. Donahue. 2021. How phenological tracking shapes species and
437 communities in non-stationary environments. *Biological Reviews* 96:2810–2827.
- 438 [52] Wolkovich, E. M., and A. K. Ettinger. 2014. Back to the future for plant phenology research. *New
439 Phytologist* 203:1021–1024.
- 440 [53] Wright, I. J., M. Westoby, P. B. Reich, J. Oleksyn, D. D. Ackerly, Z. Baruch, F. Bongers,
441 J. Cavender-Bares, T. Chapin, J. H. C. Cornellissen, M. Diemer, J. Flexas, J. Gulias, E. Garnier,
442 M. L. Navas, C. Roumet, P. K. Groom, B. B. Lamont, K. Hikosaka, T. Lee, W. Lee, C. Lusk, J. J.
443 Midgley, Ü. Niinemets, H. Osada, H. Poorter, P. Pool, E. J. Veneklaas, L. Prior, V. I. Pyankov,
444 S. C. Thomas, M. G. Tjoelker, and R. Villar. 2004. The worldwide leaf economics spectrum.
445 *Nature* 428:821–827.




Figure 1: Leaf and wood traits follow a gradient that varies from acquisitive versus conservative growth strategies, which may also include phenology. We predicted early-budbursting species would have traits associated with acquisitive growth, as they are more likely to experience greater risk of frost but reduced competition. In contrast, we expected later-budbursting species would have traits associated with conservative growth, as they experience greater competition but a more climatically benign environment.




Figure 2: Estimated species-level responses to standardized forcing (left column: a, d, g & j), chilling (middle column: b, e, h & k), and photoperiod treatments (right column: c, f, i & l) predicted by estimated trait values for height (first row: a-c), SLA (second row: d-f), log10 Seed mass (third row: g-i), and LNC (fourth row: j-l). We estimated parameters using a joint trait-phenology model, with the black line depicting the mean linear relationship between estimated trait effects and the slope of the cue response and the pink band the 50% uncertainty interval. Each set of crossed lines represents one species (with each line spanning the 50% uncertainty interval), with the species depicted in Fig 3 colored in each panel, with the acquisitive species (*Alnus incana*) shown in red, and the conservative species (*Quercus rubra*) shown in blue.




Figure 3: We expected species with traits associated with acquisitive (e.g., smaller heights and small seed mass) versus conservative (e.g., taller with larger seeds) growth strategies would have different budburst responses to phenological cues. Shown here is an example of the cue relationships with height (**a-c**) and seed mass (**d-f**) for an acquisitive species, *Alnus incana* shown in red, and a conservative species, *Quercus rubra*, shown in blue. **a**, The effect of height on budburst timing was smaller in response to forcing cues, but larger in response to both **b**, chilling and **c**, photoperiod. In contrast, seed mass had a negligible effect on **d**, forcing and **f**, photoperiod responses, **e**, but a greater response to chilling. Points represent the raw data and the coloured bands the 50% uncertainty intervals of the model estimates and points individual trait measurements.




Figure 4: Traits differed in the relative magnitude of their species-level and study-level variation, with only **a**, the height model estimating greater study-level variation than species-level, while our **b**, specific leaf area, **c**, leaf nitrogen content, and **d**, seed mass models all estimated higher species-level variation. Shown here are the posterior densities for the species-level variation, shown in red, and study-level variation, shown in yellow. For comparison, we show all x-axes spanning the minimum to maximum variances across all four traits. The histograms depict the full distribution of the study and species-level variance, where each bin is normalized by the total count of the posterior estimate.