

<sub>1</sub> Supplementary Material: Budburst timing within a functional  
<sub>2</sub> trait framework

<sub>3</sub>

<sub>4</sub> **Figures & Tables**




Figure S1: Our initial aim was to include all species from Morales-Castilla *et al.* (2024), however trait data was only available for a subset of these species. Data was cleaned extensively, with each step depicted by a box, and subset to only include traits for adult trees growing under natural conditions for which we had a complete suite of trait values.




Figure S2: A projection of tree traits across the first and second principle component axis. Arrows represent the direction of vectors for the six functional traits with complete trait data. Points represent the 26 species for which complete trait data was available.




Figure S3: We obtained data from the TRY and BIEN plant trait databases and the OSPREE database of plant phenology experiments for temperate tree species. Following our cleaning of the data and selection of uncorrelated traits, our final dataset included 37 species from 24 unique datasources for the trait data and 34 unique studies for the budburst data. Our data is focused on temperate ecosystems globally with most data originating from North America and Europe.




Figure S4: For most traits, we found the species-level model posterior distributions from our joint model of trait effects on budburst cues were well-aligned with the raw data for our 37 woody deciduous plant species. Four functional traits—(a) height, (b) SLA, (c) seed mass, and (d) LNC—were modeled individually, with the calculated trait value being used to jointly model species' responses to standardized chilling, forcing, and photoperiod cues. Model posteriors are shown in black, with the thicker line depicting the 50% uncertainty interval and the thinner black line the 95% uncertainty interval. The raw data is shown in red and the species-level means from the raw data are denoted as a purple stars.

Table S1: Bibliographic information for trait data sources from both BIEN and TRY trait databases. Datasets without references or incomplete references are denoted below as ‘unreferenced’.

Database	Reference	Trait name	Unit	No. observations	No. Species
BIEN	Mchugh <i>et al.</i> (2015)	Height	m	26	8
BIEN	Marx <i>et al.</i> (2016)	Height	m	2	2
BIEN	Price <i>et al.</i> (2014)	Height	m	27	19
BIEN	Unreferenced	Height	m	18	16
BIEN	Kleyer <i>et al.</i> (2008)	Height	m	90	19
BIEN	Unreferenced	Height	m	10	10
BIEN	Moles, Angela; unreferenced	Height	m	21	14
BIEN	Reams, Greg; unreferenced	Height	m	47036	19
BIEN	Grime, Hodgson, & Hunt; unreferenced	Height	m	5	5
BIEN	Unreferenced	Height	m	8	5
BIEN	Pérez-de Lis <i>et al.</i> (2017)	Height	m	18	1
BIEN	Robinson <i>et al.</i> (2015)	Height	m	120	1
BIEN	Anderson-teixeira <i>et al.</i> (2015)	Height	m	20	1
TRY	Bond-Lamberty <i>et al.</i> (2002)	Height	m	2	1
TRY	Unpublished	Height	m	275	3
TRY	Wright <i>et al.</i> (2004)	Height	m	28	19
TRY	Prentice <i>et al.</i> (2011)	Height	m	2	2
TRY	Schweingruber & Landolt (2010)	Height	m	21	21
TRY	Unpublished	Height	m	35	2
TRY	Moles <i>et al.</i> (2004)	Height	m	5	5
TRY	Cavender-Bares <i>et al.</i> (2006)	Height	m	1	1
TRY	Diaz <i>et al.</i> (2004)	Height	m	11	10
TRY	Craine <i>et al.</i> (2009)	LNC	mg/g	287	12
TRY	Wilson <i>et al.</i> (2000)	LNC	mg/g	44	2
TRY	Wenxuan <i>et al.</i> (2012)	LNC	mg/g	7	4
TRY	Yahan <i>et al.</i> (2013)	LNC	mg/g	7	3
TRY	Wright <i>et al.</i> (2004)	LNC	mg/g	65	32
TRY	Prentice <i>et al.</i> (2011)	LNC	mg/g	3	2
TRY	Vergutz <i>et al.</i> (2012)	LNC	mg/g	120	20
TRY	Atkin <i>et al.</i> (2015)	LNC	mg/g	24	8
TRY	Marie <i>et al.</i> (2015)	LNC	mg/g	72	22
TRY	Cornelissen <i>et al.</i> (2003)	LNC	mg/g	2	1
TRY	Unpublished	LNC	mg/g	3216	37
TRY	Wang <i>et al.</i> (2017)	LNC	mg/g	6	2
BIEN	Marx <i>et al.</i> (2016)	Seed mass	mg	3	3
BIEN	Unreferenced	Seed mass	mg	4	2
BIEN	Liu <i>et al.</i> (2018)	Seed mass	mg	250	37
BIEN	Ameztegui <i>et al.</i> (2017)	Seed mass	mg	12	12
BIEN	Paine <i>et al.</i> (2015)	Seed mass	mg	12	7
TRY	Wilson <i>et al.</i> (2000)	SLA	mm <sup>2</sup> mg-1	44	2
TRY	Unpublished	SLA	mm <sup>2</sup> mg-1	204	3
TRY	Wright <i>et al.</i> (2004)	SLA	mm <sup>2</sup> mg-1	93	33
TRY	Prentice <i>et al.</i> (2011)	SLA	mm <sup>2</sup> mg-1	2	2
TRY	Kleyer <i>et al.</i> (2008)	SLA	mm <sup>2</sup> mg-1	102	18
TRY	Unpublished	SLA	mm <sup>2</sup> mg-1	83	2
TRY	Atkin <i>et al.</i> (2015)	SLA	mm <sup>2</sup> mg-1	40	11
TRY	Marie <i>et al.</i> (2015)	SLA	mm <sup>2</sup> mg-1	86	23
TRY	Cornelissen <i>et al.</i> (2003)	SLA	mm <sup>2</sup> mg-1	615	14
TRY	Unpublished	SLA	mm <sup>2</sup> mg-1	6307	37
TRY	Wang <i>et al.</i> (2017)	SLA	mm <sup>2</sup> mg-1	6	2
TRY	Shipley & Vu (2002)	SLA	mm <sup>2</sup> mg-1	20	2
TRY	Cavender-Bares <i>et al.</i> (2006)	SLA	mm <sup>2</sup> mg-1	42	2
TRY	Unpublished	SLA	mm <sup>2</sup> mg-1	1	1
TRY	Diaz <i>et al.</i> (2004)	SLA	mm <sup>2</sup> mg-1	11	10

Table S2: Summary of estimates from our model of height ( $n = 42781$ ) using data from 37 focal species. Values represent the mean estimate for model parameters as well as the lower and upper limits (25%–75%) of the 50% uncertainty interval and the lower and upper limits (5%–95%) of the 90% uncertainty interval (UI). The strength of the relationship can be assessed by comparing across UI, with parameters with intervals that cross zero depicting weak relationships.

	mean	5%	25%	75%	95%
$\mu_{grand.trait}$	13.3	10.0	11.9	14.6	16.5
$\mu_{k,g}$	32.1	27.7	30.3	33.9	36.6
$\mu_{force}$	-10.7	-15.2	-12.5	-8.8	-6.2
$\mu_{chill}$	-4.0	-11.1	-6.5	-1.1	2.8
$\mu_{photo}$	1.0	-2.7	-0.5	2.5	4.7
$\beta_{trait.force}$	0.2	-0.1	0.0	0.3	0.5
$\beta_{trait.chill}$	-0.5	-1.0	-0.7	-0.3	-0.1
$\beta_{trait.photo}$	-0.2	-0.5	-0.4	-0.1	0.0
$\sigma_{species}$	6.0	4.9	5.5	6.5	7.3
$\sigma_{study}$	7.5	5.7	6.6	8.3	9.8
$\sigma_{trait}$	5.4	5.4	5.4	5.4	5.4
$\sigma_{pheno}$	15.1	11.8	13.6	16.4	18.6
$\sigma_{force}$	5.0	3.3	4.2	5.7	7.0
$\sigma_{chill}$	8.8	5.8	7.2	10.0	12.7
$\sigma_{photo}$	3.5	2.2	2.9	4.1	5.3
$\sigma_d$	14.2	13.8	14.	14.3	14.6

Table S3: Summary of estimates from our model of seed mass ( $n = 281$ ) using data from 37 focal species. Values represent the mean estimate for model parameters as well as the lower and upper limits (25%–75%) of the 50% uncertainty interval and the lower and upper limits (5%–95%) of the 90% uncertainty interval (UI). The strength of the relationship can be assessed by comparing across UI, with parameters with intervals that cross zero depicting weak relationships.

	mean	5%	25%	75%	95%
$\mu_{grand.trait}$	1.9	1.0	1.5	2.2	2.7
$\mu_{k,g}$	31.4	27.1	29.6	33.1	35.8
$\mu_{force}$	-8.2	-10.8	-9.2	-7.1	-5.6
$\mu_{chill}$	-9.4	-14.0	-11.2	-7.5	-4.8
$\mu_{photo}$	-1.3	-3.4	-2.1	-0.4	0.8
$\beta_{trait.force}$	-0.3	-1.4	-0.8	0.1	0.9
$\beta_{trait.chill}$	-1.1	-2.9	-1.8	-0.4	0.7
$\beta_{trait.photo}$	-0.6	-1.5	-0.9	-0.2	0.4
$\sigma_{species}$	1.6	1.3	1.5	1.7	2.0
$\sigma_{study}$	1.0	0.8	0.9	1.0	1.1
$\sigma_{trait}$	0.2	0.2	0.2	0.3	0.3
$\sigma_{pheno}$	14.8	11.2	13.3	16.3	18.6
$\sigma_{force}$	4.9	3.4	4.2	5.5	6.7
$\sigma_{chill}$	10.7	7.1	8.8	12.1	15.4
$\sigma_{photo}$	3.6	2.3	3.0	4.1	5.1
$\sigma_d$	14.1	13.7	14.0	14.3	14.5

Table S4: Summary of estimates from our model of specific leaf area (SLA,  $n = 7656$ ) using data from 37 focal species. Values represent the mean estimate for model parameters as well as the lower and upper limits (25%–75%) of the 50% uncertainty interval and the lower and upper limits (5%–95%) of the 90% uncertainty interval (UI). The strength of the relationship can be assessed by comparing across UI, with parameters with intervals that cross zero depicting weak relationships.

	mean	5%	25%	75%	95%
$\mu_{grand.trait}$	16.8	14.5	15.8	17.8	19.3
$\mu_{k,g}$	31.3	27.2	29.6	33.0	35.6
$\mu_{force}$	-11.4	-16.2	-13.1	-9.6	-7.2
$\mu_{chill}$	-16.7	-24.6	-19.8	-13.3	-9.1
$\mu_{photo}$	1.9	-2.4	0.2	3.6	5.8
$\beta_{trait.force}$	0.2	-0.1	0.1	0.3	0.4
$\beta_{trait.chill}$	0.3	-0.1	0.2	0.5	0.7
$\beta_{trait.photo}$	-0.2	-0.5	-0.3	-0.1	0.0
$\sigma_{species}$	7.8	6.4	7.1	8.4	9.4
$\sigma_{study}$	3.3	2.0	2.6	3.8	5.1
$\sigma_{trait}$	6.2	6.1	6.1	6.2	6.3
$\sigma_{pheno}$	13.9	10.7	12.4	15.3	17.6
$\sigma_{force}$	5.0	3.3	4.2	5.6	7.0
$\sigma_{chill}$	10.6	7.2	8.9	12.0	14.8
$\sigma_{photo}$	3.5	2.3	2.9	3.9	5.0
$\sigma_d$	14.2	13.8	14.0	14.3	14.6

Table S5: Summary of estimates from our model of leaf nitrogen content (LNC,  $n = 3853$ ) using data from 37 focal species. Values represent the mean estimate for model parameters as well as the lower and upper limits (25%–75%) of the 50% uncertainty interval and the lower and upper limits (5%–95%) of the 90% uncertainty interval (UI). The strength of the relationship can be assessed by comparing across UI, with parameters with intervals that cross zero depicting weak relationships.

	mean	5%	25%	75%	95%
$\mu_{grand.trait}$	22.6	20.4	21.7	23.5	24.9
$\mu_{k,g}$	31.1	27.1	29.4	32.8	35.3
$\mu_{force}$	-19.3	-27.9	-22.9	-15.8	-10.5
$\mu_{chill}$	-27.1	-38.5	-31.7	-22.4	-15.4
$\mu_{photo}$	-9.4	-17.0	-12.5	-6.2	-1.9
$\beta_{trait.force}$	0.5	0.1	0.3	0.6	0.8
$\beta_{trait.chill}$	0.7	0.2	0.5	0.9	1.2
$\beta_{trait.photo}$	0.3	0.0	0.2	0.4	0.6
$\sigma_{species}$	5.1	4.2	4.7	5.5	6.2
$\sigma_{study}$	3.6	2.2	2.9	4.1	5.3
$\sigma_{trait}$	5.1	5.0	5.1	5.2	5.2
$\sigma_{pheno}$	14.0	10.9	12.7	15.3	17.4
$\sigma_{force}$	4.6	3.0	3.8	5.2	6.6
$\sigma_{chill}$	8.9	6.1	7.5	10.1	12.5
$\sigma_{photo}$	3.6	2.4	3.0	4.1	5.0
$\sigma_d$	14.2	13.8	14.0	14.3	14.6



## 5 Stan model code

```

6
7 data {
8   int<lower = 1> n_spec;
9   int<lower = 1> N;
10  int<lower = 1, upper = n_spec> trait_species[N];
11  int<lower = 1> n_study;
12  int<lower = 1, upper = n_study> study[N];
13  vector[N] yTraiti;
14
15  int<lower = 1> Nph;
16  int<lower = 1, upper = n_spec> phenology_species[Nph];
17  vector[Nph] yPhenoi;
18  vector[Nph] forcei;
19  vector[Nph] chilli;
20  vector[Nph] photoi;
21 }
22
23 parameters{
24
25   real mu_grand;
26   vector[n_spec] muSp;
27   vector[n_study] muStudy;
28   real<lower = 0> sigma_traity;
29   real<lower = 0> sigma_sp;
30   real<lower = 0> sigma_study;
31
32   real alphaForceSp[n_spec];
33   real muForceSp;
34   real<lower = 0> sigmaForceSp;
35   real alphaChillSp[n_spec];
36   real muChillSp;
37   real<lower = 0> sigmaChillSp;
38   real alphaPhotoSp[n_spec];
39   real muPhotoSp;
40   real<lower = 0> sigmaPhotoSp;
41   real alphaPhenoSp[n_spec];
42   real muPhenoSp;
43   real<lower = 0> sigmaPhenoSp;
44   real betaTraitxForce;
45   real betaTraitxChill;
46   real betaTraitxPhoto;
47   real<lower = 0> sigmaphenoy;
48 }
49
50 transformed parameters{
51
52   vector[N] y_hat;
53   vector[n_spec] mu_grand_sp;
54
55   real betaForceSp[n_spec];

```

```

56   real betaPhotoSp[n_spec];
57   real betaChillSp[n_spec];
58
59   for(i in 1:n_spec){
60     mu_grand_sp[i] = mu_grand + muSp[i];
61   }
62   for (i in 1:N){
63     y_hat[i] = mu_grand + muSp[trait_species[i]] + muStudy[study[i]];
64   }
65
66   for (isp in 1:n_spec){
67     betaForceSp[isp] = alphaForceSp[isp] + betaTraitxForce * (mu_grand_sp[isp]);
68   }
69   for (isp in 1:n_spec){
70     betaPhotoSp[isp] = alphaPhotoSp[isp] + betaTraitxPhoto * (mu_grand_sp[isp]);
71   }
72   for (isp in 1:n_spec){
73     betaChillSp[isp] = alphaChillSp[isp] + betaTraitxChill * (mu_grand_sp[isp]);
74   }
75 }
76
77 model{
78   yTraiti ~ normal(y_hat, sigma_traity);
79   muSp ~ normal(0, sigma_sp);
80   muStudy ~ normal(0, sigma_study);
81   mu_grand ~ normal(20,10);
82   sigma_sp ~ normal(4,5);
83   sigma_study ~ normal(2,5);
84   sigma_traity ~ normal(3,5);
85
86   for (i in 1:Nph){
87     yPhenoi[i] ~ normal(alphaPhenoSp[phenology_species[i]] +
88       betaForceSp[phenology_species[i]] * forcei[i] +
89       betaPhotoSp[phenology_species[i]] * photoi[i] +
90       betaChillSp[phenology_species[i]] * chilli[i],
91       sigmapheno_y);
92   }
93
94   alphaPhenoSp ~ normal(muPhenoSp, sigmaPhenoSp);
95   alphaForceSp ~ normal(muForceSp, sigmaForceSp);
96   alphaChillSp ~ normal(muChillSp, sigmaChillSp);
97   alphaPhotoSp ~ normal(muPhotoSp, sigmaPhotoSp);
98
99   muPhenoSp ~ normal(40,10);
100  sigmaPhenoSp ~ normal(5,5);
101
102  sigmapheno_y ~ normal(10,5);
103
104  muForceSp ~ normal(-15,10);
105  sigmaForceSp ~ normal(5,5);
106
107  muChillSp ~ normal(-15,10);

```

```

108     sigmaChillSp ~ normal(5,5);
109
110     muPhotoSp ~ normal(-15,10);
111     sigmaPhotoSp ~ normal(5,5);
112
113     betaTraitxForce ~ normal(0,1);
114     betaTraitxPhoto ~ normal(0,1);
115     betaTraitxChill ~ normal(0,1);
116
117 }
118

```

## References

- Ameztegui, A., Paquette, A., Shipley, B., Heym, M., Messier, C. & Gravel, D. (2017) Shade tolerance and the functional trait : demography relationship in temperate and boreal forests. *Functional Ecology* **31**, 821–830.
- Anderson-teixeira, K.J., Mcgarvey, J.C., Muller-landau, H.C., Park, J.Y., Gonzalez-akre, E.B., Herrmann, V., Bennett, A.C., So, C.V., Bourg, N.A., Thompson, J.R., McMahon, S.M. & Mcshea, W.J. (2015) Size-related scaling of tree form and function in a mixed-age forest. *Functional Ecology* **29**, 1587–1602.
- Atkin, O., Bloomfield, K., Reich, P., Tjoelker, M., Asner, G., Bonal, D., Bönisch, G., Bradford, M., Cernusak, L., Cosio, E., Creek, D., K.Y., C., Domingues, T., Dukes, J., Egerton, J., Evans, J., Farquhar, G., Fyllas, N., Gauthier, P., Gloor, E., Gimeno, T., Griffin, K., Guerrieri, R., Heskell, M., Huntingford, C., Ishida, F., Kattge, J., Lambers, H., Liddell, M., Lloyd, J., Lusk, C., Martin, R., Maksimov, A., Maximov, T., Malhi, Y., Medlyn, B., Meir, P., Mercado, L., Mirotchnick, N., Ng, D., Niinemets, Ü., O’Sullivan, O., Phillips, O., Poorter, L., Poot, P., Prentice, I., Salinas, N., Rowland, L., Ryan, M., Sitch, S., Slot, M., Smith, N., Turnbull, M., VanderWel, M., Valladares, F., Veneklaas, E., Weerasinghe, L., Wirth, C., Wright, I., Wythers, K., Xiang, J., Xiang, S. & Zaragoza-Castells, J. (2015) Global variability in leaf respiration in relation to climate, plant functional types and leaf traits. *New Phytologist* **206**, 614–636.
- Bond-Lamberty, B., Wang, C. & Gower, S.T. (2002) Aboveground and belowground biomass and sapwood area allometric equations for six boreal tree species of northern Manitoba. *Canadian Journal of Forest Research* **32**, 1441–1450.
- Cavender-Bares, J., Keen, A. & Miles, B. (2006) Phylogenetic structure of floridian plant communities depends on taxonomic and spatial scale. *Ecology* **87**, 109–122.
- Cornelissen, J.H.C., Cerabolini, B., Castro-Diez, P., Villar-Salvador, P., Montserrat-Marti, G., Puyravaud, J.P., Maestro, M., Werger, M.J.A. & Aerts, R. (2003) Functional traits of woody plants: correspondence of species rankings between field adults and laboratory-grown seedlings? *Journal of Vegetation Science* **14**, 311–322.
- Craine, J.M., Elmore, A.J., Aidar, M.P.M., Bustamante, M., Dawson, T.E., Hobbie, E.A., Kahmen, A., Mack, M.C., Mclauchlan, K.K., Michelsen, A., Nardoto, G.B., Pardo, L.H., Penuelas, J., Reich, P.B., Schuur, E.A.G., Stock, W.D., Templer, P.H., Virginia, R.A., Welker, J.M. & Wright, I.J. (2009) Global patterns of foliar nitrogen isotopes and their relationships with climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New Phytologist* **183**, 980–992.
- Diaz, S., Hodgson, J.G., Thompson, K., Cabido, M., Cornelissen, J.H.C., Jalili, A., Montserrat-Marti, G., Grime, J.P., Zarrinkamar, F., Asri, Y., Band, S.R., Basconcelo, S., Castro-Diez, P., Funes, G., Hamzehee, B., Khoshnevi, M., Pérez-Harguindeguy, N., Pérez-Rontomé, M.C., Shirvany, F.A., Vendramini, F., Yazdani, S., Abbas-Azimi, R., Bogaard, A., Boustani, S., Charles, M., Dehghan, M., de Torres-Espuny, L., Falczuk, V., Guerrero-Campo, J., Hynd, A., Jones, G., Kowsary, E., Kazemi-Saeed, F., Maestro-Martinez, M., Romo-Diez, A., Shaw, S., Siavash, B., Villar-Salvador, P. & Zak, M.R. (2004) The plant traits that drive ecosystems : Evidence from three continents. *Journal of Vegetation Science* **15**, 295–304.
- Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M., Poschlod, P., Groenendaal, J.M.V., Klime, L., Klimesova, J., Klotz, S., Rusch, G.M., Hermy, M., Adriaens, D., Boedeltje, G., Bossuyt, B., Dannemann, A., Endels, P., Götzenberger, L., Hodgson, J.G., Jackel, A.k., Kühn, I., Kunzmann, D., Ozinga, W.A., Römermann, C., Stadler, M., Schlegelmilch, J., Steendam, H.J., Tackenberg, O., Wilmann, B., Cornelissen, J.H.C., Eriksson, O., Garnier, E. & Peco, B. (2008) The LEDA Traitbase : a database of life-history traits of the Northwest European flora. *Journal of Ecology* **96**, 1266–1274.

166 Liu, K., Eastwood, R.J. and Flynn, S., Turner, R. & Stuppy, W. (2018) Kew database.

167 Marie, V., Wright, I.J., Prentice, I.C., Batjes, N.H., Bhaskar, R., van Bodegom, P.M., Cornwell, W.K.,  
 168 Ellsworth, D., Niinemets, Ü., Ordonez, A., Reich, P.B. & Santiago, L.S. (2015) Global effects of soil  
 169 and climate on leaf photosynthetic traits and rates. *Global Ecology and Biogeography* **24**, 706–717.

170 Marx, H.E., Giblin, D.E., Dunwiddie, P.W. & Tank, D.C. (2016) Deconstructing Darwin’s Naturaliza-  
 171 tion using community phylogenetics and functional traits. *Diversity and Distributions* **22**, 318–331.

172 Mchugh, N., Edmondson, J.L., Gaston, K.J., Leake, J.R. & Sullivan, O.S.O. (2015) Modelling short-  
 173 rotation coppice and tree planting for urban carbon management – a citywide analysis. *Journal of*  
 174 *Applied Ecology* **52**, 1237–1245.

175 Moles, A.T., Falster, D.S., Leishman, M.R. & Westoby, M. (2004) Small-seeded species produce more  
 176 seeds per square metre of canopy per year, but not per individual per lifetime. *Journal of Ecology*  
 177 **92**, 384–396.

178 Morales-Castilla, I., Davies, T.J., Legault, G., Buonaiuto, D.M., Chamberlain, C.J., Ettinger, A.K.,  
 179 Garner, M., Jones, F.A.M., Loughnan, D., Pearse, W.D., Sodhi, D. & Wolkovich, E.M. (2024)  
 180 Phylogenetic estimates of species-level phenology improve ecological forecasting. *Nature Climate*  
 181 *Change* **14**, 989–995.

182 Paine, C.E.T., Amissah, L., Auge, H., Baraloto, C., Baruffol, M., Bourland, N., Bruelheide, H.,  
 183 Dainou, K., de Gouvenain, R.C., Doucet, J.L., Doust, S., Fine, P.V.A., Fortunel, C., Haase, J., Holl,  
 184 K.D., Jactel, H., Li, X., Kitajima, K., Koricheva, J., Martínez-Garza, C., Messier, C., Paquette, A.,  
 185 Philipson, C., Piotto, D., Poorter, L., Posada, J.M., Potvin, C., Rainio, K., Russo, S.E., Ruiz-jaen,  
 186 M., Scherer-lorenzen, M., Webb, C.O., Wright, S.J., Zahawi, R.A. & Hector, A. (2015) Globally ,  
 187 functional traits are weak predictors of juvenile tree growth , and we do not know why. *Journal of*  
 188 *Ecology* **103**, 978–989.

189 Pérez-de Lis, G., Olano, J.M., Rozas, V., Rossi, S., Vázquez-Ruiz, R.A. & García-Gonzalez, I. (2017)  
 190 Environmental conditions and vascular cambium regulate carbon allocation to xylem growth in  
 191 deciduous oaks. *Functional Ecology* **31**, 592–603.

192 Prentice, I.C., Meng, T., Wang, H., Harrison, S.P., Ni, J. & Wang, G. (2011) Evidence of a universal  
 193 scaling relationship for leaf CO<sub>2</sub> drawdown along an aridity gradient. *New Phytologist* **190**, 169–180.

194 Price, C.A., Wright, I.J., Ackerly, D.D., Niinemets, Ü., Reich, P.B. & Veneklaas, E.J. (2014) Are leaf  
 195 functional traits ‘invariant’ with plant size and what is ‘invariance’ anyway? *Functional Ecology* **28**,  
 196 1330–1343.

197 Robinson, K.M., Hauzy, C., Loeuille, N. & Albrechtsen, B.R. (2015) Relative impacts of environmental  
 198 variation and evolutionary history on the nestedness and modularity of tree–herbivore networks.  
 199 *Ecology and Evolution* **5**, 2898–2915.

200 Schweingruber, F. & Landolt, W. (2010) The xylem database.

201 Shipley, B. & Vu, T.T. (2002) Dry matter content as a measure of dry matter concentration in plants  
 202 and their parts. *New Phytologist* **153**, 359–364.

203 Vergutz, L., Manzoni, S., Porporato, A., Novais, R. & Jackson, R. (2012) A Global Database of  
 204 Carbon and Nutrient Concentrations of Green and Senesced Leaves. *Oak Ridge National Laboratory*  
 205 *Distributed Active Archive Center* **Oak Ridge, Tennessee, U.S.A.**

206 Wang, H., Harrison, S.P., Prentice, I.C., Yang, Y., Bai, F., Furstenau Togashi, H., Wang, M., Zhou,  
 207 S. & Ni, J. (2017) The China Plant Trait Database. *PANGAEA* .

- 208 Wenxuan, H., Yahan, C., Fang-Jie, Z., Tang, L., Rongfeng, J. & Fusuo, Z. (2012) Floral, climatic and  
 209 soil pH controls on leaf ash content in China's terrestrial plants. *Global Ecology and Biogeography*  
 210 **21**, 376–382.
- 211 Wilson, K.B., Baldocchi, D.D. & Hanson, P.J. (2000) Spatial and seasonal variability of photosynthetic  
 212 parameters and their relationship to leaf nitrogen in a deciduous forest. *Tree Physiology* **20**, 565–578.
- 213 Wright, I.J., Westoby, M., Reich, P.B., Oleksyn, J., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-  
 214 Bares, J., Chapin, T., Cornellissen, J.H.C., Diemer, M., Flexas, J., Gulias, J., Garnier, E., Navas,  
 215 M.L., Roumet, C., Groom, P.K., Lamont, B.B., Hikosaka, K., Lee, T., Lee, W., Lusk, C., Midgley,  
 216 J.J., Niinemets, Ü., Osada, H., Poorter, H., Pool, P., Veneklaas, E.J., Prior, L., Pyankov, V.I.,  
 217 Thomas, S.C., Tjoelker, M.G. & Villar, R. (2004) The worldwide leaf economics spectrum. *Nature*  
 218 **428**, 821–827.
- 219 Yahan, C., Wenxuan, H., Luying, T., Zhiyao, T. & Jingyun, F. (2013) Leaf nitrogen and phosphorus  
 220 concentrations of woody plants differ in responses to climate, soil and plant growth form. *Ecography*  
 221 **36**, 178–184.