

<sup>1</sup> Woody plant phenological responses are strongly associated  
<sup>2</sup> with key functional traits

<sup>3</sup> Deirdre Loughnan<sup>1</sup>, Faith A M Jones<sup>1,2</sup>, Geoffrey Legault<sup>1</sup>, Daniel Buonaiuto<sup>3,4,5</sup>,  
Catherine Chamberlain<sup>3,4,6</sup>, Ailene Ettinger<sup>7</sup>, Mira Garner<sup>1</sup>, Ignacio Morales-Castilla<sup>8,9</sup>,  
Darwin Sodhi<sup>1</sup> and E M Wolkovich<sup>1,3,4</sup>

<sup>4</sup> August 15, 2024

<sup>5</sup> <sup>1</sup> Department of Forest and Conservation, Faculty of Forestry, University of British Columbia, 2424  
<sup>6</sup> Main Mall Vancouver, BC Canada V6T 1Z4.

<sup>7</sup> <sup>2</sup> Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sci-  
<sup>8</sup> ences, 901 83 Umeå, Sweden

<sup>9</sup> <sup>3</sup> Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

<sup>10</sup> <sup>4</sup> Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts,  
<sup>11</sup> USA;

<sup>12</sup> <sup>5</sup> Department of Environmental Conservation, University of Massachusetts, Amherst, 160 Holdsworth  
<sup>13</sup> Way, Amherst, MA, USA

<sup>14</sup> <sup>6</sup> The Nature Conservancy, 334 Blackwell St Ste 300, Durham, NC, USA

<sup>15</sup> <sup>7</sup> The Nature Conservancy of Washington, 74 Wall Street, Seattle, WA USA

<sup>16</sup> <sup>8</sup> GloCEE—Global Change Ecology and Evolution Group, Department of Life Sciences, University of  
<sup>17</sup> Alcalá, Alcalá de Henares, Spain

<sup>18</sup> <sup>9</sup> Edificio Ciencias, Campus Universitario 28805 Alcalá de Henares, Madrid, Spain

<sup>20</sup> Corresponding Author: Deirdre Loughnan deirdre.loughnan@ubc.ca

<sup>21</sup> Running title: Budburst responses associated with traits

<sup>31</sup> **Summary**

<sup>32</sup> Species-level variation in phenology—the timing of recurring life history events—can vary seasonally  
<sup>33</sup> with changes in climatic risk, light, and nutrients. This favours acquisitive growth strategies early in  
<sup>34</sup> the spring season and conservative growth strategies under the more climatically benign, but compet-  
<sup>35</sup> itive, environment later in the season. This framework infers suites of traits that may co-vary with  
<sup>36</sup> phenologies, but high trait variability across environments makes this challenging to test. Here, we  
<sup>37</sup> combine a new joint modeling approach to accommodate this variability with global data on plant  
<sup>38</sup> traits and budburst responses in controlled environment experiments. We find that earlier species—  
<sup>39</sup> which are generally most responsive to anthropogenic warming—are generally shorter with denser,  
<sup>40</sup> lower nitrogen leaves. These results suggest warming may reshape the trait structure of plant com-  
<sup>41</sup> munities, and could help improve predictions of how growth strategies and phenologies together may  
<sup>42</sup> shift with continued climate change.

<sup>43</sup>

<sup>44</sup> Key Words: Budburst, spring phenology, functional traits, trees, climate change, forest communities

<sup>45</sup> **Introduction**

<sup>46</sup> The timing of life history events—phenology—can shape both ecosystem services and community  
<sup>47</sup> dynamics. Spring phenology, for example, defines the start and overall length of the growing season—  
<sup>48</sup> shaping forest carbon storage and species interactions<sup>3,9,18</sup>. Shifts in phenology with climate change  
<sup>49</sup> across systems<sup>31,45</sup> have thus led to growing concerns over their possible impacts.

<sup>50</sup>

<sup>51</sup> Predicting these changes requires understanding the drivers of phenology both at a proximate scale—  
<sup>52</sup> the environmental triggers of phenology each year, such as temperature and daylength—and at an  
<sup>53</sup> ultimate scale, where long-term environmental pressures may select for different phenologies across  
<sup>54</sup> species e.g., certain species are early or late relative to other species each year,<sup>33,51</sup>. At the proximate  
<sup>55</sup> level, environmental conditions throughout the winter and spring cause species to start growth at dif-  
<sup>56</sup> ferent times. This is well documented for the start of growth in woody plants each year<sup>15,24</sup>. Similar  
<sup>57</sup> environmental conditions appear to trigger spring phenological events across taxa, including in the  
<sup>58</sup> timing of egg laying in birds<sup>11,12</sup> and the advance of spawning in amphibians<sup>23,43</sup>, but current work  
<sup>59</sup> provides limited insights into the drivers of species differences<sup>6,15,24</sup>.

<sup>60</sup>

<sup>61</sup> At the ultimate level, species phenologies may vary due to changing pressures across the growing season.  
<sup>62</sup> Species that start growth early often risk high tissue loss—due to frost damage<sup>2,37</sup> or high herbivore  
<sup>63</sup> apparency<sup>46</sup>—but benefit from higher resource availability<sup>21,34</sup>. In contrast, later species face greater  
<sup>64</sup> biotic pressures, especially from high competition for resources<sup>27,52</sup>. For plants, this variation in early  
<sup>65</sup> to late season growth, may mirror the stressors from early to late successional communities, and may  
<sup>66</sup> similarly shape phenology<sup>24</sup>.

<sup>67</sup>

<sup>68</sup> Different pressures could shape a number of species attributes related to their growth strategies, includ-  
<sup>69</sup> ing phenology. Species with earlier phenology may produce cheaper tissues that are easily replaced if  
<sup>70</sup> damaged<sup>36</sup>, while species with later phenology may benefit from investing in tissues that infer greater  
<sup>71</sup> resource retention<sup>16</sup>. Differences in traits, and trade-offs in allocation of resources to growth and tissue  
<sup>72</sup> quality, can be related to a broader framework of species growth strategies and functional traits<sup>52</sup> (Fig  
<sup>73</sup> 1), where species range from acquisitive (fast) to more conservative (slow) growth<sup>5,53</sup>.

<sup>74</sup>

<sup>75</sup> Globally, gradients from acquisitive to conservative strategies appear common, and form the foun-  
<sup>76</sup> dation of the leaf economic and the wood economic spectra<sup>5,13,47,48,53</sup>, but they can make limited  
<sup>77</sup> predictions of trait variability<sup>44</sup>. As a result, highly variable traits like phenology are often excluded  
<sup>78</sup> from trait studies, leaving the relationships between broader trait syndromes and phenology largely

79 unknown. Understanding these relationships is critical to forecasting community dynamics and re-  
80 sponses to climate change. To date, studies have generally only examined the relationship between  
81 traits and phenology within a single site, as reviewed by Wolkovich and Cleland<sup>50</sup> and Wolkovich and  
82 Donahue<sup>51</sup>, where the problem of proximate drivers causing phenological variation can be more easily  
83 ignored. Adding phenology to broader trait frameworks becomes more complex when high levels of  
84 variation occurs across large spatial and temporal ranges.

85

86 Consistently defining early to late phenology is possible using the underlying cues that predict gradi-  
87 ents in phenology which do not generally vary strongly across space and time,<sup>7,15,20</sup>. For many plants,  
88 early species generally have small in magnitude responses to all three major cues of spring leafout:  
89 warm spring temperatures (forcing), cool winter temperatures (chilling) and daylength (photoperiod).  
90 In contrast, later species have larger responses to chilling and/or photoperiod<sup>15,24</sup>, and likely larger  
91 forcing responses.

92

93 Studies of spring phenology in temperate forests may provide the best opportunity to integrate phe-  
94 nology into functional trait research. In addition to how well we understand the environmental cues  
95 that trigger early versus late leafout, spring in many forests includes strong gradients in potential se-  
96 lective environments (Fig 1). Based on trade-offs between early and late spring phenologies, we predict  
97 acquisitive species to be shorter, with leaf traits favourable to higher light availability and tolerance  
98 of late spring frost (high specific leaf area, SLA, and leaf nitrogen content, LNC; Fig 1). Such species  
99 should exhibit early phenology, with small cue responses. Canopy species that budburst later via larger  
100 cue responses, when competition for soil resources is greater, would then have traits associated with  
101 conservative growth—taller with denser wood<sup>25</sup>—with leaf traits suited for more variable light (low  
102 SLA and LNC, Fig 1). Seed size may similarly be predicted from this acquisitive to conservative con-  
103 tinuum, as acquisitive species produce smaller seeds and conservative species produce larger—better  
104 provisioned—seeds (Fig 1).

105

106 To test our predicted relationships between budburst responses to environmental cues and common  
107 functional traits (height, SLA, seed mass, and LNC), we merged available data from trait databases—  
108 BIEN<sup>29</sup> and TRY<sup>22</sup>—with budburst data from the OSPREE database of controlled environment stud-  
109 ies<sup>14</sup>. We developed a hierarchical Bayesian joint model that predicts phenological responses to forcing,  
110 chilling and photoperiod treatments based on species-level trait values, while allowing additional varia-  
111 tion due to species. This approach takes a step towards predicting variation via species traits instead of  
112 species identity (when traits explain a significant portion of the variation, species identity will explain  
113 only a small amount), which could help forecast species phenological responses based on trait values  
114 alone.

115

## 116 Methods

117 We merged three major databases for our analysis. We gathered phenological data from the OSPREE  
118 database<sup>14</sup>, which contains budburst data for woody, deciduous species from experiments of forcing,  
119 chilling and photoperiod. We updated this database since its initial publication, the methods of which  
120 are discussed by Morales-Castilla et al.<sup>32</sup>. We gathered trait data from TRY and BIEN (v. 4.0)<sup>22,29</sup>,  
121 both of which are large trait databases that include plant trait data across many individuals, species,  
122 and studies (Table S1). We obtained data from both databases on 5 December, 2018, with an updated  
123 version of the TRY data obtained 10 April, 2019. We focused our search for trait data on the subset of  
124 234 OSPREE species used in Morales-Castilla et al.<sup>32</sup>. Using the BIEN R package, version 1.2.5<sup>29</sup>, we  
125 downloaded trait data for the 94 species available, for which there were 13 traits. The TRY database  
126 included data for 10 traits for 96 of our focal species<sup>22</sup>. Given our focus on phenology of adult trees, we  
127 included trait data from adult individuals with a minimum height of 1.38 m. We further removed all

128 data from experiments or from plants growing in non-natural habitats. We also grouped trait values  
 129 where appropriate, for example categorizing trait values for “SLA”, “SLA with petioles”, and “SLA  
 130 without petioles” as simply “SLA” in our analysis (see Table S1). Duplicated data in both the TRY  
 131 and BIEN datasets were also removed ( $n = 434905$ ). Based on our selection criteria, our final dataset  
 132 included data for 11 traits from 91 of the species also represented in the OSPREE database, with each  
 133 species differing in the number and types of traits measured.  
 134

135 For our analysis, we only include species for which we had a complete trait profile (i.e., all traits mea-  
 136 sured for all species). We initially considered six commonly measured traits—SLA, leaf dry matter  
 137 content (LDMC), height, seed mass, stem specific density (SSD), and LNC—for which 26 species had at  
 138 least one trait measurement for each trait. We then used a principle component analysis to understand  
 139 trait correlations and adjusted which traits we included. A PCA of our six initial traits identified high  
 140 correlations between SLA and LDMC, and between height and SSD (see Supplementary material).  
 141 The first principal component explained 32% of variation while the second explained 24.2% of the  
 142 variation (Fig. S1). By excluding one trait from each of these highly correlated trait pairs (specifically  
 143 LDMC and SSD) we increased the number of species in our dataset from the 26 species with six traits,  
 144 to 37 species for which we had complete datasets for four traits. The data for these 37 species were  
 145 from 24 unique studies (height  $n = 47781$ , seed mass  $n = 281$ , LNC  $n = 3853$ , SLA  $n = 7656$ ). We  
 146 subsampled height measurements to reduce the influence the 13 most frequently measured tree species  
 147 had on our height model. Since these 13 species were measured 19 times more frequently than other  
 148 species, for each species, we randomly sampled 3000 height measurements.  
 149

## 150 Joint model of trait and phenology

151 To understand connections between phenology and species traits, we developed and then parameterized  
 152 a joint model for each trait: height, SLA, LNC, and seed mass. Our model is a joint model insofar  
 153 as it involves two types of data—trait observations and phenological observations—that arise from  
 154 shared latent processes. In particular, we assume that species “true” trait values determine observed  
 155 trait values across different studies (trait sub-model), and separately, that the same “true” trait values  
 156 interact with phenological cues (forcing, chilling, and photoperiod) to determine observed phenology,  
 157 specifically day of year of budburst (phenology sub-model). Below we describe the two sub-models,  
 158 noting which parameters are shared across sub-models and which are independent.

### 159 Trait sub-model

160 The trait sub-model describes the processes that determine trait observations for 1 to  $n$  species across  
 161 each of the 1 to  $m$  studies in our trait dataset (TRY and BIEN data). We use hierarchical modeling to  
 162 partitions trait variation by measurement error, species identity, and study identity. In particular, we  
 163 assume that a trait observation for species  $i$  from study  $j$ ,  $Y_{\text{trait}_{i,j}}$ , has the following normal distribution:

$$Y_{\text{trait}_{i,j}} \sim \mathcal{N}(\mu_{i,j}, \sigma_m^2) \quad (1)$$

with

$$\mu_{i,j} = \alpha_{\text{grand trait}} + \alpha_{\text{trait}_i} + \alpha_{\text{study}_j} \quad (2)$$

164 where  $\alpha_{\text{trait}_i}$  and  $\alpha_{\text{study}_j}$  are elements of the normal random vectors:

$$\begin{aligned} \boldsymbol{\alpha}_{\text{trait}} &= \{\alpha_{\text{trait}_1}, \dots, \alpha_{\text{trait}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{trait}} \sim \mathcal{N}(0, \sigma_{\text{trait}}^2) \\ \boldsymbol{\alpha}_{\text{study}} &= \{\alpha_{\text{study}_1}, \dots, \alpha_{\text{study}_m}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{study}} \sim \mathcal{N}(0, \sigma_{\text{study}}^2) \end{aligned} \quad (3)$$

165 The latent parameter  $\alpha_{\text{grand trait}}$  represents a trait value that is independent of study and species,  
 166  $\alpha_{\text{species},i}$  and  $\alpha_{\text{study},j}$  are species- and study-level offsets from that trait value,  $\sigma_m^2$  is measurement error,  
 167 and  $\sigma_{\text{trait}}^2$  and  $\sigma_{\text{study}}^2$  represent species and study-level variances in trait values. Of these parameters,  
 168  $\alpha_{\text{trait}}$  are shared by the phenology sub-model.

## 169 Phenology sub-model

170 The phenology sub-model describes the processes that determine phenological observations for 1 to  
 171  $n$  species, specifically the timing (day of year) of budburst from the updated OSPREE dataset. We  
 172 assume that an observation of budburst day for species  $k$  under set  $g$  of chilling, forcing, and photoper-  
 173 iod treatments, which we  $z$ -scored to allow direct comparison of cues  $(c_g, f_g, p_g)$ ,  $Y_{\text{pheno}_{k,g}}$ , has the  
 174 following normal distribution:

$$Y_{\text{pheno}_{k,g}} \sim \mathcal{N}(\mu_{k,g}, \sigma_d^2) \quad (4)$$

with

$$\mu_{k,g} = \alpha_{\text{pheno},k} + \beta_{\text{chill}_k} \cdot c_g + \beta_{\text{force}_k} \cdot f_g + \beta_{\text{photo}_k} \cdot p_g \quad (5)$$

and

$$\begin{aligned} \beta_{\text{chill},k} &= \alpha_{\text{chill},k} + \beta_{\text{trait.chill}} \cdot \alpha_{\text{trait},k} \\ \beta_{\text{force},k} &= \alpha_{\text{force},k} + \beta_{\text{trait.force}} \cdot \alpha_{\text{trait},k} \\ \beta_{\text{photo},k} &= \alpha_{\text{photo},k} + \beta_{\text{trait.photo}} \cdot \alpha_{\text{trait},k} \end{aligned} \quad (6)$$

175 where  $\alpha_{\text{pheno}_k}$ ,  $\alpha_{\text{chill}_k}$ ,  $\alpha_{\text{force}_k}$ , and  $\alpha_{\text{photo}_k}$  are elements of the normal random vectors:

$$\begin{aligned} \boldsymbol{\alpha}_{\text{pheno}} &= \{\alpha_{\text{pheno}_1}, \dots, \alpha_{\text{pheno}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{pheno}} \sim \mathcal{N}(\mu_{\text{pheno}}, \sigma_{\text{pheno}}^2) \\ \boldsymbol{\alpha}_{\text{chill}} &= \{\alpha_{\text{chill}_1}, \dots, \alpha_{\text{chill}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{chill}} \sim \mathcal{N}(\mu_{\text{chill}}, \sigma_{\text{chill}}^2) \\ \boldsymbol{\alpha}_{\text{force}} &= \{\alpha_{\text{force}_1}, \dots, \alpha_{\text{force}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{force}} \sim \mathcal{N}(\mu_{\text{force}}, \sigma_{\text{force}}^2) \\ \boldsymbol{\alpha}_{\text{photo}} &= \{\alpha_{\text{photo}_1}, \dots, \alpha_{\text{photo}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{photo}} \sim \mathcal{N}(\mu_{\text{photo}}, \sigma_{\text{photo}}^2) \end{aligned} \quad (7)$$

176 Parameter  $\alpha_{\text{pheno},k}$  represents the day of budburst for species  $k$  without the influence of chilling, forc-  
 177 ing, or photoperiod treatments. Average day of budburst (independent of treatments) is  $\mu_{\text{pheno}}$ , and  
 178  $\sigma_{\text{pheno}}^2$  is the variance across species. The parameters  $\alpha_{\text{chill},k}$ ,  $\alpha_{\text{force},k}$ , and  $\alpha_{\text{photo},k}$  represent the trait-  
 179 independent responses of species  $k$  to chilling, forcing, and photoperiod treatments respectively, each  
 180 with an associated mean ( $\mu_{\text{chill}}$ ,  $\mu_{\text{force}}$ ,  $\mu_{\text{photo}}$ ) and variance ( $\sigma_{\text{chill}}^2$ ,  $\sigma_{\text{force}}^2$ ,  $\sigma_{\text{photo}}^2$ ) across species. The  
 181 effect of the species trait value,  $\alpha_{\text{trait},k}$  (parameter shared with trait sub-model above), on its responses  
 182 to chilling, forcing, and photoperiod are described by parameters  $\beta_{\text{trait.chill}}$ ,  $\beta_{\text{trait.force}}$ ,  $\beta_{\text{trait.photo}}$ . Fi-  
 183 nally,  $\sigma_d^2$  is the variance arising from measurement error.

184  
 185 We chose weakly informative priors, and validated them using a series of prior predictive checks.  
 186 The model was coded in the Stan programming language, fit using the rstan package version 3.3.6<sup>41</sup>,  
 187 with 1,000 iterations per chain across 4 chains (4,000 total sampling iterations), and all models met  
 188 basic diagnostic checks, including no divergences, high effective sample size ( $n_{\text{eff}}$ ), and  $\hat{R}$  close to 1,  
 189 fitting the data well (Fig S2). Here we present our model estimates as the means and 90% posterior  
 190 uncertainty intervals.

191 **Results**

192 Across traits, height, SLA, and LNC strongly related to chilling, forcing and photoperiod treatments  
193 ( $\beta_{\text{chill}_k}$ ,  $\beta_{\text{force}_k}$ , and  $\beta_{\text{photo}_k}$ , Fig 2 a-f & j-l), but the direction of these relationships only showed  
194 consistent trends for LNC (Fig. 2 j-l). As we predicted, height was related to chilling ( $\beta_{\text{chill}_k}$ ) and pho-  
195 toperiod ( $\beta_{\text{photo}_k}$ ), with taller species having larger responses to cues (-0.5 m per standardized chilling;  
196 90% uncertainty interval (UI): -1.0, -0.1 and -0.2 m per standardized photoperiod; 90% UI: -0.5, 0.0,  
197 Fig 2 a-c, Table S2). As illustrated for one characteristically acquisitive species, *Alnus incana*, and  
198 one characteristically conservative species, *Quercus rubra* (Fig S3), the cue relationships with height  
199 led to generally later budburst ( $\mu_{k,g}$ ; Fig. 3). In contrast, seed mass had the smallest responses, with  
200 no relationship between seed mass and any cue (Fig. 2 g-i, Fig 3 d-f, & Table S3).

201  
202 Of our leaf traits, we found that species SLA related to photoperiod ( $\beta_{\text{photo}_k}$ , -0.2 mm<sup>2</sup>/mg per stan-  
203 dardized photoperiod; 90% UI: -0.4, 0.0, Fig. 2 f, Table S4), but did not strongly predict responses  
204 to chilling ( $\beta_{\text{chill}_k}$ ) or forcing treatments ( $\beta_{\text{force}_k}$ , Fig. 2 d and e). Thus, species with more acquisitive  
205 growth strategies (thin leaves and a lower investment in leaf mass that leads to large SLA values),  
206 had larger responses to photoperiod, contrary to our predictions (Fig. 2 f). For LNC, we found that  
207 species that produce leaves with high nitrogen content, which relates generally to high photosynthetic  
208 rates and acquisitive growth, show smaller responses to cues (Fig. 2 j-l). These findings are in line  
209 with our predictions that high LNC species (acquisitive) would be less responsive to chilling (0.7 mg/g  
210 per standardized chilling; 90% UI: 0.3, 1.2, Table S5), but we also found high LNC species to be less  
211 responsive to photoperiod (0.3 mg/g per standardized photoperiod; 90% UI: 0, 0.6) and to forcing (0.5  
212 mg/g per standardized forcing; 90% UI: 0.1, 0.9, Fig 2 j-l & Fig S3 d-f).

213  
214 Across our models, we found species-level variation across traits ( $\sigma_{\text{trait}}^2$ ) were comparable to or greater  
215 than variation across studies ( $\sigma_{\text{study}}^2$ , Fig 4). The magnitude of study-level variation ( $\sigma_{\text{study}}^2$ ) that we  
216 found, however, suggests that models using large trait databases that fail to separate out study from  
217 species-level variation ( $\sigma_{\text{trait}}^2$ ) may poorly estimate species traits. Variation across studies was greatest  
218 for height (with  $\sigma_{\text{study}}^2$  of 7.5 m compared to 5.9 m for  $\sigma_{\text{trait}}^2$ , Fig 4a). For seed mass and LNC study-  
219 level variation was less than that of the species-level variation, with estimates of 1 mg for study-level  
220 variation versus 1.6 for species-level variation in seed mass and estimates of 3.6 mg g<sup>-1</sup> for study-level  
221 variation and 5.1 mg g<sup>-1</sup> for the species-level variation in LNC (Fig 4c and d). At the lowest end,  
222 study-level variation in SLA was approximately half the value of the species-level variation (3.3 mm<sup>2</sup>  
223 mg<sup>-1</sup> versus 7.8 mm<sup>2</sup> mg<sup>-1</sup> for  $\sigma_{\text{study}}^2$  and  $\sigma_{\text{trait}}^2$ , respectively, Fig 4b).

224

225 **Discussion**

226 We found species traits influenced the timing of budburst in response to the three primary cues of  
227 spring phenology: chilling, forcing and photoperiod. These trait effects were associated with earlier  
228 or later phenology following well-established gradients in growth strategies predicted by functional  
229 trait frameworks<sup>5,13,47,48,53</sup>: early species tended to have traits associated with fast and acquisitive  
230 strategies while later species had traits associated with conservative, slower strategies. We found the  
231 largest budburst responses occurred for traits related to resource acquisition and structure, with SLA,  
232 LNC, and height all showing large responses across our three cues. In contrast, our one reproductive  
233 trait—seed mass—showed a smaller response. Our results provide a major step forward in integrating  
234 phenology into broader trait syndromes that shape species growth strategies, and support previous  
235 findings from more local scales that found strong relationships between height and species phenol-  
236 ogy<sup>39,40,42</sup>. Our findings also suggest other traits—such as seed mass—show no relationship with  
237 phenology in our more global analysis.

238

**239 Effects of phenology-trait relationships on community assembly**

240 Our findings suggest the changing pressures across the early growing season may affect the temporal  
241 assembly of communities. Strong abiotic pressures alongside weak competition early in the season were  
242 associated with early-budbursting species with acquisitive traits (shorter heights and low LNC) that  
243 allow faster return on resource investments<sup>5,17,47</sup>. These traits should allow early species to more easily  
244 replace tissue if lost to frost or other abiotic disturbances, and benefit from greater light availability in  
245 the open canopy of many temperate forests in the early spring. In contrast, later-budbursting species  
246 had traits associated with greater competitive abilities and slower growth Fig. 2,<sup>5,17,47</sup>, which may  
247 help them compete for soil and light resources when most other species are already growing. These  
248 traits can be linked to other ecological processes and species characteristics, such as species succes-  
249 sional position, as illustrated by the differences between early and late successional species (e.g., *Alnus*  
250 *incana* and *Quercus rubra*; Fig 2).

251 The traits with cue responses that deviated from our expectations also offer novel insights into the  
252 tradeoffs between traits and environmental cues. All of our traits are associated with numerous aspects  
253 of species growth, and may be adaptive for reasons other than those we predicted. Contrary to our  
254 predictions, we found large responses to forcing for short trees, which could prevent frost damage or  
255 xylem cavitation under a late spring frost<sup>10,30</sup> and influence annual cambial meristem growth<sup>26</sup>. Simi-  
256 larly, the lack of a response to chilling or forcing by high SLA individuals could be driven by other trait  
257 attributes and environmental cues—selecting for species relative growth rates or leaf longevity—and  
258 not photosynthetic potential<sup>35,47</sup>. These findings highlight the complexity of determining the drivers  
259 of species trait profiles, and further our understanding of how traits affect community dynamics under  
260 variable environments.

262

**263 Phenology-trait relationships under future climates**

264 Incorporating phenology within broader trait syndromes could aid forecasting of species and community  
265 responses to climate change. While decades of research have documented phenological shifts with  
266 anthropogenic climate change, increasing research suggests a potential connection between phenological  
267 responses to warming and performance with warming, where species that shift their phenology more  
268 also perform better<sup>8,28</sup>.

269 Our results suggest this phenology-performance relationship could be driven in part by a suite of traits  
270 that covary with phenological cues to determine how responsive species are to warming. Species with  
271 smaller responses to all cues, especially chilling and photoperiod, would tend to advance more with  
272 warming, our results suggest these species may also grow more quickly. These results could further aid  
273 in predicting the potential for invasion, as communities with similar phenologies and suites of traits,  
274 appear more susceptible to fast growing, phenologically more responsive invasive species<sup>1,38,49</sup>.

275

276 Our analytical approach and results may be especially useful to help forecast changes in forest dynamics.  
277 Identifying the trait syndromes of forest communities over spring season can aid predictions about how  
278 climate change will alter species growth and productivity. For example, our results suggest that, by  
279 favoring more phenologically responsive species (i.e., with small chilling and photoperiod responses),  
280 warming may also favor species with acquisitive growth strategies. In contrast, conservative species,  
281 which appear less phenologically responsive to changes in temperature (due to larger chilling and  
282 photoperiod responses) could therefore face greater abiotic and biotic stress<sup>19</sup>.

283 Our results could further help identify which species are most likely to be negatively impacted under  
284 future climates, and develop better strategies for climate change mitigation and conservation. Species  
285 that fail to advance phenologically with warming might experience more competition<sup>1,4</sup>, as species that  
286 begin growth increasingly earlier with warming have more time to deplete resources. In addition to  
287 altering the timing and interactions between species within a season, species trait syndromes have the

<sup>288</sup> potential to redefine the environmental conditions under which growth occurs, and as a result, shape  
<sup>289</sup> community assembly, and productivity of diverse ecological communities. By identifying the species  
<sup>290</sup> most vulnerable to climate change impacts, we can develop more effective management practices that  
<sup>291</sup> prevent the loss of ecosystem services and community diversity.

<sup>292</sup>

293    **References**

- 294 [1] Alexander, J. M., and J. M. Levine. 2019. Earlier phenology of a nonnative plant increases impacts  
295        on native competitors. PNAS 116:6199–6204.
- 296 [2] Augspurger, C. K. 2009. Spring 2007 warmth and frost : phenology , damage and refoliation in a  
297        temperate deciduous forest. Functional Ecology 23:1031–1039.
- 298 [3] Beard, K. H., K. C. Kelsey, A. J. Leffler, and J. M. Welker. 2019. The Missing Angle : Ecosystem  
299        Consequences of Phenological Mismatch. Trends in Ecology and Evolution 34:885–888.
- 300 [4] Carter, S. K., D. Saenz, and V. H. Rudolf. 2018. Shifts in phenological distributions reshape  
301        interaction potential in natural communities. Ecology Letters 21:1143–1151.
- 302 [5] Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a  
303        worldwide wood economics spectrum. Ecology Letters 12:351–366.
- 304 [6] Chuine, I., M. Bonhomme, J. M. Legave, I. García de Cortázar-Atauri, G. Charrier, A. Lacointe,  
305        and T. Améglio. 2016. Can phenological models predict tree phenology accurately in the future?  
306        The unrevealed hurdle of endodormancy break. Global change biology 22:3444–3460.
- 307 [7] Chuine, I., and P. Cour. 1999. Climatic determinants of budburst seasonality in four temperate-  
308        zone tree species. New Phytologist 143:339–349.
- 309 [8] Cleland, E. E., J. M. Allen, T. M. Crimmins, J. A. Dunne, S. Pau, S. E. Travers, E. S. Zavaleta,  
310        and E. M. Wolkovich. 2012. Phenological tracking enables positive species responses to climate  
311        change.pdf. Ecology 93:1765–1771.
- 312 [9] Cleland, E. E., I. Chuine, A. Menzel, H. A. Mooney, and M. D. Schwartz. 2007. Shifting plant  
313        phenology in response to global change. Trends in Ecology and Evolution 22:357–365.
- 314 [10] Clements, J. R., W. Fraser, J, and C. W. Yeatman. 1972. Frost Damage to White Spruce Buds.  
315        Canadian Journal of Forest Research 2:62–63.
- 316 [11] Crick, H., C. Dudley, D. E. Glue, and D. L. Thomson. 1997. UK birds are laying eggs earlier.  
317        Nature 388:526–527.
- 318 [12] D'Alba, L., P. Monaghan, and R. G. Nager. 2010. Advances in laying date and increasing popu-  
319        lation size suggest positive responses to climate change in Common Eiders Somateria mollissima  
320        in Iceland. International Journal of Avian Science 152:19–28.
- 321 [13] Díaz, S., J. Kattge, J. H. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer,  
322        C. Wirth, I. Colin Prentice, E. Garnier, G. Bönisch, M. Westoby, H. Poorter, P. B. Reich, A. T.  
323        Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright, S. N. Sheremet Ev,  
324        H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig,  
325        A. Günther, V. Falcuk, N. Rüger, M. D. Mahecha, and L. D. Gorné. 2016. The global spectrum  
326        of plant form and function. Nature 529:167–171.
- 327 [14] Ettinger, A. K., C. J. Chamberlain, I. Morales-Castilla, D. M. Buonaiuto, D. F. Flynn, T. Savas,  
328        J. A. Samaha, and E. M. Wolkovich. 2020. Winter temperatures predominate in spring phenolog-  
329        ical responses to warming. Nature Climate Change 10:1137–1142.
- 330 [15] Flynn, D. F. B., and E. M. Wolkovich. 2018. Temperature and photoperiod drive spring phenology  
331        across all species in a temperate forest community. New Phytologist 219:1353–1362.
- 332 [16] Gorné, L. D., S. Díaz, V. Minden, Y. Onoda, K. Kramer, C. Muir, S. T. Michaletz, S. Lavorel,  
333        J. Sharpe, S. Jansen, M. Slot, E. Chacon, and G. Boenisch. 2020. The acquisitive–conservative  
334        axis of leaf trait variation emerges even in homogeneous environments. Annals of Botany .

- 335 [17] Grime, J. P. 1977. Evidence for the Existence of Three Primary Strategies in Plants and Its  
336 Relevance to Ecological and Evolutionary Theory Author ( s ): J . P . Grime Source : The  
337 American Naturalist , Vol . 111 , No . 982 ( Nov . - Dec ., 1977 ), pp . 1169-1194 Published. The  
338 American Naturalist 111:1169–1194.
- 339 [18] Gu, H., Y. Qiao, Z. Xi, S. Rossi, N. G. Smith, J. Liu, and L. Chen. 2022. Warming-induced  
340 increase in carbon uptake is linked to earlier spring phenology in temperate and boreal forests.  
341 Nature Communications 13:1–8.
- 342 [19] Guy, R. D. 2014. The early bud gets to warm. New Phytologist 202:7–9.
- 343 [20] Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying  
344 dormancy requirements for Pacific Northwest tree species. Frontiers in Plant Science 6:1–12.
- 345 [21] Hufkens, K., M. A. Friedl, T. F. Keenan, O. Sonnentag, A. Bailey, J. O'keefe, and A. D. Richardson.  
346 2012. Ecological impacts of a widespread frost event following early spring leaf-out. Global  
347 Change Biology 18:2365–2377.
- 348 [22] Kattge, J., G. Bönisch, S. Díaz, S. Lavorel, I. C. Prentice, and et al. 2020. TRY plant trait  
349 database – enhanced coverage and open access. Global Change Biology 26:119–188.
- 350 [23] Kusano, T., and M. Inoue. 2008. Long-Term Trends toward Earlier Breeding of Japanese Am-  
351 phibians. Journal of Herpetology 42:608–614.
- 352 [24] Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling  
353 outweighs photoperiod in preventing precocious spring development. Global Change Biology  
354 20:170–182.
- 355 [25] Laughlin, D. C., J. J. Leppert, M. M. Moore, and C. H. Sieg. 2010. A multi-trait test of the leaf-  
356 height-seed plant strategy scheme with 133 species from a pine forest flora. Functional Ecology  
357 24:493–501.
- 358 [26] Lenz, A., G. Hoch, C. Körner, and Y. Vitasse. 2016. Convergence of leaf-out towards minimum  
359 risk of freezing damage in temperate trees. Functional Ecology 30:1480–1490.
- 360 [27] Lopez, O. R., K. Farris-Lopez, R. A. Montgomery, and T. J. Givnish. 2008. Leaf phenology in  
361 relation to canopy closure in southern Appalachian trees. American Journal of Botany 95:1395–  
362 1407.
- 363 [28] Macgregor, C. J., C. D. Thomas, D. B. Roy, M. A. Beaumont, J. R. Bell, T. Brereton, J. R. Bridle,  
364 C. Dytham, R. Fox, K. Gotthard, A. A. Hoffmann, G. Martin, I. Middlebrook, S. Nylin, P. J.  
365 Platts, R. Rasteiro, I. J. Saccheri, R. Villoutreix, C. W. Wheat, and J. K. Hill. 2019. Climate-  
366 induced phenology shifts linked to range expansions in species with multiple reproductive cycles  
367 per year. Nature Communications 10:1–10.
- 368 [29] Maitner, B. S., B. Boyle, N. Casler, R. Condit, J. Donoghue, S. M. Durán, D. Guaderrama, C. E.  
369 Hinchliff, P. M. Jørgensen, N. J. Kraft, B. McGill, C. Merow, N. Morueta-Holme, R. K. Peet,  
370 B. Sandel, M. Schildhauer, S. A. Smith, J. C. Svenning, B. Thiers, C. Violette, S. Wiser, and B. J.  
371 Enquist. 2018. The bien r package: A tool to access the Botanical Information and Ecology  
372 Network (BIEN) database. Methods in Ecology and Evolution 9:373–379.
- 373 [30] Marquis, B., Y. Bergeron, M. Simard, and F. Tremblay. 2020. Growing-season frost is a better  
374 predictor of tree growth than mean annual temperature in boreal mixedwood forest plantations.  
375 Global Change Biology 26:6537–6554.

- 376 [31] Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kübler, P. Bissolli,  
377 O. Braslavská, A. Briede, F. M. Chmielewski, Z. Crepinsek, Y. Curnel, Å. Dahl, C. Defila, A. Don-  
378 nnelly, Y. Filella, K. Jatczak, F. Måge, A. Mestre, Ø. Nordli, J. Peñuelas, P. Pirinen, V. Remišová,  
379 H. Scheifinger, M. Striz, A. Susnik, A. J. Van Vliet, F. E. Wielgolaski, S. Zach, and A. Zust. 2006.  
380 European phenological response to climate change matches the warming pattern. *Global Change  
381 Biology* 12:1969–1976.
- 382 [32] Morales-Castilla, I., T. J. Davies, G. Legault, D. M. Buonaiuto, C. J. Chamberlain, A. K. Ettinger,  
383 M. Garner, F. A. M. Jones, D. Loughnan, W. D. Pearse, D. Sodhi, and E. M. Wolkovich. 2024.  
384 Phylogenetic estimates of species-level phenology improve ecological forecasting. *Nature Climate  
385 Change* .
- 386 [33] Ovaskainen, O., S. Skorokhodova, M. Yakovleva, A. Sukhov, A. Kutenkov, N. Kutenkova,  
387 A. Shcherbakov, E. Meyke, and M. del Mar Delgado. 2013. Community-level phenological re-  
388 sponse to climate change. *PNAS* 110:13434–13439.
- 389 [34] Rathcke, B., and E. P. Lacey. 1985. Phenological patterns of terrestrial plants. *Annual Review of  
390 Ecology and Systematics* 16:179–214.
- 391 [35] Reich, P. B. 2014. The world-wide ‘ fast – slow ’ plant economics spectrum : a traits manifesto.  
392 *Journal of Ecology* 102:275–301.
- 393 [36] Reich, P. B., D. S. Ellsworth, M. B. Walters, J. M. Vose, C. Gresham, J. C. Volin, and W. D.  
394 Bowman. 1999. Generality of leaf trait relationships: A test across six biomes. *Ecology* 80:1955–  
395 1969.
- 396 [37] Sakai, A., and W. Larcher. 1987. Frost Survival of Plants: Responses and adaptation to freezing  
397 stress. Springer-Verlag, Berlin, Heidelberg.
- 398 [38] Schuster, M. J., P. D. Wragg, and P. B. Reich. 2021. Phenological niche overlap between invasive  
399 buckthorn (*Rhamnus cathartica*) and native woody species. *Forest Ecology and Management*  
400 498:119568.
- 401 [39] Segrestin, J., M.-l. Navas, and E. Garnier. 2020. Reproductive phenology as a dimension of the  
402 phenotypic space in 139 plant species from the Mediterranean. *New Phytologist* 225:740–753.
- 403 [40] Sporbert, M., D. Jakubka, S. F. Bucher, I. Hensen, M. Freiberg, K. Heubach, A. Konig, B. Nordt,  
404 C. Plos, I. Blinova, A. Bonn, B. Knickmann, T. Koubek, A. Linstadter, T. Maskova, R. Primack,  
405 C. Rosche, M. A. Shah, A.-D. Stevens, K. Teilborger, S. Trager, C. Wirth, and C. Romermann.  
406 2022. Functional traits influence patterns in vegetative and reproductive plant phenology – a  
407 multi-botanical garden study. *New Phytologist* 235:2199–2210.
- 408 [41] Stan Development Team. 2018. RStan: the R interface to Stan. R package version 2.17.3.
- 409 [42] Sun, S., and L. E. Frelich. 2011. Flowering phenology and height growth pattern are associated  
410 with maximum plant height, relative growth rate and stem tissue mass density in herbaceous  
411 grassland species. *Journal of Ecology* 99:991–1000.
- 412 [43] Tryjanowski, P., M. Rybacki, and T. Sparks. 2003. Changes in the first spawning dates of common  
413 frogs and common toads in western Poland in 1978-2002. *Annales Zoologici Fennici* 40:459–464.
- 414 [44] Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung, and J. Messier.  
415 2012. The return of the variance: Intraspecific variability in community ecology. *Trends in Ecology  
416 and Evolution* 27:244–252.

- 417 [45] Vitasse, Y., S. Ursenbacher, G. Klein, T. Bohnenstengel, Y. Chittaro, A. Delestrade, C. Monnerat,  
418 M. Rebetez, C. Rixen, N. Strelbel, B. R. Schmidt, S. Wipf, T. Wohlgemuth, N. G. Yoccoz, and  
419 J. Lenoir. 2021. Phenological and elevational shifts of plants , animals and fungi under climate  
420 change in the European Alps. *Biological Reviews* 96:1816–1835.
- 421 [46] Wainwright, C. E., E. M. Wolkovich, and E. E. Cleland. 2012. Seasonal priority effects : implications  
422 for invasion and restoration in a semi-arid system. *Journal of Applied Ecology* 49:234–241.
- 423 [47] Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*  
424 199:213–227.
- 425 [48] Westoby, M., and I. J. Wright. 2006. Land-plant ecology on the basis of functional traits. *Trends*  
426 in *Ecology and Evolution* 21:261–268.
- 427 [49] Wolkovich, E. M., and E. E. Cleland. 2011. The phenology of plant invasions: A community  
428 ecology perspective. *Frontiers in Ecology and the Environment* 9:287–294.
- 429 [50] ———. 2014. Phenological niches and the future of invaded ecosystems with climate change. *AoB*  
430 *PLANTS* 6:1–16.
- 431 [51] Wolkovich, E. M., and M. J. Donahue. 2021. How phenological tracking shapes species and  
432 communities in non-stationary environments. *Biological Reviews* 96:2810–2827.
- 433 [52] Wolkovich, E. M., and A. K. Ettinger. 2014. Back to the future for plant phenology research. *New*  
434 *Phytologist* 203:1021–1024.
- 435 [53] Wright, I. J., M. Westoby, P. B. Reich, J. Oleksyn, D. D. Ackerly, Z. Baruch, F. Bongers,  
436 J. Cavender-Bares, T. Chapin, J. H. C. Cornellissen, M. Diemer, J. Flexas, J. Gulias, E. Garnier,  
437 M. L. Navas, C. Roumet, P. K. Groom, B. B. Lamont, K. Hikosaka, T. Lee, W. Lee, C. Lusk, J. J.  
438 Midgley, Ü. Niinemets, H. Osada, H. Poorter, P. Pool, E. J. Veneklaas, L. Prior, V. I. Pyankov,  
439 S. C. Thomas, M. G. Tjoelker, and R. Villar. 2004. The worldwide leaf economics spectrum.  
440 *Nature* 428:821–827.




Figure 1: Leaf and wood traits follow a gradient that varies from acquisitive versus conservative growth strategies, which may also include phenology. We predicted early-budbursting species would have traits associated with acquisitive growth, as they are more likely to experience greater risk of frost but reduced competition. In contrast, we expected later-budbursting species would have traits associated with conservative growth, as they experience greater competition but a more climatically benign environment.




Figure 2: Estimated species-level responses to standardized forcing (left column: a, d, g & j), chilling (middle column: b, e, h & k), and photoperiod treatments (right column: c, f, i & l) predicted by estimated trait values for height (first row: a-c), SLA (second row: d-f), log<sub>10</sub> Seed mass (third row: g-i), and LNC (fourth row: j-l). We estimated parameters using a joint trait-phenology model, with the black line depicting the mean linear relationship between estimated trait effects and the slope of the cue response and the pink band the 50% uncertainty interval. Each set of crossed lines represents one species (with each line spanning the 50% uncertainty interval), with the species depicted in Fig 3 colored in each panel, with the acquisitive species (*Alnus incana*) shown in red, and the conservative species (*Quercus rubra*) shown in blue.




Figure 3: We expected species with traits associated with acquisitive (e.g., smaller heights and small seed mass) versus conservative (e.g., taller with larger seeds) growth strategies would have different budburst responses to phenological cues. Shown here is an example of the cue relationships with height (**a-c**) and seed mass (**d-f**) for an acquisitive species, *Alnus incana* shown in red, and a conservative species, *Quercus rubra*, shown in blue. **a**, The effect of height on budburst timing was smaller in response to forcing cues, but larger in response to both **b**, chilling and **c**, photoperiod. In contrast, seed mass had a negligible effect on **d**, forcing and **f**, photoperiod responses, **e**, but a greater response to chilling. Band represent the 50% uncertainty intervals of the model estimates. The coloured bands represent the 50% uncertainty intervals of the model estimates and points individual trait measurements.




Figure 4: Traits differed in the relative magnitude of their species-level and study-level variation, with only a. the height model estimating greater study-level variation than species-level, while our b. specific leaf area, c. leaf nitrogen content, and d. seed mass models all estimated higher species-level variation. Shown here are the posterior densities for the species-level variation, shown in red, and study-level variation, shown in yellow. For comparison, we show all x-axes spanning the minimum to maximum variances across all four traits. The histograms depict the full distribution of the study and species-level variance, where each bin is normalized by the total count of the posterior estimate.