

Spatial and temporal shifts in photoperiod with climate change

A.K. Ettinger, D. Buonaiuto, C. Chamberlain, I. Morales-Castilla, E. Wolkovich

August 3, 2018

1. *Introduction*

- (a) Photoperiod is a critical cue used by organisms to synchronize their activities with seasonal climatic changes (e.g., Hsu et al., 2011; Singh et al., 2017; Basler and Körner, 2012). Diverse important responses, from the timing of mating in marsupials (McAllan et al., 2006) and pheromone production in moths (Linn et al., 1996), to growth rates in salmon and the timing of budburst in woody plants (Flynn and Wolkovich, In Review.), are affected by photoperiod. Daylength is consistent across years, especially compared to other seasonal cues such as temperature and precipitation. Thus, it may act as a temporally stable signal by which organisms can optimally align events with the season (Saikkonen et al., 2012).
- (b) Despite the relative stability and predictability of daylength over time, as species undergo climate change-induced shifts in their space (i.e., shifting distributions) and time (i.e., shifting phenology), the daylength they experience will be altered. The implications of these shifts in photoperiod regime are rarely explored.
- (c) Many experiments have altered photoperiod, often interacting with temperature changes; however, photoperiod treatments in these experiments are not typically designed to be applied to climate change forecasting.
- (d) Here, we ask:
 - i. How will climate change alter the photoperiod experienced by organisms, given observed climate change-induced biological shifts, both spatially and temporally?
 - ii. What are the implications of these altered photoperiods for forecasts of climate change impacts?
 - iii. Can the large quantity of experiments altering photoperiod be applied to forecasting biological implications of climate change (i.e., do they occur at the appropriate scale)?

2. *How will climate change alter the photoperiod experienced by organisms?*

- (a) Spatial shifts in species ranges and temporal shifts in species phenology and activity will alter the photoperiods experienced by organisms under climate change.
- (b) To date, most work has focused on how spatial range shifts with climate change will affect photoperiod (Saikkonen et al., 2012), but temporal shifts are actually likely to yield bigger changes in experienced photoperiod than spatial shifts (Figure 1).
- (c) Shifts in photoperiod may vary with latitude and time of year, especially since phenology varies with latitude (Figure 2).
- (d) Photoperiod sensitivity can also vary with latitude (Howe et al., 1996; Saikkonen et al., 2012; Partanen et al., 2005; Viherä-Aarnio et al., 2006; Caffarra et al., 2011; Gauzere et al., 2017), so it is unclear how these two things interact. Cat has added a text file to the latitude analysis folder that summarizes papers that have looked at latitudinal effects on photoperiod.

3. *What are the implications of these altered photoperiods for forecasts of climate change impacts?*
 - (a) Phenology will be altered, given that daylength is known to affect vegetative growth, cell elongation, and budburst (Linkosalo and Lechowicz, 2006; Erwin, 1998; Sidaway-Lee et al., 2010; Hsu et al., 2011).
 - (b) It has been proposed that photoperiod may eventually become a limiting factor, constraining the ability of trees to shift their phenology with warming (Koerner and Basler, 2010; Vitasse and Basler, 2013; Morin et al., 2010).
 - (c) Interactions between photoperiod and forcing and chilling could result in muted or exaggerated phenological shifts, compared to what would be expected based on temperature change alone. Say something about crossing thresholds of daylength and the "external coincidence model" for photoperiod control (Bastow and Dean, 2002; Kobayashi and Weigel, 2007; Andrés and Coupland, 2012; Singh et al., 2017)?
 - (d) Effects of photoperiod on forecasting of biological impacts of climate change needs additional investigation. In some forecasting methods (e.g. species distribution modelling), the role of photoperiod is largely ignored (I think this is true? add some citations). In other cases, photoperiod is incorporated into forecasts, along with other variables such as evaporative demand, and temperature (e.g. ED Jolly et al., 2005; Medvigy et al., 2013). These models need to be more widely tested, e.g. in different ecosystems/species, and need to incorporate recent findings about the role of photoperiod in phenology.
4. *Can existing experiments be applied to forecasting?*
 - (a) In some cases, yes, experiments manipulate photoperiod at relevant scales (Figure 3, Table 1).
 - (b) However, most experiments manipulate photoperiod much more dramatically than will occur with climate change (Figures 3, 4, 5, but see (Basler and Körner, 2012)), so it is difficult to extrapolate findings. (This may not be true for all latitudes- for example high latitudes experience more dramatic changes in photoperiod across the year.)
 - (c) There is a great need to better understand exactly how photoperiod acts as a cue. The divergent effects of photoperiod observed across studies (e.g., Figure 6) suggests that photoperiod interacts with other environmental drivers, such as chilling and forcing, to affect phenology and other activities. However, exactly how it interacts with temperature to break dormancy, as well as the type of response it elicits (e.g., linear versus non-linear threshold) is unclear.
5. *Conclusions*
 - (a) Organisms may experience large changes to the photoperiod they experience, under climate change, even if they do not shift their ranges spatially.
 - (b) More studies needed with fine-scale changes in photoperiod.

To do:

1. Make Table of studies testing if photoperiod varies by latitudinal origin- cat started on this (Table 2)?
2. Update table/map to use 3 ER studies.

Random notes that may be useful:

1. Bradshaw and Holzapfel (2001) showed that the pitcher plant mosquito, *Wyeomyia smithii*, has evolved a shorter critical photoperiod in association with a longer growing season. Northern populations of this

mosquito now use a shorter day-length cue to enter winter diapause, doing so later in the fall than they did 24 years ago.

2. Decreasing day-length is the main environmental cue inducing growth cessation and bud set in many perennial plants, including poplar
 - (a) Lagercrantz U: At the end of the day: a common molecular mechanism for photoperiod responses in plants?. *J Exp Bot.* 2009, 60: 2501-2515. 10.1093/jxb/erp139.
 - (b) Howe GT, Gardner G, Hackett WP, Fournier GR: Phytochrome control of short-day-induced bud set in black cottonwood. *Physiol Plant.* 1996, 97: 95-103. 10.1111/j.1399-3054.1996.tb00484.x.
3. Response to photoperiod is under strong genetic control
 - (a) Bradshaw HD, Stettler RF: Molecular genetics of growth and development in *Populus*. IV. Mapping QTLs with large effects on growth, form, and phenology traits in a forest tree. *Genetics.* 1995, 139: 963-973.
 - (b) Keller SR, Soolanayakanahally RY, Guy RD, Silim SN, Olson MS, Tiffin P: Climate-driven local adaptation of ecophysiology and phenology in balsam poplar, *Populus balsamifera* L. (Salicaceae). *Am J Bot.* 2011, 98: 99-108. 10.3732/ajb.1000317.
 - (c) Weih M: Intensive short rotation forestry in boreal climates: present and future perspectives. *Can J Forest Res.* 2004, 34: 1369-1378. 10.1139/x04-090.

References

- Andrés, F., and G. Coupland. 2012. The genetic basis of flowering responses to seasonal cues. *Nature reviews. Genetics* 13:627.
- Basler, D., and C. Körner. 2012. Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology* 165:73–81.
- Bastow, R., and C. Dean. 2002. The molecular basis of photoperiodism. *Developmental cell* 3:461–462.
- Caffarra, A., A. Donnelly, I. Chuine, and M. B. Jones. 2011. Modelling the timing of *Betula pubescens* bud-burst. i. temperature and photoperiod: A conceptual model. *Climate Research* 46:147.
- Duputié, A., A. Rutschmann, O. Ronce, and I. Chuine. 2015. Phenological plasticity will not help all species adapt to climate change. *Global change biology* 21:3062–3073.
- Erwin, J. E. 1998. Temperature and light effects on stem elongation (plant growth regulation by physical and mechanical stimuli, for further development of horticulture in east asia). *Journal of the Japanese Society for Horticultural Science* 67:1113–1120.
- Flynn, D. F. B., and E. M. Wolkovich. In Review. Temperature and photoperiod drive spring phenology across all species in a temperate forest community. .
- Gauzere, J., S. Delzon, H. Davi, M. Bonhomme, I. G. de Cortazar-Atauri, and I. Chuine. 2017. Integrating interactive effects of chilling and photoperiod in phenological process-based models. a case study with two european tree species: *Fagus sylvatica* and *Quercus petraea*. *Agricultural and Forest Meteorology* 244:9–20.
- Howe, G. T., G. Gardner, W. P. Hackett, and G. R. Fournier. 1996. Phytochrome control of short-day-induced bud set in black cottonwood. *Physiologia Plantarum* 97:95–103.
- Hsu, C.-Y., J. P. Adams, H. Kim, K. No, C. Ma, S. H. Strauss, J. Drnevich, L. Vandervelde, J. D. Ellis, B. M. Rice, et al. 2011. Flowering locus t duplication coordinates reproductive and vegetative growth in perennial poplar. *Proceedings of the National Academy of Sciences* 108:10756–10761.

- Jolly, W. M., R. Nemani, and S. W. Running. 2005. A generalized, bioclimatic index to predict foliar phenology in response to climate. *Global Change Biology* 11:619–632.
- Kobayashi, Y., and D. Weigel. 2007. Move on up, it's time for change—mobile signals controlling photoperiod-dependent flowering. *Genes & development* 21:2371–2384.
- Koerner, C., and D. Basler. 2010. Warming, photoperiods, and tree phenology response. *Science* 329:278–278.
- Linkosalo, T., and M. J. Lechowicz. 2006. Twilight far-red treatment advances leaf bud burst of silver birch (*betula pendula*). *Tree physiology* 26:1249–1256.
- Linn, C. E., M. G. Campbell, K. R. Poole, W.-Q. Wu, and W. L. Roelofs. 1996. Effects of photoperiod on the circadian timing of pheromone response in male *trichoplusia ni*: relationship to the modulatory action of octopamine. *Journal of Insect Physiology* 42:881–891.
- McCallan, B. M., C. R. Dickman, and M. S. Crowther. 2006. Photoperiod as a reproductive cue in the marsupial genus *antechinus*: ecological and evolutionary consequences. *Biological Journal of the Linnean Society* 87:365–379.
- Medvigy, D., S.-J. Jeong, K. L. Clark, N. S. Skowronski, and K. V. Schäfer. 2013. Effects of seasonal variation of photosynthetic capacity on the carbon fluxes of a temperate deciduous forest. *Journal of Geophysical Research: Biogeosciences* 118:1703–1714.
- Morin, X., J. Roy, L. Sonié, and I. Chuine. 2010. Changes in leaf phenology of three European oak species in response to experimental climate change. *New Phytologist* 186:900–910.
- Partanen, J., H. Hänninen, and R. Häkkinen. 2005. Bud burst in Norway spruce (*picea abies*): preliminary evidence for age-specific rest patterns. *Trees* 19:66–72.
- Saikkonen, K., K. Taulavuori, T. Hyvönen, P. E. Gundel, C. E. Hamilton, I. Vänninen, A. Nissinen, and M. Helander. 2012. Climate change-driven species' range shifts filtered by photoperiodism. *Nature Climate Change* 2:239.
- Sidaway-Lee, K., E.-M. Josse, A. Brown, Y. Gan, K. J. Halliday, I. A. Graham, and S. Penfield. 2010. Spatula links daytime temperature and plant growth rate. *Current Biology* 20:1493–1497.
- Singh, R. K., T. Svystun, B. AlDahmash, A. M. Jönsson, and R. P. Bhalerao. 2017. Photoperiod-and temperature-mediated control of phenology in trees—a molecular perspective. *New Phytologist* 213:511–524.
- Viherä-Aarnio, A., R. Häkkinen, and O. Junntila. 2006. Critical night length for bud set and its variation in two photoperiodic ecotypes of *betula pendula*. *Tree physiology* 26:1013–1018.
- Vitasse, Y., and D. Basler. 2013. What role for photoperiod in the bud burst phenology of European beech. *European Journal of Forest Research* 132:1–8.

Tables

Table 1: **Growth chamber experiments and their photoperiod treatments**, compared to the spatial and temporal shifts required for organisms to experiments photoperiod changes equivalent to those treatments. For shifts in space, ‘ER’ indicates that the photoperiod treatments exceeds the change of photoperiod from moving up to 40 degrees latitudinally on June 21. For shifts in time, ‘ER’ indicates that the range of photoperiod treatments exceeds the change in daylengths at that latitude during the entire year. ‘max NA’ indicates that the maximum daylength treatment does not exist at that latitude; ‘min NA’ indicates that the minimum daylength treatment does not exist at that latitude.

idstudy	continent	lat	long	day_range	delta	space	time
ashby62_exp1	north america	42.99	-89.41	8-16	4.00	18.2	min NA (9)
basler14_exp1	europe	46.31	8.27	9.2-16	1.00	6	-22
caffarra11b_exp2	europe	52.32	-6.93	10-16	2.00	7.5	-30
devries82_exp1	europe	51.98	5.66	8-24	16.00	ER	ER
falus190_exp1	europe	46.03	10.75	9-13	4.00	16	-82
falus196_exp3	europe	38.27	15.99	9-13	4.00	21.6	-111
ghelardini10_exp1	europe	43.72	11.37	8-16	8.00	21.9	ER
heide05_exp1	europe	56.18	-4.32	10-24	14.00	ER	ER
heide08_exp1	europe	48.40	11.72	10-24	14.00	ER	ER
heide11_exp1	europe	59.67	10.67	10-20	10.00	ER	max NA (18.7)
heide12_exp1	europe	56.50	-3.06	10-24	5.00	8.9	-64
heide15_exp2	europe	56.50	-3.06	10-15	1.00	3.2	-13
heide93_exp1	europe	59.50	10.77	8-24	16.00	ER	ER
heide93a_exp1	europe	59.67	10.83	8-24	16.00	ER	ER
heide93a_exp3	europe	47.50	7.60	13-16	1.00	5.7	-18
howe95_exp1	north america	40.55	-124.10	9-24	2.00	13.1	-64
laube14a_exp1	europe	48.40	11.71	8-16	4.00	14.3	-87
myking95_exp1	europe	56.10	9.15	8-24	16.00	ER	ER
myking97_exp1	europe	59.67	10.77	12-24	12.00	ER	max NA (18.7)
nienstaedt66_exp1	north america	44.17	-103.92	8-20	12.00	ER	ER
okie11_exp1	north america	32.12	-83.12	0-12	12.00	ER	ER
partanen01_exp1	europe	61.93	26.68	6-16	10.00	ER	-105
partanen05_exp1	europe	61.82	29.32	5-20	5.00	ER	-67
partanen98_exp1	europe	60.03	23.05	8.66-12	3.34	5.1	-37
pettersen71_exp1	europe	59.66	10.77	10-24	2.00	4	-23
Sanz-Perez09_exp1	europe	40.40	-3.48	10-16	6.00	23.6	ER
skuterud94_exp1	europe	61.50	24.33	8-24	16.00	ER	ER
viheraaarnio06_exp1	europe	60.45	24.93	16-17	1.00	2.1	-12
viheraaarnio06_exp1	europe	67.73	24.93	20-21	1.00	ER	-5
viheraaarnio06_exp2	europe	60.45	24.93	15-19	4.00	5.1	-62
viheraaarnio06_exp2	europe	67.73	24.93	22-23	1.00	ER	-3
worrall67_exp 3	north america	41.31	-72.93	8-16	8.00	24.3	ER
zohner16_Exp1	europe	48.16	11.50	8-16	8.00	ER	ER

Figures




Figure 1: **Photoperiod varies with latitude and throughout the year**, such that temporal shifts in activity yield larger changes in experienced photoperiod compared with spatial shifts. Here, we show this variation at two latitudes, using hypothetical rates of spatial and temporal shifts: 30 days earlier for temporal shifts, and 0.5 degrees poleward for spatial shifts. These shifts, which are similar to observed average rates (e.g., Parmesan et. al 2006, Chen et al 2011), highlight the greater magnitude in daylength changes close to the equinox (e.g., DOY 91), versus close to the summer solstice (e.g., DOY 182).




Figure 2: **The photoperiod on the green up date (start of spring) varies over space and among years.** Hours of daylight on the date of spring green up from MODIS satellite data across North America and Europe for an average (a) and early (b) North American start of spring. The differences between the years are shown in (c).




Figure 3: **OSPREE experiments that manipulate photoperiod**, and their equivalent spatial and temporal shifts, mapped (A), and graphed (B-C). Observed rates (dashed gray lines) 16.9 kilometers per decade (or approximately 1.5 degrees in 100 years) for spatial shifts (Chen et al. 2011) and 2.3 days per decade (or 23 days in 100 years) for temporal shifts (Parmesan and Yohe 2003).




Figure 4: **Experimental treatments of daylength in the OSPREE database, shown by latitude (A) and by day of budburst (B)** for *Fagus sylvatica*. For comparison, we show the daylength when budburst occurs in its current and projected ranges (A) and in its current range only, with expected shifts in phenology (B). Estimates and projections are from Phenofit (Duputié et al., 2015)




Figure 5: Experimental treatments of daylength in the OSPREE database, shown by latitude (A) and by day of budburst (B) for *Quercus robur*. For comparison, we show the daylength when budburst occurs in its current and projected ranges (A) and in its current range only, with expected shifts in phenology (B). Estimates and projections are from Phenofit (Duputié et al., 2015).




Figure 6: Plant responses to changes in daylength vary across species and populations, and with the amount of chilling received.