

A four-step Bayesian workflow for improving ecological science

EM Wolkovich, TJ Davies, WD Pearse & M Betancourt

February 25, 2024

Abstract

Growing anthropogenic pressures have increased the need for robust predictive models. Meeting this demand requires approaches that can handle bigger data to yield forecasts that capture the variability and underlying uncertainty of ecological systems. Bayesian models are especially adept at this and are growing in use in ecology. Yet many ecologists today are not trained to take advantage of bigger ecological data and more flexible robust models. We argue this stems from a divide in training of ‘theoretical’ versus ‘empirical’ ecologists that fails to recognize the demands and needs of modern ecology. Here we describe a broadly generalizable workflow for statistical analysis centered on simulating data from models, and show how it can [reshape](#) training in ecology. Building on the increasingly computational toolkit of many ecologists, this approach represents a ground shift of not just how to build models, but how to approach model fitting to advance our science. By integrating model building and testing more fully with ecological theory, this simulation-based workflow helps fit models that are more robust and well-suited to provide new ecological insights. This in turn can help us refine where to put resources for better estimates, better models, and better forecasts.

Introduction

~~Recent years have seen an explosion in the use of Bayesian models across scientific fields (van de Schoot *et al.*, 2021; Schad *et al.*, 2021; Grinsztajn *et al.*, 2021). This comes in part from increased computational power, but mostly from new algorithms (e.g. Hamiltonian Monte Carlo, Hoffman & Gelman, 2014; Betancourt, 2019) that have made fitting and implementing Bayesian models faster, more robust and—in many ways—easier (Carpenter *et al.*, 2017). Ecologists are exploiting these advances, using the power and flexibility of Bayesian methods to help address the global challenges society is facing.~~

As ecologists work to develop global predictive models, they have developed ever larger datasets (Hampton *et al.*, 2013), but bigger data is messier data. Such data generally requires a model of both the underlying biological processes and how the measurements were made. While some fields have long used these types of models (generally in fields focused on inferring population sizes of things people want to eat or manage, Muthukumarana *et al.*, 2008; Zheng *et al.*, 2007; Trijoulet *et al.*, 2018; Strinella *et al.*, 2020), most have not. Thus many have tried to adapt what they were trained in—traditional statistical methods (e.g., F and t tests) and a strong focus on null hypothesis testing. Often they have done this by fitting multi-way interaction terms, using random effects to correct for group level factors, or comparing across a large suite of models.

But these approaches do not often align with ecology’s aims today. Beyond the reality that most traditional methods are fragile when used beyond the cleaner, simpler experiments these

methods assume (e.g. spatial, temporal and phylogenetic correlations often violate common independence assumptions), they will usually fail to produce robust, reproducible results. Multi-way interaction terms can make main effects hard to interpret and require much larger sample sizes to estimate reliably. Null hypotheses are rarely true and can lead to confusion over what is important versus ‘significant’ (Gelman & Hill, 2009; Muff *et al.*, 2022). Many model comparison and machine learning approaches prefer models whose inferences match the idiosyncrasies and biases in the available data, but don’t generalize. The disconnect between the traditional statistical tools available in ecology and the field’s fundamental theory is being made ever-clearer as the need for robust ecological forecasts grows.

Bayesian approaches provide a pathway to build powerful models that can transform how we understand our systems as large scale ecological data become increasingly available. Recognizing this, many in ecology are increasingly using Bayesian approaches, ~~but are often inadequately prepared to notice or manage the pitfalls of such models.~~ Many of these pitfalls can be avoided, by approaching analyses through specific workflows (Betancourt, 2020; Grinsztajn *et al.*, 2021; van de Schoot *et al.*, 2021), which themselves are built on a process of how to do not just statistics, but how to do science (Box, 1976).

Such approaches move away from a focus on null hypothesis testing, towards estimating effect sizes, using models calibrated (see Table 1) and better understood through simulating data at multiple steps—~~using a number of skills often reserved in ecology more for ‘theorists’ than empirical ecologists. We argue this theoretical vs empirical divide is antiquated and limits progress by isolating the insights stimulated from model building. Further, it ignores that the average modern ecologist is computational, and thus already has many of the basic skills to build bespoke models.~~

Here we provide a simplified—but powerful—workflow that builds on new insights from statistics (Betancourt, 2020; van de Schoot *et al.*, 2021) and the increasingly computational nature of ecology today. By integrating model building more fully with ecological theory and understanding—and vice versa—this approach can fit models that are more robust and better-suited to providing new ecological insights and improved predictions. ~~After outlining the workflow, we discuss how it can benefit Bayesian inference generally, has improved our own science, and suggests a new way to train in ecology.~~

A basic Bayesian workflow

Statistical analyses are designed for inference—to learn about some process, effect or behavior from data. Robust analyses yield inferences consistent with the underlying truth more often than not (quantifying this consistency is called calibration, and is a critical part of using models for inference, see Table 1). Because we do not know the ‘truth’ how we approach our analyses—and our inference—is a critical component of how we do science. ~~Doing this poorly~~ can lead science away from the type of repeatable and generalizable findings expected when inferences align with the truth. For example, an overly zealous focus on p -values has led to a replication crises in several fields, where results seem most likely the outcome of noisy data combined with a search for statistical significance through many models, effectively a garden of forking paths (Halsey *et al.*, 2015; Loken & Gelman, 2017). Common model selection approaches, including new machine learning methods, try to avoid this by comparing across models, but ~~rarely~~ generalize to provide useful forecasts.

We argue that robust analyses comes from explicitly building and challenging a model (or set of models) in an organized sequence of steps—a workflow. We outline this workflow using a Bayesian approach ~~because we believe this makes the steps more straightforward (with an example in the supplement shown in R and Stan). Our general approach, however, extends to other statistical inference methods.~~

In ecology, Bayesian methods often shift focus away from null hypothesis testing and point estimates and towards evaluating more fully a model’s output, including uncertainty—often using a specific model designed for the system and question at hand. ~~These bespoke models can be fit by applying Bayes’ theorem, which generates a *posterior* distribution from a combination of a *likelihood* and a *prior* distribution (an initial uncertainty estimate derived from basic ecological knowledge), and using iterative algorithms (e.g. MCMC, Markov Chain Monte Carlo) that provide samples that can be used to extract information from the posterior distribution (for more, see *A brief review of statistical inference using Bayesian approaches* in the Supplement).~~

~~We outline a workflow that includes~~ what we consider the major steps for Bayesian model fitting (Fig. 1). The workflow includes model development (Step 1) and calibration (Step 2), inference (Step 3) and then model improvement (Step 4). Many of these steps will be familiar to statistical ecologists, but are often overlooked, whereas other steps may appear particular to Bayesian (e.g. prior predictive checks), but are actually useful for anyone—using Bayesian models or not—to challenge their models of how the world works. Parts of this workflow could be dropped, or expanded as workflows in themselves, given other aims (see Supplement: Which workflow?). Relatedly, as we do not explain exactly how to implement each step, many of the smaller but still critical steps are omitted, including visualization, which is required at every step (and for which there are many good resources, e.g. Gabry *et al.*, 2019).

Step 1: Develop your model(s)

We start the workflow with what can feel like the biggest step—build a model (or, potentially at this step, models) that you want to fit your data to based on your aims. By developing a model designed for your biological question, data and aims, your statistical workflow naturally becomes a scientific workflow. You will more clearly see the assumptions and mechanisms in your model, which is especially valuable given how often our intuition of how models ‘work’ is wrong (Kokko, 2005).

You likely already have a model, though it may only verbal or conceptual. For this workflow, however, you’ll need to convert such models into mathematical versions (Servedio *et al.*, 2014). This step is better approached before you collect your data. After data collection, it becomes far more tempting to focus on the particular details in the data and not the latent processes and biological models from which the data were generated.

Getting to the point where this step is part of your data design and collection, however, requires starting somewhere—with some model (or models) in hand. A suite of resources for ‘generative’ or ‘narratively generative’ modeling can help (MacElreath, 2016; Betancourt, 2021b), along with two points. First, ~~you must start somewhere, so~~ know that you can and will improve on this skill. Second, as you start, ask lots of questions—and push yourself on your answers—about what you expect and what’s reasonable biologically from your model. For example, instead of simply identifying which distribution your response variable looks most similar to, ask yourself what ecological processes generate that distribution and how they influence its properties (e.g., its mean, minimum and maximum). Do you expect data below zero? Up to infinity? If not,

why not? Effective model building is about efficient brainstorming and this is a critical part of the process (see Supplement).

As you do this, you'll be generating your model—including its priors. Assigning priors generally forces you to think about your model with regards to your study system, and interrogate what's probable, possible or actually unreasonable. While many packages (e.g., `brms`, `rstanarm`, which fit a suite of pre-defined models) will automatically set default priors, assigning them yourself can quickly disabuse users of their prejudices. For example, you may not think you have a prior on how sunlight affects plant growth, until you realize your 'agnostic prior' actually allows plants to grow hundreds of meters per day.

Step 2: Check your model on simulated data

Once you have your model and its priors jotted down, you need to write up your model in a particular modeling language and check it. As with all code: just because it runs, does not mean it does what you think it does. Whether writing it out in `Stan`, where you need to be able to write out the full likelihood and set all your own priors, or using a package that writes much of the model for you (e.g. `rstanarm`), you need a way to verify the code is correct: test data.

Test data (aka 'simulated data', or 'fake data,' etc.), and the skills required to build it, are central to this workflow. With 'test data' you simulate data from your model in such a way that you can use the resulting data to test if your model code is correct (i.e., you fix values for your model parameters, then test how well your model recovers them, see the Supplement for an example). While there's no guarantee that inferences will always recover the parameter values you set, even when using the correct model, extreme disagreement is usually a good indicator that something is amiss. At the same time these simulation studies can help us understand how often our model might lead to the correct inference (see Fig. 2), and can be easily adapted into formal tests of power. As you do this, you will also be calibrating your model—seeing how close it estimates parameters you set and under what conditions.

This very basic model checking step is uncommon for many ecologists, but critical in our view. If you can simulate data from your model, then you can powerfully—and easily—answer questions related to statistical power, what effect sizes are reasonable, and—most likely—have new insights into how your model suggests the world works, all before looking at any real data. 'All models are wrong; some models are useful,' becomes much clearer when you have the power to generate data from your model under any parameter set and sample size you want. Conversely, if you cannot complete this step, you'll struggle to understand if the model fit well, and struggle further to meaningfully interpret the model output, making an apparently simple programmatic task actually encapsulate a far deeper understanding of your model.

Once done with this basic check of your code, you need to further explore the model world to which you're planning to fit your data via prior predictive checks. For these, you draw values from your **prior distribution** (usually randomly in your code) and then explore how your model performs under those draws. Seeing how this influences your resulting output reveals the extent to which your model can capture known variation in your data, and gives insight into whether your model is capable of distinguishing among competing hypotheses. It also serves as a check on the priors you're using (addressing one of the common concerns of those inexperienced with Bayesian models). How exactly to do this depends on your question, model and aims, but many guides can help you think through this (Betancourt, 2021a; Wesner & Pomeranz, 2021; Winter & Depaoli, 2023).

Step 3: Run your model on your empirical data

The next step is to run the model—you’ve now validated, test-run and have ready to go—on your exciting new empirical data. Check diagnostics so you know it’s running well and adjust until it is (this includes a suite of convergence and efficiency metrics that are well-discussed elsewhere, Betancourt, 2020; Gelman *et al.*, 2020; van de Schoot *et al.*, 2021; Gabry *et al.*, 2019). A model that doesn’t converge, or seems to suggest coefficients that are completely at odds with the data, can result from a model that was mis-specified and could never capture real-world variation, thus for big issues you may want to return to Steps 1 and 2.

This is the step many ecologists skip straight to, ourselves included. It’s easy to see the appeal: this is the inference step and where you have the opportunity to see new ecological insights! Fitting our new data to the model can feel like the moment when we’ll learn something new. But, at least in our experience, this is not always the case. When we rush to this step, that first model we fit is often followed by another, and another—perhaps because one does not converge, or the results of another do not make immediate sense. And we can get distracted from what we are actually most interested in—the inference into ecology. Following this workflow can make this step much more satisfying. Here the benefits of the workflow may become excitedly apparent: we have estimates in useful units with uncertainty we can understand ~~and explain more easily than traditional p -values~~. We can use this information to draw new conclusions, design new experiments and more—but this is also a point to stop and check your model.

Step 4: Check your model on data simulated from your empirical model output (also known as posterior retrodictive checks)

Once you have your posterior based on your model and new empirical data, it’s time to remember that it’s wrong (‘all models are wrong...’) and ask how useful it is. You can do some of this through diagnostics, such as R^2 , which compares point predictions to the observed data. With a posterior, however, you can compare an entire distribution of predictions to the observed data.

This is where simulating from your model can be especially insightful. It will not only indicate that the model isn’t adequately fitting the data but also can suggest what the problems might be. Steps 1-2 have set you up well for this, as you have a sense of what different parameter estimates do to the model, and test data provide a sense of how it works on data similar to yours. Now ~~that you have~~ the parameter estimates from your posterior you can simulate new data from them and see how that new world looks to you—called posterior retrodictive checks (or posterior predictive checks, Fig. 3). Exactly how to do these are—again—dependent on your question, model and aims (but there is lots written on this, Held *et al.*, 2010; Gelman *et al.*, 2000; Conn *et al.*, 2018).

Often here you may find big differences from your empirical data, and can start to generate hypotheses for why. For example, you may find patterns that suggest missing grouping factors (e.g., site or biome) through visualization, ~~and by grouping posteriors by that factor~~, or you may quickly realize your model predicts impossible numbers for your biological reality. You may begin to see inadequacies in your model—or potentially your data—which is one of benefits of the workflow: models don’t fail silently, they fail with a wealth of context the helps to generate new models and better experiments.

Feedbacks & workflows

~~Depending on your aims (which you set up in Step 1), you may very well find you want to tweak your model—this is all part of the workflow. At that point you return to the beginning, tweak your model, and repeat the rest of the workflow again. In this way, fitting multiple models is encouraged, but is distinct from the quest for a minimum adequate model or one ‘best’ fit. Feedbacks in this workflow are focused far more on what is biologically reasonable, and understanding the utility—and limits—of inference from your data for your model, given your goals. And there are big benefits to it.~~

This process more fully integrates mathematical modeling into statistical modeling. To complete Steps 1-2, you have to understand the underlying math of your model enough to simulate data from it. This can be challenging at first (e.g. recalling how to simulate y data for a simple linear regression is not straightforward when you never do it), but is immensely beneficial to forcing you to understand your model and its consequences. Indeed, we have found the greatest insights come not from the step we all know best—fitting the model with empirical data—but from every other step in this workflow. ~~Further, it exploits many of the benefits of Bayesian approaches, while systematically avoiding major pitfalls.~~

How this workflow helps avoid potential pitfalls of Bayesian inference

Bayesian models have many benefits, but an often-mentioned one is that ‘you can fit any model you want.’ While this is not entirely true (Gelman *et al.*, 2014; Beaumont, 2019), Bayesian modeling options can feel limitless when compared to the models ecologists can fit in popular modeling packages (e.g. `lme4`). As long as you can write out the likelihood of your desired model (and sometimes even if you can’t, Sunnåker *et al.*, 2013) and assign priors to all parameters, you can generally ‘fit’ the model. Such flexibility is incredibly powerful in ecology where data are often influenced by complex spatial or temporal patterns, non-linear processes are widespread, and common data types are non-Gaussian (e.g. counts, percent cover, etc.). ~~Fitting a bespoke model to data also yields numbers we often really want but don’t have access to in other approaches. Models can be specifically designed to estimate and report effects in relevant units (e.g., per °C of warming)—always with estimated uncertainty.~~

The valuable flexibility of Bayesian models is often mentioned as one of their greatest pitfalls: you can fit almost whatever you want, but critical parts of your model might be almost entirely unimpacted by your data. In ecological model fitting, we’re currently most often interested in parameter estimates strongly informed by our data, making this problem sound especially dangerous. In reality, however, this problem is not related to modeling, but to experimental design—and a flawed experimental design leading to low power for your model is much easier to see through this workflow compared to using traditional null hypothesis testing methods.

Priors are another major source of concern for those unfamiliar with Bayesian approaches. Often treated as the big bad wolf of Bayesian, or the unseen hand producing the model fits you get, according to some. In reality, How much priors influence your model fit is up to your model and your data. Depending on those two parts, the likelihood (influenced by your data) can easily overwhelm your priors (Fig. 2). Indeed, most work on the dangers of priors and ‘prior misspecification’ focuses on cases where you have very little data for the model you’re trying to

fit. Priors, however, can only matter more than you know when you fail to think through and check them (that is, you skip Steps 1-2).

How this workflow changed our science

As we have used this workflow, how we approach our statistical models has changed. These changes have generally been similar for each of us. We suspect they are not unique to us, our study systems or our questions. Instead, we think they represent common approaches to statistical modeling that could help the field advance, much as we believe they have helped our science advance.

Understanding nonidentifiability

Identifiability refers to when all parameters in a model can be uniquely identified with infinite data; more common in our experience is nonidentifiability. Models can be nonidentifiable in several ways, including when mathematically some parameters cannot be uniquely defined. Statistically, a kin of nonidentifiability—degeneracy—is often an outcome of the empirical data combined with the model. Degeneracy occurs when the data do not contain enough information to estimate one or more parameters uniquely (Gelman & Hill, 2009).

Nonidentifiability and degeneracy can come up in many ways in ecology—and be hard to see, especially if you rush through model fitting. But if you have to write out your model and simulate data, you may suddenly realize lots of places for nonidentifiability and degeneracies to live. For example, when species do not occur across most sites, a model including separate parameters for site and species is often degenerate, but there's no warning in packages to tell you this. We have become far better at noticing nonidentifiability and degeneracies based on this workflow—and we have adjusted how we collect data and interpret results because of it.

Know your limits

~~Once we noticed how pervasive nonidentifiability and degeneracies were we started simplifying our models. Whereas previously when we had data that qualitatively appeared complexly nested, crossed, split or twisted, we would have initially tried to fit all of these intricacies (on the intercept), we are now more slow to add these to our models. By both understanding these terms better (including the many different ways each can be modeled), and understanding them better depending on the data and model in each unique context, we now work more carefully through what to include.~~

Before using this workflow, some of us would start with complex models, then simplify them until their fits converged in our given software package. Often these were hierarchical models with many levels—for example, including every column of site, plot, transect and quadrat in our dataframe, without stopping to check how well sampled they were, or what degeneracies they might introduce. We also often fit a suite of interactions: multiple two-way interactions and the occasional three-way interaction were common fare. But in simulating data, and fitting models to real, messy, imbalanced data using the workflow we came to see how much we were asking of our data and models together. Fitting a two-way interaction with half the effect size of a main effect takes a 16X sample size, compared to fitting the main effects alone (the main effects then

average over the interactions, see Gelman & Hill, 2020, for more details). This is sobering. It's more sobering when you see it played out again and again through this workflow.

We now both add complexity and simplify based on a more careful reckoning. Usually our starting model is not simple, it often includes grouping factors that may be difficult to fit, but that we see as absolutely critical to the question, model and data at hand. We still add and consider additional grouping factors and interactions, but we do so with a careful idea of how stable the model given the data likely is with them, and we rarely fit complex three-way interactions or similar—unless we have carefully designed the model and data collection for that aim.

Looking at parameters, not p-values

Before this workflow, not all of us commonly discussed the values that parameters in our model took—things like the slope and intercept (two common model parameters) were sometimes reported, but we did not know them as well as we knew whether the p -value for the slope was < 0.05 . This changes quickly when you need to build simulated data, for example, for a phenological event (an observation of a biological event on a day within the calendar year so, ideally between 1 and 365, or 366 in a leap year) and suddenly find it's >1000 based on the parameters you put into your code. This focus on the value of parameters scales up through this and other modeling workflows. Having a better sense of parameter values across different biological contexts, model parameterizations, and time periods gives a better sense of how the biological world works, including what's reasonable, possible or wildly unrealistic.

How this workflow intersects with ecological training

This four-step workflow is a simplified version of the current best practices for Bayesian model fitting (Betancourt, 2020; van de Schoot *et al.*, 2021), but many of the skills required are not part of traditional ecological training. Writing out the math behind most statistical models enough to complete Steps 1-2 bleeds into the skillset usually reserved for those working on theory, where coding and simulating from a model are common tasks. In contrast field, lab and otherwise empirical-data based ecologists often fit models they could not simulate data from. This dichotomy seems short-sighted in our current era of bigger, messier data and greater computational methods poised to handle such messy data. Further, the increasingly computational toolkit of the modern ecologist makes it easier bridge the gap between ecological models and their underlying math.

A reasonably competent coder could easily simulate data under a complex model that they might not have the mathematical expertise to solve analytically—if doing so was part of their training and the workflows they regularly use. ~~While simulation methods may appear foreign initially, they are usually much easier to implement than the analytical derivations of traditional methods so often seen in textbooks and classes.~~ Simulating data rapidly clarifies underlying assumptions. While training in frequentist methods often includes memorizing assumptions for a particular test, or steps specifically designed to test assumptions (e.g., normal quantile plots), this workflow requires no such training. Instead it requires only the skills to identify whatever the assumptions have been encoded in their models.

~~This workflow thus requires a shift in how we approach Bayesian methods—and training in~~

~~Bayesian models in ecology. Already, uptake of new Bayesian R packages highlight that Bayesian methods are no longer the purview of only a few, and these changes come alongside advances in Bayesian workflows, algorithms, and visualizations (e.g. Betancourt, 2020; van de Schoot *et al.*, 2021; Gabry *et al.*, 2019), that ecology must adapt its training for.~~ While this is an active area, we highlight three major changes.

(1) Prior ‘beliefs’ are changing. Best practices for determining priors is an active area of statistical research (Gelman *et al.*, 2014; Gelman & Hill, 2020; Betancourt, 2021a), and training should reflect current best practices. These include that ‘non-informative priors’ are a misnomer, as they are often informative (Lemoine, 2019), and priors can easily be ‘weakly informative.’ Thus a strong focus on the dangers of priors in training can be overkill, and verges on scaremongering.

~~Current training often includes a very strong focus on mathematically convenient priors, because of the importance of conjugate priors in closed form solutions for particular posteriors. Modern algorithms, such as HMC, do not require conjugate priors.~~ Prior predictive checks provide a far more powerful way to understand how priors work within a particular model, and are more useful than rules about which priors should be fit in certain cases or memorizing which priors are conjugate (Betancourt, 2021a).

(2) ‘Random effects’ are not just random. Hierarchical models contain grouping factors, sometimes referred to as ‘random effects,’ such as species or individual. **This term**, however, is misleading, imprecise and thus no longer recommended (Gelman & Hill, 2009). In ecology, it also carries with it many older ‘rules’ of what is ‘random’ versus ‘fixed,’ including that ‘random effects are things you don’t care about’ (for example the ‘block’ effect from a randomized block design). After posterior retrodictive checks (Step 4), you might feel differently, as hierarchical effects are (by definition) drawn from an underlying distribution—meaning you can predict outside of the specific set you sampled (for example, you can predict for a new species or individual), whereas you cannot do so for most categorical ‘fixed’ effects.

(3) P-values, and null hypothesis testing in general, are easily misleading, and there’s no easy fix for that. The replication crisis, rampant in other fields, is based in part on an overly hopeful belief that p -values will separate the signal from the noise, with one easy number. In reality, small sample sizes, lack of routine reporting of interpretable effect sizes, fitting of many models without adequate explanation, poor data and code reporting habits all increase the chance of finding ‘significance’ at a level of ≤ 0.05 (Halsey *et al.*, 2015; Loken & Gelman, 2017). This reality means a similar crisis is likely lurking in ecology, especially given small sample sizes alongside a tendency to fit complicated models with multiple interactions. The answer to this, however, is not to make p -values smaller (Halsey *et al.*, 2015; Colquhoun, 2017), nor is it Bayesian approaches.

The answer is training in workflows designed for careful model building, model fitting and model interrogation informed by ecological theory and understanding—including the one we outline here. ~~This workflow depends strongly on simulating data for testing your model (Step 2), and understanding your model results (Step 4) — an area we actively under train in ecology. It makes clear how relevant and important simulation is, but the relevance of simulation extends well beyond Bayesian model fitting. Simulation approaches allow interactive learning, build intuition, and stress exploring a model in its relevant ecological context. Ecologists are much better at thinking about ecological problems than statistical ones, and grounding our approach in ecology will likely bring the best out of our statistical modelling.~~

Acknowledgements: Comments from F. Baumgarten, D. Loughnan and N. Pates improved this manuscript.

1 References

- Beaumont, M.A. (2019) Approximate Bayesian computation. *Annual review of statistics and its application* **6**, 379–403.
- Betancourt, M. (2019) The Convergence of Markov Chain Monte Carlo Methods: From the Metropolis Method to Hamiltonian Monte Carlo. *Annalen der physik* **531**.
- Betancourt, M. (2020) Towards A Principled Bayesian Workflow. https://betanalpha.github.io/assets/case_studies/principled_bayesian_workflow.html.
- Betancourt, M. (2021a) Prior modeling. https://betanalpha.github.io/assets/case_studies/prior_modeling.html.
- Betancourt, M. (2021b) (what’s the probabilistic story) modeling glory? https://betanalpha.github.io/assets/case_studies/generative_modeling.html.
- Box, G.E. (1976) Science and statistics. *Journal of the American Statistical Association* pp. 791–799.
- Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M.A., Guo, J., Li, P. & Allen, R. (2017) Stan: A probabilistic programming language. *Journal of Statistical Software* **76**, 10.18637/jss.v076.i01.
- Colquhoun, D. (2017) The reproducibility of research and the misinterpretation of p-values. *Royal Society Open Science* **4**.
- Conn, P.B., Johnson, D.S., Williams, P.J., Melin, S.R. & Hooten, M.B. (2018) A guide to Bayesian model checking for ecologists. *Ecological Monographs* **88**, 526–542.
- Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. (2019) Visualization in bayesian workflow. *Journal of the Royal Statistical Society Series a-Statistics in Society* **182**, 389–402.
- Gelman, A., Carlin, J.B., Stern, H.S., Dunson, D.B., Vehtari, A. & Rubin, D.B. (2014) *Bayesian Data Analysis*. CRC Press, New York, 3rd edn.
- Gelman, A., Goegebeur, Y., Tuerlinckx, F. & Van Mechelen, I. (2000) Diagnostic checks for discrete data regression models using posterior predictive simulations. *Journal of the Royal Statistical Society Series C-Applied Statistics* **49**, 247–268.
- Gelman, A. & Hill, J. (2009) *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge, New York.
- Gelman, A. & Hill, J. (2020) *Regression and Other Stories*. Cambridge University Press.
- Gelman, A., Vehtari, A., Simpson, D., Margossian, C.C., Carpenter, B., Yao, Y., Kennedy, L., Gabry, J., Bürkner, P.C. & Modrák, M. (2020) Bayesian workflow. arXiv.
- Grinsztajn, L., Semenova, E., Margossian, C.C. & Riou, J. (2021) Bayesian workflow for disease transmission modeling in stan. *Statistics in Medicine* **40**, 6209–6234.
- Halsey, L.G., Curran-Everett, D., Vowler, S.L. & Drummond, G.B. (2015) The fickle p value generates irreproducible results. *Nature Methods* **12**, 179–185.

-
- Hampton, S.E., Strasser, C.A., Tewksbury, J.J., Gram, W.K., Budden, A.E., Batcheller, A.L., Duke, C.S. & Porter, J.H. (2013) Big data and the future of ecology. *Frontiers in Ecology and the Environment* **11**, 156–162.
- Held, L., Schrodde, B. & Rue, H. (2010) Posterior and Cross-validatory Predictive Checks: A Comparison of MCMC and INLA. *Statistical Modelling and Regression Structures: Festschrift in Honour of Ludwig Fahrmeir* (eds. T. Kneib & G. Tutz), pp. 91–110.
- Hoffman, M.D. & Gelman, A. (2014) The No-U-Turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research* **15**, 1593–1623.
- Kokko, H. (2005) Useful ways of being wrong. *Journal of evolutionary biology* **18**, 1155–1157.
- Lemoine, N.P. (2019) Moving beyond noninformative priors: why and how to choose weakly informative priors in bayesian analyses. *Oikos* **128**, 912–928.
- Loken, E. & Gelman, A. (2017) Measurement error and the replication crisis. *Science* **355**, 584–585.
- MacElreath, R. (2016) *Statistical Rethinking*, vol. 469 pp. CRC Press, New York.
- Muff, S., Nilsen, E.B., O’Hara, R.B. & Nater, C.R. (2022) Rewriting results sections in the language of evidence. *Trends in ecology & evolution* **37**, 203–210.
- Muthukumarana, S., Schwarz, C.J. & Swartz, T.B. (2008) Bayesian analysis of mark-recapture data with travel time-dependent survival probabilities. *Canadian Journal of Statistics* **36**, 5–21.
- Schad, D.J., Betancourt, M. & Vasisht, S. (2021) Toward a principled Bayesian workflow in cognitive science. *Psychological Methods* **26**, 103–126.
- Servedio, M.R., Brandvain, Y., Dhole, S., Fitzpatrick, C.L., Goldberg, E.E., Stern, C.A., Van Cleve, J. & Yeh, D.J. (2014) Not just a theory—the utility of mathematical models in evolutionary biology. *PLoS biology* **12**, e1002017.
- Strinella, E., Scridel, D., Brambilla, M., Schano, C. & Korner-Nievergelt, F. (2020) Potential sex-dependent effects of weather on apparent survival of a high-elevation specialist. *Scientific Reports* **10**, 8386.
- Sunnåker, M., Busetto, A.G., Numminen, E., Corander, J., Foll, M. & Dessimoz, C. (2013) Approximate bayesian computation. *PLoS computational biology* **9**, e1002803.
- Trijoulet, V., Holmes, S.J. & Cook, R.M. (2018) Grey seal predation mortality on three depleted stocks in the West of Scotland: What are the implications for stock assessments? *Canadian Journal of Fisheries and Aquatic Sciences* **75**, 723–732.
- van de Schoot, R., Depaoli, S., King, R., Kramer, B., Maertens, K., Tadesse, M.C., Vannucci, M., Gelman, A., Veen, D., Willemsen, J. & Yau, C. (2021) Bayesian statistics and modelling. *Nature Reviews Methods Primers* **1**.
- Wesner, J.S. & Pomeranz, J.P.F. (2021) Choosing priors in Bayesian ecological models by simulating from the prior predictive distribution. *Ecosphere* **12**.

-
- Winter, S.D.D. & Depaoli, S. (2023) Illustrating the value of prior predictive checking for bayesian structural equation modeling. *Structural Equation Modeling-a multidisciplinary journal* .
- Zheng, C., Ovaskainen, O., Saastamoinen, M. & Hanski, I. (2007) Age-dependent survival analyzed with Bayesian models of mark-recapture data. *Ecology* **88**, 1970–1976.

2 Tables & Figures

Table 1: A set of major terms used with their simplified definitions.

<i>Term</i>	<i>Definition</i>
calibration	analyzing how often an estimate is close to the true value over an ensemble of hypothetical observations. An exact calibration would requires knowing the true data generating process which is impossible in practice. We can, however, calibrate to data simulated from the configurations of models we plan use to fit to our data (<i>Steps 1-2</i>) so we understand the models better, including their limits given data similar to ours. We emphasize simulations to calibrate model behaviors consistent with our ecological systems and understanding (e.g., working within a limited set of parameter ranges through prior predictive checks). In contrast to this approach, frequentist method are calibrated against all possible behaviors, which is not only impractical in most circumstances it's also irrelevant given that the most extreme behaviors are unlikely to manifest in reality.
degeneracy	complex uncertainties that come from a mix of sources, including, non-identified models and cases where the data cannot well inform model parameters. When the data are not informing the parameters that we care about, this highlights a measurement issue. Identifying these problems in simulation studies can highlight when we need a better experimental design (e.g., sampling for more overlapping species across sites, or changing what we measure, etc.).
non-identifiability	when all parameters in a model cannot be uniquely identified with infinite data
prior	an distribution of reasonable values for a parameter based on fundamental biological and ecological understanding, previous research, or other sources
statistical model	Mathematical approximations of the true data generating process labeled with numerical parameters. Evaluating a statistical model on observed data gives a likelihood function that quantifies how compatible different parameters are with the observed data, and hence can be used to 'fit' the best parameters. In this article, we often simplify to 'model.' See also the Supplement: What's a model?
posterior	product of the likelihood and prior; that is, a probability distribution that quantifies how compatible different model parameters are with both the observed data and the domain expertise encoded in the prior model.
workflow	a set of steps to achieve a goal, with those steps designed to help organize the process, and ideally make it more systematic




Figure 1: The four-step workflow we outline can help design models for specific ecological questions, data and aims—which makes this a statistical workflow that can naturally become a scientific workflow. It makes the step that many of us focus on—running your model on your empirical data (Step 3)—far more straightforward and insightful by using simulations both before (Step 2) and after (Step 3) it to better understand the model and data together.




Figure 2: A simple example of how to use simulated data to understand calibration issues in a mis-specified model. Here we know the true model underlying the data is $y = \alpha + \text{normal}(0, \sigma)$ where α is 5 (shown as blue vertical line) and σ is 2. The model, however, is mis-specified by a prior for α of $\text{normal}(25, 2)$ (dashed blue line), resulting in a posterior (salmon-colored histogram) not centered on the true value. In our experience it is quite rare to have a prior informed by ecological knowledge be so far off, but this is an example. How mis-calibrated the model will be depends on the data: we show examples with a sample size (N) of 5, 10 and 100 data points. (Note the change in y axis range for bottom plot.)




Figure 3: Example of a single retrodictive check from time-series data of phenological events over time. The raw data (top) looks similar to one simulated dataset (bottom), based on existing species number, their respective x data, and simulating from the parameters for each species. See ‘An example workflow’ in the Supplement for more details.