

Running title: Environmental tracking

How environmental tracking shapes species and communities in stationary and non-stationary systems

E. M. Wolkovich¹ & M. J. Donahue²

¹ *Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4 (e.wolkovich@ubc.ca)*

² *Hawaii Institute of Marine Biology, University of Hawai'i at Mānoa, Kānāeohe, HI 96744 (donahuem@hawaii.edu)*

Corresponding author: see¹ above; Ph: 604.827.5246 (no fax).

Authorship statement: EMW and MJD both conceived of the paper, performed modeling work and edited the paper, EMW additionally wrote the paper and did the literature review, while MJD additionally wrote the supplementary information on the model.

Data statement: Review, so no new primary data, but data from a comprehensive literature review will be archived in an appropriate public repository and the data DOI will be included at the end of the article.

Keywords: community assembly, global change, climate change, phenology, environmental variability

Article type: Reviews and Syntheses

Article information: Abstract: XXX words; Main text: 6,490; Figures: 5; Boxes: 4 (text in Box 1: 348; Box 2: 500; Box 3: 264, Box 4: 776); 113 references

Abstract

Climate change is reshaping the environments of all species. Predicting responses requires understanding the costs, benefits and constraints of how well species can track environmental change, and how such tracking shapes communities. Growing empirical evidence suggests that environmental tracking—how an organism shifts the timing of key life history events in response to the environment—is linked to species performance and is a structuring force of species and communities today. Here, we review current knowledge on tracking both in empirical data and through the lens of ecological theory. We provide a definition of environmental tracking that highlights both why it must be fundamentally related to fitness, and the challenges of defining it empirically. We then show how life history theory makes predictions for variation in tracking, and trade-offs with other traits, across species and environments. Finally, we examine how well basic community ecology theory can be extended to test the current paradigm that climate change should favor species with environmental tracking. We to aim provide a framework based on existing ecological theory to understand how tracking in stationary and non-stationary systems may shape species and communities and, thus, help predict the species- and community-level consequences of climate change.

1 Main text

Anthropogenic climate change is causing widespread changes in species distributions, with many species shifting in both time and space (?). Reports often focus on species shifting to higher elevations and poleward (?) and/or shifting their recurring life history events (phenology) earlier as climate warms (??). These general trends, however, hide high variability across species. A large proportion of species are not shifting at all (?), which has raised concerns about whether these species may be more vulnerable to population declines with continued warming. Such concerns come in part from increasing research that links how well species track climate change—especially through temporal shifts—to shifts in biomass, growth and other metrics related to performance (?). Tracking climate change may then be a major component to understanding and predicting the indirect effects of climate change, including population declines, with cascading effects on community and ecosystem structure.

The hypothesis that tracking predicts fitness outcomes with climate change has gained significant traction in the ecological literature focused on global change (e.g., ?) and several areas of ecological theory support it. Considering tracking as a form of plasticity, evolutionary models predict species that track will be favored in novel environmental conditions. Similarly, some models of community assembly suggest that a warming climate should open up new temporal niche space and favor species that can exploit that space (???). Yet not all studies find the purported link (e.g., ?), and there has been comparatively little work to improve predictions by formally connecting tracking to foundational ecological theory.

This disconnect could be because most ecological theory today is for stationary systems (e.g., ??). While major arenas of research such as ‘modern coexistence theory’ or population ecology are now built on assumptions of a stochastic environment, they generally still assume stationarity, where the underlying distribution of the environment is unchanged across time (i.e., constant mean and variance, ?). This assumption is common to much of the theory that underlies ecology, evolution, and myriad other research fields (e.g., ??).

Climate change upends the assumption of stationarity. By causing increases in temperature, larger pulses of precipitation, increased drought, and more storms (?), climate change has fundamentally shifted major attributes of the environment from stationary to non-stationary regimes (see Fig. 1 and Box: Environmental variability & change). This transition is reshaping ecological systems, and, while new work has aimed to adapt coexistence theory to non-stationary environments (??), there is still very little theoretical work on what such a transition may mean for communities and the species within them, despite growing empirical studies.

Here, we review current knowledge on tracking both in empirical data from the climate change impacts literature and through its related ecological theory. We provide a definition of environmental tracking that highlights why it must be fundamentally related to fitness and the complexity of defining it in empirical systems. We show how life history theory—specifically drawing on optimal control, bet-hedging and plasticity—make predictions for variation in tracking across species and environments in stationary and non-stationary systems. We then examine how well basic community ecology theory can be extended to test the current paradigm that climate change should favor species with environmental tracking.

1.1 Defining environmental tracking

While tracking is a commonly used word in the phenology and climate change literature (e.g., ???), there are few, if any, definitions of it. Most interpretations of tracking relate to how well an organism matches the timing of a life history event to the ideal timing for that event, what we refer to as ‘fundamental tracking’. Fundamental tracking thus rests on an assumption that there is a timing (an ‘ideal timing’) that yields maximum fitness, and event timings moving away from this ideal result in reduced fitness (a foundational concept of the trophic mismatch literature, ?). Yet this ideal timing is generally only clear in simplified models or in retrospect, thus species must use environmental cues to attempt to predict and match their event timing to the ideal timing across environments in both space and time. Each organism’s set of cues forms the biological basis for how a species tracks, but measuring environmental tracking requires two more components.

The first component is the environment’s variability—which aspects of the environment vary, how (e.g., temporally each year, spatially at x scale) and how much. If the varying components of the environment do not correspond closely to the organism’s cues, then the species may not track this variability. Further, under this definition, identical genotypes will have different tracking across environments, depending on the interaction of the cues and environmental variability.

Second, which aspect(s) of the environment researchers measure will determine measured environmental tracking. If researchers know the exact cue (e.g., a thermal threshold) or suite of cues (e.g., a interaction of thermal sums and daylength) and can perfectly measure these in an environment where the cue(s) varies then an organism will track the environment perfectly. If researchers measure some related attribute (e.g., mean spring temperature in place of thermal sums) or only some of the organism’s cues, then the organism will appear to track poorly (i.e., a noisier statistical relationship). If researchers measure an environmental variable that is not directly related to the cue(s) that the species actually uses, but one correlated with it (e.g., an insect tracks daylength but researchers measure temperature) then they have not measured tracking per our definition.

Accurately measuring environmental tracking thus requires a complete knowledge of an organism’s cue(s), the environment’s variability and the relationship between the actual cues and measured environmental metrics. Knowing an organism’s cues is inherently difficult, generally requiring a suite of experiments, process-based models and in-situ data to show that the model of cues is accurate. Not surprisingly then we lack this for almost all species, coming closest for some model species (e.g., *Arabidopsis thaliana*, ??), or species with very simple cues (e.g., coral *Acropora millepora*, ?) and have some basic information for some other species (e.g., the Great Tit, *Parus major*, ?).

1.2 Measuring environmental tracking

Attempting to measure environmental tracking and compare variation in it across species, space and time is a rapidly growing area of ecological research (e.g., ???). Multiple meta-

analyses now show plants' spring phenology shifts with spring or annual temperatures 4-6 days/°C on average across species (???), but also highlight high variation across species (?), even after examining multiple major climate variables (?). Variability across species appears similar when examining consumers tracking their prey (across diverse species tracking over time is 6.1 days/decade but ranges from zero to 15 days/decade, see ?).

All species-rich studies of phenology-climate relationships find high variation, including some species that do not track or track poorly (i.e., high noise surrounding observed statistical relationships). Researchers have worked to link such variation at times to the underlying cues (e.g., ?), species traits (e.g., ?) or trophic level (e.g., ?). These approaches hint at the three major classes of reasons that underlie species that do not appear to track climate (or appear to be poor trackers): (1) species do not track, as perfect environmental tracking may either not be possible or optimal for all species, (2) lack of firm biological understanding of the cues that underlie tracking, and (3) statistical artifacts that make it difficult to measure tracking robustly.

Increasingly, research has outlined statistical difficulties in measuring tracking. These issues mostly relate to the challenges of observing phenological distributions (CITES) through anthropogenically defined temporal units (e.g., day or month) in non-stationary systems, and to the complexity of climate data (more further discussion, see Box 'Statistical challenges in measuring tracking').

Even without statistical issues, translating phenological and climate data into estimates of tracking requires a firm biological understanding of an organism's cues, critical knowledge that researchers rarely have (?). Currently, 'tracking' is often measured simply as the relationship between the dates of the phenological event and a simple abiotic metric, such as mean monthly temperature (with variation in temperature derived from multiple periods of observation or induced through experiments). Simple environmental metrics, however, are almost always proxies for a more complicated underlying physiology where simple cues—such as warm temperatures—can be modified by other cues, such as photoperiod, drought or light spectra (??). Indeed, multiple studies have shown how simple correlations between phenological events and environmental variables can mask complicated relationships (??).

Modeling multivariate cues well is inherently difficult (?), especially since one cue may dominate in many conditions. For example, woody plant leafout responds strongly to warm spring temperatures, but also to cool winter temperatures to prevent leafout in mid-winter warm snaps that occur long before the last frost. Often this cool-temperature effect may be masked by sufficiently cold conditions. With warming from climate change, however, this additional trigger—which appears to vary by site, species and even inter-annual conditions (?)—may become critical (and potentially lead many phenological models to fail spectacularly in the future as additional cues come into play, see ?). In some semi-arid systems, species time growth to pulses of rain, but only when those rain events occur with cooler temperatures that indicate the start of the rainy season, and not a rare summer rainfall event in the middle of months of drought (??). Tracking in species with longer generation times may be especially complicated, as species may track low frequency climate signals and make investment choices on far longer timescales than species with shorter lifespans (?).

139 Researchers are increasingly recognizing the need to consider multiple climate variables, though
140 currently most estimates are based on long-term observational data (e.g., ??), which can lead to
141 spurious correlations without experiments to test hypothesized cues (?). Further, estimates of
142 ‘tracking’ from long-term data that are not linked to mechanistic experiments may sometimes
143 serve as proxies (i.e., environmental variables correlated with one or more actual cue that a
144 species uses) for an organism’s environmental tracking, but may not directly connect to an
145 organism’s cue(s).

146 Limited understanding of organisms’ phenological cues combined with statistical issues may
147 make many current estimates of variation in tracking less reliable than they appear, and make
148 robust quantitative analyses across species difficult (??). Yet these estimates provide the crucial
149 first step to understand variation. As estimates improve, ecologists will better capture a picture
150 of which species, when and where, do and do not track. Yet given the current difficulty of
151 measuring environmental tracking, clear testable predictions from ecological theory are perhaps
152 most critical to guide research today (?).

153 1.3 Understanding variation in environmental tracking

154 A number of research areas in ecology predict variation across species in how well they track the
155 environment. Applying these areas of research to environmental tracking, however, first requires
156 understanding phenological events. In particular, while phenological events are often coded as
157 on/off switches (e.g., a seed does/does not germinate; a coral does/does not spawn), they are
158 almost always defined by investment decisions that are part of a continuous developmental
159 process (?).

160 Phenological events are best considered as the outcome of a two-part sequential process that is
161 repeatedly observed over time. At each temporal unit, an event can either happen or not (step
162 1) and, if it happens, there is a secondary part regarding the size or degree of investment of
163 the event (step 2). This process is generally applied at the level of the individual (but it could
164 potentially apply at lower levels, for example buds on a branch, or potentially higher levels,
165 such as all the offspring from a parent). Across time, it produces an event’s distribution. After
166 starting, many events are entrained to continue based on the underlying physiological process:
167 for example, laying eggs within one clutch (here, the first part of the process is whether to
168 lay eggs or not and the second is whether to continue to invest in that process, which would
169 lead to additional eggs, which researchers then observe as number of eggs per temporal unit) or
170 flowering each growing season. In such cases, first events at the individual-level are somewhat
171 unique from the rest of the event’s distribution. In all cases, these individual-distributions scale
172 up to the population-level estimates of these events generally used by researchers (see ?, for
173 discussion of the outcomes of this scaling).

174 Considering the life history events that define part of environmental tracking as a two-part pro-
175 cess highlights that tracking is ultimately shaped by resources that species need to grow and
176 reproduce, and circles back to an organism’s fundamental tracking. This is perhaps best recog-
177 nized in the literature on trophic synchrony where there is often focus on how well consumers’
178 environmental tracking matches to the seasonal distributions of their prey (??). For example,

decades of work has studied how birds (e.g., *Parus major*) time their peak food demands—during their nesting season—to maximum prey (caterpillar) abundance (e.g., ?). Failure of environmental tracking to match prey year-to-year or over time with long-term warming has been well tied to individual-level fitness consequences in some systems (?), but not all (?), which may be due to the complexity of mechanisms that influence total fitness (??). Environmental tracking in plants and other lower trophic levels is also about resources. Alpine plant species that emerge in step with snowmelt or temperature are likely responding, at least in part, to light resources for photosynthesis. Light equally appears critical to the sequence of phenology in many temperate forests: with lower-canopy species, and younger (shorter) individuals of higher-canopy species, routinely risking frost damage to leaf out before the canopy closes and access to light becomes severely reduced (??). These ultimate controllers on tracking—which determine fundamental tracking—are then filtered through the abiotic environmental cues species use to time events. From here, predicting tracking relates to predicting which cues an organism should use: an optimal control problem.

Predicting variation in environmental tracking in stationary systems

An optimal control framing can help predict which cues an organism should have based on a consideration of the costs, benefits and constraints, in any one organism by environment system (?). First, it requires that benefits vary depending on the timing of event; this effect may be stronger in highly seasonal environments. Next, there must be a useful cue—some aspect of the environment that predicts resources or otherwise links back to the ultimate factors that shape environmental tracking (?). Some environments may inherently lack useful predictors, such as desert systems where few early-season variables seem to predict high or consistent rainfall years.

From here, the exact cue or suite of cue(s) that an organism should have depends on the cost of those cues (e.g., the machinery of monitoring temperature or daylength), the benefits of a cue (for example, how much tissue is saved by avoiding a coldsnap). Ultimately the balance of the costs of cue(s) and their benefits should determine exactly what cue(s) a species uses: apparently poor cues may occur for organisms in environments where there is both a low cost and low benefit to the cue(s). Similarly, expensive cues, such as complex multivariate ones, are possible given a high pay-off. Most in-depth studies of species' phenological cues find evidence for complex multivariate ones (?). These cues almost always appear adapted to handle unusual—though not completely uncommon—years when the simple cue alone would fail (that is, would trigger growth, reproduction or another life history event at a suboptimal time), suggesting that multivariate cues may provide a large benefit in best coupling the timing of cues to the ultimate controls (i.e., cues that couple environmental tracking strongly to fundamental tracking).

Optimal control highlights that not all species should track, but instead that tracking is based on an optimization of costs, benefits and constraints. In environments where there is no clear optimal strategy, species should bet-hedge (?). In general, species in highly variable environments, or which otherwise face high uncertainty in when to time investment decisions, should gain a substantial benefit from bet-hedging or employing other approaches that spread out risk given uncertainty (??). Assessing bet-hedging in many systems, however, requires studies of fitness over longer timescales than many current field experiments.

Constraints also shape cues and may limit tracking. Fundamental differences in life history impose constraints—for example, the type and amount of loss an organism can sustain each season is limited by its generation time and other attributes related to long-lived lifestages that yield buffered population growth (?). Additionally, constraints may arise if a species cannot closely measure relevant environmental cues (??), through unavoidable trade-offs with tracking (??), or through evolutionary pathways. Gene flow from other environments may continually push a population away from its local optimum (?), standing genetic variation limits phenotypic variation and thus can slow the evolution of optimal cues (??), deeper evolutionary history may produce co-evolved traits making it difficult for selection to act on a single trait axis (?), or other fundamental evolutionary limits to the rates of trait change and what traits are possible (?).

Predicting variation in environmental tracking in non-stationary systems

Much of life-history theory, including optimal control and bet-hedging, rests on assumptions of stationarity, thus a major open area of research is adapting life history theory to non-stationary environments. Multivariate cues may be especially robust to a non-stationary environment if they provide a tight coupling of cues to fundamental tracking, and that coupling is maintained in the non-stationary environment (?). But multivariate cues may equally be most vulnerable to failure if non-stationarity decouples the cues from fundamental tracking (?). For example, consider an organism’s whose cues evolved based on a correlation between peak prey abundance and daylength—these cues may work well in a stationary environment but fail if warming advances peak prey abundance. Predicting the outcome of non-stationarity thus relies on knowing both the full cue system of an organism, how it relates to fundamental tracking, and how both that cue system and the underlying fundamental model shift with non-stationarity.

Another area of life-history theory, that focused on plasticity, may be primed to provide insights on non-stationarity (or ‘sustained environmental change,’ see ?). Considering phenology as a trait (as we and others do, e.g., ???), environmental tracking is one type of plasticity. Researchers could thus more broadly understand environmental tracking through modeling an organism’s reaction norms (??) and understanding how cues and suites of cues—across environments—determine how fundamentally plastic an organism may be in its tracking. For example, multivariate cues should yield higher plasticity in this framework. From here, models of the role of plasticity in novel environments provide an important bridge to understanding the outcomes of non-stationarity, generally predicting non-stationarity should favor highly plastic species. This outcome, however, assumes there are no costs related to plasticity (??). If there are costs associated with plasticity, akin here directly to costs associated with tracking, then species may evolve lower tracking, because it should trade-off with other traits (?).

1.4 Tracking in multi-species environments

Plasticity theory—in contrast to much of the life-history theory discussed above (where other species are, at best, filtered into models as an aspect of the environment)—shows how critical a multi-species perspective is to understanding environmental tracking (?). In this light, tracking cannot be considered as a singular trait, but must be evaluated as part of a trait syndrome (or

mosaic of traits, ?) and selection in multi-species environments should produce communities of species where tracking trades-off with other traits.

As tracking often relates to the timing of a resource pulse, traits related to resource acquisition are likely contenders for a trade-off. Species with traits that make them poor resource competitors may need to track the environment closely to take advantage of transient periods of available resources, but will risk tissue loss to harsh environmental conditions more prevalent early in the season (e.g., frost or snow). In contrast, species with traits that make them superior resource competitors may perform well even if they track environments less closely, because their resource acquisition is not strongly constrained by competitors. Examples include under-canopy species leafing out earlier to gain access to light (?) or species with shallow roots starting growth sooner in an alpine meadow system, while species with deeper roots begin growth later (?). In such cases, tracking is akin to a competition-colonization trade-off (?), where species that track well gain priority access to resources and, thus, may co-exist with superior competitors. Research to date supports this, with several studies linking higher tracking to traits associated with being poor competitors (???). Further, many studies have found a correlation between higher tracking and ‘earlyness’ each season, which has been linked to resource acquisition traits associated with lower competitive abilities (?, see Box ‘Trait trade-offs with tracking’).

Understanding these trade-offs is clearly critical, but we also argue more theory is needed to understand the short-term dynamics of a changing environment with plastic species. Most theory predicts the outcome of a new environment, but non-stationarity in the climate today means further understanding of the trajectory to that outcome is key for understanding current trends. For example, models show how plasticity may limit standing variation and thus reduce fitness in novel environments (??). Whether such findings extend to systems transitioning from stationary to non-stationary will likely depend on how non-stationarity affects the rate of adaptation (?). Efforts to model expected outcomes given climate projections and current understanding of plasticity and genetic variation underlying event timing in some organisms provide the empirical start to this (e.g., ?), but more eco-evolutionary models that bridge this gap may prove especially useful.

Including tracking in multi-species community assembly models

Predicting how tracking may determine which species are effectively winners and losers with climate change requires integrating non-stationary environments into models of plasticity and community assembly. Recent advances in coexistence models, sometimes called ‘modern coexistence theory,’ recognize that both mechanisms independent of fluctuations in the environment (e.g., R^* and other classical niche differences) and mechanisms dependent on fluctuations in the environment (relative non-linearity and storage effect) can lead to coexistence (??). These models, which underlie much of current community ecology research (???), provide a framework to begin to model environmental tracking and non-stationarity.

How the environment is defined in most community models falls into two broad categories. In some models the environment is expressed as variation in parameters related to species. For example, in some lottery models the environment appears, effectively, as variation in birth and death rates. Building a changing environment into such models thus requires knowing how

environmental shifts filter through to species-level parameters (?). For example, ? added the temporal environment to competition models by defining interaction strength as dependent on the temporal distance between species. In other models, the environment is more specifically defined. Many of these models define the environment as a resource (e.g., many seed germination models that begin with a resource pulse each year), and thus generally model something close to fundamental tracking. Building a changing environment into these models requires knowing how the environment is changing (see Box ‘Environmental variability & change’).

Models that explicitly include the environment provide a major opportunity to predict how environmental tracking and non-stationarity determine future communities (see Box: ‘Adding tracking and non-stationarity to a common coexistence model’). Yet most current models generally examine the environment from only one of two important angles for tracking: they either represent the environment as used for species’ cues (e.g., many models of plasticity) or they represent the environment as directly affecting fitness (e.g., the storage effect model). Combining these two angles may be especially critical to understanding the costs and benefits of tracking in non-stationary environments.

Layered onto the different angles that different models take on the environment is how species responses to the environment are defined. In general, species responses to the (resource) environment can be broadly grouped into models that explicitly define when species start an event (e.g., spawning or germination) in response to the environment versus those that model the magnitude (e.g., the number of propagules or seeds) of response to the environment. Models that explicitly model when a species starts an event are often focused on situations where order of arrival is critical to predicting coexistence outcomes. Models of priority effects through niche pre-emption highlight the advantage tracking may provide when it allows species to be early (and when there is no cost to being too early): early arrivals receive a head-start advantage, by gaining priority access to resources (the environment) they can draw down the resources available to later arrivals (?).

Other models canalize species’ responses to the environment into production and investment. For example most models of inter-annual competition (much of ‘modern coexistence theory’) fall into this camp. Species produce (via offspring, tissue etc.) differentially depending on the environment each year and outcomes are mediated through density. While these models superficially may seem disconnected from timing, they critically highlight how phenology relates to production and, thus, investment in future years (in contrast, priority effect models explicitly model timing).

A model where species vary both when they start an event and how much they produce dependent on the environment would capture the important attributes of tracking—combining head-start advantages from being early with production variation based on the fitness of the environment. To our knowledge, however, most models approach these questions separately, though models of bet-hedging come closest (??). A model that explicitly models the linked decisions of when to time an event and how much to offspring/tissue to produce in the event could provide fundamental insights on the relative importance of each aspect of this process. Such a model could be adapted to address multiple questions of environmental tracking, including how these decisions may trade-off and which other traits tracking may be most strongly linked

to, as well as explicitly modeling the costs and benefits of tracking in stationary systems, a critical precursor to extending it to non-stationary systems.

Extending models to non-stationary systems is crucial to fully testing how environmental tracking relates to species persistence with climate change, and research has already begun to tackle this non-trivial challenge (???). Most work to date, however, focuses on conclusions from stationary or non-stationary systems. Thus, the transition between stationary and non-stationary is often ignored, yet we expect it may be most critical. Communities formed in stationary environment periods (or periods with environments lower non-stationarity) are effectively filtered and assembled by that environmental regime and thus produce the base variation and assembly dynamics for a shifting environment. While analytical solutions for systems transitioning from stationary to non-stationary may take time to develop (?), simulation work can provide an immediate intuition and framework to address this challenge (see Box: Adding tracking and non-stationarity to a common coexistence model).

1.5 Future directions

Growing empirical research highlights that environmental tracking is linked to species performance and, thus, may be critical to understanding the forces that assemble communities and determine species persistence—especially as anthropogenic climate change is reshaping the environment of all species. Ecological theory, including from areas of optimal control, plasticity and community assembly, is clearly primed for understanding how a variable environment can shape the formation and persistence of species and communities. To understand what advances in theory may be most useful for making predictions in the Anthropocene we need more focus on understanding the attributes of an environment shaped strongly by humans. In turn, to test theory we need more robust estimates of environmental tracking and how it fits within a mosaic of other correlated traits. To this aim, we review several major areas of research that we believe could most rapidly unite empirical and theoretical research in environmental tracking to advance the field.

How is an organism's environment changing?

Currently, much research has focused on one major shift in the climate system (earlier growing seasons), but research on multivariate environmental shifts is growing and will be critical to understanding how climate change affects an organism's whole environment. Ecologists can guide these efforts by identifying environmental shifts that are often linked (e.g., warming temperatures may drive earlier seasons and higher evaporative loss of some resources such as water). Researchers can also aim to more consistently and fully characterize the environmental distributions of their systems that appear to most drive species performance and interactions: the environment of the years of study should be clearly reported and compared against long-term and recent climate for each system.

More interdisciplinary research with climate science could also speed a fuller understanding of what shifts are and are not expected with climate change, and what climate variables are inherently correlated. Such correlations make estimating cues and other biological parameters

from long-term data especially precarious (?). But these correlations are equally critical in considering how species may view their environment and whether environmental change will couple or uncouple links between proximate cues and fundamental tracking (?).

Understanding and measuring ‘tracking’

Understanding how the environment is changing represents just one step towards robust measures of environmental tracking. Shifting environmental regimes must then be filtered through species cues. As robust estimates of these cues are currently available for few if any species, we suggest several major improvements on current methods to help interpret current trends and make comparisons more feasible.

Studies should clarify their definition of tracking, how the environment is defined and how well, or not, the underlying cue system is understood for study species. Currently, many studies examine fundamental and environmental tracking at once (e.g., ?), which is clearly helpful in advancing the field. However, the more researchers can clarify when and how they are addressing fundamental tracking versus environmental tracking the more easily we can compare across studies. Next, and relatedly, studies should define their environment: are they considering primarily the abiotic environment or measuring an environment fundamentally shaped by other species? This difference connects to fundamental versus realized niches and whether systems are primarily top-down (resources and the environment may be strongly shaped by other species) or bottom-up controlled. Finally, all researchers working on environmental tracking need to embrace their inner-physiologist, or collaborate with one. For many species there is often a related species (albeit, sometimes distantly) whose cue system has been studied. Thus, researchers should draw on the literature of their study species’ close relatives to bracket which environmental variables may represent environmental tracking and which may be proxies, and to highlight uncertainty. We expect progress will come from a balance between measures of fundamental tracking, estimating an organism’s system of cues, and measuring environmental tracking. Clear statements of what is and is not known and measured will help.

Embrace the sometimes contradictory pulls of conducting experiments to identify mechanistic cues and the multivariate climate of the real world. Clearly, we need more experiments to identify which specific aspects of the environment different species cue to and how these cues are filtered by their actual environmental regime (as outlined above and see ?). Suites of experiments, which build from identifying cues to understanding how they act when correlated, are a major gap for most organisms.

Build a model of your species’ cues and interrogate it. As research progresses in trying to estimate environmental tracking, greater progress will come from fuller and more diverse interrogations of current (and future) models. Define the framework under which you expect your cue system developed or works (e.g., bet-hedging) then test how well it works, and where it fails (see ?, for an example). Model interrogation can also use realistic environmental regimes to provide field predictions (??) or predict future species and communities. One example of this comes from in silica resurrection experiments of model organisms where future environmental regimes included a mix of regular climate projections and projections modified to test and advance understanding of environmental tracking for the study species (e.g., warmer winter and

altered photoperiod scenarios in ?).

What major traits trade-off with tracking?

Basic theory of plasticity and competition suggest that environmental tracking must trade-off with other traits to allow multi-species communities. Yet to date empirical work has mainly documented tracking, linked it to performance, or focused on how it varies between native and non-native species (???). Such work lays the groundwork that environmental tracking is important, but future empirical research should address how this trait co-occurs with other traits. Research has highlighted some traits that co-vary with tracking (e.g., ???), but to tie this empirical work to models requires more research on traits that link clearly to theory, and a fuller understanding of how tracking and other traits jointly contribute to performance under varying environments.

Traits that link to resource competition, as we focus on here (as others have as well, see ?), may be especially fruitful for greater research, but should not be the only ones considered. For example, traits related to predator tolerance or avoidance may also play a role, but have been effectively unstudied. As empirical research in this area grows, models can aid progress in understanding the outcomes of these trade-offs for community assembly.

Developing new models for tracking in stationary to non-stationary systems

As outlined above many areas of ecological and evolutionary modeling contribute to our understanding of environmental tracking. But most are limited in various ways. Community ecology models generally bifurcate in modeling differences in timing versus production amounts across species, thus studies of whether these models lead to similar or different conclusions could help predict community outcomes and advance our understanding of trade-offs. As outlined above, understanding tracking likely requires models that combine effects. This includes models that combine effects of variation in timing and production amounts and models that include environment as impacting species' cues, as well as species' fitness. Such models would explicitly allow the potential costs and benefits of tracking depending on how closely environmental tracking matches fundamental tracking.

New models will also need to examine how relaxing assumptions of closed communities (i.e., without dispersal or evolution) alter predictions. In practice, dispersal of species or individuals with traits that make them better matched to the non-stationary environment would lead to new communities that may persist longer or be continually re-assembled as long as the environment remains non-stationary. Indeed, this logic underlies the argument that invasive species may be superior trackers benefiting from how climate change has altered growing seasons (??). Evolution equally could alter predictions by allowing species traits to evolve in step with environmental change. Long-term population (e.g., ?) and resurrection studies (??), as well as field experiments (??), have repeatedly shown species can shift to earlier flowering times, higher thermal tolerances or related genetically-controlled traits that confer higher fitness in warmer climates. Yet these studies also highlight that responses can be lagged (e.g., ?), associated with reduced population viability (e.g., ?), or other factors that may constrain adaptive responses

1.6 Stationarity in the future

While most environments today are climatically non-stationary and have been for decades, the climate will return to stationarity in the future. There are many possible pathways to climatic stabilization, but almost all require first the stabilization of greenhouse gases—the subject of much policy and political debate. Once greenhouse gas emissions stabilize climate will not quickly snap back to a new stationary phase. Instead systems will slowly approach a new climatic stationarity depending on how they are effected by the earth’s multiple thermal reservoirs, and, in turn, how quickly those reservoirs stabilize. The timescale of this approach is generally expected to be on the scale of centuries, but could be much longer in certain oceanic systems (?). Thus, ecologists are—and will remain for the foreseeable future—in a research area structured by climatic non-stationarity.

As paleobiologists and evolutionary biologists often point out, climatic nonstationarity is a common part of the earth’s history (even if stationary periods—be they cold or warm (glacial and interglacial periods)—are more common. Indeed, while much of this work has examined how species survive for millions of years given large oscillations in climate (?), the periods that provide the most dramatic community reshuffling are periods shifting from stationary to non-stationary climate regimes (??). Such stories of the past are now fundamentally happening today, and ecology is challenged to understand how transitions between stationary and non-stationary environments are reshaping the species and communities we have today and will in our warmer future.

2 Acknowledgments

We thank I. Breckheimer, D. Buonaiuto, E. Cleland, J. Davies, G. Legault and two anonymous reviewers for helpful comments that improved the manuscript and the NSERC Discovery Award (RGPIN-05038 to EMW) and Canada Research Chair in Temporal Ecology (EMW) programs that provided funding.

3 Boxes

3.1 Box: Environmental variability & change

Decades of ecological research highlight how temporally variable environments shape species and their communities at multiple scales (??). In seasonal landscapes, the environment limits periods for growth each year (e.g., by temperature or drought); within-year variability in the environment (e.g., daily, hourly or finer resolution temperatures or rainfall amounts) compounds into inter-annual variability that shapes the distribution of the start and end of growing seasons. For long stretches of history this variability has been effectively stationary; that is, the underlying probability distribution that describes the start (or end) of the season (e.g., the date of the last major frost) does not change, even though the date may be dramatically different from one year to the next.

In other time periods, variability has been non-stationary in one or multiple dimensions. For example, climate in the northern hemisphere includes long warming and then cooling periods (i.e., increasing then decreasing means of the probability distribution) at the start of the Holocene, when the earth was coming out of the last glacial maximum. Anthropogenic climate change is a similar non-stationary process, with warming evident around the globe and knock-on effects for other climate metrics, such as heat extremes and the size of precipitation events.

Understanding non-stationarity in ecological systems requires first identifying which aspects of the environment have shifted—and how they have shifted with respect to one another—as the underlying distributions transition from stationary to non-stationary (Fig. 1). For example, with climate change, warming has increased mean temperatures over time, with minimum temperatures generally increasing more than maximum—this results in an underlying distribution for daily temperature where the mean is increasing through time while the within-day variance is decreasing (??). Understanding the impacts of climate change further requires recognizing that many systems can be considered stationary or non-stationary depending on the timescale and period of study. Thus, predicting the consequences of current non-stationarity in ecological systems benefits from identifying the type and scale of non-stationarity, relative to long-term trends.

3.2 Box: Statistical challenges in measuring tracking

A potentially widespread reason for observations of species that do not track is statistical bias and artifacts, including non-stationarity in units and unrecognized low power. All of these can be addressed given improved statistical approaches (e.g., ??), though such approaches may (uncomfortably) highlight how uncertain many current estimates are (?). Non-stationarity in units comes in many forms—estimates of mean days shifted per decade (i.e., days/decade, a common metric summarizing observed shifts in phenology over time in long-term datasets) depend strongly on the climate of the decade(s) studied, which is not consistent in many systems (??). Estimates based on a relevant climate variable can sometimes ameliorate this problem, but may be equally vulnerable to non-stationarity in units (e.g., ?). For example, processes that depend on thermal sums reported as days/°C will generally appear to decline with warming, as

the thermal sum of an average day has increased in most regions with climate change. Relatedly, estimates of long-term change using simple linear regression are influenced by the climate at the start of the time-series (with greater changes seen from time-series that started in unusually cold decades, such as the 1950s for much of North America). Impacts of start-years for long-term time-series can be muted by applying change-point or hinge models (e.g., ?), while metrics that move away from calendar units (e.g., day) can help address non-stationarity in units.

Low power is widespread in ecology, where even ‘long’ time-series may be far too short for robust analyses (??). Authors should be especially cautious if they find only large effects appear significant (e.g., ?), which is a well-known statistical bias associated with p-values (?). Additionally, effect sizes that are higher when climate variability is higher (for example, in temperate habitats temperature is highly variable in the spring and autumn compared to summer) may be more related to variation in statistical power than to biology (periods with higher variation yield greater variation in the predictor variable, and thus higher power). Mixed models can help better leverage understanding by pooling information across species, and often better capture uncertainty (?). We suggest mixed models should be used more widely alongside randomization and/or data-simulation approaches (e.g., ??) to better estimate and communicate uncertainty in studies. And researchers should identify what results bias may produce. For example, growing evidence suggests a potential fundamental trade-off where early species track and possess a suite of traits related to faster growth and shorter lifespans, while later species track less and possess traits related to slower growth and longer lifespans—these later species may bet-hedge more given their longer investment window. This, however, could equally be an artifact where early species use simpler cues, and, thus, their tracking is measured more accurately given current methods.

3.3 Box: Trait trade-offs with tracking

Research on phenological tracking and traits has increased greatly in recent years, with a major uptick in studies after 2010 (see SI Fig. ??). Most papers examining tracking and other traits across species have focused on plants (20/30), followed by birds and Lepidoptera (both 4/30), plankton and aphids (both 1/30). The most studied trait was how early or late a phenophase occurred (e.g., date of flowering or start of migration for a species, termed ‘earlyness’ by some authors), with earlier species tending to track more (studies included both birds and Lepidoptera, ?????). While this is an important link, it is vulnerable to statistical challenges (see Box ‘Statistical challenges in measuring tracking’). Few studies examined whether tracking correlates with resource acquisition traits; those that did generally found species with higher tracking also had traits associated with lower competitive abilities under low resources (e.g., being shallower or lacking a taproot rooted ???). These species were often also early (e.g., ??), suggesting tracking may relate to a syndrome of traits that allow species to be rapid colonizers each season, but poor competitors for resources. Indeed, previous work has documented that species with earlier phenophases tend to have resource acquisition traits associated with lower competitive abilities (e.g., they tend to be of lower height, have shallower roots, narrower diameter vessels, thinner leaves, and grow faster, reviewed in ?).

3.4 Box: Adding tracking and non-stationarity to a common coexistence model

To understand the role of environmental tracking by species in variable environments we use a simple model that allows within- and between-year dynamics to contribute to coexistence. As the model is akin to many commonly used seed germination models (?), we follow a similar terminology for ease; however the basic structure of our model could apply to other systems with one dominant (non-renewing) pulse of a limiting resource each season (e.g., water from rain or snowpack), or a multivariate resource pulse that acts effectively as one resource (e.g., nitrogen and light drawn down together over the season). In this model the environment is included between-years via variable germination, and within-years the environment is explicitly included as a resource pulse at the start of the season. We adjust the biological start time of species (τ_i for species i) to also allow species to respond to the environment dynamically through what we refer to as tracking. Here, tracking effectively moves a species intrinsic start time closer to the environmental start time in that year, resulting in a higher germination fraction (see SI for complete description and equations).

Species can co-occur via equalizing mechanisms, but they require stabilizing mechanisms to coexist. Species that is, cannot coexist given only variation in tracking—coexistence requires variation in another trait axis. As theory and empirical work suggest this trade-off may involve traits related closely to resource competition, we varied species' R^* . With variation in tracking and in R^* species can persist together as long as those species with a temporal niche advantage are also the inferior competitors (Fig. 2). That is, species that can draw resources down to a lower level and are thus the superior within-season resource competitors (lower R^*) can persist with species with that are inferior competitors but have realized biological start times closer to the environmental start time—a finding inline with currently observed empirical trade-offs (see Box ‘Trait trade-offs with tracking’). These trade-offs, however, are all environmentally dependent. They hold only so long as the environment is stationary.

We examined how trade-offs may be transformed by a non-stationary environment, by transitioning a stationary environment—in which two-species communities had persisted for 500 years—to non-stationary, via an earlier start of season (earlier timing of the resource pulse, τ_p , Fig. 2a; see SI for more details). By changing a fundamental niche axis (the distribution of the environment, an axis along which these communities were structured), we shifted one major part of the trade-off: the new non-stationary environment favored an earlier start time than the previous stationary environment. This, in turn, reshaped our two-species communities, which depended on this trade-off for persistence.

While the non-stationary environment favored higher trackers (who in turn drove the extinction of species with lower tracking values from many two-species communities) some two-species communities persisted (257 out of 1698 two-species communities persisting after end of stationary, or 15.1%, Fig. 2). These two-species communities persisted because the same fundamental trade-off between biological start time and within-season competitive ability, while narrowed, was not fully lost (Fig. 2). Taken together, these simple simulations show how non-stationarity can drive local species extinction and reshape the underlying assembly mechanisms of commu-

613 nities.

614 Our simulations support growing work that tracking should not be considered alone (?????),
615 but may be part of a larger trait syndrome. Indeed, this model trivially show that multi-
616 species communities cannot form given only variation in the temporal niche—a trade-off is
617 required. Our results thus support empirical work showing a trade-off where trackers are also
618 inferior resource competitors (??)—we show this must be the case for multi-species persistence;
619 otherwise, the species best matched to the environment would drive the other extinct.

620 Finally, our results highlight that non-stationarity may reshape the balance of equalizing versus
621 stabilizing mechanisms. As environments shifted from stationarity to non-stationarity, species
622 that co-occurred via equalizing mechanisms persisted longer. While the outcome that equalized
623 species will be more similarly affected by environmental shifts is rather obvious, it has several
624 important implications. First, it may make identifying which traits climate change promotes
625 through stabilizing mechanisms more difficult. Second, it suggests climate change—or other
626 factors that cause an environment to shift from stationary to non-stationary—may cause a
627 fundamental shift away from assembly via stabilizing mechanisms. Thus understanding the
628 prevalence of stabilizing versus equalizing mechanisms (which ecology has worked on for many
629 decades, ??) becomes critical for understanding the implications of transitions to non-stationary
630 environments.




Figure 1: Examples of non-stationarity in climate variables linked to environmental tracking: shifts before and after 1980 (a major change-point in climate for many regions) in several metrics related to the start of growing seasons (a-c) or resource pulse connected to growing season length (d). Density plots of day of 200 growing degree day units (a metric of thermal sum, here based on 0 degree base temperature using daily minima in $^{\circ}\text{C}$) in Boston, MA, USA (a), and Munich, Germany (b), first snowfree day (followed by at least 9 snowfree days) in Crested Butte, CO, USA (c) and maximum snowdepth (mm) in March (often the month before the first snowfree day) in Crested Butte, CO, USA (d). Note that (c) and (d) are likely related, with lower snowpacks leading to an earlier first snowfree day. We selected sites that have been studied for plant phenological data and included at least 80 years of daily climate data from a Global Historical Climatology Network site; we subsetting data so that there were 40 years before and after 1980 for all sites.




Figure 2: Example of how non-stationarity can reshape communities in a simple model. We shifted the environment (top panels) by changing the timing of the resource pulse from a stationary period ($\tau_p \sim \beta(10, 10)$ for the 500 years) to a nonstationary period ($\tau_p \sim \beta(5, 15)$ over the 500 years), then examined outcomes for two-species communities (bottom panels) where tracking (X axis: species 1/species 2) trades off with R^* (Y axis: species 1/species 2): each point represents one two-species community color-coded by whether both species persisted or one or more species was extirpated through 500 years of a stationary environment (bottom-left), followed by an additional 500 years of non-stationary environment (bottom-right, only two-species communities that persisted through the stationary period are shown).




Figure 3: Goober.

632 The reviewer is correct that the resource does not renew each season, we now write “one
633 dominant (non-renewing) pulse of a limiting resource each season,” (line 578) to help clarify
634 this. As we have grossly cut the text devoted to this model we do not go into great details over
635 this now, but we have added discussion about the need for more models that place a cost on
636 early arrivals throughout the manuscript (see line 317, line 345, line 454).