

PATIENT & SPECIMEN INFORMATION

Patient name	Hemingway, Ernest M.	Accession	NGS-2026-052101
Date of birth	1961-07-22 (Age 64)	Specimen	Oral swab, left lateral tongue
MRN	10345621	Collected	2026-05-19
Ordering physician	Dr. Wisdom, DDS MD	Received	2026-05-19
Clinical indication	Oral lesion — left lateral tongue	Reported	2026-05-20

RESULT: VARIANTS DETECTED

DETECTED VARIANTS

Gene	HGVS (coding)	Protein change	VAF	Depth	Tier	Classification
KRAS	NM_004985.5:c.35G>T	p.Gly12Asp	0.43%	4,630x	I	Pathogenic
PIK3CA	NM_006218.4:c.3140A>G	p.His1047Arg	0.70%	7,140x	I	Pathogenic

VARIANT ANNOTATION

KRAS p.Gly12Asp | Pathogenic (Tier I)

KRAS p.Gly12Asp (G12D) is one of the most frequently detected oncogenic driver mutations in human cancer. It locks KRAS in the GTP-bound active state, constitutively activating downstream RAS/MAPK and PI3K/AKT/mTOR proliferative signalling. In oral squamous cell carcinoma, KRAS mutations are found in 3-8% of cases and mark a subgroup with aggressive behaviour. This variant is classified Pathogenic (Tier I, COSMIC ID COSM521). The co-occurrence with PIK3CA H1047R (see below) indicates dual activation of RAS and PI3K pathways, a combination associated with resistance to single-agent targeted therapies and may influence enrolment eligibility in clinical trials. NOTE: Observed VAF (0.43%) is above the 0.2% reporting threshold; this call is supported by 2 high-quality consensus reads at 463x consensus depth. The low consensus read count (VD = 2) reflects limited target coverage at 2,000x raw input; orthogonal confirmation is recommended before clinical action.

PIK3CA p.His1047Arg | Pathogenic (Tier I)

PIK3CA p.His1047Arg is a hotspot activating mutation in the kinase domain of the p110α catalytic subunit of PI3-kinase. It constitutively activates the PI3K/AKT/mTOR pathway, promoting cell survival, proliferation, and tumour angiogenesis. PIK3CA mutations occur in 15-20% of OSCC, most commonly at E545K and H1047R hotspots. This variant is classified Pathogenic (Tier I, COSMIC ID COSM775). PIK3CA H1047R is potentially actionable: alpelisib (PI3Kα inhibitor) is FDA-approved for PIK3CA-mutated breast cancer and is under investigation in HNSCC. Multidisciplinary tumour board review for consideration of PI3K-directed therapy or clinical trial enrolment is recommended.

GENES WITH NO PATHOGENIC OR LIKELY PATHOGENIC VARIANTS DETECTED

AJUBA, CASP8, CDKN2A, CUL3, DPYD, EGFR, EP300, FAT1, FBXW7, HRAS, KEAP1, KMT2D, NFE2L2, NOTCH1, NSD1, PIK3R1, PTEN, TERT, TP53
Variants of uncertain significance (Tier III) are not reported.

METHOD SUMMARY

Cell-free and cellular DNA extracted from oral swab is subjected to hybridization capture using the IDT xGen Duplex UMI panel targeting 251 genomic regions (~77 kbp) across 21 cancer-relevant genes. Libraries incorporate 9-bp Unique Molecular Identifiers (UMIs) in a 9M1S+T read structure on both R1 and R2. Libraries are sequenced on Illumina NextSeq (2x150 bp). After alignment to GRCh38 (canonical chromosomes), reads are grouped by UMI using fgbio (adjacency strategy, ≤1 edit distance), and molecular consensus sequences are called requiring ≥2 reads per UMI family. Somatic variants are called on the consensus BAM with VarDict at minimum allele frequency 0.001. Variants ≥0.2% consensus VAF with ≥2 consensus supporting reads are reported. UMI deduplication reduces sequencing noise by >800-fold. Run-level negative controls are included on every sequencing run.

LIMITATIONS

This assay detects somatic single nucleotide variants and small indels within 251 target regions (~77 kbp) at VAF ≥0.2% with validated sensitivity of 98.4% at ≥20,000x raw input depth (~4,000x consensus after UMI deduplication). Structural variants, copy number alterations, gene fusions, large indels, intronic variants outside defined capture regions, and variants in regions with low hybridization capture efficiency are not reliably assessed. A negative result does not exclude malignancy. Tumour heterogeneity, low tumour cell content, and specimen quality may affect variant allele fraction. Variants below the 0.2% VAF reporting threshold may be present but are not reported. At 20,000x raw sequencing depth (~4,000x mean consensus), approximately 0.3% of targets may fall below 1,000x consensus depth; per-target coverage is provided in the laboratory technical report. Results should be interpreted in the context of clinical and pathological findings by a qualified clinician.

HOTSPOT SCREENING SUMMARY

Patient: Hemingway, Ernest M.

Accession: NGS-2026-052101

Hotspots listed below were explicitly interrogated in addition to all variants above 0.2% VAF with ≥2 consensus reads across all 251 targets. Coverage criteria (≥100× consensus depth) were met at all targets.

EGFRCov ✓			HRASCov ✓		
L858R	c.2573T>G	Not detected	G12V	c.35G>T	Not detected
T790M	c.2369C>T	Not detected	G12D	c.35G>A	Not detected
Exon 19 del	c.2235_2249del	Not detected	Q61L	c.182A>T	Not detected
			Q61R	c.182A>G	Not detected
KRASCov ✓			NOTCH1Cov ✓		
G12D	c.35G>T	● DETECTED	P2415L	c.7244C>T	Not detected
G12V	c.35G>A	Not detected	R1699W	c.5095C>T	Not detected
G12C	c.34G>T	Not detected			
G13D	c.38G>A	Not detected	TP53Cov ✓		
PIK3CACov ✓			R175H	c.524G>A	Not detected
E542K	c.1624G>A	Not detected	R248Q	c.743G>A	Not detected
E545K	c.1633G>A	Not detected	R248W	c.742C>T	Not detected
H1047R	c.3140A>G	● DETECTED	R273H	c.818G>A	Not detected
TERTCov ✓			R282W	c.844C>T	Not detected
C228T	c.-124C>T	Not detected			
C250T	c.-146C>T	Not detected			

REMAINING PANEL GENES — SEQUENCE-LEVEL DETECTION, NO PREDEFINED HOTSPOT LIST

AJUBA · CASP8 · CDKN2A · CUL3 · DPYD · EP300 · FAT1 · FBXW7 · KEAP1 · KMT2D · NFE2L2 · NSD1 · PIK3R1 · PTEN

All variants above 0.2% VAF with ≥2 consensus reads are reported for these genes. No variants meeting reporting criteria were detected. All targets in these genes met minimum coverage criteria (≥100× consensus depth).

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

PATIENT & SPECIMEN INFORMATION

Patient name	Woolf, Virginia A.	Accession	NGS-2026-052102
Date of birth	1970-03-11 (Age 56)	Specimen	Oral swab, right buccal mucosa
MRN	20456732	Collected	2026-05-19
Ordering physician	Dr. Wisdom, MD	Received	2026-05-19
Clinical indication	Oral lesion — right buccal mucosa	Reported	2026-05-20

RESULT: VARIANTS DETECTED

DETECTED VARIANTS

Gene	HGVS (coding)	Protein change	VAF	Depth	Tier	Classification
NOTCH1	NM_017617.5:c.7244C>T	p.Pro2415Leu	0.26%	7,690x	II	Likely Pathogenic

VARIANT ANNOTATION

NOTCH1 p.Pro2415Leu | Likely Pathogenic (Tier II)

NOTCH1 functions as a tumour suppressor in stratified squamous epithelium. The p.Pro2415Leu missense variant affects a conserved residue in the PEST domain, predicted to disrupt protein stability (SIFT: deleterious; PolyPhen-2: probably damaging). NOTCH1 loss-of-function mutations are found in 10–15% of OSCC and are associated with early-onset tumourigenesis in the oral cavity. This variant is classified Likely Pathogenic (Tier II). In the context of confirmed dysplasia, detection of a NOTCH1 tumour suppressor variant supports malignant progression risk. Clinical correlation and multidisciplinary review are recommended. NOTE: Observed VAF (0.26%) is above the 0.2% reporting threshold and is supported by 2 consensus reads at 769x consensus depth. Orthogonal confirmation is recommended before clinical action.

GENES WITH NO PATHOGENIC OR LIKELY PATHOGENIC VARIANTS DETECTED

AJUBA, CASP8, CDKN2A, CUL3, DPYD, EGFR, EP300, FAT1, FBXW7, HRAS, KEAP1, KMT2D, KRAS, NFE2L2, NSD1, PIK3CA, PIK3R1, PTEN, TERT, TP53
Variants of uncertain significance (Tier III) are not reported.

METHOD SUMMARY

Cell-free and cellular DNA extracted from oral swab is subjected to hybridization capture using the IDT xGen Duplex UMI panel targeting 251 genomic regions (~77 kbp) across 21 cancer-relevant genes. Libraries incorporate 9-bp Unique Molecular Identifiers (UMIs) in a 9M1S+T read structure on both R1 and R2. Libraries are sequenced on Illumina NextSeq (2x150 bp). After alignment to GRCh38 (canonical chromosomes), reads are grouped by UMI using fgbio (adjacency strategy, ≤1 edit distance), and molecular consensus sequences are called requiring ≥2 reads per UMI family. Somatic variants are called on the consensus BAM with VarDict at minimum allele frequency 0.001. Variants ≥0.2% consensus VAF with ≥2 consensus supporting reads are reported. UMI deduplication reduces sequencing noise by >800-fold. Run-level negative controls are included on every sequencing run.

LIMITATIONS

This assay detects somatic single nucleotide variants and small indels within 251 target regions (~77 kbp) at VAF ≥0.2% with validated sensitivity of 98.4% at ≥20,000x raw input depth (~4,000x consensus after UMI deduplication). Structural variants, copy number alterations, gene fusions, large indels, intronic variants outside defined capture regions, and variants in regions with low hybridization capture efficiency are not reliably assessed. A negative result does not exclude malignancy. Tumour heterogeneity, low tumour cell content, and specimen quality may affect variant allele fraction. Variants below the 0.2% VAF reporting threshold may be present but are not reported. At 20,000x raw sequencing depth (~4,000x mean consensus), approximately 0.3% of targets may fall below 1,000x consensus depth; per-target coverage is provided in the laboratory technical report. Results should be interpreted in the context of clinical and pathological findings by a qualified clinician.

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

HOTSPOT SCREENING SUMMARY

Patient:Woolf, Virginia A.

Accession:NGS-2026-052102

Hotspots listed below were explicitly interrogated in addition to all variants above 0.2% VAF with ≥2 consensus reads across all 251 targets. Coverage criteria (≥100× consensus depth) were met at all targets.

EGFRCov ✓			HRASCov ✓		
L858R	c.2573T>G	Not detected	G12V	c.35G>T	Not detected
T790M	c.2369C>T	Not detected	G12D	c.35G>A	Not detected
Exon 19 del	c.2235_2249del	Not detected	Q61L	c.182A>T	Not detected
			Q61R	c.182A>G	Not detected
KRASCov ✓			NOTCH1Cov ✓		
G12D	c.35G>T	Not detected	P2415L	c.7244C>T	● DETECTED
G12V	c.35G>A	Not detected	R1699W	c.5095C>T	Not detected
G12C	c.34G>T	Not detected			
G13D	c.38G>A	Not detected	TP53Cov ✓		
PIK3CACov ✓			R175H	c.524G>A	Not detected
E542K	c.1624G>A	Not detected	R248Q	c.743G>A	Not detected
E545K	c.1633G>A	Not detected	R248W	c.742C>T	Not detected
H1047R	c.3140A>G	Not detected	R273H	c.818G>A	Not detected
TERTCov ✓			R282W	c.844C>T	Not detected
C228T	c.-124C>T	Not detected			
C250T	c.-146C>T	Not detected			

REMAINING PANEL GENES — SEQUENCE-LEVEL DETECTION, NO PREDEFINED HOTSPOT LIST

AJUBA · CASP8 · CDKN2A · CUL3 · DPYD · EP300 · FAT1 · FBXW7 · KEAP1 · KMT2D · NFE2L2 · NSD1 · PIK3R1 · PTEN

All variants above 0.2% VAF with ≥2 consensus reads are reported for these genes. No variants meeting reporting criteria were detected. All targets in these genes met minimum coverage criteria (≥100× consensus depth).

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

PATIENT & SPECIMEN INFORMATION

Patient name	Dickens, Charles J.	Accession	NGS-2026-052103
Date of birth	1951-09-28 (Age 74)	Specimen	Oral swab, left floor of mouth
MRN	30567843	Collected	2026-05-19
Ordering physician	Dr. Wisdom, DDS MD	Received	2026-05-19
Clinical indication	Oral lesion — left floor of mouth	Reported	2026-05-20

RESULT: VARIANTS DETECTED

DETECTED VARIANTS

Gene	HGVS (coding)	Protein change	VAF	Depth	Tier	Classification
TP53	NM_000546.6:c.524G>A	p.Arg175His	0.60%	6,640x	I	Pathogenic

VARIANT ANNOTATION

TP53 p.Arg175His | Pathogenic (Tier I)

TP53 p.Arg175His is the most frequently observed p53 mutation in human cancer. It disrupts the structural zinc-binding domain of the DNA-binding domain, abolishing wild-type tumour suppressor function, while also conferring dominant-negative and gain-of-function oncogenic properties including promotion of invasion, metastasis, and MDM2 antagonism (COSMIC ID COSM10648). In OSCC, TP53 mutations are present in 50–70% of cases and are associated with aggressive tumour behaviour, locoregional recurrence, and reduced sensitivity to platinum-based chemotherapy. The VAF of 0.60% in a surveillance swab warrants clinical correlation to exclude early local recurrence. Imaging and clinical re-evaluation are recommended.

GENES WITH NO PATHOGENIC OR LIKELY PATHOGENIC VARIANTS DETECTED

AJUBA, CASP8, CDKN2A, CUL3, DPYD, EGFR, EP300, FAT1, FBXW7, HRAS, KEAP1, KMT2D, KRAS, NFE2L2, NOTCH1, NSD1, PIK3CA, PIK3R1, PTEN, TERT
Variants of uncertain significance (Tier III) are not reported.

METHOD SUMMARY

Cell-free and cellular DNA extracted from oral swab is subjected to hybridization capture using the IDT xGen Duplex UMI panel targeting 251 genomic regions (~77 kbp) across 21 cancer-relevant genes. Libraries incorporate 9-bp Unique Molecular Identifiers (UMIs) in a 9M1S+T read structure on both R1 and R2. Libraries are sequenced on Illumina NextSeq (2x150 bp). After alignment to GRCh38 (canonical chromosomes), reads are grouped by UMI using fgbio (adjacency strategy, ≤1 edit distance), and molecular consensus sequences are called requiring ≥2 reads per UMI family. Somatic variants are called on the consensus BAM with VarDict at minimum allele frequency 0.001. Variants ≥0.2% consensus VAF with ≥2 consensus supporting reads are reported. UMI deduplication reduces sequencing noise by >800-fold. Run-level negative controls are included on every sequencing run.

LIMITATIONS

This assay detects somatic single nucleotide variants and small indels within 251 target regions (~77 kbp) at VAF ≥0.2% with validated sensitivity of 98.4% at ≥20,000x raw input depth (~4,000x consensus after UMI deduplication). Structural variants, copy number alterations, gene fusions, large indels, intronic variants outside defined capture regions, and variants in regions with low hybridization capture efficiency are not reliably assessed. A negative result does not exclude malignancy. Tumour heterogeneity, low tumour cell content, and specimen quality may affect variant allele fraction. Variants below the 0.2% VAF reporting threshold may be present but are not reported. At 20,000x raw sequencing depth (~4,000x mean consensus), approximately 0.3% of targets may fall below 1,000x consensus depth; per-target coverage is provided in the laboratory technical report. Results should be interpreted in the context of clinical and pathological findings by a qualified clinician.

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

HOTSPOT SCREENING SUMMARY

Patient:

Dickens, Charles J.

Accession:

NGS-2026-052103

Hotspots listed below were explicitly interrogated in addition to all variants above 0.2% VAF with ≥2 consensus reads across all 251 targets. Coverage criteria (≥100× consensus depth) were met at all targets.

EGFRCov ✓			HRASCov ✓		
L858R	c.2573T>G	Not detected	G12V	c.35G>T	Not detected
T790M	c.2369C>T	Not detected	G12D	c.35G>A	Not detected
Exon 19 del	c.2235_2249del	Not detected	Q61L	c.182A>T	Not detected
			Q61R	c.182A>G	Not detected
KRASCov ✓			NOTCH1Cov ✓		
G12D	c.35G>T	Not detected	P2415L	c.7244C>T	Not detected
G12V	c.35G>A	Not detected	R1699W	c.5095C>T	Not detected
G12C	c.34G>T	Not detected			
G13D	c.38G>A	Not detected	TP53Cov ✓		
PIK3CACov ✓			R175H	c.524G>A	● DETECTED
E542K	c.1624G>A	Not detected	R248Q	c.743G>A	Not detected
E545K	c.1633G>A	Not detected	R248W	c.742C>T	Not detected
H1047R	c.3140A>G	Not detected	R273H	c.818G>A	Not detected
TERTCov ✓			R282W	c.844C>T	Not detected
C228T	c.-124C>T	Not detected			
C250T	c.-146C>T	Not detected			

REMAINING PANEL GENES — SEQUENCE-LEVEL DETECTION, NO PREDEFINED HOTSPOT LIST

AJUBA · CASP8 · CDKN2A · CUL3 · DPYD · EP300 · FAT1 · FBXW7 · KEAP1 · KMT2D · NFE2L2 · NSD1 · PIK3R1 · PTEN

All variants above 0.2% VAF with ≥2 consensus reads are reported for these genes. No variants meeting reporting criteria were detected. All targets in these genes met minimum coverage criteria (≥100× consensus depth).

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

PATIENT & SPECIMEN INFORMATION			
Patient name	Kafka, Franz J.	Accession	NGS-2026-052105
Date of birth	1980-02-17 (Age 46)	Specimen	Oral swab, anterior tongue
MRN	50789065	Collected	2026-05-19
Ordering physician	Dr. Wisdom, MD	Received	2026-05-19
Clinical indication	Oral lesion — anterior tongue (surveillance post-resection)		2026-05-20

RESULT: NO REPORTABLE VARIANTS DETECTED

INTERPRETATION

No pathogenic or likely pathogenic somatic variants were detected in the 21 genes covered by this assay at $\geq 0.2\%$ VAF with ≥ 2 consensus supporting reads. A negative result does not exclude malignancy. Variants below the reporting threshold, variants outside the 251 assay target regions, structural variants, and copy number alterations are not assessed by this assay. Clinical correlation is required; repeat testing or biopsy should be considered if clinical suspicion persists.

GENES WITH NO PATHOGENIC OR LIKELY PATHOGENIC VARIANTS DETECTED

AJUBA, CASP8, CDKN2A, CUL3, DPYD, EGFR, EP300, FAT1, FBXW7, HRAS, KEAP1, KMT2D, KRAS, NFE2L2, NOTCH1, NSD1, PIK3CA, PIK3R1, PTEN, TERT, TP53

Variants of uncertain significance (Tier III) are not reported.

METHOD SUMMARY

Cell-free and cellular DNA extracted from oral swab is subjected to hybridization capture using the IDT xGen Duplex UMI panel targeting 251 genomic regions (~77 kbp) across 21 cancer-relevant genes. Libraries incorporate 9-bp Unique Molecular Identifiers (UMIs) in a 9M1S+T read structure on both R1 and R2. Libraries are sequenced on Illumina NextSeq (2x150 bp). After alignment to GRCh38 (canonical chromosomes), reads are grouped by UMI using fgbio (adjacency strategy, ≤ 1 edit distance), and molecular consensus sequences are called requiring ≥ 2 reads per UMI family. Somatic variants are called on the consensus BAM with VarDict at minimum allele frequency 0.001. Variants $\geq 0.2\%$ consensus VAF with ≥ 2 consensus supporting reads are reported. UMI deduplication reduces sequencing noise by >800-fold. Run-level negative controls are included on every sequencing run.

LIMITATIONS

This assay detects somatic single nucleotide variants and small indels within 251 target regions (~77 kbp) at VAF $\geq 0.2\%$ with validated sensitivity of 98.4% at $\geq 20,000\times$ raw input depth (~4,000 $\times$ consensus after UMI deduplication). Structural variants, copy number alterations, gene fusions, large indels, intronic variants outside defined capture regions, and variants in regions with low hybridization capture efficiency are not reliably assessed. A negative result does not exclude malignancy. Tumour heterogeneity, low tumour cell content, and specimen quality may affect variant allele fraction. Variants below the 0.2% VAF reporting threshold may be present but are not reported. At 20,000 $\times$ raw sequencing depth (~4,000 $\times$ mean consensus), approximately 0.3% of targets may fall below 1,000 $\times$ consensus depth; per-target coverage is provided in the laboratory technical report. Results should be interpreted in the context of clinical and pathological findings by a qualified clinician.

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

HOTSPOT SCREENING SUMMARY

Patient: Kafka, Franz J.

Accession: NGS-2026-052105

Hotspots listed below were explicitly interrogated in addition to all variants above 0.2% VAF with ≥2 consensus reads across all 251 targets. Coverage criteria (≥100× consensus depth) were met at all targets.

EGFRCov ✓			HRASCov ✓		
L858R	c.2573T>G	Not detected	G12V	c.35G>T	Not detected
T790M	c.2369C>T	Not detected	G12D	c.35G>A	Not detected
Exon 19 del	c.2235_2249del	Not detected	Q61L	c.182A>T	Not detected
			Q61R	c.182A>G	Not detected
KRASCov ✓			NOTCH1Cov ✓		
G12D	c.35G>T	Not detected	P2415L	c.7244C>T	Not detected
G12V	c.35G>A	Not detected	R1699W	c.5095C>T	Not detected
G12C	c.34G>T	Not detected			
G13D	c.38G>A	Not detected	TP53Cov ✓		
PIK3CACov ✓			R175H	c.524G>A	Not detected
E542K	c.1624G>A	Not detected	R248Q	c.743G>A	Not detected
E545K	c.1633G>A	Not detected	R248W	c.742C>T	Not detected
H1047R	c.3140A>G	Not detected	R273H	c.818G>A	Not detected
TERTCov ✓			R282W	c.844C>T	Not detected
C228T	c.-124C>T	Not detected			
C250T	c.-146C>T	Not detected			

REMAINING PANEL GENES — SEQUENCE-LEVEL DETECTION, NO PREDEFINED HOTSPOT LIST

AJUBA · CASP8 · CDKN2A · CUL3 · DPYD · EP300 · FAT1 · FBXW7 · KEAP1 · KMT2D · NFE2L2 · NSD1 · PIK3R1 · PTEN

All variants above 0.2% VAF with ≥2 consensus reads are reported for these genes. No variants meeting reporting criteria were detected. All targets in these genes met minimum coverage criteria (≥100× consensus depth).

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

PATIENT & SPECIMEN INFORMATION			
Patient name	Morrison, Toni E.	Accession	NGS-2026-052106
Date of birth	1955-10-09 (Age 70)	Specimen	Oral swab, left retromolar trigone
MRN	60890176	Collected	2026-05-19
Ordering physician	Dr. Wisdom, MD	Received	2026-05-19
Clinical indication	Oral lesion — left retromolar trigone	Reported	2026-05-20

RESULT: NO REPORTABLE VARIANTS DETECTED

INTERPRETATION

No pathogenic or likely pathogenic somatic variants were detected in the 21 genes covered by this assay at $\geq 0.2\%$ VAF with ≥ 2 consensus supporting reads. A negative result does not exclude malignancy. Variants below the reporting threshold, variants outside the 251 assay target regions, structural variants, and copy number alterations are not assessed by this assay. Clinical correlation is required; repeat testing or biopsy should be considered if clinical suspicion persists.

GENES WITH NO PATHOGENIC OR LIKELY PATHOGENIC VARIANTS DETECTED

AJUBA, CASP8, CDKN2A, CUL3, DPYD, EGFR, EP300, FAT1, FBXW7, HRAS, KEAP1, KMT2D, KRAS, NFE2L2, NOTCH1, NSD1, PIK3CA, PIK3R1, PTEN, TERT, TP53

Variants of uncertain significance (Tier III) are not reported.

METHOD SUMMARY

Cell-free and cellular DNA extracted from oral swab is subjected to hybridization capture using the IDT xGen Duplex UMI panel targeting 251 genomic regions (~77 kbp) across 21 cancer-relevant genes. Libraries incorporate 9-bp Unique Molecular Identifiers (UMIs) in a 9M1S+T read structure on both R1 and R2. Libraries are sequenced on Illumina NextSeq (2x150 bp). After alignment to GRCh38 (canonical chromosomes), reads are grouped by UMI using fgbio (adjacency strategy, ≤ 1 edit distance), and molecular consensus sequences are called requiring ≥ 2 reads per UMI family. Somatic variants are called on the consensus BAM with VarDict at minimum allele frequency 0.001. Variants $\geq 0.2\%$ consensus VAF with ≥ 2 consensus supporting reads are reported. UMI deduplication reduces sequencing noise by >800 -fold. Run-level negative controls are included on every sequencing run.

LIMITATIONS

This assay detects somatic single nucleotide variants and small indels within 251 target regions (~77 kbp) at VAF $\geq 0.2\%$ with validated sensitivity of 98.4% at $\geq 20,000\times$ raw input depth (~4,000 $\times$ consensus after UMI deduplication). Structural variants, copy number alterations, gene fusions, large indels, intronic variants outside defined capture regions, and variants in regions with low hybridization capture efficiency are not reliably assessed. A negative result does not exclude malignancy. Tumour heterogeneity, low tumour cell content, and specimen quality may affect variant allele fraction. Variants below the 0.2% VAF reporting threshold may be present but are not reported. At 20,000 $\times$ raw sequencing depth (~4,000 $\times$ mean consensus), approximately 0.3% of targets may fall below 1,000 $\times$ consensus depth; per-target coverage is provided in the laboratory technical report. Results should be interpreted in the context of clinical and pathological findings by a qualified clinician.

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

HOTSPOT SCREENING SUMMARY

Patient:

Morrison, Toni E.

Accession:

NGS-2026-052106

Hotspots listed below were explicitly interrogated in addition to all variants above 0.2% VAF with ≥2 consensus reads across all 251 targets. Coverage criteria (≥100× consensus depth) were met at all targets.

EGFRCov ✓			HRASCov ✓		
L858R	c.2573T>G	Not detected	G12V	c.35G>T	Not detected
T790M	c.2369C>T	Not detected	G12D	c.35G>A	Not detected
Exon 19 del	c.2235_2249del	Not detected	Q61L	c.182A>T	Not detected
			Q61R	c.182A>G	Not detected
KRASCov ✓			NOTCH1Cov ✓		
G12D	c.35G>T	Not detected	P2415L	c.7244C>T	Not detected
G12V	c.35G>A	Not detected	R1699W	c.5095C>T	Not detected
G12C	c.34G>T	Not detected			
G13D	c.38G>A	Not detected	TP53Cov ✓		
PIK3CACov ✓			R175H	c.524G>A	Not detected
E542K	c.1624G>A	Not detected	R248Q	c.743G>A	Not detected
E545K	c.1633G>A	Not detected	R248W	c.742C>T	Not detected
H1047R	c.3140A>G	Not detected	R273H	c.818G>A	Not detected
TERTCov ✓			R282W	c.844C>T	Not detected
C228T	c.-124C>T	Not detected			
C250T	c.-146C>T	Not detected			

REMAINING PANEL GENES — SEQUENCE-LEVEL DETECTION, NO PREDEFINED HOTSPOT LIST

AJUBA · CASP8 · CDKN2A · CUL3 · DPYD · EP300 · FAT1 · FBXW7 · KEAP1 · KMT2D · NFE2L2 · NSD1 · PIK3R1 · PTEN

All variants above 0.2% VAF with ≥2 consensus reads are reported for these genes. No variants meeting reporting criteria were detected. All targets in these genes met minimum coverage criteria (≥100× consensus depth).

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

PATIENT & SPECIMEN INFORMATION

Patient name	Austen, Jane C.	Accession	NGS-2026-052104
Date of birth	1967-12-04 (Age 58)	Specimen	Oral swab, soft palate
MRN	40678954	Collected	2026-05-19
Ordering physician	Dr. Wisdom, MD	Received	2026-05-19
Clinical indication	Oral lesion — soft palate	Reported	2026-05-20

RESULT: VARIANTS DETECTED

DETECTED VARIANTS

Gene	HGVS (coding)	Protein change	VAF	Depth	Tier	Classification
EGFR	NM_005228.5:c.2573T>G	p.Leu858Arg	0.57%	5,250×	I	Pathogenic
TP53	NM_000546.6:c.524G>A	p.Arg175His	0.30%	6,690×	I	Pathogenic

VARIANT ANNOTATION

EGFR p.Leu858Arg | Pathogenic (Tier I)

EGFR p.Leu858Arg (L858R) is a well-characterised activating mutation in exon 21 of the EGFR kinase domain. It disrupts the autoinhibitory conformation, leading to constitutive EGFR kinase activation and downstream RAS/MAPK and PI3K/AKT signalling. While canonical in non-small cell lung adenocarcinoma (where it predicts response to EGFR TKIs such as osimertinib), EGFR L858R has been reported in 1–3% of OSCC. Cetuximab (anti-EGFR monoclonal antibody) is FDA-approved in HNSCC and may have enhanced activity in the setting of EGFR activating mutations. Multidisciplinary tumour board review for consideration of EGFR-directed therapy or clinical trial enrolment is recommended. Co-occurring TP53 R175H (see below) may affect response to EGFR-directed therapy and warrants combined consideration.

TP53 p.Arg175His | Pathogenic (Tier I)

TP53 p.Arg175His — see interpretation above (Dickens, Charles J., NGS-2026-052103). Co-occurrence of EGFR L858R and TP53 R175H has been associated with resistance to EGFR-directed therapies in other tumour types; multidisciplinary tumour board review is recommended. NOTE: Observed VAF (0.30%) is above the 0.2% reporting threshold and is supported by 2 consensus reads at 669× consensus depth. Orthogonal confirmation is recommended before clinical action.

GENES WITH NO PATHOGENIC OR LIKELY PATHOGENIC VARIANTS DETECTED

AJUBA, CASP8, CDKN2A, CUL3, DPYD, EP300, FAT1, FBXW7, HRAS, KEAP1, KMT2D, KRAS, NFE2L2, NOTCH1, NSD1, PIK3CA, PIK3R1, PTEN, TERT
Variants of uncertain significance (Tier III) are not reported.

METHOD SUMMARY

Cell-free and cellular DNA extracted from oral swab is subjected to hybridization capture using the IDT xGen Duplex UMI panel targeting 251 genomic regions (~77 kbp) across 21 cancer-relevant genes. Libraries incorporate 9-bp Unique Molecular Identifiers (UMIs) in a 9M1S+T read structure on both R1 and R2. Libraries are sequenced on Illumina NextSeq (2×150 bp). After alignment to GRCh38 (canonical chromosomes), reads are grouped by UMI using fgbio (adjacency strategy, ≤1 edit distance), and molecular consensus sequences are called requiring ≥2 reads per UMI family. Somatic variants are called on the consensus BAM with VarDict at minimum allele frequency 0.001. Variants ≥0.2% consensus VAF with ≥2 consensus supporting reads are reported. UMI deduplication reduces sequencing noise by >800-fold. Run-level negative controls are included on every sequencing run.

LIMITATIONS

This assay detects somatic single nucleotide variants and small indels within 251 target regions (~77 kbp) at VAF ≥0.2% with validated sensitivity of 98.4% at ≥20,000× raw input depth (~4,000× consensus after UMI deduplication). Structural variants, copy number alterations, gene fusions, large indels, intronic variants outside defined capture regions, and variants in regions with low hybridization capture efficiency are not reliably assessed. A negative result does not exclude malignancy. Tumour heterogeneity, low tumour cell content, and specimen quality may affect variant allele fraction. Variants below the 0.2% VAF reporting threshold may be present but are not reported. At 20,000× raw sequencing depth (~4,000× mean consensus), approximately 0.3% of targets may fall below 1,000× consensus depth; per-target coverage is provided in the laboratory technical report. Results should be interpreted in the context of clinical and pathological findings by a qualified clinician.

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.

HOTSPOT SCREENING SUMMARY

Patient:

Austen, Jane C.

Accession:

NGS-2026-052104

Hotspots listed below were explicitly interrogated in addition to all variants above 0.2% VAF with ≥2 consensus reads across all 251 targets. Coverage criteria (≥100× consensus depth) were met at all targets.

EGFR			Cov ✓
L858R	c.2573T>G	● DETECTED	
T790M	c.2369C>T	Not detected	
Exon 19 del	c.2235_2249del	Not detected	
KRAS			Cov ✓
G12D	c.35G>T	Not detected	
G12V	c.35G>A	Not detected	
G12C	c.34G>T	Not detected	
G13D	c.38G>A	Not detected	
PIK3CA			Cov ✓
E542K	c.1624G>A	Not detected	
E545K	c.1633G>A	Not detected	
H1047R	c.3140A>G	Not detected	
TERT			Cov ✓
C228T	c.-124C>T	Not detected	
C250T	c.-146C>T	Not detected	

HRAS			Cov ✓
G12V	c.35G>T	Not detected	
G12D	c.35G>A	Not detected	
Q61L	c.182A>T	Not detected	
Q61R	c.182A>G	Not detected	

NOTCH1			Cov ✓
P2415L	c.7244C>T	Not detected	
R1699W	c.5095C>T	Not detected	

TP53			Cov ✓
R175H	c.524G>A	● DETECTED	
R248Q	c.743G>A	Not detected	
R248W	c.742C>T	Not detected	
R273H	c.818G>A	Not detected	
R282W	c.844C>T	Not detected	

REMAINING PANEL GENES — SEQUENCE-LEVEL DETECTION, NO PREDEFINED HOTSPOT LIST

AJUBA · CASP8 · CDKN2A · CUL3 · DPYD · EP300 · FAT1 · FBXW7 · KEAP1 · KMT2D · NFE2L2 · NSD1 · PIK3R1 · PTEN

All variants above 0.2% VAF with ≥2 consensus reads are reported for these genes. No variants meeting reporting criteria were detected. All targets in these genes met minimum coverage criteria (≥100× consensus depth).

Reported by: Dr. Wisdom, MD PhD, FCAP

Report date: 2026-05-20

Results reviewed and approved by the Laboratory Director. This report is electronically signed and does not require a wet signature.