

Article

Note on the action in an elliptic orbit.

in: The quarterly journal of pure and applied mathematics | The quarterly

journal of pure and applied mathematics - 7

## Terms and Conditions

The Göttingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept there Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

### **Contact:**

Niedersächsische Staats- und Universitätsbibliothek

Digitalisierungszentrum

37070 Goettingen

Germany

Email: [gdz@sub.uni-goettingen.de](mailto:gdz@sub.uni-goettingen.de)

### **Purchase a CD-ROM**

The Goettingen State and University Library offers CD-ROMs containing whole volumes / monographs in PDF for Adobe Acrobat. The PDF-version contains the table of contents as bookmarks, which allows easy navigation in the document. For availability and pricing, please contact:

Niedersaechsische Staats- und Universitaetsbibliothek Goettingen - Digitalisierungszentrum

37070 Goettingen, Germany, Email: [gdz@sub.uni-goettingen.de](mailto:gdz@sub.uni-goettingen.de)

## NOTE ON THE ACTION IN AN ELLIPTIC ORBIT.

IF the moving particle have unit mass, the expression for the action is

$$\int v ds.$$

But, if  $p$  and  $p'$  be the perpendiculars from the foci on the tangent, we have

$$pv = h,$$

and

$$pp' = b^2.$$

Hence the action may be expressed by

$$\frac{h}{b^2} \int p' ds.$$

That is; *while the time is proportional to the area described about one focus, the action is proportional to that described about the other.*

Of course it is easy to extend this result to parabolic and hyperbolic orbits. In the case of the parabola it becomes changed to the following, *the increase of action is proportional to the increase of distance from the axis.*

Sir W. R. Hamilton in his "General Method in Dynamics" (*Philosophical Transactions*, 1834) has shown that the action in an elliptic orbit is (with the usual notation) proportional to

$$u + e \sin u,$$

but he does not give the above simple interpretation.

T.