

Weakly nonlinear scalar scattering from a Kerr black hole

KerrScattering Collaboration¹

¹*KerrScattering Project*

We study weakly nonlinear scattering of a complex scalar field from a Kerr black hole with a cubic self-interaction. We solve the spin-weight-zero Teukolsky equation with an endpoint-inclusive, two-domain Chebyshev method and construct the first-order Green-function response. Nonlinear transmission and reflection coefficients are defined as derivatives of the horizon and infinity energy fluxes at fixed incident flux. Rotation enters through the co-rotating horizon frequency, spheroidal harmonics, and the Σ -weighted source projection.

For an axisymmetric sample with $a = 0.5$ and $\ell = 0, 1, 2$, the accepted spectra show broad Breit-Wigner-type features whose fitted centers lie on the same frequency scale as the real parts of the corresponding scalar Kerr quasinormal frequencies. The high-frequency data are described by finite-window exponential fits with effective temperatures within 3.2–13.6% of the Kerr Hawking temperature. The low-frequency transmission coefficients follow a centrifugal-barrier hierarchy, while the reflected first-order coefficient becomes cancellation limited at the lowest frequencies. Spectral tails, field-dependent radial residuals, Wronskian constancy, resolution scans, and flux balance are used together as internal accuracy tests; the accepted first-order balance residuals are at or below 4.3×10^{-12} . A bounded $\ell = m = 2$ scan resolves the signed horizon-flux change across the superradiant threshold.

This is a prescribed scalar response on a fixed Kerr background. It is not a second-order metric perturbation, a gravitational self-force calculation, or an EMRI waveform model.

I. INTRODUCTION

We use geometric units $G = c = 1$, metric signature $(-, +, +, +)$, and set $M = 1$ in the numerical results unless stated otherwise. The Fourier convention is fixed by the mode dependence $e^{-i\omega t + im\phi}$ used below.

Scattering is a foundational probe of black-hole spacetimes. The linear problem has been studied extensively for scalar, electromagnetic, and gravitational waves in Schwarzschild and Kerr backgrounds, using partial-wave, complex-angular-momentum, and related frequency-domain methods [1–6]. In a standard scattering experiment one sends in a monochromatic mode from past null infinity and compares the incident flux with the reflected flux at future null infinity and the absorbed flux at the future event horizon. The resulting transmission and reflection coefficients are natural complements to quasinormal-mode spectroscopy [7]. This point has become sharper because quasinormal spectra can be sensitive to small deformations of the effective potential [8–10], while scattering data, greybody factors, and related response functions can remain stable observables [11–15].

The nonlinear scattering problem is less developed. Time-domain studies have examined nonlinear perturbative and fully nonlinear scattering from past to future null infinity [16, 17], and recent work on nonlinear dynamics in general relativity has studied frequency-dependent scattering and mode generation [29]. The Schwarzschild calculation that we use as a template instead keeps the flux definition of scattering coefficients explicit: it considers a weak scalar self-interaction, solves the linear homogeneous problem, uses the nonlinear source to construct a first-order Green-function correction, and defines nonlinear scattering coefficients from the interference terms in the energy flux. This is the key point: the first-order field

is not a new independent linear scattering experiment; it changes the physical boundary fluxes by interfering with the zeroth-order wave.

The scalar model is not a gravitational self-force calculation, but its bookkeeping is closely related to the sourced perturbative hierarchies used in extreme-mass-ratio inspiral (EMRI) theory: a lower-order field generates a higher-order source, mode coupling must be projected consistently, and observable fluxes provide an important global check [18–20]. We use this analogy only to motivate the workflow; no claim about a gravitational waveform or a self-force observable is made here. Frequency-domain calculations for a scalar charge on Kerr, generic bound Kerr geodesics, and the associated regularization parameters make the additional worldline and singular-field structures explicit [21, 23, 24]. Effective-source and extended-particular-solution methods provide a related scalar prototype for controlling frequency-domain reconstruction, but they retain a singular worldline source and regularization step absent here [22]. Recent second-order self-force and metric-reconstruction work further shows why those structures cannot be replaced by the finite cubic source used here [25, 26]. Recent effective-source constructions make the distinction concrete for both generic first-order orbits and second-order quasicircular calculations: the residual equation carries puncture data, worldline information, and slow-time source terms that are absent from the present stationary scalar problem [27, 28].

The distinction is also important when comparing this scalar prototype with nonlinear Teukolsky calculations. In the gravitational problem the second-order source contains tetrad and metric-perturbation terms and must be combined with gauge completion and, for self-force applications, a singular-field subtraction [37–39]. Our source

is instead a smooth local cubic functional of one prescribed scalar mode. We therefore use the EMRI and second-order literature to delimit the interpretation of the calculation, not to identify the scalar coefficient with a gravitational self-force observable.

This work carries the same chain to the scalar Teukolsky equation in Kerr spacetime with spin weight $s = 0$, within the standard analytic black-hole perturbation framework [33]. The aim is to keep the observable definition and the flux logic of the Schwarzschild work unchanged while replacing the radial and angular equations by their Kerr counterparts. Rotation produces three changes. First, the horizon flux is proportional to $p = \omega - m\Omega_H$, which is the origin of superradiant sign changes [30, 31]. Second, spherical harmonics are replaced by spheroidal harmonics. Third, the covariant scalar source contains $\Sigma = r^2 + a^2 \cos^2 \theta$, so a single incident mode sources several target spheroidal channels.

The paper is organized in the same order as the Schwarzschild calculation, so that each Kerr step can be compared directly with its spherical counterpart. Section II defines the nonlinear scattering problem and the flux coefficients. Section III describes the pseudo-spectral homogeneous solver, the Green-function construction, the balance law, the accuracy diagnostics, and the low- and high-frequency diagnostics. Section IV gives the main frequency-domain numerical result: Kerr nonlinear transmission curves for $\ell = 0, 1, 2$, fitted resonance parameters, and high-frequency temperatures. The main physical frequency sweep is deliberately restricted to an axisymmetric Kerr sample, $a = 0.5$ and $m = 0$. This restriction makes the comparison with the Schwarzschild flux construction as direct as possible. Non-axisymmetric self-channel runs, including a bounded superradiant check, appear separately in Sec. IV; they are not folded into the axisymmetric production curves.

The contributions of this paper are threefold. First, we derive the Kerr fixed-incident normalization of the weak cubic response, including the horizon factor p and the Σ -weighted spheroidal source projection, so that the reported quantities are energy-flux derivatives rather than unnormalized Green amplitudes. Second, we implement the endpoint-inclusive two-domain Chebyshev construction and combine spectral tails, field-dependent radial residuals, Wronskian constancy, and flux balance into an explicit acceptance protocol. Third, we use that protocol to separate robust axisymmetric resonance and tail trends from cancellation-limited low-frequency reflected coefficients, while exposing the off-diagonal target-channel structure and a bounded superradiant sign change. The resulting claims are deliberately restricted to the prescribed scalar response problem defined above.

II. NONLINEAR SCATTERING PROBLEM IN KERR SPACETIME

In Boyer-Lindquist coordinates the Kerr metric is

$$ds^2 = - \left(1 - \frac{2Mr}{\Sigma} \right) dt^2 - \frac{4Mar \sin^2 \theta}{\Sigma} dt d\phi + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2 + \frac{A \sin^2 \theta}{\Sigma} d\phi^2, \quad (1)$$

where

$$\Sigma = r^2 + a^2 \cos^2 \theta, \quad \Delta = (r - r_+)(r - r_-), \quad (2)$$

$$A = (r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta. \quad (3)$$

The horizon radii and angular velocity are

$$r_{\pm} = M \pm \sqrt{M^2 - a^2}, \quad \Omega_H = \frac{a}{r_+^2 + a^2}. \quad (4)$$

We consider a weak cubic self-interaction

$$\square \Phi + \epsilon |\Phi|^2 \Phi = 0, \quad \epsilon \in \mathbb{R}, \quad |\epsilon| \ll 1, \quad (5)$$

In four spacetime dimensions the canonical scalar has dimensions $[\Phi] = M^{-1}$, so ϵ is dimensionless in the action convention below. For a field amplitude that is not absorbed into the unit-horizon normalization, a useful local control parameter is

$$\chi_{\text{nl}} = |\epsilon| M^2 \max_{\mathcal{D}} |\Phi^{(0)}|^2 \ll 1, \quad (6)$$

where $\mathcal{D}$ is the radial and angular region that dominates the cubic source projection. The reported $T^{(1)}$ and $R^{(1)}$ are coefficients in the derivative of the fixed-incident flux with respect to ϵ at $\epsilon = 0$; a finite-coupling application must additionally check that $|\epsilon T^{(1)}| \ll |T^{(0)}|$ and the analogous reflected-flux condition. We do not interpret the present frequency-domain calculation as a test of nonlinear saturation or of the expansion beyond this perturbative regime. This equation follows from the fixed-background scalar action

$$S_{\Phi} = \int d^4x \sqrt{-g} \left[-g^{\mu\nu} \nabla_{\mu} \Phi^* \nabla_{\nu} \Phi + \frac{\epsilon}{2} |\Phi|^4 \right]. \quad (7)$$

The stress tensor used below is the Hilbert stress tensor of the same effective field theory,

$$T_{\mu\nu} = - \frac{2}{\sqrt{-g}} \frac{\delta S_{\Phi}}{\delta g^{\mu\nu}}. \quad (8)$$

For later use, the explicit tensor and the associated stationary Killing energy flux are

$$T_{\mu\nu} = \nabla_{\mu} \Phi^* \nabla_{\nu} \Phi + \nabla_{\nu} \Phi^* \nabla_{\mu} \Phi + g_{\mu\nu} \left[-\nabla_{\alpha} \Phi^* \nabla^{\alpha} \Phi + \frac{\epsilon}{2} |\Phi|^4 \right], \quad (9)$$

$$j_E^{\mu} = -T^{\mu}_{\nu} \xi^{\nu} = -T^{\mu}_t, \quad \xi = \partial_t, \quad (10)$$

$$\mathcal{F}_E^{\text{out}}(r) = \int_0^{2\pi} d\phi \int_0^{\pi} d\theta \sqrt{-g} j_E^r, \quad (11)$$

$$\mathcal{F}_E^H = - \lim_{r \rightarrow r_+} \mathcal{F}_E^{\text{out}}(r). \quad (12)$$

The reversed horizon orientation in the last definition makes positive $\mathcal{F}_E^H$ correspond to absorption when $p > 0$. The quartic term is proportional to δ^r_t in T^r_t and therefore does not itself carry radial flux; the nonlinear boundary correction comes from the interference of the zeroth- and first-order kinetic terms. We now expand the field as

$$\Phi = \Phi^{(0)} + \epsilon\Phi^{(1)} + \mathcal{O}(\epsilon^2). \quad (13)$$

The zeroth-order incident wave is a single scalar spheroidal mode,

$$\Phi^{(0)} = e^{-i\omega t + im\phi} S_{\ell m}(\theta; a\omega) R_{\ell m\omega}(r), \quad (14)$$

with angular normalization

$$\int |S_{\ell m}|^2 d\Omega = 1. \quad (15)$$

The scalar spheroidal harmonic obeys

$$\left[\frac{1}{\sin\theta} \partial_\theta \sin\theta \partial_\theta + a^2 \omega^2 \cos^2\theta - \frac{m^2}{\sin^2\theta} + A_{\ell m} \right] S_{\ell m} = 0. \quad (16)$$

The radial Teukolsky equation for $s = 0$ is [40]

$$\mathcal{L}_{\ell m\omega} R \equiv \frac{d}{dr} \left(\Delta \frac{dR}{dr} \right) + \left(\frac{K^2}{\Delta} - \lambda_{\ell m\omega} \right) R = 0, \quad (17)$$

where

$$K = (r^2 + a^2)\omega - am, \quad \lambda_{\ell m\omega} = A_{\ell m} + a^2\omega^2 - 2am\omega. \quad (18)$$

The endpoint normalization used here can be related to the analytic MST solutions, which provide independent low-frequency connection formulae [32]. The numerical results below are assessed using internal spectral convergence and flux-balance diagnostics. The tortoise coordinate is defined by

$$\frac{dr_*}{dr} = \frac{r^2 + a^2}{\Delta}, \quad (19)$$

and the horizon wave number is

$$p = \omega - m\Omega_H. \quad (20)$$

The ingoing solution is normalized to unit amplitude at the future horizon,

$$R_{\ell m\omega}^{\text{in}} \sim e^{-ipr_*}, \quad r \rightarrow r_+, \quad (21)$$

and at infinity it has the form

$$R_{\ell m\omega}^{\text{in}} \sim B^{\text{inc}} \frac{e^{-i\omega r_*}}{r} + B^{\text{ref}} \frac{e^{+i\omega r_*}}{r}, \quad r \rightarrow \infty. \quad (22)$$

With this convention the linear flux coefficients are

$$T^{(0)} = \frac{p(r_+^2 + a^2)}{\omega |B^{\text{inc}}|^2}, \quad R^{(0)} = \frac{|B^{\text{ref}}|^2}{|B^{\text{inc}}|^2}. \quad (23)$$

For clarity, these coefficients are energy-flux ratios rather than merely amplitude ratios. Let $\xi = \partial_t$ be the stationary Killing field and let $T_{\mu\nu}$ be the scalar stress tensor defined in Eq. (8). The outward-oriented Killing energy current is $j_E^\mu = -T^\mu_t$, so at a surface of constant r we use $\mathcal{F}_E^{\text{out}} = \int j_E^r \sqrt{-g} d\theta d\phi$. When quoting absorption by the future horizon, the horizon boundary orientation is reversed, so that positive absorption corresponds to positive $\mathcal{F}_E^H = -\mathcal{F}_E^{\text{out}}|_{r=r_+}$ for $p > 0$. For a single Fourier mode, the energy current is the mode frequency times the radial Klein-Gordon current, up to one common positive normalization $\mathcal{N}$. Consequently,

$$\begin{aligned} \mathcal{F}_{E,\infty}^{\text{in}} &= \mathcal{N}\omega^2 |B^{\text{inc}}|^2, \\ \mathcal{F}_{E,\infty}^{\text{ref}} &= \mathcal{N}\omega^2 |B^{\text{ref}}|^2, \\ \mathcal{F}_{E,H} &= \mathcal{N}\omega p(r_+^2 + a^2). \end{aligned} \quad (24)$$

Their ratios give Eq. (23), including the sign of the horizon flux when $p < 0$. The self-interaction contributes a term proportional to δ^r_t in the mixed stress tensor and therefore does not add a radial boundary flux; at first order, the nonlinear flux correction is the interference term between $\Phi^{(0)}$ and $\Phi^{(1)}$ derived below.

For real ω the conserved radial current gives

$$T^{(0)} + R^{(0)} = 1. \quad (25)$$

At first order,

$$\square\Phi^{(1)} = -|\Phi^{(0)}|^2\Phi^{(0)}. \quad (26)$$

Since the source has the same (ω, m) dependence, we expand

$$\Phi^{(1)} = e^{-i\omega t + im\phi} \sum_{\ell' \geq |m|} S_{\ell' m}(\theta; a\omega) R_{\ell' | \ell m\omega}^{(1)}(r). \quad (27)$$

After multiplying by Σ and projecting onto a target spheroidal harmonic, the radial source is

$$\mathcal{L}_{\ell' m\omega} R_{\ell' | \ell m\omega}^{(1)} = -Q_{\ell' \ell m}(r), \quad (28)$$

where

$$Q_{\ell' \ell m}(r) = \left[r^2 C_{\ell' \ell m}^{(0)} + a^2 C_{\ell' \ell m}^{(2)} \right] |R_{\ell m\omega}^{\text{in}}|^2 R_{\ell m\omega}^{\text{in}}, \quad (29)$$

and

$$C_{\ell' \ell m}^{(j)} = \int d\Omega \cos^j \theta S_{\ell' m}^* |S_{\ell m}|^2 S_{\ell m}, \quad j = 0, 2. \quad (30)$$

The self channel $\ell' = \ell$ is the only target channel that interferes with the linear outgoing wave in the total energy flux at order ϵ .

III. METHODS

A. Pseudo-spectral method for homogeneous solutions

The homogeneous radial equation is solved in the compact coordinate

$$z = \frac{r_+}{r}, \quad z = 0 \text{ at infinity}, \quad z = 1 \text{ at the horizon.} \quad (31)$$

As in the Schwarzschild calculation, the oscillatory endpoint phases are peeled off before Chebyshev collocation. This endpoint-first organization is also consistent with independent spectral black-hole perturbation work, where the asymptotic behavior is factored before the Chebyshev representation is truncated [34]:

$$R_{\text{in}} = e^{-ipr_*} u_{\text{in}}(z), \quad (32)$$

$$R_{\text{down}} = \frac{e^{-i\omega r_*}}{r} u_{\text{down}}(z), \quad (33)$$

$$R_{\text{up}} = \frac{e^{+i\omega r_*}}{r} u_{\text{up}}(z). \quad (34)$$

The functions u_{in} , u_{down} , and u_{up} are smooth enough on their respective subdomains that the degenerate endpoint rows of the transformed ODE can be used directly as boundary regularity conditions. The outer and inner domains are matched at a finite radius r_m by continuity of R and dR/dr .

More explicitly, after the phase peeling the collocation equation has the form

$$B_2(z)u_{,zz} + B_1(z)u_{,z} + B_0(z)u = 0. \quad (35)$$

At the exact endpoint of each branch B_2 vanishes, so the endpoint row is the regular-singular relation $B_1 u_{,z} + B_0 u = 0$. The neighboring row is replaced by the normalization $u_{\text{down}}(0) = u_{\text{up}}(0) = 1$ on the outer domain or $u_{\text{in}}(1) = 1$ on the inner domain. Thus the horizon is included as an exact collocation endpoint; no horizon truncation is introduced. Each domain uses $N+1$ Chebyshev points and a dense complex linear solve. The reference production runs use $N = 240$ on both domains, $r_m = 40M$, Gauss order $q = 192$ for the compact inner integral, and relative tolerance 10^{-11} for the oscillatory phase-channel tail. The low-frequency runs use the same endpoint formulation with the sinh map described above; all reported results are obtained in IEEE double precision and are checked by the order, matching-radius, quadrature, Wronskian, and flux scans reported below.

The angular eigenvalue is computed by diagonalizing the truncated spherical harmonic representation of $\ell(\ell+1) - (a\omega)^2 \cos^2 \theta$, retaining sixteen modes above the requested ℓ . The source projectors are evaluated with normalized spheroidal harmonics and Gauss-Legendre quadrature. For the radial Green integrals, the compact inner part is integrated in z with the Jacobian $dr = r_+/z^2 dz$, while the outer part is decomposed into

its finite phase channels in r_* before oscillatory quadrature.

The direct Teukolsky potential is long ranged, so an alternative numerical route is to transform the equation to a short-ranged generalized Sasaki-Nakamura equation [35, 36]. We retain the direct Teukolsky variable here because it keeps the covariant Σ -weighted source and the endpoint normalization in the same representation. The GSN formulation is consequently a useful independent route for future high-frequency or extended-source checks, but it is not used to define any of the production values reported below.

The scattering relation used in the code is the Kerr analogue of the Schwarzschild S-matrix relation,

$$R_{\text{in}} = B^{\text{inc}} R_{\text{down}} + B^{\text{ref}} R_{\text{up}}. \quad (36)$$

Solving this two-by-two matching system gives B^{inc} and B^{ref} and therefore the linear coefficients in Eq. (23). The residual $|T^{(0)} + R^{(0)} - 1|$ is used as the first check of the homogeneous solve.

B. Nonlinear scattering coefficients

1. The nonlinear balance law

Let $R_{\text{up}}^{(\ell)}$ denote the target-channel homogeneous solution that is purely outgoing at infinity. The Green-function solution in the self channel has outgoing amplitude at infinity and ingoing amplitude at the horizon. The conserved Wronskian is

$$W_\ell = \Delta \left(R_{\ell m \omega}^{\text{in}} \frac{dR_{\ell m \omega}^{\text{up}}}{dr} - R_{\ell m \omega}^{\text{up}} \frac{dR_{\ell m \omega}^{\text{in}}}{dr} \right), \quad (37)$$

which is independent of r for two homogeneous solutions of Eq. (17). The Green-function amplitudes are

$$A_{\text{ref}}^{(1)} = -\frac{1}{W_\ell} \int_{r_+}^{\infty} Q_{\ell m}(r) R_{\ell m \omega}^{\text{in}}(r) dr, \quad (38)$$

$$A_H^{(1)} = -\frac{1}{W_\ell} \int_{r_+}^{\infty} Q_{\ell m}(r) R_{\ell m \omega}^{\text{up}}(r) dr, \quad (39)$$

where the minus sign follows from $\mathcal{L}_{\ell m \omega} R^{(1)} = -Q_{\ell m}$. In the numerical implementation the Wronskian is evaluated from the matched homogeneous solutions and checked against its constancy. The amplitudes $A_{\text{ref}}^{(1)}$ and $A_H^{(1)}$ are quoted in the unit-horizon normalization of Eq. (21). To form scattering coefficients one must instead hold the incident wave fixed. If the zeroth-order field is rescaled by $\alpha = 1/B^{\text{inc}}$, the cubic source scales as $|\alpha|^2 \alpha$. Therefore the fixed-incident first-order boundary amplitudes are

$$A_{\text{ref,fix}}^{(1)} = \frac{A_{\text{ref}}^{(1)}}{|B^{\text{inc}}|^2 B^{\text{inc}}}, \quad A_{H,\text{fix}}^{(1)} = \frac{A_H^{(1)}}{|B^{\text{inc}}|^2 B^{\text{inc}}}. \quad (40)$$

Expanding the horizon and infinity fluxes of the fixed-incident field then gives

$$T^{(1)} = \frac{2p(r_+^2 + a^2)}{\omega |B^{\text{inc}}|^4} \text{Re } A_H^{(1)}, \quad (41)$$

$$R^{(1)} = \frac{2 \text{Re}(\overline{B^{\text{ref}}} A_{\text{ref}}^{(1)})}{|B^{\text{inc}}|^4}. \quad (42)$$

Equivalently, with the incident energy flux held fixed,

$$T^{(1)} = \frac{1}{\mathcal{F}_{E,\infty}^{\text{in}}} \left. \frac{\partial \mathcal{F}_E^H}{\partial \epsilon} \right|_{\epsilon=0}, \quad R^{(1)} = \frac{1}{\mathcal{F}_{E,\infty}^{\text{in}}} \left. \frac{\partial \mathcal{F}_{E,\infty}^{\text{ref}}}{\partial \epsilon} \right|_{\epsilon=0}. \quad (43)$$

Thus the quantities plotted below are normalized changes of the horizon and infinity energy fluxes, rather than changes of radial-wave amplitudes alone. The current conservation identity gives the first-order balance law

$$T^{(1)} + R^{(1)} = 0. \quad (44)$$

The local reason for this identity is visible directly in the radial equation. Let $J^{(1)}$ be the term linear in ϵ in the radial Klein–Gordon current. The cross-current identity gives

$$\frac{dJ^{(1)}}{dr} = -\text{Im} \left[(R_{\ell m \omega}^{\text{in}})^* Q_{\ell m}(r) \right] = 0, \quad (45)$$

because the self-channel projectors are real for a real spheroidal parameter and $Q_{\ell m} \propto |R^{\text{in}}|^2 R^{\text{in}}$. Thus the smooth cubic source changes the two boundary currents through interference, but does not inject net Killing energy into the stationary mode at this perturbative order. Equation (44) is the boundary form of this local identity, while the field residual and spectral tail remain independent discretization diagnostics. Numerically this equation is the most useful global precision diagnostic, because it tests the homogeneous normalization, the nonlinear Green integral, and the flux normalization simultaneously. The fixed-incident normalization and the appearance of the $|B^{\text{inc}}|^4$ denominators are derived explicitly in Appendix A.

2. Numerical accuracy diagnostics

The accuracy diagnostics used below are internal to the spectral calculation. Let the homogeneous equation be written at a collocation point as

$$\mathcal{R}_R = \Delta R'' + \Delta' R' + \left(\frac{K^2}{\Delta} - \lambda_{\ell m \omega} \right) R. \quad (46)$$

The reported relative residual is

$$\delta_R = \frac{|\mathcal{R}_R|}{|\Delta R''| + |\Delta' R'| + |(K^2/\Delta - \lambda_{\ell m \omega}) R|}. \quad (47)$$

Thus the denominator contains the actual local field amplitude, not only the differential-equation coefficients.

This makes the residual insensitive to an overall rescaling of the homogeneous solution. The original radial residual is evaluated on the interior collocation nodes, $0 < z < 1$; the degenerate endpoint rows are reserved for the boundary regularity conditions and are not interpreted as ordinary pointwise ODE residuals. We also monitor the Chebyshev tail

$$\eta_{\text{tail}} = \frac{\max_{N-15 \leq k \leq N} |c_k|}{\max_{0 \leq k \leq N} |c_k|}, \quad (48)$$

where c_k are the Chebyshev coefficients of the phase-peeled unknown on each subdomain. Finally, Eqs. (25) and (44) test the physical flux balance after the homogeneous matching and Green integral have both been performed.

The production data below retain the self channel $\ell' = \ell$ when quoting the first-order scattering coefficients. This is not an angular truncation of the first-order field: the cubic Kerr source in Eq. (29) does generate off-diagonal target channels. However, at order ϵ the total energy flux contains only interference between $\Phi^{(0)}$ and $\Phi^{(1)}$. Orthogonality of the spheroidal harmonics then removes every target channel except $\ell' = \ell$ from the flux correction. Off-diagonal channels are therefore required for the full first-order waveform, but not for the self-channel coefficients $T_\ell^{(1)}$ and $R_\ell^{(1)}$ reported here.

3. Low-frequency behavior

The Schwarzschild reduced radial equation admits a clean analytic low-frequency scaling argument. In the Kerr Teukolsky variable the far-zone radial normalization and the covariant source weight $r^2 C_{\ell m}^{(0)} + a^2 C_{\ell m}^{(2)}$ change the practical low-frequency behavior measured by the fixed-incident flux coefficients. Rather than importing the Schwarzschild scaling unchanged, we perform a direct Kerr low-frequency experiment for $a = 0.5$, $m = 0$, and $\ell = 0, 1, 2$.

The result is shown in Fig. 1. A log-log fit over the first four available low-frequency points gives the local slopes

$$T_\ell^{(1)} \sim \omega^{\alpha_\ell}, \quad \alpha_0 \simeq 2.50, \quad \alpha_1 \simeq 5.48, \quad \alpha_2 \simeq 8.74. \quad (49)$$

The corresponding linear slopes are approximately 2.34, 5.25, and 8.52. Thus, in the present Kerr normalization the nonlinear transmission correction follows the same centrifugal-barrier hierarchy as the linear transmission coefficient in the accessible low-frequency window. These are finite-window fitted exponents rather than analytic asymptotic indices: repeating the fit with three, four, and five points gives the ranges in Table I. The reflected coefficient is harder to extract at the lowest frequencies because the infinity-side interference integral contains a large cancellation; this is why Eq. (44) is reported alongside the coefficients.

TABLE I. Sensitivity of the low-frequency log-log slopes to the number of points retained in the fit. The baseline column uses the four-point fit quoted in the text; the ranges use three, four, and five points. Here the superscript (0) labels the linear transmission coefficient and (1) labels the first-order nonlinear coefficient. The last two columns give the largest log-space RMS residual over those fits.

ℓ	baseline ($\alpha_\ell^{(0)}, \alpha_\ell^{(1)}$)	range of $\alpha_\ell^{(0)}$	range of $\alpha_\ell^{(1)}$	$\max \sigma_{\log}^{(0)}$	$\max \sigma_{\log}^{(1)}$
0	(2.34, 2.50)	[2.31, 2.37]	[2.46, 2.50]	3.44×10^{-2}	3.33×10^{-2}
1	(5.25, 5.48)	[4.88, 5.51]	[5.03, 5.80]	2.92×10^{-1}	3.66×10^{-1}
2	(8.52, 8.74)	[7.92, 9.05]	[8.07, 9.35]	4.48×10^{-1}	5.10×10^{-1}




FIG. 1. Low-frequency Kerr diagnostics for $a = 0.5$, $m = 0$, and $\ell = 0, 1, 2$. Left: linear transmission coefficients. Right: self-channel nonlinear transmission coefficients. The fitted low-frequency slopes show that the Kerr fixed-incident coefficients decrease with the same centrifugal hierarchy in $T^{(0)}$ and $T^{(1)}$.

4. High-frequency behavior

At high frequencies the phase-peeled homogeneous amplitudes become slowly varying, while the nonlinear Green integral is controlled by rapidly oscillatory tails. The Schwarzschild calculation finds an exponential tail with a fitted temperature close to the Hawking temperature [41]. For Kerr we do not assume the Schwarzschild asymptotic integral unchanged, because the Teukolsky radial normalization and the Σ -weighted source modify the integrand. Instead we test the same one-parameter tail model,

$$T_\ell^{(1)}(\omega) \simeq A_\ell \exp \left[-\frac{\omega}{T_{\text{fit}}} \right], \quad \omega > \text{Re } \omega_{\ell 0}. \quad (50)$$

For Kerr

$$T_H = \frac{r_+ - r_-}{4\pi(r_+^2 + a^2)}. \quad (51)$$

For $a = 0.5$ the fits in Sec. IV give $\pi M T_{\text{fit}} = 0.112$, 0.104, and 0.100 for $\ell = 0, 1, 2$, close to $\pi M T_H = 0.1160$. These values should be read as finite-window effective temperatures rather than as a proof of an exact high-frequency theorem.

IV. RESULTS

Our main result on the Kerr nonlinear scattering coefficient is shown in Fig. 2. This figure is the direct Kerr analogue of the main Schwarzschild figure: it plots $T_\ell^{(1)}$ for $\ell = 0, 1, 2$, uses a linear scale to display the resonance peaks, and uses a logarithmic scale to display the high-frequency tail. The dashed lines mark the real parts of the fundamental scalar Kerr quasinormal modes computed with the local **qnm** package [42] for $s = 0$, $a = 0.5$, $m = 0$, and $n = 0$.

Three features are robust in the accepted data. First, in the axisymmetric Kerr cases studied here, $T_\ell^{(1)}$ is positive at every accepted sampled frequency in the present flux convention. For positive coupling in Eq. (5), the nonlinear correction therefore increases the absorbed flux and, by Eq. (44), decreases the reflected flux at the same order. This sign statement refers to the coefficient convention of Eqs. (41) and (42); it should not be compared to a reduced Schwarzschild field convention without translating the flux normalization.

Second, the peaks of $T_\ell^{(1)}$ lie on the frequency scale set by the real parts of the corresponding fundamental Kerr scalar QNMs [7, 42]. The fitted peak positions are $M\omega_{\text{peak}} = 0.1299, 0.3100$, and 0.5022 for $\ell = 0, 1, 2$, while

the QNM real parts are 0.1124, 0.2979, and 0.4920. The corresponding fractional offsets are approximately 16%, 4.1%, and 2.1%. To make the comparison quantitative without identifying the flux with a pole, we fit the selected samples to the four-parameter profile

$$T_\ell^{(1)}(\omega) = T_{\text{off}} + \frac{A\gamma^2}{(M\omega - \omega_p)^2 + \gamma^2}. \quad (52)$$

Here ω_p is the fitted center in units $M = 1$, and γ is the half-width parameter of the interpolation. This is the Kerr version of the Breit-Wigner-type resonance observed in the Schwarzschild nonlinear coefficients, not a claim that the flux itself is a meromorphic response. The coefficient of determination of the three fits is $R^2 = 0.9974, 0.9979$, and 0.9991 for $\ell = 0, 1, 2$, respectively. The comparison is qualitative rather than a precision QNM determination: the nonlinear coefficient is a real-frequency flux observable, whereas the QNM frequencies are poles of the analytically continued response. The quoted ω_{peak} and γ are parameters of the numerical Lorentzian interpolation used to locate a broad feature; they are not a claim that the nonlinear flux has been analytically continued to a QNM pole. The fit uses the sampled points satisfying $T_\ell^{(1)} > 0.2T_{\ell, \text{max}}^{(1)}$; the relative RMS residuals of these peak fits over the selected points are 1.36%, 1.23%, and 0.57% for $\ell = 0, 1, 2$, respectively. These residuals quantify the descriptive quality of the interpolation, not an error bar on the QNM frequencies. As a fit-window sensitivity check, we repeat the selection with thresholds from 0.15 to 0.30 of the sampled maximum, retaining at least five points in every fit. The resulting fitted centers vary by at most 7.1×10^{-5} , 3.2×10^{-4} , and 1.3×10^{-5} for $\ell = 0, 1, 2$, respectively, whereas the corresponding half-widths vary by 5.3×10^{-3} , 8.8×10^{-3} , and 4.5×10^{-4} . The center locations are therefore stable at the resolution of the sweep, while the widths should be regarded as descriptive fit parameters. The complete threshold scan is included in the reproducibility data.

Third, at high frequency the accepted tail points are well fit by the Boltzmann form Eq. (50). The fitted temperatures differ from the Kerr Hawking temperature by 3.2%, 10.1%, and 13.6% for $\ell = 0, 1, 2$, respectively. These are effective finite-window fit parameters, not an analytic derivation of a thermal spectrum. At low frequency the Kerr fixed-incident coefficients decrease with frequency in the accessible window. This differs from simply copying the Schwarzschild reduced-field statement and is therefore reported as a Kerr numerical result rather than assumed analytically.

The exponential fits use unweighted least squares for $\log T_\ell^{(1)}$ versus $M\omega$ over the windows $[0.18, 0.60]$, $[0.38, 0.85]$, and $[0.62, 1.00]$ for $\ell = 0, 1, 2$, respectively. The corresponding root-mean-square residuals in log space are 4.5×10^{-2} , 4.9×10^{-2} , and 3.8×10^{-2} . These window-dependent residuals quantify the quality of the finite-window fit; they are not estimates of a systematic error outside the computed frequency range.

TABLE II. Fitting parameters of the Kerr nonlinear transmission resonance and the high-frequency Boltzmann tail. The QNM frequencies are scalar Kerr fundamental modes with $s = 0$, $a = 0.5$, $m = 0$, and $n = 0$. All frequencies are reported in units $M = 1$.

ℓ	ω_p	γ	$\Re\omega_Q$	$ \Im\omega_Q $	πT_{fit}	ΔT
0	0.1299	0.0779	0.1124	0.1022	0.1123	3.25%
1	0.3100	0.0597	0.2979	0.0954	0.1043	10.1%
2	0.5022	0.0611	0.4920	0.0946	0.1003	13.6%

Because the effective temperature is a fit diagnostic rather than an analytic observable, we also repeat the tail fit after shifting either edge of each window while retaining at least three positive data points. The resulting window ranges and the largest logarithmic fit residual are given in Table III. The fitted values vary only at the percent level within these local windows, while their absolute offsets from T_H remain consistent with the finite-window qualification above.

For the accepted subset of the sweep, the largest first-order balance residual is 4.3×10^{-12} , 2.2×10^{-12} , and 1.2×10^{-12} for $\ell = 0, 1, 2$, respectively. The full sweep also contains rejected lowest-frequency points with residuals as large as 3.9×10^{-5} and 1.2×10^{-4} for $\ell = 1, 2$; these are caused by cancellation in the reflected interference term and are not used for precision claims. The transmitted coefficients remain smooth at those frequencies, so we retain them for the low-frequency slope measurement but explicitly separate that trend from an accuracy claim for $R^{(1)}$.

The change from the former production resolution to the active refined resolution is quantified separately in Table V. At the representative peak points the refined calculation changes the transmission coefficient by at most 3.2×10^{-5} and preserves the first-order flux balance at the 10^{-12} level. We therefore use the refined curves for the physical discussion below; the lower-resolution values are reported to make the resolution choice auditable rather than folded into an unreported systematic error.

We additionally test the self channel with $\ell = m = 2$ at both signs of the Kerr horizon frequency p and for a rapidly rotating case. Table VI records finite-resolution changes under N , r_m , and q ; these are not rigorous error bounds. The horizon amplitude is less sensitive than the reflected amplitude, while the Wronskian error remains below 6×10^{-16} . We therefore use flux balance, rather than one amplitude difference, as the acceptance criterion.

Local field diagnostic.

At the representative peak in Fig. 3, the field-dependent radial residual has maximum, 95th-percentile, and median values 2.5×10^{-10} , 3.3×10^{-12} , and 1.3×10^{-13} . The largest normalized Chebyshev tail is 2.8×10^{-14} and the Wronskian mismatch is below 3×10^{-16} . These local diagnostics are reported separately from the global flux-balance residual.




FIG. 2. Nonlinear Kerr transmission coefficients $T_\ell^{(1)}$ for $a = 0.5$, $m = 0$, and $\ell = 0, 1, 2$. The left panel uses a linear scale to show the Breit-Wigner-type peaks. The right panel uses a logarithmic scale to show the exponential high-frequency tails; dotted segments show the finite-window fits used in Table II. Dashed vertical lines mark the real parts of the fundamental scalar Kerr QNMs $\text{Re } \omega_{\ell 0}$.

TABLE III. Sensitivity of the finite-window effective temperatures to the fit window. The baseline window is the one used in Table II; the range is obtained from three nearby edge-shifted windows. The final column is the largest absolute fractional offset from the Kerr Hawking temperature among the four windows.

ℓ	baseline $[M\omega_{\min}, M\omega_{\max}]$	$\pi T_{\text{fit}}^{\text{base}}$	range of πT_{fit}	$\max \sigma_{\log}$	$\max T_{\text{fit}}/T_H - 1 $
0	[0.18, 0.60]	0.1123	[0.1112, 0.1131]	4.83×10^{-2}	4.18%
1	[0.38, 0.85]	0.1043	[0.1034, 0.1057]	4.87×10^{-2}	10.90%
2	[0.62, 1.00]	0.1003	[0.0994, 0.1012]	3.82×10^{-2}	14.37%

A. Target-channel structure

The spheroidal source projection is a genuinely Kerr-specific part of the calculation. To expose it directly, we evaluated the target channels $\ell' = 2, \dots, 6$ for the representative mode $(a, \ell, m, M\omega) = (0.5, 2, 2, 0.30)$ using $N = 288$, quadrature order $q = 224$, and Fourier-tail tolerance 10^{-11} . The amplitudes in Table VII use the unit-horizon normalization of Eq. (21); they are not fixed-incident flux coefficients. Odd target channels vanish by the even-parity angular projectors in Eq. (29), while the even channels span several orders of magnitude in the outgoing Green amplitude.

The channel scan is converged for the dominant reflected amplitudes: changing N from 256 to 288 changes $A_{\text{ref}}^{(1)}$ by 2.5×10^{-7} in the self channel and by 6.3×10^{-7} in the $\ell' = 4$ channel. The corresponding $\ell' = 4$ horizon amplitude changes by 7.1×10^{-3} , whereas the $\ell' = 6$ horizon amplitude is close to the double-precision cancellation floor and changes by order unity. For the displayed $\ell' = 6$ row, the independent diagnostic $(|B_{\text{inc}}|^2 + |B_{\text{ref}}|^2)/|B_{\text{inc}}|^2 - |B_{\text{ref}}|^2$ is 3.27×10^{13} ; this is the relevant conditioning indicator for the horizon-normalized continuation. We therefore use this experiment to establish the target-channel structure and the dominant out-

going amplitudes, not to claim uniform high precision for every off-diagonal horizon coefficient. Since off-diagonal channels do not interfere with the zeroth-order field, their energy flux first enters at $\mathcal{O}(\epsilon^2)$.

B. Non-axisymmetric superradiant check

The axisymmetric production sweep is useful for a controlled comparison with the Schwarzschild calculation, but it removes the most distinctive rotating black-hole effect because $p = \omega$ when $m = 0$. We therefore performed a bounded check with $a = 0.5$ and $\ell = m = 2$, using $N = 320$ on both spectral subdomains, quadrature order $q = 320$, and $r_m = 40M$. The threshold is $m\Omega_H = 0.2679491924$. The full ten-point scan is retained in the tracked CSV file; Table VIII lists only points satisfying the same first-order balance gate $|T^{(1)} + R^{(1)}| \leq 10^{-8}$ used for the production data. The rejected points are $M\omega = 0.20, 0.22, 0.255, 0.265, 0.28$; they lie in the near-threshold portion of the scan and are limited by cancellation in the reflected interference term, rather than by a physical discontinuity.

Figure 4 shows the signed linear and first-order horizon flux coefficients. On the accepted points below the threshold, both $T^{(0)}$ and $T^{(1)}$ are negative, while they are

TABLE IV. Numerical quality of the axisymmetric Kerr frequency sweeps. Here $\delta_0 = |T^{(0)} + R^{(0)} - 1|$ and $\delta_1 = |T^{(1)} + R^{(1)}|$. A point is counted as accepted when $\delta_1 \leq 10^{-8}$. All active frequency-sweep points use the refined resolution $N = 240$ and quadrature order $q = 192$; the earlier $N = 224$, $q = 160$ sweep is retained only as the production-resolution audit in Table V. The final column is the root-mean-square residual of the logarithmic high-frequency tail fit.

ℓ	points	$\max \delta_0$	accepted	$\max \delta_1^{\text{acc}}$	σ_{tail}
0	16	2.2×10^{-11}	16/16	4.3×10^{-12}	4.5×10^{-2}
1	16	1.9×10^{-11}	14/16	2.2×10^{-12}	4.9×10^{-2}
2	15	3.8×10^{-11}	12/15	1.2×10^{-12}	3.8×10^{-2}

TABLE V. Production-resolution audit for the axisymmetric Kerr calculation. The entries compare the former production configuration $(N, q) = (224, 160)$ with the active reference configuration $(N, q) = (240, 192)$ at fixed $r_m = 40$. Each entry in the first four columns is $|X_{240,192} - X_{224,160}|/|X_{240,192}|$ for the indicated Green-function amplitude or flux coefficient; the final column is the absolute first-order balance residual of the reference run.

case	ΔA_{ref}	ΔA_H	$\Delta T^{(1)}$	$\Delta R^{(1)}$	$ \delta_1 _{\text{ref}}$
$\ell = 0$, $M\omega = .125$	4.5×10^{-7}	7.8×10^{-7}	2.3×10^{-6}	2.3×10^{-6}	2.7×10^{-12}
$\ell = 1$, $M\omega = .34$	5.9×10^{-6}	3.3×10^{-6}	1.5×10^{-5}	1.5×10^{-5}	2.2×10^{-12}
$\ell = 2$, $M\omega = .52$	9.6×10^{-6}	7.6×10^{-6}	3.1×10^{-5}	3.1×10^{-5}	1.2×10^{-12}

positive on the accepted points above it. Thus the scalar Kerr calculation recovers the expected signed-flux change across the superradiant boundary in this representative mode. This is a validation of the horizon-frequency normalization, not a systematic survey over spin, m , or spheroidal target channels.

C. Kerr-to-Schwarzschild continuity

As a limiting-case check independent of external reference calculations, we hold $(\ell, m, M\omega) = (2, 0, 0.52)$ fixed and sweep the dimensionless spin from $a/M = 0$ to 0.5. The scalar spheroidal eigenvalue, the covariant source projection, the scattering amplitudes, and both flux coefficients are recomputed at every spin. At $a = 0$ the angular basis becomes spherical and the endpoint formulation reduces to the Schwarzschild spectral route; no radial endpoint is removed or replaced by a finite horizon cutoff. Figure 5 shows that $T^{(1)}$ and $-R^{(1)}$ vary smoothly from the $a = 0$ value, while Table IX records the balance and Wronskian diagnostics for the same points. The largest absolute first-order balance residual in this scan is 1.4×10^{-12} and the largest linear balance residual is below 1.2×10^{-11} . This experiment is a continuity and implementation check; it is not used as an external reference result or as a claim that the Kerr and Schwarzschild coefficient conventions are identical without the conversion in Appendix A.

V. DISCUSSION AND OUTLOOK

We have formulated the weakly nonlinear scattering problem for a scalar field on Kerr in a way that keeps the

observable content of the Schwarzschild Green-function calculation unchanged. The linear solution is normalized at the horizon, the incident normalization is restored through B^{inc} , and the nonlinear corrections are read from the interference terms in the physical energy fluxes. In this form the two useful balance laws, $T^{(0)} + R^{(0)} = 1$ and $T^{(1)} + R^{(1)} = 0$, remain the main global checks of the calculation. The extension is not a formal replacement of the Schwarzschild radial variable by a Kerr one: the horizon frequency $p = \omega - m\Omega_H$, spheroidal angular basis, and Σ -weighted nonlinear source all enter the normalization of the observable coefficients.

For the axisymmetric Kerr sample $a = 0.5$, $m = 0$, and $\ell = 0, 1, 2$, the present numerical diagnostics support the following qualitative picture. The nonlinear transmission coefficient develops a single dominant Breit-Wigner-type peak for each ℓ , with the peak position lying on the frequency scale of the corresponding scalar Kerr quasinormal mode. The high-frequency tail is well fit by an exponential form in the computed window, and the fitted temperature is of the same order as the Kerr Hawking temperature. At low frequency, however, the fixed-incident Teukolsky normalization and the Kerr source weight prevent a direct transcription of the Schwarzschild reduced-variable scaling; the low-frequency behavior is therefore treated here as a numerical Kerr result.

The present data also identify the current limiting numerical effect. The transmitted nonlinear coefficient is smooth across the displayed low-frequency windows, while the reflected coefficient can suffer large cancellations in the interference integral. This is why the lowest-frequency $\ell = 1, 2$ points are used for the scaling trend but are not counted as high-precision reflected-flux points in Table IV. The numerical evidence is therefore strongest for the resonance region, the high-frequency tail, and the

TABLE VI. Method-level convergence stress tests for the scalar Kerr Green integrals. Each entry is the maximum relative change in the indicated amplitude over the non-reference runs. The N scan compares $N = 224, 288$ with $N = 256$; the matching scan compares the listed alternatives with the reference r_m ; and the quadrature scan compares lower orders with $q = 192$. These non-axisymmetric self-channel tests are validation data for the numerical method and are not included in the axisymmetric production curves.

case	$\Delta A_{\text{ref}}[N]$	$\Delta A_H[N]$	$\Delta A_{\text{ref}}[r_m]$	$\Delta A_H[r_m]$	$\Delta A_{\text{ref}}[q]$	$\Delta A_H[q]$	$\max \delta_W$
$a = .5, M\omega = .24$	2.1×10^{-5}	1.1×10^{-7}	3.3×10^{-5}	5.4×10^{-8}	2.8×10^{-8}	2.5×10^{-8}	5.9×10^{-16}
$a = .5, M\omega = .30$	3.8×10^{-5}	2.8×10^{-8}	2.0×10^{-5}	4.4×10^{-8}	1.8×10^{-6}	1.7×10^{-6}	3.7×10^{-16}
$a = .9, M\omega = .58$	1.5×10^{-6}	1.2×10^{-8}	1.7×10^{-6}	9.4×10^{-9}	1.4×10^{-10}	5.1×10^{-7}	2.9×10^{-16}




FIG. 3. Internal numerical accuracy diagnostics for the axisymmetric Kerr calculation. Panels (a) and (b) use the representative peak point $a = 0.5$, $m = 0$, $\ell = 2$, and $M\omega = 0.52$ with $N = 240$ and quadrature order $q = 192$. Panel (a) shows the normalized Chebyshev coefficients of the phase-peeled homogeneous solutions on the outer and inner subdomains. Panel (b) shows the relative residual (47) after substituting the numerical solution back into the original radial Teukolsky equation. Panel (c) shows the first-order flux-balance residual over the full axisymmetric frequency sweep. The dashed line marks the acceptance threshold 10^{-8} used in Table IV.

transmitted low-frequency scaling; it is weaker for the individual low-frequency reflected coefficient before using the balance law. The representative target-channel experiment in Table VII demonstrates the matrix structure generated by the Kerr source projection and shows that the dominant reflected off-diagonal amplitudes can be resolved at the stated spectral order. A broader scan over (a, ℓ, m, ω) , uniform acceptance gates for off-diagonal horizon amplitudes, and a systematic study of the superradiant nonlinear flux remain future work.

The spin-limit audit supplies an additional implementation-level check. We fix $(\ell, m) = (2, 0)$ and $M\omega = 0.52$; the first-order coefficients change smoothly from the $a = 0$ endpoint to the production value at $a = 0.5$, while the two flux-balance laws remain satisfied. This continuity test does not replace a convention-matched Schwarzschild comparison, but it rules out a discontinuity introduced solely by the Kerr tortoise coordinate, spheroidal projection, or endpoint matching.

A. Scope and limitations

The conclusions above have a deliberately narrow interpretation. The calculation is a fixed-background scalar response problem with a prescribed cubic interaction; it is not a second-order metric perturbation, a gravitational self-force calculation, or an EMRI waveform model. The main frequency sweep is axisymmetric. The separate $m = 2$ check tests one narrow superradiant crossing but does not provide a systematic survey of superradiant sign changes. The target-channel experiment establishes the spheroidal coupling structure and the dominant reflected amplitudes, but the $\ell' = 6$ horizon coefficient is conditioning limited in double precision. The QNM comparison uses real-frequency peak locations only, and the finite-window exponential fits should not be interpreted as QNM residues or as a proof of Hawking thermality. Extending the result to generic Kerr orbits or to gravitational perturbations requires, respectively, source-harmonic and orbit-evolution modules or metric reconstruction, gauge completion, and regularization; none of these are hidden in the present scalar result.

TABLE VII. Representative Kerr target-channel matrix for an incident $(a, \ell, m, M\omega) = (0.5, 2, 2, 0.30)$ mode. The angular entries are absolute values of the two source projectors in Eq. (29); the Green amplitudes are unit-horizon first-order amplitudes. The odd channels vanish within the displayed precision.

ℓ'	$ C_{\ell'\ell m}^{(0)} $	$ C_{\ell'\ell m}^{(2)} $	$ A_{\text{ref}, \ell' \ell m}^{(1)} $	$ A_{H, \ell' \ell m}^{(1)} $
2	1.136×10^{-1}	1.034×10^{-2}	1.137×10^6	3.804×10^3
3	0	0	0	0
4	3.578×10^{-2}	5.513×10^{-3}	1.362×10^4	1.319
5	0	0	0	0
6	5.341×10^{-3}	5.364×10^{-3}	2.265×10^3	5.383×10^{-7}

Non-axisymmetric scalar Kerr flux check: $a = 0.5$, $\ell = m = 2$




FIG. 4. Bounded non-axisymmetric scalar Kerr flux check for $a = 0.5$ and $\ell = m = 2$. The dashed line marks $M\omega = m\Omega_H = 0.2679491924$. Filled circles satisfy the first-order balance gate; crosses mark scan points retained in the CSV file but rejected because of cancellation. The signed coefficients change sign across the superradiant threshold on the accepted points.

DATA AND CODE AVAILABILITY

The numerical data used in Figs. 1–5 and Tables I, II, III, IV, V, VI, VII, VIII, and IX are included in the project data directory, “results/”. The figures are generated by the scripts in “scripts/”, and the exact commands and tracked inputs are listed in “docs/prd/README.md”. The source code and frozen inputs are maintained at <https://github.com/ljq2088/KerrScattering>. The numerical checker compares the rounded entries in Tables II–VII, the spin-limit audit, the production-resolution audit, and the low-frequency slopes with the tracked CSV files. The artifact manifest records file sizes and SHA-256 hashes for the figure and CSV inputs, including the target-channel values in Table VII; a human-readable snapshot is provided in “docs/prd/reproducibility_report.md”. No external experimental data are used. In the CSV files, “status=ok” records a completed solver call; the manuscript acceptance decision is a separate postprocessing predicate on the recorded absolute first-order balance residual, $|T^{(1)} + R^{(1)}|$.

Appendix A: Flux normalizations

1. Projected Kerr source

The scalar wave equation separates only after multiplying by $\Sigma = r^2 + a^2 \cos^2 \theta$. At first order this gives

$$\Sigma \square \Phi^{(1)} = -\Sigma |\Phi^{(0)}|^2 \Phi^{(0)}. \quad (\text{A1})$$

Using

$$\begin{aligned} \Phi^{(0)} &= e^{-i\omega t + im\phi} S_{\ell m} R_{\ell m \omega}, \\ \Phi^{(1)} &= e^{-i\omega t + im\phi} \sum_{\ell'} S_{\ell' m} R_{\ell' | \ell m \omega}^{(1)}, \end{aligned} \quad (\text{A2})$$

and projecting with $S_{\ell' m}^*$ gives

$$\mathcal{L}_{\ell' m \omega} R_{\ell' | \ell m \omega}^{(1)} = - \left(r^2 C_{\ell' \ell m}^{(0)} + a^2 C_{\ell' \ell m}^{(2)} \right) |R_{\ell m \omega}^{\text{in}}|^2 R_{\ell m \omega}^{\text{in}}. \quad (\text{A3})$$

Here

$$C_{\ell' \ell m}^{(j)} = \int d\Omega \cos^j \theta S_{\ell' m}^* |S_{\ell m}|^2 S_{\ell m}, \quad j = 0, 2. \quad (\text{A4})$$

TABLE VIII. Accepted points in the bounded non-axisymmetric superradiant check. The entries are signed energy-flux coefficients for $a = 0.5$ and $\ell = m = 2$; $p = \omega - m\Omega_H$. The omitted points in the ten-point scan fail the 10^{-8} first-order balance gate.

$M\omega$	p	$T^{(0)}$	$T^{(1)}$	$ T^{(1)} + R^{(1)} $
0.18	-8.795×10^{-2}	-1.157×10^{-6}	-9.742×10^{-8}	4.788×10^{-13}
0.24	-2.795×10^{-2}	-2.373×10^{-6}	-2.153×10^{-7}	1.261×10^{-12}
0.30	3.205×10^{-2}	1.595×10^{-5}	1.582×10^{-6}	9.240×10^{-13}
0.35	8.205×10^{-2}	1.804×10^{-4}	1.960×10^{-5}	1.598×10^{-12}
0.45	1.821×10^{-1}	8.298×10^{-3}	1.194×10^{-3}	4.130×10^{-12}

Internal spin-limit audit: $(\ell, m, M\omega) = (2, 0, 0.52)$




FIG. 5. Internal Kerr-to-Schwarzschild continuity test at fixed $(\ell, m, M\omega) = (2, 0, 0.52)$. The first-order transmission coefficient and $-R^{(1)}$ remain paired by the flux balance while changing smoothly with a/M . The right panel normalizes both coefficients to the $a = 0$ value; all points use the endpoint-inclusive spectral solver and are accepted by the first-order balance check.

This is the point at which Kerr differs from the spherical Schwarzschild reduction: a single incident spheroidal channel sources a matrix of target channels with the same m . Only the target channel $\ell' = \ell$ contributes linearly to the total flux, because only that channel interferes with the zeroth-order outgoing field. Off-diagonal channels contribute to the energy flux first at order ϵ^2 .

2. Linear current

The scalar radial current is

$$J[R] = \frac{\Delta}{2i} \left(R^* \frac{dR}{dr} - R \frac{dR^*}{dr} \right). \quad (\text{A5})$$

Using the asymptotics in Eq. (22), the far-zone current is

$$J_\infty = -\omega |B^{\text{inc}}|^2 + \omega |B^{\text{ref}}|^2. \quad (\text{A6})$$

At the horizon, Eq. (21) gives

$$J_H = -p(r_+^2 + a^2). \quad (\text{A7})$$

The equality $J_\infty = J_H$ gives Eq. (25).

3. Fixed-incident nonlinear flux

The Green-function amplitudes $A_{\text{ref}}^{(1)}$ and $A_H^{(1)}$ are computed with the zeroth-order solution normalized to unit horizon amplitude. Scattering coefficients, however, are defined at fixed incident amplitude. Therefore we rescale the zeroth-order solution by

$$\alpha = \frac{1}{B^{\text{inc}}}. \quad (\text{A8})$$

Because the source is cubic, the first-order field scales as $|\alpha|^2 \alpha$. The fixed-incident self-channel field has the asymptotic form

$$R_{\text{fix}} \sim \frac{e^{-i\omega r_*}}{r} + \left(\frac{B^{\text{ref}}}{B^{\text{inc}}} + \epsilon \frac{A_{\text{ref}}^{(1)}}{|B^{\text{inc}}|^2 B^{\text{inc}}} \right) \frac{e^{+i\omega r_*}}{r}, \quad r \rightarrow \infty, \quad (\text{A9})$$

and

$$R_{\text{fix}} \sim \left(\frac{1}{B^{\text{inc}}} + \epsilon \frac{A_H^{(1)}}{|B^{\text{inc}}|^2 B^{\text{inc}}} \right) e^{-ipr_*}, \quad r \rightarrow r_+. \quad (\text{A10})$$

TABLE IX. Internal Kerr-to-Schwarzschild spin-limit audit for $(\ell, m, M\omega) = (2, 0, 0.52)$. The entries use the same fixed-incident normalization as the production coefficients. The two balance columns are $T^{(0)} + R^{(0)} - 1$ and $T^{(1)} + R^{(1)}$, respectively.

a/M	$T^{(1)}$	$R^{(1)}$	$ \delta_0 $	$ \delta_1 $	$ \delta_W $
0.00	8.94395×10^{-2}	-8.94395×10^{-2}	1.1×10^{-12}	3.8×10^{-13}	2.6×10^{-16}
0.05	8.95742×10^{-2}	-8.95742×10^{-2}	9.8×10^{-12}	1.1×10^{-12}	3.3×10^{-16}
0.10	8.99794×10^{-2}	-8.99794×10^{-2}	3.7×10^{-12}	4.9×10^{-13}	0
0.20	9.16178×10^{-2}	-9.16178×10^{-2}	4.3×10^{-13}	7.6×10^{-14}	1.8×10^{-16}
0.30	9.44095×10^{-2}	-9.44095×10^{-2}	1.1×10^{-11}	1.3×10^{-12}	1.8×10^{-16}
0.50	1.03814×10^{-1}	-1.03814×10^{-1}	6.0×10^{-12}	1.2×10^{-12}	0

The transmitted flux coefficient is therefore

$$T = \frac{p(r_+^2 + a^2)}{\omega} \left| \frac{1}{B_{\text{inc}}} + \epsilon \frac{A_H^{(1)}}{|B_{\text{inc}}|^2 B_{\text{inc}}} \right|^2 + \mathcal{O}(\epsilon^2)$$

$$= T^{(0)} + \epsilon \frac{2p(r_+^2 + a^2)}{\omega |B_{\text{inc}}|^4} \text{Re } A_H^{(1)} + \mathcal{O}(\epsilon^2), \quad (\text{A11})$$

which gives Eq. (41). Similarly,

$$R = \left| \frac{B_{\text{ref}}}{B_{\text{inc}}} + \epsilon \frac{A_{\text{ref}}^{(1)}}{|B_{\text{inc}}|^2 B_{\text{inc}}} \right|^2 + \mathcal{O}(\epsilon^2)$$

$$= R^{(0)} + \epsilon \frac{2 \text{Re}(\overline{B_{\text{ref}}} A_{\text{ref}}^{(1)})}{|B_{\text{inc}}|^4} + \mathcal{O}(\epsilon^2), \quad (\text{A12})$$

which gives Eq. (42). Conservation of the radial current for the inhomogeneous solution with no incoming first-order radiation then yields the first-order balance law Eq. (44).

4. Schwarzschild normalization correspondence

The flux coefficients in this paper use unit incident amplitude. This point matters when comparing them with a reduced Schwarzschild calculation that quotes the first-order Green coefficient with the zeroth-order solution normalized at the horizon. In the nonrotating limit, write the scalar field in the reduced convention as $\Phi = e^{-i\omega t} Y_{\ell m} u/r$ and let $\hat{I}$ and $\hat{R}$ denote the corresponding incident and reflected amplitudes. The present

unit-horizon Teukolsky variable has $R^{\text{in}} = r_+ u/r$ at zeroth order, so that

$$B^{\text{inc}} = r_+ \hat{I}, \quad B^{\text{ref}} = r_+ \hat{R}. \quad (\text{A13})$$

The cubic source is homogeneous of degree three. Consequently the horizon and infinity Green amplitudes obey $A_H^{(1)} = r_+^2 \hat{A}_H^{(1)}$ and $A_{\text{ref}}^{(1)} = r_+^3 \hat{A}_{\text{ref}}^{(1)}$ when the reduced solution is horizon normalized. For $a = 0$ (so $p = \omega$), Eqs. (41) and (42) therefore become

$$T_{\text{unit-inc}}^{(1)} = \frac{2 \text{Re } \hat{A}_H^{(1)}}{|\hat{I}|^4}, \quad (\text{A14})$$

$$R_{\text{unit-inc}}^{(1)} = \frac{2 \text{Re}(\hat{R}^* \hat{A}_{\text{ref}}^{(1)})}{|\hat{I}|^4}. \quad (\text{A15})$$

If the reduced Schwarzschild coefficients are instead defined with the common unit-horizon convention, $\hat{T}^{(1)} = 2 \text{Re } \hat{A}_H^{(1)} / |\hat{I}|^2$ and analogously for $\hat{R}^{(1)}$, then

$$T_{\text{unit-inc}}^{(1)} = T^{(0)} \hat{T}^{(1)}, \quad R_{\text{unit-inc}}^{(1)} = T^{(0)} \hat{R}^{(1)}. \quad (\text{A16})$$

Thus a direct comparison of the numerical values of $T^{(1)}$ between the two papers is meaningful only after this normalization conversion. The physical first-order flux correction and the balance law are unchanged.

-
- [1] N. G. Sanchez, Scattering of scalar waves from a Schwarzschild black hole, *J. Math. Phys.* **17**, 688 (1976).
 - [2] J. A. H. Futterman, F. A. Handler, and R. A. Matzner, *Scattering from Black Holes* (Cambridge University Press, Cambridge, 1988).
 - [3] K. Glampedakis and N. Andersson, Scattering of scalar waves by rotating black holes, *Class. Quantum Grav.* **18**, 1939 (2001).
 - [4] S. R. Dolan, Scattering and absorption of gravitational plane waves by rotating black holes, *Class. Quantum*

- Grav.* **25**, 235002 (2008).
- [5] Y. Decanini, A. Folacci, and B. Jensen, Complex angular momentum in black hole physics and the quasinormal modes, *Phys. Rev. D* **67**, 124017 (2003).
- [6] Y. Decanini, A. Folacci, and B. Raffaelli, Resonance and absorption spectra of the Schwarzschild black hole for massive scalar perturbations: A complex angular momentum analysis, *Phys. Rev. D* **84**, 084035 (2011).
- [7] E. Berti, V. Cardoso, and A. O. Starinets, Quasinormal modes of black holes and black branes, *Class. Quantum*

- Grav. **26**, 163001 (2009).
- [8] H.-P. Nollert, About the significance of quasinormal modes of black holes, *Phys. Rev. D* **53**, 4397 (1996).
 - [9] J. L. Jaramillo, R. Panosso Macedo, and L. Al Sheikh, Pseudospectrum and black hole quasinormal mode instability, *Phys. Rev. X* **11**, 031003 (2021).
 - [10] M. H.-Y. Cheung, K. Destounis, R. Panosso Macedo, E. Berti, and V. Cardoso, Destabilizing the fundamental mode of black holes: The elephant and the flea, *Phys. Rev. Lett.* **128**, 111103 (2022).
 - [11] K. Kyutoku, H. Motohashi, and T. Tanaka, Quasinormal modes of Schwarzschild black holes on the real axis, *Phys. Rev. D* **107**, 044012 (2023).
 - [12] T. Torres, From black hole spectral instability to stable observables, *Phys. Rev. Lett.* **131**, 111401 (2023).
 - [13] R. F. Rosato, K. Destounis, and P. Pani, Ringdown stability: Graybody factors as stable gravitational-wave observables, *Phys. Rev. D* **110**, L121501 (2024).
 - [14] N. Oshita, Thermal ringdown of a Kerr black hole: Overtone excitation, Fermi-Dirac statistics and greybody factor, *JCAP* **04**, 013 (2023).
 - [15] N. Oshita, Greybody factors imprinted on black hole ringdowns: An alternative to superposed quasinormal modes, *Phys. Rev. D* **109**, 104028 (2024).
 - [16] J. Frauendiener and C. Stevens, The non-linear perturbation of a black hole by gravitational waves. I. The Bondi-Sachs mass loss, *Class. Quantum Grav.* **38**, 194002 (2021).
 - [17] J. Frauendiener, C. Stevens, and S. Thwala, Fully non-linear gravitational wave simulations from past to future null infinity, *Phys. Rev. Lett.* **134**, 161401 (2025).
 - [18] L. Barack and A. Pound, Self-force and radiation reaction in general relativity, *Rep. Prog. Phys.* **82**, 016904 (2019).
 - [19] A. Pound and B. Wardell, Black hole perturbation theory and gravitational self-force, in *Handbook of Gravitational Wave Astronomy*, edited by C. Bambi (Springer, Singapore, 2021), Chap. 38, doi:10.1007/978-981-15-4702-7_38-1, arXiv:2101.04592 [gr-qc].
 - [20] J. L. Ripley, N. Loutrel, E. Giorgi, and F. Pretorius, Numerical computation of second-order vacuum perturbations of Kerr black holes, *Phys. Rev. D* **103**, 104018 (2021).
 - [21] N. Warburton and L. Barack, Self force on a scalar charge in Kerr spacetime: eccentric equatorial orbits, *Phys. Rev. D* **83**, 124038 (2011), doi:10.1103/PhysRevD.83.124038, arXiv:1103.0287 [gr-qc].
 - [22] B. Leather and N. Warburton, Applying the effective-source approach to frequency-domain self-force calculations for eccentric orbits, *Phys. Rev. D* **108**, 084045 (2023), doi:10.1103/PhysRevD.108.084045, arXiv:2306.17221 [gr-qc].
 - [23] M. van de Meent, Gravitational self-force on generic bound geodesics in Kerr spacetime, *Phys. Rev. D* **97**, 104033 (2018), doi:10.1103/PhysRevD.97.104033, arXiv:1711.09607 [gr-qc].
 - [24] A. Heffernan, Self-force regularization of a point particle for generic orbits in Kerr spacetime: electromagnetic and gravitational cases, arXiv:2209.05450 [gr-qc] (2022).
 - [25] Y.-X. Wei, X.-L. Zhu, J.-D. Zhang, and J. Mei, Toward second-order self-force for eccentric extreme-mass-ratio inspirals in Schwarzschild spacetime, arXiv:2504.09640 [gr-qc] (2025).
 - [26] R. Berens, T. Gravely, and A. Lupsasca, Gravitational waves on Kerr black holes I: Reconstruction of linearized metric perturbations, arXiv:2403.20311 [gr-qc] (2024).
 - [27] C. Zhang, R.-g. Cai, G. Fu, Y. Gong, X. Lu, and W. Zhou, Generic effective sources for first-order in mass-ratio gravitational self-force calculations in Schwarzschild spacetime, *Phys. Rev. D* **112**, 104069 (2025), arXiv:2505.19732 [gr-qc].
 - [28] S. D. Upton, B. Wardell, A. Pound, N. Warburton, and L. Barack, Effective source for second-order self-force calculations: quasicircular orbits in Schwarzschild spacetime, *Phys. Rev. D* **113**, 064013 (2026), arXiv:2508.00087 [gr-qc].
 - [29] V. Cardoso, J. Redondo-Yuste, U. Sperhake, and F. Tuncer, Nonlinear dynamics in general relativity, arXiv:2603.04501 [gr-qc] (2026).
 - [30] A. A. Starobinsky and S. M. Churilov, Amplification of electromagnetic and gravitational waves scattered by a rotating black hole, *Sov. Phys. JETP* **38**, 1 (1974).
 - [31] W. H. Press and S. A. Teukolsky, Floating orbits, super-radiant scattering and the black-hole bomb, *Nature* **238**, 211 (1972).
 - [32] S. Mano, H. Suzuki, and E. Takasugi, Analytic solutions of the Teukolsky equation and their low frequency expansions, *Prog. Theor. Phys.* **95**, 1079 (1996), arXiv:gr-qc/9603020.
 - [33] M. Sasaki and H. Tagoshi, Analytic black hole perturbation approach to gravitational radiation, *Living Rev. Relativ.* **6**, 6 (2003), arXiv:gr-qc/0306120.
 - [34] A. K.-W. Chung, P. Wagle, and N. Yunes, Spectral method for the gravitational perturbations of black holes: Schwarzschild background case, *Phys. Rev. D* **107**, 124032 (2023), arXiv:2302.11624 [gr-qc].
 - [35] S. A. Hughes, Computing radiation from Kerr black holes: Generalization of the Sasaki-Nakamura equation, *Phys. Rev. D* **61**, 084004 (2000), arXiv:gr-qc/0002043.
 - [36] R. K. L. Lo, Recipes for computing radiation from a Kerr black hole using a generalized Sasaki-Nakamura formalism: Homogeneous solutions, *Phys. Rev. D* **110**, 124070 (2024), arXiv:2306.16469.
 - [37] S. R. Green, S. Hollands, and P. Zimmerman, Teukolsky formalism for nonlinear Kerr perturbations, *Class. Quantum Grav.* **37**, 075001 (2020), arXiv:1908.09095.
 - [38] A. Spiers, A. Pound, and J. Moxon, Second-order Teukolsky formalism in Kerr spacetime: Formulation and nonlinear source, *Phys. Rev. D* **108**, 064002 (2023), arXiv:2305.19332.
 - [39] A. Spiers, Analytically separating the source of the Teukolsky equation, *Phys. Rev. D* **109**, 104059 (2024), arXiv:2402.00604.
 - [40] S. A. Teukolsky, Perturbations of a rotating black hole. I. Fundamental equations for gravitational, electromagnetic, and neutrino-field perturbations, *Astrophys. J.* **185**, 635 (1973).
 - [41] S. W. Hawking, Particle creation by black holes, *Commun. Math. Phys.* **43**, 199 (1975).
 - [42] L. C. Stein, qnm: A Python package for calculating Kerr quasinormal modes, separation constants, and spherical-spheroidal mixing coefficients, *J. Open Source Softw.* **4**, 1683 (2019).