


1) (20 points) Women's Basketball.

The score is tied as the clock winds down in the final seconds of the Cal – Stanford Women's Basketball game. A Cal player has the ball and makes a jump shot, releasing the ball from a height of h_o above the ground and a horizontal distance d_o from the basket, which is a height h_b above the ground, as shown in the diagram.

- What is the magnitude of the ball's acceleration at the highest point of its trajectory? As always, show your work and/or justify your answer.
- If the ball's trajectory is initially at an angle of θ above the horizontal, then what initial speed v_o is required so that the ball lands in the basket on its downward trajectory?
- Expressed as a function of v_o and θ , what is the smallest value of the ball's speed as it travels from the Cal player to the basket?
- A Stanford player standing a horizontal distance d_s from the basket starts to run towards the basket at the moment the ball is released. If the Stanford player starts from rest and then runs with a constant acceleration until she reaches the basket, what acceleration does she need to maintain to reach the basket at the same time that the ball does? Express your answer as a function of any combination of g , d_s , d_o , v_o and θ .



2) (20 points) Racecar driver

At a crucial moment near the end of a Nascar race, the driver of the car close behind the leader sees an opportunity and depresses the gas pedal in such a way as to make the car's velocity obey the following equation:

$$v(t) = A + Bt^2$$

where v is the car's velocity in m/s, t is the elapsed time in s, A and B are positive constants, and $t = 0$ is the exact time when the driver starts to make his move. The car is traveling along a straight piece of the racetrack for this entire problem.

- What are the physical units of the constants A and B ?
- What is the formula for the acceleration of the car as a function of time starting at $t = 0$?
- What is the formula for the position of the car along the track as a function of time starting at $t = 0$? Set the position of the car to 0 at $t = 0$.
- If the numerical values of A and B are 80 and 6, respectively (with the units you found for each of them above), then how far does the car get after 2 seconds?