

# Electron Mode in FRENSE

Luke J. Kersting

NEEP

University of Wisconsin - Madison

SNL Meeting

May 14, 2015



## Forward Mode

- Condensed History
- Secondary Particles
- Atomic Relaxation
- Simulation of hard electron transport events
  - Atomic excitation
  - Hard elastic scattering
  - Electroionization
  - Bremsstrahlung

## Adjoint Mode

- Hybrid Multigroup/Continuous-Energy Monte Carlo using Boltzmann-Fokker-Planck Equation (BFP)
- Other Possible Adjoint Methods

## MCNP

- Historically has only used a condensed-history approach with Goudsmit-Saunderson multiple scattering techniques.
- MCNP6 implemented a single-event method for energies below 1 keV, where the condensed-history method no longer holds.

## Penelope

- Implements a mixed method that simulates soft (condensed-history) events below a cutoff energy/angle and hard (single-events) above.

## EGS

- Condensed History Method
- Historically used Molière Multiple Scattering Theory
- EGS5 implemented Goudsmit-Saunderson Multiple Scattering to take into account spin and relativistic effects needed in the MeV range

## FRENSIE

- Hard events implemented using cross-sectional data from MCNP6
- Condensed history method will be chosen in conjunction with an adjoint method
- Ultimately hope to implement a mixed method for forward transport

## Current Capabilities

- Single Scattering Events from 100 GeV to 10 eV
- Elastic, Bremsstrahlung, Electroionization, Atomic Excitation
- Secondary particles created, but photons not tracked
- Atomic relaxation implemented

## Known Issues

- Absorption at low energies
- Negative energy from Electroionization

## Reaction

- There is no angular deflection.
- There are no secondary particles.
- Only energy loss needs to be taken into account

## Implementation

- Energy dependent electron energy loss are tabulated in ACE tables.
- No sampling is required for this process.

## Reaction

- There is no energy loss.
- There are no secondary particles.
- Only angular deflection needs to be taken into account.

## Implementation

- ACE tables provide histogram CDF of the outgoing angle cosine,  $\mu$ , for 14 – 16 energy groups.
- for  $\mu > 0.999999$  an analytical function,  $f(\mu)$ , derived from Molière's screening factor is used to compute the scattering angle.

$$f(\mu) = \frac{A}{(\eta + 1 - \mu)^2}$$

$$\eta(E, Z) = \frac{1}{4} \left( \frac{\alpha mc}{0.885p} \right)^2 Z^{2/3} [1.13 + 3.76(\alpha Z/\beta)^2]$$

## Reaction

- The subshell is directly sampled.
- A knock-on electron is ejected.
- The incident electron energy is reduced by the  $E_{knock} + E_{binding}$ .
- Conservation of momentum is used to find the scattering and ejection angles.

## Implementation

- ACE tables provide CDF of the knock-on energy,  $E_{knock}$ , based on the incident electron energy.
- The scattering and ejection angles are sampled independently breaking from a purely analog sampling.
- The shell vacancy is handled using atomic relaxation data.

## Conservation of Momentum

$$(p_{knock}c + p_ac)^2 = (pc)^2 + (p'c)^2 - 2pp' \cos(\theta)$$
$$\cos(\theta) = \frac{(pc)^2 + (p'c)^2 - (p_{knock}c)^2}{2pp'}$$

## Conservation of Energy

$$(E + m_e c^2) + (M_a c^2) = (E' + m_e c^2) + (E_a + M_a c^2) + (E_{knock} + m_e c^2) + E_{Binding}$$

Assume the binding energy is negligible

$$E = E' + E_{knock}$$

Solving independently you obtain:

$$\cos(\theta) = \frac{E'}{E} \frac{p}{p'} \quad \text{and} \quad \cos(\phi) = \frac{E_{knock}}{E} \frac{p}{p_{knock}}$$



The original sampling routine implemented in FRENSE sampled negative electron energies.

- ACE tables provide CDF of the knock-on energy,  $E_{knock}$ , based on the incident electron energy.
- The original implementation randomly selected whether to sample the upper or lower energy bin.
- A correlated sample must be made to avoid non physical values.

## Reaction

- A photon is ejected.
- The incident electron energy is reduced by the  $E_\gamma$ .
- The electron direction is assumed to be essentially unchanged.

## Implementation

- ACE tables provide CDF of the photon energy,  $E_\gamma$ , based on the incident electron energy.
- An analytical dipole function,  $p(\mu)$ , is used to sample the direction of the outgoing photon.

$$p(\mu)d\mu = \frac{(1 - \beta^2)}{2(1 - \beta\mu)^2}d\mu$$



## Absorption at low energies

- At energies near the cutoff (10 eV) the reaction cross section is dominated by elastic scattering (by order  $10^7$  for H barns).
- It is unlikely the electron will scatter below the cutoff energy.
- A temporary fix is to raise the cutoff energy (to 15 eV) to prevent indefinite elastic scattering.
- MCNP notes a similar problem and suggests a minimum cutoff energy of 20 eV.

## Test Problem


- A 10 keV electron delta source was modeled in cold Hydrogen.
- Surface tallies were set on concentric spheres to measure the flux.
- Radii ranged from 0.005 to 0.0025 cm for 5 spheres.
- Electron energy cutoff was set to 15 eV.
- Secondary photons were not tracked.


## Verification


- Test results were verified against MCNP6 using the ratio surface flux
- $10^6$  particles history were sampled in MCNP6.
- $10^6$  particles history were sampled in FRENSE.
- The  $3\sigma$  rule was used to look at the standard deviation of the flux ratio from the expected value of 1
- 68.27%, 95.45% and 99.73% of the ratios should be within 1, 2 and  $3\sigma$  respectively for agreement.

Radius (cm)	0.0005	0.0010	0.0015	0.0020	0.0025
1 $\sigma$	72.43%	66.84%	62.18%	57.58%	68.00%
2 $\sigma$	96.76%	93.16%	91.71%	93.43%	96.00%
3 $\sigma$	99.46%	97.89%	97.93%	98.48%	98.50%


**Table:** Surface flux ratio and the percentage of energy bins within 1, 2 and 3  $\sigma$  of the expected value














- The agreement is slightly below the  $3\sigma$  rule.
- There appears to be a tail off in the upper energy bins.
- Further testing is planned to evaluate the the single scattered spectrum.
- FRENSE code currently runs 100x slower than MCNP6 for a simple geometry.
- Modifications need to be made to speed up transport in DAG geometry by reducing the number of ray firings.
- Implementation of Root's combinational geometry would also speed up geometry transport.



## Hybrid Multigroup/Continuous-Energy BFP

- The same basic multigroup cross-section data can be used for forward and adjoint calculations.
- The adjoint transport model is nearly identical to the forward making implementation easy.

## Generalized Particle for Adjoint Transport

- The CSDA is used with Molière's multiple scattering theory.
- No secondary particles are produced. One particle enters an event and one leaves.