



- - the mean/average of CH intensity: the problem with that is that sometimes there are bright points within a CH, and this might drive the mean intensity higher
- - median intensity
- - minimum intensity
- - mean of 25% darkest pixel (ie mean of first quartile): this should be somewhat more robust than taking the minimum and would not be contaminated by bright points.



S[2,2](-4)	0.488191	S[1,3](-4)	0.170824
S[2,1](-7)	0.458501	S[1,1](-4)	0.167408
S[2,1](-6)	0.436247	dens(-26)	-0.166841
S[2,2](-5)	0.435000	D[1,2](-4)	-0.157496
speed(-27)	0.378315	D[1,1](-4)	-0.156624
speed(-28)	0.362714	D[1,1](-5)	-0.154992
S[2,1](-5)	0.318251	S[1,2](-7)	0.151728
S[2,2](-6)	0.316401	S[2,3](-4)	0.150383
speed(-26)	0.315370	ch_min	-0.147039
temp(-28)	0.254811	S[1,3](-5)	0.145891
S[1,2](-4)	0.227024	D[2,1](-5)	-0.140929
temp(-27)	0.223573	D[1,2](-5)	-0.136353
S[1,1](-6)	0.221872	temp(-26)	0.136092
S[1,2](-5)	0.216783	D[1,1](-6)	-0.135591
S[1,1](-7)	0.208836	D[2,1](-4)	-0.134799
S[1,1](-5)	0.205403	dens(-27)	-0.133042
S[2,2](-7)	0.200323	sunspot_number_change	-0.132842
ch_median	-0.199227	sunspot_number	-0.124398
ch_mean	-0.194180	S[1,3](-6)	0.118890
S[2,1](-4)	0.192222	D[1,2](-6)	-0.118502
S[1,2](-6)	0.184424	qs_median	-0.118260
mean_darkest_25	-0.177782		

- Intensity/contrast on different channels?

Xgboost

- Temporal leakage, I will need to investigate

```
--- Training Scenario: no_cme ---
```

```
RMSE: 15.42 km/s | R2: 0.974
```

```
--- Training Scenario: with_cme ---
```

```
RMSE: 25.37 km/s | R2: 0.929
```