

Exercises: duality

1. **Mohar-Poljak Theorem for MAXCUT** . Recall from the lecture that in the MAXCUT problem, we are given a graph $G = (V, E)$ with edge weights $w : E \rightarrow \mathbb{R}$ and the goal is to find a partition $V = A \uplus B$ which maximizes $w(A, B) = \sum_{a \in A, b \in B} w(ab)$.

This is equivalent to the following (non-convex) quadratic program:

$$\begin{aligned} & \text{maximize} && \frac{1}{2} \sum_{ij \in E} w(ij)(1 - x_i x_j) \\ & \text{subject to} && x_i^2 = 1 \quad \forall i \in V \end{aligned}$$

Let $n = |V|$. The *Laplacian* of (G, w) is the $n \times n$ matrix defined as follows:

$$L_{ij} = \begin{cases} \sum_{ik \in E} w(ik) & i = j \\ -w(ij) & i \neq j, ij \in E \\ 0 & \text{otherwise.} \end{cases}$$

Hence, our quadratic program can be rewritten as:

$$\begin{aligned} & \text{maximize} && \frac{1}{4} x^T L x \\ & \text{subject to} && x_i^2 = 1 \quad \forall i \in V \end{aligned}$$

Using Lagrange duality, show that the weight of maximum cut can be upper-bounded by

$$\frac{\lambda_{\max}(L)}{4} n,$$

where $\lambda_{\max}(L)$ is the largest eigenvalue of L .

2. **The sum of the largest elements of a vector.** For a given integer $r = 1, \dots, n$ define $f : \mathbb{R}^n \rightarrow \mathbb{R}$ as

$$f(x) = \sum_{i=1}^r x_{[i]},$$

where $x_{[1]} \geq x_{[2]} \geq \dots \geq x_{[n]}$ are the components of x .

- a) Prove that the constraint $f(x) \leq \alpha$ is convex.

Hint: Showing that it is equivalent to $\binom{n}{r}$ linear inequalities.

- b) Note that given $x \in \mathbb{R}^n$, the value of $f(x)$ is the optimum value of the following LP:

$$\begin{aligned} & \text{maximize} && x^T y \\ & \text{subject to} && 0 \leq y \leq 1 \\ & && \sum_{i=1}^n y_i = r. \end{aligned} \tag{1}$$

Derive the dual of the LP above and use it to emulate the constraint $f(x) \leq \alpha$ using only $O(n)$ linear inequalities.

- c) To see an application of f , extend the convex program Markovitz portfolio optimization from the lecture:

$$\begin{aligned} & \text{minimize} && x^T Q x \\ & \text{subject to} && \sum_i x_i = B \\ & && \mu^T x \geq M \\ & && x \geq 0 \end{aligned}$$

by a constraint saying that no more than 80% of the total budget can be invested in any 10% of the assets.

3. **Separating hyperplane between two polyhedra.** Formulate the following problem as an LP: find a separating hyperplane that strictly separates two polyhedra $\mathcal{P}_1 = \{x \mid Ax \leq b\}$ and $\mathcal{P}_2 = \{x \mid Cx \leq d\}$, i.e., find a hyperplane $a^T x = \gamma$ such that $a^T x \geq \gamma$ for $x \in \mathcal{P}_1$ and $a^T x \leq \gamma$ for $x \in \mathcal{P}_2$.

Hint: use LP duality to find γ and a such that

$$\inf_{x \in \mathcal{P}_1} a^T x \geq \gamma \geq \sup_{x \in \mathcal{P}_2} a^T x.$$