

# Anosov property of cyclic $\mathrm{SO}_0(2,3)$ -Higgs Bundles (arXiv:2406.08118)

张峻铭 Junming Zhang

Chern Institute of Mathematics, Nankai University

Introductory School “Geometric Structures & Discrete Group Actions”  
2025.04.15 at CIRM, Marseille



南开大学  
Nankai University

# Contents

- 1 Anosov Property and Higher (rank) Teichmüller Theory
- 2 Non-Abelian Hodge Correspondence
- 3 From Higgs Bundle to Anosov Representation

# Anosov Property and Higher (rank) Teichmüller Theory

# Classical Teichmüller Theory

Let  $S$  be a closed connected oriented surface whose genus is larger than 2. Its Teichmüller space  $\mathcal{T}(S)$  is the moduli space of marked hyperbolic structure over  $S$ .

# Classical Teichmüller Theory

Let  $S$  be a closed connected oriented surface whose genus is larger than 2. Its Teichmüller space  $\mathcal{T}(S)$  is the moduli space of marked hyperbolic structure over  $S$ .

We can view  $\mathcal{T}(S)$  as a connected component of the character variety

$$\mathrm{Hom}^{\mathrm{red}}(\pi_1(S), \mathrm{PSL}(2, \mathbb{R})) / \mathrm{PSL}(2, \mathbb{R}).$$

# Classical Teichmüller Theory

Let  $S$  be a closed connected oriented surface whose genus is larger than 2. Its Teichmüller space  $\mathcal{T}(S)$  is the moduli space of marked hyperbolic structure over  $S$ .

We can view  $\mathcal{T}(S)$  as a connected component of the character variety

$$\mathrm{Hom}^{\mathrm{red}}(\pi_1(S), \mathrm{PSL}(2, \mathbb{R})) / \mathrm{PSL}(2, \mathbb{R}).$$

Moreover,  $\mathcal{T}(S)$  is one of the two connected components which consist entirely of discrete and faithful representations. The other one is  $\mathcal{T}(\overline{S})$ , where  $\overline{S}$  denotes the surface  $S$  with the opposite orientation.

# Move to Higher Rank

Instead of focusing on representations from  $\pi_1(S)$  into  $\mathrm{PSL}(2, \mathbb{R})$ , we replace the target by a semi-simple Lie group  $G$  of higher real rank, such as

# Move to Higher Rank

Instead of focusing on representations from  $\pi_1(S)$  into  $\mathrm{PSL}(2, \mathbb{R})$ , we replace the target by a semi-simple Lie group  $G$  of higher real rank, such as

$$\mathrm{SL}(n, \mathbb{R}) \quad (n \geq 3),$$



# Move to Higher Rank

Instead of focusing on representations from  $\pi_1(S)$  into  $\mathrm{PSL}(2, \mathbb{R})$ , we replace the target by a semi-simple Lie group  $G$  of higher real rank, such as

$$\mathrm{SL}(n, \mathbb{R}) \quad (n \geq 3),$$

$$\mathrm{Sp}(2n, \mathbb{R}) \quad (n \geq 2),$$

# Move to Higher Rank

Instead of focusing on representations from  $\pi_1(S)$  into  $\mathrm{PSL}(2, \mathbb{R})$ , we replace the target by a semi-simple Lie group  $G$  of higher real rank, such as

$$\mathrm{SL}(n, \mathbb{R}) \quad (n \geq 3),$$

$$\mathrm{Sp}(2n, \mathbb{R}) \quad (n \geq 2),$$

$$\mathrm{SO}_0(p, q) \quad (\min\{p, q\} \geq 2)$$

and so on.

# Move to Higher Rank

Instead of focusing on representations from  $\pi_1(S)$  into  $\mathrm{PSL}(2, \mathbb{R})$ , we replace the target by a semi-simple Lie group  $G$  of higher real rank, such as

$$\mathrm{SL}(n, \mathbb{R}) \quad (n \geq 3),$$

$$\mathrm{Sp}(2n, \mathbb{R}) \quad (n \geq 2),$$

$$\mathrm{SO}_0(p, q) \quad (\min\{p, q\} \geq 2)$$

and so on.

In general, the set of discrete and faithful representations is only closed and NOT open.

# Higher Teichmüller Spaces

## Definition

A **higher Teichmüller space** is a subset of  $\text{Hom}^{\text{red}}(\pi_1(S), G)/G$ , which is a union of connected components that consist entirely of discrete and faithful representations.

# Higher Teichmüller Spaces

## Definition

A **higher Teichmüller space** is a subset of  $\text{Hom}^{\text{red}}(\pi_1(S), G)/G$ , which is a union of connected components that consist entirely of discrete and faithful representations.

We give two famous examples of higher Teichmüller space here.

# Higher Teichmüller Spaces

## Definition

A **higher Teichmüller space** is a subset of  $\text{Hom}^{\text{red}}(\pi_1(S), G)/G$ , which is a union of connected components that consist entirely of discrete and faithful representations.

We give two famous examples of higher Teichmüller space here.

- (Hitchin; Labourie; Fock–Goncharov) Hitchin components  $\mathcal{T}_{Hit}(S, G)$  for **real split**  $G$ , such as  $\text{SL}(n, \mathbb{R})$ ,  $\text{SO}_0(n, n+1)$ ,  $\text{Sp}(2n, \mathbb{R})$  and so on;

# Higher Teichmüller Spaces

## Definition

A **higher Teichmüller space** is a subset of  $\text{Hom}^{\text{red}}(\pi_1(S), G)/G$ , which is a union of connected components that consist entirely of discrete and faithful representations.

We give two famous examples of higher Teichmüller space here.

- (Hitchin; Labourie; Fock–Goncharov) Hitchin components  $\mathcal{T}_{Hit}(S, G)$  for **real split**  $G$ , such as  $\text{SL}(n, \mathbb{R})$ ,  $\text{SO}_0(n, n+1)$ ,  $\text{Sp}(2n, \mathbb{R})$  and so on;
- (Burger–Iozzi–Wienhard) Maximal components  $\mathcal{T}_{max}(S, G)$  for **Hermitian**  $G$ , such as  $\text{SU}(p, q)$ ,  $\text{SO}_0(2, n)$  and so on.

# Higher Teichmüller Spaces

## Definition

A **higher Teichmüller space** is a subset of  $\text{Hom}^{\text{red}}(\pi_1(S), G)/G$ , which is a union of connected components that consist entirely of discrete and faithful representations.

We give two famous examples of higher Teichmüller space here.

- (Hitchin; Labourie; Fock–Goncharov) Hitchin components  $\mathcal{T}_{\text{Hit}}(S, G)$  for **real split**  $G$ , such as  $\text{SL}(n, \mathbb{R})$ ,  $\text{SO}_0(n, n+1)$ ,  $\text{Sp}(2n, \mathbb{R})$  and so on;
- (Burger–Iozzi–Wienhard) Maximal components  $\mathcal{T}_{\text{max}}(S, G)$  for **Hermitian**  $G$ , such as  $\text{SU}(p, q)$ ,  $\text{SO}_0(2, n)$  and so on.

When  $G = \text{SL}(2, \mathbb{R})$ , the above components coincide with the classical Teichmüller space.



# Anosov Property

Anosov property plays an important role in higher Teichmüller theory. Roughly speaking, the Anosov property introduced by F. Labourie means that the representation  $\rho: \pi_1(S) \rightarrow G$  satisfies some special dynamic properties.

# Anosov Property

Anosov property plays an important role in higher Teichmüller theory. Roughly speaking, the Anosov property introduced by F. Labourie means that the representation  $\rho: \pi_1(S) \rightarrow G$  satisfies some special dynamic properties.

The original definition is highly dynamical and we will use an interpreted definition (which is proven equivalent to the original one by Kapovich–Leeb–Porti and Bochi–Potrie–Sambarino).

# Anosov Property

Anosov property plays an important role in higher Teichmüller theory. Roughly speaking, the Anosov property introduced by F. Labourie means that the representation  $\rho: \pi_1(S) \rightarrow G$  satisfies some special dynamic properties.

The original definition is highly dynamical and we will use an interpreted definition (which is proven equivalent to the original one by Kapovich–Leeb–Porti and Bochi–Potrie–Sambarino).

Also, we will assume  $G$  is a semi-simple Lie subgroup of  $\mathrm{SL}(n, \mathbb{C})$  here to avoid involving a Lie-theoretic description.

Fix a basepoint  $x_0 \in \mathbb{H}^2$ .

### Definition

$\rho: \pi_1(S) \rightarrow G$  is  $P_k$ -**Anosov** if there exist constants  $D, L > 0$  such that

$$\log \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \geq D \cdot d_{\mathbb{H}^2}(x_0, \gamma \cdot x_0) - L, \forall \gamma \in \pi_1(S),$$

where  $\sigma_i$  denotes the  $i$ -th singular value.

Fix a basepoint  $x_0 \in \mathbb{H}^2$ .

### Definition

$\rho: \pi_1(S) \rightarrow G$  is  $P_k$ -**Anosov** if there exist constants  $D, L > 0$  such that

$$\log \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \geq D \cdot d_{\mathbb{H}^2}(x_0, \gamma \cdot x_0) - L, \forall \gamma \in \pi_1(S),$$

where  $\sigma_i$  denotes the  $i$ -th singular value.

For  $g \in \mathrm{SO}_0(2, n) (n > 2)$ , it is remarkable that the singular values of  $g$  are

$$\sigma_1(g), \sigma_2(g), 1, \dots, 1, \sigma_2^{-1}(g), \sigma_1^{-1}(g).$$

Fix a basepoint  $x_0 \in \mathbb{H}^2$ .

### Definition

$\rho: \pi_1(S) \rightarrow G$  is  $P_k$ -**Anosov** if there exist constants  $D, L > 0$  such that

$$\log \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \geq D \cdot d_{\mathbb{H}^2}(x_0, \gamma \cdot x_0) - L, \forall \gamma \in \pi_1(S),$$

where  $\sigma_i$  denotes the  $i$ -th singular value.

For  $g \in \mathrm{SO}_0(2, n) (n > 2)$ , it is remarkable that the singular values of  $g$  are

$$\sigma_1(g), \sigma_2(g), 1, \dots, 1, \sigma_2^{-1}(g), \sigma_1^{-1}(g).$$

**Anosov  $\implies$  discrete + faithful.**

Fix a basepoint  $x_0 \in \mathbb{H}^2$ .

### Definition

$\rho: \pi_1(S) \rightarrow G$  is  $P_k$ -**Anosov** if there exist constants  $D, L > 0$  such that

$$\log \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \geq D \cdot d_{\mathbb{H}^2}(x_0, \gamma \cdot x_0) - L, \forall \gamma \in \pi_1(S),$$

where  $\sigma_i$  denotes the  $i$ -th singular value.

For  $g \in \mathrm{SO}_0(2, n) (n > 2)$ , it is remarkable that the singular values of  $g$  are

$$\sigma_1(g), \sigma_2(g), 1, \dots, 1, \sigma_2^{-1}(g), \sigma_1^{-1}(g).$$

**Anosov  $\implies$  discrete + faithful.**

Anosov property is open but not closed. However, all representations in the known higher Teichmüller spaces are Anosov.

# Non-Abelian Hodge Correspondence



# Higgs bundles

The Higgs bundle is a useful tool to study the higher Teichmüller space. It is usually used to give a parametrization of the higher Teichmüller space. We fix a complex structure on  $S$  such that it becomes a Riemann surface  $X$ . Let  $\mathcal{K}_X$  be its canonical line bundle.

# Higgs bundles

The Higgs bundle is a useful tool to study the higher Teichmüller space. It is usually used to give a parametrization of the higher Teichmüller space. We fix a complex structure on  $S$  such that it becomes a Riemann surface  $X$ . Let  $\mathcal{K}_X$  be its canonical line bundle.

## Definition

A  $(\mathrm{GL}(n, \mathbb{C})\text{-})$ **Higgs bundle** over  $X$  is a pair  $(\mathcal{E}, \Phi)$  consisting of the following data:

- a holomorphic vector bundle  $\mathcal{E}$  over  $X$  with  $\mathrm{rank}(\mathcal{E}) = n$ ;
- a holomorphic section  $\Phi \in H^0(X, \mathrm{End}(\mathcal{E}) \otimes \mathcal{K}_X)$  called **Higgs field**.

The non-Abelian Hodge correspondence exhibit a homeomorphism between the following moduli spaces:

$$\begin{array}{ccc}
 \{\textbf{reductive representation } \rho: \pi_1(S) \rightarrow \mathrm{GL}(n, \mathbb{C})\} / \mathrm{GL}(n, \mathbb{C}) & & \\
 \downarrow \text{associated bundle} & \uparrow \text{holonomy} & \\
 \{\textbf{reductive flat vector bundle } (E \rightarrow S, \nabla) \text{ with } \mathrm{rank}(E) = n\} / \mathcal{G} & & \\
 \uparrow \text{harmonic metric} & & \\
 \downarrow & & \\
 \{\textbf{polystable Higgs bundle } (\mathcal{E} \rightarrow X, \Phi) \text{ with } \mathrm{rank}(\mathcal{E}) = n, \deg(\mathcal{E}) = 0\} / \mathcal{G} & & 
 \end{array}$$

The non-Abelian Hodge correspondence exhibit a homeomorphism between the following moduli spaces:

$$\begin{array}{ccc}
 \{\textbf{reductive representation } \rho: \pi_1(S) \rightarrow \mathrm{GL}(n, \mathbb{C})\} / \mathrm{GL}(n, \mathbb{C}) & & \\
 \downarrow \text{associated bundle} \quad \quad \quad \uparrow \text{holonomy} & & \\
 \{\textbf{reductive flat vector bundle } (E \rightarrow S, \nabla) \text{ with } \mathrm{rank}(E) = n\} / \mathcal{G} & & \\
 \uparrow \text{harmonic metric} & & \\
 \downarrow & & \\
 \{\textbf{polystable Higgs bundle } (\mathcal{E} \rightarrow X, \Phi) \text{ with } \mathrm{rank}(\mathcal{E}) = n, \deg(\mathcal{E}) = 0\} / \mathcal{G}
 \end{array}$$

If we equip additional structure on these objects, we can get the non-Abelian Hodge correspondence for general reductive Lie group  $G$ .

# Kobayashi–Hitchin Correspondence (from Higgs to flat)

We only explain how to get a flat bundle from a Higgs bundle here. The key point is the Hitchin's self-dual equation.

# Kobayashi–Hitchin Correspondence (from Higgs to flat)

We only explain how to get a flat bundle from a Higgs bundle here. The key point is the Hitchin's self-dual equation.

## Theorem (Hitchin–Simpson)

*If  $(\mathcal{E}, \Phi)$  is a polystable Higgs bundle with  $\deg(\mathcal{E}) = 0$ , then there exists an Hermitian metric  $h$  on  $\mathcal{E}$  such that*

$$F(\nabla^h) + [\Phi, \Phi^{*h}] = 0, \quad (1)$$

*where  $\nabla^h$  is the Chern connection of the metric  $h$ ,  $F(\nabla^h)$  denotes its curvature form and  $\Phi^{*h}$  is the adjoint of  $\Phi$  with respect to  $h$ . Moreover, if  $(\mathcal{E}, \Phi)$  is stable, then such  $h$  is unique up to a constant scalar.*

# Kobayashi–Hitchin Correspondence (from Higgs to flat)

We only explain how to get a flat bundle from a Higgs bundle here. The key point is the Hitchin's self-dual equation.

## Theorem (Hitchin–Simpson)

*If  $(\mathcal{E}, \Phi)$  is a polystable Higgs bundle with  $\deg(\mathcal{E}) = 0$ , then there exists an Hermitian metric  $h$  on  $\mathcal{E}$  such that*

$$F(\nabla^h) + [\Phi, \Phi^{*h}] = 0, \quad (1)$$

*where  $\nabla^h$  is the Chern connection of the metric  $h$ ,  $F(\nabla^h)$  denotes its curvature form and  $\Phi^{*h}$  is the adjoint of  $\Phi$  with respect to  $h$ . Moreover, if  $(\mathcal{E}, \Phi)$  is stable, then such  $h$  is unique up to a constant scalar.*

If  $h$  solves (1), then

$$\nabla^h + \Phi + \Phi^{*h}$$

gives a flat connection.

# From Higgs Bundle to Anosov Representation



Higgs bundle  $\xrightarrow{\text{Hitchin's self-dual equation}}$  Anosov property?

# Example: Hitchin Component in Higgs Bundle Viewpoint

The Hitchin component for  $\mathrm{SL}(3, \mathbb{R})$  consisting of entirely the Higgs bundles of the following form:

$$\mathcal{E} = \mathcal{K}_X \oplus \mathcal{O} \oplus \mathcal{K}_X^\vee,$$

$$\Phi = \begin{pmatrix} 0 & q_2 & q_3 \\ 1 & 0 & q_2 \\ 0 & 1 & 0 \end{pmatrix},$$

where  $1: \mathcal{K}_X \rightarrow \mathcal{O} \otimes \mathcal{K}_X$  and  $1: \mathcal{O} \rightarrow \mathcal{K}_X^\vee \otimes \mathcal{K}_X$  are the natural isomorphisms and  $q_i \in H^0(X, \mathcal{K}_X^i)$ .

It corresponds to the component containing the embedding of Fuchsian representations through the unique irreducible  $\mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(3, \mathbb{R})$ .

# Special $\mathrm{SO}_0(2, 3)$ -Higgs Bundles

Below we consider the Higgs bundle whose underlying bundle is

$$\mathcal{E} = \mathcal{L}_{-2} \oplus \mathcal{L}_{-1} \oplus \mathcal{L}_0 \oplus \mathcal{L}_1 \oplus \mathcal{L}_2,$$

where  $\mathcal{L}_i$  are line bundles with  $\mathcal{L}_i \cong \mathcal{L}_{-i}^\vee$  and  $\mathcal{L}_0 \cong \mathcal{O}$  and

$$\Phi = \begin{pmatrix} 0 & 0 & 0 & \gamma & 0 \\ \tau & 0 & 0 & 0 & \gamma \\ 0 & \beta & 0 & 0 & 0 \\ 0 & 0 & \beta & 0 & 0 \\ 0 & 0 & 0 & \tau & 0 \end{pmatrix}.$$

$(\mathcal{E}, \Phi)$  is a cyclic  $\mathrm{SO}_0(2, 3)$ -Higgs bundle and we denote it by

$$\begin{array}{ccccccc} & & \gamma & & \gamma & & \\ & \swarrow & & \searrow & & \swarrow & \\ \mathcal{L}_{-2} & \xrightarrow{\tau} & \mathcal{L}_{-1} & \xrightarrow{\beta} & \mathcal{L}_0 & \xrightarrow{\beta} & \mathcal{L}_1 & \xrightarrow{\tau} & \mathcal{L}_2 \end{array}$$

If the cyclic Higgs bundle

$$\begin{array}{ccccccc}
 & & \gamma & & \gamma & & \\
 & \swarrow & & \searrow & \swarrow & \searrow & \\
 \mathcal{L}_{-2} & \xrightarrow{\tau} & \mathcal{L}_{-1} & \xrightarrow{\beta} & \mathcal{L}_0 & \xrightarrow{\beta} & \mathcal{L}_1 & \xrightarrow{\tau} & \mathcal{L}_2
 \end{array} \tag{2}$$

is (semi-)stable, then the Milnor–Wood inequality implies that

$$|\deg(\mathcal{L}_1)| \leq \deg(\mathcal{K}_X).$$

When “=” holds, the Higgs bundle corresponds to a maximal representation and maximal representations are known to be  $P_1$ -Anosov.

If the cyclic Higgs bundle

$$\begin{array}{ccccccc}
 & & \gamma & & \gamma & & \\
 & \swarrow & & \searrow & \swarrow & \searrow & \\
 \mathcal{L}_{-2} & \xrightarrow{\tau} & \mathcal{L}_{-1} & \xrightarrow{\beta} & \mathcal{L}_0 & \xrightarrow{\beta} & \mathcal{L}_1 \xrightarrow{\tau} \mathcal{L}_2
 \end{array} \tag{2}$$

is (semi-)stable, then the Milnor–Wood inequality implies that

$$|\deg(\mathcal{L}_1)| \leq \deg(\mathcal{K}_X).$$

When “=” holds, the Higgs bundle corresponds to a maximal representation and maximal representations are known to be  $P_1$ -Anosov.

Such Higgs bundle is called  $\alpha_1$ -**cyclic** if  $\tau: \mathcal{L}_1 \rightarrow \mathcal{L}_2 \otimes \mathcal{K}_X$  is an isomorphism. It is asked by Collier–Tholozan–Toulisse that

### Question

*Given a stable  $\alpha_1$ -cyclic  $\mathrm{SO}_0(2, 3)$ -Higgs bundle (2). When  $|\deg(\mathcal{L}_1)| < \deg(\mathcal{K}_X)$ , is the corresponding representation Anosov?*

From a different starting point, S. Filip considered the  $\alpha_1$ -cyclic  $\mathrm{SO}_0(2,3)$ -Higgs bundles arising from variation of Hodge structures, i.e.  $\gamma \equiv 0$ .

From a different starting point, S. Filip considered the  $\alpha_1$ -cyclic  $\mathrm{SO}_0(2,3)$ -Higgs bundles arising from variation of Hodge structures, i.e.  $\gamma \equiv 0$ .

### Theorem (Filip)

*A stable  $\mathrm{SO}_0(2,3)$ -Higgs bundle of the form*

$$\mathcal{L}_{-2} \xrightarrow{\tau} \mathcal{L}_{-1} \xrightarrow{\beta} \mathcal{L}_0 \xrightarrow{\beta} \mathcal{L}_1 \xrightarrow{\tau} \mathcal{L}_2$$

*with  $\tau$  is an isomorphism gives a  $P_2$ -Anosov representation.*

Filip proved this theorem by an **analytic method**. Inspired by his method and with some simplification, we extend his results and discover the Anosov property of a general family of  $\mathrm{SO}_0(2,3)$ -Higgs bundles.

## Theorem (Z.)

A stable  $\alpha_1$ -cyclic  $\mathrm{SO}_0(2, 3)$ -Higgs bundle

$$\mathcal{L}_{-2} \xrightarrow{\tau} \mathcal{L}_{-1} \xrightarrow{\beta} \mathcal{L}_0 \xrightarrow{\beta} \mathcal{L}_1 \xrightarrow{\tau} \mathcal{L}_2$$

with  $\deg(\mathcal{L}_1) < \deg(\mathcal{K}_X)$  gives a  $P_2$ -Anosov representation.

They form a **non-compact subset (which can go to infinity)** in the character variety.



## Theorem (Z.)

A stable  $\alpha_1$ -cyclic  $\mathrm{SO}_0(2, 3)$ -Higgs bundle

$$\mathcal{L}_{-2} \xrightarrow{\tau} \mathcal{L}_{-1} \xrightarrow{\beta} \mathcal{L}_0 \xrightarrow{\beta} \mathcal{L}_1 \xrightarrow{\tau} \mathcal{L}_2$$

with  $\deg(\mathcal{L}_1) < \deg(\mathcal{K}_X)$  gives a  $P_2$ -Anosov representation.

They form a **non-compact subset (which can go to infinity)** in the character variety.

## Remark

The trivial line bundle  $\mathcal{L}_0$  above can be replaced by an orthogonal vector bundle of rank  $n$  to get an  $\mathrm{SO}(2, n+2)$ -Higgs bundle. With the assumption of stability, the Anosov property still holds.

# Non-compact case

Lots of results in this section can be generalized when the closed hyperbolic Riemann surface  $X$  is replaced by a complete hyperbolic Riemann surface  $X := \overline{X} \setminus D$  of finite type, i.e. both  $g(\overline{X})$  and  $\#D$  are finite.

# Non-compact case

Lots of results in this section can be generalized when the closed hyperbolic Riemann surface  $X$  is replaced by a complete hyperbolic Riemann surface  $X := \overline{X} \setminus D$  of finite type, i.e. both  $g(\overline{X})$  and  $\#D$  are finite.

$D = \emptyset$ (closed surface)	$D \neq \emptyset$ (non-compact surface of finite type)

# Non-compact case

Lots of results in this section can be generalized when the closed hyperbolic Riemann surface  $X$  is replaced by a complete hyperbolic Riemann surface  $X := \overline{X} \setminus D$  of finite type, i.e. both  $g(\overline{X})$  and  $\#D$  are finite.

$D = \emptyset$ (closed surface)	$D \neq \emptyset$ (non-compact surface of finite type)
Anosov representation	almost-dominated representation
Higgs bundle	parabolic Higgs bundle

# Non-compact case

Lots of results in this section can be generalized when the closed hyperbolic Riemann surface  $X$  is replaced by a complete hyperbolic Riemann surface  $X := \overline{X} \setminus D$  of finite type, i.e. both  $g(\overline{X})$  and  $\#D$  are finite.

$D = \emptyset$ (closed surface)	$D \neq \emptyset$ (non-compact surface of finite type)
Anosov representation	almost-dominated representation
Higgs bundle	parabolic Higgs bundle

## Theorem (Z.)

*A stable  $\alpha_1$ -cyclic parabolic  $\mathrm{SO}_0(2, 3)$ -Higgs bundle satisfying specific assumption on parabolic weights and degree gives a  $P_2$ -almost-dominated representation through the non-Abelian Hodge correspondence.*

# Thank you!