

# Transform Order by Reversals : ILP vs SAT-solver

Updated on June 21, 2019 at 1:30pm

*Professor Dan Gusfield Group*

**Lei Zuo**

For Integer Linear Programming, we have the following conditions:

**1.**

For each  $k$  from 1 to  $n - 1$ :

$$NOP(k) + \sum_{(p,q): p < q} R(p, q, k) = 1$$

For an equivalent dnf, we see that there is one and only one 1 among all these values.

$$(NOP(k) \wedge \neg R_1 \wedge \neg R_2 \wedge \dots) \vee (\neg NOP(k) \wedge R_1 \wedge \neg R_2 \wedge \dots) \vee \dots$$

We want a cnf that is not too complicate. One of them is true and none of the pairs can be both true. <sup>1</sup>

$$(NOP(k) \vee R_1 \vee R_2 \vee \dots) \wedge (\neg NOP(k) \vee \neg R_1) \wedge (\neg NOP(k) \vee \neg R_2) \wedge (\neg R_1 \vee \neg R_2) \wedge \dots$$

**2**

For  $i$  from 1 to  $n$ :

$$\sum_{q=1}^n X(1, i, q) = 1$$

For an equivalent cnf, we see that there is one and only one 1 among all these values.

$$(X_1 \vee X_2 \vee X_3 \vee \dots) \wedge (\neg X_1 \vee \neg X_2) \wedge (\neg X_1 \vee \neg X_3) \wedge (\neg X_2 \vee \neg X_3) \wedge \dots$$

**3**

For  $i$  from 1 to  $n$ :

$$\sum_{p=1}^n X(n-1, i, p, Q_2(i)) = 1$$

For an equivalent cnf, we see that there is one and only one 1 among all these values.

$$(X_1 \vee X_2 \vee X_3 \vee \dots) \wedge (\neg X_1 \vee \neg X_2) \wedge (\neg X_1 \vee \neg X_3) \wedge (\neg X_2 \vee \neg X_3) \wedge \dots$$

**4**

For each node  $q$  at a level  $k$  from 2 to  $n - 1$ :

$$\sum_{p=1}^n [\sum_{i=1}^n X(k-1, i, p, q)] = 1$$

For an equivalent cnf, we see that there is one and only one 1 among all these values.

$$(X_1 \vee X_2 \vee X_3 \vee \dots) \wedge (\neg X_1 \vee \neg X_2) \wedge (\neg X_1 \vee \neg X_3) \wedge (\neg X_2 \vee \neg X_3) \wedge \dots$$

<sup>1</sup><https://math.stackexchange.com/questions/2554805/writing-in-cnf-that-only-one-statement-can-be-true>

**5**

For each node  $p$  at a level  $k$  from 2 to  $n - 1$ :

$$\sum_{q=1}^n [\sum_{i=1}^n X(k, i, p, q)] = 1$$

For an equivalent cnf, we see that there is one and only one 1 among all these values.

$$(X_1 \vee X_2 \vee X_3 \vee \dots) \wedge (\neg X_1 \vee \neg X_2) \wedge (\neg X_1 \vee \neg X_3) \wedge (\neg X_2 \vee \neg X_3) \wedge \dots$$

**6**

For each integer  $i$  from 1 to  $n$ :

$$\sum_{q=1}^n X(k-1, i, q, p) = \sum_{q=1}^n X(k, i, p, q)$$

We know that there is at most one 1 (XNOR). The tested format is in the ipynb file. Suppose  $n = 5$ .

$$(X_1 \vee X_2 \vee X_3 \vee X_4 \vee X_5 \vee \neg X_6) \wedge (X_1 \vee X_2 \vee X_3 \vee X_4 \vee X_5 \vee \neg X_7) \wedge \dots (X_6 \vee X_7 \vee X_8 \vee X_9 \vee X_{10} \vee \neg X_1) \dots$$

**7**