

Projective Bottlenecks in Passive Spectral Routing: Exact and Border Memory, Dense Low-State Limits, and Error Barriers

Lluis Eriksson

13 August 2026

Abstract

We determine the exact passive state cost of every finite routing table whose target lines lie in one complex two-plane. At distinct boundary frequencies ζ_i , a square rational-inner matrix must route one input line to prescribed target lines Y_i . Let δ be the least algebraic degree of a rational map $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ interpolating the ordered projective data (ζ_i, Y_i) . We prove the exact reduction

$$d_{\min} = \delta.$$

The upper bound is constructive: scalar Fejér–Riesz factorization lifts any coprime rational pair to an explicit 2×2 lossless network of exactly the same McMillan degree. The lower bound uses a common state-space denominator: more boundary zeros than states force the routed column to remain identically inside the target plane, where it becomes a projective rational interpolant.

The same denominator mechanism yields a quantitative result for approximate routing. If the normalized interpolation matrix M_d has smallest singular value $\sigma > 0$, then every lossless router of degree at most d has worst-node complex-projective chordal error at least

$$\frac{\sigma}{\sqrt{L(d+1) + \sigma^2}}.$$

Under a data-matrix perturbation of norm η , the bound survives with σ replaced by $(\sigma - \eta)_+$. Thus the algebraic obstruction has a finite-error consequence, not merely an exact-interpolation discontinuity. We also solve the complete zero-memory frontier: under the Bloch-sphere identification of $\mathbb{P}^1$, E_0 is the sine of half the radius of the smallest spherical cap containing the targets. A support certificate uses at most three active targets. For two targets the formula is $E_0 = \sin(\alpha/2)$, where α is their Fubini–Study angle, and the singular-value obstruction is asymptotically sharp as the targets collide.

There is also an exact positive-degree phenomenon. Define the border memory $d_{\partial} = \min\{d : E_d = 0\}$ and let $\delta(A)$ be the projective interpolation degree of the subtable indexed by A . We prove the base-point deletion law

$$d_{\partial} = \min_{B \subseteq \{1, \dots, L\}} (|B| + \delta(B^c)).$$

Thus every planar table has three rigorously separated regimes: positive error, zero but unattained infimum, and exact realization. For binary occupancies n_0, n_{∞} , border and exact memory are respectively $\min(n_0, n_{\infty})$ and $\max(n_0, n_{\infty})$; their ratio is unbounded. Universally, $E_{d_0} = 0$ at $d_0 = \lceil (L-1)/2 \rceil$, whereas below d_0 an open dense set has positive error.

This reduction reveals two competing mechanisms. For two target lines with occupancies n_0, n_{∞} , the exact cost is $\max(n_0, n_{\infty})$, a fiber multiplicity law. By contrast, for L distinct generic coplanar targets,

$$d_{\min} = \left\lceil \frac{L-1}{2} \right\rceil,$$

whereas both the span bound and the optimized constant-detector hierarchy of the preceding framework equal one. Their multiplicative underestimate is therefore unbounded. For four nodes we give the complete phase diagram: degree zero for one constant target, degree one

exactly for four distinct targets with matching ordered cross-ratio, degree three for a $3 + 1$ collision, and degree two in every remaining case. For the strict four-node fixture, an exact Sylvester certificate forces more than 21% worst-node error on every one-state router. Exact Gaussian-rational fixtures verify all strata, the quantitative obstruction, and the generic law through $L = 16$. Exact producer, replay validator, frozen evidence, and immutable release metadata are linked in the paper.

Keywords: rational inner matrix; projective interpolation; McMillan degree; passive memory; border memory; approximate routing; chordal error; cross-ratio; Fejér–Riesz factorization; Wigner–Smith delay.

1 A singular stratum with hidden memory

Let S be an $N \times N$ rational matrix, analytic in $\mathbb{D}$, unitary almost everywhere on $\mathbb{T}$, and regular at distinct nodes $\zeta_1, \dots, \zeta_L \in \mathbb{T}$. Its McMillan degree is the state dimension of a minimal conservative realization. Fix an input line $X \subset \mathbb{C}^N$ and target lines $Y_i \subset \mathbb{C}^N$. The routing task is

$$S(\zeta_i)X = Y_i, \quad 1 \leq i \leq L. \quad (1)$$

Endpoint phases and bases are free. Write $d_{\min}$ for the least McMillan degree among all such S .

The generic direct-sum law counts states by target occupancy, and its constant-detector extension gives useful lower bounds on singular target arrangements [1]. It deliberately leaves open a complete equality theory outside direct-sum position. The first unresolved stratum in that preceding framework is also the smallest geometrically nontrivial one: all Y_i lie in a fixed two-plane H . Here the span statistic is frozen, yet projective compatibility can demand arbitrarily many states.

Classical scalar rational interpolation and its state-variable realization describe minimal rational functions through finite data [2, 3, 4, 5]. Minimal boundary interpolation by scalar unimodular functions and finite Blaschke products is also well developed [6, 7, 8]. Subspace Nevanlinna interpolation and boundary matrix all-pass construction provide the matrix context [9, 10, 11]. Our contribution is not a new scalar interpolation algorithm. It is the exact identification of that scalar projective degree with the physical state count of a square passive lossless router, together with a quantitative approximation barrier for low-state routers, exact zero-memory and base-point deletion laws, the resulting generic law, unbounded detector gap, and complete four-line stratification.

2 Projective degree equals passive memory

Choose a basis of H and identify its lines with $\mathbb{P}(H) \simeq \mathbb{P}^1$. A rational map $R : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is represented by a coprime polynomial pair $[p : q]$, and $\deg R = \max(\deg p, \deg q)$. Poles of the affine ratio p/q are ordinary points of the projective map, not physical singularities.

Definition 2.1 (Projective interpolation degree). For the ordered data (ζ_i, Y_i) , let

$$\delta = \min\{\deg R : R : \mathbb{P}^1 \rightarrow \mathbb{P}(H), R(\zeta_i) = Y_i \text{ for every } i\}.$$

Common-denominator lemma. Let S be a rational-inner matrix of McMillan degree at most d , and let $F = Sx$ be any routed column. There are a polynomial χ and vector polynomial n such that

$$F(z) = \frac{n(z)}{\chi(z)}, \quad \deg \chi \leq d, \quad \deg n_j \leq d.$$

The denominator can be chosen nonzero at every regular interpolation node. Indeed, take a stable minimal realization $S(z) = D + zC(I - zA)^{-1}B$ of dimension at most d . With $\chi(z) = \det(I - zA)$,

the adjugate formula gives the stated degree bounds for every entry of Sx . Stability makes χ zero-free on $\overline{\mathbb{D}}$; more generally, after cancelling any removable common factor, it is nonzero at each node where S is regular.

Theorem 2.2 (Exact planar reduction). *If all target lines in (1) lie in one two-plane H , then*

$$\boxed{d_{\min} = \delta.} \quad (2)$$

The equality is independent of the ambient dimension and is constructive.

Proof. We first note that $\delta \leq L - 1$. Apply a projective change of target chart whose point at infinity avoids the finite target set, interpolate the resulting affine values by a polynomial of degree at most $L - 1$, and undo the chart.

For the upper bound, let $[p : q]$ be a coprime representative of degree δ . For a polynomial r of degree at most δ , set

$$r^\sharp(z) = z^\delta \overline{r(1/\bar{z})}.$$

Because p and q are coprime, they have no common zero; in particular $|p|^2 + |q|^2 > 0$ on $\mathbb{T}$. Scalar Fejér–Riesz factorization [12] supplies an outer polynomial h , zero-free in $\mathbb{D}$, such that on $\mathbb{T}$

$$|h|^2 = |p|^2 + |q|^2.$$

Equivalently, $h^\sharp h = p^\sharp p + q^\sharp q$. Therefore

$$S_2(z) = \frac{1}{h(z)} \begin{pmatrix} p(z) & -q^\sharp(z) \\ q(z) & p^\sharp(z) \end{pmatrix} \quad (3)$$

is analytic in $\mathbb{D}$ and unitary on $\mathbb{T}$. Its first column represents $[p : q]$, and

$$\det S_2 = h^\sharp/h.$$

The strict positivity above also gives $h(\zeta_i) \neq 0$ at every routing node. With reversal taken at the displayed order δ , the scalar Blaschke product $h^\sharp/h$ has degree exactly δ ; if $\deg h < \delta$, the missing reversed zeros occur at zero. For a rational-inner matrix the determinant winding equals its McMillan degree [13], so $\deg_{\text{McM}} S_2 = \delta$. Constant unitaries align the input line and the basis of H , while a block identity embeds S_2 into dimension N without adding states. Hence $d_{\min} \leq \delta$.

Conversely, take a minimum-degree interpolant S with degree d . The upper construction already gives $d \leq L - 1 < L$. A stable conservative realization of S and the common-denominator lemma above give every entry of the routed column $F = Sx$ a common denominator of degree at most d , nonzero at the boundary nodes; the corresponding numerators have degree at most d . For every linear functional ℓ annihilating H , the numerator of ℓF vanishes at all $L > d$ distinct nodes. It is therefore zero identically. Thus $F(z) \in H$ as a rational identity. In a basis of H , cancel the common polynomial factor of the two coordinate numerators of F . The resulting coprime pair has degree at most d and interpolates every Y_i , so $\delta \leq d$. This proves (2). $\square$

Remark 2.3 (Affine poles are harmless). The datum $R(z) = 1/z$ has an affine pole at zero, but the projective pair $[1 : z]$ is regular there. Formula (3) lifts this pair to an analytic inner column. Accordingly, pole tests on a chosen affine ratio cannot replace projective coprimality and Fejér–Riesz lifting.

3 Exact certificates and quantitative approximation barriers

Write $Y_i = [a_i : b_i]$ in the chosen basis of H . For $d \geq 0$, define the $L \times 2(d+1)$ matrix

$$M_d(i, :) = (b_i, b_i \zeta_i, \dots, b_i \zeta_i^d, -a_i, -a_i \zeta_i, \dots, -a_i \zeta_i^d). \quad (4)$$

A coefficient vector in $\ker M_d$ is exactly a polynomial pair (p, q) of degree at most d satisfying $[p(\zeta_i) : q(\zeta_i)] = [a_i : b_i]$, provided it has no base point at a node.

Proposition 3.1 (Fail-closed degree certificate). *The number δ is the least d for which $\ker M_d$ contains a pair (p, q) generating the unit ideal, equivalently $\gcd(p, q) = 1$. For a nonconstant pair this can be certified by $\text{Res}(p, q) \neq 0$; constant pairs are checked directly. Such a pair is an exact upper certificate. Full column rank of M_{d-1} is an exact lower certificate.*

Proof. The unit-ideal condition is precisely projective coprimality. When both polynomials form a nonconstant pair, it is equivalent to nonvanishing of the resultant; the constant pairs $[c : 0]$ and $[0 : c]$, $c \neq 0$, are checked by the unit-ideal condition itself. The claim follows from Definition 2.1 and (4). Full column rank excludes every nonzero pair of degree at most $d - 1$. $\square$

The formulation is deliberately algebraic. A kernel vector alone is not enough: a common factor can inflate its displayed degree or introduce a base point. A Bézout identity $up + vq = 1$ is an equivalent independently checkable coprimality certificate.

We now normalize (a_i, b_i) to unit norm and choose an orthonormal basis of H . For a unit vector u and a target line $Y = \mathbb{C}y$, $\|y\| = 1$, write

$$\text{dist}_{\text{ch}}(\mathbb{C}u, Y) = \sqrt{1 - |\langle u, y \rangle|^2}$$

for complex-projective chordal distance.

Theorem 3.2 (Quantitative approximate-routing obstruction). *Suppose M_d has full column rank and put $\sigma = \sigma_{\min}(M_d) > 0$. Every square rational-inner router S of McMillan degree at most d satisfies*

$$\max_{1 \leq i \leq L} \text{dist}_{\text{ch}}(S(\zeta_i)X, Y_i) \geq \frac{\sigma}{\sqrt{L(d+1) + \sigma^2}}. \quad (5)$$

If a perturbed normalized data matrix $\widehat{M}_d$ obeys $\|\widehat{M}_d - M_d\|_2 \leq \eta < \sigma$, the right-hand side remains valid with σ replaced by $\sigma - \eta$. Equivalently, the degree-constrained minimax error $E_d = \inf_{\deg_{\text{McM}} S \leq d} \max_i \text{dist}_{\text{ch}}(S(\zeta_i)X, Y_i)$ obeys the same lower bound.

Proof. Choose a unit input vector x and write $F = Sx$. On $\mathbb{T}$, $\|F\| = 1$. By the common-denominator lemma above, the two coordinates of $P_H F$ have the form $p/\chi, q/\chi$, with numerator degrees at most d . Let c collect their $2(d+1)$ coefficients. If $c = 0$, then F is orthogonal to every Y_i and the left-hand side of (5) is one, so assume $c \neq 0$.

Put $v_i = (p(\zeta_i), q(\zeta_i))$ and $e_i = \text{dist}_{\text{ch}}(\mathbb{C}F(\zeta_i), Y_i)$. Since $v_i = \chi(\zeta_i)P_H F(\zeta_i)$ and $y_i \in H$, the two-dimensional wedge identity gives

$$|(M_d c)_i| = |\det(y_i, v_i)| \leq |\chi(\zeta_i)| e_i, \quad \|v_i\| \geq |\chi(\zeta_i)| \sqrt{1 - e_i^2}.$$

Evaluation of two degree- d polynomials at a unit-modulus node has operator norm $\sqrt{d+1}$, hence $\|v_i\| \leq \sqrt{d+1} \|c\|$. If $\varepsilon = \max_i e_i < 1$, these inequalities imply

$$\|M_d c\|_2 \leq \sqrt{L(d+1)} \|c\| \frac{\varepsilon}{\sqrt{1 - \varepsilon^2}}.$$

The reverse inequality $\|M_d c\|_2 \geq \sigma \|c\|$ and cancellation of $\|c\|$ give (5); the case $\varepsilon = 1$ is immediate. Finally, Weyl's inequality gives $\sigma_{\min}(\widehat{M}_d) \geq \sigma - \eta$, proving the perturbative claim. $\square$

Corollary 3.3 (Exact-arithmetic coercivity certificate). *Let $G_d = M_d^* M_d$, $m = 2(d+1)$, and*

$$\gamma_d = \det G_d \left(\frac{m-1}{\text{tr } G_d} \right)^{m-1}.$$

If M_d has full column rank, then $\sigma^2 \geq \gamma_d > 0$, so the right-hand side of (5) is at least

$$\sqrt{\frac{\gamma_d}{L(d+1) + \gamma_d}}.$$

For Gaussian-rational nodes and targets, γ_d is an exact arithmetic certificate. Any stronger exact inequality $G_d \succeq \alpha I$ replaces γ_d by α .

Proof. If $0 < \lambda_1 \leq \dots \leq \lambda_m$ are the eigenvalues of G_d , then $\det G_d = \lambda_1 \prod_{j>1} \lambda_j$. By arithmetic–geometric mean,

$$\prod_{j>1} \lambda_j \leq \left(\frac{\sum_{j>1} \lambda_j}{m-1} \right)^{m-1} \leq \left(\frac{\text{tr } G_d}{m-1} \right)^{m-1}.$$

Thus $\sigma^2 = \lambda_1 \geq \gamma_d$. Apply Theorem 3.2. $\square$

The normalization is essential: without unit homogeneous representatives and an orthonormal basis of H , σ changes under coordinate rescaling. The theorem now controls routing error itself, rather than only persistence of an exact rank obstruction.

4 The exact zero-memory frontier

The preceding obstruction is a lower bound for general d . At $d = 0$, however, the entire minimax problem has a closed geometric solution. For a unit target representative $y_i \in H$, write

$$\rho_i = y_i y_i^* = \frac{1}{2} (I + \mathbf{r}_i \cdot \boldsymbol{\sigma}), \quad \mathbf{r}_i \in \mathbb{S}^2,$$

where $\boldsymbol{\sigma}$ is the Pauli triple. Define the smallest-cap radius

$$\phi_* = \min_{\mathbf{r} \in \mathbb{S}^2} \max_i \arccos(\mathbf{r} \cdot \mathbf{r}_i). \quad (6)$$

The spherical one-center problem and its support algorithms are classical [14]; the point here is their exact passive-routing consequence.

Theorem 4.1 (Exact zero-memory minimax law). *For every finite planar line-routing table,*

$$\boxed{E_0 = \sin(\phi_*/2)}. \quad (7)$$

Moreover, an optimizing cap center has a first-order support certificate involving at most three active target lines: the origin lies in the convex hull of their tangent projections at the center. Consequently E_0 is computable by candidate centers generated by one-, two-, and three-target supports, together with a global containment check and the usual antipodal degeneracies.

Proof. Every constant unitary can send X to an arbitrary output line, and every output line extends to a constant unitary. It is enough to optimize over output lines in H . Indeed, if a unit output vector $u \notin H$ has $P_H u \neq 0$, then $v = P_H u / \|P_H u\|$ satisfies, for every $y_i \in H$,

$$|\langle v, y_i \rangle| = \frac{|\langle u, y_i \rangle|}{\|P_H u\|} \geq |\langle u, y_i \rangle|.$$

If $P_H u = 0$, every chordal error is one and any target line is no worse. Thus projection and normalization never increase the objective. If the resulting line has Bloch vector $\mathbf{r}$, then

$$|\langle y, y_i \rangle|^2 = \text{tr}(\rho \rho_i) = \frac{1 + \mathbf{r} \cdot \mathbf{r}_i}{2}, \quad \text{dist}_{\text{ch}}(\mathbb{C}y, Y_i) = \sin\left(\frac{1}{2} \arccos(\mathbf{r} \cdot \mathbf{r}_i)\right).$$

Since the last expression is increasing in the spherical angle, (7) follows from (6).

For the support statement, maximize $\tau(\mathbf{r}) = \min_i \mathbf{r} \cdot \mathbf{r}_i$ on the sphere and let A be the active set at a maximizer $\mathbf{r}_*$. If zero were outside the convex hull of the tangent projections $P_{\mathbf{r}_*^\perp} \mathbf{r}_i$, $i \in A$, a strictly separating tangent direction would increase every active scalar product to first order. The inactive constraints have positive slack, contradicting maximality. Planar Carathéodory therefore supplies a balancing subset of at most three active projections. Enumerating those support equations and retaining only centers whose caps contain every target gives the stated finite computation. $\square$

Corollary 4.2 (Two-target frontier and sharp collision limit). *For two distinct target lines with Fubini–Study angle $\alpha = \arccos |\langle y_1, y_2 \rangle| \in (0, \pi/2]$,*

$$E_0 = \sin(\alpha/2).$$

If $t = \tan(\alpha/2)$, then

$$E_0^2 = \frac{t^2}{1+t^2}, \quad \left(\frac{\sigma_{\min}(M_0)}{\sqrt{2 + \sigma_{\min}(M_0)^2}} \right)^2 = \frac{t^2}{1+2t^2}.$$

Hence the ratio between the singular-value lower bound of Theorem 3.2 and the exact optimum tends to one as $t \downarrow 0$.

Proof. The two Bloch vectors are separated by angle 2α , so their smallest cap is centered at the geodesic midpoint and has radius α . This proves the first formula. The two normalized rows of M_0 have Gram eigenvalues $1 \pm |\langle y_1, y_2 \rangle|$, whence $\sigma_{\min}(M_0)^2 = 1 - \cos \alpha$. Substitution and the half-angle identity give the displayed rational formulas and their limit. $\square$

5 Fiber multiplicity and binary routing

Projective interpolation makes collisions transparent. If a nonconstant rational map has degree d , no fiber contains more than d distinct points. This observation becomes a state lower bound after Theorem 2.2.

Proposition 5.1 (Fiber occupancy bound). *Suppose at least two distinct target lines are used, and let $n_{\max} = \max_Y \#\{i : Y_i = Y\}$. Then*

$$d_{\min} \geq n_{\max}.$$

Proof. A constant projective map cannot hit two targets. For a nonconstant map $R = [p : q]$ of degree d , the equation $R(z) = Y$ is a nonzero polynomial equation of degree at most d . It has at most d distinct solutions. Apply this to a most occupied target and use Theorem 2.2. $\square$

For exactly two targets, the fiber obstruction is the whole answer.

Theorem 5.2 (Exact binary collision law). *If the table uses exactly two distinct target lines with occupancies n_0 and n_∞ , then*

$$d_{\min} = \max(n_0, n_\infty). \tag{8}$$

Proof. Normalize the two targets to 0 and ∞ . Let I_0, I_∞ be their node sets. The coprime pair

$$p(z) = \prod_{i \in I_0} (z - \zeta_i), \quad q(z) = \prod_{i \in I_\infty} (z - \zeta_i)$$

has degree $\max(n_0, n_\infty)$ and realizes the table. The reverse inequality is Proposition 5.1. $\square$

The binary law is the collision-dominated extreme: the constant-detector bound is sharp. The opposite extreme has all targets distinct. Then fiber multiplicity contributes only one, but global projective compatibility produces a growing cost.

Table 1: Exact planar mechanisms. The projective reduction converts every entry in the last column directly into passive state count.

Target pattern	Exact or generic degree	Governing obstruction
One target	0	constant projective map
Two targets, occupancies n_0, n_∞	exact max; border min	fiber/base-point duality
L distinct, generic	$\lceil (L-1)/2 \rceil$	projective parameter count
Exceptional table at $d_0 = \lceil (L-1)/2 \rceil$	$E_{d_0} = 0$, possibly unattained	dense exact-routing locus
Four distinct, exceptional	1 or 2	ordered cross-ratio
Four targets, pattern 3 + 1	3	triple fiber

6 The generic planar law and an unbounded gap

Lemma 6.1 (Nonempty coprime threshold locus). *Fix distinct nodes and put $d_0 = \lceil (L-1)/2 \rceil$. There is a nonempty Zariski-open set of target data for which M_{d_0} contains a coprime kernel pair. This open set meets the locus of pairwise distinct targets.*

Proof. Work first in the affine target chart $a_i/b_i = y_i$ and use the witness $y_i = \zeta_i^{d_0}$. At this point the columns of M_{d_0} are evaluations of monomials with exponents $0, \dots, 2d_0$, with exponent d_0 duplicated. Because $L \leq 2d_0 + 1$, a Vandermonde $L \times L$ minor D is nonzero. On the principal open set $U = \{D \neq 0\}$, this same pivot solves L coefficients by Gaussian elimination and expresses the remaining coefficients as free parameters with entries in the localization $\mathbb{C}[y_1, \dots, y_L]_D$. Hence $\ker M_{d_0}$ has an algebraic frame on U .

At the witness, $(z^{d_0}, 1)$ lies in the kernel. Choose the constant linear combination of the local frame that equals this vector there. Multiplying its two polynomial coordinates by a common power of D clears every parameter denominator. Their resultant is then a regular function of the target data; at the witness it is a nonzero scalar multiple of $\text{Res}(z^{d_0}, 1) = 1$. Its nonvanishing and $D \neq 0$ therefore define a nonempty Zariski-open coprime locus. The pairwise target inequalities also define a nonempty Zariski-open subset of $(\mathbb{P}^1)^L$. Their intersection is nonempty because $(\mathbb{P}^1)^L$ is irreducible. $\square$

Theorem 6.2 (Generic planar memory). *For fixed distinct nodes and L target lines in a two-plane, there is a Zariski-open dense set of ordered target data on which*

$$d_{\min} = \left\lceil \frac{L-1}{2} \right\rceil. \quad (9)$$

The open set can be intersected with the locus of pairwise distinct targets.

Proof. Put $d_0 = \lceil (L-1)/2 \rceil = \lfloor L/2 \rfloor$. For $d < d_0$, the matrix M_d has at most L columns and is generically full-column-rank. A concrete rank witness is obtained from affine data $a_i/b_i = \zeta_i^{d_0}$: an equation $p(\zeta_i) = \zeta_i^{d_0} q(\zeta_i)$ at all nodes would give L roots to a polynomial of degree at most $L-1$ and then force $p = q = 0$ at degree $d_0 - 1$. Thus the nonvanishing of a maximal minor supplies the required nonempty open set.

At $d = d_0$, M_{d_0} has more columns than rows, so it always has nonzero kernel. Lemma 6.1 supplies a nonempty open locus on which that kernel contains a coprime pair, and the lower-rank open condition above can be intersected with it. Apply Theorem 2.2 and Proposition 3.1. The scalar threshold itself is classical; the new conclusion here is its exact conversion to passive state count. $\square$

Definition 6.3 (Border memory). For $A \subseteq [L] = \{1, \dots, L\}$, let $\delta(A)$ denote the exact projective interpolation degree of the subtable indexed by A , with $\delta(\emptyset) = 0$. Define

$$d_{\partial} := \min\{d \geq 0 : E_d = 0\}.$$

Thus d_∂ is the first state count at which exact target tables lie in the closure of the degree-bounded passive class; it need not be attained.

Theorem 6.4 (Exact base-point deletion law). *Every finite planar routing table satisfies*

$$d_\partial = \min_{B \subseteq [L]} \{|B| + \delta(B^c)\}. \quad (10)$$

Consequently the complete sign-and-attainment diagram is

$$\begin{array}{c|c} d < d_\partial & E_d > 0, \\ d_\partial \leq d < \delta & E_d = 0 \text{ and the infimum is not attained,} \\ d \geq \delta & E_d = 0 \text{ and is attained.} \end{array}$$

In particular, the distinction between exact and approximate memory is itself an exactly computable projective invariant.

Proof. First suppose $E_d = 0$. Choose degree-at-most- d routers S_n whose worst-node errors tend to zero, and fix a unit vector spanning X . Project $S_n x$ orthogonally to H and use the common-denominator lemma to write its two coordinates as a polynomial pair (p_n, q_n) of degree at most d , up to a scalar denominator. For all large n this pair is not identically zero, because its values at the routing nodes approach nonzero lines in H . Normalize its coefficient vector and pass to a convergent subsequence. The limit is a nonzero pair (p, q) of degree at most d .

Let B be the set of nodes at which p and q both vanish. Write $(p, q) = g(\tilde{p}, \tilde{q})$ with the reduced pair coprime, and put $b = \deg g$ and $r = \max(\deg \tilde{p}, \deg \tilde{q})$. Distinctness of the nodes gives $|B| \leq b$, while coefficient convergence and projective continuity show that $[\tilde{p} : \tilde{q}]$ interpolates every target outside B . Therefore

$$|B| + \delta(B^c) \leq b + r = \max(\deg p, \deg q) \leq d.$$

This proves that the right side of (10) is no larger than d_∂ .

Conversely fix B , put $b = |B|$, and choose a coprime pair (p_0, q_0) of degree $r = \delta(B^c)$ interpolating the complementary subtable. If $b = 0$, the exact lift already gives $E_r = 0$. If $b > 0$, let $g(z) = \prod_{i \in B} (z - \zeta_i)$ and first choose a coprime pair (a_0, c_0) of degree at most $b - 1$ interpolating the data restricted to B ; ordinary polynomial interpolation supplies such a pair. Put $D = b + r$. Without changing its values on B , extend it to a coprime pair (a, c) of exact degree D . If $c_0 \neq 0$, take $c = c_0$ and $a = a_0 + \lambda g z^r$: only finitely many nonzero values of λ can make a vanish at a root of c_0 . If $c_0 = 0$, coprimality forces a_0 to be a nonzero constant, and $(a, c) = (a_0, g z^r)$ works. Consider

$$P_t = (1 - t)g p_0 + t a, \quad Q_t = (1 - t)g q_0 + t c.$$

At every node in B , $[P_t : Q_t] = [a : c]$ for $t \neq 0$; outside B the values converge to $[p_0 : q_0]$ as $t \rightarrow 0$. Coprimality with no common projective zero is detected by the homogeneous degree- D resultant. Since $(P_1, Q_1) = (a, c)$ is coprime and has exact degree D , this resultant is a nonzero polynomial in t . Hence a sequence $t_n \rightarrow 0$, $t_n \neq 0$, can be chosen away from its finitely many zeros, with every (P_{t_n}, Q_{t_n}) coprime. Its degree is at most $b + r$. Applying the exact same-degree lossless lift proves $E_{b+r} = 0$, and then (10).

Finally, $E_d = 0$ is monotone in d . Zero is attained exactly when an exact router of degree at most d exists, which by Theorem 2.2 is equivalent to $d \geq \delta$. The three regimes follow. $\square$

Corollary 6.5 (Finite exact certificate for border memory). *The integer d_∂ is decidable by finite exact linear algebra and polynomial gcd tests. Explicitly, for each $A \subseteq [L]$, compute $\delta(A)$ from the first interpolation matrix $M_d(A)$ whose kernel contains a coprime pair, and minimize $L - |A| + \delta(A)$. A minimizing subset and its coprime kernel pair certify the upper bound; the rank/subresultant exclusions for all smaller subset costs certify the lower bound.*

Proof. Proposition 3.1 computes every $\delta(A)$ using $0 \leq d \leq |A| - 1$, and Theorem 6.4 gives the stated minimum. All tests are exact over the coefficient field of the data. $\square$

Corollary 6.6 (Complete binary approximation phase diagram). *For two target lines with positive occupancies n_0, n_∞ ,*

$$d_\partial = \min(n_0, n_\infty), \quad d_{\min} = \max(n_0, n_\infty). \quad (11)$$

Thus $E_d > 0$ below the smaller occupancy, $E_d = 0$ but is unattained between the two occupancies, and exact routing begins at the larger occupancy. For a $1 + (L - 1)$ split the exact-to-border memory ratio is $L - 1$ and is unbounded.

Proof. Deleting all nodes carrying the less frequent target leaves a constant subtable, so Theorem 6.4 gives the upper bound $d_\partial \leq \min(n_0, n_\infty)$. For the reverse bound, fix B in (10). If the reduced interpolant on B^c is constant, B must contain every occurrence of the other target and therefore $|B|$ is at least the smaller occupancy. If it is nonconstant of degree r , each of its two fibers has at most r remaining nodes. Adding the deleted nodes shows $|B| + r \geq \max(n_0, n_\infty)$, hence certainly at least the smaller occupancy. Minimizing over B proves the first identity; the second is Theorem 5.2, and the phase statement follows from Theorem 6.4. $\square$

Theorem 6.7 (Sharp universal closure threshold). *Fix distinct nodes and put $d_0 = \lceil (L - 1)/2 \rceil$. For every ordered planar target table,*

$$\boxed{E_{d_0} = 0}. \quad (12)$$

More generally $E_d = 0$ for every $d \geq d_0$. The infimum at d_0 is attained if and only if $\delta \leq d_0$. For every $d < d_0$, by contrast, there is a Euclidean-open dense set of target tables for which $E_d > 0$. Consequently d_0 is the sharp state threshold at which degree-bounded passive routers become dense in all planar routing data.

Proof. By Theorem 6.2, the target tables admitting an exact degree- d_0 projective interpolant contain a Zariski-open dense subset of $(\mathbb{P}^1)^L$. Its complement is a proper complex algebraic set and therefore has empty Euclidean interior. Given arbitrary targets (Y_i) , choose exact-routing tables $(Y_i^{(n)})$ in that open set converging to (Y_i) . Apply the same-degree lossless lift in the proof of Theorem 2.2 to each table. The resulting degree- d_0 routers satisfy

$$\max_i \text{dist}_{\text{ch}}(S_n(\zeta_i)X, Y_i) = \max_i \text{dist}_{\text{ch}}(Y_i^{(n)}, Y_i) \longrightarrow 0,$$

which proves (12). Allowing more states cannot increase the infimum. If the infimum at d_0 is attained with value zero, the attaining router interpolates exactly, so Theorem 2.2 gives $\delta \leq d_0$; the converse is immediate. Finally, for $d < d_0$ the full-column-rank condition on M_d is a nonempty Zariski-open condition by the proof of Theorem 6.2. It is Euclidean open dense, and Theorem 3.2 supplies a strictly positive error there. $\square$

Corollary 6.8 (Exact one-state infimum without a minimizer). *For three distinct nodes and any planar target table, $E_1 = 0$. If the target multiplicity pattern is $2 + 1$, then $d_{\min} = 2$ and this zero infimum is not attained by a one-state router.*

Proof. Here $d_0 = 1$, so Theorem 6.7 gives $E_1 = 0$. The binary collision law gives $d_{\min} = 2$ for the $2 + 1$ pattern, and the attainment statement follows. For an explicit normalized example, choose a projective coordinate on the domain taking the nodes to $(0, 1, \infty)$ and choose limiting targets $(0, 0, 1)$ in an orthonormal affine chart of H . For $0 < \varepsilon < 1$,

$$R_\varepsilon(z) = \frac{\varepsilon z}{\varepsilon z + 1 - \varepsilon}$$

is a degree-one Möbius map with values $(0, \varepsilon, 1)$ and worst-node chordal error $\varepsilon/\sqrt{1 + \varepsilon^2}$. Its determinant $\varepsilon(1 - \varepsilon)$ is nonzero, so the same-degree lossless lift is valid. The errors tend to zero, whereas a nonconstant degree-one map is injective and cannot attain the repeated limiting target. $\square$




Figure 1: Left: generic exact/border memory and the binary $1 + (L - 1)$ split, whose exact memory is $L - 1$ while its border memory remains one; the shaded exact-to-border gap is unbounded. Right: exact certified worst-node chordal-error barriers for the strict four-node fixture. Degree two attains zero error, whereas degrees zero and one remain quantitatively separated.

Remark 6.9 (Why rank deficiency is not exact feasibility). At d_0 , the matrix M_{d_0} always has a nonzero kernel, but exceptional tables may have only kernel pairs with a common factor or a base point. Theorem 6.7 shows that such defective pairs are limits of coprime pairs even when no coprime pair exists at the limiting data. Thus full column rank in Theorem 3.2 is a structural separation condition, not a technical artifact: once it fails at the universal threshold, a positive error floor cannot hold for the full degree-bounded class.

For distinct lines in a two-plane, the span bound is $\dim H - 1 = 1$. Likewise, the optimized constant-detector hierarchy from [1] equals one: the kernel of a nonzero functional on H contains at most one of the distinct target lines.

Corollary 6.10 (Unbounded detector gap). *On the generic pairwise-distinct planar locus,*

$$\frac{d_{\min}}{\max\{\text{span bound}, \text{constant-detector bound}\}} = \left\lceil \frac{L - 1}{2} \right\rceil \rightarrow \infty.$$

Thus the optimized constant-detector hierarchy of [1] cannot approximate the exact passive memory within a universal multiplicative factor on this single fixed-rank stratum.

7 The complete four-line phase diagram

For four ordered distinct nodes, write

$$[z_1, z_2; z_3, z_4] = \frac{(z_1 - z_3)(z_2 - z_4)}{(z_1 - z_4)(z_2 - z_3)}$$

with the usual projective conventions at infinity. The order matters because each frequency is paired with a specified target. The projective classification below is classical/elementary; our claim is its exact consequence for square passive rational-inner routers.

Theorem 7.1 (Four-line classification). *For four target lines in one two-plane, the exact passive memory is*

$$d_{\min} = \begin{cases} 0, & \text{all four targets coincide,} \\ 1, & \text{all four are distinct and their ordered cross-ratio equals that of the nodes,} \\ 3, & \text{the target multiplicities are } 3 + 1, \\ 2, & \text{in every remaining case.} \end{cases} \quad (13)$$

Proof. A nonconstant degree-one map is a projective bijection, so it sends the four nodes to four distinct targets and preserves their ordered cross-ratio. These conditions are also sufficient by the uniqueness of a projective map through three ordered points.

If three nodes share a target and the fourth does not, a nonconstant map of degree at most two cannot work because every fiber has at most two distinct points. Polynomial interpolation gives the universal upper bound three.

It remains to show degree at most two in all other nonconstant cases. For four distinct targets with mismatched cross-ratio, normalize domain and target triples to $(0, \infty, 1)$, leaving $s \mapsto t$ with $s \neq t$. Then

$$R(z) = z(az + b), \quad a = \frac{t - s}{s(s - 1)}, \quad b = 1 - a$$

has degree two and realizes the data. For multiplicities $2 + 2$, after choosing a finite chart use

$$R(z) = c \frac{(z - z_1)(z - z_2)}{(z - z_3)(z - z_4)}.$$

For $2 + 1 + 1$, normalize the repeated nodes to $0, \infty$, the other nodes to $1, s$, and the targets to $\infty, \infty, 0, 1$. Choose $a \notin \{0, 1, s\}$ and set

$$q(z) = z, \quad p(z) = A(z - 1)(z - a), \quad A = \frac{s}{(s - 1)(s - a)}.$$

This is a coprime degree-two witness. None of these cases is constant or degree one, completing the lower bounds and the classification via Theorem 2.2. $\square$

Example 7.2 (Strict exact and approximate detector failure). Take

$$(\zeta_1, \zeta_2, \zeta_3, \zeta_4) = (1, i, -1, -i), \quad (Y_1, Y_2, Y_3, Y_4) = (0, 1, \infty, 2).$$

The ordered cross-ratios are 2 and $1/2$, respectively. Hence $d_{\min} = 2$, while both constant lower bounds equal one. An exact witness is

$$p(z) = -\frac{(z - 1)(3z - i)}{2}, \quad q(z) = z + 1, \quad \text{Res}(p, q) = -3 - i \neq 0.$$

It takes the four requested projective values exactly.

There is also a finite approximation gap below degree two. Normalize the four target representatives to unit norm. For $d = 1$, exact arithmetic gives

$$G_1 = M_1^* M_1 = \begin{pmatrix} 17/10 & 1 + 3i/10 & -9/10 & -i/10 \\ 1 - 3i/10 & 17/10 & i/10 & -9/10 \\ -9/10 & -i/10 & 23/10 & -1 - 3i/10 \\ i/10 & -9/10 & -1 + 3i/10 & 23/10 \end{pmatrix}.$$

The leading principal minors of $G_1 - \frac{3}{8}I$ are

$$\frac{53}{40}, \quad \frac{213}{320}, \quad \frac{637}{2560}, \quad \frac{213}{20480},$$

all positive. Sylvester's criterion yields $G_1 \succ \frac{3}{8}I$, and Theorem 3.2 therefore forces

$$\max_i \text{dist}_{\text{ch}}(S(\zeta_i)X, Y_i) > \sqrt{\frac{3}{67}} \approx 0.2116$$

for every router with at most one state. Likewise, $G_0 - I \succ 0$ gives the degree-zero barrier $1/\sqrt{5} \approx 0.4472$. The degree-two witness above attains zero, producing a certified finite-error staircase rather than only an exact degree jump.

8 Delay price and reproducibility

For a differentiable lossless boundary response, define the Wigner–Smith matrix

$$Q(\theta) = -iS(e^{i\theta})^* \frac{d}{d\theta} S(e^{i\theta}).$$

Jacobi’s formula gives $\text{tr}(S^{-1}\partial_\theta S) = \partial_\theta \log \det S$. Together with Potapov’s factorization and winding count, this gives

$$\int_0^{2\pi} \text{tr} Q(\theta) d\theta = 2\pi \deg_{\text{McM}} S$$

for finite rational-inner S [15, 16]. Combining it with Theorem 2.2 yields an exact action law.

Corollary 8.1 (Optimal integrated delay). *For every planar line-routing table,*

$$\inf_S \int_0^{2\pi} \text{tr} Q(\theta) d\theta = 2\pi\delta.$$

The explicit completion (3) attains the infimum.

The computational artifact uses exact arithmetic over $\mathbb{Q}(i)$. It checks the strict witness in Example 7.2; exact representatives of the constant, cross-ratio-compatible, $2+2$, $2+1+1$, and $3+1$ strata; the normalized Gram matrices and Sylvester coercivity certificates behind the degree-zero and degree-one approximation barriers; the exact two-target zero-memory formulas and their sharp-collision ratios at four rational parameters; the explicit nonattained $E_1 = 0$ degeneration at four rational values of ε ; full-column-rank exclusions below the generic threshold; coprime threshold witnesses for every $3 \leq L \leq 16$; and every nontrivial binary occupancy split through $L = 16$ (119 exact cases). For every $3 \leq L \leq 16$ it additionally checks the complete binary positive-error/nonattainment/exact phase partition and an explicit degree-one degeneration of the $1 + (L - 1)$ family, whose exact-to-border ratio is $L - 1$. The frozen validator recomputes the content hash and checks declared coverage, phase degrees, ranks, and bounds. In CI the producer is replayed from scratch and its byte-identical JSON is compared with the frozen record; this replay is the algebraic check of cross-ratios, witness values, gcds, and resultants. These calculations audit examples and certificates; they are not substitutes for the proofs.

Limitation 8.2 (Exact scope). The theorem assumes distinct frequency nodes, one input line, target lines in one common two-plane, free endpoint phases, and square finite rational-inner competitors. Repeated nodes require confluent interpolation. Fixed frames, higher-rank subspaces, lossy or active architectures, and a complete equality theory for nonplanar singular arrangements are not claimed. The quantitative obstruction is a computable lower bound, while Theorem 4.1 solves the full minimax problem at degree zero and Theorem 6.7 solves the universal zero-infimum regime $d \geq \lceil (L - 1)/2 \rceil$. More generally, Theorem 6.4 determines exactly whether E_d is positive, zero but unattained, or attained for every planar table. Determining the numerical positive values of E_d for $1 \leq d < d_\partial$ remains open.

Limitation 8.3 (Priority scope). Scalar minimal rational interpolation, the generic half-degree threshold, cross-ratio and fiber classifications, subspace Nevanlinna interpolation, Fejér–Riesz/paraunitary completion, and matrix all-pass construction predate this work; common-factor/base-point degeneration is likewise a standard algebraic mechanism. We claim the passive-memory equality (2), obtained by combining a standard exact-degree lossless lift with the escape-from-the-plane lower bound, the quantitative low-degree obstruction (5), and the exact zero-memory consequence (7), together with their physical/detector consequences, the exact base-point deletion law (10), and the sharp universal closure theorem (12). Bloch-sphere cap geometry itself is standard; we do not claim a new general rational interpolation algorithm or priority for the scalar strata.

Immutable reproducibility record. The tagged release containing the paper, exact producer, frozen validator, frozen JSON certificate, and figure is:

GitHub release: `v3.4-planar-projective-memory`

9 Conclusion

Coplanarity does not make passive routing cheap. It changes the governing invariant. Once every target lies in a two-plane, the exact state count is the degree of an ordered projective rational interpolant. Fejér–Riesz lifting turns that algebraic object into a lossless network without spending an extra state, while a common-denominator argument proves that no smaller network can escape the plane. The result exposes a generic half-linear memory law in a regime where the optimized constant-detector hierarchy reports one, classifies the first nontrivial cross-ratio strata completely, and identifies the same integer as optimal integrated delay divided by 2π . More strongly, full-rank interpolation matrices impose an explicit nonzero chordal-error floor on every low-state lossless router, stable under calibrated perturbations. The strict four-node fixture turns the exact jump into a certified 21% one-state approximation barrier. At zero memory the approximation problem is solved exactly by the smallest Bloch-sphere cap, and the singular-value bound has the correct sharp collision asymptotics for two targets. At the positive threshold $d_0 = \lceil (L - 1)/2 \rceil$, every planar table has zero minimax infimum, even when exact interpolation needs more states; below that threshold generic data retain a positive error floor. More generally, the base-point deletion formula computes the first zero-error boundary for every planar table and thereby classifies all three regimes exactly. Its binary specialization gives border memory $\min(n_0, n_\infty)$ against exact memory $\max(n_0, n_\infty)$, so approximation can save an unbounded factor of states. This sharp transition separates approximation from exact realization and gives explicit nonattained low-state optima in arbitrarily large tables. The analysis also marks the next frontier precisely: higher-dimensional singular target varieties motivate detectors that move with frequency, rather than merely more constant minors.

References

- [1] Lluís Eriksson. Exact memory of finite spectral routing tables: The direct-sum occupancy law and its singular strata. GitHub release, 2026. <https://github.com/lluise Eriksson/finite-sample-spectral-certificates/releases/tag/v2.8-ai-vixra-submission>.
- [2] Athanasios C. Antoulas and Brian D. O. Anderson. On the scalar rational interpolation problem. *IMA Journal of Mathematical Control and Information*, 3(2–3):61–88, 1986.
- [3] Brian D. O. Anderson and Athanasios C. Antoulas. Rational interpolation and state-variable realizations. *Linear Algebra and its Applications*, 137/138:479–509, 1990.
- [4] Athanasios C. Antoulas, Joseph A. Ball, Jeongook Kang, and Jan C. Willems. On the solution of the minimal rational interpolation problem. *Linear Algebra and its Applications*, 137/138:511–573, 1990.
- [5] Teresa Cortadellas Benítez, Carlos D’Andrea, and Eulàlia Montoro. Minimal solutions of the rational interpolation problem. *Revista de la Unión Matemática Argentina*, 61(2):413–429, 2020.
- [6] Christer Glader. Rational unimodular interpolation on the unit circle. *Computational Methods and Function Theory*, 6(2):481–492, 2006.
- [7] Gunter Semmler and Elias Wegert. Boundary interpolation with Blaschke products of minimal degree. *Computational Methods and Function Theory*, 6(2):493–511, 2006.
- [8] Vladimir Bolotnikov. Boundary interpolation by finite Blaschke products. In *Complex Analysis and Dynamical Systems: New Trends and Open Problems*, Trends in Mathematics, pages 39–65. Birkhäuser/Springer, Cham, 2018.
- [9] Paolo Rapisarda and Jan C. Willems. The subspace Nevanlinna interpolation problem and the most powerful unfalsified model. *Systems & Control Letters*, 32(5):291–300, 1997.
- [10] Vladimir Bolotnikov and Harry Dym. *On Boundary Interpolation for Matrix Valued Schur Functions*, volume 181 of *Memoirs of the American Mathematical Society*. American Mathematical Society, 2006. Memoir no. 856.
- [11] Agulla Surya Bharath, Devanshu Singh Gaharwar, Kumar Appaiah, and Debasattam Pal. Design of discrete-time matrix all-pass filters using subspace Nevanlinna–Pick interpolation. *Signal Processing*, 204:108839, 2023.
- [12] Michael A. Dritschel and James Rovnyak. The operator Fejér–Riesz theorem. In *A Glimpse at Hilbert Space Operators: Paul R. Halmos in Memoriam*, volume 207 of *Operator Theory: Advances and Applications*, pages 223–254. Birkhäuser Basel, 2010.
- [13] V. P. Potapov. The multiplicative structure of J -contractive matrix functions. *Trudy Moskovskogo Matematicheskogo Obshchestva*, 4:125–236, 1955. English translation in AMS Translations, Series 2, Vol. 15.
- [14] Jens Flemming. A simple linear time algorithm for smallest enclosing circles on the (hemi)sphere. *arXiv preprint arXiv:2407.19840*, 2026.
- [15] Eugene P. Wigner. Lower limit for the energy derivative of the scattering phase shift. *Physical Review*, 98:145–147, 1955.
- [16] Felix T. Smith. Lifetime matrix in collision theory. *Physical Review*, 118:349–356, 1960.