

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339882892>

Sedentary behavior and health outcomes in older adults: A systematic review

Article in AIMS Medical Science · January 2020

DOI: 10.3934/medsci.2020002

CITATIONS

16

READS

189

4 authors, including:



[Raheem J Paxton](#)

University of Alabama

89 PUBLICATIONS 2,057 CITATIONS

SEE PROFILE

Review

Sedentary behavior and health outcomes in older adults: A systematic review

Wendell C. Taylor^{1,*}, Kevin Rix², Ashley Gibson³ and Raheem J. Paxton⁴

¹ Institute for the Medical Humanities, Department of Preventive Medicine and Population Health, The University of Texas Medical Branch, Galveston, TX, USA

² Department of Health Promotion and Behavioral Sciences, School of Public Health, The University of Texas Health Science Center at Houston, Austin, TX, USA

³ Department of Graduate Studies, Cizik School of Nursing, The University of Texas Health Science Center at Houston, Houston, TX, USA

⁴ Department of Community Medicine and Population Health, Institute for Rural Health Research, University of Alabama, Tuscaloosa, AL, USA

* **Correspondence:** Email: wetaylor@utmb.edu.

Abstract: *Introduction:* Older adults (≥ 60 years old) report prolonged periods of sedentary behavior. Sedentary behavior is a potential health hazard for this priority population. Therefore, we systematically reviewed the published literature to document the relationships among sedentary behaviors and twelve health outcomes ranging from mental health to mortality. *Methods:* Major databases were searched from 2013 to 2019; 27 relevant articles were found and evaluated. In addition, we compared our findings to a previously published review. *Results:* Higher levels of sedentary behavior were related to an increased risk of all-cause mortality and adversely associated with metabolic syndrome, triglycerides/high density lipoprotein cholesterol/blood glucose, HBA1C/glucose intolerance, waist circumference, and obesity/overweight when compared to those with lower levels of sedentary behavior. Findings for blood pressure, cancer, and mental health (e.g., dementia, mild cognitive impairment, psychological well-being) were insufficient to draw conclusions or had inconsistent results. Because some sedentary behaviors were protective for mental health, we recommend a taxonomy of sedentary behaviors for older adults to provide insights into these seemingly discrepant findings. Some of our findings were similar to a prior review while other findings were different. *Conclusion:* This systematic review identified the health outcomes that were sufficiently, insufficiently, or not affected by sedentary behavior. To advance the field, we recommend better methodological quality. To improve the overall health and wellbeing of older adults, future

studies should evaluate interventions to decrease health-compromising and increase health-promoting sedentary behaviors among older adults.

Keywords: sedentary behavior; health; elderly; older adults; physical inactivity

1. Introduction

It has been estimated that by 2050 older adults (≥ 60 years old) will comprise 22% of the world's population [1]. Older adults face significant challenges related to their social, emotional, and physical health as well as other challenges such as ageism and life transitions (e.g., retirement and grandparent status) not experienced by younger people. Accelerometer data for older adults in the United States indicate that they engage in greater amounts of sedentary behavior than any other age group [2]. Globally, similar results have been reported (in more than 60 countries), whereby older adults are more likely to report sitting for four or more hours per day greater than any other age group [3].

Despite the findings that older adults report more sedentary behavior than other age group, they are underrepresented in studies examining health outcomes associated with sedentary behavior. The most recent Physical Activity Guidelines Report found limited data about the relationship between sedentary behavior, age, and mortality outcomes [4]. Furthermore, there was insufficient evidence about sedentary behavior and age related to cardiovascular disease, type 2 diabetes, and cancer [4]. Therefore, more studies are needed to clarify the relationship between sedentary behaviors and health outcomes in older adults.

In the general population, sedentary behavior has been identified as a risk factor independent of physical activity [5–8]. A central concern is the extent to which sedentary behavior affects the well-being of older adults. To address this issue, there is a need for studies that systematically and comprehensively review the literature. In a narrative review, measurement of sedentary behavior was evaluated, specific associations of sedentary time with geriatric-relevant health outcomes were presented, and interventions that target reducing sedentary time in older adults were analyzed; six health outcomes were reviewed [9]. To our knowledge, there has been only one comprehensive, systematic review of sedentary behavior and multiple health outcomes in older adults. In 2014, de Rezende et al [10] reported the associations between sedentary behaviors and twelve health outcomes in older adults: mortality, metabolic syndrome, cardiometabolic biomarkers, triglycerides, HDL cholesterol, blood pressure, plasma glucose/ Hb1Ac/ glucose intolerance, cholesterol ratio and total, other cardiometabolic biomarkers, waist circumference/waist-to-hip ratio/abdominal obesity, overweight/obesity, and cancer [10]. While this review was comprehensive, in the past six years, additional studies have been published; an update of this review is needed to advance the field. Therefore, the objectives of this systematic review were to: (1) assess the relationship among sedentary behaviors and twelve health outcomes in older adults; (2) compare our findings to previous reviews; and (3) present recommendations to advance the field.

2. Methods

2.1. Definition of sedentary behavior

Sedentary behavior is operationalized as any activity engaged in while sitting including leisure and occupational time. In addition, sedentary behavior is defined as any waking behavior accounting for energy in the range of 1.0 to 1.5 metabolic equivalents while in a sitting or reclining position [4,11]. Sedentary behaviors include television viewing and computer use [11].

2.2. Ethical considerations

In our study, there was no contact with human participants; therefore, this research was exempt from Institutional Review Board approval.

2.3. Data collection

We updated the 2014 review [10], which identified articles up to May of 2013. Therefore, we searched articles from 2013 to June 2019 (six years). Similar to the search strategy of the prior review [10], the following databases were used with the assistance of a professional librarian: Medline, Web of Science, PsycINFO, and Cumulative Index to Nursing and Allied Health Literature (CINAHL). Because of access issues, we did not search the following databases identified in the 2014 review: Excerpta Medica (EMBASE), SPORTDiscus, Literatura Latino-Americana e do Caribe em Ciências da Saúde (LILACS), and Sedentary Behavior Research Database (SBRD). However, to extend and expand our search, we added the following databases not included in the prior review: PubMed, Google Scholar, Scopus, and Academic Search Premier. We chose outcomes that were most relevant for people 60 years and older which included the outcomes chosen in the previous review [10]. Our objective was to capture “geriatric-relevant health outcomes” [9].

As in the previous review [10], the same search keywords were used: “exposures (sedentary behavior, sedentary lifestyles, sitting time, television reviewing, driving, screen-time, video game, and computer); primary outcomes (mortality, cardiovascular disease, cancer, type 2 diabetes mellitus); and secondary outcomes (accidental falls, frail elderly, obesity, metabolic syndrome, mental disorders, musculoskeletal diseases)”. Studies that were excluded: review articles, studies with adults younger than 60 years of age, and descriptive data only studies. In two subsections, there were no recent empirical data. To fill this gap and to provide a context, two systematic reviews/meta-analyses were cited as exceptions to our exclusion criteria.

2.4. Data extraction and quality assessment

Data from eligible articles were extracted by one author and reviewed independently by a second author. The tables present: Author(s), Year, Title, Journal, Hypotheses/Study Aims, Research Design, Sample Size, Ages, Country, Measures, Primary Findings, Limitations, and Strengths.

Because of its credibility and reputation of having “state of art” and “gold standard” assessments, we rated each article based on the U.S. Department of Health & Human Services, National Institutes of Health, Study Quality Assessment Tools [12]. Each research design had a specific assessment tool

to evaluate quality assessment. The research designs were controlled intervention studies (14 questions), systematic reviews and meta-analyses (8 questions), observational cohort and cross-sectional studies (14 questions), case-control studies (12 questions), before-after (pre-post) studies with no control group (12 questions), and case series studies (9 questions). For each category, the response choices were yes, no, or other (e.g., CD, cannot determine; NA, not applicable; NR, not reported). To ensure accurate reporting, each design was accompanied by a document called “guidance for assessing the quality”. This document provided guidelines, rules, and tips to answer specific questions for each research design. The final quality ratings were Good, Fair, or Poor. One of the coauthors rated each study independently. All coauthors rated the same study for consensus if there were concerns or ambiguity about any ratings.

2.5. Final selection of articles

As described in Figure 1, the search started with 10,591 potentially relevant articles. After screening articles for duplicates, review articles, meta-analyses, not relevant areas of interest, and other reasons, 27 articles were reviewed and described. Articles are presented in Tables 1 to 9. In the 2014 review, 23 articles met the inclusion criteria [10].




Figure 1. Flowchart outlining study selection of articles.

3. Results

3.1. Mortality (Table 1)

All-cause mortality is the most severe outcome associated with sedentary behavior. Across all studies, there was a consistent association between sedentary behavior and mortality [13]. Six prospective cohort studies were identified. In one prospective cohort of Spanish older adults, those who were consistently non-sedentary had a lower hazard ratio (HR = 0.74, 95% confidence ratio (CI) = 0.62–0.90) for mortality than those who were consistently sedentary [14]. Studies conducted in both older men and women yielded similar results with respect to mortality risk. Ensrud et al. [15] found that older men who were in the highest quartile of sedentary behavior had a higher risk for all-cause mortality (HR = 1.51, 95% CI: 1.10–2.08) when compared to those in the lowest quartile. Similarly, in a sample of Australian women, there was a dose response relationship between sitting and those who were not meeting physical activity guidelines [16]. In particular, the risk for mortality increased for non-active women who reported 8 or more hours of sedentary time (i.e., for 8 to 11 hours OR = 1.31 and for ≥ 11 hours OR = 1.47) when compared to those who sat less than 4 hours per day ($p < 0.05$). Similarly, in a sample of British men, each additional 30 minutes of sedentary behavior increased the risk for mortality (HR = 1.17 95% CI; 1.10 to 1.25 $p < 0.05$) [17]. Conversely, Klenk et al. [18] found that the association between sedentary behavior and mortality diminished after adjustments for several biomarkers (HR = 2.05 95% CI: 1.13, 3.73 $p < 0.05$). Furthermore, in a sample of older adults from Spain, being in the highest quartile for sedentary behavior increased the risk for mortality due to inflammatory (HR = 1.49 95% CI: 1.08, 2.06) and non-infectious inflammatory causes (HR = 1.59 95% CI: 1.08, 2.33) [19]. However, these models became non-significant when adjusting for physical activity. In examining patterns of physical activity and sedentary behavior together, high levels of sedentary behavior (> 7 hours per day), regardless of physical activity levels were associated with mortality from inflammatory (HR = 1.68 95% CI: 1.19, 2.37) and non-infectious inflammatory causes [19].

3.1.1. Our findings compared to the 2014 review for mortality

In our review, we identified six prospective cohort studies and Rezende et al. 2014 [10] described four prospective cohort studies. The findings are remarkably similar. For older adults, sedentary behavior (particularly sitting) six or more hours per day was consistently related to all-cause mortality. In the studies, the thresholds for sedentary behavior/sitting time ranged from 6, 8, 8 to 11, and 11 or more hours per day.

Table 1. Mortality, sedentary behavior, and older adults (references from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/ Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Cabanas-Sanchez, Guallar-Castillon et al., (2018); Physical activity, sitting time and mortality from inflammatory disease in older adults. <i>Frontiers in Physiology</i>	Study focused on infectious and non-infectious inflammatory causes.	Prospective Cohort study (N = 3,667) Men and women ages 60 years and older in Spain	Sedentary data were self-reported. Mortality identified using the National Death Index.	In nonlinear models, those who were sedentary > 7 hours per day had a higher mortality risk for total inflammatory and non-infectious inflammatory causes. Relationships were stronger for those who were the least active.	Sedentary time and physical activity were self-reported. No dietary data were collected. Inflammatory disease definition may not encompass all related conditions.	Prospective cohort study Spanish population National database to confirm deaths Focus on mortality due to inflammatory and non-inflammatory causes
Ensrud, Blackwell, Cauley et al., (2014); Objective measures of activity level and mortality in older men. <i>Journal of the American Geriatrics Society</i>	Study focused on developing objective measures of activity level for older adult men, related to all-cause mortality.	Prospective Cohort study (N = 2,918) Men ages 71 years and older in the United States	Accelerometer determined sedentary behavior (ACTi Graph) Deaths confirmed via death certificate and follow-up calls.	In adjusted models, men in the highest quartile for sedentary time had a higher risk for all-cause mortality. Subsequent analysis indicated that time spent sedentary was associated with non-cancer and non-CVD deaths.	Associations for women or racial and ethnic minority populations not determined	Objective assessment of behavior Cohort study Disease specific mortality
Jefferis, Parsons, Sartini et al., (2018); Objectively measured physical activity, sedentary behavior and all-cause mortality in older men: Does volume of activity matter more than pattern of accumulation? <i>Br J Sports Med</i>	To examine the association between accelerometer determined sedentary behavior, physical activity, and all-cause mortality	Prospective Cohort study (N = 1,655) Men ages 71–92years old in the United Kingdom	Accelerometer determined sedentary behavior (ACTi Graph) National database confirmed mortality	In adjusted models, those in the highest quartile for sedentary time had a higher risk for all-cause mortality when compared to men in the lowest quartile.	No measure of postural allocation Associations for women or racial and ethnic minority populations not determined	Prospective cohort study. An objective measure of sedentary behavior

Continued on next page

Reference Author(s), Year, Title, Journal	Hypotheses/ Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Klenk, Dallmeier, Denking et al., (2016); Objectively measured walking duration and sedentary behavior and four-year mortality in older people. <i>Plos ONE</i>	This study examined the effect of walking and sedentary time on mortality.	Randomly and geographically selected cohort of older adults in Germany (N = 1,271) aged 65 years or greater.	Physical activity levels were accelerometer determined sedentary behavior (ActivPAL) Mortality determined via local registration offices	Sedentary duration association between sedentary time and mortality diminished in fully adjusted models.	Limited follow-up period Slower walking speeds may have different outcomes	Use of ActivePAL to measure sedentary behavior
Leon-Munoz, Martinez-Gomez, Balboa-Castillo et al., (2013); Continued sedentariness, change in sitting time, and mortality in older adults. <i>Med Sci Sports Exerc</i>	To examine the association between continued sedentary time and changes in sedentary time with mortality	Prospective Cohort (N = 2,635) of adults aged 60 years and older.	Self-reported sedentary time in different activities National Death Index confirmed deaths	Those who were consistently sedentary had higher mortality rates when compared to those who were newly sedentary, formally sedentary, and consistently non-sedentary. Associations maintained in different SES strata.	Self-reported questionnaires Other factors may be related to mortality	Prospective cohort study Mortality confirmed via National Death Index.
Pavey, Peeters, Brown (2015); Sitting-time and 9-year all-cause mortality in older women. <i>Br J Sports Med</i>	This study looked at the relationship between total sitting-time and all-cause mortality in older women.	Prospective cohort study (N = 6,656) of Australian Women ages 76–90.	Self-reported questionnaires Mortality data confirmed by the Australian National Death Index.	In fully adjusted models there was a dose-response relationship between time spent sitting and mortality. However, only the p-value for the trend was significant.	Other factors may be related to mortality. Self-reported questionnaires	Analysis in different SES strata Confirmed mortality data Prospective cohort design

3.2. Metabolic Syndrome (Table 2)

Metabolic syndrome increases the risk of future chronic health conditions and premature mortality. Evidence from cross-sectional studies, longitudinal studies, and randomized controlled trials suggest that sedentary time, moderate-to-vigorous physical activity, and cardio-respiratory fitness are important predictors of various cardio-metabolic risk factors [20]. The main finding was that total daily time spent in sedentary behaviors was associated with increased odds of having metabolic syndrome without accounting for effect modifiers [20]. Evidence showed a 9% increase in the odds of having metabolic syndrome with each additional hour of sitting time. Findings showed a 58% (OR 1.58; 95% CI 1.01, 2.48) increased odds of metabolic syndrome when comparing the highest (> 9.49 h) and the lowest (< 6.70 h) quartiles [20].

3.2.1. Our findings compared to 2014 review for metabolic syndrome

In our review, we identified two studies published in 2015 related to sedentary behavior and metabolic syndrome. Rezende et al. described four studies [10]. The findings indicated that sedentary behavior is adversely and consistently related to metabolic syndrome.

3.3. Triglycerides/HDL cholesterol/ Blood Glucose (Table 2)

One study found statistically significant interactions between sedentary behavior and moderate-to-vigorous intensity physical activity for triglycerides ($p = 0.04$), HDL-cholesterol ($p = 0.01$), fasting blood glucose. In addition, there were increased odds of having low HDL-cholesterol with each hour of sedentary behavior time for those who accumulated 150–300 and ≥ 300 min/week of moderate-to-vigorous intensity physical activity, respectively [21]. Similarly, increased odds of 13% were found (OR 1.13; 95% CI 1.01, 1.26) for having high fasting blood glucose with each hour of sedentary time for those who accumulated ≥ 300 min/ week of moderate-to-vigorous intensity physical activity [21]. Each minute spent in sedentary behavior was significantly associated with reductions of 0.03 mmHg (95%CI: -0.05 ; -0.01) in systolic blood pressure and reductions of 0.02 mg/dL (95%CI: -0.02 ; -0.01) in HDL-cholesterol [22]. Beneficial associations of moderate-to-vigorous intensity physical activity and HDL ($\beta = 0.10$; 95%CI: 0.01; 0.18) and plasma glucose ($\beta = -0.18$; 95%CI: -0.33 ; -0.02) were observed [22]. Similarly, moderate-to-vigorous intensity physical activity was associated with higher levels of HDL ($\beta = -0.35$; 95%CI: 0.14; 0.54), but it was not related to the other cardiovascular risk markers [22]. In a randomized, sedentary behavior reduction trial, Aadahl et al. [23] found that the intervention condition experienced greater improvements in insulin (-5.9 pmol/L, $p = 0.03$), insulin resistance (-0.28 , $p = 0.03$), and waist circumference (-1.42 cm, $p = 0.01$) when compared to the control condition participants.

Table 2. Cardio-metabolic markers, cholesterol ratio and total, metabolic syndrome, triglycerides, HgA1C, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Ekbloom, Ö., Ekbloom-Bak, E., Rosengren, A., Hallsten, M., Bergström, G., Börjesson, M., & Lazzeri, C. (2015); Cardiorespiratory fitness, sedentary behavior and physical activity are independently associated with the metabolic syndrome. Results from the SCAPIS Pilot Study. <i>PLoS ONE</i>	Examine the association between sedentary behavior and metabolic syndrome	Random stratified sample of adults aged 50 to 65 years living in Sweden (N = 1111)	Accelerometer determined sedentary behavior Serum samples used to detect metabolic syndrome	In adjusted models, the highest quartile for sedentary time had higher odds for metabolic syndrome when compared to the first quartile. Sedentary time increased the odds of a larger waist circumference and higher triglyceride, and HDL cholesterol levels.	Cross-sectional design Possibility of multi-collinearity	Adjusted for dietary intake and fitness Accelerometer determined sedentary behavior Random stratified sample
Joseph, J., Echouffo-Tcheugui, J., Golden, S., Chen, H., Jenny, N., Carnethon, M., Bertoni, A. (2016); Physical activity, sedentary behaviors and the incidence of type 2 diabetes mellitus: The Multi-Ethnic Study of Atherosclerosis (MESA). <i>BMJ Open Diabetes Research & Care</i>	Examine the association between sedentary behavior and diabetes	Prospective cohort (N = 5829) of men and women from the United States aged 45–84 years old	Incident type II diabetes determined by fasting glucose Self-reported sedentary time	In adjusted models, both television time and total sedentary behavior increased the odds for type II diabetes. Possible effect modification by family history of diabetes, race, and BMI were observed.	Physical activity and sedentary behaviors were self-reported, and the time frame queried was their typical activities within the last month. Therefore, discrepancies with actual activity levels may exist and self-reported sedentary behavior	Multiethnic population Serum-based biomarkers

Continued on next page

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Wirth, K., Klenk, J., Brefka, S., Dallmeier, D., Faehling, K., Roqué, M., Denking, M. (2017); Biomarkers associated with sedentary behavior in older adults: A systematic review. <i>Ageing Research Reviews</i>	Patho-mechanisms of sedentary behavior (SB) are unclear. We conducted a systematic review to investigate the associations between SB and various biomarkers in older adults.	Electronic databases were searched (MEDLINE, EMBASE, CINAHL, AMED) up to July 2015 to identify studies with objective or subjective measures of SB, sample size ≥ 50 , mean age ≥ 60 years and accelerometer wear time ≥ 3 days. Methodological quality was appraised with the CASP tool.	Randomized controlled trials found a positive correlation for SB with BMI, neck circumference, fat mass, HbA1C, cholesterol and insulin levels, cohort studies additionally for waist circumference, leptin, C-peptide, and low-density lipoprotein and a negative correlation for HDL.	The final sample (26 articles) 63 biomarkers were detected. Most investigated markers were body mass index (BMI, $n = 15$), waist circumference (WC, $n = 15$), blood pressure ($n = 11$), triglycerides ($n = 12$) and high-density lipoprotein (HDL, $n = 15$). Some inflammation markers were identified such as interleukin-6, C-reactive protein or tumor necrosis factor alpha. There was a lack of renal, muscle or bone biomarkers.	First, we identified relatively few high quality or longitudinal studies investigating biomarkers specifically in older adults. Therefore, we were not able to conduct a meta-analysis as we anticipated.	The highlighted results of the four “risk population” studies showed associations for SB with biomarkers in the same direction as the studies performed in non-risk populations.
Figueiró, T. H., Arins, G., Santos, C., Cembranel, F., Medeiros, P. A., d'Orsi, E., & Rech, C. R. (2019); Association of objectively measured sedentary behavior and physical activity with cardiometabolic risk markers in older adults. <i>Plos ONE</i>	Examine the association between sedentary behavior and cardiometabolic markers	Cross-sectional study of Brazilians adults ($N = 425$) aged 60 years and older	Accelerometer determined sedentary behavior Waist circumference, blood pressure, and plasma-based cardio-metabolic markers	Sedentary behavior negatively and linearly associated with systolic blood pressure and HDL cholesterol. Physically inactive and sedentary individuals had greater waist circumference and lower HDL cholesterol levels than active, non-sedentary subjects.	Cross-sectional study design Confounding variables (i.e., medications and BMI) not included in the analysis	Objectively measured sedentary time Plasma-based biomarkers

Continued on next page

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Aadahl, M., Linneberg, A., Møller, T., Rosenørn, S., Dunstan, D., Witte, D., & Jørgensen, T. (2014); Motivational counseling to reduce sitting time: A community-based randomized controlled trial in adults. <i>American Journal of Preventive Medicine</i>	Examine the effectiveness of a sedentary behavior intervention	Randomized trial of 166 adults aged 18–69 years old from Denmark	Objectively measured overall sitting time; secondary measures were breaks in sitting time; anthropometric Accelerometer determined sedentary behavior (ActivPAL) Body fat percentage determined by Tanita BC-420MA Serum measures of insulin homeostasis model assessment, hepatic insulin resistance, and basal insulin secretion, glycosylated hemoglobin, total cholesterol, LDL and HDL cholesterol, and triglycerides	The reduction in sedentary time did not differ between groups. The intervention group had greater improvements in standing time than the control group. Greater improvements in insulin resistance was observed in the intervention condition when compared to the control group. No other differences were observed.	No dietary information was included Uneven sample size in the intervention and control condition.	Accelerometer determined sedentary behavior Serum biological markers Randomized design

3.3.1. Our findings compared to the 2014 review for triglycerides/high density lipoprotein cholesterol/ blood glucose

We identified three relevant studies and, in our review, with the exception of systolic blood pressure, sedentary behavior/sitting was associated with unfavorable biometric indices. In the 2014 review, for two of the three studies, the associations between time spent in sedentary behavior and high triglycerides was not statistically significant. Overall, the literature in this area is not consistent.

3.4. Blood pressure (Table 3)

In a cross-sectional descriptive study (with 306 elderly Koreans with hypertension), levels and predictors of sedentary behavior were examined [24]. The participants spent an average of 8.59 hours per day in sedentary behavior. The analysis showed that sedentary behavior was significantly predicted by a longer duration since hypertension diagnosis, greater levels of depressive symptoms, and lower levels of perceived health, vigorous-intensity physical activity, and empowerment (i.e., a composite score of sense of control, self-efficacy, problem solving, psychosocial coping, support, motivation, and decision making). These variables explained 42.6% of the variance in sedentary behavior [24].

3.4.1. Our findings compared to the 2014 review for blood pressure

In our review, we found one study related to blood pressure. However, blood pressure was not analyzed as a predictor or consequence of sedentary behavior. In the prior review, two of the three publications reported that sedentary time was associated with high blood pressure [10]. This area of research is emerging and more definitive studies are needed.

3.5. HBA1C/Glucose Intolerance (Table 2)

Sedentary behavior has been associated with glucose levels and insulin secretions. Lifestyle interventions designed to reduce sedentary behaviors such as television watching have the potential to improve glucose metabolism. Joseph et al. [25] found that both sedentary behavior (< 2 versus > 6 hours/day: OR = 1.65, 95% CI = 1.26, 2.14) and television viewing (< 2 versus > 6 hours/day: OR = 2.68, 95% CI = 1.38, 5.21) were associated with risk of type II diabetes in adjusted models. The association remained after adjusting for BMI (OR = 1.39, 95% CI = 1.06, 1.81). Some of these relationships differed by racial and ethnic identity.

3.5.1. Our findings compared to the 2014 review for HBA1C/Glucose Intolerance

Since 2013, we found one study that examined the association between sedentary behavior and type 2 diabetes. A positive association was reported between type 2 diabetes risk, leisure sedentary behavior, and television viewing, which may be more pronounced in non-Hispanics whites. In the prior review [10], four studies were found. One study reported a positive association between television viewing, sedentary time and plasma glucose; in another study, a positive relationship was found in women only; in two studies, the findings were not statistically significant [10]. Overall, the findings are mixed and inconsistent related to sedentary behavior, HBA1C/Glucose Intolerance, and diabetes. Clearly, more research is needed.

Table 3. Blood pressure, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/ Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Chang & Sok (2015); Predictors of sedentary behavior in elderly Koreans with hypertension, <i>The Journal of Nursing Research</i>	Examine predictors of sedentary behavior	Cross-sectional study (N = 306) of Korean adults aged 65 years and older	Self-reported sedentary behavior Psychosocial variables (i.e., empowerment, perceived health, depression, self-efficacy, and social support for physical activity)	Stepwise regression analysis indicated that empowerment, duration since HBP diagnosis, perceived health, vigorous-intensity physical activity, and depressive symptoms were determinants of sedentary behavior.	Self-reported measures of sedentary behavior Cross-sectional design	Validated measures

3.6. Cholesterol ratio and total (Table 2)

3.6.1. Our findings compared to the 2014 review for cholesterol ratio and total

Our review found one study with no statistically significant results. Rezende et al. 2014 reported three studies; two studies reported positive results between television viewing, high total-to-HDL cholesterol ratio and leisure-time sedentary behavior, television viewing, and cholesterol ratio [10]. A third study found no statistically significant findings between sedentary behavior and total cholesterol. Overall, two studies including the most recent one (2019) found no association and two earlier studies (2007 and 2012) found positive associations. To clarify the relationships between cholesterol and sedentary behaviors in this emerging area, more definitive research is recommended.

3.7. Other Cardio-metabolic biomarkers (Table 2)

There is a scarcity of studies investigating inflammatory biomarkers and sedentary behavior. Therefore, we included one review article as an exception to our exclusion criteria. Wirth and colleagues (2017) [26] defined sedentary activity as waking behavior with an energy expenditure ≤ 1.5 METS while in a sitting or reclining posture. C-reactive protein, which is a protein made by your liver that is sent into your bloodstream in response to inflammation, was investigated most frequently; although, restricted to four cross-sectional studies and one randomized controlled trial, only two cross-sectional studies demonstrated that sedentary behavior was positively associated with C-reactive protein [26]. Neck circumference and fat mass were positively correlated to sedentary behavior but were investigated in only one randomized controlled trial [27]. There was limited or no evidence for the other anthropometric biomarkers [26].

3.7.1. Our findings compared to the 2014 review for other cardio-metabolic biomarkers

In our review, the findings related to sedentary behavior and C-reactive protein were not consistent. Rezende et al. 2014 reported a study that found a positive relationship between sedentary behavior and C-reactive protein [10]. In three of the five studies, positive associations were found between C-reactive protein and sedentary behavior. Other cardio-metabolic biomarkers such as pericardial fat, coronary artery calcification, neck circumference, and fat mass merit further research.

3.8. Waist circumference (Table 4)

An area of great interest is the impact of sedentary behavior on the waist circumference of older adults. Three cross-sectional studies were identified; sedentary behavior had strong positive associations with waist circumference. In models adjusted for moderate-to-vigorous intensity physical activity, sedentary behavior was positively associated with both weight and waist circumference [27]. Similarly, Bann et al. [28] found that sedentary behavior was inversely associated with body mass index (BMI). In addition, less time spent in sedentary behavior and more time spent in light-intensity physical activities resulted in lower BMIs. These results were partially supported by a cross-sectional study [22].

Table 4. Waist circumference, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages	Measures	Primary Findings	Limitations	Strengths
Bann, Hire, Manini et al. (2015); Light intensity physical activity and sedentary behavior in relation to body mass index and grip strength in older adults: cross-sectional findings from the lifestyle interventions and independence for elders (LIFE) study. <i>Plos ONE</i>	Examine the association between sedentary behavior, BMI, and grip strength.	Cross-sectional study (N = 1,130) of adults aged 70–89 living in the United States.	Accelerometer determined sedentary time Jamar determined grip strength BMI and waist circumference measured by trained staff	In adjusted models, sedentary behavior was associated with BMI. Less time spent sedentary and more time spent in light-intensity physical activities was associated with BMI	Cross sectional design	Use of BMI as a measure of body size
Gennuso, Gangnon Matthews (2013); Sedentary behavior, physical activity, and markers of health in older adults. <i>Medicine and science in sports and exercise</i>	Examine the association between sedentary behavior, physical activity, and cardiometabolic markers	Population based cross-sectional study (N = 1914) of adults aged 65 years and older living in the United States	Accelerometer determined sedentary behavior Cardiometabolic markers (total, HDL, and LDL cholesterol, glucose, triglycerides, glycohemoglobin, waist circumference, blood pressure, and C-reactive protein) Physical function	In adjusted models, higher quartiles of sedentary behavior were associated with weight, BMI, waist circumference, systolic blood pressure, and C-reactive protein among participants who were insufficiently active older adults. Among sufficiently active older adults, higher quartiles of sedentary behavior were associated with waist circumference and LDL cholesterol levels. In addition, insufficiently active older adults who were less sedentary had lower odds of functional limitations.	Cross-sectional design	Accelerometer determined sedentary behavior Population based sample Serum cardiometabolic markers
Figueiró, T. H., Arins, G., Santos, C., Cembranel, F., Medeiros, P. A., d'Orsi, E., & Rech, C. R. (2019); Association of objectively measured sedentary behavior and physical activity with cardiometabolic risk markers in older adults. <i>Plos ONE</i>	Examine the association between sedentary behavior and cardiometabolic markers	Cross-sectional study of Brazilians adults (N = 425) aged 60 years and older	Accelerometer determined sedentary behavior Waist circumference, blood pressure, and plasma based cardiometabolic markers	Sedentary behavior was negatively and linearly associated with systolic blood pressure and HDL cholesterol. Physically inactive and sedentary individuals had greater waist circumference and lower HDL cholesterol levels than active, non-sedentary subjects.	Cross-sectional study design Confounding variables (i.e., medications and BMI) not included in the analysis	Objectively measured sedentary time Plasma-based biomarkers

3.8.1. Our findings compared to the 2014 review for waist circumference

In our review, the findings were consistent; in three studies, sedentary behavior was related to greater waist circumference in older adults. Similarly, in five of six studies, sedentary behavior was related to high waist circumference and high waist-to-hip ratio as reported in the prior review [10]. The only exception to the consistent findings is a study in a colorectal cancer population in which sedentary time was not associated with waist circumference. Overall, however, the research is consistent; sedentary time is a risk factor for greater waist circumference in older adults.

3.9. Obesity/Overweight (Table 5)

The Centers for Disease Control and Prevention classify individuals as obese with a body mass index (BMI) ≥ 30.0 kg/m². A systematic review study indicated that sedentary behavior was linked to increased risk of obesity in various populations [29]. We identified three cross-sectional studies in our search. In an earlier study of older adults in the United States [27], positive associations were found between sedentary behavior and BMI. Similarly, Bann et al. [28] found that sedentary behavior was inversely associated with BMI, especially among those with lower levels of light-intensity physical activity. Furthermore, these results were supported by strong associations observed between sedentary behavior and waist circumference among older adults in Portugal. In particular, for each additional 10- to 20- minutes of sedentary time, the odds of abdominal obesity increased 6.8%. The odds increased up to 48% for each additional 60 minutes of time spent in sedentary behavior [30].

3.9.1. Our findings compared to the 2014 review for obesity/overweight

In our review, we found that all three studies reported positive associations between sedentary time and obesity, measured by BMI or waist circumference. In the prior review [10], five of the six studies reported a positive association between sedentary behavior and obesity. The one exception was a study examining the association between automobile-based sedentary behavior and obesity [10]. Overall, the research is consistent in that eight of nine studies reported a positive relationship between sedentary behavior, overweight, and obesity in older adults.

Table 5. Obesity, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages	Measures	Primary Findings	Limitations	Strengths
Bann, Hire, Manini et al. (2015); Light intensity physical activity and sedentary behavior in relation to body mass index and grip strength in older adults: cross-sectional findings from the lifestyle interventions and independence for elders (LIFE) study. <i>Plos ONE</i>	Examine the association between sedentary behavior, BMI, and grip strength.	Cross-sectional study (N = 1,130) of adults aged 70 – 89 living in the United States.	Accelerometer determined sedentary time Jamar determined grip strength BMI and waist circumference measured by trained staff	In adjusted models, sedentary behavior was associated with BMI. Less time spent sedentary and more time spent in light-intensity physical activities were associated with BMI	Cross sectional design	Use of BMI as a measure of body size
Gennuso, Gangnon, Matthews (2013); Sedentary behavior, physical activity, and markers of health in older adults. <i>Medicine and science in sports and exercise</i>	Examine the association between sedentary behavior, physical activity, and cardiometabolic markers	Population based cross-sectional study (N=1914) of adults aged 65 years and older living in the United States	Accelerometer determined sedentary behavior Cardiometabolic markers (total, HDL, and LDL cholesterol, glucose, triglycerides, glycohemoglobin, waist circumference, blood pressure, and C-reactive protein) Physical function	In adjusted models, higher quartiles of sedentary behavior were associated with weight, BMI, waist circumference, systolic blood pressure, and C-reactive protein among participants who were insufficiently active older adults. Among sufficiently active older adults, higher quartiles of sedentary behavior were associated with waist circumference and LDL cholesterol levels. In addition, insufficiently active older adults who were less sedentary had lower odds of functional limitations.	Cross-sectional design	Accelerometer determined sedentary behavior Population based sample Serum cardiometabolic markers
Judice, Silva, & Sardinha (2015); Sedentary bout durations are associated with abdominal obesity in older adults. <i>The Journal of Nutrition, Health & Aging</i>	Examine the association between sedentary time and patterns of abdominal obesity	Cross-sectional study (N = 351) of older adults (aged 65–103 years) from Portugal	Accelerometer determined sedentary behavior Waist circumference measured by trained staff	No association was observed with total sedentary time and abdominal obesity. However, for bouts of sedentary time that ranged from 10 – 20 minutes up to <60 minutes, each bout significantly increased the odds of abdominal obesity.	Cross sectional design Dietary variables not considered in models	Objectively measured sedentary time

3.10. Mental health (Dementia, mild cognitive impairment, psychological well-being) (Tables 6–8)

We identified eight articles published since 2013 that examined the association between sedentary behavior and mental and cognitive health. Two of the studies focused on dementia [31,32]. In one study, the authors reported that institutionalized versus community-dwelling dementia patients had 23.5% lower daily physical activity levels ($p = 0.004$) and spent 9.3% more time in sedentary behavior ($p = 0.032$). Community-dwelling dementia patients had daily physical activity levels and sedentary time 21.6% lower ($p = 0.007$) and 8.9% longer ($p = 0.078$) than healthy older adults [31]. In another study, sedentary time and sedentary bout duration were significantly greater in dementia patients than in age- and sex-matched cognitively healthy older adults [32].

Mild cognitive impairment (MCI) is defined as cognitive decline greater than expected for age and education level, which does not interfere with independence. Mild cognitive impairment is associated with up to a 30% increased risk of developing dementia within 5 years [33]. In contrast, older adults without mild cognitive impairment develop dementia at a rate of 1% to 2% within 5 years [33]. Therefore, mild cognitive impairment is a critical phase to intervene because it is a transitional stage between healthy cognition and dementia [33]. As observed by Falck et al. 2017, providing effective strategies to maintain cognitive health during this transition period might slow the conversion to dementia [33]. We identified four articles related to mild cognitive impairment. Engeroff et al. 2018 reported that brain-derived neurotrophic factor (BDNF) levels (measure of brain plasticity) were detrimentally associated with sedentary time but beneficially related to accelerometer total activity counts and moderate-to-vigorous intensity physical activity [34]. Falck et al. [33] found that participants with probable mild cognitive impairment had lower physical activity and greater sedentary behavior compared to participants without mild cognitive impairment.

Parkinson's disease is a progressive, neurodegenerative disorder characterized by rigidity, tremor, impaired postural stability, decreased walking ability, and an increased risk of falls [35]. People with mild-to-moderate Parkinson's disease reported 75% of waking hours spent in sedentary behaviors and 18% engaged in physical activity [35].

Subjective cognitive complaints are a meaningful indicator of dementia onset or mild cognitive impairment [36]. Nemoto et al. (2018) found that moderate-to-vigorous intensity physical activity was significantly related with a lower risk of subjective cognitive complaints and that reading showed a dose-response relationship with subjective cognitive complaints [36]. The group who reported > 150 min/week physical activity and > 30 min/day reading showed 60% lower subjective cognitive complaints than the group who reported < 150 min/week physical activity and < 10 min/day reading [36].

Subjective well-being refers to personal evaluations of the degree or extent to which a person considers his/her life meaningful, satisfying, worthwhile, and rewarding [37]. A key to measure a person's wellbeing is the perception of quality of life being experienced [37]. We found two studies related to well-being and sedentary behavior. Okely et al. (2019) reported no association between wellbeing or symptoms of anxiety and sedentary time. On the other hand, symptoms of depression were positively associated with sedentary time [38]. Ku et al. (2015) found that participants who had greater frequencies of leisure-time physical activity and sedentary behaviors, especially engaging in walking, yard/gardening, group exercise, television watching, social chatting, and reading recorded greater levels of well-being. Other types of activities, such as solitary exercise, listening to radio and

playing chess/cards were not related to subjective well-being [37]. Clearly, the type of sedentary behavior makes a difference related to health outcomes.

3.10.1. Our findings compared to the 2014 review for mental health (dementia, mild cognitive impairment, psychological well-being)

In our review, we found eight articles. Dementia was related to sedentary behavior. In three of the four studies, sedentary behavior was related to mild cognitive impairment. In one study, reading was related to lower subjective cognitive complaints. Symptoms of depression were positively associated with sedentary time. In contrast, participants who watched television, chatted socially, and read, recorded greater levels of well-being. Rezende et al. 2014 presented six studies [10]. Individuals who played board games and read were less likely to develop dementia. Similarly, reading books, playing board games, craft activities, computer activities, and watching television were significantly associated with decreased odds of having mild-cognitive impairment (two studies). In contrast, two studies found that sedentary time was negatively associated with psychological well-being and the highest quartile of sitting time was negatively associated with mental health. On the other hand, one study found no association between four or more hours per day of sedentary behavior and successful aging. Considering both reviews, the type of sedentary behavior can be important in understanding the findings. A taxonomy of sedentary behaviors for older adults is needed to elucidate the divergent findings related to sedentary behavior and mental health.

3.11. Cancer (Table 9)

We found one study related to cancer [39]. The purpose of the study was to determine the impact of a mixed exercise program compared to stretching only to improve physical capacity and decrease sedentary behavior time in older adults during cancer treatment. The mixed physical exercise combined aerobic and resistance training. A greater decrease in sedentary behavior time was observed in favor of the mixed exercise program compared to the stretching only program [39].

3.11.1. Our findings compared to the 2014 review for cancer

Both reviews identified only one study related to cancer. In the Rezende et al. 2014 review [10], no association was found between time watching television or videos and renal cell carcinoma. Cancer and sedentary behavior in older adults are underdeveloped areas of research.

Table 6. Dementia, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
van Alphen et al. (2016); Older adults with dementia are sedentary for most of the day. <i>Plos ONE</i>	To objectively assess the physical activity levels of community dwelling and institutionalized ambulatory patients with dementia and to compare with the physical activity levels of cognitive healthy older adults.	The research design was a longitudinal study that examined the effects of regular physical activity on the health of dementia patients; participants were recruited via medical staff of aged care facilities. There were 83 institutionalized [mean age 83.0 (7.6)] and 37 community-dwelling dementia patients [mean age 77.3 (5.6)], and 26 healthy older adults [mean age 79.5 (5.6)]. The study was conducted in The Netherlands.	Actigraphy was used to assess physical activity levels that were based on the raw data. Sedentary behavior was classified < 100 counts/min.	Institutionalized vs. community-dwelling dementia patients had 23.5% lower daily physical activity levels ($p = .004$) and spent 9.3% more time in sedentary behavior ($p = .032$). Community-dwelling dementia patients vs healthy older adults' daily physical activity levels and sedentary time were 21.6% lower ($p = .007$) and 8.9% longer ($p = .078$).	One limitation is that all activity counts per minute represent the same physical activity intensity for healthy as well as adults with dementia. Another limitation is the selection procedure of institutions and daycare centers and homes based on existing collaborations.	The first study to objectively characterize institutionalized and community-dwelling dementia patients' physical activity and sedentary behavior levels.
Hartman et al. (2018); Dementia patients are more sedentary and less physically active than age-and sex-matched cognitively healthy older adults. <i>Dementia Geriatric Cognitive Disorders</i>	To examine physical activity and sedentary behavior characteristics of ambulatory and community-dwelling patients with dementia compared to cognitively healthy age-, sex- and weight-matched controls.	Cross-sectional study - persons with a dementia diagnosis aged > 60 years who were ambulatory and community dwelling were included. ($n = 45$, age 79.6 ± 5.9 years), and matched controls ($n = 49$, age 80.0 ± 7.7 years). Cognitively healthy controls were age, sex, and weight matched to dementia patients and had no history of cognitive impairment. The study was conducted in The Netherlands.	Dementia diagnosis was based on comprehensive clinical assessment by a physician. Mini-Mental State Examination indicated severity of cognitive impairment. Participants wore a wrist accelerometer for 7 days to assess sedentary time, sedentary bout duration and time spent in very light, light-to-moderate and moderate-to-vigorous physical activities.	Relative sedentary time and sedentary bout duration were significantly greater in dementia patients than in controls. In addition, dementia patients spent a lower percentage of their waking time in light-to-moderate and moderate-to-vigorous intensity physical activities. Secondly, older age was associated with a decline in physical activity and increase in sedentary behavior in controls; however, no such age-related changes were found in dementia patients.	The design of the study could not answer the question whether differences in physical activity and sedentary behavior are simply a consequence of dementia. Only community-dwelling patients were included; therefore, results cannot be generalized to institutionalized dementia patients. Furthermore, a potential bias related to the study enrollment should be considered. Dementia patients in the study were enrolled in an exercise trial.	An accelerometer is a validated measure of physical activity and sedentary behavior. A novel finding was that differences between dementia patients and controls remain when corrected for sleep time.

Table 7. Mild cognitive impairment, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Engeroff et al. (2018); Is objectively assessed sedentary behavior, physical activity and cardiorespiratory fitness linked to brain plasticity outcomes in old age? <i>Neuroscience</i>	Examine the association between brain plasticity outcomes and brain-derived neurotrophic factor	Cross-sectional study (N = 50) of older adults (aged 73–77 years old) in Germany	Accelerometer determined sedentary behavior Serum markers of brain metabolism and brain-derived neurotrophic factors (BDNF)	Sedentary time was associated with BDNF, but not brain metabolism measures	Small sample Cognitively healthy sample	Well controlled study Novel methods
Falck et al. (2017); Cross-sectional relationships of physical activity and sedentary behavior with cognitive function in older adults with probable mild cognitive impairment. <i>Physical Therapy</i>	To examine differences in physical activity and sedentary behavior between people with probable mild cognitive impairment and people without mild cognitive impairment and to assess whether associations of physical activity and sedentary behavior with cognitive function differed by mild cognitive impairment status.	The study was cross sectional. The sample size was 150. The mean age was 71.11 years old (7.22). The location was Vancouver, British Columbia, Canada.	Cardiorespiratory fitness Physical activity and sedentary behavior in adults dwelling in the community were measured using a wrist- worn actigraphy unit. The Montreal Cognitive Assessment categorized participants with probable mild cognitive impairment and participants without the condition. Cognitive function was indexed using the Alzheimer Disease Assessment Scale-Cognitive- Plus.	Participants with probable mild cognitive impairment had lower physical activity and higher sedentary behavior than participants without mild cognitive impairment. Higher physical activity and lower sedentary behavior were associated with better Alzheimer Disease Assessment Scale- Cognitive-Plus performance in participants without mild cognitive impairment but not in participants with probable mild cognitive impairment.	This study was cross- sectional and therefore causal associations should not be inferred. In addition, the diagnosis of mild cognitive impairment was not confirmed with a physician.	First study to report that older adults with probable mild cognitive impairment were less active and more sedentary than their cognitively healthy peers. Objective assessments of physical activity and sedentary behavior.

Continued on next page

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Walle'n et al. (2015); Levels and patterns of physical activity and sedentary behavior in elderly people with mild to moderate Parkinson's disease. <i>Physical Therapy</i>	To describe levels and patterns of physical activity and sedentary behaviors in elderly people with mild to moderate Parkinson's Disease.	This cross-sectional study involved a free-living setting and 53 men and 42 women (mean age 73.4 years; (SD-5.7 years) with mild to moderate idiopathic Parkinson's Disease. The country was Sweden.	Time spent in physical activity and sedentary behaviors were assessed for one week with accelerometers.	The main finding was that approximately 75% of all time awake was spent in sedentary behaviors; 18% was spent in physical activity. The proportion of individuals meeting the goal of achieving 150 minutes of moderate- to vigorous-intensity PA per week was 27%; 16% achieved 7,000 or more steps per day; participants spent 589 minutes daily in sedentary behaviors.	The participants were not randomly selected; therefore, generalizability is limited.	Based on objective assessment (accelerometers), described the levels and patterns of physical activity, including diurnal patterns, and estimated the extent to which physical activity recommendations were met.
Nemoto et al. (2018); The association of single and combined factors of sedentary behavior and physical activity with subjective cognitive complaints among community- dwelling older adults: Cross-sectional study. <i>Plos ONE</i>	To examine the association of type of sedentary behavior, physical activity, or their combination with subjective cognitive complaints among community dwelling older adults.	The cross-sectional study targeted independently living individuals aged 65 years and older who resided in Tsuru, Yamanashi Prefecture, Japan (n = 5,328).	Participants responded to the National Functional Survey Questionnaire (Kihon checklist) to assess subjective cognitive complaints. Do your family or friends point out your memory loss? Do you make a call by looking up phone numbers? Do you find yourself not knowing today's date? To evaluate physical activity, the Japanese version of the International Physical Activity Questionnaire Short- version (IPAQ-SV) was conducted. Sedentary behavior time was assessed as subjective average duration of television viewing and reading books or newspapers during the last seven days.	The primary findings were that moderate-to-vigorous intensity physical activity was significantly related with a lower risk of subjective cognitive complaints and that reading showed a dose- response relationship with subjective cognitive complaints. The combined group who reported >150 min/week physical activity and >30 min/ day reading showed 60% lower subjective cognitive complaints than the combined group who reported <150 min/week physical activity and <10 min/day reading.	The cross-sectional design limits causal inferences. Physical activity and sedentary behavior were assessed subjectively. Participants who developed dementia or mild cognitive impairment during the study were not necessarily completely excluded.	First study to report the association of the combined effects of sedentary time and physical activity with subjective cognitive complaints; demographic variables, health behavior, and health status were included in the analyses. The data were from a complete survey with a high response rate of 79.8% and missing values were accounted for by multiple imputation.

Table 8. Psychological well-being, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Okely et al. (2019); Positive and negative well-being and objectively measured sedentary behavior in older adults: evidence from three cohorts. <i>BMC Geriatrics</i>	To examine the association between sedentary behavior and both positive and negative effect	Three prospective (N = 698) of older adults in Scotland	Accelerometer determined sedentary behavior (ActivPAL) Validated scales for positive and negative effect, depression, and anxiety	Sedentary behavior was associated with depression symptoms	Waking time was self-reported	Accelerometer determined sedentary behavior Validated measures of mental health
Ku et al. (2015); Leisure-time physical activity, sedentary behaviors and subjective well-being in older adults: An eight-year longitudinal research. <i>Social Indicators Research</i>	Examine the association between self-reported sedentary time and subjective well-being	Longitudinal fixed cohort (N = 1,268) of adults aged 70 years or older in Taiwan.	Self-reported leisure time in sedentary behavior Subjective well-being (i.e., Life Satisfaction Index)	Television viewing, social chatting, reading, and listening to the radio were significantly associated with subjective well-being	Self-reported sedentary time	Longitudinal design

Table 9. Cancer, sedentary behavior, and older adults (References from 2013–2019).

Reference Author(s), Year, Title, Journal	Hypotheses/Study Aims	Research Design, Sample Size, Ages, Country	Measures	Primary Findings	Limitations	Strengths
Maréchal et al. (2019); Effect of a mixed-exercise program on physical capacity and sedentary behavior in older adults during cancer treatments. <i>Aging Clinical and Experimental Research</i>	To examine the impact of an intervention on increasing physical activity and reducing sedentary behavior during cancer treatment	Two-group (i.e., mixed exercise vs. stretching) pre- and post-test design (N = 14) of older patients in Canada	Self-reported sedentary time Senior fitness test – functional performance test (i.e., chair stands, Up and Go, Arm Curl, Sit and Reach, and 6-minute walk test)	Mixed exercise group had greater improvements in chair stands and global physical capacity scores when compared to those in the stretching group. No other differences were observed	Small sample Sedentary behavior was not the focus	Objective assessments of functional performance

3.12. Study quality assessment of cited literature (Table 10)

The purpose of the quality ratings was to assess risk of bias. As described earlier, we used the National Institute of Health ratings system for each research design [12]. Among 20 observational cohort and cross-sectional studies, three studies were rated as fair and seventeen studies were rated as good. Among the four other research designs [i.e., before-after (pre-post) studies with no control group; case-control studies; controlled intervention studies; and systematic reviews/meta-analyses], there were seven studies. Five were rated as fair, one as good, and one was rated poor—controlled intervention study. Most of the studies adjusted for confounders. Most of the studies were cross-sectional. More randomized controlled trial studies are needed.

Each article in our review was published in a peer-reviewed journal; therefore, only one article was rated as poor. Our quality assessment of articles probably would be different if we had included the grey literature. Given the overall quality assessment of our articles, we are confident in the findings of our review. It is not likely that risk of bias compromises our major conclusions.

Table 10. Quality ratings of cited literature based on National Institutes of Health criteria (n = 27).

Research design	Reference	Quality assessment from National Institute of Health ratings criteria (good, fair, poor)
<i>Observational Cohort and Cross-Sectional Studies</i>		
	Bann et al. 2015	Good
	Benka Wallén et al. 2015	Good
	Cabanas-Sanchez et al. 2018	Fair
	Chang et al. 2015	Fair
	Eklblom et al. 2015	Good
	Engeroff et al. 2018	Fair
	Ensrud et al. 2014	Good
	Figueiró et al. 2019	Good
	Gennuso et al. 2013	Good
	Gennuso et al. 2015	Good
	Jefferis et al. 2019	Good
	Joseph et al. 2016	Good
	Judice et al. 2015	Good
	Klenk et al. 2016	Good
	Ku et al. 2016	Good
	Leon-Munoz et al. 2013	Good
	Matthews et al. 2012	Good
	Nemoto et al. 2018	Good
	Okely et al. 2019	Good
	Pavey et al. 2015	Good
<i>Before-After (Pre-Post) Studies with No Control Group</i>		
	Hartman et al. 2018	Fair
<i>Case-Control Studies</i>		
	Falck et al. 2017	Fair
	van Alphen et al. 2016	Fair
<i>Controlled Intervention Studies</i>		
	Aadahl et al. 2019	Fair
	Maréchal et al. 2019	Poor
<i>Systematic Reviews and Meta-Analyses</i>		
	Thorp et al. 2011	Fair
	Wirth et al. 2017	Good

4. Discussion

By 2030, there will be nearly one billion older adults worldwide [40]. According to Harvey et al. (2013), almost 60% of older adults reported sitting for more than four hours per day; 65% sat in front of a screen for more than three hours per day; and over 55% reported watching television more than two hours. The previous findings were based on self-report; when measured objectively, 67% of older adults were sedentary for > 8.5 hours daily [40,41].

The most consistent associations with increased risk between sedentary behaviors and health outcomes in older adults were: all-cause mortality, metabolic syndrome, triglycerides/high density lipoprotein cholesterol/ blood glucose, HBA1C/glucose intolerance, waist circumference, and obesity/overweight. The inconsistent associations were for blood pressure, cancer, and other cardio-metabolic biomarkers. Mixed findings (positive and negative associations) were found for mental health (dementia, mild cognitive impairment, psychological well-being). A systematic review concluded that limiting sedentary time and participating in regular moderate-to-vigorous intensity physical activity promote healthy cognitive aging. Sedentary behavior was associated with lower cognitive performance, “although the attributable risk of sedentary time to all-cause dementia incidence is unclear” [31].

Our findings compared to an earlier review [10] were similar for all-cause mortality, metabolic syndrome, other cardio-metabolic biomarkers, waist circumference, obesity/overweight, cancer, and mental health (dementia, mild cognitive impairment, psychological well-being). These consistent associations may be attributable to robust results, sufficient literature to draw conclusions, and/or clear outcome measures. Our findings were dissimilar for triglycerides/high density lipoprotein cholesterol/ blood glucose, blood pressure, and HBA1C/glucose intolerance. The dissimilar findings may be accounted for by our recent update (i.e., different times for which studies were conducted), a limited number of studies in the area, or absence of consistent or robust results. Another possibility that merits further study is that heterogeneity in clinical outcomes, methods, and statistical approaches may explain dissimilar results.

Rezende et al. 2014 [10] reported detrimental effects of sedentary behaviors on physical health in older adults. However, it is noteworthy that there are studies documenting positive psychological and mental effects at least for some forms of sedentary behavior [37]. Another review concluded that some sedentary behaviors (e.g., reading or use of computers) were beneficial for older adults [9]. As noted earlier, based on our review, we recommend a taxonomy of sedentary behavior for older adults to disentangle the web of seemingly discrepant findings. Further research is needed to provide greater insights.

Based on our review and the earlier review [10], another emerging area of research for older adults is cancer and sedentary behavior; only two studies were reported. Clearly, cancer, sedentary behavior, and older adults is an understudied research area [39]. Aging is regarded as an important cause of cancer and studies show that aging and cancer together contribute to an increased risk of deconditioning [42,43]. In addition, the combination of aging and cancer is associated with an increase in sedentary behaviors. In the context of cancer treatment such as chemotherapy, patients are more likely to adopt sedentary behaviors and decrease physical activity levels that exacerbate deconditioning which hinders recovery and wellbeing [44–46]. This cycle may be particularly evident in older populations [45–47]. Unfortunately, health care professionals recommend resting and to avoid physical activity during cancer treatments believing this strategy limits cancer-related symptoms especially

cancer-related fatigue [48]. More research is required to understand better how to disrupt this health-compromising cycle and improve the health of older adults during cancer treatments.

From another perspective, the biological pathways that underlie the associations between sedentary behavior and cancer is important [49]. In 10 of 18 articles, a review found that sedentary behavior was associated with increased colorectal, endometrial, ovarian, and prostate cancer risk and cancer mortality in women [49]. The biological pathways or mechanisms that produce the association between sedentary behavior and cancer were adiposity and metabolic dysfunction. The author recommended that reducing sedentary behavior is a viable cancer control strategy [49]. For the health outcomes reported in our systematic review, more research is needed documenting biological pathways.

Limitations—The limitations of this review are a publication bias because we did search for non-published papers and the grey literature. We did not include technical reports, papers from government agencies and other scientific groups, conference proceedings, and unpublished manuscripts. However, we acknowledge that searching the grey literature is important because, typically, only positive, significant findings are published in peer-reviewed journals. Another limitation is that findings were not categorized by physical activity patterns and levels, age groups (e.g., old and very old - over 80 years of age), gender, racial and ethnic identity, or by dose and type of sedentary behavior. All studies should adjust for confounders. Because our systematic review was not a meta-analysis, we did not assess clinical, methodological, and statistical heterogeneity; differences that can influence conclusions. In a narrative review, geriatric-relevant health outcomes included physical function, cognitive function, incontinence, mental health, quality of life and wellbeing as well as sleep. In our review, we did not review research related to incontinence and sleep. Another limitation is the extent to which the literature is dependent on self-report measures. More objective assessments of sedentary behaviors are recommended.

Strengths—In spite of these limitations, our systematic review is the most recent and comprehensive study of this topic including a full description of each article and an evaluation of the quality of each article. Furthermore, we summarize the findings for twelve different health conditions ranging from mortality to mental health and then compared our findings to a previous review. Each study was evaluated for quality and risk of bias based on the National Institute of Health standardized approach for rating the quality of studies [12]. Key characteristics of each study were described and presented in both narrative and table formats.

Future research, policy, and practice – Given the growing population of older adults and the prevalence of sedentary behavior in this population, the effects of sedentary behavior both positive and negative on health outcomes merit careful study. Worldwide, the number of dementia patients has been rapidly increasing during the last few decades, and is expected to reach over 100 million in the year 2050 [36]. Age is the greatest risk factor for dementia [31]. Effective lifestyle interventions are critical because non-pharmacological therapies can be used to treat dementia. The true efficacy of these approaches is not known [31]. Therefore, to prevent dementia or cognitive decline is an urgent, global public health crisis.

To improve the methodological quality of future research, we recommend greater use of diaries and global position systems (GPS) (to provide a context) and accelerometers, which are regarded as a valid and reliable method for evaluating sedentary behavior. In addition, more subgroup analyses including gender, racial and ethnic identity, and geography would improve our understanding of sedentary behavior and health outcomes in older adults. Similarly, dose-response gradients between

sedentary behavior, physical activity, and health outcomes would advance the field. We recommend that future research include meta-analyses.

As noted in previous reviews, chronic diseases and all-cause mortality and the effects of sedentary behavior are important [9,10]. However, from a geriatric perspective, the effects of sedentary behavior on independence, quality of life, and overall impairment (e.g., mobility) are high priority concerns for older adults [9]. More research is needed in these areas.

The next stage of research is to identify modifiable determinants of sedentary behavior among older adults to inform the development of behavior change interventions to reduce health-compromising sedentary behaviors and increase physical activity to improve the health and wellbeing of older adults. An earlier review evaluated the effectiveness of sedentary behavior interventions for older adults and reported non-significant findings [9].

In terms of policy and practice, The United Kingdom advises those 65 years of age and older to minimize the time they spend being sedentary for extended periods [40]. Any proposed policies and practices should be evaluated in randomized controlled trials to evaluate the effects on mental, psychological, and physical health. Policies and practices should discourage health-compromising sedentary behaviors and promote health-enhancing sedentary behaviors (i.e., for mental and psychological health) with the objective to improve the overall wellbeing of older adults.

5. Conclusion

There is a global trend in population aging that embraces terms such as “successful aging”, “active aging,” and “healthy aging” which imply adding life to years and not just living longer (i.e., adding years to life) [37]. To achieve this objective, given the high levels of sedentary behavior among older adults, documenting the health effects of sedentary behavior is important. In our systematic review, we found that sedentary behaviors were adversely associated with six health outcomes ranging from obesity to mortality. Contrary to expectations, depending on the dose and type of sedentary behavior, older adults may benefit from particular kinds of sedentary pursuits that are stimulating and cognitively engaging. Our hope is that this review can be serve as a springboard for more rigorous studies to advance the field of sedentary behavior and health effects on older adults.

Acknowledgments

There were no sources of funding of this study. This systematic review protocol was not published in a repository.

Conflict of interest

All authors declare no conflicts of interest in this paper.

References

1. Organization, WH (2007) *Global age-friendly cities: A guide*: World Health Organization.
2. Matthews CE, Chen KY, Freedson PS, et al. (2008). Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol* 167: 875–881.

3. Hallal PC, Andersen LB, Bull FC, et al. (2012). Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet* 380: 247–257.
4. 2018 Physical Activity Guidelines Advisory Committee. *2018 Physical Activity Guidelines Advisory Committee Scientific Report*. Washington, DC: U.S. Department of Health and Human Services, 2018.
5. Katzmarzyk PT (2010) Physical activity, sedentary behavior, and health: paradigm paralysis or paradigm shift? *Diabetes* 59: 2717–2725.
6. Dunstan DW, Howard B, Healy GN, et al. (2012) Too much sitting—a health hazard. *Diabetes Res Clin Pract* 97: 368–376.
7. Owen N, Healy GN, Matthews CE, et al. (2010) Too much sitting: the population-health science of sedentary behavior. *Exerc Sport Sci Rev* 38: 105–113.
8. Tremblay MS, Colley RC, Saunders TJ, et al. (2010) Physiological and health implications of a sedentary lifestyle. *Appl Physiol Nutr Metab* 35: 725–740.
9. Copeland JL, Ashe MC, Biddle SJH, et al. (2017) Sedentary time in older adults: A critical review of measurement, associations with health, and interventions. *Br J Sports Med* 51: 1539.
10. de Rezende LFM, Rey-López JP, Matsudo VKR, et al. (2014) Sedentary behavior and health outcomes among older adults: a systematic review. *BMC Public Health* 14: 333.
11. Pate RR, O'Neill JR, Lobelo F (2008) The evolving definition of “sedentary”. *Exerc Sport Sci Rev* 36: 173–178.
12. U.S. Department of Health & Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute (2019) *Study quality assessment tools*. Accessed December 8, 2019. Available from: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>
13. Matthews CE, George SM, Moore SC, et al. (2012) Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am J Clin Nutr* 95: 437–445.
14. León-Muñoz LM, Martínez-Gómez D, Balboa-Castillo T, et al. (2013) Continued sedentariness, change in sitting time, and mortality in older adults. *Med Sci Sports Exerc* 45: 1501–1507.
15. Ensrud KE, Blackwell TL, Cauley JA, et al. (2014) Objective measures of activity level and mortality in older men. *J Am Geriatr Soc* 62: 2079–2087.
16. Pavey TG, Peeters GG, Brown WJ (2015) Sitting-time and 9-year all-cause mortality in older women. *Br J Sports Med* 49: 95–99.
17. Jefferis BJ, Parsons TJ, Sartini C, et al. (2019) Objectively measured physical activity, sedentary behaviour and all-cause mortality in older men: does volume of activity matter more than pattern of accumulation? *Br J Sports Med* 53: 1013–1020.
18. Klenk J, Dallmeier D, Denking MD, et al. (2016) Objectively measured walking duration and sedentary behaviour and four-year mortality in older people. *Plos ONE* 11: e0153779.
19. Cabanas-Sánchez V, Guallar-Castillón P, Higuera-Fresnillo S, et al. (2018) Physical activity, sitting time and mortality from inflammatory diseases in older adults. *Front Physiol* 9: 898.
20. Ekblom Ö, Ekblom-Bak E, Rosengren A, et al. (2015) Cardiorespiratory fitness, sedentary behaviour and physical activity are independently associated with the metabolic syndrome, results from the SCAPIS pilot study. *Plos ONE* 10: e0131586.
21. Gennuso KP, Gangnon RE, Thraen-Borowski KM, et al. (2015) Dose–response relationships between sedentary behaviour and the metabolic syndrome and its components. *Diabetologia* 58: 485–492.

22. Figueiró TH, Arins GCB, dos Santos CES, et al. (2019) Association of objectively measured sedentary behavior and physical activity with cardio-metabolic risk markers in older adults. *Plos ONE* 14: e0210861.
23. Aadahl M, Linneberg A, Møller TC, et al. (2014) Motivational counseling to reduce sitting time: a community-based randomized controlled trial in adults. *Am J Prev Med* 47: 576–586.
24. Chang AK, Sok SR (2015) Predictors of sedentary behavior in elderly Koreans with hypertension. *J Nurs Res* 23: 262–270.
25. Joseph JJ, Echouffo-Tcheugui JB, Golden SH, et al. (2016) Physical activity, sedentary behaviors and the incidence of type 2 diabetes mellitus: the Multi-Ethnic Study of Atherosclerosis (MESA). *BMJ Open Diabetes Res Care* 4: e000185.
26. Wirth K, Klenk J, Brefka S, et al. (2017) Biomarkers associated with sedentary behaviour in older adults: a systematic review. *Ageing Res Rev* 35: 87–111.
27. Gennuso KP, Gangnon RE, Matthews CE, et al. (2013) Sedentary behavior, physical activity, and markers of health in older adults. *Med Sci Sports Exerc* 45: 1493–1500.
28. Bann D, Hire D, Manini T, et al. (2015) Light intensity physical activity and sedentary behavior in relation to body mass index and grip strength in older adults: cross-sectional findings from the Lifestyle Interventions and Independence for Elders (LIFE) study. *Plos ONE* 10: e0116058.
29. Thorp AA, Owen N, Neuhaus M, et al. (2011) Sedentary behaviors and subsequent health outcomes in adults: a systematic review of longitudinal studies, 1996–2011. *Am J Prev Med* 41: 207–215.
30. Judice P, Silva A, Sardinha LB (2015) Sedentary bout durations are associated with abdominal obesity in older adults. *J Nutr Health Aging* 19: 798–804.
31. van Alphen HJ, Volkers KM, Blankevoort CG, et al. (2016) Older adults with dementia are sedentary for most of the day. *Plos ONE* 11: e0152457.
32. Hartman YA, Karssemeijer EG, van Diepen LA, et al. (2018) Dementia patients are more sedentary and less physically active than age-and sex-matched cognitively healthy older adults. *Dement Geriatr Cogn Disord* 46: 81–89.
33. Falck RS, Davis JC, Liu-Ambrose T (2017) What is the association between sedentary behaviour and cognitive function? A systematic review. *Br J Sports Med* 51: 800–811.
34. Engeroff T, Füzéki E, Vogt L, et al. (2018) Is objectively assessed sedentary behavior, physical activity and cardiorespiratory fitness linked to brain plasticity outcomes in old age? *Neuroscience* 388: 384–392.
35. Benka Wallén M, Franzén E, Nero H, et al. (2015) Levels and patterns of physical activity and sedentary behavior in elderly people with mild to moderate Parkinson's disease. *Phys Ther* 95: 1135–1141.
36. Nemoto Y, Sato S, Takahashi M, et al. (2018) The association of single and combined factors of sedentary behavior and physical activity with subjective cognitive complaints among community-dwelling older adults: Cross-sectional study. *Plos ONE* 13: e0195384.
37. Ku PW, Fox KR, Chen LJ (2015) Leisure-time physical activity, sedentary behaviors and subjective well-being in older adults: An eight-year longitudinal research. *Soc Indic Res* 127: 1349–1361.
38. Okely JA, Čukić I, Shaw RJ, et al. (2019) Positive and negative well-being and objectively measured sedentary behaviour in older adults: evidence from three cohorts. *BMC Geriatr* 19: 28.

39. Maréchal R, Fontvieille A, Parent-Roberge H, et al. (2019) Effect of a mixed-exercise program on physical capacity and sedentary behavior in older adults during cancer treatments. *Aging Clin Exp Res* 31:1583–1589.
40. Trends G (2003) Public health and aging: trends in aging—United States and worldwide. *Public Health* 347: 921–925.
41. Harvey J, Chastin S, Skelton D (2013) Prevalence of sedentary behavior in older adults: a systematic review. *Int J Env Res Public Health* 10: 6645–6661.
42. de Magalhães JP (2013) How ageing processes influence cancer. *Nat Rev Cancer* 13:357–365.
43. Balducci L, Fossa SD (2013) Rehabilitation of older cancer patients. *Acta Oncol* 52: 233–238.
44. Pinto BM, Eakin E, Maruyama NC (2000) Health behavior changes after a cancer diagnosis: what do we know and where do we go from here? *Ann Behav Med* 22: 38–52.
45. Dewys WD, Begg C, Lavin PT, et al. (1980) Prognostic effect of weight loss prior to chemotherapy in cancer patients. *Am J Med* 69: 491–497.
46. Mohile SG, Velarde C, Hurria A, et al. (2015) Geriatric assessment guided care processes for older adults: A delphi consensus of geriatric oncology experts. *J Natl Compr Canc Netw* 13: 1120–1130.
47. Jones LW, Eves ND, Haykowsky M, et al (2009) Exercise intolerance in cancer and the role of exercise therapy to reverse dysfunction. *Lancet Oncol* 10: 598–605.
48. Jones LW, Courneya KS (2002) Exercise discussions during cancer treatment consultations. *Cancer Pract* 10: 66–74.
49. Lynch BM (2010) Sedentary behavior and cancer: a systematic review of the literature and proposed biological mechanisms. *Cancer Epidemiol Biomarkers Prev* 19: 2691–2709.



AIMS Press

© 2020 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)