
Biliary−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
50000
100000
150000
5000
10000
15000
20000
0.00
0.25
0.50
0.75
1.00


Biliary−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
150000
100000
50000
0
20000
15000
10000
5000
1.00
0.75
0.50
0.25
0.00


Bladder−TCC nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
15000
10000
5000
0
1.00
0.75
0.50
0.25
0.00


Bladder−TCC signatures
Early raw
Late raw
Early normalised
Late normalised
value
15000
10000
5000
0
1.00
0.75
0.50
0.25
0.00


Bone−Benign nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
200
400
600
250
500
750
0.00
0.25
0.50
0.75
1.00


Bone−Benign signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
200
400
600
250
500
750
0.00
0.25
0.50
0.75
1.00


Bone−Epith nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
4000
500
1500
0.00
0.25
0.50
0.75
1.00


Bone−Epith signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
4000
500
1500
0.00
0.25
0.50
0.75
1.00


Bone−Osteosarc nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
9000
6000
3000
0
3000
2000
1000
1.00
0.75
0.50
0.25
0.00


Bone−Osteosarc signatures
Early raw
Late raw
Early normalised
Late normalised
value
9000
6000
3000
0
3000
2000
1000
1.00
0.75
0.50
0.25
0.00


Breast−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
25000
20000
15000
10000
5000
0
40000
30000
1.00
0.75
0.50
0.25
0.00


Breast−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
25000
20000
15000
10000
5000
0
40000
30000
1.00
0.75
0.50
0.25
0.00


Breast−DCIS nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
1500
1000
500
0
600
400
200
1.00
0.75
0.50
0.25
0.00


Breast−DCIS signatures
Early raw
Late raw
Early normalised
Late normalised
value
1500
1000
500
0
600
400
200
1.00
0.75
0.50
0.25
0.00


Breast−LobularCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
40000
30000
20000
10000
0
9000
6000
3000
1.00
0.75
0.50
0.25
0.00


Breast−LobularCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
40000
30000
20000
10000
0
9000
6000
3000
1.00
0.75
0.50
0.25
0.00


Cervix−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
4000
250
500
750
0.00
0.25
0.50
0.75
1.00


Cervix−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
4000
3000
2000
1000
0
750
500
250
1.00
0.75
0.50
0.25
0.00


Cervix−SCC nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
8000
6000
4000
2000
0
1.00
0.75
0.50
0.25
0.00


Cervix−SCC signatures
Early raw
Late raw
Early normalised
Late normalised
value
8000
6000
4000
2000
0
1.00
0.75
0.50
0.25
0.00


CNS−GBM nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
7500
5000
2500
0
150000
100000
50000
1.00
0.75
0.50
0.25
0.00


CNS−GBM signatures
Early raw
Late raw
Early normalised
Late normalised
value
7500
5000
2500
0
150000
100000
50000
1.00
0.75
0.50
0.25
0.00


CNS−Medullo nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
3000
2000
1000
0
1500
500
1.00
0.75
0.50
0.25
0.00


CNS−Medullo signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
500
1500
0.00
0.25
0.50
0.75
1.00


CNS−Oligo nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
5000
4000
3000
2000
1000
0
1500
500
1.00
0.75
0.50
0.25
0.00


CNS−Oligo signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
4000
5000
500
1500
0.00
0.25
0.50
0.75
1.00

CNS−PiloAstro nucleotidesubstitution1




CNS−PiloAstro signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
200
400
600
800
0.00
0.25
0.50
0.75
1.00


ColoRect−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
6e+05
4e+05
2e+05
0e+00
2000000
1500000
1000000
500000
0
1.00
0.75
0.50
0.25
0.00


ColoRect−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
6e+05
4e+05
2e+05
0e+00
2000000
1500000
1000000
500000
0
1.00
0.75
0.50
0.25
0.00


Eso−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
60000
40000
20000
0
30000
10000
1.00
0.75
0.50
0.25
0.00


Eso−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
60000
40000
20000
0
30000
10000
1.00
0.75
0.50
0.25
0.00


Head−SCC nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Head−SCC signatures
Early raw
Late raw
Early normalised
Late normalised
value
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Kidney−ChRCC nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
6000
4000
2000
0
3000
1000
1.00
0.75
0.50
0.25
0.00


Kidney−ChRCC signatures
Early raw
Late raw
Early normalised
Late normalised
value
6000
4000
2000
0
3000
1000
1.00
0.75
0.50
0.25
0.00


Kidney−RCC.clearcell nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
5000
10000
15000
20000
0.00
0.25
0.50
0.75
1.00


Kidney−RCC.clearcell signatures
Early raw
Late raw
Early normalised
Late normalised
value
20000
15000
10000
5000
0
1.00
0.75
0.50
0.25
0.00

Kidney−RCC.papillary nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
10000
7500
5000
2500
0
4000
3000
2000
1000
1.00
0.75
0.50
0.25
0.00


Kidney−RCC.papillary signatures
Early raw
Late raw
Early normalised
Late normalised
value
10000
7500
5000
2500
0
4000
3000
2000
1000
1.00
0.75
0.50
0.25
0.00


Liver−HCC nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
50000
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Liver−HCC signatures
Early raw
Late raw
Early normalised
Late normalised
value
50000
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Lung−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Lung−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Lung−SCC nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
50000
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Lung−SCC signatures
Early raw
Late raw
Early normalised
Late normalised
value
50000
40000
30000
20000
10000
0
15000
5000
1.00
0.75
0.50
0.25
0.00


Lymph−BNHL nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
10000
20000
30000
40000
5000
15000
0.00
0.25
0.50
0.75
1.00


Lymph−BNHL signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
10000
20000
30000
40000
5000
15000
0.00
0.25
0.50
0.75
1.00


Lymph−CLL nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
4000
0.00
0.25
0.50
0.75
1.00


Lymph−CLL signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
1000
2000
3000
4000
0.00
0.25
0.50
0.75
1.00


Myeloid−AML nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
500
1000
1500
2000
300
600
900
1200
0.00
0.25
0.50
0.75
1.00


Myeloid−AML signatures
Early raw
Late raw
Early normalised
Late normalised
value
2500
2000
1500
1000
500
0
1200
900
600
300
1.00
0.75
0.50
0.25
0.00


Myeloid−MPN nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
500
1000
1500
0.00
0.25
0.50
0.75
1.00


Myeloid−MPN signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
500
1000
1500
0.00
0.25
0.50
0.75
1.00


Ovary−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
30000
20000
10000
0
9000
6000
3000
1.00
0.75
0.50
0.25
0.00


Ovary−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
30000
20000
10000
0
9000
6000
3000
1.00
0.75
0.50
0.25
0.00


Panc−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
10000
20000
30000
0.00
0.25
0.50
0.75
1.00


Panc−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
10000
20000
30000
0.00
0.25
0.50
0.75
1.00


Panc−Endocrine nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
8000
6000
4000
2000
0
15000
10000
5000
1.00
0.75
0.50
0.25
0.00


Panc−Endocrine signatures
Early raw
Late raw
Early normalised
Late normalised
value
8000
6000
4000
2000
0
15000
10000
5000
1.00
0.75
0.50
0.25
0.00


Prost−AdenoCA nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
10000
5000
0
6000
4000
2000
1.00
0.75
0.50
0.25
0.00


Prost−AdenoCA signatures
Early raw
Late raw
Early normalised
Late normalised
value
10000
5000
0
6000
4000
2000
1.00
0.75
0.50
0.25
0.00


Skin−Melanoma.acral nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
0
5000
10000
15000
2500
7500
0.00
0.25
0.50
0.75
1.00


Skin−Melanoma.acral signatures
Early raw
Late raw
Early normalised
Late normalised
value
0
5000
10000
15000
2500
7500
0.00
0.25
0.50
0.75
1.00


Skin−Melanoma.cutaneous nucleotidesubstitution1
Early raw
Late raw
Early normalised
Late normalised
value
8e+05
6e+05
4e+05
2e+05
0e+00
80000
60000
40000
20000
0
1.00
0.75
0.50
0.25
0.00


Skin−Melanoma.cutaneous signatures
Early raw
Late raw
Early normalised
Late normalised
value
8e+05
6e+05
4e+05
2e+05
0e+00
80000
60000
40000
20000
0
1.00
0.75
0.50
0.25
0.00