Log ratio changes: Biliary−AdenoCA nucleotidesubstitution1

Log ratio changes: Biliary−AdenoCA signatures

Log ratio changes: Bladder−TCC nucleotidesubstitution1

Log ratio changes: Bladder−TCC signatures

Log ratio changes: Bone−Benign nucleotidesubstitution1

Log ratio changes: Bone−Benign signatures

Log ratio changes: Bone−Epith nucleotidesubstitution1

Log ratio changes: Bone−Epith signatures

Log ratio changes: Bone−Osteosarc nucleotidesubstitution1

Log ratio changes: Bone−Osteosarc signatures

Log ratio changes: Breast−AdenoCA nucleotidesubstitution1

Log ratio changes: Breast−AdenoCA signatures

Log ratio changes: Breast−DCIS nucleotidesubstitution1

Log ratio changes: Breast–DCIS signatures

Log ratio changes: Breast−LobularCA nucleotidesubstitution1

Log ratio changes: Breast−LobularCA signatures

Log ratio changes: Cervix−AdenoCA nucleotidesubstitution1

Log ratio changes: Cervix–AdenoCA signatures

Log ratio changes: Cervix−SCC nucleotidesubstitution1

Log ratio changes: Cervix−SCC signatures

Log ratio changes: CNS−GBM nucleotidesubstitution1

Log ratio changes: CNS−GBM signatures

Log ratio changes: CNS−Medullo nucleotidesubstitution1

Log ratio changes: CNS−Medullo signatures

Log ratio changes: CNS−Oligo nucleotidesubstitution1

Log ratio changes: CNS−Oligo signatures

Log ratio changes: CNS−PiloAstro nucleotidesubstitution1

Log ratio changes: CNS−PiloAstro signatures

Log ratio changes: ColoRect–AdenoCA nucleotidesubstitution1

Log ratio changes: ColoRect−AdenoCA signatures

Log ratio changes: Eso−AdenoCA nucleotidesubstitution1

Log ratio changes: Eso−AdenoCA signatures

Log ratio changes: Head−SCC nucleotidesubstitution1

Log ratio changes: Head−SCC signatures

Log ratio changes: Kidney−ChRCC nucleotidesubstitution1

Log ratio changes: Kidney−ChRCC signatures

Log ratio changes: Kidney−RCC.clearcell nucleotidesubstitution1

Log ratio changes: Kidney−RCC.clearcell signatures

Log ratio changes: Kidney−RCC.papillary nucleotidesubstitution1

Log ratio changes: Kidney−RCC.papillary signatures

Log ratio changes: Liver−HCC nucleotidesubstitution1

Log ratio changes: Liver−HCC signatures

Log ratio changes: Lung−AdenoCA nucleotidesubstitution1

Log ratio changes: Lung−AdenoCA signatures

Log ratio changes: Lung−SCC nucleotidesubstitution1

Log ratio changes: Lung−SCC signatures

Log ratio changes: Lymph−BNHL nucleotidesubstitution1

Log ratio changes: Lymph−BNHL signatures

Log ratio changes: Lymph−CLL nucleotidesubstitution1

Log ratio changes: Lymph−CLL signatures

Log ratio changes: Myeloid−AML nucleotidesubstitution1

Log ratio changes: Myeloid−AML signatures

Log ratio changes: Myeloid−MPN nucleotidesubstitution1

Log ratio changes: Myeloid−MPN signatures

Log ratio changes: Ovary−AdenoCA nucleotidesubstitution1

Log ratio changes: Ovary−AdenoCA signatures

Log ratio changes: Panc−AdenoCA nucleotidesubstitution1

Log ratio changes: Panc−AdenoCA signatures

Log ratio changes: Panc−Endocrine nucleotidesubstitution1

Log ratio changes: Panc−Endocrine signatures

Log ratio changes: Prost−AdenoCA nucleotidesubstitution1

Log ratio changes: Prost−AdenoCA signatures

Log ratio changes: Skin−Melanoma.acral nucleotidesubstitution1

Log ratio changes: Skin−Melanoma.acral signatures

Log ratio changes: Skin−Melanoma.cutaneous nucleotidesubstitution1

Log ratio changes: Skin−Melanoma.cutaneous signatures

Log ratio changes: SoftTissue−Leiomyo nucleotidesubstitution1

Log ratio changes: SoftTissue−Leiomyo signatures

Log ratio changes: SoftTissue−Liposarc nucleotidesubstitution1

Log ratio changes: SoftTissue−Liposarc signatures

Log ratio changes: Stomach−AdenoCA nucleotidesubstitution1

Log ratio changes: Stomach−AdenoCA signatures

Log ratio changes: Thy−AdenoCA nucleotidesubstitution1

Log ratio changes: Thy−AdenoCA signatures

Log ratio changes: Uterus−AdenoCA nucleotidesubstitution1

Log ratio changes: Uterus−AdenoCA signatures