



The Win Ratio and Beyond

Evolution, Applications, and Open Questions in Composite Endpoint Analysis

LU MAO

Imao@biostat.wisc.edu

Department of Biostatistics & Medical Informatics

University of Wisconsin-Madison

Outline

- **Background**
- **Censoring's impact on estimand**
 - Time frame of comparison
- **Two approaches to clarity**
 - Nonparametric: fix the time frame
 - Semiparametric: posit a time-constant feature
- **Summary and discussion**

Introduction

Composite Endpoint

- **Traditional composite endpoint**

- **Time to first event**

- Major adverse cardiac event (MACE): death, heart failure, myocardio-infarction, stroke, etc.

- **Limitations:**

- Lack of clinical priority
 - Statistical inefficiency (waste of data)

- **Hierarchical composite endpoint (HCE)**

- **Example:** Death > nonfatal MACE > six-minute walk test (6MWT)/ NYHA class

Win Ratio: Basics

- **A common HCE**


- **Proposed and popularized** by Pocock et al. ([2012](#))
- **Treatment vs control:** generalized pairwise comparisons (GPC) ([Buyse et al., 2025](#))
 - **Win-loss:** Longer survival → fewer/later nonfatal MACE → better 6MWT/NYHA
- **Effect size:** $WR = \text{wins} / \text{losses}$
 - Fold increase in probability of winning over losing

- **Alternative metrics**

- **Net benefit** (Proportion in favor): $NB = \text{wins} - \text{losses}$ ([Buyse, 2010](#))
- **Win odds:** $WO = (\text{wins} + 2^{-1}\text{ties}) / (\text{losses} + 2^{-1}\text{ties})$ ([Brunner et al., 2021](#))

Prioritized Comparison

Cut off at earlier end of follow-up




Pairwise Comparison

- **GPC Results**

- Wins: $8 / 16 = 50\%$
- Losses: $3 / 16 = 18.8\%$
- Ties: $5 / 16 = 31.2\%$


- **Effect Sizes**

- $WR = 50\% / 18.8\% = 2.67$
- $NB = 50\% - 18.8\% = 31.2\%$
- $WO = (50\% + 15.6\%) / (18.8\% + 15.6\%) = 1.72$



Stratification

- GPC within stratum and aggregate ([Dong et al., 2018, 2023](#))



$$\text{Wins: } 50\% \times 8 / (8 + 6) + 33\% \times 6 / (8 + 6) = 42.7\%$$

$$\text{Losses: } 38\% \times 8 / (8 + 6) + 22\% \times 6 / (8 + 6) = 31.1\%$$

- Overall WR: $42.7\% / 31.1\% = 1.37$

Properties and Use Cases

General Setting

- Notation

- n_1, n_0 : sample sizes in treatment and control
- W_{ij} : win-loss indicator of subject i in treatment vs j in control

$$W_{ij} = \begin{cases} 1, & \text{if subject } i \text{ in treatment wins} \\ -1, & \text{if subject } j \text{ in control wins} \\ 0, & \text{if tie} \end{cases}$$

- Wins : $\hat{w}_{1,0} = (n_1 n_0)^{-1} \sum_{i=1}^{n_1} \sum_{j=1}^{n_0} I(W_{ij} = 1)$
- Losses : $\hat{w}_{0,1} = (n_1 n_0)^{-1} \sum_{i=1}^{n_1} \sum_{j=1}^{n_0} I(W_{ij} = -1)$
- $WR = \hat{w}_{1,0} / \hat{w}_{0,1}$; $NB = \hat{w}_{1,0} - \hat{w}_{0,1}$; $WO = (1 + NB) / (1 - NB)$

Binary Case Equivalencies

- **Binary outcome, no censoring**

- $Y_i^{(a)} = 1$ (response) vs 0 (no-response) for subject i in group a ($a = 0, 1$)
- Win-loss (1-0) fractions

$$\hat{w}^{(a,1-a)} = (n_a n_{1-a})^{-1} \sum_{i=1}^{n_a} \sum_{j=1}^{n_{1-a}} Y_i^{(a)} (1 - Y_j^{(1-a)}) = \hat{p}_a (1 - \hat{p}_{1-a})$$

- $\hat{p}_a = n_a^{-1} \sum_{i=1}^{n_a} Y_i^{(a)} = \text{response rate in group } a$

- **GPC stats**

- $WR = [\hat{p}_1(1 - \hat{p}_0)] / [\hat{p}_0(1 - \hat{p}_1)] = \text{response odds ratio}$
- $NB = \hat{p}_1 - \hat{p}_0 = \text{response rate difference}$

Variance Estimation

- **Bootstrap**

- Initial proposal ([Pocock et al., 2012](#))

- **Analytic asymptotic variance**

- Asymptotic decomposition of U -statistics ([Bebu & Lachin, 2016](#); [Dong et al., 2016](#); [Luo et al., 2015](#))
- Influence function-based
- **R packages:** [WR](#), [WINS](#), [BuyseTest](#), [WinRatio](#), etc.

Null vs Alternative Hypotheses


- **Null hypothesis:** No treatment effect on any event
 - $WR = 1, NB = 0, WO = 1$ regardless of follow-up time
- **Alternative hypothesis**
 - Death and hospitalization times: *stochastically larger* in treatment than in control ([Luo et al., 2015](#); [Mao, 2019](#))
 - Stratified test: Stochastical order in each stratum
 - So $WR > 1, NB > 0, WO > 1$ regardless of follow-up time
 - Power $\rightarrow 1$ as $n = n_1 + n_0 \rightarrow \infty$

Recurrent Events (I)

- **GPC to accommodate recurrent events**
 - E.g., hospitalizations, exacerbations, infections, etc.
- **Comparison rules**
 - Death → Number of events → Time to first/last event ([Mao et al., 2022](#)) → ... (Biomarker/QoL changes)
 - H_A : Treatment stochastically delays all events

Recurrent Events (II)


◇ Censoring ■ Hospitalization ● Death



ATTRibute-CM (NEJM 2024)

- Trial data


- **Design:** Phase 3, double-blind RCT (n = 632; 2:1 randomization)
- **Population:** Transthyretin amyloid cardiomyopathy (90 % wild-type, mean age 77 yr)
- **Intervention:** Acoramidis 800 mg BID vs placebo for 30 months
- **Primary HCE:** Death → CV hospitalization → NT-proBNP → 6-min walk distance
 - Stratified by TTR genotype, NT-proBNP level, eGFR, etc.



DEFENDER (JAMA 2024)

- Trial data

- **Design:** Multicenter, open-label RCT (n = 507; 22 ICUs in Brazil)
- **Population:** Critically ill adults with an organ dysfunction (respiratory/ cardiovascular/ renal)
- **Intervention:** Dapagliflozin 10 mg daily vs standard care alone (up to 14 days)
- **Primary HCE:** Hospital mortality → initiation of KRT → ICU length of stay (28 days)



WR as Primary Endpoint (I)

Gregson et al. (2025) summarizes recent/ongoing RCTs:

Trial	Condition	Interventions	HCE
ATTRibute-CM (2024)	Transthyretin amyloid cardiomyopathy	Acoramidis vs placebo	Death → CV hosp → NT-proBNP/6MWT
COVID-PACT (2022)	Critically ill COVID-19	Full-dose AC vs standard-dose AC	Venous/arterial death → PE → DVT → ...
DAPA-MI (2024)	Clinically stable acute MI	Dapagliflozin vs placebo	CV death → non-CV death → HF hosp → ...
DEFENDER (2024)	Critical illness with organ failure	Dapagliflozin + SOC vs SOC	In-hospital death → KRT initiation → ICU LOS
EMPULSE (2022)	Acute HF (hospitalized)	Empagliflozin vs placebo	All-cause death → HF events (number/time) → KCCQ change
HEART-FID (2023)	HFrEF with iron deficiency	Ferric carboxymaltose vs placebo	All-cause death → HF hosp → 6MWT change

WR as Primary Endpoint (II)

Trial	Condition	Interventions	HCE
PARTNER 3 (2023)	Low-risk symptomatic AS	TAVR vs SAVR	Death → stroke → rehospitalization days
REDUCE LAP-HF II (2022)	HFpEF/HFmrEF	Atrial shunt device vs sham	CV death/stroke → HF hosp → KCCQ change
TRILUMINATE (2023)	Symptomatic tricuspid regurgitation	TEER vs OMT	Death/surgery → HF hosp → KCCQ change
VIP-ACS (2022)	Acute coronary syndrome	High-dose influenza vaccine vs standard	Death → MI → stroke → ...

...


Extensions and Variations

Sample Size Calculation

Estimand Issue

- Without censoring

- WR, NB, WO = functions of joint distribution of events
- Change with follow-up time ([Mao, 2024](#); [Oakes, 2016](#))



- With censoring

- Weighted average of win-loss probabilities over time
- Trial-dependent estimand ([ICH, 2020](#))

Time Restriction

- **Specify time horizon τ**
 - Imagine all subjects followed up to τ (1 yr, 3 yrs, etc.)
- **GPC under consistent time frame**
 - Estimand: $WR(\tau)$, $NB(\tau)$, $WO(\tau)$
 - Interpretation: win ratio/net benefit/win odds up to time τ
- **Estimation with censored data**
 - Inverse probability censoring weighting ([Dong et al., 2020b, 2021](#))
 - R-package: [WINS](#) ([Cui & Huang, 2023](#))
 - Multiple imputations ([T. Wang et al., 2023, 2024](#))

Regression Analysis

Win-Loss Times

DOOR

Challenges and Open Questions

Critiques

Covariate Adjustment

- **Different from regression**

- **Marginal estimands** for $\mathcal{H}^{*(1)}$ vs $\mathcal{H}^{*(0)}$, not conditioning on Z
- **Gain efficiency** when outcome-covariate model is true, otherwise still valid (robustness)
- **Standard endpoints**
 - Continuous, binary, univariate survival, etc. (Tsiatis et al., 2008; B. Wang et al., 2021; Ye et al., 2023)
 - FDA recommendation (FDA, 2023)

- **Challenges with WR**

- U -statistic structure
- Lack of likelihood structure

Interim Analysis

- **Purpose**

- Analyze interim data for evidence of efficacy/futility → stop trial early
- Univariate survival: information accrued \propto number of events

- **Challenges with WR**

- Information time not event-driven
 - Case study: RMST ([Lu & Tian, 2021](#); [Luo et al., 2019](#))
- Correlations between component events
 - *Dependent* increments

Meta Analysis

- Challenges

- Primary studies *not* reporting win-loss measures
- Primary studies over *different* time spans
- Primary studies with *different* definitions of win/loss

- WinKM: A toolkit to start

- Calculate win-loss statistics based on
 - KM estimates for OS and EFS
 - At-risk table at selected time points
 - Total event counts (reported in the CONSORT diagram or results section)

Conclusion

Summary

Acknowledgments

- Funding

- [R01HL149875](#) (11/2019 – 7/2028)
- [DMS2015526](#) (7/2020 – 6/2024)



National Heart, Lung,
and Blood Institute



- Collaborators

- [KyungMann Kim](#), [Tuo Wang](#), Gaohong Dong, Bo Huang, etc.

References

- Bebu, I., & Lachin, J. M. (2016). Large sample inference for a win ratio analysis of a composite outcome based on prioritized components. *Biostatistics*, 17(1), 178–187. <https://doi.org/10.1093/biostatistics/kxv032>
- Brunner, E., Vandemeulebroecke, M., & Mütze, T. (2021). Win odds: An adaptation of the win ratio to include ties. *Statistics in Medicine*, 40(14), 3367–3384. <https://doi.org/10.1002/sim.8967>
- Buyse, M. E. (2010). Generalized pairwise comparisons of prioritized outcomes in the two-sample problem. *Statistics in Medicine*, 29(30), 3245–3257. <https://doi.org/10.1002/sim.3923>
- Buyse, M. E., Verbeeck, J., De Backer, M., Deltuvaite-Thomas, V., Saad, E. D., & Molenberghs, G. (2025). *Handbook of generalized pairwise comparisons: Methods for patient-centric analysis*. CRC Press, Taylor & Francis Group.
- Cui, Y., & Huang, B. (2023). *WINS: The r WINS package*. <https://CRAN.R-project.org/package=WINS>
- Dong, G., Hoaglin, D. C., Huang, B., Cui, Y., Wang, D., Cheng, Y., & Gamalo-Siebers, M. (2023). The stratified win statistics (win ratio, win odds, and net benefit). *Pharmaceutical Statistics*, 22(4), 748–756. <https://doi.org/10.1002/pst.2293>

- Dong, G., Huang, B., Wang, D., Verbeeck, J., Wang, J., & Hoaglin, D. C. (2021). Adjusting win statistics for dependent censoring. *Pharmaceutical Statistics*, 20(3), 440–450. <https://doi.org/10.1002/pst.2086>
- Dong, G., Li, D., Ballerstedt, S., & Vandemeulebroecke, M. (2016). A generalized analytic solution to the win ratio to analyze a composite endpoint considering the clinical importance order among components. *Pharmaceutical Statistics*, 15(5), 430–437. <https://doi.org/10.1002/pst.1763>
- Dong, G., Mao, L., Huang, B., Gamalo-Siebers, M., Wang, J., Yu, G., & Hoaglin, D. C. (2020b). The inverse-probability-of-censoring weighting (IPCW) adjusted win ratio statistic: an unbiased estimator in the presence of independent censoring. *Journal of Biopharmaceutical Statistics*, 30(5), 882–899. <https://doi.org/10.1080/10543406.2020.1757692>
- Dong, G., Qiu, J., Wang, D., & Vandemeulebroecke, M. (2018). The stratified win ratio. *Journal of Biopharmaceutical Statistics*, 28(4), 778–796. <https://doi.org/10.1080/10543406.2017.1397007>
- FDA. (2023). *Guidance document: Adjusting for covariates in randomized clinical trials for drugs and biological products*. US Food and Drug Administration. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/adjusting-covariates-randomized-clinical-trials-drugs-and-biological-products>
- Gregson, J., Taylor, D., Owen, R., Collier, T., J. Cohen, D., & Pocock, S. (2025). Hierarchical composite outcomes and win ratio methods in cardiovascular

- trials: A review and consequent guidance. *Circulation*, 151(22), 1606–1619.
- ICH. (2020). *ICH E9 (R1) addendum on estimands and sensitivity analysis in clinical trials to the guideline on statistical principles for clinical trials, step 5*. London: European Medicines Evaluation Agency.
- Lu, Y., & Tian, L. (2021). Statistical Considerations for Sequential Analysis of the Restricted Mean Survival Time for Randomized Clinical Trials. *Statistics in Biopharmaceutical Research*, 13(2), 210–218. <https://doi.org/10.1080/19466315.2020.1816491>
- Luo, X., Huang, B., & Quan, H. (2019). Design and monitoring of survival trials based on restricted mean survival times. *Clinical Trials*, 16(6), 616–625. <https://doi.org/10.1177/1740774519871447>
- Luo, X., Tian, H., Mohanty, S., & Tsai, W. Y. (2015). An Alternative Approach to Confidence Interval Estimation for the Win Ratio Statistic. *Biometrics*, 71(1), 139–145. <https://doi.org/10.1111/biom.12225>
- Mao, L. (2019). On the Alternative Hypotheses for the Win Ratio. *Biometrics*, 75(1), 347–351. <https://doi.org/10.1111/biom.12954>
- Mao, L. (2024). Defining estimand for the win ratio: Separate the true effect from censoring. *Clinical Trials*, 21(5), 584–594.
- Mao, L., Kim, K., & Li, Y. (2022). On recurrent-event win ratio. *Statistical Methods in Medical Research*, 31(6), 1120–1134. <https://doi.org/10.1177/09622802221084134>

- Oakes, D. (2016). On the win-ratio statistic in clinical trials with multiple types of event. *Biometrika*, 103(3), 742–745. <https://doi.org/10.1093/biomet/asw026>
- Pocock, S. J., Ariti, C. A., Collier, T. J., & Wang, D. (2012). The win ratio: a new approach to the analysis of composite endpoints in clinical trials based on clinical priorities. *European Heart Journal*, 33(2), 176–182. <https://doi.org/10.1093/eurheartj/ehr352>
- Tsiatis, A. A., Davidian, M., Zhang, M., & Lu, X. (2008). Covariate adjustment for two-sample treatment comparisons in randomized clinical trials: A principled yet flexible approach. *Statistics in Medicine*, 27(23), 4658–4677. <https://doi.org/10.1002/sim.3113>
- Wang, B., Susukida, R., Mojtabai, R., Amin-Esmaeili, M., & Rosenblum, M. (2021). Model-Robust Inference for Clinical Trials that Improve Precision by Stratified Randomization and Covariate Adjustment. *Journal of the American Statistical Association*, 118(542), 1152–1163. <https://doi.org/10.1080/01621459.2021.1981338>
- Wang, T., Li, Y., & Qu, Y. (2024). Restricted time win ratio: from estimands to estimation. *Statistics in Biopharmaceutical Research*, 1–18. <https://doi.org/10.1080/19466315.2024.2332675>
- Wang, T., Zilinskas, R., Li, Y., & Qu, Y. (2023). Missing Data Imputation for a Multivariate Outcome of Mixed Variable Types. *Statistics in Biopharmaceutical Research*, 15(4), 826–837. <https://doi.org/10.1080/19466315.2023.2169753>