

Convergence rate of $X^N - X$ for McKean equations

Luis Mario Chaparro Jaquez

April 24, 2022

Todo list

type this result which is lemma 4.2 in the paper	2
this norm is wrong, should be $1 + \alpha$ on the lhs and everywhere else	4
Check this norm	6
continue from page three in notes	8
add the explanation of why to drop the diffusion term	9

Contents

1 What is this?	1
2 Some useful definitions and results	1
3 Bounds for the difference of solutions to the Kolmogorov equations	3
4 Bound for the difference of the auxiliary functions	8
5 Bound for the local time at zero of the solution to the SDEs	8
6 Convergence rate of the solution to the regularised SDE and the original	11

1 What is this?

An adaptation of [1, Proposition 3.1] for the case of ~~McKean SDEs~~ SDEs with drift in a Besov space of negative order proposed in [2] and [3].

The proof builds on a number of results presented in the sections below.

2 Some useful definitions and results

Here we present some results and definitions to refer on the text.

Definition 1: Local time at zero

For any real-valued continuous semi-martingale, the local time at zero $L_t^0(\bar{Y})$ is defined as

$$L_t^0(\bar{Y}) = \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_0^t \mathbb{1}_{\{|\bar{Y}| \leq \epsilon\}} d\langle \bar{Y} \rangle_s, \mathbb{P}\text{-a.s.} \quad (1)$$

For all $t \geq 0$.

The first result, [1, Lemma 5.1], is not necessary to prove for this particular setting since the result holds for any semi-martingale, I include it here for self-containment reasons.

Lemma 1: Bound for local time at zero for a semi-martingale

For any $\epsilon \in (0, 1)$ and any real-valued, continuous semi-martingale Z we have

$$\begin{aligned} \mathbb{E}[L_t^0(Z_s)] &\leq 4\epsilon - \mathbb{E} \left[\int_0^t \left(\mathbb{1}_{\{Z_s \in (0, \epsilon)\}} + \mathbb{1}_{\{Z_s \geq \epsilon\}} e^{1-Z_s/\epsilon} \right) dZ_s \right] \\ &\quad + \frac{1}{\epsilon} \mathbb{E} \left[\int_0^t \mathbb{1}_{\{Z > \epsilon\}} e^{1-Z_s/\epsilon} d\langle Z \rangle_s \right]. \end{aligned}$$

Let us introduce the original and regularised Kolmogorov equations.

Definition 2: Kolmogorov equations

For $\beta \in (0, 1/2)$ let $b \in C_T \mathcal{C}^{-\beta}$, $u, u^N \in C_T \mathcal{C}^{(1+\beta)+}$, and $b^N \rightarrow b$ as $N \rightarrow \infty$ in $C_T \mathcal{C}^{-\beta}$. The equations

$$\begin{cases} \partial u_i + \frac{1}{2} b_i \Delta u_i = \lambda u_i - b_i \\ u_i(T) = 0, \end{cases} \quad (2)$$

$$\begin{cases} \partial u_i^N + \frac{1}{2} b_i^N \Delta u_i^N = \lambda u_i^N - b_i^N \\ u_i^N(T) = 0. \end{cases} \quad (3)$$

are called Kolmogorov and regularised Kolmogorov equations. Here written component wise.

Lemma 2: Bounds for $\nabla u, \nabla u^N$

Let

3 Bounds for the difference of solutions to the Kolmogorov equations

We need a bound for $u - u^N$ and $\nabla u - \nabla u^N$ in L_∞ for the case in which $u \in C_T \mathcal{C}^{1+\alpha}$ for some $\alpha > \beta$ which is an adaptation of [1, Lemma 5.2].

The result builds on top of the following result:

Proposition 1: Bound for the ρ -equivalent norm of $u - u^N$

Let u, u^N be (mild) solutions to the Kolmogorov equations from ?? 2 then as $N \rightarrow \infty$

$$\|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \leq \frac{cT^{\frac{1-\beta-\alpha}{2}} \|b_i - b_i^N\|_{C_T \mathcal{C}^{-\beta}} (\|u_i\|_{C_T \mathcal{C}^{1+\alpha}} - 1)}{1 - c\rho^{\frac{\alpha+\beta-1}{2}} (\|b\|_{C_T \mathcal{C}^{-\beta}} + \lambda)} \quad (4)$$

for $\rho \geq \rho_0$, where

$$\rho_0 = 2c(\|b_i\|_{C_T \infty + \alpha} + \lambda)^{\frac{2}{\alpha+\beta+1}} \quad (5)$$

and $\lambda > 0$.

Proof. See that $u^N(T) = u(T) = 0$, and in [2], set g^N, g as b^N, b respectively. See that $b^N \rightarrow b$. Then let us reformulate the rest of the aforementioned result for $\lambda \neq 0$.

As u^N, u are mild solutions, we have

$$\begin{aligned} u_i(t) - u_i^N(t) &= P_{T-t}(u_i(T) - u_i^N(T)) \\ &\quad + \int_t^T P_{s-t}(\nabla u_i b_i - \nabla u_i^N b_i^N) ds \\ &\quad - \int_t^T P_{s-t}(\lambda u_i + b_i - \lambda u_i^N + b_i^N) ds \\ &= \int_t^T P_{s-t}(\nabla u_i b_i - \nabla u_i^N b_i^N) ds \\ &\quad - \lambda \int_t^T P_{s-t}(u_i - u_i^N) ds \\ &\quad - \int_t^T P_{s-t}(b_i - b_i^N) ds \\ &= \int_t^T P_{s-t}[(\nabla u_i b_i - \nabla u_i b_i^N) + (\nabla u_i b_i^N - \nabla u_i^N b_i^N)] ds \\ &\quad - \lambda \int_t^T P_{s-t}(u_i - u_i^N) ds \\ &\quad - \int_t^T P_{s-t}(b_i - b_i^N) ds \end{aligned}$$

$$\begin{aligned}
&= \int_t^T P_{s-t}(\nabla u_i b_i - \nabla u_i b_i^N) ds \\
&+ \int_t^T P_{s-t}(\nabla u_i b_i^N - \nabla u_i^N b_i^N) ds \\
&- \lambda \int_t^T P_{s-t}(u_i - u_i^N) ds \\
&- \int_t^T P_{s-t}(b_i - b_i^N) ds
\end{aligned}$$

Now let us compute the ρ -equivalent norm of $u - u^N$, for some $\alpha > \beta$

$$\begin{aligned}
\|u_i - u_i^N\|_{C_T C^{-\beta}}^{(\rho)} &= \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \|u(t) - u^N(t)\|_{1+\alpha} \\
&\leq \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \int_t^T P_{s-t}(\nabla u_i b_i - \nabla u_i b_i^N) ds \right\|_{1+\alpha} \\
&+ \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \int_t^T P_{s-t}(\nabla u_i b_i^N - \nabla u_i^N b_i^N) ds \right\|_{1+\alpha} \\
&- \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \lambda \int_t^T P_{s-t}(u_i - u_i^N) ds \right\|_{1+\alpha} \\
&- \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \int_t^T P_{s-t}(b_i - b_i^N) ds \right\|_{1+\alpha}.
\end{aligned}$$

Let us take each term from the right hand side of the inequality and bound them.

For the first term, using $\gamma + 2\theta = 1 + \alpha$, $\gamma = -\beta$, $\theta = \frac{1+\alpha+\beta}{2}$, $\|P_t f\|_{\gamma+2\theta} \leq ct^{-\theta} \|f\|_{\gamma}$ and $\|\nabla g\|_{\xi} \leq c\|g\|_{\xi+1}$

$$\begin{aligned}
&\sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \int_t^T P_{s-t}(\nabla u_i b_i - \nabla u_i b_i^N) ds \right\|_{1+\alpha} \\
&\leq \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \int_t^T (s-t)^{-\theta} \|\nabla u_i\|_{\alpha} \|b_i - b_i^N\|_{-\beta} \\
&\leq c \|u_i\|_{C_T C_{1+\alpha}} \|b_i - b_i^N\|_{C_T C^{-\beta}} \sup_{0 \leq t \leq T} e^{-\rho(T-t)} (T-t)^{\frac{1-\beta-\alpha}{2}} \\
&\leq c T^{\frac{1-\beta-\alpha}{2}} \|u_i\|_{C_T C_{1+\alpha}} \|b_i - b_i^N\|_{C_T C^{-\beta}}
\end{aligned}$$

For the second term, see that for $N \rightarrow \infty$, we have $\|b^N\|_{C_T C^{-\beta}} \leq 2\|b\|_{C_T C^{-\beta}}$

$$\begin{aligned}
&\sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \int_t^T P_{s-t} b_i^N (\nabla u_i - \nabla u_i^N) ds \right\|_{1+\alpha} \\
&\leq c \sup_{0 \leq t \leq T} \int_t^T (s-t)^{-\theta} e^{-\rho(T-t)} 2 \|b_i\|_{-\beta} \|\nabla u_i - \nabla u_i^N\|_{\alpha} ds
\end{aligned}$$

type this
result which
is lemma 4.2
in the paper

$$\begin{aligned}
&\leq c \|b_i\|_{C_T \mathcal{C}^{-\beta}} \|u_i - u_i^N\|_{C_T \mathcal{C}^{-\beta}}^{(\rho)} \int_t^T (s-t)^{-\theta} e^{-\rho(T-t)} ds \\
&\leq c \|b_i\|_{C_T \mathcal{C}^{-\beta}} \|u_i - u_i^N\|_{C_T \mathcal{C}^{-\beta}}^{(\rho)} \rho^{\frac{\alpha+\beta-1}{2}}
\end{aligned}$$

For the third term, which is the one that differs from the proof in [2] we need to use that $\|P_t f\|_{\gamma+2\theta} \leq ct^{-\theta} \|f\|_{\gamma}$, and in this case we have $\gamma + 2\theta = 1 + \alpha$ and $\gamma = 1 + \alpha$, so that $\theta = 0$ because $u, u^N \in C_T \mathcal{C}^{1+\alpha}$, so we will have

$$\begin{aligned}
&\sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \lambda \int_t^T P_{s-t}(u_i - u_i^N) ds \right\|_{1+\alpha} \\
&\leq c \lambda \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \int_t^T (s-t)^{-0} \|u_i - u_i^N\|_{1+\alpha} ds \\
&= c \lambda \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \int_t^T e^{-\rho(T-s)} \sup_{0 \leq s \leq T} e^{-\rho(T-s)} \|u_i - u_i^N\|_{1+\alpha} ds \\
&= c \lambda \sup_{0 \leq t \leq T} \|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \int_t^T e^{-\rho(T-s)} e^{-\rho(T-t)} ds \\
&= c \lambda \sup_{0 \leq t \leq T} \|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \int_t^T e^{-\rho(s-t)} ds \\
&= c \lambda \sup_{0 \leq t \leq T} \|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \sup_{0 \leq t \leq T} \rho^{-1} [1 - e^{-\rho(T-t)}] \\
&\leq c \lambda \sup_{0 \leq t \leq T} \|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \rho^{-1} \\
&\leq c \lambda \sup_{0 \leq t \leq T} \|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \rho^{\frac{\alpha+\beta-1}{2}}
\end{aligned}$$

And for the last term

$$\begin{aligned}
&\sup_{0 \leq t \leq T} e^{-\rho(T-t)} \left\| \int_t^T P_{T-s}(b_i - b_i^N) ds \right\|_{1+\alpha} \\
&\leq c \sup_{0 \leq t \leq T} e^{-\rho(T-t)} \int_t^T (s-t)^{-\frac{\alpha+\beta-1}{2}} \|b_i - b_i^N\|_{-\beta} ds \\
&\leq c \|b_i - b_i^N\|_{C_T \mathcal{C}^{-\beta}} \sup_{0 \leq t \leq T} e^{-\rho(T-t)} (s-t)^{-\frac{\alpha+\beta-1}{2}} \\
&\leq c T^{\frac{1-\beta-\alpha}{2}} \|b_i - b_i^N\|_{C_T \mathcal{C}^{-\beta}}
\end{aligned}$$

Putting everything together

$$\begin{aligned}
\|u_i - u_i^N\|_{C_T \mathcal{C}^{-\beta}}^{(\rho)} &\leq c T^{\frac{1-\beta-\alpha}{2}} \|u_i\|_{C_T \mathcal{C}^{1+\alpha}} \|b_i - b_i^N\|_{C_T \mathcal{C}^{-\beta}} \\
&\quad + c \|b_i\|_{C_T \mathcal{C}^{-\beta}} \|u_i - u_i^N\|_{C_T \mathcal{C}^{-\beta}}^{(\rho)} \rho^{\frac{\alpha+\beta-1}{2}} \\
&\quad - c \lambda \sup_{0 \leq t \leq T} \|u_i - u_i^N\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)} \rho^{\frac{\alpha+\beta-1}{2}}
\end{aligned}$$

$$-cT^{\frac{1-\beta-\alpha}{2}} \|b_i - b_i^N\|_{C^T \mathcal{C}^{-\beta}},$$

and finally,

$$\begin{aligned} \|u_i - u_i^N\|_{C^T \mathcal{C}^{-\beta}}^{(\rho)} (1 - c\rho^{\frac{\alpha+\beta-1}{2}} [\|b\|_{C^T \mathcal{C}^{-\beta}} + \lambda]) \\ \leq cT^{\frac{1-\beta-\alpha}{2}} \|b_i - b_i^N\|_{C^T \mathcal{C}^{-\beta}} (\|u_i\|_{C^T \mathcal{C}^{1+\alpha}} - 1) \\ \|u_i - u_i^N\|_{C^T \mathcal{C}^{-\beta}}^{(\rho)} \leq \frac{cT^{\frac{1-\beta-\alpha}{2}} \|b_i - b_i^N\|_{C^T \mathcal{C}^{-\beta}} (\|u_i\|_{C^T \mathcal{C}^{1+\alpha}} - 1)}{(1 - c\rho^{\frac{\alpha+\beta-1}{2}} [\|b\|_{C^T \mathcal{C}^{-\beta}} + \lambda])} \end{aligned}$$

As required. \square

Note that in the above we can represent the right hand side of the inequality as

$$\|u_i - u_i^N\|_{C^T \mathcal{C}^{-\beta}}^{(\rho)} \leq \frac{cT^{\frac{1-\beta-\alpha}{2}} (\|u_i\|_{C^T \mathcal{C}^{1+\alpha}} - 1)}{(1 - c\rho^{\frac{\alpha+\beta-1}{2}} [\|b\|_{C^T \mathcal{C}^{-\beta}} + \lambda])} \|b_i - b_i^N\|_{C^T \mathcal{C}^{-\beta}} \quad (6)$$

$$\|u_i - u_i^N\|_{C^T \mathcal{C}^{-\beta}}^{(\rho)} \leq c(\rho) \|b_i - b_i^N\|_{C^T \mathcal{C}^{-\beta}} \quad (7)$$

Here is the adaptation of [1, Lemma 5.2].

Proposition 2: Bounds for $\|u - u^N\|_{L^\infty}$ and $\|\nabla u - \nabla u^N\|_{L^\infty}$

Let $\beta \in (0, 1/2)$ and $b \in C^T \mathcal{C}^{-\beta}$. Let $u, u^N \in C^T \mathcal{C}^{(1+\beta)+}$ be (mild) solutions to the Kolmogorov equations from ?? 2.

Assume, by ??, that for some $\alpha > \beta$

$$\|u - u^N\|_{C^T \mathcal{C}^{1+\alpha}}^{(\rho)} \leq c(\rho) \|b - b^N\|_{C^T \mathcal{C}^{-\beta}}. \quad (8)$$

With $c(\rho)$ as in ?? and ρ_0 is large enough such that $c(\rho) > 0$ for all $\rho > \rho_0$.

Then for all $t \in [0, T]$

$$\|u^N(t) - u(t)\|_{L^\infty} \leq \kappa_\rho \|b - b^N\|_{C^T \mathcal{C}^{-\beta}} \quad (9)$$

$$\|\nabla u^N(t) - \nabla u(t)\|_{L^\infty} \leq \kappa_\rho \|b - b^N\|_{C^T \mathcal{C}^{-\beta}} \quad (10)$$

with $\kappa_\rho = c \cdot c(\rho) \cdot e^{\rho T}$.

this norm is wrong, should be $1 + \alpha$ on the lhs and everywhere else

Proof. First let us prove Eq. (9).

Let $t \in [0, T]$, and see that since $u, u^N \in C^T \mathcal{C}^{(1+\beta)+}$ there exists $\alpha > \beta$ such that $u, u^N \in C^T \mathcal{C}^{1+\alpha}$, then for any $f \in \mathcal{C}^{1+\alpha}$ we have

$$\|f\|_{C^{1+\alpha}} \leq c \left(\sup_{x \in \mathbb{R}^d} |f(x)| + \sup_{x \neq y \in \mathbb{R}^d} \frac{|\nabla f(x) - \nabla f(y)|}{|x - y|^\alpha} \right) \quad (11)$$

so we have

$$\begin{aligned}\|u^N(t) - u(t)\|_{L^\infty} &= \sup_{x \in \mathbb{R}^d} |u^N(t, x) - u(t, x)| \\ &\leq c \|u^N(t) - u(t)\|_{\mathcal{C}^{\alpha+1}}\end{aligned}\tag{12}$$

Moreover, using the (ρ) -equivalent norm

$$\|f\|_{\mathcal{C}^{1+\alpha}} = \sup_{t \in [0, T]} e^{-\rho(T-t)} \|f(t)\|_{\mathcal{C}^{1+\alpha}},\tag{13}$$

and Eq. (8) we see that

$$\begin{aligned}\|u^N - u\|_{C_T \mathcal{C}^{1+\alpha}} &= \sup_{t \in [0, T]} \|u^N - u\|_{\mathcal{C}^{1+\alpha}} \\ &= \sup_{t \in [0, T]} e^{\rho(T-t)} e^{-\rho(T-t)} \|u^N - u\|_{\mathcal{C}^{1+\alpha}} \\ &\leq e^{\rho T} \sup_{t \in [0, T]} e^{-\rho(T-t)} \|u^N - u\|_{\mathcal{C}^{1+\alpha}} \\ &= e^{\rho T} \|u^N - u\|_{C_T \mathcal{C}^{1+\alpha}}^{(\rho)}.\end{aligned}\tag{14}$$

Plugging Eq. (14) into Eq. (12)

$$\begin{aligned}\|u^N(t) - u(t)\|_{L^\infty} &\leq c \|u^N(t) - u(t)\|_{\mathcal{C}^{\alpha+1}} \\ &\leq \sup_{t \in [0, T]} c \|u^N(t) - u(t)\|_{\mathcal{C}^{\alpha+1}} \\ &= c \|u^N - u\|_{C_T \mathcal{C}^{\alpha+1}} \\ &\leq c e^{\rho T} \|u^N - u\|_{C_T \mathcal{C}^{\alpha+1}}^{(\rho)}.\end{aligned}\tag{15}$$

And finally by Eq. (8)

$$\|u^N(t) - u(t)\|_{L^\infty} \leq c \cdot c(\rho) \cdot e^{\rho T} \|b^N - b\|_{C_T \mathcal{C}^{-\beta}}\tag{16}$$

which proves Eq. (9).

For Eq. (10) recall that if $f \in \mathcal{C}^{1+\alpha}$ then $\nabla f \in \mathcal{C}^\alpha$. Also, by Bernstein inequality [3, Eqn. (9)]

$$\|\nabla f\|_\alpha \leq c \|f\|_{\infty+\alpha}.\tag{17}$$

Using the equivalent norm

$$\|f\|_{\mathcal{C}^{1+\alpha}} \leq c \left(\sup_{x \in \mathbb{R}^d} |f(x)| + \sup_{x \in \mathbb{R}^d} |\nabla f(x)| + \sup_{x \neq y \in \mathbb{R}^d} \frac{|\nabla f(x) - \nabla f(y)|}{|x - y|^\alpha} \right)\tag{18}$$

we can see that

$$\|\nabla u^N(t) - \nabla u(t)\|_{L^\infty} \leq c \|u^N(t) - u(t)\|_{\mathcal{C}^{1+\alpha}}.\tag{19}$$

And usign the same bounds that we used above for $c \|u^N(t) - u(t)\|_{\mathcal{C}^{1+\alpha}}$ this point follows. \square

4 Bound for the difference of the auxiliary functions

This is the adaptation of result [1, Lemma 5.3].

Proposition 3: Bound for $|\psi(t, x) - \psi^N(t, x)|$

Take $\rho > \rho_0$ as in ??, $N \rightarrow \infty$, κ_ρ from ?? 2, and $\beta \in (0, 1/2)$, then we have

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}} |\psi(t, x) - \psi^N(t, x)| \leq 2\kappa_\rho \|b - b^N\|_{C_T C^{-\beta}} \quad (20)$$

Proof. Recall the definition of $\psi, \phi \in C_T \mathcal{C}^1$

$$\phi(t, x) := x + u(t, x) \quad (21)$$

$$\psi(t, \cdot) = \phi^{-1}(t, \cdot). \quad (22)$$

Note that

$$u(y) = \int_0^1 \nabla u(\alpha y) y d\alpha + u(0). \quad (23)$$

From there we have

$$u(t, y) - u(t, y') = \int_0^1 \nabla u(t, \alpha(y - y'))(y - y') d\alpha \quad (24)$$

and therefore

$$|u(t, y) - u(t, y')| \geq \left(\int_0^1 |\nabla u(t, \alpha(y - y'))|^2 d\alpha \right)^{1/2} |y - y'|, \quad (25)$$

and by ?? we finally have

$$\begin{aligned} |u(t, y) - u(t, y')| &\leq \left(\frac{1}{4} \int_0^1 d\alpha \right)^{1/2} |y - y'| \\ |u(t, y) - u(t, y')|^2 &\leq \frac{1}{4} |y - y'|^2 \end{aligned} \quad (26)$$

□

Check this norm

5 Bound for the local time at zero of the solution to the SDEs

We need a bound for $\mathbb{E}[L_T^0(Y^N - Y)]$, for Sobolev spaces, this is result [1, Proposition 5.4] we present it here for the solutions to the SDE belonging to the appropriate Besov spaces.

Proposition 4: Bound for local time of solutions to the SDE

Let $b \in C_T \mathcal{C}^{-\beta}$ and $b^N \rightarrow b$ in $C_T \mathcal{C}^{-\beta}$ as $N \rightarrow \infty$ for $\beta \in (0, \frac{1}{4})$ and for any $\alpha > \beta$

Proof. Recall that Y^N, Y are solutions to the SDEs

$$Y_t = y_0 + \lambda \int_0^t u(s, \psi(s, Y_s)) ds + \int_0^t (\nabla u(s, \psi(s, Y_t)) + 1) dW_s \quad (27)$$

and

$$Y_t^N = y_0^N + \lambda \int_0^t u^N(s, \psi^N(s, Y_s^N)) ds + \int_0^t (\nabla u^N(s, \psi^N(s, Y_t^N)) + 1) dW_s \quad (28)$$

so that the difference $Y^N - Y$ is

$$\begin{aligned} Y^N - Y_t &= (y_0^N + \lambda \int_0^t u^N(s, \psi^N(s, Y_s^N)) ds + \int_0^t (\nabla u^N(s, \psi^N(s, Y_t^N)) + 1) dW_s) \\ &\quad - (y_0 + \lambda \int_0^t u(s, \psi(s, Y_s)) ds + \int_0^t (\nabla u(s, \psi(s, Y_t)) + 1) dW_s) \\ &= (y_0^N - y_0) + \lambda \int_0^t (u^N(s, \psi^N(s, Y_s^N)) - u(s, \psi(s, Y_s))) ds \\ &\quad - \int_0^t (\nabla u^N(s, \psi^N(s, Y_t^N)) - \nabla u(s, \psi(s, Y_t))) dW_s, \end{aligned} \quad (29)$$

and using ?? 1 we have the following bound

$$\begin{aligned} &\mathbb{E}[L_t^0(Y^N - Y)] \\ &\leq 4\epsilon + 1 \\ &- 2(1 + \lambda) \mathbb{E} \left[\int_0^t \left(\mathbb{1}_{\{Y_s^N - Y_s \in (0, \epsilon)\}} + \mathbb{1}_{\{Y_s^N - Y_s \geq \epsilon\}} e^{1 - \frac{Y_s^N - Y_s}{\epsilon}} \right) (u^N(s, \psi^N(s, Y_s^N)) - u(s, \psi(s, Y_s))) ds \right] \\ &+ \frac{1}{\epsilon} \mathbb{E} \left[\int_0^t \mathbb{1}_{\{Y_s^N - Y_s > \epsilon\}} e^{1 - \frac{Y_s^N - Y_s}{\epsilon}} (\nabla u^N(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s)))^2 ds \right]. \end{aligned} \quad (30)$$

For the second and third terms of Eq. (30) let us bound the factors involving the differences of u, u^N and $\nabla u, \nabla u^N$.

First, for u, u^N adding and subtracting terms and using triangle inequality we have

$$\begin{aligned} |u^N(s, \psi^N(s, Y_s^N)) - u(s, \psi(s, Y_s))| &\leq |u^N(s, \psi^N(s, Y_s^N)) - u(s, \psi^N(s, Y_s^N))| \\ &\quad + |u(s, \psi^N(s, Y_s^N)) - u(s, \psi(s, Y_s^N))| \\ &\quad + |u(s, \psi(s, Y_s^N)) - u(s, \psi(s, Y_s))|. \end{aligned} \quad (31)$$

□

continue
from page
three in
notes

The terms in the right hand side will be bounded as follows:

- For the first term, by ?? 2

$$|u^N(s, \psi^N(s, Y_s^N)) - u(s, \psi^N(s, Y_s^N))| \leq \|u^N(s) - u(s)\|_{L^\infty} \leq \kappa_\rho \|b - b^N\|_{C_T \mathcal{C}^{-\beta}}, \quad (32)$$

- for the second term, observe that u, u^N are $\frac{1}{2}$ -Lipschitz and by Section 4 we get

$$\begin{aligned} |u(s, \psi^N(s, Y_s^N)) - u(s, \psi(s, Y_s^N))| &\leq \frac{1}{2} |\psi^N(s, Y_s^N) - \psi(s, Y_s^N)| \\ &\leq \kappa_\rho \|b^N - b\|_{C_T \mathcal{C}^{-\beta}}, \end{aligned} \quad (33)$$

- and for the final term, note that ψ, ψ^N are 2-Lipschitz so that

$$|u(s, \psi(s, Y_s^N)) - u(s, \psi(s, Y_s))| \leq \frac{1}{2} |\psi(s, Y_s^N) - \psi(s, Y_s)| \leq |Y_s^N - Y_s|. \quad (34)$$

So that the following bound holds

$$|u^N(s, \psi^N(s, Y_s^N)) - u(s, \psi(s, Y_s))| \leq 2\kappa_\rho \|b^N - b\|_{C_T \mathcal{C}^{-\beta}} + |Y_s^N - Y_s|. \quad (35)$$

Now for the third term in Eq. (30) by adding and subtracting terms and using the triangle inequality

$$\begin{aligned} |\nabla u^N(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s))| &\leq |\nabla u^N(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi^N(s, Y_s^N))| \\ &\quad + |\nabla u(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s^N))| \\ &\quad + |\nabla u(s, \psi(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s))|. \end{aligned} \quad (36)$$

The terms on the right hand side will be bounded as follows:

- For the first term we use ?? 2 and we have

$$\begin{aligned} |\nabla u^N(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi^N(s, Y_s^N))| &\leq \|\nabla u^N(s) - \nabla u(s)\|_{L^\infty} \\ &\leq \kappa_\rho \|b - b^N\|_{C_T \mathcal{C}^{-\beta}}, \end{aligned} \quad (37)$$

for the second term see that $\nabla u, \nabla u^N$ are α -Hölder continuous and using Section 4 we have

$$\begin{aligned} |\nabla u(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s^N))| &\leq |\psi^N(s, Y_s^N) - \psi(s, Y_s^N)|^\alpha \|u\|_{C_T \mathcal{C}^{1+\alpha}} \\ &\leq (2\kappa_\rho \|b^N - b\|_{C_T \mathcal{C}^{-\beta}})^\alpha \|u\|_{C_T \mathcal{C}^{1+\alpha}}. \end{aligned} \quad (38)$$

Therefore we get the bound

$$\begin{aligned}
|\nabla u^N(s, \psi^N(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s))| &\leq \kappa_\rho \|b - b^N\|_{C_T \mathcal{C}^{-\beta}} \\
&\quad + 2^\alpha \kappa_\rho^\alpha \|b^N - b\|_{C_T \mathcal{C}^{-\beta}}^\alpha \|u\|_{C_T \mathcal{C}^{1+\alpha}} \quad . \quad (39) \\
&\quad + |\nabla u(s, \psi(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s))|
\end{aligned}$$

Using the bounds for Eq. (35) and Eq. (39) and the inequality

$$(x_1 + \dots + x_k)^2 \leq k(x_1 + \dots + x_k), \quad (40)$$

for some k , we get

$$\begin{aligned}
&\mathbb{E}[L_t^0(Y^N - Y)] \\
&\leq 4\epsilon + 4(1 + \lambda) \left(2\kappa_\rho \|b^N - b\|_{C_T \mathcal{C}^{-\beta}} t + \mathbb{E} \left[\int_0^t |Y_s^N - Y^N| ds \right] \right) \\
&\quad + \frac{1}{\epsilon} 3t \|b^N - b\|_{C_T \mathcal{C}^{-\beta}} \left(\kappa_\rho^2 \|b^N - b\|_{C_T \mathcal{C}^{-\beta}} + 4(2\kappa_\rho)^{2\alpha} \|u\|_{C_T \mathcal{C}^{1+\alpha}}^2 \|b^N - b\|_{C_T \mathcal{C}^{-\beta}}^{2\alpha-1} \right) \quad (41) \\
&\quad + \frac{1}{\epsilon} 3\mathbb{E} \left(\int_0^t \mathbb{1}_{\{Y_s^N - Y_s > \epsilon\}} e^{1 - \frac{Y_s^N - Y_s}{\epsilon}} |\nabla u(s, \psi(s, Y_s^N)) - \nabla u(s, \psi(s, Y_s))|^2 ds \right)
\end{aligned}$$

6 Convergence rate of the solution to the regularised SDE and the original

References

- [1] Tiziano De Angelis, Maximilien Germain, and Elena Issoglio. “A Numerical Scheme for Stochastic Differential Equations with Distributional Drift”. In: *arXiv:1906.11026 [cs, math]* (Oct. 22, 2020). arXiv: [1906.11026](#).
- [2] Elena Issoglio and Francesco Russo. “A PDE with Drift of Negative Besov Index and Related Martingale Problem”. In: (), p. 48.
- [3] Elena Issoglio and Francesco Russo. “McKean SDEs with singular coefficients”. In: *arXiv:2107.14453 [math]* (July 30, 2021). arXiv: [2107.14453](#).