

Towards a 3D likelihood analysis in VHE γ -ray astronomy: the case of H.E.S.S.

L. Mohrmann^{1*}, K. Nakashima¹, A. Specovius¹, D. Tiziani¹, S. Funk¹, C. van Eldik¹, S. Ohm^{2†}

¹ Erlangen Centre for Astroparticle Physics, Erwin-Rommel-Str. 1, 91058 Erlangen, Germany

² DESY, Platanenallee 6, 15738 Zeuthen, Germany


* Contact: lars.mohrmann@fau.de † Presenter



Construction of a model for the residual cosmic-ray background

- Description of cosmic-ray background is key element in Imaging Atmospheric Cherenkov Telescope (IACT) data analysis
→ accurate background model is pre-requisite for 3D likelihood analysis
- Construct model based on archival H.E.S.S. data taken off the Galactic plane (≈ 3240 hours of data)
- Central idea: Average rate in field-of-view coordinate system

Construction




Application of a 3D likelihood analysis to public H.E.S.S. data

- Fit spectro-morphological (3D) model templates to observed data
→ extract source morphology & spectrum simultaneously
→ obtain complete description of observed field of view
- Apply analysis to public H.E.S.S. test data set¹
→ use open-source tools *ctools*² and *Gammapy*³
- Results agree well with standard methods



RX J1713.7-3946



Validation

- Fit model normalisation and “spectral tilt” for all 4-telescope observations
→ can be well recovered
- Perform χ^2 test along slices of R.A. and Dec.
→ distribution almost flat (expectation for perfect model)

