

# GMACS Exposure Time Calculator, v2.0

Luke Schmidt<sup>1,\*</sup> and Nathaniel Peirson<sup>1</sup>

<sup>1</sup>Texas A&M University, Department of Physics & Astronomy, College Station, TX, 77843, USA

\*lschmidt@physics.tamu.edu

## ABSTRACT

This document collects the assumptions, models, and references used in the GMACS exposure time calculator. Version 2 is based on the original GMACS ETC created by Ting Li.

## Assumptions

1. Effective area of the telescope is  $368\text{ m}^2$  for full size (7 mirrors) and  $222\text{ m}^2$  for first light size (4 mirrors).
2. CCD readnoise is  $2e^- \text{ px}^{-1}$ ,  $15\mu\text{m}$  pixels.
3. GMACS consists of a blue channel (320-600 nm), and a red channel (500-1000 nm). Dichroic transition is at 558 nm.
4. All the sources, either stars or galaxies, are treated as point sources. (i.e. the angular extension in the sky only depends on the seeing.)
5. Noise is a combination of sky background and CCD read noise.
6. PSF of the object is Gaussian and the seeing is the FWHM of the PSF.
7. The extraction aperture is equal to the seeing, and the extraction is assumed to be perfect at the center.
8. SNR is calculated for every pixel. For the default  $0.7''$  slit, a resolution element is 12px (equivalent to  $3.7\text{\AA}$  for low resolution and  $1.4\text{\AA}$  for high resolution), binning options include  $1 \times 1$ ,  $2 \times 2$ ,  $3 \times 3$ , and  $4 \times 4$  pixels.

## Source Templates

1. **Star** templates are from Pickles 1998<sup>1</sup>.
2. **Extended Source** templates are from Kinney et al. 1996<sup>2</sup>. Flux below 1300 Angstrom is zero in rest frame. So the Flux for high redshift ( $z > 4$ ) at short wavelength will be also zero and thus is not correct. (For example, flux is zero in u band for an object at  $z=5$ ; in this case, SNR is set to be zero at all wavelengths)
3. **Sky backgrounds** are from Steven Villanueva et al. 2012<sup>3</sup>. You can select the sky background for different moon phases.
4. **User-defined magnitudes** are computed with SDSS filters for ugriz (<http://www.sdss3.org/instruments/camera.php#Filters>), with Johnson/Bessell filters for UBVRI from Bessell et al. (1990)<sup>4</sup>.
5. **User Submitted** sources may be sent to lschmidt@physics.tamu.edu. They should be formatted as a text file with two columns, wavelength [ $\text{\AA}$ ] and flux [ $\text{erg cm}^{-2} \text{s}^{-1} \text{\AA}^{-1}$ ] a sample file can be downloaded at [http://instrumentation.tamu.edu/gmacs/etc\\_gmacs/core/kinney/ellipticals](http://instrumentation.tamu.edu/gmacs/etc_gmacs/core/kinney/ellipticals) (this sample has a third column, the standard deviation of the flux, which is not necessary to include).

## Throughput

1. **Telescope** The primary and secondary mirrors of the GMT are assumed to be coated with Aluminum. Reflectivity values taken from in situ measurements of the Subaru 8.3 primary mirror<sup>5</sup>, <https://subarutelescope.org/Observing/Telescope/Parameters/Reflectivity/>
2. **Optics** throughput for the collimator and camera lenses are 0.60 (Blue) and 0.62 (Red).

3. **Dichroic** throughput is based on the SDSS-III BOSS dichroics that were coated by JDSU.
4. **Grating** throughput is based on low resolution VPH gratings designed by KAISER.
5. **Detectors** are assumed to be the e2v Astro Multi-2 (NIMO DD) CCD for the red channel and e2v Astro BB (NIMO std Si) CCD for the blue channel.
6. **Atmospheric extinction** is created by libRadTran<sup>6</sup> with the atmospheric parameters measured by aTmCam<sup>7</sup> at CTIO at airmass=1.0.

## Acknowledgements

The following software was used to develop the GMACS exposure time calculator. Python<sup>8</sup>, Spectres<sup>9</sup>, Astropy<sup>10,11</sup>, Bokeh<sup>12</sup>, Numpy<sup>13</sup>, and Scipy<sup>14</sup>.

## References

1. Pickles, A. J. A Stellar Spectral Flux Library: 1150-25000 Å. *Publ. Astron. Soc. Pac.* **110**, 863–878, DOI: [10.1086/316197](https://doi.org/10.1086/316197) (1998).
2. Kinney, A. L. *et al.* Template Ultraviolet to Near-Infrared Spectra of Star-forming Galaxies and Their Application to K-Corrections. *The Astrophys. J.* **467**, 38, DOI: [10.1086/177583](https://doi.org/10.1086/177583) (1996).
3. Villanueva, S., DePoy, D. L. & Marshall, J. L. Optimal resolutions for optical and NIR spectroscopy. In *Ground-based and Airborne Instrumentation for Astronomy IV*, vol. 8446 of *Proc. SPIE*, 84462V, DOI: [10.1117/12.926505](https://doi.org/10.1117/12.926505) (2012).
4. Bessell, M. S. UBVRI passbands. *Publ. Astron. Soc. Pac.* **102**, 1181–1199, DOI: [10.1086/132749](https://doi.org/10.1086/132749) (1990).
5. Okita, H., Takato, N. & Hayashi, S. S. In-situ measurement of the Subaru Telescope primary mirror reflectivity. In *Advances in Optical and Mechanical Technologies for Telescopes and Instrumentation III*, vol. 10706 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 107061U, DOI: [10.1117/12.2311899](https://doi.org/10.1117/12.2311899) (2018).
6. Mayer, B. & Kylling, A. Technical note: The libradtran software package for radiative transfer calculations - description and examples of use. *Atmospheric Chem. Phys.* **5**, 1855–1877, DOI: [10.5194/acp-5-1855-2005](https://doi.org/10.5194/acp-5-1855-2005) (2005).
7. Li, T. *et al.* aTmcam: a simple atmospheric transmission monitoring camera for sub 1% photometric precision. In *Ground-based and Airborne Instrumentation for Astronomy IV*, vol. 8446 of *Proc. SPIE*, 84462L, DOI: [10.1117/12.924792](https://doi.org/10.1117/12.924792) (2012).
8. Python Software Foundation. Python language reference, version 3.7. <https://docs.python.org/3/> (2018).
9. Carnall, A. C. SpectRes: A Fast Spectral Resampling Tool in Python. *arXiv e-prints* arXiv:1705.05165 (2017). [1705.05165](https://arxiv.org/abs/1705.05165).
10. Astropy Collaboration *et al.* Astropy: A community Python package for astronomy. *aap* **558**, A33, DOI: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068) (2013). [1307.6212](https://arxiv.org/abs/1307.6212).
11. Price-Whelan, A. M. *et al.* The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package. *aj* **156**, 123, DOI: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f) (2018).
12. Bokeh Development Team. Bokeh: Python library for interactive visualization. <https://bokeh.pydata.org> (2018).
13. Oliphant, T. E. *Guide to NumPy* (CreateSpace Independent Publishing Platform, USA, 2015), 2nd edn.
14. Jones, E., Oliphant, T., Peterson, P. *et al.* SciPy: Open source scientific tools for Python. <http://www.scipy.org/> (2001–).