

Instituto Politécnico Nacional

Escuela Superior de Cómputo

Artificial Life Robotics Laboratory
(ALIROB)

3D Cellular Automata And Pentominos

Simulations and configurations

Laura Natalia Borbolla Palacios

Winter 2018

Contents

1	Brief summary	4
1.1	Cellular Automaton	4
1.1.1	The Diffusion Rule	5
1.2	Pentominos	6
1.2.1	R-Pentomino	7
2	Experiment Results	8
2.1	The Settings	8
2.2	Pentomino By Pentomino	8
2.2.1	O-Pentomino	9
2.2.2	P-Pentomino	9
2.2.3	Q-Pentomino	10
2.2.4	R-Pentomino	13
2.2.5	S-Pentomino	16
2.2.6	T-Pentomino	21
2.2.7	U-Pentomino	21
2.2.8	V-Pentomino	23
2.2.9	W-Pentomino	23

2.2.10 X-Pentomino	26
2.2.11 Y-Pentomino	26
2.2.12 Z-Pentomino	31
2.3 Configurations found	32
2.3.1 Gliders	32
2.3.2 Oscillators	33
2.3.3 Replicator	33
2.3.4 Puffer trains	33
3 Conclusion	44
References	45
Appendices	47
A Full Pentomino Evolution	47
A.1 Pentomino P	47
A.2 Pentomino Q	47
A.3 Pentomino R	47
A.4 Pentomino S	47
A.5 Pentomino T	47

A.6	Pentomino U	47
A.7	Pentomino V	47
A.8	Pentomino W	47
A.9	Pentomino X	47
A.10	Pentomino Y	47
A.11	Pentomino Z	47

1 Brief summary

1.1 Cellular Automaton

For more information, please consult [1]–[6].

A cellular automaton is a discrete dynamical system that consists of a set of simple units (cells) that change their states at each time step, depending on the states of their neighbors (there are no actions at a distance), according to a local update rule; the cells evolve in parallel at discrete time steps and they all use the same update rule.

Also known as **CA**, the cellular automata have proved been quite useful, both as general models of complexity and as specific representations of non-linear dynamics in a great variety of fields. Despite functioning in a non-traditionally way, the **CA** can solve algorithmic problems or compute functions, all of this with the right suitable rule, of course.

Cellular automata come in a variety of shapes; the type of grid (for example, square, triangular or hexagonal cells in a two-dimensional **CA**) is one of its most fundamental properties. The neighborhood over which cells affect one another is also very important; two common neighborhoods in the case of a two-dimensional **CA** on a square grid are the Moore and the von Neumann neighborhoods.

The **elementary cellular automata** are the simplest type of cellular automaton, being a binary, nearest-neighbor, one-dimensional automaton; these **CA** have been thoroughly studied. There are 256 **ECA**.

The best known cellular automaton is the *Game of life*, discovered by J. H. Conway in 1970; this is a binary automaton with a Moore neighborhood. Over the years, it has been compiled a library of patterns with various behaviors:

- **Still life.** A fixed point pattern; the update rule keeps each cell unchanged.
- **Oscillator.** A temporally periodic pattern; the update rule may change the pattern, but after some steps, the original pattern reappears in the same location with the same orientation.
- **Spaceship.** A pattern that reappears, after some steps, although not necessarily in the same location.
- **Gun.** A pattern that, as an oscillator, periodically returns back to the initial state and emits spaceships.
- **Glider gun.** A pattern that emits gliders.

1.1.1 The Diffusion Rule

Conway's *Game of Life* inspired the scientific community and soon they started to experiment with variations of the game's rules to find new and interesting behaviors, for example, *high life*, *life 43* and *long life*; later they started experimenting with new neighborhoods, different shapes in the grids and the number of dimensions in the universe. One rule stands out of the rest: the diffusion rule; mainly because it's behavior is chaotic and with it, there has been discovered plenty of gliders, oscillators, glider guns, and puffer trains. The rule is defined as follow:

- Any living cell with less than seven neighbors dies out of loneliness.
- Any living cell with more than seven neighbors dies suffocated.
- Any living cell with seven neighbors lives onto the next generation.
- Any death cell with two living neighbors will become alive in the next generation.

1.2 Pentominos

For more information, please consult [7]–[11].

A polyomino is a finite collection of orthogonally connected cells; or, in another words, a collection of equally-sized squares that form a connected piece, meaning that each square can reach any other in it by going through adjacent squares; the order of a polyomino is the number of squares used to make it, so a fifth order polyomino (made of five squares), is called pentomino. Both terms, polyomino and pentomino, were first used by S. Golomb in 1953, in a talk to the Harvard Mathematics Club and a year later in an article. There are twelve distinct pentominoes (see Figure 1) and two naming conventions, in this paper the Conway convention (using letters *O* through *Z*) will be used; although the resemblance to the letters with this labeling scheme seems a little more strained with the other scheme, especially when using the *O* instead of *I*, it has the advantage that uses 12 consecutive letters; also, the Conway scheme tends to be used when discussing topics related to cellular automata.




Figure 1: Conway pentominoes from O through Z (left to right).

Conway used polyominoes, specially pentominoes, in his initial investigations of Life

and other cellular automata: he tracked the histories of small polyominoes as a way to explore the behavior of different cellular automata when the calculations had to be done by hand; the studies of the polyominoes led to important discoveries in Life.

1.2.1 R-Pentomino

Among the other pentominoes, the **R-pentomino** outstands; mainly because of two reasons:

- It is a methuselah¹, meaning that is a pattern that takes a large number of generations to stabilize and, at some point during its evolution, becomes much larger than it's initial configuration.
- The first glider ever observed is the glider that releases in the 69th generation.

Since all the others pentominoes stabilize in ten generations or less, this is, by far, the most active pentomino: it takes 1 103 generations and by then it's population is of 1 16.

¹M. Gardner defined methuselahs as patterns of less than ten cells that take more than fifty generations to stabilize[12].

2 Experiment Results

The experiment consisted in evolving in a three-dimensional universe, with cubes as units and an adaptation of the diffusion rule, each Conway pentomino and observing its behavior while watching out for configurations such as still lifes, oscillators gliders and especially, glider guns.

2.1 The Settings

The simulations were done in Ready, a cross-platform implementation of various reaction-diffusion systems designed to explore both, continuous and discrete cellular automata on grids and arbitrary meshes [13]. Alas, precisely because it can be used with continuous and discrete universes, Ready is not optimized for experiments of this kind, provoquing quite slow evolutions.

Each experiment started with a pentomino in the center of a 256×256 empty cube; the simulation ended when there were no more alive cells or when the cells filled the space and started to interfere one another. Since Ready does not permit to move the view when the simulation is running, each pentomino scenario was run, at least, three times in order to obtain a front, lateral and in an angle view of the evolution.

2.2 Pentomino By Pentomino

For each pentomino entry, there will be an isometric figure to illustrate the initial scenario. The pentominoes are alphabetically ordered and will be included only some moments of the evolution.




Figure 2: Isometric of the O-pentomino.

2.2.1 O-Pentomino

This is, as seen in figure 2, the simplest pentomino in the set. As observed in figure 3, in the 9^{th} generation, it splits into two identical oscillators (see figure 46), sadly, each one obstructs the way of the other and by the 21^{st} generation, all the cells are dead.

2.2.2 P-Pentomino

This pentomino (see figure 4) is quite interesting; three configurations can be seen within less than 47 generations. The first one to appear is a replicator (see figure 47), latter, the first (and one of the simplest) glider appears (see figure 41); finally, another glider appears (see figure 42).

Figure 5 shows the state of the evolution of the p-pentomino in the 42^{th} generation;




Figure 3: Evolution of the O-Pentomino, 9th generation.

the replicator can be seen in the north of the evolution; while the gliders are located in the south of the population; in 6 the replicator can be seen in the left of the figure and in 7, the gliders can be seen in the south.

The simulation of this pentomino stopped at 210 generations because space was overcrowded.

2.2.3 Q-Pentomino

This pentomino (see figure 8) shows in ten generations, two structures, one is the replicator-1 (see figure 47), and a *modified* glider: the glider-1 (see figure 41) with a satellite (see figure 43). The evolution of the 10th generation with both structures can be seen in figure 9.

As shown in 10, in the 50th generation, a mirrored glider-4 (see figure 44) emerges, both in the north and south of the population; the figures 11 and 12 show the gliders, zoomed, in the 50th and 52nd generation, respectively.

The simulation of this pentomino stopped at 109 generations because space was




Figure 4: Isometric of the P-pentomino.




Figure 5: Evolution of the P-Pentomino, 42nd generation.




Figure 6: Evolution of the P-Pentomino, zoomed replicator, 42nd generation.




Figure 7: Evolution of the P-Pentomino, zoomed gliders, 42nd generation.




Figure 8: Isometric of the Q-pentomino.

overcrowded.

2.2.4 R-Pentomino

This pentomino (see figure 13), just like it's equivalent in two dimensions, expands very quickly in a short period of time, this can be observed in figure 14, where a comparison between the evolution achieved in the same generation for different pentominoes.

The replicator (see figure 47) appears here, too; as the glider-1 (see figure 41) and glider-4 (see figure 44); the gliders can be seen in the 81st generation, in the north-east, north-west, south-east and south-west areas of figure 15; in figure 16, a pair of glider-1 can be seen easily in the left of the figure; the glider-4 is shown in the upper-left of figure 17.




Figure 9: Evolution of the Q-Pentomino, 10th generation.




Figure 10: Evolution of the Q-Pentomino, 50th generation.




Figure 11: Evolution of the Q-Pentomino, zoomed gliders, 50th generation.




Figure 12: Evolution of the Q-Pentomino, zoomed gliders, 52nd generation.




Figure 13: Isometric of the R-pentomino.

2.2.5 S-Pentomino

This pentomino (see figure 18) shows, quite fast, the replicator-1, seen in previous pentominoes first; later a clean puffer (see figure 49) emerges; four of these puffers can be observed in figure 19. This was one of the most active pentominoes; and, although the simulation had to be stopped after 131 generations. Interestingly, the growth in the population seemed more ordered, symmetrical and configuration driven: seems easier to find individual configurations in the s-pentomino than in the others. Two pairs of puffer-2 can be seen again, a pair moving towards the east, and the other towards west; in figures 20 and 21.




Figure 14: Evolution in the 52^{th} generation for the Y, W, P, X and R pentominoes respectively.




Figure 15: Evolution of the R-Pentomino, 81st generation.




Figure 16: Evolution of the R-Pentomino, zoomed glider-1, 81st generation.




Figure 17: Evolution of the R-Pentomino, zoomed glider-4, 81st generation.




Figure 18: Isometric of the S-pentomino.




Figure 19: Evolution of the S-Pentomino, 56th generation.




Figure 20: Evolution of the S-Pentomino, 57th generation.




Figure 21: Evolution of the S-Pentomino, 60th generation.

2.2.6 T-Pentomino

This pentomino (see figure 22) behaves rather differently than its equivalents in other rules (and dimensions); instead of evolving in a replicator, spaceship, pulsar or such, dies within five generations.

2.2.7 U-Pentomino

This pentomino (see figure 23), shows the puffer-1 (see figure 48) in less than 7 generations; unlike the puffer-2 (see figure 49), this puffer is not clean at all and pollutes very quickly. Interestingly enough, the population at its early stages appears to be formed by configurations that look almost like gliders and puffers found previously, but each fails a few cells. The glider-1 (see figure 41) and the replicator-1 (see figure 47) appear heavily through the evolution of the pentomino. A clean new puffer, the puffer-3 (see figure 50) and a glider, the glider-5 (see figure 45), appear in this pentomino; both, the new puffer and glider can be observed in figure 24: the glider-5 in the north and the puffer-3 in the north-east and east of the figure.




Figure 22: Isometric of the T-pentomino.




Figure 23: Isometric of the U-pentomino.




Figure 24: Evolution of the U-Pentomino, 70th generation.

2.2.8 V-Pentomino

In this pentomino (see figure 25), a pair of replicator-1 (see figure 47) appear in the early stages of its evolution. Although the replicator-1 starts by replicating itself in the z-axis, soon it starts to replicate perpendicularly to itself, as observed in figure 26.

2.2.9 W-Pentomino

This pentomino (see figure 27), unlike its predecessor, the V-pentomino, *folds* since the beginning and the two initial replicator-1 start to replicate perpendicularly since the beginning (see figure 28).




Figure 25: Isometric of the V-pentomino.




Figure 26: Evolution of the V-Pentomino, 48th generation.




Figure 27: Isometric of the W-pentomino.




Figure 28: Evolution of the W-Pentomino, 12th generation.




Figure 29: Isometric of the X-pentomino.

2.2.10 X-Pentomino

This pentomino's (see figure 29) first evolution's stages appear to be related to the oscillator-1 (see figure 46 for the oscillator and figures 30, 31, 32, 33, 34 and 35 for the pentomino's evolution). Then, as shown in figure 36, it evolves into a set of replicator-1 (see figure 47).

2.2.11 Y-Pentomino

This pentomino (see figure 37) grows quickly due to all the replicator-1 (see figure 47) present almost since the beginning of the evolution; a puffer-1 (see figure 48) before the 20th generation (see figure 38).




Figure 30: Evolution of the X-Pentomino, 2nd generation.




Figure 31: Evolution of the X-Pentomino, 3th generation.




Figure 32: Evolution of the X-Pentomino, 4th generation.




Figure 33: Evolution of the X-Pentomino, 5th generation.




Figure 34: Evolution of the X-Pentomino, 6th generation.




Figure 35: Evolution of the X-Pentomino, 7th generation.




Figure 36: Evolution of the X-Pentomino, 15th generation.




Figure 37: Isometric of the Y-pentomino.




Figure 38: Evolution of the Y-Pentomino, 16th generation.

2.2.12 Z-Pentomino

This pentomino (see figure 39),




Figure 39: Isometric of the Z-pentomino.

2.3 Configurations found

Below are presented the configurations that were found while observing the evolutions of the pentominoes; it is important to remark that, due to the fast-paced population growth in some pentominoes, it is possible that some configurations remain hidden. The configurations are listed in the order that they were found.

2.3.1 Gliders

For more information, please consult [5].

The gliders, as explained in section 1.1, are mobile particles traveling in the space. There are two types of gliders: primary and compound; the first ones cannot be decomposed into smaller mobile localizations, whereas a compound glider is made of at least




Figure 40: G1 glider in the Diffusion Rule.

two primary gliders. Some properties of the gliders are listed below:

- Volume
- Translation
- Period
- Speed
- Weight

One of the primary gliders in the Diffusion Rule is shown in figure 40.

The glider-4 (see figure 44) probably is the most simple glider; the equivalent of G1 (see figure 40) in three dimensions.


2.3.2 Oscillators

2.3.3 Replicator


2.3.4 Puffer trains

When two of this puffer-2 (see figure 49) frontally collide, a glider-4 (see figure 44) emerges.

Front



Right



Top





Figure 41: Isometric of glider-1.






Figure 42: Isometric of glider-2.

Front



Right



Top





Figure 43: Isometric of glider-3.




Figure 44: Isometric of glider-4.




Figure 45: Isometric of glider-5.




Figure 46: Isometric of oscillator-1.




Figure 47: Isometric of replicator-1.






Figure 48: Isometric of puffer-1.

Front



Right



Top





Figure 49: Isometric of puffer-2.




Figure 50: Isometric of puffer-3.

3 Conclusion

Talk about how long it took to view the simulations, even when the processor was HQ Talk about the configurations found (yayy), why it is important to find a glider gun (then, they would have "cómputo universal").

Even though the main objective was not achieved (a glider gun was not found), these experiments led to some interesting results (see section 2.3); for example, a replicator structure (see figure 47); the replicator-1, although could not be probed in this set of experiments, this configuration appears to be somehow important in the process of generating gliders; this supposition is made by the observation of the evolution of the s-pentomino population; at least four gliders were created and the pattern that appeared to create them kept replicating in the population; but the more the population grew, the more the population got in its way.

References

- [1] F. Berto and J. Tagliabue, “Cellular automata,” in *The Stanford Encyclopedia of Philosophy*, E. N. Zalta, Ed., Fall 2017, Metaphysics Research Lab, Stanford University, 2017.
- [2] J. Kari, *Cellular Automata*. University of Turku, 2013.
- [3] E. W. Weisstein, *Cellular automaton*. [Online]. Available: <http://mathworld.wolfram.com/CellularAutomaton.html>.
- [4] G. Juárez-Martínez, *Diffusion rule*, 2017. [Online]. Available: http://uncomp.uwe.ac.uk/genaro/Diffusion_Rule/diffusionLife.html.
- [5] G. Juárez-Martínez, A. Adamatzky, and H. McIntosh, “Localization dynamics in a binary two-dimensional cellular automaton: The diffusion rule,” *Journal of Cellular Automata*, 2006.
- [6] S. University, *Conway’s game of life*, 2017. [Online]. Available: <http://web.stanford.edu/~cdebs/GameOfLife/>.
- [7] “Lecture notes on computer science theoretical computer science and general issues,” in. Springer, Mar. 2014, vol. 8370, ISBN: 9783319049205. [Online]. Available: <https://books.google.com.mx/books?id=dBW6BQAAQBAJ&printsec=frontcover#v=onepage&q&f=false>.
- [8] *Polyomino*, 2019. [Online]. Available: <http://www.conwaylife.com/wiki/Polyomino>.
- [9] M. E. Coppenbarger, *Pentominoes - introduction*, 2019. [Online]. Available: <https://people.rit.edu/mecsma/Professional/Puzzles/Pentominoes/P-Intro.html>.
- [10] *R-pentomino*, 2019. [Online]. Available: <http://www.conwaylife.com/wiki/R-pentomino>.
- [11] *Methuselah*, 2019. [Online]. Available: <http://www.conwaylife.com/wiki/Methuselah>.

- [12] M. Gardner, *Wheels, Life and Other Mathematical Amusements*. W. H. Freeman, 1983, ISBN: 9780716715887.
- [13] T. Hutton, R. Munafo, A. Trevorow, T. Rokicki, and D. Wills, *Ready*, 2019. [Online]. Available: <https://github.com/GollyGang/ready>.

Appendices

A Full Pentomino Evolution

Below are the full evolutions of each pentomino.

A.1 Pentomino P

A.2 Pentomino Q

A.3 Pentomino R

A.4 Pentomino S

A.5 Pentomino T

A.6 Pentomino U

A.7 Pentomino V

A.8 Pentomino W

A.9 Pentomino X

A.10 Pentomino Y

A.11 Pentomino Z