

Notes on Nocedal and Wright's "Numerical
Optimization"
Chapter 7 – "Calculating Derivatives"

Lucas N. Ribeiro

Introduction I

- ▶ Sometimes the user can provide a function that calculates the gradient and the Hessian; sometimes not.
- ▶ When these information are not available, we can calculate them by ourselves
- ▶ There are basically three approaches to achieve that:
 - ▶ Finite differencing: Observe the response of the function of interest to small perturbations, e.g., using the central difference formula;
 - ▶ Automatic differentiation: breaks the function down into elementary operations and applies the chain rule.
 - ▶ Symbolic differentiation: algebraic specification and symbolic manipulation. Used in Maple, MATLAB, etc.
- ▶ Besides algorithms, gradients and Hessians are useful for sensitivity analysis in economics, etc.

Finite-Difference Derivative Approximations I

Let us first derive the forward-difference formula from Taylor's theorem. If f is twice continuously differentiable, then:

$$f(x + p) = f(x) + \nabla f(x)^T p + 0.5 p^T \nabla^2 f(x + tp) p \quad (1)$$

Let us assume L is a bound on the norm of $\nabla^2 f(\cdot)$. It means that

$$\exists L \mid \|\nabla^2 f\| < L$$

for whatever argument of the Hessian. Then, the norm of the last term in (1) is bounded as

$$\|0.5 p^T \nabla^2 f(x + tp) p\| < 0.5 \|p^T\| \|\nabla^2 f(x + tp)\| \|p\| \quad (2)$$

$$< (L/2) \|p\|^2 \quad (3)$$

Finite-Difference Derivative Approximations II

where we used Cauchy-Schwarz. By isolating the quadratic term in (1) and applying its bound, we get:

$$\|f(x + p) - f(x) - \nabla f(x)^T p\| \leq (L/2)\|p\|^2 \quad (4)$$

Now, set $p = \epsilon e_i$, where e_i is the i th canonical vector. Then

$$\nabla f(x)^T p = \epsilon \partial f / \partial x_i$$

Equation (4) can be rearranged as

$$\|f(x + \epsilon e_i) - f(x) - \epsilon \partial f / \partial x_i\| \leq (L/2)\epsilon^2 \quad (5)$$

The inequality above implies

$$\partial f / \partial x_i = \frac{f(x + \epsilon e_i) - f(x)}{\epsilon} + \delta \quad (6)$$

Finite-Difference Derivative Approximations III

where $|\delta| \leq (L/2)\epsilon$. When $\epsilon \rightarrow 0$, the error vanishes and the finite difference goes to the partial derivative.

A typical choice for ϵ is $\sqrt{\mathbf{u}}$, where \mathbf{u} is the unit round-off.

A more accurate approximation is the central difference formula.

Assuming the existence of second order derivative of f and Lipschitz continuity:

$$f(x + p) = f(x) + \nabla f(x)^T p + 0.5 p^T \nabla^2 f(x + tp) p \quad (7)$$

$$= f(x) + \nabla f(x)^T p + 0.5 p^T \nabla^2 f(x) p + O(\|p\|^3). \quad (8)$$

We set $p = \epsilon e_i$ and $p = -\epsilon e_i$. Then we have:

$$f(x + \epsilon e_i) = f(x) + \epsilon \frac{\partial f}{\partial x_i} + 0.5 \epsilon^2 \frac{\partial^2 f}{\partial x_i^2} + O(\epsilon^3) \quad (9)$$

$$f(x - \epsilon e_i) = f(x) - \epsilon \frac{\partial f}{\partial x_i} + 0.5 \epsilon^2 \frac{\partial^2 f}{\partial x_i^2} + O(\epsilon^3) \quad (10)$$

Finite-Difference Derivative Approximations IV

Subtracting (10) from (9) and dividing by 2ϵ gives:

$$\frac{\partial f}{\partial x_i} = \frac{f(x + \epsilon e_i) - f(x - \epsilon e_i)}{2\epsilon} + O(\epsilon^2) \quad (11)$$

- ▶ Now the error is $O(\epsilon^2)$ (unlike $O(\epsilon)$ in the forward-difference)
- ▶ More complex, though. Has to evaluate f at two different points.

Approximating a Jacobian – Now consider a function $r : \mathbb{R}^n \rightarrow \mathbb{R}^m$. The Jacobian matrix is defined as $J(x) = [\partial r_m / \partial x_n]$. Using Taylor's theorem, we can deduce that

$$\|r(x + p) - r(x) - J(x)p\| \leq (L/2)\|p\|^2$$

Finite-Difference Derivative Approximations V

Sometimes we wish an estimate for $J(x)p$ instead of the full Jacobian. In this case, we have

$$J(x)p = \frac{r(x + \epsilon p) - r(x)}{\epsilon} + O(\epsilon)$$

When we wish the full matrix, we can compute one column at a time:

$$\partial r / \partial x_i = \frac{r(x + \epsilon e_i) - r(x)}{\epsilon} + O(\epsilon)$$

Computationally efficient methods that exploit possible sparsity can be derived.

Approximating the Hessian – From Taylor's theorem:

$$\nabla f(x + p) = \nabla f(x) + \nabla^2 f(x)p + O(\|p\|^2)$$

and

$$\partial \nabla f / \partial x_i = \frac{\nabla f(x + \epsilon e_i) - \nabla f(x)}{\epsilon}$$

Finite-Difference Derivative Approximations VI

Sometimes we wish to estimate only $\nabla^2 f(x)p$ and we can achieve it by

$$\nabla^2 f(x)p \approx \frac{\nabla f(x + \epsilon p) - \nabla f(x)}{\epsilon}$$

Note that, so far, we require knowledge of the gradient vector. When it's not available we can plug our gradient approximations into that of Hessian:

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{f(x + \epsilon e_i + \epsilon e_j) - f(x + \epsilon e_i) - f(x + \epsilon e_j) + f(x)}{\epsilon^2} + O(\epsilon).$$

(12)

Automatic Differentiation

- ▶ Any function (no matter how complicated) is built by simple operation such as sums, multiplications and exponentiations.
- ▶ The basic principle of automatic differentiation is breaking down functions in these operations and applying the chain rule.