

## Analyse Mathématique

Author: CatMono

**Date:** July, 2025

Version: 0.1

## **Contents**

Preface	
Chapter	r 1 Preliminaries
1.1	Trigonometric Formulas
Chapte	r 2 Limits of Sequences and Continuity of Real Number System
2.1	Convergent Sequences
2.2	Indeterminate Form
2.3	Subsequences
2.4	Proposition on The Continuity of Real Numbers
Chapte	r 3 Limits and Continuity of Functions
3.1	Limits of Functions
3.2	Continuous Functions
3.3	Infinitesimal and Infinite Quantities
3.4	Continuous Functions on Closed Intervals
3.5	Period Three Implies Chaos
3.6	Functional Equations
Chapte	r 4 Differential
4.1	Differential and Derivative
4.2	Higher-Order Derivatives
4.3	Differential Mean Value Theorems
4.4	Theorems and Applications concerning Derivatives
4.5	Taylor's Theorem
4.6	Applications of Taylor's Theorem
Chapte	r 5 Integral
Chapte	r 6 Series of Numbers
Chapte	r 7 Series of Functions
Chapte	r 8 Power Series
Chapte	r 9 Limits and Continuity in Euclidean Spaces
9.1	Continuous Mappings
Chapte	r 10 Multi-variable Differential Calculus
10.1	Directional Derivatives and Total Differential
10.2	Differential of Vector-Valued Functions
10.3	Derivatives of Composite Mappings (Chain Rule)
10.4	Mean Value Theorem and Taylor's Formula

21

## **Preface**

This is the preface of the book...

## **Chapter 1 Preliminaries**

### 1.1 Trigonometric Formulas

### **Product-to-Sum Formulas:**

$$\sin \alpha \cos \beta = \frac{1}{2} \left[ \sin(\alpha + \beta) + \sin(\alpha - \beta) \right]$$

$$\cos \alpha \sin \beta = \frac{1}{2} \left[ \sin(\alpha + \beta) - \sin(\alpha - \beta) \right]$$

$$\cos \alpha \cos \beta = \frac{1}{2} \left[ \cos(\alpha + \beta) + \cos(\alpha - \beta) \right]$$

$$\sin \alpha \sin \beta = -\frac{1}{2} \left[ \cos(\alpha + \beta) - \cos(\alpha - \beta) \right]$$

### Sum and Difference Formulas:

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$
$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$
$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$
$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

### **Sum-to-Product Formulas:**

$$\sin \alpha + \sin \beta = 2 \sin \left(\frac{\alpha + \beta}{2}\right) \cos \left(\frac{\alpha - \beta}{2}\right)$$

$$\sin \alpha - \sin \beta = 2 \sin \left(\frac{\alpha - \beta}{2}\right) \cos \left(\frac{\alpha + \beta}{2}\right)$$

$$\cos \alpha + \cos \beta = 2 \cos \left(\frac{\alpha + \beta}{2}\right) \cos \left(\frac{\alpha - \beta}{2}\right)$$

$$\cos \alpha - \cos \beta = -2 \sin \left(\frac{\alpha + \beta}{2}\right) \sin \left(\frac{\alpha - \beta}{2}\right)$$

### **Double Angle Formulas:**

$$\sin 2\alpha = 2\sin \alpha \cos \alpha$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = 2\cos^2 \alpha - 1 = 1 - 2\sin^2 \alpha$$

$$\tan 2\alpha = \frac{2\tan \alpha}{1 - \tan^2 \alpha}$$

### Half Angle Formulas:

$$\sin \frac{\alpha}{2} = \pm \sqrt{\frac{1 - \cos \alpha}{2}}$$

$$\cos \frac{\alpha}{2} = \pm \sqrt{\frac{1 + \cos \alpha}{2}}$$


$$\tan \frac{\alpha}{2} = \frac{1 - \cos \alpha}{\sin \alpha} = \frac{\sin \alpha}{1 + \cos \alpha}$$

### **Power-Reducing Formulas:**

$$\sin^2 \alpha = \frac{1 - \cos 2\alpha}{2}$$
$$\cos^2 \alpha = \frac{1 + \cos 2\alpha}{2}$$

### **Angle Decomposition Formulas:**

$$\sin^2 \alpha - \sin^2 \beta = \sin(\alpha + \beta)\sin(\alpha - \beta)$$
$$\cos^2 \alpha - \sin^2 \beta = \cos(\alpha + \beta)\cos(\alpha - \beta)$$



### Remark

- On the gray triangle, the sum of the squares of the two numbers above is equal to the square of the number below, for instance,  $\tan^2 x + 1 = \sec^2 x$
- The three trigonometric functions in the clockwise direction have the following properties:  $\tan x = \frac{\sin x}{\cos x}$ , etc.

## Chapter 2 Limits of Sequences and Continuity of Real Number System

### 2.1 Convergent Sequences

- ¶ Convergent Sequences
- ¶ Properties of Convergent Sequences
- ¶ Cauchy's Proposition and Fitting Method

### Proposition 2.1 (Cauchy's Proposition)

Let  $\lim_{n\to\infty} x_n = l$ , then:

$$\lim_{n \to \infty} \frac{x_1 + x_2 + \dots + x_n}{n} = l.$$



- 1. In the proposition, l can be  $+\infty$  or  $-\infty$ .
- 2. Let  $\lim_{n\to\infty} x_n = l$ , then:

$$\lim_{n \to \infty} \frac{x_1 + x_2 + \dots + x_n}{n} = \lim_{n \to \infty} \sqrt[n]{x_1 x_2 \cdots x_n} = \lim_{n \to \infty} \frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}} = l.$$

It can be proved directly by Stolz's theorem. On top of that, it can also be proved by the **fitting method**.



Remark To prove  $\lim_{n\to\infty} x_n = A$ , the key is to show that  $|x_n - A|$  can be arbitrarily small. For this purpose, it is generally recommended to simplify the expression of  $x_n$  as much as possible. However, in some cases, A can also be transformed into a form similar to  $x_n$ . This method is called the fitting method. The core idea behind the method of fitting is to appropriately divide into units of 1 for analysis.

### 2.2 Indeterminate Form

- ¶ Infinitely Large Quantities and Infinitesimal Quantities
- ¶ Indeterminate Forms

### Theorem 2.1 (Stolz's Theorem

**Type**  $\frac{0}{0}$  Let  $\{a_n\}, \{b_n\}$  be two infinitesimal sequences, where  $\{a_n\}$  is also a strictly monotonic decreasing sequence. If

$$\lim_{n\to\infty} \frac{b_{n+1}-b_n}{a_{n+1}-a_n} = l \text{ (finite or } \pm \infty),$$

then

$$\lim_{n \to \infty} \frac{a_n}{b_n} = l.$$

**Type**  $\frac{*}{\infty}$  Let  $\{a_n\}$  be a strictly monotonic increasing sequence of divergent large quantities. If

$$\lim_{n \to \infty} \frac{b_{n+1} - b_n}{a_{n+1} - a_n} = l \text{ (finite or } \pm \infty),$$

then

$$\lim_{n \to \infty} \frac{a_n}{b_n} = l.$$



### Note

- 1. The inverse proposition of Stolz's Theorem does not hold.
- 2. If  $a_1$  is an undefined infinite quantity  $\infty$ , Stolz's Theorem does not hold.

## 2.3 Subsequences

- ¶ Subsequences
- $\P$  Upper Limits and Lower Limits

## 2.4 Proposition on The Continuity of Real Numbers

- ¶ Dedkind Cut
- ¶ Least Supremum Property
- ¶ Monotone Convergence Theorem
- $\P$  Bolzano-Weierstrass Theorem
- $\P$  Nested Interval Theorem
- ¶ Cauchy Criterion for Sequences
- ¶ Heine-Borel Theorem

## **Chapter 3 Limits and Continuity of Functions**

- 3.1 Limits of Functions
- 3.2 Continuous Functions
- 3.3 Infinitesimal and Infinite Quantities
- **3.4 Continuous Functions on Closed Intervals**
- 3.5 Period Three Implies Chaos
- 3.6 Functional Equations

## **Chapter 4 Differential**

### 4.1 Differential and Derivative

## **4.2 Higher-Order Derivatives**

### 4.3 Differential Mean Value Theorems

### Definition 4.1 (Extremum)

Let f(x) is defined on  $(a,b), x_0 \in (a,b)$ . If there exists  $U(x_0,\delta) \subset (a,b)$  such that  $f(x) \leqslant f(x_0)$  on it, then  $x_0$  is called a local maximum point of f, and  $f(x_0)$  is referred to as the corresponding local maximum value. The definition of the minimum value is analogous.

### Lemma 4.1 (Fermat's Lemma)

If f is differentiable at  $x_0$  which is a local extremum, then  $f'(x_0) = 0$ .

### Theorem 4.1 (Rolle's Theorem

If  $f \in C[a,b]$ ,  $f \in D(a,b)$  and f(a) = f(b), then there exists  $\xi \in (a,b)$  such that  $f'(\xi) = 0$ . Enhanced Version: If  $f \in D(a,b)$  (finite or infinite interval), and  $\lim_{x \to a^+} f(x) = \lim_{x \to b^-} f(x)$ , then there exists  $\xi \in (a,b)$  such that  $f'(\xi) = 0$ .

### Theorem 4.2 (Lagrange's Mean Value Theorem,

If  $f \in C[a, b]$ ,  $f \in D(a, b)$ , then there exists  $\xi \in (a, b)$  such that

$$f'(\xi) = \frac{f(b) - f(a)}{b - a}.$$

### Theorem 4.3 (Cauchy's Mean Value Theorem

If  $f,g\in C[a,b], f,g\in D(a,b)$  and  $g'(x)\neq 0$  for all  $x\in (a,b)$ , then there exists  $\xi\in (a,b)$  such that

$$\frac{f'(\xi)}{g'(\xi)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$$

## 4.4 Theorems and Applications concerning Derivatives

### Theorem 4.4 (Darboux's Intermediate Value Theorem for Derivatives)

If  $f(x) \in D[a, b]$ , and  $f'_+(a) \cdot f'_-(b) < 0$ , then there at least exists  $\xi \in (a, b)$  such that  $f'(\xi) = 0$ .

### Theorem 4.5 (Theorem on the Limit of Derivatives)

If  $f(x) \in C(U(x_0))$ ,  $\mathring{D}(U(x_0))$ , and  $\lim_{x \to x_0} f'(x) = A$ , then f is differentiable at  $x_0$  and  $f'(x_0) = A$ .

Remark In fact,  $\lim_{x o x_0}f'(x)=A$  has already been shown to imply that  $f\in \mathring{D}(U(x_0)).$ 

- 4.5 Taylor's Theorem
- **4.6** Applications of Taylor's Theorem

## Chapter 5 Integral

# **Chapter 6 Series of Numbers**

# **Chapter 7 Series of Functions**

# **Chapter 8 Power Series**

## **Chapter 9 Limits and Continuity in Euclidean Spaces**

## 9.1 Continuous Mappings

- Continuous Mappings on Compact Sets
- Continuous Mappings on Connected Sets

### Definition 9.1 (Connected Set)

Let S be a set of points in  $\mathbb{R}^n$ . If a continuous mapping

$$\gamma:[0,1]\to\mathbb{R}^n$$

satisfies that the range of  $\gamma([0,1])$  lies entirely within S, we call  $\gamma$  a path in S, where  $\gamma(0)$  and  $\gamma(1)$  are referred to as the starting point and ending point of the path, respectively.

If for any two points  $\mathbf{x}, \mathbf{y} \in S$ , there exists a path in S with  $\mathbf{x}$  as the starting point and  $\mathbf{y}$  as the ending point, Sis called path-connected, or equivalently, S is called a connected set.

A connected open set is called an (open) region. The closure of an (open) region is referred to as a closed region.

Remark Intuitively, this means that any two points in S can be connected by a curve lying entirely within S. Clearly, a connected subset of  $\mathbb{R}$  is an interval, and a connected subset of  $\mathbb{R}$  is compact if and only if it is a closed interval.

## **Chapter 10 Multi-variable Differential Calculus**

### 10.1 Directional Derivatives and Total Differential

#### Directional Derivative ¶

### Definition 10.1 (Directional Derivative)

Let  $U \subset \mathbb{R}^n$  be an open set,  $f: U \to \mathbb{R}^1$ , **e** is a unit vector in  $\mathbb{R}^n$ ,  $\mathbf{x}^0 \in U$ . Define

$$u(t) = f(\mathbf{x}^0 + t\mathbf{e}).$$

If the derivative of u at t = 0

$$u'(0) = \lim_{t \to 0} \frac{u(t) - u(0)}{t} = \lim_{t \to 0} \frac{f(\mathbf{x}^0 + t\mathbf{e}) - f(\mathbf{x}^0)}{t}$$

exists and is finite, it is called the directional derivative of f at  $\mathbf{x}^0$  in the direction e, denoted by  $\frac{\partial f}{\partial \mathbf{e}}(\mathbf{x}^0)$ . It is the rate of change of f at  $\mathbf{x}^0$  in the direction  $\mathbf{e}$ .

Consider the following set of unit coordinate vectors:  $\mathbf{e}_1, \mathbf{e}_2, \cdots, \mathbf{e}_n$ . Let  $\mathbf{e}_i = (0, 0, \cdots, 0, 1, 0, \cdots, 0)$ denote the standard orthonormal basis in  $\mathbb{R}^n$ , where the 1 appears in the *i*-th position. That is,

$$\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \delta_{ij} = \begin{cases} 1, & i = j, \\ 0, & i \neq j. \end{cases}$$

For a function f, the directional derivative of f at the point  $\mathbf{x}_0$  in the direction of  $\mathbf{e}_i$  is called the ith first-order **partial derivative** of f at  $\mathbf{x}^0$ , denoted by

$$\frac{\partial f}{\partial x_i}(\mathbf{x}^0)$$
 or  $D_i f(\mathbf{x}^0)$  or  $f_{x_i}(\mathbf{x}^0)$   $(i = 1, 2, \dots, n)$ .

 $\mathrm{D}_i = \frac{\partial}{\partial x_i}$  is called the ith partial differential operator ( $i=1,2,\cdots,n$ ). Let  $\mathbf{e}_i = \sum_{i=0}^n \mathbf{e}_i \cos \alpha$  be a unit vector, where  $\sum_{i=0}^n \cos^2 \alpha = 1$ . If  $\frac{\partial f}{\partial x_i}$  is continuous at  $\mathbf{x}^0$ , then the directional derivative of f at  $\mathbf{x}^0$  along the direction  $\mathbf{e}$  is given by:

$$\frac{\partial f}{\partial \mathbf{e}}(\mathbf{x}^0) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(\mathbf{x}^0) \cos \alpha_i.$$

This is the formula for expressing a directional derivative using partial derivatives.

 $ilde{Y}$  Note Let  ${f e}$  be a direction, then  $\|-{f e}\|=\|{f e}\|=1$ , which implies that  $-{f e}$  is also a direction. At this point, we have:

$$\frac{\partial f}{\partial (-\mathbf{e})}(\mathbf{x}^0) = -\frac{\partial f}{\partial \mathbf{e}}(\mathbf{x}^0).$$

### Definition 10.2 (Jacobian Matrix (Gradient))

Let

$$Jf(\mathbf{x}) = (D_1 f(\mathbf{x}), D_2 f(\mathbf{x}), \dots, D_n f(\mathbf{x})),$$

which is called the **Jacobian matrix** of the function f at the point  $\mathbf{x}$ , (a 1  $\times$  n matrix) whose counterpart is the first-order derivative of a single-variable function.

Henceforth, we represent the point x in  $\mathbb{R}^n$  and its increments h as column vectors. In this way, the differential of the function can be expressed using matrix multiplication as follows:

$$df(\mathbf{x}^0)(\mathbf{\Delta}\mathbf{x}) = Jf(\mathbf{x}^0)\mathbf{\Delta}\mathbf{x}.$$

The Jacobian matrix of the function f is also frequently denoted as grad f (or  $\nabla f$ ), that is,

$$\operatorname{grad} f(\mathbf{x}) = Jf(\mathbf{x}),$$

which is called the **gradient** of the scalar function f.

### Total Differential

### Definition 10.3 (Total Differential)

Let  $U \subset \mathbb{R}^n$  be an open set,  $f: U \to \mathbb{R}^1$ ,  $\mathbf{x}^0 \in U$ ,  $\Delta \mathbf{x} = (\Delta x_1, \Delta x_2, \cdots, \Delta x_n) \in \mathbb{R}^n$ . If

$$f(\mathbf{x}^0 + \Delta \mathbf{x}) - f(\mathbf{x}^0) = \sum_{i=1}^n A_i \Delta x_i + o(\|\Delta \mathbf{x}\|) \qquad (\|\Delta \mathbf{x}\| \to 0),$$

where  $A_1, A_2, \ldots, A_n$  are constants independent of  $\Delta \mathbf{x}$ , then the function f is said to be differentiable at the point  $\mathbf{x}^0$ , and the linear main part  $\sum_{i=1}^n A_i \Delta x_i$  is called the **total differential** of f at  $\mathbf{x}^0$ , denoted as

$$df(\mathbf{x}^0)(\mathbf{\Delta}\mathbf{x}) = \sum_{i=1}^n A_i \Delta x_i.$$

If f is differentiable at every point in the open set U, then f is called a differentiable function on U.



**Necessary Condition** If an n-variable function f is differentiable at the point  $x_0$ , then f is continuous at  $x^0$ and possesses first-order partial derivatives  $\frac{\partial f}{\partial x_i}(\mathbf{x}^0)$  at  $\mathbf{x}^0$  for  $i=1,2,\ldots,n$ , and

$$\mathbf{A} = (A_1, A_2, \dots, A_n) = Jf(\mathbf{x}^0) = (D_1 f(\mathbf{x}^0), D_2 f(\mathbf{x}^0), \dots, D_n f(\mathbf{x}^0)).$$

<sup>a</sup> However, the converse is not true.

**Sufficient Condition** Let  $U\subset\mathbb{R}^n$  be an open set, and let  $f:U\to\mathbb{R}^1$  be an n-variable function. If Jf= $(D_1f, D_2f, \dots, D_nf)$  is continuous at  $\mathbf{x}^0$  (i.e.,  $\frac{\partial f}{\partial x_i}$  is continuous at  $\mathbf{x}^0$  for  $i=1,2,\dots,n$ ), then f is differentiable at  $\mathbf{x}^0$ . However, the converse is not necessarily true.

<sup>a</sup>It is referred to as the total differential formula, and the more common form is

$$df(x_0, y_0) = \frac{\partial f}{\partial x}(x_0, y_0) dx + \frac{\partial f}{\partial y}(x_0, y_0) dy.$$



### Note 🍄

- The continuity of the derivative function at  $\mathbf{x}^0$  implies that the original function f is differentiable in some neighborhood of  $\mathbf{x}^0$ .
- In fact, this condition can be relaxed to require that one partial derivative exists at the point, while the remaining n-1partial derivative functions are continuous at that point.



**Proof** Taking a function of three variables as an example.

Assume the 3-ary function  $f: \mathbb{R}^3 \to \mathbb{R}$  meets:

- 1. There exists  $f_z(x_0, y_0, z_0)$ .
- 2. The partial derivative functions  $f_x(x, y, z)$  and  $f_y(x, y, z)$  are continuous at  $(x_0, y_0, z_0)$ , i.e. there are partial derivatives in some neighborhood of  $(x_0, y_0, z_0)$ .

Consider the total increment of f at the point  $(x_0, y_0, z_0)$ :

$$\Delta f = \underbrace{\left[f(x_0 + \Delta x, y_0 + \Delta y, z_0 + \Delta z) - f(x_0, y_0 + \Delta y, z_0 + \Delta z)\right]}_{I_1} + \underbrace{\left[f(x_0, y_0 + \Delta y, z_0 + \Delta z) - f(x_0, y_0, z_0 + \Delta z)\right]}_{I_2} + \underbrace{\left[f(x_0, y_0, z_0 + \Delta z) - f(x_0, y_0, z_0)\right]}_{I_3}.$$

For  $I_1, I_2$ , by the Lagrange's Mean Value Theorem of unary functions, there exist  $\theta_1, \theta_2 \in (0,1)$  such that

$$I_1 = f_x(x_0 + \theta_1 \Delta x, y_0 + \Delta y, z_0 + \Delta z) \Delta x,$$
  

$$I_2 = f_y(x_0, y_0 + \theta_2 \Delta y, z_0 + \Delta z) \Delta y.$$

Then by the continuity of the their partial derivatives at  $(x_0,y_0,z_0)$ , we have

$$\lim_{\Delta x, \Delta y, \Delta z \to 0} I_1 = f_x(x_0, y_0, z_0) \Delta x, \quad \lim_{\Delta x, \Delta y, \Delta z \to 0} I_2 = f_y(x_0, y_0, z_0) \Delta y.$$

They can be expressed in terms of infinitesimals( $\rho = \sqrt{\Delta x^2 + \Delta y^2 + \Delta z^2}$ ):

$$I_1 = f_x(x_0, y_0, z_0) \Delta x + \alpha_1 \Delta x, \quad \alpha_1 \to 0 (\rho \to 0),$$
  

$$I_2 = f_y(x_0, y_0, z_0) \Delta y + \alpha_2 \Delta y, \quad \alpha_2 \to 0 (\rho \to 0).$$

For  $I_3$ , by the definition of the partial derivative  $f_z(x, y, z)$  at  $(x_0, y_0, z_0)$ , we have

$$I_3 = f_z(x_0, y_0, z_0)\Delta z + \alpha_3 \Delta z, \quad \alpha_3 \to 0 (\rho \to 0).$$

Accordingly,

$$\begin{split} \Delta f &= I_1 + I_2 + I_3 \\ &= \left[ f_x(x_0, y_0, z_0) \Delta x + \alpha_1 \Delta x \right] + \left[ f_y(x_0, y_0, z_0) \Delta y + \alpha_2 \Delta y \right] + \left[ f_z(x_0, y_0, z_0) \Delta z + \alpha_3 \Delta z \right] \\ &= f_x(x_0, y_0, z_0) \Delta x + f_y(x_0, y_0, z_0) \Delta y + f_z(x_0, y_0, z_0) \Delta z + \left[ \alpha_1 \Delta x + \alpha_2 \Delta y + \alpha_3 \Delta z \right]. \end{split}$$

Apparently,

$$\lim_{\rho \to 0} \frac{\alpha_1 \Delta x + \alpha_2 \Delta y + \alpha_3 \Delta z}{\rho} = 0,$$

i.e.  $\alpha_1 \Delta x + \alpha_2 \Delta y + \alpha_3 \Delta z = o(\rho)$ . Therefore, f(x,y,z) is differentiable at  $(x_0,y_0,z_0)$ , which completes the proof.

### Ŷ Note (At some point)

- 1. Differentiable
  - $\Longrightarrow$  Continuous
  - $\Longrightarrow$  Partial derivatives exist:  $D_{\vec{u}} = \nabla f \cdot \vec{u}$
- 2. Directional Derivative
  - All directional derivatives exist  $\iff$  differentiable or continuous.
  - All directional derivatives exist and are equal  $\iff$  differentiable.
- 3. Partial Derivative
  - The continuity and existence of directional/partial derivatives are mutually exclusive.

### ¶ Higher-Order Partial Derivatives and Differential

If the first-order partial derivative of f,  $\frac{\partial f}{\partial x_i}$ , itself possesses partial derivatives, then the second-order

partial derivative of *f* is defined, and is denoted as follows(the first is also called the mixed partial derivative):

$$f_{x_i x_j} = \frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial}{\partial x_i} \left( \frac{\partial f}{\partial x_i} \right), \quad f_{x_i x_i} = \frac{\partial^2 f}{\partial x_i^2} = \frac{\partial}{\partial x_i} \left( \frac{\partial f}{\partial x_i} \right), \quad i, j = 1, 2, \dots, n.$$

Similarly, higher-order partial derivatives of order  $3, 4, \dots m, \dots$  can be defined.

The following theorem provides the conditions under which mixed partial derivatives are equal.

### Theorem 10.2 (Conditions for Equality of Mixed Partial Derivatives)

1. Let  $U \subset \mathbb{R}^2$  be an open set, and  $f: U \to \mathbb{R}$  be a function of two variables. If  $f_{xy}$  and  $f_{yx}$  are continuous at  $(x_0, y_0) \in U$ , then


$$f_{xy}(x_0, y_0) = f_{yx}(x_0, y_0).$$

2. Let  $U \subset \mathbb{R}^n$  be an open set, and  $f: U \to \mathbb{R}$  be a function of n variables. If f has partial derivatives up to order k in D, and all of them are continuous at  $\mathbf{x}^0 = (x_1^0, x_2^0, \dots, x_n^0) \in U$ , then

$$\frac{\partial^l f}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_l}}(\mathbf{x}^0) = \frac{\partial^l f}{\partial x_{i_2} \partial x_{i_1} \cdots \partial x_{i_l}}(\mathbf{x}^0) = \cdots = \frac{\partial^l f}{\partial x_{i_l} \partial x_{i_{l-1}} \cdots \partial x_{i_1}}(\mathbf{x}^0),$$

that is, the order of taking partial derivatives  $l(\leq k)$  does not affect the result.

<sup>&</sup>quot;If the condition " $f_{xy}$  and  $f_{yx}$  are continuous at  $(x_0, y_0)$ ", is not satisfied, then the conclusion " $f_{xy}(x_0, y_0) = f_{yx}(x_0, y_0)$ " does not necessarily hold.



$$\varphi(y) = f(x_0 + h, y) - f(x_0, y),$$

and

$$\psi(x) = f(x, y_0 + k) - f(x, y_0).$$

Applying the Lagrange Mean Value Theorem, we have

$$\begin{split} &[f(x_0+h,y_0+k)-f(x_0,y_0+k)]-[f(x_0+h,y_0)-f(x_0,y_0)]\\ =&\varphi(y_0+k)-\varphi(y_0)\\ =&\varphi'(y_0+\theta_1k)k\\ =&[f_y(x_0+h,y_0+\theta_1k)-f_y(x_0,y_0+\theta_1k)]k\\ =&f_{yx}(x_0+\theta_2h,y_0+\theta_1k)hk,\quad 0<\theta_1,\theta_2<1. \end{split}$$

On the other hand.

$$[f(x_0 + h, y_0 + k) - f(x_0, y_0 + k)] - [f(x_0 + h, y_0) - f(x_0, y_0)]$$

$$= [f(x_0 + h, y_0 + k) - f(x_0 + h, y_0)] - [f(x_0, y_0 + k) - f(x_0, y_0)]$$

$$= \psi(x_0 + h) - \psi(x_0)$$

$$= \psi'(x_0 + \theta_3 h) h$$

$$= [f_x(x_0 + \theta_3 h, y_0 + k) - f_x(x_0 + \theta_3 h, y_0)] h$$

$$= f_{xy}(x_0 + \theta_3 h, y_0 + \theta_4 k) hk, \quad 0 < \theta_3, \theta_4 < 1.$$

Therefore,

$$f_{xy}(x_0 + \theta_3 h, y_0 + \theta_4 k) = f_{yx}(x_0 + \theta_2 h, y_0 + \theta_1 k).$$

Since  $f_{xy}$  and  $f_{yx}$  are continuous at  $(x_0, y_0)$ , letting  $h \to 0, k \to 0$ , we obtain

$$f_{xy}(x_0, y_0) = f_{yx}(x_0, y_0).$$

By applying 10.1 and the principle of mathematical induction, one can immediately derive the following result.

Suppose z=f(x,y) has continuous partial derivatives in the domain  $U\subset\mathbb{R}^2$ . Then z is differentiable, and

$$\mathrm{d}z = \frac{\partial z}{\partial x} \mathrm{d}x + \frac{\partial z}{\partial y} \mathrm{d}y.$$

If z also has continuous second-order partial derivatives, then  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  are also differentiable, and thus  $\mathrm{d}z$  is differentiable. We call the differential of  $\mathrm{d}z$  the second-order differential of z, denoted as

$$d^2z = d(dz).$$

In general, based on the k-th order differential (d $^kz$  of z, its (k+1)-th order differential (if it exists) is defined as

$$d^{k+1}z = d(d^k z), \quad k = 1, 2, \cdots.$$

Due to the fact that for the independent variables x and y, we always have

$$d^2x = d(dx) = 0,$$
  $d^2y = d(dy) = 0,$ 

the second-order differential of z = f(x, y) is given by

$$\begin{split} \mathrm{d}^2z &= \mathrm{d}(\mathrm{d}z) = \mathrm{d}\left(\frac{\partial z}{\partial x}\mathrm{d}x + \frac{\partial z}{\partial y}\mathrm{d}y\right) \\ &= \mathrm{d}\left(\frac{\partial z}{\partial x}\right)\mathrm{d}x + \frac{\partial z}{\partial x}\mathrm{d}^2x + \mathrm{d}\left(\frac{\partial z}{\partial y}\right)\mathrm{d}y + \frac{\partial z}{\partial y}\mathrm{d}^2y \\ &= \left(\frac{\partial^2 z}{\partial x^2}\mathrm{d}x + \frac{\partial^2 z}{\partial x \partial y}\mathrm{d}y\right)\mathrm{d}x + \left(\frac{\partial^2 z}{\partial y \partial x}\mathrm{d}x + \frac{\partial^2 z}{\partial y^2}\mathrm{d}y\right)\mathrm{d}y \\ &= \frac{\partial^2 z}{\partial x^2}(\mathrm{d}x)^2 + 2\frac{\partial^2 z}{\partial x \partial y}\mathrm{d}x\mathrm{d}y + \frac{\partial^2 z}{\partial y^2}(\mathrm{d}y)^2, \end{split}$$

where  $(\mathrm{d}x)^2$  and  $(\mathrm{d}y)^2$  denote  $\mathrm{d}^2x$  and  $\mathrm{d}^2y$  respectively. If we treat  $\frac{\partial}{\partial x}$ ,  $\frac{\partial}{\partial y}$  as operators for partial differentiation and define

$$\left(\frac{\partial}{\partial x}\right)^2 = \frac{\partial^2}{\partial x^2}, \quad \left(\frac{\partial}{\partial y}\right)^2 = \frac{\partial^2}{\partial y^2}, \quad \left(\frac{\partial}{\partial x}\frac{\partial}{\partial y}\right) = \frac{\partial^2}{\partial x \partial y},$$

then the formulas for the first and second differentials can be written as

$$dz = \left(dx \frac{\partial}{\partial x} + dy \frac{\partial}{\partial y}\right) z,$$
$$d^2 z = \left(dx \frac{\partial}{\partial x} + dy \frac{\partial}{\partial y}\right)^2 z.$$

Similarly, we define

$$\left(\frac{\partial}{\partial x}\right)^p \left(\frac{\partial}{\partial y}\right)^q = \frac{\partial^{p+q}}{\partial x^p \partial y^q} = \frac{\partial^q}{\partial y^q} \left(\frac{\partial}{\partial x}\right)^p, \quad (p, q = 1, 2, \dots)$$

It is easy to use mathematical induction to prove the formula for higher-order differentials:

$$\mathrm{d}^k z = \left(\mathrm{d}x \frac{\partial}{\partial x} + \mathrm{d}y \frac{\partial}{\partial y}\right)^k z, \quad k = 1, 2, \cdots.$$

For an n-variable function  $u = f(x_1, x_2, \dots, x_n)$ , higher-order differentials can be similarly defined, and the following holds:

$$d^{k}u = \left(dx_{1}\frac{\partial}{\partial x_{1}} + dx_{2}\frac{\partial}{\partial x_{2}} + \dots + dx_{n}\frac{\partial}{\partial x_{n}}\right)^{k}u, \quad k = 1, 2, \dots$$

### 10.2 Differential of Vector-Valued Functions

Consider an n-dimensional vector-valued function defined on a domain  $U \subset \mathbb{R}^n$ :

$$f: U \to \mathbb{R}^m$$
,

$$\mathbf{x} \mapsto \mathbf{y} = f(\mathbf{x})$$

Expressed in coordinate vector form:

$$\mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix} = \begin{pmatrix} f_1(x_1, x_2, \dots, x_n) \\ f_2(x_1, x_2, \dots, x_n) \\ \vdots \\ f_m(x_1, x_2, \dots, x_n) \end{pmatrix}, \qquad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in U$$

1. If each component function  $f_i(x_1, x_2, \dots, x_n)$  ( $i = 1, 2, \dots, m$ ) is partially differentiable at  $\mathbf{x}^0$ , then the vector-valued function  $\mathbf{f}$  is differentiable at  $\mathbf{x}^0$ , and we define the matrix

$$\left(\frac{\partial f}{\partial x_j}(\mathbf{x}^0)\right)_{m \times n} = \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(\mathbf{x}^0) & \frac{\partial f_1}{\partial x_2}(\mathbf{x}^0) & \cdots & \frac{\partial f_1}{\partial x_n}(\mathbf{x}^0) \\ \frac{\partial f_2}{\partial x_1}(\mathbf{x}^0) & \frac{\partial f_2}{\partial x_2}(\mathbf{x}^0) & \cdots & \frac{\partial f_2}{\partial x_n}(\mathbf{x}^0) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1}(\mathbf{x}^0) & \frac{\partial f_m}{\partial x_2}(\mathbf{x}^0) & \cdots & \frac{\partial f_m}{\partial x_n}(\mathbf{x}^0) \end{pmatrix}$$

This matrix is called the Jacobian matrix of **f** at  $\mathbf{x}^0$ , denoted by  $f'(\mathbf{x}^0)$  (or  $Df(\mathbf{x}^0)$ ,  $J_f(\mathbf{x}^0)$ ).

For the special case m=1, i.e., n-variable scalar function  $z=f(x_1,x_2,\ldots,x_n)$ , the derivative at  $\mathbf{x}^0$  is

$$f'(\mathbf{x}^0) = \left(\frac{\partial f}{\partial x_1}(\mathbf{x}^0), \frac{\partial f}{\partial x_2}(\mathbf{x}^0), \cdots, \frac{\partial f}{\partial x_n}(\mathbf{x}^0)\right)$$

If the vector-valued function  $\mathbf{f}$  is differentiable at every point in U, then  $\mathbf{f}$  is said to be differentiable on U, and the corresponding relationship is

$$\mathbf{x} \in U \mapsto f'(\mathbf{x}) = J_f(\mathbf{x})$$

where  $f'(\mathbf{x})$  (or  $Df(\mathbf{x})$ ,  $J_f(\mathbf{x})$ ) denotes the derivative of  $\mathbf{f}$  at  $\mathbf{x}$  in U.

2. If every component function  $f_i(x_1, x_2, ..., x_n)$  (i = 1, 2, ..., m) of  $\mathbf{f}$  has continuous partial derivatives at  $\mathbf{x}^0$ , then every element of the Jacobian matrix of  $\mathbf{f}$  is continuous at  $\mathbf{x}^0$ . In this case,  $\mathbf{f}$  is said to have a continuous derivative at  $\mathbf{x}^0$  as a vector-valued function.

If the derivative of a vector-valued function f is continuous at every point in U, then f is said to have a continuous derivative on U.

3. If there exists an  $m \times n$  matrix A that depends only on  $\mathbf{x}^0$  (and not on  $\Delta \mathbf{x}$ ), such that in the neighborhood of  $\mathbf{x}^0$ ,

$$\Delta \mathbf{y} = f(\mathbf{x}^0 + \Delta \mathbf{x}) - f(\mathbf{x}^0) = A\Delta \mathbf{x} + o(\|\Delta \mathbf{x}\|)$$

(where  $\Delta \mathbf{x} = (\Delta x_1, \Delta x_2, \dots, \Delta x_n)^T$  is a column vector and  $\|\Delta \mathbf{x}\|$  denotes its norm), then f is said to be differentiable at  $\mathbf{x}^0$  as a vector-valued function, and  $A\Delta \mathbf{x}$  is called the differential of f at  $\mathbf{x}^0$ , denoted as  $d\mathbf{y}$ . If we denote  $\Delta \mathbf{x}$  by  $d\mathbf{x}$  ( $d\mathbf{x} = (dx_1, dx_2, \dots, dx_n)^T$ ), then

$$d\mathbf{v} = A d\mathbf{x}.$$

If the vector-valued function  $\mathbf{f}$  is differentiable at every point in U, then  $\mathbf{f}$  is said to be differentiable on U.

Combining the above three points, we obtain the following unified statement:

A vector-valued function f is continuous, differentiable, and has derivatives if and only if each of its coor-

dinate component functions  $f_i(x_1, x_2, \dots, x_n)$  ( $i = 1, 2, \dots, m$ ) is continuous, differentiable, and has derivatives.

## 10.3 Derivatives of Composite Mappings (Chain Rule)

Let  $U \subset \mathbb{R}^l$  and  $V \subset \mathbb{R}^n$  be open sets, and let

$$\mathbf{g}: U \to V$$
 and  $\mathbf{f}: V \to \mathbb{R}^m$ 

be mappings. If  $\mathbf{g}$  is derivative at  $\mathbf{u}^0 \in U$  and  $\mathbf{f}$  is differentiable at  $\mathbf{x}^0 = \mathbf{g}(\mathbf{u}^0)$ , then the composite mapping  $\mathbf{f} \circ \mathbf{g}$  is differentiable at  $\mathbf{u}^0$ , and:

$$J(\mathbf{f} \circ \mathbf{g})(\mathbf{u}^0) = J\mathbf{f}(\mathbf{x}^0)J\mathbf{g}(\mathbf{u}^0).$$



- 1. outer differentiable + inner derivative = total derivative
- 2. outer differentiable + inner differentiable = total differentiable

3.

Specially, define  $z=f(x,y), (x,y)\subset D_f\subset \mathbb{R}^2$ ,  $\mathbf{g}:D_g\to \mathbb{R}^2, (u,v)\mapsto (x(u,v),y(u,v))$ , and  $g(D_g)\subset D_f$ , then we have composite function

$$z = f \circ \mathbf{g} = f[x(u, v), y(u, v)], \quad (u, v) \in D_g.$$

$$\mathbb{R}^2 \xrightarrow{\mathbf{g}: \text{derivative}} \mathbb{R}^2 \xrightarrow{f: \text{differentiable}} \mathbb{R}$$

If g is derivative at  $(u_0, v_0) \in D_g$ , and f is differentiable at  $(x_0, y_0) = \mathbf{g}(u_0, v_0)$ , then  $z = f \circ \mathbf{g}$  is differentiable at  $(u_0, v_0)$ , and at the point,

$$\begin{bmatrix} \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} \end{bmatrix} = \begin{bmatrix} \frac{\partial z}{\partial x} & \frac{\partial z}{\partial y} \end{bmatrix} \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix}$$

# Proof

### 10.4 Mean Value Theorem and Taylor's Formula

### Definition 10.4 (Convex Region)

Let  $D \subseteq \mathbb{R}^n$  be a region. If every line segment connecting any two points  $\mathbf{x}_0, \mathbf{x}_1 \in D$  (denoted by  $\overline{\mathbf{x}_0}\overline{\mathbf{x}_1}$ ) is entirely contained in D, i.e., for any  $\lambda \in [0, 1]$ , we have

$$\mathbf{x}_0 + \lambda(\mathbf{x}_1 - \mathbf{x}_0) \in D,$$

then D is called a convex region.

### Theorem 10.3 (Lagrange's Mean Value Theorem)

Let f be <u>differentiable</u> on <u>a convex region</u>  $D \subseteq \mathbb{R}^n$ . For any two points  $\mathbf{a}, \mathbf{b} \in D$ , there exists a point  $\xi \in \overline{\mathbf{ab}}$  such that:

$$f(\mathbf{b}) - f(\mathbf{a}) = Jf(\xi)(\mathbf{b} - \mathbf{a}).$$

A Proof

### Theorem 10.4

Let D be a region in  $\mathbb{R}^n$ . If for any  $\mathbf{x} \in D$ , we have

$$Jf(\mathbf{x}) = 0,$$

then f is constant on D.

### $\Diamond$

### 🖋 Proof

### Theorem 10.5 (Taylor's Formula)

**Lagrange's Remainder** Let  $D \subseteq \mathbb{R}^n$  be a convex region, and let  $f: D \to \mathbb{R}$  have m+1 continuous partial derivatives. For  $\mathbf{x}^0 = (x_1^0, x_2^0, \dots, x_n^0) \in D$  and  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in D$ , there exists  $\xi \in \overline{\mathbf{x}^0 \mathbf{x}}$  such that:

$$f(\mathbf{x}) = f(\mathbf{x}^0) + \sum_{k=1}^{m} \frac{1}{k!} \left( \sum_{i=1}^{n} (x_i - x_i^0) \frac{\partial}{\partial x_i} \right)^k f(\mathbf{x}^0) + \frac{1}{(m+1)!} \left( \sum_{i=1}^{n} (x_i - x_i^0) \frac{\partial}{\partial x_i} \right)^{m+1} f(\xi)$$

**Peano's Remainder** Let  $D \subseteq \mathbb{R}^n$  be a convex region, and let  $f: D \to \mathbb{R}$  have m continuous partial derivatives. Then:

$$f(\mathbf{x}) = f(\mathbf{x}^0) + \sum_{k=1}^m \frac{1}{k!} \sum_{i_1, i_2, \dots, i_k = 1}^n \frac{\partial^k f}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} (\mathbf{x}^0) \prod_{j=1}^k (x_{i_j} - x_{i_j}^0) + R_m(\mathbf{x} - \mathbf{x}^0),$$

where 
$$R_m(\mathbf{x} - \mathbf{x}^0) = O(\|\mathbf{x} - \mathbf{x}^0\|^{m+1})$$
 or  $o(\|\mathbf{x} - \mathbf{x}^0\|^m)$ , as  $\|\mathbf{x} - \mathbf{x}^0\| \to 0$ .

In applications, particularly important is the expression of the first three terms in Taylor's formula, which is given as (let  $x_1 - x_1^0$  be denoted by  $\Delta x_1$ , and similarly for other variables;  $\Delta \mathbf{x} = (\Delta x_1, \Delta x_2, \dots, \Delta x_n)$ ):

$$f(\mathbf{x}) = f(\mathbf{x}^0) + Jf(\mathbf{x}^0)(\Delta \mathbf{x}) + \frac{1}{2!}(\Delta \mathbf{x})Hf(\mathbf{x}^0)(\Delta \mathbf{x})^{\mathrm{T}} + \cdots,$$

where the matrix

$$Hf(\mathbf{x}^{0}) = \begin{bmatrix} \frac{\partial^{2} f}{\partial x_{1}^{2}} & \frac{\partial^{2} f}{\partial x_{1} \partial x_{2}} & \cdots & \frac{\partial^{2} f}{\partial x_{1} \partial x_{n}} \\ \frac{\partial^{2} f}{\partial x_{2} \partial x_{1}} & \frac{\partial^{2} f}{\partial x_{2}^{2}} & \cdots & \frac{\partial^{2} f}{\partial x_{2} \partial x_{n}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^{2} f}{\partial x_{n} \partial x_{1}} & \frac{\partial^{2} f}{\partial x_{n} \partial x_{2}} & \cdots & \frac{\partial^{2} f}{\partial x_{n}^{2}} \end{bmatrix}_{\mathbf{x}^{0}}$$

is called the **Hessian matrix** of the function f. It is an  $n \times n$  symmetric matrix.

# **Chapter 11 Multiple Integrals**

## **Bibliography**

- [1] 徐森林, 薛春华. 数学分析. 第一版. 清华大学出版社, 2005.
- [2] 陈纪修,於崇华. 数学分析. 第三版. 高等教育出版社, 2019.
- [3] 常庚哲, 史济怀. 数学分析教程. 第三版. 中国科学技术大学出版社, 2012.
- [4] 裴礼文. 数学分析中的典型问题与方法. 第三版. 高等教育出版社, 2021.
- [5] 汪林. 数学分析中的问题与反例. 第一版. 高等教育出版社, 2015.
- [6] 谢惠民, 恽自求, 易法槐, 钱定边. 数学分析习题课讲义. 第二版. 高等教育出版社, 2019.
- [7] Walter Rudin. Principles of Mathematical Analysis. Third Edition. McGraw-Hill, 1976.
- [8] 菲赫金哥尔茨. 微积分学教程. 第八版. 高等教育出版社, 2006.