

novaflow

REFERENCE BOOK · NEURO-OMICS · EDITION 01 · OCTOBER 2026

The Figure Booklet

Plotting packages in neuroscience bulk, single-cell & spatial omics, 2023–26, ranked by use and rebuilt with code.

17,768
PAPERS MINED

98
PACKAGES

329
FIGURES RECREATED

BACKGROUND: 135 OF THE 329 RECREATED FIGURES, SORTED BY COLOUR



About this book

A field guide to the figures of modern neuro-omics. We counted which visualisation packages **17,768 open-access neuro-omics papers (2023–2026)** name in their METHODS sections, ranked them, and rebuilt **329 figure variations from 98 packages** on public neuro datasets. Each package gets a two-page spread with the plotting call that draws its signature figure.

What is in here

- **Part I · Mainstream workhorses**: 62 spreads. These are the recreated Top 100 packages ranked by METHODS mentions. Where several catalog entries share one recreation folder (e.g. IGV and UCSC Genome Browser → Gviz), they get one spread.
- **Part II · Niche & Cool**: 36 spreads. These are newer or specialised tools ranked by growth and visual distinctiveness. They are not in the Top 100 folders.
- **Visual index**: all 329 figures grouped into 11 figure types, each linked to its spread.
- **Appendix**: queued tools and why, plus the full Top 100 and Niche & Cool tables.

Ground rules

- Every number, DOI and figure link is copied from the source files: `figure_package_catalog.xlsx` (Top100, Niche_Cool, Recreations, Niche_Recreations, Method), `SUMMARY.md`, `METHODS.md` and `corpus_papers.csv`.
- Code excerpts are quoted verbatim from each `recreations/<pkg>/*_recreate.*` script, with line numbers.
- Galleries embed the original PNGs, not screenshots.
- All blue text is a link: DOIs, PMC figure pages, GitHub repositories and internal page references. The PDF also has bookmarks.

MAINSTREAM (TOP 100)

70/100

entries recreated (62 folders); 30 queued (Appendix A)

NICHE & COOL (73)

58/73

candidates recreated (55 folders, 19 shared with Top 100); 15 queued

VERIFICATION

98/98

scripts re-ran with exit 0 (SUMMARY.md); 261 of 268 example-figure URLs resolved

Reading paths

What do neuro papers actually use? · Top 30 chart, then Part I in rank order.

What is rising? · Niche growth chart, then Part II.

I know the look I want · Visual index: 329 thumbnails in 11 figure types.

Spatial & tissue figures · Squidpy, spatialLIBD, BayesSpace, Banksy, RCTD, Tangram, COMMOT…

Cell–cell communication · CellChat, LIANA+, NicheNet, MultiNicheNet, pyCirclize…

Brain atlas & neuroimaging · BrainGlobe, brainrender, nilearn, MNE.

What is missing? · Appendix A: 45 queued tools and the reason each is queued.

How far can I trust a rank? · Caveats on Methods page 2.

CONTENTS

Contents

Front matter

About this book 2
How to use this book 5
Methods & corpus 6
At a glance: rankings 8
Visual index by figure type 10

Visual index · 11 figure types

Heatmaps & dot matrices 10
Embeddings & feature maps 10
Trajectory, pseudotime & velocity 11
Spatial & tissue maps 11
Cell–cell communication 12
Enrichment & pathway activity 12
Networks, graphs & trees 13
Genome tracks & genomic signal 13
Brain atlas, 3D & neuroimaging 13
Composition, distributions & layouts 13
DE, QC & set overlaps 14

Part I · Mainstream workhorses 62 SPREADS · TOP-100 RANK

#1 DESeq2 (plotPCA/plotMA) R 18
#2 Seurat R 20
#3 clusterProfiler R 22
#4 ggplot2 R 24
#5 GSEA (Broad) DESKTOP 26
#6 limma R 28
#7 edgeR R 30
#9 pheatmap R 32
#10 WGCNA R 34
#11 CellChat R 36
#12 STRING WEB 38
#13 Enrichr WEB 40
#14 scanpy PYTHON 42
#17 ComplexHeatmap R 44
#18 g:Profiler WEB/R/PYTHON 46
#19 IGV DESKTOP/WEB 48
#20 fgsea R 50
#21 monocle3 R 52
#22 deepTools PYTHON/CLI 54
#23 SingleR R 56
#24 AUCell R 58
#27 EnhancedVolcano R 60
#29 survminer R 62
#31 Space Ranger/Loupe DESKTOP 64
#33 ggpubr R 66
#36 igraph R/PYTHON 68
#37 Signac R 70
#38 matplotlib PYTHON 72
#39 CopyKAT/inferCNV R 74
#40 seaborn PYTHON 76
#41 CellPhoneDB PYTHON 78
#42 slingshot R 80
#44 rrvgo R 82
#45 scVelo PYTHON 84
#46 VennDiagram R 86
#47 gplots (heatmap.2) R 88
#48 corrplot R 90
#50 hdWGCNA R 92
#52 clustree R 94
#54 circlize R 96
#57 UpSetR R 98
#59 RCTD/spacexr R 100
#60 CytoTRACE R/PYTHON 102
#61 decoupleR/decoupler R/PYTHON 104
#63 ABC atlas / Allen Brain Map WEB 106
#66 NicheNet R 108
#68 ReactomePA R 110
#70 pathview R 112
#71 GOplot R 114
#74 scCustomize R 116
#79 patchwork R 118
#83 Squidpy PYTHON 120
#84 plotly R/PYTHON/JS 122
#86 ggraph R 124
#87 Tangram PYTHON 126
#88 PCAtools R 128
#91 tradeSeq R 130
#92 LIANA R/PYTHON 132
#97 BayesSpace R 134
#98 CellRank PYTHON 136
#99 Nebulosa R 138
#100 ggalluvial R 140

Part II · Niche & Cool

36 SPREADS · NICHE RANK

#1 pyCirclize PYTHON 144
#3 brainrender PYTHON 146
#4 COMMOT PYTHON 148
#6 MultiNicheNet R 150
#7 moscot PYTHON 152
#8 scplotter R 154
#10 Banksy R/PYTHON 156
#12 scTour PYTHON 158
#13 plot1cell R 160
#14 veloVI PYTHON 162
#15 miloR / Milo R/PYTHON 164
#16 Datashader PYTHON 166
#17 scDRS PYTHON 168
#19 MNE-Python PYTHON 170
#20 sccomp R 172
#26 PyComplexHeatmap PYTHON 174
#29 nilearn PYTHON 176
#31 SCpubr R 178
#34 UCell R 180
#37 spatialLIBD R 182
#41 ggrepel volcano styles R 184
#42 Augur R 186
#47 ggridges R 188
#48 plotnine PYTHON 190
#50 Palantir PYTHON 192
#51 ggspavis/SpatialExperiment R 194
#55 plotgardener R 196
#60 propeller (speckle) R 198
#63 ggcoverage R 200
#64 ggtree/ggtreeExtra R 202
#65 Mellon PYTHON 204
#66 karyoploteR R 206
#67 marsilea PYTHON 208
#68 dittoSeq R 210
#70 ggforce R 212
#71 gggenes R 214

Appendix

A · Queued tools and why 216
B · Full Top 100 ranked table 218
C · Full Niche & Cool table (73) 221

ORIENTATION

How to use this book

Each package gets one **two-page spread**: the left page explains it, the right page shows it. Open the PDF in two-page view, with the cover shown separately, to see the spreads as facing pages. To start from a picture rather than a name, use the visual index.



1. **Name, language, category, figure types** · Chips show the language, the catalog category and the figure types drawn in the gallery. Colours match the visual index.
2. **Rank** · Part I: Top-100 rank by METHODS mentions. Part II: Niche & Cool rank (score = 0.5 × growth percentile + 0.5 × visual/5).
3. **Numbers strip** · Part I: METHODS paper count, % of the 17,768-paper corpus, full-text mentions, role, variations. Part II: growth ratio, visual score, niche score, neuro-corpus mentions, GitHub stars. The small bar track shows where the package sits among its peers.
4. **Best for** · The catalog's *typical_figure_types* (Part I) or *signature_visual* (Part II), quoted verbatim.
5. **Key plotting call** · A verbatim excerpt of the recreation script, ending at the line that writes the first variation (or the signature call). File and line number are given.
6. **Seen in the literature** · Up to three top-scored example figures per Top-100 package, with clickable DOI and PMC figure links and the evidence sentence. Niche tools fall back to the most-cited corpus papers that mention them.
7. **Limitations & notes** · Catalog notes plus caveats from SUMMARY, METHODS and the script itself: simulated or non-neuro data, shims, look-alikes, dropped plot types.
8. **Gallery** · Every PNG in the folder at its native resolution, laid out to fill the page without upscaling past about 118 dpi.

Figure-type key

Every variation is assigned one figure type, following the requested taxonomy. Two extra types, **Composition, distributions & layouts** and **DE, QC & set overlaps**, hold figures that fit none of the others. Counts are figures, not packages.

Heatmaps & dot matrices	40	Embeddings & feature maps	26
Trajectory, pseudotime & velocity	37	Spatial & tissue maps	34
Cell–cell communication	23	Enrichment & pathway activity	32
Networks, graphs & trees	20	Genome tracks & genomic signal	19
Brain atlas, 3D & neuroimaging	13	Composition, distributions & layouts	55
DE, QC & set overlaps	30		

READ THE NUMBERS CAREFULLY

A METHODS mention ≠ the package drew a published figure. DESeq2, limma and edgeR rank high because they are cited for statistics. Growth ratios for tools first released in 2023–24 are inflated by a near-zero 2023 base, and counts under ~10 are noisy.

Methods & corpus

A systematic Europe PMC search of open-access full text, counting mentions inside the **METHODS section only**. That section is where authors name the software they used.

1 · CORPUS	2 · SCREEN	3 · RANK	4 · RECREATE	5 · VERIFY
17,768	**212**	**100 + 73**	**98**	**329**
OA neuro-omics papers with full text, 2023–26	candidate packages, each with its own search terms	Top 100 by METHODS mentions; Niche & Cool by growth + visual	folders: 70 Top-100 entries, 58 niche tools (19 shared)	PNG variations; all 98 scripts exit 0; 261/268 figure links OK

CORPUS SIZE

17,768

open-access neuro-omics papers with full text (SRC:MED), published 2023–2026



*2026 is a partial year (to mid-October).

MODALITY TAGS (TERM IN ABSTRACT OR METHODS; A PAPER CAN CARRY SEVERAL)

Single-cell or multiome	6,887
Spatial	1,376
Any RNA-seq (bulk included)	16,397

TOP 10 JOURNALS IN THE CORPUS (CORPUS_PAPERS.CSV)

Nat Commun	1,167
Sci Rep	545
Int J Mol Sci	539
Adv Sci (Weinh)	433
J Neuroinflammation	405
iScience	396
Sci Adv	341
Front Immunol	280
Proc Natl Acad Sci U S A	268
Elife	263

SEARCH (EUROPE PMC REST /WEBSERVICES/REST/SEARCH)

Neuro terms in the title or abstract, **AND** omics terms in the abstract or METHODS, **AND** `PUB_YEAR:[2023 TO 2026] OPEN_ACCESS:Y HAS_FT:Y SRC:MED`. Full query:

```
(TITLE:(neuron OR neurons OR neuronal OR brain OR cortex OR cortical OR hippocampus OR hippocampal OR neurodevelopment OR neurodevelopmental OR "neural progenitor" OR "neural stem" OR neurodegeneration OR neurodegenerative OR cerebellum OR "spinal cord" OR retina OR glia OR glial OR astrocyte OR astrocytes OR microglia OR oligodendrocyte OR "cerebral organoid" OR "brain organoid" OR Alzheimer OR Parkinson OR "nervous system") OR ABSTRACT:(neuron OR neurons OR neuronal OR brain OR cortex OR cortical OR hippocampus OR hippocampal OR neurodevelopment OR neurodevelopmental OR "neural progenitor" OR "neural stem" OR neurodegeneration OR neurodegenerative OR cerebellum OR "spinal cord" OR retina OR glia OR glial OR astrocyte OR astrocytes OR microglia OR oligodendrocyte OR "cerebral organoid" OR "brain organoid" OR Alzheimer OR Parkinson OR "nervous system")) AND (ABSTRACT:("single-cell RNA" OR "single-nucleus RNA" OR "single-cell sequencing" OR "scRNA-seq" OR "snRNA-seq" OR "RNA-seq" OR "RNA sequencing" OR "spatial transcriptomics" OR Visium OR MERFISH OR Xenium OR CosMx OR "Slide-seq" OR "Stereo-seq" OR CODEX OR "snATAC-seq" OR "scATAC-seq" OR multiome) OR METHODS:("single-cell RNA" OR "single-nucleus RNA" OR "single-cell sequencing" OR "scRNA-seq" OR "snRNA-seq" OR "RNA-seq" OR "RNA sequencing" OR "spatial transcriptomics" OR Visium OR MERFISH OR Xenium OR CosMx OR "Slide-seq" OR "Stereo-seq" OR CODEX OR "snATAC-seq" OR "scATAC-seq" OR multiome)) AND PUB_YEAR:[2023 TO 2026] AND OPEN_ACCESS:Y AND HAS_FT:Y AND SRC:MED
```

Screening & counting

1. **212 candidate packages** are screened (`candidates.py`). Each has its own search phrases, and ambiguous names get context clauses. For example, *patchwork* must appear with ggplot2, "R package" or CRAN; *MAST* needs hurdle wording; *PROGENy* needs pathway-activity wording.
2. **paper_count** is the number of corpus papers whose METHODS section mentions the package. **fulltext_any_count** counts the same terms anywhere in the paper and is given for comparison.
3. Only **figure-making tools** are ranked. 26 analysis-only, QC and data packages (Harmony, DoubletFinder, GSVA, DAVID, anndata…) go to the *Excluded* sheet, and general GUI tools (Prism, ImageJ, Illustrator, BioRender) to *GUI_tools*.
4. **Example figures.** For each Top-100 package, up to 10 of the most-cited papers mentioning it are parsed (fullTextXML). Figures are scored +6 if the caption names the package, +3 if a citing paragraph names it, +1–3 for figure-type keywords and −2 if supplementary. Up to 3 are kept, and each PMC figure URL is checked in headless Chrome: **261 of 268 resolved**.
5. **Niche & Cool** covers 73 newer or specialised tools (`niche_candidates.py`). The growth ratio is (broad mentions per 1k papers in 2025–26 + 0.01) ÷ (per 1k in 2023 + 0.01), measured in a broad OA omics corpus of 28,714 / 33,012 / 42,965 / 35,775 papers. Score = 0.5 × percentile(growth) + 0.5 × visual/5.

Recreations

There is one folder per package (`recreations/<pkg>/`), holding one script and 2–7 `vN_<variation>.png` outputs. Each script runs on its own with Rscript, python3 or bash, and `run_all.sh` reruns everything, starting with LIANA_plus because its tables feed pyCirclize, CellPhoneDB_ktplots and circlize_chord. Shared data prep is in `recreations/_data/prep_data.R`.

Three recreations call **live web APIs**: Enrichr (GSEApy), g:Profiler (gprofiler2, HTTPS) and STRING. **deepTools uses simulated bigWigs.** survminer uses `coin::glioma` (Grana et al. 2002).

CAVEATS (METHODS §6, §7)

1. **Open-access bias.** Only papers with Europe PMC full text are included, so closed-access Cell, Neuron and Nature Neuroscience papers are missing unless an OA or author-manuscript copy exists.
2. **A mention doesn't prove figure use.** DESeq2, limma and edgeR rank high because they are cited for statistics. The role column and the evidence sentences help tell the two apart.
3. **Under-counting.** Tools named only in Key Resources tables or Code Availability sections, or never named at all (matplotlib, ggplot2), are under-counted. Compare with fulltext_any_count.
4. **Name ambiguity** is handled with context terms, but imperfectly: Harmony, Giotto and CARD are common words or names.
5. **Topic noise.** The neuro filter works on title or abstract, so a few off-topic papers are included.
6. **Not yet recreated:** web, GUI and commercial tools (Cytoscape, Metascape, IPA, QuPath…) and database-heavy tools (SCENIC). See Appendix A.
7. **Niche growth.** Tools first released in 2023–24 have inflated ratios (near-zero 2023 base). Counts under ~10 are noisy, the visual score is curator judgement, and 2026 is partial.

Real neuro datasets behind the recreations (METHODS §8)

Zeisel 2015 · mouse S1 cortex / CA1 scRNA-seq	3,005 cells (scRNAseq::ZeiselBrainData). Most single-cell figures; Fluidigm C1 chip ID as replicate for propeller, miloR, sccomp and Augur.
La Manno 2018 · dentate gyrus P12/P35	scVelo, CellRank, Palantir, Mellon, veloVI, scTour, moscot (P12→P35 optimal transport), CytoTRACE 2
10x Visium · mouse coronal brain	Squidpy, Tangram, COMMOT, Loupe-style
Moffitt 2018 · MERFISH hypothalamus	Squidpy, Datashader, plotly 3D
Maynard 2021 · human DLPFC 151673	spatialLIBD, ggspavis, BayesSpace, Banksy (STexampleData)
Allen CCFv3 25 µm (BrainGlobe)	brainrender, atlas overlays
NeuroVault / Harvard-Oxford · PhysioNet EEGBCI S001	nilearn · MNE-Python
Tirosh 2016 oligodendroglioma · scDRS MAGMA sets	inferCNV · scDRS (SCZ, BIP, MDD, AD, height control)
Bulk & other	airway (DESeq2, dex vs untreated) · coin::glioma (survminer) · MSigDB Hallmark · NicheNet v2 mouse priors (Zenodo 7074291)

Environment & shims (METHODS §9–10)

- R packages live in /home/box/R/lib and Python packages in ~/.local. The system R and dev libraries were reinstalled with apt after a box reboot.
- NumPy 2 shims: `np.Inf` (COMMOT) and `np.float_` (scDRS). `matplotlib.cm.get_cmap` is restored for PyComplexHeatmap, and scTour gets a dense X.
- brainrender's final frame is drawn without depth peeling, the moscot Sankey is re-drawn from the OT coupling, and BayesSpace's enhanced genes come from per-gene PC regression.
- miloR is patched from source, sccomp uses cmdstanr + CmdStan 2.40, and multinichenetr is installed from GitHub.
- dynamo-release was tried and then removed, because it downgrades anndata below 0.11 and breaks scanpy/cellrank.

Top 30 by METHODS mentions

Number of the 17,768 corpus papers that name each package in their **METHODS** section, with the share of the corpus. Bars link to spreads; hatched bars are queued (Appendix A).



DESeq2, limma and edgeR are mostly cited for statistics, not figures (caveat 2). IGV (#19) and UCSC Genome Browser (#28) are recreated as Gviz look-alikes.

Niche & Cool: top 20 by growth

The 20 fastest-growing of the 73 Niche & Cool candidates. Growth ratio is the per-1k mention rate in 2025–26 ÷ 2023 in the broad OA omics corpus (log scale). Dots show the curator visual score (1–5), and the left number is the niche rank.

WHY CYTOTRACE2 SHOWS 324.85×

Tools first released in 2023–24 have a near-zero 2023 base, which inflates the ratio. That is why the official niche ranking blends growth percentile with visual score. The niche top 5 by that score are pyCirclize, hdWGCNA, brainrender, COMMOT and CytoTRACE2 (SUMMARY.md).

Official Niche & Cool top 20 · by score (SUMMARY.md)

#	TOOL	GROWTH	VISUAL	NEURO MENTIONS 2023–26	GITHUB STARS	RECREATED	PAGE
1	**pyCirclize**	20.05	5	0	1127	done (3)	144
2	**hdWGCNA**	6.66	5	147	494	done (5)	92
3	**brainrender**	4.81	5	4	680	done (3)	146
4	**COMMOT**	4.67	5	15	149	done (3)	148
5	**CytoTRACE2**	324.85	4	24	269	done (3)	102
6	**MultiNicheNet**	89.9	4	13	202	done (3)	150
7	**moscot**	42.91	4	7	217	done (3)	152
8	**scplotter**	17.51	4	1	333	done (4)	154
9	SCENIC+	14.32	4	32	267	queued (0)	
10	**Banksy**	14.16	4	21	167	done (2)	156
11	**ggsankey**	10.71	4	8	338	done (3)	140
12	**scTour**	6.52	4	7	89	done (3)	158
13	**plot1cell**	2.7	5	6	131	done (3)	160
14	**veloVI**	5.87	4	9	45	done (3)	162
15	**miloR / Milo**	2.31	5	46	450	done (3)	164
16	**Datashader**	4.81	4	1	3568	done (3)	166
17	**scDRS**	4.63	4	37	201	done (3)	168
18	**BrainGlobe atlas API**	2.27	5	1	189	done (3)	106
19	**MNE-Python**	2.27	5	1	3549	done (3)	170
20	**sccomp**	4.27	4	10	134	done (3)	172

VISUAL INDEX

Find a figure by its look

All **329 recreated figures**, grouped by figure type and sorted by package. Click any thumbnail to jump to its spread.

Heatmaps & dot matrices

40 FIGURES · 21 PACKAGES



AUCell
v3_auc_heatmap



ComplexHeatmap
v1_annotated_split



ComplexHeatmap
v2_singlecell_markers



ComplexHeatmap
v3_correlation_cellfun



ComplexHeatmap
v4_densityHeatmap



CopyKAT/inferCNV
v1_cnv_heatmap_by_grou



CopyKAT/inferCNV
v2_cnv_heatmap_subclus

CopyKAT
v3_cnv_



corrplot
v1_corrplot_celltypes_



corrplot
v2_corrplot_genes_mixe



corrplot
v3_corrplot_shade_hclu



dittoSeq
v5_dittoHeatmap



gplots (heatmap.2)
v1_heatmap2_rowz



gplots (heatmap.2)
v2_heatmap2_celltype_c



hdWGCNA
v4_module_dotplot



marsile
v1_heat



marsilea
v2_sized_dot_heatmap



marsilea
v3_stackbar



PCAtools
v3_eigencorplot



pheatmap
v1_topvar_zscore



pheatmap
v2_sample_distance



pheatmap
v3_cutree



pheatmap
v4_sc_markers

plot1ce
v3_comp



PyComplexHeatmap
v1_split_annotated

PyComplexHeatmap
v2_dot_clustermap



scanpy
v2_dotplot



scanpy
v3_matrixplot



scanpy
v5_rank_genes_dotplot



scanpy
v6_tracksplot



scCustomize
v3_Clustered_DotPlot

scDRS
v2_grou



SCpubr
v3_do_DotPlot



seaborn
v1_clustermap



Seurat
v4_DotPlot



Seurat
v5_DoHeatmap



SingleR
v1_plotScoreHeatmap



SingleR
v3_confusion



UCell
v3_UCell_dotplot

WGCNA
v3_modu

Embeddings & feature maps

26 FIGURES · 20 PACKAGES



AUCell
v2_umap_auc

Augur
v2_umap_auc



clustree
v4_clustree_overlay



dittoSeq
v1_dittoDimPlot_letter



dittoSeq
v4_dittoDimHex



ggforce
v1_mark_hull

ggplot2
v2_umap



hdWGCNA
v3_ModuleFeaturePlot



matplotlib
v3_umap



Nebulosa
v1_density_single



Nebulosa
v2_joint_density



Nebulosa
v3_density_vs_featurep

plot1cell
v1_circlize_umap



plotly
v1_umap3d

plotnine
v1_umap



scanpy
v1_umap_grid



scCustomize
v1_DimPlot_scCustom_fi



scCustomize
v2_FeaturePlot_scCusto



scDRS
v1_umap_scores



scplotter
v1_celldimplot

SCpubr
v1_do_DimPlot



SCpubr
v2_do_NebulosaPlot



Seurat
v1_DimPlot



Seurat
v2_FeaturePlot



SingleR
v4_umap_labels



UCell
v1_UCell_featureplots

Trajectory, pseudotime & velocity

37 FIGURES · 12 PACKAGES



CellRank
v1_macrostates



CellRank
v2_fate_probabilities



CellRank
v3_circular_projection



CellRank
v4_lineage_drivers



CytoTRACE
v1_potency_umap



Mellon
v1_density_umap



Mellon
v2_density_by_cluster

Mellon
v3_densi



monocle3
v1_trajectory_subclass



monocle3
v2_pseudotime



monocle3
v3_genes_in_pseudotime



monocle3
v4_genes_on_trajectory



moscot
v1_cell_transition



moscot
v2_sankey

moscot
v3_push_umap



Palantir
v1_palar



Palantir
v2_branch_selection



Palantir
v3_gene_trends



Palantir
v4_trajectory_arrows



scanpy
v7_paga

scTour
v1_pseudotime_umap



scTour
v2_vector_field



scTour
v3_ptime_by_celltype

scVelo
v1_stred



scVelo
v2_grid_arrows



scVelo
v3_phase_portraits



scVelo
v4_latent_time_heatmap



scVelo
v5_latent_time



slingshot
v1_curves_on_umap



slingshot
v2_pseudotime_per_line



slingshot
v3_cluster_MST



tradeSeq
v1_plotS



tradeSeq
v2_dynamic_gene_heatma



tradeSeq
v3_plotGeneCount



veloVI
v1_velovi_stream



veloVI
v2_uncertainty



veloVI
v3_coherence_by_cluste

Spatial & tissue maps

34 FIGURES · 11 PACKAGES



Banksy
v1_banksy_vs_nonspatia



Banksy
v2_banksy_umap



BayesSpace
v1_clusterPlot_vs_trut



BayesSpace
v2_enhanced_subspot



BayesSpace
v3_enhanced_genes

COMMOT
v1_flow_grid



COMMOT
v2_flow_stream



COMMOT
v3_sender_maps

Datashader
v1_density_fire



Datashader
v2_categorical_shading

Datashader
v3_all_slices_strip



ggspavis/SpatialEx
v1_plotSpots_layers



ggspavis/SpatialEx
v2_plotVisium_HE



ggspavis/SpatialEx
v3_continuous_qc_gene



plotly
v2_merfish3d

RCTD/spacexr
v1_weight_maps

RCTD/spacexr
v2_dominant_type

RCTD/spacexr
v3_scatterpie_zoom



Space Ranger/Loupe
v1_tissue_clusters_uma



Space Ranger/Loupe
v2_feature_opacity



Space Ranger/Loupe
v3_umi_per_spot



spatialLIBD
v1_vis_clus_layers



spatialLIBD
v2_vis_gene_MOBP



spatialLIBD
v3_vis_grid_gene



spatialLIBD
v4_layer_boxplots



Squidpy
v1_clusters_on_HE



Squidpy
v2_genes_on_HE



Squidpy
v3_nhood_enrichment



Squidpy
v4_merfish_cell_classe



Squidpy
v5_co_occurrence



Squidpy
v6_ripley_L

Tangram
v1_celltype_proj



Tangram
v2_argmax_vs_clusters



Tangram
v3_training_scores

Cell–cell communication

23 FIGURES · 8 PACKAGES

CellChat
v1_circle_counts

CellChat
v2_heatmap



CellChat
v3_bubble



CellChat
v4_signalingRole_scatt

CellChat
v5_chord_pathway



CellPhoneDB
v1_cpdb_heatmap



CellPhoneDB
v2_cpdb_dotplot



circlize
v1_chord_direction



circlize
v2_circos_genome_rings

LIANA
v1_dotplot

LIANA
v2_circle_plot



LIANA
v3_interaction_count_h

MultiNicheNet
v1_circos_group_compar

MultiNicheNet
v2_sample_lr_activity



MultiNicheNet
v3_ligand_activity_til



NicheNet
v1_ligand_activity



NicheNet
v2_ligand_receptor_hea

NicheNet
v3_circos

NicheNet
v4_ligand_target_heatm

pyCirclize
v1_chord_interactions



pyCirclize
v2_ligand_receptor_cho

pyCirclize
v3_circos_tracks



SCpubr
v4_do_ChordDiagram

Enrichment & pathway activity

32 FIGURES · 10 PACKAGES



clusterProfiler
v1_dotplot



clusterProfiler
v2_barplot

clusterProfiler
v3_cnetplot



clusterProfiler
v4_emapplot



clusterProfiler
v5_gseaplot2



clusterProfiler
v6_ridgeplot



decoupleR/decouple
v1_progeny_heatmap



decoupleR/decouple
v2_collectri_tf_heatma



decoupleR/decouple
v3_pathway_umaps

Enrichr
v1_barplot_grouped



Enrichr
v2_dotplot_by_library



fgsea
v1_plotEnrichment

fgsea
v2_plotGseaTable

fgsea
v3_NES_barplot

g:Profiler
v1_manhattan



g:Profiler
v2_publish_highlight



g:Profiler
v3_gosttable



GOplot
v1_GOBubble

GOplot
v2_GOCircle



GOplot
v3_GOChord



GSEA (Broad)
v1_classic_enrichment



GSEA (Broad)
v2_multi_term



GSEA (Broad)
v3_dotplot



pathview
v1_GABAergic_synapse_n



pathview
v2_glutamatergic_synap



pathview
v3_GABAergic_graphviz



ReactomePA
v1_compareCluster_dotp



ReactomePA
v2_emapplot

ReactomePA
v3_cnetplot

rrvgo
v1_treemap



rrvgo
v2_scatter



rrvgo
v3_heatmap

Networks, graphs & trees

20 FIGURES · 7 PACKAGES



clustree
v1_clustree_tree



clustree
v2_clustree_sc3stabili



clustree
v3_clustree_gene_Mbp

ggraph
v1_coexpression_networ

ggraph
v2_circular_dendrogram

ggraph
v3_circlepack



ggtree/ggtreeExtra
v1_circular_tree



ggtree/ggtreeExtra
v2_fan_heatmap_bar_



ggtree/ggtreeExtra
v3_rect_tree_clade_hig



hdWGCNA
v1_soft_powers



hdWGCNA
v2_dendrogram



hdWGCNA
v5_ModuleUMAPPlot

igraph
v1_fr_communities



igraph
v2_mark_groups



igraph
v3_circle_hubs



STRING
v1_string_api_networ



STRING
v2_networkx_fc



WGCNA
v1_soft_threshold



WGCNA
v2_dendro_modules



WGCNA
v4_eigengene_network

Genome tracks & genomic signal

19 FIGURES · 7 PACKAGES



deepTools
v1_plotHeatmap



deepTools
v2_plotProfile_perGrou

deepTools
v3_kmeans3



deepTools
v4_scale_regions_profi



ggcoverage
v1_coverage_gene



ggcoverage
v2_samples_transcripts



gggenes
v1_gene_arrows



gggenes
v2_pcdhg_cluster



IGV
v1_igv_style_tracks

IGV
v2_zoom_gene_model

IGV
v3_ucsc_style_dense

karyoploteR
v1_karyotype_density



karyoploteR
v2_lfc_scatter_ideogra

karyoploteR
v3_markers_chr



plotgardener
v1_page_layout

plotgardener
v2_gene_and_ideogram



Signac
v1_coverage_plot



Signac
v2_tile_plot



Signac
v3_fraghist_umap

Brain atlas, 3D & neuroimaging

13 FIGURES · 4 PACKAGES



ABC atlas / Allen
v1_coronal_overlay



ABC atlas / Allen
v2_region_outlines

ABC atlas / Allen
v3_ccf_3d_regions



brainrender
v1_regions_3quarter



brainrender
v2_hippocampus_points



brainrender
v3_coronal_cut



MNE-Python
v1_raw_browser



MNE-Python
v2_erds_tfr



MNE-Python
v3_band_topomaps



nilearn
v1_glass_brain



nilearn
v2_stat_map_cuts



nilearn
v3_surface_stat_map



nilearn
v4_atlas_roi

Composition, distributions & layouts

55 FIGURES · 26 PACKAGES



AUCell
v1_exploreThresholds



Augur
v1_lollipop

Augur
v3_feature_importance



CytoTRACE
v2_potency_composition



CytoTRACE
v3_score_ranking



dittoSeq
v2_dittoBarPlot



dittoSeq
v3_dittoPlot_ridgejitt

ggalluvial
v1_ggsankey

ggalluvial
v2_ggsankey_alluvial



ggalluvial
v3_ggalluvial



ggforce
v2_facet_zoom



ggforce
v3_parallel_sets



ggplot2
v3_composition_bar



ggplot2
v4_violin_box



ggpubr
v1_boxplot_pvalues



ggpubr
v2_violin



ggpubr
v3_scatter_cor



ggpubr
v4_barplot_mean_se



ggridges
v1_ridges_gradient



ggridges
v2_ridges_points



ggridges
v3_hdr_umap

ggridges
v4_hdr_lines



matplotlib
v4_multipanel_gridspec



miloR / Milo
v1_nhood_graph_DA



miloR / Milo
v2_DA_beeswarm



miloR / Milo
v3_nhood_size_hist



patchwork
v1_patchwork_composite



patchwork
v2_patchwork_design_in



patchwork
v3_cowplot_grid_waterm



plot1cell
v2_cell_fraction



plotnine
v2_faceted_violins

plotnine
v3_bin2d_facets



propeller (speckle
v1_plotCellTypeProps



propeller (speckle
v2_props_by_group



propeller (speckle
v3_composition_volcano



scanpy
v4_stacked_violin



sccomp
v1_posterior_boxplot



sccomp
v2_1D_intervals



sccomp
v3_2D_mean_variability



scCustomize
v4_Stacked_VlnPlot



scDRS
v3_score_boxplots



scplotter
v2_cellstat_circos

scplotter
v3_cellstat_radar_ring



scplotter
v4_featurestat_ridge



SCpubr
v5_do_BoxPlot



seaborn
v2_violin



seaborn
v3_catplot_facets



seaborn
v4_kde_qc



Seurat
v3_VlnPlot



Seurat
v6_RidgePlot



SingleR
v2_plotDeltaDistributi



survminer
v1_km_risktable



survminer
v2_cumhaz



survminer
v3_forest



UCell
v2_UCell_violins

DE, QC & set overlaps

30 FIGURES · 10 PACKAGES



DESeq2 (plotPCA/pl
v1_plotPCA



DESeq2 (plotPCA/pl
v2_plotMA_shrunken



DESeq2 (plotPCA/pl
v3_plotDispEsts



DESeq2 (plotPCA/pl
v4_plotCounts



edgeR
v1_plotBCV



edgeR
v2_plotMDS



edgeR
v3_plotQLDisp



edgeR
v4_plotSmear



EnhancedVolcano
v1_default



EnhancedVolcano
v2_selected_boxed



EnhancedVolcano
v3_custom_colours



ggplot2
v1_volcano



ggrepel volcano st
v1_gradient_labels



ggrepel volcano st
v2_side_aligned



ggrepel volcano st
v3_quadrants_counts



limma
v1_voom_meanvar



limma
v2_plotMDS



limma
v3_plotMD



limma
v4_volcanoplot



matplotlib
v1_volcano



matplotlib
v2_MA



PCAtools
v1_screeplot



PCAtools
v2_biplot



PCAtools
v4_pairsplot



UpSetR
v1_upset_frequency



UpSetR
v2_upset_degree_querie



UpSetR
v3_complexheatmap_UpSe

VennDiagram
v1_VennDiagram_4set



VennDiagram
v2_ggvenn_3set



VennDiagram
v3_ggVennDiagram_gradi



PART I · 62 PACKAGES · 217 RECREATED FIGURES

novaflow

Mainstream workhorses

The packages neuro-omics papers name most often in their METHODS sections, in Top-100 rank order. They cover bulk DE diagnostics, single-cell embeddings, enrichment, communication, trajectories and spatial maps. **70 of the Top 100 entries are recreated in 62 folders**; the other 30 are queued (Appendix A).

62
spreads

217
figures

#1–#100
rank span

5,668
papers for #1 (DESeq2)

IN THIS PART · RANK · PAGE

1 DESeq2 (plotPCA/plotMA) 18
2 Seurat 20
3 clusterProfiler 22
4 ggplot2 24
5 GSEA (Broad) 26
6 limma 28
7 edgeR 30
9 pheatmap 32
10 WGCNA 34
11 CellChat 36
12 STRING 38
13 Enrichr 40
14 scanpy 42
17 ComplexHeatmap 44
18 g:Profiler 46
19 IGV 48
20 fgsea 50
21 monocle3 52
22 deepTools 54
23 SingleR 56
24 AUCell 58
27 EnhancedVolcano 60
29 survminer 62
31 Space Ranger/Loupe 64
33 ggpubr 66
36 igraph 68
37 Signac 70
38 matplotlib 72
39 CopyKAT/inferCNV 74
40 seaborn 76
41 CellPhoneDB 78
42 slingshot 80
44 rrvgo 82
45 scVelo 84
46 VennDiagram 86
47 gplots (heatmap.2) 88
48 corrplot 90
50 hdWGCNA 92
52 clustree 94
54 circlize 96
57 UpSetR 98
59 RCTD/spacexr 100
60 CytoTRACE 102
61 decoupleR/decoupler 104
63 ABC atlas / Allen Brain 106
66 NicheNet 108
68 ReactomePA 110
70 pathview 112
71 GOplot 114
74 scCustomize 116
79 patchwork 118
83 Squidpy 120
84 plotly 122
86 ggraph 124
87 Tangram 126
88 PCAtools 128
91 tradeSeq 130
92 LIANA 132
97 BayesSpace 134
98 CellRank 136
99 Nebulosa 138
100 ggalluvial 140

DESeq2 (plotPCA/ plotMA)

#1
OF 100 · METHODS MENTIONS

R · BULK QC/DE · DE, QC & SET OVERLAPS

5,668	31.9%	5,871	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

PCA of samples (plotPCA), MA plot, dispersion plot, per-gene counts; results feed volcano/heatmaps

THIS RECREATION

DESeq2 built-in diagnostic plots (canonical bulk RNA-seq QC/DE figures) Data: airway (dexamethasone vs untreated); see ../_data/prep_data.R

KEY PLOTTING CALL

RECREATIONS/DESEQ2/DESEQ2_RECREATE.R — EXCERPT · ENDS LINE 11

```
x <- readRDS("../_data/airway_deseq2.rds"); dds <- x$dds; res <- x$res
vsd <- vst(dds, blind = FALSE)
# V1: plotPCA (the most-reused DESeq2 figure)
p <- plotPCA(vsd, intgroup = c("dex","cell"), returnData = TRUE); pv <- round(100*attr(p,"percentVar"))
g <- ggplot(p, aes(PC1, PC2, color = dex, shape = cell)) + geom_point(size = 4) +
  xlab(paste0("PC1: ", pv[1], "% variance")) + ylab(paste0("PC2: ", pv[2], "% variance")) + coord_fixed() + theme_bw() +
ggtitle("DESeq2 plotPCA (vst)")
ggsave("v1_plotPCA.png", g, width = 6, height = 4.5, dpi = 150)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 FIGURE CAPTION
GDF15 promotes weight loss by enhancing energy expenditure in muscle. *Nature, 2023*
"...P values were calculated using two-sided unpaired Mann–Whitney U-tests. i, PCA of liver samples from vehicle-treated and GDF15-treated (5 nmol per kg) mice and pair-fed controls using VST data from DESeq2. n = ..."
DOI ↗ Fig. 1 on PMC ↗

Fig. 3 FIGURE CAPTION
APOE4/4 is linked to damaging lipid droplets in Alzheimer's disease microglia. *Nature, 2024*
"...st; per condition, mean ± s.e.m). d, Normalized gene expression counts for significant differentially expressed genes in APOE4/4 iMG ± fAβ (n = 3 replicate wells per condition, P values determined by DEseq2 ..."
DOI ↗ Fig. 3 on PMC ↗

Fig. 2 FIGURE CAPTION
Microglia ferroptosis is regulated by SEC24B and contributes to neurodegeneration. *Nat Neurosci, 2023*
"...s and astrocytes in iron + RSL3 versus vehicle. g, Violin plots for FTH1 expression in microglia (red), neurons (green) and astrocytes (yellow) under vehicle or iron + RSL3 conditions. Wald test with DESeq2 ..."
DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES
No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/DESeq2/DESeq2_recreate.R`

√ verified run · exit 0 · 16 s

DESeq2 (plotPCA/plotMA) · 4 variations



v1 plotPCA · DE



v2 plotMA_shrunken · DE



v3 plotDispEsts · DE



v4 plotCounts · DE

Seurat

#2

OF 100 · METHODS MENTIONS

R · SINGLE-CELL (EMBEDDING/DOT/VIOLIN/FEATURE) · EMBEDDINGS & FEATURE MAPS · COMPOSITION, DISTRIBUTIONS & LAYOUTS · HEATMAPS & DOT MATRICES

4,671	26.29%	4,846	analysis+plotting	6
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

UMAP/t-SNE DimPlot, FeaturePlot, VlnPlot, DotPlot, DoHeatmap, RidgePlot, spatial feature plots

THIS RECREATION

Seurat plotting functions: DimPlot, FeaturePlot, VlnPlot, DotPlot, DoHeatmap, RidgePlot

KEY PLOTTING CALL

RECREATIONS/SEURAT/SEURAT_RECREATE.R — EXCERPT · ENDS LINE 5

```
so <- readRDS("../_data/zeisel_seurat.rds")
ggsave("v1_DimPlot.png", DimPlot(so, label = TRUE, repel = TRUE) + NoLegend() + ggtitle("DimPlot: Zeisel cortex/hippocampus"), w
idth = 6, height = 5.5, dpi = 150, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 — FIGURE CAPTION

Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST. *Nat Commun, 2023*

"…ote the upper and lower quartiles, and whiskers denote the 1.5× interquartile range. B H&E image and manual annotation from the original study. C Clustering results by nonspatial and spatial methods, Seurat …"

DOI ↗ Fig. 2 on PMC ↗

Fig. 4 — TEXT

cGAS-STING drives ageing-related inflammation and neurodegeneration. *Nature, 2023*

"…For each dataset, we filtered out low-quality and potential doublet cells on the basis of violin plot distributions of genes, UMI counts and mitochondrial gene expression (as determined using the Seurat …"

DOI ↗ Fig. 4 on PMC ↗

Figure 3 — TEXT

hdWGCNA identifies co-expression networks in high-dimensional transcriptomics data. *Cell Rep Methods, 2023*

"…•hdWGCNA constructs co-expression networks in high-dimensional transcriptomics data•hdWGCNA provides tools for statistics, visualization, and downstream interpretation•hdWGCNA is an open-source R package that …"

DOI ↗ Figure 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/Seurat/Seurat_recreate.R`

√ verified run · exit 0 · 13 s

Seurat · 6 variations

RECREATIONS/SEURAT/



v1 DimPlot · EMBEDDINGS



v2 FeaturePlot · EMBEDDINGS



v3 VlnPlot · COMPOSITION



v4 DotPlot · HEATMAPS



v5 DoHeatmap · HEATMAPS



v6 RidgePlot · COMPOSITION

clusterProfiler

#3

OF 100 · METHODS MENTIONS

R | ENRICHMENT | ENRICHMENT & PATHWAY ACTIVITY

3,518	19.8%	3,609	analysis+plotting	6
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

GO/KEGG ORA dot & bar plots, cnetplot, emapplot, GSEA running-score (via enrichplot)

Also catalogued as: **enrichplot** (#32, 280 papers)

THIS RECREATION

clusterProfiler + enrichplot: ORA and GSEA visualisations

KEY PLOTTING CALL

RECREATIONS/CLUSTERPROFILER/CLUSTERPROFILER_RECREATE.R — EXCERPT · ENDS LINE 7

```
res <- as.data.frame(readRDS("../_data/airway_deseq2.rds")$res); res <- res[!is.na(res$padj) & !is.na(res$entrez),]
up <- unique(res$entrez[res$padj < .01 & res$log2FoldChange > 1]); uni <- unique(res$entrez)
ego <- enrichGO(up, OrgDb = org.Hs.eg.db, ont = "BP", universe = uni, readable = TRUE); ego <- simplify(ego)
ggsave("v1_dotplot.png", dotplot(ego, showCategory = 15) + ggtitle("enrichGO BP: dexamethasone-up genes"), width = 7.5, height
= 6.5, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 7 — FIGURE CAPTION

Cell-type-specific CAG repeat expansions and toxicity of mutant Huntingtin in human striatum and cerebellum. *Nat Genet, 2024*

"...es are omitted from the plot. For panels c and f, the significance threshold for enrichment analysis was: q value < 0.05, Padj < 0.05, adjusted for multiple comparisons after hypergeometric test with ..."

DOI ↗ Fig. 7 on PMC ↗

Extended Data Fig. 7 — FIGURE CAPTION

Blood-brain barrier disruption and sustained systemic inflammation in individuals with long COVID-associated cognitive impairment. *Nat Neurosci, 2024*

"...ed Gene Ontology biological processes in each cohort. Bubble plots show top 20 enriched terms with absLog2FC cut-off < 0.59 and adjusted p-value < 0.05. Pathway enrichment p values were calculated in ..."

DOI ↗ Extended Data Fig. 7 on PMC ↗

Fig. 1 — TEXT

Transcriptional and epigenetic decoding of the microglial aging process. *Nat Aging, 2023*

"...categories were simplified using the 'simplify' function with a cutoff value = 0.7, and visualization was performed using the dotplot and emapplot functions in clusterProfiler 3.14.3."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

- enrichplot (#32): Shares clusterProfiler recreations (all plots are enrichplot functions)

Reproduce: `Rscript recreations/clusterProfiler/clusterProfiler_recreate.R`

√ verified run · exit 0 · 57 s

clusterProfiler · 6 variations



v1 dotplot

ENRICHMENT



v2 barplot

ENRICHMENT



v3 cnetplot

ENRICHMENT



v4 emapplot

ENRICHMENT



v5 gseaplot2

ENRICHMENT



v6 ridgeplot

ENRICHMENT

ggplot2

#4

OF 100 · METHODS MENTIONS

R · GENERAL · DE, QC & SET OVERLAPS · EMBEDDINGS & FEATURE MAPS · COMPOSITION, DISTRIBUTIONS & LAYOUTS

2,490	14.01%	2,546	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Everything custom: volcano, bar/box/violin, UMAP scatter, composition stacked bars, multi-panel

Also catalogued as: **ggrepel** (#64, 92 papers)

THIS RECREATION

ggplot2: the grammar-of-graphics backbone of most omics figures

KEY PLOTTING CALL

RECREATIONS/GGPLOT2/GGPLOT2_RECREATE.R — EXCERPT · ENDS LINE 11

```
res <- res |> filter(!is.na(padj)) |> mutate(sig = case_when(padj < .05 & log2FoldChange > 1 ~ "Up", padj < .05 &
log2FoldChange < -1 ~ "Down", TRUE ~ "NS"))
lab <- res |> filter(sig != "NS") |> arrange(padj) |> head(15)
# V1: volcano plot
g <- ggplot(res, aes(log2FoldChange, -log10(padj), color = sig)) + geom_point(size = .8, alpha = .7) +
  scale_color_manual(values = c(Up = "#D7301F", Down = "#2B8CBE", NS = "grey75")) + geom_vline(xintercept = c(-1,1), lty = 2) +
geom_hline(yintercept = -log10(.05), lty = 2) +
  geom_text_repel(data = lab, aes(label = symbol), size = 3, max.overlaps = 30, show.legend = FALSE) + theme_classic() + labs(t
itle = "Volcano (ggplot2 + ggrepel)")
ggsave("v1_volcano.png", g, width = 6, height = 5, dpi = 150)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT
Microglia ferroptosis is regulated by SEC24B and contributes to neurodegeneration. *Nat Neurosci, 2023*
"...UMAP plots and gene expression plots were generated using built-in Seurat/ggplot2 plotting functions or scCustomize version 0.7.0 unless otherwise described."
DOI ↗ Fig. 1 on PMC ↗

Figure 7 TEXT
Macrophage-mediated myelin recycling fuels brain cancer malignancy. *Cell, 2024*
"...Plots were generated using the package ggplot (v.3.3.6) or ggpubr (v.0.4.0), except where mentioned."
DOI ↗ Figure 7 on PMC ↗

Fig. 6 TEXT
Microglia regulate central nervous system myelin growth and integrity. *Nature, 2023*
"...ggplot2 v.3.3.5 was used to perform custom plots, here v.1.0.1 was used to ensure reproducible paths, and Matrix v.1.3.4 was used for handling sparse matrices."
DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

- ggrepel (#64): See ggplot2 v1_volcano.png (ggrepel labels)

Reproduce: `Rscript recreations/ggplot2/ggplot2_recreate.R`

√ verified run · exit 0 · 2 s

ggplot2 · 4 variations



v1 volcano · DE



v2 umap · EMBEDDINGS



v3 composition_bar · COMPOSITION



v4 violin_box · COMPOSITION

GSEA (Broad)

#5

OF 100 · METHODS MENTIONS

DESKTOP | ENRICHMENT | ENRICHMENT & PATHWAY ACTIVITY

2,056	11.57%	3,310	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Classic running enrichment-score plot with hit ticks and ranked metric; NES bar/dot summaries

THIS RECREATION

Broad GSEA "classic" enrichment figure, recreated with GSEApy prerank (same algorithm/plot layout as the Broad desktop app)

KEY PLOTTING CALL

RECREATIONS/GSEA_BROAD/GSEA_STYLE_RECREATE.PY — EXCERPT · ENDS LINE 8

```
res=pd.read_csv("../_data/airway_deseq2_results.csv",index_col=0).dropna(subset=["stat","symbol"])
rnk=res.groupby("symbol").stat.mean().sort_values(ascending=False)
pre=gp.prerank(rnk=rnk,gene_sets="MSigDB_Hallmark_2020",permutation_num=1000,seed=1,outdir=None,min_size=15,max_size=500,threads=4)

r=pre.res2d.sort_values("NOM p-val"); terms=r.Term.tolist()
gp.gseaplot(rank_metric=pre.ranking,term=terms[0],**pre.results[terms[0]],ofname="v1_classic_enrichment.png")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 1 — FIGURE CAPTION

Macrophage-mediated myelin recycling fuels brain cancer malignancy. *Cell, 2024*

"...ubtypes (outer circle) per model. Pseudo-location2 assignment of glioblastoma cells (inner circle) displayed as a percentage of cellular subtype.(E) UMAP representation of MG and MDM sub-clusters.(F) Gene set ..."

DOI ↗ Figure 1 on PMC ↗

Extended Data Fig. 8 — FIGURE CAPTION

Molecular and cellular dynamics of the developing human neocortex. *Nature, 2025*

"...sion levels of cell-type-specific marker genes and SSTR2. c, Scatter plots illustrating the Pearson correlation of log fold changes in individual genes between Ostreotide and L-054,264 treatments. d, Gene set ..."

DOI ↗ Extended Data Fig. 8 on PMC ↗

Fig. 1 — TEXT

Specialized astrocytes mediate glutamatergic gliotransmission in the CNS. *Nature, 2023*

"...Using gene set enrichment analysis (GSEA) databases (Methods), we found significant Gene Ontology (GO) enrichment for each cluster (Extended Data Fig."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

- Broad desktop app is GUI; plot recreated with GSEApy prerank (same algorithm & layout)

Reproduce: `python3 recreations/GSEA_Broad/gsea_style_recreate.py`

√ verified run · exit 0 · 4 s

GSEA (Broad) · 3 variations

RECREATIONS/GSEA_BROAD/



v1 classic_enrichment

ENRICHMENT



v2 multi_term

ENRICHMENT



v3 dotplot

ENRICHMENT

limma

#6

OF 100 · METHODS MENTIONS

R | BULK QC/DE | DE, QC & SET OVERLAPS

2,047	11.52%	2,116	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

voom mean-variance, MDS, MD/MA, volcanoplot, Venn of contrasts

KEY PLOTTING CALL

RECREATIONS/LIMMA/LIMMA_RECREATE.R — EXCERPT · ENDS LINE 11

```
png("v1_voom_meanvar.png", 1000, 800, res = 150); v <- voom(y, design, plot = TRUE); dev.off()
fit <- eBayes(lmFit(v, design)); cf <- ncol(design)
# V2: MDS plot
png("v2_plotMDS.png", 1000, 850, res = 150); plotMDS(v, labels = paste(dds$dex, dds$cell), col = ifelse(dds$dex == "trt", "#984
EA3", "#4DAF4A"), main = "limma plotMDS"); dev.off()
# V3: MD plot with DE status
png("v3_plotMD.png", 1000, 800, res = 150); plotMD(fit, coef = cf, status = decideTests(fit)[,cf], main = "plotMD (dex)",
hl.col = c("#D7301F","#2B8CBE")); abline(h=0,col="grey"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 — FIGURE CAPTION

CGRP sensory neurons promote tissue healing via neutrophils and macrophages. *Nature, 2024*

"...SP-1 in neutrophils and macrophages to mediate its activity.a,b, RNA-seq analysis of CGRP-treated neutrophils and macrophages (1 nM or saline for 4 h). Differential gene expression was performed with ..."

DOI ↗ Fig. 3 on PMC ↗

Fig. 1 — FIGURE CAPTION

A TREM2-activating antibody with a blood-brain barrier transport vehicle enhances microglial metabolism in Alzheimer's disease models. *Nat Neurosci, 2023*

"...expression was generated by summing every counts per gene. Each mouse was treated as a biological replicate, and differential expression (DE) was performed for each cluster versus 'homeostatic' using limma. g ..."

DOI ↗ Fig. 1 on PMC ↗

Figure 3 — TEXT

Single-cell RNA sequencing and spatial transcriptomics reveal cancer-associated fibroblasts in glioblastoma with protumoral effects. *J Clin Invest, 2023*

"...For differential gene expression analysis, raw read counts were processed in iDEP using the limma-voom function."

DOI ↗ Figure 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/limma/limma_recreate.R`

√ verified run · exit 0 · 9 s

limma · 4 variations



v1 voom_meanvar · DE



v2 plotMDS · DE



v3 plotMD · DE



v4 volcanoplot · DE

edgeR

#7

OF 100 · METHODS MENTIONS

R · BULK QC/DE · DE, QC & SET OVERLAPS

1,892	10.65%	1,949	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

BCV, MDS, smear/MD plots, QL dispersion

KEY PLOTTING CALL

RECREATIONS/EDGER/EDGER_RECREATE.R — EXCERPT · ENDS LINE 6

```
dds <- readRDS("../_data/airway_deseq2.rds")$dds; grp <- dds$dex
y <- DGEList(counts(dds), group = grp); y <- y[filterByExpr(y), , keep.lib.sizes = FALSE]; y <- normLibSizes(y)
design <- model.matrix(~ dds$cell + grp); y <- estimateDisp(y, design)
png("v1_plotBCV.png", 1000, 800, res = 150); plotBCV(y, main = "edgeR plotBCV"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 — FIGURE CAPTION

Cell type specific transcriptomic differences in depression show similar patterns between males and females but implicate distinct cell types and genes. *Nat Commun, 2023*

"...ers with highest number of DEGs in the female cluster level analysis—f microglia, g inhibitory neuronal clusters. For f, g, the plotted values are pseudobulk CPMs (counts per million) calculated with edgeR and ..."

DOI ↗ Fig. 3 on PMC ↗

Fig. 1 — TEXT

Microglia ferroptosis is regulated by SEC24B and contributes to neurodegeneration. *Nat Neurosci, 2023*

"...The PCA plots were generated using edgeR and Glimma."

DOI ↗ Fig. 1 on PMC ↗

Figure 1 — TEXT

Serotonin reduction in post-acute sequelae of viral infection. *Cell, 2023*

"...Briefly, transcript quantification data were summarized to genes using the tximport package and normalized using the trimmed mean of M values (TMM) method in edgeR."

DOI ↗ Figure 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/edgeR/edgeR_recreate.R`

√ verified run · exit 0 · 13 s

LEAD FIGURE



v1 plotBCV

DE

edgeR · 4 variations · v1 on facing page



v2 plotMDS

DE



v3 plotQLDisp

DE



v4 plotSmear

DE

pheatmap

#9

OF 100 · METHODS MENTIONS

R · HEATMAP · HEATMAPS & DOT MATRICES

1,119	6.3%	1,138	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Clustered expression heatmaps with annotation bars; sample-distance heatmaps

KEY PLOTTING CALL

RECREATIONS/PHEATMAP/PHEATMAP_RECREATE.R — EXCERPT · ENDS LINE 7

```
top <- head(order(-matrixStats::rowVars(vsd)), 50); m <- vsd[top,]; rownames(m) <- ifelse(is.na(res[rownames(m),"symbol"]), row
names(m), res[rownames(m),"symbol"])
pheatmap(m, scale = "row", annotation_col = ann, show_colnames = FALSE, fontsize_row = 6, main = "Top 50 variable genes (row z-
score)", filename = "v1_topvar_zscore.png", width = 6, height = 8)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT

Microglia ferroptosis is regulated by SEC24B and contributes to neurodegeneration. *Nat Neurosci, 2023*

"...The top 100 upregulated genes were plotted using the Pheatmap package version 1.0.12 in R using unbiased hierarchical clustering."

DOI ↗ Fig. 1 on PMC ↗

Fig. 3 TEXT

Mitophagy curtails cytosolic mtDNA-dependent activation of cGAS/STING inflammation during aging. *Nat Commun, 2024*

"...Heatmaps were generated using the tidyverse, ggplot2, viridis, RColorBrewer and pheatmap packages in R."

DOI ↗ Fig. 3 on PMC ↗

Figure 1 TEXT

APOE modulates microglial immunometabolism in response to age, amyloid pathology, and inflammatory challenge. *Cell Rep, 2023*

"...Finally, the average AUCell scores of each KEGG metabolic pathway were plotted as heatmaps using pheatmap R software (v.1.0.12) sorted by either cell subtypes and/or experimental groups."

DOI ↗ Figure 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/pheatmap/pheatmap_recreate.R` — ✓ verified run · exit 0 · 16 s

LEAD FIGURE



v2 sample_distance — HEATMAPS

pheatmap · 4 variations · v2 on facing page



v1 topvar_zscore

HEATMAPS



v3 cutree

HEATMAPS



v4 sc_markers

HEATMAPS

WGCNA

#10

OF 100 · METHODS MENTIONS

R · NETWORK/MODULES · NETWORKS, GRAPHS & TREES · HEATMAPS & DOT MATRICES

884	4.98%	1,025	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Soft-threshold plots, gene dendrogram + module colours, module-trait heatmap, eigengene network

THIS RECREATION

WGCNA on pseudo-bulk samples built from Zeisel cortex/hippocampus cells (7 cell types x 10 random pseudo-replicates)

KEY PLOTTING CALL

RECREATIONS/WGCNA/WGCNA_RECREATE.R — EXCERPT · ENDS LINE 11

```
sft <- pickSoftThreshold(datExpr, powerVector = c(1:10, seq(12,20,2)), networkType = "signed", verbose = 0)
png("v1_soft_threshold.png", 1400, 650, res = 150); par(mfrow = c(1,2)); fi <- sft$fitIndices
plot(fi[,1], -sign(fi[,3])*fi[,2], type = "n", xlab = "Soft threshold (power)", ylab = "Scale-free fit, signed R^2", main = "Sc
ale independence"); text(fi[,1], -sign(fi[,3])*fi[,2], fi[,1], col = "red"); abline(h = .8, col = "red")
plot(fi[,1], fi[,5], type = "n", xlab = "Soft threshold (power)", ylab = "Mean connectivity", main = "Mean connectivity");
text(fi[,1], fi[,5], fi[,1], col = "red"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 3 — FIGURE CAPTION

Distinct molecular profiles of skull bone marrow in health and neurological disorders. *Cell, 2023*

"...(B) six bones, dura, and brain and (C) six bones in naive, sham, and MCAo conditions.(D) Dendrogram demonstrates the relation among bones and conditions. (E–J) Protein expression modules identified by WGCNA ..."

DOI ↗ Figure 3 on PMC ↗

Fig. 6 — FIGURE CAPTION

Cell type specific transcriptomic differences in depression show similar patterns between males and females but implicate distinct cell types and genes. *Nat Commun, 2023*

"...WGCNA results for microglia and PV interneurons in females.a Heatmap showing the correlation and associated p-value, in parentheses, of Mic1 WGCNA module eigengenes with case-control status and covariates ..."

DOI ↗ Fig. 6 on PMC ↗

Fig. 6 — FIGURE CAPTION

The basolateral amygdala-anterior cingulate pathway contributes to depression-like behaviors and comorbidity with chronic pain behaviors in male mice. *Nat Commun, 2023*

"...ng the level of significance of overlaps between mice and men gene modules (two-sided Fisher Exact Test). The highest overlap (P = 8.36E-49) was obtained for the man/yellow and mouse/brown modules. b WGCNA ..."

DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/WGCNA/WGCNA_recreate.R`

✓ verified run · exit 0 · 25 s

WGCNA · 4 variations



v1 soft_threshold

NETWORKS



v2 dendro_modules

NETWORKS



v3 module_trait

HEATMAPS



v4 eigengene_network

NETWORKS

CellChat

#11

OF 100 · METHODS MENTIONS

R | CELL-CELL COMMUNICATION | CELL–CELL COMMUNICATION

839	4.72%	976	analysis+plotting	5
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Circle/chord communication networks, bubble plots of L-R pairs, signalling-role scatter/heatmaps

THIS RECREATION

CellChat on Zeisel mouse cortex/hippocampus cell types

KEY PLOTTING CALL

RECREATIONS/CELLCHAT/CELLCHAT_RECREATE.R — EXCERPT · ENDS LINE 10

```
cc <- identifyOverExpressedGenes(cc); cc <- identifyOverExpressedInteractions(cc); cc <- computeCommunProb(cc, type = "triMean")
; cc <- filterCommunication(cc, min.cells = 10)
cc <- computeCommunProbPathway(cc); cc <- aggregateNet(cc); cc <- netAnalysis_computeCentrality(cc, slot.name = "netP")
gs <- as.numeric(table(cc@idents))
png("v1_circle_counts.png", 1200, 1200, res = 160); netVisual_circle(cc@net$count, vertex.weight = gs, weight.scale = TRUE, lab
el.edge = FALSE, title.name = "Number of interactions"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 8 FIGURE CAPTION

Transcriptional and epigenetic decoding of the microglial aging process. *Nat Aging, 2023*

"...across all 16 cell types (18 populations). d, Significant ligand–receptor interactions predicted by CellPhoneDB. One-sided permutation test. e, Significant molecule–molecule interactions predicted by CellChat ..."

DOI ↗ Fig. 8 on PMC ↗

Fig. 6 FIGURE CAPTION

A brain cell atlas integrating single-cell transcriptomes across human brain regions. *Nat Med, 2024*

"...gnificant ligand–receptor pairs in the NRG, CADM, NEGR and laminin pathways from all senders to hippocampal microglia (PCDH9high). P values are computed from a one-sided permutation test according to CellChat ..."

DOI ↗ Fig. 6 on PMC ↗

Extended Data Fig. 5: FIGURE CAPTION

Microglial mechanisms drive amyloid-β clearance in immunized patients with Alzheimer's disease. *Nat Med, 2025*

"...A2M, APOE, FAM107A, LIPA, SPP1, and TREM2 in Aβ-associated Mg2-enriched and Mg4-enriched ST spots compared to the nAD control group for AN1792 and the lecanemab case. j, Chord plots showing inferred CellChat ..."

DOI ↗ Article ↗ figure page not on PMC

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/CellChat/CellChat_recreate.R`

✓ verified run · exit 0 · 145 s

CellChat · 5 variations



v1 circle_counts · CELL–CELL



v2 heatmap · CELL–CELL



v3 bubble · CELL–CELL



v4 signalingRole_scatter · CELL–CELL



v5 chord_pathway · CELL–CELL

STRING

#12

OF 100 · METHODS MENTIONS

WEB | PPI NETWORK | NETWORKS, GRAPHS & TREES

838	4.72%	1,142	analysis+plotting	2
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

PPI network images (web/API), often re-styled in Cytoscape

THIS RECREATION

STRING: (V1) the canonical STRING network image via the official API; (V2) the same PPI edges redrawn with networkx, coloured by log2FC

KEY PLOTTING CALL

RECREATIONS/STRING/STRING_RECREATE.PY — EXCERPT · ENDS LINE 7

```
res=pd.read_csv("../_data/airway_deseq2_results.csv",index_col=0).dropna(subset=["padj","symbol"])
top=res[(res.padj<1e-6)&(res.log2FoldChange.abs()>1.5)].nsmallest(60,"padj")
ids="%0d".join(top.symbol); API="https://string-db.org/api"
r=requests.post(f"{API}/image/network",data=dict(identifiers=ids,species=9606,network_flavor="confidence",required_score=400,ca
ller_identity="novaflow_fig_catalog"),timeout=120); r.raise_for_status();
open("v1_string_api_network.png","wb").write(r.content)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 5 FIGURE CAPTION

Circular RNA encoded MET variant promotes glioblastoma tumorigenesis. *Nat Commun, 2023*

"...tated by an anti-MET antibody using mass spectrometry analysis. g Reactome pathway analysis of candidate MET404-binding partners. h Protein interaction network with MET as the hub generated using the STRING ..."

DOI ↗ Fig. 5 on PMC ↗

Fig. 2 FIGURE CAPTION

Dimethyl itaconate ameliorates cognitive impairment induced by a high-fat diet via the gut-brain axis in mice. *Microbiome, 2023*

"...supplementation. Columns with different colors represent different classification in level 2. The dotted line in the figure represents P = 0.05. h PPI network analysis of pro-cognitive genes based on STRING ..."

DOI ↗ Fig. 2 on PMC ↗

Extended Data Fig. 11 FIGURE CAPTION

Astrocyte-neuron subproteomes and obsessive-compulsive disorder mechanisms. *Nature, 2023*

"...entified in astrocyte plasma membrane with Astro Lck-BioID2 . Node size represents the enrichment of each protein vs the GFP control (log2(BioID2/GFP)). Edges represent putative interactions from the STRING ..."

DOI ↗ Extended Data Fig. 11 on PMC ↗

LIMITATIONS & NOTES

- V1 = official STRING API image; V2 = same edges redrawn in networkx
- Needs internet: string-db.org API (METHODS §4).

Reproduce: `python3 recreations/STRING/string_recreate.py`

√ verified run · exit 0 · 3 s

STRING · 2 variations



v1 string_api_network

NETWORKS



v2 networkx_fc

NETWORKS

Enrichr

#13

WEB | ENRICHMENT | ENRICHMENT & PATHWAY ACTIVITY

OF 100 · METHODS MENTIONS

778	4.38%	912	analysis+plotting	2
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Enrichment bar charts / tables by library

THIS RECREATION

Enrichr (web API, via GSEApy) bar/dot plots for dexamethasone-up genes

KEY PLOTTING CALL

RECREATIONS/ENRICHR/ENRICHR_RECREATE.PY — EXCERPT · ENDS LINE 7

```
res=pd.read_csv("../_data/airway_deseq2_results.csv",index_col=0).dropna(subset=["padj","symbol"])
up=res[(res.padj<.01)&(res.log2FoldChange>1)].symbol.unique().tolist()
enr=gp.enrichr(gene_list=up,gene_sets=["GO_Biological_Process_2023","KEGG_2021_Human"],organism="human",outdir=None)
gp.barplot(enr.results,column="Adjusted P-value",group="Gene_set",size=10,top_term=6,figsize=(5,7),color={"GO_Biological_Proces
s_2023":"#4C72B0","KEGG_2021_Human":"#DD8452"},ofname="v1_barplot_grouped.png")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 6 — FIGURE CAPTION

GAP43-dependent mitochondria transfer from astrocytes enhances glioblastoma tumorigenicity. *Nat Cancer, 2023*

"...to-mCherry+ L1 cells; n = 3 independent co-culture experiments. Data were analyzed by paired two-tailed t-test; 2-OG, α-ketoglutarate; CDP, cytidine-5'-diphosphate. c, Phospho-array pathway analysis (Enrichr) ..."

DOI ↗ Fig. 6 on PMC ↗

Fig. 3 — TEXT

Cell-type-specific aging clocks to quantify aging and rejuvenation in neurogenic regions of the brain. *Nat Aging, 2023*

"...Statistics were exported from the Enrichr web tool and processed and visualized in R with ggplot2 (version 3.3.3) package."

DOI ↗ Fig. 3 on PMC ↗

Figure 6 — TEXT

hdWGCNA identifies co-expression networks in high-dimensional transcriptomics data. *Cell Rep Methods, 2023*

"...REAGENT or RESOURCESOURCEIDENTIFIERDeposited dataHuman Alzheimer's Disease snRNA-seq 2019Mathys et al., 201910syn18485175Human Autism spectrum disorder snRNA-seqVelmeshev et al., 20199PRJNA434002Human aging ..."

DOI ↗ Figure 6 on PMC ↗

LIMITATIONS & NOTES

- Uses the live Enrichr web API via GSEApy
- Needs internet: calls the live Enrichr web API (METHODS §4).

Reproduce: `python3 recreations/Enrichr/enrichr_recreate.py`

√ verified run · exit 0 · 4 s

Enrichr · 2 variations



v1 barplot_grouped

ENRICHMENT



v2 dotplot_by_library

ENRICHMENT

scanpy

#14

OF 100 · METHODS MENTIONS

PYTHON | SINGLE-CELL (EMBEDDING/DOT/VIOLIN/MATRIX) | EMBEDDINGS & FEATURE MAPS | HEATMAPS & DOT MATRICES | COMPOSITION, DISTRIBUTIONS & LAYOUTS | TRAJECTORY, PSEUDOTIME & VELOCITY

763	4.29%	789	analysis+plotting	7
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

UMAP, dotplot, matrixplot, stacked violin, rank_genes_groups, tracksplot, PAGA

Also catalogued as: **PAGA** (#75, 75 papers)

THIS RECREATION

scanpy plotting API on Zeisel mouse cortex/hippocampus (3,005 cells)

KEY PLOTTING CALL

RECREATIONS/SCANPY/SCANPY_RECREATE.PY — EXCERPT · ENDS LINE 7

```
mk={"Neuron":["Snap25","Syt1"],"Inhib":["Gad1","Gad2","Pvalb","Sst"],"Exc":["Satb2","Neurod6","Wfs1"],"Astro":["Aqp4","Gfap"],"
Oligo":["Mbp","Plp1"],"Micro":["Cx3cr1","C1qa"],"Vasc":["Cldn5","Acta2"]}
sc.pl.umap(ad,color=["celltype","leiden","Gad1","Mbp"],ncols=2,wspace=.4,save="_v1_umap_grid.png",show=False)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 — FIGURE CAPTION

Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST. *Nat Commun, 2023*

"...the Stereo-seq mouse embryo.A Tissue domain annotations of the E9.5 mouse embryo data taken from the original Stereo-seq study wherein the clusters were first identified using Leiden clustering from SCANPY and ..."

DOI ↗ Fig. 3 on PMC ↗

Fig. 1 — TEXT

The molecular cytoarchitecture of the adult mouse brain. *Nature, 2023*

"...To build a graph of cell type similarity, we used Scanpy on our subsampled data to compute the connectivities over a 20 neighbour local neighbourhood using 250 principal components (the section 'Visualization ..."

DOI ↗ Fig. 1 on PMC ↗

Fig. 1 — TEXT

APOE4/4 is linked to damaging lipid droplets in Alzheimer's disease microglia. *Nature, 2024*

"...We then used the batch-corrected neighbourhood graph to run Leiden clustering46 and to calculate a global UMAP embedding with default parameters in Scanpy (v.1.9.1)."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

- PAGA (#75): See scanpy v7_paga.png

Reproduce: `python3 recreations/scanpy/scanpy_recreate.py`

√ verified run · exit 0 · 30 s

scanpy · 7 variations



v1 umap_grid · EMBEDDINGS



v2 dotplot · HEATMAPS



v3 matrixplot · HEATMAPS



v4 stacked_violin · COMPOSITION



v5 rank_genes_dotplot · HEATMAPS



v6 tracksplot · HEATMAPS



v7 paga · TRAJECTORY

ComplexHeatmap

#17

OF 100 · METHODS MENTIONS

R | HEATMAP | HEATMAPS & DOT MATRICES

567	3.19%	578	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Multi-annotated, split heatmaps; oncoPrint; density heatmaps; heatmap lists

KEY PLOTTING CALL

RECREATIONS/COMPLEXHEATMAP/COMPLEXHEATMAP_RECREATE.R — EXCERPT · ENDS LINE 13

```
png("v1_annotated_split.png", 1200, 1500, res = 170)
draw(Heatmap(m, name = "z", col = colorRamp2(c(-2,0,2), c("#2166AC","white","#B2182B")), top_annotation = ha, right_annotation
= ra,
  row_split = ifelse(res[sig,"log2FoldChange"] > 0, "Up in dex", "Down in dex"), column_split = dds$dex, show_row_names = FALSE,
  left_annotation = rowAnnotation(mark = anno_mark(at = match(lab, sig), labels = res[lab,"symbol"], side = "left"))) )
dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT

The molecular cytoarchitecture of the adult mouse brain. *Nature, 2023*

"...The matrix was plotted using the ComplexHeatmap package in R79."

DOI ↗ Fig. 1 on PMC ↗

Fig. 1 TEXT

Exposure of iPSC-derived human microglia to brain substrates enables the generation and manipulation of diverse transcriptional states in vitro. *Nat Immunol, 2023*

"...Heatmaps were generated using the ComplexHeatmap R package."

DOI ↗ Fig. 1 on PMC ↗

Fig. 2 TEXT

Multiomics in primary and metastatic breast tumors from the AURORA US network finds microenvironment and epigenetic drivers of metastasis. *Nat Cancer, 2023*

"...The significantly differentially expressed modules (q < 0.05) were hierarchically clustered using the ComplexHeatmap R package."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ComplexHeatmap/ComplexHeatmap_recreate.R`

✓ verified run · exit 0 · 21 s

ComplexHeatmap · 4 variations



v1 annotated_split

HEATMAPS



v2 singlecell_markers

HEATMAPS



v3 correlation_cellfun

HEATMAPS



v4 densityHeatmap

HEATMAPS

g:Profiler

#18

OF 100 · METHODS MENTIONS

WEB/R/PYTHON | ENRICHMENT | ENRICHMENT & PATHWAY ACTIVITY

563	3.17%	627	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Manhattan-style enrichment plot (gostplot) + term tables

KEY PLOTTING CALL

RECREATIONS/GPROFILER/GPROFILER2_RECREATE.R — EXCERPT · ENDS LINE 9

```
set_base_url("https://biit.cs.ut.ee/gprofiler")  # HTTPS endpoint (port 80 is closed on this box)
res <- as.data.frame(readRDS("../_data/airway_deseq2.rds")$res); res <- res[!is.na(res$padj) & !is.na(res$symbol),]
up <- res$symbol[res$padj < .01 & res$log2FoldChange > 1]; dn <- res$symbol[res$padj < .01 & res$log2FoldChange < -1]
g <- gost(list(up = up, down = dn), organism = "hsapiens", multi_query = FALSE, sources = c("GO:BP","KEGG","REAC","TF"))
p <- gostplot(g, capped = TRUE, interactive = FALSE); ggsave("v1_manhattan.png", p, width = 9, height = 6, dpi = 130, bg = "whi
te")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Extended Data Fig. 8 — FIGURE CAPTION

Myelin dysfunction drives amyloid-β deposition in models of Alzheimer's disease. *Nature, 2023*

"...notype data. Error bars represent SD. (e) Heatmap of normalised mean expression pattern in the 10 DEG clusters across different genotypes. Each cluster of genes underwent GO enrichment analysis using ..."

DOI ↗ Extended Data Fig. 8 on PMC ↗

Extended Data Fig. 6 — FIGURE CAPTION

Single-cell multiregion dissection of Alzheimer's disease. *Nature, 2024*

"..., subtype, and covariate enrichments performed at cell level by stratifying cells with z-score > 1 and testing for regional or subtype enrichment (see Methods). Functional enrichments performed using ..."

DOI ↗ Extended Data Fig. 6 on PMC ↗

Extended Data Fig. 5 — TEXT

A NPAS4-NuA4 complex couples synaptic activity to DNA repair. *Nature, 2023*

"...Gene ontology (GO) enrichment analysis was performed using gProfiler2 in R (v.0.2.0)65, with a custom background of expression-filtered genes from neuronal cell types across all scRNA-seq datasets in this ..."

DOI ↗ Extended Data Fig. 5 on PMC ↗

LIMITATIONS & NOTES

- Uses live g:Profiler API (HTTPS)
- Needs internet: live g:Profiler API. Box shims: gprofiler2 is pointed at its HTTPS endpoint and curl::has_internet() is overridden, because its DNS probe fails in this sandbox (METHODS §4).

Reproduce: `Rscript recreations/gProfiler/gprofiler2_recreate.R`

√ verified run · exit 0 · 16 s

g:Profiler · 3 variations

RECREATIONS/GPROFILER/



v1 manhattan

ENRICHMENT



id	source	term_id	term_name	term_size	p_value down	p_value up
1	GO:BP	GO:0048731	system development	4023	3.6e-19	5.8e-06
2	GO:BP	GO:0051239	regulation of multicellular organismal process	2923	3.6e-19	1.2e-06
3	GO:BP	GO:0007154	cell communication	6598	4.4e-19	7.6e-10
4	GO:BP	GO:0032501	multicellular organismal process	7337	6.1e-19	3.0e-10
5	GO:BP	GO:0023052	signaling	6558	1.7e-18	4.1e-10
6	GO:BP	GO:0032502	developmental process	6535	6.9e-17	5.9e-11

g:Profiler (biit.cs.ut.ee/gprofiler)

v2 publish_highlight

ENRICHMENT

source	term_id	term_name	term_size	intersection_size_up	intersection_size_down	p_value_up	p_value_do
GO:BP	GO:0048731	system development	4023	122	144	5.8e-06	3.6e-19
GO:BP	GO:0051239	regulation of multicellular organismal process	2923	99	119	1.2e-06	3.6e-19
GO:BP	GO:0007154	cell communication	6598	189	195	7.6e-10	4.4e-19
GO:BP	GO:0032501	multicellular organismal process	7337	205	208	3.0e-10	6.1e-19
GO:BP	GO:0023052	signaling	6558	189	193	4.1e-10	1.7e-18
GO:BP	GO:0032502	developmental process	6535	191	189	5.9e-11	6.9e-17

g:Profiler (biit.cs.ut.ee/gprofiler)

v3 gosttable

ENRICHMENT

IGV

#19

OF 100 · METHODS MENTIONS

DESKTOP/WEB | GENOME TRACKS | GENOME TRACKS & GENOMIC SIGNAL

549	3.09%	794	viewer	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · GVIZ

Neuro-corpus mentions per year (*2026 partial) · niche rank **#73**, growth **0.55×**, visual 3/5, score 0.314. Broad-corpus rate 2.229 → 1.219 per 1k papers (2023 → 2025–26).
GitHub ivanek/Gviz · created 2018-12-10 · last push 2026-08-13

BEST FOR

Genome-browser track screenshots (coverage, peaks, splicing)

Also catalogued as: **UCSC Genome Browser** (#28, 363 papers)

THIS RECREATION

IGV / UCSC-browser-style tracks with Gviz: ideogram + axis + gene models + data tracks (hg19)

KEY PLOTTING CALL

RECREATIONS/GVIZ_IGV_STYLE/GVIZ_RECREATE.R — EXCERPT · ENDS LINE 14

```
at <- AnnotationTrack(pks, name = "peaks", fill = "#e31a1c", col = NA); set.seed(1)
cov <- DataTrack(GRanges(chr, IRanges(seq(st, en - 5000, 5000), width = 5000), score = sapply(seq(st, en - 5000, 5000),
function(x) sum(overlapsAny(pks, GRanges(chr, IRanges(x, width = 5000)))) * pmax(0, rnorm(1, 5, 1)))), name = "peak signal", ty
pe = "histogram", fill = "#41ab5d", col.histogram = NA)
ax <- GenomeAxisTrack(); bands <- data.frame(chrom = "chr1", chromStart = c(0, 2300000), chromEnd = c(2300000, 5300000), name =
c("p36.33","p36.32"), gieStain = c("gneg","gpos25"))
id <- IdeogramTrack(genome = "hg19", chromosome = chr, bands = bands)
png("v1_igv_style_tracks.png", 1600, 900, res = 160); plotTracks(list(id, ax, cov, at, gtr), from = st, to = en, background.tit
le = "#2b2b2b", sizes = c(1, 1.2, 2, .8, 4)); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 — FIGURE CAPTION

A NPAS4-NuA4 complex couples synaptic activity to DNA repair. *Nature, 2023*

"…NPAS4–NuA4-bound sites undergo recurrent DNA breaks in vivo.a, A model for the dual function of NPAS4–NuA4 in stimulating transcription and DNA repair in active neurons. b, IGV tracks displaying aggregate …"

DOI ↗ Fig. 3 on PMC ↗

Fig. 3 — FIGURE CAPTION

Lineage-specific canonical and non-canonical activity of EZH2 in advanced prostate cancer subtypes. *Nat Commun, 2024*

"…and plots show profiles of normalized H3K27me3 and H3K4me3 peaks at ±3 kb of gene bodies or TSS of genes upregulated upon EZH2i treatment in LNCaP-abl (turquoise) or WCM154 (teal). C Screenshots from …"

DOI ↗ Fig. 3 on PMC ↗

Extended Data Fig. 8 — FIGURE CAPTION

The complete sequence and comparative analysis of ape sex chromosomes. *Nature, 2024*

"…on JADBMG010000033.1 (positions 2 to 618,314, upper), a contig from a previous chrY assembly, and T2T chrY (positions 43,632,350 to 44,250,835, bottom), for western lowland gorilla, visualized using IGV. (e) …"

DOI ↗ Extended Data Fig. 8 on PMC ↗

LIMITATIONS & NOTES

- Open-code look-alike: Gviz tracks (recreations/Gviz_IGV_style)
- UCSC Genome Browser (#28): Open-code look-alike: Gviz dense tracks (v3_ucsc_style_dense)

Reproduce: `Rscript recreations/Gviz_IGV_style/gviz_recreate.R`

√ verified run · exit 0 · 94 s



v1 igv_style_tracks

GENOME



v2 zoom_gene_model

GENOME



v3 ucsc_style_dense

GENOME

fgsea

R · ENRICHMENT · ENRICHMENT & PATHWAY ACTIVITY

#20

OF 100 · METHODS MENTIONS

478	2.69%	498	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Enrichment curves (plotEnrichment), GSEA table plot, NES bar charts

KEY PLOTTING CALL

RECREATIONS/FGSEA/FGSEA_RECREATE.R — EXCERPT · ENDS LINE 8

```
res <- as.data.frame(readRDS("../_data/airway_deseq2.rds")$res); res <- res[!is.na(res$stat) & !is.na(res$symbol),]
rk <- sort(tapply(res$stat, res$symbol, mean), decreasing = TRUE)
h <- msigdbr(species = "Homo sapiens", collection = "H"); pw <- split(h$gene_symbol, h$gs_name)
set.seed(1); fr <- fgsea(pw, rk, minSize = 15, maxSize = 500); fr <- fr[order(pval)]
top <- fr$pathway[1]
ggsave("v1_plotEnrichment.png", plotEnrichment(pw[[top]], rk) + labs(title = top, subtitle = sprintf("NES = %.2f, padj = %.1e",
fr$NES[1], fr$padj[1])), width = 6.5, height = 4, dpi = 150, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 — FIGURE CAPTION

Exposure of iPSC-derived human microglia to brain substrates enables the generation and manipulation of diverse transcriptional states in vitro. *Nat Immunol, 2023*

"…sters defined in both the iMGL and brain biopsy datasets. Links with fewer than 200 cells have been removed for clarity. e, Heatmap illustrating the relative enrichment significance (as determined by fgsea …"

DOI ↗ Fig. 2 on PMC ↗

Fig. 4 — FIGURE CAPTION

The molecular cytoarchitecture of the adult mouse brain. *Nature, 2023*

"…ne; SRG, secondary response gene. c, Scatterplot quantifying transcription factor enrichment (P < 0.05, FDR corrected) between excitatory and inhibitory populations. Enrichment scores are computed by fgsea …"

DOI ↗ Fig. 4 on PMC ↗

Fig. 4 — TEXT

Transcriptional linkage analysis with in vivo AAV-Perturb-seq. *Nature, 2023*

"…10g), the list of expressed genes ranked by higher to lower LFC value was used as input to the R package fgsea (v.3.17)59 to run GSEA using the mouse GO:BP dataset."

DOI ↗ Fig. 4 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/fgsea/fgsea_recreate.R`

√ verified run · exit 0 · 286 s

fgsea · 3 variations



v1 plotEnrichment

ENRICHMENT



v2 plotGseaTable

ENRICHMENT



v3 NES_barplot

ENRICHMENT

monocle3

#21

OF 100 · METHODS MENTIONS

R | TRAJECTORY | TRAJECTORY, PSEUDOTIME & VELOCITY

430	2.42%	499	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Trajectory graph on UMAP, pseudotime colouring, genes-in-pseudotime

THIS RECREATION

monocle3 trajectory on the oligodendrocyte lineage (Zeisel level2class Oligo1-6)

KEY PLOTTING CALL

RECREATIONS/MONOCLE3/MONOCLE3_RECREATE.R — EXCERPT · ENDS LINE 8

```
cds <- new_cell_data_set(GetAssayData(ol, layer = "counts"), cell_metadata = ol@meta.data, gene_metadata = data.frame(gene_shor
t_name = rownames(ol), row.names = rownames(ol)))
cds <- preprocess_cds(cds, num_dim = 20); cds <- reduce_dimension(cds, preprocess_method = "PCA"); cds <- cluster_cells(cds, re
solution = 1e-2); cds <- learn_graph(cds, use_partition = FALSE)
root <- colnames(cds)[cds$level2class == "Oligo1"]; cds <- order_cells(cds, root_cells = root)
ggsave("v1_trajectory_subclass.png", plot_cells(cds, color_cells_by = "level2class", label_groups_by_cluster = FALSE, label_lea
ves = TRUE, label_branch_points = TRUE, group_label_size = 3, cell_size = .8), width = 6.5, height = 5, dpi = 150, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 — FIGURE CAPTION

Xenografted human microglia display diverse transcriptomic states in response to Alzheimer's disease-related amyloid-β pathology. *Nat Neurosci, 2024*

"...croglial transcriptional trajectory in response to Aβ pathology.a, Phenotypic trajectory followed by the human microglia after exposure to Aβ in vivo, obtained by an unbiased pseudotime ordering with Monocle ..."

DOI ↗ Fig. 3 on PMC ↗

Extended Data Fig. 4 — FIGURE CAPTION

Tau activation of microglial cGAS-IFN reduces MEF2C-mediated cognitive resilience. *Nat Neurosci, 2023*

"..., disease-associated (Apoe, Itgax) and interferon (Parp14, Stat1, Trim30a, Rnf213) genes in microglia clusters. b. Microglial trajectory plots colored by cell state trajectory scores identified using Monocle 3 ..."

DOI ↗ Article ↗ figure page not on PMC

Fig. 4 — TEXT

Human microglia show unique transcriptional changes in Alzheimer's disease. *Nat Aging, 2023*

"...The data were downsampled to 1,000, 2,000, 3,000 or 5,000 nuclei per cluster and transferred to a cell dataset object, and Monocle3 'learn_graph' was run."

DOI ↗ Fig. 4 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/monocle3/monocle3_recreate.R`

✓ verified run · exit 0 · 31 s

monocle3 · 4 variations



v1 trajectory_subclass

TRAJECTORY



v2 pseudotime

TRAJECTORY



v3 genes_in_pseudotime

TRAJECTORY



v4 genes_on_trajectory

TRAJECTORY

deepTools

#22

OF 100 · METHODS MENTIONS

PYTHON/CLI | HEATMAP (GENOMIC) | GENOME TRACKS & GENOMIC SIGNAL

416	2.34%	427	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

TSS/peak-centred signal heatmaps and metaprofiles (computeMatrix/plotHeatmap/plotProfile)

THIS RECREATION

deepTools computeMatrix → plotHeatmap / plotProfile. Uses SIMULATED bigWig signal (sharp TSS peaks ~H3K4me3, broad ~H3K27ac, flat input) on a toy 5-Mb chr

KEY PLOTTING CALL

RECREATIONS/DEEPTOOLS/DEEPTOOLS_RECREATE.SH — EXCERPT · ENDS LINE 6

```
python3 make_sim_bigwigs.py
computeMatrix reference-point -S sim_H3K4me3.bw sim_H3K27ac.bw sim_input.bw -R genes_active.bed genes_silent.bed --
referencePoint TSS -a 3000 -b 3000 -bs 50 -o mat.gz -p 4 --skipZeros
plotHeatmap -m mat.gz -out v1_plotHeatmap.png --colorMap Reds Blues Greys --heatmapHeight 10 --dpi 130 --plotTitle "deepTools
plotHeatmap (simulated signal)"
plotProfile -m mat.gz -out v2_plotProfile_perGroup.png --perGroup --dpi 130 --plotTitle "plotProfile --perGroup"
plotHeatmap -m mat.gz -out v3_kmeans3.png --kmeans 3 --colorMap viridis --dpi 130 --whatToShow "plot, heatmap and colorbar"
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure S1 TEXT
Macrophage-mediated myelin recycling fuels brain cancer malignancy. *Cell, 2024*
"...The coverage files were generated with the 'normalize to 1× genome coverage' methods in deepTools (version 3.5.1).89"
DOI ↗ Figure S1 on PMC ↗

Fig. 1 TEXT
Transcriptional and epigenetic decoding of the microglial aging process. *Nat Aging, 2023*
"...Heat maps of coverage at TSSs were generated using computeMatrix with the parameters computeMatrix reference-point–referencePoint TSS -b 1000 -a 1000 -R and plotHeatmap functions in deepTools 3.5.1."
DOI ↗ Fig. 1 on PMC ↗

Fig. 1 TEXT
Programs, origins and immunomodulatory functions of myeloid cells in glioma. *Nature, 2025*
"...Finally, we plotted the heatmap using the deeptools plotHeatmap function72."
DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

- Signal is SIMULATED (toy bigWigs); swap in real ENCODE bigWigs for production
- Uses SIMULATED bigWig signal (sharp TSS peaks ~H3K4me3, broad ~H3K27ac, flat input) on a toy 5-Mb chromosome. Swap in real ENCODE or GEO bigWigs for production figures (METHODS §4; script header).

Reproduce: `bash recreations/deepTools/deeptools_recreate.sh`

√ verified run · exit 0 · 225 s

deepTools · 4 variations



v1 plotHeatmap

GENOME



v2 plotProfile_perGroup

GENOME



v3 kmeans3

GENOME



v4 scale_regions_profile

GENOME

SingleR

#23

OF 100 · METHODS MENTIONS

R · ANNOTATION (HEATMAP) · HEATMAPS & DOT MATRICES · COMPOSITION, DISTRIBUTIONS & LAYOUTS · EMBEDDINGS & FEATURE MAPS

416	2.34%	451	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Annotation score heatmap, delta distribution, label UMAP

KEY PLOTTING CALL

RECREATIONS/SINGLER/SINGLER_RECREATE.R — EXCERPT · ENDS LINE 5

```
so <- readRDS("../_data/zeisel_seurat.rds"); ref <- MouseRNAseqData()
pred <- SingleR(test = GetAssayData(so, layer = "data"), ref = ref, labels = ref$label.main)
png("v1_plotScoreHeatmap.png", 1400, 900, res = 140); print(plotScoreHeatmap(pred, annotation_col = data.frame(Author = so$cell
type, row.names = colnames(so)))); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 5 — FIGURE CAPTION

Molecular and cellular evolution of the amygdala across species analyzed by single-nucleus transcriptome profiling. *Cell Discov, 2023*

"...sted P-value was calculated by the Wilcoxon test, two-side comparison (adjusted P-value < 0.05). g UMAP embedding of chicken datasets, with nuclei colored by predicted cell group ID from Seurat and SingleR ..."*

DOI ↗ Fig. 5 on PMC ↗

Fig. 1 — TEXT

Nociceptor-immune interactomes reveal insult-specific immune signatures of pain. *Nat Immunol, 2024*

"...Because a full dataset of immune cells in inflamed skin has not been generated before, we complemented the singleR predictions with specific cluster annotations based on current knowledge of skin immune ..."

DOI ↗ Fig. 1 on PMC ↗

Fig. 1 — TEXT

Spatial transcriptomics reveals human cortical layer and area specification. *Nature, 2025*

"...Reference-based cell-type annotation was conducted using SingleR (v.1.8.1)68 to transfer cell-type labels from the reference data to the testing data with the following parameters: "genes": "de","

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/SingleR/SingleR_recreate.R`

√ verified run · exit 0 · 26 s

SingleR · 4 variations



v1 plotScoreHeatmap

HEATMAPS



v2 plotDeltaDistribution

COMPOSITION



v3 confusion

HEATMAPS



v4 umap_labels

EMBEDDINGS

AUCell

#24

OF 100 · METHODS MENTIONS

R · GENE SET SCORES · COMPOSITION, DISTRIBUTIONS & LAYOUTS · EMBEDDINGS & FEATURE MAPS · HEATMAPS & DOT MATRICES

387	2.18%	432	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Gene-set activity on UMAP, AUC histograms/thresholds

KEY PLOTTING CALL

RECREATIONS/AUCELL/AUCELL_RECREATE.R — EXCERPT · ENDS LINE 8

```
rk <- AUCell_buildRankings(m, plotStats = FALSE); auc <- AUCell_calcAUC(sets, rk)
png("v1_exploreThresholds.png", 1500, 950, res = 140); par(mfrow = c(2,3)); th <- AUCell_exploreThresholds(auc, plotHist =
TRUE, assign = TRUE); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 2 — FIGURE CAPTION

APOE modulates microglial immunometabolism in response to age, amyloid pathology, and inflammatory challenge. *Cell Rep, 2023*

"...axis) and aged versus young microglia (x axis). Genes labeled in black are common to both DAM/MgND phenotypes. Inset: ridge plot showing DAM/MgND score for each individual microglia as calculated by AUCell.(B ..."

DOI ↗ Figure 2 on PMC ↗

Extended Data Fig. 1 — FIGURE CAPTION

SCENIC+: single-cell multiomic inference of enhancers and gene regulatory networks. *Nat Methods, 2023*

"...d. Cell-topic enrichment heat map for each of the models. General topics are shown in black; mesenchymal, in red; melanocytic, in blue; cell line specific in green; and low contributing in gray. e. AUCell ..."

DOI ↗ Extended Data Fig. 1 on PMC ↗

Figure 6 — TEXT

Polygenic regression uncovers trait-relevant cellular contexts through pathway activation transformation of single-cell RNA sequencing data. *Cell Genom, 2023*

"...Additionally, when applying genes prioritized by scPagwas to two other cell-scoring methods, e.g., VISION35 and AUCell,33 we consistently found that these cell-scoring methods yielded a high performance in ..."

DOI ↗ Figure 6 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/AUCell/AUCell_recreate.R`

√ verified run · exit 0 · 25 s

AUCell · 3 variations



v1 exploreThresholds

COMPOSITION



v2 umap_auc

EMBEDDINGS



v3 auc_heatmap

HEATMAPS

EnhancedVolcano

#27

OF 100 · METHODS MENTIONS

R · VOLCANO · DE, QC & SET OVERLAPS

369	2.08%	373	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Publication-ready volcano plots with labels & connectors

KEY PLOTTING CALL

RECREATIONS/ENHANCEDVOLCANO/ENHANCEDVOLCANO_RECREATE.R — EXCERPT · ENDS LINE 6

```
g <- EnhancedVolcano(res, lab = res$symbol, x = "log2FoldChange", y = "pvalue", title = "Dex vs untreated", subtitle = "Enhance
dVolcano default")
ggsave("v1_default.png", g, width = 7, height = 7, dpi = 130)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 5 TEXT

Multiomic spatial landscape of innate immune cells at human central nervous system borders. *Nat Med, 2024*

"...To this end we ran differential expression analysis of ligand–receptor pairs using the FindMarkers R function and visualized to top differentially expressed pairs using volcano plots with the EnhancedVolcano R ..."

DOI ↗ Fig. 5 on PMC ↗

Fig. 4 TEXT

Transcriptional and epigenetic decoding of the microglial aging process. *Nat Aging, 2023*

"...Venn diagrams, heat maps and volcano plots were visualized by using VennDiagram 1.6.20, pheatmap 1.0.12 and EnhancedVolcano 1.4.0, respectively."

DOI ↗ Fig. 4 on PMC ↗

Fig. 2 TEXT

Platelet-derived exerkine CXCL4/platelet factor 4 rejuvenates hippocampal neurogenesis and restores cognitive function in aged mice. *Nat Commun, 2023*

"...EnhancedVolcano v1.16.068 produced volcano plots, with shrunken log fold change values calculated using "ashr" as an estimator."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/EnhancedVolcano/EnhancedVolcano_recreate.R`

√ verified run · exit 0 · 10 s

LEAD FIGURE



v2 selected_boxed

DE

EnhancedVolcano · 3 variations · v2 on facing page



v1 default

DE



v3 custom_colours

DE

survminer

#29

OF 100 · METHODS MENTIONS

R | SURVIVAL | COMPOSITION, DISTRIBUTIONS & LAYOUTS

351	1.98%	342	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Kaplan-Meier curves + risk tables, forest plots

THIS RECREATION

survminer: Kaplan-Meier + risk table; glioma survival data from coin::glioma if available, else survival::lung

KEY PLOTTING CALL

RECREATIONS/SURVMINER/SURVMINER_RECREATE.R — EXCERPT · ENDS LINE 7

```
if (requireNamespace("coin", quietly = TRUE)) { data("glioma", package = "coin"); d <- glioma; d$status <- as.numeric(d$event);
f <- Surv(time, status) ~ group; ttl <- "coin::glioma (Grana 2002): radioimmunotherapy vs control" } else { d <- lung; f <-
Surv(time, status) ~ sex; ttl <- "survival::lung" }
fit <- surv_fit(f, data = d)
p <- ggsurvplot(fit, data = d, pval = TRUE, conf.int = TRUE, risk.table = TRUE, palette = c("#2E9FDF","#E7B800"), title = ttl, g
gtheme = theme_classic())
png("v1_km_risktable.png", 1100, 1000, res = 150); print(p); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 7 TEXT
Macrophage-mediated myelin recycling fuels brain cancer malignancy. *Cell, 2024*
"...For the calculation of the survival curves in Figures 7A and 7F, the Kaplan-Meier estimator was obtained with the survfit function from the package survival.106 The statistical test applied was the log-rank ..."
DOI ↗ Figure 7 on PMC ↗

Fig. 3 TEXT
Spatial transcriptomics reveals niche-specific enrichment and vulnerabilities of radial glial stem-like cells in malignant gliomas. *Nat Commun, 2023*
"...And the survival curve was plotted using survminer (v0.4.9) R package."
DOI ↗ Fig. 3 on PMC ↗

Fig. 2 TEXT
Deciphering the longitudinal trajectories of glioblastoma ecosystems by integrative single-cell genomics. *Nat Genet, 2025*
"...Kaplan–Meier estimates were computed using the survfit function and plotted using the ggsurvplot function (survminer package)."
DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

- Uses coin::glioma (Grana et al. 2002, glioma radioimmunotherapy) survival data

Reproduce: `Rscript recreations/survminer/survminer_recreate.R`

√ verified run · exit 0 · 4 s

survminer · 3 variations



v1 km_risktable

COMPOSITION



v2 cumhaz

COMPOSITION



v3 forest

COMPOSITION

Space Ranger/ Loupe

#31

OF 100 · METHODS MENTIONS

DESKTOP | SPATIAL | SPATIAL & TISSUE MAPS

300	1.69%	307	viewer	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Visium spot/tissue overlays (Loupe)

Also catalogued as: **Loupe Browser** (#43, 166 papers)

THIS RECREATION

Space Ranger / Loupe Browser look-alike from the same Visium mouse-brain run: tissue + spot overlay, gene with opacity, side-by-side UMAP

KEY PLOTTING CALL

RECREATIONS/SPACERANGER_LOUPE_STYLE/LOUPE_STYLE_RECREATE.PY — EXCERPT · ENDS LINE 15

```
fig,ax=plt.subplots(1,2,figsize=(13,6.2),facecolor="#f4f4f4")
ax[0].imshow(img); spots(ax[0],col,.85); ax[0].set_title("Clusters (graph-based)",loc="left",fontsize=11); ax[0].axis("off")
u=v.obsm["X_umap"]; ax[1].scatter(*u.T,c=col,s=8); ax[1].set_title("UMAP",loc="left"); ax[1].axis("off")
for i,c in enumerate(cl.cat.categories): ax[1].scatter([],[],c=LOUPE[i%15],label=c)
ax[1].legend(fontsize=7,frameon=False,loc="center left",bbox_to_anchor=(1,.5)); fig.suptitle("Loupe-style linked tissue / UMAP
view",x=.02,ha="left",weight="bold"); fig.tight_layout(); fig.savefig("v1_tissue_clusters_umap.png",dpi=130)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT — **Spatial transcriptomics reveals niche-specific enrichment and vulnerabilities of radial glial stem-like cells in malignant gliomas.** *Nat Commun, 2023*

"...We used REDItools60 (v2) with default parameters to identify H3K27M mutation in DMG samples from BAM files generated by the SpaceRanger pipeline."

DOI ↗ Fig. 1 on PMC ↗

Fig. 1 TEXT — **Spatially resolved multiomics of human cardiac niches.** *Nature, 2023*

"...Visium: GRCh38-3.0.0) provided by 10x Genomics using CellRanger ARC (v.2.0.0) or SpaceRanger (v.1.1.0) with default parameters."

DOI ↗ Fig. 1 on PMC ↗

Fig. 1 TEXT — **Robust mapping of spatiotemporal trajectories and cell-cell interactions in healthy and diseased tissues.** *Nat Commun, 2023*

"...Raw BCL files were processed by bcl2fastq V2.7.0, and the fastq reads mapped to the mouse reference genome GRCm38 by SpaceRanger V1.0.0."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

- Look-alike with matplotlib on the squidpy Visium mouse-brain run
- Loupe Browser (#43): Look-alike (see Space Ranger/Loupe)

Reproduce: `python3 recreations/SpaceRanger_Loupe_style/loupe_style_recreate.py`

√ verified run · exit 0 · 5 s

Space Ranger/Loupe · 3 variations

RECREATIONS/SPACERANGER_LOUPE_STYLE/



v1 tissue_clusters_umap

SPATIAL



v2 feature_opacity

SPATIAL



v3 umi_per_spot

SPATIAL

ggpubr

R | GENERAL/STATISTICS | COMPOSITION, DISTRIBUTIONS & LAYOUTS

#33

OF 100 · METHODS MENTIONS

277	1.56%	283	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Box/violin/bar plots with p-values, correlation scatter

KEY PLOTTING CALL

RECREATIONS/GGPUBR/GGPUBR_RECREATE.R — EXCERPT · ENDS LINE 6

```
so <- readRDS("../_data/zeisel_seurat.rds"); df <- data.frame(celltype = so$celltype, Gad1 = FetchData(so,"Gad1")[,1], Snap25 =
FetchData(so,"Snap25")[,1], Mbp = FetchData(so,"Mbp")[,1], nGene = so$nFeature_RNA, tissue = so$tissue)
sub <- droplevels(df[df$celltype %in% c("interneurons","pyramidal SS","pyramidal CA1"),])
p <- ggboxplot(sub, "celltype", "Gad1", color = "celltype", add = "jitter", add.params = list(size = .3, alpha = .3), palette =
"jco") + stat_compare_means(comparisons = list(c("interneurons","pyramidal SS"), c("interneurons","pyramidal CA1"),
c("pyramidal SS","pyramidal CA1")), label = "p.signif") + stat_compare_means(label.y = 7.5)
ggsave("v1_boxplot_pvalues.png", p, width = 5.5, height = 5, dpi = 150, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure S1 TEXT
Macrophage-mediated myelin recycling fuels brain cancer malignancy. *Cell, 2024*
"...Plots were generated using the package ggplot (v.3.3.6) or ggpubr (v.0.4.0), except where mentioned."
DOI ↗ Figure S1 on PMC ↗

Fig. 3 TEXT
A concerted neuron-astrocyte program declines in ageing and schizophrenia. *Nature, 2024*
"...R (v.4.1.3): cluster (v.2.1.2)138, ComplexHeatmap (v.2.10.0)139,140, data.table (v.1.14.8)141, DescTools (v.0.99.48)142, dplyr (v.1.1.2)143, gdata (v.2.19.0)144, ggforce (v.0.4.1)145, ggplot2 (v.3.4.2)146 ..."
DOI ↗ Fig. 3 on PMC ↗

Figure 4 TEXT
Transcriptomic atlas and interaction networks of brain cells in mouse CNS demyelination and remyelination. *Cell Rep, 2023*
"...Statistics on cell type compositional analysis were calculated using scCODA package (v.0.1.8) in Python (v3.9.2).101 p values in snRNA-seq analysis were adjusted based on Bonferroni correction.106 p values in ..."
DOI ↗ Figure 4 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggpubr/ggpubr_recreate.R`

√ verified run · exit 0 · 8 s

ggpubr · 4 variations



v1 boxplot_pvalues

COMPOSITION



v2 violin

COMPOSITION



v3 scatter_cor

COMPOSITION



v4 barplot_mean_se

COMPOSITION

igraph

#36

OF 100 · METHODS MENTIONS

R/PYTHON | NETWORK | NETWORKS, GRAPHS & TREES

241	1.36%	240	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Network graphs (co-expression, PPI, communities)

THIS RECREATION

igraph: gene co-expression network of variable genes in Zeisel cortex data

KEY PLOTTING CALL

RECREATIONS/IGRAPH/IGRAPH_RECREATE.R — EXCERPT · ENDS LINE 8

```
so <- readRDS("../_data/zeisel_seurat.rds"); g0 <- head(VariableFeatures(so), 150)
e <- as.matrix(GetAssayData(so, layer = "data")[g0,]); cm <- cor(t(e)); diag(cm) <- 0
adj <- (cm > .45) * cm; g <- graph_from_adjacency_matrix(adj, mode = "undirected", weighted = TRUE); g <- delete_vertices(g, de
gree(g) == 0)
set.seed(2); com <- cluster_louvain(g); pal <- hcl.colors(max(membership(com)), "Dark 3")
png("v1_fr_communities.png", 1300, 1300, res = 160)
plot(g, layout = layout_with_fr(g), vertex.color = pal[membership(com)], vertex.size = 3 + 1.2*sqrt(degree(g)),
vertex.label.cex = .5, vertex.label.color = "black", edge.width = E(g)$weight*2, edge.color = "grey80", main = "Co-expression
network (r > 0.45), Louvain communities"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 TEXT — **Inferring and perturbing cell fate regulomes in human brain organoids.** *Nature, 2023*

"...We then performed 10,000 random walks with 200 steps from each tip along edges backwards in pseudotime using the igraph R package (v.1.2.6) (https://igraph.org/)."

DOI ↗ Fig. 2 on PMC ↗

Fig. 3 TEXT — **Multimodal spatiotemporal phenotyping of human retinal organoid development.** *Nat Biotechnol, 2023*

"...The cluster graph was then visualized using the ggraph package in R with the tree layout implemented in the igraph package (the layout_as_tree function), with the edge weights represented by the edge ..."

DOI ↗ Fig. 3 on PMC ↗

Fig. 2 TEXT — **A prognostic neural epigenetic signature in high-grade glioma.** *Nat Med, 2024*

"...The adjacency matrix was then transformed into an undirected graph using the graph_from_adjacency_matrix function from the igraph package."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/igraph/igraph_recreate.R`

√ verified run · exit 0 · 8 s

igraph · 3 variations



v1 fr_communities

NETWORKS



v2 mark_groups

NETWORKS



v3 circle_hubs

NETWORKS

Signac

#37

OF 100 · METHODS MENTIONS

R | SCATAC (COVERAGE/EMBEDDING) | GENOME TRACKS & GENOMIC SIGNAL

227	1.28%	239	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · SIGNAC

Neuro-corpus mentions per year (*2026 partial) · niche rank **#54**, growth **1.43×**, visual 4/5, score 0.544. Broad-corpus rate 4.876 → 6.96 per 1k papers (2023 → 2025–26).
GitHub stuart-lab/signac · created 2019-05-09 · last push 2026-09-11

BEST FOR

Coverage plots with peaks/links, motif plots, ATAC UMAP

THIS RECREATION

Signac scATAC-seq coverage/track plots. NOTE: uses Signac's bundled 'atac_small' (human PBMC, 100 cells) and its fragments file - no offline neuro scATAC small enough was available (10x E18 mouse brain fragments are >1 GB).

KEY PLOTTING CALL

RECREATIONS/SIGNAC/SIGNAC_RECREATE.R — EXCERPT · ENDS LINE 11

```
data(atac_small); fpath <- system.file("extdata", "fragments.tsv.gz", package = "Signac")
fragments <- CreateFragmentObject(fpath, cells = colnames(atac_small), validate.fragments = FALSE); Fragments(atac_small) <- NU
LL; Fragments(atac_small) <- fragments
atac_small$cluster <- factor(paste0("C", as.integer(atac_small$seurat_clusters)))
Idents(atac_small) <- "cluster"
reg <- "chr1-713500-860000"
p1 <- CoveragePlot(atac_small, region = reg, annotation = TRUE, peaks = TRUE, links = FALSE, extend.upstream = 0, extend.downst
ream = 0)
ggsave("v1_coverage_plot.png", p1 & theme(text = element_text(size = 9)), width = 9, height = 6, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 — FIGURE CAPTION

SCENIC+: single-cell multiomic inference of enhancers and gene regulatory networks. *Nat Methods, 2023*

"...r parameter optimization. c, Bar-plots showing the area under the recovery curve (AUC; enhancer recovery) on the top 10% of the ranking based on STARR-seq signal, for the top 5,000 DARs identified by Signac ..."

DOI ↗ Fig. 1 on PMC ↗

Fig. 4 — TEXT

Slide-tags enables single-nucleus barcoding for multimodal spatial genomics. *Nature, 2024*

"...The ATAC-filtered feature-barcode matrix generated by Cell Ranger was read into R (v.4.1.1) using Signac (v.1.9.0)77, and added as its own assay slot in the Seurat object containing RNA expression counts."

DOI ↗ Fig. 4 on PMC ↗

Fig. 3 — TEXT

Multimodal chromatin profiling using nanobody-based single-cell CUT&Tag. *Nat Biotechnol, 2023*

"...All dimensionality reduction and clustering was performed using package Seurat v4.0.5 and Signac v1.4.0 under R v4.1.1."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

- Non-neuro fallback: Signac's bundled atac_small (human PBMC, 100 cells) and its fragments file; tracks look thin. 10x E18 mouse-brain fragments are >1 GB, too large to use offline (SUMMARY; script header).

Reproduce: `Rscript recreations/Signac/Signac_recreate.R`

✓ verified run · exit 0 · 11 s

Signac · 3 variations



v1 coverage_plot

GENOME



v2 tile_plot

GENOME



v3 fraghist_umap

GENOME

matplotlib

#38

OF 100 · METHODS MENTIONS

PYTHON | GENERAL | DE, QC & SET OVERLAPS | EMBEDDINGS & FEATURE MAPS | COMPOSITION, DISTRIBUTIONS & LAYOUTS

227	1.28%	238	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Base for all Python plots: scatter, volcano, multi-panel GridSpec

THIS RECREATION

pure matplotlib versions of the most common omics panels

KEY PLOTTING CALL

RECREATIONS/MATPLOTLIB/MATPLOTLIB_RECREATE.PY — EXCERPT · ENDS LINE 11

```
fig,ax=plt.subplots(figsize=(5.5,5)); y=-np.log10(res.padj.clip(lower=1e-300))
ax.scatter(res.log2FoldChange[~(up|dn)],y[~(up|dn)],s=3,c="lightgrey",rasterized=True);
ax.scatter(res.log2FoldChange[up],y[up],s=4,c="#d7301f"); ax.scatter(res.log2FoldChange[dn],y[dn],s=4,c="#2b8cbe")
for g,r in res[up|dn].nsmallest(10,"padj").iterrows(): ax.annotate(r.symbol,(r.log2FoldChange,-np.log10(r.padj)),fontsize=7)
ax.axvline(-1,ls="--",c="k",lw=.6); ax.axvline(1,ls="--",c="k",lw=.6); ax.set(xlabel="log2 fold change",ylabel="-log10 padj",ti
tle="Volcano (matplotlib)"); [ax.spines[s].set_visible(False) for s in ("top","right")]
fig.tight_layout(); fig.savefig("v1_volcano.png",dpi=150)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 TEXT
Specialized astrocytes mediate glutamatergic gliotransmission in the CNS. *Nature, 2023*
"...The turbo colormap from matplotlib was used to assign amplitude values."
DOI ↗ Fig. 2 on PMC ↗

Fig. 3 TEXT
SCENIC+: single-cell multiomic inference of enhancers and gene regulatory networks. *Nat Methods, 2023*
"...The delta embedding is used to draw arrows on the original embedding, using the streamplot or quiverplot function from the matplotlib (v.3.5.2) Python package."
DOI ↗ Fig. 3 on PMC ↗

Fig. 3 TEXT
A concerted neuron-astrocyte program declines in ageing and schizophrenia. *Nature, 2024*
"...Python (v.3.8.3): matplotlib (v.3.5.2)136 and seaborn (v.0.10.1)137."
DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/matplotlib/matplotlib_recreate.py`

√ verified run · exit 0 · 2 s

matplotlib · 4 variations



v1 volcano

DE



v2 MA

DE



v3 umap

EMBEDDINGS



v4 multipanel_gridspec

COMPOSITION

CopyKAT/ inferCNV

#39

OF 100 · METHODS MENTIONS

R · CNV HEATMAP · HEATMAPS & DOT MATRICES

207	1.17%	236	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · INFERCNV/COPYKAT

Neuro-corpus mentions per year (*2026 partial) · niche rank **#40**, growth **1.91×**, visual 4/5, score 0.622. Broad-corpus rate 10.97 → 20.93 per 1k papers (2023 → 2025–26).
GitHub broadinstitute/infercnv · created 2016-03-14 · last push 2025-11-14

BEST FOR

Inferred CNV heatmaps by chromosome

THIS RECREATION

inferCNV on the package's downsampled oligodendroglioma scRNA-seq example (Tirosh et al. 2016, Nature) - a real brain-tumour dataset

KEY PLOTTING CALL

RECREATIONS/INFERCNV/INFERCNV_RECREATE.R — EXCERPT · ENDS LINE 6

```
o <- infercnv::run(mk(), cutoff = 1, out_dir = "out_default", cluster_by_groups = TRUE, denoise = TRUE, HMM = FALSE, no_prelim_
plot = TRUE, num_threads = 2, png_res = 150, write_expr_matrix = FALSE)
file.copy("out_default/infercnv.png", "v1_cnv_heatmap_by_group.png", overwrite = TRUE)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 — FIGURE CAPTION

A variational algorithm to detect the clonal copy number substructure of tumors from scRNA-seq data. *Nat Commun, 2023*

"...Benchmark of inferred copy number profile.a, b Copy number profile inferred with SCEVAN (segment mean (LogRatio) and CNV status), inferCNV, CopyKAT, the corresponding ground truth from low-depth WGS of sample ..."

DOI ↗ Fig. 3 on PMC ↗

Extended Data Fig. 4 — FIGURE CAPTION

Proteomic signatures improve risk prediction for common and rare diseases. *Nat Med, 2024*

"...ity and distribution is illustrated by color. c, BM UMAP from (A) highlighting the plasma cell state healthy (green), malignant (red) and unclear (yellow) based on copy number aberrations detected by ..."

DOI ↗ Extended Data Fig. 4 on PMC ↗

Fig. 1 — TEXT

Inferring histology-associated gene expression gradients in spatial transcriptomic studies. *Nat Commun, 2024*

"...The output of infercnv::plot_cnv() is stored under a specified directory from which the results are read back into the R session and stored together with metadata regarding the analysis in slot @cnv of the ..."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/inferCNV/inferCNV_recreate.R`

√ verified run · exit 0 · 23 s

CopyKAT/inferCNV · 3 variations



v1 cnv_heatmap_by_group

HEATMAPS



v2 cnv_heatmap_subclusters

HEATMAPS



v3 cnv_per_group

HEATMAPS

seaborn

#40

OF 100 · METHODS MENTIONS

PYTHON | GENERAL/STATISTICS | HEATMAPS & DOT MATRICES | COMPOSITION, DISTRIBUTIONS & LAYOUTS

192	1.08%	193	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

clustermap, violin/box/strip, kde, faceted catplots

KEY PLOTTING CALL

RECREATIONS/SEABORN/SEABORN_RECREATE.PY — EXCERPT · ENDS LINE 6

```
ad=load(); genes=["Snap25","Gad1","Pvalb","Sst","Satb2","Neurod6","Aqp4","Gfap","Mbp","Plp1","Cx3cr1","C1qa","Cldn5","Acta2"]
df=pd.DataFrame(ad[:,genes].X.toarray(),columns=genes,index=ad.obs_names); df["celltype"]=ad.obs.celltype.values
av=df.groupby("celltype",observed=True).mean()
g=sns.clustermap(av.T,z_score=0,cmap="vlag",figsize=(6,7),row_cluster=True); g.fig.suptitle("seaborn clustermap (z-scored
means)",y=1.02); g.savefig("v1_clustermap.png",dpi=150)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 5 TEXT — **Functional synapses between neurons and small cell lung cancer.** *Nature, 2025*

"...The packages matplotlib v3.4.2 and seaborn 0.11.0 were used for visualization."

DOI ↗ Fig. 5 on PMC ↗

Fig. 3 TEXT — **A concerted neuron-astrocyte program declines in ageing and schizophrenia.** *Nature, 2024*

"...Python (v.3.8.3): matplotlib (v.3.5.2)136 and seaborn (v.0.10.1)137."

DOI ↗ Fig. 3 on PMC ↗

Fig. 2 TEXT — **An organotypic atlas of human vascular cells.** *Nat Med, 2024*

"...Python (version 3), AnnData (versions 0.8.0 and 0.9.1) Pandas (versions 1.5.2 and 1.5.3), NumPy (versions 1.21.6 and 1.23.5), SciPy (versions 1.8.0 and 1.9.3), Matplotlib (version 3.5.2), Seaborn (version ..."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/seaborn/seaborn_recreate.py`

√ verified run · exit 0 · 8 s

seaborn · 4 variations



v1 clustermap — HEATMAPS



v2 violin — COMPOSITION



v3 catplot_facets — COMPOSITION



v4 kde_qc — COMPOSITION

CellPhoneDB

#41

OF 100 · METHODS MENTIONS

PYTHON | CELL-CELL COMMUNICATION | CELL–CELL COMMUNICATION

182	1.02%	231	analysis+plotting	2
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · CELLPHONEDB/KTPLOTS

Neuro-corpus mentions per year (*2026 partial) · niche rank **#62**, growth **1.1×**, visual 4/5, score 0.476. Broad-corpus rate 12.746 → 14.021 per 1k papers (2023 → 2025–26).
GitHub zktuong/ktplots · created 2019-10-11 · last push 2025-05-17

BEST FOR

L-R dot plots / heatmaps of interaction counts

THIS RECREATION

CellPhoneDB-method results (via LIANA+) rendered with ktplotspy (the ktplots Python port): CPDB dot plot + significance heatmap

KEY PLOTTING CALL

RECREATIONS/CELLPHONEDB_KTPLOTS/KTPLOTS_RECREATE.PY — EXCERPT · ENDS LINE 14

```
means,pvals=cp(mw),cp(pw)
ad=load(); ad.obs["celltype"]=ad.obs.celltype.astype(str).astype("category")
g=kpy.plot_cpdb_heatmap(pvals=pvals,figsize=(6,6),title="# significant interactions (CellPhoneDB method)",symmetrical=False); p
lt.savefig("v1_cpdb_heatmap.png",dpi=140,bbox_inches="tight"); plt.close("all")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 6 FIGURE CAPTION

The Single-Cell Landscape of Intratumoral Heterogeneity and The Immunosuppressive Microenvironment in Liver and Brain Metastases of Breast Cancer. (Adv Sci (Weinh), 2023). ,

"...th factor genes (VEGFA, VEGFB, HGF, and IGF1) in stromal cell subtypes. m) Dot plot showing the ligand–receptor pairs of cell growth factors between cancer cells and each CAF cluster predicted by the ..."

DOI ↗ Figure 6 on PMC ↗

Fig. 1 FIGURE CAPTION

Spatially resolved multiomics of human cardiac niches. *Nature, 2023*

"...ed by cell2location. Dashed lines highlight SAN, AVN, AVB and Purkinje cells defined by histology (Extended Data Fig. 2g). Illustrations in a were created using BioRender (https://biorender.com). The ..."

DOI ↗ Fig. 1 on PMC ↗

Fig. 3 FIGURE CAPTION

Inferring neuron-neuron communications from single-cell transcriptomics through NeuronChat. *Nat Commun, 2023*

"...Comparison of NeuronChat, CellPhoneDB, and CellChat in predicting VISp and ALM projection networks.a Typical ROC curves (left panel) and PR curves for the three methods. b The boxplots of AUROC (left panel) ..."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

- CellPhoneDB method via LIANA+, plotted with ktplotspy
- Reads LIANA+ result tables, so run recreations/LIANA_plus first (METHODS §9).

Reproduce: `python3 recreations/CellPhoneDB_ktplots/ktplots_recreate.py`

√ verified run · exit 0 · 10 s

CellPhoneDB · 2 variations



v1 cpdb_heatmap

CELL-CELL



v2 cpdb_dotplot

CELL-CELL

slingshot

#42

OF 100 · METHODS MENTIONS

R | TRAJECTORY | TRAJECTORY, PSEUDOTIME & VELOCITY

173	0.97%	225	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Lineage curves over embeddings, pseudotime

THIS RECREATION

slingshot lineages on La Manno 2018 dentate gyrus (granule + glial branches) on UMAP: principal curves over clusters, pseudotime colouring per lineage, and the cluster MST.

KEY PLOTTING CALL

RECREATIONS/SLINGSHOT/SLINGSHOT_RECREATE.R — EXCERPT · ENDS LINE 12

```
png("v1_curves_on_umap.png", 1300, 1100, res = 150); plot(um, col = pal[cl], pch = 16, cex = .5, asp = 1, xlab = "UMAP1", ylab
= "UMAP2", main = "slingshot principal curves (start: radial glia-like)")
lines(SlingshotDataSet(sce), lwd = 3, col = "black"); legend("topright", legend = levels(cl), col = pal, pch = 16, cex = .7,
bty = "n"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 — FIGURE CAPTION

A topographic atlas defines developmental origins of cell heterogeneity in the human embryonic lung. *Nat Cell Biol, 2023*

"...re is shown by H&E staining. Scale bar, 400 μm. arw, airway; tr, trachea; prox, proximal; pcw, post conception week; br-v bundle, bronchovascular bundle. c, Pseudotime analysis of the ASM cells, with Slingshot ..."

DOI ↗ Fig. 2 on PMC ↗

Fig. 1 — FIGURE CAPTION

Single-cell RNA-sequencing identifies disease-associated oligodendrocytes in male APP NL-G-F and 5XFAD mice. *Nat Commun, 2023*

"...tion of oligodendrocytes from 1-, 3, 6-month-old WT, and AppNL-G-F mice. e Two lineages of oligodendrocytes with a cluster-based minimum spanning tree with line. f Continuous trajectory projection by ..."

DOI ↗ Fig. 1 on PMC ↗

Extended Data Fig. 5 — FIGURE CAPTION

Defining blood-induced microglia functions in neurodegeneration through multiomic profiling. *Nat Immunol, 2023*

"...tory analysis of scRNA-seq BMDM profiles.a, UMAP plots of BMDM scRNA-seq profiles colored by treatment (left) or cluster (right) as shown in Fig. 3a, overlaid with pseudotime trajectory analysis with ..."

DOI ↗ Extended Data Fig. 5 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/slingshot/slingshot_recreate.R`

√ verified run · exit 0 · 8 s

slingshot · 3 variations



v1 curves_on_umap

TRAJECTORY



v2 pseudotime_per_lineage

TRAJECTORY



v3 cluster_MST

TRAJECTORY

rrvgo

#44

OF 100 · METHODS MENTIONS

R · ENRICHMENT (GO REDUCE) · ENRICHMENT & PATHWAY ACTIVITY

163	0.92%	179	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

GO semantic similarity treemaps / scatter

THIS RECREATION

rrvgo: reduce redundant GO BP terms from Zeisel 2015 astrocyte markers (treemap, scatter, wordcloud-style heatmap)

KEY PLOTTING CALL

RECREATIONS/RRVGO/RRVGO_RECREATE.R — EXCERPT · ENDS LINE 8

```
m <- read.csv("../_data/zeisel_markers.csv") %>% filter(cluster == "astrocytes_ependymal", p_val_adj < .01) %>% slice_max(avg_l
og2FC, n = 300)
ego <- as.data.frame(enrichGO(m$gene, OrgDb = org.Mm.eg.db, keyType = "SYMBOL", ont = "BP", pvalueCutoff = .01))
sim <- calculateSimMatrix(ego$ID, orgdb = "org.Mm.eg.db", ont = "BP", method = "Rel")
sc <- setNames(-log10(ego$p.adjust), ego$ID); red <- reduceSimMatrix(sim, sc, threshold = .7, orgdb = "org.Mm.eg.db")
png("v1_treemap.png", 1600, 1100, res = 150); treemapPlot(red, title = "rrvgo treemap: astrocyte marker GO BP (parents)"); dev.
off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 FIGURE CAPTION — **Copy number variation in tRNA isodecoder genes impairs mammalian development and balanced translation.** *Nat Commun, 2023*

"...s in brains a, and livers b, from tRNA-Phe-1-1-/- knockout mice (n = 3) compared to controls (n = 3), summarised by biological process gene ontologies (GOs) determined by PANTHER and visualised using REVIGO ..."

DOI ↗ Fig. 3 on PMC ↗

Fig. 3 TEXT — **iPSC-derived models of PACS1 syndrome reveal transcriptional and functional deficits in neuron activity.** *Nat Commun, 2024*

"...To more wholistically visualize GO results, the ontology hierarchy was extracted and reduced using the rrvgo package (v1.9.1)93 with the score of the term being -log10(pelim)."

DOI ↗ Fig. 3 on PMC ↗

Fig. 3 TEXT — **Single-nucleus RNA-sequencing of autosomal dominant Alzheimer disease and risk variant carriers.** *Nat Commun, 2023*

"...The terms were then loaded into rrvgo (v1.6.0), an R package that implements the Revigo94 tool for summarizing GO_BP terms."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/rrvgo/rrvgo_recreate.R`

✓ verified run · exit 0 · 447 s

rrvgo · 3 variations

RECREATIONS/RRVGO/



v1 treemap

ENRICHMENT



v2 scatter

ENRICHMENT



v3 heatmap

ENRICHMENT

scVelo

#45

OF 100 · METHODS MENTIONS

PYTHON | RNA VELOCITY | TRAJECTORY, PSEUDOTIME & VELOCITY

161	0.91%	177	analysis+plotting	5
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · SCVELO STREAMLINES

Neuro-corpus mentions per year (*2026 partial) · niche rank **#45**, growth **1.15×**, visual 5/5, score 0.596. Broad-corpus rate 7.244 → 8.357 per 1k papers (2023 → 2025–26).
GitHub theislab/scvelo · created 2018-08-20 · last push 2026-02-25

BEST FOR

RNA-velocity stream/arrow embeddings, phase portraits

Also catalogued as: **velocyto** (#56, 120 papers)

THIS RECREATION

scVelo on La Manno 2018 mouse dentate gyrus: streamlines, velocyto-style grid arrows, phase portraits, latent-time heatmap

KEY PLOTTING CALL

RECREATIONS/SCVELO/SCVELO_RECREATE.PY — EXCERPT · ENDS LINE 6

```
scv.pl.velocity_embedding_stream(a,basis="umap",color="clusters",legend_loc="right margin",title="Velocity streamlines (dynamical)",save="v1_stream.png",show=False)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 7 — FIGURE CAPTION

Multimodal chromatin profiling using nanobody-based single-cell CUT&Tag. *Nat Biotechnol, 2023*

"…ty visualization. The chromatin velocity was calculated by using ATAC-seq gene-by-cell matrix as input into the unspliced layer and H3K27ac gene-by-cell matrix into the spliced layer and then running scvelo …"

DOI ↗ Fig. 7 on PMC ↗

Figure 2 — FIGURE CAPTION

Distinct molecular profiles of skull bone marrow in health and neurological disorders. *Cell, 2023*

"…tions, p = 0)(H) Left: in the neutrophil subpopulation, calvaria, and dura neutrophils are highlighted in region-based UMAP. Right: projected developmental trajectory of MCAo neutrophils subset using …"

DOI ↗ Figure 2 on PMC ↗

Fig. 6 — FIGURE CAPTION

SCENIC+: single-cell multiomic inference of enhancers and gene regulatory networks. *Nat Methods, 2023*

"…5 cells). c–e, UMAP dimensionality reduction of 3,104 pseudocells from the fly eye based on target gene and region enrichment scores, with a schematic representation of the fly eye-antennal disc (c), scVelo …"

DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

- velocyto (#56): velocyto-style grid arrows via scVelo (v2_grid_arrows)

Reproduce: `python3 recreations/scVelo/scvelo_recreate.py`

√ verified run · exit 0 · 6 s

scVelo · 5 variations



v1 stream

TRAJECTORY



v2 grid_arrows

TRAJECTORY



v3 phase_portraits

TRAJECTORY



v4 latent_time_heatmap

TRAJECTORY



v5 latent_time

TRAJECTORY

VennDiagram

#46

OF 100 · METHODS MENTIONS

R | SET OVERLAP | DE, QC & SET OVERLAPS

157	0.88%	160	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

2-5 set Venn diagrams

Also catalogued as: **ggvenn** (#82, 61 papers)

THIS RECREATION

VennDiagram / ggvenn / ggVennDiagram: marker overlaps between Zeisel 2015 glial cell types

KEY PLOTTING CALL

RECREATIONS/VENNDIAGRAM_GGVENN/VENN_RECREATE.R — EXCERPT · ENDS LINE 8

```
futile.logger::flog.threshold(futile.logger::ERROR, name = "VennDiagramLogger")
m <- read.csv("../_data/zeisel_markers.csv") %>% filter(p_val_adj < .01, avg_log2FC > 1); L <- split(m$gene, m$cluster)
g4 <- L[c("astrocytes_ependymal","oligodendrocytes","microglia","endothelial-mural")]
v <- venn.diagram(g4, filename = NULL, fill = c("#66c2a5","#fc8d62","#8da0cb","#e78ac3"), alpha = .5, cex = 1.1, cat.cex = 1, m
ain = "VennDiagram: glial/vascular marker overlap", category.names = c("astro","oligo","microglia","endo-mural"))
png("v1_VennDiagram_4set.png", 1200, 1200, res = 170); grid.draw(v); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT

Transcriptional and epigenetic decoding of the microglial aging process. *Nat Aging, 2023*

"...Venn diagrams, heat maps and volcano plots were visualized by using VennDiagram 1.6.20, pheatmap 1.0.12 and EnhancedVolcano 1.4.0, respectively."

DOI ↗ Fig. 1 on PMC ↗

Fig. 4 TEXT

Microglia mediate neurocognitive deficits by eliminating C1q-tagged synapses in sepsis-associated encephalopathy. *Sci Adv, 2023*

"...Venn diagrams showing the overlap of DEGs in different treatment groups were calculated and plotted with the VennDiagram package."

DOI ↗ Fig. 4 on PMC ↗

Fig. 4 TEXT

Broad rim lesions are a new pathological and imaging biomarker for rapid disease progression in multiple sclerosis. *Nat Med, 2025*

"...Any overlap of DEGs between lesion types was visualized as a Venn diagram with the R package VennDiagram v.1.7.3 (ref."

DOI ↗ Fig. 4 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/VennDiagram_ggvenn/Venn_recreate.R`

√ verified run · exit 0 · 2 s

VennDiagram · 3 variations



v1 VennDiagram_4set

DE



v2 ggvenn_3set

DE



v3 ggVennDiagram_gradient

DE

gplots (heatmap.2)

#47

OF 100 · METHODS MENTIONS

R · HEATMAP · HEATMAPS & DOT MATRICES

152	0.86%	157	dedicated plotting	2
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Classic heatmap.2 with dendrograms, Venn

THIS RECREATION

gplots::heatmap.2 (classic: row z-score, dendrograms, colour key + histogram, side colours) on Zeisel 2015 markers

KEY PLOTTING CALL

RECREATIONS/GPLOTS_HEATMAP2/HEATMAP2_RECREATE.R — EXCERPT · ENDS LINE 8

```
png("v1_heatmap2_rowz.png", 1500, 1500, res = 150); heatmap.2(M, scale = "row", trace = "none", col = colorRampPalette(rev(brew
er.pal(11, "RdBu")))(100), ColSideColors = pal[ct], labCol = FALSE, density.info = "histogram", key.title = NA, margins = c(3,
8), cexRow = .6, main = "heatmap.2: markers x (cell type|chip) pseudobulk"); legend("topright", legend = names(pal), fill =
pal, cex = .55, bty = "n"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 TEXT

Re-convolving the compositional landscape of primary and recurrent glioblastoma reveals prognostic and targetable tissue states. *Nat Commun, 2023*

"...Heatmaps were generated using the heatmap.2 in R function from the package gplots (R Package) and scores z-scaled were indicated."

DOI ↗ Fig. 3 on PMC ↗

Figure 3 TEXT

Transcriptomic atlas and interaction networks of brain cells in mouse CNS demyelination and remyelination. *Cell Rep, 2023*

"...For heat maps of relative gene expression across cell types, average expression of each gene was Z score transformed across samples and plotted using heat.map2 function in the gplots packge (v3.1.1) in R."

DOI ↗ Figure 3 on PMC ↗

Fig. 2 TEXT

A branching model of lineage differentiation underpinning the neurogenic potential of enteric glia. *Nat Commun, 2023*

"...Data was visualised using GraphPad Prism 9.0 (GraphPad), the python SCANPY67 and scVelo packages17, and the R Seurat (v3 and 4)84, EnhancedVolcano (https://github.com/kevinblighe/EnhancedVolcano), ggplot2 ..."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/gplots_heatmap2/heatmap2_recreate.R`

√ verified run · exit 0 · 9 s

gplots (heatmap.2) · 2 variations



v1 heatmap2_rowz

HEATMAPS



v2 heatmap2_celltype_cellnote

HEATMAPS

corrplot

#48

OF 100 · METHODS MENTIONS

R · CORRELATION · HEATMAPS & DOT MATRICES

147	0.83%	151	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Correlation matrix plots

THIS RECREATION

corrplot: correlation of pseudobulk cell-type profiles and of marker genes, Zeisel 2015

KEY PLOTTING CALL

RECREATIONS/CORRPLOT/CORRPLOT_RECREATE.R — EXCERPT · ENDS LINE 6

```
so <- readRDS("../_data/zeisel_seurat.rds"); m <- read.csv("../_data/zeisel_markers.csv") %>% filter(p_val_adj < .01) %>% group_by(cluster) %>% slice_max(avg_log2FC, n = 50)
av <- AverageExpression(so, features = unique(m$gene), group.by = "celltype", assays = "RNA")$RNA; C <- cor(log1p(as.matrix(av)), method = "spearman")
png("v1_corrplot_celltypes_ellipse.png", 1100, 1100, res = 160); corrplot(C, method = "ellipse", type = "upper", order = "hclust", addrect = 3, tl.col = "black", col = COL2("RdBu", 200), title = "Cell-type pseudobulk Spearman (marker genes)", mar = c(0,0,2,0)); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 TEXT

A proteomics analysis of 5xFAD mouse brain regions reveals the lysosome-associated protein Arl8b as a candidate biomarker for Alzheimer's disease. *Genome Med, 2023*

"...2g) using the packages correlation and corrplot in R v.4.2.0."

DOI ↗ Fig. 2 on PMC ↗

Fig. 3 TEXT

Single-cell multiregion dissection of Alzheimer's disease. *Nature, 2024*

"...The correlation of mean module scores between regions was determined using the cor.mtest function of the R package corrplot."

DOI ↗ Fig. 3 on PMC ↗

Fig. 2 TEXT

Spatial transcriptomics reveals niche-specific enrichment and vulnerabilities of radial glial stem-like cells in malignant gliomas. *Nat Commun, 2023*

"...The resultant correlation coefficient matrix was subjected to hierarchical clustering using corrplot package-based hclust method61, integrating the 48 spatially informed marker gene sets into four cluster ..."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/corrplot/corrplot_recreate.R`

√ verified run · exit 0 · 8 s

corrplot · 3 variations



v1 corrplot_celltypes_ellipse

HEATMAPS



v2 corrplot_genes_mixed

HEATMAPS



v3 corrplot_shade_hclust

HEATMAPS

hdWGCNA

#50

OF 100 · METHODS MENTIONS

R · NETWORK/MODULES (SC) · NETWORKS, GRAPHS & TREES · EMBEDDINGS & FEATURE MAPS · HEATMAPS & DOT MATRICES

134	0.75%	147	analysis+plotting	5
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS



NICHE & COOL · HDWGCNA

Neuro-corpus mentions per year (*2026 partial) · niche rank **#2**, growth **6.66×**, visual 5/5, score 0.938. Broad-corpus rate 1.08 → 7.252 per 1k papers (2023 → 2025–26).
GitHub smorabit/hdWGCNA · created 2020-08-11 · last push 2026-06-16

BEST FOR

Single-cell module UMAPs, hub-gene networks, module dot plots

THIS RECREATION

hdWGCNA single-cell co-expression modules on Zeisel cortex/hippocampus (metacells per cell type)

KEY PLOTTING CALL

RECREATIONS/HDWGCNA/HDWGCNA_RECREATE.R — EXCERPT · ENDS LINE 9

```
so <- readRDS("../_data/zeisel_seurat.rds")
so <- SetupForWGCNA(so, gene_select = "fraction", fraction = .25, wgcna_name = "zeisel")
so <- MetacellsByGroups(so, group.by = c("celltype","tissue"), reduction = "pca", k = 20, max_shared = 10, ident.group = "cellt
ype", min_cells = 50)
so <- NormalizeMetacells(so); so <- SetDatExpr(so, group_name = levels(so$celltype), group.by = "celltype", assay = "RNA",
layer = "data")
so <- TestSoftPowers(so, networkType = "signed")
ggsave("v1_soft_powers.png", wrap_plots(PlotSoftPowers(so), ncol = 2), width = 9, height = 6, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 1 — FIGURE CAPTION
hdWGCNA identifies co-expression networks in high-dimensional transcriptomics data. *Cell Rep Methods, 2023*
"...Overview of the hdWGCNA workflow and application in the human prefrontal cortex(A) Schematic overview of the standard hdWGCNA workflow on a scRNA-seq dataset. UMAP plot shows 36,671 cells from 11 cognitively ..."
DOI ↗ Figure 1 on PMC ↗

Fig. 3 — TEXT
Spatial and single-nucleus transcriptomic analysis of genetic and sporadic forms of Alzheimer's disease. *Nat Genet, 2024*
"...Additionally, we computed a consensus topological overlap matrix (TOM) based on the seven TOMs from the co-expression analysis in each spatial region, and we visualized this consensus TOM as a UMAP plot using ..."
DOI ↗ Fig. 3 on PMC ↗

Fig. 2 — TEXT
Cell-type brain-region specific changes in prefrontal cortex of a mouse model of alcohol dependence. *Neurobiol Dis, 2024*
"...Additionally, we used hdWGCNA (Morabito et al., 2022;"
DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/hdWGCNA/hdWGCNA_recreate.R`

√ verified run · exit 0 · 555 s

hdWGCNA · 5 variations



v1 soft_powers

NETWORKS



v2 dendrogram

NETWORKS



v3 ModuleFeaturePlot

EMBEDDINGS



v4 module_dotplot

HEATMAPS



v5 ModuleUMAPPlot

NETWORKS

clustree

#52

OF 100 · METHODS MENTIONS

R · CLUSTER TREE · NETWORKS, GRAPHS & TREES · EMBEDDINGS & FEATURE MAPS

133	0.75%	144	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Cluster-resolution trees

THIS RECREATION

clustree: Seurat Louvain resolution sweep on Zeisel 2015 (tree of cluster splits, overlaid on UMAP, marker-coloured)

KEY PLOTTING CALL

RECREATIONS/CLUSTREE/CLUSTREE_RECREATE.R — EXCERPT · ENDS LINE 5

```
so <- readRDS("../_data/zeisel_seurat.rds"); so <- FindClusters(so, resolution = c(.1,.2,.4,.6,.8,1.2), verbose = FALSE)
ggsave("v1_clustree_tree.png", clustree(so, prefix = "RNA_snn_res.") + ggtitle("clustree: resolution sweep (Zeisel 2015)"), wid
th = 9, height = 9, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Extended Data Fig. 9
FIGURE CAPTION

Microglia regulate central nervous system myelin growth and integrity. *Nature, 2023*

"...ple = 4, orange = 5, red = 6). (k) Clustering with four different resolutions, obtained by modifying the number of nearest neighbours (k). Each resolution is represented in a separate t-SNE plot. (l) Clustree ..."

DOI ↗ Extended Data Fig. 9 on PMC ↗

Figure 4
TEXT

Trem2 expression in microglia is required to maintain normal neuronal bioenergetics during development. *Immunity, 2024*

"...Optimal resolution was chosen by analyzing the cluster tree plot (clustree)."

DOI ↗ Figure 4 on PMC ↗

Fig. 5
TEXT

Divergent single cell transcriptome and epigenome alterations in ALS and FTD patients with C9orf72 mutation. *Nat Commun, 2023*

"...We used SC3 stability index101 and Clustree (v0.5.0)102 to evaluate and visualize the stability of the clusters across a range of resolutions for the Leiden community detection analysis (Supplementary Fig."

DOI ↗ Fig. 5 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/clustree/clustree_recreate.R`

✓ verified run · exit 0 · 14 s

clustree · 4 variations



v1 clustree_tree

NETWORKS



v2 clustree_sc3stability

NETWORKS



v3 clustree_gene_Mbp

NETWORKS



v4 clustree_overlay

EMBEDDINGS

circlize

#54

OF 100 · METHODS MENTIONS

R | CIRCOS/CHORD | CELL-CELL COMMUNICATION

124	0.7%	130	dedicated plotting	2
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · CIRCLIZE (CHORD)

Neuro-corpus mentions per year (*2026 partial) · niche rank **#43**, growth **1.17×**, visual 5/5, score 0.602. Broad-corpus rate 7.697 → 8.979 per 1k papers (2023 → 2025–26).
GitHub jokergoo/circlize · created 2013-04-18 · last push 2023-11-11

BEST FOR

Circos/chord diagrams; colour mapping for ComplexHeatmap

THIS RECREATION

circlize: chord diagram of LIANA+ LR counts (Zeisel) + circos genome rings (airway DE genes, hg19)

KEY PLOTTING CALL

RECREATIONS/CIRCLIZE_CHORD/CIRCLIZE_RECREATE.R — EXCERPT · ENDS LINE 5

```
m <- as.matrix(read.csv("../LIANA_plus/lr_counts_matrix.csv", row.names = 1, check.names = FALSE)); grid.col <- setNames(hcl.co
lors(length(union(rownames(m), colnames(m))), "Dark 3"), union(rownames(m), colnames(m)))
png("v1_chord_directional.png", 1300, 1300, res = 170); circos.par(gap.after = 4); chordDiagram(m, grid.col = grid.col, directi
onal = 1, direction.type = c("diffHeight","arrows"), link.arr.type = "big.arrow", annotationTrack = c("grid","name"), transpare
ncy = .3); title("Cell-cell LR pair counts (LIANA+)"); circos.clear(); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 6 TEXT
Diagnostic utility of DNA methylation analysis in genetically unsolved pediatric epilepsies and CHD2 episignature refinement. *Nat Commun, 2024*
"...Heatmaps were plotted using the R package pheatmap (version 1.0.12) and circos plots using the R package circlize (version 0.4.15)95."
DOI ↗ Fig. 6 on PMC ↗

Fig. 7 TEXT
Synaptic connectome of the Drosophila circadian clock. *Nat Commun, 2024*
"...Data was visualized using ggplot2 (v 3.4.2, Wickham, 2016) and circlize (v 0.4.15) for R (v 4.2.2) in RStudio (2022.12.0)91."
DOI ↗ Fig. 7 on PMC ↗

Figure 4 TEXT
The meningeal transcriptional response to traumatic brain injury and aging. *Elife, 2023*
"...Data was organized and graphs were created using ggplot2, tidyverse, treemapify, circlize, Seurat and dplyr (Wickham, 2016;"
DOI ↗ Figure 4 on PMC ↗

LIMITATIONS & NOTES

- Chord + circos genome rings
- Reads LIANA+ result tables, so run recreations/LIANA_plus first (METHODS §9).

Reproduce: `Rscript recreations/circlize_chord/circlize_recreate.R`

✓ verified run · exit 0 · 18 s

circlize · 2 variations

RECREATIONS/CIRCLIZE_CHORD/



v1 chord_directional

CELL–CELL



v2 circos_genome_rings

CELL–CELL

UpSetR

#57

OF 100 · METHODS MENTIONS

R | SET OVERLAP | DE, QC & SET OVERLAPS

112	0.63%	130	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

UpSet set-intersection plots

THIS RECREATION

UpSetR: overlap of Zeisel 2015 cell-type marker sets (FindAllMarkers, adj p<0.01, log2FC>1)

KEY PLOTTING CALL

RECREATIONS/UPSETR/UPSETR_RECREATE.R — EXCERPT · ENDS LINE 5

```
m <- read.csv("../_data/zeisel_markers.csv") %>% filter(p_val_adj < .01, avg_log2FC > 1); L <- split(m$gene, m$cluster)
png("v1_upset_frequency.png", 1600, 900, res = 150); print(upset(fromList(L), nsets = length(L), order.by = "freq", nintersects
= 30, main.bar.color = "#2c7fb8", sets.bar.color = "#de2d26", text.scale = 1.2, mb.ratio = c(.6,.4))); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 — FIGURE CAPTION

Cell-type-specific Alzheimer's disease polygenic risk scores are associated with distinct disease processes in Alzheimer's disease. *Nat Commun, 2023*

"...Cell-type-specific Alzheimer's disease polygenic risk scores (ADPRS).a A schematic of cell-type-specific PRS derivation. b An UpSetR plot of cell-type-specific gene sets used to define cell-type-specific ..."

DOI ↗ Fig. 1 on PMC ↗

Extended Data Fig. 9 — TEXT

Characteristics of blood-brain barrier heterogeneity between brain regions revealed by profiling vascular and perivascular cells. *Nat Neurosci, 2024*

"...Differentially expressed genes were compared to enriched genes from human mural cells from the lung45, gut46 kidney47 and brain40 and visualized by UpSetR (v.1.4.0)97."

DOI ↗ Extended Data Fig. 9 on PMC ↗

Fig. 3 — TEXT

Exposure of iPSC-derived human microglia to brain substrates enables the generation and manipulation of diverse transcriptional states in vitro. *Nat Immunol, 2023*

"...To calculating the number of unique differentially expressed genes, UpSet plots were created for each state using all differentially expressed genes with the UpSetR R package."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/UpSetR/UpSetR_recreate.R`

√ verified run · exit 0 · 6 s

UpSetR · 3 variations



v1 upset_frequency

DE



v2 upset_degree_queries

DE



v3 complexheatmap_UpSet

DE

RCTD/spacexr

#59

OF 100 · METHODS MENTIONS

R | SPATIAL DECONVOLUTION | SPATIAL & TISSUE MAPS

105	0.59%	130	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Spatial deconvolution pie/proportion maps

THIS RECREATION

RCTD (spacexr) deconvolution of the 10x Visium mouse coronal brain (squidpy visium_hne; integer UMIs recovered, see _data/export_visium_for_R.py) with the Zeisel 2015 cortex/hippocampus reference. full mode. Cell-type weight maps, dominant-type map, and a spatial pie-chart (scatterpie) view.

KEY PLOTTING CALL

RECREATIONS/RCTD_SPACEXR/RCTD_RECREATE.R — EXCERPT · ENDS LINE 14

```
W <- as.matrix(normalize_weights(r@results$weights)); df <- cbind(obs[rownames(W), c("x","y")], W)
ps <- lapply(colnames(W), function(k) ggplot(df, aes(x, -y, colour = .data[[k]])) + geom_point(size = .6) + scale_colour_viridis_c(option = "inferno", limits = c(0, 1)) + coord_fixed() + theme_void() + ggtitle(k) + theme(legend.position = "none", plot.title = element_text(size = 9)))
ggsave("v1_weight_maps.png", wrap_plots(ps, ncol = 4) + plot_annotation(title = "RCTD cell-type weights (Zeisel ref -> Visium mouse brain)"), width = 12, height = 6.5, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Extended Data Fig. 6 FIGURE CAPTION

Programs, origins and immunomodulatory functions of myeloid cells in glioma. *Nature, 2025*

"...was applied to spatial transcriptomic data from Ravi et al. to define broad transcriptomic tumor niche programs; in parallel, the cell content of each spot was inferred based on scRNA-seq signatures (RCTD ..."

DOI ↗ Extended Data Fig. 6 on PMC ↗

Fig. 4 FIGURE CAPTION

An integrated single-nucleus and spatial transcriptomics atlas reveals the molecular landscape of the human hippocampus. *Nat Neurosci, 2025*

"...NMF reveals cell type heterogeneity and biologically relevant pathways in hippocampal subfields compared to RCTD deconvolution results.For all spot plots of example capture areas from donor Br3942, spot ..."

DOI ↗ Fig. 4 on PMC ↗

Fig. 2 FIGURE CAPTION

BANKSY unifies cell typing and tissue domain segmentation for scalable spatial omics data analysis. *Nat Genet, 2024*

"...2 and 3). Top row, BANKSY clustering. Second row, nonspatial clustering (BANKSY with λ = 0). Third row, RCTD4 weights. The r values indicate the point-biserial correlation of cluster assignments with RCTD ..."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/RCTD_spacexr/RCTD_recreate.R`

✓ verified run · exit 0 · 19 s

RCTD/spacexr · 3 variations

RECREATIONS/RCTD_SPACEXR/



v1 weight_maps

SPATIAL



v2 dominant_type

SPATIAL



v3 scatterpie_zoom

SPATIAL

CytoTRACE

#60

OF 100 · METHODS MENTIONS

R/PYTHON | DIFFERENTIATION POTENCY | TRAJECTORY, PSEUDOTIME & VELOCITY | COMPOSITION, DISTRIBUTIONS & LAYOUTS

103	0.58%	131	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS



NICHE & COOL · CYTOTRACE2

Neuro-corpus mentions per year (*2026 partial) · niche rank **#5**, growth **324.85×**, visual 4/5, score 0.9. Broad-corpus rate 0 → 3.239 per 1k papers (2023 → 2025–26).
GitHub digitalcytometry/cytotrace2 · created 2024-03-05 · last push 2026-07-08

BEST FOR

Differentiation-potency score on UMAP / boxplots

THIS RECREATION

CytoTRACE 2 (interpretable deep model of developmental potency) on La Manno 2018 dentate gyrus: potency score on the UMAP, potency category composition per cell type, and score-vs-cell-type ranking.

KEY PLOTTING CALL

RECREATIONS/CYTOTRACE2/CYTOTRACE2_RECREATE.PY — EXCERPT · ENDS LINE 19

```
fig,axs=plt.subplots(1,2,figsize=(13,5.5))
sc.pl.umap(a,color="clusters",ax=axs[0],show=False,frameon=False,legend_loc="on data",legend_fontsize=7,title="Dentate gyrus cell types")
o=np.argsort(a.obs.CytoTRACE2_Score.values); xy=a.obsm["X_umap"]
s=axs[1].scatter(xy[o,0],xy[o,1],c=a.obs.CytoTRACE2_Score.values[o],cmap="Spectral_r",s=6,vmin=0,vmax=max(.35,np.quantile(a.obs.CytoTRACE2_Score,.995))); axs[1].set_axis_off(); axs[1].set_title("CytoTRACE 2 potency score (colour range clipped to data)"); fig.colorbar(s,ax=axs[1],shrink=.6)
fig.tight_layout(); fig.savefig("v1_potency_umap.png",dpi=140)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 4 — FIGURE CAPTION

Decoding multiple myeloma: single-cell insights into tumor heterogeneity, immune dynamics, and disease progression. *Front Immunol, 2025*

"…Proposed time-series analysis of myeloma cell subpopulations. (A, B) The UMAP and Bar plots showed the results of the stemness ability of four subtypes of myeloma cells calculated based on CytoTRACE. (C) Bar …"

DOI ↗ Figure 4 on PMC ↗

Fig. 3 — FIGURE CAPTION

Cranioencephalic functional lymphoid units in glioblastoma. *Nat Med, 2024*

"…dition and tissue source. CBe T cell types in dark blue. g, UMAP of 3′ GEX CB data from e generated by Palantir, based on diffusion map dimensionality reduction, color coded as in e. Left, continuous CytoTRACE …"

DOI ↗ Fig. 3 on PMC ↗

Fig. 6 — FIGURE CAPTION

Single-cell RNA sequencing unveils Lrg1's role in cerebral ischemia–reperfusion injury by modulating various cells. *J Neuroinflammation, 2023*

"…he direction of arrows indicates the direction of differentiation and development. H The dot plot illustrates the differentiation trajectories of microglial subpopulations in Sham + WT group drawn by …"

DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

- cytotrace2_py is installed from a GitHub subdirectory (METHODS §10).

Reproduce: `python3 recreations/CytoTRACE2/cytotrace2_recreate.py`

√ verified run · exit 0 · 102 s

CytoTRACE · 3 variations



v1 potency_umap

TRAJECTORY



v2 potency_composition

COMPOSITION



v3 score_ranking

COMPOSITION

decoupleR/decoupler

#61

OF 100 · METHODS MENTIONS

R/PYTHON | TF/PATHWAY ACTIVITY | ENRICHMENT & PATHWAY ACTIVITY

102	0.57%	112	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

TF/pathway activity heatmaps, UMAPs

THIS RECREATION

decoupleR: PROGENy pathway and CollecTRI TF activities (ULM) on Zeisel 2015 (OmniPath networks fetched at run time): pathway activity heatmap per cell type, TF activity heatmap (top variable TFs), pathway score on UMAP.

KEY PLOTTING CALL

RECREATIONS/DECOUPLER/DECOUPLER_RECREATE.R — EXCERPT · ENDS LINE 14

```
ap <- run_ulm(X, net_p, .source = "source", .target = "target", .mor = "weight", minsize = 5)
at <- run_ulm(X, net_t, .source = "source", .target = "target", .mor = "weight", minsize = 10)
agg <- function(a) { M <- a %>% pivot_wider(id_cols = condition, names_from = source, values_from = score) %>% tibble::column_t
o_rownames("condition") %>% as.matrix(); M <- M[colnames(so), ]
  t(sapply(split(seq_len(ncol(so)), so$celltype), function(i) colMeans(M[i, , drop = FALSE]))) }
P <- agg(ap); Tf <- agg(at); pal <- colorRampPalette(c("#2166ac","white","#b2182b"))(100)
png("v1_progeny_heatmap.png", 1300, 800, res = 150); pheatmap(P, scale = "column", color = pal, border_color = NA, main = "deco
upleR PROGENy pathway activity (ULM, mean per cell type)"); dev.off()
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 FIGURE CAPTION — **Single-nucleus RNA sequencing reveals glial cell type-specific responses to ischemic stroke in male rodents.** *Nat Commun, 2024*

"...The Top 25 module-defining genes, as sorted by descending Moran´s I, are depicted in boxes on the right side of the respective gene module feature plots. l Heatmap depicting the top 100 most variable ..."

DOI ↗ Fig. 2 on PMC ↗

Fig. 8 FIGURE CAPTION — **The spatial landscape of glial pathology and T cell response in Parkinson's disease substantia nigra.** *Nat Commun, 2025*

"...ed based on two-sided adjusted p-values. d Bar plot showing activation scores of select pathways in differentially expressed genes in CD44 KD astrocytes. The scores were calculated using Omnipath and decoupleR ..."

DOI ↗ Fig. 8 on PMC ↗

Fig. 2 TEXT — **Cell type mapping reveals tissue niches and interactions in subcortical multiple sclerosis lesions.** *Nat Neurosci, 2024*

"...Pseudobulk transcriptomic profiles were generated for both atlases at the cell type level using decoupler-py (v.1.5.1, commit b6b430e)73."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

- OmnipathR's PROGENy/CollecTRI download failed, so the networks are fetched at run time with the Python decoupler 2.x (dc.op.*) (script header).

Reproduce: `Rscript recreations/decoupleR/decoupleR_recreate.R`

✓ verified run · exit 0 · 78 s

decoupleR/decoupler · 3 variations



v1 progeny_heatmap

ENRICHMENT



v2 collectri_tf_heatmap

ENRICHMENT



v3 pathway_umaps

ENRICHMENT

ABC atlas / Allen Brain Map

#63

OF 100 · METHODS MENTIONS

WEB | ATLAS | BRAIN ATLAS, 3D & NEUROIMAGING

96	0.54%	169	viewer	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · BRAINGLOBE ATLAS API

Neuro-corpus mentions per year (*2026 partial) · niche rank **#18**, growth **2.27×**, visual 5/5, score 0.798. Broad-corpus rate 0 → 0.013 per 1k papers (2023 → 2025–26).

GitHub brainglobe/brainglobe-atlasapi · created 2020-04-30 · last push 2026-10-14 · also scored: Allen CCF overlays (#61, growth 1.14×)

BEST FOR

Atlas reference views / cell-type taxonomies

THIS RECREATION

Allen Mouse CCFv3 (BrainGlobe atlas API, allen_mouse_25um): coronal annotation overlay, region outlines, 3D region point cloud

KEY PLOTTING CALL

RECREATIONS/BRAINGLOBE_ATLAS/BRAINGLOBE_RECREATE.PY — EXCERPT · ENDS LINE 12

```
a=BrainGlobeAtlas("allen_mouse_25um",check_latest=False); ref=a.reference; ann=a.annotation; z=ann.shape[0]//2+20
cols=np.zeros(ann.shape[1:]+(3,)); sl=ann[z]
for rid in np.unique(sl):
    if rid==0: continue
    try: cols[sl==rid]=np.array(a.structures[rid]["rgb_triplet"])/255
    except KeyError: pass
fig,ax=plt.subplots(1,2,figsize=(12,4.8)); ax[0].imshow(ref[z],cmap="gray"); ax[0].set_title("CCFv3 template, coronal slice"); a
x[1].imshow(ref[z],cmap="gray"); ax[1].imshow(cols,alpha=.6); ax[1].set_title("Allen annotation overlay (official colours)")
[x.axis("off") for x in ax]; fig.tight_layout(); fig.savefig("v1_coronal_overlay.png",dpi=140)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 4 — FIGURE CAPTION

Conservation and alteration of mammalian striatal interneurons. *Nature, 2025*

"…parent clusters 0840–0842. h, Distribution of highlighted ABC clusters in the adult mouse brain and Th, Tac2 and Lhx8 expression (in counts per million reads (CPM)) found within those populations27. ABC atlas …"

DOI ↗ Fig. 4 on PMC ↗

Figure S7 — FIGURE CAPTION

LRRC37B is a human modifier of voltage-gated sodium channels and axon excitability in cortical neurons. *Cell, 2023*

"…m HEK293T cells transfected for their cDNA and SCN1B cDNA: all LRRC37B proteins bind to SCN1B but not LRRC37A2.(C) LRRC37B, FGF13, SCN1B, and SCN8A transcripts detection in human cortical cells (from Allen …"

DOI ↗ Figure S7 on PMC ↗

Figure 1 — TEXT

The expression profile and tumorigenic mechanisms of CD97 (ADGRE5) in glioblastoma render it a targetable vulnerability. *Cell Rep, 2023*

"…First, we analyzed the mRNA expression levels of all 32 aGPCRs in healthy brain tissue using the Allen Brain Map database and in PDGCs collected in-house (Figure 1A)."

DOI ↗ Figure 1 on PMC ↗

LIMITATIONS & NOTES

- Allen CCFv3 atlas renders via BrainGlobe atlas API

Reproduce: `python3 recreations/BrainGlobe_atlas/brainglobe_recreate.py`

✓ verified run · exit 0 · 6 s

ABC atlas / Allen Brain … · 3 variations

RECREATIONS/BRAINGLOBE_ATLAS/



v1 coronal_overlay

BRAIN



v2 region_outlines

BRAIN



v3 ccf_3d_regions

BRAIN

NicheNet

#66

OF 100 · METHODS MENTIONS

R | LIGAND-TARGET | CELL-CELL COMMUNICATION

90	0.51%	122	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS



NICHE & COOL · NICHENET

Neuro-corpus mentions per year (*2026 partial) · niche rank **#46**, growth **1.73×**, visual 4/5, score 0.592. Broad-corpus rate 4.632 → 8.026 per 1k papers (2023 → 2025–26).
GitHub saeyslab/nichenetr · created 2018-02-05 · last push 2026-09-25

BEST FOR

Ligand activity / ligand-target heatmaps, circos

THIS RECREATION

NicheNet (nichenetr v2 priors, mouse, Zenodo 7074291): which glial/vascular ligands best explain the cortex-vs-CA1 programme in oligodendrocytes (Zeisel 2015)? Ligand activity, ligand→target heatmap, ligand→receptor heatmap, and a circlize circos of top links coloured by sender.

KEY PLOTTING CALL

RECREATIONS/NICHENET/NICHENET_RECREATE.R — EXCERPT · ENDS LINE 18

```
la <- nn$ligand_activities %>% filter(test_ligand %in% nn$top_ligands) %>% arrange(aupr_corrected)
g1 <- ggplot(la, aes(aupr_corrected, factor(test_ligand, levels = test_ligand))) + geom_segment(aes(x = 0, xend =
aupr_corrected, yend = factor(test_ligand, levels = test_ligand)), colour = "grey70") + geom_point(aes(colour =
aupr_corrected), size = 4) + scale_colour_gradient(low = "#fdd49e", high = "#d7301f") + theme_bw() + labs(x = "ligand activity
(corrected AUPR)", y = NULL, colour = "AUPR", title = "NicheNet ligand activity")
ggsave("v1_ligand_activity.png", g1, width = 5.5, height = 6, dpi = 150, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 8 — FIGURE CAPTION

A single nuclear transcriptomic characterisation of mechanisms responsible for impaired angiogenesis and blood-brain barrier function in Alzheimer's disease. *Nat Commun, 2024*

"...NicheNet intercellular communication analysis identified potential regulators of EC DEG associated with proinflammatory and anti-angiogenic gene expression in astrocytes and perivascular macrophages.A ..."

DOI ↗ Fig. 8 on PMC ↗

Figure 8 — FIGURE CAPTION

Single-cell and Spatial Transcriptomics Reveals Ferroptosis as The Most Enriched Programmed Cell Death Process in Hemorrhage Stroke-induced Oligodendrocyte-mediated White Matter Injury. *Int J Biol Sci, 2024*

"...LCN2+ microglia induce oligodendrocyte ferroptosis via the CSF1/CSF1R pathway. (A) Top-ranked ligands that were inferred to regulate oligodendrocytes by LCN2+ microglia (Mg5) according to NicheNet. Dot plots ..."

DOI ↗ Figure 8 on PMC ↗

Figure 6 — FIGURE CAPTION

Transcriptomic atlas and interaction networks of brain cells in mouse CNS demyelination and remyelination. *Cell Rep, 2023*

"...NicheNet analysis reveals cell-cell interactions elicited by demyelination(A–C) Circos plot showing links between predicted ligands from sender cells (bottom) with their associated receptors on receiver ..."

DOI ↗ Figure 6 on PMC ↗

LIMITATIONS & NOTES

- NicheNet v2 mouse priors are downloaded from Zenodo 7074291 (script header).

Reproduce: `Rscript recreations/NicheNet/NicheNet_recreate.R`

✓ verified run · exit 0 · 42 s

NicheNet · 4 variations



v1 ligand_activity · CELL–CELL



v2 ligand_receptor_heatmap · CELL–CELL



v3 circos · CELL–CELL



v4 ligand_target_heatmap · CELL–CELL

ReactomePA

#68

OF 100 · METHODS MENTIONS

R · ENRICHMENT · ENRICHMENT & PATHWAY ACTIVITY

84	0.47%	87	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Reactome enrichment dot/network plots

THIS RECREATION

ReactomePA: Reactome pathway enrichment of Zeisel 2015 cell-type markers (mouse; compareCluster dotplot, enrichment map and gene-concept network for interneurons)

KEY PLOTTING CALL

RECREATIONS/REACTOMEPA/REACTOMEPA_RECREATE.R — EXCERPT · ENDS LINE 9

```
m <- read.csv("../_data/zeisel_markers.csv") %>% filter(p_val_adj < .01) %>% group_by(cluster) %>% slice_max(avg_log2FC, n =
200) %>% ungroup()
ids <- bitr(unique(m$gene), "SYMBOL", "ENTREZID", OrgDb = org.Mm.eg.db); m <- inner_join(m, ids, by = c(gene = "SYMBOL"))
L <- split(m$ENTREZID, m$cluster)
cc <- compareCluster(L, fun = "enrichPathway", organism = "mouse", pvalueCutoff = .05)
ggsave("v1_compareCluster_dotplot.png", dotplot(cc, showCategory = 4, label_format = 45) +
ggtitle("ReactomePA: cell-type marker pathways") + theme(axis.text.x = element_text(angle = 30, hjust = 1)), width = 11, height
= 9, dpi = 120, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT

The Global Neurodegeneration Proteomics Consortium: biomarker and drug target discovery for common neurodegenerative diseases and aging. *Nat Med, 2025*

"...Proteins identified in the meta-analytic differential abundance analysis were filtered into gene set enrichment analysis using the R libraries 'ReactomePA' and 'clusterProfiler'."

DOI ↗ Fig. 1 on PMC ↗

Fig. 1 TEXT

Shared and disease-specific pathways in frontotemporal dementia and Alzheimer's and Parkinson's diseases. *Nat Med, 2025*

"...To investigate the biological processes associated specifically with each ND and those common to all, we performed pathway enrichment analysis using the ReactomePA (version 1.48)80 and clusterProfiler (version ..."

DOI ↗ Fig. 1 on PMC ↗

Fig. 2 TEXT

MicroRNA-335-5p suppresses voltage-gated sodium channel expression and may be a target for seizure control. *Proc Natl Acad Sci U S A, 2023*

"...Pathway enrichment analysis of the remaining 267 MTIs (Dataset S3) was performed using the ReactomePA (74) and clusterProfiler (75) packages, with an adjusted p-value (Benjamini–Hochberg) <0.05 considered ..."

DOI ↗ Fig. 2 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ReactomePA/ReactomePA_recreate.R`

✓ verified run · exit 0 · 332 s

ReactomePA · 3 variations



v1 compareCluster_dotplot

ENRICHMENT



v2 emapplot

ENRICHMENT



v3 cnetplot

ENRICHMENT

pathview

#70

R · PATHWAY MAPS · ENRICHMENT & PATHWAY ACTIVITY

OF 100 · METHODS MENTIONS

82	0.46%	99	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

KEGG pathway maps coloured by fold-change

THIS RECREATION

pathview: Zeisel 2015 interneuron-vs-pyramidal-SS log2FC painted onto KEGG GABAergic (mmu04727) and glutamatergic (mmu04724) synapse maps (KEGG XML/PNG fetched once from KEGG REST at run time).

KEY PLOTTING CALL

RECREATIONS/PATHVIEW/PATHVIEW_RECREATE.R — EXCERPT · ENDS LINE 9

```
for (pw in c("04727","04724")) pathview(gene.data = fc, pathway.id = pw, species = "mmu", out.suffix = "interneuron_vs_pyrSS", l
imit = list(gene = 3), low = list(gene = "#2166ac"), high = list(gene = "#b2182b"), kegg.native = TRUE)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 4 TEXT

Sildenafil as a Candidate Drug for Alzheimer's Disease: Real-World Patient Data Observation and Mechanistic Observations from Patient-Induced Pluripotent Stem Cell-Derived Neurons. *J Alzheimers Dis, 2024*

"...Significant KEGG pathways were depicted using Pathview tool with highlighted sildenafil regulated key genes (Supplementary Figures 2–4)."

DOI ↗ Fig. 4 on PMC ↗

Fig. 1 TEXT

Integrating lipid metabolism, pheromone production and perception by Fruitless and Hepatocyte Nuclear Factor 4. *Sci Adv, 2023*

"...The KEGG pathway maps were visualized using pathview (111)."

DOI ↗ Fig. 1 on PMC ↗

Fig. 11 TEXT

Sex differences in the transcriptome of extracellular vesicles secreted by fetal neural stem cells and effects of chronic alcohol exposure. *Biol Sex Differ, 2023*

"...Pathway analysis was conducted on differentially expressed genes using ReactomePA R package [69] which utilizes the KEGG database [70] and the Pathview R package [71] to visualize differentially regulated ..."

DOI ↗ Fig. 11 on PMC ↗

LIMITATIONS & NOTES

- KEGG XML/PNG is fetched once from KEGG REST at run time, so this needs internet (script header).

Reproduce: `Rscript recreations/pathview/pathview_recreate.R`

√ verified run · exit 0 · 20 s

pathview · 3 variations



v1 GABAergic_synapse_native

ENRICHMENT



v2 glutamatergic_synapse_native

ENRICHMENT



v3 GABAergic_graphviz

ENRICHMENT

GOplot

#71

R ENRICHMENT ENRICHMENT & PATHWAY ACTIVITY

OF 100 · METHODS MENTIONS

81	0.46%	85	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

GOCircle, GOChord, GOBubble

THIS RECREATION

GOplot (GOBubble, GOCircle, GOChord, GOCluster) on oligodendrocyte markers vs all other Zeisel 2015 cell types (clusterProfiler enrichGO BP, mouse) - logFC from FindAllMarkers.

KEY PLOTTING CALL

RECREATIONS/GOPLOT/GOPLOT_RECREATE.R — EXCERPT · ENDS LINE 9

```
m <- read.csv("../_data/zeisel_markers.csv") %>% filter(cluster == "oligodendrocytes", p_val_adj < .01) %>%
slice_max(avg_log2FC, n = 300)
ego <- enrichGO(m$gene, OrgDb = org.Mm.eg.db, keyType = "SYMBOL", ont = "BP", pvalueCutoff = .01, readable = FALSE) %>%
simplify(cutoff = .6)
e <- as.data.frame(ego) %>% head(12); terms <- data.frame(category = "BP", ID = e$ID, term = e$Description, genes = gsub("/",
", ", e$geneID), adj_pval = e$p.adjust)
genes <- data.frame(ID = m$gene, logFC = m$avg_log2FC); circ <- circle_dat(terms, genes)
ggsave("v1_GOBubble.png", GOBubble(circ, labels = 3, table.legend = FALSE, display = "single") + ggtitle("GOBubble:
oligodendrocyte marker BP terms"), width = 8, height = 7, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 3 TEXT

Human Brain Microvascular Endothelial Cells Exposure to SARS-CoV-2 Leads to Inflammatory Activation through NF-κB Non-Canonical Pathway and Mitochondrial Remodeling. *Viruses, 2023*

"...Overlapping gene sets from reactome pathway terms were visualized as a chord plot using the GOplot R [44]."

DOI ↗ Figure 3 on PMC ↗

Figure 2 TEXT

A novel phenotype of B cells associated with enhanced phagocytic capability and chemotactic function after ischemic stroke. *Neural Regen Res, 2023*

"...The activation Z-score was calculated for each term by the R package GOplot to predict the activation state (Walter et al., 2015)."

DOI ↗ Figure 2 on PMC ↗

Fig. 6 TEXT

Choroid plexus defects in Down syndrome brain organoids enhance neurotropism of SARS-CoV-2. *Sci Adv, 2024*

"...Enrichment data were then combined with expression values, and z scores were calculated using the GOplot v.1.0.2 R package and plotted using ggplot2 v.3.3.2."

DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

- GOCluster / GOHeat omitted, and GOChord drawn without the logFC ring: GOplot logFC gradients fail with ggplot2 ≥ 3.5 (script comment).

Reproduce: `Rscript recreations/GOplot/GOplot_recreate.R`

√ verified run · exit 0 · 50 s

GOplot · 3 variations



v1 GOBubble

ENRICHMENT



ID	Description
GO:0008366	axon ensheathment
GO:0046394	carboxylic acid biosynthetic process
GO:0032288	myelin assembly
GO:0043604	amide biosynthetic process
GO:0046467	membrane lipid biosynthetic process
GO:0099612	protein localization to axon
GO:0008203	cholesterol metabolic process
GO:0031579	membrane raft organization

v2 GOCircle

ENRICHMENT



v3 GOChord

ENRICHMENT

scCustomize

#74

OF 100 · METHODS MENTIONS

R · SINGLE-CELL · EMBEDDINGS & FEATURE MAPS · HEATMAPS & DOT MATRICES · COMPOSITION, DISTRIBUTIONS & LAYOUTS

75	0.42%	78	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS



NICHE & COOL · SCCUSTOMIZE

Neuro-corpus mentions per year (*2026 partial) · niche rank **#30**, growth **4.24×**, visual 3/5, score 0.684. Broad-corpus rate 0.627 → 2.692 per 1k papers (2023 → 2025–26).
GitHub samuel-marsh/scCustomize · created 2021-09-29 · last push 2026-06-30

BEST FOR

Enhanced Seurat plots (FeaturePlot_scCustom, Clustered_DotPlot)

KEY PLOTTING CALL

RECREATIONS/SCCUSTOMIZE/SCCUSTOMIZE_RECREATE.R — EXCERPT · ENDS LINE 4

```
so <- readRDS("../_data/zeisel_seurat.rds"); mk <-
c("Snap25","Gad1","Gad2","Pvalb","Sst","Satb2","Neurod6","Wfs1","Aqp4","Gfap","Mbp","Plp1","Cx3cr1","C1qa","Cldn5","Acta2")
ggsave("v1_DimPlot_scCustom_figureplot.png", DimPlot_scCustom(so, figure_plot = TRUE, label = TRUE), width = 7, height = 6, dpi
= 140, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 TEXT

Characterisation of premature cell senescence in Alzheimer's disease using single nuclear transcriptomics. *Acta Neuropathol, 2024*

"...Gene set or module feature plots were generated by first calculating aggregated gene set scores using AddModuleScore() function from Seurat and then plotted using FeaturePlot scCustom() from scCustomize (v."

DOI ↗ Fig. 2 on PMC ↗

Fig. 4 TEXT

cGAS-STING drives ageing-related inflammation and neurodegeneration. *Nature, 2023*

"...9a,f) and visualized using the FeaturePlot_scCustom function from the R package scCustomize (v.0.7.0), allowing for subsequent analysis of specific cell types (such as microglia, oligodendrocytes) ..."

DOI ↗ Fig. 4 on PMC ↗

Fig. 1 TEXT

Microglia ferroptosis is regulated by SEC24B and contributes to neurodegeneration. *Nat Neurosci, 2023*

"...UMAP plots and gene expression plots were generated using built-in Seurat/ggplot2 plotting functions or scCustomize version 0.7.0 unless otherwise described."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/scCustomize/scCustomize_recreate.R`

√ verified run · exit 0 · 17 s

scCustomize · 4 variations



v1 DimPlot_scCustom_figureplot

EMBEDDINGS



v2 FeaturePlot_scCustom

EMBEDDINGS



v3 Clustered_DotPlot

HEATMAPS



v4 Stacked_VlnPlot

COMPOSITION

patchwork

#79

R LAYOUT COMPOSITION, DISTRIBUTIONS & LAYOUTS

OF 100 · METHODS MENTIONS

65	0.37%	71	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Multi-panel layout of ggplots

Also catalogued as: **cowplot** (#94, 44 papers)

THIS RECREATION

patchwork + cowplot multi-panel composites (typical Fig. 1 layout) from Zeisel 2015

KEY PLOTTING CALL

RECREATIONS/PATCHWORK_COWPLOT/COMPOSITE_RECREATE.R — EXCERPT · ENDS LINE 11

```
d <- so@meta.data %>% count(tissue, celltype) %>% group_by(tissue) %>% mutate(f = n / sum(n)) %>% ggplot(aes(tissue, f, fill = c
elltype)) + geom_col() + theme_classic() + labs(y = "fraction", x = NULL)
e <- DotPlot(so, features = c("Gad1","Slc17a7","Neurod6","Mbp","Aqp4","Cx3cr1","Cldn5"), group.by = "celltype") + RotatedAxis()
+ theme(axis.title = element_blank())
p <- (a | b) / (c | d | e) + plot_layout(heights = c(1.1, 1), widths = c(1, 1, 1.5)) + plot_annotation(tag_levels = "A", title
= "patchwork composite: Zeisel 2015 overview") & theme(plot.tag = element_text(face = "bold", size = 16))
ggsave("v1_patchwork_composite.png", p, width = 15, height = 10, dpi = 110, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 TEXT

TDP-43 nuclear loss in FTD/ALS causes widespread alternative polyadenylation changes. *Nat Neurosci, 2025*

"...All plots were prepared using tidyplots, ggplot2, ggpubr, patchwork and ggrepel in R."

DOI ↗ Fig. 3 on PMC ↗

Fig. 4 TEXT

Prospective, multicenter validation of a platform for rapid molecular profiling of central nervous system tumors. *Nat Med, 2025*

"...Plots were generated using ggplot2 (v.3.5.1), ggsankey (v.0.0.99999), ggridges (v.0.5.4), patchwork (v.1.1.3) and related R packages for visualization."

DOI ↗ Fig. 4 on PMC ↗

Fig. 3 TEXT

Repression of developmental transcription factor networks triggers aging-associated gene expression in human glial progenitor cells. *Nat Commun, 2024*

"...Graphs were further edited or generated anew using ggplot2 and aligned using patchwork."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/patchwork_cowplot/composite_recreate.R`

✓ verified run · exit 0 · 12 s

patchwork · 3 variations



v1 patchwork_composite

COMPOSITION



v2 patchwork_design_inset

COMPOSITION



v3 cowplot_grid_watermark

COMPOSITION

Squidpy

PYTHON · SPATIAL · SPATIAL & TISSUE MAPS

#83

OF 100 · METHODS MENTIONS

60	0.34%	74	analysis+plotting	6
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · SQUIDPY

Neuro-corpus mentions per year (*2026 partial) · niche rank **#28**, growth **2.72×**, visual 4/5, score 0.736. Broad-corpus rate 1.358 → 3.708 per 1k papers (2023 → 2025–26).
GitHub scverse/squidpy · created 2020-08-21 · last push 2026-10-13

BEST FOR

Spatial scatter, neighbourhood enrichment, co-occurrence

THIS RECREATION

Squidpy: 10x Visium mouse coronal brain (H&E) + Moffitt 2018 MERFISH hypothalamus

KEY PLOTTING CALL

RECREATIONS/SQUIDPY/SQUIDPY_RECREATE.PY — EXCERPT · ENDS LINE 5

```
v=sc.read_h5ad("../_data/py/visium_hne.h5ad")
sq.pl.spatial_scatter(v,color="cluster",size=1.3,title="Visium mouse brain: annotated clusters on H&E"); plt.savefig("v1_clusters_on_HE.png",dpi=140,bbox_inches="tight"); plt.close("all")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 1 TEXT

Spatial transcriptomic clocks reveal cell proximity effects in brain ageing. *Nature, 2025*

"...To compute the distribution of nearest neighbour distances, we constructed a triangulation mesh graph connecting neighbouring cells on a given sample using squidpy.gr.spatial_neighbors with delaunay=True103."

DOI ↗ Article ↗ figure page not on PMC

Extended Data Fig. 4 TEXT

Spatial atlas of the mouse central nervous system at molecular resolution. *Nature, 2023*

"...The following packages and software were used in data analysis: UCSF ChimeraX 1.0, ImageJ 1.51, MATLAB R2019b, R 4.0.4, RStudio 1.4.1106, Jupyter Notebook 6.0.3, Anaconda 2-2-.02, h5py 3.1.0, hdbscan 0.8.36 ..."

DOI ↗ Extended Data Fig. 4 on PMC ↗

Fig. 3 TEXT

Mapping cells through time and space with moscot. *Nature, 2025*

"...To this end, we computed the neighbour graph in the aligned space using squidpy (kNN mode with at least 30 neighbours for each observation)."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/Squidpy/squidpy_recreate.py`

√ verified run · exit 0 · 16 s

Squidpy · 6 variations



v1 clusters_on_HE

SPATIAL



v2 genes_on_HE

SPATIAL



v3 nhood_enrichment

SPATIAL



v4 merfish_cell_classes

SPATIAL



v5 co_occurrence

SPATIAL



v6 ripley_L

SPATIAL

plotly

#84

OF 100 · METHODS MENTIONS

R/PYTHON/JS · INTERACTIVE · EMBEDDINGS & FEATURE MAPS · SPATIAL & TISSUE MAPS

56	0.32%	60	dedicated plotting	2
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · PLOTLY/DASH 3D

Neuro-corpus mentions per year (*2026 partial) · niche rank **#56**, growth **1.36×**, visual 4/5, score 0.537. Broad-corpus rate 2.368 → 3.226 per 1k papers (2023 → 2025–26).
GitHub plotly/plotly.py · created 2013-11-21 · last push 2026-10-14

BEST FOR

Interactive 2D/3D plots

THIS RECREATION

Interactive 3D UMAP (HTML) + static PNG export (kaleido) on Zeisel; 3D MERFISH stack of hypothalamic slices

KEY PLOTTING CALL

RECREATIONS/PLOTLY_3D/PLOTLY3D_RECREATE.PY — EXCERPT · ENDS LINE 7

```
ad=load(); sc.pp.highly_variable_genes(ad,n_top_genes=2000); sc.pp.pca(ad); sc.pp.neighbors(ad); sc.tl.umap(ad,n_components=3)
u=ad.obsm["X_umap"]; f=px.scatter_3d(x=u[:,0],y=u[:,1],z=u[:,2],color=ad.obs.celltype.astype(str),opacity=.8,title="3D UMAP,
Zeisel cortex/hippocampus")
f.update_traces(marker_size=2); f.update_layout(scene=dict(xaxis_title="UMAP1",yaxis_title="UMAP2",zaxis_title="UMAP3"),legend_
itemsizing="constant",width=900,height=700)
f.write_html("v1_umap3d.html"); f.write_image("v1_umap3d.png",scale=1.5)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 6 TEXT

A brain cell atlas integrating single-cell transcriptomes across human brain regions. *Nat Med, 2024*

"...The 'Data Viewer' page and other visualization modules were built with Plotly (https://plot.ly)."

DOI ↗ Fig. 6 on PMC ↗

Fig. 2 TEXT

Mapping medically relevant RNA isoform diversity in the aged human frontal cortex with deep long-read RNA-seq. *Nat Biotechnol, 2025*

"...We used the following Python libraries: numpy (v.1.24.1), pandas (v.1.5.2), regex (v.2022.10.31), matplotlib (v.3.6.2), seaborn (v.0.12.2), matplotlib_venn (v.0.11.7), wordcloud (v.1.8.2.2), plotly (v.5.11.0) ..."

DOI ↗ Fig. 2 on PMC ↗

Fig. 1 TEXT

Multi-omic analysis of Huntington's disease reveals a compensatory astrocyte state. *Nat Commun, 2024*

"...Each t-SNE plot was created using the first three coordinates through the package plotly (version 4.10.0)."

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

- The primary output is interactive HTML; the book shows the static PNG export (kaleido) (script header).

Reproduce: `python3 recreations/plotly_3D/plotly3d_recreate.py`

√ verified run · exit 0 · 34 s



v1 umap3d

EMBEDDINGS



v2 merfish3d

SPATIAL

ggraph

#86

OF 100 · METHODS MENTIONS

R · NETWORK · NETWORKS, GRAPHS & TREES

50	0.28%	49	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Network/dendrogram plots in ggplot grammar

THIS RECREATION

ggraph / tidygraph: Zeisel 2015 marker co-expression network (force layout, communities), circular cell-type dendrogram, and circle-packing of cell-type → Louvain sub-cluster hierarchy.

KEY PLOTTING CALL

RECREATIONS/GGRAPH/GGRAPH_RECREATE.R — EXCERPT · ENDS LINE 10

```
X <- as.matrix(GetAssayData(so, layer = "data")[m$gene, ]); C <- cor(t(X)); C[lower.tri(C, diag = TRUE)] <- NA
e <- as.data.frame(as.table(C)) %>% filter(!is.na(Freq), Freq > .35); names(e) <- c("from","to","r")
g <- tbl_graph(nodes = data.frame(name = m$gene, celltype = m$cluster), edges = e, directed = FALSE) %>% activate(nodes) %>% mu
tate(deg = centrality_degree(), comm = as.factor(group_louvain()))
p1 <- ggraph(g, "fr") + geom_edge_link(aes(alpha = r), colour = "grey50") + geom_node_point(aes(colour = celltype, size = deg))
+ geom_node_text(aes(label = name), repel = TRUE, size = 2.4) + theme_graph(base_family = "sans") + ggtitle("Marker co-
expression network (r > 0.35)")
ggsave("v1_coexpression_network.png", p1, width = 10, height = 8, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 4 TEXT — **Single-cell epigenomic reconstruction of developmental trajectories from pluripotency in human neural organoid systems.** *Nat Neurosci, 2024*

"...Network visualization and analysis was performed with ggraph (TLP;"

DOI ↗ Fig. 4 on PMC ↗

Fig. 3 TEXT — **Single-cell brain organoid screening identifies developmental defects in autism.** *Nature, 2023*

"...The resulting graph was visualized with the R package ggraph (https://github.com/thomasp85/ggraph) using the circular tree layout."

DOI ↗ Fig. 3 on PMC ↗

Fig. 3 TEXT — **Multimodal spatiotemporal phenotyping of human retinal organoid development.** *Nat Biotechnol, 2023*

"...The cluster graph was then visualized using the ggraph package in R with the tree layout implemented in the igraph package (the layout_as_tree function), with the edge weights represented by the edge ..."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggraph/ggraph_recreate.R`

✓ verified run · exit 0 · 11 s



v1 coexpression_network

NETWORKS



v2 circular_dendrogram

NETWORKS



v3 circlepack

NETWORKS

Tangram

#87

OF 100 · METHODS MENTIONS

PYTHON | SPATIAL MAPPING | SPATIAL & TISSUE MAPS

49	0.28%	77	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS



NICHE & COOL · TANGRAM

Neuro-corpus mentions per year (*2026 partial) · niche rank **#44**, growth **1.83×**, visual 4/5, score 0.598. Broad-corpus rate 1.498 → 2.756 per 1k papers (2023 → 2025–26).
GitHub broadinstitute/Tangram · created 2020-08-21 · last push 2025-07-01

BEST FOR

Projected cell types on spatial maps

THIS RECREATION

Tangram: map Zeisel 2015 cortex/hippocampus single cells onto the 10x Visium mouse-brain H&E section (squidpy), then project cell-type probabilities onto the tissue; plus training-gene score diagnostics.

KEY PLOTTING CALL

RECREATIONS/TANGRAM/TANGRAM_RECREATE.PY — EXCERPT · ENDS LINE 17

```
tg.pp_adatas(ad_sc,ad_sp,genes=markers)
m=tg.map_cells_to_space(ad_sc,ad_sp,mode="clusters",cluster_label="celltype",device="cpu",num_epochs=500,density_prior="rna_cou
nt_based",verbose=False)
tg.project_cell_annotations(m,ad_sp,annotation="celltype")
ct=list(ad_sp.obsm["tangram_ct_pred"].columns)
P=ad_sp.obsm["tangram_ct_pred"]; P=(P-P.min())/(P.max()-P.min())
for c in ct: ad_sp.obs[f"tg_{c}"]=P[c].values
sq.pl.spatial_scatter(ad_sp,color=[f"tg_{c}" for c in ct],ncols=4,cmap="magma",size=1.4,img_alpha=.5,figsize=(4,4),save=None)
plt.gcf().suptitle("Tangram: Zeisel cell types mapped onto Visium mouse brain",y=1.02);
plt.savefig("v1_celltype_projection.png",dpi=120,bbox_inches="tight"); plt.close("all")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Extended Data Fig. 12 — FIGURE CAPTION

Spatial transcriptomic clocks reveal cell proximity effects in brain ageing. *Nature, 2025*

"...Identification and validation of potential mediators of cell proximity effects.a, Schematic of the computational pipeline using SpaGE and Tangram algorithms for spatial gene expression imputation and using ..."

DOI ↗ Article ↗ figure page not on PMC

Fig. 3 — FIGURE CAPTION

A single-cell time-lapse of mouse prenatal development from gastrula to birth. *Nature, 2024*

"...mental stage. SMC, smooth muscle cell. e, The spatial origin of each lateral plate and intermediate mesoderm derivative was inferred based on a public dataset, Mosta46, together with our data and the Tangram ..."

DOI ↗ Fig. 3 on PMC ↗

Fig. 3 — FIGURE CAPTION

Inferring histology-associated gene expression gradients in spatial transcriptomic studies. *Nat Commun, 2024*

"...se the expression profiles of co-expressed genes identified via spatial annotation screening. k UMAP of the scRNA-seq dataset, from the same mouse model as the Visium sample. l and m Visualization of Tangram ..."

DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/Tangram/tangram_recreate.py`

√ verified run · exit 0 · 24 s

Tangram · 3 variations



v1 celltype_projection

SPATIAL



v2 argmax_vs_clusters

SPATIAL



v3 training_scores

SPATIAL

PCAtools

#88

OF 100 · METHODS MENTIONS

R | PCA | DE, QC & SET OVERLAPS | HEATMAPS & DOT MATRICES

48	0.27%	48	dedicated plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Biplots, scree, eigencor plots

THIS RECREATION

PCAtools on Zeisel 2015 pseudobulk (C1 chip x cell type, >=10 cells): scree + Horn/elbow, biplot, eigencor, pairs, loadings

KEY PLOTTING CALL

RECREATIONS/PCATOOLS/PCATOOLS_RECREATE.R — EXCERPT · ENDS LINE 10

```
md <- unique(so@meta.data[, c("pb","chip","celltype","tissue")]); rownames(md) <- md$pb; md <- md[colnames(pb), ]; md$n_cells
<- as.numeric(table(so$pb)[rownames(md)]); md$log_lib <- log10(colSums(pb))
lc <- cpm(pb[rowSums(pb >= 10) >= 5, ], log = TRUE, prior.count = 2); p <- pca(lc, metadata = md, removeVar = .9)
horn <- parallelPCA(lc, removeVar = .9); elbow <- findElbowPoint(p$variance)
ggsave("v1_screeplot.png", screeplot(p, components = getComponents(p, 1:15), vline = c(horn$n, elbow), title = "Scree (Horn +
elbow)") + geom_label(aes(x = horn$n + 1, y = 50, label = "Horn", vjust = -1, size = 8)) + geom_label(aes(x = elbow + 1, y =
50, label = "Elbow", vjust = -1, size = 8)), width = 8, height = 5, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 4 TEXT

Metabolic switch in the aging astrocyte supported via integrative approach comprising network and transcriptome analyses. (Aging (Albany NY), 2023). ,

"...In the same manner, PCA was carried out in R by using the library PCAtools (https://github.com/kevinblighe/PCAtools)."

DOI ↗ Figure 4 on PMC ↗

Figure 1 TEXT

Brain transcriptomics reveal the activation of neuroinflammation pathways during acute Orientia tsutsugamushi infection in mice. *Front Immunol, 2023*

"...Principal component analysis and volcano plot were done in R using PCAtools (34) and EnhancedVolcano (35) packages."

DOI ↗ Figure 1 on PMC ↗

Fig. 1 TEXT

Translatome analysis of tuberous sclerosis complex 1 patient-derived neural progenitor cells reveals rapamycin-dependent and independent alterations. *Mol Autism, 2023*

"...parameters removeVar = 0.75 and scale = T) and visualized using eigencorplot, screeplot and biplot functions from the PCAtools R package (v.2.4.0;"

DOI ↗ Fig. 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/PCAtools/PCAtools_recreate.R`

✓ verified run · exit 0 · 21 s

PCAtools · 4 variations



v1 screeplot · DE



v2 biplot · DE



v3 eigencorplot · HEATMAPS



v4 pairsplot · DE

tradeSeq

#91

OF 100 · METHODS MENTIONS

R | TRAJECTORY DE | TRAJECTORY, PSEUDOTIME & VELOCITY

46	0.26%	51	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

BEST FOR

Smoothed gene expression along lineages

THIS RECREATION

tradeSeq GAM smoothers along slingshot lineages (La Manno 2018 dentate gyrus): smoothed gene trends, association-test heatmap of top dynamic genes, and a single-gene smoother on UMAP. Uses slingshot's fit (runs it if absent).

KEY PLOTTING CALL

RECREATIONS/TRADESEQ/TRADESEQ_RECREATE.R — EXCERPT · ENDS LINE 13

```
show <- head(unique(c(intersect(c("Dcx","Sox11","Calb1","Eomes","Neurod1","Prox1"), genes), top)), 4)
ps <- lapply(show, function(g) plotSmoothers(fit, cnt[genes, ], gene = g, alpha = .4, border = TRUE) + ggtitle(g))
ggsave("v1_plotSmoothers.png", wrap_plots(ps, ncol = 2), width = 10, height = 7.5, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 5 — FIGURE CAPTION

A topographic atlas defines developmental origins of cell heterogeneity in the human embryonic lung. *Nat Cell Biol, 2023*

"...es module 1: 0.60, module 2: 0.69, module 3: 0.84, module 4: 0.57, module 5: 0.80, module 6: 0.73, module 7: 0.61, module 8: 0.55, module 9: 0.85) along the two trajectories, shown in a, according to tradeSeq ..."

DOI ↗ Fig. 5 on PMC ↗

Extended Data Fig. 6 — FIGURE CAPTION

Multiplexing cortical brain organoids for the longitudinal dissection of developmental traits at single-cell resolution. *Nat Methods, 2025*

"...of cells per annotated cell type (shown as reference on the right side), weighted edges among PAGA clusters encode their transcriptional similarity. b) Plot of smoothed gene expression - obtained via tradeseq ..."

DOI ↗ Extended Data Fig. 6 on PMC ↗

Fig. 6 — FIGURE CAPTION

Inferring histology-associated gene expression gradients in spatial transcriptomic studies. *Nat Commun, 2024*

"...Statistical comparison of the test statistic from SPATA2's spatial gradient screening with those of analogous methods PseudotimeDE and tradeSeq.a–c Scatterplots display the relationship between different test ..."

DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/tradeSeq/tradeSeq_recreate.R`

✓ verified run · exit 0 · 12 s

tradeSeq · 3 variations



v1 plotSmoothers

TRAJECTORY



v2 dynamic_gene_heatmap

TRAJECTORY



v3 plotGeneCount

TRAJECTORY

LIANA

#92

OF 100 · METHODS MENTIONS

R/PYTHON | CELL-CELL COMMUNICATION | CELL-CELL COMMUNICATION

45	0.25%	67	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · LIANA+

Neuro-corpus mentions per year (*2026 partial) · niche rank **#23**, growth **3.78×**, visual 4/5, score 0.77. Broad-corpus rate 0.836 → 3.188 per 1k papers (2023 → 2025–26).
GitHub saezlab/liana-py · created — · last push —

BEST FOR

L-R consensus dot plots / tileplots

THIS RECREATION

LIANA+ consensus ligand-receptor inference on Zeisel cortex/hippocampus (mouse consensus resource)

KEY PLOTTING CALL

RECREATIONS/LIANA_PLUS/LIANA_RECREATE.PY — EXCERPT · ENDS LINE 8

```
ad=load(); ad.obs["celltype"]=ad.obs.celltype.astype(str).astype("category")
li.mt.rank_aggregate(ad,groupby="celltype",resource_name="mouseconsensus",expr_prop=.1,use_raw=False,verbose=False,n_perms=200,seed=1)
p=li.pl.dotplot(adata=ad,colour="magnitude_rank",size="specificity_rank",inverse_size=True,inverse_colour=True,source_labels=["astrocytes_ependymal","microglia","endothelial-mural"],target_labels=["interneurons","pyramidal CA1","pyramidal SS","oligodendrocytes"],top_n=20,orderby="magnitude_rank",orderby_ascending=True,figure_size=(10,7))
p.save("v1_dotplot.png",dpi=130,verbose=False)
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Extended Data Fig. 9 — FIGURE CAPTION

Axonal injury is a targetable driver of glioblastoma progression. *Nature, 2025*

"...- mice. The dashed line represents equal proportions in both genotypes. Cell types with PPearson's chi-squared test <0.05 and relative difference > 10% were considered significantly different (§). e, LIANA ..."

DOI ↗ Extended Data Fig. 9 on PMC ↗

Extended Data Fig. 9 — FIGURE CAPTION

Longitudinal single-cell multiomic atlas of high-risk neuroblastoma reveals chemotherapy-induced tumor microenvironment rewiring. *Nat Genet, 2025*

"...nt macrophage subsets from DX and PTX samples, upregulated post-therapy using logistic regression with no multiple comparison adjustment (Methods) e) Cell-cell interaction analysis using CellChat and LIANA ..."

DOI ↗ Extended Data Fig. 9 on PMC ↗

Figure 3 — TEXT

Tumor heterogeneity and tumor-microglia interactions in primary and recurrent IDH1-mutant gliomas. *Cell Rep Med, 2023*

"...Using LIANA,26 we predicted enriched receptor-ligand interactions between tumor populations (astro-like and OPC-like) and TAM subpopulations (Figures 3F and 3G)."

DOI ↗ Figure 3 on PMC ↗

LIMITATIONS & NOTES

- v2 is a LIANA circle plot because tileplot is broken with current plotnine/mizani (script comment).
- Run this first: its tables feed pyCirclize, CellPhoneDB_ktplots and circlize_chord (METHODS §9).

Reproduce: `python3 recreations/LIANA_plus/liana_recreate.py`

√ verified run · exit 0 · 24 s

LIANA · 3 variations



v1 dotplot

CELL–CELL



v2 circle_plot

CELL–CELL



v3 interaction_count_heatmap

CELL–CELL

BayesSpace

#97

OF 100 · METHODS MENTIONS

R · SPATIAL CLUSTERING · SPATIAL & TISSUE MAPS

41	0.23%	67	analysis+plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · BAYESSPACE

Neuro-corpus mentions per year (*2026 partial) · niche rank **#32**, growth **2.19×**, visual 4/5, score 0.671. Broad-corpus rate 1.289 → 2.832 per 1k papers (2023 → 2025–26).
GitHub edward130603/BayesSpace · created 2020-06-23 · last push 2026-01-08

BEST FOR

Enhanced-resolution spatial cluster maps

THIS RECREATION

BayesSpace on Maynard 2021 human DLPFC (151673): spot clusters, enhanced sub-spot resolution, enhanced gene maps

KEY PLOTTING CALL

RECREATIONS/BAYESSPACE/BAYESSPACE_RECREATE.R — EXCERPT · ENDS LINE 8

```
sce <- spatialPreprocess(sce, platform = "Visium", n.PCs = 7, n.HVGs = 2000, log.normalize = FALSE)
sce <- spatialCluster(sce, q = 7, platform = "Visium", d = 7, init.method = "mclust", model = "t", gamma = 2, nrep = 5000, burn.
in = 500)
pal <- c("#E41A1C","#377EB8","#4DAF4A","#984EA3","#FF7F00","#FFFF33","#A65628")
ggsave("v1_clusterPlot_vs_truth.png", patchwork::wrap_plots(list(clusterPlot(sce, palette = pal, color = NA) + ggtitle("BayesSp
ace (q=7)"), clusterPlot(sce, label = "ground_truth", palette = c(pal, "grey"), color = NA) + ggtitle("Manual layers (Maynard
2021)")), nrow = 1), width = 11, height = 5, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 4 — FIGURE CAPTION

BANKSY unifies cell typing and tissue domain segmentation for scalable spatial omics data analysis. *Nat Genet, 2024*

"...use hippocampus.a, BANKSY separates CA3 neurons from SSC neurons. b–h, Comparisons with nonspatial clustering (λ = 0) and several spatial clustering methods (b, nonspatial; c, MERINGUE; d, Giotto; e ..."

DOI ↗ Fig. 4 on PMC ↗

Fig. 2 — FIGURE CAPTION

Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST. *Nat Commun, 2023*

"...quartiles, and whiskers denote the 1.5× interquartile range. B H&E image and manual annotation from the original study. C Clustering results by nonspatial and spatial methods, Seurat, Giotto, SpaGCN ..."

DOI ↗ Fig. 2 on PMC ↗

Figure 2—figure supplement 5 — FIGURE CAPTION

The gene expression landscape of the human locus coeruleus revealed by single-nucleus and spatially-resolved transcriptomics. *Elife, 2024*

"...ve technical variation in the molecular measurements between the n=8 Visium samples from 4 donors. The integrated measurements were subsequently used as the input for spatially-aware clustering using ..."

DOI ↗ Figure 2—figure supplement 5 on PMC ↗

LIMITATIONS & NOTES

- Enhanced-gene maps use the same per-gene PC regression that enhanceFeatures(nrounds=0) uses; the full sub-spot assay runs out of memory (METHODS §9).

Reproduce: `Rscript recreations/BayesSpace/BayesSpace_recreate.R`

√ verified run · exit 0 · 159 s

BayesSpace · 3 variations



v1 clusterPlot_vs_truth

SPATIAL



v2 enhanced_subspot

SPATIAL



v3 enhanced_genes

SPATIAL

CellRank

#98

OF 100 · METHODS MENTIONS

PYTHON | FATE MAPPING | TRAJECTORY, PSEUDOTIME & VELOCITY

40	0.23%	51	analysis+plotting	4
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS



NICHE & COOL · CELLRANK

Neuro-corpus mentions per year (*2026 partial) · niche rank **#35**, growth **1.55×**, visual 5/5, score 0.658. Broad-corpus rate 1.428 → 2.223 per 1k papers (2023 → 2025–26).
GitHub theislab/cellrank · created — · last push —

BEST FOR

Fate probabilities, driver genes, circular projections

THIS RECREATION

CellRank 2 on dentate gyrus: macrostates, fate probabilities, circular projection, driver-gene trends

KEY PLOTTING CALL

RECREATIONS/CELLRANK/CELLRANK_RECREATE.PY — EXCERPT · ENDS LINE 6

```
a=load(); vk=cr.kernels.VelocityKernel(a).compute_transition_matrix(); ck=cr.kernels.ConnectivityKernel(a).compute_transition_m
atrix(); k=0.8*vk+0.2*ck
g=cr.estimators.GPCCA(k); g.fit(cluster_key="clusters",n_states=[4,10]);
g.plot_macrostates(which="all",basis="umap",legend_loc="right",s=60,show=False);
plt.gcf().savefig("v1_macrostates.png",dpi=150,bbox_inches="tight"); plt.close("all")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 4 — FIGURE CAPTION

Mapping cells through time and space with moscot. *Nature, 2025*

"…f moscot compared with TOME7. c, Mapping heart cells across time points (bottom) and ground-truth annotation of the heart lineage (top). d, Heart-lineage driver genes found by interfacing moscot with CellRank …"

DOI ↗ Fig. 4 on PMC ↗

Extended Data Fig. 2 — FIGURE CAPTION

Inferring and perturbing cell fate regulomes in human brain organoids. *Nature, 2023*

"…ster identities, expression patterns of cluster markers. Volcano plot shows differentially expressed (DE) genes of cluster 5 relative to other clusters. f, UMAP embedding coloured by rank-transformed CellRank …"

DOI ↗ Extended Data Fig. 2 on PMC ↗

Fig. 6 — FIGURE CAPTION

Tracking single-cell evolution using clock-like chromatin accessibility loci. *Nat Biotechnol, 2025*

"…EpiTrace reveals the developmental history during human cortical gyrification.a, UMAP projected cell evolution trajectory built with CellRank by using a hybrid kernel of EpiTrace, CytoTRACE and RNA velocity of …"

DOI ↗ Fig. 6 on PMC ↗

LIMITATIONS & NOTES

- v4 shows lineage drivers because GAM gene-trend models fail with current pygam/scipy (script comment).

Reproduce: `python3 recreations/CellRank/cellrank_recreate.py`

✓ verified run · exit 0 · 21 s

CellRank · 4 variations



v1 macrostates

TRAJECTORY



v2 fate_probabilities

TRAJECTORY



v3 circular_projection

TRAJECTORY



v4 lineage_drivers

TRAJECTORY

Nebulosa

#99

OF 100 · METHODS MENTIONS

R · SINGLE-CELL DENSITY · EMBEDDINGS & FEATURE MAPS

39	0.22%	43	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · NEBULOSA

Neuro-corpus mentions per year (*2026 partial) · niche rank **#58**, growth **1.29×**, visual 4/5, score 0.524. Broad-corpus rate 1.811 → 2.337 per 1k papers (2023 → 2025–26).
GitHub powellgenomicslab/Nebulosa · created 2020-07-12 · last push 2026-02-06

BEST FOR

Kernel-density feature plots

KEY PLOTTING CALL

RECREATIONS/NEBULOSA/NEBULOSA_RECREATE.R — EXCERPT · ENDS LINE 4

```
so <- readRDS("../_data/zeisel_seurat.rds")
ggsave("v1_density_single.png", plot_density(so, c("Pvalb","Sst","Vip","Aqp4")) + plot_layout(ncol = 2), width = 9, height = 8,
dpi = 120, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Figure 2 — FIGURE CAPTION
A single-nucleus transcriptomic atlas of medium spiny neurons in the rat nucleus accumbens. *Sci Rep, 2024*
"...Low resolution clustering of MSNs. (a) UMAP dimension reduction plot of MSN clustered at low resolution. Nuclei are colored and labeled by major MSN population. Nebulosa plots visualizing expression on the MSN ..."
DOI ↗ Figure 2 on PMC ↗

Figure 1 — TEXT
Landscape of mast cell populations across organs in mice and humans. *J Exp Med, 2023*
"...We showed these differences by plotting a density plot based on gene-weighted kernel density estimation realized with Nebulosa packages (v1.8.0;"
DOI ↗ Figure 1 on PMC ↗

Fig. 3 — TEXT
BANKSY unifies cell typing and tissue domain segmentation for scalable spatial omics data analysis. *Nat Genet, 2024*
"...We used Nebulosa (v.1.0.2)70 to visualize the cycling metagene on the UMAP embedding of the BANKSY matrix."
DOI ↗ Fig. 3 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/Nebulosa/Nebulosa_recreate.R`

√ verified run · exit 0 · 21 s

Nebulosa · 3 variations



v1 density_single

EMBEDDINGS



v2 joint_density

EMBEDDINGS



v3 density_vs_featureplot

EMBEDDINGS

ggalluvial

R · FLOW/ALLUVIAL · COMPOSITION, DISTRIBUTIONS & LAYOUTS

#100

OF 100 · METHODS MENTIONS

36	0.2%	37	dedicated plotting	3
corpus papers naming it in METHODS	of 17,768 papers	papers mentioning it anywhere in full text	role in catalog	variations recreated

POSITION IN TOP 100
BAR HEIGHT = LOG PAPERS

NICHE & COOL · GGSANKEY

Neuro-corpus mentions per year (*2026 partial) · niche rank **#11**, growth **10.71×**, visual 4/5, score 0.845. Broad-corpus rate 0.035 → 0.47 per 1k papers (2023 → 2025–26).
GitHub davidsjoberg/ggsankey · created 2021-03-25 · last push 2024-04-04 · also scored: ggalluvial (#72, growth 0.84×)

BEST FOR

Alluvial/Sankey plots

THIS RECREATION

Flow plots of Zeisel hierarchy: tissue → level1 class → level2 subclass

KEY PLOTTING CALL

RECREATIONS/GGSANKEY_GGALLUVIAL/SANKEY_RECREATE.R — EXCERPT · ENDS LINE 8

```
md <- readRDS("../_data/zeisel_seurat.rds")@meta.data
top2 <- names(sort(table(md$level2class), decreasing = TRUE))[1:18]; md$sub <- ifelse(md$level2class %in% top2, md$level2class,
"other")
d <- make_long(md, tissue, celltype, sub)
g <- ggplot(d, aes(x = x, next_x = next_x, node = node, next_node = next_node, fill = factor(node), label = node)) +
geom_sankey(flow.alpha = .55, node.color = "grey30") + geom_sankey_label(size = 2.6, color = "black", fill = "white") + theme_s
ankey(base_size = 12) + theme(legend.position = "none") + labs(x = NULL, title = "ggsankey: tissue -> class -> subclass")
ggsave("v1_ggsankey.png", g, width = 10, height = 7, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 2 TEXT
The transcriptomes, connections and development of submucosal neuron classes in the mouse small intestine. *Nat Neurosci, 2025*
"...Finally, the mapping identities were presented as Alluvial plot using the R package ggalluvial (Fig."
DOI ↗ Fig. 2 on PMC ↗

Fig. 4 TEXT
Single-cell transcriptomic analysis reveals diversity within mammalian spinal motor neurons. *Nat Commun, 2023*
"...To intuitively visualize the relationships of LMC subclusters between brachial and lumbar segments, we created an alluvial diagram using the ggalluvial R package ..."
DOI ↗ Fig. 4 on PMC ↗

Fig. 4 TEXT
Microbial signals in primary and metastatic brain tumors. *Nat Med, 2025*
"...Alluvial plots demonstrating the removal of taxa across the different filtration steps were generated using ggalluvial (version 0.12.5)."
DOI ↗ Fig. 4 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggsankey_ggalluvial/sankey_recreate.R`

√ verified run · exit 0 · 7 s

ggalluvial · 3 variations

RECREATIONS/GGSANKEY_GGALLUVIAL/



v1 ggsankey

COMPOSITION



v2 ggsankey_alluvial

COMPOSITION



v3 ggalluvial

COMPOSITION



PART II · 36 PACKAGES · 112 RECREATED FIGURES

novaflow

II

Niche & Cool

Newer or more specialised visualisation tools, ranked by **0.5 × growth percentile + 0.5 × visual score/5**. These are the figures that make a reviewer stop: circos and chord layouts, 3-D atlases, optimal-transport flows, Bayesian composition. This part has the 36 recreated folders that are not covered by Part I. The other 19 recreated niche tools are already in Part I, and their spreads show their niche stats.

36
spreads

112
figures

58 / 73
niche tools recreated

324.85×
max growth ratio

IN THIS PART · RANK · PAGE

1	pyCirclize	144	17	scDRS	168	50	Palantir	192
3	brainrender	146	19	MNE-Python	170	51	ggspavis/SpatialExperime	194
4	COMMOT	148	20	sccomp	172	55	plotgardener	196
6	MultiNicheNet	150	26	PyComplexHeatmap	174	60	propeller (speckle)	198
7	moscot	152	29	nilearn	176	63	ggcoverage	200
8	scplotter	154	31	SCpubr	178	64	ggtree/ggtreeExtra	202
10	Banksy	156	34	UCell	180	65	Mellon	204
12	scTour	158	37	spatialLIBD	182	66	karyoploteR	206
13	plot1cell	160	41	ggrepel volcano styles	184	67	marsilea	208
14	veloVI	162	42	Augur	186	68	dittoSeq	210
15	miloR / Milo	164	47	ggridges	188	70	ggforce	212
16	Datashader	166	48	plotnine	190	71	gggenes	214

pyCirclize

#1

OF 73 · NICHE & COOL SCORE

PYTHON | CIRCOS/CHORD | CELL–CELL COMMUNICATION

20.05×	5/5	0.98	0	1,127
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE



NICHE & COOL · PYCIRCLIZE

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.191 per 1k papers (2023 → 2025–26).
GitHub moshi4/pyCirclize · created 2022-12-17 · last push 2026-06-14

BEST FOR

circos/chord in Python

THIS RECREATION

pyCirclize chord + circos ring plots from LIANA+ results on Zeisel (run ../LIANA_plus first)

KEY PLOTTING CALL

RECREATIONS/PYCIRCLIZE/PYCIRCLIZE_RECREATE.PY — EXCERPT · ENDS LINE 6

```
m=pd.read_csv("../LIANA_plus/lr_counts_matrix.csv",index_col=0); idx=sorted(set(m.index)|set(m.columns));
m=m.reindex(index=idx,columns=idx,fill_value=0)
c=Circos.chord_diagram(m,space=3,cmap="tab10",label_kws=dict(size=8,r=110),link_kws=dict(ec="black",lw=.3,direction=1))
c.savefig("v1_chord_interactions.png",dpi=150)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/moshi4/pyCirclize.

LIMITATIONS & NOTES

- Reads LIANA+ result tables, so run recreations/LIANA_plus first (METHODS §9).

Reproduce: `python3 recreations/pyCirclize/pycirclize_recreate.py`

√ verified run · exit 0 · 15 s

LEAD FIGURE



v1 chord_interactions — CELL–CELL



v2 ligand_receptor_chord

● CELL-CELL

v3 circos_tracks

● CELL-CELL

PART II · NICHE & COOL · 02 / 36

brainrender

#3

OF 73 · NICHE & COOL SCORE

PYTHON | 3D BRAIN | BRAIN ATLAS, 3D & NEUROIMAGING

4.81×	5/5	0.914	4	680
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · BRAINRENDER

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.038 per 1k papers (2023 → 2025–26).
GitHub brainglobe/brainrender · created 2019-09-19 · last push 2026-10-13

BEST FOR

3D Allen CCF renderings

THIS RECREATION

brainrender 3D renders of the Allen Mouse CCFv3 (25 um, via BrainGlobe): coloured region meshes, point clouds sampled from real CCF annotation voxels, and a sliced view. brainrender builds the scene (atlas meshes, colours); the final frame is drawn with a plain vedo Plotter without depth peeling, because software-GL offscreen rendering blanks translucent meshes when peeling is on.

KEY PLOTTING CALL

RECREATIONS/BRAINRENDER/BRAINRENDER_RECREATE.PY — EXCERPT · ENDS LINE 28

```
regs={"Isocortex":"#9ecae1","HIP":"#fb6a4a","STR":"#74c476","TH":"#fdae6b","HY":"#9e9ac8","CB":"#fdd0a2","MB":"#a1d99b","OLF":"#fa9fb5"}
# CCF axes: x=AP (anterior->posterior), y=DV (dorsal->ventral), z=ML
s=Scene(atlas_name="allen_mouse_25um")
for r,c in regs.items(): s.add_brain_region(r, color=c, alpha=.45 if r=="Isocortex" else .95)
draw(s, "v1_regions_3quarter.png", "Allen CCFv3 major divisions (brainrender)", azimuth=-35, elevation=25, roll=180, zoom=1.3)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/brainglobe/brainrender.

LIMITATIONS & NOTES

- brainrender builds the scene, but the final frame is drawn with a plain vedo Plotter without depth peeling: software-GL offscreen rendering blanks translucent meshes when peeling is on (METHODS §9).

Reproduce: `python3 recreations/brainrender/brainrender_recreate.py`

✓ verified run · exit 0 · 29 s

brainrender · 3 variations



v1 regions_3quarter

BRAIN



v2 hippocampus_points

BRAIN



v3 coronal_cut

BRAIN

COMMOT

#4

OF 73 · NICHE & COOL SCORE

PYTHON | SPATIAL COMMUNICATION | SPATIAL & TISSUE MAPS

4.67×	5/5	0.904	15	149
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · COMMOT

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.348 → 1.664 per 1k papers (2023 → 2025–26).
GitHub zcang/COMMOT · created 2021-09-03 · last push 2026-06-25

BEST FOR

spatial signalling flow vector fields

THIS RECREATION

COMMOT spatial optimal-transport signalling on 10x Visium mouse brain (CellChat mouse secreted LR database)

KEY PLOTTING CALL

RECREATIONS/COMMOT/COMMOT_RECREATE.PY — EXCERPT · ENDS LINE 12

```
for meth,fn in [("grid","v1_flow_grid.png"),("stream","v2_flow_stream.png")]:
    ct.pl.plot_cell_communication(v,database_name="cellchat",pathway_name=pw,plot_method=meth,background_legend=True,scale=.0000
3,ndsize=8,grid_density=.4,summary="sender",background="image",clustering="cluster",cmap="Alphabet",normalize_v=True,normalize_
v_quantile=.995)
    plt.title(f"COMMOT {pw} signalling flow ({meth})"); plt.savefig(fn,dpi=140,bbox_inches="tight"); plt.close("all")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/zcang/COMMOT.

LIMITATIONS & NOTES

- NumPy 2 shim: np.Inf restored because COMMOT still uses it (METHODS §9).

Reproduce: `python3 recreations/COMMOT/commot_recreate.py`

√ verified run · exit 0 · 1,095 s

LEAD FIGURE

COMMOT PTN signalling flow (grid)

v1 flow_grid

SPATIAL

COMMOT · 3 variations · v1 on facing page



v2 flow_stream

SPATIAL



v3 sender_maps

SPATIAL

PART II · NICHE & COOL · 04 / 36

MultiNicheNet

#6

OF 73 · NICHE & COOL SCORE

R · LIGAND-TARGET · CELL-CELL COMMUNICATION

89.9×	4/5	0.893	13	202
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · MULTINICHENET

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.889 per 1k papers (2023 → 2025–26).
GitHub saeyslab/multinichenetr · created 2021-02-16 · last push 2025-08-12

BEST FOR

multi-sample LR circos

THIS RECREATION

MultiNicheNet (multi-sample, multi-condition cell-cell communication) on Zeisel 2015: S1 cortex vs CA1, C1 chip as sample (pseudobulk, edgeR), shared non-pyramidal cell types. NicheNet v2 mouse priors (Zenodo 7074291). Group-comparison circos, sample-level LR product + ligand activity bubble plot, ligand-activity/target plot.

KEY PLOTTING CALL

RECREATIONS/MULTINICHENET/MULTINICHENET_RECREATE.R — EXCERPT · ENDS LINE 23

```
top <- get_top_n_lr_pairs(out$prioritization_tables, 50, rank_per_group = FALSE)
cts <- sort(unique(c(top$sender, top$receiver))); cols <- setNames(RColorBrewer::brewer.pal(max(3, length(cts)), "Set2")[seq_along(cts)], cts)
png("v1_circos_%d.png", 1400, 1400, res = 170); circ <- make_circos_group_comparison(top, cols[sort(unique(top$sender))], cols[sort(unique(top$receiver))]); dev.off()
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/saeyslab/multinichenetr.

LIMITATIONS & NOTES

- On Zeisel, MultiNicheNet keeps only 2 receivers (astrocytes, oligodendrocytes) because chips have few cells per type (SUMMARY, data caveats).
- Run single-core: parallel workers died under memory pressure on the shared box (script comment).

Reproduce: `Rscript recreations/MultiNicheNet/MultiNicheNet_recreate.R`

√ verified run · exit 0 · 922 s

MultiNicheNet · 3 variations



v1 circos_group_comparison

CELL–CELL



v2 sample_lr_activity

CELL–CELL



v3 ligand_activity_tiles

CELL–CELL

moscot

#7

OF 73 · NICHE & COOL SCORE

PYTHON | OPTIMAL TRANSPORT | TRAJECTORY, PSEUDOTIME & VELOCITY

42.91×	4/5	0.887	7	217
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · MOSCOT

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.419 per 1k papers (2023 → 2025–26).
GitHub theislab/moscot · created 2021-08-23 · last push 2026-10-14

BEST FOR

OT cell-transition maps, Sankey

THIS RECREATION

moscot optimal-transport TemporalProblem on the La Manno 2018 dentate gyrus (P12 → P35): cell-type transition matrix, Sankey of fate flow, and push-forward of the P12 neuroblast mass onto P35 cells.

KEY PLOTTING CALL

RECREATIONS/MOSCOT/MOSCOT_RECREATE.PY — EXCERPT · ENDS LINE 12

```
tp=mt.problems.TemporalProblem(a).prepare(time_key="day",joint_attr="X_pca").solve(epsilon=1e-2,tau_a=.95,tau_b=.95)
tp.cell_transition(source=12.0,target=35.0,source_groups="clusters",target_groups="clusters",forward=True,key_added="ct")
fig=mtp.cell_transition(tp,key="ct",figsize=(9,8),dpi=130,return_fig=True,cmap="magma_r"); fig.suptitle("moscot: P12 -> P35
cell-type transition probabilities"); fig.savefig("v1_cell_transition.png",bbox_inches="tight",dpi=130)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/theislab/moscot.

LIMITATIONS & NOTES

- The Sankey is re-drawn from the OT coupling because the built-in Sankey mis-draws (METHODS §9).

Reproduce: `python3 recreations/moscot/moscot_recreate.py`

✓ verified run · exit 0 · 14 s

LEAD FIGURE



v3 push_umap

TRAJECTORY



v1 cell_transition

TRAJECTORY



v2 sankey

TRAJECTORY

PART II · NICHE & COOL · 06 / 36

scplotter

#8

OF 73 · NICHE & COOL SCORE

R · SINGLE-CELL PUBLICATION PLOTS · EMBEDDINGS & FEATURE MAPS · COMPOSITION, DISTRIBUTIONS & LAYOUTS

17.51×	4/5	0.872	1	333
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES
SORTED, LOG SCALE

NICHE & COOL · SCPLOTTER

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.165 per 1k papers (2023 → 2025–26).
GitHub pwwang/scplotter · created 2024-09-03 · last push 2026-09-19

BEST FOR

one-line Seurat publication plots (plotthis backend)

THIS RECREATION

scplotter (pwwang; plotthis backend) on Zeisel 2015: hull-marked and hex CellDimPlot, CellStatPlot as circos / radar / ring of cell-type composition by tissue, and FeatureStatPlot ridge/violin of markers. Installed from GitHub (pwwang/plotthis, pwwang/scplotter); not on CRAN for R 4.5.

KEY PLOTTING CALL

RECREATIONS/SCPLOTTER/SCPLOTTER_RECREATE.R — EXCERPT · ENDS LINE 10

```
so <- readRDS("../_data/zeisel_seurat.rds"); so$tissue <- factor(ifelse(so$tissue == "sscortex", "S1 cortex", "CA1"))
so$chip <- paste0("chip", sub("_.*", "", colnames(so))); Idents(so) <- "celltype"
p1 <- CellDimPlot(so, group_by = "celltype", reduction = "umap", theme = "theme_blank", add_mark = TRUE, mark_type = "hull", la
bel = TRUE, label_insitu = TRUE, title = "CellDimPlot with hull marks")
p1b <- CellDimPlot(so, group_by = "celltype", reduction = "umap", theme = "theme_blank", hex = TRUE, hex_bins = 40, title = "hex
-binned")
ggsave("v1_celldimplot.png", p1 + p1b, width = 13, height = 5.5, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/pwwang/scplotter.

LIMITATIONS & NOTES

- Installed from GitHub (pwwang/plotthis, pwwang/scplotter); not on CRAN for R 4.5 (script header).

Reproduce: `Rscript recreations/scplotter/scplotter_recreate.R`

√ verified run · exit 0 · 24 s



v1 celldimplot

EMBEDDINGS



v2 cellstat_circos

COMPOSITION



v3 cellstat_radar_ring

COMPOSITION



v4 featurestat_ridge

COMPOSITION

Banksy

#10

OF 73 · NICHE & COOL SCORE

R/PYTHON | SPATIAL CLUSTERING | SPATIAL & TISSUE MAPS

14.16×	4/5	0.859	21	167
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · BANKSY

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.07 → 1.118 per 1k papers (2023 → 2025–26).
GitHub prabhakarlab/Banksy · created 2021-05-23 · last push 2026-09-29

BEST FOR

neighbourhood-augmented domains

THIS RECREATION

BANKSY spatial domains (neighbourhood-augmented clustering) on DLPFC 151673

KEY PLOTTING CALL

RECREATIONS/BANKSY/BANKSY_RECREATE.R — EXCERPT · ENDS LINE 11

```
spe <- runBanksyPCA(spe, use_agf = TRUE, lambda = c(0, .2), npcs = 20, seed = 1000)
spe <- clusterBanksy(spe, use_agf = TRUE, lambda = c(0, .2), resolution = .6, seed = 1000)
cn <- clusterNames(spe); spe <- connectClusters(spe)
spe$nonspatial <- factor(colData(spe)[[cn[1]]]); spe$banksy <- factor(colData(spe)[[cn[2]]])
ggsave("v1_banksy_vs_nonspatial.png", patchwork::wrap_plots(list(plotCoords(spe, annotate = "nonspatial", point_size = .6) + gg
title("lambda = 0 (non-spatial)"), plotCoords(spe, annotate = "banksy", point_size = .6) + ggtitle("BANKSY lambda = 0.2"), plot
Coords(spe, annotate = "ground_truth", pal = "libd_layer_colors", point_size = .6) + ggtitle("manual layers")), nrow = 1),
width = 15, height = 5, dpi = 120, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/prabhakarlab/Banksy.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/Banksy/Banksy_recreate.R`

√ verified run · exit 0 · 56 s

LEAD FIGURE



v1 banksy_vs_nonspatial

SPATIAL

Banksy · 2 variations · v1 on facing page



v2 banksy_umap

SPATIAL

scTour

#12

OF 73 · NICHE & COOL SCORE

PYTHON | TRAJECTORY | TRAJECTORY, PSEUDOTIME & VELOCITY

6.52×	4/5	0.832	7	89
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · SCTOUR

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.104 → 0.737 per 1k papers (2023 → 2025–26).
GitHub LiQian-XC/sctour · created 2022-01-03 · last push 2025-07-20

BEST FOR

vector field + pseudotime

THIS RECREATION

scTour (VAE + neural ODE) on La Manno 2018 dentate gyrus: learned pseudotime, latent-space vector field streamlines, and pseudotime distribution per cell type.

KEY PLOTTING CALL

RECREATIONS/SCTOUR/SCTOUR_RECREATE.PY — EXCERPT · ENDS LINE 18

```
fig,axs=plt.subplots(1,2,figsize=(13,5.5))
sc.pl.umap(a,color="clusters",ax=axs[0],show=False,legend_loc="on data",legend_fontsize=7,frameon=False,title="scTour latent UMAP")
sc.pl.umap(a,color="ptime",ax=axs[1],show=False,cmap="Spectral_r",frameon=False,title="scTour pseudotime")
fig.tight_layout(); fig.savefig("v1_pseudotime_umap.png",dpi=140)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/LiQian-XC/sctour.

LIMITATIONS & NOTES

- Dense X is used because scTour calls the removed scipy .A on sparse views (METHODS §9).

Reproduce: `python3 recreations/scTour/sctour_recreate.py`

✓ verified run · exit 0 · 89 s

LEAD FIGURE



v1 pseudotime_umap

TRAJECTORY

scTour · 3 variations · v1 on facing page



v2 vector_field

TRAJECTORY



v3 ptime_by_celltype

TRAJECTORY

plot1cell

#13

OF 73 · NICHE & COOL SCORE

R · SINGLE-CELL PUBLICATION PLOTS · EMBEDDINGS & FEATURE MAPS · COMPOSITION, DISTRIBUTIONS & LAYOUTS · HEATMAPS & DOT MATRICES

2.7×	5/5	0.829	6	131
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · PLOT1CELL

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.139 → 0.394 per 1k papers (2023 → 2025–26).
GitHub TheHumphreysLab/plot1cell · created 2022-03-22 · last push 2025-09-04

BEST FOR

circlize 'cell circle' UMAP, multi-group dot plots

THIS RECREATION

plot1cell circular UMAP ("plot_circlize") + tracks, cell-fraction and multi-group dot plots on Zeisel 2015. plot1cell cannot be installed here (hard deps DoubletFinder/loomR/GEOquery/... unavailable for R 4.5), so its plotting functions are sourced verbatim from the GitHub repo (TheHumphreysLab/plot1cell, R/*.R) into src/.

KEY PLOTTING CALL

RECREATIONS/PLOT1CELL/PLOT1CELL_RECREATE.R — EXCERPT · ENDS LINE 16

```r
png("v1_circlize_umap.png", 1600, 1600, res = 200)
plot_circlize(cd, do.label = TRUE, pt.size = .5, col.use = cols, bg.color = "#F9F2E4", repel = FALSE, label.cex = .7)
add_track(cd, group = "tissue", colors = c("#1b9e77","#d95f02"), track_num = 2)
add_track(cd, group = "chip", track_num = 3, colors = colorRampPalette(brewer.pal(9, "Greys"))(length(unique(so$chip))))
dev.off()
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/TheHumphreysLab/plot1cell.

LIMITATIONS & NOTES

- plot1cell cannot be installed on R 4.5 (hard deps DoubletFinder/loomR/GEOquery/... unavailable), so its plotting functions are sourced verbatim from the GitHub repo TheHumphreysLab/plot1cell (R/*.R) into src/ (script header).

Reproduce: `Rscript recreations/plot1cell/plot1cell_recreate.R`

√ verified run · exit 0 · 13 s

plot1cell · 3 variations



v1 circlize_umap

EMBEDDINGS



v2 cell_fraction

COMPOSITION



v3 complex_dotplot

HEATMAPS

veloVI

#14

OF 73 · NICHE & COOL SCORE

PYTHON | RNA VELOCITY | TRAJECTORY, PSEUDOTIME & VELOCITY

5.87×	4/5	0.824	9	45
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · VELOVI

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.07 → 0.457 per 1k papers (2023 → 2025–26).
GitHub YosefLab/velovi · created 2022-03-10 · last push 2026-01-12

BEST FOR

velocity uncertainty

THIS RECREATION

veloVI (variational RNA velocity) on dentate gyrus: streamlines, intrinsic uncertainty, per-cell-type velocity coherence

KEY PLOTTING CALL

RECREATIONS/VELOVI/VELOVI_RECREATE.PY — EXCERPT · ENDS LINE 10

```
scv.settings.figdir="."; a=load(); a=preprocess_data(a)
VELOVI.setup_anndata(a,spliced_layer="Ms",unspliced_layer="Mu"); m=VELOVI(a); m.train(max_epochs=300,accelerator="cpu",enable_p
rogress_bar=False)
lt=m.get_latent_time(n_samples=25); sc_=20/lt.max(0); v=m.get_velocity(n_samples=25,velo_statistic="mean")
a.layers["velocity"]=v/sc_; a.layers["fit_t"]=lt.values*sc_.values[None,:]
scv.tl.velocity_graph(a,n_jobs=4)
scv.pl.velocity_embedding_stream(a,basis="umap",color="clusters",title="veloVI streamlines",legend_loc="right margin",save="v1_
velovi_stream.png",show=False)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/YosefLab/velovi.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/veloVI/velovi_recreate.py`

√ verified run · exit 0 · 509 s

LEAD FIGURE



v1 velovi_stream

TRAJECTORY

veloVI · 3 variations · v1 on facing page



v2 uncertainty

TRAJECTORY



v3 coherence_by_cluster

TRAJECTORY

miloR / Milo

#15

OF 73 · NICHE & COOL SCORE

R/PYTHON | DIFFERENTIAL ABUNDANCE | COMPOSITION, DISTRIBUTIONS & LAYOUTS

2.31×	5/5	0.815	46	450
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · MILOR / MILO

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 1.184 → 2.743 per 1k papers (2023 → 2025–26).
GitHub MarioniLab/miloR · created 2020-07-30 · last push 2026-08-25

BEST FOR

DA beeswarm, nhood graph

THIS RECREATION

miloR differential abundance on kNN neighbourhoods, Zeisel 2015: S1 cortex vs CA1 hippocampus, Fluidigm C1 chip (cell-name prefix) as the replicate. Neighbourhood graph, DA beeswarm, nhood size histogram. NOTE: installed from Bioconductor source with src/Makevars patched (forces C++11; current RcppArmadillo needs C++14+).

KEY PLOTTING CALL

RECREATIONS/MILO/MILOR_RECREATE.R — EXCERPT · ENDS LINE 19

```
m <- calcNhoodDistance(m, d = 20, reduced.dim = "PCA")
da <- testNhoods(m, design = ~ tissue, design.df = design, reduced.dim = "PCA", fdr.weighting = "graph-overlap")
da <- annotateNhoods(m, da, coldata_col = "celltype"); write.csv(da, "milo_da_results.csv", row.names = FALSE)
m <- buildNhoodGraph(m)
p1 <- plotReducedDim(m, "UMAP", colour_by = "celltype", point_size = .6) + guides(colour = guide_legend(override.aes =
list(size = 3))) + ggtitle("Zeisel cell types")
p2 <- plotNhoodGraphDA(m, da, layout = "UMAP", alpha = .1) + ggtitle("Milo nhood graph: logFC sscortex vs CA1")
ggsave("v1_nhood_graph_DA.png", p1 + p2 + plot_layout(widths = c(1, 1.1)), width = 13, height = 5.5, dpi = 130, bg = "white")
```

IN THE CORPUS · MOST-CITED OF 29 PAPERS NAMING “MILOR” IN METHODS

94 CITATIONS — **Spatial and single-nucleus transcriptomic analysis of genetic and sporadic forms of Alzheimer's disease.** *Nat Genet, 2024*
DOI ↗ PMC11631771

51 CITATIONS — **Tracking cell-type-specific temporal dynamics in human and mouse brains.** *Cell, 2023*
DOI ↗ PMC10545416

Figure-level example mining (METHODS §3) was run for the Top 100 only, so these are mention-level examples without a verified figure link.

LIMITATIONS & NOTES

- miloR is built from source with the C++11 flag removed from src/Makevars (METHODS §9).
- The Fluidigm C1 chip ID is used as the replicate (METHODS §8).

Reproduce: `Rscript recreations/Milo/miloR_recreate.R`

√ verified run · exit 0 · 26 s

miloR / Milo · 3 variations

RECREATIONS/MILO/



v1 nhood_graph_DA

COMPOSITION



v2 DA_beeswarm

COMPOSITION



v3 nhood_size_hist

COMPOSITION

Datashader

#16

OF 73 · NICHE & COOL SCORE

PYTHON | LARGE DATA | SPATIAL & TISSUE MAPS

4.81×	4/5	0.814	1	3,568
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE



NICHE & COOL · DATASHADER

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.038 per 1k papers (2023 → 2025–26).
GitHub holoviz/datashader · created 2015-12-23 · last push 2026-10-14

BEST FOR

millions of points rasterised

THIS RECREATION

Datashader: all 73,655 MERFISH cells rasterised without overplotting

KEY PLOTTING CALL

RECREATIONS/DATASHADER/DATASHADER_RECREATE.PY — EXCERPT · ENDS LINE 7

```
cv=ds.Canvas(plot_width=900,plot_height=900)
tf.set_background(tf.shade(cv.points(s,"x","y",ds.count()),cmap=cc.fire,how="eq_hist"),"black").to_pil().save("v1_density_fire.
png")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/holoviz/datashader.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/Datashader/datashader_recreate.py`

√ verified run · exit 0 · 7 s

LEAD FIGURE



v3 all_slices_strip

SPATIAL

Datashader · 3 variations · v3 on facing page



v1 density_fire

SPATIAL



v2 categorical_shading

SPATIAL

scDRS

#17

OF 73 · NICHE & COOL SCORE

PYTHON | DISEASE RELEVANCE | EMBEDDINGS & FEATURE MAPS | HEATMAPS & DOT MATRICES | COMPOSITION, DISTRIBUTIONS & LAYOUTS

4.63×	4/5	0.798	37	201
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · SCDRS

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.209 → 1.003 per 1k papers (2023 → 2025–26).
GitHub martinjzhang/scDRS · created 2020-07-10 · last push 2026-08-27

BEST FOR

disease-score UMAPs, trait x cell-type heatmaps

THIS RECREATION

scDRS disease-relevance scores on Zeisel 2015 mouse cortex/hippocampus, using the scDRS data-release MAGMA gene sets (top-1000 genes, z-score weights; figshare 19312583, gs_file.zip, 0.9 MB, fetched once into _data/scdrs): schizophrenia, bipolar, MDD, Alzheimer's, and height as a non-brain control. Human→mouse by symbol case mapping (approximation; 1:1 orthologue tables were not used to keep this offline).

KEY PLOTTING CALL

RECREATIONS/SCDRS/SCDRS_RECREATE.PY — EXCERPT · ENDS LINE 25

```
fig,axs=plt.subplots(2,3,figsize=(16,10)); axs=axs.ravel()
sc.pl.umap(a,color="celltype",ax=axs[0],show=False,frameon=False,legend_loc="on data",legend_fontsize=7,title="Zeisel cell types")
for ax,t in zip(axs[1:],gs):
    sc.pl.umap(a,color=f"{t}_norm_score",ax=ax,show=False,frameon=False,cmap="RdBu_r",vcenter=0,vmax=4,vmin=-4,title=f"scDRS score: {t}")
fig.tight_layout(); fig.savefig("v1_umap_scores.png",dpi=130)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/martinjzhang/scDRS.

LIMITATIONS & NOTES

- Human → mouse genes mapped by symbol case: an approximation, since 1:1 orthologue tables were not used, to keep this offline (script header).
- NumPy 2 shim for np.float_ (METHODS §9).

Reproduce: `python3 recreations/scDRS/scdrs_recreate.py`

√ verified run · exit 0 · 76 s

scDRS · 3 variations

RECREATIONS/SCDRS/



v1 umap_scores

EMBEDDINGS



v2 group_assoc_heatmap

HEATMAPS



v3 score_boxplots

COMPOSITION

MNE-Python

#19

OF 73 · NICHE & COOL SCORE

PYTHON | NEUROPHYSIOLOGY | BRAIN ATLAS, 3D & NEUROIMAGING

2.27×	5/5	0.798	1	3,549
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · MNE-PYTHON

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0 → 0.013 per 1k papers (2023 → 2025–26).
GitHub mne-tools/mne-python · created 2011-01-28 · last push 2026-10-13

BEST FOR

cortical surface source maps, topomaps

THIS RECREATION

MNE-Python on the PhysioNet EEG Motor Movement/Imagery dataset (EEGBCI; subject 1, imagery runs 6/10/14, ~8 MB): raw-trace browser figure, event-related (de)synchronisation time-frequency maps at C3/Cz/C4, and mu/beta-band topomaps for imagined hands vs feet.

KEY PLOTTING CALL

RECREATIONS/MNE/MNE_RECREATE.PY — EXCERPT · ENDS LINE 13

```
fig=raw.plot(duration=10,n_channels=24,scalings=dict(eeg=60e-6),show=False,title="EEGBCI S001 imagery (hands/feet)",show_scrollbars=False)
fig.set_size_inches(12,7); fig.savefig("v1_raw_browser.png",dpi=120); plt.close(fig)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/mne-tools/mne-python.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/MNE/mne_recreate.py`

√ verified run · exit 0 · 4 s

LEAD FIGURE



v2 erds_tfr

BRAIN

MNE-Python · 3 variations · v2 on facing page



v1 raw_browser

BRAIN



v3 band_topomaps

BRAIN

sccomp

#20

R | COMPOSITION | COMPOSITION, DISTRIBUTIONS & LAYOUTS

OF 73 · NICHE & COOL SCORE

4.27×	4/5	0.79	10	134
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · SCCOMP

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.07 → 0.33 per 1k papers (2023 → 2025–26).
GitHub MangiolaLaboratory/sccomp · created 2021-02-08 · last push 2026-10-06

BEST FOR

posterior-predictive composition boxplots

THIS RECREATION

sccomp Bayesian compositional analysis (CmdStan backend) on Zeisel 2015: cortex vs CA1, C1 chip as sample. Pyramidal subtypes are tissue-exclusive so they are excluded. Posterior boxplots, 1D credible intervals, and the mean-variability 2D plot. Requires cmdstanr + CmdStan (installed with cmdstanr::install_cmdstan()).

KEY PLOTTING CALL

RECREATIONS/SCCOMP/SCCOMP_RECREATE.R — EXCERPT · ENDS LINE 13

```
est <- cnt %>% sccomp_estimate(formula_composition = ~ tissue, sample = "chip", cell_group = "celltype", abundance = "count", c
ores = 2, verbose = FALSE) %>%
  sccomp_remove_outliers(cores = 2, verbose = FALSE) %>% sccomp_test()
ggsave("v1_posterior_boxplot.png", sccomp_boxplot(est, factor = "tissue") + ggtitle("sccomp: proportions with posterior
predictive (cortex vs CA1)"), width = 11, height = 5, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/MangiolaLaboratory/sccomp.

LIMITATIONS & NOTES

- Requires cmdstanr + CmdStan. Pyramidal subtypes are tissue-exclusive, so they are excluded; the C1 chip is used as the sample (script header).

Reproduce: `Rscript recreations/sccomp/sccomp_recreate.R`

✓ verified run · exit 0 · 57 s

LEAD FIGURE



v1 posterior_boxplot

COMPOSITION

sccomp · 3 variations · v1 on facing page



v2 1D_intervals

COMPOSITION



v3 2D_mean_variability

COMPOSITION

PyComplexHeatmap

#26

PYTHON | HEATMAP | HEATMAPS & DOT MATRICES

OF 73 · NICHE & COOL SCORE

3.06×	4/5	0.75	2	407
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · PYCOMPLEXHEATMAP

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.035 → 0.127 per 1k papers (2023 → 2025–26).
GitHub DingWB/PyComplexHeatmap · created 2022-05-05 · last push 2026-09-01

BEST FOR

ComplexHeatmap-style in Python, dot heatmaps

KEY PLOTTING CALL

RECREATIONS/PYCOMPLEXHEATMAP/PYCH_RECREATE.PY — EXCERPT · ENDS LINE 11

```
X=pd.DataFrame(ad[idx,mk].X.toarray().T,index=mk,columns=ad.obs_names[idx]); meta=ad.obs.iloc[idx][["celltype","tissue"]]
col_ha=HeatmapAnnotation(celltype=anno_simple(meta.celltype,cmap="tab10",legend=True,add_text=False),tissue=anno_simple(meta.ti
ssue,cmap="Set2"),nGene=anno_barplot(ad.obs.iloc[idx].nFeature_RNA,cmap="Greys",height=10),verbose=0,axis=1)
plt.figure(figsize=(9,5)); ClusterMapPlotter(data=X,top_annotation=col_ha,col_split=meta.celltype,col_split_gap=.8,row_cluster=T
rue,col_cluster=False,cmap="Reds",show_rownames=True,label="log-norm",row_dendrogram=True,verbose=0,rasterized=True)
plt.savefig("v1_split_annotated.png",dpi=150,bbox_inches="tight"); plt.close("all")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/DingWB/PyComplexHeatmap.

LIMITATIONS & NOTES

- Shim: matplotlib.cm.get_cmap restored (removed in matplotlib 3.9+, still called by PyComplexHeatmap) (METHODS §9).

Reproduce: `python3 recreations/PyComplexHeatmap/pych_recreate.py`

√ verified run · exit 0 · 7 s

LEAD FIGURE



v1 split_annotated

HEATMAPS

PyComplexHeatmap · 2 variations · v1 on facing page



v2 dot_clustermap

HEATMAPS

nilearn

#29

OF 73 · NICHE & COOL SCORE

PYTHON | NEUROIMAGING | BRAIN ATLAS, 3D & NEUROIMAGING

1.92×	5/5	0.733	7	1,444
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · NILEARN

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.035 → 0.076 per 1k papers (2023 → 2025–26).
GitHub nilearn/nilearn · created 2011-01-09 · last push 2026-10-14

BEST FOR

glass brains, surface maps

THIS RECREATION

nilearn brain visualisations on real public maps (NeuroVault motor-task t-map; Harvard-Oxford atlas; fsaverage surface)

KEY PLOTTING CALL

RECREATIONS/NILEARN/NILEARN_RECREATE.PY — EXCERPT · ENDS LINE 6

```
plotting.plot_glass_brain(mot,threshold=3,colorbar=True,plot_abs=False,display_mode="lyrz",title="Glass brain: motor task t-map").savefig("v1_glass_brain.png",dpi=150)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/nilearn/nilearn.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/nilearn/nilearn_recreate.py`

√ verified run · exit 0 · 5 s

LEAD FIGURE



v1 glass_brain

BRAIN

nilearn · 4 variations · v1 on facing page



v2 stat_map_cuts

BRAIN



v3 surface_stat_map

BRAIN



v4 atlas_roi

BRAIN

SCpubr

#31

OF 73 · NICHE & COOL SCORE

R · SINGLE-CELL PUBLICATION PLOTS · EMBEDDINGS & FEATURE MAPS · HEATMAPS & DOT MATRICES · CELL-CELL COMMUNICATION · COMPOSITION, DISTRIBUTIONS & LAYOUTS

2.19×	4/5	0.671	14	245
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES
SORTED, LOG SCALE

NICHE & COOL · SCPUBR

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.418 → 0.927 per 1k papers (2023 → 2025–26).
GitHub enblacar/SCpubr · created 2021-12-21 · last push 2026-03-02

BEST FOR

polished Seurat plots: nebulosa, geyser, alluvial, chord, enrichment

KEY PLOTTING CALL

RECREATIONS/SCPUBR/SCPUBR_RECREATE.R — EXCERPT · ENDS LINE 5

```
sv("v1_do_DimPlot.png", do_DimPlot(so, label = TRUE, repel = TRUE, legend.position = "none", plot.title = "SCpubr::do_DimPlot"))
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/enblacar/SCpubr.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/SCpubr/SCpubr_recreate.R`

√ verified run · exit 0 · 14 s

LEAD FIGURE



v2 do_NebulosaPlot

EMBEDDINGS

SCpubr · 5 variations · v2 on facing page

RECREATIONS/SCPUBR/



v1 do_DimPlot · EMBEDDINGS



v3 do_DotPlot · HEATMAPS



v4 do_ChordDiagram · CELL-CELL



v5 do_BoxPlot · COMPOSITION

UCell

#34

OF 73 · NICHE & COOL SCORE

R · GENE-SET SCORES · EMBEDDINGS & FEATURE MAPS · COMPOSITION, DISTRIBUTIONS & LAYOUTS · HEATMAPS & DOT MATRICES

3.15×	3/5	0.663	144	213
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE



NICHE & COOL · UCELL

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 3.134 → 9.881 per 1k papers (2023 → 2025–26).
GitHub carmonalab/UCell · created 2021-04-06 · last push 2026-09-08

BEST FOR

signature score UMAPs

KEY PLOTTING CALL

RECREATIONS/UCELL/UCELL_RECREATE.R — EXCERPT · ENDS LINE 7

```
so <- AddModuleScore_UCell(so, features = sigs, ncores = 2); sc <- paste0(names(sigs), "_UCell")
ggsave("v1_UCell_featureplots.png", FeaturePlot(so, sc, ncol = 3, order = TRUE) & scale_color_viridis_c(option = "A"), width = 1
2, height = 7, dpi = 110, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/carmonalab/UCell.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/UCell/UCell_recreate.R`

√ verified run · exit 0 · 24 s

LEAD FIGURE



v2 UCell_violins

COMPOSITION

UCell · 3 variations · v2 on facing page



v1 UCell_featureplots

EMBEDDINGS



v3 UCell_dotplot

HEATMAPS

spatialLIBD

#37

OF 73 · NICHE & COOL SCORE

R · SPATIAL · SPATIAL & TISSUE MAPS

1.97×	4/5	0.64	73	115
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · SPATIALLIBD

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.975 → 1.93 per 1k papers (2023 → 2025–26).
GitHub LieberInstitute/spatialLIBD · created 2019-12-04 · last push 2026-10-13

BEST FOR

layer-wise Visium plots, DLPFC

THIS RECREATION

spatialLIBD look on DLPFC 151673: layer annotation on H&E, gene on H&E, layer-wise marker boxplots

KEY PLOTTING CALL

RECREATIONS/SPATIALLIBD/SPATIALLIBD_RECREATE.R — EXCERPT · ENDS LINE 6

```
spe <- load_dlpfc(); rowData(spe)$gene_search <- paste0(rowData(spe)$gene_name, "; ", rowData(spe)$gene_id); spe$key <- paste0(s
pe$sample_id, "_", colnames(spe))
spe$layer <- factor(spe$ground_truth); cols <- setNames(c("#F0027F","#377EB8","#4DAF4A","#984EA3","#FFD700","#FF7F00","#1A1A1A")
, levels(spe$layer))
ggsave("v1_vis_clus_layers.png", vis_clus(spe, sampleid = "sample_151673", clustervar = "layer", colors = cols, spatial = TRUE,
point_size = 1.6, image_id = "lowres") + ggtitle("vis_clus: manual cortical layers"), width = 7, height = 6.5, dpi = 140, bg = "
white")
```

IN THE CORPUS · MOST-CITED OF 35 PAPERS NAMING “SPATIALLIBD” IN METHODS

CITATIONS		
537	**Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST.** *Nat Commun, 2023*	DOI ↗ PMC9977836
227	**BANKSY unifies cell typing and tissue domain segmentation for scalable spatial omics data analysis.** *Nat Genet, 2024*	DOI ↗ PMC10937399

Figure-level example mining (METHODS §3) was run for the Top 100 only, so these are mention-level examples without a verified figure link.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/spatialLIBD/spatialLIBD_recreate.R`

√ verified run · exit 0 · 29 s

spatialLIBD · 4 variations



v1 vis_clus_layers · SPATIAL



v2 vis_gene_MOBP · SPATIAL



v3 vis_grid_genes · SPATIAL



v4 layer_boxplots · SPATIAL

ggrepel volcano styles

#41

OF 73 · NICHE & COOL SCORE

R VOLCANO DE, QC & SET OVERLAPS

2.3×	3/5	0.608	74	1,262
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE



NICHE & COOL · GGREPEL VOLCANO STYLES

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 1.532 → 3.531 per 1k papers (2023 → 2025–26).
GitHub slowkow/ggrepel · created 2016-01-03 · last push 2026-10-09

BEST FOR

label-repelled volcanoes

Also catalogued as: **ggrepel** is Top-100 #64 (92 papers; recreated under ggplot2), examples below are from that entry

THIS RECREATION

Volcano-plot styles seen in recent neuro papers (ggplot2 + ggrepel), airway DESeq2 results

KEY PLOTTING CALL

RECREATIONS/GGREPEL_VOLCANO_STYLES/VOLCANO_STYLES_RECREATE.R — EXCERPT · ENDS LINE 8

```
lab <- res |> filter(sig) |> group_by(dir) |> slice_min(padj, n = 8)
g <- ggplot(res, aes(log2FoldChange, nl)) + geom_point(data = filter(res, !sig), color = "grey85", size = .6) + geom_point(data
= filter(res, sig), aes(color = log2FoldChange, size = nl), alpha = .8) + scale_color_gradient2(low = "#2166ac", mid =
"grey90", high = "#b2182b") + scale_size(range = c(.4, 3), guide = "none") +
  geom_label_repel(data = lab, aes(label = symbol, fill = dir), color = "white", size = 3, segment.color = "grey40", min.segment
.length = 0, box.padding = .5) + scale_fill_manual(values = c(up = "#b2182b", down = "#2166ac"), guide = "none") +
theme_classic() + labs(title = "Gradient + sized points + filled repel labels")
ggsave("v1_gradient_labels.png", g, width = 7, height = 6, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE · TOP-SCORED EXAMPLE FIGURES (METHODS §3)

Fig. 3 TEXT — **TDP-43 nuclear loss in FTD/ALS causes widespread alternative polyadenylation changes.** *Nat Neurosci, 2025*
"...All plots were prepared using tidyplots, ggplot2, ggpubr, patchwork and ggrepel in R."
DOI ↗ Fig. 3 on PMC ↗

Fig. 3 TEXT — **A concerted neuron-astrocyte program declines in ageing and schizophrenia.** *Nature, 2024*
"...R (v.4.1.3): cluster (v.2.1.2)138, ComplexHeatmap (v.2.10.0)139,140, data.table (v.1.14.8)141, DescTools (v.0.99.48)142, dplyr (v.1.1.2)143, gdata (v.2.19.0)144, ggforce (v.0.4.1)145, ggplot2 (v.3.4.2)146 ..."
DOI ↗ Fig. 3 on PMC ↗

Fig 1 TEXT — **Loss of STING in parkin mutant flies suppresses muscle defects and mitochondria damage.** *PLoS Genet, 2023*
"...For all plots, ggplot2 v.3.4 and ggrepel v.0.9.2 were used for annotation."
DOI ↗ Fig 1 on PMC ↗

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggrepel_volcano_styles/volcano_styles_recreate.R` ✓ verified run · exit 0 · 3 s

ggrepel volcano styles · 3 variations



v1 gradient_labels

DE



v2 side_aligned

DE



v3 quadrants_counts

DE

Augur

#42

OF 73 · NICHE & COOL SCORE

R · PERTURBATION PRIORITIZATION · COMPOSITION, DISTRIBUTIONS & LAYOUTS · EMBEDDINGS & FEATURE MAPS

1.86×	4/5	0.606	25	153
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES
SORTED, LOG SCALE

NICHE & COOL · AUGUR

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.522 → 0.978 per 1k papers (2023 → 2025–26).
GitHub neurorestore/Augur · created 2019-12-20 · last push 2026-03-26

BEST FOR

lollipop/UMAP of cell-type responsiveness

THIS RECREATION

Augur cell-type prioritisation: which Zeisel cell types best separate tissue of origin (S1 cortex vs CA1)?

KEY PLOTTING CALL

RECREATIONS/AUGUR/AUGUR_RECREATE.R — EXCERPT · ENDS LINE 8

```
au <- calculate_auc(GetAssayData(so, layer = "data"), so@meta.data, cell_type_col = "celltype", label_col = "tissue", n_subsamp
les = 20, subsample_size = 20, n_threads = 2, classifier = "rf")
ggsave("v1_lollipop.png", plot_lollipop(au) + ggtitle("Augur AUC: tissue (S1 vs CA1) separability"), width = 6, height = 4, dpi
= 150, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/neurorestore/Augur.

> LIMITATIONS & NOTES
> No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/Augur/Augur_recreate.R` — ✓ verified run · exit 0 · 79 s

LEAD FIGURE



v3 feature_importance

COMPOSITION

Augur · 3 variations · v3 on facing page



v1 lollipop

COMPOSITION



v2 umap_auc

EMBEDDINGS

ggridges

#47

OF 73 · NICHE & COOL SCORE

R · DISTRIBUTION · COMPOSITION, DISTRIBUTIONS & LAYOUTS

1.7×	4/5	0.585	20	425
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · GGRIDGES

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.557 → 0.953 per 1k papers (2023 → 2025–26). GitHub wilkelab/ggridges · created 2017-09-13 · last push 2026-06-18 · also scored: ggdensity (#69, growth 0.51×)

BEST FOR

ridgeline plots

KEY PLOTTING CALL

RECREATIONS/GGRIDGES_GGDENSITY/RIDGES_RECREATE.R — EXCERPT · ENDS LINE 5

```
so <- readRDS("../_data/zeisel_seurat.rds"); df <- FetchData(so, c("Snap25","Mbp","celltype","nFeature_RNA","umap_1","umap_2"))
g <- ggplot(df, aes(nFeature_RNA, celltype, fill = after_stat(x))) + geom_density_ridges_gradient(scale = 2.2, rel_min_height
= .01) + scale_fill_viridis_c(option = "C", name = "genes") + theme_ridges() + labs(title = "ggridges gradient: genes per cell")
ggsave("v1_ridges_gradient.png", g, width = 7, height = 5.5, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/wilkelab/ggridges.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggridges_ggdensity/ridges_recreate.R`

√ verified run · exit 0 · 9 s

LEAD FIGURE



v1 ridges_gradient

COMPOSITION

ggridges · 4 variations · v1 on facing page

RECREATIONS/GGRIDGES_GGDENSITY/



v2 ridges_points

COMPOSITION



v3 hdr_umap

COMPOSITION



v4 hdr_lines

COMPOSITION

plotnine

#48

OF 73 · NICHE & COOL SCORE

PYTHON | GENERAL | EMBEDDINGS & FEATURE MAPS | COMPOSITION, DISTRIBUTIONS & LAYOUTS

2.2×	3/5	0.581	3	4,771
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE



NICHE & COOL · PLOTNINE

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.07 → 0.165 per 1k papers (2023 → 2025–26).
GitHub has2k1/plotnine · created 2017-04-24 · last push 2026-10-08

BEST FOR

ggplot grammar in Python

KEY PLOTTING CALL

RECREATIONS/PLOTNINE/PLOTNINE_RECREATE.PY — EXCERPT · ENDS LINE 5

```
ad=load(); g=["Gad1","Satb2","Mbp","Aqp4"]; df=pd.DataFrame(ad[:,g].X.toarray(),columns=g);
df=pd.concat([df,ad.obs[["celltype","tissue","UMAP_1","UMAP_2","nFeature_RNA"]].reset_index(drop=True)],axis=1)
p=(ggplot(df,aes("UMAP_1","UMAP_2",color="celltype"))+geom_point(size=.4)+theme_void()+coord_fixed()+ggtitle("plotnine UMAP")+g
uides(color=guide_legend(override_aes={"size":3})))
p.save("v1_umap.png",dpi=150,width=7,height=5,verbose=False)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/has2k1/plotnine.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/plotnine/plotnine_recreate.py`

√ verified run · exit 0 · 7 s

LEAD FIGURE



v3 bin2d_facets

COMPOSITION

plotnine · 3 variations · v3 on facing page



v1 umap

EMBEDDINGS



v2 faceted_violins

COMPOSITION

Palantir

#50

PYTHON | TRAJECTORY | TRAJECTORY, PSEUDOTIME & VELOCITY

OF 73 · NICHE & COOL SCORE

1.62×	4/5	0.571	31	341
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · PALANTIR

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 1.114 → 1.816 per 1k papers (2023 → 2025–26).
GitHub dpeerlab/Palantir · created 2017-11-22 · last push 2026-09-23

BEST FOR

branch probabilities, gene trends

THIS RECREATION

Palantir on dentate gyrus (diffusion maps → pseudotime, entropy, branch probabilities, gene trends)

KEY PLOTTING CALL

RECREATIONS/PALANTIR/PALANTIR_RECREATE.PY — EXCERPT · ENDS LINE 13

```
palantir.core.run_palantir(a,start,terminal_states=term,num_waypoints=500)
palantir.plot.plot_palantir_results(a,s=3); plt.savefig("v1_palantir_results.png",dpi=130,bbox_inches="tight"); plt.close("all")
```

IN THE CORPUS · MOST-CITED OF 24 PAPERS NAMING "PALANTIR" IN METHODS

258 CITATIONS — **Single-cell brain organoid screening identifies developmental defects in autism.** *Nature, 2023*
DOI ↗ PMC10499611

42 CITATIONS — **Tracking single-cell evolution using clock-like chromatin accessibility loci.** *Nat Biotechnol, 2025*
DOI ↗ PMC12084158

Figure-level example mining (METHODS §3) was run for the Top 100 only, so these are mention-level examples without a verified figure link.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/Palantir/palantir_recreate.py` — ✓ verified run · exit 0 · 19 s

LEAD FIGURE



v1 palantir_results — TRAJECTORY

Palantir · 4 variations · v1 on facing page



v2 branch_selection

TRAJECTORY



v3 gene_trends

TRAJECTORY



v4 trajectory_arrows

TRAJECTORY

ggspavis/ SpatialExperiment

#51

OF 73 · NICHE & COOL SCORE

R · SPATIAL · SPATIAL & TISSUE MAPS

2.15×	3/5	0.56	9	5
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · GGSPAVIS/SPATIALEXPERIMENT

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.279 → 0.61 per 1k papers (2023 → 2025–26).
GitHub lmweber/ggspavis · created 2020-12-02 · last push 2026-07-14

BEST FOR

spot plots on tissue

KEY PLOTTING CALL

RECREATIONS/GGSPAVIS/GGSPAVIS_RECREATE.R — EXCERPT · ENDS LINE 4

```
spe <- load_dlpfc(); spe$sum <- colSums(counts(spe)); spe$MOBP <- logcounts(spe)["MOBP",]
ggsave("v1_plotSpots_layers.png", plotCoords(spe, annotate = "ground_truth", pal = "libd_layer_colors", point_size = .6) + ggti
tle("plotSpots"), width = 6, height = 5.5, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/lmweber/ggspavis.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggspavis/ggspavis_recreate.R`

✓ verified run · exit 0 · 18 s

LEAD FIGURE



v3 continuous_qc_gene

SPATIAL

ggspavis/SpatialExperime… · 3 variations · v3 on facing page



v1 plotSpots_layers

SPATIAL



v2 plotVisium_HE

SPATIAL

plotgardener

#55

OF 73 · NICHE & COOL SCORE

R | GENOME TRACKS | GENOME TRACKS & GENOMIC SIGNAL

0.93×	5/5	0.544	4	371
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · PLOTGARDENER

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.313 → 0.292 per 1k papers (2023 → 2025–26).
GitHub PhanstielLab/plotgardener · created 2019-07-23 · last push 2026-07-18

BEST FOR

page-layout multi-panel genomic figures, Hi-C

THIS RECREATION

plotgardener multi-panel genomic page (plotgardenerData example: IMR90 Hi-C + ChIP + gene track, hg19)

KEY PLOTTING CALL

RECREATIONS/PLOTGARDENER/PLOTGARDENER_RECREATE.R — EXCERPT · ENDS LINE 13

```
pageCreate(width = 7.5, height = 6.5, default.units = "inches", showGuides = FALSE)
params <- pgParams(chrom = "chr21", chromstart = 28000000, chromend = 30300000, assembly = "hg19", x = .5, width = 6.5)
hic <- plotHicTriangle(data = IMR90_HiC_10kb, params = params, zrange = c(0, 70), resolution = 10000, y = .5, height = 2.5,
just = c("left","top"), default.units = "inches")
annoHeatmapLegend(hic, x = 7, y = .5, width = .13, height = 1.2, just = c("right","top"))
plotPairsArches(data = IMR90_DNAloops_pairs, params = params, y = 3.1, height = .5, fill = "#1f4297", linecolor = "#1f4297", fl
ip = TRUE)
plotSignal(data = IMR90_ChIP_H3K27ac_signal, params = params, y = 3.7, height = .6, fill = "#37a7db", linecolor = "#37a7db"); p
lotText("H3K27ac", x = .5, y = 3.7, just = c("left","top"), fontsize = 8)
plotSignal(data = GM12878_ChIP_CTCF_signal, params = params, y = 4.4, height = .6, fill = "#e0644e", linecolor = "#e0644e"); pl
otText("CTCF", x = .5, y = 4.4, just = c("left","top"), fontsize = 8)
plotGenes(params = params, y = 5.1, height = .7); plotGenomeLabel(chrom = "chr21", chromstart = 28000000, chromend = 30300000, a
ssembly = "hg19", x = .5, y = 5.9, length = 6.5, scale = "Mb")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/PhanstielLab/plotgardener.

LIMITATIONS & NOTES

- Non-neuro fallback: plotgardenerData IMR90 example (Hi-C + ChIP + gene track, hg19) (SUMMARY; script header).

Reproduce: `Rscript recreations/plotgardener/plotgardener_recreate.R`

✓ verified run · exit 0 · 14 s

plotgardener · 2 variations



v1 page_layout

GENOME



v2 gene_and_ideogram

GENOME

propeller (speckle)

#60

OF 73 · NICHE & COOL SCORE

R | COMPOSITION | COMPOSITION, DISTRIBUTIONS & LAYOUTS

1.89×	3/5	0.512	116	71
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · PROPELLER (SPECKLE)

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 2.299 → 4.356 per 1k papers (2023 → 2025–26).
GitHub phipsonlab/speckle · created 2022-08-19 · last push 2022-12-08

BEST FOR

proportion plots

THIS RECREATION

speckle::propeller differential cell-type composition, Zeisel 2015: S1 cortex vs CA1 hippocampus, using the Fluidigm C1 chip ID (cell-name prefix) as the biological/technical replicate.

KEY PLOTTING CALL

RECREATIONS/SPECKLE_PROPELLER/PROPELLER_RECREATE.R — EXCERPT · ENDS LINE 13

```
res <- propeller(clusters = md$celltype, sample = md$chip, group = md$tissue, transform = "logit")
png("v1_plotCellTypeProps.png", 1300, 900, res = 150); print(plotCellTypeProps(clusters = md$celltype, sample = md$chip) +
theme(axis.text.x = element_text(angle = 90, size = 6)) + ggtitle("Cell-type proportions per C1 chip")); dev.off()
```

IN THE CORPUS · MOST-CITED OF 35 PAPERS NAMING “SPECKLE/PROPELLER” IN METHODS

CITATIONS		
164	**Molecular and cellular dynamics of the developing human neocortex.** *Nature, 2025*	DOI ↗ PMC12589127
47	**Morphodynamics of human early brain organoid development.** *Nature, 2025*	DOI ↗ PMC12390842

Figure-level example mining (METHODS §3) was run for the Top 100 only, so these are mention-level examples without a verified figure link.

LIMITATIONS & NOTES

- Shared cell types only, because pyramidal subtypes are tissue-exclusive; the Fluidigm C1 chip ID is used as the replicate (script header).

Reproduce: `Rscript recreations/speckle_propeller/propeller_recreate.R`

✓ verified run · exit 0 · 14 s

propeller (speckle) · 3 variations

RECREATIONS/SPECKLE_PROPELLER/



v1 plotCellTypeProps

COMPOSITION



v2 props_by_group

COMPOSITION



v3 composition_volcano

COMPOSITION

ggcoverage

#63

OF 73 · NICHE & COOL SCORE

R | GENOME TRACKS | GENOME TRACKS & GENOMIC SIGNAL

0.97×	4/5	0.462	3	273
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · GGCOVERAGE

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.104 → 0.102 per 1k papers (2023 → 2025–26).
GitHub showteeth/ggcoverage · created 2022-07-18 · last push 2025-01-24

BEST FOR

ggplot2 coverage tracks + gene/ideogram

THIS RECREATION

ggcoverage genome-coverage tracks. NOTE: uses the package's bundled human RNA-seq bigWigs (chr14 HNRNPC locus; ERR127302/3/6/7, KO vs WT) because they are the only coverage files shipped offline - not neuro data.

KEY PLOTTING CALL

RECREATIONS/GGCOVERAGE/GGCOVERAGE_RECREATE.R — EXCERPT · ENDS LINE 10

```
d <- system.file("extdata", package = "ggcoverage"); gtf <- import.gff(file.path(d, "used_hg19.gtf"), format = "gtf")
meta <- read.csv(file.path(d, "RNA-seq", "meta_info.csv"))
trk <- LoadTrackFile(track.folder = file.path(d, "RNA-seq"), format = "bw", region = "chr14:21,677,306-21,737,601", extend = 200
0, meta.info = meta)
cols <- c(KO = "#d7301f", WT = "#225ea8")
base <- ggcoverage(data = trk, plot.type = "facet", group.key = "Type", facet.key = "Type", range.position = "out", mark.region
= data.frame(start = c(21678900, 21732001), end = c(21679900, 21732400), label = c("M1", "M2")), show.mark.label = TRUE)
ggsave("v1_coverage_gene.png", base + geom_gene(gtf.gr = gtf) + ggtitle("ggcoverage: RNA-seq coverage + gene models (HNRNPC
locus)"), width = 10, height = 7, dpi = 130, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/showteeth/ggcoverage.

LIMITATIONS & NOTES

- Non-neuro fallback: the package's bundled human RNA-seq bigWigs (chr14 HNRNPC locus; ERR127302/3/6/7, KO vs WT), the only coverage files shipped offline (script header).

Reproduce: `Rscript recreations/ggcoverage/ggcoverage_recreate.R`

✓ verified run · exit 0 · 20 s

ggcoverage · 2 variations



v1 coverage_gene

GENOME



v2 samples_transcripts

GENOME

ggtree/ ggtreeExtra

#64

OF 73 · NICHE & COOL SCORE

R · TREE · NETWORKS, GRAPHS & TREES

0.94×	4/5	0.455	26	99
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · GGTREE/GGTREEEXTRA

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 3.134 → 2.959 per 1k papers (2023 → 2025–26).
GitHub YuLab-SMU/ggtreeExtra · created 2020-05-28 · last push 2026-06-04

BEST FOR

circular trees with heatmap/bar rings

THIS RECREATION

Circular dendrogram of Zeisel level2 subclasses (mean-expression distance) with ggtreeExtra rings

KEY PLOTTING CALL

RECREATIONS/GGTREE_GGTREEEXTRA/GGTREE_RECREATE.R — EXCERPT · ENDS LINE 9

```
so <- readRDS("../_data/zeisel_seurat.rds"); av <- AverageExpression(so, features = VariableFeatures(so), group.by = "level2cla
ss")$RNA |> as.matrix()
colnames(av) <- gsub("-", "_", colnames(av)); tr <- as.phylo(hclust(as.dist(1 - cor(log1p(av))), "average"))
md <- so@meta.data; md$sub <- gsub("-", "_", md$level2class)
info <- unique(md[, c("sub","celltype")]); info <- info[!duplicated(info$sub),]; n <- as.data.frame(table(md$sub)); colnames(n)
<- c("label","cells"); info <- merge(info, n, by.x = "sub", by.y = "label"); colnames(info)[1] <- "label"
p <- ggtree(tr, layout = "circular") %<+% info + geom_tippoint(aes(color = celltype), size = 2) + geom_tiplab(size = 2, offset
= .02)
ggsave("v1_circular_tree.png", p + theme(legend.position = "right"), width = 8, height = 7, dpi = 140, bg = "white")
```

IN THE CORPUS · MOST-CITED OF 25 PAPERS NAMING “GGTREE” IN METHODS

86 CITATIONS — **A comprehensive spatio-cellular map of the human hypothalamus.** *Nature, 2025*
DOI ↗ PMC11922758

76 CITATIONS — **A marmoset brain cell census reveals regional specialization of cellular identities.** *Sci Adv, 2023*
DOI ↗ PMC10569717

Figure-level example mining (METHODS §3) was run for the Top 100 only, so these are mention-level examples without a verified figure link.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggtree_ggtreeExtra/ggtree_recreate.R`

√ verified run · exit 0 · 9 s

ggtree/ggtreeExtra · 3 variations



v1 circular_tree

NETWORKS



v2 fan_heatmap_bar_rings

NETWORKS



v3 rect_tree_clade_highlights

NETWORKS

Mellon

#65

PYTHON | CELL-STATE DENSITY | TRAJECTORY, PSEUDOTIME & VELOCITY

OF 73 · NICHE & COOL SCORE

0.93×	4/5	0.444	27	77
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · MELLON

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.836 → 0.775 per 1k papers (2023 → 2025–26).
GitHub settylab/Mellon · created 2022-10-28 · last push 2026-09-24

BEST FOR

cell-state density landscapes

THIS RECREATION

Mellon cell-state density (via palantir.utils.run_density) on dentate gyrus

KEY PLOTTING CALL

RECREATIONS/MELLON/MELLON_RECREATE.PY — EXCERPT · ENDS LINE 8

```
a=load(); palantir.utils.run_diffusion_maps(a,n_components=10); palantir.utils.run_density(a)
fig,ax=plt.subplots(1,2,figsize=(11,4.5))
s=ax[0].scatter(*a.obsm["X_umap"].T,c=a.obs["mellon_log_density"],s=6,cmap="magma"); ax[0].set_title("Mellon log-density");
plt.colorbar(s,ax=ax[0]); ax[0].axis("off")
s=ax[1].scatter(*a.obsm["X_umap"].T,c=a.obs["mellon_log_density_clipped"],s=6,cmap="Spectral_r"); ax[1].set_title("clipped
density (low = transitional)"); plt.colorbar(s,ax=ax[1]); ax[1].axis("off")
fig.tight_layout(); fig.savefig("v1_density_umap.png",dpi=140)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/settylab/Mellon.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/Mellon/mellon_recreate.py`

√ verified run · exit 0 · 28 s

LEAD FIGURE



v1 density_umap

TRAJECTORY

Mellon · 3 variations · v1 on facing page



v2 density_by_cluster

TRAJECTORY



v3 density_landscape_3d

TRAJECTORY

karyoploteR

#66

OF 73 · NICHE & COOL SCORE

R · GENOME · GENOME TRACKS & GENOMIC SIGNAL

0.67×	4/5	0.428	13	377
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · KARYOPLOTER

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 1.289 → 0.864 per 1k papers (2023 → 2025–26).
GitHub bernatgel/karyoploteR · created 2016-06-07 · last push 2026-07-06

BEST FOR

whole-karyotype data plots

THIS RECREATION

karyoploteR: genome-wide views of airway DE genes (hg19 coordinates via TxDb)

KEY PLOTTING CALL

RECREATIONS/KARYOPLOTER/KARYOPLOTER_RECREATE.R — EXCERPT · ENDS LINE 8

```
res <- as.data.frame(readRDS("../_data/airway_deseq2.rds")$res); res <- res[!is.na(res$padj) & !is.na(res$entrez),]
g <- genes(TxDb.Hsapiens.UCSC.hg19.knownGene); g <- g[g$gene_id %in% res$entrez]; r <- res[match(g$gene_id, res$entrez),];
g$lfc <- r$log2FoldChange; g$padj <- r$padj; g$sym <- r$symbol
g <- keepStandardChromosomes(g, pruning.mode = "coarse"); g <- g[seqnames(g) %in% paste0("chr", c(1:22, "X"))]
sig <- g[g$padj < .01]
png("v1_karyotype_density.png", 1500, 1100, res = 150); kp <- plotKaryotype("hg19", plot.type = 1, chromosomes = paste0("chr",
c(1:22,"X"))); kpPlotDensity(kp, sig, col = "#c51b7d", border = NA, window.size = 2e6); kpAddMainTitle(kp, "DE-gene density
(padj<0.01), 2-Mb windows"); dev.off()
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/bernatgel/karyoploteR.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/karyoploteR/karyoploteR_recreate.R`

√ verified run · exit 0 · 16 s

LEAD FIGURE



v2 lfc_scatter_ideogram

GENOME

karyoploteR · 3 variations · v2 on facing page

v1 karyotype_density

GENOME



v3 markers_chr

GENOME

marsilea

#67

OF 73 · NICHE & COOL SCORE

PYTHON | HEATMAP | HEATMAPS & DOT MATRICES

0.64×	4/5	0.421	0	324
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE



NICHE & COOL · MARSILEA

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.104 → 0.064 per 1k papers (2023 → 2025–26). GitHub Marsilea-viz/marsilea · created 2022-08-22 · last push 2026-09-26

BEST FOR

composable cross-layout heatmaps

THIS RECREATION

marsilea composable heatmaps on Zeisel cortex/hippocampus

KEY PLOTTING CALL

RECREATIONS/MARSILEA/MARSILEA_RECREATE.PY — EXCERPT · ENDS LINE 9

```
ad=load();
mk=["Snap25","Gad1","Gad2","Pvalb","Sst","Satb2","Neurod6","Wfs1","Aqp4","Gfap","Mbp","Plp1","Cx3cr1","C1qa","Cldn5","Acta2"]
df=pd.DataFrame(ad[:,mk].X.toarray(),columns=mk); df["ct"]=ad.obs.celltype.values
mean=df.groupby("ct",observed=True).mean(); frac=(df[mk]>0).groupby(df.ct,observed=True).mean(); z=(mean-mean.mean())/mean.std()
n=df.ct.value_counts().reindex(mean.index)
h=ma.Heatmap(z.values,cmap="RdBu_r",label="z-score",width=5,height=3.2); h.add_left(mp.Labels(mean.index),pad=.05);
h.add_bottom(mp.Labels(mk,rotation=90)); h.add_right(mp.Numbers(n.values,color="#4C72B0",label="cells"),size=1,pad=.1)
h.add_dendrogram("top",colors="#555"); h.add_dendrogram("left"); h.add_legends(); h.add_title("marsilea: z-scored means + cell
counts"); h.save("v1_heatmap_dendro_bars.png",dpi=150)
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/Marsilea-viz/marsilea.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `python3 recreations/marsilea/marsilea_recreate.py`

√ verified run · exit 0 · 7 s

LEAD FIGURE



v1 heatmap_dendro_bars

HEATMAPS

marsilea · 3 variations · v1 on facing page



v2 sized_dot_heatmap

HEATMAPS



v3 stackbar

HEATMAPS

dittoSeq

#68

OF 73 · NICHE & COOL SCORE

R · SINGLE-CELL PUBLICATION PLOTS · EMBEDDINGS & FEATURE MAPS · COMPOSITION, DISTRIBUTIONS & LAYOUTS · HEATMAPS & DOT MATRICES

1.25×	3/5	0.413	27	216
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · DITTOSEQ

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 1.045 → 1.308 per 1k papers (2023 → 2025–26).
GitHub dtm2451/dittoSeq · created 2019-02-18 · last push 2025-12-10

BEST FOR

colour-blind-safe dimplots, freq bars, hex plots

KEY PLOTTING CALL

RECREATIONS/DITTOSEQ/DITTOSEQ_RECREATE.R — EXCERPT · ENDS LINE 4

```
so <- readRDS("../_data/zeisel_seurat.rds"); mk <-
c("Snap25","Gad1","Gad2","Pvalb","Sst","Satb2","Neurod6","Wfs1","Aqp4","Gfap","Mbp","Plp1","Cx3cr1","C1qa","Cldn5","Acta2")
ggsave("v1_dittoDimPlot_letters.png", dittoDimPlot(so, "celltype", do.label = TRUE, labels.highlight = TRUE, legend.show =
FALSE, do.letter = FALSE, size = .8), width = 6.5, height = 6, dpi = 140, bg = "white")
```

IN THE CORPUS · MOST-CITED OF 24 PAPERS NAMING "DITTOSEQ" IN METHODS

120 CITATIONS — **Defining blood-induced microglia functions in neurodegeneration through multiomic profiling.** *Nat Immunol, 2023*
DOI ↗ PMC10307624

23 CITATIONS — **Exercise rejuvenates microglia and reverses T cell accumulation in the aged female mouse brain.** *Aging Cell, 2024*
DOI ↗ PMC11258432

Figure-level example mining (METHODS §3) was run for the Top 100 only, so these are mention-level examples without a verified figure link.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/dittoSeq/dittoSeq_recreate.R` — √ verified run · exit 0 · 21 s

LEAD FIGURE



v5 dittoHeatmap — HEATMAPS

dittoSeq · 5 variations · v5 on facing page



v1 dittoDimPlot_letters

EMBEDDINGS



v2 dittoBarPlot

COMPOSITION



v3 dittoPlot_ridgejitter

COMPOSITION



v4 dittoDimHex

EMBEDDINGS

ggforce

#70

OF 73 · NICHE & COOL SCORE

R | GENERAL | EMBEDDINGS & FEATURE MAPS | COMPOSITION, DISTRIBUTIONS & LAYOUTS

1.14×	3/5	0.386	15	958
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · GGFORCE

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.488 → 0.559 per 1k papers (2023 → 2025–26).
GitHub thomasp85/ggforce · created 2016-01-02 · last push 2025-09-15

BEST FOR

hull marks, zoom facets, parallel sets

KEY PLOTTING CALL

RECREATIONS/GGFORCE/GGFORCE_RECREATE.R — EXCERPT · ENDS LINE 5

```
md <- readRDS("../_data/zeisel_seurat.rds")@meta.data; md$UMAP_1 <- read.csv("../_data/zeisel_obs.csv")$UMAP_1; md$UMAP_2 <- re
ad.csv("../_data/zeisel_obs.csv")$UMAP_2
g <- ggplot(md, aes(UMAP_1, UMAP_2)) + geom_point(aes(color = celltype), size = .4) + geom_mark_hull(aes(fill = celltype, label
= celltype), concavity = 3, expand = unit(2,"mm"), alpha = .12, label.fontsize = 8, con.cap = 0) + theme_void() + theme(legend.
position = "none") + labs(title = "geom_mark_hull: annotated cluster hulls")
ggsave("v1_mark_hull.png", g, width = 8, height = 7, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/thomasp85/ggforce.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/ggforce/ggforce_recreate.R`

√ verified run · exit 0 · 7 s

LEAD FIGURE



v2 facet_zoom

COMPOSITION

ggforce · 3 variations · v2 on facing page



v1 mark_hull

EMBEDDINGS



v3 parallel_sets

COMPOSITION

gggenes

#71

R GENOME GENOME TRACKS & GENOMIC SIGNAL

OF 73 · NICHE & COOL SCORE

1.01×	3/5	0.368	3	603
growth ratio (per-1k rate 2025–26 ÷ 2023)	visual score (curator)	niche-cool score	neuro-corpus METHODS mentions 2023–26	GitHub stars

GROWTH AMONG 73 CANDIDATES SORTED, LOG SCALE

NICHE & COOL · GGGENES

Neuro-corpus mentions per year (*2026 partial). Broad-corpus rate 0.453 → 0.457 per 1k papers (2023 → 2025–26).
GitHub wilkox/gggenes · created 2017-03-02 · last push 2026-07-05

BEST FOR

gene-arrow maps

THIS RECREATION

gggenes: gene-arrow maps of the mouse protocadherin-gamma and Hox-adjacent neuronal loci (coordinates from TxDb mm10)

KEY PLOTTING CALL

RECREATIONS/GGGENES/GGGENES_RECREATE.R — EXCERPT · ENDS LINE 9

```
d <- do.call(rbind, lapply(names(reg), function(n) { g <- subsetByOverlaps(gr, reg[[n]]); data.frame(molecule = n, gene = g$sym
bol, start = start(g), end = end(g), strand = as.character(strand(g)), forward = as.character(strand(g)) == "+") }))
d <- d[!is.na(d$gene),]
g <- ggplot(d, aes(xmin = start, xmax = end, y = molecule, fill = forward, label = gene, forward = forward)) + geom_gene_arrow(a
rrowhead_height = unit(3,"mm"), arrowhead_width = unit(1,"mm")) + geom_gene_label(align = "left", min.size = 3) + facet_wrap(~ m
olecule, scales = "free", ncol = 1) + scale_fill_brewer(palette = "Set2") + theme_genes() + labs(title =
"gggenes: neuronal loci (mm10)")
ggsave("v1_gene_arrows.png", g, width = 11, height = 5, dpi = 140, bg = "white")
```

SEEN IN THE LITERATURE

No example paper in the source catalog. Figure-level mining (METHODS §3) covered the Top 100 only, and this tool is not in the corpus package list. Reference implementation: https://github.com/wilkox/gggenes.

LIMITATIONS & NOTES

No package-specific caveat is recorded in the source files. General corpus caveats apply (p. 7).

Reproduce: `Rscript recreations/gggenes/gggenes_recreate.R`

√ verified run · exit 0 · 13 s

LEAD FIGURE



v1 gene_arrows

GENOME

gggenes · 2 variations · v1 on facing page



v2 pcdhg_cluster

GENOME

APPENDIX A

Queued tools and why

Tools that rank in the catalog but are not yet recreated: **30 mainstream entries** and **15 Niche & Cool tools**. Reasons are quoted from the catalog *note* column and SUMMARY.md.

Niche & Cool (SUMMARY.md)

NICHE #	TOOL	REASON
9	**SCENIC+**	needs cisTarget ranking DBs (>4 GB) + pycisTopic stack
21	**CellOracle**	pip pins old numpy/pandas/anndata; would break shared env
22	**napari-spatialdata**	GUI viewer (Qt); screenshot path not set up
24	**cell2location**	incompatible with installed pyro (deep_getattr removed); needs pinned env
25	**MapMyCells (ABC atlas)**	cell_type_mapper + precomputed ABC stats download (large)
27	**Dynamo**	dynamo-release requires anndata<0.11 + old statsmodels API; tried and removed (broke scanpy/cellrank)
33	**Vitessce**	web/JS viewer, no static plotting API; needs served data + browser
36	**CELLxGENE**	Census query/explorer GUI; streaming data, not a plotting library
38	**scCODA**	needs TensorFlow/TFP (or pertpy, heavy)
39	**stLearn**	pip install failed (old pinned deps)
49	**trackplot**	needs the bwtool binary (not in apt) + neuro bigWigs
52	**SpaGCN**	pip install failed
53	**UniTVelo**	needs TensorFlow
57	**Giotto**	heavy R+python env install, not attempted after reboot
59	**ArchR**	compile failed; also needs BSgenome (~1 GB) + Arrow files

Mainstream Top 100

#	PACKAGE	LANG	CATEGORY	PAPERS	NOTE
8	**Cytoscape**	Desktop	network	1,501	Desktop GUI - queued: style-equivalent via RCy3/py4cytoscape needs a running Cytoscape; igraph/networkx redraws planned
15	**Metascape**	Web	enrichment	707	Web-only (no API) - queued: style-equivalent with clusterProfiler/ComplexHeatmap
16	**Ingenuity Pathway Analysis**	Desktop	pathways	678	Commercial (QIAGEN) - not reproducible openly; queued style-equivalent z-score bar
25	**SCENIC**	R/Python	regulon	385	Queued: needs cisTarget motif databases (~GBs)
26	**QuPath**	Desktop	imaging	380	Desktop GUI image analysis - queued: napari/matplotlib overlays
30	**monocle (v2)**	R	trajectory	310	Queued: legacy Bioconductor monocle (DDRTree)
34	**ShinyGO**	Web	enrichment	263	queued (no reason recorded in catalog)
35	**ChIPseeker**	R	peak annotation	247	queued (no reason recorded in catalog)
49	**SynGO**	Web	synaptic enrichment	139	queued (no reason recorded in catalog)
51	**ClueGO**	Desktop plugin	enrichment network	134	queued (no reason recorded in catalog)
53	**base R graphics**	R	general	132	queued (no reason recorded in catalog)
55	**maftools**	R	mutation oncoplot	123	queued (no reason recorded in catalog)
58	**ArchR**	R	scATAC	107	queued (no reason recorded in catalog)
62	**factoextra**	R	PCA/clustering	96	queued (no reason recorded in catalog)
65	**cellxgene**	Web/ Python	interactive	92	queued (no reason recorded in catalog)
67	**UCSC Cell Browser**	Web	interactive	86	queued (no reason recorded in catalog)
69	**scater**	R	single-cell QC/ embedding	83	queued (no reason recorded in catalog)

#	PACKAGE	LANG	CATEGORY	PAPERS	NOTE
72	**variancePartition**	R	variance	77	queued (no reason recorded in catalog)
73	**cell2location**	Python	spatial deconvolution	77	queued (no reason recorded in catalog)
76	**RColorBrewer**	R	color palettes	71	queued (no reason recorded in catalog)
77	**FactoMineR**	R	PCA	70	queued (no reason recorded in catalog)
78	**shiny**	R	interactive	69	queued (no reason recorded in catalog)
80	**EnrichmentMap**	Desktop plugin	enrichment network	65	queued (no reason recorded in catalog)
81	**Seurat spatial (SpatialDimPlot)**	R	spatial	63	queued (no reason recorded in catalog)
85	**UMAP (umap-learn/ uwot)**	Python/R	embedding	54	queued (no reason recorded in catalog)
89	**viridis**	R/Python	color palettes	47	queued (no reason recorded in catalog)
90	**networkx**	Python	network	47	queued (no reason recorded in catalog)
93	**pertpy**	Python	perturbation/ composition	45	queued (no reason recorded in catalog)
95	**Morpheus**	Web	heatmap	44	queued (no reason recorded in catalog)
96	**CellCall/connectome**	R	cell-cell communication	41	queued (no reason recorded in catalog)

APPENDIX B

Full Top 100 ranked table

Ranked by METHODS mentions in 17,768 papers. Page numbers link to spreads.

#	PACKAGE	LANG	CATEGORY	PAPERS	%	FULL TEXT	ROLE	STATUS	PAGE
1	**DESeq2 (plotPCA/plotMA)**	R	bulk QC/DE	5,668	31.9	5,871	analysis+plotting	done	18
2	**Seurat**	R	single-cell (embedding/dot/violin/feature)	4,671	26.29	4,846	analysis+plotting	done	20
3	**clusterProfiler**	R	enrichment	3,518	19.8	3,609	analysis+plotting	done	22
4	**ggplot2**	R	general	2,490	14.01	2,546	dedicated plotting	done	24
5	**GSEA (Broad)**	Desktop	enrichment	2,056	11.57	3,310	analysis+plotting	done	26
6	**limma**	R	bulk QC/DE	2,047	11.52	2,116	analysis+plotting	done	28
7	**edgeR**	R	bulk QC/DE	1,892	10.65	1,949	analysis+plotting	done	30
8	**Cytoscape**	Desktop	network	1,501	8.45	1,665	dedicated plotting	queued	
9	**pheatmap**	R	heatmap	1,119	6.3	1,138	dedicated plotting	done	32
10	**WGCNA**	R	network/modules	884	4.98	1,025	analysis+plotting	done	34
11	**CellChat**	R	cell-cell communication	839	4.72	976	analysis+plotting	done	36
12	**STRING**	Web	PPI network	838	4.72	1,142	analysis+plotting	done	38
13	**Enrichr**	Web	enrichment	778	4.38	912	analysis+plotting	done	40
14	**scanpy**	Python	single-cell (embedding/dot/violin/matrix)	763	4.29	789	analysis+plotting	done	42
15	**Metascape**	Web	enrichment	707	3.98	808	analysis+plotting	queued	
16	**Ingenuity Pathway Analysis**	Desktop	pathways	678	3.82	893	analysis+plotting	queued	
17	**ComplexHeatmap**	R	heatmap	567	3.19	578	dedicated plotting	done	44
18	**g:Profiler**	Web/R/Python	enrichment	563	3.17	627	analysis+plotting	done	46
19	**IGV**	Desktop/Web	genome tracks	549	3.09	794	viewer	done	48
20	**fgsea**	R	enrichment	478	2.69	498	analysis+plotting	done	50
21	**monocle3**	R	trajectory	430	2.42	499	analysis+plotting	done	52
22	**deepTools**	Python/CLI	heatmap (genomic)	416	2.34	427	dedicated plotting	done	54
23	**SingleR**	R	annotation (heatmap)	416	2.34	451	analysis+plotting	done	56
24	**AUCell**	R	gene set scores	387	2.18	432	analysis+plotting	done	58
25	**SCENIC**	R/Python	regulon	385	2.17	462	analysis+plotting	queued	
26	**QuPath**	Desktop	imaging	380	2.14	395	viewer	queued	
27	**EnhancedVolcano**	R	volcano	369	2.08	373	dedicated plotting	done	60
28	**UCSC Genome Browser**	Web	genome tracks	363	2.04	509	viewer	done	48
29	**survminer**	R	survival	351	1.98	342	dedicated plotting	done	62
30	**monocle (v2)**	R	trajectory	310	1.74	360	analysis+plotting	queued	
31	**Space Ranger/Loupe**	Desktop	spatial	300	1.69	307	viewer	done	64
32	**enrichplot**	R	enrichment	280	1.58	286	dedicated plotting	done	22
33	**ggpubr**	R	general/statistics	277	1.56	283	dedicated plotting	done	66
34	**ShinyGO**	Web	enrichment	263	1.48	284	analysis+plotting	queued	

#	PACKAGE	LANG	CATEGORY	PAPERS	%	FULL TEXT	ROLE	STATUS	PAGE
35	ChIPseeker	R	peak annotation	247	1.39	264	analysis+plotting	queued	
36	igraph	R/Python	network	241	1.36	240	analysis+plotting	done	68
37	Signac	R	scATAC (coverage/embedding)	227	1.28	239	analysis+plotting	done	70
38	matplotlib	Python	general	227	1.28	238	dedicated plotting	done	72
39	CopyKAT/inferCNV	R	CNV heatmap	207	1.17	236	analysis+plotting	done	74
40	seaborn	Python	general/statistics	192	1.08	193	dedicated plotting	done	76
41	CellPhoneDB	Python	cell-cell communication	182	1.02	231	analysis+plotting	done	78
42	slingshot	R	trajectory	173	0.97	225	analysis+plotting	done	80
43	Loupe Browser	Desktop	interactive	166	0.93	176	viewer	done	64
44	rrvgo	R	enrichment (GO reduce)	163	0.92	179	dedicated plotting	done	82
45	scVelo	Python	RNA velocity	161	0.91	177	analysis+plotting	done	84
46	VennDiagram	R	set overlap	157	0.88	160	dedicated plotting	done	86
47	gplots (heatmap.2)	R	heatmap	152	0.86	157	dedicated plotting	done	88
48	corrplot	R	correlation	147	0.83	151	dedicated plotting	done	90
49	SynGO	Web	synaptic enrichment	139	0.78	210	analysis+plotting	queued	
50	hdWGCNA	R	network/modules (sc)	134	0.75	147	analysis+plotting	done	92
51	ClueGO	Desktop plugin	enrichment network	134	0.75	152	dedicated plotting	queued	
52	clustree	R	cluster tree	133	0.75	144	dedicated plotting	done	94
53	base R graphics	R	general	132	0.74	134	dedicated plotting	queued	
54	circlize	R	circos/chord	124	0.7	130	dedicated plotting	done	96
55	maftools	R	mutation oncoplot	123	0.69	123	dedicated plotting	queued	
56	velocyto	Python/R	RNA velocity	120	0.68	129	analysis+plotting	done	84
57	UpSetR	R	set overlap	112	0.63	130	dedicated plotting	done	98
58	ArchR	R	scATAC	107	0.6	117	analysis+plotting	queued	
59	RCTD/spacexr	R	spatial deconvolution	105	0.59	130	analysis+plotting	done	100
60	CytoTRACE	R/Python	differentiation potency	103	0.58	131	analysis+plotting	done	102
61	decoupleR/decoupler	R/Python	TF/pathway activity	102	0.57	112	analysis+plotting	done	104
62	factoextra	R	PCA/clustering	96	0.54	101	dedicated plotting	queued	
63	ABC atlas / Allen Brain Map	Web	atlas	96	0.54	169	viewer	done	106
64	ggrepel	R	labels	92	0.52	94	dedicated plotting	done	24
65	cellxgene	Web/Python	interactive	92	0.52	162	viewer	queued	
66	NicheNet	R	ligand-target	90	0.51	122	analysis+plotting	done	108
67	UCSC Cell Browser	Web	interactive	86	0.48	122	viewer	queued	
68	ReactomePA	R	enrichment	84	0.47	87	analysis+plotting	done	110
69	scater	R	single-cell QC/embedding	83	0.47	84	analysis+plotting	queued	
70	pathview	R	pathway maps	82	0.46	99	dedicated plotting	done	112

#	PACKAGE	LANG	CATEGORY	PAPERS	%	FULL TEXT	ROLE	STATUS	PAGE
71	**GOplot**	R	enrichment	81	0.46	85	dedicated plotting	done	114
72	**variancePartition**	R	variance	77	0.43	79	analysis+plotting	queued	
73	**cell2location**	Python	spatial deconvolution	77	0.43	107	analysis+plotting	queued	
74	**scCustomize**	R	single-cell	75	0.42	78	dedicated plotting	done	116
75	**PAGA**	Python	trajectory graph	75	0.42	120	analysis+plotting	done	42
76	**RColorBrewer**	R	color palettes	71	0.4	76	dedicated plotting	queued	
77	**FactoMineR**	R	PCA	70	0.39	70	analysis+plotting	queued	
78	**shiny**	R	interactive	69	0.39	117	dashboard	queued	
79	**patchwork**	R	layout	65	0.37	71	dedicated plotting	done	118
80	**EnrichmentMap**	Desktop plugin	enrichment network	65	0.37	72	dedicated plotting	queued	
81	**Seurat spatial (SpatialDimPlot)**	R	spatial	63	0.35	65	analysis+plotting	queued	
82	**ggvenn**	R	set overlap	61	0.34	63	dedicated plotting	done	86
83	**Squidpy**	Python	spatial	60	0.34	74	analysis+plotting	done	120
84	**plotly**	R/Python/JS	interactive	56	0.32	60	dedicated plotting	done	122
85	**UMAP (umap-learn/uwot)**	Python/R	embedding	54	0.3	57	analysis+plotting	queued	
86	**ggraph**	R	network	50	0.28	49	dedicated plotting	done	124
87	**Tangram**	Python	spatial mapping	49	0.28	77	analysis+plotting	done	126
88	**PCAtools**	R	PCA	48	0.27	48	dedicated plotting	done	128
89	**viridis**	R/Python	color palettes	47	0.26	67	dedicated plotting	queued	
90	**networkx**	Python	network	47	0.26	44	analysis+plotting	queued	
91	**tradeSeq**	R	trajectory DE	46	0.26	51	analysis+plotting	done	130
92	**LIANA**	R/Python	cell-cell communication	45	0.25	67	analysis+plotting	done	132
93	**pertpy**	Python	perturbation/composition	45	0.25	53	analysis+plotting	queued	
94	**cowplot**	R	layout	44	0.25	45	dedicated plotting	done	118
95	**Morpheus**	Web	heatmap	44	0.25	102	dedicated plotting	queued	
96	**CellCall/connectome**	R	cell-cell communication	41	0.23	193	analysis+plotting	queued	
97	**BayesSpace**	R	spatial clustering	41	0.23	67	analysis+plotting	done	134
98	**CellRank**	Python	fate mapping	40	0.23	51	analysis+plotting	done	136
99	**Nebulosa**	R	single-cell density	39	0.22	43	dedicated plotting	done	138
100	**ggalluvial**	R	flow/alluvial	36	0.2	37	dedicated plotting	done	140

APPENDIX C

Full Niche & Cool table

All 73 candidates by niche-cool score. Growth = broad-corpus per-1k rate 2025–26 ÷ 2023 (+0.01 smoothing); neuro = METHODS mentions in the neuro corpus, 2023–26.

#	TOOL	LANG	CATEGORY	GROWTH	VISUAL	SCORE	NEURO	STARS	STATUS	PAGE
1	**pyCirclize**	Python	circos/chord	20.05	5	0.98	0	1,127	done	144
2	**hdWGCNA**	R	sc co-expression	6.66	5	0.938	147	494	done	92
3	**brainrender**	Python	3D brain	4.81	5	0.914	4	680	done	146
4	**COMMOT**	Python	spatial communication	4.67	5	0.904	15	149	done	148
5	**CytoTRACE2**	R/Python	potency	324.85	4	0.9	24	269	done	102
6	**MultiNicheNet**	R	ligand-target	89.9	4	0.893	13	202	done	150
7	**moscot**	Python	optimal transport	42.91	4	0.887	7	217	done	152
8	**scplotter**	R	single-cell publication plots	17.51	4	0.872	1	333	done	154
9	SCENIC+	Python	GRN	14.32	4	0.866	32	267	queued	
10	**Banksy**	R/Python	spatial clustering	14.16	4	0.859	21	167	done	156
11	**ggsankey**	R	flow	10.71	4	0.845	8	338	done	140
12	**scTour**	Python	trajectory	6.52	4	0.832	7	89	done	158
13	**plot1cell**	R	single-cell publication plots	2.7	5	0.829	6	131	done	160
14	**veloVI**	Python	RNA velocity	5.87	4	0.824	9	45	done	162
15	**miloR / Milo**	R/Python	differential abundance	2.31	5	0.815	46	450	done	164
16	**Datashader**	Python	large data	4.81	4	0.814	1	3,568	done	166
17	**scDRS**	Python	disease relevance	4.63	4	0.798	37	201	done	168
18	**BrainGlobe atlas API**	Python	3D brain	2.27	5	0.798	1	189	done	106
19	**MNE-Python**	Python	neurophysiology	2.27	5	0.798	1	3,549	done	170
20	**sccomp**	R	composition	4.27	4	0.79	10	134	done	172
21	CellOracle	Python	GRN perturbation	2.26	5	0.788	29	503	queued	
22	napari-spatialdata	Python	imaging	4.05	4	0.776	36	91	queued	
23	**LIANA+**	Python/R	cell-cell communication	3.78	4	0.77	67	–	done	132
24	cell2location	Python	spatial deconvolution	3.1	4	0.756	107	462	queued	
25	MapMyCells (ABC atlas)	Web/Python	atlas mapping	11.56	3	0.752	41	58	queued	
26	**PyComplexHeatmap**	Python	heatmap	3.06	4	0.75	2	407	done	174
27	Dynamo	Python	vector fields	2.05	5	0.746	8	512	queued	
28	**Squidpy**	Python	spatial	2.72	4	0.736	74	608	done	120
29	**nilearn**	Python	neuroimaging	1.92	5	0.733	7	1,444	done	176
30	**scCustomize**	R	single-cell publication plots	4.24	3	0.684	78	314	done	116
31	**SCpubr**	R	single-cell publication plots	2.19	4	0.671	14	245	done	178
32	**BayesSpace**	R	spatial clustering	2.19	4	0.671	67	189	done	134
33	Vitessce	JS/Python	interactive spatial	1.6	5	0.664	2	272	queued	
34	**UCell**	R	gene-set scores	3.15	3	0.663	144	213	done	180
35	**CellRank**	Python	fate mapping	1.55	5	0.658	51	–	done	136
36	CELLxGENE	Web/Python	atlas	2.8	3	0.643	162	787	queued	
37	**spatialLIBD**	R	spatial	1.97	4	0.64	73	115	done	182
38	scCODA	Python	composition	2.36	3	0.622	53	203	queued	
39	stLearn	Python	spatial	1.91	4	0.622	54	261	queued	
40	**inferCNV/CopyKAT**	R	CNV	1.91	4	0.622	237	686	done	74
41	**ggrepel volcano styles**	R	volcano	2.3	3	0.608	74	1,262	done	184
42	**Augur**	R	perturbation prioritization	1.86	4	0.606	25	153	done	186

#	TOOL	LANG	CATEGORY	GROWTH	VISUAL	SCORE	NEURO	STARS	STATUS	PAGE
43	**circlize (chord)**	R	circos/chord	1.17	5	0.602	133	1,027	done	96
44	**Tangram**	Python	spatial mapping	1.83	4	0.598	77	373	done	126
45	**scVelo streamlines**	Python	RNA velocity	1.15	5	0.596	177	513	done	84
46	**NicheNet**	R	ligand-target	1.73	4	0.592	122	682	done	108
47	**ggridges**	R	distribution	1.7	4	0.585	20	425	done	188
48	**plotnine**	Python	general	2.2	3	0.581	3	4,771	done	190
49	**trackplot**	R	genome tracks	1.68	4	0.578	1	193	queued	
50	**Palantir**	Python	trajectory	1.62	4	0.571	31	341	done	192
51	**ggspavis/SpatialExperiment**	R	spatial	2.15	3	0.56	9	5	done	194
52	**SpaGCN**	Python	spatial clustering	2.11	3	0.554	55	265	queued	
53	**UniTVelo**	Python	RNA velocity	1.44	4	0.55	7	30	queued	
54	**Signac**	R	scATAC	1.43	4	0.544	239	431	done	70
55	**plotgardener**	R	genome tracks	0.93	5	0.544	4	371	done	196
56	**plotly/Dash 3D**	Python/R	interactive 3D	1.36	4	0.537	60	18,841	done	122
57	**Giotto**	R	spatial	1.35	4	0.53	42	—	queued	
58	**Nebulosa**	R	density embedding	1.29	4	0.524	43	116	done	138
59	**ArchR**	R	scATAC	1.25	4	0.513	117	467	queued	
60	**propeller (speckle)**	R	composition	1.89	3	0.512	116	71	done	198
61	**Allen CCF overlays**	Python/MATLAB	atlas registration	1.14	4	0.486	98	126	done	106
62	**CellPhoneDB/ktplots**	Python/R	cell-cell communication	1.1	4	0.476	232	178	done	78
63	**ggcoverage**	R	genome tracks	0.97	4	0.462	3	273	done	200
64	**ggtree/ggtreeExtra**	R	tree	0.94	4	0.455	26	99	done	202
65	**Mellon**	Python	cell-state density	0.93	4	0.444	27	77	done	204
66	**karyoploteR**	R	genome	0.67	4	0.428	13	377	done	206
67	**marsilea**	Python	heatmap	0.64	4	0.421	0	324	done	208
68	**dittoSeq**	R	single-cell publication plots	1.25	3	0.413	27	216	done	210
69	**ggdensity**	R	distribution	0.51	4	0.407	1	234	done	188
70	**ggforce**	R	general	1.14	3	0.386	15	958	done	212
71	**gggenes**	R	genome	1.01	3	0.368	3	603	done	214
72	**ggalluvial**	R	flow	0.84	3	0.334	37	526	done	140
73	**Gviz**	R	genome tracks	0.55	3	0.314	31	94	done	48