ENGSCI 355 Project 1

ANDREW JACKSON LOGAN WU SCOTT SUNG

Department of Engineering Science

AUGUST 14, 2017 The University of Auckland

1 Formulation

Our model can be conceptualised as two, linked linear programs. The first is a roster of shifts, and the second one is a cyclical network representing ward occupancies during the 42 days of the roster.

1.a Parameters

 $S = \{A, P, N, Z, X, O\}$; Set of shift types $W = \{1, ..., 6\}$; Set of weeks in the roster cycle $D = \{Mon, ..., Sun\}$; Set of days in a week

Note that for continuity, a dummy week 0 and dummy day Sun_{dummy} also exist but are not part of the sets W and D.

1.b Decision Variables

X is the array of binary variables determining if a type of shift belongs to a given week and day in the roster:

$$x_{s,w,d} \in \{0,1\} \quad \forall s \in S, \ w \in W \cup \{0\}, \ d \in D \cup \{\text{Sun}_{\text{dummy}}\}$$

Y determines if a given week in the roster is the night shift week. Note the night shift also includes the final three days of the preceding week:

$$y_w \in \{0,1\} \quad \forall w \in W \cup \{\text{Sun}_{\text{dummv}}\}$$

V denotes whether registrars are forced to take a weekend off:

$$v_w \in \{0,1\} \quad \forall w \in W \cup \{\operatorname{Sun}_{\operatorname{dummy}}\}$$

1.c Constraints

Create a dummy week 0 that is equal to the final week.

$$x_{s,0,d} = x_{s|W|d} \quad \forall s \in S, \ d \in D \tag{1}$$

Create a dummy day 0 that is equal to Sunday of the previous week, allowing wrap-around from Sunday to Monday.

$$x_{s,w-1,|D|} = x_{s,w,0} \quad \forall s \in S, \ w \in W$$
 (2)

Ensure every slot has a shift assigned by summing over all shift types, except for the night-shift week which must have two.

$$\sum_{s \in S} x_{s,w,d} - y_w = 1 \quad \forall w \in W, \ d \in D$$
(3)

Every day must have a single registrar assigned to each A, P and N shift.

$$\sum_{w \in W} x_{s,w,d} = 1 \quad \forall s \in \{A, P, N\}, \ d \in D$$

$$\tag{4}$$

Every P shift must follow an A shift, except for Sunday where an A must follow.

$$x_{P,w,d} = x_{A,w,d-1} \quad \forall w \in W, \ d \in D$$
 (5)

$$x_{A,w,d} = x_{A,w,d-1} \quad \forall w \in W, \ d \in \{Sun\}$$
 (6)

Setting up the night shift: only one week can be the full 'night shift' week.

$$\sum_{w \in W} y_w = 1 \tag{7}$$

The Friday to Sunday before the full night shift week are also night shifts.

$$x_{N,w-1,d} - y_w = 0 \quad \forall w \in W, \ d \in \{Fri, Sat, Sun\}$$
 (8)

The Monday to Thursday of the full night shift week are night shifts.

$$x_{N,w,d} - y_w = 0 \quad \forall w \in W, \ d \in \{\text{Mon, Tue, Wed, Thu}\}$$
 (9)

The Friday to Sunday of the full night shift week are sleep shifts.

$$x_{Z,w,d} - y_w = 0 \quad \forall w \in W, \ d \in \{\text{Fri, Sat, Sun}\}$$
 (10)

Ensure no one else has a sleep shift and is slacking off!

$$\sum_{w \in W. \ d \in D} x_{Z,w,d} = \text{n.o. allowed rests (i.e., 3)}$$
(11)

To give people weekends off: no weekdays are allowed to be taken off:

$$x_{X w d} = 0 \quad \forall w \in W, \ d \in \{Mon, \dots, Fri\}$$
 (12)

Weekends must be taken off if scheduled as a 'weekend off' (but may be taken off on other weekends):

$$x_{X,w,d} \ge 0 \quad \forall w \in W, \ d \in \{\text{Sat}, \text{Sun}\}$$
 (13)

No two consecutive weekends can pass without a weekend off being forced.

$$v_w + v_{w-1} \ge 1 \quad \forall w \in W \tag{14}$$

1.d Objective Parameters

There are several new definitions that are particular to this section of the formulation.

 $WA = \{\text{lime, navy, yellow}\};$ Set of wards $R = W \times D;$ Set of all days in the roster, to simplify r + 1 and r - 1 subscripts $\rho = \text{Poisson constant for discharges}$

1.e Objective Variables

A is a unimodular matrix determining whether a ward is admitting on a certain day.

$$a_{wa,r} \in \{0, 1\} \quad \forall wa \in WA, r \in R$$

O is a matrix of each ward's mean expected occupancy for each day in the roster.

$$o_{wa,r} \ge 0 \quad \forall wa \in WA, \ r \in R$$

P is the mean daily admission rate for each day of the week.

$$p_d \in Z^+ \quad \forall d \in D$$

 Δ is the maximum pairwise difference between all wards for a given day of the roster.

$$\delta_r \ge 0 \quad \forall r \in R$$

1.f Objective Constraints and Objective Function

These constraints calculate the expected mean occupancy of each ward throughout the 42-day roster. % is the modulo function and $\lceil x \rceil$ rounds up to the nearest integer.

Control whether a ward is admitting on a given day of the 6-week cycle. The absolutely insane subscripts adjust for starting week offsets; i.e. on week one, day one, a registrar who started on week 6 will actually be looking at week six, day one of the roster.

$$a_{wa,(r+1)\%|W\times D|+1} = \sum_{re \in \{1...n_{\text{registrars}}\}} x_{\mathbf{A},\lceil\frac{r+|D|s_{wa},re-1}{|D|}} \mathbf{1}\%|W|+1,(r-1)\%|D|+1} \quad \forall wa \in WA, \ r \in R$$
 (15)

Calculate the expected occupancy based on the discounted occupancy from the previous day, plus the previous day's admissions if the ward was admitting. We used a mean stay of 4.5 days to model discharges as a Poisson process with a rate of 0.22.

$$o_{wa,r\%|R|+1} = o_{wa,r} \times (1-\rho) + a_{wa,r} \times \dot{p}_{(r-1)\%|D|+1} \quad \forall wa \in WA, \ r \in R$$
 (16)

For each day calculate the maximum pairwise difference, between the most and least occupied wards.

$$\delta_r \ge o_{wa,r} - o_{wb,r} \quad \forall r \in W \times D, \ wa \in WA, \ wb \in WA$$
 (17)

Minimize the maximum pairwise occupant difference, summing over all days in the roster.

$$\min \sum_{r \in R} \delta_r \tag{18}$$