Arithmetic Function Analogs for K-Tuples

Logan McBroom

Abstract

We encode the essential information of k-tuples in some extensions of common arithmetic functions. Analogs of some fundamental relationships between those common functions are shown to hold for our extensions. The basic analytic properties of some of these functions are derived.

1 K-Tuples

1.1 Basic Properties

Definition 1.1. *Define $\mathbb{K}$, the set of **k-tuples**, by*

$$\mathbb{K} := \mathscr{P}(\mathbb{Z}).$$

*We say a k-tuple is **singular** if it contains exactly one element.*

Some authors forbid infinite k-tuples. We denote the number of elements in a set S by $\#S$. The motivation for much of this paper is to extend a definition or result pertaining to integers to the k-tuples. That is, we view traditional number theoretic concepts involving an integer k as being about the singular k-tuple $\{0\}$, and attempt to extend that result in a natural way to nonsingular k-tuples. We also refer to such singular definitions and results collectively as "singular number theory".

Definition 1.2. *For $A, B \in \mathbb{K}$, we define*

$$A + B := \{a + b : a \in A, b \in B\}.$$

For $k \in \mathbb{Z}$ and $A \in \mathbb{K}$, define for convenience $k + A := \{k\} + A$. That is, $k + A$ is the translation of A by k.

We are often concerned with congruence between elements of k-tuples, but the usual definition of congruence is insufficient, as it describes only

exact incidence between numbers relative to a modulus. We are instead concerned with the contents of one k-tuple being contained in another relative to a modulus. The following definition handles the "modulus" aspect of that need, and the containment will be handled by $\subset$.

Definition 1.3. *Given a k-tuple A and a natural number n, define the* ***periodization*** *of A with respect to n by*

$$|A|_n := \{a + kn : a \in A, k \in \mathbb{Z}\}$$

Define $\mathbb{K}_p \subset \mathbb{K}$, the set of ***periodic k-tuples****, to be the set of all such periodizations.*

Note that $|A|_n$ is periodic with period n, but there may be a smaller value over which the periodization is periodic.

1.2 Natural K-Tuples

The greatest common divisor function has no obvious analog in $\mathbb{K}$. We are still interested, though, in whether each element of a k-tuple is coprime to a given natural number.

Definition 1.4. *Define $\varkappa(n) : \mathbb{N} \to \mathbb{K}_p$ by:*

$$\varkappa(n) := \{k \in \mathbb{Z} : (k, n) = 1\}.$$

If $\varkappa(n) = C$ for some n, we say that C is a ***natural k-tuple****.*

In other words, $\varkappa(n)$ is the set for which χ_0, the principle Dirichlet character mod n, is the indicator. Note that if $\varkappa(n) = C$ for some n, C is periodic with period n, though again, it may possess a smaller period than n as will be seen in Proposition 1.7. For a k-tuple A, we can then speak of each element of A being coprime to some n by saying $A \subset \varkappa(n)$.

Proposition 1.5. *For any natural numbers m and n*

$$\varkappa(mn) = \varkappa(m) \cap \varkappa(n).$$

Proof. For $m, n \in \mathbb{N}$, we have

$$\begin{aligned}
\varkappa(mn) &= \{k \in \mathbb{Z} : (k, mn) = 1\} \\
&= \{k \in \mathbb{Z} : (k, m) = 1 \text{ and } (k, n) = 1\} \\
&= \{k \in \mathbb{Z} : (k, m) = 1\} \cap \{k \in \mathbb{Z} : (k, n) = 1\} \\
&= \varkappa(m) \cap \varkappa(n).
\end{aligned}$$

□

Corollary 1.6. *For any natural numbers m and n, and any k-tuple A, we have*

$$A \subset \varkappa(mn) \Leftrightarrow A \subset \varkappa(m) \text{ and } A \subset \varkappa(n).$$

Proof. Let $m, n \in \mathbb{N}$ and let $A \in \mathbb{K}$. Then by the Proposition,

$$\begin{aligned} & A \subset \varkappa(mn) \\ \Leftrightarrow & A \subset (\varkappa(m) \cap \varkappa(n)) \\ \Leftrightarrow & A \subset \varkappa(m) \text{ and } A \subset \varkappa(n). \end{aligned}$$

□

Proposition 1.7. *For any prime p and natural number k, we have*

$$\varkappa(p^k) = \varkappa(p).$$

Proof. This is immediate from $(n, p^k) = 1 \Leftrightarrow (n, p) = 1$. □

Proposition 1.8. *For any natural number n with prime factorization*

$$n = \prod_{p_i | n} p_i^{a_i},$$

we have

$$\varkappa(n) = \varkappa\left(\prod_{p_i | n} p_i\right).$$

Proof. We have by Propositions 1.5 and 1.7,

$$\begin{aligned} \varkappa(n) &= \varkappa\left(\prod_{p_i | n} p_i^{a_i}\right) \\ &= \bigcap_{p_i | n} \varkappa(p_i^{a_i}) \\ &= \bigcap_{p_i | n} \varkappa(p_i) \\ &= \varkappa\left(\prod_{p_i | n} p_i\right). \end{aligned}$$

□

1.3 The Totient Function

We now extend Euler's totient function φ, which first requires a conceptual reinterpreting of the function. Instead of counting which natural numbers up to n are relatively prime to n, we can think of φ as the sliding of a singular k-tuple over the elements of $\varkappa(n)$ over one of its periods. The output of the function will be the number of times the singular k-tuple sat upon an element of $\varkappa(n)$. With this perspective, we may now slide a non-singular k-tuple, e.g. $A = \{0, 2\}$, in the same fashion, and count the occurences of all elements of A being over an element of $\varkappa(6)$. We graph this process below. Rather than graphing beyond 6, we rely on the periodicity of $\varkappa$ and move points beyond 6 back to 0.

0	•		•				
		•				•	
1		•		•			
		•				•	
2			•		•		
		•				•	
3				•		•	
		•				•	
4	•				•		
		•				•	
5		•				•	
		•				•	

There is only a single occurrence, on table 5, of complete containment by $\varkappa(6)$. The result is then one, and we are conceptually prepared for the following definition.

Definition 1.9 (K-Tuple Totient Function)**.**
For any k-tuple A and natural number n, define φ_A by:

$$\varphi_A(1) := 1$$
$$\varphi_A(n) := \#\{k \in \mathbb{Z} : 0 \leq k < n \text{ and } k + A \subset \varkappa(n)\}$$

Theorem 1.10. *For any k-tuple A, φ_A is a multiplicative function.*

Proof. Let A be a k-tuple and let m and n be coprime natural numbers. For each natural number a, define the set $P_A(a)$ by

$$P_A(a) = \{k \in \mathbb{N} : 0 \leq k < a \text{ and } k + A \subset \varkappa(a)\}$$

By the definition of φ_A, $\#P_A(a) = \varphi_A(a)$.

Now let $\Gamma : P_A(mn) \longmapsto P_A(m) \times P_A(n)$ be given by

$$\Gamma(a) = (a \bmod m, a \bmod n)$$

where we take $a \bmod k$ to mean the unique integer on $[0, k)$ congruent to a modulo k.

We must first show this mapping does not take an element of $P_A(mn)$ to a pair not in $P_A(m) \times P_A(n)$. Given $k \in P_A(mn)$, we have $k < mn$ and $k + A \subset \varkappa(mn)$. By Corrolary 1.5, $k + A \subset \varkappa(m)$ and $k + A \subset \varkappa(n)$. But $\varkappa(m)$ is periodic with period m, so $a \bmod m + A \subset \varkappa$. We also have $0 \leq a \bmod m < m$, so $a \bmod m \in P_A(m)$ and likewise $a \bmod n \in P_A(n)$. Thus, the mapping is well defined.

Let $a_1,\ a_2 \in P(mn)$ such that $\Gamma(a_1) = \Gamma(a_2)$. Then $a_1 \equiv a_2 \pmod{m}$ and $a_1 \equiv a_2 \pmod{n}$. Because m and n are coprime, $a_1 \equiv a_2 \pmod{mn}$. Because a_1 and a_2 are also less than mn, $a_1 = a_2$. Thus, Γ is injective.

Now let $(b, c) \in P(m) \times P(n)$. Then $b < m$, $c < n$, $b + A \subset \varkappa(m)$, and $c + A \subset \varkappa(n)$. Because m and n are coprime, there exists exactly one a satisfying $a \equiv b \pmod{m}$, $a \equiv c \pmod{n}$, and $0 \leq a < mn$ by the Chinese Remainder Theorem. Because $b + A \subset \varkappa(m)$ and $a \equiv b \pmod{m}$, $a + A \subset \varkappa(m)$, and likewise $a + A \subset \varkappa(n)$. Then by Corrolary 1.6, $a + A \subset \varkappa(mn)$, so $a \in P_A(mn)$. Thus, $\Gamma(a) = (b, c)$, so Γ is surjective.

Finally, Γ is injective and surjective, so Γ is a bijection, and

$$\begin{aligned}
\varphi_s(mn) &= \#P(mn) \\
&= \#(P(m) \times P(n)) \\
&= \#P(m) \cdot \#P(n) \\
&= \varphi_s(m) \cdot \varphi_s(n).
\end{aligned}$$

□

1.4 The Unit Function

The singular unit function $u(n)$ is defined by $u(n) = 1$ for all n. It is surprising that such a trivial function has a non-trivial extension to k-tuples.

Definition 1.11. *For any k-tuple A and prime p, define $u_A(p)$ by:*

$$u_A(p) = \#\{k \in |A|_p : 0 \leq k < p\}$$

We extend u_A to all natural numbers by complete multiplicativity.

We have immediately from complete multiplicativity that $u_A(1) = 1$. The definition for primes tells us that u_A gives the number of residues the elements of A fill modulo p. The definition of u_A on primes is used in the first Hardy and Littlewood conjecture, discussed later, and so is not entirely new. It's usage in the following theorem to describe extensions of φ is also known, at least in the well studied cases.

Theorem 1.12. *For any k-tuple A, prime p, and natural number k we have:*

$$\varphi_A(p^k) = p^k - u_A(p)p^{k-1}.$$

Proof. By definition

$$\begin{aligned} u_A(p) &= \#\{m \in |A|_p : 0 \leq m < p\} \\ \Rightarrow p - u_A(p) &= \#\{m \in \mathbb{Z} : 0 \leq m < p \text{ and } m \notin |A|_p\}. \end{aligned}$$

Now, as m takes on each residue modulo p, so does $-m$, so

$$\begin{aligned} p - u_A(p) &= \#\{m \in \mathbb{Z} : 0 \leq m < p \text{ and } -m \notin |A|_p\} \\ &= \#\{m \in \mathbb{Z} : 0 \leq m < p \text{ and } (\forall\ a \in A,\ m + a \not\equiv 0 \pmod{p})\}. \end{aligned}$$

For any m satisfying $m+a \not\equiv 0 \pmod{p}$, $m+jp$ will satisfy that condition for any integer j. So by extending the upper bound of the range condition to ap we have a times as many values of m satisfying both conditions. Using this with $a = p^{k-1}$ we have

$$p^k - u_A(p)p^{k-1} = \#\{m \in \mathbb{Z} : 0 \leq m < p^k \text{ and } (\forall\ a \in A,\ m + a \not\equiv 0 \pmod{p})\}.$$

Now, $\varkappa(p)$ contains exactly the integers on the interval $[1, p-1]$, so we may replace the latter condition to obtain

$$p^k - u_A(p)p^{k-1} = \#\{m \in \mathbb{Z} : 0 \leq m < p^k \text{ and } m + A \subset \varkappa(p)\}.$$

Then by Theorem 1.7 we have

$$\begin{aligned} p^k - u_A(p)p^{k-1} &= \#\{m \in \mathbb{Z} : 0 \leq m < p^k \text{ and } m + A \subset \varkappa(p^k)\} \\ &= \varphi_A(p^k). \end{aligned}$$

□

Theorem 1.13 (Totient Product Formula)**.**
For any k-tuple A and natural number n, we have:

$$\varphi_A(n) = n \prod_{p|n} \left(1 - \frac{u_A(p)}{p}\right).$$

Proof. Let A be a k-tuple and n be a natural number. Then n has a factorization, say

$$n = \prod_{p_i|n} p_i^{a_i}.$$

Then we have by Theorems 1.10 and 1.12,

$$\begin{aligned}
\varphi_A(n) &= \varphi_A\left(\prod_{p_i|n} p_i^{a_i}\right) \\
&= \prod_{p_i|n} \varphi_A(p_i^{a_i}) \\
&= \prod_{p_i|n} \left(p_i^{a_i} - u_A(p_i)p_i^{a_i-1}\right) \\
&= n \prod_{p_i|n} \left(1 - \frac{u_A(p_i)}{p_i}\right).
\end{aligned}$$

□

Theorem 1.14. *For any k-tuple A, the totient function has the following properties:*

(a) $\varphi_A(mn) = \varphi_A(m)\varphi_A(n)(d/\varphi_A(d))$, *where* $d = (m, n)$.

(b) $\varphi_A(mn) = \varphi_A(m)\varphi_A(n)$ *if* $(m, n) = 1$.

(c) $a|b$ *implies* $\varphi_A(a)|\varphi_A(b)$.

Proof. The proof of these follows nearly verbatim the singular case, which is shown in Theorem 2.5 of [1].

□

The following theorem shows the existence of a k-tuple with a desired unit function, along with some bounds. Uniqueness is not guaranteed, even among k-tuples containing 0. For example, the sequence $2, 1, 3$ is satisfied by the k-tuples $\{0, 12, 21\}$ and $\{0, 21, 27\}$.

Theorem 1.15. *Let p_n be the nth prime number. For any sequence a_n where $0 < a_n \leq p_n$, and any natural number N, there exists a k-tuple A such that*

1. *For all $a \in A$, $0 \leq a < p_N\#$ (the Nth primorial)*
2. *$\#A = \max_{i<N}\{a_i\} = M$.*
3. *For $n \leq N$, $u_A(p_n) = a_n$*

Proof. Let $A_1 = \{0\}$. Now for each i from 2 to M, let Q_i be the set of primes up to p_N for which u_{A_i} is smaller than needed. By the Chinese Remainder Theorem there exists a unique k with $0 \leq k < p_N\#$ that solves $k \equiv i \pmod{q}$ for all $q \in Q_i$, and $k \equiv 0 \pmod{p_i}$ for each prime $p_i \leq p_N$ which is not in Q_i. Let $A_i = A_{i-1} \cup k$.

All conditions are satisfied by A_M:

1. Each element added was on $[0, p_N\#)$.
2. Each A_i contains i elements, so $\#A_M = M$.
3. For each prime p_n up to p_N, exactly the first a_n residues have been filled by elements of A_M, that is, $u_{A_M}(p_n) = a_n$.

□

1.5 Dirichlet Products and the Möbius Function

For any arithmetic functions f and g we denote their Dirichlet product by

$$(f * g)(n) := \sum_{ab=n} f(a)g(b).$$

We denote the Dirichlet inverse of an arithmetic function f by f^{-1}. We define the arithmetic function N by $N(n) = n$.

Definition 1.16. *For any k-tuple A, define μ_A to be the unique Dirichlet inverse of u_A.*

By definition u_A is completely multiplicative, so by Theorem 2.17 in [1], we have its Dirichlet inverse to be

$$\mu_A(n) = u_A^{-1}(n) = \mu(n)u_A(n).$$

We have also from Theorem 2.18 of [1] that for any multiplicative function f and natural number n, that

$$\sum_{d|n} \mu(d) f(d) = \prod_{p|n} (1 - f(p)), \tag{1}$$

which is used to prove the following.

Theorem 1.17. *We have*

$$u_A * \varphi_A = N.$$

Proof. Applying equation (1) with $f(n) = u_A(n)/n$ to the totient product formula, we have

$$\begin{aligned}
\varphi_A(n) &= n \prod_{p|n} \left(1 - \frac{u_A(p)}{p}\right) \\
&= n \sum_{d|n} \mu(d) \frac{u_A(d)}{d} \\
&= \sum_{d|n} \mu_A(d) \frac{n}{d} \\
&= (\mu_A * N)(n).
\end{aligned}$$

Taking the Dirichlet product of both sides by u_A, we obtain the theorem. □

This result, while simple, is the reason for our remaining analysis. Intuitively, φ_A should encode information about relative primality in the k-tuple sense. The function N is neutral in this regard, so we should expect $u_A = N * \varphi_A^{-1}$ to retain some aspect of that information. As will be seen later, a normalization of u_A also allows the trivial application of many of the common tools of analytic number theory.

1.6 Admissibility

Definition 1.18. *We say a k-tuple A is* ***admissible*** *if, for each prime p, we have*

$$u_A(p) < p.$$

Then a k-tuple is admissible if it does not fill all of the residue classes modulo any prime. We are now equipped to state the prime k-tuple conjecture.

Conjecture 1.19 (Prime K-Tuple Conjecture)**.** *For any finite, admissible k-tuple A, there exists an infinite k-tuple B such that $A + B \subset \mathbb{P}$.*

Admissibility is required, for if a k-tuple A was inadmissible by prime p, for $b \in B$ large enough that each element of $b + A$ is greater than p, we must have some element of A filling the 0 residue. That element is then divisible by p, $b + A \not\subset \mathbb{P}$, and it cannot be for any b beyond that bound. Thus, there can be only finitely many integers at which $b + A \subset \mathbb{P}$ and no valid B exists.

The finiteness of A is also required. For example, let A be the set of all primorials. This fills at most i residues of each prime p_i and so is admissible. We cannot have negatives in B, for $2 \in A$ is the minimum prime. We cannot have 0 or 1 in B, because $A + B$ would contain 6 and 4 respectively. If B contains $b > 1$, b is divisible by some prime p, and the primorials will eventually be divisible by p. Then for some element $a \in A$ divisible by p, $p|a + b$, so $A + B \not\subset \mathbb{P}$. Thus, there are no integers b such that $b + A \subset \mathbb{P}$. Not only can B not be infinite in this case, it cannot contain any integer.

If the conjecture holds, there exist infinite k-tuples A with arbitrary large finite B such that $A + B \subset \mathbb{P}$. Indeed, given a finite admissible k-tuple B, we apply the conjecture and find such an A. The question of adding two infinite k-tuples and obtaining a prime k-tuple is handled in greater generality in [2], the results of which we state in less generality as the following theorem. It is often said that the Prime K-Tuple Conjecture feels intuitively true, so it is somewhat alarming that it holds as a consequence the following result, which seems intuitively impossible.

Theorem 1.20. *Assume the Prime K-Tuple Conjecture. Then there exist infinite k-tuples A and B such that $A + B \subset \mathbb{P}$.*

2 Analytic Results

The function u_A is convenient to work with in the setting of arithmetic functions in that it takes on only integer values and gives the relationship $u_A * \phi_A = N$. Transporting it to the domain of Dirichlet series, though, poses a challenge. The function $\mathcal{D}(u_A)(s) = \sum_{n \geq 1} u_A(n) n^{-s}$ permits an Euler product, and that product can be written as a power of ζ, a finite product, and the series $\mathcal{D}((\#A)^{\Omega})$, where $\Omega(n)$ is the number of primes (counting multiplicity) which compose n. While the final portion of this form has reasonable convergence properties, it will be the cause of much of the complexity in analyzing u_A. That portion, though, does not seem to have much to do with the additive patterns we seek to analyze; rather

it describes that u_A should grow approximately with the number of primes in its composition. The following adjustment, then, gives an arithmetic function which contains the same essential information, but which has far simpler analytic properties.

Definition 2.1. *Let A be a k-tuple. For each n, define $w_A(n)$ by*

$$w_A(n) = \frac{u_A(n)}{(\#A)^{\Omega(n)}}.$$

This is equivalent to defining $w_A(p) = u_A(p)/\#A$ on primes and extending it by complete multiplicativity as with u_A, but we may also show this directly: because Ω is completely additive, $(\#A)^{\Omega}$ is completely multiplicative, so w_A is as well. Note also that for primes p with $\max(A) \leq p$, $u_A(p) = \#A$, so w_A is eventually 1 on primes. We will work with both $\mathcal{D}(u_A)$ and $\mathcal{D}(w_A)$.

2.1 Zeta Function Extensions

Definition 2.2. *Let A be a k-tuple, and let $s \in \mathbb{C}$. For $\Re(s) > 1$, define*

$$\zeta_A(s) := \mathcal{D}(w_A)(s) = \sum_{n \geq 1} \frac{w_A(n)}{n^s}.$$

Clearly $\zeta_{\{0\}} = \zeta$. We have that w_A is completely multiplicative, so ζ_A permits the Euler product

$$\zeta(s) = \prod_p (1 - w_A(p)p^{-s})^{-1}.$$

But we know for sufficiently large primes, $w_A(p) = 1$, so letting S_A be the set of primes on which $w_A(p) \neq 1$, we have

$$\zeta_A(s) = \zeta(s) \prod_{p \in S_A} \frac{1 - p^{-s}}{1 - w_A(p)p^{-s}}.$$

Let $M_A(s)$ denote the finite product above and call this the modulator for A. This form immediately yields the analytic continuation of ζ_A via the continuation of ζ to the entire plane, except at the pole contributed by ζ at $s = 1$, and one infinite (but well behaved) set of poles for each $p \in S_A$. These poles are at the solutions of $0 = 1 - w_A(p)p^{-s}$, which are found, for any $k \in \mathbb{Z}$, at $s_{p,k} = \frac{\log(w_A(p)) + 2\pi k i}{\log(p)}$. We have $\mathrm{Res}(\zeta, 1) = 1$, so

$$\mathrm{Res}(\zeta_A, 1) = M_A(1) = \prod_{p \in S_A} \frac{p - 1}{p - w_A(p)}. \tag{2}$$

Proposition 2.3. *For any admissible k-tuple, we have*

$$\frac{1}{x}\sum_{n\leq x} w_A(n) \sim M_A(1).$$

Proof. Every modulator pole has real part less than one. Thus, we can apply the Wiener-Ikehara theorem (1.1 in [3]) and have

$$\frac{1}{n}\sum_{n\leq x} w_A(n) \sim \mathrm{Res}(\zeta_A, 1) = M_A(1).$$

□

To find the residues of any pole of the modulator, we first show that no two poles among those listed coincide. Clearly none of those given by the modulator coincide with $s = 1$, so we need only check that no poles among the modulator coincide. Suprisingly, though, this will walk us directly into the four exponentials conjecture.

Conjecture 2.4 (Four exponentials conjecture)**.** *For $1 \leq i, j \leq 2$, let $\lambda_{ij} \in \mathbb{C}$ such that each $exp(\lambda_{ij})$ is algebraic. Suppose that λ_{11} and λ_{12} are linearly independent, and that λ_{21} and λ_{22} are linearly independent, both in the rationals. Then*

$$\lambda_{11}\lambda_{22} \neq \lambda_{12}\lambda_{21}.$$

Lemma 2.5. *Assume the four exponentials conjecture. Then we have*

$$s_{p_1,k_1} = s_{p_2,k_2} \implies (p_1, k_1) = (p_2, k_2).$$

Proof. Assume $s_{p_1,k_1} = s_{p_2,k_2}$, and assume for contradiction $p_1 \neq p_2$ Then their real parts coincide and we have $\log a_1/\log p_1 = \log a_2/\log p_2$, which gives

$$\log a_1 \log p_2 = \log a_2 \log p_1. \tag{3}$$

Each a_i and p_i is rational and thus algebraic, so the exponential of each logarithm is algebraic.

To show the linear independence of $\log p_1$ and $\log p_2$, assume there are $r, s \in \mathbb{Q}$ such that $r \log p_1 + s \log p_2 = 0$. Then $p_1^r p_2^s = 1$, which is only possible if $r = s = 0$.

If $\log a_1$ and $\log a_2$ are dependent, then there are $r, s \in \mathbb{Q}$, not both 0, such that $r \log a_1 + s \log a_2 = 0$. Without loss of generality, assume $r \neq 0$. Then $a_1 = a_2^{-s/r}$. Applying this to equation (3), we have

$$\begin{aligned} -\frac{s}{r} \log p_2 &= \log p_1 \\ \implies p_2^{-s/r} &= p_1, \end{aligned}$$

which is a contradiction. Thus, $\log a_1$ and $\log a_2$ are linearly independent.

Finally, we may apply the four exponentials conjecture, which contradicts (3). Thus, $p_1 = p_2$. Showing $k_1 = k_2$ is trivial. □

The values of a_i in the above proof are not entirely arbitrary, coming from w_A. As such, there may be some yet undiscovered way of showing this result holds without the four exponentials conjecture. We continue to assume the conjecture as we now find the residues of the poles associated with the modulator.

Theorem 2.6.

$$Res(\zeta_A, s_{p,k}) = \frac{1 - p^{-s_{p,k}}}{\log p} \zeta(s_{p,k}) \prod_{\substack{q \in S \\ q \neq p}} \frac{1 - q^{-s_{p_k}}}{1 - w_A(q) q^{-s_{p,k}}}.$$

Proof. By the lemma, each pole of ζ_A is simple. That is, for any $s_{p,k}$ we have $\zeta_A(s) = H_{p,k}(s) \cdot g(s)$, where

$$g(s) = \frac{1}{1 - w_A(p) p^{-s}},$$

and

$$H_{p,k}(s) = \zeta(s)(1 - p^{-s}) \prod_{\substack{q \in S \\ q \neq p}} \frac{1 - q^{-s_{p_k}}}{1 - w_A(q) q^{-s_{p,k}}},$$

where $H_{p,k}$ is holomorphic at $s_{p,k}$. Now, note that by the definition of $s_{p,k}$, $1 - w_A(p) p^{-s_{p,k}} = 0$. Using this, we have

$$\begin{aligned} \mathrm{Res}((1 - w_A(p) p^{-s})^{-1}, s_{p,k}) &= \frac{1}{\frac{d}{ds}(1 - w_A(p) p^{-s})|_{s_{p,k}}} \\ &= \frac{1}{w_A(p) \log(p) p^{-s_{p,k}}} \\ &= \frac{1}{\log(p)}, \end{aligned}$$

which gives

$$\mathrm{Res}(\zeta_A, s_{p,k}) = \frac{H_{p,k}(s_{p,k})}{\log(p)}$$

as was needed. □

2.2 The First Hardy-Littlewood Conjecture

Definition 2.7. *For any admissible k-tuple A, define*

$$\pi_A(x) = \#\{n \leq x : n + A \subset \mathbb{P}\}.$$

Definition 2.8. *For $n > 0$, define*

$$\mathrm{Li}_n(x) = \int_2^\infty \frac{\mathrm{dt}}{\log^n(t)}.$$

We call this the nth (offset) logarithmic integral.

Conjecture 2.9 (First Hardy-Littlewood conjecture)**.** *Let A be an admissible k-tuple. Then*

$$\pi_A(x) \sim C_A \mathrm{Li}_{\#A}(x),$$

where

$$C_A = \prod_p \frac{1 - u_A(p)p^{-1}}{(1 - p^{-1})^{\#A}}.$$

This, of course, implies the prime k-tuple conjecture. It also implies that the exact structure of a k-tuple does not determine its asymptotic behavior, but rather that behavior is entirely decided by u_A. For example, $\{0, 2\}$ and $\{0, 4\}$ produce the same u_A. This yields an equivalence relation.

Definition 2.10. *Given k-tuples A and B, we say A and B are* ***residually equivalent*** *if we have $u_A = u_B$, and we denote this by $A \sim B$.*

The following asymptotic expansion is well known:

$$\mathrm{Li}_1(x) \sim \frac{x}{\log(x)} \sum_{k \geq 0} \frac{k!}{\log^k(x)}. \tag{4}$$

The simpler estimate $\frac{x}{\log x}$ given in the prime number theorem is then the truncation of this expansion to its first term. Then we should expect to analagously use the truncation of the asymptotic expansion of Li_n in attempting to gain traction in the first Hardy-Littlewood conjecture.

Proposition 2.11. *TODO: this result needs additional care due to the meaning of asymptotic expansion, though it is correct. We have the asymptotic expansion of* Li_n *to be*

$$\mathrm{Li}_n(x) \sim \frac{x}{(n-1)!\log x} \sum_{k \geq n-1} \frac{k!}{\log^k(x)}.$$

Proof. First, apply integration by parts to Li_n with $dv = dx/(x \log^n(x))$ and $u = x$ to obtain

$$\mathrm{Li}_n(x) = \frac{-x}{(n-1)\log^{n-1} x} + \frac{1}{n-1}\mathrm{Li}_{n-1}(x) \tag{5}$$

for $n > 1$. We have the result for $n = 1$: it is exactly equation (4). Assume then for induction the result holds for some n. Then applying equation (5) we have

$$\begin{aligned}
\mathrm{Li}_{n+1}(x) &= \frac{-x}{n\log^n x} + \frac{1}{n}\mathrm{Li}_n(x) \\
&\sim \frac{-x}{n\log^n x} + \frac{x}{n!\log x} \sum_{k \geq n-1} \frac{k!}{\log^k x} \\
&= \frac{x}{n!\log x}\left(\frac{-(n-1)!}{\log^{n-1} x} + \sum_{k \geq n-1} \frac{k!}{\log^k x}\right) \\
&= \frac{x}{n!\log x} \sum_{k \geq n} \frac{k!}{\log^k x}.
\end{aligned}$$

□

All this is to say: if the first Hardy-Littlewood conjecture is correct, we should attempt to show, following in the steps of the prime number theorem, that

$$\pi_A(x) \sim \frac{C_A x}{\log^{\#A} x}.$$

2.3 K-Tuple Constants

For each admissible constellation A, conjecture 2.9 gives an associated constant C_A. This section aims to explore these constants in our context. Just as we describe ζ_A in terms of ζ and a finite modulator, we may describe $\mathcal{D}(u_A)$ as

$$\mathcal{D}(u_A)(s) = \mathcal{D}((\#A)^{\Omega})(s) \prod_{p \in S_A} \frac{1 - \#A p^{-s}}{1 - u_A(p)p^{-s}}.$$

We then multiply and divide by $\zeta^{\#A}$ and obtain

$$\mathcal{D}(u_A)(s) = \zeta^{\#A}(s) \prod_p \frac{(1-p^{-s})^{\#A}}{1-\#Ap^{-s}} \prod_{p \in S_A} \frac{1-\#Ap^{-s}}{1-u_A(p)p^{-s}}.$$

This final step is taken because the central product now has reasonable convergence properties. We refer to this central product as $\Omega_A(s)$, and the finite product once again as the modulator, this time denoted $N_A(s)$, so that

$$\mathcal{D}(u_A)(s) = \zeta^{\#A}(s)\Omega_A(s)N_A(s) \tag{6}$$

Lemma 2.12. *For any admissible k-tuple A, $\Omega_A(s)$ converges for s with $\Re(s) > 1/2$.*

Proof. Let A be an admissible k-tuple. Fix $s = \sigma + it$ with $\sigma > 1/2$. Let

$$\Omega_{A,r} = \prod_{p>r} \frac{(1-p^{-s})^{\#A}}{1-\#Ap^{-s}}.$$

We will show the convergence of $\Omega_{A,r}$ at s for $r = todo$, from which the result follows. We have

$$\begin{aligned}\log \Omega_{A,r}(s) &= \log \prod_{p>r} \frac{(1-p^{-s})^{\#A}}{1-\#Ap^{-s}} \\ &= \sum_{p>r} \left(\#A \log(1-p^{-s}) - \log(1-\#Ap^{-s})\right).\end{aligned}$$

Expanding the logs into power series gives

$$\begin{aligned}\log \Omega_{A,r}(s) &= \sum_{p>r} \left(-\#A \sum_{k\geq 1} \frac{p^{-sk}}{k} + \sum_{k\geq 1} \frac{(\#Ap^{-s})^k}{k} \right) \\ &= \sum_{p>r} \sum_{k\geq 2} \left(\frac{(\#A)^k - \#A}{k} p^{-sk} \right).\end{aligned}$$

Note the cancellation of the $k = 1$ term. Now,

$$|\log \Omega_{A,r}(s)| \leq \sum_{p>r} \sum_{k\geq 2} \left(\#Ap^{-\sigma}\right)^k.$$

For each $p > r$ the k sum is a convergent geometric series. We have also for $p > r$ that $1 - \#Ap^{-\sigma} > 1/2$, so

$$\begin{aligned} |\log \Omega_{A,r}(s)| &= \sum_{p>r} \frac{(\#Ap^{-\sigma})^2}{1 - \#Ap^{-\sigma}} \\ &\leq 2(\#A)^2 \sum_{p>r} p^{-2\sigma} \\ &\leq 2(\#A)^2 \sum_{n>r} n^{-2\sigma}. \end{aligned}$$

We have $\sigma > 1/2$, so this is a convergent p-series. Thus, $\log \Omega_{A,r}$ converges absolutely, and exponentiating gives the result. □

We do not need the full domain of convergence of the prior result; our goal was to justify evaluating $\Omega_A(1)$ and obtaining the following.

Corollary 2.13. *For any admissible k-tuple A, we have*

$$C_A = \Omega_A(1) N_A(1),$$

and

$$Res(u_A, 1) = C_A Res(\zeta^{\#A}, 1).$$

The Laurent expansion of ζ is known, so the residues above are computable. The constants Ω_A are determined entirely by $\#A$, and thus hold no information about the structure of A. Then it seems reasonable to expect $N_A(1)$, along with $M_A(1)$, to contain some trace of the structural information. The following table gives these values for some admissible, non-residually-equivalent k-tuples, where the p column gives those primes over which the product ranges. The second data row gives these values for any odd prime p.

A	p	$M_A(1)$	$N_A(1)$
$\{0, 2\}$	2	$2/3$	$1/2$
$\{0, 2p\}$	2,p	$\frac{4(p-1)}{3(2p-1)}$	$\left(\frac{p-1}{2p}\right)^2$
$\{0, 30\}$	2,3,5	$64/135$	$4/15$
$\{0, 2, 6\}$	2,3	$18/35$	$2/9$
$\{0, 4, 10\}$	2,3,5	$2^3 3^3 / 5 \cdot 7 \cdot 13$	$2^7 / 3^3 5^2$
$\{0, 4, 28\}$	2,3,7	$2^2 3^4 / 5 \cdot 7 \cdot 19$	$2^4 3 / 5 \cdot 7^2$
$\{0, 2, 6, 8\}$	2,3,5	?	?
$\{0, 2, 6, 30\}$	2,3,5,7	?	?

3 Dump zone

3.1 Unplaced results

Theorem 3.1. *For any admissible k-tuple A, we have*

$$u_A(n) \leq \varphi(n).$$

Proof. Let $A \in \mathbb{K}$, and let n be a natural number with factorization

$$n = \prod_{p_i | n} p_i^{a_i}.$$

Then

$$\begin{aligned} u_A(n) &= u_A\left(\prod_{p_i|n} p_i^{a_i}\right) \\ &= \prod_{p_i|n} u_A(p_i)^{a_i} \\ &\leq \prod_{p_i|n} (p_i - 1)^{a_i} \\ &\leq \prod_{p_i|n} (p_i - 1) \cdot p_i^{a_i - i} \\ &= \varphi(n). \end{aligned}$$

□

TODO The proof for the asymptotic order of the sum over mangoldt times v is cool, but is there an easier way?

Definition 3.2.

$$\Psi_A(x) = \sum_{n \leq x} \Lambda(n) w_A(n).$$

Theorem 3.3.

$$\Psi_A(x) = \Psi(x) - \textit{something}$$

3.2 Notes/TODO

1. Write program to compute N and M for large data set of k-tuples. We could try to order them by N or M values; do they produce the same order?

2. Write out the proof for psiA in terms of psi, not sure if it's needed but it's cool.

3. Rewrite the stupid deleted proof of the uA L series converging for real over one half.

4. We need to understand where the HLC constants are coming from. The denominator seems to be accounting for the Omega growth, is there a way to express it in terms of w? Yes, we have $C_A = (\zeta^{\#A}/L(s, u_A))(s)$, and the L-series for u_A and v_A are related.

$$L(s, v_A) = \zeta(s)M_A(s),$$

$$L(s, u_A) = \zeta^{\#A}(s)M'_A(s)L(s, (\#A)^{\Omega}),$$

   where M'

5. Compute the ratio of the $0, 2$ and $0, 6$ HLC constants. Answer: $1/2$ (just requires cancellations) Can we get a formula for any ratio?

6. Double check Granville actually implies the thing. Oh, just noticed he is the pretentious theory guy.

7. We should check if any results require finiteness on k-tuple

8. Is there a meaning to u or v on composites outside of comp-mult? It would be more convincing to define them by a property and then prove comp-mult.

9. Where in the argument for prime number theorem iff chebyshev relation does the argument break down for our functions?

10. φ_A appears to be equivalent to a discrete convolution. Is that worth thinking about? Can we use that to improve the main proofs? What else can be done with this?

11. Scan for AB instead of A$\vee$ B - old note, what does this even mean

12. Old note - implicit conversion from int to K should probably give $\{0, k\}$, this makes products an extension. This will probably require rereading the entire document though. This breaks addition being an extension. We could add modulus and two converter ops for integers that give completely full and empty As.

References

[1] Tom M. Apostol. *Introduction to Analytic Number Theory*. Springer, 1976. ISBN: 9788184895216.

[2] Andrew Granville. *A note on the sums of primes*. Tech. rep. University of Toronto, 1990.

[3] Jaap Korevaar. "The Wiener-Ikehara Theorem by Complex Analysis". In: *Proceedings of the American Mathematical Society* 134.4 (2005).